

(19) AUSTRALIAN PATENT OFFICE

(54) Title
N-acetylcysteine compositions and methods for the treatment and prevention of drug toxicity

(51)⁶ International Patent Classification(s)

A61K 31/167	31/7072
(2006.01)	20060101ALI2005100
A61K 31/198	8BMEP A61P
(2006.01)	29/00
A61K 31/223	20060101ALI2005122
(2006.01)	0BMJP A61P
A61K 31/7072	31/04
(2006.01)	20060101ALI2005122
A61P 29/00 (2006.01)	0BMJP A61P
A61P 31/04 (2006.01)	31/12
A61P 31/12 (2006.01)	20060101ALI2005122
A61P 31/18 (2006.01)	0BMJP A61P
A61P 39/00 (2006.01)	31/18
A61P 39/02 (2006.01)	20060101ALI2005122
A61P 43/00 (2006.01)	0BMJP A61P
A61K 31/167	39/00
20060101AFI2005122	20060101ALI2005122
0BMJP A61K	0BMJP A61P
31/198	39/02
20060101ALI2005100	20060101ALI2005122
8BMEP A61K	0BMJP A61P
31/223	43/00
20060101ALI2005122	20060101ALI2005122
0BMJP A61K	0BMJP
	PCT/US02/09442

(21) Application No: 2002255942 (22) Application Date: 2002.03.28

(87) WIPO No: WO02/078627

(30) Priority Data

(31) Number	(32) Date	(33) Country
09/833,228	2001.04.11	US
60/280,600	2001.03.30	US

(43) Publication Date : 2002.10.15

(43) Publication Journal Date : 2003.04.03

(71) Applicant(s)
The Board of Trustees of the Leland Stanford Junior University

(72) Inventor(s)
De rosa, Stephen C., Andrus, James, Herzenberg, Leonard A., Herzenberg, Leonore A.

(74) Agent/Attorney
Phillips Ormonde & Fitzpatrick, Level 22 367 Collins Street, Melbourne, VIC, 3000

(56) Related Art
Gogu et al. vol 59, 1996, 1323-1329
US 5475757 A

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

(10) International Publication Number
WO 02/078627 A3

(51) International Patent Classification⁷: A61K 31/195, 31/10, 31/70

(21) International Application Number: PCT/US02/09442

(22) International Filing Date: 28 March 2002 (28.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/280,600 30 March 2001 (30.03.2001) US
09/833,228 11 April 2001 (11.04.2001) US

(71) Applicant: BIOADVANTEX PHARMA INC. [US/CA]; 6175 Kenway Drive, Mississauga Ontario L5T 2L3 (CA).

(72) Inventors: HERZENBERG, Leonard, A.; 876 Cedro Way, Stanford, CA 94503 (US). DE ROSA, Stephen, C.; 3 Pooks Hill Road, Apartment 118, Bethesda, MD 20814 (US). ANDRUS, James; 201 Bradbury Lane, Redwood City, CA 94061 (US). HERZENBERG, Leonore A.; 876 Cedro Way, Stanford, CA 94503 (US).

(74) Agent: RZUCIDLO, Eugene, C.; Greenberg Traurig LLP, 885 Third Avenue, New York, NY 10022 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
18 September 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/078627 A3

(54) Title: N-ACETYLCYSTEINE COMPOSITIONS AND METHODS FOR THE TREATMENT AND PREVENTION OF DRUG TOXICITY

(57) Abstract: The invention provides pharmaceutical compositions for the treatment or prevention of the toxic effects of therapeutic agents and methods of treating or preventing such toxicity using a toxicity reducing amount of N-acetylcysteine either alone or in combination with a therapeutically effective amount or, to achieve its therapeutic advantages, an amount larger than what is customarily given as a therapeutically effective amount, of a therapeutic agent. The invention also provides pharmaceutical compositions for the treatment or prevention of the toxic effects of therapeutic agents and methods of treating or preventing such toxicity using a toxicity reducing amount of N-acetylcysteine either alone or in combination with a therapeutically effective amount or, to achieve its therapeutic advantages, an amount larger than what is customarily given as a therapeutically effective amount, of a therapeutic agent whose side effects are made worse by increased oxidative stress or treatment related decreases in subject cysteine/glutathione levels or are otherwise relieved by administration of NAC.

**N-ACETYL CYSTEINE COMPOSITIONS AND METHODS FOR THE
TREATMENT AND PREVENTION OF DRUG TOXICITY**

5

Field of the Invention

The present invention relates to compositions for the treatment or prevention of drug toxicity in mammals including humans which may result from the administration of therapeutic agents and to methods of treating or preventing such toxicity. The compositions 10 of this invention comprise a toxicity-reducing amount of N-acetylcysteine (NAC) alone or in combination with a therapeutically effective amount or, to achieve its therapeutic advantages, an amount larger than what is customarily given as a therapeutically effective amount, of a therapeutic agent, preferably in combination with a pharmaceutically acceptable carrier. The method of treating or preventing drug toxicity in humans comprises administering a toxicity-reducing amount of NAC alone or in combination with a therapeutically effective amount or, 15 to achieve its therapeutic advantages, an amount larger than what is customarily given as a therapeutically effective amount, of a therapeutic agent, preferably in combination with a pharmaceutically acceptable carrier.

GOVERNMENT SUPPORT

20 This work is supported at least in part by grants from the N.I.H. CA42509-14. The government may have certain rights in this invention.

BACKGROUND OF THE INVENTION

Glutathione (GSH), a tripeptide that is normally found in all animal cells and most 25 plants and bacteria at relatively high (1-10 millimolar) concentrations, helps to protect cells against oxidative damage that would otherwise be caused by free radicals and reactive oxidative intermediates (ROIs) produced during cell metabolism or as the results of, for example, drug overdose. Glutathione is itself the major scavenger of reactive oxidative intermediates present in all eukaryotic forms of life and is generally required to protect cells against damage by oxidants. Glutathione reduces (and thereby detoxifies) intracellular 30 oxidants and is consumed by this reaction. Glutathione is oxidized to the disulfide linked dimer (GSSG), which is actively pumped out of cells and becomes largely unavailable for reconversion to reduced glutathione. Thus, unless glutathione is resynthesized through other pathways, utilization of this compound is associated with a reduction in the amount of glutathione available. The antioxidant effects of glutathione are also mediated less directly by 35 the role of this compound in maintaining other antioxidants in reduced form. Thus,

5 pharmaceutical compounds that replenish or elevate glutathione levels work, at least in part, through enhancement of the defense mechanisms seemingly utilized to normally protect tissue from ROI mediated damage.

Glutathione depletion has been implicated in the pathology of a number of diseases including infection by human immunodeficiency virus (HIV). In HIV infection, 10 cysteine/glutathione depletion is known to impair T-cell function and is associated with impaired survival of subjects with less than 200 CD4 T-cells per μ l blood.

Drug toxicity is a very widespread problem. Cysteine/glutathione depletion and oxidative stress (See U.S. Patent No. 4,757,063) intensify drug toxicity effects and have been implicated in the mechanism of drug toxicity reactions.

15 For example, acetaminophen is known to act to deplete cysteine/glutathione and cause a variety of drug toxicity symptoms. Acetaminophen, also known as paracetamol and N-acetyl-p-aminophenol, is one of the most widely used pharmaceutical analgesic and antipyretic agents in the world. It is contained in over 100 products and is commonly found in the U.S. as immediate release tablets and as extended-release preparations. Various children's 20 chewable, suspension, and elixir formulations that contain acetaminophen are prevalent. Acetaminophen is also found as a component of combination drugs, such as propoxyphene/acetaminophen and oxycodone acetaminophen.

Acetaminophen continues to be the most commonly encountered substance in toxic ingestions. In many cases, acetaminophen overdoses are unintentional and are undiagnosed 25 until after substantial damage has already occurred. Repeated administration of acceptable size doses of acetaminophen can produce toxicity symptoms. As discussed by Donovan (1999) Academic Emergency Med. 6:1079-1082, methods for detecting post-ingestion blood levels of acetaminophen suffer from poor predictive values. Even in the simple case of a single acute ingestion, patients with no discernible risk factors for liver injury and low blood 30 levels of acetaminophen still develop toxicity and even die.

Many companies package acetaminophen under different trade names, resulting in inadvertent overdosing by less sophisticated patients and parents who do not read the information on the packaging. In addition, cold remedies and other over-the-counter preparations often contain acetaminophen, which is listed among a series of generic drug

5 names that are difficult for patients and parents to read. Therefore, patients often are unaware
of the amount of acetaminophen that they have received. Children are especially vulnerable
to accidental exposure due to their smaller size, the presence of acetaminophen in multiple
over-the-counter remedies, and a reluctance to administer aspirin and other NSAIDs to
children for fever due to the risk of Reye's Syndrome and renal tubular injury. The
10 antipyretic value of acetaminophen clearly has been demonstrated and hence acetaminophen
is widely used in hospitals for this purpose. However, acetaminophen may not be the
antipyretic agent of choice under circumstances where renal or hepatic function is in danger
of being compromised.

It is well established that large acetaminophen overdose causes hepatotoxicity and, in
15 some cases, nephrotoxicity in humans and in experimental animals. Acute overdosage of
acetaminophen results in dose-dependent and potentially fatal hepatic necrosis as well as (in
rare cases) renal tubular necrosis and hypoglycemia. Acetaminophen is rapidly absorbed from
the stomach and small intestine and is normally metabolized by conjugation in the liver to
nontoxic agents, which are then eliminated in the urine. In acute overdoses, or when
20 maximum daily doses are exceeded over a prolonged period, the normal pathways of
metabolism become saturated.

Excess acetaminophen is metabolized in the liver via the mixed function oxidase P450
system to a toxic, N-acetyl-p-benzoquinone-Imine (NAPQI). NAPQI has an extremely short
half-life and is rapidly conjugated with glutathione, a sulfhydryl donor, and removed from the
25 system. Under conditions of excessive NAPQI formation or reduced glutathione stores,
NAPQI is free to bind to vital proteins and the lipid bilayer of hepatocytes. This results in
hepatocellular death and subsequent centrilobular liver necrosis. Immunohistochemical
studies have suggested that NAPQI-protein adducts appear even at sub-hepatotoxic
30 acetaminophen doses and before depletion of total hepatic glutathione which may be related
to rare cases of hypersensitivity. In addition, decreased intracellular cysteine/glutathione can
contribute to cell death via mechanisms that do not involve NAPQI.

The direct cost of acetaminophen overdose has been estimated to be \$87 million
annually. Effective protocols have been developed and tested to stratify risk and treat patients
who present soon after a single large dose of acetaminophen. However, many patients present
35 after a delay long enough to metabolize all the acetaminophen, after two or more ingestions

5 over several hours, or after several days of excessive self-medication. Under these circumstances it is difficult for the clinician to estimate the risk of adverse outcome before hepatic or renal injury occurs. See, for example, Bond and Hite (1999) Acad. Emerg. Med. 6:1115-1120; and Donovan (1999) Acad. Emerg. Med. 6:1079-1082. However, early treatment of acetaminophen overdosage is considered to be crucial, and vigorous supportive 10 therapy is essential when intoxication is severe.

Nucleoside reverse transcriptase inhibitors (Notes), of which the pyrimidine nucleoside analogue azidothymidine (AZT, zidovudine), is a common example, are often given in combination therapies with other anti-retroviral drugs to treat HIV. Long-term therapy with AZT is commonly associated with dose-dependent hematologic toxicity which 15 manifests as low erythrocyte counts and elevated mean red cell volume, and with muscle fiber toxicity, particularly in patients with advanced HIV disease. Some studies indicate that AZT's toxic interactions result from the generation of reactive oxygen species (ROIs) that react with and deplete intracellular glutathione levels. See de la Asuncion, et al (1998) J. Clin. Invest. 102(1): 4-9; Gogu et al. (1991) Exp. Hematol. 19(7): 649-652; Gogu and 20 Agrawal (1996) Life Sci. 59 (16): 1323-1329; Prakash et al (1997) Arch. Biochem. Biophys. 343 (2): 173-80.

Results have shown that acetaminophen usage, which lowers glutathione levels 25 exacerbates AZT toxicity. Richman, et al. (1987) N. Eng. J. Med. 317: 192-97. De Rosa et al. recently have shown that treatment with NAC which increases glutathione levels decreases the toxicity. De Rosa et al., submitted to JA MA for publication.

Evidence from in vitro and animal studies supports this conclusion. AZT treatment 30 caused oxidative damage to mitochondrial DNA (including increased mitochondrial lipoperoxidation) and increased levels of oxidized glutathione in skeletal muscle in mice. See de la Asuncion, et al. (1998) J. Clin. Invest. 102(1): 4-9. NAC and the anti-oxidant Vitamins C and E have been shown to prevent this AZT-induced toxicity. See; Gogu, et al (1991) Exp. Hematol. 19, 649-52; and Gogu and Agrawal (1996) Life Chem. Rep. 4, 1-35. Furthermore, 35 AZT treatment intensifies glutathione depletion in HIV-TAT transgenic mice (see Prakash, 0., et al. Arch. Biochem. Biophys. (1997) 343: 173-80) where expression of the TAT protein has been shown to deplete glutathione by decreasing glutathione biosynthesis (see Choi, J., et al., (2000) J. Biol. Chem. 275 (5): 3693-98) and the activity of antioxidant enzymes (Flores et

5 al. (1993) Proc. Nat. Acad. Sci. 90 (16): 7632-36). Studies with TAT-transgenic mice also show that AZT toxicity is enhanced in this glutathione-depleted environment. See Prakash et al. (1997) Arch. Biochem. Biophys. 343, 173-80.

10 Early clinical trials of AZT efficacy in HIV disease revealed an association between AZT toxicity and the use of acetaminophen. See, e.g., Richrnan, et al. (1987) New Eng. J. Med. 317(4): 192-97. Although the mechanism of this toxic reaction is not fully understood, acetaminophen does not impair AZT detoxification since no difference in AZT's rate of destruction has been observed. Since acetaminophen is known to deplete glutathione, the potentially harmful effect of co-administering acetaminophen and AZT appears to be mediated through glutathione depletion. Thus, in conditions where glutathione already is 15 depleted, such as in later stages of HIV disease, detoxification of acetaminophen (which can be expected to further deplete glutathione stores in the liver and elsewhere) would increase the potential for AZT toxicity.

20 Long-term antibiotic usage also often produces drug toxicity reactions. Toxicity reactions for a given antibiotic are a function of its mechanism of action and the pathway(s) by which it is metabolically degraded.

25 Scientists have long sought to identify agents that will be generally effective in combating drug toxicity reactions. Protective agents for drug overdose have been extensively studied. A known method of treatment for acetaminophen overdose is the administration of sulphydryl compounds. L-methionine, L-cysteine, and either the purified L-enantiomer or the racemate mixture of N-acetylcysteine are known to have a protective action in animals. 30 Methionine and another sulphydryl compound, cysteamine, have been reported to provide some protection. Also, cimetidine, dimethyl sulfoxide, and ethanol have been shown to inhibit acetaminophen bioactivation. N-acetylcysteine has been shown to be effective in humans when given orally. Early administration of compounds supplying sulphydryl groups 35 (0 to 10 hours after acetaminophen ingestion) may prevent or minimize hepatic and renal injury in cases of acetaminophen overdose. NAC is now used by many physicians for treatment of hepatic failure of any etiology, whether known or unknown, and is the accepted antidote for cyclophosphamide poisoning. NAC also is used to prevent toxicity due to radiation therapy contrast material in patients undergoing such treatment. The mechanisms through which NAC prevents or reverses toxicity are mainly thought to involve glutathione

2002255942 23 Oct 2006

replenishment. However, additional mechanisms through which NAC works directly via the cysteine molecule itself are not excluded.

Recently we have shown that treatment with NAC decreases AZT's hematologic toxicity in 5 subjects taking AZT. It appears that NAC provides cysteine needed to redress the excessive sulfur loss that occurs in HIV disease and specifically to replenish intracellular glutathione. This in turn helps to restore the reducing power necessary for deoxynucleotide synthesis and to bring the size of the deoxynucleotide pool, and hence the rate of cell division, back to normal range. This decreases the AZT-mediated inhibition of erythroid development, helps to improve 10 the overall metabolism and stability of erythrocytes and their progenitors (for example, by enabling optimal functioning of the glucose-6-phosphate dehydrogenase and other energy-supplying pathways). In addition, it improves the cell's ability to withstand the production of oxidants induced by the introduction of drugs (such as AZT) and the internal production of molecules (such as TNF and HIV-TAT) that trigger intracellular oxidant production.

15

Improved formulations and methods to prevent drug toxicity reactions during long term therapy are of particular interest in view of the considerable loss of life attributable to, and cost of treatment of, such reactions. The present invention addresses this problem.

20

The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

25

Throughout the description and the claims of this specification the word "comprise" and variations of the word, such as "comprising" and "comprises" is not intended to exclude other additives, components, integers or steps.

30

SUMMARY OF THE INVENTION

Pharmaceutical compositions and methods of treatment are provided for serial or co-administration of NAC with any therapeutic agent to decrease drug toxicity, including those drugs whose side effects are made worse by decreases in subject intracellular 35 cysteine/glutathione levels or increased oxidative stress, or whose side effects are otherwise relieved by administration of NAC. The serial or co-administration of NAC makes a

therapeutic agent safer at currently recognised standard therapeutic doses and allows increased doses of that therapeutic agent to be administered for its beneficial purposes without toxic side effects.

5

According to one embodiment of the invention, there is provided a pharmaceutical composition useful for treating or preventing the toxic effects of therapeutic agents in mammals; comprising:

(a) a toxicity-reducing amount of N-acetylcysteine, a pharmaceutically acceptable salt of N-acetylcysteine, or a pharmaceutically acceptable derivative of N-acetylcysteine in combination with a pharmaceutically acceptable excipient or carrier,

10 wherein a unit dose of N-acetylcysteine comprises at least 1.5 mg N-acetylcysteine per kg of body weight and not more than 2000 mg N-acetylcysteine, and

(b) a therapeutic or greater amount of a therapeutic agent that produces oxidative stress;

15 wherein the composition does not comprise diallyl sulfide, diallyl sulfone or diallyl sulfoxide.

According to another embodiment of the invention, there is provided a method of treating or preventing the toxic effect of therapeutic agents in mammals, the method comprising the step of administering the composition comprising:

20 (a) a toxicity-reducing amount of N-acetylcysteine, a pharmaceutically acceptable salt of N-acetylcysteine, or a pharmaceutically acceptable derivative of N-acetylcysteine in combination with a pharmaceutically acceptable excipient or carrier,

25 wherein a unit dose of N-acetylcysteine comprises at least 1.5 mg N-acetylcysteine per kg of body weight and not more than about 2000 mg N-acetylcysteine; and

(b) a therapeutic or greater amount of a therapeutic agent that produces oxidative stress;

wherein the composition does not comprise diallyl sulfide, diallyl sulfone or diallyl sulfoxide.

30

5 therapeutic agent, in combination with a pharmaceutically acceptable excipient or carrier. In another embodiment, such compositions comprise at least about 1 mg N-acetylcysteine or a pharmaceutically-acceptable salt or derivative thereof. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof,
10 wherein the therapeutic agent is at least one antibiotic or antiviral agent. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof wherein the therapeutic agent is at least one antibiotic or antiviral agent that produces oxidative stress. In another embodiment, such compositions comprise a
15 therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the therapeutic agent is at least one antibiotic or antiviral agent that produces treatment-related decreases in subject cysteine/glutathione levels. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of
20 NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the therapeutic agent is an analgesic, antipyretic or other therapeutic compound. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the therapeutic agent is an analgesic, antipyretic or other therapeutic
25 compound that produces oxidative stress. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the therapeutic agent is an analgesic, antipyretic or other therapeutic compound that produces treatment-related decreases in subject cysteine/glutathione levels. In another embodiment,
30 such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the compound is a paracetamol derivative, which includes phenacetin and its active metabolite acetaminophen. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent in combination with a toxicity-reducing
35 amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein each dosage unit contains a standard dose of acetaminophen and at least about 1 mg of NAC. In another embodiment, such compositions comprise a therapeutic amount of a therapeutic agent

5 in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the therapeutic agent is an anti-retroviral agent. In another embodiment, such compositions comprise a therapeutically effective amount of AZT in combination with a toxicity-reducing amount of NAC to decrease key aspects of AZT toxicity in human HIV patients, e.g., macrocytic anemia.

10 In another embodiment, a method of treating or preventing the toxic effects of therapeutic agents in mammals comprises administering a toxicity reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, alone or in combination with a normal dosage or a greater than normal dosage of a therapeutic agent, in combination with a pharmaceutically acceptable carrier. In another embodiment, the method comprises 15 administering at least about 1 mg NAC or a pharmaceutically acceptable salt or derivative thereof to a patient receiving nutrition parenterally. In another embodiment, the method comprises administering a therapeutic amount of at least one antibiotic or antiviral agent in combination with NAC or a pharmaceutically acceptable salt or derivative thereof. In another embodiment, the method comprises administering a therapeutic amount of at least one 20 antibiotic or antiviral agent in combination with NAC or a pharmaceutically acceptable salt or derivative thereof wherein the antibiotic or antiviral agent produces oxidative stress. In another embodiment, the method comprises administering a therapeutic amount of at least one antibiotic or antiviral agent in combination with a toxicity-reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof, wherein the antibiotic or antiviral 25 agent produces treatment-related decreases in subject cysteine/glutathione levels. In another embodiment, the method comprises administering a therapeutic amount of an anti-retroviral agent in combination with a toxicity reducing amount of NAC. In another embodiment, the method comprises administering a therapeutic amount of an anti-retroviral agent in combination with a toxicity-reducing amount of NAC wherein the antiretroviral agent is 30 AZT. In another embodiment, the method comprises administering a therapeutic amount of AZT serially or in combination with a toxicity-reducing amount of NAC perinatally. In another embodiment, the method comprises administering a therapeutic amount of AZT serially or in combination with a toxicity-reducing amount of NAC neonatally. In another embodiment, the method comprises administering a therapeutic amount of AZT for suspected 35 or known exposure to the HW virus serially or in combination with a toxicity-reducing amount of NAC. In another embodiment, the method comprises administering a therapeutic

5 amount of at least one analgesic, antipyretic or other therapeutic compound in combination with a toxicity reducing amount of NAC or a pharmaceutically acceptable salt or derivative thereof. In another embodiment, the method comprises administering a therapeutic amount of at least one analgesic, antipyretic or other therapeutic compound in combination with a toxicity-reducing amount of NAC, wherein the compound produces oxidative stress. In
10 another embodiment, the method comprises administering a therapeutic amount of at least one analgesic, antipyretic or other therapeutic compound in combination with a toxicity-reducing amount of NAC, wherein the compound produces treatment-related decreases in subject cysteine/glutathione levels. In another embodiment, the method comprises administering a therapeutic amount of at least one analgesic, antipyretic or other therapeutic
15 compound in combination with a toxicity-reducing amount of NAC wherein the compound is a paraminophenol derivative, which includes phenacetin and its active metabolite acetaminophen. In another embodiment, the method comprises administering a therapeutic amount of acetaminophen in a standard dosage serially or in combination with at least about 1 mg per dose N-acetylcysteine to treat a fever or pain in a patient who responds normally to
20 acetaminophen. In another embodiment, the method comprises administering a greater than standard dosage of acetaminophen serially or in combination with at least about 4 mg/kg N-acetylcysteine to treat a fever nonresponsive to the standard dose of acetaminophen. In another embodiment, the method comprises administering a standard dosage of acetaminophen serially or in combination with at least about 1 mg per dose N-acetylcysteine
25 to treat patients who require acetaminophen during their exposure to radiation contrast agents.

The combined compositions and methods provide protection against toxic drug reactions, particularly where overdosing may be inadvertent, undiagnosable, or even therapeutically desirable if the toxic side effects could be removed.

30

DETAILED DESCRIPTION OF THE INVENTION

The invention described herein provides for the serial or co- administration of any therapeutically active agent with NAC to relieve the toxic effects of such therapy in mammals including where toxic effects of such therapy in mammals may be due to oxidative stress or treatment-related decreases in subject cysteine/glutathione levels or are otherwise relieved by
35 administration of NAC.

5 The pharmaceutical compositions and methods of treatment above described include
providing the pharmaceutical compositions in oral, parenteral or suppository form for oral,
parenteral or rectal administration. It is preferred that the NAC be substantially free of
sulfones or other chemicals that interfere with the metabolism of the co-administered drug,
e.g., acetaminophen, in its bioactive form. It is also preferred that the NAC be substantially
10 free of its oxidized form, di-n-acetylcysteine. It is preferred that the therapeutic agent, serially
or co-administered, be in any form in which it is typically available and the composition
should be prepared in a manner that substantially prevents oxidation of the NAC during
preparation or storage.

15 In one aspect of the invention, the combined formulation is administered to
individuals having an increased susceptibility for acetaminophen toxicity. With regard to
acetaminophen toxicity in adults, hepatotoxicity may occur after ingestion of a single dose of
more than about 7.5 to 10 g of acetaminophen. However, alcoholics and individuals taking
isoniazid have elevated P-450 2E1 levels and can have increased susceptibility for
acetaminophen toxicity. The risk of hepatotoxicity with single or serial doses of
20 acetaminophen may also be increased in patients regularly taking other hepatic enzyme-
inducing agents, especially barbiturates or other anticonvulsants. In addition, individuals
taking classes of antibiotics that reduce GSH, including sulfas, chloramphenicol, macrolides
and fluoroquinolones, may have increased risk of hepatotoxicity with acetaminophen. Also,
although most acetaminophen is metabolized through the glucuronidation pathway in adults,
25 some is converted to a reactive intermediate that is detoxified in a GSH-dependent reaction.
See Thomas, S.H. (1993) *Pharmacol. Ther.* 60: 91-120. Pre-adolescent children, particularly
below school age, lack the normal glucuronidation pathway and rely to a greater degree than
adults on GSH for detoxification of acetaminophen, increasing their susceptibility to
acetaminophen-induced toxicity. Further, patients receiving parenteral nutrition who may not
30 receive an adequate dietary source of cysteine or whose condition may result in oxidative
stress will have an increased susceptibility to acetaminophen toxicity. Increased susceptibility
can also result from a predisposition to GSH deficiency, due to HIV and other infections,
metabolic diseases, e.g., diabetes, chronic diseases, sepsis, hepatic insufficiency and other
physiologic oxidative stress.

35 With respect to acetaminophen, the formulations of the present invention find use as
anti-pyretic and analgesic agents, and are suitable for medical indications treatable with

5 acetaminophen alone. An improvement in product safety is provided by the inclusion of NAC, which substantially prevents' the possibility of accidental or inadvertent overdosage. The combined use of acetaminophen and NAC enhances the desired effects of acetaminophen while preventing its side effects, e.g. administering acetaminophen with NAC can allow acetaminophen dosing at higher levels, which would otherwise carry an unacceptable risk of

10 toxicity. The co-formulation with NAC can enhance the safety of acetaminophen administration in severe illness and hence decrease patient morbidity and mortality due to acetaminophen-induced hepatic and renal injury. NAC administration may provide an additional benefit since it tends to reduce the negative effects of tumor necrosis factor alpha (TNF- α) and similar-acting cytokines.

15 NAC replenishes glutathione following acetaminophen overdose, which otherwise leads to a fatal depletion of glutathione in the liver. This non-toxic drug enters cells readily and replenishes the intracellular cysteine required to produce glutathione, thus leading to an increase in glutathione levels. It is important to note that NAC does not interfere with the peripheral action of acetaminophen as an analgesic or anti-pyretic, and thus its presence in

20 the combined formulations of the invention does not decrease the potency of the therapeutic agent. It may be noted that the effectiveness of NAC depends on the presence of the reduced form, which may, for example, liberate the reduced form of glutathione from homo- and hetero-disulfide derivatives in thiol-disulfide exchange reactions. The unit dose of acetaminophen in the provided formulation may be higher than the conventional dose in the

25 absence of NAC. A typical unit dosage may be a solution suitable for oral or intravenous administration; an effervescent tablet suitable for dissolving in water, fruit juice, or carbonated beverage and administered orally; a tablet taken from two to six times daily, or one time-release capsule or tablet taken once or twice a day and containing a proportionally higher content of active ingredient, etc. The time-release effect may be obtained by capsule

30 materials that dissolve at different pH values, by capsules that release slowly by osmotic pressure, or by any other known means of controlled release. Unit dosage forms may be provided wherein each dosage unit, for example, teaspoonful, tablespoonful, gel capsule, tablet or suppository, contains a predetermined amount of the compositions of the present invention. Similarly, unit dosage forms for injection or intravenous administration may

35 comprise the compound of the present invention in a composition as a solution in sterile water, normal saline or another pharmaceutically acceptable carrier. The specifications for the

5 unit dosage forms of the present invention depend on the effect to be achieved and the intended recipient.

It is preferred that the pharmaceutical compositions according to the present invention contained from about 80 mg to about 2000 mg of acetaminophen per dosage unit, particularly from about 650 mg to about 2000 mg per dosage unit. The amount of NAC per dosage unit is 10 preferably from 1 mg to 25000 mg, preferably at least 3 mg to 2,000 mg per dosage unit for oral administration, and 20-20,000 mg for parenteral.

The acetaminophen present in orally administrable solid unit doses will usually be at least about 80 mg (for pediatric doses), 325 mg, 500 mg and 650 mg, and may be as high as about 2000 mg, more usually not more than about 1500 mg together with effective amounts 15 of NAC. Suppositories are formulated in the manner well known in the art and usually comprise at least about 120 mg, 125 mg, 325 mg, 500 mg and 650 mg acetaminophen per dosage unit and may be as high as about 2000 mg, more usually not more than about 1500 mg together with effective amounts of NAC. Oral liquid dosage forms usually comprise at least about 100 mg/ml, 120 mg/2.5 ml, 120 mg/S ml, 160 mg/S ml, 165 mg/S ml, 325 mg/S 20 ml acetaminophen, and may be as high as about 2000 mg, more usually not more than about 1500 mg together with effective amounts of NAC.

The unit dose of NAC, in combination with any one of the above doses of acetaminophen, or alone for the treatment of acute hepatic injury in the absence of toxic levels of acetaminophen, will usually comprise at least about 1.5 mg/kg to a maximum 25 amount of 70 mg/kg (for pediatric doses), usually at least about 500 mg (for adult doses); and usually not more than about 2,000 mg at the physician's discretion. Patients on therapy known to deplete cysteine/glutathione or produce oxidative stress may benefit from higher amounts of NAC.

Over-the-counter NAC can be variably produced and packaged. Because the 30 production and packaging methods generally do not guard against oxidation, the NAC can be significantly contaminated with bioactive oxidation products. These may be particularly important in view of data indicating that the oxidized form of NAC has effects counter to those reported for NAC and is bioactive at doses roughly 10-100 fold less than NAC (see Sarnstrand et al (1999) J. Pharmacol. Exp. Ther. 288: 1174-84).

5 The distribution of the oxidation states of NAC as a thiol and disulfide depends on the
oxidation/reduction potential. The half-cell potential obtained for the NAC thiol/disulfide pair
is about +63 mV, indicative of its strong reducing activity among natural compounds (see
Noszal et al. (2000) J. Med. Chem. 43:2176-2182). In a preferred embodiment of the
10 invention, the preparation and storage of the formulation is performed in such a way that the
reduced form of NAC is the primary form administered to the patient. Maintaining NAC
containing formulations in solid form is preferable for this purpose. When in solution, NAC
containing formulations are preferably stored in a brown bottle that is vacuum sealed. Storage
in cool dark environments is also preferred.

15 The determination of reduced and oxidized species present in a sample may be
determined by various methods known in the art, for example with capillary electrophoresis,
HPLC, etc. as described by Chassaing et al. (1999) J Chromatogr B Biomed Sci Appl
735(2):219-27.

20 The compositions of the present invention may be administered orally, parenterally,
or rectally in dosage unit formulations containing conventional nontoxic pharmaceutically
acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein
includes subcutaneous injections, intravenous, intramuscular, intrastemal injection, or
infusion techniques. Topical administration may also involve the use of transdermal
administration such as transdermal patches or iontophoresis devices which are prepared
according to techniques and procedures well known in the art.

25 Solid dosage forms for oral administration may include capsules, tablets, pills,
powders, granules and gels. In such solid dosage forms, the active compounds may be
admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms
may also comprise, as in normal practice, additional substances other than inert diluents, e.g.,
lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the
30 dosage forms may also comprise buffering agents. Tablets and pills can additionally be
prepared with enteric coatings.

35 Suppositories for rectal administration of the drug composition, such as for treating
pediatric fever etc., can be prepared by mixing the drug with a suitable nonirritating excipient
such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but
liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

5 Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions, may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1, 3-butanediol. Among the acceptable vehicles and solvents that
10 may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

The compositions of this invention can further include conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral application which do not deleteriously react with the active compounds. Suitable
15 pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil; fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethylcellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, e.g.,
20 lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds. For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. Aqueous suspensions may contain substances which
25 increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol and/or dextran. Optionally, the suspension may also contain stabilizers.

The composition, if desired, can also contain minor amounts of wetting or emulsifying agents or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The
30 composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc.

The therapeutically active agent of the present invention can be formulated per se or
35 in salt form. Pharmaceutically acceptable salts include, but are not limited to, those formed

5 with free amino groups such as those derived from hydrochloric, phosphoric, sulfuric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of compounds in the compositions of the present invention which will be
10 effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. See, for example, Goodman and Gilman; The Physician's Desk Reference, Medical Economics Company, Inc., Oradell, N.J., 1995; and to Drug Facts and Comparisons, Facts and Comparisons, Inc., St. Louis, Mo., 1993. As the addition of NAC does not affect the
15 therapeutic efficacy of acetaminophen, it is generally not necessary to adjust the dosage from what would ordinarily be administered for acetaminophen alone, and in fact the dose may be raised due to the increased safety of the present formulations. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the
20 practitioner and each patient's circumstances.

The present invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of
25 pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

In another embodiment of the invention, formulations of NAC are provided for the treatment of acute hepatic failure of unknown etiology, for example for the treatment of acetaminophen or other drug toxicity where the serum levels of the toxic drug indicate non-toxic levels where there may be an increased risk for toxicity due to oxidative stress or other aspects of the patient's condition. The NAC formulations for treatment of such patients will utilize a formulation as described above.

Many such patients having severe hepatic failure also have serum levels of acetaminophen within the accepted non-toxic range, due to various factors. Such factors may

5 include the lapsed time before presentation of the toxic effects, for example in incidences of
intentional overdose.

Inadvertent acetaminophen toxicity has also been reported with normally accepted
dosing regimens when associated with certain viral infections, toxic exposure, and drug use,
such as sulfa, chloramphenicol and erythromycin. Many of these patients present with non-
10 toxic acetaminophen levels, but toxicity still results from oxidative stress and other drug-
related effects on hepatocytes.

Another group of patients are under oxidative stress or otherwise have severe illness
that makes them more susceptible than normal to acetaminophen toxicity. Such patients may
15 include, for example, patients with septic shock, distributive shock, hemorrhagic shock, acute
respiratory distress syndrome, organ failure, and closed head injury. Patients in this group are
routinely treated with acetaminophen as an anti-pyretic and mild analgesic, which may
inadvertently result in severe hepatic or renal damage.

Where a range of values is provided, it is understood that each intervening value, to
the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between
20 the upper and lower limit of that range and any other stated or intervening value in that stated
range is encompassed within the invention. The upper and lower limits of these smaller
ranges which may independently be included in the smaller ranges is also encompassed
within the invention, subject to any specifically excluded limit in the stated range. Where the
stated range includes one or both of the limits, ranges excluding either both of those included
25 limits are also included in the invention.

Unless defined otherwise, all technical and scientific terms used herein have the same
meaning as commonly understood by one of ordinary skill in the art to which this invention
belongs. Although any methods and materials similar or equivalent to those described herein
30 can also be used in the practice or testing of the present invention, the preferred methods and
materials are now described. All publications mentioned herein are incorporated herein by
reference to disclose 'and describe the methods and/or materials in connection with which the
publications are cited.

5 It must be noted that as used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. All technical and scientific terms used herein have the same meaning.

10 The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

EXAMPLES

15 The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. We present AZT and acetaminophen as examples of the model for toxicity we propose. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, 20 temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1

25 A child or adult having a fever can be treated with a formulation comprising 15 mg/kg acetaminophen and 1 mg/kg of either the purified L-enantiomer or the racemate mixture composed of equal proportions of the D- and L- isomers of NAC administered either serially or co-administered three or four times a day up to the highest tolerable dose, given that there will be individual variability in the ability to tolerate NAC. The dose of NAC may vary from 30 0.03 to four times the amount of acetaminophen taken.

Example 2

A child or adult having a fever that does not respond to acetaminophen at a standard dose level can be treated with a formulation comprising 15-50 mg/kg acetaminophen and 3-130 mg/kg of either the purified L-enantiomer or the racemate mixture composed of equal

5 proportions of the D- and L- isomers of NAC administered either serially or co-administered
three or four times a day up to the highest tolerable dose, given that there will be individual
variability in the ability to tolerate NAC.

Example 3

A normally healthy child or adult having a low fever can be treated for several days at
10 normal acetaminophen dosage levels with a formulation comprising 80 mg per tablet
acetaminophen and 3 mg per tablet of either the purified L-enantiomer or the racemate
mixture composed of equal proportions of the D- and L- isomers of NAC administered either
serially or co-administered three or four times a day up to the highest tolerable dose, given
that there will be individual variability in the ability to tolerate NAC, the number of tablets to
15 vary according to the patient's circumstances and body weight.

Example 4

A child or adult receiving nutrition parenterally can be treated to supplement the
nutrition formula with a formulation comprising 3 mg or more of either the purified
20 Lenantiomer or the racemate mixture composed of equal proportions of the D- and L-isomers
of NAC administered three or four times a day up to the highest tolerable dose, given that
there will be individual variability in the ability to tolerate NAC, variability in the amount of
the cysteine source in the total parenteral nutrition formulation and the variability of the
concomitantly administered drugs.

Example 5

25 A child or adult having liver failure or liver damage or having elevated liver enzymes
and fever who is already compromised can be treated with a cold, pain, antipyretic or other
formulation comprising between 4 and 10 mg/kg acetaminophen and up to 130 mg/kg of
either the purified L-enantiomer or the racemate mixture composed of equal proportions of
the D- and L- isomers of NAC administered either serially or coadministered three or four
30 times a day up to the highest tolerable dose given that there will be individual variability in
the ability to tolerate NAC.

Example 6

A patient having HIV infection can be treated with a formulation comprising a
therapeutically effective amount of AZT as part of a multi-drug antiviral regimen and a

5 toxicity-reducing amount of either the purified L-enantiomer or the racemate mixture composed of equal proportions of the D- and L- isomers of NAC administered either serially or co-administered two, three or four times a day up to the highest tolerable dose, given that there will be individual variability in the ability to tolerate NAC. It is preferred that NAC be formulated at high doses as an effervescent tablet or in granular form in a single dose packet
10 to be dissolved in water to prevent untoward stomach effects. This dosage of NAC is sufficient to decrease key aspects of AZT toxicity in human HIV patients.

Example 7

AZT and NAC can be administered perinatally and neonatally to prevent vertical transmission of the HIV virus to the child. Before delivery, the mother is treated with a
15 therapeutically effective amount of AZT and a toxicity-reducing amount of either the purified L-enantiomer or the racemate mixture composed of equal proportions of the D- and L-isomers of NAC administered either serially or co-administered two, three, or four times a day to the highest tolerable dose, given that there will be individual variability in the ability to tolerate NAC. Dosage is continued to the child after delivery to prevent oxidative stress
20 and other AZT toxicity. It is preferred that NAC be formulated at high doses as an effervescent tablet or in granular form in a single dose packet to be dissolved in water to prevent untoward stomach effects. This dosage of NAC is sufficient to decrease key aspects of AZT toxicity in human HIV patients.

Example 8

25 A patient with newly-diagnosed or suspected exposure to HIV can be treated with a formulation comprising a therapeutically effective amount of AZT or other similar antiretroviral drug as normally used and a toxicity-reducing amount of either the purified L-enantiomer or the racemate mixture composed of equal proportions of the D- and L-isomers of NAC administered either serially or co-administered two, three or four times a day up to the
30 highest tolerable dose, given that there will be individual variability in the ability to tolerate NAC. It is preferred that NAC be formulated at high doses as an effervescent tablet or in granular form in a single dose packet to be dissolved in water to prevent untoward stomach effects. This dosage of NAC is sufficient to decrease key aspects of AZT toxicity in human HIV patients.

5 While the present invention has been described with reference to the specific
embodiments thereof, it should be understood by those skilled in the art that various changes
may be made and equivalents may be substituted without departing from the true spirit and
scope of the Invention. In addition, many modifications may be made to adapt a particular
situation, material, composition of matter, process, process step or steps, to the objective,
10 spirit and scope of the present invention. All such modifications are intended to be within the
scope of the claims appended hereto.

What is claimed is:

1. A pharmaceutical composition useful for treating or preventing the toxic effects of therapeutic agents in mammals, comprising:
 - 5 (a) a toxicity-reducing amount of N-acetylcysteine, a pharmaceutically acceptable salt of N-acetylcysteine, or a pharmaceutically acceptable derivative of N-acetylcysteine in combination with a pharmaceutically acceptable excipient or carrier,
wherein a unit dose of N-acetylcysteine comprises at least 1.5 mg N-acetylcysteine per kg of body weight and not more than 2000 mg N-acetylcysteine, and
 - 10 (b) a therapeutic or greater amount of a therapeutic agent that produces oxidative stress;
wherein the composition does not comprise diallyl sulfide, diallyl sulfone or diallyl sulfoxide.
2. A pharmaceutical composition according to claim 1 wherein each dosage unit contains at least 1 mg N-acetylcysteine or a pharmaceutically-acceptable salt or derivative thereof.
- 15 3. A pharmaceutical composition according to claim 1 or 2, wherein the therapeutic agent is at least one antibiotic or antiviral agent.
- 20 4. A pharmaceutical composition according to any one of claims 1 to 3 wherein the therapeutic agent is at least one antibiotic or antiviral agent that produces oxidative stress.
5. A pharmaceutical composition according to any one of claims 1 to 3, wherein the therapeutic agent is at least one antibiotic or antiviral agent that produces treatment-related decreases
25 in subject cysteine/glutathione levels.
6. A pharmaceutical composition according to claim 1 or 2 wherein the therapeutic agent is at least one analgesic, antipyretic or other therapeutic compound.
- 30 7. A pharmaceutical composition according to claim 1 or 2, wherein the therapeutic agent is at least one analgesic, antipyretic or other therapeutic compound, wherein the compound produces oxidative stress.
8. A pharmaceutical composition according to claim 1 or 2 wherein the therapeutic agent is at
35 least one analgesic, antipyretic or other therapeutic compound, wherein the compound produces treatment-related decreases in subject cysteine/glutathione levels.
9. A pharmaceutical composition according to claim 6, wherein the compound is acetaminophen.

23 Oct 2006
2002255942

30

10. A pharmaceutical composition according to claim 9, wherein each dosage unit contains a standard dose of acetaminophen and at least about 1 mg of N-acetylcysteine.
11. A pharmaceutical composition according to claim 1 or 2, wherein the therapeutic agent is an anti-retroviral agent.
12. A pharmaceutical composition according to claim 11, wherein the anti-retroviral agent is AZT.
13. A method of treating or preventing the toxic effects of therapeutic agents in mammals, the method comprising the step of administering a composition comprising:
 - (a) a toxicity-reducing amount of N-acetylcysteine, a pharmaceutically acceptable salt of N-acetylcysteine, or a pharmaceutically acceptable derivative of N-acetylcysteine in combination with a pharmaceutically acceptable excipient or carrier,
wherein a unit dose of N-acetylcysteine comprises at least about 1.5 mg N-acetylcysteine per kg of body weight and not more than about 2000 mg N-acetylcysteine, and
 - (b) a therapeutic or greater amount of a therapeutic agent that produces oxidative stress;
wherein the composition does not comprise diallyl sulfide, diallyl sulfone or diallyl sulfoxide.
14. A method according to claim 13, wherein at least about 1 mg N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof is administered to a patient receiving nutrition parenterally.
15. A method according to claim 13, wherein a therapeutic amount of at least one antibiotic or antiviral agent is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof.
16. A method according to claim 13, wherein a therapeutic amount of at least one antibiotic or antiviral agent is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof, wherein the agent produces oxidative stress.

35

17. A method according to claim 13, wherein a therapeutic amount of at least one antibiotic or antiviral agent is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof, wherein the agent produces treatment-related decreases in subject cysteine/glutathione levels.

18. A method according to claim 13, wherein the therapeutic agent is an anti-retroviral agent.
19. A method according to claim 18, wherein the anti-retroviral agent is AZT.

5 20. A method according to claim 19, wherein AZT is administered serially or in combination with a toxicity-reducing amount of N-acetylcysteine perinatally.

21. A method according to claim 19, wherein AZT is administered serially or in combination with a toxicity-reducing amount of N-acetylcysteine neonatally.

22. A method according to claim 19, wherein AZT is administered for suspected
10 or known exposure to the HIV virus serially or in combination with a toxicity-reducing amount of N-acetylcysteine.

23. A method according to claim 13, wherein a therapeutic amount of at least one analgesic, antipyretic or other therapeutic compound is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof.

15 24. A method according to claim 13, wherein a therapeutic amount of at least one analgesic, antipyretic or other therapeutic compound is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof, wherein the compound produces oxidative stress.

25. A method according to claim 13, wherein a therapeutic amount of at least one analgesic, antipyretic or other therapeutic compound is administered in combination with N-acetylcysteine or a pharmaceutically acceptable salt or derivative thereof, wherein the compound produces treatment-related decreases in subject cysteine/glutathione levels.

26. A method according to claim 23, wherein the compound is acetaminophen.

27. A method according to claim 26, wherein the acetaminophen is administered
25 in a standard dosage serially or in combination with at least about 1 mg per dose N-acetylcysteine to treat a fever or pain in a patient who responds normally to acetaminophen.

28. A method according to claim 26, wherein the acetaminophen is administered in a greater than standard dosage serially or in combination with at least about 4 mg/kg N-acetylcysteine to treat a fever nonresponsive to the standard dose of acetaminophen.

2002255942 16 Jun 2005

29. A method according to claim 26, wherein the acetaminophen is administered in a standard dosage serially or in combination with at least about 1 mg per dose N-acetylcysteine to treat patients who require acetaminophen during their exposure to radiation contrast agents.

5 30. A pharmaceutical composition according to claim 1 substantially as hereinbefore described, with reference to any of the Examples.

31. A method according to claim 13 substantially as hereinbefore described, with reference to any of the Examples.

10

Dated: 15 June 2005

PHILLIPS ORMONDE & FITZPATRICK

Agents for: BIOADVANTEX PHARMA INC.

15 