



(22) Date de dépôt/Filing Date: 1990/08/08

(41) Mise à la disp. pub./Open to Public Insp.: 1991/02/11

(45) Date de délivrance/Issue Date: 2002/03/26

(30) Priorité/Priority: 1989/08/10 (P 39 26 468.8) DE

(51) Cl.Int.⁵/Int.Cl.⁵ C07C 327/46, A01N 37/18

(72) Inventeurs/Inventors:

Leyendecker, Joachim, DE;
Kardorff, Uwe, DE;
Kuenast, Christoph, DE;
Hofmeister, Peter, DE;
Neubauer, Hans-Juergen, DE;
Krieg, Wolfgang, DE


(73) Propriétaire/Owner:

BASF AKTIENGESELLSCHAFT, DE

(74) Agent: ROBIC

(54) Titre : CYCLOPROPANETHIOCARBOXAMIDES

(54) Title: CYCLOPROPANETHIOCARBOXAMIDES




(57) Abrégé/Abstract:

Cyclopropanethiocarboxamides I (see formula I) (R^1 , R^2 = H, CN, NO_2 , halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy; R^3 , R^4 = H, C_1 - C_4 -alkyl; n = 0 or 1). The compounds I are suitable as pesticides.



ABSTRACT OF THE DISCLOSURE: Cyclopropanethiocarboxamides I



(R¹, R² = H, CN, NO₂, halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy; R³, R⁴ = H, C₁-C₄-alkyl; n = 0 or 1).

5

The compounds I are suitable as pesticides.

10

15

20

25


30

35

40

Cyclopropanethiocarboxamides

The present invention relates to novel cyclopropanethiocarboxamides of the general formula I



5 where

R^1 and R^2 are each hydrogen, cyano, nitro, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, and the alkyl and alkoxy radicals may be partially or completely halogenated,

10 R^3 and R^3 are each hydrogen or C_1 - C_4 -alkyl and n is 0 or 1.

The present invention furthermore relates to a process for the preparation of these compounds, their use for pest control and agents which contain these compounds as active substances.

15 DE-A 36 28 082 discloses N-[2-(4-phenoxyphenoxy)-ethyl]-cyclopropanecarboxamides as pesticidal compounds.

However, the action of these known pesticides on the pests and the duration of their action is satisfactory only to a limited extent.

20 It is an object of the present invention to provide novel pesticides with which the pests can be controlled better than in the past.

We have found that this object is achieved by the cyclopropanethiocarboxamides of the general formula I which were defined at the outset.

25 The substituents in the novel compounds I have the following specific meanings:

R^1 is hydrogen, cyano or nitro;

halogen, including preferably fluorine, chlorine or bromine;


30 straight-chain or branched C_1 - C_4 -alkyl, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, preferably methyl, ethyl, isopropyl, isobutyl or sec-butyl;

straight-chain or branched C₁-C₄-alkoxy, preferably methoxy, ethoxy or isopropoxy;
 partially or completely halogenated C₁-C₄-alkyl, preferably straight-chain or branched C₁-C₄-fluoro- or C₁-C₄-chloroalkyl, such as fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, trichloromethyl or 2,2,2-trichloroethyl;
 partially or completely halogenated C₁-C₄-alkoxy, preferably straight-chain or branched C₁-C₄-fluoro- or C₁-C₄-chloroalkoxy, such as trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, pentafluoroethoxy or trichloromethoxy;
 R² is preferably in the 4-position of the phenyl ring and has the same meanings as R¹; and
 R³ and R⁴ are each hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, preferably methyl or ethyl.

Particularly suitable compounds I are those in which R³ and R⁴ are each hydrogen and n is 0. Preferred compounds I are described in the Examples.

The novel compounds I can be prepared by conventional methods of thiocarboxamide synthesis (cf. C. Ferri, Reaktionen der organischen Synthese, Georg Thieme Verlag, Stuttgart 1978, page 549 et seq.), for example by sulfurization of the corresponding cyclopropanecarboxamides with P₄S₁₀ or with Lawesson reagent (2,4-bis-(4-methoxyphenyl)-2,4-dithioxo-1,3,2,4-dithiadiphosphetane).

The compounds I are particularly advantageously obtained by reacting a 4-phenoxyphenoxyalkylamine II



with a cyclopropanedithiocarboxylate III



where R⁵ is a short-chain alkyl, preferably C₁-C₆ alkyl, especially methyl.

The starting compounds II and III are preferably used in a stoichiometric ratio, but in some cases an excess of one or other component, for example not more than 10%, may be preferable.

5 The reaction usually takes place at an adequate rate at above -30°C . In general, the temperature is from -30 to 130°C , in particular from -10 to 80°C , preferably from 0 to 50°C .

10 It is advantageously carried out in a solvent or diluent at atmospheric pressure. A lower or higher pressure is possible but generally has no advantages.

15 Suitable solvents or diluents are aliphatic and aromatic hydrocarbons and chlorohydrocarbons, such as petroleum ether, n-pentane, n-hexane, hexane isomer mix-
tures, benzene, toluene, xylene, gasoline, dichloro-
methane, chloroform, carbon tetrachloride, 1,2-dichloro-
ethane and chlorobenzene, ethers, such as diethyl ether,
di-n-butyl ether, methyl tert-butyl ether, tetrahydro-
furan and dioxane, esters, such as ethyl acetate,
20 ketones, such as acetone, methyl ethyl ketone and methyl isopropyl ketone, nitriles, such as acetonitrile and propionitrile, alcohols, such as methanol, ethanol, n-propanol and isopropanol, and aprotic dipolar solvents, such as dimethylformamide, dimethyl sulfoxide and pyrid-
25 ine. Mixtures of these substances can also be used as solvents and diluents.

30 Some of the 4-phenoxyphenoxyalkylamines II required for preparation of the compounds I are known from Houben/Weyl, Vol. VI/3, Methoden der organischen Chemie, Thieme Verlag, 1965, page 85 et seq.; those which are unknown can be prepared by the methods described there.

35 The cyclopropanedithiocarboxylates III additionally required are known from C.R. Acad. Sci., Ser. C, 274, (1972), 642 or can be prepared by the method described there. They are obtained, for example, by subjecting the cyclopropyl bromide, magnesium and carbon

disulfide to a Grignard reaction and then reacting the resulting Grignard compound with an alkyl iodide, preferably methyl iodide.


Some of the novel compounds I are obtained in the form of colorless or slightly brownish oils, which can be freed from the final volatile constituents by prolonged heating to moderately elevated temperatures under reduced pressure (incipient distillation) and can be purified in this manner. If the compounds of the formula I are obtained in crystalline form, they can be purified by recrystallization.

The novel compounds I may contain one or more centers of asymmetry. They are obtained as racemates in most preparation processes but can, if desired, also be separated into the pure isomers by conventional methods, for example by chromatography over an optically active adsorbent.

In contrast to most of the pesticides known to date, which are contact or ingested poisons and thus kill, incapacitate or repel the animals, the compounds of the formula I intervene in the hormone system of the animal organism. In the case of insects, for example, the transformation to the imago, the laying of viable eggs and the development of normal laid eggs is disturbed and the sequence of generations is thus interrupted. The novel agents are virtually completely nontoxic to vertebrates.

Preparation Example

[N-[4-(3-Fluorophenoxy)-phenoxy]-ethyl]-cyclopropanethiocarboxamide



A solution of 2.4 g (9.7 mmol) of [4-(3-fluorophenoxy)-phenoxy]-ethylamine, 1.3 g (9.7 mmol) of methyl cyclopropanedithiocarboxylate and 40 ml of toluene were stirred for 14 hours at 20°C, after which the readily

volatile substances were distilled off under reduced pressure. The residue was recrystallized from ethyl acetate/n-hexane. The product was obtained in the form of colorless crystals in a yield of 75%.


5 ^1H -NMR spectrum (300 MHz, d^6 -DMSO, tetramethylsilane as standard [0 ppm]):

0.68-1.2 ppm (m, 4 H); 2.05-2.30 ppm (m, 1 H); 3.92 ppm (t, 2 H); 4.21 ppm (t, 2H); 6.58-7.42 ppm (m, 8 H); 10.38 ppm (s, 1 H).

10 Further physical data are shown in the Table below, which lists other compounds which were prepared, or can be prepared, in a similar manner.

Table 1

Novel cyclopropanethiocarboxamides



No.	n	R1	R2	R3	R4	mp. [°C]	IR data [cm ⁻¹]	Yield [%]
1	0	F	H	H	H	69	1507, 1487, 1248, 1240, 1209 1121, 962, 848, 841	75
2	0	H	H	H	H	60-62	1504, 1488, 1247, 1235, 1194, 994, 832	73
3	0	H	4-F	H	H	65	1504, 1250, 1231, 1192, 1064, 994, 827	50
4	0	Cl	H	H	H	101	1505, 1472, 1464, 1216, 1192, 1056, 995, 967, 842	62
5	0	Br	H	H	H	56- 57	1505, 1489, 1468, 1229, 1222, 1058, 994	83
6	0	Cl	4-F	H	H	99-100	1507, 1493, 1463, 1247, 1209, 1054, 998, 845, 784	68
7	0	CH3	H	H	H	oil	1502, 1486, 1464, 1328, 1256, 1240, 1208, 1059, 996	81
8	0	CF3	H	H	H	oil	1503, 1491, 1450, 1328, 1221, 1191, 1170, 1126, 1064, 914	90
9	0	C2H5	H	H	H	oil	1502, 1485, 1465, 1447, 1328, 1233, 1207, 1059, 996	87
10	0	OCH3	H	H	H			
11	0	Cl	4-Cl	H	H			
12	0	OCF3	H	H	H			
13	0	H	4-Cl	H	H			
14	0	H	4-Br	H	H			
15	0	NO2	H	H	H			
16	0	F	H	CH3	H			
17	0	H	H	CH3	H			

Table 1 (contd.)

No.	n	R1	R2	R3	R4	mp. [°C]	IR data [cm ⁻¹]	Yield [%]
18	0	H	4-F	CH3	H			
19	0	Cl	H	CH3	H			
20	0	Br	H	CH3	H			
21	0	Cl	4-F	CH3	H			
22	0	CH3	H	CH3	H			
23	0	CF3	H	CH3	H			
24	0	C2H5	H	CH3	H			
25	0	OCH3	H	CH3	H			
26	0	Cl	4-Cl	CH3	H			
27	0	OCF3	H	CH3	H			
28	0	H	4-Cl	CH3	H			
29	0	H	4-Br	CH3	H			
30	0	NO2	H	CH3	H			
31	0	F	H	H	CH3			
32	0	H	H	H	CH3			
33	0	H	4-F	H	CH3			
34	0	Cl	H	H	CH3			
35	0	Br	H	H	CH3			
36	0	Cl	4-F	H	CH3			
37	0	CH3	H	H	CH3			
38	0	CF3	H	H	CH3			
39	0	C2H5	H	H	CH3			

Table 1 (contd.)

No.	n	R ¹	R ²	R ³	R ⁴	mp. [°C]	IR data [cm ⁻¹]	Yield [%]
40	O	OCH ₃	H	H	CH ₃			
41	O	Cl	4-Cl	H	CH ₃			
42	O	OCF ₃	H	H	CH ₃			
43	O	H	4-Cl	H	CH ₃			
44	O	H	4-Br	H	CH ₃			
45	O	NO ₂	H	H	CH ₃			

The cyclopropanethiocarboxamides are suitable for effectively combating pests such as insects, arachnids and nematodes. They may be used as pesticides in crop protection and in the hygiene, stores protection and veterinary sector.

5

Examples of injurious insects belonging to the Lepidoptera order are *Agrotis ypsilon*, *Agrotis segetum*, *Alabama argillacea*, *Anticarsia gemmatilis*, *Argyresthia conjugella*, *Autographa gamma*, *Bupalus piniarius*, *Cacoecia murinana*, *Capua reticulana*, *Cheimatobia brumata*, *Choristoneura*
 10 *fumiferana*, *Choristoneura occidentalis*, *Cirphis unipuncta*, *Cydia pomonella*, *Dendrolimus pini*, *Diaphania nitidalis*, *Diatraea grandiosella*, *Earias insulana*, *Elasmopalpus lignosellus*, *Eupoecilia ambiguella*, *Evetria bouliana*, *Feltia subterranea*, *Galleria mellonella*, *Grapholita funebrana*, *Grapholita molesta*, *Heliothis armigera*, *Heliothis virescens*, *Heliothis*
 15 *zea*, *Hellula undalis*, *Hibernia defoliaria*, *Hyphantria cunea*, *Hyponomeuta malinellus*, *Keifferia lycopersicella*, *Lambdina fiscellaria*, *Laphygma exigua*, *Leucoptera coffeella*, *Leucoptera scitella*, *Lithocolletis blanchardella*, *Lobesia botrana*, *Loxostege sticticalis*, *Lymantria dispar*, *Lymantria monacha*, *Lyonetia clerkella*, *Malacosoma neustria*, *Mamestra*
 20 *brassicae*, *Orgyia pseudotsugata*, *Ostrinia nubilalis*, *Panolis flamea*, *Pectinophora gossypiella*, *Peridroma saucia*, *Phalera bucephala*, *Phthorimaea operculella*, *Phyllocnistis citrella*, *Pieris brassicae*, *Plathypena scabra*, *Plutella xylostella*, *Pseudoplusia includens*, *Phyacionia frustrana*, *Scrobipalpula absoluta*, *Sitotroga cerealella*, *Sparganothis pilleriana*, *Spodoptera*
 25 *frugiperda*, *Spodoptera littoralis*, *Spodoptera litura*, *Thaumatopoea pityocampa*, *Tortrix viridana*, *Trichoplusia ni* and *Zeiraphera canadensis*.

Examples from the Coleoptera order are *Agrilus sinuatus*, *Agriotes lineatus*, *Agriotes obscurus*, *Amphimallus solstitialis*, *Anisandrus dispar*,
 30 *Anthonomus grandis*, *Anthonomus pomorum*, *Atomaria linearis*, *Blastophagus piniperda*, *Blitophaga undata*, *Bruchus rufimanus*, *Bruchus pisorum*, *Bruchus lentis*, *Byctiscus betulae*, *Cassida nebulosa*, *Cerotoma trifurcata*, *Ceuthorrhynchus assimilis*, *Ceuthorrhynchus napi*, *Chaetocnema tibialis*, *Conoderus vespertinus*, *Crioceris asparagi*, *Diabrotica longicornis*,
 35 *Diabrotica 12-punctata*, *Diabrotica virgifera*, *Epilachna varivestis*, *Epitrix hirtipennis*, *Eutinobothrus brasiliensis*, *Hylobius abietis*, *Hypera brunneipennis*, *Hypera postica*, *Ips typographus*, *Lema bilineata*, *Lema melanopus*, *Leptinotarsa decemlineata*, *Limonius californicus*, *Lissorhoptrus oryzophilus*, *Melanotus communis*, *Meligethes aeneus*, *Melolontha*
 40 *hippocastani*, *Melolontha melolontha*, *Onlema oryzae*, *Ortiorrhynchus sulcatus*, *Ortiorrhynchus ovatus*, *Phaedon cochleariae*, *Phyllotreta chrysocephala*, *Phyllophaga* sp., *Phyllopertha horticola*, *Phyllotreta nemorum*, *Phyllotreta striolata*, *Popillia japonica*, *Sitona lineatus* and *Sitophilus granaria*.

Examples from the Diptera order are *Aedes aegypti*, *Aedes vexans*,
Anastrepha ludens, *Anopheles maculipennis*, *Ceratitis capitata*, *Chrysomya*
bezziana, *Chrysomya hominivorax*, *Chrysomya macellaria*, *Contarinia*
sorghicola, *Cordylobia anthropophaga*, *Culex pipiens*, *Dacus cucurbitae*,
5 *Dacus oleae*, *Dasineura brassicae*, *Fannia canicularis*, *Gasterophilus*
intestinalis, *Glossia morsitans*, *Haematobia irritans*, *Haplodiplosis*
equestris, *Hylemyia platura*, *Hypoderma lineata*, *Liriomyza sativae*,
Liriomyza trifolii, *Lucilia caprina*, *Lucilia cuprina*, *Lucilia sericata*,
Lycoria pectoralis, *Mayetiola destructor*, *Musca domestica*, *Muscina*
10 *stabulans*, *Oestrus ovis*, *Oscinella frit*, *Pegomya hysocyami*, *Phorbia*
antiqua, *Phorbia brassicae*, *Phorbia coarctata*, *Rhagoletis cerasi*, *Rhago-*
letis pomonella, *Tabanus bovinus*, *Tipula oleracea* and *Tipula paludosa*.

Examples from the Thysanoptera order are *Frankliniella fusca*,
15 *Frankliniella occidentalis*, *Frankliniella tritici*, *Scirtothrips citri*,
Thrips oryzae, *Thrips palmi* and *Thrips tabaci*.

Examples from the Hymenoptera order are *Athalia rosae*, *Atta cephalotes*,
Atta sexdens, *Atta texana*, *Hoplocampa minuta*, *Hoplocampa testudinea*,
20 *Monomorium pharaonis*, *Solenopsis geminata* and *Solenopsis invicta*.

Examples from the Heteroptera order are *Acrosternum hilare*, *Blissus*
leucopterus, *Cyrtopeltis notatus*, *Dysdercus cingulatus*, *Dysdercus*
intermedius, *Eurygaster integriceps*, *Euchistus impictiventris*,
25 *Leptoglossus phyllopus*, *Lygus lineolaris*, *Lygus pratensis*, *Nezara*
viridula, *Piesma quadrata*, *Solubea insularis* and *Thyanta perditor*.

Examples from the Homoptera order are *Acyrtosiphon onobrychis*, *Adelges*
laricis, *Aphidula nasturtii*, *Aphis fabae*, *Aphis pomi*, *Aphis sambuci*,
30 *Brachycaudus cardui*, *Brevicoryne brassicae*, *Cerosipha gossypii*, *Dreyfusia*
nordmanniana, *Dreyfusia piceae*, *Dyasphis radicola*, *Dysaulacorthum*
pseudosolani, *Empoasca fabae*, *Macrosiphum avenae*, *Macrosiphum euphorbiae*,
Macrosiphon rosae, *Megoura viciae*, *Metopolophium dirhodum*, *Myzodes*
persicae, *Myzus cerasi*, *Nilaparvata lugens*, *Pemphigus bursarius*,
35 *Perkinsiella saccharicida*, *Phorodon humuli*, *Psylla mali*, *Psylla piri*,
Rhopalomyzus ascalonicus, *Rhopalosiphum maidis*, *Sappaphis mala*, *Sappaphis*
mali, *Schizaphis graminum*, *Schizoneura lanuginosa*, *Trialeurodes*
vaporariorum and *Viteus vitifolii*.

40 Examples from the Isoptera order are *Calotermes flavicollis*, *Leucotermes*
flavipes, *Reticulitermes lucifugus* and *Termes natalensis*.

Examples from the Orthoptera order are *Acheta domestica*, *Blatta orientalis*, *Blattella germanica*, *Forficula auricularia*, *Gryllotalpa gryllotalpa*, *Locusta migratoria*, *Melanoplus birittatus*, *Melanoplus femur-rubrum*, *Melanoplus mexicanus*, *Melanoplus sanguinipes*, *Melanoplus*
 5 *spretus*, *Nomadacris septemfasciata*, *Periplaneta americana*, *Schistocerca americana*, *Schistocerca peregrina*, *Stauronotus maroccanus* and *Tachycines asynamorus*.

Examples from the Acarina order are *Amblyomma americanum*, *Amblyomma*
 10 *variegatum*, *Argas persicus*, *Boophilus annulatus*, *Boophilus decoloratus*, *Boophilus microplus*, *Brevipalpus phoenicis*, *Bryobia praetiosa*, *Dermacentor silvarum*, *Eotetranychus carpini*, *Eriophyes sheldoni*, *Hyalomma truncatum*, *Ixodes ricinus*, *Ixodes rubicundus*, *Ornithodoros moubata*, *Otobius megnini*, *Paratetranychus pilosus*, *Permanyssus gallinae*, *Phyllocaptrata oleivora*,
 15 *Polyphagotarsonemus latus*, *Psoroptes ovis*, *Rhipicephalus appendiculatus*, *Rhipicephalus evertsi*, *Sarcoptes scabiei*, *Tetranychus cinnabarinus*, *Tetranychus kanzawai*, *Tetranychus pacificus*, *Tetranychus telarius* and *Tetranychus urticae*.


20 Examples from the nematodes class are root-knot nematodes, e.g., *Meloidogyne hapla*, *Meloidogyne incognita* and *Meloidogyne javanica*, cyst-forming nematodes, e.g., *Globodera rostochiensis*, *Heterodera avenae*, *Heterodera glycinae*, *Heterodera schachtii* and *Heterodera trifolii*, and stem and leaf
 25 eelworms, e.g., *Belonolaimus longicaudatus*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Heliocotylenchus multicinctus*, *Longidorus elongatus*, *Radopholus similis*, *Rotylenchus robustus*, *Trichodorus primitivus*, *Tylenchorhynchus claytoni*, *Tylenchorhynchus dubius*, *Pratylenchus neglectus*, *Pratylenchus penetrans*, *Pratylenchus curvatus* and *Pratylenchus goodeyi*.

30 The active ingredients may be applied for instance as such or together with inert additives in the form of formulations or application forms prepared therefrom, e.g., directly sprayable solutions, powders, suspensions, dispersions, emulsions, oil dispersions, pastes, dusts,
 35 broadcasting agents, or granules by spraying, atomizing, dusting, broadcasting or watering. The forms of application depend entirely on the purpose for which the agents are being used, but they must ensure as fine a distribution of

11a

the active ingredients according to the invention as possible.

For the preparation of solutions, emulsions, pastes and oil dispersions to be sprayed direct, mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, further coal-tar oils, and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons such as benzene, toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes and their derivatives such as methanol, ethanol, propanol,



butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, etc., and strongly polar solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone, water, etc. are suitable.

5

Aqueous formulations may be prepared from emulsion concentrates, pastes, oil dispersions or wettable powders by adding water. To prepare emulsions, pastes and oil dispersions the ingredients as such or dissolved in an oil or solvent may be homogenized in water by means of wetting or dispersing agents, adherents or emulsifiers. Concentrates which are suitable for dilution with water may be prepared from active ingredient, wetting agent, adherent, emulsifying or dispersing agent and possibly solvent or oil.

Examples of surfactants are: alkali metal, alkaline earth metal and ammonium salts of ligninsulfonic acid, naphthalenesulfonic acids, phenolsulfonic acids, alkylaryl sulfonates, alkyl sulfates, and alkyl sulfonates, alkali metal and alkaline earth metal salts of dibutyl-naphthalenesulfonic acid, lauryl ether sulfate, fatty alcohol sulfates, alkali metal and alkaline earth metal salts of fatty acids, salts of sulfated hexadecanols, heptadecanols, and octadecanols, salts of sulfated fatty alcohol glycol ethers, condensation products of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensation products of naphthalene or naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ethers, ethoxylated isooctyl-phenol, ethoxylated octylphenol and ethoxylated nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methyl cellulose.

Powders, dusts and broadcasting agents may be prepared by mixing or grinding the active ingredients with a solid carrier.

35 Examples of formulations are given below.

I. 5 parts by weight of compound no. 1 is intimately mixed with 95 parts by weight of particulate kaolin. A dust is obtained containing 5% by weight of the active ingredient.

40

II. 30 parts by weight of compound no. 2 is intimately mixed with a mixture consisting of 92 parts by weight of powdered silica gel and 8 parts by weight of paraffin oil which has been sprayed onto the surface of this silica gel. A formulation of the active ingredient is obtained having good adherence.

III. 10 parts by weight of compound no. 3 is dissolved in a mixture consisting of 90 parts by weight of xylene, 6 parts by weight of the adduct of 8 to 10 moles of ethylene oxide and 1 mole of oleic acid-N-monoethanolamide, 2 parts by weight of the calcium salt of dodecylbenzenesulfonic acid, and 2 parts by weight of the adduct of 40 moles of ethylene oxide and 1 mole of castor oil.

IV. 20 parts by weight of compound no. 4 is dissolved in a mixture consisting of 60 parts by weight of cyclohexanone, 30 parts by weight of 10 isobutanol, 5 parts by weight of the adduct of 7 moles of ethylene oxide and 1 mole of isooctylphenol, and 5 parts by weight of the adduct of 40 moles of ethylene oxide and 1 mole of castor oil.

V. 80 parts by weight of compound no. 5 is well mixed with 3 parts by 15 weight of the sodium salt of diisobutyl-naphthalene-alpha-sulfonic acid, 10 parts by weight of the sodium salt of a lignin-sulfonic acid obtained from a sulfite waste liquor, and 7 parts by weight of powdered silica gel, and triturated in a hammer mill.

20 Granules, e.g., coated, impregnated or homogeneous granules, may be prepared by bonding the active ingredients to solid carriers. Examples of solid carriers are mineral earths such as silicic acid, silica gels, silicates, talc, kaolin, attapul-gus clay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium 25 sulfate, magnesium oxide, ground plastics, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, and ureas, and vegetable products such as grain flours, bark meal, wood meal, and nutshell meal, cellulosic powders, etc.

30 The formulations generally contain from 0.1 to 95, and preferably 0.5 to 90, % by weight of active ingredient.

The active ingredient concentrations in the finished formulations may vary over a wide range. Generally, they are from 0.0001 to 10, and preferably 35 from 0.01 to 1, %.

The active ingredients may also successfully be used in the ultra-low-volume (ULV) method, where it is possible to apply formulations containing more than 95wt% of active ingredient, or even the active ingredient with- 40 out additives.

In the open, the amount of active ingredient applied is for example from 0.02 to 10, particularly from 0.1 to 2, kg/ha.

There may be added to the active ingredients (if desired, immediately before use (tankmix)) oils of various types, herbicides, fungicides, other pesticides and bactericides. These agents may be added to the active ingredients according to the invention in a weight ratio of from 1:10 to 5 10:1.

Use examples

The concentrations at which the compounds investigated achieved 100% or 10 80% kill or inhibition are the minimum concentrations in each case (action threshold).

The purity of the active ingredients was >95%.

15 The following formulations were used:

- a) a 0.1% solution of the active ingredient in acetone, which was further diluted with acetone according to the stated dosage rates;
- 20 b) a 10% emulsion of the active ingredient in a mixture consisting of 70wt% of cyclohexanone, 20wt% of Nekanil® LN ($\cong$ Lutensol AP6, a spreadersticker with an emulsifying and dispersing action based on ethoxylated alkylphenols) and 10wt% of Emulphor® EL ($\cong$ Emulan® EL, an emulsifier based on ethoxylated fatty alcohols). The concentrations
25 given in the examples were obtained by diluting the formulated active ingredient with water.

Example A

30 Ovicidal action on *Dysdercus intermedius* (cotton stainer)

Pieces of double-sided adhesive tape (about 0.8 cm) were stuck to the top edge of plastic plant markers. 24 hours before commencement of the experiment, eggs of the cotton stainer contained in a vessel were attached to
35 the adhesive strips by dipping the markers into the vessel. The eggs were then dipped for 5 seconds into aqueous formulations of the active ingredients and excess liquid was allowed to drip off onto filter paper, care being taken to prevent the eggs coming into contact with the paper.

40 The markers were then placed in plastic trays (adhesive strip at the top). Half a roll of absorbent cotton was moistened with water and placed in each beaker to prevent drying out, and the trays were covered with a glass plate. Assessment took place after 8 days (control bugs hatched).

At an active ingredient concentration of 0.1 to 10 ppm, compounds 1, 3, 4, 5, 6 and 7 achieved 80 to 100% kill.

Example B

5

Dysdercus intermedius (cotton stainer); breeding experiment

The experiment was carried out in 1 liter jars containing 200 g of sterile quartz sand to which 25 ml of aqueous active ingredient formulations had been admixed. Approx. 20 larvae of the third stage were then introduced into each jar. The food proffered was swollen cottonseed, which was changed once a week. The sand was also moistened once a week. The experiment was run until the subsequent generation hatched. The action in percentage kill was then assessed.

15

At an active ingredient concentration of 0.01 to 1 ppm, compounds 1 to 7 and 9 achieved 100% kill.

Example C

20

Breeding experiment with *Musca domestica* (housefly)

The experiment was run in 100 ml plastic beakers. 25ml of a dry feed mix (1 g of bran, 250 g of yeast powder and 35 g of fishmeal) was introduced into the beakers, the active ingredients were added together with 25 ml of a solution of milk and sugar (1 liter of milk and 42 cm³ of sugar). 20 larvae in the first larval stage were then placed in each beaker. Perforated lids were then placed on the beakers. The experiment was run until the flies in a control experiment (without active ingredient) hatched.

30

At a concentration of 20 to 40 ppm, compounds 1, 3, 4, 5, 6, 7 and 9 achieved 100% kill.

Example D

35

Prodenia litura; breeding experiment on a treated nutrient medium

Glass Petri dishes 10 cm in diameter were treated with acetonc formulations of the active ingredients (a); after the solvent had evaporated, 5 caterpillars in the fourth larval stage were placed in the dishes, over which a cover was then placed. After 4 hours the kill rate was ascertained and 5 surviving caterpillars (10 to 12 mm long) were placed in 250 ml beakers. These beakers contained about 100 ml of a standard nutrient medium (3.1 liters of water, 80 g of agar, 137 g of brewers' yeast, 515 g of corn meal, 130 g of wheat germ and conventional additives and vita-

liquid, and perforated transparent lids were placed on the beakers. The experiment was run until the moths in a control experiment (without active ingredient) emerged.

5 At an active ingredient concentration of 0.002 to 0.004 ppm, compounds 1 and 3 to 9 achieved 80 to 100% kill.

10

15

20

25


30

35

40

CLAIMS


1. A cyclopropanethiocarboxamide of the general formula (I):



where R^1 and R^2 are hydrogen, cyano, nitro, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, the alkyl and alkoxy radicals being partially or completely halogenated; R^3 and R^4 are hydrogen or C_1 - C_4 -alkyl; and n is 0 or 1.

2. A cyclopropanethiocarboxamide of the formula (I) as set forth in claim 1, where R^3 and R^4 are hydrogen and n is 0.

3. A process for the manufacture of a cyclopropanethiocarboxamide of the formula (I) as set forth in claim 1, wherein a 4-phenoxyphenoxyalkylamine of the general formula (II):



where R^1 , R^2 , R^3 , R^4 and n are defined as in claim 1, is reacted with a cyclopropanedithiocarboxylate of the general formula (III):



where R^5 is a short-chain alkyl.

4. The use of a cyclopropanethiocarboxamide of the formula (I) as set forth in claim 1 for combating pests.

5. A pesticide containing at least one cyclopropane-thiocarboxamide of the formula (I) as set forth in claim 1 and inert additives.

6. A pesticide as set forth in claim 5, containing from 0.1 to 99 wt% of a cyclopropanethiocarboxamide of the formula (I).

7. A process for combating pests, wherein an effective
10 amount of a cyclopropanethiocarboxamide of the formula (I)
as set forth in claim 1 is allowed to act on pests or their
habitat.

