

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199734360 B2**
(10) Patent No. **725670**

(54) Title
Progestogen-anti-progestogen regimens

(51)⁶ International Patent Classification(s)
A61K 031/58 A61K 031/57
A61K 031/565

(21) Application No: **199734360**

(22) Application Date: **1997.06.23**

(87) WIPO No: **W097/49407**

(30) Priority Data

(31) Number	(32) Date	(33) Country
96201744	1996.06.25	EP

(43) Publication Date : **1998.01.14**

(43) Publication Journal Date : **1998.03.12**

(44) Accepted Journal Date : **2000.10.19**

(71) Applicant(s)
Akzo Nobel N.V.

(72) Inventor(s)
Herman Jan Tijmen Coelingh Bennink; Pieter M. Verboost

(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

(56) Related Art
WO 93/21927
WO 96/15794
WO 94/04156

OPI DATE 14/01/98 APPLN. ID 34360/97
AOJP DATE 12/03/98 PCT NUMBER PCT/EP97/03288



AU9734360

INTE

(51) International Patent Classification ⁶: A61K 31/58, 31/57, 31/565		A1	(11) International Publication Number: WO 97/49407 (43) International Publication Date: 31 December 1997 (31.12.97)
(21) International Application Number: PCT/EP97/03288 (22) International Filing Date: 23 June 1997 (23.06.97) (30) Priority Data: 96201744.8 25 June 1996 (25.06.96) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): COELINGH BENNINK, Herman, Jan, Tjimen [NL/NL]; Melvill van Carnebeeklaan 38, NL-3971 BE Driebergen (NL). VERBOST, Pieter, M. [NL/NL]; Plantsoen 15, NL-5384 ET Heesch (NL). (74) Agent: KRAAK, Hajo; P.O. Box 20, NL-5340 BH Oss (NL).		(81) Designated States: AU, BR, CA, CN, CZ, HU, JP, KR, MX, NO, NZ, PL, RU, SG, TR, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: PROGESTOGEN-ANTI-PROGESTOGEN REGIMENS (57) Abstract Disclosed is a contraceptive and/or HRT (hormone replacement therapy) kit comprising sequential daily dosage units each containing as the sole contraceptively effective ingredient of a progestogen, or as effective ingredient for HRT a progestogen with or without an estrogen or an estrogen only, and further two or more dosage units comprising an anti-progestogen. One of the anti-progestogen is administered at the beginning and the others regularly divided throughout the cycle, preferably one only in the middle of the cycle.			

PROGESTOGEN-ANTI-PROGESTOGEN REGIMENS

- 5 The invention relates generally to progestogen-anti-progestogen regimens for use in contraception and hormone replacement therapy, and more specifically for contraception to progestogen-anti-progestogen regimens involving only the administration of a progestogen and an anti-progestogen.
- 10 It has been known for some time that contraception can be achieved by the oral administration of sufficient quantities of a progestogen to a female of child-bearing age. Contraceptive preparations that minimize the incidence of menstrual spotting, break through bleeding, variations in menstrual cycle length and amenorrhea are preferred. It is further preferred to use contraceptive regimens that minimize the
- 15 amounts of estrogens and progestogens used. Preparations that fulfil many of these requirements are disclosed in WO 93/21927, wherein a contraceptive regimen free from estrogens is described, the active ingredient being a progestational agent and intermittently an anti-progestogen. The regimen used is a regimen wherein only levonorgestel is administered as the progestogen, except that on days 1, 30, 60, 90,
- 20 120, 150, and 180 a dosage of the anti-progestogen RU 486 is administered. In fact the regimen is a progestogen-only regimen, interrupted by anti-progestogen administration at the beginning of each cycle. Although this regimen is a considerable improvement over existing regimens comprising estrogens, the bleeding profile is still not perfect since it recurs slowly after an almost bleeding-free interval, and further
- 25 improvement is therefor desirable.

"Progestogen-only pills" are a preferred method of contraception for breast-feeding mothers, older women, women for whom estrogen is contraindicated, women who are hypertensive, and women who develop migraine headaches when taking a combined

30 pill (i.e. one containing an estrogen and progestogen component). See, e.g. "Contraception for women over the age of 35", IPPF Medical Bulletin, 22: 3-4 (1988) and P.W. Howie "The progestogen-only pill", Brit. J. Obstet. Gynaecol., 92: 1001-2 (1985).

While different progestogen-only regimens have been described, they are still associated with incomplete ovulation inhibition, and relatively high failure rates. Vessey et al "Progestogen-only oral contraception. Findings in a large prospective study with special reference to effectiveness", Brit. J. Family Planning, 292: 526-30 (1986). It has been suggested to increase the daily dosage of progestogen in order to induce complete ovulation inhibition, however such an increase in dosage also increases the frequency of intermenstrual bleeding (i.e. "spotting"), which is clearly not desired. E. Diczfalussy et al, Progestogens in Therapy, p. 150 (Raven Press, NY 1983). Moreover, a high prevalence of functional ovarian cysts have been reported with progestogen only contraceptive regimens, which resolve after discontinuation of the progestogen-only contraceptive. Fotherby, K. "The Progestogen-pill", in: Filshie et al eds. Contraception: Science and Practice, pp. 94-108 (1989), and Howie, supra. A need exists for a progestogen-only contraceptive regimen which more effectively inhibits ovulation, while still not increasing the frequency of intermenstrual bleeding, or leading to persistent functional ovarian cysts. The solution to this need by adding intermittently an anti-progestogen needs further elaboration.

Surprisingly it has now been found that apart from administering an anti-progestagen at the beginning of a cycle, by selecting one or more additional days during the cycle, preferably one day in the middle of the cycle, on which the anti-progestogen is administered, whereas over the rest of an entire menstrual cycle (e.g. 28 days) desogestrel or 3-ketodesogestrel is administered as the progestogen at certain specified dosages, complete ovulation inhibition is achieved, while retaining good cycle control and almost completely decreasing the amount of spotting.

The invention thus includes a drug delivery system for contraceptive use containing daily oral dosage units, each unit containing a progestogen, and two or more units comprising an anti-progestogen, one of which is administered at the end and the others orderly divided through the cycle (if one: in the middle of the cycle).

The invention also includes a drug delivery system for HRT (hormone replacement therapy) containing daily oral dosage units, each unit comprising a progestogen with or without an estrogen or an estrogen only, and two or more dosage units comprising an anti-progestogen, one of which is preferably administered at the beginning and the others orderly divided through the cycle (if one: in the middle of the cycle).

In general terms the invention relates to a contraceptive and/or HRT (hormone replacement therapy) kit comprising sequential daily dosage units for oral administration each comprising as the sole contraceptively effective ingredient a progestogen, or as effective ingredient for HRT a progestogen with or without an estrogen or an estrogen alone, and further two or more units comprising an anti-progestogen when used for a multi-phase hormonal treatment cycle comprising administration of one dosage unit, comprising the anti-progestogen, at the beginning or the end of the cycle and one or more other dosage units, comprising the anti-progestogen, orderly divided through the cycle.

If desired the kits may contain placebo pills to bridge two periods of administration of active ingredients. This is usual for contraceptive regimens containing less than 28 dosage units, in order to obtain a kit still having 28 pills (the usual cycle).

According to another embodiment of this invention there is provided the use of daily dosage units consisting essentially a progestogen, and of an anti-progestogen, in the preparation of a drug delivery system, said drug delivery system characterized by consisting of daily dosage units containing only a progestogenic compound as therapeutically effective ingredient and two to seven dosage units comprising anti-progestogen, one of which is administered at the beginning, and the other or others divided regularly throughout the cycle.

According to a further embodiment of this invention there is provided the use of oral daily dosage units consisting essentially of an estrogen and a progestogen, and of an anti-progestogen, in the preparation of a drug delivery system, said drug delivery system characterized by consisting of daily dosage units containing progestogen and/or estrogen as therapeutically effective ingredient and two to seven dosage units comprising an anti-progestogen, one of which is administered at the beginning and the other or others divided regularly throughout the cycle.

According to another embodiment of this invention there is provided a method to provide contraception or hormone replacement therapy to a woman by sequential administration of daily dosage units, at least two of which dosage units are comprising an anti-progestogen, one of which anti-progestogen comprising dosage units is administered at the beginning or the end of the sequential administration and one or more of the other anti-progestogen comprising dosage units is administered orderly divided through the sequence of administrations.

The invention also includes a pharmaceutical product (i.e. the dosage units or the package containing the dosage units), a method of using the product, and a process of manufacturing the pharmaceutical product.

The invention also includes a method of providing contraception and/or HRT for a pre-, peri-, or post-menopausal women involving administering to the woman the above-mentioned regimens.



Progestogen for use with the invention are 3-keto-desogestrel (etonogestrel), desogestrel, gestodene, levonorgestel, norgestrel and other progestogen commonly used for contraception and HRT. Desogestrel has the chemical name 13-ethyl-11-methylene-18,19-di-nor-17 α -pregn-4-en-20-yn-17-ol, and is the preferred progestogen. Desogestrel is believed to be metabolized in the body into 3-ketodesogestrel. Preferred the dosage units contain 75 μ g of desogestrel or 3-ketodesogestrel, or amount of other progestogen having the equivalent effect as 75 μ g of desogestrel. Based on practically applied doses, levonorgestrel, desogestrel, and 3-keto-desogestrel are relatively equipotent in progestogenic activity. Gestodene is approximately 1.5 times as potent as these compounds. Norgestrel is about one-half as potent as levonorgestrel.

The anti-progestogen can be an inhibitor of progesterone synthesis, such as epostane, azastene or trilostane (Creange, Contraception 24, 289, 1981; Drugs of the Future 7,



661, 1982, van der Spuy et al., Contraception 35, 111, 1987; US patent 3296255) or a progesterone receptor antagonist, or any such pharmaceutically suitable agent that counteracts the normal biological activity of progesterone, such as antibodies or ligands bindable to progestogens or to the progesterone receptor.

- 5 A suitable anti-progestogen is a progesterone receptor antagonist. For example RU486, Onapristone, Org 31710 [(6 α ,11 β ,17 β)-11-(4-dimethylaminophenyl)-6-methyl-4',5'-dihydrospiro[estra-4,9-diene-17,2'-(3'H)-furan]-3-one], and Org 33628 [11 β ,17 α)-11-(4-acetylphenyl)-17,23-epoxy-19,24-dinorchola-4,9,20-trien-3-one] are particularly suitable in the practice of the present invention.
- 10 Suitable amounts of anti-progestogen are for example 0.1 to 300 mg, and preferably 0.5 to 150 mg of Org 31710 or such amounts of other anti-progestogens which have equivalent activity. The anti-progestogen is administered at the beginning of the cycle, preferably on day 1, and in the middle of the cycle, preferably day 14. More generally, the first of the anti-progestogen dosage units may be administered between days 1 and
- 15 3 (the beginning of the cycle), and the second between days 12 and 16 (the middle of the cycle). In the case of more than one additional day of anti-progestagen administration, these additional days may be, e.g., in the middle of the remaining phases, i.e. anti-progestagen then is administered once a week. Maximally, the anti-progestagen can be administered every 4 days, i.e. for a cyclus of 28 days the number
- 20 of dosage units comprising anti-progestagen is two to seven. Every 5-7 days is preferred, but it is most preferred that anti-progestagen be administered only one time additionally, i.e. in total twice a month. Particularly in the embodiments in which anti-progestagen is administered more than one time additionally, it is preferred to administer it simultaneously with progestagen, i.e. the anti-progestagen does not
- 25 substitute the progestagen but comes in addition. It is important that the anti-progestagen is administered only one day at a row. For HRT regimens it is desired for the anti-progestogen to be dosed in as low an amount as is capable of inhibiting bleeding, without inducing menses.
- 30 In the method of this invention the contraceptive and/or HRT kit consists of four phases, in which the first phase is a single dosage unit comprising an anti-progestogen and optionally a progestogen, the second phase containing progestogen, estrogen or a

mixture thereof, the third phase comprising an anti-progestogen and optionally a progestogen, and the fourth phase containing progestogen, estrogen or a mixture thereof.

- 5 Preferably the dosage of the anti-progestogen in the third and optional further phases is lower than the dosage of the anti-progestogen in the first phase. More preferably the first phase consists of 10 to 150 mg of Org 31710, and most preferably 25 mg of Org 31710, and the third phase (and any optional subsequent phase) consists of 0.5 to 25 mg, more preferably of 2.5 to 12.5 mg, and even more preferably of 5 mg of Org
10 31710. In the case of a more than four-phase regimen involving more than two times administration of anti-progestagen, the first phase dosage will generally be the same as above, but it is possible to further decrease the dosage in the subsequent phases.

For HRT regimens dosage units comprising the anti-progestogen units as above and
15 units comprising an estrogen, progestogen or mixtures thereof are envisaged. Estrogens which can be used include 17 β -estradiol and ethinyl estradiol. Mestranol (17- α -ethinyl estradiol 3-methylether) and conjugated estrogens are also useful estrogens. Suitable amounts of ethinyl estradiol per dosage unit are between 0.005 and 0.1 mg. Of course amounts having equivalent activity of other estrogens can also be
20 used. As an approximation, 1 mg of 17 β -estradiol is equivalent in estrogenic activity to 0.015 mg of ethinyl estradiol and 0.030 mg of mestranol.

The progestogen and anti-progestogen are incorporated into dosage units for oral administration. The term "dosage unit" generally refers to physically discrete units
25 suitable as unitary dosages for humans, each containing a predetermined quantity of active material calculated to produce the desired effect, for instance tablets, pills, powders, suppositories, capsules and the like.

Methods and compositions for making such dosage units are well-known to those skilled in the art. For example, conventional techniques for making tablets and pills,
30 containing active ingredients, are described in the standard reference, Gennaro et al., Remington's Pharmaceutical Sciences, (18th ed., Mack Publishing Company, 1990, see especially Part 8: Pharmaceutical Preparations and Their Manufacture).

For making dosage units, e.g. tablets, the use of conventional additives, e.g. fillers, colorants, polymeric binders and the like is contemplated. In general any pharmaceutically acceptable additive which does not interfere with the function of the active compounds can be used in the one or more of the compositions.

- 5 Suitable carriers with which the compositions can be administered include lactose, starch, cellulose derivatives and the like used in suitable amounts. Lactose is a preferred carrier. Mixtures of carriers can also be used.

- 10 A process of manufacturing the kit of the invention comprises mixing predetermined quantities of progestogen (for instance desogestrel, 3-ketodesogestrel, or mixtures thereof) or anti-progestogen (for instance Org 31710) with predetermined quantities of excipients and converting the mixture into dosage units containing progestogen or anti-progestogen.

- 15 For HRT regimens unit dosages may be obtained by mixing predetermined quantities of progestogen and estrogen.

Preferred kits contain a total of 20 to 32 of said daily sequential dosage units.

Converting the mixture into dosage units generally involves moulding the mixture into a tablet, filling a capsule with a dried mixture, or filling a capsule with a wet mixture.

- 20 A preferred process of manufacturing the pharmaceutical product according to the invention involves incorporating the desired dosages of contraceptive steroid (for example desogestrel, 3-ketodesogestrel, or mixtures thereof) into tablets by techniques such as wet granulation tableting techniques. The package containing the dosage units will contain between 7 and 180, preferably 28, dosage units.

- 25 While the description hereinbefore refers to daily dosage units making up the anticonceptive or HRT kit, the method of treatment according to the invention extends to any other suitable form of administration. Therefore, the invention relates to pharmaceutical preparations capable of releasing progestagen and anti-progestagen, 30 optionally simultaneously, in accordance with the regimen described above. Thus the administration in four or more phases of anti-progestagen and progestagen can be effectuated not only through tablets, but also by means of an implant system, a vaginal ring, injectable systems, transdermal systems, or by any combination thereof

(particularly: the progestagen administered as a tablet, the anti-progestagen in the form of a non-oral system such as an implant).

The invention is further explained by the following illustrative examples.

5

Example I

The following coated tablets intended for once daily administration were made:

Composition (per tablet):

10

Tablet I

<u>Compound</u>	<u>Amount (mg/tablet)</u>
desogestrel	0.075
corn starch	6.500
15 povidone	1.950
stearic acid	0.650
colloidal silicone dioxide	0.650
dl- α -tocopherol	0.080
lactose	qsad 65.000

20

Coating layer: (filmcoat-dry)

<u>Compound</u>	<u>Amount (mg/tablet)</u>
hydroxypropylmethylcellulose	0.75
25 polyethylene glycol 400	0.15
titanium dioxide	0.1125
talc	0.1875

Tablet II

	<u>Compound</u>	<u>Amount (mg/tablet)</u>
	Org 31710	25
5	corn starch	6.500
	povidone	1.950
	stearic acid	0.650
	colloidal silicone dioxide	0.650
	dl- α -tocopherol	0.080
10	lactose	qsad 65.000

The same coating layer as for tablet I was used.

Tablet III

15	<u>Compound</u>	<u>Amount (mg/tablet)</u>
	Org 31710	5
	corn starch	6.500
	povidone	1.950
20	stearic acid	0.650
	colloidal silicone dioxide	0.650
	dl- α -tocopherol	0.080
	lactose	qsad 65.000

25 The same coating layer as for tablet I was used.

The tablets were packed in push-through packs as follows:

- 1 Tablet II
13 Tablets I
30 1 Tablet III
13 Tablets I

The push through packs are placed in folding cartons, which are additionally sealed in aluminum sachets.

5 Example II

The tablets of Example I, along with similar tablets containing 0.030 and 0.050 mg of desogestrel were tested in healthy female volunteers in a non-public, double-blind randomized study. Ovulation was completely inhibited with the tablets of Example I.

- 10 Furthermore, the use of the tablets of Example I also had a low percentage of bleeding and spotting days in comparison to conventional regimens of desogestrel.

The claims defining the invention are as follows:

1. A contraceptive and/or HRT (hormone replacement therapy) kit comprising sequential daily dosage units each containing as the sole contraceptively effective ingredient a progestogen, or as effective ingredient for HRT a progestogen with or without an estrogen or an estrogen only, and
 5 further two dosage units comprising an anti-progestogen when used for a multi-phase hormonal treatment cycle comprising administration of one dosage unit, comprising the anti-progestogen, at the beginning or the end of the cycle and one or more other dosage units, comprising the anti-progestogen, orderly divided through the cycle.
2. The contraceptive and/or HRT kit when used according to claim 1 wherein said kit
 10 contains a total of 20 to 32 dosage units.
3. The contraceptive and/or HRT kit when used according to claim 1 or 2 consisting of four phases, in which the first phase is a single dosage unit containing an anti-progestogen and optionally a progestogen, the second phase containing progestogen, estrogen or a mixture thereof, the third phase containing an anti-progestogen and optionally a progestogen, and the fourth phase containing
 15 progestogen, estrogen or a mixture thereof.
4. The contraceptive and/or HRT kit when used according to claim 1 consisting of four phases, in which the first phase is a single dosage unit containing an anti-progestogen and optionally a progestogen, the second phase consists of 9-15 dosage units containing progestogen, estrogen or a mixture thereof, the third phase is a single dosage unit containing an anti-progestogen and optionally
 20 a progestogen, and the fourth phase consists of 9-15 dosage units containing progestogen, estrogen or a mixture thereof.
5. A contraceptive and/or HRT kit when used according to claim 1 consisting of four to eight phases in which the odd-numbered phases all comprise a single dosage unit containing an anti-progestogen and optionally a progestogen and the even-numbered phases all consist of 8-15 dosage
 25 units containing progestogen, estrogen or a mixture thereof.
6. The contraceptive and/or HRT kit when used according to claim 3, 4 or 5, further comprising placebo dosage units.
7. The contraceptive and/or HRT kit when used according to any one of claims 3 to 6, wherein the dosage of the anti-progestogen in the third phase and, if present, subsequent phases, is
 30 lower than the dosage of the anti-progestogen in the first phase.
8. The contraceptive and/or HRT kit when used according to any one of claims 1-7, wherein the progestogen is desogestrel present in a quantity of 75µg per dosage unit and the anti-progestogen is Org 31710 present in a quantity of 25mg in the first phase and in a quantity of 2.5 to 12.5mg in the third and subsequent phases.
9. The use of daily dosage units consisting essentially a progestogen, and of an anti-progestogen, in the preparation of a drug delivery system, said drug delivery system characterized by
 consisting of daily dosage units containing only a progestogenic compound as therapeutically effective



ingredient and two to seven dosage units comprising anti-progestogen, one of which is administered at the beginning, and the other or others divided regularly throughout the cycle.

10. The use of oral daily dosage units consisting essentially of an estrogen and a progestogen, and of an anti-progestogen, in the preparation of a drug delivery system, said drug delivery system characterized by consisting of daily dosage units containing progestogen and/or estrogen as therapeutically effective ingredient and two to seven dosage units comprising an anti-progestogen, one of which is administered at the beginning and the other or others divided regularly throughout the cycle.

11. A process of manufacturing a kit according to any one of claims 1-7 comprising mixing predetermined quantities of a progestogen, and if required estrogen, with predetermined quantities of excipient and converting the mixture into dosage units each comprising the progestogen and optionally the estrogen or the mixture of the progestogen and the estrogen, and mixing predetermined quantities of an anti-progestogen with predetermined quantities of excipients and optionally predetermined quantities of a progestogen and converting the mixture into dosage units each comprising the anti-progestogen, and packaging a plurality of said dosage units into a kit.

12. A method to provide contraception or hormone replacement therapy to a woman by sequential administration of daily dosage units, at least two of which dosage units are comprising an anti-progestogen, one of which anti-progestogen comprising dosage units is administered at the beginning or the end of the sequential administration and one or more of the other anti-progestogen comprising dosage units is administered orderly divided through the sequence of administrations.

13. A contraceptive and/or hormone replacement therapy kit substantially as herein described with reference to Example I when used for a multi-phase hormonal treatment cycle comprising administration of one dosage unit, comprising the anti-progestogen, at the beginning or the end of the cycle and one or more other dosage units, comprising the anti-progestogen, orderly divided through the cycle.

14. The use according to claim 10 wherein the drug delivery system is substantially as herein described with reference to Example I.

15. A process of manufacturing a kit according to claim 1 which process is substantially as herein described with reference to Example I.

16. A kit when made by the process of claim 11 or claim 15.

Dated 14 March, 2000

Akzo Nobel N.V.

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

