



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/08/04
(87) Date publication PCT/PCT Publication Date: 2018/02/08
(85) Entrée phase nationale/National Entry: 2019/02/04
(86) N° demande PCT/PCT Application No.: US 2017/045446
(87) N° publication PCT/PCT Publication No.: 2018/027106
(30) Priorités/Priorities: 2016/08/04 (US62/370,754);
2017/07/20 (US62/534,733); 2017/08/03 (US62/540,639)

(51) Cl.Int./Int.Cl. *C07H 21/02* (2006.01),
A61K 48/00 (2006.01)
(71) Demandeur/Applicant:
ARROWHEAD PHARMACEUTICALS, INC., US
(72) Inventeurs/Inventors:
LI, ZHEN, US;
ZHU, RUI, US;
WOODDELL, CHRISTINE I., US;
GIVEN, BRUCE D., US;
PEI, TAO, US;
LEWIS, DAVID L., US;
ALMEIDA, LAUREN J., US;
...
(74) Agent: NORTON ROSE FULBRIGHT CANADA

(54) Titre : AGENT ARNI CONTRE L'INFECTION PAR LE VIRUS DE L'HEPATITE B
(54) Title: RNAI AGENTS FOR HEPATITIS B VIRUS INFECTION

(57) **Abrégé/Abstract:**

Described are compositions and methods for inhibition of Hepatitis B virus gene expression. RNA interference (RNAi) agents for inhibiting the expression of Hepatitis B virus gene are described. The HBV RNAi agents disclosed herein may be targeted to cells, such as hepatocytes, for example, by using conjugated targeting ligands. Pharmaceutical compositions comprising one or more HBV RNAi agents optionally with one or more additional therapeutics are also described. Delivery of the described HBV RNAi agents to infected liver in vivo provides for inhibition of HBV gene expression and treatment of diseases and conditions associated with HBV infection.

(72) Inventeurs(suite)/Inventors(continued): ROZEMA, DAVID B., US; WAKEFIELD, DARREN H., US

(74) Agent(suite/continued): LLP/S.E.N.C.R.L., S.R.L.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
08 February 2018 (08.02.2018)



(10) International Publication Number
WO 2018/027106 A3

(51) International Patent Classification:

C07H 21/02 (2006.01) A61K 48/00 (2006.01)

(21) International Application Number:

PCT/US2017/045446

(22) International Filing Date:

04 August 2017 (04.08.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/370,754	04 August 2016 (04.08.2016)	US
62/534,733	20 July 2017 (20.07.2017)	US
62/540,639	03 August 2017 (03.08.2017)	US

(71) Applicant: **ARROWHEAD PHARMACEUTICALS, INC.** [US/US]; 225 South Lake Avenue, Suite 1050, Pasadena, CA 91101 (US).

(72) Inventors: **LI, Zhen**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US). **ZHU, Rui**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US). **WOODDELL, Christine I.**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US). **GIVEN, Bruce D.**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US). **PEI, Tao**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US). **LEWIS, David L.**; 2930 Lakeland Avenue, Madison, WI 53704 (US). **ALMEIDA, Lauren J.**; 490 Togstad Glenn, Madison, WI 53711 (US). **ROZEMA, David B.**; 4361 Gils Way, Cross Plains, WI 53528 (US). **WAKEFIELD, Darren H.**; 5236 Stoneman Drive, Fitchburg, WI 53711 (US).

(74) Agent: **TEIGEN, Robert Michael**; Arrowhead Pharmaceuticals, Inc., 502 S. Rosa Road, Madison, WI 53719 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
— with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

15 March 2018 (15.03.2018)

(54) Title: RNAi AGENTS FOR HEPATITIS B VIRUS INFECTION

(57) Abstract: Described are compositions and methods for inhibition of Hepatitis B virus gene expression. RNA interference (RNAi) agents for inhibiting the expression of Hepatitis B virus gene are described. The HBV RNAi agents disclosed herein may be targeted to cells, such as hepatocytes, for example, by using conjugated targeting ligands. Pharmaceutical compositions comprising one or more HBV RNAi agents optionally with one or more additional therapeutics are also described. Delivery of the described HBV RNAi agents to infected liver *in vivo* provides for inhibition of HBV gene expression and treatment of diseases and conditions associated with HBV infection.



WO 2018/027106 A3

RNAi Agents for Hepatitis B Virus Infection

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority from United States Provisional Patent Application Serial No. 62/540,639, filed on August 3, 2017, United States Provisional Patent Application Serial No. 62/534,733, filed on July 20, 2017, and United States Provisional Patent Application Serial No. 62/370,754, filed on August 4, 2016, the contents of each of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

Disclosed herein are RNA interference (RNAi) agents for inhibition of Hepatitis B Virus gene expression, compositions that include HBV RNAi agents, and methods of use thereof.

BACKGROUND

The Hepatitis B Virus (HBV) is a strict hepatotropic, double-stranded DNA containing virus. Although DNA is the genetic material, the replication cycle involves a reverse transcription step to copy a pregenomic RNA into DNA. Hepatitis B Virus is classified as one member of the Hepadnaviruses and belongs to the family of Hepadnaviridae. The primary infection of adult humans with Hepatitis B Virus causes an acute hepatitis with symptoms of organ inflammation, fever, jaundice and increased liver transaminases in blood. Those patients that are not able to overcome the virus infection suffer a chronic disease progression over many years with increased risk of developing cirrhotic liver or liver cancer. Perinatal transmission from Hepatitis B Virus-infected mothers to newborns also leads to chronic hepatitis.

Upon uptake by hepatocytes, the nucleocapsid is transferred to the nucleus and DNA is released. There, the DNA strand synthesis is completed and gaps repaired to give the covalently closed circular (ccc) supercoiled DNA of 3.2kb. The cccDNA serves as a template for transcription of five major viral mRNAs, which are 3.5, 3.5, 2.4, 2.1 and 0.7 kb long. All mRNAs are 5'-capped and polyadenylated at the 3'-end. There is sequence overlap at the 3'-end between all five mRNAs.

One 3.5 kb mRNA serves as template for core protein and polymerase production. In addition, the same transcript serves as a pre-genomic replication intermediate and allows the viral

polymerase to initiate the reverse transcription into DNA. Core protein is needed for nucleocapsid formation. The other 3.5 kb mRNA encodes pre-core, the secretable e-antigen (HBeAg). In the absence of replication inhibitors, the abundance of e-antigen in blood correlates with Hepatitis B Virus replication in liver and serves as an important diagnostic marker for monitoring the disease progression.

The 2.4 and 2.1 kb mRNAs carry the open reading frames ("ORF") pre-S1, pre-S2 and S for expression of viral large, medium and small surface antigen. The s-antigen is associated with infectious, complete particles. In addition, blood of infected patients also contain non-infectious particles derived from s-antigen alone, free of genomic DNA or polymerase. The function of these particles is not fully understood. The complete and lasting depletion of detectable s-antigen in blood is considered as a reliable indicator for Hepatitis B Virus clearance.

The 0.7 kb mRNA encodes the X protein. This gene product is important for efficient transcription of viral genes and also acts as a transactivator on host gene expression. The latter activity seems to be important for hepatocyte transformation during development of liver cancer.

Patients with detectable s-antigen, e-antigen, and/or viral DNA in the blood for more than 6 months are considered chronically infected. Nucleoside analogs as inhibitors of reverse transcriptase activity are typically the first treatment option for many patients. Administration of lamivudine, tenofovir, and/or entecavir has been shown to suppress Hepatitis B Virus replication, sometimes to undetectable levels, with improvement of liver function and reduction of liver inflammation typically seen as the most important benefits. However, only few patients achieve complete and lasting remission after the end of treatment. Furthermore, the Hepatitis B Virus develops drug resistance with increasing duration of treatment. This is especially difficult for patients co-infected with Hepatitis B and Human Immunodeficiency Virus (HIV). Both viruses are susceptible to nucleoside analogue drugs and may co-develop resistance.

A second treatment option is the administration of interferon-alpha. Here, patients receive high doses of interferon-alpha over a period of 6 months. The Asian genotype B gives very poor response rates. Co-infection with Hepatitis D Virus (HDV) or Human Immunodeficiency Virus

has been shown to render interferon-alpha therapy completely ineffective. Patients with strong liver damage and heavy fibrotic conditions are not qualified for interferon-alpha therapy.

Certain Hepatitis B Virus-specific RNA interference (RNAi) agents have been previously shown to inhibit expression of HBV gene expression. For example, U.S. Patent Application Publication No. 2013/0005793, to Chin et al., which is incorporated herein by reference in its entirety, discloses certain double-stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of Hepatitis B Virus gene.

SUMMARY

There exists a need for novel Hepatitis B Virus (HBV)-specific RNA interference (RNAi) agents (also herein termed RNAi agent, RNAi trigger, or trigger) that are able to selectively and efficiently inhibit the expression of an Hepatitis B Virus (HBV) gene. Further, there exists a need for combinations of novel HBV-specific RNAi agents for the treatment of HBV infection and prevention of diseases associated with HBV.

Described herein are HBV gene-specific RNAi agents able to selectively and efficiently decrease expression of an HBV gene. The described HBV RNAi agents can be used in methods for therapeutic treatment and/or prevention of symptoms and diseases associated with HBV infection, including but not limited to chronic liver diseases/disorders, inflammations, fibrotic conditions, proliferative disorders (including cancers, such as hepatocellular carcinoma), Hepatitis D Virus (HDV) infection, and acute HBV infection. In some embodiments, the HBV RNAi agents can be used in methods for therapeutic treatment and/or prevention of symptoms and diseases associated with chronic HBV infection and/or HDV infection. Such methods comprise administration of one or more HBV RNAi agents as described herein to a subject, e.g., a human or animal subject.

Additionally, described herein are compositions comprising one or more of the disclosed HBV RNAi agents that are able to selectively and efficiently decrease expression of an HBV gene. The compositions comprising one or more HBV RNAi agents can be administered to a subject, such as a human or animal subject, for the treatment and/or prevention of symptoms and diseases associated with HBV infection.

Each HBV RNAi agent disclosed herein includes at least a sense strand and an antisense strand. The sense strand and the antisense strand can be partially, substantially, or fully complementary to each other. The length of the RNAi agent sense and antisense strands described herein each can be 16 to 30 nucleotides in length. In some embodiments, the sense and antisense strands are independently 17 to 26 nucleotides in length. In some embodiments, the sense and antisense strands are independently 19 to 26 nucleotides in length. In some embodiments, the sense and antisense strands are independently 21 to 26 nucleotides in length. In some embodiments, the sense and antisense strands are independently 21 to 24 nucleotides in length. The sense and antisense strands can be either the same length or different lengths. The HBV RNAi agents disclosed herein have been designed to include antisense strand sequences that are at least partially complementary to a sequence in the HBV genome that is conserved across the majority of known serotypes of HBV. The RNAi agents described herein, upon delivery to a cell expressing HBV, inhibit the expression of one or more HBV genes *in vivo* or *in vitro*.

15 An HBV RNAi agent includes a sense strand (also referred to as a passenger strand) that includes a first sequence, and an antisense strand (also referred to as a guide strand) that includes a second sequence. A sense strand of the HBV RNAi agents described herein includes a core stretch having at least about 85% identity to a nucleotide sequence of at least 16 consecutive nucleotides in an HBV mRNA. In some embodiments, the sense strand core nucleotide stretch having at least about 85% identity to a sequence in an HBV mRNA is 16, 17, 18, 19, 20, 21, 22, or 23 nucleotides in length. An antisense strand of an HBV RNAi agent comprises a nucleotide sequence having at least about 85% complementarity over a core stretch of at least 16 consecutive nucleotides to a sequence in an HBV mRNA and the corresponding sense strand. In some embodiments, the antisense strand core nucleotide sequence having at least about 85% complementarity to a sequence in an HBV mRNA or the corresponding sense strand is 16, 17, 18, 19, 20, 21, 22, or 23 nucleotides in length.

Examples of HBV RNAi agent sense strands and antisense strands that can be used in HBV RNAi agents are provided in Tables 3 and 4. Examples of HBV RNAi agent duplexes are provided in Table 5. Examples of 19-nucleotide core stretch sequences that consist of or are included in the sense strands and antisense strands of HBV RNAi agents disclosed herein, are provided in Table 2.

In some embodiments, one or more HBV RNAi agents are delivered to target cells or tissues using any oligonucleotide delivery technology known in the art. Nucleic acid delivery methods include, but are not limited to, by encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres, proteinaceous vectors or Dynamic Polyconjugates (DPCs) (see, for example WO 2000/053722, WO 2008/0022309, WO 2011/104169, and WO 2012/083185, each of which is incorporated herein by reference). In some embodiments, an HBV RNAi agent is delivered to target cells or tissues by covalently linking the RNAi agent to a targeting group. In some embodiments, the targeting group can include a cell receptor ligand, such as an asialoglycoprotein receptor (ASGPr) ligand. In some embodiments, an ASGPr ligand includes or consists of a galactose derivative cluster. In some embodiments, a galactose derivative cluster includes an N-acetyl-galactosamine trimer or an N-acetyl-galactosamine tetramer. In some embodiments, a galactose derivative cluster is an N-acetyl-galactosamine trimer or an N-acetyl-galactosamine tetramer.

A targeting group can be linked to the 3' or 5' end of a sense strand or an antisense strand of an HBV RNAi agent. In some embodiments, a targeting group is linked to the 3' or 5' end of the sense strand. In some embodiments, a targeting group is linked to the 5' end of the sense strand. In some embodiments, a targeting group is linked to the RNAi agent via a linker.

A targeting group, with or without a linker, can be linked to the 5' or 3' end of any of the sense and/or antisense strands disclosed in Tables 2, 3, and 4. A linker, with or without a targeting group, can be attached to the 5' or 3' end of any of the sense and/or antisense strands disclosed in Tables 2, 3, and 4.

In some embodiments, described herein are compositions that include one or more HBV RNAi agents having the duplex sequences disclosed in Table 5.

In some embodiments, described herein are compositions that include a combination or cocktail of at least two HBV RNAi agents having different nucleotide sequences. In some embodiments, the two or more different HBV RNAi agents are each separately and independently linked to targeting groups. In some embodiments, the two or more different HBV RNAi agents are each linked to targeting groups comprised of N-acetyl-galactosamines. In some embodiments, when two or more RNAi agents are included in a composition, each of the RNAi agents is linked to

the same targeting group. In some embodiments, when two or more RNAi agents are included in a composition, each of the RNAi agents is linked to different targeting groups, such as targeting groups having different chemical structures.

- 5 In some embodiments, targeting groups are linked to the HBV RNAi agents without the use of an additional linker. In some embodiments, the targeting group is designed having a linker readily present to facilitate the linkage to an HBV RNAi agent. In some embodiments, when two or more RNAi agents are included in a composition, the two or more RNAi agents may be linked to the targeting groups using the same linkers. In some embodiments, when two or more
- 10 RNAi agents are included in a composition, the two or more RNAi agents are linked to the targeting groups using different linkers.

- In some embodiments, described herein are compositions that include a combination of at least two HBV RNAi agents having different sequences, wherein each HBV RNAi agent targets a
- 15 different location or different region of an HBV gene. In some embodiments, described herein are compositions that include a combination of at least two HBV RNAi agents, wherein each HBV RNAi agent is designed to target a different HBV transcript (for example, a composition that includes two HBV RNAi agents, wherein the first HBV RNAi agent includes an antisense strand that is at least partially complementary to a nucleotide sequence located in the S ORF of
- 20 an HBV gene, while the second HBV RNAi agent includes an antisense strand that is at least partially complementary to a nucleotide sequence located in the X ORF of an HBV gene). As used herein, an RNAi agent that includes an antisense strand at least partially complementary to a nucleotide sequence located in the S ORF targets a portion of the HBV genome of SEQ ID NO:1 between positions 1-1307 and 3185-3221. As used herein, an RNAi agent that includes
- 25 an antisense strand at least partially complementary to a nucleotide sequence located in the X ORF targets a portion of the HBV genome of SEQ ID NO:1 between positions 1308-1930.

- HBV mRNA is known to be polycistronic, resulting in the translation of multiple polypeptides, and separate mRNAs overlap in RNA sequence, therefore a single RNAi agent targeting an
- 30 HBV gene may result in inhibition of most or all HBV transcripts. However, while not wishing to be bound to any theory, it is hypothesized that a composition that includes two or more HBV RNAi agents targeting different locations or regions of an HBV gene (and, in particular, two or more HBV RNAi agents wherein one HBV RNAi agent targets the S ORF and a second HBV RNAi agent targets the X ORF) may provide for additional advantages over a

composition that includes only a single HBV RNAi agent, such as (a) ensuring that all HBV viral transcripts are targeted (i.e., 3.5 kb pre-genomic RNA; 3.5 kb pre-core mRNA; 2.4 kb pre-S1 mRNA; 2.1 kb pre-S2/S mRNA; 0.7 kb X mRNA; as well as any S-antigen expressing mRNAs produced from integrated HBV DNA); (b) serving to expand the genotype coverage
 5 to potentially address a larger patient population; and/or (c) potentially decreasing the viral resistance due to mutations in the siRNA binding site.

In some embodiments, described herein are compositions that include a combination of one HBV RNAi agent that targets the S ORF of an HBV RNA (i.e., having an antisense strand that
 10 targets the S transcripts (S, pre-S1, and pre-S2), the pregenomic RNA (core and polymerase), and the pre-core transcripts (HBeAg) of an HBV genome), and one HBV RNAi agent that targets the X ORF of an HBV RNA (i.e., having an antisense strand that targets the X transcript of an HBV genome, the S transcripts (S, pre-S1, and pre-S2), the pregenomic RNA (core and polymerase), and the pre-core transcripts (HBeAg) of an HBV genome). In some embodiments,
 15 the compositions described herein include at least one HBV RNAi agent that contains a sequence that targets the S ORF of an HBV gene, and a second HBV RNAi agent that contains a sequence that targets the X ORF of an HBV gene.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising
 20 administering one or more HBV RNAi agents having an antisense strand comprising the sequence of any of the sequences in Table 3.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising administering one or more HBV RNAi agents having a sense strand comprising the sequence
 25 of any of the sequences in Table 4.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising administering one or more HBV RNAi agents having an antisense strand comprising the sequence of any of the sequences in Table 3, and a sense strand comprising the sequence of
 30 any of the sequences in Table 4 that is at least partially complementary to the antisense strand.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising administering one or more HBV RNAi agents having an antisense strand that consists of the sequence of any of the sequences in Table 3, and a sense strand that consists of the sequence

of any of the sequences in Table 4 that is at least partially complementary to the antisense strand.

Disclosed herein are methods for inhibiting expression of an HBV gene in a cell, the method comprising administering one or more HBV RNAi agents having the duplex structure of Table 5.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering one or more HBV RNAi agents having an antisense strand comprising the sequence of any of the sequences in Table 3.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering one or more HBV RNAi agents having a sense strand comprising the sequence of any of the sequences in Table 4.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering one or more HBV RNAi agents having an antisense strand comprising the sequence of any of the sequences in Table 3, and a sense strand comprising the sequence of any of the sequences in Table 4 that is at least partially complementary to the antisense strand.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering one or more HBV RNAi agents having an antisense strand that consists of the sequence of any of the sequences in Table 3, and a sense strand that consists of the sequence of any of the sequences in Table 4 that is at least partially complementary to the antisense strand.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering one or more HBV RNAi agents having the duplex structure of Table 5.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising administering (i) an HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 3, and (ii) a second HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in
5 Table 2 or Table 3.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering (i) an HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the
10 sequences in Table 2 or Table 3, and (ii) a second HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 3.

Disclosed herein are methods for inhibiting expression of an HBV gene, the method comprising administering (i) a first HBV RNAi agent having an antisense strand comprising or consisting
15 of the sequence of any of the sequences in Table 2 or Table 3 and a sense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 4 that is at least partially complementary to the antisense strand of the first HBV RNAi agent, and (ii) a second HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 3 and a sense strand comprising or consisting of the sequence of
20 any of the sequences in Table 2 or Table 4 that is at least partially complementary to the antisense strand of the second HBV RNAi agent.

Disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection, the method comprising administering (i) a first HBV
25 RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 3 and a sense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 4 that is at least partially complementary to the antisense strand of the first HBV RNAi agent, and (ii) a second HBV RNAi agent having an antisense strand comprising or consisting of the sequence of any of the sequences in Table 2 or
30 Table 3 and a sense strand comprising or consisting of the sequence of any of the sequences in Table 2 or Table 4 that is at least partially complementary to the antisense strand of the second HBV RNAi agent.

In some embodiments, an HBV RNAi agent disclosed herein comprises:

- a. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AUUGAGAGAAGUCCACCAC (SEQ ID NO: 7), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAAU (SEQ ID NO: 34); or
- b. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UUUGAGAGAAGUCCACCAC (SEQ ID NO: 8), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAA (SEQ ID NO: 35); or
- c. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AAUUGAGAGAAGUCCACCA (SEQ ID NO: 12), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAAUU (SEQ ID NO: 39); or
- d. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCA (SEQ ID NO: 13), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAUA (SEQ ID NO: 40); or
- e. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCC (SEQ ID NO: 17), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAAUUUUCU (SEQ ID NO: 44); or
- f. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCC (SEQ ID NO: 18), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAAUUUUCA (SEQ ID NO: 45); or
- g. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGC (SEQ ID NO: 22), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3

- nucleobases from the sequence (5'→3') GCUGUAGGCAUAAAUUGGU (SEQ ID NO: 49); or
- h. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGC (SEQ ID NO: 23), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GCUGUAGGCAUAAAUUGGA (SEQ ID NO: 50); or
- i. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GACCAAUUUAUGCCUACAG (SEQ ID NO: 27), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUC (SEQ ID NO: 54); or
- j. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAG (SEQ ID NO: 28), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUU (SEQ ID NO: 55); or
- k. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAG (SEQ ID NO: 29), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 56).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising an HBV RNAi agent.

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two or more HBV RNAi agents, wherein a first HBV RNAi agent comprises:

- i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AAUUGAGAGAAGUCCACCA (SEQ ID NO: 12), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAAUU (SEQ ID NO: 39); or

- ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCA (SEQ ID NO: 13), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAUA (SEQ ID NO: 40);

and wherein a second HBV RNAi agent comprises:

- i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GACCAAUUUAUGCCUACAG (SEQ ID NO: 27), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUC (SEQ ID NO: 54); or
- ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAG (SEQ ID NO: 28), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUU (SEQ ID NO: 55); or
- iii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAG (SEQ ID NO: 29), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 56).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two or more HBV RNAi agents, wherein a first HBV RNAi agent comprises:

- i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCC (SEQ ID NO: 17), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAUUUUCU (SEQ ID NO: 44); or
- ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCC (SEQ ID NO: 18), and a sense strand that comprises the nucleobase sequence differing by 0, 1,

2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAUUAUUA (SEQ ID NO: 45);

and wherein a second HBV RNAi agent comprises:

- i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GACCAAUUUAUGCCUACAG (SEQ ID NO: 27), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUC (SEQ ID NO: 54); or
- ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAG (SEQ ID NO: 28), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUU (SEQ ID NO: 55); or
- iii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAG (SEQ ID NO: 29), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 56).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two or more HBV RNAi agents, wherein a first HBV RNAi agent comprises:

- i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AAUUGAGAGAAGUCCACCA (SEQ ID NO: 12), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAUUA (SEQ ID NO: 39); or
- ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCA (SEQ ID NO: 13), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAUA (SEQ ID NO: 40);

and wherein a second HBV RNAi agent comprises an antisense strand having a sequence that is at least partially complementary to a portion of the X ORF of an HBV mRNA.

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two or more HBV RNAi agents, wherein a first HBV RNAi agent comprises:

- 5 i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCC (SEQ ID NO: 17), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAUUAUUCU (SEQ ID NO: 44); or
- 10 ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCC (SEQ ID NO: 18), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAUUAUUCU (SEQ ID NO: 45);
- 15 and wherein a second HBV RNAi agent comprises an antisense strand having a sequence that is at least partially complementary to a portion of the X ORF of an HBV mRNA:

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two or more HBV RNAi agents, wherein a first
20 HBV RNAi agent comprises an antisense strand having a sequence that is at least partially complementary to a portion of the S ORF of an HBV mRNA, and wherein a second HBV RNAi agent comprises:

- 25 i) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GACCAAUUUAUGCCUACAG (SEQ ID NO: 27), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUC (SEQ ID NO: 54); or
- 30 ii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAG (SEQ ID NO: 28), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUU (SEQ ID NO: 55); or
- iii) an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAG (SEQ ID NO: 29);

29), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 56).

- 5 In some embodiments, an HBV RNAi agent disclosed herein comprises:
- a. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGCCUUAU (SEQ ID NO: 149); or
 - b. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3
10 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGCCU (SEQ ID NO: 150); or
 - c. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGC (SEQ ID NO: 151); or
 - d. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3
15 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCCUU (SEQ ID NO: 152); or
 - e. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID
20 NO: 154); or
 - f. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACG (SEQ ID NO: 160); or
 - g. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3
25 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162); or
 - h. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCCUU (SEQ ID NO: 163); or
 - 30 i. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACGA (SEQ ID NO: 170); or

- j. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171); or
- k. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCUU (SEQ ID NO: 172); or
- l. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCCU (SEQ ID NO: 173); or
- m. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCAUU (SEQ ID NO: 174); or
- n. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175); or
- o. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCUU (SEQ ID NO: 178); or
- p. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCACUU (SEQ ID NO: 179); or
- q. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCACC (SEQ ID NO: 180); or
- r. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 181); or
- s. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCUU (SEQ ID NO: 182); or
- t. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCUU (SEQ ID NO: 183); or

- u. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCUC (SEQ ID NO: 184); or
- v. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCUU (SEQ ID NO: 185); or
- w. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCUU (SEQ ID NO: 186); or
- x. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCU (SEQ ID NO: 187); or
- y. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188); or
- z. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAGCC (SEQ ID NO: 189); or
- aa. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCU (SEQ ID NO: 190); or
- bb. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCU (SEQ ID NO: 191); or
- cc. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCG (SEQ ID NO: 192); or
- dd. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCG (SEQ ID NO: 193); or
- ee. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGG (SEQ ID NO: 194);

and wherein the HBV RNAi agent further comprises a sense strand at least partially complementary to the respective antisense strand.

In some embodiments, an HBV RNAi agent disclosed herein comprises:

- 5 a. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGCCUUAU (SEQ ID NO: 149); or
- b. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGCCU (SEQ
10 ID NO: 150); or
- c. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGC (SEQ ID
 NO: 151); or
- d. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3
15 nucleobases from the sequence (5'→3') UGAAA AUUGAGAGAAGUCCUU (SEQ ID
 NO: 152); or
- e. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID
 NO: 154); or
- 20 f. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACG (SEQ ID
 NO: 160); or
- g. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID
25 NO: 162); or
- h. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCCUU (SEQ
 ID NO: 163); or
- i. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3
30 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACGA (SEQ
 ID NO: 170); or
- j. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAA AUUGAGAGAAGUCCAC (SEQ ID
 NO: 171); or

- k. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCUU (SEQ ID NO: 172); or
- l. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCCU (SEQ ID NO: 173); or
- m. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCAUU (SEQ ID NO: 174); or
- n. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175); or
- o. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCUU (SEQ ID NO: 178); or
- p. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCACUU (SEQ ID NO: 179); or
- q. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCACC (SEQ ID NO: 180); or
- r. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 181); or
- s. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCUU (SEQ ID NO: 182); or
- t. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCUU (SEQ ID NO: 183); or
- u. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCUC (SEQ ID NO: 184); or

- v. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCUU (SEQ ID NO: 185); or
- w. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCU (SEQ ID NO: 186); or
- x. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCU (SEQ ID NO: 187); or
- y. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188); or
- z. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AACCAAUUUAUGCCUACAGCC (SEQ ID NO: 189); or
- aa. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCU (SEQ ID NO: 190); or
- bb. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCU (SEQ ID NO: 191); or
- cc. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') ACCAAUUUAUGCCUACAGCCG (SEQ ID NO: 192); or
- dd. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCCAAUUUAUGCCUACAGCCG (SEQ ID NO: 193); or.
- ee. an antisense strand that consists of the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGGG (SEQ ID NO: 194);

and wherein the HBV RNAi agent further comprises a sense strand at least partially complementary to the respective antisense strand.

In some embodiments, an HBV RNAi agent disclosed herein comprises:

- i. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscCfaAfuUfuAfuGfcCfuAfcAfgGfccsusuAu (SEQ ID NO: 61); or
- 5 ii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscCfaAfuUfuAfuGfcCfuAfcAfgGfcscsu (SEQ ID NO: 62); or
- iii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgGfccsu (SEQ ID NO: 63); or
- 10 iv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgGfsc (SEQ ID NO: 64); or
- v. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgusu (SEQ ID NO: 68); or
- 15 vi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscscaauUfuAfuGfcCfuacagcsc (SEQ ID NO: 85); or
- vii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusugagAfgAfaGfuCfcaccacsg (SEQ ID NO: 94); or
- 20 viii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgAfaGfuCfcAfcCfaCfgsa (SEQ ID NO: 98); or
- 25 ix. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuuuauGfcCfuAfcAfgcsc (SEQ ID NO: 102); or
- x. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuuuauGfcCfuAfcAfgcusu (SEQ ID NO: 103); or
- 30 xi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuuuauGfcCfuAfcAfgccsu (SEQ ID NO: 104); or

- xii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsCsCfaAfuuuauGfcCfuAfcAfgccusu (SEQ ID NO: 105); or
- 5 xiii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') cPrpusAfsCsCfaAfuUfuAfuGfcCfuAfcAfgusu (SEQ ID NO: 107); or
- xiv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') cPrpusAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg (SEQ ID NO: 108); or
- 10 xv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgagaaGfuCfcAfcCfausu (SEQ ID NO: 109); or
- xvi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgagaaGfuCfcAfcCfacsg (SEQ ID NO: 110); or
- 15 xvii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgagaaGfuCfcAfcCfacsusu (SEQ ID NO: 111); or
- xviii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgagaaGfuCfcAfcCfacsgsa (SEQ ID NO: 112); or
- 20 xix. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfsusUfgAfgagaaGfuCfcAfcCfacusu (SEQ ID NO: 120); or
- 25 xx. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu (SEQ ID NO: 125);
- xxi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc (SEQ ID NO: 126); or
- 30 xxii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asGfsasAfaAfuUfgAfgAfgAfaGfuCfcacusu (SEQ ID NO: 127); or

- xxiii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asGfsasAfaAfuUfgAfgAfaGfuCfcacsc (SEQ ID NO: 128); or
- xxiv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu (SEQ ID NO: 129); or
- xxv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc (SEQ ID NO: 130); or
- xxvi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu (SEQ ID NO: 131); or
- xxvii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu (SEQ ID NO: 132); or
- xxviii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusc (SEQ ID NO: 133); or
- xxix. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu (SEQ ID NO: 134); or
- xxx. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu (SEQ ID NO: 135); or
- xxxi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgscsc (SEQ ID NO: 136); or
- xxxii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgscsc (SEQ ID NO: 137); or
- xxxiii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgscsc (SEQ ID NO: 138); or

- xxxiv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsu (SEQ ID NO: 139); or
- xxxv. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsg (SEQ ID NO: 140); or
- xxxvi. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsc (SEQ ID NO: 141); or
- xxxvii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfUfAfuGfcCfuAfcAfgusu (SEQ ID NO: 142); or
- xxxviii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfgCfsc (SEQ ID NO: 143); or
- xxxix. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu (SEQ ID NO: 144); or
- xl. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu (SEQ ID NO: 145); or
- xli. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') asCfscAfaUfuUfaUfgCfcUfaCfaGfccsg (SEQ ID NO: 146); or
- xlii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usCfscAfaUfuUfaUfgCfcUfaCfaGfccsg (SEQ ID NO: 147); or
- xliii. an antisense strand that comprises the sequence differing by 0, 1, 2 or 3 nucleotides from the sequence (5'→3') usAfscsCfaAfuUfuAfuGfcCfuAfcAfggsg (SEQ ID NO: 148);

wherein a, g, c and u are 2'-O-methyl (2'-OMe) modified nucleotides; Af, Cf, Gf, and Uf are 2'-fluoro modified nucleotides; s is a phosphorothioate internucleoside linkage and the remaining nucleotide monomers are linked by phosphodiester bonds; and cPrpu is 5'-cyclopropyl phosphonate-2'-O-methyl modified nucleotide; and wherein the HBV RNAi agent

further comprises a sense strand at least partially complementary to the respective antisense strand.

In some embodiments, an HBV RNAi agent disclosed herein comprises:

- 5 i. an antisense strand that consists of the sequence (5'→3')
usAfscCfaAfuUfuAfuGfcCfuAfcAfgGfccsusuAu (SEQ ID NO: 61); or
- ii. an antisense strand that consists of the sequence (5'→3')
usAfscCfaAfuUfuAfuGfcCfuAfcAfgGfcscsu (SEQ ID NO: 62); or
- iii. an antisense strand that consists of the sequence (5'→3')
10 usAfscsCfaAfuUfuAfuGfcCfuAfcAfgGfccsu (SEQ ID NO: 63); or
- iv. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgGfsc (SEQ ID NO: 64); or
- v. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgusu (SEQ ID NO: 68); or
- 15 vi. an antisense strand that consists of the sequence (5'→3')
usAfscscaauUfuAfuGfcCfuacagcsc (SEQ ID NO: 85); or
- vii. an antisense strand that consists of the sequence (5'→3')
usAfsusugagAfgAfaGfuCfcaccacsg (SEQ ID NO: 94); or
- viii. an antisense strand that consists of the sequence (5'→3')
20 usAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfgsa (SEQ ID NO: 98); or
- ix. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuuuauGfcCfuAfcAfgcsc (SEQ ID NO: 102); or
- x. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuuuauGfcCfuAfcAfgcusu (SEQ ID NO: 103); or
- 25 xi. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuuuauGfcCfuAfcAfgccsu (SEQ ID NO: 104); or
- xii. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuuuauGfcCfuAfcAfgccusu (SEQ ID NO: 105); or
- xiii. an antisense strand that consists of the sequence (5'→3')
30 cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgusu (SEQ ID NO: 107); or
- xiv. an antisense strand that consists of the sequence (5'→3')
cPrpusAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg (SEQ ID NO: 108); or
- xv. an antisense strand that consists of the sequence (5'→3')
usAfsusUfgAfgagaaGfuCfcAfcCfausu (SEQ ID NO: 109); or

- xvi. an antisense strand that consists of the sequence (5'→3')
usAfsusUfgAfgagaaGfuCfcAfcCfacsg (SEQ ID NO: 110); or
- xvii. an antisense strand that consists of the sequence (5'→3')
usAfsusUfgAfgagaaGfuCfcAfcCfacusu (SEQ ID NO: 111); or
- 5 xviii. an antisense strand that consists of the sequence (5'→3')
usAfsusUfgAfgagaaGfuCfcAfcCfacsgsa (SEQ ID NO: 112); or
- xix. an antisense strand that consists of the sequence (5'→3')
usAfsusUfgAfgagaaGfuCfcAfcCfacusu (SEQ ID NO: 120); or
- xx. an antisense strand that consists of the sequence (5'→3')
10 asGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu (SEQ ID NO: 125);
- xxi. an antisense strand that consists of the sequence (5'→3')
asGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc (SEQ ID NO: 126); or
- xxii. an antisense strand that consists of the sequence (5'→3')
asGfsasAfaAfuUfgAfgAfgAfaGfuCfcacusu (SEQ ID NO: 127); or
- 15 xxiii. an antisense strand that consists of the sequence (5'→3')
asGfsasAfaAfuUfgAfgAfgAfaGfuCfcacsc (SEQ ID NO: 128); or
- xxiv. an antisense strand that consists of the sequence (5'→3')
usGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu (SEQ ID NO: 129); or
- xxv. an antisense strand that consists of the sequence (5'→3')
20 usGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc (SEQ ID NO: 130); or
- xxvi. an antisense strand that consists of the sequence (5'→3')
asCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu (SEQ ID NO: 131); or
- xxvii. an antisense strand that consists of the sequence (5'→3')
asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu (SEQ ID NO: 132); or
- 25 xxviii. an antisense strand that consists of the sequence (5'→3')
asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusc (SEQ ID NO: 133); or
- xxix. an antisense strand that consists of the sequence (5'→3')
usCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu (SEQ ID NO: 134); or
- xxx. an antisense strand that consists of the sequence (5'→3')
30 usCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu (SEQ ID NO: 135); or
- xxxi. an antisense strand that consists of the sequence (5'→3')
cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsc (SEQ ID NO: 136); or
- xxxii. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgscsc (SEQ ID NO: 137); or

- xxxiii. an antisense strand that consists of the sequence (5'→3')
cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgscsc (SEQ ID NO: 138); or
- xxxiv. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsu (SEQ ID NO: 139); or
- 5 xxxv. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsg (SEQ ID NO: 140); or
- xxxvi. an antisense strand that consists of the sequence (5'→3')
asAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsc (SEQ ID NO: 141); or
- xxxvii. an antisense strand that consists of the sequence (5'→3')
10 usAfscsCfaAfuUfUfAfuGfcCfuAfcAfgusu (SEQ ID NO: 142); or
- xxxviii. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfgCfsc (SEQ ID NO: 143); or
- xxxix. an antisense strand that consists of the sequence (5'→3')
asCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu (SEQ ID NO: 144); or
- 15 xl. an antisense strand that consists of the sequence (5'→3')
usCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu (SEQ ID NO: 145); or
- xli. an antisense strand that consists of the sequence (5'→3')
asCfscAfaUfuUfaUfgCfcUfaCfaGfccsg (SEQ ID NO: 146); or
- xlii. an antisense strand that consists of the sequence (5'→3')
20 usCfscAfaUfuUfaUfgCfcUfaCfaGfccsg (SEQ ID NO: 147); or
- xliii. an antisense strand that consists of the sequence (5'→3')
usAfscsCfaAfuUfuAfuGfcCfuAfcAfggsg (SEQ ID NO: 148);

wherein a, g, c and u are 2'-O-methyl (2'-OMe) modified nucleotides; Af, Cf, Gf, and Uf are 2'-fluoro modified nucleotides; s is a phosphorothioate internucleoside linkage and the
25 remaining nucleotide monomers are linked by phosphodiester bonds; and cPrpu is 5'-cyclopropyl phosphonate-2'-O-methyl modified nucleotide; and wherein the HBV RNAi agent further comprises a sense strand at least partially complementary to the respective antisense strand.

- 30 In some embodiments, an HBV RNAi agent disclosed herein comprises:
- a. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UUGCCUGUAGGCAUAAAUUGGUAUT (SEQ ID NO: 275); or

- b. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUAUGCCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 276); or
- c. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 278); or
- d. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGUGGUGGACUUCUCUCAAUU (SEQ ID NO: 285); or
- e. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGUGGUGGACUUCUCUCAUA (SEQ ID NO: 289); or
- f. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292); or
- g. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294); or
- h. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UCGUGGUGGACUUCUCUCAAUU (SEQ ID NO: 300); or
- i. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); or
- j. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GCUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 303); or
- k. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 304); or
- l. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UGGUGGACUUCUCUCAAUUU (SEQ ID NO: 306); or

- m. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAAUAUU (SEQ ID NO: 307); or
- 5 n. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AAUGGUGGACUUCUCUCAAUAUU (SEQ ID NO: 308); or
- o. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAAUUUUCU (SEQ ID NO: 318); or
- 10 p. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 319); or
- q. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGACUUCUCUCAAUUUUCA (SEQ ID NO: 320); or
- 15 r. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCA (SEQ ID NO: 321); or
- s. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GCUGUAGGCAUAAAUUGGU (SEQ ID NO: 322); or
- 20 t. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 323); or
- u. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GAGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 324); or
- 25 v. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GCUGUAGGCAUAAAUUGGA (SEQ ID NO: 325); or
- 30 w. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 326); or

- x. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 327); or
 - y. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328); or
 - z. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUU (SEQ ID NO: 329); or
 - aa. an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 330); or
 - bb. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 331); or
 - cc. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 332); or
 - dd. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 333); or
 - ee. a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CCCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 334);
- and wherein the HBV RNAi agent further comprises an antisense strand at least partially complementary to the respective antisense strand.

In some embodiments, an HBV RNAi agent disclosed herein comprises:

- a. a sense strand that consists of the nucleobase sequence (5'→3') UUGCCUGUAGGCAUAAAUUGGUAUT (SEQ ID NO: 275); or
- b. a sense strand that consists of the nucleobase sequence (5'→3') UUAUAGCCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 276); or
- c. a sense strand that consists of the nucleobase sequence (5'→3') CUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 278); or

- d. a sense strand that consists of the nucleobase sequence (5'→3')
CGUGGUGGACUUCUCUCAAUU (SEQ ID NO: 285); or
- e. a sense strand that consists of the nucleobase sequence (5'→3')
CGUGGUGGACUUCUCUCAUA (SEQ ID NO: 289); or
- 5 f. a sense strand that consists of the nucleobase sequence (5'→3')
CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292); or
- g. a sense strand that consists of the nucleobase sequence (5'→3')
GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294); or
- h. a sense strand that consists of the nucleobase sequence (5'→3')
10 UCGUGGUGGACUUCUCUCAAUU (SEQ ID NO: 300); or
- i. a sense strand that consists of the nucleobase sequence (5'→3')
GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); or
- j. a sense strand that consists of the nucleobase sequence (5'→3')
GCUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 303); or
- 15 k. a sense strand that consists of the nucleobase sequence (5'→3')
GGCUGUAGGCAUAAAUUGGUAUU (SEQ ID NO: 304); or
- l. a sense strand that consists of the nucleobase sequence (5'→3')
UGGUGGACUUCUCUCAUAUU (SEQ ID NO: 306); or
- m. a sense strand that consists of the nucleobase sequence (5'→3')
20 GUGGUGGACUUCUCUCAUAUU (SEQ ID NO: 307); or
- n. a sense strand that consists of the nucleobase sequence (5'→3')
AAUGGUGGACUUCUCUCAUAUU (SEQ ID NO: 308); or
- o. a sense strand that comprises the nucleobase sequence (5'→3')
GGACUUCUCUCAAUUUUCU (SEQ ID NO: 318); or
- 25 p. a sense strand that consists of the nucleobase sequence (5'→3')
GGUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 319); or
- q. a sense strand that consists of the nucleobase sequence (5'→3')
GGACUUCUCUCAAUUUUCA (SEQ ID NO: 320); or
- r. a sense strand that consists of the nucleobase sequence (5'→3')
30 GUGGACUUCUCUCAAUUUUCA (SEQ ID NO: 321); or
- s. a sense strand that consists of the nucleobase sequence (5'→3')
GCUGUAGGCAUAAAUUGGU (SEQ ID NO: 322); or
- t. a sense strand that consists of the nucleobase sequence (5'→3')
GGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 323); or

- u. a sense strand that consists of the nucleobase sequence (5'→3')
GAGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 324); or
- v. a sense strand that consists of the nucleobase sequence (5'→3')
GCUGUAGGCAUAAAUUGGA (SEQ ID NO: 325); or
- 5 w. a sense strand that consists of the nucleobase sequence (5'→3')
GGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 326); or
- x. a sense strand that consists of the nucleobase sequence (5'→3')
AGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 327); or
- y. a sense strand that consists of the nucleobase sequence (5'→3')
10 CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328); or
- z. a sense strand that consists of the nucleobase sequence (5'→3')
GGCUGUAGGCAUAAAUUGGUU (SEQ ID NO: 329); or
- aa. an antisense strand that comprises the nucleobase sequence (5'→3')
AGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 330); or
- 15 bb. a sense strand that consists of the nucleobase sequence (5'→3')
AGGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 331); or
- cc. a sense strand that consists of the nucleobase sequence (5'→3')
CGGCUGUAGGCAUAAAUUGGU (SEQ ID NO: 332); or
- dd. a sense strand that consists of the nucleobase sequence (5'→3')
20 CGGCUGUAGGCAUAAAUUGGA (SEQ ID NO: 333); or
- ee. a sense strand that consists of the nucleobase sequence (5'→3')
CCCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 334);

and wherein the HBV RNAi agent further comprises an antisense strand at least partially complementary to the respective antisense strand.

25

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAUAUU (SEQ ID NO: 307); and wherein a second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID NO: 154), and a sense strand that comprises the

nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175), and a sense strand that consists of the nucleobase sequence (5'→3') GUGGUGGACUUCUCUCAUAU (SEQ ID NO: 307); and wherein a second HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID NO: 154), and a sense strand that consists of the nucleobase sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that consists of the nucleobase sequence (5'→3') GUGGACUUCUCUCAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188), and a sense strand that consists of the nucleobase sequence (5'→3') CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that consists of the nucleobase sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that consists of the nucleobase sequence (5'→3') GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAUAUU (SEQ ID NO: 307).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein a first HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3')
 5 AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that consists of the nucleobase sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein a second HBV RNAi agent comprises an antisense strand that consists of the nucleobase sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that consists of the nucleobase sequence (5'→3')
 10 GUGGUGGACUUCUCUCAUAU (SEQ ID NO: 307).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the
 15 nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence
 20 (5'→3') GUGGUGGACUUCUCUCAUAU (SEQ ID NO: 307); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID NO: 154), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3')
 25 CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the
 30 nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence

(5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the

nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAUAU (SEQ ID NO: 307).

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UAUUGAGAGAAGUCCACCACUU (SEQ ID NO: 175), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAUAU (SEQ ID NO: 307); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGUU (SEQ ID NO: 154), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CUGUAGGCAUAAAUUGGUA (SEQ ID NO: 292), and wherein the sense strand of the first HBV RNAi agent and the second HBV RNAi agent are conjugated to a targeting ligand comprising N-acetyl-galactosamine.

20

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCG (SEQ ID NO: 188), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') CGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 328), and wherein the sense strand of the

30

first HBV RNAi agent and the second HBV RNAi agent are conjugated to a targeting ligand comprising N-acetyl-galactosamine.

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GGCUGUAGGCAUAAAUUGGUA (SEQ ID NO: 294), and wherein the sense strand of the first HBV RNAi agent and the second HBV RNAi agent are conjugated to a targeting ligand comprising N-acetyl-galactosamine.

20

In some embodiments, disclosed herein are compositions for inhibiting expression of an HBV gene in a cell, the composition comprising two HBV RNAi agents, wherein all or substantially all of the nucleotides in the sense strand are modified and/or all or substantially all of the nucleotides in the antisense strand in the first and/or second HBV RNAi agent are modified nucleotides, and wherein the first HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') AGAAAAUUGAGAGAAGUCCAC (SEQ ID NO: 171), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGACUUCUCUCAAUUUUCU (SEQ ID NO: 302); and wherein the second HBV RNAi agent comprises an antisense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') UACCAAUUUAUGCCUACAGCC (SEQ ID NO: 162), and a sense strand that comprises the nucleobase sequence differing by 0, 1, 2 or 3 nucleobases from the sequence (5'→3') GUGGUGGACUUCUCUCAUAUU (SEQ ID NO: 307), and wherein the sense strand of the

first HBV RNAi agent and the second HBV RNAi agent are conjugated to a targeting ligand comprising N-acetyl-galactosamine.

5 In some embodiments, disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection comprising administering to a subject in need thereof an effective amount of AD04872 and an effective amount of AD05070. In some embodiments, the ratio of AD04872 to AD05070 administered to a subject in need thereof is about 2:1. In some embodiments, the ratio of AD04872 to AD05070 administered to a subject in need thereof is about 3:1. In some embodiments, the ratio of AD04872 to AD05070
10 administered to a subject in need thereof is about 1:1. In some embodiments, the ratio of AD04872 to AD05070 administered to a subject in need thereof is about 4:1. In some embodiments, the ratio of AD04872 to AD05070 administered to a subject in need thereof is about 5:1. In some embodiments, the ratio of AD04872 to AD05070 administered to a subject in need thereof is about 1:2.

15 In some embodiments, about 1 mg/kg (mpk) of AD04872 and about 1 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 1.5 mg/kg of AD04872 and about 1.5 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 2.0 mg/kg of AD04872 and about 1.0 mg/kg of AD05070 are administered
20 to a subject in need thereof. In some embodiments, about 3.0 mg/kg of AD04872 and about 1.0 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 3.2 mg/kg of AD04872 and about 0.8 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 2.7 mg/kg of AD04872 and about 1.3 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 4.0 mg/kg
25 of AD04872 and about 1.0 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about 3.3 mg/kg of AD04872 and about 1.7 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, between about 0.05 and about 5 mg/kg of AD04872 and between about 0.05 and about 5 mg/kg of AD05070 are administered to a subject in need thereof. In some embodiments, about AD04872 and about AD05070 are
30 administered separately (e.g., in separate injections). In some embodiments, the respective dose of AD04872 and the respective dose of AD05070 are administered together (e.g., in the same injection). In some embodiments, the respective dose of AD04872 and the respective dose of AD05070 are prepared in a single pharmaceutical composition.

In some embodiments, disclosed herein are methods of treatment of an HBV infection or prevention of diseases or symptoms caused by an HBV infection comprising administering to a subject in need thereof an effective amount of AD04872 and an effective amount of AD04776. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject
 5 in need thereof is about 2:1. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject in need thereof is about 3:1. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject in need thereof is about 4:1. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject in need thereof is about 1:1. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject
 10 in need thereof is 5:1. In some embodiments, the ratio of AD04872 to AD04776 administered to a subject in need thereof is 1:2.

In some embodiments, about 1 mg/kg (mpk) of AD04872 and about 1 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 1.5 mg/kg of AD04872
 15 and about 1.5 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 2.0 mg/kg of AD04872 and about 1.0 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 3.0 mg/kg of AD04872 and about 1.0 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 3.2 mg/kg of AD04872 and about 0.8 mg/kg of AD04776 are administered to a subject
 20 in need thereof. In some embodiments, about 2.7 mg/kg of AD04872 and about 1.3 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 4.0 mg/kg of AD04872 and about 1.0 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, about 3.3 mg/kg of AD04872 and about 1.7 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, between about 0.05 and
 25 about 5 mg/kg of AD04872 and between about 0.05 and about 5 mg/kg of AD04776 are administered to a subject in need thereof. In some embodiments, the respective doses of AD04872 and AD04776 are administered separately (e.g., in separate injections). In some embodiments, the respective doses of AD04872 and AD04776 are administered together (e.g., in the same injection). In some embodiments, the respective doses of AD04872 and AD04776
 30 are prepared in a single pharmaceutical composition.

In some embodiments, disclosed herein are methods of treatment of an HBV infection or prevention of disease or symptoms caused by an HBV infection comprising administering to a subject in need thereof an effective amount of AD04872 and an effective amount of AD04982.

In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is about 2:1. In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is about 3:1. In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is about 4:1. In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is about 1:1. In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is about 5:1. In some embodiments, the ratio of AD04872 to AD04982 administered to a subject in need thereof is 1:2.

10 In some embodiments, about 1 mg/kg (mpk) of AD04872 and about 1 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 1.5 mg/kg of AD04872 and about 1.5 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 2.0 mg/kg of AD04872 and about 1.0 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 3.0 mg/kg of AD04872 and about 1.0

15 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 3.2 mg/kg of AD04872 and about 0.8 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 2.7 mg/kg of AD04872 and about 1.3 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, about 4.0 mg/kg of AD04872 and about 1.0 mg/kg of AD04982 are administered to a subject in need thereof. In

20 some embodiments, about 3.3 mg/kg of AD04872 and about 1.7 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, between about 0.05 and about 5 mg/kg of AD04872 and between about 0.05 and about 5 mg/kg of AD04982 are administered to a subject in need thereof. In some embodiments, the respective doses of AD04872 and AD04982 are administered separately (e.g., in separate injections). In some embodiments, the


25 respective doses of AD04872 and AD04982 are administered together (e.g., in the same injection). In some embodiments, the respective doses of AD04872 and AD04982 are prepared in a single pharmaceutical composition.


In some embodiments, disclosed herein are methods of treatment of an HBV infection or


30 prevention of disease or symptoms caused by an HBV infection comprising administering to a subject in need thereof an effective amount of AD04580 and an effective amount of AD04585. In some embodiments, the ratio of AD04580 to AD04585 administered to a subject in need thereof is about 2:1. In some embodiments, the ratio of AD04580 to AD04585 administered to a subject in need thereof is about 3:1. In some embodiments, the ratio of AD04580 to AD04585

administered to a subject in need thereof is about 4:1. In some embodiments, the ratio of AD04580 to AD04585 administered to a subject in need thereof is about 5:1. In some embodiments, the ratio of AD04580 to AD04585 administered to a subject in need thereof is about 1:1. In some embodiments, the ratio of AD04580 to AD04585 administered to a subject in need thereof is about 1:2. In some embodiments, about 1 mg/kg (mpk) of AD04580 and about 1 mg/kg of AD04585 are administered to a subject in need thereof. In some embodiments, about 1.5 mg/kg of AD04580 and about 1.5 mg/kg of AD04585 are administered to a subject in need thereof. In some embodiments, between about 0.05 and about 5 mg/kg of AD04580 and between about 0.05 and about 5 mg/kg of AD04585 are administered to a subject in need thereof.


In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD05070 linked to (NAG37)s shown as a sodium salt having the structure represented by the following:










In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD05070 linked to (NAG25)s shown as a sodium salt having the structure represented by the following:













In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD05070 linked to (NAG37)s shown as a free acid having the structure represented by the following:













In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD04580 linked to (NAG31)s shown as a sodium salt having the structure represented by the following:













In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD04585 linked to (NAG25)s shown as a sodium salt having the structure represented by the following:













In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD04872 linked to (NAG37)s shown as a sodium salt having the structure represented by the following:













In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD04872 linked to (NAG25)s shown as a sodium salt having the structure represented by the following:












In some embodiments, an HBV RNAi agent disclosed herein consists of or comprises AD04872 linked to (NAG37)s shown as a free acid having the structure represented by the following:









In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or more additional (i.e., second, third, etc.) therapeutics. A second therapeutic can be another HBV RNAi agent (e.g., a HBV RNAi agent which targets a different sequence within an HBV genome). An additional therapeutic can also be a small molecule drug, antibody, antibody
5 fragment, and/or vaccine. The HBV RNAi agents, with or without the one or more additional therapeutics, can be combined with one or more excipients to form pharmaceutical compositions.

In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or
10 more additional therapeutics, wherein the additional therapeutic is a nucleoside inhibitor or nucleotide inhibitor. In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or more additional therapeutics, wherein the additional therapeutic entecavir, tenofovir, tenofovir alafenamide, tenofovir disoproxil, lamivudine, or another antiviral therapeutic. In some embodiments, the described HBV RNAi agent(s) are optionally
15 combined with one or more additional therapeutics, wherein the additional therapeutic is an interferon. In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or more additional therapeutics, wherein the additional therapeutic is interferon-alpha. In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or more HBV additional therapeutics, wherein the additional therapeutic is an HBV vaccine.

20

In some embodiments, the described HBV RNAi agent(s) are optionally combined with one or more additional therapeutics in a single dosage form (i.e., a cocktail included in a single injection). In some embodiments, the described HBV RNAi agent(s) may be administered separately from one or more optional additional therapeutics. In some embodiments, the
25 described HBV RNAi agent(s) are administered to a subject in need thereof via subcutaneous injection, and the one or more optional additional therapeutics are administered orally, which together provide for a treatment regimen for diseases and conditions associated with HBV infection. In some embodiments, the described HBV RNAi agent(s) are administered to a subject in need thereof via subcutaneous injection, and the one or more optional additional
30 therapeutics are administered via a separate subcutaneous injection.

In some embodiments, disclosed herein are compositions for delivering an HBV RNAi agent to a liver cell *in vivo*, the composition including an HBV RNAi agent conjugated or linked to a targeting group. In some embodiments, the targeting group is an asialoglycoprotein receptor

ligand. In some embodiments, compositions for delivering an HBV RNAi agent to a liver cell *in vivo* are described, the composition including an HBV RNAi agent linked to an N-acetyl-galactosamine targeting ligand.

- 5 In some embodiments, one or more of the described HBV RNAi agents are administered to a mammal in a pharmaceutically acceptable carrier or diluent. In some embodiments, the mammal is a human.

The use of Hepatitis B Virus RNAi agent(s) provides methods for therapeutic and/or
10 prophylactic treatment of diseases/disorders which are associated with HBV infection. The described HBV RNAi agents mediate RNA interference to inhibit the expression of one or more genes necessary for replication and/or pathogenesis of Hepatitis B Virus. In particular, for example, HBV RNAi agents may inhibit viral polymerase, core protein, surface antigen, e-antigen and/or the X protein, in a cell, tissue or mammal. HBV RNAi agents can be used to
15 treat HBV infection. HBV RNAi agents can also be used to treat or prevent chronic liver diseases/disorders, inflammations, fibrotic conditions and proliferative disorders, like cancers, associated with HBV infection. In some embodiments, the methods further comprise treatment of Hepatitis D Virus (HDV) in the subject. Such methods comprise administration of HBV RNAi agent to a human being or animal infected with HBV. Further, compositions for delivery
20 of HBV RNAi agents to liver cells *in vivo* are described.

The pharmaceutical compositions comprising one or more HBV RNAi agents can be administered in a number of ways depending upon whether local or systemic treatment is desired. Administration can be, but is not limited to, intravenous, intraarterial, subcutaneous,
25 intraperitoneal, subdermal (e.g., via an implanted device), and intraparenchymal administration. In some embodiments, the pharmaceutical compositions described herein are administered by subcutaneous injection.

The described HBV RNAi agents and/or compositions can be used in methods for therapeutic
30 treatment of HBV infection or disease or conditions caused by HBV infection. Such methods include administration of an HBV RNAi agent as described herein to a subject, e.g., a human or animal subject.

As used herein, the terms “oligonucleotide” and “polynucleotide” mean a polymer of linked nucleosides each of which can be independently modified or unmodified.

As used herein, an “RNAi agent” or “RNAi trigger” means a composition that contains an RNA or RNA-like (e.g., chemically modified RNA) oligonucleotide molecule that is capable of degrading or inhibiting translation of messenger RNA (mRNA) transcripts of a target mRNA in a sequence specific manner. As used herein, RNAi agents may operate through the RNA interference mechanism (i.e., inducing RNA interference through interaction with the RNA interference pathway machinery (RNA-induced silencing complex or RISC) of mammalian cells), or by any alternative mechanism(s) or pathway(s). While it is believed that RNAi agents, as that term is used herein, operate primarily through the RNA interference mechanism, the disclosed RNAi agents are not bound by or limited to any particular pathway or mechanism of action. RNAi agents disclosed herein are comprised of a sense strand and an antisense strand, and include, but are not limited to: short interfering RNAs (siRNAs), double-stranded RNAs (dsRNA), micro RNAs (miRNAs), short hairpin RNAs (shRNA), and dicer substrates. The antisense strand of the RNAi agents described herein is at least partially complementary to the mRNA being targeted. RNAi agents may be comprised of modified nucleotides and/or one or more non-phosphodiester linkages.

As used herein, the terms “silence,” “reduce,” “inhibit,” “down-regulate,” or “knockdown” when referring to expression of a given gene, mean that the expression of the gene, as measured by the level of RNA transcribed from the gene or the level of polypeptide, protein or protein subunit translated from the mRNA in a cell, group of cells, tissue, organ, or subject in which the gene is transcribed, is reduced when the cell, group of cells, tissue, organ, or subject is treated with oligomeric compounds, such as RNAi agents, described herein as compared to a second cell, group of cells, tissue, organ, or subject that has not or have not been so treated.

As used herein, the term “sequence” or “nucleotide sequence” mean a succession or order of nucleobases or nucleotides, described with a succession of letters using standard nomenclature.

As used herein, a “nucleotide base,” or “nucleobase” is a heterocyclic pyrimidine or purine compound, which is a standard constituent of all nucleic acids, and includes the bases that form the nucleotides adenine (A), guanine (G), cytosine (C), thymine (T), and uracil (U). A

nucleobase may further be modified to include, without limitation, universal bases, hydrophobic bases, promiscuous bases, size-expanded bases, and fluorinated bases.

As used herein, and unless otherwise indicated, the term “complementary,” when used to describe a first nucleotide sequence (*e.g.*, RNAi agent sense strand or targeted mRNA) in relation to a second nucleotide sequence (*e.g.*, RNAi agent antisense strand or a single-stranded antisense oligonucleotide), means the ability of an oligonucleotide or polynucleotide including the first nucleotide sequence to hybridize (form base pair hydrogen bonds under mammalian physiological conditions (or similar conditions *in vitro*)) and form a duplex or double helical structure under certain conditions with an oligonucleotide or polynucleotide including the second nucleotide sequence. Complementary sequences include Watson-Crick base pairs or non-Watson-Crick base pairs and include natural or modified nucleotides or nucleotide mimics, at least to the extent that the above hybridization requirements are fulfilled. Sequence identity or complementarity is independent of modification. For example, a and Af are complementary to U (or T) and identical to A for the purposes of determining identity or complementarity.

As used herein, “perfectly complementary” or “fully complementary” means that all (100%) of the bases in a contiguous sequence of a first polynucleotide will hybridize with the same number of bases in a contiguous sequence of a second polynucleotide. The contiguous sequence may comprise all or a part of a first or second nucleotide sequence.

As used herein, “partially complementary” means that in a hybridized pair of nucleobase sequences, at least 70%, but not all, of the bases in a contiguous sequence of a first polynucleotide will hybridize with the same number of bases in a contiguous sequence of a second polynucleotide.

As used herein, “substantially complementary” means that in a hybridized pair of nucleobase sequences, at least about 85%, but not all, of the bases in a contiguous sequence of a first polynucleotide will hybridize with the same number of bases in a contiguous sequence of a second polynucleotide. The terms “complementary,” “fully complementary,” and “substantially complementary” herein may be used with respect to the base matching between the sense strand and the antisense strand of a double-stranded RNAi agent, between the antisense strand of an RNAi agent and a sequence of a target mRNA, or between a single-stranded antisense oligonucleotide and a sequence of a target mRNA.


As used herein, the term “substantially identical” or “substantially identity” as applied to nucleic acid sequence means that a nucleic acid sequence comprises a sequence that has at least about 85% sequence identity or more, preferably at least 90%, at least 95%, or at least 99%, compared to a reference sequence. Percentage of sequence identity is determined by comparing two optimally aligned sequences over a comparison window. The percentage is calculated by determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. The inventions disclosed herein encompasses nucleotide sequences substantially identical to those disclosed herein, e.g., in Tables 2, 3, and 4. In some embodiments, the sequences disclosed herein are exactly identical, or at least about 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% percent identical to those disclosed herein, e.g., in Tables 1, 2, 3 and 4.

As used herein, the terms “treat,” “treatment,” and the like, mean the methods or steps taken to provide relief from or alleviation of the number, severity, and/or frequency of one or more symptoms of a disease or condition in a subject.

20

As used herein, the phrase “introducing into a cell,” when referring to an oligomeric compound, means functionally delivering the oligomeric compound into a cell. The phrase “functional delivery,” means that delivering the oligomeric compound to the cell in a manner that enables the oligomeric compound to have the expected biological activity, e.g., sequence-specific inhibition of gene expression.

25

Unless stated otherwise, use of the symbol  as used herein means that any group or groups may be linked thereto that is in accordance with the scope of the inventions described herein.

As used herein, the term “isomers” refers to compounds that have identical molecular formulae, but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers.” Stereoisomers that are not mirror images of one another are termed “diastereoisomers,” and stereoisomers that are non-superimposable mirror images are termed

30

“enantiomers,” or sometimes optical isomers. A carbon atom bonded to four non-identical substituents is termed a “chiral center.”

As used herein, unless specifically identified in a structure as having a particular conformation,
5 for each structure in which asymmetric centers are present and thus give rise to enantiomers, diastereomers, or other stereoisomeric configurations, each structure disclosed herein is intended to represent all such possible isomers, including their optically pure and racemic forms. For example, the structures disclosed herein are intended to cover mixtures of diastereomers as well as single stereoisomers.

10

As used in a claim herein, the phrase “consisting of” excludes any element, step, or ingredient not specified in the claim. When used in a claim herein, the phrase “consisting essentially of” limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s) of the claimed invention.

15

The person of ordinary skill in the art would readily understand and appreciate that the compounds and compositions disclosed herein may have certain atoms (e.g., N, O, or S atoms) in a protonated or deprotonated state, depending upon the environment in which the compound or composition is placed. Accordingly, as used herein, the structures disclosed herein envisage
20 that certain functional groups, such as, for example, OH, SH, or NH, may be protonated or deprotonated. The disclosure herein is intended to cover the disclosed compounds and compositions regardless of their state of protonation based on the environment (such as pH), as would be readily understood by the person of ordinary skill in the art.

25 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in
30 their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

DETAILED DESCRIPTION

5 Described herein are RNAi agents for inhibiting expression of Hepatitis B Virus (HBV) (referred to herein as HBV RNAi agents or HBV RNAi triggers). Each HBV RNAi agent comprises a sense strand and an antisense strand. The sense strand and the antisense strand each can be 16 to 30 nucleotides in length. In some embodiments, the sense and antisense strands each can be 17 to 26 nucleotides in length. The sense and antisense strands can be either
10 the same length or they can be different lengths. In some embodiments, the sense and antisense strands are each independently 17 to 26 nucleotides in length. In some embodiments, the sense and antisense strands are each independently 17-21 nucleotides in length. In some embodiments, both the sense and antisense strands are each 21-26 nucleotides in length. In some embodiments, the sense strand is about 19 nucleotides in length while the antisense strand
15 is about 21 nucleotides in length. In some embodiments, the sense strand is about 21 nucleotides in length while the antisense strand is about 23 nucleotides in length. In some embodiments, both the sense and antisense strands are each 26 nucleotides in length. In some embodiments, the RNAi agent sense and antisense strands are each independently 17, 18, 19, 20, 21, 22, 23, 24, 25, or 26 nucleotides in length. In some embodiments, a double-stranded
20 RNAi agent has a duplex length of about 16, 17, 18, 19, 20, 21, 22, 23 or 24 nucleotides. This region of perfect or substantial complementarity between the sense strand and the antisense strand is typically 15-25 (e.g., 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25) nucleotides in length and occurs at or near the 5' end of the antisense strand (e.g., this region may be separated from the 5' end of the antisense strand by 0, 1, 2, 3, or 4 nucleotides that are not perfectly or
25 substantially complementary).

The sense strand and antisense strand each contain a core stretch sequence that is 16 to 23 nucleobases in length. An antisense strand core stretch sequence is 100% (perfectly) complementary or at least about 85% (substantially) complementary to a nucleotide sequence
30 (sometimes referred to, e.g., as a target sequence) present in the HBV mRNA target. A sense strand core stretch sequence is 100% (perfectly) complementary or at least about 85% (substantially) complementary to a core stretch sequence in the antisense strand, and thus the sense strand core stretch sequence is perfectly identical or at least about 85% identical to a

nucleotide sequence (target sequence) present in the HBV mRNA target. A sense strand core stretch sequence can be the same length as a corresponding antisense core sequence or it can be a different length. In some embodiments, the antisense strand core stretch sequence is 16, 17, 18, 19, 20, 21, 22, or 23 nucleotides in length. In some embodiments, the sense strand core stretch sequence is 16, 17, 18, 19, 20, 21, 22, or 23 nucleotides in length.

Examples of sense and antisense strand nucleotide sequences used in forming HBV RNAi agents are provided in Tables 3 and 4. Examples of RNAi agent duplexes, that include the nucleotide sequences in Tables 3 and 4, are provided in Table 5.

10

The HBV RNAi agent sense and antisense strands anneal to form a duplex. A sense strand and an antisense strand of an HBV RNAi agent may be partially, substantially, or fully complementary to each other. Within the complementary duplex region, the sense strand core stretch sequence is at least about 85% complementary or 100% complementary to the antisense core stretch sequence. In some embodiments, the sense strand core stretch sequence contains a sequence of at least 16, at least 17, at least 18, at least 19, at least 20, or at least 21 nucleotides that is at least about 85% or 100% complementary to a corresponding 16, 17, 18, 19, 20, or 21 nucleotide sequence of the antisense strand core stretch sequence (i.e., the sense strand and antisense core stretch sequences of an HBV RNAi agent have a region of at least 16, at least 17, at least 18, at least 19, at least 20, or at least 21 nucleotides that is at least 85% base paired or 100% base paired.).

20

In some embodiments, the antisense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the antisense strand sequences in Table 2 or Table 3. In some embodiments, the sense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the sense strand sequences in Table 2 or Table 4.

25

The length of the HBV RNAi agent sense and antisense strands described herein are independently 16 to 30 nucleotides in length. In some embodiments, the sense and antisense strands are independently 17 to 26 nucleotides in length. In some embodiments, the sense and antisense strands are 19-26 nucleotides in length. In some embodiments, the described RNAi agent sense and antisense strands are independently 17, 18, 19, 20, 21, 22, 23, 24, 25, or 26 nucleotides in length. The sense and antisense strands can be either the same length or they can be different lengths. In some embodiments, a sense strand and an antisense strand are each 26

30

nucleotides in length. In some embodiments, a sense strand is 23 nucleotides in length and an antisense strand is 21 nucleotides in length. In some embodiments, a sense strand is 22 nucleotides in length and an antisense strand is 21 nucleotides in length. In some embodiments, a sense strand is 21 nucleotides in length and an antisense strand is 21 nucleotides in length. In some embodiments, a sense strand is 19 nucleotides in length and an antisense strand is 21 nucleotides in length.

The sense strand and/or the antisense strand may optionally and independently contain an additional 1, 2, 3, 4, 5, or 6 nucleotides (extension) at the 3' end, the 5' end, or both the 3' and 5' ends of the core sequences. The antisense strand additional nucleotides, if present, may or may not be complementary to the corresponding sequence in an HBV mRNA. The sense strand additional nucleotides, if present, may or may not be identical to the corresponding sequence in an HBV mRNA. The antisense strand additional nucleotides, if present, may or may not be complementary to the corresponding sense strand's additional nucleotides, if present.

As used herein, an extension comprises 1, 2, 3, 4, 5, or 6 nucleotides at the 5' and/or 3' end of the sense strand core stretch sequence and/or antisense strand core stretch sequence. The extension nucleotides on a sense strand may or may not be complementary to nucleotides, either core stretch sequence nucleotides or extension nucleotides, in the corresponding antisense strand. Conversely, the extension nucleotides on an antisense strand may or may not be complementary to nucleotides, either core stretch sequence nucleotides or extension nucleotides, in the corresponding sense strand. In some embodiments, both the sense strand and the antisense strand of an RNAi agent contain 3' and 5' extensions. In some embodiments, one or more of the 3' extension nucleotides of one strand base pairs with one or more 5' extension nucleotides of the other strand. In other embodiments, one or more of 3' extension nucleotides of one strand do not base pair with one or more 5' extension nucleotides of the other strand. In some embodiments, an HBV RNAi agent has an antisense strand having a 3' extension and a sense strand having a 5' extension.

In some embodiments, an HBV RNAi agent comprises an antisense strand having a 3' extension of 1, 2, 3, 4, 5, or 6 nucleotides in length. In other embodiments, an HBV RNAi agent comprises an antisense strand having a 3' extension of 1, 2, or 3 nucleotides in length. In some embodiments, one or more of the antisense strand extension nucleotides comprise uracil or thymidine nucleotides or nucleotides which are complementary to a corresponding HBV

mRNA sequence. In some embodiments, a 3' antisense strand extension includes or consists of, but is not limited to: AUA, UGCUU, CUG, UG, UGCC, CUGCC, CGU, CUU, UGCCUA, CUGCCU, UGCCU, UGAUU, GCCUAU, T, TT, U, UU (each listed 5' to 3').

- 5 In some embodiments, the 3' end of the antisense strand may include additional abasic nucleosides (Ab). In some embodiments, Ab or AbAb may be added to the 3' end of the antisense strand.

10 In some embodiments, an HBV RNAi agent comprises an antisense strand having a 5' extension of 1, 2, 3, 4, or 5 nucleotides in length. In other embodiments, an HBV RNAi agent comprises an antisense strand having a 5' extension of 1 or 2 nucleotides in length. In some embodiments, one or more of the antisense strand extension nucleotides comprises uracil or thymidine nucleotides or nucleotides which are complementary to a corresponding HBV mRNA sequence. In some embodiments, the 5' antisense strand extension includes or consists of, but is no limited to, UA, TU, U, T, UU, TT, CUC (each listed 5' to 3'). An antisense strand
15 may have any of the 3' extensions described above in combination with any of the 5' antisense strand extensions described, if present.

In some embodiments, an HBV RNAi agent comprises a sense strand having a 3' extension of
20 1, 2, 3, 4, or 5 nucleotides in length. In some embodiments, one or more of the sense strand extension nucleotides comprises adenosine, uracil, or thymidine nucleotides, AT dinucleotide, or nucleotides which correspond to nucleotides in the HBV mRNA sequence. In some embodiments, the 3' sense strand extension includes or consists of, but is not limited to: T, UT, TT, UU, UUT, TTT, or TTTT (each listed 5' to 3').

25

In some embodiments, the 3' end of the sense strand may include additional abasic nucleosides. In some embodiments, UUAb, UAb, or Ab may be added to the 3' end of the sense strand. In some embodiments, the one or more abasic nucleosides added to the 3' end of the sense strand may be inverted (invAb). In some embodiments, one or more inverted abasic nucleosides may
30 be inserted between the targeting ligand and the nucleobase sequence of the sense strand of the RNAi agent. In some embodiments, the inclusion of one or more inverted abasic nucleosides at or near the terminal end or terminal ends of the sense strand of an RNAi agent may allow for enhanced activity or other desired properties of an RNAi agent.

In some embodiments, an HBV RNAi agent comprises a sense strand having a 5' extension of 1, 2, 3, 4, 5, or 6 nucleotides in length. In some embodiments, one or more of the sense strand extension nucleotides comprise uracil or adenosine nucleotides or nucleotides which correspond to nucleotides in the HBV mRNA sequence. In some embodiments, the sense strand 5' extension can be, but is not limited to: CA, AUAGGC, AUAGG, AUAG, AUA, A, AA, AC, GCA, GGCA, GGC, UAUCA, UAUC, UCA, UAU, U, UU (each listed 5' to 3'). A sense strand may have a 3' extension and/or a 5' extension.

In some embodiments, the 5' end of the sense strand may include an additional abasic nucleoside (Ab) or nucleosides (AbAb). In some embodiments, the one or more abasic nucleosides added to the 5' end of the sense strand may be inverted (invAb). In some embodiments, one or more inverted abasic nucleosides may be inserted between the targeting ligand and the nucleobase sequence of the sense strand of the RNAi agent. In some embodiments, the inclusion of one or more inverted abasic nucleosides at or near the terminal end or terminal ends of the sense strand of an RNAi agent may allow for enhanced activity or other desired properties of an RNAi agent.

Examples of nucleotide sequences used in forming HBV RNAi agents are provided in Tables 3 and 4. In some embodiments, an HBV RNAi agent antisense strand includes a nucleotide sequence of any of the sequences in Table 3. In some embodiments, an HBV RNAi agent antisense strand includes the sequence of nucleotides 1-17, 2-15, 2-17, 1-18, 2-18, 1-19, 2-19, 1-20, 2-20, 1-21, 2-21, 1-22, 2-22, 1-23, 2-23, 1-24, 2-24, 1-25, 2-25, 1-26, or 2-26 of any of the sequences in Table 3. In some embodiments, an HBV RNAi agent sense strand includes the nucleotide sequence of any of the sequences in Table 4. In some embodiments, an HBV RNAi agent sense strand includes the sequence of nucleotides 1-18, 1-19, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-26, 2-19, 2-20, 2-21, 2-22, 2-23, 2-24, 2-25, 2-26, 3-20, 3-21, 3-22, 3-23, 3-24, 3-25, 3-26, 4-21, 4-22, 4-23, 4-24, 4-25, 4-26, 5-22, 5-23, 5-24, 5-25, 5-26, 6-23, 6-24, 6-25, 6-26, 7-24, 7-25, 7-25, 8-25, 8-26 of any of the sequences in Table 4.

In some embodiments, the sense and antisense strands of the RNAi agents described herein contain the same number of nucleotides. In some embodiments, the sense and antisense strands of the RNAi agents described herein contain different numbers of nucleotides. In some embodiments, the sense strand 5' end and the antisense strand 3' end of an RNAi agent form a blunt end. In some embodiments, the sense strand 3' end and the antisense strand 5' end of an

RNAi agent form a blunt end. In some embodiments, both ends of an RNAi agent form blunt ends. In some embodiments, neither end of an RNAi agent is blunt-ended. As used herein a blunt end refers to an end of a double stranded RNAi agent in which the terminal nucleotides of the two annealed strands are complementary (form a complementary base-pair). In some

5 embodiments, the sense strand 5' end and the antisense strand 3' end of an RNAi agent form a frayed end. In some embodiments, the sense strand 3' end and the antisense strand 5' end of an RNAi agent form a frayed end. In some embodiments, both ends of an RNAi agent form a frayed end. In some embodiments, neither end of an RNAi agent is a frayed end. As used herein a frayed end refers to an end of a double stranded RNAi agent in which the terminal nucleotides

10 of the two annealed strands from a pair (i.e. do not form an overhang) but are not complementary (i.e. form a non-complementary pair). As used herein, an overhang is a stretch of one or more unpaired nucleotides at the end of one strand of a double stranded RNAi agent. The unpaired nucleotides may be on the sense strand or the antisense strand, creating either 3' or 5' overhangs. In some embodiments, the RNAi agent contains: a blunt end and a frayed end,

15 a blunt end and 5' overhang end, a blunt end and a 3' overhang end, a frayed end and a 5' overhang end, a frayed end and a 3' overhang end, two 5' overhang ends, two 3' overhang ends, a 5' overhang end and a 3' overhang end, two frayed ends, or two blunt ends.

A nucleotide base (or nucleobase) is a heterocyclic pyrimidine or purine compound which is a

20 constituent of all nucleic acids and includes adenine (A), guanine (G), cytosine (C), thymine (T), and uracil (U). As used herein, the term "nucleotide" can include a modified nucleotide (such as, for example, a nucleotide mimic, abasic site (Ab), or a surrogate replacement moiety). Modified nucleotides, when used in various polynucleotide or oligonucleotide constructs, may preserve activity of the compound in cells while at the same time increasing the serum stability

25 of these compounds, and can also minimize the possibility of activating interferon activity in humans upon administering of the polynucleotide or oligonucleotide construct.

In some embodiments, an HBV RNAi agent is prepared or provided as a salt, mixed salt, or a free-acid. In some embodiments, an HBV RNAi agent is prepared as a sodium salt. Such forms

30 are within the scope of the inventions disclosed herein.

Modified Nucleotides

In some embodiments, an HBV RNAi agent contains one or more modified nucleotides. As used herein, a "modified nucleotide" is a nucleotide other than a ribonucleotide (2'-hydroxyl

nucleotide). In some embodiments, at least 50% (e.g., at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100%) of the nucleotides are modified nucleotides. As used herein, modified nucleotides include, but are not limited to, deoxyribonucleotides, nucleotide mimics, abasic nucleotides (represented herein as Ab), 2'-modified nucleotides, 3' to 3' linkages (inverted) nucleotides (represented herein as invdN, invN, invn, invAb), non-natural base-comprising nucleotides, bridged nucleotides, peptide nucleic acids (PNAs), 2',3'-seco nucleotide mimics (unlocked nucleobase analogues, represented herein as NUNA or NUNA), locked nucleotides (represented herein as NLNA or NLNA), 3'-O-methoxy (2' internucleoside linked) nucleotides (represented herein as 3'-OMen), 2'-F-Arabino nucleotides (represented herein as NfANA or NfANA), 5'-Me, 2'-fluoro nucleotide (represented herein as 5Me-Nf), morpholino nucleotides, vinyl phosphonate deoxyribonucleotides (represented herein as vpdN), vinyl phosphonate containing nucleotides, and cyclopropyl phosphonate containing nucleotides (cPrpN). 2'-modified nucleotides (i.e. a nucleotide with a group other than a hydroxyl group at the 2' position of the five-membered sugar ring) include, but are not limited to, 2'-O-methyl nucleotides (represented herein as a lower case letter 'n' in a nucleotide sequence), 2'-deoxy-2'-fluoro nucleotides (represented herein as Nf, also represented herein as 2'-fluoro nucleotide), 2'-deoxy nucleotides (represented herein as dN), 2'-methoxyethyl (2'-O-2-methoxyethyl) nucleotides (represented herein as NM or 2'-MOE), 2'-amino nucleotides, and 2'-alkyl nucleotides. It is not necessary for all positions in a given compound to be uniformly modified. Conversely, more than one modification may be incorporated in a single HBV RNAi agent or even in a single nucleotide thereof. The HBV RNAi agent sense strands and antisense strands may be synthesized and/or modified by methods known in the art. Modification at one nucleotide is independent of modification at another nucleotide.

Modified nucleobases include synthetic and natural nucleobases, such as 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, (e.g., 2-aminopropyladenine, 5-propynyluracil, or 5-propynylcytosine), 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-alkyl (e.g., 6-methyl, 6-ethyl, 6-isopropyl, or 6-n-butyl) derivatives of adenine and guanine, 2-alkyl (e.g., 2-methyl, 2-ethyl, 2-isopropyl, or 2-n-butyl) and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, cytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-sulfhydryl, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and

guanines, 5-halo (e.g., 5-bromo), 5-trifluoromethyl, and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, and 3-deazaadenine.

5 In some embodiments, all or substantially all of the nucleotides of an RNAi agent are modified nucleotides. As used herein, an RNAi agent wherein substantially all of the nucleotides present are modified nucleotides is an RNAi agent having four or fewer (i.e., 0, 1, 2, 3, or 4) nucleotides in both the sense strand and the antisense strand being ribonucleotides. As used herein, a sense strand wherein substantially all of the nucleotides present are modified nucleotides is a sense strand having two or fewer (i.e., 0, 1, or 2) nucleotides in the sense strand being ribonucleotides. As used herein, an antisense sense strand wherein substantially all of the nucleotides present are modified nucleotides is an antisense strand having two or fewer (i.e., 0, 1, or 2) nucleotides in the sense strand being ribonucleotides. In some embodiments, one or more nucleotides of an RNAi agent is a ribonucleotide.

15

Modified Internucleoside Linkages

In some embodiments, one or more nucleotides of an HBV RNAi agent are linked by non-standard linkages or backbones (i.e., modified internucleoside linkages or modified backbones). In some embodiments, a modified internucleoside linkage is a non-phosphate-containing covalent internucleoside linkage. Modified internucleoside linkages or backbones include, but are not limited to, 5'-phosphorothioate groups (represented herein as a lower case "s"), chiral phosphorothioates, thiophosphates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, alkyl phosphonates (e.g., methyl phosphonates or 3'-alkylene phosphonates), chiral phosphonates, phosphinates, phosphoramidates (e.g., 3'-amino phosphoramidate, aminoalkylphosphoramidates, or thionophosphoramidates), thionoalkyl-phosphonates, thionoalkylphosphotriesters, morpholino linkages, boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of boranophosphates, or boranophosphates having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. In some embodiments, a modified internucleoside linkage or backbone lacks a phosphorus atom. Modified internucleoside linkages lacking a phosphorus atom include, but are not limited to, short chain alkyl or cycloalkyl inter-sugar linkages, mixed heteroatom and alkyl or cycloalkyl inter-sugar linkages, or one or more short chain heteroatomic or heterocyclic inter-sugar linkages. In some embodiments, modified internucleoside backbones include, but are not limited to, siloxane backbones, sulfide backbones, sulfoxide backbones,

sulfone backbones, formacetyl and thioformacetyl backbones, methylene formacetyl and thioformacetyl backbones, alkene-containing backbones, sulfamate backbones, methyleneimino and methylenehydrazino backbones, sulfonate and sulfonamide backbones, amide backbones, and other backbones having mixed N, O, S, and CH₂ components.

5

In some embodiments, a sense strand of an HBV RNAi agent can contain 1, 2, 3, 4, 5, or 6 phosphorothioate linkages, an antisense strand of an HBV RNAi agent can contain 1, 2, 3, 4, 5, or 6 phosphorothioate linkages, or both the sense strand and the antisense strand independently can contain 1, 2, 3, 4, 5, or 6 phosphorothioate linkages. In some embodiments,
10 a sense strand of an HBV RNAi agent can contain 1, 2, 3, or 4 phosphorothioate linkages, an antisense strand of an HBV RNAi agent can contain 1, 2, 3, or 4 phosphorothioate linkages, or both the sense strand and the antisense strand independently can contain 1, 2, 3, or 4 phosphorothioate linkages.

15 In some embodiments, an HBV RNAi agent sense strand contains at least two phosphorothioate internucleoside linkages. In some embodiments, the at least two phosphorothioate internucleoside linkages are between the nucleotides at positions 1-3 from the 3' end of the sense strand. In some embodiments, the at least two phosphorothioate internucleoside linkages are between the nucleotides at positions 1-3, 2-4, 3-5, 4-6, 4-5, or 6-8 from the 5' end of the
20 sense strand. In some embodiments, an HBV RNAi agent antisense strand contains four phosphorothioate internucleoside linkages. In some embodiments, the four phosphorothioate internucleoside linkages are between the nucleotides at positions 1-3 from the 5' end of the sense strand and between the nucleotides at positions 19-21, 20-22, 21-23, 22-24, 23-25, or 24-26 from the 5' end. In some embodiments, an HBV RNAi agent contains at least two
25 phosphorothioate internucleoside linkages in the sense strand and three or four phosphorothioate internucleoside linkages in the antisense strand.

In some embodiments, an HBV RNAi agent contains one or more modified nucleotides and one or more modified internucleoside linkages. In some embodiments, a 2'-modified
30 nucleoside is combined with modified internucleoside linkage.

HBV RNAi Agents

In some embodiments, the HBV RNAi agents disclosed herein target an HBV gene at or near the positions of the HBV genome shown in the following Table 1. In some embodiments, the

antisense strand of an HBV RNAi agent disclosed herein includes a core stretch sequence that is fully, substantially, or at least partially complementary to a target HBV 19-mer sequence disclosed in Table 1.

- 5 **Table 1.** Example 19-mer HBV cDNA target sequences for HBV RNAi agents (taken from Hepatitis B virus (subtype ADW2), genotype A, complete genome GenBank AM282986.1 (SEQ ID NO:1)).

SEQ ID No.	HBV 19-mer Target Sequences (5' → 3')	Genome Position of SEQ ID NO: 1	Region of HBV Gene Targeted
2	GTGGTGGACTTCTCTCAAT	256-274	S ORF
3	TGGTGGACTTCTCTCAATT	257-275	S ORF
4	GGACTTCTCTCAATTTTCT	261-279	S ORF
5	GCTGTAGGCATAAATTGGT	1780-1798	X ORF
6	CTGTAGGCATAAATTGGTC	1781-1799	X ORF

- 10 In some embodiments, an HBV RNAi agent includes an antisense strand wherein position 19 of the antisense strand (5' → 3') is capable of forming a base pair with position 1 of a 19-mer target sequence disclosed in Table 1. In some embodiments, an HBV RNAi agent includes an antisense strand wherein position 1 of the antisense strand (5' → 3') is capable of forming a base pair with position 19 of the 19-mer target sequence disclosed in Table 1.

- 15 In some embodiments, an HBV RNAi agent includes an antisense strand wherein position 2 of the antisense strand (5' → 3') is capable of forming a base pair with position 18 of the 19-mer target sequence disclosed in Table 1. In some embodiments, an HBV RNAi agent includes an antisense strand wherein positions 2 through 18 of the antisense strand (5' → 3') are capable of forming base pairs with each of the respective complementary bases located at positions 18 through 2 of the 19-mer target sequence disclosed in Table 1.

In some embodiments, the HBV RNAi agents include core 19-mer nucleotide sequences shown in the following Table 2.

Table 2. HBV RNAi agent antisense strand and sense strand core stretch sequences (N=any nucleotide).

SEQ ID NO:	Antisense Sequence (5' → 3') (19-mer)	SEQ ID NO:	Sense Sequence (5' → 3') (19-mer)	Genome Position of SEQ ID NO: 1
7	AUUGAGAGAAAGUCCACCAC	34	GUGGUGGACUUCUCUCAAU	256-274
8	UUUGAGAGAAAGUCCACCAC	35	GUGGUGGACUUCUCUCAAA	256-274
9	AUUGAGAGAAAGUCCACCAN	36	NUGGUGGACUUCUCUCAAU	256-274
10	UUUGAGAGAAAGUCCACCAN	37	NUGGUGGACUUCUCUCAAA	256-274
11	NUUGAGAGAAAGUCCACCAN	38	NUGGUGGACUUCUCUCAAN	256-274
12	AAUUGAGAGAAAGUCCACCA	39	UGGUGGACUUCUCUCAAUU	257-275
13	UAUUGAGAGAAAGUCCACCA	40	UGGUGGACUUCUCUCAAU	257-275
14	AAUUGAGAGAAAGUCCACCN	41	NGGUGGACUUCUCUCAAUU	257-275
15	UAUUGAGAGAAAGUCCACCN	42	NGGUGGACUUCUCUCAAU	257-275
16	NAUUGAGAGAAAGUCCACCN	43	NGGUGGACUUCUCUCAUN	257-275
17	AGAAAUUUGAGAGAAAGUCC	44	GGACUUCUCUCAAUUUUCU	261-279
18	UGAAAUUUGAGAGAAAGUCC	45	GGACUUCUCUCAAUUUUCA	261-279
19	AGAAAUUUGAGAGAAAGUCN	46	NGACUUCUCUCAAUUUUCU	261-279
20	UGAAAUUUGAGAGAAAGUCN	47	NGACUUCUCUCAAUUUUCA	261-279
21	NGAAAUUUGAGAGAAAGUCN	48	NGACUUCUCUCAAUUUUCN	261-279
22	ACCAAUUUAUGCCUACAGC	49	GCUGUAGGCAUAAAUUGGU	1780-1798
23	UCCAAUUUAUGCCUACAGC	50	GCUGUAGGCAUAAAUUGGA	1780-1798
24	ACCAAUUUAUGCCUACAGN	51	NCUGUAGGCAUAAAUUGGU	1780-1798
25	UCCAAUUUAUGCCUACAGN	52	NCUGUAGGCAUAAAUUGGA	1780-1798
26	NCCAAUUUAUGCCUACAGN	53	NCUGUAGGCAUAAAUUGGN	1780-1798
27	GACCAAUUUUAUGCCUACAG	54	CUGUAGGCAUAAAUUGGUC	1781-1799
28	AACCAAUUUUAUGCCUACAG	55	CUGUAGGCAUAAAUUGGUU	1781-1799
29	UACCAAUUUUAUGCCUACAG	56	CUGUAGGCAUAAAUUGGUA	1781-1799

SEQ ID NO:	Antisense Sequence (5' → 3') (19-mer)	SEQ ID NO:	Sense Sequence (5' → 3') (19-mer)	Genome Position of SEQ ID NO: 1
30	GACCAAUUUAUGCCUACAN	57	NUGUAGGCAUAAAUUGGUC	1781-1799
31	AACCAAUUUAUGCCUACAN	58	NUGUAGGCAUAAAUUGGUU	1781-1799
32	UACCAAUUUAUGCCUACAN	59	NUGUAGGCAUAAAUUGGUA	1781-1799
33	NACCAAUUUAUGCCUACAN	60	NUGUAGGCAUAAAUUGGUN	1781-1799

The HBV RNAi agent sense strands and antisense strands that comprise or consist of the nucleotide sequences in Table 2 can be modified nucleotides or unmodified nucleotides. In some embodiments, the HBV RNAi agents having the sense and antisense strand sequences that comprise or consist of the nucleotide sequences in Table 2 are all or substantially all modified nucleotides.

In some embodiments, the antisense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the antisense strand sequences in Table 2. In some embodiments, the sense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the sense strand sequences in Table 2.

Modified HBV RNAi agent antisense strand sequences, as well as their underlying unmodified sequences, are provided in Table 3. Modified HBV RNAi agent sense strands, as well as their underlying unmodified sequences, are provided in Table 4. In forming HBV RNAi agents, each of the nucleotides in each of the unmodified sequences listed in Tables 3 and 4 may be a modified nucleotide.

As used herein (including in Tables 3 and 4), the following notations are used to indicate modified nucleotides, targeting groups, and linking groups. As the person of ordinary skill in the art would readily understand, unless otherwise indicated by the sequence, that when present in an oligonucleotide, the monomers are mutually linked by 5'-3'-phosphodiester bonds:

	A	=	adenosine-3'-phosphate;
	C	=	cytidine-3'-phosphate;
	G	=	guanosine-3'-phosphate;
25	U	=	uridine-3'-phosphate
	n	=	any 2'-OMe modified nucleotide
	a	=	2'-O-methyladenosine-3'-phosphate
	as	=	2'-O-methyladenosine-3'-phosphorothioate
	c	=	2'-O-methylcytidine-3'-phosphate
30	cs	=	2'-O-methylcytidine-3'-phosphorothioate
	g	=	2'-O-methylguanosine-3'-phosphate
	gs	=	2'-O-methylguanosine-3'-phosphorothioate
	t	=	2'-O-methyl-5-methyluridine-3'-phosphate
	ts	=	2'-O-methyl-5-methyluridine-3'-phosphorothioate
35	u	=	2'-O-methyluridine-3'-phosphate
	us	=	2'-O-methyluridine-3'-phosphorothioate

	Nf	=	any 2'-fluoro modified nucleotide
	Af	=	2'-fluoroadenosine-3'-phosphate
	Afs	=	2'-fluoroadenosine-3'-phosphorothioate
	Cf	=	2'-fluorocytidine-3'-phosphate
5	Cfs	=	2'-fluorocytidine-3'-phosphorothioate
	Gf	=	2'-fluoroguanosine-3'-phosphate
	Gfs	=	2'-fluoroguanosine-3'-phosphorothioate
	Tf	=	2'-fluoro-5'-methyluridine-3'-phosphate
	Tfs	=	2'-fluoro-5'-methyluridine-3'-phosphorothioate
10	Uf	=	2'-fluorouridine-3'-phosphate
	Ufs	=	2'-fluorouridine-3'-phosphorothioate
	dN	=	any 2'-deoxyribonucleotide
	dT	=	2'-deoxythymidine-3'-phosphate
	N _{UNA}	=	2',3'-seco nucleotide mimics (unlocked nucleobase analogs)
15	N _{LNA}	=	locked nucleotide
	Nf _{ANA}	=	2'-F-Arabino nucleotide
	NM	=	2'-methoxyethyl nucleotide
	AM	=	2'-methoxyethyladenosine-3'-phosphate
	AMs	=	2'-methoxyethyladenosine-3'-phosphorothioate
20	TM	=	2'-methoxyethylthymidine-3'-phosphate
	TMs	=	2'-methoxyethylthymidine-3'-phosphorothioate
	R	=	ribitol
	(invdN)	=	any inverted deoxyribonucleotide (3'-3' linked nucleotide)
	(invAb)	=	inverted (3'-3' linked) abasic deoxyribonucleotide, see Table 6
25	(invAb)s	=	inverted (3'-3' linked) abasic deoxyribonucleotide-5'-phosphorothioate, see Table 6
	(invn)	=	any inverted 2'-OMe nucleotide (3'-3' linked nucleotide)
	s	=	phosphorothioate linkage
	vpdN	=	vinyl phosphonate deoxyribonucleotide
30	(5Me-Nf)	=	5'-Me, 2'-fluoro nucleotide
	cPrp	=	cyclopropyl phosphonate, see Table 6
	epTcPr	=	see Table 6
	epTM	=	see Table 6

35 The person of ordinary skill in the art would readily understand that the terminal nucleotide at the 3' end of a given oligonucleotide sequence would typically have a hydroxyl (-OH) group at the respective 3' position of the given monomer instead of a phosphate moiety *ex vivo*. Thus, for example, as shown above in the structure representation of AD05070, above, the "g" modified nucleotide on the terminal 3' end of the antisense strand of AM06606-AS has a

40 hydroxyl group positioned at its 3' position. Unless expressly indicated otherwise herein, such

understandings of the person of ordinary skill in the art are used when describing the HBV RNAi agents and compositions of HBV RNAi agents disclosed herein.

Targeting groups and linking groups include the following, for which their chemical structures
5 are provided below in Table 6: (PAZ), (NAG13), (NAG13)s, (NAG18), (NAG18)s, (NAG24),
(NAG24)s, (NAG25), (NAG25)s, (NAG26), (NAG26)s, (NAG27), (NAG27)s, (NAG28),
(NAG28)s, (NAG29), (NAG29)s, (NAG30), (NAG30)s, (NAG31), (NAG31)s, (NAG32),
(NAG32)s, (NAG33), (NAG33)s, (NAG34), (NAG34)s, (NAG35), (NAG35)s, (NAG36),
(NAG36)s, (NAG37), (NAG37)s, (NAG38), (NAG38)s, (NAG39), (NAG39)s. Each sense
10 strand and/or antisense strand can have any targeting groups or linking groups listed above, as
well as other targeting or linking groups, conjugated to the 5' and/or 3' end of the sequence.

Table 3. HBV RNAi Agent antisense strand sequences.

AS Strand ID	Modified sequence (5' → 3')	SEQ ID NO.	Unmodified sequence (5' → 3')	SEQ ID NO.
AM03508-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgGfccusuAu	61	UACCAAUUUAUGCCUACAGGCCUUAU	149
AM04441-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgGfccsu	62	UACCAAUUUAUGCCUACAGGCCU	150
AM04442-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgGfccsu	63	UACCAAUUUAUGCCUACAGGCCU	150
AM04443-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgGfsc	64	UACCAAUUUAUGCCUACAGGC	151
AM04661-AS	usGfsugaAfgCfGfaaguGfcAfcacsusu	65	UGUGAAGCGAAGUGCACACUU	152
AM04768-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgCfcsuscegc	66	UACCAAUUUAUGCCUACAGCCUCCGC	153
AM04769-AS	vpusApscCfaAfuUfuAfuGfcCfuAfcAfgCfcsuscgc	67	UACCAAUUUAUGCCUACAGCCUCCGC	153
AM05011-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgusu	68	UACCAAUUUAUGCCUACAGUU	154
AM05012-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfggsc	69	UACCAAUUUAUGCCUACAGGC	151
AM05013-AS	vpusApscCfaAfuUfuAfuGfcCfuAfcAfgGfsc	70	UACCAAUUUAUGCCUACAGGC	151
AM05014-AS	vpusApscCfaAfuUfuAfuGfcCfuAfcAfgusu	71	UACCAAUUUAUGCCUACAGUU	154
AM05052-AS	asUfsusGfaGfaGfaAfgUfcCfaCfcAfcGfsa	72	AUUGAGAGAAGUCCACCACGA	155
AM05053-AS	asUfsusGfaGfaGfaAfgUfcCfaCfcAfcgsa	73	AUUGAGAGAAGUCCACCACGA	155
AM05054-AS	asUfsusGfaGfaGfaAfgUfcCfaCfcAfcusu	74	AUUGAGAGAAGUCCACCACUU	156
AM05055-AS	vpusUfsusGfaGfaGfaAfgUfcCfaCfcAfcGfsa	75	UUUGAGAGAAGUCCACCACGA	157
AM05056-AS	asAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg	76	AAUUGAGAGAAGUCCACCACG	158
AM05057-AS	asAfsusUfgAfgAfgAfaGfuCfcAfcCfacsg	77	AAUUGAGAGAAGUCCACCACG	158
AM05058-AS	asAfsusUfgAfgAfgAfaGfuCfcAfcCfausu	78	AAUUGAGAGAAGUCCACCACUU	159
AM05060-AS	vpusAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg	79	UAUUGAGAGAAGUCCACCACG	160
AM05351-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgGfsu	80	UACCAAUUUAUGCCUACAGGU	161
AM05608-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgsusu	81	UACCAAUUUAUGCCUACAGUU	154
AM05609-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgcsc	82	UACCAAUUUAUGCCUACAGCC	162
AM05610-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgccusu	83	UACCAAUUUAUGCCUACAGCCUU	163
AM05611-AS	usApscCfaAfuUfuAfuGfcCfuAfcAfgccusc	84	UACCAAUUUAUGCCUACAGCCUC	164

AM05612-AS	usAfscscaauUfuAfuGfcCfuacagcsc	85	UACCAAUUUAUGCCUACAGCC	162
AM05613-AS	usAfscscaauUfuAfuGfcCfuacagccusu	86	UACCAAUUUAUGCCUACAGCCUU	163
AM05614-AS	usAfscscaauUfuAfuGfcCfuacagccusc	87	UACCAAUUUAUGCCUACAGCCUC	164
AM05618-AS	asUfsusgagaGfaAfgUfcCfaccacusu	88	AUUGAGAGAAGUCCACCACUU	156
AM05621-AS	usUfsusGfaGfaGfaAfgUfcCfaCfcAfcusu	89	UUUGAGAGAAGUCCACCACUU	165
AM05623-AS	asUfsusGfaGfaGfaAfgUfcCfaCfcAfcggusu	90	AUUAGAGAGAAGUCCACCACGGUU	166
AM05626-AS	asUfsusgagaGfaAfgUfcCfaccacggusu	91	AUUGAGAGAAGUCCACCACGGUU	166
AM05628-AS	asUfsusGfaGfaGfaAfgUfcCfaCfcAfcgagsu	92	AUUGAGAGAAGUCCACCACGAGU	167
AM05631-AS	usAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg	93	UAUUGAGAGAAGUCCACCACG	160
AM05632-AS	usAfsusugagAfgAfaGfuCfcaccacsg	94	UAUUGAGAGAAGUCCACCACG	160
AM05633-AS	usAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfgsu	95	UAUUGAGAGAAGUCCACCACGUU	168
AM05634-AS	usAfsusugagAfgAfaGfuCfcaccacgag	96	UAUUGAGAGAAGUCCACCACGAG	169
AM05635-AS	usAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg	97	UAUUGAGAGAAGUCCACCACGAG	169
AM05637-AS	usAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfgsa	98	UAUUGAGAGAAGUCCACCACGA	170
AM05638-AS	usAfsusugagAfgAfaGfuCfcaccacgsa	99	UAUUGAGAGAAGUCCACCACGA	170
AM05747-AS	asGfsasAfaAfuugagAfgAfaGfuCfcAfsc	100	AGAAAUUGAGAGAAGUCCAC	171
AM05849-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgusu	101	UACCAAUUUAUGCCUACAGUU	154
AM05850-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgcsc	102	UACCAAUUUAUGCCUACAGCC	162
AM05851-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgcusu	103	UACCAAUUUAUGCCUACAGCUU	172
AM05852-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgccsu	104	UACCAAUUUAUGCCUACAGCCU	173
AM05853-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgccusu	105	UACCAAUUUAUGCCUACAGCCUU	163
AM05854-AS	usAfscsCfaAfuuuauGfcCfuAfcAfgccusc	106	UACCAAUUUAUGCCUACAGCCUC	164
AM05855-AS	cPrpusAfscsCfaAfuUfuAfuGfcCfuAfcAfgusu	107	UACCAAUUUAUGCCUACAGUU	154
AM05860-AS	cPrpusAfsusUfgAfgAfgAfaGfuCfcAfcCfaCfsg	108	UAUUGAGAGAAGUCCACCACG	160
AM05862-AS	usAfsusUfgAfgagaaGfuCfcAfcCfausu	109	UAUUGAGAGAAGUCCACCACUU	174
AM05863-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacsg	110	UAUUGAGAGAAGUCCACCACG	160
AM05864-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacusu	111	UAUUGAGAGAAGUCCACCACUU	175
AM05865-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacsgsa	112	UAUUGAGAGAAGUCCACCACGA	170

AM05867-AS	vpusAfsusUfgAfgagaaGfuCfcAfcCfaCfsg	113	UAUUGAGAGAAGUCCACCACG	160
AM05873-AS	usUfsusGfaGfagaagUfcCfaCfcAfcusu	114	UUUGAGAGAAGUCCACCACUU	165
AM05874-AS	usUfsusGfaGfagaagUfcCfaCfcAfcgsa	115	UUUGAGAGAAGUCCACCACGA	157
AM05875-AS	usUfsusGfaGfagaagUfcCfaCfcAfcgusu	116	UUUGAGAGAAGUCCACCACGUU	176
AM05876-AS	usUfsusGfaGfagaagUfcCfaCfcAfcgasg	117	UUUGAGAGAAGUCCACCACGAG	177
AM05877-AS	cPrpusUfsusGfaGfaGfaAfgUfcCfaCfcAfcusu	118	UUUGAGAGAAGUCCACCACUU	165
AM06074-AS	cPrpusAfsusUfgAfgagaaGfuCfcAfcCfacusu	119	UAUUGAGAGAAGUCCACCACUU	175
AM06142-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacusu	120	UAUUGAGAGAAGUCCACCACUU	175
AM06143-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacgusu	121	UAUUGAGAGAAGUCCACCACGUU	168
AM06144-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacuus(invAb)	122	UAUUGAGAGAAGUCCACCACUU	175
AM06145-AS	usAfsusUfgAfgagaaGfuCfcAfcCfacgasg	123	UAUUGAGAGAAGUCCACCACGAG	169
AM06222-AS	usAfsusUfgAfgAfaGfuCfcAfcCfacusu	124	UAUUGAGAGAAGUCCACCACUU	175
AM06281-AS	asGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu	125	AGAAAUUUGAGAGAAGUCCUU	178
AM06282-AS	asGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc	126	AGAAAUUUGAGAGAAGUCCAC	171
AM06283-AS	asGfsasAfaAfuUfgAfgAfgAfaGfuCfcacusu	127	AGAAAUUUGAGAGAAGUCCACUU	179
AM06284-AS	asGfsasAfaAfuUfgAfgAfgAfaGfuCfcacsc	128	AGAAAUUUGAGAGAAGUCCACC	180
AM06285-AS	usGfsasAfaAfuUfgAfgAfgAfaGfuCfcusu	129	UGAAAUUUGAGAGAAGUCCUU	152
AM06286-AS	usGfsasAfaAfuUfgAfgAfgAfaGfuCfcasc	130	UGAAAUUUGAGAGAAGUCCAC	181
AM06299-AS	asCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu	131	ACCAAUUUAUGCCUACAGCUU	182
AM06300-AS	asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu	132	ACCAAUUUAUGCCUACAGCCUU	183
AM06301-AS	asCfscsAfaUfuUfaUfgCfcUfaCfaGfccusc	133	ACCAAUUUAUGCCUACAGCCUC	184
AM06302-AS	usCfscsAfaUfuUfaUfgCfcUfaCfaGfcusu	134	UCCAAUUUAUGCCUACAGCUU	185
AM06303-AS	usCfscsAfaUfuUfaUfgCfcUfaCfaGfccusu	135	UCCAAUUUAUGCCUACAGCCUU	186
AM06463-AS	cPrpusAfsesCfaAfuUfuAfuGfcCfuAfcAfgscsc	136	UACCAAUUUAUGCCUACAGCC	162
AM06464-AS	usAfsesCfaAfuUfuAfuGfcCfuAfcAfgscsc	137	UACCAAUUUAUGCCUACAGCC	162
AM06465-AS	cPrpusAfsesCfaAfuUfuAfuGfcCfuAfcAfgscsc	138	UACCAAUUUAUGCCUACAGCC	162
AM06604-AS	usAfsesCfaAfuUfuAfuGfcCfuAfcAfgesu	139	UACCAAUUUAUGCCUACAGCU	187
AM06606-AS	usAfsesCfaAfuUfuAfuGfcCfuAfcAfgcsg	140	UACCAAUUUAUGCCUACAGCG	188

AM06608-AS	asAfscsCfaAfuUfuAfuGfcCfuAfcAfgcsc	141	AACCAAUUUUGCCUACAGCC	189
AM06611-AS	usAfscsCfaAfuUfuAfuGfcCfuAfcAfgusu	142	UACCAAUUUUGCCUACAGUU	154
AM06612-AS	usAfscsCfaAfuUfuAfuGfcCfuAfcAfgCfsc	143	UACCAAUUUUGCCUACAGCC	162
AM06614-AS	asCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu	144	ACCAAUUUUGCCUACAGCCU	190
AM06616-AS	usCfscAfaUfuUfaUfgCfcUfaCfaGfcCfsu	145	UCCAAUUUUGCCUACAGCCU	191
AM06618-AS	asCfscAfaUfuUfaUfgCfcUfaCfaGfccsg	146	ACCAAUUUUGCCUACAGCCG	192
AM06620-AS	usCfscAfaUfuUfaUfgCfcUfaCfaGfccsg	147	UCCAAUUUUGCCUACAGCCG	193
AM06751-AS	usAfscsCfaAfuUfuAfuGfcCfuAfcAfgggsg	148	UACCAAUUUUGCCUACAGGG	194

Table 4. HBV RNAi agent sense strand sequences.

Strand ID	Modified sequence (5' → 3')	SEQ ID NO.	Unmodified sequence (5' → 3')	SEQ ID NO.
AM04444-SS	(NAG25)uusgscucguagGfCfAfuaaaauugguaus(inv dT)	195	UUGCCUGUAGGCAUAAAUUGGUAUT	275
AM04445-SS	(NAG25)uaauusgscucguagGfCfAfuaaaauuggu(inv dA)	196	UAUAUGCCUGUAGGCAUAAAUUGGUA	276
AM04767-SS	(NAG25)gcfgagsgcucguagGfCfAfuaaaauuggTM(inv dA)	197	GCGGAGGCUGUAGGCAUAAAUUGGTA	277
AM05010-SS	(NAG25)scsuguagGfCfAfuaaaauugguaus(inv Ab)	198	CUGUAGGCAUAAAUUGGUUU	278
AM05015-SS	(NAG25)sgscucguagGfCfAfuaaaauugguas(inv Ab)	199	GCCUGUAGGCAUAAAUUGGUA	279
AM05016-SS	(NAG25)sgscucguagGfCfAfuaaaauuggus(inv dA)	200	GCCUGUAGGCAUAAAUUGGUA	279
AM05017-SS	(NAG25)sgscucguagGfCfAfuaaaauugguAMs(inv Ab)	201	GCCUGUAGGCAUAAAUUGGUA	279
AM05018-SS	(NAG25)sgscucguagGfCfAfuaaaauuggTMAMs(inv Ab)	202	GCCUGUAGGCAUAAAUUGGTA	280
AM05019-SS	(NAG25)sasacuguagGfCfAfuaaaauugguas(inv Ab)	203	AACUGUAGGCAUAAAUUGGUA	281
AM05034-SS	(NAG25)suscguggugGfAfCfuucucuaas(inv Ab)	204	UCGUGGUGGACUUCUCUCAAU	282
AM05046-SS	(NAG25)sasaguggugGfAfCfuucucuaas(inv Ab)	205	AAGUGGUGGACUUCUCUCAAU	283
AM05047-SS	(NAG25)suscguggugGfAfCfuucucucaAMTMs(inv Ab)	206	UCGUGGUGGACUUCUCUCAAT	284
AM05048-SS	(NAG25)scsgugguggAfCfUfucucuaaas(inv Ab)	207	CGUGGUGGACUUCUCUCAAUU	285
AM05049-SS	(NAG25)sasaugguggAfCfUfucucuaaas(inv Ab)	208	AAUGGUGGACUUCUCUCAAUU	286
AM05050-SS	(NAG25)scsgugguggAfCfUfucucucaAMTMs(inv Ab)	209	CGUGGUGGACUUCUCUCAATT	287
AM05051-SS	(NAG25)sgsgacuucuCfUfCfaauuuucaaas(inv Ab)	210	GGACUUCUCUCAAUUUUCUAA	288
AM05063-SS	(NAG25)scsgugguggAfCfUfucucuaaas(inv Ab)	211	CGUGGUGGACUUCUCUCAAUU	289
AM05064-SS	(NAG25)suscguggugGfAfCfuucucuaaas(inv Ab)	212	UCGUGGUGGACUUCUCUCAAA	290
AM05346-SS	(NAG31)sasccuguagGfCfAfuaaaauugguas(inv Ab)	213	ACCUGUAGGCAUAAAUUGGUA	291
AM05347-SS	(NAG31)s(inv Ab)scuguagGfCfAfuaaaauugguas(inv Ab)	214	CUGUAGGCAUAAAUUGGUA	292
AM05606-SS	(NAG25)s(inv Ab)scuguagGfCfAfuaaaauugguas(inv Ab)	215	CUGUAGGCAUAAAUUGGUA	292
AM05607-SS	(NAG37)s(inv Ab)scuguagGfCfAfuaaaauugguas(inv Ab)	216	CUGUAGGCAUAAAUUGGUA	292
AM05615-SS	(NAG25)s(inv Ab)sacuguagGfCfAfuaaaauugguas(inv Ab)	217	ACUGUAGGCAUAAAUUGGUA	293

Strand ID	Modified sequence (5' → 3')	SEQ ID NO.	Unmodified sequence (5' → 3')	SEQ ID NO.
AM05616-SS	(NAG25)sgsgcuguagGfCfAfuaauugguas(invAb)	218	GGCUGUAGGCAUAAAUUGGUA	294
AM05617-SS	(NAG37)sasaguggugGfAfCfuucucucaaus(invAb)	219	AAGUGGUGGACUUCUCUCAAU	283
AM05620-SS	(NAG25)sasaguggugGfAfCfuucucucaaaas(invAb)	220	AAGUGGUGGACUUCUCUCAAA	295
AM05622-SS	(NAG25)scscguggugGfAfCfuucucucaaus(invAb)	221	CCGUGGUGGACUUCUCUCAAU	296
AM05624-SS	(NAG25)s(invAb)sccguggugGfAfCfuucucucaaus(invAb)	222	CCGUGGUGGACUUCUCUCAAU	296
AM05627-SS	(NAG25)scsucguggugGfAfCfuucucucaaus(invAb)	223	CUCGUGGUGGACUUCUCUCAAU	297
AM05629-SS	(NAG25)s(invAb)sguggugGfAfCfuucucucaaus(invAb)	224	GUGGUGGACUUCUCUCAAU	298
AM05630-SS	(NAG25)s(invAb)sguggugGfAfCfuucucucaauusu(invAb)	225	GUGGUGGACUUCUCUCAAUU	299
AM05636-SS	(NAG25)suscguggugAfCfUfuucucucaaus(invAb)	226	UCGUGGUGGACUUCUCUCAAUU	300
AM05639-SS	(NAG25)s(invAb)suggugGfCfUfuucucucaaus(invAb)	227	UGGUGGACUUCUCUCAAUU	301
AM05640-SS	(NAG37)s(invAb)suggugAfCfUfuucucucaaus(invAb)	228	UGGUGGACUUCUCUCAAUU	301
AM05746-SS	(NAG25)sgsuggacuuCfUfCfucaauuuucus(invAb)	229	GUGGACUUCUCUCAAUUUUCU	302
AM05856-SS	(NAG25)s(invAb)scuguagGfCfAfuaauugguausu(invAb)	230	CUGUAGGCAUAAAUUGGUUUU	278
AM05857-SS	(NAG25)s(invAb)sgcuguagGfCfAfuaauugguausu(invAb)	231	GCUGUAGGCAUAAAUUGGUUUU	303
AM05858-SS	(NAG25)s(invAb)sggcuguagGfCfAfuaauugguausu(invAb)	232	GGCUGUAGGCAUAAAUUGGUUUU	304
AM05859-SS	(NAG25)s(invAb)saacuguagGfCfAfuaauugguausu(invAb)	233	AACUGUAGGCAUAAAUUGGUUUU	305
AM05868-SS	(NAG25)s(invAb)uggugGfCfUfuucucucaauusu(invAb)	234	UGGUGGACUUCUCUCAAUUUU	306
AM05869-SS	(NAG25)s(invAb)sguggugAfCfUfuucucucaauusu(invAb)	235	GUGGUGGACUUCUCUCAAUUUU	307
AM05870-SS	(NAG25)sasaguggugAfCfUfuucucucaauusu(invAb)	236	AAUGGUGGACUUCUCUCAAUUUU	308
AM05871-SS	(NAG25)scsguggugGfCfUfuucucucaauusu(invAb)	237	CGUGGUGGACUUCUCUCAAUUUU	309
AM05872-SS	(NAG31)scsguggugGfCfUfuucucucaauas(invAb)	238	CGUGGUGGACUUCUCUCAAUUUU	289
AM05879-SS	(NAG25)s(invAb)saaguggugGfAfCfuucucucaaus(invAb)	239	AAGUGGUGGACUUCUCUCAAUUUU	283
AM05880-SS	(NAG25)s(invAb)sguggugGfAfCfuucucucaaausu(invAb)	240	GUGGUGGACUUCUCUCAAAUUU	310
AM05881-SS	(NAG25)s(invAb)scguggugGfAfCfuucucucaaausu(invAb)	241	CGUGGUGGACUUCUCUCAAAUUU	311

Strand ID	Modified sequence (5' → 3')	SEQ ID NO.	Unmodified sequence (5' → 3')	SEQ ID NO.
AM05882-SS	(NAG25)sasaguggGfAfCfuucucuaaasu(invAb)	242	AAGUGGUGGACUUCUCUCAAUU	312
AM05883-SS	(NAG25)suscguggGfAfCfuucucuaaasu(invAb)	243	UCGUGGUGGACUUCUCUCAAUU	313
AM06146-SS	(NAG37)s(invAb)sgugggAfCfUfucucuaaasu(invAb)	244	GUGGUGGACUUCUCUCAAUU	307
AM06147-SS	(NAG37)s(invAb)scgugggAfCfUfucucuaaasu(invAb)	245	CGUGGUGGACUUCUCUCAAUU	309
AM06148-SS	(NAG37)s(invAb)scucguggAfCfUfucucuaaasu(invAb)	246	CUCGUGGUGGACUUCUCUCAAUA	314
AM06149-SS	(NAG37)s(invAb)scucguggAfCfUfucucuaaasu(invAb)	247	CUCGUGGUGGACUUCUCUCAAUU	315
AM06150-SS	(NAG37)s(invAb)sggcugGfCfAfuaaauugguas(invAb)	248	GGCUGUAGGCAUAAAUUGGUA	294
AM06151-SS	(NAG37)s(invAb)sgaggcugagGfCfAfuaaauugguas(invAb)	249	GAGGCUGUAGGCAUAAAUUGGUA	316
AM06152-SS	(NAG37)s(invAb)sgaggcugagGfCfAfuaaauugguas(invAb)	250	GAGGCUGUAGGCAUAAAUUGGUAUU	317
AM06287-SS	(NAG37)s(invAb)sggacuuCfUfCfuaauuuucus(invAb)	251	GGACUUCUCUCAAUUUUCU	318
AM06288-SS	(NAG37)s(invAb)sgggacuuCfUfCfuaauuuucus(invAb)	252	GUGGACUUCUCUCAAUUUUCU	302
AM06289-SS	(NAG37)s(invAb)sggugacuuCfUfCfuaauuuucus(invAb)	253	GGUGGACUUCUCUCAAUUUUCU	319
AM06290-SS	(NAG37)s(invAb)sggacuuCfUfCfuaauuuucas(invAb)	254	GGACUUCUCUCAAUUUUCA	320
AM06291-SS	(NAG37)s(invAb)sgggacuuCfUfCfuaauuuucas(invAb)	255	GUGGACUUCUCUCAAUUUUCA	321
AM06304-SS	(NAG37)s(invAb)sgcuguaGfGfCfauaauuggus(invAb)	256	GCUGUAGGCAUAAAUUGGU	322
AM06305-SS	(NAG37)s(invAb)sggcuguaGfGfCfauaauuggus(invAb)	257	GGCUGUAGGCAUAAAUUGGU	323
AM06306-SS	(NAG37)s(invAb)sgaggcuguaGfGfCfauaauuggus(invAb)	258	GAGGCUGUAGGCAUAAAUUGGU	324
AM06307-SS	(NAG37)s(invAb)sgcuguaGfGfCfauaauuggas(invAb)	259	GCUGUAGGCAUAAAUUGGA	325
AM06308-SS	(NAG37)s(invAb)sggcuguaGfGfCfauaauuggas(invAb)	260	GGCUGUAGGCAUAAAUUGGA	326
AM06603-SS	(NAG37)s(invAb)sagcuguaGfCfAfuaaauugguas(invAb)	261	AGCUGUAGGCAUAAAUUGGUA	327
AM06605-SS	(NAG37)s(invAb)scgcuguaGfCfAfuaaauugguas(invAb)	262	CGCUGUAGGCAUAAAUUGGUA	328
AM06607-SS	(NAG37)s(invAb)sggcuguaGfCfAfuaaauugguas(invAb)	263	GGCUGUAGGCAUAAAUUGGUU	329
AM06609-SS	(NAG37)s(invAb)scuguaGfCfAfuaaauugguas(invAb)	264	CUGUAGGCAUAAAUUGGUAUU	278
AM06610-SS	(NAG37)s(invAb)scuGfuAfgGfCfAfuaUfgGfuasuus(invAb)	265	CUGUAGGCAUAAAUUGGUAUU	278

Strand ID	Modified sequence (5' → 3')	SEQ ID NO.	Unmodified sequence (5' → 3')	SEQ ID NO.
AM06613-SS	(NAG37)s(invAb)saggcuguaGfGcfauaaauggus(invAb)	266	AGGCUGUAGGCAUAAAUUGGU	330
AM06615-SS	(NAG37)s(invAb)saggcuguaGfGcfauaaauggas(invAb)	267	AGGCUGUAGGCAUAAAUUGGA	331
AM06617-SS	(NAG37)s(invAb)scggcuguaGfGcfauaaauggus(invAb)	268	CGGCUGUAGGCAUAAAUUGGU	332
AM06619-SS	(NAG37)s(invAb)scggcuguaGfGcfauaaauggas(invAb)	269	CGGCUGUAGGCAUAAAUUGGA	333
AM06750-SS	(NAG37)s(invAb)scgcuguagGfCfAfuaaaugguas(invAb)	270	CCCUGUAGGCAUAAAUUGGUA	334
AM06752-SS	(NAG37)csgcuguagGfCfAfuaaaugguas(invAb)	271	CGCUGUAGGCAUAAAUUGGUA	328
AM06753-SS	(NAG37)cscgcuguagGfCfAfuaaaugguas(invAb)	272	CCCUGUAGGCAUAAAUUGGUA	334
AM06776-SS	(NAG25)s(invAb)sgggacuuCfUcfucaauuuucus(invAb)	273	GUGGACUUCUCUCACAUUUUUCU	302
AM06777-SS	(NAG25)s(invAb)scgcuguagGfCfAfuaaaugguas(invAb)	274	CGCUGUAGGCAUAAAUUGGUA	328

The HBV RNAi agents described herein are formed by annealing an antisense strand with a sense strand. A sense strand containing a sequence listed in Table 4 can be hybridized to any antisense strand containing a sequence listed in Table 3, provided the two sequences have a region of at least about 85% complementarity over a contiguous 16, 17, 18, 19, 20, or 21 nucleotide sequence.

In some embodiments, the antisense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the antisense strand sequences in Table 3. In some embodiments, the sense strand of an HBV RNAi agent disclosed herein differs by 0, 1, 2, or 3 nucleotides from any of the sense strand sequences in Table 4.

In some embodiments, an HBV RNAi agent antisense strand comprises a nucleotide sequence of any of the sequences in Table 3. In some embodiments, an HBV RNAi agent antisense strand comprises the sequence of nucleotides (from 5' end → 3' end) 1-17, 2-17, 1-18, 2-18, 1-19, 2-19, 1-20, 2-20, 1-21, 2-21, 1-22, 2-22, 1-23, 2-23, 1-24, 2-24, 1-25, 2-25, 1-26, or 2-26 of any of the sequences in Table 3.

In some embodiments, an HBV RNAi agent sense strand comprises the nucleotide sequence of any of the sequences in Table 4. In some embodiments, an HBV RNAi agent sense strand comprises the sequence of nucleotides (from 5' end → 3' end) 1-17, 2-17, 3-17, 4-17, 1-18, 2-18, 3-18, 4-18, 1-19, 2-19, 3-19, 4-19, 1-20, 2-20, 3-20, 4-20, 1-21, 2-21, 3-21, 4-21, 1-22, 2-22, 3-22, 4-22, 1-23, 2-23, 3-23, 4-23, 1-24, 2-24, 3-24, 4-24, 1-25, 2-25, 3-25, 4-25, 1-26, 2-26, 3-26, or 4-26 of any of the sequences in Table 4.

For the HBV RNAi agents disclosed herein, the nucleotide at position 1 of the antisense strand (from 5' end → 3' end) can be perfectly complementary to an HBV gene, or can be non-complementary to an HBV gene. In some embodiments, the nucleotide at position 1 of the antisense strand (from 5' end → 3' end) is a U, A, or dT. In some embodiments, the nucleotide at position 1 of the antisense strand (from 5' end → 3' end) forms an A:U or U:A base pair with the sense strand.

In some embodiments, an HBV RNAi agent antisense strand comprises the sequence of nucleotides (from 5' end → 3' end) 2-18 or 2-19 of any of the antisense strand sequences in

Table 3. In some embodiments, an HBV RNAi sense strand comprises the sequence of nucleotides (from 5' end → 3' end) 1-17 or 1-18 of any of the sense strand sequences in Table 4.

- 5 In some embodiments, an HBV RNAi agent includes (i) an antisense strand comprising the sequence of nucleotides (from 5' end → 3' end) 2-18 or 2-19 of any of the antisense strand sequences in Table 3, and (ii) a sense strand comprising the sequence of nucleotides (from 5' end → 3' end) 1-17 or 1-18 of any of the sense strand sequences in Table 4.
- 10 A sense strand containing a sequence listed in Table 4 can be hybridized to any antisense strand containing a sequence listed in Table 3 provided the two sequences have a region of at least about 85% complementarity over a contiguous 16, 17, 18, 19, 20, or 21 nucleotide sequence. Representative sequence pairings are exemplified by the Duplex ID Nos. shown in Table 5.
- 15 In some embodiments, an HBV RNAi agent comprises of any of the Duplex ID Nos. presented herein. In some embodiments, an HBV RNAi agent consists of any of the Duplex ID Nos. presented herein. In some embodiments, an HBV RNAi agent comprises the sense strand and/or the antisense strand nucleotide sequences of any of the Duplex ID Nos. presented herein. In some embodiments, an HBV RNAi agent comprises the sense strand and antisense strand
20 nucleotide sequences of any of the Duplex ID Nos. presented herein and a targeting group and/or linking group wherein the targeting group and/or linking group is covalently linked (i.e. conjugated) to the sense strand or the antisense strand. In some embodiments, an HBV RNAi agent comprises the sense strand and antisense strand modified nucleotide sequences of any of the Duplex ID Nos. presented herein. In some embodiments, an HBV RNAi agent comprises
25 the sense strand and antisense strand modified nucleotide sequences of any of the Duplex ID Nos. presented herein and a targeting group and/or linking group wherein the targeting group and/or linking group is covalently linked to the sense strand or the antisense strand.

- 30 In some embodiments, an HBV RNAi agent comprises an antisense strand and a sense strand having the nucleotide sequences of any of the antisense strand/sense strand duplexes of Table 5, and further comprises an asialoglycoprotein receptor ligand targeting group.

In some embodiments, an HBV RNAi agent comprises an antisense strand and a sense strand having the nucleotide sequences of any of the antisense strand and/or sense strand nucleotide sequences of any of the duplexes of Table 5, and further comprises a targeting group selected from the group consisting of (PAZ), (NAG13), (NAG13)s, (NAG18), (NAG18)s, (NAG24),
 5 (NAG24)s, (NAG25), (NAG25)s, (NAG26), (NAG26)s, (NAG27), (NAG27)s, (NAG28), (NAG28)s, (NAG29), (NAG29)s, (NAG30), (NAG30)s, (NAG31), (NAG31)s, (NAG32), (NAG32)s, (NAG33), (NAG33)s, (NAG34), (NAG34)s, (NAG35), (NAG35)s, (NAG36), (NAG36)s, (NAG37), (NAG37)s.

10 In some embodiments, an HBV RNAi agent comprises an antisense strand and a sense strand having the modified nucleotide sequences of any of the antisense strand and/or sense strand nucleotide sequences of any of the duplexes of Table 5.

In some embodiments, an HBV RNAi agent comprises an antisense strand and a sense strand
 15 having the modified nucleotide sequences of any of the antisense strand and/or sense strand nucleotide sequences of any of the duplexes of Table 5, and further comprises an asialoglycoprotein receptor ligand targeting group.

In some embodiments, an HBV RNAi agent comprises any of the duplexes of Table 5.
 20

In some embodiments, an HBV RNAi agent consists of any of the duplexes of Table 5.

Table 5. Examples of HBV RNAi agent duplexes.

Duplex ID	Antisense Strand ID	Sense Strand ID	Duplex ID	Antisense Strand ID	Sense Strand ID
AD03498	AM03508-AS	AM04445-SS	AD04426	AM05623-AS	AM05622-SS
AD03499	AM04441-AS	AM04444-SS	AD04427	AM05623-AS	AM05624-SS
AD03500	AM04442-AS	AM04444-SS	AD04428	AM05626-AS	AM05622-SS
AD03501	AM04443-AS	AM04444-SS	AD04429	AM05626-AS	AM05624-SS
AD03738	AM04768-AS	AM04767-SS	AD04430	AM05628-AS	AM05627-SS
AD03739	AM04769-AS	AM04767-SS	AD04431	AM05054-AS	AM05629-SS
AD03967	AM04443-AS	AM05010-SS	AD04432	AM05054-AS	AM05630-SS
AD03968	AM05011-AS	AM05010-SS	AD04433	AM05631-AS	AM05048-SS
AD03969	AM04443-AS	AM05015-SS	AD04434	AM05632-AS	AM05048-SS
AD03970	AM05011-AS	AM05019-SS	AD04435	AM05633-AS	AM05048-SS
AD03971	AM05012-AS	AM05015-SS	AD04436	AM05635-AS	AM05048-SS
AD03972	AM04443-AS	AM05016-SS	AD04437	AM05634-AS	AM05048-SS
AD03973	AM04443-AS	AM05017-SS	AD04438	AM05637-AS	AM05636-SS
AD03974	AM04443-AS	AM05018-SS	AD04439	AM05638-AS	AM05636-SS
AD03975	AM05013-AS	AM05015-SS	AD04440	AM05058-AS	AM05639-SS
AD03976	AM05014-AS	AM05019-SS	AD04441	AM05057-AS	AM05639-SS
AD03977	AM05013-AS	AM05017-SS	AD04442	AM05057-AS	AM05640-SS
AD03978	AM05013-AS	AM04444-SS	AD04511	AM05747-AS	AM05746-SS
AD04001	AM05052-AS	AM05034-SS	AD04570	AM05011-AS	AM05856-SS
AD04002	AM05053-AS	AM05034-SS	AD04571	AM05849-AS	AM05856-SS
AD04003	AM05054-AS	AM05046-SS	AD04572	AM05850-AS	AM05856-SS
AD04004	AM05052-AS	AM05047-SS	AD04573	AM05851-AS	AM05857-SS
AD04005	AM05055-AS	AM05064-SS	AD04574	AM05852-AS	AM05857-SS
AD04006	AM05056-AS	AM05048-SS	AD04575	AM05853-AS	AM05858-SS
AD04007	AM05057-AS	AM05048-SS	AD04576	AM05854-AS	AM05858-SS
AD04008	AM05058-AS	AM05049-SS	AD04577	AM05011-AS	AM05859-SS
AD04009	AM05056-AS	AM05050-SS	AD04578	AM05850-AS	AM05858-SS
AD04010	AM05060-AS	AM05063-SS	AD04579	AM05014-AS	AM05347-SS
AD04176	AM05351-AS	AM05346-SS	AD04580	AM05855-AS	AM05347-SS
AD04177	AM04443-AS	AM05347-SS	AD04581	AM05860-AS	AM05063-SS
AD04178	AM05011-AS	AM05347-SS	AD04583	AM05862-AS	AM05868-SS
AD04412	AM05011-AS	AM05606-SS	AD04584	AM05863-AS	AM05868-SS
AD04413	AM05011-AS	AM05607-SS	AD04585	AM05864-AS	AM05869-SS
AD04414	AM05608-AS	AM05606-SS	AD04586	AM05865-AS	AM05869-SS
AD04415	AM05011-AS	AM05615-SS	AD04587	AM05862-AS	AM05870-SS
AD04416	AM05609-AS	AM05616-SS	AD04588	AM05863-AS	AM05871-SS
AD04417	AM05610-AS	AM05616-SS	AD04590	AM05867-AS	AM05063-SS
AD04418	AM05611-AS	AM05616-SS	AD04591	AM05860-AS	AM05872-SS
AD04419	AM05612-AS	AM05616-SS	AD04592	AM05054-AS	AM05879-SS
AD04420	AM05613-AS	AM05616-SS	AD04593	AM05873-AS	AM05880-SS
AD04421	AM05614-AS	AM05616-SS	AD04594	AM05874-AS	AM05880-SS
AD04422	AM05054-AS	AM05617-SS	AD04595	AM05875-AS	AM05881-SS
AD04423	AM05618-AS	AM05046-SS	AD04596	AM05876-AS	AM05881-SS
AD04425	AM05621-AS	AM05620-SS	AD04597	AM05873-AS	AM05882-SS

Duplex ID	Antisense Strand ID	Sense Strand ID
AD04598	AM05874-AS	AM05883-SS
AD04599	AM05877-AS	AM05620-SS
AD04734	AM06074-AS	AM05869-SS
AD04771	AM06142-AS	AM06146-SS
AD04772	AM06143-AS	AM06147-SS
AD04773	AM06144-AS	AM06146-SS
AD04774	AM06145-AS	AM06148-SS
AD04775	AM06145-AS	AM06149-SS
AD04776	AM05850-AS	AM06150-SS
AD04777	AM05854-AS	AM06151-SS
AD04778	AM05854-AS	AM06152-SS
AD04822	AM06222-AS	AM06146-SS
AD04823	AM05609-AS	AM06150-SS
AD04871	AM06281-AS	AM06287-SS
AD04872	AM06282-AS	AM06288-SS
AD04873	AM06283-AS	AM06288-SS
AD04874	AM06284-AS	AM06289-SS
AD04875	AM06285-AS	AM06290-SS
AD04876	AM06286-AS	AM06291-SS
AD04881	AM06299-AS	AM06304-SS
AD04882	AM06300-AS	AM06305-SS
AD04883	AM06301-AS	AM06306-SS
AD04884	AM06302-AS	AM06307-SS
AD04885	AM06303-AS	AM06308-SS
AD04962	AM05864-AS	AM06146-SS
AD04963	AM05855-AS	AM05607-SS
AD04981	AM06463-AS	AM06150-SS
AD04982	AM06464-AS	AM06150-SS
AD04983	AM06465-AS	AM06150-SS
AD05069	AM06604-AS	AM06603-SS
AD05070	AM06606-AS	AM06605-SS
AD05071	AM06608-AS	AM06607-SS
AD05072	AM05011-AS	AM06609-SS
AD05073	AM06611-AS	AM06610-SS
AD05074	AM06612-AS	AM06150-SS
AD05075	AM06614-AS	AM06613-SS
AD05076	AM06616-AS	AM06615-SS
AD05077	AM06618-AS	AM06617-SS
AD05078	AM06620-AS	AM06619-SS
AD05147	AM06751-AS	AM06750-SS
AD05148	AM06606-AS	AM06752-SS
AD05149	AM06751-AS	AM06753-SS
AD05164	AM06282-AS	AM06776-SS
AD05165	AM06606-AS	AM06777-SS

In some embodiments, an HBV RNAi agent is prepared or provided as a salt, mixed salt, or a free-acid. The RNAi agents described herein, upon delivery to a cell expressing an HBV gene, inhibit or knockdown expression of one or more HBV genes *in vivo*.

5 Targeting Groups, Linking Groups, and Delivery Vehicles

In some embodiments, an HBV RNAi agent is conjugated to one or more non-nucleotide groups including, but not limited to a targeting group, linking group, delivery polymer, or a delivery vehicle. The non-nucleotide group can enhance targeting, delivery or attachment of the RNAi agent. Examples of targeting groups and linking groups are provided in Table 6. The
 10 non-nucleotide group can be covalently linked to the 3' and/or 5' end of either the sense strand and/or the antisense strand. In some embodiments, an HBV RNAi agent contains a non-nucleotide group linked to the 3' and/or 5' end of the sense strand. In some embodiments, a non-nucleotide group is linked to the 5' end of an HBV RNAi agent sense strand. A non-nucleotide group may be linked directly or indirectly to the RNAi agent via a linker/linking
 15 group. In some embodiments, a non-nucleotide group is linked to the RNAi agent via a labile, cleavable, or reversible bond or linker.

In some embodiments, a non-nucleotide group enhances the pharmacokinetic or biodistribution properties of an RNAi agent or conjugate to which it is attached to improve
 20 cell- or tissue-specific distribution and cell-specific uptake of the conjugate. In some embodiments, a non-nucleotide group enhances endocytosis of the RNAi agent.

Targeting groups or targeting moieties enhance the pharmacokinetic or biodistribution properties of a conjugate to which they are attached to improve cell-specific distribution and
 25 cell-specific uptake of the conjugate. A targeting group can be monovalent, divalent, trivalent, tetravalent, or have higher valency. Representative targeting groups include, without limitation, compounds with affinity to cell surface molecule, cell receptor ligands, hapten, antibodies, monoclonal antibodies, antibody fragments, and antibody mimics with affinity to cell surface molecules. In some embodiments, a targeting group is linked to an RNAi agent
 30 using a linker, such as a PEG linker or one, two, or three abasic and/or ribitol (abasic ribose) groups. In some embodiments, a targeting group comprises a galactose derivative cluster.

The HBV RNAi agents described herein may be synthesized having a reactive group, such as an amine group, at the 5'-terminus. The reactive group may be used to subsequently attach a targeting moiety using methods typical in the art.

5 In some embodiments, a targeting group comprises an asialoglycoprotein receptor ligand. In some embodiments, an asialoglycoprotein receptor ligand includes or consists of one or more galactose derivatives. As used herein, the term galactose derivative includes both galactose and derivatives of galactose having affinity for the asialoglycoprotein receptor that is equal to or greater than that of galactose. Galactose derivatives include, but are not limited to: galactose,
 10 galactosamine, N-formylgalactosamine, N-acetyl-galactosamine, N-propionyl-galactosamine, N-n-butanoyl-galactosamine, and N-iso-butanoylgalactos-amine (see for example: Iobst, S.T. and Drickamer, K. *J.B.C.* 1996, 271, 6686). Galactose derivatives, and clusters of galactose derivatives, that are useful for *in vivo* targeting of oligonucleotides and other molecules to the liver are known in the art (see, for example, Baenziger and Fiete, 1980, *Cell*, 22, 611-620; Connolly et al., 1982, *J. Biol. Chem.*, 257, 939-945). Galactose derivatives have been used to target molecules to hepatocytes *in vivo* through their binding to the asialoglycoprotein receptor (ASGPr) expressed on the surface of hepatocytes. Binding of ASGPr ligands to the ASGPr(s) facilitates cell-specific targeting to hepatocytes and endocytosis of the molecule into hepatocytes. ASGPr ligands can be monomeric (e.g., having a single galactose derivative) or
 20 multimeric (e.g., having multiple galactose derivatives). The galactose derivative or galactose derivative cluster may be attached to the 3' or 5' end of the RNAi polynucleotide using methods known in the art. The preparation of targeting groups, such as galactose derivative clusters, is described in, for example, U.S. Patent Application Serial Nos. 15/452,324 and 15/452,423, the contents of both of which are incorporated herein in their entirety.

25 As used herein, a galactose derivative cluster comprises a molecule having two to four terminal galactose derivatives. A terminal galactose derivative is attached to a molecule through its C-1 carbon. In some embodiments, the galactose derivative cluster is a galactose derivative trimer (also referred to as tri-antennary galactose derivative or tri-valent galactose derivative). In
 30 some embodiments, the galactose derivative cluster comprises N-acetyl-galactosamines. In some embodiments, the galactose derivative cluster comprises three N-acetyl-galactosamines. In some embodiments, the galactose derivative cluster is a galactose derivative tetramer (also

referred to as tetra-antennary galactose derivative or tetra-valent galactose derivative). In some embodiments, the galactose derivative cluster comprises four N-acetyl-galactosamines.

As used herein, a galactose derivative trimer contains three galactose derivatives, each linked to a central branch point. As used herein, a galactose derivative tetramer contains four galactose derivatives, each linked to a central branch point. The galactose derivatives can be attached to the central branch point through the C-1 carbons of the saccharides. In some embodiments, the galactose derivatives are linked to the branch point via linkers or spacers. In some embodiments, the linker or spacer is a flexible hydrophilic spacer, such as a PEG group (see, for example, U.S. Patent No. 5,885,968; Biessen et al. J. Med. Chem. 1995 Vol. 39 p. 1538-1546). In some embodiments, the PEG spacer is a PEG₃ spacer. The branch point can be any small molecule which permits attachment of three galactose derivatives and further permits attachment of the branch point to the RNAi agent. An example of branch point group is a di-lysine or di-glutamate. Attachment of the branch point to the RNAi agent can occur through a linker or spacer. In some embodiments, the linker or spacer comprises a flexible hydrophilic spacer, such as, but not limited to, a PEG spacer. In some embodiments, the linker comprises a rigid linker, such as a cyclic group. In some embodiments, a galactose derivative comprises or consists of N-acetyl-galactosamine. In some embodiments, the galactose derivative cluster is comprised of a galactose derivative tetramer, which can be, for example, an N-acetyl-galactosamine tetramer.

In some embodiments, pharmaceutical compositions for delivering an HBV RNAi agent to a liver cell *in vivo* are described. Such pharmaceutical compositions can include, for example, an HBV RNAi agent conjugated to a galactose derivative cluster. In some embodiments, the galactose derivative cluster is comprised of a galactose derivative trimer, which can be, for example, an N-acetyl-galactosamine trimer, or galactose derivative tetramer, which can be, for example, an N-acetyl-galactosamine tetramer.

Targeting groups include, but are not limited to, (PAZ), (NAG13), (NAG13)s, (NAG18), (NAG18)s, (NAG24), (NAG24)s, (NAG25), (NAG25)s, (NAG26), (NAG26)s, (NAG27), (NAG27)s, (NAG28), (NAG28)s, (NAG29), (NAG29)s, (NAG30), (NAG30)s, (NAG31), (NAG31)s, (NAG32), (NAG32)s, (NAG33), (NAG33)s, (NAG34), (NAG34)s, (NAG35), (NAG35)s, (NAG36), (NAG36)s, (NAG37), (NAG37)s, (NAG38), (NAG38)s, (NAG39), and

(NAG39)s. Other targeting groups, including galactose cluster targeting ligands, are known in the art.

In some embodiments, a linking group is conjugated to the RNAi agent. The linking group facilitates covalent linkage of the agent to a targeting group or delivery polymer or delivery vehicle. The linking group can be linked to the 3' or the 5' end of the RNAi agent sense strand or antisense strand. In some embodiments, the linking group is linked to the RNAi agent sense strand. In some embodiments, the linking group is conjugated to the 5' or 3' end of an RNAi agent sense strand. In some embodiments, a linking group is conjugated to the 5' end of an RNAi agent sense strand. Examples of linking groups, include, but are not limited to: reactive groups such as primary amines and alkynes, alkyl groups, abasic nucleosides, ribitol (abasic ribose), and/or PEG groups.





A linker or linking group is a connection between two atoms that links one chemical group (such as an RNAi agent) or segment of interest to another chemical group (such as a targeting group or delivery polymer) or segment of interest via one or more covalent bonds. A labile linkage contains a labile bond. A linkage may optionally include a spacer that increases the distance between the two joined atoms. A spacer may further add flexibility and/or length to the linkage. Spacers may include, but are not be limited to, alkyl groups, alkenyl groups, alkynyl groups, aryl groups, aralkyl groups, aralkenyl groups, and aralkynyl groups; each of which can contain one or more heteroatoms, heterocycles, amino acids, nucleotides, and saccharides. Spacer groups are well known in the art and the preceding list is not meant to limit the scope of the description.


Any of the HBV RNAi agent nucleotide sequences listed in Tables 3 and 4, whether modified or unmodified, may contain 3' or 5' targeting group and/or linking group. Any of the HBV RNAi agent sequences listed in Table 3 and 4 which contain a 3' or 5' targeting group and/or linking group, may alternatively contain no 3' or 5' targeting group and/or linking group, or may contain a different 3' or 5' targeting group and/or linking group including, but not limited to, those depicted in Table 3. Any of the HBV RNAi agent duplexes listed in Table 5, whether modified or unmodified, may further comprise a targeting group and/or linking group, including, but not limited to, those depicted in Table 3, and the targeting group or linking group

may be attached to the 3' or 5' terminus of either the sense strand or the antisense strand of the HBV RNAi agent duplex.


Examples of targeting groups and linking groups are provided in Table 6. Table 4 provides
5 several embodiments of HBV RNAi agent sense strands having a targeting group or linking group linked to the 5' or 3' end.

Table 6. Structures representing various modified nucleotides, targeting groups, and linking groups.


 <p>vpdT</p>	 <p>5Me-Gf</p>
 <p>cPrpTM</p>	 <p>cPrpu</p>



When positioned internally on oligonucleotide:




When positioned internally on oligonucleotide:




When positioned at the 3' terminal end of oligonucleotide:


linkage towards 5' end of
oligonucleotide






(invAb)









(PAZ)









(NAG13)




































In each of the above structures in Table 6, NAG comprises an N-acetyl-galactosamine or another ASGPr ligand, as would be understood by a person of ordinary skill in the art to be attached in view of the structures above and description provided herein. For example, in some


5 embodiments, NAG in the structures provided in Table 6 is represented by the following structure:




(N-acetyl-galactosamine)

Each (NAGx) may be attached to an HBV RNAi agent via a phosphate group (as in (NAG25), (NAG30), and (NAG31)), or a phosphorothioate group, (as is (NAG25)s, (NAG29)s, (NAG30)s, (NAG31)s, or (NAG37)s), or another linking group.

5



Phosphate group



Phosphorothioate group

Other linking groups known in the art may be used.

10

Delivery Vehicles

In some embodiments, a delivery vehicle may be used to deliver an RNAi agent to a cell or tissue. A delivery vehicle is a compound that improves delivery of the RNAi agent to a cell or tissue. A delivery vehicle can include, or consist of, but is not limited to: a polymer, such as an amphipathic polymer, a membrane active polymer, a peptide, a melittin peptide, a melittin-like peptide (MLP), a lipid, a reversibly modified polymer or peptide, or a reversibly modified membrane active polyamine.

In some embodiments, the RNAi agents can be combined with lipids, nanoparticles, polymers, liposomes, micelles, DPCs or other delivery systems available in the art. The RNAi agents can also be chemically conjugated to targeting groups, lipids (including, but not limited to cholesterol and cholesteryl derivatives), nanoparticles, polymers, liposomes, micelles, DPCs (see, for example WO 2000/053722, WO 2008/0022309, WO 2011/104169, and WO 2012/083185, WO 2013/032829, WO 2013/158141, each of which is incorporated herein by reference), or other delivery systems available in the art.

25

Pharmaceutical Compositions and Formulations

The HBV RNAi agents disclosed herein may be prepared as pharmaceutical compositions or formulations. In some embodiments, pharmaceutical compositions include at least one HBV RNAi agent. These pharmaceutical compositions are particularly useful in the inhibition of the expression of the target mRNA in a target cell, a group of cells, a tissue, or an organism. The pharmaceutical compositions can be used to treat a subject having a disease or disorder that

30

would benefit from reduction in the level of the target mRNA, or inhibition in expression of the target gene. The pharmaceutical compositions can be used to treat a subject at risk of developing a disease or disorder that would benefit from reduction of the level of the target mRNA or an inhibition in expression the target gene. In one embodiment, the method includes
5 administering an HBV RNAi agent linked to a targeting ligand as described herein, to a subject to be treated. In some embodiments, one or more pharmaceutically acceptable excipients (including vehicles, carriers, diluents, and/or delivery polymers) are added to the pharmaceutical compositions including an HBV RNAi agent, thereby forming a pharmaceutical formulation suitable for *in vivo* delivery to a human.

10

The pharmaceutical compositions that include an HBV RNAi agent and methods disclosed herein may decrease the level of the target mRNA in a cell, group of cells, group of cells, tissue, or subject, including: administering to the subject a therapeutically effective amount of a herein described HBV RNAi agent, thereby inhibiting the expression of a target mRNA in
15 the subject.

In some embodiments, the described pharmaceutical compositions including an HBV RNAi agent are used for treating or managing clinical presentations associated with HBV infection. In some embodiments, a therapeutically or prophylactically effective amount of one or more
20 of pharmaceutical compositions is administered to a subject in need of such treatment, prevention or management. In some embodiments, administration of any of the disclosed HBV RNAi agents can be used to decrease the number, severity, and/or frequency of symptoms of a disease in a subject.

25 The described pharmaceutical compositions including an HBV RNAi agent can be used to treat at least one symptom in a subject having a disease or disorder that would benefit from reduction or inhibition in expression of HBV mRNA. In some embodiments, the subject is administered a therapeutically effective amount of one or more pharmaceutical compositions including an HBV RNAi agent thereby treating the symptom. In other embodiments, the subject is
30 administered a prophylactically effective amount of one or more HBV RNAi agents, thereby preventing the at least one symptom.

The route of administration is the path by which an HBV RNAi agent is brought into contact with the body. In general, methods of administering drugs and nucleic acids for treatment of a mammal are well known in the art and can be applied to administration of the compositions described herein. The HBV RNAi agents disclosed herein can be administered via any suitable route in a preparation appropriately tailored to the particular route. Thus, herein described pharmaceutical compositions can be administered by injection, for example, intravenously, intramuscularly, intracutaneously, subcutaneously, intraarticularly, or intraperitoneally. In some embodiments, there herein described pharmaceutical compositions via subcutaneous injection.

10

The pharmaceutical compositions including an HBV RNAi agent described herein can be delivered to a cell, group of cells, tumor, tissue, or subject using oligonucleotide delivery technologies known in the art. In general, any suitable method recognized in the art for delivering a nucleic acid molecule (in vitro or in vivo) can be adapted for use with a herein described compositions. For example, delivery can be by local administration, (e.g., direct injection, implantation, or topical administering), systemic administration, or subcutaneous, intravenous, intraperitoneal, or parenteral routes, including intracranial (e.g., intraventricular, intraparenchymal and intrathecal), intramuscular, transdermal, airway (aerosol), nasal, oral, rectal, or topical (including buccal and sublingual) administration. In certain embodiments, the compositions are administered by subcutaneous or intravenous infusion or injection.

20

Accordingly, in some embodiments, the herein described pharmaceutical compositions may comprise one or more pharmaceutically acceptable excipients. In some embodiments, the pharmaceutical compositions described herein can be formulated for administration to a subject.

25

As used herein, a pharmaceutical composition or medicament includes a pharmacologically effective amount of at least one of the described therapeutic compounds and one or more pharmaceutically acceptable excipients. Pharmaceutically acceptable excipients (excipients) are substances other than the Active Pharmaceutical ingredient (API, therapeutic product, e.g., HBV RNAi agent) that are intentionally included in the drug delivery system. Excipients do not exert or are not intended to exert a therapeutic effect at the intended dosage. Excipients may act to a) aid in processing of the drug delivery system during manufacture, b) protect,

30

support or enhance stability, bioavailability or patient acceptability of the API, c) assist in product identification, and/or d) enhance any other attribute of the overall safety, effectiveness, of delivery of the API during storage or use. A pharmaceutically acceptable excipient may or may not be an inert substance.

5

Excipients include, but are not limited to: absorption enhancers, anti-adherents, anti-foaming agents, anti-oxidants, binders, buffering agents, carriers, coating agents, colors, delivery enhancers, delivery polymers, dextran, dextrose, diluents, disintegrants, emulsifiers, extenders, fillers, flavors, glidants, humectants, lubricants, oils, polymers, preservatives, saline, salts, solvents, sugars, suspending agents, sustained release matrices, sweeteners, thickening agents, tonicity agents, vehicles, water-repelling agents, and wetting agents.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline. It should be stable under the conditions of manufacture and storage and should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, and sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile

powders for the preparation of sterile injectable solutions, methods of preparation include vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

- 5 Formulations suitable for intra-articular administration can be in the form of a sterile aqueous preparation of the drug that can be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems can also be used to present the drug for both intra-articular and ophthalmic administration.

10

- The active compounds can be prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. Liposomal suspensions can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

- 20 The HBV RNAi agents can be formulated in compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the disclosure are dictated by and directly dependent on the unique characteristics of the active compound and the therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

- 30 A pharmaceutical composition can contain other additional components commonly found in pharmaceutical compositions. Such additional components include, but are not limited to: anti-pruritics, astringents, local anesthetics, or anti-inflammatory agents (e.g., antihistamine, diphenhydramine, etc.). It is also envisioned that cells, tissues or isolated organs that express or comprise the herein defined RNAi agents may be used as “pharmaceutical compositions.”

As used herein, “pharmacologically effective amount,” “therapeutically effective amount,” or simply “effective amount” refers to that amount of an RNAi agent to produce a pharmacological, therapeutic or preventive result.

5 Generally, an effective amount of an active compound will be in the range of from about 0.1 to about 100 mg/kg of body weight/day, e.g., from about 1.0 to about 50 mg/kg of body weight/day. In some embodiments, an effective amount of an active compound will be in the range of from about 0.25 to about 5 mg/kg of body weight per dose. In some embodiments, an effective amount of an active ingredient will be in the range of from about 0.5 to about 3
10 mg/kg of body weight per dose. The amount administered will also likely depend on such variables as the overall health status of the patient, the relative biological efficacy of the compound delivered, the formulation of the drug, the presence and types of excipients in the formulation, and the route of administration. Also, it is to be understood that the initial dosage administered can be increased beyond the above upper level in order to rapidly achieve the
15 desired blood-level or tissue level, or the initial dosage can be smaller than the optimum.

For treatment of disease or for formation of a medicament or composition for treatment of a disease, the pharmaceutical compositions described herein including an HBV RNAi agent can be combined with an excipient or with a second therapeutic agent or treatment including, but
20 not limited to: a second or other RNAi agent, a small molecule drug, an antibody, an antibody fragment, and/or a vaccine.

The described HBV RNAi agents, when added to pharmaceutically acceptable excipients or adjuvants, can be packaged into kits, containers, packs, or dispensers. The pharmaceutical
25 compositions described herein may be packaged in pre-filled syringes or vials.

Methods of Treatment and Inhibition of Expression

The HBV RNAi agents disclosed herein can be used to treat a subject (e.g., a human or mammal) having a disease or disorder that would benefit from administration of the compound.
30 In some embodiments, the RNAi agents disclosed herein can be used to treat a subject (e.g., a human) having a disease or disorder that would benefit from reduction or inhibition in expression of HBV mRNA. The subject is administered a therapeutically effective amount of any one or more RNAi agents. The subject can be a human, patient, or human patient. The

subject may be an adult, adolescent, child, or infant. The described pharmaceutical compositions including an HBV RNAi agent can be used to provide methods for the therapeutic treatment of diseases. Such methods include administration of a pharmaceutical composition described herein to a human being or animal.

5

In some embodiments, the HBV RNAi agents described herein are used to treat a subject infected with HBV. In some embodiments, the described HBV RNAi agents are used to treat at least one symptom in a subject having a HBV infection. The subject is administered a therapeutically effective amount of any one or more of the described RNAi agents.

10

In some embodiments, the subject has both a HBV infection and a HDV infection. In some embodiments, the HBV RNAi agents described herein are used to treat a subject infected with both HBV and HDV. In some embodiments, the described HBV RNAi agents are used to treat at least one symptom in a subject having a HBV or a HDV infection. The subject is administered a therapeutically effective amount of any one or more of the described RNAi agents.

15

In some embodiments, the HBV RNAi agents are used to treat or manage a clinical presentation wherein a subject infected with HBV. The subject is administered a therapeutically or effective amount of one or more of the HBV RNAi agents or HBV RNAi agent-containing compositions described herein. In some embodiments, the method comprises administering a composition comprising an HBV RNAi agent described herein to a subject to be treated.

20

In some embodiments, the gene expression level and/or mRNA level of an HBV gene in a subject to whom a described HBV RNAi agent is administered is reduced by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater than 99% relative to the subject prior to being administered the HBV RNAi agent or to a subject not receiving the HBV RNAi agent. The gene expression level and/or mRNA level in the subject may be reduced in a cell, group of cells, and/or tissue of the subject. In some embodiments, the expressed protein level of an HBV gene in a subject to whom a described HBV RNAi agent has been administered is reduced by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater than 99% relative to the subject

25

30

prior to being administered the HBV RNAi agent or to a subject not receiving the HBV RNAi agent. The protein level in the subject may be reduced in a cell, group of cells, tissue, blood, and/or other fluid of the subject. For example, in some embodiments, the amount or level of Hepatitis B surface antigen (HBsAg) in a subject to whom a described HBV RNAi agent has been administered is reduced by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater than 99% relative to the subject prior to being administered the HBV RNAi agent or to a subject not receiving the HBV RNAi agent. In some embodiments, the amount or level of Hepatitis B e-antigen (HBeAg) in a subject to whom a described HBV RNAi agent has been administered is reduced by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater than 99% relative to the subject prior to being administered the HBV RNAi agent or to a subject not receiving the HBV RNAi agent. In some embodiments, the amount or level of serum HBV DNA in a subject to whom a described HBV RNAi agent has been administered is reduced by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater than 99% relative to the subject prior to being administered the HBV RNAi agent or to a subject not receiving the HBV RNAi agent. A reduction in the presence of serum HBV DNA, HBV gene expression, HBV mRNA, or HBV protein amounts or levels may be assessed by methods known in the art. Reduction or decrease in HBV mRNA amount or level, expressed protein amount or level, and/or serum HBV DNA amount or level, are collectively referred to herein as a reduction or decrease in HBV or inhibiting or reducing the expression of HBV.

Cells and Tissues and non-Human organisms

Cells, tissues, and non-human organisms that include at least one of the HBV RNAi agents described herein is contemplated. The cell, tissue, or non-human organism is made by delivering the RNAi agent to the cell, tissue, or non-human organism.

The above provided embodiments and items are now illustrated with the following, non-limiting examples.

EXAMPLES

Example 1. Synthesis of HBV RNAi agents.

HBV RNAi agent duplexes shown in Table 5 were synthesized in accordance with the following:

A. *Synthesis.* The sense and antisense strands of the HBV RNAi agents were synthesized according to phosphoramidite technology on solid phase used in oligonucleotide synthesis. Depending on the scale, either a MerMade96E® (Bioautomation), a MerMade12® (Bioautomation), or an OP Pilot 100 (GE Healthcare) was used. Syntheses were performed on a solid support made of controlled pore glass (CPG, 500 Å or 600Å, obtained from Prime Synthesis, Aston, PA, USA). All RNA and 2'-modified phosphoramidites were purchased from Thermo Fisher Scientific (Milwaukee, WI, USA). Specifically, the following 2'-O-methyl phosphoramidites were used: (5'-O-dimethoxytrityl-N⁶-(benzoyl)-2'-O-methyl-adenosine-3'-O-(2-cyanoethyl-N,N-diisopropylamino) phosphoramidite, 5'-O-dimethoxy-trityl-N⁴-(acetyl)-2'-O-methyl-cytidine-3'-O-(2-cyanoethyl-N,N-diisopropyl-amino) phosphoramidite, (5'-O-dimethoxytrityl-N²-(isobutyryl)-2'-O-methyl-guanosine-3'-O-(2-cyanoethyl-N,N-diisopropylamino) phosphoramidite, and 5'-O-dimethoxytrityl-2'-O-methyl-uridine-3'-O-(2-cyanoethyl-N,N-diisopropylamino) phosphoramidite. The 2'-deoxy-2'-fluoro-phosphoramidites carried the same protecting groups as the 2'-O-methyl amidites. The abasic (3'-O-dimethoxytrityl-2'-deoxyribose-5'-O-(2-cyanoethyl-N,N-diisopropylamino) phosphoramidites were purchased from ChemGenes (Wilmington, MA, USA). Targeting ligand containing phosphoramidites were dissolved in anhydrous dichloromethane or anhydrous acetonitrile (50 mM), while all other amidites were dissolved in anhydrous acetonitrile (50 mM) and molecular sieves (3Å) were added. 5-Benzylthio-1H-tetrazole (BTT, 250 mM in acetonitrile) or 5-Ethylthio-1H-tetrazole (ETT, 250 mM in acetonitrile) was used as activator solution. Coupling times were 12 min (RNA), 15 min (targeting ligand), 90 sec (2'OMe), and 60 sec (2'F). In order to introduce phosphorothioate linkages, a 100 mM solution of 3-phenyl 1,2,4-dithiazoline-5-one (POS, obtained from PolyOrg, Inc., Leominster, MA, USA) in anhydrous Acetonitrile was employed.

B. *Cleavage and deprotection of support bound oligomer.* After finalization of the solid phase synthesis, the dried solid support was treated with a 1:1 volume solution of 40 wt. % methylamine in water and 28% ammonium hydroxide solution (Aldrich) for 1.5 hours at 30°C. The solution was evaporated and the solid residue was reconstituted in water (see below).

C. *Purification.* Crude oligomers were purified by anionic exchange HPLC using a TSKgel SuperQ-5PW 13 μ m column and Shimadzu LC-8 system. Buffer A was 20 mM Tris, 5 mM EDTA, pH 9.0 and contained 20% Acetonitrile and buffer B was the same as buffer A with the addition of 1.5 M sodium chloride. UV traces at 260 nm were recorded. Appropriate
5 fractions were pooled then run on size exclusion HPLC using a GE Healthcare XK 26/40 column packed with Sephadex G-25 fine with a running buffer of filtered DI water or 100mM ammonium bicarbonate, pH 6.7 and 20% Acetonitrile.

D. *Annealing.* Complementary strands were mixed by combining equimolar RNA
10 solutions (sense and antisense) in 1 \times Phosphate-Buffered Saline (Corning, Cellgro) to form the RNAi agents. Some RNAi agents were lyophilized and stored at -15 to -25°C . Duplex concentration was determined by measuring the solution absorbance on a UV-Vis spectrometer in 1 \times Phosphate-Buffered Saline. The solution absorbance at 260 nm was then multiplied by a conversion factor and the dilution factor to determine the duplex concentration. Unless
15 otherwise stated, all conversion factor was 0.037 mg/(mL \cdot cm). For some experiments, a conversion factor was calculated from an experimentally determined extinction coefficient.

Example 2. pHBV model mice.

Six to eight-week-old female NOD.CB17-Prkdcscid/NcrCrl (NOD-SCID) mice were
20 transiently transfected *in vivo* with MC-HBV1.3 by hydrodynamic tail vein injection (Yang PL et al. "Hydrodynamic injection of viral DNA: a mouse model of acute hepatitis B virus infection," *PNAS USA* 2002 Vol. 99: p. 13825-13830), administered 30 to 45 days prior to administration of an HBV RNAi agent or control. MC-HBV1.3 is a plasmid-derived minicircle that contains the same terminally redundant human hepatitis B virus sequence HBV1.3 as in
25 plasmid pHBV1.3 and in the HBV1.3.32 transgenic mice (GenBank accession #V01460) (Guidotti LG et al., "High-level hepatitis B virus replication in transgenic mice," *J Virol* 1995 Vol. 69, p6158-6169.). 5 or 10 μ g MC-HBV1.3 in Ringer's Solution in a total volume of 10% of the animal's body weight was injected into mice via tail vein to create pHBV model of chronic HBV infection. The solution was injected through a 27-gauge needle in 5-7 seconds
30 as previously described (Zhang G et al., "High levels of foreign gene expression in hepatocytes after tail vein injection of naked plasmid DNA." *Human Gene Therapy* 1999 Vol. 10, p1735-1737.). At pre-dose (either day 1 pre-dose, day -1, or day -2), Hepatitis B surface antigen

(HBsAg) HBsAg expression levels in serum were measured by ELISA and the mice were grouped according to average HBsAg expression levels.

Analyses: At various times, before and after administration of HBV RNAi agents, serum
 5 HBsAg, serum HBeAg, serum HBV DNA, or liver HBV RNA may be measured. HBV expression levels were normalized to pre-administration expression levels and to control mice injected with phosphate buffered saline (“PBS”).

i) Serum collection: Mice were anesthetized with 2-3% isoflurane and blood samples
 10 were collected from the submandibular area into serum separation tubes (Sarstedt AG & Co., Nümbrecht, Germany). Blood was allowed to coagulate at ambient temperature for 20 min. The tubes were centrifuged at 8,000 ×g for 3 min to separate the serum and stored at 4°C.

ii) Serum Hepatitis B surface antigen (HBsAg) levels: Serum was collected and diluted
 15 10 to 8000-fold in PBS containing 5% nonfat dry milk. Secondary HBsAg standards diluted in the nonfat milk solution were prepared from serum of ICR mice (Harlan Sprague Dawley) that had been transfected with 10 µg HBsAg-expressing plasmid pRc/CMV-HBs (Aldevron, Fargo, ND). HBsAg levels were determined with a GS HBsAg EIA 3.0 kit (Bio-Rad Laboratories, Inc., Redmond, WA) as described by the manufacturer. Recombinant HBsAg protein, ayw
 20 subtype, also diluted in nonfat milk in PBS, was used as a primary standard (Aldevron).

HBsAg expression for each animal was normalized to the control group of mice injected with PBS in order to account for the non-treatment related decline in expression of MC-HBV1.3. First, the HBsAg level for each animal at a time point was divided by the pre-
 25 treatment level of expression in that animal in order to determine the ratio of expression “normalized to pre-treatment”. Expression at a specific time point was then normalized to the control group by dividing the “normalized to pre-treatment” ratio for an individual animal by the average “normalized to pre-treatment” ratio of all mice in the normal PBS control group.

iii) Serum Hepatitis B e-antigen (HBeAg) levels: HBeAg analysis was performed with
 30 the HBeAg enzyme linked immunosorbent assay (ELISA) as described by the manufacturer (DiaSorin) using serum diluted 4- to 20-fold in 5% nonfat dry milk. The amount of antigen

was determined in the linear range of the assay and quantitated against HBeAg protein standards (Fitzgerald Industries International, catalog # 30-AH18, Acton, MA).

5 HBeAg expression for each animal was normalized to the control group of mice injected with PBS in order to account for the non-treatment related decline in expression of MC-HBV1.3. For evaluation of HBeAg in serum, HBeAg is analyzed from pooled group or subgroup serum samples. First, the HBeAg level for each pooled group or subgroup was divided by the pre-treatment level of expression in the same group or subgroup in order to determine the ratio of expression “normalized to pre-treatment”. Expression at a specific time
10 point was then normalized to the control group by dividing the “normalized to pre-treatment” ratio for a group or subgroup by the average “normalized to pre-treatment” ratio of all samples from the normal PBS control group.

iv) Serum HBV DNA levels: Equal volumes of serum from mice in a group or subgroup
15 were pooled to a final volume of 100 μ L. DNA was isolated from serum samples using the QIAamp MinElute Virus Spin Kit (Qiagen, Valencia, CA) following the manufacturer’s instructions. Sterile 0.9% saline was added to each sample to a final volume of 200 μ L. Serum samples were added to tubes containing buffer and protease. Carrier RNA was added to aid in the isolation of small amounts of DNA. 1 ng of pHCR/UbC-SEAP plasmid DNA (Wooddell
20 CI, et al. "Long-term RNA interference from optimized siRNA expression constructs in adult mice." *Biochem Biophys Res Commun* (2005) 334, 117-127) was added as a recovery control. After incubating 15 min at 56°C, nucleic acids were precipitated from the lysates with ethanol and the entire solution applied to a column. After washing, the samples were eluted into a volume of 50 μ L Buffer AVE.

25

The number of copies of HBV genomes in DNA isolated from the pHBV mouse model serum was determined by qPCR. Plasmid pSEAP-HBV353-777, encoding a short segment of the HBV genome within the S gene (bases 353-777 of GenBank accession #V01460), was used to create a six log standard curve. Samples with recovery of DNA below 2 standard deviations
30 from the average, based on detection of pHCR/UbC-SEAP were omitted. TaqMan chemistry-based primers and probes with fluor/ZEN/IBFQ are utilized.

qPCR assays were performed on a 7500 Fast or StepOne Plus Real-Time PCR system (Life Technologies). For evaluation of HBV DNA in serum, DNA was isolated from singlet or duplicate purification steps from pooled group serum samples. Quantitations of HBV DNA and recovery control plasmid were determined by qPCR reactions performed in triplicate. The probes to quantitate HBV and pHCR/UbC-SEAP were included in each reaction.

Example 3. *HBV RNAi agents in pHBV model mice.*

The pHBV mouse model described in Example 2, above, was used. At day 1, each mouse was administered a single subcutaneous injection of 200 μ l containing 2 mg/kg (mpk) of an HBV RNAi agent formulated in phosphate buffered saline ("PBS"), or 200 μ l of phosphate buffered saline without an HBV RNAi agent, to be used as a control. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The HBV RNAi agents tested included those having the duplex numbers shown in Table 7, below. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

Serum was collected on day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 7. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 3 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
PBS	1.000 \pm 0.185	1.000 \pm 0.288	1.000 \pm 0.540	1.000 \pm 0.326
AD04178	0.164 \pm 0.043	0.206 \pm 0.044	0.293 \pm 0.050	0.348 \pm 0.099
AD04579	0.083 \pm 0.028	0.099 \pm 0.022	0.112 \pm 0.022	0.138 \pm 0.056
AD04580	0.048 \pm 0.007	0.073 \pm 0.012	0.085 \pm 0.012	0.126 \pm 0.014
AD04570	0.241 \pm 0.076	0.294 \pm 0.071	0.276 \pm 0.068	0.474 \pm 0.092
AD04572	0.190 \pm 0.040	0.279 \pm 0.011	0.323 \pm 0.049	0.441 \pm 0.046
AD04573	0.333 \pm 0.143	0.505 \pm 0.106	0.361 \pm 0.060	0.444 \pm 0.068
AD04574	0.291 \pm 0.032	0.650 \pm 0.056	0.388 \pm 0.048	0.485 \pm 0.070
AD04575	0.397 \pm 0.189	0.514 \pm 0.234	0.574 \pm 0.204	0.689 \pm 0.207
AD04419	0.262 \pm 0.038	0.174 \pm 0.042	0.258 \pm 0.064	0.311 \pm 0.089

AD04578	0.210 ± 0.056	0.235 ± 0.033	0.298 ± 0.035	0.336 ± 0.049
---------	-------------------	-------------------	-------------------	-------------------

- RNAi agents AD04178, AD04579, AD04580, AD04570, AD04572, AD04573, AD04574, AD04575, AD04419, and AD04578 were each designed to have antisense strand sequences at least partially complementary to the X open reading frame at positions 1781-1789 of the HBV genome shown in Tables 1 and 2, above. Each of the HBV RNAi agents showed substantial reduction in HBsAg as compared to the PBS control across all measured time points. For example, AD04580 showed greater than 95% reduction in s-antigen levels at day 8 (0.048 ± 0.007 HBsAg level) when normalized to pre-treatment and PBS control.
- Additionally, serum HBV DNA levels were determined for the PBS, AD04579, and AD04580 groups from serum samples collected on days 8, 15, 22, 29, 36, 43 and 50, pursuant to the procedure set forth in Example 2, above. Serum from each group was pooled and then DNA was isolated from the serum in duplicate isolations. Data are presented in the following Table:
- Table 8.** Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 3 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
PBS	1.0000 ± 0.1185	1.0000 ± 0.0591	1.0000 ± 0.0322	1.0000 ± 0.0597
AD04579	0.1541 ± 0.0070	0.1776 ± 0.0027	0.1810 ± 0.0450	0.3738 ± 0.0302
AD04580	0.0921 ± 0.0253	0.0869 ± 0.0117	0.1444 ± 0.0755	0.0950 ± 0.0026
Group	Day 36	Day 43	Day 50	
PBS	1.0000 ± 0.1625	1.0000 ± 0.0055	1.0000 ± 0.1484	
AD04579	0.9670 ± 0.1247	0.7643 ± 0.1334	0.6299 ± 0.1319	
AD04580	0.4949 ± 0.0096	0.4350 ± 0.0344	0.6819 ± 0.0266	

- The data in Table 8 indicate that both RNAi agents examined provided a substantial reduction in HBV DNA levels compared to the PBS group, with AD04580 achieving slightly greater than 1 log knockdown at nadir (e.g., 0.0869 ± 0.0117 average serum DNA level at day 15).

Example 4. HBV RNAi agents in pHBV model mice.

- The pHBV mouse model described in Example 2, above, was used. At day 1, each mouse was given a single subcutaneous administration of 200 μ l containing 2 mg/kg (mpk) of an HBV

RNAi agent formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent to be used as a control. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The HBV RNAi agents administered included those listed in Table 9, below. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

Serum was collected on day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 9. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 4 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
PBS	1.000 \pm 0.085	1.000 \pm 0.235	1.000 \pm 0.171	1.000 \pm 0.099
AD04010	0.229 \pm 0.141	0.165 \pm 0.091	0.142 \pm 0.085	0.116 \pm 0.076
AD04581	0.379 \pm 0.042	0.221 \pm 0.066	0.135 \pm 0.040	0.112 \pm 0.050
AD04591	0.285 \pm 0.101	0.145 \pm 0.064	0.086 \pm 0.024	0.081 \pm 0.026
AD04434	0.295 \pm 0.041	0.191 \pm 0.008	0.147 \pm 0.016	0.187 \pm 0.049
AD04583	0.488 \pm 0.018	0.545 \pm 0.037	0.511 \pm 0.086	0.663 \pm 0.112
AD04584	0.392 \pm 0.136	0.337 \pm 0.073	0.364 \pm 0.075	0.515 \pm 0.155
AD04585	0.099 \pm 0.016	0.042 \pm 0.014	0.030 \pm 0.009	0.044 \pm 0.014
AD04586	0.222 \pm 0.056	0.107 \pm 0.034	0.074 \pm 0.016	0.106 \pm 0.039
AD04588	0.255 \pm 0.065	0.205 \pm 0.021	0.185 \pm 0.021	0.207 \pm 0.024
AD04438	0.265 \pm 0.106	0.113 \pm 0.045	0.091 \pm 0.031	0.130 \pm 0.038

RNAi agents AD04010, AD04581, AD04591, AD04434, AD04583, AD04584, AD04585, AD04586, AD04588, and AD04438 were designed to have antisense strand sequences that are at least partially complementary to the S open reading frame at positions 257-275 of the HBV genome, as shown in Tables 1 and 2. The HBV RNAi agents shown in Table 9, directly above, each showed substantial reduction in HBsAg as compared to the PBS control across all measured time points. For example, AD04585 exhibited approximately a 90% reduction of HBsAg at day 8, a 95% reduction at day 15, a 97% reduction at day 22, and a 95% reduction at day 29.

Additionally, serum HBV DNA levels were determined for the PBS, AD04585 groups from serum samples collected on days 8, 15, 22, 29, 36, 43 and 50, pursuant to the procedure set forth in Example 2, above. Serum from each group was pooled and then DNA was isolated from the serum in duplicate isolations. Data are presented in the following Table:

Table 10. Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 4 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
PBS	1.000 ± 0.248	1.000 ± 0.089	1.000 ± 0.195	1.000 ± 0.180
AD04585	0.901 ± 0.183	0.225 ± 0.003	0.187 ± 0.023	0.191 ± 0.004
Group	Day 36	Day 43	Day 50	
PBS	1.000 ± 0.018	1.000 ± 0.033	1.000 ± 0.778	
AD04585	0.209 ± 0.017	0.171 ± 0.019	0.305 ± 0.010	

The data in Table 10 indicate that HBV RNAi agent AD04585 provided a reduction in HBV DNA levels compared to the PBS group.

Example 5. Dose response and combinations of HBV RNAi Agents in pHBV model mice.

The pHBV mouse model described in Example 2, above, was used. The mice were divided into various groups including those set forth in Table 11, below, and the mice were given 200 µl subcutaneous injections pursuant to the dosing regimen set forth in Table 11:

Table 11. Dosing groups of pHBV mice for Example 5.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS (no RNAi agent)	Single injection on day 1
B	3.0 mg/kg AD04585	Single injection on day 1
C	3.0 mg/kg AD04585	Injection on day 1, day 8, and day 15 (i.e., three weekly injections)
D	3.0 mg/kg AD04580	Single injection on day 1
E	3.0 mg/kg AD04580	Injection on day 1, day 8, and day 15 (i.e., three weekly injections)
F	1.0 mg/kg AD4585 + 1.0 mg/kg AD04580	Injection on day 1, and another injection on day 22
G	1.0 mg/kg AD4585 + 1.0 mg/kg AD04580	Injection on day 1, day 8, day 15, and day 43
H	1.5 mg/kg AD4585 + 1.5 mg/kg AD04580	Injection on day 1, day 22, and day 43
I	1.5 mg/kg AD4585 + 1.5 mg/kg AD04580	Injection on day 1, day 8, day 15, and day 43

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 11. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

Serum was collected on day 8, day 15, day 22, day 29, day 36, day 43, day 50, and day 57, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 12. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 5 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 \pm 0.162	1.000 \pm 0.138	1.000 \pm 0.083	1.000 \pm 0.204
B	0.060 \pm 0.015	0.010 \pm 0.003	0.006 \pm 0.002	0.007 \pm 0.002
C	0.087 \pm 0.014	0.004 \pm 0.001	0.001 \pm 0.0003	0.0002 \pm 0.0001

D	0.026 ± 0.009	0.035 ± 0.013	0.037 ± 0.014	0.046 ± 0.006
E	0.023 ± 0.005	0.002 ± 0.001	0.001 ± 0.0003	0.001 ± 0.0004
F	0.063 ± 0.046	0.083 ± 0.051	0.086 ± 0.016	0.027 ± 0.006
G	0.062 ± 0.011	0.022 ± 0.008	0.009 ± 0.003	0.008 ± 0.002
H	0.055 ± 0.015	0.062 ± 0.002	0.072 ± 0.013	0.011 ± 0.001
I	0.031 ± 0.006	0.008 ± 0.001	0.003 ± 0.0004	0.003 ± 0.0003
Group	Day 36	Day 43	Day 50	Day 57
A	1.000 ± 0.211	1.000 ± 0.189	1.000 ± 0.179	1.000 ± 0.062
B	0.013 ± 0.005	0.027 ± 0.004	0.026 ± 0.004	0.057 ± 0.012
C	0.001 ± 0.0002	0.002 ± 0.001	0.008 ± 0.004	0.020 ± 0.015
D	0.116 ± 0.019	0.214 ± 0.056	0.263 ± 0.046	0.404 ± 0.030
E	0.003 ± 0.0001	0.007 ± 0.001	0.012 ± 0.002	0.033 ± 0.011
F	0.029 ± 0.003	0.065 ± 0.005	0.064 ± 0.004	0.161 ± 0.033
G	0.014 ± 0.008	0.039 ± 0.011	0.018 ± 0.008	0.046 ± 0.008
H	0.017 ± 0.005	0.039 ± 0.008	0.007 ± 0.001	0.013 ± 0.003
I	0.007 ± 0.001	0.020 ± 0.002	0.005 ± 0.001	0.011 ± 0.002

HBV RNAi agents AD04580 and AD04585 each individually showed a reduction in HBsAg as compared to the PBS control across all measured time points. Furthermore, combination treatment of AD04585 and AD04580, which as noted in the Examples above target different regions of the HBV genome, also showed reduction in HBsAg as compared to the PBS control across all measured time points.

Additionally, serum HBV DNA levels were determined for each of the groups in Table 11 from serum samples collected on days 8, 15, 22, 29, and 36, pursuant to the procedure set forth in Example 2, above. Serum from each group was pooled and then DNA was isolated from the serum in duplicate reactions. Data are presented in the following Table:

Table 13. Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 5 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 ± 0.063	1.000 ± 0.059	1.000 ± 0.372	1.000 ± 0.237
B	0.267 ± 0.003	0.043 ± 0.016	0.038 ± 0.008	0.044 ± 0.004
C	0.236 ± 0.016	0.023 ± 0.001	0.004 ± 0.001	0.002 ± 0.000
D	0.058 ± 0.016	0.085 ± 0.017	0.252 ± 0.071	0.217 ± 0.009

E	0.056 ± 0.002	0.0009 ± 0.0004	0.0005 ± 0.0002	0.003 ± 0.002
F	0.298 ± 0.013	0.351 ± 0.032	0.823 ± 0.127	0.217 ± 0.007
G	0.276 ± 0.035	0.112 ± 0.020	0.061 ± 0.002	0.073 ± 0.002
H	0.232 ± 0.012	0.213 ± 0.028	0.403 ± 0.047	0.079 ± 0.005
I	0.092 ± 0.026	0.055 ± 0.000	0.002 ± 0.003	0.010 ± 0.004
Group	Day 36			
A	1.000 ± 0.024			
B	0.046 ± 0.007			
C	0.003 ± 0.000			
D	0.319 ± 0.034			
E	0.002 ± 0.000			
F	0.122 ± 0.004			
G	0.047 ± 0.006			
H	0.056 ± 0.003			
I	0.021 ± 0.007			

The data in Table 13 indicate that the RNAi agents examined, both individually and in combination, provided a reduction in HBV DNA levels compared to the PBS group. Re-dosing or increasing the dose amount yielded additional HBV DNA reductions.

5

Example 6. HBV RNAi agents in pHBV mice: dose response and combination studies.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 14, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 14:

10

Table 14. Dosing groups of pHBV mice for Example 6.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS (no RNAi agent)	Single injection on day 1
B	4.0 mg/kg AD04981	Single injection on day 1
C	1.0 mg/kg AD04981	Single injection on day 1
D	2.0 mg/kg AD04981	Single injection on day 1
E	1.0 mg/kg AD04963	Single injection on day 1
F	2.0 mg/kg AD04963	Single injection on day 1
G	3.0 mg/kg AD04872	Single injection on day 1
H	3.0 mg/kg AD04872 + 1.0 mg/kg AD04981	Single injection on day 1
I	3.0 mg/kg AD04872 + 1.0 mg/kg AD04963	Single injection on day 1

J	3.0 mg/kg AD04872 + 2.0 mg/kg AD04981	Single injection on day 1
---	--	---------------------------

- Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 14. Each of the HBV RNAi agents included
- 5 N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).
- 10 Serum was collected on day -1 prior to administration, and then on day 8, day 15, day 22, day 29, and day 36, and serum HBsAg levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table 15, with Average HBsAg reflecting the normalized average value of HBsAg:
- 15 **Table 15.** Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 6 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
A	1.000 \pm 0.068	1.000 \pm 0.183	1.000 \pm 0.181
B	0.085 \pm 0.020	0.068 \pm 0.005	0.089 \pm 0.014
C	0.283 \pm 0.039	0.343 \pm 0.055	0.436 \pm 0.004
D	0.161 \pm 0.052	0.137 \pm 0.036	0.190 \pm 0.068
E	0.182 \pm 0.040	0.233 \pm 0.023	0.436 \pm 0.029
F	0.078 \pm 0.024	0.093 \pm 0.015	0.167 \pm 0.028
G	0.066 \pm 0.030	0.013 \pm 0.002	0.010 \pm 0.002
H	0.033 \pm 0.012	0.016 \pm 0.005	0.020 \pm 0.005
I	0.040 \pm 0.011	0.028 \pm 0.003	0.032 \pm 0.007
J	0.035 \pm 0.010	0.019 \pm 0.002	0.021 \pm 0.001
Group	Day 29	Day 36	
A	1.000 \pm 0.032	1.000 \pm 0.141	
B	0.148 \pm 0.016	0.194 \pm 0.047	
C	0.622 \pm 0.041	0.741 \pm 0.132	
D	0.234 \pm 0.055	0.280 \pm 0.071	
E	0.623 \pm 0.116	0.782 \pm 0.114	
F	0.259 \pm 0.014	0.368 \pm 0.068	

G	0.010 ± 0.003	0.009 ± 0.004
H	0.022 ± 0.005	0.024 ± 0.009
I	0.065 ± 0.014	0.087 ± 0.015
J	0.031 ± 0.0001	0.044 ± 0.002

The HBV RNAi agents tested showed a reduction in HBsAg as compared to the PBS control across all measured time points. Furthermore, combination treatment of AD04872 (which includes an antisense strand sequence that is at least partially complementary to the S ORF at positions 261-279 of the HBV genome, as shown in Tables 1 and 2) and either AD04981 or AD04963 (both of which include antisense strand sequences that are at least partially complementary to the X ORF at positions 1781-1799 of the HBV genome, as shown in Tables 1 and 2), which are shown in Groups H, I, and J of Example 6, illustrate that combination treatment of two RNAi agents targeting, one which targets in the S ORF, and the other which targets in the X ORF of the HBV genome, similarly showed reduction in HBsAg compared to the PBS control across all measured time points.

Additionally, Serum Hepatitis B e-antigen (HBeAg) levels were also assessed. Samples from the mice in each respective group were first pooled, and the resulting serum samples were assayed in singlet. Data from the experiment is shown in the following Table:

Table 16. Average HBeAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 6.

Group	Day 8	Day 15	Day 22	Day 29	Day 36
A	1.000	1.000	1.000	0.183	1.000
B	0.138	0.180	0.274	0.005	0.089
C	0.316	0.376	0.588	0.055	0.436
D	0.167	0.250	0.262	0.036	0.190
E	0.301	0.327	0.447	0.023	0.436
F	0.167	0.172	0.305	0.015	0.167
G	0.275	0.135	0.158	0.002	0.010
H	0.080	0.053	0.094	0.005	0.020
I	0.165	0.124	0.185	0.003	0.032
J	0.120	0.057	0.101	0.002	0.021

As shown in Table 16, the combination AD04872 (which targets the S ORF of the HBV genome) with either AD04981 or AD04963 (both of which target the X ORF of the HBV

genome), showed a further reduction in HBeAg levels relative to administering AD04872 alone.

Example 7. *HBV RNAi Agents in pHBV mice: additional dose response and combination studies.*

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 17, below, and each mouse was administered a single 200 μ l subcutaneous injection pursuant to the dosing regimen set forth in Table 17:

Table 17. Dosing groups of pHBV mice for Example 7.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS (no RNAi agent)	Single injection on day 1
B	4.0 mg/kg AD04776	Single injection on day 1
C	1.0 mg/kg AD04982	Single injection on day 1
D	2.0 mg/kg AD04982	Single injection on day 1
E	1.0 mg/kg AD04776	Single injection on day 1
F	2.0 mg/kg AD04776	Single injection on day 1
G	3.0 mg/kg AD04872	Single injection on day 1
H	3.0 mg/kg AD04872 + 1.0 mg/kg AD04982	Single injection on day 1
I	3.0 mg/kg AD04872 + 2.0 mg/kg AD04982	Single injection on day 1

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 17. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Four (4) mice in each group were tested on day -1 and day 8 (n=4), and then one mouse per group was euthanized for histological evaluation. Three (3) mice in each group were tested at day 22 and day 29 (n=3).

Serum was collected on day -1 prior to administration, and then on day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table 18:

5

Table 18. Average HBsAg levels normalized to pre-treatment (day -1) and PBS control in pHBV mice following administration of HBV RNAi agents from Example 7 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 ± 0.347	1.000 ± 0.278	1.000 ± 0.194	1.000 ± 0.318
B	0.117 ± 0.069	0.085 ± 0.039	0.148 ± 0.045	0.198 ± 0.049
C	0.519 ± 0.058	0.375 ± 0.012	0.422 ± 0.046	0.525 ± 0.037
D	0.342 ± 0.062	0.255 ± 0.046	0.272 ± 0.122	0.314 ± 0.068
E	0.279 ± 0.057	0.245 ± 0.032	0.374 ± 0.121	0.304 ± 0.035
F	0.224 ± 0.018	0.161 ± 0.009	0.310 ± 0.016	0.482 ± 0.053
G	0.029 ± 0.010	0.005 ± 0.001	0.004 ± 0.001	0.006 ± 0.001
H	0.016 ± 0.005	0.004 ± 0.001	0.010 ± 0.006	0.015 ± 0.008
I	0.026 ± 0.012	0.008 ± 0.001	0.010 ± 0.002	0.015 ± 0.005

10 The HBV RNAi agents tested showed a reduction in HBsAg as compared to the PBS control across all measured time points.

15 Additionally, Serum Hepatitis B e-antigen (HBeAg) levels were also assessed. Samples from the mice in each respective group were first pooled, and the resulting serum samples were assayed in singlet. Data from the experiment is shown in the following Table:

Table 19. Average HBeAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 7.

Group	Day 8	Day 15	Day 22	Day 29	Day 36
A	1.000	1.000	1.000	1.000	1.000
B	0.193	0.213	0.260	0.307	0.464
C	0.471	0.424	0.562	0.513	0.705
D	0.335	0.310	0.411	0.442	0.500
E	0.381	0.368	0.355	0.564	0.483
F	0.275	0.255	0.370	0.495	0.449
G	0.323	0.218	0.205	0.250	0.190
H	0.124	0.102	0.099	0.156	0.156
I	0.081	0.059	0.045	0.063	0.086

Table 19-1. Average HBeAg fold knockdown normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 7.

Group	Day 8 (Fold KD)	Day 15 (Fold KD)	Day 22 (Fold KD)	Day 29 (Fold KD)	Day 36 (Fold KD)
A	1.0	1.0	1.0	1.0	1.0
B	5.2	4.7	3.8	3.3	2.2
C	2.1	2.4	1.8	2.0	1.4
D	3.0	3.2	2.4	2.3	2.0
E	2.6	2.7	2.8	1.8	2.1
F	3.6	3.9	2.7	2.0	2.2
G	3.1	4.6	4.9	4.0	5.3
H	8.1	9.8	10.1	6.4	6.4
I	12.3	17.0	22.3	15.7	11.6

Table 19-1 reflects the fold knockdown ratio of HBeAg compared to control, which is calculated as normalized HBeAg level of the control (PBS) group/normalized HBeAg level of the respected RNAi agent(s) group (i.e., 1.000/HBeAg level). The data in Table 19-1 indicate that the combination of AD04872 (which, as noted above, includes an antisense strand sequence that is at least partially complementary to the S ORF at positions 261-279 of the HBV genome) with AD04982 (which includes an antisense strand sequence that is at least partially complementary to the X ORF at positions 1781-1799 of the HBV genome), showed a further reduction in HBeAg levels relative to administering the individual RNAi agents alone (*See, e.g.,* Tables 19 and 19-1 for Groups H and I). Further, the data from this Example also show that the combination of AD04872 with AD04982 resulted in fold decrease of HBeAg greater than the sum of the fold decrease of HBeAg in AD04872 and AD04982 administered individually. For example, Group I (which is the administration of 3.0 mg/kg AD04872 + 2.0 mg/kg AD04982) resulted in a fold decrease of HBeAg at day 15 of 17.0, which is greater than the sum of the fold decrease for Group G (3.0 mg/kg AD04872) of 4.6 plus the fold decrease for Group D (2.0 mg/kg AD04982) of 3.2.

Further, serum HBV DNA levels were determined for each of the groups in Table 17 from serum samples collected on days -1, 8, 15, 22, 29, and 36, pursuant to the procedure set forth in Example 2, above. Serum HBV DNA was isolated from each animal at each time point. Data are presented in the following Table:

Table 20. Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 7 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 ± 0.493	1.000 ± 0.358	1.000 ± 0.424	1.000 ± 0.387
B	0.224 ± 0.150	0.263 ± 0.185	0.335 ± 0.204	0.449 ± 0.108
C	0.358 ± 0.207	0.428 ± 0.073	0.433 ± 0.220	0.474 ± 0.090
D	0.516 ± 0.163	0.523 ± 0.264	0.244 ± 0.123	0.241 ± 0.085
E	0.601 ± 0.388	0.319 ± 0.125	0.279 ± 0.138	0.506 ± 0.525
F	0.363 ± 0.128	0.374 ± 0.197	0.275 ± 0.146	0.385 ± 0.141
G	0.071 ± 0.032	0.022 ± 0.009	0.015 ± 0.015	0.025 ± 0.005
H	0.069 ± 0.070	0.018 ± 0.014	0.019 ± 0.020	0.022 ± 0.001
I	0.044 ± 0.024	0.033 ± 0.016	0.017 ± 0.012	0.022 ± 0.014
Group	Day 36			
A	1.000 ± 0.326			
B	0.603 ± 0.068			
C	0.509 ± 0.163			
D	0.543 ± 0.079			
E	0.444 ± 0.407			
F	0.721 ± 0.043			
G	0.058 ± 0.030			
H	0.047 ± 0.021			
I	0.058 ± 0.051			

5

The data in Table 20 indicate that the RNAi agents examined, both individually and in combination, provided a reduction in HBV DNA levels compared to the PBS group, and further show that the combination of AD04872 (which targets the S ORF) and AD04982 (which targets the X ORF) reduces serum HBV DNA to a similar degree as an equal amount of AD04872 alone.

10

Example 8. HBV RNAi Agents in pHBV mice: further dose response and combination studies.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 21, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 21:

15

Table 21. Dosing groups of pHBV mice for Example 8.

Group	RNAi Agent and Dose	Dosing Regimen	Number of Animals (n)
1	PBS (no RNAi agent)	Single injection on day 1	4
2A	4.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1	4
2B	4.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1	4
3A	3.3 mg/kg AD04872 + 1.7 mg/kg AD05070	Single injection on day 1	4
3B	3.3 mg/kg AD04872 + 1.7 mg/kg AD05070	Single injection on day 1	4
4A	3.2 mg/kg AD04872 + 0.8 mg/kg AD05070	Single injection on day 1	4
4B	3.2 mg/kg AD04872 + 0.8 mg/kg AD05070	Single injection on day 1	4
5A	2.7 mg/kg AD04872 + 1.3 mg/kg AD05070	Single injection on day 1	4
5B	2.7 mg/kg AD04872 + 1.3 mg/kg AD05070	Single injection on day 1	4
6A	4.0 mg/kg AD05070	Single injection on day 1	4
6B	4.0 mg/kg AD05070	Single injection on day 1	4
7A	1.7 mg/kg AD05070	Single injection on day 1	4
7B	1.7 mg/kg AD05070	Single injection on day 1	4
8A	0.8 mg/kg AD05070	Single injection on day 1	4
8B	0.8 mg/kg AD05070	Single injection on day 1	4
9	1.7 mg/kg AD05148	Single injection on day 1	4
10	2.7 mg/kg AD04872	Single injection on day 1	3
11	1.7 mg/kg AD05147	Single injection on day 1	3
12	4.0 mg/kg AD04872	Single injection on day 1	3
13	1.7 mg/kg AD05149	Single injection on day 1	3

Additionally, the mice are scheduled to be euthanized pursuant to the following schedule:

- Day 11: Euthanize 2 mice from groups 2A, 3A, 4A, 5A, 6A, 7A and 8A, and euthanize one mouse from group 9.
- Day 14: Euthanize 2 mice from groups 2A, 3A, 4A, 5A, 6A, 7A, and 8A.
- Day 21: Euthanize 2 mice from groups 2B, 3B, 4B, 5B, 6B, 7B, and 8B.

- Day 28: Euthanize 2 mice from groups 1, 2B, 3B, 4B, 5B, 6B, 7B, and 8B, and all mice (4) from groups 10 and 12.

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 21. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. As shown in Table 14 above, four (4) mice in each group were tested (n=4), except for groups 10, 11, 12 and 13, in which three mice were tested (n=3).

Serum was collected on day -1 prior to administration, and on days 8, 14, 21 and 28, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 22. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 8 (standard deviation reflected as (+/-)).

Group Number	Day 8	Day 14	Day 21	Day 28
1	1.000 \pm 0.089	1.000 \pm 0.087	1.000 \pm 0.132	1.000 \pm 0.138
2A	0.009 \pm 0.003	0.005 \pm 0.001		
2B	0.006 \pm 0.003	0.002 \pm 0.001	0.004 \pm 0.001	0.005 \pm 0.001
3A	0.032 \pm 0.021	0.009 \pm 0.004		
3B	0.028 \pm 0.027	0.008 \pm 0.006	0.012 \pm 0.005	0.015 \pm 0.005
4A	0.036 \pm 0.020	0.012 \pm 0.006		
4B	0.029 \pm 0.025	0.010 \pm 0.008	0.015 \pm 0.005	0.022 \pm 0.004
5A	0.027 \pm 0.014	0.008 \pm 0.002		
5B	0.027 \pm 0.013	0.007 \pm 0.003	0.019 \pm 0.004	0.031 \pm 0.005
6A	0.058 \pm 0.035	0.069 \pm 0.039		
6B	0.117 \pm 0.058	0.079 \pm 0.047	0.145 \pm 0.082	0.135 \pm 0.061
7A	0.189 \pm 0.100	0.084 \pm 0.029		
7B	0.099 \pm 0.010	0.147 \pm 0.025	0.267 \pm 0.048	0.345 \pm 0.063
8A	0.355 \pm 0.099	0.366 \pm 0.069		
8B	0.271 \pm 0.058	0.334 \pm 0.060	0.464 \pm 0.055	0.624 \pm 0.053
9	0.239 \pm 0.148	0.179 \pm 0.127	0.309 \pm 0.213	0.345 \pm 0.225
10	0.018 \pm 0.009	0.005 \pm 0.003	0.005 \pm 0.002	0.007 \pm 0.003
11	0.129 \pm 0.068	0.138 \pm 0.060	0.239 \pm 0.092	0.315 \pm 0.119

12	0.033 ± 0.022	0.002 ± 0.001	0.002 ± 0.001	0.002 ± 0.0004
13	0.200 ± 0.093	0.239 ± 0.114	0.367 ± 0.123	0.477 ± 0.125

The HBV RNAi agents tested, both alone and in combination, showed a substantial reduction in HBsAg as compared to the PBS control across all measured time points.

5 **Example 9. RNAi agent delivery.**

The pHBV mouse model described in Example 2, above, was used. At day 1, each mouse was administered a single subcutaneous injection of 200 μ l containing 10 mg/kg (mpk) of an HBV RNAi agent formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, to be used as a control. The HBV RNAi agents tested included those having the duplex numbers shown in Table 23, below, which each included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

15

Serum was collected prior to administration, and then on day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

20

Table 23. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 9 (standard deviation reflected as (+/-)).

RNAi agent	HBsAg in serum at nadir (norm. fraction)	%KD at nadir	Day of nadir
PBS	1.000	N/A	N/A
AD03498	0.087 ± 0.016	91.3%	8
AD03499	0.069 ± 0.011	93.1%	15
AD03500	0.095 ± 0.031	90.5%	8
AD03501	0.046 ± 0.020	95.4%	15

Each of the HBV RNAi agents shown in Table 23, above, included an antisense strand sequence that is at least partially complementary to the X ORF at positions 1781-1799 of the HBV genome. Each of the RNAi agents showed a significant knockdown compared to PBS control.

5

Example 10. HBV RNAi Agents in pHBV mice: further combination studies.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 24, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 24:

10

Table 24. Dosing groups of pHBV mice for Example 10.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS Group I (no RNAi agent)	Single injection on day 1 and day 22
B	PBS Group II (no RNAi agent)	Single injection on day 1 and day 22
C	3.0 mg/kg AD04585	Single injection on day 1, day 22, day 50, and day 64
D	3.0 mg/kg AD04771	Single injection on day 1 and day 22
E	3.0 mg/kg AD04580	Single injection on day 1, day 22, day 50, and day 64
F	3.0 mg/kg AD04776	Single injection on day 1 and day 22
G	1.5 mg/kg AD04585 + 1.5 mg/kg AD04580	Single injection on day 1, day 22, day 50, and day 64
H	1.5 mg/kg AD04771 + 1.5 mg/kg AD04776	Single injection on day 1 and day 22
I	2.0 mg/kg AD04771 + 1.0 mg/kg AD04776	Single injection on day 1 and day 22
J	2.25 mg/kg AD04771 + 0.75 mg/kg AD04776	Single injection on day 1 and day 22

Each mouse was given a subcutaneous administration of 200 µl containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 µl of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 24. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

15

Serum was collected prior to administration, and then on day -1, day 8, day 15, day 22, day 29, day 36, day 43, day 50, day 57, and day 64. Serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the

5 experiment is shown in the following:

Table 25. Average HBsAg levels normalized to pre-treatment and PBS control (Group A used as control) in pHBV mice following administration of HBV RNAi agents from Example 10 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
A	1.000 ± 0.146	1.000 ± 0.095	1.000 ± 0.202
B	0.931 ± 0.161	1.091 ± 0.156	1.132 ± 0.259
C	0.071 ± 0.050	0.031 ± 0.022	0.024 ± 0.013
D	0.134 ± 0.035	0.130 ± 0.024	0.119 ± 0.028
E	0.015 ± 0.001	0.041 ± 0.012	0.087 ± 0.015
F	0.197 ± 0.081	0.308 ± 0.138	0.476 ± 0.156
G	0.029 ± 0.015	0.069 ± 0.029	0.094 ± 0.016
H	0.191 ± 0.057	0.315 ± 0.094	0.420 ± 0.126
I	0.153 ± 0.050	0.194 ± 0.076	0.233 ± 0.116
J	0.155 ± 0.059	0.177 ± 0.067	0.316 ± 0.117
Group	Day 29	Day 36	Day 43
A	1.000 ± 0.182	1.000 ± 0.287	1.000 ± 0.298
B	1.417 ± 0.414	1.166 ± 0.248	
C	0.007 ± 0.005	0.004 ± 0.003	0.006 ± 0.001
D	0.048 ± 0.023	0.036 ± 0.020	0.052 ± 0.027
E	0.014 ± 0.006	0.021 ± 0.011	0.026 ± 0.011
F	0.246 ± 0.081	0.244 ± 0.097	0.179 ± 0.061
G	0.023 ± 0.009	0.027 ± 0.009	0.037 ± 0.013
H	0.200 ± 0.080	0.185 ± 0.081	0.194 ± 0.055
I	0.141 ± 0.082	0.133 ± 0.051	0.151 ± 0.082
J	0.133 ± 0.064	0.102 ± 0.039	0.129 ± 0.050
Group	Day 50	Day 57	Day 64
A	1.000 ± 0.296	1.000 ± 0.394	1.000 ± 0.395
B			
C	0.015 ± 0.0001	0.002 ± 0.001	0.004 ± 0.001
D			
E	0.052 ± 0.015	0.009 ± 0.002	0.018 ± 0.007
F			
G	0.076 ± 0.020	0.012 ± 0.003	0.020 ± 0.007
H			
I			

J			
---	--	--	--

HBV RNAi agents AD04585 and AD04771 were designed to have antisense strand sequences that are at least partially complementary to the S open reading frame at positions 257-275 of the HBV genome, as shown in Tables 1 and 2. HBV RNAi agents AD04580 and AD04776 were designed to have antisense strand sequences that are at least partially complementary to the X open reading frame at positions 1781-1799 of the HBV genome, as shown in Tables 1 and 2. The HBV RNAi agents tested, both alone and in combination, showed a reduction in HBsAg as compared to the PBS control across all measured time points. Each subsequent dose further reduced the nadir of HBsAg reduction.

Additionally, serum HBV DNA levels were determined for Group C (3.0 mg/kg AD04585), Group E (3.0 mg/kg AD04580), and Group G (1.5 mg/kg AD04585 + 1.5 mg/kg AD04580) in Table 24, from serum samples collected on days -1, 8, 15, 22, 29, and 36, 43 and 50 pursuant to the procedure set forth in Example 2, above. Serum HBV DNA was isolated for each animal at each of these time points. Data are presented in the following Table:

Table 26. Average Serum HBV DNA levels normalized to pre-treatment and PBS controls (both PBS groups A and B) in pHBV mice following administration of HBV RNAi agents from Example 10 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A/B (PBS)	1.000 ± 0.316	1.000 ± 0.427	1.000 ± 0.428	1.000 ± 0.475
C	0.172 ± 0.151	0.142 ± 0.079	0.252 ± 0.132	0.072 ± 0.086
E	0.024 ± 0.015	0.042 ± 0.037	0.449 ± 0.184	0.053 ± 0.048
G	0.093 ± 0.053	0.083 ± 0.037	0.370 ± 0.153	0.211 ± 0.060
Group	Day 36	Day 43	Day 50	
A/B (PBS)	1.000 ± 0.623	1.000 ± 0.532	1.000 ± 0.532	
C	0.044 ± 0.020	0.104 ± 0.033	0.156 ± 0.016	
E	0.012 ± 0.004	0.061 ± 0.031	0.161 ± 0.019	
G	0.048 ± 0.022	0.147 ± 0.010	0.295 ± 0.041	

The data in Table 26 indicate that the HBV RNAi agents examined, both individually and in combination, provided a reduction in HBV DNA levels compared to the PBS group.

Example 11. HBV RNAi Agents in pHBV mice: combination studies.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 27, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 27:

5

Table 27. Dosing groups of pHBV mice for Example 11.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS (no RNAi agent)	Single injection on day 1
B	3.0 mg/kg AD04962	Single injection on day 1
C	3.0 mg/kg AD04963	Single injection on day 1
D	1.5 mg/kg AD04962 + 1.5 mg/kg AD04963	Single injection on day 1
E	2.0 mg/kg AD04962 + 1.0 mg/kg AD04963	Single injection on day 1
F	2.25 mg/kg AD04962 + 0.75 mg/kg AD04963	Single injection on day 1
G	1.5 mg/kg AD04962 + 1.5 mg/kg AD04963	Single injection on day 1
H	3.0 mg/kg AD04962 + 3.0 mg/kg AD04963	Single injection on day 1
I	1.5 mg/kg AD04962 + 1.5 mg/kg AD04963	Single injection on day 1
J	4.5 mg/kg AD04962 + 4.5 mg/kg AD04963	Single injection on day 1
K	3.0 mg/kg AD04872	Single injection on day 1
L	3.0 mg/kg AD04882	Single injection on day 1
M	3.0 mg/kg AD04885	Single injection on day 1

Each mouse was given a subcutaneous administration of 200 µl containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 µl of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 24. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

15

Serum was collected on day -1 prior to administration, and then on day 8, day 15, day 22, day 29, and day 36 (except for Group L (AD04882) and Group M (AD04885), and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

5

Table 28. Average HBsAg normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 11 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
A	1.000 ± 0.048	1.000 ± 0.144	1.000 ± 0.083
B	0.125 ± 0.025	0.083 ± 0.014	0.063 ± 0.016
C	0.019 ± 0.005	0.035 ± 0.008	0.052 ± 0.009
D	0.054 ± 0.013	0.079 ± 0.009	0.108 ± 0.021
E	0.099 ± 0.025	0.098 ± 0.053	0.142 ± 0.050
F	0.070 ± 0.015	0.103 ± 0.036	0.140 ± 0.020
G	0.041 ± 0.021	0.012 ± 0.008	0.021 ± 0.013
H	0.020 ± 0.006	0.044 ± 0.010	0.062 ± 0.019
I	0.077 ± 0.017	0.019 ± 0.004	0.004 ± 0.001
J	0.012 ± 0.002	0.021 ± 0.001	0.032 ± 0.002
K	0.045 ± 0.014	0.013 ± 0.005	0.008 ± 0.005
L	0.106 ± 0.020	0.176 ± 0.044	0.215 ± 0.082
M	0.275 ± 0.029	0.378 ± 0.080	0.572 ± 0.043
Group	Day 29	Day 36	
A	1.000 ± 0.209	1.000 ± 0.270	
B	0.079 ± 0.020	0.096 ± 0.007	
C	0.087 ± 0.014	0.164 ± 0.026	
D	0.176 ± 0.014	0.292 ± 0.030	
E	0.223 ± 0.082	0.373 ± 0.150	
F	0.213 ± 0.020	0.328 ± 0.034	
G	0.031 ± 0.013	0.078 ± 0.064	
H	0.97 ± 0.028	0.160 ± 0.060	
I	0.008 ± 0.001	0.002 ± 0.0003	
J	0.044 ± 0.008	0.069 ± 0.009	
K	0.011 ± 0.007	0.011 ± 0.009	
L	0.299 ± 0.009		
M	0.792 ± 0.057		

10 RNAi agent AD04962 was designed to have an antisense strand sequence that is at least partially complementary to the S open reading frame at positions 257-275 of the HBV genome, as shown in Tables 1 and 2. RNAi agent AD04872 was designed to have an antisense strand

sequence that is at least partially complementary to the S open reading frame at positions 261-279 of the HBV genome, as shown in Tables 1 and 2. RNAi agent AD04963 was designed to have an antisense strand sequence that is at least partially complementary to the X open reading frame at positions 1781-1799 of the HBV genome, as shown in Tables 1 and 2. RNAi agents
 5 AD04882 and AD04885 were designed to have antisense strand sequences that are at least partially complementary to the X open reading frame at positions 1780-1798 of the HBV genome, as shown in Tables 1 and 2. The HBV RNAi agents shown in Table 9, directly above, each showed a reduction in HBsAg as compared to the PBS control across all measured timepoints, both individually and in combination. Re-dosing yielded additional HBsAg
 10 reduction.

Additionally, Serum Hepatitis B e-antigen (HBeAg) levels were also assessed for all groups except Groups L and M. Samples from the mice in each respective group were first pooled, and the resulting serum samples were assayed in singlet. Data from the experiment is shown
 15 in the following Table:

Table 29. Average HBeAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 11.

Group	Day 8	Day 22	Day 29	Day 36
A	1.000	1.000	1.000	1.000
B	0.425	0.291	0.371	0.365
C	0.152	0.170	0.328	0.356
D	0.266	0.249	0.456	0.440
E	0.278	0.295	0.589	0.561
F	0.306	0.291	0.718	0.522
G	0.183	0.138	0.291	0.249
H	0.091	0.131	0.315	0.238
I	0.183	0.052	0.069	0.036
J	0.089	0.114	0.190	0.236
K	0.458	0.172	0.322	0.207

20 Further, serum HBV DNA levels were determined for each of the groups in Table 27 from serum samples collected on days 8, 15, 22, and 29, pursuant to the procedure set forth in Example 2, above. Serum HBV DNA was isolated from each animal at each time point. Data are presented in the following Table:

Table 30. Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 7 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 ± 0.232	1.000 ± 0.463	1.000 ± 0.272	1.000 ± 0.205
B	0.577 ± 0.219	0.222 ± 0.064	0.196 ± 0.055	0.261 ± 0.117
C	0.165 ± 0.051	0.070 ± 0.042	0.142 ± 0.105	0.228 ± 0.174
D	0.343 ± 0.125	0.307 ± 0.091	0.300 ± 0.092	0.356 ± 0.032
E	0.262 ± 0.033	0.216 ± 0.018	0.227 ± 0.028	0.279 ± 0.090
F	0.320 ± 0.134	0.332 ± 0.208	0.344 ± 0.209	0.338 ± 0.211
G	0.231 ± 0.036	0.034 ± 0.024	0.069 ± 0.039	0.077 ± 0.020
H	0.229 ± 0.101	0.155 ± 0.121	0.148 ± 0.079	0.215 ± 0.035
I	0.281 ± 0.129	0.109 ± 0.071	0.023 ± 0.019	0.011 ± 0.009
J	0.078 ± 0.050	0.061 ± 0.020	0.074 ± 0.029	0.056 ± 0.030
K	0.314 ± 0.064	0.119 ± 0.043	0.076 ± 0.067	0.078 ± 0.095
L	0.295 ± 0.077	0.305 ± 0.101	0.213 ± 0.088	0.186 ± 0.084
M	0.515 ± 0.247	0.505 ± 0.293	0.488 ± 0.318	0.478 ± 0.267

- 5 The data in Table 30 indicate that the RNAi agents examined, both individually and in combination, provided a reduction in HBV DNA levels compared to the PBS group. Re-dosing yielded addition reduction of HBV DNA.

Example 12. HBV RNAi Agents in pHBV mice.

- 10 The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 31, below, and each mouse was administered a single 200 μ l subcutaneous injection pursuant to the dosing regimen set forth in Table 31:

Table 31. Dosing groups of pHBV mice for Example 12.

Group	RNAi Agent and Dose	Dosing Regimen
A	PBS (no RNAi agent)	Single injection on day 1
B	2.0 mg/kg AD04871	Single injection on day 1
C	2.0 mg/kg AD04872	Single injection on day 1
D	2.0 mg/kg AD04874	Single injection on day 1
E	2.0 mg/kg AD04875	Single injection on day 1
F	2.0 mg/kg AD04876	Single injection on day 1
G	2.0 mg/kg AD04881	Single injection on day 1
H	2.0 mg/kg AD04883	Single injection on day 1

I	2.0 mg/kg AD04884	Single injection on day 1
---	-------------------	---------------------------

- Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 24. Each of the HBV RNAi agents included
- 5 N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).
- 10 Serum was collected prior to administration, and then on day 8, day 15, and day 22. Group A (PBS), Group B (2.0 mg/kg AD04871), Group C (2.0 mg/kg AD04872), Group D (2.0 mg/kg AD04874), Group E (2.0 mg/kg AD04875), and Group F (2.0 mg/kg AD04876) also had serum collected on day 29, day 36, day 43, and day 50. Serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from
- 15 the experiment is shown in the following Table:

Table 32. Average HBsAg normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 12 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
A	1.000 \pm 0.132	1.000 \pm 0.089	1.000 \pm 0.080	1.000 \pm 0.098
B	0.102 \pm 0.034	0.041 \pm 0.021	0.049 \pm 0.033	0.048 \pm 0.031
C	0.153 \pm 0.064	0.064 \pm 0.032	0.063 \pm 0.034	0.042 \pm 0.017
D	0.123 \pm 0.022	0.049 \pm 0.017	0.039 \pm 0.010	0.023 \pm 0.001
E	0.190 \pm 0.075	0.094 \pm 0.038	0.107 \pm 0.061	0.081 \pm 0.051
F	0.190 \pm 0.031	0.076 \pm 0.035	0.084 \pm 0.038	0.049 \pm 0.024
G	0.159 \pm 0.047	0.216 \pm 0.057	0.235 \pm 0.151	
H	0.508 \pm 0.078	0.666 \pm 0.131	0.543 \pm 0.048	
I	0.279 \pm 0.087	0.357 \pm 0.078	0.614 \pm 0.156	
Group	Day 36	Day 43	Day 50	
A	1.000 \pm 0.065	1.000 \pm 0.242	1.000 \pm 0.224	
B	0.054 \pm 0.038	0.064 \pm 0.030	0.092 \pm 0.025	
C	0.049 \pm 0.017	0.054 \pm 0.015	0.085 \pm 0.010	
D	0.037 \pm 0.004	0.037 \pm 0.010	0.065 \pm 0.012	
E	0.126 \pm 0.077	0.125 \pm 0.063	0.170 \pm 0.079	

F	0.089 ± 0.044	0.082 ± 0.034	0.115 ± 0.028
G			
H			
I			

HBV RNAi agents AD04871, AD04872, AD04874, AD04875, and AD04876 were each designed to have antisense strand sequences that are at least partially complementary to the S open reading frame at positions 261-279 of the HBV genome, as shown in Tables 1 and 2.

Each of these HBV RNAi agents should a substantial reduction in HBsAg compared to PBS control. For example, a single 2 mg/kg dose of each of AD04871 (Group B), AD04872 (Group C) and AD04874 (Group D), and AD04876 (Group F), exhibited a greater than 90% reduction in HBsAg for each of the timepoints measured from day 15 through day 43 compared to control. HBV RNAi agents AD04881, AD04883, AD04884 were each designed to have antisense strand sequences that are at least partially complementary to the X open reading frame at positions 1780-1798 of the HBV genome, as shown in Tables 1 and 2.

Example 13. Dose response and combinations of HBV RNAi Agents in X Region Knockout model mice.

As an alternative means in assessing the effects of the combination of an RNAi agent that includes an antisense strand sequence that is at least partially complementary to a region located in the S ORF of an HBV mRNA, and a second RNAi agent that includes an antisense strand sequence that is at least partially complementary to a region located in the X ORF of an HBV mRNA, a plasmid was generated that included the HBV genome with a knockout of the binding site for HBV RNAi agents that target positions 1780 and 1781, as shown in Tables 1 and 2 (hereinafter referred to as X Region Knockout mice). This model was generated by mutating ten (10) bases in the pHBV1.3 plasmid within the binding site of these RNAi agents. The remainder of the HBV mRNA, including the S-region, remained functional. Thus, in this HBV mouse model, inclusion of an HBV RNAi agent having an antisense strand that targets positions 1780 and 1781 of the HBV genome disclosed herein is expected to be ineffective in silencing expression.

The mice were divided into various groups including those set forth in Table 33, below, and the mice were given 200 µl subcutaneous injections pursuant to the dosing regimen set forth in the following Table:

Table 33. Dosing groups of X Region Knockout mice for Example 13.

Group	RNAi Agent and Dose	Dosing Regimen	Number of Animals (n)
1	PBS (no RNAi agent)	Single injection on day 1	4
2	2.0 mg/kg AD04585 + 1.0 mg/kg AD04963	Single injection on day 1	4
3	2.0 mg/kg AD04872 + 1.0 mg/kg AD04963	Single injection on day 1	4
4	2.5 mg/kg AD04585 + 0.5 mg/kg AD04963	Single injection on day 1	4
5	2.5 mg/kg AD04872 + 0.5 mg/kg AD04963	Single injection on day 1	4
6	3.0 mg/kg AD04963	Single injection on day 15	1

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 33. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

Serum was collected on day 5, day 8, day 15, day 22, and day 29 and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Serum was also collected for Groups 1 through 5 on days 36 and 43. Data from the experiment is shown in the following Table 34:

Table 34. Average HBsAg normalized to pre-treatment and PBS control in X Region Knockout mice following administration of HBV RNAi agents from Example 13 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
1	1.000 \pm 0.186	1.000 \pm 0.165	1.000 \pm 0.132
2	0.061 \pm 0.034	0.041 \pm 0.035	0.030 \pm 0.015
3	0.020 \pm 0.011	0.007 \pm 0.003	0.003 \pm 0.002
4	0.063 \pm 0.039	0.022 \pm 0.011	0.029 \pm 0.013

5	0.027 ± 0.014	0.003 ± 0.003	0.001 ± 0.001
6	0.948	1.360	1.652
	Day 29	Day 36	Day 43
1	1.000 ± 0.059	1.000 ± 0.044	1.000 ± 0.045
2	0.051 ± 0.029	0.062 ± 0.029	
3	0.004 ± 0.003	0.008 ± 0.003	0.018 ± 0.007
4	0.040 ± 0.022	0.061 ± 0.030	
5	0.002 ± 0.001	0.003 ± 0.002	0.014 ± 0.006
6	1.831		

As expected, Group 6, which was a single dose of 3.0 mg/kg of HBV RNAi agent AD04963 and includes an antisense strand that is at least partially complementary to the X open reading frame at positions 1781-1799 of the HBV genome, was unable to provide knockdown of HBsAg. Additionally, each of Groups 2 through 5 provided substantial knockdown of HBsAg compared to PBS control, with both Group 3 and Group 5 exhibiting a greater than 2 log reduction in HBsAg at nadir (day 22).

Example 14. Dose response and combinations of HBV RNAi Agents in X Region Knockout model mice.

The X Region Knockout mouse model described in Example 13, above, was used. Mice were divided into various groups including those set forth in Table 31, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 35:

Table 35. Dosing groups of X Region Knockout mice for Example 14.

Group	RNAi Agent and Dose	Dosing Regimen
1	PBS (no RNAi agent)	Single injection on day 1
2	2.0 mg/kg AD04872	Single injection on day 1
3	2.0 mg/kg AD04872 + 0.7 mg/kg AD05070	Single injection on day 1
4	2.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1
5	2.0 mg/kg AD04872 + 2.0 mg/kg AD05070	Single injection on day 1

Each mouse was given a subcutaneous administration of 200 µl containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 µl of phosphate buffered saline

without an HBV RNAi agent, as set forth in Table 35. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group shown in Table 35 were tested (n=3).

Serum was collected on day 1 (pre-dose), day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 36. Average HBsAg levels normalized to pre-treatment and PBS control in X Region Knockout mice from Example 14.

Group	Day 8	Day 15	Day 22	Day 29
1	1.000 ± 0.120	1.000 ± 0.255	1.000 ± 0.224	1.000 ± 0.143
2	0.104 ± 0.104	0.009 ± 0.009	0.005 ± 0.004	0.005 ± 0.003
3	0.076 ± 0.041	0.010 ± 0.009	0.006 ± 0.005	0.005 ± 0.005
4	0.036 ± 0.008	0.002 ± 0.001	0.001 ± 0.001	0.002 ± 0.001
5	0.019 ± 0.017	0.003 ± 0.002	0.003 ± 0.001	0.004 ± 0.000

Table 36 shows that HBV RNAi agent AD04872 administered alone, and the combination of AD04872 (which includes an antisense strand that is at least partially complementary to the S open reading frame at positions 261-279 of the HBV genome) and AD05070 (which includes an antisense strand that is at least partially complementary to the X open reading frame at positions 1781-1799 of the HBV genome), provided significant knockdown of HBsAg compared to PBS control across each of the time points measured. Addition of 0.7 mg/kg to 2 mg/kg HBV RNAi agent AD05070 for which there was a mutated target site in this X Region Knockout model did not diminish the activity of the 2 mg/kg HBV RNAi agent AD04872.

Additionally, serum HBV DNA levels were determined from serum samples collected on days 8, 15, and 22 pursuant to the procedure set forth in Example 2, above. Serum from each group was pooled and then DNA was isolated from the serum in singlet. Data are presented in the following Table:

Table 37. Average Serum HBV DNA levels normalized to pre-treatment and PBS controls in X Region Knockout mice following administration of HBV RNAi agents from Example 14 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
1	1.000 ± 0.007	1.000 ± 0.011	1.000 ± 0.066
2	0.225 ± 0.019	0.022 ± 0.001	0.036 ± 0.001
3	0.151 ± 0.002	0.029 ± 0.001	0.042 ± 0.003
4	0.140 ± 0.006	0.016 ± 0.000	0.018 ± 0.000
5	0.069 ± 0.002	0.018 ± 0.003	0.043 ± 0.002

- 5 Addition of 0.7 mg/kg to 2 mg/kg HBV RNAi agent AD05070 for which there was a mutated target site in this X Region Knockout model did not diminish the activity of the 2 mg/kg HBV RNAi agent AD04872.

Example 15. HBV RNAi agents in pHBV mice.

- 10 The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups including those set forth in Table 38, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 38:

Table 38. Dosing groups of pHBV mice for Example 15.

Group	RNAi Agent and Dose	Dosing Regimen
1	PBS (no RNAi agent)	Single injection on day 1
2	2.0 mg/kg AD04776	Single injection on day 1
3	2.0 mg/kg AD05069	Single injection on day 1
4	2.0 mg/kg AD05070	Single injection on day 1
5	2.0 mg/kg AD05071	Single injection on day 1
6	2.0 mg/kg AD05073	Single injection on day 1
7	2.0 mg/kg AD05074	Single injection on day 1
8	2.0 mg/kg AD05075	Single injection on day 1
9	2.0 mg/kg AD05076	Single injection on day 1
10	2.0 mg/kg AD05077	Single injection on day 1
11	2.0 mg/kg AD05078	Single injection on day 1
12	3.0 mg/kg AD04872 + 1.0 mg/kg AD04776	Single injection on day 1

13	3.0 mg/kg AD04872 + 1.0 mg/kg AD05069	Single injection on day 1
14	3.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1
15	3.0 mg/kg AD04872 + 1.0 mg/kg AD05071	Single injection on day 1
16	3.0 mg/kg AD04872 + 1.0 mg/kg AD05073	Single injection on day 1
17	3.0 mg/kg AD04872 + 1.0 mg/kg AD05074	Single injection on day 1
18	3.0 mg/kg AD04872 + 1.0 mg/kg AD05075	Single injection on day 1
19	3.0 mg/kg AD04872 + 1.0 mg/kg AD05076	Single injection on day 1
20	3.0 mg/kg AD04872 + 1.0 mg/kg AD05077	Single injection on day 1
21	3.0 mg/kg AD04872 + 1.0 mg/kg AD05078	Single injection on day 1

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 38. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).

Serum was collected on day -1 prior to administration, and then on day 8, day 15, day 22, day 29, day 36, day 43, and day 50. Serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table 39, with Average HBsAg reflecting the normalized average value of HBsAg:

Table 39. Average HBsAg normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 15.

Group	Day 8	Day 15	Day 22	Day 29
1	1.000 \pm 0.119	1.000 \pm 0.047	1.000 \pm 0.080	1.000 \pm 0.027
2	0.339 \pm 0.076	0.414 \pm 0.126	0.385 \pm 0.067	0.450 \pm 0.075
3	0.240 \pm 0.096	0.361 \pm 0.078	0.446 \pm 0.073	0.508 \pm 0.114

4	0.081 ± 0.026	0.127 ± 0.031	0.223 ± 0.057	0.330 ± 0.112
5	0.452 ± 0.020	0.431 ± 0.126	0.373 ± 0.079	0.383 ± 0.080
6	0.375 ± 0.181	0.632 ± 0.192	0.463 ± 0.117	0.567 ± 0.159
7	0.325 ± 0.032	0.438 ± 0.125	0.393 ± 0.056	0.443 ± 0.096
8	0.155 ± 0.031	0.322 ± 0.019	0.333 ± 0.077	0.463 ± 0.043
9	0.245 ± 0.063	0.467 ± 0.090	0.477 ± 0.045	0.562 ± 0.049
10	0.120 ± 0.062	0.173 ± 0.029	0.289 ± 0.019	0.331 ± 0.042
11	0.128 ± 0.042	0.172 ± 0.046	0.179 ± 0.015	0.215 ± 0.049
12	0.040 ± 0.015	0.014 ± 0.004	0.014 ± 0.006	0.015 ± 0.004
13	0.050 ± 0.020	0.015 ± 0.011	0.017 ± 0.008	0.022 ± 0.009
14	0.020 ± 0.011	0.011 ± 0.006	0.015 ± 0.006	0.023 ± 0.004
15	0.043 ± 0.005	0.013 ± 0.005	0.010 ± 0.002	0.011 ± 0.004
16	0.021 ± 0.017	0.008 ± 0.004	0.012 ± 0.003	0.011 ± 0.001
17	0.032 ± 0.011	0.009 ± 0.003	0.007 ± 0.002	0.008 ± 0.0003
18	0.023 ± 0.014	0.010 ± 0.006	0.009 ± 0.006	0.009 ± 0.004
19	0.025 ± 0.006	0.010 ± 0.004	0.009 ± 0.002	0.010 ± 0.003
20	0.061 ± 0.013	0.027 ± 0.006	0.020 ± 0.003	0.029 ± 0.006
21	0.061 ± 0.050	0.013 ± 0.010	0.012 ± 0.005	0.018 ± 0.006
Group	Day 36	Day 43	Day 50	
1	1.000 ± 0.031	1.000 ± 0.114	1.000 ± 0.112	
2	0.617 ± 0.116	0.643 ± 0.154	0.665 ± 0.199	
3	0.638 ± 0.067	0.743 ± 0.015	0.792 ± 0.115	
4	0.472 ± 0.121	0.515 ± 0.126	0.689 ± 0.167	
5	0.591 ± 0.159	0.604 ± 0.086	0.709 ± 0.115	
6	0.717 ± 0.136	0.686 ± 0.194	0.781 ± 0.301	
7	0.586 ± 0.069	0.775 ± 0.143	0.747 ± 0.095	
8	0.666 ± 0.066	0.803 ± 0.096	0.856 ± 0.180	
9	0.801 ± 0.047	0.667 ± 0.055	0.765 ± 0.208	
10	0.640 ± 0.059	0.667 ± 0.034	0.742 ± 0.133	
11	0.429 ± 0.063	0.383 ± 0.005	0.497 ± 0.060	
12	0.037 ± 0.013	0.044 ± 0.012	0.056 ± 0.014	
13	0.046 ± 0.011	0.055 ± 0.010	0.070 ± 0.010	
14	0.054 ± 0.016	0.070 ± 0.018	0.096 ± 0.012	
15	0.029 ± 0.011	0.032 ± 0.015	0.051 ± 0.020	
16	0.033 ± 0.005	0.038 ± 0.007	0.062 ± 0.004	
17	0.021 ± 0.002	0.031 ± 0.004	0.061 ± 0.005	
18	0.034 ± 0.014	0.047 ± 0.016	0.079 ± 0.017	
19	0.028 ± 0.005	0.037 ± 0.006	0.060 ± 0.011	
20	0.070 ± 0.009	0.063 ± 0.018	0.097 ± 0.018	
21	0.040 ± 0.012	0.066 ± 0.007	0.120 ± 0.036	

RNAi agents AD04776, AD05069, AD05070, AD05071, AD05073, and AD05074 were each designed to have an antisense strand sequence that is at least partially complementary to the X open reading frame at positions 1781-1799 of the HBV genome, as shown in Tables 1 and 2.

RNAi agents AD05075, AD05076, AD05077, and AD05078 were each designed to have antisense strand sequences that are at least partially complementary to the X open reading frame at positions 1780-1798 of the HBV genome, as shown in Tables 1 and 2.

- 5 Table 39 shows that HBV RNAi agents AD04776, AD05069, AD05070, AD05071, AD05073, and AD05074 administered alone or their combination with AD04872 (which includes an antisense strand that is at least partially complementary to the S open reading from at positions 261-279 of the HBV genome) provided significant knockdown of HBsAg compared to PBS control across each of the time points measured.

10

Example 16. HBV RNAi Agents in pHBV mice: dose response and combination studies.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 40, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 40:

15

Table 40. Dosing groups of pHBV mice for Example 16.

Group	RNAi Agent and Dose	Dosing Regimen
1	PBS (no RNAi agent)	Single injection on day 1
2	3.2 mg/kg AD04872	Single injection on day 1
3	3.2 mg/kg AD04872	Single injection on day 1 and day 22
4	3.0 mg/kg AD04872 + 0.8 mg/kg AD05070	Single injection on day 1
5	3.0 mg/kg AD04872 + 0.8 mg/kg AD05070	Single injection on day 1 and day 22
6	3.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1
7	3.0 mg/kg AD04872 + 1.0 mg/kg AD05070	Single injection on day 1 and day 22
8	2.7 mg/kg AD04872 + 1.3 mg/kg AD05070	Single injection on day 1
9	2.7 mg/kg AD04872 + 1.3 mg/kg AD05070	Single injection on day 1 and day 22
10	2.0 mg/kg AD04872 + 2.0 mg/kg AD04776	Single injection on day 1 and day 22
11	0.8 mg/kg AD05070	Single injection on day 1 and day 22
12	1.3 mg/kg AD05070	Single injection on day 1 and day 22

Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 40. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Six (6) mice in each group were tested (n=6).

Serum was collected prior to administration, and then on day 8, day 15, day 22, and day 29, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table 41:

Table 41. Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 16 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
1	1.000 \pm 0.117	1.000 \pm 0.213	1.000 \pm 0.169	1.000 \pm 0.130
2	0.050 \pm 0.018	0.015 \pm 0.007	0.011 \pm 0.005	0.009 \pm 0.006
3	0.051 \pm 0.037	0.014 \pm 0.011	0.010 \pm 0.006	0.002 \pm 0.001
4	0.029 \pm 0.018	0.010 \pm 0.006	0.011 \pm 0.006	0.010 \pm 0.005
5	0.022 \pm 0.003	0.007 \pm 0.001	0.009 \pm 0.003	0.001 \pm 0.001
6	0.027 \pm 0.012	0.007 \pm 0.004	0.008 \pm 0.005	0.011 \pm 0.005
7	0.028 \pm 0.012	0.010 \pm 0.005	0.009 \pm 0.005	0.001 \pm 0.000
8	0.033 \pm 0.016	0.016 \pm 0.008	0.020 \pm 0.009	0.021 \pm 0.011
9	0.034 \pm 0.025	0.015 \pm 0.011	0.018 \pm 0.013	0.003 \pm 0.002
10	0.038 \pm 0.021	0.015 \pm 0.005	0.019 \pm 0.004	0.003 \pm 0.001
11	0.446 \pm 0.143	0.376 \pm 0.120	0.474 \pm 0.149	0.338 \pm 0.123
12	0.307 \pm 0.111	0.257 \pm 0.122	0.236 \pm 0.057	0.138 \pm 0.031

The HBV RNAi agents tested, both individually and in combination, showed a reduction in HBsAg as compared to the PBS control across all measured time points. HBsAg expression was further reduced in all groups that were re-dosed on day 22.

Additionally, Serum Hepatitis B e-antigen (HBeAg) levels were also assessed. For the day 8 measurement, the serum samples for all six mice in each group were pooled, and the resulting samples were assayed in singlet. For the day -1, day 15, day 22, and day 29 measurements,

the six mice from each group were paired within each group and their respective serum samples were pooled, forming three subgroups for each group. The serum samples for each of the three subgroups for each group were then assayed. Data from the experiment is shown in the following Table 42:

5

Table 42. Average HBeAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 16 (standard deviation for days 15, 22, and 29 reflected as (+/-)).

Group	Day 8	Day 15	Day 22	Day 29
1	1.000	1.000 ± 0.011	1.000 ± 0.170	1.000 ± 0.173
2	0.510	0.308 ± 0.031	0.217 ± 0.021	0.226 ± 0.035
3	0.488	0.301 ± 0.065	0.283 ± 0.081	0.147 ± 0.030
4	0.213	0.216 ± 0.067	0.192 ± 0.029	0.141 ± 0.048
5	0.192	0.211 ± 0.053	0.216 ± 0.088	0.047 ± 0.016
6	0.176	0.163 ± 0.022	0.238 ± 0.069	0.117 ± 0.011
7	0.165	0.175 ± 0.046	0.215 ± 0.061	0.028 ± 0.012
8	0.128	0.166 ± 0.065	0.386 ± 0.284	0.167 ± 0.118
9	0.172	0.171 ± 0.037	0.244 ± 0.052	0.032 ± 0.010
10	0.180	0.211 ± 0.012	0.283 ± 0.034	0.034 ± 0.001
11	0.634	0.594 ± 0.082	0.840 ± 0.152	0.271 ± 0.029
12	0.486	0.441 ± 0.066	0.804 ± 0.096	0.214 ± 0.039

10 The HBV RNAi agents tested, both individually and in combination, showed a reduction in HBeAg as compared to the saline control across all measured time points. HBeAg expression was further reduced in all groups that were re-dosed on day 22.

Further, serum HBV DNA levels were determined for each of the groups in Table 40 from serum samples collected on days -1, 8, 15, and 22, pursuant to the procedure set forth in Example 2, above. Serum from each pair of mice was pooled and then DNA was isolated from each serum pool in a single isolation. Data are presented in the following Table:

20 **Table 43.** Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 16 (standard deviation reflected as (+/-)).

Group	Day 8	Day 15	Day 22
1	1.000 ± 0.122	1.000 ± 0.299	1.000 ± 0.241

2	0.312 ±0.016	0.126 ± 0.008	0.087 ± 0.018
3	0.264 ± 0.065	0.081 ± 0.023	0.073 ± 0.028
4	0.321 ± 0.254	0.120 ± 0.066	0.134 ± 0.101
5	0.319 ± 0.081	0.108 ± 0.038	0.098 ± 0.051
6	0.260 ± 0.095	0.068 ± 0.010	0.076 ± 0.031
7	0.170 ± 0.028	0.082 ± 0.013	0.062 ± 0.018
8	0.188 ± 0.020	0.192 ± 0.160	0.307 ± 0.309
9	0.242 ± 0.003	0.100 ± 0.042	0.075 ± 0.028
10	0.322 ± 0.028	0.159 ± 0.025	0.086 ± 0.016
11	1.124 ± 0.142	0.742 ± 0.127	0.807 ± 0.192
12	1.004 ± 0.144	0.541 ± 0.340	0.569 ± 0.060

The HBV RNAi agents tested, both individually and in combination, showed a reduction in serum HBV DNA as compared to the saline control across all measured time points except in groups 11 and 12 that had no reduction in serum HBV DNA at Day 8.

5

Example 17. HBV RNAi Agents in in pHBV mice.

The pHBV mouse model described in Example 2, above, was used. Mice were divided into various groups as set forth in Table 44, below, and each mouse was administered a single 200 µl subcutaneous injection pursuant to the dosing regimen set forth in Table 44:

10

Table 44. Dosing groups of pHBV mice for Example 17.

Group	RNAi Agent and Dose	Dosing Regimen
1	PBS (no RNAi agent)	Single injection on day 1
2	5 mg/kg AD04585 + 1 mg/kg AD04963	Single injection on day 1
3	5 mg/kg AD04872 + 1 mg/kg AD04963	Single injection on day 1
4	5 mg/kg AD04585 + 1 mg/kg AD04963	Single injection on day 1 and day 8
5	5 mg/kg AD04872 + 1 mg/kg AD04963	Single injection on day 1 and day 8
6	2.5 mg/kg AD04585 + 0.5 mg/kg AD04963	Single injection on day 1
7	2.0 mg/kg AD04585 + 1.0 mg/kg AD04963	Single injection on day 1
8	2.5 mg/kg AD04872 + 0.5 mg/kg AD04963	Single injection on day 1
9	2.0 mg/kg AD04872 + 1.0 mg/kg AD04963	Single injection on day 1
10	5 mg/kg AD04872 + 1 mg/kg AD04981	Single injection on day 1

11	2.5 mg/kg AD04872 + 0.5 mg/kg AD04981	Single injection on day 1 and day 8
12	2.5 mg/kg AD04872 + 0.5 mg/kg AD04981	Single injection on day 1
13	2 mg/kg AD04872 + 1 mg/kg AD04981	Single injection on day 1
14	2.5 mg/kg AD04585 + 0.5 mg/kg AD04981	Single injection on day 1
15	2 mg/kg AD04585 + 1 mg/kg AD04981	Single injection on day 1
16	0.5 mg/kg AD04981	Single injection on day 1

- Each mouse was given a subcutaneous administration of 200 μ l containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or 200 μ l of phosphate buffered saline without an HBV RNAi agent, as set forth in Table 44. Each of the HBV RNAi agents included
- 5 N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area. Three (3) mice in each group were tested (n=3).
- 10 Serum was collected prior to administration, and then on day 8, day 14, day 21, and day 29 and day 36, and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table 45:
- 15 **Table 45.** Average HBsAg levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 17 (standard deviation reflected as (+/-)).

Group	Day 8	Day 14	Day 21	Day 29
1	1.000 \pm 0.068	1.000 \pm 0.125	1.000 \pm 0.152	1.000 \pm 0.110
2	0.058 \pm 0.033	0.059 \pm 0.022	0.085 \pm 0.023	0.158 \pm 0.021
3	0.025 \pm 0.009	0.014 \pm 0.006	0.015 \pm 0.008	0.026 \pm 0.015
4	0.032 \pm 0.007	0.005 \pm 0.001	0.006 \pm 0.002	0.014 \pm 0.002
5	0.024 \pm 0.009	0.003 \pm 0.001	0.001 \pm 0.0004	0.001 \pm 0.0005
6	0.063 \pm 0.020	0.077 \pm 0.013	0.131 \pm 0.011	0.214 \pm 0.026
7	0.041 \pm 0.018	0.059 \pm 0.017	0.091 \pm 0.016	0.140 \pm 0.045
8	0.070 \pm 0.008	0.046 \pm 0.016	0.043 \pm 0.009	0.055 \pm 0.012
9	0.043 \pm 0.006	0.027 \pm 0.003	0.064 \pm 0.017	0.064 \pm 0.014
10	0.015 \pm 0.008	0.005 \pm 0.003	0.005 \pm 0.003	0.005 \pm 0.003

11	0.047 ± 0.014	0.005 ± 0.003	0.003 ± 0.002	0.003 ± 0.003
12	0.062 ± 0.006	0.025 ± 0.007	0.027 ± 0.005	0.033 ± 0.005
13	0.092 ± 0.029	0.050 ± 0.021	0.050 ± 0.022	0.054 ± 0.0019
14	0.310 ± 0.180	0.056 ± 0.010	0.081 ± 0.010	0.112 ± 0.0018
15	0.304 ± 0.044	0.083 ± 0.021	0.115 ± 0.013	0.165 ± 0.025
16	1.667 ± 0.217	0.416 ± 0.163	0.341 ± 0.179	0.511 ± 0.0011
Group	Day 36			
1	1.000 ± 0.225			
2				
3	0.049 ± 0.019			
4				
5	0.004 ± 0.0004			
6				
7				
8	0.081 ± 0.010			
9	0.108 ± 0.026			
10	0.009 ± 0.004			
11	0.005 ± 0.003			
12	0.060 ± 0.014			
13	0.094 ± 0.027			
14				
15				
16	0.634 ± 0.005			

The HBV RNAi agent combinations tested showed a reduction in HBsAg as compared to the saline control across all measured time points. Combinations containing AD04872 showed greater reductions than the equivalent combinations with AD04585 in place of AD04872.

5

Additionally, serum HBV DNA levels were determined for serum samples collected on days 8, 14, 21, and 29 pursuant to the procedure set forth in Example 2, above. Serum HBV DNA was isolated from each animal at each time point. Data are presented in the following Table 46:

10

Table 46. Average Serum HBV DNA levels normalized to pre-treatment and PBS control in pHBV mice following administration of HBV RNAi agents from Example 17 (standard deviation reflected as (+/-)).

Group	Day 8	Day 14	Day 21	Day 29
1	1.000 ± 0.280	1.000 ± 0.269	1.000 ± 0.418	1.000 ± 0.383
2	0.136 ± 0.068	0.192 ± 0.071	0.173 ± 0.032	0.292 ± 0.039
3	0.097 ± 0.034	0.068 ± 0.016	0.076 ± 0.034	0.131 ± 0.061

4	0.061 ± 0.039	0.002 ± 0.001	0.003 ± 0.001	0.019 ± 0.013
5	0.068 ± 0.025	0.003 ± 0.002	0.0009 ± 0.0003	0.0009 ± 0.0003
6	0.354 ± 0.299	0.345 ± 0.187	0.522 ± 0.234	0.509 ± 0.106
7	0.103 ± 0.064	0.291 ± 0.025	0.203 ± 0.043	0.203 ± 0.015
8	0.336 ± 0.142	0.185 ± 0.071	0.183 ± 0.065	0.162 ± 0.064
9	0.198 ± 0.055	0.093 ± 0.023	0.118 ± 0.054	0.143 ± 0.032
10	0.122 ± 0.071	0.024 ± 0.026	0.023 ± 0.020	0.014 ± 0.017
11	0.160 ± 0.069	0.016 ± 0.023	0.003 ± 0.001	0.005 ± 0.004
12	0.158 ± 0.039	0.120 ± 0.044	0.100 ± 0.049	0.091 ± 0.034
13	0.190 ± 0.038	0.169 ± 0.025	0.066 ± 0.015	0.081 ± 0.015
14	0.434 ± 0.136	0.318 ± 0.104	0.144 ± 0.094	0.240 ± 0.029
15	0.358 ± 0.185	0.287 ± 0.108	0.279 ± 0.080	0.303 ± 0.038
16	0.713 ± 0.085	0.674 ± 0.140	0.496 ± 0.128	0.590 ± 0.093

The HBV RNAi agent combinations tested showed a reduction in serum HBV DNA as compared to the saline control across all measured time points. Combinations containing AD04872 showed greater reductions than the equivalent combinations with AD04585 in place of AD04872. These greater reductions were observed at Day 22 and Day 29.

Example 18. HBV RNAi Agents in a HBV-infected Humanized Mouse Model.

For this study, Male FRG® (genotype Fah $^{-/-}$ Rag2 $^{-/-}$ Il2rg $^{-/-}$ triple knockout mice on a C57BL/6 background (Yecuris) were transplanted with human hepatocytes when they were 1-2 months old. The human hepatocytes were allowed to repopulate the liver for approximately 6 months with periodic NTBC treatment to discourage growth of mouse hepatocytes. At 9 months of age the mice were given an intravenous inoculation of 4×10^8 genomes/kg HBV genotype C, which infected the human hepatocytes. After 2-3 months, serum HBV DNA levels reached a plateau indicating the human hepatocytes were maximally infected (mouse hepatocytes cannot be infected by HBV). Mice were one year old at the start of treatment with HBV RNAi agents, thus nearing the end of their life span.

Pre-treatment serum samples were taken on day -10 and day -3. Beginning on day 1, each mouse was administered an oral daily gavage with 0.01 mg/kg Entecavir dissolved in water to inhibit HBV replication. Daily dosing of Entecavir continued until the day mice were euthanized. Entecavir administration was expected to reduce serum HBV DNA in chronically infected human patients, but not reduce HBsAg.

Mice were divided into various groups including those set forth in Table 47, below:

Table 47. Dosing groups of HBV-infected FRG humanized model mice for Example 18.

Group	RNAi Agent and Dose	Dosing Regimen	Terminal Day
A- mouse 1	PBS (no RNAi agent)	Single injection on day 1	Euthanized day 21 (unhealthy animal)
A- mouse 2	PBS (no RNAi agent)	Single injection on day 1 and day 29	Euthanized day 36
B- mouse 1	4.0 mg/kg AD04872 + 2.0 mg/kg AD05070	Single injection on day 1 and day 29	Euthanized day 36
B- mouse 2	4.0 mg/kg AD04872 + 2.0 mg/kg AD05070	Single injection on day 1 and day 29	Euthanized day 40
C- mouse 1	4.5 mg/kg AD04872 + 1.5 mg/kg AD05070	Single injection on day 1	Euthanized day 15
C- mouse 2	4.5 mg/kg AD04872 + 1.5 mg/kg AD05070	Single injection on day 1 and day 29	Euthanized day 36
C- mouse 3	4.5 mg/kg AD04872 + 1.5 mg/kg AD05070	Single injection on day 1 and day 29	Euthanized day 40

Each mouse was also given a subcutaneous administration of 100 μ l per 20 grams body weight containing the amount of HBV RNAi agent(s) formulated in phosphate buffered saline, or an equal volume of phosphate buffered saline without an HBV RNAi agent, on day 1 and on day 29 (if still alive on day 29), pursuant to the schedule as set forth in Table 47, directly above. Each of the HBV RNAi agents included N-acetyl-galactosamine targeting ligands conjugated to the 5'-terminal end of the sense strand, as shown in Tables 4 and 5. The injections were performed between the skin and muscle (i.e. subcutaneous injections) into the loose skin over the neck and shoulder area.

Serum was collected on day 8, day 15, day 22, day 29, day 36, and day 40 and serum Hepatitis B surface antigen (HBsAg) levels were determined pursuant to the procedure set forth in Example 2, above. Data from the experiment is shown in the following Table:

Table 48. Average HBsAg levels normalized to pre-treatment (day -3) for each individual HBV-infected humanized FRG model mouse from Example 18.

Group	Day 8	Day 15	Day 22	Day 29	Day 36	Day 40
A-1	0.830	0.828	0.932	0.858	1.107	
A-2	1.303	1.328				
B-1	0.548	0.314	0.272	0.207	0.138	
B-2	0.592	0.337	0.243	0.215	0.160	0.175

C-1	0.643	0.460	0.415	0.251	0.164	
C-2	0.353	0.228	0.182	0.172	0.224	0.216
C-3	0.814	0.674				

Additionally, serum HBV DNA levels were determined from serum samples collected on days -10, -3, 8, 15, 22, 29, 36, and 40, pursuant to the procedure set forth in Example 2, above. Data are presented in the following Table 49:

5

Table 49. Serum HBV DNA levels normalized to the average of pre-treatment day -10 and day -3 for each HBV-infected FRG humanized mouse following administration of HBV RNAi agents from Example 14.

Group	Day -10	Day -3	Day 8	Day 15	Day 22	Day 29	Day 36	Day 40
A-1	0.883	1.117	0.072	0.038	0.015	0.027	0.060	
A-2	1.070	0.930	0.130	0.075				
B-1	1.538	0.462	0.032	0.017	0.011	0.006	0.010	
B-2	1.350	0.650	0.042	0.018	0.012	0.007	0.008	0.007
C-1	1.348	0.652	0.041	0.020	0.016	0.005	0.004	
C-2	1.030	0.970	0.031	0.015	0.006	0.011	0.008	0.008

10 As expected, administration of Entecavir reduced viral replication in both the absence and presence of HBV RNAi agents.

OTHER EMBODIMENTS

15 It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

20

Claims:

1. An RNAi agent for inhibiting expression of a Hepatitis B Virus gene comprising an antisense strand and a sense strand, wherein the antisense strand comprises a sequence
5 having at least about 85% identity to any of the antisense strand sequences provided in Table 2 or Table 3.
2. The RNAi agent of claim 1, wherein the antisense strand comprises at least 16 contiguous nucleotides of any of the antisense sequences provided in Table 2 or Table 3.
3. The composition of claim 1, wherein the antisense strand comprises the nucleotides at
10 positions 2 through 18 (5'→3') of any of the antisense strand sequences provided in Table 2 or Table 3.
4. The RNAi agent of claim 1 or claim 2, wherein at least one nucleotide of the sense strand and/or at least one nucleotide of the antisense strand of the RNAi agent is a modified nucleotide and/or has a modified internucleoside linkage.
- 15 5. The RNAi agent of any of claims 1-4, wherein the sense strand and the antisense strand are each 17 to 30 nucleotides in length.
6. The RNAi agent of any of claims 1-5, wherein the sense strand comprises at least 16 contiguous nucleotides of any of the sense strand sequences provided in Table 2 or Table 4, and wherein the sense strand has a region of at least about 85% complementarity over
20 the contiguous 16 nucleotides to the antisense strand.
7. The RNAi agent of any of claims 1-6, wherein all or substantially all of the nucleotides in both the sense strand and the antisense strand of the RNAi agent are modified nucleotides.
8. The RNAi agent of claim 1, wherein the antisense strand comprises the nucleotide
25 sequence of any of the modified sequences provided in Table 3.
9. The RNAi agent of claim 1 or claim 8, wherein the sense strand comprises the nucleotide sequence of any of the modified sense sequences provided in Table 4.
10. A composition for inhibiting the expression of a HBV gene comprising an RNAi agent of any of claims 1-9, wherein the RNAi agent is conjugated to a targeting ligand.
- 30 11. The composition of claim 10, wherein the targeting ligand comprises N-acetyl-galactosamine.
12. The composition of claim 11, wherein the targeting ligand is (NAG13), (NAG13)s, (NAG18), (NAG18)s, (NAG24), (NAG24)s, (NAG25), (NAG25)s, (NAG26), (NAG26)s, (NAG27), (NAG27)s, (NAG28), (NAG28)s, (NAG29), (NAG29)s,

- (NAG30), (NAG30)s, (NAG31), (NAG31)s, (NAG32), (NAG32)s, (NAG33), (NAG33)s, (NAG34), (NAG34)s, (NAG35), (NAG35)s, (NAG36), (NAG36)s, (NAG37), (NAG37)s, (NAG38), (NAG38)s, (NAG39), or (NAG39)s.
13. The composition of any of claims 10-12, wherein the targeting ligand is conjugated to the sense strand of the RNAi agent.
14. The composition of any of claims 10-13, wherein the composition further comprises a pharmaceutically acceptable excipient.
15. The composition of any of claims 10-14, wherein the RNAi agent has the duplex structure of AD03498; AD03499; AD03500; AD03501; AD03738; AD03739; AD03967; AD03968; AD03969; AD03970; AD03971; AD03972; AD03973; AD03974; AD03975; AD03976; AD03977; AD03978; AD04001; AD04002; AD04003; AD04004; AD04005; AD04006; AD04007; AD04008; AD04009; AD04010; AD04176; AD04177; AD04178; AD04412; AD04413; AD04414; AD04415; AD04416; AD04417; AD04418; AD04419; AD04420; AD04421; AD04422; AD04423; AD04425; AD04426; AD04427; AD04428; AD04429; AD04430; AD04431; AD04432; AD04433; AD04434; AD04435; AD04436; AD04437; AD04438; AD04439; AD04440; AD04441; AD04442; AD04511; AD04570; AD04571; AD04572; AD04573; AD04574; AD04575; AD04576; AD04577; AD04578; AD04579; AD04580; AD04581; AD04583; AD04584; AD04585; AD04586; AD04587; AD04588; AD04590; AD04591; AD04592; AD04593; AD04594; AD04595; AD04596; AD04597; AD04598; AD04599; AD04734; AD04771; AD04772; AD04773; AD04774; AD04775; AD04776; AD04777; AD04778; AD04822; AD04823; AD04871; AD04872; AD04873; AD04874; AD04875; AD04876; AD04881; AD04882; AD04883; AD04884; AD04885; AD04962; AD04963; AD04981; AD04982; AD04983; AD05069; AD05070; AD05071; AD05072; AD05073; AD05074; AD05075; AD05076; AD05077; AD05078; AD05147; AD05148; AD05149; AD05164; or AD05165.
16. The composition of claim 15, wherein the RNAi agent has the duplex structure of AD04580; AD04585; AD04776; AD04872; AD04962; AD04963; AD04982; or AD05070.
17. The composition of any of claims 10-16, wherein the composition further comprises a second RNAi agent comprising a sense strand and an antisense strand for inhibiting expression of a Hepatitis B Virus gene.

18. The composition of claim 17, wherein the second RNAi agent comprises an antisense sequence having at least about 85% identity to any of the antisense strand sequences provided in Table 2 or Table 3.
19. The composition of claim 18, wherein the antisense strand of the second RNAi agent comprises at least 16 contiguous nucleotides of any of the antisense sequences provided in Table 2 or Table 3.
20. The composition of claim 19, wherein the antisense strand of the second RNAi agent comprises the nucleotides at positions 2 through 18 (5' → 3') of any of the antisense strand sequences provided in Table 2 or Table 3.
21. The composition of any of claims 17-20, wherein the first RNAi agent targets the S open reading frame (ORF) of an HBV gene, and the second RNAi agent targets the X open reading frame (ORF) of an HBV gene.
22. The composition of any of claims 17-20, wherein the first RNAi agent targets the X ORF of an HBV gene, and the second RNAi agent targets the S ORF of an HBV gene.
23. The composition of any of claims 17-21, wherein the first RNAi agent is selected from the group consisting of: AD04001; AD04002; AD04003; AD04004; AD04005; AD04006; AD04007; AD04008; AD04009; AD04010; AD04422; AD04423; AD04425; AD04426; AD04427; AD04428; AD04429; AD04430; AD04431; AD04432; AD04433; AD04434; AD04435; AD04436; AD04437; AD04438; AD04439; AD04440; AD04441; AD04442; AD04511; AD04581; AD04583; AD04584; AD04585; AD04586; AD04587; AD04588; AD04590; AD04591; AD04592; AD04593; AD04594; AD04595; AD04596; AD04597; AD04598; AD04599; AD04734; AD04771; AD04772; AD04773; AD04774; AD04775; AD04822; AD04871; AD04872; AD04873; AD04874; AD04875; AD04876; AD04962; and AD05164; and the second RNAi agent is selected from the group consisting of: AD03498; AD03499; AD03500; AD03501; AD03738; AD03739; AD03967; AD03968; AD03969; AD03970; AD03971; AD03972; AD03973; AD03974; AD03975; AD03976; AD03977; AD03978; AD04176; AD04177; AD04178; AD04412; AD04413; AD04414; AD04415; AD04416; AD04417; AD04418; AD04419; AD04420; AD04421; AD04570; AD04571; AD04572; AD04573; AD04574; AD04575; AD04576; AD04577; AD04578; AD04579; AD04580; AD04776; AD04777; AD04778; AD04823; AD04881; AD04882; AD04883; AD04884; AD04885; AD04963; AD04981; AD04982; AD04983; AD05069; AD05070; AD05071; AD05072; AD05073; AD05074; AD05075; AD05076; AD05077; AD05078; AD05147; AD05148; AD05149; and AD05165.

24. The composition of any of claims 17-21 or 23, wherein the first RNAi agent is AD04872 and the second RNAi agent is AD05070.
25. The composition of any of claims 17-21 or 23, wherein the first RNAi agent is AD04872 and the second RNAi agent is AD04982.
- 5 26. The composition of any of claims 17-21 or 23, wherein the first RNAi agent is AD04872 and the second RNAi agent is AD04776.
27. The composition of any of claims 17-21, wherein the first RNAi agent is AD04585 and the second RNAi agent is AD04580.
28. The composition of any of claims 10-27, further comprising one or more additional
10 therapeutics.
29. The composition of claim 28, wherein the additional therapeutic is lamivudine, tenofovir, tenofovir alafenamide, tenofovir disoproxil, or entecavir.
30. The composition of claim 28, wherein the additional therapeutic is interferon.
31. A method for inhibiting expression of a Hepatitis B Virus gene in a subject, comprising
15 administering to the subject the RNAi agent of any of claims 1-9.
32. A method for inhibiting expression of a Hepatitis B Virus gene in a subject, comprising administering to the subject the composition of any of claims 10-30.
33. The method of claim 31 or claim 32, wherein the subject has or is at risk of having an HBV infection.
- 20 34. A method for treating HBV infection and/or a disease, disorder, or condition associated with HBV infection in a subject, comprising administering to a subject an effective amount of the RNAi agent of any of claims 1-9.
35. A method for treating HBV infection and/or a disease, disorder, or condition associated with HBV infection in a subject, comprising administering to a subject an effective
25 amount of the composition of any of claims 10-30.
36. The method of claim 32, 33, or 35, wherein the composition comprises an effective amount of two RNAi agents for inhibiting the expression of a Hepatitis B Virus gene.
37. The method of any of any of claims 32, 33, 35, or 36, wherein the effective amount of the composition is sufficient to reduce the level of HBsAg in a subject by at least about
30 40% relative to the subject's HBsAg expression level prior to administration of the composition.
38. The method of any of any of claims 32, 33, 35, or 36, wherein the effective amount of the composition is sufficient to reduce the level of HBeAg in a subject by at least about

- 40% relative to the subject's HBeAg expression level prior to administration of the composition.
39. The method of any of any of claims 32, 33, 35, or 36, wherein the effective amount of the composition is sufficient to reduce the serum HBV DNA level in a subject by at least about 40% relative to the subject's serum HBV DNA level prior to administration of the composition.
40. The method of claims 34 or 35, wherein the disease, disorder, or condition associated with HBV infection is a chronic liver disease or disorder, liver inflammation, fibrotic condition, a proliferative disorder, hepatocellular carcinoma, Hepatitis D virus infection, or acute HBV infection.
41. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the first RNAi agent and the second RNAi agent are administered in a ratio of about 1:1.
42. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the first RNAi agent and the second RNAi agent are administered in a ratio of about 2:1.
43. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the first RNAi agent and the second RNAi agent are administered in a ratio of about 3:1.
44. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the first RNAi agent and the second RNAi agent are administered in a ratio of about 4:1.
45. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the first RNAi agent and the second RNAi agent are administered in a ratio of about 5:1.
46. The method of any of claims 32, 33, 35, 36, 37, 38, 39, or 40, wherein the effective amount of the first RNAi agent is between about 0.5 mg/kg and about 5 mg/kg, and the effective amount of the second RNAi agent is between about 0.5 mg/kg and about 5 mg/kg.
47. The method of claim 31, wherein the effective amount of the RNAi agent is between about 0.5 mg/kg and about 5 mg/kg.
48. Use of the RNAi agent of any of claims 1-9 for the treatment of HBV infection.
49. Use of the composition of any of claims 10-30 for the treatment of HBV infection.
50. Use of the composition of any of claims 10-30 for the manufacture of a medicament for treatment of HBV infection.
51. An RNAi agent for inhibiting expression of a Hepatitis B Virus gene comprising an antisense strand and a sense strand, wherein the RNAi agent has a duplex structure comprising a sense strand or an antisense strand that is identical or substantially identical

- to the respective sense or antisense strand in the duplex structure from the group consisting of: AD03498; AD03499; AD03500; AD03501; AD03738; AD03739; AD03967; AD03968; AD03969; AD03970; AD03971; AD03972; AD03973; AD03974; AD03975; AD03976; AD03977; AD03978; AD04001; AD04002; AD04003; AD04004; AD04005; AD04006; AD04007; AD04008; AD04009; AD04010; AD04176; AD04177; AD04178; AD04412; AD04413; AD04414; AD04415; AD04416; AD04417; AD04418; AD04419; AD04420; AD04421; AD04422; AD04423; AD04425; AD04426; AD04427; AD04428; AD04429; AD04430; AD04431; AD04432; AD04433; AD04434; AD04435; AD04436; AD04437; AD04438; AD04439; AD04440; AD04441; AD04442; AD04511; AD04570; AD04571; AD04572; AD04573; AD04574; AD04575; AD04576; AD04577; AD04578; AD04579; AD04580; AD04581; AD04583; AD04584; AD04585; AD04586; AD04587; AD04588; AD04590; AD04591; AD04592; AD04593; AD04594; AD04595; AD04596; AD04597; AD04598; AD04599; AD04734; AD04771; AD04772; AD04773; AD04774; AD04775; AD04776; AD04777; AD04778; AD04822; AD04823; AD04871; AD04872; AD04873; AD04874; AD04875; AD04876; AD04881; AD04882; AD04883; AD04884; AD04885; AD04962; AD04963; AD04981; AD04982; AD04983; AD05069; AD05070; AD05071; AD05072; AD05073; AD05074; AD05075; AD05076; AD05077; AD05078; AD05147; AD05148; AD05149; AD05164; or AD05165.
52. The RNAi agent of claim 51, wherein the RNAi agent has a duplex structure comprising a sense strand that is identical or substantially identical to the respective sense in the duplex structure from the group consisting of: AD03498; AD03499; AD03500; AD03501; AD03738; AD03739; AD03967; AD03968; AD03969; AD03970; AD03971; AD03972; AD03973; AD03974; AD03975; AD03976; AD03977; AD03978; AD04001; AD04002; AD04003; AD04004; AD04005; AD04006; AD04007; AD04008; AD04009; AD04010; AD04176; AD04177; AD04178; AD04412; AD04413; AD04414; AD04415; AD04416; AD04417; AD04418; AD04419; AD04420; AD04421; AD04422; AD04423; AD04425; AD04426; AD04427; AD04428; AD04429; AD04430; AD04431; AD04432; AD04433; AD04434; AD04435; AD04436; AD04437; AD04438; AD04439; AD04440; AD04441; AD04442; AD04511; AD04570; AD04571; AD04572; AD04573; AD04574; AD04575; AD04576; AD04577; AD04578; AD04579; AD04580; AD04581; AD04583; AD04584; AD04585; AD04586; AD04587; AD04588; AD04590; AD04591; AD04592; AD04593; AD04594; AD04595; AD04596; AD04597; AD04598; AD04599; AD04734; AD04771; AD04772; AD04773; AD04774; AD04775; AD04776; AD04777; AD04778;

- AD04822; AD04823; AD04871; AD04872; AD04873; AD04874; AD04875; AD04876;
AD04881; AD04882; AD04883; AD04884; AD04885; AD04962; AD04963; AD04981;
AD04982; AD04983; AD05069; AD05070; AD05071; AD05072; AD05073; AD05074;
AD05075; AD05076; AD05077; AD05078; AD05147; AD05148; AD05149; AD05164;
5 or AD05165.
53. The RNAi agent of claim 52, wherein the RNAi agent has a duplex structure comprising
an antisense strand that is identical or substantially identical to the respective antisense
strand in the duplex structure from the group consisting of: AD03498; AD03499;
AD03500; AD03501; AD03738; AD03739; AD03967; AD03968; AD03969; AD03970;
10 AD03971; AD03972; AD03973; AD03974; AD03975; AD03976; AD03977; AD03978;
AD04001; AD04002; AD04003; AD04004; AD04005; AD04006; AD04007; AD04008;
AD04009; AD04010; AD04176; AD04177; AD04178; AD04412; AD04413; AD04414;
AD04415; AD04416; AD04417; AD04418; AD04419; AD04420; AD04421; AD04422;
AD04423; AD04425; AD04426; AD04427; AD04428; AD04429; AD04430; AD04431;
15 AD04432; AD04433; AD04434; AD04435; AD04436; AD04437; AD04438; AD04439;
AD04440; AD04441; AD04442; AD04511; AD04570; AD04571; AD04572; AD04573;
AD04574; AD04575; AD04576; AD04577; AD04578; AD04579; AD04580; AD04581;
AD04583; AD04584; AD04585; AD04586; AD04587; AD04588; AD04590; AD04591;
AD04592; AD04593; AD04594; AD04595; AD04596; AD04597; AD04598; AD04599;
20 AD04734; AD04771; AD04772; AD04773; AD04774; AD04775; AD04776; AD04777;
AD04778; AD04822; AD04823; AD04871; AD04872; AD04873; AD04874; AD04875;
AD04876; AD04881; AD04882; AD04883; AD04884; AD04885; AD04962; AD04963;
AD04981; AD04982; AD04983; AD05069; AD05070; AD05071; AD05072; AD05073;
AD05074; AD05075; AD05076; AD05077; AD05078; AD05147; AD05148; AD05149;
25 AD05164; or AD05165.
54. The RNAi agent of claim 53, wherein the RNAi agent has a duplex structure comprising
a sense strand and an antisense strand that are both identical or substantially identical to
the respective sense and antisense strand in the duplex structure from the group consisting
of: AD03498; AD03499; AD03500; AD03501; AD03738; AD03739; AD03967;
30 AD03968; AD03969; AD03970; AD03971; AD03972; AD03973; AD03974; AD03975;
AD03976; AD03977; AD03978; AD04001; AD04002; AD04003; AD04004; AD04005;
AD04006; AD04007; AD04008; AD04009; AD04010; AD04176; AD04177; AD04178;
AD04412; AD04413; AD04414; AD04415; AD04416; AD04417; AD04418; AD04419;

- AD04420; AD04421; AD04422; AD04423; AD04425; AD04426; AD04427; AD04428;
 AD04429; AD04430; AD04431; AD04432; AD04433; AD04434; AD04435; AD04436;
 AD04437; AD04438; AD04439; AD04440; AD04441; AD04442; AD04511; AD04570;
 AD04571; AD04572; AD04573; AD04574; AD04575; AD04576; AD04577; AD04578;
 5 AD04579; AD04580; AD04581; AD04583; AD04584; AD04585; AD04586; AD04587;
 AD04588; AD04590; AD04591; AD04592; AD04593; AD04594; AD04595; AD04596;
 AD04597; AD04598; AD04599; AD04734; AD04771; AD04772; AD04773; AD04774;
 AD04775; AD04776; AD04777; AD04778; AD04822; AD04823; AD04871; AD04872;
 AD04873; AD04874; AD04875; AD04876; AD04881; AD04882; AD04883; AD04884;
 10 AD04885; AD04962; AD04963; AD04981; AD04982; AD04983; AD05069; AD05070;
 AD05071; AD05072; AD05073; AD05074; AD05075; AD05076; AD05077; AD05078;
 AD05147; AD05148; AD05149; AD05164; or AD05165
55. The RNAi agent of claims 51-54, wherein the substantial identity is exactly, at least, or
 about 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%,
 15 or 99%.
56. The RNAi agent of any of claims 51-54, wherein the substantial identity is at least, or
 about 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%,
 or 99%.
57. The RNAi agent of any of claims 51-54, wherein the substantial identity is about 85%,
 20 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%.
58. The RNAi agent of any of claims 51-54, wherein the substantial identity is at least about
 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%.
59. A composition comprising the RNAi agent of any of claims 51-58.
60. A composition comprising at least two RNAi agents of any of claims 51-58.
- 25 61. The RNAi agent of any of claims 1-9 or 51-58, wherein the RNAi agent comprises at
 least one overhang.
62. The RNAi agent of any of claim 61, wherein the RNAi agent comprises an overhang at
 the 3' end of the antisense strand.
63. The RNAi agent of claim 62, wherein the RNAi agent comprises an overhang at the 3'
 30 end of the antisense strand and the 3' end of the sense strand.
64. The RNAi agent of any of claims 1-9 or 51-58, wherein the RNAi agent comprises one
 or two blunt ends.

65. The RNAi agent of any of claims 1-9 or 51-58, wherein the RNAi agent comprises one or two frayed ends.
66. The RNAi agent of any of claims 1-9, 51-58, or 61-65, wherein the sense strand comprises at least one inverted abasic nucleoside.

5