

(12) STANDARD PATENT

(11) Application No. AU 2008279509 B2

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Stabilized immune modulatory RNA (SIMRA) compounds

(51) International Patent Classification(s)
C07H 21/02 (2006.01) **A61K 48/00** (2006.01)

(21) Application No: **2008279509** (22) Date of Filing: **2008.07.07**

(87) WIPO No: **WO09/014887**

(30) Priority Data

| | | |
|-------------------|-------------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/981,161 | 2007.10.19 | US |
| 60/957,195 | 2007.08.22 | US |
| 60/948,529 | 2007.07.09 | US |
| 61/015,284 | 2007.12.20 | US |

(43) Publication Date: **2009.01.29**

(44) Accepted Journal Date: **2011.07.21**

(71) Applicant(s)
Idera Pharmaceuticals, Inc.

(72) Inventor(s)
Agrawal, Sudhir;Kandimalla, Ekambar R.;Wang, Daqing;Bhagat, Lakshmi;Lan, Tao

(74) Agent / Attorney
Freehills Patent & Trade Mark Attorneys, Level 43 101 Collins Street, Melbourne, VIC, 3000

(56) Related Art
Lan et al. (2007) Proc. Natl. Acad. Sci. USA 104(34): 13750-13755
WO 2007/117686 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 January 2009 (29.01.2009)

PCT

(10) International Publication Number
WO 2009/014887 A2

(51) International Patent Classification:
C07H 21/02 (2006.01) *A61K 39/00* (2006.01)
A61K 31/7105 (2006.01) *A61K 39/395* (2006.01)

AGRAWAL, Sudhir [—/US]; 61 Lamplighter Drive, Shrewsbury, MA 01545 (US).

(21) International Application Number:
PCT/US2008/069335

(74) Agent: **ZUCCHERO, Joseph, C.**; KLEOWN & ZUCCHERO, LLP, 500 West Cummings Park, Suite 1200, Woburn, MA 01801 (US).

(22) International Filing Date: 7 July 2008 (07.07.2008)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English

English

(26) Publication Language:

English

(30) Priority Data:

| | | |
|------------|-------------------------------|----|
| 60/948,529 | 9 July 2007 (09.07.2007) | US |
| 60/957,195 | 22 August 2007 (22.08.2007) | US |
| 60/981,161 | 19 October 2007 (19.10.2007) | US |
| 61/015,284 | 20 December 2007 (20.12.2007) | US |

(71) **Applicant (for all designated States except US): IDERA PHARMACEUTICALS, INC. [US/US]; 167 Sidney Street, Cambridge, MA 02139 (US).**

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TI, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) **Inventors; and**
(75) **Inventors/Applicants (for US only): KANDIMALLA, Ekambar, R.** [US/US]; 6 Candlewood Lane, Southboro, MA 01772 (US). **LAN, Tao** [CN/US]; 995 Massachusetts Avenue, Apt #206, Arlington, MA 02476 (US). **WANG, Daqing** [US/US]; 7 Selfridge Road, Bedford, MA 01730 (US). **BHAGAT, Lakshmi** [IN/US]; 1550 Worcester Street, Unit 605, Framingham, MA 01702 (US).

Published:

- without international search report and to be republished upon receipt of that report

WO 2009/014887 A2

(54) Title: STABILIZED IMMUNE MODULATORY RNA (SIMRA) COMPOUNDS

(57) Abstract: The invention relates to the therapeutic use of novel stabilized oligoribonucleotides as immune modulatory agents for immune therapy applications. Specifically, the invention provides novel RNA-based oligoribonucleotides with improved nucleic acid and RNase stability and that have immune modulatory activity through TLR7 and/or TLR8.

STABILIZED IMMUNE MODULATORY RNA (SIMRA) COMPOUNDS

(Attorney Docket No. IDR-046PC)

BACKGROUND OF THE INVENTIONRelated Applications

[0001] This application claims the benefit of US Provisional Application Serial No. 60/948,529, filed on July 9, 2007; US Provisional Application Serial No. 60/957,195, filed on August 22, 2007; US Provisional Application Serial No. 60/981,161, filed on October 19, 2007; and US Provisional Application Serial No. 61/015,284, filed on December 20, 2007. The contents of these applications are incorporated herein by reference in their entirety.

Field of the Invention

[0002] The invention relates generally to the field of immunology and immunotherapy applications using oligoribonucleotides as immune modulatory agents. More particularly, the invention relates to immune modulatory RNA compositions and methods of use thereof for modulating the immune response through Toll-like receptor 8 (TLR8), Toll-like receptor 7 (TLR7) and TLR7 and TLR8.

Summary of the Related Art

[0003] The immune response involves both an innate and an adaptive response based upon the subset of cells involved in the response. For example, the T helper (Th) cells involved in classical cell-mediated functions such as delayed-type hypersensitivity and activation of cytotoxic T lymphocytes (CTLs) are Th1 cells, whereas the Th cells involved as helper cells for B-cell activation are Th2 cells. The type of immune response is influenced by the cytokines and chemokines produced in response to antigen exposure. Cytokines provide a means for controlling the immune response by affecting the balance of T helper 1 (Th1) and T helper 2 (Th2) cells, which directly affects the type of immune response that occurs. If the balance is toward higher numbers of Th1 cells, then a cell-mediated immune response occurs, which includes activation of cytotoxic T cells (CTLs). When the balance is toward higher numbers of Th2 cells, then a humoral or antibody immune response occurs. Each of these immune response results in a different set of cytokines being secreted from Th1 and Th2 cells. Differences in the cytokines secreted by Th1 and Th2 cells may be the result of the different biological functions of these two T cell subsets.

[0004] Th1 cells are involved in the body's innate response to antigens (e.g. viral infections, intracellular pathogens, and tumor cells). The initial response to an antigen can be the secretion of IL-12 from antigen presenting cells (e.g. activated macrophages and dendritic cells) and the concomitant activation of Th1 cells. The result of activating Th1 cells is a secretion of certain cytokines (e.g. IL-2, IFN-gamma and other cytokines) and a concomitant activation of antigen-specific CTLs. Th2 cells are known to be activated in response to bacteria, parasites, antigens, and allergens and may mediate the body's adaptive immune response (e.g. immunoglobulin production and eosinophil activation) through the secretion of certain cytokines (e.g. IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13 and other cytokines) and chemokines. Secretion of certain of these cytokines may result in B-cell proliferation and an increase in antibody production. In addition, certain of these cytokines may stimulate or inhibit the release of other cytokines (e.g. IL-10 inhibits IFN- γ secretion from Th1 cells and IL-12 from dendritic cells). Ultimately, the balance between Th1 and Th2 cells and the cytokines and chemokines released in response to selected stimulus can have an important role in how the body's immune system responds to disease. For example, IFN- α may inhibit hepatitis C, and MIP-1 α and MIP-1 β (also known as CCL3 and CCL4 respectively) may inhibit HIV-1 infection. Optimal balancing of the Th1/Th2 immune response presents the opportunity to use the immune system to treat and prevent a variety of diseases.

[0005] The Th1 immune response can be induced in mammals for example by introduction of bacterial or synthetic DNA containing unmethylated CpG dinucleotides, which immune response results from presentation of specific oligonucleotide sequences (e.g. unmethylated CpG) to receptors on certain immune cells known as pattern recognition receptors (PRRs). Certain of these PRRs are Toll-like receptors (TLRs).

[0006] TLRs are intimately involved in inducing the innate immune response in response to microbial infection. In vertebrates, TLRs consist of a family of ten proteins (TLR1 to TLR10) that are known to recognize pathogen associated molecular patterns. Of the ten, TLR3, 7, 8, and 9 are known to localize in endosomes inside the cell and recognize nucleic acids (DNA and RNA) and small molecules such as nucleosides and nucleic acid metabolites. TLR3 and TLR9 are known to recognize nucleic acid such as dsRNA and unmethylated CpG dinucleotide present in viral and bacterial and synthetic DNA, respectively. Bacterial DNA has been shown to activate the immune system and to generate antitumor activity (Tokunaga T et al., J. Natl. Cancer Inst. (1984) 72:955-962; Shimada S, et al., Jpn. J. cancer Res., 1986, 77, 808-816; Yamamoto S, et al., Jpn. J. Cancer Res., 1986, 79,

866-73; Messina, J, et al., *J. Immunol.* (1991) 147:1759-1764). Other studies using antisense oligonucleotides containing CpG dinucleotides have shown stimulation of an immune response (Zhao Q, et al., *Biochem.Pharmacol.* 1996, 26, 173-82). Subsequent studies showed that TLR9 recognizes unmethylated CpG motifs present in bacterial and synthetic DNA (Hemmi H, et al., *Nature.* (2000) 408:740-5). Other modifications of CpG-containing phosphorothioate oligonucleotides can also affect their ability to act through TLR9 and modulate the immune response (see, e.g., Zhao et al., *Biochem. Pharmacol.* (1996) 51:173-182; Zhao et al., *Biochem Pharmacol.* (1996) 52:1537-1544; Zhao et al., *Antisense Nucleic Acid Drug Dev.* (1997) 7:495-502; Zhao et al., *Bioorg. Med. Chem. Lett.* (1999) 9:3453-3458; Zhao et al., *Bioorg. Med. Chem. Lett.* (2000) 10:1051-1054; Yu et al., *Bioorg. Med. Chem. Lett.* (2000) 10:2585-2588; Yu et al., *Bioorg. Med. Chem. Lett.* (2001) 11:2263-2267; and Kandimalla et al., *Bioorg. Med. Chem.* (2001) 9:807-813). In addition, structure activity relationship studies have allowed identification of synthetic motifs and novel DNA-based structures that induce specific immune response profiles that are distinct from those resulting from unmethylated CpG dinucleotides. (Kandimalla ER, et al., *Proc Natl Acad Sci U S A.* (2005) 102:6925-30. Kandimalla ER, et al., *Proc Natl Acad Sci U S A.* (2003) 100:14303-8. Cong YP, et al., *Biochem Biophys Res Commun.* (2003) 310:1133-9. Kandimalla ER, et al., *Biochem Biophys Res Commun.* (2003) 306:948-53. Kandimalla ER, et al., *Nucleic Acids Res.* (2003) 31:2393-400. Yu D, et al., *Bioorg Med Chem.* (2003) 11:459-64. Bhagat L, et al., *Biochem Biophys Res Commun.* (2003) 300:853-61. Yu D, et al., *Nucleic Acids Res.* (2002) 30:4460-9. Yu D, et al., *J Med Chem.* (2002) 45:4540-8. Yu D, et al., *Biochem Biophys Res Commun.* (2002) 297:83-90. Kandimalla ER, et al., *Bioconjug Chem.* (2002) 13:966-74. Yu D, K et al., *Nucleic Acids Res.* (2002) 30:1613-9. Yu D, et al., *Bioorg Med Chem.* (2001) 9:2803-8. Yu D, et al., *Bioorg Med Chem Lett.* (2001) 11:2263-7. Kandimalla ER, et al., *Bioorg Med Chem.* (2001) 9:807-13. Yu D, et al., *Bioorg Med Chem Lett.* (2000) 10:2585-8, Putta MR, et al., *Nucleic Acids Res.* (2006) 34:3231-8). However, until recently, natural ligands for TLR7 and TLR8 were unknown.

[0007] It has been shown that TLRs 7 and 8 recognize viral and synthetic single-stranded RNAs and small molecules, including a number of nucleosides (Diebold, S.S., et al., *Science* v: 303, 1529-1531 (2004). Diebold et al. (*Science*, 303:1529-1531 (2004)) show that the IFN- α response to influenza virus requires endosomal recognition of influenza genomic RNA and signaling by means of TLR7 and MyD88 and identify ssRNA as a ligand for TLR7. Certain synthetic compounds, the imidazoquinolones, imiquimod (R-837) and resiquimod (R-

848) are ligands of TLR7 and TLR8 (Hemmi H et al., (2002) *Nat Immunol* 3:196-200; Jurk M et al., (2002) *Nat Immunol* 3:499). In addition, certain guanosine analogs, such as 7-deaza-G, 7-thia-8-oxo-G (TOG), and 7-allyl-8-oxo-G (loxoribine), have been shown to activate TLR7 at high concentrations (Lee J et al., *Proc Natl Acad Sci USA*. 2003, 100:6646-51). However, these small molecules, eg. imiquimod, are also known to act through other receptors (Schon MP, et al., (2006) *J. Invest Dermatol.*, 126, 1338-47)

[0008] The lack of any known specific ssRNA motif for TLR7 or TLR8 recognition and the potentially wide range of stimulatory ssRNA molecules suggest that TLR7 and TLR8 can recognize both self and viral RNA. Recently it was shown that certain GU-rich oligoribonucleotides are immunostimulatory and act through TLR7 and TLR8 (Heil et al. *Science*, 303: 1526-1529 (2004); Lipford et al. WO03/086280; Wagner et al. WO98/32462) when complexed with N-[1-(2,3-Dioleyloxy)propyl]-N,N,N trimethylammoniummethylsulfate (DOTAP) or other lipid agents. However, RNA molecules have been used for many years, for example as ribozymes and, more recently, siRNA and microRNA, and RNA employed as ribozymes, siRNA, and microRNA contain GU dinucleotides. In addition, a number these RNA molecules have been shown to elicit immune responses through TLR stimulation in the presence of lipids (Kariko et al., *Immunity* (2005) 23:165-75; Ma Z et al., *Biochem Biophys Res Commun.*, (2005) 330, 755-9). However, the instability of these RNA molecules has hindered progress in using and applying these molecules in many areas (e.g. prevention and treatment of human disease).

[0009] Oligonucleotides and oligodeoxynucleotides containing a ribose or deoxyribose sugar have been used in a wide variety of fields, including but not limited to diagnostic probing, PCR priming, antisense inhibition of gene expression, siRNA, microRNA, aptamers, ribozymes, and immunotherapeutic agents based on Toll-like Receptors (TLRs). More recently, many publications have demonstrated the use of oligodeoxynucleotides as immune modulatory agents and their use alone or as adjuvants in immunotherapy applications for many diseases, such as allergy, asthma, autoimmunity, cancer and infectious disease.

[0010] The fact that DNA oligonucleotides are recognized by TLR9, while RNA oligonucleotides are recognized by TLR7 and/or TLR8 is most likely due to differences in the structural conformations between DNA and RNA. However, the chemical differences between DNA and RNA also make DNA far more chemically and enzymatically stable than RNA.

[0011] RNA is rapidly degraded by ubiquitous extracellular ribonucleases (RNases) which ensure that little, if any, self-ssRNA reaches the antigen-presenting cells. Exonuclease degradation of nucleic acids is predominantly of 3'-nuclease digestion with a smaller percentage through 5'-exonuclease action. In addition to exonuclease digestion, RNA can also be degraded by endonuclease activity of RNases. RNA-based molecules have so far had to be complexed with lipids to provide stability against nucleases.

[0012] While providing an essential function of preventing autoimmune reactivity, these ribonucleases also present a substantial problem for any synthetic ssRNA molecule designed to be exploited for immunotherapy, as ribonucleases will rapidly degrade both synthetic and natural ssRNA. To overcome this hurdle, protection of ssRNA molecules from degradation has been attempted by encapsulating the ssRNA in liposomes, condensing it with polyethylenimine, or complexing it to molecules such as N-[1-(2,3 dioleyloxy)-propyl]-N,N,N-trimethylammonium methyl-sulfate (DOTAP). However, these protective measures are secondary measures applied to a still unstable ssRNA, and the effects of these protective measures on the *in vivo* efficacy and immune modulatory activity of ssRNA (natural or synthetic) remain unclear.

[0013] Agrawal et al. (11/697,422) describe a novel class of SIMRA compositions. However, a challenge remains to develop additional compounds that retain the naked RNA such that it continues to be recognized as a ligand for TLR7 and/or TLR8, while improving its stability such that it can be made to be a useful *in vivo* molecule. Ideally, this challenge might be met through the design of inherently stable RNA-based molecules that can act as new immunotherapeutic agents, which will find use in a number of clinically relevant applications, such as improving the effects of vaccination when co-administered or treating and/or preventing diseases when invoking or enhancing an immune response is beneficial, for example cancer, autoimmune disorders, airway inflammation, inflammatory disorders, infectious diseases, skin disorders, allergy, asthma or diseases caused by pathogens.

BRIEF SUMMARY OF THE INVENTION

[0014] In a first aspect, the invention provides novel stabilized immune modulatory RNA (“SIMRA”) compounds, further defined below, and their use for inducing and/or enhancing an immune response. The novel chemical entities according to the invention provide immune response inducing and/or enhancing compounds that are substantially more effective at inducing an immune response and substantially less susceptible to degradation. The methods according to the invention enable using SIMRA to modify the cytokine and/or chemokine profile for immunotherapy applications.

[0015] In one embodiment of the first aspect, the invention provides a SIMRA compound as an agonist for TLR8.

[0016] In another embodiment of the first aspect, the invention provides a SIMRA compound as an agonist for TLR7 and TLR8.

[0017] In a further embodiment of the first aspect, the invention provides a SIMRA compound as an agonist for TLR7.

[0018] In a further embodiment of the first aspect, the invention provides a SIMRA compound as an adjuvant.

[0019] In a second aspect, the invention provides pharmaceutical compositions. These compositions comprise any one of the SIMRA compositions of the invention and a physiologically acceptable or pharmaceutically acceptable carrier.

[0020] In a third aspect, the invention provides a method for generating an immune response in a vertebrate, the method comprising administering to the vertebrate at least one of the SIMRA compounds according to the invention in a pharmaceutically effective amount.

[0021] In a fourth aspect, the invention provides a method for therapeutically treating a vertebrate having a disease or disorder where inducing and/or enhancing an immune response would be beneficial, for example cancer, autoimmune disorders, airway inflammation, inflammatory disorders, infectious diseases, skin disorders, allergy, asthma or diseases caused by pathogens, such method comprising administering to the patient having such a disorder or disease at least one of the SIMRA compounds according to the invention in a pharmaceutically effective amount.

[0022] In a fifth aspect, the invention provides a method for preventing a disease or disorder in a vertebrate where inducing and/or enhancing an immune response would be

08 Apr 2011

2008279509

beneficial, for example cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, infectious disease, skin disorders, allergy, asthma or diseases caused by a pathogen, such method comprising administering to a vertebrate that is susceptible to such a disorder or disease at least one of the SIMRA compounds according to the invention in a pharmaceutically effective amount.

[0023] In a sixth aspect, the invention provides a method of isolating cells capable of producing cytokine or chemokines (e.g. immune cells, PBMCs), culturing such cells under standard cell culture conditions, ex vivo treating such cells with at least one of the SIMRA compounds of the invention such that the isolated cells produce or secrete increased levels of cytokines or chemokines, and administering or re-administering the treated cells to a patient in need of cytokine or chemokine therapy for the prevention or treatment of disease.

[0024] In a further embodiment of this aspect of the invention, the patient in need of cytokine or chemokine therapy for prevention or treatment of disease is administered the isolated, SIMRA-treated cells in combination with one or more SIMRA compounds.

[0024a] In one embodiment of the first aspect, the invention provides a stabilized immune modulatory RNA (SIMRA) compound selected from the group consisting of
5'-UG₁CUG₁CUUUUG₁-X-G₁UUUUCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-Z-G₁UUUCCG₁UCG₁U-5';
5'-UG₂CUGCUUGUG-X-GUGUUCGUCG₂U-5';
5'-UG₂CUG₂CUUG₂UG₂-X-G₂UG₂UUCG₂UCG₂U-5';
5'-UG₂CUG₂CUUG₂UG₂-M-G₂UG₂UUCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂U-5';
5'-UGC₁UGC₁UUGUG-X-GUGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁UUC₁UG-X-GUC₁UUC₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-M-GUUUC₁C₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-m-GUUUC₁C₁GUC₁GU-5';
5'-UGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGU-5';
5'-YUGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGUY-5';

2008279509 08 Apr 2011

5'-UGUUGUGUGA₁C-X-CA₁GUGUGUUGU-5';
5'-UG₂CUG₂CUUG₂UG₂-m-G₂UG₂UUCG₂UCG₂U-5';
5'-UGC₁UGC₁UUGUG-m-GUGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-X-GUUUC₁C₁GUC₁GU-5';
5'-YUGACGACGCUU-X-UUCGCAGCAGUY-5';
5'-YUGACGACUGCU-X-UCGUCAGCAGUY-5';
5'-YUGACGACUUGC-X-CGUUCAGCAGUY-5';
5'-YUGCGCGAACUU-X₃-UUCAAGCGCGUY-5';
5'-AG₁UG₁UUUUCUG₁-X-G₁UCUUUJUG₁UG₁A-5';
5'-UG₁UUG₁UUUG₁UG₁-X-G₁UG₁UUUG₁UUG₁U-5';
5'-UG₁AUG₁AAG₁CUU-X-UUCG₁AAG₁UAG₁U-5';
5'-UGUUCGAACAC-X-CACAAGCUUGU-5';
5'-UUGACGUUGAC-X-CAGUUGCAGUU-5';
5'-UUGACGUUGAC-Z-CAGUUGCAGUU-5';
5'-UUGACGUUGAC-M-CAGUUGCAGUU-5';
5'-UUGACGUUGAC-m-CAGUUGCAGUU-5';
5'-GUGCCUGAUGA-X-AGUAGUCCGUG-5';
5'-CCGAUGCCGAC-X-CAGCCGUAGCC-5';
5'-CCGAUGCAUCG-X-GCUACGUAGCC-5';
5'-GUGCCUGAUGA-Z-AGUAGUCCGUG-5';
5'-GUGCCUGAUGA-M-AGUAGUCCGUG-5';
5'-GUGCCUGAUGA-m-AGUAGUCCGUG-5';
5'-CCGAUGCCGAC-Z-CAGCCGUAGCC-5';
5'-CCGAUGCCGAC-M-CAGCCGUAGCC-5';
5'-CCGAUGCCGAC-m-CAGCCGUAGCC-5';
5'-CCGAUGCAUCG-Z-GCUACGUAGCC-5';
5'-CCGAUGCAUCG-M-GCUACGUAGCC-5';
5'-CCGAUGCAUCG-m-GCUACGUAGCC-5';
5'-AAAAAAA-X-AAAAAAA-5';
5'-YCACUGUUGAGA-X-AGAGUUGUCACY-5';
5'-YAACUGUUGACC-X-CCAGUUGUCAAY-5';
5'-YCAACGACCUGU-X-UGUCCAGCAACY-5';
5'-CACUG₁UUG₁AG₁A-X-AG₁AG₁UUG₁UCAC-5';

2008279509 08 Apr 2011

5'-AACUG₁UUG₁ACC-X-CCAG₁UUG₁UCAA-5';
5'-CAACG₁ACCUG₁U-X-UG₁UCCAG₁CAAC-5';
5'-AUGCUGCGCUG-X-GUCGCGUCGUA-5';
5'-AUGCUGCGCUG-Z-GUCGCGUCGUA-5';
5'-AUGCUGCGCUG-m-GUCGCGUCGUA-5';
5'-CCGAUGCAUCG-X-GCUACGUAGCC-5';
5'-AACGAACCGAC-X-CAGCCAAGCAA-5';
5'-YCAACGACCUGU-X-UGUCCAGCAACY-5';
5'-LCAACGACCUGU-X-UGUCCAGCAACL-5';
5'-CGUUGUGAUGA-X-AGUAGUGUUGC-5';
5'-ACGAUUGUGAC-X-CAGUGUUAGCA-5';
5'-ACUUUGACGAU-X-UAGCAGUUUCA-5';
5'-CGAUGCGAUGA-X-AGUAGCGUAGC-5';
5'-ACGUCUGACGA-X-AGCAGUCUGCA-5';
5'-AACUGCUGGAU-X-UAGGUCGUCAA-5';
5'-UUGGACUCCAG-X-GACCUUCAGGUU-5';
5'-UCGACUUCCAG-X-GACCUUCAGCU-5';
5'-CCGACUUGGAC-X-CAGGUUCAGCC-5';
5'-AAGACUGAACU-X-UCAAGUCAGAA-5';
5'-YUG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UY-5';
5'-YUG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UY-5';
5'-YUG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UY-5';
5'-LUG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UL-5';
5'-LUG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UL-5';
5'-LUG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UL-5';
5'-L₁UG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UL₁-5';
5'-L₁UG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UL₁-5';
5'-L₁UG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UL₁-5';
5'-G₁UCCUUG₁CUUG₁-m-G₁UUCG₁UUCCUG₁-5';
5'-UUCUG₁CUUCUG₁-m-G₁UCUUCG₁UCUU-5';
5'-UG₁CUG₁CCUUUG₁-M-G₁UUUCCG₁UCG₁U-5';
5'-UG₁CCUUUG₁CUG₁-M-G₁UCG₁UUUCCG₁U-5';
5'-UG₁CUG₁CUUCUG₁-M-G₁UCUUCG₁UCG₁U-5';

2008279509 08 Apr 2011

5'-UG₁C₁UG₁C₁C₁UUUG₁-X-G₁UUUC₁C₁G₁UC₁G₁U-5';
5'-UG₁C₁UG₁C₁UUC₁UG₁-X-G₁UC₁UUC₁G₁UC₁G₁U-5';
5'-YUGCUGCUUCUG-6Eg-M-6Eg-GUCUUCGUCGUY-5';
5'-UGGCUGACGC-X-CGCAGUUUCGGU-5';
5'-UGCUGCUUGAA-X-AAGUUCGUCGU-5';
5'-UGACGAUGAGU-X-UGAGUAGCAGU-5';
5'-UGAUGAGGACU-X-UCAGGAGUAGU-5';
5'-UGUUGAGGAAC-X-CAAGGAGUUGU-5';
5'-UGCGAGACUGC-X-CGUCAGAGCGU-5';
5'-GUUGAACGACU-X-UCAGCAAGUUG-5';
5'-UGACUGAUGAC-X-CAGUAGUCAGU-5';
5'-UGUUGAACGAC-X-CAGCAAGUUGU-5';
5'-UGAGCGUGAAC-X-CAAGUGCGAGU-5';
5'-UG₁G₁CUUG₁ACG₁C-X-CG₁CAG₁UUCG₁G₁U-5';
5'-UG₁ACG₁AUG₁AG₁U-X-UG₁AG₁UAG₁CAG₁U-5';
5'-UG₁UUG₁AG₁G₁AAC-X-CAAG₁G₁AG₁UUG₁U-5';
5'-UG₁UUG₁AACG₁AC-X-CAG₁CAAG₁UUG₁U-5';
5'-YUGCAGACUGC-X-CGUCAGAGCGUY-5';
5'-YUGAGCGUGAAC-X-CAAGUGCGAGU-5';
5'-YUUGAGCUGGAC-X-CAGGUCGAGUUY-5';
5'-YGUUGAGGAACU-X-UCAAGGAGUUGY-5';
5'-YUGAUGAAGCUU-X-UUCGAAGUAGUY-5';
5'-YUUGACGAUGAG-X-GAGUAGCAGUUY-5';
5'-YUUGUUGAACGA-X-AGCAAGUUGUUY-5';
5'-YUUGAACGACUU-X-UUCAGCAAGUUY-5';
5'-YUGAUGGAACGA-X-AGCAAGGUAGUY-5';
5'-UG₁CUG₁CCUUUG₁-M-G₁UUUCCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-Y-M-Y-G₁UUUCCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-3Eg-M-3Eg-G₁UUUCCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-4Eg-M-4Eg-G₁UUUCCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-6Eg-M-6Eg-G₁UUUCCG₁UCG₁U-5';
5'-YUGCUGCUUGUG-Y-M-Y-GUGUUCGUCGUY-5';
5'-YUGCUGCUUGUG-3Eg-M-3Eg-GUGUUCGUCGUY-5';

2008279509 08 Apr 2011

5'-YUGCUGCUUGUG-4Eg-M-4Eg-GGUUCGUCGUY-5';
5'-UGAACGUUGAA-X-AAGUUCGAAGU-5';
5'-UGAACGUGAAC-X-CAAGUGCAAGU-5';
5'-YUGC₄UGC₄UUGUG-X-G₁UC₄UUC₄G₁UC₄G₁U-5';
5'-UG₁C₄UG₁C₄UUC₄UG₁-X-G₁UC₄UUC₄G₁UC₄G₁U-5';
5'-YUG₁ACG₁AUG₁AG₁U-X-UG₁AG₁UAG₁CAG₁UY-5';
5'-G₁UCCUUG₁CUUG₁-X₁-G₁UUCG₁UUCCUG₁-5';
5'-UUCUG₁CUUCUG₁-X₁-G₁UCUUCG₁UCUU-5';
5'-UG₁CCUUUG₁CUG₁-X₂-G₁UCG₁UUUCCG₁U-5';
5'-UG₁CUG₁CUUCUG₁-X₂-G₁UCUUCG₁UCG₁U-5';
5'-PUUCUG₁CUUCUG₁-m-G₁UCUUCG₁UCUUP-5';
5'-PUG₁CCUUUG₁CUG₁-M-G₁UCG₁UUUCCG₁UP-5';
5'-UG₁CUG₁CUUC₁UG₁-X₁-G₁UC₁UUCG₁UCG₁U-5';
5'-UG₁CUG₁CUUC₁UG₁-X₃-G₁UC₁UUCG₁UCG₁U-5';
5'-C₂UGAACG₂UUGU-X-UGUUC₂GAAGUC₂-5';
5'-CU₂GAAGCU₂U₂GU₂-X-U₂GU₂U₂CGAACGU₂C-5';
5'-YAACUG₂UUG₂ACC-X-CCAG₂UUG₂UCAAY-5';
5'-L₁UG₂G₂CUUG₂ACG₂C-X-CG₂CAG₂UUCG₂G₂UL₁-5';
5'-YUGACUGCGACU-X-UCAGCGUCAGUY-5';
5'-YUGCGAACGCUU-X-UUCGCAAGCGUY-5';
5'-YUGCGACUGACU-X₃-UCAGUCAGCGUY-5';
5'-YUGCGCUGAACU-X₃-UCAAGUCGCGUY-5';
5'-YUGCUGACGACU-X₃-UCAGCAGUCGUY-5';
5'-YUGCUUGAACGC-X₃-CGCAAGUUCGUY-5';
5'-YUUGCUGAACGC-X₃-CGCAAGUCGUUY-5';
5'-UG₁CUG₁CCUUUG₁-Y-X-Y-G₁UUUCCG₁UCG₁U-5';
5'-YUGUUGUGUGAC-X-CAGUGUGUUGUY-5';
5'-YUGCUGCCUUUG-X-GUUUCCGUCGUY-5';
5'-YUGUUGUGUGAC-Z-CAGUGUGUUGUY-5';
5'-EUUGAACGACUU-X-UUCAGCAAGUUE-5';
5'-EUGUUGUGUGAC-X-CAGUGUGUUGUE-5';
5'-EUGCUGCCUUUG-X-GUUUCCGUCGUE-5';
5'-EUGCUGCUGCUG-X-GUCGUCGUCGUE-5';

2008279509 08 Apr 2011

5'-U₁GCUGCUUGUG-X-GGUUCGUCGU₁-5';
5'-U₁GCU₁GCUUGUG-X-GGUUCGU₁CGU₁-5';
5'-U₁GCU₁GCU₁U₁GU₁G-X-GU₁GU₁U₁CGU₁CGU₁-5';
5'-UG₁CUG₁CUUCUG₁-6Eg-M-6Eg-G₁UCUUCG₁UCG₁U-5';
5'-LUGCUGCCUUUG-m-GUUUCCGUCGUL-5';
5'-YUUGACGUUGAC-m-CAGUUGCAGUUY-5';
5'-YGUGCCUGAUGA-X-AGUAGUCCGUGY-5';
5'-AUGCUGCGCUG-M-GUCGCGUCGUA-5';
5'-UG₂CUG₂CUUG₂UG₂-m-G₂UG₂UUCG₂UCG₂U-5';
5'-UGC₁UGC₁UUGUG-m-GGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-X-GUUUC₁C₁GUC₁GU-5';
5'-GAUUGUGACGU-X-UGCAGUGUUAG-5';
5'-CUGAACGUUGU-X-UGUUCGAAGUC-5';
5'-UG₂CUG₂CUUG₂UG₂-M-G₂UG₂UUCG₂UCG₂U-5';
5'-UG₃CUG₃CUUCUG₃-X-G₃UCUUCG₃UCG₃U-5';
5'-UG₃CUG₃CCUUUG₃-m-G₃UUUCCG₃UCG₃U-5';
5'-YUGACGAUGAGU-X-UGAGUAGCAGUY-5';
5'-UGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGU-5';
5'-UGC₁UGC₁C₁UUUG-m-GUUUC₁C₁GUC₁GU-5';
5'-YUGCUGCUUGAA-X-AAGUUCGUCGUY-5';
5'-YUUGACUGAUGA-X-AGUAGUCAGUUY-5';
5'-UGCUGCUUUUG-X-GUUUUCGUCGU-5'; and
5'-UGUUGUUUGUG-X-GGUUUUGUUGU-5',
wherein G₁ = 7-deaza-rG; G₂ = ara-G; G₃ = 7-deaza-ara-G; C₁ = ara-C; C₂ = 2' -F-C; C₄ = 5-methyl-C; A₁ = ara-A; U₁ = ara-U; U₂ = 2' -F-U; M = cis,cis-cyclohexanetriol linker; m = cis,trans-cyclohexanetriol; Z = 1,3,5 -pentane triol linker; X = glycerol linker; X₁ = 1,2,4 -butane triol linker; X₂ = cyanuric acid; X₃ = isobutanetriol linker; Y = 1,3-propanediol; L = 1,5-pentanediol; L₁ = 1',2'-dideoxyribose; 6Eg = hexaethylene glycol linker; 3Eg = triethylene glycol linker; 4Eg = tetraethylene glycol linker; P = phosphorothioate; E = ethylene diol.

[0024b] In a further embodiment, the invention provides use of a SIMRA compound as described herein or a composition as described herein for the preparation of a medicament for generating an immune response in a vertebrate.

08 Apr 2011

2008279509

[0024c] In a further embodiment, the invention provides use of a SIMRA compound as described herein or a composition as described herein for the preparation of a medicament for therapeutically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial.

[0024d] In a further embodiment, the invention provides use of a SIMRA compound as described herein or a composition as described herein for the preparation of a medicament for prophylactically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial.

As used herein, except where the context requires otherwise, the term “comprise” and variations of the term, such as “comprising”, “comprises” and “comprised”, are not intended to exclude other additives, components, integers or steps.

Reference to any prior art in the specification is not, and should not be taken as, an acknowledgment, or any form of suggestion, that this prior art forms part of the common general knowledge in Australia or any other jurisdiction or that this prior art could reasonably be expected to be ascertained, understood and regarded as relevant by a person skilled in the art.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] SIMRA compounds of the invention were synthesized according to Example 1. Figure 1 is a synthetic scheme for the parallel synthesis of SIMRA compounds of the invention. DMTr = 4,4'-dimethoxytrityl; CE = cyanochethyl.

[0026] Figure 2A depicts NF- κ B activity in HEK293 cells expressing human TLR8 that were treated and analyzed according to example 2. Briefly, the HEK293 cells were stimulated with 150 μ g/ml of agonists of TLR8 for 18 hr, and the levels of NF- κ B were determined using SEAP (secreted form of human embryonic alkaline phosphatase) assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0027] Figure 2B depicts NF- κ B activity in HEK293 cells expressing human TLR8 that were treated and analyzed according to example 2. Briefly, the HEK293 cells were stimulated with 150 μ g/ml of agonists of TLR8 for 20 hr and the levels of NF- κ B were determined using SEAP assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0028] Figure 2C – 2E depict NF- κ B activity in HEK293 cells expressing human TLR8 that were treated and analyzed according to example 2. Briefly, the HEK293 cells were stimulated with 150 μ g/ml of agonists of TLR8 for 18 hr, and the levels of NF- κ B were determined using SEAP assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0029] Figure 2F depicts NF- κ B activity in HEK293 cells expressing human TLR8 that were treated and analyzed according to example 2. Briefly, HEK293 cells expressing human TLR8 were stimulated with 0, 20, 50, 100, 200, or 300 μ g/ml of agonists for 18 hr. The levels of NF- κ B were determined using SEAP assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0030] Figure 2G depicts NF- κ B activity in HEK293 cells expressing human TLR7 that were treated and analyzed according to example 2. Briefly, HEK293 cells expressing human TLR7 were stimulated with 0, 20, 50, 100, 200, or 300 μ g/ml of agonists for 18 hr. The levels of NF- κ B were determined using SEAP assay. The data demonstrate that

administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0031] Figures 2H, 2J and 2L depict NF- κ B activity in HEK293 cells expressing human TLR8 that were treated and analyzed according to example 2. Briefly, HEK293 cells expressing human TLR8 were stimulated with 0, 20, 50, 100, 200, or 300 μ g/ml of agonists for 18 hr. The levels of NF- κ B were determined using SEAP assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response profile.

[0032] Figures 2I, 2K and 2M depict NF- κ B activity in HEK293 cells expressing human TLR7 that were treated and analyzed according to example 2. Briefly, HEK293 cells expressing human TLR7 were stimulated with 0, 20, 50, 100, 200, or 300 μ g/ml of agonists for 18 hr. The levels of NF- κ B were determined using SEAP assay. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated immune response.

[0033] Figures 3A – 3C depict cytokine and chemokine concentrations from human PBMCs that were treated and analyzed according to example 3. Briefly, the PBMCs were isolated from freshly obtained healthy human volunteer's blood and cultured with 50 μ g/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0034] Figures 4A – 4C depict cytokine and chemokine concentrations from human PBMCs that were treated and analyzed according to example 3. Briefly, the PBMCs were isolated from freshly obtained healthy human volunteer's blood and cultured with 200 μ g/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0035] Figures 4D – 4AA depict cytokine and chemokine concentrations from human PBMCs that were treated and analyzed according to example 3. Briefly, the PBMCs were isolated from freshly obtained healthy human volunteer's blood and cultured with increasing concentrations of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by

Luminex multiplex assay cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct, dose-dependent, TLR-mediated cytokine and chemokine profile.

[0036] Figures 5A - 5C depict cytokine and chemokine concentrations from human plasmacytoid dendritic cells (pDCs) that were isolated, treated, and analyzed according to example 3. Briefly, the pDCs were isolated from freshly obtained healthy human volunteer's blood PBMCs and cultured with 50 µg/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay for cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0037] Figure 5D depicts cytokine and chemokine concentrations from human plasmacytoid dendritic cells (pDCs) that were isolated, treated, and analyzed according to example 3. Briefly, the pDCs were isolated from freshly obtained healthy human volunteer's blood PBMCs and cultured with a 50 µg/ml or 200 µg/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay for cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0038] Figures 6A, 6B, 6C, 6E and 6F depict cytokine and chemokine concentrations from human plasmacytoid dendritic cells (pDCs) that were treated and analyzed according to example 3. Briefly, the pDC were isolated from freshly obtained healthy human volunteer's blood PBMCs and cultured with 200 µg/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay for cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0039] Figure 6D depicts cytokine and chemokine concentrations from human plasmacytoid dendritic cells (pDCs) that were treated and analyzed according to example 3. Briefly, the pDC were isolated from freshly obtained healthy human volunteer's blood PBMCs and cultured with 100 µg/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay for cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0040] Figures 7A – 7C depict cytokine and chemokine concentrations from human myeloid dendritic cells (mDCs) that were treated and analyzed according to example 3. Briefly, the mDCs were isolated from freshly obtained healthy human volunteer's blood PBMCs and cultured with 50 µg/ml dose of TLR7/8 agonists for 24 hr, and supernatants were collected and analyzed by Luminex multiplex assay for cytokine and chemokine levels. The data demonstrate that administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0041] Figures 8A and 8B depict serum cytokine induction in C57BL/6 mice (n=3) 2 hours after they were treated and analyzed according to example 4. Briefly, the C57BL/6 mice were injected subcutaneously with 25 mg/kg dose of TLR7/8 agonists, and 2 hours after administration of the agonist, serum was analyzed for cytokine and chemokine levels, and IL-12 levels are presented. The data demonstrate that in vivo administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0042] Figures 9A and 9B depict serum cytokine induction in BALB/c mice (n=3) 2 hours after they were treated and analyzed according to example 4. Briefly, the BALB/c mice injected subcutaneously with 25 mg/kg dose of TLR7/8 agonists, and 2 hours after administration of the agonist, serum was analyzed for cytokine and chemokine levels, and IL-12 levels are presented. The data demonstrate that in vivo administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0043] Figures 9C - 9F depict serum cytokine induction in BALB/c mice (n=3) 2 hours after they were treated and analyzed according to example 4. Briefly, the BALB/c mice injected subcutaneously with 10 mg/kg or 25 mg/kg dose of TLR7/8 agonists, and 2 hours after administration of the agonist, serum was analyzed for cytokine and chemokine levels, and IL-12 levels are presented. The data demonstrate that in vivo administration of a SIMRA according to the invention generates a distinct TLR-mediated cytokine and chemokine profile.

[0044] Figures 10A -10H depict serum stability of exemplar SIMRA compounds from Table 2 that were treated according to example 5. Briefly, approximately 0.5 OD of exemplar SIMRA compounds were individually incubated in 1% human serum in PBS for 30 minute at 37 °C. At the end of the 30 minute incubation, the SIMRA compound was

analyzed on anion-exchange HPLC to determine the percentage of full-length SIMRA compound that remained as compared to the amount of SIMRA compound present before serum treatment.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0045] The invention relates to the therapeutic use of oligoribonucleotides as immune modulatory agents for immunotherapy applications. Specifically, the invention provides RNA-based oligonucleotides with improved in vivo stability that modulate the immune response through TLR7 alone, TLR7 and TLR8 or TLR8 alone (SIMRA compounds). By initiating diverse innate and acquired immune response mechanisms, for example through activation of dendritic cells and other antigen-presenting cells with stable agonists of TLR7 and/or TLR8, or SIMRA compounds, the resulting cytokine profile can lead to the destruction of pathogens, infected cells or tumor cells and development of antigen-specific antibody and CTL responses. Thus, the invention provides a diverse set of SIMRA compounds, each having its own unique immune regulatory characteristics. In this way, the scope and nature of the immune response can be customized for distinct medical indications by providing the SIMRA compound having the desired set of immune modulatory characteristics for that indication. The issued patents, patent applications, and references that are cited herein are hereby incorporated by reference to the same extent as if each was specifically and individually indicated to be incorporated by reference. In the event of inconsistencies between any teaching of any reference cited herein and the present specification, the latter shall prevail for purposes of the invention.

[0046] The invention provides methods for using SIMRA compounds to enhance the immune response. Such methods will find use in immunotherapy applications such as, but not limited to, treatment of cancer, autoimmune disorders, asthma, respiratory allergies, food allergies, skin allergies, and bacteria, parasitic, and viral infections in adult and pediatric human and veterinary applications. Thus, the invention further provides novel SIMRA compounds having optimal levels of immune modulatory effect for immunotherapy and methods for making and using such compounds. In addition, SIMRA compounds of the invention are useful as adjuvants or in combination with an agent useful for treating the disease or condition that does not diminish the immune modulatory effect of the SIMRA compound for prevention and treatment of diseases.

DEFINITIONS

[0047] The term "2'-substituted ribonucleoside" or "2'-substituted arabinoside" generally includes ribonucleosides or arabinonucleosides in which the hydroxyl group at the 2' position of the pentose moiety is substituted to produce a 2'-substituted or 2'-O-substituted

ribonucleoside. In certain embodiments, such substitution is with a lower hydrocarbyl group containing 1-6 saturated or unsaturated carbon atoms, with a halogen atom, or with an aryl group having 6-10 carbon atoms, wherein such hydrocarbyl, or aryl group may be unsubstituted or may be substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carboalkoxy, or amino groups. Arabinonucleosides of the invention include, but are not limited to, arabino-G, arabino-C, arabino-U, arabino-A. Examples of 2'-O-substituted ribonucleosides or 2'-O-substituted-arabinosides include, without limitation 2'-amino, 2'-fluoro, 2'-allyl, 2'-O-alkyl and 2'-propargyl ribonucleosides or arabinosides, 2'-O-methylribonucleosides or 2'-O-methylarabinosides and 2'-O-methoxyethoxyribonucleosides or 2'-O-methoxyethoxyarabinosides.

[0048] The term "3'", when used directionally, generally refers to a region or position in a polynucleotide or oligonucleotide 3' (toward the 3' position of the sugar) from another region or position in the same polynucleotide or oligonucleotide.

[0049] The term "5'", when used directionally, generally refers to a region or position in a polynucleotide or oligonucleotide 5' (toward the 5' position of the sugar) from another region or position in the same polynucleotide or oligonucleotide.

[0050] The term "about" generally means that the exact number is not critical. Thus, the number of ribonucleoside residues in the oligoribonucleotides is not critical, and oligoribonucleotides having one or two fewer ribonucleoside or arabinonucleoside residues, or from one to several additional ribonucleoside or arabinonucleoside residues are contemplated as equivalents of each of the embodiments described above.

[0051] The term "adjuvant" generally refers to a substance which, when added to an immunogenic agent such as vaccine or antigen, enhances or potentiates an immune response to the agent in the recipient host upon exposure to the mixture.

[0052] The term "airway inflammation" generally includes, without limitation, inflammation in the respiratory tract caused by infectious allergens, including asthma.

[0053] The term "allergen" generally refers to an antigen or antigenic portion of a molecule, usually a protein, which elicits an allergic response upon exposure to a subject. Typically the subject is allergic to the allergen as indicated, for instance, by the wheal and flare test or any method known in the art. A molecule is said to be an allergen even if only a small subset of subjects exhibit an allergic (e.g., IgE) immune response upon exposure to the molecule.

[0054] The term "allergy" generally includes, without limitation, food allergies, respiratory allergies, and skin allergies.

[0055] The term "antigen" generally refers to a substance that is recognized and selectively bound by an antibody or by a T cell antigen receptor. Antigens may include but are not limited to peptides, proteins, nucleosides, nucleotides, and combinations thereof. Antigens may be natural or synthetic and generally induce an immune response that is specific for that antigen.

[0056] The term "autoimmune disorder" generally refers to disorders in which "self" antigen undergo attack by the immune system.

[0057] Blocking 3' or 5' degradation or "cap" or "capping" means that the 3' or 5' end of the oligoribonucleotide is attached to another molecule (e.g. linker or other non-RNA nucleotide) to sufficiently inhibit nuclease degradation (e.g. 3' exonuclease degradation).

[0058] The term "carrier" generally encompasses any excipient, diluent, filler, salt, buffer, stabilizer, solubilizer, oil, lipid, lipid containing vesicle, microspheres, liposomal encapsulation, or other material well known in the art for use in pharmaceutical formulations. It will be understood that the characteristics of the carrier, excipient, or diluent will depend on the route of administration for a particular application. The preparation of pharmaceutically acceptable formulations containing these materials is described in, e.g., *Remington's Pharmaceutical Sciences*, 18th Edition, ed. A. Gennaro, Mack Publishing Co., Easton, PA, 1990.

[0059] The term "co-administration" generally refers to the administration of at least two different substances sufficiently close in time to modulate an immune response. Co-administration includes simultaneous administration of at least two different substances.

[0060] The term "complementary" generally means having the ability to hybridize to a nucleic acid. Such hybridization is ordinarily the result of hydrogen bonding between complementary strands, preferably to form Watson-Crick or Hoogsteen base pairs, although other modes of hydrogen bonding, as well as base stacking can also lead to hybridization.

[0061] The term "immune modulatory oligoribonucleotide" generally refers to an oligoribonucleotide that induces or represses an immune response when administered to a vertebrate, such as a fish, fowl or mammal.

[0062] The term "in combination with" generally means in the course of treating the same disease in the same patient, and includes administering a SIMRA compound and an agent useful for treating the disease or condition that does not diminish the immune modulatory effect of the SIMRA compound in any order, including simultaneous administration or co-administration, as well as temporally spaced order from a few seconds up to several days apart. Such combination treatment may also include more than a single administration of a SIMRA compound, and/or independently the agent. The administration of the SIMRA compound and the agent may be by the same or different routes.

[0063] The term "individual" or "subject" generally refers to a mammal, such as a human. Mammals generally include, but are not limited to, humans, non-human primates, rats, mice, cats, dogs, horses, cattle, cows, pigs, sheep, and rabbits.

[0064] The term "linear synthesis" generally refers to a synthesis that starts at one end of the immune modulatory oligoribonucleotide and progresses linearly to the other end. Linear synthesis permits incorporation of either identical or non-identical (in terms of length, base composition and/or chemical modifications incorporated) monomeric units into the immune modulatory oligoribonucleotides.

[0065] The term "linker" generally refers to any moiety that can be attached to an oligoribonucleotide by way of covalent or non-covalent bonding through a sugar, a base, or the backbone. The linker can be used to attach two or more nucleosides or can be attached to the 5' and/or 3' terminal nucleotide in the oligoribonucleotide. Such linker can be either a non-nucleotidic linker or a nucleotidic linker.

[0066] The term "modified nucleoside" generally is a nucleoside that includes a modified heterocyclic base, a modified sugar moiety, or any combination thereof. In some embodiments, the modified nucleoside is a non-natural pyrimidine or purine nucleoside, as herein described. For purposes of the invention, a modified nucleoside, a pyrimidine or purine analog or non-naturally occurring pyrimidine or purine can be used interchangeably and refers to a nucleoside that includes a non-naturally occurring base and/or non-naturally occurring sugar moiety. For purposes of the invention, a base is considered to be non-natural if it is not guanine, cytosine, adenine or uracil. In some embodiments, the modified nucleoside is a 2'-substituted ribonucleoside an arabinonucleoside or a 2'-deoxy-2'-substituted-arabinoside that can be substituted into selected positions of the oligoribonucleotide to improve stability without interfering with TLR7 or TLR8 activity.

[0067] The term "modulation" or "stimulation" generally refers to change, such as an increase in a response or qualitative difference in a response, which can arise from eliciting and/or enhancement of a response.

[0068] The term "non-nucleotidic linker" generally refers to a chemical moiety other than a nucleotidic linkage that can be attached to an oligoribonucleotide by way of covalent or non-covalent bonding. Preferably such non-nucleotidic linker is from about 2 angstroms to about 200 angstroms in length, and may be either in a cis or trans orientation.

[0069] The term "nucleotidic linkage" generally refers to a chemical linkage to join two nucleosides through their sugars (e.g. 3'-3', 2'-3', 2'-5', 3'-5') consisting of a phosphate, non-phosphate, charged, or neutral group (e.g., phosphodiester, phosphorothioate or phosphorodithioate) between adjacent nucleosides.

[0070] The term "peptide" generally refers to polypeptides that are of sufficient length and composition to affect a biological response, e.g., antibody production or cytokine activity whether or not the peptide is a hapten. The term "peptide" may include modified amino acids (whether or not naturally or non-naturally occurring), where such modifications include, but are not limited to, phosphorylation, glycosylation, pegylation, lipidization, and methylation.

[0071] The terms "pharmaceutically acceptable" or "physiologically acceptable" generally refer to a material that does not interfere with the effectiveness of a compound according to the invention and that is compatible with a biological system such as a cell, cell culture, tissue, or organism. Preferably, the biological system is a living organism, such as a vertebrate.

[0072] The term a "pharmaceutically effective amount" generally refers to an amount sufficient to affect a desired biological effect, such as a beneficial result. Thus, a "pharmaceutically effective amount" will depend upon the context in which it is being administered. A pharmaceutically effective amount may be administered in one or more prophylactic or therapeutic administrations.

[0073] The term "SIMRA" generally refers to stabilized immune modulatory RNA compounds which are recognized as ligands by TLR7 and/or TLR8, wherein the compounds may contain single-stranded RNA (ssRNA) and/or double-stranded RNA (dsRNA), and modifications to protect or stabilize its 3' ends (e.g. by blocking 3' degradation or by capping the 3' ends or by linking the 3' ends of two or more oligoribonucleotides), provided that the SIMRA is or would be more stable *in vivo* than an unmodified oligoribonucleotide and, thus,

affect its immune modulatory capabilities. The SIMRA may contain modified oligoribonucleotides. The SIMRA compound may also contain modifications to protect its 5' ends (e.g., by blocking 5' degradation or capping the 5' ends) to further improve the stability of the oligoribonucleotide(s). The SIMRA can be linear or branched, with nucleic acids being polymers of ribonucleosides linked through, for example, phosphodiester, phosphorothioate, or alternate linkages. A SIMRA may consist of a purine (adenine (A) or guanine (G) or derivatives thereof (e.g. 7-deaza-G, arabino-G and arabino-A)) or pyrimidine (cytosine (C) or uracil (U), or derivatives thereof (e.g. arabino-C and arabino-U)) base covalently attached to a ribose sugar residue or a derivative thereof.

[0074] The term "treatment" generally refers to an approach intended to obtain a beneficial or desired result, which may include alleviation of symptoms, or delaying or ameliorating a disease progression.

[0075] The term "viral disease" generally refers to a disease that has a virus as its etiologic agent, including but not limited to hepatitis B, hepatitis C, influenza, acquired immunodeficiency syndrome (AIDS), and herpes zoster.

[0076] In a first aspect, the invention provides novel SIMRA compounds. The present inventors have discovered that modification of an immune modulatory oligoribonucleotide to protect its 3' end (e.g. by blocking 3' degradation or capping the 3' end or by linking the 3' ends of two or more oligoribonucleotides) surprisingly affects its immune modulatory capabilities. In addition, it has been determined that this protection surprisingly improves the stability of the oligoribonucleotides, removing the need for lipid association or other means of protection. Further, blocking 5' degradation or capping the 5' end in addition to or in combination with protecting the 3'-end can also improve the stability of the oligoribonucleotide.


[0077] In the present invention activation of TLR8 and induction of unique immune responses (e.g. changes in cytokine and/or chemokine profiles) with novel SIMRA compounds is demonstrated. Moreover, the incorporation of certain chemical modification(s) in such human TLR8 activating RNAs can also activate TLR7, resulting in distinct immune response(s) and a change in cytokine and/or chemokine profiles. Thus, the present inventors have surprisingly discovered that through activation of TLR8 and/or TLR7 cytokine and/or chemokine profiles associated therewith can be modulated by using modified chemical


structures, including modified bases, modified sugars, backbone, linkers, linkages, and/or caps as part of an immune modulatory oligoribonucleotide.

[0078] In one embodiment, the invention provides an immune modulatory compound comprising at least two RNA-based oligonucleotides linked at their 3' ends, or an internucleoside linkage or a functionalized nucleobase or sugar to a non-nucleotidic linker. Such embodiment of the invention may have at least one accessible 5' end, which may be capped or uncapped. It has been determined that this structure provides further stability (e.g. inhibition of exonuclease activity) to the SIMRA compounds without the need for lipid association or other protection. An “accessible 5' end” means that the 5'-terminus of the SIMRA is not modified in such a way as to prevent the SIMRA compound from modulating an immune response through TLR7 and/or TLR8.

[0079] In another embodiment of this aspect of the invention comprises at least two oligoribonucleotides, wherein the immune modulatory compound has a structure including, but not limited to, those as detailed in Formulas I – X in Table 1.

Table 1: Oligoribonucleotide Formulas I – X





[0080] Domains A, B, C, and D may be independently from about 2 to about 35 ribonucleotides, and in some embodiments from about 2 to about 20, or from about 2 to about 12, or from about 2 to about 11 or from about 2 to about 8 ribonucleotides in length.

Domains A, B, C, and/or D may or may not be identical. Domains A, B, C, and D may independently be 5'-3' or 2'-5' RNA having or not having a self-complementary domain, a homo or hetero ribonucleotide sequence, or a linker. "n" may be from 1 to an unlimited number.

[0081] "X" is a linker joining or capping Domains A, B, C, and/or D that may be through a 3' or 5' linkage, a phosphate group, a nucleobase, a non-RNA nucleotide, or a non-nucleotidic linker that may be aliphatic, aromatic, aryl, cyclic, chiral, achiral, a peptide, a carbohydrate, a lipid, a fatty acid, mono- tri- or hexapolyethylene glycol, or a heterocyclic moiety, or combinations thereof.

[0082] In a further embodiment, the invention provides a SIMRA compound comprising at least two oligoribonucleotides linked by a non-nucleotidic linker, wherein the

sequences of the immune modulatory oligoribonucleotides may be at least partially self-complementary. As would be recognized by one skilled in the art, the complementary sequence of the oligoribonucleotides allows for intermolecular hydrogen bonding thereby giving the oligoribonucleotides secondary structure. Additional oligoribonucleotides can bind together thereby creating a chain, or multimers, of oligoribonucleotides according to the invention.

[0083] Similar considerations apply to intermolecular base pairing between immune modulatory oligoribonucleotides of different base sequence. Thus, where a plurality of immune modulatory oligoribonucleotides is used together, the plurality of immune modulatory oligoribonucleotides may, but need not, include sequences that are at least partially complementary to one another. In one embodiment the plurality of immune modulatory oligoribonucleotides includes an immune modulatory oligoribonucleotide having a first sequence and an immune modulatory oligoribonucleotide having a second sequence, wherein the first sequence and the second sequence are at least 50 percent complementary. For example, as between two 8-mers that are at least 50 percent complementary, they may form 4, 5, 6, 7, or 8 G-C, A-U, and/or G-U wobble basepairs. Such basepairs may, but need not necessarily, involve bases located at either end of the complementary immune modulatory oligoribonucleotides. The degree of complementarity may depend on the alignment between immune modulatory oligoribonucleotides, and such alignment may or may not include single- or multiple-nucleoside overhangs. In other embodiments, the degree of complementarity is at least 60 percent, at least 70 percent, at least 80 percent, at least 90 percent, or even 100 percent.


[0084] As would be recognized by one skilled in the art, the depicted immune modulatory compounds may have secondary structure because the sequences of the domains are complementary allowing for intermolecular hydrogen bonding. Moreover, as can be imagined from Formulas I through X, additional linked RNA-based oligonucleotides can bind through intermolecular hydrogen bonding thereby creating a chain, or multimers, wherein any number of linked RNA-based oligonucleotides may be incorporated.

[0085] In another embodiment, the invention provides an immune modulatory compound comprising at least two RNA-based oligonucleotides linked at their 3' or 5' ends, or through an internucleoside linkage or a functionalized nucleobase or sugar to a non-nucleotidic linker, and wherein a linker (e.g. cap) is attached to at least one 5' end. It has been determined that this structure provides further stability (e.g. inhibition of exonuclease

activity) to the SIMRA compounds. The 5'-terminus of the SIMRA is not modified in such a way as to prevent the SIMRA compound from modulating an immune response through TLR7 and/or TLR8.

[0086] In some embodiments, the oligoribonucleotides each independently have from about 2 to about 35 ribonucleoside residues. Thus in certain embodiments the oligoribonucleotide can independently be 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or 35 ribonucleotides long. Preferably the oligoribonucleotide is from about 4 to about 30 ribonucleoside residues, more preferably from about 4 to about 20 ribonucleoside residues or from about 4 to about 11 ribonucleoside residues. In some embodiments, the immune modulatory oligoribonucleotides comprise oligoribonucleotides having from about 1 to about 18, or from about 1 to about 11, or from about 5 to about 14 ribonucleoside residues. In some embodiments, one or more of the oligoribonucleotides have 11 ribonucleotides or from about 8 to about 14 ribonucleotides or from about 10 to about 12 ribonucleotides. In the context of immune modulatory oligoribonucleotides, preferred embodiments have from about 1 to about 35 ribonucleotides, preferably from about 5 to about 26 ribonucleotides, more preferably from about 13 to about 26 ribonucleotides. Preferably, the immune modulatory oligoribonucleotide comprises at least one phosphodiester, phosphorothioate, or phosphorodithioate interribonucleoside linkage.

[0087] In exemplar embodiments each ribonucleoside unit includes a heterocyclic base and a pentofuranosyl, trehalose, arabinose, 2'-deoxy-2'-substituted arabinose, 2'-O-substituted ribose or arabinose, or hexose sugar group. The ribonucleoside residues can be coupled to each other by any of the numerous known interribonucleoside linkages. Such interribonucleoside linkages include, without limitation, phosphodiester, phosphorothioate, phosphorodithioate, alkylphosphonate, alkylphosphonothioate, phosphotriester, phosphoramidate, siloxane, carbonate, carboalkoxy, acetamidate, carbamate, morpholino, borano, thioether, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphorothioate, and sulfone interribonucleoside linkages. Possible sites of conjugation for the ribonucleotide are indicated in Formula XI, below, wherein B represents a heterocyclic base.



[0088] The SIMRA compounds of the invention can include naturally occurring ribonucleosides, modified ribonucleosides, or mixtures thereof.

[0089] In the present invention, novel SIMRA compounds are recognized by human TLR8 and incorporation of certain chemical modification(s) in such human TLR8 activating RNAs can cause them to be recognized by human TLR7 and induce immune responses. Such chemical modifications include, but are not limited to, guanine analogues such as 7-deaza-G, ara-G, 6-thio-G, Inosine, Iso-G, loxoribine, TOG(7-thio-8-oxo)-G, 8-bromo-G, 8-hydroxy-G, 5-aminoformycin B, Oxoformycin, 7-methyl-G, 9-p-chlorophenyl-8-aza-G, 9-phenyl-G, 9-hexyl-guanine, 7-deaza-9-benzyl-G, 6-Chloro-7-deazaguanine, 6-methoxy-7-deazaguanine, 8-Aza-7-deaza-G(PPG), 2-(Dimethylamino)guanosine, 7-Methyl-6-thioguanosine, 8-Benzylxylguanosine, 9-Deazaguanosine, and 1-(B-D-ribofuranosyl)-2-oxo-7-deaza-8-methyl-purine. Chemical modifications also include, but are not limited to, adenine analogues such as 9-benzyl-8-hydroxy-2-(2-methoxyethoxy)adenine, 2-Amino-N2-O-, methyladenosine, 8-Aza-7-deaza-A, 7-deaza-A, ara-A, Vidarabine, 2-Aminoadenosine, N1-Methyladenosine, 8-Azaadenosine, 5-Iidotubercidin. Chemical modifications also include, but are not limited to, cytosine and uracil analogues such as pseudouridine, ara-C, ara-U, 5-methylcytidine, 4-thiouridine, N4-ethyluridine, zebularine, 5-aminoallyluridine, N3-methyluridine, 5-fluorouridine.

[0090] The “immune modulatory oligoribonucleotides” according to the invention are SIMRA compounds that comprise at least two oligoribonucleotides linked covalently or non-covalently at their 3'- or 2'-ends or functionalized ribose or functionalized ribonucleobase via a non-nucleotidic or a nucleotidic linker. Several examples of linkers are set forth below. Non-covalent linkages include, but are not limited to, electrostatic interaction, hydrophobic interactions, π -stacking interactions and hydrogen bonding.

[0091] In yet other embodiments, the non-nucleotidic linker is an organic moiety having functional groups that permit attachment to the oligoribonucleotide. Such attachment preferably is by a stable covalent linkage. As a non-limiting example, the linker may be attached to any suitable position on the nucleotide. In some preferred embodiments, the linker is attached to the 3'-hydroxyl. In such embodiments, the linker preferably comprises a hydroxyl functional group, which preferably is attached to the 3'-hydroxyl by means of a phosphate-based linkage like, phosphodiester, phosphorothioate, phosphorodithioate, methylphosphonate or non-phosphate-based linkages.

[0092] In some embodiments, the non-nucleotidic linker is a small molecule, macromolecule or biomolecule, including, without limitation, polypeptides, antibodies, lipids, antigens, allergens, and oligosaccharides. In some other embodiments, the non-nucleotidic linker is a small molecule. For purposes of the invention, a small molecule is an organic moiety having a molecular weight of less than 1,000 Da. In some embodiments, the small molecule has a molecular weight of less than 750 Da.

[0093] In some embodiments, the small molecule is an aliphatic or aromatic hydrocarbon, either of which optionally can include, either in the linear chain connecting the oligoribonucleotides or appended to it, one or more functional groups including, but not limited to, hydroxy, amino, thiol, thioether, ether, amide, thioamide, ester, urea, or thiourea. The small molecule can be cyclic or acyclic. Examples of small molecule linkers include, but are not limited to, amino acids, carbohydrates, cyclodextrins, adamantane, cholesterol, haptens and antibiotics. However, for purposes of describing the non-nucleotidic linker, the term "small molecule" is not intended to include a nucleoside.

[0094] In some embodiments, the non-nucleotidic linker is an alkyl linker or amino linker. The alkyl linker may be branched or unbranched, cyclic or acyclic, substituted or unsubstituted, saturated or unsaturated, chiral, achiral or racemic mixture. The alkyl linkers can have from about 2 to about 18 carbon atoms. In some embodiments such alkyl linkers have from about 3 to about 9 carbon atoms. Some alkyl linkers include one or more functional groups including, but not limited to, hydroxy, amino, thiol, thioether, ether, amide, thioamide, ester, urea, and thioether. Such alkyl linkers can include, but are not limited to, 1-propanol, 1,2 propanediol, 1,3 propanediol, 1,2,3, propanetriol, triethylene glycol, hexaethylene glycol, polyethylene glycol linkers (e.g. [-O-CH₂-CH₂-]_n (n= 1-9)), methyl linkers, ethyl linkers, propyl linkers, butyl linkers, or hexyl linkers. In some embodiments, such alkyl linkers may include peptides or amino acids.

[0095] In some embodiments, the non-nucleotidic linker may include, but are not limited to, those listed in Table 2.





Table 2: Representative Non-nucleotidic Linkers

Table 2: Continued


cis-1,3,5-Cyclohexanetriol




1,3-Di(hydroxyethoxy)-2-hydroxyl-propane




cis-1,3,5-Tri(hydroxymethyl)cyclohexane




1,3-Di(hydroxypropoxy)-2-hydroxyl-propane




1,3,5-Trihydroxyl-benzene




2-Deoxy-D-ribose




3,5-Di(hydroxymethyl)phenol




1,2,4-Trihydroxyl-benzene




1,3,5-Tri(hydroxymethyl)benzene




D-Galactoal

Table 2: Continued1,6-anhydro- β -D-Glucose


4,6-Nitropyrogallol




1,3,5-Tris(2-hydroxyethyl)-Cyanuric acid



Gallic acid




3,5,7-Trihydroxyflavone

Table 2: Continued


Ethylene glycol




1,3-Propanediol




1,2-Propanediol




1,5-Pentanediol




2,4-Pentanediol




1,6-Hexanediol




1,4-Butanediol




1,2-Hexanediol




1,3-Butanediol




1,5-Hexanediol



2,3-Butanediol



2,5-Hexanediol



1,4-Butanediol





Table 2: Continued

Table 2: Continued


Cis, cis-cyclohexanetriol linker



Cis, trans-cyclohexanetriol linker



1,3,4- Isobutanetriol



Cyanuric Acid

[0096] In some embodiments, the small molecule linker is glycerol or a glycerol homolog of the formula $\text{HO}-(\text{CH}_2)_o-\text{CH}(\text{OH})-(\text{CH}_2)_p-\text{OH}$, wherein o and p independently are integers from 1 to about 6, from 1 to about 4, or from 1 to about 3. In some other embodiments, the small molecule linker is a derivative of 1,3-diamino-2-hydroxypropane. Some such derivatives have the formula

$\text{HO}-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHC}(\text{O})-(\text{CH}_2)_n-\text{OH}$, wherein m is an integer from 0 to about 10, from 0 to about 6, from 2 to about 6, or from 2 to about 4.

[0097] Some non-nucleotidic linkers according to the invention permit attachment of more than two oligoribonucleotides, as depicted in Table 1. For example, the small molecule linker glycerol has three hydroxyl groups to which oligoribonucleotides may be covalently attached. Some immune modulatory oligoribonucleotides according to the invention, therefore, comprise more than two oligoribonucleotides (e.g., a Domain C and so on, the additional domains comprise oligoribonucleotides as defined above for Domains A, B, C, and D) linked at their 3' ends to a non-nucleotidic linker.

[0098] In a further embodiment of this aspect of the invention, a SIMRA may contain three or more oligoribonucleotides linked at their 3' or 5' ends, or through an internucleoside linkage or a functionalized nucleobase or sugar to two or more linkers, as depicted in Table 1.

The oligoribonucleotides of this aspect of the invention may have the same or different sequences. The linkers of this aspect of the invention may be the same or different.

[0099] The immune modulatory oligoribonucleotides of the invention may conveniently be synthesized using an automated synthesizer and phosphoramidite approach. In some embodiments, the immune modulatory oligoribonucleotides are synthesized by a linear synthesis approach.

[00100] An alternative mode of synthesis is "parallel synthesis", in which synthesis proceeds outward from a central linker moiety (see Figure 1). A solid support attached linker can be used for parallel synthesis, as is described in U.S. Patent No. 5,912,332. Alternatively, a universal solid support (such as phosphate attached controlled pore glass) support can be used.

[00101] Parallel synthesis of immune modulatory oligoribonucleotides has several advantages over linear synthesis: (1) parallel synthesis permits the incorporation of identical monomeric units; (2) unlike in linear synthesis, both (or all) the monomeric units are synthesized at the same time, thereby the number of synthetic steps and the time required for the synthesis is the same as that of a monomeric unit; and (3) the reduction in synthetic steps improves purity and yield of the final immune modulatory oligoribonucleotide product.

[00102] At the end of the synthesis by either linear synthesis or parallel synthesis protocols, the immune modulatory oligoribonucleotides may conveniently be deprotected with concentrated ammonia solution or as recommended by the phosphoramidite supplier, if a modified nucleoside is incorporated. The product immune modulatory oligoribonucleotide is preferably purified by reversed phase HPLC, detritylated, desalting and dialyzed.

[00103] Table 3 shows RNA- based immune modulatory oligoribonucleotides according to the invention. Unless otherwise specified, all nucleosides are ribonucleosides.

Table 3. Stabilized RNA-based Immune Modulatory Oligonucleotide (SIMRA) Sequences

| SIMRA# (SEQ ID NO.) | Sequences and Modification |
|------------------------|---|
| 1 | 5'-YUUCUG ₁ CUUCUG ₁ -X-G ₁ UCUUCG ₁ UCUUY-5' |
| 2 | 5'-UG ₁ CUG ₁ CCUUUG ₁ -X-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 3 | 5'-UG ₁ CUG ₁ CCUUUG ₁ -Z-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 4 | 5'-G ₁ UCCUUUG ₁ CUG ₁ -X-G ₁ UCG ₁ UUUCCUG ₁ -5' |
| 5 | 5'-L ₁ UGCUGCUUGUG-X-GUGUUCGUCGUL ₁ -5' |
| 6 | 5'-LUGCUGCCUUUG-m-GUUUCCGUCGUL-5' |
| 7 | 5'-G ₁ UCCUUG ₁ CUUG ₁ -M-G ₁ UUCG ₁ UUCCUG ₁ -5' |
| 8 | 5'-UUCUG ₁ CUUCUG ₁ -M-G ₁ UCUUCG ₁ UCUU-5' |
| 9 | 5'-G ₁ UCCUUUG ₁ CUG ₁ -m-G ₁ UCG ₁ UUUCCUG ₁ -5' |
| 10 | 5'-YUUGACGUUGAC-m-CAGUUGCAGUUY-5' |
| 11 | 5'-YGUGCCUGAUGA-X-AGUAGUCCGUGY-5' |
| 12 | 5'-AUGCUGCGCUG-M-GUCGCGUCGUA-5' |
| 13 | 5'-UGCUGCUUG ₂ UG-X-GUG ₂ UUCGUCGU-5' |
| 14 | 5'-UG ₂ CUGCUUGUG-X-GUGUUCGUCG ₂ U-5' |
| 15 | 5'-UG ₂ CUG ₂ CUUG ₂ UG ₂ -X-G ₂ UG ₂ UUCG ₂ UCG ₂ U-5' |
| 16 | 5'-UG ₂ CUG ₂ CUUG ₂ UG ₂ -m-G ₂ UG ₂ UUCG ₂ UCG ₂ U-5' |
| 17 | 5'-UG ₂ CUG ₂ CUUG ₂ UG ₂ -M-G ₂ UG ₂ UUCG ₂ UCG ₂ U-5' |
| 18 | 5'-UG ₂ CUG ₂ CCUUUG ₂ -M-G ₂ UUUCCG ₂ UCG ₂ U-5' |
| 19 | 5'-UG ₂ CUG ₂ CCUUUG ₂ -m-G ₂ UUUCCG ₂ UCG ₂ U-5' |
| 20 | 5'-UG ₂ CUG ₂ CCUUUG ₂ -X-G ₂ UUUCCG ₂ UCG ₂ U-5' |
| 21 | 5'-UGC ₁ UGC ₁ UUGUG-X-GUGUUC ₁ GUC ₁ GU-5' |
| 22 | 5'-UGC ₁ UGC ₁ UUGUG-m-GUGUUC ₁ GUC ₁ GU-5' |
| 23 | 5'-UGC ₁ UGC ₁ UUGUG-M-GUGUUC ₁ GUC ₁ GU-5' |
| 24 | 5'-UGC ₁ UGC ₁ UUC ₁ UG-X-GUC ₁ UUC ₁ GUC ₁ GU-5' |
| 25 | 5'-UGC ₁ UGC ₁ C ₁ UUUG-M-GUUUC ₁ C ₁ GUC ₁ GU-5' |
| 26 | 5'-UGC ₁ UGC ₁ C ₁ UUUG-m-GUUUC ₁ C ₁ GUC ₁ GU-5' |
| 27 | 5'-UGC ₁ UGC ₁ C ₁ UUUG-X-GUUUC ₁ C ₁ GUC ₁ GU-5' |
| 28 | 5'-UGCUGCU ₁ U ₁ CU ₁ G-X-GU ₁ CU ₁ U ₁ CGUCGU-5' |
| 29 | 5'-YUGCUGCU ₁ U ₁ CU ₁ G-X-GU ₁ CU ₁ U ₁ CGUCGUY-5' |
| 30 | 5'-UGUUGUGUGA ₁ C-X-CA ₁ GUGUGUUGU-5' |
| 31 | 5'-UG ₂ CUG ₂ CUUG ₂ UG ₂ -m-G ₂ UG ₂ UUCG ₂ UCG ₂ U-5' |

| | |
|----|---|
| 32 | 5'-UGC ₁ UGC ₁ UUGUG-m-GGUUC ₁ GUC ₁ GU-5' |
| 33 | 5'-UGC ₁ UGC ₁ C ₁ UUUG-X-GUUUC ₁ C ₁ GUC ₁ GU-5' |
| 34 | 5'-GAUUGUGACGU-X-UGCAGUGUAG-5' |
| 35 | 5'-CUGAACGUUGU-X-UGUUCGAAGUC-5' |
| 36 | 5'-UG ₂ CUG ₂ CUUG ₂ UG ₂ -M-G ₂ UG ₂ UUCG ₂ UCG ₂ U-5' |
| 37 | 5'-YUGCUGCUUGUG-X-GGUUCGUCGUY-5' |
| 38 | 5'-UG ₁ CUG ₁ CUUCUG ₁ -X-G ₁ UCUUCG ₁ UCG ₁ U-5' |
| 39 | 5'-UG ₃ CUG ₃ CUUCUG ₃ -X-G ₃ UCUUCG ₃ UCG ₃ U-5' |
| 40 | 5'-UG ₃ CUG ₃ CCUUUG ₃ -m-G ₃ UUUCCG ₃ UCG ₃ U-5' |
| 41 | 5'-YUGACGAUGAGU-X-UGAGUAGCAGUY-5' |
| 42 | 5'-UGCUGCU ₁ U ₁ CU ₁ G-X-GU ₁ CU ₁ U ₁ CGUCGU-5' |
| 43 | 5'-UGC ₁ UGC ₁ C ₁ UUUG-m-GGUUC ₁ C ₁ GUC ₁ GU-5' |
| 44 | 5'-UG ₁ CUG ₁ CUUCUG ₁ -6Eg-M-6Eg-G ₁ UCUUCG ₁ UCG ₁ U-5' |
| 45 | 5'-YUGACGACGCUU-X-UUCGCAGCAGUY-5' |
| 46 | 5'-YUGACGACUGCU-X-UCGUCAGCAGUY-5' |
| 47 | 5'-YUGACGACUUGC-X-CGUUCAGCAGUY-5' |
| 48 | 5'-YUGCGCGAACUU-X ₃ -UUCAAGCGCGUY-5' |
| 49 | 5'-AG ₁ UG ₁ UUUCUG ₁ -X-G ₁ UCUUUUG ₁ UG ₁ A-5' |
| 50 | 5'-UG ₁ CUG ₁ CUUUUG ₁ -X-G ₁ UUUUCG ₁ UCG ₁ U-5' |
| 51 | 5'-UG ₁ UUG ₁ UUUG ₁ UG ₁ -X-G ₁ UG ₁ UUUG ₁ UUG ₁ U-5' |
| 52 | 5'-UG ₁ AUG ₁ AAG ₁ CUU-X-UUCG ₁ AAG ₁ UAG ₁ U-5' |
| 53 | 5'-YUGCUGCUUGAA-X-AAGUUCGUCGUY-5' |
| 54 | 5'-YUUGACUGAUGA-X-AGUAGUCAGUUY-5' |
| 55 | 5'-UGCUGCUUUUG-X-GUUUUCGUCGU-5' |
| 56 | 5'-UGUUGUUUGUG-X-GGUUUUGUUGU-5' |
| 57 | 5'-UGUUCGAACAC-X-CACAAGCUUGU-5' |
| 58 | 5'-UUGACGUUGAC-X-CAGUUGCAGUU-5' |
| 59 | 5'-UUGACGUUGAC-Z-CAGUUGCAGUU-5' |
| 60 | 5'-UUGACGUUGAC-M-CAGUUGCAGUU-5' |
| 61 | 5'-UUGACGUUGAC-m-CAGUUGCAGUU-5' |
| 62 | 5'-GUGCCUGAUGA-X-AGUAGUCCGUG-5' |
| 63 | 5'-CCGAUGCCGAC-X-CAGCCGUAGCC-5' |
| 64 | 5'-CCGAUGCAUCG-X-GCUACGUAGCC-5' |
| 65 | 5'-GUGCCUGAUGA-Z-AGUAGUCCGUG-5' |

| | |
|----|---|
| 66 | 5'-GUGCCUGAUGA-M-AGUAGUCCGUG-5' |
| 67 | 5'-GUGCCUGAUGA-m-AGUAGUCCGUG-5' |
| 68 | 5'-CCGAUGCCGAC-Z-CAGCCGUAGCC-5' |
| 69 | 5'-CCGAUGCCGAC-M-CAGCCGUAGCC-5' |
| 70 | 5'-CCGAUGCCGAC-m-CAGCCGUAGCC-5' |
| 71 | 5'-CCGAUGCAUCG-Z-GCUACGUAGCC-5' |
| 72 | 5'-CCGAUGCAUCG-M-GCUACGUAGCC-5' |
| 73 | 5'-CCGAUGCAUCG-m-GCUACGUAGCC-5' |
| 74 | 5'-AGCACAAACUGU-X-UGUCAACACGA-5' |
| 75 | 5'-AAAAAAAAAA-X-AAAAAAAAAA-5' |
| 76 | 5'-YCACUGUUGAGA-X-AGAGUUGUCACY-5' |
| 77 | 5'-YACUGUUGACC-X-CCAGUUGUCAAY-5' |
| 78 | 5'-YCAACGACCUGU-X-UGUCCAGCAACY-5' |
| 79 | 5'-CACUG ₁ UUG ₁ AG ₁ A-X-AG ₁ AG ₁ UUG ₁ UCAC-5' |
| 80 | 5'-AACUG ₁ UUG ₁ ACC-X-CCAG ₁ UUG ₁ UCAA-5' |
| 81 | 5'-CAACG ₁ ACCUG ₁ U-X-UG ₁ UCCAG ₁ CAAC-5' |
| 82 | 5'-AUGCUGCGCUG-X-GUCGCGUCGUA-5' |
| 83 | 5'-AUGCUGCGCUG-Z-GUCGCGUCGUA-5' |
| 84 | 5'-AUGCUGCGCUG-m-GUCGCGUCGUA-5' |
| 85 | 5'-AACUGUUGACC-X-CCAGUUGUCAA-5' |
| 86 | 5'-CACUGUUGAGA-X-AGAGUUGUCAC-5' |
| 87 | 5'-GCACACUUGUU-X-UUGUUCACACG-5' |
| 88 | 5'-UGUUGUGUGAC-X-CAGUGUGUUGU-5' |
| 89 | 5'-CCGAUGCAUCG-X-GCUACGUAGCC-5' |
| 90 | 5'-AACGAACCGAC-X-CAGCCAAGCAA-5' |
| 91 | 5'-YCAACGACCUGU-X-UGUCCAGCAACY-5' |
| 92 | 5'-LCAACGACCUGU-X-UGUCCAGCAACL-5' |
| 93 | 5'-CGUUGUGAUGA-X-AGUAGUGUUGC-5' |
| 94 | 5'-ACGAUUGUGAC-X-CAGUGUUAGCA-5' |
| 95 | 5'-ACUUUGACGAU-X-UAGCAGUUUCA-5' |
| 96 | 5'-CGAUGCGAUGA-X-AGUAGCGUAGC-5' |
| 97 | 5'-ACGUCUGACGA-X-AGCAGUCUGCA-5' |
| 98 | 5'-AACUGCUGGAU-X-UAGGUCGUCAA-5' |
| 99 | 5'-UUGGACUCCAG-X-GACCUCAGGUU-5' |

| | |
|-----|---|
| 100 | 5'-UCGACUUCCAG-X-GACCUUCAGCU-5' |
| 101 | 5'-CCGACUUGGAC-X-CAGGUUCAGCC-5' |
| 102 | 5'-AAGACUGAACU-X-UCAAGUCAGAA-5' |
| 103 | 5'-YUG ₂ CUG ₂ CCUUUG ₂ -X-G ₂ UUUCCG ₂ UCG ₂ UY-5' |
| 104 | 5'-YUG ₂ CUG ₂ CCUUUG ₂ -M-G ₂ UUUCCG ₂ UCG ₂ UY-5' |
| 105 | 5'-YUG ₂ CUG ₂ CCUUUG ₂ -m-G ₂ UUUCCG ₂ UCG ₂ UY-5' |
| 106 | 5'-LUG ₂ CUG ₂ CCUUUG ₂ -X-G ₂ UUUCCG ₂ UCG ₂ UL-5' |
| 107 | 5'-LUG ₂ CUG ₂ CCUUUG ₂ -M-G ₂ UUUCCG ₂ UCG ₂ UL-5' |
| 108 | 5'-LUG ₂ CUG ₂ CCUUUG ₂ -m-G ₂ UUUCCG ₂ UCG ₂ UL-5' |
| 109 | 5'-L ₁ UG ₂ CUG ₂ CCUUUG ₂ -X-G ₂ UUUCCG ₂ UCG ₂ UL ₁ -5' |
| 110 | 5'-L ₁ UG ₂ CUG ₂ CCUUUG ₂ -M-G ₂ UUUCCG ₂ UCG ₂ UL ₁ -5' |
| 111 | 5'-L ₁ UG ₂ CUG ₂ CCUUUG ₂ -m-G ₂ UUUCCG ₂ UCG ₂ UL ₁ -5' |
| 112 | 5'-G ₁ UCCUUG ₁ CUUG ₁ -m-G ₁ UUCG ₁ UCCUG ₁ -5' |
| 113 | 5'-UUCUG ₁ CUUCUG ₁ -m-G ₁ UCUUCG ₁ UCUU-5' |
| 114 | 5'-UG ₁ CUG ₁ CCUUUG ₁ -M-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 115 | 5'-UG ₁ CCUUUG ₁ CUG ₁ -M-G ₁ UCG ₁ UUUCCG ₁ U-5' |
| 116 | 5'-UG ₁ CUG ₁ CUUCUG ₁ -M-G ₁ UCUUCG ₁ UCG ₁ U-5' |
| 117 | 5'-UG ₁ C ₁ UG ₁ C ₁ C ₁ UUUG ₁ -X-G ₁ UUUC ₁ C ₁ G ₁ UC ₁ G ₁ U-5' |
| 118 | 5'-UG ₁ C ₁ UG ₁ C ₁ UUC ₁ UG ₁ -X-G ₁ UC ₁ UUC ₁ G ₁ UC ₁ G ₁ U-5' |
| 119 | 5'-YUGCUGCUUCUG-6Eg-M-6Eg-GUCUUCGUCGUY-5' |
| 120 | 5'-UGGCUUGACGC-X-CGCAGUUCGGU-5' |
| 121 | 5'-UGCUGCUUGAA-X-AAGUUCGUCGU-5' |
| 122 | 5'-UGACGAUGAGU-X-UGAGUAGCAGU-5' |
| 123 | 5'-UGAUGAGGACU-X-UCAGGAGUAGU-5' |
| 124 | 5'-UGUUGAGGAAC-X-CAAGGAGUJUGU-5' |
| 125 | 5'-UGCGAGACUGC-X-CGUCAGAGCGU-5' |
| 126 | 5'-GUUGAACGACU-X-UCAGCAAGUUG-5' |
| 127 | 5'-UGACUGAUGAC-X-CAGUAGUCAGU-5' |
| 128 | 5'-UGUUGAACGAC-X-CAGCAAGUUGU-5' |
| 129 | 5'-UGAGCGUGAAC-X-CAAGUGCGAGU-5' |
| 130 | 5'-UG ₁ G ₁ CUUG ₁ ACG ₁ C-X-CG ₁ CAG ₁ UUCG ₁ G ₁ U-5' |
| 131 | 5'-UG ₁ ACG ₁ AUG ₁ AG ₁ U-X-UG ₁ AG ₁ UAG ₁ CAG ₁ U-5' |
| 132 | 5'-UG ₁ UUG ₁ AG ₁ G ₁ AAC-X-CAAG ₁ G ₁ AG ₁ UUG ₁ U-5' |
| 133 | 5'-UG ₁ UUG ₁ AACG ₁ AC-X-CAG ₁ CAAG ₁ UUG ₁ U-5' |

| | |
|-----|---|
| 134 | 5'-YUGCGAGACUGC-X-CGUCAGAGCGUY-5' |
| 135 | 5'-YUGAGCGUGAAC-X-CAAGUGCGAGUY-5' |
| 136 | 5'-YUUGAGCUGGAC-X-CAGGUCGAGUUY-5' |
| 137 | 5'-YGUUGAGGAACU-X-UCAAGGAGUUGY-5' |
| 138 | 5'-YUGAUGAAGCUU-X-UUCGAAGUAGUY-5' |
| 139 | 5'-YUUGACGAUGAG-X-GAGUAGCAGUUY-5' |
| 140 | 5'-YUUGUUGAACGA-X-AGCAAGUUGUUY-5' |
| 141 | 5'-YUUGAACGACUU-X-UUCAGCAAGUUY-5' |
| 142 | 5'-YUGAUGGAACGA-X-AGCAAGGUAGUY-5' |
| 143 | 5'-UG ₁ CUG ₁ CCUUUJUG ₁ -M-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 144 | 5'-UG ₁ CUG ₁ CCUUUJUG ₁ -Y-M-Y-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 145 | 5'-UG ₁ CUG ₁ CCUUUJUG ₁ -3Eg-M-3Eg-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 146 | 5'-UG ₁ CUG ₁ CCUUUJUG ₁ -4Eg-M-4Eg-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 147 | 5'-UG ₁ CUG ₁ CCUUUJUG ₁ -6Eg-M-6Eg-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 148 | 5'-YUGCUGCUUGUG-Y-M-Y-GUGUUCGUCGUY-5' |
| 149 | 5'-YUGCUGCUUGUG-3Eg-M-3Eg-GUGUUCGUCGUY-5' |
| 150 | 5'-YUGCUGCUUGUG-4Eg-M-4Eg-GUGUUCGUCGUY-5' |
| 151 | 5'-CAACGAACCCU-X-UCCCAAGCAAC-5' |
| 152 | 5'-UGCUGCUGCUG-X-GUCGUCGUCGU-5' |
| 153 | 5'-UGAACGUUGAA-X-AAGUUCGAAGU-5' |
| 154 | 5'-UGAACGUGAAC-X-CAAGUGCAAGU-5' |
| 155 | 5'-YUGC ₄ UGC ₄ UUGUG-X-GUGUUCG ₄ GUC ₄ GUY-5' |
| 156 | 5'-UG ₁ C ₄ UG ₁ C ₄ UUC ₄ UG ₁ -X-G ₁ UC ₄ UUC ₄ G ₁ UC ₄ G ₁ U-5' |
| 157 | 5'-YUG ₁ ACG ₁ AUG ₁ AG ₁ U-X-UG ₁ AG ₁ UAG ₁ CAG ₁ UY-5' |
| 158 | 5'-G ₁ UCCUUG ₁ CUUG ₁ -X ₁ -G ₁ UUCG ₁ UUCCUG ₁ -5' |
| 159 | 5'-UUCUG ₁ CUUCUG ₁ -X ₁ -G ₁ UCUUCG ₁ UCUU-5' |
| 160 | 5'-UG ₁ CCUUUJUG ₁ CUG ₁ -X ₂ -G ₁ UCG ₁ UUUCCG ₁ U-5' |
| 161 | 5'-UG ₁ CUG ₁ CUUCUG ₁ -X ₂ -G ₁ UCUUCG ₁ UCG ₁ U-5' |
| 162 | 5'-PUUCUG ₁ CUUCUG ₁ -m-G ₁ UCUUCG ₁ UCUUP-5' |
| 163 | 5'-PUG ₁ CCUUUJUG ₁ CUG ₁ -M-G ₁ UCG ₁ UUUCCG ₁ UP-5' |
| 164 | 5'-UG ₁ CUG ₁ CUUC ₁ UG ₁ -X ₁ -G ₁ UC ₁ UUCG ₁ UCG ₁ U-5' |
| 165 | 5'-UG ₁ CUG ₁ CUUC ₁ UG ₁ -X ₃ -G ₁ UC ₁ UUCG ₁ UCG ₁ U-5' |
| 166 | 5'-C ₂ UGAACG ₂ UUGU-X-UGUUC ₂ GAAGUC ₂ -5' |
| 167 | 5'-CU ₂ GAAGCU ₂ U ₂ GU ₂ -X-U ₂ GU ₂ U ₂ CGAAGU ₂ C-5' |

| | |
|-----|---|
| 168 | 5'-YAACUG ₂ UUG ₂ ACC-X-CCAG ₂ UUG ₂ UCAA _Y -5' |
| 169 | 5'-L ₁ UG ₂ G ₂ CUUG ₂ ACG ₂ C-X-CG ₂ CAG ₂ UUCG ₂ G ₂ UL ₁ -5' |
| 170 | 5'-YUGACCGUGACU-X-UGACGCUGACUY-5' |
| 171 | 5'-YUGACUGCGACU-X-UCAGCGUCAGUY-5' |
| 172 | 5'-YUGCGAACGCUU-X-UUCGCAAGCGUY-5' |
| 173 | 5'-YUGCGACUGACU-X ₃ -UCAGUCAGCGUY-5' |
| 174 | 5'-YUGCGCUGAACU-X ₃ -UCAAGUCGCGUY-5' |
| 175 | 5'-YUGCUGACGACU-X ₃ -UCAGCAGUCGUY-5' |
| 176 | 5'-YUGCUUGAACGC-X ₃ -CGCAAGUUCGUY-5' |
| 177 | 5'-YUUGCUGAACGC-X ₃ -CGCAAGUCGUUY-5' |
| 178 | 5'-UG ₁ CUG ₁ CCUUUG ₁ -Y-X-Y-G ₁ UUUCCG ₁ UCG ₁ U-5' |
| 179 | 5'-YUGUUGUGUGAC-X-CAGUGUGUUGUY-5' |
| 180 | 5'-YUGCUGCCUUUG-X-GUUUCCGUCGUY-5' |
| 181 | 5'-YUGCUGCUGCUG-X-GUCGUCGUCGUY-5' |
| 182 | 5'-YUGUUGUGUGAC-Z-CAGUGUGUUGUY-5' |
| 183 | 5'-EUUGAACGACUU-X-UUCAGCAAGUUE-5' |
| 184 | 5'-EUGUUGUGUGAC-X-CAGUGUGUUGUE-5' |
| 185 | 5'-EUGCUGCCUUUG-X-GUUUCCGUCGUE-5' |
| 186 | 5'-EUGCUGCUGCUG-X-GUCGUCGUCGUE-5' |
| 187 | 5'-U ₁ GCUGCUUGUG-X-GUGUUCGUCGU ₁ -5' |
| 188 | 5'-U ₁ GCU ₁ GCUUGUG-X-GUGUUCGU ₁ CGU ₁ -5' |
| 189 | 5'-U ₁ GCU ₁ GCU ₁ U ₁ GU ₁ G-X-GU ₁ GU ₁ U ₁ CGU ₁ CGU ₁ -5' |

G_1 = 7-deaza-rG; G_2 = ara-G; G_3 = 7-deaza-ara-G; C_1 = ara-C; C_2 = 2' -F-C; C_4 = 5-methyl-C;
 A_1 = ara-A; U_1 = ara-U; U_2 = 2'-F-U; M = cis,cis-cyclohexanetriol linker; m = cis,trans-cyclohexanetriol; Z = 1,3,5 -pentane triol linker; X = glycerol linker; X_1 = 1,2,4 -butane triol linker; X_2 = cyanuric acid; X_3 = isobutanetriol linker; Y = 1,3-propanediol; L = 1,5-pentanediol; L_1 = 1',2'-dideoxyribose; 6Eg = hexaethylene glycol linker; 3Eg = triethylene glycol linker; 4Eg = tetraethylene glycol linker; P = phosphorothioate; E = ethylene diol.

[00104] In a second aspect, the invention provides pharmaceutical formulations comprising a SIMRA compound according to the invention and a pharmaceutically acceptable carrier.

[00105] In a third aspect, the invention provides methods for generating TLR7 and/or TLR8 mediated immune responses in a vertebrate, such methods comprising administering to the vertebrate a SIMRA compound according to the invention. In some embodiments, the vertebrate is a mammal. In preferred embodiments, SIMRA compound is administered to a vertebrate in need of immune modulation.

[00106] In a fourth aspect, the invention provides methods for therapeutically treating a patient having a disease or disorder, such methods comprising administering to the patient a SIMRA compound according to the invention. In various embodiments, the disease or disorder to be treated is one in which an immune modulation may be desirable. For example, but not limited to, cancer, an autoimmune disorder, infectious disease, airway inflammation, inflammatory disorders, allergy, asthma, or a disease caused by a pathogen. Pathogens include bacteria, parasites, fungi, viruses, viroids and prions.

[00107] In a fifth aspect, the invention provides methods for preventing a disease or disorder, such methods comprising administering to the patient SIMRA compound according to the invention. In various embodiments, the disease or disorder to be prevented is one in which an immune modulation may be desirable. For example, but not limited to, cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, infectious disease, allergy, asthma, or a disease caused by a pathogen. Pathogens include bacteria, parasites, fungi, viruses, viroids, and prions.

[00108] In a sixth aspect the invention provides a method of preventing or treating a disorder, such methods comprises isolating cells capable of producing cytokines or chemokines including, but not limited to, immune cells, B cells, T-regulatory cells, B-cells, PBMCs, pDCs and lymphoid cells; culturing such cells under standard cell culture conditions, treating such cells ex vivo with a SIMRA such that the isolated cells produce or secrete increased levels of cytokines or chemokines, and administering or re-administering the treated cells to a patient in need of cytokine or chemokine therapy for the prevention or treatment of disease. This aspect of the invention would be in accordance with standard adoptive cellular immunotherapy techniques to produce activated immune cells.

[00109] In some embodiments of this aspect of the invention, the cells capable of producing cytokines or chemokines may be isolated from subjects with or without a disease or disorder. Such isolation may include identification and selection and could be performed using standard cell isolation procedures, including those set forth in the specific examples

below. Such isolated cells are cultured according to standard cell culturing procedures and using standard cell culture conditions, which may include the culturing procedures and conditions set forth in the specific examples below. In a further aspect of this embodiment of the invention, the isolated cells are cultured in the presence of at least one SIMRA, in an amount and for a time period sufficient to induce, increase or enhance the production and/or secretion of cytokines and/or chemokines as compared to the isolated cells cultured in the absence of such one or more SIMRA. Such time may be from minutes, to hours, to days. Such isolated, SIMRA-treated cells may find use following re-administration to the donor or administration to a second histologically compatible patient, wherein such donor or second patient are in need of induced, increased or enhanced production and/or secretion of cytokines and/or chemokines. For example, re-administration to a donor or administration to a second patient having cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, infectious disease, allergy, asthma, or a disease caused by a pathogen. Such re-administration or administration may be accomplished using various modes, including catheter or injection administration or any other effective route. This aspect of the invention may also find use in patients who may have a limited or incomplete ability to mount an immune response or are immune compromised (e.g. patient infected with HIV and bone marrow transplant patients). This aspect of the invention may also find use in combination with SIMRA administration to the patient administered or re-administered the isolated, SIMRA-treated cells.

[00110] In any of the methods according to the invention, the SIMRA compound can variously act by producing direct immune modulatory effects alone or in combination with any other agent useful for treating or preventing the disease or condition that does not diminish the immune modulatory effect of the SIMRA compound. In any of the methods according to the invention, the agent(s) useful for treating or preventing the disease or condition includes, but is not limited to, vaccines, antigens, antibodies, preferably monoclonal antibodies, cytotoxic agents, allergens, antibiotics, siRNA, microRNA, antisense oligonucleotides, TLR agonist (e.g. agonists of TLR9 and/or agonists of TLR7 and/or agonists of TLR8), chemotherapeutic agents (both traditional chemotherapy and modern targeted therapies), targeted therapeutic agents, activated cells, peptides, proteins, gene therapy vectors, peptide vaccines, protein vaccines, DNA vaccines, adjuvants, and co-stimulatory molecules (e.g. cytokines, chemokines, protein ligands, trans-activating factors, peptides or peptides comprising modified amino acids), or combinations thereof. For

example, in the treatment of cancer, it is contemplated that the SIMRA compound may be administered in combination with one or more chemotherapeutic compound, targeted therapeutic agent and/or monoclonal antibody. Alternatively, the agent can include DNA vectors encoding for antigen or allergen. Alternatively, the SIMRA compounds can be administered in combination with other adjuvants to enhance the specificity or magnitude of the immune response to the SIMRA compound.

[00111] In any of the methods according to the invention, administration of SIMRA compound, alone or in combination with any other agent, can be by any suitable route, including, without limitation, parenteral, mucosal delivery, oral, sublingual, transdermal, topical, inhalation, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, by gene gun, dermal patch or in eye drop or mouthwash form. Administration of the therapeutic compositions of SIMRA compound can be carried out using known procedures using a pharmaceutically effective amount and for periods of time effective to reduce symptoms or surrogate markers of the disease. For example, a pharmaceutically effective amount of a SIMRA compound for treating a disease and/or disorder could be that amount necessary to alleviate or reduce the symptoms, or delay or ameliorate a tumor, cancer, or bacterial, viral or fungal infection. A pharmaceutically effective amount for use as a vaccine adjuvant could be that amount useful for boosting a subject's immune response to a vaccine or antigen. In the context of administering a composition that modulates an immune response to a co-administered antigen, a pharmaceutically effective amount of a SIMRA compound and antigen is an amount sufficient to achieve the desired modulation as compared to the immune response obtained when the antigen is administered alone. The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular oligonucleotide being administered, the size of the subject, or the severity of the disease or condition. One of ordinary skill in the art can empirically determine the pharmaceutically effective amount of a particular oligonucleotide without necessitating undue experimentation.

[00112] When administered systemically, the therapeutic composition is preferably administered at a sufficient dosage to attain a blood level of SIMRA compound from about 0.0001 micromolar to about 10 micromolar. For localized administration, much lower concentrations than this may be effective, and much higher concentrations may be tolerated. Preferably, a total dosage of SIMRA compound ranges from about 0.001 mg per patient per day to about 200 mg per kg body weight per day. It may be desirable to administer

simultaneously, or sequentially a therapeutically effective amount of one or more of the therapeutic compositions of the invention to an individual as a single treatment episode.

[00113] The SIMRA compound may optionally be linked to one or more allergens and/or antigens (self or foreign), an immunogenic protein or peptide, such as keyhole limpet hemocyanin (KLH), cholera toxin B subunit, or any other immunogenic carrier protein. SIMRA can also be used in combination with other compounds (e.g. adjuvants) including, without limitation, TLR agonists (e.g. TLR2 agonists and TLR9 agonists), Freund's incomplete adjuvant, KLH, monophosphoryl lipid A (MPL), alum, and saponins, including QS-21 and imiquimod, or combinations thereof.

[00114] The methods according to this aspect of the invention are useful for model studies of the immune system. The methods are also useful for the prophylactic or therapeutic treatment of human or animal disease. For example, the methods are useful for pediatric and veterinary vaccine applications.

[00115] The examples below are intended to further illustrate certain exemplar embodiments of the invention, and are not intended to limit the scope of the invention.

EXAMPLES

Example 1.

Immune Modulatory Oligoribonucleotide Synthesis

[00116] The immune modulatory oligoribonucleotides were chemically synthesized using phosphoramidite chemistry on automated DNA/RNA synthesizer. N-acetyl protected (Except U) 2'-O-TBDMS RNA monomers, A, G, C and U, were purchased from Sigma-Aldrich. 7-deaza-G, inosine was purchased from ChemGenes Corporation. 0.25M 5-ethylthio-1H-tetrazole, PAC- anhydride Cap A and Cap B were purchased from Glen Research. 3% trichloroacetic acid (TCA) in dichloromethane (DCM) and 5% 3H-1,2-Benzodithiole-3-one-1,1-dioxide (Beaucage reagent) were made in house.

[00117] Immune modulatory oligoribonucleotides were synthesized at 1-2 μ M scale using a standard RNA synthesis protocol.

Cleavage and base deprotection

[00118] Immune modulatory oligoribonucleotides were cleaved from solid support and the protecting groups of exo-cyclic-amines were removed in methylamine and ammonium hydroxide solution. The resulting solution was dried completely in a SpeedVac.

IE HPLC Purification

[00119] Immune modulatory oligoribonucleotides were purified by ion exchange HPLC.

Using Dionex DNAPac 100 column. Crude immune modulatory oligoribonucleotide solution was injected into HPLC. Above gradient is performed and the fractions were collected. All fractions containing more than 90% desired product were mixed, and then the solution was concentrated to almost dry by RotoVap. RNase-free water was added to make final volume of 10ml.

C-18 Reversed Phase Desalting

[00120] tC-18 Sep-Pak cartridge purchased from Waters was washed by passing with 10ml of acetonitrile followed by 10 ml of 0.5 M sodium acetate through the cartridge. Then 10 ml of immune modulatory oligoribonucleotide solution was loaded on to the cartridge. Then 15 ml of water was used to wash out the salt. The immune modulatory oligoribonucleotide was finally eluted using 1 ml of 50% acetonitrile in water. The solution was placed in SpeedVac for 30 minutes. The remaining solution was filtered through a 0.2 micron filter and then was lyophilized. The solid was then re-dissolved in RNase free water

to make the desired concentration. The final solution was stored below 0 °C. Oligoribonucleotides were analyzed for purity by Capillary Electrophoresis, Ion Exchange HPLC and PAGE analysis, and for molecular mass by MALDI-ToF mass spectrometry.

Example 2.

Protocols for assays with HEK293 cells expressing TLRs

[00121] HEK293 or HEK293XL /human TLR7 or HEK293 or HEK293XL /human TLR8 cells (Invivogen, San Diego, CA) were cultured in 48-well plates in 250 µl/well DMEM supplemented with 10% heat-inactivated FBS in a 5% CO₂ incubator.

Reporter gene transformation

[00122] HEK293 or HEK293XL cells stably expressing human TLR7 or TLR8 (Invivogen, San Diego, CA) were cultured in 48-well plates in 250 µl/well DMEM supplemented with 10% heat-inactivated FBS in a 5% CO₂ incubator. At 80% confluence, cultures were transiently transfected with 400 ng/ml of SEAP (secreted form of human embryonic alkaline phosphatase) reporter plasmid (pNifty2-Seap) (Invivogen) in the presence of 4 µl/ml of lipofectamine (Invitrogen, Carlsbad, CA) in culture medium. Plasmid DNA and lipofectamine were diluted separately in serum-free medium and incubated at room temperature for 5 minutes. After incubation, the diluted DNA and lipofectamine were mixed and the mixtures were incubated at room temperature for 20 minutes. Aliquots of 25 µl of the DNA/lipofectamine mixture containing 100 ng of plasmid DNA and 1 µl of lipofectamine were added to each well of the cell culture plate, and the cultures were continued for 4 hours.

IMO-treatment

[00123] After transfection, medium was replaced with fresh culture medium. The HEK293 or HEK29XL cells expressing human TLR7 or TLR8 were stimulated with 0, 20, 50, 100, 150, 200, or 300 µg/ml of agonists of TLR7 or TLR8, SIMRAs, and the cultures were continued for 18 hours - 20 hours. At the end of SIMRA treatment, 30 µl of culture supernatant was taken from each treatment and used for SEAP assay following manufacturer's protocol (Invivogen).

SEAP (secreted form of human embryonic alkaline phosphatase) assay

[00124] Briefly, culture supernatants were incubated with p-nitrophenyl phosphate substrate and the yellow color generated was measured by a plate reader at 405 nm. The data are shown as fold increase in NF- κ B activity over PBS control. (Putta MR et al, Nucleic Acids Res., 2006, 34:3231-8).

Example 3.

Human cell culture protocols

Human PBMC isolation

[00125] Peripheral blood mononuclear cells (PBMCs) from freshly drawn, healthy volunteer blood (CBR Laboratories, Boston, MA) were isolated by Ficoll density gradient centrifugation method (Histopaque-1077, Sigma).

Human pDC isolation

[00126] Peripheral blood mononuclear cells (PBMCs) from freshly drawn healthy volunteer blood (CBR Laboratories, Boston, MA) were isolated by Ficoll density gradient centrifugation method (Histopaque-1077, Sigma). pDCs were isolated from PBMCs by positive selection using the BDCA4 cell isolation kits (Miltenyi Biotec) according to the manufacturer's instructions.

Human mDC isolation

[00127] Peripheral blood mononuclear cells (PBMCs) from freshly drawn healthy volunteer blood (CBR Laboratories, Boston, MA) were isolated by Ficoll density gradient centrifugation method (Histopaque-1077, Sigma). Myeloid dendritic cells (mDCs) were isolated from PBMCs by positive selection using the BDCA4 cell isolation kits (Miltenyi Biotec) according to the manufacturer's instructions.

Multiplex Cytokine Assays

[00128] Human PBMCs were plated in 48-well plates using 5×10^6 cells/ml. pDCs were plated in 96-well dishes using 1×10^6 cells/ml. The SIMRAs dissolved in DPBS (pH 7.4; Mediatech) were added to a final concentration of 20, 50, 100, 200 or 300 μ g/ml or as indicated in the figures to the cell cultures. The cells were then incubated at 37 °C for 24 hr and the supernatants were collected for luminex multiplex or ELISA assays. The experiments were performed in triplicate wells. The levels of IFN- α , IL-6, or TNF- α were measured by sandwich ELISA. The required reagents, including cytokine antibodies and standards, were purchased from PharMingen.

[00129] Luminex multiplex assays were performed using Biosource human multiplex cytokine assay kits on Luminex 100/200 instrument and the data were analyzed using StarStation software supplied by Applied Cytometry Systems (Sacramento, CA).

Example 4

***In vivo* cytokine secretion in mouse model treated with TLR9 agonist compounds**

[00130] C57BL/6 mice and BALB/c mice, 5-6 weeks old, were obtained from Taconic Farms, Germantown, NY and maintained in accordance with Idera Pharmaceutical's IACUC approved animal protocols. Mice (n=3) were injected subcutaneously (s.c) with individual stabilized immune modulatory RNA-based oligonucleotides from Table 3 at 25 mg/kg (single dose). Serum was collected by retro-orbital bleeding 2 hr after immune modulatory oligonucleotide administration and cytokine and chemokine levels were determined by sandwich ELISA or Luminex multiplex assays. The results are shown in Figures 8A, 8B, 9A and 9B and demonstrate that *in vivo* administration of SIMRA oligonucleotides according to the invention generates unique cytokine and chemokine profiles. All reagents, including cytokine and chemokine antibodies and standards were purchased from PharMingen. (San Diego, CA).

Example 5

Serum stability assay

[00131] Approximately 0.5 OD of exemplar SIMRA compounds from Table 3 was individually incubated in 1% human serum in PBS for 30 minute at 37 °C. Following 30 minutes of incubation in 1% human serum, the SIMRA compound was analyzed on anion-exchange HPLC to determine the percentage of full-length SIMRA compound that remained as compared to the amount of SIMRA compound present before serum treatment. The results are shown in Figures 10A — 10H and demonstrate that chemical modifications according to the invention made to RNA-based compounds can enhance their stability.

EQUIVALENTS

[00132] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention and appended claims.

2008279509 08 Apr 2011

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A stabilized immune modulatory RNA (SIMRA) compound selected from the group consisting of
5'-UG₁CUG₁CUUUUG₁-X-G₁UUUUCG₁UCG₁U-5';
5'-UG₁CUG₁CCUUUG₁-Z-G₁UUUCCG₁UCG₁U-5';
5'-UG₂CUGCUUGUG-X-GUGUUCGUCG₂U-5';
5'-UG₂CUG₂CUUG₂UG₂-X-G₂UG₂UUCG₂UCG₂U-5';
5'-UG₂CUG₂CUUG₂UG₂-M-G₂UG₂UUCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂U-5';
5'-UG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂U-5';
5'-UGC₁UGC₁UUGUG-X-GUGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁UUGUG-M-GUGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁UUC₁UG-X-GUC₁UUC₁GUC₁GU-5';
15 5'-UGC₁UGC₁C₁UUUG-M-GUUUC₁C₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-m-GUUUC₁C₁GUC₁GU-5';
5'-UGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGU-5';
5'-YUGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGU-Y-5';
5'-UGUUGUGUGA₁C-X-CA₁GUGUGUJUGU-5';
20 5'-UG₂CUG₂CUUG₂UG₂-m-G₂UG₂UUCG₂UCG₂U-5';
5'-UGC₁UGC₁UUGUG-m-GUGUUC₁GUC₁GU-5';
5'-UGC₁UGC₁C₁UUUG-X-GUUUC₁C₁GUC₁GU-5';
5'-YUGACGACGCUU-X-UUCGCAGCAGUY-5';
5'-YUGACGACUGCU-X-UCGUCAGCAGUY-5';
25 5'-YUGACGACUJUGC-X-CGUUCAGCAGUY-5';
5'-YUGCGCGAACUU-X₃-UUCAAGCGCGUY-5';
5'-AG₁UG₁UUUUCUG₁-X-G₁UCUUUUG₁UG₁A-5';
5'-UG₁UUG₁UUUG₁UG₁-X-G₁UG₁UUUG₁UUG₁U-5';
5'-UG₁AUG₁AAG₁CUU-X-UUCG₁AAG₁UAG₁U-5';
30 5'-UGUUCGAACAC-X-CACAAGCUJUGU-5';
5'-UUGACGUUGAC-X-CAGUUGCAGUU-5';
5'-UUGACGUUGAC-Z-CAGUUGCAGUU-5';
5'-UUGACGUUGAC-M-CAGUUGCAGUU-5';

2008279509 08 Apr 2011

5'-UUGACGUUGAC-m-CAGUUGCAGUU-5';
5'-GUGCCUGAUGA-X-AGUAGUCCGUG-5';
5'-CCGAUGCCGAC-X-CAGCCGUAGCC-5';
5'-CCGAUGCAUCG-X-GCUACGUAGCC-5';
5'-GUGCCUGAUGA-Z-AGUAGUCCGUG-5';
5'-GUGCCUGAUGA-M-AGUAGUCCGUG-5';
5'-GUGCCUGAUGA-m-AGUAGUCCGUG-5';
5'-CCGAUGCCGAC-Z-CAGCCGUAGCC-5';
5'-CCGAUGCCGAC-M-CAGCCGUAGCC-5';
5'-CCGAUGCCGAC-m-CAGCCGUAGCC-5';
5'-CCGAUGCAUCG-Z-GCUACGUAGCC-5';
5'-CCGAUGCAUCG-M-GCUACGUAGCC-5';
5'-CCGAUGCAUCG-m-GCUACGUAGCC-5';
5'-AAAAAAAAAAA-X-AAAAAAAAAAA-5';
5'-YCACUGUUGAGA-X-AGAGUUGUCACY-5';
5'-YAACUGUUGACC-X-CCAGUUGUCAAY-5';
5'-YCAACGACCUGU-X-UGUCCAGCAACY-5';
5'-CACUG₁UUG₁AG₁A-X-AG₁AG₁UUG₁UCAC-5';
5'-AACUG₁UUG₁ACC-X-CCAG₁UUG₁UCAA-5';
5'-CAACG₁ACCUG₁U-X-UG₁UCCAG₁CAAC-5';
5'-AUGCUGCGCUG-X-GUCGCGUCGUA-5';
5'-AUGCUGCGCUG-Z-GUCGCGUCGUA-5';
5'-AUGCUGCGCUG-m-GUCGCGUCGUA-5';
5'-CCGAUGCAUCG-X-GCUACGUAGCC-5';
5'-AACGAACCGAC-X-CAGCCAAGCAA-5';
5'-YCAACGACCUGU-X-UGUCCAGCAACY-5';
5'-LCAACGACCUGU-X-UGUCCAGCAACL-5';
5'-CGUUGUGAUGA-X-AGUAGUGUUGC-5';
5'-ACGAUUGUGAC-X-CAGUGUUAGCA-5';
5'-ACUUUGACGAU-X-UAGCAGUUUCA-5';
5'-CGAUGCGAUGA-X-AGUAGCGUAGC-5';
5'-ACGUCUGACGA-X-AGCAGUCUGCA-5';
5'-AACUGCUGGAU-X-UAGGUCGUCAA-5';
5'-UUGGACUCCAG-X-GACCUCAGGUU-5';

2008279509 08 Apr 2011

5'-UCGACUUCCAG-X-GACCUUCAGCU-5';
 5'-CCGACUUGGAC-X-CAGGUUCAGCC-5';
 5'-AAGACUGAACU-X-UCAAGUCAGAA-5';
 5'-YUG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UY-5';
 5'-YUG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UY-5';
 5'-YUG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UY-5';
 5'-LUG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UL-5';
 5'-LUG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UL-5';
 5'-LUG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UL-5';
 5'-L₁UG₂CUG₂CCUUUG₂-X-G₂UUUCCG₂UCG₂UL₁-5';
 5'-L₁UG₂CUG₂CCUUUG₂-M-G₂UUUCCG₂UCG₂UL₁-5';
 5'-L₁UG₂CUG₂CCUUUG₂-m-G₂UUUCCG₂UCG₂UL₁-5';
 5'-G₁UCCUUG₁CUUG₁-m-G₁UUCG₁UUCCUG₁-5';
 5'-UUCUG₁CUUCUG₁-m-G₁UCUUCG₁UCUU-5';
 5'-UG₁CUG₁CCUUUG₁-M-G₁UUUCCG₁UCG₁U-5';
 5'-UG₁CCUUUG₁CUG₁-M-G₁UCG₁UUUCCG₁U-5';
 5'-UG₁CUG₁CUUCUG₁-M-G₁UCUUCG₁UCG₁U-5';
 5'-UG₁C₁UG₁C₁C₁UUUG₁-X-G₁UUUC₁C₁G₁UC₁G₁U-5';
 5'-UG₁C₁UG₁C₁UUC₁UG₁-X-G₁UC₁UUC₁G₁UC₁G₁U-5';
 5'-YUGCUGCUUCUG-6Eg-M-6Eg-GUCUUCGUCGU-5';
 5'-UGGUUGACGC-X-CGCAGUUCGGU-5';
 5'-UGCUGCUUGAA-X-AAGUUCGUCGU-5';
 5'-UGACGAUGAGU-X-UGAGUAGCAGU-5';
 5'-UGAUGAGGACU-X-UCAGGAGUAGU-5';
 5'-UGUUGAGGAAC-X-CAAGGAGUUGU-5';
 5'-UGCGAGACUGC-X-CGUCAGAGCGU-5';
 5'-GUUGAACGACU-X-UCAGCAAGUUG-5';
 5'-UGACUGAUGAC-X-CAGUAGUCAGU-5';
 5'-UGUUGAACGAC-X-CAGCAAGUUGU-5';
 5'-UGAGCGUGAAC-X-CAAGUGCGAGU-5';
 5'-UG₁G₁CUUG₁ACG₁C-X-CG₁CAG₁UUCG₁G₁U-5';
 5'-UG₁ACG₁AUG₁AG₁U-X-UG₁AG₁UAG₁CAG₁U-5';
 5'-UG₁UUG₁AG₁G₁AAC-X-CAAG₁G₁AG₁UUG₁U-5';
 5'-UG₁UUG₁AACG₁AC-X-CAG₁CAAG₁UUG₁U-5';

2008279509 08 Apr 2011

5'-YUGCGAGACUGC-X-CGUCAGAGCGUY-5';
 5'-YUGAGCGUGAAC-X-CAAGUGCGAGUY-5';
 5'-YUUGAGCUGGAC-X-CAGGUCGAGUUY-5';
 5'-YGUUGAGGAACU-X-UCAAGGAGUUGY-5';
 5'-YUGAUGAAGCUU-X-UUCGAAGUAGUY-5';
 5'-YUUGACGAUGAG-X-GAGUAGCAGUUY-5';
 5'-YUUGUUGAACGA-X-AGCAAGUJUGUUY-5';
 5'-YUUGAACGACUU-X-UUCAGCAAGUUY-5';
 5'-YUGAUGGAACGA-X-AGCAAGGUAGUY-5';
 5'-UG₁CUG₁CCUUUG₁-M-G₁UUUCCG₁UCG₁U-5';
 5'-UG₁CUG₁CCUUUG₁-Y-M-Y-G₁UUUCCG₁UCG₁U-5';
 5'-UG₁CUG₁CCUUUG₁-3Eg-M-3Eg-G₁UUUCCG₁UCG₁U-5';
 5'-UG₁CUG₁CCUUUG₁-4Eg-M-4Eg-G₁UUUCCG₁UCG₁U-5';
 5'-UG₁CUG₁CCUUUG₁-6Eg-M-6Eg-G₁UUUCCG₁UCG₁U-5';
 5'-YUGCUGCUCUUGUG-Y-M-Y-GUGUUCGUCGUY-5';
 5'-YUGCUGCUCUUGUG-3Eg-M-3Eg-GUGUUCGUCGUY-5';
 5'-YUGCUGCUCUUGUG-4Eg-M-4Eg-GUGUUCGUCGUY-5';
 5'-UGAACGUUGAA-X-AAGUUCGAAGU-5';
 5'-UGAACGUGAAC-X-CAAGUGCAAGU-5';
 5'-YUGC₄UGC₄UUGUG-X-GUGUUC₄GUC₄GUY-5';
 5'-UG₁C₄UG₁C₄UUC₄UG₁-X-G₁UC₄UUC₄G₁UC₄G₁U-5';
 5'-YUG₁ACG₁AUG₁AG₁U-X-UG₁AG₁UAG₁CAG₁UY-5';
 5'-G₁UCCUUG₁CUUG₁-X₁-G₁UUCG₁UUCCUG₁-5';
 5'-UUCUG₁CUUCUG₁-X₁-G₁UCUUCG₁UCUU-5';
 5'-UG₁CCUUUG₁CUG₁-X₂-G₁UCG₁UUUCCG₁U-5';
 5'-UG₁CUG₁CUUCUG₁-X₂-G₁UCUUCG₁UCG₁U-5';
 5'-PUUCUG₁CUUCUG₁-m-G₁UCUUCG₁UCUUP-5';
 5'-PUG₁CCUUUG₁CUG₁-M-G₁UCG₁UUUCCG₁UP-5';
 5'-UG₁CUG₁CUUC₁UG₁-X₁-G₁UC₁UUCG₁UCG₁U-5';
 5'-UG₁CUG₁CUUC₁UG₁-X₃-G₁UC₁UUCG₁UCG₁U-5';
 5'-C₂UGAACG₂UUGU-X-UGUUC₂GAAGUC₂-5';
 5'-CU₂GAAGCU₂U₂GU₂-X-U₂GU₂U₂CGAACGU₂C-5';
 5'-YAACUG₂UUG₂ACC-X-CCAG₂UUG₂UCAAY-5';
 5'-L₁UG₂G₂CUUG₂ACG₂C-X-CG₂CAG₂UUCG₂G₂UL₁-5';

08 Apr 2011

2008279509

15

5'-YUGACUGCGACU-X-UCAGCGUCAGUY-5';
 5'-YUGCGAACGCUU-X-UUCGCAAGCGUY-5';
 5'-YUGCGACUGACU-X₃-UCAGUCAGCGUY-5';
 5'-YUGCGCUGAACU-X₃-UCAAGUCGCGUY-5';
 5'-YUGCUGACGACU-X₃-UCAGCAGUCGUY-5';
 5'-YUGCUUGAACGC-X₃-CGCAAGUUCGUY-5';
 5'-YUUGCUGAACGC-X₃-CGCAAGUCGUUY-5';
 5'-UG₁CUG₁CCUUUG₁-Y-X-Y-G₁UUUCCG₁UCG₁U-5';
 5'-YUGUUGUGUGAC-X-CAGUGUGUUGUY-5';
 5'-YUGCUGCCUUUG-X-GUUUCCGUCGUY-5';
 5'-YUGUUGUGUGAC-Z-CAGUGUGUUGUY-5';
 5'-EUUGAACGACUU-X-UUCAGCAAGUUE-5';
 5'-EUGUUGUGUGAC-X-CAGUGUGUUGUE-5';
 5'-EUGCUGGCCUUUG-X-GUUUCCGUCGUE-5';
 5'-EUGCUGCUGCUG-X-GUCGUCGUCGUE-5';
 5'-U₁GCUGCUUGUG-X-GUGUUCGUCGU₁-5';
 5'-U₁GCU₁GCUUGUG-X-GUGUUCGU₁CGU₁-5';
 5'-U₁GCU₁GCU₁U₁GU₁G-X-GU₁GU₁U₁CGU₁CGU₁-5';
 5'-UG₁CUG₁CUUCUG₁-6Eg-M-6Eg-G₁UCUUCG₁UCG₁U-5';
 5'-LUGCUGCCUUUG-m-GUUUCCGUCGUL-5';
 5'-YUUGACGUUGAC-m-CAGUUGCAGUUY-5';
 5'-YGUGCCUGAUGA-X-AGUAGUCCGUGY-5';
 5'-AUGCUGCGCUG-M-GUCGCGUCGUA-5';
 5'-UG₂CUG₂CUUG₂UG₂-m-G₂UG₂UUCG₂UCG₂U-5';
 5'-UGC₁UGC₁UUGUG-m-GUGUUC₁GUC₁GU-5';
 5'-UGC₁UGC₁C₁UUUG-X-GUUUC₁C₁GUC₁GU-5';
 5'-GAUUGUGACGU-X-UGCAGUGUUAG-5';
 5'-CUGAACGUUGU-X-UGUUCGAAGUC-5';
 5'-UG₂CUG₂CUUG₂UG₂-M-G₂UG₂UUCG₂UCG₂U-5';
 5'-UG₃CUG₃CUUCUG₃-X-G₃UCUUCG₃UCG₃U-5';
 5'-UG₃CUG₃CCUUUG₃-m-G₃UUUCCG₃UCG₃U-5';
 5'-YUGACGAUGAGU-X-UGAGUAGCAGUY-5';
 5'-UGCUGCU₁U₁CU₁G-X-GU₁CU₁U₁CGUCGU-5';
 5'-UGC₁UGC₁C₁UUUG-m-GUUUC₁C₁GUC₁GU-5';

08 Apr 2011
2008279509

5' -YUGCUGCUUGAA-X-AAGUUCGUCGUY-5';
5' -YUUGACUGAUGA-X-AGUAGUCAGUUY-5';
5' -UGCUGCUUUUG-X-GUUUUCGUCGU-5'; and
5' -UGUUGUUUGUG-X-GUGUUUGUUGU-5',
wherein $G_1 = 7$ -deaza-rG; $G_2 =$ ara-G; $G_3 = 7$ -deaza-ara-G; $C_1 =$ ara-C; $C_2 = 2'$ -F-C; $C_4 =$ 5-methyl-C; $A_1 =$ ara-A; $U_1 =$ ara-U; $U_2 = 2'$ -F-U; M = cis,cis-cyclohexanetriol linker; m = cis,trans-cyclohexanetriol; Z = 1,3,5 -pentane triol linker; X = glycerol linker; $X_1 =$ 1,2,4 -butane triol linker; $X_2 =$ cyanuric acid; $X_3 =$ isobutanetriol linker; Y = 1,3-propanediol; L = 1,5-pentanediol; $L_1 = 1',2'$ -dideoxyribose; 6Eg = hexaethylene glycol linker; 3Eg = triethylene glycol linker; 4Eg = tetraethylene glycol linker; P = phosphorothioate; E = ethylene diol.

2. A composition comprising the SIMRA compound of claim 1 and a physiologically acceptable carrier.
3. A method for generating an immune response in a vertebrate comprising administering a SIMRA compound according to claim 1 or a composition according to claim 2.
4. A method for therapeutically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial comprising administering a SIMRA compound according to claim 1 or a composition according to claim 2.
5. The method according to claim 4 wherein the disease or disorder is cancer, an autoimmune disorder, airway inflammation, inflammatory disorder, infectious disease, skin disorder, allergy, asthma or a disease caused by a pathogen.
6. The method according to claim 4 or 5, further comprising administering one or more chemotherapeutic compounds or a targeted therapeutic agent.
7. The method according to claim 4 or 5, further comprising administering an antibody.
8. The method according to any one of claims 4 to 7, further comprising administering a vaccine selected from the group consisting of a DNA vaccine, a protein vaccine and a peptide vaccine.
9. The method according to any one of claims 4 to 7, further comprising administering an antigen.

08 Apr 2011
2008279509

10. The method according to any one of claims 4 to 9, further comprising administering an adjuvant.
11. A method for prophylactically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial comprising administering a SIMRA compound according to claim 1 or a composition according to claim 2.
12. The method according to claim 11 wherein the disease or disorder is cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, infectious disease, skin disorders, allergy, asthma or a disease caused by a pathogen in a vertebrate.
13. The method according to claim 11 or 12, further comprising administering one or more chemotherapeutic compounds or a targeted therapeutic agent.
14. The method according to claim 11 or 12, further comprising administering an antibody.
15. The method according to any one of claims 11 to 14, further comprising administering a vaccine selected from the group consisting of DNA vaccine, a protein vaccine or a peptide vaccine.
16. The method according to any one of claims 12 to 14, further comprising administering an antigen.
17. The method according to any one of claims 12 to 16, further comprising administering an adjuvant.
18. The method according to claim 10 or 17, wherein the adjuvant is an alum based adjuvant.
19. Use of a SIMRA compound according to claim 1 or a composition according to claim 2 for the preparation of a medicament for generating an immune response in a vertebrate.
20. Use of a SIMRA compound according to claim 1 or a composition according to claim 2 for the preparation of a medicament for therapeutically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial.
21. Use of a SIMRA compound according to claim 1 or a composition according to claim 2 for the preparation of a medicament for prophylactically treating a vertebrate having a disease or disorder where modulating an immune response would be beneficial.

08 Apr 2011

2008279509

22. A use according to claim 20 or 21 wherein the disease or disorder is cancer, an autoimmune disorder, airway inflammation, inflammatory disorder, infectious disease, skin disorder, allergy, asthma or a disease caused by a pathogen.
23. A SIMRA compound according to claim 1, substantially as hereinbefore described.
- 5 24. A composition according to claim 2, substantially as hereinbefore described.
25. A method according to any one of claims 3, 4 or 11, substantially as hereinbefore described.
26. Use according to any one of claims 19 to 21, substantially as hereinbefore described.

Parallel Synthesis of SIMRA Compounds





Figure 1

Activation of HEK293 cells expressing hTLR8**Figure 2A.**




Figure 2B.




Figure 2C.




Figure 2D.




Figure 2E.




Figure 2F.




Figure 2G.




Figure 2H.




Figure 2I.




Figure 2J.





Figure 2K.

**Figure 2L.**




Figure 2M.




Figure 3A.




Figure 3B.




Figure 3C.




Figure 4A.




Figure 4B.







Figure 4C.




Figures 4D – 4I.




Figures 4J – 4O.



Figures 4P – 4U.



Figures 4V – 4AA.

Cytokines/chemokines induced by TLR7/8 agonists in human pDC cultures**Figure 5A.**




Figure 5B.




Figure 5C.





Figure 5D.

Cytokines/chemokines induced by TLR7/8 agonists in human pDC cultures**Figure 6A.**




Figure 6B.




Figure 6C.




Figure 6D.




Figure 6E.





Figure 6F.

Cytokines/chemokines induced by TLR7/8 agonists in human mDC cultures

**Figure 7A.**




Figure 7B.




Figure 7C.




Figure 8A.




Figure 8B.




Figure 9A.




Figure 9B.




Figure 9C.




Figure 9D.




Figure 9E.




Figure 9F.




Figure 10A.




Figure 10B.




Figure 10C.




Figure 10D.

Before incubation in serum




Figure 10E.




Figure 10F.




Figure 10G.




Figure 10H.