



(22) Date de dépôt/Filing Date: 2002/09/17

(41) Mise à la disp. pub./Open to Public Insp.: 2003/03/20

(30) Priorité/Priority: 2001/09/20 (01 121 780.9) EP

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/55, C12N 9/80, C12Q 1/34

(71) Demandeur/Applicant:
F. HOFFMANN-LA ROCHE AG, CH


(72) Inventeurs/Inventors:
SHAO, ZHIXIN, DE;
SCHMUCK, RAINER, DE;
KRATZSCH, PETER, DE;
KENKLIES, JANET, DE;
WEISSER, HARALD, DE

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : VARIANTS D'UNE CREATINASE DE TYPE ERWINIA

(54) Title: VARIANTS OF AN ERWINIA-TYPE CREATINASE

Expression vector comprising a creatinase gene



(57) **Abrégé/Abstract:**

The present invention relates to mutated genes encoding variants of an Erwinia- type creatinase (EC 3.5.3.3, alternative name: creatine amidinohydrolase), to the creatinase variants encoded by these genes, and to different applications of these variants of creatinase, particularly to their use for determining the creatinine and/or creatine concentration in a sample.

Abstract

The present invention relates to mutated genes encoding variants of an Erwinia-type creatinase (EC 3.5.3.3, alternative name: creatine amidinohydrolase), to the creatinase variants encoded by these genes, and to different applications of these variants of creatinase, particularly to their use for determining the creatinine and/or creatine concentration in a sample.

Variants of an Erwinia-type creatinase

The present invention relates to mutated genes encoding variants of an Erwinia-type creatinase (EC 3.5.3.3, alternative name: creatine amidinohydrolase), to the creatinase variants encoded by these genes, and to different applications of these variants of creatinase, particularly to their use for determining the creatinine and/or creatine concentration in a sample.

Background of the Invention

Creatine and creatinine are usually found in blood and urine, and can be used as very important parameters for diagnosing uremia, chronic nephritis, acute nephritis, gigantism, tonic muscular dystrophy and some other related diseases. For such routine diagnostic purposes, creatinine and creatine concentrations in blood and urine need to be determined quickly and accurately (Schumacher, G., et al., Ann Biol Clin 51 (1993) 815-9, Fujita, T., et al., Clin Chem 39 (1993) 2130-6, Spencer, K., Ann Clin Biochem 23 (1986) 1-25).

Generally, creatinase catalyses the hydrolysis of creatine (e.g. as generated from creatinine through creatininase) to urea and sarcosine. In the presence of sarcosine oxidase, sarcosine is oxidized and hydrogen peroxide is generated. The hydrogen peroxide formed is determined by a standard procedure for measurement of hydrogen peroxide, e.g. by a colorimetric procedure catalyzed by horse-radish peroxidase. Through these coupled reactions, creatinine and/or creatine in a sample can be quickly and quantitatively determined (Kopetzki, E., et al., Clin Chem 40 (1994) 688-704, Beyer, C., et al., Clin Chem 39 (1993) 1743-4).

Enzymes with creatinase activity have been found in different microorganisms, including Arthrobacter (Kaplan, A. Szabo, L. L., Mol Cell Biochem 3 (1974) 17-25, Kaplan, A. Naugler, D., Mol Cell Biochem 3 (1974) 9-15, Nishiya, Y., et al., Mol Gen Genet 257 (1998) 581-6), Bacillus (Suzuki, K., et al., Journal of Fermentation & Bioengineering 76 (1993) 77-81), Alcaligenes (Matsuda, Y., et al., Chem. Pharm. Bull. 34 (1986) 2155-2160), Pseudomonas (Hong, M. C., et al., Biochem Genet 36 (1998) 407-15) and Flavobacterium (Koyama, Y., et al., Agric Biol Chem 54 (1990) 1453-7).

Several creatinase enzymes obtained from different microorganisms have been industrially produced and used as reagents for clinical tests. However, even with state-of-the-art creatinase enzymes and/or with state-of-the-art creatine assays, the creatinase enzyme is generally the major limiting factor for or in such assays.

Important criteria for broad industrial, especially diagnostic applicability, comprise solubility, conductivity (in case of electronic signal generation and measurement), stability, and enzymatic activity with creatine as substrate. Obviously, stability (e.g., short-term heat stability or long-term storage stability) and enzymatic activity are especially relevant or critical in routine applications. Schumann et al. (Schumann, J., et al., Biol Chem Hoppe Seyler 374 (1993) 427-344) showed that "extrinsic factors", such as DTE, bovine serum albumin (BSA) and glycerol, can improve the stability of *P. putida* creatinase. But these extrinsic factors are generally not compatible with the conditions for a coupled creatinine and/or creatine assay system using creatininase, creatinase, sarcosine oxidase and peroxidase.

All the native or wild-type creatinase enzymes known from the art exhibit certain limitations with regard to thermostability and their K_m value for creatine. For example, the enzyme derived from the *Bacillus* is unstable at any temperature of more than 40°C (see patent US 4,420,562). The creatinase from *Pseudomonas putida* has both, rather low storage stability (Schumann, J., et al., supra) and a rather high K_m value (see patent EP 0 291 055), and, consequently, the enzyme is not suitable for creatine determination in a liquid formulation as required in the coupled creatine assay mentioned above using also sarcosine oxidase and peroxidase as auxiliary reagents.

Creatinases isolated from *Corynebacterium* (see patent DE 2 659 878), *Micrococcus*, or *Bacillus* (see patent DE 3 024 915) have been found unstable at elevated temperatures and, therefore are not suitable for the formulation of long-term stable liquid assay reagents, either. A creatinase from *Alcaligenes* has been reported to have a K_m value of 13 mM and to be stable at a temperature of 45° C for 30 minutes (see patent US 5,451,520).

There are few literature reports about engineered creatinases (Schumann, J., et al., Protein Sci 2 (1993) 1612-20) used random and site-directed mutagenesis to introduce point mutations into *P. putida* creatinase in order to generate mutants with an altered stability profile. Three point mutants with A109V, V355M, or V182I, one double mutant with A109V and V355M, and one triple mutant with all three substitutions in *P. putida* creatinase were generated and compared to the wild-type enzyme regarding their physical and enzymological properties. Physicochemical measurements showed that the mutations exhibit only a small increase in overall stability. Even the best mutant, however, is not stable at elevated temperature, e.g. temperatures of 45° C and above. The catalytic properties are in the range of the wild-type enzyme or are inferior.

EP 0 790 303 describes a process of generating *Alcaligenes faecalis* creatinase variants with lower K_m values as compared to the wild-type enzyme using the in vivo mutagenesis approach in *E. coli* strain XL1-Red (Stratagene, Cat. Nr. 200 129). The K_m values of the creatinase variants were found to range from 4.5 mM to 9.0 mM.

It can be summarized that the attempts known from the art, e.g., as discussed above, either aiming at the isolation and purification of creatinase from various microorganisms or at the genetic improvement of such enzymes, to date still have led to enzymes lacking important industrial properties.

A great demand and clinical need therefore exists for mutant forms of creatinase with improved properties, especially regarding improved stability and/or lower K_m -value to its substrate creatine.

It was the task of the present invention to provide new mutants or variants of creatinase with significantly lower K_m -value or significantly improved stability, or both.

Surprisingly it has been found that it is possible to provide such creatinase enzymes by mutating one or several amino acids of an *Erwinia*-type creatinase at precisely defined

positions. Such relevant positions are defined and given as amino acid positions corresponding to amino acid positions of the wild-type creatinase isolated from *Erwinia* sp. DSM 97-934.

Summary of the invention

The present invention relates to a variant of an *Erwinia*-type creatinase modified relative to a wild-type creatinase having creatinase activity, characterized in that said variant comprises at least one amino acid substitutions at a position corresponding to a position selected from the group of positions consisting of N130, M203, I278, I304, and F395 of the amino acid sequence shown in SEQ ID NO:2.

The variant creatinases according to the present invention exhibit improved properties as compared to the respective wild-type enzymes. Improved variants are described, exhibiting either individually or as a combination of such improved properties, like lower conductivity, improved stability, or lower K_m -value for creatine.

The present invention also relates to a reagent for determination of creatine comprising a creatinase as described in the present invention. It especially relates to a reagent, which comprises a creatinase variant and also comprises a sarcosine oxidase as well as reagents for the detection of hydrogen peroxide.

Due to the significantly improved properties as compared to wild-type or mutated enzymes known in the art, the variant creatinase according to the present invention is used with great advantage in a method for detection of creatine.

Detailed description of the invention

In a first embodiment the present invention relates to a variant of an *Erwinia*-type creatinase modified relative to a wild-type creatinase having creatinase activity, characterized in that said variant comprises at least one amino acid substitutions at a position corresponding to a

-5-


position selected from the group of positions consisting of N130, M203, I278, I304, and F395 of the amino acid sequence shown in SEQ ID NO:2.

The present invention relates to improved variants of an *Erwinia*-type creatinase. An „*Erwinia*-type creatinase" is defined by at least 85 % sequence identity on the amino acid level as compared to SEQ ID NO:2.

The terminology used, for example N130, indicates that the amino acid asparagine (N) of position 130 in SEQ ID NO:2 is referred to. If the amino acid position is followed by one or more amino acids (as described by single letter code) for example, N130 D, E this indicates that the amino acid arginine found in wild-type position 130 has been substituted either by aspartic acid (D) or glutamic acid (E).

Creatinase (creatine amidinohydrolase, EC3.5.3.3) catalyzes the hydrolytic cleavage of creatine to sarcosine and urea. An enzyme with this catalytic activity is defined in the art as having creatinase activity.

The use of creatinase enzymes is highly important in medical diagnostics because it is used to assess creatinine as well as creatine concentrations. Especially important is the assessment of creatinine clearance, which is used to monitor kidney function. The enzyme creatinase catalyzes the second step in such a coupled creatinine-creatine assay (Siedel, J., et al., Clinical Chemistry 30 (1984) 968-969).



As is obvious from the above reaction schemes creatinase may be used in the detection of creatine alone or in the combined detection of creatinine and creatine. The hydrogen peroxide (H_2O_2) generated upon the action of sarcosidase is measured. Preferably a dye system is used comprising a coupler reagent (e.g. as above 4-AA = 4-Amino-2,3-dimethyl-1-phenyl-3-pyrazoline-5-one) and a color generating substance (e.g. as above TBMB = 2,4,6-tribromo-3-hydroxybenzoic acid) which upon action of peroxidase produces a color signal (benzoquinone) proportional to the peroxide concentration and thus proportional to the concentration of creatinine and creatine in the sample analyzed.

The present invention is based on the surprising finding that variants of wild-type creatinase can be provided, which represent significant improvements as compared to the corresponding wild-type enzyme. The improvements according to the present invention are described based on the amino acid composition and numbering of the creatinase enzyme as isolated from *Erwinia* species (DSM 97-934). The wild-type enzyme from this *Erwinia* species has been cloned, sequenced (SEQ ID NO:1) and the protein sequence found to correspond to SEQ ID NO:2.

It has been found that amino acid substitutions in amino acid positions F59, N113, N203, I278, I304, and F395 of SEQ ID NO:2 are particularly relevant to produce enzyme variants with improved properties. Comparative sequence analysis using amino acid sequence motives of 7 amino acids in length and comprising the above identified critical amino acid position in their center, revealed that corresponding positions can be identified in creatinases isolated from various other organisms. Obviously, the mutations described in the present invention can also be used to modify corresponding sequence positions of other known and yet un-identified creatinases. The term "corresponding to a position" is used to indicate that creatinase enzymes and variants thereof may also be found or generated comprising additional amino acids or lacking amino acids, which upon sequence alignment to SEQ ID NO:2 results in a different absolute number for the corresponding sequence position or motive.

The multiple alignment and comparison of *Erwinia* creatinase with other related creatinases has been performed with the PileUp program of GCG Package Version 10.2 (Genetics Computer Group, Inc.). PileUp creates a multiple sequence alignment using a simplification of the progressive alignment method of Feng, D. F. Doolittle, R. F., J Mol Evol 25 (1987) 351-60, and the scoring matrixes for identical, similar, or different amino acid residues are defined accordingly. This process begins with the pairwise alignment of the two most similar (creatinase) sequences, producing a cluster of two aligned sequences. This cluster can then be aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences can be aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments that include increasingly dissimilar sequences and clusters, until all sequences have been included in the final pairwise alignment.

A variant of an *Erwinia*-type creatinase in the sense of the present invention is a modified creatinase enzyme, which relative to its wild-type sequence comprises at least one amino acid substitution at a position, which corresponds to a position selected from the group of positions consisting for N130, N203, I278, I304, and F395 of SEQ ID NO:2.

In case the wild-type enzyme does not comprise a tryptophane(W) residue in position 59, the preferred creatinase variant may additionally be modified to comprise W in position 59. A preferred *Erwinia*-type creatinase variant according to the present invention therefore contains the amino acid substitution F59W and at least one further amino acid substitution selected from the group of positions corresponding to N130, N203, I278, I304, and F395 of SEQ ID NO:2.

The positions corresponding to the above mentioned positions of SEQ ID NO:2 from related creatinase enzymes can easily be found by sequence alignment using the following sequence motives: 56 – 62 (SEQ ID NO:3), 127 – 133 (SEQ ID NO:4), 200 – 206 (SEQ ID NO:5), 275 – 281 (SEQ ID NO:6), 301 – 307 (SEQ ID NO:7), and 392 – 398 (SEQ ID NO:8). These positions correspond if six out of seven amino acids are identical.

A preferred variant of an Erwinia-type creatinase according to the present invention comprises a mutation in an amino acid position corresponding to position N130 of SEQ ID NO:2. Preferably the amino acid residues introduced by substitution are polar residues or charged residues. Preferred are creatinase variants comprising the substitution N130E or N130D, the variant comprising N130D being most preferred.

A preferred variant of an Erwinia-type creatinase according to the present invention comprises a mutation in an amino acid position corresponding to position M203 of SEQ ID NO:2. Preferably the amino acid residues introduced by substitution are aliphatic residues. Preferred are creatinase variants comprising the substitution M203V, M203I or M203L, the variant comprising M203V being most preferred.

A preferred variant of an Erwinia-type creatinase according to the present invention comprises a mutation in an amino acid position corresponding to position I278 of SEQ ID NO:2. Preferably the amino acid residues introduced by substitution are small polar residues or small hydrophobic residues. Preferred are creatinase variants comprising the substitution I278T, I278S, I278V or I278G, the variant comprising I278T being most preferred.

A preferred variant of an Erwinia-type creatinase according to the present invention comprises a mutation in an amino acid position corresponding to position I304 of SEQ ID NO:2. Preferably the amino acid residues introduced by substitution are aliphatic residues. Preferred are creatinase variants comprising the substitution I304L or I304V, the variant comprising I304L being most preferred.

A preferred variant of an Erwinia-type creatinase according to the present invention comprises a mutation in an amino acid position corresponding to position F395 of SEQ ID NO:2. Preferably the amino acid residues introduced by substitution are aliphatic residues. Preferred are creatinase variants comprising the substitution F395L, F395I, or F395V, the variant comprising F395L being most preferred.

In a further preferred embodiment the variant of an Erwinia-type creatinase according to the present invention comprises at least two substitutions which corresponds to a position selected from the group of positions consisting of N130, N203, I278, I304, and F395 of SEQ ID NO:2.

In a preferred embodiment a variant of an Erwinia-type creatinase according to the present invention comprises a substitution of at least one of the amino acids corresponding to positions N130 or I278.

Substitutions of amino acids at positions corresponding to 130 or 278 of SEQ ID NO:2 have been found extremely important to increase the stability of an Erwinia-type creatinase variant.

A further preferred Erwinia-type creatinase variant comprises at least one substitution at an amino acid position corresponding to position 130 by D or E (aspartic or glutamic acid) or position 278 by threonine (T). A variant comprising substitutions at both the above discussed amino acid positions represents a further preferred embodiment.

The term "stability" in general is used to describe many different properties like pH-tolerance, salt tolerance or thermal inactivation. According to the present invention stability relates to heat or to storage stability. Heat or storage stability is easily analyzed using appropriate model systems.

In a further preferred embodiment the variant of an Erwinia-type creatinase according to the present invention is characterized in that said variant exhibits improved long-term storage stability.

Storage stability relates to the long term stability of reagents, e.g. to stability under storage for several months or a few years. The long-term storage stability is assessed by treating a solution containing creatinase for 21 days at 35° C. Stability in this model system is known to correspond to a storage stability of at least 18 months using storage temperatures between 4 and 8° C. A creatinase variant is considered stable if after storage according to this model at

least 90 % of the original enzymatic activity are still found. With the improved variants according to the present invention at least 90 % of the original enzymatic activity are left and can be measured by routine procedures.

Heat stability is assessed in a short-term stress model, for example by incubation of the enzyme at 56° C for 20 minutes. Such a short-term stress model is very important to decide, whether reagents may be shipped at ambient temperature or whether cooling is required. A creatinase or a variant thereof is defined as meeting these arbitrary stability criteria (i.e., the short-term thermal stress stability criteria) if after 20 minutes at 56° C at least 30 % of the original creatinase activity is left. A preferred variant of creatinase according to the present invention exhibits improved stability at 56°C. I. e., after incubation at 56° C for 20 minutes at least 30 % of the original activity are left.

Variants of an Erwinia-type creatinase have been found, which exhibit improvements both with respect to both short-term thermal stress, i.e. heat stability as well as to long-term storage stability. Such variants exhibiting improved stability, both under heat stress and under long-term storage conditions represent a further preferred embodiment of the present invention.

In a further preferred embodiment according to the present invention the variant of an Erwinia-type creatinase is characterized in that it comprises an amino acid substitution in at least one of the positions corresponding to I304 or F395 of SEQ ID NO:2.

As is known from the state of the art (see above) the catalytic property for cleavage of creatine of the various creatinases so far at hand is considered critical. One of these catalytic properties is most easily expressed in terms of Km-values.

The Km-value of an enzyme reflects both the binding affinity between enzyme and substrate and the rate at which the enzyme-bound substrate is converted to the corresponding reaction product. The higher the Km-value, the lower is the catalytic efficiency of an enzyme. It surprisingly has been found that amino acid positions corresponding to position 304 and 395

of SEQ ID NO:2 are very important in order to generate variants of creatinase with improved, i.e., lower K_m -values. In a preferred embodiment the variant of an Erwinia-type creatinase comprises at least one amino acid substitution at an amino acid position corresponding to 304 by the amino acid L, or V, or position 395 by the amino acid L, I, or V.

An Erwinia-type creatinase variant with lower K_m -value for creatine comprising substitutions at both the amino acid positions corresponding to 304 and 395 of SEQ ID NO:2 represents a further preferred embodiment.

A variant creatinase which is characterized by having a lower K_m -value in the enzymatic cleavage of creatinine represents a further preferred embodiment. The K_m -value of a creatinase variant is either assessed in comparison to the corresponding wild-type enzyme or given in absolute terms as K_m -value. The K_m -value is determined by routine procedures, for example using the reagents as described in Example 7.

An Erwinia-type creatinase according to the present invention preferably exhibits a K_m -value which compared to the corresponding wild-type enzyme is 80% or less, more preferred 65% or less and most preferred 50% or less of the K_m -value as determined for the corresponding wild-type enzyme.

An Erwinia-type creatinase according to the present invention in absolute terms preferably exhibits a K_m -value of less than 10 mM, more preferred of less than 4 mM and most preferred of less than 2.5 mM.

It has been found that some of the improvements in terms of stability go to the expense of the K_m -value (see Table 4). Interestingly and surprisingly, it has been also been found that by including the appropriate "catalytic property mutations" as discussed above, into a variant which exhibits improved stability properties, creatinase variants can be designed with advantageous properties in both respects. In a preferred embodiment the Erwinia-type creatinase variant is characterized in that said variant comprises at least one amino acid

substitution at a position corresponding to a position in the amino acid sequence shown in SEQ ID NO: 1, selected from the positions N130 and I278 as well as at least one substitution at a position corresponding to a position of SEQ ID NO:2 selected from the positions I304 and F395

An Erwinia-type creatinase variant comprising substitutions in the position corresponding to N130, M203 and I278 of SEQ ID NO:2 represents a further preferred embodiment according to the present invention. Preferably this variant comprises N130D, M203V and I278T.

In a preferred embodiment the variant creatinase according to the present invention exhibits both improved stability and a lower Km-value in the enzymatic cleavage of creatine as compared to the wild-type creatinase obtained from Erwinia spec. (DSM97-934).

A creatinase variant according to the present invention can e.g., be produced by starting from a creatinase gene as isolated from Erwinia spec. DSM 97-934 as well as by starting from a homologous sequence. In the context of this application the term "homologous" is meant to comprise wild-type creatinases as isolated from other microorganisms, provided that the sequence homology as compared to SEQ ID NO:2 is at least 85 %. With other words, after appropriate alignment using the PileUp program, at least 85 % of the amino acids of that creatinase are identical to the amino acids described in SEQ ID NO:2.

It will be understood that variations of DNA and amino acid sequences naturally exists, or may be intentionally introduced using methods known in the art. These variations may result in up to 15 % amino acid differences in the overall sequence, due to deletions, substitutions, insertions, inversions or additions of one or more amino acid residues in said sequence as compared to SEQ ID NO:2. Such amino acid substitutions may be made, for example, on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; amino acids with uncharged polar head groups or nonpolar head groups having

-13-

similar hydrophilicity values include the following: leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, tyrosine. Other contemplated variations include salts and esters of the aforementioned polypeptides, as well as precursors of the aforementioned polypeptides, for example, precursors having N-terminal substitution such as methionine, N-formylmethionine used as leader sequences. Such variations may be made without necessarily departing from the scope and the spirit of the present invention.

Procedures for isolating a creatinase gene from a microorganism as well as operations and methods used for the purification, characterization, and cloning of relevant DNA are well known in the art (e.g., EP 0 790 303 and Ausubel, F., et al. in "Current Protocols in Molecular Biology" (1994) Wiley Verlag).

Based on the identification of critically important amino acid positions as described in the present invention the skilled artisan now can easily produce further appropriate variants of creatinase.

The present invention further includes an expression vector comprising a nucleic acid sequence of the invention operably linked to a promoter sequence capable of directing its expression in a host cell. Preferred vectors are plasmids such as pKK177-3HB. A vector containing the wild-type *Erwinia* creatinase gene is shown in Figure 1 (pCT-WT).

Expression vectors useful in the present invention typically contain an origin of replication, a promoter located in front (i.e., upstream of) the DNA sequence and followed by the DNA sequence coding for all or part of creatinase variants. The DNA sequence coding for all or part of the creatinase variants is followed by transcription termination sequences and the remaining vector. The expression vectors may also include other DNA sequences known in the art, for example, stability leader sequences which provide for stability of the expression product, secretory leader sequences which provide for secretion of the expression product, sequences which allow expression of the structural gene to be modulated (e.g., by the presence or absence of nutrients or other inducers in the growth medium), marking sequences which are

-14-

capable of providing phenotypic selection in transformed host cells, and the sequences which provide sites for cleavage by restriction endonucleases. The characteristics of the actual expression vector used must be compatible with the host cell which is to be employed. For example, when cloning in an E. coli cell system, the expression vector should contain promoters isolated from the genome of E. coli cells (e.g., tac, lac, or trp). Suitable origins of replication in E. coli various hosts include, for example, a ColE1 plasmid replication origin. Suitable promoter include, for example, the tac, lac, and trp. It is also preferred that the expression vector include a sequence coding for a selectable marker. The selectable marker is preferably antibiotic resistance. As selectable markers, ampicillin resistance, or kanamycin resistance may be conveniently employed. All of these materials are known in the art and are commercially available.

Suitable expression vectors containing the desired coding and control sequences may be constructed using standard recombinant DNA techniques known in the art, many of which are described in Sambrook et al. (Sambrook, J., et al. - in "Molecular Cloning: A Laboratory Manual" (1989) -, Cold Spring Harbour, NY, Cold Spring Harbour Laboratory Press).

The present invention additionally concerns host cells containing and expression vector which comprises a DNA sequence coding for all or part of the mutant creatinase. The host cells preferably contain an expression vector which comprises all or part of one of the DNA sequences having one or more mutations. Further preferred are the host cells containing an expression vector comprising one or more regulatory DNA sequences capable of directing the replication and/or the expression of, and operatively linked to a DNA sequence coding for, all or part of mutant creatinase. Suitable host cells include, for example, E. coli HB101 (ATCC 33694) available from Pomega (2800 Woods Hollow Road, Madison, WI, USA), XL1-Blue MRF available from Stratagene (11011 North Torrey Pine Road, La Jolla, CA, USA) and the like.

Expression vectors may be introduced into host cells by various methods known in the art. For example, transformation of host cells with expression vectors can be carried out by

polyethylene glycol mediated protoplast transformation method (Sambrook, et al. 1989). However, other methods for introducing expression vectors into host cells, for example, electroporation, biolistic injection, or protoplast fusion, can also be employed.

Once an expression vector containing creatinase variants has been introduced into an appropriate host cell, the host cell may be cultured under conditions permitting expression of the desired creatinase variants. Host cells containing an expression vector which contains a DNA sequence coding for all or part of the mutant creatinase may also be identified by one or more of the following general approaches: DNA hybridization, the presence or absence of marker gene functions, assessment of the level of transcription as measured by the production of creatinase mRNA transcripts in the host cell, and detection of the gene product immunologically, but preferably by identification through enzyme assay (colorimetric detection, etc.).

The present invention also teaches methods for screening of appropriate creatinase variants. The assay conditions chosen to assess enzymatic activity, are adapted to ensure that the expected small enhancements brought about e.g., by a single amino acid substitution, can be measured. This has been accomplished by adjusting the assay conditions such that the wild type (or parent) enzyme activity is close to the lower detection limit. One mode of selection or screening of appropriate mutants with lower K_m -values is given in Example 3. Any change or improvement as compared over the wild-type enzyme this way can be clearly detected. Methods to assess long term storage stability as well as short term thermal stress stability are also described in detail in the Examples section.

By creating a replica plate of the clones to be assayed, the clones can be subjected to an array of procedures without loss of viability. For example, many intracellular enzymes require cell lysis before they can be assayed. This procedure could be performed in a screen, and when positives are found, they can be re-grown from the replica plate.

-16-

Furthermore, multiple measurements can be made on each sample to account for variability in protein expression levels or to check other key enzyme properties. Additionally, these assays are often done in much the same way the enzyme is traditionally assayed (e.g. spectrophotometrically), making the assay a straight-forward technique that can be implemented quickly.

It should, of course, be understood that not all expression vectors and DNA regulatory sequences will function equally well to express the DNA sequences of the present invention. Neither will all host cells function equally well with the same expression system. However, one of ordinary skill in the art may make a selection among expression vectors, DNA regulatory sequences, and host cells using the guidance provided herein without undue experimentation and without departing from the scope of the present invention.

It is preferred that the polypeptides of the present invention are obtained by production in host cells expressing a DNA sequence coding the mutant creatinase. However, the polypeptides of the present invention may be obtained by *in vitro* translation of the mRNA encoded by a DNA sequence coding for the mutant creatinase. For example, the DNA sequences may be synthesized using PCR as described above and inserted into a suitable expression vector, which in turn may be used to *in vitro* transcription/translation system.

In a preferred embodiment the present invention relates to a method for production of an *Erwinia*-type creatinase variant, the method comprising the steps of cloning a wild-type enzyme, introducing mutations, selecting as compared to the wild-type enzyme variants with improved thermostability or improved creatinase activity, and recombinantly producing said variant.

The polypeptides produced in these manners may then be isolated and purified to some degree using various protein purification techniques. For example, chromatographic procedures such as ion exchange chromatography, gel filtration chromatography and affinity chromatography may be employed.

-17-

The polypeptides of the present invention have been defined by means of determined DNA and deduced amino acid sequencing. Due to the degeneracy nature of the genetic code, which results from there being more than one codon for most of the amino acid residues and stop signals, other DNA sequences which encode the same amino acid sequence as given in SEQ ID NO:2 may be used for the production of the polypeptides of the present invention.

The invention also contemplates a process for producing a creatinase variant of the current invention comprising culturing a host cell of the invention under conditions suitable for production of the mutant creatinase of the invention. For bacterial host cells, typical culture conditions are liquid medium containing the appropriate antibiotic and induction agent. Cultures are shaken or stirred at temperature suitable for optimal production of enzyme, e.g., about 25°C to about 37°C. Typical appropriate antibiotics include ampicillin, kanamycin, chloroamphenicol, tetracycline and the like. Typical induction agents include IPTG, glucose, lactose and the like.

As discussed, improved variants of creatinase are very important in order to produce appropriate reagents for creatinine and/or creatine assays, e.g. as used in the clinical diagnostic routine. Especially a liquid reagent comprising an improved creatinase meeting improved stability criteria is highly import. In a preferred embodiment the present invention relates to a reagent for determination of creatine (or for the determination of creatinine or for the simultaneous determination of both analytes) comprising an Erwinia-type creatinase variant according to the present invention.

In clinical routine nowadays in the determination of an analyte as few reagents as possible, requiring as little handling as possible have to be provided to facilitate measurement for the customer and to reduce the chances for erroneous handling. In a further preferred embodiment the present invention therefore relates to a reagent comprising a creatinase variant according to the present invention and also comprising a sarcosine oxidase and reagents for the detection of hydrogen peroxide.

-18-

One of the major applications of the improved creatinase variants in this invention is for the measurement of creatinine and/or creatine in clinical samples. Due to their enhanced overall stability and relatively small K_m values for creatine, assays system using these improved creatinase variants e.g., as liquid reagents, are more economic and more robust. In a further preferred embodiment the present invention relates to a method for detection of creatine, the method comprising the steps of a) incubating a sample to be analyzed with b) a reagent comprising a variant of an Erwinia-type creatinase according to any of claims 1 to 7, a sarcosine oxidase and reagents for detection of hydrogen peroxide and c) correlating the hydrogen peroxide generated to the concentration of creatinine in said sample.

The reagents for the determination of creatine according to the present invention contains each ingredient in appropriate amounts. Preferred ranges are: creatinase 0.01- 100 U/ml, sarcosine oxidase 1 - 50 U/ml, peroxidase 0.01 - 30 U/ml, a coupler reagent at 0.1 - 10 mM, and a color generating substance at 0.1 - 50 mM. More preferred ranges are: creatinase 1- 50 U/ml, sarcosine oxidase 1 - 50 U/ml, peroxidase 0.1 - 10 U/ml, a coupler reagent at 0.1 - 10 mM, and a color generating substance at 1 - 30 mM. The buffer system used preferably is a potassium phosphate buffer of 0.1 M with a pH of 7.5 to 8.0 or a 0.05 M TAP-buffer of pH 8.0 to 8.5 (TAP = 3-[(tris(hydroxymethyl)methyl]amino]propane sulfonic acid).

The creatinase from Erwinia and variants thereof exhibit an additional important property, i.e. a much lower conductivity. These creatinases can also be used in biosensors similar to those described by Kim et al. (Kim, E. J., et al., Anal. Chim. Acta 394 (1999) 225-231) and by Khan et al. (Khan, G. F. Wernet, W., Anal. Chim. Acta 351 (1997) 151-158) for online monitoring creatinine/creatine concentration in a sample or a reactor.

The improved creatinase variants of the present invention are described by making reference to substitutions of amino acids from SEQ ID NO:2, the creatinase isolated from Erwinia sp. (DSM 97-934).

Sequence comparisons to other known creatinases revealed that the corresponding enzymes isolated from Actinobacillus or Alcaligenes strains are more than 90 % identical to SEQ ID NO:2. Even more surprisingly, we found that the 7-mer peptide sequence motives of SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8 are identical in these

creatinases. The same type of substitution(s) as described in the present invention is(are) used to improve these enzymes.

In a preferred embodiment the present invention relates to a variant of *Erwinia* creatinase, which compared to the corresponding wild-type enzyme as isolated from DSM (97-934), is characterized in that said variant comprises at least one amino acid substitution at a position corresponding to a position selected from the group of positions consisting of F59, N130, M203, I278, I304, and F395 of the amino acid sequence shown in SEQ ID NO:2.

In a further preferred embodiment the present invention relates to a variant of an *Alcaligenes* creatinase which compared to the corresponding wild-type enzyme is characterized in that said variant comprises at least one amino acid substitution at a position corresponding to a position selected from the group of positions consisting of N130, M203, I278, I304, and F395 of the amino acid sequence shown in SEQ ID NO:2.

In a further preferred embodiment the present invention relates to a variant of an *Actinobacillus* creatinase which compared to the corresponding wild-type enzyme is characterized in that said variant comprises at least one amino acid substitution at a position corresponding to a position selected from the group of positions consisting of N130, M203, I278, I304, and F395 of the amino acid sequence shown in SEQ ID NO:2.

Preferably, the above described variants of an *Erwinia* creatinase, an *Alcaligenes* creatinase or of an *Actinobacillus* creatinase in comparison to the corresponding wild-type sequence, comprise 20 amino acid substitutions or less, more preferred 15 amino acid substitutions or less, even more preferred 10 amino acid substitutions or less, and most preferred 6 amino acid substitutions or less.

In the following examples, all reagents, restriction enzymes, and other materials were obtained from Roche Diagnostics, unless specified from other commercial sources, and used according to the indication by suppliers. Operations employed for the purification, characterization and cloning of DNA are well known in the art and can be adapted from published literatures.

The following examples, references, sequence listing and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended

claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Figures

Figure 1 This figure shows an expression vector comprising the gene of SEQ ID NO: 1 coding for the full length creatinase of SEQ ID NO:2. The expression vector is known in the art as pCT.

Figure 2 This figure shows the short term thermal stress stability of the wild-type creatinase isolated from *Erwinia* sp. (DSM 97-934) and of variants according to the present invention. The stability is assessed by calculating a "Thermo-Index". This index is obtained by calculating the residual enzymatic activity after short term heat stress in relation to the activity without heat stress.

Example 1: Cloning and expression of Erwinia DSM97-934 creatinase gene in E. coli

The creatinase gene was amplified from *Erwinia* DSM97-934 genome with PCR. Both the amplified DNA fragment and the pKK177-3HB plasmid were digested with the restriction enzymes Eco RI and Hin dIII. The digested products were gel purified and ligated. An aliquot of the ligation reaction mixture was used to transform competent *E. coli* cells, for example *E. coli* HB101 cells. The transformants were subsequently selected on LB plates containing ampicillin. To assay, individual colonies were chosen, grown over night in LB medium (cf. Sambrook et al., 1989, supra) containing ampicillin and subjected to screening.

Example 2: Mutating the wild-type creatinase from Erwinia sp.

In order to improve the enzymes properties the following mutagenesis methods were used to generate creatinase variants: mutagenic PCR, saturation mutagenesis and site directed mutagenesis.

-21-

Mutagenic PCR is a method to introduce random point mutations into the selected DNA. This method was performed according to the protocol described by Cadwell and Joyce. (Cadwell, R. C. Joyce, G. F., PCR Methods Appl 3 (1994) 136-40) with some modifications.

The mutagenic PCR was set up as following:

Template DNA	61 fmol (= 40 ng pCT-WT)
Forward Primer	40 pmol
Reverse Primer	40 pmol
10 x Taq Puffer	10 µl (without Mg ²⁺ ; Roche # 1 699 105)
MgCl ₂	7 mM
MnCl ₂	0 – 0.8 mM
dATP, dGTP	0,2 mM (Roche # 1 969 064)
dCTP, dTTP	1 mM (Roche # 1 969 064)
Taq DNA polymerase	5 U(Roche # 1 418 432)
H ₂ O	added to total volume of 100 µl

The PCR cycling was done with the following conditions:

95 °C	5 min	
95 °C	1 min	
46/70 °C	1 min	30 cycles

-22-

72 °C	2 min	
4 °C	time span variable, set as appropriate	

Two generations of mutagenic PCR were performed. First generation was carried out with plasmid pCT-WT, primer ECF21 (5' CAG GAA ACA GAA TTC ATG ACT 3') /HIR21 (5' CCA AAA CAG CCA AGC TTT CAG 3') and an annealing temperature of 46°C. Second round of mutagenic PCR was carried out with plasmid pCT1m24, primer CtextF (5' CAT CGG CTC GTA TAA TGT GTG G 3')/CtextR (5' GCC AAA ACA GCC AAG CTT TCA G 3') and an annealing temperature of 70°C. Each round of mutagenic PCR was performed with several MnCl₂ concentrations to obtain different mutagenic rates and therefore different rates of inactive clones.

Mutagenic PCR products were purified using ion exchange techniques (Roche High Pure™ PCR Product Purification Kit # 28104) and were eluted in H₂O. The purified PCR products and plasmid pKK177-3HB were digested with restriction enzymes Eco RI and HindIII, purified by preparative gel electrophoresis (1% agarose/TAE) and gel extraction with QIAquick Gel Extraction Kit (Quiagen Cat.# 28706). Eco RI/HindIII-digested PCR products were ligated into Eco RI/Hind III-digested plasmid vector pKK177-3HB using T4 DNA ligase (Roche # 481 220) according to the manual. Ligation reactions were introduced into E. coli HB101 according to the manual for high-efficiency transformation by electroporation (Current protocols in molecular biology, chapter 1.8.4) or into Epicurian Coli® XL1-Blue super competent cells (Stratagene # 200236) according to the suppliers manual. Transformants were plated on LB-Agar with 100 µg/ml ampicillin and grown at 37 °C, 14 h. The pool with about 50 % active clones were chosen for screening. In each evolution generation, about 3 000 clones were screened.

Saturation Mutagenesis is a method to randomly introduce any of the 20 canonic amino acids into a defined position of the protein using a PCR like method with defined primers

-23-

containing a random three base stretch. The whole plasmid is amplified using two complementary primers. The parental plasmid (without the mutation) is removed by enzymatic cleavage using Dpn I. For mutagenesis of creatinase genes, QuikChange™ Site-Directed Mutagenesis Kit (Stratagene # 200518) was used according to the suppliers manual. Plasmids with mutagenized creatinase genes were introduced into E. coli HB 101 or E. coli XL1-Blue as described above. About 200 clones were screened in each saturation mutagen experiment to ensure sufficient statistical diversity.

To combine several useful mutations in one creatinase mutant the method of site directed mutagenesis was also used. The principle and protocol is similar to saturation mutagenesis, except that defined primers containing the mutated triplet were used.

Example 3: Screening and assay of Erwinia-creatinase and variants thereof

The activity assay for creatinase and its variants was performed by using creatine as substrate and a coupled enzymatic reaction according to Guder, W. G., et al., J Clin Chem Clin Biochem 24 (1986) 889-902.

The reaction mixture was set up by combining the following reagents:

90 mM creatine in 0.1 M potassium phosphate buffer, pH7.8	2.5 ml
30 mM 4-Amino-2,3-dimethyl-1-phenyl-3-pyrozoline-5-one	0.05 ml
143,5 mM 2,4,6-tribrom-3-hydroxybenzoic acid	0.1 ml
sarcosine oxidase (150 U/ml)	0.1 ml
peroxidase (500 U/ml)	0.02 ml
creatinase solution (crude extract)	0.01 ml

-24-

The assay for creatinase was carried out at 25°C and at the wavelength of 546 nm for 10 min. ΔE was calculated from the measurement points of 6 to 10 min.

Screening clones of creatinase and its variants was carried out according to the above assay procedure.

Each screening step was performed in 96-well microtiterplates. Colonies were picked into these plates and grown for 24 h in 200 μ l of LB-medium (cf. Sambrook et al., 1989, supra). After cell lysis by addition of 5 μ l detergent (B-PER™ Bacterial Protein Extraction Reagent; PIERCE #78248) per well, 10 μ l aliquots of the crude extracts were assayed.

In order to screen creatinase variants with lower K_m values, creatinase activity of the crude enzyme extract thus obtained was determined. Using an activity determination reagent having only the 15th part of the above creatine substrate concentration, the creatinase activity of the same samples was again determined. Any clone wherein the ratio of the two kinds of the activity measures (activity with 1/15 substrate concentration /activity obtained by conventional method) increased beyond that of a wild type creatinase was selected as a candidate mutant having a lower K_m value. For example, the activity screening was done using an optimal creatine concentration (90 mM) and a reduced creatine concentration (6 mM). Comparison of the activity ratios of the two with that obtained using a wild type creatinase leads to the identification of the potentially positive creatinase variants having smaller K_m values.

In order to screen for thermostable creatinase variants, the crude extract of each creatinase variant was used to determine initial and residual creatinase activity before and after heat treatment (e.g. 56° C, 30 min). Comparison of the activity ratios of the two experiments of the activity ratio obtained using a wild type creatinase leads potentially positive creatinase variants with enhanced thermostability.

Example 4: Mutants of Erwinia creatinase with improved properties

-25-

The above mentioned screening efforts led to several creatinase variants with enhanced thermostabilities, improved solubilities, lower K_m values and/or lower conductivities. These variants are shown in Table 1. The sequences of the primers used in each generation are shown in Table 2.

Table 1: Examples of creatinase mutants generated

Mutant	DNA and amino acid sequence, respectively	Primers used
CT1m24	SEQ ID NO: 9 and SEQ ID NO: 10	ECF21 and HIR 21
CT2m9	SEQ ID NO: 11 and SEQ ID NO: 12	CTextF and CTextR
CT2m10	SEQ ID NO: 13 and SEQ ID NO: 14	CTextF and CTextR
CT2m28	SEQ ID NO: 15 and SEQ ID NO: 16	CTextF and CTextR
CTsd2	SEQ ID NO: 19 and SEQ ID NO: 20	CTf395lF and CTf395lR
CTqc2	SEQ ID NO: 17 and SEQ ID NO: 18	CT59F and CT59R
CTsd7	SEQ ID NO: 21 and SEQ ID NO: 22	CTf59wF and CTf59wR

Table 2 The sequences of the used primers

ECF21	CAG GAA ACA GAA TTC ATG ACT (SEQ ID NO: 23)
HIR21	CCA AAA CAG CCA AGC TTT CAG (SEQ ID NO: 24)
CTextF	CAT CGG CTC GTA TAA TGT GTG G (SEQ ID NO: 25)
CTextR	GCC AAA ACA GCC AAG CTT TCA G (SEQ ID NO: 26)
CTf395lF	CAT TAC CGG ATT CCC CCT GGG GCC TGA GCA CAA C (SEQ ID NO: 27)
CTf395lR	GTT GTG CTC AGG CCC CAG GGG GAA TCC GGT AAT G (SEQ ID NO: 28)
CT59F	GCA TCA ATT ACT ACT CTG GAN NNC TGT ACT GCT ATT TCG GCC GC (SEQ ID NO: 29)
CT59R	GCG GCC GAA ATA GCA GTA CAG NNN TCC AGA GTA GTA ATT GAT GC (SEQ ID NO: 30)
CTf59wF	GCA TCA ATT ACT ACT CTG GAT GGC TGT ACT GCT ATT TCG GCC GC (SEQ ID NO: 31)
CTf59wR	GCG GCC GAA ATA GCA GTA CAG CCA TCC AGA GTA GTA ATT GAT GC (SEQ ID NO: 32)

Example 5: Genetic characterization of Erwinia creatinase mutants with improved properties

The creatinase mutants with improved properties were genetically characterized, and their DNA and corresponding amino acid sequences are given in SEQ ID NO: 9 - 22. Table 3 summarizes the mutations as identified for each of these creatinase mutant.

Table 3: Amino acid substitutions of the creatinase variants with improved properties

Mutant	Mutations
CT1m24	N130D, I278T
CT2m9	N130D, M203V, I278T
CT2m10	N130D, I278T, F395L
CT2m28	N130D, I304L, I278T
CTqc2	F59W, N130D, I278T
CTsd2	N130D, M203V, I278T, F395L
CTsd7	F59W, N130D, M203V, I278T, F395L

Example 6 Purification of an Erwinia-type creatinase

The method for obtaining the purified creatinase and creatinase variants as selected above from the cell cultures may be any known method, such as the following. After the cells obtained by culturing in a nutrient medium were recovered, the cell pellets were homogenized in 20 mM Tris-HCl (pH 8) using high pressure cell disruption at a pressure of 900 bar to give a crude cell extract. The resulting enzyme extract was then subjected to ion exchange chromatography on DEAE-Sepharose fast flow using a 0.1 – 0.35 M NaCl gradient for elution. After this separation step, the resulting creatinase fraction may be further separated and purified by e.g. phenylsepharose column chromatography (Pharmacia Biotech) to give a

-28-

standard purified enzyme product. Usually, the end product thus obtained is at least 90 % pure and shows one predominant band in SDS-PAGE corresponding to creatinase. In case SDS PAGE should occasionally reveal a lower degree of purity the purification steps are repeated.

Example 7 Characterization of Erwinia-creatinase and of variants thereof

The characterizations of Erwinia-creatinase and its variants included K_M and short term thermal stress determination as well as determination of the residual activity after incubation at 35° C for 21 days in CreaPlus buffer.

a) K_M determination

For K_M determination, creatinase activity was determined according to the assay procedure described in Example 3 using six different concentrations of creatine substrate: 90 mM, 45 mM, 22.5 mM, 11.3 mM, 5.6 mM and 2.8 mM.

b) Assessment of short term thermal stress stability

The thermoinactivation of the cloned wild type Erwinia creatinase and its thermostable mutants was done with purified enzyme or enzyme mutants. The enzyme was stressed in concentrated form at an enzyme concentration of 30 U/ml in 0.1 M potassium phosphate buffer (pH 7.8). The (residual) creatinase activity was determined from appropriate dilutions made from this stock solution, as described by Guder et al. supra. The creatinase assay was set up as follows:

0.1 M creatine in 0.1 M potassium phosphate buffer, pH 7.8	2.5 ml
30 mM 4-Amino-2,3-dimethyl-1-phenyl-3-pyrazoline-5-one	0.05 ml
143,5 mM 2,4,6-tribromo-3-hydroxybenzoic acid	0.1 ml

-29-

sarcosine oxidase (150 U/ml)	0.1 ml
peroxidase (500 U/ml)	0.02 ml
creatinase (dilutions ranging from 0.01 to 5.0 U/ml)	0.01 ml

The assay was carried out at 25°C and at the wavelength of 546 nm for 10 min. ΔE was calculated from the measurement points of 6 to 10 min.

After the initial enzyme activities were determined, all samples were subjected to thermoinactivation by incubating in 0.1 M potassium phosphate buffer at 56°C for 20 minutes. The residual activity of each heat-treated sample was determined as described above. An thermo-index (initial activity/residual activity) for each tested sample was used for the enzyme thermostability evaluation. As shown in Table 4 (or Figure 4), all mutants CT1m24, CT2m10, CT2m28, CTsd2, CTqc2, CT2m9 and CTsd7 are more thermostable than the wild type *Erwinia* creatinase.

Table 4 summarizes the characterization results K_M and T_M values of the novel creatinase mutants compared to native creatinase from *Erwinia* sp.

Table 4: Characteristics of creatinase and creatinase mutants

Mutant	K_M	Thermo-Index
Wildtyp	4,3 mM	0,7%
CT1m24	8,2 mM	6.3%
CT2m10	3,4 mM	3.9%
CT2m28	4,0 mM	3.9%
CTsd2	2,1 mM	39.7%
CTqc2	6,6 mM	50.8%
CT2m9	6,3 mM	62.9%

-30-

CTsd7	2,0 mM	69.2%
-------	--------	-------

As shown in the above table, the novel creatinase mutants CTsd2 and CTsd7 of the present invention have much lower K_m values and much enhanced thermostability as compared to the wild type *Erwinia* creatinase.

Example 8 Application of an improved creatinase variant of a determination for creatinine/creatine concentration

The improved creatinase variant of the present invention is according to standard procedures used in the determination of a creatine concentration upon combining it with a sacrosine oxidase, a peroxidase and other compositions for detection of hydrogen peroxide.

Moreover, if a creatininase is also used, the concentration of creatinine can also be determined using the improved creatinase variants according to the present invention. In case in the sample investigated creatine as well as creatinine are present, the sum of both analytes is measured.

The reagent for the determination of creatine according to the present invention contains each the ingredients in an optimized concentration. An example is a reagent comprising creatinase at 16 U/ml, sacrosine oxidase at 9.8 U/ml, peroxidase at 0.81 U/ml, a coupler reagent at 0.65 mM, and a color generating substance at 7.0 mM. The buffer system used here is a potassium phosphate buffer with a concentration at 0.1 M and pH7.8. The assay usually is carried out at 25°C and color development is measured at the wavelength of 546 nm for 10 minutes. ΔE is calculated from the measurement points in the linear region and used for determining creatine concentration.

List of References

- Ausubel, F., et al. in "Current Protocols in Molecular Biology" (1994) Wiley Verlag
- Beyer, C., et al., Clin Chem 39 (1993) 1743-4
- Cadwell, R. C. Joyce, G. F., PCR Methods Appl 3 (1994) 136-40
- Feng, D. F. Doolittle, R. F., J Mol Evol 25 (1987) 351-60
- Fujita, T., et al., Clin Chem 39 (1993) 2130-6
- Guder, W. G., et al., J Clin Chem Clin Biochem 24 (1986) 889-902.
- Hong, M. C., et al., Biochem Genet 36 (1998) 407-15
- Kaplan, A. Naugler, D., Mol Cell Biochem 3 (1974) 9-15
- Kaplan, A. Szabo, L. L., Mol Cell Biochem 3 (1974) 17-25
- Khan, G. F. Wernet, W., Anal. Chim. Acta 351 (1997) 151-158
- Kim, E. J., et al., Anal. Chim. Acta 394 (1999) 225-231
- Kopetzki, E., et al., Clin Chem 40 (1994) 688-704
- Koyama, Y., et al., Agric Biol Chem 54 (1990) 1453-7
- Matsuda, Y., et al., Chem. Pharm. Bull. 34 (1986) 2155-2160
- Nishiya, Y., et al., Mol Gen Genet 257 (1998) 581-6
- Sambrook, J., et al. - in "Molecular Cloning: A Laboratory Manual" (1989) -, Cold Spring Harbour, NY, Cold Spring Harbour Laboratory Press
- Schumacher, G., et al., Ann Biol Clin 51 (1993) 815-9
- Schumann, J., et al., Biol Chem Hoppe Seyler 374 (1993) 427-34
- Schumann, J., et al., Protein Sci 2 (1993) 1612-20
- Siedel, J., et al., Clinical Chemistry 30 (1984) 968-969
- Spencer, K., Ann Clin Biochem 23 (1986) 1-25

-32-

Suzuki, K., et al., Journal of Fermentation & Bioengineering 76 (1993) 77-81

DE 2 659 878

DE 3 024 915

EP 0 291 055

EP 0 790 303

US 4,420,562

US 5,451,520

EP 790 303

-33-

SEQUENCE LISTING

<110> Roche Diagnostics GmbH
F. HOFFMANN-LA ROCHE AG

<120> Creatinase variants

<130>20981 EP

<140>

<141>

<160> 32

<170> PatentIn Ver. 2.1

<210> 1

<211> 1215

<212> DNA

<213> Erwinia sp.

<220>

<221> CDS

<222> (1)..(1215)

<400> 1

atg act gac gac atg ttg cac gtg atg aaa tgg cac aat ggt gag aag 48

Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
1 5 10 15

gaa tat tcc ccc ttt tcc gat gcc gag atg acg cgc cgc cag agt gac 96

Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
20 25 30

gtg cgg cgc tgg atg gcc gaa aac gac gtc gac gct gcg ctg ttc acc 144

Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
35 40 45

tcc tat cat tgc atc aat tac tac tct gga ttc ctg tac tgc tat ttc 192

Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Phe Leu Tyr Cys Tyr Phe
50 55 60

ggc cgc aaa tac ggc atg gtc atc gac cag gac cat gcc acg acc atc 240

Gly Arg Lys Tyr Gly Met Val Ile Asp Gln Asp His Ala Thr Thr Ile
65 70 75 80

tcg gcc ggc atc gat ggc ggt cag ccc tgg cgc cgt agc ttc ggc gac 288

Ser Ala Gly Ile Asp Gly Gly Gln Pro Trp Arg Arg Ser Phe Gly Asp
85 90 95

-34-

aac atc acc tat acg gac tgg cgc cgc gac aac ttc tac cag gcc gtc	336
Asn Ile Thr Tyr Thr Asp Trp Arg Arg Asp Asn Phe Tyr Gln Ala Val	
100 105 110	
cgc caa ctc acc ccc ggc gcc agg cgc atc ggt atc gag ttc gat cac	384
Arg Gln Leu Thr Pro Gly Ala Arg Arg Ile Gly Ile Glu Phe Asp His	
115 120 125	
gtg aac ctt gac ttc cgc cgc acg ctc gaa gag gcg ctg ccc ggc gtc	432
Val Asn Leu Asp Phe Arg Arg Thr Leu Glu Glu Ala Leu Pro Gly Val	
130 135 140	
gag ttc gtc gat atc ggt caa ccg tcg atg tgg atg cgc acg gtc aag	480
Glu Phe Val Asp Ile Gly Gln Pro Ser Met Trp Met Arg Thr Val Lys	
145 150 155 160	
tcg ctc gaa gag cag aag ctg atc cgc gag ggt gcc cgt atc tgc gac	528
Ser Leu Glu Glu Gln Lys Leu Ile Arg Glu Gly Ala Arg Ile Cys Asp	
165 170 175	
gtc ggc ggt gcc gcc tgc gtt gcc gcc gtc aag gcc ggc gtt ccg gag	576
Val Gly Gly Ala Ala Cys Val Ala Ala Val Lys Ala Gly Val Pro Glu	
180 185 190	
cac gag gtc gcg atc gcc aca acc aat gcg atg gtc cgc gag atc gcc	624
His Glu Val Ala Ile Ala Thr Thr Asn Ala Met Val Arg Glu Ile Ala	
195 200 205	
aag tcg ttc ccc ttc gtc gaa ctg atg gac acc tgg acc tgg ttc cag	672
Lys Ser Phe Pro Phe Val Glu Leu Met Asp Thr Trp Thr Trp Phe Gln	
210 215 220	
tcg ggc atc aac acc gac ggc gcc cac aat ccg gtg acc aac cgc atc	720
Ser Gly Ile Asn Thr Asp Gly Ala His Asn Pro Val Thr Asn Arg Ile	
225 230 235 240	
gtg caa tcg ggc gat atc ctg tcg ctc aac acg ttc ccg atg atc ttc	768
Val Gln Ser Gly Asp Ile Leu Ser Leu Asn Thr Phe Pro Met Ile Phe	
245 250 255	
ggc tac tac acg gcg ctg gag cga acg ctg ttc tgc gac cac gtc gac	816
Gly Tyr Tyr Thr Ala Leu Glu Arg Thr Leu Phe Cys Asp His Val Asp	
260 265 270	

-35-

gat gcc agc ctt gac atc tgg gag aag aac gtc gcc gtg cac cgc cgc 864
Asp Ala Ser Leu Asp Ile Trp Glu Lys Asn Val Ala Val His Arg Arg
275 280 285
ggc ctc gaa ctc atc aag ccg ggt gcg cgc tgc aag gat atc gcc atc 912
Gly Leu Glu Leu Ile Lys Pro Gly Ala Arg Cys Lys Asp Ile Ala Ile
290 295 300
gaa ctc aac gag atg tac cgg gag tgg gat ctg ctg aag tac cgc tcc 960
Glu Leu Asn Glu Met Tyr Arg Glu Trp Asp Leu Leu Lys Tyr Arg Ser
305 310 315 320
ttc ggc tac ggc cac tcc ttc ggc gtg ctc tcc cac tac tac ggc cgc 1008
Phe Gly Tyr Gly His Ser Phe Gly Val Leu Ser His Tyr Tyr Gly Arg
325 330 335
gag gcg ggc gtg gaa ctg cgc gag gac atc gat acc gtg ctg cag ccc 1056
Glu Ala Gly Val Glu Leu Arg Glu Asp Ile Asp Thr Val Leu Gln Pro
340 345 350
ggc atg gtg gtc tcc atg gag ccg atg gtg atg ttg cca gaa ggc gct 1104
Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala
355 360 365
ccc ggc gcg ggc ggc tac cgc gag cac gac atc ctg atc gtg ggg gaa 1152
Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu
370 375 380
gac ggc gca gag aac att acc gga ttc ccc ttc ggg cct gag cac aac 1200
Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Phe Gly Pro Glu His Asn
385 390 395 400
atc atc cgc aac tga 1215
Ile Ile Arg Asn
405

<210> 2

<211> 404

<212> PRT

<213> Erwinia sp.

<400> 2

Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
1 5 10 15

-36-

Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp
			20					25					30		
Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr
		35					40					45			
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe
	50					55					60				
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile
65					70					75					80
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp
				85				90						95	
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val
			100					105					110		
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His
		115					120					125			
Val	Asn	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val
	130					135					140				
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys
145					150					155					160
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp
				165				170						175	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu
			180					185					190		
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala
		195					200					205			
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln
	210					215					220				
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile
225					230					235					240
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe
				245				250						255	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp
			260					265					270		
Asp	Ala	Ser	Leu	Asp	Ile	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg
		275					280					285			
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile
	290					295					300				
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser
305					310					315					320
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg
				325					330					335	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro
			340					345					350		
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala
		355					360					365			
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu
	370					375					380				
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn
385					390					395					400
Ile	Ile	Arg	Asn												

<210> 3

<211> 7

<212> PRT

-37-

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide 1

<400> 3

Tyr Ser Gly Xaa Leu Tyr Cys
1 5

<210> 4

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide 2

<400> 4

Asp His Val Xaa Leu Asp Phe
1 5

<210> 5

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide 3

<400> 5

Thr Asn Ala Xaa Val Arg Glu
1 5

<210> 6

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide 4

<400> 6

Ser Leu Asp Xaa Trp Glu Lys
1 5

<210> 7

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

-38-

<223> Description of Artificial Sequence:peptide 5

<400> 7

Asp	Ile	Ala	Xaa	Glu	Leu	Asn
1				5		

<210> 8

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide 6

<400> 8

Gly	Phe	Pro	Xaa	Gly	Pro	Glu
1				5		

<210> 9

<211> 1215

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:variant CT1m24

<220>

<221> CDS

<222> (1)..(1215)

<400> 9

atg	act	gac	gac	atg	ttg	cac	gtg	atg	aaa	tgg	cac	aat	ggt	gag	aag	48
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys
1				5					10					15	

gaa	tat	tcc	ccc	ttt	tcc	gat	gcc	gag	atg	acg	cgc	cgc	cag	agt	gac	96
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp
			20					25					30		

gtg	cgg	cgc	tgg	atg	gcc	gaa	aac	gac	gtc	gac	gct	gcg	ctg	ttc	acc	144
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr
			35				40				45				

tcc	tat	cat	tgc	atc	aac	tac	tac	tct	gga	ttc	ctg	tac	tgc	tat	ttc	192
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe
	50					55					60				

ggc	cgc	aaa	tac	ggc	atg	gtc	atc	gac	cag	gac	cat	gcc	acg	acc	atc	240
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-39-

Gly 65	Arg	Lys	Tyr	Gly	Met 70	Val	Ile	Asp	Gln	Asp 75	His	Ala	Thr	Thr	Ile 80	
tcg	gcc	ggc	atc	gat	ggc	ggt	cag	ccc	tgg	cgc	cgt	agc	ttc	ggc	gac	288
Ser	Ala	Gly	Ile	Asp 85	Gly	Gly	Gln	Pro	Trp 90	Arg	Arg	Ser	Phe	Gly 95	Asp	
aac	atc	acc	tat	acg	gac	tgg	cgc	cgc	gac	aac	ttc	tac	cag	gcc	gtc	336
Asn	Ile	Thr	Tyr 100	Thr	Asp	Trp	Arg	Arg 105	Asp	Asn	Phe	Tyr	Gln 110	Ala	Val	
cgc	caa	ctc	acc	ccc	ggc	gcc	agg	cgc	atc	ggt	atc	gag	ttc	gat	cac	384
Arg	Gln 115	Leu	Thr	Pro	Gly	Ala	Arg 120	Arg	Ile	Gly	Ile	Glu 125	Phe	Asp	His	
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gcg	ctg	ccc	ggc	gtc	432
Val 130	Asp	Leu	Asp	Phe	Arg	Arg 135	Thr	Leu	Glu	Glu 140	Ala	Leu	Pro	Gly	Val	
gag	ttc	gtc	gat	atc	ggt	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480
Glu 145	Phe	Val	Asp	Ile	Gly 150	Gln	Pro	Ser	Met	Trp 155	Met	Arg	Thr	Val	Lys 160	
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggt	gcc	cgt	atc	tgc	gac	528
Ser	Leu	Glu	Glu	Gln 165	Lys	Leu	Ile	Arg	Glu 170	Gly	Ala	Arg	Ile	Cys 175	Asp	
gtc	ggc	ggt	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576
Val	Gly	Gly	Ala 180	Ala	Cys	Val	Ala 185	Ala	Val	Lys	Ala	Gly 190	Val	Pro	Glu	
cac	gag	gtc	gcg	atc	gcc	aca	acc	aat	gcg	atg	gtc	cgc	gag	atc	gcc	624
His	Glu 195	Val	Ala	Ile	Ala	Thr	Thr 200	Asn	Ala	Met	Val	Arg 205	Glu	Ile	Ala	
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672
Lys 210	Ser	Phe	Pro	Phe	Val	Glu 215	Leu	Met	Asp	Thr	Trp 220	Thr	Trp	Phe	Gln	
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720
Ser 225	Gly	Ile	Asn	Thr	Asp 230	Gly	Ala	His	Asn 235	Pro	Val	Thr	Asn	Arg	Ile 240	
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768

-40-

Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe		
				245					250					255			
ggc	tac	tac	acg	gcg	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp		
			260					265					270				
gat	gcc	agc	ctt	gac	acc	tgg	gag	aag	aac	gtc	gcc	gtg	cac	cgc	cgc	864	
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg		
		275					280					285					
ggc	ctc	gaa	ctc	atc	aag	ccg	ggt	gcg	cgc	tgc	aag	gat	atc	gcc	atc	912	
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile		
	290					295					300						
gaa	ctc	aac	gag	atg	tac	cgg	gag	tgg	gat	ctg	ctg	aag	tac	cgc	tcc	960	
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser		
305					310					315					320		
ttc	ggc	tac	ggc	cac	tcc	ttc	ggc	gtg	ctc	tcc	cac	tac	tac	ggc	cgc	1008	
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg		
				325					330					335			
gag	gcg	ggc	gtg	gaa	ctg	cgc	gag	gac	atc	gat	acc	gtg	ctg	cag	ccc	1056	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro		
			340					345					350				
ggc	atg	gtg	gtc	tcc	atg	gag	ccg	atg	gtg	atg	ttg	cca	gaa	ggc	gct	1104	
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala		
		355					360					365					
ccc	ggc	gcg	ggc	ggc	tac	cgc	gag	cac	gac	atc	ctg	atc	gtg	ggg	gaa	1152	
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu		
		370				375					380						
gac	ggc	gca	gag	aac	att	acc	gga	ttc	ccc	ttc	ggg	cct	gag	cac	aac	1200	
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn		
385					390				395						400		
atc	atc	cgc	aac	tga												1215	
Ile	Ile	Arg	Asn														
				405													

<210> 10

-41-

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CT1m24

<400> 10

Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys
1				5				10						15	
Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp
			20					25					30		
Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr
			35				40					45			
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe
	50					55					60				
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile
65					70					75					80
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp
				85					90					95	
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val
			100					105					110		
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His
		115					120					125			
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val
	130						135				140				
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys
145					150					155					160
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp
				165					170					175	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu
			180					185					190		
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala
		195					200					205			
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln
	210					215					220				
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile
225					230					235					240
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe
			245						250					255	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp
			260					265					270		
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg
		275					280					285			
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile
	290					295					300				
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser
305					310					315					320
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg
			325						330					335	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro
			340					345					350		
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala
		355					360					365			
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu
	370					375					380				
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn

```
<210> 11
<211> 1212
<212> DNA
<213> Artificial Sequence
```

```
<220>
<221> CDS
<222> (1)..(1212)
```

<400> 11																
atg	act	gac	gac	atg	ttg	cac	gtg	atg	aaa	tgg	cac	aat	ggt	gag	aag	48
Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys	
1				5					10					15		
gaa	tat	tcc	ccc	ttt	tcc	gat	gcc	gag	atg	acg	cgc	cgc	cag	agt	gac	96
Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp	
			20					25					30			
gtg	cgg	cgc	tgg	atg	gcc	gaa	aac	gac	gtc	gac	gct	gcg	ctg	ttc	acc	144
Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr	
		35					40					45				
tcc	tat	cat	tgc	atc	aac	tac	tac	tct	gga	ttc	ctg	tac	tgc	tat	ttc	192
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe	
	50					55					60					
ggc	cgc	aaa	tac	ggc	atg	gtc	atc	gac	cag	gac	cat	gcc	acg	acc	atc	240
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile	
65					70					75					80	
tcg	gcc	ggc	atc	gat	ggc	ggg	cag	ccc	tgg	cgc	cgt	agc	ttc	ggc	gac	288
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp	
				85					90					95		
aac	atc	acc	tat	acg	gac	tgg	cgc	cgc	gac	aac	ttc	tac	cag	gcc	gtc	336
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val	
			100					105					110			
cgc	caa	ctc	acc	ccc	ggc	gcc	agg	cgc	atc	ggt	atc	gaq	ttc	qat	cac	384

-43-

Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His		
	115						120					125					
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gcg	ctg	ccc	ggc	gtc	432	
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val		
	130					135					140						
gag	ttc	gtc	gat	atc	ggc	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480	
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys		
145					150					155					160		
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggc	gcc	cgt	atc	tgc	gac	528	
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp		
				165					170					175			
gtc	ggc	ggc	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu		
			180					185					190				
cac	gag	gtc	gag	atc	gcc	aca	acc	aat	gag	gtg	gtc	cgc	gag	atc	gcc	624	
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Val	Val	Arg	Glu	Ile	Ala		
	195						200					205					
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672	
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln		
	210					215					220						
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720	
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile		
225					230					235					240		
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768	
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe		
				245					250					255			
ggc	tac	tac	acg	gag	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp		
			260					265					270				
gat	gcc	agc	ctt	gac	acc	tgg	gag	aag	aac	gtc	gcc	gtg	cac	cgc	cgc	864	
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg		
	275						280					285					
ggc	ctc	gaa	ctc	atc	aag	ccg	ggc	gag	cgc	tgc	aag	gat	atc	gcc	atc	912	

-44-

Gly Leu Glu Leu Ile Lys Pro Gly Ala Arg Cys Lys Asp Ile Ala Ile
 290 295 300
 gaa ctc aac gag atg tac cgg gag tgg gat ctg ctg aag tac cgc tcc 960
 Glu Leu Asn Glu Met Tyr Arg Glu Trp Asp Leu Leu Lys Tyr Arg Ser
 305 310 315 320
 ttc ggc tac ggc cac tcc ttc ggc gtg ctc tcc cac tac tac ggc cgc 1008
 Phe Gly Tyr Gly His Ser Phe Gly Val Leu Ser His Tyr Tyr Gly Arg
 325 330 335
 gag gcg ggc gtg gaa ctg cgc gag gac atc gat acc gtg ctg cag ccc 1056
 Glu Ala Gly Val Glu Leu Arg Glu Asp Ile Asp Thr Val Leu Gln Pro
 340 345 350
 ggc atg gtg gtc tcc atg gag ccg atg gtg atg ttg cca gaa ggc gct 1104
 Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala
 355 360 365
 ccc ggc gcg ggc ggc tac cgc gag cac gac atc ctg atc gtg ggg gaa 1152
 Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu
 370 375 380
 gac ggc gca gag aac att acc gga ttc ccc ttc ggg cct gag cac aac 1200
 Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Phe Gly Pro Glu His Asn
 385 390 395 400
 atc atc cgc aac 1212
 Ile Ile Arg Asn

<210> 12

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CT2m9

<400> 12

Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
 1 5 10 15

Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
 20 25 30

Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
 35 40 45

Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Phe Leu Tyr Cys Tyr Phe

-45-

50					55					60					
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile
65					70					75					80
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp
				85					90					95	
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val
			100					105					110		
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His
		115					120					125			
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val
	130					135					140				
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys
145					150					155					160
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp
				165					170					175	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu
			180					185					190		
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Val	Val	Arg	Glu	Ile	Ala
		195					200					205			
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln
	210					215					220				
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile
225					230					235					240
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe
				245					250					255	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp
			260					265					270		
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg
		275					280					285			
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile
	290					295					300				
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser
305					310					315					320
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg
				325					330					335	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro
			340					345					350		

-46-

Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala
 355 360 365

Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu
 370 375 380

Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Phe Gly Pro Glu His Asn
 385 390 395 400

Ile Ile Arg Asn

<210> 13
 <211> 1212
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:variant CT2m10

<220>
 <221> CDS
 <222> (1)..(1212)

<400> 13
 atg act gac gac atg ttg cac gtg atg aaa tgg cac aat ggt gag aag 48
 Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
 1 5 10 15
 gaa tat tcc ccc ttt tcc gat gcc gag atg acg cgc cgc cag agt gac 96
 Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
 20 25 30
 gtg cgg cgc tgg atg gcc gaa aac gac gtc gac gct gcg ctg ttc acc 144
 Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
 35 40 45
 tcc tat cat tgc atc aac tac tac tct gga ttc ctg tac tgc tat ttc 192
 Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Phe Leu Tyr Cys Tyr Phe
 50 55 60
 ggc cgc aaa tac ggc atg gtc atc gac cag gac cat gcc acg acc atc 240
 Gly Arg Lys Tyr Gly Met Val Ile Asp Gln Asp His Ala Thr Thr Ile
 65 70 75 80
 tcg gcc ggc atc gat ggc ggt cag ccc tgg cgc cgt agc ttc ggc gac 288

-47-

Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp	
				85					90					95		
aac	atc	acc	tat	acg	gac	tgg	cgc	cgc	gac	aac	ttc	tac	cag	gcc	gtc	336
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val	
			100					105					110			
cgc	caa	ctc	acc	ccc	ggc	gcc	agg	cgc	atc	ggg	atc	gag	ttc	gat	cac	384
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His	
		115					120					125				
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gcg	ctg	ccc	ggc	gtc	432
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val	
	130					135					140					
gag	ttc	gtc	gat	atc	ggg	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys	
145					150					155					160	
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggg	gcc	cgt	atc	tgc	gac	528
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp	
				165					170					175		
gtc	ggc	ggg	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu	
			180					185					190			
cac	gag	gtc	gcg	atc	gcc	aca	acc	aat	gcg	atg	gtc	cgc	gag	atc	gcc	624
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala	
		195					200					205				
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln	
	210					215					220					
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile	
225					230					235				240		
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe	
				245					250					255		
ggc	tac	tac	acg	gcg	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816

-48-

Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp		
			260					265					270				
gat gcc agc ctt gac acc tgg gag aag aac gtc gcc gtg cac cgc cgc 864																	
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg		
		275					280					285					
ggc ctc gaa ctc atc aag ccg ggt gcg cgc tgc aag gat atc gcc atc 912																	
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile		
	290					295					300						
gaa ctc aac gag atg tac cgg gag tgg gat ctg ctg aag tac cgc tcc 960																	
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser		
305					310					315					320		
ttc ggc tac ggc cac tcc ttc ggc gtg ctc tcc cac tac tac ggc cgc 1008																	
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg		
				325					330					335			
gag gcg ggc gtg gaa ctg cgc gag gac atc gat acc gtg ctg cag ccc 1056																	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro		
			340					345					350				
ggc atg gtg gtc tcc atg gag ccg atg gtg atg ttg cca gaa ggc gct 1104																	
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala		
		355					360					365					
ccc ggc gcg ggc ggc tac cgc gag cac gac atc ctg atc gtg ggg gaa 1152																	
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu		
	370					375					380						
gac ggc gca gag aac att acc gga ttc ccc ctc ggg cct gag cac aac 1200																	
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Leu	Gly	Pro	Glu	His	Asn		
385					390					395					400		
atc atc cgc aac 1212																	
Ile Ile Arg Asn																	

<210> 14

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CT2m10

<400> 14

Met 1	Thr	Asp	Asp	Met 5	Leu	His	Val	Met	Lys 10	Trp	His	Asn	Gly	Glu 15	Lys
Glu	Tyr	Ser	Pro 20	Phe	Ser	Asp	Ala	Glu 25	Met	Thr	Arg	Arg	Gln 30	Ser	Asp
Val	Arg	Arg 35	Trp	Met	Ala	Glu	Asn 40	Asp	Val	Asp	Ala	Ala 45	Leu	Phe	Thr
Ser	Tyr 50	His	Cys	Ile	Asn	Tyr 55	Tyr	Ser	Gly	Phe	Leu 60	Tyr	Cys	Tyr	Phe
Gly 65	Arg	Lys	Tyr	Gly	Met 70	Val	Ile	Asp	Gln	Asp 75	His	Ala	Thr	Thr	Ile 80
Ser	Ala	Gly	Ile	Asp 85	Gly	Gly	Gln	Pro	Trp 90	Arg	Arg	Ser	Phe	Gly 95	Asp
Asn	Ile	Thr	Tyr 100	Thr	Asp	Trp	Arg	Arg 105	Asp	Asn	Phe	Tyr	Gln 110	Ala	Val
Arg	Gln	Leu 115	Thr	Pro	Gly	Ala	Arg 120	Arg	Ile	Gly	Ile	Glu 125	Phe	Asp	His
Val	Asp 130	Leu	Asp	Phe	Arg	Arg 135	Thr	Leu	Glu	Glu	Ala 140	Leu	Pro	Gly	Val
Glu 145	Phe	Val	Asp	Ile	Gly 150	Gln	Pro	Ser	Met	Trp 155	Met	Arg	Thr	Val	Lys 160
Ser	Leu	Glu	Glu	Gln 165	Lys	Leu	Ile	Arg	Glu 170	Gly	Ala	Arg	Ile	Cys 175	Asp
Val	Gly	Gly	Ala 180	Ala	Cys	Val	Ala 185	Ala	Val	Lys	Ala	Gly	Val 190	Pro	Glu
His	Glu	Val 195	Ala	Ile	Ala	Thr	Thr 200	Asn	Ala	Met	Val	Arg 205	Glu	Ile	Ala
Lys	Ser 210	Phe	Pro	Phe	Val	Glu 215	Leu	Met	Asp	Thr	Trp 220	Thr	Trp	Phe	Gln
Ser 225	Gly	Ile	Asn	Thr	Asp 230	Gly	Ala	His	Asn 235	Pro	Val	Thr	Asn	Arg	Ile 240
Val	Gln	Ser	Gly	Asp 245	Ile	Leu	Ser	Leu	Asn 250	Thr	Phe	Pro	Met	Ile 255	Phe
Gly	Tyr	Tyr	Thr 260	Ala	Leu	Glu	Arg	Thr 265	Leu	Phe	Cys	Asp	His 270	Val	Asp
Asp	Ala	Ser 275	Leu	Asp	Thr	Trp	Glu 280	Lys	Asn	Val	Ala	Val 285	His	Arg	Arg
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile

-50-

290	295	300
Glu Leu Asn Glu Met Tyr Arg Glu Trp Asp Leu Leu Lys Tyr Arg Ser 305 310 315 320		
Phe Gly Tyr Gly His Ser Phe Gly Val Leu Ser His Tyr Tyr Gly Arg 325 330 335		
Glu Ala Gly Val Glu Leu Arg Glu Asp Ile Asp Thr Val Leu Gln Pro 340 345 350		
Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala 355 360 365		
Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu 370 375 380		
Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Leu Gly Pro Glu His Asn 385 390 395 400		
Ile Ile Arg Asn		

<210> 15
 <211> 1215
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:variant CT2m28

<220>
 <221> CDS
 <222> (1)..(1215)

<400> 15	
atg act gac gac atg ttg cac gtg atg aaa tgg cac aat ggt gag aag	48
Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys	
1 5 10 15	
gaa tat tcc ccc ttt tcc gat gcc gag atg acg cgc cgc cag agt gac	96
Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp	
20 25 30	
gtg cgg cgc tgg atg gcc gaa aac gac gtc gac gct gcg ctg ttc acc	144
Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr	
35 40 45	
tcc tat cat tgc atc aac tac tac tct gga ttc ctg tac tgc tat ttc	192

-51-

Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe	
50						55					60					
ggc	cgc	aaa	tac	ggc	atg	gtc	atc	gac	cag	gac	cat	gcc	acg	acc	atc	240
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile	
65					70				75						80	
tcg	gcc	ggc	atc	gat	ggc	ggc	cag	ccc	tgg	cgc	cgt	agc	ttc	ggc	gac	288
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp	
				85					90					95		
aac	atc	acc	tat	acg	gac	tgg	cgc	cgc	gac	aac	ttc	tac	cag	gcc	gtc	336
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val	
			100					105					110			
cgc	caa	ctc	acc	ccc	ggc	gcc	agg	cgc	atc	ggc	atc	gag	ttc	gat	cac	384
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His	
		115					120					125				
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gag	ctg	ccc	ggc	gtc	432
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val	
	130					135					140					
gag	ttc	gtc	gat	atc	ggc	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys	
145					150					155					160	
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggc	gcc	cgt	atc	tgc	gac	528
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp	
				165					170					175		
gtc	ggc	ggc	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu	
			180					185					190			
cac	gag	gtc	gag	atc	gcc	aca	acc	aat	gag	atg	gtc	cgc	gag	atc	gcc	624
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala	
	195						200					205				
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln	
	210					215					220					
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720

[illegible]

-53-

Ile Ile Arg Asn

405

<210> 16

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CT2m28

<400> 16

Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys
1				5				10						15	
Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp
			20					25					30		
Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr
			35				40					45			
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Phe	Leu	Tyr	Cys	Tyr	Phe
	50					55					60				
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile
65					70					75					80
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp
				85					90					95	
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val
			100					105					110		
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His
		115					120					125			
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val
	130					135					140				
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys
145					150					155					160
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp
				165					170					175	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu
			180					185					190		
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala
		195					200					205			
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln
	210					215					220				
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile
225					230					235					240
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe
				245					250					255	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp
			260					265					270		
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg
		275					280					285			
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Leu
	290					295					300				
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser
305					310					315					320
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg
				325					330					335	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro
			340					345					350		

-54-

Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala
		355					360					365			
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu
	370					375					380				
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn
385					390					395					400
Ile	Ile	Arg	Asn												

<210> 17

<211> 1215

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:variant CTqc2

<220>

<221> CDS

<222> (1)..(1215)

<400> 17

atg	act	gac	gac	atg	ttg	cac	gtg	atg	aaa	tgg	cac	aat	ggt	gag	aag	48
Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys	
1				5					10					15		
gaa	tat	tcc	ccc	ttt	tcc	gat	gcc	gag	atg	acg	cgc	cgc	cag	agt	gac	96
Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp	
			20					25					30			
gtg	cgg	cgc	tgg	atg	gcc	gaa	aac	gac	gtc	gac	gct	gcg	ctg	ttc	acc	144
Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr	
			35				40					45				
tcc	tat	cat	tgc	atc	aac	tac	tac	tct	gga	tgg	ctg	tac	tgc	tat	ttc	192
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Trp	Leu	Tyr	Cys	Tyr	Phe	
	50					55					60					
ggc	cgc	aaa	tac	ggc	atg	gtc	atc	gac	cag	gac	cat	gcc	acg	acc	atc	240
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile	
65					70				75						80	
tcg	gcc	ggc	atc	gat	ggc	ggt	cag	ccc	tgg	cgc	cgt	agc	ttc	ggc	gac	288
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp	
				85					90					95		
aac	atc	acc	tat	acg	gac	tgg	cgc	cgc	gac	aac	ttc	tac	cag	gcc	gtc	336

-55-

Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val	
		100						105					110			
cgc	caa	ctc	acc	ccc	ggc	gcc	agg	cgc	atc	ggt	atc	gag	ttc	gat	cac	384
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His	
	115						120					125				
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gcg	ctg	ccc	ggc	gtc	432
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val	
	130					135					140					
gag	ttc	gtc	gat	atc	ggt	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys	
145					150					155					160	
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggt	gcc	cgt	atc	tgc	gac	528
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp	
				165					170					175		
gtc	ggc	ggt	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu	
			180				185						190			
cac	gag	gtc	gcg	atc	gcc	aca	acc	aat	gcg	atg	gtc	cgc	gag	atc	gcc	624
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala	
	195						200					205				
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln	
	210					215					220					
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile	
225					230					235					240	
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe	
				245					250					255		
ggc	tac	tac	acg	gcg	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp	
			260				265						270			
gat	gcc	agc	ctt	gac	acc	tgg	gag	aag	aac	gtc	gcc	gtg	cac	cgc	cgc	864

-56-

Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg		
		275					280					285					
ggc	ctc	gaa	ctc	atc	aag	ccg	ggt	gcg	cgc	tgc	aag	gat	atc	gcc	atc	912	
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile		
	290					295					300						
gaa	ctc	aac	gag	atg	tac	cgg	gag	tgg	gat	ctg	ctg	aag	tac	cgc	tcc	960	
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser		
305					310					315					320		
ttc	ggc	tac	ggc	cac	tcc	ttc	ggc	gtg	ctc	tcc	cac	tac	tac	ggc	cgc	1008	
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg		
				325					330					335			
gag	gcg	ggc	gtg	gaa	ctg	cgc	gag	gac	atc	gat	acc	gtg	ctg	cag	ccc	1056	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro		
			340					345					350				
ggc	atg	gtg	gtc	tcc	atg	gag	ccg	atg	gtg	atg	ttg	cca	gaa	ggc	gct	1104	
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala		
	355						360					365					
ccc	ggc	gcg	ggc	ggc	tac	cgc	gag	cac	gac	atc	ctg	atc	gtg	ggg	gaa	1152	
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu		
	370					375					380						
gac	ggc	gca	gag	aac	att	acc	gga	ttc	ccc	ttc	ggg	cct	gag	cac	aac	1200	
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn		
385					390					395					400		
atc	atc	cgc	aac	tga												1215	
Ile	Ile	Arg	Asn														
			405														

<210> 18

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CTqc2

<400> 18

Met	Thr	Asp	Asp	Met	Leu	His	Val	Met	Lys	Trp	His	Asn	Gly	Glu	Lys		
1				5				10						15			
Glu	Tyr	Ser	Pro	Phe	Ser	Asp	Ala	Glu	Met	Thr	Arg	Arg	Gln	Ser	Asp		
			20					25					30				

-57-

Val	Arg	Arg	Trp	Met	Ala	Glu	Asn	Asp	Val	Asp	Ala	Ala	Leu	Phe	Thr
		35					40					45			
Ser	Tyr	His	Cys	Ile	Asn	Tyr	Tyr	Ser	Gly	Trp	Leu	Tyr	Cys	Tyr	Phe
	50					55					60				
Gly	Arg	Lys	Tyr	Gly	Met	Val	Ile	Asp	Gln	Asp	His	Ala	Thr	Thr	Ile
65					70					75					80
Ser	Ala	Gly	Ile	Asp	Gly	Gly	Gln	Pro	Trp	Arg	Arg	Ser	Phe	Gly	Asp
				85					90					95	
Asn	Ile	Thr	Tyr	Thr	Asp	Trp	Arg	Arg	Asp	Asn	Phe	Tyr	Gln	Ala	Val
			100					105					110		
Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His
		115						120					125		
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val
	130						135				140				
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys
145					150					155					160
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp
				165					170					175	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu
			180					185					190		
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Met	Val	Arg	Glu	Ile	Ala
		195					200					205			
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln
	210					215					220				
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile
225					230					235					240
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe
				245					250					255	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp
			260					265					270		
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg
		275					280					285			
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile
	290					295					300				
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser
305					310					315					320
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg
				325					330					335	
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro
			340					345					350		
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala
		355					360					365			
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu
	370					375					380				
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Phe	Gly	Pro	Glu	His	Asn
385					390					395					400
Ile	Ile	Arg	Asn												

<210> 19

<211> 1212

<212> DNA

<213> Artificial Sequence

-58-

<220>

<223> Description of Artificial Sequence:variant CTsd2

<220>

<221> CDS

<222> (1)..(1212)

<400> 19

```

atg act gac gac atg ttg cac gtg atg aaa tgg cac aat ggt gag aag      48
Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
  1          5          10          15

gaa tat tcc ccc ttt tcc gat gcc gag atg acg cgc cgc cag agt gac      96
Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
          20          25          30

gtg cgg cgc tgg atg gcc gaa aac gac gtc gac gct gcg ctg ttc acc     144
Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
          35          40          45

tcc tat cat tgc atc aac tac tac tct gga ttc ctg tac tgc tat ttc     192
Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Phe Leu Tyr Cys Tyr Phe
          50          55          60

ggc cgc aaa tac ggc atg gtc atc gac cag gac cat gcc acg acc atc     240
Gly Arg Lys Tyr Gly Met Val Ile Asp Gln Asp His Ala Thr Thr Ile
          65          70          75          80

tcg gcc ggc atc gat ggc ggt cag ccc tgg cgc cgt agc ttc ggc gac     288
Ser Ala Gly Ile Asp Gly Gly Gln Pro Trp Arg Arg Ser Phe Gly Asp
          85          90          95

aac atc acc tat acg gac tgg cgc cgc gac aac ttc tac cag gcc gtc     336
Asn Ile Thr Tyr Thr Asp Trp Arg Arg Asp Asn Phe Tyr Gln Ala Val
          100          105          110

cgc caa ctc acc ccc ggc gcc agg cgc atc ggt atc gag ttc gat cac     384
Arg Gln Leu Thr Pro Gly Ala Arg Arg Ile Gly Ile Glu Phe Asp His
          115          120          125

gtg gac ctt gac ttc cgc cgc acg ctc gaa gag gcg ctg ccc ggc gtc     432
Val Asp Leu Asp Phe Arg Arg Thr Leu Glu Glu Ala Leu Pro Gly Val
          130          135          140

gag ttc gtc gat atc ggt caa ccg tcg atg tgg atg cgc acg gtc aag     480

```

-59-

Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys	
145					150					155					160	
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggt	gcc	cgt	atc	tgc	gac	528
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp	
				165					170					175		
gtc	ggc	ggt	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu	
			180					185					190			
cac	gag	gtc	gcg	atc	gcc	aca	acc	aat	gcg	gtg	gtc	cgc	gag	atc	gcc	624
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Val	Val	Arg	Glu	Ile	Ala	
		195					200					205				
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln	
	210					215					220					
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile	
225					230					235					240	
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe	
				245					250					255		
ggc	tac	tac	acg	gcg	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp	
			260					265					270			
gat	gcc	agc	ctt	gac	acc	tgg	gag	aag	aac	gtc	gcc	gtg	cac	cgc	cgc	864
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg	
		275					280					285				
ggc	ctc	gaa	ctc	atc	aag	ccg	ggt	gcg	cgc	tgc	aag	gat	atc	gcc	atc	912
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile	
	290					295					300					
gaa	ctc	aac	gag	atg	tac	cgg	gag	tgg	gat	ctg	ctg	aag	tac	cgc	tcc	960
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser	
305					310					315					320	
ttc	ggc	tac	ggc	cac	tcc	ttc	ggc	gtg	ctc	tcc	cac	tac	tac	ggc	cgc	1008

```

<400> 20
Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
  1             5             10             15

Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
          20             25             30

Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
          35             40             45

Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Phe Leu Tyr Cys Tyr Phe
  50             55             60

Gly Arg Lys Tyr Gly Met Val Ile Asp Gln Asp His Ala Thr Thr Ile
  65             70             75             80

Ser Ala Gly Ile Asp Gly Gly Gln Pro Trp Arg Arg Ser Phe Gly Asp
          85             90             95

Asn Ile Thr Tyr Thr Asp Trp Arg Arg Asp Asn Phe Tyr Gln Ala Val
          100             105             110

```

-61-

Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His	115	120	125
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val	130	135	140
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys	145	150	155
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp	165	170	175
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu	180	185	190
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Val	Val	Arg	Glu	Ile	Ala	195	200	205
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln	210	215	220
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile	225	230	235
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe	245	250	255
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp	260	265	270
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg	275	280	285
Gly	Leu	Glu	Leu	Ile	Lys	Pro	Gly	Ala	Arg	Cys	Lys	Asp	Ile	Ala	Ile	290	295	300
Glu	Leu	Asn	Glu	Met	Tyr	Arg	Glu	Trp	Asp	Leu	Leu	Lys	Tyr	Arg	Ser	305	310	315
Phe	Gly	Tyr	Gly	His	Ser	Phe	Gly	Val	Leu	Ser	His	Tyr	Tyr	Gly	Arg	325	330	335
Glu	Ala	Gly	Val	Glu	Leu	Arg	Glu	Asp	Ile	Asp	Thr	Val	Leu	Gln	Pro	340	345	350
Gly	Met	Val	Val	Ser	Met	Glu	Pro	Met	Val	Met	Leu	Pro	Glu	Gly	Ala	355	360	365
Pro	Gly	Ala	Gly	Gly	Tyr	Arg	Glu	His	Asp	Ile	Leu	Ile	Val	Gly	Glu	370	375	380
Asp	Gly	Ala	Glu	Asn	Ile	Thr	Gly	Phe	Pro	Leu	Gly	Pro	Glu	His	Asn	385	390	395

-62-

Ile Ile Arg Asn

<210> 21

<211> 1212

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:variant CTsd7

<220>

<221> CDS

<222> (1)..(1212)

<400> 21

atg act gac gac atg ttg cac gtg atg aaa tgg cac aat ggt gag aag	48
Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys	
1 5 10 15	
gaa tat tcc ccc ttt tcc gat gcc gag atg acg cgc cgc cag agt gac	96
Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp	
20 25 30	
gtg cgg cgc tgg atg gcc gaa aac gac gtc gac gct gcg ctg ttc acc	144
Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr	
35 40 45	
tcc tat cat tgc atc aat tac tac tct gga tgg ctg tac tgc tat ttc	192
Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Trp Leu Tyr Cys Tyr Phe	
50 55 60	
ggc cgc aaa tac ggc atg gtc atc gac cag gac cat gcc acg acc atc	240
Gly Arg Lys Tyr Gly Met Val Ile Asp Gln Asp His Ala Thr Thr Ile	
65 70 75 80	
tcg gcc ggc atc gat ggc ggt cag ccc tgg cgc cgt agc ttc ggc gac	288
Ser Ala Gly Ile Asp Gly Gly Gln Pro Trp Arg Arg Ser Phe Gly Asp	
85 90 95	
aac atc acc tat acg gac tgg cgc cgc gac aac ttc tac cag gcc gtc	336
Asn Ile Thr Tyr Thr Asp Trp Arg Arg Asp Asn Phe Tyr Gln Ala Val	
100 105 110	
cgc caa ctc acc ccc ggc gcc agg cgc atc ggt atc gag ttc gat cac	384

-63-

Arg	Gln	Leu	Thr	Pro	Gly	Ala	Arg	Arg	Ile	Gly	Ile	Glu	Phe	Asp	His		
	115						120					125					
gtg	gac	ctt	gac	ttc	cgc	cgc	acg	ctc	gaa	gag	gcg	ctg	ccc	ggc	gtc	432	
Val	Asp	Leu	Asp	Phe	Arg	Arg	Thr	Leu	Glu	Glu	Ala	Leu	Pro	Gly	Val		
	130					135					140						
gag	ttc	gtc	gat	atc	ggc	caa	ccg	tcg	atg	tgg	atg	cgc	acg	gtc	aag	480	
Glu	Phe	Val	Asp	Ile	Gly	Gln	Pro	Ser	Met	Trp	Met	Arg	Thr	Val	Lys		
145					150					155					160		
tcg	ctc	gaa	gag	cag	aag	ctg	atc	cgc	gag	ggc	gcc	cgt	atc	tgc	gac	528	
Ser	Leu	Glu	Glu	Gln	Lys	Leu	Ile	Arg	Glu	Gly	Ala	Arg	Ile	Cys	Asp		
				165					170					175			
gtc	ggc	ggc	gcc	gcc	tgc	gtt	gcc	gcc	gtc	aag	gcc	ggc	gtt	ccg	gag	576	
Val	Gly	Gly	Ala	Ala	Cys	Val	Ala	Ala	Val	Lys	Ala	Gly	Val	Pro	Glu		
			180					185					190				
cac	gag	gtc	gag	atc	gcc	aca	acc	aat	gag	gtg	gtc	cgc	gag	atc	gcc	624	
His	Glu	Val	Ala	Ile	Ala	Thr	Thr	Asn	Ala	Val	Val	Arg	Glu	Ile	Ala		
	195						200					205					
aag	tcg	ttc	ccc	ttc	gtc	gaa	ctg	atg	gac	acc	tgg	acc	tgg	ttc	cag	672	
Lys	Ser	Phe	Pro	Phe	Val	Glu	Leu	Met	Asp	Thr	Trp	Thr	Trp	Phe	Gln		
	210					215					220						
tcg	ggc	atc	aac	acc	gac	ggc	gcc	cac	aat	ccg	gtg	acc	aac	cgc	atc	720	
Ser	Gly	Ile	Asn	Thr	Asp	Gly	Ala	His	Asn	Pro	Val	Thr	Asn	Arg	Ile		
225					230					235					240		
gtg	caa	tcg	ggc	gat	atc	ctg	tcg	ctc	aac	acg	ttc	ccg	atg	atc	ttc	768	
Val	Gln	Ser	Gly	Asp	Ile	Leu	Ser	Leu	Asn	Thr	Phe	Pro	Met	Ile	Phe		
				245					250					255			
ggc	tac	tac	acg	gag	ctg	gag	cga	acg	ctg	ttc	tgt	gac	cac	gtc	gac	816	
Gly	Tyr	Tyr	Thr	Ala	Leu	Glu	Arg	Thr	Leu	Phe	Cys	Asp	His	Val	Asp		
			260					265					270				
gat	gcc	agc	ctt	gac	acc	tgg	gag	aag	aac	gtc	gcc	gtg	cac	cgc	cgc	864	
Asp	Ala	Ser	Leu	Asp	Thr	Trp	Glu	Lys	Asn	Val	Ala	Val	His	Arg	Arg		
	275						280					285					
ggc	ctc	gaa	ctc	atc	aag	ccg	ggc	gag	cgc	tgc	aag	gat	atc	gcc	atc	912	

-64-

Gly Leu Glu Leu Ile Lys Pro Gly Ala Arg Cys Lys Asp Ile Ala Ile
 290 295 300
 gaa ctc aac gag atg tac cgg gag tgg gat ctg ctg aag tac cgc tcc 960
 Glu Leu Asn Glu Met Tyr Arg Glu Trp Asp Leu Leu Lys Tyr Arg Ser
 305 310 315 320
 ttc ggc tac ggc cac tcc ttc ggc gtg ctc tcc cac tac tac ggc cgc 1008
 Phe Gly Tyr Gly His Ser Phe Gly Val Leu Ser His Tyr Tyr Gly Arg
 325 330 335
 gag gcg ggc gtg gaa ctg cgc gag gac atc gat acc gtg ctg cag ccc 1056
 Glu Ala Gly Val Glu Leu Arg Glu Asp Ile Asp Thr Val Leu Gln Pro
 340 345 350
 ggc atg gtg gtc tcc atg gag ccg atg gtg atg ttg cca gaa ggc gct 1104
 Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala
 355 360 365
 ccc ggc gcg ggc ggc tac cgc gag cac gac atc ctg atc gtg ggg gaa 1152
 Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu
 370 375 380
 gac ggc gca gag aac att acc gga ttc ccc ctg ggg cct gag cac aac 1200
 Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Leu Gly Pro Glu His Asn
 385 390 395 400
 atc atc cgc aac 1212
 Ile Ile Arg Asn

<210> 22

<211> 404

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:variant CTsd7

<400> 22

Met Thr Asp Asp Met Leu His Val Met Lys Trp His Asn Gly Glu Lys
 1 5 10 15
 Glu Tyr Ser Pro Phe Ser Asp Ala Glu Met Thr Arg Arg Gln Ser Asp
 20 25 30
 Val Arg Arg Trp Met Ala Glu Asn Asp Val Asp Ala Ala Leu Phe Thr
 35 40 45
 Ser Tyr His Cys Ile Asn Tyr Tyr Ser Gly Trp Leu Tyr Cys Tyr Phe

-65-

50		55		60											
Gly 65	Arg	Lys	Tyr	Gly 70	Met	Val	Ile	Asp	Gln	Asp 75	His	Ala	Thr	Thr	Ile 80
Ser	Ala	Gly	Ile	Asp 85	Gly	Gly	Gln	Pro	Trp 90	Arg	Arg	Ser	Phe	Gly 95	Asp
Asn	Ile	Thr	Tyr 100	Thr	Asp	Trp	Arg	Arg 105	Asp	Asn	Phe	Tyr	Gln 110	Ala	Val
Arg	Gln	Leu 115	Thr	Pro	Gly	Ala	Arg 120	Arg	Ile	Gly	Ile	Glu 125	Phe	Asp	His
Val	Asp 130	Leu	Asp	Phe	Arg	Arg 135	Thr	Leu	Glu	Glu	Ala 140	Leu	Pro	Gly	Val
Glu 145	Phe	Val	Asp	Ile	Gly 150	Gln	Pro	Ser	Met	Trp 155	Met	Arg	Thr	Val	Lys 160
Ser	Leu	Glu	Glu	Gln 165	Lys	Leu	Ile	Arg	Glu 170	Gly	Ala	Arg	Ile	Cys 175	Asp
Val	Gly	Gly	Ala 180	Ala	Cys	Val	Ala	Ala 185	Val	Lys	Ala	Gly	Val 190	Pro	Glu
His	Glu	Val 195	Ala	Ile	Ala	Thr	Thr 200	Asn	Ala	Val	Val	Arg 205	Glu	Ile	Ala
Lys	Ser	Phe	Pro	Phe	Val	Glu 215	Leu	Met	Asp	Thr	Trp 220	Thr	Trp	Phe	Gln
Ser 225	Gly	Ile	Asn	Thr	Asp 230	Gly	Ala	His	Asn	Pro 235	Val	Thr	Asn	Arg	Ile 240
Val	Gln	Ser	Gly	Asp 245	Ile	Leu	Ser	Leu	Asn 250	Thr	Phe	Pro	Met	Ile 255	Phe
Gly	Tyr	Tyr	Thr 260	Ala	Leu	Glu	Arg	Thr 265	Leu	Phe	Cys	Asp	His 270	Val	Asp
Asp	Ala	Ser	Leu	Asp 275	Thr	Trp	Glu 280	Lys	Asn	Val	Ala	Val 285	His	Arg	Arg
Gly	Leu	Glu	Leu	Ile	Lys	Pro 295	Gly	Ala	Arg	Cys	Lys 300	Asp	Ile	Ala	Ile
Glu 305	Leu	Asn	Glu	Met	Tyr 310	Arg	Glu	Trp	Asp	Leu 315	Leu	Lys	Tyr	Arg	Ser 320
Phe	Gly	Tyr	Gly	His 325	Ser	Phe	Gly	Val 330	Leu	Ser	His	Tyr	Tyr 335	Gly	Arg
Glu	Ala	Gly	Val 340	Glu	Leu	Arg	Glu	Asp 345	Ile	Asp	Thr	Val	Leu	Gln	Pro 350

-66-

Gly Met Val Val Ser Met Glu Pro Met Val Met Leu Pro Glu Gly Ala
 355 360 365

Pro Gly Ala Gly Gly Tyr Arg Glu His Asp Ile Leu Ile Val Gly Glu
 370 375 380

Asp Gly Ala Glu Asn Ile Thr Gly Phe Pro Leu Gly Pro Glu His Asn
 385 390 395 400

Ile Ile Arg Asn

<210> 23
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:primer ECF21

<400> 23
 caggaaacag aattcatgac t 21

<210> 24
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:primer HIR21

<400> 24
 ccaaaacagc caagctttca g 21

<210> 25
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:primer CTextF

<400> 25
 catcggtcg tataatgtgt gg 22

<210> 26

-67-

<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer CTextR

<400> 26
gccaaaacag ccaagctttc ag 22

<210> 27
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer CTf3951F

<400> 27
cattaccgga ttccccctgg ggcctgagca caac 34

<210> 28
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer CTf3951R

<400> 28
gttgtgctca ggccccaggg ggaatccggt aatg 34

<210> 29
<211> 44
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer CT59F

<400> 29
gcatcaatta ctactctgga nnnctgtact gctatttcgg ccgc 44

<210> 30
<211> 44
<212> DNA
<213> Artificial Sequence

-68-

<220>

<223> Description of Artificial Sequence:primer CT59R

<400> 30

gcggccgaaa tagcagtaca gnnntccaga gtagtaattg atgc

44

<210> 31

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CTf59wF

<400> 31

gcatcaatta ctactctgga tggctgtact gctatttcgg ccgc

44

<210> 32

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CTf59wR

<400> 32

gcggccgaaa tagcagtaca gccatccaga gtagtaattg atgc

44

Patent Claims

1. A variant of an Erwinia-type creatinase modified relative to a wild-type creatinase having creatinase activity, characterized in that said variant comprises at least one amino acid substitutions at a position corresponding to a position of the amino acid sequence shown in SEQ ID NO: 2 selected from the group of positions consisting of N130, M203, I278, I304, and F395.
2. The variant creatinase according to claim 1, further characterized in that said variant comprises at least one substitution at an amino acid position corresponding to position N130 or I278.
3. The variant creatinase according to claim 2, further characterized in that said variant exhibits improved stability at 56° C.
4. The variant creatinase according to claim 1, further characterized in that said variant comprises at least one amino acid substitution at an amino acid position corresponding to I304 or F395.
5. The variant creatinase according to claim 4, further characterized in that said variant exhibits a lower Km-value in the enzymatic cleavage of creatine.
6. The variant creatinase according to any of claims 1 to claim 5, further characterized in that said variant comprises at least one amino acid substitution at a position corresponding to a position in the amino acid sequence shown in SEQ ID NO:2, selected from the positions N130 and I278 as well as at least one substitution selected from the positions I304 and F395.
7. The variant creatinase of claim 6, further characterized in that said variant exhibits both improved stability at 56° C and improved catalytic activity in the enzymatic cleavage of creatine.

8. A reagent for determination of creatine comprising an Erwinia-type creatinase variant according to any of the preceding claims.
9. A reagent according to claim 8, further characterized in that said reagent also comprises a sarcosine oxidase and reagents for the detection of hydrogen peroxide.
10. Method for detection of creatine, the method comprising the steps of a) incubating a sample to be analyzed with b) a reagent comprising a variant of an Erwinia-type creatinase according to any of claims 1 to 7, a sarcosine oxidase, a peroxidase, and reagents for detection of hydrogen peroxide and c) correlating the hydrogen peroxide generated to the concentration of creatine in said sample.
11. Method for detection of creatinine, the method comprising the steps of a) incubating a sample to be analyzed with b) a reagent comprising a variant of an Erwinia-type creatinase according to any of claims 1 to 7, a creatininase, a sarcosine oxidase, a peroxidase and reagents for detection of hydrogen peroxide and c) correlating the hydrogen peroxide generated to the concentration of creatinine in said sample.
12. Method for production of an Erwinia-type creatinase variant, the method comprising the steps of cloning a wild-type enzyme, introducing mutations, selecting as compared to the wild-type enzyme variants with improved thermostability or improved creatinase activity, and recombinantly producing said variant.

Application number / numéro de demande: 2404293


Figures: 2

Pages: _____

Unscannable items
received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de préparation des dossiers au
10^{ème} étage)

1/2

Figure 1: Expression vector comprising a creatinase gene

Expression vector comprising a creatinase gene

