



(86) Date de dépôt PCT/PCT Filing Date: 1993/09/24
(87) Date publication PCT/PCT Publication Date: 1994/04/14
(45) Date de délivrance/Issue Date: 2009/03/10
(85) Entrée phase nationale/National Entry: 1995/03/22
(86) N° demande PCT/PCT Application No.: AU 1993/000491
(87) N° publication PCT/PCT Publication No.: 1994/007921
(30) Priorité/Priority: 1992/09/25 (AU PL 4973)

(51) Cl.Int./Int.Cl. *C07K 16/00* (2006.01),
A61K 39/395 (2006.01), *C07K 16/08* (2006.01),
C07K 16/18 (2006.01), *C07K 16/46* (2006.01),
C12N 15/13 (2006.01), *C12P 21/08* (2006.01),
G01N 33/53 (2006.01), *G01N 33/577* (2006.01)

(72) Inventeurs/Inventors:
HUDSON, PETER J., AU;
LAH, MARIA, AU;
KORTT, ALEX A., AU;
IRVING, ROBERT A., AU;
ATWELL, JOHN L., AU;

...

(73) Propriétaire/Owner:


(54) Titre : POLYPEPTIDE LIANT CIBLE
(54) Title: TARGET BINDING POLYPEPTIDE

DELETION LINKER CONSTRUCTS

DISTANCE




54 Å




36 Å



18 Å



0 Å



(57) Abrégé/Abstract:

A target-binding polypeptide having (a) a stable core polypeptide region (SCR); and (b) at least one target-binding region (TBR), in which the target-binding region(s) are covalently attached to the SCR and which have optionally been subjected to a maturation

(72) Inventeurs(suite)/Inventors(continued): MALBY, ROBYN L., AU; POWER, BARBARA E., AU; COLMAN, PETER M., AU

(73) Propriétaires(suite)/Owners(continued): AVIPEP PTY LIMITED, AU

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(57) Abrégé(suite)/Abstract(continued):

step in order to modify the specificity, the affinity or the avidity of binding to the target. The polypeptides may self associate to form stable dimers, aggregates or arrays. The polypeptides of the invention have utility in the diagnostic, therapeutic, predictive or preventative fields of the pharmaceutical and health care industries, as well as more general application in the detection and assay of chemical entities.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 15/12, C12P 21/08 C12N 15/10, 15/11, 15/12 C12N 15/13	A1	(11) International Publication Number: WO 94/07921 (43) International Publication Date: 14 April 1994 (14.04.94)
(21) International Application Number: PCT/AU93/00491 (22) International Filing Date: 24 September 1993 (24.09.93) (30) Priority data: PL 4973 25 September 1992 (25.09.92) AU (71) Applicant (for all designated States except US): COMMON-WEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, ACT 2600 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only) : HUDSON, Peter, John [AU/AU]; 6 Cabena Street, Donvale, VIC 3111 (AU). LAH, Maria [AU/AU]; 33 Clarendon Street, Thornbury, VIC 3071 (AU). KORTT, Alex, Andrew [AU/AU]; 23 Upland Road, Strathmore, VIC 3041 (AU). IRVING, Robert, Alexander [AU/AU]; 1 Honeysuckle Avenue, Mulgrave, VIC 3170 (AU). ATWELL, John, Leslie [AU/AU]; 7 Glenwerri Court, Vermont South, VIC 3133 (AU). MALBY, Robyn, Louise [AU/AU]; Unit 7, 5 Barkly Street, Brunswick, VIC 3056 (AU). POWER, Barbara, Elaine [NZ/AU]; 6 Cabena Street, Donvale, VIC 3111 (AU). COLMAN, Peter, Malcolm [AU/AU]; 30a Hawthorn Glen, Hawthorn, VIC 3122 (AU).		(74) Agent: SANTER, Vivien, B.; Griffith Hack & Co., 509 St. Kilda Road, Melbourne, VIC 3004 (AU). (81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published 2145278 <i>With international search report.</i>
(54) Title: TARGET BINDING POLYPEPTIDE (57) Abstract A target-binding polypeptide having (a) a stable core polypeptide region (SCR); and (b) at least one target-binding region (TBR), in which the target-binding region(s) are covalently attached to the SCR and which have optionally been subjected to a maturation step in order to modify the specificity, the affinity or the avidity of binding to the target. The polypeptides may self associate to form stable dimers, aggregates or arrays. The polypeptides of the invention have utility in the diagnostic, therapeutic, predictive or preventative fields of the pharmaceutical and health care industries, as well as more general application in the detection and assay of chemical entities.		

TARGET BINDING POLYPEPTIDE

This invention relates to the construction, application and production of novel polypeptides with enhanced or modified binding activity or specificity to haptens and antigens.

The invention also relates to the construction, modification and selection of recombinant antibody-like molecules derived from expression of libraries of surface presenting antigen- or hapten-binding moieties, and to uses of these molecules.

The polypeptides of the invention have utility in the diagnostic, therapeutic, predictive or preventative fields of the pharmaceutical and health care industries, as well as more general application in the detection and assay of chemical entities.

Background of the Invention

Antibodies are protein molecules which possess a binding affinity for a target antigen or hapten. Due to the specificity of the binding interaction, antibodies are commonly used as diagnostic and therapeutic reagents. Monoclonal antibodies are derived from a pure cell line such as hybridoma cells; however, the hybridoma technology is expensive, time-consuming to maintain and limited in scope. It is not possible to produce monoclonal antibodies, much less antibodies of the appropriate affinity, to a complete range of antigens.

Antibody genes or fragments thereof can be cloned and expressed in *E. coli* in a biologically functional form. Antibodies and antibody fragments can also be produced by recombinant DNA technology using either bacterial or mammalian cells. In the Fab region of an antibody, the combination of the two heavy and light chains provides six variable surface loops at the extremity of the molecule.

These loops in the outer domain (Fv) are termed complementarity-determining-regions (CDRs), and provide the specificity of binding of the antibody to its antigenic target. This binding function is localised to the variable domains of the antibody molecule, which are located at the amino-terminal end of both the heavy and light chains. This is illustrated in Figure 1. The variable regions of some antibodies remain non-covalently associated (as $V_H V_L$ dimers, termed Fv regions) even after proteolytic cleavage from the native antibody molecule, and retain much of their antigen recognition and binding capabilities. Methods of manufacture of two-chain Fv substantially free of constant region are disclosed in US-4,642,334

Recombinant Fv fragments are prone to dissociation, and therefore some workers have chosen to covalently link the two domains to form a construct designated scFv, in which two peptides with binding domains (usually antibody heavy and light variable regions) are joined by a linker peptide connecting the C-terminus of one domain to the N-terminus of the other, so that the relative positions of the antigen binding domains are consistent with those found in the original antibody (see Figure 1).

Methods of manufacture of covalently linked Fv fragments are disclosed in US-4,946,778 and US-5,132,405. Further heterogeneity can be achieved by the production of bifunctional and multifunctional agents (Huston et al U.S. Patent No. 5,091,513, and Ladner et al U.S. Patent No. 4,816,397).

The construction of scFv libraries is disclosed for example in European Patent Application No. 239400 and U.S. Patent No. 4,946,778. However, single-chain Fv libraries are limited in size because of problems inherent in the cloning of a single DNA molecule encoding the scFv. Non-scFv libraries, such as V_H or Fab libraries, are also

known, (Ladner and Guterman WO 90/02809), and may be used with a phage system for surface expression (Ladner et al WO 88/06630 and Bonnert et al at WO 92/01047).

For use in antibody therapy, monoclonal
5 antibodies, which are usually of mouse origin, have limited use unless they are first "humanised", because they elicit an antigenic response on administration to humans. The variable domains of an antibody consist of a β -sheet
10 framework with six hypervariable regions (CDRs) which fashion the antigen-binding site. Humanisation consists of substituting mouse sequences that provide the binding affinity, particularly the CDR loop sequences, into a human variable domain structure. The murine CDR loop regions can therefore provide the binding affinities for the required
15 antigen. Recombinant antibody "humanisation" by grafting of CDRs is disclosed by Winter et al (EP-239400).

The expression of diverse recombinant human antibodies by the use of expression/combinatorial systems has been described. (Marks et al, J. Mol. Biol. 1991 222
20 581-597). Recent developments in methods for the expression of peptides and proteins on the surface of filamentous phage (McCafferty et al, Nature 1991 348 552; Clackson et al, J. Mol. Biol., 1991 352 624-28) offer the potential for the selection, improvement and development of
25 these reagents as diagnostics and therapeutics. The use of modified bacteriophage genomes for the expression, presentation and pairing of cloned heavy and light chain genes of both mouse and human origins has been described (Hoogenboom et al, Nucl. Acids. Res., 19 4133-4137; Marks
30 et al 1991 op.cit. and Bonnert et al, WPI Acc. No. 92-056862/07)

Receptor molecules, whose expression is the result of the receptor-coding gene library in the expressing organism, may also be displayed in the same way

2145278

4

(Lerner and Sorge, WO 90/14430). The cell surface expression of single chain antibody domains fused to a cell surface protein is disclosed by Ladner et al WO 88/06630.

Affinity maturation is a process whereby the binding specificity, affinity or avidity of an antibody can be modified. A number of laboratory techniques have been devised whereby amino acid sequence diversity is created by the application of various mutation strategies, either on the entire antibody fragment or on selected regions such as the CDRs. Mutation to change enzyme specific activity has also been reported. The person skilled in the art will be aware of a variety of methods for achieving random or site-directed mutagenesis, and for selecting molecules with a desired modification. Mechanisms to increase diversity and to select specific antibodies by the so called "chain shuffling" technique, ie. the reassortment of a library of one chain type e.g. heavy chain, with a fixed complementary chain, such as light chain, have also been described (Kang et al, Proc. Natl. Acad. Sci. USA, 1991 88 4363-466; Hoogenboom et al, Nucl. Acid Res., 1991 19 4133-4137; Marks et al, Bio/Technology, 1992 10 779-783).

In order to overcome the problems of human reactions to murine sequences in any part of the V-domains, framework or constant regions of the antibodies, recombinant human antibody-gene libraries may be constructed from a variety of human tissues, including peripheral blood lymphocytes (Winter and Milstein Nature, 1991 349 293). Adult humans will already have been subjected to antigenic stimulation, and therefore the capacity of the pre-immunised adult B-cell population to recognise as wide a range of antigens is diminished compared to the naive B-cell population, and is reflected in the restricted populations of antibody mRNA molecules.

- 5 -

Thus in order to access as wide a range of antigen-binding potential as possible, one of the tissues of choice is foetal peripheral blood, which being naive has a higher proportion of IgM antibody molecules than adult blood, (approximately 70% compared to the 30% for IgG), and provides the ideal source of genetic material for the construction of an antibody library destined for maturation (evolution) to a breadth where a wide range of antigens can be bound.

10 Summary of the Invention

The present invention therefore includes within its scope:

- 1) The identification and construction of novel recombinant target binding polypeptides;
- 2) Modification of such reagents to alter their performance, for example by mechanisms involving the mutation of their DNA coding regions; and
- 3) Further changing these reagents either at the genetic or the protein level, by reassortment of their subcomponents.

According to a first aspect, the invention provides a recombinant target-binding polypeptide having:

- a) a stable core polypeptide region (SCR); and
- b) at least one target-binding region (TBR).

in which the target-binding region(s) has optionally been subjected to a maturation step in order to modify the specificity, the affinity or the avidity of binding to the target.

We have been able to design and construct polypeptides according to the invention in which the specificity, affinity or avidity of binding is modified,

without the necessity for performing a maturation step. For example this has been done using immunoglobulin (Ig) and CD8.

We describe the construction of monovalent target
5 binding polypeptides in which the TBR is covalently linked to a SCR. The SCR is preferably formed by association of two covalently linked Ig-like domains of the Ig superfamily such as two antibody variable domains or CD8 domains.

We also show how polyfunctional target binding
10 polypeptides can be produced by forming separate or overlapping TBRs on a SCR. We have also shown that the Ig-like domains of members of the Ig superfamily can be constructed as SCRs and joined non-covalently to produce bifunctional or polyfunctional target-binding polypeptides.
15 We describe how to design amino acid sequences which can covalently link Ig-like domains and thereby direct self association to form stable dimers, aggregates or arrays preferably with bifunctional or polyfunctional specificity.

The target-binding region is able to bind a
20 target molecule, which may be a chemical entity of any type. For example, the target may be a small molecule such as a pesticide or a drug, a hormone such as a steroid, an amino acid, a peptide, or a polypeptide; an antigen, such as a bacterial, viral or cell surface antigen; antibodies
25 or other members of the Ig superfamily; a tumour marker, a growth factor, etc. The skilled person will readily be able to select a wide variety of targets of interest.

Where the polypeptide of the invention is to be used for *in vitro* diagnostic purposes, the core polypeptide
30 region may be any suitable protein. However, where the polypeptide of the invention is intended for use *in vivo*, the core polypeptide region should preferably be non-antigenic. Thus any normal human protein of the type which is present in serum or displayed on cell surfaces, and is

generally tolerated, would be suitable. Certain domains of normal cell-surface proteins can be produced in soluble form and, by the methods of this invention, have their affinity properties enhanced or modified. For human proteins which have as their natural target T-cell surface proteins, the soluble fragments become potential immunomodulatory therapeutic reagents especially useful for transplantation. Many of these fragments will possess homology to proteins of the immunoglobulin superfamily.

In particularly preferred embodiments of the invention, the target is selected from the group consisting of glycophorin or other red blood cell surface proteins, influenza virus neuraminidase; viral antigens such as hepatitis B antigen, and the gp40 protein of HIV; tumour markers, cell surface proteins such as CD28 and CD4; transforming growth factor α (TGF- α); and leukaemia inhibitory factor (LIF). For both diagnostic and therapeutic applications, it is particularly useful if the target binding region has more than one specificity. It is especially preferred that the target binding polypeptide possesses affinity to more than one target; this affinity is provided by separate or overlapping surfaces, thus forming a bifunctional or polyfunctional reagent. It is envisaged that bifunctional or polyfunctional reagents can also be formed by covalent or non-covalent attachment of individual target binding polypeptides, optionally using a linker polypeptide.

In a second aspect, the invention provides a DNA construct encoding the target binding polypeptide.

In a third aspect, the invention provides a method for producing a DNA construct encoding a target binding polypeptide of the invention, comprising the step of subjecting DNA encoding a target binding polypeptide to one or more cycles of mutagenesis and selection to obtain a

sub-population of DNA molecules encoding target binding polypeptides having modified characteristics of affinity, specificity, or avidity.

Preferably the DNA encoding the target binding polypeptide is present in a replication-competent element or display vector, ie. a vector which is self-replicating, optionally when present in a suitable host. The display vector is preferably selected from the group consisting of bacteriophage, filamentous bacteriophages such as Fd, viruses, bacteria, yeast, slime moulds, or mammalian cells.

Mutagenesis may be either random or site-directed, and the person skilled in the art will be aware of many suitable methods for carrying out this step. One or more target binding regions of the target binding polypeptide may be subjected to mutagenesis.

A preferred mutation system for use in the invention utilises specific mutator strains of *Escherichia coli*, designated *mutD* and *mutT1* (R. Fowler et al, J. Bacteriol., 1986 167 130). These particular mutator strains permit transfection with phage, making them especially useful for the purposes of the invention.

In a preferred embodiment, the method of producing the target binding polypeptide comprises the steps of:

- a) isolating DNA encoding the framework structure of one or more desired target-binding polypeptides by means of the polymerase chain reaction;
- b) optionally subjecting the DNA to mutagenesis in order to induce mutations in one or more target binding regions of the target binding polypeptide;
- c) inserting the DNA into one or more display vectors;

- d) selecting a sub-population of display vectors displaying target binding polypeptides of desired specificity, avidity or affinity;
- 5 e) subjecting the selected sub-population to one or more cycles of mutagenesis and selection in order to obtain a sub-population of display vectors displaying target binding polypeptides having modified characteristics of affinity, specificity or
- 10 avidity; and
- f) inserting DNA encoding the modified target binding polypeptides into a high level expression vector.

15 Selection of the sub-population of display vectors may be achieved by a variety of conventional methods such as target binding, fluorescence-activated cell sorting, or exploitation of the biotin-avidin or biotin-streptavidin systems. A particularly preferred method is

20 affinity selection on an insoluble support such as immunotubes; this has been found to be especially convenient.

It will be appreciated that the invention therefore also provides a method of producing the target

25 binding polypeptide, by transferring the high level expression vector described above into an appropriate expression host, expressing the target binding polypeptide, and isolating the protein thus produced.

It will also be clearly understood that the

30 target binding regions and the stable core polypeptide may be different regions of the same molecule, or may be derived from different molecules.

Types of target binding polypeptide constructs which are contemplated by the invention include modified

2145278

10

antibodies or antibody fragments, scFv fragments comprising an association link to permit continuous reassortment, modified CD8 molecules, for example single chain CD8, and combinations of antibody molecules or fragments thereof with CD8 or other molecules related to the immunoglobulin superfamily such as the individual domains of the MHC Class I and II molecules. For example, the $\alpha 3$ domain of MHC Class I binds to CD8, and therefore soluble versions of $\alpha 3$ become potential immunomodulatory reagents. Preferred constructs utilizing CD8 include:

- a) Single-chain molecules in which the V-like domains only of the α and β subunits are linked, and
- b) Molecules in which the N-terminal amino acid has been altered from lysine to serine, in order to alter the charge balance of the signal peptides, thus enabling bacterial expression without adversely affecting biological activity.

Target binding polypeptides may include covalently attached polypeptide tails which can be TBRs or which may permit non-covalent association to other TBPs.

Although the following description refers in some examples specifically to IgG type antibodies and their fragments, it will be clearly understood that the invention is also applicable to other types of antibody molecule, such as IgM and IgA.

The DNA sequence encoding the target binding polypeptide may be cloned into any vector which will allow display of the polypeptide on bacteriophage or cell surfaces. Preferred vectors include pHFA, whose construction is described in International Patent Application No. PCT/AU93/00228, and its structure is illustrated in Figure 4. Preferred bacterial hosts for protein expression are *E. coli* and *Bacillus subtilis*.

Detailed Description of the Drawings

The invention will be described in detail by reference to the following non-limiting examples and to the drawings, in which:

5 Figure 1 illustrates the structure of antibodies and their fragments:

10 a) This shows the polypeptide chain structure of a typical IgG antibody molecule, which is composed of two identical heavy and two identical light chains, each divided into variable (V) and constant (C) domains. The whole IgG molecule has two identical antigen binding surfaces termed Fv regions, which are formed by the pairing of V_H and V_L chains. The combination of V_H and V_L provides 6 loops, termed complementarity-determining regions (CDRs),
15 at the extremity of the molecule and these provide the antigen binding surface and thereby the binding specificity of the antibody to its target antigen.

 b) an Fab antibody fragment comprises one light and a portion of one heavy chain.

20 c) A single-chain scFv is shown as V_H and V_L domains joined by a peptide linker between the C-terminus of V_H to the N-terminus of V_L . Both Fab and Fv fragments are expected to have the same antigen binding surface as the parent antibody.

25 Figure 2 shows antibody fragments such as Fab and scFv molecules displayed on the surface of filamentous Fd bacteriophage by covalent fusion to either the minor coat protein at the tip of the phage, the gene III protein or as fusions with the major, gene VIII coat protein. For
30 display of Fab molecules, only one of the chains (Heavy or Light) is anchored to the phage coat protein, and the other chain is provided in soluble form in the host cell periplasm. The Fd bacteriophage are still viable, although fusions on the gene III protein reduce infectivity into

host cells.

Figure 3 shows how pools (libraries) of heavy and light variable chains can be constructed into a Fd phage display vector with one of the chains fused to either the gene III protein or gene VIII protein of the phage. The display vector is transfected into host cells to generate a dual-combinatorial library. Each host cell produces viable Fd phage in which the antibody fragment is displayed on the phage surface and the gene encoding the antibody is packaged with the viral genome. Affinity purification of the phage is based on affinity to a target antigen, and allows simultaneous recovery of the gene encoding the antibody from the viable phage. Phagemid display vectors can improve transformation yields, but require helper phage to assemble viable progeny. Alternative strategies include the construction of hierarchical libraries in which one chain is held constant and displayed with a library of the second chain to select the highest affinity paired chains. More complex libraries can be constructed using gene pools on separate display vectors and then cross-transfecting host cells. Gene recovery will depend on the relative packing efficiency of the two vectors.

Figure 4 shows the structure of the phagemid vector pHFA. This vector has the ability in suppressor strains of *E. coli* to express cloned antibodies as fusions with the gene III protein on the surface of the Fd phage, whereas in non-suppressor strains the cloned antibody genes are expressed as soluble products. The lacZ promoter allows induction of expression with IPTG, and the FLAG tail, which is expressed as a fusion with the antibody, is used for detection of synthesis and affinity purification.

Figure 5 shows the series of scFv NC10 deletion linker constructs and the theoretical minimum distance (in

Angstroms) spanned by the polypeptide linker.

Figure 6 shows the DNA sequence of synthetic oligonucleotide duplexes encoding peptide linkers of different lengths that were inserted into appropriately restricted pPOW-scFv NC10.

Figure 7 shows the analysis of synthesised scFv NC10 proteins from the $V_H.15.V_L$, $V_H.10.V_L$, $V_H.5.V_L$ and $V_H.V_L$ from uninduced (lanes 1-4) and induced (lanes 5-8) respectively on a Coomassie gel (upper panel) and a Western blot (lanes 9-16 lower panel) of the Coomassie gel probed with the anti-FLAG, M2 antibody (IBI, New Haven, CT) followed by goat anti-mouse horse radish peroxidase conjugate (Sigma) as the second antibody and detected by enhanced chemiluminescence (Amersham).

Figure 8 shows a summary of observed scFv NC10 associations and activity to its target antigen.

Figure 9 shows the similarity in structure between an antibody Fv fragment and a CD8 α chain heterodimer.

a) This is a ribbon drawing of a V_H or V_L molecule showing the CDR loops numbered 1-6 and the structurally conserved framework regions as ribbons.

b) This is a ribbon drawing of two CD8 α chains. The regions corresponding to antibody CDR loops are shown at the top of the molecule and numbered.

The homodimer is oriented with the molecular dyad axis situated vertically in the plane of the page. CVR-like loops from the top surface of the molecule as shown, and the CDR 1-like, CDR 2-like and CDR 3-like loops are labelled 1, 2 and 3 respectively for one sub-unit, and 1', 2' and 3' for the other sub-unit. The C-termini extend from the bottom of the molecule. The loops forming the dimer interface are the CDR 3-like loops (top) and C-C' loops (bottom).

Figure 10 shows a Coomassie stained SDS-PAGE gel of the synthesised scCD8 in pPOW using *E. coli* host cells pop2136 showing whole cell lysates. Arrows show the

positions of the fused and mature (cleaved signal sequence) scCD8. In this figure:

	Lane 1	uninduced cells containing pPOW-scCD8
5	Lane 2	4 hours post-induction of pPOW-scCD8
	Lane 3	uninduced cells containing pPOW-Lys→Ser scCD8
	Lane 4	4 hours post-induction of pPOW-Lys→Ser scCD8
10	Lane 5	pre-stained molecular weight markers.

Figure 11 illustrates a scheme whereby individual antibody genes can be affinity matured. Individual genes can be selected from phage display libraries, and then subjected to rounds of *in vivo* or *in vitro* mutations. The affinity matured antibody fragments are then selected for their ability to bind antigen, prior to further rounds of mutation or high level gene expression. Entire antibody libraries can be increased in complexity by cyclic rounds of mutation prior to selection of individual phage via panning or affinity selection.

Figure 12 shows results of ELISA screening of colonies subjected to mutation for affinity maturation.

Figure 13 shows an example of the p569 vector (a gift from W. Nellen), a shuttle vector for expression in *D. discoideum*. The vector has the alpha L fucosidase promoter and signal sequence, a multi-cloning site, a transcription terminator and the transposon Tn903 for selection by G418.

Figure 14 shows the tertiary structure depicted as the polypeptide backbone of NC10 scFv fragments complexed with two influenza neuraminidase subunits solved at 3 Angstroms resolution by X-ray diffraction analysis. The linker polypeptide joining the heavy and light chain variable regions is not depicted in this figure. In the crystal structure two Fv fragments are associated back-to-back to dimerise two different neuraminidase subunits. In the context of the scFv fragments the dimeric Fv module can be considered a bifunctional reagent.

Figure 15 shows a model of two neuraminidase tetramers which are bound together by four NC10 scFv dimers in solution as resolved by electron microscopy.

5 Figure 16 shows a model of bifunctional Fv molecules dimerised back-to-back and are closely associated without steric interference. In this model, the C-terminus of the heavy chain can be directly linked to the -13 residue of the light chain variable region with minimum reorganisation of the remaining polypeptide backbone.

10 Figure 17 shows a schematic representation of non-covalently and covalently joined scFv dimers respectively.

Sequence I.D. 1 shows an example of a mouse Ly-2⁺Ly-3 V domains construct designed for bacterial expression.

Sequence I.D. 2 shows an example of a human single-chain CD8 construct designed for expression in a bacterial secretion vector such as pPOW.

20 Sequence I.D. 3 shows an example of a mouse MHC α 3 domain designed for expression.

Sequence I.D. 4 shows the DNA sequence of the linkerless 1C3 scFv in pHFA.

25 Sequence I.D. 5 shows the DNA sequence of the anti influenza NC10 scFv with the pel B secretion signal and the FLAG C-terminal peptide.

Sequence I.D. 6 shows the DNA sequence of the first 1443 bases of the anti-glycophorin 1C3Fab fragment in pHFA ready for ligation post PCR amplification for ligation into p569.

30 Preferred embodiments of the invention include the following:

1. The structure of the target binding polypeptides may be based on scFv molecules in which one TBR is formed by six surface polypeptide loops to provide contact region to antigen, and hence specificity. In a particularly preferred embodiment, the TBR may be formed by four CDR

2145278

16

loops for contact with antigen to provide sufficient contact area and affinity (Figure 14). Our results using NC10 indicate that it is feasible to randomly mutate these polypeptide sequences to modify target affinity.

5 2. Bifunctional or polyfunctional reagents can be produced by covalent linkage of individual target binding polypeptides. The covalent linkage can be provided by polypeptide chains in the manner of single-chain Fv molecules. Specificities can be linked by joining together
10 individual proteins, regions which have a propensity for dimerisation or aggregation. Thus it is not necessary for example to link two single-chain Fv fragments in their normal orientation by an additional polypeptide chain, but this can be achieved by linking Heavy and Light chains of
15 differing specificity, or Heavy to Heavy chains, Light to Light chains which can then associate to form functional dimers or aggregates. Of course the method can also be used to join V domains with required specificity to other protein domains, including the Immunoglobulin-like domains
20 derived from CD8, T-cell receptor fragments, or MHC fragments.

In a preferred embodiment the covalent joining of two Ig-like domains, such as the heavy and light antibody variable regions, can be produced with or without a linker
25 polypeptide. For monomeric scFv fragments the polypeptide linker covalently joins the V_H and V_L domains between the carboxyl terminal of one variable (V) domain to the amino terminal of the other V domain without compromising the fidelity of the scFv binding site. The scFv may be
30 assembled in either domain order as a V_H -linker- V_L fusion protein or as a V_L -linker- V_H fusion protein. The linker should preferably be hydrophilic in nature to prevent it from associating with hydrophobic V domain surfaces. The lengths of the linker may be less than 25 amino acid

residues with a preferred size established by empirical selection. A preferred linker sequence consists of pentameric units of Gly4Ser in which the serine residues enhance the hydrophilic characteristics of the peptide backbone, while the glycyl residues give the linker enough flexibility to adopt a range of conformations around the V domains. In a special preferred embodiment the covalent association of polymeric Fv fragments can be produced without an additional linker polypeptide by removal of a segment of one Ig-domain at the junction sequence. These linker-minus constructs are referred to herein as tightly coupled domains (TCDs). The number of amino acids to be removed can be determined either empirically or with the aid of protein design considerations. Figure 16 depicts the association of two Fv molecules "back-to-back" as TCDs and in which the two TBRs are at opposite ends of the molecule thus forming a bifunctional reagent capable of cross-linking two target molecules. In this example, preferably up to 13 amino acids are removed for close association. The resultant molecule has a propensity for oligomerisation, at least to dimers, with a close but not sterically inhibited interaction between the Ig-like domains. It will be appreciated that polypeptide tails can be added at the free amino and carboxyl termini to increase the number of TBRs on the molecule. It will also be obvious that the missing polypeptide sequences that had been removed at the junction of Ig-like domains can be replaced, in whole or in part, by providing the polypeptide sequences attached to another position in the Ig-like domains. We anticipate that these molecules will be capable of forming two-dimensional arrays thereby providing a bifunctional surface. It will be appreciated that these arrays will have special application as biological coating devices.

3. The complete three-dimensional structure of mature human or mouse CD8, comprising heterodimeric α and β chains, is not yet known. Predictions from a crystal structure of homodimeric human CD8 α suggests that the α chains are similar in topology to antibody V_L domains (Leahy et al, Cell, 1992 68 1145-1162). We have constructed single-chain variants of mouse CD8 α/β heterodimer for expression using bacterial secretion vectors, and similar results would be expected using human CD8. Native human or mouse CD8 molecules are presumed to have affinity only for MHC Class I molecules. We predict that random library approaches, such as those described in the Examples herein, will enable scCD8 molecules to be used as a stable framework for the production of target binding polypeptides. By this process, scCD8 molecules can be used as antibody mimics. Furthermore, the scCD8 molecules can be further modified in the size and conformation of CDR-equivalent loop structure to provide a framework for less than six CDR loops in the contact surface. In a particularly preferred embodiment, we envisage a stable protein framework capable of providing four or five CDR loops in the contact region. We also envisage the strategy to apply to other Ig domains. For example the immunoglobulin-like domains of MHC Class I and II can be expressed in soluble form and when modified can be used as immunomodulatory reagents.

4. Modifications to target binding polypeptides such as those described above can be based on mutation of the coding region, by the use of library selection and modification strategies such as those shown in Figure 3, to mature a single TBP or TBP library of low affinity and wide specificity to enhance the range of target molecules which it recognises, but more importantly to produce a range of

binding affinities for each member of the library, the individual DNA coding regions of which may be easily selected and isolated by modifications of known methodologies. It is envisaged that such a library will
5 comprise antibody-like fragments, or any other peptide which shows an affinity for a ligand or another protein, enzyme or receptor. This may also include a stable core polypeptide which is not in itself antigenic, but may be modified by the addition of CDR loops or peptides with an
10 affinity for specified ligands by grafting the coding regions by recombinant DNA techniques. It can also be seen that a change or changes to the framework regions may result in a change of conformation of the protein such that an altered binding surface is presented, with binding
15 properties different from those of the parent molecule.

The most trivial example includes the construction of expression libraries that produce recombinant antibody fragments (including single-chain Fv fragments) with predetermined target binding specificity.
20 *In vitro* mutation and affinity maturation provide means of presenting the binding molecule such that the appropriate coding regions are selected and retained. Presentation vectors which will allow continual reassortment of the binding domains (which in this example as a preferred
25 embodiment will encompass V_H and V_L domains) subsequent to each of the selection steps shown in Figure 3 can suitably be used, for example pHFA.

5. The invention may be used for the construction and selection of a wide range of receptors, receptor-like
30 molecules and molecules constructed with mutations in potentially critical regions for both binding, structural integrity and biological activity. Initially phage surface presentation after expression and phage rescue from *E. coli* is used to monitor the efficacy of this approach, but other

systems such as the eukaryotic systems are also expression competent. Yeast (*Saccharomyces cerevisiae*) has been shown to express the V_H of NC41, a monoclonal antibody directed against influenza virus neuraminidase, under the control of the alpha mating factor promoter, and the slime mould *Dictyostelium discoideum* is able to express recombinant proteins including both V_H and the scFv of NC10.

6. The specific selection of target-binding polypeptides able to bind to the specified antigens (which may include LIF, TGF- α , glycophorin, cell surface markers or other cell specific surface proteins), is made possible as a result of the presentation on the display vector, for example the presentation on the phage surface of these peptides fused to the Gene III product. Having selected the appropriate phages, they are then subjected to rounds of mutation, as shown in Figure 5.

In the following examples, the *mutD* and *mutT1* mutator strains of *E. coli* are used to induce mutations at random throughout the molecule. This is done by transformation of these *E. coli* strains with the plasmid DNA by any of the standard techniques that appear in the literature; the preferred method is by electroporation. Alternatively the recombinant phage may be transfected into the mutator strains by standard transfection methods. After rounds of growth of these plasmid/phage-bearing *E. coli*, the phage may be rescued by standard techniques with a helper phage, and can then be used in antigen-binding assays to determine the effects of various mutations on the binding affinity.

These mutations are not confined to base substitutions in the DNA, but may also encompass the addition of peptides to the structure of the molecule such that the number, size and location of the binding regions in the molecule is altered. A single domain binding unit

with these additions will show binding characteristics of substantially altered affinity if not specificity. The correlation between mutation at specified sites and the binding affinity may then be used to design novel CDR loops and framework regions for target binding polypeptides with therapeutic and diagnostic potential.

Also included in the scope of the invention is the expression of recombinant proteins from recombinant cells under the direct control of the antigen, or some other ligand which is responsible for the first step in the process towards controlled expression of the "antibody genes".

7. Bifunctional or polyfunctional reagents can be selected using the library technology described above. Target binding polypeptides may be displayed for affinity selection by attachment through a polypeptide tail. Selection based on affinity to two or more different target antigens or haptens will select a single molecule which has two binding surfaces at different positions of the same molecule. The binding surfaces can be overlapping. To construct a library for selection of bifunctional or polyfunctional reagents, the strategy of site specific and random mutagenesis applied to two or more surfaces of the protein molecules may be used. In the case of single-chain Fv or CD8, the preferred regions for mutation will be the CDR loops and their opposite counterpart loops at the other end of the scFv molecule. In the case of Fab molecules, the preferred mutations will be at CDR loops and the opposite counterpart loops at the other end of the constant domains.

Unless otherwise specifically stated, all standard methods referred to herein are to be found in "Molecular Cloning-A Laboratory Manual" Sambrook et al 1990.

2145278

22

Example 1Construction and Expression of Target
Binding Polypeptides as Single Chain
Fv Fragments Using Polypeptide
Linkers of Different Lengths

5 A parent scFv fragment of NC10 (a monoclonal
antibody that recognises the neuraminidase (NA) molecule on
the N9 strain of influenza virus) was designed, constructed
and expressed in *E. coli* (Sequence I.D. 5). The amino
terminal secretion signal PelB directed the synthesised
10 protein into the *E. coli* periplasm where it became
associated with the insoluble membrane fraction. An
octapeptide (FLAG; IBI USA) tail was fused to the carboxyl-
terminal of the scFv and was used to monitor the scFv
through subsequent purification procedures. This reagent
15 is bifunctional with specificity to both neuraminidase and
anti-FLAG antibodies.

 The scFv NC10 protein was purified by
solubilization of the *E. coli* membrane fraction with
guanidinium hydrochloride followed by column
20 chromatography. Size exclusion HPLC of purified scFv NC10
showed that the scFv fragment emerged in two peaks
corresponding in size to monomers (27 kDa) and dimers (54
kDa). Furthermore, the monomeric form bound to N9 NA to
form a complex of ~320 kDa while the dimeric form bound to
25 N9 NA to form a complex of ~640 kDa. The 320 kDa complex
could consist of four scFv molecules binding to a single NA
molecule, while the 640 kDa complex could consist of four
scFv molecules binding to two NA molecules. Electron
microscopy confirmed the tight coupling of two
30 neuraminidase tetramers by four bifunctional scFv dimers
(Figure 15). High resolution electron microscopy was
performed on the tern N9 (avian) strain of influenza
neuraminidase complexed with scFv constructs of the NC10 of
the Mab, where the molecular complexes were stained

- 23 -

(contrasted) either with potassium phospho-tungstate at pH 7.0 or with uranyl acetate at pH 4.0.

Based on our previous extensive experience of imaging molecular complexes of the same N9NA with
5 monoclonal Fabs (32/3, NC35 and NC41) and with whole monoclonal IgGs (32/3, NC41 and NC10), we were able to interpret the N9Na-scFv complex images as closed structures of pairs of neuraminidase heads coupled together face-to-face by four bridging scFv dimers in such a manner as to
10 maintain four-fold point-group symmetry of this densely packed molecular complex (Figure 15). This image interpretation of the N9Na-scFv molecular complex is directly compatible with the observed molecular weight of the complex in solution of
15 $M_r \sim 610,000$.

X-ray diffraction analysis of crystals in which the scFv is complexed with neuraminidase (Figure 14) demonstrates a close association between two scFv molecules related by a two-fold axis of rotation. Two possible
20 dimeric conformations are possible. In the first instance the V_H and V_L domains encoded by a single polypeptide chain with additional peptide tails form a bifunctional scFv which associates non-covalently with the separate scFv molecule (Figure 17). In the second instance, the V_H and
25 V_L domains forming the antigen binding surface (the TBR) in each Fv are non-covalently associated and the two Fvs are covalently joined by the linker polypeptide (Figure 17).

Molecular modelling studies (Figure 14) indicate the distance between the V_H and V_L domains of a non-
30 covalently associated dimer would be at least 35Å, whereas the distance between the V_H and V_L domains of a covalent dimer would be less than 25Å. Given the 3.8Å (0.38nm) distance between adjacent peptide bonds and the distance

lengths that the linkers can theoretically span (54Å, 36Å, 18Å and 0Å for the 15, 10, 5 and 0 residue linker pPOW-scFv NC10 constructs respectively) we examined the type of scFv-NA complexes formed when scFv proteins with different linker lengths bind to antigen.

A series of scFv NC10 proteins with shortened linker lengths were constructed (Figure 5). The first pPOW-scFv NC10 construct has a polypeptide linker consisting of three pentameric Gly4Ser units (this pPOW-scFv NC10 construct was referred to as the 15 residue linker, $V_H.15.V_L$). The deletion linker mutants were constructed by sequentially removing each of these pentameric units to form constructs with two, one and zero units (referred to as the $V_H.10.V_L$, $V_H.5.V_L$ and $V_H.V_L$ residue linker pPOW-scFv NC10 constructs respectively). Furthermore, a scFv NC10 construct was made by deleting the first β -strand of the V_L domain (the first 13 amino acids) so that the carboxyl-terminal of the V_H domain joined directly to the V_L domain ($V_H.-13.V_L$).

20

Detailed Construction of pPOW-scFv NC10 with Shortened Linker Lengths

The pPOW-scFv NC10 construct was digested successively with *Bst*E II (New England Biolabs) and *Sac* I (Pharmacia) according to manufacturers' specifications and the polypeptide linker released. The restricted linkerless pPOW-scFv NC10 DNA was electroeluted from an 0.8% agarose gel and the DNA concentrated by precipitation with 0.3M Na acetate and 2.5 volumes of ethanol. Synthetic oligonucleotides were phosphorylated at their 5' termini by incubating at 37°C for 30 min with 0.5 units of T4 polynucleotide kinase (Pharmacia) and 1mM ATP in One-Phor-All Buffer PLUSTM (Pharmacia). Pairs of complementary

30

phosphorylated oligonucleotide primers (Figure 6) were premixed in equimolar ratios to form DNA duplexes encoding single chain linkers of altered lengths. These duplexes were ligated into the *BstE* II-*Sac* I restricted pPOW-scFv NC10 plasmid using an Amersham ligation kit. A slightly different approach was required to make the V_H -13. V_L construct. An oligonucleotide primer (Figure 6) spanning the deleted V_L domain was constructed and used in conjunction with a FLAG specific oligonucleotide (Figure 6) to amplify by PCR a V_H -13. V_L fragment of the scFv NC10. The amplification product was digested with *BstE* II and *EcoR* I and ligated into similarly digested pPOW-scFv NC10 plasmid using an Amersham ligation kit. The ligation mixtures were purified by extraction with an equal volume of phenol/chloroform and precipitated with 0.3 M Na acetate and 2.5 volumes of ethanol. The ligated DNA was resuspended in 20 ml H₂O and 5 ml of the sample was transformed into *E. coli* DH5a (*supE44*, *hsdR17*, *recA1*, *endA1*, *gyrA96*, *thi-1*, *relA1*) and LE392 (*supE44*, *supF58*, *hsdR14*, *lacY1*, *galK2*, *galT22*, *metB1*, *trpR55*). Cells were shaken in 1 ml of LB medium for 1 hr and plated onto 2xYT medium with 100 mg/ml ampicillin. Recombinant clones were identified by PCR screening with oligonucleotides directed to the PelB leader and FLAG sequences of the pPOW vector. The DNA sequence of the shortened linker regions were verified by sequencing double-stranded DNA using Sequenase 2.0 (United States Biochemical).

Protein Expression of the scFv NC10 Proteins with Shortened Linkers

Transformed LE392 were grown overnight at 30°C in SB medium and diluted 1:10 to inoculate fresh SB medium. Cultures were grown at 30°C with shaking until the absorbance at 600nm (A_{600}) was approximately four. The

2145278

26

temperature was raised to 42°C for the remainder of the induction period (which continued for 4 hr until the A_{600} ~7). Cells were recovered by centrifugation (Beckman JA10 6,000 rpm for 15 min) and the supernatant fraction removed.

5 The cell pellet was resuspended in 10% of the original volume in 20% sucrose, 10 mM Tris.HCl. pH7.5 and left on ice for 5 min. EDTA was added to a final volume of 5 mM and the mixture incubated on ice for a further 10 min and centrifuged as before to pellet the cells. The supernatant

10 was discarded and the cell pellet resuspended in H_2O , the mixture was recentrifuged and the supernatant containing the periplasmic proteins removed. The resulting cell pellet was resuspended in H_2O and lysed by sonication (six 30 sec bursts for large scale preparations and one 30 sec

15 burst for small scale preparations) and kept on ice for 5 min. After centrifugation the aqueous phase was recovered as the solubilized cytoplasmic fraction while the pellet contained the insoluble membrane-associated fraction. To verify scFv NC10 expression total cell lysate from

20 individual clones were analysed by SDS-PAGE under reducing conditions and Western blotting using the anti-FLAG monoclonal antibody, M2 (Figure 7). Single positive bands migrating at ~28, 29, 31 and 32 kDa were observed (Figure 7, lanes 13-16) which correlate with the anticipated Mr of

25 the scFv NC10-FLAG fusion protein synthesised by pPOW-scFv NC10 constructs with 0, 5, 10 and 15 residue linkers respectively. ScFv NC10 proteins with 0, 5, and 10 residue linkers showed the same characteristics as the 15 residue linker. The scFv NC10-FLAG fusion proteins were associated

30 with the insoluble membrane fraction of *E. coli*, approximately half of which could be solubilized by treating with guanidinium hydrochloride.

The soluble products were purified by gel filtration and chromatography on Mono-Q or on an affinity

matrix containing an antibody specific for the tail moiety. The pure products were characterized by SDS-PAGE, size exclusion HPLC (SE-HPLC), ultracentrifugal analysis, binding activity towards the parent antigen (influenza virus neuraminidase), electron microscopy of the complexes formed between the antigen and reagent. Cross linking experiments confirmed the size of the products. The properties are summarized in Figure 8.

Example 2

Construction of mouse and human scCD8

The α and β chains containing only the V-like domains of mouse CD8 were amplified separately by PCR with VentTM polymerase using primers containing homology to the V-like domains (using available database sequences) and with additional nucleotides encoding the (Gly₄Ser)₃ linker (Sequence I.D. 1). After annealing the two separate domains the products were extended using dNTPs and polymerase. The scCD8 gene was amplified using new primers containing MscI site at the 3' end and SalI site at the 5' end. The single chain product was digested with MscI and SalI then cloned into MscI and SalI digested pPOW vector. High level protein synthesis was obtained in *E. coli* host cell strain pop2136. An N-terminal modification was designed to increase the synthesis of correctly cleaved product which was achieved by changing the N-terminal residue of mouse CD8 α chain from Lysine to Serine (the Human CD8 α chain N-terminal residue is a Serine). The synthesis of scCD8 in pPOW using *E. coli* host cell strain pop2136, showing whole cell lysates, can be seen in Figure 10. The synthesised scCD8 product was detected by anti-CD8 antibodies that only recognise protein in the conformationally correct form.

Human scCD8

The DNA encoding the V-like domain of the mature α chain protein was amplified by PCR using TaqTM polymerase and primers containing homology to the V-like domain (using available database sequences) with additional nucleotides encoding the (Gly,Ser)₃ linker and incorporating restriction enzyme sites *MscI* and *BamHI* (Sequence I.D. 2). The V-like domain of the CD8 β chain was amplified by PCR directly from DNA isolated from blood using primers containing *BamHI* and *EcoRI* restriction enzyme sites. The two individual products were digested with the appropriate enzymes then ligated into *MscI* and *EcoRI* digested pPOW vector.

The DNA sequence of each of the single chain CD8 constructs was confirmed by double stranded DNA sequencing. The nucleotide sequence can be seen in Sequence I.D. 1 and 2. In this example the vector directs the synthesis of a scCD8 with a C-terminal peptide tail for diagnostic and coupling applications, including affinity purification.

Preferred techniques to monitor the biological activity of the scCD8 product include:

- a) Direct measurement of protein binding affinity for example using biosensor technology or ultracentrifugation using binding to whole cells, cell surface molecules or their fragments such as β 2 microglobulin or the α 3 domains of the MHC class I molecule.
- b) measurement of binding to the MHC class I molecules expressed in RMA-S cells (peptide loaded) using the C-terminal peptide tails as diagnostic markers.
- c) an interference of function assay such as monitoring changes to the peptide induced

dose-dependent effect on IL2 production during T-cell activation.

Example 3

Construction of Linkerless
AntiGlycophorin 1C3

5

The parent 1C3 antibody and scFv derivatives are disclosed in the International Patent Application No. PCT/AU93/00228.

Oligonucleotide N2034

10

(5'-ACGTAGGTCACCGTCGCCTCCGACATCGTCATGTCACAGTCTCCATCCTCC-3') was synthesised to have complementarity to the last 15 bases at the 3' end of 1C3 V_H coupled directly to the first 30 bases of the 1C3 V_L 5' sequence without any intervening linker sequence.

15

Oligonucleotide N2035

(5'-TTTATAATCTGCGGCCGCCCCGATTAATTTC-3') was synthesised to have complementarity to the 1C3 V_L sequence on the opposite strand around the Not I site near the 3' end.

20

The two oligonucleotides were used with 1C3 template DNA in a Polymerase Chain Reaction to produce a 1C3 product of 3'-V_H sequence juxtaposed to 5'-V_L sequence flanked by Bst EII and Not I restriction endonuclease sites.

25

After incubation of the PCR product with restriction endonucleases Bst EII and Not I, the resultant fragment was ligated with vector pHFA containing the 1C3 scFv sequence previously digested with Bst EII and Not I to remove the intervening sequence. The ligated product was used to transform *E. coli* strain TG1. Transformant colonies containing inserts were verified as containing the DNA sequence as shown in Sequence I.D. 4.

30

This gene construct was expressed in this vector and related vectors when transferred to a non-suppressor

2145278

30

E. coli strain and induced with IPTG, or by transferring the 1C3 coding region to the thermoinducible expression vector pPOW.

5 Example 4 Mutation with mutator strains of *E. coli*

10 The NC10 scFv plasmid coding for the expression of the recombinant antineuraminidase antibody NC10 scFv was electroporated into *E. coli* mutD. Mutants were produced by
15 subjecting the samples to the mutation cycle shown in Figure 11. They were grown for 50 generations in exponential phase (to induce mutation of the phasmid DNA) in YT+AMP+TET and then rescued with the helper phage. The rescued phage was applied to the immunotubes previously
20 coated with 10µg/ml of the antigen, non-binding phage removed by washing with PBS etc and the specifically bound phage eluted with 100mM triethylamine, collected into 0.5 volumes of 1M Tris-Hydroxymethylmethyllamine-HCl pH7.5 and then transfected into mutD cells by standard methods,
25 (unless otherwise specifically stated all standard methods referred to herein are to be found in "Molecular Cloning-A Laboratory Manual" Sambrook et al, 1990) and again grown through 50 generations whilst maintaining the cells in the logarithmic phase of growth. After an appropriate number
30 of rounds of mutation selection which in this example is three the phage titres are in the region of 10^7 - 10^8 phage/ml. After the final panning step, eluted phage were transfected into *E. coli* TG1 cells and plated onto YT+AMP+Glucose plates and then each of the isolated
35 colonies grown before phage rescue and analysis by ELISA on "flu" virus or glycophorin. The colonies which exhibited non-wild-type levels of ELISA activity, were then amplified, the DNA sequenced and the phage transfected into *E. coli* HB2151 cells available from the American Type Culture Collection, for soluble expression. The phage were

- 31 -

transfected into HB2151 by the standard methods and the selected individual colonies of each phage sample grown in YT+AMP(100µg/ml) prior to induction with 1mM IPTG (isopropylthiogalactoside) for 4 to 16h at 37°C, with or
 5 without subsequent incubation at 4°C for 16h. The culture supernatant and the extracts of periplasm, cell membranes and cell cytoplasm were collected and analysed for the recombinant gene expression as described (Power et al, Gene 1992 113 95-99).

10 Example 5

The recombinant 1C3 scFv (a glycophorin-binding antibody coding region) in the phagemid pHFA prepared as described in International Patent Application No. PCT/AU93/00228 was subjected to random mutation in the *mutD*
 15 *E. coli* as discussed in Example 4, and the selection protocol similarly applied, with the exceptions that the selection involved coating the solid phase matrix (ELISA plate, Immuntube,TM or latex bead) with glycophorin A from a 10µg/ml solution in PBS. The results of the ELISA
 20 screening for selection of individual colonies is illustrated in Figure 12. Competitive ELISA assays, using detection with anti-FLAG antibody, were performed on selected colonies after mutation, and Table 1 shows the increases in relative affinity of the expressed proteins
 25 for the antigen asialoglycophorin.

Table 1

	recombinant cloned scFv	Mutation	Affinity nM (off rate)
	1C3	wt	62
30	1C3.A13		40
	1C3.B7		29

Example 6

A scFv library in the phagemid vector pHEN (Medical Research Council, U.K.) was transferred into the *mutD* strains of *E. coli* and treated as for Examples 4 or 5 for the mutation, detection and selection of scFv with modified binding ability. Selecting for glycophorin binders. To increase the range of glycophorin-binding antibodies available the naive scFv library was used as the starting point for this maturation and affinity selection of phage displayed scFvs. Two of the unique anti-glycophorin scFvs that were selected from the naive scFv phage display library, have the deduced amino acid sequence shown for the region of their Vk4 chains that were subsequently shown to be mutated are shown in Table 2.

15

Table 2 (SEQ ID NOS: 3 AND 4)

FTASTGDVPDRFSGSGSGTDFTLRISLQAEDVAVYYCQQASVFP (SEQ ID NO:3)

CIYWNPDSPDRFSGSGSGTDFTLRISLQAEDVAVYYCQQASVFP (SEQ ID NO:4)

20

Affinity maturation of each of these molecules was achieved by using the mutation (*mut D5*) affinity selection cycle, as we show in Figure 11, and the changes that result to a subset of the mutated molecules is shown in Table 3.

25

Table 3

	scFv selected from Naive Library	Mutation A.A (position)	Affinity μ M (off rate)
5	A9	wt	48
	A9.5	S-V (98)	15
	A9.13	G-D (63)	24
		V-S (64)	0.06
10	E3	wt	18
	E3.1	S-Q (82)	9
	E3.2	S-T (83)	2
	C12	wt	11
	C12.1	G-Q (48)	35
	C12.2	L-G (15)	1

15 Example 7

Expression of the antibody-fragment coding regions in *D. discoideum* is from the vector pAV1 which has been constructed from the parent vector p569 (a gift from W. Nellen, Max-Planck Institute, Munich, Germany) and the V_H coding region of the NC41 V_H as described below. The vector p569 is shown in Figure 13; this is one of a family of vectors that are *E. coli/D. discoideum* shuttle vectors using the α -L fucosidase promoter and signal sequence for the secretion of the expressed "ligand binding peptide" to the cell surface. Table 4 shows the results from the immunodot-blot of the expression of the Influenza NC41 V_H FLAG detected by the antiFLAG antibody (M2).

The V_H coding sequence of the monoclonal antibody NC41 was amplified by Polymerase Chain Reaction using the oligonucleotide sequences:

N849

5' CCTTGCCTGCAGGTCGACCTATGGACAGGTGCAGCTGCAGCAG 3'

2145278

34

N863

5' TTACCATGGTTACTTGACCTTAATCAGCAGGACAAATGAAATAAATTTATCATCAT
CATCTTTATAATC 5'

5

10 N849 contains sequence complementary to the N-terminus of the NC41V_H coding region together with the α -L-fucosidase signal sequence and cleavage site, as well as Sall restriction site suitable for cloning into the expression vector p569.

N863 contains sequence complementary to the FLAG coding sequence of the NC41V_H, together with a transmembrane hydrophobic sequence, an NcoI restriction site for cloning and a translation stop codon.

15 DNA of the vector pAV 569 (a gift from W. Nellen, Max-Planck Institute, Martinsreid, Germany) was digested with the restriction enzyme NcoI and Sall, and the cut vector was purified by the standard techniques of agarose electrophoresis and phenol extraction.

20 The PCR amplified and restriction digested NC41V_H FLAG was ligated into the vector and the mixture was transformed into *E. coli*. Recombinant colonies were selected on ampicillin-containing YT plates and recombinant plasmids were recovered, purified and identified using
25 standard techniques. The recombinant plasmid is designated pAV1.

The recombinant plasmid pAV1 was transformed into vegetative cells of *D. discoideum* by the feeding method disclosed in GB-2159821, by Friendlender and Mella.

30 Recombinant *D. discoideum* were selected using the antibiotic G418 at 10 μ g/ml on DMB medium. Recombinant *D. discoideum* amoebae were grown in 2ml cultures of DMB medium containing 10 μ g/ml G418. After growth for 48 hours

at 22°C, dot blot analysis was performed on 100µl aliquots of the culture supernatant, and an anti-FLAG antibody was used to detect the presence of the NC10V_H FLAG antibody fragment in the culture. The results are summarised in Table 4.

Table 4

EXPRESSION OF NC41V_H-FLAG IN *D. DISCOIDEUM*
Immunodetection Dot Blot with Anti-FLAG Antibody

<u><i>D. discoideum</i> clone #</u>		<u>Detection by Antibody</u>
	Negative Control	-
	Negative Control	-
15	Positive Control	+++
	9C Recombinant <i>D. discoideum</i>	+
	9D "	-
	9E "	++
	9F "	-
20	9G "	++
	8B "	+++
	8C "	-
	8D "	+++
	8E "	+++
25	8F "	-
	8G "	+++
	7D "	+
	7E "	-
	7F "	+
30	7G "	-
- = no reaction		
+++ = strong reaction		
+ = weak reaction		

Example 8

Construction of the recombinant vector containing the 1C3Fab for expression in *D. discoideum*. The structure of the parent vector is shown in Figure 13 (p569), and was modified by removal of a *Bam*HI/*Bgl*III fragment by restriction digestion and religation, leaving unique *Xba*I and *Ssp*I sites for the cloning of the antibody coding regions that were constructed by the polymerase chain reaction with the primers:

5' CAGGTCGACTCTAGAGTATGGGAGGTGAGGCTTCTCGAG 3'

5' AAATTATAATTATTATCATCATCATCTTTATAATC 3'

15

and the 1C3Fab coding region (see Sequence I.D. 6) as template. This Fab is a polyfunctional polypeptide as it combines binding activities (TBRs) for glycophorin; and antiFLAG and anti EEF antibodies. Restriction digestion of the PCR products was followed by standard purification, ligation and transformation protocols for construction in *E. coli*. Transformation of *D. discoideum* is effected by feeding transformed *E. coli* (see GB-2159821A) or by standard methods by those skilled in the art as published in the literature such as by calcium phosphate crystals (Nellen et al, Mol. Cell. Biol., 1984 4 2890-2898) or electroporation (Howard et al, 1988, 16 2613-2623) with selection on G418 (geneticin).

30

It will be clearly understood that the invention in its general aspects is not limited to the specific details referred to hereinabove.

File: B:\POWLY23V.SEQ
Description: JLApelB Ly-2+Ly-3 V domains linked MscI-SalI with FLAG tail
From base: 1
To base: 849
Total bases: 849

```
      M K Y   L L P T   A A A   G L L   L L A A   Q P A
1  ATGAAATACC TATTGCCTAC GGCAGCCGCT GGATTGTTAT TACTCGCTGC CCAACCAGCG

      M A K   P Q A P   E L R   I F P   K K M D   A E L
61 ATGGCCAAGC CACAGGCACC CGAACTCCGA ATCTTTCCAA AGAAAATGGA CGCCGAACTT

      G Q K   V D L V   C E V   L G S   V S Q G   C S W
121 GGTCAGAAGG TGGACCTGGT ATGTGAAGTG TTGGGGTCCG TTTCGCAAGG ATGCTCTTGG

      L F Q   N S S S   K L P   Q P T   F V V Y   M A S
181 CTCTTCCAGA ACTCCAGCTC CAAACTCCCC CAGCCCACCT TCGTTGTCTA TATGGCTTCA

      S H N   K I T W   D E K   L N S   S K L F   S A M
241 TCCCACAACA AGATAACGTG GGACGAGAAG CTGAATTCGT CGAAACTGTT TTCTGCCATG

      R D T   N N K Y   V L T   L N K   F S K E   N E G
301 AGGGACACGA ATAATAAGTA CGTTCTCACC CTGAACAAGT TCAGCAAGGA AAACGAAGGC

      Y Y F   C S V I   S N S   V M Y   F S S V   V P V
361 TACTATTTCT GCTCAGTCAT CAGCAACTCG GTGATGTACT TCAGTTCTGT CGTGCCAGTC

      L Q G   G G G S   G G G   G S G   G G G S   L I Q
421 CTTCAGGGTG GCGGAGGCTC AGGCGGTGGT GGATCAGGTG GCGGCGGATC TCTCATTCAG

      T P S   S L L V   Q T N   H T A   K M S C   E V K
481 ACCCCTTCGT CCCTGCTGGT TCAAACCAAC CATACGGCAA AGATGTCCTG TGAGGTAAAA

      S I S   K L T S   I Y W   L R E   R Q D P   K D K
541 AGCATCTCTA AGTTAACAAG CATCTACTGG CTGCGGGAGC GCCAGGACCC CAAGGACAAG

      Y F E   F L A S   W S S   S K G   V L Y G   E S V
601 TACTTTGAGT TCCTGGCCTC CTGGAGTTCT TCCAAAGGAG TTTTGTATGG TGAAAGTGTG

      D K K   R N I I   L E S   S D S   R R P F   L S I
661 GACAAGAAAA GAAATATAAT TCTTGAGTCT TCAGACTCAA GACGGCCCTT TCTCAGTATC

      M N V   K P E D   S D F   Y F C   A T V G   S P K
721 ATGAATGTGA AGCCAGAGGA CAGTGACTTC TACTTCTGCG CGACGGTTGG GAGCCCCAAG

      M V F   G T G T   K L T   V V D   Y K D D   D D K
781 ATGGTCTTTG GGACAGGGAC GAAGCTGACT GTGGTTGATT ACAAGGACGA CGATGACAAG

      * S T
841 TAGTCGACA
```

SEQUENCE I.D. 1

2145278

38

File: B:\CD8ACD8B.SEQ
Description: Human single chain CD8 in pPOW (pelB CD8a and CD8b v dom)
From base: 1
To base: 822
Total bases: 822

```
      M K Y L L P T A A A G L L L L A A Q P A
1  ATGAAATACC TATTGCCTAC GGCAGCCGCT GGATTGTTAT TACTCGCTGC CCAACCAGCG

      M A S Q F R V S P L D R T W N L G E T V
61 ATGGCCAGCC AGTTCCGGGT GTCGCCGCTG GATCGGACCT GGAACCTGGG CGAGACAGTG

      E L K C Q V L L S N P T S G C S W L F Q
121 GAGCTGAAGT GCCAGGTGCT GCTGTCCAAC CCGACGTCGG GCTGCTCGTG GCTCTTCCAG

      P R G A A A S P T F L L Y L S Q N K P K
181 CCGCGCGGCG CCGCCGCCAG TCCCACCTTC CTCCTATACC TCTCCCAAAA CAAGCCCAAG

      A A E G L D T Q R F S G K R L G D T F V
241 GCGGCCGAGG GGCTGGACAC CCAGCGGTTC TCGGGCAAGA GGTTGGGGGA CACCTTCGTC

      L T L S D F R R E N E G Y Y F C S A L S
301 CTCACCCTGA GCGACTTCCG CCGAGAGAAC GAGGGCTACT ATTTCTGCTC GGCCCTGAGC

      N S I M Y F S H F V P V F L P A G G R G
361 AACTCCATCA TGTACTTCAG CCACTTCGTG CCGGTCTTCC TGCCAGCGGG CGGCCGCGGT

      S G G G G S G G G G S L Q Q T P A Y I K
421 TCAGGTGGAG GTGGATCCGG AGGCGGTGGA TCTCTCCAGC AGACCCCTGC ATACATAAAG

      V Q T N K M V M L S C E A K I S L S N M
481 GTGCAAACCA ACAAGATGGT GATGCTGTCC TGCAGAGGCTA AAATCTCCCT CAGTAACATG

      R I Y W L R Q R Q A P S S D S H H E F L
541 CGCATCTACT GGCTGAGACA GCGCCAGGCA CCGAGCAGTG ACAGTCACCA CGAGTTCCTG

      A L W D S A K G T I H G E E V E Q E K I
601 GCCCTCTGGG ATTCCGCAAA AGGGACTATC CACGGTGAAG AGGTGGAACA GGAGAAGATA

      A V F R D A S R F I L N L T S V K P E D
661 GCTGTGTTTC GGGATGCAAG CCGGTTTATT CTCAATCTCA CAAGCGTGAA GCCGGAAGAC

      S G I Y F C M I V G S P E L T F G K G T
721 AGTGGCATCT ACTTCTGCAT GATCGTCGGG AGCCCCGAGC TGACCTTCGG GAAGGGAACI

      Q L S V V D Y K D D D D K *
781 CAGCTGAGTG TGTTGATTA CAAGGACGAC GATGACAAGT AG
```

SEQUENCE I.D. 2

File: B:\MHCA3.SEQ
Description: MHCI a3 H-2K domain in pPOW pelB Msc-EcoRI (no FLAG)
From base: 1
To base: 371
Total bases: 371

```
      M K Y L L P T A A A G L L L L A A Q P A  
1  ATGAAATACC TATTGCCTAC GGCAGCCGCT GGATTGTTAT TACTCGCTGC CCAACCAGCG  
  
      M A K A H V T H H R R P E G D V T L R C  
61 ATGGCCAAGG CCCATGTCAC CCATCACCGC AGACCTGAAG GTGATGTCAC CCTGAGGTGC  
  
      W A L G F Y P A D I T L T W Q L N G D E  
121 TGGGCCCTGG GCTTCTACCC TGCTGACATC ACCCTGACCT GGCAGTTGAA TGGGGACGAG  
  
      L T Q E M E L V E T R P A G D G T F Q K  
181 CTGACCCAGG AAATGGAGCT TGTGGAGACC AGGCCTGCAG GGGATGGAAC CTTCCAGAAG  
  
      W A S V V V P L G K E Q K Y T C H V E H  
241 TGGGCATCTG TGGTGGTGCC TCTTGGGAAG GAGCAGAAGT ACACATGCCA TGTGGAACAT  
  
      E G L P E P L T L R W G K E E P P S S T  
301 GAGGGGCTGC CTGAGCCCCT CACCCTGAGA TGGGGCAAGG AGGAGCCTCC TTCATCCACC  
  
      K * N  
361 AAGTAGAATT C
```

SEQUENCE I.D. 3

2145278

40

Linkerless 1C3 as Constructed in pHFA.

From Hind III site in pHFA to start of gene 3 sequence.

```

1  aag ctt gca tgc aaa ttc tat ttc aag gag aca gtc ata ATG AAA TAC
    L  L  P  T  A  A  A  G  L  L  L  L  A  A  Q  P
49  CTA TTG CCT ACG GCA GCC GCT GGA TTG TTA TTA CTC GCG GCC CAG CCG
    A  M  A  E  V  K  L  Q  E  S  G  G  G  P  V  Q
97  GCC ATG GCC GAG GTG AAG CTG CAG GAG TCT GGA GGT GGC CCG GTA CAA
    P  G  G  S  L  K  L  S  C  A  A  S  G  F  D  F
145  CCT GGA GGA TCC CTG AAA CTC TCC TGT GCA GCC TCA GGA TTC GAT TTT
    S  R  Y  W  M  N  W  V  R  R  A  P  G  K  G  L
193  AGT AGA TAC TGG ATG AAT TGG GTC CGG CGG GCT CCA GGG AAG GGG CTA
    E  W  I  G  E  I  N  Q  Q  S  S  T  I  N  Y  S
241  GAG TGG ATT GGA GAA ATT AAT CAA CAA AGC AGT ACG ATA AAC TAT TCG
    P  P  L  K  D  K  F  I  I  S  R  D  N  A  K  S
289  CCA CCT CTG AAG GAT AAA TTC ATC ATC TCC AGA GAC AAC GCC AAA AGT
    T  L  Y  L  Q  M  N  K  V  R  S  E  D  T  A  L
337  ACG CTG TAC CTG CAA ATG AAC AAA GTG AGA TCT GAG GAC ACA GCC CTT
    Y  Y  C  A  R  L  S  L  T  A  A  G  F  A  Y  W
385  TAT TAT TGT GCA AGA CTT TCT CTT ACT GCG GCA GGG TTT GCT TAC TGG
    G  Q  G  T  L  V  T  V  A  S  D  I  V  M  S  Q
433  GGC CAA GGG ACT CTG GTC ACC GTC GCC TCC GAC ATC GTC ATG TCA CAG
    S  P  S  S  L  A  V  S  V  G  E  K  V  T  M  S
481  TCT CCA TCC TCC CTG GCT GTG TCA GTA GGA GAG AAG GTC ACT ATG AGC
    C  R  S  S  Q  S  L  F  N  S  R  T  R  K  N  Y
529  TGC AGA TCC AGT CAG AGT CTG TTC AAC AGT AGA ACC CGA AAG AAC TAC
    L  T  W  Y  Q  Q  K  P  G  Q  S  P  K  P  L  I
577  TTG ACT TGG TAC CAG CAG AAA CCA GGG CAG TCT CCT AAA CCG CTG ATC
    Y  W  A  S  T  R  E  S  G  V  P  D  R  F  T  G
625  TAC TGG GCA TCC ACT AGG GAA TCT GGG GTC CCT GAT CGC TTC ACA GGC
    S  G  S  G  T  D  F  T  L  T  I  S  S  V  Q  A
673  AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGT GTG CAG GCT
    E  D  L  A  D  Y  Y  C  K  Q  S  Y  N  L  R  T
721  GAA GAC CTG GCA GAT TAT TAC TGC AAG CAA TCT TAT AAT CTT CGG ACG
    F  G  G  G  T  K  L  E  I  N  R  A  A  A  D  Y
769  TTC GGT GGA GGC ACC AAG CTG GAA ATT AAT CGG GCG GCC GCA GAT TAT
    K  D  D  D  D  K  *  A  A  *  T  V  E  S  C  L
817  AAA GAT GAT GAT GAT AAA TAG GCC GCA TAG ACT GTT GAA AGT TGT TTA
    A  K
865  GCA AAA.....
```

SEQUENCE I.D. 4

SUBSTITUTE SHEET (Rule 26)

pelB signal sequence

M K Y L L P T A A A G L L L L A A Q P A
ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCG

H1 Pst1 H10
M A Q V Q L Q Q S G A E L V K P G A S V
ATGGCGCAGGTGCAGCTGCAGCAGTCTGGGGCTGAACTGGTGAAGCCTGGGGCCTCAGTG

H20 H30
R M S C K A S G Y T F T N Y N M Y W V K
AGGATGTCCTGCAAGGCTTCTGGCTACACATTTACCAATTACAACATGTACTGGGTAAAA

H40 H50 H52A
Q S P G Q G L E W I G I F Y P G N G D T
CAGTCACCTGGACAGGGCCTGGAGTGGATTGGAATTTTTTATCCAGGAAATGGTGATACT

H60 H70
S Y N Q K F K D K A T L T A D K S S N T
TCCTACAATCAGAAGTTCAAAGACAAGGCCACATTGACTGCTGACAAATCCTCCAACACA

H80 H82A H82C H90
A Y M Q L S S L T S E D S A V Y Y C A R
GCCTACATGCAGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTACTGTGCAAGA

H100 H100E BstE2
S G G S Y R Y D G G F D Y W G Q G T T V
TCGGGGGGCTCCTATAGATACGACGGAGGCTTTGACTACTGGGGCCAAGGGACCACGGTC

H110 linker L1
T V S G G G G S G G G G S G G G G S D I
ACCGTCTCCGGTGGTGGTGGTTCGGGTGGTGGTGGTTCGGGTGGTGGTTCGGATATC

SacI L10 L20
E L T Q T T S S L S A S L G D R V T I S
GAGCTCACACAGACTACATCCTCCCTGTCTGCCTCTCTGGGAGACAGAGTCACCATCAGT

L30 L40
C R A S O D I S N Y L N W Y Q Q N P D G
TGCAGGGCAAGTCAGGACATTAGTAATTATTTAAACTGGTATCAACAGAATCCAGATGGA

L50 L60
T V K L L I Y Y T S N L H S E V P S R F
ACTGTAAACTCCTGATCTACTACACATCAAATTTACTCAGAAGTCCCATCACGGTTC

L70 L80
S G S G S G T D Y S L T I S N L E Q E D
AGTGGCAGTGGGTCTGGAACAGATTATTCTCTCACCATTAGCAACCTGGAACAAGAAGAT

L90 L100
I A T Y F C Q Q D F T L P F T F G G G T
ATTGCCACTTACTTTTGCCAACAGGATTTTACGCTTCCGTTACGTTTCGGAGGGGGGACC

XhoI FLAG EcoRI
K L E I R D Y K D D D D K * *
AAGCTCGAGATAAGAGACTACAAAGACGATGACGATAAATAATAAGAATTC

SEQUENCE I.D. 5

SUBSTITUTE SHEET (Rule 26)

File: A:\1C3FAB.SEQ
Description: anti-glycophorin 1C3 Fab
From base: 1
To base: 1443
Total bases: 1443

```
1  aaaaaagcGG CCCAGCCGGC CATGGCCGAG GTGAGGCTTC TCGAGTCTGG AGGTGGCCCCG
61  GTACAACCTG GAGGATCCCT GAAACTCTCC TGTGCAGCCT CAGGATTCGA TTTTAGTAGA
121 TACTGGATGA ATTGGgtcCG GCGGGCTCCA GGAAGGGGC TAGAGTGGAT TGGAGAAATT
181 AATCAACAAA GCAGTACGAT AAATATTTCG CCACCTCTGA AGGATAAATT CATCATCTCC
241 AGAGACAACG CCAAAGTAC GCTGTACCTG CAAATGAACA AAGTGAGATC TGAGGACACA
301 GCCCTTTATT ATTGTGCAAG ACTTTCTCTT ACTGCGGCAG GGTTTGCTTA CTGGGGCCAA
361 GGGACTCTGG TCACTGTCTC TGCAGCCAAA ACGACACCCC CATCTGTCTA TCCACTGGCC
421 CCTGGATCTG CTGCCCAAAC TAACTCCATG GTGACCCTGG GATGCCTGGT CAAGGGCTAT
481 TTCCCTGAGC CAGTGACAGT GACCTGGAAC TCTGGATCCC TGTCCAGCGG TGTGCACACC
541 TTCCCAGCTG TCCTGCAGTC TGACCTCTAC ACTCTGAGCA GCTCAGTGAC TGTCCCCTCC
601 AGCACCTGGC CCAGCGAGAC CGTCACCTGC AACGTTGCCC ACCCGGCCAG CAGCACCAAG
661 GTGGACAAGA AAATTgaaga attttaatta aaacatggaa ataaaGTGAA ACAAAGCACT
721 ATTGCACTGG CACTCTTACC GTTACTGTTT ACCCCGGTAA CCAAAGCCGA CATCGTCATG
781 TCACAGTCTC CATCCTCCCT GGCTGTGTCA GTAGGAGAGA AGGTCACTAT GAGCTGCAGA
841 TCCAGTCAGA GTCTGTTCAA CAGTAGAACC CGAAAGAACT ACTTGACTTG GTACCAGCAG
901 AAACCAGGGC AGTCTCCTAA ACCGCTGATC TACTGGGCAT CCACTAGGGA ATCTGGGGTC
961 CCTGATCGCT TCACAGGCAG TGGATCTGGG ACAGATTTCA CTCTCACCAT CAGCAGTGTG
1021 CAGGCTGAAG ACCTGGCAGA TTATTACTGC AAGCAATCTT ATAATCTTCG GACGTTCCGT
1081 GGAGGCACCA AGCTGGAAAT TAAACGGGCT GATGCTGCAG TATCCATCTT CCCACCATCC
1141 AGTGAGCAGT TAACATCTGG ATCTGGAGGT GCCTCAGTCG TGTGCTTCTT GAACAACTTC
1201 TACCCCAAAG ACATCAATGT CAAGTGGAAG ATTGATGGCA GTGAACGACA AAATGGCGTC
1261 CTGAACAGTT GGA CTGATCA GGACAGCAAA GACAGCACCT ACAGCATGAG CAGCACCTTC
1321 ACGTTGACCA AGGACGAGTA TGAACGACAT AACAGCTATA CCTGTGAGGC CACTCACAAG
1381 ACATCAACTT CACCCATTGT CAAGAGCTTC AACAGGggaG AGTGTgcggc cgcagattat
1441 aaa
```

SEQUENCE I.D. 6

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A polypeptide which has specificity for a target molecule and which can self-associate to form stable dimers or multimers which can bind to the target molecule, the polypeptide comprising an immunoglobulin-like V_L domain linked to an immunoglobulin-like V_H domain, wherein said V_L domain and said V_H domain are:

- (a) directly linked, or
- (b) linked via a peptide linker that consists of a chain of up to 10 amino acids.

2. The polypeptide according to claim 1, wherein said peptide linker is a hydrophilic peptide.

3. The polypeptide according to claim 1, wherein said peptide linker is a chain of up to 10 amino acids.

4. The polypeptide according to claim 1, wherein said V_H domain and said V_L domain are directly linked.

5. The polypeptide according to claim 2, wherein said peptide linker is 5 or 10 amino acids.

6. The polypeptide according to claim 5, wherein said peptide linker is Gly₄Ser or (Gly₄Ser)₂.

7. The polypeptide according to claim 2, wherein said peptide linker connects the carboxyl terminal of the V_H domain to the amino terminal of the V_L domain.

8. The polypeptide according to claim 2, wherein said peptide linker connects the carboxyl terminal of the V_L domain to the amino terminal of the V_H domain.

9. The polypeptide according to claim 4, wherein up to 13 amino acids of the V_L domain or the V_H domain at the junction of said V_L and said V_H domains are deleted.

10. The polypeptide according to claim 9, wherein up to 13 amino acids of the V_L domain or the V_H domain immediately following the junction of said V_L and said V_H domains are deleted.

11. The polypeptide according to claim 1, wherein said V_H and said V_L domains have different binding specificities.

12. A dimer or multimer comprising two or more polypeptides according to any one of claims 1 to 11.

13. The dimer or multimer according to claim 12, which is bifunctional.

14. The multimer according to claim 12, which is polyfunctional.

15. The polypeptide according to any one of claims 1 to 11, or a dimer or multimer according to any one of claims 12 to 14, in which the V_L domain, the V_H domain, or the V_L and V_H domains is capable of binding to a polypeptide, wherein said polypeptide is a glycophorin, a red blood cell surface

protein, influenza virus neuraminidase, a viral antigen, an antibody, a member of the Ig superfamily, transforming growth factor- α (TGF α), a tumour marker, a cell surface protein, or leukaemia inhibitory factor (LIF).

16. A composition consisting essentially of dimeric or multimeric polypeptides together with a carrier, wherein said dimeric or multimeric polypeptides are formed by non-covalent association of monomeric units, wherein

(a) each of said monomeric units comprises an immunoglobulin-like V_L domain covalently linked to an immunoglobulin-like V_H domain and

(b) said V_L domain and said V_H domain are directly linked, or

(c) said V_L domain and said V_H domain are linked via a peptide linker that consists of a chain of up to 10 amino acids.

17. The composition according to claim 16, wherein said composition consisting essentially of dimeric units.

18. The composition according to claim 16, wherein said composition consisting essentially of multimeric units.

19. The composition according to claim 17, wherein said V_H and V_L domains in said monomeric units have different binding specificities and wherein said dimeric units are bifunctional.

20. The composition according to claim 18, wherein said V_H and V_L domains in said monomeric units have different binding

specificities and said multimeric units are bifunctional or polyfunctional.

21. A polynucleotide molecule encoding a polypeptide according to any one of claims 1 to 11.

22. A DNA construct comprising the polynucleotide of claim 21, in which the polynucleotide is DNA.

23. A vector comprising the polynucleotide molecule according to claim 21.

24. A host cell transformed with a vector according to claim 23.

25. A method of producing a polypeptide, comprising culturing the host cell according to claim 24 and then isolating said polypeptide.

26. A method for producing a polynucleotide encoding a polypeptide having modified characteristics of affinity, specificity or avidity comprising the step of subjecting DNA encoding the polypeptide according to any one of claims 1 to 11 to affinity maturation to obtain a sub-population of polynucleotides encoding said polypeptide having modified characteristics of affinity, specificity or avidity.

27. The method according to claim 26, wherein said DNA encoding said polypeptide is present in a replication-competent element or display vector.

28. The method according to claim 27, wherein the display vector is a bacteriophage, filamentous bacteriophage, a virus, a bacteria, a yeast, a slime mould or a mammalian cell.

29. The method according to claim 28, wherein the display vector is pHFA.

30. The method according to claim 29, wherein the DNA is subjected to mutagenesis using a mutator strain of *Escherichia coli*.

31. A method of producing a polypeptide having modified characteristics of affinity, specificity or avidity, comprising:

(a) isolating DNA encoding framework regions of one or more of the polypeptides according to any one of claims 1 to 11 by means of polymerase chain reaction;

(b) inserting said DNA into one or more display vectors;

(c) selecting a sub-population of display vectors displaying said one or more polypeptides;

(d) subjecting said sub-population to affinity maturation in order to obtain a sub-population of display vectors displaying polypeptides having modified affinity, specificity or avidity;

(e) inserting DNA encoding said polypeptides having modified affinity, specificity or avidity into a high level expression vector;

(f) transferring said high level expression vector into an appropriate expression host cell;

(g) expressing the polypeptide having said modified affinity, specificity or avidity; and

(h) isolating the polypeptide thus produced.

32. A pharmaceutical composition comprising the polypeptide according to any one of claims 1 to 11, or the dimer or multimer according to any one of claims 12 to 14, together with a pharmaceutically acceptable carrier.

33. A pharmaceutical composition comprising the composition according to any one of claims 16 to 20, wherein said carrier is a pharmaceutically acceptable carrier.

34. A diagnostic reagent comprising the polypeptide according to any one of claims 1 to 11, the dimer or multimer according to any one of claims 12 to 14 or the composition according to any one of claims 16 to 20, together with a diluent.

35. Use of the pharmaceutical composition according to claim 32 or 33 for the manufacture of a medicament.

36. Use of the pharmaceutical composition according to claim 32 or 33 for the manufacture of a medicament capable of binding to a polypeptide, wherein said polypeptide is a glycophorin, a red blood cell surface protein, influenza virus neuraminidase, a viral antigen, an antibody, a member of the Ig superfamily, transforming growth factor- α (TGF α), a tumour marker, a cell surface protein, or leukaemia inhibitory factor (LIF).

37. A diagnostic kit for use in the detection of a polypeptide, wherein said polypeptide is a glycophorin, a red blood cell surface protein, influenza virus neuraminidase, a viral antigen, an antibody, a member of the Ig superfamily, transforming growth factor- α (TGF α), a tumour marker, a cell surface protein, or leukaemia inhibitory factor (LIF) in a sample comprising carrier means being compartmentalized to receive in close confinement therein one or more containers comprising:

(a) a first container containing a polypeptide which can self-associate to form stable dimers, multimers, aggregates or arrays comprising an immunoglobulin-like V_L domain linked to an immunoglobulin-like V_H domain, wherein said V_L domain and said V_H domain are directly linked; and optionally including at least two elements selected from the group of:

- (i) a receptacle for sample collection;
- (ii) a solid support for immobilization of any polypeptides present within a sample;
- (iii) a contained amount of a blocking solution comprising a protein, a buffer and a preservative, wherein said protein will bind to any unoccupied sites of the solid support;
- (iv) contained amount of a wash solution comprising a buffer and a preservative;
- (v) a contained amount of a signal developing reagent whereby said reagent will react with said antibody to generate a signal as a positive result;
- (vi) a contained amount of a standard solution of said polypeptide which can self-associate to form stable dimers, multimers, aggregates or arrays comprising

an immunoglobulin-like V_L domain linked to an immunoglobulin-like V_H domain, wherein said V_L domain and said V_H domain are directly linked, or one which is substantially identical to said polypeptide, in a buffer; and
(vii) instructions for use.

38. A diagnostic kit for use in the detection of a polypeptide, wherein said polypeptide is a glycoporphin, a red blood cell surface protein, influenza virus neuraminidase, a viral antigen, an antibody, a member of the Ig superfamily, transforming growth factor- α (TGF α), a tumour marker, a cell surface protein, or leukaemia inhibitory factor (LIF) in a sample comprising carrier means being compartmentalized to receive in close confinement therein one or more containers comprising:

(a) a first container containing a polypeptide which can self-associate to form stable dimers, multimers, aggregates or arrays comprising an immunoglobulin-like V_L domain linked to an immunoglobulin-like V_H domain, wherein said V_L domain and said V_H domain are linked via a peptide linker that consists of a chain of up to 10 amino acids; and optionally including at least two elements selected from the group of:

- (i) a receptacle for sample collection;
- (ii) a solid support for immobilization of any polypeptides present within a sample;
- (iii) a contained amount of a blocking solution comprising a protein, a buffer and a preservative, wherein said protein will bind to any unoccupied sites of the solid support;

- (iv) a contained amount of a wash solution comprising a buffer and a preservative;
- (v) a contained amount of a signal developing reagent whereby said reagent will react with said antibody to generate a signal as a positive result;
- (vi) a contained amount of a standard solution of said polypeptide which can self-associate to form stable dimers, multimers, aggregates or arrays comprising an immunoglobulin-like V_L domain linked to an immunoglobulin-like V_H domain, wherein said V_L domain and said V_H domain are linked via a peptide linker that consists of a chain of up to 10 amino acids, or one which is substantially identical to said polypeptide, in a buffer; and
- (vii) instructions for use.

1/18




FIGURE 1

2/18




FIGURE 2

3/18




FIGURE 3

4/18




FIGURE 4

5/18




FIGURE 5

b/18

DELETION LINKER CONSTRUCTS

<u>Construct</u>	<u>Complementary Oligonucleotide Pair</u>
((Gly4)Ser)3	5' GTC ACC GTC TCC (GGT GGT GGT GGT TCG)3 GAT ATC GAG CT 3' 3' G CAG AGG (CCA CCA CCA CCA AGC)3 CTA TAG C 5'
((Gly4)Ser)2	5' GTC ACC GTC TCC (GGT GGT GGT GGT TCG)2 GAT ATC CAG CT 3' 3' G CAG AGG (CCA CCA CCA CCA AGC)2 CTA TAG C 5'
(Gly4)Ser)	5' GTC ACC GTC TCC GGT GGT GGT GGT TCG GAT ATC GAG CT 3' 3' G CAG AGG CCA CCA CCA CCA AGC CTA TAG C 5'
-	5' GTC ACC GTC TCC GAT ATC GAG CT 3' 3' G CAG AGG CTA TAG C 5'
-13.VL	5' GGG ACC ACG GTC ACC GTC TCC TCA GCC TCT CTG GGA GAC AGA GTC ACC 5' GTC GAC GAA TTC TTA TTA TTT ATC GTC ATC ATC TTT GTA GTC 3'

FIGURE 6

7/18




FIGURE 7

8/ 18

DELETION LINKER CONSTRUCTS

<u>Linker Unit</u>	<u>Observed Forms</u>	<u>Activity</u>
((Gly4)Ser) ₃	Monomers & dimers	++
((Gly4)Ser) ₂	Dimers	++
(Gly4)Ser	Dimers	++
-	Multimers	+
-13.VL Δ	Multimers	

FIGURE 8

9/18




FIGURE 9a

10/ 18




FIGURE 9b

11/ 18




FIGURE 10

12/18




FIGURE 11

13/18

ELISA on Mutated 1C3scFv Culture S/N Selection of Individual Colonies




FIGURE 12

14/18




FIGURE 13

15/18




FIGURE 14

16/18




FIGURE 15

17/18




FIGURE 16

18/18


Formation of Dimeric scFvs


FIGURE 17

DELETION LINKER CONSTRUCTS

DISTANCE




54 Å



36 Å



18 Å



0 Å

