



(43) International Publication Date
11 June 2015 (11.06.2015)

(10) International Publication Number
WO 2015/083008 A1

(51) International Patent Classification:

A61K 31/517 (2006.01) *A61P 35/00* (2006.01)
A61K 31/519 (2006.01) *A61P 35/02* (2006.01)
A61K 31/52 (2006.01) *A61P 35/04* (2006.01)
A61K 45/06 (2006.01) *A61P 43/00* (2006.01)

(21) International Application Number:

PCT/IB2014/003121

(22) International Filing Date:

5 December 2014 (05.12.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/912,515 5 December 2013 (05.12.2013) US
61/974,665 3 April 2014 (03.04.2014) US
62/035,777 11 August 2014 (11.08.2014) US

(71) Applicant: **ACERTA PHARMA B.V.** [NL/NL]; Pivot
Park Room RK 2111, Molenweg 79, NL-5349 AC Oss
(NL).

(72) Inventors: **HAMDY, Ahmed**; 640 Henry Cowell Drive,
Santa Cruz, CA 95060 (US). **ROTHBAUM, Wayne**; 101
Central Park West, PHC, New York, NY 10023 (US).
IZUMI, Raquel; 3437 Brittan Avenue, San Carlos, CA
94070 (US). **LANNUTTI, Brian**; 627 Glencrest Place, So-
lana Beach, CA 92075 (US). **COVEY, Todd**; 3437 Brittan
Avenue, San Carlos, CA 94070 (US). **ULRICH, Roger**;
22525 SE 46th Place, Sammamish, WA 98075 (US).
JOHNSON, Dave; 4442 54th Avenue SW, Seattle, WA
98116 (US). **BARF, Tjeerd**; St. Luciastraat 7, NL-5371

AS Ravenstein (NL). **KAPTEIN, Allard**; Nonnenstraat
14, NL-5301 BJ Zaltmommel (NL).

(74) Agent: **HANSEN, Norbert**; Maiwald Patentanwalts
GmbH, Elisenhof, Elisenstrasse 3, 80335 München (DE).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: THERAPEUTIC COMBINATION OF A PI3K INHIBITOR AND A BTK INHIBITOR

(57) Abstract: In some embodiments, the invention includes a therapeutic combination of a phosphoinositide 3-kinase (PI3K) inhibitor, including PI3K inhibitors selective for the γ - and δ -isoforms and selective for both γ - and δ -isoforms, and a Bruton's tyrosine kinase (BTK) inhibitor. In some embodiments, the invention includes therapeutic methods of using a BTK inhibitor and a PI3K- δ inhibitor to treat solid tumor cancers by modulation of the tumor microenvironment, including macrophages, monocytes, mast cells, helper T cells, cytotoxic T cells, regulatory T cells, natural killer cells, myeloid-derived suppressor cells, regulatory B cells, neutrophils, dendritic cells, and fibroblasts.



WO 2015/083008 A1

THERAPEUTIC COMBINATION OF A PI3K INHIBITOR AND A BTK INHIBITOR

FIELD OF THE INVENTION

[001] A therapeutic combination of a phosphoinositide 3-kinase (PI3K) inhibitor and a Bruton's Tyrosine Kinase (BTK) inhibitor and uses of the therapeutic combination are disclosed herein.

BACKGROUND OF THE INVENTION

[002] PI3K kinases are members of a unique and conserved family of intracellular lipid kinases that phosphorylate the 3'-OH group on phosphatidylinositols or phosphoinositides. PI3K kinases are key signaling enzymes that relay signals from cell surface receptors to downstream effectors. The PI3K family comprises 15 kinases with distinct substrate specificities, expression patterns, and modes of regulation. The class I PI3K kinases (p110 α , p110 β , p110 δ , and p110 γ) are typically activated by tyrosine kinases or G-protein coupled receptors to generate PIP3, which engages downstream effectors such as those in the Akt/PDK1 pathway, mTOR, the Tec family kinases, and the Rho family GTPases.

[003] The PI3K signaling pathway is known to be one of the most highly mutated in human cancers. PI3K signaling is also a key factor in disease states including hematologic malignancies, non-Hodgkin lymphoma (such as diffuse large B-cell lymphoma), allergic contact dermatitis, rheumatoid arthritis, osteoarthritis, inflammatory bowel diseases, chronic obstructive pulmonary disorder, psoriasis, multiple sclerosis, asthma, disorders related to diabetic complications, and inflammatory complications of the cardiovascular system such as acute coronary syndrome. The role of PI3K in cancer has been discussed, for example, in J. A. Engleman, *Nat. Rev. Cancer* **2009**, 9, 550-562. The PI3K- δ and PI3K- γ isoforms are preferentially expressed in normal and malignant leukocytes.

[004] The delta (δ) isoform of class I PI3K (PI3K- δ) is involved in mammalian immune system functions such as T-cell function, B-cell activation, mast cell activation, dendritic cell function, and neutrophil activity. Due to its role in immune system function, PI3K- δ is also involved in a number of diseases related to undesirable immune response such as allergic reactions, inflammatory diseases, inflammation mediated angiogenesis, rheumatoid arthritis,

auto-immune diseases such as lupus, asthma, emphysema and other respiratory diseases. The gamma (γ) isoform of class I PI3K (PI3K- γ) is also involved in immune system functions and plays a role in leukocyte signaling and has been implicated in inflammation, rheumatoid arthritis, and autoimmune diseases such as lupus.

[005] Downstream mediators of the PI3K signal transduction pathway include Akt and mammalian target of rapamycin (mTOR). One important function of Akt is to augment the activity of mTOR, through phosphorylation of TSC2 and other mechanisms. mTOR is a serine-threonine kinase related to the lipid kinases of the PI3K family and has been implicated in a wide range of biological processes including cell growth, cell proliferation, cell motility and survival. Disregulation of the mTOR pathway has been reported in various types of cancer.

[006] In view of the above, PI3K inhibitors are prime targets for drug development, as described in J. E. Kurt and I. Ray-Coquard, *Anticancer Res.* **2012**, 32, 2463-70. Several PI3K inhibitors are known, including those that are PI3K- δ inhibitors, PI3K- γ inhibitors, and PI3K- δ,γ inhibitors.

[007] Bruton's Tyrosine Kinase (BTK) is a Tec family non-receptor protein kinase expressed in B cells and myeloid cells. The function of BTK in signaling pathways activated by the engagement of the B cell receptor (BCR) and FCER1 on mast cells is well established. Functional mutations in BTK in humans result in a primary immunodeficiency disease characterized by a defect in B cell development with a block between pro- and pre-B cell stages. The result is an almost complete absence of B lymphocytes, causing a pronounced reduction of serum immunoglobulin of all classes. These findings support a key role for BTK in the regulation of the production of auto-antibodies in autoimmune diseases.

[008] Other diseases with an important role for dysfunctional B cells are B cell malignancies. The reported role for BTK in the regulation of proliferation and apoptosis of B cells indicates the potential for BTK inhibitors in the treatment of B cell lymphomas. BTK inhibitors have thus been developed as potential therapies, as described in O. Cruz et al., *OncoTargets and Therapy* **2013**, 6, 161-176.

[009] In many solid tumors, the supportive microenvironment (which may make up the majority of the tumor mass) is a dynamic force that enables tumor survival. The tumor

microenvironment is generally defined as a complex mixture of “cells, soluble factors, signaling molecules, extracellular matrices, and mechanical cues that promote neoplastic transformation, support tumor growth and invasion, protect the tumor from host immunity, foster therapeutic resistance, and provide niches for dominant metastases to thrive,” as described in Swartz et al., *Cancer Res.*, **2012**, 72, 2473. Although tumors express antigens that should be recognized by T cells, tumor clearance by the immune system is rare because of immune suppression by the microenvironment. Addressing the tumor cells themselves with e.g. chemotherapy has also proven to be insufficient to overcome the protective effects of the microenvironment. New approaches are thus urgently needed for more effective treatment of solid tumors that take into account the role of the microenvironment.

[0010] The present invention provides the unexpected finding that the combination of a PI3K inhibitor with a BTK inhibitor is effective in the treatment of any of several types of cancers such as leukemia, lymphoma and solid tumor cancers.

SUMMARY OF THE INVENTION

[0011] In an embodiment, the invention provides a composition comprising a PI3K inhibitor and a BTK inhibitor in combination. This composition is typically a pharmaceutical composition.

[0012] In an embodiment, the invention provides a composition comprising a PI3K- γ inhibitor and a BTK inhibitor in combination. This composition is typically a pharmaceutical composition.

[0013] In an embodiment, the invention provides a composition comprising a PI3K- δ inhibitor and a BTK inhibitor in combination. This composition is typically a pharmaceutical composition.

[0014] In an embodiment, the invention provides a composition comprising a PI3K- γ,δ inhibitor and a BTK inhibitor in combination. This composition is typically a pharmaceutical composition.

[0015] In an embodiment, the invention provides a pharmaceutical composition comprising a PI3K inhibitor and a BTK inhibitor in combination for use in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0016] In an embodiment, the invention provides a pharmaceutical composition comprising a PI3K- γ inhibitor and a BTK inhibitor in combination for use in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0017] In an embodiment, the invention provides a pharmaceutical composition comprising a PI3K- δ inhibitor and a BTK inhibitor in combination for use in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0018] In an embodiment, the invention provides a pharmaceutical composition comprising a PI3K- γ,δ inhibitor and a BTK inhibitor in combination for use in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0019] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K inhibitor and a composition comprising a BTK inhibitor. These compositions are typically both pharmaceutical compositions. The kit is for co-administration of the PI3K inhibitor and the BTK inhibitor, either simultaneously or separately.

[0020] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- γ inhibitor and a composition comprising a BTK inhibitor. These compositions are typically both pharmaceutical compositions. The kit is for co-administration of the PI3K- γ inhibitor and the BTK inhibitor, either simultaneously or separately.

[0021] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- δ inhibitor and a composition comprising a BTK inhibitor. These compositions are typically both pharmaceutical compositions. The kit is for co-administration of the PI3K- δ inhibitor and the BTK inhibitor, either simultaneously or separately.

[0022] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- γ,δ inhibitor and a composition comprising a BTK inhibitor. These compositions are typically both pharmaceutical compositions. The kit is for co-administration of the PI3K- γ,δ inhibitor and the BTK inhibitor, either simultaneously or separately.

[0023] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K inhibitor and a composition comprising a BTK inhibitor for use in the treatment of leukemia, lymphoma or a solid tumor cancer. The compositions are typically both

pharmaceutical compositions. The kit is for use in co-administration of the PI3K inhibitor and the BTK inhibitor, either simultaneously or separately, in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0024] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- γ inhibitor and a composition comprising a BTK inhibitor for use in the treatment of leukemia, lymphoma or a solid tumor cancer. The compositions are typically both pharmaceutical compositions. The kit is for use in co-administration of the PI3K- γ inhibitor and the BTK inhibitor, either simultaneously or separately, in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0025] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- δ inhibitor and a composition comprising a BTK inhibitor for use in the treatment of leukemia, lymphoma or a solid tumor cancer. The compositions are typically both pharmaceutical compositions. The kit is for use in co-administration of the PI3K- δ inhibitor and the BTK inhibitor, either simultaneously or separately, in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0026] In an embodiment, the invention provides a kit comprising a composition comprising a PI3K- γ,δ inhibitor and a composition comprising a BTK inhibitor for use in the treatment of leukemia, lymphoma or a solid tumor cancer. The compositions are typically both pharmaceutical compositions. The kit is for use in co-administration of the PI3K- γ,δ inhibitor and the BTK inhibitor, either simultaneously or separately, in the treatment of leukemia, lymphoma or a solid tumor cancer.

[0027] In an embodiment, the invention provides a method of treating leukemia, lymphoma or a solid tumor cancer in a subject, comprising co-administering to a mammal in need thereof a therapeutically effective amount of a PI3K inhibitor and a BTK inhibitor.

[0028] In an embodiment, the invention provides a method of treating leukemia, lymphoma or a solid tumor cancer in a subject, comprising co-administering to a mammal in need thereof a therapeutically effective amount of a PI3K- γ inhibitor and a BTK inhibitor.

[0029] In an embodiment, the invention provides a method of treating leukemia, lymphoma or a solid tumor cancer in a subject, comprising co-administering to a mammal in need thereof a therapeutically effective amount of a PI3K- δ inhibitor and a BTK inhibitor.

[0030] In an embodiment, the invention provides a method of treating leukemia, lymphoma or a solid tumor cancer in a subject, comprising co-administering to a mammal in need thereof a therapeutically effective amount of a PI3K- γ,δ inhibitor and a BTK inhibitor.

[0031] The BTK inhibitor, in one specific embodiment, is a compound of Formula (XVIII), or a pharmaceutically acceptable salt thereof. The PI3K inhibitor, in one specific embodiment, is a PI3K- δ inhibitor, in particular a compound of Formula IX, or a pharmaceutically acceptable salt thereof. In one specific embodiment, the BTK inhibitor is a compound of Formula (XVIII) or a pharmaceutically acceptable salt thereof, and the PI3K inhibitor is a PI3K- δ inhibitor, in particular a compound of Formula IX, or a pharmaceutically acceptable salt thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings.

[0033] FIG. 1 illustrates the sensitivity of the TMD8 diffuse large B cell lymphoma (DLBCL) cell line to individual treatment with the BTK inhibitor of Formula XVIII (“Tested Btk Inhibitor”) and the PI3K inhibitor of Formula IX (“Tested PI3K Inhibitor”) and combined treatment with Formula XVIII and Formula IX (“Btki + PI3Ki”) at different concentrations. The concentration of the first agent in the combination (the BTK inhibitor) and the concentration of the individual agents is given on the x-axis, and the concentration of the added PI3K inhibitor in combination with the BTK inhibitor is given in the legend.

[0034] FIG. 2 illustrates the sensitivity of the MINO mantle cell lymphoma cell to individual treatment with the BTK inhibitor of Formula XVIII (“Tested Btk Inhibitor”) and the PI3K inhibitor of Formula IX (“Tested PI3K Inhibitor”) and combined treatment with Formula XVIII and Formula IX (“Btki + PI3Ki”) at different concentrations. The concentration of the first agent in the combination (the BTK inhibitor) and the concentration of the individual agents is given on

the x-axis, and the concentration of the added PI3K inhibitor in combination with the BTK inhibitor is given in the legend.

[0035] FIG. 3 illustrates the proliferative activity in primary mantle cell lymphoma cells of Formula XVIII (“Tested Btki”) and Formula IX (“Tested PI3Ki”). The percentage viability of cells (“% viability”, y-axis) is plotted versus the concentration of the Formula XVIII (“[Tested Btk Inhibitor]”, x-axis). The concentration of the individual BTK and PI3K inhibitors (i.e. not in combination) are also given on the x-axis.

[0036] FIG. 4 illustrates the interaction index of the combination of the BTK inhibitor of Formula XVIII and the PI3K inhibitor of Formula IX in primary mantle cell lymphoma cells.

[0037] FIG. 5 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include Maver-1 (B cell lymphoma, mantle), Jeko (B cell lymphoma, mantle), CCRF (B lymphoblast, acute lymphoblastic leukemia), and SUP-B15 (B lymphoblast, acute lymphoblastic leukemia). The dose-effect curves for these cell lines are given in FIG. 6, FIG. 7, FIG. 8, and FIG. 9. ED25, ED50, ED75, and ED90 refer to the effective doses causing 25%, 50%, 75%, and 90% of the maximum biological effect (proliferation).

[0038] FIG. 6 illustrates the dose-effect curves obtained for the tested Maver-1 cell line (B cell lymphoma, mantle) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0039] FIG. 7 illustrates the dose-effect curves obtained for the tested Jeko cell line (B cell lymphoma, mantle) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0040] FIG. 8 illustrates the dose-effect curves obtained for the tested CCRF cell line (B lymphoblast, acute lymphoblastic leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis

(“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0041] FIG. 9 illustrates the dose-effect curves obtained for the tested SUP-B15 cell line (B lymphoblast, acute lymphoblastic leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0042] FIG. 10 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include Jeko (B cell lymphoma, mantle cell lymphoma) and SU-DHL-4 (activated B cell like (ABC) diffuse large B cell lymphoma). The dose-effect curves for these cell lines are given in FIG. 11 and FIG. 12.

[0043] FIG. 11 illustrates the dose-effect curves obtained for the tested Jeko cell line (B cell lymphoma, mantle) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0044] FIG. 12 illustrates the dose-effect curves obtained for the tested SU-DHL-4 cell line (diffuse large B cell lymphoma, ABC) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0045] FIG. 13 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include CCRF (B lymphoblast, acute lymphoblastic leukemia), SUP-B15 (B lymphoblast, acute lymphoblastic leukemia), JVM-2 (prolymphocytic leukemia), Ramos (Burkitt’s lymphoma), and Mino (mantle cell lymphoma). The dose-effect curves for these cell lines are given in FIG. 14, FIG. 15, FIG. 16, and FIG. 17. No dose-effect curve is given for Ramos (Burkitt’s lymphoma) because of negative slope.

[0046] FIG. 14 illustrates the dose-effect curves obtained for the tested CCRF cell line (B lymphoblast, acute lymphoblastic leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0047] FIG. 15 illustrates the dose-effect curves obtained for the tested SUP-B15 cell line (B lymphoblast, acute lymphoblastic leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0048] FIG. 16 illustrates the dose-effect curves obtained for the tested JVM-2 cell line (prolymphocytic leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0049] FIG. 17 illustrates the dose-effect curves obtained for the tested Mino cell line (mantle cell lymphoma) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0050] FIG. 18 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include Raji (B lymphocyte, Burkitt's lymphoma), SU-DHL-1 (DLBCL-ABC), and Pfeiffer (follicular lymphoma). The dose-effect curves for these cell lines are given in FIG. 19, FIG. 20, and FIG. 21.

[0051] FIG. 19 illustrates the dose-effect curves obtained for the tested Raji cell line (B lymphocyte, Burkitt's lymphoma) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0052] FIG. 20 illustrates the dose-effect curves obtained for the tested SU-DHL-1 cell line (DLBCL-ABC) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0053] FIG. 21 illustrates the dose-effect curves obtained for the tested Pfeiffer cell line (follicular lymphoma) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0054] FIG. 22 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include Ly1 (Germinal center B-cell like diffuse large B-cell lymphoma, DLBCL-GCB), Ly7 (DLBCL-GCB), Ly19 (DLBCL-GCB), SU-DHL-2 (Activated B-cell like diffuse large B-cell lymphoma, DLBCL-ABC), and DOHH2 (follicular lymphoma, FL). The dose-effect curves for these cell lines are given in FIG. 23, FIG. 24, FIG. 25, and FIG. 26, except for the Ly19 cell line, which is not graphed because of a negative slope.

[0055] FIG. 23 illustrates the dose-effect curves obtained for the tested Ly1 cell line (DLBCL-GCB) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0056] FIG. 24 illustrates the dose-effect curves obtained for the tested Ly7 cell line (DLBCL-GCB) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0057] FIG. 25 illustrates the dose-effect curves obtained for the tested DOHH2 cell line (FL) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0058] FIG. 26 illustrates the dose-effect curves obtained for the tested SU-DHL-2 cell line (DLBCL-ABC) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0059] FIG. 27 illustrates the synergy observed in certain cell lines when Formula (XVIII) and Formula (IX) are combined. The tested cell lines include U937 (histiocytic lymphoma and/or myeloid), K562 (leukemia, myeloid, and/or chronic myelogenous leukemia), Daudi (human Burkitt's lymphoma), and SU-DHL-6 (DLBCL-GCB and/or peripheral T-cell lymphoma, PTCL). The dose-effect curves for these cell lines are given in FIG. 28, FIG. 29, FIG. 30, and FIG. 31.

[0060] FIG. 28 illustrates the dose-effect curves obtained for the tested U937 cell line (histiocytic lymphoma and/or myeloid) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0061] FIG. 29 illustrates the dose-effect curves obtained for the tested K562 cell line (leukemia, myeloid, and/or chronic myelogenous leukemia) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0062] FIG. 30 illustrates the dose-effect curves obtained for the tested Daudi cell line (human Burkitt's lymphoma) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0063] FIG. 31 illustrates the dose-effect curves obtained for the tested SU-DHL-6 cell line (DLBCL-GCB and/or PTCL) using combined dosing of the BTK inhibitor of Formula (XVIII) (“Inh.1”) and the PI3K- δ inhibitor of Formula (IX) (“Inh.3”). The y-axis (“Effect”) is given in units of Fa (fraction affected) and the x-axis (“Dose”) is given in linear units of μM .

[0064] FIG. 32 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include SU-DHL-6 (DLBCL-GCB or PTCL), TMD-8 (DLBCL-ABC), HBL-1 (DLBCL-ABC), and Rec-1 (follicular lymphoma). The dose-effect curves for these cell lines are given in FIG. 34, FIG. 35, FIG. 36, and FIG. 37.

[0065] FIG. 33 illustrates the synergy observed in certain cell lines when the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are combined. The tested cell lines include SU-DHL-6 (DLBCL-GCB or PTCL), TMD-8 (DLBCL-ABC), HBL-1 (DLBCL-ABC), and Rec-1 (follicular lymphoma). All corresponding CIs are shown for each of the combinations tested as listed on the x axis.

[0066] FIG. 34 illustrates the dose-effect curves obtained for the tested SU-DHL-6 cell line (DLBCL-GCB or PTCL) cell line using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0067] FIG. 35 illustrates the dose-effect curves obtained for the tested TMD-8 cell line (DLBCL-ABC) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0068] FIG. 36 illustrates the dose-effect curves obtained for the tested HBL-1 cell line (DLBCL-ABC) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0069] FIG. 37 illustrates the dose-effect curves obtained for the tested Rec-1 cell line (follicular lymphoma) using combined dosing of the BTK inhibitor of Formula (XVIII) ("Inh.1") and the PI3K- δ inhibitor of Formula (IX) ("Inh.3"). The y-axis ("Effect") is given in units of Fa (fraction affected) and the x-axis ("Dose") is given in linear units of μ M.

[0070] FIG. 38 illustrates tumor growth suppression in an orthotopic pancreatic cancer model. Mice were dosed orally with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the

PI3K- δ inhibitor of Formula (IX), or a combination of both drugs. The statistical p-value (presumption against null hypothesis) is shown for each tested single agent and for the combination against the vehicle.

[0071] FIG. 39 illustrates the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K- δ inhibitor of Formula (IX), or a combination of both inhibitors on myeloid tumor-associated macrophages (TAMs) in pancreatic tumor-bearing mice.

[0072] FIG. 40 illustrates the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K- δ inhibitor of Formula (IX), or a combination of both inhibitors on myeloid-derived suppressor cells (MDSCs) in pancreatic tumor-bearing mice.

[0073] FIG. 41 illustrates the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K- δ inhibitor of Formula (IX), or a combination of both inhibitors on regulatory T cells (Tregs) in pancreatic tumor-bearing mice.

[0074] FIG. 42 illustrates the effects of vehicle on flux at two timepoints, as a control for comparison with FIG. 100, in the ID8 syngeneic orthotropic ovarian cancer model.

[0075] FIG. 43 illustrates the effects of the BTK inhibitor of Formula (XVIII) on flux at two timepoints, for comparison with FIG. 99, in the ID8 syngeneic orthotropic ovarian cancer model.

[0076] FIG. 44 illustrates tumor response to treatment with the BTK inhibitor of Formula (XVIII) correlates with a significant reduction in immunosuppressive tumor associated lymphocytes in tumor-bearing mice, in comparison to a control (vehicle).

[0077] FIG. 45 illustrates that treatment with the BTK inhibitor of Formula (XVIII) impairs ID8 ovarian cancer growth in the syngeneic murine model in comparison to a control (vehicle).

[0078] FIG. 46 illustrates that treatment with the BTK inhibitor of Formula (XVIII) induces a tumor response that correlates with a significant reduction in total B cells in tumor-bearing mice.

[0079] FIG. 47 illustrates that treatment with the BTK inhibitor of Formula (XVIII) induces a tumor response that correlates with a significant reduction in B regulatory cells (Bregs) in tumor-bearing mice.

[0080] FIG. 48 illustrates that treatment with the BTK inhibitor of Formula (XVIII) induces a tumor response that correlates with a significant reduction in immunosuppressive tumor

associated Tregs.

[0081] FIG. 49 illustrates that treatment with the BTK inhibitor of Formula (XVIII) induces a tumor response that correlates with an increase in CD8⁺ T cells.

[0082] FIG. 50 illustrates the effects on tumor volume of vehicle (measured in mm³) of the BTK inhibitor of Formula (XVIII), a combination of the BTK inhibitor of Formula (XVIII) and gemcitabine (“Gem”), and gemcitabine alone.

[0083] FIG. 51 illustrates the effects on the amount of CD8⁺ T cells, given as a percentage of cells expressing the T cell receptor (CD3), of the BTK inhibitor of Formula (XVIII), a combination of the BTK inhibitor of Formula (XVIII) and gemcitabine (“Gem”), and gemcitabine alone.

[0084] FIG. 52 illustrates the effects on the percentage of CD4⁺, CD25⁺, and FoxP3⁺ T regulatory cells (“Tregs”), given as a percentage of cells expressing the T cell receptor (CD3), of the BTK inhibitor of Formula (XVIII), a combination of the BTK inhibitor of Formula (XVIII) and gemcitabine (“Gem”), and gemcitabine alone.

[0085] FIG. 53 illustrates the effects on the percentage of CD11b⁺, LY6C^{low}, F4/80⁺, and Csf1r⁺ tumor-associated macrophages (“TAMs”), given as a percentage of cells expressing the T cell receptor (CD3), of the BTK inhibitor of Formula (XVIII), a combination of the BTK inhibitor of Formula (XVIII) and gemcitabine (“Gem”), and gemcitabine alone.

[0086] FIG. 54 illustrates the effects on the percentage of Gr1⁺ and LY6C^{hi}, F4/80⁺, and Csf1r⁺ myeloid-derived suppressor cells (“MDSCs”), given as a percentage of cells expressing the T cell receptor (CD3), of the BTK inhibitor of Formula (XVIII), a combination of the BTK inhibitor of Formula (XVIII) and gemcitabine (“Gem”), and gemcitabine alone.

[0087] FIG. 55 illustrates representative photomicrographs and comparison of maximal thrombus size in laser injured arterioles of VWF HA1 mutant mice infused with human platelets in the absence or presence of various BTK inhibitors. Representative photomicrographs are given as a comparison of maximal thrombus size in laser-injured arterioles (1 μM concentrations shown).

[0088] FIG. 56 illustrates a quantitative comparison obtained by *in vivo* analysis of early thrombus dynamics in a humanized mouse laser injury model using three BTK inhibitors at a

concentration 1 μM .

[0089] FIG. 57 illustrates the effect of the tested BTK inhibitors on thrombus formation. The conditions used were N=4, 3 mice per drug; anti-clotting agents < 2000 μM^2 . In studies with ibrutinib, 48% MCL bleeding events were observed with 560 mg QD and 63% CLL bleeding events were observed with 420 mg QD, where bleeding event is defined as subdural hematoma, ecchymoses, GI bleeding, or hematuria.

[0090] FIG. 58 illustrates the effect of the concentration of the tested BTK inhibitors on thrombus formation.

[0091] FIG. 59 illustrates the results of GPVI platelet aggregation studies of Formula XVIII ($\text{IC}_{50} = 1.15 \mu\text{M}$) and Formula XX-A (ibrutinib, $\text{IC}_{50} = 0.13 \mu\text{M}$).

[0092] FIG. 60 illustrates the results of GPVI platelet aggregation studies of Formula XVIII and Formula XX-A (ibrutinib).

DETAILED DESCRIPTION OF THE INVENTION

[0093] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs.

[0094] The terms “co-administration” and “administered in combination with” as used herein, encompass administration of two or more agents (such as at least one PI3K inhibitor and at least one BTK inhibitor) to a subject (such as a human or a mammal), so that both agents and/or their metabolites are present in the subject at the same time. Agents are also referred to as active ingredients, or active pharmaceutical ingredients, or drugs. Co-administration includes simultaneous administration in separate compositions (also referred to as concurrent administration), administration at different times in separate compositions, or administration in a composition in which both agents are present. Simultaneous administration in separate compositions and administration in a composition in which both agents are present are preferred.

[0095] The term “effective amount” or “therapeutically effective amount” refers to that amount of a compound or combination of compounds as described herein that is sufficient to effect the intended application including, but not limited to, disease treatment. A therapeutically effective amount may vary depending upon the intended application (*in vitro* or *in vivo*), or the subject and

disease condition being treated (*e.g.*, the weight, age and gender of the subject), the severity of the disease condition, the manner of administration, *etc.* which can readily be determined by one of ordinary skill in the art. The term also applies to a dose that will induce a particular response in target cells, (*e.g.*, the reduction of platelet adhesion and/or cell migration). The specific dose will vary depending on the particular compounds chosen, the dosing regimen to be followed, whether the compound is administered in combination with other compounds, timing of administration, the tissue to which it is administered, and the physical delivery system in which the compound is carried.

[0096] A “therapeutic effect” as that term is used herein, encompasses a therapeutic benefit and/or a prophylactic benefit. A prophylactic effect includes delaying or eliminating the appearance of a disease or condition, delaying or eliminating the onset of symptoms of a disease or condition, slowing, halting, or reversing the progression of a disease or condition, or any combination thereof.

[0097] The term “pharmaceutically acceptable salt” refers to salts derived from a variety of organic and inorganic counter ions known in the art. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and phosphoric acid. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid and salicylic acid. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese and aluminum. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins. Specific examples include isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine. In selected embodiments, the pharmaceutically acceptable base addition salt is chosen from ammonium, potassium, sodium, calcium, and magnesium salts.

[0098] “Pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and inert ingredients. The use of such pharmaceutically acceptable carriers or pharmaceutically acceptable excipients for active pharmaceutical ingredients is well known in the art. Except insofar as any conventional pharmaceutically acceptable carrier or pharmaceutically acceptable excipient is incompatible with the active pharmaceutical ingredient, its use in the therapeutic compositions of the invention is contemplated. Supplementary active pharmaceutical ingredients, such as other drugs, can also be incorporated into the described compositions and methods.

[0099] “Prodrug” is intended to describe a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound described herein. Thus, the term "prodrug" refers to a precursor of a biologically active compound that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject, but is converted *in vivo* to an active compound, for example, by hydrolysis. The prodrug compound often offers the advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see, *e.g.*, Bundgaard, H., Design of Prodrugs (1985) (Elsevier, Amsterdam). The term “prodrug” is also intended to include any covalently bonded carriers, which release the active compound *in vivo* when administered to a subject. Prodrugs of an active compound, as described herein, may be prepared by modifying functional groups present in the active compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to yield the active parent compound. Prodrugs include, for example, compounds wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the active compound is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetates, formates and benzoate derivatives of an alcohol, various ester derivatives of a carboxylic acid, or acetamide, formamide and benzamide derivatives of an amine functional group in the active compound.

[00100] The term “*in vivo*” refers to an event that takes place in a subject's body.

[00101] The term “*in vitro*” refers to an event that takes place outside of a subject's body. *In vitro* assays encompass cell-based assays in which cells alive or dead are employed and may also encompass a cell-free assay in which no intact cells are employed.

[00102] Unless otherwise stated, the chemical structures depicted herein are intended to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds where one or more hydrogen atoms is replaced by deuterium or tritium, or wherein one or more carbon atoms is replaced by ^{13}C - or ^{14}C -enriched carbons, are within the scope of this invention.

[00103] When ranges are used herein to describe, for example, physical or chemical properties such as weight or chemical formulae, all combinations and subcombinations of ranges and specific embodiments therein are intended to be included. Use of the term “about” when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error), and thus the number or numerical range may vary. The variation is typically from 0% to 10%, preferably from 0% to 10%, more preferably from 0% to 5% of the stated number or numerical range. The term “comprising” (and related terms such as “comprise” or “comprises” or “having” or “including”) encompasses those embodiments such as, for example, an embodiment of any composition of matter, method or process that “consist of” or “consist essentially of” the described features.

[00104] “Alkyl” refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to ten carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{10}$ alkyl and $(\text{C}_1\text{-C}_{10})$ alkyl). Whenever it appears herein, a numerical range such as “1 to 10” refers to each integer in the given range - *e.g.*, “1 to 10 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the definition is also intended to cover the occurrence of the term “alkyl” where no numerical range is specifically designated. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, iso-butyl, sec-butyl isobutyl, tertiary butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, nonyl and decyl. The alkyl moiety may be attached to the rest of the molecule by a single bond, such as for example, methyl (Me), ethyl (Et), *n*-propyl (Pr), 1-methylethyl (iso-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl

(*t*-butyl) and 3-methylhexyl. Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted by one or more of substituents which are independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where *t* is 1 or 2), -S(O)_tOR^a (where *t* is 1 or 2), -S(O)_tN(R^a)₂ (where *t* is 1 or 2), or PO₃(R^a)₂ where each R^a is independently hydrogen, unsubstituted alkyl, fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00105] “Alkylaryl” refers to an -(alkyl)aryl radical where aryl and alkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for aryl and alkyl respectively.

[00106] “Alkylhetaryl” refers to an -(alkyl)hetaryl radical where hetaryl and alkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for aryl and alkyl respectively.

[00107] “Alkylheterocycloalkyl” refers to an -(alkyl) heterocycyl radical where alkyl and heterocycloalkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for heterocycloalkyl and alkyl respectively.

[00108] An “alkene” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon double bond, and an “alkyne” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon triple bond. The alkyl moiety, whether saturated or unsaturated, may be branched, straight chain, or cyclic.

[00109] “Alkenyl” refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one double bond, and having from two to ten carbon atoms (*i.e.*, C₂-C₁₀ alkenyl and (C₂-C₁₀)alkenyl). Whenever it appears herein, a numerical range such as “2 to 10” refers to each integer in the given range - *e.g.*, “2 to 10 carbon atoms” means that the alkenyl group may consist of 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms. The alkenyl moiety may be attached to the rest of the molecule by a single bond, such as for example, ethenyl (*i.e.*, vinyl), prop-1-enyl (*i.e.*, allyl), but-1-enyl,

pent-1-enyl and penta-1,4-dienyl. Unless stated otherwise specifically in the specification, an alkenyl group is optionally substituted by one or more substituents which are independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00110] “Alkenyl-cycloalkyl” refers to an (alkenyl)cycloalkyl radical where alkenyl and cycloalkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for alkenyl and cycloalkyl respectively.

[00111] “Alkynyl” refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one triple bond, having from two to ten carbon atoms (*i.e.* C₂-C₁₀ alkynyl and (C₂-C₁₀)alkynyl). Whenever it appears herein, a numerical range such as “2 to 10” refers to each integer in the given range - *e.g.*, “2 to 10 carbon atoms” means that the alkynyl group may consist of 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms. The alkynyl may be attached to the rest of the molecule by a single bond, for example, ethynyl, propynyl, butynyl, pentynyl and hexynyl. Unless stated otherwise specifically in the specification, an alkynyl group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00112] “Alkynyl-cycloalkyl” refers to an -(alkynyl)cycloalkyl radical where alkynyl and cycloalkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for alkynyl and cycloalkyl respectively.

[00113] “Carboxaldehyde” refers to a -(C=O)H radical.

[00114] “Carboxyl” refers to a -(C=O)OH radical.

[00115] “Cyano” refers to a -CN radical.

[00116] “Cycloalkyl” refers to a monocyclic or polycyclic radical that contains only carbon and hydrogen, and may be saturated, or partially unsaturated. Cycloalkyl groups include groups having from 3 to 10 ring atoms (*i.e.* C₂-C₁₀ cycloalkyl and (C₂-C₁₀)cycloalkyl). Whenever it appears herein, a numerical range such as “3 to 10” refers to each integer in the given range - *e.g.*, “3 to 10 carbon atoms” means that the cycloalkyl group may consist of 3 carbon atoms and greater, up to and including 10 carbon atoms. Illustrative examples of cycloalkyl groups include, but are not limited to the following moieties: cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, norbornyl, and the like. Unless stated otherwise specifically in the specification, a cycloalkyl group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00117] “Cycloalkyl-alkenyl” refers to a -(cycloalkyl)alkenyl radical where cycloalkyl and alkenyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for cycloalkyl and alkenyl, respectively.

[00118] “Cycloalkyl-heterocycloalkyl” refers to a -(cycloalkyl)heterocycloalkyl radical where cycloalkyl and heterocycloalkyl are as disclosed herein and which are optionally substituted by

one or more of the substituents described as suitable substituents for cycloalkyl and heterocycloalkyl, respectively.

[00119] “Cycloalkyl-heteroaryl” refers to a -(cycloalkyl)heteroaryl radical where cycloalkyl and heteroaryl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for cycloalkyl and heteroaryl, respectively.

[00120] The term “alkoxy” refers to the group -O-alkyl, including from 1 to 8 carbon atoms of a straight, branched, cyclic configuration and combinations thereof attached to the parent structure through an oxygen. Examples include, but are not limited to, methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy and cyclohexyloxy. “Lower alkoxy” refers to alkoxy groups containing one to six carbons.

[00121] The term “substituted alkoxy” refers to alkoxy wherein the alkyl constituent is substituted (*i.e.*, -O-(substituted alkyl)). Unless stated otherwise specifically in the specification, the alkyl moiety of an alkoxy group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00122] The term “alkoxycarbonyl” refers to a group of the formula (alkoxy)(C=O)- attached through the carbonyl carbon wherein the alkoxy group has the indicated number of carbon atoms. Thus a C₁-C₆ alkoxycarbonyl group is an alkoxy group having from 1 to 6 carbon atoms attached through its oxygen to a carbonyl linker. “Lower alkoxycarbonyl” refers to an alkoxycarbonyl group wherein the alkoxy group is a lower alkoxy group.

[00123] The term “substituted alkoxycarbonyl” refers to the group (substituted alkyl)-O-C(O)- wherein the group is attached to the parent structure through the carbonyl functionality. Unless stated otherwise specifically in the specification, the alkyl moiety of an alkoxycarbonyl group is

optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-OC(O)N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)C(O)N(R^a)_2$, $N(R^a)C(NR^a)N(R^a)_2$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00124] “Acyl” refers to the groups (alkyl)-C(O)-, (aryl)-C(O)-, (heteroaryl)-C(O)-, (heteroalkyl)-C(O)- and (heterocycloalkyl)-C(O)-, wherein the group is attached to the parent structure through the carbonyl functionality. If the R radical is heteroaryl or heterocycloalkyl, the hetero ring or chain atoms contribute to the total number of chain or ring atoms. Unless stated otherwise specifically in the specification, the alkyl, aryl or heteroaryl moiety of the acyl group is optionally substituted by one or more substituents which are independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-OC(O)N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)C(O)N(R^a)_2$, $N(R^a)C(NR^a)N(R^a)_2$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00125] “Acyloxy” refers to a $R(C=O)O-$ radical wherein “R” is alkyl, aryl, heteroaryl, heteroalkyl or heterocycloalkyl, which are as described herein. If the R radical is heteroaryl or heterocycloalkyl, the hetero ring or chain atoms contribute to the total number of chain or ring atoms. Unless stated otherwise specifically in the specification, the “R” of an acyloxy group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-OC(O)N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)C(O)N(R^a)_2$, $N(R^a)C(NR^a)N(R^a)_2$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

$S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00126] “Amino” or “amine” refers to a $-N(R^a)_2$ radical group, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl, unless stated otherwise specifically in the specification. When a $-N(R^a)_2$ group has two R^a substituents other than hydrogen, they can be combined with the nitrogen atom to form a 4-, 5-, 6- or 7-membered ring. For example, $-N(R^a)_2$ is intended to include, but is not limited to, 1-pyrrolidinyl and 4-morpholinyl. Unless stated otherwise specifically in the specification, an amino group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, $-OR^a$, $-SR^a$, $-OC(O)R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-OC(O)N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)C(O)N(R^a)_2$, $N(R^a)C(NR^a)N(R^a)_2$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00127] The term “substituted amino” also refers to N-oxides of the groups $-NHR^d$, and NR^dR^d each as described above. N-oxides can be prepared by treatment of the corresponding amino group with, for example, hydrogen peroxide or m-chloroperoxybenzoic acid.

[00128] “Amide” or “amido” refers to a chemical moiety with formula $-C(O)N(R)_2$ or $-NHC(O)R$, where R is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), each of which moiety may itself be optionally substituted. The R_2 of $-N(R)_2$ of the amide may optionally be taken together with the nitrogen to which it is attached to form a 4-, 5-, 6- or 7-membered ring. Unless stated otherwise specifically in the specification, an amido group is optionally substituted independently by one or more of the substituents as described herein for alkyl, cycloalkyl, aryl, heteroaryl, or heterocycloalkyl. An amide may be an amino acid or a peptide molecule attached to a compound of Formula (I), thereby forming a prodrug. The

procedures and specific groups to make such amides, including the use of protecting groups, are known to those of skill in the art and can readily be found in seminal sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, N.Y., 1999.

[00129] “Aromatic” or “aryl” or “Ar” refers to an aromatic radical with six to ten ring atoms (*e.g.*, C₆-C₁₀ aromatic or C₆-C₁₀ aryl) which has at least one ring having a conjugated pi electron system which is carbocyclic (*e.g.*, phenyl, fluorenyl, and naphthyl). Bivalent radicals formed from substituted benzene derivatives and having the free valences at ring atoms are named as substituted phenylene radicals. Bivalent radicals derived from univalent polycyclic hydrocarbon radicals whose names end in “-yl” by removal of one hydrogen atom from the carbon atom with the free valence are named by adding “-idene” to the name of the corresponding univalent radical, *e.g.*, a naphthyl group with two points of attachment is termed naphthylidene. Whenever it appears herein, a numerical range such as “6 to 10” refers to each integer in the given range; *e.g.*, “6 to 10 ring atoms” means that the aryl group may consist of 6 ring atoms, 7 ring atoms, *etc.*, up to and including 10 ring atoms. The term includes monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of ring atoms) groups. Unless stated otherwise specifically in the specification, an aryl moiety is optionally substituted by one or more substituents which are independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00130] “Aralkyl” or “arylalkyl” refers to an (aryl)alkyl-radical where aryl and alkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for aryl and alkyl respectively.

[00131] “Ester” refers to a chemical radical of formula -COOR, where R is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and

heteroalicyclic (bonded through a ring carbon). The procedures and specific groups to make esters, including the use of protecting groups, are known to those of skill in the art and can readily be found in seminal sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, N.Y., 1999. Unless stated otherwise specifically in the specification, an ester group is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, trifluoromethyl, trifluoromethoxy, nitro, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00132] “Fluoroalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more fluoro radicals, as defined above, for example, trifluoromethyl, difluoromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, and the like. The alkyl part of the fluoroalkyl radical may be optionally substituted as defined above for an alkyl group.

[00133] “Halo,” “halide,” or, alternatively, “halogen” is intended to mean fluoro, chloro, bromo or iodo. The terms “haloalkyl,” “haloalkenyl,” “haloalkynyl,” and “haloalkoxy” include alkyl, alkenyl, alkynyl and alkoxy structures that are substituted with one or more halo groups or with combinations thereof. For example, the terms “fluoroalkyl” and “fluoroalkoxy” include haloalkyl and haloalkoxy groups, respectively, in which the halo is fluorine.

[00134] “Heteroalkyl,” “heteroalkenyl,” and “heteroalkynyl” refer to optionally substituted alkyl, alkenyl and alkynyl radicals and which have one or more skeletal chain atoms selected from an atom other than carbon, e.g., oxygen, nitrogen, sulfur, phosphorus or combinations thereof. A numerical range may be given - e.g., C₁-C₄ heteroalkyl which refers to the chain length in total, which in this example is 4 atoms long. A heteroalkyl group may be substituted with one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, nitro, oxo, thioxo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -

OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂,
 N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2),
 -S(O)_tN(R^a)₂ (where t is 1 or 2), or PO₃(R^a)₂, where each R^a is independently hydrogen, alkyl,
 fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl,
 heteroaryl or heteroarylalkyl.

[00135] “Heteroalkylaryl” refers to an -(heteroalkyl)aryl radical where heteroalkyl and aryl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for heteroalkyl and aryl, respectively.

[00136] “Heteroalkylheteroaryl” refers to an -(heteroalkyl)heteroaryl radical where heteroalkyl and heteroaryl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for heteroalkyl and heteroaryl, respectively.

[00137] “Heteroalkylheterocycloalkyl” refers to an -(heteroalkyl)heterocycloalkyl radical where heteroalkyl and heterocycloalkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for heteroalkyl and heterocycloalkyl, respectively.

[00138] “Heteroalkylcycloalkyl” refers to an -(heteroalkyl)cycloalkyl radical where heteroalkyl and cycloalkyl are as disclosed herein and which are optionally substituted by one or more of the substituents described as suitable substituents for heteroalkyl and cycloalkyl, respectively.

[00139] “Heteroaryl” or “heteroaromatic” or “HetAr” refers to a 5- to 18-membered aromatic radical (*e.g.*, C₅-C₁₃ heteroaryl) that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur, and which may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system. Whenever it appears herein, a numerical range such as “5 to 18” refers to each integer in the given range - *e.g.*, “5 to 18 ring atoms” means that the heteroaryl group may consist of 5 ring atoms, 6 ring atoms, etc., up to and including 18 ring atoms. Bivalent radicals derived from univalent heteroaryl radicals whose names end in “-yl” by removal of one hydrogen atom from the atom with the free valence are named by adding “-idene” to the name of the corresponding univalent radical - *e.g.*, a pyridyl group with two points of attachment is a pyridylidene. A N-containing “heteroaromatic” or “heteroaryl” moiety refers to an aromatic group in which at least one of the skeletal atoms of the ring is a nitrogen atom. The polycyclic

heteroaryl group may be fused or non-fused. The heteroatom(s) in the heteroaryl radical are optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heteroaryl may be attached to the rest of the molecule through any atom of the ring(s). Examples of heteroaryls include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzindolyl, 1,3-benzodioxolyl, benzofuranyl, benzoaxazolyl, benzo[*d*]thiazolyl, benzothiadiazolyl, benzo[*b*][1,4]dioxepinyl, benzo[*b*][1,4]oxazinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzoxazolyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzofurazanyl, benzothiazolyl, benzothienyl(benzothiophenyl), benzothieno[3,2-*d*]pyrimidinyl, benzotriazolyl, benzo[4,6]imidazo[1,2-*a*]pyridinyl, carbazolyl, cinnolinyl, cyclopenta[*d*]pyrimidinyl, 6,7-dihydro-5*H*-cyclopenta[4,5]thieno[2,3-*d*]pyrimidinyl, 5,6-dihydrobenzo[*h*]quinazolinyl, 5,6-dihydrobenzo[*h*]cinnolinyl, 6,7-dihydro-5*H*-benzo[6,7]cyclohepta[1,2-*c*]pyridazinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furazanyl, furanonyl, furo[3,2-*c*]pyridinyl, 5,6,7,8,9,10-hexahydrocycloocta[*d*]pyrimidinyl, 5,6,7,8,9,10-hexahydrocycloocta[*d*]pyridazinyl, 5,6,7,8,9,10-hexahydrocycloocta[*d*]pyridinyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, 5,8-methano-5,6,7,8-tetrahydroquinazolinyl, naphthyridinyl, 1,6-naphthyridinonyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 5,6,6a,7,8,9,10,10a-octahydrobenzo[*h*]quinazolinyl, 1-phenyl-1*H*-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyranyl, pyrrolyl, pyrazolyl, pyrazolo[3,4-*d*]pyrimidinyl, pyridinyl, pyrido[3,2-*d*]pyrimidinyl, pyrido[3,4-*d*]pyrimidinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, quinazolinyl, quinoxalinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, 5,6,7,8-tetrahydroquinazolinyl, 5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-*d*]pyrimidinyl, 6,7,8,9-tetrahydro-5*H*-cyclohepta[4,5]thieno[2,3-*d*]pyrimidinyl, 5,6,7,8-tetrahydropyrido[4,5-*c*]pyridazinyl, thiazolyl, thiadiazolyl, thiapyranyl, triazolyl, tetrazolyl, triazinyl, thieno[2,3-*d*]pyrimidinyl, thieno[3,2-*d*]pyrimidinyl, thieno[2,3-*c*]pyridinyl, and thiophenyl (*i.e.* thienyl). Unless stated otherwise specifically in the specification, a heteroaryl moiety is optionally substituted by one or more substituents which are independently: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, nitro, oxo, thioxo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -OC(O)N(R^a)₂, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)C(O)N(R^a)₂, N(R^a)C(NR^a)N(R^a)₂, -N(R^a)S(O)_tR^a (where *t* is 1 or 2), -S(O)_tOR^a (where *t* is

1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00140] Substituted heteroaryl also includes ring systems substituted with one or more oxide (-O-) substituents, such as, for example, pyridinyl *N*-oxides.

[00141] "Heteroarylalkyl" refers to a moiety having an aryl moiety, as described herein, connected to an alkylene moiety, as described herein, wherein the connection to the remainder of the molecule is through the alkylene group.

[00142] "Heterocycloalkyl" refers to a stable 3- to 18-membered non-aromatic ring radical that comprises two to twelve carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. Whenever it appears herein, a numerical range such as "3 to 18" refers to each integer in the given range - *e.g.*, "3 to 18 ring atoms" means that the heterocycloalkyl group may consist of 3 ring atoms, 4 ring atoms, *etc.*, up to and including 18 ring atoms. Unless stated otherwise specifically in the specification, the heterocycloalkyl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems. The heteroatoms in the heterocycloalkyl radical may be optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heterocycloalkyl radical is partially or fully saturated. The heterocycloalkyl may be attached to the rest of the molecule through any atom of the ring(s). Examples of such heterocycloalkyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazoliny, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranlyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless stated otherwise specifically in the specification, a heterocycloalkyl moiety is optionally substituted by one or more substituents which independently are: alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, hydroxy, halo, cyano, nitro, oxo, thioxo, trimethylsilanyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-OC(O)N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)C(O)N(R^a)_2$, $N(R^a)C(NR^a)N(R^a)_2$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is

1 or 2), $-S(O)_tN(R^a)_2$ (where t is 1 or 2), or $PO_3(R^a)_2$, where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl or heteroarylalkyl.

[00143] “Heterocycloalkyl” also includes bicyclic ring systems wherein one non-aromatic ring, usually with 3 to 7 ring atoms, contains at least 2 carbon atoms in addition to 1-3 heteroatoms independently selected from oxygen, sulfur, and nitrogen, as well as combinations comprising at least one of the foregoing heteroatoms; and the other ring, usually with 3 to 7 ring atoms, optionally contains 1-3 heteroatoms independently selected from oxygen, sulfur, and nitrogen and is not aromatic.

[00144] “Isomers” are different compounds that have the same molecular formula.

“Stereoisomers” are isomers that differ only in the way the atoms are arranged in space - *i.e.*, having a different stereochemical configuration. “Enantiomers” are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a “racemic” mixture. The term “(±)” is used to designate a racemic mixture where appropriate.

“Diastereoisomers” are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other. The absolute stereochemistry is specified according to the Cahn-Ingold-Prelog R-S system. When a compound is a pure enantiomer the stereochemistry at each chiral carbon can be specified by either *R* or *S*. Resolved compounds whose absolute configuration is unknown can be designated (+) or (-) depending on the direction (dextro- or levorotatory) which they rotate plane polarized light at the wavelength of the sodium D line. Certain of the compounds described herein contain one or more asymmetric centers and can thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that can be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)-. The present chemical entities, pharmaceutical compositions and methods are meant to include all such possible isomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (*R*)- and (*S*)-isomers can be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers.

[00145] “Enantiomeric purity” as used herein refers to the relative amounts, expressed as a percentage, of the presence of a specific enantiomer relative to the other enantiomer. For example, if a compound, which may potentially have an (*R*)- or an (*S*)-isomeric configuration, is present as a racemic mixture, the enantiomeric purity is about 50% with respect to either the (*R*)- or (*S*)-isomer. If that compound has one isomeric form predominant over the other, for example, 80% (*S*)- and 20% (*R*)-, the enantiomeric purity of the compound with respect to the (*S*)-isomeric form is 80%. The enantiomeric purity of a compound can be determined in a number of ways known in the art, including but not limited to chromatography using a chiral support, polarimetric measurement of the rotation of polarized light, nuclear magnetic resonance spectroscopy using chiral shift reagents which include but are not limited to lanthanide containing chiral complexes or the Pirkle alcohol, or derivatization of a compounds using a chiral compound such as Mosher's acid followed by chromatography or nuclear magnetic resonance spectroscopy.

[00146] “Moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[00147] “Nitro” refers to the -NO₂ radical.

[00148] “Oxa” refers to the -O- radical.

[00149] “Oxo” refers to the =O radical.

[00150] “Tautomers” are structurally distinct isomers that interconvert by tautomerization. “Tautomerization” is a form of isomerization and includes prototropic or proton-shift tautomerization, which is considered a subset of acid-base chemistry. “Prototropic tautomerization” or “proton-shift tautomerization” involves the migration of a proton accompanied by changes in bond order, often the interchange of a single bond with an adjacent double bond. Where tautomerization is possible (e.g. in solution), a chemical equilibrium of tautomers can be reached. An example of tautomerization is keto-enol tautomerization. A specific example of keto-enol tautomerization is the interconversion of pentane-2,4-dione and 4-hydroxypent-3-en-2-one tautomers. Another example of tautomerization is phenol-keto tautomerization. A specific example of phenol-keto tautomerization is the interconversion of pyridin-4-ol and pyridin-4(1*H*)-one tautomers.

[00151] The terms “enantiomerically enriched,” “enantiomerically pure” and “non-racemic,” as used herein, refer to compositions in which the percent by weight of one enantiomer is greater than the amount of that one enantiomer in a control mixture of the racemic composition (e.g., greater than 1:1 by weight). For example, an enantiomerically enriched preparation of the (S)-enantiomer, means a preparation of the compound having greater than 50% by weight of the (S)-enantiomer relative to the (R)-enantiomer, such as at least 75% by weight, such as at least 80% by weight. In some embodiments, the enrichment can be significantly greater than 80% by weight, providing a “substantially enantiomerically enriched,” “substantially enantiomerically pure” or a “substantially non-racemic” preparation, which refers to preparations of compositions which have at least 85% by weight of one enantiomer relative to the other enantiomer, such as at least 90% by weight, and such as at least 95% by weight. The terms “diastereomerically enriched” and “diastereomerically pure,” as used herein, refer to compositions in which the percent by weight of one diastereomer is greater than the amount of that one diastereomer in a control mixture of diastereomers. In some embodiments, the enrichment can be significantly greater than 80% by weight, providing a “substantially diastereomerically enriched” or “substantially diastereomerically pure” preparation, which refers to preparations of compositions which have at least 85% by weight of one diastereomer relative to other diastereomers, such as at least 90% by weight, and such as at least 95% by weight.

[00152] In preferred embodiments, the enantiomerically enriched composition has a higher potency with respect to therapeutic utility per unit mass than does the racemic mixture of that composition. Enantiomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred enantiomers can be prepared by asymmetric syntheses. See, for example, Jacques, *et al.*, *Enantiomers, Racemates and Resolutions*, Wiley Interscience, New York, 1981; and Eliel, *Stereochemistry of Carbon Compounds*, McGraw-Hill, NY, 1962.

[00153] A “leaving group or atom” is any group or atom that will, under selected reaction conditions, cleave from the starting material, thus promoting reaction at a specified site. Examples of such groups, unless otherwise specified, include halogen atoms and mesyloxy, p-nitrobenzensulphonyloxy and tosyloxy groups.

[00154] “Protecting group” is intended to mean a group that selectively blocks one or more reactive sites in a multifunctional compound such that a chemical reaction can be carried out selectively on another unprotected reactive site and the group can then be readily removed after the selective reaction is complete. A variety of protecting groups are disclosed, for example, in T. H. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Third Edition, John Wiley & Sons, New York, 1999.

[00155] “Solvate” refers to a compound in physical association with one or more molecules of a pharmaceutically acceptable solvent.

[00156] “Substituted” means that the referenced group may have attached one or more additional moieties individually and independently selected from, for example, acyl, alkyl, alkylaryl, cycloalkyl, aralkyl, aryl, carbohydrate, carbonate, heteroaryl, heterocycloalkyl, hydroxy, alkoxy, aryloxy, mercapto, alkylthio, arylthio, cyano, halo, carbonyl, ester, thiocarbonyl, isocyanato, thiocyanato, isothiocyanato, nitro, oxo, perhaloalkyl, perfluoroalkyl, phosphate, silyl, sulfinyl, sulfonyl, sulfonamidyl, sulfoxyl, sulfonate, urea, and amino, including mono- and di-substituted amino groups, and protected derivatives thereof. The substituents themselves may be substituted, for example, a cycloalkyl substituent may itself have a halide substituent at one or more of its ring carbons. The term “substituted” also means that one or more hydrogens on the designated atom/atoms is/are replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible if such combinations result in stable compounds. “Stable compound” or “stable structure” is defined as a compound or structure that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. The terms “optionally substituted” and “may optionally be substituted” means optional substitution with the specified groups, radicals or moieties.

[001] “Sulfanyl” refers to groups that include -S-(optionally substituted alkyl), -S-(optionally substituted aryl), -S-(optionally substituted heteroaryl) and -S-(optionally substituted heterocycloalkyl).

[002] “Sulfinyl” refers to groups that include -S(O)-H, -S(O)-(optionally substituted alkyl), -S(O)-(optionally substituted amino), -S(O)-(optionally substituted aryl), -S(O)-(optionally substituted heteroaryl) and -S(O)-(optionally substituted heterocycloalkyl).

[003] “Sulfonyl” refers to groups that include -S(O₂)-H, -S(O₂)-(optionally substituted alkyl), -S(O₂)-(optionally substituted amino), -S(O₂)-(optionally substituted aryl), -S(O₂)-(optionally substituted heteroaryl), and -S(O₂)-(optionally substituted heterocycloalkyl).

[004] “Sulfonamidyl” or “sulfonamido” refers to a -S(=O)₂-NRR radical, where each R is selected independently from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). The R groups in -NRR of the -S(=O)₂-NRR radical may be taken together with the nitrogen to which it is attached to form a 4-, 5-, 6- or 7-membered ring. A sulfonamido group is optionally substituted by one or more of the substituents described for alkyl, cycloalkyl, aryl, heteroaryl, respectively.

[005] “Sulfoxyl” refers to a -S(=O)₂OH radical.

[006] “Sulfonate” refers to a -S(=O)₂-OR radical, where R is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). A sulfonate group is optionally substituted on R by one or more of the substituents described for alkyl, cycloalkyl, aryl, heteroaryl, respectively.

[007] Compounds of the invention also include crystalline and amorphous forms of those compounds, including, for example, polymorphs, pseudopolymorphs, solvates, hydrates, unsolvated polymorphs (including anhydrides), conformational polymorphs, and amorphous forms of the compounds, as well as mixtures thereof. “Crystalline form” and “polymorph” are intended to include all crystalline and amorphous forms of the compound, including, for example, polymorphs, pseudopolymorphs, solvates, hydrates, unsolvated polymorphs (including anhydrides), conformational polymorphs, and amorphous forms, as well as mixtures thereof, unless a particular crystalline or amorphous form is referred to.

Co-administration of compounds

[008] An embodiment of the invention is a composition, such as a pharmaceutical composition, comprising a combination of a PI3K inhibitor and a BTK inhibitor. Another embodiment is a kit containing both components formulated into separate pharmaceutical compositions, which are formulated for co-administration.

[009] Another embodiment of the invention is a method of treating a disease or condition in a subject, in particular a hyperproliferative disorder like leukemia, lymphoma or a solid tumor cancer in a subject, comprising co-administering to the subject in need thereof a therapeutically effective amount of a combination of a PI3K inhibitor and a BTK inhibitor. The pharmaceutical composition comprising the combination, and the kit, are both for use in treating such disease or condition.

[0010] In an exemplary embodiment, the solid tumor cancer is selected from the group consisting of breast, lung, colorectal, thyroid, bone sarcoma and stomach cancers.

[0011] In an exemplary embodiment, the leukemia is selected from the group consisting of acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), and acute lymphoblastic leukemia (ALL).

[0012] In a preferred embodiment, the PI3K inhibitor is a PI3K- γ inhibitor.

[0013] In another preferred embodiment, the PI3K inhibitor is a PI3K- δ inhibitor.

[0014] In another preferred embodiment, the PI3K inhibitor is a PI3K- γ,δ inhibitor.

[0015] In a particularly preferred embodiment, the PI3K inhibitor is a PI3K- δ inhibitor. This PI3K- δ inhibitor is more preferably a compound of Formula VIII, even more preferably the compound of Formula IX.

[0016] The BTK inhibitor is preferably a compound of Formula XVII, even more preferably the compound of Formula XVIII.

[0017] In one specific embodiment, the PI3K inhibitor is a PI3K- δ inhibitor and the BTK inhibitor is a compound of Formula XVII, even more preferably the compound of Formula

XVIII. In a specifically preferred embodiment, the PI3K inhibitor is the compound of Formula IX and the BTK inhibitor is the compound of Formula XVIII. One or both of said inhibitors may also be in the form of a pharmaceutically acceptable salt.

[0018] In an exemplary embodiment, the PI3K inhibitor is a PI3K inhibitor selective for δ -PI3K, γ -PI3K, or γ,δ -PI3K isoforms .

[0019] The combination may be administered by any route known in the art. In an exemplary embodiment, the combination of the PI3K inhibitor, which is preferably selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, with the BTK inhibitor is administered by oral, intravenous, intramuscular, intraperitoneal, subcutaneous or transdermal means. In one embodiment, the administration is by injection.

[0020] In an exemplary embodiment, the PI3K inhibitor, which is preferably selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, is in the form of a pharmaceutically acceptable salt.

[0021] In an exemplary embodiment, the BTK inhibitor is in the form of a pharmaceutically acceptable salt.

[0022] In an exemplary embodiment, the PI3K inhibitor, which is preferably selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, is administered to the subject before administration of the BTK inhibitor.

[0023] In an exemplary embodiment, the PI3K inhibitor, which is preferably selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, is administered concurrently with the administration of the BTK inhibitor.


[0024] In an exemplary embodiment, the PI3K inhibitor, which is preferably selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, is administered to the subject after administration of the BTK inhibitor.

[0025] In an exemplary embodiment, the subject is a mammal, such as a human.

PI3K Inhibitors

[0026] In particular, it is one of the PI3K inhibitors described in more detail in the following paragraphs. Preferably, it is a PI3K inhibitor selected from the group consisting of PI3K- γ inhibitor, PI3K- δ inhibitor, and PI3K- γ,δ inhibitor. In one specific embodiment, it is a PI3K- δ inhibitor. In a preferred embodiment, it is a compound of Formula IX or a pharmaceutically acceptable salt thereof.

[0027] In an exemplary embodiment, the PI3K inhibitor, which may preferably be selected from the group consisting of PI3K- γ inhibitor, PI3K- δ inhibitor, and PI3K- γ,δ inhibitor, is a compound selected from the structures disclosed in U.S. Patent Nos. 8,193,182 and 8,569,323, and U.S. Patent Application Publication Nos. 2012/0184568 A1, 2013/0344061 A1, and 2013/0267521 A1. In an exemplary embodiment, the PI3K inhibitor is a compound of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt thereof,

wherein:

Cy is aryl or heteroaryl substituted by 0 or 1 occurrences of R³ and 0, 1, 2, or 3 occurrences of R⁵;

W_b⁵ is CR⁸, CHR⁸, or N;

R⁸ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfonamido, halo, cyano, hydroxyl or nitro;

B is hydrogen, alkyl, amino, heteroalkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl, each of which is substituted with 0, 1, 2, 3, or 4 occurrences of R²;

each R^2 is independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkoxy, amido, amino, acyl, acyloxy, alkoxycarbonyl, sulfonamido, halo, cyano, hydroxyl, nitro, phosphate, urea, or carbonate;

X is $-(CH(R^9))_z-$;

Y is $-N(R^9)-C(=O)-$, $-C(=O)-N(R^9)-$, $-C(=O)-N(R^9)-(CHR^9)-$, $-N(R^9)-S(=O)-$, $-S(=O)-N(R^9)-$, $S(=O)_2-N(R^9)-$, $-N(R^9)-C(=O)-N(R^9)$ or $-N(R^9)S(=O)_2-$;

z is an integer of 1, 2, 3, or 4;

R^3 is alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, fluoroalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfinyl, sulfonyl, sulfoxide, sulfone, sulfonamido, halo, cyano, aryl, heteroaryl, hydroxyl, or nitro;


each R^5 is independently alkyl, alkenyl, alkynyl, cycloalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfonamido, halo, cyano, hydroxyl, or nitro;

each R^9 is independently hydrogen, alkyl, cycloalkyl, heterocyclyl, or heteroalkyl; or two adjacent occurrences of R^9 together with the atoms to which they are attached form a 4- to 7-membered ring;

W_d is heterocyclyl, aryl, cycloalkyl, or heteroaryl, each of which is substituted with one or more R^{10} , R^{11} , R^{12} or R^{13} , and

R^{10} , R^{11} , R^{12} and R^{13} are each independently hydrogen, alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkoxy, heterocycloxy, amido, amino, acyl, acyloxy, alkoxycarbonyl, sulfonamido, halo, cyano, hydroxyl, nitro, phosphate, urea, carbonate or $NR'R''$ wherein R' and R'' are taken together with nitrogen to form a cyclic moiety.

[0028] In an exemplary embodiment, the PI3K inhibitor, PI3K- γ inhibitor, PI3K- δ inhibitor, or PI3K- γ,δ inhibitor is a compound of Formula (I-1):




Formula (I-1)

or a pharmaceutically acceptable salt thereof,

wherein:

B is a moiety of Formula (II):



Formula (II) ;

W_c is aryl, heteroaryl, heterocycloalkyl, or cycloalkyl;


q is an integer of 0, 1, 2, 3, or 4;


X is a bond or $-(CH(R^9))_z-$, and z is an integer of 1, 2, 3 or 4;

Y is a bond, $-N(R^9)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-C(=O)-$, $-C(=O)(CHR^9)_z-$, $-N(R^9)-C(=O)-$,
 $-N(R^9)-C(=O)NH-$ or $-N(R^9)C(R^9)_2-$;

z is an integer of 1, 2, 3, or 4;

W_d is:






X_1 , X_2 and X_3 are each independently C, CR^{13} or N; and X_4 , X_5 and X_6 are each independently N, NH, CR^{13} , S or O;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, alkoxy, amido, alkoxy carbonyl, sulfonamido, halo, cyano, or nitro;


R^2 is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, heteroarylalkyl, alkoxy, amino, halo, cyano, hydroxy or nitro; $R_{sup.3}$ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, alkoxy, amido, amino, alkoxycarbonyl sulfonamido, halo, cyano, hydroxy or nitro; and each instance of R^9 is independently hydrogen, alkyl, or heterocycloalkyl.

[0029] In an exemplary embodiment, the PI3K inhibitor is a compound of Formula (III), also known as (*S*)-3-(1-((9*H*-purin-6-yl)amino)ethyl)-8-chloro-2-phenylisoquinolin-1(2*H*)-one,



or a pharmaceutically acceptable salt thereof.


[0030] In an exemplary embodiment, the PI3K inhibitor is a compound of Formula (IV), also known as (*S*)-3-amino-*N*-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazolin-2-yl)ethyl)pyrazine-2-carboxamide,



Formula (IV)

or a pharmaceutically acceptable salt thereof.

[0031] In an exemplary embodiment, the PI3K inhibitor is a compound selected from the structures disclosed in U.S. Patent Nos. 8,193,199 and 8,586,739. In an exemplary embodiment, the PI3K inhibitor is a compound of Formula (V):



Formula (V)

or a pharmaceutically-acceptable salt thereof, wherein:

X¹ is C(R⁹) or N;

X² is C(R₁₀) or N;

Y is N(R¹¹), O or S;

Z is CR⁸ or N;

n is 0, 1, 2 or 3;

R¹ is a direct-bonded or oxygen -linked saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R² substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl;

R² is selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, OS(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a; or R² is selected from C₁₋₆alkyl, phenyl, benzyl, heteroaryl, heterocycle, —(C₁₋₃alkyl)heteroaryl, —(C₁₋₃alkyl)heterocycle, —O(C₁₋₃alkyl)heteroaryl, —O(C₁₋₃alkyl)heterocycle, —NR^a(C₁₋₃alkyl)heteroaryl, —NR^a(C₁₋₃alkyl)heterocycle, —(C₁₋₃alkyl)phenyl, —O(C₁₋₃alkyl)phenyl and —NR^a(C₁₋₃alkyl)phenyl all of which are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₄haloalkyl, OC₁₋₄alkyl, Br, Cl, F, I and C₁₋₄alkyl;

R³ is selected from H, halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aNR^aR^a, —NR^aC₂₋₆alkylOR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₆haloalkyl, OC₁₋₆alkyl, Br, Cl, F, I and C₁₋₆alkyl;

R⁴ is, independently, in each instance, halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl or C₁₋₄haloalkyl;

R⁵ is, independently, in each instance, H, halo, C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC₁₋₄alkyl, C₁₋₄alkyl, C₁₋₃haloalkyl,

OC₁₋₄alkyl, NH₂, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl; or both R⁵ groups together form a C₃₋₆spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, OC₁₋₄alkyl, C₁₋₄alkyl, C₁₋₃haloalkyl, OC₁₋₄alkyl, NH₂, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl;

R⁶ is selected from H, halo, C₁₋₆alkyl, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a;

R⁷ is selected from H, halo, C₁₋₆alkyl, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a;

R⁸ is selected from H, C₁₋₆haloalkyl, Br, Cl, F, I, OR^a, NR^aR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₆haloalkyl, OC₁₋₆alkyl, Br, Cl, F, I and C₁₋₆alkyl;

R⁹ is selected from H, halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^aC(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(O)NR^aR^aN(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylOR^a, —NR^aC₂₋₆alkylOR^a; or R⁹ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1,

2, 3 or 4 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a;

R¹⁰ is H, C₁₋₃alkyl, C₁₋₃haloalkyl, cyano, nitro, CO₂R^a, C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —S(=O)R^b, S(=O)₂R^b or S(=O)₂NR^aR^a;

R¹¹ is H or C₁₋₄alkyl;

R^a is independently, at each instance, H or R^b; and

R^b is independently, at each instance, phenyl, benzyl or C₁₋₆alkyl, the phenyl, benzyl and C₁₋₆alkyl being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alkyl, C₁₋₃haloalkyl, —OC₁₋₄alkyl, —NH₂, —NHC₁₋₄alkyl, —N(C₁₋₄alkyl)C₁₋₄alkyl.

[0032] In a preferred embodiment, X¹ is C(R⁹). In a further preferred embodiment, X¹ is C(R⁹) and X² is N. In a further embodiment, X¹ is C(R⁹) and X² is C(R¹⁰).

[0033] In a preferred embodiment, R¹ is phenyl substituted by 0 or 1 R² substituents, and the phenyl is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl. In one specific embodiment, R¹ is unsubstituted phenyl. In a further specific embodiment, R¹ is phenyl substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl.

[0034] In one specific embodiment, R¹ is selected from 2-methylphenyl, 2-chlorophenyl, 2-trifluoromethylphenyl, 2-fluorophenyl and 2-methoxyphenyl.

[0035] In a further preferred embodiment, R¹ is phenoxy.

[0036] In a further preferred embodiment, R¹ is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 1, 2, 3 or 4

atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R² substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl.

[0037] In one specific embodiment, R¹ is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 0 or 1 R² substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl.

[0038] In a further specific embodiment, R¹ is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl.

[0039] In one specific embodiment, R¹ is an unsubstituted unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S.

[0040] In a further specific embodiment, R¹ is selected from pyridyl and pyrimidinyl.

[0041] In one embodiment, R³ is selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₆haloalkyl, OC₁₋₆alkyl, Br, Cl, F, I and C₁₋₆alkyl.

[0042] In one preferred embodiment, R³ is H.

[0043] In another preferred embodiment, R^3 is selected from F, Cl, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, OC_{1-6} alkyl, Br, Cl, F, I and C_{1-6} alkyl.

[0044] In one embodiment, R^5 is, independently, in each instance, H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a C_{3-6} spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[0045] In one preferred embodiment, R^5 is H.

[0046] In another preferred embodiment, one R^5 is S-methyl, the other is H.

[0047] In another preferred embodiment, at least one R^5 is halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[0048] In a preferred embodiment, R^6 is H.

[0049] In another preferred embodiment, R^6 is F, Cl, cyano or nitro.

[0050] In a preferred embodiment, R^7 is H.

[0051] In another preferred embodiment, R^7 is F, Cl, cyano or nitro.

[0052] In a preferred embodiment, R^8 is selected from H, CF_3 , C_{1-3} alkyl, Br, Cl and F.

[0053] In one specific embodiment, R^8 is H.

[0054] In another specific embodiment, R^8 is selected from CF_3 , C_{1-3} alkyl, Br, Cl and F.

[0055] In a preferred embodiment, R^9 is H.

[0056] In another embodiment, R^9 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, — $C(=O)R^a$, — $C(=O)OR^a$, — $C(=O)NR^aR^a$, — $C(=NR^a)NR^aR^a$, — OR^a , — $OC(=O)R^a$, —

OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₂₋₆alkylOR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₂₋₆alkylOR^a.


[0057] In one embodiment, R⁹ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a.

[0058] In one preferred embodiment, R¹⁰ is H.

[0059] In another preferred embodiment, R¹⁰ is cyano, nitro, CO₂R^a, C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, S(=O)R^b, S(=O)₂R^b or S(=O)₂NR^aR^a.

[0060] In one preferred embodiment, R¹¹ is H.

[0061] In another exemplary embodiment, the PI3K inhibitor is a compound of Formula (VI):



or a pharmaceutically-acceptable salt thereof, wherein:

X^1 is $C(R^9)$ or N;

X^2 is $C(R^{10})$ or N;

Y is $N(R^{11})$, O or S;

Z is CR^8 or N;

R^1 is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl;

R^2 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}$ alkyl NR^aR^a , $-OC_{2-6}$ alkyl OR^a , $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alkyl NR^aR^a and $-NR^aC_{2-6}$ alkyl OR^a ; or R^2 is selected from C_{1-6} alkyl, phenyl, benzyl, heteroaryl, heterocycle, $-(C_{1-3}$ alkyl)heteroaryl, $-(C_{1-3}$ alkyl)heterocycle, $-O(C_{1-3}$ alkyl)heteroaryl, $-O(C_{1-3}$ alkyl)heterocycle, $-NR^a(C_{1-3}$ alkyl)heteroaryl, $-NR^a(C_{1-3}$ alkyl)heterocycle, $-(C_{1-3}$ alkyl)phenyl, $-O(C_{1-3}$ alkyl)phenyl

and $\text{—NR}^a(\text{C}_{1-3}\text{alkyl})\text{phenyl}$ all of which are substituted by 0, 1, 2 or 3 substituents selected from $\text{C}_{1-4}\text{haloalkyl}$, $\text{OC}_{1-4}\text{alkyl}$, Br, Cl, F, I and $\text{C}_{1-4}\text{alkyl}$;

R^3 is selected from H, halo, $\text{C}_{1-4}\text{haloalkyl}$, cyano, nitro, —C(=O)R^a , —C(=O)OR^a , $\text{C(=O)NR}^a\text{R}^a$, $\text{C(=NR}^a\text{)NR}^a\text{R}^a$, —OR^a , —OC(=O)R^a , $\text{—OC(=O)NR}^a\text{R}^a$, $\text{—OC(=O)N(R}^a\text{)S(=O)}_2\text{R}^a$, $\text{—OC}_{2-6}\text{alkylNR}^a\text{R}^a$, $\text{—OC}_{2-6}\text{alkylOR}^a$, —SR^a , —S(=O)R^a , $\text{—S(=O)}_2\text{R}^a$, $\text{—S(=O)}_2\text{NR}^a\text{R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)OR}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)NR}^a\text{R}^a$, NR^aR^a , $\text{—N(R}^a\text{)C(=O)R}^a$, $\text{—N(R}^a\text{)C(=O)OR}^a$, $\text{—N(R}^a\text{)C(=O)NR}^a\text{R}^a$, $\text{—N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{—N(R}^a\text{)S(=O)}_2\text{R}^a$, $\text{—N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{—NR}^a\text{C}_{2-6}\text{alkylOR}^a$, $\text{C}_{1-6}\text{alkyl}$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $\text{C}_{1-6}\text{alkyl}$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $\text{C}_{1-6}\text{haloalkyl}$, $\text{OC}_{1-6}\text{alkyl}$, Br, Cl, F, I and $\text{C}_{1-6}\text{alkyl}$;

R^5 is, independently, in each instance, H, halo, $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$, or $\text{C}_{1-6}\text{alkyl}$ substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, $\text{OC}_{1-4}\text{alkyl}$, $\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-3}\text{haloalkyl}$, $\text{OC}_{1-4}\text{alkyl}$, NH_2 , $\text{NHC}_{1-4}\text{alkyl}$, $\text{N(C}_{1-4}\text{alkyl)C}_{1-4}\text{alkyl}$; or both R^5 groups together form a C_{3-6} -spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, $\text{OC}_{1-4}\text{alkyl}$, $\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-3}\text{haloalkyl}$, $\text{OC}_{1-4}\text{alkyl}$, NH_2 , $\text{NHC}_{1-4}\text{alkyl}$, $\text{N(C}_{1-4}\text{alkyl)C}_{1-4}\text{alkyl}$;

R^6 is selected from H, halo, $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$, cyano, nitro, —C(=O)R^a , —C(=O)OR^a , $\text{—C(=O)NR}^a\text{R}^a$, $\text{—C(=NR}^a\text{)NR}^a\text{R}^a$, —S(=O)R^a , $\text{—S(=O)}_2\text{R}^a$, $\text{—S(=O)}_2\text{NR}^a\text{R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)OR}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)NR}^a\text{R}^a$;

R^7 is selected from H, halo, $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$, cyano, nitro, —C(=O)R^a , —C(=O)OR^a , $\text{—C(=O)NR}^a\text{R}^a$, $\text{—C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{—S(=O)R}^a\text{S(=O)}_2\text{R}^a$, $\text{—S(=O)}_2\text{NR}^a\text{R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)OR}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)NR}^a\text{R}^a$;

R^8 is selected from H, $\text{C}_{1-6}\text{haloalkyl}$, Br, Cl, F, I, OR^a , NR^aR^a , $\text{C}_{1-6}\text{alkyl}$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $\text{C}_{1-6}\text{alkyl}$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $\text{C}_{1-6}\text{haloalkyl}$, $\text{OC}_{1-6}\text{alkyl}$, Br, Cl, F, I and $\text{C}_{1-6}\text{alkyl}$;

R^9 is selected from H, halo, $\text{C}_{1-4}\text{haloalkyl}$, cyano, nitro, —C(=O)R^a , —C(=O)OR^a , $\text{—C(=O)NR}^a\text{R}^a$, $\text{—C(=NR}^a\text{)NR}^a\text{R}^a$, —OR^a , —OC(=O)R^a , $\text{OC(=O)NR}^a\text{R}^a$, $\text{—OC(=O)N(R}^a\text{)S(=O)}_2\text{R}^a$, $\text{—OC}_{2-6}\text{alkylNR}^a\text{R}^a$, $\text{—OC}_{2-6}\text{alkylOR}^a$, —SR^a , —S(=O)R^a , $\text{—S(=O)}_2\text{R}^a$, $\text{—S(=O)}_2\text{NR}^a\text{R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)R}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)OR}^a$, $\text{—S(=O)}_2\text{N(R}^a\text{)C(=O)NR}^a\text{R}^a$, $\text{—NR}^a\text{R}^a$, $\text{—N(R}^a\text{)C(=O)R}^a$, $\text{—N(R}^a\text{)C(=O)OR}^a$, —

$N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^a$, $-NR^aC_{2-6}alkylOR^a$; or R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$;

R^{10} is H, $C_{1-3}alkyl$, $C_{1-3}haloalkyl$, cyano, nitro, CO_2R^a , $C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-S(=O)R^b$, $S(=O)_2R^b$ or $S(=O)_2NR^aR^a$; $-R^{11}$ is H or $C_{1-4}alkyl$;

R^a is independently, at each instance, H or R^b ; and

R^b is independently, at each instance, phenyl, benzyl or $C_{1-6}alkyl$, the phenyl, benzyl and $C_{1-6}alkyl$ being substituted by 0, 1, 2 or 3 substituents selected from halo, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $-OC_{1-4}alkyl$, $-NH_2$, $-NHC_{1-4}alkyl$, $-N(C_{1-4}alkyl)C_{1-4}alkyl$.

[0062] In a preferred embodiment, X^1 is $C(R^9)$. In a further preferred embodiment, X^1 is $C(R^9)$ and X^2 is N. In a further embodiment, X^1 is $C(R^9)$ and X^2 is $C(R^{10})$.

[0063] In a preferred embodiment, R^1 is phenyl substituted by 0 or 1 R^2 substituents, and the phenyl is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl. In one specific embodiment, R^1 is unsubstituted phenyl. In a further specific embodiment, R^1 is phenyl substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl.

[0064] In one specific embodiment, R^1 is selected from 2-methylphenyl, 2-chlorophenyl, 2-trifluoromethylphenyl, 2-fluorophenyl and 2-methoxyphenyl.

[0065] In a further preferred embodiment, R^1 is phenoxy.

[0066] In a further preferred embodiment, R^1 is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl.

[0067] In one specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl.

[0068] In a further specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl.

[0069] In one specific embodiment, R^1 is an unsubstituted unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S.

[0070] In a further specific embodiment, R^1 is selected from pyridyl and pyrimidinyl.

[0071] In one embodiment, R^3 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $C_{1-6}haloalkyl$, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$.

[0072] In one preferred embodiment, R^3 is H.

[0073] In another preferred embodiment, R^3 is selected from F, Cl, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $C_{1-6}haloalkyl$, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$.

[0074] In one embodiment, R^5 is, independently, in each instance, H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, or $C_{1-6}alkyl$ substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a $C_{3-6}spiroalkyl$ substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[0075] In one preferred embodiment, R^5 is H.

[0076] In another preferred embodiment, one R^5 is S-methyl, the other is H.

[0077] In another preferred embodiment, at least one R^5 is halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[0078] In a preferred embodiment, R^6 is H.

[0079] In another preferred embodiment, R^6 is F, Cl, cyano or nitro.

[0080] In a preferred embodiment, R^7 is H.

[0081] In another preferred embodiment, R^7 is F, Cl, cyano or nitro.

[0082] In a preferred embodiment, R^8 is selected from H, CF_3 , C_{1-3} alkyl, Br, Cl and F.

[0083] In one specific embodiment, R^8 is H.

[0084] In another specific embodiment, R^8 is selected from CF_3 , C_{1-3} alkyl, Br, Cl and F.

[0085] In a preferred embodiment, R^9 is H.

[0086] In another embodiment, R^9 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, —
 $C(=O)R^a$, — $C(=O)OR^a$, — $C(=O)NR^aR^a$, — $C(=NR^a)NR^aR^a$, — OR^a , — $OC(=O)R^a$, —
 $OC(=O)NR^aR^a$, — $OC(=O)N(R^a)S(=O)_2R^a$, — $OC_{2-6}alkylNR^aR^a$, — $OC_{2-6}alkylOR^a$, — SR^a , —
 $S(=O)R^a$, — $S(=O)_2R^a$, — $S(=O)_2NR^aR^a$, — $S(=O)_2N(R^a)C(=O)R^a$, — $S(=O)_2N(R^a)C(=O)OR^a$,
— $S(=O)_2N(R^a)C(=O)NR^aR^a$, — NR^aR^a , — $N(R^a)C(=O)R^a$, — $N(R^a)C(=O)OR^a$, —
 $N(R^a)C(=O)NR^aR^a$, — $N(R^a)C(=NR^a)NR^aR^a$, — $N(R^a)S(=O)_2R^a$, — $N(R^a)S(=O)_2NR^aR^a$, —
 $NR^aC_{2-6}alkylNR^aR^a$, — $NR^aC_{2-6}alkylOR^a$, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle,
wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by
0, 1, 2 or 3 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, — $C(=O)R^a$, —
 $C(=O)OR^a$, — $C(=O)NR^aR^a$, — $C(=NR^a)NR^aR^a$, — OR^a , — $OC(=O)R^a$, — $OC(=O)NR^aR^a$, —
 $OC(=O)N(R^a)S(=O)_2R^a$, — $OC_{2-6}alkylOR^a$, — SR^a , — $S(=O)R^a$, — $S(=O)_2R^a$, — $S(=O)_2NR^aR^a$,
— $S(=O)_2N(R^a)C(=O)R^a$, — $S(=O)_2N(R^a)C(=O)OR^a$, — $S(=O)_2N(R^a)C(=O)NR^aR^a$, — NR^aR^a ,
— $N(R^a)C(=O)R^a$, — $N(R^a)C(=O)OR^a$, — $N(R^a)C(=O)NR^aR^a$, — $N(R^a)C(=NR^a)NR^aR^a$, —
 $N(R^a)S(=O)_2R^a$, — $N(R^a)S(=O)_2NR^aR^a$, — $NR^aC_{2-6}alkylNR^aR^a$, — $NR^aC_{2-6}alkylOR^a$.


[0087] In one embodiment, R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$.

[0088] In one preferred embodiment, R^{10} is H.

[0089] In another preferred embodiment, R^{10} is cyano, nitro, CO_2R^a , $C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $S(=O)R^b$, $S(=O)_2R^b$ or $S(=O)_2NR^aR^a$.

[0090] In one preferred embodiment, R^{11} is H.

[0091] In another exemplary embodiment, the PI3K inhibitor is a compound of Formula (VII):



or a pharmaceutically-acceptable salt thereof, wherein:

X¹ is C(R⁹) or N;

X² is C(R¹⁰) or N;

Y is N(R¹¹), O or S;

Z is CR⁸ or N; R¹ is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R² substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl;

R² is selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a; or R² is selected from C₁₋₆alkyl, phenyl, benzyl, heteroaryl, heterocycle, —(C₁₋₃alkyl)heteroaryl, —(C₁₋₃alkyl)heterocycle, —O(C₁₋₃alkyl)heteroaryl, —O(C₁₋₃alkyl)heterocycle, —NR^a(C₁₋₃alkyl)heteroaryl, —NR^a(C₁₋₃alkyl)heterocycle, —(C₁₋₃alkyl)phenyl, —O(C₁₋₃alkyl)phenyl and —NR^a(C₁₋₃alkyl)phenyl all of which are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₄haloalkyl, OC₁₋₄alkyl, Br, Cl, F, I and C₁₋₄alkyl;

R³ is selected from H, halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR¹, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₂₋₆alkylOR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₆haloalkyl, OC₁₋₆alkyl, Br, Cl, F, I and C₁₋₆alkyl;

R^5 is, independently, in each instance, H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a C_{3-6} -spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$;

R^6 is selected from H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^aS(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;

R^7 is selected from H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^aS(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;

R^8 is selected from H, C_{1-6} haloalkyl, Br, Cl, F, I, OR^a , NR^aR^a , C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, OC_{1-6} alkyl, Br, Cl, F, I and C_{1-6} alkyl;

R^9 is selected from H, halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^8$, $-OC(=O)R^8$, $-OC(=O)NR^2R^8$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^8$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$; or R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected

from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a;

R¹⁰ is H, C₁₋₃alkyl, C₁₋₃haloalkyl, cyano, nitro, CO₂R^a, C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —S(=O)R^b, S(=O)₂R^b or S(=O)₂NR^aR^a;

R¹¹ is H or C₁₋₄alkyl;

R^a is independently, at each instance, H or R^b; and

R^b is independently, at each instance, phenyl, benzyl or C₁₋₆alkyl, the phenyl, benzyl and C₁₋₆alkyl being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alkyl, C₁₋₃haloalkyl, —OC₁₋₄alkyl, —NH₂, —NHC₁₋₄alkyl, —N(C₁₋₄alkyl)C₁₋₄alkyl.

[0092] In a preferred embodiment, X¹ is C(R⁹). In a further preferred embodiment, X¹ is C(R⁹) and X² is N. In a further embodiment, X¹ is C(R⁹) and X² is C(R¹⁰).

[0093] In a preferred embodiment, R¹ is phenyl substituted by 0 or 1 R² substituents, and the phenyl is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl. In one specific embodiment, R¹ is unsubstituted phenyl. In a further specific embodiment, R¹ is phenyl substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl.

[0094] In one specific embodiment, R¹ is selected from 2-methylphenyl, 2-chlorophenyl, 2-trifluoromethylphenyl, 2-fluorophenyl and 2-methoxyphenyl.

[0095] In a further preferred embodiment, R¹ is phenoxy.

[0096] In a further preferred embodiment, R^1 is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[0097] In one specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[0098] In a further specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[0099] In one specific embodiment, R^1 is an unsubstituted unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S.

[00100] In a further specific embodiment, R^1 is selected from pyridyl and pyrimidinyl.

[00101] In one embodiment, R^3 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$.

[00102] In one preferred embodiment, R^3 is H.

[00103] In another preferred embodiment, R^3 is selected from F, Cl, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, OC_{1-6} alkyl, Br, Cl, F, I and C_{1-6} alkyl.

[00104] In one embodiment, R^5 is, independently, in each instance, H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a C_{3-6} spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[00105] In one preferred embodiment, R^5 is H.

[00106] In another preferred embodiment, one R^5 is S-methyl, the other is H.

[00107] In another preferred embodiment, at least one R^5 is halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$.

[00108] In a preferred embodiment, R^6 is H.

[00109] In another preferred embodiment, R^6 is F, Cl, cyano or nitro.

[00110] In a preferred embodiment, R^7 is H.

[00111] In another preferred embodiment, R^7 is F, Cl, cyano or nitro.

[00112] In a preferred embodiment, R^8 is selected from H, CF_3 , C_{1-3} alkyl, Br, Cl and F.

[00113] In one specific embodiment, R^8 is H.

[00114] In another specific embodiment, R^8 is selected from CF_3 , C_{1-3} alkyl, Br, Cl and F.

[00115] In a preferred embodiment, R^9 is H.

[00116] In another embodiment, R^9 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $—C(=O)R^a$, $—C(=O)OR^a$, $—C(=O)NR^aR^a$, $—C(=NR^a)NR^aR^a$, $—OR^a$, $—OC(=O)R^a$, $—OC(=O)NR^aR^a$, $—OC(=O)N(R^a)S(=O)_2R^a$, $—OC_{2-6}alkylNR^aR^a$, $—OC_{2-6}alkylOR^a$, $—SR^a$, $—S(=O)R^a$, $—S(=O)_2R^a$, $—S(=O)_2NR^aR^a$, $—S(=O)_2N(R^a)C(=O)R^a$, $—S(=O)_2N(R^a)C(=O)OR^a$, $—S(=O)_2N(R^a)C(=O)NR^aR^a$, $—NR^aR^a$, $—N(R^a)C(=O)R^a$, $—N(R^a)C(=O)OR^a$, $—N(R^a)C(=O)NR^aR^a$, $—N(R^a)C(=NR^a)NR^aR^a$, $—N(R^a)S(=O)_2R^a$, $—N(R^a)S(=O)_2NR^aR^a$, $—NR^aC_{2-6}alkylNR^aR^a$, $—NR^aC_{2-6}alkylOR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $—C(=O)R^a$, $—C(=O)OR^a$, $—C(=O)NR^aR^a$, $—C(=NR^a)NR^aR^a$, $—OR^a$, $—OC(=O)R^a$, $—OC(=O)NR^aR^a$, $—OC(=O)N(R^a)S(=O)_2R^a$, $—OC_{2-6}alkylOR^a$, $—SR^a$, $—S(=O)R^a$, $—S(=O)_2R^a$, $—S(=O)_2NR^aR^a$, $—S(=O)_2N(R^a)C(=O)R^a$, $—S(=O)_2N(R^a)C(=O)OR^a$, $—S(=O)_2N(R^a)C(=O)NR^aR^a$, $—NR^aR^a$, $—N(R^a)C(=O)R^a$, $—N(R^a)C(=O)OR^a$, $—N(R^a)C(=O)NR^aR^a$, $—N(R^a)C(=NR^a)NR^aR^a$, $—N(R^a)S(=O)_2R^a$, $—N(R^a)S(=O)_2NR^aR^a$, $—NR^aC_{2-6}alkylNR^aR^a$, $—NR^aC_{2-6}alkylOR^a$.


[00117] In one embodiment, R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $—C(=O)R^a$, $—C(=O)OR^a$, $—C(=O)NR^aR^a$, $—C(=NR^a)NR^aR^a$, $—OR^a$, $—OC(=O)R^a$, $—OC(=O)NR^aR^a$, $—OC(=O)N(R^a)S(=O)_2R^a$, $—OC_{2-6}alkylNR^aR^a$, $—OC_{2-6}alkylOR^a$, $—SR^a$, $—S(=O)R^a$, $—S(=O)_2R^a$, $—S(=O)_2NR^aR^a$, $—S(=O)_2N(R^a)C(=O)R^a$, $—S(=O)_2N(R^a)C(=O)OR^a$, $—S(=O)_2N(R^a)C(=O)NR^aR^a$, $—NR^aR^a$, $—N(R^a)C(=O)R^a$, $—N(R^a)C(=O)OR^a$, $—N(R^a)C(=O)NR^aR^a$, $—N(R^a)C(=NR^a)NR^aR^a$, $—N(R^a)S(=O)_2R^a$, $—N(R^a)S(=O)_2NR^aR^a$, $—NR^aC_{2-6}alkylNR^aR^a$ and $—NR^aC_{2-6}alkylOR^a$.

[00118] In one preferred embodiment, R^{10} is H.

[00119] In another preferred embodiment, R^{10} is cyano, nitro, CO_2R^a , $C(=O)NR^aR^a$, $—C(=NR^a)NR^aR^a$, $—S(=O)_2N(R^a)C(=O)R^a$, $—S(=O)_2N(R^a)C(=O)OR^a$, $—S(=O)_2N(R^a)C(=O)NR^aR^a$, $S(=O)R^b$, $S(=O)_2R^b$ or $S(=O)_2NR^aR^a$.

In one preferred embodiment, R¹¹ is H.

[00120] In another exemplary embodiment, the PI3K inhibitor is a compound of Formula (VIII):



or a pharmaceutically-acceptable salt thereof, wherein:

X¹ is C(R⁹) or N;

X² is C(R¹⁰) or N;

Y is N(R¹¹), O or S;

Z is CR⁸ or N;

R¹ is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R² substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C₁₋₄alkyl, OC₁₋₄alkyl, OC₁₋₄haloalkyl, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl and C₁₋₄haloalkyl;

R² is selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a—C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC_{2,6}alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —

$N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$; or R^2 is selected from $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl, heterocycle, $-(C_{1-3}alkyl)heteroaryl$, $-(C_{1-3}alkyl)heterocycle$, $-O(C_{1-3}alkyl)heteroaryl$, $-O(C_{1-3}alkyl)heterocycle$, $-NR^a(C_{1-3}alkyl)heteroaryl$, $-NR^a(C_{1-3}alkyl)heterocycle$, $-(C_{1-3}alkyl)phenyl$, $-O(C_{1-3}alkyl)phenyl$ and $-NR^a(C_{1-3}alkyl)phenyl$ all of which are substituted by 0, 1, 2 or 3 substituents selected from $C_{1-4}haloalkyl$, $OC_{1-4}alkyl$, Br, Cl, F, I and $C_{1-4}alkyl$;

R^3 is selected from H, halo, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $C_{1-6}haloalkyl$, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$;

R^5 is, independently, in each instance, H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, or $C_{1-6}alkyl$ substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a C_{3-6} -spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$;

R^6 is selected from H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;

R^7 is selected from H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;

R^8 is selected from H, $C_{1-6}haloalkyl$, Br, Cl, F, I, OR^a , NR^aR^a , $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $C_{1-6}haloalkyl$, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$;

R^9 is selected from H, halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$; or R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$;

R^{10} is H, $C_{1-3}alkyl$, $C_{1-3}haloalkyl$, cyano, nitro, CO_2R^a , $C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$ or $S(=O)_2NR^aR^a$;

R^{11} is H or $C_{1-4}alkyl$;

R^a is independently, at each instance, H or R^b ; and

R^b is independently, at each instance, phenyl, benzyl or C_{1-6} alkyl, the phenyl, benzyl and C_{1-6} alkyl being substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} alkyl, C_{1-3} haloalkyl, $-OC_{1-4}$ alkyl, $-NH_2$, $-NHC_{1-4}$ alkyl, $-N(C_{1-4}alkyl)C_{1-4}alkyl$.

[00121] In a preferred embodiment, X^1 is $C(R^9)$. In a further preferred embodiment, X^1 is $C(R^9)$ and X^2 is N. In a further embodiment, X^1 is $C(R^9)$ and X^2 is $C(R^{10})$.

[00122] In a preferred embodiment, R^1 is phenyl substituted by 0 or 1 R^2 substituents, and the phenyl is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl. In one specific embodiment, R^1 is unsubstituted phenyl. In a further specific embodiment, R^1 is phenyl substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[00123] In one specific embodiment, R^1 is selected from 2-methylphenyl, 2-chlorophenyl, 2-trifluoromethylphenyl, 2-fluorophenyl and 2-methoxyphenyl.

[00124] In a further preferred embodiment, R^1 is phenoxy.

[00125] In a further preferred embodiment, R^1 is a direct-bonded or oxygen-linked saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[00126] In one specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl.

[00127] In a further specific embodiment, R^1 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the ring is substituted by 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl and C_{1-4} haloalkyl.

[00128] In one specific embodiment, R^1 is an unsubstituted unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S.

[00129] In a further specific embodiment, R^1 is selected from pyridyl and pyrimidinyl.

[00130] In one embodiment, R^3 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^a$, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, OC_{1-6} alkyl, Br, Cl, F, I and C_{1-6} alkyl.

[00131] In one preferred embodiment, R^3 is H.

[00132] In another preferred embodiment, R^3 is selected from F, Cl, C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C_{1-6} alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from C_{1-6} haloalkyl, OC_{1-6} alkyl, Br, Cl, F, I and C_{1-6} alkyl.

[00133] In one embodiment, R^5 is, independently, in each instance, H, halo, C_{1-6} alkyl, C_{1-4} haloalkyl, or C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl; or both R^5 groups together form a C_{3-6} spiroalkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, OC_{1-4} alkyl, C_{1-4} alkyl, C_{1-3} haloalkyl, OC_{1-4} alkyl, NH_2 , NHC_{1-4} alkyl, $N(C_{1-4}$ alkyl) C_{1-4} alkyl.

[00134] In one preferred embodiment, R⁵ is H.

[00135] In another preferred embodiment, one R⁵ is S-methyl, the other is H.

[00136] In another preferred embodiment, at least one R⁵ is halo, C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, OC₁₋₄alkyl, C₁₋₄alkyl, C₁₋₃haloalkyl, OC₁₋₄alkyl, NH₂, NHC₁₋₄alkyl, N(C₁₋₄alkyl)C₁₋₄alkyl.

[00137] In a preferred embodiment, R⁶ is H.

[00138] In another preferred embodiment, R⁶ is F, Cl, cyano or nitro.

[00139] In a preferred embodiment, R⁷ is H.

[00140] In another preferred embodiment, R⁷ is F, Cl, cyano or nitro.

[00141] In a preferred embodiment, R⁸ is selected from H, CF₃, C₁₋₃alkyl, Br, Cl and F.

[00142] In one specific embodiment, R⁸ is H.

[00143] In another specific embodiment, R⁸ is selected from CF₃, C₁₋₃alkyl, Br, Cl and F.

[00144] In a preferred embodiment, R⁹ is H.

[00145] In another embodiment, R⁹ is selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₂₋₆alkylOR^a, C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a,

$-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{OR}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{NR}^a)\text{NR}^a\text{R}^a$, $-\text{N}(\text{R}^a)\text{S}(=\text{O})_2\text{R}^a$, $-\text{N}(\text{R}^a)\text{S}(=\text{O})_2\text{NR}^a\text{R}^a$, $-\text{NR}^a\text{C}_{2-6}\text{alkylNR}^a\text{R}^a$, $-\text{NR}^a\text{C}_{2-6}\text{alkylOR}^a$.


[00146] In one embodiment, R^9 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, $\text{C}_{1-4}\text{haloalkyl}$, cyano, nitro, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{NR}^a\text{R}^a$, $-\text{OR}^a$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{OC}(=\text{O})\text{N}(\text{R}^a)\text{S}(=\text{O})_2\text{R}^a$, $-\text{OC}_{2-6}\text{alkylNR}^a\text{R}^a$, $-\text{OC}_{2-6}\text{alkylOR}^a$, $-\text{SR}^a$, $-\text{S}(=\text{O})\text{R}^a$, $-\text{S}(=\text{O})_2\text{R}^a$, $-\text{S}(=\text{O})_2\text{NR}^a\text{R}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{OR}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{NR}^a\text{R}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{OR}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{N}(\text{R}^a)\text{C}(=\text{NR}^a)\text{NR}^a\text{R}^a$, $-\text{N}(\text{R}^a)\text{S}(=\text{O})_2\text{R}^a$, $-\text{N}(\text{R}^a)\text{S}(=\text{O})_2\text{NR}^a\text{R}^a$, $-\text{NR}^a\text{C}_{2-6}\text{alkylNR}^a\text{R}^a$ and $-\text{NR}^a\text{C}_{2-6}\text{alkylOR}^a$.

[00147] In one preferred embodiment, R^{10} is H.

[00148] In another preferred embodiment, R^{10} is cyano, nitro, CO_2R^a , $\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{NR}^a\text{R}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{OR}^a$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^a)\text{C}(=\text{O})\text{NR}^a\text{R}^a$, $\text{S}(=\text{O})\text{R}^b$, $\text{S}(=\text{O})_2\text{R}^b$ or $\text{S}(=\text{O})_2\text{NR}^a\text{R}^a$.


[00149] In one preferred embodiment, R^{11} is H.

[00150] In an exemplary preferred embodiment, the PI3K inhibitor is a compound of Formula (IX):




which is (*S*)-*N*-(1-(7-fluoro-2-(pyridin-2-yl)quinolin-3-yl)ethyl)-9*H*-purin-6-amine, or a pharmaceutically-acceptable salt thereof. In an exemplary embodiment, the PI3K- δ inhibitor or PI3K- δ inhibitor is (*S*)-*N*-(1-(7-fluoro-2-(pyridin-2-yl)quinolin-3-yl)ethyl)-9*H*-purin-6-amine or a pharmaceutically-acceptable salt thereof.

[00151] In an exemplary embodiment, the PI3K inhibitor is a PI3K- δ inhibitor which is a compound of Formula (X):



which is (*S*)-*N*-(1-(6-fluoro-3-(pyridin-2-yl)quinoxalin-2-yl)ethyl)-9*H*-purin-6-amine, or a pharmaceutically-acceptable salt thereof.


[00152] In an exemplary embodiment, the PI3K inhibitor is a PI3K- δ inhibitor, which is a compound of Formula (XI):



which is (*S*)-*N*-(1-(2-(3,5-difluorophenyl)-8-fluoroquinolin-3-yl)ethyl)-9*H*-purin-6-amine, or a pharmaceutically-acceptable salt thereof.

[00153] In an exemplary embodiment, the PI3K inhibitor or PI3K- δ inhibitor is (*S*)-*N*-(1-(2-(3,5-difluorophenyl)-8-fluoroquinolin-3-yl)ethyl)-9*H*-purin-6-amine or a pharmaceutically-acceptable salt thereof.


[00154] In an exemplary embodiment, the PI3K inhibitor is a PI3K- δ inhibitor which is a compound of Formula (XII):



which is (*S*)-3-(1-((9*H*-purin-6-yl)amino)ethyl)-2-(pyridin-2-yl)quinoline-8-carbonitrile, or a pharmaceutically-acceptable salt thereof

[00155] In an exemplary embodiment, the PI3K inhibitor or PI3K- δ inhibitor is (*S*)-3-(1-((9*H*-purin-6-yl)amino)ethyl)-2-(pyridin-2-yl)quinoline-8-carbonitrile or a pharmaceutically-acceptable salt thereof.


[00156] In an exemplary embodiment, the PI3K inhibitor is a PI3K- δ inhibitor which is a compound of Formula (XIII):



which is (*S*)-*N*-(1-(5,7-difluoro-2-(pyridin-2-yl)quinolin-3-yl)ethyl)-9*H*-purin-6-amine, or a pharmaceutically-acceptable salt thereof

[00157] In an exemplary embodiment, the PI3K inhibitor or PI3K- δ inhibitor is (*S*)-*N*-(1-(5,7-difluoro-2-(pyridin-2-yl)quinolin-3-yl)ethyl)-9*H*-purin-6-amine or a pharmaceutically-acceptable salt thereof.

[00158] In an exemplary embodiment, the PI3K inhibitor is a compound selected from the structures disclosed in U.S. Patent Nos. 7,932,260 and 8,207,153. In an exemplary embodiment, the PI3K inhibitor is a compound of Formula (XIV):



wherein

X and Y, independently, are N or CH;

Z is N—R⁷ or O;

R¹ are the same and are hydrogen, halo, or C₁₋₃alkyl;

R² and R³, independently, are hydrogen, halo, or C₁₋₃alkyl;

R⁴ is hydrogen, halo, OR^a, CN, C₂₋₆alkynyl, C(=O)R^a, C(=O)NR^aR^b, C₃₋₆heterocycloalkyl, C₁₋₃alkyleneC₃₋₆heterocycloalkyl, OC₁₋₃alkyleneOR^a, OC₁₋₃alkyleneNR^aR^b, OC₁₋₃alkyleneC₃₋₆cycloalkyl, OC₃₋₆heterocycloalkyl, OC₁₋₃alkyleneC≡CH, or OC₁₋₃alkyleneC(=O)NR^aR^b;

R⁵ is C₁₋₃alkyl, CH₂CF₃, phenyl, CH₂C≡CH, C₁₋₃alkyleneOR^c, C₁₋₄alkyleneNR^aR^b, or C₁₋₄alkyleneNHC(=O)OR^a,

R⁶ is hydrogen, halo, or NR^aR^b;

R⁷ is hydrogen or R⁵ and R⁷ are taken together with the atoms to which they are attached to form a five- or six-membered saturated ring;

R⁸ is C₁₋₃alkyl, halo, CF₃, or CH₂C₃₋₆heterocycloalkyl;

n is 0, 1, or 2;


R^a is hydrogen, C₁₋₄alkyl, or CH₂C₆H₅;

R^b is hydrogen or C₁₋₃alkyl; and

R^c is hydrogen, C₁₋₃alkyl, or halo,

wherein when the R¹ groups are different from hydrogen, R² and R⁴ are the same; or a pharmaceutically acceptable salt, or prodrug, or solvate (e.g., hydrate) thereof.

[00159] In a preferred embodiment, the PI3K inhibitor is an enantiomer of Formula (XIV), as shown in Formula (XV):




wherein X, Y, Z, R¹ through R⁸, R^a, R^b, R^c, and n are as defined above for Formula (XIV).

[00160] In various embodiments exhibiting increased potency relative to other compounds, R⁸ is C₁₋₃alkyl, F, Cl, or CF₃. Alternatively, in such embodiments, n is 0 (such that there is no R⁸ substituent).

[00161] In further embodiments exhibiting increased potency, X is N and Y is CH. Alternatively, X and Y may also both be CH.

[00162] In preferred embodiments, Z is N—R⁷, and the bicyclic ring system containing X and Y is:



[00163] In further embodiments exhibiting increased potency, R⁶ is hydrogen, halo, or NH₂. Preferably, R⁶ is hydrogen.


[00164] In preferred embodiments exhibiting such increased potency, n is 0 or 1; R⁸ (if n is 1) is C₁₋₃alkyl, F, Cl, or CF₃; R⁶ is hydrogen; X is N and Y is CH or X and Y are both CH; Z is NH;

R^1 are the same and are hydrogen, halo, or C_{1-3} alkyl; and R^2 and R^3 , independently, are hydrogen, halo, or C_{1-3} alkyl. Preferably, R^1 , R^2 , and R^3 are hydrogen.


[00165] Unexpectedly, potency against PI3K- δ is conserved when R^1 is the same. In structural formulae (XIV) and (XV), R^2 and R^4 may differ provided that R^1 is H. When R^1 is H, free rotation is unexpectedly permitted about the bond connecting the phenyl ring substituent to the quinazoline ring, and the compounds advantageously do not exhibit atropisomerism (i.e., multiple diastereomer formation is avoided). Alternatively, R^2 and R^4 can be the same such that the compounds advantageously do not exhibit atropisomerism.

[00166] As used with respect to Formula (XIV) and Formula (XV), the term “alkyl” is defined as straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, e.g., methyl, ethyl, and straight chain and branched propyl and butyl groups. The terms “ C_{1-3} alkylene” and “ C_{1-4} alkylene” are defined as hydrocarbon groups containing the indicated number of carbon atoms and one less hydrogen than the corresponding alkyl group. The term “ C_{2-6} alkynyl” is defined as a hydrocarbon group containing the indicated number of carbon atoms and a carbon-carbon triple bond. The term “ C_{3-6} cycloalkyl” is defined as a cyclic hydrocarbon group containing the indicated number of carbon atoms. The term “ C_{2-6} heterocycloalkyl” is defined similarly as cycloalkyl except the ring contains one or two heteroatoms selected from the group consisting of O, NR^a , and S. The term “halo” is defined as fluoro, bromo, chloro, and iodo.


[00167] In other preferred embodiments, R^1 is hydrogen, fluoro, chloro, methyl, or



and R^2 is hydrogen, methyl, chloro, or fluoro; R^3 is hydrogen or fluoro; R^6 is NH_2 , hydrogen, or fluoro; R^7 is hydrogen or R^5 and R^7 are taken together to form




R^8 is methyl, trifluoromethyl, chloro, or fluoro; R^4 is hydrogen, fluoro, chloro, OH, OCH_3 , $OCH_2C\equiv CH$, $O(CH_2)_2N(CH_3)_2$, $C(=O)CH_3$, $C\equiv CH$, CN, $C(=O)NH_2$, $OCH_2C(=O)NH_2$, $O(CH_2)_2OCH_3$, $O(CH_2)_2N(CH_3)_2$,



and R^5 is methyl, ethyl, propyl, phenyl, CH_2OH , $CH_2OCH_2C_6H_5$, CH_2CF_3 , $CH_2OC(CH_3)_3$, $CH_2C\equiv CH$, $(CH_2)_3N(C_2H_5)_2$, $(CH_2)_3NH_2$, $(CH_2)_4NH_2$, $(CH_2)_3NHC(=O)OCH_2C_6H_5$, or $(CH_2)_4NHC(=O)OCH_2C_6H_5$; R^c is hydrogen, methyl, fluoro, or bromo; and n is 0 or 1.

[00168] In a preferred embodiment, the PI3K inhibitor is a PI3K- δ inhibitor of Formula (XVI):



which is (*S*)-2-(1-((9*H*-purin-6-yl)amino)propyl)-5-fluoro-3-phenylquinazolin-4(3*H*)-one (other names: 4(3*H*)-quinazolinone, 5-fluoro-3-phenyl-2-[(1*S*)-1-(9*H*-purin-6-ylamino)propyl], and 5-fluoro-3-phenyl-2-[(1*S*)-1-[(7*H*-purin-6-yl)amino]propyl} quinazolin-4(3*H*)-one) or a pharmaceutically-acceptable salt thereof.

[00169] In a preferred embodiment, the PI3K inhibitor or PI3K- δ inhibitor is (S)-2-(1-((9H-purin-6-yl)amino)propyl)-5-fluoro-3-phenylquinazolin-4(3H)-one or a pharmaceutically-acceptable salt thereof.


[00170] In an embodiment, the PI3K inhibitor or PI3K- δ inhibitor is 4(3H)-quinazolinone, 5-fluoro-3-phenyl-2-[(1S)-1-(9H-purin-6-ylamino)propyl]-5-fluoro-3-phenyl-2-[(1S)-1-[(7H-purin-6-yl)amino]propyl]quinazolin-4(3H)-one or a pharmaceutically-acceptable salt thereof

[00171] Other PI3K inhibitors suitable for use in the described combination with a BTK inhibitor also include, but are not limited to, those described in, for example, U.S. Patent No. 8,193,182 and U.S. Published Application Nos. 2013/0267521; 2013/0053362; 2013/0029984; 2013/0029982; 2012/0184568; and 2012/0059000.

BTK Inhibitors

[00172] The BTK inhibitor may be any BTK inhibitor known in the art. In particular, it is one of the BTK inhibitors described in more detail in the following paragraphs. Preferably, it is a compound of Formula XVII or a pharmaceutically acceptable salt thereof. In one specific embodiment, it is a compound of Formula XVIII or a pharmaceutically acceptable salt thereof.

[00173] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XVII):



Formula (XVII)

or a pharmaceutically acceptable salt thereof,

wherein:

X is CH, N, O or S;

Y is C(R₆), N, O or S;

Z is CH, N or bond;

A is CH or N;

B₁ is N or C(R₇);

B₂ is N or C(R₈);

B₃ is N or C(R₉);

B₄ is N or C(R₁₀);

R₁ is R₁₁C(=O), R₁₂S(=O), R₁₃S(=O)₂ or (1-6C)alkyl optionally substituted with R₁₄;

R₂ is H, (C₁₋₃)alkyl or (C₃₋₇)cycloalkyl;

R₃ is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl); or

R₂ and R₃ form, together with the N and C atom they are attached to, a (3-7C)heterocycloalkyl optionally substituted with one or more fluorine, hydroxyl, (1-3C)alkyl, (1-3C)alkoxy or oxo;

R₄ is H or (C₁₋₃)alkyl;

R₅ is H, halogen, cyano, (C₁₋₄)alkyl, (C₁₋₃)alkoxy, (C₃₋₆)cycloalkyl, any alkyl group of which is optionally substituted with one or more halogen; or R₅ is (C₆₋₁₀)aryl or (C₂₋₆)heterocycloalkyl;

R₆ is H or (C₁₋₃)alkyl; or

R₅ and R₆ together may form a (C₃₋₇)cycloalkenyl or (C₂₋₆)heterocycloalkenyl, each optionally substituted with (C₁₋₃)alkyl or one or more halogens;

R₇ is H, halogen, CF₃, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R₈ is H, halogen, CF₃, (C₁₋₃)alkyl or (C₁₋₃)alkoxy; or

R₇ and R₈ together with the carbon atoms they are attached to, form (C₆₋₁₀)aryl or (C₁₋₉)heteroaryl;

R₉ is H, halogen, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R₁₀ is H, halogen, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R₁₁ is independently selected from the group consisting of (C₁₋₆)alkyl, (C₂₋₆)alkenyl and (C₂₋₆)alkynyl, where each alkyl, alkenyl or alkynyl is optionally substituted with one or more substituents selected from the group consisting of hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl and (C₃₋₇)heterocycloalkyl; or R₁₁ is (C₁₋₃)alkyl-C(O)-S-(C₁₋₃)alkyl; or

R₁₁ is (C₁₋₅)heteroaryl optionally substituted with one or more substituents selected from the group consisting of halogen or cyano;

R₁₂ and R₁₃ are independently selected from the group consisting of (C₂₋₆)alkenyl or (C₂₋₆)alkynyl, both optionally substituted with one or more substituents selected from the group consisting of hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl and (C₃₋₇)heterocycloalkyl; or a (C₁₋₅)heteroaryl optionally substituted with one or more substituents selected from the group consisting of halogen and cyano; and

R₁₄ is independently selected from the group consisting of halogen, cyano, (C₂₋₆)alkenyl and (C₂₋₆)alkynyl, both optionally substituted with one or more substituents selected from the group consisting of hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₄)alkylamino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl, (C₁₋₅)heteroaryl and (C₃₋₇)heterocycloalkyl;

with the proviso that:

0 to 2 atoms of X, Y, Z can simultaneously be a heteroatom;

when one atom selected from X, Y is O or S, then Z is a bond and the other atom selected from X, Y can not be O or S;

when Z is C or N then Y is C(R₆) or N and X is C or N;

0 to 2 atoms of B₁, B₂, B₃ and B₄ are N;

with the terms used having the following meanings:

(C₁₋₂)alkyl means an alkyl group having 1 to 2 carbon atoms, being methyl or ethyl,

(C₁₋₃)alkyl means a branched or unbranched alkyl group having 1-3 carbon atoms, being methyl, ethyl, propyl or isopropyl;

(C₁₋₄)alkyl means a branched or unbranched alkyl group having 1-4 carbon atoms, being methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl, (C₁₋₃)alkyl groups being preferred;

(C₁₋₅)alkyl means a branched or unbranched alkyl group having 1-5 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl and isopentyl, (C₁₋₄)alkyl groups being preferred. (C₁₋₆)alkyl means a branched or unbranched alkyl group having 1-6 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl and n-hexyl. (C₁₋₅)alkyl groups are preferred, (C₁₋₄)alkyl being most preferred;

(C₁₋₂)alkoxy means an alkoxy group having 1-2 carbon atoms, the alkyl moiety having the same meaning as previously defined;

(C₁₋₃)alkoxy means an alkoxy group having 1-3 carbon atoms, the alkyl moiety having the same meaning as previously defined. (C₁₋₂)alkoxy groups are preferred;

(C₁₋₄)alkoxy means an alkoxy group having 1-4 carbon atoms, the alkyl moiety having the same meaning as previously defined. (C₁₋₃)alkoxy groups are preferred, (C₁₋₂)alkoxy groups being most preferred;

(C₂₋₄)alkenyl means a branched or unbranched alkenyl group having 2-4 carbon atoms, such as ethenyl, 2-propenyl, isobutenyl or 2-butenyl;

(C₂₋₆)alkenyl means a branched or unbranched alkenyl group having 2-6 carbon atoms, such as ethenyl, 2-butenyl, and n-pentenyl, (C₂₋₄)alkenyl groups being most preferred;

(C₂₋₄)alkynyl means a branched or unbranched alkynyl group having 2-4 carbon atoms, such as ethynyl, 2-propynyl or 2-butylnyl;

(C₂₋₆)alkynyl means a branched or unbranched alkynyl group having 2-6 carbon atoms, such as ethynyl, propynyl, n-butylnyl, n-pentylnyl, isopentylnyl, isohexynyl or n-hexynyl. (C₂₋₄)alkynyl

- groups are preferred; (C₃₋₆)cycloalkyl means a cycloalkyl group having 3-6 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;
- (C₃₋₇)cycloalkyl means a cycloalkyl group having 3-7 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl;
- (C₂₋₆)heterocycloalkyl means a heterocycloalkyl group having 2-6 carbon atoms, preferably 3-5 carbon atoms, and one or two heteroatoms selected from N, O and/or S, which may be attached via a heteroatom if feasible, or a carbon atom; preferred heteroatoms are N or O; also preferred are piperidine, morpholine, pyrrolidine and piperazine; with the most preferred (C₂₋₆)heterocycloalkyl being pyrrolidine; the heterocycloalkyl group may be attached via a heteroatom if feasible;
- (C₃₋₇)heterocycloalkyl means a heterocycloalkyl group having 3-7 carbon atoms, preferably 3-5 carbon atoms, and one or two heteroatoms selected from N, O and/or S. Preferred heteroatoms are N or O; preferred (C₃₋₇) heterocycloalkyl groups are azetidiny, pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl; more preferred (C₃₋₇)heterocycloalkyl groups are piperidine, morpholine and pyrrolidine; and the heterocycloalkyl group may be attached via a heteroatom if feasible;
- (C₃₋₇)cycloalkoxy means a cycloalkyl group having 3-7 carbon atoms, with the same meaning as previously defined, attached via a ring carbon atom to an exocyclic oxygen atom;
- (C₆₋₁₀)aryl means an aromatic hydrocarbon group having 6-10 carbon atoms, such as phenyl, naphthyl, tetrahydronaphthyl or indenyl; the preferred (C₆₋₁₀)aryl group is phenyl;
- (C₁₋₅)heteroaryl means a substituted or unsubstituted aromatic group having 1-5 carbon atoms and 1-4 heteroatoms selected from N, O and/or S; the (C₁₋₅)heteroaryl may optionally be substituted; preferred (C₁₋₅)heteroaryl groups are tetrazolyl, imidazolyl, thiadiazolyl, pyridyl, pyrimidyl, triazinyl, thienyl or furyl, a more preferred (C₁₋₅)heteroaryl is pyrimidyl;
- (C₁₋₉)heteroaryl means a substituted or unsubstituted aromatic group having 1-9 carbon atoms and 1-4 heteroatoms selected from N, O and/or S; the (C₁₋₉)heteroaryl may optionally be substituted; preferred (C₁₋₉)heteroaryl groups are quinoline, isoquinoline and indole;
- [(C₁₋₄)alkyl]amino means an amino group, monosubstituted with an alkyl group containing 1-4 carbon atoms having the same meaning as previously defined; preferred [(C₁₋₄)alkyl]amino group is methylamino;

di[(C₁₋₄)alkyl]amino means an amino group, disubstituted with alkyl group(s), each containing 1-4 carbon atoms and having the same meaning as previously defined; preferred di[(C₁₋₄)alkyl]amino group is dimethylamino;

halogen means fluorine, chlorine, bromine or iodine;

(C₁₋₃)alkyl-C(O)-S-(C₁₋₃)alkyl means an alkyl-carbonyl-thio-alkyl group, each of the alkyl groups having 1 to 3 carbon atoms with the same meaning as previously defined;

(C₃₋₇)cycloalkenyl means a cycloalkenyl group having 3-7 carbon atoms, preferably 5-7 carbon atoms; preferred (C₃₋₇)cycloalkenyl groups are cyclopentenyl or cyclohexenyl; cyclohexenyl groups are most preferred;

(C₂₋₆)heterocycloalkenyl means a heterocycloalkenyl group having 2-6 carbon atoms, preferably 3-5 carbon atoms; and 1 heteroatom selected from N, O and/or S; preferred (C₂₋₆)heterocycloalkenyl groups are oxycyclohexenyl and azacyclohexenyl group.

In the above definitions with multifunctional groups, the attachment point is at the last group.

When, in the definition of a substituent, is indicated that "all of the alkyl groups" of said substituent are optionally substituted, this also includes the alkyl moiety of an alkoxy group.

A circle in a ring of Formula (XVII) indicates that the ring is aromatic.

Depending on the ring formed, the nitrogen, if present in X or Y, may carry a hydrogen.

[00174] In an exemplary embodiment of Formula (XVII), B₁ is C(R₇); B₂ is C(R₈); B₃ is C(R₉); B₄ is C(R₁₀); R₇, R₉, and R₁₀ are each H; and R₈ is hydrogen or methyl.

[00175] In an exemplary embodiment of Formula (XVII), the ring containing X, Y and Z is selected from the group consisting of pyridyl, pyrimidyl, pyridazyl, triazinyl, thiazolyl, oxazolyl and isoxazolyl.

[00176] In an exemplary embodiment of Formula (XVII), the ring containing X, Y and Z is selected from the group consisting of pyridyl, pyrimidyl and pyridazyl.

[00177] In an exemplary embodiment of Formula (XVII), the ring containing X, Y and Z is selected from the group consisting of pyridyl and pyrimidyl.

[00178] In an exemplary embodiment of Formula (XVII), the ring containing X, Y and Z is pyridyl.

[00179] In an exemplary embodiment of Formula (XVII), R₅ is selected from the group consisting of hydrogen, fluorine, methyl, methoxy and trifluoromethyl.

[00180] In an exemplary embodiment of Formula (XVII), R₅ is hydrogen.

[00181] In an exemplary embodiment of Formula (XVII), R₂ and R₃ together form a heterocycloalkyl ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, homopiperidiny and morpholiny, optionally substituted with one or more of fluoro, hydroxyl, (C₁₋₃)alkyl and (C₁₋₃)alkoxy.

[00182] In an exemplary embodiment of Formula (XVII), R₂ and R₃ together form a heterocycloalkyl ring selected from the group consisting of azetidiny, pyrrolidiny and piperidiny.

[00183] In an exemplary embodiment of Formula (XVII), R₂ and R₃ together form a pyrrolidiny ring.

[00184] In an exemplary embodiment of Formula (XVII), R₁ is independently selected from the group consisting of (C₁₋₆)alkyl, (C₂₋₆)alkenyl or (C₂₋₆)alkynyl, each optionally substituted with one or more substituents selected from the group consisting of hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl] amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl and (C₃₋₇)heterocycloalkyl.

[00185] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X is N; Y and Z are CH; R₅ is CH₃; A is N; R₂, R₃ and R₄ are H; and R₁ is CO-CH₃.

[00186] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X and Y are N; Z is CH; R₅ is CH₃; A is N; R₂, R₃ and R₄ are H; and R₁ is CO-CH₃.

[00187] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X and Y are N; Z is CH; R₅ is CH₃; A is CH; R₂ and R₃ together form a piperidiny ring; R₄ is H; and R₁ is CO-ethenyl.


[00188] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X, Y and Z are CH; R₅ is H; A is CH; R₂ and R₃ together form a pyrrolidiny ring; R₄ is H; and R₁ is CO-propynyl.

[00189] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X, Y and Z are CH; R₅ is CH₃; A is CH; R₂ and R₃ together form a piperidinyl ring; R₄ is H; and R₁ is CO-propynyl.

[00190] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X and Y are N; Z is CH; R₅ is H; A is CH; R₂ and R₃ together form a morpholinyl ring; R₄ is H; and R₁ is CO-ethenyl.

[00191] In an exemplary embodiment of Formula (XVII), B₁, B₂, B₃ and B₄ are CH; X and Y are N; Z is CH; R₅ is CH₃; A is CH; R₂ and R₃ together form a morpholinyl ring; R₄ is H; and R₁ is CO-propynyl.


[00192] In an exemplary preferred embodiment, the BTK inhibitor is a compound of Formula (XVIII):



which is (*S*)-4-(8-amino-3-(1-(but-2-ynoyl)pyrrolidin-2-yl)imidazo[1,5-a]pyrazin-1-yl)-*N*-(pyridin-2-yl)benzamide, or a pharmaceutically-acceptable salt thereof. The preparation of this compound is described at Example 6 of International Patent Application Publication No. WO 2013/010868. The preparation of this compound and related structures are described in the Examples of International Patent Application Publication No. WO 2013/010868.

[00193] In an exemplary embodiment, the BTK inhibitor is (*S*)-4-(8-amino-3-(1-(but-2-ynoyl)pyrrolidin-2-yl)imidazo[1,5-a]pyrazin-1-yl)-*N*-(pyridin-2-yl)benzamide or pharmaceutically-acceptable salt thereof.

[00194] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XIX) or a pharmaceutically-acceptable salt of a compound of Formula (XIX):



Formula (XIX)

In Formula (XIX) the substituents are defined as:

X is CH, N, O or S;

Y is C(R₆), N, O or S;

Z is CH, N or bond;

A is CH or N;

B₁ is N or C(R₇);

B₂ is N or C(R₈);

B₃ is N or C(R₉);

B₄ is N or C(R₁₀);

R₁ is R₁₁C(O), R₁₂S(O), R₁₃SO₂ or (C₁₋₆)alkyl optionally substituted with R₁₄;

R₂ is H, (C₁₋₃)alkyl or (C₃₋₇)cycloalkyl;

R₃ is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl; or

R_2 and R_3 form, together with the N and C atom they are attached to, a (C₃₋₇)heterocycloalkyl optionally substituted with one or more fluorine, hydroxyl, (C₁₋₃)alkyl, (C₁₋₃)alkoxy or oxo;

R_4 is H or (C₁₋₃)alkyl;

R_5 is H, halogen, cyano, (C₁₋₄)alkyl, (C₁₋₃)alkoxy, (C₃₋₆)cycloalkyl; all alkyl groups of R_5 are optionally substituted with one or more halogen; or R_5 is (C₆₋₁₀)aryl or (C₂₋₆)heterocycloalkyl;

R_6 is H or (C₁₋₃)alkyl; or R_5 and R_6 together may form a (C₃₋₇)cycloalkenyl, or (C₂₋₆)heterocycloalkenyl; each optionally substituted with (C₁₋₃)alkyl, or one or more halogen;

R_7 is H, halogen, CF₃, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R_8 is H, halogen, CF₃, (C₁₋₃)alkyl or (C₁₋₃)alkoxy; or

R_7 and R_8 together with the carbon atoms they are attached to, form (6-10C)aryl or (1-5C)heteroaryl;

R_9 is H, halogen, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R_{10} is H, halogen, (C₁₋₃)alkyl or (C₁₋₃)alkoxy;

R_{11} is independently selected from a group consisting of (C₁₋₆)alkyl, (C₂₋₆)alkenyl and (C₂₋₆)alkynyl each alkyl, alkenyl or alkynyl optionally substituted with one or more groups selected from hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl or (C₃₋₇)heterocycloalkyl, or

R_{11} is (C₁₋₃)alkyl-C(O)-S-(C₁₋₃)alkyl; or

R_{11} is (C₁₋₅)heteroaryl optionally substituted with one or more groups selected from halogen or cyano.

R_{12} and R_{13} are independently selected from a group consisting of (C₂₋₆)alkenyl or (C₂₋₆)alkynyl both optionally substituted with one or more groups selected from hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl, or (C₃₋₇)heterocycloalkyl; or

(C₁₋₅)heteroaryl optionally substituted with one or more groups selected from halogen or cyano;

R_{14} is independently selected from a group consisting of halogen, cyano or (C₂₋₆)alkenyl or (C₂₋₆)alkynyl both optionally substituted with one or more groups selected from hydroxyl, (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl, [(C₁₋₄)alkyl]amino, di[(C₁₋₄)alkyl]amino, (C₁₋₃)alkoxy, (C₃₋₇)cycloalkoxy, (C₆₋₁₀)aryl, (C₁₋₅)heteroaryl or (C₃₋₇)heterocycloalkyl;

with the proviso that

- 0 to 2 atoms of X, Y, Z can simultaneously be a heteroatom;
- when one atom selected from X, Y is O or S, then Z is a bond and the other atom selected from X, Y can not be O or S;
- when Z is C or N then Y is C(R₆) or N and X is C or N;
- 0 to 2 atoms of B₁, B₂, B₃ and B₄ are N;

with the terms used having the following meanings:

- (C₁₋₃)alkyl means a branched or unbranched alkyl group having 1-3 carbon atoms, being methyl, ethyl, propyl or isopropyl;
- (C₁₋₄)alkyl means a branched or unbranched alkyl group having 1-4 carbon atoms, being methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl and *tert*-butyl, (C₁₋₃)alkyl groups being preferred;
- (C₁₋₆)alkyl means a branched or unbranched alkyl group having 1-6 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, *n*-pentyl and *n*-hexyl. (C₁₋₅)alkyl groups are preferred, (C₁₋₄)alkyl being most preferred;
- (C₁₋₂)alkoxy means an alkoxy group having 1-2 carbon atoms, the alkyl moiety having the same meaning as previously defined;
- (C₁₋₃)alkoxy means an alkoxy group having 1-3 carbon atoms, the alkyl moiety having the same meaning as previously defined, with (C₁₋₂)alkoxy groups preferred;
- (C₂₋₃)alkenyl means an alkenyl group having 2-3 carbon atoms, such as ethenyl or 2-propenyl;
- (C₂₋₄)alkenyl means a branched or unbranched alkenyl group having 2-4 carbon atoms, such as ethenyl, 2-propenyl, isobutenyl or 2-butenyl;
- (C₂₋₆)alkenyl means a branched or unbranched alkenyl group having 2-6 carbon atoms, such as ethenyl, 2-butenyl, and *n*-pentenyl, with (C₂₋₄)alkenyl groups preferred, and (C₂₋₃)alkenyl groups even more preferred;
- (C₂₋₄)alkynyl means a branched or unbranched alkynyl group having 2-4 carbon atoms, such as ethynyl, 2-propynyl or 2-butynyl;
- (C₂₋₃)alkynyl means an alkynyl group having 2-3 carbon atoms, such as ethynyl or 2-propynyl;
- (C₂₋₆)alkynyl means a branched or unbranched alkynyl group having 2-6 carbon atoms, such as ethynyl, propynyl, *n*-butynyl, *n*-pentynyl, isopentynyl, isohexynyl or *n*-hexynyl, with (C₂₋₄)alkynyl groups preferred, and (2-3C)alkynyl groups more preferred;

- (C₃₋₆)cycloalkyl means a cycloalkyl group having 3-6 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;
- (C₃₋₇)cycloalkyl means a cycloalkyl group having 3-7 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl;
- (C₂₋₆)heterocycloalkyl means a heterocycloalkyl group having 2-6 carbon atoms, preferably 3-5 carbon atoms, and one or two heteroatoms selected from N, O and/or S, which may be attached via a heteroatom if feasible, or a carbon atom; preferred heteroatoms are N or O; preferred groups are piperidine, morpholine, pyrrolidine and piperazine; a most preferred (C₂₋₆)heterocycloalkyl is pyrrolidine; and the heterocycloalkyl group may be attached via a heteroatom if feasible;
- (C₃₋₇)heterocycloalkyl means a heterocycloalkyl group having 3-7 carbon atoms, preferably 3-5 carbon atoms, and one or two heteroatoms selected from N, O and/or S; preferred heteroatoms are N or O; preferred (C₃₋₇) heterocycloalkyl groups are azetidiny, pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl; more preferred (C₃₋₇)heterocycloalkyl groups are piperidine, morpholine and pyrrolidine; even more preferred are piperidine and pyrrolidine; and the heterocycloalkyl group may be attached via a heteroatom if feasible;
- (C₃₋₇)cycloalkoxy means a cycloalkyl group having 3-7 carbon atoms, with the same meaning as previously defined, attached via a ring carbon atom to an exocyclic oxygen atom;
- (C₆₋₁₀)aryl means an aromatic hydrocarbon group having 6-10 carbon atoms, such as phenyl, naphthyl, tetrahydronaphthyl or indenyl; the preferred (C₆₋₁₀)aryl group is phenyl;
- (C₁₋₅)heteroaryl means a substituted or unsubstituted aromatic group having 1-5 carbon atoms and 1-4 heteroatoms selected from N, O and/or S, wherein the (C₁₋₅)heteroaryl may optionally be substituted.; preferred (C₁₋₅)heteroaryl groups are tetrazolyl, imidazolyl, thiadiazolyl, pyridyl, pyrimidyl, triazinyl, thienyl or furyl, and the more preferred (C₁₋₅) heteroaryl is pyrimidyl;
- [(C₁₋₄)alkyl]amino means an amino group, monosubstituted with an alkyl group containing 1-4 carbon atoms having the same meaning as previously defined; the preferred [(C₁₋₄)alkyl]amino group is methylamino;
- di[(C₁₋₄)alkyl]amino means an amino group, disubstituted with alkyl group(s), each containing 1-4 carbon atoms and having the same meaning as previously defined; the preferred di[(C₁₋₄)alkyl]amino group is dimethylamino;

halogen means fluorine, chlorine, bromine or iodine;

(C₁₋₃)alkyl-C(O)-S-(C₁₋₃)alkyl means an alkyl-carbonyl-thio-alkyl group, each of the alkyl groups having 1 to 3 carbon atoms with the same meaning as previously defined;

(C₃₋₇)cycloalkenyl means a cycloalkenyl group having 3-7 carbon atoms, preferably 5-7 carbon atoms; preferred (C₃₋₇)cycloalkenyl groups are cyclopentenyl or cyclohexenyl; and cyclohexenyl groups are most preferred;

(C₂₋₆)heterocycloalkenyl means a heterocycloalkenyl group having 2-6 carbon atoms, preferably 3-5 carbon atoms; and 1 heteroatom selected from N, O and/or S; the preferred (C₂₋₆) heterocycloalkenyl groups are oxycyclohexenyl and azacyclohexenyl groups.

In the above definitions with multifunctional groups, the attachment point is at the last group.


When, in the definition of a substituent, is indicated that "all of the alkyl groups" of said substituent are optionally substituted, this also includes the alkyl moiety of an alkoxy group.

A circle in a ring of Formula (XIX) indicates that the ring is aromatic.

Depending on the ring formed, the nitrogen, if present in X or Y, may carry a hydrogen.

[00195] In one embodiment the invention provides a compound according to Formula XIX, wherein B₁ is C(R₇); B₂ is C(R₈); B₃ is C(R₉) and B₄ is C(R₁₀).

[00196] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XX):



Formula (XX)

or a pharmaceutically acceptable salt thereof,

wherein:

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;

Z is $\text{C}(=\text{O})$, $\text{OC}(=\text{O})$, $\text{NRC}(=\text{O})$, $\text{C}(=\text{S})$, $\text{S}(=\text{O})_x$, $\text{OS}(=\text{O})_x$ or $\text{NRS}(=\text{O})_x$, where x is 1 or 2;


R^7 and R^8 are each independently H; or R^7 and R^8 taken together form a bond;

R^6 is H; and


R is H or $\text{C}_1\text{-C}_6$ alkyl.

[00197] In an exemplary embodiment, the BTK inhibitor is ibrutinib or a pharmaceutically-acceptable salt thereof. In an exemplary embodiment, the BTK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one. In an exemplary embodiment, the BTK inhibitor is 1-[(3R)-3-[4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]piperidin-1-yl]prop-2-en-1-one. In an exemplary embodiment, the BTK inhibitor is (S)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one. In an exemplary embodiment, the BTK inhibitor has the structure of Formula (XX-A), or an enantiomer thereof, or a pharmaceutically acceptable salt, solvate, hydrate, cocrystal, or prodrug thereof.

Formula (XX-A)



[00198] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXI):



Formula (XXI)

or a pharmaceutically acceptable salt thereof,

wherein:

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$ or $NRS(=O)_x$, where x is 1 or 2;

R^7 and R^8 are each H; or R^7 and R^8 taken together form a bond;

R^6 is H; and

R is H or C_1 - C_6 alkyl.

[00199] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXII):



Formula (XXII)

or a pharmaceutically acceptable salt thereof,

wherein:

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$ or $NRS(=O)_x$, where x is 1 or 2;

R^7 and R^8 are each H; or R^7 and R^8 taken together form a bond;

R^6 is H; and

R is H or C_1 - C_6 alkyl.

[00200] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXIII):



Formula (XXIII)

or a pharmaceutically acceptable salt thereof,

wherein:

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$ or $NRS(=O)_x$, where x is 1 or 2;

R^7 and R^8 are each H; or R^7 and R^8 taken together form a bond;

R^6 is H; and

R is H or C_1 - C_6 alkyl.

[00201] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXIV):



Formula (XXIV)

or a pharmaceutically acceptable salt thereof,

wherein:

Q^1 is aryl¹, heteroaryl¹, cycloalkyl, heterocyclyl, cycloalkenyl, or heterocycloalkenyl, any of which is optionally substituted by one to five independent G^1 substituents;

R^1 is alkyl, cycloalkyl, bicycloalkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heterocyclyl, or heterobicycloalkyl, any of which is optionally substituted by one or more independent G^{11} substituents;

G^1 and G^{41} are each independently halo, oxo, $-CF_3$, $-OCF_3$, $-OR^2$, $-NR^2R^3(R^{3a})_{j1}$, $-C(O)R^2$, $-CO_2R^2$, $-CONR^2R^3$, $-NO_2$, $-CN$, $-S(O)_{j1}R^2$, $-SO_2NR^2R^3$, $NR^2(C=O)R^3$, $NR^2(C=O)OR^3$, $NR^2(C=O)NR^2R^3$, $NR^2S(O)_{j1}R^3$, $-(C=S)OR^2$, $-(C=O)SR^2$, $-NR^2(C=NR^3)NR^{2a}R^{3a}$, $-NR^2(C=NR^3)OR^{2a}$, $-NR^2(C=NR^3)SR^{3a}$, $-O(C=O)OR^2$, $-O(C=O)NR^2R^3$, $-O(C=O)SR^2$, $-S(C=O)OR^2$, $-S(C=O)NR^2R^3$, C_{0-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy C_{1-10} alkyl, C_{1-10} alkoxy C_{2-10} alkenyl, C_{1-10} alkoxy C_{2-10} alkynyl, C_{1-10} alkylthio C_{1-10} alkyl, C_{1-10} alkylthio C_{2-10} alkenyl, C_{1-10} alkylthio C_{2-10} alkynyl, cyclo C_{3-8} alkyl, cyclo C_{3-8} alkenyl, cyclo C_{3-8} alkyl C_{1-10} alkyl, cyclo C_{3-8} alkenyl C_{1-10} alkyl, cyclo C_{3-8} alkyl C_{2-10} alkenyl, cyclo C_{3-8} alkenyl C_{2-10} alkenyl, cyclo C_{3-8} alkyl C_{2-10} alkynyl, cyclo C_{3-8} alkenyl C_{2-10} alkynyl, heterocyclyl- C_{0-10} alkyl, heterocyclyl- C_{2-10} alkenyl, or heterocyclyl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, oxo, $-CF_3$, $-OCF_3$, $-OR^{222}$, $-NR^{222}R^{333}(R^{333a})_{j1a}$, $-C(O)R^{222}$, $-CO_2R^{222}$, $-CONR^{222}R^{333}$, $-NO_2$, $-CN$, $-S(O)_{j1a}R^{222}$, $-SO_2NR^{222}R^{333}$, $NR^{222}(C=O)R^{333}$, $NR^{222}(C=O)OR^{333}$, $NR^{222}(C=O)NR^{222}R^{333}$, $NR^{222}S(O)_{j1a}R^{333}$, $-(C=S)OR^{222}$, $-(C=O)SR^{222}$, $-NR^{222}(C=NR^{333})NR^{222a}R^{333a}$, $-NR^{222}(C=NR^{333})OR^{222a}$, $-NR^{222}(C=NR^{333})SR^{333a}$, $-O(C=O)OR^{222}$, $-O(C=O)NR^{222}R^{333}$, $-O(C=O)SR^{222}$, $-S(C=O)OR^{222}$, or $-S(C=O)NR^{222}R^{333}$ substituents; or $-(X^1)_n-(Y^1)_m-R^4$; or aryl- C_{0-10} alkyl, aryl- C_{2-10} alkenyl, or aryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{222}$, $-NR^{222}R^{333}(R^{333a})_{j2a}$, $-C(O)R^{222}$, $-CO_2R^{222}$, $-CONR^{222}R^{333}$, $-NO_2$, $-CN$, $-S(O)_{j2a}R^{222}$, $-SO_2NR^{222}R^{333}$, $NR^{222}(C=O)R^{333}$, $NR^{222}(C=O)OR^{333}$, $NR^{222}(C=O)NR^{222}R^{333}$, $NR^{222}S(O)_{j2a}R^{333}$, $-(C=S)OR^{222}$, $-(C=O)SR^{222}$, $-NR^{222}(C=NR^{333})NR^{222a}R^{333a}$, $-NR^{222}(C=NR^{333})OR^{222a}$, $-NR^{222}(C=NR^{333})SR^{333a}$, $-O(C=O)OR^{222}$, $-O(C=O)NR^{222}R^{333}$, $-O(C=O)SR^{222}$, $-S(C=O)OR^{222}$, or $-S(C=O)NR^{222}R^{333}$ substituents; or hetaryl- C_{0-10} alkyl, hetaryl- C_{2-10} alkenyl, or hetaryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{222}$, $-NR^{222}R^{333}(R^{333a})_{j3a}$, $-C(O)R^{222}$, $-CO_2R^{222}$, $-CONR^{222}R^{333}$, $-NO_2$, $-CN$, $-S(O)_{j3a}R^{222}$, $-SO_2NR^{222}R^{333}$,


$\text{NR}^{222}(\text{C}=\text{O})\text{R}^{333}$, $\text{NR}^{222}(\text{C}=\text{O})\text{OR}^{333}$, $\text{NR}^{222}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$, $\text{NR}^{222}\text{S}(\text{O})_{j3a}\text{R}^{333}$, $-(\text{C}=\text{S})\text{OR}^{222}$,
 $-(\text{C}=\text{O})\text{SR}^{222}$, $-\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{NR}^{222}\text{aR}^{333}\text{a}$, $-\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{OR}^{222a}$,
 $\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{SR}^{333}\text{a}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{222}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{222}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{222}$,
 or $-\text{S}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$ substituents;

G^{11} is halo, oxo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{21}$, $-\text{NR}^{21}\text{R}^{31}(\text{R}^{3a1})_{j4}$, $-\text{C}(\text{O})\text{R}^{21}$, $-\text{CO}_2\text{R}^{21}$, $-\text{CONR}^{21}\text{R}^{31}$, $-\text{NO}_2$,
 $-\text{CN}$, $-\text{S}(\text{O})_{j4}\text{R}^{21}$, $-\text{SO}_2\text{NR}^{21}\text{R}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{R}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{OR}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$,
 $\text{NR}^{21}\text{S}(\text{O})_{j4}\text{R}^{31}$, $-(\text{C}=\text{S})\text{OR}^{21}$, $-(\text{C}=\text{O})\text{SR}^{21}$, $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{NR}^{2a1}\text{R}^{3a1}$, $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{OR}^{2a1}$,
 $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{SR}^{3a1}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{21}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{21}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{21}$,
 $-\text{S}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$, $-\text{P}(\text{O})\text{OR}^{21}\text{OR}^{31}$, $\text{C}_{0-10}\text{alkyl}$, $\text{C}_{2-10}\text{alkenyl}$, $\text{C}_{2-10}\text{alkynyl}$, $\text{C}_{1-10}\text{alkoxyC}_{1-10}\text{alkyl}$,
 $\text{C}_{1-10}\text{alkoxyC}_{2-10}\text{alkenyl}$, $\text{C}_{1-10}\text{alkoxyC}_{2-10}\text{alkynyl}$, $\text{C}_{1-10}\text{alkylthioC}_{1-10}\text{alkyl}$, $\text{C}_{1-10}\text{alkylthioC}_{2-10}\text{alkenyl}$,
 $\text{C}_{1-10}\text{alkylthioC}_{2-10}\text{alkynyl}$, $\text{cycloC}_{3-8}\text{alkyl}$, $\text{cycloC}_{3-8}\text{alkenyl}$, $\text{cycloC}_{3-8}\text{alkylC}_{1-10}\text{alkyl}$, $\text{cycloC}_{3-8}\text{alkenylC}_{1-10}\text{alkyl}$,
 $\text{cycloC}_{3-8}\text{alkylC}_{2-10}\text{alkenyl}$, $\text{cycloC}_{3-8}\text{alkylC}_{2-10}\text{alkynyl}$, $\text{cycloC}_{3-8}\text{alkenylC}_{2-10}\text{alkynyl}$,
 $\text{heterocyclyl-C}_{0-10}\text{alkyl}$, $\text{heterocyclyl-C}_{2-10}\text{alkenyl}$, or $\text{heterocyclyl-C}_{2-10}\text{alkynyl}$, any of
 which is optionally substituted with one or more independent halo, oxo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j4a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$,
 $-\text{S}(\text{O})_{j4a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{OR}^{3331}$,
 $\text{NR}^{2221}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}\text{S}(\text{O})_{j4a}\text{R}^{3331}$, $-(\text{C}=\text{S})\text{OR}^{2221}$, $-(\text{C}=\text{O})\text{SR}^{2221}$,
 $\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{NR}^{222a1}\text{R}^{333a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{OR}^{222a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{SR}^{333a1}$,
 $-\text{O}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{P}(\text{O})\text{OR}^{2221}\text{OR}^{3331}$,
 or $-\text{S}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$ substituents; or $\text{aryl-C}_{0-10}\text{alkyl}$, $\text{aryl-C}_{2-10}\text{alkenyl}$, or $\text{aryl-C}_{2-10}\text{alkynyl}$, any of which is optionally substituted with one or more independent halo, $-\text{CF}_3$,
 $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j5a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$,
 $-\text{S}(\text{O})_{j5a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{OR}^{3331}$,
 $\text{NR}^{2221}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}\text{S}(\text{O})_{j5a}\text{R}^{3331}$, $-(\text{C}=\text{S})\text{OR}^{2221}$, $-(\text{C}=\text{O})\text{SR}^{2221}$,
 $\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{NR}^{222a1}\text{R}^{333a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{OR}^{222a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{SR}^{333a1}$,
 $-\text{O}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{P}(\text{O})\text{OR}^{2221}\text{R}^{3331}$, or
 $-\text{S}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$ substituents; or $\text{hetaryl-C}_{0-10}\text{alkyl}$, $\text{hetaryl-C}_{2-10}\text{alkenyl}$, or $\text{hetaryl-C}_{2-10}\text{alkynyl}$, any of which is optionally substituted with one or more independent halo, $-\text{CF}_3$,
 $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j6a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$,
 $-\text{S}(\text{O})_{j6a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{OR}^{3331}$,

$\text{NR}^{2221}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}\text{S}(\text{O})_{j6a}\text{R}^{3331}$, $-(\text{C}=\text{S})\text{OR}^{2221}$, $-(\text{C}=\text{O})\text{SR}^{2221}$, -
 $\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{NR}^{222a1}\text{R}^{333a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{OR}^{222a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{SR}^{333a1}$, -
 $\text{O}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{P}(\text{O})\text{OR}^{2221}\text{OR}^{3331}$,
 or $-\text{S}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$ substituents; or G^{11} is taken together with the carbon to which it is
 attached to form a double bond which is substituted with R^5 and G^{111} ;
 R^2 , R^{2a} , R^3 , R^{3a} , R^{222} , R^{222a} , R^{333} , R^{333a} , R^{21} , R^{2a1} , R^{31} , R^{3a1} , R^{2221} , R^{222a1} , R^{3331} , and R^{333a1} are
 each independently equal to C_{0-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy C_{1-10} alkyl, C_{1-10}
 alkoxy C_{2-10} alkenyl, C_{1-10} alkoxy C_{2-10} alkynyl, C_{1-10} alkylthio C_{1-10} alkyl, C_{1-10} alkylthio C_{2-10}
 alkenyl, C_{1-10} alkylthio C_{2-10} alkynyl, cyclo C_{3-8} alkyl, cyclo C_{3-8} alkenyl, cyclo C_{3-8} alkyl C_{1-10}
 alkyl, cyclo C_{3-8} alkenyl C_{1-10} alkyl, cyclo C_{3-8} alkyl C_{2-10} alkenyl, cyclo C_{3-8} alkenyl C_{2-10}
 alkenyl, cyclo C_{3-8} alkyl C_{2-10} alkynyl, cyclo C_{3-8} alkenyl C_{2-10} alkynyl, heterocyclyl- C_{0-10} alkyl,
 heterocyclyl- C_{2-10} alkenyl, or heterocyclyl- C_{2-10} alkynyl, any of which is optionally
 substituted by one or more G^{111} substituents; or aryl- C_{0-10} alkyl, aryl- C_{2-10} alkenyl, or aryl- C_{2-10}
 alkynyl, hetaryl- C_{0-10} alkyl, hetaryl- C_{2-10} alkenyl, or hetaryl- C_{2-10} alkynyl, any of which is
 optionally substituted by one or more G^{111} substituents; or in the case of $-\text{NR}^2\text{R}^3(\text{R}^{3a})_{j1}$ or -
 $\text{NR}^{222}\text{R}^{333}(\text{R}^{333a})_{j1a}$ or $-\text{NR}^{222}\text{R}^{333}(\text{R}^{333a})_{j2a}$ or $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j3a}$ or $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j4a}$
 or $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j5a}$ or $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j6a}$, R^2 and R^3 or R^{222} and R^{333} or R^{2221} and
 R^{3331} taken together with the nitrogen atom to which they are attached form a 3-10 membered
 saturated ring, unsaturated ring, heterocyclic saturated ring, or heterocyclic unsaturated ring,
 wherein said ring is optionally substituted by one or more G^{111} substituents;
 X^1 and Y^1 are each independently -O-, $-\text{NR}^7$ -, $-\text{S}(\text{O})_{j7}$ -, $-\text{CR}^5\text{R}^6$ -, $-\text{N}(\text{C}(\text{O})\text{OR}^7)$ -, $-\text{N}(\text{C}(\text{O})\text{R}^7)$ -,
 $-\text{N}(\text{SO}_2\text{R}^7)$ -, $-\text{CH}_2\text{O}$ -, $-\text{CH}_2\text{S}$ -, $-\text{CH}_2\text{N}(\text{R}^7)$ -, $-\text{CH}(\text{NR}^7)$ -, $-\text{CH}_2\text{N}(\text{C}(\text{O})\text{R}^7)$ -, $-\text{CH}_2\text{N}(\text{C}(\text{O})\text{OR}^7)$ -,
 $-\text{CH}_2\text{N}(\text{SO}_2\text{R}^7)$ -, $-\text{CH}(\text{NHR}^7)$ -, $-\text{CH}(\text{NHC}(\text{O})\text{R}^7)$ -, $-\text{CH}(\text{NHSO}_2\text{R}^7)$ -, $-\text{CH}(\text{NHC}(\text{O})\text{OR}^7)$ -,
 $-\text{CH}(\text{OC}(\text{O})\text{R}^7)$ -, $-\text{CH}(\text{OC}(\text{O})\text{NHR}^7)$ -, $-\text{CH}=\text{CH}$ -, $-\text{C.ident.C}$ -, $-\text{C}(=\text{NOR}^7)$ -, $-\text{C}(\text{O})$ -, -
 $\text{CH}(\text{OR}^7)$ -, $-\text{C}(\text{O})\text{N}(\text{R}^7)$ -, $-\text{N}(\text{R}^7)\text{C}(\text{O})$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})_2$ -, $-\text{OC}(\text{O})\text{N}(\text{R}^7)$ -, -
 $\text{N}(\text{R}^7)\text{C}(\text{O})\text{N}(\text{R}^7)$ -, $-\text{NR}^7\text{C}(\text{O})\text{O}$ -, $-\text{S}(\text{O})\text{N}(\text{R}^7)$ -, $-\text{S}(\text{O})_2\text{N}(\text{R}^7)$ -, $-\text{N}(\text{C}(\text{O})\text{R}^7)\text{S}(\text{O})$ -, -
 $\text{N}(\text{C}(\text{O})\text{R}^7)\text{S}(\text{O})_2$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})\text{N}(\text{R}^7)$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{N}(\text{R}^7)$ -, $-\text{C}(\text{O})\text{N}(\text{R}^7)\text{C}(\text{O})$ -, -
 $\text{S}(\text{O})\text{N}(\text{R}^7)\text{C}(\text{O})$ -, $-\text{S}(\text{O})_2\text{N}(\text{R}^7)\text{C}(\text{O})$ -, $-\text{OS}(\text{O})\text{N}(\text{R}^7)$ -, $-\text{OS}(\text{O})_2\text{N}(\text{R}^7)$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})\text{O}$ -, -
 $\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{O}$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})\text{C}(\text{O})$ -, $-\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{C}(\text{O})$ -, $-\text{SON}(\text{C}(\text{O})\text{R}^7)$ -, $-\text{SO}_2\text{N}(\text{C}(\text{O})\text{R}^7)$ -, -
 $\text{N}(\text{R}^7)\text{SON}(\text{R}^7)$ -, $-\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)$ -, $-\text{C}(\text{O})\text{O}$ -, $-\text{N}(\text{R}^7)\text{P}(\text{OR}^8)\text{O}$ -, $-\text{N}(\text{R}^7)\text{P}(\text{OR}^8)$ -, -
 $\text{N}(\text{R}^7)\text{P}(\text{O})(\text{OR}^8)\text{O}$ -, $-\text{N}(\text{R}^7)\text{P}(\text{O})(\text{OR}^8)$ -, $-\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)\text{O}$ -, $-\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)$ -, -

$\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{O})(\text{OR}^8)\text{O}-$, $-\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})_2-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{OR}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{SO}_2\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{O}-$, $-\text{CH}(\text{R}^7)\text{S}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{OR}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{SO}_2\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{C}(=\text{NOR}^7)-$, $-\text{CH}(\text{R}^7)\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{CH}(\text{OR}^7)-$, $-\text{CH}(\text{R}^7)\text{C}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})_2-$, $-\text{CH}(\text{R}^7)\text{OC}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{C}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{NR}^7\text{C}(\text{O})\text{O}-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})_2\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{S}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{S}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{C}(\text{O})\text{N}(\text{R}^7)\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})\text{N}(\text{R}^7)\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{S}(\text{O})_2\text{N}(\text{R}^7)\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{OS}(\text{O})\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{OS}(\text{O})_2\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{S}(\text{O})_2\text{C}(\text{O})-$, $-\text{CH}(\text{R}^7)\text{SON}(\text{C}(\text{O})\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{SO}_2\text{N}(\text{C}(\text{O})\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{SON}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)-$, $-\text{CH}(\text{R}^7)\text{C}(\text{O})\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{P}(\text{OR}^8)\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{P}(\text{OR}^8)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{P}(\text{O})(\text{OR}^8)\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{R}^7)\text{P}(\text{O})(\text{OR}^8)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)\text{O}-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)-$, $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{O})(\text{OR}^8)\text{O}-$, or $-\text{CH}(\text{R}^7)\text{N}(\text{C}(\text{O})\text{R}^7)\text{P}(\text{OR}^8)-$;

or X^1 and Y^1 are each independently represented by one of the following structural formulas:



R^{10} , taken together with the phosphinamide or phosphonamide, is a 5-, 6-, or 7-membered aryl, heteroaryl or heterocyclyl ring system;

R^5 , R^6 , and G^{111} are each independently a C_{0-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy C_{1-10} alkyl, C_{1-10} alkoxy C_{2-10} alkenyl, C_{1-10} alkoxy C_{2-10} alkynyl, C_{1-10} alkylthio C_{1-10} alkyl, C_{1-10} alkylthio C_{2-10} alkenyl, C_{1-10} alkylthio C_{2-10} alkynyl, cyclo C_{3-8} alkyl, cyclo C_{3-8} alkenyl, cyclo C_{3-8} alkyl C_{1-10} alkyl, cyclo C_{3-8} alkenyl C_{1-10} alkyl, cyclo C_{3-8} alkyl C_{2-10} alkenyl, cyclo C_{3-8} alkenyl C_{2-10} alkenyl, cyclo C_{3-8} alkyl C_{2-10} alkynyl, cyclo C_{3-8} alkenyl C_{2-10} alkynyl, heterocyclyl- C_{0-10} alkyl, heterocyclyl- C_{2-10} alkenyl, or heterocyclyl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{77}$, $-NR^{77}R^{87}$, $-C(O)R^{77}$, $-CO_2R^{77}$, $-CONR^{77}R^{87}$, $-NO_2$, $-CN$, $-S(O)_{j5a}R^{77}$, $-SO_2NR^{77}R^{87}$, $NR^{77}(C=O)R^{87}$, $NR^{77}(C=O)OR^{87}$, $NR^{77}(C=O)NR^{78}R^{87}$, $NR^{77}S(O)_{j5a}R^{87}$, $-(C=S)OR^{77}$, $-(C=O)SR^{77}$, $-NR^{77}(C=NR^{87})NR^{78}R^{88}$, $-NR^{77}(C=NR^{87})OR^{78}$, $-NR^{77}(C=NR^{87})SR^{78}$, $-O(C=O)OR^{77}$, $-O(C=O)NR^{77}R^{87}$, $-O(C=O)SR^{77}$, $-S(C=O)OR^{77}$, $-P(O)OR^{77}OR^{87}$, or $-S(C=O)NR^{77}R^{87}$ substituents; or aryl- C_{0-10} alkyl, aryl- C_{2-10} alkenyl, or aryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{77}$, $-NR^{77}R^{87}$, $-C(O)R^{77}$, $-CO_2R^{77}$, $-CONR^{77}R^{87}$, $-NO_2$, $-CN$, $-S(O)_{j5a}R^{77}$, $-SO_2NR^{77}R^{87}$, $NR^{77}(C=O)R^{87}$, $NR^{77}(C=O)OR^{87}$, $NR^{77}(C=O)NR^{78}R^{87}$, $NR^{77}S(O)_{j5a}R^{87}$, $-(C=S)OR^{77}$, $-(C=O)SR^{77}$, $-NR^{77}(C=NR^{87})NR^{78}R^{88}$, $-NR^{77}(C=NR^{87})OR^{78}$, $-NR^{77}(C=NR^{87})SR^{78}$, $-O(C=O)OR^{77}$, $-O(C=O)NR^{77}R^{87}$, $-O(C=O)SR^{77}$, $-S(C=O)OR^{77}$, $-P(O)OR^{77}R^{87}$, or $-S(C=O)NR^{77}R^{87}$ substituents; or hetaryl- C_{0-10} alkyl, hetaryl- C_{2-10} alkenyl, or hetaryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{77}$, $-NR^{77}R^{87}$, $-C(O)R^{77}$, $-CO_2R^{77}$, $-CONR^{77}R^{87}$, $-NO_2$, $-CN$, $-S(O)_{j5a}R^{77}$, $-SO_2NR^{77}R^{87}$, $NR^{77}(C=O)R^{87}$, $NR^{77}(C=O)OR^{87}$, $NR^{77}(C=O)NR^{78}R^{87}$, $NR^{77}S(O)_{j5a}R^{87}$, $-(C=S)OR^{77}$, $-(C=O)SR^{77}$, $-NR^{77}(C=NR^{87})NR^{78}R^{88}$, $-NR^{77}(C=NR^{87})OR^{78}$, $-NR^{77}(C=NR^{87})SR^{78}$, $-O(C=O)OR^{77}$, $-O(C=O)NR^{77}R^{87}$, $-O(C=O)SR^{77}$, $-S(C=O)OR^{77}$, $-P(O)OR^{77}OR^{87}$, or $-S(C=O)NR^{77}R^{87}$ substituents; or R^5 with R^6 taken together with the respective carbon atom to which they are attached, form a 3-10 membered saturated or unsaturated ring, wherein said ring is optionally substituted with R^{69} ; or R^5 with R^6 taken together with the respective carbon atom to which they are attached, form a 3-10 membered saturated or unsaturated heterocyclic ring, wherein said ring is optionally substituted with R^{69} ;

R^7 and R^8 are each independently H, acyl, alkyl, alkenyl, aryl, heteroaryl, heterocyclyl or cycloalkyl, any of which is optionally substituted with one or more G^{111} substituents;

R^4 is H, alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, cycloalkenyl, or heterocycloalkenyl, any of which is optionally substituted by one or more G^{41} substituents;


R^{69} is equal to halo, $-OR^{78}$, $-SH$, $-NR^{78}R^{88}$, $-CO_2R^{78}$, $-CONR^{78}R^{88}$, $-NO_2$, $-CN$, $-S(O)_jR^{78}$, $-SO_2NR^{78}R^{88}$, $C_{0-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, $C_{1-10}alkoxyC_{1-10}alkyl$, $C_{1-10}alkoxyC_{2-10}alkenyl$, $C_{1-10}alkoxyC_{2-10}alkynyl$, $C_{1-10}alkylthioC_{1-10}alkyl$, $C_{1-10}alkylthioC_{2-10}alkenyl$, $C_{1-10}alkylthioC_{2-10}alkynyl$, $cycloC_{3-8}alkyl$, $cycloC_{3-8}alkenyl$, $cycloC_{3-8}alkylC_{1-10}alkyl$, $cycloC_{3-8}alkenylC_{1-10}alkyl$, $cycloC_{3-8}alkylC_{2-10}alkenyl$, $cycloC_{3-8}alkenylC_{2-10}alkenyl$, $cycloC_{3-8}alkylC_{2-10}alkynyl$, $cycloC_{3-8}alkenylC_{2-10}alkynyl$, heterocyclyl- $C_{0-10}alkyl$, heterocyclyl- $C_{2-10}alkenyl$, or heterocyclyl- $C_{2-10}alkynyl$, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-OR^{778}$, $-SO_2NR^{778}R^{888}$, or $-NR^{778}R^{888}$ substituents; or aryl- $C_{0-10}alkyl$, aryl- $C_{2-10}alkenyl$, or aryl- $C_{2-10}alkynyl$, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-OR^{778}$, $C_{1-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, halo $C_{1-10}alkyl$, halo $C_{2-10}alkenyl$, halo $C_{2-10}alkynyl$, $-COOH$, $C_{1-4}alkoxycarbonyl$, $-CONR^{778}R^{888}$, $-SO_2NR^{778}R^{888}$, or $-NR^{778}R^{888}$ substituents; or hetaryl- $C_{0-10}alkyl$, hetaryl- $C_{2-10}alkenyl$, or hetaryl- $C_{2-10}alkynyl$, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-OR^{778}$, $C_{1-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, halo $C_{1-10}alkyl$, halo $C_{2-10}alkenyl$, halo $C_{2-10}alkynyl$, $-COOH$, $C_{1-4}alkoxycarbonyl$, $-CONR^{778}R^{888}$, $-SO_2NR^{778}R^{888}$, or $-NR^{778}R^{888}$ substituents; or mono($C_{1-6}alkyl$)amino $C_{1-6}alkyl$, di($C_{1-6}alkyl$)amino $C_{1-6}alkyl$, mono(aryl)amino $C_{1-6}alkyl$, di(aryl)amino $C_{1-6}alkyl$, or $-N(C_{1-6}alkyl)-C_{1-6}alkyl-aryl$, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-OR^{778}$, $C_{1-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, halo $C_{1-10}alkyl$, halo $C_{2-10}alkenyl$, halo $C_{2-10}alkynyl$, $-COOH$, $C_{1-4}alkoxycarbonyl$, $-CONR^{778}R^{888}$, $-SO_2NR^{778}R^{888}$, or $-NR^{778}R^{888}$ substituents; or in the case of $-NR^{78}R^{88}$, R^{78} and R^{88} taken together with the nitrogen atom to which they are attached form a 3-10 membered saturated ring, unsaturated ring, heterocyclic saturated ring, or heterocyclic unsaturated ring, wherein said ring is optionally substituted with one or more independent halo, cyano, hydroxy, nitro, $C_{1-10}alkoxy$, $-SO_2NR^{778}R^{888}$, or $-NR^{778}R^{888}$ substituents;

R^{77} , R^{78} , R^{87} , R^{88} , R^{778} , and R^{888} are each independently $C_{0-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, $C_{1-10}alkoxyC_{1-10}alkyl$, $C_{1-10}alkoxyC_{2-10}alkenyl$, $C_{1-10}alkoxyC_{2-10}alkynyl$, $C_{1-10}alkylthioC_{1-10}alkyl$, $C_{1-10}alkylthioC_{2-10}alkenyl$, $C_{1-10}alkylthioC_{2-10}alkynyl$, $cycloC_{3-8}alkyl$, $cycloC_{3-8}alkenyl$, $cycloC_{3-8}alkylC_{1-10}alkyl$, $cycloC_{3-8}alkenylC_{1-10}alkyl$, $cycloC_{3-8}alkylC_{2-10}alkenyl$,


cycloC₃₋₈alkenylC₂₋₁₀alkenyl, cycloC₃₋₈alkylC₂₋₁₀alkynyl, cycloC₃₋₈alkenylC₂₋₁₀alkynyl, heterocyclyl-C₀₋₁₀alkyl, heterocyclyl-C₂₋₁₀alkenyl, heterocyclyl-C₂₋₁₀alkynyl, C₁₋₁₀alkylcarbonyl, C₂₋₁₀alkenylcarbonyl, C₂₋₁₀alkynylcarbonyl, C₁₋₁₀alkoxycarbonyl, C₁₋₁₀alkoxycarbonylC₁₋₁₀alkyl, monoC₁₋₆alkylaminocarbonyl, diC₁₋₆alkylaminocarbonyl, mono(aryl)aminocarbonyl, di(aryl)aminocarbonyl, or C₁₋₁₀alkyl(aryl)aminocarbonyl, any of which is optionally substituted with one or more independent halo, cyano, hydroxy, nitro, C₁₋₁₀alkoxy, -SO₂N(C₀₋₄alkyl)(C₀₋₄alkyl), or -N(C₀₋₄alkyl)(C₀₋₄alkyl) substituents; or aryl-C₀₋₁₀alkyl, aryl-C₂₋₁₀alkenyl, or aryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -O(C₀₋₄alkyl), C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CON(C₀₋₄alkyl)(C₀₋₁₀alkyl), -SO₂N(C₀₋₄alkyl)(C₀₋₄alkyl), or -N(C₀₋₄alkyl)(C₀₋₄alkyl) substituents; or hetaryl-C₀₋₁₀alkyl, hetaryl-C₂₋₁₀alkenyl, or hetaryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -O(C₀₋₄alkyl), C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CON(C₀₋₄alkyl)(C₀₋₄alkyl), -SO₂N(C₀₋₄alkyl)(C₀₋₄alkyl), or -N(C₀₋₄alkyl)(C₀₋₄alkyl) substituents; or mono(C₁₋₆alkyl)aminoC₁₋₆alkyl, di(C₁₋₆alkyl)aminoC₁₋₆alkyl, mono(aryl)aminoC₁₋₆alkyl, di(aryl)aminoC₁₋₆alkyl, or -N(C₁₋₆alkyl)-C₁₋₆alkyl-aryl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -O(C₀₋₄alkyl), C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CON(C₀₋₄alkyl)(C₀₋₄alkyl), -SO₂N(C₀₋₄alkyl)(C₀₋₄alkyl), or -N(C₀₋₄alkyl)(C₀₋₄alkyl) substituents; and

n, m, j1, j1a, j2a, j3a, j4, j4a, j5a, j6a, j7, and j8 are each independently equal to 0, 1, or 2.

[00202] In an exemplary embodiment, the BTK inhibitor is a compound selected from the structures disclosed in U.S. Patent Nos. 8,450,335 and 8,609,679, and U.S. Patent Application Publication Nos. 2010/0029610 A1, 2012/0077832 A1, 2013/0065879 A1, 2013/0072469 A1, and 2013/0165462 A1. In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXV) or Formula (XXVI):



Formula (XXV)



Formula (XXVI)

or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted group selected from phenyl, a 3-7 membered saturated or partially unsaturated carbocyclic ring, an 8-10 membered bicyclic saturated, partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 4-7 membered saturated or partially unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 7-10 membered bicyclic saturated or partially unsaturated heterocyclic ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Ring B is an optionally substituted group selected from phenyl, a 3-7 membered saturated or partially unsaturated carbocyclic ring, an 8-10 membered bicyclic saturated, partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 4-7 membered saturated or partially unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 7-10 membered bicyclic saturated or partially unsaturated

heterocyclic ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

R^1 is a warhead group;

R^y is hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, C_{1-4} aliphatic, C_{1-4} haloaliphatic, $-\text{OR}$, $-\text{C}(\text{O})\text{R}$, or $-\text{C}(\text{O})\text{N}(\text{R})_2$;

each R group is independently hydrogen or an optionally substituted group selected from C_{1-6} aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

W^1 and W^2 are each independently a covalent bond or a bivalent C_{1-3} alkylene chain wherein one methylene unit of W^1 or W^2 is optionally replaced by $-\text{NR}^2-$, $-\text{N}(\text{R}^2)\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^2)-$, $-\text{N}(\text{R}^2)\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^2)-$, $-\text{O}-$, $-\text{C}(\text{O})-$, $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{SO}_2-$;

R^2 is hydrogen, optionally substituted C_{1-6} aliphatic, or $-\text{C}(\text{O})\text{R}$, or:

R^2 and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated, partially unsaturated, or aromatic fused ring, or:

R^2 and R^y are taken together with their intervening atoms to form a 4-7 membered partially unsaturated or aromatic fused ring;

m and p are independently 0-4; and

R^x and R^v are independently selected from $-\text{R}$, halogen, $-\text{OR}$, $-\text{O}(\text{CH}_2)_q\text{OR}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{SO}_2\text{R}$, $-\text{SO}_2\text{N}(\text{R})_2$, $-\text{SOR}$, $-\text{C}(\text{O})\text{R}$, $-\text{CO}_2\text{R}$, $-\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{NRC}(\text{O})\text{R}$, $-\text{NRC}(\text{O})\text{NR}_2$, $-\text{NRSO}_2\text{R}$, or $-\text{N}(\text{R})_2$, wherein q is 1-4; or:

R^x and R^1 when concurrently present on Ring B are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, $-\text{CN}$, or C_{1-6} aliphatic; or

R^v and R^1 when concurrently present on Ring A are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with

a warhead group and 0-3 groups independently selected from oxo, halogen, —CN, or C₁₋₆ aliphatic.

[00203] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXV) or Formula (XXVI), wherein:

Ring A is an optionally substituted group selected from phenyl, a 3-7 membered saturated or partially unsaturated carbocyclic ring, an 8-10 membered bicyclic saturated, partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 4-7 membered saturated or partially unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 7-10 membered bicyclic saturated or partially unsaturated heterocyclic ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Ring B is an optionally substituted group selected from phenyl, a 3-7 membered saturated or partially unsaturated carbocyclic ring, an 8-10 membered bicyclic saturated, partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 4-7 membered saturated or partially unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 7-10 membered bicyclic saturated or partially unsaturated heterocyclic ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

R¹ is -L-Y, wherein:

L is a covalent bond or a bivalent C₁₋₈ saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one, two, or three methylene units of L are optionally and independently replaced by cyclopropylene, —NR—, —N(R)C(O)—, —C(O)N(R)—, —N(R)SO₂—, —SO₂N(R)—, —O—, —C(O)—, —OC(O)—, —C(O)O—, —S—, —SO—, —SO₂—, —C(=S)—, —C(=NR)—, —N=N—, or —C(=N₂)—;

Y is hydrogen, C₁₋₆ aliphatic optionally substituted with oxo, halogen, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with at 1-4 groups independently selected from -Q-Z, oxo, NO₂, halogen, CN, or C₁₋₆ aliphatic, wherein:

Q is a covalent bond or a bivalent C₁₋₆ saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by —NR—, —S—, —O—, —C(O)—, —SO—, or —SO₂—; and

Z is hydrogen or C₁₋₆ aliphatic optionally substituted with oxo, halogen, or CN;

R^y is hydrogen, halogen, —CN, —CF₃, C₁₋₄ aliphatic, C₁₋₄ haloaliphatic, —OR, —C(O)R, or —C(O)N(R)₂;

each R group is independently hydrogen or an optionally substituted group selected from C₁₋₆ aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

W¹ and W² are each independently a covalent bond or a bivalent C₁₋₃ alkylene chain wherein one methylene unit of W¹ or W² is optionally replaced by —NR²—, —N(R²)C(O)—, —C(O)N(R²)—, —N(R²)SO₂—, —SO₂N(R²)—, —O—, —C(O)—, —OC(O)—, —C(O)O—, —S—, —SO— or —SO₂—;

R² is hydrogen, optionally substituted C₁₋₆ aliphatic, or —C(O)R, or:

R² and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered partially unsaturated or aromatic fused ring; or

R² and R^y are taken together with their intervening atoms to form a 4-6 membered saturated, partially unsaturated, or aromatic fused ring;

m and p are independently 0-4; and

R^x and R^v are independently selected from $-R$, halogen, $-OR$, $-O(CH_2)_qOR$, $-CN$, $-NO_2$, $-SO_2R$, $-SO_2N(R)_2$, $-SOR$, $-C(O)R$, $-CO_2R$, $-C(O)N(R)_2$, $-NRC(O)R$, $-NRC(O)NR_2$, $-NRSO_2R$, or $-N(R)_2$, or:


R^x and R^1 when concurrently present on Ring B are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, $-CN$, or C_{1-6} aliphatic; or

R^v and R^1 when concurrently present on Ring A are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, $-CN$, or C_{1-6} aliphatic.

As defined generally above, Ring A is an optionally substituted group selected from phenyl, a 3-7 membered saturated or partially unsaturated carbocyclic ring, an 8-10 membered bicyclic saturated, partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 4-7 membered saturated or partially unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, a 7-10 membered bicyclic saturated or partially unsaturated heterocyclic ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, Ring A is an optionally substituted phenyl group. In some embodiments, Ring A is an optionally substituted naphthyl ring or an optionally substituted bicyclic 8-10 membered heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain other embodiments, Ring A is an optionally substituted 3-7 membered carbocyclic ring. In yet other embodiments, Ring A is an optionally substituted 4-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

In certain embodiments, Ring A is substituted as defined herein. In some embodiments, Ring A is substituted with one, two, or three groups independently selected from halogen, R^o , or $-(CH_2)_{0-4}OR^o$, or $-O(CH_2)_{0-4}R^o$, wherein each R^o is an alkyl or aryl group. Exemplary substituents on Ring A include Br, I, Cl, methyl, $-CF_3$, $-C\equiv CH$, $-OCH_2phenyl$, $-OCH_2(fluorophenyl)$, or $-OCH_2pyridyl$.

[00204] In an exemplary embodiment, the BTK inhibitor is CC-292. In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXVII):




which is *N*-(3-((5-fluoro-2-((4-(2-methoxyethoxy)phenyl)amino)pyrimidin-4-yl)amino)phenyl)acrylamide, or a pharmaceutically acceptable salt thereof, in an exemplary embodiment a hydrochloride salt or besylate salt thereof. The preparation of this compound is described in U.S. Patent Application Publication No. 2010/0029610 A1 at Example 20. The preparation of the hydrochloride salt or besylate salt of this compound is described in U.S. Patent Application Publication No. 2012/0077832 A1.

[00205] In an exemplary embodiment, the BTK inhibitor is *N*-(3-((5-fluoro-2-((4-(2-methoxyethoxy)phenyl)amino)pyrimidin-4-yl)amino)phenyl)acrylamide or a pharmaceutically acceptable salt thereof, or a hydrochloride salt thereof. The preparation of this compound is described in U.S. Patent Application Publication No. 2012/0077832 A1.

[00206] In an exemplary embodiment, the BTK inhibitor is (*N*-(3-(5-fluoro-2-(4-(2-methoxyethoxy)phenylamino)pyrimidin-4-ylamino)phenyl)acrylamide), or a pharmaceutically acceptable salt thereof, or a besylate salt thereof. The preparation of this compound is described in U.S. Patent Application Publication No. 2010/0029610 A1 at Example 20. The preparation of its besylate salt is described in U.S. Patent Application Publication No. 2012/0077832 A1.

[00207] In an exemplary embodiment, the BTK inhibitor is a compound of Formula (XXVIII):



or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the BTK inhibitor is the hydrochloride salt of a compound of Formula (XXVIII). The preparation of this compound is described in International Patent Application Publication No. WO 2013/081016 A1. In an exemplary embodiment, the BTK inhibitor is 6-amino-9-(1-(but-2-ynoyl)pyrrolidin-3-yl)-7-(4-phenoxyphenyl)-7,9-dihydro-8*H*-purin-8-one or a pharmaceutically acceptable salt thereof, or a hydrochloride salt thereof.

[00208] In an exemplary embodiment, the BTK inhibitor is 6-amino-9-[(3*R*)-1-(2-butynoyl)-3-pyrrolidinyl]-7-(4-phenoxyphenyl)-7,9-dihydro-8*H*-purin-8-one or a pharmaceutically acceptable salt thereof, or a hydrochloride salt thereof. The preparation of this compound is described in International Patent Application Publication No. WO 2013/081016 A1.

[00209] In an exemplary embodiment, the BTK inhibitor is 6-amino-9-[(3*S*)-1-(2-butynoyl)-3-pyrrolidinyl]-7-(4-phenoxyphenyl)-7,9-dihydro-8*H*-purin-8-one or a pharmaceutically acceptable salt thereof, or a hydrochloride salt thereof. The preparation of this compound is described in International Patent Application Publication No. WO 2013/081016 A1.

[00210] BTK inhibitors suitable for use in the described combination with a PI3K inhibitor, the PI3K inhibitor in selected embodiments being selected from the group consisting of a PI3K- γ

inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor also include, but are not limited to, those described in, for example, International Patent Application Publication Nos. WO 2013/010868; WO 2012/158843; WO 2012/135944; WO 2012/135937; U.S. Patent Application Publication No. 2011/0177011; and U.S. Patent Nos. 8,501,751; 8,476,284; 8,008,309; 7,960,396; 7,825,118; 7,732,454; 7,514,444; 7,459,554; 7,405,295; and 7,393,848.

Pharmaceutical Compositions

[00211] In one embodiment, the invention provides a pharmaceutical composition comprising a combination of a PI3K inhibitor and a BTK inhibitor. In selected embodiments, the PI3K inhibitor is selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor. Said pharmaceutical composition typically also comprises at least one pharmaceutically acceptable excipient.

[00212] Said pharmaceutical composition is in one embodiment for use in the treatment of the diseases and conditions described below. In particular, it is for use in the treatment of hyperproliferative disorders.

[00213] In selected embodiments, the invention provides a pharmaceutical composition comprising a combination of a PI3K inhibitor and a BTK inhibitor for treating solid tumor cancers, lymphomas and leukemia. In selected embodiments, the PI3K inhibitor is selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor.

[00214] In selected embodiments, the invention provides a pharmaceutical composition comprising a combination of a PI3K inhibitor, including a PI3K inhibitor selected from the group consisting of a PI3K- γ , a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, and a BTK inhibitor for the treatment of disorders such as hyperproliferative disorder including but not limited to cancer such as acute myeloid leukemia, thymus, brain, lung, squamous cell, skin, eye, retinoblastoma, intraocular melanoma, oral cavity and oropharyngeal, bladder, gastric, stomach, pancreatic, bladder, breast, cervical, head, neck, renal, kidney, liver, ovarian, prostate, colorectal, esophageal, testicular, gynecological, thyroid, CNS, PNS, AIDS-related (*e.g.*, lymphoma and Kaposi's sarcoma) or viral-induced cancer. In some embodiments, said pharmaceutical composition is for the treatment of a non-cancerous hyperproliferative disorder such as benign

hyperplasia of the skin (*e.g.*, psoriasis), restenosis, or prostate (*e.g.*, benign prostatic hypertrophy (BPH)).

[00215] The invention further provides a pharmaceutical composition comprising a combination of a PI3K inhibitor and a BTK inhibitor for the prevention of blastocyte implantation in a mammal.

[00216] The invention also provides a pharmaceutical composition comprising a combination of a PI3K inhibitor and a BTK inhibitor for treating a disease related to vasculogenesis or angiogenesis in a mammal which can manifest as tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, inflammatory bowel disease, atherosclerosis, skin diseases such as psoriasis, eczema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

[00217] The pharmaceutical compositions are typically formulated to provide a therapeutically effective amount of a combination of a PI3K inhibitor, including a PI3K inhibitor selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, and BTK inhibitor as the active ingredients, or a pharmaceutically acceptable salt, ester, prodrug, solvate, or hydrate thereof. Where desired, the pharmaceutical compositions contain a pharmaceutically acceptable salt and/or coordination complex thereof, and one or more pharmaceutically acceptable excipients, carriers, including inert solid diluents and fillers, diluents, including sterile aqueous solution and various organic solvents, permeation enhancers, solubilizers and adjuvants.

[00218] The pharmaceutical compositions are administered as a combination of a PI3K inhibitor, including a PI3K inhibitor selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor, and a BTK inhibitor. Where desired, other agent(s) may be mixed into a preparation or both components may be formulated into separate preparations for use in combination separately or at the same time. A kit containing both components formulated into separate preparations for said use is also provided by the invention.

[00219] The weight ratio of the PI3K inhibitor to the BTK inhibitor in the combination is typically with the range from 0.01 to 100, preferably from 2.5:1 to 1:2.5, and more preferably about 1:1.

[00220] In selected embodiments, the concentration of each of the PI3K and BTK inhibitors provided in the pharmaceutical compositions of the invention is independently less than, for example, 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v or v/v of each of the BTK or PI3K inhibitors.

[00221] In selected embodiments, the concentration of each of the PI3K and BTK inhibitors provided in the pharmaceutical compositions of the invention is independently greater than 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19.75%, 19.50%, 19.25% 19%, 18.75%, 18.50%, 18.25% 18%, 17.75%, 17.50%, 17.25% 17%, 16.75%, 16.50%, 16.25% 16%, 15.75%, 15.50%, 15.25% 15%, 14.75%, 14.50%, 14.25% 14%, 13.75%, 13.50%, 13.25% 13%, 12.75%, 12.50%, 12.25% 12%, 11.75%, 11.50%, 11.25% 11%, 10.75%, 10.50%, 10.25% 10%, 9.75%, 9.50%, 9.25% 9%, 8.75%, 8.50%, 8.25% 8%, 7.75%, 7.50%, 7.25% 7%, 6.75%, 6.50%, 6.25% 6%, 5.75%, 5.50%, 5.25% 5%, 4.75%, 4.50%, 4.25% 4%, 3.75%, 3.50%, 3.25% 3%, 2.75%, 2.50%, 2.25%, 2%, 1.75%, 1.50%, 1.25% 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v, or v/v of each of the BTK or PI3K inhibitors.

[00222] In selected embodiments, the concentration of each of the PI3K and BTK inhibitors of the invention is independently in the range from about 0.0001% to about 50%, about 0.001% to about 40%, about 0.01% to about 30%, about 0.02% to about 29%, about 0.03% to about 28%, about 0.04% to about 27%, about 0.05% to about 26%, about 0.06% to about 25%, about 0.07% to about 24%, about 0.08% to about 23%, about 0.09% to about 22%, about 0.1% to about 21%, about 0.2% to about 20%, about 0.3% to about 19%, about 0.4% to about 18%, about 0.5% to about 17%, about 0.6% to about 16%, about 0.7% to about 15%, about 0.8% to about 14%, about 0.9% to about 12% or about 1% to about 10% w/w, w/v or v/v. v/v of each of the BTK or PI3K inhibitors.

[00223] In selected embodiments, the concentration of each of the PI3K and BTK inhibitors of the invention is independently in the range from about 0.001% to about 10%, about 0.01% to about 5%, about 0.02% to about 4.5%, about 0.03% to about 4%, about 0.04% to about 3.5%, about 0.05% to about 3%, about 0.06% to about 2.5%, about 0.07% to about 2%, about 0.08% to about 1.5%, about 0.09% to about 1%, about 0.1% to about 0.9% w/w, w/v or v/v of each of the BTK or PI3K inhibitors.

[00224] In selected embodiments, the amount of each of the PI3K and BTK inhibitors of the invention is independently equal to or less than 10 g, 9.5 g, 9.0 g, 8.5 g, 8.0 g, 7.5 g, 7.0 g, 6.5 g, 6.0 g, 5.5 g, 5.0 g, 4.5 g, 4.0 g, 3.5 g, 3.0 g, 2.5 g, 2.0 g, 1.5 g, 1.0 g, 0.95 g, 0.9 g, 0.85 g, 0.8 g, 0.75 g, 0.7 g, 0.65 g, 0.6 g, 0.55 g, 0.5 g, 0.45 g, 0.4 g, 0.35 g, 0.3 g, 0.25 g, 0.2 g, 0.15 g, 0.1 g, 0.09 g, 0.08 g, 0.07 g, 0.06 g, 0.05 g, 0.04 g, 0.03 g, 0.02 g, 0.01 g, 0.009 g, 0.008 g, 0.007 g, 0.006 g, 0.005 g, 0.004 g, 0.003 g, 0.002 g, 0.001 g, 0.0009 g, 0.0008 g, 0.0007 g, 0.0006 g, 0.0005 g, 0.0004 g, 0.0003 g, 0.0002 g or 0.0001 g.

[00225] In selected embodiments, the amount of each of the PI3K and BTK inhibitors of the invention is independently more than 0.0001 g, 0.0002 g, 0.0003 g, 0.0004 g, 0.0005 g, 0.0006 g, 0.0007 g, 0.0008 g, 0.0009 g, 0.001 g, 0.0015 g, 0.002 g, 0.0025 g, 0.003 g, 0.0035 g, 0.004 g, 0.0045 g, 0.005 g, 0.0055 g, 0.006 g, 0.0065 g, 0.007 g, 0.0075 g, 0.008 g, 0.0085 g, 0.009 g, 0.0095 g, 0.01 g, 0.015 g, 0.02 g, 0.025 g, 0.03 g, 0.035 g, 0.04 g, 0.045 g, 0.05 g, 0.055 g, 0.06 g, 0.065 g, 0.07 g, 0.075 g, 0.08 g, 0.085 g, 0.09 g, 0.095 g, 0.1 g, 0.15 g, 0.2 g, 0.25 g, 0.3 g, 0.35 g, 0.4 g, 0.45 g, 0.5 g, 0.55 g, 0.6 g, 0.65 g, 0.7 g, 0.75 g, 0.8 g, 0.85 g, 0.9 g, 0.95 g, 1 g, 1.5 g, 2 g, 2.5, 3 g, 3.5, 4 g, 4.5 g, 5 g, 5.5 g, 6 g, 6.5 g, 7 g, 7.5 g, 8 g, 8.5 g, 9 g, 9.5 g or 10 g.

[00226] Each of the PI3K and BTK inhibitors according to the invention is effective over a wide dosage range. For example, in the treatment of adult humans, dosages independently range from 0.01 to 1000 mg, from 0.5 to 100 mg, from 1 to 50 mg per day, and from 5 to 40 mg per day are examples of dosages that may be used. The exact dosage will depend upon the route of administration, the form in which the compound is administered, the gender and age of the subject to be treated, the body weight of the subject to be treated, and the preference and experience of the attending physician.

[00227] Efficacy of the compounds and combinations of compounds described herein in treating, preventing and/or managing the indicated diseases or disorders can be tested using various animal models known in the art. Efficacy in treating, preventing and/or managing asthma can be assessed using the ova induced asthma model described, for example, in Lee *et al.*, J. Allergy Clin. Immunol. 118(2):403-9 (2006). Efficacy in treating, preventing and/or managing arthritis (*e.g.*, rheumatoid or psoriatic arthritis) can be assessed using the autoimmune animal models described in, for example, Williams *et al.*, Chem Biol, 17(2):123-34 (2010), WO 2009/088986, WO 2009/088880, and WO 2011/008302. Efficacy in treating, preventing and/or managing psoriasis can be assessed using transgenic or knockout mouse model with targeted mutations in epidermis, vasculature or immune cells, mouse model resulting from spontaneous mutations, and immuno-deficient mouse model with xenotransplantation of human skin or immune cells, all of which are described, for example, in Boehncke *et al.*, Clinics in Dermatology, 25: 596-605 (2007). Efficacy in treating, preventing and/or managing fibrosis or fibrotic conditions can be assessed using the unilateral ureteral obstruction model of renal fibrosis, which is described, for example, in Chevalier *et al.*, Kidney International 75:1145-1152 (2009); the bleomycin induced model of pulmonary fibrosis described in, for example, Moore *et al.*, Am. J. Physiol. Lung. Cell. Mol. Physiol. 294:L152-L160 (2008); a variety of liver/biliary fibrosis models described in, for example, Chuang *et al.*, Clin. Liver Dis.12:333-347 (2008) and Omenetti *et al.*, Laboratory Investigation 87:499-514 (2007) (biliary duct-ligated model); or any of a number of myelofibrosis mouse models such as described in Varicchio *et al.*, Expert Rev. Hematol. 2(3):315-334 (2009). Efficacy in treating, preventing and/or managing scleroderma can be assessed using a mouse model induced by repeated local injections of bleomycin described, for example, in Yamamoto *et al.*, J. Invest. Dermatol. 112: 456-462 (1999). Efficacy in treating, preventing and/or managing dermatomyositis can be assessed using a myositis mouse model induced by immunization with rabbit myosin as described, for example, in Phyanagi *et al.*, Arthritis & Rheumatism, 60(10): 3118-3127 (2009). Efficacy in treating, preventing and/or managing lupus can be assessed using various animal models described, for example, in Ghoreishi *et al.*, Lupus, 19: 1029-1035 (2009); Ohl *et al.*, Journal of Biomedicine and Biotechnology, Article ID 432595 (2011); Xia *et al.*, Rheumatology, 50:2187-2196 (2011); Pau *et al.*, PLoS ONE, 7(5):e36761 (2012); Mustafa *et al.*, Toxicology, 290:156-168 (2011); Ichikawa *et al.*, Arthritis and Rheumatism, 62(2): 493-503 (2012); Ouyang *et al.*, J. Mol. Med.

(2012); Rankin *et al.*, Journal of Immunology, 188:1656-1667 (2012). Efficacy in treating, preventing and/or managing Sjögren's syndrome can be assessed using various mouse models described, for example, in Chiorini *et al.*, J. Autoimmunity, 33: 190-196 (2009).

[00228] To explore the role of PI3K signaling in diffuse large B-cell lymphoma ("DLBCL"), several DLBCL cell lines of varying molecular profiles may be utilized. In an exemplary embodiment, a cellular growth inhibition assay used five cell lines, including four GCB (SU-DHL-4, SU-DHL-6, OCI-LY-8 and WSU-DLCL-2) and one ABC (Ri-1) subtype. In an exemplary embodiment, a cellular growth inhibition assay used five cell lines that were OCI-LY-3, OCI-LY-7, Pfeiffer, Toledo and U2932. In an exemplary embodiment, evidence of PI3K pathway inhibition is measured by reduction in phospho (p)-AKT. In an exemplary embodiment, the kinetics of pathway modulation was characterized by examination of phosphorylation of AKT, PRAS40 and S6 following a time-course of treatment by a PI3K-inhibitor in selected cell lines. In one embodiment, upon B-cell receptor stimulation via antibody-induced crosslinking, some cell lines exhibited enhanced AKT phosphorylation.

[00229] In an exemplary embodiment, the combination effect of a PI3K inhibitor with a BTK inhibitor was observed in a cellular growth inhibition assay in the SU-DHL-4 cell line and in the OCI-LY-8 cell line with BCR crosslinking.

[00230] In one embodiment, provided herein is a method of treating, preventing and/or managing asthma. As used herein, "asthma" encompasses airway constriction regardless of the cause. Common triggers of asthma include, but are not limited to, exposure to an environmental stimulants (e.g., allergens), cold air, warm air, perfume, moist air, exercise or exertion, and emotional stress. Also provided herein is a method of treating, preventing and/or managing one or more symptoms associated with asthma. Examples of the symptoms include, but are not limited to, severe coughing, airway constriction and mucus production.

[00231] Described below are non-limiting exemplary pharmaceutical compositions and methods for preparing the same.

Pharmaceutical Compositions for Oral Administration

[00232] In selected embodiments, the invention provides a pharmaceutical composition for oral administration containing the combination of a PI3K and BTK inhibitor, and a pharmaceutical excipient suitable for oral administration.

[00233] In selected embodiments, the invention provides a solid pharmaceutical composition for oral administration containing: (i) an effective amount of each of a PI3K and BTK inhibitor in combination and (ii) a pharmaceutical excipient suitable for oral administration. In selected embodiments, the composition further contains (iii) an effective amount of a fourth compound.

[00234] In selected embodiments, the pharmaceutical composition may be a liquid pharmaceutical composition suitable for oral consumption. Pharmaceutical compositions of the invention suitable for oral administration can be presented as discrete dosage forms, such as capsules, cachets, or tablets, or liquids or aerosol sprays each containing a predetermined amount of an active ingredient as a powder or in granules, a solution, or a suspension in an aqueous or non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such dosage forms can be prepared by any of the methods of pharmacy, but all methods include the step of bringing the active ingredient(s) into association with the carrier, which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient(s) with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. For example, a tablet can be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with an excipient such as, but not limited to, a binder, a lubricant, an inert diluent, and/or a surface active or dispersing agent. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

[00235] The invention further encompasses anhydrous pharmaceutical compositions and dosage forms since water can facilitate the degradation of some compounds. For example, water may be added (*e.g.*, 5%) in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms of the invention which contain lactose can be made anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected. An anhydrous pharmaceutical composition may be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions may be packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastic or the like, unit dose containers, blister packs, and strip packs.

[00236] Each of the PI3K and BTK inhibitors active ingredients can be combined in an intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier can take a wide variety of forms depending on the form of preparation desired for administration. In preparing the compositions for an oral dosage form, any of the usual pharmaceutical media can be employed as carriers, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, and the like in the case of oral liquid preparations (such as suspensions, solutions, and elixirs) or aerosols; or carriers such as starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents can be used in the case of oral solid preparations, in some embodiments without employing the use of lactose. For example, suitable carriers include powders, capsules, and tablets, with the solid oral preparations. If desired, tablets can be coated by standard aqueous or nonaqueous techniques.

[00237] Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, microcrystalline cellulose, and mixtures thereof.

[00238] Examples of suitable fillers for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof.

[00239] Disintegrants may be used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Too much of a disintegrant may produce tablets which disintegrate in the bottle. Too little may be insufficient for disintegration to occur, thus altering the rate and extent of release of the active ingredients from the dosage form. Thus, a sufficient amount of disintegrant that is neither too little nor too much to detrimentally alter the release of the active ingredient(s) may be used to form the dosage forms of the compounds disclosed herein. The amount of disintegrant used may vary based upon the type of formulation and mode of administration, and may be readily discernible to those of ordinary skill in the art. About 0.5 to about 15 weight percent of disintegrant, or about 1 to about 5 weight percent of disintegrant, may be used in the pharmaceutical composition. Disintegrants that can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums or mixtures thereof.

[00240] Lubricants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (*e.g.*, peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, or mixtures thereof. Additional lubricants include, for example, a silica gel, a coagulated aerosol of synthetic silica, or mixtures thereof. A lubricant can optionally be added, in an amount of less than about 1 weight percent of the pharmaceutical composition.

[00241] When aqueous suspensions and/or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents,

coloring matter or dyes and, if so desired, emulsifying and/or suspending agents, together with such diluents as water, ethanol, propylene glycol, glycerin and various combinations thereof.

[00242] The tablets can be uncoated or coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed. Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

[00243] Surfactants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, hydrophilic surfactants, lipophilic surfactants, and mixtures thereof. That is, a mixture of hydrophilic surfactants may be employed, a mixture of lipophilic surfactants may be employed, or a mixture of at least one hydrophilic surfactant and at least one lipophilic surfactant may be employed.

[00244] A suitable hydrophilic surfactant may generally have an HLB value of at least 10, while suitable lipophilic surfactants may generally have an HLB value of or less than about 10. An empirical parameter used to characterize the relative hydrophilicity and hydrophobicity of non-ionic amphiphilic compounds is the hydrophilic-lipophilic balance ("HLB" value). Surfactants with lower HLB values are more lipophilic or hydrophobic, and have greater solubility in oils, while surfactants with higher HLB values are more hydrophilic, and have greater solubility in aqueous solutions. Hydrophilic surfactants are generally considered to be those compounds having an HLB value greater than about 10, as well as anionic, cationic, or zwitterionic compounds for which the HLB scale is not generally applicable. Similarly, lipophilic (*i.e.*, hydrophobic) surfactants are compounds having an HLB value equal to or less than about 10. However, HLB value of a surfactant is merely a rough guide generally used to enable formulation of industrial, pharmaceutical and cosmetic emulsions.

[00245] Hydrophilic surfactants may be either ionic or non-ionic. Suitable ionic surfactants include, but are not limited to, alkylammonium salts; fusidic acid salts; fatty acid derivatives of amino acids, oligopeptides, and polypeptides; glyceride derivatives of amino acids,

oligopeptides, and polypeptides; lecithins and hydrogenated lecithins; lysolecithins and hydrogenated lysolecithins; phospholipids and derivatives thereof; lysophospholipids and derivatives thereof; carnitine fatty acid ester salts; salts of alkylsulfates; fatty acid salts; sodium docusate; acylactylates; mono- and di-acetylated tartaric acid esters of mono- and di-glycerides; succinylated mono- and di-glycerides; citric acid esters of mono- and di-glycerides; and mixtures thereof.

[00246] Within the aforementioned group, ionic surfactants include, by way of example: lecithins, lysolecithin, phospholipids, lysophospholipids and derivatives thereof; carnitine fatty acid ester salts; salts of alkylsulfates; fatty acid salts; sodium docusate; acylactylates; mono- and di-acetylated tartaric acid esters of mono- and di-glycerides; succinylated mono- and di-glycerides; citric acid esters of mono- and di-glycerides; and mixtures thereof.

[00247] Ionic surfactants may be the ionized forms of lecithin, lysolecithin, phosphatidylcholine, phosphatidylethanolamine, phosphatidylglycerol, phosphatidic acid, phosphatidylserine, lysophosphatidylcholine, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysophosphatidic acid, lysophosphatidylserine, PEG-phosphatidylethanolamine, PVP-phosphatidylethanolamine, lactic esters of fatty acids, stearyl-2-lactylate, stearyl lactylate, succinylated monoglycerides, mono/diacetylated tartaric acid esters of mono/diglycerides, citric acid esters of mono/diglycerides, cholesteryl, caproate, caprylate, caprate, laurate, myristate, palmitate, oleate, ricinoleate, linoleate, linolenate, stearate, lauryl sulfate, teracecyl sulfate, docusate, lauroyl carnitines, palmitoyl carnitines, myristoyl carnitines, and salts and mixtures thereof.

[00248] Hydrophilic non-ionic surfactants may include, but not limited to, alkylglucosides; alkylmaltosides; alkylthioglucosides; lauryl macrogolglycerides; polyoxyalkylene alkyl ethers such as polyethylene glycol alkyl ethers; polyoxyalkylene alkylphenols such as polyethylene glycol alkyl phenols; polyoxyalkylene alkyl phenol fatty acid esters such as polyethylene glycol fatty acids monoesters and polyethylene glycol fatty acids diesters; polyethylene glycol glycerol fatty acid esters; polyglycerol fatty acid esters; polyoxyalkylene sorbitan fatty acid esters such as polyethylene glycol sorbitan fatty acid esters; hydrophilic transesterification products of a polyol with at least one member of the group consisting of glycerides, vegetable oils, hydrogenated vegetable oils, fatty acids, and sterols; polyoxyethylene sterols, derivatives, and analogues

thereof; polyoxyethylated vitamins and derivatives thereof; polyoxyethylene-polyoxypropylene block copolymers; and mixtures thereof; polyethylene glycol sorbitan fatty acid esters and hydrophilic transesterification products of a polyol with at least one member of the group consisting of triglycerides, vegetable oils, and hydrogenated vegetable oils. The polyol may be glycerol, ethylene glycol, polyethylene glycol, sorbitol, propylene glycol, pentaerythritol, or a saccharide.

[00249] Other hydrophilic-non-ionic surfactants include, without limitation, PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG-15 stearate, PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, and poloxamers.

[00250] Suitable lipophilic surfactants include, by way of example only: fatty alcohols; glycerol fatty acid esters; acetylated glycerol fatty acid esters; lower alcohol fatty acids esters; propylene glycol fatty acid esters; sorbitan fatty acid esters; polyethylene glycol sorbitan fatty acid esters; sterols and sterol derivatives; polyoxyethylated sterols and sterol derivatives; polyethylene glycol alkyl ethers; sugar esters; sugar ethers; lactic acid derivatives of mono- and di-glycerides; hydrophobic transesterification products of a polyol with at least one member of the group consisting of glycerides, vegetable oils, hydrogenated vegetable oils, fatty acids and sterols; oil-soluble vitamins/vitamin derivatives; and mixtures thereof. Within this group, preferred

lipophilic surfactants include glycerol fatty acid esters, propylene glycol fatty acid esters, and mixtures thereof, or are hydrophobic transesterification products of a polyol with at least one member of the group consisting of vegetable oils, hydrogenated vegetable oils, and triglycerides.

[00251] In an exemplary embodiment, the composition may include a solubilizer to ensure good solubilization and/or dissolution of the compound of the present invention and to minimize precipitation of the compound of the present invention. This can be especially important for compositions for non-oral use - *e.g.*, compositions for injection. A solubilizer may also be added to increase the solubility of the hydrophilic drug and/or other components, such as surfactants, or to maintain the composition as a stable or homogeneous solution or dispersion.

[00252] Examples of suitable solubilizers include, but are not limited to, the following: alcohols and polyols, such as ethanol, isopropanol, butanol, benzyl alcohol, ethylene glycol, propylene glycol, butanediols and isomers thereof, glycerol, pentaerythritol, sorbitol, mannitol, transcitol, dimethyl isosorbide, polyethylene glycol, polypropylene glycol, polyvinylalcohol, hydroxypropyl methylcellulose and other cellulose derivatives, cyclodextrins and cyclodextrin derivatives; ethers of polyethylene glycols having an average molecular weight of about 200 to about 6000, such as tetrahydrofurfuryl alcohol PEG ether (glycofurol) or methoxy PEG; amides and other nitrogen-containing compounds such as 2-pyrrolidone, 2-piperidone, ϵ -caprolactam, N-alkylpyrrolidone, N-hydroxyalkylpyrrolidone, N-alkylpiperidone, N-alkylcaprolactam, dimethylacetamide and polyvinylpyrrolidone; esters such as ethyl propionate, tributylcitrate, acetyl triethylcitrate, acetyl tributyl citrate, triethylcitrate, ethyl oleate, ethyl caprylate, ethyl butyrate, triacetin, propylene glycol monoacetate, propylene glycol diacetate, ϵ -caprolactone and isomers thereof, δ -valerolactone and isomers thereof, β -butyrolactone and isomers thereof; and other solubilizers known in the art, such as dimethyl acetamide, dimethyl isosorbide, N-methyl pyrrolidones, monoctanoin, diethylene glycol monoethyl ether, and water.

[00253] Mixtures of solubilizers may also be used. Examples include, but not limited to, triacetin, triethylcitrate, ethyl oleate, ethyl caprylate, dimethylacetamide, N-methylpyrrolidone, N-hydroxyethylpyrrolidone, polyvinylpyrrolidone, hydroxypropyl methylcellulose, hydroxypropyl cyclodextrins, ethanol, polyethylene glycol 200-100, glycofurol, transcitol, propylene glycol, and dimethyl isosorbide. Particularly preferred solubilizers include sorbitol, glycerol, triacetin, ethyl alcohol, PEG-400, glycofurol and propylene glycol.

[00254] The amount of solubilizer that can be included is not particularly limited. The amount of a given solubilizer may be limited to a bioacceptable amount, which may be readily determined by one of skill in the art. In some circumstances, it may be advantageous to include amounts of solubilizers far in excess of bioacceptable amounts, for example to maximize the concentration of the drug, with excess solubilizer removed prior to providing the composition to a patient using conventional techniques, such as distillation or evaporation. Thus, if present, the solubilizer can be in a weight ratio of 10%, 25%, 50%, 100%, or up to about 200% by weight, based on the combined weight of the drug, and other excipients. If desired, very small amounts of solubilizer may also be used, such as 5%, 2%, 1% or even less. Typically, the solubilizer may be present in an amount of about 1% to about 100%, more typically about 5% to about 25% by weight.

[00255] The composition can further include one or more pharmaceutically acceptable additives and excipients. Such additives and excipients include, without limitation, detackifiers, anti-foaming agents, buffering agents, polymers, antioxidants, preservatives, chelating agents, viscomodulators, tonicifiers, flavorants, colorants, odorants, opacifiers, suspending agents, binders, fillers, plasticizers, lubricants, and mixtures thereof.

[00256] In addition, an acid or a base may be incorporated into the composition to facilitate processing, to enhance stability, or for other reasons. Examples of pharmaceutically acceptable bases include amino acids, amino acid esters, ammonium hydroxide, potassium hydroxide, sodium hydroxide, sodium hydrogen carbonate, aluminum hydroxide, calcium carbonate, magnesium hydroxide, magnesium aluminum silicate, synthetic aluminum silicate, synthetic hydrocalcite, magnesium aluminum hydroxide, diisopropylethylamine, ethanolamine, ethylenediamine, triethanolamine, triethylamine, triisopropanolamine, trimethylamine, tris(hydroxymethyl)aminomethane (TRIS) and the like. Also suitable are bases that are salts of a pharmaceutically acceptable acid, such as acetic acid, acrylic acid, adipic acid, alginic acid, alkanesulfonic acid, amino acids, ascorbic acid, benzoic acid, boric acid, butyric acid, carbonic acid, citric acid, fatty acids, formic acid, fumaric acid, gluconic acid, hydroquinosulfonic acid, isoascorbic acid, lactic acid, maleic acid, oxalic acid, para-bromophenylsulfonic acid, propionic acid, p-toluenesulfonic acid, salicylic acid, stearic acid, succinic acid, tannic acid, tartaric acid, thioglycolic acid, toluenesulfonic acid, uric acid, and the like. Salts of polyprotic acids, such as

sodium phosphate, disodium hydrogen phosphate, and sodium dihydrogen phosphate can also be used. When the base is a salt, the cation can be any convenient and pharmaceutically acceptable cation, such as ammonium, alkali metals and alkaline earth metals. Example may include, but not limited to, sodium, potassium, lithium, magnesium, calcium and ammonium.

[00257] Suitable acids are pharmaceutically acceptable organic or inorganic acids. Examples of suitable inorganic acids include hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, nitric acid, boric acid, phosphoric acid, and the like. Examples of suitable organic acids include acetic acid, acrylic acid, adipic acid, alginic acid, alkanesulfonic acids, amino acids, ascorbic acid, benzoic acid, boric acid, butyric acid, carbonic acid, citric acid, fatty acids, formic acid, fumaric acid, gluconic acid, hydroquinosulfonic acid, isoascorbic acid, lactic acid, maleic acid, methanesulfonic acid, oxalic acid, para-bromophenylsulfonic acid, propionic acid, p-toluenesulfonic acid, salicylic acid, stearic acid, succinic acid, tannic acid, tartaric acid, thioglycolic acid, toluenesulfonic acid and uric acid.

Pharmaceutical Compositions for Injection

[00258] In selected embodiments, the invention provides a pharmaceutical composition for injection containing the combination of the PI3K and BTK inhibitors and a pharmaceutical excipient suitable for injection. Components and amounts of agents in the compositions are as described herein.

[00259] The forms in which the compositions of the present invention may be incorporated for administration by injection include aqueous or oil suspensions, or emulsions, with sesame oil, corn oil, cottonseed oil, or peanut oil, as well as elixirs, mannitol, dextrose, or a sterile aqueous solution, and similar pharmaceutical vehicles.

[00260] Aqueous solutions in saline are also conventionally used for injection. Ethanol, glycerol, propylene glycol and liquid polyethylene glycol (and suitable mixtures thereof), cyclodextrin derivatives, and vegetable oils may also be employed. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, for the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid and thimerosal.

[00261] Sterile injectable solutions are prepared by incorporating the combination of the PI3K and BTK inhibitors in the required amounts in the appropriate solvent with various other ingredients as enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, certain desirable methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Pharmaceutical Compositions for Topical Delivery

[00262] In some embodiments, the invention provides a pharmaceutical composition for transdermal delivery containing the combination of the PI3K and BTK inhibitors and a pharmaceutical excipient suitable for transdermal delivery.

[00263] Compositions of the present invention can be formulated into preparations in solid, semi-solid, or liquid forms suitable for local or topical administration, such as gels, water soluble jellies, creams, lotions, suspensions, foams, powders, slurries, ointments, solutions, oils, pastes, suppositories, sprays, emulsions, saline solutions, dimethylsulfoxide (DMSO)-based solutions. In general, carriers with higher densities are capable of providing an area with a prolonged exposure to the active ingredients. In contrast, a solution formulation may provide more immediate exposure of the active ingredient to the chosen area.

[00264] The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients, which are compounds that allow increased penetration of, or assist in the delivery of, therapeutic molecules across the stratum corneum permeability barrier of the skin. There are many of these penetration-enhancing molecules known to those trained in the art of topical formulation. Examples of such carriers and excipients include, but are not limited to, humectants (*e.g.*, urea), glycols (*e.g.*, propylene glycol), alcohols (*e.g.*, ethanol), fatty acids (*e.g.*, oleic acid), surfactants (*e.g.*, isopropyl myristate and sodium lauryl sulfate), pyrrolidones, glycerol monolaurate, sulfoxides, terpenes (*e.g.*, menthol), amines, amides, alkanes, alkanols, water,

calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

[00265] Another exemplary formulation for use in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the combination of the PI3K and BTK inhibitors in controlled amounts, either with or without another agent.

[00266] The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, *e.g.*, U.S. Patent Nos. 5,023,252; 4,992,445 and 5,001,139. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Pharmaceutical Compositions for Inhalation

[00267] Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described supra. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be inhaled directly from the nebulizing device or the nebulizing device may be attached to a face mask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices that deliver the formulation in an appropriate manner.

Other Pharmaceutical Compositions

[00268] Pharmaceutical compositions may also be prepared from compositions described herein and one or more pharmaceutically acceptable excipients suitable for sublingual, buccal, rectal, intraosseous, intraocular, intranasal, epidural, or intraspinal administration. Preparations for such pharmaceutical compositions are well-known in the art. See, *e.g.*, Anderson, Philip O.; Knoben, James E.; Troutman, William G, eds., Handbook of Clinical Drug Data, Tenth Edition, McGraw-

Hill, 2002; and Pratt and Taylor, eds., Principles of Drug Action, Third Edition, Churchill Livingston, N.Y., 1990.

[00269] Administration of the combination of the PI3K and BTK inhibitors or pharmaceutical composition of these compounds can be effected by any method that enables delivery of the compounds to the site of action. These methods include oral routes, intraduodenal routes, parenteral injection (including intravenous, intraarterial, subcutaneous, intramuscular, intravascular, intraperitoneal or infusion), topical (*e.g.*, transdermal application), rectal administration, via local delivery by catheter or stent or through inhalation. The combination of compounds can also be administered intraadiposally or intrathecally.

[00270] The compositions of the invention may also be delivered via an impregnated or coated device such as a stent, for example, or an artery-inserted cylindrical polymer. Such a method of administration may, for example, aid in the prevention or amelioration of restenosis following procedures such as balloon angioplasty. Without being bound by theory, compounds of the invention may slow or inhibit the migration and proliferation of smooth muscle cells in the arterial wall which contribute to restenosis. A compound of the invention may be administered, for example, by local delivery from the struts of a stent, from a stent graft, from grafts, or from the cover or sheath of a stent. In some embodiments, a compound of the invention is admixed with a matrix. Such a matrix may be a polymeric matrix, and may serve to bond the compound to the stent. Polymeric matrices suitable for such use, include, for example, lactone-based polyesters or copolyesters such as polylactide, polycaprolactoneglycolide, polyorthoesters, polyanhydrides, polyaminoacids, polysaccharides, polyphosphazenes, poly(ether-ester) copolymers (*e.g.* PEO-PLLA); polydimethylsiloxane, poly(ethylene-vinylacetate), acrylate-based polymers or copolymers (*e.g.*, polyhydroxyethyl methylmethacrylate, polyvinyl pyrrolidinone), fluorinated polymers such as polytetrafluoroethylene and cellulose esters. Suitable matrices may be nondegrading or may degrade with time, releasing the compound or compounds. The combination of the PI3K and BTK inhibitors may be applied to the surface of the stent by various methods such as dip/spin coating, spray coating, dip-coating, and/or brush-coating. The compounds may be applied in a solvent and the solvent may be allowed to evaporate, thus forming a layer of compound onto the stent. Alternatively, the compound may be located in the body of the stent or graft, for example in microchannels or micropores. When implanted, the

compound diffuses out of the body of the stent to contact the arterial wall. Such stents may be prepared by dipping a stent manufactured to contain such micropores or microchannels into a solution of the compound of the invention in a suitable solvent, followed by evaporation of the solvent. Excess drug on the surface of the stent may be removed via an additional brief solvent wash. In yet other embodiments, compounds of the invention may be covalently linked to a stent or graft. A covalent linker may be used which degrades *in vivo*, leading to the release of the compound of the invention. Any bio-labile linkage may be used for such a purpose, such as ester, amide or anhydride linkages. The combination of the PI3K and BTK inhibitors may additionally be administered intravascularly from a balloon used during angioplasty. Extravascular administration of the combination of the PI3K and BTK inhibitors via the pericard or via adventitial application of formulations of the invention may also be performed to decrease restenosis.

[00271] Exemplary parenteral administration forms include solutions or suspensions of active compound in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired.

[00272] The invention also provides kits. The kits include each of the PI3K and BTK inhibitors, either alone or in combination in suitable packaging, and written material that can include instructions for use, discussion of clinical studies and listing of side effects. Such kits may also include information, such as scientific literature references, package insert materials, clinical trial results, and/or summaries of these and the like, which indicate or establish the activities and/or advantages of the composition, and/or which describe dosing, administration, side effects, drug interactions, or other information useful to the health care provider. Such information may be based on the results of various studies, for example, studies using experimental animals involving *in vivo* models and studies based on human clinical trials. The kit may further contain another agent. In selected embodiments, the PI3K and BTK inhibitors and the agent are provided as separate compositions in separate containers within the kit. In selected embodiments, the PI3K and BTK inhibitors and the agent are provided as a single composition within a container in the kit. Suitable packaging and additional articles for use (*e.g.*, measuring cup for liquid preparations, foil wrapping to minimize exposure to air, and the like) are known in the art and may be included in the kit. Kits described herein can be provided, marketed and/or promoted to

health providers, including physicians, nurses, pharmacists, formulary officials, and the like. Kits may also, in selected embodiments, be marketed directly to the consumer.

Dosages and Dosing Regimens

[00273] The amounts of the combination of the PI3K and BTK inhibitors administered will be dependent on the mammal being treated, the severity of the disorder or condition, the rate of administration, the disposition of the compounds and the discretion of the prescribing physician. However, an effective dosage of a PI3K or a BTK inhibitor is in the range of about 0.001 to about 100 mg per kg body weight per day, such as about 1 to about 35 mg/kg/day, in single or divided doses. For a 70 kg human, this would amount to about 0.05 to 7 g/day, such as about 0.05 to about 2.5 g/day. In some embodiments, an effective dosage of a BTK inhibitor disclosed herein, either alone or administered in combination with a PI3K inhibitor, is in the range of about 1 mg to about 300 mg, about 10 mg to about 250 mg, about 20 mg to about 225 mg, about 25 mg to about 200 mg, about 10 mg to about 200 mg, about 20 mg to about 150 mg, about 30 mg to about 120 mg, about 10 mg to about 90 mg, about 20 mg to about 80 mg, about 30 mg to about 70 mg, about 40 mg to about 60 mg, about 45 mg to about 55 mg, about 48 mg to about 52 mg, about 50 mg to about 150 mg, about 60 mg to about 140 mg, about 70 mg to about 130 mg, about 80 mg to about 120 mg, about 90 mg to about 110 mg, about 95 mg to about 105 mg, about 150 mg to about 250 mg, about 160 mg to about 240 mg, about 170 mg to about 230 mg, about 180 mg to about 220 mg, about 190 mg to about 210 mg, about 195 mg to about 205 mg, or about 198 to about 202 mg. In some embodiments, an effective dosage of a BTK inhibitor disclosed herein, either alone or in combination with a PI3K inhibitor, is about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, or about 250 mg. In some embodiments a BTK inhibitor disclosed herein, is administered either alone or administered in combination with a PI3K inhibitor, in a single dose, while in other embodiments a BTK inhibitor disclosed herein, is administered either alone or in combination with a PI3K inhibitor, b.i.d. (twice a day).

[00274] In some embodiments, an effective dosage of a BTK inhibitor disclosed herein, either alone or administered in combination with a PI3K inhibitor, is in the range of about 0.01 mg/kg to about 4.3 mg/kg, about 0.15 mg/kg to about 3.6 mg/kg, about 0.3 mg/kg to about 3.2 mg/kg, about 0.35 mg/kg to about 2.85 mg/kg, about 0.15 mg/kg to about 2.85 mg/kg, about 0.3 mg to

about 2.15 mg/kg, about 0.45 mg/kg to about 1.7 mg/kg, about 0.15 mg/kg to about 1.3 mg/kg, about 0.3 mg/kg to about 1.15 mg/kg, about 0.45 mg/kg to about 1 mg/kg, about 0.55 mg/kg to about 0.85 mg/kg, about 0.65 mg/kg to about 0.8 mg/kg, about 0.7 mg/kg to about 0.75 mg/kg, about 0.7 mg/kg to about 2.15 mg/kg, about 0.85 mg/kg to about 2 mg/kg, about 1 mg/kg to about 1.85 mg/kg, about 1.15 mg/kg to about 1.7 mg/kg, about 1.3 mg/kg mg to about 1.6 mg/kg, about 1.35 mg/kg to about 1.5 mg/kg, about 2.15 mg/kg to about 3.6 mg/kg, about 2.3 mg/kg to about 3.4 mg/kg, about 2.4 mg/kg to about 3.3 mg/kg, about 2.6 mg/kg to about 3.15 mg/kg, about 2.7 mg/kg to about 3 mg/kg, about 2.8 mg/kg to about 3 mg/kg, or about 2.85 mg/kg to about 2.95 mg/kg. In some embodiments, an effective dosage of a BTK inhibitor disclosed herein, either alone or administered in combination with a PI3K inhibitor, is about 0.35 mg/kg, about 0.7 mg/kg, about 1 mg/kg, about 1.4 mg/kg, about 1.8 mg/kg, about 2.1 mg/kg, about 2.5 mg/kg, about 2.85 mg/kg, about 3.2 mg/kg, or about 3.6 mg/kg. In some embodiments a BTK inhibitor disclosed herein, is administered either alone or in combination with a PI3K inhibitor, in a single dose, while in other embodiments a BTK inhibitor disclosed herein, is administered either alone or in combination with a PI3K inhibitor, b.i.d. (twice a day).

[00275] In some embodiments, an effective dosage of a PI3K inhibitor disclosed herein, either alone or administered in combination with a BTK inhibitor, is in the range of about 1 mg to about 300 mg, about 10 mg to about 250 mg, about 20 mg to about 225 mg, about 25 mg to about 200 mg, about 1 mg to about 50 mg, about 5 mg to about 45 mg, about 10 mg to about 40 mg, about 15 mg to about 35 mg, about 20 mg to about 30 mg, about 23 mg to about 28 mg, about 50 mg to about 150 mg, about 60 mg to about 140 mg, about 70 mg to about 130 mg, about 80 mg to about 120 mg, about 90 mg to about 110 mg, or about 95 mg to about 105 mg, about 98 mg to about 102 mg, about 150 mg to about 250 mg, about 160 mg to about 240 mg, about 170 mg to about 230 mg, about 180 mg to about 220 mg, about 190 mg to about 210 mg, about 195 mg to about 205 mg, or about 198 to about 207 mg. In some embodiments, an effective dosage of a PI3K inhibitor disclosed herein, either alone or administered in combination with a BTK inhibitor, is about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, or about 250 mg. In some embodiments the PI3K inhibitor disclosed herein, is administered either alone or in combination with a BTK inhibitor, in a single dose, while in other embodiments a PI3K inhibitor

disclosed herein, is administered either alone or in combination with a BTK inhibitor, b.i.d. (twice a day).

[00276] In some embodiments, an effective dosage of a PI3K inhibitor disclosed herein, either alone or administered in combination with a BTK inhibitor, is in the range of about 0.01 mg/kg to about 4.3 mg/kg, about 0.15 mg/kg to about 3.6 mg/kg, about 0.3 mg/kg to about 3.2 mg/kg, about 0.35 mg/kg to about 2.85 mg/kg, about 0.01 mg/kg to about 0.7 mg/kg, about 0.07 mg/kg to about 0.65 mg/kg, about 0.15 mg/kg to about 0.6 mg/kg, about 0.2 mg/kg to about 0.5 mg/kg, about 0.3 mg/kg to about 0.45 mg/kg, about 0.3 mg/kg to about 0.4 mg/kg, about 0.7 mg/kg to about 2.15 mg/kg, about 0.85 mg/kg to about 2 mg/kg, about 1 mg/kg to about 1.85 mg/kg, about 1.15 mg/kg to about 1.7 mg/kg, about 1.3 mg/kg to about 1.6 mg/kg, about 1.35 mg/kg to about 1.5 mg/kg, about 1.4 mg/kg to about 1.45 mg/kg, about 2.15 mg/kg to about 3.6 mg/kg, about 2.3 mg/kg to about 3.4 mg/kg, about 2.4 mg/kg to about 3.3 mg/kg, about 2.6 mg/kg to about 3.15 mg/kg, about 2.7 mg/kg to about 3 mg/kg, about 2.8 mg/kg to about 3 mg/kg, or about 2.85 mg/kg to about 2.95 mg/kg. In some embodiments, an effective dosage of a PI3K inhibitor disclosed herein, either alone or administered in combination with a BTK inhibitor, is about 0.4 mg/kg, about 0.7 mg/kg, about 1 mg/kg, about 1.4 mg/kg, about 1.8 mg/kg, about 2.1 mg/kg, about 2.5 mg/kg, about 2.85 mg/kg, about 3.2 mg/kg, or about 3.6 mg/kg. In some embodiments a PI3K inhibitor disclosed herein, is administered either alone or in combination with a BTK inhibitor, in a single dose, while in other embodiments a PI3K inhibitor disclosed herein, is administered either alone or in combination with a BTK inhibitor, b.i.d. (twice a day).

[00277] In some embodiments, 10 to 200 mg BID including 50, 60, 70, 80, 90, 100 or 150 mg BID, for the BTK inhibitor, and 10 to 300 mg BID including 25, 50, 75, 100, 150 or 200 mg BID for the PI3K inhibitor. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect - *e.g.*, by dividing such larger doses into several small doses for administration throughout the day.

[00278] In selected embodiments, the combination of the PI3K and BTK inhibitors is administered in a single dose. In selected embodiments, such administration will be by injection - *e.g.*, intravenous injection, in order to introduce the agents quickly. However, other routes may

be used as appropriate. A single dose of the combination of the PI3K and BTK inhibitors may also be used for treatment of an acute condition.

[00279] In selected embodiments, the combination of the PI3K and BTK inhibitors is administered in multiple doses. Dosing may be about once, twice, three times, four times, five times, six times, or more than six times per day. Dosing may be about once a month, once every two weeks, once a week, or once every other day. In other embodiments, the combination of the PI3K and BTK inhibitors is administered about once per day to about 6 times per day. In another embodiment the administration of the combination of the PI3K and BTK inhibitors continues for less than about 7 days. In yet another embodiment the administration continues for more than about 6, 10, 14, 28 days, two months, six months, or one year. In some cases, continuous dosing is achieved and maintained as long as necessary.

[00280] Administration of the agents of the invention may continue as long as necessary. In selected embodiments, the combination of the PI3K and BTK inhibitors is administered for more than 1, 2, 3, 4, 5, 6, 7, 14, or 28 days. In some embodiments, the combination of the PI3K and BTK inhibitors is administered for less than 28, 14, 7, 6, 5, 4, 3, 2, or 1 day. In selected embodiments, the combination of the PI3K and BTK inhibitors is administered chronically on an ongoing basis - *e.g.*, for the treatment of chronic effects.

[00281] An effective amount of the combination of the PI3K and BTK inhibitors may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, including rectal, buccal, intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, orally, topically, or as an inhalant.

Methods of Treatment

[00282] In selected embodiments, the invention provides a method of treating a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a PI3K inhibitor (or a PI3K- γ inhibitor, PI3K- δ inhibitor, or PI3K- γ,δ inhibitor) and BTK inhibitor, or a pharmaceutically acceptable salt or ester, prodrug, solvate or hydrate of either or both the PI3K inhibitor (or a PI3K- γ inhibitor, PI3K- δ inhibitor, or PI3K- γ,δ inhibitor) or the BTK inhibitor. In selected embodiments, the method relates to the

treatment of cancer such as non-Hodgkin's lymphomas (such as diffuse large B-cell lymphoma), acute myeloid leukemia, thymus, brain, lung, squamous cell, skin, eye, retinoblastoma, intraocular melanoma, oral cavity and oropharyngeal, bladder, gastric, stomach, pancreatic, bladder, breast, cervical, head, neck, renal, kidney, liver, ovarian, prostate, colorectal, bone (*e.g.*, metastatic bone), esophageal, testicular, gynecological, thyroid, CNS, PNS, AIDS-related (*e.g.* lymphoma and Kaposi's sarcoma), viral-induced cancers such as cervical carcinoma (human papillomavirus), B-cell lymphoproliferative disease and nasopharyngeal carcinoma (Epstein-Barr virus), Kaposi's Sarcoma and primary effusion lymphomas (Kaposi's sarcoma herpesvirus), hepatocellular carcinoma (hepatitis B and hepatitis C viruses), and T-cell leukemias (Human T-cell leukemia virus-1), and T-cell leukemias (Human T-cell leukemia virus-1), B cell acute lymphoblastic leukemia, Burkitt's leukemia, juvenile myelomonocytic leukemia, hairy cell leukemia, Hodgkin's disease, multiple myeloma, mast cell leukemia, or mastocytosis. In selected embodiments, the method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (*e.g.*, psoriasis), restenosis, or prostate conditions (*e.g.*, benign prostatic hypertrophy (BPH)).

[00283] In selected embodiments, the invention provides a method of treating an inflammatory, immune, or autoimmune disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a PI3K inhibitor (or a PI3K- γ inhibitor, PI3K- δ inhibitor, or PI3K- γ,δ inhibitor) and BTK inhibitor, or a pharmaceutically acceptable salt or ester, prodrug, solvate or hydrate of either or both the PI3K inhibitor (or a PI3K- γ inhibitor, PI3K- δ inhibitor, or PI3K- γ,δ inhibitor) or the BTK inhibitor. In selected embodiments, the invention also provides a method of treating a disease selected from the group consisting of tumor angiogenesis, chronic inflammatory disease, rheumatoid arthritis, atherosclerosis, inflammatory bowel disease, skin diseases such as psoriasis, eczema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma and melanoma, ulcerative colitis, atopic dermatitis, pouchitis, spondylarthritis, uveitis, Behcets disease, polymyalgia rheumatica, giant-cell arteritis, sarcoidosis, Kawasaki disease, juvenile idiopathic arthritis, hidratenitis suppurativa, Sjögren's syndrome, psoriatic arthritis, juvenile rheumatoid arthritis, ankylosing spondylitis, Crohn's Disease, lupus, and lupus nephritis.

[00284] In selected embodiments, the invention provides a method of treating a solid tumor

cancer with a composition including a combination of a PI3K inhibitor, including a PI3K- γ or PI3K- δ inhibitor, and a BTK inhibitor, wherein the dose is effective to inhibit signaling between the solid tumor cells and at least one microenvironment selected from the group consisting of macrophages, monocytes, mast cells, helper T cells, cytotoxic T cells, regulatory T cells, natural killer cells, myeloid-derived suppressor cells, regulatory B cells, neutrophils, dendritic cells, and fibroblasts. In selected embodiments, the invention provides a method of treating pancreatic cancer, breast cancer, ovarian cancer, melanoma, lung cancer, head and neck cancer, and colorectal cancer using a combination of a BTK inhibitor and a PI3K inhibitor, wherein the dose is effective to inhibit signaling between the solid tumor cells and at least one microenvironment selected from the group consisting of macrophages, monocytes, mast cells, helper T cells, cytotoxic T cells, regulatory T cells, natural killer cells, myeloid-derived suppressor cells, regulatory B cells, neutrophils, dendritic cells, and fibroblasts. Efficacy of the compounds and combinations of compounds described herein in treating, preventing and/or managing the indicated diseases or disorders can be tested using various models known in the art. For example, models for determining efficacy of treatments for pancreatic cancer are described in Herreros-Villanueva, et al. *World J. Gastroenterol.* **2012**, *18*, 1286-1294. Models for determining efficacy of treatments for breast cancer are described e.g. in A. Fantozzi, *Breast Cancer Res.* **2006**, *8*, 212. Models for determining efficacy of treatments for ovarian cancer are described e.g. in Mullany et al., *Endocrinology* **2012**, *153*, 1585-92; and Fong et al., *J. Ovarian Res.* **2009**, *2*, 12. Models for determining efficacy of treatments for melanoma are described e.g. in Damsky et al., *Pigment Cell & Melanoma Res.* **2010**, *23*, 853-859. Models for determining efficacy of treatments for lung cancer are described e.g. in Meuwissen et al., *Genes & Development*, **2005**, *19*, 643-664. Models for determining efficacy of treatments for lung cancer are described e.g. in Kim, *Clin. Exp. Otorhinolaryngol.* **2009**, *2*, 55-60; and Sano, *Head Neck Oncol.* **2009**, *1*, 32. Models for determining efficacy of treatments for colorectal cancer, including the CT26 model, are described below in the examples.

[00285] Efficacy of the compounds and combinations of compounds described herein in treating, preventing and/or managing other indicated diseases or disorders described here can also be tested using various models known in the art. Efficacy in treating, preventing and/or managing asthma can be assessed using the ova induced asthma model described, for example, in Lee et al., *J. Allergy Clin. Immunol.* **2006**, *118*, 403-9. Efficacy in treating, preventing and/or

managing arthritis (e.g., rheumatoid or psoriatic arthritis) can be assessed using the autoimmune animal models described in, for example, Williams et al., *Chem. Biol.* **2010**, *17*, 123-34, WO 2009/088986, WO 2009/088880, and WO 2011/008302. Efficacy in treating, preventing and/or managing psoriasis can be assessed using transgenic or knockout mouse model with targeted mutations in epidermis, vasculature or immune cells, mouse model resulting from spontaneous mutations, and immuno-deficient mouse model with xenotransplantation of human skin or immune cells, all of which are described, for example, in Boehncke et al., *Clinics in Dermatology*, **2007**, *25*, 596-605. Efficacy in treating, preventing and/or managing fibrosis or fibrotic conditions can be assessed using the unilateral ureteral obstruction model of renal fibrosis, which is described, for example, in Chevalier et al., *Kidney International* **2009**, *75*, 1145-1152; the bleomycin induced model of pulmonary fibrosis described in, for example, Moore et al., *Am. J. Physiol. Lung. Cell. Mol. Physiol.* **2008**, *294*, L152-L160; a variety of liver/biliary fibrosis models described in, for example, Chuang et al., *Clin. Liver Dis.* **2008**, *12*, 333-347 and Omenetti et al., *Laboratory Investigation*, **2007**, *87*, 499-514 (biliary duct-ligated model); or any of a number of myelofibrosis mouse models such as described in Varicchio et al., *Expert Rev. Hematol.* **2009**, *2*, 315-334. Efficacy in treating, preventing and/or managing scleroderma can be assessed using a mouse model induced by repeated local injections of bleomycin described, for example, in Yamamoto et al., *J. Invest. Dermatol.* **1999**, *112*, 456-462. Efficacy in treating, preventing and/or managing dermatomyositis can be assessed using a myositis mouse model induced by immunization with rabbit myosin as described, for example, in Phyanagi et al., *Arthritis & Rheumatism*, **2009**, *60(10)*, 3118-3127. Efficacy in treating, preventing and/or managing lupus can be assessed using various animal models described, for example, in Ghoreishi et al., *Lupus*, **2009**, *19*, 1029-1035; Ohl et al., *J. Biomed. & Biotechnol.*, Article ID 432595 (2011); Xia et al., *Rheumatology*, **2011**, *50*, 2187-2196; Pau et al., *PLoS ONE*, **2012**, *7(5)*, e36761; Mustafa et al., *Toxicology*, **2011**, *90*, 156-168; Ichikawa et al., *Arthritis & Rheumatism*, **2012**, *62(2)*, 493-503; Rankin et al., *J. Immunology*, **2012**, *188*, 1656-1667. Efficacy in treating, preventing and/or managing Sjögren's syndrome can be assessed using various mouse models described, for example, in Chiorini et al., *J. Autoimmunity*, **2009**, *33*, 190-196.

[00286] In selected embodiments, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, comprising the step of administering a

therapeutically effective dose of a BTK inhibitor, or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof. In a preferred embodiment, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is Formula (XVIII), or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof. In a preferred embodiment, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is Formula (XVIII), or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof, further comprising the step of administering a therapeutically effective dose of an anticoagulant or antiplatelet agent.

[00287] In a preferred embodiment, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is Formula (XVIII), or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof, further comprising the step of administering a therapeutically effective dose of an anticoagulant or antiplatelet agent, wherein the anticoagulant or antiplatelet agent is selected from the group consisting of clopidogrel, prasugrel, ticagrelor, ticlopidine, warfarin, acenocoumarol, dicumarol, phenprocoumon, heparin, low molecular weight heparin, fondaparinux, and idraparinux.

[00288] In selected embodiments, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is Formula (XVIII), and wherein the cancer is selected from the group consisting of bladder cancer, squamous cell carcinoma including head and neck cancer, pancreatic ductal adenocarcinoma (PDA), pancreatic cancer, colon carcinoma, mammary carcinoma, breast cancer, fibrosarcoma, mesothelioma, renal cell carcinoma, lung carcinoma, thymoma, prostate cancer, colorectal cancer, ovarian cancer, acute myeloid leukemia, thymus cancer, brain cancer, squamous cell cancer, skin cancer, eye cancer, retinoblastoma, melanoma, intraocular melanoma, oral cavity and oropharyngeal cancers, gastric cancer, stomach cancer, cervical cancer, head, neck, renal cancer, kidney cancer, liver cancer, ovarian cancer, prostate cancer, colorectal cancer, esophageal cancer, testicular cancer, gynecological cancer, thyroid cancer, acquired immune deficiency syndrome (AIDS)-related

cancers (e.g., lymphoma and Kaposi's sarcoma), viral-induced cancer, glioblastoma, esophageal tumors, hematological neoplasms, non-small-cell lung cancer, chronic myelocytic leukemia, diffuse large B-cell lymphoma, esophagus tumor, follicle center lymphoma, head and neck tumor, hepatitis C virus infection, hepatocellular carcinoma, Hodgkin's disease, metastatic colon cancer, multiple myeloma, non-Hodgkin's lymphoma, indolent non-Hogkin's lymphoma, ovary tumor, pancreas tumor, renal cell carcinoma, small-cell lung cancer, stage IV melanoma, chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia (ALL), mature B-cell ALL, follicular lymphoma, mantle cell lymphoma, and Burkitt's lymphoma.

[00289] In selected embodiments, the invention provides a method of treating a cancer in a human with a history of thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof. In selected embodiments, the invention provides a method of treating a cancer in a human sensitive to platelet-mediated thrombosis, method of treating a cancer in a human with a history of thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is a compound of Formula (XVIII) or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof.

[00290] In selected embodiments, the BTK inhibitor and the anticoagulant or the antiplatelet agent are administered sequentially. In selected embodiments, the BTK inhibitor and the anticoagulant or the antiplatelet agent are administered concomitantly. In selected embodiments, the BTK inhibitor is administered before the anticoagulant or the antiplatelet agent. In selected embodiments, the BTK inhibitor is administered after the anticoagulant or the antiplatelet agent.

[00291] Preferred anti-platelet and anticoagulant agents for use in the methods of the present invention include, but are not limited to, cyclooxygenase inhibitors (e.g., aspirin), adenosine diphosphate (ADP) receptor inhibitors (e.g., clopidogrel and ticlopidine), phosphodiesterase inhibitors (e.g., cilostazol), glycoprotein IIb/IIIa inhibitors (e.g., abciximab, eptifibatide, and tirofiban), adenosine reuptake inhibitors (e.g., dipyridamole), and acetylsalicylic acid (aspirin). In other embodiments, examples of anti-platelet agents for use in the methods of the present invention include anagrelide, aspirin/extended-release dipyridamole, cilostazol,

clopidogrel, dipyridamole, prasugrel, ticagrelor, ticlopidine, vorapaxar, tirofiban HCl, eptifibatide, abciximab, argatroban, bivalirudin, dalteparin, desirudin, enoxaparin, fondaparinux, heparin, lepirudin, apixaban, dabigatran etexilate mesylate, rivaroxaban, and warfarin.

Combinations of PI3K and BTK Inhibitors

[00292] Non-limiting, exemplary embodiments of combinations of the PI3K inhibitors and BTK inhibitors described above are given in the following numbered paragraphs 1 to 50. The disclosure encompassed herein should in no way be construed as being limited to these examples, but rather should be construed to encompass any and all variations which become evident as a result of the teachings provided herein.

[00293] In an embodiment, the PI3K inhibitor and BTK inhibitor are provided in a PI3K inhibitor to BTK inhibitor ratio (by mass) selected from the group consisting of 0.01 to 1, 0.05 to 1, 0.1 to 1, 0.5 to 1, 1 to 1, 2 to 1, 5 to 1, 10 to 1, 20 to 1, and 100 to 1.

[00294] While preferred embodiments of the invention are shown and described herein, including the above embodiments, such embodiments are provided by way of example only and are not intended to otherwise limit the scope of the invention. Various alternatives to the described embodiments of the invention may be employed in practicing the invention.

EXAMPLES

[00295] The embodiments encompassed herein are now described with reference to the following examples. These examples are provided for the purpose of illustration only and the disclosure encompassed herein should in no way be construed as being limited to these examples, but rather should be construed to encompass any and all variations which become evident as a result of the teachings provided herein.

Example 1 – Synergistic Combination of a BTK inhibitor and a PI3K- δ inhibitor

[00296] Ficoll purified mantle cell lymphoma (MCL) cells (2×10^5) isolated from bone marrow or peripheral blood were treated with each drug alone and with six equimolar concentrations of a BTK inhibitor (Formula XVIII) and a PI3K- δ inhibitor (Formula IX) ranging from 0.01 nM to 10 μ M on 96-well plates in triplicate. Plated cells were then cultured in HS-5 conditioned media at

37°C with 5% CO₂. After 72 hours of culture, cell viability was determined using an (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) (MTS) assay (Cell Titer 96, Promega). Viability data were used to generate cell viability curves for each drug alone and in combination for each sample. The potential synergy of the combination of the BTK inhibitor of Formula XVIII and the PI3K- δ inhibitor of Formula IX at a given equimolar concentration was determined using the median effect model as described in Chou and Talalay, *Adv Enzyme Regul.* **1984**, 22, 27-55. The statistical modeling was run in R using a script that utilizes the median effect model as described in Lee et al., *J. Biopharm. Stat.* **2007**, 17, 461-80. A value of 1, less than 1, and greater than 1 using R defines an additive interaction, synergistic and antagonistic, respectively. The Lee et al. method calculates a 95% confidence interval for each data point. For each viability curve, to be considered synergistic, a data point must have an interaction index below 1 and the upper confidence interval must also be below 1. In order to summarize and demonstrate collective synergy results, an interaction dot blot was generated for the primary patient samples.

[00297] A similar approach was utilized to study diffuse large B cell lymphoma (DLBCL) (TMD8) and MCL (MINO) cell lines. Cells were treated with each drug alone and with six equimolar concentrations of the BTK inhibitor of Formula XVIII and the PI3K- δ inhibitor of Formula IX ranging from 0.003 nM to 1.0 μ M (for TMD8) or 0.03 nM to 10 μ M (for MINO) on 96-well plates in triplicate. Plated cells were then cultured in standard conditioned media plus FBS at 37°C with 5% CO₂. After 72 hours of culture, viability was determined using an MTS assay (Cell Titer 96, Promega). Viability data were used to generate cell viability curves for each drug alone and in combination for each sample. The results of the experiments described in this example are shown in FIG. 1, FIG. 2, FIG. 3, and FIG. 4.

Example 2 – Synergistic Combination of a BTK inhibitor and a PI3K- δ inhibitor

[00298] Combination experiments were performed to determine the synergistic, additive, or antagonistic behavior of drug combinations using the Chou/Talalay method/algorithm by defining combination indexes for drug combinations. Information about experimental design for evaluation of synergy is described in e.g. Chou and Talalay, *Adv. Enzyme Regul.* **1984**, 22, 27-55 and more generally in e.g. Greco et al., *Pharmacol. Rev.* **1995**, 47, 331-385. The study was performed using the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula

(IX). Single agent activities were first determined in the various cell lines and subsequently, the combination indexes were established using equimolar ratios taking the single agent drug EC50s into consideration. For individual agents that displayed no single agent activity, equimolar ratios were used at fixed concentrations to establish combination indexes. The readout from 72 hour proliferation assays using Cell TiterGlo (ATP content of remaining cells) determined the fraction of cells that were effected as compared to untreated cells ($F_a = \text{fraction affected} = (1 - ((\text{cells} + \text{inhibitor}) - \text{background signal}) / ((\text{cells} + \text{DMSO}) - \text{background signal}))$).

[00299] The combination index obtained was ranked according to Table 1.

TABLE 1. Combination Index (CI) Ranking Scheme

Range of CI	Description
<0.1	Very strong synergism
0.1-0.3	Strong synergism
0.3-0.7	Synergism
0.7-0.85	Moderate synergism
0.85-0.9	Slight synergism
0.9-1.1	Nearly additive
1.1-1.2	Slight antagonism
1.2-1.45	Moderate antagonism
1.45-3.3	Antagonism
3.3-10	Strong antagonism
>10	Very strong antagonism

[00300] The detailed results of the cell line studies for the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) are given in FIG. 5 to FIG. 37. The results of the cell line studies are summarized in Table 2.

TABLE 2. Summary of results of the combination of a BTK inhibitor with a PI3K- δ inhibitor (S = synergistic, A = additive, X = no effect).

Cell Line	Indication	ED25	ED50	ED75	ED90
Raji	Burkitt's	S	S	S	S
Ramos	Burkitt's	X	X	X	X
Daudi	Burkitt's	S	S	S	S
Mino	MCL	S	S	S	S
Pfeiffer	iNHL	S	S	S	S
DOHH	iNHL	S	S	S	S
REC-1	iNHL	S	S	A	A
U937	Myeloid	S	S	S	S
K562	CML	X	X	X	X
SU-DHL-1	ABC	S	A	X	X
SU-DHL-2	ABC	S	S	S	S
HBL-1	ABC	S	S	S	S
TMD8	ABC	S	S	S	S
LY19	GCB	X	X	X	X
LY7	GCB	S	S	S	S
LY1	GCB	X	X	X	X
SU-DHL-6	GCB	S	S	S	S
SupB15	B-ALL	S	S	S	S
CCRF	B-ALL	S	A/S	X	X

Example 3 – BTK Inhibitory Effects on Solid Tumor Microenvironment in an Orthotopic Pancreatic Cancer Model

[00301] An orthotopic pancreatic cancer model was used to investigate the therapeutic efficacy of the combination of the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) through treatment of the solid tumor microenvironment. Mice were dosed orally with 15 mg/kg of Formula (XVIII), 15 mg/kg of Formula (IX), or a combination of 15 mg/kg of both drugs.

[00302] Cell line derived from KrasG12D;Trp53R172H;Pdx1-Cre (KPC) mice were orthotopically implanted into the head of the pancreas after 35 passages. Based on the mice background from where the cell lines were generated, 1×10^6 cells were injected in C57BL/6 mice. Throughout the experiment, animals were provided with food and water ad libitum and subjected to a 12-h dark/light cycle. Animal studies were performed in accordance with the U.S. Public Health Service "Guidelines for the Care and Use of Laboratory Animals" (IACUC). After

euthanization, pancreatic tumors were dissected out, weighed and single cell suspensions were prepared for flow cytometry analysis.

[00303] Results of the experiments are shown in FIG. 38, which illustrates tumor growth suppression in the orthotopic pancreatic cancer model. The statistical p-value (presumption against null hypothesis) is shown for each tested single agent and for the combination against the vehicle. The results show that all three treatments provide statistically significant reductions in tumor volume in the pancreatic cancer model.

[00304] Additional results of the experiments relating to treatment of the tumor microenvironment are shown in FIG. 39 to FIG. 41. FIG. 39 shows the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K inhibitor of Formula (IX), or a combination of both drugs on myeloid tumor-associated macrophages (TAMs) in pancreatic tumor-bearing mice. FIG. 40 illustrates the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K inhibitor of Formula (IX), or a combination of both inhibitors on myeloid-derived suppressor cells (MDSCs) in pancreatic tumor-bearing mice. FIG. 41 illustrates the effects of oral dosing with 15 mg/kg of the BTK inhibitor of Formula (XVIII), 15 mg/kg of the PI3K inhibitor of Formula (IX), or a combination of both inhibitors on regulatory T cells (Tregs) in pancreatic tumor-bearing mice. The results shown in FIG. 39 to FIG. 41 demonstrate that administration of the BTK inhibitor of Formula (XVIII) and the combination of the BTK inhibitor of Formula (XVIII) and the PI3K inhibitor of Formula (IX) reduce immunosuppressive tumor associated myeloid cells and Tregs in pancreatic tumor-bearing mice. Overall, BTK inhibition with Formula (XVIII) or a combination of Formula (XVIII) and Formula (IX) significantly reduced tumor burden in an aggressive orthotopic PDA model, decreased immature myeloid infiltrate, reduced the number of tumor associated macrophages, and reduced the number of immunosuppressive Tregs, demonstrating a strong effect on the tumor microenvironment.

Example 4 – BTK Inhibitory Effects on Solid Tumor Microenvironment in an Ovarian Cancer Model

[00157] The ID8 syngeneic orthotropic ovarian cancer murine model was used to investigate the therapeutic efficacy of the BTK inhibitor of Formula (XVIII) through treatment of the solid tumor microenvironment. Human ovarian cancer models, including the ID8 syngeneic

orthotropic ovarian cancer model and other animal models, are described in Fong and Kakar, *J. Ovarian Res.* **2009**, 2, 12; Greenaway et al., *Gynecol. Oncol.* **2008**, 108, 385-94; Urzua et al., *Tumour Biol.* **2005**, 26, 236-44; Janat-Amsbury et al., *Anticancer Res.* **2006**, 26, 3223-28; Janat-Amsbury et al., *Anticancer Res.* **2006**, 26, 2785-89. Animals were treated with vehicle or Formula (XVIII), 15 mg/kg/BID given orally. The results of the study are shown in FIG. 42, FIG. 43, FIG. 44, FIG. 45, FIG. 46, FIG. 47, FIG. 48, and FIG. 49.

[00158] FIG. 42 and FIG. 43 demonstrate that the BTK inhibitor of Formula (XVIII) impairs ID8 ovarian cancer growth in the ID8 syngeneic murine model. FIG. 44 shows that tumor response to treatment with the BTK inhibitor of Formula (XVIII) correlates with a significant reduction in immunosuppressive tumor-associated lymphocytes in tumor-bearing mice. FIG. 45 shows treatment with the BTK inhibitor of Formula (XVIII) impairs ID8 ovarian cancer growth (through reduction in tumor volume) in the syngeneic murine model. FIG. 46 and FIG. 47 show that the tumor response induced by treatment with the BTK inhibitor of Formula (XVIII) correlates with a significant reduction in immunosuppressive B cells in tumor-bearing mice. FIG. 48 and FIG. 49 show that the tumor response induced by treatment with the BTK inhibitor of Formula (XVIII) correlates with a significant reduction in immunosuppressive tumor associated Tregs and an increase in CD8⁺ T cells.

[00159] The results shown in FIG. 42 to FIG. 49 illustrate the surprising efficacy of the BTK inhibitor of Formula (XVIII) in modulating tumor microenvironment in a model predictive of efficacy as a treatment for ovarian cancer in humans.

Example 5 – BTK Inhibitory Effects on Solid Tumor Microenvironment Through Modulation of Tumor-Infiltrating MDSCs and TAMs

[00160] A study was performed to observe potential reduction in tumor burden through modulation of tumor infiltrating MDSCs and TAMs using the BTK inhibitor of Formula (XVIII) and/or gemcitabine (“Gem”). In this study, KPC derived mouse pancreatic cancer cells (KrasG12D;Trp53R172H;Pdx1-Cre) were injected into the pancreases. Animals were treated with (1) vehicle; (2) Formula (XVIII), 15 mg/kg/BID given orally; (3) gemcitabine 15 mg/kg intravenous (IV) administered every 4 days for 3 injections; or (4) Formula (XVIII), 15 mg/kg/BID given orally with together with gemcitabine, 15 mg/kg IV administered every 4 days for 3 injections.

[00161] Single cell suspensions from tumor samples. Mouse tumor tissue was collected and stored in PBS/0.1% soybean trypsin inhibitor prior to enzymatic dissociation. Samples were finely minced with a scissors and mouse tissue was transferred into DMEM containing 1.0 mg/ml collagenase IV (Gibco), 0.1% soybean trypsin inhibitor, and 50 U/ml DNase (Roche) and incubated at 37C for 30 min. with constant stirring while human tissue was digested in 2.0 mg/ml collagenase IV, 1.0 mg/ml hyaluronidase, 0.1% soybean trypsin inhibitor, and 50 U/ml DNase for 45 minutes. Suspensions were filtered through a 100 micron filter and washed with FACS buffer (PBS/0.5% BSA/2.0 mM EDTA) prior to staining. Two million total cells were stained with antibodies as indicated. Intracellular detection of FoxP3 was achieved following permeabilization with BD Perm Buffer III (BD Biosciences) and eBioscience Fix/Perm respectively. Following surface staining, samples were acquired on a BD Fortessa and analyzed using FlowJo (Treestar) software.

[00162] In FIG. 50, the reduction in tumor size upon treatment is shown. The effects on particular cell subsets are shown in the flow cytometry data presented in FIG. 51, FIG. 52, FIG. 53, and FIG. 54.

[00163] The results shown in FIG. 50 to FIG. 54 illustrate reduction in tumor burden by modulating the tumor infiltrating MDSCs and TAMs, which affects Treg and CD8⁺ T cell levels, through inhibition of BTK using Formula (XVIII).

Example 6 – Effects of BTK Inhibitors on Thrombosis

[00164] Clinical studies have shown that targeting the BCR signaling pathway by inhibiting BTK produces significant clinical benefit (Byrd, et al., *N. Engl. J. Med.* **2013**, 369(1), 32-42, Wang, et al., *N. Engl. J. Med.* **2013**, 369(6), 507-16). However, in these studies, bleeding has been reported in up to 50% of ibrutinib-treated patients. Most bleeding events were of grade 1-2 (spontaneous bruising or petechiae) but, in 5% of patients, they were of grade 3 or higher after trauma. These results are reflected in the prescribing information for ibrutinib, where bleeding events of any grade, including bruising and petechiae, were reported in approximately half of patients treated with ibrutinib (IMBRUVICA package insert and prescribing information, revised July 2014, U.S. Food and Drug Administration).

[00165] Constitutive or aberrant activation of the BCR signaling cascade has been implicated in the propagation and maintenance of a variety of B cell malignancies. Small molecule inhibitors

of BTK, a protein early in this cascade and specifically expressed in B cells, have emerged as a new class of targeted agents. There are several BTK inhibitors, including Formula XXVII (CC-292), and Formula XX-A (PCI-32765, ibrutinib), in clinical development. Importantly, early stage clinical trials have found ibrutinib to be particularly active in chronic lymphocytic leukemia (CLL) and mantle cell lymphoma (MCL), suggesting that this class of inhibitors may play a significant role in various types of cancers (Aalipour and Advani, *Br. J. Haematol.* **2013**, *163*, 436-43). However, their effects are not limited to leukemia or lymphomas as platelets also rely on the Tec kinases family members BTK and Tec for signal transduction in response to various thrombogenic stimuli (Oda, et al., *Blood* **2000**, *95*(5), 1663-70; Atkinson, et al. *Blood* **2003**, *102*(10), 3592-99). In fact, both Tec and BTK play an important role in the regulation of phospholipase C γ 2 (PLC γ 2) downstream of the collagen receptor glycoprotein VI (GPVI) in human platelets. In addition, BTK is activated and undergoes tyrosine phosphorylation upon challenge of the platelet thrombin receptor, which requires the engagement of α IIb β 3 integrin and PI3K activity (Laffargue, et al., *FEBS Lett.* **1999**, *443*(1), 66-70). It has also been implicated in GPIb α -dependent thrombus stability at sites of vascular injury (Liu, et al., *Blood* **2006**, *108*(8), 2596-603). Thus, BTK and Tec are involved in several processes important in supporting the formation of a stable hemostatic plug, which is critical for preventing significant blood loss in response to vascular injury. Hence, the effects of the BTK inhibitor of Formula (XVIII) and ibrutinib were evaluated on human platelet-mediated thrombosis by utilizing the *in vivo* human thrombus formation in the VWF HA1 mice model described in Chen et al. *Nat. Biotechnol.* **2008**, *26*(1), 114-19.

[00166] Administration of anesthesia, insertion of venous and arterial catheters, fluorescent labeling and administration of human platelets (5×10^8 /ml), and surgical preparation of the cremaster muscle in mice have been previously described (Chen et al. *Nat Biotechnol.* **2008**, *26*(1), 114-19). Injury to the vessel wall of arterioles (~40–65 μ m diameter) was performed using a pulsed nitrogen dye laser (440 nm, Photonic Instruments) applied through a 20 $\times$ water-immersion Olympus objective (LUMPlanFI, 0.5 numerical aperture (NA)) of a Zeiss Axiotech vario microscope. Human platelet and wall interactions were visualized by fluorescence microscopy using a system equipped with a Yokogawa CSU-22 spinning disk confocal scanner, iXON EM camera, and 488 nm and 561 nm laser lines to detect BCECF-labeled and rhodamine-labeled platelets, respectively (Revolution XD, Andor Technology). The extent of thrombus

formation was assessed for 2 min after injury and the area (μm^2) of coverage determined (Image IQ, Andor Technology). For the Formula (XVIII), Formula (XXVII) (CC-292), and Formula (XX-A) (ibrutinib) inhibition studies, the BTK inhibitors were added to purified human platelets for 30 min before administration.

[00167] The *in vivo* thrombus effects of the BTK inhibitors, Formula (XVIII), Formula (XXVII) (CC-292), and Formula (XX-A) (ibrutinib), were evaluated on human platelet-mediated thrombosis by utilizing the *in vivo* human thrombus formation in the VWF HA1 mice model, which has been previously described (Chen et al. *Nat Biotechnol.* **2008**, 26(1), 114-19). Purified human platelets were preincubated with various concentrations of the BTK inhibitors (0.1 μM , 0.5 μM , or 1 μM) or DMSO and then administered to VWF HA1 mice, followed by laser-induced thrombus formation. The BTK inhibitor-treated human platelets were fluorescently labeled and infused continuously through a catheter inserted into the femoral artery. Their behavior in response to laser-induced vascular injury was monitored in real time using two-channel confocal intravital microscopy (Furie and Furie, *J. Clin. Invest.* **2005**, 115(12), 2255-62). Upon induction of arteriole injury untreated platelets rapidly formed thrombi with an average thrombus size of $6,450 \pm 292 \text{ mm}^2$ (mean $\pm$ s.e.m.), as shown in FIG. 55 and FIG. 56. Similarly, Formula (XVIII) (1 μM) treated platelets formed a slightly smaller but not significantly different thrombi with an average thrombus size of $5733 \pm 393 \text{ mm}^2$ (mean $\pm$ s.e.m.). In contrast, a dramatic reduction in thrombus size occurred in platelets pretreated with 1 μM of Formula XX-A (ibrutinib), $2600 \pm 246 \text{ mm}^2$ (mean $\pm$ s.e.m.), resulting in a reduction in maximal thrombus size by approximately 61% compared with control ($P > 0.001$) (FIG. 55 and 57). Similar results were obtained with platelets pretreated with 500 nM of Formula (XVIII) or ibrutinib: thrombus size of $5946 \pm 283 \text{ mm}^2$, and $2710 \pm 325 \text{ mm}^2$ respectively. These initial results may provide some mechanistic background and explanation on the reported 44% bleeding related adverse event rates in the Phase III RESONATETM study comparing ibrutinib with ofatumumab. The results obtained for Formula XXVII (CC-292) were similar to that for Formula XX-A (ibrutinib), as shown in FIG. 55, 56, and 57. The effect of the BTK inhibitor concentration is shown in FIG. 58. These results demonstrate the surprising advantage of the BTK inhibitor of Formula (XVIII), which does not interfere with thrombus formation, while the BTK inhibitors of Formula XXVII (CC-292) and Formula XX-A (ibrutinib) interfere with thrombus formation.

[00168] The objective of this study was to evaluate *in vivo* thrombus formation in the presence

of BTK inhibitors. *In vivo* testing of novel antiplatelet agents requires informative biomarkers. By utilizing a genetic modified mouse von Willebrand factor (VWFR1326H) model that supports human but not mouse platelet-mediated thrombosis, we evaluated the effects of Formula (XVIII), Formula XXVII (CC-292), and Formula XX-A (ibrutinib) on thrombus formation. These results show that Formula (XVIII) had no significant effect on human platelet-mediated thrombus formation while Formula XX-A (ibrutinib) was able to limit this process, resulting in a reduction in maximal thrombus size by 61% compared with control. Formula XXVII (CC-292) showed an effect similar to Formula XX-A (ibrutinib). These results, which show reduced thrombus formation for ibrutinib at physiologically relevant concentrations, may provide some mechanistic background for the Grade ≥ 3 bleeding events (eg, subdural hematoma, gastrointestinal bleeding, hematuria and postprocedural hemorrhage) that have been reported in $\leq 6\%$ of patients treated with Formula XX-A (ibrutinib).

[00169] GPVI platelet aggregation was measured for Formula (XVIII) and Formula XX-A (ibrutinib). Blood was obtained from untreated humans, and platelets were purified from plasma-rich protein by centrifugation. Cells were resuspended to a final concentration of 350,000/ μ L in buffer containing 145 mmol/L NaCl, 10 mmol/L HEPES, 0.5 mmol/L Na_2HPO_4 , 5 mmol/L KCl, 2 mmol/L MgCl_2 , 1 mmol/L CaCl_2 , and 0.1% glucose, at pH 7.4. Stock solutions of Convulxin (CVX) GPVI were prepared on the day of experimentation and added to platelet suspensions 5 minutes (37 °C, 1200 rpm) before the induction of aggregation. Aggregation was assessed with a Chronolog Lumi-Aggregometer (model 540 VS; Chronolog, Havertown, PA) and permitted to proceed for 6 minutes after the addition of agonist. The results are reported as maximum percent change in light transmittance from baseline with platelet buffer used as a reference. The results are shown in FIG. 59.

[00170] In FIG. 60, the results of CVX-induced (250 ng/mL) human platelet aggregation results before and 15 min after administration of the BTK inhibitors to 6 healthy individuals are shown.

[00171] The results depicted in FIG. 59 and FIG. 60 indicate that the BTK inhibitor of Formula XX-A (ibrutinib) significantly inhibits GPVI platelet aggregation, while the BTK inhibitor of Formula (XVIII) does not, further illustrating the surprising benefits of the latter compound.

Example 7 – Study of a BTK Inhibitor and a Combination of a BTK Inhibitor and a PI3K Inhibitor in Canine Lymphoma

[00172] Canine B cell lymphoma exists as a pathological entity that is characterized by large anaplastic, centroblastic or immunoblastic lymphocytes with high proliferative grade, significant peripheral lymphadenopathy and an aggressive clinical course. While some dogs respond initially to prednisone, most canine lymphomas progress quickly and must be treated with combination therapies, including cyclophosphamide, vincristine, doxorubicin, and prednisone (CHOP), or other cytotoxic agents. In their histopathologic features, clinical course, and high relapse rate after initial treatment, canine B cell lymphomas resemble diffuse large B cell lymphoma (DLBCL) in humans. Thus, responses of canine B cell lymphomas to experimental treatments are considered to provide proof of concept for therapeutic candidates in DLBCL.

[00173] In this example, companion dogs with newly diagnosed or relapsed/refractory LSA were enrolled on a veterinary clinical trial of the BTK inhibitor of Formula (XVIII) (“Arm 1”) or the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) (“Arm 2”). Enrollment has completed for Arm 1 and is ongoing for Arm 2. With approximately 1/3 of Arm 2 subjects treated, the preliminary results show that combined treatment with the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) may have greater efficacy than treatment with the BTK inhibitor of Formula (XVIII) alone in aggressive lymphoma.

[00174] Twenty-one dogs were treated in Arm 1 with the BTK inhibitor of Formula (XVIII) at dosages of 2.5 mg/kg once daily to 20 mg/kg twice daily. Intra-subject dose escalation was allowed. Six of the 11 dogs that initiated at 2.5 or 5 mg/kg once daily were escalated and completed the study with dosages of 10 mg/kg twice daily. Among all the dose cohorts, 8 dogs had shrinkage of target lesions >20%; the best tumor responses were between 45-49% reduction in the sum of target lesions in two dogs. Complete responses (“CR”, disappearance of all evidence of disease per evaluator judgment; and absence of new lesions) were not observed in Arm 1.

[00175] In the combination phase of the study (Arm 2), 7 dogs have been treated with 10 mg/kg the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) at 2.5 or 3.5 mg/kg, on a twice daily schedule. To date, 4 dogs had shrinkage of target lesions > 20%; and the best tumor responses were between 58-65% reduction in the sum of target lesions, with one

sustained CR observed. Initial reductions in the sum of target lesions were observed to deepen during the course of therapy in 4 of the 7 dogs. A summary of the results is presented in Table 5.

TABLE 5. Summary of the results of the canine lymphoma study.

Response Metric	Formula (XVIII) and Formula (IX)^a	Formula (XVIII) monotherapy
Sum LD ^b decreased by $\geq 20\%$	4/7 (57.1 %)	8/21 (38.1 %)
Sum LD ^b decreased by $\geq 30\%$ (PR)	2/7 (28.6 %)	6/21 (28.6 %)
CR by investigator evaluation	1/7 (14.3 %)	0/21
Median time on study (all subjects)	25 days	24 days
Median time to best response	21 days	7 days

^a Arm 2 is still recruiting subjects

^b LD, longest diameter of up to 5 target lesion


[00305] These preliminary data suggest that in companion dogs with naturally occurring B cell lymphomas, treatment with the combination of the BTK inhibitor of Formula (XVIII) and the PI3K- δ inhibitor of Formula (IX) may provide increased biological activity (tumor shrinkage and stable disease) and may possibly lead to deeper responses than treatment with the BTK inhibitor of Formula (XVIII) alone. Although the available data represent only 1/3 of the planned Arm 2 population, the extended response time (median time to best response) and observation of a CR among the few dogs treated to date may be evidence of synergy between Formula (XVIII) and Formula (IX) in this highly aggressive disease.

CLAIMS

We claim:

1. A method of treating a hyperproliferative disorder in a subject, comprising co-administering to a subject in need thereof a therapeutically effective amount of a phosphoinositide 3-kinase (PI3K) inhibitor, or of a pharmaceutically acceptable salt thereof, in combination with a Bruton's tyrosine kinase (BTK) inhibitor, or of a pharmaceutically acceptable salt thereof.
2. The method of Claim 1, wherein the PI3K inhibitor is selected from the group consisting of a PI3K- γ inhibitor, a PI3K- δ inhibitor, and a PI3K- γ,δ inhibitor.
3. The method of claim 2, wherein the PI3K inhibitor is a PI3K- γ,δ inhibitor.
4. The method of any one of claims 1 to 3, wherein the hyperproliferative disorder is leukemia, lymphoma, or a solid tumor cancer.
5. The method of any one of Claims 1 to 4, wherein the solid tumor cancer is selected from the group consisting of breast, lung, colorectal, thyroid, bone sarcoma and stomach cancers.
6. The method of any one of Claims 1 to 5, wherein the combination of the PI3K inhibitor with the BTK inhibitor is administered by oral, intravenous, intramuscular, intraperitoneal, subcutaneous or transdermal means.
7. The method of any one of Claims 1 to 6, wherein the PI3K inhibitor and/or BTK inhibitor is in the form of a pharmaceutically acceptable salt.
8. The method of any one of Claims 1 to 7, wherein the PI3K inhibitor is administered to the subject before administration of the BTK inhibitor.
9. The method of any one of Claims 1 to 7, wherein the PI3K inhibitor is administered concurrently with the administration of the BTK inhibitor.

10. The method of any one of Claims 1 to 7, wherein the PI3K inhibitor is administered to the subject after administration of the BTK inhibitor.
11. The method of any one of Claims 1 to 10, wherein the PI3K inhibitor is:



or a pharmaceutically-acceptable salt thereof, wherein:

X^1 is $C(R^9)$ or N;

X^2 is $C(R_{10})$ or N;

Y is $N(R^{11})$, O or S;

Z is CR^8 or N;

n is 0, 1, 2 or 3;

R^1 is a direct-bonded or oxygen-linked saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0 or 1 R^2 substituents, and the ring is additionally substituted by 0, 1, 2 or 3 substituents independently selected from halo, nitro, cyano, C_{1-4} alkyl, OC_{1-4} alkyl, OC_{1-4} haloalkyl, NHC_{1-4} , $N(C_{1-4}alkyl)C_{1-4}alkyl$ and C_{1-4} haloalkyl;

R^2 is selected from halo, C_{1-4} haloalkyl, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^a$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $OS(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-$

- $N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$; or R^2 is selected from $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl, heterocycle, $-(C_{1-3}alkyl)heteroaryl$, $-(C_{1-3}alkyl)heterocycle$, $-O(C_{1-3}alkyl)heteroaryl$, $-O(C_{1-3}alkyl)heterocycle$, $-NR^a(C_{1-3}alkyl)heteroaryl$, $-NR^a(C_{1-3}alkyl)heterocycle$, $-(C_{1-3}alkyl)phenyl$, $-O(C_{1-3}alkyl)phenyl$ and $-NR^a(C_{1-3}alkyl)phenyl$ all of which are substituted by 0, 1, 2 or 3 substituents selected from $C_{1-4}haloalkyl$, $OC_{1-4}alkyl$, Br, Cl, F, I and $C_{1-4}alkyl$;
- R^3 is selected from H, halo, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^a$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^2$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^a$, $-N(R^a)C(=O)OR^a$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from $C_{1-6}haloalkyl$, $OC_{1-6}alkyl$, Br, Cl, F, I and $C_{1-6}alkyl$;
- R^4 is, independently, in each instance, halo, nitro, cyano, $C_{1-4}alkyl$, $OC_{1-4}alkyl$, $OC_{1-4}haloalkyl$, $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$ or $C_{1-4}haloalkyl$;
- R^5 is, independently, in each instance, H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, or $C_{1-6}alkyl$ substituted by 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$; or both R^5 groups together form a $C_{3-6}spiroalkyl$ substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, OH, $OC_{1-4}alkyl$, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $OC_{1-4}alkyl$, NH_2 , $NHC_{1-4}alkyl$, $N(C_{1-4}alkyl)C_{1-4}alkyl$;
- R^6 is selected from H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;
- R^7 is selected from H, halo, $C_{1-6}alkyl$, $C_{1-4}haloalkyl$, cyano, nitro, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^a$, $-S(=O)_2N(R^a)C(=O)OR^a$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$;
- R^8 is selected from H, $C_{1-6}haloalkyl$, Br, Cl, F, I, OR^a , NR^aR^a , $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle, wherein the $C_{1-6}alkyl$, phenyl, benzyl, heteroaryl and heterocycle

are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₆haloalkyl, OC₁₋₆alkyl, Br, Cl, F, I and C₁₋₆alkyl;


R⁹ is selected from H, halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^aC(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(O)NR^aR^aN(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a, —NR^aC₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle, wherein the C₁₋₆alkyl, phenyl, benzyl, heteroaryl and heterocycle are additionally substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylOR^a, —NR^aC₂₋₆alkylOR^a; or R⁹ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, but containing no more than one O or S, wherein the available carbon atoms of the ring are substituted by 0, 1 or 2 oxo or thioxo groups, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from halo, C₁₋₄haloalkyl, cyano, nitro, —C(=O)R^a, —C(=O)OR^a, —C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —OR^a, —OC(=O)R^a, —OC(=O)NR^aR^a, —OC(=O)N(R^a)S(=O)₂R^a, —OC₂₋₆alkylNR^aR^a, —OC₂₋₆alkylOR^a, —SR^a, —S(=O)R^a, —S(=O)₂R^a, —S(=O)₂NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —NR^aR^a, —N(R^a)C(=O)R^a, —N(R^a)C(=O)OR^a, —N(R^a)C(=O)NR^aR^a, —N(R^a)C(=NR^a)NR^aR^a, —N(R^a)S(=O)₂R^a, —N(R^a)S(=O)₂NR^aR^a, —NR^aC₂₋₆alkylNR^aR^a and —NR^aC₂₋₆alkylOR^a;

R¹⁰ is H, C₁₋₃alkyl, C₁₋₃haloalkyl, cyano, nitro, CO₂R^a, C(=O)NR^aR^a, —C(=NR^a)NR^aR^a, —S(=O)₂N(R^a)C(=O)R^a, —S(=O)₂N(R^a)C(=O)OR^a, —S(=O)₂N(R^a)C(=O)NR^aR^a, —S(=O)R^b, S(=O)₂R^b or S(=O)₂NR^aR^a;

R¹¹ is H or C₁₋₄alkyl;


R^a is independently, at each instance, H or R^b ; and R^b is independently, at each instance, phenyl, benzyl or C_{1-6} alkyl, the phenyl, benzyl and C_{1-6} alkyl being substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} alkyl, C_{1-3} haloalkyl, $-OC_{1-4}$ alkyl, $-NH_2$, $-NHC_{1-4}$ alkyl, $-N(C_{1-4}alkyl)C_{1-4}alkyl$.

12. The method of Claim 11, wherein the PI3K inhibitor is:



or a pharmaceutically acceptable salt thereof.

13. The method of any one of Claims 1 to 10, wherein the PI3K inhibitor is:



or an enantiomer, a mixture of enantiomers, or a mixture of two or more diastereomers thereof, or a pharmaceutically acceptable salt thereof, wherein

Cy is aryl or heteroaryl substituted by 0 or 1 occurrence of R^3 and 0, 1, 2, or 3 occurrence(s) of R^5 ;

W_b is CR^8 , CHR^8 , or N;

R^8 is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfonamido, halo, cyano, hydroxyl or nitro; B is hydrogen, alkyl, amino, heteroalkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl, each of which is substituted with 0, 1, 2, 3, or 4 occurrence(s) of R^2 ;

each R^2 is independently alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkoxy, amido, amino, acyl, acyloxy, alkoxycarbonyl, sulfonamido, halo, cyano, hydroxyl, nitro, phosphate, urea, or carbonate;

X is $-(CH(R^9))_z-$;

Y is $-N(R^9)-C(=O)-$, $-C(=O)-N(R^9)-$, $-C(=O)-N(R^9)-(CHR^9)-$, $-N(R^9)-S(=O)-$, $-S(=O)-N(R^9)-$, $-S(=O)_2-N(R^9)-$, $-N(R^9)-C(=O)-N(R^9)$ or $-N(R^9)S(=O)_2-$;

z is an integer of 1, 2, 3, or 4;

R^3 is alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, fluoroalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfinyl, sulfonyl, sulfoxide, sulfone, sulfonamido, halo, cyano, aryl, heteroaryl, hydroxyl, or nitro;


each R^5 is independently alkyl, alkenyl, alkynyl, cycloalkyl, heteroalkyl, alkoxy, amido, amino, acyl, acyloxy, sulfonamido, halo, cyano, hydroxyl, or nitro;

each R^9 is independently hydrogen, alkyl, cycloalkyl, heterocyclyl, or heteroalkyl; or two adjacent occurrences of R^9 together with the atoms to which they are attached form a 4- to 7-membered ring;

W_d is heterocyclyl, aryl, cycloalkyl, or heteroaryl, each of which is substituted with one or more R^{10} , R^{11} , R^{12} or R^{13} , and


R^{10} , R^{11} , R^{12} and R^{13} are each independently hydrogen, alkyl, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkoxy, heterocycloxy, amido, amino, acyl, acyloxy, alkoxycarbonyl, sulfonamido, halo, cyano, hydroxyl, nitro, phosphate, urea, carbonate or $NR'R''$ wherein R' and R'' are taken together with nitrogen to form a cyclic moiety.

14. The method of Claim 13, wherein the PI3K inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

B is:




wherein W_c is aryl, heteroaryl, heterocycloalkyl, or cycloalkyl, and







q is an integer of 0, 1, 2, 3, or 4;

X is a bond or $-(CH(R^9))_z-$, and z is an integer of 1;

Y is $-N(R^9)-$;

W_d is:





;


X₁, X₂ and X₃ are each independently C, CR¹³ or N; and X₄, X₅ and X₆ are each independently N, NH, CR¹³, S or O;

R¹ is hydrogen, alkyl, alkenyl, alkynyl, alkoxy, amido, alkoxy carbonyl, sulfonamido, halo, cyano, or nitro;

R² is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, heteroarylalkyl, alkoxy, amino, halo, cyano, hydroxy or nitro; R³ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, alkoxy, amido, amino, alkoxy carbonyl sulfonamido, halo, cyano, hydroxy or nitro; and


each instance of R⁹ is independently hydrogen, alkyl, or heterocycloalkyl.

15. The method of Claim 14, wherein the PI3K inhibitor is



or a pharmaceutically acceptable salt thereof.


16. The method of Claim 14, wherein the PI3K inhibitor is



Formula (IV)

or a pharmaceutically acceptable salt thereof.

17. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

X is CH, N, O or S;

Y is C(R₆), N, O or S;

Z is CH, N or bond;

A is CH or N;

B₁ is N or C(R₇);

B₂ is N or C(R₈);

B₃ is N or C(R₉);

B₄ is N or C(R₁₀);

R₁ is R₁₁C(O), R₁₂S(O), R₁₃SO₂ or (1-6C)alkyl optionally substituted with R₁₄;

R₂ is H, (1-3C)alkyl or (3-7C)cycloalkyl;

R₃ is H, (1-6C)alkyl or (3-7C)cycloalkyl); or

R₂ and R₃ form, together with the N and C atom they are attached to, a (3- 7C)heterocycloalkyl optionally substituted with one or more fluorine, hydroxyl, (1-3C)alkyl, (1-3C)alkoxy or oxo;

R₄ is H or (1-3C)alkyl;

R₅ is H, halogen, cyano, (1-4C)alkyl, (1-3C)alkoxy, (3-6C)cycloalkyl, any alkyl group of which is optionally substituted with one or more halogen; or R₅ is (6-10C)aryl or (2-6C)heterocycloalkyl;

R₆ is H or (1-3C)alkyl; or

R₅ and R₆ together may form a (3-7C)cycloalkenyl, or (2-6C)heterocycloalkenyl; each optionally substituted with (1-3C)alkyl, or one or more halogen;

R₇ is H, halogen, CF₃, (1-3C)alkyl or (1-3C)alkoxy;

R₈ is H, halogen, CF₃, (1-3C)alkyl or (1-3C)alkoxy; or

R₇ and R₈ together with the carbon atoms they are attached to, form (6-10C)aryl or (1-9C)heteroaryl;

R₉ is H, halogen, (1-3C)alkyl or (1-3C)alkoxy;

R₁₀ is H, halogen, (1-3C)alkyl or (1-3C)alkoxy;

R₁₁ is independently selected from a group consisting of (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl each alkyl, alkenyl or alkynyl optionally substituted with one or more groups selected from hydroxyl, (1-4C)alkyl, (3-7C)cycloalkyl, [(1-4C)alkyl]amino, di[(1-4C)alkyl]amino, (1-3C)alkoxy, (3-7C)cycloalkoxy, (6-10C)aryl or (3-7C)heterocycloalkyl; or R₁₁ is (1-3C)alkyl-C(O)-S-(1-3C)alkyl; or


R₁₁ is (1-5C)heteroaryl optionally substituted with one or more groups selected from halogen or cyano;

R₁₂ and R₁₃ are independently selected from a group consisting of (2-6C)alkenyl or (2-6C)alkynyl both optionally substituted with one or more groups selected from hydroxyl, (1-4C)alkyl, (3-7C)cycloalkyl, [(1-4C)alkyl]amino, di[(1-4C)alkyl]amino, (1-3C)alkoxy, (3-7C)cycloalkoxy, (6-10C)aryl or (3-7C)heterocycloalkyl; or

(1-5C)heteroaryl optionally substituted with one or more groups selected from halogen or cyano; and

R₁₄ is independently selected from a group consisting of halogen, cyano or (2-6C)alkenyl or (2-6C)alkynyl both optionally substituted with one or more groups selected from hydroxyl, (1-4C)alkyl, (3-7C)cycloalkyl, (1-4C)alkylamino, di[(1-4C)alkyl]amino, (1-3C)alkoxy, (3-7C)cycloalkoxy, (6-10C)aryl, (1-5C)heteroaryl or (3-7C)heterocycloalkyl.

18. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$, $NRS(=O)_x$, where x is 1 or 2;

R_7 and R_8 are each H; or R_7 and R_8 taken together form a bond;

R_6 is H; and

R is H or C_1 - C_6 alkyl.

19. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$, $NRS(=O)_x$, where x is 1 or 2;

R_7 and R_8 are each H; or R_7 and R_8 taken together form a bond;

R_6 is H; and

R is H or C_1 - C_6 alkyl.

20. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$, $NRS(=O)_x$, where x is 1 or 2;

R_7 and R_8 are each H; or R_7 and R_8 taken together form a bond;

R_6 is H; and

R is H or C_1 - C_6 alkyl.

21. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof, wherein

L_a is CH_2 , O, NH or S;

Ar is a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl;

Y is an optionally substituted group selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl;


Z is $C(=O)$, $OC(=O)$, $NRC(=O)$, $C(=S)$, $S(=O)_x$, $OS(=O)_x$, $NRS(=O)_x$, where x is 1 or 2;

R_7 and R_8 are each H; or R_7 and R_8 taken together form a bond;

R_6 is H; and

R is H or C_1 - C_6 alkyl.

22. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically acceptable salt thereof,

wherein:

Q^1 is aryl, heteroaryl, cycloalkyl, heterocyclyl, cycloalkenyl, or heterocycloalkenyl, any of which is optionally substituted by one to five independent G^1 substituents;

R^1 is alkyl, cycloalkyl, bicycloalkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heterocyclyl, or heterobicycloalkyl, any of which is optionally substituted by one or more independent G^{11} substituents;

G^1 and G^{41} are each independently halo, oxo, $-CF_3$, $-OCF_3$, $-OR^2$, $-NR^2R^3(R^{3a})_{j1}$, $-C(O)R^2$, $-CO_2R^2$, $-CONR^2R^3$, $-NO_2$, $-CN$, $-S(O)_{j1}R^2$, $-SO_2NR^2R^3$, $NR^2(C=O)R^3$, $NR^2(C=O)OR^3$, $NR^2(C=O)NR^2R^3$, $NR^2S(O)_{j1}R^3$, $-(C=S)OR^2$, $-(C=O)SR^2$, $-NR^2(C=NR^3)NR^{2a}R^{3a}$, $-NR^2(C=NR^3)OR^{2a}$, $-NR^2(C=NR^3)SR^{3a}$, $-O(C=O)OR^2$, $-O(C=O)NR^2R^3$, $-O(C=O)SR^2$, $-S(C=O)OR^2$, $-S(C=O)NR^2R^3$, $C_{0-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, $C_{1-10}alkoxyC_{1-10}alkyl$, $C_{1-10}alkoxyC_{2-10}alkenyl$, $C_{1-10}alkoxyC_{2-10}alkynyl$, $C_{1-10}alkylthioC_{1-10}alkyl$, $C_{1-10}alkylthioC_{2-10}alkenyl$, $C_{1-10}alkylthioC_{2-10}alkynyl$, $cycloC_{3-8}alkyl$, $cycloC_{3-8}alkenyl$, $cycloC_{3-8}alkylC_{1-10}alkyl$, $cycloC_{3-8}alkenylC_{1-10}alkyl$, $cycloC_{3-8}alkylC_{2-10}alkenyl$, $cycloC_{3-8}alkenylC_{2-10}alkenyl$, $cycloC_{3-8}alkylC_{2-10}alkynyl$, $cycloC_{3-8}alkenylC_{2-10}alkynyl$, $heterocyclyl-C_{0-10}alkyl$, $heterocyclyl-C_{2-10}alkenyl$, or $heterocyclyl-C_{2-10}alkynyl$, any of which is optionally substituted with one or more independent halo, oxo, $-CF_3$, $-OCF_3$, $-OR^{222}$, $-NR^{222}R^{333}(R^{333a})_{j1a}$, $-C(O)R^{222}$, $-CO_2R^{222}$, $-CONR^{222}R^{333}$, $-NO_2$, $-CN$, $-S(O)_{j1a}R^{222}$, $-SO_2NR^{222}R^{333}$, $NR^{222}(C=O)R^{333}$, $NR^{222}(C=O)OR^{333}$, $NR^{222}(C=O)NR^{222}R^{333}$, $NR^{222}S(O)_{j1a}R^{333}$, $-(C=S)OR^{222}$, $-(C=O)SR^{222}$, $-NR^{222}(C=NR^{333})NR^{222a}R^{333a}$, $-NR^{222}(C=NR^{333})OR^{222a}$, $-NR^{222}(C=NR^{333})SR^{333a}$, $-O(C=O)OR^{222}$, $-O(C=O)NR^{222}R^{333}$, $-O(C=O)SR^{222}$, $-S(C=O)OR^{222}$, or $-S(C=O)NR^{222}R^{333}$ substituents; or $-(X^1)_n-(Y^1)_m-R^4$; or aryl- $C_{0-10}alkyl$, aryl- $C_{2-10}alkenyl$, or aryl- $C_{2-10}alkynyl$, any of which is optionally substituted with one or more independent halo, $-CF_3$, $-OCF_3$, $-OR^{222}$, $-NR^{222}R^{333}(R^{333a})_{j2a}$, $-C(O)R^{222}$, $-CO_2R^{222}$, $-CONR^{222}R^{333}$, $-NO_2$, $-CN$, $-S(O)_{j2a}R^{222}$, $-SO_2NR^{222}R^{333}$, $NR^{222}(C=O)R^{333}$, $NR^{222}(C=O)OR^{333}$, $NR^{222}(C=O)NR^{222}R^{333}$, $NR^{222}S(O)_{j2a}R^{333}$, $-(C=S)OR^{222}$, $-(C=O)SR^{222}$, $-NR^{222}(C=NR^{333})NR^{222a}R^{333a}$, $-NR^{222}(C=NR^{333})OR^{222a}$, $-NR^{222}(C=NR^{333})SR^{333a}$, $-O(C=O)OR^{222}$, $-O(C=O)NR^{222}R^{333}$, $-O(C=O)SR^{222}$, $-S(C=O)OR^{222}$, or $-S(C=O)NR^{222}R^{333}$ substituents; or hetaryl- $C_{0-10}alkyl$, hetaryl- $C_{2-10}alkenyl$, or hetaryl- $C_{2-10}alkynyl$, any of which is optionally

substituted with one or more independent halo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{222}$, $-\text{NR}^{222}$, $\text{R}^{333}(\text{R}^{333a})_{j3a}$, $-\text{C}(\text{O})\text{R}^{222}$, $-\text{CO}_2\text{R}^{222}$, $-\text{CONR}^{222}\text{R}^{333}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_{j3a}\text{R}^{222}$, $-\text{SO}_2\text{NR}^{222}\text{R}^{333}$, $\text{NR}^{222}(\text{C}=\text{O})\text{R}^{333}$, $\text{NR}^{222}(\text{C}=\text{O})\text{OR}^{333}$, $\text{NR}^{222}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$, $\text{NR}^{222}\text{S}(\text{O})_{j3a}\text{R}^{333}$, $-(\text{C}=\text{S})\text{OR}^{222}$, $-(\text{C}=\text{O})\text{SR}^{222}$, $-\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{NR}^{222a}\text{R}^{333a}$, $-\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{OR}^{222a}$, $-\text{NR}^{222}(\text{C}=\text{NR}^{333})\text{SR}^{333a}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{222}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{222}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{222}$, or $-\text{S}(\text{C}=\text{O})\text{NR}^{222}\text{R}^{333}$ substituents;

G^{11} is halo, oxo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{21}$, $-\text{NR}^{21}\text{R}^{31}(\text{R}^{3a1})_{j4}$, $-\text{C}(\text{O})\text{R}^{21}$, $-\text{CO}_2\text{R}^{21}$, $-\text{CONR}^{21}\text{R}^{31}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_{j4}\text{R}^{21}$, $-\text{SO}_2\text{NR}^{21}\text{R}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{R}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{OR}^{31}$, $\text{NR}^{21}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$, $\text{NR}^{21}\text{S}(\text{O})_{j4}\text{R}^{31}$, $-(\text{C}=\text{S})\text{OR}^{21}$, $-(\text{C}=\text{O})\text{SR}^{21}$, $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{NR}^{2a1}\text{R}^{3a1}$, $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{OR}^{2a1}$, $-\text{NR}^{21}(\text{C}=\text{NR}^{31})\text{SR}^{3a1}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{21}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{21}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{21}$, $-\text{S}(\text{C}=\text{O})\text{NR}^{21}\text{R}^{31}$, $-\text{P}(\text{O})\text{OR}^{21}\text{OR}^{31}$, $\text{C}_{0-10}\text{alkyl}$, $\text{C}_{2-10}\text{alkenyl}$, $\text{C}_{2-10}\text{alkynyl}$, $\text{C}_{1-10}\text{alkoxyC}_{1-10}\text{alkyl}$, $\text{C}_{1-10}\text{alkoxyC}_{2-10}\text{alkenyl}$, $\text{C}_{1-10}\text{alkoxyC}_{2-10}\text{alkynyl}$, $\text{C}_{1-10}\text{alkylthioC}_{1-10}\text{alkyl}$, $\text{C}_{1-10}\text{alkylthioC}_{2-10}\text{alkenyl}$, $\text{C}_{1-10}\text{alkylthioC}_{2-10}\text{alkynyl}$, $\text{cycloC}_{3-8}\text{alkyl}$, $\text{cycloC}_{3-8}\text{alkenyl}$, $\text{cycloC}_{3-8}\text{alkylC}_{1-10}\text{alkyl}$, $\text{cycloC}_{3-8}\text{alkenylC}_{1-10}\text{alkyl}$, $\text{cycloC}_{3-8}\text{alkylC}_{2-10}\text{alkenyl}$, $\text{cycloC}_{3-8}\text{alkenylC}_{2-10}\text{alkenyl}$, $\text{cycloC}_{3-8}\text{alkylC}_{2-10}\text{alkynyl}$, $\text{cycloC}_{3-8}\text{alkenylC}_{2-10}\text{alkynyl}$, $\text{heterocyclyl-C}_{0-10}\text{alkyl}$, $\text{heterocyclyl-C}_{2-10}\text{alkenyl}$, or $\text{heterocyclyl-C}_{2-10}\text{alkynyl}$, any of which is optionally substituted with one or more independent halo, oxo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j4a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_{j4a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{OR}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}\text{S}(\text{O})_{j4a}\text{R}^{3331}$, $-(\text{C}=\text{S})\text{OR}^{2221}$, $-(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{NR}^{222a1}\text{R}^{333a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{OR}^{222a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{SR}^{333a1}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{P}(\text{O})\text{OR}^{2221}\text{OR}^{3331}$, or $-\text{S}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$ substituents; or $\text{aryl-C}_{0-10}\text{alkyl}$, $\text{aryl-C}_{2-10}\text{alkenyl}$, or $\text{aryl-C}_{2-10}\text{alkynyl}$, any of which is optionally substituted with one or more independent halo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j5a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_{j5a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{R}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{OR}^{3331}$, $\text{NR}^{2221}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $\text{NR}^{2221}\text{S}(\text{O})_{j5a}\text{R}^{3331}$, $-(\text{C}=\text{S})\text{OR}^{2221}$, $-(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{NR}^{222a1}\text{R}^{333a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{OR}^{222a1}$, $-\text{NR}^{2221}(\text{C}=\text{NR}^{3331})\text{SR}^{333a1}$, $-\text{O}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{O}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$, $-\text{O}(\text{C}=\text{O})\text{SR}^{2221}$, $-\text{S}(\text{C}=\text{O})\text{OR}^{2221}$, $-\text{P}(\text{O})\text{OR}^{2221}\text{R}^{3331}$, or $-\text{S}(\text{C}=\text{O})\text{NR}^{2221}\text{R}^{3331}$ substituents; or $\text{hetaryl-C}_{0-10}\text{alkyl}$, $\text{hetaryl-C}_{2-10}\text{alkenyl}$, or $\text{hetaryl-C}_{2-10}\text{alkynyl}$, any of which is optionally substituted with one or more independent halo, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{2221}$, $-\text{NR}^{2221}\text{R}^{3331}(\text{R}^{333a1})_{j6a}$, $-\text{C}(\text{O})\text{R}^{2221}$, $-\text{CO}_2\text{R}^{2221}$, $-\text{CONR}^{2221}\text{R}^{3331}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_{j6a}\text{R}^{2221}$, $-\text{SO}_2\text{NR}^{2221}\text{R}^{3331}$,


NR²²²¹(C=O)R³³³¹, NR²²²¹(C=O)OR³³³¹, NR²²²¹(C=O)NR²²²¹R³³³¹, NR²²²¹S(O)_{j6a}R³³³¹,
 -(C=S)OR²²²¹, -(C=O)SR²²²¹, -NR²²²¹(C=NR³³³¹)NR^{222a1}R^{333a1}, -NR²²²¹(C=NR³³³¹)OR^{222a1},
 -NR²²²¹(C=NR³³³¹)SR^{333a1}, -O(C=O)OR²²²¹, -O(C=O)NR²²²¹R³³³¹, -O(C=O)SR²²²¹,
 -S(C=O)OR²²²¹, -P(O)OR²²²¹OR³³³¹, or -S(C=O)NR²²²¹R³³³¹ substituents; or G¹¹¹ is taken
 together with the carbon to which it is attached to form a double bond which is substituted with
 R⁵ and G¹¹¹;

R², R^{2a}, R³, R^{3a}, R²²², R^{222a}, R³³³, R^{333a}, R²¹, R^{2a1}, R³¹, R^{3a1}, R²²²¹, R^{222a1}, R³³³¹, and R^{333a1} are
 each independently equal to C₀₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₁₋₁₀alkoxyC₁₋₁₀alkyl, C₁₋₁₀
 alkoxyC₂₋₁₀alkenyl, C₁₋₁₀alkoxyC₂₋₁₀alkynyl, C₁₋₁₀alkylthioC₁₋₁₀alkyl, C₁₋₁₀alkylthioC₂₋₁₀
 alkenyl, C₁₋₁₀alkylthioC₂₋₁₀alkynyl, cycloC₃₋₈alkyl, cycloC₃₋₈alkenyl, cycloC₃₋₈alkylC₁₋₁₀alkyl,
 cycloC₃₋₈alkenylC₁₋₁₀alkyl, cycloC₃₋₈alkylC₂₋₁₀alkenyl, cycloC₃₋₈alkenylC₂₋₁₀alkenyl, cycloC₃₋₈
 alkylC₂₋₁₀alkynyl, cycloC₃₋₈alkenylC₂₋₁₀alkynyl, heterocyclyl-C₀₋₁₀alkyl, heterocyclyl-C₂₋₁₀
 alkenyl, or heterocyclyl-C₂₋₁₀alkynyl, any of which is optionally substituted by one or more
 G¹¹¹ substituents; or aryl-C₀₋₁₀alkyl, aryl-C₂₋₁₀alkenyl, or aryl-C₂₋₁₀alkynyl, hetaryl-C₀₋₁₀alkyl,
 hetaryl-C₂₋₁₀alkenyl, or hetaryl-C₂₋₁₀alkynyl, any of which is optionally substituted by one or
 more G¹¹¹ substituents; or in the case of -NR²R³(R^{3a})_{j1} or -NR²²²R³³³(R^{333a})_{j1a} or
 -NR²²²R³³³(R^{333a})_{j2a} or -NR²²²¹R³³³¹(R^{333a1})_{j3a} or -NR²²²¹R³³³¹(R^{333a1})_{j4a} or -NR²²²¹R³³³¹(R^{333a1})_{j5a}
 or -NR²²²¹R³³³¹(R^{333a1})_{j6a}, R² and R³ or R²²² and R³³³ or R²²²¹ and R³³³¹ taken together with the
 nitrogen atom to which they are attached form a 3-10 membered saturated ring, unsaturated ring,
 heterocyclic saturated ring, or heterocyclic unsaturated ring, wherein said ring is optionally
 substituted by one or more G¹¹¹ substituents;

X¹ and Y¹ are each independently -O-, -NR⁷-, -S(O)_{j7}-, -CR⁵R⁶-, -N(C(O)OR⁷)-, -N(C(O)R⁷)-,
 -N(SO₂R⁷)-, -CH₂O-, -CH₂S-, -CH₂N(R⁷)-, -CH(NR⁷)-, -CH₂N(C(O)R⁷)-, -CH₂N(C(O)OR⁷)-,
 -CH₂N(SO₂R⁷)-, -CH(NHR⁷)-, -CH(NHC(O)R⁷)-, -CH(NHSO₂R⁷)-, -CH(NHC(O)OR⁷)-,
 -CH(OC(O)R⁷)-, -CH(OC(O)NHR⁷)-, -CH=CH-, -C.ident.C-, -C(=NOR⁷)-, -C(O)-, -CH(OR⁷)-,
 -C(O)N(R⁷)-, -N(R⁷)C(O)-, -N(R⁷)S(O)-, -N(R⁷)S(O)₂- -OC(O)N(R⁷)-, -N(R⁷)C(O)N(R⁷)-,
 -NR⁷C(O)O-, -S(O)N(R⁷)-, -S(O)₂N(R⁷)-, -N(C(O)R⁷)S(O)-, -N(C(O)R⁷)S(O)₂-,
 -N(R⁷)S(O)N(R⁷)-, -N(R⁷)S(O)₂N(R⁷)-, -C(O)N(R⁷)C(O)-, -S(O)N(R⁷)C(O)-, -S(O)₂N(R⁷)C(O)-,
 -OS(O)N(R⁷)-, -OS(O)₂N(R⁷)-, -N(R⁷)S(O)O-, -N(R⁷)S(O)₂O-, -N(R⁷)S(O)C(O)-,
 -N(R⁷)S(O)₂C(O)-, -SON(C(O)R⁷)-, -SO₂N(C(O)R⁷)-, -N(R⁷)SON(R⁷)-, -N(R⁷)SO₂N(R⁷)-,

-C(O)O-, -N(R⁷)P(OR⁸)O-, -N(R⁷)P(OR⁸)-, -N(R⁷)P(O)(OR⁸)O-, -N(R⁷)P(O)(OR⁸)-,
 -N(C(O)R⁷)P(OR⁸)O-, -N(C(O)R⁷)P(OR⁸)-, -N(C(O)R⁷)P(O)(OR⁸)O-, -N(C(O)R⁷)P(OR⁸)-,
 -CH(R⁷)S(O)-, -CH(R⁷)S(O)₂-, -CH(R⁷)N(C(O)OR⁷)-, -CH(R⁷)N(C(O)R⁷)-, -CH(R⁷)N(SO₂R⁷)-,
 -CH(R⁷)O-, -CH(R⁷)S-, -CH(R⁷)N(R⁷)-, -CH(R⁷)N(C(O)R⁷)-, -CH(R⁷)N(C(O)OR⁷)-,
 -CH(R⁷)N(SO₂R⁷)-, -CH(R⁷)C(=NOR⁷)-, -CH(R⁷)C(O)-, -CH(R⁷)CH(OR⁷)-,
 -CH(R⁷)C(O)N(R⁷)-, -CH(R⁷)N(R⁷)C(O)-, -CH(R⁷)N(R⁷)S(O)-, -CH(R⁷)N(R⁷)S(O)₂-,
 -CH(R⁷)OC(O)N(R⁷)-, -CH(R⁷)N(R⁷)C(O)N(R⁷)-, -CH(R⁷)NR⁷C(O)O-, -CH(R⁷)S(O)N(R⁷)-,
 -CH(R⁷)S(O)₂N(R⁷)-, -CH(R⁷)N(C(O)R⁷)S(O)-, -CH(R⁷)N(C(O)R⁷)S(O)-,
 -CH(R⁷)N(R⁷)S(O)N(R⁷)-, -CH(R⁷)N(R⁷)S(O)₂N(R⁷)-, -CH(R⁷)C(O)N(R⁷)C(O)-,
 -CH(R⁷)S(O)N(R⁷)C(O)-, -CH(R⁷)S(O)₂N(R⁷)C(O)-, -CH(R⁷)OS(O)N(R⁷)-,
 -CH(R⁷)OS(O)₂N(R⁷)-, -CH(R⁷)N(R⁷)S(O)O-, -CH(R⁷)N(R⁷)S(O)₂O-, -CH(R⁷)N(R⁷)S(O)C(O)-,
 -CH(R⁷)N(R⁷)S(O)₂C(O)-, -CH(R⁷)SON(C(O)R⁷)-, -CH(R⁷)SO₂N(C(O)R⁷)-,
 -CH(R⁷)N(R⁷)SON(R⁷)-, -CH(R⁷)N(R⁷)SO₂N(R⁷)-, -CH(R⁷)C(O)O-, -CH(R⁷)N(R⁷)P(OR⁸)O-,
 -CH(R⁷)N(R⁷)P(OR⁸)-, -CH(R⁷)N(R⁷)P(O)(OR⁸)O-, -CH(R⁷)N(R⁷)P(O)(OR⁸)-,
 -CH(R⁷)N(C(O)R⁷)P(OR⁸)O-, -CH(R⁷)N(C(O)R⁷)P(OR⁸)-, -CH(R⁷)N(C(O)R⁷)P(O)(OR⁸)O-, or
 -CH(R⁷)N(C(O)R⁷)P(OR⁸)-; or

X¹ and Y¹ are each independently represented by one of the following structural formulas:



R¹⁰, taken together with the phosphinamide or phosphonamide, is a 5-, 6-, or 7-membered aryl, heteroaryl or heterocyclyl ring system;

R⁵, R⁶, and G¹¹¹ are each independently a C₀₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₁₋₁₀alkoxyC₁₋₁₀alkyl, C₁₋₁₀alkoxyC₂₋₁₀alkenyl, C₁₋₁₀alkoxyC₂₋₁₀alkynyl, C₁₋₁₀alkylthioC₁₋₁₀alkyl, C₁₋₁₀alkylthioC₂₋₁₀alkenyl, C₁₋₁₀alkylthioC₂₋₁₀alkynyl, cycloC₃₋₈alkyl, cycloC₃₋₈alkenyl, cycloC₃₋₈alkylC₁₋₁₀alkyl, cycloC₃₋₈alkenylC₁₋₁₀alkyl, cycloC₃₋₈alkylC₂₋₁₀alkenyl, cycloC₃₋₈alkenylC₂₋₁₀alkenyl, cycloC₃₋₈alkylC₂₋₁₀alkynyl, cycloC₃₋₈alkenylC₂₋₁₀alkynyl, heterocyclyl-C₀₋₁₀alkyl, heterocyclyl-C₂₋₁₀alkenyl, or heterocyclyl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, -CF₃, -OCF₃, -OR⁷⁷, -NR⁷⁷R⁸⁷, -C(O)R⁷⁷, -CO₂R⁷⁷, -CONR⁷⁷R⁸⁷, -NO₂, -CN, -S(O)_{j5a}R⁷⁷, -SO₂NR⁷⁷R⁸⁷, NR⁷⁷(C=O)R⁸⁷, NR⁷⁷(C=O)OR⁸⁷, NR⁷⁷(C=O)NR⁷⁸R⁸⁷, NR⁷⁷S(O)_{j5a}R⁸⁷, -(C=S)OR⁷⁷, -(C=O)SR⁷⁷, -NR⁷⁷(C=NR⁸⁷)NR⁷⁸R⁸⁸, -NR⁷⁷(C=NR⁸⁷)OR⁷⁸, -NR⁷⁷(C=NR⁸⁷)SR⁷⁸, -O(C=O)OR⁷⁷, -O(C=O)NR⁷⁷R⁸⁷, -O(C=O)SR⁷⁷, -S(C=O)OR⁷⁷, -P(O)OR⁷⁷OR⁸⁷, or -S(C=O)NR⁷⁷R⁸⁷ substituents; or aryl-C₀₋₁₀alkyl, aryl-C₂₋₁₀alkenyl, or aryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, -CF₃, -OCF₃, -OR⁷⁷, -NR⁷⁷R⁸⁷, -C(O)R⁷⁷, -CO₂R⁷⁷, -CONR⁷⁷R⁸⁷, -NO₂, -CN, -S(O)_{j5a}R⁷⁷, -SO₂NR⁷⁷R⁸⁷, NR⁷⁷(C=O)R⁸⁷, NR⁷⁷(C=O)OR⁸⁷, NR⁷⁷(C=O)NR⁷⁸R⁸⁷, NR⁷⁷S(O)_{j5a}R⁸⁷, -(C=S)OR⁷⁷, -(C=O)SR⁷⁷, -NR⁷⁷(C=NR⁸⁷)NR⁷⁸R⁸⁸, -NR⁷⁷(C=NR⁸⁷)OR⁷⁸, -NR⁷⁷(C=NR⁸⁷)SR⁷⁸, -O(C=O)OR⁷⁷, -O(C=O)NR⁷⁷R⁸⁷, -O(C=O)SR⁷⁷, -S(C=O)OR⁷⁷, -P(O)OR⁷⁷R⁸⁷, or -S(C=O)NR⁷⁷R⁸⁷ substituents; or hetaryl-C₀₋₁₀alkyl, hetaryl-C₂₋₁₀alkenyl, or hetaryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, -CF₃, -OCF₃, -OR⁷⁷, -NR⁷⁷R⁸⁷, -C(O)R⁷⁷, -CO₂R⁷⁷, -CONR⁷⁷R⁸⁷, -NO₂, -CN, -S(O)_{j5a}R⁷⁷, -SO₂NR⁷⁷R⁸⁷, NR⁷⁷(C=O)R⁸⁷, NR⁷⁷(C=O)OR⁸⁷, NR⁷⁷(C=O)NR⁷⁸R⁸⁷, NR⁷⁷S(O)_{j5a}R⁸⁷, -(C=S)OR⁷⁷, -(C=O)SR⁷⁷, -NR⁷⁷(C=NR⁸⁷)NR⁷⁸R⁸⁸, -NR⁷⁷(C=NR⁸⁷)OR⁷⁸, -NR⁷⁷(C=NR⁸⁷)SR⁷⁸, -O(C=O)OR⁷⁷, -O(C=O)NR⁷⁷R⁸⁷, -O(C=O)SR⁷⁷, -S(C=O)OR⁷⁷, -P(O)OR⁷⁷OR⁸⁷, or -S(C=O)NR⁷⁷R⁸⁷ substituents; or R⁵ with R⁶ taken together with the respective carbon atom to which they are attached, form a 3-10 membered saturated or unsaturated ring, wherein said ring is optionally substituted with R⁶⁹; or R⁵ with R⁶ taken together with the respective carbon atom to which they are attached, form a 3-10 membered saturated or unsaturated heterocyclic ring, wherein said ring is optionally substituted with R⁶⁹;

R⁷ and R⁸ are each independently H, acyl, alkyl, alkenyl, aryl, heteroaryl, heterocyclyl or cycloalkyl, any of which is optionally substituted by one or more G¹¹¹ substituents;


R⁴ is H, alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, cycloalkenyl, or heterocycloalkenyl, any of which is optionally substituted by one or more G⁴¹ substituents;

R⁶⁹ is equal to halo, -OR⁷⁸, -SH, -NR⁷⁸R⁸⁸, -CO₂R⁷⁸, -CONR⁷⁸R⁸⁸, -NO₂, -CN, -S(O)₈R⁷⁸, -SO₂NR⁷⁸R⁸⁸, C₀₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₁₋₁₀alkoxyC₁₋₁₀alkyl, C₁₋₁₀alkoxyC₂₋₁₀alkenyl, C₁₋₁₀alkoxyC₂₋₁₀alkynyl, C₁₋₁₀alkylthioC₁₋₁₀alkyl, C₁₋₁₀alkylthioC₂₋₁₀alkenyl, C₁₋₁₀alkylthioC₂₋₁₀alkynyl, cycloC₃₋₈alkyl, cycloC₃₋₈alkenyl, cycloC₃₋₈alkylC₁₋₁₀alkyl, cycloC₃₋₈alkenylC₁₋₁₀alkyl, cycloC₃₋₈alkylC₂₋₁₀alkenyl, cycloC₃₋₈alkenylC₂₋₁₀alkenyl, cycloC₃₋₈alkylC₂₋₁₀alkynyl, cycloC₃₋₈alkenylC₂₋₁₀alkynyl, heterocyclyl-C₀₋₁₀alkyl, heterocyclyl-C₂₋₁₀alkenyl, or heterocyclyl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -OR⁷⁷⁸, -SO₂NR⁷⁷⁸R⁸⁸⁸, or -NR⁷⁷⁸R⁸⁸⁸ substituents; or aryl-C₀₋₁₀alkyl, aryl-C₂₋₁₀alkenyl, or aryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -OR⁷⁷⁸, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CONR⁷⁷⁸R⁸⁸⁸, -SO₂NR⁷⁷⁸R⁸⁸⁸, or -NR⁷⁷⁸R⁸⁸⁸ substituents; or hetaryl-C₀₋₁₀alkyl, hetaryl-C₂₋₁₀alkenyl, or hetaryl-C₂₋₁₀alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -OR⁷⁷⁸, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CONR⁷⁷⁸R⁸⁸⁸, -SO₂NR⁷⁷⁸R⁸⁸⁸, or -NR⁷⁷⁸R⁸⁸⁸ substituents; or mono(C₁₋₆alkyl)aminoC₁₋₆alkyl, di(C₁₋₆alkyl)aminoC₁₋₆alkyl, mono(aryl)aminoC₁₋₆alkyl, di(aryl)aminoC₁₋₆alkyl, or -N(C₁₋₆alkyl)-C₁₋₆alkyl-aryl, any of which is optionally substituted with one or more independent halo, cyano, nitro, -OR⁷⁷⁸, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, haloC₁₋₁₀alkyl, haloC₂₋₁₀alkenyl, haloC₂₋₁₀alkynyl, -COOH, C₁₋₄alkoxycarbonyl, -CONR⁷⁷⁸R⁸⁸⁸, -SO₂NR⁷⁷⁸R⁸⁸⁸, or -NR⁷⁷⁸R⁸⁸⁸ substituents; or in the case of -NR⁷⁸R⁸⁸, R⁷⁸ and R⁸⁸ taken together with the nitrogen atom to which they are attached form a 3-10 membered saturated ring, unsaturated ring, heterocyclic saturated ring, or heterocyclic unsaturated ring, wherein said ring is optionally substituted with one or more independent halo, cyano, hydroxy, nitro, C₁₋₁₀alkoxy, -SO₂NR⁷⁷⁸R⁸⁸⁸, or -NR⁷⁷⁸R⁸⁸⁸ substituents;


R^{77} , R^{78} , R^{87} , R^{88} , R^{778} , and R^{888} are each independently C_{0-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy C_{1-10} alkyl, C_{1-10} alkoxy C_{2-10} alkenyl, C_{1-10} alkoxy C_{2-10} alkynyl, C_{1-10} alkylthio C_{1-10} alkyl, C_{1-10} alkylthio C_{2-10} alkenyl, C_{1-10} alkylthio C_{2-10} alkynyl, cyclo C_{3-8} alkyl, cyclo C_{3-8} alkenyl, cyclo C_{3-8} alkyl C_{1-10} alkyl, cyclo C_{3-8} alkenyl C_{1-10} alkyl, cyclo C_{3-8} alkyl C_{2-10} alkenyl, cyclo C_{3-8} alkenyl C_{2-10} alkenyl, cyclo C_{3-8} alkyl C_{2-10} alkynyl, cyclo C_{3-8} alkenyl C_{2-10} alkynyl, heterocyclyl- C_{0-10} alkyl, heterocyclyl- C_{2-10} alkenyl, heterocyclyl- C_{2-10} alkynyl, C_{1-10} alkylcarbonyl, C_{2-10} alkenylcarbonyl, C_{2-10} alkynylcarbonyl, C_{1-10} alkoxycarbonyl, C_{1-10} alkoxycarbonyl C_{1-10} alkyl, mono C_{1-6} alkylaminocarbonyl, di C_{1-6} alkylaminocarbonyl, mono(aryl)aminocarbonyl, di(aryl)aminocarbonyl, or C_{1-10} alkyl(aryl)aminocarbonyl, any of which is optionally substituted with one or more independent halo, cyano, hydroxy, nitro, C_{1-10} alkoxy, $-SO_2N(C_{0-4}alkyl)(C_{0-4}alkyl)$, or $-N(C_{0-4}alkyl)(C_{0-4}alkyl)$ substituents; or aryl- C_{0-10} alkyl, aryl- C_{2-10} alkenyl, or aryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-O(C_{0-4}alkyl)$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, halo C_{1-10} alkyl, halo C_{2-10} alkenyl, halo C_{2-10} alkynyl, $-COOH$, C_{1-4} alkoxycarbonyl, $-CON(C_{0-4}alkyl)(C_{0-10}alkyl)$, $-SO_2N(C_{0-4}alkyl)(C_{0-4}alkyl)$, or $-N(C_{0-4}alkyl)(C_{0-4}alkyl)$ substituents; or hetaryl- C_{0-10} alkyl, hetaryl- C_{2-10} alkenyl, or hetaryl- C_{2-10} alkynyl, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-O(C_{0-4}alkyl)$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, halo C_{1-10} alkyl, halo C_{2-10} alkenyl, halo C_{2-10} alkynyl, $-COOH$, C_{1-4} alkoxycarbonyl, $-CON(C_{0-4}alkyl)(C_{0-4}alkyl)$, $-SO_2N(C_{0-4}alkyl)(C_{0-4}alkyl)$, or $-N(C_{0-4}alkyl)(C_{0-4}alkyl)$ substituents; or mono(C_{1-6} alkyl)amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino C_{1-6} alkyl, mono(aryl)amino C_{1-6} alkyl, di(aryl)amino C_{1-6} alkyl, or $-N(C_{1-6}alkyl)-C_{1-6}alkyl-aryl$, any of which is optionally substituted with one or more independent halo, cyano, nitro, $-O(C_{0-4}alkyl)$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, halo C_{1-10} alkyl, halo C_{2-10} alkenyl, halo C_{2-10} alkynyl, $-COOH$, C_{1-4} alkoxycarbonyl, $-CON(C_{0-4}alkyl)(C_{0-4}alkyl)$, $-SO_2N(C_{0-4}alkyl)(C_{0-4}alkyl)$, or $-N(C_{0-4}alkyl)(C_{0-4}alkyl)$ substituents; and

n , m , $j1$, $j1a$, $j2a$, $j3a$, $j4$, $j4a$, $j5a$, $j6a$, $j7$, and $j8$ are each independently equal to 0, 1, or 2.

23. The method of any one of Claims 1 to 10, wherein the BTK inhibitor is:



or a pharmaceutically-acceptable salt thereof, and the PI3K inhibitor is:




or a pharmaceutically-acceptable salt thereof.

24. The method of paragraphs 1, 2, or 3, wherein the BTK inhibitor is (*S*)-4-(8-amino-3-(1-(but-2-ynoyl)pyrrolidin-2-yl)imidazo[1,5-a]pyrazin-1-yl)-*N*-(pyridin-2-yl)benzamide or a pharmaceutically-acceptable salt thereof, and the PI3K inhibitor or PI3K- δ inhibitor is (*S*)-*N*-(1-(7-fluoro-2-(pyridin-2-yl)quinolin-3-yl)ethyl)-9*H*-purin-6-amine or a pharmaceutically-acceptable salt thereof.

25. The method of any one of claims 1 to 23, which is a method of treating a cancer in a


human comprising administering a therapeutically effective dose of a BTK inhibitor and a PI3K inhibitor in combination, wherein the dose is effective to inhibit signaling between the cells of the cancer and at least one microenvironment selected from the group consisting of macrophages, monocytes, mast cells, helper T cells, cytotoxic T cells, regulatory T cells, natural killer cells, myeloid-derived suppressor cells, regulatory B cells, neutrophils, dendritic cells, and fibroblasts.

26. The method of Claim 24, wherein the cancer is a solid tumor cancer.
27. The method of Claim 25, wherein the solid tumor cancer is selected from the group consisting of colon carcinoma, pancreatic carcinoma, breast cancer, lung cancer, colorectal cancer, thyroid cancer, bone sarcoma, and stomach cancer.
28. The method of any one of Claims 24-26, wherein the dose is further effective to increase immune system recognition and rejection of the cancer by the human.
29. A PI3K inhibitor and a BTK inhibitor in combination, for use in treating a hyperproliferative disorder.
30. A pharmaceutical composition comprising a PI3K inhibitor and a BTK inhibitor in combination, and at least one pharmaceutically acceptable excipient.
31. The pharmaceutical composition of claim 29, for use in treating a hyperproliferative disorder.
32. A kit comprising a pharmaceutical composition comprising a PI3K inhibitor and a pharmaceutical composition comprising a BTK inhibitor, for co-administration of the PI3K inhibitor and the BTK inhibitor, either simultaneously or separately.
33. A method of treating a cancer in a human sensitive to platelet-mediated thrombosis comprising the step of administering a therapeutically effective dose of a PI3K inhibitor and a BTK inhibitor, wherein the BTK inhibitor is:



or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof.

34. The method of Claim 32, wherein the PI3K inhibitor is:




or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof.

35. The method of any of Claims 32-33, further comprising the step of administering a therapeutically effective dose of an anticoagulant or antiplatelet agent.

36. The method of Claim 34, wherein the anticoagulant or antiplatelet agent is selected from the group consisting of clopidogrel, prasugrel, ticagrelor, ticlopidine, warfarin,

acenocoumarol, dicumarol, phenprocoumon, heparin, low molecular weight heparin, fondaparinux, and idraparinux.

37. The method of any of Claims 32-35, wherein the cancer is selected from the group consisting of bladder cancer, squamous cell carcinoma including head and neck cancer, pancreatic ductal adenocarcinoma (PDA), pancreatic cancer, colon carcinoma, mammary carcinoma, breast cancer, fibrosarcoma, mesothelioma, renal cell carcinoma, lung carcinoma, thymoma, prostate cancer, colorectal cancer, ovarian cancer, acute myeloid leukemia, thymus cancer, brain cancer, squamous cell cancer, skin cancer, eye cancer, retinoblastoma, melanoma, intraocular melanoma, oral cavity and oropharyngeal cancers, gastric cancer, stomach cancer, cervical cancer, head, neck, renal cancer, kidney cancer, liver cancer, ovarian cancer, prostate cancer, colorectal cancer, esophageal cancer, testicular cancer, gynecological cancer, thyroid cancer, acquired immune deficiency syndrome (AIDS)-related cancer, viral-induced cancer, glioblastoma, esophageal tumors, hematological neoplasms, non-small-cell lung cancer, chronic myelocytic leukemia, diffuse large B-cell lymphoma, esophagus tumor, follicle center lymphoma, head and neck tumor, hepatitis C virus infection, hepatocellular carcinoma, Hodgkin's disease, metastatic colon cancer, multiple myeloma, non-Hodgkin's lymphoma, indolent non-Hodgkin's lymphoma, ovary tumor, pancreas tumor, renal cell carcinoma, small-cell lung cancer, stage IV melanoma, chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia (ALL), mature B-cell ALL, follicular lymphoma, mantle cell lymphoma, and Burkitt's lymphoma.
38. A method of treating a cancer in a human with a history of thrombosis, comprising the step of administering a therapeutically effective dose of a BTK inhibitor, wherein the BTK inhibitor is:



or a pharmaceutically-acceptable salt, cocrystal, hydrate, solvate, or prodrug thereof.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17

18/60




FIG. 18




FIG. 19

20/60




FIG. 20




FIG. 21

22/60




FIG. 22




FIG. 23




FIG. 24

25/60




FIG. 25

26/60




FIG. 26

27/60




FIG. 27




FIG. 28




FIG. 29

30/60




FIG. 30




FIG. 31

32/60




FIG. 32




FIG. 33

34/60




FIG. 34




FIG. 35

36/60




FIG. 36

37/60




FIG. 37

38/60




FIG. 38

39/60




FIG. 39

40/60




FIG. 40

41/60




FIG. 41

42/60




FIG. 42

43/60




FIG. 43

44/60




FIG. 44

45/60




FIG. 45

46/60




FIG. 46

47/60




FIG. 47




FIG. 48




FIG. 49

50/60




FIG. 50

51/60




FIG. 51

52/60




FIG. 52

53/60




FIG. 53

54/60




FIG. 54




FIG. 55




FIG. 56




FIG. 57




FIG. 58




FIG. 59

60/60




FIG. 60

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2014/003121

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K31/517 A61K31/519 A61K31/52 A61K45/06 A61P35/00 A61P35/02 A61P35/04 A61P43/00 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, CHEM ABS Data, MEDLINE, EMBASE, BIOSIS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/153514 A2 (PHARMACYCLICS INC [US]; BUGGY JOSEPH J [US]; ELIAS LAURENCE [US]; FYFE) 8 December 2011 (2011-12-08)	1-38
Y	abstract paragraph [0004] - paragraph [0006] paragraph [0309] paragraph [0322] - paragraph [0323] claims 46-48, 82-84, 117 -----	1-38
X	JIANGCHUAN TAO: "Combined treatment of BTK and PI3K inhibitors synergistically disrupts BCR-signaling, overcomes microenviroment-mediated survival and drug resistance in mantle cell lymphoma", AACR ANNUAL MEETING 2013, 10 April 2013 (2013-04-10), pages 1-2, XP055129363, the whole document ----- -/-	1-38
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 21 April 2015	Date of mailing of the international search report 28/04/2015	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Taylor, Mark	

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2014/003121

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2008/118468 A1 (AMGEN INC [US]) 2 October 2008 (2008-10-02) cited in the application abstract page 1, line 4 - line 6 page 9, line 25 - page 10, line 10 page 31, lines 1-2 and 10-16 example 45 claims 1-10 -----	1-38
Y	WO 2013/010868 A1 (MSD OSS BV [NL]; BARF TJEERD A [NL]; JANS CHRISTIAAN GERARDUS JOHANNES) 24 January 2013 (2013-01-24) cited in the application abstract page 1, line 6 - line 9 page 2, line 35 - page 4, line 28 page 21, line 5 - line 24 page 22, line 15 - line 20 example 6 page 105, line 12 - line 13 claims 13-17 -----	1-38
X,P	WO 2014/071109 A1 (INFINITY PHARMACEUTICALS INC [US]) 8 May 2014 (2014-05-08) abstract paragraph [0540] - paragraph [0543] paragraph [0570] example 20 claims 1-5, 7, 56, 57 -----	1-38

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2014/003121

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011153514	A2	08-12-2011	
		AU 2011261185 A1	10-01-2013
		CA 2800913 A1	08-12-2011
		CL 2012003381 A1	22-03-2013
		CN 103153311 A	12-06-2013
		EA 201270793 A2	30-08-2013
		EP 2575818 A2	10-04-2013
		JP 2013527249 A	27-06-2013
		KR 20130086957 A	05-08-2013
		SG 186077 A1	30-01-2013
		US 2012087915 A1	12-04-2012
		US 2012100138 A1	26-04-2012
		US 2012183535 A1	19-07-2012
		US 2013195852 A1	01-08-2013
		US 2013202611 A1	08-08-2013
		US 2013273030 A1	17-10-2013
		US 2013310402 A1	21-11-2013
		US 2014194446 A1	10-07-2014
		US 2015031710 A1	29-01-2015
		US 2015031711 A1	29-01-2015
		WO 2011153514 A2	08-12-2011
WO 2008118468	A1	02-10-2008	
		AU 2008231304 A1	02-10-2008
		BR PI0809141 A2	26-08-2014
		CA 2681136 A1	02-10-2008
		CN 101715453 A	26-05-2010
		CR 11053 A	23-10-2009
		CR 11582 A	10-01-2011
		EA 200901276 A1	30-04-2010
		EP 2137186 A1	30-12-2009
		IL 200867 A	30-11-2014
		JP 5527761 B2	25-06-2014
		JP 2010522179 A	01-07-2010
		JP 2014058526 A	03-04-2014
		KR 20100015795 A	12-02-2010
		NZ 579834 A	30-03-2012
		UA 98955 C2	10-07-2012
		US 2009137581 A1	28-05-2009
		US 2012220585 A1	30-08-2012
		US 2012220586 A1	30-08-2012
		US 2015045361 A1	12-02-2015
		WO 2008118468 A1	02-10-2008
WO 2013010868	A1	24-01-2013	
		AU 2012285987 A1	06-02-2014
		CA 2841886 A1	24-01-2013
		CL 2014000130 A1	22-08-2014
		CN 103889987 A	25-06-2014
		CO 6940411 A2	09-05-2014
		CR 20140030 A	03-06-2014
		DO P2014000008 A	30-04-2014
		EA 201490300 A1	30-05-2014
		EC SP14013217 A	31-03-2015
		EP 2734522 A1	28-05-2014
		JP 2014520870 A	25-08-2014
		KR 20140036324 A	25-03-2014
		MA 35348 B1	01-08-2014
		PE 16812014 A1	14-11-2014
		US 2014155385 A1	05-06-2014
		WO 2013010868 A1	24-01-2013

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2014/003121

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014071109	A1	08-05-2014	NONE