



(86) Date de dépôt PCT/PCT Filing Date: 2009/09/25
(87) Date publication PCT/PCT Publication Date: 2010/04/01
(45) Date de délivrance/Issue Date: 2017/08/29
(85) Entrée phase nationale/National Entry: 2011/03/14
(86) N° demande PCT/PCT Application No.: GB 2009/002310
(87) N° publication PCT/PCT Publication No.: 2010/035012
(30) Priorités/Priorities: 2008/09/26 (GB0817704.0);
2009/03/26 (GB0905314.1)

(51) Cl.Int./Int.Cl. *C07K 16/46* (2006.01),
C07K 16/18 (2006.01), *C07K 16/24* (2006.01),
C07K 16/28 (2006.01)

(72) Inventeurs/Inventors:
ADAMS, RALPH, GB;
HANCOCK, LAURA, GB;
HEYWOOD, SAM PHILIP, GB

(73) Propriétaire/Owner:
UCB PHARMA S.A., BE

(74) Agent: ROBIC

(54) Titre : PROTEINES DE FUSION D'ANTICORPS MULTIVALENTES
(54) Title: MULTIVALENT ANTIBODY FUSION PROTEINS

(57) **Abrégé/Abstract:**

A multivalent antibody fusion protein which comprises an immunoglobulin moiety, for example a Fab or Fab' fragment, with a first specificity for an antigen of interest, and further comprises two single domain antibodies (dAb) with specificity for a second antigen of interest, wherein the two single domain antibodies are linked by a disulfide bond. There is also provided particular dual specificity antibody fusion proteins comprising a Fab or Fab' fragment and one or more single domain antibodies which may be stabilised by a disulfide bond therebetween.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
1 April 2010 (01.04.2010)

PCT

(10) International Publication Number
WO 2010/035012 A9

(51) International Patent Classification:

C07K 16/46 (2006.01) C07K 16/28 (2006.01)
C07K 16/18 (2006.01) C07K 16/24 (2006.01)

(21) International Application Number:

PCT/GB2009/002310

(22) International Filing Date:

25 September 2009 (25.09.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0817704.0 26 September 2008 (26.09.2008) GB
0905314.1 26 March 2009 (26.03.2009) GB

(71) Applicant (for all designated States except US): **UCB PHARMA S.A.** [BE/BE]; 60 Allée de la recherche, B-1070 Brussels (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ADAMS, Ralph** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). **HANCOCK, Laura** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). **HEYWOOD, Sam, Philip** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).

(74) Agent: **THOMPSON, John**; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(48) Date of publication of this corrected version:

17 March 2011

(15) Information about Correction:

see Notice of 17 March 2011

(54) Title: BIOLOGICAL PRODUCTS

(57) Abstract: A multivalent antibody fusion protein which comprises an immunoglobulin moiety, for example a Fab or Fab' fragment, with a first specificity for an antigen of interest, and further comprises two single domain antibodies (dAb) with specificity for a second antigen of interest, wherein the two single domain antibodies are linked by a disulfide bond. There is also provided particular dual specificity antibody fusion proteins comprising a Fab or Fab' fragment and one or more single domain antibodies which may be stabilised by a disulfide bond therebetween.



WO 2010/035012 A9

MULTIVALENT ANTIBODY FUSION PROTEINS

The present invention relates to new dual specificity antibody fusion proteins. Such antibodies comprise a first specificity to an antigen of interest, and a second specificity for a second antigen of interest, for example a serum carrier protein for use in extending their *in vivo* serum half-life. Methods for the production of such molecules and pharmaceutical compositions comprising them are also provided.

The high specificity and affinity of antibodies makes them ideal diagnostic and therapeutic agents, particularly for modulating protein:protein interactions. Advances in the field of recombinant antibody technology have resulted in the production of antibody fragments, such as Fv, Fab, Fab' and F(ab')₂ fragments and other antibody fragments. These smaller molecules retain the antigen binding activity of whole antibodies and can also exhibit improved tissue penetration and pharmacokinetic properties in comparison to whole immunoglobulin molecules. Indeed, antibody fragments are proving to be versatile therapeutic agents, as seen by the recent success of products such as ReoPro® and Lucentis®. Whilst such fragments appear to exhibit a number of advantages over whole immunoglobulins, they also suffer from an increased rate of clearance from serum since they lack the Fc domain that imparts a long lifetime *in vivo* (Medasan *et al.*, 1997, J. Immunol. 158:2211-2217).

Antibodies with dual specificity, *i.e.* which bind to two different antigens have been previously described (for reviews, see Segal *et al.*, 1999, Curr. Opin. Immunol. 11:558-562; Plückthun & Pack, 1997, Immunotechnology, 3:83-105; Fischer and Leger, 2007, Pathobiology, 74, 3-14). Dual specificity antibodies are also described in WO02/02773, US2007065440, US2006257406, US2006106203 and US2006280734. Previous approaches to making hetero-bispecific antibody-based molecules have generally employed chemical cross-linking or protein engineering techniques. Chemical cross-linking suffers from poor yields of hetero- and homo-dimer formation and the requirement for their subsequent chromatographic separation. Protein engineering approaches have either been highly elaborate (*e.g.* knobs-into-holes engineering; Ridgway *et al.*, 1996, Protein Eng. 9(7):617-621) or have used molecules with inappropriate stability characteristics (*e.g.* diabodies, scFv). In some cases bispecific antibodies can also suffer from steric hindrance problems such that both antigens cannot bind simultaneously to each antibody arm.

Single variable domain antibodies also known as single domain antibodies or dAbs, correspond to the variable regions of either the heavy (VH) or light (VL) chain of an antibody. Murine single-domain antibodies were described by Ward *et al.*, 1989, Nature, 341, 544-546. Human and 'camelised' human single domain antibodies have also been described (Holt *et al.*, 2003, *Trends in Biotechnology*, 21, 484-490). Single domain antibodies have also been obtained from the camelids (camels and llamas) and cartilaginous fish (wobbegong and nurse sharks). These organisms have evolved high affinity single V-like domains (called VhH in camelids and V-NAR in

sharks), mounted on an Fc-equivalent constant domain framework as an integral and crucial component of their immune system (see Holliger & Hudson, for a review; 2005, Nature Biotechnology, 23(9):1126-1136).

Single domain antibody-enzyme fusions have been described in EP0368684.

5 Single domain-effector group fusions have also been described in WO2004/058820 which comprise a single variable domain. Dual variable domain immunoglobulins have been described in WO2007/024715. Dual specific ligands comprising two single domain antibodies with differing specificities have been described in EP1517921.

10 Means to improve the half-life of antibody fragments, such as Fv, Fab, Fab', F(ab')₂ and other antibody fragments, are known. One approach has been to conjugate the fragment to polymer molecules. Thus, the short circulating half-life of Fab', F(ab')₂ fragments in animals has been improved by conjugation to polyethylene glycol (PEG; see, for example, WO98/25791, WO99/64460 and WO98/37200).

15 Another approach has been to modify the antibody fragment by conjugation to an agent that interacts with the FcRn receptor (see, for example, WO97/34631). Yet another approach to extend half-life has been to use polypeptides that bind serum albumin (see, for example, Smith *et al.*, 2001, Bioconjugate Chem. 12:750-756; EP0486525; US6267964; WO04/001064; WO02/076489; and WO01/45746).

20 However, there still remains a need to produce antigen-binding immunoglobulin proteins that have a long *in vivo* half-life, as an alternative to those that have a long half life because they interact with the FcRn receptor, without being chemically modified by conjugation to PEG, or being conjugated to human serum albumin.

A variety of proteins exist in plasma and include thyroxine-binding protein, transthyretin, α 1-acid glycoprotein, transferrin, fibrinogen and albumin, or a fragment
25 of any thereof. Serum carrier proteins circulate within the body with half-lives measured in days, for example, 5 days for thyroxine-binding protein or 2 days for transthyretin (Bartalena & Robbins, 1993, Clinics in Lab. Med. 13:583-598), or 65 hours in the second phase of turnover of iodinated α 1-acid glycoprotein (Bree *et al.*,
30 1986, Clin. Pharmacokin. 11:336-342). Data from Gitlin *et al.* (1964, J. Clin. Invest. 10:1938-1951) suggest that in pregnant women, the half-life of α 1-acid glycoprotein is 3.8 days, 12 days for transferrin and 2.5 days for fibrinogen. Serum albumin is an abundant protein in both vascular and extravascular compartments with a half-life in
35 man of about 19 days (Peters, 1985, Adv Protein Chem. 37:161-245). This is similar to the half-life of IgG1, which is about 21 days (Waldeman & Strober, 1969, Progr. Allergy, 13:1-110).

The present invention provides improved dual specificity antibody fusion proteins which can be produced recombinantly and are capable of binding two antigens simultaneously, in particular two distinct/different antigens.

40 Thus, the present invention provides dual specificity antibody fusion proteins which comprise an immunoglobulin moiety, for example a Fab or Fab' fragment, with

a first specificity for an antigen of interest, and further comprise a single domain antibody (dAb) with specificity for a second antigen of interest, in particular where the first antigen and second antigen are different entities.

5 Multivalent as employed herein is intended to refer to an entity that has two or more binding sites, for example two or three binding sites such as two binding sites. Each binding site may bind the same epitope or different epitopes on the same antigen, or may bind different (distinct) antigens.

The present invention also provides dual specificity antibody fusion proteins which comprise an immunoglobulin moiety, for example a Fab or Fab' fragment, with
10 a first specificity for an antigen of interest, and further comprise at least one single domain antibody with specificity for a second antigen of interest.

A dual specificity antibody fusion of the invention will be capable of selectively binding to two antigens of interest.

In one embodiment the first and second antigen are the same antigen.

15 In one embodiment, an antigen of interest bound by the Fab or Fab' fragment may be a cell-associated protein, for example a cell surface protein on cells such as bacterial cells, yeast cells, T-cells, endothelial cells or tumour cells, or it may be a soluble protein. Antigens of interest may also be any medically relevant protein such as those proteins upregulated during disease or infection, for example receptors and/or
20 their corresponding ligands. Particular examples of cell surface proteins include adhesion molecules, for example integrins such as β 1 integrins e.g. VLA-4, E-selectin, P selectin or L-selectin, CD2, CD3, CD4, CD5, CD7, CD8, CD11a, CD11b, CD18, CD19, CD20, CD23, CD25, CD33, CD38, CD40, CD45, CDW52, CD69, CD134 (OX40), ICOS, BCMP7, CD137, CD27L, CDCP1, DPCR1, DPCR1,
25 dudulin2, FLJ20584, FLJ40787, HEK2, KIAA0634, KIAA0659, KIAA1246, KIAA1455, LTBP2, LTK, MAL2, MRP2, nectin-like2, NKCC1, PTK7, RAIG1, TCAM1, SC6, BCMP101, BCMP84, BCMP11, DTD, carcinoembryonic antigen (CEA), human milk fat globulin (HMFG1 and 2), MHC Class I and MHC Class II antigens, and VEGF, and where appropriate, receptors thereof.

30 Soluble antigens include interleukins such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, IL-12, IL-16 or IL-17, viral antigens for example respiratory syncytial virus or cytomegalovirus antigens, immunoglobulins, such as IgE, interferons such as interferon α , interferon β or interferon γ , tumour necrosis factor- α , tumor necrosis factor- β , colony stimulating factors such as G-CSF or GM-CSF, and platelet derived
35 growth factors such as PDGF- α , and PDGF- β and where appropriate receptors thereof. Other antigens include bacterial cell surface antigens, bacterial toxins, viruses such as influenza, EBV, HepA, B and C, bioterrorism agents, radionuclides and heavy metals, and snake and spider venoms and toxins.

40 In one embodiment, the antibody fusion protein of the invention may be used to functionally alter the activity of the antigen of interest. For example, the antibody

fusion protein may neutralize, antagonize or agonise the activity of said antigen, directly or indirectly.

In one embodiment, a second antigen of interest bound by the single domain antibody or antibodies in the dual specificity antibody fusion proteins of the invention may be a cell-associated protein, for example a cell surface protein on cells such as bacterial cells, yeast cells, T-cells, endothelial cells or tumour cells, or it may be a soluble protein. Antigens of interest may also be any medically relevant protein such as those proteins upregulated during disease or infection, for example receptors and/or their corresponding ligands. Particular examples of cell surface proteins include adhesion molecules, for example integrins such as $\beta 1$ integrins e.g. VLA-4, E-selectin, P selectin or L-selectin, CD2, CD3, CD4, CD5, CD7, CD8, CD11a, CD11b, CD18, CD19, CD20, CD23, CD25, CD33, CD38, CD40, CD45, CDW52, CD69, CD134 (OX40), ICOS, BCMP7, CD137, CD27L, CDCP1, DPCR1, DPCR1, dudulin2, FLJ20584, FLJ40787, HEK2, KIAA0634, KIAA0659, KIAA1246, KIAA1455, LTBP2, LTK, MAL2, MRP2, nectin-like2, NKCC1, PTK7, RAIG1, TCAM1, SC6, BCMP101, BCMP84, BCMP11, DTD, carcinoembryonic antigen (CEA), human milk fat globulin (HMFG1 and 2), MHC Class I and MHC Class II antigens, and VEGF, and where appropriate, receptors thereof.

Soluble antigens include interleukins such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, IL-12, IL-16 or IL-17, viral antigens for example respiratory syncytial virus or cytomegalovirus antigens, immunoglobulins, such as IgE, interferons such as interferon α , interferon β or interferon γ , tumour necrosis factor- α , tumor necrosis factor- β , colony stimulating factors such as G-CSF or GM-CSF, and platelet derived growth factors such as PDGF- α , and PDGF- β and where appropriate receptors thereof. Other antigens include bacterial cell surface antigens, bacterial toxins, viruses such as influenza, EBV, HepA, B and C, bioterrorism agents, radionuclides and heavy metals, and snake and spider venoms and toxins.

Other antigens which may be bound by the single domain antibody or antibodies include serum carrier proteins, polypeptides which enable cell-mediated effector function recruitment and nuclide chelator proteins.

Thus, in one example the present invention provides dual specificity antibody fusion proteins which comprise an immunoglobulin moiety with a first specificity for an antigen of interest, and further comprise a single domain antibody with specificity for a second protein, the latter providing the ability to recruit effector functions, such as complement pathway activation and/or effector cell recruitment. Further, fusion proteins of the present invention may be used to chelate radionuclides by virtue of a single domain antibody which binds to a nuclide chelator protein. Such fusion proteins are of use in imaging or radionuclide targeting approaches to therapy.

Accordingly, in one example there is provided an isolated dual specificity antibody fusion protein comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one dAb which has

specificity for a recruitment polypeptide, said dAb providing the ability to recruit cell-mediated effector function(s), directly or indirectly, by binding to said recruitment polypeptide.

The recruitment of effector function may be direct in that effector function is associated with a cell, said cell bearing a recruitment molecule on its surface. Indirect recruitment may occur when binding of a dAb to a recruitment molecule causes release of, for example, a factor which in turn may directly or indirectly recruit effector function, or may be via activation of a signalling pathway. Examples include TNF α , IL2, IL6, IL8, IL17, IFN γ , histamine, C1q, opsonin and other members of the classical and alternative complement activation cascades, such as C2, C4, C3-convertase, and C5 to C9.

As used herein, 'a recruitment polypeptide' includes a Fc γ R such as Fc γ RI, Fc γ RII and Fc γ RIII, a complement pathway protein such as, but without limitation, C1q and C3, a CD marker protein (Cluster of Differentiation marker) such as, but without limitation, CD68, CD115, CD16, CD80, CD83, CD86, CD56, CD64, CD3, CD4, CD8, CD28, CD45, CD19, CD20 and CD22. Further recruitment polypeptides which are CD marker proteins include CD1, CD1d, CD2, CD5, CD8, CD9, CD10, CD11, CD11a, CD11b, CD11c, CD13, CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD26, CD27, CD28, CD29, CD30, CD31, CD32, CD33, CD34, CD35, CD36, CD37, CD38, CD40, CD43, CD44, CD45, CD46, CD49, CD49a, CD49b, CD49c, CD49d, CD52, CD53, CD54, CD55, CD56, CD58, CD59, CD61, CD62, D62E, CD62L, CD62P, CD63, CD64, CD66e, CD68, CD70, CD71, CD72, CD79, CD80, CD81, CD82, CD83, CD84, CD85, CD86, CD88, CD89, CD90, CD94, CD95, CD98, CD106, CD114, CD116, CD117, CD118, CD120, CD122, CD130, CD131, CD132, CD133, CD134, CD135, CD137, CD138, CD141, CD142, CD143, CD146, CD147, CD151, CD152, CD153, CD154, CD155, CD162, CD164, CD169, CD184, CD206, CD209, CD257, CD278, CD281, CD282, CD283 and CD304, or a fragment of any thereof which retains the ability to recruit cell-mediated effector function either directly or indirectly. A recruitment polypeptide also includes immunoglobulin molecules such as IgG1, IgG2, IgG3, IgG4, IgE and IgA which possess effector function.

In one embodiment, the second protein for which the dAb has specificity is a complement pathway protein, with C1q being particularly preferred.

In a preferred embodiment, the second protein for which the dAb has specificity is a CD marker protein, with CD68, CD80, CD86, CD64, CD3, CD4, CD8 CD45, CD16 and CD35 being particularly preferred.

Accordingly also provided is an isolated dual specificity antibody fusion protein comprising an antibody fragment with specificity for an antigen of interest, said fragment being fused to at least one dAb which has specificity for a CD molecule selected from the group consisting of CD68, CD80, CD86, CD64, CD3, CD4, CD8 CD45, CD16 and CD35.

In one embodiment the single domain antibody or antibodies provide an extended half-life to the immunoglobulin moiety with the first specificity.

Accordingly, in one embodiment there is provided a dual specificity antibody fusion protein comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody which has specificity for a serum carrier protein, a circulating immunoglobulin molecule, or CD35/CR1, said single domain antibody providing an extended half-life to the antibody fragment with specificity for said antigen of interest by binding to said serum carrier protein, circulating immunoglobulin molecule or CD35/CR1.

In one embodiment there is provided an isolated dual specificity antibody fusion protein comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody which has specificity for a serum carrier protein, a circulating immunoglobulin molecule, or CD35/CR1, said single domain antibody providing an extended half-life to the antibody fragment with specificity for said antigen of interest by binding to said serum carrier protein, circulating immunoglobulin molecule or CD35/CR1.

As used herein, 'serum carrier proteins' include thyroxine-binding protein, transthyretin, α 1-acid glycoprotein, transferrin, fibrinogen and albumin, or a fragment of any thereof.

As used herein, a 'circulating immunoglobulin molecule' includes IgG1, IgG2, IgG3, IgG4, sIgA, IgM and IgD, or a fragment of any thereof.

CD35/CR1 is a protein present on red blood cells which have a half life of 36 days (normal range of 28 to 47 days; Lanaro *et al.*, 1971, Cancer, 28(3):658-661).

In a preferred embodiment, the second protein for which the dAb has specificity is a serum carrier protein, with a human serum carrier protein being particularly preferred. In a most preferred embodiment, the serum carrier protein is human serum albumin.

Accordingly provided is a dual specificity antibody fusion protein comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody which has specificity for human serum albumin.

In one embodiment the present invention provides an isolated dual specificity antibody fusion protein comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody which has specificity for human serum albumin.

In one embodiment, the antibody fragment with specificity for an antigen of interest is a Fab fragment. In another embodiment, the antibody fragment with specificity for an antigen of interest is a Fab' fragment.

Thus, in one most preferred embodiment, the antibody fusion proteins of the invention are translation fusion proteins, *i.e.* genetic fusions, the sequence of each of which is encoded by an expression vector. Alternatively, the antibody fusion protein

components have been fused using chemical means, *i.e.* by chemical conjugation or chemical cross-linking. Such chemical means are known in the art.

In one example, the antibody fragments are Fab' fragments which possess a native or a modified hinge region. Where the antibody fragment for use in preparing a dual specificity antibody fusion protein of the invention is a Fab' fragment, said fragment is generally extended at the C-terminus of the heavy chain by one or more amino acids. Thus, an antibody fusion of the invention can comprise a Fab' fragment translation fused (or chemically fused) to a dAb, directly or via a linker. Further, examples of suitable antibody Fab' fragments include those described in WO2005003170 and WO2005003171.

In another example, the antibody fragments are Fab fragments. Thus, an antibody fusion of the invention can comprise a Fab fragment translation fused (or chemically fused) to a linker sequence which in turn is translation fused (or chemically fused) to a dAb. Preferably, the Fab fragment is a Fab fragment which terminates at the interchain cysteines, as described in WO2005/003169.

The antibody Fab or Fab' fragments of use in the present invention can be from any species but are preferably derived from a monoclonal antibody, a human antibody, or are humanised fragments. An antibody fragment for use in the present invention can be derived from any class (e.g. IgG, IgE, IgM, IgD or IgA) or subclass of immunoglobulin molecule and may be obtained from any species including for example mouse, rat, shark, rabbit, pig, hamster, camel, llama, goat or human.

In one embodiment, the antibody Fab or Fab' fragment is a monoclonal, fully human, humanized or chimeric antibody fragment. In one embodiment the antibody Fab or Fab' fragments are fully human or humanised.

Monoclonal antibodies may be prepared by any method known in the art such as the hybridoma technique (Kohler & Milstein, *Nature*, 1975, 256, 495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today*, 1983, 4, 72) and the EBV-hybridoma technique (Cole *et al.*, "Monoclonal Antibodies and Cancer Therapy", pp. 77-96, Alan R. Liss, Inc., 1985).

Antibodies for use in the invention may also be generated using single lymphocyte antibody methods by cloning and expressing immunoglobulin variable region cDNAs generated from single lymphocytes selected for the production of specific antibodies by, for example, the methods described by Babcook, J. *et al.*, *Proc. Natl. Acad. Sci. USA*, 1996, 93(15), 7843-7848, WO 92/02551, WO2004/051268 and WO2004/106377.

Humanized antibodies are antibody molecules from non-human species having one or more complementarity determining regions (CDRs) from the non-human species and a framework region from a human immunoglobulin molecule (see, for example, US 5,585,089).

The antibodies for use in the present invention can also be generated using various phage display methods known in the art and include those disclosed by

Brinkman *et al.*, *J. Immunol. Methods*, 1995, 182, 41-50; Ames *et al.*, *J. Immunol. Methods*, 1995, 184, 177-186; Kettleborough *et al.* *Eur. J. Immunol.*, 1994, 24, 952-958; Persic *et al.*, *Gene*, 1997 187, 9-18; and Burton *et al.*, *Advances in Immunology*, 1994, 57, 191-280; WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; and WO 95/20401; and US 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743; and 5,969,108. Also, transgenic mice, or other organisms, including other mammals, may be used to generate humanized antibodies.

Fully human antibodies are those antibodies in which the variable regions and the constant regions (where present) of both the heavy and the light chains are all of human origin, or substantially identical to sequences of human origin, not necessarily from the same antibody. Examples of fully human antibodies may include antibodies produced for example by the phage display methods described above and antibodies produced by mice in which the murine immunoglobulin variable and/or constant region genes have been replaced by their human counterparts eg. as described in general terms in EP0546073 B1, US 5,545,806, US 5,569,825, US 5,625,126, US 5,633,425, US 5,661,016, US5,770,429, EP 0438474 B1 and EP0463151 B1.

The antibody Fab or Fab' fragment starting material for use in the present invention may be obtained from any whole antibody, especially a whole monoclonal antibody, using any suitable enzymatic cleavage and/or digestion techniques, for example by treatment with pepsin. Alternatively, or in addition the antibody starting material may be prepared by the use of recombinant DNA techniques involving the manipulation and re-expression of DNA encoding antibody variable and/or constant regions. Standard molecular biology techniques may be used to modify, add or delete amino acids or domains as desired. Any alterations to the variable or constant regions are still encompassed by the terms 'variable' and 'constant' regions as used herein.

The antibody fragment starting material may be obtained from any species including for example mouse, rat, rabbit, hamster, camel, llama, goat or human. Parts of the antibody fragment may be obtained from more than one species, for example the antibody fragments may be chimeric. In one example, the constant regions are from one species and the variable regions from another. The antibody fragment starting material may also be modified. In another example, the variable region of the antibody fragment has been created using recombinant DNA engineering techniques. Such engineered versions include those created for example from natural antibody variable regions by insertions, deletions or changes in or to the amino acid sequences of the natural antibodies. Particular examples of this type include those engineered variable region domains containing at least one CDR and, optionally, one or more framework amino acids from one antibody and the remainder of the variable region domain from a second antibody. The methods for creating and manufacturing these antibody fragments are well known in the art (see for example, Boss *et al.*, US

4,816,397; Cabilly et al., US 6,331,415; Shrader et al., WO 92/02551; Ward et al., 1989, Nature, 341, 544; Orlandi et al., 1989, Proc.Natl.Acad.Sci. USA, 86, 3833; Riechmann et al., 1988, Nature, 322, 323; Bird et al, 1988, Science, 242, 423; Queen et al., US 5,585,089; Adair, WO91/09967; Mountain and Adair, 1992, Biotechnol. Genet. Eng. Rev, 10, 1-142; Verma et al., 1998, Journal of Immunological Methods, 216, 165-181).

In the present invention each single domain antibody fused to the Fab or Fab' fragment may linked directly or via a linker.

Linked directly are employed herein is intended to refer to the fact that the "last" amino acid of the Fab or Fab' is joined by a peptide bond to the "first" amino acid of the single domain antibody(or indeed vice versa)

Examples of suitable linker regions for linking a dAb to a Fab or Fab' include, but are not limited to, flexible linker sequences and rigid linker sequences. Flexible linker sequences include those disclosed in Huston *et al.*, 1988, PNAS 85:5879-5883; Wright & Deonarain, Mol. Immunol., 2007, 44(11):2860-2869; Alfthan *et al.*, Prot. Eng., 1995, 8(7):725-731; Luo *et al.*, J. Biochem., 1995, 118(4):825-831; Tang *et al.*, 1996, J. Biol. Chem. 271(26):15682-15686; and Turner *et al.*, 1997, JIMM 205, 42-54 (see Table 1 for representative examples).

Table 1. Flexible linker sequences

SEQ ID NO:	SEQUENCE
1	SGGGGSE
2	DKTHTS
3	(S)GGGGS
45	(S)GGGGSGGGGS
46	(S)GGGGSGGGGS
47	(S)GGGGSGGGGS
48	(S)GGGGSGGGGS
4	AAAGSG-GASAS
5	AAAGSG-XGGGS-GASAS
49	AAAGSG-XGGGSXGGGS -GASAS
50	AAAGSG- XGGGSXGGGSXGGGS -GASAS
51	AAAGSG- XGGGSXGGGSXGGGSXGGGS-GASAS
6	AAAGSG-XS-GASAS
7	PGGNRGTTTTRRPATTTGSSPGPTQSHY
8	ATTGSSPGPT
9	ATTGS
-	GS

10

10	EPSGPISTINSPPSKESHKSP
11	GTVAAPSVFIFPPSD
12	GGGGIAPSMVGGGGS
13	GGGGKVEGAGGGGGS
14	GGGGSMKSHDGGGGS
15	GGGGNLITIVGGGGS
16	GGGGVVPSPGPGGGS
17	GGEKSIPGGGGS
18	RPLSYRPPFPFGFPSVRP
19	YPRSIYIRRRHPSPLTT
20	TPSHLSHILPSFGLPTFN
21	RPVSPFTFPRLSNSWLPA
22	SPA AHFPRSIPRPGPIRT
23	APGPSAPSHRSLPSRAFG
24	PRNSIHFLHPLLVA PLGA
25	MPSLSGVLQVRYLSPPDL
26	SPQYPSPLTLTLPPHPSL
27	NPSLNPPSYLHRAPSRIS
28	LPWRTSLLPSLPLRRRP
29	PPLFAKGPVGLLSRSFPP
30	VPPAPVVSLRSAHARPPY
31	LRPTPPRVRSYTCCPTP-
32	PNVAHVLP LLTVPWDNLR
33	CNPLLPLCARSPAVRTEP

S) is optional in sequence3 and 45 to 48.

Examples of rigid linkers include the peptide sequences GAPAPAAPAPA (SEQ ID NO:34), PPPP (SEQ ID NO:35) and PPP.

- 5 In one embodiment the peptide linker is an albumin binding peptide. Examples of albumin binding peptides are provided in WO 2007/106120 and include:

Table 2

SEQ ID NO:	SEQUENCE
208	DLCLRDWGCLW
209	DICLPRWGCLW
210	MEDICLPRWGCLWGD
211	QRLMEDICLPRWGCLWEDDE
212	QGLIGDICLPRWGCLWGRSV

11

213	QGLIGDICLPRWGCLWGRSVK
214	EDICLPRWGCLWEDD
215	RLMEDICLPRWGCLWEDD
216	MEDICLPRWGCLWEDD
217	MEDICLPRWGCLWED
218	RLMEDICLARWGCLWEDD
219	EVRSFCTRWPAEKSCKPLRG
220	RAPESFVCYWETICFERSEQ
221	EMCYFPGICWM

In one embodiment, an antibody hinge sequence or part thereof is used as a linker, eg. the upper hinge sequence. Typically, antibody Fab' fragments for use in the present invention possess a native or a modified hinge region. Such hinge regions are used as a natural linker to the dAb moiety. The native hinge region is the hinge region normally associated with the C_H1 domain of the antibody molecule. A modified hinge region is any hinge that differs in length and/or composition from the native hinge region. Such hinges can include hinge regions from any other species, such as human, mouse, rat, rabbit, hamster, camel, llama or goat hinge regions. Other modified hinge regions may comprise a complete hinge region derived from an antibody of a different class or subclass from that of the C_H1 domain. Thus, for instance, a C_H1 domain of class γ 1 may be attached to a hinge region of class γ 4. Alternatively, the modified hinge region may comprise part of a natural hinge or a repeating unit in which each unit in the repeat is derived from a natural hinge region.

In a further alternative, the natural hinge region may be altered by converting one or more cysteine or other residues into neutral residues, such as alanine, or by converting suitably placed residues into cysteine residues. By such means the number of cysteine residues in the hinge region may be increased or decreased. In addition other characteristics of the hinge can be controlled, such as the distance of the hinge cysteine(s) from the light chain interchain cysteine, the distance between the cysteines of the hinge and the composition of other amino acids in the hinge that may affect properties of the hinge such as flexibility *e.g.* glycines may be incorporated into the hinge to increase rotational flexibility or prolines may be incorporated to reduce flexibility. Alternatively combinations of charged or hydrophobic residues may be incorporated into the hinge to confer multimerisation properties, see for example, Richter *et al.*, 2001, Prot. Eng. 14(10):775-783 for use of charged or ionic tails, *e.g.*, acidic tails as linkers and Kostelny *et al.*, 1992, J. Immunol. 5(1):1547-1553 for leucine zipper sequences. Other modified hinge regions may be entirely synthetic and

may be designed to possess desired properties such as length, composition and flexibility.

A number of modified hinge regions have already been described for example, in US5,677,425, US6642356, WO9915549, WO2005003170, WO2005003169, WO2005003170, WO9825971 and WO2005003171. Such hinges generally follow on from the CH1 region, but may also be incorporated onto the end of constant region of a light chain kappa or lambda fragment; see Table 3 for examples.

Table 3. Hinge linker sequences

SEQ ID NO:	SEQUENCE
36	DKTHTCAA
37	DKTHTCPPCPA
38	DKTHTCPPCPATCPPCPA
39	DKTHTCPPCPATCPPCPATCPPCPA
40	DKTHTCPPCPAGKPTLYNSLVMSDTAGTCY
41	DKTHTCPPCPAGKPTHVNVSVVMAEVDGTCY
42	DKTHTCCVECPCPA
43	DKTHTCPRCPEPKSCDTPPPCPRCPA
44	DKTHTCPSCPA

Single variable domains also known as single domain antibodies or dAbs for use in the present invention can be generated using methods known in the art and include those disclosed in WO2005118642, Ward *et al.*, 1989, Nature, 341, 544-546 and Holt *et al.*, 2003, *Trends in Biotechnology*, 21, 484-490. In one embodiment a single domain antibody for use in present invention is a heavy chain variable domain (VH) or a light chain domain (VL). Each light chain domain may be either of the kappa or lambda subgroup. Methods for isolating VH and VL domains have been described in the art, see for example EP0368684 and Ward *et al.*, *supra*. Such domains may be derived from any suitable species or antibody starting material. In one embodiment the single domain antibody may be derived from a rodent, a human or other species. In one embodiment the single domain antibody is humanised.

In one embodiment the single domain antibody is derived from a phage display library, using the methods described in for example, WO2005/118642, Jespers *et al.*, 2004, *Nature Biotechnology*, 22, 1161-1165 and Holt *et al.*, 2003, *Trends in Biotechnology*, 21, 484-490. Preferably such single domain antibodies are fully human but may also be derived from other species. In one embodiment the single variable domain is chimeric in that the framework is human or substantially human in origin and the CDR(s) is/are of non-human origin. It will be appreciated that the sequence of the single domain antibody once isolated may be modified to improve the

characteristics of the single domain antibody, for example solubility, as described in Holt *et al.*, *supra*.

Substantially human as employed herein is intended to refer that the human character of the original material is retained, which may be relevant to immunogenicity. Substantially human material would include wherein one amino acid in the framework sequence is added deleted or replaced by another amino acid.

In one embodiment the dAb is a human sequence obtained from scFv phage-display or from a transgenic Humouse™ or Velocimouse™ or a humanised rodent.

In one embodiment, the dAb is obtained from a human or humanised rodent, a camelid or a shark. Such a dAb will preferably be humanised. In one example the single domain antibody is a VHH domain based on camelid immunoglobulins as described in EP0656946. In one example, a camel or a llama is immunised with an antigen of interest and blood collected when the titre is appropriate. The gene encoding the dAb may be cloned by single cell PCR, or the B cell(s) encoding the dAb may be immortalised by EBV transformation, or by fusion to an immortal cell line.

As described herein above, the present invention provides dual specificity antibody fusion proteins comprising an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody, directly or via a linker, which has specificity for a second antigen of interest.

Accordingly, in one embodiment, the antibody fragment, eg. Fab or Fab' fragment is fused at the N-terminus of the heavy or the light chain variable region to a dAb directly or via a linker. Alternatively, the antibody Fab or Fab' fragment is fused at the C-terminus of the heavy or light chain to a dAb directly or via a linker. In another embodiment the heavy and light chains of the antibody Fab or Fab' fragment are each fused at the C-terminus to a dAb directly or via a linker. The linkage can be a chemical conjugation but is most preferably a translation fusion, *i.e.* a genetic fusion where the sequence of each is encoded in sequence by an expression vector.

Typically the N-terminus of the single domain antibody will be fused to the C-terminus of the heavy or light chain of the Fab or Fab' fragment, directly or via a linker, and where the single domain antibody is fused to the N-terminus of the Fab or Fab' it will be fused via its C-terminus, optionally via a linker.

In one embodiment the present invention provides a dual specificity antibody fusion protein comprising or consisting of an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to a single domain antibody at the N-terminus of the heavy or light chain which has specificity for a second antigen of interest.

In one embodiment the present invention provides a dual specificity antibody fusion protein comprising or consisting of an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to a single domain

antibody at the C-terminus of the heavy or light chain which has specificity for a second antigen of interest.

In one embodiment the present invention provides a dual specificity antibody fusion protein comprising or consisting of an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to at least one single domain antibody at the C-terminus of the heavy or light chain which has specificity for a second antigen of interest.

In one embodiment the present invention provides a dual specificity antibody fusion protein comprising or consisting of an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to two single domain antibodies wherein each single domain antibody is fused in linear sequence to each other, optionally via a linker and the resulting single domain antibody fusion is fused to the C-terminus of the light chain or the heavy chain of the Fab or Fab' fragment.

In one embodiment the present invention provides a dual specificity antibody fusion protein comprising or consisting of an antibody Fab or Fab' fragment with specificity for an antigen of interest, said fragment being fused to two single domain antibodies wherein one single domain antibody is fused to the C-terminus of the light chain of the Fab or Fab' fragment and the other single domain antibody is fused to the C-terminus of the heavy chain of the Fab or Fab' fragment, said single domain antibodies having specificity for a second antigen of interest.

In one embodiment where the heavy and light chains of the Fab or Fab' fragment each comprise a single domain antibody at the C-terminus the two single domain antibodies are identical i.e. have the same binding specificity for the same antigen. In one example, they bind the same epitope on the same antigen. For example the single domain antibodies may both be the same VH dAb, the same VHH dAb or the same VL dAb.

Preferably where the heavy and light chains of the Fab or Fab' fragment each comprise a single domain antibody at the C-terminus the two single domain antibodies are a complementary VH/VL pair which bind the antigen co-operatively i.e. they are a complementary VH/VL pair which have the same binding specificity. Typically they will be a VH/VL pair derived from the same antibody.

In one embodiment, where the dual specificity antibody fusion protein of the present invention comprises two single domain antibodies which are a complementary VH/VL pair, the VH single domain antibody is fused to the C-terminus of the heavy chain constant region (CH1) and the VL single domain antibody is fused to the C-terminus of the light chain constant region (C kappa or C lambda). In one embodiment, where the dual specificity antibody fusion protein of the present invention comprises two single domain antibodies which are a complementary VH/VL pair, the VL single domain antibody is fused to the C-terminus of the heavy chain constant region (CH1) and the VH single domain antibody is fused to the C-terminus of the light chain constant region (C kappa or C lambda).

In one embodiment, where the dual specificity antibody fusion protein of the present invention comprises two single domain antibodies which are linked by one or more disulfide bonds for example two single domain antibodies which are a complementary VH/VL pair linked by one or more (such as 1 or 2) disulfide bonds, such as the VH single domain antibody is fused to the C-terminus of the heavy chain constant region (CH1) and the VL single domain antibody is fused to the C-terminus of the light chain constant region (C kappa or C lambda). Alternatively the VL single domain antibody is fused to the C-terminus of the heavy chain constant region (CH1) and the VH single domain antibody is fused to the C-terminus of the light chain constant region (C kappa or C lambda).

The disulfide bond is thought to provide additional stabilisation to the construct, which may be advantageous.

In one or more embodiments the disulfide bond between the heavy and light chain such as between the CH domain and CL or CK domain is not present, for example because one or more cysteines which form the bond are replaced. Said one or more cysteines may be replaced with, for example serine.

In one or more embodiments an interchain disulfide bond between the heavy and light chain between the CH domain and CL or CK domain is present.

In one embodiment there is provided a F(ab)₂ fragment comprising one, two, three or four single domain antibodies, for example a two separate VH/VL pairs which may be directed to the same or different antigens.

In dual specificity fusion proteins of the present invention the single domain antibody or antibodies bind to a second antigen, different from that bound by the Fab or Fab' fragment component.

In one example the dAbs for use in the present invention exhibit specificity for a complement pathway protein, a CD marker protein or an FcγR. In this case the dAb is preferably specific for a CD molecule. Most preferably, the dAb exhibits specificity for a CD molecule selected from the group consisting of CD68, CD80, CD86, CD64, CD3, CD4, CD8 CD45, CD16 and CD35.

In a preferred example the dAbs for use in the present invention exhibit specificity for a serum carrier protein, a circulating immunoglobulin molecule, or CD35/CR1, the serum carrier protein preferably being a human serum carrier protein such as thyroxine-binding protein, transthyretin, α1-acid glycoprotein, transferrin, fibrinogen or serum albumin. Most preferably, the dAb exhibits specificity for human serum albumin. Thus, in one example, a rabbit, mouse, rat, camel or a llama is immunised with a serum carrier protein, a circulating immunoglobulin molecule, or CD35/CR1 (e.g. human serum albumin) and blood collected when the titre is appropriate. The gene encoding the dAb may be cloned by single cell PCR, or the B cell(s) encoding the dAb may be immortalised by EBV transformation, or by fusion to an immortal cell line. Alternatively the single domain antibody may be obtained by phage display as described herein above.

In one embodiment the single domain antibody or antibodies bind human serum albumin. In one embodiment the single domain antibody or antibodies bind human serum albumin, murine serum albumin and rat serum albumin.

5 In one embodiment the single domain antibody which binds serum albumin is a dAb provided in WO2005/118642 (see for example figures 1c and 1d) or a VHH provided in WO2004/041862 or a humanised nanobody described in, for example table III of WO2006/122787.

10 In one embodiment a single domain antibody which binds human serum albumin for use in the present invention is a heavy chain VH single domain antibody which comprises at least one of a CDR having the sequence given in Figure 5 (e) SEQ ID NO:56 or Figure 5 (k) SEQ ID NO:62 for CDR-H1, a CDR having the sequence given in Figure 5(f) SEQ ID NO:57 or Figure 5 (l) SEQ ID NO:63 for CDR-H2 and a CDR having the sequence given in Figure 5 (g) SEQ ID NO:58 or Figure 5 (m) SEQ ID NO:64 for CDR-H3.

15 In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a heavy chain VH antibody, wherein at least two of CDR-H1, CDR-H2 and CDR-H3 of the VH domain are selected from the following: the sequence given in SEQ ID NO:56 or SEQ ID NO:62 for CDR-H1, the sequence given in SEQ ID NO:57 or SEQ ID NO:63 for CDR-H2 and the sequence
20 given in SEQ ID NO:58 or SEQ ID NO:64 for CDR-H3. For example, the single domain antibody may comprise a VH domain wherein CDR-H1 has the sequence given in SEQ ID NO:56 and CDR-H2 has the sequence given in SEQ ID NO:57. Alternatively, the single domain antibody may comprise a VH domain wherein CDR-H1 has the sequence given in SEQ ID NO:56 and CDR-H3 has the sequence given in
25 SEQ ID NO:58. For the avoidance of doubt, it is understood that all permutations are included.

In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a heavy chain VH single domain antibody, wherein the VH domain comprises the sequence given in SEQ ID NO:56 for CDR-
30 H1, the sequence given in SEQ ID NO:57 for CDR-H2 and the sequence given in SEQ ID NO:58 for CDR-H3.

In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a heavy chain VH single domain antibody, wherein the VH domain comprises the sequence given in SEQ ID NO:62 for CDR-
35 H1, the sequence given in SEQ ID NO:63 for CDR-H2 and the sequence given in SEQ ID NO:64 for CDR-H3.

In one embodiment a single domain antibody which binds human serum albumin for use in the present invention is a humanised heavy chain VH single domain antibody, dAbH1, having the sequence given in Figure 5 (a) (SEQ ID NO:52).
40 An example of a suitable CH1-dAbH1 fusion comprising a G₄S linker is given in Figure 6 (SEQ ID NO:68).

In one embodiment the single domain antibody which binds human serum albumin for use in the present invention is a humanised heavy chain VH single domain antibody, dAbH2, having the sequence given in Figure 5 (c) (SEQ ID NO:54). An example of a suitable CH1-dAbH2 fusion comprising a G₄S linker is given in Figure 6 (SEQ ID NO:69).

The residues in antibody variable domains are conventionally numbered according to a system devised by Kabat *et al.* This system is set forth in Kabat *et al.*, 1987, in Sequences of Proteins of Immunological Interest, US Department of Health and Human Services, NIH, USA (hereafter “Kabat *et al. (supra)*”). This numbering system is used in the present specification except where otherwise indicated.

The Kabat residue designations do not always correspond directly with the linear numbering of the amino acid residues. The actual linear amino acid sequence may contain fewer or additional amino acids than in the strict Kabat numbering corresponding to a shortening of, or insertion into, a structural component, whether framework or complementarity determining region (CDR), of the basic variable domain structure. The correct Kabat numbering of residues may be determined for a given antibody by alignment of residues of homology in the sequence of the antibody with a “standard” Kabat numbered sequence.

The CDRs of the heavy chain variable domain are located at residues 31-35 (CDR-H1), residues 50-65 (CDR-H2) and residues 95-102 (CDR-H3) according to the Kabat numbering system. However, according to Chothia (Chothia, C. and Lesk, A.M. J. Mol. Biol., 196, 901-917 (1987)), the loop equivalent to CDR-H1 extends from residue 26 to residue 32. Thus ‘CDR-H1’, as used herein, comprises residues 26 to 35, as described by a combination of the Kabat numbering system and Chothia’s topological loop definition.

The CDRs of the light chain variable domain are located at residues 24-34 (CDR-L1), residues 50-56 (CDR-L2) and residues 89-97 (CDR-L3) according to the Kabat numbering system.

In one embodiment a single domain antibody which binds human serum albumin for use in the present invention is a light chain VL single domain antibody which comprises at least one of a CDR having the sequence given in Figure 5 (h) SEQ ID NO:59 or Figure 5 (n) SEQ ID NO:65 for CDR-L1, a CDR having the sequence given in Figure 5(i) SEQ ID NO:60 or Figure 5 (o) SEQ ID NO:66 for CDR-L2 and a CDR having the sequence given in Figure 5 (j) SEQ ID NO:61 or Figure 5 (p) SEQ ID NO:67 for CDR-L3.

In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a light chain VL antibody, wherein at least two of CDR-L1, CDR-L2 and CDR-L3 of the VL domain are selected from the following: the sequence given in SEQ ID NO:59 or SEQ ID NO:65 for CDR-L1, the sequence given in SEQ ID NO:60 or SEQ ID NO:66 for CDR-L2 and the sequence given in SEQ ID NO:61 or SEQ ID NO:67 for CDR-L3. For example, the domain

antibody may comprise a VL domain wherein CDR-L1 has the sequence given in SEQ ID NO:59 and CDR-L2 has the sequence given in SEQ ID NO:60.

Alternatively, the domain antibody may comprise a VL domain wherein CDR-L1 has the sequence given in SEQ ID NO:59 and CDR-L3 has the sequence given in SEQ ID NO:61. For the avoidance of doubt, it is understood that all permutations are included.

In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a light chain VL domain antibody, wherein the VL domain comprises the sequence given in SEQ ID NO:59 for CDR-L1, the sequence given in SEQ ID NO:60 for CDR-L2 and the sequence given in SEQ ID NO:61 for CDR-L3.

In another embodiment a single domain antibody which binds human serum albumin for use in the present invention is a light chain VL domain antibody, wherein the VL domain comprises the sequence given in SEQ ID NO:65 for CDR-L1, the sequence given in SEQ ID NO:66 for CDR-L2 and the sequence given in SEQ ID NO:67 for CDR-L3.

In one embodiment a single domain antibody which binds human serum albumin for use in the present invention is a humanised light chain VL single domain antibody, dAbL1, having the sequence given in Figure 5 (b) (SEQ ID NO:53). An example of a suitable CH1-dAbL1 fusion and a Ck1-dAbL1 fusion both comprising a G₄S linker is given in Figure 6 (SEQ ID NO:70 and SEQ ID NO:72).

In one embodiment a single domain antibody which binds human serum albumin for use in the present invention is a humanised light chain VL single domain antibody, dAbL2, having the sequence given in Figure 5 (d) (SEQ ID NO:55). An example of a suitable CH1-dAbL2 fusion and a Ck1-dAbL2 fusion both comprising a G₄S linker is given in Figure 6 (SEQ ID NO:71 and SEQ ID NO:73).

In one embodiment where the heavy and light chains of the Fab or Fab' fragment each comprise a single domain antibody at the C-terminus and the two single domain antibodies are a complementary VH/VL pair which bind the antigen co-operatively as described herein above, the VH dAb is dAbH1 (SEQ ID NO:52) and the VL dAb is dAbL1 (SEQ ID NO:53).

In one embodiment where the heavy and light chains of the Fab or Fab' fragment each comprise a single domain antibody at the C-terminus the two single domain antibodies are a complementary VH/VL pair which bind the antigen co-operatively as described herein above, the VH dAb is dAbH2 (SEQ ID NO:54) and the VL dAb is dAbL2 (SEQ ID NO:55).

In another aspect, the present invention provides albumin binding antibodies or fragments thereof containing one or more of the CDRs provided herein above and in Figure 5 (e-p), in particular comprising a CDRH1 with the sequence shown in SED ID NO. 56, a CDRH2 with the sequence shown in SED ID NO. 57, a CDRH3 with the sequence shown in SED ID NO. 58, a CDRL1 with the sequence shown in SED

ID NO. 59, a CDRL2 with the sequence shown in SED ID NO. 60, and/or a CDRL3 with the sequence shown in SED ID NO. 61. In one embodiment the albumin binding antibodies or fragments comprise a CDRH1 with the sequence shown in SED ID NO. 62, a CDRH2 with the sequence shown in SED ID NO. 63, a CDRH3 with the sequence shown in SED ID NO. 64, a CDRL1 with the sequence shown in SED ID NO. 65, a CDRL2 with the sequence shown in SED ID NO. 66, and/or a CDRL3 with the sequence shown in SED ID NO. 67. Said CDRs may be incorporated into any suitable antibody framework and into any suitable antibody format. Such antibodies include whole antibodies and functionally active fragments or derivatives thereof which may be, but are not limited to, monoclonal, humanised, fully human or chimeric antibodies. Accordingly, such albumin binding antibodies may comprise a complete antibody molecule having full length heavy and light chains or a fragment thereof and may be, but are not limited to Fab, modified Fab, Fab', F(ab')₂, Fv, single domain antibodies, scFv, bi, tri or tetra-valent antibodies, Bis-scFv, diabodies, triabodies, tetrabodies and epitope-binding fragments of any of the above (see for example Holliger and Hudson, 2005, Nature Biotech. 23(9):1126-1136; Adair and Lawson, 2005, Drug Design Reviews - Online 2(3), 209-217). The methods for creating and manufacturing these antibody fragments are well known in the art (see for example Verma et al., 1998, Journal of Immunological Methods, 216, 165-181). Multi-valent antibodies may comprise multiple specificities or may be monospecific (see for example WO 92/22853 and WO05/113605). It will be appreciated that this aspect of the invention also extends to variants of these albumin binding antibodies.

It will be appreciated that such albumin binding antibodies, in particular single domain antibodies may be conjugated to any other antibody or protein or other molecule, as desired or used in any other suitable context. In one example the single domain antibodies dAbH1, dAbL1, dAbH2, dAbL2 as described above and shown in Figure 5 (a-d) may be incorporated into any suitable antibody format or used as single domain antibodies in any suitable context, such as a fusion or conjugate.

In one embodiment antibodies of this aspect of the invention comprise the sequence given in Figure 5(e) for CDR-H1, the sequence given in Figure 5(f) for CDR-H2 and the sequence given in Figure 5(g) for CDR-H3.

In one embodiment antibodies of this aspect of the invention comprise the sequence given in Figure 5(k) for CDR-H1, the sequence given in Figure 5(l) for CDR-H2 and the sequence given in Figure 5(m) for CDR-H3.

In one embodiment antibodies of this aspect of the invention comprise the sequence given in Figure 5(h) for CDR-L1, the sequence given in Figure 5(i) for CDR-L2 and the sequence given in Figure 5(j) for CDR-L3.

In one embodiment antibodies of this aspect of the invention comprise the sequence given in Figure 5(n) for CDR-L1, the sequence given in Figure 5(o) for CDR-L2 and the sequence given in Figure 5(p) for CDR-L3.

In the antibody formats below each of the sequences from the sequence listing herein may be located in the position corresponding to the natural position or a non-natural position. Natural position will be for the relevant sequence in the listing labelled CDRH1 position H1, for the relevant sequence in the listing labelled CDRH2 position H2, for the relevant sequence in the listing labelled CDRH3 position H3, for the relevant sequence in the listing labelled CDRL1 position L1, for the relevant sequence in the listing labelled CDRL2 position L2, and for the relevant sequence in the listing labelled CDRL3 position L3. Combinations thereof are also envisaged such as H1 and H2, H1 and H3, H1 and L1, H1 and L2, H1 and L3, H2 and L1, H2 and L2, H2 and L3, H2 and H3, H3 and L1, H3 and L2, H3 and L3, H1 and H2 and H3, H1 and H2 and L1, H1 and H2 and L2, H1 and H2 and L3, H2 and H3 and L1, H2 and H3 and L2, H2 and H3 and L3, H3 and L1 and L2, H3 and L1 and L3, H3 and L2 and L3, L1 and L2 and L3, H1 and H2 and H3 and L1, H1 and H2 and H3 and L2, H1 and H2 and H3 and L3, H2 and H3 and L1 and L2, H2 and H3 and L1 and L3, and H2 and H3 and L2 and L3, H3 and L1 and L2 and L3, H1 and H2 and H3 and L1 and L2, H1 and H2 and H3 and L2 and L3, H1 and H2 and H3 and L1 and L3, L1 and L2 and L3 and H1 and H2, L1 and L2 and L3 and H1 and H3, L1 and L2 and L3 and H2 and H3, H1 and H2 and H3 and L1 and L2 and L3.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 88 to 93.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 94 to 99.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 100 to 105.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 106 to 111.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 112 to 117.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 118 to 123.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 124 to 129.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 130 to 135.

5 In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 136 to 141.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 142 to 147.

10 In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 148 to 153.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 154 to 159.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 160 to 165.

20 In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 166 to 171.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 172 to 177.

25 In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 178 to 183.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 184 to 189.

In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 190 to 195.

35 In one embodiment the antibody fusion protein of the disclosure comprises a sequence, for example 1, 2, 3, 4, 5 or 6 sequence(s) selected from Sequence ID NO: 196 to 201.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 202.

40 In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 203.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID Nos: 202 and 203.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 204.

5 In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 205.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID Nos: 204 and 205.

10 In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 206.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 207.

In one embodiment the antibody fusion protein of the disclosure comprises Sequence ID No: 206 and 207.

15 Where the single domain antibody or antibodies of the dual specificity fusion protein of the present invention bind to albumin the binding affinity of the single domain antibody for albumin will be sufficient to extend the half-life of the Fab or Fab' *in vivo*. It has been reported that an affinity for albumin of less than or equal to 2.5 μ M affinity will extend half-life in vivo (Nguyen, A. *et al* (2006) Protein
20 Engineering, Design & Selection, 19(7), 291-297). The single domain antibody molecules of the present invention preferably have a binding affinity suited to their purpose and the antigen to which they bind. In one example the single domain antibodies have a high binding affinity, for example picomolar. In one example the single domain antibodies have a binding affinity for antigen which is nanomolar or
25 micromolar. Affinity may be measured using any suitable method known in the art, including BIAcore as described in the Examples herein using natural or recombinant antigen.

Preferably the single domain antibody molecules of the present invention which bind albumin have a binding affinity of about 2 μ M or better. In one
30 embodiment the single domain antibody molecule of the present invention has a binding affinity of about 1 μ M or better. In one embodiment the single domain antibody molecule of the present invention has a binding affinity of about 500nM or better. In one embodiment the single domain antibody molecule of the present invention has a binding affinity of about 200nM or better. In one embodiment the
35 domain antibody molecule of the present invention has a binding affinity of about 1nM or better. It will be appreciated that the affinity of single domain antibodies provided by the present invention and known in the art may be altered using any suitable method known in the art. The present invention therefore also relates to variants of the domain antibody molecules of the present invention, which have an
40 improved affinity for albumin. Such variants can be obtained by a number of affinity maturation protocols including mutating the CDRs (Yang *et al.*, J. Mol. Biol., 254,

392-403, 1995), chain shuffling (Marks *et al.*, Bio/Technology, 10, 779-783, 1992), use of mutator strains of *E. coli* (Low *et al.*, J. Mol. Biol., 250, 359-368, 1996), DNA shuffling (Patten *et al.*, Curr. Opin. Biotechnol., 8, 724-733, 1997), phage display (Thompson *et al.*, J. Mol. Biol., 256, 77-88, 1996) and sexual PCR (Crameri *et al.*,
5 Nature, 391, 288-291, 1998). Vaughan *et al.* (*supra*) discusses these methods of affinity maturation.

The single domain antibody or antibodies of the dual specificity fusion protein may be provided as monomers, dimers or trimers, as required. The desired product may be obtained by adjusting the downstream processing steps the material is
10 subjected to. In one embodiment the processed material is provided as a substantially homogenous monomer. In one embodiment the processed material is provided a substantially homogenous dimer. In one embodiment the processed material is provided as a substantially homogenous trimer.

The present invention also provides an isolated DNA sequence encoding a
15 dual specificity antibody fusion protein of the present invention. The DNA sequences of the present invention may comprise synthetic DNA, for instance produced by chemical processing, cDNA, genomic DNA or any combination thereof.

DNA sequences which encode the dual specificity antibody fusion protein of the present invention can be obtained by methods well known to those skilled in the
20 art. For example, DNA sequences coding for part or all of the antibody fragments, linkers and/or dAbs may be synthesised as desired from the determined DNA sequences or on the basis of the corresponding amino acid sequences.

Standard techniques of molecular biology may be used to prepare DNA sequences coding for the dual specificity antibody fusion protein of the present
25 invention. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate.

The present invention further relates to a cloning or expression vector comprising one or more DNA sequences of the present invention. Accordingly,
30 provided is a cloning or expression vector comprising one or more DNA sequences encoding a dual specificity antibody fusion protein of the present invention. In one preferred embodiment, the cloning or expression vector comprises a single DNA sequence encoding the entire dual specificity antibody fusion protein. Thus, the cloning or expression vector comprises DNA encoded transcription units in sequence
35 such that a translation fusion protein is produced.

Indeed, it will be understood by those skilled in the art that a fusion protein of the invention can have the dAb at the N-terminus or the C-terminus and thus, the dAb DNA encoded transcription unit will be first or last, respectively, within the DNA sequence encoding the translation fusion. Thus, a translation fusion may comprise an
40 N-terminal dAb and a C-terminal Fab or Fab'. Further, a translation fusion may comprise an N-terminal Fab or Fab' and a C-terminal dAb.

It will be appreciated that the heavy chain and light chain of the Fab or Fab' may be incorporated into the same or different vectors. In one embodiment one vector may comprise a translation fusion comprising a Fab or Fab' heavy chain and a C-terminal dAb and another vector may comprise a translation fusion comprising a Fab or Fab' light chain and a C-terminal dAb.

For example, where the desire is to produce a dual specificity antibody fusion protein with the dAb moiety at the N-terminal end of the antibody fragment, the vector will comprise DNA transcription units in sequence order; a DNA transcription unit encoding the dAb moiety, optionally a DNA transcription unit encoding a linker sequence, and a DNA transcription unit encoding an antibody fragment. Where the desire is to produce a dual specificity antibody fusion protein with the dAb moiety at the C-terminal end of the antibody fragment, the vector will comprise DNA transcription units in sequence order; a DNA transcription unit encoding an antibody fragment, optionally a DNA transcription unit encoding a linker sequence, and a DNA transcription unit encoding dAb moiety with specificity for a serum carrier protein, a circulating immunoglobulin molecule, or CD35/CR1, for example, human serum albumin. Thus, a translation fusion of the invention can be in different configurations including, for example but without limitation, dAb-linker-Fab, Fab-linker-dAb, dAb-Fab, Fab-dAb, Fab'-dAb, dAb-Fab', dAb-linker Fab', Fab'-linker-dAb. Where two vectors are used for example, the first may comprise the heavy chain of a Fab or Fab' fused to a dAb and the second may comprise the light chain of a Fab or Fab' fused to a dAb.

DNA code for an antibody fragment comprised within a translation fusion of the invention can be incorporated into a vector as a transcription unit in configurations as known to the person skilled in the art, for example a transcription unit can comprise code for the light chain followed by the heavy chain code, or vice versa; see, in particular, *Humphreys et al.*, 2002, *Protein Expression and Purification*, 26:309-320.

Preferably, a vector according to the present invention comprises an appropriate leader sequence, such as an antibody leader sequence. Such leader sequences are well known in the art.

General methods by which the vectors may be constructed, transfection and transformation methods and culture methods are well known to those skilled in the art. In this respect, reference is made to "Current Protocols in Molecular Biology", 1999, F. M. Ausubel (ed), Wiley Interscience, New York and the Maniatis Manual produced by Cold Spring Harbor Publishing.

Also provided is a host cell comprising one or more cloning or expression vectors comprising one or more DNA sequences encoding a dual specificity antibody fusion protein of the present invention. Any suitable host cell/vector system may be used for expression of the DNA sequences encoding the dual specificity antibody fusion protein. Bacterial, for example *E. coli*, and other microbial systems may be used or eukaryotic, for example mammalian, host cell expression systems may also be

used. Suitable mammalian host cells include NS0, CHO, myeloma or hybridoma cells. Accordingly in one embodiment the fusion protein of the present invention is expressed in *E.coli*. In another embodiment the fusion protein of the present invention is expressed in mammalian cells.

5 The present invention also provides a process for the production of a dual specificity antibody fusion protein comprising culturing a host cell comprising a vector of the present invention under conditions suitable for the expression of protein from the DNA sequence encoding said dual specificity antibody fusion protein. The invention further provides methods for isolating the dual specificity antibody fusion

10 protein.

On production, a dual specificity antibody fusion protein of the present invention may be purified, where necessary, using any suitable method known in the art. For example, but without limitation, chromatographic techniques such as ion exchange, size exclusion, protein G or hydrophobic interaction chromatography may

15 be used.

The size of a dual specificity antibody fusion protein may be confirmed by conventional methods known in the art such as size exclusion chromatography and non-reducing SDS-PAGE. Such techniques can be used, for example to confirm that the protein has not dimerised and/or does not have a portion missing, e.g. the dAb

20 portion. If dimers are detected and a homogenous monomeric product is required then the monomeric dual specificity antibody fusion protein may be purified away from the dimeric species using conventional chromatography techniques as described above.

Dual specificity antibody fusion proteins of the invention are useful in the

25 treatment of diseases or disorders including inflammatory diseases and disorders, immune disease and disorders, fibrotic disorders and cancers.

The term “inflammatory disease” or “disorder” and “immune disease or disorder” includes rheumatoid arthritis, psoriatic arthritis, still's disease, Muckle Wells disease, psoriasis, Crohn's disease, ulcerative colitis, SLE (Systemic Lupus

30 Erythematosus), asthma, allergic rhinitis, atopic dermatitis, multiple sclerosis, vasculitis, Type I diabetes mellitus, transplantation and graft-versus-host disease.

The term “fibrotic disorder” includes idiopathic pulmonary fibrosis (IPF), systemic sclerosis (or scleroderma), kidney fibrosis, diabetic nephropathy, IgA nephropathy, hypertension, end-stage renal disease, peritoneal fibrosis (continuous

35 ambulatory peritoneal dialysis), liver cirrhosis, age-related macular degeneration (ARMD), retinopathy, cardiac reactive fibrosis, scarring, keloids, burns, skin ulcers, angioplasty, coronary bypass surgery, arthroplasty and cataract surgery.

The term “cancer” includes a malignant new growth that arises from epithelium, found in skin or, more commonly, the lining of body organs, for example:

40 breast, ovary, prostate, lung, kidney, pancreas, stomach, bladder or bowel. Cancers

tend to infiltrate into adjacent tissue and spread (metastasise) to distant organs, for example: to bone, liver, lung or the brain.

Thus, according to a further aspect of the invention, there is provided a pharmaceutical composition which comprises an antibody fusion of the invention in
5 association with one or more pharmaceutically acceptable carriers, excipients or diluents. Also provided is the use of an antibody fusion protein of the invention for the manufacture of a medicament for the treatment of a disease or disorder. Most preferably, the disease or disorder is an inflammatory disease or disorder.

Pharmaceutical compositions according to the invention may take a form
10 suitable for oral, buccal, parenteral, subcutaneous, nasal, topical, ophthalmic or rectal administration, or a form suitable for administration by inhalation or insufflation.

Where appropriate, for example if the single domain antibody or antibodies of the antibody fusion protein bind to albumin, it may be desirable to pre-formulate the dual specificity fusion protein with human or recombinant serum albumin, using any
15 suitable method known in the art.

Where the pharmaceutical formulation is a liquid, for example a solution or suspension then the formulation may further comprise albumin, for example human serum albumin, in particular recombinant albumin such as recombinant human serum albumin. Suitable amounts may be in the range of less than 2% w/w of the total
20 formulation, in particular less than 1, 0.5, or 0.1% w/w. This may assist in stabilizing the antibody component in the formulation. The pharmaceutical composition may be lyophilized for reconstitution later, with an aqueous solvent.

In one embodiment there is provided a unit dose container, such as a vial, comprising a lyophilized "antibody" according to the invention.

25 For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methyl cellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogenphosphate); lubricants (e.g.
30 magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before
35 use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles or preservatives. The preparations may also contain buffer salts, flavouring agents, colouring agents or sweetening agents, as appropriate.

Preparations for oral administration may be suitably formulated to give
40 controlled release of the active compound.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

The bispecific antibodies of the invention may be formulated for parenteral administration by injection, e.g. by bolus injection or infusion. Formulations for injection may be presented in unit dosage form, e.g. in glass ampoules or multi-dose containers, e.g. glass vials. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising, preserving and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

In addition to the formulations described above, the bispecific antibodies of the invention may also be formulated as a depot preparation. Such long-acting formulations may be administered by implantation or by intramuscular injection.

For nasal administration or administration by inhalation, the compounds according to the present invention may be conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of a suitable propellant, e.g. dichlorodifluoromethane, fluorotrichloromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas or mixture of gases.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack or dispensing device may be accompanied by instructions for administration.

For topical administration the compounds according to the present invention may be conveniently formulated in a suitable ointment containing the active component suspended or dissolved in one or more pharmaceutically acceptable carriers. Particular carriers include, for example, mineral oil, liquid petroleum, propylene glycol, polyoxyethylene, polyoxypropylene, emulsifying wax and water. Alternatively, the compounds according to the present invention may be formulated in a suitable lotion containing the active component suspended or dissolved in one or more pharmaceutically acceptable carriers. Particular carriers include, for example, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, benzyl alcohol, 2-octyldodecanol and water.

In one embodiment the formulation is provided as a formulation for topical administrations including inhalation.

Suitable inhalable preparations include inhalable powders, metering aerosols containing propellant gases or inhalable solutions free from propellant gases. Inhalable powders according to the disclosure containing the active substance may consist solely of the abovementioned active substances or of a mixture of the abovementioned active substances with physiologically acceptable excipient.

These inhalable powders may include monosaccharides (e.g. glucose or arabinose), disaccharides (e.g. lactose, saccharose, maltose), oligo- and

polysaccharides (e.g. dextrans), polyalcohols (e.g. sorbitol, mannitol, xylitol), salts (e.g. sodium chloride, calcium carbonate) or mixtures of these with one another. Mono- or disaccharides are suitably used, the use of lactose or glucose, particularly but not exclusively in the form of their hydrates.

5 Particles for deposition in the lung require a particle size less than 10 microns, such as 1-9 microns for example from 0.1 to 5 μm , in particular from 1 to 5 μm . The particle size of the active ingredient (such as the antibody or fragment) is of primary importance.

10 The propellant gases which can be used to prepare the inhalable aerosols are known in the art. Suitable propellant gases are selected from among hydrocarbons such as n-propane, n-butane or isobutane and halohydrocarbons such as chlorinated and/or fluorinated derivatives of methane, ethane, propane, butane, cyclopropane or cyclobutane. The abovementioned propellant gases may be used on their own or in mixtures thereof.

15 Particularly suitable propellant gases are halogenated alkane derivatives selected from among TG 11, TG 12, TG 134a and TG227. Of the abovementioned halogenated hydrocarbons, TG134a (1,1,1,2-tetrafluoroethane) and TG227 (1,1,1,2,3,3,3-heptafluoropropane) and mixtures thereof are particularly suitable.

20 The propellant-gas-containing inhalable aerosols may also contain other ingredients such as cosolvents, stabilisers, surface-active agents (surfactants), antioxidants, lubricants and means for adjusting the pH. All these ingredients are known in the art.

25 The propellant-gas-containing inhalable aerosols according to the invention may contain up to 5 % by weight of active substance. Aerosols according to the invention contain, for example, 0.002 to 5 % by weight, 0.01 to 3 % by weight, 0.015 to 2 % by weight, 0.1 to 2 % by weight, 0.5 to 2 % by weight or 0.5 to 1 % by weight of active ingredient.

30 Alternatively topical administrations to the lung may also be by administration of a liquid solution or suspension formulation, for example employing a device such as a nebulizer, for example, a nebulizer connected to a compressor (e.g., the Pari LC-Jet Plus(R) nebulizer connected to a Pari Master(R) compressor manufactured by Pari Respiratory Equipment, Inc., Richmond, Va.).

35 The antibody formats of the invention can be delivered dispersed in a solvent, e.g., in the form of a solution or a suspension. It can be suspended in an appropriate physiological solution, e.g., saline or other pharmacologically acceptable solvent or a buffered solution. Buffered solutions known in the art may contain 0.05 mg to 0.15 mg disodium edetate, 8.0 mg to 9.0 mg NaCl, 0.15 mg to 0.25 mg polysorbate, 0.25 mg to 0.30 mg anhydrous citric acid, and 0.45 mg to 0.55 mg sodium citrate per 1 ml of water so as to achieve a pH of about 4.0 to 5.0. A suspension can employ, for
40 example, lyophilised antibody.

The therapeutic suspensions or solution formulations can also contain one or more excipients. Excipients are well known in the art and include buffers (e.g., citrate buffer, phosphate buffer, acetate buffer and bicarbonate buffer), amino acids, urea, alcohols, ascorbic acid, phospholipids, proteins (e.g., serum albumin), EDTA, sodium chloride, liposomes, mannitol, sorbitol, and glycerol. Solutions or suspensions can be encapsulated in liposomes or biodegradable microspheres. The formulation will generally be provided in a substantially sterile form employing sterile manufacture processes.

This may include production and sterilization by filtration of the buffered solvent/solution used for the for the formulation, aseptic suspension of the antibody in the sterile buffered solvent solution, and dispensing of the formulation into sterile receptacles by methods familiar to those of ordinary skill in the art.

Nebulizable formulation according to the present disclosure may be provided, for example, as single dose units (e.g., sealed plastic containers or vials) packed in foil envelopes. Each vial contains a unit dose in a volume, e.g., 2 ml, of solvent/solution buffer.

The antibodies formats of the present disclosure are thought to be suitable for delivery via nebulisation.

For ophthalmic administration the compounds according to the present invention may be conveniently formulated as microionized suspensions in isotonic, pH-adjusted sterile saline, either with or without a preservative such as a bactericidal or fungicidal agent, for example phenylmercuric nitrate, benzylalkonium chloride or chlorhexidine acetate. Alternatively, for ophthalmic administration compounds may be formulated in an ointment such as petrolatum. For rectal administration the compounds according to the present invention may be conveniently formulated as suppositories. These can be prepared by mixing the active component with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and so will melt in the rectum to release the active component. Such materials include, for example, cocoa butter, beeswax and polyethylene glycols.

The quantity of a compound of the invention required for the prophylaxis or treatment of a particular condition will vary depending on the compound chosen and the condition of the patient to be treated. In general, however, daily dosages may range from around 10 ng/kg to 1000 mg/kg, typically from 100 ng/kg to 100 mg/kg, e.g. around 0.01 mg/kg to 40 mg/kg body weight for oral or buccal administration, from around 10 ng/kg to 50 mg/kg body weight for parenteral administration, and from around 0.05 mg to around 1000 mg, e.g. from around 0.5 mg to around 1000 mg, for nasal administration or administration by inhalation or insufflation.

Preferred features of each embodiment of the invention are as for each of the other embodiments *mutatis mutandis*.

Comprising in the context of the present specification is intended to meaning including.

Where technically appropriate embodiments of the invention may be combined.

5 Embodiments are described herein as comprising certain features/elements. The disclosure also extends to separate embodiments consisting or consisting essentially of said features/elements.

 The invention will now be described with reference to the following examples, which are merely illustrative and should not in any way be construed as limiting the
10 scope of the present invention.

 The present description relates to a multivalent antibody fusion protein which comprises an Fab or Fab' fragment, with a first specificity for an antigen of interest, and further comprises two single domain antibodies (dAb) with specificity for a second antigen of interest, wherein the two single domain antibodies are linked by a disulfide bond and wherein
15 the two single domain antibodies are a complementary V_H/V_L pair and the V_H dAb is directly or indirectly connected to the C-terminus of the Fab or Fab' heavy chain and the V_L dAb is directly or indirectly connected to the C-terminus of the Fab or Fab' light chain.

 The present description also relates to a pharmaceutical composition comprising the multivalent antibody fusion protein as defined herein in association with one or more
20 pharmaceutically acceptable carriers, excipients or diluents.

 The present description also relates to the multivalent antibody fusion protein as defined herein, for use in therapy.

List of Figures:

25 Figure 1: Diagrammatic representation of Fab-dAbs where the dAb is at the C-terminus

 Figure 2A: Diagrammatic representation of Fab-didAbs

 Figure 2B: Diagrammatic representation of Fab-didAbs with additional disulfide stabilisation between the dAbs.

30 Figure 3: SDS PAGE analysis of FabA-dAbL3 (CK-SG₄SE) (1) and FabA-dAbL3 (CK-G[APAPA]₂) (2).

 Figure 4: Western blot analysis of FabA-dAbL3 (CK-SG₄SE) (1) and FabA-dAbL3 (CK-G[APAPA]₂) (2).

30a

Figure 4a: SDS PAGE of FabB-didAbs

Lane M = SeeBlue markers

Lanes 1 & 2 = IgG control

Lane 3 = FabB

5 Lane 4 = FabB-didAb, -dAbL1 (CK-G4Sx2) & dAbH1 (CH1-G4Sx2)

Lane 5 = FabB-didAb, -dAbL2 (CK-G4Sx2) & dAbH2 (CH1-G4Sx2)

Figure 5: Sequences of domain antibodies dAbH1, dAbH2, dAbL1 and dAbL2 and the CDRs derived from each of those antibodies.

Figure 6: FabB-dAb constructs comprising FabB heavy or light chain variable domain
10 fused to a domain antibody.

Figure 7: Fab'A heavy and light chain sequences and FabA heavy chain sequence..

Figure 8a, 8b & 8c: Murinised Fab-didAb amino acid sequences.

Figure 8a shows the amino acid sequence of CDRs in various murine dAbs.

Figure 8b shows the amino acid sequence of mFabD-mdidAb:


15 dAbL1(CK-G4Sx2)

dAbH1(CH1-G4Sx2)

dAbL2(CK-G4Sx2) &

dAbH2(CH1-G4Sx2)

Figure 8c shows the amino acid sequence of mFabD-mdidAb:



dAbL1(CK-G4Sx2) &
dAbH1(CH1-G4Sx2)mFabC-mdAbH1
dAbL2(CK-G4Sx2) &
dAbH2(CH1-G4Sx2)

- 5 Figure 9 shows SDS PAGE of FabB-didAbs
Lanes 1 & 4 are Fab'B
Lanes 2 & 5 are FabB-didAb, -dAbL1(CK-G4Sx2) & -dAbH1(CH1-G4Sx2)
Lanes 3 & 6 are FabB-didAb, -dAbL2(CK-G4Sx2) & -dAbH2(CH1-G4Sx2)
- 10 Figure 10 shows a diagrammatic representation of a Thermofluor thermal stability assay.
Figure 11 shows a plot of HAS-FITC signal/HAS-FITC mixes bound to activated mouse T cells.
Figure 12 shows a plot of an aggregation stability assay.
Figure 13 shows in vivo concentration profiles over time after subcutaneous and
15 intravenous dosing
Figure 14A, B and C show certain CD4+ cell and CD8+ cell readouts
Figure 15 shows SDS-PAGE analysis of FabB-645Fv
Figure 16 shows size exclusion analysis of FabB-645Fv
Figure 17 shows thermograms of FabB-645Fv with various linker lengths.
20 Figure 18 shows SDS-PAGE analysis of certain FabB constructs
Figure 19 shows size exclusion analysis of various FabB-645Fv constructs
Figures 20 to 24 show sequences for certain formats.

KEY

- 25 -645Fv equates to didAbL1 and H1 (the linker used for each dAB will be the same unless indicated otherwise).
648Fv equates to didAbL2 and H2 (the linker used for each dAB will be the same unless indicated otherwise).
-645dsFv equates to didAbL1 and H1 (the linker used for each dAB will be the same unless indicated otherwise) wherein L1 and H1 are stabilised by a
30 disulfide bond.
-648dsFv equates to didAbL2 and H2 (the linker used for each dAB will be the same unless indicated otherwise) wherein L2 and H3 are stabilised by a disulfide bond.
35 FabΔ are Fabs which lack the interchain cysteine bond (ie between CH and CL or CK)

Experimental:

Abbreviations: unless the context indicates otherwise “m” as a pre-fix is intended to refer to murine.

Unless the context indicates otherwise “h” as a pre-fix is intended to refer to human.

- 5 Fab A, Fab B, Fab C and Fab D components may be provided below in different formats.

EXAMPLE 1. Production of a dAb specific for human serum albumin

- 10 An in-frame DNA encoded transcription unit encoding a dAb with specificity for human serum albumin was produced using recombinant DNA technology.

Where desired an in-frame DNA encoded transcription unit encoding a dAb with specificity for a recruitment protein can be produced using recombinant DNA technology.

15 **EXAMPLE 2. Production of antibody fragment**

- For fusion of a dAb to the C-terminus of the light chain, DNA was synthesised encoding a human kappa light chain constant region (with the Km3 allotype of the kappa constant region), a peptide linker and a dAb and cloned as a SacI-PvuII restriction fragment into the UCB-Celltech in-house expression vector pTTOD(Fab) (a derivative of pTTO-1, described in Popplewell *et al.*, Methods Mol. Biol. 2005; 20 308: 17-30) which contains DNA encoding the human gamma-1 CH1 constant region. This gave rise to a dicistronic gene arrangement consisting of the gene for the humanised light chain fused via a linker to a dAb followed by the gene for the humanised heavy chain Fab fragment, both under the control of the tac promoter. 25 Also encoded is a unique BspEI site upstream of the Gly4Ser linker, or an AscI site upstream of the Ala-Pro-rich linker.

- For fusion of a dAb the C-terminus of the heavy chain, DNA was synthesised encoding a human CH1 fragment (of the $\gamma 1$ isotype) followed by a linker encoding sequence and a dAb. This was subcloned as an ApaI-EcoRI restriction fragment into 30 the UCB-Celltech in-house expression vector pTTOD(Fab) (a derivative of pTTO-1, described in Popplewell *et al.*, above) which contains DNA encoding the human gamma-1 CH1 constant region. This gave rise to a dicistronic gene arrangement consisting of the gene for the humanised light chain a non-coding intergenic sequence and followed by a heavy chain fused via a linker to a dAb, both under the control of 35 the tac promoter. The recombinant expression plasmid was transformed into the *E. coli* strain W3110 in which expression is induced by addition of IPTG. Expression experiments were performed at small scale initially (5ml culture volumes) with addition of 200uM IPTG at OD(600nm) of approx. 0.5, cells were harvested 2 hours post induction and extracted overnight at 30°C in Tris/EDTA. Clarified extracts were 40 used for affinity analysis by Biacore. Constructs giving promising expression yields and activities were selected for fermentation.

Methods applicable to the following Examples

In the following examples the antibody chain to which the dAb is fused is denoted either as CK or LC for the cKappa light chain and as CH1 or HC for the heavy chain constant domain, CH1.

Construction of FabA-dAb fusion plasmids for expression in E.coli

Fab-dAb fusion proteins were constructed by fusing dAbL3 or dAbH4 to the C-terminus of the constant region of either the light or heavy chain of FabA. A flexible (SGGGGSE (SEQ ID NO:1)) or a rigid (G(APAPA)₂ (SEQ ID NO: 34)) linker was used to link the dAb to the cKappa region (SEQ ID NO:75) whereas the linker DKTHTS (SEQ ID NO:2) was used to link the dAb to the CHI region (SEQ ID NO:76). The DNA sequence coding for the constant region-dAb fusion was manufactured synthetically as fragments to enable sub-cloning into the FabA sequence of the in-house pTTOD vector.

Light chain-dAb fusions were constructed by sub-cloning the *SacI*-*ApaI* fragment of the synthesized genes, encoding a C-terminal cKappa fused to either dAbL3 or dAbH4 via either a (SGGGGSE (SEQ ID NO:1)) or a rigid (G(APAPA)₂ (SEQ ID NO: 34)) linker, into the corresponding sites of a plasmid capable of expressing FabA.

Heavy chain-dAb fusions were constructed by sub-cloning the *ApaI*-*EcoRI* fragment of the synthesised genes, encoding a C-terminal CHI fused to either dAbL3 or dAbH4 via a DKTHTS linker, into the corresponding sites of a plasmid capable of expressing FabA.

Fab' A is derived from an IL-1 beta binding antibody, the heavy and light chain sequences of which are provided in SEQ ID NOs:74 and 75 respectively as shown in Figure 7. In Fab' A where the light chain has a dAb attached, the hinge of the heavy chain was altered to DKTHTS even where no dAb is attached to the heavy chain (SEQ ID NO:76).

FabA comprises the same light chain sequence (SEQ ID NO:75) and a truncated heavy chain sequence which terminates at the interchain cysteine (SEQ ID NO:77). dAbL3 and dAbH4 are light and heavy chain domain antibodies respectively which bind human serum albumin.

Construction of FabA-didAb fusion plasmids for expression in E.coli

FabA-didAb with dAbL3 or dAbH4 on both light and heavy chains were constructed by sub-cloning the *ApaI*-*EcoRI* fragment coding for CH1-dAb fusions into the existing Fab-dAb plasmids where the dAb is fused to the light chain via the flexible linker.

Construction of FabB-dAb fusion plasmids for expression in mammalian cells

The FabB-dAbs, FabB-dAbH1 (CH1-G₄Sx2), FabB-dAbH2 (CH1-G₄Sx2), FabB-dAbL1 (CH1-G₄Sx2), FabB-dAbL2 (CH1-G₄Sx2) were all assembled by PCR then cloned into a mammalian expression vector under the control of the HCMV-MIE promoter and SV40E polyA sequence. These were paired with a similar vector
 5 containing the FabB light chain for expression in mammalian cells (see below).

FabB is derived from an antibody which binds a cell surface co-stimulatory molecule. dAbH1, dAbH2, dAbL1 and dAbL2 were obtained as described in Example 3.

10 **Mammalian expression of FabB-dAbs and dAbs**

HEK293 cells were transfected with the heavy and light chain plasmids using Invitrogen's 293fectin transfection reagent according to the manufacturer's instructions. Briefly, 2µg heavy chain plasmid + 2µg light chain plasmid was incubated with 10µl 293fectin + 340µl OptiMem media for 20mins at RT. The
 15 mixture was then added to 5x10⁶ HEK293 cells in suspension and incubated for 4 days with shaking at 37°C.

Biacore

Binding affinities and kinetic parameters for the interactions of Fab-dAb constructs
 20 were determined by surface plasmon resonance (SPR) conducted on a Biacore T100 using CM5 sensor chips and HBS-EP (10mM HEPES (pH7.4), 150mM NaCl, 3mM EDTA, 0.05% v/v surfactant P20) running buffer. Fab-dAb samples were captured to the sensor chip surface using either a human F(ab')₂-specific goat Fab (Jackson ImmunoResearch, 109-006-097) or an in-house generated anti human CH1
 25 monoclonal antibody. Covalent immobilisation of the capture antibody was achieved by standard amine coupling chemistry.

Each assay cycle consisted of firstly capturing the Fab-dAb using a 1 min injection, before an association phase consisting of a 3 min injection of antigen, after which
 30 dissociation was monitored for 5 min. After each cycle, the capture surface was regenerated with 2 x 1 min injections of 40mM HCl followed by 30s of 5mM NaOH. The flow rates used were 10µl/min for capture, 30µl/min for association and dissociation phases, and 10µl/min for regeneration.

For kinetic assays, a titration of antigen (for human serum albumin typically 62.5nM-
 35 2µM, for IL-1β 1.25-40nM) was performed, a blank flow-cell was used for reference subtraction and buffer-blank injections were included to subtract instrument noise and drift.

Kinetic parameters were determined by simultaneous global-fitting of the resulting sensorgrams to a standard 1:1 binding model using Biacore T100 Evaluation software.

In order to test for simultaneous binding, 3 min injections of either separate 5 μ M HSA or 100nM IL-1 β , or a mixed solution of 5 μ M HSA and 100nM IL-1 β were injected over the captured Fab-dAb.

5 **Fab-dAb Purification from *E.coli***

Periplasmic extraction

E.coli pellets containing the Fab-dAbs within the periplasm were re-suspended in original culture volume with 100mM Tris/HCl, 10mM EDTA pH 7.4. These suspensions were then incubated at 4°C for 16 hours at 250rpm. The re-suspended
10 pellets were centrifuged at 10000xg for 1 hour at 4°C. The supernatants were removed and 0.45 μ m filtered.

Protein-G capture

The Fab-dAbs were captured from the filtered supernatant by Protein-G chromatography. Briefly the supernatants were applied, with a 20 minute residence
15 time, to a Gammabind Plus Sepharose (GE Healthcare) column equilibrated in 20mM phosphate, 150mM NaCl pH7.1. The column was washed with 20mM phosphate, 150mM NaCl pH7.1 and the bound material eluted with 0.1M glycine/HCl pH2.8. The elution peak was collected and pH adjusted to ~pH5 with 1M sodium acetate. The pH adjusted elutions were concentrated and diafiltered into 50mM sodium acetate
20 pH4.5 using a 10k MWCO membrane.

Ion Exchange

The Fab-dAbs were further purified by cation exchange chromatography at pH4.5 with a NaCl elution gradient. Briefly the diafiltered Protein-G eluates were applied to
25 a Source15S (GE Healthcare) column equilibrated in 50mM sodium acetate pH4.5. The column was washed with 50mM sodium acetate pH4.5 and the bound material eluted with a 20 column volume linear gradient from 0 to 1M NaCl in 50mM sodium acetate pH4.5. Third column volume fractions were collected through out the gradient. The fractions were analysed by A280 and SDS-PAGE and relevant fractions
30 pooled.

Gel filtration

If required the Fab-dAbs were further purified by gel filtration. Briefly the FabA-dAbL3 (CK-SG₄SE) pooled ion exchange elution fractions were applied to a
35 Superdex200 (GE Healthcare) column equilibrated in 50mM sodium acetate, 125mM NaCl pH 5.0 and eluted with an isocratic gradient of 50mM sodium acetate, 125mM NaCl pH 5.0. 1/120 column volume fractions were collected through out the gradient. The fractions were analysed by A280 and SDS-PAGE and relevant fractions pooled. For Fab-dAbs that did not undergo gel filtration, the pooled ion exchange elution
40 fractions were concentrated and diafiltered into 50mM sodium acetate, 125mM NaCl pH 5.0 using a 10k MWCO membrane.

SDS-PAGE

Samples were diluted with water where required and then to 10 μ l was added 10 μ L 2X sample running buffer. For non-reduced samples, 2 μ L of 100mM NEM was added at this point, for reduced samples 2 μ L of 10X reducing agent was added. The sample
 5 were vortexed, incubated at 85°C for 5 mins, cooled and centrifuged at 12500 rpm for 30secs. The prepared samples were loaded on to a 4-20% acrylamine Tris/Glycine SDS gel and run for 100mins at 125V. The gels were either transferred onto PVDF membranes for Western blotting or stained with Coomassie Blue protein stain.

Western Blotting

Gels were transferred to PVDF membranes in 12mM Tris, 96mM glycine pH8.3 for 16 hours at 150mA. The PVDF membrane was block for 1hr with 2% MarvelTM in PBS + 0.1% Tween20 (Blocking buffer)
anti-light chain
 15 HRP-rabbit anti-human kappa light chains, 1/5000 dilution in blocking buffer for 1hr.
anti-heavy chain
 mouse anti-human heavy chain, 1/7000 dilution in blocking buffer for 1hr. Followed by HRP-goat anti-mouse, 1/2000 dilution in blocking buffer for 1hr.
anti-His tag
 20 rabbit anti-His6, 1/1000 dilution in blocking buffer for 1hr. Followed by HRP-goat anti-rabbit IgG, 1/1000 dilution in blocking buffer for 1hr.
 All blots were washed 6 times with 100ml PBS + 0.1% Tween20 for 10 minutes per wash. The blots were developed with either ECL reagent for 1min before being exposed to Amersham Hyperfilm, or metal enhanced DAB reagent for 20-30 minutes
 25 followed by water.

High temperature reverse phase HPLC

Samples (2 μ g) were analysed on a 2.1mm C8 Poroshell column at 80°C, with a flow rate of 2ml/min and a gradient of 18–38% B over 4mins. A = 0.1% TFA in H₂O
 30 B = 0.065% TFA in 80:20 IPA:MeOH. Detection is by absorption at 214nm.

ELISA

The yields of Fab-dAb were measured using a sandwich ELISA. Briefly, the Fab-dAb was captured with an anti-CH1 antibody then revealed with an anti-kappa-HRP.
 35

FACS

Samples (mFabD-didAb's) were incubated with 5 μ g/ml FITC (fluorescein isothiocyanate) labelled HSA for 45 min. The sample/HSA-FITC incubations were then added to activated mouse CD4⁺ T-cells and incubated for a further 45 min. The
 40 cells were washed with PBS and the cell associated fluorescence measured by FACS (fluorescence activated cell sorting).

EXAMPLE 3**Generating anti-albumin antibodies**

½ lop rabbits were immunised with recombinant chromatopure human serum albumin (purchased from Jackson). Rabbits received 3 immunisations of 100ug HSA protein subcutaneously, the first immunisation in complete Freund's adjuvant and subsequent immunisations in incomplete Freund's. Antibodies 1 and 2, 646, 647, and 649 which bind human, mouse and rat serum albumin were isolated using the methods described in WO04/051268. Genes for the heavy chain variable domain (VH) and light chain variable domain (VL) of antibodies 1 and 2 were isolated and sequenced following cloning via reverse transcription PCR.

The light chain grafted sequences were sub-cloned into the rabbit light chain expression vector pVRbck which contains the DNA encoding the rabbit C-Kappa constant region. The heavy chain grafted sequences were sub-cloned into the rabbit heavy chain expression vector pVRbHFab, which contains the DNA encoding the rabbit Fab' heavy chain constant region. Plasmids were co-transfected into CHO cells and the antibodies produced screened for albumin binding and affinity (Table 1). Transfections of CHO cells were performed using the Lipofectamine™ 2000 procedure according to manufacturer's instructions (Invitrogen, catalogue No. 11668).

Generating Humanised domain antibodies dAbL1, dAbH1, dAbL2 and dAbH2

Humanised VL and VH regions were designed using human V-region acceptor frameworks and donor residues in the framework regions. One grafted VL region (L1 (SEQ ID NO:53) and L2 (SEQ ID NO:55)) and one VH region (H1 (SEQ ID NO:52) and H2 (SEQ ID NO:54)) were designed for each of antibodies 1 and 2 respectively and genes were built by oligonucleotide assembly and PCR mutagenesis. The grafted domain antibodies and their CDRs are shown in Figure 5.

Table 1: Affinities of anti-albumin antibodies

	as rabbit Fab		as humanised IgG		
	HSA	MurineSA	HumanSA	MurineSA	RatSA
	nM	nM	nM	nM	nM
Antibody 1 (Antibody 645)	0.31	2.6	0.82	2.9	7.9
Antibody 2 (Antibody 648)	0.33	12	0.13	23	54
Antibody 646	0.14	1.6	0.57	1.7	4.5
Antibody 647	0.60	3.6	1.3	26	10
Antibody 649	0.54	13	0.32	17	44

EXAMPLE 4: Analysis of FabB-dAbs expressed in mammalian cells

FabB-dAb constructs were produced as described in the methods and the supernatants from the transfected HEK293 cells containing the FabB-dAbs were tested directly in BIAcore.

- 5 Kinetic analysis was conducted to assess the interaction of HSA with FabB-dAb constructs. These consisted of either dAbL1, dAbH2 or dAbL3 fused to the C-terminus of CH1 of FabB (See Figure 6). The FabB-dAbL1 has a higher affinity for HSA, $K_D = 170\text{nM}$, than FabB-dAbL3, $K_D = 392\text{nM}$. The FabB-dAbH2 was shown to possess the poorest affinity towards HSA, $K_D = 1074\text{nM}$, see Table 2.

10

Table 2

Construct	$k_a (\times 10^4 \text{M}^{-1} \text{s}^{-1})$	$k_d (\times 10^{-3} \text{s}^{-1})$	$K_D (\times 10^{-9} \text{M})$
FabB-dAbL1 (CH1-G ₄ Sx2)	1.91 ± 0.74	2.18 ± 1.21	170 ± 78
FabB-dAbH2 (CH1-G ₄ Sx2)	2.66 ± 0.39	29 ± 4.76	1074 ± 42
FabB-dAbL3 (CH1-G ₄ Sx2)	2.63 ± 0.39	9.87 ± 1.63	392 ± 119

Affinity and kinetic parameters determined for the binding of HSA to FabBs fused to dAbL1, dAbH2 or dAbL3. The data shown are mean values $\pm$ SEM. (For FabB-dAbL1 and FabB-dAbH2 n=4. For FabB-dAbL3 n=2).

15

SDS-PAGE and western blotting of the FabB-dAb proteins confirmed that the FabB-dAbs produced were of the expected size.

EXAMPLE 5: Analysis of FabB-didAbs expressed in mammalian cells

- 20 FabB-didAb constructs were produced as described in the methods and the supernatants from the transfected HEK293 cells containing the didAbs tested directly in BIAcore.

Further analysis was performed using didAb constructs in which single dAbs were fused to both heavy and light C-termini of Fab. Constructs in which the didAb was derived from a natural heavy and light variable domain pairing showed a marked improvement in affinity compared to the single dAb alone (table 2 and 3). The didAb fusion consisting of two identical dAbL1s showed no improvement in affinity over that seen for the single dAbL1 (data not shown).

25

Table 3

Construct	k_a ($\times 10^4 \text{M}^{-1} \text{s}^{-1}$)	k_d ($\times 10^{-3} \text{s}^{-1}$)	K_D ($\times 10^{-9} \text{M}$)
FabB-didAb, -dAbL1 (CK-G ₄ Sx2) & dAbH1 (CH1-G ₄ Sx2)	1.78	0.16	9
FabB-didAb, -dAbL2 (CK-G ₄ Sx2) & dAbH2 (CH1-G ₄ Sx2)	0.54	0.21	39

Affinity and kinetic parameters determined for the binding of HSA to FabBs fused to both dAbL1 & dAbH1 or dAbL2 & dAbH2.

- 5 SDS-PAGE of the FabB-didAb proteins confirmed that the FabB-didAbs expressed well and were of the expected size (See Figure 4a). Note this SDS PAGE gel is total protein expressed by the cell.

EXAMPLE 6

10 Analysis of purified FabA-dAbs

- Plasmids for expression of the Fab-dAbs, Fab'A-dAbL3 (CK-SG₄SE) Fab'A-dAbL3 (CK-G[APAPA]₂) in E.coli were constructed as described in the methods. The Fab-dAbs were expressed into the periplasm of the E.coli and purified to homogeneity as described in the methods. The purity of the Fab-dAbs were assessed by high
- 15 temperature reverse phase HPLC, SDS-PAGE and Western blotting. The Fab-dAbs were also assessed for antigen binding by Biacore.

High temperature reverse phase HPLC

- High temperature reverse phase HPLC as performed as described in the methods gave
- 20 quantitative analysis of all species contained in FabA-dAbL3 (CK-SG₄SE) and FabA-dAbL3 (CK-G[APAPA]₂). The percentage of each species present is shown in table 4.

Table 4: Quantification of species present in Fab-dAb batches

Species	Fab'A-dAbL3 (CK-SG ₄ SE)	Fab'A-dAbL3 (CK-G[APAPA] ₂)
Peak 1	0.6%	1.8%
Peak 2	0.6%	0.0%
Peak 3	1.0%	0.3%
Peak 4	0.9%	0.8%
Fab-dAb peak	85.5%	92.9%
Di Fab-dAb peak	11.5%	4.2%

SDS-PAGE

Fab-dAb samples were prepared under non-reduced and reduced conditions and run on a gel as described in the methods. The gel was Coomassie stained. The banding profile of both Fab-dAb samples, Fab'A-dAbL3 (CK-SG₄SE) and Fab'A-dAbL3 (CK-G[APAPA]₂), corresponds well to the profile observed by high temperature reverse phase HPLC (figure 3).

Western Blot

Fab-dAb samples were subjected to non-reduced SDS-PAGE followed by western blot analysis with anti-light chain and anti-heavy chain antibodies as described in the methods. This confirmed that the dAb was on the light chain of the Fab and that the heavy chain was unmodified in both samples (figure 4). It also demonstrates that all bands detected by coomassie stained, non-reduced SDS PAGE are Fab-dAb related products.

Biacore

Kinetic analysis by SPR as described in the methods was used to assess the binding of human serum albumin to Fab'A-dAbL3 (CK-SG₄SE) and Fab'A-dAbL3 (CK-G[APAPA]₂). The results in table 5 demonstrate that both constructs are able to bind human serum albumin with a similar affinity (K_D) of approximately 1 μ M.

Table 5

Construct	k_a ($\times 10^4 \text{M}^{-1} \text{s}^{-1}$)	k_d ($\times 10^{-2} \text{s}^{-1}$)	K_D ($\times 10^{-9} \text{M}$)
Fab'A-dAbL3 (CK- SG ₄ SE)	3.44	1.42	411
Fab'A-dAbL3 (CK- G[APAPA] ₂)	9.61	2.85	296

Further kinetic analysis demonstrated that all the fusion constructs retained the interaction characteristics of the original FabA towards IL-1 β , table 6, with only minor differences seen in the kinetic and affinity parameters.

Table 6

Construct	k_a ($\times 10^5 \text{M}^{-1} \text{s}^{-1}$)	k_d ($\times 10^{-5} \text{s}^{-1}$)	K_D ($\times 10^{-12} \text{M}$)
Fab'A-dAbL3 (CK- SG ₄ SE)	1.90	4.21	221
Fab'A-dAbL3 (CK- G[APAPA] ₂)	2.17	3.99	184
Fab'A	2.02	6.46	320

The potential for each construct to bind simultaneously to both human serum albumin and the IL-1 β antigen was assessed by capturing each construct to the sensor chip surface, before performing either separate 3 min injections of 5 μ M human serum albumin or 100nM IL-1 β , or a mixed solution of both 5 μ M human serum albumin and 100nM IL-1 β . For each Fab-dAb construct the response seen for the combined

HSA/IL-1 β solution was almost identical to the sum of the responses of the independent injections, see table 7. This shows that the Fab-dAbs are capable of simultaneous binding to both IL-1 β and human serum albumin, and that binding of either IL-1 β or human serum albumin does not inhibit the interaction of the other.

- 5 The original FabA bound only to IL-1 β , with negligible binding to human serum albumin.

Table 7

Construct	Analyte	Binding (RU)	
Fab'A-dAbL3 (CK- SG ₄ SE)	HSA + IL-1 β	37.6	
	HSA	13.2	(37.9)
	IL-1 β	24.7	
Fab'A-dAbL3 (CK- G[APAPA] ₂)	HSA + IL-1 β	61.9	
	HSA	30.7	(63.6)
	IL-1 β	32.9	
Fab'A	HSA + IL-1 β	30.3	
	HSA	1.3	(30.0)
	IL-1 β	28.7	

- 10 The table above shows the binding response (RU) seen for each construct after separate injections of HSA or IL-1 β , or injection of premixed HSA and IL-1 β . In each case the final concentration was 5 μ M for HSA and 100nM for IL-1 β . The sum of the individual HSA and IL-1 β responses is shown in parentheses.

EXAMPLE 7 FabA didAbs

15 **Expression of FabA-didAbs in *E.coli***

- FabA-dAbs and FabA-didAb fusions terminating with a C-terminal histidine tag (HIS6 tag) were expressed in *Escherichia coli*. After periplasmic extraction, dAb fusion proteins were purified via the C-terminal His6 tag. Fab expression was analysed by Western blotting of a non-reduced gel with anti-CH1 and anti-cKappa antibodies. FabA-dAb and FabA-didAb were expressed as full-length proteins and were shown to react to both antibody detection reagents.
- 20

Analysis of FabA-didAbs expressed in *E.coli*

- Further analysis was conducted to characterise the binding of HSA to FabA constructs to which one or more dAbs were fused. Binding assays were performed on a variety of constructs in which dAbL3 or dAbH4 fused to either the light or heavy chain of the FabA (see Table 8 for details of the constructs and summary of the binding data). Although constructs carrying only dAbH4, on either the light or heavy chain, were seen to bind HSA with comparatively poor affinity ($\approx$ 9 μ M and 3 μ M respectively), higher affinity binding was observed for constructs carrying dAbL3, either as a single
- 25
- 30

fusion (on either light or heavy chain) or partnered with a second dAb (dAbL3 or dAbH4) on the opposing chain.

Table 8

Construct	$k_a (\times 10^4 \text{M}^{-1} \text{s}^{-1})$	$k_d (\times 10^{-3} \text{s}^{-1})$	$K_D (\times 10^{-9} \text{M})$
FabA	-	-	nb
FabA-dAbL3 (LC-SG4SE)	4.46	16.2	363
FabA-dAbH4 (LC SG4SE)	-	-	9142
FabA-dAbL3 (HC-DKTHTS)	8.24	15.4	187
FabA-dAbH4 (HC-DKTHTS)	-	-	2866
FabA-didAb, -dAbL3 (LC-SG4SE) & -dAbL3 (HC-DKTHTS)	3.00	15.1	502
FabA-didAb, -dAbL3 (LC-SG4SE) & -dAbH4 (HC-DKTHTS)	4.36	16.3	373

- 5 Affinity and kinetic parameters determined for the binding of HSA to FabAs carrying dAbL3 or dAbH4 on either light chain (LC) or heavy chain (HC) or both as indicated. No binding (nb) of HSA to the original FabA was detected. The interaction kinetics for the binding of HSA to the FabA with (dAbH4 on HC) or (dAbH4 on LC), were too rapid to determine, therefore affinity (KD) was determined from steady-state
10 binding.

EXAMPLE 8

Expression and purification of FabB-didAbs

Mammalian expression

- 15 Prior to transfection CHO-XE cells were washed in Earls Balanced Salts Solution (EBSS), pelleted and resuspended in EBSS at 2×10^8 cells/ml. Heavy and light chain plasmids were added to the cells at a total concentration of 400ug. Optimised electrical parameters for 800μl cells/DNA mix on the in-house electroporator were used for transfection. Transfected cells were directly transferred to 1L CD-CHO
20 media supplied with glutamax, HT and antimycotic antibiotic solution. Cells were incubated, shaking at 37°C for 24 hours and then shifted to 32°C. Sodium Butyrate 3mM was added on day 4. Supernatants were harvested on day 14 by centrifugation at 1500xg to remove cells. Expression levels were determined by ELISA.

25 **Mammalian expression supernatant concentration**

The mammalian supernatants containing ~55μg/ml of FabB-didAb as assessed by ELISA were concentrated from 1.8L to 200ml using a Minisette concentrator fitted with a 10kDa molecular weight cut off polyethersulphone (PES) membrane.

Protein-G purification

The concentrated supernatants were applied to a Gammabind Plus Sepharose (GE Healthcare) column equilibrated in 20mM phosphate, 150mM NaCl pH7.1. The column was washed with 20mM phosphate, 150mM NaCl pH7.1 and the bound
5 material eluted with 0.1M glycine/HCl pH2.7. The elution peak was collected and pH adjusted to ~pH7 with 2M Tris/HCl pH8.8. The pH adjusted elutions were concentrated to 1mg/ml and diafiltered into 20mM phosphate, 150mM NaCl pH7.1 using a 10kD molecular weight cut off PES membrane.

10 SDS-PAGE

Samples were diluted with water where required and then to 26μl was added 10μL 4X LDS sample running buffer. For non-reduced samples, 4μL of 100mM NEM was added and for reduced samples 4μL of 10X reducing agent was added. The samples were vortexed, incubated at 85°C for 5 mins, cooled and centrifuged at 12500 rpm for
15 30secs. The prepared samples were loaded on to a 4-20% acrylamine Tris/Glycine SDS gel and run for 110mins at 125V. The gels were stained with Coomassie Blue protein stain.

ELISA

20 The yields of Fab-didAb were measured using a sandwich ELISA. Briefly, the Fab-didAb was captured with an anti-CH1 antibody then revealed with an anti-kappa-HRP.

SDS-PAGE

25 FabB and FabB-didAb samples were prepared under non-reduced and reduced conditions and separated on a gel and stained as described in the methods. See Figure 9.

EXAMPLE 9**30 Thermofluor thermal stability assay on FabB-Fv**

Samples (1μl of sample at ~1mg/ml, 8μl of PBS and 1μl of 30x stock of Sypro orange fluorescent dye) were run in quadruplicate in 384 well plates. The plate is heated from 20-99°C using a 7900HT fast real-time PCR system and the fluorescence (excitation at 490nm, emission at 530nm) measured. The results are shown in Table
35 D and Figure 10.

Table 9

	T_m °C (Fab)	T_m °C (Fv)
FabB-didAb, -dAbL1(CK-G ₄ Sx2) & -dAbL1(CH1-G ₄ Sx2)	81.9 ± 0.6	68.5 ± 0.5
FabB-didAb, -dAbL2(CK-G ₄ Sx2) & -dAbL2(CH1-G ₄ Sx2)	82.4 ± 0.2	70.6 ± 0.8

EXAMPLE 10**Aggregation stability assay of FabB-Fv**

- 5 Samples at 1mg/ml in PBS were incubated at 25°C with vortexing at 1400rpm. The absorbance is measured at 595nm. This absorbance is due to light scattered by particles and can be correlated with sample aggregation. Both FabB-645Fv (G₄Sx2) and FabB-648Fv (G₄Sx2) are as resistant to aggregation as FabB alone. They are all more resistant to aggregation than the IgG control. (Figure 12)

10

EXAMPLE 11**pH dependency of Fab-Fv binding to HSA**

- Binding affinities for the interactions of Fab-Fv constructs with HSA were determined as described in the methods except that the running buffers at pH5.0, 5.5, 6.0 and 7.0
15 were created by mixing 40mM citric acid, 150mM NaCl, 3mM EDTA, 0.05% v/v surfactant P20 and 80mM disodium hydrogen phosphate, 150mM NaCl, 3mM EDTA, 0.05% v/v surfactant P20 to give the desired pH.

- The affinity of FabB-645Fv (G₄Sx2) for HSA is unaffected by pH from 7.4 (standard assay pH) to 5.0. The affinity of FabB-648Fv (G₄Sx2) for HSA is affected by pH and
20 there is approximately a 10 fold loss in affinity between pH7.4 and pH5.0.

Table 10

	K_D (x10⁻⁹M)			
	pH7.0	pH6.0	pH5.5	pH5.0
FabB-645Fv (G ₄ Sx2)	13.3	12.5	10.7	7.1
FabB-648Fv (G ₄ Sx2)	3.3	11.1	24.1	47.8

EXAMPLE 12**In vivo murine PK of FabB-Fv**

- 25 The pharmacokinetics of FabB-645Fv (G₄Sx2) and FabB-648Fv (G₄Sx2) in male BALB/c mouse were determined following a single administration at 10mg/kg either subcutaneously (sc) or intravenously (iv). Six mice were dosed for each construct and route of administration. Serial blood samples (30 µL) were collected from the tail vein at the following time points: 1, 4, 8, 24, 48, 72, 102 and 168 hours following

- subcutaneous administration and 30 minutes, 1, 8, 24, 48, 72, 96 and 168 hours following intravenous administration. The collected blood was dispensed into a Sarstedt microvette CB300Z with clot activator for serum separation, and left at room temperature for at least 20 minutes. The microvette was then centrifuged at 20°C at 10,000 rpm for 5 minutes. Serum was removed and stored frozen prior to analysis. The concentration of FabB-645Fv (G₄Sx2) or FabB-648Fv (G₄Sx2) in serum samples was assessed by ELISA. Briefly Nunc Maxisorb Immunomodule Plates were coated with hOX40-Fc in PBS and blocked with 1% BSA in PBS. Serum samples and standards were diluted in 1% BSA in PBS and applied to the plate for 1 hour. The plate was washed with PBS and the revealing antibody of goat anti-human kappa HRP conjugate applied in 1% BSA in PBS for 1 hour. The plate was washed and then developed with TMB substrate followed by stopping with 2.5M sulphuric acid. The absorbance at 630nm was measured and the concentrations determined from the standard curve.
- Both FabB-645Fv (G₄Sx2) and FabB-648Fv (G₄Sx2) have extended half-life in plasma, Figure 13. The half-lives for FabB-645Fv (G₄Sx2) are 71h sc and 62h iv and for FabB-648Fv (G₄Sx2) are 25h sc and 30h iv.

EXAMPLE 13

In vivo efficacy study of FabB-Fv

- A study to investigate if FabB-645Fv and FabB-648Fv are efficacious *in vivo* was undertaken. Briefly this involved steady state dosing in HuSCID mice and the read out was the prevention of T cell engraftment.
- CB17 SCID mice were dosed with a loading dose subcutaneously on day -2 of 2.475mg/kg FabB-645Fv or FabB-648Fv or FabB-PEG40k or PBS. On every subsequent day up to and including day 10 they were dosed with a maintenance dose subcutaneously of 0.75mg/kg FabB-645Fv or FabB-648Fv or FabB-PEG40k or PBS. Each dosing group consisted of 9-10 mice. On day -1 all the mice were treated with 0.87mg/mouse of rat anti-murine TM-β1 antibody to abrogate natural killer cell activity. On day 0 all the mice received an inter peritoneal injection of 8x10⁶ human peripheral blood mononuclear cells. On day 14 the mice are sacrificed and the blood, spleen and a peritoneal lavage were taken. The samples were analysed by FACS for CD4⁺ and CD8⁺ T cells. The data sets were analysed by one way Anova with Dunnett's post test comparison. All the test constructs FabB-645Fv, FabB-648Fv and FabB-PEG40k were equally efficacious in all the compartments tested, i.e. blood peritoneum and spleen. Figures 14A, B and C.

EXAMPLE 14

FabB-645Fv mutations to change the affinity of 645Fv for albumin

- Point mutations were introduced into selected residues in the CDRs of the heavy chain of the 645Fv portion of FabB-645dsFv (S3xG₄S) by mutagenic PCR. For example

I50A is a replacement of Ile 50 with Ala. The various mutations are given in Table 11 below. The affinity of the Fab-645Fv mutants for human albumin was assessed by BIAcore as described in the methods. All the mutations had either unchanged or reduced affinity for human albumin.

5 **Table 11**

Fv heavy mutation	Albumin	ka (1/Ms)	kd (1/s)	KD (nM)
I50A	HSA	3.12E+04	1.90E-03	60.9
T56A	HSA	4.65E+04	3.78E-04	8.12
T95A	HSA	2.81E+04	2.64E-03	94.0
V96A	HSA	2.81E+04	6.42E-04	22.9
P97A	HSA	4.60E+04	1.26E-02	275
G98A	HSA	4.73E+04	2.71E-04	5.73
Y99A	HSA	4.71E+04	4.79E-04	10.2
S100A	HSA	3.94E+04	1.44E-03	36.6
T100aA	HSA	3.60E+05	1.86E-02	51.6
Y100cA	HSA	1.23E+04	1.07E-03	87.0
I50A and T95A	HSA	2.12E+04	9.94E-03	468
I50A and G98A	HSA	1.79E+04	6.96E-03	389
I50A and Y99A	HSA			>3500
T56A and T95A	HSA	2.84E+04	8.57E-04	30.1
T56A and G98A	HSA	2.40E+04	3.68E-03	153
T56A and Y99A	HSA	2.24E+04	1.49E-02	664

EXAMPLE 15

1-5 Gly4Ser linker length between Fab and Fv

Construction of FabB-645Fv fusion plasmids for expression in mammalian cells

10 The FabB-645Fv's with either a SGGGGS, SGGGGS SGGGGS, SGGGGS SGGGGS SGGGGS, SGGGGS SGGGGS SGGGGS SGGGGS or SGGGGS SGGGGS SGGGGS SGGGGS SGGGGS linker between the C-termini of the Fab and the N-termini of the Fv were assembled by PCR then cloned into a mammalian expression vectors under the control of the HCMV-MIE promoter and SV40E polyA
 15 sequence. The relevant heavy and light chain plasmids were paired for expression in mammalian cells.

Mammalian expression of FabB-645Fv (1-5xG₄S)

HEK293 cells were transfected with the heavy and light chain plasmids using
 Invitrogen's 293fectin transfection reagent according to the manufacturer's
 20 instructions. Briefly, 24μg heavy chain plasmid + 24μg light chain plasmid was incubated with 120μl 293fectin + 4080μl Optimem media for 20mins at RT. The mixture was then added to 60x10⁶ HEK293 cells in 60mL suspension and incubated for 4 days with shaking at 37°C. All the constructs were equally well expressed.

Protein-G purification

The mammalian expression suspensions were clarified by centrifugation and the supernatants were concentrated to ~1.8mL using 10kDa molecular weight cut off centrifugation concentrators. The concentrated supernatants were centrifuged at 16000xg for 10 min to remove any precipitate and then 1.5mL was loaded onto 1ml HiTrap Protein-G columns (GE Healthcare) at 1ml/min. The columns were washed with 20mM phosphate, 40mM NaCl pH7.4 and bound material eluted with 0.1M glycine/HCl pH2.7. The elution peak (2mL) was collected and pH adjusted to ~pH5 with 250μL of 1M sodium acetate. The pH adjusted elutions were diafiltered into 20mM phosphate, 150mM NaCl pH7.1 using 10kDa molecular weight cut off centrifugation concentrators and concentrated to ~250μL. All the constructs had similar purification profiles and the final concentrations were 0.5-1.1mg/ml.

Affinity of FabB-645Fv (1-5xG₄S) for albumin

The affinities of the purified FabB-645Fv (1-5xG₄S) constructs for human and mouse albumin were determined as described in the Methods. The different linker lengths of the Fv of 1 to 5 xGly4Ser between the C-termini of the Fab and the N-termini of the Fv had no affect on the affinity of the 645Fv for either human or mouse albumin.

Table 12

	Albumin	KD (nM)	Albumin	KD (nM)
FabB-645Fv (1xG ₄ S)	Human	8.77	Mouse	2.18
FabB-645Fv (2xG ₄ S)	Human	6.72	Mouse	8.01
FabB-645Fv (3xG ₄ S)	Human	9.87	Mouse	8.92
FabB-645Fv (4xG ₄ S)	Human	7.90	Mouse	7.24
FabB-645Fv (5xG ₄ S)	Human	3.90	Mouse	6.09

SDS-PAGE analysis of purified FabB-645Fv (1-5xG₄S)

FabB-645Fv (1-5xG₄S) samples were prepared under non-reduced and reduced conditions and separated on a gel and stained as described in the methods. See Figure 15.

Size exclusion analysis of purified FabB-645Fv (1-5xG₄S)

FabB-645Fv (1-5xG₄S) samples were analysed for size on a Superdex200 10/300GL Tricorn column (GE Healthcare) developed with an isocratic gradient of 20mM phosphate 150mM NaCl pH7.4 at 1ml/min.

A linker length between the C-termini of the Fab and the N-termini of the Fv of either 1xG₄S or 2xG₄S reduces the amount of monomer FabB-645Fv whilst increasing the amount of dimer and higher multimers. The amount of monomer is least for the 1xG₄S linker length. A linker length between the C-termini of the Fab and the N-

termini of the Fv of either 3xG₄S, 4xG₄S or 5xG₄S increased the amount of monomer FabB-645Fv whilst decreasing the amount of dimer and higher multimers with the levels being similar for all three linker lengths. Figure 16.

5 **Table 13**

	Monomer	Dimer	High Multimers
FabB-645Fv (1xG ₄ S)	5 %	47 %	48 %
FabB-645Fv (2xG ₄ S)	27 %	38 %	36 %
FabB-645Fv (3xG ₄ S)	51 %	32 %	17 %
FabB-645Fv (4xG ₄ S)	55 %	30 %	15 %
FabB-645Fv (5xG ₄ S)	51 %	31 %	18 %

Thermofluor thermal stability analysis of purified FabB-645Fv (1-5xG₄S)

Samples (1µl of sample at ~1mg/ml, 8µl of PBS and 1µl of 30x stock of Sypro orange fluorescent dye) were run in quadruplicate in 384 well plates. The plate is heated
 10 from 20-99°C using a 7900HT fast real-time PCR system and the fluorescence (excitation at 490nm, emission at 530nm) measured. The results are shown in Table 14 and Figure 17.

Table 14

	Tm °C (Fab)	Tm °C (Fv)
FabB-645Fv (1xG ₄ S)	82.8 ± 0.6	67.4 ± 0.4
FabB-645Fv (2xG ₄ S)	83.4 ± 0.3	68.7 ± 0.3
FabB-645Fv (3xG ₄ S)	83.4 ± 0.3	69.5 ± 0.6
FabB-645Fv (4xG ₄ S)	83.8 ± 0.3	71.3 ± 1.0
FabB-645Fv (5xG ₄ S)	83.8 ± 0.4	72.0 ± 0.7

15

EXAMPLE 16

Disulphide stabilisation of the Fv in a Fab-Fv

FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), FabΔB-645dsFv (2xG₄S) and FabΔB-648dsFv (2xG₄S) fusion plasmids for expression in mammalian cells

20 Point mutations were introduced into the FabB-645Fv (2xG₄S) and FabB-648Fv (2xG₄S) DNA sequences at selected residues in the framework region of both the heavy chain and the light chain of the Fv by mutagenic PCR. The mutations introduced to create an interchain disulphide bond between the heavy and light chains of the Fv were heavy chain G44C and light chain G100C. As well as adding the
 25 cysteins to create the interchain disulphide bond in the Fv, the natural interchain disulphide between the heavy chain and light chain of the Fab was removed by mutagenic PCR by changing the cysteines to serines. Fvs that contain an interchain disulphide bond were termed dsFv, Fabs that lack an interchain disulphide bond were

termed Fab Δ . The DNA for all these constructs was then cloned into a mammalian expression vectors under the control of the HCMV-MIE promoter and SV40E polyA sequence. The relevant heavy and light chain plasmids were paired for expression in mammalian cells.

5

Mammalian expression of FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), Fab Δ B-645dsFv (2xG₄S) and Fab Δ B-648dsFv (2xG₄S)

HEK293 cells were transfected with the heavy and light chain plasmids using Invitrogen's 293fectin transfection reagent according to the manufacturer's instructions. Briefly, 24 μ g heavy chain plasmid + 24 μ g light chain plasmid was incubated with 120 μ l 293fectin + 4080 μ l Optimem media for 20mins at RT. The mixture was then added to 60x10⁶ HEK293 cells in 60mL suspension and incubated for 4 days with shaking at 37°C. All the constructs were equally well expressed.

Protein-G purification of FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), Fab Δ B-645dsFv (2xG₄S) and Fab Δ B-648dsFv (2xG₄S)

The mammalian expression suspensions were clarified by centrifugation and the supernatants were concentrated to ~1.8mL using 10kDa molecular weight cut off centrifugation concentrators. The concentrated supernatants were centrifuged at 16000xg for 10 min to remove any precipitate and then 1.5mL was loaded onto 1ml HiTrap Protein-G columns (GE Healthcare) at 1ml/min. The columns were washed with 20mM phosphate, 40mM NaCl pH7.4 and bound material eluted with 0.1M glycine/HCl pH2.7. The elution peak (2mL) was collected and pH adjusted to ~pH5 with 250 μ L of 1M sodium acetate. The pH adjusted elutions were diafiltered into 20mM phosphate, 150mM NaCl pH7.1 using 10kDa molecular weight cut off centrifugation concentrators and concentrated to ~250 μ L. All the constructs had similar purification profiles and the final concentrations were 0.5-0.8mg/ml.

Affinity of FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), Fab Δ B-645dsFv (2xG₄S) and Fab Δ B-648dsFv (2xG₄S) for albumin

The affinities of the purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S) Fab Δ B-645dsFv (2xG₄S), Fab Δ B-648dsFv (2xG₄S) constructs for human and mouse albumin were determined as described in the Methods. The disulphide stabilisation of the Fv had no affect or slightly increased the affinity of the Fv for both human or mouse albumin.

35

Table 15

	Albumin	KD (nM)	Albumin	KD (nM)
FabB-645Fv (2xG ₄ S)	Human	17.5	Mouse	24.7
FabB-645dsFv (2xG ₄ S)	Human	12.6	Mouse	14.0

FabΔB-645dsFv (2xG ₄ S)	Human	8.3	Mouse	12.2
FabB-648Fv (2xG ₄ S)	Human	9.4	Mouse	42.4
FabB-648dsFv (2xG ₄ S)	Human	3.1	Mouse	59.6
FabΔB-648dsFv (2xG ₄ S)	Human	8.3	Mouse	59.8

SDS-PAGE analysis of purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), FabΔB-645dsFv (2xG₄S) and FabΔB-648dsFv (2xG₄S)

Purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S) FabΔB-645dsFv (2xG₄S),
 5 FabΔB-648dsFv (2xG₄S) samples were prepared under non-reduced and reduced conditions and separated on a gel and stained as described in the methods. See Figure 18.

Size exclusion analysis of purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), FabΔB-645dsFv (2xG₄S) and FabΔB-648dsFv (2xG₄S)

Purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S) FabΔB-645dsFv (2xG₄S),
 10 FabΔB-648dsFv (2xG₄S) samples were analysed for size on a Superdex200 10/300GL Tricorn column (GE Healthcare) developed with an isocratic gradient of 20mM phosphate 150mM NaCl pH7.4 at 1ml/min.

15 The introduction of an interchain disulphide bond into the Fv of either a 645Fv or 648Fv increased the amount of monomer Fab-Fv species compared with the Fab-Fv in which the Fv did not have an inter chain disulphide. The removal of the natural interchain disulphide bond from the Fab part of a Fab-Fv had only a small effect on the amount of monomer species present. Figure 19.

20

Table 16

	Monomer	Dimer	High Multimers
FabB-645Fv (2xG ₄ S)	26 %	38 %	35 %
FabB-645dsFv (2xG ₄ S)	43 %	21 %	37 %
FabΔB-645dsFv (2xG ₄ S)	40 %	25 %	34 %
FabB-648dsFv (2xG ₄ S)	50 %	26 %	24 %
FabΔB-648dsFv (2xG ₄ S)	55 %	24 %	20 %

Thermofluor thermal stability analysis of purified FabB-645dsFv (2xG₄S), FabB-648dsFv (2xG₄S), FabΔB-645dsFv (2xG₄S) and FabΔB-648dsFv (2xG₄S)

25 Samples (1μl of sample at ~1mg/ml, 8μl of PBS and 1μl of 30x stock of Sypro orange fluorescent dye) were run in quadruplicate in 384 well plates. The plate is heated from 20-99°C using a 7900HT fast real-time PCR system and the fluorescence (excitation at 490nm, emission at 530nm) measured.

30 The introduction of an interchain disulphide bond into the Fv part of a Fab-Fv of either a 645Fv or 648Fv increased the thermal stability of the Fv compared with the

Fab-Fv in which the Fv did not have an inter chain disulphide. The removal of the natural interchain disulphide bond from the Fab part of a Fab-Fv decreased the thermal stability of the Fab part of the Fab-Fv

5 **Table 17**

	T_m °C (Fab)	T_m °C (Fv)
FabB-645Fv (2xG ₄ S)	81.9 ± 0.6	68.5 ± 0.5
FabB-645dsFv (2xG ₄ S)	83.6 ± 0.3	71.6 ± 0.3
FabΔB-645dsFv (2xG ₄ S)	79.5 ± 0.1	70.8 ± 0.6
FabB-648Fv (2xG ₄ S)	82.4 ± 0.2	70.6 ± 0.8
FabB-648dsFv (2xG ₄ S)	82.8 ± 0.3	75.0 ± 0.6
FabΔB-648dsFv (2xG ₄ S)	n.d.	73.6 ± 0.8

n.d. = not determined. The analysis software was unable to resolve this inflection point.

Biacore Method For FabD

10 Binding affinities and kinetic parameters for the interactions of Fab-dAb and Fab-didAb constructs were determined by surface plasmon resonance (SPR) conducted on a Biacore T100 using CM5 sensor chips and HBS-EP (10mM HEPES (pH7.4), 150mM NaCl, 3mM EDTA, 0.05% v/v surfactant P20) running buffer. Human Fab samples were captured to the sensor chip surface using either a human F(ab')₂-
 15 specific goat Fab (Jackson ImmunoResearch, 109-006-097) or an in-house generated anti human CH1 monoclonal antibody. Murine Fab samples were captured using a murine F(ab')₂-specific goat Fab (Jackson ImmunoResearch, 115-006-072). Covalent immobilisation of the capture antibody was achieved by standard amine coupling chemistry.

20 Each assay cycle consisted of firstly capturing the Fab-dAb or Fab-didAb construct using a 1 min injection, before an association phase consisting of a 3 min injection of antigen, after which dissociation was monitored for 5 min. After each cycle, the capture surface was regenerated with 2 x 1 min injections of 40mM HCl followed by
 25 30s of 5mM NaOH. The flow rates used were 10μl/min for capture, 30μl/min for association and dissociation phases, and 10μl/min for regeneration.

For kinetic assays, a titration of antigen (for human or mouse serum albumin typically 62.5nM-2μM, for IL-1β 1.25-40nM, for cell surface receptor D 20-1.25nM) was performed, a blank flow-cell was used for reference subtraction and buffer-blank
 30 injections were included to subtract instrument noise and drift.

Kinetic parameters were determined by simultaneous global-fitting of the resulting sensorgrams to a standard 1:1 binding model using Biacore T100 Evaluation software.

In order to test for simultaneous binding, 3 min injections of either separate 5 μ M HSA or 100nM IL-1 β , or a mixed solution of 5 μ M HSA and 100nM IL-1 β were injected over the captured Fab-dAb. Simultaneous binding of albumin and cell surface receptor D was assessed in the same manner using final concentrations of 2 μ M HSA or MSA and 20nM murine cell surface receptor D.

EXAMPLE 17

Mammalian expression of mFabC-mdidAbs and mFabD-mdidAbs

HEK293 cells were transfected with the heavy and light chain plasmids using Invitrogen's 293fectin transfection reagent according to the manufacturer's instructions. Briefly, 2 μ g heavy chain plasmid + 2 μ g light chain plasmid was incubated with 10 μ l 293fectin + 340 μ l Optimem media for 20mins at RT. The mixture was then added to 5x10⁶ HEK293 cells in suspension and incubated for 6 days with shaking at 37°C.

ELISA

The yields of mFab-mdidAb were measured using a sandwich ELISA. Briefly, the mFab-mdidAb was captured with an anti-CH1 antibody then revealed with an anti-kappa-HRP.

Table 18

	ELISA expression (ug/mL)
mFabD-mdidAb, -dAbL1(CK-G ₄ Sx2) & -dAbH1(CH1-G ₄ Sx2)	44
mFabD-mdidAb, -dAbL2(CK-G ₄ Sx2) & -dAbH2(CH1-G ₄ Sx2)	35
mFabC-mdidAb, -dAbL1(CK-G ₄ Sx2) & -dAbH1(CH1-G ₄ Sx2)	11
mFabC-mdidAb, -dAbL2(CK-G ₄ Sx2) & -dAbH2(CH1-G ₄ Sx2)	14

EXAMPLE 18

Further kinetic analysis was conducted to assess the interactions of serum albumin and human OX40 to the purified FabB-didAb, -dAbL1(CK-G₄Sx2) & -dAbH1(CH1-G₄Sx2) and FabB-didAb, -dAbL2(CK-G₄Sx2) & -dAbH2(CH1-G₄Sx2) fusions (Table 19). Both FabB-didAb, -dAbL1(CK-G₄Sx2) & -dAbH1(CH1-G₄Sx2) and FabB-didAb, -dAbL2(CK-G₄Sx2) & -dAbH2(CH1-G₄Sx2) retained the affinity for human OX40 of the original FabB (Table 20).

The potential for the FabB-didAb, -dAbL1(CK-G₄Sx2) & -dAbH1(CH1-G₄Sx2) and FabB-didAb, -dAbL2(CK-G₄Sx2) & -dAbH2(CH1-G₄Sx2) constructs to bind

simultaneously to both human or mouse serum albumin and human OX40 was assessed by capturing each Fab-didAb construct to the sensor chip surface, before performing either separate 3 min injections of 2 μ M albumin (human or mouse) or 50nM human OX40, or a mixed solution of both 2 μ M albumin and 50nM OX40.

- 5 HSA binding was seen for both Fab-didAb constructs. For each Fab-didAb construct the response seen for the combined albumin/OX40 solution was almost identical to the sum of the responses of the independent injections (summarised in table 21). This shows that the Fab-didAbs are capable of simultaneous binding to both OX40 and serum albumin. The original FabB bound only OX40, with no significant binding to
- 10 either human or mouse albumin.

Table 19

Construct	Albumin	k_a (x10 ⁴ M ⁻¹ s ⁻¹)	k_d (x10 ⁻⁵ s ⁻¹)	K_D (x10 ⁻⁹ M)
FabB-didAb, -dAbL1(CK-G4Sx2) & -dAbH1(CH1-G4Sx2	HSA	1.65	2.06	12.5
FabB-didAb, -dAbL2(CK-G4Sx2) & -dAbH2(CH1-G4Sx2	HSA	1.80	1.24	6.92
FabB-didAb, -dAbL1(CK-G4Sx2) & -dAbH1(CH1-G4Sx2	MSA	1.83	1.82	9.94
FabB-didAb, - dAbL2(CK-G4Sx2) & -dAbH2(CH1-G4Sx2	MSA	nd	nd	-

Affinity and kinetic parameters determined for HSA and MSA binding to Fab-didAb fusions.

15

Table 20

Construct	k_a (x10 ⁵ M ⁻¹ s ⁻¹)	k_d (x10 ⁻⁵ s ⁻¹)	K_D (x10 ⁻¹² M)
FabB	2.92	22.6	775
FabB-didAb, -dAbL1(CK-G4Sx2) & -dAbH1(CH1-G4Sx2	3.58	8.54	238
FabB-didAb, -dAbL2(CK-G4Sx2) & -dAbH2(CH1-G4Sx2	3.27	13.6	415

Affinity and kinetic parameters for hOX40-Fc binding to FabB and FabB-didAb fusions.

Table 21

Construct	Analyte	Binding (RU)	
FabB	HSA	2.5	
	MSA	-2.5	
	OX40	89.5	
	HSA + OX40	90.1	(92)
	MSA + OX40	86.5	(87)
FabB-didAb, -dAbL1(CK-G4Sx2) & -dAbH1(CH1-G4Sx2)	HSA	109.1	
	MSA	93.3	
	OX40	73.7	
	HSA + OX40	186.1	(182.8)
	MSA + OX40	170.3	(167)
FabB-didAb, -dAbL2(CK-G4Sx2) & -dAbH2(CH1-G4Sx2)	HSA	50.9	
	MSA	2.4	
	OX40	52.9	
	HSA + OX40	104.2	(103.8)
	MSA + OX40	54.9	(55.3)

The table above shows the binding response (RU) seen for each construct after separate injections of HSA or MSA or hOX40-Fc, or injection of premixed albumin and hOX40-Fc. In each case the final concentration was 2 μ M albumin HSA and 50nM hOX40-Fc. The sum of the individual albumin and hOX40-Fc responses is shown in parentheses.

EXAMPLE 19

Further kinetic analysis was conducted to assess the interactions of serum albumin and murine cell surface receptor D to mFabD-mdidAb, -mdAbL1(CK-G₄Sx2) & mdAbH1(CH1-G₄Sx2) and mFabD-mdidAb, -mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2) (Table 22). Both mFabD-mdidAbs showed relatively high affinity binding to HSA (K_D = 2.78nM and 8.97nM respectively). mFabD-mdidAb, -

mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2) also bound MSA with a similar affinity ($K_D = 22\text{nM}$), however no binding to MSA was seen for mFabD-mdidAb, -mdAbL1(CK-G₄Sx2) & mdAbH1(CH1-G₄Sx2). Both mFabD-mdidAbs retained the affinity for murine cell surface receptor D of the original mFabD (Table 23).

5

The potential for mFabD-mdidAb, -mdAbL1(CK-G₄Sx2) & mdAbH1(CH1-G₄Sx2) and mFabD-mdidAb, -mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2) to bind simultaneously to both human or mouse serum albumin and murine cell surface receptor D was assessed by capturing each mFab-mdidAb construct to the sensor chip surface, before performing either separate 3 min injections of 2 μM albumin (human or mouse) or 20nM murine cell surface receptor D, or a mixed solution of both 2 μM albumin and 20nM cell surface receptor D. Again HSA binding was seen for both mFab-mdidAb constructs whereas only mFabD-mdidAb, -mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2) bound MSA. For each mFab-mdidAb construct the response seen for the combined albumin/ cell surface receptor D solution was almost identical to the sum of the responses of the independent injections (summarised in table 24). This shows that the mFab-mdidAbs are capable of simultaneous binding to both cell surface receptor D and serum albumin. The original mFabD bound only cell surface receptor D, with no significant binding to either human or mouse albumin.

20

Table 22

Construct	Albumin	$k_a (\times 10^4 \text{M}^{-1} \text{s}^{-1})$	$k_d (\times 10^{-5} \text{s}^{-1})$	$K_D (\times 10^{-9} \text{M})$
mFabD-mdidAb, -mdAbL1(CK-G ₄ Sx2) & mdAbH1(CH1-G ₄ Sx2)	HSA	1.01	2.82	2.78
mFabD-mdidAb, -mdAbL2(CK-G ₄ Sx2) & mdAbH2(CH1-G ₄ Sx2)	HSA	1.19	10.69	8.97
mFabD-mdidAb, -mdAbL1(CK-G ₄ Sx2) & mdAbH1(CH1-G ₄ Sx2)	MSA	-	-	-
mFabD-mdidAb, -mdAbL2(CK-G ₄ Sx2) & mdAbH2(CH1-G ₄ Sx2)	MSA	1.03	22.73	22.06

Affinity and kinetic parameters determined for HSA and MSA binding to mFabD-mdidAb, -mdAbL1(CK-G₄Sx2) & mdAbH1(CH1-G₄Sx2) and mFabD-mdidAb, -mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2).

25

Table 23

Construct	k_a ($\times 10^5 \text{M}^{-1} \text{s}^{-1}$)	k_d ($\times 10^{-5} \text{s}^{-1}$)	K_D ($\times 10^{-12} \text{M}$)
mFabD	1.98	2.50	126
mFabD-mdidAb, -mdAbL1(CK-G ₄ Sx2) & mdAbH1(CH1-G ₄ Sx2)	2.01	4.67	233
mFabD-mdidAb, -mdAbL2(CK-G ₄ Sx2) & mdAbH2(CH1-G ₄ Sx2)	3.62	6.36	176

5 Affinity and kinetic parameters for murine cell surface receptor D-Fc binding to mFabD, mFabD-mdidAb, -mdAbL1(CK-G₄Sx2) & mdAbH1(CH1-G₄Sx2) and mFabD-mdidAb, -mdAbL2(CK-G₄Sx2) & mdAbH2(CH1-G₄Sx2).

Table 24

Construct	Analyte	Binding (RU)	
mFabD	receptor D	61.3	
	HSA	0.9	
	MSA	-1.1	
	receptor D + HSA	62.9	(62.2)
	receptor D + MSA	59.2	(60.2)
mFabD-mdidAb, -mdAbL1(CK-G ₄ Sx2) & mdAbH1(CH1-G ₄ Sx2)	receptor D	39.8	
	HSA	59.9	
	MSA	-0.6	
	receptor D + HSA	101.2	(99.7)
	receptor D + MSA	39.9	(39.2)
mFabD-mdidAb, -mdAbL2(CK-G ₄ Sx2) & mdAbH2(CH1-G ₄ Sx2)	receptor D	42.6	
	HSA	61.9	
	MSA	43.5	
	receptor D + HSA	105.3	(104.5)
	receptor D + MSA	86.3	(86.1)

The table above shows the binding response (RU) seen for each construct after separate injections of HSA or MSA or murine cell surface receptor D-Fc, or injection of premixed albumin and murine cell surface receptor D-Fc. In each case the final concentration was 2 μ M albumin HSA and 20nM murine cell surface receptor D-Fc.

- 5 The sum of the individual albumin and murine cell surface receptor D-Fc responses is shown in parentheses.


EXAMPLE 20

- Further analysis was conducted to assess the simultaneous interaction of mFabD-
10 mdidAb, -mdAbL1(CK-G₄Sx₂) & mdAbH1(CH1-G₄Sx₂) or mFabD-mdidAb, -
mdAbL2(CK-G₄Sx₂) & mdAbH2(CH1-G₄Sx₂) with serum albumin and murine cell
surface receptor D expressed on the cell surface. Both mFabD-mdidAbs were capable
of binding FITC labelled HSA and cell surface receptor X expressed on the cell
surface of activated murine T-cells simultaneously (figure 11). mFabD was capable
15 of binding cell surface receptor X expressed on the cell surface of activated murine T-
cells, data not shown, but did not bind FITC labelled HSA.

CLAIMS:

1. A multivalent antibody fusion protein which comprises an Fab or Fab' fragment, with a first specificity for an antigen of interest, and further comprises two single domain antibodies (dAb) with specificity for a second antigen of interest, wherein the two single domain antibodies are linked by a disulfide bond and wherein the two single domain antibodies are a complementary V_H/V_L pair and the V_H dAb is directly or indirectly connected to the C-terminus of the Fab or Fab' heavy chain and the V_L dAb is directly or indirectly connected to the C-terminus of the Fab or Fab' light chain.
2. The multivalent antibody fusion protein of claim 1, wherein the first antigen and second antigen are different entities.
3. The multivalent antibody fusion protein of claim 1 or 2, wherein the second antigen is albumin.
4. The multivalent antibody fusion protein of claim 3, wherein the albumin is human serum albumin.
5. The multivalent antibody fusion protein of claim 4, wherein the V_H single domain antibody comprises the sequence given in SEQ ID NO: 56 for CDRH1, the sequence given in SEQ ID NO: 57 for CDRH2 and the sequence given in SEQ ID NO: 58 for CDRH3 and the V_L single domain antibody comprises the sequence given in SEQ ID NO: 59 for CDRL1, the sequence given in in SEQ ID NO: 60 for CDRL2, and the sequence given in SEQ ID NO: 61 for CDRL3.
6. The multivalent antibody fusion protein of claim 4, wherein the V_H single domain antibody comprises the sequence given in SEQ ID NO: 62 for CDRH1, the sequence given in SEQ ID NO: 63 for CDRH2 and the sequence given in SEQ ID NO: 64 for CDRH3 and the V_L single domain antibody comprises the sequence given in SEQ ID NO: 65 for CDRL1, the sequence given in in SEQ ID NO: 66 for CDRL2, and the sequence given in SEQ ID NO: 67 for CDRL3.
7. A pharmaceutical composition comprising the multivalent antibody fusion protein as defined in any one of claims 1 to 6 in association with one or more pharmaceutically acceptable carriers, excipients or diluents.
8. The multivalent antibody fusion protein as defined in any one of claims 1 to 6, for use in therapy.



1/28

Figure 1




dAbL or dAbH are linked to the C-terminus of the constant region of either the light or heavy chain via a linker (—).

Variable region of light chain  or heavy chain  . Constant regions cKappa  and CH1  . Domain antibody fragments dAbL  and dAbH 

2/28

Figure 2A**Figure 2B**

dAbL and dAbH are linked to the C-terminus of the constant region of each chain so that a LC-dAbL or LC-dAbH fusion is paired with either the HC-dAbL or HC-dAbH.

Variable region of light chain  or heavy chain . Constant regions cKappa  and CH1 . Domain antibody fragments, dAbL  and dAbH. 

Disulfide bond —

3/28



Figure 3

Figure 4

5/28


Figure 4a

Figure 5**a) dAbH1**

EVQLLES GGGLVQPGGSLRLSCAVSGIDLSNYAINWVRQAPGKGLEWIGIIWA
SGTTFYATWAKGRFTISR DSTTVYLQMNSLRAEDTAVYYCARTVPGYSTAPY
FDLWGQGTLVTVSS (SEQ ID NO:52)

b) dAbL1

DIVMTQSPSSVSASVGDRVTITCQSSPSVWSNFLSWYQQKPGKAPKLLIYEAS
KLTSGVPSRFKSGSGTDFTLTISSLQPEDFATYYCGGGYSSISDTTFGGGTKV
EIK (SEQ ID NO:53)

c) dAbH2

EVQLVES GGGLVQPGGSLRLSCAVSGFSLSR YAMTWVRQAPGKGLEWIGTIT
TGGNTNYANWAKGRFTISK DSTTVYLQMNSLRAEDTAVYYCARGGYVSYA
DATELSLWGQGTLVTVSS (SEQ ID NO:54)

d) dAbL2

DIVMTQSPSTLSASVGDRVTITCQASQSIGSR LAWYQQKPGKAPKLLIYYAST
VASGVPSRFKSGSGTEFTLTISLQPD DFATYYCQSYDYSSSSSYAFGGGTKV
EIK (SEQ ID NO:55)

dAbH1

- e) CDRH1: GIDLSNYAIN (SEQ ID NO:56)
- f) CDRH2: IIWASGTTFYATWAKG (SEQ ID NO:57)
- g) CDRH3: TVPGYSTAPYFDL (SEQ ID NO:58)

dAbL1

- h) CDRL1: QSSPSVWSNFLS (SEQ ID NO:59)
- i) CDRL2: EASKLTS (SEQ ID NO:60)
- j) CDRL3: GGGYSSISDTT (SEQ ID NO:61)

dAbH2

- k) CDRH1: GFSLSR YAMT (SEQ ID NO:62)
- l) CDRH2: TITTTGGNTNYANWAKG (SEQ ID NO:63)
- m) CDRH3: GGYVSYADATELSL (SEQ ID NO:64)

dAbL2

- n) CDRL1: QASQSIGSR LA (SEQ ID NO:65)
- o) CDRL2: YASTVAS (SEQ ID NO:66)
- p) CDRL3: QSYDYSSSSSYA (SEQ ID NO:67)

Figure 6**FabB-dAbH1 (CH1-G₄Sx2)**

FAB-B HEAVY CHAIN VARIABLE DOMAIN +

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF
 PAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCGGG
GSGGGGSEVQLLESGGLVQPGGSLRLSCAVSGIDLSNYAINWVRQAPGKGL
 EWIGIIWASGTTFYATWAKGRFTISRDTTVYQLQMNSLRAEDTAVYYCARTV
 PGYSTAPYFDLWGQGTLLVTVSS (SEQ ID NO:68)

FabB-dAbH2 (CH1-G₄Sx2)

FAB-B HEAVY CHAIN VARIABLE DOMAIN +

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF
 PAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCGGG
GSGGGGSEVQLVESGGLVQPGGSLRLSCAVSGFSLRYAMTWVRQAPGKG
 LEWIGTITTGGNTNYANWAKGRFTISKDSTTVYQLQMNSLRAEDTAVYYCAR
 GGYVSYADATELSLWGQGTLLVTVSS (SEQ ID NO:69)

FabB-dAbL1 (CH1-G₄Sx2)

FAB-B HEAVY CHAIN VARIABLE DOMAIN +

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF
 PAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCGGG
GSGGGGSDIVMTQSPSSVSASVGDRVITTCQSSPSVWSNFLSWYQQKPGKAP
 KLLIYEASKLTSGVPSRFGSGSGTDFTLTISSLQPEDFATYYCGGGYSSISDTT
 FGGGTKVEIK (SEQ ID NO:70)

FabB-dAbL2 (CH1-G₄Sx2)

FAB-B HEAVY CHAIN VARIABLE DOMAIN +

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF
 PAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCGGG
GSGGGGSDIVMTQSPSTLSASVGDRVITTCQASQSIGSLAWYQQKPGKAPKL
 LIYYASTVASGVPSRFGSGSGTEFTLTISSLQPDDFATYYCQSYDYSSSSSYA
 FGGGTKVEIK (SEQ ID NO:71)

FabB-dAbL1 (CK-G₄Sx2)

FABB- LIGHT CHAIN VARIABLE DOMAIN +

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNS
 QESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRG
 ECGGGGSGGGGSDIVMTQSPSSVSASVGVDRVITTCQSSPSVWSNFLSWYQQK

8/28

Figure 6 continued

PGKAPKLLIYEASKLTSGVPSRFBKSGSGTDFTLTISLQPEDFATYYCGGGYS
SISDTTFGGGGTKVEIK (SEQ ID NO:72)

FabB-dAbL2 (CK-G₄Sx2)

FABB- LIGHT CHAIN VARIABLE DOMAIN +

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNS
QESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRG
ECGGGGSGGGGSDIVMTQSPSTLSASVGDRVTITCQASQSIGSRLAWYQQKP
GKAPKLLIYYASTVASGVPSRFBKSGSGGTEFTLTISLQPDDFATYYCQSYDYS
SSSSYAFGGGGTKVEIK (SEQ ID NO:73)

Figure 7:**Fab'A heavy chain**

MKKTAIAIAVALAGFATVAQAEVQLVESGGGLVQPGGSLRLSCAFSGFSLST
SGVGVGWVRQAPGKGLEWVAHIWWDGDESYNPSLKTQFTISKDTSKNTVYL
QMNSLRAEDTAVYYCARNRYDPPWFVDWGQGTLLTVSSASTKGPSVFPLAP
SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS
SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKTCDKTHTCPPCPA (SEQ
ID NO:74)

FabA light chain

MKKTAIAIAVALAGFATVAQADIQMTQSPSSLSASVGDRVTITCRASQDISN
YLSWYQQKPGKAPKLLIYYTSKLHSGVPSRFSGSGSGTDYTLTISSLQPEDFAT
YYCQQGKMLPWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLN
NFYPREAKVQWKVDNAVQSGNSQESVTEQDSKDYSLSTLTLSKADYEK
HKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:75)

Fab'A heavy chain (modified hinge linker)

MKKTAIAIAVALAGFATVAQAEVQLVESGGGLVQPGGSLRLSCAFSGFSLST
SGVGVGWVRQAPGKGLEWVAHIWWDGDESYNPSLKTQFTISKDTSKNTVYL
QMNSLRAEDTAVYYCARNRYDPPWFVDWGQGTLLTVSSASTKGPSVFPLAP
SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS
SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKTCDKTHTS (SEQ ID
NO:76)

FabA heavy chain

MKKTAIAIAVALAGFATVAQAEVQLVESGGGLVQPGGSLRLSCAFSGFSLST
SGVGVGWVRQAPGKGLEWVAHIWWDGDESYNPSLKTQFTISKDTSKNTVYL
QMNSLRAEDTAVYYCARNRYDPPWFVDWGQGTLLTVSSASTKGPSVFPLAP
SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS
SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKTC (SEQ ID NO:77)

Bold text in sequences of Figure 7 is signal sequence.

10/28

Figure 8a**mdAbH1**

QVQLKESGPGLVAPSQSL SITCTVSGIDLSNYAINWVRQPPGKGLEWIGIIWAS
GTTFYATWAKGRFSISRSTTVFLKMNSLQTDDTARYYCARTVPGYSTAPYF
DLWGQGTLVTVSS (SEQ ID NO:78)

CDRH1 GIDLSNYAIN (SEQ ID NO:56)
CDRH2 IIWASGTTFYATWAKG (SEQ ID NO:57)
CDRH3 TVPGYSTAPYFDL (SEQ ID NO:58)

mdAbL1

DIVMTQSPSSLTVTAGEKVTMSCQSSPSVWSNFLSWYQQKPGQPPKLLIYEAS
KLTSGVPDRFKGSGSGTDFTLTISSVQAEDLATYYCGGGYSSISDTTFGGGK
LEIKR (SEQ ID NO:79)

CDRL1 QSSPSVWSNFLS (SEQ ID NO:59)
CDRL2 EASKLTS (SEQ ID NO:60)
CDRL3 GGGYSSISDTT (SEQ ID NO:61)

mdAbH2

QVQLKESGPGLVAPSQSL SITCTVSGFSLSR YAMTWVRQPPGKGLEWIGTITT
GGNTNYANWAKGRFSISKDSTQVFLKMNSLQTDDTAMYYCARGGYVSYAD
ATELSLWGQGTLVTVSS (SEQ ID NO:80)

CDRH1 GFSLSR YAMT (SEQ ID NO:62)
CDRH2 TITTTGGNTNYANWAKG (SEQ ID NO:63)
CDRH3 GGYVSYADATELSL (SEQ ID NO:64)

mdAbL2

DIVMTQSPSSLSASLGERVTITCQASQSIGSR LAWYQQKPGKTPKLLIFYASTV
ASGVPSRFRKGSGSGTEFSLTISSLQPEDFATYYCQSYDYSSSSSYAFGGGKLEI
KR (SEQ ID NO:81)

CDRL1 QASQSIGSR LA (SEQ ID NO:65)
CDRL2 YASTVAS (SEQ ID NO:66)
CDRL3 QSYDYSSSSSYA (SEQ ID NO:67)

11/28

Figure 8b**mFabD-mdidAb, -dAbL1(CK-G₄Sx2) & -dAbH1(CH1-G₄Sx2)****Heavy chain variable domain +**

AKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHT
 FPAVLQSDLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDKKIVPRDCGGG
GSGGGGSQVQLKESGPGLVAPSQSLSITCTVSGIDLSNYAINWVRQPPGKGLE
 WIGIIWASGTTFYATWAKGRFSISRSTTVFLKMNSLQTDDTARYYCARTVP
 GYSTAPYFDLWGQGTLVTVSSDYKDDDDK (SEQ ID NO:82)

Light chain variable domain +

TDAAPTVSIFPPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQNGVLN
 SWTDQDSKDSTYSMSSTLTTLTKDEYERHNSYTCEATHKTSTSPIVKSFNRENEC
GGGGSGGGGSDIVMTQSPSSLTVTAGEKVTMSCQSSPSVWSNFLSWYQQKPG
 QPPKLLIYEASKLTSGVPSRFRKGSGSGTDFLTITSSVQAEDLATYYCGGGYSSI
 SDTTFGGGTKLEIKREQKLISEEDL (SEQ ID NO:83)

mFabD-mdidAb, -dAbL2(CK-G₄Sx2) & -dAbH2(CH1-G₄Sx2)**Heavy chain variable domain +**

AKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHT
 FPAVLQSDLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDKKIVPRDCGGG
GSGGGGSQVQLKESGPGLVAPSQSLSITCTVSGFSLSRYAMTWVRQPPGKGL
 EWIGTITTGGNTNYANWAKGRFSISKDSTQVFLKMNSLQTDDTAMYYCARG
 GYVSYADATELSLWGQGTLVTVSSDYKDDDDK (SEQ ID NO:84)

Light chain variable domain +

TDAAPTVSIFPPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQNGVLN
 SWTDQDSKDSTYSMSSTLTTLTKDEYERHNSYTCEATHKTSTSPIVKSFNRENEC
GGGGSGGGGSDIVMTQSPSSLSASLGERVTITCQASQSIGSRLAWYQQKPGKT
 PKLLIFYASTVASGVPSRFRKGSGSGTEFSLTISLQPEDFATYYCQSYDYSSSSS
 YAFGGGTKLEIKREQKLISEEDL (SEQ ID NO:85)

The C-termini of the dAbL are terminated with a FLAG tag and the C-termini of the dAbH are terminated with aMyc tag.

12/28

Figure 8c**mFabC-mdidAb, -dAbL1(CK-G₄Sx2) & -dAbH1(CH1-G₄Sx2)mFabC-mdAbH1****Heavy chain variable domain +**

AKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHT
 FPAVLQSDLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDKKIVPRDCGGG
GSGGGGSQVQLKESGPGLVAPSQSLSITCTVSGIDLSNYAINWVRQPPGKGLE
 WIGIIWASGTTFYATWAKGRFSISRSTTVFLKMNSLQTDDTARYYCARTVP
 GYSTAPYFDLWGQGTLLTVSSSDYKDDDDK (SEQ ID NO:86)

Light chain variable domain +

TDAAPTVSIFPPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQNGVLN
 SWTDQDSKDSTYSMSSTLTTLTKDEYERHNSYTCEATHKTSTSPIVKSFNRECE
GGGGSGGGGSDIVMTQSPSSLTVTAGEKVTMSCQSSPSVWSNFLSWYQQKPG
 QPPKLLIYEASKLTSGVPDRFKGSGSGTDFTLTISSVQAEDLATYYCGGGYSSI
 SDTTFGGGTKLEIKREQKLISEEDL (SEQ ID NO:87)



mFabC-mdidAb, -dAbL2(CK-G₄Sx2) & -dAbH2(CH1-G₄Sx2)**Heavy chain variable domain +**

AKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHT
 FPAVLQSDLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDKKIVPRDCGGG
GSGGGGSQVQLKESGPGLVAPSQSLSITCTVSGFSLSRyamTWVRQPPGKGL
 EWIGTITTGGNTNYANWAKGRFSISKDSTQVFLKMNSLQTDDTAMYYCARG
 GYVSYADATELSLWGQGTLLTVSSSDYKDDDDK (SEQ ID NO:88)

Light chain variable domain +

TDAAPTVSIFPPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQNGVLN
 SWTDQDSKDSTYSMSSTLTTLTKDEYERHNSYTCEATHKTSTSPIVKSFNRECE
GGGGSGGGGSDIVMTQSPSSLSASLGERVTITCQASQSIGSLAWYQQKPGKT
 PKLLIFYASTVASGVPSRFRKSGSGGTEFSLTISSLQPEDFATYYCQSYDYSSSSS
 YAFGGGTKLEIKREQKLISEEDL (SEQ ID NO:89)

13/28

Figure 9**Figure 10**

14/28

Figure 11






Figure 12




15/28

Figure 13

16/28


Figure 14A



P<0.01, *P<0.001

17/28


Figure 14B



P<0.01, *P<0.001


18/28

Figure 14C




P<0.01, *P<0.001

19/28

Figure 15SDS-PAGE analysis of FabB-645Fv (1-5xG₄S)


20/28

Figure 16
Size exclusion analysis of FabB-645Fv (1-5xG₄S)




21/28

Figure 17
Thermograms of FabB-645Fv (1-5xG₄S)



22/28

Figure 18

- 1 = FabB-645Fv (2xG₄S)
2 = FabB-648Fv (2xG₄S)
3 = FabΔB-645dsFv (2xG₄S)
4 = FabΔB-648dsFv (2xG₄S)
5 = FabB-645dsFv (2xG₄S)
6 = FabB-648dsFv (2xG₄S)

23/28

Figure 19

Size exclusion analysis of FabB-645Fv (2xG₄S), FabB-645dsFv (2xG₄S), FabΔB-645dsFv (2xG₄S)




Figure 20
646Fv

CDRH1	GIDLSNYAIN (SEQ ID NO: 222)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 223)
CDRH3	TVPGYSTAPYFDL (SEQ ID NO: 90)
CDRL1	QSSPSVWSDFLS (SEQ ID NO: 91)
CDRL2	GASTLAS (SEQ ID NO: 92)
CDRL3	GGGYSSISD TT (SEQ ID NO: 93)

647Fv

CDRH1	GFTLSNNYWMC (SEQ ID NO: 94)
CDRH2	CIYTGDDGTAYTSWAKG (SEQ ID NO: 95)
CDRH3	SGGSYYDYVFIL (SEQ ID NO: 96)
CDRL1	QASQSLGNRLA (SEQ ID NO: 97)
CDRL2	RASTLAS (SEQ ID NO: 98)
CDRL3	QCTYIGSKMGA (SEQ ID NO: 99)

648Fv is the same as a didAb of dAbH2, dAbL2

649Fv

CDRH1	GFSFSGNYWIC (SEQ ID NO: 100)
CDRH2	CIFTADGDTAYTSWAKG (SEQ ID NO: 101)
CDRH3	SGGSAFDYVFIL (SEQ ID NO: 102)
CDRL1	QASQSIG NRLG (SEQ ID NO: 103)
CDRL2	RASTLES (SEQ ID NO: 104)
CDRL3	QCTYIGKLMGA (SEQ ID NO: 105)

645HeavyI50AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 106)
CDRH2	AIWASGTTFYATWAKG (SEQ ID NO: 107)
CDRH3	TVPGYSTAPYFDL (SEQ ID NO: 108)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 109)
CDRL2	EASKLTS (SEQ ID NO: 110)
CDRL3	GGGYSSISDTT (SEQ ID NO: 111)

Figure 21**645HeavyT56AFv**

CDRH1	GIDLSNYAIN (SEQ ID NO: 112)
CDRH2	IIWASGATFYATWAKG (SEQ ID NO: 113)
CDRH3	TVPGYSTAPYFDL (SEQ ID NO: 114)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 115)
CDRL2	EASKLTS (SEQ ID NO: 116)
CDRL3	GGGYSSISDTT (SEQ ID NO: 117)

645HeavyT95AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 118)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 119)
CDRH3	AVPGYSTAPYFDL (SEQ ID NO: 120)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 121)
CDRL2	EASKLTS (SEQ ID NO: 122)
CDRL3	GGGYSSISDTT (SEQ ID NO: 123)

645HeavyV96AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 124)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 125)
CDRH3	TAPGYSTAPYFDL (SEQ ID NO: 126)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 127)
CDRL2	EASKLTS (SEQ ID NO: 128)
CDRL3	GGGYSSISDTT (SEQ ID NO: 129)

645HeavyP97AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 130)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 131)
CDRH3	TVAGYSTAPYFDL (SEQ ID NO: 132)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 133)
CDRL2	EASKLTS (SEQ ID NO: 134)
CDRL3	GGGYSSISDTT (SEQ ID NO: 135)

645HeavyG98AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 136)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 137)
CDRH3	TVPAYSTAPYFDL (SEQ ID NO: 138)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 139)
CDRL2	EASKLTS (SEQ ID NO: 140)
CDRL3	GGGYSSISDTT (SEQ ID NO: 141)

Figure 22**645HeavyY99AFv**

CDRH1	GIDLSNYAIN (SEQ ID NO: 142)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 143)
CDRH3	TVPGASTAPYFDL (SEQ ID NO: 144)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 145)
CDRL2	EASKLTS (SEQ ID NO: 146)
CDRL3	GGGYSSISDTT (SEQ ID NO: 147)

645HeavyS100AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 148)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 149)
CDRH3	TVPGYATAPYFDL (SEQ ID NO: 150)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 151)
CDRL2	EASKLTS (SEQ ID NO: 152)
CDRL3	GGGYSSISDTT (SEQ ID NO: 153)

645HeavyT100aAFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 154)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 155)
CDRH3	TVPGYSAAPYFDL (SEQ ID NO: 156)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 157)
CDRL2	EASKLTS (SEQ ID NO: 158)
CDRL3	GGGYSSISDTT (SEQ ID NO: 159)

645HeavyP100cAFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 160)
CDRH2	IIWASGTTFYATWAKG (SEQ ID NO: 161)
CDRH3	TVPGYSTAAYFDL (SEQ ID NO: 162)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 163)
CDRL2	EASKLTS (SEQ ID NO: 164)
CDRL3	GGGYSSISDTT (SEQ ID NO: 165)

645HeavyI50A+T95AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 166)
CDRH2	AIWASGTTFYATWAKG (SEQ ID NO: 167)
CDRH3	AVPGYSTAPYFDL (SEQ ID NO: 168)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 169)
CDRL2	EASKLTS (SEQ ID NO: 170)
CDRL3	GGGYSSISDTT (SEQ ID NO: 171)

Figure 23**645HeavyI50A+G98AFv**

CDRH1	GIDLSNYAIN (SEQ ID NO: 172)
CDRH2	AIWASGTTFYATWAKG (SEQ ID NO: 173)
CDRH3	TVPAYSTAPYFDL (SEQ ID NO: 174)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 175)
CDRL2	EASKLTS (SEQ ID NO: 176)
CDRL3	GGGYSSISDTT (SEQ ID NO: 177)

645HeavyI50A+Y99AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 178)
CDRH2	AIWASGTTFYATWAKG (SEQ ID NO: 179)
CDRH3	TVPGASTAPYFDL (SEQ ID NO: 180)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 181)
CDRL2	EASKLTS (SEQ ID NO: 182)
CDRL3	GGGYSSISDTT (SEQ ID NO: 183)

645HeavyT56A+T95AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 184)
CDRH2	IIWASGATFYATWAKG (SEQ ID NO: 185)
CDRH3	AVPGYSTAPYFDL (SEQ ID NO: 186)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 187)
CDRL2	EASKLTS (SEQ ID NO: 188)
CDRL3	GGGYSSISDTT (SEQ ID NO: 189)

645HeavyT56A+G98AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 190)
CDRH2	IIWASGATFYATWAKG (SEQ ID NO: 191)
CDRH3	TVPAYSTAPYFDL (SEQ ID NO: 192)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 193)
CDRL2	EASKLTS (SEQ ID NO: 194)
CDRL3	GGGYSSISDTT (SEQ ID NO: 195)

645HeavyT56A+Y99AFv

CDRH1	GIDLSNYAIN (SEQ ID NO: 196)
CDRH2	IIWASGATFYATWAKG (SEQ ID NO: 197)
CDRH3	TVPGASTAPYFDL (SEQ ID NO: 198)
CDRL1	QSSPSVWSNFLS (SEQ ID NO: 199)
CDRL2	EASKLTS (SEQ ID NO: 200)
CDRL3	GGGYSSISDTT (SEQ ID NO: 201)

Figure 24

645dsFv - is a disulphide stabilised Fv. The disulphide can be introduced into the Fv by the double mutation of heavy chain G44C and light chain G100C

Heavy chain Fv

EVQLLES GGGLVQP GGSRLRLSCAVSGIDLSNYAINWVRQAPGKC LEWIGIIWA
SGTTFYATWAKGRFTISR DSTTVYLQMNSLRAEDTAVYYCARTVPGYSTAPY
FDLWGQGTLVTVSS (SEQ ID NO: 202)

Light chain Fv

DIVMTQSPSSVSASVGDRVTITCQSSPSVWSNFLSWYQQKPGKAPKLLIYEAS
KLTSGVPSRFRKGSGSGTDFTLTISLQPEDFATYYCGGGYSSISDTTFGCGTKV
EIK (SEQ ID NO: 203)

648dsFv - is a disulphide stabilised Fv. The disulphide can be introduced into the Fv by the double mutation of heavy chain 44C and light chain 100C

Heavy chain Fv

EVQLVES GGGLVQP GGSRLRLSCAVSGFSLSR YAMTWVRQAPGKC LEWIGTIT
TGGNTNYANWAKGRFTISK DSTTVYLQMNSLRAEDTAVYYCARGGYVSYA
DATELSLWGQGTLVTVSS (SEQ ID NO: 204)

Light chain Fv

DIVMTQSPSTLSASVGDRVTITCQASQSIGSR LAWYQQKPGKAPKLLIYYAST
VASGVPSRFRKGSGSGTEFTLTISLQPD DFATYYCQSYDYSSSSSYAFGCGTKV
EIK (SEQ ID NO: 205)

Fv stabilisation or solubility mutations - mutations that increase the stability, solubility or expressability of the Fv. For example heavy chain A84D.

Heavy chain of 645Fv

EVQLLES GGGLVQP GGSRLRLSCAVSGIDLSNYAINWVRQAPGK GLEWIGIIWA
SGTTFYATWAKGRFTISR DSTTVYLQMNSLRDEDTAVYYCARTVPGYSTAPY
FDLWGQGTLVTVSS (SEQ ID NO: 206)

Light chain of 645Fv

DIVMTQSPSSVSASVGDRVTITCQSSPSVWSNFLSWYQQKPGKAPKLLIYEAS
KLTSGVPSRFRKGSGSGTDFTLTISLQPEDFATYYCGGGYSSISDTTFGGGTKV
EIK (SEQ ID NO: 207)