

(21) **2 670 776**

(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2007/11/29
(87) Date publication PCT/PCT Publication Date: 2008/06/05
(85) Entrée phase nationale/National Entry: 2009/05/25
(86) N° demande PCT/PCT Application No.: US 2007/085933
(87) N° publication PCT/PCT Publication No.: 2008/067465
(30) Priorité/Priority: 2006/11/29 (US60/867,764)

(51) *Cl.Int./Int.Cl. C07D 401/04* (2006.01),
A61K 31/513 (2006.01), *A61P 9/00* (2006.01)


(71) **Demandeur/Applicant:**
TAKEDA PHARMACEUTICAL COMPANY LIMITED, JP

(72) **Inventeurs/Inventors:**
KELLY, RON C., US;
KOZTECKI, LIEN H., US

(74) **Agent:** SMART & BIGGAR

(54) Titre : POLYMORPHES DE SEL DE SUCCINATE DE 2-[6-(3-AMINO-PIPERIDIN-1-YL)-3-METHYL-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YLMETHY]-4-FLUOR-BENZONITRILE ET LEURS PROCEDES D'UTILISATION

(54) Title: POLYMORPHS OF SUCCINATE SALT OF 2-[6-(3-AMINO-PIPERIDIN-1-YL)-3-METHYL-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YLMETHY]-4-FLUOR-BENZONITRILE AND METHODS OF USE THEREFOR



(57) Abrégé/Abstract:

Compositions comprising the succinate salt of 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile (referred to herein as Compound I) which has the formula: (I) wherein the Compound I is present in one or more polymorphic forms. Also provided are novel methods for the preparation of the polymorphs of Compound I, and kits and articles of manufacture of the compositions, and methods of using the compositions to treat various diseases.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2008 (05.06.2008)

PCT

(10) International Publication Number
WO 2008/067465 A1

(51) International Patent Classification:

C07D 401/04 (2006.01) **A61P 9/00** (2006.01)
A61K 31/513 (2006.01)

(21) International Application Number:

PCT/US2007/085933

(22) International Filing Date:

29 November 2007 (29.11.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/867,764 29 November 2006 (29.11.2006) US

(71) **Applicant** (for all designated States except US): **TAKEDA PHARMACEUTICAL COMPANY LIMITED** [JP/JP];
1-1, Doshomachi 4-chome Chuo-ku, Osaka-shi, Osaka
541-0045 (JP).

(72) Inventors; and

(75) **Inventors/Applicants** (for US only): **KELLY, Ron C.** [US/US]; 1413 Via Vista, San Mateo, CA 94404 (US).
KOZTECKI, Lien H. [US/US]; 1528 1/2 Main Street,
Lafayette, Indiana 47901 (US).

(74) **Agents:** **WEITZ, David J.** et al.; Takeda San Diego, Inc.,
10410 Science Center Drive, San Diego, CA 92121 (US).


(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) **Title:** POLYMORPHS OF SUCCINATE SALT OF 2-[6-(3-AMINO-PIPERIDIN-1-YL)-3-METHYL-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YLMETHYL]-4-FLUOR-BENZONITRILE AND METHODS OF USE THEREFOR



(57) **Abstract:** Compositions comprising the succinate salt of 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile (referred to herein as Compound I) which has the formula: (I) wherein the Compound I is present in one or more polymorphic forms. Also provided are novel methods for the preparation of the polymorphs of Compound I, and kits and articles of manufacture of the compositions, and methods of using the compositions to treat various diseases.

WO 2008/067465 A1


POLYMORPHS OF SUCCINATE SALT OF 2-[6-(3-AMINO-PIPERIDIN-1-YL)-3-METHYL-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YLMETHYL]-4-FLUOR-BENZONITRILE AND METHODS OF USE THEREFOR

FIELD OF THE INVENTION

[0001] The present invention relates generally to polymorphic forms of the succinic acid salt of 2-[6-(3-amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile, (referred to herein as “Compound I”) and methods for their preparation. The present invention also relates to pharmaceutical compositions, kits and articles of manufacture comprising polymorphs of Compound I, and methods of their use.

DESCRIPTION OF RELATED ART

[0002] Compound I, which has the formula:



is a DPP-IV inhibitor that is described in U.S. Patent Application Ser. No. 11/080,992 filed March 15, 2005 (*see* Compound 34). Its dosing, administration and biological activities are described in U.S. Patent Application Ser. No. 11/531,671 filed September 13, 2006. U.S. Patent Application Ser. No. 11/080,992 and Ser. No. 11/531,671 are incorporated herein by reference in their entirety.

[0003] Dipeptidyl peptidase IV (IUBMB Enzyme Nomenclature EC.3.4.14.5) (referred herein as “DPP-IV”) is a type II membrane protein and a non-classical serine aminodipeptidase that removes Xaa-Pro dipeptides from the amino terminus (N-terminus) of polypeptides and proteins. DPP-IV is constitutively expressed on epithelial and endothelial cells of a variety of different tissues (*e.g.*, intestine, liver, lung, kidney and placenta), and is also found in body fluids. DPP-IV is also expressed on circulating T-lymphocytes and has been shown to be synonymous with the cell-surface antigen, CD-26. DPP-IV has been implicated in a number of human disease states, including, but are not limit to, diabetes, particularly type II diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG),

metabolic acidosis, ketosis, appetite regulation and obesity; autoimmune diseases such as inflammatory bowel disease, multiple sclerosis and rheumatoid arthritis; AIDS; and cancers.

[0004] DPP-IV inhibitors are believed to be useful agents for the prevention, delay of progression, and/or treatment of conditions mediated by DPP-IV.

SUMMARY OF THE INVENTION

[0005] The present invention provides novel polymorphic forms of Compound I and methods of preparing these polymorphic forms, as well as compositions comprising one or more of the novel polymorphs. For ease of reference, the different polymorphs described herein are referred to consistently throughout the application as Forms A through G, and Amorphous Form.

Form A

[0006] In one embodiment, the present invention relates to a polymorph of Compound I, referred to herein as Form A. Based on its physical properties, Form A is a crystalline form.

[0007] Form A may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form A):

- (a) may be crystallized from a solvent selected from the group consisting of (i) acetone, (ii) acetonitrile, (iii) dichloromethane, (iv) 1, 4-dioxane, (v) dimethylformamide, (vi) ethanol, (vii) ethylacetate, (viii) diethylether, (ix) hexane; (x) methanol, (xi) isopropanol, (xii) tetrahydrofuran, (xiii) toluene, (xiv) trifluoroethanol, (xv) water, (xvi) acetonitrile:water (85:15), (xvii) ethanol:water (95:5), (xviii) isopropanol:water (88:12), and (xix) tetrahydrofuran:water (9:1);
- (b) may be precipitated by adding a miscible antisolvent to Compound I dissolved in a solvent, wherein the solvent/antisolvent system is selected from the group consisting of (i) dimethylformamide/acetonitrile, (ii) dimethylformamide/toluene, (iii) dimethylformamide/ethylacetate, (iv) dimethylformamide/isopropanol, (v) methanol/acetonitrile, (v) methanol/dichloromethane, (vi) trifluoroethanol/isopropanol, (vii)

- trifluoroethanol/acetonitrile, (viii) trifluoroethanol/ethylacetate, (ix) water/acetonitrile, and (x) water/tetrahydrofuran;
- (c) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418 \text{ \AA}$) comprising diffraction peaks at about 11.31, 11.91, and 22.32 degree 2-theta ($^{\circ}2\theta$), more particularly, comprising five or more diffraction peaks selected from the group consisting of diffraction peaks at about 4.80, 11.31, 11.91, 12.86, 14.54, 15.81, 16.83, 17.59, 18.11, 19.26, 19.52, 20.32, 21.04, 21.80, 22.32, 23.42, 23.83, 24.78, 25.28, 25.84, 26.14, 26.63, 27.62, 27.84, 28.14, 29.39, 29.87, 30.27, 31.60, 31.88, 32.44, 33.86, 34.51, 35.87, 36.36, 37.31, 38.64, and 39.49 $^{\circ}2\theta$, even more particularly, comprising diffraction peaks at about 11.31, 11.91, 19.26, 21.04, and 22.32 $^{\circ}2\theta$;
 - (d) has a Fourier transform infra-red absorption spectrum comprising absorption bands at about 3141, 2953, 2934, 2266, 1699, 1657, 1450, and 1206 wave number (cm^{-1}), and more particularly absorption bands at about 3141, 2953, 2934, 2266, 2225, 1699, 1657, 1450, 1206, 886, 760, 685, 594, and 516 cm^{-1} ;
 - (e) has a Raman spectrum comprising Raman peaks at about 2954, 2935, 2225, 1698, 1659, and 1607 cm^{-1} , and more particularly comprising Raman peaks at about 3068, 2954, 2935, 2225, 1698, 1659, 1607, 1586, 1223, 1180, 901, 780, 751, 669, and 516 cm^{-1} ;
 - (f) a differential scanning calorimetry spectrum having an endotherm centered at about 195 $^{\circ}\text{C}$; and
 - (g) insignificant weight loss when heated from 25 $^{\circ}\text{C}$ to 165 $^{\circ}\text{C}$.

Form B

[0008] Form B may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form B):

- (a) may be crystallized from a solvent selected from the group consisting of (i) isopropanol, (ii) ethanol:water (95:5), (iii) isopropanol:water (88:12), (iii) tetrahydrofuran:water (9:1), and (iv) water;
- (b) may be precipitated by adding dioxane to Compound I dissolved in water;

- (c) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418\text{\AA}$) which comprises characteristic peaks at about 12.51, 18.83, and 24.46 $^{\circ}2\theta$;
- (d) has a differential scanning calorimetry thermogram comprising a broad endotherm at about 100 $^{\circ}\text{C}$, and two small endotherms at about 138 $^{\circ}\text{C}$ and about 163 $^{\circ}\text{C}$, and another endotherm at about 193 $^{\circ}\text{C}$;
- (e) has a significant weight loss ($>2.0\%$) when heated from about 25 $^{\circ}\text{C}$ to 175 $^{\circ}\text{C}$; and
- (f) converts to Form A when dissolved in water and recovered by evaporation of the solvent.

Form C

[0009] Form C may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form C):

- (a) may be crystallized from an isopropanol solution of Compound I; and
- (b) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418\text{ \AA}$) which comprises diffraction peaks at about 5.44 ± 0.2 and 6.07 ± 0.2 $^{\circ}2\theta$.

Form D

[0010] Form D may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form D):

- (a) may be precipitated by adding a miscible antisolvent to a solution of Compound I; e.g., adding acetonitrile to a water solution of Compound I;
- (b) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418\text{ \AA}$) comprising diffraction peaks at about 12.19, 22.88, and 24.33 $^{\circ}2\theta$; alternatively, a diffraction pattern comprising one diffraction peak at about 24.33 $^{\circ}2\theta$ and four other peaks selected from the group consisting of diffraction peaks at about 12.19, 14.04, 16.71, 17.75, 18.86, 19.96, 22.08, 22.88, 23.27, 25.02, 25.49, 26.03, and 27.99 $^{\circ}2\theta$, and more particularly, a diffraction pattern comprising one diffraction peak at about 24.33 $^{\circ}2\theta$ and four other peaks selected from the group consisting of diffraction peaks at about 12.19, 16.71, 22.08, 22.88, and 23.27 $^{\circ}2\theta$;

- (c) has a differential scanning calorimetry thermogram having a broad, noisy endotherm centered at about 88 °C, and two others endotherms at about 107 °C and 192 °C; and
- (d) has a weight loss of >20% when heated from 20 °C to 85 °C due to water loss.

Form E

[0011] Form E may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form E):

- (a) may be precipitated by adding a miscible antisolvent to a solution of Compound I; e.g., by adding acetonitrile to a water solution of Compound I;
- (b) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418 \text{ \AA}$) comprising two diffraction peaks at about 21.27 and 17.15 °2 θ , and three other diffraction peaks selected from the group consisting of peaks at about 11.90, 12.66, 13.10, 13.59, 13.94, 17.54, 22.03, 22.61, 24.06, 24.70, 26.31, 27.34, and 31.10 °2 θ , and more particularly a diffraction pattern comprising diffraction peaks at about 13.10, 13.94, 17.15, and 21.27 °2 θ ;
- (c) has a differential scanning calorimetry thermogram having two small endotherms at about 59 °C and about 75 °C, a forked endotherm having peaks at about 107 °C, 110 °C and 114 °C, and another endotherm at about 192 °C; and
- (d) has a weight loss of >2% when heated from 25 °C to 85 °C.

Form F

[0012] Form F may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form F):

- (a) may be crystallized from a tetrahydrofuran and water solution of Compound I; and
- (b) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418 \text{ \AA}$) which comprises diffraction peaks at about 12.39, 20.63, 26.03, and 30.05 °2 θ .

Form G

[0013] Form G may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Form G):

- (a) may be formed by crystallization from a tetrahydrofuran and water solution of Compound I; and
- (b) has an X-ray powder diffraction pattern ($\text{CuK}\alpha$, $\lambda=1.5418 \text{ \AA}$) comprising three or more diffraction peaks selected from a group consisting of peaks at about 13.22, 14.23, 18.62, 19.77, 24.36, 25.06, and 30.71 $^{\circ}2\theta$.

Amorphous Form

[0014] The amorphous form ("Amorphous Form") may be characterized as having one or more of the following physical characteristics (it being noted that a composition need not necessarily exhibit all of these characteristics in order to indicate the presence of Amorphous Form):

- (a) may be formed by lyophilizing a water solution of Compound I;
- (b) has an XRPD spectrum characterized by a large halo with no discernable peak;
- (c) has a differential scanning calorimetry spectrum showing the glass transitional temperature at 82 $^{\circ}\text{C}$, an exotherm at about 138 $^{\circ}\text{C}$ and an endotherm at about 199 $^{\circ}\text{C}$.

[0015] Methods by which the above referenced analyses were performed in order to identify these physical characteristics are described in the Example 3.

Compositions Comprising Compound I

[0016] The present invention relates to compositions comprising Compound I, wherein Compound I is present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. It is noted that other crystalline and amorphous forms of Compound I may also be present in the composition.

[0017] In one variation, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%,

97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. The composition may optionally be a pharmaceutical composition. The pharmaceutical composition may optionally further include one or more pharmaceutical carriers.

Kits and Articles of Manufacture Comprising Compound I

[0018] The invention also provided are kits and other articles of manufacture comprising a composition that comprises Compound I, wherein Compound I is present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. In one variation, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. The composition in the kits and articles of manufacture may optionally be a pharmaceutical composition. The pharmaceutical composition may optionally further include one or more pharmaceutical carriers.

[0019] In regard to each of the above embodiments including a pharmaceutical composition, the pharmaceutical composition may be formulated in any manner where at least a portion of Compound I is present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. Optionally, a portion of Compound I is present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form for a period of time subsequent to administration of the pharmaceutical formulation to a human.

Methods of Making Form A through Form G and Amorphous Form

[0020] Various methods are also provided for making Form A through Form G, and Amorphous Form. Various methods are also provided for manufacturing pharmaceutical compositions, kits and other articles of manufacture comprising one or more of Form A through Form G, and Amorphous Form.

Methods of Using Form A through Form G and Amorphous Form

[0021] Methods of using a pharmaceutical composition, kit and other article of manufacture comprising one or more of Form A through Form G, and Amorphous Form to treat various diseases mediated by DPP-IV are also provided.

[0022] In one embodiment, the present invention relates to a method of inhibiting dipeptidyl peptidases comprising administering a composition where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I.

[0023] In another embodiment, the present invention relates to a method of inhibiting dipeptidyl peptidases in a subject (*e.g.*, human body) with Compound I by administering Compound I where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form, when the compound is administered. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I.

[0024] In another embodiment, the present invention relates to a method of inhibiting dipeptidyl peptidases in a subject (*e.g.*, human body) with Compound I by administering Compound I where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, and Amorphous Form for a period of time after the compound has been administered to a subject. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I.

[0025] In still another embodiment, the present invention provides a method of treating a disease state for which dipeptidyl peptidases possesses activity that contributes to the pathology and/or symptomology of the disease state, comprising administering to a subject (*e.g.*, human body) a composition where greater than 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is

present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form when administered. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I.

[0026] In still another embodiment, the present invention provides a method of treating a disease state for which dipeptidyl peptidases possesses activity that contributes to the pathology and/or symptomology of the disease state, comprising causing a composition to be present in a subject (*e.g.*, human body) where greater than 0.1%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form, for a period of time after the composition has been administered to a subject. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I.

[0027] In another embodiment, a method is provided for preventing, delaying the of progression, and/or treating conditions mediated by DPP-IV, in particular diabetes and more particularly, type 2 diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation and obesity; autoimmune diseases such as inflammatory bowel disease, multiple sclerosis and rheumatoid arthritis; AIDS, cancers, and others.

[0028] In each instance where it is stated that Compound I may be present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form, it is intended for the invention to encompass compositions where only one form is present, where two forms are present (all combinations) and where three, four or more forms are present (all combinations).

BRIEF DESCRIPTION OF THE FIGURES

[0029] Figure 1 is a characteristic x-ray powder diffraction (XRPD) spectrum of Form A.

[0030] Figure 2 is a characteristic thermogravimetric analysis (TGA) thermogram of Form A.

[0031] Figure 3 is a characteristic differential scanning calorimetric (DSC) thermogram of Form A.

[0032] Figure 4A is a characteristic Fourier transform infrared absorption (FT-IR) spectrum (4000-400 wavenumbers (cm⁻¹)) of Form A.

[0033] Figure 4B is an expansion (4000-2500 cm⁻¹) of the FT-IR spectrum of Figure 4A.

[0034] Figure 4C is an expansion (2500-1500 cm⁻¹) of the FT-IR spectrum of Figure 4A.

[0035] Figure 4D is an expansion (1600-400 cm⁻¹) of the FT-IR spectrum of Figure 4A.

[0036] Figure 5A is a characteristic FT-Raman spectrum (3600-0 cm⁻¹) of Form A.

[0037] Figure 5B is an expansion (3600-1500 cm⁻¹) of the FT-Raman spectrum of Figure 5A.

[0038] Figure 5C is an expansion (1500-500 cm⁻¹) of the FT-Raman spectrum of Figure 5A.

[0039] Figure 5D is an expansion (500-100 cm⁻¹) of the FT-Raman spectrum of Figure 5A.

[0040] Figure 6 is a solution proton nuclear magnetic resonance (¹H NMR) spectrum of Form A.

[0041] Figure 7 is a solid state Carbon-13 nuclear magnetic resonance (¹³C NMR) spectrum of Form A.

[0042] Figure 8 is a characteristic XRPD spectrum of a mixture of Form A and Form B (Form A+B).

[0043] Figure 9 is a characteristic TGA thermogram of Form A+B.

[0044] Figure 10 is the TGA thermogram of a TG-IR analysis of Form A+B.

[0045] Figure 11 is the linked (~5 minutes delay) IR spectrum of the TG-IR analysis of Form A+B. The bottom spectrum is the IR analysis of the volatiles evolved at ~5 minutes from the start of the TG analysis (Figure 10). The top spectrum is a reference spectrum of water.

[0046] Figure 12 is a characteristic DSC thermogram of Form A+B.

[0047] Figure 13 is a moisture sorption and desorption isotherm of Form A+B.


[0048] Figure 14 is a characteristic XRPD spectrum of Form C.

[0049] Figure 15 is a characteristic XRPD spectrum of Form D.

- [0050] Figure 16 is a characteristic TGA thermogram of Form D.
- [0051] Figure 17 is a TGA thermogram of a TG-IR analysis of Form D.
- [0052] Figure 18 is the linked IR spectrum (~5 minutes delay) of the TG-IR analysis of Form D. The top spectrum is the IR spectrum of the volatiles evolved at ~5 minutes from the start of the TG analysis (Figure 17). The bottom spectrum is a reference spectrum of water.
- [0053] Figure 19 is a characteristic DSC thermogram of Form D.
- [0054] Figure 20 is a characteristic XRPD spectrum of Form E.
- [0055] Figure 21 is a characteristic TGA thermogram of Form E.
- [0056] Figure 22 is a characteristic DSC thermogram of Form E.
- [0057] Figure 23 is a moisture sorption and desorption isotherm of Form E.
- [0058] Figure 24 is a characteristic XRPD spectrum of a mixture of Form A and Form F (Form A+F).
- [0059] Figure 25 is a characteristic XRPD spectrum of a mixture of Form A, Form E and Form G (Form A+E+G).
- [0060] Figure 26 is a characteristic XRPD spectrum of Amorphous Form.
- [0061] Figure 27 is a characteristic TGA thermogram of Amorphous Form.
- [0062] Figure 28 is a characteristic DSC thermogram of Amorphous Form.
- [0063] Figure 29 is a DSC thermogram for assessing the glass transitional temperature of Amorphous Form.
- [0064] Figure 30 is a moisture sorption and desorption isotherm of Amorphous Form.
- [0065] Figure 31 shows XRPD spectra of Amorphous Form before (top) and after (middle) a moisture sorption/desorption analysis, and an XRPD pattern of Form A (bottom).

DETAILED DESCRIPTION OF THE INVENTION

- [0066] The present invention is directed to novel polymorphs, and compositions comprising the succinic acid salt of 2-[6-(3-amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2*H*-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile (Compound 1) of the formula



where at least a portion of Compound I is present in the composition in a form selected from the group consisting of crystalline forms, Forms A to G, and an amorphous form, Amorphous Form.

[0067] Also provided are kits and other articles of manufacture with compositions comprising Compound I where at least a portion of Compound I is present in the composition in a form selected from the group consisting of crystalline Forms A to G, and Amorphous Form.

[0068] Various methods are also provided including methods of making each of the disclosed forms; methods for manufacturing pharmaceutical compositions comprising Compound I where at least a portion of Compound I is present in the composition in a form selected from the group consisting of crystalline Forms A to G, and Amorphous Form; and methods of using compositions comprising Compound I where at least a portion of Compound I is present in the composition in a form selected from the group consisting of crystalline Forms A to G, and Amorphous Form.

[0069] As one will appreciate, depending on how a composition comprising a given compound is produced and then, once produced, how the composition is stored and manipulated, will influence the crystalline content of the composition. Accordingly, it is possible for a composition to comprise no crystalline content or may comprise higher concentrations of crystalline content.

[0070] It is further noted that a compound may be present in a given composition in one or more different polymorphic forms, as well as optionally also being present as an amorphous material. This may be the result of (a) physically mixing two or more different polymorphic forms; (b) having two or more different polymorphic forms be generated from crystallization conditions; (c) having all or a portion of a given polymorphic form convert into another polymorphic form; (d) having all or a portion of a compound in an amorphous state convert into two or more polymorphic forms; as well as for a host of other reasons.

[0071] As can be seen, depending on how a composition comprising a compound is prepared, the percentage, by weight, of that compound in a given polymorphic form can vary from 0% to 100%. According to the present invention, compositions are provided where greater than 0.1%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% or more of Compound I (by weight) is present in the composition in a form selected from the group consisting of Forms A to G, and Amorphous Form.

Definitions

[0072] “Crystalline”, as the term is used herein, refers to a material, which may be hydrated and/or solvated, and has sufficient ordering of the chemical moiety to exhibit a discernable diffraction pattern by XRPD or other diffraction techniques. Often, a crystalline material that is obtained by direct crystallization of a compound dissolved in a solution or by interconversion of crystals obtained under different crystallization conditions, will have crystals that contain the solvent used in the crystallization, termed a crystalline solvate. Also, the specific solvent system and physical embodiment in which the crystallization is performed, collectively termed crystallization conditions, may result in the crystalline material having physical and chemical properties that are unique to the crystallization conditions, generally due to the orientation of the chemical moieties of the compound with respect to each other within the crystal and/or the predominance of a specific polymorphic form of the compound in the crystalline material.

[0073] Depending upon the polymorphic form(s) of the compound that are present in a composition, various amounts of the compound in an amorphous solid state may also be present, either as a side product of the initial crystallization, and/or a product of degradation of the crystals comprising the crystalline material. Thus, crystalline, as the term is used herein, contemplates that the composition may include amorphous content; the presence of the crystalline material among the amorphous material being detectably among other methods by the composition having a discernable diffraction pattern.

[0074] The amorphous content of a crystalline material may be increased by grinding or pulverizing the material, which is evidenced by broadening of diffraction and other spectral lines relative to the crystalline material prior to grinding. Sufficient grinding and/or pulverizing may broaden the lines relative to the crystalline material prior to grinding to the extent that the XRPD or other crystal specific spectrum may become undiscernable, making the material substantially amorphous or quasi-amorphous.

[0075] Continued grinding would be expected to increase the amorphous content and further broaden the XRPD pattern with the limit of the XRPD pattern being so broadened that it can no longer be discerned above noise. When the XRPD pattern is broadened to the limit of being indiscernible, the material may be considered no longer a crystalline material, but instead be wholly amorphous. For material having increased amorphous content and wholly amorphous material, no peaks should be observed that would indicate grinding produces another form.

[0076] “Amorphous”, as the term is used herein, refers to a composition comprising a compound that contains too little crystalline content of the compound to yield a discernable pattern by XRPD or other diffraction techniques. Glassy materials are a type of amorphous material. Glassy materials do not have a true crystal lattice, and technically resembling very viscous non-crystalline liquids. Rather than being true solids, glasses may better be described as quasi-solid amorphous material.

[0077] “Broad” or “broadened”, as the term is used herein to describe spectral lines, including XRPD, NMR and IR spectroscopy, and Raman spectroscopy lines, is a relative term that relates to the line width of a baseline spectrum. The baseline spectrum is often that of an unmanipulated crystalline form of a specific compound as obtained directly from a given set of physical and chemical conditions, including solvent composition and properties such as temperature and pressure. For example, broadened can be used to describe the spectral lines of a XRPD spectrum of ground or pulverized material comprising a crystalline compound relative to the material prior to grinding. In materials where the constituent molecules, ions or atoms, as solvated or hydrated, are not tumbling rapidly, line broadening is indicative of increased randomness in the orientation of the chemical moieties of the compound, thus indicative of an increased amorphous content. When comparisons are made between crystalline materials obtained via different crystallization conditions, broader spectral lines indicate that the material producing the relatively broader spectral lines has a higher level of amorphous material.

[0078] “About” as the term is used herein, refers to an estimate that the actual value falls within $\pm 5\%$ of the value cited.

[0079] “Forked” as the term is used herein to describe DSC endotherms and exotherms, refers to overlapping endotherms or exotherms having distinguishable peak positions.

Preparation and Characterization of the Polymorphs

A. Preparation of Compound I

[0080] Various methods may be used to synthesize Compound I. A representative method for synthesizing Compound I is provided in Example 1. It is noted, however, that other synthetic routes may also be used to synthesize Compound I including those disclosed in U.S. Patent Application Serial No. 11/080,992, filed March 15, 2005, the disclosure of which is hereby incorporated by reference in its entirety.

B. Preparation of the Polymorphs of Compound I

[0081] General methods for precipitating and crystallizing a compound may be applied to prepare the various polymorphs described herein. These general methods are known to those skilled in the art of synthetic organic chemistry and pharmaceutical formulation, and are described, for example, by J. March, *“Advanced Organic Chemistry: Reactions, Mechanisms and Structure,”* 4th Ed. (New York: Wiley-Interscience, 1992).

[0082] In general, a given polymorph of a compound may be obtained by direct crystallization of the compound or by crystallization of the compound followed by inter-conversion from another polymorphic form or from an amorphous form. Depending on the method by which a compound is crystallized, the resulting composition may contain different amounts of the compound in crystalline form as opposed to as an amorphous material. Also, the resulting composition may contain differing mixtures of different polymorphic forms of the compound.

[0083] Compositions comprising a higher percentage of crystalline content (*e.g.*, forming crystals having fewer lattice defects and proportionately less glassy material) are generally prepared when using conditions that favor slower crystal formation, including slow solvent evaporation and those affecting kinetics. Crystallization conditions may be appropriately adjusted to obtain higher quality crystalline material as necessary. Thus, for example, if poor crystals are formed under an initial set of crystallization conditions, the solvent temperature may be reduced and ambient pressure above the solution may be increased relative to the initial set of crystallization conditions in order to slow down crystallization.

[0084] Precipitation of a compound from solution, often affected by rapid evaporation of solvent, is known to favor the compound forming an amorphous solid as opposed to crystals. A compound in an amorphous state may be produced by rapidly evaporating

solvent from a solvated compound, or by grinding, pulverizing or otherwise physically pressurizing or abrading the compound while in a crystalline state.

[0085] Compound I as prepared by the method described in Example 1 may be used as the starting material for preparation of all other polymorphic forms. The methods for testing the solubility of Compound I are described in Example 2, and the solubilities of Compound I in various solvents are summarized in Table A of Example 2. Poor solubility was observed in most solvents with the exceptions of dimethyl formamide (DMF), methanol, trifluoroethanol (TFE), water, and water:solvent mixtures.

[0086] Methods by which the various polymorphic forms may be prepared are described in Subsection A, Example 3. Specific methods by which Forms A-G, and Amorphous Form may be prepared are described in Tables B and C of Example 3. Methods useful for forming Form A, Form A+B, Form C, Form D, Form E, Form A+F, Form A+E+G, and Amorphous Form are described in Examples 3-10, respectively.

C. Polymorphs of Compound I

[0087] Seven crystalline forms and one amorphous solid were identified by conducting a polymorph screen (Example 3).

[0088] Described herein are Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form of Compound I. Where possible, the results of each test for each different polymorph are provided. Forms A, C, D and E were prepared as pure forms. Forms B, F, and G were prepared as mixtures with Form A.

[0089] Various tests were performed in order to physically characterize the polymorphs of Compound I including X-ray powder diffraction (XRPD), differential scanning calorimetry (DSC), thermogravimetric analysis (TGA), hot stage microscopy, Fourier transform infrared spectroscopy (FT-IR), Fourier transform Raman spectrometry, linked thermogravimetric-infrared spectroscopy (TG-IR), solution proton nuclear magnetic resonance (^1H -NMR), solid state ^{13}C carbon nuclear magnetic resonance (^{13}C -NMR), and moisture sorption and desorption analysis (M S/Des). Detailed experimental conditions for each of the analytical techniques are described in Example 3, Subsection B. The characterization of Forms A-G and Amorphous Form are described in Examples 11-18. Methods for testing the stabilities of the various forms of Compound I, and the conditions under which the polymeric forms interconvert are describe in Examples 19-25.

1. Form A

[0090] Form A may be prepared by crystallization from the various solvents and under the various crystallization conditions used during the polymorph screen (e.g., fast and slow evaporation, cooling of saturated solutions, slurries, and solvent/antisolvent additions).

Tables B and C of Example 3 summarize the procedures by which Form A was prepared. For example, Form A was obtained by room temperature slurry of an excess amount of Compound I in acetone, acetonitrile, dichloromethane, 1,4-dioxane, diethyl ether, hexane, methanol, isopropanol, water, ethylacetate, tetrahydrofuran, toluene, or other like solvents on a rotating wheel for approximately 5 or 7 days. The solids were collected by vacuum filtration, and air dried in the hood. Also, Form A was precipitated from a methanol solution of Compound I by slow evaporation (SE).

[0091] Form A was characterized by XRPD, TGA, hot stage microscopy, IR, Raman spectroscopy, solution ^1H -NMR, and solid state ^{13}C -NMR.

[0092] Figure 1 shows a characteristic XRPD spectrum ($\text{CuK}\alpha$, $\lambda=1.5418\text{\AA}$) of Form A. The XRPD pattern confirmed that Form A was crystalline. Major X-Ray diffraction lines expressed in $^{\circ}2\theta$ and their relative intensities are summarized in Table 1.

Table 1. Characteristic XRPD Peaks ($\text{CuK}\alpha$) of Form A

Peak No.	2θ ($^{\circ}$)	d-spacing	Intensity	I/I _o
1	4.80	18.40	91	3
2	11.31	7.82	2382	80
3	11.91	7.43	1416	48
4	12.86	6.88	266	9
5	14.54	6.09	1009	34
6	15.81	5.60	1457	49
7	16.83	5.26	522	18
8	17.59	5.04	299	10
9	18.11	4.89	91	3
10	19.26	4.60	816	27
11	19.52	4.54	642	22
12	20.32	4.37	178	6
13	21.04	4.22	996	34
14	21.80	4.07	174	6
15	22.32	3.98	2969	100
16	23.42	3.80	127	4
17	23.83	3.73	1025	35
18	24.78	3.59	181	6
19	25.28	3.52	684	23
20	25.84	3.45	120	4
21	26.14	3.41	352	12

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
22	26.63	3.34	477	16
23	27.62	3.23	342	12
24	27.84	3.20	303	10
25	28.14	3.17	274	9
26	29.39	3.04	132	4
27	29.87	2.99	255	9
28	30.27	2.95	183	6
29	31.60	2.83	163	5
30	31.88	2.80	166	6
31	32.44	2.76	106	4
32	33.86	2.65	123	4
33	34.51	2.60	141	5
34	35.87	2.50	305	10
35	36.36	2.47	133	4
36	37.31	2.41	146	5
37	38.64	2.33	129	4
38	39.49	2.28	90	3

[0093] The above set of XRPD peak positions or a subset thereof can be used to identify Form A. One subset comprises peaks at about 11.31, 11.91, 12.86, 14.54, 15.81, 16.83, 17.59, 19.26, 19.52, 21.04, 22.32, 26.63, and 29.87 °2 θ . Another subset comprises peaks at about 11.31, 11.91, 19.26, 21.04, and 22.32 °2 θ ; the peaks of this subset show no shoulder peaks or peak split greater than 0.2 °2 θ . Another subset comprises peaks at about 11.31, 11.91 and 22.32 °2 θ .

[0094] Figure 2 is a TGA thermogram of Form A. TGA analysis showed that Form A exhibited insignificant weight loss when heated from 25 °C to 165 °C; this result is indicative that Form A was an anhydrous solid.

[0095] Figure 3 shows a characteristic DSC thermogram of Form A. DSC analysis showed a single endothermic event occurred at approximately 195 °C (peak maximum). This endothermic event was confirmed by hot stage microscopy which showed the melting of Form A, which onset around 177 °C and the melting point estimated to be at approximately 184 °C.

[0096] Figure 4 (A-D) shows a characteristic FT-IR spectrum of Form A. The major bands expressed in reciprocal wavelengths (wavenumber in cm⁻¹) are positioned at about 3815, 3736, 3675, 3460, 3402, 3141, 3098, 3068, 3049, 2953, 2934, 2854, 2760, 2625, 2536, 2481, 2266, 2225, 2176, 1990, 1890, 1699, 1657, 1638, 1626, 1609, 1586, 1553, 1517, 1492, 1478, 1450, 1419, 1409, 1380, 1351, 1327, 1289, 1271, 1236, 1206, 1180, 1158, 1115, 1087, 1085, 1064, 1037, 1027, 971, 960, 951, 926, 902, 886, 870, 831, 820,

806, 780, 760, 740, 728, 701, 685, 668, 637, 608, 594, 567, 558, and 516 cm^{-1} (values rounded to the nearest whole number). This unique set of IR absorption bands or a subset thereof can be used to identify Form A. One such subset comprises absorption bands at about 3141, 3098, 3068, 3049, 2953, 2934, 2854, 2266, 2225, 1699, 1657, 1609, 1586, 1553, 1517, 1492, 1478, 1450, 1380, 1351, 1327, 1236, 1206, 1115, 1063, 902, 886, 870, 820, 780, 760, 685, 608, 594, and 516 cm^{-1} . Another subset comprises absorption bands at about 3141, 2953, 2934, 2854, 2266, 2225, 1699, 1657, 1450, 1206, 886, 760, 685, 594, and 516 cm^{-1} . Yet another subset comprises absorption bands at about 3141, 2953, 2934, 2266, 1699, 1657, 1450, and 1206 cm^{-1} .

[0097] Figure 5 (A-D) shows a characteristic Raman spectrum of Form A. The major Raman bands expressed in reciprocal wavelengths (wavenumber in cm^{-1}) are positioned at about 3100, 3068, 3049, 2977, 2954, 2935, 2875, 2855, 2787, 2263, 2225, 2174, 1698, 1659, 1626, 1607, 1586, 1492, 1478, 1451, 1439, 1409, 1400, 1382, 1351, 1328, 1290, 1281, 1271, 1237, 1223, 1213, 1180, 1155, 1134, 1115, 1084, 1063, 1035, 971, 952, 926901, 868, 805, 780, 759, 740, 727, 701, 686, 669, 609, 594, 566, 558, 516, 487, 479, 433, 418, 409, 294, 274, 241, 218, 191 and 138 cm^{-1} (values rounded to the nearest whole number). This unique set of Raman bands or a subset thereof may be used to identify Form A. One such subset comprises Raman bands at about 2954, 2935, 2225, 1698, 1659, and 1607 cm^{-1} . Another subset comprises Raman bands at about 3068, 2954, 2935, 2225, 1698, 1659, 1607, 1586, 1223, 1180, 901, 780, 759, 669, and 516 cm^{-1} . Yet another subset comprises Raman bands at about 3100, 3068, 2225, 1698, 1659, 1607, 1586, 1351, 1237, 1223, 1180, 1155, 1134, 1115, 1063, 952, 926, 901, 868, 805, 780, 759, 740, 669, 609, and 516 cm^{-1} .

[0098] Form A was further characterized by solution ^1H NMR and solid-state ^{13}C carbon NMR. The spectra are reported in Figures 6 and 7, respectively. Chemical assignments were not performed; however, the spectra are consistent with the known chemical structure of Compound I.

[0099] A relative humidity stressing study (Example 19) showed that after storing under approximately 58% and 97% relative humidity for 29 days, Form A did not exhibit form changes as evidenced by XRPD. Additional relative humidity stressing studies, studies showed that Form B and Amorphous Form also converted to Form A.

[0100] Based on the above available characterization data, Form A was anhydrous and stable at ambient conditions.

2. Form B

[0101] Form B was prepared as a mixture with Form A (hereinafter refers to as “Form A+B”) (Example 5). Form A+B was characterized by XRPD, TGA, DSC, hot stage microscopy, TG-IR, and moisture sorption/desorption analysis.

[0102] The XRPD pattern of Form A+B displayed in Figure 8 confirmed that Form A+B was crystalline. The characteristic diffraction lines expressed in 2θ ($\text{CuK}\alpha$, $\lambda=1.5418\text{\AA}$) and their relative intensity are summarized in Table 2 below.

Table 2. Characteristic XRPD Peaks ($\text{CuK}\alpha$) of Form A+B

Peak No.	2θ (°)	d-spacing	Intensity	I/I ₀
1	4.69	18.82	146	10
2	9.53	9.28	71	5
3	11.32	7.81	1121	74
4	11.87	7.45	605	40
5	12.51	7.07	483	32
6	13.29	6.66	91	6
7	14.05	6.30	130	9
8	14.53	6.09	371	25
9	15.83	5.59	1062	70
10	16.92	5.24	646	43
11	17.60	5.03	156	10
12	18.83	4.71	279	18
13	19.19	4.62	907	60
14	19.59	4.53	478	32
15	20.33	4.36	92	6
16	21.10	4.21	646	43
17	21.79	4.08	259	17
18	22.44	3.96	1513	100
19	23.80	3.74	542	36
20	24.46	3.64	160	11
21	25.27	3.52	505	33
22	26.12	3.41	349	23
23	26.74	3.33	520	34
24	27.93	3.19	262	17
25	28.80	3.10	177	12
26	29.41	3.03	182	12
27	30.02	2.97	166	11
28	31.34	2.85	139	9
29	31.96	2.80	194	13
30	33.69	2.66	86	6
31	34.37	2.61	148	10
32	35.90	2.50	126	8

[0103] This unique set of XRPD peaks or a subset thereof can be used to identify Form

A+B of Compound I. A subset set of peaks that is characteristic of Form B was obtained by subtracting the XRPD peak positions of Form A from those of Form A+B. This subset which may be used to identify the presence of Form B comprises peaks positioned at about 12.51, 18.83 and 24.46 °2 θ .

[0104] DSC analysis (Figure 12) showed a broad endotherm at about 100 °C, followed by small endotherms at 138 °C and 163 °C and a major endotherm at 193 °C. These events suggested a slow desolvation/dehydration of Form A+B, and then melting of the desolvated Form B (163 °C) and Form A (193 °C).

[0105] These melting events were confirmed by hot stage microscopy which also showed portion of the particles melted at approximately 179 °C and the rest of the particles melted at 188 °C (micrographs not included).

[0106] The TGA analysis (Figure 9) indicated a 2.0% weight loss when Form A+B was heated from 25 °C to 98 °C and an additional 0.7% weight loss when heating continued from 98 °C to 175 °C.

[0107] TG-IR (Figures 10 and 11) analysis revealed that the weight loss shown by TGA (Figure 10) was due to the volatilization of water (Figure 11). A 2.4% weight loss was consistent with approximately 0.5 to 1.0 moles of water. Since Form A was showed to be anhydrous, these results suggest that Form B was a hydrated solid form.

[0108] Moisture sorption/desorption analysis (Figure 13 and Example 23) showed a 0.7% weight loss on equilibration at 5% relative humidity. A reversible weight change of 1.2% weight was observed on sorption/desorption. XRPD analysis of the material remaining after the moisture sorption analysis was consistent with Form A.

[0109] Form B may be converted to Form A under dehydrating conditions. When Form A+B was stressed at 58% and 88% relative humidity for 23 days, the solids remained were Form A (Example 19). When Form A+B was exposed to ambient conditions for two days and in a vacuum oven at ambient temperature for one day, Form A+B converted to Form A (Example 20). When Form A+B was slurried in water and the water subsequently removed by evaporation, Form A+B converted to Form A (Example 21).

[0110] Based on the characterization data collected, Form B was a crystalline, hydrated solid.

3. Form C

[0111] Form C was prepared by precipitation from an IPA solution of Compound I by

slow cooling (Example 5). XRPD confirmed that Form C was crystalline.

[0112] The characteristic XRPD spectrum (CuK α , $\lambda=1.5418\text{\AA}$) of Form C is shown in Figure 14. The major diffraction peaks expressed in $^{\circ}2\theta$ and their relative intensities are summarized in Table 3 below.

Table 3. Characteristic XRPD Peaks (CuK α) of Form C.

Peak No.	2θ ($^{\circ}$)	d-spacing	Intensity	I/I _o
1	4.39	20.13	143	16
2	5.44	16.24	885	100
3	6.07	14.56	318	36
4	11.63	7.60	136	15
5	12.87	6.87	540	61
6	15.72	5.63	210	24
7	17.12	5.17	419	47
8	19.18	4.62	281	32
9	20.13	4.41	133	15
10	21.43	4.14	161	18
11	22.61	3.93	247	28
12	23.66	3.76	343	39
13	25.42	3.50	135	15

[0113] This unique set of XRPD peak positions or a subset thereof can be used to identify Form C. Particularly useful for identification of Form C are peaks at about 5.44 and 6.07 $^{\circ}2\theta$ because these peaks did not exhibit a shoulder or peak split greater than 0.2 $^{\circ}2\theta$.

4. Form D

[0114] Form D was precipitated by adding acetonitrile (a miscible antisolvent) to a water solution of Compound I (Example 6). The precipitates were characterized by solution ^1H NMR, XRPD, TGA, DSC, TG-IR, and hot stage microscopy.

[0115] Solution ^1H NMR data (spectrum not included) showed that Form D had the same chemical structure as Compound I.

[0116] The characteristic XRPD spectrum (CuK α) of Form D is shown in Figure 15. The XRPD diffraction pattern confirmed that Form D was crystalline. The major diffraction peaks expressed in $^{\circ}2\theta$ and their relative intensities are summarized in Table 4.

Table 4. Characteristic XRPD Peaks (CuK α) of Form D

Peak No.	2θ ($^{\circ}$)	d-spacing	Intensity	I/I _o
1	4.06	21.73	110	8
2	12.19	7.25	787	59
3	13.45	6.58	123	9

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
4	13.61	6.50	120	9
5	14.04	6.30	351	26
6	15.91	5.57	123	9
7	16.71	5.30	1344	100
8	17.75	4.99	516	38
9	18.59	4.77	83	6
10	18.86	4.70	210	16
11	19.45	4.56	124	9
12	19.63	4.52	227	17
13	19.96	4.44	541	40
14	20.46	4.34	75	6
15	21.89	4.06	334	25
16	22.08	4.02	687	51
17	22.88	3.88	190	14
18	23.27	3.82	528	39
19	23.51	3.78	188	14
20	24.33	3.65	433	32
21	24.76	3.59	291	22
22	25.02	3.56	289	22
23	25.49	3.49	217	16
24	26.03	3.42	200	15
25	26.59	3.35	76	6
26	26.89	3.31	86	6
27	27.15	3.28	261	19
28	27.99	3.18	276	21
29	28.23	3.16	167	12
30	29.17	3.06	145	11
31	29.33	3.04	87	6
32	29.76	3.00	201	15
33	31.75	2.82	152	11
34	32.11	2.79	147	11
35	33.02	2.71	67	5
36	33.81	2.65	106	8
37	36.81	2.44	68	5
38	37.10	2.42	67	5
39	38.61	2.33	87	6

[0117] This above set of XRPD peak positions or a subset thereof can be used to identify Form D of Compound I. Particularly useful for identifying Form D are the subsets of peaks that comprise a peak located at about 24.33°2 θ and four other peaks selected from the group consisting of peaks at about 12.19, 14.04, 16.71, 17.75, 18.86, 19.96, 22.08, 22.88, 23.27, 25.02, 25.49, 26.03, and 27.99°2 θ . More particularly useful are the subsets of peaks that comprise a peak located at about 24.33°2 θ and four other peaks selected from the group consisting of peaks at about 12.19, 16.71, 22.08, 22.88, and

23.27 °2 θ . Other subsets of peaks that are useful for identifying Form D are those comprising the peaks at about 12.19, 22.88 and 24.33 °2 θ , ± 0.2 °2 θ .

[0118] Form D was formed when the Amorphous Form was vapor stressed (Example 22). TGA analysis (Figure 16) showed that Form D exhibited an approximately 8% weight loss on equilibration, followed by a 21.9% weight loss from 25 °C to 85 °C. The approximately 30% total weight loss was consistent with approximately 11 moles of water.

[0119] Further, TG-IR analysis (Figures 17 and 18) confirms that the weight loss from 20 °C to 100 °C was due to volatilization of water.

[0120] DSC analysis of Form D (Figure 19) showed multiple, broad endotherms with an apparent peak maximum at 88 °C, followed by endotherms at approximately 107 °C and 192 °C. The data suggested that the solid desolvated/dehydrated, melted and possibly converted to Form A with the subsequent melting of Form A upon continued heating.

[0121] Hot stage analysis showed the final melting of the solid at 192 °C (micrograph not included) which further substantiated that Form D converted to Form A which has a melting temperature of about 192 °C.

[0122] In summary, the characterization data indicated that Form D was a crystalline, hydrated solid.

5. Form E

[0123] Form E was precipitated by the evaporation of a H₂O/ACN solvent/antisolvent solution of Compound I (Example 7). The precipitates were characterized by solution ¹H NMR, XRPD, TGA, DSC, TG-IR, hot stage microscopy, and moisture sorption/desorption analysis.

[0124] Solution ¹H NMR data (spectrum not included) showed that Form E possessed the same chemical structure as Compound I.

[0125] The characteristic XRPD spectrum (CuK α , $\lambda=1.5418\text{\AA}$) of Form E is shown in Figure 20. The XRPD diffraction pattern confirmed that Form E was crystalline. The major diffraction peaks expressed in °2 θ , and their relative intensities are summarized in Table 5 below.

Table 5. Characteristic XRPD Peaks (CuK α) of Form E

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
1	11.90	7.43	292	16
2	12.17	7.27	121	7

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
3	12.66	6.99	187	10
4	13.10	6.75	727	39
5	13.59	6.51	266	14
6	13.94	6.35	676	37
7	16.78	5.28	306	17
8	17.15	5.17	1026	55
9	17.54	5.05	450	24
10	18.18	4.88	196	11
11	19.02	4.66	274	15
12	20.03	4.43	239	13
13	21.27	4.17	1849	100
14	22.03	4.03	659	36
15	22.61	3.93	744	40
16	23.02	3.86	121	7
17	24.06	3.70	476	26
18	24.70	3.60	270	15
19	25.60	3.48	145	8
20	25.95	3.43	285	15
21	26.31	3.38	219	12
22	26.70	3.34	191	10
23	27.34	3.26	336	18
24	28.08	3.18	247	13
25	28.41	3.14	128	7
26	28.61	3.12	241	13
27	29.00	3.08	242	13
28	29.70	3.01	114	6
29	31.10	2.87	314	17
30	33.76	2.65	118	6
31	34.48	2.60	95	5
32	35.04	2.56	127	7
33	35.58	2.52	122	7
34	37.65	2.39	117	6
35	38.10	2.36	93	5
36	38.42	2.34	93	5

[0126] The unique set of XRPD peak positions above or a subset thereof can be used to identify Form E. Such subsets of peaks may comprise any five peaks selecting from the group consisting of peaks at about 11.90, 12.66, 13.10, 13.59, 13.94, 17.15, 17.54, 21.27, 22.03, 22.61, 24.06, 24.70, 26.31, 27.34, and 31.10 °2 θ . Other subsets of peaks that may be used to identify Form E include two peaks at about 21.27 and 17.15 °2 θ , and three other peaks selected from the group consisting of peaks at about 11.90, 12.66, 13.10, 13.59, 13.94, 17.54, 22.03, 22.61, 24.06, 24.70, 26.31, 27.34, and 31.10 °2 θ . Another subset of peaks that may be used for identifying Form E comprises peaks located at about 13.10,

13.94, 17.15, 21.27, 26.31 and 27.34 °2θ.

[0127] TGA analysis (Figure 21) showed that Form E exhibited a 2.6% weight loss upon heating from 25 °C to 85 °C.

[0128] DSC analysis (Figure 22) showed two small endotherms at 59 °C and 75 °C (dehydration), a forked endotherm with peaks at 107 °C, 110 °C, and 114 °C, followed by a single endotherm at 192 °C.

[0129] Hot stage analysis of Form E (micrographs not included) showed changes in birefringence at 59 °C and 81 °C, which was consistent with dehydration of the solid. Hot stage microscopy further showed that the approximate end of melt occurred at about 185 °C.

[0130] Moisture sorption/desorption data (Figure 23 and Example 24) showed that Form E under dehydrating conditions convert to Form A. A 1.3% weight loss was observed on equilibration to 5% relative humidity. The sample gained approximately 5% weight on sorption from a 5-95% relative humidity change. A loss of approximately 6% weight was observed on desorption to 5% relative humidity. XRPD analysis on the post moisture sorption/desorption sample showed a mixture of Form A and Form E.

[0131] In summary, the characterization data indicated that Form E was a crystalline, hydrated solid.

6. Form F

[0132] Form F was prepared as a mixture with Form A (Form A+F) by precipitation from a THF:water (9:1) solution of Compound I under slow cooling crystallization conditions (Example 8). Precipitates of Form A+F was characterized by XRPD.

[0133] The characteristic XRPD spectrum (CuKα, λ=1.5418Å) of Form A+F is shown in Figure 24. The XRPD pattern confirmed that the material was crystalline. The major diffraction peaks expressed in °2θ and their relative intensities are summarized in Table 6.

Table 6. Characteristic XRPD Peaks (CuKα) of Form A+F

Peak No.	2θ (°)	d-spacing	Intensity	I/I _o
1	11.37	7.78	420	61
2	11.93	7.41	335	49
3	12.39	7.14	298	44
4	12.91	6.85	77	11
5	14.54	6.09	267	39
6	15.86	5.58	351	51
7	16.88	5.25	358	52
8	17.54	5.05	128	19

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
9	19.39	4.57	289	42
10	20.63	4.30	258	38
11	21.05	4.22	314	46
12	22.34	3.98	683	100
13	23.98	3.71	355	52
14	24.71	3.60	158	23
15	25.29	3.52	150	22
16	26.03	3.42	127	19
17	26.63	3.35	92	13
18	27.80	3.21	113	17
19	30.05	2.97	81	12
20	31.86	2.81	78	11

[0134] The unique set of XRPD peaks above or a subset of peaks thereof can be used to identify Form A+F of Compound I. A subset of peaks which are characteristic of Form F alone was which may be used to identify Form F was obtained by subtracting the XRPD peaks of Form A from those of Form A+F; this subset comprises peaks at about 12.39, 20.63, 26.03, and 30.05 °2 θ and may be used to identify the presence of Form E

7. Form G

[0135] Form G was prepared as a mixture with Form A and Form E (Form A+E+G) by precipitation from a THF:water (9:1) solution of Compound I by slurring at ambient conditions (Example 9). The precipitant was characterized by XRPD and was confirmed to be crystalline. The characteristic XRPD spectrum (CuK α) of Form A+E+G is shown in Figure 25. The major diffraction peaks expressed in °2 θ and their relative intensities are summarized in Table 7.

Table 7. Characteristic XRPD Peaks (CuK α) of Form A+E+G

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
1	4.60	19.21	88	16
2	9.27	9.53	57	10
3	11.37	7.78	144	26
4	12.01	7.37	96	17
5	12.21	7.24	93	17
6	12.64	7.00	334	60
7	13.22	6.69	293	52
8	13.91	6.36	50	9
9	14.23	6.22	228	41
10	14.59	6.07	69	12
11	15.50	5.71	142	25
12	15.87	5.58	86	15

Peak No.	2 θ (°)	d-spacing	Intensity	I/I _o
13	16.81	5.27	426	76
14	17.33	5.11	79	14
15	17.71	5.00	62	11
16	18.08	4.90	137	24
17	18.62	4.76	230	41
18	19.20	4.62	259	46
19	19.77	4.49	128	23
20	20.42	4.35	273	49
21	20.99	4.23	81	14
22	21.28	4.17	139	25
23	21.76	4.08	560	100
24	22.33	3.98	376	67
25	23.44	3.79	419	75
26	23.79	3.74	95	17
27	24.36	3.65	223	40
28	24.69	3.60	93	17
29	25.06	3.55	281	50
30	25.39	3.51	182	33
31	25.77	3.45	70	13
32	25.95	3.43	107	19
33	26.33	3.38	64	11
34	26.83	3.32	107	19
35	27.15	3.28	78	14
36	27.54	3.24	124	22
37	28.07	3.18	93	17
38	28.69	3.11	176	31
39	29.13	3.06	53	9
40	29.87	2.99	130	23
41	30.71	2.91	98	18
42	32.75	2.73	110	20

[0136] The unique set of XRPD peaks above or a subset thereof can be used to identify Form A+E+G of Compound I. A subset of peaks that is characteristic of Form G was obtained by subtracting the XRPD peaks of Form A and Form E from the XRPD spectrum of Form A+E+G; this subset comprised peaks positioned at about 13.22, 14.23, 18.62, 19.77, 24.36, 25.06, and 30.71 °2 θ . This subset of characteristic peaks of Form G may be used for identifying the presence of Form G.

8. Amorphous Form

[0137] The Amorphous Form of Compound I was prepared by lyophilization of an aqueous solution of Compound I (Example 10). The residue material was characterized by XRPD and the resulting XRPD spectrum displayed in Figure 26. The XRPD spectrum shows a broad halo with no specific peaks present, which confirms that the material is

amorphous. The material was further characterized by TGA, DSC, hot stage microscopy, and moisture sorption analysis.

[0138] TGA analysis (Figure 27) showed a 1.8% weight loss from 25 °C to 95 °C, which was likely due to loss of residual solvent.

[0139] DSC analysis (Figure 28) showed a slightly concave baseline up to an exotherm at 130 °C (recrystallization), followed by an endotherm at 194 °C, which results from the melting of Form A. Hot stage microscopy confirmed these recrystallization and melting events (micrographs not included). An approximate glass transition was observed (Figure 29) at an onset temperature of 82 °C.

[0140] Moisture sorption/desorption data (Figure 30 and Example 25) showed a 1.0% weight loss on equilibration at 5% relative humidity. Approximately 8% of weight was gained up to 65% relative humidity. Approximately 7% of weight was lost at 75% relative humidity. This is likely due to the recrystallization of the amorphous material to a crystalline solid. A 4.4% weight gain was observed on sorption from 75% to 95% relative humidity. Approximately 4.7% weight was lost on desorption from 95% to 5% relative humidity. The solid material remaining after the moisture sorption analysis was determined to be Form A by XRPD (Figure 31).

[0141] Stressing Amorphous Form in acetone or ethanol vapors resulted in Form A (Example 22). Stressing Amorphous Form at 58% and 88% relative humidity resulted in Form A (Figure 19). Stressing Amorphous Form in water vapor resulted in a mixture of Forms A and D (Example 22).

Indications for Use of Compound I

[0142] The present invention also relates to methods to alter, preferably to reduce DPP-IV activity within a subject by administering Compound I in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, Amorphous Form, and combinations thereof.

[0143] DPP-IV is believed to contribute to the pathology and/or symptomology of several different diseases such that reduction of the activity of DPP-IV in a subject through inhibition may be used to therapeutically address these disease states. Examples of various diseases that may be treated using Compound I of the present invention are described herein. It is noted that additional diseases beyond those disclosed herein may be later identified as the biological roles that DPP-IV plays in various pathways becomes

more fully understood.

[0144] Compound I may be used to treat or prevent diabetes and obesity related conditions. DPP-IV has been shown to be the primary degrading enzyme of GLP-1 (7-36), *in vivo*. GLP-1 (7-36) has multiple actions *in vivo* including the stimulation of insulin secretion, inhibition of glucagon secretion, the promotion of satiety, and the slowing of gastric emptying. The actions of GLP-1 (7-36) are believed to be beneficial in the prevention and treatment of type II diabetes and potentially obesity. DPP-IV degrades GLP-1 (7-36) efficiently to GLP-1 (9-36), which has been speculated to act as a physiological antagonist to GLP-1 (7-36). Inhibiting DPP-IV *in vivo* is therefore believed to be useful for potentiating endogenous levels of GLP-1 (7-36) and attenuating the formation of its antagonist GLP-1 (9-36). Thus, Compound I are believed to be useful in preventing, delaying the progression, and/or treating diabetes and more particularly, type 2 diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation and obesity.

[0145] Compound I may be used as immuno-suppressants (or cytokine release suppressant drugs). DPP-IV expression is increased in T-cells upon mitogenic or antigenic stimulation (Mattem, T., et al., *Scand. J. Immunol.*, **1991**, 33, 737). It has been reported that inhibitors of DPP-IV and antibodies to DPP-IV suppress the proliferation of mitogen-stimulated and antigen-stimulated T-cells in a dose-dependant manner (Schon, E., et al., *Biol. Chem.*, **1991**, 372, 305). Various other functions of T-lymphocytes such as cytokine production, IL-2 mediated cell proliferation and B-cell helper activity have been shown to be dependent on DPP-IV activity (Schon, E., et al., *Scand. J. Immunol.*, **1989**, 29, 127). DPP-IV inhibitors, based on boroProline, (Flentke, G. R., et al., *Proc. Nat. Acad. Sci. USA*, **1991**, 88, 1556) although unstable, were effective at inhibiting antigen-induced lymphocyte proliferation and IL-2 production in murine CD4⁺ T-helper cells. Such boronic acid inhibitors have been shown to have an effect *in vivo* in mice causing suppression of antibody production induced by immune challenge (Kubota, T. et al., *Clin. Exp. Immun.*, 1992, 89, 192). The role of DPP-IV in regulating T lymphocyte activation may also be attributed, in part, to its cell-surface association with the transmembrane phosphatase, CD45. DPP-IV inhibitors or non-active site ligands may possibly disrupt the CD45-DPP-IV association. CD45 is known to be an integral component of the T-cell signaling apparatus. It has been reported that DPP-IV is essential for the penetration and

infectivity of HIV-1 and HIV-2 viruses in CD4⁺ T-cells (Wakselman, M., Nguyen, C., Mazaleyrat, J.-P., Callebaut, C., Krust, B., Hovanessian, A. G., Inhibition of HIV-1 infection of CD26⁺ but not CD26⁻ cells by a potent cyclopeptidic inhibitor of the DPP-IV activity of CD26, Abstract P.44 of the 24th European Peptide Symposium 1996).

Additionally, DPP-IV has been shown to associate with the enzyme adenosine deaminase (ADA) on the surface of T-cells (Kameoka, J., et al., *Science*, 193, 26 466). ADA deficiency causes severe combined immunodeficiency disease (SCID) in humans. This ADA-CD26 interaction may provide clues to the pathophysiology of SCID. It follows that inhibitors of DPP-IV may be useful immunosuppressants (or cytokine release suppressant drugs) for the treatment of, among other things, organ transplant rejection; autoimmune diseases such as inflammatory bowel disease, multiple sclerosis and rheumatoid arthritis; and the treatment of AIDS.

[0146] Compound I may be used for treating various cancers. It has been shown that lung endothelial cell DPP-IV is an adhesion molecule for lung-metastatic rat breast and prostate carcinoma cells (Johnson, R. C., et al., *J. Cell. Biol.*, **1993**, 121, 1423). DPP-IV is known to bind to fibronectin and some metastatic tumor cells are known to carry large amounts of fibronectin on their surface. Potent DPP-IV inhibitors may be useful as drugs to prevent metastases of, for example, breast and prostate tumors to the lungs.

[0147] Compound I may be useful as agents to treat dermatological diseases such as psoriasis and lichen planus. High levels of DPP-IV expression have also been found in human skin fibroblast cells from patients with psoriasis, rheumatoid arthritis (RA) and lichen planus (Raynaud, F., et al., *J. Cell. Physiol.*, **1992**, 151, 378). Therefore, DPP-IV inhibitors may be useful as agents to treat these conditions.

[0148] Compound I may be useful as a male contraceptive agent, and for treating female infertility and amenorrhea. High DPP-IV activity has been found in tissue homogenates from patients with benign prostate hypertrophy and in prostatosomes. These are prostate derived organelles important for the enhancement of sperm forward motility (Vanhoof, G., et al., *Eur. J. Clin. Chem. Clin. Biochem.*, **1992**, 30, 333). DPP-IV inhibitors may also act to suppress sperm motility and therefore act as a male contraceptive agent. Conversely, DPP-IV inhibitors have been implicated as novel for treatment of infertility, and particularly human female infertility due to polycystic ovary syndrome (PCOS, Stein-Leventhal syndrome) which is a condition characterized by thickening of the ovarian capsule and formation of multiple follicular cysts. It results in

infertility and amenorrhea.

[0149] Compound I may be used to modulate cleavage of various cytokines (stimulating hematopoietic cells), growth factors and neuropeptides. It was discovered that inhibitors of DPP-IV are useful for stimulating the growth and differentiation of hematopoietic cells in the absence of exogenously added cytokines or other growth factors or stromal cells. Stimulated hematopoietic cells are useful for the treatment of disorders that are characterized by a reduced number of hematopoietic cells or their precursors *in vivo*. Such conditions occur frequently in patients who are immunosuppressed, for example, as a consequence of chemotherapy and/or radiation therapy for cancer.

[0150] Compound I may be useful in the treatment of short stature due to growth hormone deficiency (Dwarfism). DPP-IV in human plasma has been shown to cleave N-terminal Tyr-Ala from growth hormone-releasing factor and cause inactivation of the hormone. Accordingly, inhibiting DPP-IV will modulate its effect and may promote GH-dependent tissue growth or re-growth.

[0151] Compound I may be useful for the regulation or normalization of neurological disorders. DPP-IV can cleave neuropeptides and has been shown to modulate the activity of neuroactive peptides substance P, neuropeptide Y and CLIP (Mentlein, R., Dahms, P., Grandt, D., Kruger, R., Proteolytic processing of neuropeptide Y and peptide YY by DPP-IV, *Regul. Pept.*, 49, 133, 1993; Wetzel, W., Wagner, T., Vogel, D., Demuth, H.-U., Balschun, D., Effects of the CLIP fragment ACTH 20-24 on the duration of REM sleep episodes, *Neuropeptides*, 31, 41, 1997). Thus, inhibiting DPP-IV would lessen its damaging effect on the neuropeptides.

[0152] Compositions, according to the present invention, may be administered, or coadministered with other active agents. These additional active agents may include, for example, one or more other pharmaceutically active agents. Coadministration in the context of this invention is intended to mean the administration of more than one therapeutic agent, one of which includes Compound I. Such co-administration may also be coextensive, that is, occurring during overlapping periods of time or may be sequential, that is, occurring during non-overlapping periods of time. Examples of co-administration of Compound I with other active ingredients in a combination therapy are described in U.S. Patent Application Ser. No. 11/531,671, filed September 13, 2006, the disclosure of which is expressly incorporated herein by reference in its entirety.

[0153] For oncology indications, Compound I may be administered in conjunction with

other agents to inhibit undesirable and uncontrolled cell proliferation. Examples of other anti-cell proliferation agents that may be used in conjunction with Compound I include, but are not limited to, retinoid acid and derivatives thereof, 2-methoxyestradiol, ANGIOSTATINTM protein, ENDOSTATINTM protein, suramin, squalamine, tissue inhibitor of metalloproteinase-1, tissue inhibitor of metalloproteinase-2, plasminogen activator inhibitor-1, plasminogen activator inhibitor-2, cartilage-derived inhibitor, paclitaxel, platelet factor 4, protamine sulfate (clupeine), sulfated chitin derivatives (prepared from queen crab shells), sulfated polysaccharide peptidoglycan complex (sp-pg), staurosporine, modulators of matrix metabolism, including for example, proline analogs ((l-azetidine-2-carboxylic acid (LACA)), cishydroxyproline, d,l-3,4-dehydroproline, thiaproline, beta.-aminopropionitrile fumarate, 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone, methotrexate, mitoxantrone, heparin, interferons, 2 macroglobulin-serum, chimp-3, chymostatin, beta.-cyclodextrin tetradecasulfate, eponemycin; fumagillin, gold sodium thiomalate, d-penicillamine (CDPT), beta.-1-anticollagenase-serum, alpha.2-antiplasmin, bisantrene, lobenzarit disodium, n-2-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA", thalidomide; angostatic steroid, carboxyaminoimidazole; metalloproteinase inhibitors such as BB94. Other anti-angiogenesis agents that may be used include antibodies, preferably monoclonal antibodies against these angiogenic growth factors: bFGF, aFGF, FGF-5, VEGF isoforms, VEGF-C, HGF/SF and Ang-1/Ang-2. Ferrara N. and Alitalo, K. "Clinical application of angiogenic growth factors and their inhibitors" (1999) Nature Medicine 5:1359-1364.

[0154] In another embodiment, a therapeutic method is provided that comprises administering Compound I. In another embodiment, a method of inhibiting cell proliferation is provided that comprises contacting a cell with an effective amount of Compound I. In another embodiment, a method of inhibiting cell proliferation in a patient is provided that comprises administering to the patient a therapeutically effective amount of Compound I.

[0155] In another embodiment, a method of treating diabetes and related conditions, including, but are not limited to, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation, obesity, in a patient is provided that comprises administering to the patient a therapeutically effective amount of Compound I.

[0156] In another embodiment, a method of treating a condition in a patient which is

known to be mediated by DPP-IV, or which is known to be treated by DPP-IV inhibitors, comprising administering to the patient a therapeutically effective amount of Compound I. In another embodiment, a method is provided for using Compound I in order to manufacture a medicament for use in the treatment of disease state which is known to be mediated by DPP-IV, or which is known to be treated by DPP-IV inhibitors.

[0157] In another embodiment, a method is provided for treating a disease state for which DPP-IV possesses activity that contributes to the pathology and/or symptomology of the disease state, the method comprising: administering Compound I to a subject such that the free base form of Compound I is present in the subject in a therapeutically effective amount for the disease state.

[0158] The present invention relates generally to a method comprising administering between 1 mg/day and 250 mg/day of Compound I to a patient, optionally between 2.5 mg and 200 mg of Compound I, optionally between 2.5 mg and 150 mg of Compound I, and optionally between 5 mg and 100 mg of Compound I (in each instance based on the molecular weight of the free base form of Compound I). Specific dosage amounts that may be used include, but are not limited to 2.5 mg, 5 mg, 6.25 mg, 10 mg, 20 mg, 25 mg, 50 mg, 75 mg and 100 mg of Compound I per day. It is noted that the dosage may be administered as a daily dose or weekly dose, once daily or multiple doses per day. It is noted that Compound I may be administered in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. However, the dosage amounts and ranges provided herein are always based on the molecular weight of the free base form of Compound I.

[0159] Compound I may be administered by any route of administration. In particular embodiments, however, the method of the present invention is practiced by administering Compound I orally.

Pharmaceutical Compositions Comprising Compound I Where At Least One of Form A, Form B, Form C, Form D, Form E, Form F, Form G, or Amorphous Form Is Present

[0160] Compound I may be used in various pharmaceutical compositions where at least a portion of Compound I is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. The pharmaceutical composition should contain a sufficient quantity

of Compound I to reduce dipeptidyl peptidases activity *in vivo* sufficiently to provide the desired therapeutic effect. Such pharmaceutical compositions may comprise Compound I present in the composition in a range of between 0.005% and 100% (weight/weight), optionally 0.1-95%, and optionally 1-95%.

[0161] In particular embodiments, the pharmaceutical compositions comprise at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I in a form selected from a group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, Amorphous Form, and a mixture thereof. In another embodiment, a particular polymorphic form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, Amorphous Form may comprise at least 0.1%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of the total amount of Compound I (weight/weight) in the pharmaceutical composition.

[0162] In addition to Compound I, the pharmaceutical composition may comprise one or more additional components that do not deleteriously affect the use of Compound I. For example, the pharmaceutical compositions may include, in addition to Compound I, conventional pharmaceutical carriers; excipients; diluents; lubricants; binders; wetting agents; disintegrating agents; glidants; sweetening agents; flavoring agents; emulsifying agents; solubilizing agents; pH buffering agents; perfuming agents; surface stabilizing agents; suspending agents; and other conventional, pharmaceutically inactive agents. In particular, the pharmaceutical compositions may comprise lactose, mannitol, glucose, sucrose, dicalcium phosphate, magnesium carbonate, sodium saccharin, carboxymethylcellulose, magnesium stearate, calcium stearate, sodium crosscarmellose, talc, starch, natural gums (*e.g.*, gum acaciagelatin), molasses, polyvinylpyrrolidone, celluloses and derivatives thereof, povidone, crospovidones acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, biocompatible polymers, such as collagen, ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others such agents.

[0163] Pharmaceutical compositions according to the present invention may be adapted for administration by any of a variety of routes. For example, pharmaceutical compositions according to the present invention can be administered orally, parenterally, intraperitoneally, intravenously, intraarterially, topically, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example, by catheter or stent),

subcutaneously, intraadiposally, intraarticularly, or intrathecally, optionally in a slow release dosage form. In particular embodiments, the pharmaceutical compounds are administered orally, by inhalation or by injection subcutaneously, intramuscularly, intravenously or directly into the cerebrospinal fluid.

[0164] In general, the pharmaceutical compositions of the present invention may be prepared in a gaseous, liquid, semi-liquid, gel, or solid form, and formulated in a manner suitable for the route of administration to be used.

[0165] Compositions according to the present invention are optionally provided for administration to humans and animals in unit dosage forms or multiple dosage forms, such as tablets, capsules, pills, powders, dry powders for inhalers, granules, sterile parenteral solutions or suspensions, oral solutions or suspensions, oil-water emulsions, sustained release formulations, such as, but not limited to, implants and microencapsulated delivery systems, containing suitable quantities of Compound I. Methods of preparing such dosage forms are known in the art, and will be apparent to those skilled in this art; for example, see *Remington's Pharmaceutical Sciences*, 19th Ed. (Easton, Pa.: Mack Publishing Company, 1995).

[0166] Unit-dose forms, as used herein, refers to physically discrete units suitable for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of Compound I sufficient to produce the desired therapeutic effect, in association with a pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes, and individually packaged tablets or capsules. Unit-dose forms may be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules, or bottles of pints or gallons. Hence, multiple dose form may be viewed as a multiple of unit-doses that are not segregated in packaging.

[0167] In general, the total amount of Compound I in a pharmaceutical composition according to the present invention should be sufficient to a desired therapeutic effect. This amount may be delivered as a single per day dosage, multiple dosages per day to be administered at intervals of time, or as a continuous release dosage form. It is noted that Compound I may be administered in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, Amorphous Form, and a mixture

thereof. Compound I may advantageously be used when administered to a patient at a daily dose of between 1 mg/day and 250 mg/day of Compound I, optionally between 2.5 mg and 200 mg of Compound I, optionally between 2.5 mg and 150 mg of Compound I, and optionally between 5 mg and 100 mg of Compound I (in each instance based on the molecular weight of the free base form of Compound I). Specific dosage amounts that may be used include, but are not limited to 2.5 mg, 5 mg, 6.25 mg, 10 mg, 20 mg, 25 mg, 50 mg, 75 mg, and 100 mg of Compound I per day. It may be desirable for Compound I to be administered one time per day. Accordingly, pharmaceutical compositions of the present invention may be in the form of a single dose form comprising between 1 mg/day and 250 mg/day of Compound I, optionally between 2.5 mg and 200 mg of Compound I, optionally between 2.5 mg and 150 mg of Compound I, and optionally between 5 mg and 100 mg of Compound I. In specific embodiments, the pharmaceutical composition comprises 2.5 mg, 5 mg, 6.25 mg, 10 mg, 20 mg, 25 mg, 50 mg, 75 mg or 100 mg of Compound I.

A. Formulations for Oral Administration

[0168] Oral pharmaceutical dosage forms may be as a solid, gel or liquid where at least a portion of Compound I is present in the composition in a form selected from the group consisting of one or more of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form.

[0169] In certain embodiments, Compound I is provided as solid dosage forms. Examples of solid dosage forms include, but are not limited to pills, tablets, troches, capsules, granules, and bulk powders. More specific examples of oral tablets include compressed, chewable lozenges, troches and tablets that may be enteric-coated, sugar-coated or film-coated. Examples of capsules include hard or soft gelatin capsules. Granules and powders may be provided in non-effervescent or effervescent forms. The powders may be prepared by lyophilization or by other suitable methods.

[0170] The tablets, pills, capsules, troches and the like may optionally contain one or more of the following ingredients, or compounds of a similar nature: a binder; a diluent; a disintegrating agent; a lubricant; a glidant; a coloring agent; a sweetening agent; a flavoring agent; and a wetting agent.

[0171] Examples of binders that may be used include, but are not limited to, microcrystalline cellulose, gum tragacanth, glucose solution, acacia mucilage, gelatin solution, sucrose, and starch paste.

[0172] Examples of diluents that may be used include, but are not limited to, lactose, sucrose, starch, kaolin, salt, mannitol, and dicalcium phosphate.

[0173] Examples of disintegrating agents that may be used include, but are not limited to, crosscarmellose sodium, sodium starch glycolate, alginic acid, corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose.

[0174] Examples of lubricants that may be used include, but are not limited to, talc, starch, magnesium or calcium stearate, lycopodium and stearic acid.

[0175] Examples of glidants that may be used include, but are not limited to, colloidal silicon dioxide.

[0176] Examples of coloring agents that may be used include, but are not limited to, any of the approved certified water soluble FD and C dyes, mixtures thereof; and water insoluble FD and C dyes suspended on alumina hydrate.

[0177] Examples of sweetening agents that may be used include, but are not limited to, sucrose, lactose, mannitol and artificial sweetening agents such as sodium cyclamate and saccharin, and any number of spray-dried flavors.

[0178] Examples of flavoring agents that may be used include, but are not limited to, natural flavors extracted from plants such as fruits and synthetic blends of compounds that produce a pleasant sensation, such as, but not limited to peppermint and methyl salicylate.

[0179] Examples of wetting agents that may be used include, but are not limited to, propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate, and polyoxyethylene lauryl ether.

[0180] Examples of anti-emetic coatings that may be used include, but are not limited to, fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates.

[0181] Examples of film coatings that may be used include, but are not limited to, hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

[0182] When the dosage form is a pill, tablet, torches, or the like, Compound I may optionally be provided in a composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an enteric coating that maintains its integrity in the stomach and releases the active compound in the intestine. The composition may also be formulated in combination with an antacid or other such ingredient.

[0183] When the dosage unit form is a capsule, it may optionally additionally comprise

a liquid carrier such as a fatty oil. In addition, dosage unit forms may optionally additionally comprise various other materials that modify the physical form of the dosage unit, for example, coatings of sugar and other enteric agents.

[0184] Compound I may also be administered as a component of an elixir, emulsion, suspension, microsuspension, syrup, wafer, sprinkle, chewing gum or the like. A syrup may optionally comprise, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes, colorings, and flavors.

[0185] Alternatively, liquid or semi-solid oral formulations may be prepared by dissolving or dispersing the active compound or salt in vegetable oils, glycols, triglycerides, propylene glycol esters (e.g. propylene carbonate) and other such carriers, and encapsulating these solutions or suspensions in hard or soft gelatin capsule shells. Other useful formulations include those set forth in U.S. Pat. Nos. Re 28,819 and 4,358,603.

[0186] Examples of oral formulations that may be used to administer Compound I has been described in U.S. Patent Application Ser. No. 11/531,671, filed September 13, 2006, the disclosure of which is herein expressly incorporated by reference in its entirety.

[0187] Exemplary tablet formulations are provided below. It is noted that the examples are, by way of illustration but not limitation. It is also noted that Compound I is present in the formulation in a form selected from the group consisting of one or more of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form. It is also noted that the formulations provided herein may be varied as is known in the art.

12.5 mg of Compound I (weight of free base form) per tablet

Core Tablet Formulation

(1) Compound I	17.0 mg
(2) Lactose Monohydrate, NF, Ph, Eur (FOREMOST 316 FAST FLO)	224.6 mg
(3) Microcrystalline Cellulose, NF, Ph, Eur (AVICEL PH 102)	120.1 mg
(4) Croscarmellose Sodium, NF, Ph, Eur (AC-DO-SOL)	32.0 mg
(5) Colloidal Silicon Dioxide, NF, Ph, Eur (CAB-O-SIL M-5P)	3.2 mg
(6) Magnesium Stearate, NF, Ph, Eur (MALLINCKRODT, Non-bovine Hyqual)	3.2 mg
TOTAL (per tablet)	400.0 mg

Film Coat (12.0 mg in total)

- (1) Opadry II 85F18422, White – Portion 1 (COLORCON)
- (2) Opadry II 85F18422, White – Portion 2 (COLORCON)
- (3) Opadry II 85F18422, White – Portion 3 (COLORCON)

25 mg of Compound I (weight of free base form) per tablet

Core Tablet Formulation

(1) Compound I	34.0 mg
(2) Lactose Monohydrate, NF, Ph, Eur (FOREMOST 316 FAST FLO)	207.6 mg
(3) Microcrystalline Cellulose, NF, Ph, Eur (AVICEL PH 102)	120.1 mg
(4) Croscarmellose Sodium, NF, Ph, Eur (AC-DO-SOL)	32.0 mg
(5) Colloidal Silicon Dioxide, NF, Ph, Eur (CAB-O-SIL M-5P)	3.2 mg
(6) Magnesium Stearate, NF, Ph, Eur (MALLINCKRODT, Non-bovine Hyqual)	3.2 mg
TOTAL (per tablet)	400.0 mg

Film Coat (12.0 mg in total)

- (1) Opadry II 85F18422, White – Portion 1 (COLORCON)
- (2) Opadry II 85F18422, White – Portion 2 (COLORCON)
- (3) Opadry II 85F18422, White – Portion 3 (COLORCON)

50 mg of Compound I (weight of free base form) per tablet

Core Tablet Formulation

(1) Compound I	68.0 mg
(2) Lactose Monohydrate, NF, Ph, Eur (FOREMOST 316 FAST FLO)	173.6 mg
(3) Microcrystalline Cellulose, NF, Ph, Eur (AVICEL PH 102)	120.1 mg
(4) Croscarmellose Sodium, NF, Ph, Eur (AC-DO-SOL)	32.0 mg
(5) Colloidal Silicon Dioxide, NF, Ph, Eur (CAB-O-SIL M-5P)	3.2 mg
(6) Magnesium Stearate, NF, Ph, Eur (MALLINCKRODT, Non-bovine Hyqual)	3.2 mg
TOTAL (per tablet)	400.0 mg

Film Coat (12.0 mg in total)

- (1) Opadry II 85F18422, White – Portion 1 (COLORCON)

(2) Opadry II 85F18422, White – Portion 2 (COLORCON)

(3) Opadry II 85F18422, White – Portion 3 (COLORCON)

B. Injectables, solutions and emulsions

[0188] Compound I present in a form or a mixture of forms selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form may be formulated for parenteral administration. Parenteral administration generally characterized by injection, either subcutaneously, intramuscularly or intravenously. Implantation of a slow-release or sustained-release system, such that a constant level of dosage is maintained (see, e.g., U.S. Pat. No. 3,710,795) is also contemplated herein. The percentage of active compound contained in such parenteral compositions is highly dependent on the route of administration and the indication of disease to be treated.

[0189] Injectables may be prepared in any conventional form. These formulations include, but are not limited to, sterile solutions, suspensions, microsuspensions, and emulsions ready for injection, and solid forms, e.g., lyophilized or other powders including hypodermic tablets, ready to be combined with a carrier just prior to use. Generally, the resulting formulation may be a solution, microsuspension, suspension and emulsion. The carrier may be an aqueous, non-aqueous liquid, or a solid vehicle that can be suspended in liquid.

[0190] Examples of carriers that may be used in conjunction with injectables according to the present invention include, but are not limited to water, saline, dextrose, glycerol or ethanol. The injectable compositions may also optionally comprise minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents, stabilizers, solubility enhancers, and other such agents, such as for example, sodium acetate, sorbitan monolaurate, triethanolamine oleate and cyclodextrins.

[0191] When administered intravenously, examples of suitable carriers include, but are not limited to physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene glycol and mixtures thereof.

[0192] Examples of pharmaceutically acceptable carriers that may optionally be used in parenteral preparations include, but are not limited to aqueous vehicles, nonaqueous vehicles, antimicrobial agents, isotonic agents, buffers, antioxidants, local anesthetics,

suspending and dispersing agents, emulsifying agents, sequestering or chelating agents and other pharmaceutically acceptable substances.

[0193] Examples of aqueous vehicles that may optionally be used include Sodium Chloride Injection, Ringers Injection, Isotonic Dextrose Injection, Sterile Water Injection, Dextrose and Lactated Ringers Injection.

[0194] Examples of nonaqueous parenteral vehicles that may optionally be used include fixed oils of vegetable origin, cottonseed oil, corn oil, sesame oil and peanut oil.

[0195] Antimicrobial agents in bacteriostatic or fungistatic concentrations may be added to parenteral preparations, particularly when the preparations are packaged in multiple-dose containers and thus designed to be stored and multiple aliquots to be removed. Examples of antimicrobial agents that may be used include phenols or cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl p-hydroxybenzoic acid esters, thimerosal, benzalkonium chloride and benzethonium chloride.

[0196] Examples of isotonic agents that may be used include sodium chloride and dextrose. Examples of buffers that may be used include phosphate and citrate. Examples of antioxidants that may be used include sodium bisulfate. Examples of local anesthetics that may be used include procaine hydrochloride. Examples of suspending and dispersing agents that may be used include sodium carboxymethylcellulose, hydroxypropyl methylcellulose and polyvinylpyrrolidone. Examples of emulsifying agents that may be used include Polysorbate 80 (TWEEN 80). A sequestering or chelating agent of metal ions includes EDTA.

[0197] Pharmaceutical carriers may also optionally include ethyl alcohol, polyethylene glycol and propylene glycol for water miscible vehicles and sodium hydroxide, hydrochloric acid, citric acid or lactic acid for pH adjustment.

[0198] The concentration of Compound I in the parenteral formulation may be adjusted so that an injection administers a pharmaceutically effective amount sufficient to produce the desired pharmacological effect. The exact concentration of Compound I and/or dosage to be used will ultimately depend on the age, weight and condition of the patient or animal as is known in the art.

[0199] Unit-dose parenteral preparations may be packaged in an ampoule, a vial or a syringe with a needle. All preparations for parenteral administration should be sterile, as is known and practiced in the art.

[0200] Injectables may be designed for local and systemic administration. Typically a

therapeutically effective dosage is formulated to contain a concentration of at least about 0.1% w/w up to about 90% w/w or more, preferably more than 1% w/w of Compound I to the treated tissue(s). Compound I may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment will be a function of the location of where the composition is parenterally administered, the carrier and other variables that may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test data. It is to be noted that concentrations and dosage values may also vary with the age of the individual treated. It is to be further understood that for any particular subject, specific dosage regimens may need to be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations. Hence, the concentration ranges set forth herein are intended to be exemplary and are not intended to limit the scope or practice of the claimed formulations.

[0201] Compound I may optionally be suspended in micronized or other suitable form or may be derivatized to produce a more soluble active product or to produce a prodrug. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for ameliorating the symptoms of the disease state and may be empirically determined.

C. Powders

[0202] Compound I in a form or a mixture of forms selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form may be prepared as powders, which can be reconstituted for administration as solutions, emulsions and other mixtures. The powders may also be formulated as solids or gels.

[0203] Powders of Compound I may be prepared by grinding, spray drying, lyophilization and other techniques that are well known in the art. Sterile, lyophilized powder may be prepared by dissolving Compound I in a sodium phosphate buffer solution containing dextrose or other suitable excipient. Subsequent sterile filtration of the solution followed by lyophilization under standard conditions known to those of skill in the art provides the desired formulation. Briefly, the lyophilized powder may optionally be prepared by dissolving dextrose, sorbitol, fructose, corn syrup, xylitol, glycerin, glucose, sucrose or other suitable agent, about 1-20%, preferably about 5 to 15%, in a suitable

buffer, such as citrate, sodium or potassium phosphate or other such buffer known to those of skill in the art at, typically, about neutral pH. Then, Compound I is added to the resulting mixture, preferably above room temperature, more preferably at about 30-35 °C, and stirred until it dissolves. The resulting mixture is diluted by adding more buffer to a desired concentration. The resulting mixture is sterile filtered or treated to remove particulates and to insure sterility, and apportioned into vials for lyophilization. Each vial may contain a single dosage or multiple dosages of Compound I.

D. Topical administration

[0204] Compound I present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form may also be administered as topical mixtures. Topical mixtures may be used for local and systemic administration. The resulting mixture may be a solution, suspension, microsuspension, emulsions or the like and are formulated as creams, gels, ointments, emulsions, solutions, elixirs, lotions, suspensions, tinctures, pastes, foams, aerosols, irrigations, sprays, suppositories, bandages, dermal patches or any other formulations suitable for topical administration.

[0205] Compound I may be formulated for topical applications to the respiratory tract. These pulmonary formulations can be in the form of an aerosol, solution, emulsion, suspension, microsuspension for a nebulizer, or as a microfine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will typically have diameters of less than 50 microns, preferably less than 10 microns. Examples of aerosols for topical application, such as by inhalation are disclosed in U.S. Pat. Nos. 4,044,126, 4,414,209, and 4,364,923, which describe aerosols for delivery of a steroid useful for treatment inflammatory diseases, particularly asthma.

[0206] Compound I may also be formulated for local or topical application, such as for topical application to the skin and mucous membranes, such as in the eye, in the form of gels, creams, and lotions and for application to the eye or for intracisternal or intraspinal application. Topical administration is contemplated for transdermal delivery and also for administration to the eyes or mucosa, or for inhalation therapies. Nasal solutions or suspensions of Compound I alone or in combination with other pharmaceutically acceptable excipients can also be administered.

E. Formulations for other routes of administration

[0207] Depending upon the disease state being treated, Compound I present in one form or a mixture of forms selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form may be formulated for other routes of administration, such as topical application, transdermal patches, and rectal administration, may also be used. For example, pharmaceutical dosage forms for rectal administration are rectal suppositories, capsules and tablets for systemic effect. Rectal suppositories are used herein mean solid bodies for insertion into the rectum that melt or soften at body temperature releasing one or more pharmacologically or therapeutically active ingredients. Pharmaceutically acceptable substances utilized in rectal suppositories are bases or vehicles and agents to raise the melting point. Examples of bases include cocoa butter (theobroma oil), glycerin-gelatin, carbowax, (polyoxyethylene glycol) and appropriate mixtures of mono-, di- and triglycerides of fatty acids. Combinations of the various bases may be used. Agents to raise the melting point of suppositories include spermaceti and wax. Rectal suppositories may be prepared either by the compressed method or by molding. The typical weight of a rectal suppository is about 2 to 3 gm. Tablets and capsules for rectal administration may be manufactured using the same pharmaceutically acceptable substance and by the same methods as for formulations for oral administration.

Kits and Articles of Manufacture Comprising Compound I Polymorphs

[0208] The present invention is also directed to kits and other articles of manufacture for treating diseases associated with dipeptidyl peptidases. It is noted that diseases are intended to cover all conditions for which the dipeptidyl peptidases possesses activity that contributes to the pathology and/or symptomology of the condition.

[0209] In one embodiment, a kit is provided that comprises a pharmaceutical composition comprising Compound I where greater than 0.1%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form; and instructions for use of the kit. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I. The instructions may indicate the disease state for which the composition is to be administered, storage information, dosing

information and/or instructions regarding how to administer the composition. The kit may also comprise packaging materials. The packaging material may comprise a container for housing the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.

[0210] In another embodiment, an article of manufacture is provided that comprises a pharmaceutical composition comprising Compound I where greater than 0.1%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97% or 99% of Compound I (by weight) is present in the composition in a form selected from the group consisting of Form A, Form B, Form C, Form D, Form E, Form F, Form G, and Amorphous Form; and packaging materials. Optionally, the composition comprises at least 0.1%, 0.25%, 0.5%, 1%, 5%, 10%, 25%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% of Compound I. The packaging material may comprise a container for housing the composition. The container may optionally comprise a label indicating the disease state for which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition. The kit may also optionally comprise additional components, such as syringes for administration of the composition. The kit may comprise the composition in single or multiple dose forms.


[0211] It is noted that the packaging material used in kits and articles of manufacture according to the present invention may form a plurality of divided containers such as a divided bottle or a divided foil packet. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material, for example a paper or cardboard box, a glass or plastic bottle or jar, a re-sealable bag (for example, to hold a "refill" of tablets for placement into a different container), or a blister pack with individual doses for pressing out of the pack according to a therapeutic schedule. The container that is employed will depend on the exact dosage form involved, for example a conventional cardboard box would not generally be used to hold a liquid suspension. It is feasible that more than one container can be used together in a single package to market a single dosage form. For example, tablets may be contained in a bottle that is in turn contained within a box. Typically the kit includes directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (*e.g.*, oral, topical, transdermal and parenteral), are administered at different dosage intervals, or

when titration of the individual components of the combination is desired by the prescribing physician.



[0212] One particular example of a kit according to the present invention is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process recesses are formed in the plastic foil. The recesses have the size and shape of individual tablets or capsules to be packed or may have the size and shape to accommodate multiple tablets and/or capsules to be packed. Next, the tablets or capsules are placed in the recesses accordingly and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are individually sealed or collectively sealed, as desired, in the recesses between the plastic foil and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

[0213] Another specific embodiment of a kit is a dispenser designed to dispense the daily doses one at a time in the order of their intended use. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical counter that indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.


EXAMPLES

Example 1: Preparation of 2-[6-(3-amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile succinate (Compound I)


[0214] Compound I may be prepared by the follow synthetic route (Scheme 1)

**A. Preparation of 4-fluoro-2-methylbenzonitrile (Compound B)**

[0215] Compound **B** was prepared by refluxing a mixture of 2-bromo-5-fluoro-toluene (Compound **A**) (3.5 g, 18.5 mmol) and CuCN (2 g, 22 mmol) in DMF (100 mL) for 24 hours. The reaction was diluted with water and extracted with hexane. The organics were dried over MgSO₄ and the solvent removed to give product **B** (yield 60%). ¹H-NMR (400 MHz, CDCl₃): δ 7.60 (dd, J=5.6, 8.8 Hz, 1H), 6.93-7.06 (m, 2H), 2.55 (s, 3H).


B. Preparation of 2-bromomethyl-4-fluorobenzonitrile (Compound C)

[0216] Compound **C** was prepared by refluxing a mixture of 4-fluoro-2-methylbenzonitrile (Compound **B**) (2 g, 14.8 mmol), N-bromosuccinimide (NBS) (2.64 g, 15 mmol) and azo-bis-isobutyronitrile (AIBN) (100 mg) in CCl_4 under nitrogen for 2 hours. The reaction was cooled to room temperature. The solid was removed by filtration. The organic solution was concentrated to give the crude product the form of an oil, which was used in the next step without further purification. $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 7.68 (dd, J = 5.2, 8.4 Hz, 1H), 7.28 (dd, J = 2.4, 8.8 Hz, 1H), 7.12 (m, 1H), 4.6 (s, 2H).

C. Preparation of 2-(6-chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile (Compound D)


[0217] Compound **E** was prepared by stirring a mixture of crude 3-methyl-6-chlorouracil **D** (0.6 g, 3.8 mmol), 2-bromomethyl-4-fluorobenzonitrile (0.86 g, 4 mmol) and K_2CO_3 (0.5 g, 4 mmol) in DMSO (10 mL) at 60° C for 2 hours. The reaction was diluted with water and extracted with EtOAc. The organics were dried over MgSO_4 and the solvent removed. The residue was purified by column chromatography. 0.66 g of the product was obtained (yield: 60%). $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 7.73 (dd, J =7.2, 8.4Hz, 1H), 7.26 (d, J =4.0Hz, 1H), 7.11-7.17 (m, 1H), 6.94 (dd, J =2.0, 9.0 Hz, 1H), 6.034 (s, 2H), 3.39 (s, 3H). MS (ES) $[m+H]$ calc'd for $\text{C}_{13}\text{H}_9\text{ClFN}_3\text{O}_2$, 293.68; found 293.68.

D. Preparation of 2-(6-chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile (Compound F)



[0218] Compound **F** was prepared by mixing and stirring 2-(6-chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile (Compound **E**) (300 mg, 1.0 mmol), (*R*)-3-amino-piperidine dihydrochloride (266 mg, 1.5 mmol) and sodium bicarbonate (500 mg, 5.4 mmol) in a sealed tube in EtOH (3 mL) at 100 °C for 2 hrs. The final compound was obtained as trifluoroacetate (TFA) salt after HPLC purification. ¹H-NMR (400 MHz, CD₃OD): δ. 7.77-7.84 (m, 1H), 7.16-7.27 (m, 2H), 5.46 (s, 1H), 5.17-5.34 (ABq, 2H, J = 35.2, 15.6 Hz), 3.33-3.47 (m, 2H), 3.22 (s, 3H), 2.98-3.08 (m, 1H), 2.67-2.92 (m, 2H), 2.07-2.17 (m, 1H), 1.82-1.92 (m, 1H), 1.51-1.79 (m, 2H). MS (ES) [m+H] calc'd for C₁₈H₂₀FN₅O₂, 357.38; found, 357.38.

E. Preparation of Compound I: the succinic acid salt of 2-(6-Chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile



[0219] The TFA salt prepared in the above step (Example 1, Step D) was suspended in DCM, and then washed with saturated Na₂CO₃. The organic layer was dried and removed in vacuo. The benzonitrile product (approximately 10 mg) was dissolved in MeOH (1 mL) and to which succinic acid in THF (1.05 equivalents) was added. The solutions were allowed to stand for three days open to the air. If a precipitate formed, the solid was collected by filtration. If no solid formed, the mixture was concentrated in vacuo, and the succinate salt was obtained after removing the solvent. ¹H-NMR (400 MHz, CD₃OD): δ. 7.77-7.84 (m, 1H), 7.12-7.26 (m, 2H), 5.47 (s, 1H), 5.21-5.32 (ABq, 2H, J = 32.0, 16.0

Hz), 3.35-3.5 (m, 2H), 3.22 (s, 3H), 3.01-3.1 (m, 1H), 2.69-2.93 (m, 2H), 2.07-2.17 (m, 1H), 1.83-1.93 (m, 1H), 1.55-1.80 (m, 2H). MS (ES) [m+H] calc'd for C₁₈H₂₀FN₅O₂, 357.38; found, 357.38.

[0220] Compound I such prepared was found to be crystalline as determined by x-ray powder diffraction analysis (Figure 1). The crystal material was designated Form A.

Example 2: Approximate Solubility of Compound I in Different Solvents

[0221] Compound I prepared by the method described in Example 1 was used for the solubility study. Solvents and other reagents were of ACS or HPLC grade and were used as received.

[0222] A weighed sample (typically about 20-25 mg) of Compound I was treated at ambient temperature with aliquots (typically $\geq 50 \mu\text{L}$) of the test solvent with sonication between additions. Solvents were either reagent or HPLC grade. Solubility was estimated from the total volume of solvent used to obtain a clear solution, as determined by visual inspection. Solubility is expressed as “less than” when dissolution was not observed. If dissolution occurred after the last aliquot was added then solubility is expressed as “greater than or equal to” ($\geq$). The actual solubilities may be greater than determined, due to over addition of solvent (large aliquots) or slow dissolution rates. Approximate solubilities of Compound I are summarized in Table A. The solubility values have been rounded to the nearest whole number and are reported to the nearest mg/mL.

Table A. Approximate Solubilities of Compound I

Solvent	Solubility (mg/mL)^a
Acetone	2
Acetonitrile (ACN)	<1
Dichloromethane (DCM)	<1
Dimethyl Formamide (DMF)	68
1,4-Dioxane	<1
Ethanol (EtOH)	2
Ethyl Acetate (EtOAc)	<1
di-Ethyl ether	<1
Hexanes	<1
2-Propanol (IPA)	<1
Methanol (MeOH)	20
Tetrahydrofuran (THF)	<1
Toluene	<1
Trifluoroethanol (TFE)	>200
Water (H ₂ O)	51

Solvent	Solubility (mg/mL) ^a
ACN:H ₂ O (85:15)	101
EtOH:H ₂ O (95:5)	5
IPA:H ₂ O (88:12)	11

- a) Approximate solubilities are calculated based on the total solvent used to give a solution; actual solubilities may be greater because of the volume of the solvent portions utilized or a slow rate of dissolution. Solubilities are reported to the nearest mg/mL.

Example 3. Polymorph Screen

[0223] Compound I as prepared by the method described in Example 1 was used as the starting material for the polymorph screen. Solvents and other reagents were of ACS or HPLC grade and were used as received.

A. Sample Generation.

[0224] Solids for form identification were prepared via the following methods from Compound I.

1. Fast Evaporation (FE)

[0225] A solution of Compound I was prepared in test solvents. The sample was placed in the hood, uncovered, to evaporate under ambient conditions. The solids were analyzed by XRPD for form identification.

2. Slow Evaporation (SE)

[0226] A solution of Compound I was prepared in test solvents. The sample was placed in the hood, covered with foil rendered with pinholes, to evaporate under ambient conditions. The solids were analyzed by XRPD for form identification.

3. Room Temperature (RT) Slurries

[0227] An excess amount of Compound I was slurried in test solvent on a rotating wheel for approximately 5 or 7 days. The solids were typically collected by vacuum filtration, air dried in the hood, and analyzed by XRPD for form identification.

4. Elevated Temperature Slurries

[0228] Excess Compound I was slurried in test solvents at 47 °C on a shaker block for approximately 5 days. The solids were collected by vacuum filtering, dried in the hood, and then analyzed by XRPD for form identification.

5. Slow Cooling Crystallization (SC)

[0229] A saturated or near saturated solution of Compound I was prepared at elevated temperature. The samples were filtered through warmed 0.2 μm filters into warmed vials. The heat source was turned off and the samples slowly cooled to ambient temperature. If precipitation did not occur within a day the samples were placed in the refrigerator. The samples were transferred to a freezer if precipitation did not occur within several days. The solids were collected by decanting the solvent or vacuum filtration, dried in the hood and analyzed by XRPD for form identification.

6. Crash Cooling Crystallization (CC)

[0230] A saturated or near saturated solution of Compound I was prepared at elevated temperature. The samples were filtered through warmed 0.2 μm filters into warmed vials then rapidly cooled in an acetone/dry ice or ice bath. If precipitation did not occur after several minutes the samples were placed in the refrigerator or freezer. Solids were collected by decanting solvent or vacuum filtration, dried in the hood, and then analyzed by XRPD. Samples that did not precipitate under subambient conditions after several days were evaporated in the hood and analyzed by XRPD for form identification.

7. Solvent/Antisolvent Crystallization (S/AS)

[0231] A solution of Compound I was prepared in test solvent. A miscible antisolvent was added with a disposable pipette. Precipitate was collected by vacuum filtration or decanting solvent. The samples were stored under subambient conditions if precipitation did not occur. If solids were not observed after several days the samples were evaporated in the hood. Collected solids were analyzed by XRPD for form identification.

8. Relative Humidity (RH) Stressing Experiments

[0232] Samples of Compound I were placed uncovered in approximately 58%, 88%, and 97% relative humidity jars. The samples were stored in the jars for approximately 8 days. The solids were collected and analyzed by XRPD for form identification.

9. Lyophilization

[0233] Compound I was dissolved in water in a glass vial. The solution was frozen by swirling the vial in an acetone/dry ice bath. The frozen sample was placed on the lyophilizer until all of the frozen solvent was removed. The solids were collected and analyzed by XRPD for form identification.

10. Grinding Experiments

[0234] Aliquots of Compound I were ground manually with a mortar and pestle as a dry solid and a wet paste in water. The samples were ground for approximately three minutes. The solids were collected and analyzed by XRPD for form identification.

11. Dehydration Experiments

[0235] Hydrated samples of Compound I were dehydrated at ambient conditions (2 days) and in an ambient temperature vacuum oven (1 day). The solids were collected and analyzed by XRPD for form identification.

12. Vapor Stress Experiments

[0236] Amorphous Compound I was placed in acetone, ethanol, and water vapor chambers for up to eight days. The solids were collected and analyzed by XRPD for form identification.

B. Sample Characterization.

[0237] The following analytical techniques and combination thereof were used determine the physical properties of the solid phases prepared.

1. X-Ray Powder Diffraction (XRPD)

[0238] X-ray powder diffraction (XRPD) analyses were performed using an Inel XRG-3000 diffractometer equipped with a CPS (Curved Position Sensitive) detector with a 2θ (2θ) range of 120° . Real time data were collected using Cu-K α radiation starting at approximately $4^\circ 2\theta$ at a resolution of $0.03^\circ 2\theta$. The tube voltage and amperage were set to 40 kV and 30 mA, respectively. The pattern is displayed from 2.5 to $40^\circ 2\theta$. Samples were prepared for analysis by packing them into thin-walled glass capillaries. Each capillary was mounted onto a goniometer head that is motorized to permit spinning of the capillary during data acquisition. The samples were analyzed for approximately 5 minutes. Instrument calibration was performed using a silicon reference standard. Peak picking was performed using the automatic peak picking in the Shimadzu XRD-6000 Basic Process version 2.6. The files were converted to Shimadzu format before performing the peak picking analysis. Default parameters were used to select the peaks.

2. Thermogravimetric Analysis (TGA)

[0239] Thermogravimetric (TG) analyses were performed using a TA Instruments 2950 thermogravimetric analyzer. Each sample was placed in an aluminum sample pan and

inserted into the TG furnace. The furnace was first equilibrated at 25 °C, then heated under nitrogen at a rate of 10 °C/min, up to a final temperature of 350 °C. Nickel and Alumel™ were used as the calibration standards.

3. Differential Scanning Calorimetry (DSC)

[0240] Differential scanning calorimetry (DSC) was performed using a TA Instruments differential scanning calorimeter 2920. The sample was placed into an aluminum DSC pan, and the weight accurately recorded. The pan was covered with a lid and then crimped. The sample cell was equilibrated at 25 °C and heated under a nitrogen purge at a rate of 10 °C/min, up to a final temperature of 350 °C. Indium metal was used as the calibration standard. Reported temperatures are at the transition maxima.

[0241] For studies of the glass transition temperature (T_g) of the amorphous material, the sample cell was equilibrated at ambient temperature, then heated under nitrogen at a rate of 20 °C/min, up to 100 °C. The sample cell was then allowed to cool and equilibrate at -20 °C. It was again heated at a rate of 20 °C/min up to 100 °C and then cooled and equilibrated at -20 °C. The sample cell was then heated at 20 °C/min up to a final temperature of 350 °C. The T_g is reported from the onset point of the transition.

4. Hot Stage Microscopy.

[0242] Hot stage microscopy was performed using a Linkam hot stage (model FTIR 600) mounted on a Leica DM LP microscope. The samples were prepared between two cover glasses and observed using a 20× objective with crossed polarizers and first order compensator. Each sample was visually observed as the stage was heated. Images were captured using a SPOT Insight™ color digital camera with SPOT Software v. 3.5.8. The hot stage was calibrated using USP melting point standards.

5. Thermogravimetric-Infrared (TG-IR)

[0243] Thermogravimetric infrared (TG-IR) analyses were acquired on a TA Instruments thermogravimetric (TG) analyzer model 2050 interfaced to a Magna 560® Fourier transform infrared (FT-IR) spectrophotometer (Thermo Nicolet) equipped with an Ever-Glo mid/far IR source, a potassium bromide (KBr) beamsplitter, and a deuterated triglycine sulfate (DTGS) detector. The TG instrument was operated under a flow of helium at 90 and 10 cc/min for the purge and balance, respectively. Each sample was placed in a platinum sample pan, inserted into the TG furnace, accurately weighed by the instrument, and the furnace was heated from ambient temperature to 250 °C at a rate of 20

°C/min. The TG instrument was started first, immediately followed by the FT-IR instrument. Each IR spectrum represents 32 co-added scans collected at a spectral resolution of 4 cm^{-1} . A background scan was collected before the beginning of the experiment. Wavelength calibration was performed using polystyrene. The TG calibration standards were nickel and Alumel™. Volatiles were identified from a search of the High Resolution Nicolet TGA Vapor Phase spectral library.

6. Fourier Transform Infrared Spectroscopy (FT-IR)

[0244] Infrared spectra were acquired on a Magna-IR 560[®] or 860[®] Fourier transform infrared (FT-IR) spectrophotometer (Thermo Nicolet) equipped with an Ever-Glo mid/far IR source, an extended range potassium bromide (KBr) beamsplitter, and a deuterated triglycine sulfate (DTGS) detector. A diffuse reflectance accessory (the Collector™, Thermo Spectra-Tech) was used for sampling. Each spectrum represents 256 co-added scans collected at a spectral resolution of 4 cm^{-1} . Sample preparation consisted of physically mixing the sample with KBr and placing the sample into a 13-mm diameter cup. A background data set was acquired on a sample of KBr. A Log 1/R (R = reflectance) spectrum was acquired by taking a ratio of these two data sets against each other. Wavelength calibration was performed using polystyrene. Automatic peak picking was performed using Omnic version 7.2.

7. Fourier Transform Raman Spectroscopy (FT-Raman)

[0245] FT-Raman spectra were acquired on a Raman accessory module interfaced to a Magna 860[®] Fourier transform infrared (FT-IR) spectrophotometer (Thermo Nicolet). This module uses an excitation wavelength of 1064 nm and an indium gallium arsenide (InGaAs) detector. Approximately 0.5 W of Nd:YVO₄ laser power was used to irradiate the sample. The samples were prepared for analysis by placing the material in a glass tube and positioning the tube in a gold-coated tube holder in the accessory. A total of 256 sample scans were collected from at a spectral resolution of 4 cm^{-1} , using Happ-Genzel apodization. Wavelength calibration was performed using sulfur and cyclohexane. Automatic peak picking was performed using Omnic version 7.2.

8. Solid State Nuclear Magnetic Resonance Spectroscopy (¹³C-NMR)

[0246] The solid-state ¹³C cross polarization magic angle spinning (CP/MAS) NMR spectrum was acquired at ambient temperature on a Varian^{UNITY} INOVA-400 spectrometer (Larmor frequencies: ¹³C = 100.542 MHz, ¹H = 399.799 MHz). The sample was packed

into a 4 mm PENCIL type zirconia rotor and rotated at 12 kHz at the magic angle. The spectrum was acquired with phase modulated (SPINAL-64) high power ^1H decoupling during the acquisition time using a ^1H pulse width of 2.2 μs (90°), a ramped amplitude cross polarization contact time of 5 ms, a 30 ms acquisition time, a 10 second delay between scans, a spectral width of 45 kHz with 2700 data points, and 100 co-added scans. The free induction decay (FID) was processed using Varian VNMR 6.1C software with 32768 points and an exponential line broadening factor of 10 Hz to improve the signal-to-noise ratio. The first three data points of the FID were back predicted using the VNMR linear prediction algorithm to produce a flat baseline. The chemical shifts of the spectral peaks were externally referenced to the carbonyl carbon resonance of glycine at 176.5 ppm.

9. Solution Nuclear Magnetic Resonance Spectroscopy (^1H -NMR)

[0247] The solution ^1H NMR spectrum was acquired at ambient temperature with a Varian^{UNITY} INOVA-400 spectrometer at a ^1H Larmor frequency of 399.803 MHz. The sample was dissolved in methanol. The spectrum was acquired with a ^1H pulse width of 8.4 μs , a 2.50 second acquisition time, a 5 second delay between scans, a spectral width of 6400 Hz with 32000 data points, and 40 co-added scans. The free induction decay (FID) was processed using Varian VNMR 6.1C software with 65536 points and an exponential line broadening factor of 0.2 Hz to improve the signal-to-noise ratio. The spectrum was referenced to internal tetramethylsilane (TMS) at 0.0 ppm.

10. Moisture Sorption/Desorption Analysis

[0248] Moisture sorption/desorption data were collected on a VTI SGA-100 Vapor Sorption Analyzer. Sorption and desorption data were collected over a range of 5% to 95% relative humidity (RH) at 10% RH intervals under a nitrogen purge. Samples were not dried prior to analysis. Equilibrium criteria used for analysis were less than 0.0100% weight change in 5 minutes, with a maximum equilibration time of 3 hours if the weight criterion was not met. Data were not corrected for the initial moisture content of the samples. NaCl and PVP were used as calibration standards.

[0249] Crystallization experiments performed for the polymorph screen are summarized in Tables B and C. Table B summarizes the crystallization experiments in various solvent. Table C summarized the crystallization experiments in various solvents/antisolvents.

Table B. Crystallization Experiments of Compound I from Solvents

Solvent	Conditions^a	XRPD Form^b	Color/Morphology
Acetone	FE	A	white, aggregates of birefringent aciculars
		A	white + yellow, needles
	SE	A	light yellow, needles + blades in large rosettes
		-	light yellow, spherulite
	partial dissolution of the light yellow spherulite precipitated from acetone by SE	A	white, needles
	RT slurry, 7 days	A	white, needles
	SC (60 °C)	A	white, tiny needles
	CC	qty. too small	white, tiny needles/spherulites
ACN	FE	A-peaks, PO	white, needles/rosettes
	SE	A	white, needles
	RT slurry, 7 days	A	white, needles
	SC (60 °C)	A	white, blades/spherulites
DCM	RT slurry, 7 days	A	white, needles
1,4-Dioxane	FE	low crystallinity A	yellow, needles, glass
	RT slurry, 7 days	A	white, needles
	SC (60 °C)	A	white, needles
DMF	FE	A	off white, needles
EtOH	FE	A	off white, needles/rosettes + unknown morphology
		A	off white, needles, spherulites
	SE	A	white, needles/spherulites
	SC (60 °C)	A	off white, spherulites
	CC-evaporated	A	white, needles, spherulites
EtOAc	RT slurry, 7 days	A	white, needles
	SC (60 °C)	A	white, needles/spherulites
di-Ethyl ether	RT slurry, 7 days	A	white, needles
Hexanes	RT slurry, 7 days	A	white, needles
MeOH	FE	A	white, irregularly shaped, w/birefringent acicular regions
		A	off white, needles, spherulites
	SE	A	white, needles
	RT slurry, 5 days	A	white, tiny needles
	SC (60 °C)	A	yellow, spherulites of tiny needles

Solvent	Conditions ^a	XRPD Form ^b	Color/Morphology
	CC	A	white, needles
IPA	RT slurry, 7 days	A	white, needles
	SC (60 °C)	Pattern C	white, needles
		C - peaks	
		low crystallinity A	white, spherulites
		low crystallinity A+B	white, spherulites
		low crystallinity A+B	white, needles, spherulites
		low crystallinity A+B	white, unknown, spheres
		A	white, spheres
		low crystallinity material	white, needles + unknown
		low crystallinity material	white, unknown, spheres
		low crystallinity A+B	white, unknown
		low crystallinity A+B	white, needles, spheres
		low crystallinity material	white, needles + unknown
		low crystallinity A+B	white, needles
THF	RT slurry, 7 days	A	white, needles
	SC (60 °C)	low crystallinity A	white, tiny needles
Toluene	RT slurry, 7 days	A	white, needles
TFE	FE	A	off white, needles, glass
	SE	A	off white, needles
Water	FE	A	white, needles/rosettes
		A	white, off white edges, needles/spherulites
		A	white, brown edges, needles/spherulites
	SE	A	off white, some pink spots, needles
		A	white w/brown edges, needles
		A	white w/brown edges, needles
	RT slurry, 5 days	A	off white, needles
	SC (60 °C)	A	off white, needles/spherulites
Water	CC	A+B+peaks	white, needles/spherulites
	lyophilization	amorphous	-
ACN:H ₂ O (85:15)	FE	A	white, aggregates of birefringent aciculars
	SE	A	off white, needles/spherulites
EtOH:H ₂ O (95:5)	FE	A+1 peak	white + light yellow, needles/rosettes
	SE	A	white, needles

Solvent	Conditions ^a	XRPD Form ^b	Color/Morphology
	RT slurry, 7 days	A	white, needles
	elevated temp. slurry	A	white, light yellow, small sticky orange-ish spot, needles
	CC	A+B	white, blades, needles
	CC	A+B	white, blades, needles
		A+B	off white, blades/spherulites
IPA:H ₂ O (88:12)	FE	low crystallinity A+B	white, fine blades, aciculars, birefringent
	SE	A+B	white, pink spots, needles
		A+B	white, needles
	partial SE	A+B	white, needles
	RT slurry, 5 days	A	white, tiny needles
	SC (60 °C)	A+B	white, needles, spherulites
		A+B	white, needles
	CC	A+B	white, needles/spherulites
THF:H ₂ O (9:1)	RT slurry, 13 days	A+E+G peaks	white, needles
	elevated temp. slurry	A	light yellow, white, needles
	SC (60 °C)	A+B	white, needles/blades
		A+B	white needles, spherulites
		A+B	
		A+minor E	off white, needles
		A+F peaks	white, needles
	SC (60 °C), 70 °C, 1 day	A+F peaks	white, glass, needles
	damp 2333-34-03	A+E+G peaks	white, needles
	SC (60 °C), 70 °C, 3 days	A+F peaks	off white, glass + needles
-	Post MB of a sample prepared by CC from EtOH:H ₂ O (95:5)	A+B	-
	Post MB of a lyophilized sample from H ₂ O solution	low crystallinity A	-
	Post MB of a sample crystallized from H ₂ O/ACN	A+small quantity of E	-
	ground (dry) ~3 minutes	A	-
-	ground (wet) ~3 minutes	A	-
	Form A+B, under vacuum, overnight	A	off white, needles

Solvent	Conditions ^a	XRPD Form ^b	Color/Morphology
	Form A+B, at ambient, ~2 days	A	off white, needles

a) FE = fast evaporation; SE = slow evaporation; RT = room temperature; SC = slow cool;

CC = crash cool, MB = moisture sorption/desorption analysis

b) qty = quantity; PO = preferred orientation

Table C. Crystallization Experiments of Compound I in Various Solvent/Antisolvent

Solvent	Antisolvent	Form	Color/Morphology
DMF	ACN	A	off white, needles/spherulites
	toluene	A	white, needles/spherulites
	EtOAc	A	white, spherulites
	IPA	A	white, needles/spherulites
MeOH	ACN	A ^a	white + yellow, needles
	DCM	A ^a	white + yellow, needles
TFE	IPE	A	white, tiny spherulites
	ACN	A	off white, needles, spherulites
	IPA	A	white, tiny spherulites
	EtOAc	A	white, tiny spherulites
H ₂ O	ACN	D ^a	off white, needles
		A ^a	white, brown edges, needles/blades
		E ^a	off white, needles
		A ^a	white + light yellow, needles
		A ^a	white, off white at edges, needles/spherulites
		A ^a	white w/brown edges, needles
		A ^a	white w/brown edges, needles
		A ^a	off white, brown at edges, needles/spherulites
		A ^a	white, needles
		A ^a	white, needles
		A	white/off white, needles
	dioxane	A+B ^a	off white, needles
	THF	A ^a	light yellow solid/paste, needles

^a precipitated by evaporation of solvent

Example 4. Preparation of Form A+B

[0250] Approximately 40 mg of Compound I starting material was dissolved in 4 mL of isopropanol:water (88:12). The vial was covered with a piece of aluminum foil rendered with five pinholes for slow evaporation. Solids yielding Form A+B were recovered after 14 days.

Example 5. Preparation of Form C

[0251] A slurry of Compound I starting material (40 mg) in IPA (8 mL) was stirred on a hot plate set to 60 °C, 300 rpm for approximately 5 hours. The slurry was filtered through a warmed 0.2 µm nylon filter into a warmed vial, and the hot plate was turned off to slow cool the solution. The sample was stored at ambient temperature for approximately one day, then transferred to a refrigerator for approximately 3 days. A clear solution was observed. After 3 days in the refrigerator, the sample was transferred to a freezer for approximately 5 days. The solids were collected by vacuum filtration and dried in the hood.

Example 6. Preparation of Form D

[0252] A sample of Compound I starting material (40 mg) was dissolved in water (400 µl). The solution was filtered through a 0.2 µm nylon filter into a clean vial. Approximately half a disposable pipette full of acetonitrile was added to the water solution. No precipitation was observed. The sample was placed into a refrigerator for approximately 12 days. Precipitates were not observed. The solution was placed in the hood, uncovered, to evaporate. Solids yielding Form D were recovered after two days.

Example 7. Preparation of Form E

[0253] A sample of Compound I (40 mg) was dissolved in water (400 µl). The solution was filtered through a 0.2 µm nylon filter into a clean vial. Approximately ¾ a disposable pipette full of acetonitrile was added to the water solution. The sample was agitated by hand. No precipitation was observed. The solution was placed in the hood, uncovered, to evaporate. Solids were recovered after 10 days.

Example 8. Preparation of Form A+F

[0254] A sample of Compound I (51 mg) was dissolved in 800 µl of a THF:water (9:1) solution. The sample was stirred on a hot plate set at 60 °C, 300 rpm for approximately one hour. The hot plate was turned off to allow the clear solution to slow cool. After the sample reached room temperature, it was placed in the refrigerator to induce precipitation. Precipitates were observed after approximately two days. The solvent was decanted and the solids allowed to air dry.

Example 9. Preparation of Form A+E+G

[0255] A slurry of Compound I (79 mg) was prepared in 1 mL of THF:water (9:1). The sample was slurried at ambient conditions for 13 days. The solids were collected by filtration and air dried. The resulting solids yielded a mixture of Forms A+E+G.

Example 10. Preparation of Amorphous Form

[0256] A sample of Compound I (40 mg) was dissolved in 1000 µl of water. The solution was filtered through a 0.2 µm nylon filter into a clean vial then frozen in a dry ice/acetone bath. The vials were covered with a Kimwipe then placed on a lyophilizer overnight. The resulting solids yielded Amorphous Form.

Example 11. Characterization of Form A

[0257] Material prepared by the procedure of Example 1 was designated as Form A. The material was characterized by XRPD, TGA, DSC, hot stage microscopy, FT-IR, FT-Raman, ¹H NMR, and ¹³C NMR. The analyses were conducted according to the procedures outlined in Section B of Example 3.

[0258] The characteristic spectra and thermograms for Form A are reported in Figures 1-7. The characterization data are summarized in Table D.

Table D. Characterization Data of Form A of Compound I

Sample Source	Technique	Analysis/Result
Prepared by procedure of Example 1	XRPD	crystalline (Figure 1)
	TGA	insignificant weight loss from 25 °C to 165 °C (Figure 2)
	DSC	endotherm 195 °C (Figure 2)
	Hot stage microscopy	T=26 °C T=34 °C T=89 °C T=110 °C T=177 °C, melt onset T=184 °C, approximate melting point
	IR	Figure 4 A-D
	Raman	Figure 5 A-D
	¹ H NMR	consistent with chemical structure (Figure 6)
	¹³ C NMR	consistent with chemical structure (Figure 7)

T=temperature

Example 12. Characterization of Form A+B

[0259] Form A+B material was prepared according to Example 4.

[0260] The material was characterized by XRPD, TGA, DSC, hot stage microscopy, TG-IR and moisture sorption/desorption analysis. The analyses were conducted according to the procedures outlined in Section B of Example 3.

[0261] The characteristic spectra and thermograms for Form A+B are reported in Figures 8-13. The characterization data of Form A+B are summarized in Table E.

Table E. Characterization Data of Form A+B

Sample Source	Technique	Analysis/Result
Crystallized from IPA:H ₂ O (88:12), by slow evaporation (SE)	XRPD	Crystalline (Figure 8)
	TGA	2.0% weight loss from 25 °C to 98 °C 0.7% weight loss from 98 °C to 175 °C (Figure 9)
	DSC	broad endotherm 100 °C small endotherm 138 °C, 163 °C endotherm 193 °C (Figure 12)
	Hot Stage	T=26 °C, birefringence with extinction T=99 °C, no changes observed T=145 °C, no changes observed T=165 °C, no changes observed T=179 °C, some particles appear to be melting T=185 °C, particles appear to be melting T=188 °C, possible end of melt, approximate melting point T=233 °C, sample appears brown, suggesting decomposition T=302 °C, sample brown, cooled to approximately 30 °C, no recrystallization
	TG-IR	2.4% weight loss from 21 °C to 105 °C and water by IR (Figures 10-11)
Crystallized from EtOH:H ₂ O (95:5) by crash cooling (CC)	MB	0.7% weight loss on equilibration at 5% RH 1.2% weight gain from 5-95% RH 1.2% weight lost from 95-5% RH (Figure 13)

T=temperature, RH=relative humidity, and MB = moisture sorption/desorption analysis

Example 13. Characterization of Form C

[0262] Form C material was prepared according to Example 5. The material was characterized by XRPD. The XRPD spectrum of Form C is reported in Figure 14.

Example 14. Characterization of Form D

[0263] Form D material was prepared according to Example 6.

[0264] The material was characterized by ¹H NMR, XRPD, TGA, DSC, TG-IR, and hot stage microscopy. The characterization analyses were conducted according to the procedures outlined in Section B of Example 3. The characteristic spectra and thermograms for Form D are reported in Figures 15-19. The characterization data of Form D are summarized in Table F.

Table F. Characterization Data of Form D

Sample Source	Technique	Analysis/Result
H ₂ O/ACN evaporated	XRPD	crystalline (Figure 15)
	TGA	~8% weight lost on equilibration, and 21.9% weight loss from 25 °C to 85 °C (Figure 16)
	DSC	broad, noisy endotherm at 88 °C, endotherm at 107 °C and 192 °C (Figure 19)
	TG-IR	24.0% weight loss when heated from 20 °C to 100 °C due to evaporation of water (Figures 17-18)
	Hot stage microscopy	T = 25 °C, birefringence with extinction T = 84 °C, no changes T = 90 °C, no changes T = 115 °C, no changes T = 184 °C, small particles appear to be melting T = 188 °C, melt appears to continue T = 192 °C, approximate end of melt T = 218 °C, condensation on the cover slip T = 230 °C, melt appears to have evaporated and condensed on the cover slip. Cooled to ambient, no recrystallization
	¹ H NMR	consistent with chemical structure

T=temperature, MB = moisture sorption/desorption analysis

Example 15. Characterization of Form E

[0265] Form E material was prepared according to Example 7.

[0266] The material was characterized by ¹H NMR, XRPD, TGA, DSC, TG-IR, and hot stage microscopy, moisture sorption/desorption analysis (m s/des). The characterization analyses were conducted according to the procedures outlined in Section B of Example 3.

[0267] The characteristic spectra and thermograms for Form D are reported in Figures 20-23. The moisture sorption and desorption isotherm of Form E is reported in Figure 23. The characterization data of Form E are summarized in Table G.

Table G. Characterization Data of Form E

Sample Source	Technique	Analysis/Result
H ₂ O/ACN evaporated	XRPD	Crystalline (Figure 20)
	TGA	2.6% weight loss from 25 °C to 85 °C (Figure 21)
	DSC	small endotherms at 59 °C, 75 °C; forked endotherms at 107 °C, 110 °C, 114 °C; and endotherm at 192 °C (Figure 22)
	Hot stage microscopy	T=26 °C, birefringence with extinction T=59 °C, changes in birefringence T=81 °C, changes in birefringence T=124 °C, no changes, refocused T=169 °C, needles appear to be forming T=179 °C, more needles, possible melt onset T=183 °C, continued melt T=185 °C, approximate end of melt T=208 °C, sample appears brown, decomposition T=250 °C, sample appears brown
	MB	1.3% weight loss on equilibration, 5.4% weight gain from 5-95% RH, and 5.5% weight lost from 95-5% RH (Figure 23).
	¹ H NMR	consistent with chemical structure

T=temperature, RH=relative humidity, MB = moisture sorption/desorption analysis

Example 16. Characterization of Form A+F

[0268] Form A+F material was prepared according to Example 8. The material was characterized by XRPD. The resulted XRPD spectrum of Form A+F is reported in Figure 24.

Example 17. Characterization of Form G

[0269] Form A+E+G material was prepared according to Example 9. The material was characterized by XRPD. The resulted XRPD spectrum of Form A+E+G is reported in Figure 25.

Example 18. Characterization of Amorphous Form

[0270] The amorphous form of Compound I was prepared by according to Example 10.

[0271] Amorphous Form material was characterized by XRPD TGA, DSC, hot stage microscopy, and moisture sorption/desorption analysis. The glass transition temperature of Form A was evaluated using the procedure outlined in Example 3, Section B-3. The material resulted after the moisture sorption/desorption analysis was characterized by XRPD.

[0272] The characteristic spectra and thermograms of Amorphous Form are reported in Figures 26-29. The moisture sorption and desorption isotherm of Amorphous Form is reported in Figure 30. The DSC thermogram for the determination of glass transition temperature of Amorphous Form is reported in Figure 31. The characterization data of Amorphous Form is summarized in Table H below.

Table H. Characterization Data of Amorphous Form

Sample Source	Technique	Analysis/Result
Lyophilized from Compound I in water	XRPD	Amorphous (Figure 26)
	TGA	1.8% weight loss from 25 °C to 95 °C (Figure 27)
	DSC	slightly concave baseline to exotherm 130 °C endotherm 194 °C (Figure 28)
		approximate glass transition at 82 °C, exotherm at 138 °C and endotherm 199 °C (Figure 29)
	Hot stage microscopy	T=25 °C, some particles appear birefringent with extinction T=100 °C no changes T=116 °C, appears to be recrystallizing T=131 °C, appears to be recrystallizing T=145 °C, no changes T=174 °C, possible start of melt T=185 °C, continued melt T=188 °C, possible end of melt T=207 °C, no change
	MB	1.0% weight lost on equilibration at 5% RH, 4.4% weight gain from 5-95% RH, and 4.7% weight lost from 95-5% RH (Figure 30)
	Post MB XRPD	Crystalline, Form A (Figure 31)

T=temperature, RH=relative humidity, MB = moisture sorption/desorption analysis

Example 19: Relative Humidity Stressing Experiments

[0273] Aliquots of Form A of Compound I were stored under approximately 50% and 97% relative humidity for 29 days. An aliquot of a mixture of Form A+B was stressed at 88% relative humidity for 23 days. Amorphous Form was stressed at 58% and 88% relative humidity for 23 days. The solids remaining were characterized by XRPD and determined to be Form A. The result is summarized in Table I.

Table I. Relative Humidity Stressing Experiments of Polymorphic Form A and Form A+B of Compound I

% RH	Initial Form	Days	Form
97	A	29	A
58	A	29	A
88	A+B	23	A
58	amorphous	23	A
88	amorphous	23	A

Example 20: Dehydration Experiment

[0274] Aliquots of Form A+B of Compound I were stored at ambient condition for two days and in an ambient temperature vacuum over for one day. The solids were collected and analyzed by XRPD and confirmed to be Form A.

Example 21: Slurry Interconversion Studies

[0275] Samples of mixture of Form A and Form B of Compound I were slurried in water and THF:water (9:1), for up to 13 days. The result solid materials isolated were characterized by X-Ray powder diffraction and determined to be either Form A or Form A+B. The result is summarized in Table J.

Table J. Slurry Experiments of Form A+B of Compound I

Solvent	Initial Forms	Days (approximate)	Final Form
Water	A+B	went into solution, evaporated	A
THF:H ₂ O (9:1)	A+B	13	A+B

Example 22: Vapor Stress of Amorphous Solid of Compound I

[0276] Solids of Amorphous Form of Compound I were stressed in acetone and ethanol vapor stress chambers for 4 days. Other aliquots of the solids were stressed in a water vapor chamber for 4 days and 8 days. The remaining solids were analyzed by XRPD. The amorphous solids convert to Form A after exposure in acetone and ethanol and convert to a mixture of Form A and Form D after exposure to water vapor. The results of these experiments are summarized in Table K.

Table K. Vapor Stress of Amorphous Solid of Compound I

Solvent	Days	Color/morphology	Form
Acetone	4	off white, irregular	A
EtOH	4	off white, irregular	A
Water	4	off white, irregular, almost liquid/drops	A+D
	8	white, irregular, appear "damp"	low crystallinity

Example 23: Moisture Sorption/Desorption Study on Form A+B.

[0277] Moisture sorption and desorption study was conducted on a sample of Form A+B. The sample was prepared by crashed cooling crystallization (Example 3, section A.6) of the polymorphic forms from a ethanol:water (95:5) solution of Compound 1. The moisture sorption and desorption study was conducted according to the procedures outlined in Example 3, section B.10. The data collected is plotted in Figure 13 and summarized in Table L.

Table L. Moisture Sorption/Desorption of Form A+B

Elapsed Time min	Weight mg	Weight % chg	Samp Temp °C	Samp RH %
0.1	3.184	0.000	25.03	62.77
91.5	3.163	-0.665	25.05	5.14
137.8	3.178	-0.184	25.04	14.88
165.5	3.184	0.004	25.07	25.03
179.7	3.187	0.098	25.06	34.82
196.2	3.191	0.208	25.06	44.88
208.7	3.193	0.284	25.06	54.95
219.7	3.195	0.347	25.08	65.00
231.9	3.197	0.406	25.05	75.01
243.8	3.199	0.466	25.09	84.92
256.6	3.202	0.563	25.08	94.67
265.2	3.200	0.510	25.08	85.01

Elapsed Time min	Weight mg	Weight % chg	Samp Temp °C	Samp RH %
274.2	3.199	0.456	25.08	75.35
285.2	3.197	0.406	25.04	64.94
296.5	3.195	0.353	25.07	55.02
307.3	3.194	0.299	25.07	45.07
319.3	3.192	0.237	25.06	34.92
338.5	3.189	0.149	25.09	25.11
359.7	3.185	0.020	25.09	14.96
445.9	3.163	-0.658	25.10	4.83

Example 24: Moisture Sorption/Desorption Study on Form E.

[0278] Moisture sorption and desorption study was conducted on a sample of Form E. The sample was prepared by solvent/antisolvent crystallization (Example 3, section A.7) by the addition of acetonitrile to Compound I in water. The moisture sorption and desorption study was conducted according to the procedures outlined in Example 3, section B.10. The data collected is plotted in Figure 23 and summarized in Table M.

Table M. Moisture Sorption/Desorption of Form E

Elapsed Time min	Weight mg	Weight % chg	Samp Temp °C	Samp RH %
0.1	3.704	0.000	24.77	35.22
138.3	3.657	-1.268	24.77	5.23
154.5	3.660	-1.187	24.78	14.89
170.5	3.664	-1.074	24.78	24.87
184.5	3.669	-0.947	24.78	34.83
203.4	3.676	-0.755	24.78	44.87
253.5	3.695	-0.223	24.79	55.04
281.1	3.716	0.325	24.78	65.05
301.2	3.726	0.608	24.78	74.87
322.2	3.756	1.410	24.79	84.67
415.8	3.860	4.216	24.78	94.73
452.4	3.764	1.632	24.77	85.18
471.9	3.739	0.949	24.78	75.25
487.4	3.727	0.616	24.78	65.16
499.6	3.718	0.398	24.78	55.05
512.2	3.713	0.236	24.78	45.03
526.0	3.708	0.112	24.79	35.13
543.5	3.703	-0.013	24.79	25.02
557.2	3.700	-0.102	24.79	14.94
730.6	3.655	-1.322	24.77	5.03

Example 25: Moisture Sorption/Desorption Study of Amorphous Form.


[0279] Moisture sorption and desorption study was conducted on a sample of Amorphous Form. The sample was prepared by lyophilization of a solution of Compound I in water (Example 3, section A.9). The moisture sorption and desorption study was conducted according to the procedures outlined in Example 3, section B.10. The data collected is plotted in Figure 29 and summarized in Table N.

Table N. Moisture Sorption/Desorption of Amorphous Form

Elapsed Time min	Weight mg	Weight % chg	Samp Temp °C	Samp RH %
0.1	2.720	0.000	25.04	41.98
84.0	2.694	-0.975	25.06	5.09
113.6	2.704	-0.589	25.06	14.98
158.4	2.716	-0.166	25.05	24.92
277.5	2.743	0.841	25.04	34.90
392.7	2.785	2.393	25.03	45.05
470.8	2.844	4.540	25.02	54.99
508.0	2.910	6.988	25.02	64.90
654.6	2.732	0.437	25.02	74.82
703.8	2.745	0.922	25.03	84.78
806.9	2.813	3.400	25.02	94.59
877.9	2.748	1.010	25.02	85.31
957.5	2.725	0.176	25.02	74.92
1039.8	2.713	-0.280	25.03	65.07
1092.3	2.705	-0.545	25.04	55.11
1144.4	2.699	-0.776	25.05	45.07
1179.2	2.695	-0.912	25.05	35.06
1210.7	2.692	-1.026	25.05	25.13
1243.6	2.689	-1.148	25.04	15.06
1278.0	2.686	-1.269	25.05	4.97


What is claimed is:

1. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form A characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern (CuK α) comprising diffraction peaks at about 11.31, 11.91, and 22.32 degree 2-theta ($^{\circ}2\theta$);
 - (ii) a Raman spectrum comprising absorption bands expressed in reciprocal wavelengths at about 2954, 2935, 2225, 1698, 1659, and 1607 cm^{-1} ;
 - (iii) an IR spectrum comprising absorption bands expressed in reciprocal wavelengths at about 3141, 2953, 2934, 2266, 1699, 1657, 1450, and 1206 cm^{-1} ; and
 - (iv) a differential scanning calorimetry spectrum having an endotherm centered at about 193 $^{\circ}\text{C}$.
2. A composition comprising Compound I having the formula




wherein a portion of Compound I is present as polymorphic Form A characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern (CuK α) comprising diffraction peaks at about 11.31, 11.91, 19.26, 21.04, and 22.32 $^{\circ}2\theta$;
- (ii) a Raman spectrum comprising absorption bands expressed in reciprocal wavelengths at about 3068, 2954, 2935, 2225, 1698, 1659, 1607, 1586, 1223, 1180, 901, 780, 751, 669, and 516 cm^{-1} ;

- (iii) an IR spectrum comprising absorption bands expressed in reciprocal wavelengths at about 3141, 2953, 2934, 2266, 2225, 1699, 1657, 1450, 1206, 886, 760, 685, 594, and 516 cm^{-1} ; and
 - (iv) a differential scanning calorimetry spectrum having an endotherm centered at about $193\text{ }^{\circ}\text{C}$.
3. The composition according to either claim 1 or claim 2, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form A.
4. The composition according to either claim 1 or claim 2, wherein greater than 1% of Compound I (by weight) is present in the composition as Form A.
5. The composition according to either claim 1 or claim 2, wherein greater than 50% of Compound I (by weight) is present in the composition as Form A.
6. The composition according to either claim 1 or claim 2, wherein greater than 90% of Compound I (by weight) is present in the composition as Form A.
7. A method for preparing a composition of Compound I according to any one of claims 1-6 comprising crystallizing Compound I from a solvent system selected from the group consisting of: (i) acetone, (ii) acetonitrile, (iii) dichloromethane, (iv) 1,4-dioxane, (v) dimethylformamide, (vi) methanol, (vii) ethylacetate, (viii) diethylether, (ix) hexane; (x) methanol, (xi) isopropanol, (xii) tetrahydrofuran, (xiii) toluene, (xiv) trifluoroethanol, (xv) water, (xvi) acetonitrile:water (85:15), (xvii) ethanol:water (95:5), (xviii) isopropanol:water (88:12), and (xix) tetrahydrofuran:water (9:1).
8. A method for preparing a composition of Compound I according to any one of claims 1-6 comprising adding a miscible antisolvent to Compound I dissolved in a solvent, wherein the solvent/antisolvent are selected from the group consisting of (i) dimethylformamide/acetonitrile, (ii) dimethylformamide/toluene, (iii) dimethylformamide/ethylacetate, (iv) dimethylformamide/isopropanol, (v) methanol/acetonitrile, (vi) methanol/dichloromethane, (vii) trifluoroethanol/isopropanol, (viii) trifluoroethanol/ethylacetate, (ix) water/acetonitrile, and (x) water/tetrahydrofuran.


9. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form B characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern (CuK α) comprising diffraction peaks at about 12.5, 18.83 and 24.46 $^{\circ}2\theta$; and
 - (ii) a differential scanning calorimetry spectrum comprising a broad endotherm at about 100 $^{\circ}\text{C}$, and two other endotherms at about 138 $^{\circ}\text{C}$ and about 163 $^{\circ}\text{C}$.
10. The composition according to claim 9, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form B.
11. The composition according to claim 9, wherein greater than 1% of Compound I (by weight) is present in the composition as Form B.
12. The composition according to claim 9, wherein greater than 50% of Compound I (by weight) is present in the composition as Form B.
13. The composition according to claim 9, wherein greater than 90% of Compound I (by weight) is present in the composition as Form B.
14. A method for preparing a composition of Compound I according to any one of claims 9-13 comprising crystallizing Compound I from a solvent selected from the group consisting of: (i) isopropanol, (ii) ethanol and water (95:5), (iii) isopropanol and water (88:12), (iv) tetrahydrofuran and water (9:1), and (v) water.
15. A method for preparing a composition of Compound I according to any one of claims 9-13 comprising adding dioxane to Compound I dissolved in water.

16. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form C characterized by an X-ray powder diffraction pattern (CuK α) comprising diffraction peaks at about 5.44 and 6.07 $^{\circ}2\theta$.

17. The composition according to claim 16, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form C.


18. The composition according to claim 16, wherein greater than 1% of Compound I (by weight) is present in the composition as Form C.

19. The composition according to claim 16, wherein greater than 50% of Compound I (by weight) is present in the composition as Form C.

20. The composition according to claim 16, wherein greater than 90% of Compound I (by weight) is present in the composition as Form C.

21. A method for preparing a composition of Compound I according to any one of claims 16-20 comprising crystallizing Compound I from an isopropanol solution of Compound I.


22. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form D characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern (CuK α) comprising one diffraction peak at about 24.33 °2 θ , and four other diffraction peaks selected from the group consisting of peaks at about 12.19, 16.71, 22.08, 22.88, and 23.27 °2 θ ; and
- (ii) a differential scanning calorimetry spectrum comprising a broad endotherm centered at about 88 °C and two other endotherms at about 107 °C and 192 °C.

23. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form D characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern (CuK α) comprising one diffraction peak at about 24.33 °2 θ , and four other diffraction peaks selected from the group consisting of peaks at about 12.19, 14.04, 16.71, 17.75, 18.86, 19.96, 22.08, 22.88, 23.27, 25.02, 25.49, 26.03, and 27.99 °2 θ ; and
- (ii) a differential scanning calorimetry spectrum having a broad endotherm centered at about 88 °C, and two others endotherms at about 107 °C and about 192 °C.

24. The composition according to either claim 22 or claim 23, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form D.


25. The composition according to either claim 22 or claim 23, wherein greater than 1% of Compound I (by weight) is present in the composition as Form D.

26. The composition according to either claim 22 or claim 23, wherein greater than 50% of Compound I (by weight) is present in the composition as Form D.

27. The composition according to either claim 22 or claim 23, wherein greater than 90% of Compound I (by weight) is present in the composition as Form D.

28. A method for preparing a composition of Compound I according to anyone of claims 22-27 comprising adding acetonitrile to Compound I dissolved in water.


29. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form E characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern ($\text{CuK}\alpha$) comprising diffraction peaks at about 13.10, 13.94, 17.15, and 21.27 $^{\circ}2\theta$; and
- (ii) a differential scanning calorimetry spectrum comprising a forked endotherm that comprises peaks at about 107 $^{\circ}\text{C}$, about 110 $^{\circ}\text{C}$ and about 114 $^{\circ}\text{C}$.

30. A composition comprising Compound I having the formula




wherein a portion of Compound I is present as polymorphic Form E characterized by one or more physical properties selected from the group consisting of:

- (i) an X-ray powder diffraction pattern ($\text{CuK}\alpha$) comprising two diffraction peaks at about 21.27 and 17.15 $^{\circ}2\theta$, and three diffraction peaks selected

- from the group consisting of peaks at about 11.90, 12.66, 13.10, 13.59, 13.94, 17.54, 22.03, 22.61, 24.06, 24.70, 26.31, 27.34, and 31.10 °2 θ ; and
- (ii) a differential scanning calorimetry spectrum comprising a forked endotherm that comprises peaks at about 107 °C, about 110 °C and about 114 °C.


31. The composition according to either claim 29 or claim 30, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form E.
32. The composition according to either claim 29 or claim 30, wherein greater than 1% of Compound I (by weight) is present in the composition as Form E.
33. The composition according to either claim 29 or claim 30, wherein greater than 50% of Compound I (by weight) is present in the composition as Form E.
34. The composition according to either claim 29 or claim 30, wherein greater than 90% of Compound I (by weight) is present in the composition as Form E.
35. A method for preparing a composition of Compound I according to anyone of claims 29-34 comprising: adding acetonitrile to Compound I dissolved in water.
36. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as polymorphic Form F characterized by an X-ray powder diffraction pattern (CuK α) comprising diffraction peaks at about 12.39, 20.63, 26.03, and 30.05 °2 θ .

37. The composition according to claim 36, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form F.

38. The composition according to claim 36, wherein greater than 1% of Compound I (by weight) is present in the composition as Form F.
39. The composition according to claim 36, wherein greater than 50% of Compound I (by weight) is present in the composition as Form F.
40. The composition according to claim 36, wherein greater than 90% of Compound I (by weight) is present in the composition as Form F.
41. A method of preparing a composition of Compound I according to any one of claims 36-40 comprising: crystallizing Compound I from tetrahydrofuran and water solution of Compound I.
42. A composition comprising Compound I having the formula




wherein a portion of Compound I is present as polymorphic Form G characterized by an X-ray powder diffraction pattern (CuK α) comprising five or more diffraction peaks selected from the group consisting of diffraction peaks at about 13.22, 14.23, 18.62, 19.77, 24.36, 25.06, and 30.71 $^{\circ}2\theta$.

43. The composition according to claim 42, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Form G.
44. The composition according to claim 42, wherein greater than 1% of Compound I (by weight) is present in the composition as Form G.
45. The composition according to claim 42, wherein greater than 50% of Compound I (by weight) is present in the composition as Form G.

46. The composition according to claim 42, wherein greater than 90% of Compound I (by weight) is present in the composition as Form G.

47. A method for preparing a composition of Compound I according to any one of claims 42-46 comprising: crystallizing Compound I from a tetrahydrofuran and water solution of Compound I.

48. A composition comprising Compound I having the formula



wherein a portion of Compound I is present as Amorphous Form, characterized by one or more physical properties selected from the group consisting of:

- (i) an XRPD diffraction pattern (CuK α) characterized by a broad halo with no discernable diffraction peak; and
- (ii) a glass transition temperature at about 82 °C and an exotherm at about 138 °C.

49. The composition according to claim 48, wherein greater than 0.1% of Compound I (by weight) is present in the composition as Amorphous Form.

50. The composition according to claim 48, wherein greater than 1% of Compound I (by weight) is present in the composition as Amorphous Form.

51. The composition according to claim 48, wherein greater than 50% of Compound I (by weight) is present in the composition as Amorphous Form.

52. The composition according to claim 48, wherein greater than 90% of Compound I (by weight) is present in the composition as Amorphous Form.

53. A method for preparing a composition of Compound I according to any one of claims 48-52 comprising: lyophilizing a water solution of Compound I.

54. A pharmaceutical composition comprising a composition of Compound I according to any one of claims 1-6, 9-13, 16-20, 22-27, 29-34, 36-40, 42-46, and 48-52; and one or more pharmaceutical carriers.

55. The pharmaceutical composition according to claim 54 which is adapted for administration via a route selected from the group consisting of oral, parenteral, topical, transdermal, and pulmonary.

56. The pharmaceutical composition according to claim 54 which is in an oral dosage form selected from the group consisting of pills, tablets, capsules, emulsions, suspensions, microsuspensions, wafers, sprinkles, chewing gum, powders, lyophilized powders, granules, and troches.

57. The pharmaceutical composition according to claim 54, wherein the polymorphic form of the Compound I is at least partially preserved for a period of time following administration.

58. A kit comprising:
a pharmaceutical composition according to any one of claims 54-57; and
instructions which comprise one or more forms of information selected from the group consisting of indicating a disease state for which the composition is to be administered, storage information for the composition, dosing information and instructions regarding how to administer the composition.

59. The kit according to claim 58, wherein the kit comprises the composition in a multiple dose form.

60. An article of manufacture comprising:
a composition according to any one of claims 1-6, 9-13, 16-20, 22-27, 29-34, 36-40, 42-46, and 48-52; and
packaging materials.

61. The article of manufacture according to claim 60, wherein the packaging material comprises a container for housing the composition.

62. The article of manufacture according to claim 61, wherein the container comprises a label indicating one or more members of the group consisting of a disease state for which the composition is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition.

63. The article of manufacture according to claim 60, wherein the composition is in a multiple dose form.

64. Use of a composition according to any one of claims 1-6, 9-13, 16-20, 22-27, 29-34, 36-40, 42-46, and 48-52 in the manufacture of a medicament for the inhibition of dipeptidyl peptidases.

65. Use of a composition according to any one of claims 1-6, 9-13, 16-20, 22-27, 29-34, 36-40, 42-46, and 48-52 in the manufacture of a medicament for the treatment of a disease state for which dipeptidyl peptidases possess activity that contributes to the pathology and/or symptomology of the disease state.

66. The medicament according to claim 65, wherein the disease state is selected from the group consisting of type I diabetes, type II diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation and obesity; autoimmune diseases such as inflammatory bowel disease, multiple sclerosis, psoriasis, and rheumatoid arthritis; AIDS; and cancers.

67. The medicament according to claim 65, wherein the disease state is selected from the group consisting of type II diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation and obesity.

68. The medicament according to claim 65, wherein the disease state is type II diabetes mellitus.


FIGURE 1


FIGURE 2


FIGURE 3


FIGURE 4A


FIGURE 4B


FIGURE 4C


FIGURE 4D


FIGURE 5A


FIGURE 5B


FIGURE 5C


FIGURE 5D


FIGURE 6


FIGURE 7


FIGURE 8


FIGURE 9


FIGURE 10


FIGURE 11


FIGURE 12


FIGURE 13


FIGURE 14


FIGURE 15


FIGURE 16


FIGURE 17


FIGURE 18


FIGURE 19


FIGURE 20


FIGURE 21

FIGURE 22

DSC





FIGURE 23


FIGURE 24


FIGURE 25


FIGURE 26


FIGURE 27

FIGURE 28

DSC





FIGURE 29


FIGURE 30

FIGURE 31