

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 April 2010 (15.04.2010)

(10) International Publication Number
WO 2010/042887 A2

PCT

(51) **International Patent Classification:**
C07D 403/10 (2006.01) C07F 5/02 (2006.01)
C07F 3/06 (2006.01)

(74) **Agent:** DELANEY, Karoline, A.; Knobbe, Martens, Olson & Bear, LLP, 2040 Main Street, 14th Floor, Irvine, CA 92614 (US).

(21) **International Application Number:**
PCT/US2009/060267

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) **International Filing Date:**
9 October 2009 (09.10.2009)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
61/104,469 10 October 2008 (10.10.2008) US

(71) **Applicant** (for all designated States except US): **TRIUS THERAPEUTICS** [US/US]; 6310 Nancy Ridge Dr., Suite 101, San Diego, CA 92121 (US).

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **COSTELLO, Carrie, A.** [US/US]; 14148 Bahama Cove, Troy, NY (US). **SIMSON, Jaqueline, A.** [US/US]; Cohoes, New York, NY (US). **DUGUID, Robert, J.** [US/US]; 6310 Nancy Ridge Dr., Suite 101, Glenmont, NY (US). **PHILLIPSON, Douglas** [-/US]; 14148 Bahama Cove, Del Mar, CA 92014 (US).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))



WO 2010/042887 A2

(54) **Title:** METHODS FOR PREPARING OXAZOLIDINONES AND COMPOSITIONS CONTAINING THEM

(57) **Abstract:** Methods of preparing a class of oxazolidinones useful to impede bacterial growth are disclosed.

METHODS FOR PREPARING OXAZOLIDINONES AKD COMPOSITIONS
CONTAINING THEM

BACKGROUND OF THE INVENTION


[0001] Oxazolidinones as a chemical class find widespread use as pharmaceutical agents for the therapy and prophylaxis of such medical conditions as bacterial infections and atherosclerosis. The utility of these compounds has spurred efforts to find efficient routes to synthesize them, such as the copper-catalyzed cross coupling disclosed in US 20070049759. US 20070155798, which is hereby incorporated by reference in its entirety, recently disclosed potentially anti-bacterial oxazolidinones that feature substituted pyridinyl phenyl moieties. These moieties were initially incorporated by synthetic routes involving tin-based couplings, which because of the toxicity of any residual tin compounds is not desirable for pharmaceutical use. Accordingly, a need exists for synthetic routes to substituted (pyridinyl)phenyl oxazolidinones that does not involve use of tin reagents.

Field of the Invention

[0002] Novel methods are useful in the preparation of oxazolidinone-containing compounds.

SUMMARY OF THE INVENTION

[0003] A method of synthesizing a compound of the structure




wherein

R is H,


R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom,

comprises treating a compound having the structure




wherein R^2 is selected from the group consisting of optionally substituted benzyl and optionally substituted C_1-C_6 alkyl, with a strong base or an organolithium salt and then addition of glycidyl butyrate to the resulting anion under conditions to make




[0004] In some aspects, the treating step is performed in the presence of a facilitating compound, such as 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*i*)-pyrimidinone.

[0005] In some embodiments, the method includes an additional step comprising reacting




with $POCl_3$, $POCl(OBn)_2$, or $P(N-iPr)_2(O-tBu)_2$ under conditions form




wherein R' is $PO(OH)_2$.

[0006] The method may also comprise treating the compound of the structure




where R' is PO(OH)₂ with a base under conditions to form the compound of the structure




wherein R'' is a pharmaceutically acceptable salt of PO(OH)₂. In some aspects, the base is a sodium-containing base. In some aspects, R'' is PO₃N_a₂


[0007] A separate method of making an intermediate, or an additional step before the steps above, comprises coupling a first intermediate of the structure



wherein X is a leaving group such as selected from the group consisting of Cl, Br, I, and trifluoromethanesulfonate, with a second intermediate of the structure




wherein Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted C₁-C₆ alcohols and dihydropyridine alcohols, and wherein R³ and R⁴ together may form a ring, under conditions to produce the compound of the structure




[0008] In some aspects, the coupling is carried out in the presence of a palladium complex such as phosphine ligand bound to palladium, for example, dichlorobis(triphenylphosphine)palladium(II), tetrakis(triphenylphosphine)palladium(0), or $\text{Pd}_2(\text{dba})_3$

[0009] A separate method of making an intermediate, or an additional step before the coupling step above, comprises


a) treating an aryl halide of structure 5a




wherein X^1 is leaving group, with a strong base such as n-butyl lithium and then reacting a resulting anion with a trialkylboric acid ester under conditions to form



b) treating the aryl halide of structure 5a with a palladium catalyst such as $\text{PdCl}_2(\text{dppf})_2$ and a dipinacolate ester of diboronic acid under conditions to form




[0010] In some embodiments, **Y** is selected from the group consisting of OfB(OH)_2 , BF_3 , and




[0011] In some embodiments, **Het** is selected from the group consisting of optionally-substituted pyrrole, furan, piperazine, piperidine, imidazole, 1,2,4-triazol, 1,2,3-triazol, tetrazole, pyrazole, pyrrolidine, oxazole, isoxazole, oxadiazole, pyridin, pyrimidine, thiazole or pyrazine, such as an optionally-substituted tetrazolyl group, for example 2-methyl-tetrazol-5-yl.

[0012] In some embodiments, the method further comprises treating the compound of the structure




with a glycidyl ester such as glycidyl butyrate. In some aspects the glycidyl ester has *R* stereochemistry, such as *R*-(-)-glycidyl butyrate. This treating step may be carried out in the presence of lithium hexamethyldisilazide.


[0013] Compounds made from the processes described herein include



and



[0014] In some embodiments, a compound of the formula has the structure:




wherein;

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

R^2 is selected from the group consisting of optionally substituted benzyl and optionally substituted C_1 - C_6 alkyl, and

Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom.

[0015] In some embodiments, a compound of the formula has the following structure:




wherein

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

R^2 is selected from the group consisting of optionally substituted benzyl and optionally substituted C_1 - C_6 alkyl, and

Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted Ci-C₆ mono and dihydric alcohols, and wherein R³ and R⁴ together may form a ring.

[0016] In some aspects, a composition comprises the compound herein such as prepared in accordance with the processes herein and a dimer having the following structure or a pharmaceutically acceptable salt of the dimer



wherein R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,


Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom.

[0017] In some aspects, R^{1a} is F and R^{1b} is H and Het is 2-methyl-tetrazol-5-yl.

[0018] In further embodiments a composition comprises the compound herein such as prepared in accordance with the processes above, wherein the composition lacks tin impurities.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

[0019] Methods are provided for synthesizing substituted (pyridinyl)phenyl-oxazolidinones



1

wherein Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom such as optionally-substituted tetrazolyl, oxazolyl, triazolyl, oxadiazolyl, thiazolyl, and isoxazolyl moieties. In some aspects, Het is an optionally-substituted tetrazolyl such as 2-methyl-tetrazol-5-yl.


R^{1a} and R^{1b} are independently selected from H and F, provided that at least one is F, and

R is selected from H, $PO(OH)_2$, and pharmaceutically acceptable salts of $PO(OH)_2$.

[0020] Unless otherwise specified, technical terms here take their usual meanings, specifically those specified in the McGraw-Hill Dictionary of Scientific and Technical Terms, 6th edition.


[0021] In some embodiments, methods include synthesizing substituted N-(pyridinyl)aryloxazolidinones by the following route (**[0022]** Scheme 1)

[0022] Scheme 1



[0023] In Scheme 1, a first intermediate (4) is coupled in Rxn 1 with a second intermediate (6) to afford a coupling product (7), which in Rxn 2 is then treated with a glycidyl ester to afford compound (1).

Scheme 2



[0024] In Scheme 2, intermediate 6 may be formed by treatment of intermediate 5a with 2 equivalents of a strong base such as a Ci-C₆ alkyl lithium for example n-butyl lithium or t-butyl lithium followed by the addition of the appropriate electrophile such as ZnCl₂ or B(OR)₃ i.e., Ci-C₆ trialkoxyboronate such as triisopropyl boronate. Aqueous workup of the resulting reaction mixture where the electrophile is a trialkoxyborate ester yields the boronic acid 6a. If the dianion of 5a is treated with a cyclic boronate ester then the cyclic boronic acid ester 6b can be isolated. Further, if the electrophile is ZnCl₂ then the zinc reagent 6c can be isolated. Alternatively, the boronic acids may be prepared by the Miyaura

boration procedure (Miyaura *Top. Curr. Chem.* 2002, 219, 11-59). In this reaction, a diester of diboronic acid such as dipinacolate ester of diboronic acid is coupled to an arylhalide (5a) using a palladium catalyst. The resulting boronic acid ester 6b can be hydrolyzed with aqueous acid to the boronic acid 6a. Further, the trifluoroborate derivative 6d can be formed from the boronic acid 6a by treatment with KF and/or KHF_2 .


[0025] In the above schemes, X is a leaving group. In some embodiments, X is selected from Cl, Br, I, and trifluoromethanesulfonate.

[0026] X^1 is a leaving group. In some embodiments, X^1 is a halogen such as Cl, Br, or I.

[0027] Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom, including optionally-substituted pyrrole, furan, piperazine, piperidine, imidazole, 1,2,4-triazol, 1,2,3-triazol, tetrazole, pyrazole, pyrrolidine, oxazole, isoxazole, oxadiazole, pyridin, pyrimidine, thiazole or pyrazine. In some aspects, Het is optionally-substituted tetrazolyl or 2-methyl-tetrazol-5-yl. In some embodiments, Het is unsubstituted or has 1 or 2 substituents.

[0028] R^{1a} and R^{1b} are independently selected from H and F, provided that at least one is F;

Y is selected from ZnCl , BF_3 , and BR^3R^4 , wherein R^3 and R^4 are independently selected from OH and optionally-substituted $\text{C}_1\text{-C}_6$ mono and dihydric alcohols, and wherein R^3 and R^4 together may form a ring. In some embodiments, Y is $\text{B}(\text{OH})_2$ or pinacolatoborate,



such as $\text{B}(\text{OH})_2$. $\text{C}_1\text{-C}_6$ mono and dihydric alcohols may be optionally substituted with $\text{C}_1\text{-C}_4$ alkyl. A Negishi reaction may be performed to form compounds wherein Y is ZnCl (Negishi: *Chem. Ind.* 1988, 33, 381-407).

[0029] In some embodiments, Het may be unsubstituted or optionally substituted with one or more substituents, for example, independently selected from the group consisting of halogen, hydroxy, amino, $\text{C}_1\text{-}_4$ alkylamino, $\text{di}(\text{C}_1\text{-}_4 \text{ alkyl})\text{amino}$, cyano, nitro, $\text{C}_1\text{-}_4$ alkyl, $\text{C}_1\text{-}_4$

alkoxy, C₁₋₄ acyl, C₁₋₄ thioalkyl, C₁₋₄ thiooxoalkyl, halogen substituted C₁₋₄ alkyl and halogen substituted C₁₋₄ alkoxy.

[0030] Also in Scheme 1, R² is optionally substituted benzyl or optionally substituted C₁-C₆ alkyl. In some embodiments, benzyl and C₁-C₆ alkyl are unsubstituted or independently optionally substituted with halogen or alkoxy such as C₁-C₄ alkyloxy. In some embodiments, R² is benzyl and [0071] R is H.

[0031] Suitable catalysts for cross-coupling reaction Rxn 1 are palladium complexes, for example palladium phosphine complexes or dichlorobis(triphenylphosphine)-palladium(II), tetrakis(triphenylphosphine)palladium(0), and that prepared *in situ* from Pd₂(dba)₃ (dba = benzylideneacetone) in the presence of PCy₃. The proportion of Pd complex to substrates to be coupled is not critical, but approximately 1 mole % (relative to either 4 or 6) has been found to be useful.


[0032] Cyclization to produce the oxazolidinone ring is effected in Rxn 2 by treating 7 with a strong base, such as lithium hexamethyldisilazide or an organolithium salt, such as n-butyl lithium, in the presence of 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (DMPU), followed by a glycidyl ester such as an R-(-)-glycidyl ester, for example, butyrate, to afford compound 1 (R = H). One embodiment uses lithium hexamethyldisilazide as the base, and THF as the solvent, with DMPU present to facilitate the reaction, at a temperature between about 0°C and about 30°C, and at a stoichiometry of 7 to glycidyl ester of about 1:1 on a molar basis.

[0033] If desired, compound 1 (R = H) can further be converted to the dihydrogen phosphate, for example, by treatment with POCl₃, according to well-known methods. For example, compound 1 (R = H) can be treated with POCl₃ followed by an aqueous quench or in a two step process using a protected form of phosphorous oxychloride such as: POCl(OBn)₂ where the first step prepares the phosphate triester and the second step removes the protecting group (for example H₂ZPd-C to remove the benzyl esters). Alternatively, the 5-hydroxymethyl-oxazolidinone can be treated with P(N-iPr)₂(O-tBu)₂ followed by oxidation with an oxidizing reagent such as mCPBA followed by treatment with base or aqueous acid to remove the tert-butyl esters).

[0034] The resulting dihydrogen phosphate compound 1 ($R = PO(OH)_2$) can further be converted to a pharmaceutically acceptable salt such as the disodium salt of compound 1 ($R = PO(O)_2 2Na$) by reaction with NaOMe or other suitable sodium-containing base.

[0035] Those skilled in the art of medicinal chemistry will appreciate that the term "pharmaceutically acceptable salt" refers to salts formed with such biologically compatible cations and/or anions, as appropriate. Such cations include those of metallic elements, such as sodium, lithium, potassium, magnesium, aluminum, calcium, zinc, and quaternary cations of organic nitrogenous bases, such as N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, N-methylglucamine and procaine-salts. Such anions include those of inorganic acids, such as hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, perchloric, fumaric, acetic, propionic, succinic, glycolic, formic, lactic, maleic, tartaric, citric, palmoic, malonic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, fumaric, toluenesulfonic, methanesulfonic, naphthalene-2-sulfonic, benzenesulfonic hydroxynaphthoic, hydroiodic, malic, steroic, tannic and similar acids.

[0036] Oxazolidinones prepared by the methods herein differ from the oxazolidinones that have been prepared in accordance with the US 20070155798 method. Oxazolidinones made in accordance to the process described herein do not contain tin impurities as no tin-containing reactants are used. In addition, in some embodiments, a dimer impurity has been observed, for example, in batches in which phosphorus oxychloride ($POCl_3$) was used to convert hydroxyl to the dihydrogen phosphate. Specifically, a molecule of TR-701 reacts with a molecule of phosphate ester containing at least one P-Cl bond to form the dimer, such as having the following structure.



[0037] The impurity is present in some detectable quantity and is present in less than about 10% by weight of the composition, and in some cases less than about 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1%, such as less than 0.1% or 0.05%. Thus, in some embodiments,

compositions comprise an oxazolidinone as prepared in accordance with the process herein and a dimer. In some embodiments compositions comprise an oxazolidinone lacking any tin impurities.

[0038] Oxazolidinones prepared by the methods herein are useful as medicaments, and particularly for impeding the growth of bacteria, as is disclosed in detail in US 20070155798, which has been incorporated by reference in its entirety.

[0039] The terms "approximately", "about", and "substantially" as used herein represent an amount close to the stated amount that still performs the desired function or achieves the desired result. For example, the terms "approximately", "about" and "substantially" may refer to an amount that is within less than 10% of, within less than 5% of, within less than 1% of, within less than 0.1% of, and within less than 0.01% of the stated amount.

Examples

[0040] The practice of the inventive method is illustrated by the following non-limiting example.

Experimental and Analytical Data

[0041] Reagents were purchased from commercial sources and were used as received. Proton nuclear magnetic resonance spectra were obtained on a Bruker AVANCE 300 spectrometer at 300 MHz or an AVANCE 500 spectrometer at 500 MHz with tetramethylsilane used as an internal reference. Carbon nuclear magnetic resonance spectra were obtained on a Bruker AVANCE 500 spectrometer at 125 MHz with the solvent peak used as the reference. Phosphorus nuclear magnetic resonance spectra were obtained on a Bruker AVANCE 500 spectrometer at 202 MHz with phosphoric acid used as the reference. Fluorine nuclear magnetic resonance spectra were obtained on a Bruker AVANCE 300 spectrometer at 282 MHz. Mass spectra were obtained on a Finnigan AQA spectrometer with electrospray ionization. Thin-layer chromatography (TLC) was performed using Whatman No. 4500-101 (Diamond No. MK6F silica gel 60 A) plates. Visualization of TLC plates was performed using UV light (254 nm) or potassium permanganate stain. HPLC analyses were obtained on a Varian Prostar HPLC equipped with a Waters SunFire C18 column (150 x 4.60

mm, 3.5 μm) or Waters XBridge C18 column (75 mm x 4.6 mm x 2.5 μm) using the methods below with the detector at the specified wavelength.

[0042] Method A (Waters SunFire C18 Column)

Time (min)	Flow (mL/min)	% A	% B
0.0	1.0	98.0	2.0
15.0	1.0	5.0	95.0
25.0	1.0	5.0	95.0
27.0	1.0	98.0	2.0
30.0	1.0	98.0	2.0

A = water with 0.05% v/v trifluoroacetic acid

B = acetonitrile with 0.05% v/v trifluoroacetic acid

Wavelength = 254 nm

[0043] Method B (Waters XBridge C18 Column)

1. Time (min)	2. Flow (mL/min)	% A	% B
0.0	1.0	98.0	2.0
15.0	1.0	5.0	95.0
25.0	1.0	5.0	95.0
27.0	1.0	98.0	2.0
30.0	1.0	98.0	2.0

A = 87% 25 mM ammonium bicarbonate solution in water/13% acetonitrile

B = acetonitrile

Wavelength = 254 nm

[0044] Method C (Waters SunFire C18 Column)

3. Time (min)	4. Flow (mL/min)	% A	% B
0.0	1.0	98.0	2.0
15.0	1.0	5.0	95.0

25.0	1.0	5.0	95.0
27.0	1.0	98.0	2.0
30.0	1.0	98.0	2.0

A = water with 0.05% v/v trifluoroacetic acid

B = acetonitrile with 0.05% v/v trifluoroacetic acid

Wavelength = 240 nm

Example 1: Preparation of 5-Bromo-2-(2H-tetrazol-5-yl)pyridine, 3

[0045] To a 22-L, three-neck, round-bottom flask equipped with an overhead stirrer, nitrogen inlet/outlet, thermocouple and heating mantle was charged 5-bromo-2-cyanopyridine (799 g, 4.37 mol, 1 weight), *N,N*-dimethylformamide (6.4 L, 8 volumes), ammonium chloride (350.3 g, 6.55 mol, 1.5 equivalents), and sodium azide (425.7 g, 6.55 mol, 1.5 equivalents) while stirring. The internal reactor temperature set-point was adjusted to 85°C (Target temperature is 90°C). The temperature set-point was reached after 45 minutes, and the reaction continued to self-heat to 94°C over 40 minutes. The reaction was judged to be complete after 1 hour by HPLC analysis by complete consumption of the starting material with an assay of 76.7% (AUC) of the tetrazole ammonium salt. The mixture was cooled and filtered at room temperature. The reactor and wet cake were washed with 2-propanol (3.2 L, 4 volumes) and dried under high vacuum at ambient temperature to afford the tetrazole ammonium salt as an off-white solid (847.9 g, 80% yield, 89.9% AUC). A differential scanning calorimetry experiment was conducted on the tetrazole ammonium salt to understand its thermal stability. The salt melted at approximately 228°C and had an energetic decomposition at approximately 270°C.

Example 2: Preparation of 5-Bromo-2-(2-methyl-2H-tetrazol-5-yl)pyridine, 4 (X = Br)

[0046] To a 22-L, four-neck, round-bottom flask equipped with an overhead stirrer, nitrogen inlet/outlet, and thermocouple placed in an ice/brine bath was charged the tetrazole ammonium salt (835.0 g, 3.44 mol, 1 weight), tetrahydrofuran (7.5 L, 9 volumes), *N,N*-dimethylformamide (2.5 L, 3 volumes) and sodium hydroxide powder (343.5 g, 8.59 mol, 2.5 equivalents) while stirring. The internal reactor temperature was allowed to reach 12°C, whereupon iodomethane (1.22 kg, 8.59 mol, 2.5 equivalents) was added dropwise over 50

minutes, maintaining the reaction temperature below 20°C. After 20 minutes addition time, due to a rapid increase in temperature, the addition was paused and the reaction continued to self-heat from 15-20°C over ten minutes. The remainder of the addition was completed at constant temperature (18°C). Upon completion of the addition, the ice/brine bath was removed and the reactor was equipped with a water condenser and a heating mantle. The internal reactor temperature was adjusted to 40°C, however the reaction continued to self-heat to 48°C. The reaction was judged to be complete after 6 hours by HPLC analysis by complete consumption of the starting material. The reaction mixture was cooled to room temperature overnight for convenience. The THF was removed by distillation, and water (8.35 L, 10 volumes) was charged to the reactor. The slurry was stirred for 30 minutes and filtered by vacuum filtration and the reactor and filter cake were washed with water (4.2 L, 5 volumes) to afford crude 4/Nl isomer mixture as a peach colored solid (500.7 g, 61% yield, 3.85: 1 4:Nl).

[0047] The solids (500.7 g) were dissolved in CH₂Cl₂ (2.5 L, 5 volumes) to which 6 N aqueous HCl (7.5 L, 15 volumes) was added. The biphasic mixture was stirred and the layers were separated. At this point, the desired product is in the aqueous HCl layer. The CH₂Cl₂ layer was washed with 6 N aqueous HCl (4.5 L, 3 x 3 volumes) until <5% AUC 4 was present by HPLC analysis. The combined 6 N HCl extracts were transferred to a reactor and the pH was adjusted to 10.6 with 50% aqueous NaOH (-3.2 L) while maintaining the internal temperature below 40°C. The solids were isolated by vacuum filtration and the reactor and filter cake were rinsed with water (1 L, 2 volumes) to afford crude 4 as a yellow/orange solid (322.4 g, 64% recovery, 39% yield, 93.5% AUC 4, 4.1% AUC N-I isomer) as confirmed by HPLC and ¹H NMR analyses.

[0048] The crude 4 was further purified by an isopropyl acetate (IPAc) reslurry (1.61 L, 5 volumes) at 50°C for 1 hour. Upon cooling to room temperature, the solids were filtered and the reactor and filter cake were washed with additional IPAc (500 mL, 1.6 volumes) to afford purified 4 as a off-white/yellow solid (275.5 g, 85% recovery, 33% yield, 98.2% AUC) as confirmed by HPLC and ¹H NMR analyses. DSC analysis of 4 showed a decomposition exotherm at approximately 245°C.

Example 3: Preparation of benzyl (4-bromo-3-fluorophenyl)carbamate, 5

[0049] To a 12-L, three-neck, round-bottom flask equipped with an overhead stirrer, nitrogen inlet/outlet, addition funnel and thermocouple was charged 4-bromo-3-fluoroaniline (800.0 g, 4.21 mol, Matrix lot # Q13H), THF (6.4 L, 8 vol), and solid sodium bicarbonate (530.5 g, 6.32 mol, 1.5 eq). The addition funnel was charged with benzyl chloroformate (861.9 g, 5.05 mol, 1.2 eq), which was added dropwise to the reactor over 70 minutes. The temperature of the reaction was maintained below 20°C with an ice water bath. The batch was aged 1 hour at room temperature at which point HPLC analysis indicated that the reaction was complete. The reaction mixture was transferred to a 22-L flask and the mixture was diluted with water (6.4 L, 8 vol). The two-phase mixture was warmed to 50°C and held at temperature for 16 hours to quench the excess benzyl chloroformate. The mixture was transferred hot to a separatory funnel to remove the lower aqueous phase. A rag layer was observed which was taken with the aqueous layer. The THF layer was filtered through Whatman #1 filter paper to remove some particulates, and the mixture was transferred back to a 22-L flask equipped for distillation. Heptane was added in portions and distilled to remove the THF. (Note that it is best to distill some of the THF out first before adding the first amount of heptane.) A total of 26.5 L of heptane was added, and the total distillate collected was 25 L. At this point, the pot temperature had reached 97.7°C and the distillate coming over contained 0.9% THF by ¹H NMR analysis. The mixture was cooled to room temperature and the thick white slurry was filtered. The filter cake was washed with heptane (4 L). The product was dried in a vacuum oven at 40°C to give 1257.0 g of intermediate 5 (92% yield). The HPLC assay was 98.3% (AUC).

Example 4: Preparation of 4-(Benzyloxycarbonylamino)-2-fluorophenylboronic acid 6
(R^{1a} = F. R^{1b} = H. R² = Bz. Y = B(OH)₂)

[0050] A 22-L, three-neck, round-bottom flask was equipped with an overhead stirrer, temperature probe, 2-L addition funnel, and a nitrogen inlet adapter. The flask was charged with intermediate 5 (1.00 kg, 3.08 mol, AMRI lot # CAR-L-18(3)), THF (10 L, 10 vol) and triisopropyl borate (638.2 g, 3.39 mol, 1.1 eq.). The mixture was stirred and cooled to -72°C in a dry ice/acetone bath. The addition funnel was charged in portions with 2.5 M *n*-butyllithium (2.59 L, 6.48 mol, 2.1 eq.), which was added dropwise to the reaction over approximately 2 hours. The maximum temperature during the addition was -65°C. The

reaction was deemed complete by HPLC analysis. The acetone was removed from the cooling bath, and the reaction was quenched with 20% aqueous ammonium chloride solution (5.5 L), allowing the reaction to warm to -1°C. The phases were separated and the THF layer was evaporated to dryness. The crude product was reslurried in 3:2 ethanol/water (10 L, 10 vol) at room temperature for 1 hour. The mixture was filtered and the filter cake was rinsed with 3:2 ethanol/water (2 x 2 L). The product was dried in a vacuum oven at room temperature to give 592.8 g of intermediate 6 (66% yield) that was 89.8% (AUC) by HPLC analysis (Method A). The material was much less pure by ¹⁹F NMR analysis and HPLC analysis at 240 nm (Method C).

[0051] Later development of this process used 2.5 volumes of CH₂Cl₂ to reslurry the crude product in place of 3:2 ethanol/water, which removed the *des-bromo* by-product, which was the impurity observed in the ¹⁹F NMR spectrum and the HPLC at 240 nm.

Example 5: Preparation of benzyl (4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)carbamate. 7 (Het = 2-methyltetrazol-5-yl. RIa= F. RIb = H. R2 = Bz)(Ref: JAS-G-96) (Ref: CAR-L-93. DUG-AF-202)

[0052] To a 5-L, three-neck, round-bottom flask was charged 4 (200.0 g, 0.833 mol) followed by 1,4-dioxane (3 L, 15 vol). Crude compound 6 (361.2 g, 1.249 mol, 1.5 equiv.), Pd₂(dba)₃ (11.44 g, 0.0125 mol, 0.015 equiv.), and PCy₃ (7.0 g, 0.025 mol, 0.03 equiv.) was charged and degassed with nitrogen for 30 minutes. A solution of K₂CO₃ (195.7 g, 1.7 equiv.) in water (800 mL, 4 vol) was charged, and the reaction was heated to 70 °C. The reaction was complete after 1 hour with 0.5 area % of 4 remaining. The reaction was cooled to 50 °C, and Darco G-60 (40 g, 0.2 wt) was added and stirred for 30 minutes. Celite 545 (40 g, 0.2 wt) was charged and then the reaction was filtered through Celite 545 (100 g, 0.5 wt) wetted with water (300 mL). The hot filtration into the water from the Celite caused precipitation of the product. Tetrahydrofuran (1.2 L, 6 vol) and brine (600 mL, 3 vol) were added, and the product re-dissolved at room temperature. The phase split was accomplished cleanly (Vmax = 28 volumes). The dioxane was concentrated and ethanol (1 L, 5 vol) was added and concentrated. Then the product was reslurried in ethanol: water (4:1, 2 L, 10 vol) at 70°C, cooled to room temperature over 3 hours, filtered and washed with ethanol (2 x 400 mL). Compound 7 was isolated in 87% yield (292.6 g) with a purity of 97.7 % (AUC) by

HPLC analysis. The ^1H NMR and ^{19}F NMR indicated the presence of one compound. Pd analysis showed 135 ppm Pd was in the product.

[0053] The intermediate 7 was recrystallized from ethyl acetate to further reduce the level of palladium. Intermediate 7 (130 g) and ethyl acetate (3.9 L, 30 volumes) were charged to a 5-L, three-neck, round-bottom flask. The slurry was warmed to 75°C at which point the solids dissolved. The hot solution was filtered to remove any palladium black (0.2- to 0.45- μ filters the best) and returned to a clean 5-L flask. The ethyl acetate solution was distilled at atmospheric pressure to remove 2.2 L of the ethyl acetate (b.p. 77-78°C). The solution was cooled to 22°C and the resulting slurry was filtered. The flask and filter cake were washed with ethyl acetate (3 x 130 mL) of ethyl acetate. The purified intermediate 7 was dried in a vacuum oven at 50°C to give 110.5 g of intermediate 7 (85% recovery). The HPLC assay of the purified intermediate 7 was 98.5% (AUC). The palladium level in the purified product was 6 ppm. The mother liquor was evaporated to recover 18 g of crude product (14% recovery, 2254 ppm Pd).

Example 6: Preparation of (R)-3-(4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)-5-hydroxymethyl oxazolidin-2-one, 1 (R = H), also referred to as "TR-700"

[0054] A 5-L, three-neck, round-bottom flask was equipped with an overhead stirrer, a thermocouple, a 500-mL addition funnel and a nitrogen-inlet adapter. The flask was dried with a heat gun under a flow of nitrogen to an internal temperature of 60°C. The flask was charged with intermediate 7 (110.0 g, 0.272 mol, AMRI lot # DUG-AF-202Q) and anhydrous THF (2.2 L, 20 vol). The slurry was stirred and a light green solution formed. The addition funnel was charged with 1.0 M lithium hexamethyldisilazide (299 mL, 0.286 mol, 1.05 eq.). The LiHMDS solution was added dropwise to the solution of intermediate 7 over approximately 25 minutes. A red solution formed. The solution was stirred one hour at room temperature and then DMPU (34.9 g, 0.272 mol, 1 eq) was added, and the mixture turned to a yellow slurry. The batch was cooled in an ice bath to 5.7°C. R-(-)-Glycidyl butyrate (41.25 g, 0.286 mol, 1.05 eq) was then added in one portion. The mixture was stirred in the ice bath for 0.5 hour and then was warmed to room temperature and stirred overnight. The reaction formed a tan slurry at this point, and HPLC analysis after 15 hours indicated that there was approximately 87% TR-700, 1.6% intermediate 7, and approximately 7% of the butyrate ester

of TR-700. A small amount of sodium methoxide in methanol (11 mL, 0.1 vol) was added, and the batch was stirred for 1 hour to remove the residual ester. The in-process HPLC analysis at this point showed there was approximately 90.7% TR-700 and 0.2% of the butyrate ester. The reaction was quenched by the addition of 10% w/w ammonium chloride solution (1.1 L, 10 vol). A modest exothermic event from 22°C to 25°C was observed upon addition of the ammonium chloride solution. The two-phase mixture was distilled to a pot temperature of 70°C (atmospheric pressure) to remove approximately 2.2 L of the THF. This formed a thick slurry which is diluted with water (550 mL, 5 volumes). The slurry was cooled to room temperature (23.6°C) and was filtered. The filter cake was washed with water (1.1 L, 10 vol) and methanol (550 mL, 5 vol) to give TR-700 as a white solid. The wet cake was dried overnight in a vacuum oven at 50°C to give 89.7 g of TR-700 (89% yield) that was 97.8% (AUC) by HPLC analysis. The TR-700 was further purified by reslurrying in 2.7 L (30 vol) of 4:1 methanol/water at 70°C, cooling to 23°C, filtering and washing with methanol (180 ml). This removed some of the over-alkylated product that is observed. The purified TR-700 was recovered in 96% yield (85% overall yield), and the purity was improved to 98.4% (AUC) by HPLC analysis. The palladium content was 10 ppm.

Example 7: Preparation of (R)-3-(4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)-5-hydroxymethyl oxazolidin-2-one dihydrogen phosphate 1 (R = PO(OH)₂)
also referred to as "TR-701FA"

[0055] A 5-L, jacketed round-bottom flask was equipped with an overhead, mechanical stirrer, addition funnel, thermocouple, nitrogen inlet, and a circulating chiller unit. The flask was charged with TR-700 (70.0 g, 0.189 mol), THF (1.4 L, 20 vol), and triethylamine (58.2 g, 0.575 mol, 3 eq). The slurry was stirred and the jacket temperature was set to 0°C. The addition funnel was charged with phosphorus oxychloride (87.0 g, 0.567 mol, 3 eq) in THF (70 mL, 1 vol). Once the internal temperature reached 1°C, the POCl₃ solution was added dropwise over 44 minutes. The maximum internal temperature was 2.2°C. The mixture was stirred for 3 hours at 1-2°C at which point HPLC analysis indicated that <0.5% of the TR-700 remained. A 5-L, three-neck, round-bottom flask equipped with a Teflon diaphragm pump was charged with water (1.4 L, 20 vol) and was cooled to 3.8°C in an ice, salt water bath. The reaction mixture was pumped into the quench water subsurface over 1

hour. The maximum temperature during the quench was 11.9°C. The reactor and pump lines were rinsed with water (-210 mL) into the quench vessel. The yellow slurry was stirred overnight. The slurry was filtered through Whatman paper, and the filter cake was rinsed with water (700 mL, 10 vol) and methanol (700 mL, 10 vol). The product was dried at room temperature in a vacuum oven until a constant weight was obtained. The yield of crude TR-701FA was 81.6 g (96%), and the purity by HPLC analysis (Method B) was 95.3% (AUC).

Example 8: Preparation of (R)-3-(4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)-5-hydroxymethyl oxazolidin-2-one phosphate, disodium salt 1 (R = PO₃.2Na) also referred to as "TR-701"

[0056] Crude **1** (R = PO(OH)₂) (60.0 g, 0.133 mol) was charged to a 2-L reactor. Methanol (720 mL, 12 vol) was added and the slurry was stirred at room temperature. The 25% sodium methoxide in methanol (86.1 g, 0.398 mol, 3 eq) was added dropwise over 13 minutes. The reaction temperature increased from 20.4°C to 26.8°C during the addition of sodium methoxide. The slurry was stirred one hour at room temperature and then was filtered. The reactor and filter cake were rinsed with methanol (300 mL, 5 vol) and acetone (300 mL, 5 vol). The product was dried in a vacuum oven at 50-60°C to give 65.3 g of crude TR-701 (99% yield). The crude product was dissolved in water (653 mL, 10 vol) to give a straw colored solution. The solution was stirred with Darco G-60 carbon (3.3 g, 0.05 wt) at room temperature for 30 minutes. The pH of the slurry was 7.2, so 5-10 mL of 2 N NaOH was added to the solution to raise the pH to 11. The slurry was filtered through Celite 545 (65 g, wetted with water). Some black color passed through. The filtrate was refiltered through a 0.45-μ filter, but some carbon passed through again. The filtrate was added dropwise to acetone (2.6 L, 40 vol), and the resulting slurry was stirred overnight for convenience. The slurry was then filtered, rinsed with acetone (650 mL), and dried in a vacuum oven at 50°C to give 46.9 g of **1** (R = PO₂Na) (71% yield) that was gray in color. The HPLC purity of this material was 99.0% (AUC), but since it was gray, it was re-dissolved in water (470 mL). The aqueous solution was pH 9.6, so sodium hydroxide solution was added to raise the pH to 10. The solution was then filtered through a 0.45-μ filter to remove color. The filtrate was added dropwise to acetone (1.88 L). The white slurry was filtered and was washed with acetone (470 mL). After drying the product, the TR-701 weighed 43.2 g (66% overall yield). The

HPLC purity (Method B) was 99.6% (AUC). The other analyses conducted on this lot of **1** ($R = PO_2Na$) are shown in Table 1.

Example 9: Preparation of Purified R)-3-(4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)-5-hydroxymethyl oxazolidin-2-one dihydrogen phosphate, 1 (R = PO(OH)₂)

[0057] A 3-L, three-neck, round-bottom flask was charged with crude **1** ($R = PO(OH)_2$) (99.8 g, 0.222 mol, AMRI lot # 8AK0242C) and water (1 L, 10 vol). The pH of this slurry was 2.05. A fresh solution of 1 M sodium hydroxide solution was prepared by dissolving 50.9% aqueous sodium hydroxide (39.3 g, 0.50 mol) in a total volume of 0.5 L of water. The 1 M sodium hydroxide solution (444 mL, 0.444 mol, 2 eq) was added dropwise to the free acid slurry. At pH 5.7, the solids dissolved, even though only a little over half the sodium hydroxide solution had been added. At the end of the addition the pH was 8.57. Darco G-60 carbon (5.1 g, 0.05 wt) was added to the solution and the mixture was stirred for 1 hour at room temperature. The slurry was filtered through Whatman #1 filter paper to remove the bulk carbon, and then through a 0.45- μ filter to remove the fines. The straw-colored filtrate was added dropwise to a 12-L round-bottom flask containing acetone (4 L, 40 vol). The resulting slurry was stirred for 1 hour at room temperature, was filtered and washed with acetone (500 mL, 5 vol). The wet cake was loaded into a 3-L round-bottom flask and was allowed to dry under a nitrogen purge overnight.

[0058] The disodium salt **1** ($R = PO_2 2Na$) was re-dissolved in water (1 L, 10 vol) and then was filtered through Whatman #1 filter paper when black flecks were observed in the solution. The filtrate was diluted with THF (1 L, 10 vol). The pH of the aqueous THF solution was 9.57. Freshly prepared 2 M hydrochloric acid solution (222 mL, 0.444 mol, 2 eq.) was added dropwise to pH 1.34. The product did not precipitate until approximately 170 mL of the 2 M HCl solution was added. The yellow slurry was filtered and rinsed with water (500 mL, 5 vol) and methanol (500 mL, 5 vol). The filter cake cracked as it dried, so it was smoothed out before adding the rinse solvents. The product was dried in a vacuum oven at 60°C for 19.5 hours to give 79.3 g of **1** ($R = P(OH)_2$) (80% yield). HPLC analysis (Method B): 99.5% (AUC) $t_R = 5.6$ min. ¹H and ³¹P NMR analyses were consistent with the assigned structure. The level of residual THF by NMR analysis was 1600 ppm, and the palladium level

was 11 ppm. Since extended drying did not remove the THF, future batches were made with use of ethanol as the antisolvent.

Example 10: Isolation of bis([(5R)-3-(3-fluoro-4-[6-(2-methyl-2H-tetrazol-5-yl)pyridin-3-yl]phenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl} dihydrogen diphosphate (the dimer of 1)

[0059] Crude 1 from example 8 was dissolved in phosphate buffer and chromatographed on a Gilson preparative HPLC system. The mobile phase was a linear gradient of water and acetonitrile t+0 was 100% H₂O and T=20 was 100% acetonitrile. Fractions were analyzed using analytical HPLC. Those fractions found to be enriched in the Dimer were pooled providing a solution containing over 60% Dimer. Further purification of the Dimer enriched fractions was accomplished on a semi preparative HPLC. This yielded pure dimer: accurate mass (m/z 883; calcd. For C₃₄H₃₁F₂N₁₂O₁ 1P₂ = 883.1679, found 883.1658, Δ = 2.4 ppm m/z 905 calcd. for C₃₄H₃₀F₂N₁₀ 1P₂Na = 905.1498; found 905.1484, Δ = 1.6 ppm) confirming the formula for this compound.


Table 1. Analysis of TR-701 (R)-3-(4-(2-(2-methyltetrazol-5-yl)pyridin-5-yl)-3-fluorophenyl)-5-hydroxymethyl oxazolidin-2-one, 1 (R = H)

Test	Result
Appearance	White to Off-white
¹ H NMR	Conforms
³¹ P NMR	Conforms
Retention Time	5.18 min
MS	<i>m/z</i> 371
HPLC Purity	99.6*
HPLC Impurities	Dimer, 0.09%*
Copper Content	<1 ppm
Palladium Content	1 ppm
Sodium Content	8.34%
Water Content	5.5%
Specific Rotation	-34.9°

XRPD	Amorphous
Particle Size	1-300 μm

WHAT IS CLAIMED IS:

1. A method of synthesizing a compound of the structure




wherein

R is H,


R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom,


comprising treating a compound having the structure




wherein R² is selected from the group consisting of optionally substituted benzyl and optionally substituted C₁-C₆ alkyl, with a strong base or an organolithium salt and then addition of glycidyl butyrate to the resulting anion under conditions to make



2. The method of claim 1 further comprising reacting




with POCl_3 , $\text{POCl}(\text{OBn})_2$, or $\text{P}(\text{N-iPr})_2(\text{O-tBu})_2$ under conditions form




wherein R' is $\text{PO}(\text{OH})_2$.

3. The method of claim 2 further comprising treating the compound of the structure




where R' is $\text{PO}(\text{OH})_2$ with a base under conditions to form the compound of the structure




wherein R'' is a pharmaceutically acceptable salt of $\text{PO}(\text{OH})_2$.

4. The method of claim 1 further comprising before said treating step: coupling a first intermediate of the structure




wherein X is a leaving group, with a second intermediate of the structure




wherein Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted C₁-C₆ mono and dihydric alcohols, and wherein R³ and R⁴ together may form a ring,


under conditions to produce the compound of the structure




5. The method of claim 4 further comprising before said coupling step:
a) treating an aryl halide of structure 5a




wherein X¹ is leaving group, with a strong base and then reacting a resulting anion with a trialkylboric acid ester under conditions to form



- b) treating the aryl halide of structure 5a with a palladium catalyst and a dipinacolate ester of diboronic acid under conditions to form



6. The method of claim 5,
 wherein the strong base is n-butyl lithium; or
 wherein the palladium catalyst is PdCl₂(dppf)₂.
7. A method of synthesizing a compound of the structure



wherein:


R² is selected from the group consisting of optionally substituted benzyl and optionally substituted Ci-C₆ alkyl,

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F, and


Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom,

comprising:

coupling a first intermediate of the structure




wherein X is a leaving group, with a second intermediate of the structure




wherein

Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted Ci-C₆ mono and dihydric alcohols, and wherein R³ and R⁴ together may form a ring

under conditions to produce the compound of the structure



8. A method of making the second intermediate of the structure



wherein


R² is selected from the group consisting of optionally substituted benzyl and optionally substituted Ci-C₆ alkyl,

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F, and


Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted Ci-C₆ mono and dihydric alcohols, and wherein R³ and R⁴ together may form a ring;

comprising:

a) treating an aryl halide of structure 5a and



wherein X^1 is leaving group, with a strong base and then reacting a resulting anion with $ZnCl_2$ or a trialkylboric acid ester under conditions to form




b) treating the aryl halide of structure 5a with a palladium catalyst and a diester of diboronic acid.

9. The method of claim 8,
 - wherein the strong base is n-butyl lithium; or
 - wherein the palladium catalyst is $PdCl_2(dppf)_2$.
10. The method of claim 1
 - wherein the treating step is performed in the presence of a facilitating compound.
11. The method of claim 10
 - wherein the facilitating compound is 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*i*)-pyrimidinone.
12. The method of claim 4, wherein X is selected from the group consisting of Cl, Br, I, and trifluoromethanesulfonate.
13. The method of claim 7, wherein the coupling is carried out in the presence of a palladium complex.
14. The method of claim 13, wherein the palladium complex includes a phosphine ligand bound to palladium.

15. The method of claim 14, wherein the palladium complex is selected from the group consisting of dichlorobis(triphenylphosphine)palladium(II), tetrakis(triphenylphosphine)palladium(0), and $\text{Pd}_2(\text{dba})_3$

16. The method of claim 15, wherein the palladium complex is $\text{Pd}_2(\text{dba})_3$.

17. The method of claim 5, wherein Y is selected from the group consisting of $\text{B}(\text{OH})_2$, BF_3 , and



18. The method of claim 17, wherein Y is $\text{B}(\text{OH})_2$.

19. The method of claim 1, wherein Het is selected from the group consisting of optionally-substituted pyrrole, furan, piperazine, piperidine, imidazole, 1,2,4-triazol, 1,2,3-triazol, tetrazole, pyrazole, pyrrolidine, oxazole, isoxazole, oxadiazole, pyridin, pyrimidine, thiazole or pyrazine.

20. The method of claim 19, where Het is an optionally-substituted tetrazolyl group.


21. The method of claim 20 wherein Het is 2-methyl-tetrazol-5-yl.

22. The method of claim 21 wherein Het is 2-methyl-tetrazol-5-yl and X is Br.

23. The method of claim 3,

wherein the base is a sodium-containing base and wherein R" is PO_3Na_2

24. The method of claim 7, further comprising treating the compound of the structure




with a glycidyl ester.

25. The method of claim 24, wherein the glycidyl ester is glycidyl butyrate.

26. The method of claim 24, wherein the glycidyl ester has *R* stereochemistry.


27. The method of claim 24, wherein the glycidyl ester is *R*-(-)-glycidyl butyrate.

28. The method of claim 24, wherein the treating of the compound of the structure




with a glycidyl ester is carried out in the presence of lithium hexamethyldisilazide.


29. The method of claim 1, wherein the compound of the structure




is




30. The method of claim 2, wherein the compound of the structure




is




31. The method of claim 21, wherein the compound of the structure



is



32. A compound of the formula:




wherein;

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

R^2 is selected from the group consisting of optionally substituted benzyl and optionally substituted C_1-C_6 alkyl, and

Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom.

33. A compound of the formula:




wherein

R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

R² is selected from the group consisting of optionally substituted benzyl and optionally substituted Ci-C₆ alkyl, and

Y is selected from the group consisting of ZnCl, BF₃, and BR³R⁴, wherein R³ and R⁴ are independently selected from the group consisting of OH and optionally-substituted C₁-C₆ mono and dihydric alcohols, and wherein R³ and R⁴ together may form a ring.

34. A composition comprising
the compound prepared in accordance with the process of Claim 2; and
a dimer having the following structure or a pharmaceutically acceptable salt



wherein R^{1a} and R^{1b} are independently selected from H and F, provided that at least one of R^{1a} and R^{1b} is F,

Het is an optionally-substituted five- or six-membered heterocycle comprising at least one N, O, or S atom.

35. The composition of claim 34
wherein R^{1a} is F and R^{1b} is H and Het is 2-methyl-tetrazol-5-yl.
36. A composition comprising
the compound prepared in accordance with the process of Claim 1, wherein the composition lacks tin impurities.
37. A composition comprising
the compound prepared in accordance with the process of Claim 2, wherein the composition lacks tin impurities.