

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
WO 03/013580 A1

- (51) International Patent Classification<sup>7</sup>: A61K 38/17, 45/00, G01N 33/50, A61P 3/00
- (21) International Application Number: PCT/IB02/03404
- (22) International Filing Date: 31 July 2002 (31.07.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/310,088 3 August 2001 (03.08.2001) US
- (71) Applicant (for all designated States except US): GENSET S.A. [FR/FR]; Intellectual Property Department, 24, rue Royale, F-75008 Paris (FR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): LUCAS, John [US/US]; 339 The Valley Road, Concord, MA 01742 (US). DIALYNAS, Deno [US/US]; 3425 Lebon Drive, #813, San Diego, CA 92122 (US). BRIGGS, Kristen [US/US]; 2454 Mango Way, Del Mar, CA 92014 (US).
- (74) Common Representative: GENSET S.A.; c/o De Luca, Giampiero, Serono International S.A., Intellectual Property, 12, Chemin des Aulx, 1228 Plan-les-Ouates (CH).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/013580 A1

(54) Title: TR XIDATIN AGONISTS AND ANTAGONISTS TREATMENT OF METABOLIC DISORDERS

(57) Abstract: The present invention relates to the field of metabolic research, in particular the discovery of compounds effective for reducing body mass and useful for treating obesity-related diseases and disorders. The obesity-related diseases or disorders envisioned to be treated by the methods of the invention include, but are not limited to, hyperlipidemia, atherosclerosis, insulin resistance, diabetes, and hypertension. In particular, the invention provides for methods of identifying and using AGONISTS and ANTAGONISTS of XIDATIN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity.

## XIDATIN AGONISTS AND ANTAGONISTS TREATMENT OF METABOLIC DISORDERS.

### Field of the Invention

The present invention relates to the field of metabolic research, in particular the discovery of compounds effective for reducing body mass and maintaining weight loss and useful for treating obesity-related diseases and disorders. The obesity-related diseases or disorders envisioned to be treated by the methods of the invention include, but are not limited to, hyperlipidemia, atherosclerosis, insulin resistance, diabetes, and hypertension. The present invention additionally relates elsewhere to the field of metabolic research, in particular the discovery of compounds effective for increasing body mass and useful for treating disorders associated with excessive weight loss. Applicant reserves the right to exclude any of the aforesaid obesity-related diseases or disorders. The disorders associated with excessive weight loss and envisioned to be treated by the methods of the invention include, but are not limited to, cachexia, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, and anorexia. Applicant reserves the right to exclude any of the aforesaid disorders associated with excessive weight loss.

In particular, the invention provides for methods of identifying and using AGONISTS and ANTAGONISTS of XIDATIN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity.

### Background of the Invention

The following discussion is intended to facilitate the understanding of the invention, but is not intended nor admitted to be prior art to the invention.

Obesity is a public health problem that is serious, widespread, and increasing. In the United States, 20 percent of the population is obese; in Europe, a slightly lower percentage is obese (Friedman (2000) Nature 404:632-634). Obesity is associated with increased risk of hypertension, cardiovascular disease, diabetes, and cancer as well as respiratory complications and osteoarthritis (Kopelman (2000) Nature 404:635-643). Even modest weight loss ameliorates these associated conditions.

Recently it was shown that particular carboxyl-terminal fragments of the full-length ACRP30 (mouse) and APM1 (human) polypeptides have unexpected effects *in vitro* and *in vivo*, including utility for weight reduction, prevention of weight gain, and control of blood glucose levels (Fruebis et al (2001) Proc Natl Acad Sci USA 98:2005-10). The effects of ACRP30 fragment administration in mammals also include reduction of elevated free fatty acid levels including elevated free fatty acid levels caused by administration of epinephrine, *i.v.* injection of "intralipid", or administration of a high fat test meal, as well as increased fatty acid oxidation in muscle cells, and weight reduction in mammals consuming a normal or high fat/high sucrose diet.

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification referenced in this application are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

5

### Summary of the Invention

APM1 belongs to an expanding family of related secreted polypeptides that includes among others C2P, ZADJ-2 and ZADJ-7. These polypeptides have in common the structure: signal peptide, N-terminally disposed unique region, collagen-like region, and globular C-terminal C1q homology domain. APM1, C2P, ZADJ-2 and ZADJ-7 further share an NGLXXD amino acid motif  
10 C-terminally disposed within the globular domain within a loop implicated in receptor binding, wherein said receptor is XIDATIN. Fragments of APM1, C2P, ZADJ-2 and ZADJ-7 polypeptide comprising the globular domain are herein referred to as gAPM1, gC2P, gZADJ-2 and gZADJ-7. It is further taken to be understood herein that LIGAND refers to a composition consisting essentially of or consisting of *in vitro* or *in vivo* self-assembling homotrimer comprised of gAPM1, gC2P,  
15 gZADJ-2, or gZADJ-7 polypeptide fragment.

XIDATIN is a member of the Tumor Necrosis Factor Receptor Super Family (TNFRSF) and is a Type I transmembrane protein. The instant invention is based on XIDATIN as receptor for LIGAND that mediates effects, including utility for weight reduction, maintenance of weight loss, prevention of weight gain, increased insulin sensitivity, and control of blood glucose levels in  
20 humans and other mammals. These effects in mammals of XIDATIN engagement by LIGAND also include reduction of elevated free fatty acid levels including elevated free fatty acid levels including elevated free fatty acid levels caused by administration of epinephrine, *i.v.* injection of "intralipid", or administration of a high fat test meal, as well as increased fatty acid oxidation in muscle cells, and weight reduction in mammals consuming a normal or high fat/high sucrose diet. More specifically,  
25 the present invention is directed to XIDATIN to which LIGAND binds and through which LIGAND mediates said effects.

In particular, the invention provides for methods of identifying and using AGONISTS and ANTAGONISTS of XIDATIN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity, as well as to pharmaceutical and  
30 physiologically acceptable compositions comprising said XIDATIN AGONISTS or ANTAGONISTS and methods of administering said pharmaceutical and physiologically acceptable compositions in order to increase or reduce body weight, maintain weight loss, or to treat obesity-related diseases and disorders. Assays for identifying AGONISTS and ANTAGONISTS of obesity-related activity are also part of the invention.

Preferably said XIDATIN AGONIST or ANTAGONIST is a compound selected from the group consisting of polypeptide, polypeptide fragment, peptide, proein, antibody, carbohydrate, lipid, small molecular weight organic compound and small molecular weight inorganic compound.

Preferably said XIDATIN AGONIST or ANTAGONIST is a compound that selectively  
5 binds to the extracellular domain of XIDATIN.

In other embodiment, said XIDATIN AGONIST or ANTAGONIST is a compound that selectively binds to the intracellular domain of a polypeptide comprising the extracellular domain of XIDATIN.

The present invention also provides a method of assaying test compounds to identify a test  
10 compound that binds to XIDATIN polypeptide. The method comprises contacting XIDATIN polypeptide with a test compound and to determine the extent of binding of the test compound to said XIDATIN polypeptide. The method further comprises determining whether such test compounds are AGONISTS or ANTAGONISTS of XIDATIN polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of XIDATIN  
15 polypeptide or on the activity of XIDATIN polypeptide.

The present invention also relates to diagnostic methods of identifying individuals or non-human animals having elevated or reduced levels of XIDATIN products, which individuals are likely to benefit from therapies to suppress or enhance XIDATIN expression, respectively, and to methods of identifying individuals or non-human animals at increased risk for developing, or present  
20 state of having, certain diseases/disorders associated with XIDATIN abnormal expression or biological activity.

The present invention provides for methods of identifying AGONISTS of XIDATIN polypeptide biological activity comprising contacting a small molecule compound with XIDATIN polypeptides and measuring XIDATIN polypeptide biological activity in the presence and absence  
25 of these small molecules. The present invention further provides for methods of identifying ANTAGONISTS of XIDATIN polypeptide biological activity comprising contacting a small molecule compound with XIDATIN polypeptides and measuring XIDATIN polypeptide biological activity in the presence and absence of these small molecules. These small molecules can be a naturally occurring medicinal compound or derived from combinatorial chemical libraries.

30 The present invention also relates to pharmaceutical or physiologically acceptable compositions comprising, an active agent, including AGONIST or ANTAGONIST of the present invention.

In a first aspect, the invention is directed to XIDATIN AGONISTS, wherein said AGONIST is an antibody that specifically binds XIDATIN, a compound excluding said XIDATIN antibody

(e.g., small organic or inorganic compound, protein, peptide, carbohydrate, lipid), or a LIGAND polypeptide or fragment thereof.

In a further preferred embodiment, the invention is directed to a XIDATIN AGONIST, wherein said AGONIST is an antibody that specifically binds XIDATIN. More preferably the  
5 invention is directed to said XIDATIN antibody, wherein said XIDATIN antibody binds XIDATIN and manifests LIGAND activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein.

In a further preferred embodiment, the invention is directed to a XIDATIN AGONIST, wherein said AGONIST is a compound excluding said XIDATIN antibody. More preferably the  
10 invention is directed to said compound, wherein said compound binds XIDATIN and manifests LIGAND activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. Further more preferably the invention is directed to said compound, wherein said compound manifests LIGAND activity  
15 exclusive of binding to XIDATIN, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. Further more preferably the invention is directed to said compound, wherein said compound increases XIDATIN expression.

In a further preferred embodiment, the invention is directed to a XIDATIN AGONIST that selectively binds to a polypeptide comprising the extracellular domain of XIDATIN.

20 In a further preferred embodiment, the invention is directed to a XIDATIN AGONIST, wherein said AGONIST is LIGAND, and wherein it is understood that LIGAND refers to a composition consisting essentially of or consisting of *in vitro* or *in vivo* self-assembling homotrimer comprised of gAPM1, gC2P, gZADJ-2, or gZADJ-7 polypeptide fragment. More preferably the invention is directed to said LIGAND, wherein said LIGAND binds XIDATIN and elicits biological  
25 activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. More preferably the invention is directed to said LIGAND, wherein said LIGAND induces, enhances, or potentiates said biological activity exclusive of binding to XIDATIN. In preferred embodiment, said homotrimer is comprised of preferred gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment.

30 APM1. Preferred gAPM1 polypeptide fragment is selected from amino acids **18-244**, **34-244**, 49-244, 56-244, 59-244, 66-244, 69-244, 78-244, 85-244, 93-244, 101-244, 102-244, 103-244, 104-244, 107-244, **110-244** or **113-244**, wherein said numbering of said amino acids within APM1 amino acid sequence is understood to be taken from said APM1 amino acid sequence presented in Table 2. Less preferred gAPM1 fragments are indicated in bold.

35 C2P. Preferred gC2P polypeptide fragment is selected from amino acids **20-333**, 25-333, 43-333, 45-333, 46-333, 50-333, 53-333, 61-333, 67-333, 74-333, 75-333, 77-333, 81-333, 82-333,

86-333, 89-333, 95-333, 100-333, 104-333, 113-333, 116-333, 125-333, 128-333, 140-333, 160-333, 164-333, 179-333, 182-333, 185-333, 188-333, 191-333, 193-333, or **202-333**, wherein said numbering of said amino acids within C2P amino acid sequence is understood to be taken from said C2P amino acid sequence presented in Table 2. Less preferred gC2P fragments are indicated in  
5 bold.

ZADJ-2. Preferred gZADJ-2 polypeptide fragment is selected from amino acids **16-285**, **25-285**, **26-285**, **29-285**, **30-285**, 91-285, 93-285, 97-285, 98-285, 99-285, 105-285, 109-285, 112-285, 120-285, 126-285, 127-285, 130-285, 132-285, 133-285, 134-285, or **150-285**, wherein said numbering of said amino acids within ZADJ-2 amino acid sequence is understood to be taken from  
10 said ZADJ-2 amino acid sequence presented in Table 2. Less preferred gZADJ-2 fragments are indicated in bold.

ZADJ-7. Preferred gZADJ-7 polypeptide fragment is selected from amino acids **31-303**, **39-303**, 78-303, 81-303, 84-303, 85-303, 88-303, 91-303, 97-303, 99-303, 109-303, 117-303, 118-303, 127-303, 139-303, 142-303, **155-303**, or **162-303**, wherein said numbering of said amino acids  
15 within ZADJ-7 amino acid sequence is understood to be taken from said ZADJ-7 amino acid sequence presented in Table 2. Less preferred gZADJ-7 fragments are indicated in bold.

More preferred LIGAND is APM1.

In a further preferred embodiment, said AGONIST is able to lower circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

20 Further preferred AGONISTS are those that significantly stimulate muscle lipid or free fatty acid oxidation as compared to untreated cells. Further preferred AGONISTS are those that cause C2C12 cells differentiated in the presence of said AGONISTS to undergo at least 10%, 20%, 30%, 35%, or 40% more oleate oxidation as compared to untreated cells.

Further preferred AGONISTS are those that increase by at least 10%, 20%, 30%, 35%, or  
25 40% leptin uptake in a liver cell line [preferably BPRCL mouse liver cells (ATCC CRL-2217)] as compared to untreated cells.

Further preferred AGONISTS are those that significantly reduce the postprandial increase in plasma free fatty acids or triglycerides, particularly following a high fat meal.

Further preferred AGONISTS are those that significantly reduce or eliminate ketone body  
30 production, particularly following a high fat meal.

Further preferred AGONISTS are those that increase glucose uptake in skeletal muscle cells.

Further preferred AGONISTS are those that increase glucose uptake in adipose cells.

Further preferred AGONISTS are those that increase glucose uptake in neuronal cells.

Further preferred AGONISTS are those that increase glucose uptake in red blood cells.

Further preferred AGONISTS are those that increase glucose uptake in the brain.

Further preferred AGONISTS are those that significantly reduce the postprandial increase in plasma glucose following a meal, particularly a high carbohydrate meal.

Further preferred AGONISTS are those that significantly prevent the postprandial increase  
5 in plasma glucose following a meal, particularly a high fat or a high carbohydrate meal.

Further preferred AGONISTS are those that improve insulin sensitivity.

Further preferred said AGONISTS are those that decrease body mass, wherein said decrease in body mass is comprised of a change in mass of the subcutaneous adipose tissue.

Further preferred said AGONISTS are those that decrease body mass, wherein said decrease  
10 in body mass is comprised of a change in mass of the visceral (omental) adipose tissue.

In a second aspect, the invention features a pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of, said AGONIST described in the first aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

In a third aspect, the invention features a method of reducing body mass comprising  
15 providing or administering to individuals in need of reducing body mass said pharmaceutical or physiologically acceptable composition described in the second aspect.

In a fourth aspect, the invention features a method of preventing or treating an obesity-related disease or disorder comprising providing or administering to an individual in need of such treatment said pharmaceutical or physiologically acceptable composition described in the second  
20 aspect. Preferably, said obesity-related disease or disorder is selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions,  
25 retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by said XIDATIN AGONIST of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human.

In related aspects, embodiments of the present invention includes methods of causing or  
30 inducing a desired biological response in an individual comprising the steps of: providing or administering to an individual a composition comprising AGONIST, wherein said biological response is selected from the group consisting of:

- (a) lowering circulating (either blood, serum, or plasma) levels (concentration) of free fatty acids;

- (b) lowering circulating (either blood, serum or plasma) levels (concentration) of glucose;
- (c) lowering circulating (either blood, serum or plasma) levels (concentration) of triglycerides;
- 5 (d) stimulating muscle lipid or free fatty acid oxidation;
- (c) increasing leptin uptake in the liver or liver cells;
- (e) reducing the postprandial increase in plasma free fatty acids, particularly following a high fat meal;
- (f) reducing or eliminating ketone body production, particularly following a high fat
- 10 meal;
- (g) increasing tissue sensitivity to insulin, particularly muscle, adipose, liver or brain,

and further wherein said biological response is significantly greater than, or at least 10%, 20%, 30%, 35%, or 40% greater than that observed in the absence of treatment; or alternatively wherein said biological response is greater than a transient response; or alternatively wherein said biological

15 response is sustained. In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or

20 physiologically acceptable composition can be used as a method to control blood glucose in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or

25 physiologically acceptable composition can be used as a method to control body weight in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or

30 physiologically acceptable composition can be used as a method to control body weight in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or

physiologically acceptable composition can be used as a method to control blood glucose in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) alone, without combination of insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) alone, without combination of insulin therapy.

5 In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control body weight in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) alone, without combination of insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or  
10 physiologically acceptable composition can be used as a method to control body weight in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) alone, without combination of insulin therapy.

In a further preferred embodiment, the present invention may be used in complementary therapy of NIDDM patients to improve their weight or glucose control in combination with an  
15 insulin secretagogue or an insulin sensitising agent. Preferably, the insulin secretagogue is 1,1-dimethyl-2-(2-morpholino phenyl)guanidine fumarate (BTS67582) or a sulphonylurea selected from tolbutamide, tolazamide, chlorpropamide, glibenclamide, glimepiride, glipizide and glidazide. Preferably, the insulin sensitising agent is selected from metformin, ciglitazone, troglitazone and pioglitazone.

20 The present invention further provides a method of improving the body weight or glucose control of NIDDM patients alone, without an insulin secretagogue or an insulin sensitising agent.

In a further preferred embodiment, the present invention may be used in complementary therapy of IDDM patients to improve their weight or glucose control in combination with an insulin secretagogue or an insulin sensitising agent. Preferably, the insulin secretagogue is 1,1-dimethyl-2-  
25 (2-morpholino phenyl) guanidine fumarate (BTS67582) or a sulphonylurea selected from tolbutamide, tolazamide, chlorpropamide, glibenclamide, glimepiride, glipizide and glidazide. Preferably, the insulin sensitising agent is selected from metformin, ciglitazone, troglitazone and pioglitazone.

The present invention further provides a method of improving the body weight or glucose  
30 control of IDDM patients alone, without an insulin secretagogue or an insulin sensitising agent.

In a further preferred embodiment, the present invention may be administered either concomitantly or concurrently, with the insulin secretagogue or insulin sensitising agent for example in the form of separate dosage units to be used simultaneously, separately or sequentially (either before or after the secretagogue or either before or after the sensitising agent). Accordingly, the  
35 present invention further provides for a composition of pharmaceutical or physiologically acceptable

composition and an insulin secretagogue or insulin sensitising agent as a combined preparation for simultaneous, separate or sequential use for the improvement of body weight or glucose control in NIDDM or IDDM patients.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition further provides a method for the use as an insulin sensitiser.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) without insulin therapy.

In a fifth aspect, the invention features a use of AGONIST described in the first aspect for treatment of obesity-related diseases and disorders and/or reducing body mass. Preferably, said obesity-related diseases and disorders are selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by said AGONIST of the invention include hyperlipidemia and hyperuricemia.

In a sixth aspect, the invention features a use of AGONIST described in the first aspect for the preparation of a medicament for the treatment of obesity-related diseases and disorders and/or for reducing body mass. Preferably, said obesity-related disease or disorder is selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related

disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human.

In a seventh aspect, the invention provides AGONIST of the first aspect of the invention, or a composition of the second aspect of the invention, for use in a method of treatment of the human  
5 or animal body.

In an eighth aspect, the invention features methods of reducing body weight comprising providing to an individual said pharmaceutical or physiologically acceptable composition described in the second aspect, or AGONIST described in the first aspect. Where the reduction of body weight is practiced for cosmetic purposes, the individual has a BMI of at least 20 and no more than 25. In  
10 embodiments for the treatment of obesity, the individual may have a BMI of at least 20. One embodiment for the treatment of obesity provides for the treatment of individuals with BMI values of at least 25. Another embodiment for the treatment of obesity provides for the treatment of individuals with BMI values of at least 30. Yet another embodiment provides for the treatment of individuals with BMI values of at least 40.

15 In further embodiment, the invention features methods of maintaining weight loss comprising providing to an individual said pharmaceutical or physiologically acceptable composition.

In a ninth aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the second aspect for reducing body mass and/or for treatment or  
20 prevention of obesity-related diseases or disorders. Preferably, said obesity-related disease or disorder is selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention  
25 include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human. In preferred embodiments, the identification of said individuals to be treated  
30 with said pharmaceutical or physiologically acceptable composition comprises genotyping LIGAND single nucleotide polymorphisms (SNPs) or measuring LIGAND polypeptide or mRNA levels in clinical samples from said individuals. Preferably, said clinical samples are selected from the group consisting of blood, serum, plasma, urine, and saliva.

In a tenth aspect, the invention features the pharmaceutical or physiologically acceptable  
35 composition described in the second aspect for reducing body weight for cosmetic reasons.

In an eleventh aspect, AGONIST of the invention is used in methods of treating insulin resistance comprising providing to an individual said pharmaceutical or physiologically acceptable composition described in the second aspect, or AGONIST described in the first aspect.

In a preferred aspect of the methods above and disclosed herein, the amount of AGONIST  
5 administered to an individual is sufficient to bring levels of XIDATIN activation to their normal levels (levels in individuals without obesity-related disease or disorder). "Normal levels" of XIDATIN activation may be followed using surrogate markers including circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

In a twelfth aspect, the invention is directed to a XIDATIN ANTAGONIST, wherein said  
10 ANTAGONIST is a soluble fragment of XIDATIN polypeptide, an antibody that specifically binds XIDATIN, a compound excluding said soluble fragment of XIDATIN polypeptide and said XIDATIN antibody (e.g., small molecular weight organic or inorganic compound, protein, peptide, carbohydrate, lipid), or a variant or fragment of LIGAND polypeptide.

In a further preferred embodiment, the invention is directed to a XIDATIN ANTAGONIST,  
15 wherein said ANTAGONIST is a soluble fragment of XIDATIN polypeptide. More preferably the invention is directed to purified, isolated, or recombinant soluble fragments of XIDATIN polypeptide. More preferably the invention is directed to said soluble fragment of XIDATIN polypeptide, wherein said soluble fragment binds LIGAND and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-  
20 like activity or described herein, and wherein said soluble fragment of XIDATIN polypeptide does not activate XIDATIN. Preferably said soluble fragment of XIDATIN polypeptide blocks or inhibits LIGAND binding to XIDATIN. In preferred embodiments, said soluble fragment of XIDATIN polypeptide comprises, consists essentially of, or consists of, at least 6 and not more than  
180 consecutive amino acids of SEQ ID NO:2, more preferably of amino acids comprising the  
25 extracellular domain of XIDATIN. Preferred said soluble fragment of XIDATIN comprises the extracellular domain of mature XIDATIN polypeptide. Particularly preferred soluble fragment of XIDATIN comprises amino acids 30-182, 30-185, 30-191, 30-193, 30-196, 30-199, 30-203 or 30-206 of SEQ ID NO:2, where it is understood that amino acid 30 is predicted to be the N-terminal amino acid of the mature XIDATIN polypeptide absent the putative signal peptide. In other  
30 preferred embodiments, said soluble fragment of XIDATIN polypeptide comprises an amino acid sequence at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the corresponding consecutive amino acids of SEQ ID NO:2. Further preferred embodiments include heterologous polypeptides comprising a XIDATIN polypeptide of the invention. In further preferred embodiment, a XIDATIN polypeptide of the invention is conjugated at its N- or C-  
35 terminus to an antibody Fc region or portion thereof.

In a further preferred embodiment, the invention is directed to a XIDATIN ANTAGONIST, wherein said ANTAGONIST is an antibody that specifically binds XIDATIN. More preferably the invention is directed to said XIDATIN antibody, wherein said XIDATIN antibody binds XIDATIN and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said XIDATIN antibody does not activate XIDATIN. Preferably said XIDATIN antibody blocks or inhibits LIGAND binding to XIDATIN.

In a further preferred embodiment, the invention is directed to a XIDATIN ANTAGONIST, wherein said ANTAGONIST is a compound excluding said soluble fragment of XIDATIN polypeptide and said XIDATIN antibody (e.g., small organic molecule, protein, peptide). More preferably the invention is directed to said compound, wherein said compound binds to XIDATIN and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not activate XIDATIN. Preferably said compound that binds to XIDATIN blocks or inhibits LIGAND binding to XIDATIN. Further more preferably the invention is directed to said compound, wherein said compound blocks or inhibits LIGAND activity exclusive of binding to XIDATIN, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not activate XIDATIN. Further more preferably the invention is directed to said compound, wherein said compound blocks or inhibits XIDATIN expression and wherein said compound does not have LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not activate XIDATIN.

In a further preferred embodiment, the invention is directed to a XIDATIN ANTAGONIST, wherein said ANTAGONIST is a variant or fragment of LIGAND polypeptide. More preferably the invention is directed to said variant or fragment of LIGAND polypeptide, wherein said variant or fragment of LIGAND polypeptide binds XIDATIN and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said variant or fragment of LIGAND polypeptide does not activate XIDATIN. Preferably said variant or fragment of LIGAND polypeptide blocks or inhibits LIGAND binding to XIDATIN. More preferably the invention is directed to said variant or fragment of LIGAND polypeptide, wherein said variant or fragment of LIGAND polypeptide inhibits the induction, enhancement, or potentiation of said biological activity exclusive of binding to XIDATIN.

In a further preferred embodiment, the invention is directed to a XIDATIN ANTAGONIST that selectively binds to a polypeptide comprising the extracellular domain of XIDATIN.

In a further preferred embodiment, said ANTAGONIST is able to raise circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

Further preferred said ANTAGONISTS are those that significantly inhibit muscle lipid or  
5 free fatty acid oxidation stimulated by its LIGAND. Further preferred said ANTAGONISTS are those that cause C2C12 cells differentiated in the presence of LIGAND to undergo at least 10%, 20%, 30%, 35%, or 40% less oleate oxidation as compared to untreated cells.

Further preferred said ANTAGONISTS are those that inhibit by at least 10%, 20%, 30%, 35%, or 40% the increase in leptin uptake stimulated by LIGAND polypeptide in a liver cell line  
10 [preferably BPRCL mouse liver cells (ATCC CRL-2217)] as compared to untreated cells.

Further preferred said ANTAGONISTS are those that significantly increase the postprandial increase in plasma free fatty acids, particularly following a high fat meal.

Further preferred said ANTAGONISTS are those that significantly increase ketone body production, particularly following a high fat meal.

15 Further preferred said ANTAGONISTS are those that decrease glucose uptake in skeletal muscle cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in adipose cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in neuronal  
20 cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in red blood cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in the brain stimulated by LIGAND.

25 Further preferred said ANTAGONISTS are those that significantly increase the postprandial increase in plasma glucose following a meal, particularly a high carbohydrate meal.

Further preferred said ANTAGONISTS are those that significantly facilitate the postprandial increase in plasma glucose following a meal, particularly a high fat or a high carbohydrate meal.

30 Further preferred said ANTAGONISTS are those that reduce the insulin sensitivity stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that increase body mass, wherein said increase in body mass is comprised of a change in mass of the subcutaneous adipose tissue.

Further preferred said ANTAGONISTS are those that increase body mass, wherein said increase in body mass is comprised of a change in mass of the visceral (omental) adipose tissue.

In a thirteenth aspect, the invention features a pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of, said ANTAGONIST described in  
5 the twelfth aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

In a fourteenth aspect, the invention features a method of increasing body mass comprising providing or administering to individuals in need of increasing body mass said pharmaceutical or physiologically acceptable composition described in the thirteenth aspect.

In a fifteenth aspect, the invention features a method of preventing or treating disorders  
10 associated with excessive weight loss comprising providing or administering to an individual in need of such treatment said pharmaceutical or physiologically acceptable composition described in the thirteenth aspect. Preferably said disorder is selected from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. Said disorders associated with excessive weight loss are  
15 comprised of those mediated by tumor necrosis factor (TNFalpha) alone, those mediated by TNFalpha plus one or more additional factors, and those mediated only by one or more factors exclusive of TNFalpha. Said factors include, but are not restricted to, macrophage migration inhibitory factor, interleukin 1, and interleukin 6. In preferred embodiments, said individual is a mammal, preferably a human.

20 In related aspects, embodiments of the present invention includes methods of causing or inducing a desired biological response in an individual comprising the steps of: providing or administering to an individual a composition comprising ANTAGONIST, wherein said biological response is selected from the group consisting of:

- 25
- (a) raising circulating (either blood, serum, or plasma) levels (concentration) of free fatty acids (FFA) or triglycerides (TG);
  - (b) raising circulating (either blood, serum or plasma) levels (concentration) of glucose;
  - (c) raising circulating (either blood, serum or plasma) levels (concentration) of triglycerides;
  - (d) inhibiting muscle lipid or free fatty acid oxidation;
  - 30 (c) inhibiting leptin uptake in the liver or liver cells;
  - (e) increasing the postprandial increase in plasma free fatty acids, particularly following a high fat meal; and,
  - (f) increasing or eliminating ketone body production, particularly following a high fat meal;

(g) reducing tissue sensitivity to insulin, particularly muscle, adipose, liver or brain, and further wherein said biological response is greater than a transient response; or alternatively wherein said biological response is sustained. In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method of increasing body mass in some persons with cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition further provides a method for the use as an insulin desensitiser, wherein the sensitivity of a cell or tissue to insulin is reduced.

10 In a sixteenth aspect, the invention features a method of making the XIDATIN polypeptide described in the twelfth aspect, wherein said method is selected from the group consisting of proteolytic cleavage, recombinant methodology and artificial synthesis. In a preferred embodiment, proteolytic cleavage is carried out using trypsin, plasmin, or collagenase.

In a seventeenth aspect, the invention features a use of ANTAGONIST described in the 15 twelfth aspect for the preparation of a medicament for the treatment of disorders associated with excessive weight loss and/or for increasing body mass. Preferably, said disorder is selected from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. In preferred embodiments, said individual is a mammal, preferably a human.

20 In an eighteenth aspect, the invention provides ANTAGONIST of the twelfth aspect of the invention, or a composition of the thirteenth aspect of the invention, for use in a method of treatment of the human or animal body.

In a nineteenth aspect, the invention features methods of increasing body weight comprising providing to an individual said pharmaceutical or physiologically acceptable composition described 25 in the thirteenth aspect, or ANTAGONIST described in the twelfth aspect. Where the increase of body weight is practiced for cosmetic purposes, the individual has a BMI of no greater than 25 and at least 20. In embodiments for the treatment of disorders associated with excessive weight loss, the individual may have a BMI no greater than 20. One embodiment for the treatment of disorders associated with excessive weight loss provides for the treatment of individuals with BMI values of 30 no greater than 15. Alternatively, for increasing the body weight of an individual, the BMI value should be at least 15 and no more than 20.

In a twentieth aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the thirteenth aspect for increasing body mass and/or for treatment of disorders associated with excessive weight loss. Preferably, said disorder is selected 35 from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight

loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. In preferred embodiments, said individual is a mammal, preferably a human.

In a twenty-first aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the thirteenth aspect for increasing body weight for cosmetic reasons.

In a preferred aspect of the methods above and disclosed herein, the amount of ANTAGONIST administered to an individual is sufficient to bring levels of XIDATIN activation to their normal levels (levels in healthy individuals). "Normal levels" of XIDATIN activation may be followed using surrogate markers including circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

#### Brief Description of Tables

Table 1 lists known or predicted biologic structural and functional domains for the XIDATIN polypeptide of SEQ ID NO:2 of the present invention, including the signal peptide, extracellular (EC) domain, transmembrane domain, and intracellular (IC) domain.

Table 2 lists the amino acid sequence of full-length APM1, C2P, ZADJ-2 and ZADJ-7 polypeptide. The total number of amino acids is given in parentheses. The predicted signal peptide is indicated in bold. The collagen-like region is indicated by dotted line. The region between the predicted signal peptide and the collagen-like region is the N-terminally disposed unique region. The globular C-terminal C1q homology domain is indicated by single underline. The NGLXXD amino acid motif C-terminally disposed within the globular domain is indicated by double underline. It is taken to be understood that C2P herein encompasses variants comprising the substitution of valine for methionine at position 219 and/or the substitution of methionine for valine at position 301.

#### Structure of XIDATIN Polypeptide

The full-length XIDATIN polypeptide is comprised of at least 4 distinct regions including:

1. an N-terminal putative signal peptide comprising amino acids from about amino acids 1-29 of SEQ ID NO:2;
2. an extracellular domain comprising a LIGAND binding portion and comprising amino acids from about amino acids 30-209 of SEQ ID NO:2;
3. a transmembrane domain comprising amino acids from about amino acids 210-232 of SEQ ID NO:2; and
4. an intracellular domain comprising amino acids from about amino acids 233-386 of SEQ ID NO:2.

#### Brief Description of Sequence Listing

SEQ ID NO:1 is the nucleotide sequence of cDNA with an open reading frame which location is indicated as features. When appropriate, the locations of the potential polyadenylation site and polyadenylation signal are also indicated.

SEQ ID NO:2 is the amino acid sequence of polypeptide encoded by the cDNA of SEQ ID NO:1.

The appended Sequence Listing is hereby incorporated by reference in its entirety.

#### Detailed Description

##### Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

The term "isolated" requires that the material be removed from its original environment (e.g., the natural environment if the material is naturally occurring).

The term "purified" does not require absolute purity; rather, it is intended as a relative definition. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

As used interchangeably herein, the term "polynucleotide(s)" include RNA or DNA (either single or double stranded, coding, complementary or antisense), or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or duplex form (although each of the above species may be particularly specified).

The terms "complementary" or "complement thereof" are used herein to refer to the sequences of polynucleotides that are capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region.

The terms "polypeptide" and "protein", used interchangeably herein, refer to a polymer of amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude chemical or post-expression modifications of the polypeptides of the invention, although chemical or post-expression modifications of these polypeptides may be included excluded as specific embodiments.

As used herein, the terms "recombinant polynucleotide" and "polynucleotide construct" are used interchangeably to refer to linear or circular, purified or isolated polynucleotides that have been artificially designed and which comprise at least two nucleotide sequences that are not found as contiguous nucleotide sequences in their initial natural environment. In particular, these terms mean

that the polynucleotide or cDNA is adjacent to "backbone" nucleic acid to which it is not adjacent in its natural environment.

The term "recombinant polypeptide" is used herein to refer to polypeptides that have been artificially designed and which comprise at least two polypeptide sequences that are not found as  
5 contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides which have been expressed from a recombinant polynucleotide.

As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship.

As used herein, the term "non-human animal" refers to any non-human animal, including  
10 insects, birds, rodents and more usually mammals. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

The term "domain" refers to an amino acid fragment with specific biological properties. This term encompasses all known structural and linear biological motifs.

As used herein, the term "receptor" refers to a polypeptide to which a "ligand" binds and  
15 through which said "ligand" elicits a biological response comprised of biological activities. Said receptor is preferably XIDATIN of the present invention. Said "ligand" is preferably LIGAND of the present invention. By "receptor activation" is intended "ligand"-mediated alteration of said receptor polypeptide, wherein said alteration is selected from but not limited to the group consisting of receptor alterations associated with said biological response.

20 As used herein, the term "AGONIST" refers to naturally occurring and synthetic compounds capable of inducing, enhancing, or potentiating a biological response comprised of biological activities.

As used herein, the term "ANTAGONIST" refers to naturally occurring and synthetic  
25 compounds capable of inhibiting a biological response, inhibiting the induction of a biological response, or inhibiting the potentiation of a biological response, wherein said biological response is comprised of biological activities.

Without being limited by theory, the compounds/polypeptides of the invention are capable of modulating the partitioning of dietary lipids between the liver and peripheral tissues, and are thus believed to treat "diseases involving the partitioning of dietary lipids between the liver and  
30 peripheral tissues." The term "peripheral tissues" is meant to include muscle and adipose tissue. In preferred embodiments, the compounds/polypeptides of the invention partition the dietary lipids toward or away from the muscle. In alternative preferred embodiments, the dietary lipids are partitioned toward or away from the adipose tissue. In other preferred embodiments, the dietary lipids are partitioned toward or away from the liver. In yet other preferred embodiments, the  
35 compounds/polypeptides of the invention increase or decrease the oxidation of dietary lipids,

preferably free fatty acids (FFA) by the muscle. Dietary lipids include, but are not limited to triglycerides and free fatty acids.

Preferred diseases believed to involve the partitioning of dietary lipids include obesity-related diseases and disorders such as obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. Yet other disorders of the invention include disorders associated with excessive weight loss such as cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia.

The terms "comprising", "consisting of" and "consisting essentially of" may be interchanged for one another throughout the instant application, although each retains its normal definition. The term "having" has the same meaning as "comprising" and may be replaced with either the term "consisting of" or "consisting essentially of".

#### Polypeptides of the Invention

Preferably, polypeptides of the invention are recombinantly produced using routine expression methods known in the art. The polynucleotide encoding the desired polypeptide is operably linked to a promoter into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems are used in forming recombinant polypeptides. The polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use.

Consequently, a further embodiment of the present invention is a method of making a polypeptide, said method comprising the steps of

- a) obtaining a cDNA encoding said polypeptide;
- b) inserting said cDNA in an expression vector such that the cDNA is operably linked to a promoter; and
- c) introducing said expression vector into a host cell whereby said host cell produces said polypeptide.

In one aspect of this embodiment, the method further comprises the step of isolating the polypeptide. Another embodiment of the present invention is a polypeptide obtainable by the method described in the preceding paragraph.

The expression vector is any of the mammalian, yeast, insect or bacterial expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). In preferred embodiment, 5 recombinant polypeptides of the invention are expressed in mammalian cells.

The invention is drawn, *inter alia*, to isolated, purified or recombinant polypeptides. XIDATIN polypeptides of the invention are useful for increasing (ANTAGONISTS of XIDATIN) body weight either as a cosmetic treatment or for treatment or prevention of diseases and disorders as discussed or described herein. XIDATIN polypeptides are also useful *inter alia* in screening 10 assays for AGONISTS or ANTAGONISTS of XIDATIN activity and for raising XIDATIN-specific antibodies. When used for cosmetic treatments, or for the treatment or prevention of diseases, disorders, or conditions, one or more XIDATIN polypeptides can be provided to a subject. Thus, various fragments of the full-length protein can be combined into a "cocktail" for use in the various treatment regimens. LIGAND polypeptides of the invention are useful for reducing (AGONISTS of 15 XIDATIN) body weight either as a cosmetic treatment or prevention of diseases and disorders as discussed or described herein.

The XIDATIN polypeptides of the present invention are preferably provided in an isolated form, and may be partially or substantially purified.

#### Modifying XIDATIN biological activity

20 Modifying endogenous XIDATIN biological activity is expressly contemplated by the present invention. The present invention further relates to compounds able to modulate XIDATIN biological activity and methods to use these compounds. Such compounds may interact with XIDATIN polypeptides directly or indirectly.

#### Candidate AGONISTS and ANTAGONISTS Obtained by Optical Biosensor Methods

25 Compounds interacting with a polypeptide comprising XIDATIN extracellular domain can be screened by using an Optical Biosensor as described in Edwards and Leatherbarrow (1997) and also in Szabo *et al.* (1995), the disclosures of which are incorporated by reference. This technique permits the detection of interactions between molecules in real time, without the need of labeled molecules. This technique, which is based on the surface plasmon resonance (SPR) phenomenon, is 30 presented in more detail in Example 1.

#### Compounds Modulating XIDATIN Biological Activity

Another method of screening for compounds that modulate XIDATIN biological activity is by measuring the effects of test compounds on specific biological activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or 35 as described herein, in a host cell. In one embodiment, the present invention relates to a method of

identifying an agent that alters XIDATIN activity, wherein a nucleic acid construct comprising the polynucleotide of SEQ ID NO:1 or a fragment thereof encoding full-length XIDATIN polypeptide is introduced into a mammalian host cell. The transfected mammalian host cells are maintained under conditions appropriate for expression of the encoded XIDATIN, whereby the nucleic acid is  
5 expressed. The host cells are then contacted with a compound to be assessed (an agent) and an activity of the cells is detected in the presence of the compound to be assessed, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein. Detection of a change in said activity for said transfected host cell, but not in untransfected host cell, in the presence of the agent indicates that the agent alters XIDATIN activity.  
10 In a particular embodiment, the invention relates to a method of identifying an agent which is an activator (AGONIST) of XIDATIN activity, wherein detection of an increase of said activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, in the presence of the agent indicates that the agent activates XIDATIN activity. In another particular embodiment, the invention relates to a method of  
15 identifying an agent which is an inhibitor (ANTAGONIST) of XIDATIN activity, wherein detection of a decrease of said activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, in the presence of the agent indicates that the agent inhibits XIDATIN activity.

Detection of a change in said XIDATIN activity, said activity being selected from the group  
20 consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, can be performed using a variety of techniques as described for representative activities in Examples provided herein.

In a particular embodiment a high throughput screen can be used to identify agents that activate (enhance) or inhibit XIDATIN activity (See e.g., PCT publication WO 98/45438, which  
25 disclosure is hereby incorporated by reference in its entirety).

#### Methods of Screening for Compounds Modulating XIDATIN Activity

The present invention also relates to methods of screening compounds for their ability to modulate (e.g. increase or inhibit) the activity or expression of XIDATIN. More specifically, the present invention relates to methods of testing compounds for their ability either to increase or to  
30 decrease activity of XIDATIN. The assays are performed *in vitro* or *in vivo*.

The present invention relates to a method for the screening of a candidate substance for interaction with a polypeptide comprising XIDATIN extracellular domain, said method comprising the following steps:

- a) providing said polypeptide comprising XIDATIN extracellular domain;
- 35 b) obtaining a candidate substance;

- c) bringing into contact said polypeptide with said candidate substance;
- d) detecting the complexes formed between said polypeptide and said candidate substance.

The invention further relates to a method for the production of a pharmaceutical composition comprising a method for the screening of a candidate substance that interact with a XIDATIN polypeptide, fragments or variants thereof and furthermore mixing the identified substance with a pharmaceutically acceptable carrier.

The present invention relates to a method for the screening of a candidate substance for the capacity to increase expression of XIDATIN, said method comprising the following steps:

- 10 a) isolating mRNA from cells which have or have not been contacted with said candidate substance;
- b) carrying out a Northern blot analysis with labeled cDNA probe encoding all or part of XIDATIN polypeptide;
- 15 c) wherein increased signal in cells having been contacted with said candidate substance over that of uncontacted cells is taken to indicate that said candidate substance increases expression of XIDATIN and is an AGONIST of XIDATIN activity; and
- d) wherein decreased signal in cells having been contacted with said candidate substance over that of uncontacted cells is taken to indicate that said candidate substance decreases expression of XIDATIN and is an ANTAGONIST of XIDATIN activity.
- 20

Methods of isolating mRNA and carrying out Northern blot analysis are well known to those of ordinary skill in the art.

#### Preparation of Antibody Compositions

25 Substantially pure protein or polypeptide is isolated from transfected or transformed cells containing an expression vector encoding the XIDATIN protein or a portion thereof. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the protein can then be prepared by methods well known to those of ordinary skill in the art.

30 Preferably the present invention includes monoclonal and polyclonal antibodies that specifically bind XIDATIN polypeptide fragment comprising the extracellular domain of mature XIDATIN polypeptide. Particularly preferred soluble fragment of XIDATIN comprises amino acids 30-182, 30-185, 30-191, 30-193, 30-196, 30-199, 30-203 or 30-206 of SEQ ID NO:2, where it is

understood that amino acid 30 is predicted to be the N-terminal amino acid of the mature XIDATIN polypeptide absent the putative signal peptide.

#### EXAMPLES

The following Examples are provided for illustrative purposes and not as a means of limitation. One of ordinary skill in the art would be able to design equivalent assays and methods based on the disclosure herein all of which form part of the instant invention.

##### EXAMPLE 1: Use of Biacore Technology to Detect Specific Binding of a Test Compound to Polypeptide Fragment Comprising XIDATIN Extracellular Domain

Biacore utilizes a biosensor technology for monitoring interactions between two or more molecules in real time, without the use of labels. The molecular classes that can be studied are diverse, ranging from proteins, peptides, nucleic acids, carbohydrates, and lipids to low molecular weight substances and pharmaceuticals.

The detection principle is based on the optical phenomena of surface plasmon resonance, which detects changes in refractive index close to a biosensor surface. In a typical experiment one of the interacting molecules is immobilized or captured (here, polypeptide fragment comprising XIDATIN extracellular domain) to a flexible dextran layer close to the sensor surface. The interacting partner (here, test compound) is flowed across that surface. If an interaction occurs between the two molecules, there is a resulting increase in signal due to the increase in mass at the chip surface.

Soluble polypeptide fragment comprising XIDATIN extracellular domain is attached to the sensor surface via amine coupling chemistry. The dextran is activated using N-hydroxysuccinimide and N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride for 7 minutes. Said XIDATIN polypeptide fragment is diluted in 10mM Na Acetate pH 5.0 at a concentration of 10µg/ml and injected over the activated surface for 7 minutes. The surface is then blocked for 7 minutes using ethanolamine to remove any remaining esters. A blank flow cell absent said XIDATIN polypeptide fragment is set up in parallel and used as a control surface. The running buffer is HBS-EP (0.01M HEPES pH 7.4, 0.15M NaCl, 3mM EDTA, 0.005% Surfactant P20) and the instrument temperature is 25°C.

The test compound is filtered through an Ultrafree-0.5 Centrifugal Filter Device and resuspended in HBS-EP running buffer. The test compound is then diluted 1:10 in HBS-EP and injected over the said XIDATIN polypeptide fragment surface and the blank control surface for 1 minute at a flow rate of 50 µl/min. The sensorgrams from the receptor surface and the control surface are aligned and overlaid.

To obtain the specific binding, the control surface was subtracted from the active surface comprised of said XIDATIN polypeptide fragment.

Example 2: Effect of LIGAND on Muscle Cell Fatty Acid Oxidation

C2C12 cells are differentiated in the presence or absence of 2 µg/mL LIGAND for 4 days. On day 4, oleate oxidation rates are determined by measuring conversion of 1-<sup>14</sup>C-oleate (0.2 mM) to <sup>14</sup>CO<sub>2</sub> for 90 min. This experiment can be used to screen for active polypeptides and peptides as well as AGONISTS and ANTAGONISTS or activators and inhibitors of LIGAND receptor.

The effect of LIGAND on the rate of oleate oxidation can be compared in differentiated C2C12 cells (murine skeletal muscle cells; ATCC, Manassas, VA CRL-1772) and in a hepatocyte cell line (Hepa1-6; ATCC, Manassas, VA CRL-1830). Cultured cells are maintained according to manufacturer's instructions. The oleate oxidation assay is performed as previously described (Muoio et al (1999) Biochem J 338;783-791). Briefly, nearly confluent myocytes are kept in low serum differentiation media (DMEM, 2.5% Horse serum) for 4 days, at which time formation of myotubes became maximal. Hepatocytes are kept in the same DMEM medium supplemented with 10% FCS for 2 days. One hour prior to the experiment the media is removed and 1 mL of preincubation media (MEM, 2.5% Horse serum, 3 mM glucose, 4 mM Glutamine, 25 mM Hepes, 1% FFA free BSA, 0.25 mM Oleate, 5 µg/mL gentamycin) is added. At the start of the oxidation experiment <sup>14</sup>C-Oleic acid (1µCi/mL, American Radiolabelled Chemical Inc., St. Louis, MO) is added and cells are incubated for 90 min at 37°C in the absence/presence of 2.5 µg/mL LIGAND. After the incubation period 0.75 mL of the media is removed and assayed for <sup>14</sup>C-oxidation products as described below for the muscle FFA oxidation experiment.

20 EXAMPLE 3: Effect of LIGAND on *In Vitro* Glucose Uptake by Muscle Cells

L6 Muscle cells are obtained from the European Culture Collection (Porton Down) and are used at passages 7-11. Cells are maintained in standard tissue culture medium DMEM, and glucose uptake is assessed using [<sup>3</sup>H]-2-deoxyglucose (2DG) with or without LIGAND in the presence or absence of insulin (10<sup>-8</sup> M) as has been previously described (Walker, P.S. et al. (1990) Glucose transport activity in L6 muscle cells is regulated by the coordinate control of subcellular glucose transporter distribution, biosynthesis, and mRNA transcription. JBC 265(3):1516-1523; and Kilp, A. et al. (1992) Stimulation of hexose transport by metformin in L6 muscle cells in culture. Endocrinology 130(5):2535-2544, which disclosures are hereby incorporated by reference in their entirety). Uptake of 2DG is expressed as the percentage change compared with control (no added insulin or LIGAND). Values are presented as mean ± SEM of sets of 4 wells per experiment. Differences between sets of wells are evaluated by Student's t test, probability values p<0.05 are considered to be significant.

EXAMPLE 4: Effect of LIGAND on Mice Fed a High-Fat Diet

Experiments are performed using approximately 6 week old C57Bl/6 mice (8 per group). All mice are housed individually. The mice are maintained on a high fat diet throughout each experiment. The high fat diet (cafeteria diet; D12331 from Research Diets, Inc.) has the following

composition: protein kcal% 16, sucrose kcal% 26, and fat kcal% 58. The fat is primarily composed of coconut oil, hydrogenated.

After the mice are fed a high fat diet for 6 days, micro-osmotic pumps are inserted using isoflurane anesthesia, and are used to provide LIGAND, saline, and an irrelevant peptide to the mice subcutaneously (s.c.) for 18 days. LIGAND is provided at doses of 100, 50, 25, and 2.5  $\mu\text{g}/\text{day}$  and the irrelevant peptide is provided at 10  $\mu\text{g}/\text{day}$ . Body weight is measured on the first, third and fifth day of the high fat diet, and then daily after the start of treatment. Final blood samples are taken by cardiac puncture and are used to determine triglyceride (TG), total cholesterol (TC), glucose, leptin, and insulin levels. The amount of food consumed per day is also determined for each group.

10 EXAMPLE 5: Effect of LIGAND on Plasma Free Fatty Acid in C57 BL/6 Mice

The effect of LIGAND on postprandial lipemia (PPL) in normal C57BL6/J mice is tested.

The mice used in this experiment are fasted for 2 hours prior to the experiment after which a baseline blood sample is taken. All blood samples are taken from the tail using EDTA coated capillary tubes (50  $\mu\text{L}$  each time point). At time 0 (8:30 AM), a standard high fat meal (6g butter, 6 g sunflower oil, 10 g nonfat dry milk, 10 g sucrose, 12 mL distilled water prepared fresh following Nb#6, JF, pg.1) is given by gavage (vol.=1% of body weight) to all animals.

Immediately following the high fat meal, 25 $\mu\text{g}$  a LIGAND is injected i.p. in 100  $\mu\text{L}$  saline. The same dose (25 $\mu\text{g}/\text{mL}$  in 100 $\mu\text{L}$ ) is again injected at 45 min and at 1 hr 45 min. Control animals are injected with saline (3x100 $\mu\text{L}$ ). Untreated and treated animals are handled in an alternating mode.

Blood samples are taken in hourly intervals, and are immediately put on ice. Plasma is prepared by centrifugation following each time point. Plasma is kept at  $-20^{\circ}\text{C}$  and free fatty acids (FFA), triglycerides (TG) and glucose are determined within 24 hours using standard test kits (Sigma and Wako). Due to the limited amount of plasma available, glucose is determined in duplicate using pooled samples. For each time point, equal volumes of plasma from all 8 animals per treatment group are pooled.

EXAMPLE 6: Effect of LIGAND on Plasma FFA, TG and Glucose in C57 BL/6 Mice

Briefly, 14 mice re fasted for 2 hours prior to the experiment after which a baseline blood sample is taken. All blood samples are taken from the tail using EDTA coated capillary tubes (50  $\mu\text{L}$  each time point). At time 0 (9:00AM), a standard high fat meal (see Example 6) is given by gavage (vol.=1% of body weight) to all animals. Immediately following the high fat meal, 4 mice are injected 25  $\mu\text{g}$  of LIGAND i.p. in 100 $\mu\text{L}$  saline. The same dose (25 $\mu\text{g}$  in 100 $\mu\text{L}$ ) is again injected at 45 min and at 1 hr 45 min. A second treatment group receives 3 times 50  $\mu\text{g}$  LIGAND at the same intervals. Control animals are injected with saline (3x100 $\mu\text{L}$ ). Untreated and treated animals are handled in an alternating mode.

Blood samples are immediately put on ice. Plasma is prepared by centrifugation following each time point. Plasma is kept at  $-20^{\circ}\text{C}$  and free fatty acids (FFA), triglycerides (TG) and glucose

are determined within 24 hours using standard test kits (Sigma and Wako).

EXAMPLE 7: Effect of LIGAND on FFA following Epinephrine Injection

In mice, plasma free fatty acids increase after intragastric administration of a high fat/sucrose test meal. These free fatty acids are mostly produced by the activity of lipolytic enzymes  
5 *i.e.* lipoprotein lipase (LPL) and hepatic lipase (HL). In this species, these enzymes are found in significant amounts both bound to endothelium and freely circulating in plasma. Another source of plasma free fatty acids is hormone sensitive lipase (HSL) that releases free fatty acids from adipose tissue after  $\beta$ -adrenergic stimulation. To test whether LIGAND also regulates the metabolism of free fatty acid released by HSL, mice are injected with epinephrine.

10 Two groups of mice are given epinephrine (5 $\mu$ g) by intraperitoneal injection. A treated group is injected with a LIGAND (25 $\mu$ g) one hour before and again together with epinephrine, while control animals receive saline. Plasma is isolated and free fatty acids and glucose are measured.

EXAMPLE 8: Effect of LIGAND on FFA following Intralipid Injection

Two groups of mice are intravenously (tail vein) injected with 30  $\mu$ L bolus of Intralipid-20%  
15 (Clintec) to generate a sudden rise in plasma FFAs, thus by-passing intestinal absorption. (Intralipid is an intravenous fat emulsion used in nutritional therapy). A treated group (LIGAND-treated) is injected with LIGAND (25 $\mu$ g) at 30 and 60 minutes before Intralipid is given, while control animals receive saline. Plasma is isolated and FFAs are measured as described previously. The effect of LIGAND on the decay in plasma FFAs following the peak induced by Intralipid injection is then  
20 monitored.

EXAMPLE 9: Effect of LIGAND on Weight Gain and Weight Loss of Mice and on Maintenance of Weight Loss in Mice

In the first experiment, 10-week-old male C57BL/6J mice are put on a very high fat/sucrose purified diet for 19 days to promote weight gain; the average body weight at this time is 30g. The  
25 mice are then surgically implanted with an osmotic pump (Alzet, Newark, DE) delivering either 2.5  $\mu$ g/day of LIGAND or physiological saline. The mice are continued on the high fat diet and their body weight was recorded over the following 10-day period.

Weight gain by mice treated with saline in contradistinction to weight loss by mice treated with LIGAND is taken as evidence that in this inbred strain of normal mice, a continuous infusion of  
30 a daily low dose of LIGAND can prevent weight gain caused by high fat/sucrose feeding, in a sustainable way.

Data are expressed throughout as mean  $\pm$  SEM; a p-value  $< 0.05$  is considered statistically significant. Statistical analysis is typically done using either the unpaired Student's t test or the paired Student's t test.

35 Maintenance of weight loss in mice

In order to demonstrate the ability of LIGAND to maintain weight loss, normal mice are put on a reduced calorie diet to promote weight loss. The reduced calorie diet is continued until the

mice lose 10% of their initial weight. A second group of mice are continued on the reduced calorie diet until the mice lose 20% of their initial weight. The mice are then surgically implanted with an osmotic pump (Alzet, Newark, DE) delivering either 2.5 µg/day of LIGAND or physiological saline. The mice are returned to a normal diet and their body weights are recorded over a 10-day period.

- 5 After 10 days, the outcome wherein mice treated with LIGAND have a lower weight than mice treated with saline is taken to provide evidence that treatment with LIGAND promotes the maintenance of weight loss.

EXAMPLE 10: Assessment of homotrimer formation by gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment.

- 10 Homotrimer formation by gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment is assessed using sedimentation equilibrium in analytical centrifuges, a method that determines molecular weight accurately and independently of other physical factors such as shape.

- Candidate gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment homotrimer is purified, for example using a protocol comprising a method of gel filtration such as 16/60 superdex  
 15 200 gel filtration column (Amersham). Said purified candidate gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment homotrimer protein concentration is made 3 µM in 5.7 mM phosphate (pH 7.5), 137 mM NaCl, 2.7 mM KCl. Samples are centrifuged at 8,000 rpm for 18 hours at 10°C in a Beckman XL-A analytical ultracentrifuge before absorbance is recorded. The data are fit globally, using MacNonlin PPC [Johnson ML et al., Biophys J (1981) 36:575-8; Schuster TM et al., Curr  
 20 Opin Struct Biol (1996) 6:650-8; Hensley P, Structure (1996) 4:367-73; the disclosures of which are incorporated herein by reference in their entirety] to the following equation that describes the sedimentation of a homogeneous species:  $Abs = B + A' \exp[H \times M (x^2 - x_0^2)]$  where Abs = absorbance at radius x, A' = absorbance at reference radius  $x_0$ ,  $H = (1 - \nu\rho)\omega^2/2RT$ , R = gas constant, T = temperature in Kelvin,  $\nu$  = partial specific volume = 0.71896131 mL/g,  $\rho$  = density of solvent =  
 25 1.0061 g/ml,  $\omega$  = angular velocity in radians/s, M = apparent molecular weight, and B = solvent absorbance (blank).

TABLE 1

Amino Acid Residues Comprising the Structural Domains of XIDATIN

- 30 SEQ ID NO:2 Description

SIGNAL PEPTIDE	EC DOMAIN	TRANSMEMBRANE DOMAIN	IC DOMAIN
1-29	30-209	210-232	233-386

EC, extracellular domain; IC, intracellular domain

## TABLE 2

APM1, C2P, ZADJ-2 and ZADJ-7

5 &gt;APM1 polypeptide sequence:

MLLLGAVLLLLALPGHDQETTTQGPVLLPLPKGACTGWMAGIPGHPGHNGAPGRDGRD  
GTPGEKGEKGDPGLIGPKGDIGETGVPGAEGPRGFPGIQGRKGEPGEGAYVYRSAFSVGLT  
YVTIPNMPIRFTKIFYNQONHYDGSTGKFHCNIPGLYYFAYHITVYMKDVKVSFLFKDKAM  
LFTYDQYQENNVDQASGSVLLHLEVGDQVWLQVYGEGERNGLYADNDNDSTFTGFLLYH

10 DTN (244)

&gt; C2P polypeptide sequence:

MRIWWLLLAIEICTGNINSQDTCRQGHPIGNP GHNGLPGRDGRD GAKGDKGDAGEPG  
RPGSPGKDGTSGEKGERGADGKVEAKGIKGDQGSRGSPPGKHGPKGLAGPMGEKGLRGETG  
15 PQGQKGNKGDVGP TGPEGPRGNIGPLGPTGLPGMPGPIGKPGPKGEAGPTGPQGE PGVVRGIR  
GWK GDRGEK GKIGETLVLPKSAFTVGLTVLSKFPSSDMPIKFDKILYNEFNHYDTAAGKFTC  
HIAGVYYFTYHITVFSRNVQVSLVKNGVKILHTKDAYMSSEDQASGGIVLQLKLGDEVWLQ  
VTGGERFNGLFADEDDDTTFTGFLLFSSP (333)

20 &gt;ZADJ-2 polypeptide sequence:

MIPWVLLACALPCAADPLLGA FARRDFRK GSPQLVCSLPGPQPPGPPGAPGPSGMMGRM  
GFP GKDGQDGH DGRDSDGEEGPPGRTGNRGKPGPKGKAGAI GRAGPRGPKGVNGTPGK  
HGTPGKKGPKGKKGEPGLPGPCSCGSGHTKSAFSVAVTKSYPRERLPIKFDKILMNEGGHY  
NASSGKFVCGVPGIYYFTYDITLANKHLAIGLVHNGOYRIRTFDANTGNHDTVASGSTILALK  
25 QGDEVWLQIFYSEQNGLFYDPYWTDSLFTGFLIYADQDDPNEV (285)

&gt;ZADJ-7 polypeptide sequence:

MGKEDTQETRTEPKMFVLLYVTSFAICASGQPRGNQLKGENYSPRYICSIPGLPGPPGPPG  
ANGSPGPHGRIGLPGRDGRDGRKGEKGEKGTAGLRGKTGPLGLAGEKGDQGETGKKGPIG  
30 PEGEKGEVGPPIGPPGPKGDRGEQGDPLPGVCRCSIVLKS AFSVGITTSYPEERLPIIFNKVL  
FNEGEHYNPATGKFICAFPGIYYFSYDITLANKHLAIGLVHNGOYRIKTFDANTGNHDTVASG  
STVIYLOPEDEVWLEIFFTDQNGLES DPGWADSLFSGFLLYVDTDYLD SISEDDEL (303)

What is claimed is:

1. A method of screening for an AGONIST or an ANTAGONIST of XIDATIN activity.
- 5 2. The method of Claim 1, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, insulin-like activity, free fatty acid oxidation, and weight reduction.
3. An AGONIST or an ANTAGONIST of XIDATIN activity.
4. The AGONIST or the ANTAGONIST of Claim 3, wherein said activity is selected  
10 from the group consisting of lipid partitioning, lipid metabolism, insulin-like activity, free fatty acid oxidation, and weight reduction.
5. A pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of the AGONIST or the ANTAGONIST of Claim 3.
6. A method of preventing or treating an obesity-related disease or disorder comprising  
15 providing or administering to an individual in need of such treatment the composition of Claim 5.

SEQUENCE LISTING

<110> Genset

<120> TREATMENT OF METABOLIC DISORDERS WITH XIDATIN AGONISTS AND ANTAGONISTS

<130> 135.WO1

<150> 60/310,088

<151> 2001-08-03

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1726

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (83)...(1243)

<400> 1

```

cgagaacctt tgcacgcgca caaactacgg ggacgatttc tgattgattt ttggcgcttt 60
cgatccaccc tcctcccttc tc atg gga ctt tgg gga caa agc gtc ccg acc 112
      Met Gly Leu Trp Gly Gln Ser Val Pro Thr
              1              5              10

gcc tcg agc gct cga gca ggg cgc tat cca gga gcc agg aca gcg tcg 160
Ala Ser Ser Ala Arg Ala Gly Arg Tyr Pro Gly Ala Arg Thr Ala Ser
              15              20              25

gga acc aga cca tgg ctc ctg gac ccc aag atc ctt aag ttc gtc gtc 208
Gly Thr Arg Pro Trp Leu Leu Asp Pro Lys Ile Leu Lys Phe Val Val
              30              35              40

ttc atc gtc gcg gtt ctg ctg ccg gtc cgg gtt gac tct gcc acc atc 256
Phe Ile Val Ala Val Leu Leu Pro Val Arg Val Asp Ser Ala Thr Ile
              45              50              55

ccc cgg cag gac gaa gtt ccc cag cag aca gtg gcc cca cag caa cag 304
Pro Arg Gln Asp Glu Val Pro Gln Gln Thr Val Ala Pro Gln Gln Gln
              60              65              70

agg cgc agc ctc aag gag gag gag tgt cca gca gga tct cat aga tca 352
Arg Arg Ser Leu Lys Glu Glu Glu Cys Pro Ala Gly Ser His Arg Ser
              75              80              85              90

gaa tat act gga gcc tgt aac ccg tgc aca gag ggt gtg gat tac acc 400
Glu Tyr Thr Gly Ala Cys Asn Pro Cys Thr Glu Gly Val Asp Tyr Thr
              95              100              105

att gct tcc aac aat ttg cct tct tgc ctg cta tgt aca gtt tgt aaa 448
Ile Ala Ser Asn Asn Leu Pro Ser Cys Leu Leu Cys Thr Val Cys Lys
              110              115              120

tca ggt caa aca aat aaa agt tcc tgt acc acg acc aga gac acc gtg 496
Ser Gly Gln Thr Asn Lys Ser Ser Cys Thr Thr Thr Arg Asp Thr Val
              125              130              135
    
```

tgt cag tgt gaa aaa gga agc ttc cag gat aaa aac tcc cct gag atg	544
Cys Gln Cys Glu Lys Gly Ser Phe Gln Asp Lys Asn Ser Pro Glu Met	
140	145 150
tgc cgg acg tgt aga aca ggg tgt ccc aga ggg atg gtc aag gtc agt	592
Cys Arg Thr Cys Arg Thr Gly Cys Pro Arg Gly Met Val Lys Val Ser	
155	160 165 170
aat tgt acg ccc cgg agt gac atc aag tgc aaa aat gaa tca gct gcc	640
Asn Cys Thr Pro Arg Ser Asp Ile Lys Cys Lys Asn Glu Ser Ala Ala	
	175 180 185
agt tcc act ggg aaa acc cca gca gcg gag gag aca gtg acc acc atc	688
Ser Ser Thr Gly Lys Thr Pro Ala Ala Glu Glu Thr Val Thr Thr Ile	
	190 195 200
ctg ggg atg ctt gcc tct ccc tat cac tac ctt atc atc ata gtg gtt	736
Leu Gly Met Leu Ala Ser Pro Tyr His Tyr Leu Ile Ile Ile Val Val	
	205 210 215
tta gtc atc att tta gct gtg gtt gtg gtt ggc ttt tca tgt cgg aag	784
Leu Val Ile Ile Leu Ala Val Val Val Val Gly Phe Ser Cys Arg Lys	
	220 225 230
aaa ttc att tct tac ctc aaa ggc atc tgc tca ggt ggt gga gga ggt	832
Lys Phe Ile Ser Tyr Leu Lys Gly Ile Cys Ser Gly Gly Gly Gly Gly	
	235 240 245 250
ccc gaa cgt gtg cac aga gtc ctt ttc cgg cgg cgt tca tgt cct tca	880
Pro Glu Arg Val His Arg Val Leu Phe Arg Arg Arg Ser Cys Pro Ser	
	255 260 265
cga gtt cct ggg gcg gag gac aat gcc cgc aac gag acc ctg agt aac	928
Arg Val Pro Gly Ala Glu Asp Asn Ala Arg Asn Glu Thr Leu Ser Asn	
	270 275 280
aga tac ttg cag ccc acc cag gtc tct gag cag gaa atc caa ggt cag	976
Arg Tyr Leu Gln Pro Thr Gln Val Ser Glu Gln Glu Ile Gln Gly Gln	
	285 290 295
gag ctg gca gag cta aca ggt gtg act gta gag tcg cca gag gag cca	1024
Glu Leu Ala Glu Leu Thr Gly Val Thr Val Glu Ser Pro Glu Glu Pro	
	300 305 310
cag cgt ctg ctg gaa cag gca gaa gct gaa ggg tgt cag agg agg agg	1072
Gln Arg Leu Leu Glu Gln Ala Glu Ala Glu Gly Cys Gln Arg Arg Arg	
	315 320 325 330
ctg ctg gtt cca gtg aat gac gct gac tcc gct gac atc agc acc ttg	1120
Leu Leu Val Pro Val Asn Asp Ala Asp Ser Ala Asp Ile Ser Thr Leu	
	335 340 345
ctg gat gcc tcg gca aca ctg gaa gaa gga cat gca aag gaa aca att	1168
Leu Asp Ala Ser Ala Thr Leu Glu Glu Gly His Ala Lys Glu Thr Ile	
	350 355 360
cag gac caa ctg gtg ggc tcc gaa aag ctc ttt tat gaa gaa gat gag	1216
Gln Asp Gln Leu Val Gly Ser Glu Lys Leu Phe Tyr Glu Glu Asp Glu	
	365 370 375
gca ggc tct gct acg tcc tgc ctg tga aagaatctct tcaggaaacc	1263
Ala Gly Ser Ala Thr Ser Cys Leu *	
	380 385

```

agagcttccc tcatttacct tttctcctac aaaggggaag agcctggaag aaacagtcca 1323
gtacttgacc catgccccaa caaactctac tatccaatat ggggcagctt accaatgggc 1383
ctagaacttt gttaacgcac ttggagtaat ttttatgaaa tactgcgtgt gataagcaaa 1443
cgggagaaat ttatatcaga ttcttggctg catagttata cgattgtgta ttaagggctg 1503
ttttaggcca catgcggtgg ctcatgcctg taatcccagc actttgatag gctgaggcag 1563
gtggattgct tgagctcggg agtttgagac cagcctcatc aacacagtga aactccatct 1623
caatttaaaa agaaaaaag tggtttagg atgtcattct ttgcagttct tcatcatgag 1683
acaagtcttt ttttctgctt cttatattgc aagctccatc tct 1726
    
```

```

<210> 2
<211> 386
<212> PRT
<213> Homo sapiens
    
```

```

<220>
<221> VARIANT
<222> 35
<223> Polymorphic amino acid Pro or Ser
    
```

```

<221> VARIANT
<222> 310
<223> Polymorphic amino acid Ser or Leu
    
```

```

<400> 2
Met Gly Leu Trp Gly Gln Ser Val Pro Thr Ala Ser Ser Ala Arg Ala
 1          5          10          15
Gly Arg Tyr Pro Gly Ala Arg Thr Ala Ser Gly Thr Arg Pro Trp Leu
 20          25          30
Leu Asp Pro Lys Ile Leu Lys Phe Val Val Phe Ile Val Ala Val Leu
 35          40          45
Leu Pro Val Arg Val Asp Ser Ala Thr Ile Pro Arg Gln Asp Glu Val
 50          55          60
Pro Gln Gln Thr Val Ala Pro Gln Gln Gln Arg Arg Ser Leu Lys Glu
 65          70          75          80
Glu Glu Cys Pro Ala Gly Ser His Arg Ser Glu Tyr Thr Gly Ala Cys
 85          90          95
Asn Pro Cys Thr Glu Gly Val Asp Tyr Thr Ile Ala Ser Asn Asn Leu
100          105          110
Pro Ser Cys Leu Leu Cys Thr Val Cys Lys Ser Gly Gln Thr Asn Lys
115          120          125
Ser Ser Cys Thr Thr Thr Arg Asp Thr Val Cys Gln Cys Glu Lys Gly
130          135          140
Ser Phe Gln Asp Lys Asn Ser Pro Glu Met Cys Arg Thr Cys Arg Thr
145          150          155          160
Gly Cys Pro Arg Gly Met Val Lys Val Ser Asn Cys Thr Pro Arg Ser
165          170          175
Asp Ile Lys Cys Lys Asn Glu Ser Ala Ala Ser Ser Thr Gly Lys Thr
180          185          190
Pro Ala Ala Glu Glu Thr Val Thr Thr Ile Leu Gly Met Leu Ala Ser
195          200          205
Pro Tyr His Tyr Leu Ile Ile Ile Val Val Leu Val Ile Ile Leu Ala
210          215          220
Val Val Val Val Gly Phe Ser Cys Arg Lys Lys Phe Ile Ser Tyr Leu
225          230          235          240
Lys Gly Ile Cys Ser Gly Gly Gly Gly Gly Pro Glu Arg Val His Arg
245          250          255
Val Leu Phe Arg Arg Arg Ser Cys Pro Ser Arg Val Pro Gly Ala Glu
260          265          270
Asp Asn Ala Arg Asn Glu Thr Leu Ser Asn Arg Tyr Leu Gln Pro Thr
275          280          285
Gln Val Ser Glu Gln Glu Ile Gln Gly Gln Glu Leu Ala Glu Leu Thr
290          295          300
    
```



**INTERNATIONAL SEARCH REPORT**

International Application No PCT/IB 02/03404
---

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 A61K38/17 A61K45/00 G01N33/50 A61P3/00

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
 IPC 7 A61K A61P G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)  
 EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, SEQUENCE SEARCH

<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 51645 A (GENSET SA ;YEN POTIN FRANCES (US); BIHAIN BERNARD (US); FRUEBIS JO) 19 July 2001 (2001-07-19) claims page 58, line 11 -page 59, line 15 ----	1-6
X	EP 1 033 134 A (OTSUKA PHARMA CO LTD) 6 September 2000 (2000-09-06) page 2, line 44 - line 47 ----	1-6
X	WO 00 68380 A (INCYTE GENOMICS INC ;AZIMZAI YALDA (US); YUE HENRY (US); BANDMAN O) 16 November 2000 (2000-11-16) SEQ ID NO 8 page 35, line 17 -page 37, line 5 ----- -/--	1-6

Further documents are listed in the continuation of box C.       Patent family members are listed in annex.

° Special categories of cited documents :

*A* document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E* earlier document but published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
*O* document referring to an oral disclosure, use, exhibition or other means	*Z* document member of the same patent family
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  27 November 2002	Date of mailing of the international search report  04/12/2002
---	--

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer  Böhmerova, E
--	--

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 02/03404

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 59618 A (SMITHKLINE BEECHAM CORP) 25 November 1999 (1999-11-25) page 1, line 23 - line 32 page 15, line 17 - line 20 ---	1-6
X	WO 00 73448 A (ZYMOGENETICS INC) 7 December 2000 (2000-12-07) page 86, line 21 -page 88, line 23 page 72, line 28 -page 73, line 33 ---	1-6
X	WO 00 77037 A (ASHKENAZI AVI J;GENENTECH INC) 21 December 2000 (2000-12-21) page 66, line 35 -page 67, line 3 page 87, line 21 -page 90, line 28; figure 18; examples 47,72 -----	1-6

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 02/03404

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0151645	A	19-07-2001	AU 5789901	A 24-07-2001
			BR 0107626	A 15-10-2002
			EP 1248849	A1 16-10-2002
			WO 0151645	A1 19-07-2001
			US 2002058617	A1 16-05-2002
			US 2002037849	A1 28-03-2002
			US 2002091080	A1 11-07-2002
EP 1033134	A	06-09-2000	EP 1033134	A1 06-09-2000
			US 6461821	B1 08-10-2002
			WO 9921577	A1 06-05-1999
WO 0068380	A	16-11-2000	AU 5129800	A 21-11-2000
			EP 1177296	A2 06-02-2002
			WO 0068380	A2 16-11-2000
WO 9959618	A	25-11-1999	EP 1085897	A1 28-03-2001
			EP 1083923	A1 21-03-2001
			JP 2002515452	T 28-05-2002
			WO 9959618	A1 25-11-1999
			WO 9959619	A1 25-11-1999
			US 2001009905	A1 26-07-2001
WO 0073448	A	07-12-2000	AU 5159500	A 18-12-2000
			EP 1185645	A1 13-03-2002
			WO 0073448	A1 07-12-2000
WO 0077037	A	21-12-2000	AU 5152700	A 02-01-2001
			EP 1208195	A2 29-05-2002
			WO 0077037	A2 21-12-2000
			AU 1748200	A 19-06-2000
			AU 1749800	A 04-10-2000
			AU 1749900	A 12-07-2000
			AU 2390700	A 05-02-2001
			AU 2495200	A 28-09-2000
			AU 2600800	A 28-09-2000
			AU 2879400	A 31-07-2001
			AU 2883600	A 28-09-2000
			AU 2883700	A 09-01-2001
			AU 2883900	A 30-01-2001
			AU 3107000	A 19-06-2000
			AU 3246100	A 28-12-2000
			AU 3381600	A 28-09-2000
			AU 3514400	A 28-09-2000
			AU 3774300	A 18-12-2000
			AU 5441200	A 18-12-2000
			EP 1173563	A1 23-01-2002
			EP 1159419	A1 05-12-2001
			EP 1220905	A2 10-07-2002
			EP 1210418	A1 05-06-2002
			EP 1212417	A2 12-06-2002
			EP 1135485	A2 26-09-2001
			EP 1141284	A2 10-10-2001
			EP 1135491	A2 26-09-2001
			JP 2002531092	T 24-09-2002
			WO 0053753	A2 14-09-2000
			WO 0053755	A2 14-09-2000
			WO 0153486	A1 26-07-2001

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No PCT/IB 02/03404
---

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0077037	A	WO 0053756 A2	14-09-2000
		WO 0078961 A1	28-12-2000
		WO 0104311 A1	18-01-2001
		WO 0075327 A1	14-12-2000
		WO 0053757 A2	14-09-2000
		WO 0053758 A2	14-09-2000
		WO 0073454 A1	07-12-2000
		WO 0073445 A2	07-12-2000
		WO 0032221 A2	08-06-2000
		WO 0032778 A2	08-06-2000
		WO 0055319 A1	21-09-2000
		WO 0037638 A2	29-06-2000
		WO 0105836 A1	25-01-2001
		US 2002010137 A1	24-01-2002
		AU 4851900 A	28-12-2000
		EP 1185642 A2	13-03-2002

---