

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 April 2011 (07.04.2011)

(10) International Publication Number  
**WO 2011/041154 A1**

(51) International Patent Classification:  
*C07D 513/14* (2006.01)   *A61P 3/10* (2006.01)  
*A61K 31/506* (2006.01)

Hayward, California 94545 (US). **ANDRES, Patricia** [FR/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US).

(21) International Application Number:  
PCT/US2010/049486

(74) Agents: **HUANG, Wenrong** et al.; Swiss Tanner, P.C., P.O. Box 1749, Los Altos, California 94022 (US).

(22) International Filing Date:  
20 September 2010 (20.09.2010)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English  
(26) Publication Language: English  
(30) Priority Data:  
61/247,936      1 October 2009 (01.10.2009)      US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **METABOLEX, INC.** [US/US]; 3876 Bay Center Place, Hayward, California 94545 (US).

Published:

— with international search report (Art. 21(3))



**WO 2011/041154 A1**

(54) Title: SUBSTITUTED TETRAZOL-1-YL-PHENOXYMETHYL-THIAZOL-2-YL-PIPERIDINYL-PYRIMIDINE SALTS

(57) Abstract: Crystalline salts of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl}-piperidin-1-yl}-pyrimidine, compositions thereof, methods for their preparation, and methods for their use are disclosed.

SUBSTITUTED TETRAZOL-1-YL-PHENOXYMETHYL-THIAZOL-2-YL-  
PIPERIDINYL-PYRIMIDINE SALTS

FIELD OF THE INVENTION

5 [0001] Provided are crystalline salts of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-  
phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine, compositions thereof, methods  
for their preparation, and methods for their use.

BACKGROUND OF THE INVENTION

10 [0002] WO 2008/083238 discloses 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxyethyl)-  
thiazol-2-yl]-piperidin-1-yl}-pyrimidine and its uses for the treatment of diabetes and  
metabolic disorders. PCT/US2009/038847 discloses uses of oxymethylene compounds  
including 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-  
pyrimidine in a combination therapy with a dipeptidyl peptidase IV (DPP IV) inhibitor.  
A need exists for improved therapies for the treatment of diseases relating to diabetes and  
15 metabolic disorders.

BRIEF SUMMARY OF THE INVENTION

20 [0003] In one embodiment, the present invention provides crystalline salts of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine selected  
from the group consisting of a besylate, camsylate, esylate, HBr, HCl, mesylate, sulfate,  
and tosylate salt.

[0004] In another embodiment, provided are polymorphs (Form 1 and Form 2) of 5-  
ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine  
hydrochloride.

25 [0005] In other embodiments, provided are methods for the preparation of a crystalline  
salt or polymorph described herein.

[0006] In other embodiments, provided are compositions comprising a crystalline salt or  
polymorph described herein and a pharmaceutically acceptable carrier.

[0007] In other embodiments, provided are methods for use of a crystalline salt or  
polymorph described herein to treat a disease selected from the group consisting of Type I

diabetes, Type II diabetes and metabolic syndrome, and their uses in the preparation of such medicaments.

**[0008]** In other embodiments, provided are methods for use of a crystalline salt or polymorph described herein for one or more of stimulating insulin production, stimulating

5 glucose-dependent insulin secretion, lowering blood glucose, or lowering blood triglyceride levels, and their uses in the preparation of such medicaments.

**[0009]** In other embodiments, provided are methods for use of a crystalline salt or polymorph described herein in a combination therapy with a therapeutically effective amount of a DPP IV inhibitor.

10 **[0010]** These and other aspects of the invention are further described in the text that follows.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0011]** Figure 1 shows the X-ray powder pattern of HCl salt Form I.

**[0012]** Figure 2 shows the DSC thermogram of HCl salt Form I.

15 **[0013]** Figure 3 shows the TGA thermogram of HCl salt Form I.

**[0014]** Figure 4 shows the Raman spectrum of HCl salt Form I.

**[0015]** Figure 5 shows the X-ray powder pattern of HCl salt Form II.

**[0016]** Figure 6 shows the DSC thermogram of HCl salt Form II.

**[0017]** Figure 7 shows the TGA thermogram of HCl salt Form II.

20 **[0018]** Figure 8 shows an overlay of the DSC and TGA thermograms of the besylate salt.

**[0019]** Figure 9 shows an overlay of the DSC and TGA thermograms of the camsylate salt.

25 **[0020]** Figure 10 shows an overlay of the DSC and TGA thermograms of the esylate salt.

**[0021]** Figure 11 shows an overlay of the DSC and TGA thermograms of the HBr salt.

**[0022]** Figure 12 shows an overlay of the DSC and TGA thermograms of the mesylate salt.

[0023] Figure 13 shows an overlay of the DSC and TGA thermograms of the sulfate salt.

[0024] Figure 14 shows an overlay of the DSC and TGA thermograms of the tosylate salt.

5 [0025] Figure 15 shows the X-ray powder pattern of the besylate salt.

[0026] Figure 16 shows the X-ray powder pattern of the camsylate salt.

[0027] Figure 17 shows the X-ray powder pattern of the esylate salt.

[0028] Figure 18 shows the X-ray powder pattern of the HBr salt.

[0029] Figure 19 shows the X-ray powder pattern of the mesylate salt.

10 [0030] Figure 20 shows the X-ray powder pattern of the sulfate salt.

[0031] Figure 21 shows the X-ray powder pattern of the tosylate salt.

[0032] Figure 22 shows the X-ray powder pattern of the HCl salt (pattern O).

[0033] Figure 23 shows the mean plasma concentrations of micronized MBX-2982 free-base, HCl Forms I and II, and the mesylate and tosylate salts versus time in male SD rats following single oral dose of 200 mg/kg.

#### DETAILED DESCRIPTION OF THE INVENTION

[0034] In this specification and in the claims that follow, reference will be made to a number of terms that shall be defined to have the following meanings unless specified otherwise.

20 Abbreviations

[0035] XRPD (x-ray powder diffraction); DSC (differential scanning calorimetry); TGA (thermographic analysis); besylate (bezene sulfonate salt); camsylate (camphorsulfonate salt); esylate (ethanesulfonate salt); mesylate (methanesulfonate salt); tosylate (p-toluene sulfonate salt); acetone ((CH<sub>3</sub>)<sub>2</sub>CO ); RH (relative humidity).

25 [0036] As used in the specification and claims, the singular form “a”, “an” and “the” include plural references unless the context clearly dictates otherwise.

[0037] As used herein, the term “comprising” is intended to mean that the compositions and methods include the recited elements, but not excluding others.

**[0038]** As used herein, the term "about" when used in association with a measurement, or used to modify a value, a unit, a constant, or a range of values, refers to variations of +/- 3%.

**[0039]** As used herein, the term "crystalline salt" encompasses anhydrous and 5 unsolvated crystals, as well as crystals that may be associated with varying degrees of hydrates or solvates.

**[0040]** As used herein, the term "substantially" refers to degree of variations of +/- by about 1%, about 5%, about 10%, about 15% or about 20%.

**[0041]** As used herein, the term "substantially pure" with respect to a particular 10 polymorphic form of a compound, means that the polymorph form contains about less than 30%, or about less than 20%, or about less than 15%, or about less than 10%, or about less than 5%, or about less than 1% by weight of impurities, such impurities may include other polymorphic forms of the same compound or trace solvents.

**[0042]** As used herein, the term "pharmaceutically acceptable carrier" includes any and 15 all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (See, for example, Remington's 20 Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289- 1329, incorporated herein by reference.). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated.

**[0043]** As used herein, the term "subject" refers to a mammal and includes, without 25 limitation, humans, domestic animals (e.g., dogs or cats), farm animals (cows, horses, or pigs), and laboratory animals (mice, rats, hamsters, guinea pigs, pigs, rabbits, dogs, or monkeys).

**[0044]** The term "therapeutically effective amount" refers to that amount of an active 30 ingredient that is sufficient to effect treatment, as defined below, when administered to a mammal in need of such treatment. The therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the

subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by a prescribing physician.

**[0045]** The term "treatment" or "treating" means any treatment of a disease in a mammal, including: (i) preventing the disease, that is, causing the clinical symptoms of the disease not to develop; (ii) inhibiting the disease, that is, arresting the development of clinical symptoms; and/or (iii) relieving the disease, that is, causing the regression of clinical symptoms.

**[0046]** The term "diabetes mellitus" or "diabetes" means a disease or condition that is generally characterized by metabolic defects in production and utilization of glucose that result in the failure to maintain appropriate blood sugar levels in the body. The result of these defects is elevated blood glucose, referred to as "hyperglycemia." Two major forms of diabetes are Type I diabetes and Type II diabetes. As described above, Type I diabetes is generally the result of an absolute deficiency of insulin, the hormone that regulates glucose utilization. Type II diabetes often occurs in the face of normal, or even elevated levels of insulin and can result from the inability of tissues to respond appropriately to insulin. Most Type II diabetic subjects are insulin resistant and have a relative deficiency of insulin, in that insulin secretion cannot compensate for the resistance of peripheral tissues to respond to insulin. In addition, many Type II diabetics are obese. Other types of disorders of glucose homeostasis include impaired glucose tolerance, which is a metabolic stage intermediate between normal glucose homeostasis and diabetes, and gestational diabetes mellitus, which is glucose intolerance in pregnancy in women with no previous history of Type I or Type II diabetes.

**[0047]** The term "metabolic syndrome" refers to a cluster of metabolic abnormalities including abdominal obesity, insulin resistance, glucose intolerance, diabetes, hypertension, hyperlipidemia, and dyslipidemia. These abnormalities are known to be associated with an increased risk of vascular events.

**[0048]** The term "abdominal obesity" is defined by a cutoff point of waist circumference  $\geq 102$  cm in men and  $\geq 80$  cm in women, as recommended by the third report of the national cholesterol education program expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (NCEP/ATP Panel III).

**[0049]** The term “insulin resistance” can be defined generally as a disorder of glucose metabolism. More specifically, insulin resistance can be defined as the diminished ability of insulin to exert its biological action across a broad range of concentrations producing less than the expected biologic effect (*see, e.g.*, Reaven GM, *J. Basic & Clin. Phys. & Pharm.* (1998) 9:387-406 and Flie J, *Ann Rev. Med.* (1983) 34:145-60). Insulin resistant persons have a diminished ability to properly metabolize glucose and respond poorly, if at all, to insulin therapy.

**[0050]** Manifestations of insulin resistance include insufficient insulin activation of glucose uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue and of glucose production and secretion in liver. Insulin resistance can cause or contribute to polycystic ovarian syndrome, impaired glucose tolerance, gestational diabetes, metabolic syndrome, hypertension, obesity, atherosclerosis and a variety of other disorders. Eventually, the insulin resistant individuals can progress to a point where a diabetic state is reached.

**[0051]** The term “atherosclerosis” encompasses vascular diseases and conditions that are recognized and understood by physicians practicing in the relevant fields of medicine. Atherosclerotic cardiovascular disease, coronary heart disease (also known as coronary artery disease or ischemic heart disease), cerebrovascular disease and peripheral vessel disease are all clinical manifestations of atherosclerosis and are therefore encompassed by the terms “atherosclerosis” and “atherosclerotic disease”.

**[0052]** The term “symptom” of diabetes, includes, but is not limited to, polyuria, polydipsia, and polyphagia, as used herein, incorporating their common usage. For example, “polyuria” means the passage of a large volume of urine during a given period; “polydipsia” means chronic, excessive thirst; and “polyphagia” means excessive eating.

**[0053]** Other symptoms of diabetes include, *e.g.*, increased susceptibility to certain infections (especially fungal and staphylococcal infections), nausea, and ketoacidosis (enhanced production of ketone bodies in the blood).

**[0054]** The term “complication” of diabetes includes, but is not limited to, microvascular complications and macrovascular complications. Microvascular complications are those complications that generally result in small blood vessel damage. These complications include, *e.g.*, retinopathy (the impairment or loss of vision due to blood vessel damage in the eyes); neuropathy (nerve damage and foot problems due to

blood vessel damage to the nervous system); and nephropathy (kidney disease due to blood vessel damage in the kidneys). Macrovascular complications are those complications that generally result from large blood vessel damage. These complications include, *e.g.*, cardiovascular disease and peripheral vascular disease. Cardiovascular 5 disease refers to diseases of blood vessels of the heart. See, *e.g.*, Kaplan RM, et al., “Cardiovascular diseases” in *Health and Human Behavior*, pp. 206-242 (McGraw-Hill, New York 1993). Cardiovascular disease is generally one of several forms, including, *e.g.*, hypertension (also referred to as high blood pressure), coronary heart disease, stroke, and rheumatic heart disease. Peripheral vascular disease refers to diseases of any of the 10 blood vessels outside of the heart. It is often a narrowing of the blood vessels that carry blood to leg and arm muscles.

**[0054]** The term “dyslipidemia” refers to abnormal levels of lipoproteins in blood plasma including both depressed and/or elevated levels of lipoproteins (*e.g.*, elevated levels of LDL and/or VLDL, and depressed levels of HDL).

15 **[0055]** The term “hyperlipidemia” includes, but is not limited to, the following:

- (1) *Familial Hyperchylomicronemia*, a rare genetic disorder that causes a deficiency in an enzyme, LP lipase, that breaks down fat molecules. The LP lipase deficiency can cause the accumulation of large quantities of fat or lipoproteins in the blood;
- (2) *Familial Hypercholesterolemia*, a relatively common genetic disorder caused 20 where the underlying defect is a series of mutations in the LDL receptor gene that result in malfunctioning LDL receptors and/or absence of the LDL receptors. This brings about ineffective clearance of LDL by the LDL receptors resulting in elevated LDL and total cholesterol levels in the plasma;
- (3) *Familial Combined Hyperlipidemia*, also known as multiple lipoprotein-type 25 hyperlipidemia is an inherited disorder where subjects and their affected first-degree relatives can at various times manifest high cholesterol and high triglycerides. Levels of HDL cholesterol are often moderately decreased;
- (4) *Familial Defective Apolipoprotein B-100* is a relatively common autosomal dominant genetic abnormality. The defect is caused by a single nucleotide mutation that 30 produces a substitution of glutamine for arginine, which can cause reduced affinity of

LDL particles for the LDL receptor. Consequently, this can cause high plasma LDL and total cholesterol levels;

(5) *Familial Dysbetalipoproteinemia*, also referred to as Type III


Hyperlipoproteinemia, is an uncommon inherited disorder resulting in moderate to severe elevations of serum TG and cholesterol levels with abnormal apolipoprotein E function. HDL levels are usually normal; and

(6) *Familial Hypertriglyceridemia*, is a common inherited disorder in which the concentration of plasma VLDL is elevated. This can cause mild to moderately elevated TG levels (and usually not cholesterol levels) and can often be associated with low plasma HDL levels.

**[0056]** Risk factors for hyperlipidemia include, but are not limited to, the following: (1) disease risk factors, such as a history of Type I diabetes, Type II diabetes, Cushing's syndrome, hypothyroidism and certain types of renal failure; (2) drug risk factors, which include, birth control pills; hormones, such as estrogen, and corticosteroids; certain diuretics; and various  $\beta$  blockers; (3) dietary risk factors include dietary fat intake per total calories greater than 40%; saturated fat intake per total calories greater than 10%; cholesterol intake greater than 300 mg per day; habitual and excessive alcohol use; and obesity.

**[0057]** The terms "obese" and "obesity" refers to, according to the World Health Organization, a Body Mass Index ("BMI") greater than  $27.8 \text{ kg/m}^2$  for men and  $27.3 \text{ kg/m}^2$  for women (BMI equals weight (kg)/height ( $\text{m}^2$ )). Obesity is linked to a variety of medical conditions including diabetes and hyperlipidemia. Obesity is also a known risk factor for the development of Type II diabetes (see, e.g., Barrett-Conner E, *Epidemiol. Rev.* (1989) 11:172-181; and Knowler, et al., *Am. J. Clin. Nutr.* (1991) 53:1543-1551).

**25 [0058]** The free base 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl}-piperidin-1-yl}-pyrimidine, MBX-2982, has the structure:



**[0059]** The besylate, camsylate, esylate, HBr, HCl, mesylate, sulfate, and tosylate salts of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine provided herein are found to be crystalline solids as further evidenced by their XRPD patterns. In contrast, salts formed from acetic, adipic, L-ascorbic, L-aspartic, 5 citric, formic, gentisic (2,5-dihydroxybenzoic), L-glutamic, glutaric, lactic, maleic, L- malic, phosphoric, and L-tartaric were found to be oils or gels that were difficult to handle and isolate, and would not be suitable for use in bulk preparations. The crystalline besylate, camsylate, esylate, HBr, HCl, mesylate, sulfate, and tosylates salts are therefore superior for use in preparing pharmaceutical salts of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl- 10 phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine.

**[0060]** The salts are characterized by a number of solid state techniques such as DSC and TGA. It is understood that melting point temperatures and thermograms can vary depending on instrumentation and the procedures employed, including the heating rate used. Accordingly, the temperature data and graphs disclosed herein are understood to 15 accommodate such variations.

**[0061]** The salts are also characterized by XRPD (x-ray powder diffraction). Relative intensities and peak assignments can vary depending on a number of factors, including sample preparation, mounting, and the instrument and analytical procedure and settings used to obtain the spectrum. The peak assignments described herein are intended to 20 encompass variations of plus or minus 0.5 degrees 2 theta.

**[0062]** In one embodiment, provided is a crystalline besylate salt having a DSC endotherm at about 153°C. In one aspect, the besylate salt has a DSC or TGA thermogram substantially as shown in Figure 8. In other aspects, the besylate salt has the XRPD pattern substantially as shown in Figure 15.

**[0063]** In one embodiment, provided is a crystalline camsylate salt having a DSC endotherm at about 184°C. In one aspect, the camsylate salt has a DSC or TGA thermogram substantially as shown in Figure 9. In other aspects, the camsylate salt has the XRPD pattern substantially as shown in Figure 16.

**[0064]** In one embodiment, provided is a crystalline esylate salt having a DSC endotherm at about 99°C. In one aspect, the esylate salt has a DSC or TGA thermogram

substantially as shown in Figure 10. In other aspects, the esylate salt has the XRPD pattern substantially as shown in Figure 17.

**[0065]** In one embodiment, provided is a crystalline HBr salt having a DSC endotherm at about 142°C. In one aspect, the HBr salt has a DSC or TGA thermogram substantially as shown in Figure 11. In other aspects, the HBr salt has the XRPD pattern substantially as shown in Figure 18.

**[0066]** In one embodiment, provided is a crystalline mesylate salt having a DSC endotherm at about 86°C. In one aspect, the mesylate salt has a DSC or TGA thermogram substantially as shown in Figure 12. In other aspects, the mesylate salt has the XRPD pattern substantially as shown in Figure 19.

**[0067]** In one embodiment, provided is a crystalline sulfate salt having a DSC endotherm at about 97°C. In one aspect, the sulfate salt has a DSC or TGA thermogram substantially as shown in Figure 13. In other aspects, the sulfate salt has the XRPD pattern substantially as shown in Figure 20.

**[0068]** In one embodiment, provided is a crystalline tosylate salt having a DSC endotherm at about 94°C. In one aspect, the tosylate salt has a DSC or TGA thermogram substantially as shown in Figure 14. In other aspects, the tosylate salt has the XRPD pattern substantially as shown in Figure 21.

**[0069]** In one embodiment, provided is a crystalline HCl salt. The crystalline HCl salts include the crystal “pattern O” described below and those described in the Examples (Table 6). The HCl salt may also contain a polymorph such as Form I or Form II described below or mixtures thereof. The crystalline HCl may also contain trace amounts of solvents such as acetone, ethanol, methanol, and ethyl acetate. In some aspects, the Form I polymorph provided herein is substantially pure and may contain residual acetone. In some aspects, the Form II polymorph provided herein is substantially pure and may contain residual methanol and/or ethyl acetate.

**[0070]** In one embodiment, provided is a crystalline HCl salt (pattern O) containing trace amounts of ethanol. In one aspect, the ratio of EtOH to HCl salt is approximately 1:6. In one aspect, the HCl salt has the XRPD pattern substantially as shown in Figure 22.

**[0071]** In one embodiment, provided is polymorph (Form I) HCl salt having a DSC endotherm onset at about 191°C. In one aspect, the Form I polymorph has a DSC

thermogram substantially as shown in Figure 2. In other aspects, the Form I polymorph has a TGA thermogram substantially as shown in Figure 3. In other aspects, the Form I polymorph has a Raman spectrum substantially as shown in Figure 4. In other aspects, the Form I polymorph comprises XRPD peaks at degrees 2-theta diffraction angles of 5 about 8.8, 10.8, 16.1, 17.4, 20.4, 20.9, 21.5, 21.7, 26.6, and 28.1. In still other aspects, the Form I polymorph has the XRPD pattern substantially as shown in Figure 1. Table 1 lists the observed Form 1 XRPD peaks and Table 2 lists the representative peaks, with relative intensities given in both tables. Table 3 lists the observed Raman peaks for Form I.

**[0072]** In one embodiment, provided is polymorph (Form II) HCl salt having a DSC 10 endotherm onset at about 150°C. In one aspect, the Form II polymorph has a DSC thermogram substantially as shown in Figure 6. In other aspects, the Form II polymorph has a TGA thermogram substantially as shown in Figure 7. In other aspects, the Form II polymorph comprises XRPD peaks at degrees 2-theta diffraction angles of about 7.8, 15 10.1, 12.5, 18.4, 19.0, 20.8, 23.0, and 23.5. In still other aspects, the Form II polymorph has the XRPD pattern substantially as shown in Figure 5. Table 4 lists the observed Form II XRPD peaks and Table 5 lists the representative peaks, with relative intensities given in both tables.

**[0073]** In the Tables below, the location of the peaks were automatically determined 20 using PatternMatch™ 3.0.1 software and rounded to one or two significant figures after the decimal point based upon the above criteria. Peak position variabilities are given to within  $\pm 0.10^\circ$  2 theta based upon recommendations outlined in the United States Pharmacopeia (USP 32, NF 27, Vol. 1, pg. 392, 2009) discussion of variability in x-ray 25 powder diffraction. The accuracy and precision associated with any particular measurement reported herein has not been determined. Moreover, third party measurements on independently prepared samples on different instruments may lead to variability which is greater than  $\pm 0.10^\circ$  2 theta (for example  $\pm 0.50^\circ$  2 theta or more).

**Table 1.** Observed XRPD peaks for HCl polymorph Form I.

	$^\circ 2\theta$	$d$ space (Å)	Intensity (%)
30	8.76 $\pm$ 0.10	10.092 $\pm$ 0.116	24
	10.78 $\pm$ 0.10	8.204 $\pm$ 0.077	40
	10.98 $\pm$ 0.10	8.054 $\pm$ 0.074	5
	11.34 $\pm$ 0.10	7.806 $\pm$ 0.069	3
	12.66 $\pm$ 0.10	6.994 $\pm$ 0.055	8
35	13.14 $\pm$ 0.10	6.738 $\pm$ 0.051	7

5	14.06 ± 0.10	6.299 ± 0.045	2
	14.38 ± 0.10	6.161 ± 0.043	3
	15.48 ± 0.10	5.724 ± 0.037	2
	15.88 ± 0.10	5.581 ± 0.035	8
10	16.10 ± 0.10	5.506 ± 0.034	31
	16.65 ± 0.10	5.325 ± 0.032	7
	17.08 ± 0.10	5.190 ± 0.030	5
	17.45 ± 0.10	5.082 ± 0.029	26
	18.15 ± 0.10	4.887 ± 0.027	2
15	18.30 ± 0.10	4.847 ± 0.026	3
	18.45 ± 0.10	4.808 ± 0.026	6
	18.81 ± 0.10	4.719 ± 0.025	1
	19.39 ± 0.10	4.578 ± 0.024	2
20	20.39 ± 0.10	4.355 ± 0.021	23
	20.88 ± 0.10	4.255 ± 0.020	100
	21.46 ± 0.10	4.140 ± 0.019	34
	21.76 ± 0.10	4.084 ± 0.019	20
	22.06 ± 0.10	4.029 ± 0.018	5
	22.67 ± 0.10	3.923 ± 0.017	6
25	23.49 ± 0.10	3.788 ± 0.016	2
	23.69 ± 0.10	3.757 ± 0.016	2
	24.37 ± 0.10	3.652 ± 0.015	1
	24.74 ± 0.10	3.599 ± 0.014	3
	24.99 ± 0.10	3.563 ± 0.014	2
30	25.16 ± 0.10	3.540 ± 0.014	4
	25.44 ± 0.10	3.501 ± 0.014	5
	25.67 ± 0.10	3.470 ± 0.013	4
	25.86 ± 0.10	3.446 ± 0.013	2
35	26.61 ± 0.10	3.350 ± 0.012	53
	28.06 ± 0.10	3.180 ± 0.011	22
	28.35 ± 0.10	3.148 ± 0.011	3
	28.80 ± 0.10	3.100 ± 0.011	7
	29.02 ± 0.10	3.077 ± 0.010	6
	29.37 ± 0.10	3.041 ± 0.010	6

**Table 2.** Representative XRPD peaks for HCl polymorph Form I.

	$^{\circ}2\theta$	<i>d</i> space (Å)	Intensity (%)
40	8.76 ± 0.10	10.092 ± 0.116	24
	10.78 ± 0.10	8.204 ± 0.077	40
	16.10 ± 0.10	5.506 ± 0.034	31
	17.45 ± 0.10	5.082 ± 0.029	26
	20.39 ± 0.10	4.355 ± 0.021	23
45	20.88 ± 0.10	4.255 ± 0.020	100
	21.46 ± 0.10	4.140 ± 0.019	34
	21.76 ± 0.10	4.084 ± 0.019	20
	26.61 ± 0.10	3.350 ± 0.012	53
	28.06 ± 0.10	3.180 ± 0.011	22

**Table 3.** Observed Raman peaks for HCl polymorph Form I (cm<sup>-1</sup>)

154	193	214	243
324	346	384	420
460	467	495	539
573	596	628	648
680	687	718	751
766	787	800	845
856	946	967	992
1009	1051	1068	1094
1110	1172	1203	1225
1253	1282	1300	1311
1334	1375	1394	1404
1432	1458	1474	1480
1517	1533	1591	1609
1637	2734	2869	2903
2928	2937	2967	2999
3015	3030	3063	3075

5 **Table 4.** Observed XRPD peaks for HCl polymorph Form II.

	<sup>°</sup> 2θ	d space (Å)	Intensity (%)
10	7.63 ± 0.10	11.593 ± 0.154	23
	7.79 ± 0.10	11.345 ± 0.147	43
	8.14 ± 0.10	10.857 ± 0.135	6
	9.18 ± 0.10	9.634 ± 0.106	5
	10.07 ± 0.10	8.788 ± 0.088	41
	10.35 ± 0.10	8.547 ± 0.083	15
	12.54 ± 0.10	7.059 ± 0.057	50
15	12.87 ± 0.10	6.877 ± 0.054	5
	13.24 ± 0.10	6.687 ± 0.051	18
	14.78 ± 0.10	5.994 ± 0.041	7
	14.98 ± 0.10	5.915 ± 0.040	7
	15.20 ± 0.10	5.831 ± 0.038	18
	15.61 ± 0.10	5.675 ± 0.036	3
	15.91 ± 0.10	5.569 ± 0.035	6
20	16.12 ± 0.10	5.500 ± 0.034	17
	17.69 ± 0.10	5.015 ± 0.028	28
	17.97 ± 0.10	4.936 ± 0.027	28
	18.39 ± 0.10	4.825 ± 0.026	100
	18.82 ± 0.10	4.715 ± 0.025	21
	19.01 ± 0.10	4.669 ± 0.024	85
	19.27 ± 0.10	4.605 ± 0.024	14
25	19.67 ± 0.10	4.512 ± 0.023	11
	20.14 ± 0.10	4.409 ± 0.022	9
	20.81 ± 0.10	4.268 ± 0.020	58
	21.41 ± 0.10	4.150 ± 0.019	18
	21.83 ± 0.10	4.071 ± 0.019	35

22.20 ± 0.10	4.005 ± 0.018	6
22.65 ± 0.10	3.926 ± 0.017	25
23.03 ± 0.10	3.861 ± 0.017	83
23.27 ± 0.10	3.823 ± 0.016	35
5	23.54 ± 0.10	3.780 ± 0.016
	23.84 ± 0.10	3.733 ± 0.016
	24.45 ± 0.10	3.640 ± 0.015
	24.67 ± 0.10	3.609 ± 0.014
	24.94 ± 0.10	3.570 ± 0.014
	25.46 ± 0.10	3.499 ± 0.014
10	25.87 ± 0.10	3.443 ± 0.013
	26.08 ± 0.10	3.417 ± 0.013
	26.53 ± 0.10	3.360 ± 0.012
	26.89 ± 0.10	3.315 ± 0.012
	27.23 ± 0.10	3.275 ± 0.012
	27.66 ± 0.10	3.225 ± 0.011
15	28.40 ± 0.10	3.143 ± 0.011
	29.17 ± 0.10	3.062 ± 0.010
	29.87 ± 0.10	2.991 ± 0.010

20 **Table 5.** Representative XRPD peaks for HCl polymorph Form II.

°2θ	d space (Å)	Intensity (%)
7.78 ± 0.10	11.369 ± 0.148	42
25	10.07 ± 0.10	41
	12.54 ± 0.10	50
	18.37 ± 0.10	100
	19.01 ± 0.10	86
	20.81 ± 0.10	59
	23.03 ± 0.10	84
	23.54 ± 0.10	48

[0074] The crystalline salts disclosed herein may be prepared by precipitation from organic or mixed organic solvents and may also be prepared from organic/aqueous solvents. Suitable organic solvents include acetone, acetonitrile, dichloromethane, diethyl ether, ethyl acetate, ethanol, heptane, hexane, hexafluoroisopropanol, isopropyl alcohol, isopropyl ether, methyl ethyl ketone, methanol, methyl-tert-butyl ether, 2,2,2-trifluoroethanol, and tetrahydrofuran.

[0075] The HCl salt (Form I) can generally be prepared by addition of HCl to the free base at elevated temperatures in solvent such as acetone, acetonitrile, ethanol/ethyl acetate, methanol/ethyl acetate and THF, optionally followed by further crystallizations in acetone. The HCl salt (Form II) can generally be prepared by crystallizations in methanol. The methanol solution may optionally contain other solvents such as ethyl

acetate or acetone.

**[0076]** Illustrative, non-limiting examples of such preparations are given in the Example section below. One of skill in the art, having possession of this disclosure, will be able to modify the examples to come up with alternate crystallization methods and will be able to 5 determine whether such methods are capable of producing the desired crystalline salt.

**[0077]** In accordance with one embodiment of the invention, provided are pharmaceutical compositions comprising one or more crystalline salts as described herein.

In other embodiments, provided are uses of the crystalline salts in the preparation of medicaments and their use in treating a disease selected from the group consisting of

10 Type I diabetes, Type II diabetes and metabolic syndrome. The salts may also be used in a combination therapy with a DPP IV inhibitor.

**[0078]** In one embodiment, provided is a method for one or more of stimulating insulin production, stimulating glucose-dependent insulin secretion, lowering blood glucose, or lowering blood triglyceride levels, said method comprising administering to a subject in 15 need of such treatment an effective amount of a crystalline salt as described herein.

**[0079]** The DPP IV inhibitors useful in the present invention are sitagliptin (Merck), vildagliptin (Novartis), BMS-477118 (saxagliptin) (Bristol-Myers Squibb), R1438 (aminomethylpyridine) (Roche), NVP DPP728 (Novartis), PSN9301 (Prosidion), P32/98 (isoleucine thiazolidide) (Probiodrug), GSK823093C (Denagliptin) (Glaxo Smithkline),

20 SYR-322 (Alogliptin) (Takeda), NN-7201 (NovoNordisk), ALS2-0426 (Alantos). (Green BD, Flatt PR, Bailey CJ, Dipeptidyl peptidase IB (DPP IV) inhibitors: a newly emerging drug class for the treatment of Type II diabetes, *Diabetes Vasc Dis Res* 2006, 3:159-165) Preferred DPP IV inhibitors are sitagliptin, vildagliptin, Denagliptin, saxagliptin, and alogliptin). Even more preferred DPP IV inhibitors are sitagliptin and vildagliptin.

25 Sitagliptin is an approved pharmaceutical marketed as Januvia<sup>TM</sup>, and vildagliptin is an approved pharmaceutical marked as Galvus<sup>TM</sup>.

**[0080]** The crystalline salt and DPP IV inhibitor are administered in a single dosage or in separate dosages. The single dosage is administered once a day or multiple times a day. When administered as separate dosages, the dosages can be administered once a day 30 or multiple times a day.

**[0081]** In one embodiment, when the salt and the DPP IV inhibitor are administered in a single dosage, the salt and DPP IV inhibitor are formulated as a medicament into a single pill, single table, or a single capsule. When the salt and DPP IV inhibitor are administered in separate dosages, the salt is formulated as a medicament into a pill, tablet or capsule and the DPP IV inhibitor is formulated into a separate pill or capsule.

**[0082]** When the salt and DPP IV inhibitor are administered in separate dosages, the salt can be administered first and the DPP IV inhibitor can be administered next, following administration of the salt. Alternatively, the DPP IV inhibitor can be administered first and the salt can be administered next, following administration of the DPP IV inhibitor.

10 The time between the sequential first administration and the second administration can be varied by a skilled practitioner. In one embodiment, the first administration (the salt or DPP IV inhibitor), is followed immediately by the second administration (the salt or DPP IV inhibitor). In another embodiment, the second administration is within 2 minutes, 5 minutes, 10 minutes, 15 minutes, 30 minutes, or 60 minutes, 1 hour, 2 hours, 3 hours, 4

15 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, or 12 hours following the first administration. Yet another embodiment provides for the administration to a subject the salt and/or DPP IV inhibitor in the morning followed by administration to the previously treated subject the salt and/or DPP IV inhibitor in the evening. In another embodiment, the salt and DPP IV inhibitor are preferably

20 administered once a day.

**[0083]** Another aspect of this invention provides methods of lowering blood levels of glucose in a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure blood glucose levels before and 25 after administration of the salt and DPP IV inhibitor. Blood glucose levels are easily measured by numerous commercially available glucose monitoring devices that measure blood glucose from samples of blood or urine, or as taught herein. Blood glucose can also be measured by commercially available glucometers that do not require blood or urine samples.

30 **[0084]** Another aspect of this invention provides methods of lowering blood levels of insulin in a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the

mammal. The method further comprises steps to measure blood insulin levels before and after administration of the salt and a DPP IV inhibitor. Blood insulin levels are easily measured by well-known insulin monitoring assays that measure insulin from samples of blood or urine, or as taught herein.

5   **[0085]** In another aspect, this invention provides methods of increasing blood levels of incretins in a subject by administering the salt and a DPP IV inhibitor. The incretins are GLP-1 and GIP. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure blood incretin levels before and after administration of the salt and a DPP IV inhibitor. Blood  
10   incretin levels are easily measured by well-known incretin monitoring assays that, or as taught herein.

**[0086]** Yet another aspect of this invention provides methods of lowering blood triglyceride levels in a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure blood triglycerides levels before and after administration of the salt and DPP IV inhibitor. Blood triglyceride levels are easily measured by numerous commercially available devices that measure blood triglyceride levels from samples of blood.

20   **[0087]** A further aspect of this invention provides methods of lowing gastric emptying in a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure blood incretin levels before and after administration of the salt and a DPP IV inhibitor. Blood incretin levels are easily measured by well-known incretin monitoring assays, or as taught herein.

25   **[0088]** Another aspect of this invention provides methods of increasing insulin production in the islet cells of a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure insulin production in islet cells or the beta cells of the pancreas before and after administration of the salt and a  
30   DPP IV inhibitor. The insulin production of islets and beta cells are easily measured by well-known assays, or as taught herein.

**[0089]** In yet another aspect, this invention provides methods of preserving islet function in a subject by administering the salt and a DPP IV inhibitor. The method comprises administering an effective amount of the salt and DPP IV inhibitor to the mammal. The method further comprises steps to measure the function of islets' or beta cell's ability to produce insulin before and after administration of the salt and a DPP IV inhibitor. The insulin production of islets and beta cells are easily measured by well-known assays, or as taught herein.

**[0090]** A therapeutically effective amount of the salt and DPP IV inhibitor can be used for the preparation of one or more pharmaceutical compositions useful for treating Type II diabetes and/or lowering the plasma level of glucose. In addition, a therapeutically effective amount of the salt and a DPP IV inhibitor can be used for the preparation of one or more pharmaceutical compositions useful for treating other indications that include diabetes as a component, such as metabolic syndrome, as well as indications that can be improved as a result of increased insulin production (such as the early stages of Type I diabetes).

**[0091]** The compositions of the invention can include the salt and optionally DPP IV inhibitors, pharmaceutically acceptable salts thereof, or a hydrolysable precursor thereof. In general, the salt is mixed with suitable carriers or excipient(s) in a therapeutically effective amount. By a "therapeutically effective dose", "therapeutically effective amount", or, interchangeably, "pharmacologically acceptable dose" or "pharmacologically acceptable amount", it is meant that a sufficient amount of the compound of the present invention and a pharmaceutically acceptable carrier will be present in order to achieve a desired result, *e.g.*, alleviating a symptom or complication of Type II diabetes.

**[0092]** The MBX-2982 salts that are used in the methods of the present invention can be incorporated into a variety of formulations for therapeutic administration. More particularly, the salts can be formulated into pharmaceutical compositions by combination with appropriate, pharmaceutically acceptable carriers, excipients, or diluents, and can be formulated into preparations in solid or semi-solid forms such as tablets, capsules, pills, powders, granules, dragees, gels, ointments, suppositories, inhalants. Administration can be achieved in various ways, including oral, buccal, rectal, intradermal, and transdermal administration. Moreover, the salt can be administered in a local rather than systemic

manner, in a depot or sustained release formulation. In addition, the salts can be administered in a liposome.

**[0093]** Suitable formulations for use in the present invention are found in *Remington's Pharmaceutical Sciences* (Mack Publishing Company (1985) Philadelphia, PA, 17th ed.),

5 which is incorporated herein by reference. The pharmaceutical compositions described herein can be manufactured in a manner that is known to those of skill in the art, *i.e.*, by means of conventional mixing, granulating, dragee-making, levigating, emulsifying, encapsulating, or entrapping processes. The following methods and excipients are merely exemplary and are in no way limiting.

10 **[0094]** For oral administration, the salt can be formulated readily by combining with pharmaceutically acceptable carriers that are well known in the art. Pharmaceutical preparations for oral use can be obtained by mixing the compounds with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable  
15 excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone. If desired, disintegrating agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar,  
20 or alginic acid or the salt thereof such as sodium alginate.

**[0095]** Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions can be used, which can optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to  
25 the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

**[0096]** Pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture  
30 with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or

liquid polyethylene glycols. In addition, stabilizers can be added. All formulations for oral administration should be in dosages suitable for such administration.

**[0097]** For buccal administration, the compositions can take the form of tablets or lozenges formulated in conventional manner.

5   **[0098]** For administration by inhalation, the salts according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas, or from propellant-free, dry-powder inhalers. In the case of a pressurized aerosol the dosage 10 unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

15   **[0099]** The salts can also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter, carbowaxes, polyethylene glycols or other glycerides, all of which melt at body temperature, yet are solidified at room temperature.

20   **[0100]** In addition to the formulations described previously, the salts can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the salts can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

25   **[0101]** Alternatively, other delivery systems for hydrophobic pharmaceutical compounds can be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. In a presently preferred embodiment, long-circulating, *i.e.*, stealth liposomes can be employed. Such liposomes are generally described in Woodle, et al., U.S. Patent No. 5,013,556. The compounds of the present invention can also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; 30 and 4,008,719.

[0102] Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in a therapeutically effective amount. The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician. Determination of an effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

[0103] The amount of salt that can be combined with a carrier material to produce a single dosage form will vary depending upon the disease treated, the mammalian species, and the particular mode of administration. However, as a general guide, suitable unit doses for the salts can, for example, preferably contain between 0.1 mg to about 1000 mg of the active compound. A preferred unit dose is between 1 mg to about 500 mg. A more preferred unit dose is between 1 mg to about 300mg. Another preferred unit dose is between 1 mg to about 100 mg. Other preferred unit doses include 25, 50, 100, 150, 200, 250, 300, 350, 400, 450, and 500 mg. The unit doses can also be administered 1, 2, 3, 4, 5 or 6 times a day, preferably 1 or 2 times per day, or more preferably once a day so that the total dosage, for example, for a 70 kg adult is in the range of 0.001 to about 15 mg per kg weight of subject per administration. A preferred dosage is about 0.5 to about 10 mg, about 0.5 to about 7.5 mg, about 0.5 to about 5 mg, about 0.5 to about 4 mg, about 0.5 to about 3 mg, about 0.5 to about 2 mg, or about 0.5 to about 1 mg per kg weight of subject per administration, and such therapy can extend for a number of weeks or months, and in some cases, years. It will be understood, however, that the specific dose level for any particular subject will depend on a variety of factors including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the individual being treated; the time and route of administration; the rate of excretion; other drugs that have previously been administered; and the severity of the particular disease undergoing therapy, as is well understood by those of skill in the area.

[0104] A typical dosage can be one 1 mg to about 500 mg, or a 25, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, or 500 mg tablet taken once a day or as a time-release capsule or tablet taken once a day and containing a proportionally higher content of active ingredient. The time-release effect can be obtained by capsule materials that

dissolve at different pH values, by capsules that release slowly by osmotic pressure, or by any other known means of controlled release.

**[0105]** In addition, the present invention provides for kits with unit doses of the salt and/or DPP IV inhibitor, either in oral or injectable doses. In addition to the containers containing the unit doses will be an informational package insert describing the use and attendant benefits of the drugs in treating diabetes, obesity, hyperlipidemia, atherosclerosis and metabolic syndrome, and/or their respective related symptoms, complications and disorders.

**[0106]** The following examples are provided to further illustrate certain aspects of the present invention and to aid those of skill in the art in practicing the invention. These examples are not meant to limit the scope of the invention.

## EXAMPLES

**[0107]** The following abbreviations are used in the Examples and throughout the application.

ACN	Acetonitrile
DCM	Dichloromethane
DEE	Diethyl ether
EtOAc	Ethyl acetate
EtOH	Ethanol
HFIP	Hexafluoroisopropanol
IPA	Isopropyl alcohol
IPE	Isopropyl ether
MEK	Methyl ethyl ketone
MeOH	Methanol
MTBE	Methyl- <i>tert</i> -butyl ether
TFE	2,2,2-Trifluoroethanol
THF	Tetrahydrofuran
CCS	Crash cooling of a solution
FC	Fast cooling
FE	Fast evaporation
RE	Rotary evaporation
S/AS	Solvent/Anti-solvent precipitation
SC	Slow cooling
SE	Slow evaporation
VD	Vapor diffusion
VO	Vacuum Oven
DSC	Differential scanning calorimetry
NMR	Nuclear magnetic resonance spectroscopy
TG / TGA	Thermogravimetric analysis

XRPD	X-ray powder diffraction
B/E	Birefringence/extinction
NS	No solids
Ppt	Precipitation
RH	Relative humidity
RT	Room temperature

### Instrumental Techniques

**[0108]** XRPD: XRPD patterns were collected using an Inel XRG-3000 diffractometer or PANalytical X'Pert Pro diffractometer.

5      **[0109]** Inel: Inel XRG-3000 diffractometer equipped with a curved position sensitive detector with a  $2\theta$  range of  $120^\circ$ . An incident beam of Cu  $K\alpha$  radiation (40 kV, 30 mA) was used to collect data in real time at a resolution of  $0.03^\circ 2\theta$ . Prior to the analysis, a silicon standard (NIST SRM 640c) was analyzed to verify the Si 111 peak position. Samples were prepared for analysis by packing them into thin-walled glass capillaries.

10     Each capillary was mounted onto a goniometer head and rotated during data acquisition.

**[0110]** PANalytical: An incident beam of Cu  $K\alpha$  radiation was produced using an Optix long, fine-focus source. An elliptically graded multilayer mirror was used to focus the Cu  $K\alpha$  X-rays of the source through the specimen and onto the detector. Data were collected and analysed using X'Pert Pro Data Collector software (v. 2.2b). Prior to the analysis, a silicon specimen (NIST SRM 640c) was analyzed to verify the Si 111 peak position. The specimen was sandwiched between 3  $\mu\text{m}$  thick films, analyzed in transmission geometry, and rotated to optimize orientation statistics. Soller slits were used for the incident and diffracted beams to minimize axial divergence. Diffraction patterns were collected using a scanning position-sensitive detector (X'Celerator) located 240 mm from the specimen.

**[0111]** DSC: DSC was performed using a TA Instruments Q2000 differential scanning calorimeter. Temperature calibration was performed using NIST traceable indium metal. The sample was placed into an aluminum DSC pan, and the weight was accurately recorded. The pan was either covered with a lid perforated with a laser pinhole, and the lid was hermetically sealed or covered with an unperforated lid and crimped. The sample cell was equilibrated at either  $-30^\circ\text{C}$  or  $25^\circ\text{C}$  and heated under a nitrogen purge at a rate

of 10 °C/minute, up to a final temperature of 250 °C. Reported temperatures are at the transition maxima.

**[0112]** TGA: TG analyses were performed using a TA Instruments Q5000 IR thermogravimetric analyzer. Temperature calibration was performed using nickel and Alumel<sup>TM</sup>. Each sample was placed in an aluminum pan. The pan was hermetically sealed with a lid that was opened using a punching mechanism just before being inserted into the TG furnace. The furnace was heated under nitrogen at a rate of 10 °C/minute to a final temperature of 350 °C.

**[0113]** <sup>1</sup>H NMR: The solution NMR spectra were acquired with a Varian <sup>UNITY</sup> INOVA-400 spectrometer. The samples were prepared by dissolving them in CDCl<sub>3</sub> or in DMSO<sub>6</sub>.

**[0114]** FT-Raman spectroscopy: Raman spectra were acquired on a Nexus 670 FT-Raman accessory module interfaced to a Nexus 670 FT-IR spectrophotometer (Thermo Nicolet) equipped with an indium gallium arsenide (InGaAs) detector. Wavelength verification was performed using sulfur and cyclohexane. Each sample was prepared for analysis by placing the sample into a glass tube and positioning the tube in a gold-coated tube holder.

#### Crystallization Methods

**[0115]** Slow Cool (SC): Solutions containing free base and an acid of interest were prepared in various solvents at room temperature. Solids persisted and, the samples were heated to facilitate dissolution. Once a clear (solids-free) solution was obtained, the solutions were allowed to slowly cool to room temperature.

**[0116]** Volume Reduction (VR): Solutions containing free base and an acid of interest were prepared in various solvents at room temperature. No solids were seen in solution. Sample capped and left at ambient temperature for a period of hours to days. If no solids were generated, the sample was uncapped and the samples volume was reduced. Sample capped and allowed to stand under ambient temperature conditions. Once solids precipitated from solution, the solids were collected via vacuum filtration and dried.

**[0117]** Precipitation (Ppt): Solutions containing free base and an acid of interest were prepared in various solvents at room temperature. If solids persisted, the samples were either heated to facilitate dissolution or kept at ambient temperature and stirred. If a clear

solution resulted, the sample was capped and kept at ambient temperature. The samples at elevated temperature were cooled to ambient temperature. Generated solids were collected via vacuum filtration and dried.

Example 1: Preparation free base 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-

5 thiazol-2-yl]-piperidin-1-yl}-pyrimidine.

Synthesis of *tert*-butyl 4-((4-(1*H*-tetrazol-1-yl)phenoxy)methyl)thiazol-2-yl)piperidine-1-carboxylate

**[0118]** A mixture of 4-(4-chloromethyl-thiazol-2-yl)-piperidine-1-carboxylic acid *tert*-butyl ester (549 mg), 4-tetrazol-1-yl-phenol (270 mg), Cs<sub>2</sub>CO<sub>3</sub> (890 mg) in acetonitrile was heated under reflux overnight. After cooling, the reaction mixture was filtered through a pad of celite, concentrated *in vacuo*. Purification by chromatography (40-100% EtOAc/Hexanes) gave the desired product as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.01 (1H, s), 7.61 (2H, d, *J* = 8.8 Hz), 7.25 (1H, s), 7.15 (2H, d, *J* = 8.8 Hz), 5.22 (2H, s), 4.2 (2H, br), 3.17 (1H, m), 2.87 (2H, m), 2.11 (2H, m), 1.73 (2H, m), 1.46 (9H, s).

15 Synthesis of 4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl]-piperidine hydrochloride

**[0119]** To a solution of *tert*-butyl 4-((4-(1*H*-tetrazol-1-yl)phenoxy)methyl)thiazol-2-yl)piperidine-1-carboxylate (0.60 g) in methanol/dichloromethane (1.0 mL/ 1.5 mL) was added 4N HCl in dioxane (1.7 mL) at 0°C, and then stirred at room temperature for 7 hours. After removal of solvents in vacuo, a crude desired compound HCl salt was obtained as an off-white solid.

20 **[0120]** A mixture of 4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl]-piperidine hydrochloride (403 mg), 2-chloro-5-ethylpyrimidine (0.15 mL) and

diisopropylethylamine (1 mL) in isopropanol was heated at 90°C overnight, partitioned between EtOAc and water. The organic layer was washed with water / brine, and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. Purification by chromatography on silica gel ( 40-100% EtOAc/hexanes) gave the desired compound as a white solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 9.98 (1H, s), 8.24 (2H, s), 7.80 (2H, d, *J* = 8.8 Hz), 7.66 (1H, s), 7.28 (2H, d, *J* = 8.8 Hz), 5.20 (2H, s), 4.67 (2H, m), 3.32 (1H, m), 3.01 (2H, m), 2.43 (2H, q, *J* = 7.2 Hz), 2.07 (2H, m), 1.59 (2H, m), 1.11 (3H, t, *J* = 7.2 Hz) ppm.

## Example 2: Besylate salt

**[0121]** The besylate salt was generated from an acetone solution volume reduction experiment. The material crystallized as stacked plates. XRPD analysis indicated the besylate salt was crystalline. After overnight RH stress, the material remained a free-flowing powder. Thermal analysis of the stressed sample indicated a slight weight loss (0.9 wt. %) to 125°C and a sharp endotherm centered at 153°C. Associated with the endotherm was a shoulder at approximately 147°C. Rapid weight loss was observed above approximately 155°C. <sup>1</sup>H NMR spectroscopy results indicated a 1:1 salt was present.

10     <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.95 (1H, s), 8.54 (2H, br s), 7.95 (2H, m), 7.63 (2H, dt, *J* = 3, 9.2 Hz), 7.40 (3H, m), 7.29 (1H, s), 7.18 (2H, dt, *J* = 3, 9.2), 5.24 (2H, s), 4.76 (2H, m), 3.42 (3H, m), 2.61 (2H, q, *J* = 7.6 Hz), 2.33 (2H, dd, *J* = 3, 13.8 Hz), 1.97 (2H, m), 1.26 (3H, t, *J* = 7.6 Hz).

## Example 3: Camsylate salt

15     **[0122]** The camsylate salt was generated by a precipitation reaction and a slow cool in acetone. The material crystallized in an unknown morphology. XRPD analysis indicated the camsylate salt was crystalline. After overnight RH stress, the material remained a free flowing powder. Thermal analysis of the unstressed material indicated a slight weight loss (0.3 wt. %) to 125°C and a sharp endotherm centered at 184°C. Rapid weight loss, usually indicative of decomposition, was observed above approximately 180°C. <sup>1</sup>H NMR spectroscopy results indicated that a 1:1 salt had formed.

20     <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.70 (1H, s), 8.61 (2H, br s), 7.74 (2H, d, *J* = 9.2 Hz), 7.28 (1H, s), 7.18 (2H, d, *J* = 9.2 Hz), 5.24 (2H, s), 4.78 (2H, d, *J* = 13.6 Hz), 3.42 (4H, m), 2.92 (1H, d, *J* = 15.2 Hz), 2.73 (1H, m), 2.62 (2H, q, *J* = 7.6 Hz), 2.34 (3H, m), 1.99 (4H, m), 1.89 (1H, d, *J* = 18.4 Hz), 1.79 (1H, m), 1.39 (1H, m), 1.27 (3H, t, *J* = 7.6 Hz), 1.12 (3H, s), 0.86 (3H, s).

## Example 4: Esylate salt

**[0123]** The esylate salt was generated through a precipitation reaction in acetone. The esylate material existed as stacked plates and tablets. XRPD analyses indicated that the esylate salt was crystalline. After overnight stress, the material remained a free flowing powder. Thermal analysis of the stressed powder indicated a 4.8% weight loss to 125°C

and a sharp endotherm centered at 98.5°C. The endothermic event is typically indicative of a melting event. The weight loss could be indicative of a solvate or hydrate of the salt. Rapid weight loss, usually indicative of decomposition, was observed above approximately 125°C. <sup>1</sup>H NMR spectroscopy results indicated a 1:1 salt had formed.

5      <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.95 (1H, s), 8.56 (2H, br s), 7.63 (2H, d, *J* = 9.2 Hz), 7.29 (1H, s), 7.18 (2H, d, *J* = 9.2 Hz), 5.24 (2H, s), 4.79 (2H, m), 3.43 (3H, m), 2.97 (2H, q, *J* = 7.6 Hz), 2.61 (2H, q, *J* = 7.6 Hz), 2.36 (2H, m), 1.98 (2H, m), 1.39 (3H, t, *J* = 7.6 Hz), 1.27 (3H, t, *J* = 7.6 Hz).

Example 5: HBr salt

10     [0124] The HBr salt was generated through a precipitation reaction in acetone. The material crystallized in an unknown morphology. XRPD results indicated that the HBr salt was crystalline. After overnight stress, the material remained a free flowing powder. Thermal analysis of the unstressed material indicated that a wide, weak endotherm existed prior to the sample undergoing decomposition. A 1.5% wt. loss was detected prior to 15 decomposition. <sup>1</sup>H NMR spectroscopy indicated that no decomposition product was formed under ambient conditions. The salt precipitated in the presence of CDCl<sub>3</sub> and the <sup>1</sup>H NMR analysis was performed in DMSO.

1      <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 10.00 (1H, s), 8.30 (2H, s), 7.82 (2H, d, *J* = 9.2 Hz), 7.68 (1H, s), 7.30 (2H, d, *J* = 9.2 Hz), 5.22 (2H, s), 4.66 (2H, d, *J* = 13.2 Hz), 3.36 (1H, tt, *J* = 3.8, 11.6 Hz), 3.08 (2H, dt, *J* = 2.6, 12.8 Hz), 2.50 (2H, q, *J* = 7.4 Hz), 2.12 (2H, m), 1.64 (2H, m), 1.14 (3H, t, *J* = 7.4 Hz).

Example 6: Mesylate salt

20     [0125] The mesylate salt was generated in a variety of morphologies using a slow cool/volume reduction technique. XRPD results indicated that the mono-salt was 25 crystalline. After elevated RH stress, the material remained a free flowing powder. Thermal analysis of the post-stress sample indicated an endotherm centered at 86°C and a weight loss of 4.5% up to 125°C. One possible explanation is that solvent (probably water) loss occurs quickly and the sample undergoes decomposition after a melting event. <sup>1</sup>H NMR spectroscopy results confirmed a 1:1 salt was formed.

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  8.95 (1H, s), 8.55 (2H, br s), 7.63 (2H, d, *J* = 8.8 Hz), 7.29 (1H, s), 7.18 (2H, d, *J* = 9.2 Hz), 5.24 (2H, s), 4.76 (2H, m), 3.42 (3H, m), 2.88 (1H, s), 2.61 (2H, q, *J* = 7.6 Hz), 2.36 (2H, m), 1.99 (2H, m), 1.27 (3H, t, *J* = 7.6 Hz).

5 [0126] XRPD (Intel XRG-3000 diffractometer) select interlattice plane intervals: d in [Å] ( $\pm 0.1$  Å): 10.23, 6.28, 4.83, 4.75, 4.57, 4.43, 4.32, 3.49, 3.43, 3.32.

Example 7: Sulfate salt

10 [0127] The sulfate salt was generated from precipitation experiments. XRPD results indicated that the mono-salt generated crystalline material. After elevated RH stress, the material turned tacky. Thermal analysis of the unstressed sample indicated a large weight loss (9% to 125°C) was associated with an endotherm centered at 97°C. After this initial weight loss, rapid degradation of the material occurred upon further heating which may be due the loss of water, although this hypothesis was not verified. <sup>1</sup>H NMR spectroscopy results indicated that no decomposition product was formed. The salt precipitated in the presence of CDCl<sub>3</sub> and the <sup>1</sup>H NMR analysis was performed in DMSO.

15 [0128] <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>):  $\delta$  10.00 (1H, s), 8.32 (2H, s), 7.82 (2H, d, *J* = 9.2 Hz), 7.68 (1H, s), 7.30 (2H, d, *J* = 9.2 Hz), 5.22 (2H, s), 4.65 (2H, d, *J* = 13.6 Hz), 3.36 (1H, tt, *J* = 3.6, 11.2 Hz), 3.09 (2H, m), 2.48 (2H, q, *J* = 7.6 Hz), 2.12 (2H, m), 1.65 (2H, m), 1.14 (3H, t, *J* = 7.6 Hz).

Example 8: Tosylate salt

20 [0128] The tosylate salt was generated by slow cooling and volume reduction in acetone. The material crystallized in an unknown morphology. XRPD results indicated that crystalline material existed. The material remained a free flowing powder after exposure to elevated RH conditions. Thermal analysis of the material indicated minor (0.9%) weight loss to 125°C associated with a very weak endotherm centered at 94°C.

25 This wide, weak endotherm could be indicative of solvent loss. An endotherm centered at 152°C preceded a large loss in weight. <sup>1</sup>H NMR spectroscopy results confirmed a 1:1 salt had formed.

[0128] <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  8.95 (1H, s), 8.54 (2H, br s), 7.83 (2H, d, *J* = 8.0 Hz), 7.63 (2H, d, *J* = 9.2 Hz), 7.30 (1H, s), 7.19 (4H, m), 5.25 (2H, s), 4.76 (2H, m), 3.43 (3H, m), 2.61 (2H, q, *J* = 7.6 Hz), 2.36 (3H, s), 2.33 (2H, m), 1.96 (2H, m), 1.26 (3H, t, *J* = 7.6 Hz).

**[0129]** XRPD (Intel XRG-3000 diffractometer) select interlattice plane intervals: d in [Å] ( $\pm 0.1$  Å): 10.61, 8.41, 5.78, 5.17, 5.02, 4.84, 4.37, 4.22, 3.95, 3.44.

Example 9: HCl salt

**[0130]** A screen of various solvent systems for preparing the HCl salts was undertaken  
5 according to the following procedures, with the results shown in Table 6 below.

**[0131]** Evaporation Experiments: HCl salt was dissolved in a given solvent. The solution was filtered through a 0.2 µm nylon filter. For evaporation experiments at ambient, the solutions were left in open vials (fast evaporation) or covered with aluminum foil containing pinholes (slow evaporation). For evaporation experiments under vacuum  
10 (rotary evaporation), the sample was placed on the rotary evaporator at ambient or elevated temperature and the solvents evaporated to dryness.

**[0132]** Slow and Fast Cool Experiments: HCl salt was contacted with a given solvent and the sample was brought to elevated temperature in an oil bath on a hotplate. Selected samples were filtered using a 0.2 µm nylon filter. The heat source was then turned off and  
15 the hotplate and vials were allowed to cool slowly in the oil bath to ambient temperature for slow cool or placed on the lab bench for fast cool. Selected samples that did not produce solids at ambient temperature were placed in a refrigerator or freezer. Solids were recovered by vacuum filtration.

**[0133]** Crash Cool Experiments: Saturated solutions were prepared in various solvents  
20 at elevated temperature. Experiments were performed in an oil bath placed on a hotplate. The resulting solutions or slurries were rapidly filtered through a warm 0.2 µm filter into an open vial while still warm. The vial was placed into an acetone bath cooled by dry ice. Solids were collected by vacuum filtration.

**[0134]** Vapor Diffusion: Small amounts of MBX 2982 HCl were dissolved in a  
25 minimum amount of an appropriate solvent. The samples were filtered through a 0.2 µm nylon filter into a 1 dram vial. Diethyl ether was added to a 20 mL scintillation vial. The 1 dram vials were uncapped and placed into the 20 mL vials. The 20 mL vials were capped and parafilmed.

Table 6.

Solvent	Conditions	Observation	XRPD Results
Acetone: MeOH (~3:1)	FC from ~53 °C to RT (cloudy), sonicated, left standing, ~3 days	Needle-like, agglomerates, some B/E on small particles	Form II
Chloroform	SC from ~53 °C to RT (clear), sonicated, left standing, ~3 days	NS	
DCM	FE	Unknown morphology, agglomerates, B/E	Pattern E
DCM	Agitation w/stirring, RT, ~3 days <sup>a</sup>	Unknown morphology, agglomerates, no B/E	Form I
DCM/DEE	VD	Fibrous or needle-like, agglomerates, some B/E	Pattern C
EtOH:water (2:1)	Slurry w/stirring, RT, ~1 day	Unknown morphology, agglomerates, some B/E on very small particles	Form I + free base Form II
IPA	Slurry, ~60 °C, ~3 days	Unknown morphology to granule-like, some B/E	Form I
MeOH	SE	Unknown morphology, agglomerates, some B/E on smaller particles	Form I
MeOH	Slurry w/stirring, RT, ~3 days	Unknown morphology, agglomerates, no B/E	Form I
MeOH	CCS (acetone/dry ice) from ~58 °C	Unknown morphology, agglomerates, some B/E on small particles	Form I + II
MeOH/ACN	S/AS attempt (clear). Kept in freezer	NS	
MeOH/DEE	VD	Fibrous + needle-like, + opaque, agglomerates some B/E	Pattern L
MeOH:EtOAc (1:1)	SC from ~57 °C to RT	Rosettes to needle-like, agglomerates, B/E	Form II
MeOH:THF (1:1)	Slurry, RT, ~5 days	Unknown morphology, agglomerates, B/E on few small particles	Form I
MeOH:EtOAc (1:1)	Slurry, ~40 °C, ~2 days	Unknown morphology, agglomerates, some B/E on few particles	Form I
Nitromethane	Slurry, ~60 °C, ~3 days	Unknown morphology, agglomerates, some B/E on smaller particles	Form I
Nitromethane	SC from ~69 °C to RT (clear), sonicated, kept in refrigerator, 4 days	Unknown to needle-like particles, agglomerates, some B/E on smaller particles	Pattern J

TFE	RE, RT to ~40 °C (film)	Unknown morphology, generally opaque, some B/E on very small particles	Pattern H
TFE:ACN (19:1)	Slurry, ~40 °C, ~2 days	Unknown morphology to granule-like, agglomerates, some B/E on smaller particles	Form I
TFE/DEE	VD	Unknown morphology, agglomerates, some B/E on smaller particles	Form I
TFE/EtOAc	S/AS attempt (clear). Kept in freezer	Unknown morphology to rosettes, agglomerates, some B/E on smaller particles	Pattern G
THF:TFE (1:1)	SC from ~53 °C to RT (clear), sonicated, kept in freezer, ~4 days	Unknown morphology, agglomerates, some B/E on smaller particles	Pattern K
Water	Slurry, ~38 °C, ~1 day	Unknown morphology, agglomerates, some B/E on small particles	Pattern D

Example 10: HCl salt Form I

**[0135]** The free base (46.0 g) contacted with 500 mL EtOH and warmed/stirred. 9.5

mL conc. HCl added to the suspension. Sample was left for approximately 30 minutes

5 and then cooled to RT. The filtered material was vacuum dried (3 days) and identified as “pattern O” containing trace amounts of ethanol.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.95 (1H, s), 8.42 (2H, br s), 7.63 (2H, d, *J* = 9.2 Hz), 7.28 (1H, s), 7.18 (2H, d, *J* = 9.2 Hz), 5.24 (2H, s), 4.97 (2H, m), 3.46 (3H, m), 2.61 (2H, q, *J* = 7.6 Hz), 2.37 (2H, m), 1.99 (2H, m), 1.27 (3H, t, *J* = 7.6 Hz).

10 **[0136]** The solids were re-combined with original mother liquor and sample was

rapidly stirred and an additional 0.5mL conc. HCl added. Sample was heated and stirred

for approximately 30 minutes and then allowed to slowly cool to RT. The filtered

material was slurried in acetone (26.8 g/125 mL) and warmed for approximately one (1)

hour. The sample was slowly cooled to ambient temperature and slurried for six days to

15 give HCl Form I.

**[0137]** The results of thermal (DSC, TGA) characterization of HCl Form I indicated

that the material is likely unsolvated. The TGA curve indicates a ~0.2 % weight loss

between ~23 °C and ~73 °C, likely associated with residual acetone evaporation

(presence of residual acetone is seen from <sup>1</sup>H NMR). A weight loss of approximately 13.1

20 % between ~113 °C and ~210 °C was observed, followed by a sharp loss at ~302 °C

(onset) likely due to decomposition. The material may initially lose the HCl, which is accompanied by further degradation at higher temperatures. The DSC thermogram exhibited a sharp endothermic event at ~191 °C (onset) followed by immediate heat fluctuation, likely attributable to decomposition.

5 [0138] Moisture sorption results showed a ~0.3 wt% loss upon equilibration at ~5% RH. The small weight loss is comparable with the TGA loss, and is likely associated with the loss of residual acetone. A steady ~0.9 wt% gain was observed between ~5% and ~95% RH. A complete desorption occurred upon decreasing relative humidity to ~5% (~0.9 wt% loss between ~95% and ~5% RH).

10 [0139] The <sup>1</sup>H NMR showed significant shift and broadening of peak at 8.42 ppm attributable to protons of pyrimidine ring and traces of residual acetone (~0.06 moles of acetone per mole of free base).

15 <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.96 (1H, s), 8.42 (2H, br s), 7.63 (2H, d, *J* = 8.8 Hz), 7.28 (1H, s), 7.18 (2H, d, *J* = 8.8 Hz), 5.24 (2H, s), 4.98 (2H, m), 3.45 (3H, m), 2.61 (2H, q, *J* = 7.6 Hz), 2.38 (2H, m), 1.99 (2H, m), 1.27 (3H, t, *J* = 7.6 Hz).

[0140] The elemental analysis data was consistent with the material being a monohydrochloride salt: (C<sub>22</sub>H<sub>25</sub>ClN<sub>8</sub>OS) C: 54.16%; H: 5.29%; N: 22.89%; Cl: 7.40%.

20 [0141] FT-Raman spectrum: (cm<sup>-1</sup>): 154, 193, 214, 243, 324, 346, 384, 420, 460, 467, 495, 539, 573, 596, 628, 648, 680, 687, 718, 751, 766, 787, 800, 845, 856, 946, 967, 992, 1009, 1051, 1068, 1094, 1110, 1172, 1203, 1225, 1253, 1282, 1300, 1311, 1334, 1375, 1394, 1404, 1432, 1458, 1474, 1480, 1517, 1533, 1591, 1609, 1637, 2734, 2869, 2903, 2928, 2937, 2967, 2999, 3015, 3030, 3063, 3075.

Example 11: HCl salt Form I from acetone

25 [0142] To a suspension of MBX-2982 (0.9 g) in acetone (4 mL) was added one equivalent of HCl (concentrated aqueous solution) at 55°C. The suspension was stirred at 55°C for two hours and then cooled down to room temperature. The HCl salt Form I was collected by vacuum filtration as solids. Select interlattice plane intervals from the X-ray powder pattern taken from an Intel XRG-3000 diffractometer: *d* in [Å] ( $\pm 0.1$  Å): 10.09, 8.20, 5.51, 5.08, 4.36, 4.26, 4.14, 4.08, 3.35, 3.18.

30 Example 12: HCl salt Form I from ethyl acetate

[0143] To a suspension of MBX-2982 (2 g) in ethyl acetate (9 mL) was added 1.05 equivalent of HCl (1 M solution in ethyl acetate) at 55°C. The suspension was stirred at 55°C for two hours and then cooled down to room temperature. The HCl salt Form I was collected by vacuum filtration as solids. Select interlattice plane intervals from the X-ray powder pattern taken from an Intel XRG-3000 diffractometer: d in [Å] ( $\pm 0.1$  Å): 10.09, 8.22, 5.51, 5.09, 4.36, 4.26, 4.14, 4.09, 3.35, 3.18.

Example 13: HCl salt Form II:

[0144] 21.5 g pattern O material (prepared as in Example 10) was contacted with 125 mL MeOH and HCl Form II seeds. Sample was slurried for 6 days at RT to give Form II material.

[0145] XRPD analysis of HCl Form II shows that the material is crystalline.

[0146]  $^1$ H NMR analysis of HCl Form II indicated significant shift and broadening of the peak at ~8.42 ppm, attributable to protons in the pyrimidine ring, and suggest that HCl is likely positioned near the pyrimidine nitrogen atoms. In addition, the spectrum also showed traces of methanol (0.3 moles of methanol per mole of base) and residual ethyl acetate.

$^1$ H NMR (CDCl<sub>3</sub>):  $\delta$  8.98 (1H, s), 8.42 (2H, br s), 7.63 (2H, d, *J* = 8.8 Hz), 7.29 (1H, s), 7.18 (2H, d, *J* = 9.2 Hz), 5.24 (2H, s), 4.98 (2H, m), 3.47 (3H, m), 2.61 (2H, q, *J* = 7.6 Hz), 2.37 (2H, m), 1.99 (2H, m), 1.27 (3H, t, *J* = 7.6 Hz).

[0147] The results of thermal (DSC, TGA) characterization of Form II suggest that the material is likely unsolvated. The TG curve indicates a ~0.6 % weight loss between ~21 °C and ~113 °C, likely associated with residual methanol and ethyl acetate evaporation. A weight loss of approximately 14.6 % between ~113 °C and ~220 °C was observed, followed by a sharp loss at ~312 °C (onset) likely due to decomposition. The material may initially lose the HCl, which is accompanied by further degradation at higher temperatures. The DSC thermogram exhibited an endothermic event beginning ~150 °C (onset) followed by a number of heat fluctuations, likely attributable to decomposition.

Biological Example 1

[0148] Two studies were conducted in order to compare the systemic exposure and pharmacokinetics (PK) of microcrystalline MBX-2982 and salt forms of MBX-2982 in

fasted male Sprague Dawley (SD) rats following single oral gavage (PO) doses of 200 mg/kg. Four salt forms of MBX-2982 were investigated in the studies. The tested salts were HCl Form I, HCl salt Form II, mesylate salt and the tosylate salt.

**[0149] Materials and Equipment**

5 Standard: MBX-2982

Internal standard: MBX-2982, where the six phenyl ring carbon atoms are labeled as  $^{13}\text{C}$   
Blank pooled rat plasma (Bioreclamation)

Chemicals: Reagent grade

Solvents: HPLC grade

10 96-well deep plate and mat: 1 mL (Corning)

HPLC Column: Lunar C18 (2) , 5  $\mu$ , 50 x 2.1 mm I.D. (Phenomenex)

Pre-column filter: 0.2  $\mu\text{m}$  (Thermo- Fisher Scientific)

LC System: LC-20 AD Pumps and SCL-10A VP LC controller with CTC Analytics AG  
Pal Autosampler (Shimadzu Scientific Instruments, Inc.)

15 Mass Spectrometer: 4000 Q-TRAP® with Analyst 1.4.2 Software (Applied Biosystem  
Inc.)

Analytical balance: model accu-124 (Fisher Scientific)

Pipets: (Rainin Instrument, LLC)

**[0150] Sample Preparation**

20 **[0151]** Preparation of samples was performed by solvent precipitation of plasma  
proteins in a 1 mL/well 96-well plate. Standards were prepared by spiking 10  $\mu\text{L}$  of blank  
plasma with 10  $\mu\text{L}$  of standard solution (acetonitrile containing MBX-2982 at 0.01, 0.02,  
0.05, 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 50 and 100  $\mu\text{g/mL}$ ) and 300  $\mu\text{L}$  of 0.1% formic acid in  
acetonitrile containing 0.2  $\mu\text{g/mL}$  internal standard. Plasma samples were prepared by  
25 adding 10  $\mu\text{L}$  of plasma sample, 10  $\mu\text{L}$  of acetonitrile, and 300  $\mu\text{L}$  of 0.2  $\mu\text{g/mL}$  internal  
standard in 0.1% formic acid in acetonitrile. After addition of the organic solvents, all  
samples were vortexed briefly and centrifuged at 3600 rpm for 10 minutes. Some of the  
samples were diluted 1.3 to 2- fold with blank plasma. An aliquot of the supernatant (25  
30  $\mu\text{L}$ ) was transferred to a 1 mL/well 96-well plate and mixed with 200  $\mu\text{L}$  water/ACN (50/50,  
v/v) and injected into the HPLC.

**[0152]** All stock and spiking solutions were kept in polypropylene tubes and stored at approximately -80°C.

**[0153]** The suspensions for microcrystalline MBX-2982 (free base) and salt forms of MBX-2982 were prepared in 1% carboxymethylcellulose and 2% tween 80 in water

5 (w/w/v).

**[0154]** Five groups of animals were given microcrystalline and salt forms suspension doses of MBX-2982 orally at 200 mg/kg, respectively. Food was withheld the night before dose and returned nine hours after dosing. Blood samples were collected at 0.25, 0.5, 1, 2, 4, 6, 9, 24, 30 and 48 h post dose. MBX-2982 plasma concentration from these 10 samples was analyzed by a non-validated high performance liquid chromatography in conjunction with mass spectrometry (LC-MS/MS) method.

**[0155]** Plasma concentration-time data for individual animals were analyzed using WinNonlin software (Professional, version 5.0.1; Pharsight Corp.). A non-compartmental model (model 200) was used. Peak area ratios of MBX-2982 to internal standard and concentration were fitted to a quadratic equation (Calibration Curve) with 1/x weighting using a quadratic regression program in Analyst version 1.4.2 (Applied Biosystem Inc). The equations were then used to interpolate the concentrations of MBX-2982 in samples 15 from peak area to internal standard ratio.

**[0156]** The following pharmacokinetic parameters were calculated:

20 1)  $C_{max}$ : Maximum (peak) plasma concentration  
2)  $T_{max}$ : Time at which  $C_{max}$  occurred  
3)  $t_{1/2}$ : Terminal phase half-life  
4)  $AUC_{0-t}$ : Area under the concentration-time curve from time 0 to last measurable concentration  
25 5)  $AUC_{0-\infty}$ : Area under the concentration-time curve from time 0 extrapolated to infinity

**[0157]** The mean pharmacokinetic parameters of MBX-2982 in male SD rats following single 200 mg/kg PO doses are presented in Table 7. Mean plasma concentration-time 30 data and profiles are presented in Table 8 and Figure 23.

**[0158]** The results show that the tested salt forms of MBX-2982 gave higher drug exposure compared with micro MBX-2982 (free base) when dosed with equivalent

amounts. The systemic drug exposures in  $AUC_{0-t}$  increased 2.4-fold, 2.9-fold, 1.4-fold and 1.7 fold for HCl salt Form I, HCl form II, mesylate salt and tosylate salt, respectively.

**Table 7**

Mean pharmacokinetic parameters (Mean  $\pm$  SD) of MBX-2982 in Fasted male SD rats following single PO dose of 200 mg/kg

Parameter	MBX-2982	HCl Salt Form I	HCl Salt Form II	Mesylate Salt	Tosylate Salt
$t_{1/2}$ (h)	3.53 $\pm$ 0.70	4.01 $\pm$ 1.68	2.09 $\pm$	4.11 $\pm$ 1.29	3.20 $\pm$ 0.88
$T_{max}$ (h)	5.88 $\pm$ 1.55	7.25 $\pm$ 9.25	11.5 $\pm$ 12.4	4.50 $\pm$ 1.00	4.00 $\pm$ 1.63
$C_{max}$ ( $\mu$ g/mL)	17.4 $\pm$ 4.19	31.2 $\pm$ 7.41	31.8 $\pm$ 5.05	26.7 $\pm$ 4.46	26.1 $\pm$ 7.98
$AUC_{0-24}$ ( $\mu$ g*h/mL)	233 $\pm$ 64.8	454 $\pm$ 98.4	482 $\pm$ 69.4	314 $\pm$ 65.7	342 $\pm$ 117
$AUC_{0-t}$ ( $\mu$ g*h/mL)	236 $\pm$ 64.6	571 $\pm$ 249	679 $\pm$ 208	322 $\pm$ 63.6	392 $\pm$ 98.2
$AUC_{0-inf}$ ( $\mu$ g*h/mL)	1.89 $\pm$ 60.7	536 $\pm$ 29.9	469 $\pm$	323 $\pm$ 62.0	393 $\pm$ 98.1

**Table 8**

Mean concentrations ( $\mu$ g/mL; mean  $\pm$  SD) at the indicated time points of MBX-2982 in Fasted male SD rats following single PO dose of 200 mg/kg

Time (h)	MBX-2982	HCl Salt Form I	HCl Salt Form II	Mesylate Salt	Tosylate Salt
0.5	3.89 $\pm$ 0.95	9.79 $\pm$ 3.73	8.06 $\pm$ 1.61	8.52 $\pm$ 0.975	10.1 $\pm$ 2.45
1	6.70 $\pm$ 1.27	15.5 $\pm$ 3.15	16.3 $\pm$ 2.77	12.2 $\pm$ 0.839	13.0 $\pm$ 3.33
2	11.6 $\pm$ 1.61	24.2 $\pm$ 3.35	23.7 $\pm$ 4.22	18.8 $\pm$ 0.904	19.7 $\pm$ 2.55
4	15.8 $\pm$ 3.74	31.4 $\pm$ 7.10	29.2 $\pm$ 3.71	26.7 $\pm$ 4.39	23.6 $\pm$ 6.90
6	15.9 $\pm$ 4.94	28.1 $\pm$ 8.46	29.1 $\pm$ 8.13	21.8 $\pm$ 6.48	20.3 $\pm$ 11.2
9	15.0 $\pm$ 5.90	23.8 $\pm$ 6.99	23.2 $\pm$ 5.70	18.0 $\pm$ 5.18	18.6 $\pm$ 10.55
24	0.26 $\pm$ 0.28	7.43 $\pm$ 9.63	12.02 $\pm$ 9.33	0.476 $\pm$ 0.344	4.2 $\pm$ 5.51
30	0.08 $\pm$ 0.06	7.52 $\pm$ 12.63	14.9 $\pm$ 13.2	0.574 $\pm$ 0.641	3.18 $\pm$ 5.88
48	0.16 $\pm$ 0.30	1.08 $\pm$ 2.31	0.0325 $\pm$	0.082 $\pm$ 0.082	0.02 $\pm$ 0.01

[0159] In the treatment of diabetes, it is advantageous to maintain effective concentrations of drug in the blood over periods of time that permit once a day dosing.

As shown in Table 8 and Figure 23, at 24 hours post-dosing, the plasma level of micronized, free-base MBX-2982 was 0.26  $\mu$ g/ml. In contrast, at 24 hours, the plasma levels of the salt forms were 7.3  $\mu$ g/ml (HCl form I), 12.02  $\mu$ g/ml (HCl form II), 0.476  $\mu$ g/ml (mesylate), and 4.2  $\mu$ g/ml (tosylate). At 30 hours post-dose, the plasma levels of the micronized, free-base MBX-2982 was 0.08  $\mu$ g/ml, and the salt forms were 7.52  $\mu$ g/l

(HCl form I), 14.9  $\mu\text{g}/\text{ml}$  (HCl form II), 0.574  $\mu\text{g}/\text{ml}$  (mesylate), and 3.18  $\mu\text{g}/\text{ml}$  (tosylate). Thus at 30 hours post-dose, the fold difference in the plasma concentrations of the salts of MBX-2982 when compared to the micronized, free-base form of 2982 are 94 fold higher plasma concentration (HCl form I), 186 fold higher plasma concentration 5 (HCl form II), 7 fold higher plasma concentration (mesylate), and 40 fold higher plasma concentration (tosylate).

**[0160]** Modifications to the invention will be apparent to one of skill in the art given this disclosure. Such modifications and the resulting equivalents to the embodiments and examples described above are intended to be included within the scope of the following


10 claims.

## WHAT IS CLAIMED IS:


1. A crystalline salt of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl}-piperidin-1-yl}-pyrimidine selected from the group consisting of besylate, camsylate, esylate, HBr, HCl, mesylate, sulfate, and tosylate.
- 5 2. The salt of claim 1 having substantially the same X-ray powder diffraction (XRPD) pattern as shown in any one of Figures 15-21.
3. The crystalline hydrochloride salt of claim 1.
4. The crystalline hydrochloride salt of claim 3 having substantially the same XRPD pattern as shown in Figure 22.
- 10 5. The crystalline hydrochloride salt of claim 3 that is a polymorph (Form I) having a XRPD pattern comprising peaks at degrees 2-theta diffraction angles of about 8.8, 10.8, 16.1, 17.4, 20.4, 20.9, 21.5, 21.7, 26.6, and 28.1.
6. The polymorph of claim 5 having substantially the same XRPD pattern as shown in Figure 1.
- 15 7. The polymorph of claim 5 having a DSC thermogram comprising an endotherm onset at about 191°C.
8. The polymorph of claim 5 having a DSC thermogram substantially as shown in Figure 2.
9. The polymorph of claim 5 having a Raman spectrum substantially as shown Figure 4.
- 20 10. The crystalline hydrochloride salt of claim 3 that is a polymorph (Form II) having a XRPD pattern comprising peaks at degrees 2-theta diffraction angles of about 7.8, 10.1, 12.5, 18.4, 19.0, 20.8, 23.0, and 23.5.
11. The polymorph of claim 10 having substantially the same XRPD pattern as shown in Figure 5.
- 25 12. The polymorph of claim 10 having a DSC thermogram comprising an endotherm onset at about 150°C.
13. The polymorph of claim 10 having a DSC thermogram substantially as shown in Figure 6.

14. A pharmaceutical composition comprising the crystalline salt of any one of claims 1 to 12.
15. A method for preparing the polymorph (Form I) of claim 5 comprising crystallization of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl]-piperidin-1-yl}-pyrimidine hydrochloride from an acetone solution.
16. A method for preparing the polymorph (Form II) of claim 10 comprising crystallization of 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl]-piperidin-1-yl}-pyrimidine hydrochloride from a methanol solution optionally containing other solvents such as ethyl acetate or acetone.
- 10 17. A method of treating a disease selected from the group consisting of Type I diabetes, Type II diabetes and metabolic syndrome, said method comprising administering to a subject in need of such treatment an effective amount of a salt of any one of claims 1, 4, 5, or 9.
18. The method of claim 17, wherein said disease is Type II diabetes.
- 15 19. A method for one or more of stimulating insulin production, stimulating glucose-dependent insulin secretion, lowering blood glucose, or lowering blood triglyceride levels, said method comprising administering to a subject in need of such treatment an effective amount of a crystalline salt of any one of claims 1, 4, 5, or 10.
20. The method of claims 17 to 19 further comprising administering a therapeutically effective amount of a DPP IV inhibitor.
21. The method of claim 20 wherein the DPP IV inhibitor is selected from the group consisting of sitagliptin, vildagliptin, denagliptin, saxagliptin, and alogliptin.
22. The method of claim 21 wherein said DPP IV inhibitor is sitagliptin or vildagliptin.


1/23

***FIG. 1***


2/23

**FIG. 2**


3/23

**FIG. 3**

4/23


**FIG. 4**

5/23




**FIG. 5**


6/23

**FIG. 6**


7/23

**FIG. 7**


8/23

**FIG. 8**


9/23

**FIG. 9**


10/23

**FIG. 10**


11/23

**FIG. 11**


12/23

**FIG. 12**


13/23

**FIG. 13**


14/23

**FIG. 14**


15/23

**FIG. 15**


16/23

**FIG. 16**


17/23

**FIG. 17**


18/23

**FIG. 18**


19/23

**FIG. 19**


20/23

**FIG. 20**


21/23

**FIG. 21**

22/23

**FIG. 22**

23/23

**FIG. 23**

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2010/049486

**A. CLASSIFICATION OF SUBJECT MATTER**  
INV. C07D513/14 A61K31/506 A61P3/10  
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2008/083238 A2 (METABOLEX INC [US]; CHEN XIN [US]; CHENG PENG [CN]; CLEMENS L EDWARD [ ]) 10 July 2008 (2008-07-10) cited in the application example 52, abstract -----	1-22

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority, claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
11 November 2010	29/11/2010

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Sahagún Krause, H
--	---

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No  
PCT/US2010/049486

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 2008083238	A2	10-07-2008	AU 2007339772 A1	10-07-2008
			CA 2672725 A1	10-07-2008
			EA 200900896 A1	30-04-2010
			EP 2111107 A2	28-10-2009
			JP 2010514795 T	06-05-2010
			KR 20090098898 A	17-09-2009
			US 2010087465 A1	08-04-2010
			US 2010130511 A1	27-05-2010
			US 2009054475 A1	26-02-2009