



(86) Date de dépôt PCT/PCT Filing Date: 2007/02/21
(87) Date publication PCT/PCT Publication Date: 2007/08/30
(45) Date de délivrance/Issue Date: 2018/06/05
(85) Entrée phase nationale/National Entry: 2008/08/19
(86) N° demande PCT/PCT Application No.: EP 2007/051675
(87) N° publication PCT/PCT Publication No.: 2007/096387
(30) Priorités/Priorities: 2006/02/21 (DE10 2006 008 030.0);
2006/09/07 (EP06120309.7)

(51) Cl.Int./Int.Cl. *C12N 15/82* (2006.01),
A01H 5/00 (2018.01), *A01H 5/10* (2018.01),
A01H 6/20 (2018.01), *A23D 9/00* (2006.01),
C12N 15/52 (2006.01), *C12N 15/53* (2006.01),
C12N 5/10 (2006.01)

(72) Inventeurs/Inventors:
CIRPUS, PETRA, DE;
BAUER, JOERG, DE;
WETJEN, TOM, DE;
QIU, XIAO, CA;
TRUKSA, MARTIN, CA;
WU, GUOHAI, CA;
CHENG, BIFANG, CA

(73) Propriétaire/Owner:

(54) Titre : PROCEDE POUR PRODUIRE DES ACIDES GRAS POLYINSATURES
(54) Title: METHOD FOR PRODUCING POLYUNSATURATED FATTY ACIDS

(57) **Abrégé/Abstract:**

The invention relates to a method for producing eicosapentanoic acid, docosapentanoic acid and/or docohexanoic acid in transgenic plants. According to said method, the plant is provided with at least one nucleic acid sequence coding for a polypeptide with a $\Delta 6$ desaturase activity, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 6$ elongase activity, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 5$ desaturase activity, and at least one nucleic acid sequence coding for a polypeptide with a $\Delta 5$ elongase activity, the nucleic acid sequence coding for a polypeptide with a $\Delta 5$ elongase activity being modified in relation to the nucleic acid sequence in the organism from which the sequence originates, such that it is adapted to the codon use in at least one type of plant. For the production of docosahexanoic acid, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 4$ desaturase activity is also introduced into the plant.

(73) Propriétaires(suite)/Owners(continued):BASF PLANT SCIENCE GMBH, DE

(74) Agent: ROBIC

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG(19) Weltorganisation für geistiges Eigentum
Internationales Büro(43) Internationales Veröffentlichungsdatum
30. August 2007 (30.08.2007)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2007/096387 A1(51) Internationale Patentklassifikation:
C12N 15/82 (2006.01)

(21) Internationales Aktenzeichen: PCT/EP2007/051675

(22) Internationales Anmeldedatum:
21. Februar 2007 (21.02.2007)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
10 2006 008 030.0
21. Februar 2006 (21.02.2006) DE
06120309.7 7. September 2006 (07.09.2006) EP

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): BASF PLANT SCIENCE [DE/DE]; BPS - A30, 67056 Ludwigshafen (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): CIRPUS, Petra [DE/DE]; Landteilstrasse 12, 68163 Mannheim (DE). BAUER, Jörg [DE/DE]; Maxburgstrasse 6, 67117 Limburgerhof (DE). QIU, Xiao [CA/CA]; 403 Kendarline Road, Saskatoon, Saskatchewan S7N 3S5 (CA). WU, Guohai [CA/CA]; 2103 Kendarline Road, Saskatoon, Saskatchewan S7N 3S5 (CA). CHENG, Bifang [CA/CA]; Pbi, 110 Gymnasium Road, Saskatoon, Saskatchewan S7N 0W9 (CA). TRUKSA, Martin [CA/CA]; 110 Gymnasium Road, Saskatoon, Saskatchewan S7N 0W9 (CA). WETJEN, Tom [DE/DE]; Meerwiesenstrasse 21, 68163 Mannheim (DE).

(74) Anwalt: NEUEFEIND, Regina; Maiwald Patentanwalts GmbH, Elisenhof, Elisenstrasse 3, 80335 München (DE).

(81) Bestimmungsstaaten (soweit nicht anders angegeben, für jede verfügbare nationale Schutzrechtsart): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Bestimmungsstaaten (soweit nicht anders angegeben, für jede verfügbare regionale Schutzrechtsart): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Veröffentlicht:

- mit internationalem Recherchenbericht
- vor Ablauf der für Änderungen der Ansprüche geltenden Frist; Veröffentlichung wird wiederholt, falls Änderungen eintreffen
- mit dem Sequenzprotokollteil der Beschreibung in elektronischer Form getrennt veröffentlicht; auf Antrag vom Internationalen Büro erhältlich

Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

(54) Title: METHOD FOR PRODUCING POLYUNSATURATED FATTY ACIDS

(54) Bezeichnung: VERFAHREN ZUR HERSTELLUNG VON MEHRFACH UNGESÄTTIGTEN FETTSÄUREN

(57) Abstract: The invention relates to a method for producing eicosapentanoic acid, docosapentanoic acid and/or docohexanoic acid in transgenic plants. According to said method, the plant is provided with at least one nucleic acid sequence coding for a polypeptide with a $\Delta 6$ desaturase activity, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 6$ elongase activity, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 5$ desaturase activity, and at least one nucleic acid sequence coding for a polypeptide with a $\Delta 5$ elongase activity, the nucleic acid sequence coding for a polypeptide with a $\Delta 5$ elongase activity being modified in relation to the nucleic acid sequence in the organism from which the sequence originates, such that it is adapted to the codon use in at least one type of plant. For the production of docosahexanoic acid, at least one nucleic acid sequence coding for a polypeptide with a $\Delta 4$ desaturase activity is also introduced into the plant.(57) Zusammenfassung: Die vorliegende Erfindung betrifft ein Verfahren zur Herstellung von Eicosapentaensäure, Docosapentaensäure und/oder Docosahexaensäure in transgenen Pflanzen, indem in der Pflanze bereitgestellt werden mindestens eine Nukleinsäuresequenz, welche für ein Polypeptid mit einer $\Delta 6$ -Desaturase-Aktivität kodiert; mindestens eine Nukleinsäuresequenz, welche für ein Polypeptid mit einer $\Delta 6$ -Elongase-Aktivität kodiert; mindestens eine Nukleinsäuresequenz, welche für ein Polypeptid mit einer $\Delta 5$ -Desaturase-Aktivität kodiert; und mindestens eine Nukleinsäuresequenz, welche für ein Polypeptid mit einer $\Delta 5$ -Elongase-Aktivität kodiert, wobei die Nukleinsäuresequenz, welche für ein Polypeptid mit einer $\Delta 5$ -Elongase-Aktivität kodiert, gegenüber der Nukleinsäuresequenz in dem Organismus, aus dem die Sequenz stammt, dadurch verändert ist, dass sie an die Kodonverwendung in einer oder mehreren Pflanzenarten angepasst ist. Für die Herstellung von Docosahexaensäure wird zusätzlich eine oder mehrere Nukleinsäuresequenzen, die für ein Polypeptid mit einer $\Delta 4$ -Desaturase-Aktivität kodiert, in die Pflanze eingebracht.

WO 2007/096387 A1

METHOD FOR PRODUCING POLYUNSATURATED FATTY ACIDS

The present invention relates to a process for the production of eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in transgenic plants, providing in the plant at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity; at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity; at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity; and at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity,

where the nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity is modified by comparison with the nucleic acid sequence in the organism from which the sequence is derived in that it is adapted to the codon usage in one or more plant species.

In a preferred embodiment there is additionally provision of further nucleic acid sequences which code for a polypeptide having the activity of an $\omega 3$ -desaturase and/or of a $\Delta 4$ -desaturase in the plant.

The present invention relates to a process for producing eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in a transgenic plant, comprising the provision in the plant of

- at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 2;
- at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 172;
- at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 52; and

1a

- at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 65 and
- insofar as the process aims at producing docosahexaenoic acid (DHA), at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 4$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 78,

where the nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity is modified by comparison with the nucleic acid sequence in the organism from which the sequence is obtained in that it is adapted to the codon usage in the host plant species.

The present invention relates to an isolated nucleic acid molecule comprising a nucleic acid sequence as shown in SEQ ID No. 64.

The present invention relates to a recombinant nucleic acid molecule comprising:

- a) one or more copies of at least one promoter which is active in plant cells,
- b) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 2,
- c) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 52,
- d) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 172,
- e) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity and having an amino acid identity of at least 90% to

1b

SEQ ID NO: 65 and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence is obtained by being adapted to the codon usage in one or more plant species, and

f) one or more copies of at least one terminator sequence.

5 The present invention relates to a transgenic plant cell, wherein the plant cell is obtained from oilseed rape, soybean or flax, comprising the recombinant nucleic acid molecule as defined herein.

The present invention relates to a seed cell comprising the recombinant nucleic acid molecule as defined herein.

10 The present invention relates to a process for producing a transgenic plant which produces eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA) and/or docosahexaenoic acid (DHA), comprising introducing into a plant the recombinant nucleic acid molecule as defined herein and expressing the same therein.

15 The present invention relates to a use of the transgenic plant cell as defined herein, the transgenic plant as defined herein, or the transgenic seed cell as defined herein, for the production of an oil, a lipid, a fatty acid and/or a fatty acid composition.

20 The present invention relates to the use of the transgenic plant cell as defined herein, the transgenic plant as produced by the process of the present invention, or the transgenic seed cell as defined herein, for the production of an oil, a lipid, a fatty acid and/or a fatty acid composition.

The present invention relates to oil, lipid and/or fatty acid produced by a transgenic *Brassica* plant, wherein said oil, lipid and/or fatty acid comprise 60 to 85% by weight of polyunsaturated fatty acids based on the total fatty acids in the transgenic plant, wherein said polyunsaturated fatty acids comprise at least 20% by weight of eicosapentaenoic acid (EPA)
25 based on the total fatty acids in the transgenic plant in form of triacylglycerides.

The present invention relates to oil, lipid and/or fatty acid produced by a transgenic *Brassica* plant, wherein said oil, lipid and/or fatty acid comprise 60 to 85% by weight of

1c

polyunsaturated fatty acids based on the total fatty acids in the transgenic plant, wherein said polyunsaturated fatty acids comprise at least 2% by weight of docosapentaenoic acid (DPA) based on the total fatty acids in the transgenic plant in form of triacylglycerides.

5 The present invention relates to oil, lipid and/or fatty acid produced by a transgenic *Brassica* plant, wherein said oil, lipid and/or fatty acid comprise 60 to 85% by weight of polyunsaturated fatty acids based on the total fatty acids in the transgenic plant, wherein said polyunsaturated fatty acids comprise at least 4% by weight of docosahexaenoic acid (DHA) based on the total fatty acids in the transgenic plant in form of triacylglycerides.

10 The present invention relates to oil, lipid and/or fatty acid produced by a transgenic *Brassica* plant, wherein said oil, lipid and/or fatty acid comprise a total amount of at least 54% by weight of polyunsaturated ω 3-fatty acids based on the total fatty acids in the transgenic plant.

15 The present invention relates to a method for producing oil, lipid and/or fatty acid of the invention, comprising expressing in a *Brassica* plant a nucleic acid encoding a Δ 6-desaturase, a nucleic acid encoding a Δ 5-desaturase, a nucleic acid encoding a Δ 6-elongase, a nucleic acid encoding a ω 3-desaturase, a nucleic acid encoding a Δ 5-elongase, and a nucleic acid encoding a Δ 4-desaturase, wherein said nucleic acid encoding a Δ 5-elongase is codon-optimized by adapting to the codon usage of *Brassica*.

20 In a further preferred embodiment there is provision of further nucleic acid sequences which code for acyl-CoA dehydrogenase(s), acyl-ACP (acyl carrier protein) desaturase(s), acyl-ACP thioesterase(s), fatty acid acyl transferase(s), acyl-CoA; lysophospholipid acyl transferase(s), fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allene oxide synthases, hydroperoxide lyases or fatty
25 acid elongase(s) in the plant.

The invention furthermore relates to recombinant nucleic acid molecules comprising at least one nucleic acid sequence which codes for a polypeptide having a Δ 6-desaturase activity; at least one nucleic acid sequence which codes for a polypeptide having a Δ 5-desaturase activity; at least one nucleic acid sequence which codes for a polypeptide
30 having a Δ 6-elongase

WO 2007/096387

PCT/EP2007/051675

- 2 -

activity; and at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence originates in that it is adapted to the codon usage in one or more plant species.

- 5 A further part of the invention relates to oils, lipids and/or fatty acids which have been produced by the process according to the invention, and to their use.

Finally, the invention also relates to transgenic plants which have been produced by the process of the invention or which comprise a recombinant nucleic acid molecule of the invention, and to the use thereof as foodstuffs or feedstuffs.

- 10 Lipid synthesis can be divided into two sections: the synthesis of fatty acids and their binding to sn-glycerol-3-phosphate, and the addition or modification of a polar head group. Usual lipids which are used in membranes comprise phospholipids, glycolipids, sphingolipids and phosphoglycerides. Fatty acid synthesis starts with the conversion of acetyl-CoA into malonyl-CoA by acetyl-CoA carboxylase or into acetyl-ACP by acetyl transacylase. After
15 condensation reaction, these two product molecules together form acetoacetyl-ACP, which is converted via a series of condensation, reduction and dehydration reactions so that a saturated fatty acid molecule with the desired chain length is obtained. The production of the unsaturated fatty acids from these molecules is catalyzed by specific desaturases, either aerobically by means of molecular oxygen or anaerobically (regarding the fatty acid synthesis
20 in microorganisms, see F.C. Neidhardt et al. (1996) *E. coli* and *Salmonella*. ASM Press: Washington, D.C., p. 612-636 and references cited therein; Lengeler et al. (Ed.) (1999) *Biology of Prokaryotes*. Thieme: Stuttgart, New York, and the references therein, and Magnuson, K., et al. (1993) *Microbiological Reviews* 57:522-542 and the references therein). To undergo the further elongation steps, the resulting phospholipid-bound fatty acids must be
25 returned to the fatty acid CoA ester pool. This is made possibly by acyl-CoA:lysophospholipid acyltransferases. Moreover, these enzymes are capable of transferring the elongated fatty

WO 2007/096387

PCT/EP2007/051675

- 3 -

acids from the CoA esters back to the phospholipids. If appropriate, this reaction sequence can be followed repeatedly.

Furthermore, fatty acids must subsequently be transported to various modification sites and incorporated into the triacylglycerol storage lipid. A further important step during lipid
5 synthesis is the transfer of fatty acids to the polar head groups, for example by glycerol fatty acid acyltransferase (see Frentzen, 1998, Lipid, 100(4-5):161-166).

An overview of the biosynthesis of fatty acids in plants, desaturation, the lipid metabolism and the membrane transport of lipidic compounds, beta-oxidation, the modification of fatty acids, cofactors and the storage and assembly of triacylglycerol, including the references is
10 given by the following papers: Kinney (1997) Genetic Engineering, Ed.: JK Setlow, 19:149-166; Ohlrogge and Browse (1995) Plant Cell 7:957-970; Shanklin and Cahoon (1998) Annu. Rev. Plant Physiol. Plant Mol. Biol. 49:611-641; Voelker (1996) Genetic Engineering, Ed.: JK Setlow, 18:111-13; Gerhardt (1992) Prog. Lipid R. 31:397-417; Gühnemann-Schäfer & Kindl (1995) Biochim. Biophys Acta 1256:181-186; Kunau et al. (1995) Prog. Lipid Res.
15 34:267-342; Stymne et al. (1993) in: Biochemistry and Molecular Biology of Membrane and Storage Lipids of Plants, Ed.: Murata und Somerville, Rockville, American Society of Plant Physiologists, 150-158; Murphy & Ross (1998) Plant Journal. 13(1):1-16.

Depending on the desaturation pattern, two large classes of polyunsaturated fatty acids, the $\omega 6$ and the $\omega 3$ fatty acids, which differ with regard to their metabolism and their function, can be
20 distinguished.

In the text which follows, polyunsaturated fatty acids are referred to as PUFA, PUFAs, LCPUFA or LCPUFAs (poly unsaturated fatty acids, PUFA, long chain poly unsaturated fatty acids, LCPUFA).

The fatty acid linoleic acid ($18:2^{\Delta 9,12}$) acts as starting material for the $\omega 6$ metabolic pathway,
25 while the $\omega 3$ pathway proceeds via linolenic acid ($18:3^{\Delta 9,12,15}$). Linolenic acid is formed from

WO 2007/096387

PCT/EP2007/051675

- 4 -

linoleic acid by the activity of an ω 3-desaturase (Tocher et al. (1998) Prog. Lipid Res. 37: 73-117; Domergue et al. (2002) Eur. J. Biochem. 269: 4105-4113).

Mammals, and thus also humans, have no corresponding desaturase activity (Δ 12- and ω 3-desaturase) for the formation of the starting materials and must therefore take up these fatty acids (essential fatty acids) via the food. Starting with these precursors, the physiologically important polyunsaturated fatty acids arachidonic acid (= ARA, 20:4 ^{Δ 5,8,11,14}), an ω 6-fatty acid and the two ω 3-fatty acids eicosapentaenoic acid (= EPA, 20:5 ^{Δ 5,8,11,14,17}) and docosahexaenoic acid (DHA, 22:6 ^{Δ 4,7,10,13,17,19}) are synthesized via a sequence of desaturase and elongase reactions.

10 The elongation of fatty acids, by elongases, by 2 or 4 C atoms is of crucial importance for the production of C₂₀- and C₂₂-PUFAs, respectively. This process proceeds via 4 steps. The first step is the condensation of malonyl-CoA onto the fatty acid acyl-CoA by ketoacyl-CoA synthase (KCS, hereinbelow referred to as elongase). This is followed by a reduction step (ketoacyl-CoA reductase, KCR), a dehydration step (dehydratase) and a final reduction step
15 (enoyl-CoA reductase). It has been postulated that the elongase activity affects the specificity and rate of the entire process (Millar and Kunst (1997) Plant Journal 12:121-131).

Fatty acids and triacylglycerides have a multiplicity of applications in the food industry, in animal nutrition, in cosmetics and the pharmacological sector. Depending on whether they are free saturated or unsaturated fatty acids or else triacylglycerides with an elevated content of
20 saturated or unsaturated fatty acids, they are suitable for very different applications. Thus, for example, lipids with unsaturated, specifically with polyunsaturated fatty acids, are preferred in human nutrition. The polyunsaturated ω 3-fatty acids are supposed to have a positive effect on the cholesterol level in the blood and thus on the prevention of heart disease. The risk of heart disease, strokes or hypertension can be reduced markedly by adding these ω 3-fatty acids to
25 the food (Shimikawa (2001) World Rev. Nutr. Diet. 88: 100-108).

ω 3-fatty acids also have a positive effect on inflammatory, specifically on chronically inflammatory, processes in association with immunological diseases such as rheumatoid

WO 2007/096387

PCT/EP2007/051675

- 5 -

arthritis (Calder (2002) Proc. Nutr. Soc. 61: 345-358; Cleland and James (2000) J. Rheumatol. 27: 2305-2307). They are therefore added to foodstuffs, specifically to dietetic foodstuffs, or are employed in medicaments. ω 6-fatty acids such as arachidonic acid tend to have a negative effect in connection with these rheumatological diseases.

- 5 ω 3- and ω 6-fatty acids are precursors of tissue hormones, known as eicosanoids, such as the prostaglandins, which are derived from dihomog- γ -linolenic acid, arachidonic acid and eicosapentaenoic acid, and of the thromboxanes and leukotrienes, which are derived from arachidonic acid and eicosapentaenoic acid. Eicosanoids (known as the PG₂ series) which are formed from the ω 6-fatty acids, generally promote inflammatory reactions, while eicosanoids
10 (known as the PG₃ series) from ω 3-fatty acids have little or no proinflammatory effect.

Polyunsaturated long-chain ω 3-fatty acids such as eicosapentaenoic acid (= EPA, C20:5 ^{Δ 5,8,11,14,17}) or docosahexaenoic acid (= DHA, C22:6 ^{Δ 4,7,10,13,16,19}) are important components of human nutrition owing to their various roles in health aspects, including the development of the child brain, the functionality of the eyes, the synthesis of hormones and
15 other signal substances, and the prevention of cardiovascular disorders, cancer and diabetes (Poulos, A (1995) Lipids 30:1-14; Horrocks, LA and Yeo YK (1999) Pharmacol Res 40:211-225).

Owing to the present-day composition of human food, an addition of polyunsaturated ω 3-fatty acids, which are preferentially found in fish oils, to the food is particularly important. Thus,
20 for example, polyunsaturated fatty acids such as docosahexaenoic acid (= DHA, C22:6 ^{Δ 4,7,10,13,16,19}) or eicosapentaenoic acid (= EPA, C20:5 ^{Δ 5,8,11,14,17}) are added to infant formula to improve the nutritional value. There is therefore a demand for the production of polyunsaturated long-chain fatty acids.

The various fatty acids and triglycerides are mainly obtained from microorganisms such as
25 *Mortierella* or *Schizochytrium* or from oil-producing plants such as soybeans, oilseed rape, and algae such as *Cryptocodinium* or *Phaeodactylum* and others, being obtained, as a rule, in the form of their triacylglycerides (= triglycerides = triglycerols). However, they can also be

WO 2007/096387

PCT/EP2007/051675

- 6 -

obtained from animals, for example, fish. The free fatty acids are advantageously prepared by hydrolyzing the triacylglycerides. Very long-chain polyunsaturated fatty acids such as DHA, EPA, arachidonic acid (ARA, C20:4^{Δ5,8,11,14}), dihomo-γ-linolenic acid (DHGL, C20:3^{Δ8,11,14}) or docosapentaenoic acid (DPA, C22:5^{Δ7,10,13,16,19}) are, however, not synthesized in oil crops
 5 such as oilseed rape, soybeans, sunflowers and safflower. Conventional natural sources of these fatty acids are fish such as herring, salmon, sardine, redfish, eel, carp, trout, halibut, mackerel, zander or tuna, or algae.

Owing to the positive characteristics of the polyunsaturated fatty acids, there has been no lack of attempts in the past to make available genes which are involved in the synthesis of these
 10 fatty acids or triglycerides for the production of oils in various organisms with a modified content of unsaturated fatty acids. Thus, WO 91/13972 and its US equivalent describe a Δ9-desaturase. WO 93/11245 claims a Δ15-desaturase and WO 94/11516 a Δ12-desaturase. Further desaturases are described, for example, in EP-A-0 550 162, WO 94/18337, WO 97/30582, WO 97/21340, WO 95/18222, EP-A-0 794 250, Stukeley et al. (1990) J. Biol.
 15 Chem., 265: 20144-20149, Wada et al. (1990) Nature 347: 200-203 or Huang et al. (1999) Lipids 34: 649-659. However, the biochemical characterization of the various desaturases has been insufficient to date since the enzymes, being membrane-bound proteins, present great difficulty in their isolation and characterization (McKeon et al. (1981) Methods in Enzymol. 71: 12141-12147, Wang et al. (1988) Plant Physiol. Biochem., 26: 777-792).

20 As a rule, membrane-bound desaturases are characterized by being introduced into a suitable organism which is subsequently analyzed for enzyme activity by analyzing the starting materials and the products. Δ6-Desaturases are described in WO 93/06712, US 5,614,393, WO 96/21022, WO 00/21557 and WO 99/27111. The application of this enzyme for the production of fatty acids in transgenic organisms is described in WO 98/46763, WO 98/46764
 25 and WO 98/46765. The expression of various desaturases and the formation of polyunsaturated fatty acids is also described and claimed in WO 99/64616 or WO 98/46776. As regards the expression efficacy of desaturases and its effect on the formation of polyunsaturated fatty acids, it must be noted that the expression of a single desaturase as

WO 2007/096387

PCT/EP2007/051675

- 7 -

described to date has only resulted in low contents of unsaturated fatty acids/lipids such as, for example, γ -linolenic acid and stearidonic acid.

There have been a number of attempts in the past to obtain elongase genes. Millar and Kunst (1997) Plant Journal 12:121-131 and Millar et al. (1999) Plant Cell 11:825-838 describe the
5 characterization of plant elongases for the synthesis of monounsaturated long-chain fatty acids (C22:1) and for the synthesis of very long-chain fatty acids for the formation of waxes in plants (C₂₈-C₃₂). The synthesis of arachidonic acid and EPA is described, for example, in WO 01/59128, WO 00/12720, WO 02/077213 and WO 02/08401. The synthesis of
polyunsaturated C24-fatty acids is described, for example, in Tvrdik et al. (2000) J. Cell Biol.
10 149:707-718 or in WO 02/44320.

Especially suitable microorganisms for the production of PUFAs are microalgae such as Phaeodactylum tricornutum, Porphiridium species, Thraustochytrium species, Schizochytrium species or Cryptocodinium species, ciliates such as Stylonychia or Colpidium, fungi such as Mortierella, Entomophthora or Mucor and/or mosses such as Physcomitrella, Ceratodon and
15 Marchantia (R. Vazhappilly & F. Chen (1998) Botanica Marina 41: 553-558; K. Totani & K. Oba (1987) Lipids 22: 1060-1062; M. Akimoto et al. (1998) Appl. Biochemistry and Biotechnology 73: 269-278). Strain selection has resulted in the development of a number of mutant strains of the microorganisms in question which produce a series of desirable compounds including PUFAs. However, the mutation and selection of strains with an
20 improved production of a particular molecule such as the polyunsaturated fatty acids is a time-consuming and difficult process. Moreover, only limited amounts of the desired polyunsaturated fatty acids such as DPA, EPA or ARA can be produced with the aid of the abovementioned microorganisms; in addition, they are generally obtained as fatty acid mixtures. This is why recombinant methods are preferred whenever possible.

25 Higher plants comprise polyunsaturated fatty acids such as linoleic acid (C18:2) and linolenic acid (C18:3). ARA, EPA and DHA are found not at all in the seed oil of higher plants, or only in miniscule amounts (E. Ucciani: Nouveau Dictionnaire des Huiles Végétales [New

WO 2007/096387

PCT/EP2007/051675

- 8 -

Dictionary of the Vegetable Oils]. Technique & Documentation – Lavoisier, 1995. ISBN: 2-7430-0009-0). However, the production of LCPUFAs in higher plants, preferably in oil crops such as oilseed rape, linseed, sunflowers and soybeans, would be advantageous since large amounts of high-quality LCPUFAs for the food industry, animal nutrition and pharmaceutical purposes might be obtained economically. To this end, it is advantageous to introduce, into oilseeds, genes which encode enzymes of the LCPUFA biosynthesis via recombinant methods and to express them therein. These genes encode for example $\Delta 6$ -desaturases, $\Delta 6$ -elongases, $\Delta 5$ -desaturases or $\Delta 4$ -desaturases. These genes can advantageously be isolated from microorganisms and lower plants which produce LCPUFAs and incorporate them in the membranes or triacylglycerides. Thus, it has already been possible to isolate $\Delta 6$ -desaturase genes from the moss *Physcomitrella patens* and $\Delta 6$ -elongase genes from *P. patens* and from the nematode *C. elegans*.

Transgenic plants which comprise and express genes encoding LCPUFA biosynthesis enzymes and which, as a consequence, produce LCPUFAs have been described, for example, in DE-A-102 19 203 (process for the production of polyunsaturated fatty acids in plants). However, these plants produce LCPUFAs in amounts which require further optimization for processing the oils which are present in the plants. Thus, the ARA content in the plants described in DE-A-102 19 203 is only 0.4 to 2% and the EPA content only 0.5 to 1%, in each case based on the total lipid content of the plant.

To make possible the fortification of food and of feed with polyunsaturated, long-chain fatty acids, there is therefore a great need for a simple, inexpensive process for the production of polyunsaturated, long-chain fatty acids, specifically in plant systems.

One object of the invention is therefore to provide a process with which long-chain polyunsaturated fatty acids, especially eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid can be produced in large quantities and inexpensively in transgenic plants.

WO 2007/096387

PCT/EP2007/051675

- 9 -

It has now surprisingly been found that the yield of long-chain polyunsaturated fatty acids, especially eicosapentaenoic, docosapentaenoic acid and/or docosahexaenoic acid, can be increased by expressing an optimized $\Delta 5$ -elongase sequence in transgenic plants.

5 The PUFAs produced by the process of the invention comprise a group of molecules which higher animals are no longer able to synthesize and thus must consume, or which higher animals are no longer able to produce themselves in sufficient amounts and thus must consume additional amounts thereof, although they can easily be synthesized by other organisms such as bacteria.

Accordingly, the object of the invention is achieved by the process of the invention for
10 producing eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in a transgenic plant, comprising the provision in the plant of at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity; at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity; at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity; and at least one
15 nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity, where the nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity is modified by comparison with the nucleic acid sequence in the organism from which the sequence is derived in that it is adapted to the codon usage in one or more plant species. To produce DHA it is additionally necessary to provide at least one nucleic acid sequence which
20 codes for a polypeptide having a $\Delta 4$ -desaturase activity in the plant.

The “provision in the plant” means in the context of the present invention that measures are taken so that the nucleic acid sequences coding for a polypeptide having a $\Delta 6$ -desaturase activity, a polypeptide having a $\Delta 6$ -elongase activity, a polypeptide having a $\Delta 5$ -desaturase
25 activity and a polypeptide having a $\Delta 5$ -elongase activity are present together in one plant. The “provision in the plant” thus comprises the introduction of the nucleic acid sequences into the plant both by transformation of a plant with one or more recombinant nucleic acid molecules

WO 2007/096387

PCT/EP2007/051675

- 10 -

which comprise said nucleic acid sequences, and by crossing suitable parent plants which comprise one or more of said nucleic acid sequences.

The nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity is modified according to the invention by comparison with the nucleic acid sequence in the
5 organism from which the sequence originates in that it is adapted to the codon usage in one or more plant species. This means that the nucleic acid sequence has been specifically optimized for the purpose of the invention without the amino acid sequence encoded by the nucleic acid sequence having been altered thereby.

The genetic code is redundant because it uses 61 codons in order to specify 20 amino acids.

10 Therefore, most of the 20 proteinogenic amino acids are therefore encoded by a plurality of triplets (codons). The synonymous codons which specify an individual amino acid are, however, not used with the same frequency in a particular organism; on the contrary there are preferred codons which are frequently used, and codons which are used more rarely. These differences in codon usage are attributed to selective evolutionary pressures and especially the
15 efficiency of translation. One reason for the lower translation efficiency of rarely occurring codons might be that the corresponding aminoacyl-tRNA pools are exhausted and thus no longer available for protein synthesis.

In addition, different organisms prefer different codons. For this reason, for example, the expression of a recombinant DNA derived from a mammalian cell frequently proceeds only
20 suboptimally in *E. coli* cells. It is therefore possible in some cases to increase expression by replacing rarely used codons with frequently used codons. Without wishing to be bound to one theory, it is assumed that the codon-optimized DNA sequences make more efficient translation possible, and the mRNAs formed therefrom possibly have a greater half-life in the cell and therefore are available more frequently for translation. From what has been said
25 above, it follows that codon optimization is necessary only if the organism in which the nucleic acid sequence is to be expressed differs from the organism from which the nucleic acid sequence is originally derived.

WO 2007/096387

PCT/EP2007/051675

- 11 -

For many organisms of which the DNA sequence of a relatively large number of genes is known there are tables from which the frequency of use of particular codons in the respective organism can be taken. It is possible with the aid of these tables to translate protein sequences with relatively high accuracy back into a DNA sequence which comprises the codons preferred in the respective organism for the various amino acids of the protein. Tables on codon usage can be found inter alia at the following Internet address:

<http://www.kazusa.or.jp/Kodon/E.html>. In addition, several companies provide software for gene optimization, such as, for example, Entelechon (Software Leto) or Geneart (Software GeneOptimizer).

Adaptation of the sequences to the codon usage in a particular organism can take place with the aid of various criteria. On the one hand, it is possible to use for a particular amino acid always the codon which occurs most frequently in the selected organism but, on the other hand, the natural frequency of the various codons can also be taken into account, so that all the codons for a particular amino acid are incorporated into the optimized sequence according to their natural frequency. Selection of the position at which a particular base triplet is used can take place at random in this case. The DNA sequence was adapted according to the invention taking account of the natural frequency of individual codons, it also being suitable to use the codons occurring most frequently in the selected organism.

It is particularly preferred for a nucleic acid sequence from *Ostreococcus tauri* which codes for a polypeptide having a $\Delta 5$ -elongase activity, such as, for example, the polypeptide depicted in SEQ ID No. 110, to be adapted at least to the codon usage in oilseed rape, soybean and/or flax. The nucleic acid sequence originally derived from *Ostreococcus tauri* is preferably the sequence depicted in SEQ ID No. 109. The DNA sequence coding for the $\Delta 5$ -elongase is adapted in at least 20% of the positions, preferably in at least 30% of the positions, particularly preferably in at least 40% of the positions and most preferably in at least 50% of the positions to the codon usage in oilseed rape, soybean and/or flax.

The nucleic acid sequence used is most preferably the sequence indicated in SEQ ID No. 64.

WO 2007/096387

PCT/EP2007/051675

- 12 -

It will be appreciated that the invention also encompasses those codon-optimized DNA sequences which code for a polypeptide having the activity of a $\Delta 5$ -elongase and whose amino acid sequence is modified in one or more positions by comparison with the wild-type sequence but which still has substantially the same activity as the wild-type protein.

- 5 The nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity is preferably selected from the group consisting of:
 - a) nucleic acid sequences having the sequence depicted in SEQ ID No. 1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33,35,37,39 or 41, preferably having the sequence depicted in SEQ ID No. 1,
 - 10 b) nucleic acid sequences which code for the amino acid sequence indicated in SEQ ID No. 2,4,6,8,10,12,14,16,18,20,22,24,26,28,30,32,34,36,38,40 or 42, preferably in SEQ ID No. 2,
 - c) nucleic acid sequences which hybridize with the complementary strand of the nucleic acid sequences indicated a) or b) above, in particular of the nucleic acid sequence indicated in SEQ ID No. 1, under stringent conditions,
 - 15 d) nucleic acid sequences which are at least 60%, 65%, 70%, 75% or 80%, preferably at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90%, particularly preferably at least 91%, 92%, 93%, 94% or 95% and especially at least 96%, 97%, 98% or 99%, identical to the nucleic acid sequences indicated in a) or b) above, especially to the sequence indicated in SEQ ID No. 1, and
 - 20 e) nucleic acid sequences which code for an amino acid sequence and which have at least one, for example 2, 3, 4, 5, 6, 7 or 8, preferably all of the amino acid pattern indicated in SEQ ID No. 43,44,45,46,47,48,49 or 50.

Amino acid pattern means short amino acid sequences which preferably comprise less than 50, particularly preferably less than 40 and especially from 10 to 40 and even more preferably
 25 from 10 to 30 amino acids.

WO 2007/096387

PCT/EP2007/051675

- 13 -

For the present invention, the identity is ascertained preferably over the full length of the nucleotide or amino acid sequences of the invention, for example for the nucleic acid sequence indicated in SEQ ID NO: 64 over the full length of 903 nucleotides.

The nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity is preferably selected from the group consisting of:

a) nucleic acid sequences having the sequence depicted in SEQ ID No. 171, 173, 175, 177, 179, 181 or 183, especially having the sequence depicted in SEQ ID No. 171,

b) nucleic acid sequences which code for the amino acid sequence indicated in SEQ ID No. 172, 174, 176, 178, 180, 182 or 184, especially for the amino acid sequence indicated in SEQ

ID No. 172,

c) nucleic acid sequences which hybridize with the complementary strand of the nucleic acid sequences indicated a) or b) above, especially of the nucleic acid sequence indicated in SEQ ID No. 1, under stringent conditions,

d) nucleic acid sequences which are at least 60%, 65%, 70%, 75% or 80%, preferably at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90%, particularly preferably at least 91%, 92%, 93%, 94% or 95% and especially at least 96%, 97%, 98% or 99%, identical to the nucleic acid sequences indicated in a) or b) above, especially to the sequence indicated in SEQ ID No. 171, and

e) nucleic acid sequences which code for an amino acid sequence and which have at least one, for example 2, 3, 4, 5, 6, 7 or 8, preferably all of the amino acid pattern indicated in SEQ ID No. 185, 186, 187, 188, 189, 190, 191 or 192.

The nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity is in particular likewise a codon-optimized sequence according to the present invention, preferably the nucleic acid sequence depicted in SEQ ID NO: 122.

The nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity is preferably selected from the group consisting of:

a) nucleic acid sequences having the sequence depicted in SEQ ID No. 51, 53 or 55,

WO 2007/096387

PCT/EP2007/051675

- 14 -

preferably having the sequence depicted in SEQ ID No. 51,

b) nucleic acid sequences which code for the amino acid sequence indicated in SEQ ID No. 52, 54 or 56, preferably for the amino acid sequence indicated in SEQ ID No. 52,

c) nucleic acid sequences which hybridize with the complementary strand of the nucleic acid sequences indicated in a) or b) above, especially of the nucleic acid sequence indicated in SEQ ID No. 51, under stringent conditions,

d) nucleic acid sequences which are at least 60%, 65%, 70%, 75% or 80%, preferably at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90%, particularly preferably at least 91%, 92%, 93%, 94% or 95% and especially at least 96%, 97%, 98% or 99%, identical to the nucleic acid sequences indicated in a) or b) above, especially to the nucleic acid indicated under SEQ ID No. 51, and

e) nucleic acid sequences which code for an amino acid sequence which have at least one, for example 2, 3, 4, 5, 6 or 7, preferably all of the amino acid pattern indicated in SEQ ID No. 57,58,59,60,61,62 or 63.

Further suitable nucleic acid sequences can be found by the skilled worker from the literature or the well-known gene libraries such as, for example, <http://www.ncbi.nlm.nih.gov>.

In a further preferred embodiment of the process, additionally one or more nucleic acid sequences which code for a polypeptide having the activity of an ω -3-desaturase and/or of a Δ 4-desaturase are introduced into the plant.

The nucleic acid sequence which codes for a polypeptide having an ω -3-desaturase activity is preferably selected from the group consisting of:

a) nucleic acid sequences having the sequence depicted in SEQ ID No. 193 or 195, preferably the sequence depicted in SEQ ID No. 193,

b) nucleic acid sequences which code for the amino acid sequence indicated in SEQ ID No. 194,

c) nucleic acid sequences which hybridize with the complementary strand of the nucleic acid sequence indicated in SEQ ID No. 193 or 195 under stringent conditions, and

WO 2007/096387

PCT/EP2007/051675

- 15 -

d) nucleic acid sequences which are at least 60%, 65%, 70%, 75% or 80%, preferably at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90%, particularly preferably at least 91%, 92%, 93%, 94% or 95%, and especially at least 96%, 97%, 98% or 99%, identical to the sequence indicated in SEQ ID No. 193 or 195.

- 5 The ω -3-desaturase advantageously used in the process of the invention makes it possible to shift from the ω -6 biosynthetic pathway to the ω -3 biosynthetic pathway, leading to a shift from C_{18:2} to C_{18:3} fatty acids. It is further advantageous for the ω -3-desaturase to convert a wide range of phospholipids such as phosphatidylcholine (= PC), phosphatidylinositol (= PIS) or phosphatidylethanolamine (= PE). Finally, desaturation products can also be found in the
- 10 neutral lipids (= NL), that is to say in the triglycerides.

The nucleic acid sequence which codes for a polypeptide having a Δ 4-desaturase activity is preferably selected from the group consisting of:

- a) nucleic acid sequences having the sequence depicted in SEQ ID No. 77,79,81,83,85,87,89,91 or 93, preferably having the sequence depicted in SEQ ID No. 77,
- 15 b) nucleic acid sequences which code for the amino acid sequence indicated in SEQ ID No. 78,80,82,84,86,88,90,92 or 94, preferably for the amino acid sequence indicated in SEQ ID No. 78,
- c) nucleic acid sequences which hybridize with the complementary strand of the nucleic acid sequences indicated in a) or b) above, especially of the nucleic acid sequence indicated in
- 20 SEQ ID No. 77, under stringent conditions,
- d) nucleic acid sequences which are at least 60%, 65%, 70%, 75% or 80%, preferably at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90%, particularly preferably at least 91%, 92%, 93%, 94% or 95% and especially at least 96%, 97%, 98% or 99%, identical to the sequence indicated in SEQ ID No. 77, and
- 25 e) nucleic acid sequences which code for an amino acid sequence which have at least one, for example 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14, preferably all of the amino acid pattern indicated in SEQ ID No. 95,96,97,98, 99,100,101,102,103,104,105,106,107 or 108.

WO 2007/096387

PCT/EP2007/051675

- 16 -

The $\Delta 4$ -desaturase which is advantageously used in the process of the invention catalyzes the introduction of a double bond into the fatty acid docosapentaenoic acid, leading to formation of docosaheptaenoic acid.

It is advantageous for the described process of the invention additionally to introduce further nucleic acids which code for enzymes of fatty acid or lipid metabolism into the plants in addition to the nucleic acid sequences which code for polypeptides having a $\Delta 6$ -desaturase activity, a $\Delta 6$ -elongase activity, a $\Delta 5$ -desaturase activity and a $\Delta 5$ -elongase activity, and to the nucleic acid sequences which are introduced if appropriate and which code for a polypeptide having an ω -3-desaturase activity and/or a $\Delta 4$ -desaturase activity.

It is possible in principle to use all genes of fatty acid or lipid metabolism in combination with the nucleic acid sequences used in the process of the invention; genes of fatty acid or lipid metabolism selected from the group of acyl-CoA dehydrogenase(s), acyl-ACP (acyl carrier protein) desaturase(s), acyl-ACP thioesterase(s), fatty acid acyltransferase(s), acyl-CoA:lysophospholipid acyltransferases, fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allene oxide synthases, hydroperoxide lyases or fatty acid elongase(s) are preferably used in combination with the $\Delta 6$ -elongase, $\Delta 6$ -desaturase, $\Delta 5$ -desaturase and the $\Delta 5$ -elongase, and if appropriate the ω -3-desaturase and/or the $\Delta 4$ -desaturase, it being possible to use individual genes or a plurality of genes in combination.

The nucleic acids used in the process of the invention are advantageously expressed in vegetative tissues (somatic tissue). Vegetative tissue means in the context of this invention a tissue which is propagated through mitotic divisions. Tissue of this type also arises through asexual reproduction (apomixis) and propagation. Propagation is the term used when the number of individuals increases in consecutive generations. These individuals arising through asexual propagation are very substantially identical to their parents. Examples of such tissues are leaf, flower, root, stalk, runners above or below ground (side shoots, stolons), rhizomes,

WO 2007/096387

PCT/EP2007/051675

- 17 -

buds, tubers such as root tubers or stem tubers, bulb, brood bodies, brood buds, bulbuls or turion. Such tissues may also arise through pseudo vivipary, true vivipary or vivipary caused by humans. However, seeds arising through agamospermy, as are typical of Asteraceae, Poaceae or Rosaceae, are also included among the vegetative tissues in which expression
5 advantageously takes place. The nucleic acids used in the process of the invention are expressed to a small extent or not at all in generative tissue (germ line tissue). Examples of such tissues are tissues arising through sexual reproduction, i.e. meiotic cell divisions, such as, for example, seeds arising through sexual processes.

A small extent means that, compared with vegetative tissue, the expression measured at the
10 RNA and/or protein level is less than 5%, advantageously less than 3%, particularly advantageously less than 2%, most preferably less than 1; 0.5; 0.25 or 0.125%.

The nucleic acid sequences are particularly preferably expressed in the leaves of the transgenic plants. This has the advantage that the LCPUFAs produced according to the invention can be taken in by animals and humans directly by consuming the leaves, and no
15 previous processing of the plant material is necessary.

Expression of the nucleic acid sequences of the invention in the leaf can be achieved by using constitutive or leaf-specific promoters.

“Constitutive promoters” are promoters which make expression possible in a large number of, preferably in all, tissues over a substantial period during plant development, preferably
20 throughout plant development. A promoter from a plant or from a plant virus is preferably used. The promoter of the CaMV (cauliflower mosaic virus) 35S transcript (Franck et al. (1980) Cell 21: 285-294), the 19S CaMV promoter (US 5,352,605), the actin promoter from rice (McElroy et al. (1990) Plant Cell 2: 163-171), the legumin B promoter (GenBank Acc. No. X03677), the agrobacterium nopaline synthase promoter, the TR dual promoter, the
25 agrobacterium octopine synthase promoter, the ubiquitin promoter (Holtorf et al. (1995) Plant Mol. Biol. 29: 637-649), the Smas promoter, the cinnamoyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits, the pEMU promoter (Last et

WO 2007/096387

PCT/EP2007/051675

- 18 -

al. (1991) Theor. Appl. Genet. 81: 581-588), the MAS promoter (Velten et al. (1984) EMBO J. 3(12): 2723-2730), the histone H3 promoter from corn (Lepetit et al. (1992) Mol. Gen. Genet. 231: 276-285), the promoter of the nitrilase 1 gene from arabidopsis (GenBank Acc. No. U38846, nucleotides 3862-5325) and the promoter of a proline-rich protein from wheat
5 (WO 91/13991) and further promoters which mediate constitutive gene expression. The promoter of the CaMV 35S transcript is particularly preferred.

It is in principle possible to use all naturally occurring constitutive promoters with their regulatory sequences like those mentioned above for the novel process. However, it is likewise possible to use synthetic promoters in addition or alone.

10 “Leaf-specific promoters” are promoters which show a high activity in the leaf and no or only low activity in other tissues. “Low activity” means in the context of the invention that the activity in other tissues is less than 20%, preferably less than 10%, particularly preferably less than 5% and most preferably less than 3, 2 or 1% of the activity in the leaf. Examples of suitable leaf-specific promoters are the promoters of the small subunit of rubisco (Timko et al.
15 (1985) Nature 318: 579-582) and of the chlorophyll a/b-binding protein (Simpson et al. (1985) EMBO J. 4: 2723-2729).

The skilled worker is aware of further leaf-specific promoters, or he can isolate further suitable promoters with known methods. Thus, the skilled worker is able to identify leaf-specific regulatory nucleic acid elements with the aid of conventional methods of molecular
20 biology, e.g. hybridization experiments or DNA-protein binding studies. This entails for example in a first step isolating the total poly(A)⁺ RNA from leaf tissue of the desired organism from which the regulatory sequences are to be isolated, and setting up a cDNA library. In a second step, cDNA clones which are based on poly(A)⁺ RNA molecules from a non-leaf tissue are used to identify, by means of hybridization, those clones from the first
25 library whose corresponding poly(A)⁺ RNA molecules accumulate only in leaf tissue. Subsequently, these cDNAs identified in this way are used to isolate promoters which have

WO 2007/096387

PCT/EP2007/051675

- 19 -

leaf-specific regulatory elements. Further PCR-based methods for isolating suitable leaf-specific promoters are additionally available to the skilled worker.

It is, of course, also possible for the nucleic acid sequences of the present invention to be expressed in the seeds of the transgenic plants by using seed-specific promoters which are active in the embryo and/or in the endosperm. Seed-specific promoters can in principle be isolated both from dicotyledonous and from monocotyledonous plants. Preferred promoters are listed hereinafter: USP (unknown seed protein) and vicilin (*Vicia faba*) (Bäumlein et al. (1991) Mol. Gen Genet. 225(3): 459-467), napin (oilseed rape) (US 5,608,152), conlinin (flax) (WO 02/102970), acyl-carrier protein (oilseed rape) (US 5,315,001 and WO 92/18634), oleosin (*Arabidopsis thaliana*) (WO 98/45461 and WO 93/20216), phaseolin (*Phaseolus vulgaris*) (US 5,504,200), Bce4 (WO 91/13980), legume B4 (LegB4 promoter) (Bäumlein et al. (1992) Plant J. 2(2): 233-239), Lpt2 and lpt1 (barley) (WO 95/15389 and WO 95/23230), seed-specific promoters from rice, corn and wheat (WO 99/16890), Amy32b, Amy 6-6 and aleurain (US 5,677,474), Bce4 (oilseed rape) (US 5,530,149), glycinin (soybean) (EP 571 741), phosphoenolpyruvate carboxylase (soybean) (JP 06/62870), ADR 12-2 (soybean) (WO 98/08962), isocitrate lyase (oilseed rape) (US 5,689,040) or α -amylase (barley) (EP 781 849).

In a particularly preferred embodiment of the present invention, the nucleic acid sequences used, especially the nucleic acid sequence which codes for a $\Delta 5$ -elongase and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence originates by being adapted to the codon usage in one or more plant species, preferably the nucleic acid sequence described in SEQ ID NO: 64, are expressed in generative tissue, especially in the seed. Specific expression in the seed advantageously takes place by using one of the abovementioned seed-specific promoters, especially using the napin promoter. In this particularly preferred embodiment, the content of produced LCPUFAs, especially of the C22 fatty acids, in the seed oil is at least 5% by weight, advantageously at least 6, 7, 8, 9 or 10% by weight, preferably at least 11, 12, 13, 14 or 15% by weight, particularly preferably at least 16, 17, 18, 19 or 20% by weight, very particularly preferably at least 25, 30, 35 or 40% by weight, of the seed oil content. In a further particularly preferred

WO 2007/096387

PCT/EP2007/051675

- 20 -

embodiment with the nucleic acid sequence described in SEQ ID NO: 63, the content of C22 fatty acids in the seed oil is at least 8% by weight of the seed oil content.

In a further particularly preferred embodiment of the present invention, the nucleic acid sequences used, especially the nucleic acid sequence which codes for a $\Delta 5$ -elongase and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence originates by being adapted to the codon usage in one or more plant species, preferably the nucleic acid sequence described in SEQ ID NO: 64, are expressed in generative tissue, especially in the seed. Specific expression in the seed advantageously takes place by using one of the abovementioned seed-specific promoters, especially using the napin promoter. In this particularly preferred embodiment, the content of docosahexaenoic acid in the seed oil is at least 1% by weight, preferably at least 1.1, 1.2, 1.3, 1.4 or 1.5% by weight, particularly preferably at least 1.6, 1.7, 1.8 or 1.9% by weight, especially at least 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8 or 2.9% by weight, further preferably at least 3, 3.5 or 4% by weight of the seed oil content. In a further particularly preferred embodiment with the nucleic acid sequence described in SEQ ID NO: 63, the content of docosahexaenoic acid in the seed oil is at least 1.9% by weight of the seed oil content. It is known to the skilled worker in this connection that to produce docosahexaenoic acid additionally one or more nucleic acid sequences which codes for a polypeptide having the activity of a $\Delta 4$ -desaturase activity are required. A nucleic acid sequence which codes for a polypeptide having the activity of a $\Delta 4$ -desaturase activity is advantageously selected from the group consisting of nucleic acid sequences having the sequence depicted in SEQ ID No. 77, 79, 81, 83, 85, 87, 89, 91 or 93, preferably having the sequence depicted in SEQ ID No. 77.

In a further particularly preferred embodiment of the present invention, the nucleic acid sequences used, especially the nucleic acid sequence which codes for a $\Delta 5$ -elongase and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence originates by being adapted to the codon usage in one or more plant species, preferably the nucleic acid sequence described in SEQ ID NO: 64, are expressed in generative tissue, especially in the seed. Specific expression in the seed advantageously takes place by

WO 2007/096387

PCT/EP2007/051675

- 21 -

using one of the abovementioned seed-specific promoters, especially using the napin promoter. In this particularly preferred embodiment, the content of docosahexaenoic acid in the seed oil is at least 1% by weight, preferably at least 1.1, 1.2, 1.3, 1.4 or 1.5% by weight, particularly preferably at least 1.6, 1.7, 1.8 or 1.9% by weight, especially at least 2, 2.1, 2.2, 2.5, 2.6, 2.7, 2.8 or 2.9% by weight, further preferably at least 3, 3.5 or 4% by weight of the seed oil content. In this case, the content of the produced LCPUFAs, especially of the C22 fatty acids, in the seed oil is at least 5% by weight, advantageously at least 6, 7, 8, 9 or 10% by weight, preferably at least 11, 12, 13, 14 or 15% by weight, particularly preferably at least 16, 17, 18, 19 or 20% by weight, very particularly preferably at least 25, 30, 35 or 40% by weight of the seed oil content. In a further particularly preferred embodiment with the nucleic acid sequence described in SEQ ID NO: 63, the content of docosahexaenoic acid in the seed oil is at least 1.9% by weight of the seed oil content, with the content of C22 fatty acids in the seed oil being at least 8% by weight of the seed oil content.

Plant gene expression can also be achieved via a chemically inducible promoter (see a review in Gatz (1997) *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 48:89-108). Chemically inducible promoters are particularly suitable when it is desired that the gene expression takes place in a time-specific manner. Examples of such promoters are a salicylic-acid-inducible promoter (WO 95/19443), a tetracyclin-inducible promoter (Gatz et al. (1992) *Plant J.* 2, 397-404) and an ethanol-inducible promoter.

Promoters which respond to biotic or abiotic stress conditions are also suitable, for example the pathogen-induced PRP1 gene promoter (Ward et al., *Plant. Mol. Biol.* 22 (1993) 361-366), the heat-inducible tomato hsp80 promoter (US 5,187,267), the chill-inducible potato alpha-amylase promoter (WO 96/12814) or the wound-inducible pinII promoter (EP-A-0 375 091).

Other promoters which are also particularly suitable are those which bring about the plastid-specific expression, since plastids constitute the compartment in which precursors and some end products of lipid biosynthesis are synthesized. Suitable promoters, such as the viral RNA

WO 2007/096387

PCT/EP2007/051675

- 22 -

polymerase promoter, are described in WO 95/16783 and WO 97/06250, and the Arabidopsis clpP promoter, described in WO 99/46394.

It will be appreciated that the polyunsaturated fatty acids produced according to the invention can be produced not only in intact transgenic plants but also in plant cell cultures or in callous
5 cultures.

The polyunsaturated fatty acids produced in the process are advantageously bound in phospholipids and/or triacylglycerides, but may also occur as free fatty acids or else bound in the form of other fatty acid esters in the organisms. They may in this connection be present as “pure products” or else advantageously in the form of mixtures of various fatty acids or
10 mixtures of different phospholipids such as phosphatidylglycerol, phosphatidylcholine, phosphatidylethanolamine and/or phosphatidylserine and/or triacylglycerides, monoacylglycerides and/or diacylglycerides. The LCPUFAs EPA, DPA and DHA produced in the process are advantageously present in phosphatidylcholine and/or phosphatidylethanolamine and/or in the triacylglycerides. The triacylglycerides may additionally also comprise further
15 fatty acids such as short-chain fatty acids having 4 to 6 C atoms, medium-chain fatty acids having 8 to 12 C atoms or long-chain fatty acids having 14 to 24 C atoms. They preferably comprise long-chain fatty acids, particularly preferably C₂₀ or C₂₂ fatty acids.

The term “glyceride” is understood as meaning glycerol esterified with one, two or three carboxyl radicals (mono-, di- or triglyceride). “Glyceride” is also understood as meaning a
20 mixture of various glycerides. The glyceride is preferably a triglyceride. The glyceride or glyceride mixture can comprise further additions, for example free fatty acids, antioxidants, proteins, carbohydrates, vitamins and/or other substances.

A “glyceride” for the purposes of the process according to the invention is furthermore understood as meaning derivatives which are derived from glycerol. In addition to the above-
25 described fatty acid glycerides, these also include glycerophospholipids and glyceroglycolipids. Preferred examples which may be mentioned here are the glycerophospholipids such as

WO 2007/096387

PCT/EP2007/051675

- 23 -

lecithin (phosphatidylcholine), cardiolipin, phosphatidylglycerol, phosphatidylserine and alkylacylglycerophospholipids.

Phospholipids are to be understood as meaning, for the purposes of the invention, phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine, phosphatidylglycerol
5 and/or phosphatidylinositol.

The fatty acid esters with polyunsaturated C₁₈-, C₂₀- and/or C₂₂-fatty acid molecules can be isolated in the form of an oil or lipid, for example in the form of compounds such as sphingolipids, phosphoglycerides, lipids, glycolipids such as glycosphingolipids, phospholipids such as phosphatidylethanolamine, phosphatidylcholine, phosphatidylserine,
10 phosphatidylglycerol, phosphatidylinositol or diphosphatidylglycerol, monoacylglycerides, diacylglycerides, triacylglycerides or other fatty acid esters such as the acetyl-coenzyme A esters which comprise the polyunsaturated fatty acids with at least two, three or four, preferably four, five or six double bonds, from the useful plants which have been used for the preparation of the fatty acid esters; advantageously, they are isolated in the form of their
15 diacylglycerides, triacylglycerides and/or in the form of the phosphatidyl ester, especially preferably in the form of the triacylglycerides, phosphatidylcholine and/or phosphatidylethanolamine. In addition to these esters, the polyunsaturated fatty acids are also present in the plants as free fatty acids or bound in other compounds. As a rule, the various abovementioned compounds (fatty acid esters and free fatty acids) are present in the
20 organisms with an approximate distribution of 80 to 90% by weight of triglycerides, 2 to 5% by weight of diglycerides, 5 to 10% by weight of monoglycerides, 1 to 5% by weight of free fatty acids, 2 to 8% by weight of phospholipids, the total of the various compounds amounting to 100% by weight.

The LCPUFAs produced in the process of the invention are produced with a content of at least
25 4% by weight, advantageously of at least 5, 6, 7, 8, 9 or 10% by weight, preferably of at least 11, 12, 13, 14 or 15% by weight, particularly preferably of at least 16, 17, 18, 19, or 20% by weight, very particularly preferably of at least 25, 30, 35 or 40% by weight based on the total

WO 2007/096387

PCT/EP2007/051675

- 24 -

fatty acids in the transgenic plant. The fatty acids EPA, DPA and/or DHA produced in the process of the invention are moreover present with a content of in each case at least 5% by weight, preferably of in each case at least 6, 7, 8 or 9% by weight, particularly preferably of in each case at least 10, 11 or 12% by weight, most preferably of in each case at least 13, 14, 15, 16, 17, 18, 19 or 20% by weight based on the total fatty acids in the transgenic plant.

The fatty acids are advantageously produced in bound form. It is possible with the aid of the nucleic acids used in the process of the invention for these unsaturated fatty acids to be put on the sn1, sn2 and/or sn3 position of the advantageously produced triacylglycerides.

Advantageously, at least 11% of the triacylglycerides are doubly substituted (meaning on the sn1 and sn2 or sn2 and sn3 positions). Triply substituted triacylglycerides are also detectable. Since a plurality of reaction steps take place from the starting compounds linoleic acid (C18:2) and linolenic acid (C18:3), the final products of the process, such as, for example, arachidonic acid (ARA) or eicosapentaenoic acid (EPA), do not result as absolute pure products; traces or larger amounts of the precursors are always also present in the final product. If, for example, both linoleic acid and linolenic acid are present in the initial plant, the final products such as ARA or EPA and/or DPA and/or DHA are also present as mixtures. The precursors should advantageously amount to not more than 20% by weight, preferably not more than 15% by weight, particularly preferably not as 10% by weight, very particularly preferably not more than 5% by weight based on the amount of the respective final product. Advantageously, only ARA or EPA and/or DPA and/or DHA are produced in the process of the invention, bound or as free acids, as final products in a transgenic plant.

Fatty acid esters or fatty acid mixtures produced by the process of the invention advantageously comprise 6 to 15% palmitic acid, 1 to 6% stearic acid; 7-85% oleic acid; 0.5 to 8% vaccenic acid, 0.1 to 1% arachic acid, 7 to 25% saturated fatty acids, 8 to 85% monounsaturated fatty acids and 60 to 85% polyunsaturated fatty acids, in each case based on 100% and on the total fatty acid content of the organisms. Preferably at least 0.1; 0.2; 0.3; 0.4; 0.5; 0.6; 0.7; 0.8; 0.9 or 1% arachidonic acid in the total fatty acid content, are present as advantageous polyunsaturated fatty acid in the fatty acid ester or fatty acid mixtures. The fatty

WO 2007/096387

PCT/EP2007/051675

- 25 -

acid esters or fatty acid mixtures produced by the process of the invention further advantageously comprise fatty acids selected from the group of fatty acids erucic acid (13-docosaenoic acid), sterculic acid (9,10-methyleneoctadec-9-enonic acid), malvalic acid (8,9-methyleneheptadec-8-enonic acid), chaulmoogric acid (cyclopentenododecanoic acid),
 5 furan fatty acid (9,12-epoxyoctadeca-9,11-dienonic acid), vernonic acid (9,10-epoxyoctadec-12-enonic acid), taric acid (6-octadecynonic acid), 6-nonadecynonic acid, santalbic acid (11-octadecen-9-ynoic acid), 6,9-octadecenynonic acid, pyrulic acid (10-heptadecen-8-ynonic acid), crepenynic acid (9-octadecen-12-ynonic acid) 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diynonic acid, petroselenic acid (cis-6-octadecenonic acid), 9c,12t-
 10 octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid, catalpic acid (9t11t13c-octadecatrienoic acid), eleosteric acid (9c11t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), puniic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid) pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or
 15 coriolic acid (13-hydroxy-9c,11t-octadecadienonic acid). In general, the aforementioned fatty acids are advantageously present only in traces in the fatty acid esters or fatty acid mixtures produced by the process of the invention, meaning that their occurrence, based on the total fatty acid content, is less than 30%, preferably less than 25%, 24%, 23%, 22% or 21%, particularly preferably less than 20%, 15%, 10%, 9%, 8%, 7%, 6% or 5%, very particularly
 20 preferably less than 4%, 3%, 2% or 1%. In a further preferred form of the invention the occurrence of these aforementioned fatty acids, based on the total fatty acids, is less than 0.9%; 0.8%; 0.7%; 0.6% or 0.5%, particularly preferably less than 0.4%; 0.3%; 0.2%; 0.1%. The fatty acid esters or fatty acid mixtures produced by the process of the invention advantageously comprise less than 0.1% based on the total fatty acids and/or no butyric acid,
 25 no cholesterol and no nisinic acid (tetracosahexaenoic acid, C23:6^{Δ3,8,12,15,18,21}).

It is possible through the nucleic acid sequences used in the process of the invention to achieve an increase in the yield of LCPUFAs in the transgenic plants of at least 50%, advantageously of at least 80%, particularly advantageously of at least 100%, very particularly advantageously of at least 150%, compared with the non-transgenic plants.

WO 2007/096387

PCT/EP2007/051675

- 26 -

Chemically pure polyunsaturated fatty acids or fatty acid compositions can also be synthesized by the processes described above. To this end, the fatty acids or the fatty acid compositions are isolated from the plants in the known manner, for example via extraction, distillation, crystallization, chromatography or a combination of these methods. These

5 chemically pure fatty acids or fatty acid compositions are advantageous for applications in the food industry sector, the cosmetic sector and especially the pharmacological industry sector.

In principle, all dicotyledonous or monocotyledonous useful plants are suitable for the process of the invention. Useful plants mean plants which serve to produce foods for humans and animals, to produce other consumables, fibers and pharmaceuticals, such as cereals, e.g. corn,

10 rice, wheat, barley, millet, oats, rye, buckwheat; such as tubers, e.g. potato, cassava, sweet potato, yams etc.; such as sugar plants e.g. sugarcane or sugarbeet; such as legumes, e.g. beans, peas, broad bean etc.; such as oil and fat crops, e.g. soybean, oilseed rape, sunflower, safflower, flax, camolina etc., to mention only a few. Advantageous plants are selected from the group of plant families consisting of the families of Aceraceae, Actinidiaceae,

15 Anacardiaceae, Apiaceae, Arecaceae, Asteraceae, Arecaceae, Betulaceae, Boraginaceae, Brassicaceae, Bromeliaceae, Cannabaceae, Cannaceae, Caprifoliaceae, Chenopodiaceae, Convolvulaceae, Cucurbitaceae, Dioscoreaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Fagaceae, Grossulariaceae, Juglandaceae, Lauraceae, Liliaceae, Linaceae, Malvaceae, Moraceae, Musaceae, Oleaceae, Oxalidaceae, Papaveraceae, Poaceae,

20 Polygonaceae, Punicaceae, Rosaceae, Rubiaceae, Rutaceae, Scrophulariaceae, Solanaceae, Sterculiaceae and Valerianaceae.

Examples which may be mentioned are the following plants: Anacardiaceae such as the genera Pistacia, Mangifera, Anacardium, for example the genus and species Pistacia vera [pistachio], Mangifera indica (mango) or Anacardium occidentale (cashew), Asteraceae such as

25 the genera Calendula, Carthamus, Centaurea, Cichorium, Cynara, Helianthus, Lactuca, Locusta, Tagetes, Valeriana, e.g. the genus and species Calendula officinalis (common marigold), Carthamus tinctorius (safflower), Centaurea cyanus (cornflower), Cichorium intybus (chicory), Cynara scolymus (artichoke), Helianthus annuus (sunflower), Lactuca sativa,

WO 2007/096387

PCT/EP2007/051675

- 27 -

Lactuca crispa, Lactuca esculenta, Lactuca scariola L. ssp. sativa, Lactus scariola L. var.
 integrata, Lactuca scariola L. var. integrifolia, Lactuca sativa subsp. romana, Locusta
 communis, Valeriana locusta (lettuce), Tagetes lucida, Tagetes erecta or Tagetes tenuifolia
 (French marigold), Apiaceae such as the genus Daucus, e.g. the genus and species Daucus
 5 carota (carrot), Betulaceae such as the genus Corylus, e.g. the genera and species Corylus
 avellana or Corylus columna (hazelnut), Boraginaceae such as the genus Borago, e.g. the genus
 and species Borago officinalis (borage), Brassicaceae such as the genera Brassica, Camelina,
 Melanosinapis, Sinapis, Arabidopsis, e.g. the genera and species Brassica napus, Brassica
 rapa ssp. (oilseed rape), Sinapis arvensis Brassica juncea, Brassica juncea var. juncea,
 10 Brassica juncea var. crispifolia, Brassica juncea var. foliosa, Brassica nigra, Brassica
 sinapioides, Camelina sativa, Melanosinapis communis (mustard), Brassica oleracea (feed
 beet) or Arabidopsis thaliana, Bromeliaceae such as the genera Anana, Bromelia (pineapple),
 e.g. the genera and species Anana comosus, Ananas ananas or Bromelia comosa (pineapple),
 Caricaceae such as the genus Carica such as the genus and species Carica papaya (papaya),
 15 Cannabaceae such as the genus Cannabis such as the genus and species Cannabis sativa
 (hemp), Convolvulaceae such as the genera Ipomoea, Convolvulus, e.g. the genera and
 species Ipomoea batatas, Ipomoea pandurata, Convolvulus batatas, Convolvulus tiliaceus,
 Ipomoea fastigiata, Ipomoea tiliacea, Ipomoea triloba or Convolvulus panduratus (sweet
 potato, batate), Chenopodiaceae such as the genus Beta such as the genera and species Beta
 20 vulgaris, Beta vulgaris var. altissima, Beta vulgaris var. vulgaris, Beta maritima, Beta vulgaris
 var. perennis, Beta vulgaris var. conditiva or Beta vulgaris var. esculenta (sugarbeet),
 Cucurbitaceae such as the genus Cucurbita, e.g. the genera and species Cucurbita maxima,
 Cucurbita mixta, Cucurbita pepo or Cucurbita moschata (pumpkin), Elaeagnaceae such as the
 genus Elaeagnus, e.g. the genus and species Olea europaea (olive), Ericaceae such as the
 25 genus Kalmia, e.g. the genera and species Kalmia latifolia, Kalmia angustifolia, Kalmia
 microphylla, Kalmia polifolia, Kalmia occidentalis, Cistus chamaerhodendros or Kalmia
 lucida (mountain laurel), Euphorbiaceae such as the genera Manihot, Janipha, Jatropha,
 Ricinus, e.g. the genera and species Manihot utilissima, Janipha manihot, Jatropha manihot,
 Manihot aipil, Manihot dulcis, Manihot manihot, Manihot melanobasis, Manihot esculenta

WO 2007/096387

PCT/EP2007/051675

- 28 -

(cassava) or *Ricinis communis* (castor oil plant), Fabaceae such as the genera *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicago*, *Glycine*, *Dolichos*, *Phaseolus*, *Soja*, e.g. the genera and species *Pisum sativum*, *Pisum arvense*, *Pisum humile* (pea), *Albizia berteriana*, *Albizia julibrissin*, *Albizia lebbeck*, *Acacia berteriana*, *Acacia littoralis*, *Albizia berteriana*, *Albizia berteriana*, *Cathormion berteriana*, *Feuillea berteriana*, *Inga fragrans*, *Pithecellobium berterianum*, *Pithecellobium fragrans*, *Pithecolobium berterianum*, *Pseudalbizzia berteriana*, *Acacia julibrissin*, *Acacia nemu*, *Albizia nemu*, *Feuillea julibrissin*, *Mimosa julibrissin*, *Mimosa speciosa*, *Sericanrda julibrissin*, *Acacia lebbeck*, *Acacia macrophylla*, *Albizia lebbeck*, *Feuillea lebbeck*, *Mimosa lebbeck*, *Mimosa speciosa* (acacia), *Medicago sativa*, *Medicago falcata*, *Medicago varia* (alfalfa) *Glycine max* *Dolichos soja*, *Glycine gracilis*, *Glycine hispida*, *Phaseolus max*, *Soja hispida* or *Soja max* (soybean), Geraniaceae such as the genera *Pelargonium*, *Cocos*, *Oleum*, e.g. the genera and species *Cocos nucifera*, *Pelargonium grossularioides* or *Oleum cocois* (coconut), Gramineae such as the genus *Saccharum*, e.g. the genus and species *Saccharum officinarum*,

15 *Juglandaceae* such as the genera *Juglans*, *Wallia*, e.g. the genera and species *Juglans regia*, *Juglans ailanthifolia*, *Juglans sieboldiana*, *Juglans cinerea*, *Wallia cinerea*, *Juglans bixbyi*, *Juglans californica*, *Juglans hindsii*, *Juglans intermedia*, *Juglans jamaicensis*, *Juglans major*, *Juglans microcarpa*, *Juglans nigra* or *Wallia nigra* (walnut), Lauraceae such as the genera *Persea*, *Laurus*, e.g. the genera and species *Laurus nobilis* (bay), *Persea americana*, *Persea gratissima* or *Persea persea* (avocado), Leguminosae such as the genus *Arachis*. e.g. the genus and species *Arachis hypogaea* (peanut), Linaceae such as the genera *Linum*, *Adenolinum*, e.g. the genera and species *Linum usitatissimum*, *Linum humile*, *Linum austriacum*, *Linum bienne*, *Linum angustifolium*, *Linum catharticum*, *Linum flavum*, *Linum grandiflorum*, *Adenolinum grandiflorum*, *Linum lewisii*, *Linum narbonense*, *Linum perenne*, *Linum perenne*

20 var. *lewisii*, *Linum pratense* or *Linum trigynum* (flax), Lythrarieae such as the genus *Punica*, e.g. the genus and species *Punica granatum* (pomegranate), Malvaceae such as the genus *Gossypium*, e.g. the genera and species *Gossypium hirsutum*, *Gossypium arboreum*, *Gossypium barbadense*, *Gossypium herbaceum* or *Gossypium thurberi* (cotton), Musaceae such as the genus *Musa*, e.g. the genera and species *Musa nana*, *Musa acuminata*, *Musa*

WO 2007/096387

PCT/EP2007/051675

- 29 -

paradisiaca, Musa spp. (banana), Onagraceae such as the genera Camissonia, Oenothera, e.g. the genera and species Oenothera biennis or Camissonia brevipes (evening primrose), Palmae such as the genus Elaeis, e.g. the genus and species Elaeis guineensis (oil palm), Papaveraceae such as the genus Papaver, e.g. the genera and species Papaver orientale, Papaver rhoeas, 5 Papaver dubium (poppy), Pedaliaceae such as the genus Sesamum e.g the genus and species Sesamum indicum (sesame), Piperaceae such as the genera Piper, Artanthe, Peperomia, Steffensia, e.g. the genera and species Piper aduncum, Piper amalago, Piper angustifolium, Piper auritum, Piper betel, Piper cubeba, Piper longum, Piper nigrum, Piper retrofractum, Artanthe adunca, Artanthe elongata, Peperomia elongata, Piper elongatum, Steffensia elongata 10 (cayenne pepper), Poaceae such as the genera Hordeum, Secale, Avena, Sorghum, Andropogon, Holcus, Panicum, Oryza, Zea (corn), triticum, e.g. the genera and species Hordeum vulgare, Hordeum jubatum, Hordeum murinum, Hordeum secalinum, Hordeum distichon, Hordeum aegiceras, Hordeum hexastichon, Hordeum hexastichum, Hordeum irregulare, Hordeum sativum, Hordeum secalinum (barley), Secale cereale (rye), Avena 15 sativa, Avena fatua, Avena byzantina, Avena fatua var. sativa, Avena hybrida (oats), Sorghum bicolor, Sorghum halepense, Sorghum saccharatum, Sorghum vulgare, Andropogon drummondii, Holcus bicolor, Holcus sorghum, Sorghum aethiopicum, Sorghum arundinaceum, Sorghum caffrorum, Sorghum cernuum, Sorghum dochna, Sorghum drummondii, Sorghum durra, Sorghum guineense, Sorghum lanceolatum, Sorghum nervosum, 20 Sorghum saccharatum, Sorghum subglabrescens, Sorghum verticilliflorum, Sorghum vulgare, Holcus halepensis, Sorghum miliaceum, Panicum militaceum (millet), Oryza sativa, Oryza latifolia (rice), Zea mays (corn), Triticum aestivum, Triticum durum, Triticum turgidum, Triticum hybernum, Triticum macha, Triticum sativum or Triticum vulgare (wheat), Porphyridiaceae such as the genera Chroothece, Flintiella, Petrovanella, Porphyridium, 25 Rhodella, Rhodosorus, Vanhoeffenia, e.g. the genus and species Porphyridium cruentum, Proteaceae such as the genus Macadamia, e.g. the genus and species Macadamia intergrifolia (macadamia), Rubiaceae such as the genus Coffea, e.g. the genera and species Coffea spp., Coffea arabica, Coffea canephora or Coffea liberica (coffee), Scrophulariaceae such as the genus Verbascum, e.g. the genera and species Verbascum blattaria, Verbascum chaixii,

WO 2007/096387

PCT/EP2007/051675

- 30 -

Verbascum densiflorum, Verbascum lagurus, Verbascum longifolium, Verbascum lychnitis, Verbascum nigrum, Verbascum olympicum, Verbascum phlomoides, Verbascum phoenicum, Verbascum pulverulentum or Verbascum thapsus (mullein), Solanaceae such as the genera Capsicum, Nicotiana, Solanum, Lycopersicon, e.g. the genera and species Capsicum annuum, Capsicum annuum var. glabriusculum, Capsicum frutescens (pepper), Capsicum annuum (paprika), Nicotiana tabacum, Nicotiana alata, Nicotiana attenuata, Nicotiana glauca, Nicotiana langsdorffii, Nicotiana obtusifolia, Nicotiana quadrivalvis, Nicotiana repanda, Nicotiana rustica, Nicotiana sylvestris (tobacco), Solanum tuberosum (potato), Solanum melongena (aubergine), Lycopersicon esculentum, Lycopersicon lycopersicum, Lycopersicon pyriforme, Solanum integrifolium or Solanum lycopersicum (tomato), Sterculiaceae such as the genus Theobroma, e.g. the genus and species Theobroma cacao (cocoa), or Theaceae such as the genus Camellia, e.g. the genus and species Camellia sinensis (tea).

In an advantageous embodiment of the process, the useful plants used are oil fruit plants which comprise large amounts of lipid compounds, such as peanut, oilseed rape, canola, sunflower, safflower (*Carthamus tinctoria*), poppy, mustard, hemp, castor-oil plant, olive, sesame, Calendula, Punica, evening primrose, verbascum, thistle, wild roses, hazelnut, almond, macadamia, avocado, bay, pumpkin/squash, flax, soybean, pistachios, borage, trees (oil palm, coconut or walnut) or arable crops such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, Tagetes, Solanaceae plants such as potato, tobacco, egg plant and tomato, Vicia species, pea, alfalfa or bushy plants (coffee, cacao, tea), Salix species, and perennial grasses and fodder crops. Advantageous plants according to the invention are oil fruit plants such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, Calendula, Punica, evening primrose, pumpkin/squash, flax, soybean, borage, trees (oilpalm, coconut). Especially preferred are plants which are high in C18:2- and/or C18:3-fatty acids, such as sunflower, safflower, tobacco, verbascum, sesame, cotton, pumpkin/squash, poppy, evening primrose, walnut, flax, hemp or thistle. Very especially preferred plants are plants such as safflower, sunflower, poppy, evening primrose, walnut, flax, or hemp.

WO 2007/096387

PCT/EP2007/051675

- 31 -

It is also advantageous to express the nucleic acid sequences of the invention in the leaves of feed or food plants and thus to increase the content of eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in the leaves. Preferred feed plants are, for example, trefoil species such as red clover (*Trifolium pratense*), white clover (*Trifolium repens*), alsike clover (*Trifolium hybridum*), sainfoin (*Onobrychis viciifolia*), Egyptian clover (*Trifolium alexandrinum*) and Persian clover (*Trifolium resupinatum*). Preferred food plants are for instance lettuce species such as *Lactuca sativa*, *Lactuca crisper*, *Lactuca esculenta*, *Lactuca scariola* L. ssp. *sativa*, *Lactuca scariola* L. var. *integrata*, *Lactuca scariola* L. var. *integrifolia*, *Lactuca sativa* subsp. *romana*, *Locusta communis* and *Valeriana locusta*.

It is possible through the enzymatic activity of the nucleic acid sequences which are used in the process of the invention and which code for polypeptides having Δ -6-elongase, Δ -6-desaturase, Δ -5-desaturase and/or Δ -5-elongase activity, advantageously in combination with nucleic acid sequences which code for polypeptides having ω -3-desaturase and/or Δ -4-desaturase activity, and further nucleic acid sequences which code for polypeptides of fatty acid or lipid metabolism, such as further polypeptides having Δ -5-, Δ -6-, Δ -8-, Δ -12-desaturase or Δ -5-, Δ -6- or Δ -9-elongase activity, to produce a wide variety of polyunsaturated fatty acids in the process of the invention. Depending on the useful plants chosen for use in the process of the invention, mixtures of the various polyunsaturated fatty acids or individual polyunsaturated fatty acids such as EPA, DPA or DHA can be produced in free or bound form. Depending on the prevailing fatty acid composition in the starting plant (C18:2 or C18:3 fatty acids), the resulting fatty acids are derived from C18:2 fatty acids, such as GLA, DGLA or ARA or are derived from C18:3 fatty acids, such as EPA, DPA or DHA. If the only unsaturated fatty acid present in the plant used for the process is linoleic acid (LA, C18:2 ^{Δ 9,12}), the only possible products of the process are GLA, DGLA and ARA, which may be present as free fatty acids or bound. If the only unsaturated fatty acid present in the plant used in the process is α -linolenic acid (ALA, C18:3 ^{Δ 9,12,15}), for example as in flax, the only possible products of the process are SDA, ETA, EPA, DPA and/or DHA, which may be present as described above as free fatty acids or bound. It is possible to produce in a targeted manner only individual products in the plant by modifying the activity of the enzymes used in

WO 2007/096387

PCT/EP2007/051675

- 32 -

the process and involved in the synthesis Δ -6-elongase, Δ -6-desaturase, Δ -5-desaturase and/or Δ -6-elongase, advantageously in combination with further genes of lipid or fatty acid metabolism. Advantageously, only EPA, DPA or DHA or mixtures thereof are synthesized. Since the fatty acids are synthesized in biosynthesis chains, the respective final products are not present as pure substances in the organisms. Small amounts of the precursor compounds are always also present in the final product. These small amounts are less than 20% by weight, advantageously less than 15% by weight, particularly advantageously less than 10% by weight, very particularly advantageously less than 5, 4, 3, 2 or 1% by weight based on the final products EPA, DPA or DHA or mixtures thereof.

- 10 To increase the yield in the process according to the invention for the production of oils and/or triglycerides with a polyunsaturated fatty acid, content which is advantageously increased, it is advantageous to increase the amount of starting product for the synthesis of fatty acids. This can be achieved for example by introducing a nucleic acid which encodes a polypeptide with Δ 12-desaturase into the organism. This is particularly advantageous in
- 15 useful plants, such as oil-producing plants such as plants of the Brassicaceae family, such as the genus *Brassica*, for example rape; the Elaeagnaceae family, such as the genus *Elaeagnus*, for example the genus and species *Olea europaea* or the family Fabaceae, such as the genus *Glycine*, for example the genus and species *Glycine max*, which are high in oleic acid. Since these organisms have an only low linoleic acid content (Mikoklajczak et al. (1961) Journal of
- 20 the American Oil Chemical Society 38: 678 - 681) it is advantageous to use said Δ 12-desaturases for producing the starting material linolenic acid from oleic acid. It is also possible in addition for the starting fatty acids to be provided from outside, but this is less preferred for reasons of cost.

- 25 Mosses and algae are the only plant systems known to produce considerable amounts of polyunsaturated fatty acids such as arachidonic acid (ARA) and/or eicosapentaenoic acid (EPA) and/or docosahexaenoic acid (DHA). Mosses comprise PUFAs in membrane lipids, whereas algae, organisms related to algae, and some fungi also accumulate substantial amounts of PUFAs in the triacylglycerol fraction. Nucleic acid molecules isolated from

WO 2007/096387

PCT/EP2007/051675

- 33 -

strains which accumulate PUFAs also in the triacylglycerol fraction are therefore particularly advantageous for the process of the invention and thus for modifying the lipid and PUFA production system in a plant such as a useful plant such as an oil crop plant, for example oilseed rape, canola, flax, hemp, soybean, sunflower, borage. They can therefore
 5 advantageously be used in the process of the invention.

Nucleic acids used in the process of the invention are advantageously derived from plants such as algae, for example algae of the family of Prasinophyceae such as from the genera *Heteromastix*, *Mammella*, *Mantoniella*, *Micromonas*, *Nephroselmis*, *Ostreococcus*, *Prasinocladus*, *Prasinococcus*, *Pseudoscourfieldia*, *Pycnococcus*, *Pyramimonas*, *Scherffelia* or
 10 *Tetraselmis* such as the genera and species *Heteromastix longifillis*, *Mamiella gilva*, *Mantoniella squamata*, *Micromonas pusilla*, *Nephroselmis olivacea*, *Nephroselmis pyriformis*, *Nephroselmis rotunda*, *Ostreococcus tauri*, *Ostreococcus* sp. *Prasinocladus ascus*, *Prasinocladus lubricus*, *Pycnococcus provasolii*, *Pyramimonas amyliifera*, *Pyramimonas disomata*, *Pyramimonas obovata*, *Pyramimonas orientalis*, *Pyramimonas parkae*,
 15 *Pyramimonas spinefera*, *Pyramimonas* sp., *Tetraselmis apiculata*, *Tetraselmis carteriaformis*, *Tetraselmis chui*, *Tetraselmis convolutae*, *Tetraselmis desikacharyi*, *Tetraselmis gracilis*, *Tetraselmis hazeni*, *Tetraselmis impellucida*, *Tetraselmis inconspicua*, *Tetraselmis levis*, *Tetraselmis maculata*, *Tetraselmis marina*, *Tetraselmis striata*, *Tetraselmis subcordiformis*, *Tetraselmis suecica*, *Tetraselmis tetrabrachia*, *Tetraselmis tetrathele*, *Tetraselmis verrucosa*,
 20 *Tetraselmis verrucosa* fo. *rubens* or *Tetraselmis* sp. or algae from the family Euglenaceae such as from the genera *Ascoglena*, *Astasia*, *Colacium*, *Cyclidiopsis*, *Euglena*, *Euglenopsis*, *Hyalophacus*, *Khawkinea*, *Lepocinclis*, *Phacus*, *Strombomonas* or *Trachelomonas* such as the genera and species *Euglena acus*, *Euglena geniculata*, *Euglena gracilis*, *Euglena mixocylindracea*, *Euglena rostrifera*, *Euglena viridis*, *Colacium stentorium*, *Trachelomonas*
 25 *cylindrica* or *Trachelomonas volvocina*.

Further advantageous plants are algae such as *Isochrysis* or *Cryptocodinium*, algae/diatoms such as *Thalassiosira* or *Phaeodactylum*, mosses such as *Physcomitrella* or *Ceratodon* or higher plants such as the Primulaceae such as *Aleuritia*, *Calendula stellata*, *Osteospermum*

WO 2007/096387

PCT/EP2007/051675

- 34 -

spinescens or Osteospermum hyoseroides, microorganisms such as fungi such as Aspergillus, Thraustochytrium, Phytophthora, Entomophthora, Mucor or Mortierella, bacteria such as shewanella, yeasts or animals such as nematodes such as Caenorhabditis, insects, frogs, sea cucumbers or fishes. The nucleic acid sequences isolated according to the invention are

5 advantageously derived from an animal from the order of vertebrates. The nucleic acid sequences are preferably derived from the class of Vertebrata; Euteleostomi, Actinopterygii; Neopterygii; Teleostei; Euteleostei, Protacanthopterygii, Salmoniformes; Salmonidae or Oncorhynchus or Vertebrata, Amphibia, Anura, Pipidae, Xenopus or Evertebrata such as Protochordata, Tunicata, Holothuroidea, Cionidae such as Amaroucium constellatum,

10 Botryllus schlosseri, Ciona intestinalis, Molgula citrina, Molgula manhattensis, Perophora viridis or Styela partita. The nucleic acids are particularly advantageously derived from fungi, animals or from plants such as algae or mosses, preferably from the order of Salmoniformes such as of the family of Salmonidae such as of the genus Salmo, for example from the genera and species Oncorhynchus mykiss, Trutta trutta or Salmo trutta fario, from algae such as the

15 genera Mantoniella or Ostreococcus or from the diatoms such as the genera Thalassiosira or Phaeodactylum or from algae such as Cryptocodinium.

In a preferred embodiment, the process further comprises the step of obtaining a cell or a whole plant which comprises the nucleic acid sequences which are used in the process and which code for a Δ -6-desaturase, Δ -6-elongase, Δ -5-desaturase and/or Δ -5-elongase and, if

20 appropriate, nucleic acid sequences which code for an ω -3-desaturase and/or a Δ -4-desaturase, it being possible for the cell and/or the useful plant also to comprise further nucleic acid sequences of lipid or fatty acid metabolism. The nucleic acid sequences preferably used in the process are for expression advantageously incorporated into at least one gene construct and/or a vector as described hereinafter, alone or in combination with further nucleic acid sequences

25 which code for proteins of fatty acid or lipid metabolism, and finally transformed into the cell or plant. In a further preferred embodiment, this process further comprises the step of obtaining the oils, lipids or free fatty acids from the useful plants, The cell produced in this way or the useful plant produced in this way is advantageously a cell of an oil-producing plant, vegetable plant, lettuce plant, or ornamental plant or the plant itself as stated above.

WO 2007/096387

PCT/EP2007/051675

- 35 -

Growing means for the cultivation in the case of plant cells, tissue or organs on or in a nutrient medium or of the whole plant on or in a substrate, for example in hydroculture, flower pot soil or on an arable field.

For the purposes of the invention, "transgenic" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette (= gene construct) or a vector comprising the nucleic acid sequences used in the process according to the invention or a plant transformed with the nucleic acid sequences, expression cassette or vector used in the process according to the invention, all those constructions brought about by recombinant methods in which either

- 10 a) the nucleic acid sequence, or
- b) a genetic control sequence which is operably linked with the nucleic acid sequence, for example a promoter, or
- c) (a) and (b)

are not located in their natural genetic environment or have been modified by recombinant

- 15 methods, it being possible for the modification to be, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. Natural genetic environment means the natural genomic or chromosomal locus in the original organism or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The
- 20 environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, very especially preferably at least 5000 bp. A naturally occurring expression cassette – for example the naturally occurring combination of the natural promoter of the nucleic acid sequence used
- 25 proteins with corresponding $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase and $\Delta 5$ -elongase activity, advantageously in combination with nucleic acid sequences which encode proteins

WO 2007/096387

PCT/EP2007/051675

- 36 -

having ω 3-desaturase and/or Δ 4-desaturase activity – becomes a transgenic expression cassette when this expression cassette is modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in US 5,565,350 or WO 00/15815.

- 5 A "transgenic plant" for the purposes of the invention is understood as mentined above as meaning that the nucleic acids used in the process are not at their natural locus in the genome of the plant. In this case, it is possible for the nucleic acid sequences to be expressed homologously or heterologously. However, transgenic also means that, while the nucleic acids according to the invention are at their natural position in the genome of the plant, the sequence
 - 10 has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified. Transgenic is preferably understood as meaning the expression of the nucleic acids used in the process according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acid sequences takes place.
- 15 Preferred transgenic organisms are useful plants such as oil-producing plants, vegetable plants, lettuce plants or ornamental plants which are advantageously selected from the group of plant families consisting of the families of Aceraceae, Actinidiaceae, Anacardiaceae, Apiaceae, Arecaceae, Asteraceae, Arecaceae, Betulaceae, Boraginaceae, Brassicaceae, Bromeliaceae, Cannabaceae, Cannaceae, Caprifoliaceae, Chenopodiaceae, Convolvulaceae,
 - 20 Cucurbitaceae, Dioscoreaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Fagaceae, Grossulariaceae, Juglandaceae, Lauraceae, Liliaceae, Linaceae, Malvaceae, Moraceae, Musaceae, Oleaceae, Oxalidaceae, Papaveraceae, Poaceae, Polygonaceae, Punicaceae, Rosaceae, Rubiaceae, Rutaceae, Scrophulariaceae, Solanaceae, Sterculiaceae and Valerianaceae.
- 25 Host plants which are suitable for the nucleic acids, the expression cassette or the vector used in the process according to the invention are, in principle, advantageously all useful plants which are capable of synthesizing fatty acids, specifically unsaturated fatty acids, and which

WO 2007/096387

PCT/EP2007/051675

- 37 -

are suitable for the expression of recombinant genes. Examples which should be mentioned at this point are plants such as Arabidopsis, Asteraceae such as Calendula or useful plants such as soybean, peanut, castor-oil plant, sunflower, maize, cotton, flax, oilseed rape, coconut, oil palm, safflower (*Carthamus tinctorius*) or cacao bean. Further advantageous plants are
5 mentioned at other points in this application.

Microorganisms are generally used as intermediate hosts for the production of transgenic useful plants. Such utilizable intermediate host cells are detailed in: Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990).

- 10 Expression strains which can advantageously be used for this purpose are, for example, those with a lower protease activity. They are described, for example, in: Gottesman, S., Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, California (1990) 119-128.

Transgenic plants which comprise the polyunsaturated, long-chain fatty acids synthesized in
15 the process according to the invention can advantageously be marketed directly without there being any need for the oils, lipids or fatty acids synthesized to be isolated. This form of marketing is particularly advantageous.

“Plants” for the purposes of the present invention are intact plants and all plant parts, plant organs or plant parts such as leaf, stem, seeds, root, tubers, anthers, fibers, root hairs, stalks,
20 embryos, calli, cotyledons, petioles, harvested material, plant tissue, reproductive tissue and cell cultures which are derived from the actual transgenic plant and/or can be used for bringing about the transgenic plant. In this context, the seed comprises all parts of the seed such as the seed coats, epidermal cells, seed cells, endosperm or embryonic tissue.

The compounds produced in the process of the invention can, however, also be isolated from
25 the plants in the form of their oils, fat, lipids and/or free fatty acids. Polyunsaturated fatty acids produced by the process of the invention can be obtained by harvesting the plants or

WO 2007/096387

PCT/EP2007/051675

- 38 -

plant cells either from the culture in which they grow or from the field. This can take place by pressing or extracting the plant parts, preferably the plant seeds. It is possible in this connection for the oils, fats, lipids and/or free fatty acids to be obtained by pressing by so-called cold drawing or cold pressing without input of heat. To make it easier to break open
5 the plant parts, specifically the seeds, they are previously crushed, steamed or roasted. The seeds pretreated in this way can then be pressed or extracted with solvent such as warm hexane. The solvent is then removed again. It is possible in this way to isolate more than 96% of the compounds produced in the process of the invention. The products obtained in this way are then processed further, that is to say refined. This entails initially for example the plant
10 mucilage and suspended matter being removed. So-called desliming can take place enzymatically or, for example, chemically/physically by adding acid such as phosphoric acid. The free fatty acids are then removed by treatment with a base, for example sodium hydroxide solution. The resulting product is thoroughly washed with water to remove the alkali remaining in the product, and is dried. In order to remove the coloring matters still present in
15 the product, the products are subjected to a bleaching with, for example, bleaching earth or activated carbon. Finally, the product is also deodorized for example with steam.

The PUFAs or LCPUFAs produced by this process are preferably C₂₀ and/or C₂₂ fatty acid molecules having at least four double bonds in the fatty acid molecule, preferably five or six double bonds. These C₂₀ and/or C₂₂ fatty acid molecules can be isolated from the plant in the
20 form of an oil, lipid or a free fatty acid. Suitable transgenic plants are for example those mentioned above.

These oils, lipids or fatty acids of the invention comprise, as described above, advantageously 6 to 15% palmitic acid, 1 to 6% stearic acid; 7 - 85% oleic acid; 0.5 to 8% vaccenic acid, 0.1 to 1% arachic acid, 7 to 25% saturated fatty acids, 8 to 85% monounsaturated fatty acids and
25 60 to 85% polyunsaturated fatty acids, in each case based on 100% and on the total fatty acid content of the plants.

WO 2007/096387

PCT/EP2007/051675

- 39 -

Advantageous polyunsaturated, long-chain fatty acids present in the fatty acid esters or fatty acid mixtures such as phosphatidyl fatty acid esters or triacylglyceride esters are preferably at least 10; 11; 12; 13; 14; 15; 16; 17; 18; 19 or 20% by weight based on the total fatty acid content of eicosapentaenoic acid, based on the total fatty acid content, and/or at least 1; 2; 3; 4; 5 or 6% by weight of docosapentaenoic acid, based on the total fatty acid content, and/or at least 1; 2; 3; preferably at least 4; 5; 6; particularly preferably at least 7 or 8 and most preferably at least 9 or 10% by weight of docosahexaenoic acid, based on the total fatty acid content.

The fatty acid esters or fatty acid mixtures which have been produced by the process of the invention further comprise fatty acids selected from the group of fatty acids erucic acid (13-docosaic acid), sterculic acid (9,10-methylene octadec-9-enonic acid), malvalic acid (8,9-methylene heptadec-8-enonic acid), chaulmoogrinic acid (cyclopentenododecanoic acid), furan fatty acid (9,12-epoxyoctadeca-9,11-dienonic acid), vernonoic acid (9,10-epoxyoctadec-12-enonic acid), tarinic acid (6-octadecynonic acid), 6-nonadecynonic acid, santalbic acid (11-octadecen-9-ynoic acid), 6,9-octadecenynonic acid, pyrulic acid (10-heptadecen-8-ynonic acid), crepenynic acid (9-octadecen-12-ynonic acid) 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diynonic acid, petroselenic acid (cis-6-octadecenonic acid), 9c,12t-octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid, catalpic acid (9t11t13c-octadecatrienoic acid), eleosteric acid (9c11t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), punიცic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid) pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or coriolic acid (13-hydroxy-9c,11t-octadecadienonic acid). In general, the aforementioned fatty acids are advantageously present only in traces in the fatty acid esters or fatty acid mixtures produced by the process of the invention, meaning that their occurrence, based on the total fatty acid content, is less than 30%, preferably less than 25%, 24%, 23%, 22% or 21%, particularly preferably less than 20%, 15%, 10%, 9%, 8%, 7%, 6% or 5%, very particularly preferably less than 4%, 3%, 2% or 1%. In a further preferred form of the invention the occurrence of these aforementioned fatty acids, based on the total fatty acids,

WO 2007/096387

PCT/EP2007/051675

- 40 -

is less than 0.9%; 0.8%; 0.7%; 0.6% or 0.5%, particularly preferably less than 0.4%; 0.3%; 0.2%; 0.1%. The fatty acid esters or fatty acid mixtures produced by the process of the invention advantageously comprise less than 0.1% based on the total fatty acids and/or no butyric acid, no cholesterol and no nisinic acid (tetracosahexaenoic acid, C23:6^{Δ3,8,12,15,18,21}).

- 5 A further embodiment according to the invention is the use of the oils, the lipids, the fatty acids and/or the fatty acid composition, which are produced by the process of the invention, in feeding stuffs, foodstuffs, cosmetics or pharmaceuticals. The oils, lipids, fatty acids or fatty acid mixtures obtained in the process according to the invention can be used for admixture with other oils, lipids, fatty acids or fatty acid mixtures of animal origin, such as, for example,
- 10 fish oils, in the manner with which the skilled worker is familiar. These oils, lipids, fatty acids or fatty acid mixtures which are produced in this way and consist of vegetable and animal components can also be used for the preparation of feeding stuffs, foodstuffs, cosmetics or pharmaceuticals.

- The term "oil", "lipid" or "fat" is understood as meaning a fatty acid mixture comprising
- 15 unsaturated and/or saturated, preferably esterified fatty acid(s). It is preferred that the oil, fat or lipid is high in polyunsaturated free or advantageously esterified fatty acid(s), in particular linoleic acid, γ-linolenic acid, dihomo-γ-linolenic acid, arachidonic acid, α-linolenic acid, stearidonic acid, eicosatetraenoic acid, eicosapentaenoic acid, docosapentaenoic acid or docosahexaenoic acid. Preferably, the amount of unsaturated esterified fatty acids is
- 20 approximately 30%, with an amount of 50% being especially preferred and an amount of 60%, 70%, 80% or more being most preferred. The amount of the fatty acid can be determined by gas chromatography after converting the fatty acids into the methyl esters by transesterification. The oil, lipid or fat can comprise various other saturated or unsaturated fatty acids, for example calendulic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid
- 25 and the like. In particular, the amount of the various fatty acids can vary, depending on the starting plant.

WO 2007/096387

PCT/EP2007/051675

- 41 -

As described above, the polyunsaturated fatty acid esters advantageously having three, four, five or six, particularly advantageously having five or six double bonds and which have been prepared in the process advantageously take the form of fatty acid esters, for example, sphingolipid esters, phosphoglyceride esters, lipid esters, glycolipid esters, phospholipid esters, monoacylglycerol esters, diacylglycerol esters, triacylglycerol esters or other fatty acid esters, preference being given to phospholipid esters and/or triacylglycerol esters.

Starting with the polyunsaturated fatty acid esters produced thus in the process according to the invention and advantageously having at least three, four, five or six double bonds, the polyunsaturated fatty acids which are present can be liberated for example via treatment with alkali, for example with aqueous KOH or NaOH, or by acid hydrolysis, advantageously in the presence of an alcohol such as methanol or ethanol, or via enzymatic cleavage, and isolated via, for example, phase separation and subsequent acidification with, for example, H₂SO₄. However, the fatty acids can also be liberated directly without the above-described processing steps.

Substrates of the nucleic acid sequences used in the process which encode polypeptides with $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase and/or $\Delta 5$ -elongase activity and optionally nucleic acid sequences which encode polypeptides having $\omega 3$ -desaturase and/or $\Delta 4$ -desaturase activity, and/or of the further nucleic acids which are used, such as the nucleic acid sequences which encode polypeptides of the fatty acid or lipid metabolism selected from the group consisting of acyl-CoA dehydrogenase(s), acyl-ACP [= acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyl transferase(s), acyl-CoA:lysophospholipid acyltransferases, fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allene oxide synthases, hydroperoxide lyases or fatty acid elongase(s) are advantageously C₁₆-, C₁₈- or C₂₀-fatty acids. Preferably, the fatty acids converted in the process as substrates are converted in the form of their acyl-CoA esters and/or in the form of their phospholipid esters.

WO 2007/096387

PCT/EP2007/051675

- 42 -

To produce the long-chain PUFAs according to the invention, the saturated, monounsaturated C₁₆-fatty acids and/or polyunsaturated C₁₈-fatty acids must first, depending on the substrate, be desaturated and/or elongated or only deaturated by the enzymatic activity of a desaturase and/or elongase and subsequently elongated by at least two carbon atoms by an elongase.

- 5 After one elongation cycle, this enzyme activity leads either starting from C₁₆-fatty acids to C₁₈-fatty acids or starting from C₁₈-fatty acids to C₂₀-fatty acids, and after two elongation cycles starting from C₁₆-fatty acids leads to C₂₀-fatty acids. The activity of the desaturases or elongases used in the process according to the invention preferably leads to C₂₀- and/or C₂₂-fatty acids, advantageously with at least two or three double bonds in the fatty acid molecule, 10 preferably with four, five or six double bonds, especially preferably to C₂₂-fatty acids with at least five double bonds in the fatty acid molecule. Especially preferred products of the process according to the invention are eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid. The C₁₈-fatty acids with at least two double bonds in the fatty acid can be elongated by the enzymatic activity according to the invention in the form of the free fatty 15 acid or in the form of the esters, such as phospholipids, glycolipids, sphingolipids, phosphoglycerides, monoacylglycerol, diacylglycerol or triacylglycerol.

- The preferred biosynthesis site of fatty acids, oils, lipids or fats in the plants which are advantageously used is, for example, generally the seed or cell layers of the seed, so that seed-specific expression of the nucleic acids used in the process makes sense. However, it is 20 obvious that the biosynthesis of fatty acids, oils or lipids need not be limited to the seed tissue, but may also take place in a tissue specific manner in all of the remaining parts of the plant, for example in epidermal cells or in the tubers. The synthesis advantageously takes place according to the inventive process in the vegetative (somatic) tissue.

- Owing to the method according to the invention, the polyunsaturated fatty acids which are 25 produced can, in principle, be increased in two ways in the plants used in the process. Advantageously the pool of free polyunsaturated fatty acids and/or the amount of the esterified polyunsaturated fatty acids produced by the process can be increased. Advantageously, the pool of esterified polyunsaturated fatty acids in the transgenic plants is

WO 2007/096387

PCT/EP2007/051675

- 43 -

increased by the process according to the invention, advantageously in the form of the phosphatidyl esters and/or triacyl esters.

The sequences used in the process of the invention are cloned singly into expression constructs or provided on a joint recombinant nucleic acid molecule and used for introduction
5 and for expression in organisms. These expression constructs make it possible for the polyunsaturated fatty acids produced in the process of the invention to be synthesized optimally.

The nucleic acids used in the process may, after introduction into a plant or plant cell, either be located on a separate plasmid or advantageously be integrated into the genome of the host
10 cell. In the case of integration into the genome, the integration may be random or take place by recombination such that the native gene is replaced by the introduced copy, thus modulating production of the desired compound by the cell, or through use of a gene in trans, so that the gene is functionally connected to a functional expression unit which comprises at least one sequence ensuring the expression of a gene and at least one sequence ensuring the
15 polyadenylation of a functionally transcribed gene. The nucleic acid sequences are advantageously introduced into the plants via multiexpression cassettes or constructs for multiparallel expression, i.e. the nucleic acid sequences are present in a joint expression unit.

The nucleic acid construct may comprise more than one nucleic acid sequence coding for a polypeptide having the enzymatic activity of a Δ -12-desaturase, Δ -4-desaturase, Δ -5-
20 desaturase, Δ -6- desaturase, Δ -5-elongase, Δ -6- elongase, and/or ω -3- desaturase. It is also possible for a plurality of copies of a nucleic acid sequence coding for a polypeptide having the enzymatic activity of a Δ -12-desaturase, Δ -4-desaturase, Δ -5- desaturase, Δ -6- desaturase, Δ -5-elongase, Δ -6- elongase, and/or ω -3- desaturase to be present.

For the introduction, the nucleic acids used in the process are advantageously amplified and
25 ligated in the known manner. Preferably, a procedure following the protocol for Pfu DNA polymerase or a Pfu/Taq DNA polymerase mixture is followed. The primers are selected depending on the sequence to be amplified. The primers should expediently be chosen in such

WO 2007/096387

PCT/EP2007/051675

- 44 -

a way that the amplicon comprises the entire codogenic sequence from the start codon to the stop codon. After the amplification, the amplicon is expediently analyzed. For example, the analysis can be carried out by gel-electrophoretic separation with respect to quality and quantity. Thereafter, the amplicon can be purified following a standard protocol (for example

5 Qiagen). An aliquot of the purified amplicon is then available for the subsequent cloning step. Suitable cloning vectors are generally known to the skilled worker. These include, in particular, vectors which are capable of replication in microbial systems, that is to say mainly vectors which ensure efficient cloning in yeasts or fungi and which make possible the stable transformation of plants. Those which must be mentioned in particular are various binary and

10 cointegrated vector systems which are suitable for the T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they comprise at least the vir genes required for the Agrobacterium-mediated transformation and the T-DNA-delimiting sequences (T-DNA border). These vector systems preferably also comprise further cis-regulatory regions such as promoters and terminators and/or selection markers, by means of which suitably

15 transformed organisms can be identified. While in the case of cointegrated vector systems vir genes and T-DNA sequences are arranged on the same vector, binary systems are based on at least two vectors, one of which bears vir genes, but no T-DNA, while a second one bears T-DNA, but no vir genes. Owing to this fact, the last-mentioned vectors are relatively small, easy to manipulate and capable of replication both in *E. coli* and in *Agrobacterium*. These

20 binary vectors include vectors from the series pBIB-HYG, pPZP, pBecks, pGreen. In accordance with the invention, pBin19, pBI101, pBinAR, pGPTV and pCAMBIA are used by preference. An overview of the binary vectors and their use is found in Hellens et al. (2000) Trends in Plant Science 5: 446–451.

In order to prepare the vectors, the vectors can first be linearized with restriction

25 endonuclease(s) and then modified enzymatically in a suitable manner. Thereafter, the vector is purified, and an aliquot is employed for the cloning step. In the cloning step, the enzymatically cleaved and, if appropriate, purified amplificate is cloned with vector fragments which have been prepared in a similar manner, using ligase. In this context, a particular nucleic acid construct, or vector or plasmid construct, can have one or more than one

WO 2007/096387

PCT/EP2007/051675

- 45 -

codogenic gene segments. The codogenic gene segments in these constructs are preferably linked operably with regulatory sequences. The regulatory sequences include, in particular, plant sequences such as the above-described promoters and terminators. The constructs can advantageously be stably propagated in microorganisms, in particular *Escherichia coli* and

5 *Agrobacterium tumefaciens*, under selective conditions and thus make possible the transfer of heterologous DNA into plants.

The nucleic acid sequences and nucleic acid constructs used in the inventive process can be introduced into microorganisms and then into plants, advantageously using cloning vectors, and thus be used in the transformation of plants such as those which are published in and cited

10 therein: Plant Molecular Biology and Biotechnology (CRC Press, Boca Raton, Florida), Chapter 6/7, p. 71-119 (1993); F.F. White, Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, 15-38; B. Jenes et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press (1993), 128-143;

15 Potrykus, Annu. Rev. Plant Physiol. Plant Molec. Biol. (1991) 42: 205-225. Thus, the nucleic acids, nucleic acid constructs and/or vectors used in the process can be used for the recombinant modification of a broad spectrum of plants so that the latter become better and/or more efficient LCPUFA producers.

Owing to the introduction of a $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase and $\Delta 5$ -elongase

20 gene into a plant, alone or in combination with other genes, it is not only possible to increase biosynthesis flux towards the end product, but also to increase, or to create *de novo* the corresponding triacylglycerol and/or phosphatidylester composition. Likewise, the number or activity of other genes which are involved in the import of nutrients which are required for the biosynthesis of one or more fatty acids, oils, polar and/or neutral lipids, can be increased, so

25 that the concentration of these precursors, cofactors or intermediates within the cells or within the storage compartment is increased, whereby the ability of the cells to produce PUFAs, as described below, is enhanced further. By optimizing the activity or increasing the number of one or more of the $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase and/or $\Delta 5$ -elongase genes

WO 2007/096387

PCT/EP2007/051675

- 46 -

which are involved in the biosynthesis of these compounds, or by destroying the activity of one or more genes which are involved in the degradation of these compounds, it may be possible to increase the yield, production and/or production efficiency in fatty acid and lipid molecules from organisms and advantageously from plants.

- 5 The nucleic acid molecules used in the process of the invention code for proteins or parts thereof, whereas the proteins or the individual protein or parts thereof comprises an amino acid sequence which has sufficient homology to an amino acid sequence which is depicted in the sequences SEQ ID NO. 65, SEQ ID NO. 2, SEQ ID NO. 172 or SEQ ID NO. 52 and, if appropriate, SEQ ID NO. 194 or SEQ ID NO. 78, so that the proteins or parts thereof still
- 10 have a Δ -6-desaturase, Δ -6-elongase, Δ -5-desaturase and/or Δ -5-elongase activity and, if appropriate, a Δ -4-desaturase and/or ω -3-desaturase activity. The proteins or parts thereof which is/are encoded by the nucleic acid molecule/nucleic acid molecules preferably still have its/their essential enzymatic activity and the ability to participate in the metabolism of compounds necessary for constructing cell membranes or lipid bodies in organisms,
- 15 advantageously in plants, or in the transport of molecules across these membranes. The proteins encoded by the nucleic acid molecules are at least about 60% and preferably at least about 70%, 80% or 90%, and particularly preferably at least about 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequences depicted in SEQ ID NO. 65, SEQ ID NO. 2, SEQ ID NO. 172, SEQ ID
- 20 NO. 52, SEQ ID NO. 194 or SEQ ID NO. 78. Homology or homologous means in the context of the invention identity or identical.

- The homology was calculated over the entire amino acid or nucleic acid sequence region. To compare various sequences, the skilled worker has available a series of programs which are based on various algorithms. The algorithms of Needleman and Wunsch or Smith and
- 25 Waterman give particularly reliable results. The program PileUp (J. Mol. Evolution (1987) 25: 351-360; Higgins et al. (1989) CABIOS 5: 151-153) or the programs Gap and BestFit (Needleman and Wunsch (1970) J. Mol. Biol. 48: 443-453 and Smith and Waterman (1981) Adv. Appl. Math. 2: 482-489), which are part of the GCG software packet [Genetics

WO 2007/096387

PCT/EP2007/051675

- 47 -

Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711 (1991)], were used to carry out the sequence comparisons. The sequence homology data given above in % were determined over the entire sequence region using the program GAP with the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for sequence comparisons.

Essential enzymatic activity of the ω -3-desaturase, Δ -6-desaturase, Δ -6-elongase, Δ -5-elongase, Δ -4-desaturase and/or Δ -5-desaturase used in the process of the invention means that, compared with the proteins/enzymes encoded by the sequence having SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77, they still have an enzymatic activity of at least 10%, preferably of at least 20%, particularly preferably of at least 30% and most preferably of at least 40, 50 or 60%, and thus are able to participate in the metabolism of compounds necessary for synthesizing fatty acids, advantageously fatty acid esters such as phosphatidyl esters and/or triacylglyceride esters, in a plant or plant cell, or in the transport of molecules across membranes.

Nucleic acids which can be advantageously used in the process are derived from bacteria, fungi, diatoms, animals such as *Caenorhabditis* or *Oncorhynchus* or plants such as algae or mosses such as the genera *Shewanella*, *Physcomitrella*, *Thraustochytrium*, *Fusarium*, *Phytophthora*, *Ceratodon*, *Pytium irregulare*, *Mantoniella*, *Ostreococcus*, *Isochrysis*, *Aleurita*, *muscariaoides*, *Mortierella*, *Borago*, *Phaeodactylum*, *Cryptothecodinium*, specifically from the genera and species *Pytium irregulare*, *Oncorhynchus mykiss*, *Xenopus laevis*, *Ciona intestinalis*, *Thalassiosira pseudonana*, *Mantoniella squamata*, *Ostreococcus* sp., *Ostreococcus tauri*, *Euglena gracilis*, *Physcomitrella patens*, *Phytophthora infestans*, *Fusarium gramineum*, *Cryptocodium cohnii*, *Ceratodon purpureus*, *Isochrysis galbana*, *Aleurita farinosa*, *Thraustochytrium* sp., *Muscarioides viallii*, *Mortierella alpina*, *Borago officinalis*, *Phaeodactylum tricornutum*, *Caenorhabditis elegans* or particularly advantageously from *Pytium irregulare*, *Thraustochytrium* sp. and/or *Ostreococcus tauri*.

WO 2007/096387

PCT/EP2007/051675

- 48 -

It is possible additionally to use in the process of the invention nucleotide sequences which code for a Δ -12- desaturase, Δ -9- elongase or Δ -8- desaturase. The nucleic acid sequences used in the process are advantageously introduced in an expression cassette which makes expression of the nucleic acids in plants possible.

- 5 The nucleic acid sequences which code for the Δ -12- desaturase, ω -3- desaturase, Δ -9-elongase, Δ -6- desaturase, Δ -8- desaturase, Δ -6- elongase, Δ -5- desaturase, Δ -5- elongase or Δ -4- desaturase are functionally linked to one or more regulatory signals to increase the gene expression. These regulatory sequences are intended to make targeted expression of the genes possible. This may mean for example, depending on the plant, that the gene is
- 10 expressed and/or overexpressed only after induction, or that it is expressed and/or overexpressed immediately. Sequences advantageously used for the expression make constitutive expression possible, such as CaMV35S, CaMV36S, CaMV35Smas, nos, mas, ubi, stpt, lea or Super promoter. Expression preferably takes place in vegetative tissue as described above. In another preferred embodiment, the expression takes place in seeds.
- 15 These regulatory sequences are for example sequences to which inducers or repressors bind and thus regulate the expression of the nucleic acid. In addition to the regulatory sequences which are not linked in their natural locus to the nucleic acid sequences, or instead of these sequences, the natural regulation of these sequences may still be present before the actual structural genes and, if appropriate, have been genetically modified so that natural regulation
- 20 is switched off and expression of the genes is increased. The gene construct may additionally advantageously also comprise one or more so-called "enhancer sequences" functionally linked to the promoter, which make increased expression of the nucleic acid sequence possible. Additional advantageous sequences can also be inserted at the 3' end of the DNA sequences, such as further regulatory elements or terminators. Advantageous terminators are for example
- 25 viral terminators such as the 35S terminator or others. The nucleic acid sequences used in the process according to the invention may be present in one or more copies of the expression cassette (= gene construct). Preferably, only one copy of the genes is present in each expression cassette. This gene construct, or the gene constructs, can be introduced into the

WO 2007/096387

PCT/EP2007/051675

- 49 -

plant simultaneously or successively and expressed together in the host organism. In this context, the gene construct(s) can be inserted in one or more vectors and be present in the cell in free form, or else be inserted in the genome. It is advantageous for the insertion of further genes in the plant when the genes to be expressed are present together in one gene construct.

5 However, it is also possible to introduce in each case one gene construct containing a nucleic acid sequence into a plant and to cross the resulting plants with one another in order to obtain progeny which contains all gene constructs together.

In this context, the regulatory sequences or factors can, as described above, preferably have a positive effect on the gene expression of the genes introduced, thus enhancing it. Thus, an
10 enhancement of the regulatory elements, advantageously at the transcriptional level, may take place by using strong transcription signals such as promoters and/or enhancers. In addition, however, enhanced translation is also possible, for example by improving the stability of the mRNA.

To ensure the stable integration of the biosynthesis genes into the transgenic plant over a
15 plurality of generations, each of the nucleic acids which encode $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase or $\Delta 5$ -elongase and if appropriate the $\omega 3$ -desaturase or $\Delta 4$ -desaturase and which are used in the process should be expressed under the control of a separate promoter. This can be identical or different for each of the sequences. In this context, the expression cassette is advantageously constructed in such a way that a promoter is followed by a suitable
20 cleavage site for insertion of the nucleic acid to be expressed, which cleavage site is advantageously in a polylinker. If appropriate, a terminator can be positioned behind the polylinker. This sequence is repeated several times, preferably three, four, five or six times, so that up to six genes can be combined in one construct and thus introduced into the transgenic plant in order to be expressed. To express the nucleic acid sequences, the latter are inserted
25 behind the promoter via the suitable cleavage site, for example in the polylinker.

Advantageously, each nucleic acid sequence has its own promoter and, if appropriate, its own terminator. However, it is also possible to insert a plurality of nucleic acid sequences behind a promoter and, if appropriate, before a terminator. Here, the insertion site, or the sequence, of

WO 2007/096387

PCT/EP2007/051675

- 50 -

the inserted nucleic acids in the expression cassette is not of critical importance, that is to say a nucleic acid sequence can be inserted at the first or last position in the cassette without the expression being substantially influenced by the position. In an advantageous embodiment, different promoters such as, for example, the USP, LegB4 or DC3 promoter, and different
5 terminators can be used in the expression cassette. In a further advantageous embodiment, identical promoters such as the CaMV35S promoter can also be used.

As described above, the transcription of the genes which have been introduced should advantageously be terminated by suitable terminators at the 3' end of the biosynthesis genes which have been introduced (behind the stop codon). An example of a sequence which can be
10 used in this context is the OCS 1 or the 35SCaMV terminator. As is the case with the promoters, different terminator sequences should be used here for each gene.

As described above, the gene construct can also comprise further genes to be introduced into the organisms. It is possible and advantageous to introduce into the host plants, and to express therein, regulatory genes such as genes for inductors, repressors or enzymes which, owing to
15 their enzyme activity, engage in the regulation of one or more genes of a biosynthesis pathway. These genes can be of heterologous or of homologous origin. Moreover, further biosynthesis genes of the fatty acid or lipid metabolism can advantageously be present in the nucleic acid construct, or gene construct or alternatively, these genes can also be present on one further or more further nucleic acid constructs. A biosynthesis gene of the fatty acid or
20 lipid metabolism which is preferably chosen is one or more genes selected from the group of acyl-CoA dehydrogenase(s), acyl-ACP [= acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyl- transferase(s), acyl-CoA:lysophospholipid acyltransferases, fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases,
25 triacylglycerol lipases, allenoxide synthases, hydroperoxide lyases or fatty acid elongase(s) or combinations thereof. Especially advantageous nucleic acid sequences are biosynthesis genes of the fatty acid or lipid metabolism selected from the group of the acyl-

WO 2007/096387

PCT/EP2007/051675

- 51 -

CoA:lysophospholipid acyltransferase, $\Delta 8$ -desaturase, $\Delta 9$ -desaturase, $\Delta 12$ -desaturase and/or $\Delta 9$ -elongase.

In this context, the abovementioned nucleic acids or genes can be cloned into expression cassettes, like those mentioned above, in combination with other elongases and desaturases and used for transforming plants with the aid of *Agrobacterium*.

The term “vector” used in this description relates to a nucleic acid molecule which is capable of transporting another nucleic acid to which it is bound. One type of vector is a “plasmid”, a circular double-stranded DNA loop into which additional DNA segments can be ligated. A further type of vector is a viral vector, it being possible for additional DNA segments to be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they have been introduced (for example bacterial vectors with bacterial replication origin). Other vectors are advantageously integrated into the genome of a host cell when they are introduced into the host cell, and thus replicate together with the host genome. Moreover, certain vectors can govern the expression of genes with which they are in operable linkage. These vectors are referred to in the present context as “expression vectors”. Usually, expression vectors which are suitable for DNA recombination techniques take the form of plasmids. In the present description, “plasmid” and “vector” can be used exchangeably since the plasmid is the form of vector which is most frequently used. However, the invention is also intended to cover other forms of expression vectors, such as viral vectors, which exert similar functions. Furthermore, the term vector is also intended to encompass other vectors with which the skilled worker is familiar, such as phages, viruses such as SV40, CMV, TMV, transposons, IS elements, phasmids, phagemids, cosmids, linear or circular DNA.

The recombinant expression vectors advantageously used in the process comprise the nucleic acid sequences or the above-described gene construct used in the process in a form which is suitable for expressing the nucleic acids used in a host cell, which means that the recombinant expression vectors comprise one or more regulatory sequences, which are selected on the basis of the host cells to be used for the expression, which regulatory sequence(s) is/are linked

WO 2007/096387

PCT/EP2007/051675

- 52 -

operably with the nucleic acid sequence to be expressed. In a recombinant expression vector, “linked operably” means that the nucleotide sequence of interest is bound to the regulatory sequence(s) in such a way that the expression of the nucleotide sequence is made possible and they are bound to each other in such a way that both sequences carry out the predicted

5 function which is ascribed to the sequence (for example in an in-vitro transcription/translation system, or in a host cell if the vector is introduced into the host cell). The term “regulatory sequence” is intended to comprise promoters, enhancers and other expression control elements (for example polyadenylation signals). These regulatory sequences are described, for example, in Goeddel: Gene Expression Technology: Methods in Enzymology 185, Academic

10 Press, San Diego, CA (1990), or see: Gruber and Crosby, in: Methods in Plant Molecular Biology and Biotechnology, CRC Press, Boca Raton, Florida, Ed.: Glick and Thompson, Chapter 7, 89-108, including the references cited therein. Regulatory sequences comprise those which govern the constitutive expression of a nucleotide sequence in many types of host cell and those which govern the direct expression of the nucleotide sequence only in specific

15 host cells under specific conditions. The skilled worker knows that the design of the expression vector can depend on factors such as the choice of host cell to be transformed, the desired degree of expression of the protein and the like.

The recombinant expression vectors used can be designed for the expression of the nucleic acid sequences used in the process in such a way that they can be transformed into prokaryotic

20 intermediate hosts and finally, after introduction into the plants, make expression of the genes possible therein. This is advantageous because on account of simplicity, intermediate steps in vector construction are frequently carried out in microorganisms. For example, the Δ -6-desaturates, Δ -6-elongase, Δ -5-desaturate and/or Δ -5-elongase genes can be expressed in bacterial cells, insect cells (using baculovirus expression vectors), yeast cells and other fungal

25 cells (see Romanos, M.A., et al. (1992) Yeast 8:423-488; van den Hondel, C.A.M.J.J., et al. (1991) “Heterologous gene expression in filamentous fungi”, in: More Gene Manipulations in Fungi, J.W. Bennet & L.L. Lasure, Editors, pp. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J., & Punt, P.J. (1992) “Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, Peberdy, J.F., et al., Editors,

WO 2007/096387

PCT/EP2007/051675

- 53 -

pp. 1-28, Cambridge University Press: Cambridge), Algae (Falciatore et al. (1999) *Marine Biotechnology*.1: (3):239-251), ciliates, with vectors following a transformation process as described in WO 98/01572, and preferably in cells of multicellular plants (see Schmidt, R. and Willmitzer, L. (1988) "High efficiency *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* leaf and cotyledon explants" *Plant Cell Rep.*:538-586; 5 *Plant Molecular Biology and Biotechnology*, C Press, Boca Raton, Florida, chapter 6/7, pp. 71-119 (1993); F.F. White, B. Jenes et al., *Techniques for Gene Transfer in: Transgenic Plants*, vol. 1, Engineering and Utilization, Editors.: Kung and R. Wu, Academic Press (1993), 128-43; Potrykus (1991) *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42: 205-225 10 (and references cited therein)). Suitable hosts are what are further discussed in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). The recombinant expression vector may alternatively be transcribed and translated in vitro for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes usually takes place with vectors which comprise 15 constitutive or inducible promoters which control the expression of fusion or non-fusion proteins. Typical fusion expression vectors are inter alia pGEX (Pharmacia Biotech Inc; Smith, D.B., and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ), of which glutathione S-transferase (GST), maltose E-binding protein and protein A, respectively, are fused to the recombinant 20 target protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors are inter alia pTrc (Amann et al. (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression by the pTrc vector is based on transcription by host RNA 25 polymerase from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector is based on transcription from a T7-gn10-lac fusion promoter which is mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is provided by the host strains BL21 (DE3) or HMS174 (DE3) from a resident λ prophage which harbors a T7 gn1

WO 2007/096387

PCT/EP2007/051675

- 54 -

gene under transcription control of the lacUV 5 promoter.

Other vectors suitable in prokaryotic organisms are known to the skilled worker; these vectors are for example in *E. coli* pLG338, pACYC184, the pBR series such as pBR322, the pUC series such as pUC18 or pUC19, the M113mp series, pKC30, pRep4, pHS1, pHS2, pPLc236, 5 pMBL24, pLG200, pUR290, pIN-III113-B1, λ gt11 or pBdCI, in streptomyces pIJ101, pIJ364, pIJ702 or pIJ361, in bacillus pUB110, pC194 or pBD214, in corynebacterium pSA77 or pAJ667.

In a further embodiment, the expression vector is a yeast expression vector. Examples of vectors for expression in the yeast *S. cerevisiae* include pYeDesaturasecl (Baldari et al. 10 (1987) Embo J. 6:229-234), pMFa (Kurjan and Herskowitz (1982) Cell 30:933-943), pJRY88 (Schultz et al. (1987) Gene 54:113-123) and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and processes for constructing vectors suitable for use in other fungi, such as the filamentous fungi, are described in detail in: van den Hondel, C.A.M.J.J., & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular 15 Genetics of fungi, J.F. Peberdy et al., editors, pp. 1-28, Cambridge University Press: Cambridge, or in: More Gene Manipulations in Fungi (J.W. Bennet & L.L. Lasure, Editors, pp. 396-428: Academic Press: San Diego). Further suitable yeast vectors are for example pAG-1, YEpl6, YEpl3 or pEMBLYe23.

Alternatively, the nucleic acid sequences used in the process of the invention can be expressed 20 in insect cells using baculovirus expression vectors. Baculovirus vectors available for expressing proteins in cultured insect cells (e.g. Sf9 cells) include the pAc series (Smith et al. (1983) Mol. Cell Biol. 3:2156-2165) and the pVL series (Lucklow and Summers (1989) Virology 170:31-39).

The above mentioned vectors provide only a small survey of possible suitable vectors. Further 25 plasmids are known to the skilled worker and are described for example in: Cloning Vectors (Editors Pouwels, P.H. et al., Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). Further suitable expression systems for prokaryotic and eukaryotic cells see in

WO 2007/096387

PCT/EP2007/051675

- 55 -

chapters 16 and 17 of Sambrook, J., Fritsch, E.F., and Maniatis, T., Molecular Cloning: A Laboratory Manual, 2nd edition, Cold spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

5 The genes used in the process can also be expressed in single-celled plant cells (such as algae), see Falciatore et al. (1999) Marine Biotechnology 1 (3):239-251 and references cited therein, and in plant cells from higher plants (for example spermatophytes such as arable crops). Examples of plant expression vectors comprise those which are described in detail in: Becker, D., Kemper, E., Schell, J., and Masterson, R. (1992) Plant Mol. Biol. 20:1195-1197; and Bevan, M.W. (1984) Nucl. Acids Res. 12:8711-8721; Vectors for Gene Transfer in
10 Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, p. 15-38.

A plant expression cassette preferably comprises regulatory sequences which are capable of governing the expression of genes in plant cells and are linked operably so that each sequence can fulfil its function, such as transcriptional termination, for example polyadenylation
15 signals. Preferred polyadenylation signals are those which are derived from *Agrobacterium tumefaciens* T-DNA, such as gene 3 of the Ti plasmid pTiACH5 (Gielen et al., (1984) EMBO J. 3 835 et seq.), which is known as octopine synthase, or functional equivalents thereof, but all other terminators which are functionally active in plants are also suitable.

Since the regulation of plant gene expression is very often not limited to the transcriptional
20 level, a plant expression cassette preferably comprises other sequences which are linked operably, such as translation enhancers, for example the overdrive sequence, which enhances the tobacco mosaic virus 5' - untranslated leader sequence, which increases the protein/RNA ratio (Gallie et al. (1987) Nucl. Acids Research 15:8693-8711).

As described above, the plant gene expression must be linked operably with a suitable
25 promoter which controls gene expression. Advantageously utilizable promoters are constitutive promoters (Benfey et al., EMBO J. (1989) 8: 2195-2202), such as those which are derived from plant viruses, such as 35S CaMV (Franck et al. (1980) Cell 21: 285-294), 19S

WO 2007/096387

PCT/EP2007/051675

- 56 -

CaMV (see also US 5352605 and WO 84/02913), or plant promoters, such as the promoter of the Rubisco small subunit, which is described in US 4,962,028.

Other preferred sequences for use for functional connection in plant gene expression cassettes are targeting sequences which are necessary for guiding the gene product into its appropriate cellular compartment, for example into the vacuoles, the cell nucleus, all types of plastids such as amyloplasts, chloroplasts, chromoplasts, the extracellular space, the mitochondria, the endoplasmic reticulum, oil bodies, peroxisomes and other compartments of plant cells; (see a review in Kermode (1996) Crit. Rev. Plant Sci. 15(4): 284-423 and literature cited therein).

Vector DNA can be introduced into prokaryotic or eukaryotic cells via traditional transformation or transfection techniques. The terms "transformation" and "transfection", conjugation and transduction as used in the present context are intended to encompass the multiplicity of prior-art methods for introducing heterologous nucleic acids (for example DNA) into a host cell, including calcium phosphate or calcium chloride coprecipitation, DEAE-dextran-mediated transfection, lipofection, natural competence, chemically mediated transfer, electroporation or particle bombardment. Suitable methods for the transformation or transfection of host cells, including plant cells, can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual., 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) and other laboratory manuals such as Methods in Molecular Biology, 1995, Vol. 44, Agrobacterium protocols, Ed: Gartland and Davey, Humana Press, Totowa, New Jersey.

The term "nucleic acid (molecule)" as used herein comprises in an advantageous embodiment additionally the untranslated sequence located at the 3' end and at the 5' end of the coding gene region: at least 500, preferably 200, particularly preferably 100 nucleotides of the sequence upstream of the 5' end of the coding region and at least 100, preferably 50, particularly preferably 20 nucleotides of the sequence downstream of the 3' end of the coding gene region. An "isolated" nucleic acid molecule is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. An "isolated" nucleic

WO 2007/096387

PCT/EP2007/051675

- 57 -

acid preferably has no sequences which naturally flank the nucleic acid in the genomic DNA of the organism from which the nucleic acid is derived (e.g. sequences located at the 5' and 3' ends of the nucleic acid). In various embodiments, the isolated Δ -6-desaturase, Δ -6-elongase or Δ -5-desaturase and, if appropriate, the ω -3-desaturase or Δ -4-desaturase molecule used in the process may for example comprise less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in the genomic DNA of the cell from which the nucleic acid is derived.

The nucleic acid molecules used in the process can be isolated by using standard techniques of molecular biology and the sequence information provided herein. It is also possible for example with the aid of comparative algorithms to identify a homologous sequence or homologous, conserved sequence regions at the DNA or amino acid level. These can be used as hybridization probe in standard hybridization techniques (as described for example in Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd edition, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) for isolating further nucleic acid sequences useful in the process. The nucleic acid molecule used in the process, or parts thereof, can moreover be isolated by polymerase chain reaction, in which case oligonucleotide primers based on this sequence or on parts thereof are used (e.g. a nucleic acid molecule comprising the complete sequence or a part thereof can be isolated by polymerase chain reaction using oligonucleotide primers which have been constructed on the basis of this identical sequence). For example, mRNA can be isolated from cells (e.g. by the guanidinium thiocyanate extraction method of Chirgwin et al. (1979) Biochemistry 18:5294-5299) and cDNA can be prepared with the aid of reverse transcriptase (e.g. Moloney MLV reverse transcriptase obtainable from Gibco/BRL, Bethesda, MD or AMV reverse transcriptase, obtainable from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide primers for amplification by means of polymerase chain reaction can be constructed on the basis of one of the sequences shown in SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77 or with the aid of the amino acid sequences depicted in SEQ ID NO. 65, SEQ ID NO. 2, SEQ ID NO. 172, SEQ ID

WO 2007/096387

PCT/EP2007/051675

- 58 -

NO. 52, SEQ ID NO. 194 or SEQ ID NO. 78. A nucleic acid of the invention can be amplified by standard PCR amplification techniques using cDNA or alternatively genomic DNA as template and suitable oligonucleotide primers. The nucleic acid amplified in this way can be cloned into a suitable vector and characterized by DNA sequence analysis. Oligonucleotides
5 can be prepared by standard synthetic methods, for example using an automatic DNA synthesizer.

Homologs of the Δ -5-elongase, ω -3-desaturase, Δ -6-desaturase, Δ -6-elongase, Δ -4-desaturase or Δ -5-desaturase nucleic acid sequences used, having the sequence SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77, mean for
10 example allelic variants having at least about 40, 50 or 60%, preferably at least about 60 or 70%, more preferably at least about 70 or 80%, 90% or 95% and even more preferably at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity or homology to one of the nucleotide sequences shown in SEQ ID NO. 64, 66, 68 or 70, to one of the nucleotide sequences shown in SEQ ID NO. 1, 3, 5, 7, 9, 11, 13, 15,
15 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 or 41, to one of the nucleotide sequences shown in SEQ ID NO. 171, 173, 175, 177, 179, 181 or 183, to one of the nucleotide sequences shown in SEQ ID NO. 51, 53 or 55, to one of the nucleotide sequences shown in SEQ ID NO. 193 or 195 or to one of the nucleotide sequences shown in or SEQ ID NO. 77, 79, 81, 83, 85, 87, 89, 91 or 93, especially the nucleotide sequences shown in SEQ ID NO. 64, SEQ ID
20 NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77, or their homologs, derivatives or analogs or parts thereof. Also included are isolated nucleic acid molecules of a nucleotide sequence which hybridize for example under stringent conditions to one of the nucleotide sequences shown in SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77 or a part thereof. A part means in this
25 connection according to the invention that at least 25 base pairs (= bp), 50 bp, 75 bp, 100 bp, 125 bp or 150 bp, preferably at least 175 bp, 200 bp, 225 bp, 250 bp, 275 bp or 300 bp, particularly preferably 350 bp, 400 bp, 450 bp, 500 bp or more base pairs are used for the hybridization. It is also possible advantageously to use the complete sequence. Allelic variants

WO 2007/096387

PCT/EP2007/051675

- 59 -

comprise in particular functional variants which can be obtained by deletion, insertion or substitution of nucleotides from the sequence depicted in SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77, but where the enzyme activity of the proteins encoded thereby is substantially retained for the insertion.

- 5 Nucleic acid molecules advantageous for the process of the invention can be isolated on the basis of their homology to the ω -3-desaturase, Δ -6-desaturase, Δ -5-desaturase, Δ -5-elongase, Δ -4-desaturase and/or Δ -6-elongase nucleic acid sequences disclosed herein by using the sequences or a part thereof as hybridization probe in standard hybridization techniques under stringent hybridization conditions. It is possible in this connection for example to use isolated
- 10 nucleic acid molecules which are at least 15 nucleotides long and hybridize under stringent conditions with the nucleic acid molecules which comprise a nucleotide sequence of SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77. It is also possible to use nucleic acid molecules having at least 25, 50, 100, 250 or more nucleotides.
- 15 The term "hybridizes under stringent conditions" as used herein is intended to describe hybridization and washing conditions under which nucleic acid sequences which are at least 60% mutually homologous normally remain hybridized together. The conditions are preferably such that sequences which are at least about 65%, preferably at least about 70% and particularly preferably at least about 75% or more mutually homologous normally remain
- 20 hybridized together. These stringent conditions are known to the skilled worker and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-restrictive example of stringent hybridization conditions are hybridizations in 6 x sodium chloride/sodium citrate (= SSC) at about 45°C, followed by one or more washing steps in 0.2 x SSC, 0.1% SDS at 50 to 65°C. The skilled worker is aware
- 25 that these hybridization conditions differ depending on the type of nucleic acid and, for example organic solvents are present, in relation to the temperature and the concentration of the buffer. The temperature for example under "standard hybridization conditions" is, depending on the type of nucleic acid, between 42°C and 58°C in aqueous buffer with a

WO 2007/096387

PCT/EP2007/051675

- 60 -

concentration of 0.1 to 5 x SSC (pH 7.2). If organic solvent, for example 50% formamide, is present in the abovementioned buffer, the temperature under standard conditions is about 42°C. The hybridization conditions for DNA:DNA hybrids are preferably for example 0.1 x SSC and 20°C to 45°C, preferably 30°C to 45°C. The hybridization conditions for DNA:RNA hybrids are preferably for example 0.1 x SSC and 30°C to 55°C, preferably 45°C to 55°C. The aforementioned hybridization temperatures are determined for example for a nucleic acid with a length of about 100 bp (= base pairs) and a G + C content of 50% in the absence of formamide. The skilled person knows how the necessary hybridization conditions can be determined on the basis of textbooks such as the abovementioned or from the following textbooks Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989; Hames and Higgins (editors) 1985, "Nucleic Acids Hybridization: A Practical Approach", IRL Press at Oxford University Press, Oxford; Brown (editor) 1991, "Essential Molecular Biology: A Practical Approach", IRL Press at Oxford University Press, Oxford.

In order to determine the percentage of homology (= identity) of two amino acid sequences (for example one of the sequences of SEQ ID NO. 65, SEQ ID NO. 2, SEQ ID NO. 172, SEQ ID NO. 52, SEQ ID NO. 194 or SEQ ID NO. 78) or of two nucleic acids (for example SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77), the sequences are written one under the other in order to be able to compare them optimally (for example, gaps may be introduced into the sequence of a protein or of a nucleic acid in order to generate optimal alignment with the other protein or the other nucleic acid). Then, the amino acid radicals or nucleotides at the corresponding amino acid positions or nucleotide positions are compared. If a position in a sequence is occupied by the same amino acid radical or the same nucleotide as the corresponding position in another sequence, then the molecules are homologous at this position (i.e. amino acid or nucleic acid "homology" as used in the present context corresponds to amino acid or nucleic acid "identity"). The percentage of homology between the two sequences is a function of the number of identical positions which the sequences share (i.e. % homology = number of identical positions/total number of positions x 100). The programs and algorithms used to determine the homology are described above.

WO 2007/096387

PCT/EP2007/051675

- 61 -

An isolated nucleic acid molecule which codes for an ω -3-desaturase, Δ -6-desaturase, Δ -5-desaturase, Δ -5-elongase, Δ -4-desaturase and/or Δ -6-elongase which is used in the process and which is homologous to a protein sequence of SEQ ID NO. 65, SEQ ID NO. 2, SEQ ID NO. 172, SEQ ID NO. 52, SEQ ID NO. 194 or SEQ ID NO. 78 can be generated by

5 introducing one or more nucleotide substitutions, additions or deletions into a nucleotide sequence of SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID NO. 193 or SEQ ID NO. 77, so that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations may be introduced into one of the sequences of SEQ ID NO. 64, SEQ ID NO. 1, SEQ ID NO. 171, SEQ ID NO. 51, SEQ ID

10 NO. 193 or SEQ ID NO. 77 by standard techniques such as site-specific mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions are preferably produced at one or more of the predicted nonessential amino acid residues. In a "conservative amino acid substitution" the amino acid residue is replaced by an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art.

15 These families include amino acids with basic side chains (e.g. lysine, arginine, histidine), acidic side chains (e.g. aspartic acid, glutamic acid), uncharged polar side chains (e.g. glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g. alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g. threonine, valine, isoleucine) and aromatic side chains (e.g.

20 tyrosine, phenylalanine, tryptophan, histidine). A predicted nonessential amino acid residue in an ω -3-desaturase, Δ -6-desaturase, Δ -5-desaturase, Δ -5-elongase, Δ -4-desaturase or Δ -6-elongase is thus preferably replaced by another amino acid residue from the same side-chain family. An alternative possibility in another embodiment is to introduce the mutations randomly over the whole or a part of the ω -3-desaturase-, Δ -6-desaturase-, Δ -5-desaturase-, Δ -5-elongase-, Δ -4-desaturase- or Δ -6-elongase-encoding sequence, e.g. by saturation

25 mutagenesis, and the resulting mutants can be screened for the ω -3-desaturase, Δ -6-desaturase, Δ -5-desaturase, Δ -5-elongase, Δ -4-desaturase or Δ -6-elongase activity described herein in order to identify mutants which have retained the ω -3-desaturase, Δ -6-desaturase, Δ -5-desaturase, Δ -5-elongase, Δ -4-desaturase or Δ -6-elongase activity. The encoded protein can

be recombinantly expressed after the mutagenesis, and the activity of the protein can be determined for example by using the assays described herein.

The invention is illustrated in greater detail by the examples which follow, which are not to be construed as limiting.

5 The following table shows the sequence identifiers as used in the priority application of February 21, 2006, with the German application number 102006008030.0, and the corresponding sequence identifiers in this subsequent application. The nucleic acid sequence identified by SEQ ID NO: 1 of the priority application corresponds for example to the nucleic acid sequence
10 identified by SEQ ID NO: 64 of the subsequent application.

Table of concordance of sequence identifiers of the priority application and the sequence identifiers in the subsequent application:

SEQ ID NO: Priority application German application number 102006008030.0	SEQ ID NO: this subsequent application	Organism
1	64	Ostreococcus tauri
2	65	Ostreococcus tauri
3	1	Phytium irregulare
4	2	Phytium irregulare
5	171	Traustochytrium sp.
6	172	Traustochytrium sp.

WO 2007/096387

PCT/EP2007/051675

- 63 -

7	51	Thraustochytrium ssp.
8	52	Thraustochytrium ssp.
9	193	Phytophthora infestans
10	194	Phytophthora infestans
11	77	Traustochytrium sp.
12	78	Traustochytrium sp.
13	109	Ostreococcus tauri
n.a.	110	Ostreococcus tauri
14	122	Ostreococcus tauri
n.a.	123	Ostreococcus tauri
15	143	Ostreococcus tauri
16	144	Ostreococcus tauri
17	161	Cauliflower mosaic virus
18	162	Cauliflower mosaic virus
19	163	Thalassiosira pseudonana
20	164	Thalassiosira pseudonana

Examples

WO 2007/096387

PCT/EP2007/051675

- 64 -

Example 1: General cloning methods

The cloning methods such as, for example, restriction cleavages, agarose gel electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linkage of DNA fragments, transformation of *Escherichia coli* cells, bacterial

5 cultures and the sequence analysis of recombinant DNA were carried out as described by Sambrook et al. (1989) (Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6).

Example 2: Sequence analysis of recombinant DNA

Recombinant DNA molecules were sequenced with an ABI laser fluorescence DNA sequencer by the method of Sanger (Sanger et al. (1977) Proc. Natl. Acad. Sci. USA 74:

10 5463-5467). Fragments resulting from a polymerase chain reaction were sequenced and verified to avoid polymerase errors in constructs to be expressed.

Example 3: Cloning of genes from *Ostreococcus tauri*

It was possible by searching for conserved regions in an *Ostreococcus tauri* sequence database (genomic sequences) in each case a sequence coding for a protein having Δ -5-elongase

15 activity or Δ -6-elongase activity. These are the following sequences:

Gene name	SEQ ID	Amino acids
OtELO1.1, (Δ -6-Elongase)	SEQ ID NO. 143	292
OtELO2.1, (Δ -5-Elongase)	SEQ ID NO. 109	300

OtElo2.1 shows greatest similarity to an elongase from *Danio rerio* (GenBank AAN77156;

WO 2007/096387

PCT/EP2007/051675

- 65 -

approx. 26% identity), whereas OtElo1.1 shows greatest similarity to the elongase from *Physcomitrella* (PSE) (approx. 36% identity) (alignments were carried out with the tBLASTn algorithm (Altschul et al. (1990) J. Mol. Biol. 215: 403 - 410)).

The cloning of the elongases was carried out as follows:

5 40 ml of an *Ostreococcus tauri* culture in the stationary phase were spun down and resuspended in 100 µl of double-distilled water and stored at -20°C. The corresponding genomic DNAs were amplified by the PCR method. The corresponding primer pairs were selected so that they harbored the yeast consensus sequence for high-efficiency translation (Kozak (1986) Cell 44: 283-292) beside the start codon. Amplification of the OtElo DNAs
10 was carried out in each case with 1 µl of thawed cells, 200 µM dNTPs, 2.5 U Taq polymerase and 100 pmol of each primer in a total volume of 50 µl. The conditions for the PCR were as follows: first denaturation at 95°C for 5 minutes, followed by 30 cycles at 94°C for 30 seconds, 55°C for 1 minute and 72°C for 2 minutes, and a final elongation step at 72°C for 10 minutes.

15 Example 4: Optimization of elongase genes from *Ostreococcus tauri*

Elongases from the organism *Ostreococcus tauri* were isolated as described in example 3. In order to achieve an increase in the content of C22 fatty acids, the sequences SEQ ID No. 143 (Δ6-elongase) and SEQ ID No. 109 (coding for a protein identified by SEQ ID No. 110) (Δ5-elongase) were adapted to the codon usage in oilseed rape, flax and soybean. For this
20 purpose, the amino acid sequence of the Δ6-elongase and of the Δ5-elongase (SEQ ID NO. 144 for the Δ6-elongase; SEQ ID NO. 65 for the Δ5-elongase) was back-translated to obtain degenerate DNA sequences. These DNA sequences were adapted by means of the GeneOptimizer program (from Geneart, Regensburg) to the codon usage in oilseed rape, soybean and flax, taking account of the natural frequency of individual codons. The optimized
25 sequences obtained in this way, which are indicated in SEQ ID NO. 64 (Δ5-elongase) and SEQ ID NO. 122 (coding for a protein identified by SEQ ID NO. 123) (Δ6-elongase) were

WO 2007/096387

PCT/EP2007/051675

- 66 -

synthesized in vitro.

Example 5: Cloning of expression plasmids for heterologous expression in yeasts

To characterize the function of the optimized nucleic acid sequences, the open reading frames of the respective DNAs were cloned downstream of the galactose-inducible GAL1 promoter of pYES2.1/V5-His-TOPO (Invitrogen), resulting in the plasmids pOTE1.2 (comprising the
5 Δ 6-elongase sequence) and pOTE2.2 (comprising the Δ 5-elongase sequence).

Overview of the elongase sequences cloned into the yeast vector pYES2.1/V5-His-TOPO:

Gene name	SEQ ID	Amino acids
pOTE1.1, (Δ -6-elongase)	SEQ ID NO. 143	292
pOTE1.2, (Δ -6-elongase)	SEQ ID NO. 122	292, codon-optimized
pOTE2.1, (Δ -5-elongase)	SEQ ID NO. 109	300
pOTE2.2, (Δ -5-elongase)	SEQ ID NO. 64	300, codon-optimized

The *Saccharomyces cerevisiae* strain 334 was transformed by electroporation (1500 V) with
10 the vectors pOTE1.2 and pOTE2.2 and with the comparative constructs pOTE1.1 and pOTE2.1 which comprise the natural nucleic acid sequence coding for the Δ 6-elongase and Δ 5-elongase, respectively. A yeast transformed with the empty vector pYES2 was used as control. The transformed yeasts were selected on complete minimal medium (CMdum) agar plates with 2% glucose but without uracil. After the selection, three transformants in each case
15 were selected for further functional expression.

To express the Ot elongases, initially precultures composed of in each case 5 ml of CMdum liquid medium with 2% (w/v) raffinose but without uracil were inoculated with the selected

WO 2007/096387

PCT/EP2007/051675

- 67 -

transformants and incubated at 30°C, 200 rpm for 2 days. 5 ml of CMdum liquid medium (without uracil) with 2% raffinose were then inoculated with the precultures to an OD₆₀₀ of 0.05. Moreover, 0.2 mM γ -linolenic acid (GLA) was added in each case to the yeast culture transformed with pOTE1.1 and pOTE1.2. On the basis of the activity of OtELO1.1, an
5 elongation of the γ -linolenic acid to the 20:3 fatty acid is to be expected. Respectively 0.2 mM arachidonic acid and eicosapentaenoic acid were added in each case to the yeast culture transformed with pOTE2.1 and pOTE2.2. Corresponding to the activity of OtELO2.1, it is to be expected that the fatty acids ARA and EPA will be elongated respectively to the 22:4 and 22:5 fatty acids. Expression was induced by adding 2% (w/v) galactose. The cultures were
10 incubated at 20°C for a further 96 h.

Example 6: Expression of OtELO2.2 (as depicted in SEQ ID NO: 64) and OtELO1.2 (as in SEQ ID NO: 122) in yeasts

Yeasts transformed as in example 5 with the plasmids pYES2, pOTE1.2 and pOTE2.1 were analyzed in the following way:

15 The yeast cells from the main cultures were harvested by centrifugation (100 x g, 5 min, 20°C) and washed with 100 mM NaHCO₃, pH 8.0, in order to remove remaining medium and fatty acids. Fatty acid methyl esters (FAMES) were prepared from the yeast cell sediments by acidic methanolysis. For this purpose, the cell sediments were incubated with 2 ml of 1 N methanolic sulfuric acid and 2% (v/v) dimethoxypropane at 80°C for 1 h. The FAMES were
20 extracted by extraction twice with petroleum ether (PE). To remove underivatized fatty acids, the organic phases were washed once each with 2 ml of 100 mM NaHCO₃, pH 8.0 and with 2 ml of distilled water. The PE phases were then dried with Na₂SO₄, evaporated under argon and taken up in 100 μ l of PE. The samples were separated on a DB-23 capillary column (30 m, 0.25 mm, 0.25 μ m, Agilent) in a Hewlett-Packard 6850 gas chromatograph with flame
25 ionization detector. The conditions for the GLC analysis were as follows: the oven temperature was programmed from 50°C to 250°C at a rate of 5°C/min and finally 10 min at 250°C (holding).

WO 2007/096387

PCT/EP2007/051675

- 68 -

The signals were identified by comparing the retention times with appropriate fatty acid standards (Sigma). The methodology is described for example in Napier and Michaelson (2001) *Lipids* 36(8): 761-766; Sayanova et al. (2001) *Journal of Experimental Botany* 52(360): 1581-1585, Sperling et al. (2001) *Arch. Biochem. Biophys.* 388(2): 293-298 and
 5 Michaelson et al. (1998) *FEBS Letters* 439(3): 215-218. The results of the analyses are depicted in table 1.

It was possible to confirm the appropriate activities both for pOTE1.1/pOTE1.2 and for pOTE2.1/2.2. The optimized sequence (respectively pOTE1.2 and pOTE2.2) showed activity in both cases. Synthesis of γ -linolenic acid could be increased only slightly by pOTE1.2
 10 compared with the wild-type sequence. By contrast, it was possible to observe for pOTE2.2 surprisingly both an increase in the activity and an alteration in the specificity (table 1). In this connection, the activity for elongation of EPA had virtually doubled, while the elongation of ARA had more than quadrupled. It was thus possible with the optimization of the sequence of the $\Delta 5$ -elongase from *Ostreococcus tauri* to increase the yield of the precursors of DHA 6-fold
 15 in yeast with the same amount of substrate.

Example 7: Cloning expression plasmids for the seed-specific expression in plants

The following general conditions described apply to all subsequent experiments unless described otherwise.

pBin19, pBI101, pBinAR, pGPTV, pCAMBIA or pSUN are preferably used for the following
 20 examples in accordance with the invention. An overview of the binary vectors and their use can be found in Hellens et al, *Trends in Plant Science* (2000) 5: 446–451. A pGPTV derivative as described in DE10205607 was used. This vector differs from pGPTV by an additionally inserted *AscI* restriction cleavage site.

Starting point of the cloning procedure was the cloning vector pUC19 (Maniatis et al.). In the
 25 first step, the conlinin promoter fragment was amplified using the following primers:

Cnl1 C 5': gaattcggcgcgccgagctcctcgagcaacgggtccggcggtatagagttgggtaattcga

WO 2007/096387

PCT/EP2007/051675

- 69 -

Cnl1 C 3': cccgggatcgatgccggcagatctccaccatttttggtggtgat

Composition of the PCR mix (50 µl):

5.00 µl template cDNA

5 5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂

5.00 µl of 2mM dNTP

1.25 µl of each primer (10 pmol/µl)

0.50 µl of Advantage polymerase (Clontech)

10 PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

Number of cycles: 35

15

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme EcoRI and then for 12 hours at 25°C with the restriction enzyme SmaI. The cloning vector pUC19 was incubated in the same manner. Thereafter, the PCR product and the 2668 bp cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1-C was verified by sequencing.

20

In the next step, the OCS terminator (Genbank Accession V00088; De Greve, H., et al. (1982) J. Mol. Appl. Genet. 1 (6): 499-511) was amplified from the vector pGPVT-USP/OCS (DE 102 05 607) using the following primers:

25

WO 2007/096387

PCT/EP2007/051675

- 70 -

OCS_C 5': aggcctccatggcctgcttaatgagatatgcgagacgcc

OCS_C 3': cccgggccggacaatcagtaaattgaacggag

Composition of the PCR mix (50 µl):

- 5 5.00 µl template cDNA
5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂
5.00 µl of 2mM dNTP
1.25 µl of each primer (10 pmol/µl)
0.50 µl of Advantage polymerase (Clontech)

10

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

- 15 Number of cycles: 35

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme StuI and then for 12 hours at 25°C with the restriction enzyme SmaI. The vector pUC19-Cn11-C was incubated for 12 hours at 25°C with the restriction enzyme SmaI. Thereafter, the PCR product and the cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cn11-C_OCS was verified by sequencing.

- 25 In the next step, the Cn11-B promoter was amplified by PCR by means of the following primers:

WO 2007/096387

PCT/EP2007/051675

- 71 -

Cnl1-B 5': aggcctcaacggttcggcggtatag

Cnl1-B 3': cccgggggtaacgctagcgggcccgatcggatcccatttttggtggtgattggttct

Composition of the PCR mix (50 µl):

- 5 5.00 µl template cDNA
5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂
5.00 µl of 2mM dNTP
1.25 µl of each primer (10 pmol/µl)
0.50 µl of Advantage polymerase (Clontech)

10

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

- 15 Number of cycles: 35

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme StuI and then for 12 hours at 25°C with the restriction enzyme SmaI. The vector pUC19-Cnl1-C was incubated for 12 hours at 25°C with the restriction enzyme SmaI. Thereafter, the PCR product and the cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1-C_Cnl1B_OCS was verified by sequencing.

- 25 In a further step, the OCS terminator for Cnl1B was inserted. To this end, the PCR was carried out using the following primers:

WO 2007/096387

PCT/EP2007/051675

- 72 -

OCS2 5': aggcctcctgctttaatgagatatgcgagac

OCS2 3': cccgggcggacaatcagtaaattgaacggag

Composition of the PCR mix (50 µl):

- 5 5.00 µl template cDNA
- 5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂
- 5.00 µl of 2mM dNTP
- 1.25 µl of each primer (10 pmol/µl)
- 0.50 µl of Advantage polymerase (Clontech)

10

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

- 15 Number of cycles: 35

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme StuI and then for 12 hours at 25°C with the restriction enzyme SmaI. The vector pUC19-

Cnl1C_Cnl1B_OCS was incubated for 12 hours at 25°C with the restriction enzyme SmaI.

- 20 Thereafter, the PCR product and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1-C_Cnl1B_OCS2 was verified by sequencing.

- 25 In the next step, the Cnl1-A promoter is amplified by PCR using the following primers:

WO 2007/096387

PCT/EP2007/051675

- 73 -

Cnl1-B 5': aggcctcaacggttccggcggtatagag

Cnl1-B 3': aggccttctagactgcaggcggccgcccgcatttttggtggtgattggt

Composition of the PCR mix (50 µl):

- 5 5.00 µl template cDNA
- 5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂
- 5.00 µl of 2mM dNTP
- 1.25 µl of each primer (10 pmol/µl)
- 0.50 µl of Advantage polymerase (Clontech)

10

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

- 15 Number of cycles: 35

The PCR product was incubated for 2 hours at 37°C with the restriction enzyme StuI. The vector pUC19-Cnl1-C was incubated for 12 hours at 25°C with the restriction enzyme SmaI. Thereafter, the PCR product and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1C_Cnl1B_Cnl1A_OCS2 was verified by sequencing.

In a further step, the OCS terminator for Cnl1A was inserted. To this end, the PCR was carried out with the following primers:

25

WO 2007/096387

PCT/EP2007/051675

- 74 -

OCS2 5': ggcctcctgctttaatgagatatgcga

OCS2 3': aagcttggcgcgcccagctcgtcgacggacaatcagtaaattgaacggaga

Composition of the PCR mix (50 µl):

- 5 5.00 µl template cDNA
5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂
5.00 µl of 2mM dNTP
1.25 µl of each primer (10 pmol/µl)
0.50 µl of Advantage polymerase (Clontech)

10

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

- 15 Number of cycles: 35

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme StuI and then for 2 hours at 37°C with the restriction enzyme HindIII. The vector pUC19-

Cnl1C_Cnl1B_Cnl1A_OCS2 was incubated for 2 hours at 37°C with the restriction enzyme

- 20 StuI and for 2 hours at 37°C with the restriction enzyme HindIII. Thereafter, the PCR product and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated.

The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-

- 25 Cnl1-C_Cnl1B_Cnl1A_OCS3 was verified by sequencing.

WO 2007/096387

PCT/EP2007/051675

- 75 -

In the next step, the plasmid pUC19-Cnl1C_Cnl1B_Cnl1A_OCS3 was used for cloning the $\Delta 6$ -, $\Delta 5$ -desaturase and $\Delta 6$ -elongase. To this end, the *Phytium irregulare* $\Delta 6$ -desaturase (WO02/26946) was amplified using the following PCR primers:

- 5 D6Des(Pir) 5': agatctatggtggacctcaagcctggagtg
D6Des(Pir) 3': ccatggcccgggttacatcgctgggaactcggtgat

Composition of the PCR mix (50 μ l):

- 5.00 μ l template cDNA
10 5.00 μ l 10x buffer (Advantage polymerase) + 25mM $MgCl_2$
5.00 μ l of 2mM dNTP
1.25 μ l of each primer (10 pmol/ μ l)
0.50 μ l of Advantage polymerase (Clontech)

- 15 PCR reaction conditions:

Annealing temperature: 1 min 55°C
Denaturation temperature: 1 min 94°C
Elongation temperature: 2 min 72°C
Number of cycles: 35

20

- The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme Bg/II and then for 2 hours at 37°C with the restriction enzyme NcoI. The vector pUC19-Cnl1C_Cnl1B_Cnl1A_OCS3 was incubated for 2 hours at 37°C with the restriction enzyme Bg/II and for 2 hours at 37°C with the restriction enzyme NcoI. Thereafter, the PCR product
25 and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA

WO 2007/096387

PCT/EP2007/051675

- 76 -

fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_d6Des(Pir) was verified by sequencing.

- 5 In the next step, the plasmid pUC19-Cnl1_d6Des(Pir) was used for cloning the *Thraustochytrium* ssp. $\Delta 5$ -desaturase (WO02/26946). To this end, the *Thraustochytrium* ssp. $\Delta 5$ -desaturase was amplified using the following PCR primers:

D5Des(Tc) 5': gggatccatgggcaagggcagcgagggccg

D5Des(Tc) 3': ggcgccgacaccaagaagcaggactgagatatac

10

Composition of the PCR mix (50 μ l):

5.00 μ l template cDNA

5.00 μ l 10x buffer (Advantage polymerase) + 25mM MgCl₂

5.00 μ l of 2mM dNTP

- 15 1.25 μ l of each primer (10 pmol/ μ l)

0.50 μ l of Advantage polymerase (Clontech)

PCR reaction conditions:

Annealing temperature: 1 min 55°C

- 20 Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

Number of cycles: 35

- 25 The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme BamHI and then for 2 hours at 37°C with the restriction enzyme *EcoRV*. The vector pUC19-

WO 2007/096387

PCT/EP2007/051675

- 77 -

Cnl1_d6Des(Pir) was incubated for 2 hours at 37°C with the restriction enzyme BamHI and for 2 hours at 37°C with the restriction enzyme EcoRV. Thereafter, the PCR product and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_d6Des(Pir)_d5Des(Tc) was verified by sequencing.

In the next step, the plasmid pUC19-Cnl1_d6Des(Pir)_d5Des(Tc) was used for cloning the *Physcomitrella patens* $\Delta 6$ -elongase (WO01/59128), for which purpose the latter was amplified using the following PCR primers:

D6Elo(Pp) 5': gcggccgcatggaggtcgtggagagattctacggtg

D6Elo(Pp) 3': gcaaaaggagctaaaactgagtgatctaga

Composition of the PCR mix (50 μ l):

5.00 μ l template cDNA

5.00 μ l 10x buffer (Advantage polymerase) + 25mM MgCl₂

5.00 μ l of 2mM dNTP

1.25 μ l of each primer (10 pmol/ μ l)

0.50 μ l of Advantage polymerase (Clontech)

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

Elongation temperature: 2 min 72°C

Number of cycles: 35

WO 2007/096387

PCT/EP2007/051675

- 78 -

The PCR product was first incubated for 2 hours at 37°C with the restriction enzyme NotI and then for 2 hours at 37°C with the restriction enzyme XbaI. The vector pUC19-Cnl1_d6Des(Pir)_d5Des(Tc) was incubated for 2 hours at 37°C with the restriction enzyme NotI and for 2 hours at 37°C with the restriction enzyme XbaI. Thereafter, the PCR product and cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of the Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp) was verified by sequencing.

10 The binary vector for the transformation of plants was prepared starting from pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp). To this end, pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp) was incubated for 2 hours at 37°C with the restriction enzyme AscI. The vector pGPTV was treated in the same manner. Thereafter, the fragment from pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp) and the cleaved pGPTV
15 vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pGPTV-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp) was verified by sequencing.

20 A further construct, pGPTV-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co), was used. To this end, the amplification was carried out with the following primers, starting from pUC19-Cnl1C_OCS:

Cnl1_OCS 5': gtcgatcaacgggtccggcggtatagagttg

Cnl1_OCS 3': gtcgatcggacaatcagtaaattgaacggaga

25

Composition of the PCR mix (50 µl):

WO 2007/096387

PCT/EP2007/051675

- 79 -

5.00 µl template cDNA

5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂

5.00 µl of 2mM dNTP

1.25 µl of each primer (10 pmol/µl)

5 0.50 µl of Advantage polymerase (Clontech)

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

10 Elongation temperature: 2 min 72°C

Number of cycles: 35

The PCR product was incubated for 2 hours at 37°C with the restriction enzyme Sall. The vector pUC19 was incubated for 2 hours at 37°C with the restriction enzyme Sall. Thereafter, the PCR product and the cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_OCS was verified by sequencing.

20 In a further step, the *Calendula officinalis* Δ12-desaturase gene (WO01/85968) was cloned into pUC19-Cnl1_OCS. To this end, d12Des(Co) was amplified with the following primers:

D12Des(Co) 5': agatctatgggtgcaggcggtcgaatgc

D12Des(Co) 3': ccatggttaaatcttattacgatacc

25 Composition of the PCR mix (50 µl):

WO 2007/096387

PCT/EP2007/051675

- 80 -

5.00 µl template cDNA

5.00 µl 10x buffer (Advantage polymerase) + 25mM MgCl₂

5.00 µl of 2mM dNTP

1.25 µl of each primer (10 pmol/µl)

5 0.50 µl of Advantage polymerase (Clontech)

PCR reaction conditions:

Annealing temperature: 1 min 55°C

Denaturation temperature: 1 min 94°C

10 Elongation temperature: 2 min 72°C

Number of cycles: 35

The PCR product was incubated for 2 hours at 37°C with the restriction enzyme Bg/II and thereafter for 2 hours at the same temperature with NcoI. The vector pUC19-Cnl1_OCS was
15 incubated in the same manner. Thereafter, the PCR fragment and the cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_D12Des(Co) was
20 verified by sequencing.

The plasmid pUC19-Cnl1_D12Des(Co) and the plasmid pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp) were incubated for 2 hours at 37°C with the restriction enzyme Sall. Thereafter, the vector fragment and the cleaved vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA
25 was purified by means of Qiagen Gel Purification Kit following the manufacturer's instructions. Thereafter, vector and vector fragment were ligated. The Rapid Ligation Kit from

WO 2007/096387

PCT/EP2007/051675

- 81 -

Roche was used for this purpose. The resulting plasmid pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co) was verified by sequencing.

The binary vector for the transformation of plants was prepared starting from pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co). To this end,
5 pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co) was incubated for 2 hours at 37°C with the restriction enzyme *AscI*. The vector pGPTV was treated in the same manner. Thereafter, the fragment from pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co) and the cleaved pGPTV vector were separated by agarose gel electrophoresis and the corresponding DNA fragments were excised. The DNA was purified by means of Qiagen Gel
10 Purification Kit following the manufacturer's instructions. Thereafter, vector and PCR product were ligated. The Rapid Ligation Kit from Roche was used for this purpose. The resulting plasmid pGPTV-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co) was verified by sequencing.

A further example of the use of seed-specific expression constructs is the Napin promoter.
15 Preparation of these expression constructs in the vectors pGPTV or pSUN is described in Wu et al. (2005) Nat. Biotech. 23:1013-1017.

A further vector suitable for plant transformation is pSUN2. This vector was used in combination with the Gateway system (Invitrogen, Karlsruhe) in order to increase the number of expression cassettes present in the vector to more than four. For this purpose, the Gateway
20 cassette A was inserted into the vector pSUN2 in accordance with the manufacturer's instructions, as described below:

The pSUN2 vector (1 µg) was incubated with the restriction enzyme EcoRV at 37° for 1 h. The Gateway cassette A (Invitrogen, Karlsruhe) was then ligated into the cut vector using the Rapid Ligation kit from Roche, Mannheim. The resulting plasmid was transformed into
25 *E. coli* DB3.1 cells (Invitrogen). The isolated plasmid pSUN-GW was then verified by sequencing.

WO 2007/096387

PCT/EP2007/051675

- 82 -

In the second step, the expression cassette was cut out of pUC19-Cnl1_d6Des(Pir)_d5Des(Tc)_D6Elo(Pp)_D12Des(Co) using AscI and ligated into the likewise treated vector pSUN-GW. The plasmid obtained in this way pSUN-4G was used for further gene constructs.

- 5 For this purpose, firstly a pENTR clone was modified in accordance with the manufacturer's instructions (Invitrogen). The plasmid pENTR1A (Invitrogen) was incubated with the restriction enzyme EcoRI at 37° for 1 h and then treated with Klenow enzyme and with a 1 µM dNTP mix for 30 min, and subsequently the AscI adapter (5'-ggcgcgcc; phosphorylated at the 5' end, double-stranded) was ligated into the pENTR1A vector. Genes were inserted as
10 described above stepwise into the Cnl cassette in these modified and transferred via AscI into the pENTR vector, resulting in the pENTR-Cnl vector.

- In a further step, the pSUN-8G construct was prepared. For this purpose, 5' and 3' primers for the genes with the SEQ ID NOs: 1, 3, 5 and 7 with the restriction cleavage sites described above and with the first and in each case last 20 nucleotides of the open reading frame were
15 produced and amplified with the standard conditions (see above) and ligated into the pENTR-Cnl vector, which was subsequently subjected to a recombination reaction with the pSUN-4G vector in accordance with the manufacturer's instructions.

The construct pSUN-8G was prepared in this way and was transformed into Brassica juncea and Brassica napus. The seeds of the transgenic plants were analyzed by gas chromatography.

- 20 A further construct which was used for transformation of B. juncea and B. napus was the construct pSUN-9G. This construct was prepared according to Wu et al. (2005) Nat. Biotech. 23:1013-1017 with the napin promoter. In a modification of Wu et al. 2005, the coding sequence of OtELO2.2 was inserted in the described manner instead of the gene OmELO. The resulting construct pSUN-9G was then transformed into B. juncea and B. napus.

- 25 Example 8: Lipid extraction from plant material

WO 2007/096387

PCT/EP2007/051675

- 83 -

The effect of the genetic modification in plants on the production of a desired compound (such as a fatty acid) can be determined by growing the modified plant under suitable conditions (such as those described above) and analyzing the medium and/or the cellular components for the elevated production of the desired product (i.e. of the lipids or a fatty acid). These analytical techniques are known to the skilled worker and comprise spectroscopy, thin-layer chromatography, various types of staining methods, enzymatic and microbiological methods and analytical chromatography such as high-performance liquid chromatography (see, for example, Ullman, Encyclopedia of Industrial Chemistry, Vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon A. et al. (1987) "Applications of HPLC in Biochemistry" in: Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 17; Rehm et al. (1993) Biotechnology, Vol. 3, Chapter III: "Product recovery and purification", p. 469-714, VCH: Weinheim; Belter, P.A. et al. (1988) Bioseparations: downstream processing for Biotechnology, John Wiley and Sons; Kennedy, J.F., and Cabral, J.M.S. (1992) Recovery processes for biological Materials, John Wiley and Sons; Shaeiwitz, J.A., and Henry, J.D. (1988) Biochemical Separations, in: Ullmann's Encyclopedia of Industrial Chemistry, Vol. B3; Chapter 11, p. 1-27, VCH: Weinheim; and Dechow, F.J. (1989) Separation and purification techniques in biotechnology, Noyes Publications).

In addition to the abovementioned methods, plant lipids are extracted from plant material as described by Cahoon et al. (1999) Proc. Natl. Acad. Sci. USA 96 (22):12935-12940 and Browse et al. (1986) Analytic Biochemistry 152:141-145. The qualitative and quantitative analysis of lipids or fatty acids is described by Christie, William W., Advances in Lipid Methodology, Ayr/Scotland: Oily Press (Oily Press Lipid Library; 2); Christie, William W., Gas Chromatography and Lipids. A Practical Guide - Ayr, Scotland: Oily Press, 1989, Repr. 1992, IX, 307 pp. (Oily Press Lipid Library; 1); "Progress in Lipid Research, Oxford: Pergamon Press, 1 (1952) - 16 (1977) under the title: Progress in the Chemistry of Fats and Other Lipids CODEN.

In addition to measuring the end product of the fermentation, it is also possible to analyze other components of the metabolic pathways which are used for the production of the desired

WO 2007/096387

PCT/EP2007/051675

- 84 -

compound, such as intermediates and by-products, in order to determine the overall production efficiency of the compound. The analytical methods comprise measuring the amount of nutrients in the medium (for example sugars, hydrocarbons, nitrogen sources, phosphate and other ions), measuring the biomass composition and the growth, analyzing the production of conventional metabolites of biosynthetic pathways and measuring gases which are generated during the fermentation. Standard methods for these measurements are described in *Applied Microbial Physiology; A Practical Approach*, P.M. Rhodes and P.F. Stanbury, Ed., IRL Press, p. 103-129; 131-163 and 165-192 (ISBN: 0199635773) and references cited therein.

One example is the analysis of fatty acids (abbreviations: FAME, fatty acid methyl ester; GC-MS, gas liquid chromatography/mass spectrometry; TAG, triacylglycerol; TLC, thin-layer chromatography).

The unambiguous detection for the presence of fatty acid products can be obtained by analyzing recombinant organisms using analytical standard methods: GC, GC-MS or TLC, as described on several occasions by Christie and the references therein (1997, in: *Advances on Lipid Methodology*, Fourth Edition: Christie, Oily Press, Dundee, 119-169; 1998, *Gaschromatographie-Massenspektrometrie-Verfahren* [Gas chromatography/mass spectrometric methods], *Lipide* 33:343-353).

The material to be analyzed can be disrupted by sonication, grinding in a glass mill, liquid nitrogen and grinding or via other applicable methods. After disruption, the material must be centrifuged. The sediment is resuspended in distilled water, heated for 10 minutes at 100°C, cooled on ice and recentrifuged, followed by extraction for one hour at 90°C in 0.5 M sulfuric acid in methanol with 2% dimethoxypropane, which leads to hydrolyzed oil and lipid compounds, which give transmethylated lipids. These fatty acid methyl esters are extracted in petroleum ether and finally subjected to a GC analysis using a capillary column (Chrompack, WCOT Fused Silica, CP-Wax-52 CB, 25 m, 0.32 mm) at a temperature gradient of between 170°C and 240°C for 20 minutes and 5 minutes at 240°C. The identity of the resulting fatty

WO 2007/096387

PCT/EP2007/051675

- 85 -

acid methyl esters must be defined using standards which are available from commercial sources (i.e. Sigma).

Plant material is initially homogenized mechanically by comminuting in a pestle and mortar to make it more amenable to extraction.

- 5 This is followed by heating at 100°C for 10 minutes and, after cooling on ice, by resedimentation. The cell sediment is hydrolyzed for one hour at 90°C with 1 M methanolic sulfuric acid and 2% dimethoxypropane, and the lipids are transmethylated. The resulting fatty acid methyl esters (FAMES) are extracted in petroleum ether. The extracted FAMES are analyzed by gas liquid chromatography using a capillary column (Chrompack, WCOT Fused
10 Silica, CP-Wax-52 CB, 25 m, 0.32 mm) and a temperature gradient of from 170°C to 240°C in 20 minutes and 5 minutes at 240°C. The identity of the fatty acid methyl esters is confirmed by comparison with corresponding FAME standards (Sigma). The identity and position of the double bond can be analyzed further by suitable chemical derivatization of the FAME mixtures, for example to give 4,4-dimethoxyazolin derivatives (Christie, 1998) by means of
15 GC-MS.

Example 9: Use of the optimized $\Delta 5$ -elongase (as depicted in SEQ ID NO: 64) from *Ostreococcus tauri* for constructs for constitutive expression

- Transformation vectors based on pGPTV-35S, a plasmid based on pBIN19-35S (Bevan M. (1984) Nucl. Acids Res. 18:203), were produced for the transformation of plants. For this
20 purpose, firstly an expression cassette consisting of the promoter element CaMV35S (SEQ ID No. 161) and the 35S terminator (SEQ ID NO. 162; Franck, A. et al. (1980) Cell 21 (1): 285-294) was assembled in a pUC vector. This entailed the promoter being inserted via the Sall/XbaI restriction cleavage sites and the terminator via the BamHI/SmaI restriction cleavage sites. In addition, a polylinker with the XhoI cleavage site was attached to the
25 terminator ('triple ligation'). The resulting plasmid pUC19-35S was then employed for cloning PUFA genes. In parallel, the open reading frames of the $\Delta 6$ -desaturase (SEQ ID NO. 1), of the $\Delta 5$ -desaturase (SEQ ID NO. 51) and $\Delta 6$ -elongase (SEQ ID NO. 171) sequences

WO 2007/096387

PCT/EP2007/051675

- 86 -

were inserted via the EcoRV cleavage site into pUC19-35S vectors. The resulting plasmids pUC-D6, pUC-D5, pUC-E6(Tc) were used to construct the binary vector pGPTV-35S_D6D5E6(Tc). For this purpose, the vector pGPTV was digested with the enzyme Sall, the plasmid pUC-D6 was digested with Sall/XhoI, and the correct fragments were ligated.

5 The resulting plasmid pGPTV-D6 was then digested with Sall, the plasmid pUC-D5 was digested with Sall/XhoI, and the correct fragments were ligated. The resulting plasmid pGPTV-D6-D5 was then digested once more with Sall, the plasmid pUC-E6(Tc) with Sall/XhoI, and the correct fragments were ligated. These sequential cloning steps resulted in the binary vector pGPTV-D6D5E6(Tc), which was employed for the transformation.

10 In a further procedure, the sequence of d6Elo(Tp) (SEQ ID NO. 163) was inserted into the vector pUC19-35S instead of the sequence d6Elo(Tc). The resulting plasmid pUC-E6(Tp) was used to prepare the binary vector pGPTV-35S_D6D5E6(Tp).

In a further procedure, the open reading frame of ω 3Des (SEQ ID NO. 193) was cloned into pUC19-35S. The resulting plasmid pUC- ω 3Pi was transferred via Sall/XhoI into the binary
15 vectors pGPTV-D6D5E6(Tc) and pGPTV-D6D5E6(Tp). The resulting vectors pGPTV-D6D5E6(Tc) ω 3Pi and pGPTV-D6D5E6(Tp) ω 3Pi were employed for the plant transformation.

In a further procedure, the open reading frame of the optimized Δ 5-elongase from *Ostreococcus tauri* (SEQ ID No. 64) and the open reading frame of the Δ 4-desaturase from
20 *Thraustochytrium* sp. (SEQ ID No. 77) was cloned into pUC19-35S. The resulting plasmids pUC-E5 and pUC-D4 were then transferred via Sall/XhoI in accordance with the above statements into the vector pGPTV-D6D5E6(Tp) ω 3Pi. The resulting vector pGPTV-D6D5E6(Tp) ω 3PiE5D4 was employed for the plant transformation.

All the binary vectors were transformed into *E. coli* DH5 α cells (Invitrogen) in accordance
25 with the manufacturer's instructions. Positive clones were identified by PCR, and plasmid DNA was isolated (Qiagen Dneasy).

WO 2007/096387

PCT/EP2007/051675

- 87 -

Example 10: Transformation of the constitutive binary vectors into plants

a) Generation of transgenic *Brassica napus* and *Brassica juncea* plants. The protocol for the transformation of oilseed rape plant was used (modification of Moloney et al. (1992) Plant Cell Reports 8:238-242)

5 The binary vector pGPTV-D6D5E6(Tp) ω 3PiE5D4 was transformed in *Agrobacterium tumefaciens* C58C1:pGV2260 (Deblaere et al. (1984) Nucl. Acids. Res. 13: 4777-4788). A 1:50 dilution of an overnight culture of a positively transformed agrobacterial colony in Murashige-Skoog medium (Murashige and Skoog (1962) Physiol. Plant. 15: 473) supplemented with 3% sucrose (3MS medium) was used for the transformation of

10 *Orychophragmus violaceus*. Petioles or hypocotyls of freshly germinated sterile plants (in each case approx. 1 cm²) were incubated with a 1:50 agrobacterial dilution for 5-10 minutes in a Petri dish. This is followed by 3 days of coincubation in the dark at 25°C on 3MS medium supplemented with 0.8% Bacto agar. Thereafter, the cultivation was continued with 16 hours light/8 hours dark and a weekly rhythm on MS medium supplemented with 500 mg/l Claforan

15 (cefotaxime-sodium), 50 mg/l kanamycin, 20 μ M benzylaminopurine (BAP) and 1.6 g/l glucose. Growing shoots were transferred to MS medium supplemented with 2% sucrose, 250 mg/l Claforan and 0.8% Bacto agar. If no roots had developed after three weeks, 2-indolebutyric acid was added to the medium as growth hormone for rooting.

Regenerated shoots were obtained on 2MS medium with kanamycin and Claforan, then, after

20 rooting, transferred into soil and, after cultivation, grown for two weeks in a controlled-environment cabinet or in the greenhouse, allowed to flower, mature seeds were harvested and analyzed for elongase expression such as Δ 6-elongase activity or for Δ 5- or Δ 6-desaturase activity by means of lipid analyses. In this manner, lines with elevated contents of polyunsaturated C20- and C22-fatty acids were identified.

25 b) Generation of transgenic *Orychophragmus violaceus* plants

WO 2007/096387

PCT/EP2007/051675

- 88 -

The protocol for the transformation of oilseed rape plants was used (modification of Moloney et al. (1992) Plant Cell Reports 8:238-242) as described under a).

To generate transgenic plants, the binary vector pGPTV-D6D5E6(Tp) ω 3PiE5D4 was transformed into *Agrobacterium tumefaciens* C58C1:pGV2260 (Deblaere et al. (1984) Nucl. Acids. Res. 13: 4777-4788). A 1:50 dilution of an overnight culture of a positively transformed *Agrobacterium* colony in Murashige-Skoog medium (Murashige and Skoog (1962) Physiol. Plant, 15: 473) with 3% sucrose (3MS medium) was used to transform *Orychophragmus violaceus*. Petioles or hypocotyls of freshly germinated sterile plants (each about 1 cm²) were incubated with a 1:50 agrobacterial dilution in a Petri dish for 5-10 minutes. This is followed by coincubation on 3MS medium with 0.8% Bacto agar in the dark at 25°C for 3 days. The cultivation was then continued with 16 hours light/8 hours dark and in a weekly rhythm on MS medium with 500 mg/l Claforan (cefotaxime sodium), 15 mg/l kanamycin, 20 μ M benzylaminopurine (BAP) and 1.6 g/l glucose. Growing shoots were transferred to MS medium with 2% sucrose, 250 mg/l Claforan and 0.8% Bacto agar. If no roots had developed after three weeks, 2-indolebutyric acid was added to the medium as growth hormone for rooting.

Regenerated shoots were obtained on 2MS medium with kanamycin and Claforan and, after rooting, transferred to soil and, after cultivation, grown for two weeks in a controlled environment cabinet or in a greenhouse, allowed to flower, and mature seeds were harvested and examined by lipid analyses for elongase expression such as Δ -6-elongase activity or Δ -5- or Δ -6-desaturase activity. Lines with increased contents of polyunsaturated C20 and C22 fatty acids were identified in this way.

c) Transformation of *Arabidopsis thaliana* plants

The protocol of Bechthold et al. (1993) C.R. Acad. Sci. Ser. III Sci. Vie. 316: 1194-1199 was used.

To generate transgenic plants, the generated binary vector pGPTV-D6D5E6(Tp) ω 3PiE5D4 was transformed into *Agrobacterium tumefaciens* C58C1:pMP90 (Deblaere et al. (1984)

WO 2007/096387

PCT/EP2007/051675

- 89 -

Nucl. Acids. Res. 13: 4777-4788) and, in accordance with the protocol of Bechthold et al. (1993), flowers of *Arabidopsis thaliana* cv. Columbia 0 were dipped in an agrobacterial solution with OD₆₀₀=1.0. The procedure was repeated again two days later. Seeds from these flowers were then placed on agar plates with ½ MS, 2% sucrose and 50 mg/l kanamycin.

5 Green seedlings were then transferred to soil.

Example 11: Analysis of plant material of transgenic *Orychophragmus* or *Arabidopsis* plants

Extraction of leaf material of transgenic *Orychophragmus violaceus* and *Arabidopsis thaliana* plants transformed with pGPTV-D6D5E6(Tp)ω3PiE5D4 and the gas chromatography analysis was carried out as described in example 8. Table 2 shows the results of the analyses.

10 The various fatty acids are indicated in percent by weight. It was possible to show that long-chain polyunsaturated fatty acids were synthesized by both different plant species. It was surprisingly possible with the optimized sequence of the Δ5-elongase (as depicted in SEQ ID NO: 64) from *Ostreococcus tauri* to obtain a distinctly higher yield of DHA than reported for example by Robert et al. (2005) Functional Plant Biology 32: 473-479 for *Arabidopsis*
15 *thaliana* with 1.5% DHA. It was possible for the first time to achieve a synthesis of long-chain polyunsaturated fatty acids for *Orychophragmus violaceus*.

Example 12: Analysis of seeds of transgenic *Brassica juncea* lines

Extraction of seeds of transgenic *Brassica juncea* plants transformed with pSUN-9G, and the gas chromatography analysis was carried out as described in example 8. Table 6 shows the
20 results of the analyses. The various fatty acids are indicated in percent area. As in Wu et al. 2005 it was possible to show the synthesis of long-chain polyunsaturated fatty acids (PUFA). Surprisingly, the use of the modified elongase sequence OtELO2.2 such as the nucleic acid sequence described by SEQ ID NO: 64 resulted in a drastic increase in the content of C22 fatty acids. In total, the seed oil contained about 8% by weight % polyunsaturated C22 fatty
25 acids. Specifically, the content of the fatty acid docosahexaenoic acid (DHA) in the seed oil was 1.9% by weight %, representing an increase by a factor of 10 compared with Wu et al. 2005.

WO 2007/096387

PCT/EP2007/051675

- 90 -

Example 13: Detailed analysis of the lipid classes and position analysis of leaf material from *O. violaceus*

About 1 g of leaf tissue was heated in 4 ml of isopropanol at 95°C for 10 minutes, homogenized by Polytron and shaken after addition of 1.5 ml of chloroform. The samples
5 were centrifuged, the supernatant was collected, and the pellet was extracted again with isopropanol:chloroform 1:1 (v/v). The two extracts were combined, dried and dissolved in chloroform. The lipid extract was prefractionated on a silica prepsep column (Fisher Scientific, Nepean, Canada) into neutral lipids, glycolipids and phospholipids, eluting with chloroform:acetic acid 100:1 (v/v), acetone:acetic acid 100:1 (v/v) and
10 methanol:chloroform:water 100:50:40 (v/v/v), respectively. These fractions were further fractionated on silica G-25 thin-layer chromatography plates (TLC; Macherey-Nagel, Düren, Germany). Neutral lipids were developed with hexane:diethyl ether:acetic acid (70:30:1), glycolipids with chloroform:methanol:ammonia (65:25:4 v/v/v) and phospholipids with chloroform:methanol:ammonia:water (70:30:4:1 v/v/v/v). The individual lipid classes were
15 identified after spraying with primulin under UV light, removed by scraping off the plates and either used for direct transmethylation or extracted by a suitable solvent for further analysis.

It was possible by the disclosed methods for the various lipid classes (neutral lipids, phospholipids and galactolipids) to be fractionated and analyzed separately. The glycolipids were additionally examined for the position of the individual fatty acids.

20 a) Regiospecific analysis of the triacylglycerides (TAG)

Three to five mg of the TLC-purified TAG were dried under nitrogen in a glass tube, resuspended in aqueous buffer by brief ultrasound treatment (1 M Tris pH 8; 2.2% CaCl₂ (w/v); 0.05% bile salts (w/v)) and incubated at 40°C for 4 minutes. After addition of 0.1 ml of a solution of pancreatic lipase (10 mg/ml in water), the samples were vigorously vortexed for
25 3 minutes, and the digestion was stopped by adding 1 ml of ethanol and 1.5 ml of 4 M HCl. The partly digested TAGs were extracted twice with diethyl ether, washed with water, dried and dissolved in a small volume of chloroform. Monoacylglycerols (MAG) were separated

WO 2007/096387

PCT/EP2007/051675

- 91 -

from the free fatty acids and undigested TAGs on a TLC plate as described above for neutral lipids. The point corresponding to the MAGs was analyzed by GC and represented the sn-2 position of the TAGs. The distribution of the fatty acids to the remaining sn-1 and sn-3 positions was calculated by the following formula: $sn-1 + sn-3 = (TAG \times 3 - MAG)/2$.

- 5 This position analysis of the triacylglycerides revealed in this case that EPA and DHA are present in similar concentrations in the sn-2 and sn-1/3 positions, while ARA is to be found overall only in small amounts in the triacylglycerides, and here mainly in the sn-2 position (Tab. 3).

b) Stereospecific analysis of phospholipids

- 10 Fractionated and extracted phosphatidylglycol (PG), phosphatidylethanolamine (PE) and phosphatidylcholine (PC) were dried under N₂ and resuspended in 0.5 ml of borate buffer (0.5M, pH 7.5, containing 0.4 mM CaCl₂). After a brief ultrasound treatment, 5U of phospholipase A2 from the venom of *Naja mossambica* (Sigma P-7778) and 2 ml of diethyl ether were added and the samples were vortexed at room temperature for 2 hours. The ether
15 phase was dried, the digestion was stopped with 0.3 ml of 1M HCl, and the reaction mixture was extracted with chloroform:methanol (2:1 v/v). The digested phospholipids were separated by TLC in chloroform:methanol:ammonia:water (70:30:4:2 v/v/v/v) and points which corresponded to the liberated free fatty acids and lysophospholipids were removed by scraping and directly transmethyated.
- 20 Positional analysis of the phospholipids showed an accumulation of EPA and DHA in the sn-2 position of phosphatidylcholine (PC), while DHA was similarly distributed in sn-1 and sn-2 position in phosphatidylethanolamine (PE). Only traces of, or no, ARA was to be found in both phospholipids (Tab. 4). The concentrations of EPA and DHA in phosphatidylglycerol were lower than in the other investigated phospholipids, with accumulation in the sn-2
25 position also to be observed in this lipid class (Tab. 4, PG).

c) Stereospecific analysis of glycolipids

WO 2007/096387

PCT/EP2007/051675

- 92 -

The galactolipids were investigated as a further polar lipid class. Galactolipids are found in the membranes of plastids and form the main components there.

TLC-purified monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG) were dried under nitrogen and dissolved in 0.5 ml of diethyl ether. Then 25 units of the lipase from *Rhizopus arrhizus* (Sigma 62305), resuspended in 2 ml of borate buffer (50 mM, pH 7.5 containing 2 mM CaCl_2), were added, and the samples were vortexed at room temperature for 2 hours. The ether phase was dried and the digestion was stopped by adding 0.3 ml of 1M HCl, and the lipids were extracted with 4 ml of chloroform:methanol (2:1 v/v). After drying, the digested galactolipids were in a small volume of chloroform:methanol (2:1 v/v) and developed twice on a precoated silica TLC plate, firstly with chloroform:methanol: ammonia:water (70:30:4:1 v/v/v/v) to about two thirds the height of the plate, followed by complete development in hexane:diethyl ether:acetic acid (70:30:1). The points which corresponded to the liberated free fatty acids and the lysogalactolipids were identified after spraying with primulin, scraped off and transmethyalted directly for GC analysis.

It was possible to find VLCPUFA in these lipids too, with an accumulation of EPA in the sn-2 position being observed. DHA was to be found only in the digalactodiacylglycerols (DGDG) and was undetectable in the monogalactodiacylglycerols (MGDG) (Table 5). The distribution of VLCPUFA in galactolipids, a compartment in which these fatty acids were not expected, shows the dynamics of the synthesis and the later transformation. VLCPUFA in polar lipids are of particular nutritional value because they can be absorbed better in the intestines of mammals than the neutral lipids.

WO 2007/096387

PCT/EP2007/051675

- 93 -

Table 1: Test of the optimized sequences of pOTE1.1 and pOTE2.1 in yeast. The conversion rates were determined in accordance with the substrate conversions. A distinct rise in activity was achievable with the optimized sequence in plasmid pOTE2.2.

5

Conversion rates of the <i>Ostreococcus tauri</i> elongases					
	Genes	Substrate Product	GLA 20:3	ARA 22:4	EPA 22:5
pOTE1.1	d6Elongase(Ot)		21.1		
pOTE1.2	d6Elongase(Ot)_opt		25.6		
pOTE2.1	d5Elongase(Ot)			7.3	35.9
pOTE2.2	d5Elongase(Ot)_opt			32.7	63.1

WO 2007/096387

PCT/EP2007/051675

- 94 -

Table 2: Gas chromatographic analysis of leaf material of *Orychophragmus violaceus* and *Arabidopsis thaliana*. The individual fatty acids are indicated in percent area.

Fatty acid composition of leaf material of <i>Orychophragmus violaceus</i>											
Fatty acids	16:0	16:3	18:1	18:2	GLA	18:3	18:4	ARA	EPA	DPA	DHA
Control	20.9	8.5	3.3	16.0	0.0	47.4	0.0	0.0	0.0	0.0	0.0
Transgene	21.3	8.2	5.2	5.2	4.2	23.1	5.0	0.6	13.5	2.7	4.5

Fatty acid composition of leaf material of <i>Arabidopsis thaliana</i>											
Fatty acids	16:0	16:3	18:1	18:2	GLA	18:3	18:4	ARA	EPA	DPA	DHA
Control	12.8	10.0	3.5	14.2	0.0	54.6	0.0	0.0	0.0	0.0	0.0
Transgene	19.3	8.5	5.0	4.6	6.4	31.0	4.4	0.0	6.3	1.5	6.3

Table 3: Regiospecific analysis of the triacylglycerides from leaf material from transgenic O. violaceus plants.

TAG	16:0	18:0	18:1n-7	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3	22:5n-3	22:6n-3
wt	25.12	3.03	5.06	18.53		44.72							
sn-2	1.42	0.76	6.79	27.62		62.03							
sn-1+3	36.97	4.17	4.19	13.98		36.07							
Transgene	22.63	3.12	3.46	2.35	9.51	6.37	13.03	0.74	0.83	3.87	24.96	2.22	4.15
sn-2	1.62	0.64	8.33	5.15	16.21	10.88	19.84	0.17	1.38	1.98	24.82	3.27	3.02
sn-1+3	33.13	4.36	1.02	0.96	6.16	4.11	9.63	1.02	0.55	4.90	25.03	1.69	4.72

Table 4: Stereospecific analysis of the phospholipids from leaf material from transgenic O. violaceus plants.

PG	16:0	16:1	18:0	18:1n-9	18:1n-7	18:2n-9	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3	22:5n-3	22:6n-3
WT	27.96	20.04	4.11	2.89	0.90		21.82	0.00	21.56							
sn-1	17.26	0.53	2.61	3.82	1.91		39.01	0.00	34.44							
sn-2	38.66	39.56	5.62	1.96	0.00		4.62	0.00	8.69							
Transgene	27.15	24.70	3.08	4.62	1.20	0.00	15.15	1.53	17.94	1.40	0.00	0.00	0.45	2.18	0.10	0.58
sn-1	21.16	3.61	4.23	7.52	2.14		27.40	0.50	31.57	0.81			0.38	1.24	0.00	0.33
sn-2	33.15	45.79	1.94	1.71	0.27		2.90	2.57	4.30	2.00			0.51	3.13	0.27	0.83

PE	16:0	16:1	18:0	18:1n-9	18:1n-7	18:2n-9	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3	22:5n-3	22:6n-3
WT	37.49	0.00	6.62	4.35	1.37		19.28		29.95							
sn-1	54.22	0.00	7.74	3.39	3.42		12.64		13.71							
sn-2	20.77	0.00	5.51	5.31	0.00		25.93		46.18							
Transgene	31.78	0.81	5.84	3.08	2.20	0.85	5.57	11.25	11.34	7.38	0.00	0.00	2.88	9.41	1.90	4.90
sn-1	50.17	0.33	10.86	3.22	4.94	0.35	2.63	3.27	3.59	2.31	0.56		4.42	6.18	0.38	4.19
sn-2	13.40	1.29	0.83	2.95	0.00	1.35	8.50	19.23	19.10	12.45	0.00		1.34	12.64	3.41	5.61

PC	16:0	16:1	18:0	18:1n-9	18:1n-7	18:2n-9	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3	22:5n-3	22:6n-3
WT	27.67	0.84	6.38	8.56	1.80		21.75		33.01							
sn-1	48.05	0.44	8.65	5.05	3.41		14.52		18.04							
sn-2	7.28	1.24	4.11	12.06	0.18		28.97		47.98							
Transgene	21.00	0.00	8.01	10.02	2.86	1.25	3.77	11.63	5.60	12.11	0.50	0.00	4.34	11.16	3.76	3.70
sn-1	45.35	0.00	14.71	5.08	5.70	0.31	3.23	3.09	4.58	2.65	0.61	0.08	4.01	8.32	0.41	1.18
sn-2	3.36	0.00	1.30	14.96	0.02	2.20	4.31	20.18	6.62	21.56	0.38	0.00	4.66	13.99	7.12	6.22

Table 5: Stereospecific analysis of the galactolipids from leaf material from transgenic O. violaceus plants.

MGDG	16:0	16:1	16:2	16:3	18:0	18:1n-9	18:1n-7	18:2n-9	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3
wt	264	013	123	3072	033	036	026	381		6052						
sn-1	000	005	000	7.11	035	031	041	460		8730						
sn-2	534	021	255	5434	031	039	012	301		3374						
tr	416	020	108	3381	093	073	052	003	188	4482	273	004	030	050	508	
sn-1	122	029	054	479	151	115	093	280	014	8019	000	008	017	087	386	
sn-2	7.11	011	161	6282	034	031	011	047	362	946	548	000	043	014	631	

DG DG	16:0	16:1	16:2	16:3	18:0	18:1n-9	18:1n-7	18:2n-9	18:2n-6	18:3n-6	18:3n-3	18:4n-3	20:3n-6	20:4n-6	20:4n-3	20:5n-3
wt	17.67	019	038	215	161	051	094	556		7071						
sn-1	1684	025	050	252	221	055	175	607	000	6874						
sn-2	1850	012	027	178	101	046	013	505		7268						
tr	1850	000	000	262	284	136	139	000	355	5466	000	000	000	218	536	
sn-1	2274	017	023	048	455	171	232	922	023	5606	027	000	000	036	123	
sn-2	1427	000	000	477	112	100	046	333	688	5326	000	000	000	401	949	

Table 6: Gas chromatographic determination of the fatty acids from seeds of transgenic Brassica juncea plants transformed with the construct pSUN-9G in percent by weight. WT describes the unmodified wild-type control.

	Lipid Profile(%)						
	16:0	18:0	18:1	18:2	18:3	18:4	20:0
BJ223_PUFA184_MKP71_581A	4.4	3.0	22.5	16.9	27.0	3.2	0.6
BJ223_PUFA184_MKP71_581A	4.7	3.9	17.9	10.6	29.5	4.0	0.9
BJ223_PUFA184_MKP71_581A	4.4	3.0	18.9	13.8	30.5	3.2	0.7
BJ223_PUFA184_MKP71_581A	4.6	3.3	20.5	13.2	29.8	3.3	0.8

20:3 (8,11,14)	20:3 (11,14,17)	20:4 (ARA) (5,8,11,14)	20:4 (ETE4) (8,11,14,17)	20:5 (EPA) (5,8,11,14,17)	22:1	22:4	22:5	22:6
1,1	0.5	3.1	0.6	4.6	0.0	1.5	2.0	1.5
2,0	0.9	4.2	1.0	4.1	0.0	3.1	3.5	1.3
1,3	0.7	4.1	0.5	4.5	0.0	2.7	2.8	1.6
1,4	0.6	3.0	0.6	4.4	0.0	2.4	2.5	1.6

CLAIMS

1. A process for producing eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in a transgenic plant, comprising the provision in the plant of
 - 5 • at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 2;
 - at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 172;
 - 10 • at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 52; and
 - at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 65 and
 - 15 • insofar as the process aims at producing docosahexaenoic acid (DHA), at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 4$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 78,
 - 20 where the nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity is modified by comparison with the nucleic acid sequence in the organism from which the sequence is obtained in that it is adapted to the codon usage in the host plant species.
 - 25
2. The process as claimed in claim 1, where the nucleic acid sequence which codes for the polypeptide having a $\Delta 5$ -elongase activity is adapted at least to the codon usage in oilseed rape, soybean and/or flax.
3. The process as claimed in either of claims 1 or 2, where the nucleic acid sequence

which codes for the polypeptide having a $\Delta 5$ -elongase activity is adapted taking into account the natural frequency of individual codons.

4. The process as claimed in any one of claims 1 to 3, where the modified nucleic acid sequence is the nucleic acid sequence indicated in SEQ ID NO. 64.
5. The process as claimed in any one of claims 1 to 4, where the nucleic acid sequences are expressed under the control of a seed-specific promoter.
6. The process as claimed in claim 5, where the promoter is the USP, vicilin, napin, Glp, SBP, peroxiredoxin, legumin, Fad3, conlinin or oleosin promoter.
7. The process as claimed in claim 6, where the content of polyunsaturated C22 fatty acids in the seed oil of the transgenic plant is 5% by weight or more of the seed oil content.
8. The process as claimed in any one of claims 1 to 7, where additionally one or more nucleic acid sequences coding for a polypeptide having the activity of an $\omega 3$ -desaturase and/or a $\Delta 4$ -desaturase are provided in the plant.
9. The process as claimed in claim 8, where the content of docosaheptaenoic acid in the seed oil of the transgenic plant is 1% by weight or more of the seed oil content.
10. The process as claimed in any one of claims 1 to 9, where the eicosapentaenoic acid, docosapentaenoic acid and/or docosaheptaenoic acid is present in the plant mainly bound as ester in phospholipids or triacylglycerides.
11. The process as claimed in any one of claims 1 to 10, where the plant is an oil-producing plant.
12. The process as claimed in claim 11, wherein the oil-producing plant is *Brassica napus*, *Brassica juncea* or *Glycine max*.
13. The process as claimed in any one of claims 1 to 12, further comprising obtaining the

eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid in the form of their oils, lipids or free fatty acids from the plant.

14. A recombinant nucleic acid molecule comprising:
- a) one or more copies of at least one promoter which is active in plant cells,
 - 5 b) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 2,
 - c) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -desaturase activity and having an amino acid identity of at least 90% to
10 SEQ ID NO: 52,
 - d) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 6$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 172,
 - 15 e) at least one nucleic acid sequence which codes for a polypeptide having a $\Delta 5$ -elongase activity and having an amino acid identity of at least 90% to SEQ ID NO: 65 and which is modified by comparison with the nucleic acid sequence in the organism from which the sequence is obtained by being adapted to the codon usage in one or more plant species, and
 - f) one or more copies of at least one terminator sequence.
- 20 15. The recombinant nucleic acid molecule as claimed in claim 14, further comprising a nucleic acid sequence coding for a polypeptide having a $\Delta 4$ -desaturase activity and having an amino acid identity of at least 90% to SEQ ID NO: 78.
16. The recombinant nucleic acid molecule as claimed in claim 14 or 15, where the modified nucleic acid sequence is the nucleic acid sequence indicated in SEQ ID

NO: 64.

17. The recombinant nucleic acid molecule as claimed in any one of claims 14 to 16, additionally comprising one or more nucleic acid sequence/s coding for a polypeptide having the activity
 - 5 a) of an ω 3-desaturase
 - b) of a Δ 4-desaturase which is mandatory for the production of DHA or
 - c) of an ω 3-desaturase and a Δ 4-desaturase which is mandatory for the production of DHA.
- 10 18. A transgenic plant cell, wherein the plant cell is obtained from oilseed rape, soybean or flax, comprising the recombinant nucleic acid molecule as defined in any one of claims 14 to 17.
- 15 19. A process for producing a transgenic plant which produces eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA) and/or docosahexaenoic acid (DHA), comprising introducing into a plant the recombinant nucleic acid molecule as defined in any one of claims 14 to 16 and expressing the same therein.
20. The process according to claim 19, wherein the recombinant nucleic acid molecule additionally comprises one or more nucleic acid/s as defined in claim 17.
21. The process according to claim 19 or 20, wherein the plant is an oil-producing plant being *Brassica napus*, *Brassica juncea* or *Glycine max*.
- 20 22. A seed cell comprising the recombinant nucleic acid molecule as defined in any one of claims 14 to 16.
23. The seed cell according to claim 22, wherein the recombinant nucleic acid molecule additionally comprises one or more nucleic acid molecule/s as defined in claim 17.

24. The seed cell according to claim 22 or 23, wherein the seed cell is obtained from an oil producing plant being *Brassica napus*, *Brassica juncea* or *Glycine max*.
25. Use of the transgenic plant cell as defined in claim 18, the transgenic plant as produced by the process of any one of claims 19 to 21, or the transgenic seed cell as defined in any one of claims 22 to 24, for the production of an oil, a lipid, a fatty acid and/or a fatty acid composition.
26. The use according to claim 25, wherein eicosapentaenoic acid, docosapentaenoic acid and/or docosahexaenoic acid is/are produced.