




AU9221801

(12) PATENT ABRIDGMENT (11) Document No. AU-B-21801/92
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 654249

(Modified Examination)


- (54) Title
SUBSTITUTED SILYL ALCOHOLS
- International Patent Classification(s)
(51)^s **C07F 007/08**
- (21) Application No. : **21801/92** (22) Application Date : **05.06.92**
- (87) PCT Publication Number : **WO92/22557**
- (30) Priority Data
- (31) Number (32) Date (33) Country
712302 07.06.91 US UNITED STATES OF AMERICA
- (43) Publication Date : **12.01.93**
- (44) Publication Date of Accepted Application : **27.10.94**
- (71) Applicant(s)
ABBOTT LABORATORIES
- (72) Inventor(s)
JOSEPH E CELEBUSKI
- (74) Attorney or Agent
SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001
- (57) Claim

1. An alcoholic compound of the formula:



wherein R_1 , R_2 and R_3 are independently selected from the group consisting of aryl, substituted aryl, substituted aryl, and substituted or unsubstituted sterically bulky alkyl; and n is an integer from 2 to about 20.

9. An ester compound of the formula:



wherein R_1 , R_2 and R_3 are independently selected from the group consisting of aryl, substituted aryl, and substituted or unsubstituted sterically bulky alkyl; R_4 is lower alkyl; and n is an integer from 2 to about 20.

OPI DATE 12/01/93 APPLN. ID 21801/92
AOJP DATE 11/03/93 PCT NUMBER PCT/US92/04723



AU9221801

(PCT)

(51) International Patent Classification ⁵ : C07F 7/08, 7/18	A1	(11) International Publication Number: WO 92/22557 (43) International Publication Date: 23 December 1992 (23.12.92)
(21) International Application Number: PCT/US92/04723 (22) International Filing Date: 5 June 1992 (05.06.92) (30) Priority data: 712,302 7 June 1991 (07.06.91) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD D-377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventor: CELEBUSKI, Joseph, E. ; 5080 Fox Lane, Gurnee, IL 60031 (US). (74) Agents: GORMAN, Edward, Hoover, Jr. et al.; Abbott Laboratories, CHAD 377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> 654249
(54) Title: SUBSTITUTED SILYL ALCOHOLS		
(57) Abstract <p>Novel silyl alcohols having bulky substituents bonded to the silicon, and the silyl group attached to a carbon include the preferred 2-silyl-ethan-1-ols. A method for synthesizing silyl substituted alcohols includes hydrosilation of a vinylic ester, especially vinyl acetate, followed by hydrolysis in mild base. The silyl alcohols are useful in preparing phosphorylating reagents for phosphorylating an oligonucleotide. The phosphorylated intermediate bearing the silyl group may be separated from failure product on the basis of bulky substituents on the silyl protecting group, which is later removed, e.g. by fluoride ion.</p>		

SUBSTITUTED SILYL ALCOHOLS

The invention relates generally to silyl alcohols, their synthesis, and their use. More specifically, the invention relates to specific silyl alcohols and to a method for synthesizing silyl alcohols having the silicon atom bonded to a carbon rather than to the 5 hydroxyl. The invention also relates to reagents and methods for phosphorylating oligonucleotides, and to intermediate compounds and methods useful for purification of phosphorylated oligonucleotides.

This application is related to co-owned and co-pending applications Ser. Nos. 07/712,001 and 07/712,020 filed concurrently herewith, each of which is incorporated 10 herein by reference.

Background

Chemically synthesized oligonucleotides have been used in hybridization assays for some time, and by now are fairly routine. However, for uses which imitate biological processes, e.g., hybridizations of nucleic acid probes on a template followed by ligation, 15 the normal 5' hydroxyl terminus must be converted to a phosphate to provide the proper substrate for a ligase. Methods of phosphorylating include enzymatic and synthetic as described below. The present invention describes a particular synthetic method, wherein silyl substituted alcohols are useful reagents.

Synthesis of silyl substituted alcohols has been previously achieved by oxidation of 20 organoboranes. The organoboranes are in turn prepared by the Grignard reaction or by hydrocarbon of vinyl-and allyl-silanes. This technique is described in Kumada, et al. *J. Organometal. Chem.* 6:490-495 (1966) and Seyferth, *J. Am. Chem. Soc.* 81:1844 (1959). This technique is useful only when the requisite vinyl or allylsilanes can be synthesized or obtained commercially. However, if the desired vinyl silane is commercially unavailable 25 or difficult to synthesize this method is not useful.

Alpha silyl esters have been prepared by reacting a chlorosilane and an alpha-bromo ester with zinc under Reformatsky conditions. See Feesenden, et al. *J. Org. Chem.* 32:3535 (1967).

An important drawback of these synthesis methods is the side reactions which can 30 occur leading to undesirable products and decreasing the yields. In conventional processes for hydrolyzing silyl substituted esters to the corresponding silyl substituted alcohol, a carbanion intermediate is generally formed. With β -silyl substituted alcohols, fragmentation to the silanol and an olefin can occur; with alpha silyl substituted alcohols, a Brook rearrangement to



give a silyl protected ether will occur. Thus, in these carbanion intermediates there is a strong tendency for an elimination reaction whereby the silicon atom shifts to the oxygen atom to form the R_3SiOH byproduct. This tendency is especially pronounced when the reaction is performed in strong base and when

5 groups substituted on the silicon are particularly bulky.

Hydrosilation, the addition of H and silyl compounds across the double bond of an olefin, has also been described in the literature. See Collman, et al. **Principles and Applications of Organotransition Metal Chemistry**, University Science Books (1980) p. 384-389 and Pegram, et al. *Carbohydrate*

10 *Research* 184:276 (1988). In a particularly relevant hydrosilation reaction, Salimgareeva, et al., *Zh. Obshch. Khim* 48(4):930-31 (1978)(Russian) (see also C.A. 89:146961y) report hydrosilation of vinyl acetate with dimethylsilane. This reaction resulted in two silyl substituted products: a monoacetate and a diacetate. The reference fails to describe synthesis of any silyl alcohol or the use

15 or synthesis of any bulky silyl substituted compound.

Honda, et al. *Tetrahedron Letters*, 22(22): 2093-2096 (1981) describe a β -silyl substituted ethanol wherein the silyl group bears two phenyl and one methyl substituent. Honda, et al. used this compound to prepare a phosphorylating agent which places a protected phosphate group between

20 nucleotides in oligonucleotide synthesis. The substituted silyl protecting group can be removed to give a silyl fluoride compound, ethylene and the phosphate. The substituted silyl ethanol was obtained by reduction of the bisphenylmethyl silyl acetate with $LiAlH_4$ according to a modification of the procedure of Gerlach, *Helv Chim. Acta*, 60:3039 (1977).

25 Other silyl substituted ethanols have been described in the literature, but primarily include alkyl substituted silyl groups. Examples of such silyl ethanols and their literature citations are found in the following table.

TABLE 1			
$\begin{array}{c} \text{R} \\ \\ \text{R}' - \text{Si} - \text{CH}_2 - \text{CH}_2 - \text{O H} \\ \\ \text{R}'' \end{array}$			
= GENERAL STRUCTURE			
<u>R</u>	<u>R'</u>	<u>R''</u>	<u>Literature Citation</u>
isopropyl	isopropyl	isopropyl	CA111(11):97352n
methyl	methyl	propenyl	CA105(13):115112s CA108(17):150554w
methyl	n-butyl	n-butyl	CA88(3):23391j; CA83(11):97563k; CA78(15):93640g; & CA78(13):84526x
methyl	methyl	t-butyl	CAOLD (prior to 1967)
ethyl	ethyl	ethyl	CA111(11):973552n; CA98(9):72207u; CA87(15):117488c; CA85(21):154709e; CA80(11):59132z; & CA77(18):120049a
propyl	propyl	propyl	CA103(19):160573n
phenyl	phenyl	methyl	Honda, et al., <i>Tetrahedron Letters</i> , 22(22):2093-2096 (1981)

Triphenyl silane (not the alcohol) has been described by Lesage, et al. *J. Org. Chem* 55:5413 (1990) as a useful reducing agent.

In addition to the method of Honda, et al. (See above), several methods for phosphorylating the 5' terminus of an oligonucleotide are known. Initially, enzymatic methods using polynucleotide kinase were employed after the oligonucleotide was synthesized and removed from the solid support. Others have taught methods and reagents for chemically phosphorylating a synthesized oligonucleotide prior to its removal from the solid support. Some of these are described below.

Kondo, et al. *Nucl. Acids Res. Symposium Series* 16:161-164 (1985) describe phosphotriester (1) and phosphoramidite (2) reagents for phosphorylating 5' termini. Phosphorylation is achieved by preparing a special diphosphorylated (3' - 5') nucleotide which is added as the last nucleotide in the chain. The 3' phosphate is linked via the phosphotriester or phosphoramidite to the extending nucleotide chain. The 5' phosphate is protected with a protecting group which is ultimately removed.

Uhlmann, et al. *Tetrahedron Letters* 27(9): 1023-1026 (1986) describe a phosphoramidite phosphorylating reagent using a p-nitrophenylethyl

group as a blocking group. They mention that the hydrophobic p-nitrophenylethyl is advantageous in that phosphorylated compounds can be separated from non-phosphorylated compounds by reversed phase HPLC.

Uhlmann, et al, however, used only hexamers to which the p-nitrophenylethyl "handle" was attached. A similar approach using a p-nitrophenylethyl handle with 20-mers is described by G. Zon in chapter 14 of *HPLC in Biotechnology*, (W.S. Hancock, ed), J. Wiley & Sons, New York, NY. pp 359-363 (1990). The purification results obtained by Zon with this method are marginal.

10 Marugg, et al. *Nucl. Acids Res.* 12(22):8639-8651 (1984) describe a new phosphorylating agent, 2-cyano-1,1-dimethylethoxy dichlorophosphine. This agent has the alleged advantage of being removed under just basic conditions.

Himmelsbach, et al. *Tetrahedron Letters* 23(46):4793-4796 (1982) describe a new phosphorylating agent, bis-(p-nitrophenylethyl) phosphoromonochloridate. Van der Marel, et al. *Tetrahedron Letters*, 22(19):1463-1466 (1981) describe a morpholino phosphoro bis-3-nitro-1, 2, 4-triazolidate.

Horn, et al. *Tetrahedron Letters* 27 (39):4705-4708 (1986) describe a phosphorylating reagent including a 4, 4' dimethoxytrityl group which, upon release, can be used to monitor the efficiency of phosphorylation. This disclosure appears to be quite similar to that of EP-A-304 215 and to the commercially available Clontech product known as 5' Phosphate-On.

Lipshutz, et al. *Tetrahedron Letters* 30(51): 7149-7152 (1989) ("Lipshutz 1989") and Lipshutz, et al. *Tetrahedron Letters* 21:3343-3346 (1980) ("Lipshutz 1980") and Von Peter Sieber, *Helvetica Chimica Acta* 60:2711 (1977) all disclose the use of fluoride in the removal of a silyl protecting group. In this regard, they are similar to Honda, et al. (See above).


While each of the above reagents and methods are adequate for phosphorylating synthesized oligonucleotides, each has draw backs as well. For example, each of the recited references discloses a method for removing the phosphate blocking group to generate the native 5' phosphate. Some (e.g. Horn, et al.) describe a blocking agent having a detectable characteristic (eg. color) by which the extent of phosphorylation can be monitored. While the extent of phosphorylation can be monitored by this means, it provides no means for purification. Uhlmann, et al. suggest that the hydrophobic p-nitrophenylethyl group can be used prior to cleavage to separate phosphorylated hexamers by HPLC. The protected hexamers cited by Uhlmann, having a relatively low

molecule/protecting group mass ratio, are generally too short to provide specificity necessary in hybridization assays.

However, none of the references teach phosphorylating/ blocking reagents comprising silyl substitutes. Further, none suggest that the silyl protecting group
5 can be used to purify phosphorylated nucleotides from unphosphorylated failure product. The present invention seeks to overcome these disadvantages.


SUMMARY OF THE INVENTION

In a first aspect, the invention relates to an alcoholic compound of the
10 formula:



wherein R_1 , R_2 and R_3 are independently selected from the group consisting of
15 aryl, substituted aryl, and substituted or unsubstituted sterically bulky alkyl; and n is an integer from 2 to about 20, more often 2 to about 6, and ideally 2. Where any of R_1 , R_2 and R_3 are aryl, substituted aryl or aralkyl, the aromatic portions will generally comprise rings having at least 5 carbon atoms. Phenyl, naphthyl, methoxyphenyl, tolyl and triphenylmethyl are illustrative groups. Where any of
20 R_1 , R_2 and R_3 are alkyl or substituted alkyl, they will comprise sterically bulky alkyl having at least 4 carbons in a branching and/or cyclic chain. Illustrative groups include t-butyl, neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl.

25 In another aspect, the invention relates to esters, particularly acetates, of the formula:



wherein R_1 , R_2 and R_3 are independently selected as for the alcohol above; R_4 is
30 lower alkyl, especially methyl; and n is an integer from 2 to about 20, more often 2 to about 6, and ideally 2. Preferred R_1 , R_2 and R_3 substituents are the same as for the alcohol above.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a chromatogram showing separation of a phosphorylated oligonucleotide (peak 4 at 15.5 min) ^{for} from the failure products (peak 1 at 8.4 min). The chromatogram was generated from a Waters μ Bondapak™ C18 column, 3.9 mm x 150 mm flowing at 1.5 mL/min. Solvent A was 100mM Triethylammonium Acetate and solvent B was Acetonitrile. Solvents were mixed according to a linear gradient table such that the ratio of A:B was as follows: At time=0, 90:10; at time=15 min, 60:40; at time=25, 60:40; and at time=30, 90:10. Detection was in absorbance units at 260 nm. (See Example ^{9a} 10a.)

10 Figure 2 is a chromatogram showing separation of deprotected, phosphorylated oligonucleotide (peak 1 at 8.4 min) from other products (eg silylfluorides). The conditions are the same as in Figure 1. (See Example ^{9b} 10b.)

DETAILED DESCRIPTION15 A. General Definitions

In general, terms like "alkyl", "alkenyl" and "aryl" have the meanings usually attributed to them by persons skilled in the art of organic chemistry. For example, alkyl refers generally to monovalent straight or branched aliphatic radicals which may be derived from alkanes by the removal of one hydrogen, and have the general formula C_nH_{2n+1} . Alkyl groups may have from 1 to about 30 carbons, more practically 1 to about 15 or 20. "Lower alkyl" refers to alkyls having from 1 to about 6 carbons. Examples of lower alkyl include CH_3- , CH_3CH_2- , $CH_3CH(CH_3)-$, and $CH_3(CH_2)_4-$. As used herein, "alkyl" includes cycloalkyl as well as straight alkyl. Thus, cyclohexyl and others are included.

25 "Alkenyl" refers to monovalent straight or branched aliphatic radicals which may be derived from alkenes by the removal of one hydrogen, and have the general formula C_nH_{2n-1} . Alkenyl substituents may have from 1 to about 30 carbons, more practically 1 to about 20. "Lower alkenyl" refers to alkenyls having from 1 to about 6 carbons. "Olefinic" is a synonym for alkenyl.

30 As used herein, "alkylene" refers to a divalent straight or branched chain spacer group containing less than 30 carbon atoms, including but not limited to, $-CH_2-$, $-CH(CH_3)-$, $-CH(C_2H_5)-$, $-CH(CH_3)CH_2-$, $-(CH_2)_3-$, and the like. Generally, an alkylene spacer group is aliphatic.

35 "Aryl" refers to a monovalent radical derived from aromatic hydrocarbons by the removal of one hydrogen. Aryl substituents have ring structures, such as those of phenyl and naphthyl. Typically, aryl substituents are planar with the π electron clouds of each carbon remaining on opposite sides of the plane.



Although alkyl, alkenyl and aryl are generally limited to groups having no atoms other than carbon and hydrogen (ie. no heteroatoms), the invention is not so limited. Heteroatoms, especially oxygen and sulfur, can be present in "R" groups to form "oxa" and "thia" analogs, respectively. However, because of the anticipated elimination, it is desirable to avoid oxa analogs having an oxygen atom 2 carbons removed from the point of monovalency where the R group is attached to the molecule of interest. Exemplary oxa analogs include alkoxy, such as t-butoxy, isopropoxy and ethoxy, phenoxy and ether substituents.

As used here, "substituted" refers to the presence of moieties covalently bonded to the "R" groups, including, but not limited to, halide (especially Br and Cl), nitro, lower alkoxy (having from 1-6 carbon atoms, especially methoxy and ethoxy), lower alkyl (having from 1-6 carbon atoms, especially methyl and ethyl), hydroxy, and amino (protecting group may be required). Subject to constraints imposed by the desired solubility, and hydrophobicity of the desired compound, and by the steric constraints of organic chemistry principles, the substituting groups may be placed anywhere, and in any number, on the R group. Some specific substitutions include: Alkaryl, which refers to a monovalent aryl radical bearing alkyl substituents where the aryl radical includes the point of monovalency (eg. toluyl); and Aralkyl, which refers to monovalent alkyl radicals bearing aryl substituents. In this latter case, the alkyl radical includes the point of monovalency. Benzyl is an example of an aralkyl group.

As used herein, "sterically bulky" refers to substituents groups which occupy a relatively large volume. Aryl groups having five or more carbons are considered "sterically bulky", as are substituted aryl groups. Alkyl and alkenyl groups are "sterically bulky" when they possess at least 4 carbons and are arranged in a branched configuration, the more branches, the bulkier. Any alkyl occupying a volume equal to or larger than t-butyl; and any aryl occupying a volume equal to or larger than phenyl, is considered "sterically bulky". Thus, neopentyl, neohexyl and others meet this description.


"Hydrophobic" refers generally to compounds which are relatively insoluble in aqueous solutions and will not substantially mix with water. Specifically, a compound is deemed hydrophobic if it has a partition coefficient of 0.51 or greater to octanol in a water/octanol partitioning test.

35 B. Silyl Alcohol Synthesis

Silyl alcohols prepared by any method may be useful in the inventions described below. 2-silyl-ethan-1-ols (or β -silylethanol or silapropanols) are

Catalytic hydrosilation can be performed under the following conditions. The molar ratio of acetate to silane can range from about 30:1 to about 1:2, and is preferably about 1:1. Intermediate ratios, such as 10:1 or 2:1, are contemplated as well. The metal catalyst may be present in mole percentages ranging from about 5 0.01% to about 3%, preferably between about 0.2% and about 2%. Lower percentages may require longer reaction times or higher temperatures. For $\{\text{RhCl}(\text{CO})_2\}_2$ an optimal mole % is between about 0.25% and about 1.0%. For other catalysts, the optimal concentrations can be obtained from the literature or from routine experimentation. The reaction is best run at room temperature for 10 about 50-70 hours, preferably not longer than 2 weeks. It may, however, proceed more quickly at elevated temperatures; for example, in less than 24 hours at 82 C. The principle reagents should be present at a concentration ranging from neat in vinyl acetate to 4 M in toluene; preferably about 1 M in toluene. Other reaction conditions for this catalytic step can be found in Collman, et al., which is 15 incorporated by reference.

In hydrosilation, two major products result because the silyl substituent may bond to either side of the double bond to give both 1- and 2- substituted products. If necessary, these can be separated and purified by chromatography, for example, silica based chromatography such as flash column or HPLC. 20 However, in a specific instance, purification is greatly simplified. When vinyl acetate is used as the ester, two products are again obtained as follows:



25 Upon hydrolysis in mild aqueous or alcoholic base, the acetate is converted to an alcohol. However, the 1-silyl substituted alcohol is unstable and spontaneously undergoes a Brook rearrangement (A.G. Brook, *Accounts Chemical Research*, 7:77 (1974)) to give:



The 2-silyl substituted alcohol does not undergo this rearrangement. Since the 2-silyl substituted product behaves as an alcohol, while the silyl ether behaves as an ether, the two products are easily separated on the basis of these properties using silica gel chromatography, especially HPLC. As will be seen from the examples
5 which follow, this hydrolysis reaction can be run in the same vessel without any intermediate purification of the acetate.

The conditions of hydrolysis preferably are carefully controlled. It will be recalled that 2- or β - substituted intermediates in the anionic form will undergo fragmentation to the silanol as described in the Background section. However,
10 reaction conditions can be selected which will minimize the formation of the undesired product. First, a mild base is selected, preferably one having a pK_b between about 3 and 8. Acceptable bases include the sodium or potassium salts of $H_2BO_3^-$, HPO_4^{2-} , SO_3^{2-} , HCO_3^- and CO_3^{2-} . A mild base, having only a weak tendency to dissociate, tends to keep anionic species protonated moreso than strong
15 bases or hydrides (eg. $LiAlH_4$) taught in the prior art.

The base should be present in a base:acetate molar ratio of from about 0.01:1 to about 3:1, preferably between 0.1:1 and 2.5:1, most preferably between 1:1 and 2:1. The reaction generally takes from 0.5 to 24 h, but preferably takes about 1 hour. In addition, solvents can be selected to minimize
20 the formation of undesired product. For example, the solubility of the base in a particular solvent will affect its strength. It is desirable to have a substantial amount of the base insoluble so as to buffer the ionization equilibrium going on in solution. Also, protic solvents are preferred over aprotic solvents, due to their ability to quench the formation of anionic species. Suitable protic solvents include
25 water, methanol and ethanol. Although the reaction will work in an aqueous medium, it is preferable to use methanol as the solvent.


Silyl alcohols synthesized by this method, as well as silyl alcohols synthesized by other processes, find utility in the synthesis of phosphorylating reagents and protecting agents as described in a later section.

30 C. Silyl Alcohols:

While the above-described method may be used to synthesize many silyl substituted alcohols, one class of silyl alcohols is of particular interest. Previous methods have not been known to synthesize silyl alcohols having three large, bulky
35 groups bonded to the silicon. This is because the principal prior art method of synthesis-- ie., via vinyl silanes as taught by Kumada, et al and Seyferth, *supra*- requires vinyl silane reagents appropriately substituted with the necessary

bulky groups. Presumably due to steric considerations, bulky vinyl silanes are not readily available or easily synthesized. Although triphenyl silane is known (see Lesage, et al. *supra*), such a bulky silane has not been associated with a vinyl radical to make the bulky vinyl silane.

- 5 However, sterically bulky silyl alcohols can be made by the above described method, and have the general formula:



- 10 wherein R₁, R₂ and R₃ are independently selected from sterically bulky groups like aryl (eg. phenyl and naphthyl), substituted aryl (eg. methoxyphenyl, or nitrophenyl) aralkyl (eg. triphenylmethyl), alkaryl and alkyl or substituted alkyl having at least 4 carbons in a branched chain (eg. t-butyl, neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl). In the formula above, n
15 is an integer from 2 to about 20, usually 2 to about 6 and most preferably 2. Exemplary compounds are listed in the table below, although this is by no means an exhaustive list.

TABLE 2: Illustrative Novel Silyl Alcohols

20

R ₁	R ₂	R ₃	n
phenyl	phenyl	phenyl	2
phenyl	phenyl	phenyl	6
phenyl	phenyl	t-butyl	2
phenyl	t-butyl	t-butyl	2
phenyl	naphthyl	neopentyl	2
t-butyl	t-butyl	neopentyl	2
phenyl	naphthyl	t-butyl	2
phenyl	t-butyl	neohexyl	2
phenyl	phenyl	phenyl	3
phenyl	phenyl	t-butyl	3
t-butyl	t-butyl	phenyl	3
phenyl	naphthyl	neopentyl	3
t-butyl	t-butyl	neopentyl	3
phenyl	naphthyl	t-butyl	3
phenyl	t-butyl	neohexyl	3

For reasons which will become apparent, substituted bulky groups preferably are substituted with nonpolar substituents.

- 25 As mentioned, silyl alcohols find utility in preparing phosphorylating agents and protecting agents. These are described in detail below.

D. Phosphorylating Reagents


Many types of reagents can phosphorylate-- ie. put a phosphate group on the end of-- an oligonucleotide. Generally these reagents are classified as
 5 phosphotriester reagents, phosphonate reagents (Hydrogen or Alkyl) and phosphoramidite reagents. The mechanisms by which each of these reagents phosphorylate an oligonucleotide is described in the literature.

A novel phosphorylating reagent is represented by the formula:



wherein R_5 , R_6 and R_7 are independently selected from H, alkyl, aryl, substituted alkyl, substituted aryl, oxa and thia analogs of alkyl, aryl, substituted alkyl and substituted aryl, and halogen; and wherein Q represents a moiety selected from the
 15 group consisting of phosphoramidites, alkyl phosphonates, hydrogen phosphonates and phosphotriesters.


For a phosphoramidite, Q has the formula:



and R_8 is generally selected from the group consisting of 2-cyanoethyl, methyl, ethyl, 2-alkylsulfonyl ethyl, 2-(p-nitrophenyl)ethyl, 2-(9-fluorenyl)ethyl, 2-(2-anthraquinonyl)ethyl, 2-alkylthioethyl, 2-arylthioethyl, 2-trihalomethylethyl, 2-phenylethyl and 2-(2-naphthyl)ethyl. R_9 and R_{10} are
 25 generally selected independently from H, or straight or branched alkyl having from 1-6 carbons. In a very common phosphoramidite moiety, R_8 is 2-cyanoethyl while R_9 and R_{10} are both isopropyl.

The novel silyl phosphoramidite may be prepared in a conventional manner by reacting a chlorophosphoramidite with a silyl substituted alcohol. See, eg.,
 30 Koster *Tetrahedron Letters*, 24:5843 (1983) which is incorporated herein by reference. Here, it is preferred to use a 2-silyl-ethan-1-ol. The reaction conditions are well known from the literature.

For a phosphotriester reagent, Q has the formula:



wherein Y is hydroxyl, or alkoxy.



- 5 For hydrogen phosphonate or alkyl phosphonate reagents, Q has the above formula, but Y is H or alkyl, respectively.

E. Methods Using Phosphorylating Agents

10 The above-described phosphoramidite, phosphotriester and phosphonate reagents can be used in a method for phosphorylating an oligonucleotide, particularly an oligonucleotide synthesized on a solid support. It will be realized by those of ordinary skill in the art that a single nucleoside could equally well be phosphorylated in this manner, as could longer polynucleotides. For simplicity, it will be understood that the term "oligonucleotide" will include structures having
15 from one to several hundred nucleoside subunits.

Many methods are known in the literature for synthesizing oligonucleotides and the particular method employed is not relevant to the present invention. Generally, however, automated synthesis is preferred and may be performed using commercial instruments such as an ABI 380A Synthesizer or a Milligen 8700
20 Synthesizer.


The reaction steps employed by such automated synthesizers are generally known in the art and need not be repeated here. However, it will be noted that when a phosphoramidite or H phosphonate reagent is used, the resulting intermediate is a trivalent phosphite. It is subsequently oxidized to the
25 biologically useful pentavalent phosphate. This oxidation step is readily achieved using, for example iodine, in the automated synthesis process.

A major advantage of the present invention is that the phosphorylation step can be accomplished in the same instrument as synthesis, without removal of the oligonucleotide from the support. Alternatively, oligonucleotides synthesized by
30 other methods (eg. enzymatic) may be phosphorylated by the methods of the present invention, provided the amino and hydroxy functions present can be protected.

While known methods of phosphorylating have been described in the Background section, none use silyl reagents. Any of the phosphorylating reagents
35 prepared in the preceding section, may be used to phosphorylate an oligonucleotide

according to the invention. The methods and conditions are conventional, although the reagents are not. The examples provide further details but the method generally comprises reacting the 5' hydroxyl of an oligonucleoside with a phosphorylating reagent described above, ultimately to form a phosphodiester protected by the silyl group.

The silyl protected, phosphorylated intermediate has the structure:



where R_5 , R_6 and R_7 are selected as before; Z is H or OH; and BASE represents one of the nucleic acid bases A, C, G, T or U, or analogs thereof. The terminal nucleoside may be attached at its 3' carbon to a support (in the case of phosphorylating a single nucleoside) or, more likely, to a string of one or more other nucleosides (to form an oligonucleotide). Generally, such a string of nucleosides will be connected via phosphodiester linkages, although other linkages are possible (eg. alkyl phosphonate neutral probes). Obviously, where Z is H, the nucleoside is a deoxyribonucleoside; where Z is OH, it is a ribonucleoside. Analogs of the bases A, C, G, T or U are compounds which, when incorporated into an oligonucleotide, will still permit Watson-Crick base pairing with their respective complementary base. Some exemplary base analogs are published in the USPTO Official Gazette at 1114 OG 43, which is incorporated herein by reference.

While the silyl protecting group must be removed for biological use (eg. template guided ligation) the protected intermediate also has utility. The silyl group, particularly if it is endowed with bulky, hydrophobic substituents R_5 , R_6 and R_7 , is useful as a "handle" for purifying and separating phosphorylated oligonucleotides from unphosphorylated failure product by chromatography, eg. HPLC. Provided the R groups are sufficiently hydrophobic, the oligonucleotide bearing the silyl protecting group is easily differentiable from the unphosphorylated, unprotected oligonucleotide, even when the oligonucleotides approach 50-mer lengths. Of course, shorter lengths are also easily separated.

This goes a step beyond the known trityl protecting groups which are useful to monitor phosphorylation success, but not to separate or purify product.

If desired, a deprotecting step may follow phosphorylation and/or separation to yield the 5' terminal phosphate. The deprotecting step is done by any
5 useful method to yield the desired phosphate. A preferred method, especially useful when the silyl substituent is β to the oxygen as above, involves reacting the protected phosphodiester with fluoride ion to give the silyl fluoride, ethylene and the terminal phosphate. Tetrabutylammonium fluoride (TBAF) is a useful fluoride ion for removing the silyl protecting group. This reaction is driven by the release
10 of ethylene when the phosphorylating reagent above is used. See, eg Grob, *Helv. Chim. Acta*, 38:594 (1955). It is for this reason that 2-silyl-ethan-1-ols (β silylethanol) are preferred silyl alcohol reagents (they have two carbons between the silicon and the oxygen of the phosphodiester, thus permitting the Grob elimination of ethylene). Any other length will not be removed in the deprotection
15 step as easily as the β silyl-ethanol derivative.

"Protecting" group and "deprotecting" steps refer to the silapropyl substituent attached to the oxygen of the phosphate. This group may or may not afford "protection" in the usual sense from subsequent reactions that would affect the oxygen atom. However, the term is used as a synonym for "handle" because of
20 the ability to separate phosphorylated oligonucleotide from unphosphorylated failure product using the silapropyl group, and because of the subsequent removal of the group to give the desired phosphate.

The inventions herein described will be better understood in view of the following examples which are intended to be illustrative and non-limiting.


25

EXAMPLES

A. Preparation of Silyl Alcohols:

5 EXAMPLE 1:

a) Preparation of 1,1,1-Triphenyl-3-acetoxy-1-silapropane (3)



- 10 A solution of 3.69 mL (40 mmol) of vinyl acetate (1), 10.42 g (40 mmol) of triphenylsilane (2), and 77.8 mg (0.25 mmol) of $\text{Rh}_2\text{Cl}_2(\text{CO})_4$ in 40 mL of toluene was stirred at room temperature under N_2 for a total of 63 h. Several runs of the reaction at this scale had unpredictable induction periods, followed by rapid heat evolution. Scaleup of this reaction should be done with a cooling bath
- 15 close at hand. The very dark reaction mixture was treated with 5 g of decolorizing charcoal, and the mixture boiled briefly. After cooling, the mixture was filtered through a 1 cm pad of Celite™ with filtrate and washings being collected. Solvent was evaporated and the remaining residue was vacuum dried. At this point, the crude material was carried on to the hydrolysis step. NMR analysis showed an a:β
- 20 ratio of 1:1.57. The following protocol was carried out for compound identification purposes. A 100 mg sample of crude material was flash chromatographed using 4 % EtOAc in cyclohexane on a 25 mm I.D. x 150 mm long silica gel column. This afforded 29 mg of (3) after recrystallization from MeOH, mp 67-68 C.

25

IR: (CDCl_3 , cm^{-1}) 3070 (m), 1728 (vs), 1425 (vs), 1249 (vs)

MS: (DCI/ NH_3) m/e 364 ($\text{M}+\text{NH}_4$)

NMR: (300 MHz, CD_2Cl_2) δ 7.6-7.3 (m, 15H, phenyl), 4.22 (B₂ of A₂B₂, 2H, CH₂O), 1.87 (s, 3H, CH₃), 1.86 (A₂ of A₂B₂, 2H, CH₂Si)


- 30 ¹³C NMR: (75 MHz, CDCl_3) δ 171.1 (C=O), 135.5 (meta), 134 (ipso), 129.7 (para), 128 (ortho), 62.1 (CH₂O), 21 (Me), 14.4 (CH₂Si)

Elemental Analysis: Calc'd for $\text{C}_{22}\text{H}_{22}\text{O}_2\text{Si}$; C: 76.26; H: 6.40

-17-

Found; C: 76.45; H: 6.37

b) Preparation of 1,1,1-Triphenyl-1-silapropene-3-ol (4)



5

The crude (3) was dissolved in 100 mL MeOH, and 10.0 g of K_2CO_3 was added all at once. The reaction was complete after 1 h of stirring at room temperature. The solids were filtered off, and the filtrate was concentrated. The concentrated residue was partitioned between 100/100 mL $\text{H}_2\text{O}/\text{EtOAc}$. After solvent removal from the organic layer, the residue was vacuum dried. Flash chromatography (18% EtOAc in cyclohexane, $R_f=0.32$) using a 41 mm I.D. x 150 mm long silica gel column afforded 3.42 g (28%) of (4). Recrystallization from cyclohexane gave the analytical sample as a snow-white solid, mp 96-97 C.

15

IR: (CDCl_3 , cm^{-1}) 3616 (m), 2970 (m), 1429 (vs)

MS: (FAB/DMF-KI) m/e 343 (M+K)

NMR: (300 MHz, CD_3OD) δ 7.55-7.3 (m, 15H, phenyl), 3.73 (B_2 of A_2B_2 , 2H, CH_2O), 1.78 (A_2 of A_2B_2 , 2H, CH_2Si)

20 ^{13}C NMR: (75 MHz, CDCl_3) δ 135.5 (meta), 134.4 (ipso), 129.6 (para), 128 (ortho), 59.8 (CH_2O), 18.7 (CH_2Si)

Elemental Analysis: Calc'd for $\text{C}_{20}\text{H}_{20}\text{OSi} \cdot 0.2 \text{H}_2\text{O}$; C:77.98; H: 6.67

Found; C:77.92; H:6.62

25 EXAMPLE 2:

a) Preparation of 1,1-Dimethyl-1-phenyl-3-acetoxy-1-silapropene

To a solution of 6.13 mL (40 mmol) of PhMe_2SiH and 3.69 mL of vinyl acetate in 40 mL of toluene was added 61.3 mg (0.16 mmol) of $\text{Rh}_2\text{Cl}_2(\text{CO})_4$. Immediately, the reaction evolved heat and gas. Within 5 min, the golden yellow reaction had turned dark brown in color. After 1 h, the reaction was complete. The reaction was worked up as in example 1a to give 8.39 g of crude adduct. Proton NMR analysis showed an

30

-18-

a:b addition ratio of 1.44:1.0. A 100 mg sample was purified by flash chromatography as in example 1a to give 28 mg of the title compound as a colorless oil.

- 5 IR: (CDCl₃, cm⁻¹) 2960 (m), 1724 (vs), 1426 (m), 1255 (vs)
MS: (DCI/NH₃) m/e 240 (M+ NH₄)
NMR: (300 MHz, CDCl₃) δ 7.6-7.3 (m, 5H, phenyl), 4.18 (B₂ of A₂B₂, 2H, CH₂O), 1.99 (s, 3H, Me), 1.25 (A₂ of A₂B₂, 2H, CH₂Si), 0.35 (s, 6H, SiMe)
13C NMR: (75MHz, CDCl₃) δ 171.1 (CO), 138 (ipso), 133.4 (meta), 129.2 (para), 127.9 (ortho), 62.3 (CH₂O), 21.1 (Me), 16.5 (CH₂Si), -2.9 (SiMe)
10 Elemental Analysis: Calc'd for C₁₂H₁₈O₂Si; C: 64.82; H: 8.16
Found; C: 65.02; H: 8.07

b) Preparation of 1,1-Dimethyl-1-phenyl-1-silapropane-3-ol

- 15 The remaining 8.29 g of crude product from part b, above, was worked up as in the case of example 1 to give 1.64 g of 1,1-dimethyl-1-phenyl-1-silapropane-3-ol as a colorless oil, 23 % overall.


- IR: (CDCl₃, cm⁻¹) 3616 (m), 2960 (m), 1425 (m), 1251 (s)
20 MS: (DCI/NH₃) m/e 198 (M+NH₄)
NMR: (300 MHz, CDCl₃) δ 7.6-7.3 (m, 5H, phenyl), 3.75 (B₂ of A₂B₂, 2H, CH₂O), 1.49 (s, 1.2H, OH), 1.22 (A₂ of A₂B₂, 2H, CH₂Si), 0.33 (s, 6H, SiMe)
13C NMR: (75 MHz, CDCl₃) δ 138.5 (ipso), 133.4 (meta), 129 (para), 127.8 (ortho), 59.9 (CH₂O), 21.1 (CH₂Si), -2.8 (SiMe)
25 Elemental Analysis: Calc'd for C₁₀H₁₆OSi·0.1 H₂O; C: 65.92; H: 8.99
Found; C: 65.95; H: 8.97

EXAMPLE 3:

- a) Preparation of 1,1,1-Triethyl-3-acetoxy-1-silapropane To a
30 solution of 6.39 mL (40 mmol) of Et₃SiH and 3.69 mL (40 mmol) of vinyl acetate in 40 mL of toluene is added 61.3 mg (0.16 mmol) of Rh₂Cl₂(CO)₄.
Caution: the reaction evolves heat and gas. Within about 5 min, the reaction mixture darkens in color. Reaction is judged complete by TLC analysis (10% EtOAc in cyclohexane) after 1 h. The reaction is worked up and purified, if
35 desired, as in example 1a.

b) Preparation of 1,1,1-Triethyl-1-silapropane-3-ol () The crude product from part a) can be worked up as in the case of triphenylsilylethanol (example 1) to give the 1,1,1-triethyl-1-silapropane-3-ol.

5 EXAMPLE 4: Preparation of 1,1,1-Triphenyl-1-silaheptane-7-ol (5)



(5)

10 The acetate of 5-hexen-1-ol is prepared by refluxing 4.8 mL (40 mmol) of the alcohol in 15/15 mL of pyridine/acetic anhydride for 4 h. The solvents are removed in vacuo, and the residue is thoroughly vacuum dried. The crude acetate is dissolved in 40 mL of toluene, and 10.42 g of triphenylsilane is added, followed by 77.8 mg (0.25 mmol) of Rh₂Cl₂(CO)₄. The reaction is stirred at room
15 temperature under N₂ for 24 h, during which time the reaction turns dark brown in color. Some quantities of the isomer 2-methyl-1,1,1 triphenylsilahexan-6-ol can be expected. If necessary, the isomers can be separated by chromatography. Workup as in example 1a, followed by base hydrolysis as in 1b, affords the title compound, (5).

20

B. Preparation of Phosphorylating Reagents:

EXAMPLE 5: Preparation of 2-Trimethylsilylethyl-2-cyanoethyl-N,N-diisopropylaminophosphoramidite (1)


25

To a solution of 573 μ L (4 mmol) of 2-trimethylsilylethanol (commercially available from Aldrich Chemical, Milwaukee, WI; or prepared in a manner analogous to example 2, above) and 1.39 mL (8 mmol) of *i*-Pr₂NEt in 8 mL of THF at 0 C was added 892 μ L (4 mmol) of 2-cyanoethyl-N,N-
30 diisopropylaminochloro-phosphoramidite all at once. The reaction became very cloudy almost immediately. The ice bath was removed, and the reaction stirred to room temperature overnight, for a total of 19 h. After filtration to remove *i*-Pr₂NEt-HCl, the THF was evaporated. The residue was partitioned between 50/50 mL EtOAc/0.1 M Na₂CO₃, pH 12. After phase separation and solvent removal of
35 organic phase, the residue was vacuum dried. Flash chromatography using 12%

-20-

EtOAc in cyclohexane on a 150 mm x 25 mm ID column afforded 573.8 mg (78%) of the title compound as a water-white viscous oil, $R_f = 0.65$ in 15% EtOAc in cyclohexane.

- 5 MS: (DCI, NH_3) 319 (M+H), 291 (M-HCN)
 NMR: (CD_2Cl_2) δ 3.9-3.62 (m, 4H), 3.56 (dsept, 2H, $J_{\text{CH}}=7.0$ Hz, $J_{\text{PH}}=10.0$ Hz, NH), 2.59 (t, 2H, $J=6.2$ Hz, CH_2CN), 1.15 (dd, 12H, $J_{\text{CH}}=7.0$, $J_{\text{PH}}=2.2$ Hz, Me), 0.97 (tq, 2H, $J=8.0, 0.7$ Hz, CH_2Si), 0.03 (s, 9H, SiMe)
- 10 **EXAMPLE 6: Preparation of 2-Triphenylsilylethyl-2-cyanoethyl-N,N-diisopropylaminophosphoramidite (6)**



- 15 To a solution of 3.04 g (10 mmol) of (4), 4.18 mL (24 mmol) of $i\text{-Pr}_2\text{NEt}$, and 5 mg of 4,4-dimethylaminopyridine in 15 mL of THF at 0 C was added 2.68 mL (12 mmol) of 2-cyanoethyl-N,N-diisopropylaminochlorophosphoramidite (5) all at once. A white precipitate formed almost immediately. Reaction was complete after 30 min at 0 C. After solvent removal, the residue was partitioned between
- 20 100/100 mL 0.1 M $\text{Na}_2\text{CO}_3/\text{EtOAc}$, and the phases separated. The aqueous phase was re-extracted with 50 mL EtOAc, and the combined organic phases were concentrated and vacuum dried. Flash chromatography (10% EtOAc in cyclohexane) using a 41 mm I.D. x 150 mm long silica gel column gave 3.35 g of (6) (66%) after vacuum drying overnight as a viscous colorless oil. This
- 25 material gradually crystallized in a -20 C freezer over the course of several weeks. During the chromatography, 100 μL NEt_3 was added to each fraction, in

order to minimize the effects of adventitious acid in the fraction tubes or in the silica gel used for flash chromatography.

IR: (film, cm^{-1}) 2962 (m), 1426 (m)

5 MS: (DCI/NH_3) m/e 505 (M+H)

NMR: (300 MHz, CD_3CN) δ 7.6-7.3 (m, 15H, phenyl), 3.9-3.7 (m, 2H, CH_2O), 3.66 (dt, 2H, $J_{\text{CH}}=5.9$ Hz, $J_{\text{PH}}=7.7$ Hz, CH_2O), 3.51 (dsept, 2H, $J_{\text{CH}}=6.6$ Hz, $J_{\text{PH}}=9.9$ Hz, NH), 2.54 (t, 2H, $J=5.5$ Hz, CH_2CN), 1.87 (br t, 2H, $J=6.3$ Hz, CH_2Si), 1.07 (dd, 12H, $J_{\text{CH}}=6.6$ Hz, $J_{\text{PH}}=29.4$ Hz, Me)

10 ^{13}C NMR: (75 MHz, CD_3CN) δ 136.3 (meta), 135.5 (ipso), 130.7 (para), 129 (ortho), 117.7 (CN), 61.1 (d, $J_{\text{PC}}=18.3$ Hz, CH_2O), 59.3 (d, $J_{\text{PC}}=18.3$ Hz, CH_2O), 43.6 (d, $J_{\text{PC}}=12.2$ Hz, NCH), 24.8 (virtual t, $J_{\text{PC}}=7.3$ Hz, Me), 21 (d, $J_{\text{PC}}=7.3$ Hz, CH_2CN), 17.2 (d, $J_{\text{PC}}=7.3$ Hz, CH_2Si)

^{31}P NMR: (202 MHz, CD_3CN) δ 145.6

15

EXAMPLE 7: Preparation of 2-Triethylsilylethyl-2-cyanoethyl-N,N-diisopropylaminophosphoramidite

Example 5 is repeated except the product of example 3b is used as the starting compound to produce the title compound.

20

EXAMPLE 8: Preparation of 2-bismethylphenylsilylethyl-2-cyanoethyl-N,N-diisopropylaminophosphoramidite

Example 6 is repeated except the product of example 2b is used as the starting compound to produce the title compound.

25


EXAMPLE 9: Preparation of Triphenylsilylethyl H-phosphonate-DBU reagent

To a solution of N-methylmorpholine (89 equiv), triazole (33 equiv) and PCl_3 (10 equiv) is added triphenylsilylethanol at 0 C. The reaction is stirred at this
30 temperature for 2.5 h. The reaction is then quenched by addition of 100mM 1,5-diazabicyclo [5.4.0] undec-5-ene (DBU)-bicarbonate, and the phases are separated. The organic phase is stripped to dryness in vacuo, and the crude H-phosphonate-DBU is purified by chromatography.

C. Preparation of Phosphorylated, Protected Oligonucleotides and Deprotection Thereof:


EXAMPLE 10:

5 a) Use of (6) in Automated Phosphorylation of DNA



10 The phosphoramidite (6) (example 6, above) was used to phosphorylate a 25-
 mer oligonucleotide at the 1 μ mol level using an ABI (Foster City, CA) 380A DNA
 Synthesizer. The phosphoramidite couplings were run using the synthesis
 program from the manufacturer except that the "wait" time (time of contact of
 phosphoramidite solution with support) and "wash" time are both doubled. The
 preparative HPLC run, showing separation of the failure sequences from full-
 15 length oligo, is shown in Figure 1.

b) Deprotection of Phosphorylated Oligonucleotide



The collected material (7) from part a) was dried in vacuo, then ethanol precipitated. The purified DNA was then desilylated using 100/100 μ L of DMSO/1.0 M Tetra-n-butyl ammonium Fluoride (TBAF) (Aldrich, Milwaukee, WI) . The reaction was performed in a 68 C heating block for 3.5 h. The reaction was diluted to 500 μ L with 300 μ L of water, and the reaction was desalted by passage down a NAP-5 column (Pharmacia, Piscataway, NJ). The 1.0 mL eluate was dried in vacuo, then was ethanol precipitated to give purified, terminally phosphorylated DNA. HPLC analysis of this material is shown in Figure 2.

10

EXAMPLE 11:

Example 10 is repeated except the phosphoramidite reagent of example 5 is used in place of the phosphoramidite reagent of example 6.

15 **EXAMPLE 12:**

Example 10 is repeated except the phosphoramidite reagent of example 7 is used in place of the phosphoramidite reagent of example 6.

EXAMPLE 13:20

Example 10 is repeated except the phosphoramidite reagent of example 8 is used in place of the phosphoramidite reagent of example 6.


EXAMPLE 14: Phosphorylation by the H-phosphonate method

The reagent from Example 9 is used to phosphorylate an oligonucleotide using the general reaction protocol and conditions of Froehler, et al., *Tetrahedron Letters*, 27:469-472 (1986) except the coupling reagent is adamantoyl chloride and the capping reagent is β -cyanoethyl hydrogen phosphonate. After adamantoyl chloride catalyzed coupling of the 5'-hydroxyoligonucleotide with the triphenylsilylethyl hydrogen phosphonate is complete, all H-phosphonate linkages in the oligonucleotide are oxidized with iodine to the phosphodiester oxidation state. The oligonucleotide obtained can be separated on HPLC in the same manner as DNA of identical sequence which is prepared using phosphoramidite chemistry. This material may be desilylated in the same fashion as the phosphoramidite-prepared oligonucleotide.

35


The claims defining the invention are as follows:

1. An alcoholic compound of the formula:



wherein R_1 , R_2 and R_3 are independently selected from the group consisting of aryl, substituted aryl, ~~substituted aryl~~, and substituted or unsubstituted sterically bulky alkyl; and n is an integer from 2 to about 20.

2. The compound of claim 1 wherein R_1 , R_2 and R_3 are independently aryl, or substituted aryl.
3. The compound of claim 2 wherein R_1 , R_2 and R_3 are each phenyl.
4. The compound of claim 1 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of bulky alkyl, substituted bulky alkyl, bulky aralkyl or substituted bulky aralkyl.
5. The compound of claim 4 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of t-butyl, neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl.
6. The compound of claim 1 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of phenyl, substituted phenyl, naphthyl, triphenylmethyl, t-butyl neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl.
7. The compound of claim 1 wherein n is from 2 to about 6.
8. The compound of claim 7 wherein n is 2.
9. An ester compound of the formula:



wherein R_1 , R_2 and R_3 are independently selected from the group consisting of aryl, substituted aryl, and substituted or unsubstituted sterically bulky alkyl; R_4 is lower alkyl; and n is an integer from 2 to about 20.

10. The compound of claim 9 wherein R_1 , R_2 and R_3 are independently aryl, or substituted aryl.
11. The compound of claim 10, wherein R_1 , R_2 and R_3 are each phenyl.
12. The compound of claim 9 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of bulky alkyl, substituted bulky alkyl, bulky aralkyl or substituted bulky aralkyl.
13. The compound of claim 12 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of t-butyl, neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl.

14. The compound of claim 9 wherein R_1 , R_2 and R_3 are independently selected from the group consisting of phenyl, substituted phenyl, naphthyl, triphenylmethyl, t-butyl, neopentyl, neohexyl, cyclohexyl, 3-pentyl and 3-ethyl-3-pentyl.
15. The compound of claim 9 wherein n is from 2 to about 6.
16. The compound of claim 15 wherein n is 2.
17. The compound of claim 9 wherein R_4 is methyl.

Dated 16 June, 1994

Abbott Laboratories

**Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON**

1/2




FIG. 1

2/2




FIG. 2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/04723

A. CLASSIFICATION OF SUBJECT MATTER

IPC(S) : C07F 7/08 C07F 7/18
US CL : 556/449, 556/436

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 556/440

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 3,943,157 (Henrick et al.) 09 March 1976 See entire document.	1-12
A	US, A, 4,180,682 (Cohen et al.) 25 December 1979 See entire document.	1-12
A	US, A, 4,191,842 (Cohen et al.) 04 March 1980 see entire document.	1-12
A	US, A, 4,310,465 (Olson et al.) 12 January 1982 See entire document.	1-12
A	US, A, 4,882,451 (Yoshida et al.) 21 November 1989 See entire document.	1-12

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 02 SEPTEMBER 1992	Date of mailing of the international search report 13 NOV 1992
Name and mailing address of the ISA/ Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. NOT APPLICABLE	Authorized officer <i>Arthur Prescott</i> ARTHUR PRESCOTT Telephone No. (703) 308-4631