

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 May 2009 (22.05.2009)

PCT


(10) International Publication Number
WO 2009/063250 A2

- (51) International Patent Classification:
A61K 8/44 (2006.01) A61Q 5/12 (2006.01)
A61K 8/46 (2006.01) A61Q 5/02 (2006.01)
A61Q 19/10 (2006.01) C11D 1/94 (2006.01)
- (21) International Application Number:
PCT/GB2008/051071
- (22) International Filing Date:
17 November 2008 (17.11.2008)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0722550.1 16 November 2007 (16.11.2007) GB
- (71) Applicant (for all designated States except US): **INNOSPEC LIMITED** [GB/GB]; Innospec Manufacturing Park, Oil Sites Road, Ellesmere Port Cheshire CH65 4EY (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **COTRELL, Philip** [US/US]; 410 Wildwood Drive, Salisbury, North Carolina 28146 (US).
- (74) Agent: **PIDGGEON, Robert**; 15 Clare Road, Halifax Yorkshire HX1 2HY (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report



WO 2009/063250 A2

(54) Title: COMPOSITION



(57) Abstract: An aqueous shear-thinning personal care composition comprises : (a) at least 3 wt% of a sulfonate compound of general formula (I): where R¹ represents a C₈₋₂₂ alkyl group; R² represents a C₁₋₄ alkyl group; each of R³, R⁴ and R⁵ independently represents a hydrogen atom or a C₁₋₄ alkyl group; and M⁺ represents a sodium or potassium cation; (b) at least 3 wt% of an amphoteric or zwitterionic surfactant; (c) at least 10 wt% of water; and (d) at least 0.5 wt% of an additional component in the form of particles or droplets suspended in the composition. The composition is stable in an aqueous environment and can be formulated with a wide range of additional components.

COMPOSITION

This invention relates to a personal care composition, for example a liquid or gel body wash, shower gel, handwash, skin cream, hair conditioner or moisturiser.

US 5,612,307 discloses an aqueous liquid cleansing and moisturising composition comprising: a) a surface active agent selected from anionic, nonionic, zwitterionic and cationic, surface active agents, soap and mixtures thereof; and b) a benefit agent;

wherein the benefit agent and surface active agent are separate but combinedly dispensable from a single packaging means in a predetermined ratio as discrete domains, the domains having one dimension of at least about 1000 microns.

An advantage of the invention of US 5,612,307 is said to be that it leads to improved deposition of benefit agents from a surface active agent containing aqueous liquid composition during use. The surface active agent and benefit agent are separated in the composition, i.e. they do not directly contact one another in the composition. This avoids adverse interactions which may occur between these two components and which may result in ineffective deposition of the benefit agent.

The composition of US 5,612,307 is said to be suitable for cleansing and "moisturising", "conditioning" or "protection" of the skin. The benefit agent is included in the composition to moisturise, condition and/or protect the skin. By "benefit agent" is meant a substance that softens the skin (stratum corneum) and keeps it soft by

retarding the decrease of its water content and/or protects the skin.

Preferred benefit agents of US 5,612,307 include:
5 silicones and siloxanes; fats and oils; waxes; hydrocarbons; higher fatty acids; higher alcohols; esters; essential oils; lipids; vitamins; sunscreens; and phospholipids.

10 Where adverse interactions between the benefit agent and surface active are likely to be particularly acute, US 5,612,307 states that the benefit agent may be incorporated in the compositions in a carrier.

15 The surface active agent of US 5,612,307 is preferably present at a level of from 1 to 35 wt %, preferably 3 to 30 wt %.

One preferred anionic detergent for use in the invention
20 of US 5,612,307 is a fatty acyl ethane sulfonate of formula $\text{RCO}_2\text{CH}_2\text{CH}_2\text{SO}_3\text{M}$, where R is an alkyl or alkenyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably at least three quarters of the RCO
25 groups have 12 to 18 carbon atoms and may be derived from coconut, palm or a coconut/palm blend.


It is also preferable in US 5,612,307 that the composition includes from 0.5 to 15 wt % of a cosurfactant agent with
30 skin-mildness benefits. Suitable materials are zwitterionic detergents identified in US 5,612,307.

A favoured fatty acyl ethane sulfonate is "SCI" (sodium cocyl isethionate). SCI is commercially available and is a successful product but the scope it offers formulators is limited by its low solubility in water; its low hydrolytic stability, and the lack of stability in its pH value, over time.

The present invention is based on the use of a different fatty acyl sulfonate salt, and its advantageous co-
10 formulation with amphoteric or zwitterionic surfactants, in aqueous environments.

In accordance with a first aspect of the present invention there is provided an aqueous shear-thinning personal care
15 composition comprising:

(a) at least 3 wt% of a sulfonate compound of general formula



20

where R¹ represents a C₈₋₂₂ alkyl group;

R² represents a C₁₋₄ alkyl group;

each of R³, R⁴ and R⁵ independently represents a hydrogen
25 atom or a C₁₋₄ alkyl group; and

M⁺ represents a cation;

30

(b) at least 3 wt% of an amphoteric or zwitterionic surfactant;

(c) at least 10 wt% of water; and

(d) at least 0.5 wt% of an additional component in the form of particles or droplets suspended in the composition.

5 There may be further components present in the composition. Percentages given herein are percentages on total weight of the composition, including any such further components, for example when there is a plurality of compounds (a); and/or a plurality of surfactants (b).

10

Many additional components are described hereafter, some of which may go into solution and some of which remain in the composition as particles or droplets. The stipulation of the definition of the invention is that there must be
15 at least one additional component which is in the form of particles or droplets suspended in the composition. The compositions of the present invention have an exceptional ability to be formulated with further components and in particular to suspend particles or droplets and we regard
20 this as an essential element of the invention.

Preferably R^1 represents a C_{10-20} alkyl group, most preferably a C_{12-18} alkyl group.

25 Preferably R^2 represents a C_{1-4} alkyl group in which a propyl or butyl group is straight-chain. Preferably R^2 represents an n-propyl, ethyl or, most preferably, a methyl group.

30 Preferably R^3 represents a hydrogen atom.

Preferably one of R^4 and R^5 represents a hydrogen atom and the other represents a hydrogen atom or a C_{1-4} alkyl group,

Preferably one of R⁴ and R⁵ represents a hydrogen atom or a C₁₋₄ alkyl group in which a propyl or butyl group is straight-chain. Preferably one of R⁴ and R⁵ represents an n-propyl, ethyl or methyl group or, most preferably, a hydrogen atom. Most preferably both R⁴ and R⁵ represent hydrogen atoms.

Preferably M⁺ represents an ammonium cation or, most preferably, a metal cation. Suitable metal cations include alkali metal cations, for example sodium and potassium cations, and alkaline earth metal cations, for example calcium and magnesium cations. Preferably M⁺ represents a potassium cation, or, especially, a sodium cation.

15


Of course, a plurality of compounds (a) may be present.

Component (b) may include amphoteric or zwitterionic surfactants selected from betaines, for example alkyl betaines, alkylamidopropyl betaines, alkyamidopropyl hydroxy sultaines, alkylampho acetates, alkylamphodi-acetates, alkylamphopropionates, alkylamphodipropionates, alkyliminodipropionates and alkyliminodiacetate.

25 Of course, a plurality of surfactants (b) may be present.

Amphoteric or zwitterionic surfactants may include those which have an alkyl or alkenyl group of 7 to 22 carbon atoms and comply with an overall structural formula:


30



where R¹ is alkyl or alkenyl of 7 to 22 carbon atoms; R² and R³ are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 6 carbon atoms; m is 2 to 4; n is 0 or 1; X is alkylene of 1 to 6 carbon atoms optionally substituted with hydroxyl; and Y is -CO₂ or -SO₃.


Amphoteric or zwitterionic surfactants may include simple betaines of formula:

10



and amido betaines of formula:


15




where m is 2 or 3.

In both formulae R¹, R² and R³ are as defined previously. R¹ may, in particular, be a mixture of C₁₂ and C₁₄ alkyl groups derived from coconut so that at least half, preferably at least three quarters, of the groups R¹ has 10 to 14 carbon atoms. R² and R³ are preferably methyl.

Amphoteric or zwitterionic surfactants may include sulphobetaines of formula:




or



5


where m is 2 or 3, or variants of these in which
 $-\text{(CH}_2\text{)}_3\text{SO}_3^-$ is replaced by



10


where R^1 , R^2 and R^3 in these formulae are as defined
 previously.

15 Amphoteric or zwitterionic surfactants may include
 amphoacetates and diamphoacetates. Amphoacetates
 generally conform to the following formula:



20

Diamphoacetates generally conform to the following
 formula:



where R is an aliphatic group of 8 to 22 carbon atoms and M is a cation such as sodium, potassium, ammonium, or substituted ammonium.

5

Compounds such as oleylbetaine and cocoamidopropylhydroxy-sultaine are regarded as very safe for the human body and for the eyes. Also preferred in some embodiments are sodium lauroamphoacetate, sodium cocoamphoacetate, 10 disodium lauroamphoacetate, and disodium cocoampho-diacetate.

Preferably, in a composition in accordance with the present invention, component (a) is present in an amount 15 of up to 30% by total composition weight.

Preferably in a composition in accordance with the present invention, component (b) is present in an amount of up to 30% by total composition weight.

20

Preferably in a composition in accordance with the present invention, component (c) is present in an amount of from 20% to 95% by total composition weight, preferably from 50 to 90%.

25

In one embodiment the composition of the present invention is a free-flowing liquid.

In one embodiment a composition in accordance with the 30 invention is a shear thinning composition.

A composition in accordance with the present invention may suitably have a viscosity of from 250 to 300,000 mPa.s,

measured at 25°C using a Brookfield rotational viscometer equipped with an appropriate spindle at a rotation speed of from 0.1 revolutions per minute (rpm) to about 60 rpm. Examples of appropriate spindles include the Brookfield
5 RVT Viscometer Helipath T-Bar Spindles B and C (used in the examples which follow). At low shear rates (rotation speeds) the viscosity as measured may be up to 300,000 mPa.s. At high shear rates (rotation speeds) the viscosity as measured may be as low as 250 mPa.s.

10

The compositions of the invention may be pourable liquids or semi-liquids e.g..

The compositions of the invention may be soft gels and may
15 suitably be of viscosity in the range 800 to 20,000 mPa.s, measured under the conditions stated above.

The compositions of the invention may be firmer gels and may be of viscosity in the range 3000 to 100,000 mPa.s,
20 measured under the conditions stated above.

Surprisingly, the aqueous compositions of the invention have desirable non-Newtonian properties and excellent suspending properties, being capable of maintaining solid
25 or non-solid water-insoluble particles or droplets in suspension. Under the conditions of testing which have been applied in relation to the invention, no sedimentation or separation whatsoever of the water-insoluble particles or droplets has developed over time.
30 Furthermore the compositions of the invention produce an excellent rich creamy foam which is stable, and is mild to the skin and eyes.

Without being bound by theory, it is believed possible that the following microstructural conditions are achieved in the present invention.

5 The compositions of the present invention are believed to exhibit a so-called lamellar phase structure comprising spherulites in suspension.

By "lamellar phase" is intended a hydrated solid phase or
10 a liquid crystal phase in which several double layers are arranged in a parallel network, separated by layers of water or of an aqueous solution. In respect of the spherulites, these are multilamellar vesicles comprising several layers of surface-active agents arranged
15 concentrically and generally ranging from 0.1 to 50 microns in size.

More specifically, the corresponding phase structure may, we believe, be a birefringent solution, optically
20 characteristic of a non-isotropic mesomorphic phase. By the term "water-insoluble particles" are intended solid or non-solid entities which are not solubilized in the aqueous medium of the subject composition. These, more particularly, are solid particles or suspended droplets.

25 However the more significant point is not the microstructure per se but the fact that the components together produce a composition with excellent overall or macroscopic properties.

30 Important advantages of the compositions of the invention over many other compositions, and in over particular over compositions containing fatty acyl ethane sulfonates, such

as SCI, is that they are hydrolytically stable and pH-stable, and that they can be formulated with many further ingredients, including particulates and droplets, without adverse interactions being found.

5

The composition may contain an electrolyte. An electrolyte is suitably present in an amount of from 0.5% to 25%, preferably 1% to 15%, preferably 2 to 10%, by total composition weight. The concentration thereof may be
10 adjusted to produce the desired formulation.

Suitable as an electrolyte are ionic compounds, for example metal salts selected from sodium chloride, sodium sulfate, potassium chloride, potassium sulfate, sodium
15 phosphate, disodium phosphate, potassium phosphate, dipotassium phosphate, sodium isethionate, sodium methyl isethionate and sodium lactate; and non-metallic salts, for example sodium citrate, ammonium chloride, ammonium citrate and ammonium lactate.

20

Preferred electrolytes include sodium and ammonium chloride.

Said additional component (d) is suitably one or more
25 benefit agents in the form of particles or droplets. By "benefit agent" we can adopt the definition of US 2003/0180246A, that is, a "benefit agent" is any active ingredient that is to be delivered into the skin or hair, or onto the skin or hair, or both, at a desired location.
30 For example, a benefit agent may be a substance that softens the skin (stratum corneum) and keeps it soft by retarding the decrease of its water content; and/or a substance which protects the skin; and/or an exfoliating

material, for example a water insoluble particulate or suspended abrasive.

Preferred benefit agents include:

- 5 • a) silicone oils, gums and modifications thereof such as linear and cyclic polydimethylsiloxanes, amino, alkyl alkylaryl and aryl silicone oils;
- b) fats and oils including natural fats and oils such as jojoba, soyabean, rice bran, avocado, almond,
10 olive, sesame, persic, castor, coconut, mink oils; cacao fat, beef tallow, lard; hardened oils obtained by hydrogenating the aforementioned oils; and synthetic mono, di and triglycerides such as myristic acid glyceride and 2-ethylhexanoic acid glyceride;
- 15 • c) waxes such as carnauba, spermaceti, beeswax, lanolin and derivatives thereof;
- d) hydrophobic plant extracts;
- e) hydrocarbons such as liquid paraffins, petroleum jelly, microcrystalline wax, ceresin, squalene,
20 squalane, and mineral oil;
- f) higher fatty acids such as lauric, myristic, palmitic, stearic, behenic, oleic, linoleic linolenic, lanolic, isostearic and poly unsaturated fatty acids (PUPA) acids;
- 25 • g) higher alcohols such as lauryl, cetyl, steryl, oleyl, behenyl, cholesterol and 2-hexadecanol alcohol;
- h) esters (fatty acid esters, fatty alcohol esters) including cetyl octanoate, myristyl lactate, cetyl
30 lactate, isopropyl myristate, myristyl myristate, isopropyl palmitate, isopropyl adipate, butyl stearate, decyl oleate, cholesterol isostearate, glycerol monostearate, glycerol distearate, glycerol

tristearate, alkyl lactate for example lauryl lactate, alkyl citrate and alkyl tartrate;

- i) essential oils such as fish oils, fragrance oils, mentha, jasmine, camphor, white cedar, bitter orange peel, ryu, turpentine, cinnamon, bergamont, citrus unshiu, calamus, pine, lavender, bay, clove, hiba, eucalyptus, lemon, starflower, thyme, peppermint, rose, sage, menthol, cineole, eugenol, citral, citronelle, borneol, linalool, geraniol, evening primrose, camphor, thymol, spirantol, pinene, limonene and terpenoid oils;
- j) lipids such as cholesterol, ceramides, sucrose esters and pseudo-ceramides as described in European Patent Specification No. 556 957;
- k) vitamins such as vitamin A and E, and vitamin alkyl esters, including those vitamin C alkyl esters;
- l) sunscreens such as octyl methoxyl cinnamate (Parsol MCX) and butyl methoxy benzoylmethane (Parsol 1789);
- m) phospholipids; and
- n) one or more of scrubbing beads, anti-dandruff agents, anti-UV actives, mica, silica, pigments, natural and synthetic waxes, zinc oxide, titanium dioxide.

25

Other examples of suitable benefit agents include: depigmentation agents; reflectants; UV absorbers, thickening agents; detangling/wet combing agents; film forming polymers; humectants; amino acids and their derivatives; antimicrobial agents; anti-acne agents; anti-aging agents; antiseptics; analgesics; local anesthetics; anti-hair loss agents; hair growth inhibitor agents; inflammation inhibitors; proteins; deodorants and anti-

30

perspirants; agents for treatment of dandruff, seborrheic dermatitis and psoriasis; skin emollients and skin moisturizers; hair conditioners; hair softeners; hair moisturizers; vitamins; tanning agents; skin lightening agents; antifungals such as antifungals for foot preparations; depilating agents; counterirritants; hemorrhoidals; insecticides; pigments or opacifying agents, moisturizing beads, natural abrasives, synthetic abrasives such as polyoxyethylene beads, mineral oils, petrolatum, silicone oils, polyalkylsiloxanes, polyalkarylsiloxanes, sunscreens, reflectants and the like; and mixtures thereof.

Examples of suitable reflectants include mica, alumina, calcium silicate, glycol dioleate, glycol distearate, silica, sodium magnesium fluorosilicate, and mixtures thereof.

Examples of suitable UV absorbers include benzophenone, bornelone, PABA (Para Amino Benzoic Acid), butyl PABA, cinnamidopropyl trimethyl ammonium chloride, disodium distyrylbiphenyl disulfonate, potassium methoxycinnamate, and mixtures thereof.

The benefit agent (or benefit agents, in total) is preferably present in amount of from 0.1 to 50%, preferably from 1 to 40%, preferably from 1.5 to 30%, and most preferably from 2 to 20% by weight based on the composition weight.

30

A structurant may be added to the phase comprising the surface active agent, suitably in an amount of from 0.01%,

preferably from 0.1%, more preferably from 1%, by weight based on the composition weight.

A structurant may be added to the phase comprising the surface active agent, suitably in an amount of up to 15%, preferably up to 12%, more preferably up to 10%, and most preferably up to 5%, by weight based on the composition weight.


The primary purpose of a structurant is to assist in the formation of a lamellar phase. It is believed that a lamellar phase, if achieved, may enable the compositions to suspend particles more readily while still maintaining good shear thinning properties. The lamellar phase also provides consumers with desired rheology ("heaping").

Suitable structurants include swelling clays, for example laponite; fatty acids and derivatives thereof, in particular, fatty acid esters; cross-linked polyacrylates such as Carbopol (TM) (polymers available from Goodrich); acrylates and copolymers thereof; polyvinylpyrrolidone and copolymers thereof; polyethylene imines; salts such as sodium chloride and ammonium sulphate; sucrose esters; gellants; and mixtures thereof. Examples of fatty acids which may be used are C₁₀-C₂₂ acid (e.g. lauric, oleic etc.), isostearic acid, linoleic acid, linolenic acid, ricinoleic acid, elaidic acid, arichidonic acid, myristoleic acid and palmitoleic acid. Ester derivatives include propylene glycol isostearate, propylene glycol oleate, glyceryl isostearate, glyceryl oleate and polyglyceryl diisostearate. Monoglyceride polyglycol ethers are suitable structurants.

Of the clays, particularly preferred are synthetic hectorite (laponite) clay used in conjunction with an electrolyte salt capable of causing the clay to thicken. Suitable electrolytes include alkali and alkaline earth salts such as halides, ammonium salts and sulphates.

The surface active agent phase may also comprise a thickening agent, ie a material which maintains the viscosity of this phase as the shear rate thereof is increased during use. Suitable materials include cross-linked polyacrylates such as Carbopol (TM) (polymers available from Goodrich); natural gums including alginates, guar, xanthan and polysaccharide derivatives including carboxy methyl cellulose and hydroxypropyl guar; propylene glycols and propylene glycol oleates; salts such as sodium chloride and ammonium sulphate; glycerol tallowates; and mixtures thereof.

Thickeners may also be added to the benefit agent in order to achieve the required viscosity during use. Preferred thickeners for the benefit agent include fumed silica; polyethylene; alkyl silicone waxes; aluminium silicate; lanesterol; natural and synthetic waxes; fatty acids and derivatives thereof, in particular, fatty acid monoglyceride polyglycol ethers; higher fatty alcohols; petrolatum; narogel; polyammonium stearate; hydrotalcites; and mixtures thereof. Hydrotalcites are materials of general formula



where M is a divalent metal ion e.g. Mg^{2-} ; N is a trivalent metal ion e.g. Al^{3+} ; X is an exchangeable anion

e.g. CO_3^- , NO_3^- ; stearate, cinnamate; m is the number of divalent metal ions; and n is the number of trivalent metal ions.

5 Furthermore, the benefit agent may also function as a carrier to deliver efficacy agents to skin treated with the compositions of the invention. This route is particularly useful for delivering efficacy agents which are difficult to deposit onto the skin or those which
10 suffer detrimental interactions with other components in the composition. In such cases the carrier is often a silicone or hydrocarbon oil which is non solubilised/micellised by the surface active phase and in which the efficacy agent is relatively soluble. Examples
15 of such efficacy agents include anti-viral agents; hydroxycaprylic acids; pyrrolidone; carboxylic acids; 3,4,4'-trichlorocarbanilide; benzoyl peroxide; perfumes; essential oils; germicides and insect repellants such as 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300);
20 salicylic acid; willow extract, N,N-dimethyl m-toluamide (DEET); and mixtures thereof.

The composition may contain one or more additional anionic surface-active agents, in addition to the acyl sulphonate
25 salt defined above; preferably selected from salts of fatty acids; alkali metal salts of mono- or dialkyl sulfates; mono- or dialkyl ether sulfates; lauryl ether sulfates; alkyl sulfonates; alkyl aryl sulfonates; primary alkane disulfonates; alkene sulfonates; hydroxyalkane
30 sulfonates; alkyl glyceryl ether sulfonates; alpha-olefinsulfonates; alkyl phosphates; sulfonates of alkylphenolpolyglycol ethers; salts of alkyl sulfopolycarboxylic acid esters; alkyl sulfosuccinates and

salts thereof, alkyl ether sulfosuccinates and salts thereof, non-acylated alkyl isethionates; fatty acid taurates; acyl taurates; products of condensation of fatty acids with oxy- and aminoalkanesulfonic acids; sulfated derivatives of fatty acids and polyglycols; alkyl and acyl 5 sarcosinates; sulfoacetates; alkyl phosphates; alkyl phosphate esters; acyl lactates; alkanolamides of sulfated fatty acids and salts of lipoamino acids. Particularly exemplary salts of the above, where applicable, are the 10 sodium, potassium, ammonium, magnesium and triethylamine salts.

In the case of sodium lauryl ether sulfate, LESNa, this is a surface-active agent widely used in shampoos by reason 15 of its foaming properties. It is a cheap, conventional, washing base.

The composition may contain one or more non-ionic surface-active agents, preferably selected from the following: 20 reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene oxide (for example alkyl (C₆-C₂₂) phenol-ethylene 25 oxide condensates, the condensation products of aliphatic (C₈ -C₁₈) primary or secondary linear or branched alcohols with ethylene oxide, and products made by condensation of ethylene oxide with the reaction products of propylene oxide and ethylenediamine); long chain tertiary amine 30 oxides, long chain tertiary phosphine oxides and dialkyl sulphoxides; alkyl amido amine oxides; alkyl tertiary phosphine oxides; alkoxyalkyl amines; sorbitan; sorbitan esters; sucrose esters; sugar amides, such as a

polysaccharide amide; lactobionamides; and alkyl polysaccharide nonionic surfactants, for example alkylpolyglycosides. Preferred non-ionic surfactants are sucroglycerides and ethoxylated fatty alcohols, especially those derived from lauryl, cetylstearyl, stearyl, cetyl and oleocetyl alcohols.

A non-ionic surface active agent, when present, preferably comprises up to 15% wt of the composition, preferably up to 10% wt.

Representative solid particles include active materials or agents such as those used for hair treatments, e.g., zinc pyrithione, or any abrasive materials which may be of natural or synthetic origin. In particular, these are polycarbonates, polypropylenes, polyethylenes, alumina, calcite and clays. Such particles generally have a crystal size ranging from about 1 to 600 microns and preferably from about 10 to 400 microns.

In the case of the suspended droplets, these are preferably droplets of at least one vegetable oil, essential oil and/or, more particularly, silicone oil.

The silicone oils which are well suited according to the present invention include the polyalkylsiloxanes, polyalkylarylsiloxanes and mixtures thereof. The preferred polyalkylsiloxanes are, especially, polydimethylsiloxanes such as dimethicone whose viscosity may range from about 20 mPa.s to 50 Pa.s at 25°C, pure or mixed with cyclomethicone. Particularly exemplary of the polyalkylarylsiloxanes are the polyphenyldimethylsiloxanes.

In particular, polydiorganosiloxanes such as polydimethylsiloxanes having a molecular weight of less than or equal to 3,000,000 and polydimethyl-
5 diphenylsiloxanes of molecular weight of about 600,000 are especially well suited according to the invention. The size of the oil droplets within the compositions of the invention advantageously ranges from about 0.5 to 50 microns.

10

The formulations according to the invention may contain approximately 0.5% to 8% and, preferably, about 1.5% to 4.5% by weight of water-insoluble particles.

15 Specific formulations of the invention may include these:

- (a) as additional anionic agent LESNa and as co-surface-active agent lauryl alcohol containing two moles of ethylene oxide per mole of lauric alcohol and as electrolyte NaCl in a lauryl alcohol 2EO/LESNa
20 mole ratio of from 2.1 to 4.3,
- (b) as additional anionic agent, a sodium lipoamino acid and the lipoproteol LC0® mixed with the same lauryl alcohol containing two moles of ethylene oxide per mole of lauric acid and having a lauryl alcohol
25 2EO/sodium lipoamino acid mole ratio ranging from about 1.2 to 6.7,
- (c) as additional anionic agent LESNa, and as co-surface-active agent oleylbetaine and as electrolyte NaCl, in an oleylbetaine/LESNa mole ratio of from 1.1
30 to 3.7.

Generally, the compositions according to the invention contain an anionic surface-active agent mixed with at least one cosurface-active agent in a cosurface-active agent/anionic surface-active agent mole ratio equal to or
5 greater than 1.

A benefit of the invention is the ability to suspend oil and/or emollient particles in a non-Newtonian composition. The following oil/emollients may optionally be suspended
10 in the compositions of the invention. Various classes of oils are set forth below.

Vegetable oils: Arachis oil, castor oil, cocoa butter, coconut oil, corn oil, cotton seed oil, olive oil, palm
15 kernel oil, rapeseed oil, safflower seed oil, sesame seed oil and soybean oil.

Esters: Butyl myristate, cetyl palmitate, decyloleate, glyceryl laurate, glyceryl ricinoleate, glyceryl stearate,
20 glyceryl isostearate, hexyl laurate, isobutyl palmitate, isocetyl stearate, isopropyl isostearate, isopropyl laurate, isopropyl linoleate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, propylene glycol monolaurate, propylene glycol ricinoleate, propylene
25 glycol stearate, and propylene glycol isostearate.

Animal Fats: acetylated lanolin alcohols, lanolin, lard, mink oil and tallow.

30 Other examples of oil/emollients include mineral oil, petrolatum, silicone oil such as dimethyl polysiloxane, lauryl and myristyl lactate.

The emollient/oil is generally used in an amount from about 0 to 20%, preferably 0 to 15% by wt. of the composition.

5 In addition, the compositions of the invention may include optional ingredients as follows:

- 10 o Organic solvents, such as ethanol; auxiliary thickeners, sequestering agents, such as tetrasodium ethylenediaminetetraacetate (EDTA), EHDP or mixtures in an amount of 0.01 to 1%, preferably 0.01 to 0.05%; and coloring agents, opacifiers and pearlizers such as zinc stearate, magnesium stearate, TiO₂, EGMS (ethylene glycol monostearate) or Lytron 621
15 (Styrene/Acrylate copolymer); all of which are useful in enhancing the appearance or cosmetic properties of the product.
- 20 o Antimicrobials such as 2-hydroxy-4,2'4' trichloro-diphenyl-ether (DP300); preservatives, for example dimethyloldimethyl-hydantoin (Glydant XL1000), parabens, sorbic acid etc.
- 25 o Coconut acyl mono- or diethanol amides as suds boosters, and strongly ionizing salts such as sodium chloride and sodium sulfate may also be used to advantage.
- 30 o Antioxidants such as, for example, butylated hydroxytoluene (BHT) may be used advantageously in amounts of about 0.01% or higher if appropriate.

- o Cationic conditioners which may be used include Quatrisoft LM-200 Polyquaternium-24, Merquat Plus 3330-Polyquaternium 39; Jellner Polyquaternium 10; Merquat 550; and Jaguar® type conditioners.
- 5
- o Deflocculating polymers.
 - o Chelating agents including but not limited to EDTA, EDTA salts, NTA, methylenediamine disuccinate and
- 10 salts of methylenediamine disuccinate.

Compositions of the invention may be formulated as products for washing the skin, for example, bath or shower gels, hand washing compositions or facial washing liquids; pre- and post-shaving products; rinse-off, wipe-off and

15 leave-on skin care products; products for washing the hair and for dental use.

In accordance with a second aspect of the present invention there is provided the use of a composition of the invention, as defined above, in the formulation of a personal care product with improved stability, in an aqueous environment.

20

In accordance with a third aspect of the present invention there is provided the use of a composition of the invention, as defined above, in the formulation of a personal care product which has improved stability in an aqueous environment compared with a corresponding compound

25

30 in which each of R^2 to R^5 is a hydrogen atom.

Suitably the composition, and in particular the defined sulfonated compound, is resistant to breakdown in an

aqueous environment. Preferably it has essentially no effect on the pH of the product, over time. It has excellent lathering properties and is very mild to skin and eyes.

5

Weight percentage values herein refer to the total complement of compounds of a named type, e.g. amphoteric or zwitterionic surfactants in total; or benefit agents in total.

10

Where a percentage value is given for a component it refers to the active content of that component.

The invention will now be further described, by way of example, with reference to the following examples.

Examples 1 to 5 were carried out to assess the suitability of certain isethionate salts for use in the present invention. Examples 6 - 14 relate to certain compositions which are within the scope of the present invention.

In the examples abbreviations used are as follows:

SCMI - sodium cocoylmethylisethionate

SCI - sodium cocoyl isethionate

25 SLS - sodium lauryl sulfate

SLMI - sodium lauroylmethylisethionate

SLES - sodium laureth sulfate

SCEI - sodium cocoyl ethyl isethionate

CAPB - cocamidopropyl betaine

30 CMEA - cocamide monoethanolamine.

Example 1

In a preliminary test the stability of SCMI in 10 wt% aqueous solutions adjusted to have initial pH values respectively of 4.5, 5.5, 6.5, 7.5, 8.5 and 9.5, each held at 48.9°C for an extended period of days, were tested for the amount of SCMI present over the test period.

The pH values were read using a standard electrode pH meter (Fisher Accumet XL-25). The electrode was placed in the composition (no further dilution) and the pH recorded.

The results are shown in Fig. 1. It can be seen that the percentage of SCMI does not change from Day 0 to Day 30, at any of these pH values.

Example 2

In a preliminary test 10 wt% solutions of SCMI, adjusted to have initial pH values respectively of 4.5, 5.5, 6.5, 7.5, 8.5 and 9.5, each held at 48.9°C for an extended period of days, were tested for their pH stability over the extended period. The pH testing was as described above.

25

The results are shown in Fig. 2. It can be seen that the pH values were substantially unchanged from Day 0 to Day 30 at all these pH values, except for a small pH reduction in the first few days, for the sample held at the highest pH value, 9.5.

Example 3

In a preliminary test based on the Zein protocol the skin irritancy properties of a number of compounds were tested, including two sulfate surfactants SLS and SLES (known to have significant irritancy), distilled water, SCI (known to have low irritancy) and SCMI. The results are shown in Fig. 3. SCMI and SLMI were shown to have low irritancy, similar to that shown by SCI; and significantly lower than the sulfates.

The Zein protocol is an in-vitro test to measure skin irritancy of formulations. The method makes use of the correlation between binding ability of surfactants to proteins and the damage the surfactant causes to the skin. The de-naturation of epidermal protein is a key mechanism in the development of observable damage to the skin by surfactants. Zein protein, an insoluble protein extracted from corn kernel, is used as a model for epidermal protein and the solubility of the Zein protein in surfactant solutions is believed to be a reliable guide for the skin irritancy caused by the surfactant. The test involves establishing the amount of Zein protein which can be solubilised by surfactant. 5 g of Zein protein may be dispersed in 40 cm³ of surfactant solution (at 2 wt% concentration). The mixture is shaken for 1 hour at 35°C, and immediately centrifuged to remove any non-solubilised Zein. The amount solubilised is estimated from the solution nitrogen content using micro-Kjeldahl method (making allowance for any nitrogen in the compound tested).

Example 4

Simple solubility tests were carried out.

10 wt% of SCI shaken with deionised water gave a milky
5 white suspension. The solubility limit of SCI in water is
quoted as 0.5 wt%.

10 wt% of SCMI added to deionised water immediately gave a
clear solution.

10 wt% of SCEI added to deionised water immediately gave a
10 clear solution.

Example 5

Foaming tests were performed using a capped measuring
15 cylinder which was charged with an aliquot of the stated
surfactant leaving room for foam to form. Foam heights in
the graduated cylinder were measured at the start and
after 10 inversions of the cylinder. The operation was
carried out in the same manner for each surfactant.

20

100% SLES gave a foam of medium height but which was also
open and "watery".

80 wt% SLS / 20 wt% CAPB gave a foam which was high but
25 open and "watery".

100% SCMI gave a foam of medium height but which was rich
and creamy.

30 80% SCMI / 20% CAPB gave a foam of medium height but which
was rich and creamy.

Examples 6 to 14

Formulated compositions containing canola oil as benefit agent were prepared and tested for viscosity and physical stability over a one-month period. The results are set out in the table below.

The formulations of Examples 15 and 16 were prepared by progressive mixing and blending of the components, in the order given. The pH was adjusted to between 5.2 and 5.8 using citric acid after blending of the preceding
5 components (in the respective tables), but before blending in of the fragrance, dye and preservative.

Example 15

The following moisturizing and conditioning shampoo was prepared.

Ingredients	Tradename (Supplier)	%w/w
Water		to 100
Trisodium		
Ethylenediamine	Natrlquest E30	
disuccinate	(Innospec)	0.15
Sodium Chloride		4.00
Guar		
Hydroxypropyltrimonium	Activsoft C-17	
Chloride	(Innospec)	0.2
Disodium	Miranol C2M Conc.	
Cocoamphodipropionate	(Rhodia)	7.5
Sodium Methyl Cocoyl	Pureact WS Conc.	
Taurate	(Innospec)	7.5
Guar Gum	Activsoft S (Innospec)	0.5
Glycerin		1.00
	Pureact SLMI-85	
SLMI	(Innospec)	12.5
	Aminol CM Flakes	
Cocamide MEA	(Innospec)	3.5
Quaternium 75	Condicare CT (Innospec)	2.00
	Dow Corning 200 Fluid	
Dimethicone	(Dow Corning)	4.00
	Mirataine BET C-30	
Cocamidopropyl Betaine	(Rhodia)	10.00
		pH 5.2 -
Citric Acid (50% soln)		5.8
Preservative		trace
Fragrance		trace
Dye(s)		trace

The viscosity of the conditioning shampoo of Example 15 was tested under different shear rates, on a viscometer with a T-Bar C Spindle, at 22°C. The results were as follows:

5	0.5 RPM = 160,000cps
	1.0 RPM = 90,000cps
	2.5 RPM = 42,400cps

Example 16

The following foaming body lotion was prepared.

Ingredients	Tradename (Supplier)	%w/w to 100
Water		to 100
Trisodium		
Ethylenediamine	Natrlquest E30	
disuccinate	(Innospec)	0.15
Sodium Chloride		3.6
Guar		
Hydroxypropyltrimonium	Activsoft C-17	
Chloride	(Innospec)	0.18
Disodium	Miranol C2M Conc.	
Cocoamphodipropionate	(Rhodia)	6.7
Sodium Methyl Cocoyl	Pureact WS Conc.	
Taurate	(Innospec)	6.7
Guar Gum	Activsoft S (Innospec)	0.45
Glycerin		1.00
	Pureact SLMI-85	
SLMI	(Innospec)	11.2
	Aminol CM Flakes	
Cocamide MEA	(Innospec)	3.1
Canola Oil	Rita Canola Oil (Rita)	9.00
	Mirataine BET C-30	
Cocamidopropyl Betaine	(Rhodia)	10.00
Trideceth-7 Carboxylic		
Acid	Pureact TA (Innospec)	3.5
		to pH 5.2 -
Citric Acid (50% soln)		5.8
Preservative		trace
Fragrance		trace
Dye(s)		trace

The viscosity of the foaming body lotion of Example 16 was tested under different shear rates, on a viscometer with a T-Bar C Spindle, at 22°C. The results were as follows:


5	0.5 RPM = 240,000cps
	1.0 RPM = 140,000cps
	2.5 RPM = 64,000cps

CLAIMS

1. An aqueous shear-thinning personal care composition comprising:

5

(a) at least 3 wt% of a sulfonate compound of general formula



10

where R¹ represents a C₈₋₂₂ alkyl group;

R² represents a C₁₋₄ alkyl group;

each of R³, R⁴ and R⁵ independently represents a hydrogen atom or a C₁₋₄ alkyl group; and

15

M⁺ represents a cation;

(b) at least 3 wt% of an amphoteric or zwitterionic surfactant;

20

(c) at least 10 wt% of water; and

(d) at least 0.5 wt% of an additional component in the form of particles or droplets suspended in the composition.

25

2. A composition as claimed in claim 1 wherein R² represents a methyl group.

3. A composition as claimed in claim 1 or 2 wherein each of R³, R⁴ and R⁵ represents a hydrogen atom.

30

4. A composition as claimed in any preceding claim wherein component (a) is present in an amount of up to 30 wt%.
5. A composition as claimed in any preceding claim wherein component (b) is present in an amount of up to 30 wt%.
6. A composition as claimed in any preceding claim wherein component (c) is present in an amount of from 20% to 95% by total composition weight.
7. A composition as claimed in any preceding claim wherein component (d) is present in an amount from 0.1 wt% to 50 wt%.
8. A composition as claimed in any preceding claim, wherein said additional component (d) is a benefit agent.
9. A composition as claimed in any preceding claim, wherein the composition further includes a benefit agent which is in solution in the composition.
10. A composition as claimed in any preceding claim, including an electrolyte.
11. A composition as claimed in claim 8 wherein the electrolyte is present in an amount from 0.5 wt% to 25 wt%.
12. A composition as claimed in any preceding claim, formulated as a product selected from the following: a bath or shower gel; a hand washing composition; a facial washing composition; a pre-, during- or post-shaving product; a skin moisturiser; a hair gel; a medicinal skin treatment product; a shampoo; a hair conditioner; a dental product.


13. Use of a composition as claimed in any preceding claim, in the formulation of a personal care product which is stable in an aqueous environment.

5

14. Use of a composition as claimed in any of claims 1 to 12, in the formulation of a personal care product which has improved stability in an aqueous environment compared with a corresponding compound in which each of R^2 to R^5 is a
10 hydrogen atom.

1/3

Figure 1
Sodium Acyl Alkyl Isethionate
Activity @ 48.9°C



2/3

Figure 2
10% Sodium Acyl Alkyl Isethionate pH Stability @ 48.9°C




Figure 3
14 Day Cumulative Skin Irritation
0.50% Active
Mean Cumulative Score

