

[19] Patents Registry
The Hong Kong Special Administrative Region
香港特別行政區
專利註冊處

[11] 1249050 B
EP 3266449 B1

[12] **STANDARD PATENT (R) SPECIFICATION**
轉錄標準專利說明書

[21] Application no. 申請編號
18108881.5

[22] Date of filing 提交日期
09.07.2018

[51] Int. Cl.
A61K 9/20 (2006.01) A61K 9/28 (2006.01)
A61K 9/48 (2006.01) A61K 45/06 (2006.01)
A61K 9/00 (2006.01) A61K 47/40 (2006.01)
A61K 31/40 (2006.01) A61K 31/41 (2006.01)
A61K 31/401 (2006.01) A61P 15/08 (2006.01)

[54] ORAL FORMULATIONS OF PYRROLIDINE DERIVATIVES
吡咯烷衍生物的口服製劑

[30] Priority 優先權
17.12.2013 EP 13197606
[43] Date of publication of application 申請發表日期
26.10.2018
[45] Date of publication of grant of patent 批予專利的發表日期
17.01.2020
EP Application no. & date 歐洲專利申請編號及日期
EP 17188131.1 15.12.2014
EP Publication no. & date 歐洲專利申請發表編號及日期
EP 3266449 10.01.2018
Date of grant in designated patent office 指定專利當局批予專利日期
27.02.2019

[73] Proprietor 專利所有人
OBSEVA S.A.
12, CHEMIN DES AULX
1228 PLAN-LES-OUATES, GENEVE
SWITZERLAND
[72] Inventor 發明人
CHOLLET, André
POHL, Olivier
[74] Agent and / or address for service 代理人及/或送達地址
CHINA PATENT AGENT (HONG KONG) LIMITED
22/F, Great Eagle Centre, 23 Harbour Road
Wanchai
HONG KONG



(11) **EP 3 266 449 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
27.02.2019 Bulletin 2019/09

(21) Application number: **17188131.1**

(22) Date of filing: **15.12.2014**

(51) Int Cl.:
A61K 9/20 (2006.01) **A61K 9/28** (2006.01)
A61K 9/48 (2006.01) **A61K 45/06** (2006.01)
A61K 9/00 (2006.01) **A61K 47/40** (2006.01)
A61K 31/40 (2006.01) **A61K 31/41** (2006.01)
A61K 31/401 (2006.01) **A61P 15/08** (2006.01)

(54) **ORAL FORMULATIONS OF PYRROLIDINE DERIVATIVES**

ORALE FORMULIERUNGEN VON PYRROLYDINDERIVATEN

FORMULES ORALES DE DÉRIVÉS DE PYRROLYDINE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(30) Priority: **17.12.2013 EP 13197606**

(43) Date of publication of application:
10.01.2018 Bulletin 2018/02

(60) Divisional application:
19150037.0

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
14820814.3 / 3 082 773

(73) Proprietor: **ObsEva S.A.**
1228 Plan-les-Ouates/Geneve (CH)

(72) Inventors:
• **CHOLLET, André**
1228 Plan-Les-Ouates (CH)
• **POHL, Olivier**
1228 Plan-Les-Ouates (CH)

(74) Representative: **KATZAROV S.A.**
European Patent Attorneys
12, Avenue des Morgines
1213 Petit-Lancy (CH)

(56) References cited:
WO-A1-02/074741 WO-A1-2004/005249
WO-A2-2004/076407

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

TECHNICAL FIELD

[0001] The present invention relates to solid oral formulations comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and the use of said formulations in the treatment and/or prevention of preterm labor, premature birth, dysmenorrhea and embryo implantation failure due to uterine contractions. The present invention is furthermore related to processes for their preparation.

BACKGROUND OF THE INVENTION

[0002] Oxytocin (OT) is a cyclic nona-peptide that mediates its physiological actions through activation of the oxytocin receptor (OT-R), a cell membrane receptor belonging to the class of G protein-coupled receptors that is similar to arginine vasopressin receptors. One important action of OT is to cause the contraction of the uterus of mammals during labor. Repeated, concerted and regular contraction of the uterus will cause the dilatation of the cervix, the rupture of foetal membranes and lead to expulsion of the foetus. Premature labor is when these contractions occur before the normal term of pregnancy. Preterm increase of uterine activity is the most common expression of preterm labor.

[0003] Premature labor leads to undesired premature birth, a serious health problem that remains the major cause of perinatal mortality and severe morbidity, especially respiratory distress syndrome, intraventricular haemorrhage, bronchopulmonary dysplasia and necrotising enterocolitis that are far more common in preterm than in term infants. Long-term impairments such as cerebral palsy, visual impairment and hearing loss are also more common in preterm infants. Nowadays, preterm birth remains the leading cause of infant mortality and morbidity in industrialized nations, where, despite the significant improvements in obstetrical medicine, it is causing high costs for neonatal intensive care of premature babies. The actual costs are even higher to society when taking into consideration the healthcare provision of preterm childbirth-related ailments, such as respiratory distress syndrome, heart conditions, cerebral palsy, epilepsy, and severe learning disabilities. The management of preterm labor represents a significant problem in the field of obstetrics.

[0004] The OT/OT-R system plays a vital role in initiating labor in mammals, in particular in humans. The density of OT-R increases markedly in the myometrium before the onset and during labor. Also it is thought that the local OT peptide hormone concentration increases markedly before parturition in human. The high circulating concentrations of progesterone induce uterine quiescence while the uterus acquires contractile ability. Shortly before term, plasma progesterone concentrations fall, OT-R expression in the uterus increases markedly, OT is released and uterine contractile activity increases. At term, the contractions rise to a crescendo, resulting in delivery as a result of two interacting positive feedback loops. The first is a local uterine loop: within the uterus itself, contractile prostaglandins are produced and released in response to OT and uterine contractions. These prostaglandins may play a further role in cervical ripening and weakening of fetal membranes. The second loop involves the hypothalamus: in response to uterine contractions and vaginal and cervical distension, magnocellular oxytocin neurons in the hypothalamus increase their activity resulting in the release of OT from their axon terminals in the posterior pituitary. The released OT acts upon the uterus both to stimulate the further production of prostaglandins and to contribute further to the contractions of the uterus.

[0005] Therefore, blocking the effect of OT by antagonizing OT-R might represent an attractive modality for the treatment of diseases related to the OT-R activity, in particular preterm labor, premature birth and dysmenorrhea.

[0006] Tocolytic, i.e. uterus relaxing agents, have been used in clinical studies for the pharmaceutical treatment of preterm labor. Most of these agents are used off-label. They have shown very limited efficacy, if any, in prolonging gestation and without clear demonstration of improvement of neonate outcome. Current tocolytics are very often associated with unwanted adverse effects on women, foetus or neonate. Such tocolytics include beta-2-adrenergic agonists, prostaglandin synthesis inhibitors, magnesium sulfate, nitric acid donors and calcium channel blockers. Beta-2-adrenergic agonists such as ritodrine or terbutaline cause a number of cardiovascular and metabolic side effects including maternal tachycardia, palpitations, hypotension, altered thyroid function and fetal and neonatal hypoglycaemia, tachycardia. Ritodrine is no longer FDA approved. The calcium channel blocker nifedipine is also a medicine that is used to try to stop contractions. Some of the side effects that may occur include facial flushing, headache, nausea, palpitations, and lightheadedness. The total prostaglandin synthesis inhibitor (NSAID) indomethacin has been used. It can also have serious effects on the fetus: constriction of ductus arteriosus, pulmonary hypertension, decrease in renal function with oligohydramnios, intraventricular hemorrhage, hyperbilirubinemia, necrotizing enterocolitis. Maternal side effects include abdominal discomfort, nausea, vomiting, depression and dizzy spells for the mother. Another NSAID is sulindac that has a side effect profile similar to indomethacin. For magnesium sulfate, meta-analyses have failed to support it as a tocolytic agent. Women reported important side effects such as flushing, lethargy, headache, muscle weakness, pulmonary edema and cardiac arrest. A newborn that has been exposed to magnesium sulfate may show lethargy, hypotonia,

respiratory depression, bone problems, osteopenia and fractures. Recently, the FDA is advising healthcare professionals against using magnesium sulfate injection for longer than 5-7 days to stop preterm labor in women.

[0007] Atosiban, a dual vasopressin V1a receptor and OT-R antagonist is marketed in EU and used to stop contractions and delay preterm delivery by a few days. The principal drawback to the use of peptide antagonists like Atosiban is the

problem of low oral bioavailability resulting from intestinal degradation. Hence, they must be administered parenterally.

[0008] The development of orally active small molecule antagonists that are selective for the OT-R is expected to overcome these problems. Pyrrolidine derivatives being OT-R antagonists are disclosed in WO 01/72705, WO 02/102799, WO 2002/074741, and WO 2004/005249.

[0009] Thus, in the management of preterm labor and premature birth, there is a need for an oral formulation which is convenient to administer, which is suitable for providing a fast onset of action and which provides a good bioavailability of a compound being an OT-R antagonist.

SUMMARY OF THE INVENTION

[0010] The present invention provides a dispersible tablet comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and at least one or more pharmaceutically acceptable excipients.

[0011] The invention also provides said dispersible tablet, for use in the treatment and/or prevention of disorders selected from the group comprising preterm labor, premature birth, embryo implantation failure due to uterine contractions, dysmenorrhea, premature ejaculation, sexual dysfunction, endometriosis, infertility, benign prostatic hyperplasia, neuropsychiatric disorders, autism, social behavior disorders, psycho-social stress, and/or cardiovascular disorders.

[0012] Also provided is a process for the preparation of said dispersible tablet characterized by a step of wet granulation.

[0013] The invention further provides a kit comprising said dispersible tablet, and information for use thereof.

BRIEF DESCRIPTION OF THE FIGURES

[0014]

Figure 1: Plasma concentration profiles of solid oral formulations in the dog. Figure 1A shows the plasma concentration (ng/ml) vs. time profile of formulations 1 (10% granules), 2 (5.8% granules), 3 (dispersible tablets) and 4 (conventional tablets) over the time period from 0 to 72h. Figure 1B shows an enlargement of Figure 1A for the period from 0 to 12h.

Figure 2: Individual plasma concentration profiles of formulation 3 (dispersible tablets) in the dog. Plasma concentration of the compound of formula (3Z,5S) is measured (ng/ml) for each dog (n=5) for the time period from 0 to 12h.

Figure 3: Individual plasma concentration profiles of formulation 4 (conventional tablets) in the dog. Plasma concentration of the compound of formula (3Z,5S) is measured (ng/ml) for each dog (n=5) for the time period from 0 to 12h.

DETAILED DESCRIPTION OF THE INVENTION

[0015] Generally, the present invention relates to a solid oral formulation comprising a compound of formula 5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, its geometrical isomers, its optically active forms as enantiomers, diastereoisomers, mixtures of these, its racemate forms as well as active metabolites thereof, and at least one or more pharmaceutically acceptable excipients.

[0016] Preferably, the present invention relates to a solid oral formulation comprising a compound of formula (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients.

[0017] Even more preferably, the present invention relates to a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients.

[0018] The compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime also named (3Z,5S) herein is produced by methods such as those disclosed for example in WO2004/005249 and WO2005/082848.

[0019] Usually, said compound is synthesized and obtained in isomeric mixtures (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime comprising (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.

[0020] In case the isomer Z is preferred, then the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime synthesized and obtained in isomeric mixtures is purified according to methods disclosed in PCT/EP2014/066075.

[0021] Thus, the purity of the compound (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime in said isomeric mixtures is at least 85% to 100%, preferably 85% to 99.9%, more preferably 90% to 99.9%, and even more preferably 95% to 99.9%.

[0022] Alternatively, the present invention relates to a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, provided in substantially pure form, and at least one or more pharmaceutically acceptable excipients.

[0023] As used herein, the term "substantially pure" refers to a compound provided in a form which is substantially free of other compounds. Examples of said "other compounds" include (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one, (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one oxime, (3R,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-3-methoxyamino-pyrrolidine, (3S,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-3-methoxyamino-pyrrolidine, (3Z,5S)-5-(O-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and (3E,5S)-5-(O-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.


[0024] Most preferably, the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof is substantially free of the compound of formula (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.

[0025] Even more preferably, the purity of a substantially pure form compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, is at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.2%, at least 99.3%, at least 99.4%, at least 99.5%, at least 99.6%, at least 99.7%, at least 99.8%, at least 99.9% or at least 100% and is therefore substantially free of compound of formula (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.


[0026] Even more preferably, the purity of the substantially pure form compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, is at least in the range of 85% to 100%, preferably 85% to 99.9%, more preferably 90% to 99.9%, and even more preferably in the range of 95% to 99.9%.

[0027] As used herein, the term "active metabolite thereof" refers to a product produced through metabolism in the body, or *in vitro*, of a specified compound, i.e. in the present case (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and which exhibits the same biological activity as (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.

[0028] Active metabolites of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime may be identified using routine techniques known in the art and their activities determined using tests such as those described herein. Such metabolites may result for example from the oxidation, glucuronidation or other conjugation, hydrolysis, reduction and the like, of the administered Z form. Accordingly, the invention includes active metabolites of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, including compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such metabolite may also be produced *in vitro* by oxidation, reduction, hydrolysis, glucuronidation or other conjugation transformation of the corresponding (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime. Examples of active metabolites of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, include compounds whose structures are shown below:




5




10

15




20

25




30

35




40

45



50

55



[0029] A compound which, upon administration to the recipient, is capable of being converted into a compound of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof as described above, is known as a "prodrug". A prodrug may, for example, be converted within the body, e. g. by hydrolysis in the blood, into its active form that has medical effects. Pharmaceutical acceptable prodrugs are described in T. Higuchi and V. Stella, Prodrugs as Novel Delivery Systems, Vol. 14 of the A. C. S. Symposium Series (1976); "Design of Prodrugs" ed. H. Bundgaard, Elsevier, 1985; and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, which disclosures are incorporated herein by reference.

[0030] As used herein, the term "solid oral formulation" refers to a tablet, a dispersible tablet, a fast dissolving tablet, a quick dissolving tablet, a fast melt tablet, a mouth-dissolving tablet, a melt-in mouth tablet, an orodispersible tablet, a lyophilised unit, a porous tablet, a conventional tablet, a coated tablet, an uncoated tablet, a gastro-resistant tablet, an effervescent tablet, a soluble tablet, a chewable tablet, an oral lyophilisate, a powder, an oral powder, a pellet, a capsule and/or a granule. Preferably, the solid oral formulation is a tablet, more preferably, a dispersible tablet.

[0031] As used herein, the term "dispersible tablet" includes a disintegrating tablet that is swallowed, or intended to be disintegrated rapidly in water and to be swallowed.

[0032] As used herein, "pharmaceutically acceptable excipients" includes any carriers, diluents, adjuvants, vehicles, preserving agents, antioxidant agents, fillers, bulking agent, glidant, buffering agents, thickening agents, disintegrating agents, lubricants, binders, wetting agents, sweeteners, flavouring agent, taste-masking agents, emulsifying agents, suspending agents, solvents, dispersion media, coatings, antibacterial agents, anti-oxidants, antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well-known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the solid oral formulation is contemplated. Supplementary active ingredients can also be incorporated into the solid oral formulation as suitable therapeutic combinations.

[0033] Preferably, the dispersible tablet of the invention comprises a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and at least one or more pharmaceutically acceptable excipients selected from the group comprising a disintegrant, a wetting agent, a carrier, a lubricant, a binder, a diluent, a sweetener, and/or a taste-masking agent.

[0034] Thus, the present invention relates to a dispersible tablet comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and a disintegrant. For example, the "disintegrant" is selected from one of the group comprising sodium croscarmellose, crospovidone, sodium alginate, colloidal magnesium-aluminum silicate, calcium silicate, sodium starch glycolate, acrylic acid derivatives, microcrystalline cellulose, sodium carboxymethyl cellulose, calcium carboxymethyl cellulose, modified cellulose gum, cross-linked povidone, alginic acid and alginates, pregelatinised starch, modified corn starch and combination thereof. Preferably, the "disintegrant" is selected from the group comprising sodium croscarmellose, crospovidone and combination thereof. More preferably, the "disintegrant" is sodium croscarmellose.

[0035] The present invention alternatively relates to a dispersible tablet comprising a compound of formula (3Z,SS)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and a wetting agent. For example, the "wetting agent" is selected from the group comprising poloxamer, sodium lauryl sulphate, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene stearate, sorbitan fatty acid esters and combination thereof. Preferably, the "wetting agent" is selected from the group comprising poloxamer, sodium lauryl sulfate and combination thereof. More preferably, the "wetting agent" is poloxamer 188.

[0036] Furthermore, the present invention relates to a dispersible tablet comprising a compound of formula (3Z,SS)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and a carrier. For example, the "carrier" is selected from the group comprising calcium silicate, calcium carbonate, calcium phosphate, tribasic calcium phosphate, lactose, starch, modified starch, sugars, celluloses, cellulose derivatives, polymethacrylates, chitin, chitosan and combination thereof. Preferably, the "carrier" is selected from the group comprising calcium silicate, calcium carbonate, calcium phosphate and combination thereof. More preferably, the carrier is calcium silicate.

[0037] The present invention also relates to a dispersible tablet comprising a compound of formula (3Z,SS)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or a disintegrant, and/or a wetting agent, and/or a carrier. Preferably, said dispersible tablet comprises a compound of formula (3Z,SS)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or sodium croscarmellose, and/or poloxamer 188, and/or calcium silicate.

[0038] The present invention also relates to a dispersible tablet comprising a compound of formula (3Z,SS)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and at least one or more pharmaceutically acceptable excipients in an amount effective to provide a tablet that releases between about 90 to 100% of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime. A classical Pharmacopeia compliant dissolution test was performed. As shown in the example, a rapid dissolution profile of the dispersible tablet of 200mg is observed at 15 min wherein the concentration of (3Z,5S) in water is between 90% to 100%

of the initial concentration value (Table 30).

[0039] Preferably the one or more pharmaceutical acceptable excipients includes at least one disintegrant. More preferably, the disintegrant is selected from the group consisting of sodium croscarmellose, crospovidone and a combination thereof. More preferably, the "disintegrant" is sodium croscarmellose.

[0040] For example, the "binder" is selected from the group comprising polyvinylpyrrolidone, cross-linked PVP, cellulose or cellulose derivatives such as hydroxypropylmethyl cellulose (HPMC), carboxymethylcellulose sodium, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, methyl cellulose, carboxyethylcellulose, calcium, guar gum, tragacanth, polyvinylacetates, gelatin, pregelatinised starch, starch, polyvinylalcohols, alginic acid, sodium alginate, sorbitol, glucose, magnesium aluminium silicate, dextrin, polyethylene glycol, polymethacrylates and combination thereof.

[0041] For example, the "diluent" is selected from the group comprising microcrystalline cellulose, lactose monohydrate, lactose, compressible sugar, sugar, dextrose, mannitol, dextrin, maltodextrin, sorbitol, xylitol, sodium chloride, calcium carbonate, magnesium carbonate, calcium phosphate, calcium sulphate, magnesium oxide, kaolin, powdered cellulose, pregelatinized starch, starch, barium sulphate, magnesium trisilicate, aluminium hydroxide and combination thereof.

[0042] For example, the "sweetener" is sodium saccharine, sucrose, sucralose, aspartame, sorbitol or combination thereof.

[0043] For example, the "lubricant" is selected from the group comprising glycerol dibehenate, glycerol tribehenate, magnesium stearate, calcium stearate, talc, sodium stearyl fumarate, sodium behenate, stearic acid, cethyl alcohol, polyoxyethylene glycol, leucine, sodium benzoate, stearates, talc, polyethylene glycol, glyceryl monostearate, glyceryl palmitostearate, liquid paraffin, poloxamer, sodium lauryl sulphate, magnesium lauryl sulphate, hydrogenated castor oil, colloidal silicon dioxide, palmitostearate, stearic acid, zinc stearate, stearyl alcohol, hydrogenated vegetable oil and combination thereof.

[0044] The present invention also relates to a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients, wherein the concentration of said compound and/or active metabolite thereof, is comprised between about 1% and 50% w/w.

Preferably, the concentration of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof is 10-40% w/w, 20-30% w/w, about 20% w/w.

[0045] As used herein, the term "about" applies to numeric values and refers to a range of numbers that one of skill in the art would consider equivalent to the recited values. For example, "about 20% w/w" refers to the range 15%-25%w/w.

[0046] The present invention also relates to a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients, wherein said formulation comprises about 10mg to about 500mg of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof.

[0047] Preferably, said formulation comprises about 20-400mg or 40-200mg of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof. Preferably, said formulation in the form of a dispersible tablet comprises about 50mg or 200mg of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.

[0048] Advantageously, the present invention provides a solid oral formulation, which is i) convenient to administer, ii) suitable for providing a fast onset of action and which provides a good bioavailability of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof. As used herein, the term "Tmax" refers to the time to reach the peak plasma concentration (Cmax) of a drug after administration wherein the concentration is the amount of the drug in a given volume of plasma, expressed in ng/ml in the examples.

[0049] As used herein, the term "onset of action" refers to the time required after administration of a drug to become effective.

[0050] As shown in the examples, the solid oral formulations of the present invention, in particular in the form of a dispersible tablet, have the advantage of being suitable for providing a rapid onset of action. The maximum concentration in blood of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof, is reached at a time Tmax less than 5h, preferably less than 4h, more preferably less than 3h, less than 2h, less than 1.5h, even more preferably less than 1h following administration of said solid oral formulation.

[0051] Also, the maximum concentration in blood of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof, is reached at a time Tmax between 0.5 to 4 hours, 0.5 to 2 hours, preferably 0.5 to 1.5 hours, more preferably at a time between 0.5 to 1 hour following administration of said solid oral formulation. Preferably, said solid oral formulation is a dispersible tablet.

[0052] Noteworthy, at a time of 0.5 hour following administration of the solid oral formulation, the concentration in

blood of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof, is at least 25%, at least 35%, at least 40%, at least 45%, at least 55%, at least 65%, at least 75%, or at least 85% of C_{max}. Preferably, said solid oral formulation is a dispersible tablet.

[0053] Also, at a time of 0.5 hour following administration of the solid oral formulation, the concentration in blood of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof, is comprised between 35% to 100%, 45% to 100%, 55% to 100% of C_{max}, preferentially 57% to 92% of C_{max} indicating that the solid oral formulation is suitable for providing a rapid onset of action. Preferably, said solid oral formulation is a dispersible tablet.

As shown in the examples, at a time of 0.5 hour following administration of the dispersible tablet, the concentration in blood of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, is comprised between 59% to 100% of C_{max} in animal (Table 17), preferentially 57% to 92% of C_{max} in human subjects (Table 22).

[0054] Thus, the present invention provides a solid oral formulation, preferably a dispersible tablet that is suitable for providing a rapid onset of action, which is crucial for the management of preterm labor and premature birth.

[0055] In particular, it has been shown that the maximum concentration of the active substance (3Z,5S) is detected rapidly at about 4 hours, 2 hours, 1.5 hours, or 1 hour following administration of the dispersible tablet of the present invention.

[0056] Furthermore, the solid oral formulation of the present invention is characterized by a bioavailability of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or active metabolite thereof, comprised between 50-100%, and/or 50-99%. Preferably, said bioavailability is comprised between 75-100%, or 75-99%, more preferably, between 80-100%, or 80-99%.

As used herein, the term "bioavailability" (F%) refers to the fraction of an administered dose of a product that reaches the systemic circulation. By definition, when the product is administered intravenously, its bioavailability is 100%. When the product is administered via other routes, its bioavailability generally decreases.

[0057] As shown in the examples, the bioavailability (F%) of the solid oral formulations of the present invention, is comprised between 58% and 90% (Table 16). In particular, the bioavailability of the solid oral formulation in the form of a dispersible tablet is comprised between 80% and 100% (Table 18, 102% in table 18 is indicated as 100% based on the standard deviation).

[0058] In particular, it has been shown that the bioavailability of the active substance (3Z,5S) is about 89% following administration of the dispersible tablet of the present invention (Table 16). Advantageously, patients administered with the solid oral formulation of the present invention will benefit from a fast onset of action and/or a good bioavailability.

[0059] Also, the present invention provides a solid oral formulation comprising:

20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof;

1-20% by weight of calcium silicate;

0.1-20% by weight of PVP30K;

0.01-5% by weight of poloxamer 188;

0.5-20% by weight of sodium croscarmellose;

1-90% by weight of microcrystalline cellulose 112;

1-90% by weight of lactose monohydrate;

0.01-0.5% by weight of sodium saccharine; and

0.1-10% by weight of glycerol dibehenate.

[0060] Alternatively, the present invention provides a dispersible tablet comprising 20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and 0.5-20% by weight of a disintegrant. Preferably said disintegrant is sodium croscarmellose.

[0061] Also alternatively provided is a dispersible tablet comprising 20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and 1-20% by weight of a carrier. Preferably, said carrier is calcium silicate.

[0062] Alternatively, it further provides a dispersible tablet comprising 20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and 0.01-5% by weight of a wetting agent. Preferably, said wetting agent is poloxamer 188.

[0063] Preferably, said solid oral formulation consists of:

20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof,

5% by weight of calcium silicate,

1% by weight of PVP30K,
 2% by weight of Poloxamer 188,
 5% by weight of Sodium croscarmellose,
 15% by weight of Microcrystalline cellulose 112,
 47.8% by weight of Lactose monohydrate,
 0.2% by weight of Sodium saccharine and
 4% by weight of Glycerol dibehenate.

[0064] Whilst the compound (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and/or the active metabolite thereof may be used as the sole active ingredient of the solid oral formulation, it is also possible for the compound to be used in combination with at least one or more further active compounds. Such further active compounds may be further compounds according to the invention, or other active compounds selected from the group comprising calcium channel blockers, magnesium sulfate, selective prostaglandin modulators, beta-2-adrenergic agonists, beta-3-adrenergic receptor agonists, and/or corticosteroids.

[0065] Preferably, corticosteroids are selected from the group comprising Betamethasone and Dexamethasone, and/or salts thereof. These corticosteroids are given before birth to accelerate a preterm fetus' lung development and maturation to prevent respiratory distress syndrome (RDS) and other related complications following premature birth.

[0066] Alternatively, the solid oral formulation of the invention can be administered concomitantly or separately with at least one compound selected from the group comprising calcium channel blockers (such as nifedipine), magnesium sulfate, prostaglandin receptors modulators (such as agonists or antagonists of either EP1 or EP2 or EP3 or EP4 or FP receptors), prostaglandin synthesis inhibitors (such as indomethacin, nimesulide, sulindac, rofecoxib, celecoxib), beta-2-adrenergic agonists (such as ritodrine, terbutaline, salbutamol), beta-3-adrenergic receptor agonists, nitric acid donors (such as glyceryl trinitrate) and/or corticosteroids (such as dexamethasone, betamethasone).

[0067] As used herein, the term "concomitantly" refers to the administration of a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, which is then immediately followed by the administration of at least one compound selected from the group disclosed *supra*.

[0068] As used herein, the term "separately" encompasses sequential or subsequent administration and refers to the administration of a solid oral formulation of the invention comprising the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, followed by a time period of discontinuance, which is then followed by the administration of at least one compound disclosed *supra*.

[0069] The compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, is an oxytocin receptor antagonist.

[0070] As used herein, the term "oxytocin receptor antagonist" refers to a compound that functions by inhibiting (partially or completely) or blocking the oxytocin receptor (OT-R), thereby preventing activation of the receptor by oxytocin.

[0071] Generally, the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, is a vasopressin V1a receptor antagonist.

[0072] As used herein, the term "vasopressin V1a receptor antagonist" refers to a compound that functions by inhibiting (partially or completely) or blocking the vasopressin V1a receptor (also known as Arginine vasopressin receptor 1A), thereby preventing activation of the receptor by vasopressin. Vasopressin V1a receptor is one of the three major receptor types for the peptide hormone arginine vasopressin, the others being V1b and V2 receptors.

[0073] Hence, the present invention relates to a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients, wherein said compound is an oxytocin receptor antagonist and/or a vasopressin V1a receptor antagonist.

[0074] Disorders associated with the oxytocin receptor activity and/or vasopressin V1a receptor activity are selected from the non-limiting group comprising preterm labor, premature birth, embryo implantation failure due to uterine contractions, dysmenorrhea, premature ejaculation, sexual dysfunction, endometriosis, infertility, benign prostatic hyperplasia, neuro-psychiatric disorders, autism, social behavior disorders, psycho-social stress, and/or cardiovascular disorders.

[0075] The term "preterm labor" referring also to "premature labor", shall mean expulsion from the uterus of a viable infant before the normal end of gestation, or more particularly, onset of labor with effacement and dilation of the cervix before the 37th week of gestation. It may or may not be associated with vaginal bleeding or rupture of the membranes.

[0076] The term "dysmenorrhea" refers to a condition characterized by cyclic pain associated with menses during ovulatory cycles. The pain is thought to result from uterine contractions and ischemia.

[0077] The term "sexual dysfunction" refers to any disturbance or variation in the four phases of sexual response: excitement phase, plateau phase, orgasmic phase and resolution phase characterizing the human sexual response.

[0078] The term "neuro-psychiatric disorders" as used herein refers to mental disorders attributable to diseases of the

nervous system, e.g. depression, obsessive-compulsive disorder and others.

[0079] The term "social behavior disorders" as used herein refers to emotional disturbance, inappropriate types of behavior or feelings, pervasive mood of unhappiness or depression and a range of perceived difficulties to build or maintain satisfactory interpersonal relationships. The term "psycho-social stress" as used herein refers to a condition resulting from a perceived threat to the social status, social esteem, self-worth, respect or acceptance within a group, and that lead to development of a stress response in the body and physical symptoms.

[0080] Assisted reproduction technologies are methods applied in humans for the treatment of infertility and in animals for producing pregnancies. Infertility, which affects about 10% of human pairs worldwide, may be treated by in vitro fertilization and embryo transfer (IVF-ET) or in less complicated cases, by artificial insemination. Generally, a success of an embryo transfer is dependent on uterine receptivity, an entity that is defined as an ability of uterus to provide optimal conditions mandating proper implantation and embryo development. Basic components of uterine receptivity are uterine contractile activity and the condition of endometrium.

[0081] Uterine contractions occurring during the embryo transfer may expel embryos from the uterus towards vagina or oviducts, which may be a cause of unsuccessful treatment, or in latter case a cause of extra uterine pregnancy, a serious, potentially life-threatening complication.

[0082] Hence, the present invention provides a solid oral formulation, for use in the treatment and/or prevention of disorders selected from the group comprising preterm labor, premature birth, dysmenorrhea, premature ejaculation, sexual dysfunction, endometriosis, embryo implantation failure due to uterine contractions, infertility, benign prostatic hyperplasia, neuro-psychiatric disorders, autism, social behaviour disorders, psycho-social stress, and/or cardiovascular disorders. Preferably, said solid oral formulation is a dispersible tablet.

[0083] Preferably, the present invention provides a solid oral formulation for use in the treatment and/or prevention of preterm labor, premature birth, dysmenorrhea and embryo implantation failure due to uterine contractions.

[0084] The present invention also provides a process for the preparation of a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyl-oxime, and/or an active metabolite thereof, characterized by a step of wet granulation.

[0085] In wet granulation, granules are formed by the addition on powder particles of a liquid such as water, ethanol and isopropanol, either alone or in combination.

[0086] Preferably, the present invention provides a process for the preparation of a tablet comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyl-oxime, and/or an active metabolite thereof, characterized by a step of wet granulation. More preferably, said tablet is a dispersible tablet.

[0087] Alternatively, the present invention provides a process for the preparation of a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyl-oxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients, characterized by the steps of:

- (i) mixing the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyl-oxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients;
- (ii) wet-granulating;
- (iii) sieving the granules
- (iv) blending with a lubricant such as glycerol dibehenate; and
- (v) compressing the mixture obtained in step (iv) to form a tablet.

Preferably, said tablet is a dispersible tablet.

[0088] The tablet cores may vary in shape and be, for example, round, oval, oblong, cylindrical or any other suitable shape.

[0089] The present invention also provides a kit comprising a solid oral formulation comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyl-oxime, and/or an active metabolite thereof, and at least one or more pharmaceutically acceptable excipients, and information for use thereof. The information contains instructions to administer the oral formulation to a subject in need thereof.

[0090] Generally, in the present invention the subject in need thereof is preferably a mammal, most preferably a human, more preferably a woman.

[0091] For practical reason, the solid oral formulation of the present invention can be packaged in a unit dose. As used herein, the term "unit dose" refers to a solid oral formulation that is dispensed in a package ready to administer to the patient. Each unit dose contains a predetermined quantity of active product calculated to produce the desired therapeutic effect, in association with at least one or more suitable pharmaceutical excipients.

EXAMPLES

Example 1: Purification of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime

1.1 Synthesis of (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime

[0092] In the present invention, the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime was obtained as a crude isomeric mixture comprising (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime.


[0093] Synthetic pathways of compounds used in the invention are for example those described in WO2004005249 and WO2005082848.

[0094] The compound "(3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime" used herein is also defined as "(4Z,2S)-2-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidine-4-one-O-methyloxime" depending on the nomenclature used.

[0095] The compound (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime can also be prepared following stages 1 to 6 as described below:

Stage 1: Preparation of 4-(2-methylphenyl)benzoic acid


[0096]



[0097] A solution of potassium carbonate (0.908Kg, 6.57 mol, 2.06 wt) in water (2.20L, 5.0vol) was charged to a slurry of 4-bromobenzoic acid (0.441 Kg, 2.19 mol, 1.0 wt) in water (4.41L, 15.0vol) at 15 to 25°C. The resulting slurry was stirred at 15 to 25°C and degassed three times using a vacuum-nitrogen purge cycle. Tetrakis(triphenylphosphine)palladium(0) (0.022Kg, 0.019 mol, 0.05 wt) was charged and the vacuum-nitrogen purge cycle repeated. A solution of o-tolylboronic acid (0.313Kg, 2.30 mol, 0.707 wt) in methanol (3.53L, 8.0 vol) was degassed three times, using a vacuum-nitrogen purge cycle, and then charged to the 4-bromobenzoic acid slurry at 15 to 25°C. The reaction mixture was heated to and maintained at reflux (71 to 78°C) until reaction completion (The reaction is considered complete at 95% conversion), as determined by ¹H NMR analysis (d6-DMSO), typically 1.5 to 2.5 hours. The reaction mixture was concentrated to 15vol under vacuum at 40 to 45°C. Toluene (4.41L, 10.0 vol) and tetrahydrofuran (4.41L, 10.0vol) were added to the residue, the resulting mixture stirred vigorously and acidified to pH 1 with hydrochloric acid (6M, 2.00L, 4.5vol). The contents were stirred vigorously for 30 to 60 minutes and the layers separated. Toluene (2.20L, 5.0vol) and tetrahydrofuran (2.20L, 5.0 vol) were added to the aqueous phase and the mixture stirred for 5 to 10 minutes. The layers were separated, the combined organic phases filtered and concentrated to 10.0vol under vacuum at 35 to 40°C. Toluene (4.41L, 10.0 vol) was added to the residue and the resultant concentrated under vacuum at 35 to 40°C. The tetrahydrofuran content of the resulting slurry was determined by ¹H NMR analysis (d6-DMSO) (Pass level: ≤1.0%w/w tetrahydrofuran with respect to toluene). The slurry was cooled to and aged at 0 to 5°C for 30 to 60 minutes, the solid collected by filtration and the filter-cake washed with toluene (2.20L, 5.0 val). The solid was dried in a vacuum oven at 35 to 40°C to give 4-(2-methylphenyl)benzoic acid [0.438Kg, 94.1%th, 99.3%w/w, ¹H NMR (d6-DMSO) concordant with structure] as a pale yellow solid.

Stage 2: Preparation of 4-(2-methylphenyl)benzoic acid chloride


[0098]



[0099] Thionyl chloride (0.300L, 4.11 mol, 0.685 vol) was added to a slurry of 4-(2-methylphenyl)benzoic acid (0.435Kg, 2.05 mol, 1.0 wt) in toluene (4.35L, 10.0 vol) at 10 to 25°C and the mixture heated to and maintained at 75 to 80°C until complete by ¹H NMR analysis (d6-benzene), typically 4 to 5 hours. Reaction completion was accompanied by the formation of a hazy solution. The resultant was concentrated to 5.0 vol by removal of toluene under reduced pressure at 35 to 45°C. Toluene (2.18L, 5.0 vol) was added to the concentrate and the mixture concentrated to 4.0 vol by removal of toluene under reduced pressure at 35 to 45°C. The resultant was filtered through glass microfibre paper and the filter-cake washed with toluene (0.44L, 1.0 vol). The toluene solution of 4-(2-methylphenyl)benzoic acid chloride [0.439Kg, 92.8%th, 100.9%w/w, ¹H NMR (d6-benzene) concordant with structure] was used directly in Stage 3.

Stage 3: Preparation of (4R)-4-hydroxy-1-[(2'-methyl-1,1'-biphenyl-4-yl)-carbonyl]-L-proline


[0100]



[0101] A solution of potassium carbonate (0.526Kg, 3.81 mol, 1.2 wt) in water (0.57L, 1.3 vol) was charged to a solution of 4-hydroxy-L-proline (0.274Kg, 2.09 mol, 0.625 wt) in tetrahydrofuran (2.20L, 5.0 vol) and water (0.44L, 1.0 vol) at 15 to 25°C followed by a line rinse of water (0.44L, 1.0 vol). The mixture was cooled to 0 to 5°C with rapid stirring and a solution of 4-(2-methylphenyl)benzoic acid chloride (0.438Kg, 1.90 mol, 1.0 wt) in toluene (2.19L, 5.0 vol) charged at that temperature followed by a line rinse of toluene (0.44L, 1.0 vol). The reaction mixture was warmed to 15 to 25°C over 1 to 2 hours and stirred at this temperature until judged complete by TLC analysis. Water (2.20L, 5.0 vol) was charged to the reaction mixture at 15 to 25°C and the layers separated. The aqueous phase was acidified to pH 5 to 6 with aq. hydrochloric acid (6M, 0.66L, 1.5 vol) and then to pH1 with aq. hydrochloric acid (2M, 0.88L, 2.0vol) at 15 to 25°C. The mixture was cooled to and aged at 0 to 5°C for 30 to 60 minutes, the precipitated solid collected by filtration, the filter-cake washed with water (2x 1.75L, 2x 4.0 vol) and toluene (0.88L, 2.0 vol) and pulled dry on the filter for 12 to 24 hours. The collected solid was dried under vacuum at 40 to 45°C until the water content by KF was ≤0.2%w/w to afford (4R)-4-hydroxy-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-L-proline [0.599Kg, 97.0%th, 136.8%w/w, ¹H NMR (d₆-DMSO) concordant with structure] as an off-white solid.

Stage 4: Preparation of 1-(2'-methyl-1,1'-biphenyl-4-yl)carbonyl-4-oxo-L-proline


[0102]



[0103] Triethylamine (1.80L, 13.56 mol, 3.0 vol) was charged to a solution of (4R)-4-hydroxy-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-L-proline (0.598Kg, 1.84 mol, 1.0 wt) in dimethyl sulfoxide (4.42L, 7.4 vol) at 15 to 20°C. Pyridine-sulphur trioxide complex (0.879Kg, 5.52 mol, 1.47 wt) was charged portion-wise at 15 and 25°C and the reaction mixture stirred at that temperature until reaction completion, as determined by TLC analysis (typically 1 to 3 hours). The reaction was quenched with aq. hydrochloric acid (3M, 4.80L, 8.0 vol) at 0 to 30°C, tetrahydrofuran (3.00L, 5.0 vol) and heptanes (0.60L, 1.0 vol) charged, the layers separated and the aqueous phase extracted with tetrahydrofuran (2x 3.00L, 2x 5.0 vol). The combined organic phases were washed with aq. hydrochloric acid (1 M, 2x 1.20L, 2x 2.0 vol) and saturated sodium chloride solution (2x 1.20L, 2x 2.0 vol), the aqueous washes combined and back-extracted with tetrahydrofuran (2x 0.60L, 2x 1.0 vol). The combined organics were dried over magnesium sulphate (1.794Kg, 3.0 wt), filtered, the filtercake washed with tetrahydrofuran (0.60L, 1.0 vol) and the filtrates concentrated under vacuum at 40 to 45°C to give a pale brown foam. Ethyl acetate (6.00L, 10.0 vol) was charged to the foam, the contents stirred for 5 to 10 minutes to reach dissolution and the solvent removed under vacuum at 40 to 45°C. This was repeated using ethyl acetate (6.00L, 5.0 vol) until tetrahydrofuran was not detected by ^1H NMR analysis (d_6 -DMSO). The residue was slurried in ethyl acetate (4.80L, 8.0 vol), activated carbon (0.084Kg, 0.14 wt) added followed by a line rinse of ethyl acetate (3.00L, 5.0 vol), the resultant heated to and maintained at 70 to 80°C for 20 to 30 minutes, cooled to 40 to 55°C and filtered through glass microfibre paper. The filter-cake was washed with ethyl acetate (1.50L, 2.5 vol) and the combined filtrates and wash concentrated to 2.5 to 3.5 vol under vacuum at 40 to 45°C. Crystallisation commenced during the concentration. The concentrate was transferred to a suitable vessel with a line rinse of ethyl acetate (0.30L, 0.5 vol) and heated to 70 to 80°C. Additional ethyl acetate (0.30L, 0.5 vol) was added as necessary to achieve dissolution. Heptanes (1.80L, 3.0 vol) was added at 70 to 80°C and the contents allowed to cool to between 15 and 25°C over 1 to 2 hours. The slurry was further cooled to and aged at 0 to 5°C for 2 to 3 hours, filtered and the filtercake washed with ethyl acetate:heptanes (1:1, 0.60L, 1.0 vol) at 0 to 5°C followed by heptanes (3.0L, 2.5 vol). The collected solid was dried under vacuum at 40 to 45°C to give 1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-4-oxo-L-proline [0.444Kg, 74.7%th, 74.2%w/w, ^1H NMR (d_6 -DMSO) concordant with structure] as an off-white solid.

Stage 5: Preparation of (4Z/E)-4-methoxyimino-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-L-proline

[0104]




[0105] Triethylamine (0.40L, 2.85 mol, 0.92 vol) was added to a solution of 1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-4-oxo-L-proline (0.434Kg, 1.34 mol, 1.0 wt) in dichloromethane (4.40L, 10.0 vol) at 10 to 25°C followed by a line rinse of dichloromethane (0.43L, 1.0 vol). Methoxylamine hydrochloride (0.130Kg, 1.56 mol, 0.30 wt) was added portionwise at 10 to 25°C followed by a line rinse of dichloromethane (0.43L, 1.0 vol) and the reaction mixture stirred at 10 to 25°C

until reaction completion, as determined by TLC analysis (typically 3 to 5 hours, TLC eluent: dichloromethane:methanol:acetic acid (90:10:1); uv visualization). The solvent was removed under vacuum at 35 to 40°C, the resultant dissolved in ethyl acetate (4.40L, 10.0 vol) and washed with aq. hydrochloric acid (1 M, 2x 2.20L, 2x 5.0 vol). The acidic washes were back extracted with ethyl acetate (2.20L, 5.0 vol), the combined organic phases washed with sat. aq. sodium chloride solution (3.10L, 7.0 vol), dried over magnesium sulfate (0.300Kg, 0.69 wt), filtered and the filtercake washed with ethyl acetate (2.20L, 5.0vol). The filtrate and washes were combined and concentrated under vacuum at 35 to 40°C to afford 4-methoxyimino-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-L-proline [0.476Kg, 100.6%th, 109.6%w/w, ¹H NMR (CDCl₃) concordant with structure] as an off-white solid.

Stage 6: Preparation of (4Z/E, 2S)-methyl-1-[(2'-methyl-1,1'-biphenyl-4-yl)-carbonyl]-4-methoxyimino pyrrolidine-2-carboxylate


[0106]



Potassium carbonate (0.476Kg, 3.44 mol, 1.0 wt) was added to a solution of 4-methoxyimino-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]-L-proline (0.475Kg, 1.35 mol, 1.0 wt) in acetone (4.75L, 10.0 vol) and the mixture cooled to 0 to 10°C. Dimethyl sulfate (0.128L, 1.35 mol, 0.27 vol) was added at 0 to 15°C and the mixture stirred at 15 to 25°C until reaction completion, as determined by TLC analysis, typically 3 to 16 hours. The solvent was removed under vacuum at 40 to 45°C and the resultant partitioned between ethyl acetate (3.80L, 8.0 vol) and water (3.80L, 8.0 vol). The layers were separated, the organic phase washed with sat. aq. sodium chloride solution (2.85L, 6.0 vol), dried over sodium sulfate (0.953Kg, 2.0 wt) and filtered. The filter-cake was washed with ethyl acetate (0.48L, 1.0 vol) and the combined filtrate and wash concentrated under vacuum at 40 to 45°C. Excess ethyl acetate was removed by azeotropic distillation with tetrahydrofuran (2x 0.95L, 2x 2.0 vol) under vacuum at 40 to 45°C to give (4Z/E, 2S)-methyl-1-[(2'-methyl-1,1'-biphenyl-4-yl)-carbonyl]-4-methoxyimino pyrrolidine-2-carboxylate [0.492Kg, 99.6%th, 103.6%w/w, ¹H NMR (CDCl₃) concordant with structure] as a viscous brown oil.

Stage 7: Preparation of (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime

[0108]



Lithium borohydride (0.049Kg, 2.26 mol, 0.1 wt) was added portionwise under nitrogen to a stirred solution of (4Z/E, 2S)-methyl-1-[(2'-methyl-1,1'-biphenyl-4-yl)-carbonyl]-4-methoxyimino pyrrolidine-2-carboxylate (0.492Kg, 1.34 mol, 1.0 wt) in tetrahydrofuran (2.31L, 4.7 vol) and methanol (2.31L, 4.7 vol) at 0 to 30°C. The mixture was stirred at 15 to 25°C to reaction completion, as determined by TLC analysis (Eluent: ethyl acetate; Visualisation: ninhydrin), typically

2 to 6 hours. The reaction mixture was quenched with water (0.40L, 0.8 vol) at 15 to 25°C and stirred at 15 to 25°C for 16 to 20 hours. The resultant was concentrated under vacuum at 40 to 45°C and the residue partitioned between water (2.46L, 5.0 vol) and ethyl acetate (4.92L, 10.0 vol). The layers were separated, the organic phase washed sequentially with aq. hydrochloric acid (1M, 2.46L, 5.0 vol), sat. aq. sodium hydrogen carbonate solution (2.46L, 5.0 vol) and sat. aq. sodium chloride solution (2.46L, 5.0 vol). The organic phase was dried over magnesium sulfate (0.985Kg, 2.0 wt), filtered and the filter-cake washed with ethyl acetate (0.50L, 1.0 vol). The combined filtrate and wash were concentrated under vacuum to give a crude isomeric mixture comprising (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and (3E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime [0.395Kg, 86.9%th, 80.3%w/w, ¹H NMR (CDCl₃) concordant with structure; 82.0% area by HPLC, 71.4:28.6 Z/E ratio] as a viscous brown oil. The oil was dissolved in toluene (0.40L, 1.0vol, with respect to weight of product) and stored until required.

1.2 Dry flash chromatography of crude (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime


[0110] A dry flash chromatography purification of the crude isomeric mixture obtained following the protocol described above was attempted using different elution conditions. A crude mixture of (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime concentrated to dryness was re-dissolved in 2 volume toluene and loaded onto a pad of SiO₂ (5 wt) prior to elution using 25 volume fractions of eluent.

Fractions 1-5: eluted with pure toluene

Fractions 6-10: eluted with Toluene / MeOH 1% vol/vol

Fractions 10 to 15: eluted with Toluene / MeOH 2% vol/vol

Schematic TLC profile of the collected fractions



[0111] The Z and E forms are shown by shaded spots. Fractions 8 to 13 were combined and concentrated to dryness. The results show a recovery of 75%. There was no improvement in the E/Z ratio. A minor gain of about 4% area in purity of the isomeric mixture (E+Z) was observed before and after dry-flash chromatography (Table 1).

Table 1: Comparative impurity profile before and after dry-flash chromatography

	% area			
	Impurity at RRT 0.7	E+Z-isomers	Impurity at RRT 1.08	RRT 1.12 (Ar-ArCH ₂ OH)
Before dry flash	4.6	91.3	<0.5	4.1
After dry-flash	2.5	95.6	<0.5	0.7
RRT: Relative retention time				

[0112] The dry-flash chromatography of the crude isomeric mixture does not allow the purification of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime. The E/Z ratio pre and post dry-flash remain in the range of 30/70 to 40/60. Furthermore, such an approach should be considered on the basis of the scale at which the operation has to be carried out. On a 20 L scale, this operation would not be a time saving approach.

1.3 Assessment toward crystallization of the pure Z from the crude isomeric mixture

[0113] The first part of the assessment toward crystallisation of the pure (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime from the crude mixture (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime, has been looking at solubility and possible crystallisation conditions of the pure (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime. The results of the solubility/crystallisation tests carried out on 15 mg scale are reported in Table 2 below

Table 2: Qualitative solubility data for (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime

Solvent	Dissolves in:	Comment
heptanes	--	insoluble in 20 vol
toluene	2 vol cold	
DIPE	40 vol hot	
THF	4 vol cold	
tBuOH	6 vol hot	
MIBK	4 vol hot	
IPA	4 vol hot	

[0114] The initial solubility screen showed that pure (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime isomer is soluble in a range of solvents. On the basis of the above results, crystallisation by addition of anti-solvent was examined and the results reported in Table 3. The anti-solvent was added to a warm solution ca 40-50°C and allowed to cool to room temperature.

[0115] In particular, the water (anti-solvent) was added to a warm (40-50°C) solution of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime in IPA until cloudiness was reached and the mixture was allowed to cool to room temperature.

Table 3: Crystallisation via addition of anti-solvent

Solvent	Antisolvent	Comment
toluene 20 vol	heptanes 39 vol	oils out
THF 10 vol	heptanes 40 vol	oils out
tBuOH 10 vol	water 20 vol	oils out
MIBK 10 vol	heptanes 40 vol	oils out
IPA 20 vol	water 160 vol	very fine solid, oils out on standing
IPA 8 vol	water 18 vol	very fine solid, oils out on standing
DMSO 10 vol	water 12 vol	gel
NMP 10 vol	water 28 vol	oils out
MeOH 10 vol	water 10 vol	oils out
DMSO 20 vol	water 16 vol	oils out
acetone 10 vol	water 10 vol	oils out
DCM 10 vol	heptanes 50 vol	oils out

[0116] The IPA/water crystallisation conditions were applied to a crude isomeric mixture. The toluene solution was first concentrated to dryness prior to dissolution in IPA (8 vol) and addition of water (18 vol). Unfortunately, this resulted in material de-mixing as oil.

[0117] In another experiment, the antisolvent was added to a solution of crude (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime (90.4% area purity, contained 0.5% w/w toluene and

3.7%w/w THF) at room temperature until cloudiness was reached and the mixture was left to stand at room temperature (Table 4).

Table 4: Crystallisation by addition of water at 18-22°C

Solvent	Antisolvent	Comment
MeOH 5 vol	water 3 vol	oils out
DMSO 5 vol	water 3 vol	oils out

[0118] At this point of the investigation, no suitable conditions of crystallisation of the pure (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime or allowing isolation of solid containing (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime have been identified.

[0119] Further crystallisation attempts were carried out using crude isomeric mixture of (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime. In all cases, the volume of solvents was smaller than what used previously and based only on a single solvent. The crude material (E/Z ratio 33:67 and purity (E+Z) 79.52 %area) used for this crystallisation was concentrated to a foam (Table 5).

Table 5: crystallisation from single solvent at lower volume

Material	Solvent	Ageing in freezer	Ageing in fridge
'Pure Z'	Ethyl Acetate 1.8 vol	Crystallises re-dissolves as warms	Stays in solution, with and without seeding after 2 days.
Crude		Does not crystallize with or without seeding.	n/a
'Pure Z'	Diethylether 2.3 vol	On addition of ether at 18-22°C starts to dissolve then crashes out again. Recovery 70% Used for seeding	n/a
Crude		Oils Re-dissolves as warms	Crystallises recovery 41 % E/Z ratio 40/60 purity 85.4%area. (mother liquors E/Z ratio 20/80 purity 62.1%area). Seeds not used.
'Pure Z'	TBME 2.3 vol	Oils Re-dissolves as warms	Stays in solution, with and without seeding after 2 days.
Crude		Oils Re-dissolves as warms	Stays in solution, with and without seeding after 2 days.

[0120] Crystallisation using ethyl acetate followed by aging in a freezer overnight gave crystallisation using the pure (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime material, but quickly re-dissolved as the sample warmed. No crystals were observed using crude material in ethyl acetate even when seeds were added.

[0121] Crystallisation using diethylether followed by aging in a fridge gave crystallisation using the crude (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime material. The solid was collected in 41 % recovery. Unfortunately, the collected solid had a slighter poorer E/Z ratio than the input material and a slightly higher chemical purity.

[0122] TBME as solvent for both pure Z and crude gave oiling after aging in freezer, and stayed in solution after aging in the fridge with and without seeds.

[0123] Suitable crystallization conditions of the crude isomeric mixture allowing improvement of the Z/E ratio and of the purity of the isomeric mixture (E+Z) have not been found.

1.4 Substantially pure form of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime

1.4.1 Small scale purification

[0124] The isolation procedure in substantially pure form of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime was performed by chromatography using a Biotage system (Biotage AB, SE-751 03 Uppsala, Sweden) of the crude isomeric mixture isolated after reduction of the oxime ester (Stage 7 of Example 1).

[0125] Five distinct batches (No. 020, 180, 062, 068, 076) of the crude isomeric mixture were purified by Biotage chromatography. Furthermore, different conditions were used regarding batches No. 068 and 076. Purification was performed with a 5%w/w spike of oxime methyl ester added (No. 068), and with an overloaded Biotage column (No. 076).

[0126] Each chromatography was run using Biotage 40M cartridges (40 g silica) which had been pre-flushed with toluene. Toluene:MeOH (99:1 v/v) was then eluted and collected in 100 ml fractions (total volume 4 L), followed by a flush of toluene:MeOH (96:4 v/v).

[0127] Fractions were analysed by TLC (eluent: ethylacetate) to determine which fractions could be discarded and which fractions contained Z isomer. These Z fractions were then analyzed by HPLC. The pass criteria for a fraction was >96% Z isomer and <1.2% E isomer.

[0128] Surprisingly, the purification through Biotage chromatography of various batches was very efficient as the substantially pure form of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one O-methyloxime is purified at 99.4% (Batches No.020, No.062, No.068) and at 99.2% (Batches No.180, No.076). In particular, the Biotage chromatography in presence of oxime ester removes 5%w/w oxime ester without detriment to recovery or quality (Batch No.068) and a 25 % overcharge of the Biotage column does not cause a decrease in yield or quality(batch No.076).

Table 6: efficiency of the Biotage chromatography

Batch No.	Input % E/Z	Output % E/Z	yield of Z isomer
020	3.0 g 85.7% area purity % E/Z: 30.5/69.5	Pure Z-fractions: 1.0 g 98.8% area purity % E/Z: 0.6/99.4	33%
180	2.0 g 92.0% area purity % E/Z: 32.8/67.2	Pure Z-fractions 0.9 g 99.6%area purity % E/Z: 0.8/99.2	45%
062	3.0 g 83.5% area purity % E/Z: 32.7/67.3	Pure Z-fractions 1.3 g 99.8%area purity % E/Z: 0.6/99.4 Mixture: 1.2 g 91.0%area purity % E/Z: 69.6/30.4	43% 11%

(continued)

Batch No.	Input % E/Z	Output % E/Z	yield of Z isomer
068	3.0g spiked with ~5% ester ~78% area purity % E/Z: 32.7/67.3	Pure Z fractions: 1.2 g 99.8%area purity % E/Z: 0.6/99.4	40%
		Mixture: 0.6 g 98.8%area purity % E/Z: 27.9/72.1	14%
		Pure E fractions: 1.1 g 70.7%area purity % E/Z: 98.7/1.3 (19.3% ester)	N/A
076	3.8g 83.5%area purity % E/Z: 32.7/67.3	Pure Z fractions 1.4 g 99.8%area purity % E/Z: 0.8/99.2	37%
		Mixture: 1.8 g 95.0%area purity % E/Z: 63.6/36.4	17%

1.4.2 Large scale purification

[0129] Various batches of crude (3Z/E,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime (0.392kg, 1.16 mol, 1.0 wt) were charged to a Biotage 150L SIM unit as an approximate 50%w/w solution in toluene and purified using 1% methanol in toluene (150L) followed by 2% methanol in toluene (50L), fraction size 5.0L. The collected fractions were analysed by TLC¹⁵ and HPLC analyses, as appropriate. The fractions that were deemed to contain clean (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime (criteria: Z-isomer ≥96.00% area, E-isomer ≤ 1.20% area) were combined and concentrated under vacuum at 40 to 45°C. Absolute ethanol (2x 2L) was added to the residue and the solution concentrated under vacuum at 40 to 45°C until the foamy solid could be manipulated. The desired product, (3Z, 5S)-1-[(biphenyl-4-yl-carbonyl)-5-hydroxymethyl]pyrrolidine-3-one-O-methyloxime (0.089Kg, 22.7%w/w, ¹H NMR (CDCl₃) concordant with structure, 99.3%area by HPLC, 98,4:0.9 Z/E ratio was obtained as an off-white to light brown solid.

Table 7: Summary of purification of different batches of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime in substantially pure form.

Batch No.	Input (kg)	Output (kg)	Yield (%w/w)	% Z form (% area)	% E form (% area)
12	0.392	0.089	22.8	98.65	0.85
116	0.392	0.114	29	98.34	0.89
120	0.441	0.081	18.4	97.90	1.81
122	0.380	0.094	24.3	98.52	1.14
124	0.387	0.096	25.3	98.89	0.73
126	0.390	0.132	33.8	98.40	0.95
128	0.526	0.010	2	98.20	0.83

EP 3 266 449 B1

(continued)

Batch No.	Input (kg)	Output (kg)	Yield (%w/w)	% Z form (% area)	% E form (% area)
130	0.453	0.086	19	98.46	1.23
132	0.440	0.082	19.3	98.86	0.85
134	0.39	0.144	36.9	98.73	0.96
138	0.273	0.098	35.9	98.92	0.66
140	0.463	0.059	13.1	98.52	1.13
142	0.462	0.084	18.4	99.37	0.48
144	0.442	0.126	29	99.1	0.68
146	0.409	0.135	33.5	99.21	0.46
148	0.460	0.107	23.8	99.13	0.65
150	0.409	0.071	18	98.92	0.66
152	0.392	0.054	14.3	98.82	0.76
156	0.445	0.039	8.8	98.64	0.87
158	0.392	0.06	15.3	98.73	0.63
162	0.435	0.150	34.5	98.94	0.79
164	0.434	0.192	44.2	99.21	0.58
166	0.415	0.074	17.8	98.79	0.73
174	0.518	0.108	20.8	99.11	0.64
176	0.342	0.072	21	98.88	0.77
178	0.415	0.074	17.8	99.07	0.71
180	0.353	0.174	49.3	99.03	0.82
182	0.270	0.178	65.9	99.10	0.53

[0130] Appropriate batches of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime (2.713kg, 1.0 wt) isolated from the Biotage chromatography were combined and dissolved in absolute ethanol (5.16L, 2.0 vol) at 15 to 25°C, clarified by filtration through glass micro fibre paper and an absolute ethanol wash (0.50L, 0.2 vol) applied to the filter. The combined filtrates were concentrated portion wise under vacuum at 40 to 45°C. The resultant was transferred to drying trays and dried under vacuum at 30°C for 24 hours. The oven temperature was then increased incrementally from 30 to 40°C over 80 hours. The level of residual solvent was determined by ¹H NMR analysis (CDCl₃) and when found to be <1.0%w/w the solid was passed through a 500µm aperture sieve. The solid was returned to the oven and dried at 40 to 42°C until the solvent level was <0.40%w/w to afford (3Z, 5S)-1-[(biphenyl-4-yl-carbonyl)-5-hydroxy-methyl]-pyrrolidine-3-one-O-methyloxime (2.633Kg, 97.1%w/w, ¹H NMR (CDCl₃) concordant with structure, 98.65% area by HPLC.

[0131] The combination procedure is summarized below:

Input: 2.713kg
Output: 2.633kg
Yield: 97.1%w/w

Example 2: Capsule oral formulation

2.1 Bulk preparation.

[0132] Excipients were weighed directly into a beaker, which was transferred into a thermostatic water bath until all excipients were molten at 60°C. Then, always under controlled temperature, small aliquots of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime were added until all the drug was dissolved or

dispersed, under magnetic stirring, helix mixer or homogenizer.

2.2 Liquid filling capsules

[0133] The semi-solid bulk was maintained at 60°C during the liquid filling of the capsule shells. The filling step was performed at 60°C (both dosing pump and feeder) with an automatic lab-scale capsule-filler machine. The machine was set to the correct dosage by weighing the filled capsules.

Composition of active capsules (Table 8):

[0134]

Table 8:

Component	mg/capsule	
(3Z,5S)	30.0	300.0
Gelucire 50/13TM	33.4	334.0
LabrasolTM	16.6	166.0
Capsule, gelatine	Size 00	Size 00

Composition of placebo capsules (Table 9):

[0135]

Table 9:

Component	mg/capsule	
Gelucire 50/13TM	63.4	634.0
LabrasolTM	16.6	166.0
Capsule, gelatine	Size 00	Size 00

Example 3: Granules oral formulation

Granules 10% were prepared by hot melt granulation (Table 10):

[0136] Hot melt granulation was conducted in a high shear granulator MiMiPro (Procept) using different set-ups depending on the batch size. The general method of manufacture is characterized by the steps of:

- i) heat granulator water jacket to melting of waxy binder;
- ii) screen powders into the heated bowls and mix gently;
- iii) add waxy binder and mix gently; allow wax to soften;
- iv) granulate for few minutes; rest the material for few minutes and granulate again if necessary and
- v) cool and screen the granules.

[0137] In particular, the jacketed vessel was used at 65°C when Gelucire 50/13 was used as binder. Cover temperature was set at 5°C lower than the jacketed vessel's. Duration of phases depend on the desired particle size distribution and on the batch size.

Table 10:

Component	Amount w/w
(3Z,5S)	10.0%
Saccharose	10.0%
Acdisol	5.0%
Gelucire 50/13	15.0%
Sodium Saccharine	0.2%
Lactose anhydrous	59.8%

[0138] Hot melt granulation is an alternative technique of granulation: unlike the traditional use of aqueous or organic solvents as binders, in this process the agglomeration is obtained through the addition of a molten binder or a solid binder, which melts during the process and remains as a constituent of the formulation.

Composition of granules 5.8% prepared by spray-drying (Table 11):

[0139] The spray-drying process produces porous /hollow particles and amorphous forms of the sprayed material. This approach is used when dissolution rate improvement is required. The spray-drying process consists of four steps: atomisation of feed solution into a spray camera, spray-air contact involving flow and mixing, drying of sprayed droplets at elevated temperatures and separation of dried product from air.

[0140] The granules 5.8% were prepared by spray-drying of (3Z,5S), in presence of HP- β -CD (hydroxypropyl- β -cyclodextrin) in hydroethanolic solution. (3Z,5S) is an amorphous material that forms clumps of particles in water, which reduces the drug dissolution rate. Therefore, a hydrophilic excipient, spray-dried together with (3Z,5S) was used to improve the dissolution rate of the drug by preventing the aggregation in water. HP- β -CD was selected as hydrophilic excipients.

[0141] HP- β -CD spray dried product was obtained from the hydroalcoholic solution both using the Mini AirPro or Buchi equipment. The feeding solution was prepared by mixing an equal volume of a HP- β -CD (100g in 200 ml) aqueous solution and an (3Z,5S) (24 g in 200ml) ethanolic solution that was left for 24 hours under agitation at room temperature. The spray-drying conditions in the fluid bed Mini AirPro were: blower speed 1m³/min, nozzle pressure 1bar, liquid speed 3, inlet air temperature 70°C.

[0142] Spray-dried materials presented very poor flow properties precluding their use for sachet filling. To obtain an easy handling powder, dry granulation and ethanolic wet granulation were the processes applied to the (3Z,5S)-HP- β -CD spray-dried material. The dissolution rate of (3Z,5S) after spray-drying with HP- β -CD was almost instantaneous, in 15 minutes almost all drug was dissolved.

[0143] The granulation process by ethanolic wet granulation was performed on the (3Z,5S)-HP β CD spray-dried material. It did not modify the dissolution rate of the drug when compared to the (3Z,5S)-HP β CD spray-dried material.

[0144] It produced a material that showed almost instantaneous dissolution of the drug (Table 11).

Table 11:

Component	Amount w/w
(3Z,5S)	5.8%
Saccharose	10.0%
Acdisol	3.0%
Sodium saccharine	0.2%
HP- β -CD	24.2%
Poloxamer 188	2.0%
Avicel RC 591	10.0%
Avicel PH 112	10.0%
Lactose monohydrate	34.8%

Example 4: conventional tablet oral formulation

[0145] A solution containing (3Z,5S) dissolved in Labrasol:Ethanol 3:1 v/v was prepared by heating at 45°C and adding stepwise the required amount of drug. Zeopharm 600 was dried for 2 hours into a vacuum oven at 50°C. The addition of the solution on Zeopharm 600 bed was carried out into a 1900 ml bowl at 5 ml/min. The granulator was set as follows: impeller at 900 rpm, chopper at 3500 rpm, and cover temperature at 80° C. To remove most of the solvent, the material was left overnight at room temperature, and 3.5 hours in a vacuum oven at 50°C. As the granule-adsorbate particle size was slightly high, the material was sieve-milled first through a 1.5 mm sieve and then through a 1 mm sieve. The production yield, including the milling step, was 89.68%. The granule-adsorbate (87.5%) was then mixed with AcDiSol (4%), Compritol 888 ATO (3.5%), GL100 (0.2%) and Zeopharm 600 (4.8%) for 20 minutes at 22 r.p.m in the Turbula mixer. The final blend for tableting possessed a good (3Z,5S) content uniformity. The conventional tablets were produced by compression of the granules using the eccentric tableting machine EK-0.

[0146] Conventional tablets of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, were prepared using carriers of calcium silicate (Table 12).

Table 12:

Component	Amount w/w
(3Z,5S)	35%
Acdisol	4%
Calcium silicate (Zeopharm 600)	32.7%
Compritol ATO 888	3.5%
Labrasol	24.6%
Rx GL 100	0.2%

Example 5: Dispersible tablet oral formulation

[0147] A 850 g batch granulate was produced and then compressed into tablets. A 5000 ml vessel set-up was used for this preparation. Calcium silicate was vacuum dried prior to use. The wet granulation was conducted with (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime and all excipients (with the exception of the lubricant) in a high shear granulator at room temperature with only the cover of the granulator heated (65°C).

[0148] In the granulation process after a gentle pre-mix phase, an alternation of liquid addition and mixing phases of few minutes duration was set-up.

[0149] A total amount of 80 ml of ethanol was necessary to obtain suitable granules. A final milling and/or sieving step was necessary to obtain a better granule size distribution. The resulting granules were blended with the lubricant before compression.

[0150] A single punch eccentric tableting machine was used.

[0151] The compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, and/or an active metabolite thereof, is in the form of dispersible tablets containing 50 or 200 mg of active drug substance (Table 13).

Table 13:

Component	Amount			Function
	mg	mg	% (w/w)	
(3Z,5S)	50.0	200.00	20.0	Active drug substance
Calcium silicate	12.5	50.0	5.0	Carrier
PVP 30K	2.5	10.0	1.0	Binder
Poloxamer 188	5.0	20.0	2.0	Wetting agent
Sodium croscarmellose	12.5	50.0	5.0	Disintegrant

(continued)

Component	Amount			Function
	mg	mg	% (w/w)	
Microcrystalline cellulose 112	37.5	150.0	15.0	Diluent
Lactose monohydrate	119.5	478.0	47.8	Diluent
Sodium saccharine	0.5	2.0	0.2	Sweetener
Glycerol dibehenate	10.0	40.0	4.0	Lubricant
Total	250.0	1000.0		-

Example 6: Dimensions of the tablets

[0152] For example, the shape and dimensions of the tablets are the following:

Table 14:

Tablet or dispersible tablet	Shape and dimensions
50 mg	Capsule shape; 14 x 6 mm or 13 x 6 mm
200 mg	Capsule shape; 22 x 9 mm or 19 x 9 mm

Example 7: Pharmacokinetics studies in the dog

[0153] A pharmacokinetic study was conducted to measure the plasma concentration of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime following oral administration of said compound to female Beagle dogs. Following the protocol (Table 15), 5 dogs were administered with an oral formulation composed of liquid filled capsules ("Reference capsule"), 5 dogs were administered with an oral formulation composed of granules 10% (formulation 1), 5 dogs were administered with an oral formulation composed of granules 5.8% (formulation 2), 5 dogs were administered with a dispersible tablet (formulation 3), 5 dogs were administered with a conventional tablet (formulation 4). Also 5 dogs were administered by the IV route of administration ("Reference IV") a solution of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime at 15mg/kg. Blood sampling was performed at 0, 0.25, 0.5, 1, 2, 4, 6, 8, 24, 48 and 72h.

Table 15:

Group	Reference IV	Reference Capsule	Formulation 1	Formulation 2	Formulation 3	Formulation 4
Formulation	Solution of (3Z,5S)	Liquid filled capsules	Granules 10%	Granules 5.8%	Dispersible tablet 200mg	Conventional tablet 200mg
Dose (mg/kg)	15mg/kg	300 mg/dog	20	20	23.4(3.4) (200mg/dog)	23.3 (3.2) (200mg/dog)
Volume of administration		1 capsule size 00	5ml/kg	5ml/kg	50ml/dog	50ml/dog

7.1 Plasma concentration profiles of solid oral formulations in the dog

[0154] Figure 1A shows the plasma concentration vs. time profile of the different formulations over the time period from 0 to 72h. Figure 1B shows an enlargement of Figure 1A for the time period from 0 to 12h. The curve corresponding to formulation 3 (dispersible tablet) shows that the maximum concentration of the active substance is detected rapidly at about 0.5-1 hour following its administration. In comparison, the maximum concentration of the active substance is detected at about 2-4 hours following administration of formulation 4 (conventional tablet).

[0155] Individual plasma profile for each dog are presented on Figure 2 (formulation 3), and Figure 3 (formulation 4)

to show the inter-animal variability.

7.2 Pharmacokinetic parameters of solid oral formulations in the dog

[0156] In pharmacokinetics, the bioavailability is measured by calculating the area under the curve (AUC) of the product concentration vs. time profile. The absolute bioavailability compares the bioavailability of the product in systemic circulation following oral administration with the bioavailability of the product following intravenous administration.

[0157] "D" as used herein refers to the dose that is the amount of drug administered.

[0158] "C_{max}" as used herein refers to the peak plasma concentration of a drug after administration wherein the concentration is the amount of drug in a given volume of plasma.

[0159] "T_{max}" as used herein refers to the time to reach C_{max}.

[0160] "T_{1/2}" as used herein refers to the elimination half-life as the time required for the concentration of the drug to reach half of its original value.

[0161] "AUC" as used herein refers to the area under the curve that is the integral of the concentration-time curve (after a single dose or in steady state).

[0162] "F%" as used herein refers to the bioavailability that is the systemically available fraction of a drug. The index of bioavailability after oral administration is calculated by the following equation using the AUC found after i.v. administration:

$$F\% = (AUC_{OS}/AUC_{IV}) \times (Dose_{IV}/Dose_{OS}) \times 100$$

Table 16 :

Group	Reference IV	Reference Capsule	1	2	3	4
Formulation	IV	Capsule	Granules 10%	Granules 5.8%	Dispersible tablet	Conventional tablet
Route of administration	IV	OS	OS	OS	OS	OS
Dose (mg/kg)	15	34.4 (±4.0)	20	20	25.7 (±2.3)	23.3 (±3.2)
C (0.25h) (ng/ml)	-	13 (±15)	3590 (±1450)	4523 (±2757)	5108 (±1955)	88 (±54)
C _{max} (ng/ml)	17071 (±6162)	15243 (±3194)	7488 (±2236)	8722 (±2036)	13966 (±3217)	9245 (±2869)
T _{max} (h)	-	4 (4-4)	1 (0.5-2)	0.5 (0.5-0.5)	1 (0.5-1)	2 (2-4)
AUC (h*ng/ml)	38684 (±2310)	80139 (±18793)	32775 (±9120)	29893 (±7483)	58952 (±5770)	53563 (±7570)
T _{1/2} (h)	9.5 (±4.4)	7.1 (±1.2)	5.9 (±3.9)	4.7 (±0.9)	6.6 (±0.6)	5.0 (±1.7)
F (%)	-	90 (±15)	64 (±18)	58 (±15)	89 (±9)	89 (±13)

Numbers in brackets represent the standard deviation.

[0163] Formulations 1 and 2 (granules) showed equivalent responses and were characterized by a favorable fast compound absorption (Table 16). Their absolute bioavailability was good (about 60%), even though lower than the other formulations tested, but associated with a moderate variability between animals. Dispersible tablets Formulation 3 showed a rate of absorption comparable to granules followed by a decay comparable to the Reference Capsule formulation. A very high compound exposure highlighted by the absolute bioavailability (higher than after administration of granules and comparable to the liquid filled capsule reference formulation) was observed. Conventional tablets Formulation 4 gave a delayed absorption compared to Formulation 3 with a roughly similar bioavailability.

[0164] Formulation 3 appears overall the most suitable for the indication of preterm labor. By comparison with the Reference Capsule formulation (300 mg/dog, liquid filled capsules), the results highlight a faster compound absorption

EP 3 266 449 B1

(about 37% of the amount found at C_{max} already found at the first sampling time, 0.25 h), C_{max} reached at earlier time (median T_{max} = 1 h), terminal elimination rate comparable (T_{1/2} of about 7 h) and overall more uniform responses between animals. Formulation 3 exposure was equivalent to the Reference Capsule formulation and also absolute bioavailability was equivalent to that of the Reference Capsule formulation (89% vs. 90%).

7.3 Individual plasma concentration profiles of Formulation 3 in the dog

[0165]

Route of administration: Oral
Administered dose of (3Z,5S): 200 mg/dog
Formulation 3: 200 mg dispersible tablet
Dose regimen: Single

Table 17:

Sampling time (h)	Animal No. (3Z,5S) plasma concentrations in ng/mL				
	6F	7F	8F	9F	10F
0 (pre-dose)	1.5	2.3	*	3.4	*
0.25	8122	3965	2937	4994	5522
0.5	14107	9206	6913	16555	9070
1	18285	11934	11575	9137	11479
2	8175	7289	8341	7476	7942
4	5151	5188	6589	5660	4924
6	2508	2807	3434	2849	3027
8	1343	1626	2456	1819	3389
24	70	166	422	355	36
48	5	8.7	12	22	12
72	3.7	2	2.1	2.3	3.4
* = Below the lower limit of quantification (1 ng/mL)					

[0166] Remarkably, at 0.5 hour time point following administration of the dispersible tablet, the concentration in blood of the compound of formula (3Z,5S) is comprised between 59% to 100% of C_{max} indicating that said formulation is suitable for providing a rapid onset of action (Table 17).

7.4 Individual pharmacokinetic parameters of Formulation 3 in the dog

[0167]

Table 18:

PK parameter	Animal No.				
	6F	7F	8F	9F	10F
C _{max} (ng/ml)	18285	11934	11575	16555	11479
T _{max} (h)	1	1	1	0.5	1
C (0.25h) (ng/ml)	8122	3965	2937	4994	5522
AUC (h*ng/ml)	56242	52929	67865	61111	56615
T _{1/2} (h)	6	5.8	5.7	6.6	7.2

EP 3 266 449 B1

(continued)

PK parameter	Animal No.				
	6F	7F	8F	9F	10F
F (%)	85	80	102	92	85

[0168] The maximum concentration in blood of the compound of formula (3Z,5S) is reached at a time T_{max} between 0.5 to 1 hour following administration of Formulation 3 (dispersible tablet). In addition, Formulation 3 is characterized by a bioavailability of the compound of formula (3Z,5S) comprised between 80-100% (Table 18).

7.5 Individual plasma concentration profiles of Formulation 4 in the dog

[0169]

Table 19:

Sampling time (h)	Animal No. (3Z,5S) plasma concentrations in ng/mL				
	1F	2F	3F	4F	5F
0 (pre-dose)	*	*	*	*	*
0.25	152	17	53	116	104
0.5	2217	2379	1680	2926	1945
1	3907	3031	3241	4339	4482
2	7759	5761	6671	9227	13492
4	5171	5646	9983	7085	7848
6	2806	4822	4307	3856	3185
8	1527	4195	3051	2713	2211
24	87	78	53	92	172
48	1.8	3.1	1.9	4.5	4.5
72	*	*	*	1.2	1.6
* = Below the lower limit of quantification (1 ng/mL)					

[0170] The maximum concentration in blood of the compound of formula (3Z,5S) is reached at a time T_{max} between 2 to 4 hours following administration of Formulation 4 (conventional tablet).

[0171] At 0.5 hour time point following administration of Formulation 4 (conventional tablet), the concentration in blood of the compound of formula (3Z,5S) is comprised between 14% to 41% of C_{max} (Table 19) a value markedly lower than for formulation 3 which is characterized by a higher concentration of the compound of formula (3Z,5S) comprised between 59% to 100% of C_{max} (Table 17).

7.6 Individual pharmacokinetic parameters of Formulation 2 in the dog

[0172]

Table 20:

PK parameter	Animal No.				
	11F	12F	13F	14F	15F
C _{max} (ng/ml)	7029	9050	9897	6347	11287
T _{max} (h)	0.5	0.5	0.5	0.5	0.5
C (0.25h) (ng/ml)	3521	1275	4512	4445	8865

(continued)

PK parameter	Animal No.				
	11F	12F	13F	14F	15F
AUC (h.ng/ml)	21062	34881	39126	23931	30466
T _{1/2} (h)	3.3	5.3	4.5	5.1	5.4
F (%)	41	68	76	46	59

[0173] Formulation 2 (granule 5.8%) is characterized by a bioavailability of the compound of formula (3Z,5S) comprised between 41-76% (Table 20) markedly lower than for formulation 3 which is characterized by a very high bioavailability comprised between 80-100% (Table 18).

[0174] Thus, the dispersible tablet formulation 3 displays characteristics that are suitable for providing a fast onset of action and high bioavailability for the treatment of preterm labor. In contrast conventional tablet (formulation 4) or granules (formulation 2) do not meet the pharmacokinetic requirements for treating preterm labor.

Example 8: Pharmacokinetics studies in Human

Study protocol

[0175] Twelve healthy Caucasian women aged 54-62 years (mean 58.3 years), with a weight of 51 to 67 kg (mean 60.6 kg $\pm$ 5.1) and a body mass index ranging between 19.4-25.5 kg/m² (mean 23.12 kg/m² $\pm$ 2.05) were enrolled. They were administered, on three separate treatment periods of one week, either two consecutive (3Z,5S) doses of 600mg/day (administered to subjects using 3 dispersible tablets of 200mg in 150ml of water) or two intramuscular injections of 12 mg/day betamethasone or both drugs in combination.

Betamethasone (Célestène®, Schering-Plough, France) was administered by intramuscular injection of 12 mg/3ml into the gluteus muscle, which is a recommended dose for antenatal betamethasone in preterm labor for the prevention of respiratory distress syndrome in neonates.

[0176] One subject (Subject S6) was withdrawn from the study after the first treatment period, due to high pre-dose blood pressure in the second treatment period. Therefore, pharmacokinetic parameters of the treatment with (3Z,5S) dispersible tablets were calculated for 12 subjects, whereas pharmacokinetic parameters of the combination treatment were assessed for 11 participants.

[0177] During each treatment period of one week, blood samples were collected for the analysis of (3Z,5S) and betamethasone at time points 0 (pre-dose) and at 0.25, 0.5, 0.75, 1, 1.5, 2, 3, 4, 6, 8, 10, 12, 16, 20 and 24 hours after first dose on day 1. The plasma was prepared and stored below -20°C. All samples were analysed using validated LC-MS/MS methods. For the analysis of (3Z,5S), the lower limit of quantification was 1.0 ng/ml. For betamethasone analyses, the lower limit of quantification was 0.5 ng/ml.

Pharmacokinetic parameters were estimated by non-compartmental methods using the Phoenix® WinNonLin® version 6.3 (Pharsight). The following pharmacokinetic parameters were calculated for (3Z,5S) during each treatment period: measured maximum concentration (C_{max}), time to C_{max} (T_{max}), and area under the plasma concentration-time curve (AUC).

8.1 Individual plasma concentration profiles of (3Z, 5S) dispersible tablet (Formulation 3) in Human

[0178] Tables 21 to 23 show plasma concentration profiles of (3Z,5S) and pharmacokinetic parameters concerning (3Z,5S) which was administered to subjects using 3 dispersible tablets of 200mg in 150ml of water.

Table 21:

Time	Subject S Plasma concentration of (3Z,5S) in ng/ml											
	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12
0.25	1190	270	335	867	574	234	481	407	1730	1610	409	1990
0.5	3020	1170	1310	3120	2020	860	1550	1510	2630	2640	2510	2690
0.75	3010	1450	1810	2760	2220	1400	2090	1440	2700	2120	2350	3070
1	3100	2410	1810	3160	1830	1560	1440	1450	2670	2180	1940	2900

(continued)

	Subject S Plasma concentration of (3Z,5S) in ng/ml											
Time	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12
1.5	3280	3380	1650	4070	1600	1770	1790	1370	2570	3190	2990	3600
2	3170	3150	1520	3820	1550	1940	2440	2100	3050	2980	4100	3630
3	3060	3360	1540	3070	1560	2060	2280	2060	2660	2980	3410	3570
4	2870	2600	2300	2940	1990	3060	2290	2160	2730	2770	3300	3460
6	2400	2200	1910	1910	1580	2000	1800	1670	2280	1780	1960	1980
8	1710	1870	1800	1940	1310	1670	1560	1290	1690	1550	1620	1760
10	1510	1700	1620	1560	1130	1490	1370	973	1570	1430	1490	1490
12	1420	1870	1720	1630	949	1380	1250	794	1510	1400	1650	1410
16	1000	1040	1030	1170	640	981	762	586	928	844	1020	866
20	804	868	718	978	554	812	654	444	698	818	827	699
24	765	680	567	843	385	792	501	327	555	560	759	612

Table 22:

	Subject S % of Cmax											
Time	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12
0.25	36%	8%	15%	21%	26%	8%	20%	19%	57%	50%	10%	55%
0.5	92%	35%	57%	77%	91%	28%	64%	70%	86%	83%	61%	74%
0.75	92%	43%	79%	68%	100%	46%	86%	67%	89%	66%	57%	85%
1	95%	71%	79%	78%	82%	51%	59%	67%	88%	68%	47%	80%
1.5	100%	100%	72%	100%	72%	58%	73%	63%	84%	100%	73%	99%
2	97%	93%	66%	94%	70%	63%	100%	97%	100%	93%	100%	100%
3	93%	99%	67%	75%	70%	67%	93%	95%	87%	93%	83%	98%
4	88%	77%	100%	72%	90%	100%	94%	100%	90%	87%	80%	95%
6	73%	65%	83%	47%	71%	65%	74%	77%	75%	56%	48%	55%
8	52%	55%	78%	48%	59%	55%	64%	60%	55%	49%	40%	48%
10	46%	50%	70%	38%	51%	49%	56%	45%	51%	45%	36%	41%
12	43%	55%	75%	40%	43%	45%	51%	37%	50%	44%	40%	39%
16	30%	31%	45%	29%	29%	32%	31%	27%	30%	26%	25%	24%
20	25%	26%	31%	24%	25%	27%	27%	21%	23%	26%	20%	19%
24	23%	20%	25%	21%	17%	26%	21%	15%	18%	18%	19%	17%

[0179] Following administration of the dispersible tablet of the invention, the maximum concentration C_{max} in blood of the compound of formula (3Z,5S) is reached at a time T_{max} between 0.5 to 4 hours. In particular, T_{max} is in the range of 0.5 to 2 hours for 9 subjects, and in the range of 0.5 to 1.5 hours for 5 subjects.

[0180] Of note, at 0.5 hour time point, the concentration in blood of the compound of formula (3Z,5S) is comprised between 55% to 95% of C_{max} for 10 subjects, preferentially between 57% to 92% of C_{max}, indicating that the solid oral formulation is rapidly absorbed and suitable for providing a rapid onset of pharmacological action. For subjects S2 and S6, the plasma concentration of (3Z,5S) was below 55% at 0.5h, respectively 35% and 28% of C_{max}. However, it was respectively 71% and 51% of C_{max} at 1h following administration of the dispersible tablet which concentration is suitable

for the management of preterm labor.

8.2 Individual pharmacokinetic parameters of (3Z,5S) dispersible tablet (Formulation 3) in Human

[0181]

Table 23:

	Subject S						
	S1	S2	S3	S4	S5	S6	S7
C _{max} ng/ml	3280	3380	2300	4070	2220	3060	2440
T _{max} hour	1.5	1.5	4	1.5	0.75	4	2
AUC ng.h/ml	38557	39042	32421	41209	24814	33388	29867

	Subject S				
	S8	S9	S10	S11	S12
C _{max} ng/ml	2160	3050	3190	4100	3630
T _{max} hour	4	2	1.5	2	2
AUC ng.h/ml	24085	36317	34853	38944	38467

8.3 Plasma concentration profiles in Human for a combination of (3Z,5S) and betamethasone

[0182] Tables 24 to 26 show plasma concentration profiles of (3Z,5S) and pharmacokinetic parameters concerning (3Z,5S) administered in combination with Betamethasone.

Table 24: Plasma concentration of (3Z,5S)

	Subject S Plasma concentration of (3Z,5S) in ng/ml										
Time	S1	S2	S3	S4	S5	S7	S8	S9	S10	S11	S12
0.25	41.8	597	322	486	264	536	359	396	1150	934	934
0.5	210	3240	978	1890	1250	2510	1100	2420	2590	2040	1980
0.75	407	5290	1130	2210	1780	2910	1610	2570	2490	2060	3010
1	593	5320	1100	2290	1580	2600	1730	2250	2350	1930	3020
1.5	1050	4940	1350	3550	2010	2620	1860	2300	3130	1920	3430
2	1050	4750	2180	4230	2080	3290	2400	2270	3520	4170	3550
3	1480	4480	3500	3620	2280	3540	2520	2340	3150	4440	3780
4	2040	4130	3500	3880	2350	3430	2540	2310	3340	4020	3550
6	1440	2510	1710	2290	2050	2710	1600	2910	2350	2690	2670
8	1470	1950	1530	2180	1680	2180	1250	2280	1820	2300	1970
10	1570	2090	1250	2120	1270	1960	981	2120	1620	1840	1750
12	1300	1870	1430	1930	1040	1750	1000	1970	1430	1630	2130
16	521	1140	1000	1330	693	935	550	959	869	951	1010
20	512	1100	943	1240	481	831	471	825	746	786	877
24	418	828	712	1020	353	569	383	605	533	650	657

Table 25: % of Cmax

	Subject S										
	% of Cmax										
Time	S1	S2	S3	S4	S5	S7	S8	S9	S10	S11	S12
0.25	2%	11%	9%	11%	11%	15%	14%	14%	33%	21%	25%
0.5	10%	61%	28%	45%	53%	71%	43%	83%	74%	46%	52%
0.75	20%	99%	32%	52%	76%	82%	63%	88%	71%	46%	80%
1	29%	100%	31%	54%	67%	73%	68%	77%	67%	43%	80%
1.5	51%	93%	39%	84%	86%	74%	73%	79%	89%	43%	91%
2	51%	89%	62%	100%	89%	93%	94%	78%	100%	94%	94%
3	73%	84%	100%	86%	97%	100%	99%	80%	89%	100%	100%
4	100%	78%	100%	92%	100%	97%	100%	79%	95%	91%	94%
6	71%	47%	49%	54%	87%	77%	63%	100%	67%	61%	71%
8	72%	37%	44%	52%	71%	62%	49%	78%	52%	52%	52%
10	77%	39%	36%	50%	54%	55%	39%	73%	46%	41%	46%
12	64%	35%	41%	46%	44%	49%	39%	68%	41%	37%	56%
16	26%	21%	29%	31%	29%	26%	22%	33%	25%	21%	27%
20	25%	21%	27%	29%	20%	23%	19%	28%	21%	18%	23%
24	20%	16%	20%	24%	15%	16%	15%	21%	15%	15%	17%

[0183] Following administration of the combination (3Z,5S) dispersible tablet and betamethasone, the maximum concentration Cmax in blood of the compound of formula (3Z,5S) is reached at a time Tmax between 1 to 6 hours. In particular, Tmax is in the range of 2 to 4 hours for 9 subjects, and in the range of 2 to 3 hours for 6 subjects.

[0184] At 0.5 hour time point, the concentration in blood of the compound of formula (3Z,5S) is comprised between 43% to 83% of Cmax for 10 subjects indicating that the solid oral formulation administered in combination with betamethasone is rapidly absorbed and suitable for providing a rapid onset of pharmacological action. For subjects S1 and S3, the plasma concentration of (3Z,5S) was below 43% at 0.5h, respectively 10% and 28% of Cmax. However, it was respectively 51% and 62% of Cmax at 2h following administration of the dispersible tablet.

8.4 Pharmacokinetic parameters in Human for a combination of (3Z,5S) Formulation 3 and betamethasone

[0185]

Table 26:

Subject S	S1	S2	S3	S4	S5	S7	S8	S9	S10	S11	S12
Cmax ng/ml	2040	5320	3500	4230	2350	3540	2540	2910	3520	4440	3780
Tmax hour	4	1	3	2	4.03	3	4	6	2	3	3
AUC ng.h/ml	24060	50306	34518	47627	28322	42044	25974	39688	38208	43314	44082

Example 9: The manufacturing process of dispersible tablets

[0186] The manufacturing process of the dispersible tablets of the present invention comprises the following steps:

- (i) Preparing a mixture consisting of 20% of (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime, 5% of calcium silicate, 1% of PVP 30K, 2% of Poloxamer 188, 5% of sodium croscarmellose, 15% of Microcrystalline cellulose 112, 47.8% of lactose monohydrate, 0.2 % sodium saccharine, in weight based on the total weight of the tablet;

EP 3 266 449 B1

(ii) Wet granulating in presence of ethanol and vacuum drying;

(iii) Sieving the granules;

(iv) Blending the granules with 4% of glycerol dibehenate in weight based on the total weight of the tablet;

(v) Tableting.

[0187] The wet granulation is preferably conducted in a high shear granulator at room temperature with a minimal amount of ethanol equivalent to at least 7.4% (in weight based on the total weight of the tablet). The vacuum drying is performed at room temperature

[0188] A sieving step is applied on the resulting granules. Sieved glycerol dibehenate is blended with the granules. The final blend is compressed with an eccentric or rotary tablet press and adapted punches for the targeted dispersible tablet strength.

Example 10: Stability study of dispersible tablet (Formulation 3)

[0189] The determination of the (3Z,5S) content after dissolution of dispersible tablets is performed by HPLC using the following parameters:

Apparatus: USP II paddle apparatus

Dissolution medium: 0.5% Sodium Lauryl Sulfate in water

Dissolution medium volume: 900 ml

Dissolution medium temperature: 37°C +/- 0.5°C

Rotation speed: 50 rpm

Sampling time: 15, 30, 45, 60 and 120 min

Sampling volume: 3 ml

Separative technique: PALL Acrodisc PSF GxG/Glass 1.0 µm

[0190] The paddle assembly is arranged so that the bottom of the paddle was 2.5 cm ± 0.2 cm from the inside bottom of the flask. The appropriate volume of dissolution medium is poured into each one of the six dissolution vessels. The medium is equilibrated at 37.0°C ± 0.5°C. A (3Z,5S) dispersible tablet is inserted into each vessel. The paddles rotation was controlled at 50 rpm. At the designated time-points, 3 ml of medium is taken from a zone midway between the surface of the dissolution medium and the top of the blade of the paddle. Then, the sample is filtered through a PALL Acrodisc PSF GxG/Glass 1.0 µm directly into an HPLC vial for analysis.

[0191] Stability data are available at 1, 2 and 6 months storage time for (3Z,5S) dispersible tablets packaged in Alu/Alu blister packs. Parameters such as tablet appearance, (3Z,5S) content in % compared to the initial content value, disintegration time and dissolution were assessed (Tables 28 and 29).

[0192] The initial concentration value of (3Z,5S) measured by HPLC with a dispersible tablet is 90.0-110.0%.

Table 28: Dispersible tablet of 50mg

Storage condition	Storage Time (months)	Tablet appearance	Water content (%w/w)	(3Z,5S) Content (% initial value)	Disintegr. Time (sec)	% Dissolved after 60 min
Initial	0	White capsule shaped	4.5	102.9	42	101
5°C	1	White capsule shaped	4.7	102.4	19	100
	2	White capsule shaped	4.6	102.2	17	100
	6	White capsule shaped	4.8	103.3	26	99

EP 3 266 449 B1

(continued)

Storage condition	Storage Time (months)	Tablet appearance	Water content (%w/w)	(3Z,5S) Content (% initial value)	Disintegr. Time (sec)	% Dissolved after 60 min
25°C/ 60%RH	1	White capsule shaped	4.6	102.2	16	97
	2	White capsule shaped	4.6	103.8	22	97
	6	Off-white capsule shaped	4.7	105.7	28	99
40°C/ 75%RH	1	White capsule shaped	np	np	np	np
	2	Off-white capsule shaped	4.5	99.9	50	95
	6	Yellowish capsule shaped*	4.6	102.1	113	94
np = not performed						

Table 29: Dispersible tablet of 200mg

Storage condition	Storage Time (month)	Tablet appearance	Water content (%w/w)	(3Z,5S) Content (% initial value)	Disintegr. Time (sec)	% Dissolved after 60 min
Initial	0	White capsule shaped	4.5	98.1	102	93
5°C	1	White capsule shaped	4.0	97.7	26	99
	2	White capsule shaped	4.3	97.7	26	96
	6	White capsule shaped	4.1	99.7	28	99
25°C/ 60%RH	1	White capsule shaped	4.0	98.0	26	96
	2	White capsule shaped	4.2	98.9	28	100
	6	Off-white capsule shaped	4.1	99.5	40	98
40°C/ 75%RH	1	White capsule shaped	np	np	np	np
	2	Off-white capsule shaped	4.2	96.6	107	98
	6	Yellowish capsule shaped*	4.1	98.3	170	96
np = not performed						

[0193] A rapid disintegration of the tablet is observed following dissolution in the range of 20 to 40 sec at 25°C/60%RH.

[0194] The disintegration property of the dispersible tablet is likely due to its disintegrant component, for example,

Sodium croscarmellose that promotes the breakup or disintegration of the tablet when placed in an aqueous environment and supports a fast dissolution profile. In addition, the wetting agent, for example, Poloxamer 188, facilitates the water uptake during the disintegration and assists the drug dissolution. Furthermore, the carrier agent selected with a large surface area, for example Calcium silicate, was also identified as a potential benefit for the disintegration of the tablet.

Table 30: Dissolution profile of 200mg dispersible tablet and content of (3Z,5S) at 15, 30, 45, 60, and 120 min.

Storage Condition	Time (Months)	15 min (3Z,5S) Content %	30 min (3Z,5S) Content %	45 min (3Z,5S) Content %	60 min (3Z,5S) Content %	120 min (3Z,5S) Content %
Initial	0	89	94	96	97	99
5°C/ AmbH	1	91	95	96	97	99
	3	92	95	97	97	99
	6	95	98	98	100	101
25°C/60% RH	1	93	95	96	97	98
	3	94	97	98	98	100
	6	97	99	100	100	102
40°C/75% RH	1	96	98	99	99	100
	3	95	97	97	98	99
	6	95	97	98	98	99

[0195] A rapid dissolution profile of the tablet is observed for the different storage conditions. In particular, at 25°C/60%RH, the content of (3Z,5S) measured at 15 min is between 90 to 100% of the initial value.

Claims

1. A dispersible tablet comprising a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime in substantially pure form, and one or more pharmaceutically acceptable excipients, for use in a method of treating infertility, wherein the dispersible tablet is administered to a human patient undergoing an embryo transfer procedure.
2. The dispersible tablet according to claim 1, wherein the one or more pharmaceutically acceptable excipients comprise one or more of a disintegrant, a wetting agent, a carrier, a lubricant, a binder, a diluent, a sweetener, and a taste-masking agent.
3. The dispersible tablet according to claim 2, wherein the carrier is calcium silicate, calcium carbonate, calcium phosphate, tribasic calcium phosphate, lactose, starch, modified starch, a sugar, a cellulose, a cellulose derivative, a polymethacrylate, chitin, chitosan, or a combination thereof.
4. The dispersible tablet according to claim 2, wherein the binder is polyvinylpyrrolidone (PVP), cross-linked PVP, cellulose, a cellulose derivative, hydroxypropylmethyl cellulose (HPMC), carboxymethylcellulose sodium, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, methyl cellulose, carboxyethylcellulose calcium, guar gum, tragacanth, a polyvinylacetate, gelatin, pregelatinised starch, starch, a polyvinylalcohol, alginic acid, sodium alginate, sorbitol, glucose, magnesium aluminium silicate, dextrin, polyethylene glycol, a polymethacrylate, or a combination thereof.
5. The dispersible tablet according to claim 2, wherein the wetting agent is a poloxamer, sodium lauryl sulphate, a polyoxyethylene sorbitan fatty acid ester, polyoxyethylene stearate, a sorbitan fatty acid ester, or a combination thereof.
6. The dispersible tablet according to claim 2, wherein the disintegrant is sodium croscarmellose, crospovidone, sodium alginate, colloidal magnesium-aluminum silicate, calcium silicate, sodium starch glycolate, an acrylic acid derivative, microcrystalline cellulose, sodium carboxymethyl cellulose, calcium carboxymethyl cellulose, modified cellulose

gum, cross-linked povidone, alginic acid, an alginate, pregelatinised starch, modified corn starch, or a combination thereof.

7. The dispersible tablet according to claim 2, wherein the diluent is microcrystalline cellulose, lactose monohydrate, lactose, compressible sugar, sugar, dextrose, mannitol, dextrin, maltodextrin, sorbitol, xylitol, sodium chloride, calcium carbonate, magnesium carbonate, calcium phosphate, calcium sulphate, magnesium oxide, kaolin, powdered cellulose, pregelatinized starch, starch, barium sulphate, magnesium trisilicate, aluminium hydroxide, or a combination thereof.

8. The dispersible tablet according to claim 2, wherein the sweetener is sodium saccharine.

9. The dispersible tablet according to claim 2, wherein the lubricant is glycerol dibehenate, glycerol tribehenate, magnesium stearate, calcium stearate, talc, sodium stearyl fumarate, sodium behenate, stearic acid, cethyl alcohol, polyoxyethylene glycol, leucine, sodium benzoate, a stearate, polyethylene glycol, glyceryl monostearate, glyceryl palmitostearate, liquid paraffin, a poloxamer, sodium lauryl sulphate, magnesium lauryl sulphate, hydrogenated castor oil, colloidal silicon dioxide, a palmitostearate, zinc stearate, stearyl alcohol, hydrogenated vegetable oil, or a combination thereof.

10. The dispersible tablet according to claim 1, wherein the dispersible tablet comprises:

20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime;

1-20% by weight of calcium silicate;

0.1-20% by weight of PVP30K;

0.01-5% by weight of poloxamer 188;

0.5-20% by weight of sodium croscarmellose;

1-90% by weight of microcrystalline cellulose 112;

1-90% by weight of lactose monohydrate;

0.01-0.5% by weight of sodium saccharine; and

0.1-10% by weight of glycerol dibehenate.

11. The dispersible tablet according to claim 11, wherein the dispersible tablet consists essentially of:

20% by weight of a compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime;

5% by weight of calcium silicate;

1% by weight of PVP30K;

2% by weight of poloxamer 188;

5% by weight of sodium croscarmellose;

15% by weight of microcrystalline cellulose 112;

47.8% by weight of lactose monohydrate;

0.2% by weight of sodium saccharine; and

4% by weight of glycerol dibehenate.

12. The dispersible tablet according to any one of claims 1 to 11, wherein the purity of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime is at least in the range of from 85% to 99.9%.

13. The dispersible tablet according to any one of claims 1 to 11, wherein the purity of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime is from 90% to 99.9%.

14. The dispersible tablet according to any one of claims 1 to 11, wherein the purity of the compound of formula (3Z,5S)-5-(hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-one-O-methyloxime is from 95% to 99.9%.

15. The dispersible tablet according to any one of claims 1 to 14, wherein the method comprises *in vitro* fertilization (IVF) thereby producing an embryo, followed by transfer of the embryo (IVF-ET) to the patient.

Patentansprüche

1. Dispergierbare Tablette, mit einer Verbindung der Formel (3Z, 5S) -5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim in einer im Wesentlichen reinen Form, und einem oder mehreren pharmazeutisch annehmbaren Hilfsstoffen, zur Verwendung in einem Verfahren zur Behandlung von Unfruchtbarkeit, worin die dispergierbare Tablette an menschlichen Patienten, die sich ein Embryotransferverfahren unterziehen, verabreicht wird.
2. Dispergierbare Tablette gemäss Anspruch 1, worin der eine oder die mehreren pharmazeutisch annehmbare(n) Hilfsstoff(e) umfassen eines oder mehreren aus: einem Sprengmittel, einem Netzmittel, einem Träger, einem Schmiermittel, einem Bindemittel, einem Verdünnungsmittel, einem Süssungsmittel und einem Geschmacksmaskierungsmittel.
3. Dispergierbare Tablette gemäss Anspruch 2, worin der Träger Calciumsilikat, Calciumcarbonat, Calciumphosphat, tribasisches Calciumphosphat, Lactose, Stärke, modifizierte Stärke, Zucker, eine Cellulose, ein Cellulosederivat, ein Polymethacrylat, Chitin, Chitosan oder eine Kombination davon ist.
4. Dispergierbare Tablette gemäss Anspruch 2, worin das Bindemittel Polyvinylpyrrolidon (PVP), vernetzte PVP, Cellulose, ein Cellulosederivat, Hydroxypropylmethylcellulose (HPMC), Natriumcarboxymethylcellulose, Ethylcellulose, Hydroxyethylcellulose, Hydroxypropylcellulose, Methylcellulose, Carboxyethylcellulose Calcium, Guar gummi, Tragacanth, ein Polyvinylacetat, Gelatine, vorgelatinierte Stärke, Stärke, in Polyvinylalkohol, Alginsäure, Natriumalginat, Sorbit, Glucose, Magnesiumaluminiumsilikat, Dextrin, Polyethylenglykol, ein Polymethacrylat oder eine Kombination davon ist.
5. Dispergierbare Tablette gemäss Anspruch 2, worin das Benetzungsmittel ein Poloxamer, Natriumlaurylsulfat, ein Polyoxyethylensorbitanfettsäureester, Polyoxyethylenstearat, ein Sorbitanfettsäureester oder eine Kombination davon ist.
6. Dispergierbare Tablette gemäss Anspruch 2, worin das Sprengmittel Natriumcroscarmellose, Crospovidon, Natriumalginat, kolloidales Magnesiumaluminiumsilikat, Calciumsilicat, Natriumstärkeglykolat, ein Acrylsäurederivat, mikrokristalline Cellulose, Natriumcarboxymethylcellulose, Calciumcarboxymethylcellulose, modifizierter Cellulosegummi, vernetztes Povidon, Alginsäure, ein Alginat, vorgelatinierte Stärke, modifizierte Maisstärke oder eine Kombination davon ist.
7. Dispergierbare Tablette gemäss Anspruch 2, worin das Verdünnungsmittel mikrokristalline Cellulose, Lactose-Monohydrat, Lactose, kompressiblen Zucker, Zucker, Dextrose, Mannit, Dextrin, Maltodextrin, Sorbit, Xylit, Natriumchlorid, Calciumcarbonat, Magnesiumcarbonat, Calciumphosphat, Calciumsulfat, Magnesiumoxid, Kaolin, pulverförmige Cellulose, vorgelatinisierte Stärke, Stärke, Bariumsulfat, Magnesiumtrisilicat, Aluminiumhydroxid oder eine Kombination davon ist.
8. Dispergierbare Tablette gemäss Anspruch 2, worin das Süssungsmittel Natriumsaccharin ist.
9. Dispergierbare Tablette gemäss Anspruch 2, worin das Gleitmittel Glycerindibehenat, Glycerintribehenat, Magnesiumstearat, Calciumstearat, Talk, Natriumstearylformarat, Natriumbehenat, Stearinsäure, Cethylalkohol, Polyoxyethylenglykol, Leucin, Natriumbenzoat, ein Stearat, Polyethylenglykol, Glycerylmonostearat, Glycerylpalmitostearat, flüssiges Paraffin, ein Poloxamer, Natriumlaurylsulfat, Magnesiumlaurylsulfat, hydriertes Ricinusöl, kolloidales Siliciumdioxid, ein Palmitostearat, Zinkstearat, Stearylalkohol, hydriertes Pflanzenöl oder eine Kombination davon ist.
10. Dispergierbare Tablette gemäss Anspruch 1, worin die dispergierbare Tablette umfasst:
 - 20 Gew.-% einer Verbindung der Formel (3Z, 5S)-5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl)-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim;
 - 1-20 Gew.-% Calciumsilikat;
 - 0,1-20 Gew.-% PVP30K;
 - 0,01-5 Gew.-% Poloxamer 188;
 - 0,5-20 Gew.-% Natriumcroscarmellose;
 - 1-90 Gew.-% mikrokristalline Cellulose 112;
 - 1-90 Gew.-% Lactose-Monohydrat;

0,01-0,5 Gew.-% Natriumsaccharin; und
0,1-10 Gew.-% Glycerindibehenat.

11. Dispergierbare Tablette gemäss Anspruch 1, worin die dispergierbare Tablette im Wesentlichen besteht aus:

20 Gew.-% einer Verbindung der Formel (3Z, 5S)-5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim;
5 Gew.-% Calciumsilikat;
1 Gew.-% PVP30K;
2 Gew.-% Poloxamer 188;
5 Gew.-% Natriumcroscarmellose;
15 Gew.-% mikrokristalline Cellulose 112;
47,8 Gew.-% Lactose-Monohydrat;
0,2 Gew.-% Natriumsaccharin; und
4 Gew.-% Glycerindibehenat.

12. Dispergierbare Tablette gemäss irgendeinem der Ansprüche 1 bis 11, worin die Reinheit der Verbindung der Formel (3Z, 5S)-5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim mindestens innerhalb des Bereichs von 85% bis 99.9% liegt.

13. Dispergierbare Tablette gemäss irgendeinem der Ansprüche 1 bis 11, worin die Reinheit der Verbindung der Formel (3Z, 5S)-5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim mindestens innerhalb des Bereichs von 90% bis 99.9% liegt.

14. Dispergierbare Tablette gemäss irgendeinem der Ansprüche 1 bis 11, worin die Reinheit der Verbindung der Formel (3Z, 5S)-5-(Hydroxymethyl)-1-[(2'-methyl-1,1'-biphenyl-4-yl)carbonyl]pyrrolidin-3-on-O-methyloxim mindestens innerhalb des Bereichs von 95% bis 99.9% liegt.

15. Dispergierbare Tablette gemäss irgendeinem der Ansprüche 1 bis 14, worin das Verfahren eine *In-vitro*-Fertilisation (IVF) umfasst, wodurch ein Embryo produziert wird, gefolgt von einem Embryotransfer (IVF-ET) zur Patientin.

Revendications

1. Comprimé dispersible comprenant un composé de formule (3Z, 5S) -5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényl-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime sous forme essentiellement pure et un ou plusieurs excipients pharmaceutiquement acceptables, à usage dans un procédé de traitement de l'infertilité, dans lequel le comprimé dispersible est administré à un patient humain subissant une procédure de transfert d'embryon.

2. Comprimé dispersible selon la revendication 1, dans lequel le un ou les plusieurs excipients pharmaceutiquement acceptables comprennent un ou plusieurs parmi un désagréant, un agent mouillant, un support, un lubrifiant, un liant, un diluant, un édulcorant et un agent de masquage de goût.

3. Comprimé dispersible selon la revendication 2, dans lequel le support est le silicate de calcium, le carbonate de calcium, le phosphate de calcium, le phosphate de calcium tribasique, le lactose, l'amidon, l'amidon modifié, un sucre, une cellulose, un dérivé de cellulose, un polyméthacrylate, la chitine, le chitosan ou une combinaison de ceux-ci.

4. Comprimé dispersible selon la revendication 2, dans lequel le liant est la polyvinylpyrrolidone (PVP), la PVP réticulée, la cellulose, un dérivé de cellulose, l'hydroxypropylméthylcellulose (HPMC), la carboxyméthylcellulose sodique, l'éthylcellulose, l'hydroxyéthylcellulose, l'hydroxypropylcellulose, la méthylcellulose, la carboxyéthylcellulose calcique, la gomme de guar, la gomme adragante, les acétate de polyvinyle, la gélatine, l'amidon prégélatinisé, l'amidon, les polyvinylalcools, l'acide alginique, l'alginate de sodium, le sorbitol, le glucose, le silicate de magnésium et d'aluminium, la dextrine, le polyéthylèneglycol, un polyméthacrylate ou une combinaison de ceux-ci.

5. Comprimé dispersible selon la revendication 2, dans lequel l'agent mouillant est un poloxamère, le laurylsulfate de sodium, un ester d'acides gras de polyoxyéthylène sorbitane, le stéarate de polyoxyéthylène, un ester d'acides gras de sorbitane ou une combinaison de ceux-ci.

6. Comprimé dispersible selon la revendication 2, dans lequel le désintégrant est la croscarmellose au sodium, la crospovidone, l'alginate de sodium, le silicate de magnésium et d'aluminium colloïdal, le silicate de calcium, le glycolate d'amidon sodique, un dérivé d'acide acrylique, la cellulose microcristalline, la carboxyméthylcellulose sodique, la carboxyméthylcellulose au calcium, la gomme de cellulose modifiée, la povidone réticulée, l'acide alginique, un alginat, l'amidon prégélatinisé, l'amidon de maïs modifié ou une combinaison de ceux-ci.
7. Comprimé dispersible selon la revendication 2, dans lequel le diluant est la cellulose microcristalline, le lactose monohydraté, le lactose, le sucre compressible, le sucre, le dextrose, le mannitol, la dextrine, la maltodextrine, le sorbitol, le xylitol, le chlorure de sodium, le carbonate de calcium, le carbonate de magnésium, le phosphate de calcium, le sulfate de calcium, l'oxyde de magnésium, le kaolin, la cellulose en poudre, l'amidon prégélatinisé, l'amidon, le sulfate de baryum, le trisilicate de magnésium, l'hydroxyde d'aluminium ou une combinaison de ceux-ci.
8. Comprimé dispersible selon la revendication 2, dans lequel l'édulcorant est la saccharine sodique.
9. Comprimé dispersible selon la revendication 2, dans lequel le lubrifiant est le dibéhenate de glycérol, le tribéhenate de glycérol, le stéarate de magnésium, le stéarate de calcium, le talc, le fumarate de stéaryle de sodium, le béhénate de sodium, l'acide stéarique, l'alcool cétylique, le polyoxyéthylène glycol, la leucine, le benzoate de sodium, un stéarate, le polyéthylèneglycol, le monostéarate de glycéryle, le palmito-stéarate de glycéryle, la paraffine liquide, le poloxamère, le laurylsulfate de sodium, le laurylsulfate de magnésium, l'huile de ricin hydrogénée, le dioxyde de silicium colloïdal, un palmito-stéarate, le stéarate de zinc, l'alcool stéarylique, l'huile végétale hydrogénée ou une combinaison de ceux-ci.
10. Comprimé dispersible selon la revendication 1, dans lequel le comprimé dispersible comprend:
- 20% en poids d'un composé de formule (3Z, 5S)-5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényle-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime ;
 1-20% en poids de silicate de calcium ;
 0,1 à 20% en poids de PVP30K;
 0,01-5% en poids de poloxamère 188;
 0,5 à 20% en poids de croscarmellose de sodium;
 1-90% en poids de cellulose microcristalline 112;
 1-90% en poids de lactose monohydraté;
 0,01-0,5% en poids de saccharine de sodium; et
 0,1 à 10% en poids de dibéhenate de glycérol.
11. Comprimé dispersible selon la revendication 11, dans lequel le comprimé dispersible consiste essentiellement en:
- 20% en poids d'un composé de formule (3Z, 5S)-5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényle-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime,
 5% en poids de silicate de calcium ;
 1% en poids de PVP30K ;
 2% en poids de poloxamère 188 ;
 5% en poids de croscarmellose de sodium ;
 15% en poids de cellulose microcristalline 112 ;
 47,8% en poids de lactose monohydraté ;
 0,2% en poids de saccharine de sodium ; et
 4% en poids de dibéhenate de glycérol.
12. Comprimé dispersible selon l'une quelconque des revendications 1 à 11, dans lequel la pureté du composé de formule (3Z, 5S)-5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényle-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime est entre 85% à 99.9%.
13. Comprimé dispersible selon l'une quelconque des revendications 1 à 11, dans lequel la pureté du composé de formule (3Z, 5S)-5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényle-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime est entre 90% à 99.9%.
14. Comprimé dispersible selon l'une quelconque des revendications 1 à 11, dans lequel la pureté du composé de formule (3Z, 5S)-5-(hydroxyméthyl)-1-[(2'-méthyl-1,1'-biphényle-4-yl)carbonyl]pyrrolidin-3-one-O-méthylloxime est

entre 95% à 99.9%.

15. Comprimé dispersible selon l'une quelconque des revendications 1 à 14, dans lequel le procédé comprend une fécondation *in vitro* (FIV) produisant ainsi un embryon, suivi par le transfert de l'embryon (FIVETE) chez le patient.

5

10

15

20

25

30

35

40

45

50

55

Fig. 1A





Fig. 1B



—◆— Reference IV, 15 mg/kg,

—△— Reference capsule, 300 mg/dog

—■— Formulation 1, 20 mg/kg (10% granules)

—▲— Formulation 2, 20 mg/kg (5.8% granules)

—○— Formulation 3, 200 mg/dog (Dispersible tablets)

—●— Formulation 4, 200 mg/dog (Conventional tablets)

Fig. 2





Fig. 3



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0172705 A [0008]
- WO 02102799 A [0008]
- WO 2002074741 A [0008]
- WO 2004005249 A [0008] [0018] [0093]
- WO 2005082848 A [0018] [0093]
- EP 2014066075 W [0020]

Non-patent literature cited in the description

- **T. HIGUCHI ; V. STELLA.** Prodrugs as Novel Delivery Systems. A. C. S. *Symposium Series*, 1976, vol. 14 [0029]
- Design of Prodrugs. Elsevier, 1985 [0029]
- Bioreversible Carriers in Drug Design. American Pharmaceutical Association and Pergamon Press, 1987 [0029]