



(86) Date de dépôt PCT/PCT Filing Date: 2000/07/11
(87) Date publication PCT/PCT Publication Date: 2001/02/15
(85) Entrée phase nationale/National Entry: 2001/06/04
(86) N° demande PCT/PCT Application No.: EP 00/06590
(87) N° publication PCT/PCT Publication No.: WO 01/10443
(30) Priorité/Priority: 1999/08/10 (9918885.6) GB

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/4745, A61K 9/52, A61K 47/38

(71) Demandeur/Applicant:
PHARMACIA & UPJOHN S.P.A., IT


(72) Inventeurs/Inventors:
MARTINI, ALESSANDRO, IT;
MUGGETTI, LORENA, IT;
CIVAROLI, PAOLA, IT

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : FORMULATIONS PHARMACEUTIQUES EN CAPSULES D'HYDROXYPROPYL-METHYLCELLULOSE
(54) Title: PHARMACEUTICAL FORMULATIONS IN HYDROXYPROPYLMETHYLCELLULOSE CAPSULES

(57) **Abrégé/Abstract:**

An oral pharmaceutical formulation which comprises, in a hydroxypropylmethylcellulose capsule, a camptothecin analogue dispersed or solubilised in a semi-solid matrix. These formulations are useful for the treatment of neoplastic diseases.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 February 2001 (15.02.2001)

PCT

(10) International Publication Number
WO 01/10443 A1

(51) International Patent Classification⁷: **A61K 31/4745**,
9/52, 47/38

(21) International Application Number: **PCT/EP00/06590**

(22) International Filing Date: **11 July 2000 (11.07.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
9918885.6 **10 August 1999 (10.08.1999)** **GB**

(71) Applicant (for all designated States except US): **PHAR-**
MACIA & UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1.2,
I-20152 Milano (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CIVAROLI, Paola**
[IT/IT]; Via Giuseppe Frua, 20, I-20100 Milano (IT).
MUGGETTI, Lorena [IT/IT]; Via Trento, 38, I-20035
Meda (IT). **MARTINI, Alessandro** [IT/IT]; Via Desiderio
da Settignano, 14, I-20100 Milano (IT).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- *With international search report.*
- *Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/10443 A1

(54) Title: **PHARMACEUTICAL FORMULATIONS IN HYDROXYPROPYLMETHYLCELLULOSE CAPSULES**

(57) Abstract: An oral pharmaceutical formulation which comprises, in a hydroxypropylmethylcellulose capsule, a camptothecin analogue dispersed or solubilised in a semi-solid matrix. These formulations are useful for the treatment of neoplastic diseases.

PHARMACEUTICAL FORMULATIONS IN HYDROXYPROPYLMETHYLCELLULOSE CAPSULES

5

FIELD OF THE INVENTION

The present invention relates to oral dosage forms for camptothecin (CPT) analogues, such as, for example, (S)-[1,4'-Bipiperidine]-1'-carboxylic acid, 4,11-Diethyl-3,4,12,14-tetradhydro-4-hydroxy-3,14-dioxo-1H-pyrano [3',4':6,7]indolizino [1,2-b] quinolin-9-yl ester, monohydrochloride, trihydrate also known as irinotecan hydrochloride trihydrate or CPT-11, in hydroxypropylmethylcellulose capsules.

15 BACKGROUND OF THE INVENTION

CPT-11 is a water-soluble pro-drug of SN-38 (7-ethyl-10-hydroxy-CPT), a biologically potent derivative of the anticancer agent CPT, a topoisomerase I inhibitor. CPT-11 was proven efficacious in a variety of malignancies. Its clinical efficacy has been highlighted by the fact that CPT-11 is the first new active drug to obtain marketing approval in forty years for colorectal cancer.

The intravenous drug form of CPT-11 is currently used for the treatment of colorectal cancer.

25 It is well known that parenteral administration of antitumor drugs is associated with some intrinsic disadvantages and drawbacks, e.g., patient discomfort or the requirement for the patient to travel to the physician's office for drug administration, with obvious results in patient inconvenience.

30 Thus the need has arisen to find oral formulations of antitumoral drugs that would allow longer dosing regimens, as with continuous infusion, but without the inconvenience or the discomfort of the patient.

Classic oral formulations are, for example, solid oral dosage forms, that are medication delivery systems presented as solid

WO 01/10443

PCT/EP00/06590

2

dose units readily administered by mouth. The group includes tablets, capsules, cachets and pills, as well as bulk or unit-dose powders and granules. The group constitutes the most popular form of presentation, and tablets and capsules account for the
5 greatest number of preparations in this category.

It has long been known in the pharmaceutical industries that capsules are a convenient form for the oral administration of a variety of active agents because of their relative ease of manufacture (compared with other dosage forms such as tablets),
10 flexibility of size and dose. Capsules have traditionally been used for powder or granule formulations, but, in recent years, capsules have been adapted to contain the active ingredient in the form of paste, semi-solid or liquid formulation.

Since, for example, CPT-11 is classified as a class I cytotoxic
15 agent, any form of leakage from the dosage form would present a safety concern.

Therefore, formulations as tablets or powder-filled capsules are not as safe and user-friendly as semi-solid filled capsules, since the risks of leakage of the active ingredient from the unit
20 dosage form, both during manufacturing and distribution, is extremely high.

Thus, in light of the above-mentioned problem about the safe handling of CPT-11, it is desirable to formulate CPT-11 as a semi-solid dispersion or solution to be filled into capsules.

25 In particular, a thermoplastic hot-melt type capsule formulation would result in enhanced stability and minimization of leakage concerns.

Furthermore, it is known that the presence of certain active principles and/or of certain ingredients in the filling medium
30 can promote cross-linking in the capsule gelatin shell with the passage of time and/or under stressed conditions. When cross-linking occurs, the gelatin shell becomes less soluble in aqueous media. Cross-linking causes retardation of the disintegration of the capsule shell, and thus retardation of the dissolution of the
35 capsule contents, relative to identical capsules, which have not

been subjected to aging or storing at stressed conditions (i.e. temperature and humidity).

Thus, when a filled gelatin capsule contains an ingredient, which promotes cross-linking in the gelatin shell, such as, e.g. a
5 polyethylenglycol (PEG), it is challenging to prepare a formulation, which does not show retarded disintegration and/or dissolution when the formulation grows old.

There is therefore a need to prepare a safe and stable oral encapsulated formulation containing a CPT analogue which does not
10 present neither leakage nor stability problems.

The present invention fulfills such a need by providing stable oral pharmaceutical formulations of CPT analogues which avoid chemical interaction between the active ingredient and/or the excipients with capsule shells and maintain the dissolution
15 performances of the formulations with aging.

DESCRIPTION OF THE INVENTION

In one aspect of this invention there is provided an oral pharmaceutical formulation, which comprises a camptothecin
20 analogue dispersed or solubilised in a semi-solid matrix, filled into a hydroxypropyl-methylcellulose (HPMC) capsule.

The camptothecin analogue of the present invention is selected from the group comprising: CPT-11, topotecan, SN-38, SN-22, 9-amino-20(S)-CPT and 9-nitro-20(S)-CPT.

25 In particular, the present invention provides an oral pharmaceutical formulation, which comprises CPT-11, dispersed or solubilized into a semi-solid matrix, filled into a hydroxypropylmethylcellulose capsule.

A suitable semi-solid matrix for the formulation according to
30 this invention can be, e.g., a polyethylenglycol (PEG) in the molecular weight range between 400 and 20000, preferably between 1000 and 4000 and specifically 3000, optionally in combination with suitable excipients for semi-solid matrix compositions.

Suitable excipients can be, for example, pharmaceutical or food-
35 grade oils, e.g. soya or fractionated coconuts oils; surfactants,

e.g. polysorbates; poloxamers, i.e. polyoxyethylene-polyoxypropylene copolymers; solubilising agent, e.g. ethanol and triacetin; natural or synthetic glycerides, e.g. fractionated medium chain glycerides or saturated polyglycol glycerides; or
5 phospholipids.

The formulations according to the invention may be prepared by means of conventional techniques well known to one of ordinary skill in the art.

Typically, the semi-solid matrix to be filled into capsules may
10 be prepared by adding the camptothecin analogue to a molten homogeneous mixture of the excipients. This is then followed by thorough mixing of the molten mass and capsule filling using standard techniques. Hydroxypropylmethylcellulose capsules are chosen as primary packaging for such a formulation.

15 For example, the amount of CPT-11 may be in the range of from about 0.2 to about 200 mg, preferably from about 20 to about 100 mg per unit dose.

Generally, a semi-solid matrix formulation is a dispersion or a solution of the active ingredient in thermosoftening hot melt
20 inert carrier prepared by mixing or homogenization.

The obtained semi-solid matrix is therefore filled into capsules as liquid using fluid-filling pumps and allowed to solidify at ambient temperature. The major advantage of semi-solid formulations is the safety during manufacturing, being the drug
25 dispersed or dissolved in a liquid mass. At ambient condition such a formulation is solid, providing better chemical stability and minimizing leakage problems.

Selection of suitable excipients for semi-solid matrix formulations is commonly based on physico-chemical, rheological
30 and thermal properties, compatibility with drug and capsule shell as well as on the required drug release profile and bioavailability characteristics of the final dosage form.

In general, thermosoftening materials should have a melting temperature in the range of about 30°C to 70°C and a quite rapid
35 solidification time at ambient condition to avoid any leakage

phenomena from capsule after filling and/or during the sealing operation.

Drugs or excipients containing aldehyde groups, or producing aldehyde in decomposition, promote cross-linking in conventional capsules shells made from gelatin, forming a thin insoluble membrane that may delay dissolution.

Moreover, in semi-solid matrix formulations any interaction between the components and the capsule shell is magnified in comparison to powder filled capsules, being the surface contact area between the active ingredients/excipients and the gelatin shell higher.

It can be considered as an unexpected result the fact that the formulations of the present invention are able to guarantee the maintenance of the physico-chemical characteristics of the formulations during manufacturing and storage and to overcome the undesired cross-linking effect, that is practically highlighted by a reduced drug release profile from the dosage forms, especially with aging.

Dissolution rate test results, recovered during accelerated stability studies, as reported in the Examples 1 to 5, clearly demonstrate the increased stability shown by CPT-11 formulations placed into hydroxypropylmethylcellulose capsules according to the invention, when compared with the same CPT-11 formulations placed into conventional hard gelatin capsules.

The following examples are given with the purpose to better illustrate the invention but in no way they must be considered as a limitation of the scope of the invention itself.

It is to be understood that, although the examples reported in the description consider the use of CPT-11 as a representative compound of the camptothecin analogue compounds, the formulation approach according to the invention may be analogously applicable to other camptothecin analogues.

EXAMPLE 1

In the following table 1, the dissolution rate test results, recovered during an accelerated stability study, on Batch ND1216

WO 01/10443

PCT/EP00/06590

6

(CPT-11 semi-solid matrix formulation in PEG1000 - 100 mg CPT-11/capsule - Hard Gelatin Capsule as primary packaging) are shown.

5 The results are expressed as percent labeled amount (Release %) and its relative standard deviation percent (r.s.d. %) of the active dissolved after 60 minutes in simulated gastric fluid (SGF) without pepsin.

table 1

Storage (Time/conditions)	Release (%)	r.s.d. (%)	Dissolution medium
Initial	100.2	0.3	SGF without pepsin
1 month / 55°C	57.0	64.0	SGF without pepsin
1 month / 40°C+75%R.H.	74.6	20.3	SGF without pepsin

10

R.H. = Relative Humidity

As evident from the above—tabulated data, the release performances of the active from hard gelatin capsules decrease
15 with aging. No chemical degradation of the active, that could justify such a reduction in the dissolution release performances, was observed.

EXAMPLE 2

20 In the following table 2, the dissolution rate test results, recovered during an accelerated stability study, on Batch D54G01 (CPT-11 semi-solid matrix formulation in PEG1000 - 50 mg CPT-11/capsule - Hard Gelatin Capsule as primary packaging) are shown.

25 The results are expressed as percent labeled amount (Release %) and its relative standard deviation percent (r.s.d. %) of the active dissolved after 60 minutes in simulated gastric fluid (SGF) without pepsin.

WO 01/10443

PCT/EP00/06590

7

Table 2

Storage (Time/conditions)	Release (%)	r.s.d. (%)	Dissolution medium
Initial	100.1	0.4	SGF without pepsin
1 month / 40°C+75%R.H.	56.0	67.4	SGF without pepsin
1 month / 40°C+75%R.H.	97.5 (1)	2.2	SGF without pepsin

R.H. = Relative Humidity

5 (1)Dissolution performed on 'capsule content'

As evident from the above-tabulated data, the release performances of the active from hard gelatin capsules decrease with aging. No chemical degradation of the active, that could
10 justify such reduction in the dissolution release performances, was observed.

The good dissolution performance of the formulation stored for 1 month at 40°C 75% R.H., by removing the content from the gelatin shell before the dissolution testing, is the clear
15 demonstration that the reduction in the release profile is exclusively due to the partial insolubilization of the capsule shell induced by the cross-linking process.

20 EXAMPLE 3

In the following table 3, the dissolution rate test results, recovered during an accelerated stability study on Batch ND1283 (CPT-11 semi-solid matrix formulation in PEG3000 - 100 mg CPT-11/capsule - Hard Gelatin Capsule as primary packaging) are
25 shown.

The results are expressed as percent labeled amount (Release %) and as minimum and maximum release percent value of the active dissolved after 60 minutes in simulated gastric fluid (SGF) with and without pepsin.

WO 01/10443

PCT/EP00/06590

8

The use of pepsin is allowed by regulatory authorities in order to overcome the cross-linking issue of hard gelatin capsules. In fact, if the cross-linked gelatin is disrupted by the mechanical forces of gastric emptying or is broken down by digestive enzymes, its formation would not affect absorption and bioavailability of the active. The United States Food and Drug Administration (FDA) / Industry Gelatin Capsule Working Group concluded that formation of an insoluble membrane could be considered to have a negligible impact on drug bioavailability if the capsules dissolved during the 'two-tiered dissolution test' which employs a methods containing digestive enzymes.

Table 3

Storage (Time/conditions)	Release (%)	Minimum maximum (%) value	and release Dissolution medium
Initial	92.1	87.9 - 97.4	SGF without pepsin
1 month / 55°C	9.8	5.2 - 16.8	SGF without pepsin
1 month / 40°C+75%R.H.	6.3	3.7 - 11.5	SGF without pepsin
1 month / 55°C	89.6	91.2 - 97.4	SGF with pepsin
1 month / 40°C+75%R.H.	91.9	89.2 - 96.1	SGF with pepsin
3 month / 40°C+75%R.H.	69.6	37.6 - 89.9	SGF with pepsin

R.H. = Relative Humidity

It is clearly evident from the above-tabulated data that the addition of pepsin to the dissolution medium is not sufficient to overcome the problem. After 3 month storage at 40°C and 75%R.H., not even the addition of pepsin to the dissolution medium has allowed obtaining a complete release of the active from the formulation. No chemical degradation of the active, that could justify such reduction in the dissolution release performances, was observed.

EXAMPLE 4

In the following table 4, the dissolution rate test results, recovered during an accelerated stability study, on Batch ND1651 (CPT-11 semi-solid matrix formulation in PEG3000 - 50 mg CPT-11/capsule - Hydroxypropylmethylcellulose Capsule as primary packaging) are shown.

The results are expressed as percent labeled amount (Release %) and its relative standard deviation percent (r.s.d. %) of the active dissolved after 60 minutes in simulated gastric fluid (SGF) without pepsin.

Table 4

Storage (Time/conditions)	Release (%)	r.s.d. (%)	Dissolution medium
Initial	101.5	2.5	SGF without pepsin
1 month / 55°C	100.2	10.5	SGF without pepsin
3 month / 40°+75%R.H.	106.5	1.9	SGF without pepsin

It is clearly evident from the above-tabulated data that there is not any influence of aging on the release performances of the active from the HPMC formulation. The same unit dosage strength (i.e. 50 mg CPT-11/capsule), packed into hard gelatin capsule, showed reduced dissolution behavior already after 1 month storage at 40°C + 75% R.H. (as above shown in Example 2), while a 3 month storage at the same condition has no effect at all on the dissolution performance of the formulation manufactured into hydroxypropylmethylcellulose capsule. No chemical degradation of the active was observed.

EXAMPLE 5

In the following table 5, the dissolution rate test results, recovered during an accelerated stability study, on Batch ND1655 (CPT-11 semi-solid matrix formulation in PEG3000 - 100 mg CPT-

10

11/capsule - Hydroxypropylmethylcellulose Capsule as primary packaging) are shown.

The results are expressed as percent labeled amount (Release %) and its relative standard deviation percent (r.s.d. %) of the active dissolved after 60 minutes in simulated gastric fluid (SGF) without pepsin.

Table 5

Storage (Time/conditions)	Release (%)	r.s.d. (%)	Dissolution medium
Initial	90.3	4.1	SGF without pepsin
1 month / 55°C	90.2	1.6	SGF without pepsin
1 month / 40°C+75%R.H.	94.6	2.0	SGF without pepsin
3 month / 40°C+75%R.H.	92.3	2.0	SGF without pepsin

It is clearly evident from the above-tabulated data that there is not any influence of aging on the release performances of the active from the HPMC encapsulated formulation. The same unit dosage strength (i.e. 100 mg CPT-11/capsule), packed into hard gelatin capsule, showed reduced dissolution behavior already after 1 month storage at 40°C + 75% R.H. (as above shown in Example 3), while a 3 month storage at the same condition has no effect at all on the dissolution performance of the formulation manufactured into hydroxypropylmethylcellulose capsule. No chemical degradation of the active was observed.

20

EXAMPLE 6

In this example, typical semi-solid filling matrices, suitable for hydroxypropylmethylcellulose capsules, are shown. The below mentioned formulae are not intended to be exhaustive or to limit anyway the scope of the invention itself.

25

WO 01/10443

PCT/EP00/06590

11

FORMULA A	<u>mg per capsule</u>	<u>mg per capsule</u>
CPT-11	50 mg	100 mg
Polyethylenglycol3000	575 mg	525 mg
Fill weight per capsule	625 mg	625 mg

5

Hydroxypropylmethylcellulose capsules size 0

FORMULA B	<u>mg per capsule</u>	<u>mg per capsule</u>
CPT-11	50 mg	100 mg
10 Polyethylenglycol1000	575 mg	525 mg
Fill weight per capsule	625 mg	625 mg

Hydroxypropylmethylcellulose capsules size 0

15 FORMULA C	<u>mg per capsule</u>	<u>mg per capsule</u>
CPT-11	50 mg	100 mg
Polyethylenglycol3000	520 mg	470 mg
Triacetin	50 mg	50 mg
Polysorbate80	5 mg	5 mg
20 Fill weight per capsule	625 mg	625 mg

Hydroxypropylmethylcellulose capsules size 0

FORMULA D	<u>mg per capsule</u>	<u>mg per capsule</u>
25 CPT-11	50 mg	100 mg
Polyethylenglycol3000	520 mg	470 mg
Polyethylenglycol400	50 mg	50 mg
Polysorbate80	5 mg	5 mg
Fill weight per capsule	625 mg	625 mg

30

Hydroxypropylmethylcellulose capsules size 0

FORMULA E	<u>mg per capsule</u>	<u>mg per capsule</u>
CPT-11	50 mg	100 mg
35 Polyethylenglycol3000	525 mg	475 mg
Gelucire44/14	50 mg	50 mg

WO 01/10443

PCT/EP00/06590

12

Fill weight per capsule 625 mg 625 mg

Hydroxypropylmethylcellulose capsules size 0

- 5 Another aspect of the present invention provides a method for treating a tumor in a mammal, including a human, which comprises administering to said mammal an oral pharmaceutical formulation of a camptothecin analogue as described in the present invention.
- 10 The formulations according to the invention are useful for treating neoplastic diseases, reducing tumor burden, preventing or treating metastasis of neoplasms and preventing recurrences of tumor/neoplastic growths in mammals.
- In particular, formulations according to the present invention,
- 15 wherein the camptothecin analogue is CPT-11, are suitable for the treatment of colorectal cancer.

CLAIMS

1. An oral pharmaceutical formulation which comprises, in a hydroxypropylmethylcellulose capsule, a camptothecin analogue dispersed or solubilised in a semi-solid matrix.
5
2. A formulation according to claim 1, wherein the camptothecin analogue is selected from CPT-11, topotecan, SN-38, SN-22, 9-amino-20(S)-CPT and 9-nitro-20(S)-CPT.
10
3. A formulation according to claim 1 wherein the camptothecin analogue is CPT-11.
4. A formulation according to claim 3 which comprises from 0.2 to 200 mg per unit dose of CPT-11.
15
5. A formulation according to claim 4 which comprises from 20 to 100 mg per unit dose of CPT-11.
- 20 6. A formulation according to any one of the preceding claims wherein the semi-solid matrix is a polyethyleneglycol with a molecular weight of from 400 to 20000.
7. A formulation according to claim 6, wherein the polyethyleneglycol has a molecular weight of from 1000 to 4000.
25
8. A formulation according to claim 7 wherein the polyethyleneglycol has a molecular weight of 3000.
30
9. A formulation according to any one of the preceding claims which further comprises an excipient for the semi-solid matrix.
- 35 10. A formulation according to claim 9 wherein the excipient is selected from a food-grade oil, a surfactant, a poloxamer, a solubilising agent, a natural or synthetic glyceride and a phospholipid.

WO 01/10443

PCT/EP00/06590

14

11. An oral pharmaceutical formulation as defined in claim 1 for use in a method of treatment of the human or animal body by therapy.

5 12. A formulation as claimed in claim 11 for use in the treatment of neoplastic diseases, the reduction of tumors, the prevention or treatment of neoplasm metastasis or the prevention of recurrences of tumor or neoplastic growths in mammals.

10

13. A formulation as defined in claim 3, for use in the treatment of colorectal cancer.