



(86) **Date de dépôt PCT/PCT Filing Date:** 2007/10/18  
(87) **Date publication PCT/PCT Publication Date:** 2008/04/24  
(45) **Date de délivrance/Issue Date:** 2018/06/19  
(85) **Entrée phase nationale/National Entry:** 2009/04/15  
(86) **N° demande PCT/PCT Application No.:** GB 2007/003983  
(87) **N° publication PCT/PCT Publication No.:** 2008/047134  
(30) **Priorité/Priority:** 2006/10/18 (GB0620729.4)

(51) **Cl.Int./Int.Cl. C07K 16/24** (2006.01),  
**A61K 39/395** (2006.01), **A61P 35/00** (2006.01),  
**A61P 37/06** (2006.01), **C12N 15/13** (2006.01)

(72) **Inventeurs/Inventors:**  
ADAMS, RALPH, GB;  
POPPELWELL, ANDREW GEORGE, GB;  
RAPECKI, STEPHEN EDWARD, GB

(73) **Propriétaire/Owner:**  
UCB BIOPHARMA SPRL, BE

(74) **Agent:** ROBIC

(54) **Titre : MOLECULES D'ANTICORPS QUI SE LIENT A IL-17A ET A IL-17F**  
(54) **Title: ANTIBODY MOLECULES WHICH BIND IL-17A AND IL-17F**

(57) **Abrégé/Abstract:**

The invention relates to antibody molecules having specificity for antigenic determinants of both IL-17A and IL-17F, therapeutic uses of the antibody molecules and methods for producing said antibody molecules.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 April 2008 (24.04.2008)

PCT

(10) International Publication Number  
**WO 2008/047134 A3**

## (51) International Patent Classification:

*C07K 16/24* (2006.01)      *A61P 37/06* (2006.01)  
*A61K 39/395* (2006.01)    *A61P 35/00* (2006.01)  
*C12N 15/13* (2006.01)

## (21) International Application Number:

PCT/GB2007/003983

(22) International Filing Date: 18 October 2007 (18.10.2007)

(25) Filing Language: English

(26) Publication Language: English

## (30) Priority Data:

0620729.4      18 October 2006 (18.10.2006)      GB

(71) Applicant (for all designated States except US): **UCB PHARMA S.A.** [BE/BE]; 60 Allée de la Recherche, B-1070 Brussels (BE).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): **ADAMS, Ralph** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). **POPPEWELL, Andrew, George** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). **RAPECKI, Stephen, Edward** [GB/GB]; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).

(74) Agent: **THOMPSON, John**; UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(88) Date of publication of the international search report:

12 June 2008

(54) Title: ANTIBODY MOLECULES WHICH BIND IL-17A AND IL-17F

(57) Abstract: The invention relates to antibody molecules having specificity for antigenic determinants of both IL-17A and IL-17F, therapeutic uses of the antibody molecules and methods for producing said antibody molecules.

WO 2008/047134 A3

## Antibody molecules which bind IL-17A and IL-17F

The present invention relates to antibody molecules having specificity for antigenic determinants of both IL-17A and IL-17F. The present invention also relates to the therapeutic  
5 uses of the antibody molecules and methods for producing them.

Interleukin 17 (IL-17), also known as CTLA-8 or IL-17A, is a pro-inflammatory cytokine which stimulates the secretion of a wide range of other cytokines from various non-immune cells. IL-17A is capable of inducing the secretion of IL-6, IL-8, PGE2, MCP-1 and G-CSF by adherent cells like fibroblasts, keratinocytes, epithelial and endothelial cells and is  
10 also able to induce ICAM-1 surface expression, proliferation of T cells, and growth and differentiation of CD34+ human progenitors into neutrophils when cocultured in the presence of irradiated fibroblasts (Fossiez *et al.*, 1998, *Int.Rev.Immunol.* 16, 541-551). IL-17A is predominantly produced by activated memory T cells and acts by binding to a ubiquitously distributed cell surface receptor (IL-17R) (Yao *et al.*, 1997, *Cytokine*, 9, 794-800). It may also  
15 act through binding to a complex of IL-17RA and IL-17RC (Toy *et al.*, 2006, *J. Immunol.* 177(11);36-39). IL-17 producing T cells called 'TH17 cells' have been implicated in the pathogenesis of certain cancers (Weaver *et al.*, 2006, *Immunity*, 24, 677-688; Langowski *et al.*, 2006, 442, 461-465; Iwakura and Ishigame, 2006, *J.Clin.Invest.* 116, 5, 1218-1222).

A number of homologues of IL-17 have been identified which have both similar and  
20 distinct roles in regulating inflammatory responses. For a review of IL-17 cytokine/receptor families see Dumont, 2003, *Expert Opin. Ther. Patents*, 13, 287-303. One such homologue is IL-17F, also known as IL-24 and ML-1, which is around 55% identical to IL-17A and is thought to share the same receptors as IL-17A (Kolls and Linden 2004, *Immunity*, 21, 467-476; Hymowitz, *et al.*, 2001, *EMBO J.* 20(19), 5332-5341; Kuestner *et al.*, 2007, *Journal of*  
25 *Immunology*, 179, 5462-5473).

Both IL-17A and IL-17F can form both homodimeric and heterodimeric proteins, all of which are produced by activated human CD4+ T cells (Wright *et al.*, 2007, *J Biol Chem.* 282 (18), 13447-13455).

IL-17 may contribute to a number of diseases mediated by abnormal immune  
30 responses, such as rheumatoid arthritis and air-way inflammation, as well as organ transplant rejection and antitumour immunity. Inhibitors of IL-17 activity are well known in the art for example a murine IL-17R:human Fc fusion protein, a murine soluble IL-17R and an anti-IL-

17 monoclonal antibody have been used to demonstrate the role of IL-17 in various models of rheumatoid arthritis (Lubberts *et al.*, J.Immunol. 2001,167, 1004-1013; Chabaud *et al.*, Arthritis Res. 2001, 3, 168-177). In addition, neutralising polyclonal antibodies have been used to reduce peritoneal adhesion formation (Chung *et al.*, 2002, J.Exp.Med., 195, 1471-  
5 1478). Rat derived anti-human IL-17 antibodies were described in WO04/106377. A humanised anti-IL-17 antibody with an affinity of around 220pM was described in WO2006/054059. A monoclonal anti-IL-17 fully human antibody with an affinity of around 188pM was described in WO2006/013107. Antibodies which bind IL-17F and IL-17A/IL-17F heterodimers were described in WO2006/088833. Antibodies which specifically bind the  
10 IL-17A/IL-17F heterodimer were described in WO2005/010044.

IL-17F antagonism has been associated with protection against asthma (Kawaguchi *et al.*, 2006, J.Allergy Clin. Immunol. 117(4); 795-801) and IL-17F is also thought to play a role in arthritis pathology (Lubberts 2003, Current Opinion in Investigational Drugs, 4 (5), 572-577).

15 Accordingly dual antagonists of IL-17A and IL-17F may be more effective than a sole antagonist in treating IL-17 mediated diseases. Antibodies which bind IL-17A and IL-17F were described in WO2007/106769 published 20.9.07.

We have been able to demonstrate that it is possible to isolate an antibody which is capable of binding to both IL-17A and IL-17F and is capable of neutralising the activity of  
20 both isoforms of IL-17. Hence the present invention provides an anti-IL-17 antibody which is capable of binding to both IL-17A and IL-17F. In particular, the antibody of the present invention is capable of specifically binding to both IL-17A and IL-17F i.e. the antibody does not bind to other isoforms of IL-17. Preferably the antibody of the present invention also binds the IL-17A/IL-17F heterodimer. Preferably, the antibody of the present invention  
25 neutralises the activity of both IL-17A and IL-17F. In one embodiment the antibody of the present invention also neutralises the activity of the IL-17A/IL-17F heterodimer. The antibodies of the present invention therefore have the advantageous property that they can inhibit the biological activity of both IL-17A and IL-17F. Accordingly, the present invention also provides the use of such antibodies in the treatment of and/or prophylaxis of a disease  
30 mediated by either or both of IL-17A or IL-17F such as autoimmune or inflammatory disease or cancer.

As used herein, the term 'neutralising antibody' describes an antibody that is capable of neutralising the biological signalling activity of both IL-17A and IL17F for example by blocking binding of IL-17A and IL17F to one or more of their receptors and by blocking binding of the IL-17A/IL-17F heterodimer to one or more of its receptors. It will be appreciated that the term 'neutralising' as used herein refers to a reduction in biological signalling activity which may be partial or complete. Further, it will be appreciated that the extent of neutralisation of IL-17A and IL-17F activity by the antibody may be the same or different. In one embodiment the extent of neutralisation of the activity of the IL-17A/IL-17F heterodimer may be the same or different as the extent of neutralisation of IL-17A or IL-17F activity.

In one embodiment the antibodies of the present invention specifically bind to IL-17A and IL-17F. Specifically binding means that the antibodies have a greater affinity for IL-17A and IL-17F polypeptides (including the IL-17A/IL-17F heterodimer) than for other polypeptides. Preferably the IL-17A and IL-17F polypeptides are human. In one embodiment the antibody also binds cynomolgus IL-17F.

IL-17A or IL-17F polypeptides or a mixture of the two or cells expressing one or both of said polypeptides can be used to produce antibodies which specifically recognise both polypeptides. The IL-17 polypeptides (IL-17A and IL-17F) may be 'mature' polypeptides or biologically active fragments or derivatives thereof which preferably include the receptor binding site. Preferably the IL-17 polypeptides are the mature polypeptides. IL-17 polypeptides may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems or they may be recovered from natural biological sources. In the present application, the term "polypeptides" includes peptides, polypeptides and proteins. These are used interchangeably unless otherwise specified. The IL-17 polypeptide may in some instances be part of a larger protein such as a fusion protein for example fused to an affinity tag. Antibodies generated against these polypeptides may be obtained, where immunisation of an animal is necessary, by administering the polypeptides to an animal, preferably a non-human animal, using well-known and routine protocols, see for example Handbook of Experimental Immunology, D. M. Weir (ed.), Vol 4, Blackwell Scientific Publishers, Oxford, England, 1986). Many warm-blooded animals, such as rabbits, mice, rats, sheep, cows or pigs may be immunized. However, mice, rabbits, pigs and rats are generally preferred.

Antibodies for use in the present invention include whole antibodies and functionally active fragments or derivatives thereof and may be, but are not limited to, monoclonal, multi-valent, multi-specific, humanized or chimeric antibodies, domain antibodies e.g. VH, VL, VHH, single chain antibodies, Fab fragments, Fab' and F(ab')<sub>2</sub> fragments and epitope-binding  
5 fragments of any of the above. Other antibody fragments include those described in International patent applications WO2005003169, WO2005003170 and WO2005003171. Antibody fragments and methods of producing them are well known in the art, see for example Verma *et al.*, 1998, Journal of Immunological Methods, 216, 165-181; Adair and Lawson, 2005. Therapeutic antibodies. *Drug Design Reviews - Online* 2(3):209-217.

10 Antibodies for use in the present invention include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.* molecules that contain an antigen binding site that specifically binds an antigen. The immunoglobulin molecules of the invention can be of any class (*e.g.* IgG, IgE, IgM, IgD and IgA) or subclass of immunoglobulin molecule.

15 Monoclonal antibodies may be prepared by any method known in the art such as the hybridoma technique (Kohler & Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, 1983, Immunology Today, 4:72) and the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, pp77-96, Alan R Liss, Inc., 1985).

20 Antibodies for use in the invention may also be generated using single lymphocyte antibody methods by cloning and expressing immunoglobulin variable region cDNAs generated from single lymphocytes selected for the production of specific antibodies by for example the methods described by Babcook, J. *et al.*, 1996, Proc. Natl. Acad. Sci. USA 93(15):7843-78481; WO92/02551; WO2004/051268 and International Patent Application  
25 number WO2004/106377.

Humanized antibodies are antibody molecules from non-human species having one or more complementarity determining regions (CDRs) from the non-human species and a framework region from a human immunoglobulin molecule (see, *e.g.* US 5,585,089; WO91/09967).

30 Chimeric antibodies are those antibodies encoded by immunoglobulin genes that have been genetically engineered so that the light and heavy chain genes are composed of immunoglobulin gene segments belonging to different species. These chimeric antibodies are

likely to be less antigenic. Bivalent antibodies may be made by methods known in the art (Milstein *et al.*, 1983, Nature 305:537-539; WO 93/08829, Traunecker *et al.*, 1991, EMBO J. 10:3655-3659). Multi-valent antibodies may comprise multiple specificities or may be monospecific (see for example WO 92/22853 and WO05/113605).

5 The antibodies for use in the present invention can also be generated using various phage display methods known in the art and include those disclosed by Brinkman *et al.* (in J. Immunol. Methods, 1995, 182: 41-50), Ames *et al.* (J. Immunol. Methods, 1995, 184:177-186), Kettleborough *et al.* (Eur. J. Immunol. 1994, 24:952-958), Persic *et al.* (Gene, 1997 187 9-18), Burton *et al.* (Advances in Immunology, 1994, 57:191-280) and WO 90/02809; WO 10 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and US 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108. Techniques for the production of single chain antibodies, such as those described in US 4,946,778 can also be adapted to produce single chain antibodies which bind to IL-17A and IL-17F. Also, 15 transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The residues in antibody variable domains are conventionally numbered according to a system devised by Kabat *et al.* This system is set forth in Kabat *et al.*, 1987, in Sequences of Proteins of Immunological Interest, US Department of Health and Human Services, NIH, 20 USA (hereafter "Kabat *et al. (supra)*"). This numbering system is used in the present specification except where otherwise indicated.

The Kabat residue designations do not always correspond directly with the linear numbering of the amino acid residues. The actual linear amino acid sequence may contain fewer or additional amino acids than in the strict Kabat numbering corresponding to a 25 shortening of, or insertion into, a structural component, whether framework or complementarity determining region (CDR), of the basic variable domain structure. The correct Kabat numbering of residues may be determined for a given antibody by alignment of residues of homology in the sequence of the antibody with a "standard" Kabat numbered sequence.

30 The CDRs of the heavy chain variable domain are located at residues 31-35 (CDR-H1), residues 50-65 (CDR-H2) and residues 95-102 (CDR-H3) according to the Kabat numbering system. However, according to Chothia (Chothia, C. and Lesk, A.M. J. Mol.

Biol., 196, 901-917 (1987)), the loop equivalent to CDR-H1 extends from residue 26 to residue 32. Thus 'CDR-H1', as used herein, comprises residues 26 to 35, as described by a combination of the Kabat numbering system and Chothia's topological loop definition.

The CDRs of the light chain variable domain are located at residues 24-34 (CDR-L1),  
5 residues 50-56 (CDR-L2) and residues 89-97 (CDR-L3) according to the Kabat numbering system.

In one embodiment the present invention provides a neutralising antibody having specificity for human IL-17A and human IL-17F, comprising a heavy chain, wherein the variable domain of the heavy chain comprises at least one of a CDR having the sequence  
10 given in SEQ ID NO:1 for CDR-H1, a CDR having the sequence given in SEQ ID NO:2 for CDR-H2 and a CDR having the sequence given in SEQ ID NO:3 for CDR-H3.

In another embodiment the present invention provides a neutralising antibody having specificity for human IL-17A and human IL-17F, comprising a heavy chain, wherein at least two of CDR-H1, CDR-H2 and CDR-H3 of the variable domain of the heavy chain are  
15 selected from the following: the sequence given in SEQ ID NO:1 for CDR-H1, the sequence given in SEQ ID NO:2 for CDR-H2 and the sequence given in SEQ ID NO:3 for CDR-H3. For example, the antibody may comprise a heavy chain wherein CDR-H1 has the sequence given in SEQ ID NO:1 and CDR-H2 has the sequence given in SEQ ID NO:2. Alternatively, the antibody may comprise a heavy chain wherein CDR-H1 has the sequence given in SEQ  
20 ID NO:1 and CDR-H3 has the sequence given in SEQ ID NO:3, or the antibody may comprise a heavy chain wherein CDR-H2 has the sequence given in SEQ ID NO:2 and CDR-H3 has the sequence given in SEQ ID NO:3. For the avoidance of doubt, it is understood that all permutations are included.

In another embodiment the present invention provides a neutralising antibody having  
25 specificity for human IL-17A and human IL-17F, comprising a heavy chain, wherein the variable domain of the heavy chain comprises the sequence given in SEQ ID NO:1 for CDR-H1, the sequence given in SEQ ID NO:2 for CDR-H2 and the sequence given in SEQ ID NO:3 for CDR-H3.

In one embodiment the present invention provides a neutralising antibody having  
30 specificity for human IL-17A and human IL-17F, comprising a light chain, wherein the variable domain of the light chain comprises at least one of a CDR having the sequence given

in SEQ ID NO:4 for CDR-L1, a CDR having the sequence given in SEQ ID NO:5 for CDR-L2 and a CDR having the sequence given in SEQ ID NO:6 for CDR-L3.

5 In another embodiment the present invention provides a neutralising antibody which specifically binds human IL-17A and human IL-17F, comprising a heavy chain and a light chain wherein the heavy chain variable domain comprises the sequence given in SEQ ID NO:9 or a sequence at least 95% identical thereto and the light chain variable domain comprises the sequence given in SEQ ID NO:7 or a sequence at least 95% identical thereto.

10 In another embodiment the present invention provides a neutralising antibody which specifically binds human IL-17A and human IL-17F, comprising a heavy chain and a light chain wherein the heavy chain variable domain comprises the sequence given in SEQ ID NO:9 or a sequence at least 95% identical thereto and the light chain variable domain comprises the sequence given in SEQ ID NO:7 or a sequence at least 95% identical thereto, wherein the heavy chain comprises the CDRH1, CDRH2 and CDRH3 identified in SEQ ID NO: 1, 2 and 3 respectively and wherein the antibody has an affinity for IL-17A of better than 20pM and an affinity for IL-17F of better than 2nM.

15 In another embodiment the present invention provides an isolated DNA sequence encoding the heavy and/or light chain(s) of an antibody according to the neutralising antibody as defined herein.

20 In another embodiment the present invention provides an isolated DNA molecule encoding the heavy and/or light chain(s) of the neutralising antibody as defined herein.

In another embodiment the present invention provides an isolated DNA molecule encoding the heavy and the light chain(s) of the neutralising antibody as defined herein.

In another embodiment the present invention provides a cloning or expression vector comprising one or more DNA sequences as defined herein.

25 In another embodiment the present invention provides a cloning or expression vector comprising one or more DNA molecules as defined herein.

In another embodiment the present invention provides a cloning or expression vector comprising the isolated DNA molecule as defined herein.

7a

In another embodiment the present invention provides a host cell comprising one or more cloning or expression vectors as defined herein.

5 In another embodiment the present invention provides a process for the production of the antibody as defined herein, comprising culturing the host cell and isolating the antibody.

In another embodiment the present invention provides a pharmaceutical composition comprising the antibody defined herein, in combination with one or more of a pharmaceutically acceptable excipient, diluent or carrier.

10 In another embodiment the present invention provides an antibody or the pharmaceutical composition both defined herein, for use in the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder  
15 disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma , graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or  
20 hypochlorhydria.

In another embodiment the present invention provides a use of the antibody according to the invention or the pharmaceutical composition according to the invention, for the treatment or prophylaxis of a pathological disorder being a viral infection, a bacterial  
25 infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and

7b

peripheral nervous system, an autoimmune disorder, pancreatitis, trauma , graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydia.

5 In another embodiment the present invention provides a use of the antibody or the pharmaceutical composition both defined herein, for the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder  
10 disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma , graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or  
15 hypochlorhydia.

In another embodiment the present invention provides a use of the antibody or the pharmaceutical composition both defined herein, for the preparation of a medicament for the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an  
20 endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune  
25 disorder, pancreatitis, trauma , graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydia.

In another embodiment the present invention provides a use of the antibody according to the invention or the pharmaceutical composition according to the invention, for  
30 the preparation of a medicament for the treatment or prophylaxis of a pathological disorder

7c

being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical  
5 adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma , graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydria.

10 In another embodiment the present invention provides a neutralising antibody having specificity for human IL-17A and human IL-17F, comprising a light chain, wherein at least two of CDR-L1, CDR-L2 and CDR-L3 of the variable domain of the light chain are selected from the following: the sequence given in SEQ ID NO:4 for CDR-L1, the sequence given in  
15 SEQ ID NO:5 for CDR-L2 and the sequence given in SEQ ID NO:6 for CDR-L3. For example, the antibody may comprise a light chain wherein CDR-L1 has the sequence given in SEQ ID NO:4 and CDR-L2 has the sequence given in SEQ ID NO:5. Alternatively, the antibody may comprise a light chain wherein CDR-L1 has the sequence given in SEQ ID NO:4 and CDR-L3 has the sequence given in SEQ ID NO:6, or the antibody may comprise a light chain wherein CDR-L2 has the sequence given in SEQ ID NO:5 and CDR-L3 has the  
20 sequence given in SEQ ID NO:6. For the avoidance of doubt, it is understood that all permutations are included.

In another embodiment the present invention provides a neutralising antibody having specificity for human IL-17A and human IL-17F, comprising a light chain, wherein the variable domain comprises the sequence given in SEQ ID NO:4 for CDR-L1, the sequence  
25 given in SEQ ID NO:5 for CDR-L2 and the sequence given in SEQ ID NO:6 for CDR-L3.

The antibody molecules of the present invention preferably comprise a complementary light chain or a complementary heavy chain, respectively.


Hence in one embodiment, an antibody according to the present invention comprises a heavy chain, wherein the variable domain of the heavy chain comprises the sequence given  
30 in SEQ ID NO:1 for CDR-H1, the sequence given in SEQ ID NO:2 for CDR-H2 and the

7d

sequence given in SEQ ID NO:3 for CDR-H3 and a light chain wherein the variable domain of the light chain comprises the sequence given in SEQ ID NO:4 for CDR-L1, the sequence given in SEQ ID NO:5 for CDR-L2 and the sequence given in SEQ ID NO:6 for CDR-L3.

5 It will be appreciated that one or more amino acid substitutions may be made to the CDRs provided by the present invention without significantly altering the ability of the antibody to bind to IL-17A and IL-17F and to neutralise IL-17A and IL-17F activity. The effect of any amino acid substitutions on binding and neutralisation can be readily tested by one skilled in the art, for example by using the methods described herein. Accordingly, the

---



present invention provides an antibody comprising one or more CDRs selected from CDRH-1 (SEQ ID NO:1), CDRH-2 (SEQ ID NO:2), CDRH-3 (SEQ ID NO:3), CDRL-1 (SEQ ID NO:4), CDRL-2 (SEQ ID NO:5) and CDRL-3 (SEQ ID NO:6) in which one or more amino acids in one or more of the CDRs has been substituted with another amino acid. It will also  
5 be appreciated that the length of one or more of the CDRs may be altered without significantly altering the ability of the antibody to bind to IL-17A and IL-17F and to neutralise IL-17A and IL-17F activity.

In one embodiment, an antibody of the present invention comprises a heavy chain, wherein the variable domain of the heavy chain comprises three CDRs wherein the sequence  
10 of CDRH-1 has at least 60% identity or similarity to the sequence given in SEQ ID NO:1, CDRH-2 has at least 60% identity or similarity to the sequence given in SEQ ID NO:2 and/or CDRH-3 has at least 60% identity or similarity to the sequence given in SEQ ID NO:3. In another embodiment, an antibody of the present invention comprises a heavy chain, wherein the variable domain of the heavy chain comprises three CDRs wherein the sequence of  
15 CDRH-1 has at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:1, CDRH-2 has at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:2 and/or CDRH-3 has at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:3.

"Identity", as used herein, indicates that at any particular position in the aligned  
20 sequences, the amino acid residue is identical between the sequences. "Similarity", as used herein, indicates that, at any particular position in the aligned sequences, the amino acid residue is of a similar type between the sequences. For example, leucine may be substituted for isoleucine or valine. Other amino acids which can often be substituted for one another include but are not limited to:

- 25 - phenylalanine, tyrosine and tryptophan (amino acids having aromatic side chains);
- lysine, arginine and histidine (amino acids having basic side chains);
- aspartate and glutamate (amino acids having acidic side chains);
- asparagine and glutamine (amino acids having amide side chains); and
- cysteine and methionine (amino acids having sulphur-containing side chains). Degrees  
30 of identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing. Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of

Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

- 5 In another embodiment, an antibody of the present invention comprises a light chain, wherein the variable domain of the light chain comprises three CDRs wherein the sequence of CDRL-1 has at least 60% identity or similarity to the sequence given in SEQ ID NO:4, CDRL-2 has at least 60% identity or similarity to the sequence given in SEQ ID NO:5 and/or CDRL-3 has at least 60% identity or similarity to the sequence given in SEQ ID NO:6. In
- 10 another embodiment, an antibody of the present invention comprises a light chain, wherein the variable domain of the heavy chain comprises three CDRs wherein the sequence of CDRL-1 has at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:4, CDRL-2 has at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:5 and/or CDRL-3 has at least 70%, 80%, 90%, 95% or
- 15 98% identity or similarity to the sequence given in SEQ ID NO:6.

In one embodiment the antibody provided by the present invention is a monoclonal antibody.

In one embodiment the antibody provided by the present invention is a chimeric antibody.

- 20 In one embodiment the antibody provided by the present invention is a CDR-grafted antibody molecule comprising one or more of the CDRs provided in SEQ ID NOS:1 to 6. As used herein, the term 'CDR-grafted antibody molecule' refers to an antibody molecule wherein the heavy and/or light chain contains one or more CDRs (including, if desired, one or more modified CDRs) from a donor antibody (e.g. a murine monoclonal antibody) grafted
- 25 into a heavy and/or light chain variable region framework of an acceptor antibody (e.g. a human antibody). For a review, see Vaughan *et al*, Nature Biotechnology, 16, 535-539, 1998. In one embodiment rather than the entire CDR being transferred, only one or more of the specificity determining residues from any one of the CDRs described herein above are transferred to the human antibody framework (see for example, Kashmiri *et al.*, 2005,
- 30 Methods, 36, 25-34). In one embodiment only the specificity determining residues from one or more of the CDRs described herein above are transferred to the human antibody

framework. In another embodiment only the specificity determining residues from each of the CDRs described herein above are transferred to the human antibody framework.

When the CDRs or specificity determining residues are grafted, any appropriate acceptor variable region framework sequence may be used having regard to the class/type of the donor antibody from which the CDRs are derived, including mouse, primate and human framework regions. Preferably, the CDR-grafted antibody according to the present invention has a variable domain comprising human acceptor framework regions as well as one or more of the CDRs or specificity determining residues described above. Thus, provided in one embodiment is a neutralising CDR-grafted antibody wherein the variable domain comprises human acceptor framework regions and non-human donor CDRs.

Examples of human frameworks which can be used in the present invention are KOL, NEWM, REI, EU, TUR, TEL, LAY and POM (Kabat *et al.*, *supra*). For example, KOL and NEWM can be used for the heavy chain, REI can be used for the light chain and EU, LAY and POM can be used for both the heavy chain and the light chain. Alternatively, human germline sequences may be used; these are available at: <http://vbase.mrc-cpe.cam.ac.uk/>

In a CDR-grafted antibody of the present invention, the acceptor heavy and light chains do not necessarily need to be derived from the same antibody and may, if desired, comprise composite chains having framework regions derived from different chains.

The preferred framework region for the heavy chain of the CDR-grafted antibody of the present invention is derived from the human sub-group VH3 sequence 1-3 3-07 together with JH4. Accordingly, provided is a neutralising CDR-grafted antibody comprising at least one non-human donor CDR wherein the heavy chain framework region is derived from the human subgroup sequence 1-3 3-07 together with JH4. The sequence of human JH4 is as follows: (YFDY)WGQGTLVTVSS . The YFDY motif is part of CDR-H3 and is not part of framework 4 (Ravetch, J.V. *et al.*, 1981, *Cell*, 27, 583-591).

The preferred framework region for the light chain of the CDR-grafted antibody of the present invention is derived from the human germline sub-group VK1 sequence 2-1-(1) L4 together with JK1. Accordingly, provided is a neutralising CDR-grafted antibody comprising at least one non-human donor CDR wherein the light chain framework region is derived from the human subgroup sequence VK1 2-1-(1) L4 together with JK1. The JK1 sequence is as

follows: (WT)FGQGTKVEIK. The WT motif is part of CDR-L3 and is not part of framework 4 (Hieter, P.A., *et al.*, 1982, *J. Biol. Chem.*, 257, 1516-1522).

Also, in a CDR-grafted antibody of the present invention, the framework regions need not have exactly the same sequence as those of the acceptor antibody. For instance, unusual  
5 residues may be changed to more frequently-occurring residues for that acceptor chain class or type. Alternatively, selected residues in the acceptor framework regions may be changed so that they correspond to the residue found at the same position in the donor antibody (see Reichmann *et al.*, 1998, *Nature*, 332, 323-324). Such changes should be kept to the minimum necessary to recover the affinity of the donor antibody. A protocol for selecting residues in  
10 the acceptor framework regions which may need to be changed is set forth in WO 91/09967.

Preferably, in a CDR-grafted antibody molecule of the present invention, if the acceptor heavy chain has the human VH3 sequence 1-3 3-07 together with JH4, then the acceptor framework regions of the heavy chain comprise, in addition to one or more donor CDRs, a donor residue at at least position 94 (according to Kabat *et al.*, (supra)). Accordingly,  
15 provided is a CDR-grafted antibody, wherein at least the residue at position 94 of the variable domain of the heavy chain is a donor residue.

Preferably, in a CDR-grafted antibody molecule according to the present invention, if the acceptor light chain has the human sub-group VK1 sequence 2-1-(1) L4 together with JK1, then no donor residues are transferred i.e. only the CDRs are transferred. Accordingly,  
20 provided is a CDR-grafted antibody wherein only the CDRs are transferred to the donor framework.

Donor residues are residues from the donor antibody, i.e. the antibody from which the CDRs were originally derived.

In one embodiment, an antibody of the present invention comprises a heavy chain,  
25 wherein the variable domain of the heavy chain comprises the sequence given in SEQ ID NO:9 (gH9).

In another embodiment, an antibody of the present invention comprises a heavy chain, wherein the variable domain of the heavy chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:9. In one embodiment, an antibody  
30 of the present invention comprises a heavy chain, wherein the variable domain of the heavy

chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:9.

In one embodiment, an antibody of the present invention comprises a light chain, wherein the variable domain of the light chain comprises the sequence given in SEQ ID NO:7  
5 (gL7).

In another embodiment, an antibody of the present invention comprises a light chain, wherein the variable domain of the light chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:7. In one embodiment the antibody of the present invention comprises a light chain, wherein the variable domain of the  
10 light chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:7.

In one embodiment an antibody of the present invention comprises a heavy chain, wherein the variable domain of the heavy chain comprises the sequence given in SEQ ID NO:9 and a light chain, wherein the variable domain of the light chain comprises the sequence  
15 given in SEQ ID NO:7.

In another embodiment of the invention, the antibody comprises a heavy chain and a light chain, wherein the variable domain of the heavy chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:9 and the variable domain of the light chain comprises a sequence having at least 60% identity or similarity to  
20 the sequence given in SEQ ID NO:7. Preferably, the antibody comprises a heavy chain, wherein the variable domain of the heavy chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:9 and a light chain, wherein the variable domain of the light chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID  
25 NO:7.

As described herein above, the antibody molecule of the present invention may comprise a complete antibody molecule having full length heavy and light chains or a fragment thereof, such as a domain antibody e.g. VH, VL, VHH, Fab, modified Fab, Fab', F(ab')<sub>2</sub>, Fv or scFv fragment.

30 The constant region domains of the antibody molecule of the present invention, if present, may be selected having regard to the proposed function of the antibody molecule, and

in particular the effector functions which may be required. For example, the constant region domains may be human IgA, IgD, IgE, IgG or IgM domains. In particular, human IgG constant region domains may be used, especially of the IgG1 and IgG3 isotypes when the antibody molecule is intended for therapeutic uses and antibody effector functions are  
5 required. Alternatively, IgG2 and IgG4 isotypes may be used when the antibody molecule is intended for therapeutic purposes and antibody effector functions are not required, e.g. for simply blocking IL-17 activity. For example IgG4 molecules in which the serine at position 241 has been changed to proline as described in Angal *et al.*, Molecular Immunology, 1993, 30 (1), 105-108 may be used. Particularly preferred is the IgG4 constant domain comprising  
10 this change.

In one embodiment the antibody heavy chain comprises a CH1 domain and the antibody light chain comprises a CL domain, either kappa or lambda.

In a preferred embodiment the antibody provided by the present invention is a neutralising antibody having specificity for human IL-17A and human IL-17F in which the  
15 heavy chain constant region comprises the human IgG4 constant region in which the serine at position 241 has been substituted by proline as described in Angal *et al.*, *supra*. Accordingly, the present invention provides an antibody in which the heavy chain comprises or consists of the sequence given in SEQ ID NO:15.

In one embodiment of the invention, the antibody comprises a heavy chain, wherein  
20 the heavy chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:15. Preferably, the antibody comprises a heavy chain, wherein the heavy chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:15.

In one embodiment an antibody molecule according to the present invention  
25 comprises a light chain comprising the sequence given in SEQ ID NO:11.

In one embodiment of the invention, the antibody comprises a light chain, wherein the light chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:11. Preferably, the antibody comprises a light chain, wherein the light chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or  
30 similarity to the sequence given in SEQ ID NO:11.

In one embodiment the present invention provides an antibody in which the heavy chain comprises or consists of the sequence given in SEQ ID NO:15 and the light chain comprises or consists of the sequence given in SEQ ID NO:11.

In one embodiment of the invention, the antibody comprises a heavy chain and a light  
5 chain, wherein the heavy chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:15 and the light chain comprises a sequence having at least 60% identity or similarity to the sequence given in SEQ ID NO:11. Preferably, the antibody comprises a heavy chain, wherein the heavy chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence  
10 given in SEQ ID NO:15 and a light chain, wherein the light chain comprises a sequence having at least 70%, 80%, 90%, 95% or 98% identity or similarity to the sequence given in SEQ ID NO:11.

Also provided by the present invention is a specific region or epitope of human IL-17A and/or a specific region or epitope of human IL-17F and/or a specific region or epitope of human  
15 IL-17A/F heterodimer which is bound by an antibody provided by the present invention, in particular an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and/or the light chain sequence gL7 (SEQ ID NO:7).

The specific region or epitope of the human IL-17A polypeptide and the specific region or epitope of the human IL-17F polypeptide and the specific region or epitope of the human IL-  
20 17A/F heterodimer can be identified by any suitable epitope mapping method known in the art in combination with any one of the antibodies provided by the present invention. Examples of such methods include screening peptides of varying lengths derived from IL-17A and IL-17F for binding to the antibody of the present invention with the smallest fragment that can specifically bind to the antibody containing the sequence of the epitope recognised by the antibody. The IL-  
25 17 peptides may be produced synthetically or by proteolytic digestion of the appropriate IL-17 polypeptide. Peptides that bind the antibody can be identified by, for example, mass spectrometric analysis. In another example, NMR spectroscopy can be used to identify the epitope bound by an antibody of the present invention. Once identified, the epitopic fragment which binds an antibody of the present invention can be used, if required, as an immunogen to  
30 obtain additional neutralising antibodies which bind the same epitope.

Antibodies which cross-block the binding of an antibody according to the present invention, in particular, an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7), may be similarly useful in neutralising IL-17A and IL-17F activity. Accordingly, the present invention also provides a neutralising  
5 antibody which binds human IL-17A and human IL-17F, which cross-blocks the binding of any one of the antibodies described above to human IL-17A and/or human IL-17F and/or human IL-17A/F heterodimer and/or is cross-blocked from binding IL-17A and/or IL-17F and/or human IL-17A/F heterodimer by any one of those antibodies. In one embodiment, such an antibody binds to the same epitope as an antibody described herein above. In another  
10 embodiment the cross-blocking neutralising antibody binds to an epitope which borders and/or overlaps with the epitope bound by an antibody described herein above. In another embodiment the cross-blocking neutralising antibody of this aspect of the invention does not bind to the same epitope as an antibody of the present invention or an epitope that borders and/or overlaps with said epitope.

15 Cross-blocking antibodies can be identified using any suitable method in the art, for example by using competition ELISA or BIAcore where binding of the cross blocking antibody to human IL-17A and/or human IL-17F prevents the binding of an antibody of the present invention or vice versa.

In one embodiment there is provided a neutralising antibody which binds to human  
20 IL-17A and human IL-17F, which cross-blocks the binding of an antibody whose heavy chain comprises the sequence gH9 (SEQ ID NO:9) and whose light chain comprises the sequence gL7 (SEQ ID NO:7) to human IL-17A and to human IL-17F. In one embodiment the cross-blocking antibodies provided by the present invention inhibit the binding of an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7  
25 (SEQ ID NO:7) to IL-17A by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95% and to IL-17F by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95%.

In one embodiment there is provided a neutralising antibody which binds to human  
30 IL-17A and human IL-17F, which cross-blocks the binding of an antibody whose heavy chain comprises the sequence gH9 (SEQ ID NO:9) and whose light chain comprises the sequence

gL7 (SEQ ID NO:7) to human IL-17A and to human IL-17F and to human IL-17A/F heterodimer. In one embodiment the cross-blocking antibodies provided by the present invention inhibit the binding of an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7) to IL-17A by greater than 80%,  
5 preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95% and to IL-17F by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95% and to IL-17A/F heterodimer to IL-17F by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95%.

10 In one embodiment there is provided a neutralising antibody which binds to human IL-17A and human IL-17F, which cross-blocks the binding of an antibody whose heavy chain comprises the sequence gH9 (SEQ ID NO:9) and whose light chain comprises the sequence gL7 (SEQ ID NO:7) to human IL-17A or to human IL-17F or human IL-17A/F heterodimer. In one embodiment the cross-blocking antibodies provided by the present invention inhibit the  
15 binding of an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7) to IL-17A or IL-17F or IL-17A/F by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95%.

Alternatively or in addition, neutralising antibodies according to this aspect of the  
20 invention may be cross-blocked from binding to human IL-17A and human IL-17F by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7). Also provided therefore is a neutralising antibody molecule which binds to human IL-17A and to human IL-17F which is cross-blocked from binding human IL-17A and human IL-17F by an antibody comprising the heavy chain sequence gH9  
25 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7). In one embodiment the neutralising antibodies provided by this aspect of the invention are inhibited from binding to human IL-17A and human IL-17F by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7) by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably  
30 by greater than 95%.

In another embodiment there is provided a neutralising antibody molecule which binds to human IL-17A and to human IL-17F which is cross-blocked from binding human IL-17A and human IL-17F and IL-17A/F heterodimer by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7). In one  
5 embodiment the neutralising antibodies provided by this aspect of the invention are inhibited from binding to human IL-17A and human IL-17F and human IL-17A/F heterodimer by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7) by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95%.

10 Also provided therefore is a neutralising antibody molecule which binds to human IL-17A and to human IL-17F which is cross-blocked from binding human IL-17A or human IL-17F or human IL-17A/F by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7). In one embodiment the neutralising antibodies provided by this aspect of the invention are inhibited from binding to human IL-  
15 17A or human IL-17F or human IL-17A/F by an antibody comprising the heavy chain sequence gH9 (SEQ ID NO:9) and the light chain sequence gL7 (SEQ ID NO:7) by greater than 80%, preferably by greater than 85%, more preferably by greater than 90%, even more preferably by greater than 95%.

The antibody molecule of any aspect of the present invention preferably has a high  
20 binding affinity, preferably nanomolar, even more preferably picomolar. It will be appreciated that the binding affinity of an antibody according to the present invention for human IL-17A may be different from the binding affinity of the same antibody for human IL-17F and/or the IL-17A/F heterodimer. In one example the antibody molecule of the present invention has an affinity for IL-17A that is greater than its affinity for IL-17F. In one  
25 example the antibody molecule of the present invention has an affinity for IL-17A which is at least 10 fold greater than its binding affinity for IL-17F. In one example the antibody molecule of the present invention has an affinity for IL-17A which is at least 50 fold greater than its binding affinity for IL-17F. In one example the antibody molecule of the present invention has an affinity for IL-17A which is at least 100 fold greater than its binding affinity  
30 for IL-17F. In one example the antibody molecule of the present invention has an affinity for IL-17F that is greater than its affinity for IL-17A. In one example the antibody molecule of the present invention has an affinity for IL-17A that is the same as its affinity for IL-17F. In

one example the antibody molecule of the present invention has a picomolar affinity for IL-17A and a nanomolar affinity for IL-17F. In one example the antibody molecule of the present invention has a nanomolar affinity for IL-17F and a picomolar affinity for IL-17A. In one example the antibody molecule of the present invention has a nanomolar affinity for both  
5 IL-17A and IL-17F. In one example the antibody molecule of the present invention has a picomolar affinity for both IL-17A and IL-17F.

Preferably the antibody molecule of the present invention has a binding affinity for IL-17A of better than 10nM. In one embodiment the antibody molecule of the present invention has a binding affinity for IL-17A of better than 500 pM. In one embodiment the antibody  
10 molecule of the present invention has a binding affinity for IL-17A of better than 100 pM. In one embodiment the antibody molecule of the present invention has a binding affinity for IL-17A of better than 20pM. In one embodiment the antibody of the present invention has an affinity for IL-17A of 16pM.

Preferably the antibody molecule of the present invention has a binding affinity for IL-  
15 17F of better than 10nM. In one embodiment the antibody of the present invention has an affinity for IL-17F of better than 2 nM. In one embodiment the antibody of the present invention has an affinity for IL-17F of 1.75 nM.

Preferably the antibody molecule of the present invention has a binding affinity for IL-17A/F heterodimer of better than 10nM. In one embodiment the antibody molecule of the present invention has a binding affinity for IL-17A/F heterodimer of better than 500 pM. In  
20 one embodiment the antibody molecule of the present invention has a binding affinity for IL-17A/F heterodimer of better than 150 pM. In one embodiment the antibody molecule of the present invention has a binding affinity for IL-17A/F heterodimer of 116pM.

In one embodiment the antibody molecule of the present invention has a binding  
25 affinity for cynomolgus IL-17F of better than 2nM. In one embodiment the antibody molecule of the present invention has a binding affinity for cynomolgus IL-17F of 1.03nM.

It will be appreciated that the affinity of antibodies provided by the present invention may be altered using any suitable method known in the art. The present invention therefore also relates to variants of the antibody molecules of the present invention, which have an  
30 improved affinity for IL-17A and/or IL-17F. Such variants can be obtained by a number of affinity maturation protocols including mutating the CDRs (Yang *et al.*, *J. Mol. Biol.*, 254,

392-403, 1995), chain shuffling (Marks *et al.*, *Bio/Technology*, 10, 779-783, 1992), use of mutator strains of *E. coli* (Low *et al.*, *J. Mol. Biol.*, 250, 359-368, 1996), DNA shuffling (Patten *et al.*, *Curr. Opin. Biotechnol.*, 8, 724-733, 1997), phage display (Thompson *et al.*, *J. Mol. Biol.*, 256, 77-88, 1996) and sexual PCR (Cramer *et al.*, *Nature*, 391, 288-291, 1998).  
5 Vaughan *et al.* (*supra*) discusses these methods of affinity maturation.

In one embodiment the antibody molecules of the present invention neutralise IL-17A and IL-17F activity, for example in the *in vitro* assays described in the Examples. In one embodiment the present invention provides a neutralising antibody having specificity for human IL-17A and IL-17F which is capable of inhibiting the activity of 0.8nM human IL-  
10 17A by 50% at a concentration of less than 5nM and the activity of 4.2nM IL-17F by 50% at a concentration of less than 12nM said inhibitory activity being measured on the IL-17A or IL-17F induced release of IL-6 from Hela cells. In one embodiment the concentration of antibody which inhibits IL-17A by 50% is less than 3nM. In one embodiment the concentration of antibody which inhibits IL-17F by 50% is less than 11nM. In one  
15 embodiment the human IL-17A and human IL-17F used in the assay are recombinant human IL-17A and IL-17F. In one embodiment the neutralising antibody is a humanised or fully human antibody.

If desired an antibody for use in the present invention may be conjugated to one or more effector molecule(s). It will be appreciated that the effector molecule may comprise a single  
20 effector molecule or two or more such molecules so linked as to form a single moiety that can be attached to the antibodies of the present invention. Where it is desired to obtain an antibody fragment linked to an effector molecule, this may be prepared by standard chemical or recombinant DNA procedures in which the antibody fragment is linked either directly or via a coupling agent to the effector molecule. Techniques for conjugating such effector  
25 molecules to antibodies are well known in the art (see, Hellstrom *et al.*, *Controlled Drug Delivery*, 2nd Ed., Robinson *et al.*, eds., 1987, pp. 623-53; Thorpe *et al.*, 1982, *Immunol. Rev.*, 62:119-58 and Dubowchik *et al.*, 1999, *Pharmacology and Therapeutics*, 83, 67-123). Particular chemical procedures include, for example, those described in WO 93/06231, WO 92/22583, WO 89/00195, WO 89/01476 and WO03031581. Alternatively, where the effector  
30 molecule is a protein or polypeptide the linkage may be achieved using recombinant DNA procedures, for example as described in WO 86/01533 and EP0392745.

The term effector molecule as used herein includes, for example, antineoplastic agents, drugs, toxins, biologically active proteins, for example enzymes, other antibody or antibody fragments, synthetic or naturally occurring polymers, nucleic acids and fragments thereof e.g. DNA, RNA and fragments thereof, radionuclides, particularly radioiodide, radioisotopes, 5 chelated metals, nanoparticles and reporter groups such as fluorescent compounds or compounds which may be detected by NMR or ESR spectroscopy.

Examples of effector molecules may include cytotoxins or cytotoxic agents including any agent that is detrimental to (*e.g.* kills) cells. Examples include combrestatins, dolastatins, epothilones, staurosporin, maytansinoids, spongistatins, rhizoxin, halichondrins, roridins, 10 hemiasterlins, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof.

15 Effector molecules also include, but are not limited to, antimetabolites (*e.g.* methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.* mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), 20 anthracyclines (*e.g.* daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.* dactinomycin (formerly actinomycin), bleomycin, mithramycin, anthramycin (AMC), calicheamicins or duocarmycins), and anti-mitotic agents (*e.g.* vincristine and vinblastine).

Other effector molecules may include chelated radionuclides such as  $^{111}\text{In}$  and  $^{90}\text{Y}$ ,  $\text{Lu}^{177}$ , Bismuth $^{213}$ , Californium $^{252}$ , Iridium $^{192}$  and Tungsten $^{188}$ /Rhenium $^{188}$ ; or drugs such as 25 but not limited to, alkylphosphocholines, topoisomerase I inhibitors, taxoids and suramin.

Other effector molecules include proteins, peptides and enzymes. Enzymes of interest include, but are not limited to, proteolytic enzymes, hydrolases, lyases, isomerases, transferases. Proteins, polypeptides and peptides of interest include, but are not limited to, immunoglobulins, toxins such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin, a 30 protein such as insulin, tumour necrosis factor,  $\alpha$ -interferon,  $\beta$ -interferon, nerve growth factor, platelet derived growth factor or tissue plasminogen activator, a thrombotic agent or an anti-angiogenic agent, *e.g.* angiostatin or endostatin, or, a biological response modifier such

as a lymphokine, interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), nerve growth factor (NGF) or other growth factor and immunoglobulins.

Other effector molecules may include detectable substances useful for example in  
5 diagnosis. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive nuclides, positron emitting metals (for use in positron emission tomography), and nonradioactive paramagnetic metal ions. See generally U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics. Suitable enzymes include horseradish  
10 peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; suitable prosthetic groups include streptavidin, avidin and biotin; suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride and phycoerythrin; suitable luminescent materials include luminol; suitable bioluminescent materials include luciferase, luciferin, and  
15 aequorin; and suitable radioactive nuclides include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{111}\text{In}$  and  $^{99}\text{Tc}$ .

In another example the effector molecule may increase the half-life of the antibody *in vivo*, and/or reduce immunogenicity of the antibody and/or enhance the delivery of an antibody across an epithelial barrier to the immune system. Examples of suitable effector molecules of this type include polymers, albumin, albumin binding proteins or albumin  
20 binding compounds such as those described in WO05/117984.

Where the effector molecule is a polymer it may, in general, be a synthetic or a naturally occurring polymer, for example an optionally substituted straight or branched chain polyalkylene, polyalkenylene or polyoxyalkylene polymer or a branched or unbranched polysaccharide, e.g. a homo- or hetero- polysaccharide.

25 Particular optional substituents which may be present on the above-mentioned synthetic polymers include one or more hydroxy, methyl or methoxy groups.

Particular examples of synthetic polymers include optionally substituted straight or branched chain poly(ethyleneglycol), poly(propyleneglycol) poly(vinylalcohol) or derivatives thereof, especially optionally substituted poly(ethyleneglycol) such as  
30 methoxypoly(ethyleneglycol) or derivatives thereof.

Particular naturally occurring polymers include lactose, amylose, dextran, glycogen or derivatives thereof.

“Derivatives” as used herein is intended to include reactive derivatives, for example thiol-selective reactive groups such as maleimides and the like. The reactive group may be  
5 linked directly or through a linker segment to the polymer. It will be appreciated that the residue of such a group will in some instances form part of the product as the linking group between the antibody fragment and the polymer.

The size of the polymer may be varied as desired, but will generally be in an average molecular weight range from 500Da to 50000Da, preferably from 5000 to 40000Da and more  
10 preferably from 20000 to 40000Da. The polymer size may in particular be selected on the basis of the intended use of the product for example ability to localize to certain tissues such as tumors or extend circulating half-life (for review see Chapman, 2002, Advanced Drug Delivery Reviews, 54, 531-545). Thus, for example, where the product is intended to leave the circulation and penetrate tissue, for example for use in the treatment of a tumour, it may  
15 be advantageous to use a small molecular weight polymer, for example with a molecular weight of around 5000Da. For applications where the product remains in the circulation, it may be advantageous to use a higher molecular weight polymer, for example having a molecular weight in the range from 20000Da to 40000Da.

Particularly preferred polymers include a polyalkylene polymer, such as a  
20 poly(ethyleneglycol) or, especially, a methoxypoly(ethyleneglycol) or a derivative thereof, and especially with a molecular weight in the range from about 15000Da to about 40000Da.

In one example antibodies for use in the present invention are attached to poly(ethyleneglycol) (PEG) moieties. In one particular example the antibody is an antibody fragment and the PEG molecules may be attached through any available amino acid side-  
25 chain or terminal amino acid functional group located in the antibody fragment, for example any free amino, imino, thiol, hydroxyl or carboxyl group. Such amino acids may occur naturally in the antibody fragment or may be engineered into the fragment using recombinant DNA methods (see for example US 5,219,996; US 5,667,425; WO98/25971). In one  
example the antibody molecule of the present invention is a modified Fab fragment wherein  
30 the modification is the addition to the C-terminal end of its heavy chain one or more amino acids to allow the attachment of an effector molecule. Preferably, the additional amino acids

form a modified hinge region containing one or more cysteine residues to which the effector molecule may be attached. Multiple sites can be used to attach two or more PEG molecules.

Preferably PEG molecules are covalently linked through a thiol group of at least one cysteine residue located in the antibody fragment. Each polymer molecule attached to the modified antibody fragment may be covalently linked to the sulphur atom of a cysteine residue located in the fragment. The covalent linkage will generally be a disulphide bond or, in particular, a sulphur-carbon bond. Where a thiol group is used as the point of attachment appropriately activated effector molecules, for example thiol selective derivatives such as maleimides and cysteine derivatives may be used. An activated polymer may be used as the starting material in the preparation of polymer-modified antibody fragments as described above. The activated polymer may be any polymer containing a thiol reactive group such as an  $\alpha$ -halocarboxylic acid or ester, e.g. iodoacetamide, an imide, e.g. maleimide, a vinyl sulphone or a disulphide. Such starting materials may be obtained commercially (for example from Nektar, formerly Shearwater Polymers Inc., Huntsville, AL, USA) or may be prepared from commercially available starting materials using conventional chemical procedures. Particular PEG molecules include 20K methoxy-PEG-amine (obtainable from Nektar, formerly Shearwater; Rapp Polymere; and SunBio) and M-PEG-SPA (obtainable from Nektar, formerly Shearwater).

In one embodiment, the antibody is a modified Fab fragment which is PEGylated, *i.e.* has PEG (poly(ethyleneglycol)) covalently attached thereto, *e.g.* according to the method disclosed in EP 0948544 [see also "Poly(ethyleneglycol) Chemistry, Biotechnical and Biomedical Applications", 1992, J. Milton Harris (ed), Plenum Press, New York, "Poly(ethyleneglycol) Chemistry and Biological Applications", 1997, J. Milton Harris and S. Zalipsky (eds), American Chemical Society, Washington DC and "Bioconjugation Protein Coupling Techniques for the Biomedical Sciences", 1998, M. Aslam and A. Dent, Grove Publishers, New York; Chapman, A. 2002, Advanced Drug Delivery Reviews 2002, 54:531-545]. In one example PEG is attached to a cysteine in the hinge region. In one example, a PEG modified Fab fragment has a maleimide group covalently linked to a single thiol group in a modified hinge region. A lysine residue may be covalently linked to the maleimide group and to each of the amine groups on the lysine residue may be attached a methoxypoly(ethyleneglycol) polymer having a molecular weight of approximately 20,000

Da. The total molecular weight of the PEG attached to the Fab fragment may therefore be approximately 40,000 Da.

In one embodiment, the present invention provides a neutralising antibody molecule having specificity for human IL-17A and human IL-17F, which is a modified Fab fragment  
5 having a heavy chain comprising the sequence given in SEQ ID NO:9 and a light chain comprising the sequence given in SEQ ID NO:7 and having at the C-terminal end of its heavy chain a modified hinge region containing at least one cysteine residue to which an effector molecule is attached. Preferably the effector molecule is PEG and is attached using the methods described in (WO98/25971 and WO2004072116) whereby a lysyl-maleimide group  
10 is attached to the cysteine residue at the C-terminal end of the heavy chain, and each amino group of the lysyl residue has covalently linked to it a methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da. The total molecular weight of the PEG attached to the antibody is therefore approximately 40,000Da.

In another example effector molecules may be attached to antibody fragments using  
15 the methods described in International patent applications WO2005/003169, WO2005/003170 and WO2005/003171.

The present invention also provides an isolated DNA sequence encoding the heavy and/or light chain(s) of an antibody molecule of the present invention. Preferably, the DNA sequence encodes the heavy or the light chain of an antibody molecule of the present  
20 invention. The DNA sequence of the present invention may comprise synthetic DNA, for instance produced by chemical processing, cDNA, genomic DNA or any combination thereof.

DNA sequences which encode an antibody molecule of the present invention can be obtained by methods well known to those skilled in the art. For example, DNA sequences coding for part or all of the antibody heavy and light chains may be synthesised as desired  
25 from the determined DNA sequences or on the basis of the corresponding amino acid sequences.

DNA coding for acceptor framework sequences is widely available to those skilled in the art and can be readily synthesised on the basis of their known amino acid sequences.

Standard techniques of molecular biology may be used to prepare DNA sequences  
30 coding for the antibody molecule of the present invention. Desired DNA sequences may be

synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate.

Examples of suitable sequences are provided in SEQ ID NO:8; SEQ ID NO:10; SEQ ID NO:13; SEQ ID NO:14; SEQ ID NO:17 and SEQ ID NO:18. Nucleotides 1-57 in SEQ ID NO 18 and 1-60 in SEQ ID NO 14 encode the signal peptide sequence from mouse antibody B72.3 (Whittle *et al.*, 1987, Protein Eng. 1(6) 499-505.) which is cleaved to give a neutralising antibody molecule of the present invention.

The present invention also relates to a cloning or expression vector comprising one or more DNA sequences of the present invention. Accordingly, provided is a cloning or expression vector comprising one or more DNA sequences encoding an antibody of the present invention. Preferably, the cloning or expression vector comprises two DNA sequences, encoding the light chain and the heavy chain of the antibody molecule of the present invention, respectively. Preferably, a vector according to the present invention comprises the sequences given in SEQ ID NO:14 and SEQ ID NO:18. Nucleotides 1-57 in SEQ ID NO 18 and 1-60 in SEQ ID NO 14 encode the signal peptide sequence from mouse antibody B72.3 (residues 1-19 in SEQ ID NO: 16 and 1-20 in SEQ ID NO:12 respectively) which is most preferably cleaved to give a neutralising antibody molecule of the present invention.

General methods by which the vectors may be constructed, transfection methods and culture methods are well known to those skilled in the art. In this respect, reference is made to "Current Protocols in Molecular Biology", 1999, F. M. Ausubel (ed), Wiley Interscience, New York and the Maniatis Manual produced by Cold Spring Harbor Publishing.

Also provided is a host cell comprising one or more cloning or expression vectors comprising one or more DNA sequences encoding an antibody of the present invention. Any suitable host cell/vector system may be used for expression of the DNA sequences encoding the antibody molecule of the present invention. Bacterial, for example *E. coli*, and other microbial systems may be used or eukaryotic, for example mammalian, host cell expression systems may also be used. Suitable mammalian host cells include CHO, myeloma or hybridoma cells.

The present invention also provides a process for the production of an antibody molecule according to the present invention comprising culturing a host cell containing a

vector of the present invention under conditions suitable for leading to expression of protein from DNA encoding the antibody molecule of the present invention, and isolating the antibody molecule.

The antibody molecule may comprise only a heavy or light chain polypeptide, in which case only a heavy chain or light chain polypeptide coding sequence needs to be used to 5 transfect the host cells. For production of products comprising both heavy and light chains, the cell line may be transfected with two vectors, a first vector encoding a light chain polypeptide and a second vector encoding a heavy chain polypeptide. Alternatively, a single vector may be used, the vector including sequences encoding light chain and heavy chain 10 polypeptides.

As the antibodies of the present invention are useful in the treatment and/or prophylaxis of a pathological condition, the present invention also provides a pharmaceutical or diagnostic composition comprising an antibody molecule of the present invention in combination with one or more of a pharmaceutically acceptable excipient, diluent or carrier. 15 Accordingly, provided is the use of an antibody according to the present invention for the manufacture of a medicament. The composition will usually be supplied as part of a sterile, pharmaceutical composition that will normally include a pharmaceutically acceptable carrier. A pharmaceutical composition of the present invention may additionally comprise a pharmaceutically-acceptable adjuvant.

20 The present invention also provides a process for preparation of a pharmaceutical or diagnostic composition comprising adding and mixing the antibody molecule of the present invention together with one or more of a pharmaceutically acceptable excipient, diluent or carrier.

The antibody molecule may be the sole active ingredient in the pharmaceutical or 25 diagnostic composition or may be accompanied by other active ingredients including other antibody ingredients, for example anti-TNF, anti-IL-1 $\beta$ , anti-T cell, anti-IFN $\gamma$  or anti-LPS antibodies, or non-antibody ingredients such as xanthines.

The pharmaceutical compositions preferably comprise a therapeutically effective amount of the antibody of the invention. The term "therapeutically effective amount" as used 30 herein refers to an amount of a therapeutic agent needed to treat, ameliorate or prevent a targeted disease or condition, or to exhibit a detectable therapeutic or preventative effect. For

any antibody, the therapeutically effective amount can be estimated initially either in cell culture assays or in animal models, usually in rodents, rabbits, dogs, pigs or primates. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for  
5 administration in humans.

The precise therapeutically effective amount for a human subject will depend upon the severity of the disease state, the general health of the subject, the age, weight and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities and tolerance/response to therapy. This amount can be determined by routine  
10 experimentation and is within the judgement of the clinician. Generally, a therapeutically effective amount will be from 0.01 mg/kg to 50 mg/kg, preferably 0.1 mg/kg to 20 mg/kg. Pharmaceutical compositions may be conveniently presented in unit dose forms containing a predetermined amount of an active agent of the invention per dose.

Compositions may be administered individually to a patient or may be administered in  
15 combination (*e.g.* simultaneously, sequentially or separately) with other agents, drugs or hormones.

The dose at which the antibody molecule of the present invention is administered depends on the nature of the condition to be treated, the extent of the inflammation present and on whether the antibody molecule is being used prophylactically or to treat an existing  
20 condition.

The frequency of dose will depend on the half-life of the antibody molecule and the duration of its effect. If the antibody molecule has a short half-life (*e.g.* 2 to 10 hours) it may be necessary to give one or more doses per day. Alternatively, if the antibody molecule has a long half life (*e.g.* 2 to 15 days) it may only be necessary to give a dosage once per day, once  
25 per week or even once every 1 or 2 months.

The pharmaceutically acceptable carrier should not itself induce the production of antibodies harmful to the individual receiving the composition and should not be toxic. Suitable carriers may be large, slowly metabolised macromolecules such as proteins, polypeptides, liposomes, polysaccharides, polylactic acids, polyglycolic acids, polymeric  
30 amino acids, amino acid copolymers and inactive virus particles.

Pharmaceutically acceptable salts can be used, for example mineral acid salts, such as hydrochlorides, hydrobromides, phosphates and sulphates, or salts of organic acids, such as acetates, propionates, malonates and benzoates.

Pharmaceutically acceptable carriers in therapeutic compositions may additionally  
5 contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents or pH buffering substances, may be present in such compositions. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries and suspensions, for ingestion by the patient.

10 Preferred forms for administration include forms suitable for parenteral administration, e.g. by injection or infusion, for example by bolus injection or continuous infusion. Where the product is for injection or infusion, it may take the form of a suspension, solution or emulsion in an oily or aqueous vehicle and it may contain formulatory agents, such as suspending, preservative, stabilising and/or dispersing agents. Alternatively, the  
15 antibody molecule may be in dry form, for reconstitution before use with an appropriate sterile liquid.

Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals. However, it is preferred that the compositions are adapted for administration to human subjects.

20 The pharmaceutical compositions of this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, transcutaneous (for example, see WO 98/20734), subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, intravaginal or rectal routes. Hyposprays may also be used to administer the pharmaceutical  
25 compositions of the invention. Typically, the therapeutic compositions may be prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared.

Direct delivery of the compositions will generally be accomplished by injection, subcutaneously, intraperitoneally, intravenously or intramuscularly, or delivered to the  
30 interstitial space of a tissue. The compositions can also be administered into a lesion. Dosage treatment may be a single dose schedule or a multiple dose schedule.

It will be appreciated that the active ingredient in the composition will be an antibody molecule. As such, it will be susceptible to degradation in the gastrointestinal tract. Thus, if the composition is to be administered by a route using the gastrointestinal tract, the composition will need to contain agents which protect the antibody from degradation but which release the antibody once it has been absorbed from the gastrointestinal tract.

A thorough discussion of pharmaceutically acceptable carriers is available in Remington's Pharmaceutical Sciences (Mack Publishing Company, N.J. 1991).

It is also envisaged that the antibody of the present invention will be administered by use of gene therapy. In order to achieve this, DNA sequences encoding the heavy and light chains of the antibody molecule under the control of appropriate DNA components are introduced into a patient such that the antibody chains are expressed from the DNA sequences and assembled *in situ*.

The present invention also provides an antibody molecule for use in the control of inflammatory diseases. Preferably, the antibody molecule can be used to reduce the inflammatory process or to prevent the inflammatory process.

The present invention also provides the antibody molecule of the present invention for use in the treatment or prophylaxis of a pathological disorder that is mediated by IL-17A and/or IL-17F or is associated with an increased level of IL-17A and/or IL-17F. Preferably, the pathological condition is selected from the group consisting of infections (viral, bacterial, fungal and parasitic), endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, inflammatory bowel disease, Ulcerative colitis, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, surgical adhesions, stroke, Type I Diabetes, Lyme arthritis, meningoencephalitis, immune mediated inflammatory disorders of the central and peripheral nervous system such as multiple sclerosis and Guillain-Barré syndrome, other autoimmune disorders, pancreatitis, trauma (surgery), graft-versus-host disease, transplant rejection, cancer (both solid tumours such as melanomas, hepatoblastomas, sarcomas, squamous cell carcinomas, transitional cell cancers, ovarian cancers and hematologic malignancies and in particular acute myelogenous leukaemia, chronic myelogenous leukemia, gastric cancer and colon cancer), heart disease including ischaemic

---

diseases such as myocardial infarction as well as atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis and hypochlorhydria.

The present invention also provides an antibody molecule according to the present invention for use in the treatment or prophylaxis of pain.

5 The present invention further provides the use of an antibody molecule according to the present invention in the manufacture of a medicament for the treatment or prophylaxis of a pathological disorder that is mediated by IL-17A and/or IL-17F or associated with an increased level of IL-17A and/or IL-17F. Preferably the pathological disorder is rheumatoid arthritis or multiple sclerosis.

10 The present invention further provides the use of an antibody molecule according to the present invention in the manufacture of a medicament for the treatment or prophylaxis of pain.

An antibody molecule of the present invention may be utilised in any therapy where it is desired to reduce the effects of IL-17A and/or IL-17F in the human or animal body. IL-  
15 17 A and/or IL-17F may be circulating in the body or may be present in an undesirably high level localised at a particular site in the body, for example a site of inflammation.

An antibody molecule according to the present invention is preferably used for the control of inflammatory disease, autoimmune disease or cancer.

The present invention also provides a method of treating human or animal subjects  
20 suffering from or at risk of a disorder mediated by IL-17A and/or IL-17F, the method comprising administering to the subject an effective amount of an antibody molecule of the present invention.

An antibody molecule according to the present invention may also be used in diagnosis, for example in the *in vivo* diagnosis and imaging of disease states involving IL-  
25 17A and/or IL-17F.

The present invention is further described by way of illustration only in the following examples, which refer to the accompanying Figures, in which:

Figure:

- 1 a) Light chain V region of antibody CA028\_0496 (SEQ ID NO:7)  
b) Heavy chain V region of antibody CA028\_0496 (SEQ ID NO:9)  
c) CDRH1 (SEQ ID NO:1), CDRH2 (SEQ ID NO:2), CDRH3 (SEQ ID NO:3),  
5 CDRL1 (SEQ ID NO:4), CDRL2 (SEQ ID NO:5), CDRL3 (SEQ ID NO:6) of  
antibody CA028\_496.  
d) Light chain of antibody CA028\_496 (SEQ ID NO:11).  
e) Heavy chain of antibody CA028\_496 (SEQ ID NO:15).  
10 f) DNA encoding light chain of antibody CA028\_496 including signal sequence  
(SEQ ID NO:14).  
g) DNA encoding heavy chain of antibody CA028\_496 including signal sequence  
(SEQ ID NO:18)

Figure 2 a) The effect of antibody CA028\_0496 (designated Ab#496 in legend) on human IL-  
15 17 induced IL-6 production from Hela cells. b) The effect of antibody CA028\_0496  
(designated Ab#496 in legend) on human IL-17F induced IL-6 production from Hela cells

#### *DNA manipulations and general methods*

*E. coli* strain INV $\alpha$ F' (Invitrogen) was used for transformation and routine culture  
20 growth. DNA restriction and modification enzymes were obtained from Roche Diagnostics  
Ltd. and New England Biolabs. Plasmid preparations were performed using Maxi Plasmid  
purification kits (QIAGEN, catalogue No. 12165). DNA sequencing reactions were  
performed using the ABI Prism Big Dye terminator sequencing kit (catalogue No. 4304149)  
and run on an ABI 3100 automated sequencer (Applied Biosystems). Data was analysed  
25 using the program AutoAssembler (Applied Biosystems). Oligonucleotides were obtained  
from Invitrogen. The concentration of IgG was determined using IgG assembly ELISA.

#### *IL-17 isoforms*

Recombinant IL-17A and IL-17F were purchased from R&D Systems.

Recombinant IL-17A/F heterodimer was produced by linking IL-17A and IL-17F using a GS linker. The heterodimer had the following sequence (SEQ ID NO:19)

MGITIPRNPGPCNSEDKNFPRTVMVNLNIHNRNTNTNPKRSSDYNNRSTSPWNLHRN  
 5 EDPERYPSVIWEAKCRHLGCINADGNVDYHMNSVPIQQEILVLRREPPHCPNSFRLEK  
 ILVSVGCTCVTPIVHHVAGGGGSGGGGSGGGGSGGGGSRKIPKVGHTFFQKPESCP  
 PVPGGSMKLDIGIINENQRVSMRNIESRSTSPWNYTVTWDPNRYPSEVVQAQCRNL  
 GCINAQGKEDISMNSVPIQQETLVVRRKHQGC SVSFQLEKVLVTVGCTCVTPVIHHV  
 Q

10

Recombinant cynomolgus IL-17F (SEQ ID NO:20)

MRKIPKVGHTFFQKPESCPPVPEGSMKLDGTGIINENQRVSMRNIESRSTSPWNYTVTWDPN  
 RYPSEVVQAQCKHLGCINAQGKEDISMNSVPIQQETLVLRKHXQGC SVSFQLEKVLVTVGCT  
 CVTPVIHHVQ

15

The DNA sequence encoding IL-17A/F heterodimer was chemically synthesised by Entelechon GmbH and was subcloned into pET43.1a at the NdeI/XhoI sites.

The DNA sequence encoding cyno L-17F was amplified by PCR using primers that introduced NdeI and XhoI restriction sites. The PCR products were ligated into pCR4Blunt-  
 20 TOPO and sequence verified before digestion and ligation into pET43.1a at the NdeI/XhoI sites.

pET43.1a DNA encoding IL-17 isoforms was used to transfect BL21(DE3) cells and selected carbenicillin-resistant clones were grown at 37°C overnight in 2TY broth containing 2%  
 25 glucose and 50µg/ml carbenicillin. The cultures were then diluted and grown in the same medium to an OD<sub>600</sub> of 0.5-0.7, induced with 1mM IPTG and grown at 37°C for a further 4-5 hours.

Cells were harvested by centrifugation and inclusion bodies prepared from from the cells.  
 30 Inclusion bodies were solubilised in 50mM Tris-HCl, 5M guanidinium hydrochloride, 50mM NaCl, 1mM EDTA, 2mM reduced glutathione, 0.2mM oxidised glutathione, pH 8.5. IL-17 protein was refolded by dropwise addition of the solubilised protein to the above buffer

without guanidinium hydrochloride, with vigorous stirring. The final volume was chosen such that the final protein concentration was no more than 0.1mg/ml.

The refolded protein solution was concentrated if required, before buffer exchange with  
5 10mM MES pH6. The protein was then applied to a column of Sepharose SP HP equilibrated with 20mM MES pH6. Protein was eluted with a linear gradient of 0-500mM NaCl in MES pH6 over 10 column volumes. For IL-17F the gradient was extended to 600mM NaCl. In order to further purify IL-17, the relevant fraction from the Sepharose SP HP column were pooled, concentrated and diluted with 20mM CAPSO (pH10) and applied to a Mono Q  
10 column equilibrated with 20mM CAPSO. Protein was eluted with a linear gradient of 0-250mM NaCl in 20mM CAPSO over 20 column volumes. Fractions containing IL-17 were pooled and neutralised using 1M MES pH6.

#### **Example 1: Production of a neutralising anti-IL-17 antibody**

15 Female Sprague Dawly rats were immunised with recombinant human IL-17 (purchased from R & D systems). Rats received four immunisations of 20µg IL-17 in 100µl Freund's adjuvant. Antibody 225 which binds human IL-17 was isolated using the methods described in WO04/051268. Genes for the heavy chain variable domain (VH) and light chain variable domain (VL) of antibody 225 were isolated and sequenced following cloning via reverse  
20 transcription PCR.

A series of humanised VL and VH regions were designed using human V-region acceptor frameworks and by varying the number of donor residues in the framework regions. Eight grafted VL regions (gL1-8) and 9 grafted VH regions (gH1-9) were designed and genes were built by oligonucleotide assembly and PCR mutagenesis.

25 The light chain grafted sequences were sub-cloned into the human light chain expression vector pKH10.1 which contains the DNA encoding the human C-Kappa constant region (Km3 allotype). The heavy chain grafted sequences were sub-cloned into the human gamma-4 expression vector pVhg4P FL, which contains the DNA encoding the human gamma-4 constant region containing the hinge stabilising mutation S241P (Angal *et al.*,  
30 *supra*). Plasmids were co-transfected into CHO cells and the antibodies produced screened for activity in IL-17 binding and neutralisation assays. Transfections of CHO cells were

performed using the Lipofectamine™ 2000 procedure according to manufacturer's instructions (Invitrogen, catalogue No. 11668).

The most optimal graft based on expression, affinity and neutralisation potency (gL7gH9) was selected and named CA028\_0496. The V region sequences of this antibody  
5 are shown in Figure 1 (a) and (b) and in SEQ ID NOs: 7 and 9 for the light chain (gL7) and heavy chains (gH9) respectively.

The heavy chain acceptor framework is the human germline sequence VH3 1-3 3-07 with framework 4 coming from this portion of the human JH-region germline JH4. The light chain acceptor framework is the human germline sequence VK1 2-1-(1) L4, with framework  
10 4 coming from this portion of the human JK-region germline JK1.

**Example 2: Antibody CA028\_0496 neutralises IL-17 and IL-17F and IL-17A/F heterodimer**

Hela cells

The potency of antibody CA028\_0496 against human recombinant IL-17 and human  
15 recombinant IL-17F in Hela cells was tested and compared to antibody CDP435 (WO06/054059). Hela cells were obtained from the cell bank at ATCC (ATCC CCL-2). Cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% foetal calf serum, penicillin, gentamycin and glutamine.  $1 \times 10^4$  cells were plated out into 96 well flat bottomed tissue culture plates. Cells were incubated overnight and washed once in  
20 assay buffer. Either human IL-17A ( $25 \text{ ng ml}^{-1}$ ) or human IL-17F ( $125 \text{ ng ml}^{-1}$ ) was incubated in the presence of a fixed concentration of human TNF- $\alpha$  this mixture was preincubated with antibody CA028\_0496 or antibody CDP435. Cytokine plus antibody was then added to the Hela cells which were incubated overnight. The production of IL-6 in the cell culture supernatant was proportionate to the amount of IL-17A/IL-17F added to the cells. Human IL-  
25 6 levels were measured by ELISA and quantified by comparison with known standard concentrations of human IL-6.

The data (Figures 2a and 2b) indicates that antibody CA028\_0496 potently neutralised human recombinant IL-17A and also had some activity against human IL-17F. The data from these  
30 experiments indicated that antibody CA028\_0496 gave an  $\text{IC}_{50}$  of  $43 \text{ ng/ml}$  against human recombinant IL-17 ( $25 \text{ ng ml}^{-1}$ ) and  $1477 \text{ ng/ml}$  against recombinant IL-17F ( $125 \text{ ng ml}^{-1}$ ).

Accordingly, antibody CA028\_0496 gave an IC50 of 0.29M against human recombinant IL-17 (0.78nM) and 10.18nM against human recombinant IL-17F (4.16nM) in this assay (calculation based on per IgG assuming a molecular weight of 145,000 as an average IgG4 and assuming that IL-17A and IL-17F are dimers).

5

#### Human microglia cells

Human microglia cells (TCS Cellworks) were plated out in a flat bottom 96-well plate at 5,000 cells per well in a total volume of 100  $\mu$ l and left for 24 hours to attach to the plastic. At this time titrations (5, 1, 0.2 and 0.04  $\mu$ g/ml) of human recombinant IL-17A, human recombinant IL-17F, cynomolgus recombinant IL-17F and human recombinant IL-17A/F heterodimer in the presence and absence of 10 ng/ml human recombinant TNF $\alpha$  were added to wells in triplicate. Control wells contained no stimulation, IL-17A alone (100ng/ml), TNF $\alpha$  alone and IL-17A and TNF $\alpha$  together. All cytokines were added in a total volume of 110  $\mu$ l/well, making the total well volume 210  $\mu$ l. In experiments involving antibodies, cells were plated out in the same way. After 24 hours antibodies and cytokines were added at the same time to give the stated final concentrations in a total final volume of 200  $\mu$ l.

After a further 24 hours incubation at 37 $^{\circ}$ C, supernatants were harvested and frozen at -20 $^{\circ}$ C until analysis. For analysis, supernatants were diluted 1/10 and measured for IL-6 using a human IL-6 MSD kit, according to manufacturer's instructions.

All isoforms of IL-17 tested were found to be active in the assay, particularly in the presence of TNF $\alpha$ .

The potency of antibody CA028\_0496 against human recombinant IL-17A and human recombinant IL-17F, cynomolgus recombinant IL-17F and human recombinant IL-17A/F heterodimer in human microglia cells was tested in the presence of TNF $\alpha$  and compared to a control antibody and an IL-17A specific antibody using the method described above.

The control antibody had no effect on the activity of any of the cytokines tested.

Antibody CA028\_0496 had inhibitory activity against all three cytokines IL-17, IL-17F and IL-17A/F, including cynomolgus IL-17F while the IL-17A specific antibody only had inhibitory activity against IL-17A and IL-17A/F heterodimer.

5 **Example 3: Affinity of antibody CA028\_0496 (human IgG4 constant regions) for IL-17A and IL-17F**

BIA (Biomolecular Interaction Analysis) was performed using a Biacore 3000 (Biacore AB). All experiments were performed at 25 °C. Affinipure Fc Fragment goat anti-human IgG, Fc fragment specific (Jackson ImmunoResearch) was immobilised on a CM5 Sensor Chip via  
10 amine coupling chemistry to a capture level of  $\approx 6000$  response units (RUs). HBS-EP buffer (10mM HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005 % Surfactant P20, Biacore AB) was used as the running buffer with a flow rate of 10  $\mu$ l/min. A 10  $\mu$ l injection of antibody CA028\_0496 (1.81mg/ml) was used for capture by the immobilised anti-human IgG-Fc. Human IL-17A and IL-17 isoforms were titrated over the captured CA028\_0496 at doubling  
15 dilutions from 50nM to sub nM at a flow rate of 30  $\mu$ L/min. The surface was regenerated by a 30  $\mu$ L injections of 40 mM HCl, followed by one 5  $\mu$ L injection of 5 mM NaOH.

Background subtraction binding curves were double referenced and analysed using the BIAevaluation software (version 3.2) following standard procedures. Kinetic parameters  
20 were determined from the fitting algorithm.

The affinity value determined for antibody CA028\_0496 binding IL-17A was 16 pM and 1750pM for IL-17F. Antibody CA028\_0496 did not bind to the other IL-17 isoforms (IL-17 B, C, D and E). Antibody CA028\_0496 therefore specifically binds IL-17A and IL-17F.

25 **Example 4: Affinity of antibody CA028\_0496 (murine IgG1 constant regions) for IL-17A, cynomolgus IL-17F and IL-17A/F heterodimer**

BIA (Biomolecular Interaction Analysis) was performed using a Biacore 3000 (Biacore AB).

All experiments were performed at 25 °C. Affinipure F(ab')<sub>2</sub> fragment goat anti-mouse IgG, Fc fragment specific (Jackson ImmunoResearch) was immobilised on a CM5 Sensor Chip  
30

- (Biacore AB) via amine coupling chemistry to a capture level of  $\approx 6000$  response units (RUs). HBS-EP buffer (10mM HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005 % Surfactant P20, Biacore AB) was used as the running buffer with a flow rate of 10  $\mu\text{L}/\text{min}$ . A 10  $\mu\text{L}$  injection of antibody CA028\_0496 at 4 $\mu\text{g}/\text{mL}$  was used for capture by the
- 5 immobilised anti-mouse IgG, Fc. Human IL-17A, cyno IL-17F and heterodimerA/F were titrated over the captured CA028\_0496 at doubling dilutions from 25nM to sub nM at a flow rate of 30  $\mu\text{L}/\text{min}$ . The surface was regenerated at a flowrate of 10 $\mu\text{L}/\text{min}$  by a 10  $\mu\text{L}$  injection of 40 mM HCl, followed by a 5  $\mu\text{L}$  injection of 5 mM NaOH.
- 10 Double referenced background subtracted binding curves were analysed using the BIAevaluation software (version 3.2) following standard procedures. Kinetic parameters were determined from the fitting algorithm.

- Antibody CA028\_0496 had an affinity of 21pM for IL-17A, 116pM for IL-17A/F
- 15 heterodimer and 1030pM for cynomolgus IL-17F.

The scope of the claims should not be limited by the preferred embodiments set forth in the examples, but should be given the broadest interpretation consistent with the description as a whole.

**CLAIMS:**

1. A neutralising antibody which specifically binds human IL-17A and human IL-17F, comprising a heavy chain and a light chain wherein the heavy chain variable domain comprises the sequence given in SEQ ID NO:9 or a sequence at least 95% identical thereto and the light chain variable domain comprises the sequence given in SEQ ID NO:7 or a sequence at least 95% identical thereto, wherein the heavy chain comprises the CDRH1, CDRH2 and CDRH3 identified in SEQ ID NO: 1, 2 and 3 respectively and wherein the antibody has an affinity for IL-17A of better than 20pM and an affinity for IL-17F of better than 2nM.
2. The neutralising antibody according to claim 1, having the heavy chain comprising the sequence given in SEQ ID NO:9.
3. The neutralising antibody according to claim 1 or 2, wherein the light chain comprises the CDRL1, CDRL2 and CDRL3 identified in SEQ ID NO: 4, 5 and 6 respectively.
4. The neutralising antibody according to any one of claim 1 to 3, having the heavy chain comprising the sequence given in SEQ ID NO:9 and the light chain comprising the sequence given in SEQ ID NO:7.
5. The neutralising antibody according to any one of claims 1 to 4 which binds human IL-17A and human IL-17F, having the heavy chain comprising the sequence given in SEQ ID NO:15 and the light chain comprising the sequence given in SEQ ID NO:11.
6. The neutralising antibody according to any one of claims 1 to 5 wherein the antibody is a whole antibody or a functionally active fragment.
7. The neutralising antibody according to claim 6 where the antibody fragment is a domain antibody, Fab, Fab', F(ab')<sub>2</sub>, scFv or an epitope-binding fragment thereof.

8. The neutralising antibody according to claim 6 which is multi-specific.
9. The neutralising antibody according to any one of claims 1-8, wherein the antibody or  
5 fragment thereof is conjugated to one or more effector molecule(s).
10. An isolated DNA molecule encoding the heavy and the light chain(s) of the neutralising  
antibody as defined in any one of claims 1 to 9.
- 10 11. A cloning or expression vector comprising one or more DNA molecules as defined in  
claim 10.
12. A cloning or expression vector comprising the isolated DNA molecule as defined in claim  
10.
- 15 13. The vector according to claim 11 or 12, wherein the vector comprises the sequence given  
in SEQ ID NO:18.
14. The vector according to claim 11 or 12, wherein the vector comprises the sequences given  
20 in SEQ ID NO:14 and SEQ ID NO:18.
15. A host cell comprising one or more cloning or expression vectors as defined in any one of  
claims 11 to 14.
- 25 16. A process for the production of the antibody as defined in any one of claims 1 to 9,  
comprising culturing the host cell of claim 15 and isolating the antibody.
17. A pharmaceutical composition comprising the antibody according to any one of claims 1  
to 9, in combination with one or more of a pharmaceutically acceptable excipient, diluent  
30 or carrier.

18. The pharmaceutical composition according to claim 17, additionally comprising other active ingredients.
- 5 19. The antibody according to any one of claims 1 to 9 or the pharmaceutical composition according to claim 17 or claim 18, for use in the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's  
10 Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoenephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma, graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic  
15 disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydia.
20. The antibody according to claim 19 or the pharmaceutical composition according to claim 19, wherein the immune mediated inflammatory disorder of the central and peripheral  
20 nervous system is multiple sclerosis or Guillain-Barré syndrome.
21. The antibody according to claim 19 or the pharmaceutical composition according to claim 19, wherein the trauma is surgery.
- 25 22. The antibody according to claim 19 or the pharmaceutical composition according to claim 19, wherein the cancer is a solid tumour or an hematologic malignancy.
- 30 23. The antibody according to claim 22 or the pharmaceutical composition according to claim 22, wherein the solid tumour or the hematologic malignancy is melanoma, hepatoblastoma, sarcoma, squamous cell carcinoma, transitional cell cancer, ovarian

cancer, acute myelogenous leukaemia, chronic myelogenous leukemia, gastric cancer or colon cancer.

- 5 24. The antibody according to claim 19 or the pharmaceutical composition according to claim 19, wherein the heart disease is myocardial infarction.
- 10 25. Use of the antibody according to any one of claims 1 to 9 or the pharmaceutical composition according to claim 17 or claim 18, for the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma, graft-versus-host disease, transplant rejection, cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydria.
- 15
- 20 26. Use of the antibody according to any one of claims 1 to 9 or the pharmaceutical composition according to claim 17 or claim 18, for the preparation of a medicament for the treatment or prophylaxis of a pathological disorder mediated by IL-17A and/or IL-17F and being a viral infection, a bacterial infection, a fungal infection, a parasitic infection, an endotoxic shock associated with infection, arthritis, rheumatoid arthritis, asthma, pelvic inflammatory disease, Alzheimer's Disease, Crohn's disease, Peyronie's Disease, coeliac disease, gallbladder disease, Pilonidal disease, peritonitis, psoriasis, vasculitis, a surgical adhesion, stroke, Type I Diabetes, lyme arthritis, meningoencephalitis, an immune mediated inflammatory disorder of the central and peripheral nervous system, an autoimmune disorder, pancreatitis, trauma, graft-versus-host disease, transplant rejection,
- 25

cancer, heart disease, ischaemic disease, atherosclerosis, intravascular coagulation, bone resorption, osteoporosis, periodontitis or hypochlorhydria.

- 5 27. The use according to claim 25 or 26, wherein the immune mediated inflammatory disorder of the central and peripheral nervous system is multiple sclerosis or Guillain-Barré syndrome.
28. The use according to claim 25 or 26, wherein the trauma is surgery.
- 10 29. The use according to claim 25 or 26, wherein the cancer is a solid tumour or an hematologic malignancy.
- 15 30. The use according to claim 29, wherein the solid tumour or the hematologic malignancy is melanoma, hepatoblastoma, sarcoma, squamous cell carcinoma, transitional cell cancer, ovarian cancer, acute myelogenous leukaemia, chronic myelogenous leukemia, gastric cancer or colon cancer.
31. The use according to claim 25 or 26, wherein the heart disease is myocardial infarction.

1/4

Figure 1

**(a) Light Chain variable region of antibody CA028\_496 (SEQ ID NO:7)**

AIQLTQSPSSLSASVGDRVTITCRADESVTTLMHWYQQKPGKAPKLLIYLVS NRESGVPSRF  
SGSGSGTDFTLTISSLQPEDFATYYCQQTWSDPWTFGQGTKVEIKR

**(b) Heavy Chain variable region of antibody CA028\_496 (SEQ ID NO:9)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDYNMAWVRQAPGKGLEWVATITYEGRNTYYRD  
SVKGRFTISRDNKNSLYLQMNSLRAEDTAVYYCASPPQYYEGSIYRLWFAHWGQGLVTVS  
S

(c)

CDRH1: GFTFSDYNMA (SEQ ID NO:1)  
CDRH2: TITYEGRNTYYRDSVKG (SEQ ID NO:2)  
CDRH3: PPQYYEGSIYRLWFAH (SEQ ID NO:3)  
CDRL1: RADESVTTLMH (SEQ ID NO:4)  
CDRL2: LVS NRES (SEQ ID NO:5)  
CDRL3: QQTWSDPWT (SEQ ID NO:6)

**(d) Light chain of antibody CA028\_496 (SEQ ID NO:11)**

AIQLTQSPSSLSASVGDRVTITCRADESVTTLMHWYQQKPGKAPKLLIYLVS NRESGVPSRF  
SGSGSGTDFTLTISSLQPEDFATYYCQQTWSDPWTFGQGTKVEIKRTVAAPSVFIFPPSDEQ  
LKSGTASVVCLLNMFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADY  
EKHKVYACEVTHQGLSSPVTKSFNRGEC

**(e) Heavy chain of antibody CA028\_496 (SEQ ID NO:15)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDYNMAWVRQAPGKGLEWVATITYEGRNTYYRD  
SVKGRFTISRDNKNSLYLQMNSLRAEDTAVYYCASPPQYYEGSIYRLWFAHWGQGLVTVS  
SASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG  
LYSLSSVVTVPSSSLGKTYTCNVDPKPKSKVDPKRVESKYGPPCPPCPAPEFLGGPSVFLF  
PPKPKDTLMISRTPEVTCVVDVDSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSV  
LTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCL  
LVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH  
EALHNHYTQKSLSLSLGK

**(f) DNA encoding light chain of antibody CA028\_496 including signal sequence (SEQ ID NO:14)**

atgtcagttcccacacaggtgctgggcctgcttctgttgtggctcaccgatgctaggtgtgc  
catccagctgaccagagcccttctctctcagcgccagtgctcggagacagagtgactatta  
cctgcagggctgacgaaagcgtgaccacattgatgcactggtaaccaacagaagcctggcaa  
gcccccaagctcctgatctatctggtttccaatcgggagctctggagtcaccagcaggttcag  
cggcagtggttctggaactgactttaccctgacaatctcctcactccagcccgaagatttcg  
ccacctactattgccagcagacttggagcgcacccttggacatttggacagggcacaaaagtg  
gagatcaagcgtacggtagcggcccatctgtcttcatcttcccgcctctgatgagcagtt  
gaaatctggaactgcctctgttgtgtgcctgctgaataacttctatcccagagaggccaag  
tacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgtcacagagcag  
gacagcaaggacagcacctacagcctcagcagcaccctgacgctgagcaaagcagactacga

2/4

Figure 1 continued

gaaacacaaagtctacgcctgcaagtcacccatcagggcctgagctcgcccgtcaciaaaga  
gcttcaacaggggagagtgtag

(g) DNA encoding heavy chain of antibody CA028\_496 including signal sequence (SEQ ID NO:18)

atggaatggtcctgggtcttctgtttttcctttctgtcacaaccggggtgcacagcgagggt  
tcagctcgttgaatccggaggcggactcgtgcagcctgggggctccttgccgctgagctgcg  
ctgccagtggttcactttcagcgcattacaatatggcctgggtgcccaggccccaggcaag  
ggtctggagtggtggccacaattacctatgagggcagaaacacttattaccgggattcagt  
gaaagggcgatttaccatcagcagggataatgcaaagaacagtctgtacctgcagatgaact  
ctctgagagctgaggacaccgctgtctactattgtgcaagcccaccccagtagctatgagggc  
tcaatctacagattgtggtttgccattggggccagggaacactgggtgaccgtctcgagcgc  
ttctacaaagggcccatccgtcttccccctggcgcctgctccaggagcacctccgagagca  
cagccgcccctgggctgctgggtcaaggactacttccccgaaccgggtgacgggtgctggaac  
tcaggcgcctgaccagcggcgtgcacaccttcccggctgtcctacagtcctcaggactcta  
ctccctcagcagcgtgggtgaccgtgcccctccagcagcttgggcacgaagacctacacctgca  
acgtagatcacaagcccagcaacaccaagggtggacaagagagttgggtgagaggccagcacag  
ggagggagggtgtctgctggaagccaggctcagccctcctgctggacgcaccccggctgtg  
cagcccagcccagggcagcaaggcatgccccatctgtctcctcaccggaggcctctgacc  
accccactcatgcccaggagagggtcttctggatttttccaccaggctccgggcagccaca  
ggctggatgcccctaccccaggccctgcgcatcacaggggaggtgctgctgagacctgcc  
aagagccatatccgggaggacctgcccctgacctaaagcccaccccaaaggccaaactctcc  
actccctcagctcagacaccttctctcctcccagatctgagtaactcccaatcttctctctg  
cagagtccaaatatgggtccccatgcccaccatgcccaggtaagccaaccaggcctcgccc  
tccagctcaaggcgggacaggtgcccctagagtagcctgcatccagggacaggccccagccgg  
gtgctgacgcatccacctccatctctcctcagcacctgagttcctggggggaccatcagtc  
ttcctgttccccccaaaacccaaggacactctcatgatctcccggaccccctgagggtcacgtg  
cgtgggtgggtggacgtgagccaggaagaccccagaggctccagttcaactgggtacgtggatggcg  
tggagggtgcataatgccaagacaaagccgcgggaggagcagttcaacagcacgtaccgtgtg  
gtcagcgtcctcaccgtcctgcaccaggactggctgaacggcaaggagtacaagtgcaagggt  
ctccaacaaaggcctcccgtcctccatcgagaaaacctctccaaagccaaagggtgggacc  
acgggggtgagagggccacatggacagaggtcagctcggcccaccctctgcccctgggagtgac  
cgctgtgccaacctctgtccctacagggcagccccgagagccacagggtgtacacctgcccc  
catcccaggaggagatgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctac  
cccagcgacatcgccgtggagtgaggagcaatgggcagccggagacaactacaagaccac  
gcctcccgtgctggactccgacggctccttcttctctacagcaggctaaccgtggacaaga  
gcaggtggcaggaggggaatgtcttctcatgctccgtgatgcatgaggctctgcacaaccac  
tacacacagaagagcctctccctgtctctgggtaaa

Figure 2a




Figure 2b

