



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION-TREATY (PCT)

<p>(51) International Patent Classification³ : A61K 7/16, 37/16</p>	<p>A1</p>	<p>(11) International Publication Number: WO 82/ 03008 (43) International Publication Date: 16 September 1982 (16.09.82)</p>
<p>(21) International Application Number: PCT/AU82/00022 (22) International Filing Date: 4 March 1982 (04.03.82) (31) Priority Application Numbers: PE 7855 PE 9429 (32) Priority Dates: 4 March 1981 (04.03.81) 23 June 1981 (23.06.81) (33) Priority Country: AU (71) Applicants (for all designated States except US): THE UNIVERSITY OF MELBOURNE [AU/AU]; Grattan Street, Parkville, Vic. 3052 (AU). THE VICTORIAN DAIRY INDUSTRY AUTHORITY [AU/AU]; Donville Avenue, Kew, Vic. 3101 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): REYNOLDS, Eric, Charles [AU/AU]; STOREY, Elsdon [AU/AU]; MCDUGALL, Wallace, Arthur [AU/AU]; The University of Melbourne, Grattan Street, Parkville, Vic. 3052 (AU).</p>	<p>(74) Agent: SANDERCOCK, SMITH & BEADLE; 203 Riversdale Road, Hawthorn, Vic. 3122 (AU). (81) Designated States: AT (European patent), AU, BR, CH (European patent), DE (European patent), FR (European patent), GB (European patent), JP, NL (European patent), SE (European patent), US. Published <i>With international search report.</i></p>	
<p>(54) Title: CARIES INHIBITION</p> <p>(57) Abstract</p> <p>Compositions for the inhibition of caries and gingivitis, containing a protein or polypeptide particularly phosphoproteins or polyphosphopeptides including those containing the amino acid sequence (X-Y-Z)_n where X and Z are a phosphoserine, phosphothreonine, phosphotyrosine, glutamate or aspartate, Y is any amino acid and n is 1 or more. Particular examples of suitable active ingredients are sodium caseinate, calcium caseinate and phosvitin. Suitable proteins or polypeptides were tested via the dissolution rate of hydroxyapatite as measured by the rate of calcium and phosphate released from a hydroxyapatite column.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	KP	Democratic People's Republic of Korea
AU	Australia	LI	Liechtenstein
BE	Belgium	LK	Sri Lanka
BR	Brazil	LU	Luxembourg
CF	Central African Republic	MC	Monaco
CG	Congo	MG	Madagascar
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SE	Sweden
FR	France	SN	Senegal
GA	Gabon	SU	Soviet Union
GB	United Kingdom	TD	Chad
HU	Hungary	TG	Togo
JP	Japan	US	United States of America

'CARIES INHIBITION'

- 1 -

This invention relates to caries inhibition.
The invention also relates to gingivitis inhibition.

The present invention provides an orally
ingestible composition containing a caries and gingivitis
5 inhibiting amount of a protein or a polypeptide or a salt
thereof.

Preferably, the protein or polypeptide is a
phosphoprotein or a polyphosphopeptide.

Preferably, the protein or polypeptide is an
10 acidic phosphoprotein or polypeptide.

Preferably, the protein or polypeptide contains
the amino acid sequence (X-Y-Z) where X and Z are a phos-
phoserine, phosphothreonine, phosphotyrosine, glutamate or
aspartate and Y is any amino acid.

Preferably, the protein or polypeptide contains
15 a plurality of units each having the amino acid sequence
(X-Y-Z) where X, Y and Z are as stated in above.

Preferably, the protein or polypeptide contains a
group of formula (X-Y-Z)_n where X, Y and Z are as stated
20 in claim 4 and n is 1 or more.

Preferably, n is 3 or more.

Preferably, X and Z are phosphoserine.

Preferably, the protein or polypeptide is a
polyphosphoserine.

Preferably, the phosphate groups of the polyphos-
phoserine are spaced at about 6.88 Angstrom Units.

Preferably, the protein is a casein.

Preferably, the protein is alpha-s-casein.

Alternatively, the protein is phosvitin.

- 2 -

Preferably, the protein or polypeptide is in solution.

Preferably, the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 45 nmol/min under the test conditions defined herein.

1 Preferably, the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 80 nmol/min under the test conditions defined herein.

10 Preferably, the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 90 nmol/min under the test conditions defined herein.

Preferably, the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 95 nmol/min under the test conditions defined herein.

15 Preferably, the protein or polypeptide is present as up to 10% by weight.

Preferably, the protein or polypeptide is present as up to 5% by weight.

Preferably, the protein or polypeptide is present as up to 2% by weight.

20 Preferably, the composition additionally contains urea.

25 The composition of this invention may be in the form of a foodstuff, confectionery, dentifrice, tablet or comprise a pharmacologically acceptable vehicle, solution of suspension for topical application to the teeth or a mouthwash. Other modes of administering the protein or polypeptide would be acceptable if pharmacologically acceptable.

30 Of particular interest as compositions are chewing gum, breakfast foods, ice-cream and other frozen confectionery, confectionery, sweets and cakes as these are all known as caries problem foods.

Also of particular interest are dentifrices, mouth-washes and preparations for topical application to teeth.

35 α_s -casein and other caseins are obtainable from milk, phosphovitin is obtainable from egg yolks and other suitable phosphoproteins include those which are obtainable from



cereals, nuts and vegetables particularly in bran husks or sheaths. In particular, rice, wheat, oat, barley and rye brans are a source of such phosphoproteins.

53 Polypeptides of interest include polyphosphoserine, polyglutamate and polyaspartate.

10 The present invention also provides a process of inhibiting dental caries and/or tooth erosion and/or gingivitis comprising applying to the teeth with a carrier a cariostatic and/or antigingivitis agent selected from a protein or a polypeptide or a salt thereof.

The present invention also provides a first test procedure for selecting among the proteins or polypeptides that may be used those that are most effective.

TEST 1

15 The purpose of this test is to determine hydroxyapatite dissolution and in this respect since tooth enamel is largely composed of hydroxyapatite it is believed that useful comparisons can be made.

20 Double distilled, deionised water (greater than 10 MΩ/cm) was used throughout. Analytical reagent grade chemicals were obtained from Ajax Chemicals, Australia. Hydroxyapatite-spheroidal was purchased from BDH, and phosvitin from Sigma Chemical Co., Missouri, U.S.A. The milk proteins were fractionated by the method of Zittle and Custer (1), and their purity assessed by polyacrylamide gel electro-
25 phoresis using a modification of the method of Groves and Kiddy (2).

Methods

Hydroxyapatite Dissolution Rate Assay

30 A chromatography column (pharmacia K9/15) containing 1 g. of hydroxyapatite beads was used for the demineralisation assay. Tris 5 mM, pH 8.3 containing 50 mM NaCl and 20 mg/l neomycin was used as the column influent buffer at 20°C and at a rate of 1.000±0.003 ml./min. A protein solution (10 mg.
35 10 ml. of influent buffer) was applied to the column and 1 ml. fractions were collected before and after protein application

and assayed for total calcium, phosphate and protein. From these values a rate of dissolution (nmol calcium or phosphate per min) for each 1 ml. fraction was obtained.

Calcium, Phosphate and Protein Assays

5 Inorganic phosphate was measured by the method of Itaya and Ui(3) and protein by the method of Bradford (4). The determination of calcium was by atomic absorption spectrophotometry using 1% lanthanum chloride to prevent phosphate suppression.

10 Results

The proteins used for the study are listed in Table 1, they are all acidic proteins and included four phosphoproteins and three non-phosphorylated proteins from the whey fraction of bovine milk. The effect of the individual proteins on hydroxyapatite dissolution rate is shown in Table 2.

15 Table 1. Properties of various phosphorylated and non-phosphorylated acidic proteins.

Protein	Molecular Weight	Phosphoserine Content ^a	Isoelectric Point	Carbohydrate Content
Phosvitin	35,500	110	1.5	+
α_s -casein	23,613	8	4.1	-
β -casein	24,020	5	4.5	-
25 κ -casein	19,037	1	3.7	+
α -lactalbumin	14,174	-	5.1	-
β -lactoglobulin	18,362	-	5.3	-
Bovine serum albumin	66,210	-	4.7	-

30 a. Residues per mol protein

Table 2. Effect of phosphorylated and non-phosphorylated proteins on hydroxyapatite dissolution rate.

Protein	Reduction in calcium dissolution rate ^a (nmol/min)	Reduction in phosphate dissolution rate (nmol/min)	Amount of protein bound (mg)
Phosvitin	93.1± 5.4 ^b	63.8± 9.4	1.87±0.62
α _s -casein	100.1± 4.1 ^b	63.5± 3.3	5.58±0.03
β-casein	94.8±11.7 ^b	64.0±19.3	7.45±0.37
κ-casein	56.3± 8.9	33.7± 6.8	4.17±0.26
α-lactalbumin	2.7± 1.7	2.9± 0.6	0.48±0.17
β-lactoglobulin	17.1± 1.7	12.5± 1.2	1.80±0.71
Bovine serum albumin	31.6± 4.5	20.5± 3.3	2.09±0.05

a. means±SD, n = 3

b. not significantly different P>0.5

In a trial of the above test the dissolution rate of hydroxyapatite as measured by the rate of calcium and phosphate released from the hydroxyapatite column was constant over a two hour period calcium 353.6±3.9 nmol/min, phosphate 225.4±6.8 nmol/min. The dissolution rates obtained using different hydroxyapatite columns showed greater variation, calcium 354.2±23.8 nmol/min, phosphate 229.6±30.8 nmol/min, n = 11. This intercolumn variation in dissolution rate could be attributable to different column packing resulting in a different HA surface area exposed.

The effect of phosvitin on the dissolution rate of hydroxyapatite is shown in Figure 1. The protein caused an initial increase in the dissolution rate of phosphate which then decreased and approached a new steady state level; 63.8 nmol/min less than the rate prior to phosvitin application. The protein caused an immediate and marked drop in the calcium dissolution rate which then increased and approached a steady-state level 93.1 nmol/min less than that prior to phosvitin application. The amount of protein that passed through the

column was measured and from this the amount retained was calculated 1.87 mg. The dissolution rate returned to the original value after phosvitin was eluted from the column with 10 ml of eluent buffer containing 1.5 M phosphate, followed by buffer not containing phosphate.

The trace for α_s -casein was very similar to that of phosvitin except that the immediate drop in calcium dissolution rate was not as marked Fig. 2. The difference in the steady-state rates of calcium and phosphate released before and after α_s -casein application were very similar to those in Figure 1 (calcium, 100.1 nmol/min. phosphate 63.5 nmol/min).

The results obtained for β -casein (Fig. 3) were characteristic of all the other proteins tested, except for the final steady-state rates of calcium and phosphate released. All proteins (with the exception of phosvitin and α_s -casein) caused an initial increase in the dissolution rate of both calcium and phosphate which then decreased as the proteins passed out of the column. The mean differences between the steady-state dissolution rates before and after protein application, together with the amount of protein bound, for all proteins tested is presented in Table 2 above. The results show that the four phosphoproteins gave a marked reduction in the steady-state dissolution rate of HA with phosvitin, α_s -casein and β -casein all giving the same reduction in calcium and phosphate dissolution.

The results show that all the acidic proteins caused a reduction in the steady-state dissolution rate of hydroxyapatite. However, the greatest reduction was given by the four phosphoproteins; phosvitin, α_s -casein, β -casein and to a lesser extent κ -casein.

- 7 -

From the above and from other data which suggests that adjacent phosphate groups of polyphosphoserine compounds have a spacing of about 6.88 Angstrom units when in a beta-pleated sheet configuration and that calcium atoms in a hydroxyapatite surface along the c-axis will also be spaced at about 6.88 Angstrom units, we speculate that a phosphate group-calcium atom bond materially reduces hydroxyapatite dissolution.

References:

1. Zittle, C.A., Custer, J.H.: Purification and some properties of α_s -casein, J. Dairy Sci 46: 1183-1189, 1963.
2. Groves, M.L., Kiddy, C.A.: Polymorphism of γ -casein in cow's milk. Arch. Biochem. Biophys 126: 188-193, 1968.
3. Itaya, K., Ui, M.: A new micromethod for the colorimetric determination of inorganic phosphate. Clin, Chim, Acta 14: 361-366, 1966.
4. Bradford, MM.: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72:248-254, 1976.

TEST 2

For the purpose of in vivo testing, the following test procedure for determining the effect of casein and whey protein on caries incidence in the Sprague-Dawley rat.

Materials and Methods.

Forty-five, weanling, male Sprague-Dawley rats, 18 days old, bred from animals fed a fluoride free diet were used. The rats were marked for identification and then randomly distributed with respect to diet. They were housed in raised-bottom stainless-steel cages, one group of 15 per cage and fed a powdered cariogenic diet with either deionised water (control), a 2% casein solution or a 2% whey protein solution ad libitum. The cariogenic diet was a modified MIT - 200 diet shown in Table 3.

Table 3 - Composition of modified MIT - 200 cariogenic diet

	Component	% wt
5	Sucrose ^a	67
	Wheyprotein Concentrate	20
	Salt mixture ^b	3
	Cottonseed oil	3
	Cellulose ^a	6
10	Vitamin mixture ^{a, b}	1

a.. Calcium and phosphate not detectable, fluoride content of complete diet was less than 0.2 g per g dry weight.

b. Vitamin and Salt mixture described in detail elsewhere.

15 The animals were weighed daily and the amounts of powdered diet and liquid consumed over a 24 h period by each group was measured. After 42 days on the diet, the animals were killed by cervical dislocation and treated as described below.

Caries evaluation.

20 Fissure caries was assessed using the method of Konig, Marthaler and Muhlemann (1958)(5). The mandible was removed from each rat and placed in formol-saline. The jaws were sectioned and stained by the method of Konig et al (1958)(5), as described by Green and Hartles
25 (1966)(6) to provide series of 100 µm thick longitudinal mesio-distal sections of the molar teeth. Only the main fissures of the first and second molar teeth were assessed for caries.

Results

30 Diet Consumption

The relative consumption of solid and liquid diet by the three groups of rats was tested by an analysis of variance (by diet). This showed that the quantities of both solid and liquid consumed by each group were not
35 significantly different ($P > 0.75$).

Caries Analysis.

The caries data shown in Table 4 were analysed in an analysis of variance table by diet.

Table 4

5	Caries Experience Data	Number of carious fissures ^a
Animals		
Control Group	7.92±2.06	
10	Casein Group	1.87±2.50
	Whey Protein Group	4.73±3.85

a. Maximum number possible 10.

15 The animals drinking the 2% casein solution had 76.5% less carious fissures than the control animals ($p < 0.001$), and the animals consuming the 2% whey protein solution had 40.3% less carious fissures than the control group ($p < 0.01$). The correlation of caries incidence with the final weight of the rat was tested for the three groups. 20 No correlation attained significance ($p > 0.1$).

Similarly, the initial and final weights showed no correlation, nor were weight gain and caries incidence correlated.

Conclusion

25 Acidic proteins in the drinking water substantially reduced caries incidence of male Sprague-Dawley rats, however the phosphorylated protein (casein) caused a greater reduction than the non-phosphorylated whey proteins.

References:

- 30 (5) König K.G., Marthaler T.M. and Muhlemann H.R. 1958: Methodik der Kurzfristig erzeugten Rattenkaries. Dr. Zahn-Mund-u. Kieferheilk. 29, 99-127.
- (6) Green R.M. and Hartles R.L. 1966: The effect of differing high carbohydrates diets on dental caries in the 35 albino rat. Br. J. Nutr. 20, 317-323.

TEST 3

This test was to determine the effect of protein on the adsorption of the bacterium Streptococcus mutans to hydroxyapatite.

Materials and Methods

5 Hydroxyapatite discs were prepared by pressing 150 mg of hydroxyapatite (Bio-Gel HTP, Biorad Laboratories) for 5 min under 5 tons of pressure in a KBr press. The discs were hydrated then incubated with either various protein solutions or imidazole buffer (0.05M pH 7.0, containing 0.025 M NaCl). The adherence of ³H-labelled S.mutans PK1 cells was studied by incubating the pretreated discs with ³H-thymidine labelled cells (10⁹ cells/ml) suspended in the imidazole buffer. The protein solutions used were all 5 mg/ml in imidazole buffer. The proteins and polypeptides studied were α_s-casein, β-casein, κ-casein, phosphovitin, bovine serum albumin, histone III, 15 histone VIII, α-lactalbumin, β-lactoglobulin, poly-l-lysine and poly-l-glutamate. The caseins were prepared by selective precipitation and the other proteins were 20 purchased from Sigma Chemical Co., Missouri, U.S.A.

Results

The effect of pretreating hydroxyapatite discs with various protein solutions on the adherence of S.mutans cells is shown in Table 5.

25 Table 5. Effect of protein on S. mutans adherence to hydroxyapatite.

Proteins	Type	Number of S.mutans cells adsorbed (x10 ⁷)
Control	-	1.9±0.6 ^a
30 α _{s1} -casein	acidic phosphoprotein	0.5±0.3
β-casein	acidic phosphoprotein	0.4±0.4
κ-casein	acidic phosphoprotein	0.6±0.1
phosvitin	acidic phosphoprotein	0.5±0.1
BSA	acidic protein	0.6±0.1
35 histone III	basic protein	2.2±0.9
histone VIII	basic protein	2.6±0.9

- 11 -

Proteins	Type	Number of <i>S. mutans</i> cells adsorbed ($\times 10^7$)
-lactalbumin	acidic protein	1.0 \pm 0.4
-lactoglobulin	acidic protein	0.9 \pm 0.4
5 poly-lysine	basic polypeptide	3.8 \pm 1.1
poly-glutamate	acidic polypeptide	0.5 \pm 0.1

a means \pm SD, n=8

Conclusion

10 All the acidic proteins and polypeptides (especially the phosphoproteins) caused a reduction in bacterial adherence to hydroxyapatite. However, the basic proteins and polypeptides either had no effect or enhanced bacterial adherence to hydroxyapatite.

15 Having regard to the successful results obtained from using the above tests Applicants have formulated various compositions in accordance with this invention as follows. In general, the compositions contain from 0.5-20% by weight of protein or polypeptide.

20 Example 1. Flour: In a device for mixing dry substances, 1% by weight of powdered sodium caseinate was blended with flour.

25 Example 2. Cereal: A breakfast cereal was sprayed with a solution of calcium caseinate in water. The cereal flakes were then dried to produce a finished product containing 1% calcium caseinate.

Example 3. Bread: 2% by weight of calcium caseinate was added to the flour during the mixing of ingredients for the manufacture of bread.

30 Example 4. Cake mix: 1% by weight of calcium caseinate was added to the dry ingredients in the preparation of a cake mix.

Example 5. Confectionery: In the preparation of confectionery 2% by weight of calcium caseinate was added to the final mixture.

35 Example 5. Biscuit: In the preparation of a biscuit mixture 5% by weight of calcium caseinate was added to the other dry ingredients during mixing.

Example 7. Beverage: A beverage was prepared in which 1% weight of calcium caseinate had been dissolved.

Example 8. Tablet: A tablet was made containing 10% by weight of calcium caseinate together with excipients being flavouring matter and binding material.

In preparation of a typical dentifrice within the scope of this invention, the requisite salt and salts of the selected protein or polypeptide are incorporated into dentifrice compositions in any suitable manner depending on whether a powder, paste or liquid preparation is to be produced. For this purpose appropriate preparations of surface-active agents, binders, flavouring materials and other excipients required to achieve the required form of dentifrice are added.

The invention is further illustrated by the following examples: Example 9. Tooth paste: A toothpaste was prepared having the following composition:-

Calcium caseinate	5.0% by weight
Gum tragacanth	1.0% " "
Saccharin	0.1% " "
Glycerin (B.P.)	20.0% " "
Sodium lauryl sulphate	1.0% " "
Methyl parahydroxy benzoate	0.1% " "
Flavouring and colouring	1.0% " "
Dibasic calcium phosphate	35.0% " "
Water	36.8% "

Example 10. Toothpaste: A preparation as set out in example 9 was repeated but with the addition of 2% sodium fluoride in a suitable form.

Example 11. Toothpaste: A preparation as set out in example 9 was repeated but with the addition of 0.4% stannous fluoride in a suitable form

Example 12. Toothpaste: A preparation as set out in example 9 was repeated but with the addition of 0.1% mono sodium fluoro-phosphate in a suitable form.

Example 13. Tooth powder: The following preparation was made:-

5	Calcium caseinate	5.0% by weight
	Soluble saccharin	0.1% " "
	Colour agent	Trace " "
	Dibasic calcium phosphate	94.1% " "

Example 14. Tooth powder: A preparation as set out in example 13 was made but with the addition of 1% mono sodium fluorophosphate in a suitable form.

Example 15. Liquid dentifrice: A preparation was made consisting of:-

15	Sodium alginate	1.5% by weight
	Calcium caseinate	5.0% " "
	Sodium lauryl sulphate	1.0% " "
	Flavouring	Trace " "
	Colouring	Trace " "
	Water	92.5% " "

pH adjusted to 7.0

Example 16. Liquid dentifrice: As for example 15 but with 0.5% sodium fluoride added.

Example 17. Mouthwash: The following preparation was made:-

25	Sodium caseinate	2.0% by weight
	Sodium fluoride	0.5% " "
	Flavouring	Trace " "
	Colouring	Trace " "
	Water	97.5% " "

In the above, casein was used principally because of economics but in lieu phosphitin or other material might be used.

The claims form part of the disclosure of this specification.

Modifications and adaptations may be made to the above described without departing from the spirit and scope of this invention which includes every novel feature and combination of features disclosed herein.



1. An orally ingestible composition containing a caries and gingivitis inhibiting amount of a protein or a polypeptide or a salt thereof.
2. A composition as claimed in claim 1, wherein the protein or polypeptide is a phosphoprotein or a polyphosphopeptide.
3. A composition as claimed in claim 1, wherein the protein or polypeptide is an acidic phosphoprotein or polypeptide.
4. A composition as claimed in claim 1, wherein the protein or polypeptide contains the amino acid sequence (X-Y-Z) where X and Z are an phosphoserine, phosphothreonine, phosphotyrosine, glutamate or asparate and Y is any amino acid.
5. A composition as claimed in claim 4, wherein the protein or polypeptide contains a plurality of units each having the amino acid sequence (X-Y-Z) where X, Y and Z are as stated in claim 4.
6. A composition as claimed in claim 5, wherein the protein or polypeptide contains a group of formula $(X-Y-Z)_n$ where X, Y and Z are as stated in claim 4 and n is 1 or more.
7. A composition as claimed in claim 6, wherein n is 3 or more.
8. A composition as claimed in any one of claims 4-7, wherein X and Z are phosphoserine.
9. A composition as claimed in any one of claims 4-8, wherein the protein or polypeptide is a polyphosphoserine.
10. A composition as in claim 9, wherein the phosphate groups of the polyphosphoserine are spaced at about 6.88 Angstrom Units spacing.
11. A composition as claimed in any preceding claim, wherein the protein is a casein.
12. A composition as claimed in claim 11, wherein the protein is alpha-s-casein.

- 15 -

13. A composition as claimed in any one of claims 1-10, wherein the protein is phosvitin.

14. A composition as claimed in any preceding claim, wherein the protein or polypeptide is in solution.

5 15. A composition as claimed in any preceding claim, wherein the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 45 nmol/min under the test conditions defined herein.

10 16. A composition as claimed in any one of claims 1-14, wherein the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 80 nmol/min under the test conditions defined herein.

15 17. A composition as claimed in any one of claims 1-14, wherein the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 90 nmol/min under the test conditions defined herein.

20 18. A composition as claimed in any one of claims 1-14, wherein the protein or polypeptide is one exhibiting a reduction in calcium dissolution rate of at least 95 nmol/min under the test conditions defined herein.

19. A composition as claimed in any preceding claim, wherein the protein or polypeptide is present as up to 10% by weight.

25 20. A composition as claimed in any preceding claim, wherein the protein or polypeptide is present as up to 5% by weight.

21. A composition as claimed in any preceding claim, wherein the protein or polypeptide is present as up to 2% by weight.

30 22. A composition as claimed in any preceding claim, and additionally containing urea.

23. A composition as claimed in any preceding claim, in the form of a dentifrice mouthwash, tablet, lozenge or capsule.

35 24. A composition as claimed in any one of claims 1-22 in the form of a foodstuff.

SUBSTITUTE SHEET

1101



25. A composition as claimed in any one of claims 1-22 in the form of confectionery.


26. An orally ingestible composition substantially as hereinbefore described with reference to any one of the Examples.

27. A process of inhibiting dental caries and/or tooth erosion and/or gingivitis comprising applying to the teeth with a carrier a cariostatic and/or anti-gingivitis agent selected from a protein or a polypeptide or a salt thereof.

28. A process of inhibiting dental caries and/or tooth erosion and/or gingivitis comprising applying to the teeth a composition in accordance with any one of claims 1-26.

29. A composition as claimed in any preceding claim wherein the protein or polypeptide is present as 1% by weight or greater.

30. The articles, things, parts, elements steps, features, methods, processes, compounds and compositions referred to or indicated in the specification and/or claims of the application individually or collectively, and any and all combinations of any two or more of such.



INTERNATIONAL SEARCH REPORT

International Application No PCT/AU82/00022

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ¹		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int. Cl. ³ A61K 7/16, 37/16		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
IPC	A61K 7/16, 37/00 to 37/18	
US Cl	424.177	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁵		
AU:IPC as above plus A61K 7/30, 31/71, C07F 9/10, C07G 7/00; Australian Classification 87-16-22		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁵ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
X	US,A, 4225579 (KLEINBERG) 30 September 1980 (30.09.80)	
X	GB,A, 1579974 (KLEINBERG) 26 November 1980 (26.11.80) (&AU,B, 515028 & DE,A, 2727150)	
A	GB,A, 1452263 (NATIONAL PATENT DEVELOPMENT CO) 13 October 1976 (13.10.76)	
X	US,A, 2853386 (HUGHES) 23 September 1958 (23.09.58)	
A	US,A, 3966915 (CAPRINO) 29 June 1976 (29.06.76)	
A	US,A, 4230695 (ECKER) 28 October 1980 (28.10.80)	
A	US,A, 2783221 (NITSCHKE) 26 February 1957 (26.02.57)	
A	US,A, 4165312 (SCHOENENBERGER) 21 August 1979 (21.08.79)	
X	JOURNAL OF DENTAL RESEARCH, Volume 56, No. 8, issued 1977 August, H.G. Huxley, "Cariogenicity in Rats of a Diet Containing Lactalbumin or Casein as the Protein Source," see page 900.	
A	DE,A, 2152620 (SCHWEIZERISCHE FERMENT AG) 27 April 1972 (27.04.72)	
A	DE,A, 2712228 (LEIDERER) 28 September 1978 (28.09.78)	
P T O		
<p>[*] Special categories of cited documents: ¹⁵</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>¹⁷ later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ⁹	Date of Mailing of this International Search Report ⁸	
11 May 1982 (11.05.82)	17 May 1982 (17.05.82)	
International Searching Authority ¹	Signature of Authorized Officer ¹⁰	
AUSTRALIAN PATENT OFFICE	A.S. MOORE <i>A S Moore</i>	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

A	CH,A, 541967 (PACIFIC BIOCHEMICAL INC) 30 September 1973 (30.09.73)
X	GB,A, 1505513 (STOLLE RESEARCH AND DEVELOPMENT CORP) 30 March 1978 (30.03.78)

V. OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹⁰

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. Claim numbers _____, because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹², specifically:

VI. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ¹¹

This International Searching Authority found multiple inventions in this international application as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

The additional search fees were accompanied by applicant's protest.

No protest accompanied the payment of additional search fees.