




(86) Date de dépôt PCT/PCT Filing Date: 2001/04/13
(87) Date publication PCT/PCT Publication Date: 2001/10/25
(45) Date de délivrance/Issue Date: 2012/01/24
(85) Entrée phase nationale/National Entry: 2002/10/11
(86) N° demande PCT/PCT Application No.: US 2001/011967
(87) N° publication PCT/PCT Publication No.: 2001/079517
(30) Priorité/Priority: 2000/04/13 (US09/548,971)

(51) Cl.Int./Int.Cl. *C12N 15/82* (2006.01),
A01H 5/00 (2006.01), *C07K 14/415* (2006.01),
C12N 15/29 (2006.01)
(72) Inventeurs/Inventors:
LILJEGREN, SARAH, US;
YANOFSKY, MARTIN F., US
(73) Propriétaire/Owner:
THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
US
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : MAITRISE DE LA DEHISCENCE DU FRUIT DANS DES ARABIDOPSIS AU MOYEN DE GENES 1
INDEHISCENTS

(54) Title: CONTROL OF FRUIT DEHISCENCE IN ARABIDOPSIS BY INDEHISCENT1 GENES



(57) Abrégé/Abstract:

The present invention provides methods and compositions that modulate fruit dehiscence in plants.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/79517 A3(51) International Patent Classification⁷: **C12N 15/82**,
15/29, C07K 14/415, A01H 5/00(74) Agents: **HINSCH, Matthew, E.** et al.; Townsend and
Townsend and Crew LLP, Two Embarcadero Center, Eighth
Floor, San Francisco, CA 94111-3834 (US).

(21) International Application Number: PCT/US01/11967

(22) International Filing Date: 13 April 2001 (13.04.2001)

(25) Filing Language: English

(26) Publication Language: English


(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(30) Priority Data:
09/548,971 13 April 2000 (13.04.2000) US(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant: **THE REGENTS OF THE UNIVERSITY
OF CALIFORNIA** [US/US]; University of California,
San Diego Technology Transfer and Intellectual Property
Services, 9500 Gilman Drive, MC 0910, La Jolla, CA
92093-0910 (US).**Published:**

— with international search report

(72) Inventors: **LILJEGREN, Sarah**; 3268 Caminito Ameca,
La Jolla, CA 92037 (US). **YANOFSKY, Martin, F.**; 5039
Manor Ridge Lane, San Diego, CA 92130 (US).(88) Date of publication of the international search report:
30 May 2002

[Continued on next page]

(54) Title: CONTROL OF FRUIT DEHISCENCE IN ARABIDOPSIS BY INDEHISCENT GENES



(57) Abstract: The present invention provides methods and compositions that modulate fruit dehiscence in plants.



WO 01/79517 A3

WO 01/79517 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

CONTROL OF FRUIT DEHISCENCE IN *ARABIDOPSIS* BY *INDEHISCENT1* GENES

5 CROSS-REFERENCES TO RELATED APPLICATIONS

This application is a continuation-in-part of United States issued Patent No. 6,410,826. This application is also related to United States issued Patent No. 6,288,305.

10

15 STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

This invention was made with Government support under National Science Foundation Grant number IBN-9985530. The Government has certain rights in the invention.

20 FIELD OF THE INVENTION

This invention relates plant genetic engineering. In particular, the invention relates to methods and compositions that modulate fruit dehiscence in plants.

BACKGROUND OF THE INVENTION

25 Rapeseed is one of the most important oilseed crops after soybeans and cottonseed, representing 10% of the world oilseed production in 1990. Rapeseed contains 40% oil, which is pressed from the seed, leaving a high-protein seed meal of value for animal feed and nitrogen fertilizer. Rapeseed oil, also known as canola oil, is a valuable product, representing the fourth most commonly traded vegetable oil in the world.

30 Unfortunately, the yield of seed from rapeseed and related plants is limited by pod dehiscence, which is a process that occurs late in fruit development whereby the pod is opened and the enclosed seeds released. Degradation and separation of cell walls along a discrete layer of cells dividing the two halves of the pod, termed the "dehiscence

zone," result in separation of the two halves of the pod and release of the contained seeds. The dehiscence zone is a region of only one to three cells in width that extends along the entire length of the valve/replum boundary (Meakin and Roberts, *J. Exp. Botany* 41:995-1002 (1990)). As the cells in the dehiscence zone separate from one another, the valves
 5 detach from the replum, allowing seeds to be dispersed. Seed "shattering," whereby seeds are prematurely shed through dehiscence before the crop can be harvested, is a significant problem faced by commercial seed producers and represents a loss of income to the industry. Adverse weather conditions can exacerbate the process of dehiscence, resulting in greater than 50% loss of seed yield.

10 The fruit, a complex structure unique to flowering plants, mediates the maturation and dispersal of seeds. In most flowering plants, the fruit consists of the pericarp, which is derived from the ovary wall, and the seeds, which develop from fertilized ovules. *Arabidopsis*, which is typical of the more than 3000 species of the *Brassicaceae*, produces fruit in which the two carpel valves (ovary walls) are joined to
 15 the replum, a visible suture that divides the two carpels.

The plant hormone ethylene is produced by developing seeds and appears to be an important regulator of the dehiscence process. One line of evidence supporting a
 role for ethylene in regulation of dehiscence comes from studies of fruit ripening, which, like fruit dehiscence, is a process involving the breakdown of cell wall material. In fruit
 20 ripening, ethylene acts in part by activating cell wall degrading enzymes such as polygalacturonase (Theologis *et al.*, *Develop. Genetics* 14:282-295 (1993)). Moreover, in genetically modified tomato plants in which the ethylene response is blocked, such as transgenic tomato plants expressing antisense polygalacturonase, there is a significant delay in fruit ripening (Lanahan *et al.*, *The Plant Cell* 6:521-530 (1994); Smith *et al.*,
 25 *Nature* 334:724-726 (1988)).

In dehiscence, ultrastructural changes that culminate in degradation of the middle lamella of dehiscence zone cell walls weaken rapeseed pods and eventually lead to pod shatter. As in fruit ripening, hydrolytic enzymes including polygalacturonases play a role in this programmed breakdown. For example, in oilseed rape, a specific endo-
 30 polygalacturonase, RDPG1, is upregulated and expressed exclusively in the dehiscence zone late in pod development (Petersen *et al.*, *Plant Mol. Biol.* 31:517-527 (1996)).

Ethylene may regulate the activity of hydrolytic enzymes involved in the process of dehiscence as it does in fruit ripening (Meakin and Roberts, *J. Exp. Botany* 41:1003-1011 (1990)).

Yet, until now, the proteins that control the process of dehiscence, such as those regulating the relevant hydrolytic enzymes, have eluded identification.

Attempts to solve the problem of pod shatter and early fruit dehiscence over the past 20 years have focused on the breeding of shatter-resistant varieties.

5 However, these plant hybrids are frequently sterile and lose favorable characteristics that must be regained by backcrossing, which is both time-consuming and laborious. Other strategies to alleviate pod shattering include the use of chemicals such as pod sealants or mechanical techniques such as swathing to reduce wind-stimulated shattering. To date, however, a simple method for producing genetically modified plants that do not open and
10 release their seeds prematurely has not been described.

Thus, a need exists for identifying genes that regulate the dehiscence process and for developing genetically modified plant varieties in which the natural seed dispersal process is delayed. The present invention satisfies this need and provides related advantages as well.

15

SUMMARY OF THE INVENTION

This invention provides an isolated nucleic acid comprising an IND1 polynucleotide sequence encoding an IND1 polypeptide at least 70% identical to SEQ ID
20 NO:2. The isolated nucleic acid, for instance, can comprise a polynucleotide that encodes SEQ ID NO:2. In another embodiment, the isolated nucleic acid, for instance, can comprise positions from about 2765 to about 3361 of SEQ ID NO:1. For example, the nucleic acid can comprise SEQ ID NO:1.

The present invention also provides an expression cassette comprising a
25 promoter operably linked to an IND1 polynucleotide sequence, or complement thereof, encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO:1. The expression cassette, for instance, can comprise a polynucleotide that encodes SEQ ID NO:2. In another embodiment, the expression cassette, for instance, can comprise positions from about 2765 to about 3361 of SEQ ID NO:1. For example, the expression
30 cassette can comprise SEQ ID NO:1. In some embodiments, the expression cassette comprises a promoter. The promoter, for instance, can be constitutive or tissue specific. In one aspect of the invention, the promoter is a dehiscence zone specific promoter. In another aspect, the promoter can comprise positions from about 1 to about 2764 or positions from about 3362 to about 3856 of SEQ ID NO:1.

The present invention also provides a plant comprising a recombinant expression cassette comprising a promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO:1. In one aspect, the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the antisense orientation. In another aspect, the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the sense orientation. The polynucleotide sequence can further comprise a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation. In some embodiments, the plant of the invention has reduced lignification in valve margin cells. In other embodiments, the promoter is a dehiscence zone-selective regulatory element. In some of these embodiments, the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1.

The present invention also provides a method of delaying fruit dehiscence in a plant comprising suppressing expression of an IND1 nucleic acid in the plant by introducing into the plant a recombinant expression cassette comprising a promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at least 70% identical to SEQ ID NO:2. In some embodiments, the IND1 polypeptide is SEQ ID NO:2. In other embodiments, the IND1 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO:1. In one aspect of the invention, the IND1 polynucleotide comprises SEQ ID NO:1. The method can include a polynucleotide sequence encoding the IND1 polypeptide operably linked to the promoter in the antisense orientation. In another embodiment, the promoter is linked to the promoter in the sense orientation. The polynucleotide sequence can further comprise a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation. In some embodiments, the method results in a plant with reduced lignification in valve margin cells. In other embodiments, the promoter is a dehiscence zone-selective regulatory element. In some of these embodiments, the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In one aspect, the recombinant expression cassette is introduced into the plant with *Agrobacterium*.

The present invention also provides a method of delaying fruit dehiscence in a plant comprising suppressing expression of an IND1 gene in the plant by introducing into the plant a recombinant expression cassette comprising a polynucleotide sequence at

least about 70% identical to positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In one aspect of the invention, the polynucleotide sequence comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In some aspects, lignification is reduced in valve margin cells.

5

DEFINITIONS

The phrase "nucleic acid" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read through by a polymerase and do not alter expression of a polypeptide encoded by that nucleic acid.

10

The phrase "polynucleotide sequence" or "nucleic acid sequence" includes both the sense and antisense strands of a nucleic acid as either individual single strands or in the duplex. It includes, but is not limited to, self-replicating plasmids, chromosomal sequences, and infectious polymers of DNA or RNA.

15

The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a specific host cell.

20

The term "promoter" or "regulatory element" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Such promoters need not be of plant origin, for example, promoters derived from plant viruses, such as the CaMV35S promoter, can be used in the present invention.

25

As used herein, the term "dehiscence zone-selective regulatory element" refers to a nucleotide sequence that, when operatively linked to a nucleic acid molecule, confers selective expression upon the operatively linked nucleic acid molecule in a limited number of plant tissues, including the valve margin or dehiscence zone. The valve margin is the future site of the dehiscence zone and encompasses the margins of the outer replum as well as valve cells adjacent to the outer replum. The dehiscence zone,

30

which develops in the region of the valve margin, refers to the group of cells that separate during the process of dehiscence, allowing valves to come apart from the replum and the enclosed seeds to be released. Thus, a dehiscence zone-selective regulatory element, as defined herein, confers selective expression in the mature dehiscence zone, or confers selective expression in the valve margin, which marks the future site of the dehiscence zone.-

A dehiscence zone-selective regulatory element can confer specific expression exclusively in cells of the valve margin or dehiscence zone or can confer selective expression in a limited number of plant cell types including cells of the valve margin or dehiscence zone. A *SHATTERPROOF1* or *SHATTERPROOF2* (*SHP1* and *SHP2*, also designated as *AGL1* and *AGL5*, respectively) regulatory element, for example, which confers selective expression in ovules and placenta as well as in the dehiscence zone, is a dehiscence zone-selective regulatory element as defined herein. Similarly, an *IND1* regulatory element also confers selective expression in the dehiscence zone. A dehiscence zone-selective regulatory element generally is distinguished from other regulatory elements by conferring selective expression in the valve margin or dehiscence zone without conferring expression throughout the adjacent carpel valves.

It is understood that limited modifications can be made without destroying the biological function of a regulatory element and that such limited modifications can result in dehiscence zone-selective regulatory elements that have substantially equivalent or enhanced function as compared to a wild type *IND1* regulatory element. These modifications can be deliberate, as through site-directed mutagenesis, or can be accidental such as through mutation in hosts harboring the regulatory element. All such modified nucleotide sequences are included in the definition of a dehiscence zone-selective regulatory element as long as the ability to confer selective expression in the valve margin or dehiscence zone is substantially retained.

The term "plant" includes whole plants, shoot vegetative organs/structures (e.g. leaves, stems and tubers), roots, flowers and floral organs/structures (e.g. bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm, and seed coat) and fruit (the mature ovary), plant tissue (e.g. vascular tissue, ground tissue, and the like) and cells (e.g. guard cells, egg cells, trichomes and the like), and progeny of same. The class of plants that can be used in the method of the invention is generally as broad as the class of higher and lower plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants),

gymnosperms, ferns, and multicellular algae. It includes plants of a variety of ploidy levels, including aneuploid, polyploid, diploid, haploid and hemizygous.

The term "seed plant" means an angiosperm or gymnosperm. An angiosperm is a seed-bearing plant whose seeds are borne in a mature ovary (fruit). An angiosperm commonly is recognized as a flowering plant. Angiosperms are divided into two broad classes based on the number of cotyledons, which are seed leaves that generally store or absorb food. Thus, a monocotyledonous angiosperm is an angiosperm having a single cotyledon, whereas a dicotyledonous angiosperm is an angiosperm having two cotyledons. A variety of angiosperms are known including, for example, oilseed plants, leguminous plants, fruit-bearing plants, ornamental flowers, cereal plants and hardwood trees, which general classes are not necessarily exclusive. The skilled artisan will recognize that the methods of the invention can be practiced using these or other angiosperms, as desired. A gymnosperm is a seed-bearing plant with seeds not enclosed in an ovary.

The phrase "host cell" refers to a cell from any organism. Preferred host cells are derived from plants, bacteria, yeast, fungi, insects or other animals. Methods for introducing polynucleotide sequences into various types of host cells are well known in the art.

The term "delayed," as used herein in reference to the timing of seed dispersal in a fruit produced by a non-naturally occurring plant of the invention, means a significantly later time of seed dispersal as compared to the time seeds normally are dispersed from a corresponding plant at the same developmental stage expressing naturally-occurring levels of IND1. Thus, the term "delayed" is used broadly to encompass both seed dispersal that is significantly postponed as compared to the seed dispersal in a corresponding plant, and to seed dispersal that is completely precluded, such that fruits never release their seeds unless there is human or other intervention.

It is recognized that there can be natural variation of the time of seed dispersal within a plant species or variety. However, a "delay" in the time of seed dispersal in a non-naturally occurring plant of the invention readily can be identified by sampling a population of the non-naturally occurring plants and determining that the normal distribution of seed dispersal times is significantly later, on average, than the normal distribution of seed dispersal times in a population of the corresponding plant species or variety that does not contain an exogenous IND1 polynucleotide. Thus, production of non-naturally occurring plants of the invention provides a means to skew

the normal distribution of the time of seed dispersal from pollination, such that seeds are dispersed, on average, at least about 1%, 2%, 5%, 10%, 30%, 50%, 100%, 200% or 500% later than in the corresponding plant species that does not contain an exogenous nucleic acid molecule encoding an *IND1* gene product.

5 The term "suppressed" or "decreased" encompasses the absence of *IND1* protein in a plant, as well as protein expression that is present but reduced as compared to the level of *IND1* protein expression in a wild type plant. Furthermore, the term suppressed refers to *IND1* protein expression that is reduced throughout the entire domain of *IND1* expression, or to expression that is reduced in some part of the *IND1* expression domain, provided that the resulting plant is characterized by delayed seed dispersal. The
10 term "suppressed" also encompasses an amount of *IND1* protein that is equivalent to wild type *IND1* expression, but where the *IND1* protein has a reduced level of activity. As discussed above, *IND1* each contain a conserved an basic HLH domain; point mutations or gross deletions within the HLH domain that reduce the DNA-binding activity of *IND1*
15 can reduce or destroy the activity of *IND1* and, therefore, "suppress" *IND1* expression as defined herein. One skilled in the art will recognize that, preferably, *IND1* expression is essentially absent in the valve margin of a plant or the *IND1* protein is essentially non-functional.

"Increased" or "enhanced" *IND1* activity or expression of a *IND1* gene
20 refers to an augmented change in *IND1* activity. Examples of such increased activity or expression include the following. *IND1* activity or expression of the *IND1* gene is increased above the level of that in wild-type, non-transgenic control plants (i.e. the quantity of *IND1* activity or expression of the *IND1* gene is increased). *IND1* activity or expression of the *IND1* gene is in an organ, tissue or cell where it is not normally detected
25 in wild-type, non-transgenic control plants (i.e. spatial distribution of *IND1* activity or expression of the *IND1* gene is increased). *IND1* activity or expression is increased when *IND1* activity or expression of the *IND1* gene is present in an organ, tissue or cell for a longer period than in a wild-type, non-transgenic controls (i.e. duration of *IND1* activity or expression of the *IND1* gene is increased).

30 A polynucleotide sequence is "heterologous to" a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified by human action from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from

that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants.

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant, or a predecessor generation of the plant, by any means other than by a sexual cross. An exogenous nucleic acid molecule can have a naturally occurring or non-naturally occurring nucleotide sequence and can be a heterologous nucleic acid molecule derived from a different plant species than the plant into which the nucleic acid molecule is introduced or can be a nucleic acid molecule derived from the same plant species as the plant into which it is introduced. Examples of means by which this can be accomplished are described below, and include *Agrobacterium*-mediated transformation, biolistic methods, electroporation, in planta techniques, and the like.

An "*IND1* polynucleotide" is a nucleic acid sequence comprising (or consisting of) a coding region of about 50 to about 4000 nucleotides, sometimes from about 100 to about 3000 nucleotides and sometimes from about 200 to about 600 nucleotides, which hybridizes to SEQ ID NO:1 under stringent conditions (as defined below), or which encodes an *IND1* polypeptide or fragment of at least 15 amino acids thereof. *IND1* polynucleotides can also be identified by their ability to hybridize under low stringency conditions (e.g., $T_m \sim 40^\circ\text{C}$) to nucleic acid probes having the sequence of SEQ ID NO:1. SEQ ID NO:1 is an example of a *IND1* polynucleotide.

A "promoter from a *IND1* gene" or "*IND1* promoter" will typically be about 500 to about 3000 nucleotides in length, usually from about 750 to 2750. Exemplary promoter sequences are shown as SEQ ID NO:3 and SEQ ID NO:4. SEQ ID NO:3 represents the 5' untranslated region of the *IND1* and SEQ ID NO:4 represents the 3' untranslated region of *IND1*. A *IND1* promoter can also be identified by its ability to direct expression in the valve margin of fruit. In particular, the *Ind1* promoter directs expression at the valve margin of developing gynoecium just prior to fertilization (stage 13) through the maturation of the fruit (stage 17). The promoter does not provide significant expression in leaf tissue.

An "*IND1* polypeptide" is a sequence of about 50 to about 200, sometimes 100 to 190, and preferably 198 amino acid residues encoded by a *IND1* polynucleotide. *IND1* polypeptides are characterized by the presence of a basic helix-loop-helix (HLH) domain which bind specific polynucleotide sequences. For instance amino acid residues ISDDPQTVVARRRRERISEKIRILKRIVPGGAKMDTASMLDEAIRYTKFLK (SEQ

ID NO: 7) represent the HLH domain of the polypeptide shown in SEQ ID NO:2. The
HLH domain

is known in the art and is shared by other transcription factors including uncharacterized sequences represented by GenBank accession number E1283552 and 2262147 and the gene product, PIF3 (Ni *et al.* *Cell* 95:657 (1998)). The HLH domain of IND1 is therefore a DNA binding domain.

5 As used herein, a homolog of a particular IND1 gene (e.g., SEQ ID NO:1) is a second gene in the same plant type or in a different plant type, which has a polynucleotide sequence of at least 50 contiguous nucleotides which are substantially identical (determined as described below) to a sequence in the first gene. It is believed that, in general, homologs share a common evolutionary past.

10 A "polynucleotide sequence from" a particular gene is a subsequence or full length polynucleotide sequence of an IND1 gene which, when present in a transgenic plant, has the desired effect. For example, one effect is inhibition of expression of the endogenous gene driving expression of an heterologous polynucleotide. A full length sequence of a particular gene disclosed here may contain about 95%, usually at least
15 about 98% of an entire sequence shown in the Sequence Listing, below.

 The term "reproductive tissues" as used herein includes fruit, ovules, seeds, pollen, pistils, flowers, or any embryonic tissue.

 An "expression cassette" refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of a RNA or
20 polypeptide, respectively. Antisense or sense constructs that are not or cannot be translated are expressly included by this definition.

 In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical and may be "substantially identical" to a
25 sequence of the gene from which it was derived. As explained below, these variants are specifically covered by this term.

 In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same
30 polypeptide. These variants are specifically covered by the term "polynucleotide sequence from" a particular valve-margin gene, such as *IND1*. In addition, the term specifically includes sequences (e.g., full length sequences) substantially identical (determined as described below) with a IND1 gene sequence and that encode proteins that retain the function of a IND1 polypeptide.

In the case of polynucleotides used to inhibit expression of an endogenous gene, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence will typically be at least substantially identical (as determined below) to the target endogenous sequence.

5 Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

10 Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics
15 Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions
20 (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of
25 comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 25% sequence identity. Alternatively, percent identity can be any integer from 25% to 100%. More preferred embodiments include at least: 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%,
30 75%, 80%, 85%, 90%, 95%, or 99%. compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. Accordingly, IND1 sequences of the invention include nucleic acid sequences that have substantial identity to SEQ ID NO:1. IND1 sequences of the invention also include polypeptide sequences having substantial identity to SEQ ID NO:2. One of skill

will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of
5 at least 40%. Preferred percent identity of polypeptides can be any integer from 40% to 100%. More preferred embodiments include at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. Most preferred embodiments include 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74% and 75%. Polypeptides which are "substantially similar" share sequences as noted above except that
10 residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-
15 containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine,
20 lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than
25 the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C.

30 In the present invention, mRNA encoded by IND1 genes of the invention can be identified in Northern blots under stringent conditions using cDNAs of the invention or fragments of at least about 100 nucleotides. For the purposes of this disclosure, stringent conditions for such RNA-DNA hybridizations are those which include at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions.

Genomic DNA or cDNA comprising genes of the invention can be identified using the same cDNAs (or fragments of at least about 100 nucleotides) under stringent conditions, which for purposes of this disclosure, include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the cell types of the Arabidopsis fruit at maturity.

DETAILED DESCRIPTION

I. Introduction

The present invention provides methods of modulating fruit development in plants. In particular, the invention provides methods of delaying or preventing fruit dehiscence by suppressing expression of an bHLH gene such as IND1 in a plant. The invention also provides transgenic plants comprising various polynucleotides encoding an bHLH polypeptide such as IND1.

The present invention relates to the previous discovery that an *agl1 agl3* double mutant plant has a delayed seed dispersal phenotype (Liljegren *et al.*, *Nature* 404:766-770 (2000)). Loss-of-function mutations in the *SHP1* and *SHP2* genes were produced by disruptive T-DNA insertion and homologous recombination. In the resulting *shp1 shp2* double mutant plants, the dehiscence zone failed to develop normally, and the mature fruits did not undergo dehiscence. Thus, *SHP1* or *SHP2* gene expression is required for development of the dehiscence zone. These results indicate that *SHP1* and *SHP2* regulate pod dehiscence and that manipulation of *SHP1* and *SHP2* expression can allow the process of pod shatter to be controlled.

The present invention provides evidence that *IND1* is regulated by *SHP1* and *SHP2* and that expression of *IND1* modulates fruit dehiscence. The present invention also provides for methods of delaying fruit dehiscence by suppressing expression of *IND1*.

The Arabidopsis *SHP1* and *SHP2* genes encode MADS box proteins with 85% identity at the amino acid level. The *SHP1* and *SHP2* RNA expression patterns are also strikingly similar. In particular, both RNAs are specifically expressed in flowers, where they accumulate in developing carpels. In particular, strong expression of these

genes is observed in the outer replum along the valve/replum boundary (Ma *et al.*, *supra*, 1991; Savidge *et al.*, *The Plant Cell* 7:721-723 (1995); Flanagan *et al.*, *The Plant Journal* 10:343-353 (1996)). Thus, *SHP1* and *SHP2* are expressed in the valve margin, at least within the cells of the outer replum.

5 Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's
10 specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

II. Isolation of nucleic acids of the invention

15 The isolation of sequences from the genes of the invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction
20 endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs such as *IND1*, mRNA is isolated from embryos and a cDNA library that contains the gene transcripts is prepared from the mRNA.

 The cDNA or genomic library can then be screened using a probe based
25 upon the sequence of a cloned *IND1* gene such as the polynucleotides disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

 Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction
30 (PCR) technology to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying genes such as *IND1* from plant tissues are generated from comparisons of the sequences provided herein. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego

5 (1990). Appropriate primers for amplification of the genomic region of *IND1* or the *IND1* cDNA include the following primer pairs: 5'-gatgaaaatggaaaatggtatgtata-3' and 5'-gttcatcaggggtgggagttgtg-3'. The amplification conditions are typically as follows. Reaction components: 10 mM Tris-HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dATP, 200 μ M dCTP, 200 μ M dGTP, 200

10 μ M dTTP, 0.4 μ M primers, and 100 units per ml Taq polymerase. Program: 96 C for 3 min., 30 cycles of 96 C for 45 sec., 50 C for 60 sec., 72 for 60 sec, followed by 72 C for 5 min.

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, Cold Spring Harbor

15 Symp. Quant. Biol. 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

20 The genus of *IND1* nucleic acid sequences of the invention includes genes and gene products identified and characterized by analysis using nucleic acid sequences of the invention, including SEQ ID NO:1 and protein sequences of the invention, including SEQ ID NO:2. *IND1* sequences of the invention include nucleic acid sequences having substantial identity to SEQ ID NO:1. *IND1* sequences of the invention

25 also include polypeptide sequences having substantial identity to SEQ ID NO:2.

III. Use of nucleic acids of the invention

30 A. Use of nucleic acids of the invention to inhibit or suppress gene expression

The invention provides methods of modulating fruit dehiscence in a plant by introducing into a plant a recombinant expression cassette comprising a regulatory element operably linked to a HLH polynucleotide such as *IND1*. The invention also provides methods for delaying seed dispersal in a plant by suppressing expression of a

nucleic acid molecule encoding an IND1 gene product. In a transgenic plant of the invention, a nucleic acid molecule, or antisense constructs thereof, encoding an IND1 gene product can be operatively linked to an exogenous regulatory element. The invention provides, for example, a transgenic plant characterized by delayed seed
5 dispersal having an expressed nucleic acid molecule encoding an IND1 gene product, or antisense construct thereof, that is operatively linked to an exogenous constitutive regulatory element. In one embodiment, the invention provides a transgenic plant that is characterized by delayed seed dispersal due to suppression of a nucleic acid molecule encoding an IND1 ortholog. In some preferred embodiments, suppression of IND1
10 expression results in reduced lignification in valve margin cells. *See also*, United States issued Patent No. 6,410,826.

The *IND1* sequences prepared as described herein, can be used to prepare expression cassettes useful in a number of techniques, including inhibiting or suppressing expression. A number of methods can be used to inhibit gene expression in plants. For
15 instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The expression cassette is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense RNA inhibits gene expression by preventing the
20 accumulation of mRNA which encodes the enzyme of interest, see, e.g., Sheehy *et al.*, *Proc. Nat. Acad. Sci. USA*, 85:8805-8809 (1988); Pnueli *et al.*, *The Plant Cell* 6:175-186 (1994); and Hiatt *et al.*, U.S. Patent No. 4,801,340.

The antisense nucleic acid sequence transformed into plants will be substantially identical to at least a portion of the endogenous gene or genes to be
25 repressed. The sequence, however, does not have to be perfectly identical to inhibit expression. Thus, an antisense or sense nucleic acid molecule encoding only a portion of IND1 can be useful for producing a plant in which IND1 expression is suppressed. The vectors of the present invention can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to
30 the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and

homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of at least about 500 nucleotides is especially preferred.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of *IND1* genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs that are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Haseloff *et al. Nature*, 334:585-591 (1988).

Another method of suppression is sense suppression (also known as co-suppression). Introduction of expression cassettes in which a nucleic acid is configured in the sense orientation with respect to the promoter has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes see, Napoli *et al.*, *The Plant Cell* 2:279-289 (1990); Flavell, *Proc. Natl. Acad. Sci., USA* 91:3490-3496 (1994); Kooter and Mol, *Current Opin. Biol.* 4:166-171 (1993); and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

Generally, where inhibition of expression is desired, some transcription of the introduced sequence occurs. The effect may occur where the introduced sequence contains no coding sequence per se, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous

sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with
5 antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence in the expression cassette, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid
10 concurrent production of some plants that are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used.

15 In a preferred embodiment, expression of a nucleic acid of interest can be suppressed by the simultaneous expression of both sense and antisense constructs (Waterhouse *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964 (1998). See also Tabara *et al. Science* 282:430-431 (1998).

One of skill in the art will recognize that using technology based on
20 specific nucleotide sequences (e.g., antisense or sense suppression technology), families of homologous genes can be suppressed with a single sense or antisense transcript. For instance, if a sense or antisense transcript is designed to have a sequence that is conserved among a family of genes, then multiple members of a gene family can be suppressed. Conversely, if the goal is to only suppress one member of a homologous gene family,
25 then the sense or antisense transcript should be targeted to sequences with the most variance between family members.

Another means of inhibiting *IND1* function in a plant is by creation of dominant negative mutations. In this approach, non-functional, mutant *IND1* polypeptides, which retain the ability to interact with wild-type subunits are introduced
30 into a plant. A dominant negative construct also can be used to suppress *IND1* expression in a plant. A dominant negative construct useful in the invention generally contains a portion of the complete *IND1* coding sequence sufficient, for example, for DNA-binding or for a protein-protein interaction such as a homodimeric or heterodimeric protein-protein interaction but lacking the transcriptional activity of the wild type protein. For

example, a carboxy-terminal deletion mutant of AGAMOUS was used as a dominant negative construct to suppress expression of the MADS box gene AGAMOUS (Mizukami et al., Plant Cell 8:831-844 (1996)). One skilled in the art understands that, similarly, a dominant negative IND1 construct can be used to suppress IND1 expression
5 in a plant.

B. Use of nucleic acids of the invention to enhance gene expression

Isolated sequences prepared as described herein can also be used to prepare expression cassettes that enhance or increase endogenous *IND1* gene expression.
10 Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects. Enhanced expression of *IND1* polynucleotides is useful, for example, to produce plants with small fruit.

Any of a number of means well known in the art can be used to increase IND1 activity in plants. Any organ can be targeted, such as shoot vegetative
15 organs/structures (e.g. leaves, stems and tubers), roots, flowers and floral organs/structures (e.g. bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm, and seed coat) and fruit. Alternatively, one or several IND1 genes can be expressed constitutively (e.g., using the CaMV 35S promoter).

One of skill will recognize that the polypeptides encoded by the genes of
20 the invention, like other proteins, have different domains which perform different functions. Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed. As explained above, IND1 polypeptides carry a bHLH domain, which is capable of binding DNA. Thus, without being bound to any particular theory or mechanism, IND1 is likely to act as a transcriptional modulator.

25

C. Modification of endogenous IND1 genes

Methods for introducing genetic mutations into plant genes and selecting plants with desired traits are well known. For instance, seeds or other plant material can be treated with a mutagenic chemical substance, according to standard techniques. Such
30 chemical substances include, but are not limited to, the following: diethyl sulfate, ethylene imine, ethyl methanesulfonate and N-nitroso-N-ethylurea. Alternatively, ionizing radiation from sources such as, X-rays or gamma rays can be used.

Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described for

instance, in Sambrook et al., supra. Hydroxylamine can also be used to introduce single base mutations into the coding region of the gene (Sikorski, *et al.*, (1991). *Meth. Enzymol.* 194: 302-318). For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like.

5 These modifications can be used in a number of combinations to produce the final modified protein chain.

Alternatively, homologous recombination can be used to induce targeted gene modifications by specifically targeting the *IND1* gene *in vivo* (see, generally, Grewal and Klar, *Genetics* 146: 1221-1238 (1997) and Xu *et al.*, *Genes Dev.* 10: 2411-2422 (1996)). Homologous recombination has been demonstrated in plants (Puchta *et al.*, *Experientia* 50: 277-284 (1994), Swoboda *et al.*, *EMBO J.* 13: 484-489 (1994); Offringa *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 7346-7350 (1993); and Kempin *et al.* *Nature* 389:802-803 (1997)).

In applying homologous recombination technology to the genes of the invention, mutations in selected portions of an *IND1* gene sequences (including 5' upstream, 3' downstream, and intragenic regions) such as those disclosed here are made *in vitro* and then introduced into the desired plant using standard techniques. Since the efficiency of homologous recombination is known to be dependent on the vectors used, use of dicistronic gene targeting vectors as described by Mountford *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 4303-4307 (1994); and Vaulont *et al.*, *Transgenic Res.* 4: 247-255 (1995) are conveniently used to increase the efficiency of selecting for altered *IND1* gene expression in transgenic plants. The mutated gene will interact with the target wild-type gene in such a way that homologous recombination and targeted replacement of the wild-type gene will occur in transgenic plant cells, resulting in suppression of *IND1* activity.

25 Alternatively, oligonucleotides composed of a contiguous stretch of RNA and DNA residues in a duplex conformation with double hairpin caps on the ends can be used. The RNA/DNA sequence is designed to align with the sequence of the target *IND1* gene and to contain the desired nucleotide change. Introduction of the chimeric oligonucleotide on an extrachromosomal T-DNA plasmid results in efficient and specific *IND1* gene conversion directed by chimeric molecules in a small number of transformed plant cells. This method is described in Cole-Strauss *et al.*, *Science* 273:1386-1389 (1996) and Yoon *et al.*, *Proc. Natl. Acad. Sci. USA* 93: 2071-2076 (1996).

In other embodiments, the promoters derived from the *IND1* genes of the invention can be used to drive expression of heterologous genes in an valve margin-

specific manner. Suitable structural genes that could be used for this purpose include genes encoding cytotoxic proteins as discussed below.

Typically, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the IND1 genes described here. Sequences characteristic of promoter sequences can be used to identify the promoter. Sequences controlling eukaryotic gene expression have been extensively studied. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing *et al.*, in GENETIC ENGINEERING IN PLANTS, pp.221-227 (Kosage, Meredith and Hollaender, eds. (1983)).

A number of methods are known to those of skill in the art for identifying and characterizing promoter regions in plant genomic DNA (see, e.g., Jordano, *et al.*, *Plant Cell*, 1: 855-866 (1989); Bustos, *et al.*, *Plant Cell*, 1:839-854 (1989); Green, *et al.*, *EMBO J.* 7, 4035-4044 (1988); Meier, *et al.*, *Plant Cell*, 3, 309-316 (1991); and Zhang, *et al.*, *Plant Physiology* 110: 1069-1079 (1996)).

IV. Preparation of recombinant vectors

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising *et al.* *Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S

transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may direct expression of the
5 polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only in certain tissues, such as fruit, seeds, or flowers. As noted above, the promoters from the *IND1* genes described here are particularly useful for directing gene
10 expression so that a desired gene product is located in the valve margin of fruit. Other suitable promoters include those from genes such as *SHP1* or *SHP2* (Savidge, B., Rounsley, S.D., and Yanofsky, M.F. (1995) *Plant Cell* 7: 721-733). Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light.

15 If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (e.g., promoters or coding regions) from genes of the invention will typically comprise a marker gene that confers a
20 selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

IND1 nucleic acid sequences of the invention are expressed recombinantly
25 in plant cells to enhance and increase levels of endogenous *IND1* polypeptides. Alternatively, antisense or other *IND1* constructs (described above) are used to suppress *IND1* levels of expression. A variety of different expression constructs, such as expression cassettes and vectors suitable for transformation of plant cells can be prepared. Techniques for transforming a wide variety of higher plant species are well known and
30 described in the technical and scientific literature. See, e.g., Weising *et al. Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for a *IND1* polypeptide, e.g., a cDNA sequence encoding a full length protein, can be combined with cis-acting (promoter) and trans-acting (enhancer) transcriptional regulatory sequences to direct the

timing, tissue type and levels of transcription in the intended tissues of the transformed plant. Translational control elements can also be used.

The invention provides an *IND1* nucleic acid operably linked to a promoter which, in a preferred embodiment, is capable of driving the transcription of the
5 *IND1* coding sequence in plants. The promoter can be, e.g., derived from plant or viral sources. The promoter can be, e.g., constitutively active, inducible, or tissue specific. In construction of recombinant expression cassettes, vectors, transgenics, of the invention, a different promoters can be chosen and employed to differentially direct gene expression, e.g., in some or all tissues of a plant or animal. Typically, as discussed above, desired
10 promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the *IND1* genes described here.

A. *Constitutive Promoters*

A promoter fragment can be employed which will direct expression of
15 *IND1* nucleic acid in all transformed cells or tissues, e.g. as those of a regenerated plant. The term "constitutive regulatory element" means a regulatory element that confers a level of expression upon an operatively linked nucleic molecule that is relatively independent of the cell or tissue type in which the constitutive regulatory element is expressed. A constitutive regulatory element that is expressed in a plant generally is
20 widely expressed in a large number of cell and tissue types. Promoters that drive expression continuously under physiological conditions are referred to as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation.

A variety of constitutive regulatory elements useful for ectopic expression
25 in a transgenic plant are well known in the art. The cauliflower mosaic virus 35S (CaMV 35S) promoter, for example, is a well-characterized constitutive regulatory element that produces a high level of expression in all plant tissues (Odell *et al.*, *Nature* 313:810-812 (1985)). The CaMV 35S promoter can be particularly useful due to its activity in numerous diverse plant species (Benfey and Chua, *Science* 250:959-966 (1990); Futterer
30 *et al.*, *Physiol. Plant* 79:154 (1990); Odell *et al.*, *supra*, 1985). A tandem 35S promoter, in which the intrinsic promoter element has been duplicated, confers higher expression levels in comparison to the unmodified 35S promoter (Kay *et al.*, *Science* 236:1299 (1987)). Other useful constitutive regulatory elements include, for example, the cauliflower mosaic virus 19S promoter; the Figwort mosaic virus promoter; and the

nopaline synthase (nos) gene promoter (Singer *et al.*, *Plant Mol. Biol.* 14:433 (1990); An, *Plant Physiol.* 81:86 (1986)).

Additional constitutive regulatory elements including those for efficient expression in monocots also are known in the art, for example, the pEmu promoter and promoters based on the rice Actin-1 5' region (Last *et al.*, *Theor. Appl. Genet.* 81:581 (1991); Mcelroy *et al.*, *Mol. Gen. Genet.* 231:150 (1991); Mcelroy *et al.*, *Plant Cell* 2:163 (1990)). Chimeric regulatory elements, which combine elements from different genes, also can be useful for ectopically expressing a nucleic acid molecule encoding an IND1 polynucleotide (Comai *et al.*, *Plant Mol. Biol.* 15:373 (1990)).

Other examples of constitutive promoters include the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens* (see, e.g., Mengiste (1997) *supra*; O'Grady (1995) *Plant Mol. Biol.* 29:99-108); actin promoters, such as the *Arabidopsis* actin gene promoter (see, e.g., Huang (1997) *Plant Mol. Biol.* 1997 33:125-139); alcohol dehydrogenase (Adh) gene promoters (see, e.g., Millar (1996) *Plant Mol. Biol.* 31:897-904); *ACT11* from *Arabidopsis* (Huang *et al.* *Plant Mol. Biol.* 33:125-139 (1996)), *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong *et al.*, *Mol. Gen. Genet.* 251:196-203 (1996)), the gene encoding stearyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe *et al.* *Plant Physiol.* 104:1167-1176 (1994)), *GPc1* from maize (GenBank No. X15596, Martinez *et al.* *J. Mol. Biol.* 208:551-565 (1989)), *Gpc2* from maize (GenBank No. U45855, Manjunath *et al.*, *Plant Mol. Biol.* 33:97-112 (1997)), other transcription initiation regions from various plant genes known to those of skill. *See also* Holtorf *Plant Mol. Biol.* 29:637-646 (1995).

B. Inducible Promoters

Alternatively, a plant promoter may direct expression of the *IND1* nucleic acid of the invention under the influence of changing environmental conditions or developmental conditions. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light. Such promoters are referred to herein as "inducible" promoters. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) *supra*); the cold, drought, and high salt inducible promoter from potato (Kirch (1997) *Plant Mol. Biol.* 33:897-909).

Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins, are used to express the nucleic acids of the invention. For

example, the invention can use the auxin-response elements E1 promoter fragment (AuxREs) in the soybean (*Glycine max* L.) (Liu (1997) *Plant Physiol.* 115:397-407); the auxin-responsive *Arabidopsis* GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen (1996) *Plant J.* 10: 955-966); the auxin-inducible parC
 5 promoter from tobacco (Sakai (1996) 37:906-913); a plant biotin response element (Streit (1997) *Mol. Plant Microbe Interact.* 10:933-937); and, the promoter responsive to the stress hormone abscisic acid (Sheen (1996) *Science* 274:1900-1902).

Plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics, are also used to
 10 express the nucleic acids of the invention. For example, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder (1997) *Plant Cell Physiol.* 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. *IND1* coding sequence can also be under the control of, e.g., a
 15 tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) *Plant J.* 11:465-473); or, a salicylic acid-responsive element (Stange (1997) *Plant J.* 11:1315-1324; Uknes *et al.*, *Plant Cell* 5:159-169 (1993); Bi *et al.*, *Plant J.* 8:235-245 (1995)).

20 Particularly useful inducible regulatory elements include copper-inducible regulatory elements (Mett *et al.*, *Proc. Natl. Acad. Sci. USA* 90:4567-4571 (1993); Furst *et al.*, *Cell* 55:705-717 (1988)); tetracycline and chlor-tetracycline-inducible regulatory elements (Gatz *et al.*, *Plant J.* 2:397-404 (1992); Röder *et al.*, *Mol. Gen. Genet.* 243:32-38 (1994); Gatz, *Meth. Cell Biol.* 50:411-424 (1995)); ecdysone inducible regulatory
 25 elements (Christopherson *et al.*, *Proc. Natl. Acad. Sci. USA* 89:6314-6318 (1992); Kreutzweiser *et al.*, *Ecotoxicol. Environ. Safety* 28:14-24 (1994)); heat shock inducible regulatory elements (Takahashi *et al.*, *Plant Physiol.* 99:383-390 (1992); Yabe *et al.*, *Plant Cell Physiol.* 35:1207-1219 (1994); Ueda *et al.*, *Mol. Gen. Genet.* 250:533-539 (1996)); and *lac* operon elements, which are used in combination with a constitutively
 30 expressed *lac* repressor to confer, for example, IPTG-inducible expression (Wilde *et al.*, *EMBO J.* 11:1251-1259 (1992)). An inducible regulatory element useful in the transgenic plants of the invention also can be, for example, a nitrate-inducible promoter derived from the spinach nitrite reductase gene (Back *et al.*, *Plant Mol. Biol.* 17:9 (1991)) or a light-inducible promoter, such as that associated with the small subunit of RuBP

carboxylase or the LHCP gene families (Feinbaum *et al.*, *Mol. Gen. Genet.* 226:449 (1991); Lam and Chua, *Science* 248:471 (1990)).

C. Tissue-Specific Promoters

5 Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters). Tissue specific promoters are transcriptional control elements that are only active in particular cells or tissues at specific times during plant development, such as in vegetative tissues or reproductive tissues. Promoters from the *IND1* genes of the invention are particularly
10 useful for tissue-specific direction of gene expression so that a desired gene product is generated only or preferentially in embryos or seeds, as described below.

Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only (or primarily only) in certain tissues, such as vegetative tissues, e.g., roots or leaves, or reproductive tissues, such as fruit,
15 ovules, seeds, pollen, pistils, flowers, or any embryonic tissue. Reproductive tissue-specific promoters may be, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed and seed coat-specific, pollen-specific, petal-specific, sepal-specific, or some combination thereof.

The invention provides a transgenic plant that is characterized by delayed
20 seed dispersal due to expression of a nucleic acid molecule encoding an *IND1* gene product, or an antisense construct thereof, operatively linked to a dehiscence zone-selective regulatory element. The dehiscence zone-selective regulatory element can be, for example, an *SHP1* regulatory element or *SHP2* regulatory element. The *SHP1* regulatory element can be derived from the Arabidopsis *SHP1* genomic sequence
25 disclosed herein as SEQ ID NO:5 and can be, for example, a 5' regulatory sequence or intronic regulatory element. Similarly, the *SHP2* regulatory element can be derived from the Arabidopsis *SHP2* genomic sequence disclosed herein as SEQ ID NO:6 and can be, for example, a 5' regulatory sequence or intronic regulatory element.

A dehiscence zone-selective regulatory element can be derived from a
30 gene that is an ortholog of Arabidopsis *IND1* and is selectively expressed in the valve margin or dehiscence zone of a seed plant. A dehiscence zone-selective regulatory element can be derived, for example, from an *IND1* ortholog of the *Brassicaceae*, such as a *Brassica napus*, *Brassica oleracea*, *Brassica campestris*, *Brassica juncea*, *Brassica nigra* or *Brassica carinata* *IND1* ortholog. A dehiscence zone-selective regulatory

element can be derived, for example, from an *IND1* canola ortholog. A dehiscence zone-selective regulatory element also can be derived, for example, from a leguminous *IND1* ortholog, such as a soybean, pea, chickpea, moth bean, broad bean, kidney bean, lima bean, lentil, cowpea, dry bean, peanut, alfalfa, lucerne, birdsfoot trefoil, clover, stylosanthes, *lotononis bainesii*, or sainfoin *IND1* ortholog.

Dehiscence zone-selective regulatory elements also can be derived from a variety of other genes that are selectively expressed in the valve margin or dehiscence zone of a seed plant. For example, the rapeseed gene RDPG1 is selectively expressed in the dehiscence zone (Petersen *et al.*, *Plant Mol. Biol.* 31:517-527 (1996)). Thus, the RDPG1 promoter or an active fragment thereof can be a dehiscence zone-selective regulatory element as defined herein. Additional genes such as the rapeseed gene SAC51 also are known to be selectively expressed in the dehiscence zone; the SAC51 promoter or an active fragment thereof also can be a dehiscence zone-selective regulatory element of the invention (Coupe *et al.*, *Plant Mol. Biol.* 23:1223-1232 (1993)). The skilled artisan understands that a regulatory element of any such gene selectively expressed in cells of the valve margin or dehiscence zone can be a dehiscence zone-selective regulatory element as defined herein.

Additional dehiscence zone-selective regulatory elements can be identified and isolated using routine methodology. Differential screening strategies using, for example, RNA prepared from the dehiscence zone and RNA prepared from adjacent pod material can be used to isolate cDNAs selectively expressed in cells of the dehiscence zone (Coupe *et al.*, *supra*, 1993); subsequently, the corresponding genes are isolated using the cDNA sequence as a probe.

Enhancer trap or gene trap strategies also can be used to identify and isolate a dehiscence zone-selective regulatory element of the invention (Sundaresan *et al.*, *supra*, 1995; Koncz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8467-8471 (1989); Kertbundit *et al.*, *Proc. Natl. Acad. Sci. USA* 88:5212-5216 (1991); Topping *et al.*, *Development* 112:1009-1019 (1991)). Enhancer trap elements include a reporter gene such as GUS with a weak or minimal promoter, while gene trap elements lack a promoter sequence, relying on transcription from a flanking chromosomal gene for reporter gene expression. Transposable elements included in the constructs mediate fusions to endogenous loci; constructs selectively expressed in the valve margin or dehiscence zone are identified by their pattern of expression. With the inserted element as a tag, the flanking dehiscence zone-selective regulatory element is cloned using, for example, inverse polymerase chain

reaction methodology (see, for example, Aarts *et al.*, *Nature* 363:715-717 (1993); see also, Ochman *et al.*, "Amplification of Flanking Sequences by Inverse PCR," in Innis *et al.*, *supra*, 1990). The Ac/Ds transposition system of Sundaresan *et al.*, *Genes. Devel.* 9:1797-1810 (1995), can be particularly useful in identifying and isolating a dehiscence zone-selective regulatory element of the invention.

Dehiscence zone-selective regulatory elements also can be isolated by inserting a library of random genomic DNA fragments in front of a promoterless reporter gene and screening transgenic plants transformed with the library for dehiscence zone-selective reporter gene expression. The promoterless vector pROA97, which contains the npt gene and the GUS gene each under the control of the minimal 35S promoter, can be useful for such screening. The genomic library can be, for example, Sau3A fragments of *Arabidopsis thaliana* genomic DNA or genomic DNA from, for example, another Brassicaceae of interest (Ott *et al.*, *Mol. Gen. Genet.* 223:169-179 (1990); Claes *et al.*, *The Plant Journal* 1:15-26 (1991)).

Dehiscence zone-selective expression of a regulatory element of the invention can be demonstrated or confirmed by routine techniques, for example, using a reporter gene and in situ expression analysis. The GUS and firefly luciferase reporters are particularly useful for in situ localization of plant gene expression (Jefferson *et al.*, *EMBO J.* 6:3901 (1987); Ow *et al.*, *Science* 334:856 (1986)), and promoterless vectors containing the GUS expression cassette are commercially available, for example, from Clontech (Palo Alto, CA). To identify a dehiscence zone-selective regulatory element of interest such as an *IND1* regulatory element, one or more nucleotide portions of the *IND1* gene can be generated using enzymatic or PCR-based methodology (Glick and Thompson, *supra*, 1993; Innis *et al.*, *supra*, 1990); the resulting segments are fused to a reporter gene such as GUS and analyzed as described above.

Other tissue-specific promoters include seed promoters. Suitable seed-specific promoters are derived from the following genes: *MAC1* from maize (Sheridan (1996) *Genetics* 142:1009-1020); *Cat3* from maize (GenBank No. L05934, Abler (1993) *Plant Mol. Biol.* 22:10131-1038); *viviparous-1* from *Arabidopsis* (Genbank No. U93215); *atmyc1* from *Arabidopsis* (Urao (1996) *Plant Mol. Biol.* 32:571-57; Conceicao (1994) *Plant* 5:493-505); *napA* from *Brassica napus* (GenBank No. J02798, Josefsson (1987) *JBL* 26:12196-1301); and the napin gene family from *Brassica napus* (Sjodahl (1995) *Planta* 197:264-271).

A variety of promoters specifically active in vegetative tissues, such as leaves, stems, roots and tubers, can also be used to express the *IND1* nucleic acids of the invention. For example, promoters controlling patatin, the major storage protein of the potato tuber, can be used, see, e.g., Kim (1994) *Plant Mol. Biol.* 26:603-615; Martin
 5 (1997) *Plant J.* 11:53-62. The ORF13 promoter from *Agrobacterium rhizogenes* which exhibits high activity in roots can also be used (Hansen (1997) *Mol. Gen. Genet.* 254:337-343. Other useful vegetative tissue-specific promoters include: the tarin promoter of the gene encoding a globulin from a major taro (*Colocasia esculenta* L. Schott) corm protein family, tarin (Bezerra (1995) *Plant Mol. Biol.* 28:137-144); the
 10 curculin promoter active during taro corm development (de Castro (1992) *Plant Cell* 4:1549-1559) and the promoter for the tobacco root-specific gene TobRB7, whose expression is localized to root meristem and immature central cylinder regions (Yamamoto (1991) *Plant Cell* 3:371-382).

Leaf-specific promoters, such as the ribulose biphosphate carboxylase
 15 (RBCS) promoters can be used. For example, the tomato RBCS1, RBCS2 and RBCS3A genes are expressed in leaves and light-grown seedlings, only RBCS1 and RBCS2 are expressed in developing tomato fruits (Meier (1997) *FEBS Lett.* 415:91-95). A ribulose bisphosphate carboxylase promoters expressed almost exclusively in mesophyll cells in leaf blades and leaf sheaths at high levels, described by Matsuoka (1994) *Plant J.*
 20 6:311-319, can be used. Another leaf-specific promoter is the light harvesting chlorophyll a/b binding protein gene promoter, see, e.g., Shiina (1997) *Plant Physiol.* 115:477-483; Casal (1998) *Plant Physiol.* 116:1533-1538. The *Arabidopsis thaliana* myb-related gene promoter (Atmyb5) described by Li (1996) *FEBS Lett.* 379:117-121, is leaf-specific. The Atmyb5 promoter is expressed in developing leaf trichomes, stipules,
 25 and epidermal cells on the margins of young rosette and cauline leaves, and in immature seeds. Atmyb5 mRNA appears between fertilization and the 16 cell stage of embryo development and persists beyond the heart stage. A leaf promoter identified in maize by Busk (1997) *Plant J.* 11:1285-1295, can also be used.

Another class of useful vegetative tissue-specific promoters are
 30 meristematic (root tip and shoot apex) promoters. For example, the “*SHOOTMERISTEMLESS*” and “*SCARECROW*” promoters, which are active in the developing shoot or root apical meristems, described by Di Laurenzio (1996) *Cell* 86:423-433; and, Long (1996) *Nature* 379:66-69; can be used. Another useful promoter is that which controls the expression of 3-hydroxy-3- methylglutaryl coenzyme A

reductase HMG2 gene, whose expression is restricted to meristematic and floral (secretory zone of the stigma, mature pollen grains, gynoecium vascular tissue, and fertilized ovules) tissues (see, e.g., Enjuto (1995) *Plant Cell* 7:517-527). Also useful are kn1-related genes from maize and other species which show meristem-specific
5 expression, see, e.g., Granger (1996) *Plant Mol. Biol.* 31:373-378; Kerstetter (1994) *Plant Cell* 6:1877-1887; Hake (1995) *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 350:45-51. For example, the *Arabidopsis thaliana* KNAT1 promoter. In the shoot apex, KNAT1 transcript is localized primarily to the shoot apical meristem; the expression of KNAT1 in the shoot meristem decreases during the floral transition and is restricted to the cortex of
10 the inflorescence stem (see, e.g., Lincoln (1994) *Plant Cell* 6:1859-1876).

One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

15 In another embodiment, a *IND1* nucleic acid is expressed through a transposable element. This allows for constitutive, yet periodic and infrequent expression of the constitutively active polypeptide. The invention also provides for use of tissue-specific promoters derived from viruses which can include, e.g., the tobamovirus subgenomic promoter (Kumagai (1995) *Proc. Natl. Acad. Sci. USA* 92:1679-1683; the
20 rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives strong phloem-specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer (1996) *Plant Mol. Biol.* 31:1129-1139).

25

V. Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using
30 techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host

will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984).

Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Ballistic transformation techniques are described in Klein et al. *Nature* 327:70-73 (1987).

Agrobacterium tumefaciens-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype such as seedlessness. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture*, Handbook of Plant Cell Culture, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Cucumis*, *Cucurbita*, *Daucus*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Oryza*, *Panicum*, *Pennisetum*, *Persea*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Trigonella*, *Triticum*, *Vitis*, *Vigna*, and, *Zea*. A useful plant of the invention can be a dehiscent seed plant, and a particularly useful plant of the invention can be a member of the *Brassicaceae*, such as rapeseed, or a member of the *Fabaceae*, such as a soybean, pea, lentil or bean plant.

In one embodiment, the invention provides a dehiscent seed plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product in the dehiscent seed plant. As used herein, the term "dehiscent seed plant" means a plant that produces a dry dehiscent fruit, which has fruit walls that open to permit escape of the seeds contained therein. Dehiscent fruits commonly contain several seeds and include the fruits known, for example, as legumes, capsules and siliques.

In one embodiment, the invention provides a plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product, where the plant is a member of the *Brassicaceae*. The *Brassicaceae*, commonly known as the Brassicas, are a diverse group of crop plants with great economic value worldwide (see, for example, Williams and Hill, *Science* 232:1385-1389 (1986)). The *Brassicaceae* produce seed oils for margarine, salad oil, cooking oil, plastic and industrial uses; condiment mustard; leafy, stored, processed and pickled vegetables; animal fodders and green manures for soil rejuvenation. A particularly useful non-naturally occurring Brassica plant of the invention is the oilseed plant canola.

There are six major Brassica species of economic importance, each containing a range of plant forms. *Brassica napus* includes plants such as the oilseed rapes and rutabaga. *Brassica oleracea* are the cole crops such as cabbage, cauliflower, kale, kohlrabi and Brussels sprouts. *Brassica campestris* (*Brassica rapa*) includes plants such as Chinese cabbage, turnip and pak choi. *Brassica juncea* includes a variety of mustards; *Brassica nigra* is the black mustard; and *Brassica carinata* is Ethiopian mustard. The skilled artisan understands that any member of the *Brassicaceae* can be modified as disclosed herein to produce a non-naturally occurring Brassica plant characterized by delayed seed dispersal.

In a second embodiment, the invention provides a plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product, where the plant is a member of the *Fabaceae*. The *Fabaceae*, which are commonly known as members of the pea family, are plants that produce a characteristic dry dehiscent fruit known as a legume. The legume is derived from a single carpel and dehisces along the suture of the carpel margins and along the median vein. The *Fabaceae* encompass both grain legumes and forage legumes. Grain legumes include, for example, soybean (glycine), pea, chickpea, moth bean, broad bean,

kidney bean, lima bean, lentil, cowpea, dry bean and peanut. Forage legumes include alfalfa, lucerne, birdsfoot trefoil, clover, stylosanthes species, *lotononis bainessii* and sainfoin. The skilled artisan will recognize that any member of the *Fabaceae* can be modified as disclosed herein to produce a non-naturally occurring plant of the invention
5 characterized by delayed seed dispersal.

A non-naturally occurring plant of the invention characterized by delayed seed dispersal also can be a member of the plant genus *Cuphea* (family *Lythraceae*). A *Cuphea* plant is particularly valuable since *Cuphea* oilseeds contain industrially and nutritionally important medium-chain fatty acids, especially lauric acid, which is
10 currently supplied only by coconut and palm kernel oils.

A non-naturally occurring plant of the invention also can be, for example, one of the monocotyledonous grasses, which produce many of the valuable small-grain cereal crops of the world. Suppression of *IND1* expression as described above, can be useful in generating a non-naturally occurring small grain cereal plant, such as a barley,
15 wheat, oat, rye, orchard grass, guinea grass, sorghum or turf grass plant characterized by delayed seed dispersal.

VI. Additional modifications that modulate seed dispersal

It should be recognized that a plant of the invention, which contains an
20 exogenous *IND1* polynucleotide, also can contain one or more additional modifications, including naturally and non-naturally occurring modifications, that can modulate the delay in seed dispersal. For example, the plant hormone ethylene promotes fruit dehiscence, and modified expression or activity of positive or negative regulators of the ethylene response can be included in a plant of the invention (see, generally, Meakin and
25 Roberts, *J. Exp. Botany* 41:1003-1011 (1990); Ecker, *Science* 268:667-675 (1995); Chao *et al.*, *Cell* 89:1133-1144 (1997)).

Mutations in positive regulators of the ethylene response show a reduction or absence of responsiveness to treatment with exogenous ethylene. Arabidopsis mutations in positive regulators of the ethylene response include mutations in *etr*, which
30 inactivate a histidine kinase ethylene receptor (Bleeker *et al.*, *Science* 241:1086-1089 (1988); Schaller and Bleeker, *Science* 270:1809-1811 (1995)); *ers* (Hua *et al.*, *Science* 269:1712-1714 (1995)); *ein2* (Guzman and Ecker, *Plant Cell* 2:513 (1990)); *ein3* (Rothenberg and Ecker, *Sem. Dev. Biol. Plant Dev. Genet.* 4:3-13 (1993); Kieber and Ecker, *Trends Genet.* 9:356-362 (1993)); *ain1* (van der Straeten *et al.*, *Plant Physiol.*

102:401-408 (1993)); *eti* (Harpham *et al.*, *An. Bot.* 68:55 (1991)) and *ein4*, *ein5*, *ein6*, and *ein7* (Roman *et al.*, *Genetics* 139: 1393-1409 (1995)). Similar genetic functions are found in other plant species; for example, the never-ripe mutation corresponds to *etr* and confers ethylene insensitivity in tomato (Lanahan *et al.*, *The Plant Cell* 6:521-530 (1994);
5 Wilkinson *et al.*, *Science* 270:1807-1809 (1995)). A plant of the invention can include a modification that results in altered expression or activity of any such positive regulator of the ethylene response. A mutation in a positive regulator, for example, can be included in a plant of the invention and can modify the delay in seed dispersal in such plants, for example, by further postponing the delay in seed dispersal.

10 Mutations in negative regulators of the ethylene response display ethylene responsiveness in the absence of exogenous ethylene. Such mutations include those relating to ethylene overproduction, for example, the *eto1*, *eto2*, and *eto3* mutants, and those relating to constitutive activation of the ethylene signalling pathway, for example, mutations in *CTR1*, a negative regulator with sequence similarity to the Raf family of
15 protein kinases (Kieber *et al.*, *Cell* 72:427-441 (1993), which is incorporated herein by reference). A plant of the invention can include a modification that results in altered expression or activity of any such negative regulator of the ethylene response. A mutation resulting in ethylene responsiveness in the absence of exogenous ethylene, for example, can be included in a non-naturally occurring plant of the invention and can
20 modify, for example, diminish, the delay in seed dispersal.

Fruit morphological mutations also can be included in a plant of the invention. Such mutations include those in carpel identity genes such as *AGAMOUS* (Bowman *et al.*, *supra*, 1989; Yanofsky *et al.*, *supra*, 1990) and in genes required for normal fruit development such as *ETTIN*, *CRABS CLAW*, *SPATULA*, *AGL8* and
25 *TOUSLED* (Sessions *et al.*, *Development* 121:1519-1532 (1995); Alvarez and Smyth, *Flowering Newsletter* 23:12-17 (1997); and Roe *et al.*, *Cell* 75:939-950 (1993)). Thus, it is understood that a plant of the invention can include one or more additional genetic modifications, which can diminish or enhance the delay in seed dispersal.

30 VII. Expression of cytotoxic gene products

The present invention also provides a recombinant nucleic acid molecule that includes a dehiscence zone-selective regulatory element operatively linked to a nucleic acid molecule encoding a cytotoxic gene product. Further provided herein is a plant of the invention that is characterized by delayed seed dispersal due to expression of

a recombinant nucleic acid molecule having a dehiscence zone--selective regulatory element operatively linked to a nucleic acid molecule encoding a cytotoxic gene product.

A cytotoxic gene product is a gene product that causes the death of the cell in which it is expressed and, preferably, does not result in the death of cells other than the cell in which it is expressed. Thus, expression of a cytotoxic gene product from a dehiscence zone-selective regulatory element can be used to ablate the dehiscence zone without disturbing neighboring cells of the replum or valve. A variety of cytotoxic gene products useful in seed plants are known in the art including, for example, diphtheria toxin A chain polypeptides; RNase T1; Barnase RNase; ricin toxin A chain polypeptides; and herpes simplex virus thymidine kinase (tk) gene products. While the diphtheria toxin A chain, RNase T1 and Barnase RNase are preferred cytotoxic gene products, the skilled person recognizes that these, or other cytotoxic gene products can be used with a dehiscence zone-selective regulatory element to generate a non-naturally occurring seed plant characterized by delayed seed dispersal.

Diphtheria toxin is the naturally occurring toxin of *Cornibacterium diphtheriae*, which catalyzes the ADP-ribosylation of elongation factor 2, resulting in inhibition of protein synthesis and consequent cell death (Collier, *Bacteriol. Rev.* 39:54-85 (1975)). A single molecule of the fully active toxin is sufficient to kill a cell (Yamaizumi *et al.*, *Cell* 15:245-250 (1978)). Diphtheria toxin has two subunits: the diphtheria toxin B chain directs internalization to most eukaryotic cells through a specific membrane receptor, whereas the A chain encodes the toxic catalytic domain. The catalytic DT-A chain does not include a signal peptide and is not secreted. Further, any DT-A released from dead cells in the absence of the diphtheria toxin B chain is precluded from cell attachment. Thus, DT-A is cell autonomous and directs killing only of the cells in which it is expressed without apparent damage to neighboring cells. The DT-A expression cassette of Palmiter *et al.*, which contains the 193 residues of the A chain engineered with a synthetic ATG and lacking the native leader sequence, is particularly useful in the seed plants of the invention (Palmiter *et al.*, *Cell* 50:435-443 (1987); Greenfield *et al.*, *Proc. Natl. Acad. Sci., USA* 80:6853-6857 (1983)).

RNase T1 of *Aspergillus oryzae* and Barnase RNase of *Bacillus amylolique-faciens* also are cytotoxic gene products useful in the seed plants of the invention (Thorsness and Nasrallah, *Methods in Cell Biology* 50:439-448 (1995)). Barnase RNase may be more generally toxic to plants than RNase T1 and, thus, is preferred in the methods of the invention.

Ricin, a ribosome-inactivating protein produced by castor bean seeds, also is a cytotoxic gene product useful in a non-naturally occurring seed plant of the invention. The ricin toxin A chain polypeptide can be used to direct cell-specific ablation as described, for example, in Moffat *et al.*, *Development* 114:681-687 (1992). Plant
 5 ribosomes are variably susceptible to the plant-derived ricin toxin. The skilled person understands that the toxicity of ricin depends is variable and should be assessed for toxicity in the seed plant species of interest (see Olsnes and Pihl, *Molecular Action of Toxins and Viruses*, pages 51-105, Amsterdam: Elsevier Biomedical Press (1982)).

10

EXAMPLES

The following examples are offered to illustrate, but no to limit the claimed invention.

Example 1:

15

The GT140 valve margin marker (Sundaresan, V., *et al. Genes Dev.* 9, 1797-1810 (1995)) is expressed at the valve margin of the developing gynoecium just prior to fertilization (stage 13) and this pattern persists in the mature fruit (stage 17). As expression of this marker is largely absent from the valve margins of *shp1 shp2* indehiscent fruits (Liljegren, S.J., Ditta, G.S., Eshed, Y., Savidge, B., Bowman, J.L., and
 20 Yanofsky, M.F. (2000). SHATTERPROOF MADS-box genes control seed dispersal in *Arabidopsis*. *Nature* 404, 766-770), it was expected that the gene corresponding to this marker might also be involved in valve margin development and be required for fruit dehiscence.

25

To isolate flanking genomic sequence from the GT140 marker insertion site, TAIL/PCR was performed as previously described (Tsugeki, R., *et al. Plant J.* 10, 479-489 (1996)). Subsequent sequencing of the isolated PCR products demonstrated that they correspond to a fully sequenced BAC from chromosome 4, available in the public database as part of the *Arabidopsis* Genome Initiative. The GT140 insertion is located between two genes, one encoding a predicted basic helix-loop-helix (bHLH) transcription factor and the other representing a novel gene.

30

Through several lines of subsequent investigation, it was confirmed that the bHLH transcription factor (herein referred to as *IND1* as noted below) was the relevant gene (SEQ ID NO:1). Promoter/enhancer::GUS fusions of the *IND1* gene were introduced into wild-type plants and found to express GUS in an identical pattern to that of the GT140 marker line. Interestingly, approximately 25% of the transgenic lines failed

to express significant GUS activity and displayed an indehiscent phenotype. The most likely explanation of these results is that the *IND1::GUS* fusions, as well as of the endogenous *IND1* gene, were cosuppressed. Subsequent RNA blotting confirmed a down regulation of the *IND1* gene in these lines, and further RNA blotting showed, as expected,
 5 a decrease in *IND1* gene expression in *shp1 shp2* fruits.

In parallel to the studies of the GT140 valve margin marker described above, screens for Arabidopsis mutants producing indehiscent fruits were also carried out. Besides obtaining additional alleles of *SHP1* and *SHP2* through EMS mutagenesis of *shp2-1* and *shp1-1* seed stocks, indehiscent mutants that were not allelic to either *SHP1* or
 10 *SHP2*, respectively were also obtained. Because the GT140 studies suggested the possibility that one or more of these indehiscent mutants might correspond to the *IND1* gene, *IND1* from several of these mutants was cloned and sequenced. Four alleles represent independent mutant alleles of *IND1*. The strongest allele, *ind1-2*, contains a single nucleotide deletion within codon 55 that results in a frameshift and production of a
 15 truncated protein of 64 rather than 198 amino acids. The *ind1-1* and *ind1-3* alleles contain nucleotide substitutions at codons 141 and 128 that changes a leucine amino acid to a phenylalanine and an arginine to a histidine, respectively. These affected amino acids are both at conserved positions within the bHLH domain. The *ind1-4* allele contains a
 20 nucleotide substitution at codon 92 that changes a glutamine to a stop codon, causing production of a truncated protein of 91 amino acids. Since inactivation of this bHLH transcription factor prevents fruit dehiscence, the gene is referred to as *INDEHISCENT1* (*IND1*) and the mutant as, *ind1*. To date, *ind1* represents the only reported single gene mutation in Arabidopsis that specifically blocks fruit dehiscence.

25 It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

30

SEQUENCE LISTING

<110> The Regents of the University of California

<120> Control of Fruit Dehiscence in Arabidopsis by
INDEHISCENT1 Genes

<130> 40330-1956

<140> CA 2,406,388

<141> 2001-04-13

<150> US 09/548,971

<151> 2000-04-13

<160> 9

<170> PatentIn Ver. 2.1

<210> 1

<211> 3856

<212> DNA

<213> Arabidopsis thaliana

<220>

<223> INDEHISCENT1 (IND1) genomic

<220>

<221> CDS

<222> (2765) .. (3361)

<223> IND1

<400> 1

```

ctctagacca tctactatcc ggttggttgac ccttaaagct tttgaagact actagaataa 60
tgcaaatacc atatgtccat atccatcctt ttcttttgtt tgaactgaac attctaattt 120
tgtaaaagaa aaaaccttat gttaatatca ccgtaggcaa aaaaaatatc tcatcatatt 180
aaattttttat tataagatta tacatttctt cgttgtaaga gttactccaa ttgcaagtgt 240
tgtattaact aataaaaagg acgaaagtag gaagcttata attaattgat gttgcatagt 300
actggtatat tgttgatgaa tataacaagt atgaacatta atgcatgaaa cgggggtattt 360
tgtcttgaac tcattaaagg caatgtgaaa agaagatgtg aggtctcatt ttgaaaattt 420
atcttctagc tttgtcgatt ttaaattctat gaaatgaacg caacatatag aaatttcatg 480
tggacaacga catttagacg gtatcttaat tagaccgatt aattagtaat atacttatat 540
atataattag tggtgattat aagtttactt atccacttga gaattttaa aatgggcaat 600
accttaatgt cgaaagaagc cgtccccact tcgtgtaatg agttatgggg gagagatcct 660
gttaaatacgt caaataaaaac aacttaagaa ctagaaattg acaccaaaaa tcataaagag 720
aacgttgaag aagtcattta tcgtatccag ctcatatttc ctagctaaga tcaaatcaag 780
gccgttgaaa gggcttgtaa gaaaatgtcg aagaaaccgt ggggtttaga agaaagacaa 840
gaaatagaag aacaatgatg ttaaattgcc tattttggtg tataggagtt gtcaaaagag 900
gagagagaga agaaaattag gtcaaaaataa tgagcactaa aaatggagac atgtgttgag 960
taactattac aagagcgact tatgcttcct tatggcaatg atatccaaac caaagtgcaa 1020
cgctcctttt ttgccctaatt ttcgtaaagt ctctctcctt ctctcgtcctt aggaaaaacc 1080
ctagaaattt aatcccttgt tcttgatcct gctttttgag taaccatgat tttgaccaca 1140
cactatttct tctatctttt gtggtctata ggattttgct ttatatgtgt ttcttgtatt 1200
gctccgtacg tacgtatacg aattttaaag gttataacaa ggtttatata aactagcaca 1260
aatgagtcca tgaaatttgt tagcgaaaaa ggtagaaata tattgagtct ttaaaccggca 1320
atatatataa ttttgctgca aaacttagct ttaatcatga tctaatagata ttttctttta 1380
tttcctttgc caaattaatc acatgcacgg atttttggca agttatgtgt cgaattcttc 1440
cattcacaca acactaaact taattagaac tctaggaaat attttaaaat gacaacttta 1500
tcgaaaaaaa tttagttatg aaaacaattc cagaattaaa catgagctat ataatttaag 1560
ataaaatgaa gtaatatgta tatgtatgta ataacatatc tgattgcggt aaaaaaaaac 1620

```

```

atatctgatt aaattgttca tgcaggccca tgtcactatg atgtcatcac gtttttattt 1680
tcacaataac taatatatat tcaaaaaaat agttttgtca gattaaattt tttttggtgg 1740
tcagctttct ccaacctact aaactagttt ggaatgttct cttctttatt tttctttttc 1800
ttgatttctt atgtttttta tttatggaat tttaagacgg attgtttagg tcgtttctct 1860
cttttcttgt tttctaaagt tacttttgta aactcatctc ctcccaatta gacagtcaat 1920
catatagtta tcttttaata tatgtctagt tgataaaaaa aatgaaaaaa tactggtggt 1980
agttctacta atgtttgtgt aaaaaatctg atattatgaa tctaataaat ttctttgatc 2040
gtataatgtg ggttaaattt agtaattttt tacataaata agaactgtaa tgttgatgta 2100
tattggggaa tcagtatatatt agcttgggta actatacttc tggaaataact tgaagattta 2160
actatttgca aaattataat ttagtcccgaa aaaatacaga cgacgggaca cgacaacata 2220
taagcagggt tgaatcttgg aaaattttgt atacataacc tatataaata ctaatgttct 2280
ggttgggttc aaaagccttt tcaaaagtcc cattttttta attcaaggac attttacata 2340
ggaaataagt tgagtcataa aaaataatgg ttatttttgta aggttttttt tttgattaaa 2400
acgcacatat taagaagtta gttttttttc actaccaa atcaattaat ttaaaaccat 2460
gcaaccattc ataaaacaat actattaaag aatataaata atcacaaaat attaaatata 2520
cttaaaattt acatataaat ttacaaaaca tctaattaat tgaaacagaa aggaaaaggt 2580
aaaatatatc ataaaatgag acatatatcc tataaaaaaa aaatgaggca tatgaagtaa 2640
ataataagag acatgcatgt aagcattcgg ttaattaatc gagtcaaaga tatatatcag 2700
taaatacata tgtgtatatatt tctggaaaaa gaatatatat attgagaaat aagaaaagat 2760

```

```

gaaa atg gaa aat ggt atg tat aaa aag aaa gga gtg tgc gac tct tgt 2809
Met Glu Asn Gly Met Tyr Lys Lys Lys Gly Val Cys Asp Ser Cys
      1              5              10              15

```

```

gtc tgc tcc aaa agc aga tcc aac cac agc ccc aaa aga agc atg atg 2857
Val Ser Ser Lys Ser Arg Ser Asn His Ser Pro Lys Arg Ser Met Met
              20              25              30

```

```

gag cct cag cct cac cat ctc ctc atg gat tgg aac aaa gct aat gat 2905
Glu Pro Gln Pro His His Leu Leu Met Asp Trp Asn Lys Ala Asn Asp
              35              40              45

```

```

ctt ctc aca caa gaa cac gca gct ttt ctc aat gat cct cac cat ctc 2953
Leu Leu Thr Gln Glu His Ala Ala Phe Leu Asn Asp Pro His His Leu
              50              55              60

```

```

atg tta gat cca cct ccc gaa acc cta att cac ttg gac gaa gac gaa 3001
Met Leu Asp Pro Pro Pro Glu Thr Leu Ile His Leu Asp Glu Asp Glu
              65              70              75

```

```

gag tac gat gaa gac atg gat gcg atg aag gag atg cag tac atg atc 3049
Glu Tyr Asp Glu Asp Met Asp Ala Met Lys Glu Met Gln Tyr Met Ile
              80              85              90              95

```

```

gcc gtc atg cag ccc gta gac atc gac cct gcc acg gtc cct aag ccg 3097
Ala Val Met Gln Pro Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro
              100              105              110

```

```

aac cgc cgt aac gta agg ata agc gac gat cct cag acg gtg gtt gct 3145
Asn Arg Arg Asn Val Arg Ile Ser Asp Asp Pro Gln Thr Val Val Ala
              115              120              125

```

```

cgt cgg cgt cgg gaa agg atc agc gag aag atc cga att ctc aag agg 3193
Arg Arg Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg
              130              135              140

```

```

atc gtg cct ggt ggt gcg aag atg gac aca gct tcc atg ctc gac gaa 3241
Ile Val Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu
              145              150              155

```

gcc ata cgt tac acc aag ttc ttg aaa cgg cag gtg agg att ctt cag 3289
 Ala Ile Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Ile Leu Gln
 160 165 170 175

cct cac tct cag att gga gct cct atg gct aac ccc tct tac ctt tgt 3337
 Pro His Ser Gln Ile Gly Ala Pro Met Ala Asn Pro Ser Tyr Leu Cys
 180 185 190

tat tac cac aac tcc caa ccc tga tgaactacac agaagctcgc tagctagaca 3391
 Tyr Tyr His Asn Ser Gln Pro
 195

tttggtgtca tcctctcaac ctttttcatg ttgatataatt atatatagat gcataaagat 3451
 tcgatccaag attgtatggg tgttttaata ttattattct aagatatatg atgtacaatt 3511
 gtgtaccaag tttctttatc ttgatatacat atgcataaat aattggtgaa taaaaagaag 3571
 atattgattg taaacaaaaa aaagaagata ttgattgtta attaggggtt gatcattctg 3631
 tatgaaagct ttggcctgca aattaatttt cgatatatat atatatatat ggagaatata 3691
 tatcaaatac ttttttaatt tgactataat ttgtatcaat tatctgaatc tgatgagtgt 3751
 aggttatata tggattagca aaaaagaaaa caaccattat tacgcaccta cattaaaaat 3811
 catccaccaa agaagaacc atcctcaaga gggttccctc tagag 3856

<210> 2

<211> 198

<212> PRT

<213> Arabidopsis thaliana

<223> INDEHISCENT1 (IND1) protein

<400> 2

Met Glu Asn Gly Met Tyr Lys Lys Lys Gly Val Cys Asp Ser Cys Val
 1 5 10 15
 Ser Ser Lys Ser Arg Ser Asn His Ser Pro Lys Arg Ser Met Met Glu
 20 25 30
 Pro Gln Pro His His Leu Leu Met Asp Trp Asn Lys Ala Asn Asp Leu
 35 40 45
 Leu Thr Gln Glu His Ala Ala Phe Leu Asn Asp Pro His His Leu Met
 50 55 60
 Leu Asp Pro Pro Pro Glu Thr Leu Ile His Leu Asp Glu Asp Glu Glu
 65 70 75 80
 Tyr Asp Glu Asp Met Asp Ala Met Lys Glu Met Gln Tyr Met Ile Ala
 85 90 95
 Val Met Gln Pro Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn
 100 105 110
 Arg Arg Asn Val Arg Ile Ser Asp Asp Pro Gln Thr Val Val Ala Arg
 115 120 125
 Arg Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Ile
 130 135 140
 Val Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala
 145 150 155 160
 Ile Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Ile Leu Gln Pro
 165 170 175
 His Ser Gln Ile Gly Ala Pro Met Ala Asn Pro Ser Tyr Leu Cys Tyr
 180 185 190
 Tyr His Asn Ser Gln Pro
 195

<210> 3

<211> 2765

<212> DNA

<213> Arabidopsis thaliana

<220>

<223> 5' untranslated region IND1 promoter

<400> 3

```

ctctagacca tctactatcc ggttggttgac ccttaaagct tttgaagact actagaataa 60
tgcaaatacc atatgtccat atccatcctt ttcttttggt tgaactgaac attctaattt 120
tgtaaaagaa aaaaccttat gttaatatca ccgtaggcaa aaaaaatatc tcatcatatt 180
aaatttttat tataagatta tacattctct cgttgtaaga gttactccaa ttgcaagtgt 240
tgtattaact aataaaaagg acgaaagtag gaagcttata attaattgat gttgcatagt 300
actggtatat tggtgatgaa tataacaagt atgaacatta atgcatgaaa cggggtat 360
tgtcttgaac tcattaaagg caatgtgaaa agaagatgtg aggtctcatt ttgaaaattt 420
atcttctagc tttgtcgatt ttaaatctat gaaatgaacg caacatatag aaatttcatt 480
tggacaacga catttagacg gtatcttaat tagaccgatt aattagtaat atacttatat 540
atataattag tggtgattat aagtttactt atccacttga gaattttaa aatgggcaat 600
accttaatgt cgaaagaagc cgtccccact tcgtgtaatg agttatgggg gagagatcct 660
gttaaactcg caaataaaaac aacttaagaa ctagaaattg acaccaaaaa tcataaagag 720
aacgttgaag aagtcattta tcgtatccag ctcatatttc ctagctaaga tcaaatcaag 780
gccgttgaaa gggcttgtaa gaaaatgtcg aagaaaccgt ggggtttaga agaaagacaa 840
gaaatagaag aacaatgatg ttaaattgcc tattttggtg tataggagtt gtcaaaagag 900
gagagagaga agaaaattag gtcaaaataa tgagcactaa aaatggagac atgtgttgag 960
taactattac aagagcgact tatgcttctt tatggcaatg atatccaaac caaagtgcaa 1020
cgctcctttt ttgccctaatt ttcgtaaagt ctctctcctt cttcgtcctt aggaaaaacc 1080
ctagaaattt aatcccttgt tcttgatctt gctttttgag taaccatgat tttgaccaca 1140
cactatttct tctatctttt gtggtctata ggattttgct ttatatgtgt ttcttgtatt 1200
gctccgtacg tacgtatacg aattttaaag gttataacaa ggtttatata aactagcaca 1260
aatgagtcca tgaaatttgt tagcgaaaaa ggtagaaata tattgagtct ttaaaccggc 1320
atataataaa ttttgctgca aaacttagct ttaatcatga tctaatagata ttttctttaa 1380
tttcttttgc caaattaatc acatgcacgg atttttggca agttatgtgt cgaattcttc 1440
cattcacaca acactaaact taattagaac tctaggaaat atttttaa aatgacaactt 1500
tcgaaaaaaa tttagttatg aaaacaattc cagaattaaa catgagctat ataatttaag 1560
ataaaatgaa gtaatatgta tatgtatgta ataacatate tgattgcggt aaaaaaaaac 1620
atatctgatt aaattgttca tgcaggccca tgtcactatg atgtcatcac gtttttattt 1680
tcacaataac taatatatat tcaaaaaaat agttttgtca gattaaattt tttttggtgg 1740
tcagctttct ccaacctact aaactagttt ggaatgttct cttctttatt tttcttttct 1800
ttgatttctt atgtttttta tttatggaat ttaagacgg attgtttagg tcgtttctct 1860
cttttcttgt tttctaaagt tacttttgta aactcatctc ctcccaatta gacagtcaat 1920
catatagtta tcttttaata tatgtctagt tgataaaaaa aatgaaaaaa tactggtggt 1980
agttctacta atgtttgtgt aaaaaatctg atattatgaa tctaataaat ttctttgatc 2040
gtataatgtg ggtaaaattt agtaattttt tacataaata agaactgtaa tggtgatgta 2100
tattggggaa tcagtatatt agcttgggta actatacttc tggaaatact tgaagattta 2160
actatttgca aaattataat ttagtcccga aaaatacaga cgacgggaca cgacaacata 2220
taagcaggtt tgaatcttgg aaaattttgt atacataacc tatataaata ctaatgttct 2280
ggttgggttc aaaagccttt tcaaaagtcc cattttttta attcaaggac attttacata 2340
ggaaataagt tgagtcataa aaaataatgg ttattttgta aggttttttt tttgattaaa 2400
acgcacatat taagaagtta gttttttttc actaccaa atcaattaat ttaaaaccat 2460
gcaaccattc ataaaacaat actattaaag aatataaata atcacaaaat attaaataca 2520
cttaaaattt acatataaat ttacaaaaca tctaattaat tgaaacagaa aggaaaaggt 2580
aaaatatatc ataaaatgag acatatatcc tataaaaaaa aatgagggca tatgaagtaa 2640
ataataagag acatgcatgt aagcattcgg ttaattaatc gagtcaaaga tatatatcag 2700
taaatacata tgtgtatatt tctggaaaaa gaatatatat attgagaaat aagaaaagat 2760
gaaaa

```

<210> 4

<211> 496

<212> DNA

<213> Arabidopsis thaliana

<220>

<223> 3' untranslated region IND1 promoter

<400> 4

```

atgaactaca cagaagctcg ctagctagac atttggtgtc atcctctcaa cctttttcat 60
gttgatatat tatatataga tgcataaaga ttcgatccaa gattgtatgg gtgttttaat 120
attattattc taagatatat gatgtacaat tgtgtaccaa gtttctttat cttgatatca 180
tatgcataaa taattggtga ataaaaagaa gatattgatt gtaaacaaaa aaaagaagat 240
attgattggt aattagggtt tgatcattct gtatgaaagc tttggcctgc aaattaattt 300
tcgatatata tatatatata tggagaatat atatcaaata cttttttaat ttgactataa 360
tttgtatcaa ttatctgaat ctgatgagtg taggttatat atggattagc aaaaaagaaa 420
acaaccatta ttacgcacct acattaaaaa tcatccacca aagaagaaac catcctcaag 480
agggttcctt ctagag                                     496

```

<210> 5

<211> 5622

<212> DNA

<213> *Arabidopsis thaliana*

<220>

<223> SHATTERPROOF1 (SHP1) genomic

<220>

<221> modified_base

<222> (935)..(941)

<223> n = any nucleotide

<400> 5

```

agatctgcaa cagtgaaaag agaaaacaaa atggacttga agaggttttg acaatgccag 60
agataatgct tattccctaa tatgttgcca gccaaagtgt aaattggctt tttaaataatg 120
gatttctgta tcagtgggtca tatttgtgga tccaacgtat tcatcatcaa gttctcaagt 180
ttgctttcag tgcaattcta attcacacgt ttaactttaa catgcatgtc attataatta 240
cttcttcact aagacacaat acggcaaacc tttcagatta tattaatctc cataaatgaa 300
ataattaacc tcataatcaa gattcaatgt ttctaaatat atatggacaa aatttacacg 360
gaagattaga tacgtatatt agtagattta gtctttcgtt tgtgcgataa gattaaccac 420
ctcatagata gtaatatcat tgtcaaattc ctctcggttt agtcgctaaa ttgtatcttt 480
tttaagccta aaagtagtgt attcgcataat gacttatcgt cctaactttt tttttaatta 540
acaaaaaaat cgaaaagaaa ataactctgtt aaatatTTTT taagtactcc attaatgtaa 600
gtttctatTT aaaaaatgct tgaaatttga cagttatgtt caacaatttt gaatcatgag 660
cgatgtctag atactcagaa tttaatcaag atgtcttata aaatttggtg tctactcgagg 720
acccacgcaa aagaaaagac taatatgatt tttatttggg ctggatatTT ttgtagagga 780
tgaaactaag agagtgaag attcgaaatc cacaatgttc aagagagctc aaagcaaaaa 840
gaaaaatgaa gatgaaggac taaagaacaa taagcaacta cttataaccct atttccataa 900
aggattcagg tactaggaga agttgaggca agttnnnnnn nattgattca aattttcatt 960
tattttttaca atttaattca cctaagttat tatgcatttc tcatcattgg tacattttct 1020
gtatagcgta ttacatatata tgaaataaat taaatatgtc ctcacgttgc aagtagttaa 1080
tgaatgtccc cacgcaaaaa aaaatccctc caaatatgtc caccttttct tttcttttta 1140
attccaaaat taccataaac ttttggttta caaaagattt ctagaaattg aggaagatat 1200
cctaaatgat tcatgaatcc ttcaataatc tgaagtttgc gatattttcg attttcttca 1260
agagttgcca tatttgtaat ttggtgacct taaacttttt ttgataaaga gtaaacgttt 1320
tttcttaaaa gtaaaacttg attttatgtt ttagggttct agctcaactt tgtattatat 1380
ttcttgcaaa aagagttcgt taactgcatt cttcaacact ataaagtgat tatcaaaaac 1440
atcttcatga acattaagaa aaacaatatt tggtttcggg tagagcttgg ttttgcttgg 1500
cttgattcac atacccattc tagactttgg cataaatttg atacgataga gagtatctaa 1560
tggtaatgca gaagggtaaa aaaaggaaga gagaaaaggt gagaaagatt accaaaaata 1620
aggagtttca aaagatgggt ctgatgagaa acagagccca tccctctcct tttccccttc 1680
ccatgaaaga aatcggtatg tctccttca atgtcctcca cctactcttc tcttctttct 1740
ttttttcttt ctattatta accatttaat taatttcccc ttcaatttca gtttctagtt 1800
ctgtaaaaag aaaatacaca tctcacttat agatatccat atctatttat atgcatgtat 1860
agagaataaaa aaagtgtgag tttctaggta tggttgagtat gtgctgtttg gacaattggt 1920
agatgatctg tccatttttt tcttttttct tctgtgtata aatatatttg agcaciaaga 1980
aaaactaata accttctgtt ttcagcaact agggctttat aaccttcaaa gaaatattcc 2040

```

ttcaattgaa	aacccataaa	ccaaaataga	tattacaaaa	ggaaagagag	atattttcaa	2100
gaacaacata	attagaaaag	cagaagcagc	agttaagtgg	tactgagata	aatgatatag	2160
tttctcttca	agaacagttt	ctcattaccc	accttctcct	ttttgctgat	ctatcgtaat	2220
cttgagaact	caggtaaggt	tgtgaatatt	atgcaccatt	cattaaccct	aaaaataaga	2280
gatttaaaat	aatgttttct	tctttctctg	attcttgtgt	aaccaattca	tgggtttgat	2340
atgtttcttg	gttattgctt	atcaacaaag	agatttgatc	attataaagt	agattaataa	2400
ctcttaaaca	cacaaagttt	ctttatTTTT	tagttacatc	cctaattcta	gaccagaaca	2460
tggatttgat	ctatttcttg	gttatgtatc	ttgatcagga	aaagggattt	gatcatcaag	2520
attagccttc	tctctctctc	tctagatata	tttcttgaat	ttagaaatct	ttatttaatt	2580
atTTGGTgat	gtcatatatg	gatcaatgga	ggaaggtggg	agtagtcacg	acgcagagag	2640
tagcaagaaa	ctagggagag	ggaaaataga	gataaagagg	atagagaaca	caacaaatcg	2700
tcaagttact	ttctgcaaac	gacgcaatgg	tcttctcaag	aaagcttatg	aactctctgt	2760
cttgtgtgat	gccgaagttg	ccctcgtcat	cttctccact	cgtggccgtc	tctatgagta	2820
cgccaacaac	aggtacgctt	ctcctactct	atttcttgat	cttgttttct	taattttaac	2880
taaacaagat	cctagttcaa	atgataacaa	agtggggatt	gagagccaag	attaggggtt	2940
ggTTaattta	gaaaaccaga	tttcaactgt	tgatacattt	aatatctctc	tagctagatt	3000
tagtactctc	tcctctatat	atgtgtgggt	gtgtgtgtaa	gtgtgtatat	gtatgcaaat	3060
gcaagaagaa	gaagaaaaag	ttatcttgtc	ttctcaaatt	ctgatcagct	ttgaccttag	3120
tttcaactct	ttttctgcaa	atcatttgaa	cctgatgcat	gtcagtttct	acaatacact	3180
tttaattttg	acggcccatc	aaatttccta	gggtttactt	cagtgaacaa	aattgggttc	3240
ttgacacgat	ttagcatgta	tataataaaa	taggggatga	tcaagactta	tgtaacctct	3300
gtctggTgaa	actagggaca	aagtctactg	atgagttgtc	actagggatc	catttgatca	3360
tttaatccca	acaaaaatga	aacaaaattt	tgagaattta	tatgctgaag	tttttcaacc	3420
ctctttttta	aataacttta	tattatgtag	atttgtattt	agggtaatTT	gtccaactag	3480
aagtcctaaa	aatcaataaa	cacacggatg	actttgtcta	acattgtatc	agtcatcaaa	3540
tgtaaaattg	tacaaataat	gaaattaaag	atttagtctc	ttttatTTTT	tttgtttagg	3600
gtgtatatata	atatatatata	gtatatTTgt	tgcattgata	tatcaatgag	agggagagaa	3660
ctcagagaag	tgtcggaaat	taaaatggta	cgagccaatt	ggaatctctg	gcattctgag	3720
cttcatttgt	ttgttatttag	aaaaaaaaaa	aaaaaatcct	ttaaagatac	cttcatgatg	3780
acattgaatc	atgtaatatata	cacgatacat	ggctctaattc	ctcctcaaac	cctaattacc	3840
aatttcgaaa	ccataatatt	tactagtatg	tttatatatc	cttactttta	gacattgttt	3900
gtttataata	ccttgtgaat	taagaaaaaa	aaaaaaaaaac	ttgtggatct	attcaagcca	3960
tgtgttagaa	taaatttata	aatttttctcc	tcgtactggg	cagatattgg	tccaaactcc	4020
aaagccttcc	cttttcagga	aaaaaaacat	ttcgaaatta	actctaatta	atcaagaatt	4080
tcctacaatg	tatacatcta	atgttttttc	cgcgatctta	cttatttagtg	tgaggggtac	4140
aattgaaagg	tacaagaaag	cttgttccga	tgccgtcaac	cctccttccg	tcaccgaagc	4200
taatactcag	gtaccaattt	atattgtttg	attctctttg	ttttatcttc	ttcttttcat	4260
tatatatatg	atcaacaaaa	aatataacct	acaaaaagag	agagttcaag	gaaatgcatt	4320
gaaacggttt	cgttatgggtg	tttgaataca	tggatttttg	aagtactatc	agcaagaagc	4380
ctctaagctt	cggaggcaga	ttcgagatat	tcagaattca	aataggtaat	tcattaactt	4440
ttcatgaact	cttcgatttg	gtattagggtc	acttaatttg	gtgtcgggtcc	aaaagtccgc	4500
ttgtagtttt	ctttagaagt	tgttttgttt	aatgttcatg	tttacaaatt	gaaggcatat	4560
tgttggggaa	tcacttggtt	ccttgaactt	caagggaactc	aaaaacctag	aaggacgtct	4620
tgaaaaagga	atcagccgtg	tccgctccaa	aaaggtaaaa	tctacgttgc	tctctctctg	4680
tgtctctgtc	tctctctcta	tatatagtcc	cttagtttat	atagttcatc	acctttttgt	4740
gagaattttg	cagaatgagc	tgttagtggtc	agagatagag	tatatgcaga	agagggtaag	4800
aacgtttctc	ccattccaag	taattagatc	tttcttcgtc	tttgtgaggg	tttgagtttt	4860
cccataaatc	atgtgtagga	aatggagttg	caacacaata	acatgtacct	gcgagcaaag	4920
gttagccacg	ttctgttcca	aatcttaatc	tcaatatcta	ctcttttctt	cattgtataa	4980
ctaagataac	gtgaataaca	agaaaacttt	tgtttttggg	tttaatagat	agccgaaggc	5040
gccagattga	atccggacca	gcaggaatcg	agtgtgatac	aaggggacgac	agtttacgaa	5100
tccggtgtat	cttctcatga	ccagtcgcag	cattataatc	ggaactatat	tccggtgaac	5160
cttcttgaac	cgaatcagca	attctccggc	caagaccaac	ctcctcttca	acttgtgtaa	5220
ctcaaaacat	gataacttgt	ttcttcccct	cataacgatt	aagagagaga	cgagagagtt	5280
catttttatat	ttataacgcg	actgtgtatt	catagtttag	gttctaataa	tgataataac	5340
aaaactgttg	tttcttttgct	taattacatc	aacattttaa	tccaaagtTC	taaaacacgt	5400
cgagatccaa	agtttgtcat	acaagattag	acgcatacac	gatcagttaa	tagatttttaa	5460
gtgcctttta	atattttacat	atagttgcag	cttcgatttag	atcatgtcca	ccaaacactc	5520
acaattagag	acaagcaaaa	ctataaacat	tgatcataaa	atgattacaa	catgtccata	5580
aattaattat	ggattacaaa	aataaaaact	tacaaaagat	ct		5622

<210> 6
 <211> 6138
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <223> SHATTERPROOF2 (SHP2) genomic

<400> 6
 gaattcgtaa cagaatttag tgaataatat tgtaattacc aggcaaggac tctccaaacg 60
 gatagctcga atatcgttat taaagagtaa atgatccaat atgtaagcca ttgttgatca 120
 tctaacattg ttggactctc tattgctcga aatgatgcat acctaatacat ttattcagtt 180
 aactatcaag ttgcatttgt aaaaacccaaa catttaaatt cagatttgat atcacttaca 240
 gaggatagag aagcatgact ccaggcctgc atgcaacaag aaaaaggaag aaaataatgt 300
 taaaaatttg acaaatatag tgtttatattt tattatatga gacagaattt gaataaaatc 360
 ctacccaact agagcatcaa aacgttttgc aatcgcaata atgaaaccca ttttcttttt 420
 gagttttttac tcttctttca acagaaactt tctcaaactg ctttagcact gtgacgtag 480
 atatatacac aaaagcttga aatttcttca agcaaaagaa tctttgtggg agttaaggca 540
 acaagccagg taaagaatct ccaacgcatt gttacgtttt catgaacctt tttattatat 600
 gttctaagaa agaaaaaaat atctcaaagt aaacgttggg aattttctga tgaagggaaa 660
 tccaaagtct tgggttttagt atccctatga atgggtattg gaatatgttt tcgtcaaaac 720
 aaaagattct tttctttttc acaagagtta gtgatcaata acttatgcac taattaatga 780
 gattggacgt atacacaatt tgattatgat acttgagtaa aaatcacctg tcctttaatt 840
 tggaaatctc tctttcttac ccattttatat actacttctt ttcattaaaa ttaaatttca 900
 attatcaatc atcgttcaat ttgataaaga ttttaacattt tttgtcacag ggctagtaaa 960
 agcaatcttt acataattca tctttcttac atatataat tacctttttc ttcattagta 1020
 ttctatttga ttatgattat tttgtcataa agctagtaaa ttaaactc gatatgagaa 1080
 ttatattact tcacgctaata taactcttaa cacaacaaga actagtgcac attcaacttt 1140
 caaagcatat actatatatt gagaatatag accacgaaag tcaatcaaaa gacctaccag 1200
 ctctcatcaa gttctttctt gaaatgattt tgcagaattt ccaacttaat taattcgaca 1260
 tgaatgtgaa aatgtgtgtt gctcgttaag aaaattgaat agaagtacaa tgaaaatgat 1320
 gaggaatggg caaaacacaa aagagtttcc tttcgttaact acaattaatt aatgcaaact 1380
 tgagaaaggg ttcatggata atgactacac acatgattag tcattccccg tgggctctct 1440
 gctttcattt actttattag tttcatcttc tctaattata ttgtcgcata tatgatgcag 1500
 ttcttttgtc taaattacgt aatatgatgt aattaattat caaaataaat attcaaattg 1560
 ccgttggact aacctaatgt ccaagattaa gacttgaaca taagaatttt ggaaaaacta 1620
 aaccagttat aatatatact cttaaattgc catttctgaa cacaaccaa taataatata 1680
 tactatttac agtttttttt aattggcaag aacactgaaa tcttattcat tgtctcgctt 1740
 ggtagttgac aagttataac actcatattc atataacccc attctaactg tgacgacgaa 1800
 cactcatata aaccacccaa attcttagca tattagctaa atattgggtt aattggaaat 1860
 atttttttta tatataaaat gccaggtaaa tattaacgac atgcaatgta tataggagta 1920
 gggcaataaa aagaaaagga gaataaaaag ggattaccaa aaaaggaaag tttccaaaag 1980
 gtgattctga tgagaaacag agcccatacc tctctttttt cctctaaaca tgaaagaaaa 2040
 attggatggg cctccttcaa tgctctctcc ccacccaatc caaacccaac tgtcttcttt 2100
 ctttcttttt tcttctttct aatttgatat tttctaccac ttaattccaa tcaatttcaa 2160
 atttcaatct aaatgtatgc atatagaatt taattaaaag aattagggtg gtgatatttg 2220
 agaaaatgtt agaagtaatg gtccatgttc tttctttctt tttccttcta taacacttca 2280
 gtttgaaaaa aaactaccaa accttctgtt ttctgcaaat ggggtttttta atacttccaa 2340
 agaaatattc ctctaaaaga aattataaac caaacagaa accaaaaaca aaaaataaag 2400
 ttgaagcagc agttaagtgg tactgagata ataagaatag tatcttttagg ccaatgaaca 2460
 aattaactct ctcataattc atcttcccat cctcacttct ctttctttct gatataatta 2520
 atcttgctaa gccagggtatg gttattgatg atttacactt ttttttaaaa gtttcttctt 2580
 tttctccaat caaattcttc agttaatcct tataaaccat ttctttaatc caaggtgttt 2640
 gagtgcaaaa ggatttgatc tatttctctt gtgttttatac ttcagctagg gcttatagaa 2700
 atggaggggtg gtgcgagtaa tgaagtagca gagagcagca agaagatagg gagagggag 2760
 atagagataa agaggataga gaacactacg aatcgtaag tcactttctg caaacgacgc 2820
 aatggtttac tcaagaaagc ttatgagctc tctgtcttgt gtgacgctga ggttgctctt 2880
 gtcacttctt ccactcgagg ccgtctctac gagtacgcca acaacaggta cacatctttt 2940
 agctagatct tgattttgtt gaattttttt tctagaataa agtttcgact cttctgggtg 3000
 gtttttcaat ctttatggtc tctttatagt ttttttctt agtttctctg aagctcaaat 3060

```

ctctttaaaa atccccaaaa ttaggggtttg tttaaaacta gggaacccta ctttaacttc 3120
tttctcttag taaaaaagca gtgaggggtct tctctgatca ttaattagca tccccatac 3180
cttgttccag tcactttttc tccacaaatc cttataacag tatctatata tgtatctatt 3240
tatgtcagtt tgtacaagac acttcgatca atttgatgac ccatcaagtt ttatttctgc 3300
agattgatca ttaggtttcc atcatagtaa tgaaaaagta gggttcttga taaaattata 3360
ataatatata ttatttggct atataaaaaa gctatgtaga ttccttaaaa attgattcac 3420
tagggagaga ctagtaggtg tttgtcttct gacacttctc taatcttttg gtgaatcctt 3480
ttgttaaate aagaaaatga atcagggaca aagcttattg ttgagtcact taattaatca 3540
tccgatccat caatcaagaa aaataacgaa acagaaaatt ttgatttttg attgttattt 3600
tctccacttc aagttgggga cttgtcattt ccgtttttct atacgtttcc agctattaac 3660
agctcatgtt catttcacca ttttgattat ttgtctgctt tttaaagata aatgttttca 3720
aaaatattgt ttttatttgc ttggctagtt aatactataa ttgaggttga tgtatgacta 3780
taatctataa gtcaagtctc atatcatgga tctaagttaa aactagtaaa tttgtagttt 3840
caatgtgaac tttcacaacg actaaagaac tgatctgaag tttataatgg acatgactaa 3900
tttgattaac aaaagaggaa tgcattatgt atgtagaaac atgtgatata tatatgtttc 3960
tattatcaaa agtgtagtta actttcttat ttcaaaccac ctcatgcttt agtagtatct 4020
tacttttgac atttctcaac ttcagctttc cattatacaa cagcacaatg taaattactt 4080
gtatatgaat atgaaagcat aacgttatgc aaagatttct agcttttctt tttctgtttt 4140
gcaaaagatt taaaaatate atgttcttgg taaaaacata cttgcctcag ccacatatgc 4200
atgtaaategt aatgttcaaa tattaattca ggaaaaacaa agaagaagca aaattagctt 4260
ctagagtagg gaatctattg acttgacctg aaaatcactt ctttttctta aagcctagta 4320
gtgaattttt taatctaatt aggccaaaat atatactagc ctaaaatata atttggattt 4380
tgtgtcgtac ataaattggg accaattcca attaactaag agcatatgca attcaaattc 4440
tttttatttt cttctccgat ttgctacttc tttcttttgt atgttttcaa attaggatta 4500
cacttttttg gggaagtaca cattaggggtc ttctcgaact ttgattatac atatatatat 4560
atatatatat atataacttt gtgagatgtc actgttaata gataataggc aataacaata 4620
atatccaaaa aagaaggcgc aaacaaatca tatactatat ggtactgggtc cattcactat 4680
tttgtcgggt gaatttaagg tttggcgtac aaactttgtt tcaaaccctt attattccgt 4740
ctttctgtgt gttttgtata tccagaagat aaaaatatca atttctttaa cgacttcata 4800
tatatatata tatatatata tatatatatt tttctcttct ggttttagtg tttgaatcca 4860
acagttatag tttcgtgtgt ctttggtttta cttgtgggtg ttttaagttt agattttcac 4920
cgattgcate tatttacata tatagctacc acaaaaaaga ttgcatttta aaatcttttc 4980
ctttgtgtga atgttgatga agtgtgagag gaacaataga aaggtacaag aaagcttgct 5040
ccgacgccgt taaccctccg accatcaccg aagctaatac tcaggttagc ttttaattaa 5100
tacacctagc tagctagttc gttaattact taatttcttc ttcttttagt tatctgacct 5160
ttttttcacc tcttgtaaca atgatgggat cgaaattgat gaagtactat cagcaagagg 5220
cgtctaaact ccggagacag attcgggaca ttcagaattt gaacagacac attcttgggt 5280
aatctcttgg ttccttgaac ttttaaggaac tcaagaacct tgaaagtagg cttgagaaaag 5340
gaatcagtcg tgtccgatcc aagaaggtag atcactaact ctccatcaat ctcttatca 5400
ttgaatatat atccatctga ttcttgcccg ttatatattg tttttctctc cagcacgaga 5460
tgttagttgc agagattgaa tacatgcaaa aaagggtaaa agtaaaacct atcttccttc 5520
acaatgaact accctactt tattagcaac ttctctttct gatgatcatc ttttttattt 5580
tctgttgctg cttgcattgt aggaaatcga gctgcaaaac gataacatgt atctccgctc 5640
caaggtttta tacataactc tttttggcat ttttgatcat catttttttc cggtagacaa 5700
tctcttgatg tgcaaattct aaatatctct gcagattact gaaagaacag gtctacagca 5760
acaagaatcg agtgtgatac atcaagggac agtttacgag tcgggtgtta cttcttctca 5820
ccagtcgggg cagtataacc ggaattatat tgcggttaac cttcttgaac cgaatcagaa 5880
ttcctccaac caagaccaac cacctctgca acttgtttga ttcagtctaa cataagcttc 5940
tttctcagc ctgagatcga tctatagtgt cacctaaatg cggccgcgtc cctcaacatc 6000
tagtcgcaag ctgaggggaa ccactagtgt catacgaacc tccaagagac ggttacacaa 6060
acgggtacat tgttgatgtc atgtatgaca atcgcccaag taagtatcca gctgtgttca 6120
gaacgtacgt ccgaattc

```

<210> 7

<211> 51

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:IND1 basic
helix-loop-helix (bHLH) DNA binding domain

<400> 7

Ile	Ser	Asp	Asp	Pro	Gln	Thr	Val	Val	Ala	Arg	Arg	Arg	Arg	Glu	Arg
1				5					10					15	
Ile	Ser	Glu	Lys	Ile	Arg	Ile	Leu	Lys	Arg	Ile	Val	Pro	Gly	Gly	Ala
		20					25						30		
Lys	Met	Asp	Thr	Ala	Ser	Met	Leu	Asp	Glu	Ala	Ile	Arg	Tyr	Thr	Lys
		35					40					45			
Phe	Leu	Lys													
	50														

<210> 8

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer for
amplification of IND1 genomic region of cDNA

<400> 8

gatgaaaatg gaaaatggta tgtata

26

<210> 9

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer for
amplification of IND1 genomic region of cDNA

<400> 9

gttcatcagg gttgggagtt gtg

23

Claims:

1. An expression cassette comprising a promoter operably linked to an IND1 polynucleotide sequence, or a complement thereof, encoding an IND1 polypeptide, wherein the IND1 polypeptide is at least about 70% identical to SEQ ID NO: 2; the IND1 polypeptide comprises a basic helix-loop-helix (bHLH) domain; and wherein introduction of the expression cassette into a plant to suppress IND1 expression results in a plant with delayed fruit dehiscence.
2. The expression cassette of claim 1, wherein the IND1 polypeptide comprises SEQ ID NO: 2.
3. The expression cassette of claim 1, wherein the IND1 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO: 1.
4. The expression cassette of claim 3, wherein the IND1 polynucleotide comprises SEQ ID NO: 1.
5. An expression cassette comprising a promoter operably linked to a polynucleotide sequence, or a complement thereof, wherein the polynucleotide sequence is at least 60% identical to at least 200 contiguous nucleotides of a sequence encoding SEQ ID NO: 2.
6. The expression cassette of claim 5, wherein the sequence is SEQ ID NO: 1.
7. The expression cassette of claim 5 wherein the polynucleotide is at least 60% identical to a nucleotide sequence encoding SEQ ID NO: 2.
8. The expression cassette of claims 1 to 7, wherein the promoter is constitutive.
9. The expression cassette of claims 1 to 7, wherein the promoter is tissue specific.
10. The expression cassette of claim 9, wherein the promoter is a dehiscence zone specific promoter.

11. The expression cassette of claim 10, wherein the promoter comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO: 1.
12. A plant cell comprising the recombinant expression cassette of any one of claims 1 to 4.
13. A plant cell comprising the recombinant expression cassette of any one of claims 5 to 7.
14. The plant cell of claim 12 or 13, wherein the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the antisense orientation.
15. The plant cell of claim 12 or 13, wherein the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the sense orientation.
16. The plant cell of claim 15, wherein the polynucleotide sequence further comprises a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation.
17. The plant cell of claim 16, wherein the plant further comprises a second polynucleotide at least 60% identical to at least 200 contiguous nucleotides of a sequence encoding SEQ ID NO:2, wherein the second polynucleotide is operably linked to a second promoter in the antisense orientation.
18. The plant cell of any of claims 12 to 17, wherein lignification is reduced in valve margin cells of the plant compared to lignification in valve margins cells of a plant lacking the expression cassette.
19. The plant cell of any of claims 12 to 18, wherein a plant comprising the plant cell is characterized by delayed fruit dehiscence or delayed seed dispersal compared to a plant lacking the expression cassette.

20. The plant cell of any of claims 12 to 19, wherein the promoter is a dehiscence zone selective regulatory element.
21. The plant cell of claim 20, where in the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO: 1.
22. A method of delaying fruit dehiscence in a plant, the method comprising suppressing expression of an IND1 nucleic acid in the plant by introducing into the plant a recombinant expression cassette comprising a promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO : 2, wherein the IND1 polypeptide comprises a basic helix-loop-helix (bHLH) domain.
23. The method of claim 22, wherein the IND1 polypeptide comprises SEQ ID NO: 2.
24. The method of claim 22, wherein the IND1 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO: 1.
25. The method of claim 22, wherein the IND1 polynucleotide comprises SEQ ID NO: 1.
26. The method of claim 22, wherein the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the antisense orientation.
27. The method of claim 22, wherein the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the sense orientation.
28. The method of claim 27, wherein the polynucleotide sequence further comprises a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation.
29. The method of claim 22, wherein lignification is reduced in valve margin cells of the plant compared to lignification in valve margins cells of a plant lacking the expression cassette.

30. The method of claim 22, wherein the promoter is a dehiscence zone-selective regulatory element.
31. The method of claim 30, where in the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO: 1.
32. The method of claim 22, wherein the recombinant expression cassette is introduced into the plant using *Agrobacterium*.
33. A method of delaying fruit dehiscence in a plant, the method comprising:
- (i) introducing into the plant a recombinant expression cassette comprising a polynucleotide sequence at least about 70% identical to positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO: 1 thereby suppressing the expression of an IND1 gene or a polypeptide at least 60% identical to SEQ ID NO:2 in the plant; and
 - (ii) selecting a plant with delayed fruit dehiscence compared to a plant in which expression of the IND1 gene or the polypeptide is not suppressed.
34. The method of claim 33, wherein the polynucleotide sequence comprises positions from about 1 to 2764 or from about 3362 to about 3856 of SEQ ID NO: 1.
35. The method of claim 33, wherein lignification is reduced in valve margin cells of the plant compared to lignification in valve margins cells of a plant lacking the expression cassette.
36. A method for delaying fruit dehiscence in a plant, the method comprising the steps of:
- (i) treating seeds or plant material with a mutagenic chemical substance or with ionizing radiation;
 - (ii) identifying plants with a mutated IND1 gene, wherein the IND1 gene, prior to being mutated, encodes a polypeptide having at least 60% sequence identity to SEQ ID NO:2; and

(iii) selecting a plant with delayed fruit dehiscence compared to a plant in which the IND1 gene is not mutated.

37. A plant cell comprising a modified polynucleotide at least 65% identical to SEQ ID NO: 1, wherein the plant displays delayed fruit dehiscence compared to a plant lacking the modified polynucleotide.

38. The plant cell of claim 37 wherein the modified polynucleotide is at least 70% identical to SEQ ID NO: 1.

39. A method of creating a plant with delayed fruit dehiscence, the method comprising introducing into said plant a recombinant expression cassette comprising a promoter operably linked to a polynucleotide that is at least 60% identical to at least 200 contiguous nucleotides of SEQ ID NO: 1 and selecting a plant with delayed fruit dehiscence compared to a plant lacking the recombinant expression cassette.

40. The method of claim 39 comprising expressing in the plant a sense polynucleotide at least 65% identical to at least 200 contiguous nucleotides of SEQ ID NO: 1 and expressing in the plant an antisense polynucleotide at least 65% identical to at least 200 contiguous nucleotides of SEQ ID NO: 1.

