




(86) Date de dépôt PCT/PCT Filing Date: 2011/04/25
(87) Date publication PCT/PCT Publication Date: 2011/11/03
(45) Date de délivrance/Issue Date: 2018/04/10
(85) Entrée phase nationale/National Entry: 2012/10/24
(86) N° demande PCT/PCT Application No.: JP 2011/002418
(87) N° publication PCT/PCT Publication No.: 2011/135835
(30) Priorité/Priority: 2010/04/28 (JP2010-104095)

(51) Cl.Int./Int.Cl. *A01N 43/56* (2006.01),
A01N 37/18 (2006.01), *A01N 37/50* (2006.01),
A01N 39/02 (2006.01), *A01N 43/40* (2006.01),
A01N 43/54 (2006.01), *A01N 47/24* (2006.01),
A01P 3/00 (2006.01)
(72) Inventeur/Inventor:
MATSUZAKI, YUICHI, JP
(73) Propriétaire/Owner:
SUMITOMO CHEMICAL COMPANY, LIMITED, JP
(74) Agent: MARKS & CLERK

(54) Titre : COMPOSITIONS DE CONTROLE DE MALADIE DES VEGETAUX RENFERMANT UN COMPOSE
CARBOXAMIDE ET UN FONGICIDE STROBILURINE
(54) Title: PLANT DISEASE CONTROLLING COMPOSITIONS COMPRISING A CARBOXAMIDE COMPOUND AND A
STROBILURIN FUNGICIDE



(57) Abrégé/Abstract:

A plant disease control composition comprising a carboxamide compound represented by following formula (I), wherein R1 represents a hydrogen atom or a methyl group, and R2 represents a methyl group, a difluoromethyl group or a trifluoromethyl group, and one or more QoI compounds selected from group (A) consisting of dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin and N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is provided by the present invention, and this composition has excellent effect for controlling a plant disease.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 November 2011 (03.11.2011)(10) International Publication Number
WO 2011/135835 A1

(51) International Patent Classification:

A01N 43/56 (2006.01) *A01N 43/40* (2006.01)
A01N 37/18 (2006.01) *A01N 43/54* (2006.01)
A01N 37/50 (2006.01) *A01N 47/24* (2006.01)
A01N 39/02 (2006.01) *A01P 3/00* (2006.01)

(21) International Application Number:

PCT/JP2011/002418

(22) International Filing Date:

25 April 2011 (25.04.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2010-104095 28 April 2010 (28.04.2010) JP

(71) Applicant (for all designated States except US): **SUMITOMO CHEMICAL COMPANY, LIMITED** [JP/JP];
27-1, Shinkawa 2-chome, Chuo-ku, Tokyo, 1048260 (JP).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MATSUZAKI, Yuichi** [JP/JP]; 8-7-20-318, Hon-machi, Toyonaka-shi, Osaka, 5600021 (JP).(74) Agents: **NOGAWA, Shintaro** et al.; Nogawa Patent Office, Nishitenma Five Bldg., 16-3, Nishitenma 5-Chome, Kita-ku, Osaka-shi, Osaka, 5300047 (JP).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: Plant disease control composition and its use

(57) Abstract: A plant disease control composition comprising a carboxamide compound represented by following formula (I), wherein R¹ represents a hydrogen atom or a methyl group, and R² represents a methyl group, a difluoromethyl group or a trifluoromethyl group, and one or more QoI compounds selected from group (A) consisting of dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin and N-methyl- α -methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is provided by the present invention, and this composition has excellent effect for controlling a plant disease.

Description

Title of Invention: Plant Disease Controlling Compositions Comprising a Carboxamide Compound and a Strobilurin Fungicide

Technical Field

[0001] The present invention relates to a plant disease control composition and its use.

Background Art

[0002] Many compounds have been developed for controlling plant diseases and actually used (see, for example, PTL 1 and 2).

Citation List

Patent Literature

[0003] [PTL 1] : WO86/02641
[PTL 2] : WO92/12970

Summary of Invention

Technical Problem

[0004] An object of the present invention is to provide a composition having an excellent effect for controlling plant disease.


Solution to Problem

[0005] The inventor of the present invention studied for seeking a composition having an excellent effect for controlling plant disease and found that a composition comprising a carboxamide compound represented by the following formula (I) and one or more QoI compounds selected from following group (A) has an excellent effect for plant diseases and then completed the present invention.

The present invention provides the following [1] to [5].

[1] A plant disease control composition comprising a carboxamide compound represented by formula (I):

[Chem.1]



wherein

R¹ represents a hydrogen atom or a methyl group, and


R² represents a methyl group, a difluoromethyl group or a trifluoromethyl group, and one or more QoI compounds selected from group (A) consisting of dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, pi-

coxystrobin, trifloxystrobin and N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide.

[2] The plant disease control composition according to above [1], wherein the weight ratio of the carboxamide compound to the QoI compound(s) is from 0.1/1 to 10/1 of the carboxamide compound / the QoI compound(s).

[3] A method of controlling plant disease which comprises a step of treating a plant or the soil where a plant grows with an effective amount of a carboxamide compound represented by formula (I):

[Chem.2]



wherein


R¹ represents a hydrogen atom or a methyl group, and

R² represents a methyl group, a difluoromethyl group or a trifluoromethyl group, and one or more QoI compounds selected from group (A) consisting of dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin and N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide.

[4] The method of controlling plant disease according to above [3], wherein the weight ratio of the carboxamide compound to the QoI compound(s) is from 0.1/1 to 10/1 of the carboxamide compound / the QoI compound(s).

[5] The method of controlling plant disease according to above [3] or [4], wherein the plant or the soil where a plant grows is soybean or the soil where soybean grows, respectively.

According to one aspect of the invention there is provided a composition for controlling a plant fungal disease comprising carboxamide compound (1):



and a QoI compound which is dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin, N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide, or any combination thereof; wherein the weight ratio of the carboxamide compound (1) to the QoI compound is from 0.1/1 to 10/1.

According to a further aspect of the invention there is provided a method of controlling a plant fungal disease which comprises the step of treating a plant, or soil where a plant grows, with a composition as described herein.


Advantageous Effect of Invention

[0006] According to the present invention, various plant diseases can be controlled.

Description of Embodiments

[0007] The plant disease control composition of the present invention (hereinafter referred to as "composition") comprises a carboxamide compound represented by formula (I):

[Chem.3]



wherein

R¹ and R² represent the same meanings as defined in the above (hereinafter "carboxamide compound")


and one or more QoI compounds selected from group (A) consisting of dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin and N-

methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (hereinafter referred to as "QoI compound").

[0008] The "carboxamide compound" are those as described in, for example, WO86/02641 or WO92/12970, and can be prepared by the method described therein.

[0009] Particular examples of the "carboxamide compounds" are as follows:
carboxamide compound represented by formula (1):


[Chem.4]



(hereinafter referred to as "carboxamide compound (1)");

carboxamide compound represented by formula (2):


[Chem.5]



(hereinafter referred to as "carboxamide compound (2)");

carboxamide compound represented by formula (3):


[Chem.6]



(hereinafter referred to as "carboxamide compound (3)");

carboxamide compound represented by formula (4):


[Chem.7]



(hereinafter referred to as "carboxamide compound (4)");

carboxamide compound represented by formula (5):

[Chem.8]



(hereinafter referred to as "carboxamide compound (5)").

[0010] The "QoI compounds" are known compounds and described in, for example, "THE PESTICIDE MANUAL - 14th EDITION (published by BCPC) ISBN 1901396142 and WO 95/27693. These compounds can be obtained from the products containing said compound in the market or can be synthesized by publicly known methods.

[0011] The weight ratio of the "carboxamide compound" to the "QoI compound(s)" in the "composition" is usually from 0.01/1 to 500/1, and preferably from 0.1/1 to 10/1 of "carboxamide compound" / "QoI compound(s)"

[0012] Although the "composition" may be a mixture itself of a "carboxamide compound" and "QoI compound(s)", the "composition" is usually prepared by mixing a "carboxamide compound", "QoI compound(s)" and an inert carrier, and if necessary, by adding a surfactant and/or another auxiliary for formulation and by formulating the mixture into oil formulation, emulsifiable concentrate, flowable formulation, wettable powder, water dispersible granules, powder, granules, or the like. The formulation ,

which is used alone or by adding another inert component, can be used as a plant disease control agent.

The total content of a "carboxamide compound" and "QoI compound(s)" in a "composition" is usually from 0.1% to 99% by weight, preferably from 0.2% to 90% by weight, and more preferably from 1% to 80% by weight.

[0013] Examples of the solid carriers used for the formulation include fine powder or granules of, for example, mineral materials such as kaolin clay, attapulgite, bentonite, montmorillonite, acid clay, pyrophyllite, talc, diatomaceous earth and calcite; natural organic materials such as corncob powder and walnut powder; synthesized organic materials such as urea; salts such as potassium carbonate and ammonium sulfate; synthetic inorganic materials such as synthesized hydrous silicon oxide.

Examples of the liquid carriers include aromatic hydrocarbons such as xylene, alkylbenzene and methylnaphthalene; alcohols such as 2-propanol, ethylene glycol, propylene glycol and ethylene glycol mono-ethyl ether; ketones such as acetone, cyclohexanone and isophorone; vegetable oils such as soybean oil and cotton seed oil; petrolic aliphatic hydrocarbons; esters; dimethylsulfoxide; acetonitrile; and water

Examples of the surfactants include anionic surfactants such as alkyl sulfate ester salts, alkylarylsulfonate salts, dialkylsulfosuccinate salts, polyoxyethylene alkylaryl ether phosphoric acid ester salts, lignin sulfonate and naphthalene sulfonate formaldehyde polycondensed products; non-ionic surfactants such as polyoxyethylene alkyl aryl ethers, polyoxyethylene alkyl polyoxypropylene block copolymers and sorbitan fatty acid esters; and cationic surfactants such as alkyl trimethyl ammonium salts.

Examples of the other auxiliaries for formulation include water-soluble polymers such as polyvinyl alcohol and polyvinylpyrrolidone; polysaccharides such as gum arabic, alginic acid and its salt, CMC (carboxymethylcellulose) and xanthan gum; inorganic materials such as aluminum magnesium silicate and alumina sol; preservatives; coloring agents; and stabilizers such as PAP (acidic isopropyl phosphate) and BHT.

[0014] The "composition" can be also prepared by formulating a "carboxamide compound" and "QoI compound(s)" according to the method as described in the above, and then making the formulations or their diluents.

[0015] The "composition" can be used for protecting a plant from a plant disease.

[0016] Example of plant diseases which can be controlled by the "composition" include the followings.

Rice diseases: Magnaporthe grisea, Cochliobolus miyabeanus, Rhizoctonia solani, Gibberella fujikuroi;

Wheat diseases: Erysiphe graminis, Fusarium graminearum, F. avenaceum, F.

culmorum, *Microdochium nivale*, *Puccinia striiformis*, *P. graminis*, *P. recondita*, *Micronectriella nivale*, *Typhula* sp., *Ustilago tritici*, *Tilletia caries*, *Pseudocercospora herpotrichoides*, *Mycosphaerella graminicola*, *Stagonospora nodorum*, *Pyrenophora tritici-repentis*;

Barley diseases: *Erysiphe graminis*, *Fusarium graminearum*, *F. avenaceum*, *F. culmorum*, *Microdochium nivale*, *Puccinia striiformis*, *P. graminis*, *P. hordei*, *Ustilago nuda*, *Rhynchosporium secalis*, *Pyrenophora teres*, *Cochliobolus sativus*, *Pyrenophora graminea*, *Rhizoctonia solani*;

Maize diseases: *Ustilago maydis*, *Cochliobolus heterostrophus*, *Gloeocercospora sorghi*, *Puccinia polysora*, *Cercospora zeae-maydis*, *Rhizoctonia solani*;

[0017] Citrus diseases: *Diaporthe citri*, *Elsinoe fawcetti*, *Penicillium digitatum*, *P. italicum*, *Phytophthora parasitica*, *Phytophthora citrophthora*;

Apple diseases: *Monilinia mali*, *Valsa ceratosperma*, *Podosphaera leucotricha*, *Alternaria alternata* apple pathotype, *Venturia inaequalis*, *Colletotrichum acutatum*, *Phytophthora cactorum*;

Pear diseases: *Venturia nashicola*, *V. pirina*, *Alternaria alternata* Japanese pear pathotype, *Gymnosporangium haraeum*, *Phytophthora cactorum*;

Peach diseases: *Monilinia fructicola*, *Cladosporium carpophilum*, *Phomopsis* sp.;

Grape diseases: *Elsinoe ampelina*, *Glomerella cingulata*, *Uninula necator*, *Phakopsora ampelopsidis*, *Guignardia bidwellii*, *Plasmopara viticola*;

Persimmon diseases: *Gloesporium kaki*, *Cercospora kaki*, *Mycosphaerella nawae*;

Gourd diseases: *Colletotrichum lagenarium*, *Sphaerotheca fuliginea*, *Mycosphaerella melonis*, *Fusarium oxysporum*, *Pseudoperonospora cubensis*, *Phytophthora* sp., *Pythium* sp.;

Tomato diseases: *Alternaria solani*, *Cladosporium fulvum*, *Phytophthora infestans*;

Eggplant diseases: *Phomopsis vexans*, *Erysiphe cichoracearum*;

Brassicaceous vegetable diseases: *Alternaria japonica*, *Cercospora brassicae*, *Plasmodiophora brassicae*, *Peronospora parasitica*;

Welsh onion diseases: *Puccinia allii*, *Peronospora destructor*;

[0018] Soybean diseases: *Cercospora kikuchii*, *Elsinoe glycines*, *Diaporthe phaseolorum* var. *sojae*, *Septoria glycines*, *Cercospora sojina*, *Phakopsora pachyrhizi*, *Phytophthora sojae*, *Rhizoctonia solani*, *Corynespora asiicola*, *Sclerotinia sclerotiorum*;

Kidney bean diseases: *Colletotrichum lindemthianum*;

Peanut diseases: *Cercospora personata*, *Cercospora arachidicola*, *Sclerotium rolfsii*;

Pea diseases: *Erysiphe pisi*;

Potato diseases: *Alternaria solani*, *Phytophthora infestans*, *Phytophthora erythroseptica*, *Spongospora subterranean*, f. sp. *Subterranean*;

Strawberry diseases: *Sphaerotheca humuli*, *Glomerella cingulata*;

Tea diseases: *Exobasidium reticulatum*, *Elsinoe leucospila*, *Pestalotiopsis* sp., *Colletotrichum theae-sinensis*;

Tobacco diseases: *Alternaria longipes*, *Erysiphe cichoracearum*, *Colletotrichum tabacum*, *Peronospora tabacina*, *Phytophthora nicotianae*;

Rapeseed diseases: *Sclerotinia sclerotiorum*, *Rhizoctonia solani*;

Cotton diseases: *Rhizoctonia solani*;

Beet diseases: *Cercospora beticola*, *Thanatephorus cucumeris*, *Thanatephorus cucumeris*, *Aphanomyces cochlioides*;

Rose diseases: *Diplocarpon rosae*, *Sphaerotheca pannosa*, *Peronospora sparsa*;

Diseases of chrysanthemum and asteraceae: *Bremia lactuca*, *Septoria chrysanthemi-indici*, *Puccinia horiana*;

Diseases of various plants: *Pythium aphanidermatum*, *Pythium debarianum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium ultimum*, *Botrytis cinerea*, *Sclerotinia sclerotiorum*;

Radish diseases: *Alternaria brassicicola*;

Zoysia diseases: *Sclerotinia homeocarpa*, *Rhizoctonia solani*;

Banana diseases: *Mycosphaerella fijiensis*, *Mycosphaerella musicola*;

Sunflower diseases: *Plasmopara halstedii*;

Seed diseases or diseases in the initial stage of growth of various plants caused by *Aspergillus* spp., *Penicillium* spp., *Fusarium* spp., *Gibberella* spp., *Trichoderma* spp., *Thielaviopsis* spp., *Rhizopus* spp., *Mucor* spp., *Corticium* spp., *Rhoma* spp., *Rhizoctonia* spp., *Diplodia* spp., or the like;

Virus diseases of various plants mediated by *Polymixa* spp., *Olpidium* spp. or the like.

[0019] Examples of the plants for which the "composition" can be used are as follows:

Agricultural crops: maize, rice, wheat, barley, rye, oat, sorghum, cotton, soybean, peanut, backwheat, sugar beet, rapeseed, sunflower, sugar cane, tobacco, and the like;

Vegetables: Solanaceous vegetables (eggplant, tomato, green pepper, hot pepper, potato, etc.), Cucurbitaceous vegetables (cucumber, pumpkin, zucchini, watermelon, melon, squash, etc.); Cruciferous vegetables (radish, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, brown mustard, broccoli, cauliflower, etc.), Asteraceous vegetables (burdock, garland chrysanthemum, artichoke, lettuce, etc.), Liliaceous vegetables (Welsh onion, onion, garlic, asparagus, etc.), Umbelliferous vegetables (carrot, parsley, celery, parsnip, etc.), Chenopodiaceous vegetables (spinach, chard, etc.), Lamiaceous vegetables (Japanese basil, mint, basil, etc.), strawberry, sweet potato, yam, aroid, and the like;

Flowering plants;

Ornamental foliage plants;

Turf;

Fruit trees: pome fruits (apple, common pear, Japanese pear, Chinese quince, quince, etc.), stone fruits (peach, plum, nectarine, Japanese plum, cherry, apricot, prune, etc.), citrus (mandarin, orange, lemon, lime, grapefruit, etc.), nuts (chestnut, walnut, hazel nut, almond, pistachio, cashew nut, macadamia nut, etc.), berry fruits (blueberry, cranberry, blackberry, raspberry, etc.), grape, persimmon, olive, loquat, banana, coffee, date, coconut palm, and the like;

Trees other than fruit trees: tea, mulberry, flowering trees, street trees (ash tree, birch, dogwood, eucalyptus, ginkgo, lilac, maple tree, oak, poplar, cercis, Chinese sweet gum, plane tree, zelkova, Japanese arborvitae, fir tree, Japanese hemlock, needle juniper, pine, spruce, yew), and the like.

[0020] The above-described plants may be those having resistance imparted by genetic engineering technique.

[0021] Among the above plants, the "composition" is expected to have excellent controlling effect particularly to plant diseases caused in soybean.

Among the above plant diseases, soybean diseases to which especially excellent effect of the "composition" can be expected are *Rhizoctonia solani*, *Cercospora kikuchii*, *Septoria glycines*, *Corynespora asiicola*, *Phakopsora pachyrizi*, *Sclerotinia sclerotiorum*, *Cercospora sojina*, and the like.

[0022] Following compositions exemplify an embodiment of the "composition":
 a composition comprising "carboxamide compound (1)" and dimoxystrobin;
 a composition comprising "carboxamide compound (1)" and azoxystrobin;
 a composition comprising "carboxamide compound (1)" and fluoxastrobin;
 a composition comprising "carboxamide compound (1)" and pyraclostrobin;
 a composition comprising "carboxamide compound (1)" and kresoxim-methyl;
 a composition comprising "carboxamide compound (1)" and picoxystrobin;
 a composition comprising "carboxamide compound (1)" and trifloxystrobin;
 a composition comprising "carboxamide compound (1)" and N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide;
 a composition comprising "carboxamide compound (2)" and dimoxystrobin;
 a composition comprising "carboxamide compound (2)" and azoxystrobin;
 a composition comprising "carboxamide compound (2)" and fluoxastrobin;
 a composition comprising "carboxamide compound (2)" and pyraclostrobin;
 a composition comprising "carboxamide compound (2)" and kresoxim-methyl;
 a composition comprising "carboxamide compound (2)" and picoxystrobin;
 a composition comprising "carboxamide compound (2)" and trifloxystrobin;
 a composition comprising "carboxamide compound (2)" and N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide;
 a composition comprising "carboxamide compound (3)" and dimoxystrobin;

a composition comprising "carboxamide compound (3)" and azoxystrobin;
 a composition comprising "carboxamide compound (3)" and fluoxastrobin;
 a composition comprising "carboxamide compound (3)" and pyraclostrobin;
 a composition comprising "carboxamide compound (3)" and kresoxim-methyl;
 a composition comprising "carboxamide compound (3)" and picoxystrobin;
 a composition comprising "carboxamide compound (3)" and trifloxystrobin;
 a composition comprising "carboxamide compound (3)" and N-methyl-alpha -
 methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide;
 a composition comprising "carboxamide compound (4)" and dimoxystrobin;
 a composition comprising "carboxamide compound (4)" and azoxystrobin;
 a composition comprising "carboxamide compound (4)" and fluoxastrobin;
 a composition comprising "carboxamide compound (4)" and pyraclostrobin;
 a composition comprising "carboxamide compound (4)" and kresoxim-methyl;
 a composition comprising "carboxamide compound (4)" and picoxystrobin;
 a composition comprising "carboxamide compound (4)" and trifloxystrobin;
 a composition comprising "carboxamide compound (4)" and N-methyl-alpha -
 methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide;
 a composition comprising "carboxamide compound (5)" and dimoxystrobin;
 a composition comprising "carboxamide compound (5)" and azoxystrobin;
 a composition comprising "carboxamide compound (5)" and fluoxastrobin;
 a composition comprising "carboxamide compound (5)" and pyraclostrobin;
 a composition comprising "carboxamide compound (5)" and kresoxim-methyl;
 a composition comprising "carboxamide compound (5)" and picoxystrobin;
 a composition comprising "carboxamide compound (5)" and trifloxystrobin;
 a composition comprising "carboxamide compound (5)" and N-methyl-alpha -
 methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide.

- [0023] a composition comprising "carboxamide compound (1)" and dimoxystrobin in which the weight ratio of "carboxamide compound (1)" to dimoxystrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (1)" and azoxystrobin in which the weight ratio of "carboxamide compound (1)" to azoxystrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (1)" and fluoxastrobin in which the weight ratio of "carboxamide compound (1)" to fluoxastrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (1)" and pyraclostrobin in which the weight ratio of "carboxamide compound (1)" to pyraclostrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (1)" and kresoxim-methyl in which the weight ratio of "carboxamide compound (1)" to kresoxim-methyl is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (1)" and picoxystrobin in which

the weight ratio of "carboxamide compound (1)" to picoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (1)" and trifloxystrobin in which the weight ratio of "carboxamide compound (1)" to trifloxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (1)" and N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide in which the weight ratio of "carboxamide compound (1)" to N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and dimoxystrobin in which the weight ratio of "carboxamide compound (2)" to dimoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and azoxystrobin in which the weight ratio of "carboxamide compound (2)" to azoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and fluoxastrobin in which the weight ratio of "carboxamide compound (2)" to fluoxastrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and pyraclostrobin in which the weight ratio of "carboxamide compound (2)" to pyraclostrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and kresoxim-methyl in which the weight ratio of "carboxamide compound (2)" to kresoxim-methyl is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and picoxystrobin in which the weight ratio of "carboxamide compound (2)" to picoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and trifloxystrobin in which the weight ratio of "carboxamide compound (2)" to trifloxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (2)" and N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide in which the weight ratio of "carboxamide compound (2)" to N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and dimoxystrobin in which the weight ratio of "carboxamide compound (3)" to dimoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and azoxystrobin in weight ratio of "carboxamide compound (3)" to azoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and fluoxastrobin in which the weight ratio of "carboxamide compound (3)" to fluoxastrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and pyraclostrobin in which the weight ratio of "carboxamide compound (3)" to pyraclostrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and kresoxim-methyl in which the weight ratio of "carboxamide compound (3)" to kresoxim-methyl is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and picoxystrobin in which

the weight ratio of "carboxamide compound (3)" to picoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and trifloxystrobin in which the weight ratio of "carboxamide compound (3)" to trifloxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (3)" and N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide in which the weight ratio of "carboxamide compound (3)" to N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and dimoxystrobin in which the weight ratio of "carboxamide compound (4)" to dimoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and azoxystrobin in which the weight ratio of "carboxamide compound (4)" to azoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and fluoxastrobin in which the weight ratio of "carboxamide compound (4)" to fluoxastrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and pyraclostrobin in which the weight ratio of "carboxamide compound (4)" to pyraclostrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and kresoxim-methyl in which the weight ratio of "carboxamide compound (4)" to kresoxim-methyl is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and picoxystrobin in which the weight ratio of "carboxamide compound (4)" to picoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and trifloxystrobin in which the weight ratio of "carboxamide compound (4)" to trifloxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (4)" and N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide in which the weight ratio of "carboxamide compound (4)" to N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and dimoxystrobin in which the weight ratio of "carboxamide compound (5)" to dimoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and azoxystrobin in which the weight ratio of "carboxamide compound (5)" to azoxystrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and fluoxastrobin in which the weight ratio of "carboxamide compound (5)" to fluoxastrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and pyraclostrobin in which the weight ratio of "carboxamide compound (5)" to pyraclostrobin is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and kresoxim-methyl in which the weight ratio of "carboxamide compound (5)" to kresoxim-methyl is 0.1/1 to 10/1;
a composition comprising "carboxamide compound (5)" and picoxystrobin in which

the weight ratio of "carboxamide compound (5)" to picoxystrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (5)" and trifloxystrobin in which
 the weight ratio of "carboxamide compound (5)" to trifloxystrobin is 0.1/1 to 10/1;
 a composition comprising "carboxamide compound (5)" and N-methyl-alpha -
 methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide in which the weight
 ratio of "carboxamide compound (5)" to N-methyl-alpha -
 methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide is 0.1/1 to 10/1;

[0024] The method of controlling plant disease (hereinafter referred to as "controlling method") can be carried out by treating a plant or the soil where a plant grows with an effective amount of a "carboxamide compound" and "QoI compound (s)".

The part of plant to be treated is stem and leaf of a plant, seed or bulb of a plant, and the bulb means bulb, corm, rootstock, tuber, tuberous root and rhizophore.

[0025] In the "controlling method", the treatment of a plant or the soil where a plant grows with a "carboxamide compound" and "QoI compound (s)" can be carried out separately at the same timing, but the treatment is usually carried out by using a "composition" in light of convenience.

[0026] In the "controlling method", the treatment with a "carboxamide compound" and "QoI compound(s)" is, for example, stems and leaves application, soil application, roots application or seeds application.

[0027] Examples of the stems and leaves application include a treatment for surface of cultivated plant by a stem and leaves spray or a stem and tree spray.

Examples of the root application include a method of dipping a whole plant or root of a plant into a liquid containing a "carboxamide compound" and "QoI compound (s)" and a method of sticking a solid preparation comprising a "carboxamide compound", "QoI compound(s)" and a solid carrier onto root of a plant.

Examples of the soil application include a method of spraying a "composition" onto a soil, a method of mixing a "composition" with a soil and a method of irrigating a "composition" into the soil.

Examples of the seed application include a method of treating seeds or bulbs of a plant to be protected from a plant disease with a "composition". Particularly, the application can be carried out by spraying a suspension of a "composition" to the surface of seeds or the bulbs, or by spreading wettable powder, emulsifiable concentrate or flowable formulation itself or a mixture thereof with a small amount of water on the seeds or the bulbs, or by dipping the seeds into a solution of a "composition" for a prescribed time, by film coating application or pellet coating application.

[0028] The amount of a "carboxamide compound" and "QoI compound(s)" used in the "controlling method" is different depending on the kind of a plant to be treated, the kind of a plant disease to be controlled and its frequency, the kind of a formulation, timing of

treatment, method of treatment, place of treatment, weather condition, and the like.

When a "composition" is applied to stems and/or leaves of a plant or to the soil where a plant grows, the total amount of a "carboxamide compound" and "QoI compound(s)" is usually from 1g to 500g / 1000m², preferably from 2g to 200g / 1000m², and more preferably from 10g to 100g / 1000m².

When a "composition" is applied to seeds of a plant, the total amount of a "carboxamide compound" and "QoI compound(s)" is usually from 0.001g to 10g / 1kg of the seeds, and preferably from 0.01g to 1g / 1kg of the seeds.

An emulsifiable concentrate, wettable powder or flowable formulation is used usually by diluting the formulation with water and spraying the diluted formulation. In this case, the concentration of a "carboxamide compound" and "QoI compound(s)" in total of the diluted formulation is usually from 0.0005% to 2% by weight, and preferably from 0.005% to 1% by weight.

A powder formulation, granule formulation, and the like is usually used without dilution.

Example

[0029] The present invention is further explained in detail with Formulation Examples and Test Examples. However, the present invention is not limited by the following Examples.

In the following Examples, "part" means "part by weight" unless otherwise provided.

[0030] Formulation Example 1

One of "carboxamide compounds" (1) to (5) (2.5 parts), dimoxystrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0031] Formulation Example 2

One of "carboxamide compounds" (1) to (5) (2.5 parts), azoxystrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0032] Formulation Example 3

One of "carboxamide compounds" (1) to (5) (2.5 parts), fluoxastrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0033] Formulation Example 4

One of "carboxamide compounds" (1) to (5) (2.5 parts), pyraclostrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6

parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0034] Formulation Example 5

One of "carboxamide compounds" (1) to (5) (2.5 parts), kresoxim-methyl (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0035] Formulation Example 6

One of "carboxamide compounds" (1) to (5) (2.5 parts), picoxystrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0036] Formulation Example 7

One of "carboxamide compounds" (1) to (5) (2.5 parts), trifloxystrobin (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0037] Formulation Example 8

One of "carboxamide compounds" (1) to (5) (2.5 parts), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (1.25 parts), polyoxyethylene styryl phenyl ether (14 parts), calcium dodecylbenzene sulfonate (6 parts) and xylene (76.25 parts) are thoroughly mixed to give each of formulations, respectively.

[0038] Formulation Example 9

One of "carboxamide compounds" (1) to (5) (2 parts), dimoxystrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and milled by wet-milling method to give each of formulations, respectively.

[0039] Formulation Example 10

One of "carboxamide compounds" (1) to (5) (2 parts), azoxystrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0040] Formulation Example 11

One of "carboxamide compounds" (1) to (5) (2 parts), fluoxastrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0041] Formulation Example 12

One of "carboxamide compounds" (1) to (5) (2 parts), pyraclostrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0042] Formulation Example 13

One of "carboxamide compounds" (1) to (5) (2 parts), kresoxim-methyl (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0043] Formulation Example 14

One of "carboxamide compounds" (1) to (5) (2 parts), picoxystrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0044] Formulation Example 15

One of "carboxamide compounds" (1) to (5) (2 parts), trifloxystrobin (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0045] Formulation Example 16

One of "carboxamide compounds" (1) to (5) (2 parts), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (8 parts), a mixture of white carbon and polyoxyethylene alkyl ether sulfate ammonium salt (ratio by weight 1:1) (35 parts) and water (55 parts) are mixed and the mixture is milled by wet-milling method to give each of formulations, respectively.

[0046] Formulation Example 17

One of "carboxamide compounds" (1) to (5) (5 parts), dimoxystrobin (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0047] Formulation Example 18

One of "carboxamide compounds" (1) to (5) (5 parts), azoxystrobin (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the

milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0048] Formulation Example 19

One of "carboxamide compounds" (1) to (5) (5 parts), fluoxastrobin (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0049] Formulation Example 20

One of "carboxamide compounds" (1) to (5) (5 parts), pyraclostrobin (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0050] Formulation Example 21

One of "carboxamide compounds" (1) to (5) (5 parts), kresoxim-methyl (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0051] Formulation Example 22

One of "carboxamide compounds" (1) to (5) (5 parts), picoxystrobin (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0052] Formulation Example 23

One of "carboxamide compounds" (1) to (5) (5 parts), trifloxystrobin (10 parts),

sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0053] Formulation Example 24

One of "carboxamide compounds" (1) to (5) (5 parts), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (10 parts), sorbitan tri-oleate (1.5 parts) and an aqueous solution (28.5 parts) containing polyvinyl alcohol (2 parts) are mixed and the mixture is milled by wet-milling method. To the milled mixture is added an aqueous solution (45 parts) containing xanthan gum (0.05 parts) and aluminum magnesium silicate (0.1 part), and further propylene glycol (10 parts) is added to the mixture. The resultant mixture is mixed by stirring to give each of formulations, respectively.

[0054] Formulation Example 25

One of "carboxamide compounds" (1) to (5) (1 part), dimoxystrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are thoroughly mixed and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0055] Formulation Example 26

One of "carboxamide compounds" (1) to (5) (1 part), azoxystrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are thoroughly mixed and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0056] Formulation Example 27

One of "carboxamide compounds" (1) to (5) (1 part), fluoxastrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are thoroughly mixed and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0057] Formulation Example 28

One of "carboxamide compounds" (1) to (5) (1 part), pyraclostrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are thoroughly mixed and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give

each of formulations, respectively.

[0058] Formulation Example 29

One of "carboxamide compounds" (1) to (5) (1 part), kresoxim-methyl (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are thoroughly mixed and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0059] Formulation Example 30

One of "carboxamide compounds" (1) to (5) (1 part), picoxystrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are mixed well and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0060] Formulation Example 31

One of "carboxamide compounds" (1) to (5) (1 part), trifloxystrobin (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are mixed well and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0061] Formulation Example 32

One of "carboxamide compounds" (1) to (5) (1 part), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (4 parts), synthesized hydrous silicon oxide (1 part), calcium ligninsulfonate (2 parts), bentonite (30 parts) and kaolin clay (62 parts) are mixed well and milled. Water is added to the mixture and the mixture is sufficiently kneaded, granulated and then dried to give each of formulations, respectively.

[0062] Formulation Example 33

One of "carboxamide compounds" (1) to (5) (12.5 parts), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (37.5 parts), calcium ligninsulfonate (3 parts), sodium laurylsulfate (2 parts) and synthesized hydrous silicon oxide (45 parts) are thoroughly mixed and milled to give each of formulations, respectively.

[0063] Formulation Example 34

One of "carboxamide compounds" (1) to (5) (3 parts), N-methyl-alpha - methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide (2 parts), kaolin clay (85 parts) and talc (10 parts) are thoroughly mixed and milled to give each of formulations, respectively.

[0064] Test Examples using each of the "compositions" are shown in the following.

[0065] Test Example 1

A cyclohexanone solution (100microL) containing prescribed amount (weight) of test compounds was applied on seeds of soybean (variety:Natto Shoryu) (10g) by using a rotary apparatus for seed treatment (Seed dresser, manufactured by Hans-Ulrich Hege GmbH).

One day after the application, plastic pot was filled with soil contaminated by Rhizoctonia solani, and the seeds treated with the test compounds were seeded in the soil and cultivated in a glass-greenhouse for 20 days (hereinafter referred to as "treated plot").

Thereafter, the presence of disease caused by Rhizoctonia solani in the young plants which germinated from each seed was observed and disease severity was calculated by the following calculation formula (1).

On the other hand, seeds of soybean which were not treated as above were cultivated in the same way as above (hereinafter referred to as "non-treated plot") and the disease severity in "non-treated plot" was calculated in the same way as above "treated plot".

On the basis of the above disease severity in "treated plot" and "non-treated plot", efficacy in "treated plot" was evaluated by the following calculation formula (2).

The results are shown in Table 1 to Table 6.

Calculation formula (1):

Disease degree (%) =

(number of infected young plants / total number of young plants) x 100

Calculation formula (2):

Efficacy (%) =

[1 - (disease severity in "treated plot" / disease severity in "non-treated plot")] x 100

[0066] [Table 1]

"carboxamide compound (1)" [g / 100kg of seeds]	azoxystrobin [g / 100kg of seeds]	efficacy (%)
2	2	100

[0067] [Table 2]

"carboxamide compound (5)" [g / 100kg of seeds]	azoxystrobin [g / 100kg of seeds]	efficacy (%)
2	2	100

[0068]

[Table 3]

"carboxamide compound (1)" [g / 100kg of seeds]	pyraclostrobin [g / 100kg of seeds]	efficacy (%)
2	2	100

[0069] [Table 4]

"carboxamide compound (5)" [g / 100kg of seeds]	pyraclostrobin [g / 100kg of seeds]	efficacy (%)
2	2	100

[0070] [Table 5]

"carboxamide compound (1)" [g / 100kg of seeds]	N-methyl- α -methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide [g / 100kg of seeds]	efficacy (%)
2	2	100

[0071] [Table 6]

"carboxamide compound (5)" [g / 100kg of seeds]	N-methyl- α -methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide [g / 100kg of seeds]	efficacy (%)
2	2	100

[0072] Test Example 2

A plastic pot was filled a soil, and soybean seeds (variety: Natto Shoryu) were seeded in the soil and grown in a greenhouse for 14 days. Test compounds were dissolved in CEC cocktail (cyclohexanone : Solpol™ 2680X (manufactured by Toho Kagaku Kogyo) = 5:1 (by volume)) to give an emulsifiable concentrate containing total amount 5% (w/v) of the test compounds. The emulsifiable concentrate was mixed with water to give a prescribed concentration. The mixture was sprayed on leaves of the soybean so as to stick sufficiently thereto. After the spraying, the plant was air-dried, and one day after, the plant was inoculated with an aqueous suspension containing urediniospore of *Phakopsora pachyrhizi* (about 10,000/ml) by spraying the suspension. After the inoculation, the plant was left in humid circumstance at 20-23°C for one day and then cultivated in a greenhouse for 10 days (hereinafter referred to as "treated plot"). Thereafter, lesion area of *Phakopsora pachyrhizi* was investigated.

On the other hand, soybean was cultivated in the same way as above "treated plot"

except for that the plant was not treated with the mixture containing the test compounds (hereinafter referred to as "non-treated plot"), and the lesion area of *Phakopsora pachyrhizi* were investigated in the same way as above.

On the basis of the above stigmata area in the "treated plot" and "non-treated plot", efficacy in "treated plot" was evaluated according to the following calculation formula (3).

Calculation formula (3):

Efficacy (%) =

$[1 - (\text{lesion area in "treated plot"} / \text{lesion area in "non-treated plot"})] \times 100$

The results are shown in the following Table 7 to Table 18.

[0073] [Table 7]

"carboxamide compound (1)" [ppm]	trifloxystrobin [ppm]	efficacy (%)
2	2	100

[0074] [Table 8]

"carboxamide compound (5)" [ppm]	trifloxystrobin [ppm]	efficacy (%)
2	2	100

[0075] [Table 9]

"carboxamide compound (1)" [ppm]	azoxystrobin [ppm]	efficacy (%)
2	2	100

[0076] [Table 10]

"carboxamide compound (5)" [ppm]	azoxystrobin [ppm]	efficacy (%)
2	2	100

[0077] [Table 11]

"carboxamide compound (1)" [ppm]	N-methyl- α -methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide [ppm]	efficacy (%)
2	10	100

[0078]

[Table 12]

"carboxamide compound (5)" [ppm]	N-methyl- α -methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide [ppm]	efficacy (%)
2	10	100

[0079] [Table 13]

"carboxamide compound (1)" [ppm]	pyraclostrobin [ppm]	efficacy (%)
2	2	100

[0080] [Table 14]

"carboxamide compound (5)" [ppm]	pyraclostrobin [ppm]	efficacy (%)
2	2	100

[0081] [Table 15]

"carboxamide compound (1)" [ppm]	picoxystrobin [ppm]	efficacy (%)
2	2	100

[0082] [Table 16]

"carboxamide compound (5)" [ppm]	picoxystrobin [ppm]	efficacy (%)
2	2	100

[0083] [Table 17]

"carboxamide compound (1)" [ppm]	fluoxastrobin [ppm]	efficacy (%)
2	2	100

[0084] [Table 18]


"carboxamide compound (5)" [ppm]	fluoxastrobin [ppm]	efficacy (%)
2	2	100

Industrial Applicability

[0085] A plant disease control composition comprising a "carboxamide compound" represented by formula (I) and one or more QoI compounds selected from group (A) is useful for controlling plant disease.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A composition for controlling a plant fungal disease comprising carboxamide compound (1):




and

a QoI compound which is dimoxystrobin, azoxystrobin, fluoxastrobin, pyraclostrobin, kresoxim-methyl, picoxystrobin, trifloxystrobin, N-methyl-alpha-methoxyimino-2-[(2,5-dimethylphenoxy)methyl]phenylacetamide, or any combination thereof;

wherein the weight ratio of the carboxamide compound (1) to the QoI compound is from 0.1/1 to 10/1.

2. A method of controlling a plant fungal disease which comprises the step of treating a plant, or soil where a plant grows, with a composition as defined in claim 1.
3. The method according to claim 2, wherein the treated plant is soybean, or the treated soil is soil where soybeans grow.



(1)