

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2007 (08.02.2007)

PCT

(10) International Publication Number
WO 2007/016539 A2

(51) International Patent Classification:
C12N 5/08 (2006.01)

(21) International Application Number:
PCT/US2006/029873

(22) International Filing Date: 31 July 2006 (31.07.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/703,587 29 July 2005 (29.07.2005) US

(71) Applicants (for all designated States except US):
CHILDREN'S HOSPITAL MEDICAL CENTER [US/US]; 3333 Burnet Avenue, MLC 7032, Cincinnati, Ohio 45229-3039 (US). **THE STATE UNIVERSITY OF NEW YORK, STONY BROOK UNIVERSITY** [US/US]; N5002 Melville Library, SUNY At Stony Brook, Stony Brook, New York 11794-3369 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHENG, Yi** [US/US]; 6705 North Clippenger Drive, Cincinnati, Ohio 45243 (US). **NASSAR, Nicolas** [LB/US]; 18 Hawkins Path, Coram, New York 11727 (US). **SKOWRONEK, Karlheinz, R.** [US/US]; 56 Fairfields Lane, Huntington Station, New York 11746 (US).

(74) Agent: **DELANEY, Karoline A.**; Knobbe, Martens, Olson & Bear, LLP, 2040 Main Street, 14th Floor, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GTPASE INHIBITORS AND METHODS OF USE AND CRYSTAL STRUCTURE OF RAC-1 GTPASE



(57) Abstract: The preferred embodiments generally relate to methods and compositions that affect the GTP-binding activity of members of the Rho family GTPases, preferably Rac (Rac1, Rac2 and/or Rac3), such compositions include compounds that modulate the GTP/GDP exchange activity, along with uses for the compounds including screening for compounds which recognize Rac GTPase, and methods of treating pathological conditions associated or related to a Rho family GTPase, including Rac. The preferred embodiments

also relate to methods of using such compounds, or derivatives thereof, e.g., in therapeutics, diagnostics, and as research tools.

GTPase INHIBITORS AND METHODS OF USE AND CRYSTAL STRUCTURE OF RAC-1 GTPase

Cross Reference to Related Applications

[0001] This patent application claims priority under 35 U.S.C. §119(e) to U.S. Provisional Patent No. 60/703,587, filed July 29, 2005, which is incorporated herein by reference in its entirety.

Statement Regarding Federally Sponsored Research

[0002] This invention was made in part with Government support under Grant No. R01 GM60523 and No. R01 GM53943 awarded by the National Institutes of Health. The Government can have certain rights in this invention.

Background of the Invention

Field of the Invention

[0003] The present invention generally relates to methods and compositions that affect the GTP-binding activity of members of the Ras superfamily GTPases, along with uses for the compounds including screening for compounds that recognize Rac GTPase, and methods of treating pathological conditions associated or related to a Ras superfamily GTPase.

Description of the Related Art


[0004] Rho family GTPases are molecular switches that control signaling pathways regulating cytoskeleton reorganization, gene expression, cell cycle progression, cell survival, and other cellular processes (Etienne-Manneville, 2002), which is incorporated herein by reference in its entirety.

[0005] Rho family proteins constitute one of three major branches of the Ras superfamily. Rho proteins share approximately 30 percent amino acid identity with the Ras proteins. At least 14 mammalian Rho family proteins have been identified thus far, including RhoA, RhoB, RhoC, RhoD, RhoE/Rnd3, Rnd1/Rho6, Rnd2/Rho7, RhoG, Rac1, Rac2, Rac3, Cdc42, TC10, and TTF.

Summary of the Invention

[0006] The preferred embodiments provide compounds that are potent and selective inhibitors of Rho GTPases. Specifically, these compounds can be used to inhibit Rho-related Rac GTPase. These inhibitors can be used to treat diseases associated with Rac dysregulation. Furthermore, these compounds can be used to treat cancers associated with Rac dysregulation. In view of their activity, these compounds can also be used in treating other disorders responding to the inhibition of Rac.


[0007] One embodiment comprises a method for treating an indication mediated by mammalian Rho family proteins, comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (IIa):



(IIa) wherein:

R₁ to R₂ are independently selected from the group consisting of H, -X-Alk, -X-Alk-X', and -X-Y-X'; wherein X is -CR₇R₈; X' is -CHR₇R₈; Alk is a C₂-C₁₈ substituted or unsubstituted hydrocarbon chain; Y is a C₂-C₈ substituted or unsubstituted alkylene chain; R₆ is H or (C1-C4) alkyl; and R₇ and R₈ are independently selected from the group consisting of H or (C1-C4) alkyl or a salt of a compound of formula (IIa).


[0008] One embodiment comprises a method for treating an indication mediated by mammalian Rho family proteins, comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (III):



(III) wherein:

R₁₀ to R₁₂ are independently selected from the group consisting of H, halo, (C1-C4) alkyl, branched (C3-C4) alkyl, halo (C1-C4) alkyl, (C1-C4) alkoxy, NO₂, and NH₂; or a salt of a compound of formula (III).

[0009] One embodiment comprises a method for treating an indication mediated by mammalian Rho family proteins comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (IV):



(IV)

or pharmaceutically acceptable salts thereof.

[0010] One embodiment comprises a pharmaceutical composition comprising N6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine and a pharmaceutically-acceptable carrier.

[0011] One embodiment comprises a crystal of Rac-1 GTPase having a space group $P2_1$ and unit cell dimensions of about $a=41.8$ Angstroms, $b=40.0$ Angstroms, $c=52.1$ Angstroms, and $\beta=105.93^\circ$.

[0012] One embodiment comprises a crystal Rac-1 GTPase, wherein said crystal has a three-dimensional structure characterized by the atomic structure coordinates of Table 1 or Table 2.

[0013] One embodiment comprises a method of making a crystal comprising Rac-1 GTPase, wherein the crystal Rac-1 GTPase comprises at least amino acid residues 1 through 185 of the sequence of SEQ ID NO:1; wherein the crystal effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of greater than 5.0 Angstroms; and wherein said crystal has a space group $P2_1$ and unit cell dimensions of about $a=41.8$ Angstroms, $b=40.0$ Angstroms, $c=52.1$ Angstroms, and $\beta=105.93^\circ$,

[0014] said method comprising growing a crystal by hanging drop method using a buffer that comprises 20% PEG 8000, 5% Ethylene Glycol, 15% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, and 1 mM NSC23766 at 20°C; and

[0015] equilibrating the crystal in a drop comprising 18% PEG 8000, 3.5% Ethylene Glycol, 10.7% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, 14.3 mM NSC23766 over a reservoir containing a buffer comprising 25% PEG 8000, 5% Ethylene Glycol, 15% DMSO 0.1M HEPES, pH 7.0, 10 mM DTT.

[0016] One embodiment comprises a method of identifying an agent that interacts with Rac-1 GTPase comprising:

- (a) obtaining a crystal of a complex comprising Rac-1 GTPase;
- (b) obtaining the atomic coordinates of the crystal; and
- (c) using the atomic coordinates and one or more molecular modeling techniques to identify an agent that interacts with Rac-1 GTPase.

[0017] One embodiment comprises a method for evaluating the potential of a chemical entity to associate with:

[0018] a) a molecule or molecular complex comprising a binding pocket defined by structure coordinates of Rac-1 GTPase Trp-56, Lys-5, Val-7, Ser-71, Ile-36, Tyr-64, Arg-68, and Pro-73; or

[0019] b) a homologue of said molecule or molecular complex, wherein said homologue comprises a binding pocket that has a root mean square deviation from the backbone atoms of said amino acids of not more than about 1.5 Angstrom comprising the steps of:

[0020] i) employing computational means to perform a fitting operation between the chemical entity and the binding pocket of the molecule or molecular complex; and

[0021] ii) analyzing the results of said fitting operation to quantify the association between the chemical entity and the binding pocket.

[0022] One embodiment comprises a compound for inhibiting activity of Rac-1 GTPase, whose active site presents a three-dimensional structure from a crystal of Rac-1 GTPase having a space group $P2_1$ and unit cell dimensions of about $a=41.8$ Angstroms, $b=40.0$ Angstroms, $c=52.1$ Angstroms, and $\beta=105.93^\circ$.

[0023] One embodiment is directed to a crystal of Rac-1 GTPase bound to GDP.

[0024] One embodiment is directed to a crystal Rac-1 GTPase, wherein said crystal has a three-dimensional structure characterized as having atomic coordinates with a root-mean-squared deviation of $6\text{\AA}$ or less, $5\text{\AA}$ or less, $4\text{\AA}$ or less, $3\text{\AA}$ or less, $2\text{\AA}$ or less, $1\text{\AA}$ or less, relative to atoms of amino acids 30-39 and 59-73 of Table 1 or Table 2. In some such embodiments, the of amino acids 30-39 and 59-73 of Table 1 or Table 2 are peptide backbone atoms.

[0025] One embodiment is directed to atomic coordinates with a root-mean-squared deviation of 6Å or less, 5Å or less, 4Å or less, 3Å or less, 2Å or less, 1Å or less, relative to atoms of amino acids 30-39 and 59-73 of Table 1 or Table 2.

The coordinates of Claim 33, wherein the of amino acids 30-39 and 59-73 of Table 1 or Table 2 are peptide backbone atoms.

[0026] One embodiment is directed to a method of characterizing the three-dimensional structure of Rac-1 GTPase comprising: (a) obtaining a crystal of a GDP-bound Rac-1 GTPase; and (b) obtaining the atomic coordinates of the crystallized GDP-bound Rac-1 GTPase.

[0027] One embodiment is directed to a method of characterizing the three-dimensional structure of a ligand bound to a Rac-1 GTPase comprising: (a) obtaining a crystal of a ligand bound to a GDP-bound Rac-1 GTPase; and (b) obtaining the atomic coordinates of the crystallized ligand and GDP-bound Rac-1 GTPase.

[0028] One embodiment is directed to a method of identifying an agent that interacts with Rac-1 GTPase comprising: (a) obtaining a crystal of a GDP-bound Rac-1 GTPase; and (b) obtaining the atomic coordinates of the crystallized GDP-bound Rac-1 GTPase; and (c) using the atomic coordinates and one or more molecular modeling techniques to identify an agent that interacts with GDP-bound Rac-1 GTPase.

One embodiment is directed to a method of designing a Rac-1 GTPase inhibitor comprising accessing at least a portion the data of Table 1 or Table 2 and modeling the binding of one or more compounds to a Rac-1 GTPase utilizing said data. In some such embodiments, the modeling comprises predicting the likelihood of a compound to bind to Rac-1 GTPase by modeling the interaction of a compound with a Rac-1 GTPase activation site. Some such embodiments further comprise utilizing said data to identify one or more interactions between a Rac-1 GTPase and a ligand. In some such embodiments, said one or more interactions are selected from the group consisting of one or more hydrogen bonds between ligand and protein; one or more electrostatic interactions between ligand and protein; one or more hydrophobic interactions; one or more covalent bonds between ligand and protein; one or more changes in the positions of protein atoms, such as amino acid side chains upon ligand binding; one or more changes in atoms whose positions are well defined to a conformation that may maintain flexibility; one or more hydrogen bonding networks; one or more structural changes to the hydrogen bond network, one or more structural changes to the interactions with Mg²⁺, one or more

structural changes to the interactions with GDP, and one or more structural changes to the interactions with GTP.

[0029] A method identifying a Rac-1 GTPase inhibitor comprising obtaining data comprising the one or more interactions of Claim 40 and modeling a compound to interact with a Rac-1 GTPase, wherein the compound comprises one or more of the interactions of Claim 40 or improvements of one or more of the interactions of Claim 40.

[0030] One embodiment is directed to a method of designing a Rac-1 GTPase inhibitor comprising obtaining an X-ray crystal structure coordinate of a GDP-bound Rac-1 GTPase and identifying a compound that inhibits a Rac-1 GTPase activation. The method of Claim 43, comprising identifying a compound that inhibits Thr35 binding of Mg²⁺.

[0031] The method of Claim 43 or Claim 44, comprising identifying a compound that inhibits conformational change in the loop containing amino acid residues 60-64.

The method of any of Claims 43-45, comprising identifying a compound that inhibits Ala59 from displacing Mg²⁺.

The method of Claim 46, wherein said compound hydrogen bonds to Ala59.

The method of any of Claims 43-47, comprising identifying a compound that inhibits a RAC-specific GEF from binding to RAC

[0032] The method of any of Claims 43-48, comprising identifying a compound that inhibits RAC-specific GEF from dissociating from RAC.

[0033] The method of any of Claims 43-49, comprising identifying a compound that inhibits conformational change in the amino acids of switch I.

[0034] The method of any of Claims 43-50, comprising identifying a compound that inhibits conformational change in the amino acids of switch II

[0035] The method of any of claims 43-51, wherein said compound is identified as hydrogen bonding to Leu70 or Ser71, or both.

[0036] The method of any of claims 43-52, wherein said compound is identified as hydrogen bonding to Leu67, Gln74, Asp57, or Ser71, or combinations thereof.

[0037] The method of any of claims 43-53, wherein said compound is identified as being in van der Waals contact with Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 or Pro73, or combinations thereof.

[0038] One embodiment is directed to a method of identifying a proteasome inhibitor comprising accessing one or more of the coordinates of Table 1 or Table 2 and identifying a compound that occupies a position within a range of about 1.0 to 2.0 angstroms or about 2.5 to 5.0 angstroms of one or more of the coordinates of the ligand in Table 1 or Table 2.

[0039] The method of Claim 55, wherein the compound occupies a position within about 1.2 to 1.7 or about 2.6 to 3.5 angstroms of one or more of the coordinates of the ligand in at least one of Tables 1-5.

[0040] One embodiment is directed to a method of identifying a proteasome inhibitor comprising:

- (a) identifying one or more interactions between a ligand and a Rac-1 GTPase; and
- (b) changing a substituent on the ligand in order to modify the interaction between the ligand the subunit.

[0041] The method of Claim 57, wherein the ligand is NSC23766.

One embodiment is directed to a method of screening a set of one or more compounds to identify compounds most likely to bind to a Rac-1 GTPase comprising the steps of: (a) computationally determining the docking energy of each of one or more compounds in an activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site; (b) selecting one or more compounds indicated as having a likelihood of binding to said Rac-1 GTPase activation site above a threshold value.

[0042] One embodiment is directed to a method of identifying a compound that is likely to bind an activation site of a Rac-1 GTPase comprising the steps of: (a) computationally determining the docking energy of each of one or more compounds in an activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site; (b) selecting one or more compounds indicated as having a likelihood of binding to said Rac-1 GTPase activation site above a threshold value.

[0043] The method of any of Claims 59-60, wherein said activation site comprises one or more of the protein atoms of Table 1 or Table 2.

[0044] One embodiment is directed to atomic coordinates that have a root-mean-squared deviation of 6.0 Å or less, 5.5 Å or less, 5.0 Å or less, 4.5 Å or less, 4.0 Å or less, 3.5 Å or less, 3.0 Å or less, 2.5 Å or less, 2.0 Å or less, 1.7 Å or less, 1.5 Å or

less, 1.4 Å or less, 1.3 Å or less, 1.2 Å or less, 1.1 Å or less, 1.0 Å or less, 0.9 Å or less, 0.8 Å or less, 0.7 Å or less, 0.6 Å or less, 0.5 Å or less, 0.4 Å or less, 0.3 Å or less, 0.2 Å or less, or 0.1 Å or less, relative to atoms of the atomic coordinates provided in Table 1 or Table 2.

[0045] The method of any of Claims 59-61, wherein said activation site comprises protein atoms having atomic coordinates that have a root-mean-squared deviation of 6.0 Å or less, 5.5 Å or less, 5.0 Å or less, 4.5 Å or less, 4.0 Å or less, 3.5 Å or less, 3.0 Å or less, 2.5 Å or less, 2.0 Å or less, 1.7 Å or less, 1.5 Å or less, 1.4 Å or less, 1.3 Å or less, 1.2 Å or less, 1.1 Å or less, 1.0 Å or less, 0.9 Å or less, 0.8 Å or less, 0.7 Å or less, 0.6 Å or less, 0.5 Å or less, 0.4 Å or less, 0.3 Å or less, 0.2 Å or less, or 0.1 Å or less, relative to atoms of the atomic coordinates provided in Table 1 or Table 2.

[0046] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid of switch I and at least one atom from an amino acid of switch II.

[0047] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase (see SEQ ID NO:1) and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase (see SEQ ID NO:1).

[0048] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 and Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase.

[0049] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase.

[0050] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at

least one atom from an amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from Trp56 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

[0051] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

[0052] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Leu70 and Ser71 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

[0053] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Tyr64, Leu67 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

[0054] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

[0055] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 or Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-70 of Rac-1 GTPase.

[0056] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and

Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase.

[0057] The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase.

[0058] The atomic coordinates or method of any of Claims 62-76, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include the Mg²⁺ atom.

The atomic coordinates or method of any of Claims 62-76, wherein the atoms are selected from the group consisting of all atoms of a selected amino acid, backbone atoms only of a selected amino acid, and alpha carbon atoms only of a selected amino acid.

[0059] The method of any of Claims 63-78, further comprising determining the conformation of said compound in said activation site.

[0060] The method of any of Claims 63-79, wherein said step of determining the docking energy further comprises a step of generating a grid-based receptor field representation of said activation site.

[0061] The method of any of Claims 63-80, wherein said step of determining the docking energy further comprises a step of determining the optimal conformation of said compound in said activation site.

[0062] The method of any of Claims 63-81, wherein said step of determining the docking energy further comprises a step of building atoms of said compound into said activation site to thereby match said activation site.

[0063] A computer readable media comprising the atomic coordinates of any of Claims 62 or 64-78.

[0064] The computer readable media of Claim 83, comprising the atomic coordinates of the protein atoms of Table 1 or Table 2.

[0065] A computer readable media comprising the the atomic coordinates for a ligand of Table 1 or Table 2.

[0066] The computer readable media of Claim 85, wherein the ligand is NSC23766.

[0067] A computer readable media comprising at least a portion of the atomic coordinates for a GDP-bound Rac-1 GTPase.

[0068] The computer readable media of Claim 87, comprising amino acids 1-185 of Rac-1 GTPase.

[0069] The computer readable media of Claim 87, further comprising at least a portion of the atomic coordinates of a non-GDP ligand of Rac-1 GTPase.

[0070] The computer readable media of Claim 87 or Claim 88, wherein the coordinates are less than the coordinates for amino acids 1-185 of Rac-1 GTPase.

[0071] A computer system comprising the coordinates of any of Claims 62, 64-78 or 83-90.

[0072] The computer system of Claim 91, capable of generating structures and/or performing rational drug design for a proteasome/ligand complex.

Brief Description of the Drawings

[0073] The novel features of the preferred embodiments are set forth with particularity in the appended claims. The preferred embodiments, however, both as to organization and methods of operation, together with further objects and advantages thereof, can best be understood by reference to the following description, taken in conjunction with the accompanying drawings in which:

[0074] Fig. 1. shows identification of NSC23766 as an inhibitor of Rac1-Trio interaction.

[0075] Fig. 2. shows dose dependent specific inhibition of GEF interaction with Rac1 by NSC23766.

[0076] Fig. 3 shows that NSC23766 was effective in specifically inhibiting Rac1 GDP/GTP exchange stimulated by GEF.

[0077] Fig. 4. shows that NSC23766 was effective in specifically inhibiting Rac1 activation in cells.

[0078] Fig. 5. shows that NSC23766 specifically inhibited Rac GEF stimulated cell growth and transformation.

[0079] Fig. 6. shows that NSC23766 inhibited the proliferation, anchorage independent growth and invasion of PC-3 prostate cancer cells.

[0080] In the following description of the illustrated embodiments, references are made to the accompanying drawings, which form a part hereof, and in which is shown

by way of illustration various embodiments in which the invention can be practiced. It is to be understood that other embodiments can be utilized, and structural and functional changes can be made without departing from the scope of the preferred embodiments.

Detailed Description of the Preferred Embodiment

[0081] Throughout this document, all temperatures are given in degrees Celsius, and all percentages are weight percentages unless otherwise stated. The following are definitions of terms used in this specification. The initial definition provided for a group or term herein applies to that group or term throughout the present specification, individually or as part of another group, unless otherwise indicated. The following definitions, unless otherwise defined, apply to preferred embodiments:

[0082] The terms “active compounds” or “active agents” refer to any one of the agents described by formula I, II, IIa, III, IIIa or IV.

[0083] The term “alkyl” refers to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, or more, preferably 1 to 8 carbon atoms. Lower alkyl groups, that is, alkyl groups of 1 to 4 carbon atoms, are most preferred.

[0084] The term “substituted alkyl” refers to an alkyl group as defined above having at least one substituent, such as halo, amino, cyano, hydroxy, alkoxy, alkylthio, --NH(alkyl), --NH (cycloalkyl), --N(alkyl)₂, --C(=O)H, --CO₂ H, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, or heterocycle. The term “substituted alkyl” also includes an alkyl group as defined above substituted with N(substituted alkyl) or N(substituted alkyl)₂, or in other words, the groups (CH₂), NHR' and (CH₂)_n NR'R'', wherein each of R' and R'' comprises a substituted alkyl or R' and R'' together form a heterocyclo ring.

[0085] The term “alkoxy” refers to an alkyl group as defined above bonded through an oxygen (--O--). The term “alkylthio” refers to an alkyl group as defined above bonded through a sulfur (--S--).

[0086] The term “cycloalkyl” refers to fully saturated and partially unsaturated hydrocarbon rings of at least 3, preferably 3 to 9, more preferably 3 to 7, carbon atoms as well as such rings having a fused aryl ring such as indan.

[0087] The term “substituted cycloalkyl” refers to such rings having one, two or three substituents, preferably one, such as alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, cyano, amino, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, --CO₂ H, --CO₂ -

lower alkyl, aryl, heterocyclo, heteroaryl, keto, =N--OH, =N--O--lower alkyl, and a five or six membered ketal, *i.e.* 1,3-dioxolane or 1,3-dioxane.

[0088] The term "halo" refers to fluoro, chloro, bromo and iodo.

[0089] The term "aryl" refers to phenyl, 1-naphthyl and 2-naphthyl, with phenyl being preferred. The term "aryl" includes such rings having from zero, one, two or three substituents, such as alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, --CO₂ H, --(C=O)alkyl, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, --(C=O)NH₂, --(C=O)NH(alkyl), --(C=O)NH(cycloalkyl), --(C=O)N(alkyl)₂, --NH--CH₂ --CO₂ H, --NH--CH₂ --CO₂ -alkyl, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, heterocyclo, and heteroaryl.

[0090] The term "heterocyclo" refers to substituted and unsubstituted non-aromatic 3 to 7 membered monocyclic groups, 7 to 11 membered bicyclic groups, and 10 to 15 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heterocyclo group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less, and further provided that the ring contains at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups may contain only carbon atoms and can be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms can optionally be oxidized and the nitrogen atoms can optionally be quaternized. The heterocyclic group can be attached at any available nitrogen or carbon atom. The heterocyclic ring can contain one, two or three substituents, such as halo, amino, cyano, alkyl, substituted alkyl, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, alkoxy, alkylthio, hydroxy, nitro, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, --CO₂ H, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, --(C=O)NH₂, --(C=O)NH(alkyl), --(C=O)NH(cycloalkyl), --(C=O)N(alkyl)₂, --NH--CH₂ --CO₂ H, --NH--CH₂ --CO₂ -alkyl, heterocyclo, heteroaryl, keto, =N--OH, =N--O--lower alkyl, and a five or six membered ketal, *i.e.*, 1,3-dioxolane or 1,3-dioxane.

[0091] Exemplary monocyclic groups include azetidiny, pyrrolidiny, oxetany, imidazoliny, oxazolidiny, isoxazoliny, thiazolidiny, isothiazolidiny, tetrahydrofurany, piperidiny, piperaziny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolodiny, 2-oxoazepiny, azepiny, 4-piperidony, tetrahydropyrany, morpholiny, thiamorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone, 1,3-dioxolane and

tetrahydro-1, 1-dioxothienyl and the like. Exemplary bicyclic heterocyclo groups include quinuclidinyl.

[0092] The term “heteroaryl” refers to substituted and unsubstituted aromatic 5 or 6 membered monocyclic groups, 9 or 10 membered bicyclic groups, and 11 to 14 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heteroaryl group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less and each ring has at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups can contain only carbon atoms and can be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms can optionally be oxidized and the nitrogen atoms can optionally be quaternized. Heteroaryl groups which are bicyclic or tricyclic must include at least one fully aromatic ring but the other fused ring or rings can be aromatic or non-aromatic. The heteroaryl group can be attached at any available nitrogen or carbon atom of any ring. The heteroaryl ring system can contain one, two or three substituents, such as halo, amino, cyano, alkyl, substituted alkyl, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, alkoxy, alkylthio, hydroxy, nitro, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, --CO₂H, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, --(C=O)NH₂, --(C=O)N H(alkyl), --(C=O)NH(cycloalkyl), --(C=O)N(alkyl)₂, --NH--CH₂ --CO₂H, --NH--CH₂ --CO₂ -alkyl, heterocylco, and heteroaryl.

[0093] Exemplary monocyclic heteroaryl groups include pyrrolyl, pyrazolyl, pyrazolinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl and the like.

[0094] Exemplary bicyclic heteroaryl groups include indolyl, benzothiazolyl, benzodioxolyl, benzoxazolyl, benzothienyl, quinoliny, tetrahydroisoquinoliny, isoquinoliny, benzimidazolyl, benzopyranyl, indoliziny, benzofuranyl, chromonyl, coumariny, benzopyranyl, cinnoliny, quinoxaliny, indazolyl, pyrrolopyridyl, furopyridiny, dihydroisoindolyl, tetrahydroquinoliny and the like.

[0095] Exemplary tricyclic heteroaryl groups include carbazolyl, benzidolyl, phenanthrolliny, acridiny, phenanthridiny, xanthenyl and the like.

[0096] The term “substituted imidazole” refers to an imidazole, an aryl-fused imidazole such as benzimidazole, or a heteroaryl-fused imidazole such as a pyridoimidazole which contain one or two substituents, such as hydrogen, alkyl,

substituted alkyl, alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, --CO₂ H, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, --(C=O)NH₂, --(C=O)NH(alkyl), --(C=O)NH(cycloalkyl), --(C=O)N(alkyl)₂, --NH--CH₂ --CO₂ H, --NH--CH₂ --CO₂ -alkyl, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, heterocyclo, and heteroaryl.

[0097] The term “substituted triazole” refers to a triazole having at least one substituent, such as alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, --NH(alkyl), --NH(cycloalkyl), --N(alkyl)₂, --CO₂ H, --CO₂ -alkyl, cycloalkyl, substituted cycloalkyl, --(C=O)NH₂, --(C=O)NH(alkyl), --(C=O)NH(cycloalkyl), --(C=O)N(alkyl)₂, --NH--CH₂ --CO₂ H, --NH--CH₂ --CO₂ -alkyl, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, heterocyclo, and heteroaryl.

[0098] The terms “(C1 -C3) alkyl”, “(C1 -C4) alkyl”, and “(C1 -C10) alkyl”, when used alone, refer to straight chain alkyl radicals.

[0099] The terms “branched (C3 -C4) alkyl”, and “branched (C3 -C6) alkyl” refer to all alkyl isomers containing the designated number of carbon atoms, excluding the straight chain isomers.

[0100] The terms “(C1 -C4) alkoxy” and “(C1 -C7) alkoxy” refer to straight or branched chain alkoxy groups.

[0101] The term “halo (C1 -C7) alkyl” refers to a (C1 -C7) alkyl group, straight chain or branched, substituted with one or more halo groups.


[0102] The term “substituted phenyl” used alone or in combination with other terms, as in “substituted phenylthio” or “substituted phenylsulfonyl”, refers to phenyl substituted with up to three groups, such as halo, I, (C1 -C10) alkyl, branched (C3 -C6) alkyl, halo (C1 -C7) alkyl, hydroxy (C1 -C7) alkyl, (C1 -C7) alkoxy, halo (C1 -C7) alkoxy, phenoxy, phenyl, NO₂, OH, CN, (C1 -C4) alkanoyloxy, or benzyloxy.

[0103] The term “substituted phenoxy” refers to phenoxy substituted with at least one group, such as halo, I, (C1 -C10) alkyl, branched (C3 -C6) alkyl, halo (C1 -C7) alkyl, hydroxy (C1 -C7) alkyl, (C1 -C7) alkoxy, halo (C1 -C7) alkoxy, phenoxy, phenyl, NO₂, OH, CN, (C1 -C4) alkanoyloxy, or benzyloxy.

[0104] The terms “substituted naphthyl”, “substituted pyridyl” and “substituted furanyl” refer to these ring systems substituted with at least one group such as, halo, halo (C1 -C4) alkyl, CN, NO₂, (C1 -C4) alkyl, (C3 -C4) branched alkyl, phenyl, (C1 -C4) alkoxy, or halo (C1 -C4) alkoxy.

[0105] The term “unsaturated hydrocarbon chain” refers to a hydrocarbon chain containing one or two sites of unsaturation.

[0106] The preferred embodiments provide quinoline compounds of the following formula (I) or salts thereof, for use as inhibitors of RhoGTPases, especially Rac1GTPase:



(I)

[0107] wherein: R₁ to R₅ are independently: H, halo, (C1 -C4) alkyl, branched (C3-C4) alkyl, halo (C1 -C4) alkyl, (C1 -C4) alkoxy, NO₂, NH₂, -X-Alk, -X-Alk-X, -X-Y-X, -NR₆ or O-R₆, wherein

[0108] X is O, NR₆, or CR₇R₈;

[0109] Alk is a C2 -C18 saturated or unsaturated hydrocarbon chain, straight chain or branched, optionally substituted with halo, halo (C1 -C4) alkoxy, (C3 -C8) cycloalkyl, hydroxy, or acetyl;

[0110] Y is an alkylene chain 2 to 8 carbon atoms long, that optionally includes an O, S, SO, SO₂, or NR₆ group, and optionally includes a saturated or unsaturated carbocyclic ring comprising three to seven carbon atoms, and optionally is substituted with (C1 -C3) alkyl, (C2 -C4) phenyl, (C3 -C8) cycloalkyl, hydroxy, halo, or (C1 -C4) acyl; and

[0111] Ar is 1,3-benzodioxolyl fluorenyl, pyridyl substituted pyridyl, indolyl, furanyl, substituted furanyl, thienyl, optionally substituted with CH₂ or Cl, thiazolyl, cyclopentyl, 1-methylcyclopentyl, cyclohexenyl (tetrahydrophenyl), cyclohexyl (hexahydrophenyl), naphthyl, substituted naphthyl, dihydronaphthyl, tetrahydronaphthyl, or decahydronaphthyl;

[0112] R₆ is H, (C1 -C4) alkyl, or acetyl;

[0113] R₇ and R₈ are independently H, (C1 -C4) alkyl, (C1 -C4) acyl, halo, -OH, O-Y-Ar, or -NR₉ -Y-Ar; and

[0114] R₉ is H, (C1 -C4) alkyl, or acetyl.

[0115] or a salt of a compound of formula (I), provided, however, that this specifically excludes compounds that are known per se or that could be considered similar to known compounds.

[0116] Preferably at least two of R_1 to R_5 being H or CH_3 , and at least one of R_1 to R_2 is -X-Alk, -X-Alk-X or -X-Y-X, -NR₆ or O-R₆ and the rest of R_1 to R_5 are H or CH_3 ; wherein:

[0117] X is O, NR₆, or CR₇R₈;

[0118] Alk is a C2 -C18 saturated or unsaturated hydrocarbon chain, straight chain or branched, optionally substituted with halo, halo (C1 -C4) alkoxy, (C3 -C8) cycloalkyl, hydroxy, or acetyl;

[0119] Y is an alkylene chain 2 to 8 carbon atoms long, that optionally includes an O, S, SO, SO₂, or NR₆ group, and optionally includes a saturated or unsaturated carbocyclic ring comprising three to seven carbon atoms, and optionally is substituted with (C1 -C3) alkyl, (C2 -C4) phenyl, (C3 -C8) cycloalkyl, hydroxy, halo, or (C1 -C4) acyl; and


[0120] Ar is 1,3-benzodioxolyl fluorenyl, pyridyl substituted pyridyl, indolyl, furanyl, substituted furanyl, thienyl, optionally substituted with CH_2 or Cl, thiazolyl, cyclopentyl, 1-methylcyclopentyl, cyclohexenyl (tetrahydrophenyl), cyclohexyl (hexahydrophenyl), naphthyl, substituted naphthyl, dihydronaphthyl, tetrahydronaphthyl, or decahydronaphthyl;

[0121] R₆ is H, (C1 -C4) alkyl, or acetyl;

[0122] R₇ and R₈ are independently H, (C1 -C4) alkyl, (C1 -C4) acyl, halo, -OH, O-Y-Ar, or -NR₉-Y-Ar; and

[0123] R₉ is H, (C1 -C4) alkyl, or acetyl.

[0124] Preferably, the preferred embodiments provide compounds of the formula (II) or salts thereof, for use as inhibitors of RhoGTPases:



(II)

[0125] wherein:

[0126] R_1 to R_2 are independently: H, halo, (C1 -C4) alkyl, branched (C3-C4) alkyl, halo (C1 -C4) alkyl, (C1 -C4) alkoxy, NO_2 , NH_2 , -X-Alk, -X-Alk-X, -X-Y-X, -NR_6 or O-R_6 , wherein

[0127] X is O, NR_6 , or CR_7R_8 ;

[0128] Alk is a C2 -C18 saturated or unsaturated hydrocarbon chain, straight chain or branched, optionally substituted with halo, halo (C1 -C4) alkoxy, (C3 -C8) cycloalkyl, hydroxy, or acetyl;

[0129] Y is an alkylene chain 2 to 8 carbon atoms long, that optionally includes an O, S, SO, SO_2 , or NR_6 group, and optionally includes a saturated or unsaturated carbocyclic ring comprising three to seven carbon atoms, and optionally is substituted with (C1 -C3) alkyl, (C2 -C4) phenyl, (C3 -C8) cycloalkyl, hydroxy, halo, or (C1 -C4) acyl; and

[0130] Ar is 1,3-benzodioxolyl fluorenyl, pyridyl substituted pyridyl, indolyl, furanyl, substituted furanyl, thienyl, optionally substituted with CH_2 or Cl, thiazolyl, cyclopentyl, 1-methylcyclopentyl, cyclohexenyl (tetrahydrophenyl), cyclohexyl (hexahydrophenyl), naphthyl, substituted naphthyl, dihydronaphthyl, tetrahydronaphthyl, or decahydronaphthyl;


[0131] R_6 is H, (C1 -C4) alkyl, or acetyl;

[0132] R_7 and R_8 are independently H, (C1 -C4) alkyl, (C1 -C4) acyl, halo, -OH, O-Y-Ar, or -NR_9 -Y-Ar; and

[0133] R_9 is H, (C1 -C4) alkyl, or acetyl.

[0134] or a salt of a compound of formula (II).

[0135] Preferably, the preferred embodiments provide compounds of the formula (IIa) or salts thereof, for use as inhibitors of RhoGTPases:



[0136] wherein:

[0137] R_1 to R_2 are independently: H, -X-Alk, -X-Alk-X', or -X-Y-X', wherein

[0138] X is $-CR_7R_8$;

[0139] X' is $-CHR_7R_8$;

[0140] Alk is a C2 -C18 saturated or unsaturated hydrocarbon chain, straight chain or branched, optionally substituted with halo, halo (C1 -C4) alkoxy, (C3 -C8) cycloalkyl, hydroxy, or acetyl;


[0141] Y is an alkylene chain 2 to 8 carbon atoms long, that optionally includes an NR_6 group;

[0142] R_6 is H or (C1 -C4) alkyl; and

[0143] R_7 and R_8 are independently H or (C1 -C4) alkyl;

[0144] or a salt of a compound of formula (IIa).

[0145] The preferred embodiments provide compounds of the formula (III) or salts thereof, for use as inhibitors of Rho GTPases:




(III)

[0146] wherein:

[0147] R_{10} to R_{12} are independently: H, halo, (C1 -C4) alkyl, branched (C3-C4) alkyl, halo (C1 -C4) alkyl, (C1 -C4) alkoxy, NO_2 , or NH_2 ;

[0148] or a salt of a compound of formula (III).

[0149] The preferred embodiments provide compounds of the formula (IIIa) or salts thereof, for use as inhibitors of Rho GTPases:




(IIIa)

[0150] wherein:

[0151] R_{10} to R_{12} are independently: H, (C1 -C4) alkyl, or branched (C3-C4) alkyl;

[0152] or a salt of a compound of formula (IIIa).

[0153] The preferred embodiments provide compounds of the formula (IV) or salts thereof, for use as inhibitors of Rho GTPases:



(IV)

[0154] or a salt of a compound of formula (IV).

[0155] The pharmaceutical compositions of the preferred embodiments comprise a disease inhibiting and pharmaceutically acceptable amount of a compound of formula I, II, IIa, III, IIIa or IV, or N-6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine, in combination with a pharmaceutically-acceptable carrier.

[0156] The pharmaceutical compositions of the preferred embodiments comprise at least 1% by weight of a compound of formula I, II, IIa, III, IIIa or IV, or N-6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine (*e.g.*, formula IV, chemical compound 23766).

[0157] The pharmaceutical compositions of the preferred embodiments comprise a compound of formula I, II, IIa, III, IIIa or IV, or N⁶-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine (*e.g.*, formula IV, chemical compound 23766) further comprising a pharmaceutically active compound. For example, the additional pharmaceutically active compound can be a compound that is useful for inhibiting cell proliferation. For example, the additional pharmaceutically active compound can be a compound selected from the group consisting of farnesyl protein transferase inhibitors, prenyl-protein transferase inhibitors, geranylgeranyl-protein transferase inhibitors, toxins and combinations thereof.

[0158] The pharmaceutical combinations of the preferred embodiments comprise at least 1% by weight of a compound of formula I, II, IIa, III, IIIa or IV, or N⁶-

(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine (*e.g.*, formula IV, chemical compound 23766), in combination with a second pharmaceutical compound.

[0159] In another embodiment, the pharmaceutical combinations of the preferred embodiments comprise at least 1% by weight of a Rho family GTPase-regulating active compound further comprising additional cancer treatment pharmaceutical agent and, preferably, in combination with a pharmaceutically-acceptable carrier.

[0160] The pharmaceutical compositions of the preferred embodiments comprise a cancer prevention amount of a Rho family GTPase-regulating active compound in combination with a pharmaceutically-acceptable carrier.


[0161] The pharmaceutical methods of the preferred embodiments comprise administering N6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine to a subject in need of such treatment with a therapeutic amount of a compound of formula I, II, IIa, III, IIIa or IV, or of a combination described above.

[0162] Rac is one of the most important members of a family of proteins known as the Rho GTPases. This family of proteins binds to a small metabolic product called GTP, which acts as a critical regulator of Rho GTPase activity. This enables Rac to regulate a wide variety of cellular functions that span the entire gamut of a cell's life, from its initial growth and differentiation, to its movement and division, and finally to its death. They are important for gene expression, and they play crucial roles in the ability of innate immune cells to make lethal responses to bacterial infections, of skin cells to cover wounds during the healing process, of vascular cells to make new blood vessels, of cancer cells to metastasize, and of neurons to develop and make proper connections in the brain.

The inactive form of Rac-1 GTPase is the GDP-bound form. Provided herein is the three-dimensional structure of the inactive GDP-bound form, which contains conformational differences relative to the GTP-bound form of Rac-1 GTPase.


[0163] The present embodiments relate to crystalline protein corresponding to GDP-bound Rac-1 GTPase. Preferably, the crystalline protein is of sufficient quality to allow for the determination of the three-dimensional X-ray diffraction structure. The structural coordinates of a crystalline protein corresponding to Rac-1 GTPase is given in Table 1 or Table 2 below.

[0164] In one embodiment, a crystal structure of Rac-1 GTPase complexed



with (IIa) wherein:

R₁ to R₂ are independently selected from the group consisting of H, -X-Alk, -X-Alk-X', and -X-Y-X'; wherein X is -CR₇R₈; X' is -CHR₇R₈; Alk is a C₂-C₁₈ substituted or unsubstituted hydrocarbon chain; Y is a C₂-C₈ substituted or unsubstituted alkylene chain; R₆ is H or (C1 -C4) alkyl; and R₇ and R₈ are independently selected from the group consisting of H or (C1 -C4) alkyl; or a salt of a compound of formula (IIa) is obtained. In one embodiment, the compound of Formula IIa is



, Formula (IV). In the crystal structure

of Rac-1 GTPase complexed with a compound of Formula IV, the compound of Formula IV makes contacts with any of Trp-56, Lys-5, Val-7, Ser-71, Ile-36, Tyr-64, Arg-68, and Pro-73 of Rac-1 GTPase.

[0165] The embodiments also relate to methods for preparing and crystallizing the protein. Crystals of Rac-1 GTPase can be grown by a number of techniques known by those in the art, including batch crystallization, vapor diffusion (either by sitting drop or hanging drop) and by microdialysis. Seeding of the crystals in some instances is required to obtain X-ray quality crystals. Standard micro and/or macro seeding of crystals may therefore be used. An initial crystal can be allowed to grow over several hours to several months at about 4°C to about 20°C from a hanging drop. Crystals then can be subsequently grown by macroseeding from the initial crystal.

[0166] Once a crystal of the preferred embodiments is grown, X-ray diffraction data can be collected. A MAR imaging plate detector for X-ray diffraction data collection can be used for example. Crystals can be characterized by using X-rays produced in a conventional source (such as a sealed tube or a rotating anode) or using a

synchrotron source. In the example below, diffraction data was measured at National Synchrotron Light Source (Brookhaven, Upton, NY).

[0167] Methods of characterization include, but are not limited to, precession photography, oscillation photography, Laue diffraction photography, and diffractometer data collection. As exemplified below, heavy atom derivatives can be obtained by soaking crystals in stabilization solution with 1 mM Na_2OsCl_6 for about 12 hours, with 10 mM $\text{KAu}(\text{CN})_2$ for about 12 hours, with 10 mM $\text{Pb}(\text{OAc})_2$ for about 1 hour, and with 10 mM $\text{UO}_2(\text{NO}_3)_2$ for about 4 hours.

[0168] The protein itself, as well as information derived from the crystal structure can be used to analyze and modify protein activity as well as to identify compounds that interact with a catalytic domain. Once the three-dimensional structure of a crystal comprising a Rac-1 GTPase is determined, (e.g., see the coordinates in Table 1 or Table 2) a potential modulator of Rac-1 GTPase activity, can be examined through the use of computer modeling using a docking program such as, but not limited to, GRAM, DOCK, or AUTODOCK [Dunbrack et al., 1997, supra], to identify potential modulators of Rac-1 GTPase. This procedure can include computer fitting of potential modulators to Rac-1 GTPase to ascertain how well the shape and the chemical structure of the potential modulator will bind to a binding region of Rac-1 GTPase. Computer programs can also be employed to estimate the attraction, repulsion, and steric hindrance of the two binding partners. Generally the tighter the fit, the lower the steric hindrances, and the greater the attractive forces, the more potent the potential modulator since these properties are consistent with a tighter binding constant. Furthermore, the more specificity in the design of a potential drug the more likely that the drug will not interact as well with other proteins. This will minimize potential side-effects due to unwanted interactions with other proteins.

[0169] "Computer readable media" as used herein means any media which can be read and accessed directly by a computer *e.g.*, so that the media is suitable for use in the above-mentioned computer system. The media include, but are not limited to: magnetic storage media such as floppy discs, hard disc storage medium and magnetic tape; optical storage media such as optical discs or CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media.

[0170] “Computer system” as used herein means the hardware means, software means and data storage means used to analyze atomic coordinate data. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means and data storage means. Desirably a monitor is provided to visualize structure data. The data storage means may be RAM or means for accessing computer readable media of the invention. Examples of such systems are microcomputer workstations available from Silicon Graphics Incorporated and Sun Microsystems running Unix based, Windows NT or IBM OS/2 operating systems.

[0171] Some embodiments relate to methods of identifying or designing GTPase inhibitors. The methods can include the step of accessing at least a portion the data of Table 1 or Table 2, and modeling one or more compounds utilizing said data. The modeling can include, for example, predicting the likelihood of a compound to bind to the Rac-1 GTPase by modeling the interaction of a compound with a GTPase activation site of the Rac-1 GTPase. The methods further can include utilizing the data to identify one or more interactions between a Rac-1 GTPase and a ligand. For example, the interactions can include one or more hydrogen bonds between ligand and protein; one or more electrostatic interactions between ligand and protein; one or more hydrophobic interactions; one or more covalent bonds between ligand and protein; one or more changes in the positions of protein atoms; such as amino acid side chains upon ligand binding; one or more atoms whose positions are well defined versus those that may maintain flexibility in the complex; one or more hydrogen bonding networks; one or more interactions that are important to the catalytic mechanism of the protein/enzyme; one or more structural changes to the hydrogen bond network or the interactions which occur in the presence of the ligand; one or more interactions between the ligand and the protein that interfere with or prevent activation of Rac-1 GTPase; one or more key interactive forces between the ligand and the protein based on the collective data; one or more modifications to the existing ligand that enhance binding to the protein. The modification to the existing ligand can include, for example, adding the potential for electrostatic interactions, reducing steric interactions, increasing hydrophobic interactions, and increasing the surface area of the interaction. Also, the methods can include modeling a compound or compounds to include the interactions described above, or improvements of the interactions.

[0172] Some embodiments relate to methods of designing inhibitors, with or without accessing the data of Table 1 or Table 2, by identifying a compound that prevents activation of Rac-1 GTPase. Typically, a compound that prevents activation of Rac-1 GTPase acts by one or more of the following mechanisms: inhibits Thr35 binding of Mg^{2+} , inhibits conformational change in the loop containing amino acid residues 60-64, inhibits Ala59 from displacing Mg^{2+} , inhibits a RAC-specific GEF from binding to RAC, inhibits RAC-specific GEF from dissociating from RAC, inhibits conformational change in the amino acids of switch I, or inhibits conformational change in the amino acids of switch II.

[0173] New compounds can be designed which inhibit Rac-1 GTPase by binding the region between switch I and switch II on Rac-1. Some embodiments relate to methods of identifying a Rac-1 GTPase inhibitor comprising accessing one or more of the coordinates of Table 1 or Table 2 and identifying a compound that occupies a position within about 5, 4.5, 4, 3.5, 3, 2.5, 2, 1.5 or 1 angstroms of one or more of the atomic coordinates of the ligand (DRG) in said Table. For example, a Rac-1 GTPase inhibitor can be identified as a compound that occupies a position between about 5 and about 2.5 angstroms, about 4.5 and about 3 angstroms, about 3.5 and about 2.6 angstroms, or about 3.2 and about 2.6 angstroms from one or more of the atomic coordinates of amino acids Thr35, Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73, and the Mg^{2+} ion as provided in Table 1 or Table 2. For example, a Rac-1 GTPase inhibitor can be identified as a compound that is involved in hydrophobic contact, van der Waals contact, hydrogen bond formation, salt bridge formation, or other physical interaction with one or more of the atomic coordinates of amino acids Thr35, Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73, and the Mg^{2+} ion as provided in Table 1 or Table 2. In another example, a Rac-1 GTPase inhibitor can be identified as a compound that occupies a position between about 2 and about 1 angstroms, about 1.7 and about 1.2 angstroms, or about 1.6 and about 1.3 angstroms from one or more of the atomic coordinates of amino acids Thr35, Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 as provided in any Table 1 or Table 2. For example, a Rac-1 GTPase inhibitor can be identified as a compound that is covalently bound to one or more of the atomic coordinates of amino acids Thr35, Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 as provided in Table 1 or Table 2. The method can include identifying a distance between an atom of the Rac-1 GTPase and an atom of an inhibitor, and substituting the inhibitor atom with a

different atom or with a functional group to result in a modified interaction. For example, the modified interaction can be a decreased distance between the ligand and the Rac-1 GTPase, which can permit, e.g., hydrophobic interactions, van der Waals interactions, hydrogen bonding or electrostatic interactions. Thus, some aspects can relate to methods of identifying a Rac-1 GTPase inhibitor that include the steps of identifying one or more interactions between a ligand and Rac-1 GTPase; and changing a substituent on the ligand in order to modify the interaction between the ligand the subunit. An exemplary ligand for which a substituent can be modified is NSC23766.

[0174] Still some embodiments relate to methods of screening a set of one or more compounds to identify compounds most likely to bind to a Rac-1 GTPase. The methods can include the steps of: (a) computationally determining the docking energy of each of one or more compounds in a GTPase activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site; and (b) selecting one or more compounds indicated as having the highest likelihood of binding to said Rac-1 GTPase activation site.

[0175] Also, some embodiments relate to methods of identifying a compound that is likely to bind the active site of a Rac-1 GTPase, which methods can include the steps of: (a) computationally determining the docking energy of each of one or more compounds in a GTPase activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site; and (b) selecting one or more compounds indicated as having the highest likelihood of binding to said Rac-1 GTPase activation site.

[0176] Some embodiments relate to methods of screening compounds that bind to the Rac-1 GTPase so as to inhibit GTPase activity of the Rac-1 GTPase. Such methods can include predicting the likelihood of a candidate compound to bind a Rac-1 GTPase activation site. An exemplary Rac-1 GTPase activation site is a region located between switch I and switch II of the Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid of switch I and at least one amino acid of switch II. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase (see SEQ ID NO:1) and at least one amino acid selected from amino acids 56-74 of Rac-1 GTPase (see SEQ ID NO:1). Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 and Asp38 of Rac-1

GTPase and at least one amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises Trp56 of Rac-1 GTPase and at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Trp56 and Leu70 of Rac-1 GTPase and at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Trp56, Leu70 and Ser71 of Rac-1 GTPase and at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Trp56, Tyr64, Leu67 and Leu70 of Rac-1 GTPase and at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 and at least one amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 or Asp38 of Rac-1 GTPase and at least one amino acid selected from amino acids 56-70 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. Another exemplary Rac-1 GTPase activation site comprises at least one amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. An exemplary Rac-1 GTPase activation site such as those above also can comprise the Mg^{2+} atom.

[0177] Methods for predicting the likelihood of a candidate compound to bind a particular site of a protein are known in the art, and any of these known methods can be used in accordance with the teachings provided herein to evaluate compounds for their ability to bind a Rac-1 GTPase activation site so as to inhibit Rac-1 GTPase activation.

As provided herein, inhibition of Rac-1 GTPase activity can be accomplished by inhibiting Thr35 binding of Mg^{2+} , inhibiting conformational change in the loop containing amino acid residues 60-64, inhibits Ala59 from displacing Mg^{2+} , inhibiting a RAC-specific GEF from binding to RAC, inhibiting RAC-specific GEF from dissociating from RAC, inhibiting conformational change in the amino acids of switch I, or inhibiting conformational change in the amino acids of switch II. Using the three dimensional atomic structure of the Rac-1 GTPase alone or in complex with NSC23766 (see Table 1 or 2) and known modeling methods, it is possible to screen candidate compounds for their ability to bind to a Rac-1 GTPase activation site, such as those recited above.

[0178] The ability of a potential compound to bind a Rac-1 GTPase activation site can be examined through the use of computer modeling using a known docking program such as Hammerhead, FlexX, MCDOCK, ICM-dock, QXP, GOLD, CHARMM, GRAM, DOCK, or AUTODOCK (see, e.g., Walters et al., Drug Discovery Today, Vol. 3, No. 4, (1998), 160-178, and Dunbrack et al., Folding and Design, 2, (1997), 27-42). This procedure can include computer fitting of potential ligands to the Rac-1 GTPase activation site formed by the residues of the activation site, such as those recited above, to ascertain how well the shape and the chemical structure of the potential ligand will match the corresponding active site. The docking methods used in the art can function using any of a variety of different algorithms and optimizations, however generally, these methods function by indicating the goodness of fit of a compound in the binding site of the protein (e.g., the activation site of a Rac-1 GTPase). The goodness of fit can be expressed, for example, as a docking energy of a compound with a protein binding site. In such instances, docking energy can be a quantitative indicator of the likelihood of the compound to bind the Rac-1 GTPase activation site. The goodness of fit, such as docking energy, can serve as a convenient basis for evaluating candidate compounds as potential lead compounds for Rac-1 GTPase activation site binding. For example, when two or more compounds are modeled for their ability to bind to a Rac-1 GTPase activation site, a threshold or cutoff can be used to identify a subset of the two or more compounds that can be further examined as potential lead compounds. A threshold can be a designated percentage of compounds with highest likelihood of binding to a Rac-1 GTPase activation site, where such a designated percentage can be the top 1%, the top 2%, the top 3%, the top 4%, the top 5%, the top 6%, the top 7%, the top 8%, the top 9%, or the top 10%. Another threshold can be relative to a known binder of a Rac-1 GTPase activation site; for

example, a threshold can be set according to the ability of NSC23766 to bind a Rac-1 GTPase activation site. As an example, a compound predicted to bind to a Rac-1 GTPase activation site more readily than NSC23766 can be identified as a compound predicted to inhibit Rac-1 GTPase activity.

[0179] In some exemplary docking methods, a representation of a Rac-1 GTPase activation site is generated. Various types of representations of a protein binding site are known in the art, such as, for example, a grid-based field representation of a binding site. The representation of a Rac-1 GTPase activation site is typically based on the three dimensional coordinates of the atoms in the binding site. Three dimensional coordinates of the atoms in the binding site can be obtained from Table 1 or Table 2, where the atoms to be selected from Table 1 or Table 2 are selected from the atoms in the above-described exemplary Rac-1 GTPase activation sites, which will typically comprise at least one amino acid of switch I and at least one amino acid of switch II. In another example, a Rac-1 GTPase activation site representation can contain protein atoms within a designated distance from the location of a NSC23766 atom, where the designated distance can be, for example 5Å, 6Å, 7Å, 8Å, 9Å, 10Å, 11Å, 12Å, 13Å, 14Å, 15Å, 16Å, 17Å, 18Å, 19Å, or 20Å, as determined according to the atomic coordinates available in Table 1 and Table 2. A Rac-1 GTPase activation site representation also can include any water molecules observed in the Rac-1 GTPase activation site. The presence of water molecules in a Rac-1 GTPase activation site can be determined according to the atomic coordinates provided in Table 1 or Table 2. The number of atoms to include and the type of representation to be used can be determined according to the docking method used and the degree of accuracy versus computational efficiency desired.

[0180] While representations of Rac-1 GTPase activation sites can be determined according to the atomic coordinates provided in Table 1 or Table 2, one of skill in the art will recognize that the present embodiment also can encompass minor variations of the relative positions of the atoms in the representations of Rac-1 GTPase activation sites. For example, the root-mean-squared deviation (RMSD) between atoms of the activation site representation and the corresponding atoms of the atomic coordinates provided in Table 1 or Table 2 will typically be 6.0 Å or less, 5.5 Å or less, 5.0 Å or less, 4.5 Å or less, 4.0 Å or less, 3.5 Å or less, 3.0 Å or less, 2.5 Å or less, 2.0 Å or less, 1.7 Å or less, 1.5 Å or less, 1.4 Å or less, 1.3 Å or less, 1.2 Å or less, 1.1 Å or less, 1.0 Å or less, 0.9 Å or less, 0.8 Å or less, 0.7 Å or less, 0.6 Å or less, 0.5 Å or less,

0.4 Å or less, 0.3 Å or less, 0.2 Å or less, or 0.1 Å or less, as calculated for all atoms of the activation site representation or as calculated according to a subset of atoms used for the active site representation. When a subset of atoms is used to calculation the RMSD between atoms of the activation site representation and the corresponding atoms of the atomic coordinates provided in Table 1 or Table 2, the subset of atoms are typically those provided in the above-described exemplary Rac-1 GTPase activation sites. In particular, an exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid of switch I and at least one atom from an amino acid of switch II. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase (see SEQ ID NO:1) and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase (see SEQ ID NO:1). Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 and Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from Trp56 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Trp56 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Trp56, Leu70 and Ser71 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary set of atoms of

the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Trp56, Tyr64, Leu67 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 or Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-70 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation includes at least one atom from an amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. Another exemplary set of atoms of the Rac-1 GTPase activation site for use in RMSD calculation such as those above also can include the Mg^{2+} atom. In such exemplary set of atoms above, the atoms can be selected from all atoms of a selected amino acid, backbone atoms only of a selected amino acid, or alpha carbon atoms only of a selected amino acid.

[0181] The x-ray crystal structures described herein in Tables 1 and 2 report the atomic coordinates of the heretofore unknown three-dimensional structure of the GDP-bound, inactive form of Rac-1 GTPase. This three-dimensional structure of the GDP-bound form was compared with the active or GTP-bound form of Rac-1 GTPase (PDB ID# 1MH1) to understand at the molecular level the switch mechanism or conformational change involved in Rac-1 GTPase activation.

[0182] The calculated root mean square deviation (rmsd) after superposition of all C α s of both structures is 0.95 Å. Both structures superpose well outside one polypeptide spanning amino acid residues 30 to 40. This region is typically referred to as the switch I region. Another major difference between the two structures is in the way the Mg^{2+} ion is coordinated. As in other GTP-binding proteins including Ras, in the GTP-

bound form of Rac the Mg^{2+} ion is coordinated by the hydroxyl side chain group of Thr35, which is also making a hydrogen bond with the γ -phosphate. The absence of the γ -phosphate in the GDP-bound form of Rac results in the change of conformation of Thr35, which in Rac•GDP projects into the solvent. This change of conformation of Thr35 results in the change of conformation of the switch I region of Rac. For example, the ψ angle of Pro29 changes by 150° between the two conformations. Table 3 below lists the changes in distance of the C α s for the residues of switch I after superposition of both Rac-1 structures:

Table 3

<u>Residue</u>	<u>Change in Distance (Å)</u>
Gly30	3.9
Glu31	0.74
Tyr32	5.6
Ile33	4.8
Pro34	5.1
Thr35	1.6
Val36	4.0
Phe37	6.0
Asp38	2.5
Asn39	0.7

[0183] In some exemplary docking methods, the conformation of the compound in the Rac-1 GTPase activation site is calculated. Various methods for calculating the conformation of a ligand in a protein binding site are known for docking methods. These can include determining optimal conformation(s) of a compound outside of and within the active site. These also can include building the atoms into the active site such that the atoms match the binding site (sterically, chemically, electrically, or combinations thereof). Any such method can be performed in the docking methods provided herein.

[0184] Any of a variety of candidate compounds can be used in the modeling methods provided herein. Methods of predicting the ability of a candidate compound to bind to a Rac-1 GTPase activation site can be used to examine a plurality of candidate

compounds in evaluating and comparing the ability of compounds of the plurality to bind to the Rac-1 GTPase activation site. A number of exemplary public and commercial sources of libraries of compound structures are available, for example the Cambridge Structural Database (CSD), the Chemical Directory (ACD) from the company MDL (US), ZINC (Irwin and Shoichet, J. Chem. Inf. Model. (2005) 45:177-82) as well as various electronic catalogues of publicly available compounds such as the National Cancer Institute (NCI, US) catalogue, ComGenex catalogue (Budapest, Hungary), and Asinex (Moscow, Russia). Such libraries may be used to allow computer-based docking of many compounds in order to identify those with potential to interact with the Rac-1 GTPase activation site described herein.

[0185] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to block access of a RAC-specific GEF protein from interacting with the Rac-1 GTPase activation site. While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase can be activated by a RAC-specific GEF binding the Rac-1 GTPase activation site, and thus blocking access of a RAC-specific GEF to the Rac-1 GTPase activation site by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of blocking protein-protein interactions are known in the art, such as determination of steric blockage of protein-protein interaction, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0186] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to block dissociation of a RAC-specific GEF protein from the Rac-1 GTPase activation site. While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase is not completely activated by GEF binding the Rac-1 GTPase activation site until dissociation of the GEF from Rac-1, and thus blocking dissociation of a GEF from the Rac-1 GTPase activation site by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of promoting protein-protein interactions are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art,

computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0187] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to inhibit Thr35 of Rac-1 GTPase binding to Mg^{2+} . While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase activation includes Thr35 binding to Mg^{2+} , and thus blocking Thr35 binding to Mg^{2+} by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of blocking amino acid-ion interactions are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0188] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to inhibit conformational change in the loop containing amino acid residues 60-64. While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase activation includes conformational change in the loop containing amino acid residues 60-64, and thus blocking such a conformational change by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of conformational changes of protein structural elements are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0189] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to inhibit Ala59 from displacing Mg^{2+} . While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase activation includes Ala59 from displacing Mg^{2+} in exchanging GDP for GTP, and thus blocking Ala59 from displacing Mg^{2+} by a compound can increase the Rac-1 GTPase inhibiting activity of the compound by increasing the inhibition of exchange of GDP to GTP. Methods for

predicting the likelihood of blocking amino acid-ion interactions are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0190] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to inhibit conformational change in the amino acids of switch I, in particular amino acids 30-39 of SEQ ID NO:1, such as amino acids Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 and Asp38. While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase activation includes conformational change in amino acids of switch I, and thus blocking such a conformational change by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of conformational changes of protein structural elements are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0191] In some instances, methods can be used in which a determination can be made of whether or not a candidate compound is likely to inhibit conformational change in the amino acids of switch II. While not intending to be limited by the following theory, it is postulated based on the present observations that Rac-1 GTPase activation includes slight conformational change in amino acids of switch II, and thus blocking such a conformational change by a compound can increase the Rac-1 GTPase inhibiting activity of the compound. Methods for predicting the likelihood of conformational changes of protein structural elements are known in the art, and can be used in conjunction with computerized modeling methods provided herein or otherwise known in the art, computer-assisted manual examination methods provided herein or otherwise known in the art, or with manual modeling and design of candidate compounds for binding a Rac-1 GTPase activation site.

[0192] Also computer-assisted, manual examination of a Rac-1 GTPase activation site may be performed. The use of programs such as GRID (Goodford, J. Med. Chem., 28, (1985), 849-857)--a program that determines probable interaction between

molecules with various functional groups and the protein surface--may also be used to analyze the binding site to predict partial structures of ligands for the site. Computer programs can be employed to estimate the attraction, repulsion, and steric hindrance of the candidate compound and the Rac-1 GTPase activation site described herein. Generally, the tighter the fit, the fewer the steric hindrances, and the greater the attractive forces, the more potent the potential ligand since these properties are consistent with a tighter binding constant. Furthermore, the more specificity in the design of a potential ligand, the more likely it is that the ligand will not interact with other proteins as well. This will tend to minimize potential side-effects due to unwanted interactions with other proteins.

[0193] Having designed or selected possible binding compounds, these can then be screened for activity. Exemplary methods for screening binding compounds are provided herein, and include, but are not limited to those described in Macherla et al., *J. Med. Chem.* **2005** and U.S. Patent Publication No. 2005/0049294, entitled "Methods of using [3.2.0] heterocyclic compounds and analogs thereof," filed on June 18, 2004; both of which are hereby incorporated by reference in their entirety. Consequently, also provided herein are methods of evaluating a candidate compound predicted to bind to the Rac-1 GTPase activation site, where the method includes providing candidate compounds predicted to bind to the Rac-1 GTPase activation site in accordance with the methods described above, and contacting Rac-1 GTPase or a fragment thereof, with the candidate compound and determining whether the candidate compound bound to Rac-1 GTPase or fragment thereof. Also provided herein are methods of evaluating a candidate compound predicted to inhibit Rac-1 GTPase activity, where the method includes providing candidate compounds predicted to bind to Rac-1 GTPase in accordance with the methods described above, contacting a Rac-1 GTPase with the candidate compound, and determining whether the candidate compound inhibits Rac-1 GTPase activity.

[0194] The NSC23766 molecule described herein belongs to a new class of Rac-1 GTPase inhibitors, and acts by preventing activation of Rac-1 GTPase. The co-crystal structure of NSC23766 in complex with Rac-1 GTPase reveals that NSC23766 makes numerous van der Waals interactions with the switch I residue Val36 and the switch II residues Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. Hydrogen bonds were found involving the carbonyl of Leu70 and the side chain

oxygen of Ser71. Further possible hydrogen bonds involving the carbonyl of Leu67, side chain of Gln74, the carbonyl of Asp57, and the carbonyl of Ser71 were identified.

[0195] Based upon these observations, an improved inhibitor can be made by adding to or modifying NSC23766 in order to have a configuration of moieties that can hydrogen bond with the carbonyl of Leu70, the side chain oxygen of Ser71, the carbonyl of Leu67, the side chain of Gln74, the carbonyl of Asp57, and the carbonyl of Ser71. It was also observed in the crystal structure of Rac-1 GTPase without NSC23766 bound that a DMSO molecule specifically interacted with the amide main chain group of Ala59. Thus, a further addition or modification to NSC23766 can include a moiety that can hydrogen bond with amide main chain group of Ala59. Exemplary moieties that can be added as hydrogen bond donors include amine, amide and hydroxyl moieties; exemplary moieties that can be added as hydrogen bond acceptors include hydroxyl, carbonyl, and ether moieties.

[0196] Further in regard to the co-crystal observations, an improved inhibitor can be made by adding to or modifying NSC23766 in order to have a configuration of moieties having improved van der Waals and/or hydrophobic interactions with the switch I residue Val36 and the switch II residues Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase. Exemplary hydrophobic moieties that can be added include alkyl moieties, substituted alkyl moieties, aryl moieties and substituted aryl moieties.

[0197] Methods and compositions are described that affect the GTPase activity of members of the Ras superfamily, preferably Rac, such compositions include compounds that modulate the GTPase activity, along with uses for the compounds including screening for compounds which recognize Rac GTPase, and methods of treating pathological conditions associated or related to a Ras superfamily GTPase, including Rac. Another embodiment comprises binding to a Rho GTPase selected from the group consisting of Rac1, Rac2 and Rac3. Preferably, another embodiment comprises binding to a Rho GTPase that is Rac1. The preferred embodiments also relate to methods of using such compounds, or derivatives thereof, *e.g.*, in therapeutics, diagnostics, and as research tools.

[0198] Methods and compositions are described that affect the GTPase activity of members of the Ras superfamily, preferably Rac; such compositions include compounds that modulate the GTPase activity. Preferably, the indication associated with GTPase activity is selected from the group consisting of hypertension, atherosclerosis,

restenosis, cerebral ischemia, cerebral vasospasm, neuronal degeneration, spinal cord injury, cancer of the breast, colon, prostate, ovaries, brain or lung, thrombotic disorders, asthma, glaucoma, osteoporosis and erectile dysfunction.

[0199] The preferred embodiments also relate to methods of testing for and/or identifying agents that regulate Rac by measuring their effect on the ability of a Rho family GTPase-regulating active compound to regulate the action of Rac GTPase.

[0200] As used herein, the terms "Ras or Ras superfamily proteins" encompass a large family of GTP binding/GTP hydrolyzing monomeric proteins. Ras family includes the Ras, Rho, Rab, Arf, and Ran subfamilies of GTPases.

[0201] The terms "Rho GTPases" or "Rho family GTPases" refer to a subfamily of Ras superfamily and are small, membrane-bound, Ras-related GTP-binding proteins that function by binding and hydrolyzing GTP. Rho GTPases function as molecular switches, cycling between an inactive GDP-bound conformation and an active GTP-bound conformation and include RhoA, RhoB, RhoC, Cdc42, Rac1, Rac2, Rac3, TC10, RhoG, RhoD, Chp, WRCH1, TCL, and RIF.

[0202] A protein or polypeptide sequence of a Ras-related protein includes variants or fragments thereof derived from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic, or recombinant.

[0203] The terms "Rac GTPase" or "Rac protein or polypeptide" refer to Rac1, Rac2, and/or Rac3.

[0204] One embodiment provides for a method for reducing cancer cell proliferation by administering in a subject having cancer an effective amount of a Rho family GTPase-regulating active compound as defined herein.

[0205] Another embodiment provides for the use of an effective amount of a Rho family GTPase-regulating active compound as defined herein for the preparation of pharmaceutical composition for the treatment of a disease associated with abnormal cell proliferation.

[0206] Because of their cell proliferation inhibitory activity, the compounds of the preferred embodiments are suitable for treating a variety of diseases in a variety of conditions. In this regard, "treatment" or "treating" includes both therapeutic and prophylactic treatments. Accordingly, the compounds can be used at very early stages of a disease, or before early onset, or after significant progression, including metastasis. The

term "treatment" or "treating" designates in particular a reduction of the burden in a patient, such as a reduction in cell proliferation rate, a destruction of diseased proliferative cells, a reduction of tumor mass or tumor size, a delaying of tumor progression, as well as a complete tumor suppression.

[0207] Typical examples of diseases associated with abnormal cell proliferation include cancers and restenosis, for instance. The compounds of the preferred embodiments are particularly suited for the treatment of cancers, such as solid tumors or lymphoid tumors. Specific examples include leukemia, prostate cancer, ovarian cancer, pancreas cancer, lung cancer, breast cancer, liver cancer, head and neck cancer, colon cancer, bladder cancer, non-Hodgkin's lymphoma cancer and melanoma.

[0208] The active compounds of the preferred embodiments can be administered according to various routes, typically by injection, such as local or systemic injection(s). Intratumoral injections are preferred for treating existing cancers. However, other administration routes can be used as well, such as intramuscular, intravenous, intradermic, subcutaneous, *etc.* Furthermore, repeated injections can be performed, if needed, although it is believed that limited injections will be needed in view of the efficacy of the compounds.

[0209] It is contemplated that such target cells can be located within an animal or human patient, in which case a safe and effective amount of the complex, in pharmacologically acceptable form, would be administered to the patient. Generally speaking, it is contemplated that useful pharmaceutical compositions of the preferred embodiments will include the selected active compound derivative in a convenient amount, *e.g.*, from about 0.001% to about 10% (w/w) that is diluted in a pharmacologically or physiologically acceptable carrier, such as, for example, phosphate buffered saline. The route of administration and ultimate amount of material that is administered to the patient or animal under such circumstances will depend upon the intended application and will be apparent to those of skill in the art in light of the examples which follow.

[0210] Any composition chosen should be of low or non-toxicity to the cell. Toxicity for any given compound can vary with the concentration of compound used. It is also beneficial if the compound chosen is metabolized or eliminated by the body and if this metabolism or elimination is done in a manner that will not be harmfully toxic.

[0211] The examples are illustrative of the types of compounds to be used in the method claimed herein; the list is not exhaustive. Derivatives of the above compounds that fit the criteria of the claims are preferably also be considered when choosing an active compound.

[0212] The compound is preferably administered such that a therapeutically effective concentration of the compound is in contact with the affected cells of the body. The dose administered to an animal, particularly a human, in the context of the preferred embodiments is preferably sufficient to effect a therapeutic response in the animal over a reasonable period of time. The dose will be determined by the strength of the particular compound employed and the condition of the animal, as well as the body weight of the animal to be treated. The existence, nature, and extent of any adverse side effects that might accompany the administration of a particular compound also will determine the size of the dose and the particular route of administration employed with a particular patient. In general, the compounds of the preferred embodiments are therapeutically effective at low doses. The generally useful dose range is from about 0.001 mM, or less, to about 100 mM, or more. Preferably, the effective dose range is from about 0.01, 0.05, 0.1, 0.5, 0.6, 0.7, 0.8, or 0.9 mM, to about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mM. Accordingly, the compounds will be generally administered in low doses.

[0213] The compound can be administered in a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are well-known to those who are skilled in the art. The choice of carrier will be determined in part by the particular compound, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the preferred embodiments.

[0214] The compounds can be administered orally, topically, parenterally, by inhalation or spray, vaginally, rectally or sublingually in dosage unit formulations. The term "administration by injection" includes but is not limited to: intravenous, intraarticular, intramuscular, subcutaneous and parenteral injections, as well as use of infusion techniques. Dermal administration can include topical application or transdermal administration. One or more compounds can be present in association with one or more non-toxic pharmaceutically acceptable carriers and if desired other active ingredients.

[0215] Compositions intended for oral use can be prepared according to any suitable method known to the art for the manufacture of pharmaceutical compositions.

Such compositions can contain one or more agents selected from the group consisting of diluents, sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; and binding agents, for example magnesium stearate, stearic acid or talc. The tablets can be uncoated or they can be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed. These compounds can also be prepared in solid, rapidly released form.

[0216] Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

[0217] Aqueous suspensions containing the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions can also be used. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropyl-methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents can be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions can also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0218] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavoring and coloring agents, can also be present.

[0219] The compounds can also be in the form of non-aqueous liquid formulations, *e.g.*, oily suspensions which can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or peanut oil, or in a mineral oil such as liquid paraffin. The oily suspensions can contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents can be added to provide palatable oral preparations. These compositions can be preserved by the addition of an anti-oxidant such as ascorbic acid.

[0220] Compounds of the preferred embodiments can also be administrated transdermally using methods known to those skilled in the art. For example, a solution or suspension of an active agent in a suitable volatile solvent optionally containing penetration enhancing agents can be combined with additional additives known to those skilled in the art, such as matrix materials and bacteriocides. After sterilization, the resulting mixture can be formulated following known procedures into dosage forms. In addition, on treatment with emulsifying agents and water, a solution or suspension of an active agent can be formulated into a lotion or salve.

[0221] Suitable solvents for processing transdermal delivery systems are known to those skilled in the art, and include lower alcohols such as ethanol or isopropyl alcohol, lower ketones such as acetone, lower carboxylic acid esters such as ethyl acetate, polar ethers such as tetrahydrofuran, lower hydrocarbons such as hexane, cyclohexane or benzene, or halogenated hydrocarbons such as dichloromethane, chloroform, trichlorotrifluoroethane, or trichlorofluoroethane. Suitable solvents can also include mixtures of one or more materials selected from lower alcohols, lower ketones, lower carboxylic acid esters, polar ethers, lower hydrocarbons, halogenated hydrocarbons.

[0222] Suitable penetration enhancing materials for transdermal delivery system are known to those skilled in the art, and include, for example, monohydroxy or polyhydroxy alcohols such as ethanol, propylene glycol or benzyl alcohol, saturated or

unsaturated C8-C18 fatty alcohols such as lauryl alcohol or cetyl alcohol, saturated or unsaturated C8-C18 fatty acids such as stearic acid, saturated or unsaturated fatty esters with up to 24 carbons such as methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tertbutyl or monoglycerin esters of acetic acid, capronic acid, lauric acid, myristinic acid, stearic acid, or palmitic acid, or diesters of saturated or unsaturated dicarboxylic acids with a total of up to about 24 carbons such as diisopropyl adipate, diisobutyl adipate, diisopropyl sebacate, diisopropyl maleate, or diisopropyl fumarate. Additional penetration enhancing materials include phosphatidyl derivatives such as lecithin or cephalin, terpenes, amides, ketones, ureas and their derivatives, and ethers such as dimethyl isosorbide and diethyleneglycol monoethyl ether. Suitable penetration enhancing formulations can also include mixtures of one or more materials selected from monohydroxy or polyhydroxy alcohols, saturated or unsaturated C8-C18 fatty alcohols, saturated or unsaturated C8-C18 fatty acids, saturated or unsaturated fatty esters with up to 24 carbons, diesters of saturated or unsaturated dicarboxylic acids with a total of up to 24 carbons, phosphatidyl derivatives, terpenes, amides, ketones, ureas and their derivatives, and ethers.

[0223] Suitable binding materials for transdermal delivery systems are known to those skilled in the art and include polyacrylates, silicones, polyurethanes, block polymers, styrenebutadiene copolymers, and natural and synthetic rubbers. Cellulose ethers, derivatized polyethylenes, and silicates can also be used as matrix components. Additional additives, such as viscous resins or oils can be added to increase the viscosity of the matrix.

[0224] Pharmaceutical compositions of the preferred embodiments can also be in the form of oil-in-water emulsions. The oil phase can be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example, liquid paraffin or mixtures of these. Suitable emulsifying agents can be naturally-occurring gums, for example, gum acacia or gum tragacanth, naturally-occurring phosphatides, for example, soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate. The emulsions can also contain sweetening and flavoring agents. Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such

formulations can also contain a demulcent, a preservative and flavoring and coloring agents.

[0225] The compounds can also be administered in the form of suppositories for rectal or vaginal administration of the drug. These compositions can be prepared by mixing the drug with a suitable nonirritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature or vaginal temperature and will therefore melt in the rectum or vagina to release the drug. Such materials include cocoa butter and polyethylene glycols.

[0226] For all regimens of use disclosed herein for active agent, the daily oral dosage regimen will preferably be from about 0.01 to about 200 mg/Kg of total body weight. Preferably, the daily oral dosage regimen will preferably be from about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, or 5 to about 10, 50, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg/Kg of total body weight. The daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques will preferably be from 0.01 to 200 mg/Kg of total body weight. Preferably, the daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques will preferably be from about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, or 5 to about 10, 50, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg/Kg of total body weight. The daily vaginal dosage regime will preferably be from 0.01 to 200 mg/Kg of total body weight. The daily topical dosage regimen will preferably be from 0.01 to 200 mg administered between one to four times daily. The concentration for vaginal dosage and topical dosage will preferably be that required to maintain a daily dose is of from 0.1 to 200 mg/Kg. Preferably, the daily oral dosage regimen will preferably be from about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, or 5 to about 10, 50, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg/Kg of total body weight. The daily inhalation dosage regimen will preferably be from 0.01 to 10 mg/Kg of total body weight. Preferably, the daily inhalation dosage regimen will preferably be from about 0.01, 0.05, 0.1, 0.5, to about 1, 2, 3, 4, 5, or 10, mg/Kg of total body weight.

[0227] It will be appreciated by those skilled in the art that the particular method of administration will depend on a variety of factors, all of which are considered routinely when administering therapeutics. It will also be understood, however, that the specific dose level for any given patient will depend upon a variety of factors, including,

the activity of the specific compound employed, the age of the patient, the body weight of the patient, the general health of the patient, the gender of the patient, the diet of the patient, time of administration, route of administration, rate of excretion, drug combinations, and the severity of the condition undergoing therapy. It will be further appreciated by one skilled in the art that the optimal course of treatment, *i.e.*, the mode of treatment and the daily number of doses of an active agent or a pharmaceutically acceptable salt thereof given for a defined number of days, can be ascertained by those skilled in the art using conventional treatment tests.

[0228] Another aspect of the preferred embodiments relates to the regulation of biological pathways in which a GTPase is involved, particularly pathological conditions, *e.g.*, cell proliferation (*e.g.*, cancer), growth control, morphogenesis, stress fiber formation, and integrin-mediated interactions, such as embryonic development, tumor cell growth and metastasis, programmed cell death, hemostasis, leucocyte homing and activation, bone resorption, clot retraction, and the response of cells to mechanical stress. Thus, the preferred embodiments relate to all aspects of a method of modulating an activity of a Rac polypeptide comprising, administering an effective amount of an active agent, an effective amount of a compound which modulates the activity of a Rac polypeptide, or combination thereof. The activity of Rac which is modulated can include: GTP binding, GDP binding, GEF binding, GTPase activity, integrin binding, coupling or binding of Rac to receptor or effector-like molecules (such as integrins, growth factor receptors, tyrosine kinases, PI-3K, PIP-5K, *etc.*). Increasing, reducing, antagonizing, or promoting Rac can modulate the activity. The modulation of Rac can be measured by assay for GTP hydrolysis, binding to Rac-GEF, *etc.* An effective amount is any amount which, when administered, modulates the Rac activity. The activity can be modulated in a cell, a tissue, a whole organism, *in situ*, *in vitro* (test tube, a solid support, *etc.*), *in vivo*, or in any desired environment.

[0229] The modulation of oncogenic transforming activity by an active agent, or derivatives thereof, can be measured according to various known procedures. A compound can be added at any time during the method (*e.g.*, pretreatment of cells; after addition of GEF, *etc.*) to determine its effect on the oncogenic transforming activity of an active agent. Various cell lines can also be used.

[0230] Other assays for Rac-mediated signal transduction can be accomplished according to procedures known in the art, *e.g.*, as described in U.S. Pat. Nos. 5,141,851;

5,420,334; 5,436,128; and 5,482,954, all of which are incorporated herein by reference in their entirety. In addition, peptides that inhibit the interaction, *e.g.*, binding, between an active agent and a G-protein, such as Rac, can be identified.

[0231] The preferred embodiments also relate to a method of testing for and identifying an agent which modulates the activity of RacGTPase, or a biologically-active fragment thereof, or which modulates the binding between an active agent, or a biologically-active fragment thereof, and a GTPase, or a biologically-active fragment thereof, to which it binds. The method comprises contacting the active agent and Rac GTPase with an agent to be tested and then detecting the presence or amount of binding between the active agent and GTPase, or an activity of the active agent.

[0232] By modulating, it is meant that addition of the agent affects the activity or binding. The binding or activity modulation can be affected in various ways, including inhibiting, blocking, preventing, increasing, enhancing, or promoting it. The binding or activity effect does not have to be achieved in a specific way, *e.g.*, it can be competitive, noncompetitive, allosteric, sterically hindered, via cross-linking between the agent and the GEF or GTPase, *etc.* The agent can act on either the active agent or GTPase. The agent can be an agonist, an antagonist, or a partial agonist or antagonist. The presence or amount of binding can be determined in various ways, *e.g.*, directly or indirectly by assaying for an activity promoted or inhibited by the active agent, such as guanine nucleotide exchange, GTP hydrolysis, oncogenic transformation, *etc.* Such assays are described above and below, and are also known in the art. The agent can be obtained and/or prepared from a variety of sources, including natural and synthetic. It can comprise, *e.g.*, amino acids, lipids, carbohydrates, organic molecules, nucleic acids, inorganic molecules, or mixtures thereof.

[0233] The agent can be added simultaneously or sequentially. For example, the agent can be added to the active agent and then the resultant mixture can be further combined with the GTPase. The method can be carried out in liquid on isolated components, on a matrix (*e.g.*, filter paper, nitrocellulose, agarose), in cells, on tissue sections, *etc.*

[0234] The method further relates to obtaining or producing agents that have been identified according to the above-described method. The preferred embodiments also relate to products identified in accordance with such methods.

[0235] The preferred embodiments thus also relate to the treatment and prevention of diseases and pathological conditions associated with Rac-mediated signal transduction, *e.g.*, cancer, diseases associated with abnormal cell proliferation, and the like. For example, the preferred embodiments relate to a method of treating cancer comprising administering, to a subject in need of treatment, an amount of a compound effective to treat the disease, where the compound is an active agent. Treating the disease can mean, delaying its onset, delaying the progression of the disease, improving or delaying clinical and pathological signs of disease. Similarly, the method also relates to treating diseases associated with inflammation, and/or the chemotactic ability of neutrophils. A regulator compound, or mixture of compounds, can be synthetic, naturally-occurring, or a combination. A regulator compound can comprise amino acids, nucleotides, hydrocarbons, lipids, polysaccharides, *etc.* A regulator compound is preferably a regulator of Rac GTPase. To treat the disease, the compound, or mixture, can be formulated into pharmaceutical composition comprising a pharmaceutically acceptable carrier and other excipients as apparent to the skilled worker. Such composition can additionally contain effective amounts of other compounds, especially for treatment of cancer.

[0236] Based on these data, one embodiment is an improved method for treatment of tumors comprising administration of a pharmaceutically effective quantity of active agent or its pharmaceutically acceptable salts or esters, active agent analogs or their pharmaceutically acceptable salts or esters, or a combination thereof.

[0237] The compositions and preparations described preferably contain at least 0.1% of active agent. The percentage of the compositions and preparations can, of course, be varied, and can contain between about 2% and 60% of the weight of the amount administered. Preferably, the percentage of the compositions and preparations can contain between about 2, 5, 10, or 15% and 30, 35, 40, 45, 50, 55, or 60% of the weight of the amount administered. The amount of active compounds in such pharmaceutically useful compositions and preparations is such that a suitable dosage will be obtained.

[0238] The active agent form salts, which are also within the scope of the preferred embodiments. Reference to a compound of the active agent herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic and/or basic salts formed with inorganic

and/or organic acids and bases. In addition, when an active agent contains both a basic moiety, such as, but not limited to an amine or a pyridine or imidazole ring, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") can be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (*i.e.*, non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, *e.g.*, in isolation or purification steps, which can be employed during preparation. Salts of the compounds of the active agent can be formed, for example, by reacting a compound of the active agent with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

[0239] The active agent which contain a basic moiety, such as, but not limited to an amine or a pyridine or imidazole ring, can form salts with a variety of organic and inorganic acids. Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

[0240] The active agents which contain an acidic moiety, such as, but not limited to a carboxylic acid, can form salts with a variety of organic and inorganic bases. Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, hydrabamines [formed with N,N-bis(dehydroabietyl)ethylenediamine], N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-

containing groups can be quaternized with agents such as lower alkyl halides (*e.g.*, methyl, ethyl, propyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (*e.g.*, dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (*e.g.*, decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (*e.g.*, benzyl and phenethyl bromides), and others.

[0241] Prodrugs and solvates of the compounds of the preferred embodiments are also contemplated herein. The term "prodrug", as employed herein, denotes a compound which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of the active agent, and/or a salt and/or solvate thereof. Solvates of the active agent are preferably hydrates.

[0242] Active agent, and salts thereof, can exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the preferred embodiments.

[0243] All stereoisomers of the present compounds, such as those, for example, which can exist due to asymmetric carbons on any of the substituents, including enantiomeric forms (which can exist even in the absence of asymmetric carbons) and diastereomeric forms, are contemplated and within the scope of the preferred embodiments. Individual stereoisomers of the compounds of the preferred embodiments can, for example, be substantially free of other isomers, or can be admixed, for example, as racemates or with all other or other selected, stereoisomers. The chiral centers of the preferred embodiments can have the S or R configuration as defined by the IUPAC 1974 Recommendations.

[0244] When the compounds according to the preferred embodiments are in the forms of salts, they are preferably pharmaceutically acceptable salts. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable base addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pantoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic,

aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, sulphates, nitrates, phosphates, perchlorates, borates, acetates, benzoates, hydroxynaphthoates, glycerophosphates, ketoglutarates and the like. Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methylammonium, dimethylammonium, trimethylammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like. Examples of organic bases include lysine, arginine, guanidine, diethanolamine, choline and the like.

[0245] The pharmaceutically acceptable salts are prepared by reacting the active agent with 1 to 4 equivalents of a base such as sodium hydroxide, sodium methoxide, sodium hydride, potassium t-butoxide, calcium hydroxide, magnesium hydroxide and the like, in solvents like ether, THF, methanol, t-butanol, dioxane, isopropanol, ethanol, *etc.* Mixture of solvents can be used. Organic bases like lysine, arginine, diethanolamine, choline, guanidine and their derivatives *etc.* can also be used. Alternatively, acid addition salts wherever applicable are prepared by treatment with acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, p-toluenesulphonic acid, methanesulfonic acid, fonic acid, acetic acid, citric acid, maleic acid, salicylic acid, hydroxynaphthoic acid, ascorbic acid, palmitic acid, succinic acid, benzoic acid, benzenesulfonic acid, tartaric acid and the like in solvents like ethyl acetate, ether, alcohols, acetone, THF, dioxane, *etc.* Mixture of solvents can also be used.

[0246] As indicated above, a further object of the preferred embodiments relates to a pharmaceutical composition comprising at least one compound of formula I, II, IIa, III, IIIa, or IV, as defined above, and a pharmaceutically acceptable vehicle or support.

[0247] The compounds can be formulated in various forms, including solid and liquid forms, such as tablets, gel, syrup, powder, aerosol, *etc.*

[0248] The compositions of the preferred embodiments can contain physiologically acceptable diluents, fillers, lubricants, excipients, solvents, binders, stabilizers, and the like. Diluents that can be used in the compositions include but are not limited to dicalcium phosphate, calcium sulphate, lactose, cellulose, kaolin, mannitol, sodium chloride, dry starch, powdered sugar and for prolonged release tablet-hydroxy propyl methyl cellulose (HPMC). The binders that can be used in the compositions

include but are not limited to starch, gelatin and fillers such as sucrose, glucose, dextrose and lactose.

[0249] Natural and synthetic gums that can be used in the compositions include but are not limited to sodium alginate, ghatti gum, carboxymethyl cellulose, methyl cellulose, polyvinyl pyrrolidone and veegum. Excipients that can be used in the compositions include but are not limited to microcrystalline cellulose, calcium sulfate, dicalcium phosphate, starch, magnesium stearate, lactose, and sucrose. Stabilizers that can be used include but are not limited to polysaccharides such as acacia, agar, alginic acid, guar gum and tragacanth, amphotics such as gelatin and synthetic and semi-synthetic polymers such as carbomer resins, cellulose ethers and carboxymethyl chitin.

[0250] Solvents that can be used include but are not limited to Ringers solution, water, distilled water, dimethyl sulfoxide to 50% in water, propylene glycol (neat or in water), phosphate buffered saline, balanced salt solution, glycol and other conventional fluids.

[0251] The dosages and dosage regimen in which the compounds of formula I, II, IIa, III, IIIa, or IV are administered will vary according to the dosage form, mode of administration, the condition being treated and particulars of the patient being treated. Accordingly, optimal therapeutic concentrations will be best determined at the time and place through routine experimentation.

[0252] The compounds according to the preferred embodiments can also be used enterally. Orally, the compounds according to the preferred embodiments are suitable administered at the rate of 100 μ g to 100 mg per day per kg of body weight. Preferably, orally, the compounds according to the preferred embodiments are suitable administered at the rate of about 100, 150, 200, 250, 300, 350, 400, 450, or 500 μ g to about 1, 5, 10, 25, 50, 75, 100 mg per day per kg of body weight. The required dose can be administered in one or more portions. For oral administration, suitable forms are, for example, tablets, gel, aerosols, pills, dragees, syrups, suspensions, emulsions, solutions, powders and granules; a preferred method of administration consists in using a suitable form containing from 1 mg to about 500 mg of active substance. Preferably, a method of administration consists in using a suitable form containing from about 1, 2, 5, 10, 25, or 50 mg to about 100, 200, 300, 400, 500 mg of active substance.

[0253] The compounds according to the preferred embodiments can also be administered parenterally in the form of solutions or suspensions for intravenous or

intramuscular perfusions or injections. In that case, the compounds according to the preferred embodiments are generally administered at the rate of about 10 μ g to 10 mg per day per kg of body weight; a preferred method of administration consists of using solutions or suspensions containing approximately from 0.01 mg to 1 mg of active substance per ml. Preferably, the compounds according to the preferred embodiments are generally administered at the rate of about 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 μ g to 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mg per day per kg of body weight; a preferred method of administration consists of using solutions or suspensions containing approximately from 0.01, 0.02, 0.03, 0.04, or 0.5 mg to 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1 mg of active substance per ml.

[0254] The compounds of formula I, II, IIa, III, IIIa, or IV can be used in a substantially similar manner to other known anti-tumor agents for treating (both chemopreventively and therapeutically) various tumors. For the compounds of the preferred embodiments, the anti-tumor dose to be administered, whether a single dose, multiple dose, or a daily dose, will of course vary with the particular compound employed because of the varying potency of the compound, the chosen route of administration, the size of the recipient, the type of tumor, and the nature of the patient's condition. The dosage to be administered is not subject to definite bounds, but it will usually be an effective amount, or the equivalent on a molar basis of the pharmacologically active free form produced from a dosage formulation upon the metabolic release of the active drug to achieve its desired pharmacological and physiological effects. An oncologist skilled in the art of cancer treatment will be able to ascertain, without undue experimentation, appropriate protocols for the effective administration of the compounds of the preferred embodiments, such as by referring to the earlier published studies on compounds found to have anti-tumor properties.

[0255] The preferred embodiments relate to methods of treatment of disorders involving T-cells include, but are not limited to, cell-mediated hypersensitivity, such as delayed type hypersensitivity and T-cell-mediated cytotoxicity, and transplant rejection; autoimmune diseases, such as systemic lupus erythematosus, Sjogren syndrome, systemic sclerosis, inflammatory myopathies, mixed connective tissue disease, and polyarteritis nodosa and other vasculitides; immunologic deficiency syndromes, including but not limited to, primary immunodeficiencies, such as thymic hypoplasia, severe combined immunodeficiency diseases, and AIDS; leukopenia; reactive (inflammatory) proliferations

of white cells, including but not limited to, leukocytosis, acute nonspecific lymphadenitis, and chronic nonspecific lymphadenitis; neoplastic proliferations of white cells, including but not limited to lymphoid neoplasms, such as precursor T-cell neoplasms, such as acute lymphoblastic leukemia/lymphoma, peripheral T-cell and natural killer cell neoplasms that include peripheral T-cell lymphoma, unspecified, adult T-cell leukemia/lymphoma, mycosis fungoides and Szary syndrome, and Hodgkin disease.

[0256] The compounds of preferred embodiments can be used in relation to diseases of the skin. Diseases of the skin, include but are not limited to, disorders of pigmentation and melanocytes, including but not limited to, vitiligo, freckle, melasma, lentigo, nevocellular nevus, dysplastic nevi, and malignant melanoma; benign epithelial tumors, including but not limited to, seborrheic keratoses, acanthosis nigricans, fibroepithelial polyp, epithelial cyst, keratoacanthoma, and adnexal (appendage) tumors; premalignant and malignant epidermal tumors, including but not limited to, actinic keratosis, squamous cell carcinoma, basal cell carcinoma, and merkel cell carcinoma, tumors of the dermis, including but not limited to, benign fibrous histiocytoma, dermatofibrosarcoma protuberans, xanthomas, and dermal vascular tumors; tumors of cellular immigrants to the skin, including but not limited to, histiocytosis X, mycosis fungoides (cutaneous T-cell lymphoma), and mastocytosis; disorders of epidermal maturation, including but not limited to, ichthyosis; acute inflammatory dermatoses, including but not limited to, urticaria, acute eczematous dermatitis, and erythema multiforme; chronic inflammatory dermatoses, including but not limited to, psoriasis, lichen planus, and lupus erythematosus; blistering (bullous) diseases, including but not limited to, pemphigus, bullous pemphigoid, dermatitis herpetiformis, and noninflammatory blistering diseases: epidermolysis bullosa and porphyria; disorders of epidermal appendages, including but not limited to, acne vulgaris; panniculitis, including but not limited to, erythema nodosum and erythema induratum; and infection and infestation, such as verrucae, molluscum contagiosum, impetigo, superficial fungal infections, and arthropod bites, stings, and infestations.

[0257] The compounds of preferred embodiments can be used in relation to disorders arising from bone marrow cells. In normal bone marrow, the myelocytic series (polymorphonuclear cells) make up approximately 60% of the cellular elements, and the erythrocytic series, 20-30%. Lymphocytes, monocytes, reticular cells, plasma cells and megakaryocytes together constitute 10-20%. Lymphocytes make up 5-15% of normal

adult marrow. In the bone marrow, cell types are add mixed so that precursors of red blood cells (erythroblasts), macrophages (monoblasts), platelets (megakaryocytes), polymorphoneuclear leukocytes (myeloblasts), and lymphocytes (lymphoblasts) can be visible in one microscopic field. In addition, stem cells exist for the different cell lineages, as well as a precursor stem cell for the committed progenitor cells of the different lineages. The various types of cells and stages of each are known to the person of ordinary skill in the art and are found, for example, in *Immunology, Immunopathology and Immunity*, Fifth Edition, Sell *et al.* Simon and Schuster (1996), which is incorporated herein by reference in its entirety. Accordingly, the preferred embodiments are directed to disorders arising from these cells. These disorders include but are not limited to the following: diseases involving hematopoietic stem cells; committed lymphoid progenitor cells; lymphoid cells including B and T-cells; committed myeloid progenitors, including monocytes, granulocytes, and megakaryocytes; and committed erythroid progenitors. These include but are not limited to the leukemias, including B-lymphoid leukemias, T-lymphoid leukemias, undifferentiated leukemias; erythroleukemia, megakaryoblastic leukemia, monocytic; [leukemias are encompassed with and without differentiation]; chronic and acute lymphoblastic leukemia, chronic and acute lymphocytic leukemia, chronic and acute myelogenous leukemia, lymphoma, myelo dysplastic syndrome, chronic and acute myeloid leukemia, myelomonocytic leukemia; chronic and acute myeloblastic leukemia, chronic and acute myelogenous leukemia, chronic and acute promyelocytic leukemia, chronic and acute myelocytic leukemia, hematologic malignancies of monocyte-macrophage lineage, such as juvenile chronic myelogenous leukemia; secondary AML, antecedent hematological disorder; refractory anemia; aplastic anemia; reactive cutaneous angioendotheliomatosis; fibrosing disorders involving altered expression in dendritic cells, disorders including systemic sclerosis, E-M syndrome, epidemic toxic oil syndrome, eosinophilic fasciitis localized forms of scleroderma, keloid, and fibrosing colonopathy; angiomatoid malignant fibrous histiocytoma; carcinoma, including primary head and neck squamous cell carcinoma; sarcoma, including kaposi's sarcoma; fibroadenoma and phyllodes tumors, including mammary fibroadenoma; stromal tumors; phyllodes tumors, including histiocytoma; erythroblastosis; neurofibromatosis; diseases of the vascular endothelium; demyelinating, particularly in old lesions; gliosis, vasogenic edema, vascular disease, Alzheimer's and Parkinson's disease; T-cell lymphomas; B-cell lymphomas.

[0258] The compounds of preferred embodiments can be used in relation to disorders involving the spleen. Disorders involving the spleen include, but are not limited to, splenomegaly, including nonspecific acute splenitis, congestive splenomegaly, and splenic infarcts; neoplasms, congenital anomalies, and rupture. Disorders associated with splenomegaly include infections, such as nonspecific splenitis, infectious mononucleosis, tuberculosis, typhoid fever, brucellosis, cytomegalovirus, syphilis, malaria, histoplasmosis, toxoplasmosis, kala-azar, trypanosomiasis, schistosomiasis, leishmaniasis, and echinococcosis; congestive states related to partial hypertension, such as cirrhosis of the liver, portal or splenic vein thrombosis, and cardiac failure; lymphohematogenous disorders, such as Hodgkin disease, non-Hodgkin lymphomas/leukemia, multiple myeloma, myeloproliferative disorders, hemolytic anemias, and thrombocytopenic purpura; immunologic-inflammatory conditions, such as rheumatoid arthritis and systemic lupus erythematosus; storage diseases such as Gaucher disease, Niemann-Pick disease, and mucopolysaccharidoses; and other conditions, such as amyloidosis, primary neoplasms and cysts, and secondary neoplasms.

[0259] The compounds of preferred embodiments can be used in relation to disorders involving blood vessels. Disorders involving blood vessels include, but are not limited to, responses of vascular cell walls to injury, such as endothelial dysfunction and endothelial activation and intimal thickening; vascular diseases including, but not limited to, congenital anomalies, such as arteriovenous fistula, atherosclerosis, and hypertensive vascular disease, such as hypertension; inflammatory disease--the vasculitides, such as giant cell (temporal) arteritis, Takayasu arteritis, polyarteritis nodosa (classic), Kawasaki syndrome (mucocutaneous lymph node syndrome), microscopic polyangiitis (microscopic polyarteritis, hypersensitivity or leukocytoclastic angitis), Wegener granulomatosis, thromboangiitis obliterans (Buerger disease), vasculitis associated with other disorders, and infectious arteritis; Raynaud disease; aneurysms and dissection, such as abdominal aortic aneurysms, syphilitic (luetic) aneurysms, and aortic dissection (dissecting hematoma); disorders of veins and lymphatics, such as varicose veins, thrombophlebitis and phlebothrombosis, obstruction of superior vena cava (superior vena cava syndrome), obstruction of inferior vena cava (inferior vena cava syndrome), and lymphangitis and lymphedema; tumors, including benign tumors and tumor-like conditions, such as hemangioma, lymphangioma, glomus tumor (glomangioma), vascular ectasias, and bacillary angiomatosis, and intermediate-grade (borderline low-grade malignant) tumors,

such as Kaposi sarcoma and hemangioendothelioma, and malignant tumors, such as angiosarcoma and hemangiopericytoma; and pathology of therapeutic interventions in vascular disease, such as balloon angioplasty and related techniques and vascular replacement, such as coronary artery bypass graft surgery.

[0260] The compounds of preferred embodiments can be used in relation to disorders involving red cells. Disorders involving red cells include, but are not limited to, anemias, such as hemolytic anemias, including hereditary spherocytosis, hemolytic disease due to erythrocyte enzyme defects: glucose-6-phosphate dehydrogenase deficiency, sickle cell disease, thalassemia syndromes, paroxysmal nocturnal hemoglobinuria, immunohemolytic anemia, and hemolytic anemia resulting from trauma to red cells; and anemias of diminished erythropoiesis, including megaloblastic anemias, such as anemias of vitamin B12 deficiency: pernicious anemia, and anemia of folate deficiency, iron deficiency anemia, anemia of chronic disease, aplastic anemia, pure red cell aplasia, and other forms of marrow failure.

[0261] The compounds of preferred embodiments can be used in relation to disorders involving B-cells. Disorders involving B-cells include, but are not limited to precursor B-cell neoplasms, such as lymphoblastic leukemia/lymphoma. Peripheral B-cell neoplasms include, but are not limited to, chronic lymphocytic leukemia/small lymphocytic lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, Burkitt lymphoma, plasma cell neoplasms, multiple myeloma, and related entities, lymphoplasmacytic lymphoma (Waldenstrom macroglobulinemia), mantle cell lymphoma, marginal zone lymphoma (MALToma), and hairy cell leukemia.

[0262] The compounds of preferred embodiments can be used in relation to disorders related to reduced platelet number. Disorders related to reduced platelet number, thrombocytopenia, include idiopathic thrombocytopenic purpura, including acute idiopathic thrombocytopenic purpura, drug-induced thrombocytopenia, HIV-associated thrombocytopenia, and thrombotic microangiopathies: thrombotic thrombocytopenic purpura and hemolytic-uremic syndrome.

[0263] The compounds of preferred embodiments can be used in relation to disorders involving precursor T-cell neoplasms. Disorders involving precursor T-cell neoplasms include precursor T lymphoblastic leukemia/lymphoma. Disorders involving peripheral T-cell and natural killer cell neoplasms include T-cell chronic lymphocytic leukemia, large granular lymphocytic leukemia, mycosis fungoides and Szary syndrome,

peripheral T-cell lymphoma, unspecified, angioimmunoblastic T-cell lymphoma, angiocentric lymphoma (NK/T-cell lymphoma4a), intestinal T-cell lymphoma, adult T-cell leukemia/lymphoma, and anaplastic large cell lymphoma.

[0264] The compounds of preferred embodiments can be used in relation to disorders of the bone. Bone-forming cells include the osteoprogenitor cells, osteoblasts, and osteocytes. The disorders of the bone are complex because they can have an impact on the skeleton during any of its stages of development. Hence, the disorders can have variable manifestations and can involve one, multiple or all bones of the body. Such disorders include, congenital malformations, achondroplasia and thanatophoric dwarfism, diseases associated with abnormal matrix such as type 1 collagen disease, osteoporosis, Paget disease, rickets, osteomalacia, high-turnover osteodystrophy, low-turnover of aplastic disease, osteonecrosis, pyogenic osteomyelitis, tuberculous osteomyelitis, osteoma, osteoid osteoma, osteoblastoma, osteosarcoma, osteochondroma, chondromas, chondroblastoma, chondromyxoid fibroma, chondrosarcoma, fibrous cortical defects, fibrous dysplasia, fibrosarcoma, malignant fibrous histiocytoma, Ewing sarcoma, primitive neuroectodermal tumor, giant cell tumor, and metastatic tumors.

[0265] The compounds of preferred embodiments can be used in relation to disorders involving the tonsils. Disorders involving the tonsils include, but are not limited to, tonsillitis, Peritonsillar abscess, squamous cell carcinoma, dyspnea, hyperplasia, follicular hyperplasia, reactive lymphoid hyperplasia, non-Hodgkin's lymphoma and B-cell lymphoma.

[0266] The compounds of preferred embodiments can be used in relation to disorders involving the liver. Disorders involving the liver include, but are not limited to, hepatic injury; jaundice and cholestasis, such as bilirubin and bile formation; hepatic failure and cirrhosis, such as cirrhosis, portal hypertension, including ascites, portosystemic shunts, and splenomegaly; infectious disorders, such as viral hepatitis, including hepatitis A-E infection and infection by other hepatitis viruses, clinicopathologic syndromes, such as the carrier state, asymptomatic infection, acute viral hepatitis, chronic viral hepatitis, and fulminant hepatitis; autoimmune hepatitis, drug- and toxin-induced liver disease, such as alcoholic liver disease; inborn errors of metabolism and pediatric liver disease, such as hemochromatosis, Wilson disease, alpha.1-antitrypsin deficiency, and neonatal hepatitis; intrahepatic biliary tract disease, such as secondary biliary cirrhosis, primary biliary cirrhosis, primary sclerosing cholangitis, and anomalies

of the biliary tree; circulatory disorders, such as impaired blood flow into the liver, including hepatic artery compromise and portal vein obstruction and thrombosis, impaired blood flow through the liver, including passive congestion and centrilobular necrosis and peliosis hepatis, hepatic vein outflow obstruction, including hepatic vein thrombosis (Budd-Chiari syndrome) and veno-occlusive disease; hepatic disease associated with pregnancy, such as preeclampsia and eclampsia, acute fatty liver of pregnancy, and intrahepatic cholestasis of pregnancy; hepatic complications of organ or bone marrow transplantation, such as drug toxicity after bone marrow transplantation, graft-versus-host disease and liver rejection, and nonimmunologic damage to liver allografts; tumors and tumorous conditions, such as nodular hyperplasias, adenomas, and malignant tumors, including primary carcinoma of the liver and metastatic tumors.

[0267] The compounds of preferred embodiments can be used in relation to disorders involving the colon. Disorders involving the colon include, but are not limited to, congenital anomalies, such as atresia and stenosis, Meckel diverticulum, congenital aganglionic megacolon-Hirschsprung disease; enterocolitis, such as diarrhea and dysentery, infectious enterocolitis, including viral gastroenteritis, bacterial enterocolitis, necrotizing enterocolitis, antibiotic-associated colitis (pseudomembranous colitis), and collagenous and lymphocytic colitis, miscellaneous intestinal inflammatory disorders, including parasites and protozoa, acquired immunodeficiency syndrome, transplantation, drug-induced intestinal injury, radiation enterocolitis, neutropenic colitis (typhlitis), and diversion colitis; idiopathic inflammatory bowel disease, such as Crohn disease and ulcerative colitis; tumors of the colon, such as non-neoplastic polyps, adenomas, familial syndromes, colorectal carcinogenesis, colorectal carcinoma, and carcinoid tumors.

[0268] The compounds of preferred embodiments can be used in relation to disorders involving the lung. Disorders involving the lung include, but are not limited to, congenital anomalies; atelectasis; diseases of vascular origin, such as pulmonary congestion and edema, including hemodynamic pulmonary edema and edema caused by microvascular injury, adult respiratory distress syndrome (diffuse alveolar damage), pulmonary embolism, hemorrhage, and infarction, and pulmonary hypertension and vascular sclerosis; chronic obstructive pulmonary disease, such as emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis; diffuse interstitial (infiltrative, restrictive) diseases, such as pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary

eosinophilia (pulmonary infiltration with eosinophilia), Bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, including Goodpasture syndrome, idiopathic pulmonary hemosiderosis and other hemorrhagic syndromes, pulmonary involvement in collagen vascular disorders, and pulmonary alveolar proteinosis; complications of therapies, such as drug-induced lung disease, radiation-induced lung disease, and lung transplantation; tumors, such as bronchogenic carcinoma, including paraneoplastic syndromes, bronchioloalveolar carcinoma, neuroendocrine tumors, such as bronchial carcinoid, miscellaneous tumors, and metastatic tumors; pathologies of the pleura, including inflammatory pleural effusions, noninflammatory pleural effusions, pneumothorax, and pleural tumors, including solitary fibrous tumors (pleural fibroma) and malignant mesothelioma.

[0269] The active compounds can be incorporated into pharmaceutical compositions suitable for administration to a subject, *e.g.*, a human. Such compositions typically comprise the nucleic acid molecule, protein, modulator, or antibody and a pharmaceutically acceptable carrier.

[0270] As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, such media can be used in the compositions of the preferred embodiments. Supplementary active compounds can also be incorporated into the compositions. A pharmaceutical composition of the preferred embodiments is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid, buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted

with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0271] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art.

[0272] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated, each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the preferred embodiments are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[0273] As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, *e.g.*, healing of chronic conditions or in an increase in rate of healing of such conditions, or in a reduction in aberrant conditions. This includes both therapeutic and prophylactic treatments. Accordingly, the compounds can be used at very early stages of a disease, or before early onset, or after significant progression. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

[0274] In practicing the method of treatment or use of the preferred embodiments, a therapeutically effective amount of one, two, or more of the active agents of the preferred embodiments is administered to a subject afflicted with a disease or disorder related to Rho family GTPases, or to a tissue which has such disease or disorder. The active agents of the preferred embodiments can be administered in accordance with the method of the preferred embodiments either alone or in combination with other known therapies. When co-administered with one or more other therapies, the active agents of the preferred embodiments can be administered either simultaneously with the other treatment(s), or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering the active agents of the preferred embodiments in combination with the other therapy.

[0275] Generally, a therapeutically effective amount of active agent (*i.e.*, an effective dosage) ranges from about 0.001 to 5000 mg/kg body weight, more preferably about 0.01 to 1000 mg/kg body weight, more preferably about 0.01 to 500 mg/kg body weight, more preferably about 0.01 to 250 mg/kg body weight, more preferably about 0.01 to 100 mg/kg body weight, more preferably about 0.001 to 60 mg/kg body weight, more preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight.

[0276] The skilled artisan will appreciate that certain factors can influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage used for treatment can increase or decrease over the course of a particular treatment. Changes in dosage can result and become apparent from the results of diagnostic assays as described herein.

[0277] The preferred embodiments encompass one or more additional agents that modulate expression or activity of Rac GTPase. An agent can, for example, be a

small molecule. For example, such small molecules include, but are not limited to, peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (*i.e.*, including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

[0278] In one embodiment, the additional agent can be a prenylation inhibitor, such as disclosed by U.S. Pat. Nos. 6,649,638, 5,420,245; 5,574,025; 5,523,430; 5,602,098; 5,631,401; 5,705,686; 5,238,922; 5,470,832; and 6,191,147, all of which are incorporated herein by reference in their entirety.

[0279] In another embodiment, the additional agent comprises one or more inhibitor of farnesyl protein transferase (FPTase), prenyl-protein transferase or geranylgeranyl-protein transferase as described in U.S. Pat. Nos. 6,572,850; 6,458,783; 6,423,751; 6,387,926; 6,242,433; 6,191,147; 6,166,067; 6,156,746; 6,083,979; 6,011,029; 5,929,077; 5,928,924; 5,843,941; 5,786,193; 5,629,302; 5,618,964; 5,574,025; 5,567,841; 5,523,430; 5,510,510; 5,470,832; 5,447,922; 6,596,735; 6,586,461; 6,586,447; 6,579,887; 6,576,639; 6,545,020; 6,539,309; 6,535,820; 6,528,523; 6,511,800; 6,500,841; 6,495,564; 6,492,381; 6,458,935; 6,451,812; 6,441,017; 6,440,989; 6,440,974; 6,432,959; 6,426,352; 6,410,541; 6,403,581; 6,399,615; 6,387,948; 6,387,905; 6,387,903; 6,376,496; 6,372,747; 6,362,188; 6,358,968; 6,329,376; 6,316,462; 6,294,552; 6,277,854; 6,268,394; 6,265,382; 6,262,110; 6,258,824; 6,248,756; 6,242,458; 6,239,140; 6,228,865; 6,228,856; 6,225,322; 6,218,401; 6,214,828; 6,214,827; 6,211,193; 6,194,438, which are specifically incorporated herein by reference in their entirety.

[0280] A “farnesyl protein transferase inhibitor” or “FPT inhibitor” or “FTI” is defined herein as a compound which: (i) potently inhibits FPT (but generally not geranylgeranyl protein transferase I) and (ii) blocks intracellular farnesylation of ras. FPT catalyzes the addition of an isoprenyl lipid moiety onto a cysteine residue present near the carboxy-terminus of the Ras protein. This is the first step in a post-translational processing pathway that is essential for both Ras membrane-association and Ras-induced

oncogenic transformation. A number of FPT inhibitors have been reported, including a variety of peptidomimetic inhibitors as well as other small molecule inhibitors.

[0281] Farnesyl transferase inhibitors generally fall into two classes: analogs of farnesyl diphosphate; and protein substrates for farnesyl transferase. Farnesyl transferase inhibitors have been described in U.S. Pat. No. 5,756,528, U.S. Pat. No. 5,141,851, U.S. Pat. No. 5,817,678, U.S. Pat. No. 5,830,868, U.S. Pat. No. 5,834,434, and U.S. Pat. No. 5,773,455, all of which are incorporated herein by reference in their entirety. Among the farnesyl transferase inhibitors shown to be effective for inhibiting the transfer of the farnesyl moiety to Ras-related proteins are L-739,749 (a peptidomimetic analog of the C-A-A-X sequence), L-744,832 (a peptidomimetic analog of the C-A-A-X sequence), SCH 44342 (1-(4-pyridylacetyl)-4-(8-chloro-5,6 dihydro-IIH benzo [5,6] cyclohepta [1,2-b]pyridin-11-ylidene)piperidine), BZA-5B (a benzodiazepine peptidomimetic), FTI-276 (a C-A-A-X peptidomimetic), and B1086 (a C-A-A-X peptidomimetic). Administration of farnesyl transferase inhibitors (FTIs) is accomplished by standard methods known to those of skill in the art, most preferably by administration of tablets containing the FTI, and is expected to fall approximately within a range of about 0.1 mg/kg of body weight to about 20 mg/kg of body weight per day.

[0282] In another embodiment, the additional agent comprises one or more inhibitor of geranylgeranyl-protein transferase (GGT) as have been described in U.S. Pat. No. 5,470,832 (Gibbs & Graham), which is incorporated herein by reference in its entirety. These compounds can be administered to an individual in dosage amounts of between 0.5 mg/kg of body weight to about 20 mg/kg of body weight. Alternatively, one or more inhibitors of isoprenylation, including farnesyl transferase (FT) inhibitors and/or geranylgeranyl transferase inhibitors (GGT) are administered to a patient.

[0283] In another embodiment, the additional agent comprises one or more toxins such as toxins A and B from *C. difficile* and *C. sordellii* lethal toxin (LT). In addition, Rac 1 and Rac2 can be inhibited when Rho is specifically ADP ribosylated by C3 enzyme, which is one of the botulinum toxins, and Staphylococcal toxin EDIN (Narumiya, S. and Morii, S., *Cell Signal*, 5, 9-19, 1993; Sekine, A. et al., *J. Biol. Chem.*, 264, 8602-8605, 1989, all of which are incorporated herein by reference in their entirety).

[0284] It is understood that appropriate doses of small molecule agents depends upon a number of factors within the ken of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of the small molecule will vary, for example,

depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the small molecule to have upon the nucleic acid or polypeptide of the preferred embodiments. Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (*e.g.*, about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. Such appropriate doses can be determined using the assays described herein. When one or more of these small molecules is to be administered to an animal (*e.g.*, a human) in order to modulate expression or activity of a polypeptide or nucleic acid of the preferred embodiments, a physician, veterinarian, or researcher can, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

[0285] The examples disclosed below illustrated preferred embodiments and are not intended to limit the scope. It is evident to those skilled in the art that modifications or variations can be made to the preferred embodiments described herein without departing from the teachings of the present invention.

EXAMPLES

[0286] Recombinant protein production. Recombinant Trio (residues 1225-1537) containing the N-terminal DH/PH module, Rac1, Cdc42 and the p21-binding domain (PBD) of PAK1 (residues 51-135) are expressed in *E. coli* BL21 (DE3) strain as N-terminal His₆-tagged fusion proteins by using the pET expression system (Novagen). Rac1, Cdc42, Intersectin, PAK1 (PBD) and WASP (PBD) are expressed in *E. coli* DH5α strain as GST fusions by using the pGEX-KG vector. The N-terminal tagged GST or His₆ fusion proteins are purified by glutathione- or Ni²⁺-agarose affinity chromatography.

GST-Rho GTPases on glutathione beads are eluted off bound guanine nucleotides or Mg^{2+} by washing with a buffer containing 50 mM Tris-HCl, pH 7.6, 100 mM NaCl, 1 mM EDTA, and 1 mM DTT.

[0287] In vitro complex formation assay. About 0.5 μ g of His₆-tagged Trio is incubated with 0.5 μ g, EDTA-treated, GST-fused Cdc42 or Rac1 in a binding buffer containing 20 mM Tris-HCl, pH 7.6, 100 mM NaCl, 1mM DTT, 1% bovine serum albumin, 1% Triton X-100, 1mM $MgCl_2$ and 10 μ l suspended glutathione-agarose beads. ~0.75 μ g of GST-tagged Intersectin is incubated with nucleotide-free, His₆-tagged Cdc42 or Rac1 (0.25 μ g) in the binding buffer with 10 μ l suspended glutathione-agarose beads. After incubation at 4°C for 30 min under constant agitation, the glutathione beads are washed twice with the binding buffer. The amount of His₆-tagged protein co-precipitated with the GST-fusion bound beads is detected by anti-His Western blotting.

[0288] In vitro guanine nucleotide exchange assay. For these, 200 nM Rac1 loaded with mant-GDP is incubated at 25°C in an exchange buffer containing 100 mM NaCl, 5 mM $MgCl_2$, 50 mM Tris-HCl (pH 7.6), and 0.5 mM GTP in the absence or presence of 200 nM Trio. The mant-GDP fluorescence changes in the course of the exchange reactions are monitored with an excitation wavelength at 360 nm and the emission wavelength at 440 nm by a Cary Ellipse fluorescence spectrometer (Varian, Inc.).

[0289] Cell culture. NIH 3T3 fibroblasts are grown in Dulbecco's modified Dagle's medium supplemented with 10% calf serum. RWPE-1 cells are obtained from the American Type culture collection (ATCC) and are grown in keratinocyte-Serum Free medium (GIBCO-BRL) supplemented with 5 ng/ml EGF and 0.05 mg/ml bovine pituitary extract. PC-3 cells are cultured in RPM1 1640 medium (Cellgro) supplemented with 10% FBS.

[0290] Endogenous Rho GTPase activity assay. GST- or His₆-PAK1 (PBD) and GST-WASP (PBD) are expressed in Escherichia coli and purified by glutathione- or Ni^{2+} -agarose affinity chromatography. Cells are grown in log phase in a 10 cm dish, and are starved in 0.5% serum medium or indicated otherwise for 24 hrs prior to lysis in a buffer containing 20 mM Tris-HCl (pH 7.6), 100 mM NaCl, 10 mM $MgCl_2$, 1% NP-40, 10% glycerol, and 1× proteases inhibitor cocktail (Roch). Lysates are clarified and the protein concentrations are normalized. The cell lysates containing equal amount of proteins are incubated with 10 μ g GST- or His₆-fusion probes for 40 min at 4°C under

constant rotation. The beads are washed twice with the lysis buffer, and the bound-Rho GTPases are detected by anti-Rac1 (Upstate), or anti-Cdc42 (BD Transduction Laboratories) Western blotting. Quantification of the Western blots is carried out using a LAS-1000 luminescent image analyzer (Fujifilm medical system, USA, Inc.).

[0291] Immunofluorescence. After overnight serum starvation in the presence or absence of 100 μ M 23766, NIH 3T3 cells grown on cover glasses are treated with 10 nM PDGF for zero, five or ten minutes. The cells are fixed with 3.7% formaldehyde in PBS for 15 min, and permeabilized with 0.1% Trion X-100 for 20 min. The cellular actin is stained with TRITC-labeled phalloidin (Sigma) at 10 μ g/ml in PBS for 40 min at room temperature. The actin and cell morphological changes are visualized by fluorescence microscopy.

[0292] Cell growth assay. Wild type and RacL61- or various GEF-transfected NIH 3T3 cells are grown in 5 % calf serum. The cells are split in duplicate in 6-well plates at 5×10^4 cells per well and are counted daily with a hemocytometer for 4 days. The growth rate of the prostate PC-3 cells is measured by the CellTiter 96 AQueous assay (Promega). 1,500 cells/well in 200 μ l of 5 % FBS medium are plated in 96-well plates and are grown under normal conditions. Cultures are assayed in 0, 1, 2, 3, 4, or 5 days by the addition of 20 μ l of the combined MTS/PMS solution followed by incubation for one hour at 37 °C. Absorbency is measured at a wavelength of 490 nm on an automated microplate reader.

[0293] Anchorage independent growth. The prostate epithelia RWPE and PC-3 cells (1.25×10^3 per well) are grown in 0.3 % agarose in the absence or presence of different doses of compound 23766 following a published protocol (Qiu et al., 1997, which is incorporated herein by reference in its entirety). The number of colonies formed in soft agar is counted after ten days.


[0294] Cell invasion assays. The cell invasion assays are performed using 6.4-mm Biocoat Matrigel invasion chambers with 8.0-micron pore size PET membrane (Becton-Dickinson) according to the manufactory instructions. Briefly, 5×10^4 cells are resuspended in 0.5 ml of serum free culture medium and added to the upper chamber. 10 % fetal bovine serum in the culture medium is used as a chemo-attractant in the lower chamber. After the cells are incubated for overnight, the number of cell passed through the Matrigel is counted.

Results

[0295] Virtual Screening for Rac1-specific inhibitors. In the three-dimensional (3D) structure of Rac1-Tiam1 complex, Trp⁵⁶ of Rac1 is buried in a pocket formed by residues His¹¹⁷⁸, Ser1¹⁸⁴, Glu¹¹⁸³, and Ile¹¹⁹⁷ of Tiam1 and Lys⁵, Val⁷, Thr⁵⁸, and Ser⁷¹ of Rac1 (Worthylake et al., 2000, which is incorporated herein by reference in its entirety). To identify Rac1-specific inhibitors based on the structural features surrounding Trp⁵⁶, a potential inhibitor-binding pocket is created with residues of Rac1 within 6.5 angstroms of Trp⁵⁶ in the Rac1-Tiam1 monomer, including Lys⁵, Val⁷, Trp⁵⁶, and Ser⁷¹. A 3D database search is performed to identify compounds whose conformations would fit the binding pocket. In order to take the flexibility of the compounds into consideration during the screening process, the program UNITY, whose Directed Tweak algorithm allows a rapid, conformationally flexible 3D search (Hurst, 1994, which is incorporated herein by reference in its entirety), is applied.


[0296] The small molecule hits yielded by the UNITY program are next docked into the predicted binding pocket of Rac1 containing Trp⁵⁶ by using the program FlexX, an energy minimization modeling software that can quickly and flexibly dock ligand to protein binding site (Rarey et al., 1996, which is incorporated herein by reference in its entirety). Following the docking procedures, the compounds are ranked based on their predicted ability to bind the binding pocket using the program Cscore. Cscore generates a relative, consensus score based on how well the individual scoring functions of the protein-ligand complex perform (Clark et al., 2002, which is incorporated herein by reference in its entirety).

[0297] Compound NCI 23766 specifically inhibits Rac1-GEF interaction. Compounds from the virtual screening, including Compound NCI 23766, were obtained from the National Cancer Institute-Research Samples and Services from Developmental Therapeutics Program (Bethesda, Maryland). Also, Compound NCI 23766 can be synthesized as set forth herein.

Synthetic Scheme 1

[0298] Synthetic Scheme 1 follows closely the reaction conditions of Synthetic Scheme 2. In Synthetic Scheme 2, NHR_1R_2 is compound 10. NHR_1R_2 can be varied to be included within the preferred embodiments. NHR_1R_2 can be commercially available or synthesized using standard chemical methodologies. The reaction between NHR_1R_2 and Compound 9 or 9a is a standard amination reaction onto a haloaromatic ring.

[0299] The synthetic scheme described herein can be carried out using standard chemical methodologies described and referenced in standard textbooks. One may substitute other reagents known in the art which are known to be equivalent or perform a similar function. Starting material are commercially available reagents and reactions are preferably carried out in standard laboratory glassware under reaction conditions of standard temperature and pressure, except where otherwise indicated.

Synthetic Scheme 2EXPERIMENTALGeneral:

[0300] Raw materials were purchased from Aldrich, Acros, Fisher or Matrix Scientific. All solvents were ACS grade or better. Reactions were run under an atmosphere of dry nitrogen as necessary. Removal of solvents “in vacuo” refers to rotary evaporation using a Buchi apparatus at 25-50 °C and 45 Torr. Vacuum drying was done under high vacuum. All NMR spectra were recorded using a Varian-Gemini 300 spectrometer at 300 MHz for ^1H NMR using CHCl_3 (7.26 ppm) or DMSO (2.5 ppm) as a reference and at 75 MHz for ^{13}C NMR using CDCl_3 (77.0 ppm) or DMSO (39.43) as a reference.

[0301] Methyl 3-[[4-(acetylamino)phenyl]amino]but-2-enoate (3): A suspension of 4-aminoacetanilide (**2**) (253 g, 1.68 mol) and methyl acetoacetate (215 g, 1.85 mol) in MeOH (0.75 L) was heated to reflux. The resulting solution was held at reflux for 16 h and then cooled to 5 °C. The resulting off-white precipitate was filtered and washed with MTBE (3 x 200 mL) to give butenoate **3** (195 g, 47% yield). The mother liquor was concentrated in vacuo and filtered to give a second crop of **3** as pale pink solids (141 g, 34% yield, 81% overall yield). ^1H NMR (DMSO) δ 10.22 (s, 1 H), 9.97 (s, 1 H), 7.57 (d, 2 H), 7.11 (d, 2 H), 4.65 (s, 1 H), 3.56 (s, 3 H), 2.04 (s, 3 H), 1.94 (s, 3 H); ^{13}C NMR (DMSO) δ 169.64, 168.07, 159.33, 136.48, 133.55, 124.52, 119.44, 84.47, 49.77, 23.81, 19.68.

[0302] ***N*-(4-Hydroxy-2-methylquinolin-6-yl)acetamide (4):** Phenyl ether (1 L) was heated to 255 °C. Butenoate 3 (334 g, 1.35 mol) was carefully added portionwise while maintaining temperature 245-260 °C. After the addition was complete, the yellow-orange suspension was held at 255 °C for an additional 15 min. The mixture was slowly cooled to 40 °C, the solids were collected by filtration and washed with EtOAc (3 x 500 mL) followed by MeOH (3 x 500 mL) to give hydroxy quinoline 4 as yellow-orange solid (256 g, 88% yield); ¹H NMR (DMSO) δ 11.52 (br, s, 1 H), 10.07 (s, 1 H), 8.24 (s, 1 H), 7.82 (d, 1 H), 7.42 (d, 1 H), 5.84 (s, 1 H), 2.31 (s, 3 H), 2.05 (s, 1 H).

[0303] ***N*-(4-Methoxy-2-methylquinolin-6-yl)acetamide (5):** Dimethyl sulfate (294 g, 2.33 mol) was charged to a suspension of hydroxyquinoline 4 (287 g, 1.33 mol) in toluene (1.5 L) and the mixture was refluxed for 6 h. After cooling to ambient temperature, the resulting dark yellow solids were collected by filtration and washed with toluene. The dry solid was dissolved in water (2.5 L) and the pH adjusted to 14 using 35% aqueous NaOH (290 g). The resulting tan precipitate was collected by filtration, washed with copious amounts of water and dried in vacuo at 60 °C to give methoxyquinoline 5 as light tan solid (259 g, 85% yield); ¹H NMR (DMSO) δ 10.18 (s, 1 H), 8.46 (s, 1 H), 7.76 (m, 2 H), 6.84 (s, 1 H), 3.99 (s, 3 H), 2.55 (s, 3 H), 2.09 (s, 3 H); ¹³C NMR (DMSO) δ 168.36, 160.99, 158.07, 144.86, 135.87, 128.27, 122.78, 119.26, 108.80, 101.20, 55.69, 25.08, 23.97.

[0304] **2-Methylquinoline-4,6-diamine (7):** Ammonium acetate (1.3 kg) was melted and methoxyquinoline 5 (256 g, 1.11 mol) was added. The dark solution was refluxed at 135 for 4 h. After LC/MS indicated conversion of 5 (M+1 = 231) to intermediate 6 (M+1 = 216), the reaction mixture was poured into 37% HCl (2.1 L) and water (800 mL). The mixture was refluxed for 10 h and then cooled to ambient temperature for overnight. LC/MS indicated conversion of all intermediate 6 to diaminoquinoline 7 (M+1 = 174). The mixture was cooled to 5 °C and the resulting dihydrochloride salt was collected by filtration. The salt was dissolved in water (1.5 L) at 75 °C. Charcoal (13 g, Darco G-60, -100 mesh) was charged to the dark solution, the mixture was refluxed for 45 minutes and was filtered through Celite. The yellow filtrate was cooled and the pH adjusted to 14 using 35% aqueous NaOH (1 kg). The resulting precipitate was collected by filtration, washed with copious amounts of water and dried in vacuo at 60 °C to give diaminoquinoline 7 as off-white solid (136 g, 71% yield); ¹H NMR

(DMSO) δ 7.41 (m, 1 H), 6.95 (m, 2 H), 6.30 (s, 1 H), 6.03 (br, s, 2 H), 5.05 (br, s, 2 H), 2.32 (s, 3 H); ^{13}C NMR (DMSO) δ 153.42, 149.46, 144.28, 142.09, 128.84, 120.59, 118.60, 102.15, 101.11, 24.42.

[0305] N~6~-(2-chloro-6-methylpyrimidin-4-yl)-2-methylquinoline-4,6-diamine (9): Diaminoquinoline 7 (72.0 g, 0.416 mol) and 2,4-dichloro-6-methylpyrimidine (8) (67.8 g, 0.416 mol) were suspended in ethylene glycol (1 L). Addition of 37% HCl (35 mL, 0.43 mol) resulted in a yellow solution which was heated to and held at 50 °C for 4.5 h. The mixture was diluted with chilled water (1L) which resulted in a thick white paste-like precipitate and the mixture was filtered through Celite. The Celite and solid containing the product and bis-substituted by-product was slurried in water (4 L) and the Celite and insoluble by-product were removed by filtration. The filtrate pH was adjusted to 14 using 1N aqueous NaOH (1 L) resulting in precipitation of product which was removed by filtration. The damp product was transferred to a rotovap flask and dried in vacuo by azeotropic water removal with toluene (3 x 1.5 L). Product 9 was obtained as an off-white solid (33.4 g, 27% yield; Notebook reference A134-137). Another batch of 9 (7.4 g, 17% yield; Notebook reference A134-134) was similarly obtained by reaction of 7 (25.0 g, 0.144 mol) and recovery as above; MS $[M+1] = 300, 302$; ^{13}C NMR (DMSO) δ 167.51, 162.58, 159.22, 157.61, 151.05, 145.99, 133.17, 128.96, 125.39, 117.38, 114.23, 102.45, 102.26, 24.68, 23.19.

[0306] CHMC-1: A suspension of intermediate 9 (32.7 g, 0.109 mol) and diisopropylethylamine (20.0 mL, 0.115 mol) in ethylene glycol (500 mL) was heated to 90 °C to give a golden solution. 2-Amino-5-diethylaminopentane (32.0 mL, 0.165 mol) was added and the mixture was heated to and held at 110 °C for 5.5 h. The mixture was cooled to room temperature and EtOAc (750 mL) and 1N aqueous NaOH (500 mL) were added resulting in a thick white paste-like precipitate. The solid was removed by filtration through Celite and the filtrate layers were separated. The aqueous layer was twice basified using 1N NaOH (300 mL) and back-extracted using EtOAc (750 mL). The combined organic layers were washed with brine (3 x 750 mL), filtered through Celite and solvent removed in vacuo to give a brown oil (44 g). Heptane was added to the oil and allowed to sit for several days before decanting the solvent. The oil along with 12 g of crude oil from another batch were purified by silica gel (1.5 kg) flash column chromatography using EtOAc/MeOH/ NEt_3 (7:3:0.5).

[0307] Chemical compounds are examined for their ability to inhibit the Rac1 binding interaction with GEF in a complex formation assay.

[0308] Fig. 1. shows identification of NSC23766 as an inhibitor of Rac1-Trio interaction. In the upper panel of Fig. 1, the inhibitory effect of a panel of compounds predicted by Virtual Screening on Rac1 interaction with TrioN was tested in a complex formation assay. 0.5 µg of (His)₆-tagged TrioN was incubated with GST alone or nucleotide-free GST-Rac1 (2 µg) in the presence or absence of 1 mM indicated NCI compound and 10 µl suspended glutathione-agarose beads. After an incubation at 4°C for minutes, the beads associated (His)₆-TrioN were detected by anti-His Western blotting. In the lower panel of Fig. 1, the effect of the compounds on Cdc42 binding to Intersectin was determined similarly. ~1 µg of GST or GST-tagged Intersectin was incubated with the nucleotide-free, (His)₆-tagged Cdc42 (0.25 µg) under similar conditions. Data are representative of the results from four independent experiments.

[0309] For this purpose, Trio and Tiam-1, which specifically activate Rac1 but not Cdc42 (Gao et al., 2001, which is incorporated herein by reference in its entirety) and Intersectin, a Cdc42-specific GEF (Karnoub et al., 2001, which is incorporated herein by reference in its entirety), are used to assay the binding activity to their respective substrates in the presence of 1mM of each individual compound. Trio and Tiam-1 co-precipitate with GST-Rac1, but not GST or GST-Cdc42. The inhibitory effect of compound 23766 appears to be specific towards the interaction between Rac1 and its GEFs since it does not interfere with the Cdc42 binding to Intersectin nor RhoA binding to PDZ-RhoGEF (Fig. 1). Further, the inhibitory effect of compound 23766 on Rac1 is dose dependent (Fig. 2).

[0310] Fig. 2. shows dose dependent specific inhibition of GEF interaction with Rac1 by NSC23766. In Fig 2A, 0.5 µg of (His)₆-tagged TrioN was incubated with GST alone or nucleotide-free, GST-fused Cdc42 or Rac1 (2 µg) in the binding buffer containing different concentrations of NSC23766 and 10 µl suspended glutathione-agarose. After an incubation at 4 °C for 30 minutes, the beads associated (His)₆-TrioN were detected by anti-His Western blotting. The blots were quantified by densitometry analysis. The results are representative of three measurements. In Fig 2B, myc-tagged Tiam1 expressed in Cos-7 cell lysates were incubated with (His)₆-Rac1 in the presence of

increasing concentrations of NSC23766. The association of Rac1 with Tiam1 was examined by anti-His blot after anti-myc immunoprecipitation. In Fig 2C, the AU-tagged PDZ-RhoGEF was expressed in Cos-7 lysates and incubated with GST or GST-RhoA in the presence of varying concentrations of NSC23766. The RhoA associated PDZ-RhoGEF was probed with anti-AU antibody after affinity precipitation by glutathione agarose beads. In Fig 2D, (His)₆-Rac1 loaded with GTP γ S was incubated with GST-BcrGAP or GST-PAK1 (PBD) in the presence or absence of 200 μ M NSC23766 and the interaction with GSTBcrGAP or GST-PAK1 was probed by anti-His blot after affinity precipitation by glutathione agarose beads.

[0311] To determine if compound 23766 is capable of inhibiting the GEF-stimulates nucleotide exchange of Rac1, the mantGDP dissociation assays of Rac1 are carried out in the presence of increasing doses of compound 23766. In Fig.3. NSC23766 was effective in specifically inhibiting Rac1 GDP/GTP exchange stimulated by GEF. In Fig.3A, NSC23766 inhibited TrioN catalyzed GDP/GTP exchange of Rac1 in a dose dependent manner. 200 nM Rac1 loaded with mant-GDP was incubated at 25 °C in an exchange buffer containing 100 mM NaCl, 5 mM MgCl₂, 50 mM Tris-HCl (PH 7.6), and 0.5 mM GTP in the absence (top line) or presence of 100 nM TrioN. Increasing concentrations of NSC23766 were included in the exchange buffer as indicated. In Fig.3B, NSC23766 had no effect on the Intersectin-stimulated GDP/GTP exchange of Cdc42. 200 nM Cdc42 loaded with mant-GDP was incubated in the exchange buffer in the absence (top line) or presence of 100 nM Intersectin with or without 200 μ M NSC23766. In Fig.3C, the exchange reaction of RhoA catalyzed by PDZ-RhoGEF was carried out similarly in the presence or absence of 200 μ M NSC23766.

[0312] As shown in Fig. 3A, at increasing concentrations compound 23766 is able to block the mantGDP/GTP exchange catalyzed by Trio in a dose-dependent manner. On the other hand, compound 23766 has little impact on the Intersectin-stimulated mantGDP/GTP exchange of Cdc42 at similar doses (Fig. 3B), nor on the PDZ-RhoGEF-stimulated mantGDP/GTP exchange of RhoA. These results demonstrate that in vitro compounds, e.g., 23766, are able to specifically inhibit the interaction and activation of Rac1 by its GEFs.

[0313] Inhibitory effect of compound 23766 on Rac1 activity in vivo. In fibroblasts, Rac is activated by diverse stimuli including serum and PDGF (Hawkins et

al., 1995, which is incorporated herein by reference in its entirety). Rac activation in these situations is expected to be mediated by one or more Rac-specific GEFs such as Tiam1. To evaluate how compound 23766 can affect Rac activity in vivo, NIH 3T3 cells grown in 10% calf serum are treated with compound 23766 in different concentrations overnight, and the activation state of endogenous Rac1 in cells is detected by using the probe, GST-PAK (PBD) domain, that can specifically complex with Rac1-GTP. Fig. 4. shows that NSC23766 was effective in specifically inhibiting Rac1 activation in cells. In Fig. 4A, the activation states of endogenous Rac1, Cdc42 and RhoA in NIH3T3 cells with or without NSC23766 treatment were detected by the effector pull-down assays. At 80% confluency in the presence of 10% serum, NIH 3T3 cells in 100 mm dishes were treated with the indicated dosages of NSC23766 for 12 hours. Cell lysates containing similar amount of Rac1, Cdc42 or RhoA were incubated with the agarose immobilized GST-PAK1, GST-WASP or GST-Rhotekin, and the co-precipitates were subjected to anti-Rac1, Cdc42 or RhoA Western blot analysis to reveal the amount of GTP-Bound Rho proteins. In Fig. 4B, the inhibitory effect of NSC23766 on the PDGF-stimulated Rac1 activation was determined by the GST-PAK1 pull-down assay. Serum starved NIH 3T3 cells in the DMEM medium with different dosages of NSC23766 were treated with 10 nM PDGF for 2 minutes. In Fig. 4C, NSC23766 inhibited PDGF-stimulated lamellipodia formation. After overnight serum starvation in the presence or absence of 50 μ M NSC23766, Swiss 3T3 cells were treated with 10 nM PDGF for the indicated time. The cells were fixed and stained with Rhodamine-labeled phalloidin.

[0314] As shown in Fig. 4A, compound 23766 strongly inhibits Rac1 activation induced by serum. Densitometric analysis reveals that the IC_{50} of compound 23766 is about 40 μ M under these conditions. Meanwhile, the inhibitory effect of compound 23766 appears to be specific toward Rac among Rho GTPases, since the activation state of Cdc42 in these cells under serum-stimulation is unaffected by the presence of compound 23766. Interestingly, treatment with this reagent leads to a slightly increased level of RhoA-GTP in cells, consistent with previous reports suggesting that Rac1 can counter-react with RhoA activity. To examine if compound 23766 can affect Rac1 activation by PDGF stimulation, serum starved NIH 3T3 cells in the presence or absence of the compound are challenged with 10 nM PDGF for 2 minutes, and the cell lysates are assayed for the active Rac1-GTP species. Comparing with the PDGF-stimulated Rac activity in the absence of compound 23766, the cells treated with 50 μ M

23766 display a significant reduction of GTP-bound Rac (Fig. 4B), and the presence of 100 μ M 23766 leads to lower than basal level of Rac1-GTP in the cells. Thus, consistent with the *in vitro* Rac1-GEF interaction results, compound 23766 is able to specifically inhibit Rac1 activity *in vivo*.

[0315] PDGF activates Rac and induces Rac-mediated membrane ruffles and lamellipodia in fibroblasts (Hawkins et al., 1995; Ridley et al., 1992, which are incorporated herein by reference in their entirety). To evaluate the ability of compound 23766 to inhibit Rac1-mediated morphological changes, the actin cytoskeleton structures, induced by PDGF in the absence or presence of compound 23766, was examined. As shown in Fig. 4C, 10 nM PDGF potently stimulates membrane ruffling and lamellipodial formation in Swiss3T3 cells. However, in the presence of 100 μ M 23766, PDGF is only marginally effective in inducing lamellipodia at the cell edges at 5 min and completely ineffective at 10 min when the control cells that are not treated with compound 23766 displays significant lamellipodia structures. These results suggest that compound 23766 is effective in inhibiting Rac-mediated actin reorganization.

[0316] Compound 23766 specifically inhibits serum- or Trio-induced cell growth. Rho GTPase activities are important in cell growth regulation. Overexpression of dominant-negative Rac slows cell growth (Zheng et al., 1995b, which is incorporated herein by reference in its entirety). Conversely, constitutively active Rac increases growth rate of fibroblasts (Khosravi-Far et al., 1995, which is incorporated herein by reference in its entirety). Since compound 23766 is able to decrease Rac activity in NIH 3T3 cells, its effect on the growth properties of normal NIH 3T3 cells and the NIH 3T3 cells expressing constitutively active Rac1, L61Rac1 was examine.

[0317] Fig. 5. shows that NSC23766 specifically inhibited Rac GEF stimulated cell growth and transformation. In Fig.5A, wild type (WT) or L61Rac1 expressing NIH 3T3 cells were grown in 5% serum in the presence (---) or absence (—) of 100 μ M NSC23766. The cells were split in triplicate in 6-well plates at a density of 5×10^4 cells per well. The GTP-bound L61Rac1 and endogenous Rac1 of the L61Rac1-expressing cells were probed by GST-PAK1 pull-down after 12 hour treatment with increasing concentrations of NSC23766. In Fig.5B, WT or the GEF (Tiam1, Lbc or Vav) expressing NIH 3T3 cells were grown in 5% serum in the presence (---) or absence (—) of 100 μ M NSC23766, and the cell numbers were determined by daily cell counting. In Fig.5C, GST, L61Rac1, or Tiam1 transfected cells were treated with 50 μ M NSC23766

every two days. The foci numbers of the respective cells were quantified 14 days after transfection. In Fig.5D, a stable transfectant of Tiam1-expressing NIH 3T3 cells was cultured in 0.3% soft-agar medium for 14 days in the presence or absence of 100 μ M NSC23766. The number and the morphology of the colonies were examined under a microscope.

[0318] Comparison of the growth rates of the cells in the absence or presence of compound 23766 shows that compound 23766 slow the growth of wild type NIH 3T3 cells while having no effect on the growth rate of Rac1L61 expressing cells (Fig. 5A). The level of GTP-bound GST-Rac1L61 remains unchanged with or without the compound treatment, whereas the endogenous Rac activity is decreased significantly by the presence of compound 23766 (data not shown). These results suggest that the inhibitory effect of compound 23766 on cell growth correlates with its ability to inhibit cellular Rac activity.

[0319] Due to their ability to directly activate Rho GTPases, Dbl family GEFs are potent stimulators of cell proliferation. Compound 23766 is capable of inhibiting the cell growth induced by the Rac specific GEF Trio, but not that stimulated by the Rho-specific GEF Lbc, the Cdc42-specific GEF Intersectin, or the multiple Rho protein-activating GEF Vav (Fig. 5B). Thus compound 23766 is effective in specifically inhibiting cell growth caused by GEF-induced Rac activation.

[0320] Reversal of the PC-3 tumor cell phenotypes by compound 23766. Elevation of Rac1 activity is associated with cancer cell hyperproliferative and invasive properties. Next the effect of compound 23766 is tested on the growth and invasion capabilities of a prostate cancer cell line, PC-3. PC-3 cells are malignant prostate adenocarcinoma cells derived from the bone metastases of a patient with prostate cancer (Kaighn et al., 1979, which is incorporated herein by reference in its entirety). They are transforming and highly invasive (Lang et al., 2002, which is incorporated herein by reference in its entirety). The mRNA of the PTEN tumor suppressor is undetectable in these cells (Bastola et al., 2002, which is incorporated herein by reference in its entirety), and loss of PTEN has previously been correlated with Rac1 hyperactivation due to the significant increase of PIP₃ level (Liliental et al., 2000, which is incorporated herein by reference in its entirety). When the activity of endogenous Rac1 in PC-3 cells is examined by probing with GST-PAK (PBD), a ~100% higher level of GTP-bound Rac than that of the normal prostate epithelial RWPE-1 cells is observed (Fig. 5A). Consistent with the results obtained from fibroblasts, compound 23766 is able to inhibit

Rac1 activity in PC-3 cells (Fig.6A). Correlating with the decreased Rac1 activity, the proliferation rates of the compound 23766 treated PC-3 cells are inhibited by compound 23766 in a dose dependent manner (Fig. 6A). These results suggest that compound 23766 can effectively inhibit PC-3 tumor cell growth through down-regulation of Rac1 activity.

[0321] Fig. 6. shows that NSC23766 inhibited the proliferation, anchorage independent growth and invasion of PC-3 prostate cancer cells. In Fig. 6A, PC-3 cells were grown in 5% calf serum supplemented with the indicated concentrations of NSC23766. The cells were split in triplicate in 96-wells at 1.5×10^3 cells per well. Cell numbers were assayed by using CellTiter 96 AQueous cell proliferation assay kit in different days. In Fig. 6B, PC-3 and RWPE-1 prostate epithelial cells (1.25×10^3 per well) were grown in 0.3% agarose in different doses of NSC23766, and the number of colonies formed in soft agar was quantified 12 days after plating. In Fig. 6C, PC-3 cells were placed in an invasion chamber for 24 hrs at 37°C in the absence or presence of 25 μ M NSC23766. Cells invaded through Matrigel matrix were visualized with Giemasa staining.

[0322] Given that PC-3 cells contain hyperactive Rac1 activity, the ability of PC-3 cells to grow on soft agar and the effect of compound 23766 on its anchorage independent growth property can be tested. Fig. 6B shows that PC-3 cells readily form colonies ten days after being placed on soft agar, under conditions in which the normal prostate epithelia RWPE-1 cells are unable to grow. Compound 23766 efficiently blocks the colony forming activity of PC-3 cells. Approximately 10% and 1% colony-forming activities remain after treatment of the cells with 25 μ M and 50 μ M compound 23766, respectively. Moreover, the size of colonies of the treated cells appears much smaller than those of the untreated ones (Fig. 6B). PC-3 cells are reported to possess highly invasive activity (Lang et al., 2002, which is incorporated herein by reference in its entirety), which is evident in a Matri-gel invasion assay. Under similar conditions, RWPE-1 cells are non-invasive. At a dose of 25 μ M, compound 23766 significantly inhibits PC-3 cell invasion (Fig. 6C).

[0323] Taken together, these results show that the active agent is able to downregulate Rac1 activity of PC-3 tumor cells which likely results in the reversal of the proliferation, anchorage independent growth and invasion phenotypes.

Crystal Structure of Rac-1 GTPase complexed with NCI 23766Methods

[0324] **Protein Purification:** *Escherichia coli* BL21 Codon Plus (Stratagene) cells expressing His₆-tagged Rac-1 domain (residues 1 to 185 of SEQ ID NO:1) were grown at 37°C in Luria-Bertani medium to an optical Density at 600 nm of 0.8-1.0 and induced for 3 hours with 0.5 mM IPTG. The harvested cells were lysed in 20 mM Tris, pH 8.0, 500 mM NaCl, 5 mM MgCl₂ by French pressure cell. The lysate was clarified by centrifugation for 20 minutes at 16,000 rpm in a 16.250 rotor (Beckman). Clarified lysate was passed over a Ni-NTA column (Qiagen) pre-equilibrated with 5 mM Imidazole in Buffer A (20 mM Tris-HCl, pH 8.0, 500 mM NaCl). The column was washed with 20 mM Imidazole in Buffer A. After reaching baseline, the His₆-tagged Rac1 was eluted from the column with 250 mM Imidazole in Buffer A and dialyzed against 20 mM Tris-HCl, pH 8.0, 5 mM MgCl₂, 5 mM β-Mercaptoethanol and digested with thrombin (Sigma) overnight. The digested protein was passed over a Source-15Q (Sigma) equilibrated with 20 mM Tris-HCl, pH 8.0, 5 mM MgCl₂, 5 mM β-Mercaptoethanol. Rac-1 domain polypeptide comprising residues 1 through 185 of SEQ ID NO:1 was collected in the flowthrough. The flowthrough was then passed over a Superdex 200 (Amersham Biosciences) size exclusion column equilibrated with 10 mM HEPES, pH 7.5, 2 mM MgCl₂, 150 mM NaCl. The peak corresponding to Rac-1 was pooled, concentrated to 22.5 mg/mL and used in crystallization trials.

[0325] **Crystallization:** Crystals of Rac1•GDP were grown in 20% PEG 8000, 5% Ethylene Glycol, 15% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, and 1 mM NSC23766 at 20°C using the hanging drop method. Crystals of the complex appeared overnight and reached their maximum size after 2-3 days. Crystals were transferred to a drop containing 18% PEG 8000, 3.5% Ethylene Glycol, 10.7% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, 14.3 mM NSC23766 and allowed to equilibrate over a reservoir containing 25% PEG 8000, 5% Ethylene Glycol, 15% DMSO 0.1M HEPES, pH 7.0, 10 mM DTT. After 22 hrs of equilibration, the crystals were mounted and flash frozen in liquid nitrogen.

[0326] **X-ray Data Collection and Structure Solution:** A complete and redundant diffraction data set to 1.35 Å resolution was collected at the National Synchrotron Light Source (Brookhaven, Upton, NY) beamline X26C on an ADSC Quantum 4 CCD detector at a temperature of 100 K, integrated, and scaled with the

HKL2000 suite (*Otwinowski 1997 & 2003*). The Rac/NSC23766 crystals belong to space group $P2_1$ with one complex per asymmetric unit and unit cell dimensions of $a = 41.8 \text{ \AA}$, $b = 40.0 \text{ \AA}$, $c = 52.1 \text{ \AA}$ $\gamma = 105.83^\circ$. The Rac/NSC23766 structure was solved by molecular replacement using the coordinates of Rac•GDP (*K. Skowronek & N. Nassar*, unpublished observations) as the search model in the program MOLREP (*Vagin & Teplyakov 1997*). The actual model is refined using TLS groups (*Winn 2003*) in REFMAC (*Murshudov 1997*) against $1.35 \text{ \AA}$ resolution data. During refinement, an extra density in a hydrophobic pocket formed by Ile36, Trp56, Tyr64, Arg68, and Pro73 was apparent in the electron density. This extra density is not part of Rac and does not correspond to a metal ion or a water molecule. Subsequently, an NSC23766 molecule was modeled in this density and is being refined against the experimental data.

[0327] Table 1 depicts atomic coordinates of the Rac/NSC23766 structure as solved as described above and partially refined. Table 2 depicts further refined to $1.50 \text{ \AA}$ resolution, with crystallographic residuals R/Rfree of 15.1/20.5 and good stereochemistry.

[0328] It should be noted that the side chains of some residues are poorly resolved in the electron density. These are typically truncated to alanines in the listing of atomic coordinates (e.g., Table 1 or Table 2).

[0329] In addition, information regarding procedural or other details supplementary to those set forth herein, are described in cited references specifically incorporated herein by reference.

[0330] It is be evident to those skilled in the art that modifications or variations can be made to the preferred embodiment described herein without departing from the novel teachings of the present invention. All such modifications and variations are intended to be incorporated herein and within the scope of the claims.

Table 1

REMARK Written by O version 9.0.7

CRYST1	41.792	40.008	52.071	90.00	105.83	90.00			
ORIGX1	1.000000	0.000000	0.000000			0.000000			
ORIGX2	0.000000	1.000000	0.000000			0.000000			
ORIGX3	0.000000	0.000000	1.000000			0.000000			
SCALE1	0.023928	-0.000001	0.006784			0.000000			
SCALE2	0.000000	0.024995	-0.000001			0.000000			
SCALE3	0.000000	0.000000	0.019961			0.000000			
ATOM	1	N	ALA A	1	15.756	6.560	-10.373	1.00	30.78
ATOM	2	CA	ALA A	1	16.093	6.323	-8.944	1.00	29.95
ATOM	3	CB	ALA A	1	15.708	7.535	-8.107	1.00	30.54
ATOM	4	C	ALA A	1	15.397	5.066	-8.426	1.00	29.34
ATOM	5	O	ALA A	1	14.165	4.991	-8.415	1.00	29.88
ATOM	6	N	ALA A	2	16.190	4.079	-8.013	1.00	28.31
ATOM	7	CA	ALA A	2	15.653	2.839	-7.447	1.00	26.70
ATOM	8	CB	ALA A	2	16.739	1.776	-7.330	1.00	27.15
ATOM	9	C	ALA A	2	15.066	3.147	-6.080	1.00	25.29
ATOM	10	O	ALA A	2	15.671	3.887	-5.294	1.00	25.34
ATOM	11	N	ALA A	3	13.888	2.584	-5.813	1.00	23.09
ATOM	12	CA	ALA A	3	13.185	2.797	-4.545	1.00	21.06
ATOM	13	CB	ALA A	3	11.794	2.144	-4.577	1.00	21.37
ATOM	14	C	ALA A	3	13.992	2.272	-3.360	1.00	19.51
ATOM	15	O	ALA A	3	14.532	1.151	-3.409	1.00	21.10
ATOM	16	N	ILE A	4	14.103	3.097	-2.322	1.00	16.92
ATOM	17	CA	ILE A	4	14.495	2.610	-1.001	1.00	15.02
ATOM	18	CB	ILE A	4	14.648	3.775	-0.026	1.00	14.41
ATOM	19	CG1	ILE A	4	15.895	4.579	-0.429	1.00	16.51
ATOM	20	CD1	ILE A	4	16.154	5.841	0.397	1.00	16.77
ATOM	21	CG2	ILE A	4	14.740	3.274	1.433	1.00	15.96
ATOM	22	C	ILE A	4	13.388	1.675	-0.542	1.00	14.20
ATOM	23	O	ILE A	4	12.232	2.101	-0.406	1.00	15.07
ATOM	24	N	LYS A	5	13.726	0.408	-0.273	1.00	12.22
ATOM	25	CA	LYS A	5	12.723	-0.574	0.144	1.00	11.59
ATOM	26	CB	LYS A	5	13.060	-1.966	-0.389	1.00	11.41
ATOM	27	CG	LYS A	5	12.127	-3.072	0.138	1.00	11.13
ATOM	28	CD	LYS A	5	12.354	-4.367	-0.588	1.00	12.02
ATOM	29	CE	LYS A	5	11.520	-5.470	0.029	1.00	13.44
ATOM	30	NZ	LYS A	5	11.555	-6.719	-0.805	1.00	14.02
ATOM	31	C	LYS A	5	12.599	-0.604	1.665	1.00	11.00
ATOM	32	O	LYS A	5	13.549	-0.951	2.376	1.00	11.63
ATOM	33	N	CYS A	6	11.416	-0.249	2.160	1.00	10.99
ATOM	34	CA	CYS A	6	11.150	-0.184	3.592	1.00	10.47
ATOM	35	CB	CYS A	6	10.616	1.202	3.918	1.00	11.97
ATOM	36	SG	CYS A	6	10.107	1.479	5.616	1.00	12.97
ATOM	37	C	CYS A	6	10.108	-1.234	3.915	1.00	10.18
ATOM	38	O	CYS A	6	8.997	-1.192	3.382	1.00	10.84
ATOM	39	N	VAL A	7	10.453	-2.173	4.789	1.00	10.01
ATOM	40	CA	VAL A	7	9.549	-3.260	5.155	1.00	9.99
ATOM	41	CB	VAL A	7	10.261	-4.622	5.022	1.00	11.17
ATOM	42	CG1	VAL A	7	9.362	-5.747	5.495	1.00	12.37
ATOM	43	CG2	VAL A	7	10.680	-4.873	3.596	1.00	12.17
ATOM	44	C	VAL A	7	9.052	-3.010	6.579	1.00	10.17
ATOM	45	O	VAL A	7	9.833	-2.691	7.463	1.00	10.48
ATOM	46	N	VAL A	8	7.758	-3.159	6.798	1.00	10.12
ATOM	47	CA	VAL A	8	7.114	-2.817	8.057	1.00	10.09
ATOM	48	CB	VAL A	8	5.893	-1.924	7.771	1.00	10.08
ATOM	49	CG1	VAL A	8	5.184	-1.529	9.081	1.00	10.78

ATOM	50	CG2	VAL	A	8	6.324	-0.697	6.939	1.00	11.24
ATOM	51	C	VAL	A	8	6.658	-4.129	8.684	1.00	9.54
ATOM	52	O	VAL	A	8	5.908	-4.894	8.046	1.00	9.78
ATOM	53	N	VAL	A	9	7.144	-4.421	9.887	1.00	9.18
ATOM	54	CA	VAL	A	9	6.827	-5.669	10.590	1.00	9.67
ATOM	55	CB	VAL	A	9	8.027	-6.665	10.597	1.00	9.51
ATOM	56	CG1	VAL	A	9	8.500	-6.952	9.170	1.00	11.60
ATOM	57	CG2	VAL	A	9	9.183	-6.171	11.474	1.00	11.98
ATOM	58	C	VAL	A	9	6.399	-5.370	12.019	1.00	9.36
ATOM	59	O	VAL	A	9	6.596	-4.276	12.541	1.00	9.87
ATOM	60	N	GLY	A	10	5.812	-6.369	12.648	1.00	9.30
ATOM	61	CA	GLY	A	10	5.346	-6.288	14.024	1.00	10.14
ATOM	62	C	GLY	A	10	4.095	-7.136	14.205	1.00	9.38
ATOM	63	O	GLY	A	10	3.500	-7.647	13.243	1.00	9.61
ATOM	64	N	ASP	A	11	3.702	-7.283	15.466	1.00	9.39
ATOM	65	CA	ASP	A	11	2.596	-8.170	15.822	1.00	10.45
ATOM	66	CB	ASP	A	11	2.329	-8.105	17.323	1.00	10.06
ATOM	67	CG	ASP	A	11	3.288	-8.959	18.135	1.00	12.02
ATOM	68	OD1	ASP	A	11	4.248	-9.545	17.584	1.00	12.12
ATOM	69	OD2	ASP	A	11	3.029	-9.044	19.354	1.00	13.04
ATOM	70	C	ASP	A	11	1.324	-7.788	15.111	1.00	9.85
ATOM	71	O	ASP	A	11	1.101	-6.621	14.774	1.00	9.42
ATOM	72	N	GLY	A	12	0.433	-8.753	14.932	1.00	10.76
ATOM	73	CA	GLY	A	12	-0.908	-8.453	14.437	1.00	10.53
ATOM	74	C	GLY	A	12	-1.572	-7.408	15.312	1.00	10.68
ATOM	75	O	GLY	A	12	-1.433	-7.428	16.541	1.00	11.67
ATOM	76	N	ALA	A	13	-2.290	-6.488	14.674	1.00	10.27
ATOM	77	CA	ALA	A	13	-3.124	-5.478	15.320	1.00	10.10
ATOM	78	CB	ALA	A	13	-4.081	-6.090	16.347	1.00	11.47
ATOM	79	C	ALA	A	13	-2.372	-4.292	15.913	1.00	10.23
ATOM	80	O	ALA	A	13	-2.991	-3.454	16.586	1.00	11.32
ATOM	81	N	VAL	A	14	-1.073	-4.187	15.662	1.00	10.56
ATOM	82	CA	VAL	A	14	-0.355	-2.999	16.193	1.00	10.35
ATOM	83	CB	VAL	A	14	1.161	-3.227	16.365	1.00	10.51
ATOM	84	CG1	VAL	A	14	1.444	-4.399	17.307	1.00	11.36
ATOM	85	CG2	VAL	A	14	1.861	-3.405	15.042	1.00	11.73
ATOM	86	C	VAL	A	14	-0.617	-1.727	15.403	1.00	10.69
ATOM	87	O	VAL	A	14	-0.410	-0.624	15.917	1.00	11.18
ATOM	88	N	GLY	A	15	-1.108	-1.859	14.176	1.00	9.47
ATOM	89	CA	GLY	A	15	-1.432	-0.721	13.333	1.00	10.07
ATOM	90	C	GLY	A	15	-0.547	-0.518	12.119	1.00	9.10
ATOM	91	O	GLY	A	15	-0.502	0.581	11.585	1.00	9.75
ATOM	92	N	LYS	A	16	0.079	-1.580	11.616	1.00	8.57
ATOM	93	CA	LYS	A	16	0.976	-1.451	10.467	1.00	8.65
ATOM	94	CB	LYS	A	16	1.674	-2.782	10.183	1.00	8.61
ATOM	95	CG	LYS	A	16	2.485	-3.314	11.373	1.00	9.93
ATOM	96	CD	LYS	A	16	3.194	-4.658	11.106	1.00	10.00
ATOM	97	CE	LYS	A	16	2.275	-5.817	10.819	1.00	10.39
ATOM	98	NZ	LYS	A	16	1.366	-6.104	11.952	1.00	9.45
ATOM	99	C	LYS	A	16	0.227	-0.996	9.228	1.00	8.25
ATOM	100	O	LYS	A	16	0.641	-0.060	8.535	1.00	8.46
ATOM	101	N	THR	A	17	-0.875	-1.672	8.912	1.00	8.80
ATOM	102	CA	THR	A	17	-1.615	-1.343	7.710	1.00	9.41
ATOM	103	CB	THR	A	17	-2.689	-2.400	7.503	1.00	8.99
ATOM	104	OG1	THR	A	17	-2.020	-3.677	7.367	1.00	9.21
ATOM	105	CG2	THR	A	17	-3.503	-2.155	6.248	1.00	10.06
ATOM	106	C	THR	A	17	-2.204	0.070	7.800	1.00	8.93
ATOM	107	O	THR	A	17	-2.136	0.834	6.828	1.00	9.41
ATOM	108	N	CYS	A	18	-2.784	0.423	8.943	1.00	9.42
ATOM	109	CA	CYS	A	18	-3.355	1.741	9.077	1.00	10.25
ATOM	110	CB	CYS	A	18	-4.075	1.897	10.396	1.00	10.15
ATOM	111	SG	CYS	A	18	-5.693	1.128	10.433	1.00	11.45
ATOM	112	C	CYS	A	18	-2.296	2.805	8.984	1.00	10.14
ATOM	113	O	CYS	A	18	-2.576	3.871	8.414	1.00	11.24
ATOM	114	N	LEU	A	19	-1.114	2.611	9.549	1.00	10.34
ATOM	115	CA	LEU	A	19	-0.175	3.719	9.443	1.00	11.62

ATOM	116	C	LEU	A	19	0.273	3.893	8.004	1.00	11.02
ATOM	117	O	LEU	A	19	0.512	5.022	7.575	1.00	11.14
ATOM	118	CB	LEU	A	19	1.025	3.494	10.365	1.00	12.92
ATOM	119	CG	LEU	A	19	1.988	2.375	9.965	1.00	20.00
ATOM	120	CD1	LEU	A	19	3.014	2.881	8.963	1.00	15.00
ATOM	121	CD2	LEU	A	19	2.677	1.795	11.190	1.00	20.00
ATOM	122	N	LEU	A	20	0.350	2.817	7.229	1.00	10.17
ATOM	123	CA	LEU	A	20	0.685	2.943	5.809	1.00	10.15
ATOM	124	CB	LEU	A	20	1.019	1.577	5.203	1.00	10.06
ATOM	125	CG	LEU	A	20	2.331	0.991	5.722	1.00	9.84
ATOM	126	CD1	LEU	A	20	2.432	-0.485	5.309	1.00	12.20
ATOM	127	CD2	LEU	A	20	3.522	1.778	5.244	1.00	13.56
ATOM	128	C	LEU	A	20	-0.421	3.631	5.007	1.00	10.29
ATOM	129	O	LEU	A	20	-0.132	4.471	4.153	1.00	10.44
ATOM	130	N	ILE	A	21	-1.669	3.293	5.279	1.00	10.12
ATOM	131	CA	ILE	A	21	-2.789	3.912	4.590	1.00	10.14
ATOM	132	CB	ILE	A	21	-4.093	3.182	4.880	1.00	10.76
ATOM	133	CG1	ILE	A	21	-4.044	1.818	4.193	1.00	11.42
ATOM	134	CD1	ILE	A	21	-5.125	0.806	4.611	1.00	13.68
ATOM	135	CG2	ILE	A	21	-5.300	3.964	4.352	1.00	12.21
ATOM	136	C	ILE	A	21	-2.883	5.381	4.987	1.00	9.92
ATOM	137	O	ILE	A	21	-3.078	6.234	4.112	1.00	10.89
ATOM	138	N	SER	A	22	-2.730	5.697	6.259	0.50	6.37
ATOM	139	CA	SER	A	22	-2.774	7.094	6.668	0.50	6.91
ATOM	140	CB	SER	A	22	-2.711	7.193	8.182	0.50	6.73
ATOM	141	OG	ASER	A	22	-3.945	6.769	8.703	0.50	12.30
ATOM	141	OG	BSER	A	22	-2.826	8.535	8.599	0.50	11.36
ATOM	142	C	SER	A	22	-1.642	7.893	6.019	0.50	6.92
ATOM	143	O	SER	A	22	-1.841	9.031	5.573	0.50	5.72
ATOM	144	N	TYR	A	23	-0.449	7.310	5.949	1.00	10.57
ATOM	145	CA	TYR	A	23	0.681	8.011	5.355	1.00	11.21
ATOM	146	CB	TYR	A	23	1.963	7.202	5.563	1.00	11.78
ATOM	147	CG	TYR	A	23	3.203	7.835	4.958	1.00	11.21
ATOM	148	CD1	TYR	A	23	3.655	9.080	5.389	1.00	11.88
ATOM	149	CE1	TYR	A	23	4.790	9.656	4.842	1.00	11.58
ATOM	150	CZ	TYR	A	23	5.513	8.958	3.884	1.00	11.24
ATOM	151	OH	TYR	A	23	6.654	9.496	3.318	1.00	12.92
ATOM	152	CE2	TYR	A	23	5.091	7.730	3.446	1.00	11.94
ATOM	153	CD2	TYR	A	23	3.953	7.167	3.992	1.00	11.19
ATOM	154	C	TYR	A	23	0.479	8.286	3.869	1.00	12.23
ATOM	155	O	TYR	A	23	0.855	9.356	3.343	1.00	13.13
ATOM	156	N	THR	A	24	-0.100	7.321	3.172	1.00	12.75
ATOM	157	CA	THR	A	24	-0.174	7.422	1.715	1.00	13.41
ATOM	158	CB	THR	A	24	-0.089	6.044	1.027	1.00	13.28
ATOM	159	OG1	THR	A	24	-1.141	5.191	1.515	1.00	12.91
ATOM	160	CG2	THR	A	24	1.259	5.394	1.283	1.00	13.69
ATOM	161	C	THR	A	24	-1.406	8.160	1.230	1.00	14.93
ATOM	162	O	THR	A	24	-1.369	8.734	0.143	1.00	15.84
ATOM	163	N	THR	A	25	-2.477	8.181	2.022	1.00	14.62
ATOM	164	CA	THR	A	25	-3.754	8.759	1.589	1.00	15.96
ATOM	165	CB	THR	A	25	-4.890	7.711	1.573	1.00	15.67
ATOM	166	OG1	THR	A	25	-5.267	7.374	2.925	1.00	16.40
ATOM	167	CG2	THR	A	25	-4.498	6.474	0.785	1.00	15.73
ATOM	168	C	THR	A	25	-4.234	9.918	2.459	1.00	17.13
ATOM	169	O	THR	A	25	-5.197	10.593	2.089	1.00	17.10
ATOM	170	N	ASN	A	26	-3.626	10.103	3.626	1.00	18.51
ATOM	171	CA	ASN	A	26	-4.079	11.125	4.599	1.00	20.28
ATOM	172	CB	ASN	A	26	-3.885	12.548	4.040	1.00	21.61
ATOM	173	CG	ASN	A	26	-2.437	12.847	3.726	1.00	23.49
ATOM	174	OD1	ASN	A	26	-2.108	13.367	2.650	1.00	27.93
ATOM	175	ND2	ASN	A	26	-1.546	12.502	4.657	1.00	25.36
ATOM	176	C	ASN	A	26	-5.496	10.915	5.133	1.00	21.61
ATOM	177	O	ASN	A	26	-6.133	11.858	5.625	1.00	22.35
ATOM	178	N	ALA	A	27	-5.999	9.690	5.041	1.00	21.20
ATOM	179	CA	ALA	A	27	-7.270	9.364	5.665	1.00	21.90
ATOM	180	CB	ALA	A	27	-8.324	9.074	4.626	1.00	22.79

ATOM	181	C	ALA	A	27	-7.036	8.154	6.561	1.00	21.66
ATOM	182	O	ALA	A	27	-6.462	7.172	6.115	1.00	22.33
ATOM	183	N	PHE	A	28	-7.410	8.257	7.830	1.00	19.80
ATOM	184	CA	PHE	A	28	-7.389	7.090	8.715	1.00	17.99
ATOM	185	CB	PHE	A	28	-7.504	7.524	10.165	1.00	17.74
ATOM	186	CG	PHE	A	28	-7.569	6.380	11.124	1.00	16.32
ATOM	187	CD1	PHE	A	28	-6.452	5.574	11.360	1.00	16.38
ATOM	188	CE1	PHE	A	28	-6.516	4.504	12.268	1.00	14.61
ATOM	189	CZ	PHE	A	28	-7.713	4.235	12.930	1.00	16.53
ATOM	190	CE2	PHE	A	28	-8.840	5.025	12.686	1.00	16.63
ATOM	191	CD2	PHE	A	28	-8.759	6.095	11.783	1.00	15.57
ATOM	192	C	PHE	A	28	-8.544	6.155	8.342	1.00	17.65
ATOM	193	O	PHE	A	28	-9.707	6.563	8.334	1.00	17.89
ATOM	194	N	PRO	A	29	-8.237	4.897	7.982	1.00	16.19
ATOM	195	CA	PRO	A	29	-9.275	4.005	7.446	1.00	15.78
ATOM	196	CB	PRO	A	29	-8.457	3.021	6.603	1.00	14.96
ATOM	197	CG	PRO	A	29	-7.162	2.881	7.423	1.00	15.06
ATOM	198	CD	PRO	A	29	-6.889	4.281	7.933	1.00	15.67
ATOM	199	C	PRO	A	29	-10.111	3.225	8.456	1.00	16.17
ATOM	200	O	PRO	A	29	-11.017	2.496	8.064	1.00	16.59
ATOM	201	N	GLY	A	30	-9.816	3.343	9.741	1.00	16.47
ATOM	202	CA	GLY	A	30	-10.613	2.622	10.746	1.00	16.48
ATOM	203	C	GLY	A	30	-10.533	1.107	10.589	1.00	16.58
ATOM	204	O	GLY	A	30	-9.443	0.543	10.592	1.00	16.49
ATOM	205	N	GLU	A	31	-11.681	0.457	10.447	1.00	16.20
ATOM	206	CA	GLU	A	31	-11.752	-0.994	10.256	1.00	16.97
ATOM	207	CB	GLU	A	31	-12.970	-1.573	10.986	1.00	18.18
ATOM	208	CG	GLU	A	31	-13.022	-1.270	12.463	1.00	22.26
ATOM	209	CD	GLU	A	31	-11.703	-1.558	13.161	1.00	26.99
ATOM	210	OE1	GLU	A	31	-11.201	-2.698	13.072	1.00	30.47
ATOM	211	OE2	GLU	A	31	-11.164	-0.635	13.796	1.00	31.76
ATOM	212	C	GLU	A	31	-11.802	-1.400	8.780	1.00	15.98
ATOM	213	O	GLU	A	31	-11.844	-2.590	8.449	1.00	16.42
ATOM	214	N	TYR	A	32	-11.785	-0.410	7.900	1.00	16.12
ATOM	215	CA	TYR	A	32	-11.922	-0.686	6.468	1.00	17.14
ATOM	216	CB	TYR	A	32	-12.725	0.415	5.782	1.00	18.87
ATOM	217	CG	TYR	A	32	-14.165	0.415	6.240	1.00	20.99
ATOM	218	CD1	TYR	A	32	-15.095	-0.446	5.667	1.00	23.72
ATOM	219	CE1	TYR	A	32	-16.417	-0.460	6.093	1.00	26.05
ATOM	220	CZ	TYR	A	32	-16.819	0.403	7.097	1.00	24.58
ATOM	221	OH	TYR	A	32	-18.130	0.394	7.507	1.00	25.90
ATOM	222	CE2	TYR	A	32	-15.913	1.267	7.697	1.00	23.82
ATOM	223	CD2	TYR	A	32	-14.590	1.270	7.261	1.00	22.07
ATOM	224	C	TYR	A	32	-10.556	-0.856	5.848	1.00	15.47
ATOM	225	O	TYR	A	32	-10.058	-0.007	5.111	1.00	15.10
ATOM	226	N	ILE	A	33	-9.935	-1.979	6.210	1.00	14.88
ATOM	227	CA	ILE	A	33	-8.540	-2.215	5.873	1.00	14.90
ATOM	228	CB	ILE	A	33	-7.635	-1.892	7.090	1.00	15.46
ATOM	229	CG1	ILE	A	33	-8.141	-2.656	8.343	1.00	17.42
ATOM	230	CD1	ILE	A	33	-7.582	-2.237	9.684	1.00	17.18
ATOM	231	CG2	ILE	A	33	-7.558	-0.378	7.332	1.00	14.83
ATOM	232	C	ILE	A	33	-8.325	-3.670	5.511	1.00	13.45
ATOM	233	O	ILE	A	33	-9.015	-4.544	6.032	1.00	14.07
ATOM	234	N	PRO	A	34	-7.348	-3.942	4.636	1.00	11.93
ATOM	235	CA	PRO	A	34	-6.936	-5.310	4.369	1.00	11.72
ATOM	236	CB	PRO	A	34	-6.181	-5.179	3.043	1.00	11.45
ATOM	237	CG	PRO	A	34	-5.535	-3.809	3.161	1.00	12.24
ATOM	238	CD	PRO	A	34	-6.560	-2.968	3.857	1.00	11.49
ATOM	239	C	PRO	A	34	-5.994	-5.804	5.481	1.00	11.22
ATOM	240	O	PRO	A	34	-5.495	-4.999	6.281	1.00	10.92
ATOM	241	N	THR	A	35	-5.698	-7.102	5.504	1.00	10.66
ATOM	242	CA	THR	A	35	-4.644	-7.603	6.366	1.00	10.28
ATOM	243	CB	THR	A	35	-4.530	-9.132	6.220	1.00	10.32
ATOM	244	OG1	THR	A	35	-5.787	-9.740	6.564	1.00	10.83
ATOM	245	CG2	THR	A	35	-3.440	-9.719	7.103	1.00	10.76
ATOM	246	C	THR	A	35	-3.317	-6.954	5.972	1.00	9.92

ATOM	247	O	THR	A	35	-2.573	-6.502	6.855	1.00	9.20
ATOM	248	N	VAL	A	36	-3.026	-6.915	4.676	1.00	9.92
ATOM	249	CA	VAL	A	36	-1.765	-6.390	4.189	1.00	11.66
ATOM	250	CB	VAL	A	36	-0.911	-7.532	3.564	1.00	12.55
ATOM	251	CG1	VAL	A	36	0.435	-7.006	3.135	1.00	14.75
ATOM	252	CG2	VAL	A	36	-0.764	-8.749	4.538	1.00	13.86
ATOM	253	C	VAL	A	36	-2.045	-5.353	3.100	1.00	11.71
ATOM	254	O	VAL	A	36	-2.736	-5.644	2.124	1.00	12.92
ATOM	255	N	PHE	A	37	-1.555	-4.136	3.293	1.00	11.91
ATOM	256	CA	PHE	A	37	-1.630	-3.073	2.301	1.00	12.80
ATOM	257	CB	PHE	A	37	-0.957	-1.854	2.933	1.00	13.56
ATOM	258	CG	PHE	A	37	-1.143	-0.571	2.182	1.00	16.30
ATOM	259	CD1	PHE	A	37	-2.405	-0.166	1.753	1.00	19.42
ATOM	260	CE1	PHE	A	37	-2.578	1.058	1.071	1.00	20.61
ATOM	261	CZ	PHE	A	37	-1.475	1.871	0.845	1.00	17.13
ATOM	262	CE2	PHE	A	37	-0.211	1.483	1.264	1.00	17.41
ATOM	263	CD2	PHE	A	37	-0.050	0.250	1.948	1.00	18.70
ATOM	264	C	PHE	A	37	-0.893	-3.459	1.039	1.00	14.08
ATOM	265	O	PHE	A	37	0.169	-4.074	1.120	1.00	14.02
ATOM	266	N	ASP	A	38	-1.434	-3.055	-0.110	1.00	15.41
ATOM	267	CA	ASP	A	38	-0.772	-3.259	-1.404	1.00	17.12
ATOM	268	CB	ASP	A	38	-1.618	-2.627	-2.533	1.00	18.72
ATOM	269	CG	ASP	A	38	-2.911	-3.389	-2.846	1.00	23.03
ATOM	270	OD1	ASP	A	38	-3.085	-4.547	-2.409	1.00	26.95
ATOM	271	OD2	ASP	A	38	-3.757	-2.798	-3.563	1.00	28.04
ATOM	272	C	ASP	A	38	0.616	-2.577	-1.389	1.00	16.38
ATOM	273	O	ASP	A	38	0.804	-1.560	-0.735	1.00	16.07
ATOM	274	N	ASN	A	39	1.584	-3.135	-2.099	1.00	16.55
ATOM	275	CA	ASN	A	39	2.857	-2.436	-2.288	1.00	17.38
ATOM	276	CB	ASN	A	39	3.694	-3.191	-3.309	1.00	18.41
ATOM	277	CG	ASN	A	39	5.083	-2.657	-3.403	1.00	22.02
ATOM	278	OD1	ASN	A	39	5.471	-2.085	-4.423	1.00	26.34
ATOM	279	ND2	ASN	A	39	5.844	-2.809	-2.327	1.00	27.14
ATOM	280	C	ASN	A	39	2.648	-1.011	-2.788	1.00	16.04
ATOM	281	O	ASN	A	39	1.772	-0.761	-3.622	1.00	16.15
ATOM	282	N	TYR	A	40	3.440	-0.068	-2.280	1.00	14.53
ATOM	283	CA	TYR	A	40	3.306	1.337	-2.665	1.00	14.34
ATOM	284	CB	TYR	A	40	2.592	2.105	-1.544	1.00	14.12
ATOM	285	CG	TYR	A	40	2.419	3.582	-1.815	1.00	14.39
ATOM	286	CD1	TYR	A	40	1.294	4.049	-2.490	1.00	14.96
ATOM	287	CE1	TYR	A	40	1.125	5.409	-2.749	1.00	14.64
ATOM	288	CZ	TYR	A	40	2.090	6.291	-2.330	1.00	15.18
ATOM	289	OH	TYR	A	40	1.898	7.628	-2.607	1.00	16.39
ATOM	290	CE2	TYR	A	40	3.220	5.859	-1.665	1.00	14.84
ATOM	291	CD2	TYR	A	40	3.385	4.500	-1.414	1.00	14.38
ATOM	292	C	TYR	A	40	4.683	1.934	-2.873	1.00	13.60
ATOM	293	O	TYR	A	40	5.580	1.718	-2.064	1.00	14.93
ATOM	294	N	SER	A	41	4.875	2.685	-3.951	1.00	12.48
ATOM	295	CA	SER	A	41	6.077	3.495	-4.068	1.00	13.53
ATOM	296	CB	SER	A	41	7.158	2.872	-4.960	1.00	14.96
ATOM	297	OG	SER	A	41	6.766	2.869	-6.306	1.00	19.82
ATOM	298	C	SER	A	41	5.706	4.876	-4.559	1.00	12.45
ATOM	299	O	SER	A	41	4.742	5.037	-5.308	1.00	13.38
ATOM	300	N	ALA	A	42	6.485	5.856	-4.136	1.00	11.65
ATOM	301	CA	ALA	A	42	6.322	7.235	-4.575	1.00	11.45
ATOM	302	CB	ALA	A	42	5.093	7.881	-3.900	1.00	12.12
ATOM	303	C	ALA	A	42	7.555	8.004	-4.206	1.00	11.86
ATOM	304	O	ALA	A	42	8.337	7.574	-3.341	1.00	11.97
ATOM	305	N	ASN	A	43	7.743	9.139	-4.864	1.00	11.38
ATOM	306	CA	ASN	A	43	8.759	10.096	-4.456	1.00	11.29
ATOM	307	CB	ASN	A	43	9.150	11.013	-5.625	1.00	12.66
ATOM	308	CG	ASN	A	43	9.986	10.313	-6.670	1.00	14.01
ATOM	309	OD1	ASN	A	43	10.362	9.142	-6.535	1.00	15.75
ATOM	310	ND2	ASN	A	43	10.287	11.034	-7.742	1.00	16.59
ATOM	311	C	ASN	A	43	8.258	10.943	-3.315	1.00	11.75
ATOM	312	O	ASN	A	43	7.171	11.525	-3.378	1.00	11.94

ATOM	313	N	VAL	A	44	9.056	11.010	-2.257	1.00	11.62
ATOM	314	CA	VAL	A	44	8.733	11.826	-1.097	1.00	13.22
ATOM	315	CB	VAL	A	44	8.432	10.946	0.147	1.00	14.92
ATOM	316	CG1	VAL	A	44	7.320	9.959	-0.145	1.00	16.55
ATOM	317	CG2	VAL	A	44	9.626	10.198	0.556	1.00	15.99
ATOM	318	C	VAL	A	44	9.872	12.807	-0.849	1.00	13.51
ATOM	319	O	VAL	A	44	11.029	12.488	-1.103	1.00	13.42
ATOM	320	N	MET	A	45	9.541	13.990	-0.353	1.00	14.85
ATOM	321	CA	MET	A	45	10.532	15.017	-0.131	1.00	15.75
ATOM	322	CB	MET	A	45	9.877	16.407	-0.141	1.00	16.60
ATOM	323	CG	MET	A	45	9.275	16.835	-1.464	1.00	18.19
ATOM	324	SD	MET	A	45	10.418	16.742	-2.826	1.00	21.74
ATOM	325	CE	MET	A	45	11.537	18.088	-2.504	1.00	21.52
ATOM	326	C	MET	A	45	11.184	14.817	1.223	1.00	16.61
ATOM	327	O	MET	A	45	10.500	14.803	2.243	1.00	16.90
ATOM	328	N	VAL	A	46	12.506	14.689	1.220	1.00	16.86
ATOM	329	CA	VAL	A	46	13.286	14.637	2.455	1.00	18.05
ATOM	330	CB	VAL	A	46	13.720	13.181	2.816	1.00	18.32
ATOM	331	CG1	VAL	A	46	14.477	13.146	4.150	1.00	19.56
ATOM	332	CG2	VAL	A	46	12.501	12.288	2.910	1.00	18.50
ATOM	333	C	VAL	A	46	14.496	15.537	2.257	1.00	18.60
ATOM	334	O	VAL	A	46	15.222	15.388	1.288	1.00	18.55
ATOM	335	N	ASP	A	47	14.688	16.473	3.182	1.00	20.72
ATOM	336	CA	ASP	A	47	15.862	17.352	3.181	1.00	22.66
ATOM	337	CB	ASP	A	47	17.130	16.577	3.610	1.00	23.69
ATOM	338	CG	ASP	A	47	16.997	15.961	5.012	1.00	25.31
ATOM	339	OD1	ASP	A	47	16.347	16.575	5.881	1.00	28.40
ATOM	340	OD2	ASP	A	47	17.535	14.861	5.241	1.00	27.71
ATOM	341	C	ASP	A	47	16.010	18.073	1.832	1.00	23.13
ATOM	342	O	ASP	A	47	17.114	18.237	1.308	1.00	24.89
ATOM	343	N	GLY	A	48	14.866	18.474	1.274	1.00	23.26
ATOM	344	CA	GLY	A	48	14.804	19.254	0.030	1.00	23.23
ATOM	345	C	GLY	A	48	14.817	18.500	-1.290	1.00	22.89
ATOM	346	O	GLY	A	48	14.720	19.114	-2.359	1.00	23.43
ATOM	347	N	LYS	A	49	14.931	17.174	-1.225	1.00	21.63
ATOM	348	CA	LYS	A	49	15.128	16.356	-2.417	1.00	20.77
ATOM	349	CB	LYS	A	49	16.524	15.704	-2.387	1.00	21.96
ATOM	350	CG	LYS	A	49	17.693	16.691	-2.355	1.00	25.00
ATOM	351	CD	LYS	A	49	17.878	17.387	-3.687	1.00	28.52
ATOM	352	CE	LYS	A	49	18.984	18.427	-3.611	1.00	31.72
ATOM	353	NZ	LYS	A	49	19.420	18.893	-4.957	1.00	32.72
ATOM	354	C	LYS	A	49	14.071	15.257	-2.519	1.00	18.40
ATOM	355	O	LYS	A	49	13.704	14.664	-1.506	1.00	18.26
ATOM	356	N	PRO	A	50	13.598	14.964	-3.737	1.00	16.50
ATOM	357	CA	PRO	A	50	12.664	13.852	-3.850	1.00	16.28
ATOM	358	CB	PRO	A	50	12.025	14.056	-5.224	1.00	16.17
ATOM	359	CG	PRO	A	50	13.039	14.829	-6.022	1.00	16.50
ATOM	360	CD	PRO	A	50	13.882	15.601	-5.043	1.00	17.02
ATOM	361	C	PRO	A	50	13.405	12.521	-3.781	1.00	15.94
ATOM	362	O	PRO	A	50	14.446	12.339	-4.434	1.00	18.03
ATOM	363	N	VAL	A	51	12.881	11.601	-2.980	1.00	14.23
ATOM	364	CA	VAL	A	51	13.484	10.289	-2.796	1.00	13.32
ATOM	365	CB	VAL	A	51	14.018	10.128	-1.347	1.00	14.06
ATOM	366	CG1	VAL	A	51	14.630	8.741	-1.116	1.00	14.65
ATOM	367	CG2	VAL	A	51	15.039	11.224	-1.013	1.00	16.35
ATOM	368	C	VAL	A	51	12.425	9.243	-3.075	1.00	12.07
ATOM	369	O	VAL	A	51	11.325	9.309	-2.516	1.00	12.14
ATOM	370	N	ASN	A	52	12.744	8.285	-3.932	1.00	12.03
ATOM	371	CA	ASN	A	52	11.818	7.204	-4.242	1.00	12.95
ATOM	372	CB	ASN	A	52	12.243	6.548	-5.546	1.00	14.19
ATOM	373	CG	ASN	A	52	11.205	5.577	-6.094	1.00	17.09
ATOM	374	OD1	ASN	A	52	10.051	5.567	-5.665	1.00	22.79
ATOM	375	ND2	ASN	A	52	11.606	4.783	-7.073	1.00	21.92
ATOM	376	C	ASN	A	52	11.778	6.192	-3.087	1.00	12.56
ATOM	377	O	ASN	A	52	12.827	5.616	-2.720	1.00	13.33
ATOM	378	N	LEU	A	53	10.601	6.057	-2.484	1.00	11.75

ATOM	379	CA	LEU	A	53	10.391	5.213	-1.318	1.00	11.92
ATOM	380	CB	LEU	A	53	9.909	6.087	-0.151	1.00	12.50
ATOM	381	CG	LEU	A	53	9.552	5.350	1.135	1.00	11.65
ATOM	382	CD1	LEU	A	53	10.788	4.694	1.770	1.00	13.65
ATOM	383	CD2	LEU	A	53	8.882	6.340	2.104	1.00	14.03
ATOM	384	C	LEU	A	53	9.375	4.114	-1.622	1.00	11.73
ATOM	385	O	LEU	A	53	8.270	4.411	-2.109	1.00	12.99
ATOM	386	N	GLY	A	54	9.734	2.860	-1.350	1.00	11.07
ATOM	387	CA	GLY	A	54	8.833	1.725	-1.562	1.00	11.07
ATOM	388	C	GLY	A	54	8.436	1.156	-0.207	1.00	10.64
ATOM	389	O	GLY	A	54	9.300	0.945	0.648	1.00	11.82
ATOM	390	N	LEU	A	55	7.151	0.919	0.002	1.00	10.59
ATOM	391	CA	LEU	A	55	6.614	0.486	1.282	1.00	10.70
ATOM	392	CB	LEU	A	55	5.548	1.462	1.760	1.00	11.17
ATOM	393	CG	LEU	A	55	5.995	2.920	1.882	1.00	12.50
ATOM	394	CD1	LEU	A	55	4.789	3.784	2.201	1.00	14.81
ATOM	395	CD2	LEU	A	55	7.030	3.045	2.943	1.00	14.06
ATOM	396	C	LEU	A	55	6.001	-0.904	1.150	1.00	9.88
ATOM	397	O	LEU	A	55	5.134	-1.143	0.306	1.00	10.58
ATOM	398	N	TRP	A	56	6.446	-1.790	2.037	1.00	10.18
ATOM	399	CA	TRP	A	56	6.061	-3.200	2.047	1.00	10.73
ATOM	400	CB	TRP	A	56	7.266	-4.100	1.740	1.00	12.68
ATOM	401	CG	TRP	A	56	7.846	-3.871	0.371	1.00	13.71
ATOM	402	CD1	TRP	A	56	8.413	-2.719	-0.105	1.00	15.24
ATOM	403	NE1	TRP	A	56	8.836	-2.897	-1.391	1.00	17.10
ATOM	404	CE2	TRP	A	56	8.578	-4.193	-1.757	1.00	16.99
ATOM	405	CD2	TRP	A	56	7.961	-4.831	-0.658	1.00	19.16
ATOM	406	CE3	TRP	A	56	7.604	-6.180	-0.763	1.00	20.14
ATOM	407	CZ3	TRP	A	56	7.868	-6.852	-1.949	1.00	19.43
ATOM	408	CH2	TRP	A	56	8.489	-6.193	-3.031	1.00	18.67
ATOM	409	CZ2	TRP	A	56	8.853	-4.868	-2.955	1.00	18.68
ATOM	410	C	TRP	A	56	5.470	-3.549	3.410	1.00	10.65
ATOM	411	O	TRP	A	56	6.201	-3.684	4.391	1.00	11.81
ATOM	412	N	ASP	A	57	4.172	-3.756	3.441	1.00	10.06
ATOM	413	CA	ASP	A	57	3.465	-4.187	4.637	1.00	9.81
ATOM	414	CB	ASP	A	57	1.994	-3.820	4.461	1.00	10.02
ATOM	415	CG	ASP	A	57	1.179	-3.832	5.749	1.00	9.84
ATOM	416	OD1	ASP	A	57	1.699	-4.104	6.842	1.00	10.07
ATOM	417	OD2	ASP	A	57	-0.047	-3.614	5.611	1.00	10.00
ATOM	418	C	ASP	A	57	3.632	-5.700	4.751	1.00	10.58
ATOM	419	O	ASP	A	57	3.901	-6.383	3.766	1.00	11.80
ATOM	420	N	THR	A	58	3.486	-6.225	5.960	1.00	10.01
ATOM	421	CA	THR	A	58	3.585	-7.664	6.173	1.00	10.87
ATOM	422	CB	THR	A	58	4.964	-8.090	6.735	1.00	11.02
ATOM	423	OG1	THR	A	58	5.164	-7.536	8.053	1.00	11.24
ATOM	424	CG2	THR	A	58	6.097	-7.729	5.792	1.00	11.90
ATOM	425	C	THR	A	58	2.514	-8.075	7.151	1.00	10.83
ATOM	426	O	THR	A	58	1.956	-7.241	7.867	1.00	11.29
ATOM	427	N	ALA	A	59	2.239	-9.372	7.212	1.00	11.35
ATOM	428	CA	ALA	A	59	1.329	-9.920	8.199	1.00	11.95
ATOM	429	CB	ALA	A	59	0.386	-10.932	7.536	1.00	13.29
ATOM	430	C	ALA	A	59	2.173	-10.583	9.284	1.00	11.57
ATOM	431	O	ALA	A	59	2.974	-11.495	8.974	1.00	11.96
ATOM	432	N	GLY	A	60	2.009	-10.130	10.526	1.00	11.72
ATOM	433	CA	GLY	A	60	2.897	-10.527	11.600	1.00	12.73
ATOM	434	C	GLY	A	60	2.489	-11.759	12.381	1.00	12.85
ATOM	435	O	GLY	A	60	3.301	-12.278	13.147	1.00	13.80
ATOM	436	N	GLN	A	61	1.257	-12.232	12.199	1.00	12.05
ATOM	437	CA	GLN	A	61	0.766	-13.397	12.944	1.00	12.75
ATOM	438	CB	GLN	A	61	-0.732	-13.624	12.653	1.00	13.14
ATOM	439	CG	GLN	A	61	-1.633	-12.484	13.173	1.00	13.68
ATOM	440	CD	GLN	A	61	-1.918	-11.417	12.128	1.00	14.04
ATOM	441	OE1	GLN	A	61	-1.040	-11.004	11.360	1.00	14.06
ATOM	442	NE2	GLN	A	61	-3.146	-10.938	12.116	1.00	15.06
ATOM	443	C	GLN	A	61	1.556	-14.631	12.593	1.00	14.06
ATOM	444	O	GLN	A	61	2.109	-14.764	11.501	1.00	13.88

ATOM	445	N	GLU	A	62	1.565	-15.564	13.546	1.00	14.98
ATOM	446	CA	GLU	A	62	2.299	-16.825	13.410	1.00	17.27
ATOM	447	CB	GLU	A	62	2.075	-17.709	14.636	1.00	17.00
ATOM	448	CG	GLU	A	62	2.893	-18.988	14.598	1.00	21.33
ATOM	449	CD	GLU	A	62	2.667	-19.900	15.785	1.00	21.12
ATOM	450	OE1	GLU	A	62	1.627	-19.786	16.462	1.00	25.59
ATOM	451	OE2	GLU	A	62	3.548	-20.752	16.008	1.00	25.60
ATOM	452	C	GLU	A	62	1.953	-17.592	12.147	1.00	16.12
ATOM	453	O	GLU	A	62	2.821	-18.207	11.536	1.00	15.90
ATOM	454	N	ASP	A	63	0.686	-17.548	11.740	1.00	16.37
ATOM	455	CA	ASP	A	63	0.260	-18.221	10.511	1.00	17.61
ATOM	456	CB	ASP	A	63	-1.216	-17.943	10.232	1.00	19.31
ATOM	457	CG	ASP	A	63	-2.141	-18.963	10.861	1.00	24.06
ATOM	458	OD1	ASP	A	63	-1.672	-20.032	11.306	1.00	28.11
ATOM	459	OD2	ASP	A	63	-3.357	-18.680	10.879	1.00	29.29
ATOM	460	C	ASP	A	63	1.048	-17.822	9.269	1.00	15.86
ATOM	461	O	ASP	A	63	1.110	-18.562	8.302	1.00	16.35
ATOM	462	N	TYR	A	64	1.621	-16.614	9.283	1.00	14.65
ATOM	463	CA	TYR	A	64	2.334	-16.086	8.127	1.00	13.33
ATOM	464	CB	TYR	A	64	1.964	-14.616	7.880	1.00	12.64
ATOM	465	CG	TYR	A	64	0.479	-14.387	7.710	1.00	12.50
ATOM	466	CD1	TYR	A	64	-0.148	-14.596	6.492	1.00	12.77
ATOM	467	CE1	TYR	A	64	-1.531	-14.375	6.350	1.00	14.98
ATOM	468	CZ	TYR	A	64	-2.258	-13.947	7.443	1.00	13.61
ATOM	469	OH	TYR	A	64	-3.606	-13.709	7.330	1.00	15.56
ATOM	470	CE2	TYR	A	64	-1.629	-13.728	8.649	1.00	13.71
ATOM	471	CD2	TYR	A	64	-0.285	-13.953	8.774	1.00	12.14
ATOM	472	C	TYR	A	64	3.837	-16.189	8.260	1.00	13.33
ATOM	473	O	TYR	A	64	4.563	-15.686	7.415	1.00	13.05
ATOM	474	N	ASP	A	65	4.306	-16.880	9.300	1.00	13.58
ATOM	475	CA	ASP	A	65	5.748	-16.919	9.584	1.00	14.85
ATOM	476	CB	ASP	A	65	6.018	-17.727	10.861	1.00	15.32
ATOM	477	CG	ASP	A	65	5.757	-16.951	12.136	1.00	16.90
ATOM	478	OD1	ASP	A	65	5.384	-15.753	12.094	1.00	16.97
ATOM	479	OD2	ASP	A	65	5.906	-17.558	13.226	1.00	20.20
ATOM	480	C	ASP	A	65	6.570	-17.534	8.450	1.00	15.24
ATOM	481	O	ASP	A	65	7.710	-17.143	8.218	1.00	15.20
ATOM	482	N	ARG	A	66	6.003	-18.507	7.739	1.00	15.03
ATOM	483	CA	ARG	A	66	6.718	-19.130	6.623	1.00	15.74
ATOM	484	CB	ARG	A	66	6.162	-20.530	6.348	1.00	16.44
ATOM	485	CG	ARG	A	66	6.338	-21.464	7.517	1.00	17.39
ATOM	486	CD	ARG	A	66	5.662	-22.807	7.250	1.00	20.96
ATOM	487	NE	ARG	A	66	6.389	-23.591	6.258	1.00	22.46
ATOM	488	CZ	ARG	A	66	7.469	-24.325	6.534	1.00	23.29
ATOM	489	NH1	ARG	A	66	7.945	-24.376	7.776	1.00	23.36
ATOM	490	NH2	ARG	A	66	8.070	-25.000	5.568	1.00	26.82
ATOM	491	C	ARG	A	66	6.700	-18.302	5.335	1.00	16.07
ATOM	492	O	ARG	A	66	7.597	-18.415	4.503	1.00	17.45
ATOM	493	N	LEU	A	67	5.698	-17.570	5.190	1.00	15.40
ATOM	494	CA	LEU	A	67	5.552	-16.810	3.945	1.00	15.49
ATOM	495	CB	LEU	A	67	4.078	-16.706	3.580	1.00	15.84
ATOM	496	CG	LEU	A	67	3.702	-15.969	2.291	1.00	17.97
ATOM	497	CD1	LEU	A	67	4.246	-16.739	1.095	1.00	17.77
ATOM	498	CD2	LEU	A	67	2.207	-15.856	2.216	1.00	18.43
ATOM	499	C	LEU	A	67	6.174	-15.422	4.023	1.00	15.73
ATOM	500	O	LEU	A	67	6.766	-14.940	3.057	1.00	15.69
ATOM	501	N	ARG	A	68	6.056	-14.807	5.202	1.00	16.73
ATOM	502	CA	ARG	A	68	6.505	-13.442	5.402	1.00	17.12
ATOM	503	CB	ARG	A	68	6.233	-13.038	6.851	1.00	17.50
ATOM	504	CG	ARG	A	68	6.684	-11.644	7.216	1.00	14.25
ATOM	505	CD	ARG	A	68	6.307	-11.334	8.660	1.00	14.22
ATOM	506	NE	ARG	A	68	6.888	-12.277	9.614	1.00	15.63
ATOM	507	CZ	ARG	A	68	6.225	-13.190	10.317	1.00	13.45
ATOM	508	NH1	ARG	A	68	4.893	-13.350	10.240	1.00	12.81
ATOM	509	NH2	ARG	A	68	6.923	-13.947	11.139	1.00	15.55
ATOM	510	C	ARG	A	68	7.959	-13.205	4.964	1.00	17.61

ATOM	511	O	ARG	A	68	8.256	-12.166	4.383	1.00	17.22
ATOM	512	N	PRO	A	69	8.877	-14.153	5.209	1.00	17.60
ATOM	513	CA	PRO	A	69	10.254	-13.896	4.771	1.00	18.02
ATOM	514	CB	PRO	A	69	11.036	-15.080	5.357	1.00	18.83
ATOM	515	CG	PRO	A	69	10.180	-15.514	6.539	1.00	19.69
ATOM	516	CD	PRO	A	69	8.793	-15.412	5.977	1.00	18.13
ATOM	517	C	PRO	A	69	10.481	-13.723	3.251	1.00	17.20
ATOM	518	O	PRO	A	69	11.544	-13.234	2.850	1.00	17.02
ATOM	519	N	LEU	A	70	9.503	-14.098	2.414	1.00	16.34
ATOM	520	CA	LEU	A	70	9.606	-13.778	0.985	1.00	15.96
ATOM	521	CB	LEU	A	70	8.468	-14.410	0.185	1.00	16.34
ATOM	522	CG	LEU	A	70	8.755	-15.829	-0.321	1.00	19.86
ATOM	523	CD1	LEU	A	70	8.815	-16.809	0.828	1.00	22.48
ATOM	524	CD2	LEU	A	70	7.669	-16.218	-1.308	1.00	21.49
ATOM	525	C	LEU	A	70	9.608	-12.276	0.731	1.00	14.54
ATOM	526	O	LEU	A	70	9.965	-11.853	-0.361	1.00	15.20
ATOM	527	N	SER	A	71	9.188	-11.487	1.722	1.00	13.51
ATOM	528	CA	SER	A	71	9.200	-10.014	1.624	1.00	14.79
ATOM	529	CB	SER	A	71	8.211	-9.408	2.624	1.00	14.76
ATOM	530	OG	SER	A	71	6.920	-9.946	2.432	1.00	19.09
ATOM	531	C	SER	A	71	10.550	-9.367	1.891	1.00	13.58
ATOM	532	O	SER	A	71	10.753	-8.216	1.533	1.00	13.44
ATOM	533	N	TYR	A	72	11.469	-10.086	2.540	1.00	13.81
ATOM	534	CA	TYR	A	72	12.729	-9.472	2.994	1.00	13.60
ATOM	535	CB	TYR	A	72	13.329	-10.252	4.171	1.00	14.71
ATOM	536	CG	TYR	A	72	12.553	-10.168	5.464	1.00	12.94
ATOM	537	CD1	TYR	A	72	12.266	-8.937	6.064	1.00	13.82
ATOM	538	CE1	TYR	A	72	11.587	-8.877	7.270	1.00	13.38
ATOM	539	CZ	TYR	A	72	11.183	-10.045	7.884	1.00	12.61
ATOM	540	OH	TYR	A	72	10.514	-9.929	9.067	1.00	15.65
ATOM	541	CE2	TYR	A	72	11.462	-11.272	7.320	1.00	13.90
ATOM	542	CD2	TYR	A	72	12.163	-11.316	6.122	1.00	12.90
ATOM	543	C	TYR	A	72	13.852	-9.223	1.969	1.00	14.07
ATOM	544	O	TYR	A	72	14.681	-8.359	2.209	1.00	14.07
ATOM	545	N	PRO	A	73	13.950	-10.022	0.885	1.00	15.01
ATOM	546	CA	PRO	A	73	15.057	-9.728	-0.024	1.00	15.59
ATOM	547	CB	PRO	A	73	14.766	-10.640	-1.217	1.00	15.64
ATOM	548	CG	PRO	A	73	14.149	-11.831	-0.575	1.00	17.21
ATOM	549	CD	PRO	A	73	13.205	-11.212	0.435	1.00	16.17
ATOM	550	C	PRO	A	73	15.148	-8.278	-0.458	1.00	15.42
ATOM	551	O	PRO	A	73	14.136	-7.634	-0.763	1.00	15.01
ATOM	552	N	GLN	A	74	16.376	-7.776	-0.484	1.00	14.91
ATOM	553	CA	GLN	A	74	16.687	-6.420	-0.946	1.00	15.51
ATOM	554	CB	GLN	A	74	16.264	-6.223	-2.408	1.00	16.44
ATOM	555	CG	GLN	A	74	16.882	-7.245	-3.386	1.00	19.27
ATOM	556	CD	GLN	A	74	15.910	-8.340	-3.803	1.00	24.79
ATOM	557	OE1	GLN	A	74	14.722	-8.088	-4.031	1.00	27.77
ATOM	558	NE2	GLN	A	74	16.414	-9.564	-3.926	1.00	26.38
ATOM	559	C	GLN	A	74	16.139	-5.310	-0.045	1.00	14.57
ATOM	560	O	GLN	A	74	16.047	-4.151	-0.436	1.00	15.13
ATOM	561	N	THR	A	75	15.809	-5.655	1.186	1.00	13.90
ATOM	562	CA	THR	A	75	15.355	-4.625	2.129	1.00	13.47
ATOM	563	CB	THR	A	75	14.808	-5.261	3.430	1.00	13.21
ATOM	564	OG1	THR	A	75	13.662	-6.056	3.114	1.00	13.74
ATOM	565	CG2	THR	A	75	14.403	-4.203	4.454	1.00	13.03
ATOM	566	C	THR	A	75	16.460	-3.634	2.453	1.00	13.15
ATOM	567	O	THR	A	75	17.611	-4.037	2.727	1.00	14.79
ATOM	568	N	ASP	A	76	16.123	-2.347	2.456	1.00	11.58
ATOM	569	CA	ASP	A	76	17.074	-1.281	2.806	1.00	11.85
ATOM	570	CB	ASP	A	76	16.955	-0.108	1.828	1.00	11.89
ATOM	571	CG	ASP	A	76	17.315	-0.498	0.424	1.00	14.34
ATOM	572	OD1	ASP	A	76	18.452	-0.984	0.219	1.00	17.00
ATOM	573	OD2	ASP	A	76	16.471	-0.378	-0.493	1.00	13.48
ATOM	574	C	ASP	A	76	16.887	-0.771	4.228	1.00	12.51
ATOM	575	O	ASP	A	76	17.849	-0.324	4.846	1.00	13.24
ATOM	576	N	VAL	A	77	15.639	-0.758	4.737	1.00	11.39

ATOM	577	CA	VAL	A	77	15.361	-0.284	6.085	1.00	11.27
ATOM	578	CB	VAL	A	77	15.200	1.263	6.135	1.00	10.92
ATOM	579	CG1	VAL	A	77	14.118	1.771	5.187	1.00	12.49
ATOM	580	CG2	VAL	A	77	14.955	1.766	7.562	1.00	12.84
ATOM	581	C	VAL	A	77	14.122	-1.007	6.562	1.00	11.19
ATOM	582	O	VAL	A	77	13.236	-1.321	5.747	1.00	10.58
ATOM	583	N	SER	A	78	14.079	-1.311	7.848	0.50	11.65
ATOM	584	CA	SER	A	78	12.903	-1.976	8.404	0.50	12.31
ATOM	585	CB	SER	A	78	13.223	-3.368	8.931	0.50	13.27
ATOM	586	OG	ASER	A	78	12.034	-4.011	9.347	0.50	14.27
ATOM	586	OG	BSER	A	78	13.799	-4.229	7.961	0.50	14.62
ATOM	587	C	SER	A	78	12.326	-1.155	9.534	0.50	12.86
ATOM	588	O	SER	A	78	13.053	-0.618	10.368	0.50	12.62
ATOM	589	N	LEU	A	79	11.006	-1.103	9.591	1.00	11.00
ATOM	590	CA	LEU	A	79	10.301	-0.542	10.721	1.00	11.67
ATOM	591	CB	LEU	A	79	9.103	0.295	10.283	1.00	11.27
ATOM	592	CG	LEU	A	79	9.342	1.390	9.249	1.00	13.99
ATOM	593	CD1	LEU	A	79	8.087	2.201	9.048	1.00	15.38
ATOM	594	CD2	LEU	A	79	10.467	2.331	9.646	1.00	15.86
ATOM	595	C	LEU	A	79	9.782	-1.691	11.543	1.00	10.67
ATOM	596	O	LEU	A	79	8.967	-2.466	11.060	1.00	12.30
ATOM	597	N	ILE	A	80	10.265	-1.817	12.772	1.00	11.03
ATOM	598	CA	ILE	A	80	9.671	-2.773	13.715	1.00	10.63
ATOM	599	CB	ILE	A	80	10.707	-3.483	14.588	1.00	10.95
ATOM	600	CG1	ILE	A	80	11.647	-4.331	13.716	1.00	11.97
ATOM	601	CD1	ILE	A	80	12.768	-5.045	14.503	1.00	12.36
ATOM	602	CG2	ILE	A	80	9.982	-4.373	15.630	1.00	11.76
ATOM	603	C	ILE	A	80	8.685	-2.013	14.569	1.00	10.30
ATOM	604	O	ILE	A	80	9.041	-1.054	15.245	1.00	10.00
ATOM	605	N	CYS	A	81	7.416	-2.406	14.484	1.00	9.99
ATOM	606	CA	CYS	A	81	6.332	-1.674	15.116	1.00	10.50
ATOM	607	CB	CYS	A	81	5.191	-1.489	14.115	1.00	10.19
ATOM	608	SG	CYS	A	81	5.666	-0.559	12.649	1.00	11.71
ATOM	609	C	CYS	A	81	5.777	-2.453	16.303	1.00	10.21
ATOM	610	O	CYS	A	81	5.626	-3.683	16.239	1.00	9.95
ATOM	611	N	PHE	A	82	5.382	-1.727	17.344	1.00	9.79
ATOM	612	CA	PHE	A	82	4.606	-2.301	18.446	1.00	10.00
ATOM	613	CB	PHE	A	82	5.527	-2.694	19.607	1.00	10.97
ATOM	614	CG	PHE	A	82	6.103	-1.504	20.336	1.00	10.42
ATOM	615	CD1	PHE	A	82	5.450	-0.961	21.425	1.00	10.28
ATOM	616	CE1	PHE	A	82	5.961	0.171	22.084	1.00	11.10
ATOM	617	CZ	PHE	A	82	7.121	0.775	21.612	1.00	11.17
ATOM	618	CE2	PHE	A	82	7.776	0.258	20.518	1.00	11.35
ATOM	619	CD2	PHE	A	82	7.267	-0.869	19.868	1.00	11.50
ATOM	620	C	PHE	A	82	3.620	-1.232	18.880	1.00	10.12
ATOM	621	O	PHE	A	82	3.858	-0.030	18.717	1.00	10.63
ATOM	622	N	SER	A	83	2.498	-1.656	19.437	1.00	10.02
ATOM	623	CA	SER	A	83	1.511	-0.707	19.921	1.00	10.02
ATOM	624	CB	SER	A	83	0.147	-1.362	19.964	1.00	10.25
ATOM	625	OG	SER	A	83	-0.717	-0.549	20.740	1.00	11.09
ATOM	626	C	SER	A	83	1.883	-0.201	21.306	1.00	10.65
ATOM	627	O	SER	A	83	2.181	-1.004	22.203	1.00	10.75
ATOM	628	N	LEU	A	84	1.821	1.118	21.476	1.00	10.96
ATOM	629	CA	LEU	A	84	2.132	1.734	22.768	1.00	11.88
ATOM	630	CB	LEU	A	84	2.192	3.255	22.657	1.00	13.00
ATOM	631	CG	LEU	A	84	3.389	3.776	21.852	1.00	12.58
ATOM	632	CD1	LEU	A	84	3.174	5.194	21.409	1.00	14.70
ATOM	633	CD2	LEU	A	84	4.675	3.653	22.610	1.00	12.21
ATOM	634	C	LEU	A	84	1.163	1.294	23.853	1.00	11.76
ATOM	635	O	LEU	A	84	1.446	1.479	25.047	1.00	13.30
ATOM	636	N	VAL	A	85	0.034	0.708	23.458	1.00	11.66
ATOM	637	CA	VAL	A	85	-0.939	0.222	24.448	1.00	12.51
ATOM	638	CB	VAL	A	85	-2.310	0.951	24.330	1.00	12.44
ATOM	639	CG1	VAL	A	85	-2.134	2.448	24.624	1.00	14.58
ATOM	640	CG2	VAL	A	85	-2.968	0.733	22.987	1.00	13.59
ATOM	641	C	VAL	A	85	-1.073	-1.298	24.433	1.00	12.30

ATOM	642	O	VAL	A	85	-2.069	-1.842	24.915	1.00	13.43
ATOM	643	N	SER	A	86	-0.040	-1.981	23.926	1.00	12.15
ATOM	644	CA	SER	A	86	0.044	-3.431	24.003	1.00	12.42
ATOM	645	CB	SER	A	86	-0.314	-4.125	22.683	1.00	12.04
ATOM	646	OG	SER	A	86	-0.086	-5.522	22.842	1.00	15.41
ATOM	647	C	SER	A	86	1.433	-3.844	24.475	1.00	12.50
ATOM	648	O	SER	A	86	2.351	-4.063	23.678	1.00	12.54
ATOM	649	N	PRO	A	87	1.598	-4.002	25.787	1.00	12.95
ATOM	650	CA	PRO	A	87	2.869	-4.514	26.290	1.00	13.29
ATOM	651	CB	PRO	A	87	2.627	-4.615	27.808	1.00	14.21
ATOM	652	CG	PRO	A	87	1.645	-3.503	28.077	1.00	13.86
ATOM	653	CD	PRO	A	87	0.686	-3.627	26.891	1.00	13.43
ATOM	654	C	PRO	A	87	3.327	-5.837	25.679	1.00	13.26
ATOM	655	O	PRO	A	87	4.533	-6.031	25.474	1.00	13.67
ATOM	656	N	ALA	A	88	2.391	-6.731	25.355	1.00	13.08
ATOM	657	CA	ALA	A	88	2.770	-7.984	24.699	1.00	13.15
ATOM	658	CB	ALA	A	88	1.537	-8.871	24.492	1.00	14.24
ATOM	659	C	ALA	A	88	3.470	-7.721	23.363	1.00	12.54
ATOM	660	O	ALA	A	88	4.464	-8.363	23.019	1.00	13.00
ATOM	661	N	SER	A	89	2.945	-6.767	22.595	1.00	12.21
ATOM	662	CA	SER	A	89	3.536	-6.455	21.290	1.00	12.15
ATOM	663	CB	SER	A	89	2.611	-5.546	20.457	1.00	12.88
ATOM	664	OG	SER	A	89	2.641	-4.174	20.869	1.00	11.46
ATOM	665	C	SER	A	89	4.922	-5.853	21.448	1.00	11.97
ATOM	666	O	SER	A	89	5.808	-6.046	20.615	1.00	11.89
ATOM	667	N	PHE	A	90	5.106	-5.126	22.540	1.00	11.56
ATOM	668	CA	PHE	A	90	6.397	-4.531	22.869	1.00	11.82
ATOM	669	CB	PHE	A	90	6.201	-3.565	24.049	1.00	12.37
ATOM	670	CG	PHE	A	90	7.438	-2.830	24.475	1.00	13.30
ATOM	671	CD1	PHE	A	90	8.122	-2.022	23.586	1.00	14.79
ATOM	672	CE1	PHE	A	90	9.243	-1.296	24.008	1.00	15.32
ATOM	673	CZ	PHE	A	90	9.664	-1.355	25.330	1.00	15.13
ATOM	674	CE2	PHE	A	90	8.993	-2.147	26.230	1.00	15.63
ATOM	675	CD2	PHE	A	90	7.865	-2.883	25.807	1.00	15.19
ATOM	676	C	PHE	A	90	7.422	-5.613	23.178	1.00	12.20
ATOM	677	O	PHE	A	90	8.534	-5.599	22.672	1.00	12.96
ATOM	678	N	GLU	A	91	7.033	-6.572	24.006	1.00	13.27
ATOM	679	CA	GLU	A	91	7.960	-7.637	24.327	1.00	14.67
ATOM	680	CB	GLU	A	91	7.403	-8.530	25.432	1.00	15.80
ATOM	681	CG	GLU	A	91	7.325	-7.839	26.801	1.00	21.18
ATOM	682	CD	GLU	A	91	8.683	-7.312	27.296	1.00	26.68
ATOM	683	OE1	GLU	A	91	8.728	-6.172	27.812	1.00	31.58
ATOM	684	OE2	GLU	A	91	9.705	-8.023	27.166	1.00	29.55
ATOM	685	C	GLU	A	91	8.341	-8.442	23.087	1.00	13.24
ATOM	686	O	GLU	A	91	9.480	-8.870	22.961	1.00	14.31
ATOM	687	N	ASN	A	92	7.402	-8.613	22.151	1.00	11.84
ATOM	688	CA	ASN	A	92	7.744	-9.293	20.909	1.00	12.41
ATOM	689	CB	ASN	A	92	6.493	-9.684	20.145	1.00	13.20
ATOM	690	CG	ASN	A	92	5.894	-10.946	20.686	1.00	15.85
ATOM	691	OD1	ASN	A	92	6.617	-11.853	21.123	1.00	17.38
ATOM	692	ND2	ASN	A	92	4.571	-11.029	20.675	1.00	17.83
ATOM	693	C	ASN	A	92	8.718	-8.579	19.986	1.00	11.68
ATOM	694	O	ASN	A	92	9.267	-9.213	19.093	1.00	12.11
ATOM	695	N	VAL	A	93	8.950	-7.282	20.203	1.00	12.44
ATOM	696	CA	VAL	A	93	9.993	-6.580	19.454	1.00	13.31
ATOM	697	CB	VAL	A	93	10.151	-5.140	19.954	1.00	13.15
ATOM	698	CG1	VAL	A	93	11.439	-4.514	19.419	1.00	13.98
ATOM	699	CG2	VAL	A	93	8.918	-4.329	19.575	1.00	13.99
ATOM	700	C	VAL	A	93	11.312	-7.329	19.630	1.00	14.01
ATOM	701	O	VAL	A	93	12.030	-7.582	18.672	1.00	15.10
ATOM	702	N	ARG	A	94	11.611	-7.711	20.867	1.00	15.11
ATOM	703	CA	ARG	A	94	12.873	-8.371	21.160	1.00	16.16
ATOM	704	CB	ARG	A	94	13.326	-7.993	22.575	1.00	17.62
ATOM	705	CG	ARG	A	94	14.632	-8.627	22.959	1.00	21.98
ATOM	706	CD	ARG	A	94	15.816	-7.740	22.632	1.00	28.79
ATOM	707	NE	ARG	A	94	16.859	-8.516	21.966	1.00	34.14

ATOM	708	CZ	ARG	A	94	18.164	-8.362	22.164	1.00	35.82
ATOM	709	NH1	ARG	A	94	18.611	-7.460	23.024	1.00	37.30
ATOM	710	NH2	ARG	A	94	19.025	-9.129	21.504	1.00	38.38
ATOM	711	C	ARG	A	94	12.741	-9.883	20.991	1.00	15.53
ATOM	712	O	ARG	A	94	13.679	-10.550	20.556	1.00	16.61
ATOM	713	N	ALA	A	95	11.579	-10.433	21.331	1.00	13.61
ATOM	714	CA	ALA	A	95	11.399	-11.878	21.304	1.00	13.39
ATOM	715	CB	ALA	A	95	10.156	-12.269	22.125	1.00	13.49
ATOM	716	C	ALA	A	95	11.315	-12.453	19.892	1.00	13.50
ATOM	717	O	ALA	A	95	11.737	-13.575	19.626	1.00	14.22
ATOM	718	N	LYS	A	96	10.747	-11.661	18.982	1.00	13.38
ATOM	719	CA	LYS	A	96	10.404	-12.148	17.652	1.00	12.97
ATOM	720	CB	LYS	A	96	8.876	-12.266	17.478	1.00	13.16
ATOM	721	CG	LYS	A	96	8.473	-12.789	16.104	1.00	14.36
ATOM	722	CD	LYS	A	96	6.995	-13.121	16.024	1.00	15.01
ATOM	723	CE	LYS	A	96	6.677	-13.705	14.652	1.00	18.16
ATOM	724	NZ	LYS	A	96	5.205	-13.960	14.468	1.00	18.60
ATOM	725	C	LYS	A	96	10.973	-11.275	16.541	1.00	12.23
ATOM	726	O	LYS	A	96	11.650	-11.780	15.618	1.00	12.58
ATOM	727	N	TRP	A	97	10.686	-9.974	16.591	1.00	11.57
ATOM	728	CA	TRP	A	97	10.922	-9.147	15.394	1.00	11.61
ATOM	729	CB	TRP	A	97	10.047	-7.883	15.409	1.00	10.66
ATOM	730	CG	TRP	A	97	8.634	-8.324	15.386	1.00	10.77
ATOM	731	CD1	TRP	A	97	7.712	-8.217	16.400	1.00	10.78
ATOM	732	NE1	TRP	A	97	6.550	-8.851	16.035	1.00	11.23
ATOM	733	CE2	TRP	A	97	6.711	-9.401	14.788	1.00	10.82
ATOM	734	CD2	TRP	A	97	8.019	-9.110	14.357	1.00	10.00
ATOM	735	CE3	TRP	A	97	8.445	-9.577	13.101	1.00	10.71
ATOM	736	CZ3	TRP	A	97	7.569	-10.294	12.328	1.00	11.15
ATOM	737	CH2	TRP	A	97	6.282	-10.600	12.796	1.00	11.59
ATOM	738	CZ2	TRP	A	97	5.830	-10.161	14.011	1.00	11.63
ATOM	739	C	TRP	A	97	12.370	-8.859	15.123	1.00	12.18
ATOM	740	O	TRP	A	97	12.834	-9.005	13.987	1.00	11.96
ATOM	741	N	TYR	A	98	13.096	-8.446	16.161	1.00	12.70
ATOM	742	CA	TYR	A	98	14.528	-8.222	16.017	1.00	14.03
ATOM	743	CB	TYR	A	98	15.122	-7.749	17.346	1.00	16.61
ATOM	744	CG	TYR	A	98	16.633	-7.603	17.329	1.00	20.30
ATOM	745	CD1	TYR	A	98	17.274	-6.824	16.355	1.00	22.77
ATOM	746	CE1	TYR	A	98	18.662	-6.703	16.330	1.00	24.51
ATOM	747	CZ	TYR	A	98	19.411	-7.347	17.290	1.00	23.70
ATOM	748	OH	TYR	A	98	20.789	-7.224	17.270	1.00	25.67
ATOM	749	CE2	TYR	A	98	18.809	-8.115	18.270	1.00	24.56
ATOM	750	CD2	TYR	A	98	17.413	-8.242	18.284	1.00	25.03
ATOM	751	C	TYR	A	98	15.272	-9.443	15.467	1.00	13.19
ATOM	752	O	TYR	A	98	15.981	-9.316	14.483	1.00	12.70
ATOM	753	N	PRO	A	99	15.129	-10.619	16.100	1.00	13.11
ATOM	754	CA	PRO	A	99	15.862	-11.748	15.514	1.00	13.92
ATOM	755	CB	PRO	A	99	15.605	-12.915	16.491	1.00	14.48
ATOM	756	CG	PRO	A	99	14.426	-12.507	17.292	1.00	14.24
ATOM	757	CD	PRO	A	99	14.448	-10.992	17.356	1.00	13.23
ATOM	758	C	PRO	A	99	15.431	-12.083	14.087	1.00	13.28
ATOM	759	O	PRO	A	99	16.284	-12.423	13.259	1.00	14.28
ATOM	760	N	GLU	A	100	14.141	-11.995	13.765	1.00	13.11
ATOM	761	CA	GLU	A	100	13.722	-12.301	12.402	1.00	12.73
ATOM	762	CB	GLU	A	100	12.195	-12.290	12.296	1.00	13.10
ATOM	763	CG	GLU	A	100	11.737	-12.685	10.893	1.00	13.83
ATOM	764	CD	GLU	A	100	10.257	-12.894	10.778	1.00	15.02
ATOM	765	OE1	GLU	A	100	9.716	-13.813	11.423	1.00	16.97
ATOM	766	OE2	GLU	A	100	9.637	-12.123	10.023	1.00	14.34
ATOM	767	C	GLU	A	100	14.306	-11.318	11.377	1.00	12.68
ATOM	768	O	GLU	A	100	14.843	-11.720	10.320	1.00	12.94
ATOM	769	N	VAL	A	101	14.219	-10.040	11.684	1.00	12.96
ATOM	770	CA	VAL	A	101	14.710	-9.022	10.761	1.00	13.65
ATOM	771	CB	VAL	A	101	14.283	-7.602	11.213	1.00	13.29
ATOM	772	CG1	VAL	A	101	14.986	-6.503	10.385	1.00	16.16
ATOM	773	CG2	VAL	A	101	12.773	-7.457	11.127	1.00	13.75

ATOM	774	C	VAL	A	101	16.220	-9.148	10.611	1.00	14.67
ATOM	775	O	VAL	A	101	16.722	-9.083	9.509	1.00	15.13
ATOM	776	N	ARG	A	102	16.932	-9.347	11.713	1.00	15.37
ATOM	777	CA	ARG	A	102	18.398	-9.427	11.659	1.00	16.75
ATOM	778	CB	ARG	A	102	18.965	-9.395	13.087	1.00	17.79
ATOM	779	CG	ARG	A	102	20.399	-8.843	13.205	1.00	21.62
ATOM	780	CD	ARG	A	102	20.645	-7.684	12.236	1.00	23.24
ATOM	781	NE	ARG	A	102	20.443	-6.323	12.755	1.00	23.20
ATOM	782	CZ	ARG	A	102	19.984	-5.315	12.006	1.00	23.24
ATOM	783	NH1	ARG	A	102	19.599	-5.551	10.751	1.00	23.30
ATOM	784	NH2	ARG	A	102	19.860	-4.088	12.512	1.00	23.94
ATOM	785	C	ARG	A	102	18.835	-10.683	10.907	1.00	16.87
ATOM	786	O	ARG	A	102	19.842	-10.671	10.199	1.00	18.30
ATOM	787	N	HIS	A	103	18.058	-11.758	11.009	1.00	16.43
ATOM	788	CA	HIS	A	103	18.377	-12.966	10.286	1.00	17.48
ATOM	789	CB	HIS	A	103	17.492	-14.125	10.723	1.00	17.59
ATOM	790	CG	HIS	A	103	17.809	-15.400	10.004	1.00	21.53
ATOM	791	ND1	HIS	A	103	16.983	-15.945	9.049	1.00	24.85
ATOM	792	CE1	HIS	A	103	17.538	-17.042	8.559	1.00	26.32
ATOM	793	NE2	HIS	A	103	18.706	-17.213	9.148	1.00	26.17
ATOM	794	CD2	HIS	A	103	18.906	-16.193	10.047	1.00	23.34
ATOM	795	C	HIS	A	103	18.249	-12.767	8.783	1.00	16.88
ATOM	796	O	HIS	A	103	19.135	-13.160	8.025	1.00	17.24
ATOM	797	N	HIS	A	104	17.139	-12.181	8.345	1.00	16.36
ATOM	798	CA	HIS	A	104	16.870	-12.052	6.925	1.00	16.24
ATOM	799	CB	HIS	A	104	15.363	-12.094	6.675	1.00	16.12
ATOM	800	CG	HIS	A	104	14.758	-13.433	6.940	1.00	15.16
ATOM	801	ND1	HIS	A	104	14.797	-14.464	6.024	1.00	17.29
ATOM	802	CE1	HIS	A	104	14.187	-15.518	6.534	1.00	16.02
ATOM	803	NE2	HIS	A	104	13.770	-15.213	7.751	1.00	16.75
ATOM	804	CD2	HIS	A	104	14.099	-13.909	8.021	1.00	16.36
ATOM	805	C	HIS	A	104	17.466	-10.812	6.294	1.00	16.71
ATOM	806	O	HIS	A	104	17.629	-10.759	5.066	1.00	17.06
ATOM	807	N	CYS	A	105	17.770	-9.821	7.128	1.00	16.06
ATOM	808	CA	CYS	A	105	18.278	-8.524	6.690	1.00	16.59
ATOM	809	CB	CYS	A	105	17.197	-7.448	6.805	1.00	15.75
ATOM	810	SG	CYS	A	105	15.584	-7.864	6.090	1.00	17.37
ATOM	811	C	CYS	A	105	19.456	-8.140	7.573	1.00	18.00
ATOM	812	O	CYS	A	105	19.388	-7.172	8.319	1.00	17.69
ATOM	813	N	PRO	A	106	20.554	-8.899	7.482	1.00	18.97
ATOM	814	CA	PRO	A	106	21.656	-8.676	8.409	1.00	20.28
ATOM	815	CB	PRO	A	106	22.677	-9.754	8.005	1.00	20.67
ATOM	816	CG	PRO	A	106	22.345	-10.089	6.604	1.00	19.73
ATOM	817	CD	PRO	A	106	20.853	-9.998	6.545	1.00	19.37
ATOM	818	C	PRO	A	106	22.299	-7.282	8.418	1.00	22.24
ATOM	819	O	PRO	A	106	22.809	-6.871	9.462	1.00	24.24
ATOM	820	N	ASN	A	107	22.276	-6.559	7.304	1.00	23.21
ATOM	821	CA	ASN	A	107	23.015	-5.292	7.241	1.00	24.66
ATOM	822	CB	ASN	A	107	23.979	-5.303	6.062	1.00	26.10
ATOM	823	CG	ASN	A	107	25.043	-6.324	6.207	1.00	27.68
ATOM	824	OD1	ASN	A	107	25.261	-7.119	5.296	1.00	29.64
ATOM	825	ND2	ASN	A	107	25.713	-6.338	7.362	1.00	29.00
ATOM	826	C	ASN	A	107	22.139	-4.056	7.158	1.00	24.72
ATOM	827	O	ASN	A	107	22.586	-2.999	6.709	1.00	26.29
ATOM	828	N	THR	A	108	20.910	-4.195	7.632	1.00	23.11
ATOM	829	CA	THR	A	108	19.828	-3.246	7.386	1.00	22.12
ATOM	830	CB	THR	A	108	18.561	-4.066	7.011	1.00	22.14
ATOM	831	OG1	THR	A	108	18.807	-4.802	5.813	1.00	25.06
ATOM	832	CG2	THR	A	108	17.318	-3.190	6.815	1.00	24.17
ATOM	833	C	THR	A	108	19.525	-2.445	8.644	1.00	19.74
ATOM	834	O	THR	A	108	19.311	-3.049	9.695	1.00	19.19
ATOM	835	N	PRO	A	109	19.470	-1.102	8.550	1.00	16.40
ATOM	836	CA	PRO	A	109	19.031	-0.347	9.722	1.00	15.60
ATOM	837	CB	PRO	A	109	19.143	1.114	9.282	1.00	15.76
ATOM	838	CG	PRO	A	109	19.124	1.064	7.757	1.00	16.60
ATOM	839	CD	PRO	A	109	19.824	-0.211	7.423	1.00	15.88

ATOM	840	C	PRO	A	109	17.592	-0.663	10.112	1.00	15.49
ATOM	841	O	PRO	A	109	16.750	-0.973	9.252	1.00	14.97
ATOM	842	N	ILE	A	110	17.340	-0.577	11.416	1.00	14.61
ATOM	843	CA	ILE	A	110	16.030	-0.790	11.999	1.00	14.49
ATOM	844	CB	ILE	A	110	16.068	-1.905	13.054	1.00	15.01
ATOM	845	CG1	ILE	A	110	16.364	-3.245	12.393	1.00	15.73
ATOM	846	CD1	ILE	A	110	16.593	-4.392	13.368	1.00	19.39
ATOM	847	CG2	ILE	A	110	14.742	-1.941	13.870	1.00	15.94
ATOM	848	C	ILE	A	110	15.584	0.466	12.677	1.00	13.80
ATOM	849	O	ILE	A	110	16.362	1.088	13.417	1.00	14.50
ATOM	850	N	ILE	A	111	14.346	0.869	12.420	1.00	13.72
ATOM	851	CA	ILE	A	111	13.734	1.920	13.209	1.00	13.09
ATOM	852	CB	ILE	A	111	13.175	3.006	12.318	1.00	13.97
ATOM	853	CG1	ILE	A	111	14.392	3.647	11.608	1.00	18.10
ATOM	854	CD1	ILE	A	111	14.161	4.925	10.924	1.00	22.57
ATOM	855	CG2	ILE	A	111	12.345	4.022	13.119	1.00	14.53
ATOM	856	C	ILE	A	111	12.653	1.279	14.075	1.00	12.70
ATOM	857	O	ILE	A	111	11.822	0.520	13.559	1.00	12.11
ATOM	858	N	LEU	A	112	12.674	1.558	15.377	1.00	12.35
ATOM	859	CA	LEU	A	112	11.647	1.028	16.283	1.00	11.30
ATOM	860	CB	LEU	A	112	12.251	0.756	17.660	1.00	12.08
ATOM	861	CG	LEU	A	112	11.241	0.181	18.651	1.00	11.67
ATOM	862	CD1	LEU	A	112	10.703	-1.171	18.185	1.00	13.05
ATOM	863	CD2	LEU	A	112	11.879	0.073	20.040	1.00	13.53
ATOM	864	C	LEU	A	112	10.535	2.043	16.383	1.00	11.41
ATOM	865	O	LEU	A	112	10.776	3.209	16.709	1.00	11.83
ATOM	866	N	VAL	A	113	9.301	1.612	16.081	1.00	11.11
ATOM	867	CA	VAL	A	113	8.166	2.517	15.997	1.00	11.25
ATOM	868	CB	VAL	A	113	7.568	2.562	14.555	1.00	11.45
ATOM	869	CG1	VAL	A	113	6.389	3.544	14.464	1.00	12.35
ATOM	870	CG2	VAL	A	113	8.653	2.966	13.533	1.00	11.84
ATOM	871	C	VAL	A	113	7.081	2.122	16.997	1.00	10.54
ATOM	872	O	VAL	A	113	6.545	1.010	16.912	1.00	10.85
ATOM	873	N	GLY	A	114	6.795	3.003	17.953	1.00	10.72
ATOM	874	CA	GLY	A	114	5.626	2.834	18.831	1.00	10.62
ATOM	875	C	GLY	A	114	4.412	3.434	18.152	1.00	11.24
ATOM	876	O	GLY	A	114	4.420	4.609	17.765	1.00	12.45
ATOM	877	N	THR	A	115	3.365	2.633	17.972	1.00	10.04
ATOM	878	CA	THR	A	115	2.156	3.038	17.267	1.00	10.46
ATOM	879	CB	THR	A	115	1.711	1.939	16.291	1.00	10.43
ATOM	880	OG1	THR	A	115	1.467	0.771	17.063	1.00	11.87
ATOM	881	CG2	THR	A	115	2.800	1.638	15.237	1.00	11.11
ATOM	882	C	THR	A	115	1.012	3.319	18.250	1.00	10.45
ATOM	883	O	THR	A	115	1.107	2.999	19.452	1.00	11.66
ATOM	884	N	LYS	A	116	-0.075	3.847	17.703	1.00	10.99
ATOM	885	CA	LYS	A	116	-1.291	4.184	18.452	1.00	11.62
ATOM	886	CB	LYS	A	116	-1.996	2.939	18.986	1.00	11.39
ATOM	887	CG	LYS	A	116	-2.250	1.895	17.905	1.00	10.85
ATOM	888	CD	LYS	A	116	-3.303	0.882	18.343	1.00	11.77
ATOM	889	CE	LYS	A	116	-3.395	-0.273	17.383	1.00	12.92
ATOM	890	NZ	LYS	A	116	-4.345	-1.319	17.904	1.00	11.97
ATOM	891	C	LYS	A	116	-0.999	5.163	19.574	1.00	12.31
ATOM	892	O	LYS	A	116	-1.557	5.062	20.670	1.00	12.60
ATOM	893	N	LEU	A	117	-0.160	6.147	19.268	1.00	12.97
ATOM	894	CA	LEU	A	117	0.210	7.158	20.249	1.00	14.06
ATOM	895	CB	LEU	A	117	1.129	8.182	19.571	1.00	13.64
ATOM	896	CG	LEU	A	117	1.521	9.443	20.352	1.00	14.58
ATOM	897	CD1	LEU	A	117	2.293	9.043	21.565	1.00	16.08
ATOM	898	CD2	LEU	A	117	2.368	10.351	19.496	1.00	16.13
ATOM	899	C	LEU	A	117	-1.014	7.851	20.851	1.00	13.77
ATOM	900	O	LEU	A	117	-0.992	8.227	22.041	1.00	15.02
ATOM	901	N	ASP	A	118	-2.055	8.024	20.047	1.00	13.75
ATOM	902	CA	ASP	A	118	-3.300	8.677	20.499	1.00	14.43
ATOM	903	CB	ASP	A	118	-4.271	8.865	19.330	1.00	14.70
ATOM	904	CG	ASP	A	118	-4.687	7.553	18.704	1.00	13.72
ATOM	905	OD1	ASP	A	118	-3.846	6.912	18.028	1.00	14.20

ATOM	906	OD2	ASP	A	118	-5.845	7.166	18.880	1.00	14.91
ATOM	907	C	ASP	A	118	-3.968	7.927	21.662	1.00	15.41
ATOM	908	O	ASP	A	118	-4.742	8.523	22.417	1.00	16.15
ATOM	909	N	LEU	A	119	-3.656	6.645	21.832	1.00	14.12
ATOM	910	CA	LEU	A	119	-4.297	5.853	22.873	1.00	14.41
ATOM	911	CB	LEU	A	119	-4.555	4.414	22.387	1.00	14.30
ATOM	912	CG	LEU	A	119	-5.555	4.218	21.247	1.00	14.14
ATOM	913	CD1	LEU	A	119	-5.721	2.728	20.924	1.00	14.99
ATOM	914	CD2	LEU	A	119	-6.917	4.854	21.569	1.00	14.15
ATOM	915	C	LEU	A	119	-3.542	5.866	24.191	1.00	15.31
ATOM	916	O	LEU	A	119	-4.098	5.437	25.213	1.00	15.58
ATOM	917	N	ARG	A	120	-2.294	6.333	24.191	1.00	15.71
ATOM	918	CA	ARG	A	120	-1.474	6.250	25.399	1.00	17.33
ATOM	919	CB	ARG	A	120	-0.005	6.632	25.178	1.00	16.86
ATOM	920	CG	ARG	A	120	0.811	6.546	26.491	1.00	16.99
ATOM	921	CD	ARG	A	120	2.296	6.428	26.273	1.00	16.57
ATOM	922	NE	ARG	A	120	2.819	7.584	25.564	1.00	17.42
ATOM	923	CZ	ARG	A	120	3.981	7.595	24.933	1.00	15.40
ATOM	924	NH1	ARG	A	120	4.746	6.503	24.897	1.00	15.17
ATOM	925	NH2	ARG	A	120	4.357	8.708	24.306	1.00	16.70
ATOM	926	C	ARG	A	120	-2.083	7.071	26.534	1.00	19.03
ATOM	927	O	ARG	A	120	-2.005	6.669	27.705	1.00	19.26
ATOM	928	N	ASP	A	121	-2.728	8.177	26.184	1.00	21.08
ATOM	929	CA	ASP	A	121	-3.414	8.964	27.211	1.00	23.85
ATOM	930	CB	ASP	A	121	-2.857	10.401	27.262	1.00	24.79
ATOM	931	CG	ASP	A	121	-3.016	11.149	25.950	1.00	28.79
ATOM	932	OD1	ASP	A	121	-2.641	12.344	25.899	1.00	33.35
ATOM	933	OD2	ASP	A	121	-3.514	10.559	24.974	1.00	32.10
ATOM	934	C	ASP	A	121	-4.953	8.886	27.147	1.00	24.43
ATOM	935	O	ASP	A	121	-5.637	9.704	27.777	1.00	26.11
ATOM	936	N	ASP	A	122	-5.477	7.881	26.427	1.00	23.95
ATOM	937	CA	ASP	A	122	-6.928	7.600	26.266	1.00	24.46
ATOM	938	CB	ASP	A	122	-7.177	6.663	25.068	1.00	24.33
ATOM	939	CG	ASP	A	122	-8.673	6.326	24.848	1.00	24.95
ATOM	940	OD1	ASP	A	122	-9.439	7.225	24.431	1.00	27.49
ATOM	941	OD2	ASP	A	122	-9.080	5.158	25.043	1.00	24.18
ATOM	942	C	ASP	A	122	-7.456	6.928	27.521	1.00	24.87
ATOM	943	O	ASP	A	122	-6.878	5.948	27.999	1.00	25.09
ATOM	944	N	LYS	A	123	-8.581	7.436	28.026	1.00	25.34
ATOM	945	CA	LYS	A	123	-9.105	7.041	29.328	1.00	25.83
ATOM	946	CB	LYS	A	123	-10.261	7.968	29.731	1.00	26.11
ATOM	947	CG	LYS	A	123	-9.831	9.405	30.013	1.00	28.80
ATOM	948	CD	LYS	A	123	-8.859	9.464	31.181	1.00	31.10
ATOM	949	CE	LYS	A	123	-8.804	10.839	31.826	1.00	33.04
ATOM	950	NZ	LYS	A	123	-8.546	10.688	33.283	1.00	33.17
ATOM	951	C	LYS	A	123	-9.525	5.585	29.406	1.00	25.13
ATOM	952	O	LYS	A	123	-9.068	4.857	30.289	1.00	25.32
ATOM	953	N	ASP	A	124	-10.366	5.152	28.467	1.00	25.18
ATOM	954	CA	ASP	A	124	-10.836	3.769	28.418	1.00	25.02
ATOM	955	CB	ASP	A	124	-11.861	3.580	27.302	1.00	25.90
ATOM	956	CG	ASP	A	124	-12.532	2.215	27.337	1.00	29.21
ATOM	957	OD1	ASP	A	124	-13.297	1.943	28.292	1.00	32.53
ATOM	958	OD2	ASP	A	124	-12.321	1.416	26.397	1.00	33.80
ATOM	959	C	ASP	A	124	-9.685	2.798	28.231	1.00	23.79
ATOM	960	O	ASP	A	124	-9.663	1.747	28.861	1.00	23.80
ATOM	961	N	THR	A	125	-8.723	3.160	27.374	1.00	22.32
ATOM	962	CA	THR	A	125	-7.581	2.290	27.125	1.00	20.76
ATOM	963	CB	THR	A	125	-6.716	2.824	25.974	1.00	20.70
ATOM	964	OG1	THR	A	125	-7.526	2.972	24.801	1.00	22.33
ATOM	965	CG2	THR	A	125	-5.602	1.832	25.674	1.00	19.54
ATOM	966	C	THR	A	125	-6.724	2.110	28.380	1.00	20.07
ATOM	967	O	THR	A	125	-6.333	0.993	28.712	1.00	18.81
ATOM	968	N	ILE	A	126	-6.459	3.207	29.085	1.00	19.39
ATOM	969	CA	ILE	A	126	-5.681	3.144	30.321	1.00	19.62
ATOM	970	CB	ILE	A	126	-5.364	4.532	30.887	1.00	19.40
ATOM	971	CG1	ILE	A	126	-4.396	5.268	29.960	1.00	19.35

ATOM	972	CD1	ILE	A	126	-4.362	6.779	30.214	1.00	22.93
ATOM	973	CG2	ILE	A	126	-4.727	4.420	32.273	1.00	21.54
ATOM	974	C	ILE	A	126	-6.419	2.285	31.363	1.00	19.93
ATOM	975	O	ILE	A	126	-5.806	1.451	32.015	1.00	19.00
ATOM	976	N	GLU	A	127	-7.738	2.483	31.467	1.00	21.26
ATOM	977	CA	GLU	A	127	-8.583	1.729	32.397	1.00	23.21
ATOM	978	CB	GLU	A	127	-10.031	2.206	32.318	1.00	23.25
ATOM	979	CG	GLU	A	127	-10.360	3.475	33.104	1.00	25.80
ATOM	980	CD	GLU	A	127	-11.848	3.834	33.054	1.00	26.81
ATOM	981	OE1	GLU	A	127	-12.633	3.114	32.387	1.00	32.68
ATOM	982	OE2	GLU	A	127	-12.233	4.835	33.697	1.00	31.64
ATOM	983	C	GLU	A	127	-8.542	0.238	32.083	1.00	22.25
ATOM	984	O	GLU	A	127	-8.324	-0.593	32.988	1.00	22.51
ATOM	985	N	LYS	A	128	-8.732	-0.099	30.801	1.00	22.02
ATOM	986	CA	LYS	A	128	-8.708	-1.499	30.345	1.00	23.13
ATOM	987	CB	LYS	A	128	-9.106	-1.614	28.864	1.00	23.41
ATOM	988	CG	LYS	A	128	-10.556	-1.981	28.639	1.00	26.63
ATOM	989	CD	LYS	A	128	-10.900	-2.054	27.154	1.00	27.10
ATOM	990	CE	LYS	A	128	-12.342	-2.503	26.951	1.00	32.09
ATOM	991	NZ	LYS	A	128	-12.819	-2.188	25.566	1.00	34.87
ATOM	992	C	LYS	A	128	-7.382	-2.201	30.607	1.00	21.64
ATOM	993	O	LYS	A	128	-7.351	-3.373	30.992	1.00	21.72
ATOM	994	N	LEU	A	129	-6.274	-1.492	30.390	1.00	19.65
ATOM	995	CA	LEU	A	129	-4.970	-2.031	30.719	1.00	18.39
ATOM	996	CB	LEU	A	129	-3.826	-1.147	30.159	1.00	17.50
ATOM	997	CG	LEU	A	129	-3.624	-1.209	28.638	1.00	17.68
ATOM	998	CD1	LEU	A	129	-2.705	-0.084	28.182	1.00	17.69
ATOM	999	CD2	LEU	A	129	-3.061	-2.566	28.208	1.00	19.43
ATOM	1000	C	LEU	A	129	-4.818	-2.218	32.220	1.00	18.22
ATOM	1001	O	LEU	A	129	-4.308	-3.243	32.668	1.00	17.93
ATOM	1002	N	LYS	A	130	-5.281	-1.239	32.999	1.00	18.30
ATOM	1003	CA	LYS	A	130	-5.138	-1.281	34.448	1.00	19.99
ATOM	1004	CB	LYS	A	130	-5.657	0.019	35.077	1.00	19.61
ATOM	1005	CG	LYS	A	130	-5.327	0.161	36.563	1.00	22.48
ATOM	1006	CD	LYS	A	130	-5.971	1.394	37.175	1.00	23.21
ATOM	1007	CE	LYS	A	130	-5.621	1.505	38.650	1.00	28.22
ATOM	1008	NZ	LYS	A	130	-6.438	2.560	39.305	1.00	31.69
ATOM	1009	C	LYS	A	130	-5.835	-2.500	35.066	1.00	19.55
ATOM	1010	O	LYS	A	130	-5.298	-3.110	35.978	1.00	20.66
ATOM	1011	N	GLU	A	131	-7.002	-2.844	34.534	1.00	19.98
ATOM	1012	CA	GLU	A	131	-7.762	-4.027	34.993	1.00	21.82
ATOM	1013	CB	GLU	A	131	-8.989	-4.225	34.127	1.00	22.37
ATOM	1014	CG	GLU	A	131	-10.065	-3.208	34.295	1.00	25.39
ATOM	1015	CD	GLU	A	131	-11.267	-3.528	33.423	1.00	27.20
ATOM	1016	OE1	GLU	A	131	-11.173	-4.443	32.570	1.00	28.48
ATOM	1017	OE2	GLU	A	131	-12.305	-2.868	33.584	1.00	32.18
ATOM	1018	C	GLU	A	131	-6.935	-5.302	34.910	1.00	22.56
ATOM	1019	O	GLU	A	131	-7.151	-6.244	35.678	1.00	22.48
ATOM	1020	N	LYS	A	132	-6.014	-5.334	33.944	1.00	23.05
ATOM	1021	CA	LYS	A	132	-5.088	-6.449	33.722	1.00	23.56
ATOM	1022	CB	LYS	A	132	-4.911	-6.675	32.218	1.00	23.83
ATOM	1023	CG	LYS	A	132	-6.192	-6.985	31.466	1.00	24.91
ATOM	1024	CD	LYS	A	132	-5.929	-7.130	29.978	1.00	27.51
ATOM	1025	CE	LYS	A	132	-7.172	-7.624	29.260	1.00	28.92
ATOM	1026	NZ	LYS	A	132	-6.986	-7.606	27.778	1.00	31.70
ATOM	1027	C	LYS	A	132	-3.712	-6.226	34.363	1.00	23.22
ATOM	1028	O	LYS	A	132	-2.764	-7.003	34.114	1.00	23.85
ATOM	1029	N	LYS	A	133	-3.600	-5.181	35.185	1.00	23.67
ATOM	1030	CA	LYS	A	133	-2.353	-4.822	35.870	1.00	23.68
ATOM	1031	CB	LYS	A	133	-1.865	-5.956	36.793	1.00	24.89
ATOM	1032	CG	LYS	A	133	-2.776	-6.265	37.989	1.00	26.71
ATOM	1033	CD	LYS	A	133	-1.990	-6.993	39.082	1.00	30.39
ATOM	1034	CE	LYS	A	133	-2.797	-7.142	40.374	1.00	32.61
ATOM	1035	NZ	LYS	A	133	-1.962	-7.691	41.489	1.00	34.46
ATOM	1036	C	LYS	A	133	-1.263	-4.440	34.857	1.00	23.39
ATOM	1037	O	LYS	A	133	-0.074	-4.726	35.058	1.00	23.85

ATOM	1038	N	LEU	A	134	-1.691	-3.794	33.774	1.00	21.93
ATOM	1039	CA	LEU	A	134	-0.784	-3.320	32.721	1.00	20.83
ATOM	1040	CB	LEU	A	134	-1.149	-3.965	31.395	1.00	20.76
ATOM	1041	CG	LEU	A	134	-1.077	-5.492	31.338	1.00	22.55
ATOM	1042	CD1	LEU	A	134	-1.670	-5.959	30.040	1.00	22.21
ATOM	1043	CD2	LEU	A	134	0.361	-5.965	31.506	1.00	23.61
ATOM	1044	C	LEU	A	134	-0.889	-1.817	32.584	1.00	20.05
ATOM	1045	O	LEU	A	134	-1.841	-1.208	33.051	1.00	19.28
ATOM	1046	N	ILE	A	135	0.086	-1.233	31.891	1.00	18.39
ATOM	1047	CA	ILE	A	135	0.162	0.198	31.682	1.00	18.18
ATOM	1048	CB	ILE	A	135	1.028	0.856	32.786	1.00	19.88
ATOM	1049	CG1	ILE	A	135	0.682	2.322	32.955	1.00	22.25
ATOM	1050	CD1	ILE	A	135	0.371	2.670	34.398	1.00	26.92
ATOM	1051	CG2	ILE	A	135	2.501	0.581	32.588	1.00	20.81
ATOM	1052	C	ILE	A	135	0.731	0.409	30.281	1.00	16.57
ATOM	1053	O	ILE	A	135	1.407	-0.483	29.770	1.00	16.98
ATOM	1054	N	PRO	A	136	0.401	1.539	29.638	1.00	15.39
ATOM	1055	CA	PRO	A	136	1.034	1.824	28.342	1.00	14.59
ATOM	1056	CB	PRO	A	136	0.468	3.184	27.968	1.00	14.59
ATOM	1057	CG	PRO	A	136	-0.852	3.252	28.667	1.00	15.57
ATOM	1058	CD	PRO	A	136	-0.598	2.568	29.986	1.00	15.78
ATOM	1059	C	PRO	A	136	2.550	1.891	28.388	1.00	15.06
ATOM	1060	O	PRO	A	136	3.148	2.161	29.436	1.00	15.72
ATOM	1061	N	ILE	A	137	3.165	1.615	27.245	1.00	13.89
ATOM	1062	CA	ILE	A	137	4.600	1.756	27.119	1.00	13.54
ATOM	1063	CB	ILE	A	137	5.107	1.026	25.857	1.00	13.19
ATOM	1064	CG1	ILE	A	137	4.725	-0.467	25.896	1.00	14.05
ATOM	1065	CD1	ILE	A	137	5.097	-1.217	27.175	1.00	16.84
ATOM	1066	CG2	ILE	A	137	6.610	1.212	25.703	1.00	13.91
ATOM	1067	C	ILE	A	137	4.959	3.237	27.048	1.00	13.44
ATOM	1068	O	ILE	A	137	4.387	4.003	26.304	1.00	13.44
ATOM	1069	N	THR	A	138	5.933	3.618	27.867	1.00	13.55
ATOM	1070	CA	THR	A	138	6.369	5.011	27.973	1.00	13.78
ATOM	1071	CB	THR	A	138	6.930	5.314	29.379	1.00	14.14
ATOM	1072	OG1	THR	A	138	8.163	4.631	29.537	1.00	15.81
ATOM	1073	CG2	THR	A	138	5.955	4.861	30.469	1.00	15.66
ATOM	1074	C	THR	A	138	7.490	5.285	26.980	1.00	13.92
ATOM	1075	O	THR	A	138	8.163	4.354	26.514	1.00	14.34
ATOM	1076	N	TYR	A	139	7.703	6.555	26.654	1.00	15.03
ATOM	1077	CA	TYR	A	139	8.784	6.909	25.761	1.00	15.03
ATOM	1078	CB	TYR	A	139	8.822	8.424	25.553	1.00	16.21
ATOM	1079	CG	TYR	A	139	9.806	8.858	24.507	1.00	17.32
ATOM	1080	CD1	TYR	A	139	9.384	9.113	23.206	1.00	18.21
ATOM	1081	CE1	TYR	A	139	10.287	9.508	22.221	1.00	18.42
ATOM	1082	CZ	TYR	A	139	11.623	9.663	22.535	1.00	19.94
ATOM	1083	OH	TYR	A	139	12.503	10.051	21.534	1.00	21.13
ATOM	1084	CE2	TYR	A	139	12.079	9.414	23.821	1.00	19.28
ATOM	1085	CD2	TYR	A	139	11.159	9.015	24.815	1.00	18.75
ATOM	1086	C	TYR	A	139	10.168	6.360	26.207	1.00	15.12
ATOM	1087	O	TYR	A	139	10.919	5.797	25.404	1.00	14.95
ATOM	1088	N	PRO	A	140	10.540	6.530	27.497	1.00	14.94
ATOM	1089	CA	PRO	A	140	11.825	5.963	27.923	1.00	15.05
ATOM	1090	CB	PRO	A	140	11.962	6.437	29.391	1.00	15.31
ATOM	1091	CG	PRO	A	140	11.130	7.672	29.439	1.00	16.93
ATOM	1092	CD	PRO	A	140	9.921	7.297	28.591	1.00	15.16
ATOM	1093	C	PRO	A	140	11.934	4.454	27.796	1.00	14.68
ATOM	1094	O	PRO	A	140	13.010	3.925	27.532	1.00	14.99
ATOM	1095	N	GLN	A	141	10.821	3.755	27.992	1.00	14.91
ATOM	1096	CA	GLN	A	141	10.831	2.321	27.795	1.00	15.32
ATOM	1097	CB	GLN	A	141	9.504	1.709	28.260	1.00	15.84
ATOM	1098	CG	GLN	A	141	9.408	1.695	29.805	1.00	17.42
ATOM	1099	CD	GLN	A	141	8.033	1.372	30.337	1.00	21.38
ATOM	1100	OE1	GLN	A	141	7.048	1.409	29.613	1.00	19.25
ATOM	1101	NE2	GLN	A	141	7.959	1.069	31.638	1.00	24.15
ATOM	1102	C	GLN	A	141	11.100	1.970	26.329	1.00	14.71
ATOM	1103	O	GLN	A	141	11.888	1.070	26.028	1.00	14.43

ATOM	1104	N	GLY	A	142	10.450	2.704	25.432	1.00	14.86
ATOM	1105	CA	GLY	A	142	10.638	2.471	23.988	1.00	14.98
ATOM	1106	C	GLY	A	142	12.078	2.752	23.598	1.00	15.62
ATOM	1107	O	GLY	A	142	12.702	1.979	22.871	1.00	15.98
ATOM	1108	N	LEU	A	143	12.633	3.843	24.124	1.00	15.83
ATOM	1109	CA	LEU	A	143	14.029	4.179	23.832	1.00	16.50
ATOM	1110	CB	LEU	A	143	14.401	5.527	24.425	1.00	17.32
ATOM	1111	CG	LEU	A	143	15.810	6.009	24.080	1.00	19.47
ATOM	1112	CD1	LEU	A	143	16.041	6.135	22.575	1.00	21.78
ATOM	1113	CD2	LEU	A	143	15.999	7.332	24.763	1.00	20.72
ATOM	1114	C	LEU	A	143	14.988	3.100	24.327	1.00	15.76
ATOM	1115	O	LEU	A	143	15.939	2.738	23.623	1.00	15.95
ATOM	1116	N	ALA	A	144	14.747	2.576	25.529	1.00	15.60
ATOM	1117	CA	ALA	A	144	15.579	1.489	26.048	1.00	15.89
ATOM	1118	CB	ALA	A	144	15.176	1.128	27.474	1.00	16.77
ATOM	1119	C	ALA	A	144	15.525	0.242	25.171	1.00	15.42
ATOM	1120	O	ALA	A	144	16.529	-0.409	24.935	1.00	15.80
ATOM	1121	N	MET	A	145	14.329	-0.085	24.695	1.00	15.04
ATOM	1122	CA	MET	A	145	14.169	-1.232	23.817	1.00	15.55
ATOM	1123	CB	MET	A	145	12.690	-1.461	23.547	1.00	15.20
ATOM	1124	CG	MET	A	145	12.399	-2.650	22.624	1.00	17.01
ATOM	1125	SD	MET	A	145	12.861	-4.261	23.305	1.00	18.66
ATOM	1126	CE	MET	A	145	11.458	-4.668	24.331	1.00	19.29
ATOM	1127	C	MET	A	145	14.920	-1.014	22.503	1.00	15.46
ATOM	1128	O	MET	A	145	15.556	-1.943	21.992	1.00	15.89
ATOM	1129	N	ALA	A	146	14.853	0.203	21.971	1.00	15.90
ATOM	1130	CA	ALA	A	146	15.560	0.543	20.729	1.00	16.14
ATOM	1131	CB	ALA	A	146	15.283	1.978	20.323	1.00	16.29
ATOM	1132	C	ALA	A	146	17.057	0.303	20.887	1.00	17.01
ATOM	1133	O	ALA	A	146	17.708	-0.245	19.997	1.00	16.87
ATOM	1134	N	LYS	A	147	17.605	0.702	22.028	1.00	18.16
ATOM	1135	CA	LYS	A	147	19.021	0.439	22.286	1.00	19.48
ATOM	1136	CB	LYS	A	147	19.467	1.115	23.577	1.00	19.86
ATOM	1137	CG	LYS	A	147	19.450	2.619	23.513	1.00	22.38
ATOM	1138	CD	LYS	A	147	19.918	3.218	24.835	1.00	25.84
ATOM	1139	CE	LYS	A	147	19.857	4.739	24.809	1.00	27.13
ATOM	1140	NZ	LYS	A	147	20.013	5.289	26.179	1.00	29.52
ATOM	1141	C	LYS	A	147	19.341	-1.050	22.364	1.00	20.06
ATOM	1142	O	LYS	A	147	20.356	-1.501	21.828	1.00	19.99
ATOM	1143	N	GLU	A	148	18.486	-1.812	23.035	1.00	20.59
ATOM	1144	CA	GLU	A	148	18.694	-3.269	23.159	1.00	22.17
ATOM	1145	CB	GLU	A	148	17.576	-3.927	23.965	1.00	22.06
ATOM	1146	CG	GLU	A	148	17.501	-3.578	25.446	1.00	24.20
ATOM	1147	CD	GLU	A	148	16.383	-4.326	26.153	1.00	25.70
ATOM	1148	OE1	GLU	A	148	16.031	-5.439	25.711	1.00	31.32
ATOM	1149	OE2	GLU	A	148	15.838	-3.800	27.146	1.00	31.68
ATOM	1150	C	GLU	A	148	18.773	-3.977	21.809	1.00	21.83
ATOM	1151	O	GLU	A	148	19.475	-4.985	21.659	1.00	23.00
ATOM	1152	N	ILE	A	149	18.024	-3.473	20.833	1.00	20.08
ATOM	1153	CA	ILE	A	149	17.978	-4.121	19.519	1.00	19.47
ATOM	1154	CB	ILE	A	149	16.519	-4.318	18.997	1.00	18.75
ATOM	1155	CG1	ILE	A	149	15.837	-2.982	18.655	1.00	18.13
ATOM	1156	CD1	ILE	A	149	14.511	-3.149	17.895	1.00	18.17
ATOM	1157	CG2	ILE	A	149	15.700	-5.096	20.019	1.00	19.10
ATOM	1158	C	ILE	A	149	18.834	-3.408	18.492	1.00	19.23
ATOM	1159	O	ILE	A	149	18.845	-3.791	17.323	1.00	20.25
ATOM	1160	N	GLY	A	150	19.558	-2.381	18.927	1.00	18.35
ATOM	1161	CA	GLY	A	150	20.482	-1.680	18.039	1.00	17.82
ATOM	1162	C	GLY	A	150	19.745	-0.943	16.930	1.00	18.01
ATOM	1163	O	GLY	A	150	20.273	-0.782	15.825	1.00	18.92
ATOM	1164	N	ALA	A	151	18.530	-0.494	17.230	1.00	16.81
ATOM	1165	CA	ALA	A	151	17.785	0.386	16.309	1.00	16.58
ATOM	1166	CB	ALA	A	151	16.365	0.610	16.814	1.00	15.51
ATOM	1167	C	ALA	A	151	18.502	1.727	16.150	1.00	17.00
ATOM	1168	O	ALA	A	151	19.131	2.215	17.105	1.00	18.28
ATOM	1169	N	VAL	A	152	18.430	2.329	14.964	1.00	16.92

ATOM	1170	CA	VAL	A	152	19.067	3.619	14.731	1.00	17.25
ATOM	1171	CB	VAL	A	152	19.465	3.825	13.238	1.00	17.70
ATOM	1172	CG1	VAL	A	152	20.419	2.722	12.762	1.00	20.29
ATOM	1173	CG2	VAL	A	152	18.251	3.882	12.357	1.00	17.61
ATOM	1174	C	VAL	A	152	18.223	4.782	15.266	1.00	17.04
ATOM	1175	O	VAL	A	152	18.718	5.881	15.513	1.00	17.43
ATOM	1176	N	LYS	A	153	16.919	4.549	15.427	1.00	16.32
ATOM	1177	CA	LYS	A	153	16.014	5.561	15.943	1.00	15.90
ATOM	1178	CB	LYS	A	153	15.424	6.391	14.792	1.00	17.62
ATOM	1179	CG	LYS	A	153	16.374	7.355	14.069	1.00	21.36
ATOM	1180	CD	LYS	A	153	16.886	8.472	14.985	1.00	26.36
ATOM	1181	CE	LYS	A	153	15.779	9.375	15.496	1.00	29.00
ATOM	1182	NZ	LYS	A	153	16.295	10.500	16.351	1.00	31.93
ATOM	1183	C	LYS	A	153	14.867	4.871	16.657	1.00	14.89
ATOM	1184	O	LYS	A	153	14.495	3.745	16.280	1.00	14.72
ATOM	1185	N	TYR	A	154	14.316	5.562	17.655	1.00	14.38
ATOM	1186	CA	TYR	A	154	13.031	5.224	18.255	1.00	13.82
ATOM	1187	CB	TYR	A	154	13.164	5.052	19.779	1.00	14.86
ATOM	1188	CG	TYR	A	154	11.808	4.874	20.433	1.00	14.08
ATOM	1189	CD1	TYR	A	154	11.018	3.754	20.142	1.00	15.57
ATOM	1190	CE1	TYR	A	154	9.763	3.594	20.695	1.00	15.86
ATOM	1191	CZ	TYR	A	154	9.267	4.569	21.547	1.00	15.10
ATOM	1192	OH	TYR	A	154	8.019	4.379	22.110	1.00	15.01
ATOM	1193	CE2	TYR	A	154	10.016	5.700	21.845	1.00	15.83
ATOM	1194	CD2	TYR	A	154	11.279	5.852	21.272	1.00	14.37
ATOM	1195	C	TYR	A	154	12.087	6.372	17.980	1.00	13.73
ATOM	1196	O	TYR	A	154	12.419	7.544	18.252	1.00	14.62
ATOM	1197	N	LEU	A	155	10.917	6.062	17.427	1.00	12.95
ATOM	1198	CA	LEU	A	155	9.902	7.055	17.068	1.00	13.09
ATOM	1199	CB	LEU	A	155	9.914	7.303	15.564	1.00	13.61
ATOM	1200	CG	LEU	A	155	11.217	7.832	14.961	1.00	15.78
ATOM	1201	CD1	LEU	A	155	11.156	7.731	13.448	1.00	16.92
ATOM	1202	CD2	LEU	A	155	11.464	9.258	15.437	1.00	16.38
ATOM	1203	C	LEU	A	155	8.512	6.596	17.458	1.00	13.31
ATOM	1204	O	LEU	A	155	8.231	5.400	17.429	1.00	13.15
ATOM	1205	N	GLU	A	156	7.647	7.540	17.819	1.00	12.73
ATOM	1206	CA	GLU	A	156	6.245	7.258	18.145	1.00	13.69
ATOM	1207	CB	GLU	A	156	5.935	7.716	19.575	1.00	13.99
ATOM	1208	CG	GLU	A	156	6.791	6.988	20.607	1.00	14.44
ATOM	1209	CD	GLU	A	156	6.538	7.410	22.041	1.00	15.10
ATOM	1210	OE1	GLU	A	156	6.088	8.555	22.252	1.00	17.29
ATOM	1211	OE2	GLU	A	156	6.845	6.584	22.916	1.00	16.33
ATOM	1212	C	GLU	A	156	5.315	7.948	17.162	1.00	13.28
ATOM	1213	O	GLU	A	156	5.608	9.056	16.667	1.00	14.15
ATOM	1214	N	CYS	A	157	4.173	7.332	16.869	1.00	12.48
ATOM	1215	CA	CYS	A	157	3.248	7.946	15.935	1.00	12.25
ATOM	1216	CB	CYS	A	157	3.685	7.653	14.491	1.00	12.28
ATOM	1217	SG	CYS	A	157	3.426	5.920	13.944	1.00	13.16
ATOM	1218	C	CYS	A	157	1.832	7.442	16.176	1.00	11.07
ATOM	1219	O	CYS	A	157	1.635	6.467	16.913	1.00	11.61
ATOM	1220	N	SER	A	158	0.883	8.110	15.530	1.00	11.86
ATOM	1221	CA	SER	A	158	-0.529	7.714	15.475	1.00	11.57
ATOM	1222	CB	SER	A	158	-1.400	8.639	16.344	1.00	11.46
ATOM	1223	OG	SER	A	158	-2.777	8.454	16.067	1.00	12.84
ATOM	1224	C	SER	A	158	-1.030	7.770	14.043	1.00	12.35
ATOM	1225	O	SER	A	158	-0.912	8.817	13.399	1.00	13.35
ATOM	1226	N	ALA	A	159	-1.565	6.671	13.515	1.00	12.23
ATOM	1227	CA	ALA	A	159	-2.230	6.704	12.210	1.00	12.91
ATOM	1228	CB	ALA	A	159	-2.548	5.283	11.741	1.00	12.63
ATOM	1229	C	ALA	A	159	-3.520	7.513	12.248	1.00	13.19
ATOM	1230	O	ALA	A	159	-3.945	8.058	11.223	1.00	13.32
ATOM	1231	N	LEU	A	160	-4.156	7.563	13.414	1.00	13.73
ATOM	1232	CA	LEU	A	160	-5.424	8.250	13.580	1.00	14.94
ATOM	1233	CB	LEU	A	160	-6.096	7.781	14.872	1.00	15.57
ATOM	1234	CG	LEU	A	160	-7.542	8.210	15.145	1.00	17.88
ATOM	1235	CD1	LEU	A	160	-8.273	7.116	15.948	1.00	19.48

ATOM	1236	CD2	LEU	A	160	-7.551	9.552	15.882	1.00	19.96
ATOM	1237	C	LEU	A	160	-5.253	9.781	13.525	1.00	14.74
ATOM	1238	O	LEU	A	160	-5.977	10.445	12.763	1.00	16.73
ATOM	1239	N	THR	A	161	-4.291	10.314	14.271	1.00	14.73
ATOM	1240	CA	THR	A	161	-4.076	11.779	14.312	1.00	15.61
ATOM	1241	CB	THR	A	161	-3.653	12.248	15.708	1.00	15.82
ATOM	1242	OG1	THR	A	161	-2.357	11.718	16.027	1.00	15.63
ATOM	1243	CG2	THR	A	161	-4.667	11.818	16.780	1.00	16.36
ATOM	1244	C	THR	A	161	-2.985	12.248	13.357	1.00	16.70
ATOM	1245	O	THR	A	161	-2.850	13.446	13.091	1.00	17.02
ATOM	1246	N	GLN	A	162	-2.203	11.288	12.859	1.00	15.83
ATOM	1247	CA	GLN	A	162	-0.997	11.511	12.035	1.00	16.95
ATOM	1248	CB	GLN	A	162	-1.294	12.308	10.764	1.00	17.58
ATOM	1249	CG	GLN	A	162	-2.244	11.634	9.840	1.00	20.45
ATOM	1250	CD	GLN	A	162	-2.105	12.156	8.438	1.00	24.07
ATOM	1251	OE1	GLN	A	162	-2.089	11.398	7.477	1.00	25.85
ATOM	1252	NE2	GLN	A	162	-1.965	13.465	8.314	1.00	23.76
ATOM	1253	C	GLN	A	162	0.189	12.105	12.785	1.00	16.79
ATOM	1254	O	GLN	A	162	1.225	12.404	12.176	1.00	16.94
ATOM	1255	N	ARG	A	163	0.055	12.309	14.093	1.00	15.75
ATOM	1256	CA	ARG	A	163	1.174	12.792	14.891	1.00	15.88
ATOM	1257	CB	ARG	A	163	0.751	12.960	16.356	1.00	17.08
ATOM	1258	CG	ARG	A	163	1.879	13.133	17.357	1.00	20.93
ATOM	1259	CD	ARG	A	163	2.435	14.511	17.436	1.00	26.06
ATOM	1260	NE	ARG	A	163	1.382	15.517	17.404	0.75	27.25
ATOM	1261	CZ	ARG	A	163	1.616	16.822	17.415	1.00	28.54
ATOM	1262	NH1	ARG	A	163	2.865	17.266	17.462	1.00	29.24
ATOM	1263	NH2	ARG	A	163	0.602	17.670	17.354	1.00	27.89
ATOM	1264	C	ARG	A	163	2.378	11.859	14.749	1.00	15.55
ATOM	1265	O	ARG	A	163	2.258	10.640	14.947	0.50	12.27
ATOM	1266	N	GLY	A	164	3.525	12.425	14.373	1.00	14.69
ATOM	1267	CA	GLY	A	164	4.763	11.665	14.200	1.00	14.63
ATOM	1268	C	GLY	A	164	4.863	10.804	12.950	1.00	13.55
ATOM	1269	O	GLY	A	164	5.899	10.197	12.702	1.00	14.55
ATOM	1270	N	LEU	A	165	3.797	10.762	12.162	1.00	13.31
ATOM	1271	CA	LEU	A	165	3.707	9.782	11.084	1.00	13.13
ATOM	1272	CB	LEU	A	165	2.261	9.658	10.595	1.00	13.76
ATOM	1273	CG	LEU	A	165	1.980	8.553	9.573	1.00	14.86
ATOM	1274	CD1	LEU	A	165	2.309	7.157	10.106	1.00	16.12
ATOM	1275	CD2	LEU	A	165	0.535	8.633	9.149	1.00	14.31
ATOM	1276	C	LEU	A	165	4.664	10.070	9.935	1.00	13.35
ATOM	1277	O	LEU	A	165	5.419	9.185	9.497	1.00	13.32
ATOM	1278	N	LYS	A	166	4.666	11.299	9.425	1.00	13.52
ATOM	1279	CA	LYS	A	166	5.577	11.628	8.331	1.00	13.76
ATOM	1280	CB	LYS	A	166	5.261	13.010	7.784	1.00	14.69
ATOM	1281	CG	LYS	A	166	3.905	13.080	7.142	1.00	15.92
ATOM	1282	CD	LYS	A	166	3.602	14.524	6.741	1.00	19.54
ATOM	1283	CE	LYS	A	166	2.328	14.673	5.953	1.00	23.03
ATOM	1284	NZ	LYS	A	166	1.129	14.415	6.801	1.00	27.33
ATOM	1285	C	LYS	A	166	7.035	11.520	8.779	1.00	13.49
ATOM	1286	O	LYS	A	166	7.892	11.110	7.995	1.00	14.34
ATOM	1287	N	THR	A	167	7.327	11.841	10.038	1.00	13.55
ATOM	1288	CA	THR	A	167	8.671	11.722	10.581	1.00	13.77
ATOM	1289	CB	THR	A	167	8.710	12.319	11.997	1.00	14.09
ATOM	1290	OG1	THR	A	167	8.464	13.729	11.886	1.00	18.21
ATOM	1291	CG2	THR	A	167	10.041	12.057	12.697	1.00	15.51
ATOM	1292	C	THR	A	167	9.211	10.297	10.528	1.00	13.82
ATOM	1293	O	THR	A	167	10.397	10.080	10.290	1.00	13.73
ATOM	1294	N	VAL	A	168	8.341	9.311	10.743	1.00	12.93
ATOM	1295	CA	VAL	A	168	8.782	7.928	10.666	1.00	13.09
ATOM	1296	CB	VAL	A	168	7.585	6.971	10.881	1.00	12.82
ATOM	1297	CG1	VAL	A	168	7.969	5.525	10.510	1.00	13.06
ATOM	1298	CG2	VAL	A	168	7.118	7.023	12.336	1.00	13.99
ATOM	1299	C	VAL	A	168	9.458	7.652	9.316	1.00	12.52
ATOM	1300	O	VAL	A	168	10.544	7.070	9.260	1.00	12.69
ATOM	1301	N	PHE	A	169	8.807	8.071	8.244	1.00	12.32

ATOM	1302	CA	PHE	A	169	9.301	7.796	6.904	1.00	12.53
ATOM	1303	CB	PHE	A	169	8.129	7.771	5.926	1.00	12.31
ATOM	1304	CG	PHE	A	169	7.176	6.663	6.252	1.00	12.91
ATOM	1305	CD1	PHE	A	169	7.521	5.355	5.938	1.00	12.66
ATOM	1306	CE1	PHE	A	169	6.701	4.291	6.307	1.00	14.90
ATOM	1307	CZ	PHE	A	169	5.530	4.540	6.973	1.00	14.20
ATOM	1308	CE2	PHE	A	169	5.187	5.847	7.321	1.00	12.87
ATOM	1309	CD2	PHE	A	169	6.002	6.901	6.949	1.00	13.39
ATOM	1310	C	PHE	A	169	10.469	8.680	6.503	1.00	13.15
ATOM	1311	O	PHE	A	169	11.369	8.225	5.805	1.00	13.31
ATOM	1312	N	ASP	A	170	10.488	9.913	7.016	1.00	13.64
ATOM	1313	CA	ASP	A	170	11.672	10.784	6.859	1.00	14.87
ATOM	1314	CB	ASP	A	170	11.524	12.080	7.651	1.00	16.28
ATOM	1315	CG	ASP	A	170	10.543	13.048	7.058	1.00	19.88
ATOM	1316	OD1	ASP	A	170	10.119	12.892	5.896	1.00	21.00
ATOM	1317	OD2	ASP	A	170	10.202	14.002	7.798	1.00	23.66
ATOM	1318	C	ASP	A	170	12.890	10.077	7.440	1.00	14.49
ATOM	1319	O	ASP	A	170	13.957	10.021	6.826	1.00	14.33
ATOM	1320	N	GLU	A	171	12.728	9.529	8.642	1.00	13.56
ATOM	1321	CA	GLU	A	171	13.831	8.893	9.342	1.00	13.86
ATOM	1322	CB	GLU	A	171	13.515	8.776	10.840	1.00	14.47
ATOM	1323	CG	GLU	A	171	13.513	10.133	11.541	1.00	16.70
ATOM	1324	CD	GLU	A	171	14.864	10.844	11.544	1.00	22.34
ATOM	1325	OE1	GLU	A	171	15.913	10.175	11.616	1.00	25.74
ATOM	1326	OE2	GLU	A	171	14.863	12.085	11.477	1.00	26.06
ATOM	1327	C	GLU	A	171	14.221	7.564	8.716	1.00	13.90
ATOM	1328	O	GLU	A	171	15.403	7.200	8.709	1.00	14.62
ATOM	1329	N	ALA	A	172	13.246	6.838	8.180	1.00	13.38
ATOM	1330	CA	ALA	A	172	13.546	5.622	7.432	1.00	13.23
ATOM	1331	CB	ALA	A	172	12.252	5.014	6.897	1.00	14.03
ATOM	1332	C	ALA	A	172	14.495	5.956	6.286	1.00	13.28
ATOM	1333	O	ALA	A	172	15.493	5.250	6.050	1.00	14.11
ATOM	1334	N	ILE	A	173	14.178	7.033	5.573	1.00	12.65
ATOM	1335	CA	ILE	A	173	14.992	7.458	4.434	1.00	13.19
ATOM	1336	CB	ILE	A	173	14.250	8.516	3.612	1.00	12.40
ATOM	1337	CG1	ILE	A	173	13.103	7.828	2.857	1.00	14.78
ATOM	1338	CD1	ILE	A	173	12.081	8.785	2.302	1.00	13.37
ATOM	1339	CG2	ILE	A	173	15.229	9.228	2.643	1.00	13.54
ATOM	1340	C	ILE	A	173	16.370	7.940	4.916	1.00	13.03
ATOM	1341	O	ILE	A	173	17.402	7.545	4.344	1.00	13.20
ATOM	1342	N	ARG	A	174	16.382	8.744	5.977	1.00	13.62
ATOM	1343	CA	ARG	A	174	17.674	9.196	6.521	1.00	14.74
ATOM	1344	CB	ARG	A	174	17.494	10.195	7.646	1.00	15.41
ATOM	1345	CG	ARG	A	174	16.882	11.488	7.168	1.00	15.94
ATOM	1346	CD	ARG	A	174	16.629	12.458	8.320	1.00	20.70
ATOM	1347	NE	ARG	A	174	16.118	13.738	7.823	1.00	22.87
ATOM	1348	CZ	ARG	A	174	15.001	14.333	8.237	1.00	26.43
ATOM	1349	NH1	ARG	A	174	14.252	13.806	9.205	1.00	29.29
ATOM	1350	NH2	ARG	A	174	14.640	15.486	7.689	1.00	29.09
ATOM	1351	C	ARG	A	174	18.557	8.055	6.972	1.00	15.31
ATOM	1352	O	ARG	A	174	19.786	8.137	6.874	1.00	16.11
ATOM	1353	N	ALA	A	175	17.955	6.974	7.463	1.00	15.52
ATOM	1354	CA	ALA	A	175	18.750	5.836	7.925	1.00	16.05
ATOM	1355	CB	ALA	A	175	17.848	4.791	8.567	1.00	16.64
ATOM	1356	C	ALA	A	175	19.527	5.214	6.784	1.00	16.94
ATOM	1357	O	ALA	A	175	20.554	4.546	6.996	1.00	17.87
ATOM	1358	N	VAL	A	176	19.023	5.398	5.570	1.00	16.33
ATOM	1359	CA	VAL	A	176	19.637	4.805	4.402	1.00	16.94
ATOM	1360	CB	VAL	A	176	18.558	4.324	3.380	1.00	16.82
ATOM	1361	CG1	VAL	A	176	19.199	3.795	2.097	1.00	17.60
ATOM	1362	CG2	VAL	A	176	17.671	3.260	4.012	1.00	17.37
ATOM	1363	C	VAL	A	176	20.608	5.811	3.770	1.00	18.02
ATOM	1364	O	VAL	A	176	21.728	5.427	3.375	1.00	19.01
ATOM	1365	N	LEU	A	177	20.209	7.085	3.716	1.00	18.56
ATOM	1366	CA	LEU	A	177	20.959	8.112	2.960	1.00	19.99
ATOM	1367	CB	LEU	A	177	19.995	9.076	2.262	1.00	19.32

ATOM	1368	CG	LEU	A	177	19.084	8.466	1.202	1.00	19.69
ATOM	1369	CD1	LEU	A	177	18.245	9.570	0.570	1.00	21.53
ATOM	1370	CD2	LEU	A	177	19.869	7.693	0.137	1.00	20.62
ATOM	1371	C	LEU	A	177	21.971	8.922	3.771	1.00	21.89
ATOM	1372	O	LEU	A	177	22.858	9.576	3.198	1.00	22.03
ATOM	1373	N	CYS	A	178	21.826	8.919	5.089	1.00	22.74
ATOM	1374	CA	CYS	A	178	22.743	9.643	5.961	1.00	25.08
ATOM	1375	CB	CYS	A	178	21.984	10.500	6.977	1.00	25.34
ATOM	1376	SG	CYS	A	178	20.806	11.678	6.228	0.75	28.16
ATOM	1377	C	CYS	A	178	23.648	8.648	6.661	1.00	26.11
ATOM	1378	O	CYS	A	178	23.186	7.875	7.499	1.00	27.77
ATOM	1379	O6	GDP	A	191	-3.009	5.022	15.369	1.00	12.24
ATOM	1380	C6	GDP	A	191	-4.015	4.359	15.678	1.00	12.17
ATOM	1381	N1	GDP	A	191	-4.860	4.763	16.631	1.00	11.86
ATOM	1382	C2	GDP	A	191	-5.946	4.059	17.010	1.00	13.33
ATOM	1383	N3	GDP	A	191	-6.335	2.908	16.446	1.00	12.94
ATOM	1384	C4	GDP	A	191	-5.563	2.390	15.462	1.00	12.01
ATOM	1385	N2	GDP	A	191	-6.728	4.592	17.968	1.00	14.08
ATOM	1386	C5	GDP	A	191	-4.355	3.101	15.012	1.00	11.72
ATOM	1387	N7	GDP	A	191	-3.778	2.354	14.043	1.00	11.60
ATOM	1388	C8	GDP	A	191	-4.554	1.260	13.880	1.00	11.74
ATOM	1389	N9	GDP	A	191	-5.622	1.293	14.713	1.00	11.13
ATOM	1390	C1*	GDP	A	191	-6.641	0.245	14.813	1.00	12.04
ATOM	1391	C2*	GDP	A	191	-7.274	-0.146	13.495	1.00	13.49
ATOM	1392	O2*	GDP	A	191	-8.363	0.720	13.221	1.00	15.11
ATOM	1393	C3*	GDP	A	191	-7.655	-1.573	13.776	1.00	13.18
ATOM	1394	O3*	GDP	A	191	-8.833	-1.618	14.593	1.00	14.69
ATOM	1395	O4*	GDP	A	191	-5.970	-0.930	15.291	1.00	11.20
ATOM	1396	C4*	GDP	A	191	-6.533	-2.077	14.643	1.00	12.56
ATOM	1397	C5*	GDP	A	191	-5.462	-2.819	13.842	1.00	10.91
ATOM	1398	O5*	GDP	A	191	-4.919	-1.976	12.814	1.00	9.53
ATOM	1399	PA	GDP	A	191	-4.141	-2.627	11.586	1.00	9.51
ATOM	1400	O1A	GDP	A	191	-3.619	-1.488	10.785	1.00	9.65
ATOM	1401	O2A	GDP	A	191	-4.994	-3.603	10.878	1.00	10.36
ATOM	1402	O3A	GDP	A	191	-2.961	-3.377	12.352	1.00	9.19
ATOM	1403	PB	GDP	A	191	-1.946	-4.469	11.696	1.00	8.75
ATOM	1404	O1B	GDP	A	191	-0.668	-4.238	12.435	1.00	8.01
ATOM	1405	O3B	GDP	A	191	-2.522	-5.850	12.023	1.00	9.28
ATOM	1406	O2B	GDP	A	191	-1.858	-4.210	10.209	1.00	9.68
ATOM	1407	MG	MG	A	192	-2.167	-5.353	8.526	1.00	8.83
ATOM	1408	CBD	DRG	A	193	0.681	-13.450	2.776	1.00	20.00
ATOM	1409	CBB	DRG	A	193	2.267	-11.782	3.301	1.00	20.00
ATOM	1410	NBA	DRG	A	193	3.593	-11.504	3.903	1.00	20.00
ATOM	1411	CBC	DRG	A	193	3.584	-11.334	5.380	1.00	20.00
ATOM	1412	CBE	DRG	A	193	2.285	-10.824	6.039	1.00	20.00
ATOM	1413	CAZ	DRG	A	193	4.146	-10.330	3.193	1.00	20.00
ATOM	1414	CAY	DRG	A	193	3.104	-9.261	2.850	1.00	20.00
ATOM	1415	CAW	DRG	A	193	3.698	-7.876	3.101	1.00	20.00
ATOM	1416	CAV	DRG	A	193	4.865	-7.544	2.157	1.00	20.00
ATOM	1417	CAX	DRG	A	193	4.462	-6.384	1.241	1.00	20.00
ATOM	1418	NAT	DRG	A	193	5.218	-8.724	1.338	1.00	20.00
ATOM	1419	C2	DRG	A	193	5.884	-8.629	0.171	1.00	20.00
ATOM	1420	N1	DRG	A	193	7.077	-9.259	-0.012	1.00	20.00
ATOM	1421	N3	DRG	A	193	5.379	-7.889	-0.848	1.00	20.00
ATOM	1422	C4	DRG	A	193	6.016	-7.768	-2.033	1.00	20.00
ATOM	1423	CAU	DRG	A	193	5.400	-6.926	-3.158	1.00	20.00
ATOM	1424	C5	DRG	A	193	7.453	-8.603	-2.141	1.00	20.00
ATOM	1425	C6	DRG	A	193	7.779	-9.175	-1.178	1.00	20.00
ATOM	1426	NAM	DRG	A	193	8.934	-9.860	-1.330	1.00	20.00
ATOM	1427	CAJ	DRG	A	193	9.976	-9.460	-2.102	1.00	20.00
ATOM	1428	CAF	DRG	A	193	10.753	-10.423	-2.779	1.00	20.00
ATOM	1429	CAB	DRG	A	193	11.751	-9.996	-3.687	1.00	20.00
ATOM	1430	CAA	DRG	A	193	12.544	-10.894	-4.439	1.00	20.00
ATOM	1431	NAD	DRG	A	193	12.440	-12.222	-4.327	1.00	20.00
ATOM	1432	CAC	DRG	A	193	13.496	-10.415	-5.348	1.00	20.00
ATOM	1433	CAG	DRG	A	193	13.635	-9.025	-5.501	1.00	20.00

ATOM	1434	CAK	DRG	A	193	14.657	-8.464	-6.495	1.00	20.00
ATOM	1435	NAH	DRG	A	193	12.880	-8.183	-4.767	1.00	20.00
ATOM	1436	CAE	DRG	A	193	11.950	-8.599	-3.881	1.00	20.00
ATOM	1437	CAI	DRG	A	193	11.197	-7.636	-3.199	1.00	20.00
ATOM	1438	CAL	DRG	A	193	10.205	-8.083	-2.316	1.00	20.00
ATOM	1439	O	HOH	Z	1	-0.134	-5.584	8.242	1.00	8.96
ATOM	1440	O	HOH	Z	2	-4.232	-5.015	8.711	1.00	9.58
ATOM	1441	O	HOH	Z	3	-2.275	-7.057	9.673	1.00	10.94
ATOM	1442	O	HOH	Z	4	10.477	-1.304	-3.031	1.00	26.08
ATOM	1443	O	HOH	Z	5	16.317	-2.369	-2.524	1.00	29.42
ATOM	1444	O	HOH	Z	6	5.047	-8.484	10.591	1.00	12.91
ATOM	1445	O	HOH	Z	7	5.124	-6.010	17.744	1.00	11.06
ATOM	1446	O	HOH	Z	8	0.567	-9.158	20.354	1.00	21.09
ATOM	1447	O	HOH	Z	9	0.703	-11.544	16.131	1.00	12.46
ATOM	1448	O	HOH	Z	10	3.703	-12.071	16.597	1.00	22.55
ATOM	1449	O	HOH	Z	11	-0.182	-8.384	10.983	1.00	11.20
ATOM	1450	O	HOH	Z	12	-4.333	-8.769	10.467	1.00	14.76
ATOM	1451	O	HOH	Z	13	-4.936	-6.912	12.473	1.00	14.22
ATOM	1452	O	HOH	Z	14	-8.445	11.206	35.892	1.00	27.24
ATOM	1453	O	HOH	Z	15	-3.210	-2.135	20.575	1.00	12.27
ATOM	1454	O	HOH	Z	16	-1.032	2.706	13.325	1.00	11.23
ATOM	1455	O	HOH	Z	17	0.215	4.566	14.883	1.00	12.32
ATOM	1456	O	HOH	Z	18	1.314	11.622	4.713	1.00	23.82
ATOM	1457	O	HOH	Z	19	0.615	11.501	7.265	1.00	29.69
ATOM	1458	O	HOH	Z	20	-14.149	1.973	10.917	1.00	22.67
ATOM	1459	O	HOH	Z	21	-11.299	-4.865	9.719	1.00	22.28
ATOM	1460	O	HOH	Z	22	-9.705	-6.203	8.035	1.00	17.69
ATOM	1461	O	HOH	Z	23	7.468	13.145	-9.245	1.00	28.76
ATOM	1462	O	HOH	Z	24	-6.991	-5.346	11.452	1.00	15.83
ATOM	1463	O	HOH	Z	25	4.514	11.481	-2.718	1.00	12.76
ATOM	1464	O	HOH	Z	26	6.818	14.498	0.538	1.00	22.98
ATOM	1465	O	HOH	Z	27	-6.619	-9.355	9.099	1.00	17.44
ATOM	1466	O	HOH	Z	28	15.357	6.087	-3.675	1.00	27.71
ATOM	1467	O	HOH	Z	29	15.754	8.383	-5.057	1.00	25.97
ATOM	1468	O	HOH	Z	30	-1.307	-5.728	19.093	1.00	27.63
ATOM	1469	O	HOH	Z	31	-0.946	-7.181	21.049	1.00	23.44
ATOM	1470	O	HOH	Z	32	-3.074	-4.762	20.234	1.00	31.01
ATOM	1471	O	HOH	Z	33	2.733	9.521	0.192	1.00	13.56
ATOM	1472	O	HOH	Z	34	3.360	9.455	-1.336	1.00	18.57
ATOM	1473	O	HOH	Z	36	11.041	-13.275	-2.389	1.00	29.42
ATOM	1474	O	HOH	Z	37	-6.223	11.555	10.296	1.00	32.37
ATOM	1475	O	HOH	Z	38	-9.612	-4.467	12.063	1.00	31.51
ATOM	1476	O	HOH	Z	44	4.098	-4.948	-0.736	1.00	26.48
ATOM	1477	O	HOH	Z	45	2.704	-3.500	0.992	1.00	25.13
ATOM	1478	O	HOH	Z	46	2.826	-1.554	1.772	1.00	32.13
ATOM	1479	O	HOH	Z	47	-5.682	11.211	-0.359	1.00	25.76
ATOM	1480	O	HOH	Z	48	-4.127	-5.680	-0.140	1.00	32.87
ATOM	1481	O	HOH	Z	49	0.270	-2.172	-5.395	1.00	28.65
ATOM	1482	O	HOH	Z	50	-2.450	-1.603	-5.615	1.00	31.17
ATOM	1483	O	HOH	Z	51	3.268	-18.749	6.483	1.00	21.70
ATOM	1484	O	HOH	Z	52	2.161	-21.956	8.428	1.00	31.44
ATOM	1485	O	HOH	Z	53	-1.571	-18.172	6.694	1.00	17.34
ATOM	1486	O	HOH	Z	54	-0.249	-18.135	5.433	1.00	29.65
ATOM	1487	O	HOH	Z	55	2.971	2.171	-6.016	1.00	23.51
ATOM	1488	O	HOH	Z	56	7.729	11.463	5.016	1.00	17.76
ATOM	1489	O	HOH	Z	57	12.155	18.376	1.959	1.00	26.19
ATOM	1490	O	HOH	Z	58	12.594	17.158	5.114	1.00	31.33
ATOM	1491	O	HOH	Z	59	-1.569	11.596	18.927	1.00	16.57
ATOM	1492	O	HOH	Z	60	7.579	9.928	14.909	1.00	15.76
ATOM	1493	O	HOH	Z	61	19.744	-0.867	13.120	1.00	18.17
ATOM	1494	O	HOH	Z	62	8.707	10.237	18.525	1.00	17.38
ATOM	1495	O	HOH	Z	63	21.457	20.231	-3.910	1.00	19.88
ATOM	1496	O	HOH	Z	64	11.940	-14.538	15.403	1.00	19.32
ATOM	1497	O	HOH	Z	65	-3.012	1.332	32.662	1.00	17.87
ATOM	1498	O	HOH	Z	66	2.933	13.528	10.444	1.00	18.09
ATOM	1499	O	HOH	Z	67	5.511	13.684	11.621	1.00	18.83

ATOM	1500	O	HOH	Z	68	-0.341	-6.697	26.164	1.00	22.76
ATOM	1501	O	HOH	Z	69	15.508	4.752	28.264	1.00	19.65
ATOM	1502	O	HOH	Z	70	9.568	-16.322	9.992	1.00	22.23
ATOM	1503	O	HOH	Z	71	15.958	7.636	18.742	1.00	23.87
ATOM	1504	O	HOH	Z	72	-6.783	0.200	18.310	1.00	21.13
ATOM	1505	O	HOH	Z	73	11.447	9.949	19.027	1.00	19.55
ATOM	1506	O	HOH	Z	74	23.144	10.431	0.689	1.00	23.34
ATOM	1507	O	HOH	Z	75	-7.996	8.406	20.020	1.00	25.41
ATOM	1508	O	HOH	Z	76	2.608	3.969	31.316	1.00	26.88
ATOM	1509	O	HOH	Z	77	2.425	-3.227	31.617	1.00	27.72
ATOM	1510	O	HOH	Z	78	17.254	-8.666	3.011	1.00	25.39
ATOM	1511	O	HOH	Z	79	12.918	-15.190	-0.608	1.00	34.94
ATOM	1512	O	HOH	Z	80	12.067	-17.714	-1.315	1.00	32.30
ATOM	1513	O	HOH	Z	81	-10.581	2.386	3.590	1.00	25.93
ATOM	1514	O	HOH	Z	82	9.735	-14.832	-4.003	1.00	32.53
ATOM	1515	O	HOH	Z	83	11.835	-6.311	-3.598	1.00	32.48
ATOM	1516	O	HOH	Z	84	8.689	11.909	16.333	1.00	23.24
ATOM	1517	O	HOH	Z	85	-6.601	-3.539	17.808	1.00	23.91
ATOM	1518	O	HOH	Z	86	17.879	3.999	27.428	1.00	28.73
ATOM	1519	O	HOH	Z	87	12.222	-16.863	9.403	1.00	23.77
ATOM	1520	O	HOH	Z	88	5.867	8.517	27.830	1.00	26.11
ATOM	1521	O	HOH	Z	89	17.094	7.930	10.841	1.00	29.44
ATOM	1522	O	HOH	Z	90	-8.791	-1.299	17.296	1.00	26.69
ATOM	1523	O	HOH	Z	91	4.804	0.800	30.972	1.00	28.06
ATOM	1524	O	HOH	Z	92	3.461	15.172	13.591	1.00	26.08
ATOM	1525	O	HOH	Z	93	-14.826	5.257	33.876	1.00	28.68
ATOM	1526	O	HOH	Z	94	10.358	-15.459	13.503	1.00	27.42
ATOM	1527	O	HOH	Z	95	21.417	4.120	9.561	1.00	25.75
ATOM	1528	O	HOH	Z	96	-1.574	-17.006	13.344	1.00	27.33
ATOM	1529	O	HOH	Z	97	3.802	-20.213	9.738	1.00	26.02
ATOM	1530	O	HOH	Z	98	4.687	-10.827	24.294	1.00	27.00
ATOM	1531	O	HOH	Z	99	18.579	-0.462	26.775	1.00	28.17
ATOM	1532	O	HOH	Z	100	9.226	4.998	32.035	1.00	26.92
ATOM	1533	O	HOH	Z	101	-0.655	6.897	29.980	1.00	28.47
ATOM	1534	O	HOH	Z	102	6.947	10.853	20.750	1.00	25.28
ATOM	1535	O	HOH	Z	103	18.739	-13.365	14.310	1.00	28.79
ATOM	1536	O	HOH	Z	104	1.875	6.190	29.907	1.00	28.27
ATOM	1537	O	HOH	Z	105	11.693	3.613	32.212	1.00	34.50
ATOM	1538	O	HOH	Z	106	-10.191	6.741	34.392	1.00	29.28
ATOM	1539	O	HOH	Z	107	-6.667	8.319	32.568	1.00	35.78
ATOM	1540	O	HOH	Z	108	22.107	-0.247	11.857	1.00	35.64
ATOM	1541	O	HOH	Z	109	5.419	11.752	18.009	1.00	33.54
ATOM	1542	O	HOH	Z	110	18.317	4.725	18.913	1.00	28.26
ATOM	1543	O	HOH	Z	111	-19.579	1.950	5.342	1.00	33.46
ATOM	1544	O	HOH	Z	112	-0.971	-9.842	17.992	1.00	26.85
ATOM	1545	O	HOH	Z	113	18.991	1.782	28.143	1.00	32.15
ATOM	1546	O	HOH	Z	114	-8.447	10.667	8.833	1.00	31.09
ATOM	1547	O	HOH	Z	115	-7.957	-5.784	38.341	1.00	33.78
ATOM	1548	O	HOH	Z	116	23.481	8.252	-0.900	1.00	31.21
ATOM	1549	O	HOH	Z	117	16.296	-11.718	2.686	1.00	25.55
ATOM	1550	O	HOH	Z	118	-8.089	6.124	32.836	1.00	33.27
ATOM	1551	O	HOH	Z	119	-18.899	1.773	9.908	1.00	28.82
ATOM	1552	O	HOH	Z	120	-4.891	11.134	22.878	1.00	34.04
ATOM	1553	O	HOH	Z	121	-3.879	-6.962	-3.653	1.00	25.64
ATOM	1554	O	HOH	Z	122	6.294	-5.334	27.716	1.00	32.59
ATOM	1555	O	HOH	Z	123	19.061	-12.417	17.639	1.00	35.78
ATOM	1556	O	HOH	Z	124	3.653	8.065	29.157	1.00	31.42
ATOM	1557	O	HOH	Z	125	20.626	-6.832	4.898	1.00	30.71
ATOM	1558	O	HOH	Z	126	-4.089	15.494	11.914	1.00	37.36
ATOM	1559	O	HOH	Z	127	-5.505	-1.483	22.141	1.00	33.40
ATOM	1560	O	HOH	Z	128	13.760	-7.726	25.645	1.00	43.49
ATOM	1561	O	HOH	Z	129	11.225	13.014	16.204	1.00	31.87
ATOM	1562	O	HOH	Z	130	-10.901	1.977	13.994	1.00	31.78
ATOM	1563	O	HOH	Z	131	-1.232	15.529	13.872	1.00	31.18
ATOM	1564	O	HOH	Z	132	-8.158	1.006	23.180	1.00	33.63
ATOM	1565	O	HOH	Z	133	3.948	-1.576	30.415	1.00	43.32

ATOM	1566	O	HOH	Z	134	-7.606	6.346	3.782	1.00	37.67
ATOM	1567	O	HOH	Z	135	20.292	-7.467	0.209	1.00	31.70
ATOM	1568	O	HOH	Z	136	20.357	7.828	26.267	1.00	35.06
ATOM	1569	O	HOH	Z	137	20.649	-0.024	1.228	1.00	31.80
ATOM	1570	O	HOH	Z	138	-0.771	10.384	23.630	1.00	32.99
ATOM	1571	O	HOH	Z	139	17.801	-7.406	25.546	1.00	33.06
ATOM	1572	O	HOH	Z	140	-4.102	-2.547	-0.245	1.00	32.75
ATOM	1573	O	HOH	Z	141	-9.581	-5.177	30.449	1.00	32.14
ATOM	1574	O	HOH	Z	142	5.765	-21.459	11.112	1.00	32.28
ATOM	1575	O	HOH	Z	143	-7.145	-10.843	13.347	1.00	38.51
END										

Table 2

REMARK Written by O version 9.0.7

CRYST1	41.792	40.008	52.071	90.00	105.83	90.00					
ORIGX1	1.000000	0.000000	0.000000		0.000000						
ORIGX2	0.000000	1.000000	0.000000		0.000000						
ORIGX3	0.000000	0.000000	1.000000		0.000000						
SCALE1	0.023928	-0.000001	0.006784		0.000000						
SCALE2	0.000000	0.024995	-0.000001		0.000000						
SCALE3	0.000000	0.000000	0.019961		0.000000						
ATOM	1	N	ALA	A	1	15.756	6.560	-10.373	1.00	30.78	N
ATOM	2	CA	ALA	A	1	16.093	6.323	-8.944	1.00	29.95	C
ATOM	3	CB	ALA	A	1	15.708	7.535	-8.107	1.00	30.54	C
ATOM	4	C	ALA	A	1	15.397	5.066	-8.426	1.00	29.34	C
ATOM	5	O	ALA	A	1	14.165	4.991	-8.415	1.00	29.88	O
ATOM	6	N	ALA	A	2	16.190	4.079	-8.013	1.00	28.31	N
ATOM	7	CA	ALA	A	2	15.653	2.839	-7.447	1.00	26.70	C
ATOM	8	CB	ALA	A	2	16.739	1.776	-7.330	1.00	27.15	C
ATOM	9	C	ALA	A	2	15.066	3.147	-6.080	1.00	25.29	C
ATOM	10	O	ALA	A	2	15.671	3.887	-5.294	1.00	25.34	O
ATOM	11	N	ALA	A	3	13.888	2.584	-5.813	1.00	23.09	N
ATOM	12	CA	ALA	A	3	13.185	2.797	-4.545	1.00	21.06	C
ATOM	13	CB	ALA	A	3	11.794	2.144	-4.577	1.00	21.37	C
ATOM	14	C	ALA	A	3	13.992	2.272	-3.360	1.00	19.51	C
ATOM	15	O	ALA	A	3	14.532	1.151	-3.409	1.00	21.10	O
ATOM	16	N	ILE	A	4	14.103	3.097	-2.322	1.00	16.92	N
ATOM	17	CA	ILE	A	4	14.495	2.610	-1.001	1.00	15.02	C
ATOM	18	CB	ILE	A	4	14.648	3.775	-0.026	1.00	14.41	C
ATOM	19	CG1	ILE	A	4	15.895	4.579	-0.429	1.00	16.51	C
ATOM	20	CD1	ILE	A	4	16.154	5.841	0.397	1.00	16.77	C
ATOM	21	CG2	ILE	A	4	14.740	3.274	1.433	1.00	15.96	C
ATOM	22	C	ILE	A	4	13.388	1.675	-0.542	1.00	14.20	C
ATOM	23	O	ILE	A	4	12.232	2.101	-0.406	1.00	15.07	O
ATOM	24	N	LYS	A	5	13.726	0.408	-0.273	1.00	12.22	N
ATOM	25	CA	LYS	A	5	12.723	-0.574	0.144	1.00	11.59	C
ATOM	26	CB	LYS	A	5	13.060	-1.966	-0.389	1.00	11.41	C
ATOM	27	CG	LYS	A	5	12.127	-3.072	0.138	1.00	11.13	C
ATOM	28	CD	LYS	A	5	12.354	-4.367	-0.588	1.00	12.02	C
ATOM	29	CE	LYS	A	5	11.520	-5.470	0.029	1.00	13.44	C
ATOM	30	NZ	LYS	A	5	11.555	-6.719	-0.805	1.00	14.02	N
ATOM	31	C	LYS	A	5	12.599	-0.604	1.665	1.00	11.00	O
ATOM	32	O	LYS	A	5	13.549	-0.951	2.376	1.00	11.63	C
ATOM	33	N	CYS	A	6	11.416	-0.249	2.160	1.00	10.99	N
ATOM	34	CA	CYS	A	6	11.150	-0.184	3.592	1.00	10.47	C
ATOM	35	CB	CYS	A	6	10.616	1.202	3.918	1.00	11.97	C
ATOM	36	SG	CYS	A	6	10.107	1.479	5.616	1.00	12.97	S
ATOM	37	C	CYS	A	6	10.108	-1.234	3.915	1.00	10.18	C
ATOM	38	O	CYS	A	6	8.997	-1.192	3.382	1.00	10.84	O
ATOM	39	N	VAL	A	7	10.453	-2.173	4.789	1.00	10.01	N
ATOM	40	CA	VAL	A	7	9.549	-3.260	5.155	1.00	9.99	C
ATOM	41	CB	VAL	A	7	10.261	-4.622	5.022	1.00	11.17	C
ATOM	42	CG1	VAL	A	7	9.362	-5.747	5.495	1.00	12.37	C
ATOM	43	CG2	VAL	A	7	10.680	-4.873	3.596	1.00	12.17	C

ATOM	44	C	VAL	A	7	9.052	-3.010	6.579	1.00	10.17	C
ATOM	45	O	VAL	A	7	9.833	-2.691	7.463	1.00	10.48	O
ATOM	46	N	VAL	A	8	7.758	-3.159	6.798	1.00	10.12	N
ATOM	47	CA	VAL	A	8	7.114	-2.817	8.057	1.00	10.09	C
ATOM	48	CB	VAL	A	8	5.893	-1.924	7.771	1.00	10.08	C
ATOM	49	CG1	VAL	A	8	5.184	-1.529	9.081	1.00	10.78	C
ATOM	50	CG2	VAL	A	8	6.324	-0.697	6.939	1.00	11.24	C
ATOM	51	C	VAL	A	8	6.658	-4.129	8.684	1.00	9.54	C
ATOM	52	O	VAL	A	8	5.908	-4.894	8.046	1.00	9.78	O
ATOM	53	N	VAL	A	9	7.144	-4.421	9.887	1.00	9.18	N
ATOM	54	CA	VAL	A	9	6.827	-5.669	10.590	1.00	9.67	C
ATOM	55	CB	VAL	A	9	8.027	-6.665	10.597	1.00	9.51	C
ATOM	56	CG1	VAL	A	9	8.500	-6.952	9.170	1.00	11.60	C
ATOM	57	CG2	VAL	A	9	9.183	-6.171	11.474	1.00	11.98	C
ATOM	58	C	VAL	A	9	6.399	-5.370	12.019	1.00	9.36	C
ATOM	59	O	VAL	A	9	6.596	-4.276	12.541	1.00	9.87	O
ATOM	60	N	GLY	A	10	5.812	-6.369	12.648	1.00	9.30	N
ATOM	61	CA	GLY	A	10	5.346	-6.288	14.024	1.00	10.14	C
ATOM	62	C	GLY	A	10	4.095	-7.136	14.205	1.00	9.38	C
ATOM	63	O	GLY	A	10	3.500	-7.647	13.243	1.00	9.61	O
ATOM	64	N	ASP	A	11	3.702	-7.283	15.466	1.00	9.39	N
ATOM	65	CA	ASP	A	11	2.596	-8.170	15.822	1.00	10.45	C
ATOM	66	CB	ASP	A	11	2.329	-8.105	17.323	1.00	10.06	C
ATOM	67	CG	ASP	A	11	3.288	-8.959	18.135	1.00	12.02	C
ATOM	68	OD1	ASP	A	11	4.248	-9.545	17.584	1.00	12.12	O
ATOM	69	OD2	ASP	A	11	3.029	-9.044	19.354	1.00	13.04	O
ATOM	70	C	ASP	A	11	1.324	-7.788	15.111	1.00	9.85	C
ATOM	71	O	ASP	A	11	1.101	-6.621	14.774	1.00	9.42	O
ATOM	72	N	GLY	A	12	0.433	-8.753	14.932	1.00	10.76	N
ATOM	73	CA	GLY	A	12	-0.908	-8.453	14.437	1.00	10.53	C
ATOM	74	C	GLY	A	12	-1.572	-7.408	15.312	1.00	10.68	C
ATOM	75	O	GLY	A	12	-1.433	-7.428	16.541	1.00	11.67	O
ATOM	76	N	ALA	A	13	-2.290	-6.488	14.674	1.00	10.27	N
ATOM	77	CA	ALA	A	13	-3.124	-5.478	15.320	1.00	10.10	C
ATOM	78	CB	ALA	A	13	-4.081	-6.090	16.347	1.00	11.47	C
ATOM	79	C	ALA	A	13	-2.372	-4.292	15.913	1.00	10.23	C
ATOM	80	O	ALA	A	13	-2.991	-3.454	16.586	1.00	11.32	O
ATOM	81	N	VAL	A	14	-1.073	-4.187	15.662	1.00	10.56	N
ATOM	82	CA	VAL	A	14	-0.355	-2.999	16.193	1.00	10.35	C
ATOM	83	CB	VAL	A	14	1.161	-3.227	16.365	1.00	10.51	C
ATOM	84	CG1	VAL	A	14	1.444	-4.399	17.307	1.00	11.36	C
ATOM	85	CG2	VAL	A	14	1.861	-3.405	15.042	1.00	11.73	C
ATOM	86	C	VAL	A	14	-0.617	-1.727	15.403	1.00	10.69	C
ATOM	87	O	VAL	A	14	-0.410	-0.624	15.917	1.00	11.18	O
ATOM	88	N	GLY	A	15	-1.108	-1.859	14.176	1.00	9.47	N
ATOM	89	CA	GLY	A	15	-1.432	-0.721	13.333	1.00	10.07	C
ATOM	90	C	GLY	A	15	-0.547	-0.518	12.119	1.00	9.10	C
ATOM	91	O	GLY	A	15	-0.502	0.581	11.585	1.00	9.75	O
ATOM	92	N	LYS	A	16	0.079	-1.580	11.616	1.00	8.57	N
ATOM	93	CA	LYS	A	16	0.976	-1.451	10.467	1.00	8.65	C
ATOM	94	CB	LYS	A	16	1.674	-2.782	10.183	1.00	8.61	C
ATOM	95	CG	LYS	A	16	2.485	-3.314	11.373	1.00	9.93	C
ATOM	96	CD	LYS	A	16	3.194	-4.658	11.106	1.00	10.00	C
ATOM	97	CE	LYS	A	16	2.275	-5.817	10.819	1.00	10.39	C
ATOM	98	NZ	LYS	A	16	1.366	-6.104	11.952	1.00	9.45	N
ATOM	99	C	LYS	A	16	0.227	-0.996	9.228	1.00	8.25	C
ATOM	100	O	LYS	A	16	0.641	-0.060	8.535	1.00	8.46	O
ATOM	101	N	THR	A	17	-0.875	-1.672	8.912	1.00	8.80	N
ATOM	102	CA	THR	A	17	-1.615	-1.343	7.710	1.00	9.41	C
ATOM	103	CB	THR	A	17	-2.689	-2.400	7.503	1.00	8.99	C
ATOM	104	OG1	THR	A	17	-2.020	-3.677	7.367	1.00	9.21	O
ATOM	105	CG2	THR	A	17	-3.503	-2.155	6.248	1.00	10.06	C
ATOM	106	C	THR	A	17	-2.204	0.070	7.800	1.00	8.93	C
ATOM	107	O	THR	A	17	-2.136	0.834	6.828	1.00	9.41	O
ATOM	108	N	CYS	A	18	-2.784	0.423	8.943	1.00	9.42	N
ATOM	109	CA	CYS	A	18	-3.355	1.741	9.077	1.00	10.25	C

ATOM	110	CB	CYS	A	18	-4.075	1.897	10.396	1.00	10.15	C
ATOM	111	SG	CYS	A	18	-5.693	1.128	10.433	1.00	11.45	S
ATOM	112	C	CYS	A	18	-2.296	2.805	8.984	1.00	10.14	C
ATOM	113	O	CYS	A	18	-2.576	3.871	8.414	1.00	11.24	O
ATOM	114	N	LEU	A	19	-1.114	2.611	9.549	1.00	10.34	N
ATOM	115	CA	LEU	A	19	-0.175	3.719	9.443	1.00	11.62	C
ATOM	116	C	LEU	A	19	0.273	3.893	8.004	1.00	11.02	C
ATOM	117	O	LEU	A	19	0.512	5.022	7.575	1.00	11.14	O
ATOM	118	CB	LEU	A	19	1.025	3.494	10.365	1.00	12.92	C
ATOM	119	CG	LEU	A	19	1.988	2.375	9.965	1.00	20.00	C
ATOM	120	CD1	LEU	A	19	3.014	2.881	8.963	1.00	15.00	C
ATOM	121	CD2	LEU	A	19	2.677	1.795	11.190	1.00	20.00	C
ATOM	122	N	LEU	A	20	0.350	2.817	7.229	1.00	10.17	N
ATOM	123	CA	LEU	A	20	0.685	2.943	5.809	1.00	10.15	C
ATOM	124	CB	LEU	A	20	1.019	1.577	5.203	1.00	10.06	C
ATOM	125	CG	LEU	A	20	2.331	0.991	5.722	1.00	9.84	C
ATOM	126	CD1	LEU	A	20	2.432	-0.485	5.309	1.00	12.20	C
ATOM	127	CD2	LEU	A	20	3.522	1.778	5.244	1.00	13.56	C
ATOM	128	C	LEU	A	20	-0.421	3.631	5.007	1.00	10.29	C
ATOM	129	O	LEU	A	20	-0.132	4.471	4.153	1.00	10.44	O
ATOM	130	N	ILE	A	21	-1.669	3.293	5.279	1.00	10.12	N
ATOM	131	CA	ILE	A	21	-2.789	3.912	4.590	1.00	10.14	C
ATOM	132	CB	ILE	A	21	-4.093	3.182	4.880	1.00	10.76	C
ATOM	133	CG1	ILE	A	21	-4.044	1.818	4.193	1.00	11.42	C
ATOM	134	CD1	ILE	A	21	-5.125	0.806	4.611	1.00	13.68	C
ATOM	135	CG2	ILE	A	21	-5.300	3.964	4.352	1.00	12.21	C
ATOM	136	C	ILE	A	21	-2.883	5.381	4.987	1.00	9.92	C
ATOM	137	O	ILE	A	21	-3.078	6.234	4.112	1.00	10.89	O
ATOM	138	N	SER	A	22	-2.730	5.697	6.259	0.50	6.37	N
ATOM	139	CA	SER	A	22	-2.774	7.094	6.668	0.50	6.91	C
ATOM	140	CB	SER	A	22	-2.711	7.193	8.182	0.50	6.73	C
ATOM	141	OG	ASER	A	22	-3.945	6.769	8.703	0.50	12.30	O
ATOM	141	OG	BSER	A	22	-2.826	8.535	8.599	0.50	11.36	O
ATOM	142	C	SER	A	22	-1.642	7.893	6.019	0.50	6.92	C
ATOM	143	O	SER	A	22	-1.841	9.031	5.573	0.50	5.72	O
ATOM	144	N	TYR	A	23	-0.449	7.310	5.949	1.00	10.57	N
ATOM	145	CA	TYR	A	23	0.681	8.011	5.355	1.00	11.21	C
ATOM	146	CB	TYR	A	23	1.963	7.202	5.563	1.00	11.78	C
ATOM	147	CG	TYR	A	23	3.203	7.835	4.958	1.00	11.21	C
ATOM	148	CD1	TYR	A	23	3.655	9.080	5.389	1.00	11.88	C
ATOM	149	CE1	TYR	A	23	4.790	9.656	4.842	1.00	11.58	C
ATOM	150	CZ	TYR	A	23	5.513	8.958	3.884	1.00	11.24	C
ATOM	151	OH	TYR	A	23	6.654	9.496	3.318	1.00	12.92	O
ATOM	152	CE2	TYR	A	23	5.091	7.730	3.446	1.00	11.94	C
ATOM	153	CD2	TYR	A	23	3.953	7.167	3.992	1.00	11.19	C
ATOM	154	C	TYR	A	23	0.479	8.286	3.869	1.00	12.23	C
ATOM	155	O	TYR	A	23	0.855	9.356	3.343	1.00	13.13	O
ATOM	156	N	THR	A	24	-0.100	7.321	3.172	1.00	12.75	N
ATOM	157	CA	THR	A	24	-0.174	7.422	1.715	1.00	13.41	C
ATOM	158	CB	THR	A	24	-0.089	6.044	1.027	1.00	13.28	C
ATOM	159	OG1	THR	A	24	-1.141	5.191	1.515	1.00	12.91	O
ATOM	160	CG2	THR	A	24	1.259	5.394	1.283	1.00	13.69	C
ATOM	161	C	THR	A	24	-1.406	8.160	1.230	1.00	14.93	C
ATOM	162	O	THR	A	24	-1.369	8.734	0.143	1.00	15.84	O
ATOM	163	N	THR	A	25	-2.477	8.181	2.022	1.00	14.62	N
ATOM	164	CA	THR	A	25	-3.754	8.759	1.589	1.00	15.96	C
ATOM	165	CB	THR	A	25	-4.890	7.711	1.573	1.00	15.67	C
ATOM	166	OG1	THR	A	25	-5.267	7.374	2.925	1.00	16.40	O
ATOM	167	CG2	THR	A	25	-4.498	6.474	0.785	1.00	15.73	C
ATOM	168	C	THR	A	25	-4.234	9.918	2.459	1.00	17.13	C
ATOM	169	O	THR	A	25	-5.197	10.593	2.089	1.00	17.10	O
ATOM	170	N	ASN	A	26	-3.626	10.103	3.626	1.00	18.51	N
ATOM	171	CA	ASN	A	26	-4.079	11.125	4.599	1.00	20.28	C
ATOM	172	CB	ASN	A	26	-3.885	12.548	4.040	1.00	21.61	C
ATOM	173	CG	ASN	A	26	-2.437	12.847	3.726	1.00	23.49	C
ATOM	174	OD1	ASN	A	26	-2.108	13.367	2.650	1.00	27.93	O

ATOM	175	ND2	ASN	A	26	-1.546	12.502	4.657	1.00	25.36	N
ATOM	176	C	ASN	A	26	-5.496	10.915	5.133	1.00	21.61	C
ATOM	177	O	ASN	A	26	-6.133	11.858	5.625	1.00	22.35	O
ATOM	178	N	ALA	A	27	-5.999	9.690	5.041	1.00	21.20	N
ATOM	179	CA	ALA	A	27	-7.270	9.364	5.665	1.00	21.90	C
ATOM	180	CB	ALA	A	27	-8.324	9.074	4.626	1.00	22.79	C
ATOM	181	C	ALA	A	27	-7.036	8.154	6.561	1.00	21.66	C
ATOM	182	O	ALA	A	27	-6.462	7.172	6.115	1.00	22.33	O
ATOM	183	N	PHE	A	28	-7.410	8.257	7.830	1.00	19.80	N
ATOM	184	CA	PHE	A	28	-7.389	7.090	8.715	1.00	17.99	C
ATOM	185	CB	PHE	A	28	-7.504	7.524	10.165	1.00	17.74	C
ATOM	186	CG	PHE	A	28	-7.569	6.380	11.124	1.00	16.32	C
ATOM	187	CD1	PHE	A	28	-6.452	5.574	11.360	1.00	16.38	C
ATOM	188	CE1	PHE	A	28	-6.516	4.504	12.268	1.00	14.61	C
ATOM	189	CZ	PHE	A	28	-7.713	4.235	12.930	1.00	16.53	C
ATOM	190	CE2	PHE	A	28	-8.840	5.025	12.686	1.00	16.63	C
ATOM	191	CD2	PHE	A	28	-8.759	6.095	11.783	1.00	15.57	C
ATOM	192	C	PHE	A	28	-8.544	6.155	8.342	1.00	17.65	C
ATOM	193	O	PHE	A	28	-9.707	6.563	8.334	1.00	17.89	O
ATOM	194	N	PRO	A	29	-8.237	4.897	7.982	1.00	16.19	N
ATOM	195	CA	PRO	A	29	-9.275	4.005	7.446	1.00	15.78	C
ATOM	196	CB	PRO	A	29	-8.457	3.021	6.603	1.00	14.96	C
ATOM	197	CG	PRO	A	29	-7.162	2.881	7.423	1.00	15.06	C
ATOM	198	CD	PRO	A	29	-6.889	4.281	7.933	1.00	15.67	C
ATOM	199	C	PRO	A	29	-10.111	3.225	8.456	1.00	16.17	C
ATOM	200	O	PRO	A	29	-11.017	2.496	8.064	1.00	16.59	O
ATOM	201	N	GLY	A	30	-9.816	3.343	9.741	1.00	16.47	N
ATOM	202	CA	GLY	A	30	-10.613	2.622	10.746	1.00	16.48	C
ATOM	203	C	GLY	A	30	-10.533	1.107	10.589	1.00	16.58	C
ATOM	204	O	GLY	A	30	-9.443	0.543	10.592	1.00	16.49	O
ATOM	205	N	GLU	A	31	-11.681	0.457	10.447	1.00	16.20	N
ATOM	206	CA	GLU	A	31	-11.752	-0.994	10.256	1.00	16.97	C
ATOM	207	CB	GLU	A	31	-12.970	-1.573	10.986	1.00	18.18	C
ATOM	208	CG	GLU	A	31	-13.022	-1.270	12.463	1.00	22.26	C
ATOM	209	CD	GLU	A	31	-11.703	-1.558	13.161	1.00	26.99	C
ATOM	210	OE1	GLU	A	31	-11.201	-2.698	13.072	1.00	30.47	O
ATOM	211	OE2	GLU	A	31	-11.164	-0.635	13.796	1.00	31.76	O
ATOM	212	C	GLU	A	31	-11.802	-1.400	8.780	1.00	15.98	C
ATOM	213	O	GLU	A	31	-11.844	-2.590	8.449	1.00	16.42	O
ATOM	214	N	TYR	A	32	-11.785	-0.410	7.900	1.00	16.12	N
ATOM	215	CA	TYR	A	32	-11.922	-0.686	6.468	1.00	17.14	C
ATOM	216	CB	TYR	A	32	-12.725	0.415	5.782	1.00	18.87	C
ATOM	217	CG	TYR	A	32	-14.165	0.415	6.240	1.00	20.99	C
ATOM	218	CD1	TYR	A	32	-15.095	-0.446	5.667	1.00	23.72	C
ATOM	219	CE1	TYR	A	32	-16.417	-0.460	6.093	1.00	26.05	C
ATOM	220	CZ	TYR	A	32	-16.819	0.403	7.097	1.00	24.58	C
ATOM	221	OH	TYR	A	32	-18.130	0.394	7.507	1.00	25.90	O
ATOM	222	CE2	TYR	A	32	-15.913	1.267	7.697	1.00	23.82	C
ATOM	223	CD2	TYR	A	32	-14.590	1.270	7.261	1.00	22.07	C
ATOM	224	C	TYR	A	32	-10.556	-0.856	5.848	1.00	15.47	C
ATOM	225	O	TYR	A	32	-10.058	-0.007	5.111	1.00	15.10	O
ATOM	226	N	ILE	A	33	-9.935	-1.979	6.210	1.00	14.88	N
ATOM	227	CA	ILE	A	33	-8.540	-2.215	5.873	1.00	14.90	C
ATOM	228	CB	ILE	A	33	-7.635	-1.892	7.090	1.00	15.46	C
ATOM	229	CG1	ILE	A	33	-8.141	-2.656	8.343	1.00	17.42	C
ATOM	230	CD1	ILE	A	33	-7.582	-2.237	9.684	1.00	17.18	C
ATOM	231	CG2	ILE	A	33	-7.558	-0.378	7.332	1.00	14.83	C
ATOM	232	C	ILE	A	33	-8.325	-3.670	5.511	1.00	13.45	C
ATOM	233	O	ILE	A	33	-9.015	-4.544	6.032	1.00	14.07	O
ATOM	234	N	PRO	A	34	-7.348	-3.942	4.636	1.00	11.93	N
ATOM	235	CA	PRO	A	34	-6.936	-5.310	4.369	1.00	11.72	C
ATOM	236	CB	PRO	A	34	-6.181	-5.179	3.043	1.00	11.45	C
ATOM	237	CG	PRO	A	34	-5.535	-3.809	3.161	1.00	12.24	C
ATOM	238	CD	PRO	A	34	-6.560	-2.968	3.857	1.00	11.49	C
ATOM	239	C	PRO	A	34	-5.994	-5.804	5.481	1.00	11.22	C
ATOM	240	O	PRO	A	34	-5.495	-4.999	6.281	1.00	10.92	O

ATOM	241	N	THR	A	35	-5.698	-7.102	5.504	1.00	10.66	N
ATOM	242	CA	THR	A	35	-4.644	-7.603	6.366	1.00	10.28	C
ATOM	243	CB	THR	A	35	-4.530	-9.132	6.220	1.00	10.32	C
ATOM	244	OG1	THR	A	35	-5.787	-9.740	6.564	1.00	10.83	O
ATOM	245	CG2	THR	A	35	-3.440	-9.719	7.103	1.00	10.76	C
ATOM	246	C	THR	A	35	-3.317	-6.954	5.972	1.00	9.92	C
ATOM	247	O	THR	A	35	-2.573	-6.502	6.855	1.00	9.20	O
ATOM	248	N	VAL	A	36	-3.026	-6.915	4.676	1.00	9.92	N
ATOM	249	CA	VAL	A	36	-1.765	-6.390	4.189	1.00	11.66	C
ATOM	250	CB	VAL	A	36	-0.911	-7.532	3.564	1.00	12.55	C
ATOM	251	CG1	VAL	A	36	0.435	-7.006	3.135	1.00	14.75	C
ATOM	252	CG2	VAL	A	36	-0.764	-8.749	4.538	1.00	13.86	C
ATOM	253	C	VAL	A	36	-2.045	-5.353	3.100	1.00	11.71	C
ATOM	254	O	VAL	A	36	-2.736	-5.644	2.124	1.00	12.92	O
ATOM	255	N	PHE	A	37	-1.555	-4.136	3.293	1.00	11.91	N
ATOM	256	CA	PHE	A	37	-1.630	-3.073	2.301	1.00	12.80	C
ATOM	257	CB	PHE	A	37	-0.957	-1.854	2.933	1.00	13.56	C
ATOM	258	CG	PHE	A	37	-1.143	-0.571	2.182	1.00	16.30	C
ATOM	259	CD1	PHE	A	37	-2.405	-0.166	1.753	1.00	19.42	C
ATOM	260	CE1	PHE	A	37	-2.578	1.058	1.071	1.00	20.61	C
ATOM	261	CZ	PHE	A	37	-1.475	1.871	0.845	1.00	17.13	C
ATOM	262	CE2	PHE	A	37	-0.211	1.483	1.264	1.00	17.41	C
ATOM	263	CD2	PHE	A	37	-0.050	0.250	1.948	1.00	18.70	C
ATOM	264	C	PHE	A	37	-0.893	-3.459	1.039	1.00	14.08	C
ATOM	265	O	PHE	A	37	0.169	-4.074	1.120	1.00	14.02	O
ATOM	266	N	ASP	A	38	-1.434	-3.055	-0.110	1.00	15.41	N
ATOM	267	CA	ASP	A	38	-0.772	-3.259	-1.404	1.00	17.12	C
ATOM	268	CB	ASP	A	38	-1.618	-2.627	-2.533	1.00	18.72	C
ATOM	269	CG	ASP	A	38	-2.911	-3.389	-2.846	1.00	23.03	C
ATOM	270	OD1	ASP	A	38	-3.085	-4.547	-2.409	1.00	26.95	O
ATOM	271	OD2	ASP	A	38	-3.757	-2.798	-3.563	1.00	28.04	O
ATOM	272	C	ASP	A	38	0.616	-2.577	-1.389	1.00	16.38	C
ATOM	273	O	ASP	A	38	0.804	-1.560	-0.735	1.00	16.07	O
ATOM	274	N	ASN	A	39	1.584	-3.135	-2.099	1.00	16.55	N
ATOM	275	CA	ASN	A	39	2.857	-2.436	-2.288	1.00	17.38	C
ATOM	276	CB	ASN	A	39	3.694	-3.191	-3.309	1.00	18.41	C
ATOM	277	CG	ASN	A	39	5.083	-2.657	-3.403	1.00	22.02	C
ATOM	278	OD1	ASN	A	39	5.471	-2.085	-4.423	1.00	26.34	O
ATOM	279	ND2	ASN	A	39	5.844	-2.809	-2.327	1.00	27.14	N
ATOM	280	C	ASN	A	39	2.648	-1.011	-2.788	1.00	16.04	C
ATOM	281	O	ASN	A	39	1.772	-0.761	-3.622	1.00	16.15	O
ATOM	282	N	TYR	A	40	3.440	-0.068	-2.280	1.00	14.53	N
ATOM	283	CA	TYR	A	40	3.306	1.337	-2.665	1.00	14.34	C
ATOM	284	CB	TYR	A	40	2.592	2.105	-1.544	1.00	14.12	C
ATOM	285	CG	TYR	A	40	2.419	3.582	-1.815	1.00	14.39	C
ATOM	286	CD1	TYR	A	40	1.294	4.049	-2.490	1.00	14.96	C
ATOM	287	CE1	TYR	A	40	1.125	5.409	-2.749	1.00	14.64	C
ATOM	288	CZ	TYR	A	40	2.090	6.291	-2.330	1.00	15.18	C
ATOM	289	OH	TYR	A	40	1.898	7.628	-2.607	1.00	16.39	O
ATOM	290	CE2	TYR	A	40	3.220	5.859	-1.665	1.00	14.84	C
ATOM	291	CD2	TYR	A	40	3.385	4.500	-1.414	1.00	14.38	C
ATOM	292	C	TYR	A	40	4.683	1.934	-2.873	1.00	13.60	C
ATOM	293	O	TYR	A	40	5.580	1.718	-2.064	1.00	14.93	O
ATOM	294	N	SER	A	41	4.875	2.685	-3.951	1.00	12.48	N
ATOM	295	CA	SER	A	41	6.077	3.495	-4.068	1.00	13.53	C
ATOM	296	CB	SER	A	41	7.158	2.872	-4.960	1.00	14.96	C
ATOM	297	OG	SER	A	41	6.766	2.869	-6.306	1.00	19.82	O
ATOM	298	C	SER	A	41	5.706	4.876	-4.559	1.00	12.45	C
ATOM	299	O	SER	A	41	4.742	5.037	-5.308	1.00	13.38	O
ATOM	300	N	ALA	A	42	6.485	5.856	-4.136	1.00	11.65	N
ATOM	301	CA	ALA	A	42	6.322	7.235	-4.575	1.00	11.45	C
ATOM	302	CB	ALA	A	42	5.093	7.881	-3.900	1.00	12.12	C
ATOM	303	C	ALA	A	42	7.555	8.004	-4.206	1.00	11.86	C
ATOM	304	O	ALA	A	42	8.337	7.574	-3.341	1.00	11.97	O
ATOM	305	N	ASN	A	43	7.743	9.139	-4.864	1.00	11.38	N
ATOM	306	CA	ASN	A	43	8.759	10.096	-4.456	1.00	11.29	C

ATOM	307	CB	ASN	A	43	9.150	11.013	-5.625	1.00	12.66	C
ATOM	308	CG	ASN	A	43	9.986	10.313	-6.670	1.00	14.01	C
ATOM	309	OD1	ASN	A	43	10.362	9.142	-6.535	1.00	15.75	O
ATOM	310	ND2	ASN	A	43	10.287	11.034	-7.742	1.00	16.59	N
ATOM	311	C	ASN	A	43	8.258	10.943	-3.315	1.00	11.75	C
ATOM	312	O	ASN	A	43	7.171	11.525	-3.378	1.00	11.94	O
ATOM	313	N	VAL	A	44	9.056	11.010	-2.257	1.00	11.62	N
ATOM	314	CA	VAL	A	44	8.733	11.826	-1.097	1.00	13.22	C
ATOM	315	CB	VAL	A	44	8.432	10.946	0.147	1.00	14.92	C
ATOM	316	CG1	VAL	A	44	7.320	9.959	-0.145	1.00	16.55	C
ATOM	317	CG2	VAL	A	44	9.626	10.198	0.556	1.00	15.99	C
ATOM	318	C	VAL	A	44	9.872	12.807	-0.849	1.00	13.51	C
ATOM	319	O	VAL	A	44	11.029	12.488	-1.103	1.00	13.42	O
ATOM	320	N	MET	A	45	9.541	13.990	-0.353	1.00	14.85	N
ATOM	321	CA	MET	A	45	10.532	15.017	-0.131	1.00	15.75	C
ATOM	322	CB	MET	A	45	9.877	16.407	-0.141	1.00	16.60	C
ATOM	323	CG	MET	A	45	9.275	16.835	-1.464	1.00	18.19	C
ATOM	324	SD	MET	A	45	10.418	16.742	-2.826	1.00	21.74	S
ATOM	325	CE	MET	A	45	11.537	18.088	-2.504	1.00	21.52	C
ATOM	326	C	MET	A	45	11.184	14.817	1.223	1.00	16.61	C
ATOM	327	O	MET	A	45	10.500	14.803	2.243	1.00	16.90	O
ATOM	328	N	VAL	A	46	12.506	14.689	1.220	1.00	16.86	N
ATOM	329	CA	VAL	A	46	13.286	14.637	2.455	1.00	18.05	C
ATOM	330	CB	VAL	A	46	13.720	13.181	2.816	1.00	18.32	C
ATOM	331	CG1	VAL	A	46	14.477	13.146	4.150	1.00	19.56	C
ATOM	332	CG2	VAL	A	46	12.501	12.288	2.910	1.00	18.50	C
ATOM	333	C	VAL	A	46	14.496	15.537	2.257	1.00	18.60	C
ATOM	334	O	VAL	A	46	15.222	15.388	1.288	1.00	18.55	O
ATOM	335	N	ASP	A	47	14.688	16.473	3.182	1.00	20.72	N
ATOM	336	CA	ASP	A	47	15.862	17.352	3.181	1.00	22.66	C
ATOM	337	CB	ASP	A	47	17.130	16.577	3.610	1.00	23.69	C
ATOM	338	CG	ASP	A	47	16.997	15.961	5.012	1.00	25.31	C
ATOM	339	OD1	ASP	A	47	16.347	16.575	5.881	1.00	28.40	O
ATOM	340	OD2	ASP	A	47	17.535	14.861	5.241	1.00	27.71	O
ATOM	341	C	ASP	A	47	16.010	18.073	1.832	1.00	23.13	C
ATOM	342	O	ASP	A	47	17.114	18.237	1.308	1.00	24.89	O
ATOM	343	N	GLY	A	48	14.866	18.474	1.274	1.00	23.26	N
ATOM	344	CA	GLY	A	48	14.804	19.254	0.030	1.00	23.23	C
ATOM	345	C	GLY	A	48	14.817	18.500	-1.290	1.00	22.89	C
ATOM	346	O	GLY	A	48	14.720	19.114	-2.359	1.00	23.43	O
ATOM	347	N	LYS	A	49	14.931	17.174	-1.225	1.00	21.63	N
ATOM	348	CA	LYS	A	49	15.128	16.356	-2.417	1.00	20.77	C
ATOM	349	CB	LYS	A	49	16.524	15.704	-2.387	1.00	21.96	C
ATOM	350	CG	LYS	A	49	17.693	16.691	-2.355	1.00	25.00	C
ATOM	351	CD	LYS	A	49	17.878	17.387	-3.687	1.00	28.52	C
ATOM	352	CE	LYS	A	49	18.984	18.427	-3.611	1.00	31.72	C
ATOM	353	NZ	LYS	A	49	19.420	18.893	-4.957	1.00	32.72	N
ATOM	354	C	LYS	A	49	14.071	15.257	-2.519	1.00	18.40	C
ATOM	355	O	LYS	A	49	13.704	14.664	-1.506	1.00	18.26	O
ATOM	356	N	PRO	A	50	13.598	14.964	-3.737	1.00	16.50	N
ATOM	357	CA	PRO	A	50	12.664	13.852	-3.850	1.00	16.28	C
ATOM	358	CB	PRO	A	50	12.025	14.056	-5.224	1.00	16.17	C
ATOM	359	CG	PRO	A	50	13.039	14.829	-6.022	1.00	16.50	C
ATOM	360	CD	PRO	A	50	13.882	15.601	-5.043	1.00	17.02	C
ATOM	361	C	PRO	A	50	13.405	12.521	-3.781	1.00	15.94	C
ATOM	362	O	PRO	A	50	14.446	12.339	-4.434	1.00	18.03	O
ATOM	363	N	VAL	A	51	12.881	11.601	-2.980	1.00	14.23	N
ATOM	364	CA	VAL	A	51	13.484	10.289	-2.796	1.00	13.32	C
ATOM	365	CB	VAL	A	51	14.018	10.128	-1.347	1.00	14.06	C
ATOM	366	CG1	VAL	A	51	14.630	8.741	-1.116	1.00	14.65	C
ATOM	367	CG2	VAL	A	51	15.039	11.224	-1.013	1.00	16.35	C
ATOM	368	C	VAL	A	51	12.425	9.243	-3.075	1.00	12.07	C
ATOM	369	O	VAL	A	51	11.325	9.309	-2.516	1.00	12.14	O
ATOM	370	N	ASN	A	52	12.744	8.285	-3.932	1.00	12.03	N
ATOM	371	CA	ASN	A	52	11.818	7.204	-4.242	1.00	12.95	C
ATOM	372	CB	ASN	A	52	12.243	6.548	-5.546	1.00	14.19	C

ATOM	373	CG	ASN	A	52	11.205	5.577	-6.094	1.00	17.09	C
ATOM	374	OD1	ASN	A	52	10.051	5.567	-5.665	1.00	22.79	O
ATOM	375	ND2	ASN	A	52	11.606	4.783	-7.073	1.00	21.92	N
ATOM	376	C	ASN	A	52	11.778	6.192	-3.087	1.00	12.56	C
ATOM	377	O	ASN	A	52	12.827	5.616	-2.720	1.00	13.33	O
ATOM	378	N	LEU	A	53	10.601	6.057	-2.484	1.00	11.75	N
ATOM	379	CA	LEU	A	53	10.391	5.213	-1.318	1.00	11.92	C
ATOM	380	CB	LEU	A	53	9.909	6.087	-0.151	1.00	12.50	C
ATOM	381	CG	LEU	A	53	9.552	5.350	1.135	1.00	11.65	C
ATOM	382	CD1	LEU	A	53	10.788	4.694	1.770	1.00	13.65	C
ATOM	383	CD2	LEU	A	53	8.882	6.340	2.104	1.00	14.03	C
ATOM	384	C	LEU	A	53	9.375	4.114	-1.622	1.00	11.73	C
ATOM	385	O	LEU	A	53	8.270	4.411	-2.109	1.00	12.99	O
ATOM	386	N	GLY	A	54	9.734	2.860	-1.350	1.00	11.07	N
ATOM	387	CA	GLY	A	54	8.833	1.725	-1.562	1.00	11.07	C
ATOM	388	C	GLY	A	54	8.436	1.156	-0.207	1.00	10.64	C
ATOM	389	O	GLY	A	54	9.300	0.945	0.648	1.00	11.82	O
ATOM	390	N	LEU	A	55	7.151	0.919	0.002	1.00	10.59	N
ATOM	391	CA	LEU	A	55	6.614	0.486	1.282	1.00	10.70	C
ATOM	392	CB	LEU	A	55	5.548	1.462	1.760	1.00	11.17	C
ATOM	393	CG	LEU	A	55	5.995	2.920	1.882	1.00	12.50	C
ATOM	394	CD1	LEU	A	55	4.789	3.784	2.201	1.00	14.81	C
ATOM	395	CD2	LEU	A	55	7.030	3.045	2.943	1.00	14.06	C
ATOM	396	C	LEU	A	55	6.001	-0.904	1.150	1.00	9.88	C
ATOM	397	O	LEU	A	55	5.134	-1.143	0.306	1.00	10.58	O
ATOM	398	N	TRP	A	56	6.446	-1.790	2.037	1.00	10.18	N
ATOM	399	CA	TRP	A	56	6.061	-3.200	2.047	1.00	10.73	C
ATOM	400	CB	TRP	A	56	7.266	-4.100	1.740	1.00	12.68	C
ATOM	401	CG	TRP	A	56	7.846	-3.871	0.371	1.00	13.71	C
ATOM	402	CD1	TRP	A	56	8.413	-2.719	-0.105	1.00	15.24	C
ATOM	403	NE1	TRP	A	56	8.836	-2.897	-1.391	1.00	17.10	N
ATOM	404	CE2	TRP	A	56	8.578	-4.193	-1.757	1.00	16.99	C
ATOM	405	CD2	TRP	A	56	7.961	-4.831	-0.658	1.00	19.16	C
ATOM	406	CE3	TRP	A	56	7.604	-6.180	-0.763	1.00	20.14	C
ATOM	407	CZ3	TRP	A	56	7.868	-6.852	-1.949	1.00	19.43	C
ATOM	408	CH2	TRP	A	56	8.489	-6.193	-3.031	1.00	18.67	C
ATOM	409	CZ2	TRP	A	56	8.853	-4.868	-2.955	1.00	18.68	C
ATOM	410	C	TRP	A	56	5.470	-3.549	3.410	1.00	10.65	C
ATOM	411	O	TRP	A	56	6.201	-3.684	4.391	1.00	11.81	O
ATOM	412	N	ASP	A	57	4.172	-3.756	3.441	1.00	10.06	N
ATOM	413	CA	ASP	A	57	3.465	-4.187	4.637	1.00	9.81	C
ATOM	414	CB	ASP	A	57	1.994	-3.820	4.461	1.00	10.02	C
ATOM	415	CG	ASP	A	57	1.179	-3.832	5.749	1.00	9.84	C
ATOM	416	OD1	ASP	A	57	1.699	-4.104	6.842	1.00	10.07	O
ATOM	417	OD2	ASP	A	57	-0.047	-3.614	5.611	1.00	10.00	O
ATOM	418	C	ASP	A	57	3.632	-5.700	4.751	1.00	10.58	C
ATOM	419	O	ASP	A	57	3.901	-6.383	3.766	1.00	11.80	O
ATOM	420	N	THR	A	58	3.486	-6.225	5.960	1.00	10.01	N
ATOM	421	CA	THR	A	58	3.585	-7.664	6.173	1.00	10.87	C
ATOM	422	CB	THR	A	58	4.964	-8.090	6.735	1.00	11.02	C
ATOM	423	OG1	THR	A	58	5.164	-7.536	8.053	1.00	11.24	O
ATOM	424	CG2	THR	A	58	6.097	-7.729	5.792	1.00	11.90	C
ATOM	425	C	THR	A	58	2.514	-8.075	7.151	1.00	10.83	C
ATOM	426	O	THR	A	58	1.956	-7.241	7.867	1.00	11.29	O
ATOM	427	N	ALA	A	59	2.239	-9.372	7.212	1.00	11.35	N
ATOM	428	CA	ALA	A	59	1.329	-9.920	8.199	1.00	11.95	C
ATOM	429	CB	ALA	A	59	0.386	-10.932	7.536	1.00	13.29	C
ATOM	430	C	ALA	A	59	2.173	-10.583	9.284	1.00	11.57	C
ATOM	431	O	ALA	A	59	2.974	-11.495	8.974	1.00	11.96	O
ATOM	432	N	GLY	A	60	2.009	-10.130	10.526	1.00	11.72	N
ATOM	433	CA	GLY	A	60	2.897	-10.527	11.600	1.00	12.73	C
ATOM	434	C	GLY	A	60	2.489	-11.759	12.381	1.00	12.85	C
ATOM	435	O	GLY	A	60	3.301	-12.278	13.147	1.00	13.80	O
ATOM	436	N	GLN	A	61	1.257	-12.232	12.199	1.00	12.05	N
ATOM	437	CA	GLN	A	61	0.766	-13.397	12.944	1.00	12.75	C
ATOM	438	CB	GLN	A	61	-0.732	-13.624	12.653	1.00	13.14	C

ATOM	439	CG	GLN	A	61	-1.633	-12.484	13.173	1.00	13.68	C
ATOM	440	CD	GLN	A	61	-1.918	-11.417	12.128	1.00	14.04	C
ATOM	441	OE1	GLN	A	61	-1.040	-11.004	11.360	1.00	14.06	O
ATOM	442	NE2	GLN	A	61	-3.146	-10.938	12.116	1.00	15.06	N
ATOM	443	C	GLN	A	61	1.556	-14.631	12.593	1.00	14.06	C
ATOM	444	O	GLN	A	61	2.109	-14.764	11.501	1.00	13.88	O
ATOM	445	N	GLU	A	62	1.565	-15.564	13.546	1.00	14.98	N
ATOM	446	CA	GLU	A	62	2.299	-16.825	13.410	1.00	17.27	C
ATOM	447	CB	GLU	A	62	2.075	-17.709	14.636	1.00	17.00	C
ATOM	448	CG	GLU	A	62	2.893	-18.988	14.598	1.00	21.33	C
ATOM	449	CD	GLU	A	62	2.667	-19.900	15.785	1.00	21.12	C
ATOM	450	OE1	GLU	A	62	1.627	-19.786	16.462	1.00	25.59	O
ATOM	451	OE2	GLU	A	62	3.548	-20.752	16.008	1.00	25.60	O
ATOM	452	C	GLU	A	62	1.953	-17.592	12.147	1.00	16.12	C
ATOM	453	O	GLU	A	62	2.821	-18.207	11.536	1.00	15.90	O
ATOM	454	N	ASP	A	63	0.686	-17.548	11.740	1.00	16.37	N
ATOM	455	CA	ASP	A	63	0.260	-18.221	10.511	1.00	17.61	C
ATOM	456	CB	ASP	A	63	-1.216	-17.943	10.232	1.00	19.31	C
ATOM	457	CG	ASP	A	63	-2.141	-18.963	10.861	1.00	24.06	C
ATOM	458	OD1	ASP	A	63	-1.672	-20.032	11.306	1.00	28.11	O
ATOM	459	OD2	ASP	A	63	-3.357	-18.680	10.879	1.00	29.29	O
ATOM	460	C	ASP	A	63	1.048	-17.822	9.269	1.00	15.86	C
ATOM	461	O	ASP	A	63	1.110	-18.562	8.302	1.00	16.35	O
ATOM	462	N	TYR	A	64	1.621	-16.614	9.283	1.00	14.65	N
ATOM	463	CA	TYR	A	64	2.334	-16.086	8.127	1.00	13.33	C
ATOM	464	CB	TYR	A	64	1.964	-14.616	7.880	1.00	12.64	C
ATOM	465	CG	TYR	A	64	0.479	-14.387	7.710	1.00	12.50	C
ATOM	466	CD1	TYR	A	64	-0.148	-14.596	6.492	1.00	12.77	C
ATOM	467	CE1	TYR	A	64	-1.531	-14.375	6.350	1.00	14.98	C
ATOM	468	CZ	TYR	A	64	-2.258	-13.947	7.443	1.00	13.61	C
ATOM	469	OH	TYR	A	64	-3.606	-13.709	7.330	1.00	15.56	O
ATOM	470	CE2	TYR	A	64	-1.629	-13.728	8.649	1.00	13.71	C
ATOM	471	CD2	TYR	A	64	-0.285	-13.953	8.774	1.00	12.14	C
ATOM	472	C	TYR	A	64	3.837	-16.189	8.260	1.00	13.33	C
ATOM	473	O	TYR	A	64	4.563	-15.686	7.415	1.00	13.05	O
ATOM	474	N	ASP	A	65	4.306	-16.880	9.300	1.00	13.58	N
ATOM	475	CA	ASP	A	65	5.748	-16.919	9.584	1.00	14.85	C
ATOM	476	CB	ASP	A	65	6.018	-17.727	10.861	1.00	15.32	C
ATOM	477	CG	ASP	A	65	5.757	-16.951	12.136	1.00	16.90	C
ATOM	478	OD1	ASP	A	65	5.384	-15.753	12.094	1.00	16.97	O
ATOM	479	OD2	ASP	A	65	5.906	-17.558	13.226	1.00	20.20	O
ATOM	480	C	ASP	A	65	6.570	-17.534	8.450	1.00	15.24	C
ATOM	481	O	ASP	A	65	7.710	-17.143	8.218	1.00	15.20	O
ATOM	482	N	ARG	A	66	6.003	-18.507	7.739	1.00	15.03	N
ATOM	483	CA	ARG	A	66	6.718	-19.130	6.623	1.00	15.74	C
ATOM	484	CB	ARG	A	66	6.162	-20.530	6.348	1.00	16.44	C
ATOM	485	CG	ARG	A	66	6.338	-21.464	7.517	1.00	17.39	C
ATOM	486	CD	ARG	A	66	5.662	-22.807	7.250	1.00	20.96	C
ATOM	487	NE	ARG	A	66	6.389	-23.591	6.258	1.00	22.46	N
ATOM	488	CZ	ARG	A	66	7.469	-24.325	6.534	1.00	23.29	C
ATOM	489	NH1	ARG	A	66	7.945	-24.376	7.776	1.00	23.36	N
ATOM	490	NH2	ARG	A	66	8.070	-25.000	5.568	1.00	26.82	N
ATOM	491	C	ARG	A	66	6.700	-18.302	5.335	1.00	16.07	C
ATOM	492	O	ARG	A	66	7.597	-18.415	4.503	1.00	17.45	O
ATOM	493	N	LEU	A	67	5.698	-17.570	5.190	1.00	15.40	N
ATOM	494	CA	LEU	A	67	5.552	-16.810	3.945	1.00	15.49	C
ATOM	495	CB	LEU	A	67	4.078	-16.706	3.580	1.00	15.84	C
ATOM	496	CG	LEU	A	67	3.702	-15.969	2.291	1.00	17.97	C
ATOM	497	CD1	LEU	A	67	4.246	-16.739	1.095	1.00	17.77	C
ATOM	498	CD2	LEU	A	67	2.207	-15.856	2.216	1.00	18.43	C
ATOM	499	C	LEU	A	67	6.174	-15.422	4.023	1.00	15.73	C
ATOM	500	O	LEU	A	67	6.766	-14.940	3.057	1.00	15.69	O
ATOM	501	N	ARG	A	68	6.056	-14.807	5.202	1.00	16.73	N
ATOM	502	CA	ARG	A	68	6.505	-13.442	5.402	1.00	17.12	C
ATOM	503	CB	ARG	A	68	6.233	-13.038	6.851	1.00	17.50	C
ATOM	504	CG	ARG	A	68	6.684	-11.644	7.216	1.00	14.25	C

ATOM	505	CD	ARG	A	68	6.307	-11.334	8.660	1.00	14.22	C
ATOM	506	NE	ARG	A	68	6.888	-12.277	9.614	1.00	15.63	N
ATOM	507	CZ	ARG	A	68	6.225	-13.190	10.317	1.00	13.45	C
ATOM	508	NH1	ARG	A	68	4.893	-13.350	10.240	1.00	12.81	N
ATOM	509	NH2	ARG	A	68	6.923	-13.947	11.139	1.00	15.55	N
ATOM	510	C	ARG	A	68	7.959	-13.205	4.964	1.00	17.61	C
ATOM	511	O	ARG	A	68	8.256	-12.166	4.383	1.00	17.22	O
ATOM	512	N	PRO	A	69	8.877	-14.153	5.209	1.00	17.60	N
ATOM	513	CA	PRO	A	69	10.254	-13.896	4.771	1.00	18.02	C
ATOM	514	CB	PRO	A	69	11.036	-15.080	5.357	1.00	18.83	C
ATOM	515	CG	PRO	A	69	10.180	-15.514	6.539	1.00	19.69	C
ATOM	516	CD	PRO	A	69	8.793	-15.412	5.977	1.00	18.13	C
ATOM	517	C	PRO	A	69	10.481	-13.723	3.251	1.00	17.20	C
ATOM	518	O	PRO	A	69	11.544	-13.234	2.850	1.00	17.02	O
ATOM	519	N	LEU	A	70	9.503	-14.098	2.414	1.00	16.34	N
ATOM	520	CA	LEU	A	70	9.606	-13.778	0.985	1.00	15.96	C
ATOM	521	CB	LEU	A	70	8.468	-14.410	0.185	1.00	16.34	C
ATOM	522	CG	LEU	A	70	8.755	-15.829	-0.321	1.00	19.86	C
ATOM	523	CD1	LEU	A	70	8.815	-16.809	0.828	1.00	22.48	C
ATOM	524	CD2	LEU	A	70	7.669	-16.218	-1.308	1.00	21.49	C
ATOM	525	C	LEU	A	70	9.608	-12.276	0.731	1.00	14.54	C
ATOM	526	O	LEU	A	70	9.965	-11.853	-0.361	1.00	15.20	O
ATOM	527	N	SER	A	71	9.188	-11.487	1.722	1.00	13.51	N
ATOM	528	CA	SER	A	71	9.200	-10.014	1.624	1.00	14.79	C
ATOM	529	CB	SER	A	71	8.211	-9.408	2.624	1.00	14.76	C
ATOM	530	OG	SER	A	71	6.920	-9.946	2.432	1.00	19.09	O
ATOM	531	C	SER	A	71	10.550	-9.367	1.891	1.00	13.58	C
ATOM	532	O	SER	A	71	10.753	-8.216	1.533	1.00	13.44	O
ATOM	533	N	TYR	A	72	11.469	-10.086	2.540	1.00	13.81	N
ATOM	534	CA	TYR	A	72	12.729	-9.472	2.994	1.00	13.60	C
ATOM	535	CB	TYR	A	72	13.329	-10.252	4.171	1.00	14.71	C
ATOM	536	CG	TYR	A	72	12.553	-10.168	5.464	1.00	12.94	C
ATOM	537	CD1	TYR	A	72	12.266	-8.937	6.064	1.00	13.82	C
ATOM	538	CE1	TYR	A	72	11.587	-8.877	7.270	1.00	13.38	C
ATOM	539	CZ	TYR	A	72	11.183	-10.045	7.884	1.00	12.61	C
ATOM	540	OH	TYR	A	72	10.514	-9.929	9.067	1.00	15.65	O
ATOM	541	CE2	TYR	A	72	11.462	-11.272	7.320	1.00	13.90	C
ATOM	542	CD2	TYR	A	72	12.163	-11.316	6.122	1.00	12.90	C
ATOM	543	C	TYR	A	72	13.852	-9.223	1.969	1.00	14.07	C
ATOM	544	O	TYR	A	72	14.681	-8.359	2.209	1.00	14.07	O
ATOM	545	N	PRO	A	73	13.950	-10.022	0.885	1.00	15.01	N
ATOM	546	CA	PRO	A	73	15.057	-9.728	-0.024	1.00	15.59	C
ATOM	547	CB	PRO	A	73	14.766	-10.640	-1.217	1.00	15.64	C
ATOM	548	CG	PRO	A	73	14.149	-11.831	-0.575	1.00	17.21	C
ATOM	549	CD	PRO	A	73	13.205	-11.212	0.435	1.00	16.17	C
ATOM	550	C	PRO	A	73	15.148	-8.278	-0.458	1.00	15.42	C
ATOM	551	O	PRO	A	73	14.136	-7.634	-0.763	1.00	15.01	O
ATOM	552	N	GLN	A	74	16.376	-7.776	-0.484	1.00	14.91	N
ATOM	553	CA	GLN	A	74	16.687	-6.420	-0.946	1.00	15.51	C
ATOM	554	CB	GLN	A	74	16.264	-6.223	-2.408	1.00	16.44	C
ATOM	555	CG	GLN	A	74	16.882	-7.245	-3.386	1.00	19.27	C
ATOM	556	CD	GLN	A	74	15.910	-8.340	-3.803	1.00	24.79	C
ATOM	557	OE1	GLN	A	74	14.722	-8.088	-4.031	1.00	27.77	O
ATOM	558	NE2	GLN	A	74	16.414	-9.564	-3.926	1.00	26.38	N
ATOM	559	C	GLN	A	74	16.139	-5.310	-0.045	1.00	14.57	C
ATOM	560	O	GLN	A	74	16.047	-4.151	-0.436	1.00	15.13	O
ATOM	561	N	THR	A	75	15.809	-5.655	1.186	1.00	13.90	N
ATOM	562	CA	THR	A	75	15.355	-4.625	2.129	1.00	13.47	C
ATOM	563	CB	THR	A	75	14.808	-5.261	3.430	1.00	13.21	C
ATOM	564	OG1	THR	A	75	13.662	-6.056	3.114	1.00	13.74	O
ATOM	565	CG2	THR	A	75	14.403	-4.203	4.454	1.00	13.03	C
ATOM	566	C	THR	A	75	16.460	-3.634	2.453	1.00	13.15	C
ATOM	567	O	THR	A	75	17.611	-4.037	2.727	1.00	14.79	O
ATOM	568	N	ASP	A	76	16.123	-2.347	2.456	1.00	11.58	N
ATOM	569	CA	ASP	A	76	17.074	-1.281	2.806	1.00	11.85	C
ATOM	570	CB	ASP	A	76	16.955	-0.108	1.828	1.00	11.89	C

ATOM	571	CG	ASP	A	76	17.315	-0.498	0.424	1.00	14.34	C
ATOM	572	OD1	ASP	A	76	18.452	-0.984	0.219	1.00	17.00	O
ATOM	573	OD2	ASP	A	76	16.471	-0.378	-0.493	1.00	13.48	O
ATOM	574	C	ASP	A	76	16.887	-0.771	4.228	1.00	12.51	C
ATOM	575	O	ASP	A	76	17.849	-0.324	4.846	1.00	13.24	O
ATOM	576	N	VAL	A	77	15.639	-0.758	4.737	1.00	11.39	N
ATOM	577	CA	VAL	A	77	15.361	-0.284	6.085	1.00	11.27	C
ATOM	578	CB	VAL	A	77	15.200	1.263	6.135	1.00	10.92	C
ATOM	579	CG1	VAL	A	77	14.118	1.771	5.187	1.00	12.49	C
ATOM	580	CG2	VAL	A	77	14.955	1.766	7.562	1.00	12.84	C
ATOM	581	C	VAL	A	77	14.122	-1.007	6.562	1.00	11.19	C
ATOM	582	O	VAL	A	77	13.236	-1.321	5.747	1.00	10.58	O
ATOM	583	N	SER	A	78	14.079	-1.311	7.848	0.50	11.65	N
ATOM	584	CA	SER	A	78	12.903	-1.976	8.404	0.50	12.31	C
ATOM	585	CB	SER	A	78	13.223	-3.368	8.931	0.50	13.27	C
ATOM	586	OG	ASER	A	78	12.034	-4.011	9.347	0.50	14.27	O
ATOM	586	OG	BSER	A	78	13.799	-4.229	7.961	0.50	14.62	O
ATOM	587	C	SER	A	78	12.326	-1.155	9.534	0.50	12.86	C
ATOM	588	O	SER	A	78	13.053	-0.618	10.368	0.50	12.62	O
ATOM	589	N	LEU	A	79	11.006	-1.103	9.591	1.00	11.00	N
ATOM	590	CA	LEU	A	79	10.301	-0.542	10.721	1.00	11.67	C
ATOM	591	CB	LEU	A	79	9.103	0.295	10.283	1.00	11.27	C
ATOM	592	CG	LEU	A	79	9.342	1.390	9.249	1.00	13.99	C
ATOM	593	CD1	LEU	A	79	8.087	2.201	9.048	1.00	15.38	C
ATOM	594	CD2	LEU	A	79	10.467	2.331	9.646	1.00	15.86	C
ATOM	595	C	LEU	A	79	9.782	-1.691	11.543	1.00	10.67	C
ATOM	596	O	LEU	A	79	8.967	-2.466	11.060	1.00	12.30	O
ATOM	597	N	ILE	A	80	10.265	-1.817	12.772	1.00	11.03	N
ATOM	598	CA	ILE	A	80	9.671	-2.773	13.715	1.00	10.63	C
ATOM	599	CB	ILE	A	80	10.707	-3.483	14.588	1.00	10.95	C
ATOM	600	CG1	ILE	A	80	11.647	-4.331	13.716	1.00	11.97	C
ATOM	601	CD1	ILE	A	80	12.768	-5.045	14.503	1.00	12.36	C
ATOM	602	CG2	ILE	A	80	9.982	-4.373	15.630	1.00	11.76	C
ATOM	603	C	ILE	A	80	8.685	-2.013	14.569	1.00	10.30	C
ATOM	604	O	ILE	A	80	9.041	-1.054	15.245	1.00	10.00	O
ATOM	605	N	CYS	A	81	7.416	-2.406	14.484	1.00	9.99	N
ATOM	606	CA	CYS	A	81	6.332	-1.674	15.116	1.00	10.50	C
ATOM	607	CB	CYS	A	81	5.191	-1.489	14.115	1.00	10.19	C
ATOM	608	SG	CYS	A	81	5.666	-0.559	12.649	1.00	11.71	S
ATOM	609	C	CYS	A	81	5.777	-2.453	16.303	1.00	10.21	C
ATOM	610	O	CYS	A	81	5.626	-3.683	16.239	1.00	9.95	O
ATOM	611	N	PHE	A	82	5.382	-1.727	17.344	1.00	9.79	N
ATOM	612	CA	PHE	A	82	4.606	-2.301	18.446	1.00	10.00	C
ATOM	613	CB	PHE	A	82	5.527	-2.694	19.607	1.00	10.97	C
ATOM	614	CG	PHE	A	82	6.103	-1.504	20.336	1.00	10.42	C
ATOM	615	CD1	PHE	A	82	5.450	-0.961	21.425	1.00	10.28	C
ATOM	616	CE1	PHE	A	82	5.961	0.171	22.084	1.00	11.10	C
ATOM	617	CZ	PHE	A	82	7.121	0.775	21.612	1.00	11.17	C
ATOM	618	CE2	PHE	A	82	7.776	0.258	20.518	1.00	11.35	C
ATOM	619	CD2	PHE	A	82	7.267	-0.869	19.868	1.00	11.50	C
ATOM	620	C	PHE	A	82	3.620	-1.232	18.880	1.00	10.12	C
ATOM	621	O	PHE	A	82	3.858	-0.030	18.717	1.00	10.63	O
ATOM	622	N	SER	A	83	2.498	-1.656	19.437	1.00	10.02	N
ATOM	623	CA	SER	A	83	1.511	-0.707	19.921	1.00	10.02	C
ATOM	624	CB	SER	A	83	0.147	-1.362	19.964	1.00	10.25	C
ATOM	625	OG	SER	A	83	-0.717	-0.549	20.740	1.00	11.09	O
ATOM	626	C	SER	A	83	1.883	-0.201	21.306	1.00	10.65	C
ATOM	627	O	SER	A	83	2.181	-1.004	22.203	1.00	10.75	O
ATOM	628	N	LEU	A	84	1.821	1.118	21.476	1.00	10.96	N
ATOM	629	CA	LEU	A	84	2.132	1.734	22.768	1.00	11.88	C
ATOM	630	CB	LEU	A	84	2.192	3.255	22.657	1.00	13.00	C
ATOM	631	CG	LEU	A	84	3.389	3.776	21.852	1.00	12.58	C
ATOM	632	CD1	LEU	A	84	3.174	5.194	21.409	1.00	14.70	C
ATOM	633	CD2	LEU	A	84	4.675	3.653	22.610	1.00	12.21	C
ATOM	634	C	LEU	A	84	1.163	1.294	23.853	1.00	11.76	C
ATOM	635	O	LEU	A	84	1.446	1.479	25.047	1.00	13.30	O

ATOM	636	N	VAL	A	85	0.034	0.708	23.458	1.00	11.66	N
ATOM	637	CA	VAL	A	85	-0.939	0.222	24.448	1.00	12.51	C
ATOM	638	CB	VAL	A	85	-2.310	0.951	24.330	1.00	12.44	C
ATOM	639	CG1	VAL	A	85	-2.134	2.448	24.624	1.00	14.58	C
ATOM	640	CG2	VAL	A	85	-2.968	0.733	22.987	1.00	13.59	C
ATOM	641	C	VAL	A	85	-1.073	-1.298	24.433	1.00	12.30	C
ATOM	642	O	VAL	A	85	-2.069	-1.842	24.915	1.00	13.43	O
ATOM	643	N	SER	A	86	-0.040	-1.981	23.926	1.00	12.15	N
ATOM	644	CA	SER	A	86	0.044	-3.431	24.003	1.00	12.42	C
ATOM	645	CB	SER	A	86	-0.314	-4.125	22.683	1.00	12.04	C
ATOM	646	OG	SER	A	86	-0.086	-5.522	22.842	1.00	15.41	O
ATOM	647	C	SER	A	86	1.433	-3.844	24.475	1.00	12.50	C
ATOM	648	O	SER	A	86	2.351	-4.063	23.678	1.00	12.54	O
ATOM	649	N	PRO	A	87	1.598	-4.002	25.787	1.00	12.95	N
ATOM	650	CA	PRO	A	87	2.869	-4.514	26.290	1.00	13.29	C
ATOM	651	CB	PRO	A	87	2.627	-4.615	27.808	1.00	14.21	C
ATOM	652	CG	PRO	A	87	1.645	-3.503	28.077	1.00	13.86	C
ATOM	653	CD	PRO	A	87	0.686	-3.627	26.891	1.00	13.43	C
ATOM	654	C	PRO	A	87	3.327	-5.837	25.679	1.00	13.26	C
ATOM	655	O	PRO	A	87	4.533	-6.031	25.474	1.00	13.67	O
ATOM	656	N	ALA	A	88	2.391	-6.731	25.355	1.00	13.08	N
ATOM	657	CA	ALA	A	88	2.770	-7.984	24.699	1.00	13.15	C
ATOM	658	CB	ALA	A	88	1.537	-8.871	24.492	1.00	14.24	C
ATOM	659	C	ALA	A	88	3.470	-7.721	23.363	1.00	12.54	C
ATOM	660	O	ALA	A	88	4.464	-8.363	23.019	1.00	13.00	O
ATOM	661	N	SER	A	89	2.945	-6.767	22.595	1.00	12.21	N
ATOM	662	CA	SER	A	89	3.536	-6.455	21.290	1.00	12.15	C
ATOM	663	CB	SER	A	89	2.611	-5.546	20.457	1.00	12.88	C
ATOM	664	OG	SER	A	89	2.641	-4.174	20.869	1.00	11.46	O
ATOM	665	C	SER	A	89	4.922	-5.853	21.448	1.00	11.97	C
ATOM	666	O	SER	A	89	5.808	-6.046	20.615	1.00	11.89	O
ATOM	667	N	PHE	A	90	5.106	-5.126	22.540	1.00	11.56	N
ATOM	668	CA	PHE	A	90	6.397	-4.531	22.869	1.00	11.82	C
ATOM	669	CB	PHE	A	90	6.201	-3.565	24.049	1.00	12.37	C
ATOM	670	CG	PHE	A	90	7.438	-2.830	24.475	1.00	13.30	C
ATOM	671	CD1	PHE	A	90	8.122	-2.022	23.586	1.00	14.79	C
ATOM	672	CE1	PHE	A	90	9.243	-1.296	24.008	1.00	15.32	C
ATOM	673	CZ	PHE	A	90	9.664	-1.355	25.330	1.00	15.13	C
ATOM	674	CE2	PHE	A	90	8.993	-2.147	26.230	1.00	15.63	C
ATOM	675	CD2	PHE	A	90	7.865	-2.883	25.807	1.00	15.19	C
ATOM	676	C	PHE	A	90	7.422	-5.613	23.178	1.00	12.20	C
ATOM	677	O	PHE	A	90	8.534	-5.599	22.672	1.00	12.96	O
ATOM	678	N	GLU	A	91	7.033	-6.572	24.006	1.00	13.27	N
ATOM	679	CA	GLU	A	91	7.960	-7.637	24.327	1.00	14.67	C
ATOM	680	CB	GLU	A	91	7.403	-8.530	25.432	1.00	15.80	C
ATOM	681	CG	GLU	A	91	7.325	-7.839	26.801	1.00	21.18	C
ATOM	682	CD	GLU	A	91	8.683	-7.312	27.296	1.00	26.68	C
ATOM	683	OE1	GLU	A	91	8.728	-6.172	27.812	1.00	31.58	O
ATOM	684	OE2	GLU	A	91	9.705	-8.023	27.166	1.00	29.55	O
ATOM	685	C	GLU	A	91	8.341	-8.442	23.087	1.00	13.24	C
ATOM	686	O	GLU	A	91	9.480	-8.870	22.961	1.00	14.31	O
ATOM	687	N	ASN	A	92	7.402	-8.613	22.151	1.00	11.84	N
ATOM	688	CA	ASN	A	92	7.744	-9.293	20.909	1.00	12.41	C
ATOM	689	CB	ASN	A	92	6.493	-9.684	20.145	1.00	13.20	C
ATOM	690	CG	ASN	A	92	5.894	-10.946	20.686	1.00	15.85	C
ATOM	691	OD1	ASN	A	92	6.617	-11.853	21.123	1.00	17.38	O
ATOM	692	ND2	ASN	A	92	4.571	-11.029	20.675	1.00	17.83	N
ATOM	693	C	ASN	A	92	8.718	-8.579	19.986	1.00	11.68	C
ATOM	694	O	ASN	A	92	9.267	-9.213	19.093	1.00	12.11	O
ATOM	695	N	VAL	A	93	8.950	-7.282	20.203	1.00	12.44	N
ATOM	696	CA	VAL	A	93	9.993	-6.580	19.454	1.00	13.31	C
ATOM	697	CB	VAL	A	93	10.151	-5.140	19.954	1.00	13.15	C
ATOM	698	CG1	VAL	A	93	11.439	-4.514	19.419	1.00	13.98	C
ATOM	699	CG2	VAL	A	93	8.918	-4.329	19.575	1.00	13.99	C
ATOM	700	C	VAL	A	93	11.312	-7.329	19.630	1.00	14.01	C
ATOM	701	O	VAL	A	93	12.030	-7.582	18.672	1.00	15.10	O

ATOM	702	N	ARG	A	94	11.611	-7.711	20.867	1.00	15.11	N
ATOM	703	CA	ARG	A	94	12.873	-8.371	21.160	1.00	16.16	C
ATOM	704	CB	ARG	A	94	13.326	-7.993	22.575	1.00	17.62	C
ATOM	705	CG	ARG	A	94	14.632	-8.627	22.959	1.00	21.98	C
ATOM	706	CD	ARG	A	94	15.816	-7.740	22.632	1.00	28.79	C
ATOM	707	NE	ARG	A	94	16.859	-8.516	21.966	1.00	34.14	N
ATOM	708	CZ	ARG	A	94	18.164	-8.362	22.164	1.00	35.82	C
ATOM	709	NH1	ARG	A	94	18.611	-7.460	23.024	1.00	37.30	N
ATOM	710	NH2	ARG	A	94	19.025	-9.129	21.504	1.00	38.38	N
ATOM	711	C	ARG	A	94	12.741	-9.883	20.991	1.00	15.53	C
ATOM	712	O	ARG	A	94	13.679	-10.550	20.556	1.00	16.61	O
ATOM	713	N	ALA	A	95	11.579	-10.433	21.331	1.00	13.61	N
ATOM	714	CA	ALA	A	95	11.399	-11.878	21.304	1.00	13.39	C
ATOM	715	CB	ALA	A	95	10.156	-12.269	22.125	1.00	13.49	C
ATOM	716	C	ALA	A	95	11.315	-12.453	19.892	1.00	13.50	C
ATOM	717	O	ALA	A	95	11.737	-13.575	19.626	1.00	14.22	O
ATOM	718	N	LYS	A	96	10.747	-11.661	18.982	1.00	13.38	N
ATOM	719	CA	LYS	A	96	10.404	-12.148	17.652	1.00	12.97	C
ATOM	720	CB	LYS	A	96	8.876	-12.266	17.478	1.00	13.16	C
ATOM	721	CG	LYS	A	96	8.473	-12.789	16.104	1.00	14.36	C
ATOM	722	CD	LYS	A	96	6.995	-13.121	16.024	1.00	15.01	C
ATOM	723	CE	LYS	A	96	6.677	-13.705	14.652	1.00	18.16	C
ATOM	724	NZ	LYS	A	96	5.205	-13.960	14.468	1.00	18.60	N
ATOM	725	C	LYS	A	96	10.973	-11.275	16.541	1.00	12.23	C
ATOM	726	O	LYS	A	96	11.650	-11.780	15.618	1.00	12.58	O
ATOM	727	N	TRP	A	97	10.686	-9.974	16.591	1.00	11.57	N
ATOM	728	CA	TRP	A	97	10.922	-9.147	15.394	1.00	11.61	C
ATOM	729	CB	TRP	A	97	10.047	-7.883	15.409	1.00	10.66	C
ATOM	730	CG	TRP	A	97	8.634	-8.324	15.386	1.00	10.77	C
ATOM	731	CD1	TRP	A	97	7.712	-8.217	16.400	1.00	10.78	C
ATOM	732	NE1	TRP	A	97	6.550	-8.851	16.035	1.00	11.23	N
ATOM	733	CE2	TRP	A	97	6.711	-9.401	14.788	1.00	10.82	C
ATOM	734	CD2	TRP	A	97	8.019	-9.110	14.357	1.00	10.00	C
ATOM	735	CE3	TRP	A	97	8.445	-9.577	13.101	1.00	10.71	C
ATOM	736	CZ3	TRP	A	97	7.569	-10.294	12.328	1.00	11.15	C
ATOM	737	CH2	TRP	A	97	6.282	-10.600	12.796	1.00	11.59	C
ATOM	738	CZ2	TRP	A	97	5.830	-10.161	14.011	1.00	11.63	C
ATOM	739	C	TRP	A	97	12.370	-8.859	15.123	1.00	12.18	C
ATOM	740	O	TRP	A	97	12.834	-9.005	13.987	1.00	11.96	O
ATOM	741	N	TYR	A	98	13.096	-8.446	16.161	1.00	12.70	N
ATOM	742	CA	TYR	A	98	14.528	-8.222	16.017	1.00	14.03	C
ATOM	743	CB	TYR	A	98	15.122	-7.749	17.346	1.00	16.61	C
ATOM	744	CG	TYR	A	98	16.633	-7.603	17.329	1.00	20.30	C
ATOM	745	CD1	TYR	A	98	17.274	-6.824	16.355	1.00	22.77	C
ATOM	746	CE1	TYR	A	98	18.662	-6.703	16.330	1.00	24.51	C
ATOM	747	CZ	TYR	A	98	19.411	-7.347	17.290	1.00	23.70	C
ATOM	748	OH	TYR	A	98	20.789	-7.224	17.270	1.00	25.67	O
ATOM	749	CE2	TYR	A	98	18.809	-8.115	18.270	1.00	24.56	C
ATOM	750	CD2	TYR	A	98	17.413	-8.242	18.284	1.00	25.03	C
ATOM	751	C	TYR	A	98	15.272	-9.443	15.467	1.00	13.19	C
ATOM	752	O	TYR	A	98	15.981	-9.316	14.483	1.00	12.70	O
ATOM	753	N	PRO	A	99	15.129	-10.619	16.100	1.00	13.11	N
ATOM	754	CA	PRO	A	99	15.862	-11.748	15.514	1.00	13.92	C
ATOM	755	CB	PRO	A	99	15.605	-12.915	16.491	1.00	14.48	C
ATOM	756	CG	PRO	A	99	14.426	-12.507	17.292	1.00	14.24	C
ATOM	757	CD	PRO	A	99	14.448	-10.992	17.356	1.00	13.23	C
ATOM	758	C	PRO	A	99	15.431	-12.083	14.087	1.00	13.28	C
ATOM	759	O	PRO	A	99	16.284	-12.423	13.259	1.00	14.28	O
ATOM	760	N	GLU	A	100	14.141	-11.995	13.765	1.00	13.11	N
ATOM	761	CA	GLU	A	100	13.722	-12.301	12.402	1.00	12.73	C
ATOM	762	CB	GLU	A	100	12.195	-12.290	12.296	1.00	13.10	C
ATOM	763	CG	GLU	A	100	11.737	-12.685	10.893	1.00	13.83	C
ATOM	764	CD	GLU	A	100	10.257	-12.894	10.778	1.00	15.02	C
ATOM	765	OE1	GLU	A	100	9.716	-13.813	11.423	1.00	16.97	O
ATOM	766	OE2	GLU	A	100	9.637	-12.123	10.023	1.00	14.34	O
ATOM	767	C	GLU	A	100	14.306	-11.318	11.377	1.00	12.68	C

ATOM	768	O	GLU	A	100	14.843	-11.720	10.320	1.00	12.94	O
ATOM	769	N	VAL	A	101	14.219	-10.040	11.684	1.00	12.96	N
ATOM	770	CA	VAL	A	101	14.710	-9.022	10.761	1.00	13.65	C
ATOM	771	CB	VAL	A	101	14.283	-7.602	11.213	1.00	13.29	C
ATOM	772	CG1	VAL	A	101	14.986	-6.503	10.385	1.00	16.16	C
ATOM	773	CG2	VAL	A	101	12.773	-7.457	11.127	1.00	13.75	C
ATOM	774	C	VAL	A	101	16.220	-9.148	10.611	1.00	14.67	C
ATOM	775	O	VAL	A	101	16.722	-9.083	9.509	1.00	15.13	O
ATOM	776	N	ARG	A	102	16.932	-9.347	11.713	1.00	15.37	N
ATOM	777	CA	ARG	A	102	18.398	-9.427	11.659	1.00	16.75	C
ATOM	778	CB	ARG	A	102	18.965	-9.395	13.087	1.00	17.79	C
ATOM	779	CG	ARG	A	102	20.399	-8.843	13.205	1.00	21.62	C
ATOM	780	CD	ARG	A	102	20.645	-7.684	12.236	1.00	23.24	C
ATOM	781	NE	ARG	A	102	20.443	-6.323	12.755	1.00	23.20	N
ATOM	782	CZ	ARG	A	102	19.984	-5.315	12.006	1.00	23.24	C
ATOM	783	NH1	ARG	A	102	19.599	-5.551	10.751	1.00	23.30	N
ATOM	784	NH2	ARG	A	102	19.860	-4.088	12.512	1.00	23.94	N
ATOM	785	C	ARG	A	102	18.835	-10.683	10.907	1.00	16.87	C
ATOM	786	O	ARG	A	102	19.842	-10.671	10.199	1.00	18.30	O
ATOM	787	N	HIS	A	103	18.058	-11.758	11.009	1.00	16.43	N
ATOM	788	CA	HIS	A	103	18.377	-12.966	10.286	1.00	17.48	C
ATOM	789	CB	HIS	A	103	17.492	-14.125	10.723	1.00	17.59	C
ATOM	790	CG	HIS	A	103	17.809	-15.400	10.004	1.00	21.53	C
ATOM	791	ND1	HIS	A	103	16.983	-15.945	9.049	1.00	24.85	N
ATOM	792	CE1	HIS	A	103	17.538	-17.042	8.559	1.00	26.32	C
ATOM	793	NE2	HIS	A	103	18.706	-17.213	9.148	1.00	26.17	N
ATOM	794	CD2	HIS	A	103	18.906	-16.193	10.047	1.00	23.34	C
ATOM	795	C	HIS	A	103	18.249	-12.767	8.783	1.00	16.88	C
ATOM	796	O	HIS	A	103	19.135	-13.160	8.025	1.00	17.24	O
ATOM	797	N	HIS	A	104	17.139	-12.181	8.345	1.00	16.36	N
ATOM	798	CA	HIS	A	104	16.870	-12.052	6.925	1.00	16.24	C
ATOM	799	CB	HIS	A	104	15.363	-12.094	6.675	1.00	16.12	C
ATOM	800	CG	HIS	A	104	14.758	-13.433	6.940	1.00	15.16	C
ATOM	801	ND1	HIS	A	104	14.797	-14.464	6.024	1.00	17.29	N
ATOM	802	CE1	HIS	A	104	14.187	-15.518	6.534	1.00	16.02	C
ATOM	803	NE2	HIS	A	104	13.770	-15.213	7.751	1.00	16.75	N
ATOM	804	CD2	HIS	A	104	14.099	-13.909	8.021	1.00	16.36	C
ATOM	805	C	HIS	A	104	17.466	-10.812	6.294	1.00	16.71	C
ATOM	806	O	HIS	A	104	17.629	-10.759	5.066	1.00	17.06	O
ATOM	807	N	CYS	A	105	17.770	-9.821	7.128	1.00	16.06	N
ATOM	808	CA	CYS	A	105	18.278	-8.524	6.690	1.00	16.59	C
ATOM	809	CB	CYS	A	105	17.197	-7.448	6.805	1.00	15.75	C
ATOM	810	SG	CYS	A	105	15.584	-7.864	6.090	1.00	17.37	S
ATOM	811	C	CYS	A	105	19.456	-8.140	7.573	1.00	18.00	C
ATOM	812	O	CYS	A	105	19.388	-7.172	8.319	1.00	17.69	O
ATOM	813	N	PRO	A	106	20.554	-8.899	7.482	1.00	18.97	N
ATOM	814	CA	PRO	A	106	21.656	-8.676	8.409	1.00	20.28	C
ATOM	815	CB	PRO	A	106	22.677	-9.754	8.005	1.00	20.67	C
ATOM	816	CG	PRO	A	106	22.345	-10.089	6.604	1.00	19.73	C
ATOM	817	CD	PRO	A	106	20.853	-9.998	6.545	1.00	19.37	C
ATOM	818	C	PRO	A	106	22.299	-7.282	8.418	1.00	22.24	C
ATOM	819	O	PRO	A	106	22.809	-6.871	9.462	1.00	24.24	O
ATOM	820	N	ASN	A	107	22.276	-6.559	7.304	1.00	23.21	N
ATOM	821	CA	ASN	A	107	23.015	-5.292	7.241	1.00	24.66	C
ATOM	822	CB	ASN	A	107	23.979	-5.303	6.062	1.00	26.10	C
ATOM	823	CG	ASN	A	107	25.043	-6.324	6.207	1.00	27.68	C
ATOM	824	OD1	ASN	A	107	25.261	-7.119	5.296	1.00	29.64	O
ATOM	825	ND2	ASN	A	107	25.713	-6.338	7.362	1.00	29.00	N
ATOM	826	C	ASN	A	107	22.139	-4.056	7.158	1.00	24.72	C
ATOM	827	O	ASN	A	107	22.586	-2.999	6.709	1.00	26.29	O
ATOM	828	N	THR	A	108	20.910	-4.195	7.632	1.00	23.11	N
ATOM	829	CA	THR	A	108	19.828	-3.246	7.386	1.00	22.12	C
ATOM	830	CB	THR	A	108	18.561	-4.066	7.011	1.00	22.14	C
ATOM	831	OG1	THR	A	108	18.807	-4.802	5.813	1.00	25.06	O
ATOM	832	CG2	THR	A	108	17.318	-3.190	6.815	1.00	24.17	C
ATOM	833	C	THR	A	108	19.525	-2.445	8.644	1.00	19.74	C

ATOM	834	O	THR	A	108	19.311	-3.049	9.695	1.00	19.19	O
ATOM	835	N	PRO	A	109	19.470	-1.102	8.550	1.00	16.40	N
ATOM	836	CA	PRO	A	109	19.031	-0.347	9.722	1.00	15.60	C
ATOM	837	CB	PRO	A	109	19.143	1.114	9.282	1.00	15.76	C
ATOM	838	CG	PRO	A	109	19.124	1.064	7.757	1.00	16.60	C
ATOM	839	CD	PRO	A	109	19.824	-0.211	7.423	1.00	15.88	C
ATOM	840	C	PRO	A	109	17.592	-0.663	10.112	1.00	15.49	C
ATOM	841	O	PRO	A	109	16.750	-0.973	9.252	1.00	14.97	O
ATOM	842	N	ILE	A	110	17.340	-0.577	11.416	1.00	14.61	N
ATOM	843	CA	ILE	A	110	16.030	-0.790	11.999	1.00	14.49	C
ATOM	844	CB	ILE	A	110	16.068	-1.905	13.054	1.00	15.01	C
ATOM	845	CG1	ILE	A	110	16.364	-3.245	12.393	1.00	15.73	C
ATOM	846	CD1	ILE	A	110	16.593	-4.392	13.368	1.00	19.39	C
ATOM	847	CG2	ILE	A	110	14.742	-1.941	13.870	1.00	15.94	C
ATOM	848	C	ILE	A	110	15.584	0.466	12.677	1.00	13.80	C
ATOM	849	O	ILE	A	110	16.362	1.088	13.417	1.00	14.50	O
ATOM	850	N	ILE	A	111	14.346	0.869	12.420	1.00	13.72	N
ATOM	851	CA	ILE	A	111	13.734	1.920	13.209	1.00	13.09	C
ATOM	852	CB	ILE	A	111	13.175	3.006	12.318	1.00	13.97	C
ATOM	853	CG1	ILE	A	111	14.392	3.647	11.608	1.00	18.10	C
ATOM	854	CD1	ILE	A	111	14.161	4.925	10.924	1.00	22.57	C
ATOM	855	CG2	ILE	A	111	12.345	4.022	13.119	1.00	14.53	C
ATOM	856	C	ILE	A	111	12.653	1.279	14.075	1.00	12.70	C
ATOM	857	O	ILE	A	111	11.822	0.520	13.559	1.00	12.11	O
ATOM	858	N	LEU	A	112	12.674	1.558	15.377	1.00	12.35	N
ATOM	859	CA	LEU	A	112	11.647	1.028	16.283	1.00	11.30	C
ATOM	860	CB	LEU	A	112	12.251	0.756	17.660	1.00	12.08	C
ATOM	861	CG	LEU	A	112	11.241	0.181	18.651	1.00	11.67	C
ATOM	862	CD1	LEU	A	112	10.703	-1.171	18.185	1.00	13.05	C
ATOM	863	CD2	LEU	A	112	11.879	0.073	20.040	1.00	13.53	C
ATOM	864	C	LEU	A	112	10.535	2.043	16.383	1.00	11.41	C
ATOM	865	O	LEU	A	112	10.776	3.209	16.709	1.00	11.83	O
ATOM	866	N	VAL	A	113	9.301	1.612	16.081	1.00	11.11	N
ATOM	867	CA	VAL	A	113	8.166	2.517	15.997	1.00	11.25	C
ATOM	868	CB	VAL	A	113	7.568	2.562	14.555	1.00	11.45	C
ATOM	869	CG1	VAL	A	113	6.389	3.544	14.464	1.00	12.35	C
ATOM	870	CG2	VAL	A	113	8.653	2.966	13.533	1.00	11.84	C
ATOM	871	C	VAL	A	113	7.081	2.122	16.997	1.00	10.54	C
ATOM	872	O	VAL	A	113	6.545	1.010	16.912	1.00	10.85	O
ATOM	873	N	GLY	A	114	6.795	3.003	17.953	1.00	10.72	N
ATOM	874	CA	GLY	A	114	5.626	2.834	18.831	1.00	10.62	C
ATOM	875	C	GLY	A	114	4.412	3.434	18.152	1.00	11.24	C
ATOM	876	O	GLY	A	114	4.420	4.609	17.765	1.00	12.45	O
ATOM	877	N	THR	A	115	3.365	2.633	17.972	1.00	10.04	N
ATOM	878	CA	THR	A	115	2.156	3.038	17.267	1.00	10.46	C
ATOM	879	CB	THR	A	115	1.711	1.939	16.291	1.00	10.43	C
ATOM	880	OG1	THR	A	115	1.467	0.771	17.063	1.00	11.87	O
ATOM	881	CG2	THR	A	115	2.800	1.638	15.237	1.00	11.11	C
ATOM	882	C	THR	A	115	1.012	3.319	18.250	1.00	10.45	C
ATOM	883	O	THR	A	115	1.107	2.999	19.452	1.00	11.66	O
ATOM	884	N	LYS	A	116	-0.075	3.847	17.703	1.00	10.99	N
ATOM	885	CA	LYS	A	116	-1.291	4.184	18.452	1.00	11.62	C
ATOM	886	CB	LYS	A	116	-1.996	2.939	18.986	1.00	11.39	C
ATOM	887	CG	LYS	A	116	-2.250	1.895	17.905	1.00	10.85	C
ATOM	888	CD	LYS	A	116	-3.303	0.882	18.343	1.00	11.77	C
ATOM	889	CE	LYS	A	116	-3.395	-0.273	17.383	1.00	12.92	C
ATOM	890	NZ	LYS	A	116	-4.345	-1.319	17.904	1.00	11.97	N
ATOM	891	C	LYS	A	116	-0.999	5.163	19.574	1.00	12.31	C
ATOM	892	O	LYS	A	116	-1.557	5.062	20.670	1.00	12.60	O
ATOM	893	N	LEU	A	117	-0.160	6.147	19.268	1.00	12.97	N
ATOM	894	CA	LEU	A	117	0.210	7.158	20.249	1.00	14.06	C
ATOM	895	CB	LEU	A	117	1.129	8.182	19.571	1.00	13.64	C
ATOM	896	CG	LEU	A	117	1.521	9.443	20.352	1.00	14.58	C
ATOM	897	CD1	LEU	A	117	2.293	9.043	21.565	1.00	16.08	C
ATOM	898	CD2	LEU	A	117	2.368	10.351	19.496	1.00	16.13	C
ATOM	899	C	LEU	A	117	-1.014	7.851	20.851	1.00	13.77	C

ATOM	900	O	LEU	A	117	-0.992	8.227	22.041	1.00	15.02	O
ATOM	901	N	ASP	A	118	-2.055	8.024	20.047	1.00	13.75	N
ATOM	902	CA	ASP	A	118	-3.300	8.677	20.499	1.00	14.43	C
ATOM	903	CB	ASP	A	118	-4.271	8.865	19.330	1.00	14.70	C
ATOM	904	CG	ASP	A	118	-4.687	7.553	18.704	1.00	13.72	C
ATOM	905	OD1	ASP	A	118	-3.846	6.912	18.028	1.00	14.20	O
ATOM	906	OD2	ASP	A	118	-5.845	7.166	18.880	1.00	14.91	O
ATOM	907	C	ASP	A	118	-3.968	7.927	21.662	1.00	15.41	C
ATOM	908	O	ASP	A	118	-4.742	8.523	22.417	1.00	16.15	O
ATOM	909	N	LEU	A	119	-3.656	6.645	21.832	1.00	14.12	N
ATOM	910	CA	LEU	A	119	-4.297	5.853	22.873	1.00	14.41	C
ATOM	911	CB	LEU	A	119	-4.555	4.414	22.387	1.00	14.30	C
ATOM	912	CG	LEU	A	119	-5.555	4.218	21.247	1.00	14.14	C
ATOM	913	CD1	LEU	A	119	-5.721	2.728	20.924	1.00	14.99	C
ATOM	914	CD2	LEU	A	119	-6.917	4.854	21.569	1.00	14.15	C
ATOM	915	C	LEU	A	119	-3.542	5.866	24.191	1.00	15.31	C
ATOM	916	O	LEU	A	119	-4.098	5.437	25.213	1.00	15.58	O
ATOM	917	N	ARG	A	120	-2.294	6.333	24.191	1.00	15.71	N
ATOM	918	CA	ARG	A	120	-1.474	6.250	25.399	1.00	17.33	C
ATOM	919	CB	ARG	A	120	-0.005	6.632	25.178	1.00	16.86	C
ATOM	920	CG	ARG	A	120	0.811	6.546	26.491	1.00	16.99	C
ATOM	921	CD	ARG	A	120	2.296	6.428	26.273	1.00	16.57	C
ATOM	922	NE	ARG	A	120	2.819	7.584	25.564	1.00	17.42	N
ATOM	923	CZ	ARG	A	120	3.981	7.595	24.933	1.00	15.40	C
ATOM	924	NH1	ARG	A	120	4.746	6.503	24.897	1.00	15.17	N
ATOM	925	NH2	ARG	A	120	4.357	8.708	24.306	1.00	16.70	N
ATOM	926	C	ARG	A	120	-2.083	7.071	26.534	1.00	19.03	C
ATOM	927	O	ARG	A	120	-2.005	6.669	27.705	1.00	19.26	O
ATOM	928	N	ASP	A	121	-2.728	8.177	26.184	1.00	21.08	N
ATOM	929	CA	ASP	A	121	-3.414	8.964	27.211	1.00	23.85	C
ATOM	930	CB	ASP	A	121	-2.857	10.401	27.262	1.00	24.79	C
ATOM	931	CG	ASP	A	121	-3.016	11.149	25.950	1.00	28.79	C
ATOM	932	OD1	ASP	A	121	-2.641	12.344	25.899	1.00	33.35	O
ATOM	933	OD2	ASP	A	121	-3.514	10.559	24.974	1.00	32.10	O
ATOM	934	C	ASP	A	121	-4.953	8.886	27.147	1.00	24.43	C
ATOM	935	O	ASP	A	121	-5.637	9.704	27.777	1.00	26.11	O
ATOM	936	N	ASP	A	122	-5.477	7.881	26.427	1.00	23.95	N
ATOM	937	CA	ASP	A	122	-6.928	7.600	26.266	1.00	24.46	C
ATOM	938	CB	ASP	A	122	-7.177	6.663	25.068	1.00	24.33	C
ATOM	939	CG	ASP	A	122	-8.673	6.326	24.848	1.00	24.95	C
ATOM	940	OD1	ASP	A	122	-9.439	7.225	24.431	1.00	27.49	O
ATOM	941	OD2	ASP	A	122	-9.080	5.158	25.043	1.00	24.18	O
ATOM	942	C	ASP	A	122	-7.456	6.928	27.521	1.00	24.87	C
ATOM	943	O	ASP	A	122	-6.878	5.948	27.999	1.00	25.09	O
ATOM	944	N	LYS	A	123	-8.581	7.436	28.026	1.00	25.34	N
ATOM	945	CA	LYS	A	123	-9.105	7.041	29.328	1.00	25.83	C
ATOM	946	CB	LYS	A	123	-10.261	7.968	29.731	1.00	26.11	C
ATOM	947	CG	LYS	A	123	-9.831	9.405	30.013	1.00	28.80	C
ATOM	948	CD	LYS	A	123	-8.859	9.464	31.181	1.00	31.10	C
ATOM	949	CE	LYS	A	123	-8.804	10.839	31.826	1.00	33.04	C
ATOM	950	NZ	LYS	A	123	-8.546	10.688	33.283	1.00	33.17	N
ATOM	951	C	LYS	A	123	-9.525	5.585	29.406	1.00	25.13	C
ATOM	952	O	LYS	A	123	-9.068	4.857	30.289	1.00	25.32	O
ATOM	953	N	ASP	A	124	-10.366	5.152	28.467	1.00	25.18	N
ATOM	954	CA	ASP	A	124	-10.836	3.769	28.418	1.00	25.02	C
ATOM	955	CB	ASP	A	124	-11.861	3.580	27.302	1.00	25.90	C
ATOM	956	CG	ASP	A	124	-12.532	2.215	27.337	1.00	29.21	C
ATOM	957	OD1	ASP	A	124	-13.297	1.943	28.292	1.00	32.53	O
ATOM	958	OD2	ASP	A	124	-12.321	1.416	26.397	1.00	33.80	O
ATOM	959	C	ASP	A	124	-9.685	2.798	28.231	1.00	23.79	C
ATOM	960	O	ASP	A	124	-9.663	1.747	28.861	1.00	23.80	O
ATOM	961	N	THR	A	125	-8.723	3.160	27.374	1.00	22.32	N
ATOM	962	CA	THR	A	125	-7.581	2.290	27.125	1.00	20.76	C
ATOM	963	CB	THR	A	125	-6.716	2.824	25.974	1.00	20.70	C
ATOM	964	OG1	THR	A	125	-7.526	2.972	24.801	1.00	22.33	O
ATOM	965	CG2	THR	A	125	-5.602	1.832	25.674	1.00	19.54	C

ATOM	966	C	THR	A	125	-6.724	2.110	28.380	1.00	20.07	C
ATOM	967	O	THR	A	125	-6.333	0.993	28.712	1.00	18.81	O
ATOM	968	N	ILE	A	126	-6.459	3.207	29.085	1.00	19.39	N
ATOM	969	CA	ILE	A	126	-5.681	3.144	30.321	1.00	19.62	C
ATOM	970	CB	ILE	A	126	-5.364	4.532	30.887	1.00	19.40	C
ATOM	971	CG1	ILE	A	126	-4.396	5.268	29.960	1.00	19.35	C
ATOM	972	CD1	ILE	A	126	-4.362	6.779	30.214	1.00	22.93	C
ATOM	973	CG2	ILE	A	126	-4.727	4.420	32.273	1.00	21.54	C
ATOM	974	C	ILE	A	126	-6.419	2.285	31.363	1.00	19.93	C
ATOM	975	O	ILE	A	126	-5.806	1.451	32.015	1.00	19.00	O
ATOM	976	N	GLU	A	127	-7.738	2.483	31.467	1.00	21.26	N
ATOM	977	CA	GLU	A	127	-8.583	1.729	32.397	1.00	23.21	C
ATOM	978	CB	GLU	A	127	-10.031	2.206	32.318	1.00	23.25	C
ATOM	979	CG	GLU	A	127	-10.360	3.475	33.104	1.00	25.80	C
ATOM	980	CD	GLU	A	127	-11.848	3.834	33.054	1.00	26.81	C
ATOM	981	OE1	GLU	A	127	-12.633	3.114	32.387	1.00	32.68	O
ATOM	982	OE2	GLU	A	127	-12.233	4.835	33.697	1.00	31.64	O
ATOM	983	C	GLU	A	127	-8.542	0.238	32.083	1.00	22.25	C
ATOM	984	O	GLU	A	127	-8.324	-0.593	32.988	1.00	22.51	O
ATOM	985	N	LYS	A	128	-8.732	-0.099	30.801	1.00	22.02	N
ATOM	986	CA	LYS	A	128	-8.708	-1.499	30.345	1.00	23.13	C
ATOM	987	CB	LYS	A	128	-9.106	-1.614	28.864	1.00	23.41	C
ATOM	988	CG	LYS	A	128	-10.556	-1.981	28.639	1.00	26.63	C
ATOM	989	CD	LYS	A	128	-10.900	-2.054	27.154	1.00	27.10	C
ATOM	990	CE	LYS	A	128	-12.342	-2.503	26.951	1.00	32.09	C
ATOM	991	NZ	LYS	A	128	-12.819	-2.188	25.566	1.00	34.87	N
ATOM	992	C	LYS	A	128	-7.382	-2.201	30.607	1.00	21.64	C
ATOM	993	O	LYS	A	128	-7.351	-3.373	30.992	1.00	21.72	O
ATOM	994	N	LEU	A	129	-6.274	-1.492	30.390	1.00	19.65	N
ATOM	995	CA	LEU	A	129	-4.970	-2.031	30.719	1.00	18.39	C
ATOM	996	CB	LEU	A	129	-3.826	-1.147	30.159	1.00	17.50	C
ATOM	997	CG	LEU	A	129	-3.624	-1.209	28.638	1.00	17.68	C
ATOM	998	CD1	LEU	A	129	-2.705	-0.084	28.182	1.00	17.69	C
ATOM	999	CD2	LEU	A	129	-3.061	-2.566	28.208	1.00	19.43	C
ATOM	1000	C	LEU	A	129	-4.818	-2.218	32.220	1.00	18.22	C
ATOM	1001	O	LEU	A	129	-4.308	-3.243	32.668	1.00	17.93	O
ATOM	1002	N	LYS	A	130	-5.281	-1.239	32.999	1.00	18.30	N
ATOM	1003	CA	LYS	A	130	-5.138	-1.281	34.448	1.00	19.99	C
ATOM	1004	CB	LYS	A	130	-5.657	0.019	35.077	1.00	19.61	C
ATOM	1005	CG	LYS	A	130	-5.327	0.161	36.563	1.00	22.48	C
ATOM	1006	CD	LYS	A	130	-5.971	1.394	37.175	1.00	23.21	C
ATOM	1007	CE	LYS	A	130	-5.621	1.505	38.650	1.00	28.22	C
ATOM	1008	NZ	LYS	A	130	-6.438	2.560	39.305	1.00	31.69	N
ATOM	1009	C	LYS	A	130	-5.835	-2.500	35.066	1.00	19.55	C
ATOM	1010	O	LYS	A	130	-5.298	-3.110	35.978	1.00	20.66	O
ATOM	1011	N	GLU	A	131	-7.002	-2.844	34.534	1.00	19.98	N
ATOM	1012	CA	GLU	A	131	-7.762	-4.027	34.993	1.00	21.82	C
ATOM	1013	CB	GLU	A	131	-8.989	-4.225	34.127	1.00	22.37	C
ATOM	1014	CG	GLU	A	131	-10.065	-3.208	34.295	1.00	25.39	C
ATOM	1015	CD	GLU	A	131	-11.267	-3.528	33.423	1.00	27.20	C
ATOM	1016	OE1	GLU	A	131	-11.173	-4.443	32.570	1.00	28.48	O
ATOM	1017	OE2	GLU	A	131	-12.305	-2.868	33.584	1.00	32.18	O
ATOM	1018	C	GLU	A	131	-6.935	-5.302	34.910	1.00	22.56	C
ATOM	1019	O	GLU	A	131	-7.151	-6.244	35.678	1.00	22.48	O
ATOM	1020	N	LYS	A	132	-6.014	-5.334	33.944	1.00	23.05	N
ATOM	1021	CA	LYS	A	132	-5.088	-6.449	33.722	1.00	23.56	C
ATOM	1022	CB	LYS	A	132	-4.911	-6.675	32.218	1.00	23.83	C
ATOM	1023	CG	LYS	A	132	-6.192	-6.985	31.466	1.00	24.91	C
ATOM	1024	CD	LYS	A	132	-5.929	-7.130	29.978	1.00	27.51	C
ATOM	1025	CE	LYS	A	132	-7.172	-7.624	29.260	1.00	28.92	C
ATOM	1026	NZ	LYS	A	132	-6.986	-7.606	27.778	1.00	31.70	N
ATOM	1027	C	LYS	A	132	-3.712	-6.226	34.363	1.00	23.22	C
ATOM	1028	O	LYS	A	132	-2.764	-7.003	34.114	1.00	23.85	O
ATOM	1029	N	LYS	A	133	-3.600	-5.181	35.185	1.00	23.67	N
ATOM	1030	CA	LYS	A	133	-2.353	-4.822	35.870	1.00	23.68	C
ATOM	1031	CB	LYS	A	133	-1.865	-5.956	36.793	1.00	24.89	C

ATOM	1032	CG	LYS	A	133	-2.776	-6.265	37.989	1.00	26.71	C
ATOM	1033	CD	LYS	A	133	-1.990	-6.993	39.082	1.00	30.39	C
ATOM	1034	CE	LYS	A	133	-2.797	-7.142	40.374	1.00	32.61	C
ATOM	1035	NZ	LYS	A	133	-1.962	-7.691	41.489	1.00	34.46	N
ATOM	1036	C	LYS	A	133	-1.263	-4.440	34.857	1.00	23.39	C
ATOM	1037	O	LYS	A	133	-0.074	-4.726	35.058	1.00	23.85	O
ATOM	1038	N	LEU	A	134	-1.691	-3.794	33.774	1.00	21.93	N
ATOM	1039	CA	LEU	A	134	-0.784	-3.320	32.721	1.00	20.83	C
ATOM	1040	CB	LEU	A	134	-1.149	-3.965	31.395	1.00	20.76	C
ATOM	1041	CG	LEU	A	134	-1.077	-5.492	31.338	1.00	22.55	C
ATOM	1042	CD1	LEU	A	134	-1.670	-5.959	30.040	1.00	22.21	C
ATOM	1043	CD2	LEU	A	134	0.361	-5.965	31.506	1.00	23.61	C
ATOM	1044	C	LEU	A	134	-0.889	-1.817	32.584	1.00	20.05	C
ATOM	1045	O	LEU	A	134	-1.841	-1.208	33.051	1.00	19.28	O
ATOM	1046	N	ILE	A	135	0.086	-1.233	31.891	1.00	18.39	N
ATOM	1047	CA	ILE	A	135	0.162	0.198	31.682	1.00	18.18	C
ATOM	1048	CB	ILE	A	135	1.028	0.856	32.786	1.00	19.88	C
ATOM	1049	CG1	ILE	A	135	0.682	2.322	32.955	1.00	22.25	C
ATOM	1050	CD1	ILE	A	135	0.371	2.670	34.398	1.00	26.92	C
ATOM	1051	CG2	ILE	A	135	2.501	0.581	32.588	1.00	20.81	C
ATOM	1052	C	ILE	A	135	0.731	0.409	30.281	1.00	16.57	C
ATOM	1053	O	ILE	A	135	1.407	-0.483	29.770	1.00	16.98	O
ATOM	1054	N	PRO	A	136	0.401	1.539	29.638	1.00	15.39	N
ATOM	1055	CA	PRO	A	136	1.034	1.824	28.342	1.00	14.59	C
ATOM	1056	CB	PRO	A	136	0.468	3.184	27.968	1.00	14.59	C
ATOM	1057	CG	PRO	A	136	-0.852	3.252	28.667	1.00	15.57	C
ATOM	1058	CD	PRO	A	136	-0.598	2.568	29.986	1.00	15.78	C
ATOM	1059	C	PRO	A	136	2.550	1.891	28.388	1.00	15.06	C
ATOM	1060	O	PRO	A	136	3.148	2.161	29.436	1.00	15.72	O
ATOM	1061	N	ILE	A	137	3.165	1.615	27.245	1.00	13.89	N
ATOM	1062	CA	ILE	A	137	4.600	1.756	27.119	1.00	13.54	C
ATOM	1063	CB	ILE	A	137	5.107	1.026	25.857	1.00	13.19	C
ATOM	1064	CG1	ILE	A	137	4.725	-0.467	25.896	1.00	14.05	C
ATOM	1065	CD1	ILE	A	137	5.097	-1.217	27.175	1.00	16.84	C
ATOM	1066	CG2	ILE	A	137	6.610	1.212	25.703	1.00	13.91	C
ATOM	1067	C	ILE	A	137	4.959	3.237	27.048	1.00	13.44	C
ATOM	1068	O	ILE	A	137	4.387	4.003	26.304	1.00	13.44	O
ATOM	1069	N	THR	A	138	5.933	3.618	27.867	1.00	13.55	N
ATOM	1070	CA	THR	A	138	6.369	5.011	27.973	1.00	13.78	C
ATOM	1071	CB	THR	A	138	6.930	5.314	29.379	1.00	14.14	C
ATOM	1072	OG1	THR	A	138	8.163	4.631	29.537	1.00	15.81	O
ATOM	1073	CG2	THR	A	138	5.955	4.861	30.469	1.00	15.66	C
ATOM	1074	C	THR	A	138	7.490	5.285	26.980	1.00	13.92	C
ATOM	1075	O	THR	A	138	8.163	4.354	26.514	1.00	14.34	O
ATOM	1076	N	TYR	A	139	7.703	6.555	26.654	1.00	15.03	N
ATOM	1077	CA	TYR	A	139	8.784	6.909	25.761	1.00	15.03	C
ATOM	1078	CB	TYR	A	139	8.822	8.424	25.553	1.00	16.21	C
ATOM	1079	CG	TYR	A	139	9.806	8.858	24.507	1.00	17.32	C
ATOM	1080	CD1	TYR	A	139	9.384	9.113	23.206	1.00	18.21	C
ATOM	1081	CE1	TYR	A	139	10.287	9.508	22.221	1.00	18.42	C
ATOM	1082	CZ	TYR	A	139	11.623	9.663	22.535	1.00	19.94	C
ATOM	1083	OH	TYR	A	139	12.503	10.051	21.534	1.00	21.13	O
ATOM	1084	CE2	TYR	A	139	12.079	9.414	23.821	1.00	19.28	C
ATOM	1085	CD2	TYR	A	139	11.159	9.015	24.815	1.00	18.75	C
ATOM	1086	C	TYR	A	139	10.168	6.360	26.207	1.00	15.12	C
ATOM	1087	O	TYR	A	139	10.919	5.797	25.404	1.00	14.95	O
ATOM	1088	N	PRO	A	140	10.540	6.530	27.497	1.00	14.94	N
ATOM	1089	CA	PRO	A	140	11.825	5.963	27.923	1.00	15.05	C
ATOM	1090	CB	PRO	A	140	11.962	6.437	29.391	1.00	15.31	C
ATOM	1091	CG	PRO	A	140	11.130	7.672	29.439	1.00	16.93	C
ATOM	1092	CD	PRO	A	140	9.921	7.297	28.591	1.00	15.16	C
ATOM	1093	C	PRO	A	140	11.934	4.454	27.796	1.00	14.68	C
ATOM	1094	O	PRO	A	140	13.010	3.925	27.532	1.00	14.99	O
ATOM	1095	N	GLN	A	141	10.821	3.755	27.992	1.00	14.91	N
ATOM	1096	CA	GLN	A	141	10.831	2.321	27.795	1.00	15.32	C
ATOM	1097	CB	GLN	A	141	9.504	1.709	28.260	1.00	15.84	C

ATOM	1098	CG	GLN	A	141	9.408	1.695	29.805	1.00	17.42	C
ATOM	1099	CD	GLN	A	141	8.033	1.372	30.337	1.00	21.38	C
ATOM	1100	OE1	GLN	A	141	7.048	1.409	29.613	1.00	19.25	O
ATOM	1101	NE2	GLN	A	141	7.959	1.069	31.638	1.00	24.15	N
ATOM	1102	C	GLN	A	141	11.100	1.970	26.329	1.00	14.71	C
ATOM	1103	O	GLN	A	141	11.888	1.070	26.028	1.00	14.43	O
ATOM	1104	N	GLY	A	142	10.450	2.704	25.432	1.00	14.86	N
ATOM	1105	CA	GLY	A	142	10.638	2.471	23.988	1.00	14.98	C
ATOM	1106	C	GLY	A	142	12.078	2.752	23.598	1.00	15.62	C
ATOM	1107	O	GLY	A	142	12.702	1.979	22.871	1.00	15.98	O
ATOM	1108	N	LEU	A	143	12.633	3.843	24.124	1.00	15.83	N
ATOM	1109	CA	LEU	A	143	14.029	4.179	23.832	1.00	16.50	C
ATOM	1110	CB	LEU	A	143	14.401	5.527	24.425	1.00	17.32	C
ATOM	1111	CG	LEU	A	143	15.810	6.009	24.080	1.00	19.47	C
ATOM	1112	CD1	LEU	A	143	16.041	6.135	22.575	1.00	21.78	C
ATOM	1113	CD2	LEU	A	143	15.999	7.332	24.763	1.00	20.72	C
ATOM	1114	C	LEU	A	143	14.988	3.100	24.327	1.00	15.76	C
ATOM	1115	O	LEU	A	143	15.939	2.738	23.623	1.00	15.95	O
ATOM	1116	N	ALA	A	144	14.747	2.576	25.529	1.00	15.60	N
ATOM	1117	CA	ALA	A	144	15.579	1.489	26.048	1.00	15.89	C
ATOM	1118	CB	ALA	A	144	15.176	1.128	27.474	1.00	16.77	C
ATOM	1119	C	ALA	A	144	15.525	0.242	25.171	1.00	15.42	C
ATOM	1120	O	ALA	A	144	16.529	-0.409	24.935	1.00	15.80	O
ATOM	1121	N	MET	A	145	14.329	-0.085	24.695	1.00	15.04	N
ATOM	1122	CA	MET	A	145	14.169	-1.232	23.817	1.00	15.55	C
ATOM	1123	CB	MET	A	145	12.690	-1.461	23.547	1.00	15.20	C
ATOM	1124	CG	MET	A	145	12.399	-2.650	22.624	1.00	17.01	C
ATOM	1125	SD	MET	A	145	12.861	-4.261	23.305	1.00	18.66	S
ATOM	1126	CE	MET	A	145	11.458	-4.668	24.331	1.00	19.29	C
ATOM	1127	C	MET	A	145	14.920	-1.014	22.503	1.00	15.46	C
ATOM	1128	O	MET	A	145	15.556	-1.943	21.992	1.00	15.89	O
ATOM	1129	N	ALA	A	146	14.853	0.203	21.971	1.00	15.90	N
ATOM	1130	CA	ALA	A	146	15.560	0.543	20.729	1.00	16.14	C
ATOM	1131	CB	ALA	A	146	15.283	1.978	20.323	1.00	16.29	C
ATOM	1132	C	ALA	A	146	17.057	0.303	20.887	1.00	17.01	C
ATOM	1133	O	ALA	A	146	17.708	-0.245	19.997	1.00	16.87	O
ATOM	1134	N	LYS	A	147	17.605	0.702	22.028	1.00	18.16	N
ATOM	1135	CA	LYS	A	147	19.021	0.439	22.286	1.00	19.48	C
ATOM	1136	CB	LYS	A	147	19.467	1.115	23.577	1.00	19.86	C
ATOM	1137	CG	LYS	A	147	19.450	2.619	23.513	1.00	22.38	C
ATOM	1138	CD	LYS	A	147	19.918	3.218	24.835	1.00	25.84	C
ATOM	1139	CE	LYS	A	147	19.857	4.739	24.809	1.00	27.13	C
ATOM	1140	NZ	LYS	A	147	20.013	5.289	26.179	1.00	29.52	N
ATOM	1141	C	LYS	A	147	19.341	-1.050	22.364	1.00	20.06	C
ATOM	1142	O	LYS	A	147	20.356	-1.501	21.828	1.00	19.99	O
ATOM	1143	N	GLU	A	148	18.486	-1.812	23.035	1.00	20.59	N
ATOM	1144	CA	GLU	A	148	18.694	-3.269	23.159	1.00	22.17	C
ATOM	1145	CB	GLU	A	148	17.576	-3.927	23.965	1.00	22.06	C
ATOM	1146	CG	GLU	A	148	17.501	-3.578	25.446	1.00	24.20	C
ATOM	1147	CD	GLU	A	148	16.383	-4.326	26.153	1.00	25.70	C
ATOM	1148	OE1	GLU	A	148	16.031	-5.439	25.711	1.00	31.32	O
ATOM	1149	OE2	GLU	A	148	15.838	-3.800	27.146	1.00	31.68	O
ATOM	1150	C	GLU	A	148	18.773	-3.977	21.809	1.00	21.83	C
ATOM	1151	O	GLU	A	148	19.475	-4.985	21.659	1.00	23.00	O
ATOM	1152	N	ILE	A	149	18.024	-3.473	20.833	1.00	20.08	N
ATOM	1153	CA	ILE	A	149	17.978	-4.121	19.519	1.00	19.47	C
ATOM	1154	CB	ILE	A	149	16.519	-4.318	18.997	1.00	18.75	C
ATOM	1155	CG1	ILE	A	149	15.837	-2.982	18.655	1.00	18.13	C
ATOM	1156	CD1	ILE	A	149	14.511	-3.149	17.895	1.00	18.17	C
ATOM	1157	CG2	ILE	A	149	15.700	-5.096	20.019	1.00	19.10	C
ATOM	1158	C	ILE	A	149	18.834	-3.408	18.492	1.00	19.23	C
ATOM	1159	O	ILE	A	149	18.845	-3.791	17.323	1.00	20.25	O
ATOM	1160	N	GLY	A	150	19.558	-2.381	18.927	1.00	18.35	N
ATOM	1161	CA	GLY	A	150	20.482	-1.680	18.039	1.00	17.82	C
ATOM	1162	C	GLY	A	150	19.745	-0.943	16.930	1.00	18.01	C
ATOM	1163	O	GLY	A	150	20.273	-0.782	15.825	1.00	18.92	O

ATOM	1164	N	ALA	A	151	18.530	-0.494	17.230	1.00	16.81	N
ATOM	1165	CA	ALA	A	151	17.785	0.386	16.309	1.00	16.58	C
ATOM	1166	CB	ALA	A	151	16.365	0.610	16.814	1.00	15.51	C
ATOM	1167	C	ALA	A	151	18.502	1.727	16.150	1.00	17.00	C
ATOM	1168	O	ALA	A	151	19.131	2.215	17.105	1.00	18.28	O
ATOM	1169	N	VAL	A	152	18.430	2.329	14.964	1.00	16.92	N
ATOM	1170	CA	VAL	A	152	19.067	3.619	14.731	1.00	17.25	C
ATOM	1171	CB	VAL	A	152	19.465	3.825	13.238	1.00	17.70	C
ATOM	1172	CG1	VAL	A	152	20.419	2.722	12.762	1.00	20.29	C
ATOM	1173	CG2	VAL	A	152	18.251	3.882	12.357	1.00	17.61	C
ATOM	1174	C	VAL	A	152	18.223	4.782	15.266	1.00	17.04	C
ATOM	1175	O	VAL	A	152	18.718	5.881	15.513	1.00	17.43	O
ATOM	1176	N	LYS	A	153	16.919	4.549	15.427	1.00	16.32	N
ATOM	1177	CA	LYS	A	153	16.014	5.561	15.943	1.00	15.90	C
ATOM	1178	CB	LYS	A	153	15.424	6.391	14.792	1.00	17.62	C
ATOM	1179	CG	LYS	A	153	16.374	7.355	14.069	1.00	21.36	C
ATOM	1180	CD	LYS	A	153	16.886	8.472	14.985	1.00	26.36	C
ATOM	1181	CE	LYS	A	153	15.779	9.375	15.496	1.00	29.00	C
ATOM	1182	NZ	LYS	A	153	16.295	10.500	16.351	1.00	31.93	N
ATOM	1183	C	LYS	A	153	14.867	4.871	16.657	1.00	14.89	C
ATOM	1184	O	LYS	A	153	14.495	3.745	16.280	1.00	14.72	O
ATOM	1185	N	TYR	A	154	14.316	5.562	17.655	1.00	14.38	N
ATOM	1186	CA	TYR	A	154	13.031	5.224	18.255	1.00	13.82	C
ATOM	1187	CB	TYR	A	154	13.164	5.052	19.779	1.00	14.86	C
ATOM	1188	CG	TYR	A	154	11.808	4.874	20.433	1.00	14.08	C
ATOM	1189	CD1	TYR	A	154	11.018	3.754	20.142	1.00	15.57	C
ATOM	1190	CE1	TYR	A	154	9.763	3.594	20.695	1.00	15.86	C
ATOM	1191	CZ	TYR	A	154	9.267	4.569	21.547	1.00	15.10	C
ATOM	1192	OH	TYR	A	154	8.019	4.379	22.110	1.00	15.01	O
ATOM	1193	CE2	TYR	A	154	10.016	5.700	21.845	1.00	15.83	C
ATOM	1194	CD2	TYR	A	154	11.279	5.852	21.272	1.00	14.37	C
ATOM	1195	C	TYR	A	154	12.087	6.372	17.980	1.00	13.73	C
ATOM	1196	O	TYR	A	154	12.419	7.544	18.252	1.00	14.62	O
ATOM	1197	N	LEU	A	155	10.917	6.062	17.427	1.00	12.95	N
ATOM	1198	CA	LEU	A	155	9.902	7.055	17.068	1.00	13.09	C
ATOM	1199	CB	LEU	A	155	9.914	7.303	15.564	1.00	13.61	C
ATOM	1200	CG	LEU	A	155	11.217	7.832	14.961	1.00	15.78	C
ATOM	1201	CD1	LEU	A	155	11.156	7.731	13.448	1.00	16.92	C
ATOM	1202	CD2	LEU	A	155	11.464	9.258	15.437	1.00	16.38	C
ATOM	1203	C	LEU	A	155	8.512	6.596	17.458	1.00	13.31	C
ATOM	1204	O	LEU	A	155	8.231	5.400	17.429	1.00	13.15	O
ATOM	1205	N	GLU	A	156	7.647	7.540	17.819	1.00	12.73	N
ATOM	1206	CA	GLU	A	156	6.245	7.258	18.145	1.00	13.69	C
ATOM	1207	CB	GLU	A	156	5.935	7.716	19.575	1.00	13.99	C
ATOM	1208	CG	GLU	A	156	6.791	6.988	20.607	1.00	14.44	C
ATOM	1209	CD	GLU	A	156	6.538	7.410	22.041	1.00	15.10	C
ATOM	1210	OE1	GLU	A	156	6.088	8.555	22.252	1.00	17.29	O
ATOM	1211	OE2	GLU	A	156	6.845	6.584	22.916	1.00	16.33	O
ATOM	1212	C	GLU	A	156	5.315	7.948	17.162	1.00	13.28	C
ATOM	1213	O	GLU	A	156	5.608	9.056	16.667	1.00	14.15	O
ATOM	1214	N	CYS	A	157	4.173	7.332	16.869	1.00	12.48	N
ATOM	1215	CA	CYS	A	157	3.248	7.946	15.935	1.00	12.25	C
ATOM	1216	CB	CYS	A	157	3.685	7.653	14.491	1.00	12.28	C
ATOM	1217	SG	CYS	A	157	3.426	5.920	13.944	1.00	13.16	S
ATOM	1218	C	CYS	A	157	1.832	7.442	16.176	1.00	11.07	C
ATOM	1219	O	CYS	A	157	1.635	6.467	16.913	1.00	11.61	O
ATOM	1220	N	SER	A	158	0.883	8.110	15.530	1.00	11.86	N
ATOM	1221	CA	SER	A	158	-0.529	7.714	15.475	1.00	11.57	C
ATOM	1222	CB	SER	A	158	-1.400	8.639	16.344	1.00	11.46	C
ATOM	1223	OG	SER	A	158	-2.777	8.454	16.067	1.00	12.84	O
ATOM	1224	C	SER	A	158	-1.030	7.770	14.043	1.00	12.35	C
ATOM	1225	O	SER	A	158	-0.912	8.817	13.399	1.00	13.35	O
ATOM	1226	N	ALA	A	159	-1.565	6.671	13.515	1.00	12.23	N
ATOM	1227	CA	ALA	A	159	-2.230	6.704	12.210	1.00	12.91	C
ATOM	1228	CB	ALA	A	159	-2.548	5.283	11.741	1.00	12.63	C
ATOM	1229	C	ALA	A	159	-3.520	7.513	12.248	1.00	13.19	C

ATOM	1230	O	ALA	A	159	-3.945	8.058	11.223	1.00	13.32	O
ATOM	1231	N	LEU	A	160	-4.156	7.563	13.414	1.00	13.73	N
ATOM	1232	CA	LEU	A	160	-5.424	8.250	13.580	1.00	14.94	C
ATOM	1233	CB	LEU	A	160	-6.096	7.781	14.872	1.00	15.57	C
ATOM	1234	CG	LEU	A	160	-7.542	8.210	15.145	1.00	17.88	C
ATOM	1235	CD1	LEU	A	160	-8.273	7.116	15.948	1.00	19.48	C
ATOM	1236	CD2	LEU	A	160	-7.551	9.552	15.882	1.00	19.96	C
ATOM	1237	C	LEU	A	160	-5.253	9.781	13.525	1.00	14.74	C
ATOM	1238	O	LEU	A	160	-5.977	10.445	12.763	1.00	16.73	O
ATOM	1239	N	THR	A	161	-4.291	10.314	14.271	1.00	14.73	N
ATOM	1240	CA	THR	A	161	-4.076	11.779	14.312	1.00	15.61	C
ATOM	1241	CB	THR	A	161	-3.653	12.248	15.708	1.00	15.82	C
ATOM	1242	OG1	THR	A	161	-2.357	11.718	16.027	1.00	15.63	O
ATOM	1243	CG2	THR	A	161	-4.667	11.818	16.780	1.00	16.36	C
ATOM	1244	C	THR	A	161	-2.985	12.248	13.357	1.00	16.70	C
ATOM	1245	O	THR	A	161	-2.850	13.446	13.091	1.00	17.02	O
ATOM	1246	N	GLN	A	162	-2.203	11.288	12.859	1.00	15.83	N
ATOM	1247	CA	GLN	A	162	-0.997	11.511	12.035	1.00	16.95	C
ATOM	1248	CB	GLN	A	162	-1.294	12.308	10.764	1.00	17.58	C
ATOM	1249	CG	GLN	A	162	-2.244	11.634	9.840	1.00	20.45	C
ATOM	1250	CD	GLN	A	162	-2.105	12.156	8.438	1.00	24.07	C
ATOM	1251	OE1	GLN	A	162	-2.089	11.398	7.477	1.00	25.85	O
ATOM	1252	NE2	GLN	A	162	-1.965	13.465	8.314	1.00	23.76	N
ATOM	1253	C	GLN	A	162	0.189	12.105	12.785	1.00	16.79	C
ATOM	1254	O	GLN	A	162	1.225	12.404	12.176	1.00	16.94	O
ATOM	1255	N	ARG	A	163	0.055	12.309	14.093	1.00	15.75	N
ATOM	1256	CA	ARG	A	163	1.174	12.792	14.891	1.00	15.88	C
ATOM	1257	CB	ARG	A	163	0.751	12.960	16.356	1.00	17.08	C
ATOM	1258	CG	ARG	A	163	1.879	13.133	17.357	1.00	20.93	C
ATOM	1259	CD	ARG	A	163	2.435	14.511	17.436	1.00	26.06	C
ATOM	1260	NE	ARG	A	163	1.382	15.517	17.404	0.75	27.25	N
ATOM	1261	CZ	ARG	A	163	1.616	16.822	17.415	1.00	28.54	C
ATOM	1262	NH1	ARG	A	163	2.865	17.266	17.462	1.00	29.24	N
ATOM	1263	NH2	ARG	A	163	0.602	17.670	17.354	1.00	27.89	N
ATOM	1264	C	ARG	A	163	2.378	11.859	14.749	1.00	15.55	C
ATOM	1265	O	ARG	A	163	2.258	10.640	14.947	0.50	12.27	O
ATOM	1266	N	GLY	A	164	3.525	12.425	14.373	1.00	14.69	N
ATOM	1267	CA	GLY	A	164	4.763	11.665	14.200	1.00	14.63	C
ATOM	1268	C	GLY	A	164	4.863	10.804	12.950	1.00	13.55	C
ATOM	1269	O	GLY	A	164	5.899	10.197	12.702	1.00	14.55	O
ATOM	1270	N	LEU	A	165	3.797	10.762	12.162	1.00	13.31	N
ATOM	1271	CA	LEU	A	165	3.707	9.782	11.084	1.00	13.13	C
ATOM	1272	CB	LEU	A	165	2.261	9.658	10.595	1.00	13.76	C
ATOM	1273	CG	LEU	A	165	1.980	8.553	9.573	1.00	14.86	C
ATOM	1274	CD1	LEU	A	165	2.309	7.157	10.106	1.00	16.12	C
ATOM	1275	CD2	LEU	A	165	0.535	8.633	9.149	1.00	14.31	C
ATOM	1276	C	LEU	A	165	4.664	10.070	9.935	1.00	13.35	C
ATOM	1277	O	LEU	A	165	5.419	9.185	9.497	1.00	13.32	O
ATOM	1278	N	LYS	A	166	4.666	11.299	9.425	1.00	13.52	N
ATOM	1279	CA	LYS	A	166	5.577	11.628	8.331	1.00	13.76	C
ATOM	1280	CB	LYS	A	166	5.261	13.010	7.784	1.00	14.69	C
ATOM	1281	CG	LYS	A	166	3.905	13.080	7.142	1.00	15.92	C
ATOM	1282	CD	LYS	A	166	3.602	14.524	6.741	1.00	19.54	C
ATOM	1283	CE	LYS	A	166	2.328	14.673	5.953	1.00	23.03	C
ATOM	1284	NZ	LYS	A	166	1.129	14.415	6.801	1.00	27.33	N
ATOM	1285	C	LYS	A	166	7.035	11.520	8.779	1.00	13.49	C
ATOM	1286	O	LYS	A	166	7.892	11.110	7.995	1.00	14.34	O
ATOM	1287	N	THR	A	167	7.327	11.841	10.038	1.00	13.55	N
ATOM	1288	CA	THR	A	167	8.671	11.722	10.581	1.00	13.77	C
ATOM	1289	CB	THR	A	167	8.710	12.319	11.997	1.00	14.09	C
ATOM	1290	OG1	THR	A	167	8.464	13.729	11.886	1.00	18.21	O
ATOM	1291	CG2	THR	A	167	10.041	12.057	12.697	1.00	15.51	C
ATOM	1292	C	THR	A	167	9.211	10.297	10.528	1.00	13.82	C
ATOM	1293	O	THR	A	167	10.397	10.080	10.290	1.00	13.73	O
ATOM	1294	N	VAL	A	168	8.341	9.311	10.743	1.00	12.93	N
ATOM	1295	CA	VAL	A	168	8.782	7.928	10.666	1.00	13.09	C

ATOM	1296	CB	VAL	A	168	7.585	6.971	10.881	1.00	12.82	C
ATOM	1297	CG1	VAL	A	168	7.969	5.525	10.510	1.00	13.06	C
ATOM	1298	CG2	VAL	A	168	7.118	7.023	12.336	1.00	13.99	C
ATOM	1299	C	VAL	A	168	9.458	7.652	9.316	1.00	12.52	C
ATOM	1300	O	VAL	A	168	10.544	7.070	9.260	1.00	12.69	O
ATOM	1301	N	PHE	A	169	8.807	8.071	8.244	1.00	12.32	N
ATOM	1302	CA	PHE	A	169	9.301	7.796	6.904	1.00	12.53	C
ATOM	1303	CB	PHE	A	169	8.129	7.771	5.926	1.00	12.31	C
ATOM	1304	CG	PHE	A	169	7.176	6.663	6.252	1.00	12.91	C
ATOM	1305	CD1	PHE	A	169	7.521	5.355	5.938	1.00	12.66	C
ATOM	1306	CE1	PHE	A	169	6.701	4.291	6.307	1.00	14.90	C
ATOM	1307	CZ	PHE	A	169	5.530	4.540	6.973	1.00	14.20	C
ATOM	1308	CE2	PHE	A	169	5.187	5.847	7.321	1.00	12.87	C
ATOM	1309	CD2	PHE	A	169	6.002	6.901	6.949	1.00	13.39	C
ATOM	1310	C	PHE	A	169	10.469	8.680	6.503	1.00	13.15	C
ATOM	1311	O	PHE	A	169	11.369	8.225	5.805	1.00	13.31	O
ATOM	1312	N	ASP	A	170	10.488	9.913	7.016	1.00	13.64	N
ATOM	1313	CA	ASP	A	170	11.672	10.784	6.859	1.00	14.87	C
ATOM	1314	CB	ASP	A	170	11.524	12.080	7.651	1.00	16.28	C
ATOM	1315	CG	ASP	A	170	10.543	13.048	7.058	1.00	19.88	C
ATOM	1316	OD1	ASP	A	170	10.119	12.892	5.896	1.00	21.00	O
ATOM	1317	OD2	ASP	A	170	10.202	14.002	7.798	1.00	23.66	O
ATOM	1318	C	ASP	A	170	12.890	10.077	7.440	1.00	14.49	C
ATOM	1319	O	ASP	A	170	13.957	10.021	6.826	1.00	14.33	O
ATOM	1320	N	GLU	A	171	12.728	9.529	8.642	1.00	13.56	N
ATOM	1321	CA	GLU	A	171	13.831	8.893	9.342	1.00	13.86	C
ATOM	1322	CB	GLU	A	171	13.515	8.776	10.840	1.00	14.47	C
ATOM	1323	CG	GLU	A	171	13.513	10.133	11.541	1.00	16.70	C
ATOM	1324	CD	GLU	A	171	14.864	10.844	11.544	1.00	22.34	C
ATOM	1325	OE1	GLU	A	171	15.913	10.175	11.616	1.00	25.74	O
ATOM	1326	OE2	GLU	A	171	14.863	12.085	11.477	1.00	26.06	O
ATOM	1327	C	GLU	A	171	14.221	7.564	8.716	1.00	13.90	C
ATOM	1328	O	GLU	A	171	15.403	7.200	8.709	1.00	14.62	O
ATOM	1329	N	ALA	A	172	13.246	6.838	8.180	1.00	13.38	N
ATOM	1330	CA	ALA	A	172	13.546	5.622	7.432	1.00	13.23	C
ATOM	1331	CB	ALA	A	172	12.252	5.014	6.897	1.00	14.03	C
ATOM	1332	C	ALA	A	172	14.495	5.956	6.286	1.00	13.28	C
ATOM	1333	O	ALA	A	172	15.493	5.250	6.050	1.00	14.11	O
ATOM	1334	N	ILE	A	173	14.178	7.033	5.573	1.00	12.65	N
ATOM	1335	CA	ILE	A	173	14.992	7.458	4.434	1.00	13.19	C
ATOM	1336	CB	ILE	A	173	14.250	8.516	3.612	1.00	12.40	C
ATOM	1337	CG1	ILE	A	173	13.103	7.828	2.857	1.00	14.78	C
ATOM	1338	CD1	ILE	A	173	12.081	8.785	2.302	1.00	13.37	C
ATOM	1339	CG2	ILE	A	173	15.229	9.228	2.643	1.00	13.54	C
ATOM	1340	C	ILE	A	173	16.370	7.940	4.916	1.00	13.03	C
ATOM	1341	O	ILE	A	173	17.402	7.545	4.344	1.00	13.20	O
ATOM	1342	N	ARG	A	174	16.382	8.744	5.977	1.00	13.62	N
ATOM	1343	CA	ARG	A	174	17.674	9.196	6.521	1.00	14.74	C
ATOM	1344	CB	ARG	A	174	17.494	10.195	7.646	1.00	15.41	C
ATOM	1345	CG	ARG	A	174	16.882	11.488	7.168	1.00	15.94	C
ATOM	1346	CD	ARG	A	174	16.629	12.458	8.320	1.00	20.70	C
ATOM	1347	NE	ARG	A	174	16.118	13.738	7.823	1.00	22.87	N
ATOM	1348	CZ	ARG	A	174	15.001	14.333	8.237	1.00	26.43	C
ATOM	1349	NH1	ARG	A	174	14.252	13.806	9.205	1.00	29.29	N
ATOM	1350	NH2	ARG	A	174	14.640	15.486	7.689	1.00	29.09	N
ATOM	1351	C	ARG	A	174	18.557	8.055	6.972	1.00	15.31	C
ATOM	1352	O	ARG	A	174	19.786	8.137	6.874	1.00	16.11	O
ATOM	1353	N	ALA	A	175	17.955	6.974	7.463	1.00	15.52	N
ATOM	1354	CA	ALA	A	175	18.750	5.836	7.925	1.00	16.05	C
ATOM	1355	CB	ALA	A	175	17.848	4.791	8.567	1.00	16.64	C
ATOM	1356	C	ALA	A	175	19.527	5.214	6.784	1.00	16.94	C
ATOM	1357	O	ALA	A	175	20.554	4.546	6.996	1.00	17.87	O
ATOM	1358	N	VAL	A	176	19.023	5.398	5.570	1.00	16.33	N
ATOM	1359	CA	VAL	A	176	19.637	4.805	4.402	1.00	16.94	C
ATOM	1360	CB	VAL	A	176	18.558	4.324	3.380	1.00	16.82	C
ATOM	1361	CG1	VAL	A	176	19.199	3.795	2.097	1.00	17.60	C

ATOM	1362	CG2	VAL	A	176	17.671	3.260	4.012	1.00	17.37	C
ATOM	1363	C	VAL	A	176	20.608	5.811	3.770	1.00	18.02	C
ATOM	1364	O	VAL	A	176	21.728	5.427	3.375	1.00	19.01	O
ATOM	1365	N	LEU	A	177	20.209	7.085	3.716	1.00	18.56	N
ATOM	1366	CA	LEU	A	177	20.959	8.112	2.960	1.00	19.99	C
ATOM	1367	CB	LEU	A	177	19.995	9.076	2.262	1.00	19.32	C
ATOM	1368	CG	LEU	A	177	19.084	8.466	1.202	1.00	19.69	C
ATOM	1369	CD1	LEU	A	177	18.245	9.570	0.570	1.00	21.53	C
ATOM	1370	CD2	LEU	A	177	19.869	7.693	0.137	1.00	20.62	C
ATOM	1371	C	LEU	A	177	21.971	8.922	3.771	1.00	21.89	C
ATOM	1372	O	LEU	A	177	22.858	9.576	3.198	1.00	22.03	O
ATOM	1373	N	CYS	A	178	21.826	8.919	5.089	1.00	22.74	N
ATOM	1374	CA	CYS	A	178	22.743	9.643	5.961	1.00	25.08	C
ATOM	1375	CB	CYS	A	178	21.984	10.500	6.977	1.00	25.34	C
ATOM	1376	SG	CYS	A	178	20.806	11.678	6.228	0.75	28.16	S
ATOM	1377	C	CYS	A	178	23.648	8.648	6.661	1.00	26.11	C
ATOM	1378	O	CYS	A	178	23.186	7.875	7.499	1.00	27.77	O
ATOM	1379	O6	GDP	A	191	-3.009	5.022	15.369	1.00	12.24	O
ATOM	1380	C6	GDP	A	191	-4.015	4.359	15.678	1.00	12.17	C
ATOM	1381	N1	GDP	A	191	-4.860	4.763	16.631	1.00	11.86	N
ATOM	1382	C2	GDP	A	191	-5.946	4.059	17.010	1.00	13.33	C
ATOM	1383	N3	GDP	A	191	-6.335	2.908	16.446	1.00	12.94	N
ATOM	1384	C4	GDP	A	191	-5.563	2.390	15.462	1.00	12.01	C
ATOM	1385	N2	GDP	A	191	-6.728	4.592	17.968	1.00	14.08	N
ATOM	1386	C5	GDP	A	191	-4.355	3.101	15.012	1.00	11.72	C
ATOM	1387	N7	GDP	A	191	-3.778	2.354	14.043	1.00	11.60	N
ATOM	1388	C8	GDP	A	191	-4.554	1.260	13.880	1.00	11.74	C
ATOM	1389	N9	GDP	A	191	-5.622	1.293	14.713	1.00	11.13	N
ATOM	1390	C1*	GDP	A	191	-6.641	0.245	14.813	1.00	12.04	C
ATOM	1391	C2*	GDP	A	191	-7.274	-0.146	13.495	1.00	13.49	C
ATOM	1392	O2*	GDP	A	191	-8.363	0.720	13.221	1.00	15.11	O
ATOM	1393	C3*	GDP	A	191	-7.655	-1.573	13.776	1.00	13.18	C
ATOM	1394	O3*	GDP	A	191	-8.833	-1.618	14.593	1.00	14.69	O
ATOM	1395	O4*	GDP	A	191	-5.970	-0.930	15.291	1.00	11.20	O
ATOM	1396	C4*	GDP	A	191	-6.533	-2.077	14.643	1.00	12.56	C
ATOM	1397	C5*	GDP	A	191	-5.462	-2.819	13.842	1.00	10.91	C
ATOM	1398	O5*	GDP	A	191	-4.919	-1.976	12.814	1.00	9.53	O
ATOM	1399	PA	GDP	A	191	-4.141	-2.627	11.586	1.00	9.51	P
ATOM	1400	O1A	GDP	A	191	-3.619	-1.488	10.785	1.00	9.65	O
ATOM	1401	O2A	GDP	A	191	-4.994	-3.603	10.878	1.00	10.36	O
ATOM	1402	O3A	GDP	A	191	-2.961	-3.377	12.352	1.00	9.19	O
ATOM	1403	PB	GDP	A	191	-1.946	-4.469	11.696	1.00	8.75	P
ATOM	1404	O1B	GDP	A	191	-0.668	-4.238	12.435	1.00	8.01	O
ATOM	1405	O3B	GDP	A	191	-2.522	-5.850	12.023	1.00	9.28	O
ATOM	1406	O2B	GDP	A	191	-1.858	-4.210	10.209	1.00	9.68	O
ATOM	1407	MG	MG	A	192	-2.167	-5.353	8.526	1.00	8.83	MG
ATOM	1408	CBD	DRG	A	193	0.406	-14.451	2.136	1.00	20.00	C
ATOM	1409	CBB	DRG	A	193	2.146	-12.890	2.465	1.00	20.00	C
ATOM	1410	NBA	DRG	A	193	3.529	-12.690	2.960	1.00	20.00	N
ATOM	1411	CBC	DRG	A	193	3.640	-12.452	4.424	1.00	20.00	C
ATOM	1412	CBE	DRG	A	193	2.434	-11.812	5.143	1.00	20.00	C
ATOM	1413	CAZ	DRG	A	193	4.116	-11.597	2.155	1.00	20.00	C
ATOM	1414	CAY	DRG	A	193	3.136	-10.466	1.830	1.00	20.00	C
ATOM	1415	CAW	DRG	A	193	3.849	-9.122	1.968	1.00	20.00	C
ATOM	1416	CAV	DRG	A	193	4.966	-8.927	0.931	1.00	20.00	C
ATOM	1417	CAX	DRG	A	193	4.587	-7.782	-0.014	1.00	20.00	C
ATOM	1418	NAT	DRG	A	193	5.170	-10.167	0.151	1.00	20.00	N
ATOM	1419	C2	DRG	A	193	5.755	-10.178	-1.062	1.00	20.00	C
ATOM	1420	N1	DRG	A	193	6.881	-10.908	-1.294	1.00	20.00	N
ATOM	1421	N3	DRG	A	193	5.235	-9.449	-2.081	1.00	20.00	N
ATOM	1422	C4	DRG	A	193	5.792	-9.433	-3.311	1.00	20.00	C
ATOM	1423	CAU	DRG	A	193	5.162	-8.597	-4.433	1.00	20.00	C
ATOM	1424	C5	DRG	A	193	7.151	-10.382	-3.474	1.00	20.00	C
ATOM	1425	C6	DRG	A	193	7.501	-10.933	-2.508	1.00	20.00	C
ATOM	1426	NAM	DRG	A	193	8.587	-11.713	-2.703	1.00	20.00	N
ATOM	1427	CAJ	DRG	A	193	9.598	-11.432	-3.564	1.00	20.00	C

ATOM	1428	CAF	DRG	A	193	10.250	-12.483	-4.242	1.00	20.00	C
ATOM	1429	CAB	DRG	A	193	11.209	-12.178	-5.237	1.00	20.00	C
ATOM	1430	CAA	DRG	A	193	11.876	-13.169	-5.994	1.00	20.00	C
ATOM	1431	NAD	DRG	A	193	11.681	-14.479	-5.807	1.00	20.00	N
ATOM	1432	CAC	DRG	A	193	12.793	-12.809	-6.990	1.00	20.00	C
ATOM	1433	CAG	DRG	A	193	13.025	-11.443	-7.223	1.00	20.00	C
ATOM	1434	CAK	DRG	A	193	14.012	-11.010	-8.312	1.00	20.00	C
ATOM	1435	NAH	DRG	A	193	12.390	-10.511	-6.483	1.00	20.00	N
ATOM	1436	CAE	DRG	A	193	11.498	-10.812	-5.516	1.00	20.00	C
ATOM	1437	CAI	DRG	A	193	10.871	-9.762	-4.834	1.00	20.00	C
ATOM	1438	CAL	DRG	A	193	9.914	-10.089	-3.863	1.00	20.00	C
ATOM	1439	O	HOH	Z	1	-0.134	-5.584	8.242	1.00	8.96	O
ATOM	1440	O	HOH	Z	2	-4.232	-5.015	8.711	1.00	9.58	O
ATOM	1441	O	HOH	Z	3	-2.275	-7.057	9.673	1.00	10.94	O
ATOM	1442	O	HOH	Z	4	10.477	-1.304	-3.031	1.00	26.08	O
ATOM	1443	O	HOH	Z	5	16.317	-2.369	-2.524	1.00	29.42	O
ATOM	1444	O	HOH	Z	6	5.047	-8.484	10.591	1.00	12.91	O
ATOM	1445	O	HOH	Z	7	5.124	-6.010	17.744	1.00	11.06	O
ATOM	1446	O	HOH	Z	8	0.567	-9.158	20.354	1.00	21.09	O
ATOM	1447	O	HOH	Z	9	0.703	-11.544	16.131	1.00	12.46	O
ATOM	1448	O	HOH	Z	10	3.703	-12.071	16.597	1.00	22.55	O
ATOM	1449	O	HOH	Z	11	-0.182	-8.384	10.983	1.00	11.20	O
ATOM	1450	O	HOH	Z	12	-4.333	-8.769	10.467	1.00	14.76	O
ATOM	1451	O	HOH	Z	13	-4.936	-6.912	12.473	1.00	14.22	O
ATOM	1452	O	HOH	Z	14	-8.445	11.206	35.892	1.00	27.24	O
ATOM	1453	O	HOH	Z	15	-3.210	-2.135	20.575	1.00	12.27	O
ATOM	1454	O	HOH	Z	16	-1.032	2.706	13.325	1.00	11.23	O
ATOM	1455	O	HOH	Z	17	0.215	4.566	14.883	1.00	12.32	O
ATOM	1456	O	HOH	Z	18	1.314	11.622	4.713	1.00	23.82	O
ATOM	1457	O	HOH	Z	19	0.615	11.501	7.265	1.00	29.69	O
ATOM	1458	O	HOH	Z	20	-14.149	1.973	10.917	1.00	22.67	O
ATOM	1459	O	HOH	Z	21	-11.299	-4.865	9.719	1.00	22.28	O
ATOM	1460	O	HOH	Z	22	-9.705	-6.203	8.035	1.00	17.69	O
ATOM	1461	O	HOH	Z	23	7.468	13.145	-9.245	1.00	28.76	O
ATOM	1462	O	HOH	Z	24	-6.991	-5.346	11.452	1.00	15.83	O
ATOM	1463	O	HOH	Z	25	4.514	11.481	-2.718	1.00	12.76	O
ATOM	1464	O	HOH	Z	26	6.818	14.498	0.538	1.00	22.98	O
ATOM	1465	O	HOH	Z	27	-6.619	-9.355	9.099	1.00	17.44	O
ATOM	1466	O	HOH	Z	28	15.357	6.087	-3.675	1.00	27.71	O
ATOM	1467	O	HOH	Z	29	15.754	8.383	-5.057	1.00	25.97	O
ATOM	1468	O	HOH	Z	30	-1.307	-5.728	19.093	1.00	27.63	O
ATOM	1469	O	HOH	Z	31	-0.946	-7.181	21.049	1.00	23.44	O
ATOM	1470	O	HOH	Z	32	-3.074	-4.762	20.234	1.00	31.01	O
ATOM	1471	O	HOH	Z	33	2.733	9.521	0.192	1.00	13.56	O
ATOM	1472	O	HOH	Z	34	3.360	9.455	-1.336	1.00	18.57	O
ATOM	1473	O	HOH	Z	36	11.041	-13.275	-2.389	1.00	29.42	O
ATOM	1474	O	HOH	Z	37	-6.223	11.555	10.296	1.00	32.37	O
ATOM	1475	O	HOH	Z	38	-9.612	-4.467	12.063	1.00	31.51	O
ATOM	1476	O	HOH	Z	44	4.098	-4.948	-0.736	1.00	26.48	O
ATOM	1477	O	HOH	Z	45	2.704	-3.500	0.992	1.00	25.13	O
ATOM	1478	O	HOH	Z	46	2.826	-1.554	1.772	1.00	32.13	O
ATOM	1479	O	HOH	Z	47	-5.682	11.211	-0.359	1.00	25.76	O
ATOM	1480	O	HOH	Z	48	-4.127	-5.680	-0.140	1.00	32.87	O
ATOM	1481	O	HOH	Z	49	0.270	-2.172	-5.395	1.00	28.65	O
ATOM	1482	O	HOH	Z	50	-2.450	-1.603	-5.615	1.00	31.17	O
ATOM	1483	O	HOH	Z	51	3.268	-18.749	6.483	1.00	21.70	O
ATOM	1484	O	HOH	Z	52	2.161	-21.956	8.428	1.00	31.44	O
ATOM	1485	O	HOH	Z	53	-1.571	-18.172	6.694	1.00	17.34	O
ATOM	1486	O	HOH	Z	54	-0.249	-18.135	5.433	1.00	29.65	O
ATOM	1487	O	HOH	Z	55	2.971	2.171	-6.016	1.00	23.51	O
ATOM	1488	O	HOH	Z	56	7.729	11.463	5.016	1.00	17.76	O
ATOM	1489	O	HOH	Z	57	12.155	18.376	1.959	1.00	26.19	O
ATOM	1490	O	HOH	Z	58	12.594	17.158	5.114	1.00	31.33	O
ATOM	1491	O	HOH	Z	59	-1.569	11.596	18.927	1.00	16.57	O
ATOM	1492	O	HOH	Z	60	7.579	9.928	14.909	1.00	15.76	O
ATOM	1493	O	HOH	Z	61	19.744	-0.867	13.120	1.00	18.17	O

ATOM	1494	O	HOH	Z	62	8.707	10.237	18.525	1.00	17.38	O
ATOM	1495	O	HOH	Z	63	21.457	20.231	-3.910	1.00	19.88	O
ATOM	1496	O	HOH	Z	64	11.940	-14.538	15.403	1.00	19.32	O
ATOM	1497	O	HOH	Z	65	-3.012	1.332	32.662	1.00	17.87	O
ATOM	1498	O	HOH	Z	66	2.933	13.528	10.444	1.00	18.09	O
ATOM	1499	O	HOH	Z	67	5.511	13.684	11.621	1.00	18.83	O
ATOM	1500	O	HOH	Z	68	-0.341	-6.697	26.164	1.00	22.76	O
ATOM	1501	O	HOH	Z	69	15.508	4.752	28.264	1.00	19.65	O
ATOM	1502	O	HOH	Z	70	9.568	-16.322	9.992	1.00	22.23	O
ATOM	1503	O	HOH	Z	71	15.958	7.636	18.742	1.00	23.87	O
ATOM	1504	O	HOH	Z	72	-6.783	0.200	18.310	1.00	21.13	O
ATOM	1505	O	HOH	Z	73	11.447	9.949	19.027	1.00	19.55	O
ATOM	1506	O	HOH	Z	74	23.144	10.431	0.689	1.00	23.34	O
ATOM	1507	O	HOH	Z	75	-7.996	8.406	20.020	1.00	25.41	O
ATOM	1508	O	HOH	Z	76	2.608	3.969	31.316	1.00	26.88	O
ATOM	1509	O	HOH	Z	77	2.425	-3.227	31.617	1.00	27.72	O
ATOM	1510	O	HOH	Z	78	17.254	-8.666	3.011	1.00	25.39	O
ATOM	1511	O	HOH	Z	79	12.918	-15.190	-0.608	1.00	34.94	O
ATOM	1512	O	HOH	Z	80	12.067	-17.714	-1.315	1.00	32.30	O
ATOM	1513	O	HOH	Z	81	-10.581	2.386	3.590	1.00	25.93	O
ATOM	1514	O	HOH	Z	82	9.735	-14.832	-4.003	1.00	32.53	O
ATOM	1515	O	HOH	Z	83	11.835	-6.311	-3.598	1.00	32.48	O
ATOM	1516	O	HOH	Z	84	8.689	11.909	16.333	1.00	23.24	O
ATOM	1517	O	HOH	Z	85	-6.601	-3.539	17.808	1.00	23.91	O
ATOM	1518	O	HOH	Z	86	17.879	3.999	27.428	1.00	28.73	O
ATOM	1519	O	HOH	Z	87	12.222	-16.863	9.403	1.00	23.77	O
ATOM	1520	O	HOH	Z	88	5.867	8.517	27.830	1.00	26.11	O
ATOM	1521	O	HOH	Z	89	17.094	7.930	10.841	1.00	29.44	O
ATOM	1522	O	HOH	Z	90	-8.791	-1.299	17.296	1.00	26.69	O
ATOM	1523	O	HOH	Z	91	4.804	0.800	30.972	1.00	28.06	O
ATOM	1524	O	HOH	Z	92	3.461	15.172	13.591	1.00	26.08	O
ATOM	1525	O	HOH	Z	93	-14.826	5.257	33.876	1.00	28.68	O
ATOM	1526	O	HOH	Z	94	10.358	-15.459	13.503	1.00	27.42	O
ATOM	1527	O	HOH	Z	95	21.417	4.120	9.561	1.00	25.75	O
ATOM	1528	O	HOH	Z	96	-1.574	-17.006	13.344	1.00	27.33	O
ATOM	1529	O	HOH	Z	97	3.802	-20.213	9.738	1.00	26.02	O
ATOM	1530	O	HOH	Z	98	4.687	-10.827	24.294	1.00	27.00	O
ATOM	1531	O	HOH	Z	99	18.579	-0.462	26.775	1.00	28.17	O
ATOM	1532	O	HOH	Z	100	9.226	4.998	32.035	1.00	26.92	O
ATOM	1533	O	HOH	Z	101	-0.655	6.897	29.980	1.00	28.47	O
ATOM	1534	O	HOH	Z	102	6.947	10.853	20.750	1.00	25.28	O
ATOM	1535	O	HOH	Z	103	18.739	-13.365	14.310	1.00	28.79	O
ATOM	1536	O	HOH	Z	104	1.875	6.190	29.907	1.00	28.27	O
ATOM	1537	O	HOH	Z	105	11.693	3.613	32.212	1.00	34.50	O
ATOM	1538	O	HOH	Z	106	-10.191	6.741	34.392	1.00	29.28	O
ATOM	1539	O	HOH	Z	107	-6.667	8.319	32.568	1.00	35.78	O
ATOM	1540	O	HOH	Z	108	22.107	-0.247	11.857	1.00	35.64	O
ATOM	1541	O	HOH	Z	109	5.419	11.752	18.009	1.00	33.54	O
ATOM	1542	O	HOH	Z	110	18.317	4.725	18.913	1.00	28.26	O
ATOM	1543	O	HOH	Z	111	-19.579	1.950	5.342	1.00	33.46	O
ATOM	1544	O	HOH	Z	112	-0.971	-9.842	17.992	1.00	26.85	O
ATOM	1545	O	HOH	Z	113	18.991	1.782	28.143	1.00	32.15	O
ATOM	1546	O	HOH	Z	114	-8.447	10.667	8.833	1.00	31.09	O
ATOM	1547	O	HOH	Z	115	-7.957	-5.784	38.341	1.00	33.78	O
ATOM	1548	O	HOH	Z	116	23.481	8.252	-0.900	1.00	31.21	O
ATOM	1549	O	HOH	Z	117	16.296	-11.718	2.686	1.00	25.55	O
ATOM	1550	O	HOH	Z	118	-8.089	6.124	32.836	1.00	33.27	O
ATOM	1551	O	HOH	Z	119	-18.899	1.773	9.908	1.00	28.82	O
ATOM	1552	O	HOH	Z	120	-4.891	11.134	22.878	1.00	34.04	O
ATOM	1553	O	HOH	Z	121	-3.879	-6.962	-3.653	1.00	25.64	O
ATOM	1554	O	HOH	Z	122	6.294	-5.334	27.716	1.00	32.59	O
ATOM	1555	O	HOH	Z	123	19.061	-12.417	17.639	1.00	35.78	O
ATOM	1556	O	HOH	Z	124	3.653	8.065	29.157	1.00	31.42	O
ATOM	1557	O	HOH	Z	125	20.626	-6.832	4.898	1.00	30.71	O
ATOM	1558	O	HOH	Z	126	-4.089	15.494	11.914	1.00	37.36	O
ATOM	1559	O	HOH	Z	127	-5.505	-1.483	22.141	1.00	33.40	O

ATOM	1560	O	HOH	Z	128	13.760	-7.726	25.645	1.00	43.49	O
ATOM	1561	O	HOH	Z	129	11.225	13.014	16.204	1.00	31.87	O
ATOM	1562	O	HOH	Z	130	-10.901	1.977	13.994	1.00	31.78	O
ATOM	1563	O	HOH	Z	131	-1.232	15.529	13.872	1.00	31.18	O
ATOM	1564	O	HOH	Z	132	-8.158	1.006	23.180	1.00	33.63	O
ATOM	1565	O	HOH	Z	133	3.948	-1.576	30.415	1.00	43.32	O
ATOM	1566	O	HOH	Z	134	-7.606	6.346	3.782	1.00	37.67	O
ATOM	1567	O	HOH	Z	135	20.292	-7.467	0.209	1.00	31.70	O
ATOM	1568	O	HOH	Z	136	20.357	7.828	26.267	1.00	35.06	O
ATOM	1569	O	HOH	Z	137	20.649	-0.024	1.228	1.00	31.80	O
ATOM	1570	O	HOH	Z	138	-0.771	10.384	23.630	1.00	32.99	O
ATOM	1571	O	HOH	Z	139	17.801	-7.406	25.546	1.00	33.06	O
ATOM	1572	O	HOH	Z	140	-4.102	-2.547	-0.245	1.00	32.75	O
ATOM	1573	O	HOH	Z	141	-9.581	-5.177	30.449	1.00	32.14	O
ATOM	1574	O	HOH	Z	142	5.765	-21.459	11.112	1.00	32.28	O
ATOM	1575	O	HOH	Z	143	-7.145	-10.843	13.347	1.00	38.51	O
END											

References

- Bastola,D.R., Pahwa,G.S., Lin,M.F., and Cheng,P.W. (2002). Downregulation of PTEN/MMAC/TEP1 expression in human prostate cancer cell line DU145 by growth stimuli. *Mol. Cell Biochem.* 236, 75-81.
- Clark,R.D., Strizhev,A., Leonard,J.M., Blake,J.F., and Matthew,J.B. (2002). Consensus scoring for ligand/protein interactions. *J. Mol. Graph. Model.* 20, 281-295.
- Del Pozo,M.A., Price,L.S., Alderson,N.B., Ren,X.D., and Schwartz,M.A. (2000). Adhesion to the extracellular matrix regulates the coupling of the small GTPase Rac to its effector PAK. *EMBO J.* 19, 2008-2014.
- Engers,R., Zwaka,T.P., Gohr,L., Weber,A., Gerharz, C.D., and Gabbert,H.E. (2000). Tiam1 mutations in human renal-cell carcinomas. *Int. J. Cancer* 88, 369-376.
- Etienne-Manneville,S. and Hall,A. (2002). Rho GTPases in cell biology. *Nature* 420, 629-635.
- Fritz,G., Just,I., and Kaina,B. (1999). Rho GTPases are over-expressed in human tumors. *Int. J. Cancer* 81, 682-687.
- Gao,Y., Xing,J., Streuli,M., Leto,T.L., and Zheng,Y. (2001). Trp(56) of rac1 specifies interaction with a subset of guanine nucleotide exchange factors. *J. Biol. Chem.* 276, 47530-47541.
- Gruneberg,S., Wendt,B., and Klebe,G. (2001). Subnanomolar Inhibitors from Computer Screening: A Model Study Using Human Carbonic Anhydrase II Agnew. *Chem. Int. Ed Engl.* 40, 389-393.
- Guo,F., Gao,Y., Wang,L., and Zheng,Y. (2003). p19ARF-p53 tumor suppressor pathway regulates cell motility by suppression of PI3 kinase and Rac1 GTPase activities. *J. Biol. Chem.* paper in press.
- Hawkins,P.T., Eguinoa,A., Qiu,R.G., Stokoe,D., Cooke,F.T., Walters,R., Wennsstrom,S., Claesson-Welsh,L., Evans,T., Symons,M., and Stephens,L. (1995). PDGF stimulates an increase in GTP-Rac via activation of phosphoinositide 3-kinase. *Curr. Biol.* 5, 393-403.
- Hurst,T. (1994). Flexible 3D searching: the directed tweak technique. *J. Chem. Inf. Comput. Sci.* 34, 190-196.
- Kaighn,M.E., Narayan,K.S., Ohnuki,Y., Lechner,J.F., and Jones,L.W. (1979). Establishment and characterization of a human prostatic carcinoma cell line (PC-3). *Invest Urol.* 17, 16-23.


- Kamai,T., Arai,K., Tsujii,T., Honda,M., and Yoshida,K. (2001). Overexpression of RhoA mRNA is associated with advanced stage in testicular germ cell tumour. *BJU. Int.* 87, 227-231.
- Karnoub,A.E., Worthylake,D.K., Rossman,K.L., Pruitt,W.M., Campbell,S.L., Sondek,J., and Der,C.J. (2001). Molecular basis for Rac1 recognition by guanine nucleotide exchange factors. *Nat. Struct. Biol.* 8, 1037-1041.
- Kato-Stankiewicz,J., Hakimi,I., Zhi,G., Zhang,J., Serebriiskii,I., Guo,L., Edamatsu,H., Koike,H., Menon,S., Eckl,R., Sakamuri,S., Lu,Y., Chen,Q., Agarwal,S., Baumbach,W.R., Golemis,E.A., Tamanoi,F. and Khazak,V. (2002). Inhibitors of Ras/Raf-1 interaction identified by two-hybrid screening revert Ras-dependent transformation phenotypes in human cancer cells. 99, 14398-13303.
- Khosravi-Far,R., Solski,P.A., Clark,G.J., Kinch,M.S., and Der,C.J. (1995). Activation of Rac1, RhoA, and mitogen-activated protein kinases is required for Ras transformation. *Mol. Cell Biol.* 15, 6443-6453.
- Lang,S.H., Hyde,C., Reid,I.N., Hitchcock,I.S., Hart,C.A., Bryden,A.A., Villette,J.M., Stower,M.J., and Maitland,N.J. (2002). Enhanced expression of vimentin in multiple prostate cell lines in poorly differentiated and metastatic prostate carcinoma. *Prostate* 52, 253-263.
- Liliental,J., Moon,S.Y., Lesche,R., Mamillapalli,R., Li,D., Zheng,Y., Sun,H., and Wu,H. (2000). Genetic deletion of the Pten tumor suppressor gene promotes cell motility by activation of Rac1 and Cdc42 GTPases. *Curr. Bio.* 10, 401-404.
- Mira,J.P., Bernard,V., Groffen,J., Sanders,L.C., and Knaus,U.G. (2000). Endogenous, Hyperactive Rac3 controls proliferation of breast cancer cells by a p21-activated kinase-dependent pathway. *Proc. Natl. Acad. Sci. U. S. A* 97, 185-189.
- Movilla,N., Doseil,M., Zheng,Y., and Bustelo,X.R. (2001). How Vav proteins discriminate the GTPases Rac1 and RhoA from Cdc42. *Oncogene* 20, 8057-8065.
- Murshudov, G. N., Vagin, A. A., and Dodson, E. J. (1997) *Acta Crystallogr. Biol. Crystallogr.* **D53**, 240-255
- Otwinowski, Z. and Minor, W. (1997) Processing of X-ray Diffraction Data Collected in Oscillation Mode *Methods in Enzymology* **276**, 307-326
- Otwinowski, Z., Borek, D., Majewski, W., and Minor, W. (2003) Multiparametric scaling of diffraction intensities *Acta Crystallogr.* **A59**, 228-234

- Perola,E., Xu,K., Kollmeyer,T.M., Kaufmann,S.H., Prendergast,F.G., and Pang,Y.P. (2000). Successful virtual screening of a chemical database for farnesyltransferase inhibitor leads. *J. Med. Chem.* 43, 401-408.
- Qiu,R.G., Abo,A., McCormick,F., and Symons,M. (1997). Cdc42 regulates anchorage-independent growth and is necessary for Ras transformation. *Mol. Cell Biol.* 17, 3449-3458.
- Rarey,M., Kramer,B., Lengauer,T., and Klebe,G. (1996). A fast flexible docking method using an incremental construction algorithm. *J. Mol. Biol.* 261, 470-489.
- Ridley,A.J., Paterson,H.F., Johnston,C.L., Diekmann,D., and Hall,A. (1992). The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell* 70, 401-410.
- Sahai,E. (2002). Rho-GTPases and cancer. *Nature Reviews Cancer* 2, 133-142.
- Schmidt,A. and Hall,A. (2002). Guanine nucleotide exchange factors for Rho GTPases: turning on the switch. *Genes Dev.* 16, 1587-1609.
- Schmitz,A.A., Govek,E.E., Bottner,B., and Van Aelst,L. (2000). Rho GTPases: signaling, migration, and invasion. *Exp. Cell Res.* 261, 1-12.
- Schnelzer,A., Prechtel,D., Knaus,U., Dehne,K., Gerhard,M., Graeff,H., Harbeck,N., Schmitt,M., and Lengyel,E. (2000). Rac1 in human breast cancer: overexpression, mutation analysis, and characterization of a new isoform, Rac1b. *Oncogene* 19, 3013-3020.
- Stepan,V.M., Tatewaki,M., Matsushima,M., Dickinson,D.J., del Valle,J., and Todisco,A. (1999). Gastrin induces c-fos gene transcription via multiple signaling pathways. *Am. J. Physiol* 276, G415-G424.
- Suwa,H., Ohshio, G., Imamura,T., Watanabe,G., Arii,S., Imamura,M., Narumiya,S., Hiai,H., and Fukumoto,M. (1998). Overexpression of the rhoC gene correlates with progression of ductal adenocarcinoma of the pancreas. *Br. J. Cancer* 77, 147-152.
- Symons,M. (2000). Adhesion signaling: PAK meets Rac on solid ground. *Curr. Biol.* 10, R535-R537.
- Vagin, A. A. & Teplyakov, A. (1997). *J. Appl. Cryst.* 30, 1022-1025.
- Van Aelst,L. and D'Souza-Schorey,C. (1997). Rho GTPases and signaling networks. *Genes Dev.* 11, 2295-2322.

- Waszkowycz,B. (2001). Large-scale virtual screening for discovering leads in the postgenomic era. *IBM systems J.* 40, 360-376.
- Winn, M. D., Isupov, M. N., and Murshudov, G. N. (2001) Use of TLS parameters to model anisotropic displacements in macromolecular refinement *Acta Crystallogr. Biol. Crystallogr.* **D57**, 122-133
- Worthlake,D.K., Rossman,K.L., and Sondek,J. (2000). Crystal structure of Rac1 in complex with the guanine nucleotide exchange region of Tiam1. *Nature* 408, 682-688.
- Zhen,Y. (2001). Dbl family guanine nucleotide exchange factors. *Trends Biochem. Sci.* 26, 724-732.
- Zheng,Y., Olson,M.F., Hall,A., Cerione,R.A., and Toksoz,D. (1995). Direct involvement of the small GTP-binding protein Rho in Ibc oncogene function. *J. Biol. Chem.* 270, 9031-9034.

WHAT IS CLAIMED IS:

1. A method for treating an indication mediated by mammalian Rho family proteins, comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (IIa):



(IIa) wherein:

R₁ to R₂ are independently selected from the group consisting of H, -X- Alk, -X-Alk-X', and -X-Y-X'; wherein

X is -CR₇R₈;

X' is -CHR₇R₈;

Alk is a C₂-C₁₈ substituted or unsubstituted hydrocarbon chain;

Y is a C₂-C₈ substituted or unsubstituted alkylene chain;

R₆ is H or (C1 -C4) alkyl; and


R₇ and R₈ are independently selected from the group consisting of H or (C1 -C4) alkyl;

or a salt of a compound of formula (IIa).

2. The method of Claim 1, wherein Alk is substituted with halo, halo (C1 - C4) alkoxy, (C3 -C8) cycloalkyl, hydroxy, or acetyl.

3. The method of Claim 1, wherein Y is substituted with an NR₆ group.

4. A method for treating an indication mediated by mammalian Rho family proteins, comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (III):




(III) wherein:

R₁₀ to R₁₂ are independently selected from the group consisting of H, halo, (C1 -C4) alkyl, branched (C3-C4) alkyl, halo (C1 -C4) alkyl, (C1 -C4) alkoxy, NO₂, and NH₂;

or a salt of a compound of formula (III).

5. The method of Claim 4, wherein R_{10} to R_{12} are independently selected from the group consisting of H, (C1 -C4) alkyl, or branched (C3-C4) alkyl.

6. A method for treating an indication mediated by mammalian Rho family proteins comprising administering to a subject in need of such treatment a safe and effective amount of at least one compound having the formula (IV):



(IV)

or pharmaceutically acceptable salts thereof.

7. The method according to Claim 1, wherein the compound is N6-(2-((4-(diethylamino) 1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine.

8. The method according to Claim 1, wherein the mammalian Rho family GTPase is selected from the group consisting of RhoA, RhoB, RhoC, RhoD, RhoE/Rnd3, Rnd1/Rho6, Rnd2/Rho7, RhoG, Rac1, Rac2, Rac3, Cdc42, TC10, TTF/RhoH, RhoL, Chp, WRCH1, TCL, RIF, and combinations thereof.

9. The method according to Claim 8, wherein the Rho GTPase is selected from the group comprising Rac1, Rac2, Rac3, and combinations thereof.

10. The method according to Claim 9, wherein the Rho GTPase is Rac1.

11. The method according to Claim 1, wherein the compound interacts with the Rho regulatory pathway via interaction with a Rac GTPase.

12. The method according to Claim 11, wherein the Rac GTPase is selected from the group consisting of Rac1, Rac2, and Rac3.

13. The method according to Claim 11, wherein the Rac GTPase is Rac1.

14. The method according to Claim 1, wherein the indication is selected from the group consisting of leukemia, prostate cancer, ovarian cancer, pancreas cancer, lung cancer, breast cancer, liver cancer, head and neck cancer, colon cancer, bladder cancer, non-Hodgkin's lymphoma cancer and melanoma.

15. The method according to Claim 1, wherein the indication is abnormal cell proliferation.

16. The method according to Claim 1, wherein the indication is cancer cell proliferation.

17. The method according to Claim 1, wherein the indication is selected from the group consisting of hypertension, atherosclerosis, restenosis, cerebral ischemia, cerebral vasospasm, neuronal degeneration, spinal cord injury, cancer of the breast, colon, prostate, ovaries, brain or lung, thrombotic disorders, asthma, glaucoma, osteoporosis and erectile dysfunction.

18. A pharmaceutical composition comprising N6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine and a pharmaceutically-acceptable carrier.


19. The pharmaceutical composition according to Claim 18, wherein N6-(2-((4-(diethylamino)-1-methylbutyl)amino)-6-methyl-4-pyrimidinyl)-2-methyl-4,6-quinolinediamine compound is at least 1% by weight.

20. The pharmaceutical composition according to Claim 18 further comprising an additional pharmaceutically active agent.

21. The pharmaceutical composition according to Claim 20, wherein the second pharmaceutically active agent is selected from the group consisting of farnesyl protein transferase inhibitors, prenyl-protein transferase inhibitors, geranylgeranyl-protein transferase inhibitors, toxins and combinations thereof.

22. A crystal of Rac-1 GTPase having a space group $P2_1$ and unit cell dimensions of about $a=41.8$ Angstroms, $b=40.0$ Angstroms, $c=52.1$ Angstroms, and $\beta=105.93^\circ$.

23. The crystal of Claim 22, wherein Rac-1 GTPase is complexed with a compound having the formula (IIa):



(IIa) wherein:

R_1 to R_2 are independently selected from the group consisting of H, -X-Alk, -X-Alk-X', and -X-Y-X'; wherein

X is $-CR_7R_8$;

X' is $-CHR_7R_8$;

Alk is a C₂–C₁₈ substituted or unsubstituted hydrocarbon chain;

Y is a C₂–C₈ substituted or unsubstituted alkylene chain;


R₆ is H or (C1 -C4) alkyl; and

R₇ and R₈ are independently selected from the group consisting of H or (C1 -C4) alkyl;

or a salt of a compound of formula (IIa).

24. A crystal Rac-1 GTPase, wherein said crystal has a three-dimensional structure characterized by the atomic structure coordinates of Table 1.

25. The crystal of Claim 24, wherein Rac-1 GTPase is complexed with a compound having the formula (IIa):



(IIa) wherein:

R₁ to R₂ are independently selected from the group consisting of H, -X- Alk, -X-Alk-X', and -X-Y-X'; wherein

X is -CR₇R₈;

X' is -CHR₇R₈;

Alk is a C₂–C₁₈ substituted or unsubstituted hydrocarbon chain;

Y is a C₂–C₈ substituted or unsubstituted alkylene chain;

R₆ is H or (C1 -C4) alkyl; and

R₇ and R₈ are independently selected from the group consisting of H or (C1 -C4) alkyl;

or a salt of a compound of formula (IIa).

26. A method of making a crystal comprising Rac-1 GTPase, wherein the crystal Rac-1 GTPase comprises amino acid residues 1 through 185 of the sequence of SEQ ID NO:1; wherein the crystal effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of greater than 5.0 Angstroms; and wherein said crystal has a space group P2₁ and unit cell dimensions of about a=41.8 Angstroms, b=40.0 Angstroms, c=52.1 Angstroms, and β=105.93°.

said method comprising growing a crystal by hanging drop method using a buffer that comprises 20% PEG 8000, 5% Ethylene Glycol, 15% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, and 1 mM NSC23766 at 20°C; and

equilibrating the crystal in a drop comprising 18% PEG 8000, 3.5% Ethylene Glycol, 10.7% DMSO, 0.1 M HEPES, pH 7.0, 10 mM DTT, 14.3 mM NSC23766 over a reservoir containing a buffer comprising 25% PEG 8000, 5% Ethylene Glycol, 15% DMSO, 0.1M HEPES, pH 7.0, and 10 mM DTT.

27. A method of identifying an agent that interacts with Rac-1 GTPase comprising:

- (a) obtaining a crystal of a complex comprising Rac-1 GTPase;
- (b) obtaining the atomic coordinates of the crystal; and
- (c) using the atomic coordinates and one or more molecular modeling techniques to identify an agent that interacts with Rac-1 GTPase.

28. A method for evaluating the potential of a chemical entity to associate with:

a) a molecule or molecular complex comprising a binding pocket defined by structure coordinates of Rac-1 GTPase Trp-56, Lys-5, Val-7, Ser-71, Ile-36, Tyr-64, Arg-68, and Pro-73; or

b) a homologue of said molecule or molecular complex, wherein said homologue comprises a binding pocket that has a root mean square deviation from the backbone atoms of said amino acids of not more than about 1.5 Angstrom comprising the steps of:

- (i) employing computational means to perform a fitting operation between the chemical entity and the binding pocket of the molecule or molecular complex; and
- (ii) analyzing the results of said fitting operation to quantify the association between the chemical entity and the binding pocket.

29. A compound for inhibiting activity of Rac-1 GTPase, whose active site presents a three-dimensional structure from a crystal of Rac-1 GTPase having a space group $P2_1$ and unit cell dimensions of about $a=41.8$ Angstroms, $b=40.0$ Angstroms, $c=52.1$ Angstroms, and $\beta=105.93^\circ$.

30. A crystal of Rac-1 GTPase bound to GDP.

31. A crystal Rac-1 GTPase, wherein said crystal has a three-dimensional structure characterized as having atomic coordinates with a root-mean-squared deviation of 6Å or less, 5Å or less, 4Å or less, 3Å or less, 2Å or less, 1Å or less, relative to atoms of amino acids 30-39 and 59-73 of Table 1 or Table 2.

32. The crystal of Claim 31, wherein the of amino acids 30-39 and 59-73 of Table 1 or Table 2 are peptide backbone atoms.

33. Atomic coordinates with a root-mean-squared deviation of 6Å or less, 5Å or less, 4Å or less, 3Å or less, 2Å or less, 1Å or less, relative to atoms of amino acids 30-39 and 59-73 of Table 1 or Table 2.

34. The coordinates of Claim 33, wherein the of amino acids 30-39 and 59-73 of Table 1 or Table 2 are peptide backbone atoms.

35. A method of characterizing the three-dimensional structure of Rac-1 GTPase comprising:

(a) obtaining a crystal of a GDP-bound Rac-1 GTPase; and

(b) obtaining the atomic coordinates of the crystallized GDP-bound Rac-1 GTPase.

36. A method of characterizing the three-dimensional structure of a ligand bound to a Rac-1 GTPase comprising:

(a) obtaining a crystal of a ligand bound to a GDP-bound Rac-1 GTPase; and

(b) obtaining the atomic coordinates of the crystallized ligand and GDP-bound Rac-1 GTPase.

37. A method of identifying an agent that interacts with Rac-1 GTPase comprising:

(a) obtaining a crystal of a GDP-bound Rac-1 GTPase; and

(b) obtaining the atomic coordinates of the crystallized GDP-bound Rac-1 GTPase; and

(c) using the atomic coordinates and one or more molecular modeling techniques to identify an agent that interacts with GDP-bound Rac-1 GTPase.

38. A method of designing a Rac-1 GTPase inhibitor comprising accessing at least a portion the data of Table 1 or Table 2 and modeling the binding of one or more compounds to a Rac-1 GTPase utilizing said data.

39. The method of Claim 1, wherein said modeling comprises predicting the likelihood of a compound to bind to Rac-1 GTPase by modeling the interaction of a compound with a Rac-1 GTPase activation site.

40. The method of Claim 1, further comprising utilizing said data to identify one or more interactions between a Rac-1 GTPase and a ligand.

41. The method of Claim 40, wherein said one or more interactions are selected from the group consisting of one or more hydrogen bonds between ligand and protein; one or more electrostatic interactions between ligand and protein; one or more hydrophobic interactions; one or more covalent bonds between ligand and protein; one or more changes in the positions of protein atoms, such as amino acid side chains upon ligand binding; one or more changes in atoms whose positions are well defined to a conformation that may maintain flexibility; one or more hydrogen bonding networks; one or more structural changes to the hydrogen bond network, one or more structural changes to the interactions with Mg^{2+} , one or more structural changes to the interactions with GDP, and one or more structural changes to the interactions with GTP.

42. A method identifying a Rac-1 GTPase inhibitor comprising obtaining data comprising the one or more interactions of Claim 40 and modeling a compound to interact with a Rac-1 GTPase, wherein the compound comprises one or more of the interactions of Claim 40 or improvements of one or more of the interactions of Claim 40.

43. A method of designing a Rac-1 GTPase inhibitor comprising obtaining an X-ray crystal structure coordinate of a GDP-bound Rac-1 GTPase and identifying a compound that inhibits a Rac-1 GTPase activation.

44. The method of Claim 43, comprising identifying a compound that inhibits Thr35 binding of Mg^{2+} .

45. The method of Claim 43 or Claim 44, comprising identifying a compound that inhibits conformational change in the loop containing amino acid residues 60-64.

46. The method of Claim 43, comprising identifying a compound that inhibits Ala59 from displacing Mg^{2+} .

47. The method of Claim 46, wherein said compound hydrogen bonds to Ala59.

48. The method of Claim 43, comprising identifying a compound that inhibits a RAC-specific GEF from binding to RAC

49. The method of Claim 43, comprising identifying a compound that inhibits RAC-specific GEF from dissociating from RAC.

50. The method of Claim 43, comprising identifying a compound that inhibits conformational change in the amino acids of switch I.

51. The method of Claim 43, comprising identifying a compound that inhibits conformational change in the amino acids of switch II

52. The method of Claim 43, wherein said compound is identified as hydrogen bonding to Leu70 or Ser71, or both.

53. The method of Claim 43, wherein said compound is identified as hydrogen bonding to Leu67, Gln74, Asp57, or Ser71, or combinations thereof.

54. The method of Claim 43, wherein said compound is identified as being in van der Waals contact with Val36, Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 or Pro73, or combinations thereof.

55. A method of identifying a proteasome inhibitor comprising accessing one or more of the coordinates of Table 1 or Table 2 and identifying a compound that occupies a position within a range of about 1.0 to 2.0 angstroms or about 2.5 to 5.0 angstroms of one or more of the coordinates of the ligand in Table 1 or Table 2.

56. The method of Claim 55, wherein the compound occupies a position within about 1.2 to 1.7 or about 2.6 to 3.5 angstroms of one or more of the coordinates of the ligand in at least one of Tables 1-5.

57. A method of identifying a proteasome inhibitor comprising:

(a) identifying one or more interactions between a ligand and a Rac-1 GTPase; and

(b) changing a substituent on the ligand in order to modify the interaction between the ligand the subunit.

58. The method of Claim 57, wherein the ligand is NSC23766.

59. A method of screening a set of one or more compounds to identify compounds most likely to bind to a Rac-1 GTPase comprising the steps of:

(a) computationally determining the docking energy of each of one or more compounds in an activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site;

(b) selecting one or more compounds indicated as having a likelihood of binding to said Rac-1 GTPase activation site above a threshold value.

60. A method of identifying a compound that is likely to bind an activation site of a Rac-1 GTPase comprising the steps of:

(a) computationally determining the docking energy of each of one or more compounds in an activation site of a Rac-1 GTPase, wherein each respective docking energy indicates the likelihood of said compound binding to said Rac-1 GTPase activation site;

(b) selecting one or more compounds indicated as having a likelihood of binding to said Rac-1 GTPase activation site above a threshold value.

61. The method of any of Claims 59-60, wherein said activation site comprises one or more of the protein atoms of Table 1 or Table 2.

62. Atomic coordinates that have a root-mean-squared deviation of 6.0 Å or less, 5.5 Å or less, 5.0 Å or less, 4.5 Å or less, 4.0 Å or less, 3.5 Å or less, 3.0 Å or less, 2.5 Å or less, 2.0 Å or less, 1.7 Å or less, 1.5 Å or less, 1.4 Å or less, 1.3 Å or less, 1.2 Å or less, 1.1 Å or less, 1.0 Å or less, 0.9 Å or less, 0.8 Å or less, 0.7 Å or less, 0.6 Å or less, 0.5 Å or less, 0.4 Å or less, 0.3 Å or less, 0.2 Å or less, or 0.1 Å or less, relative to atoms of the atomic coordinates provided in Table 1 or Table 2.

63. The method of Claim 59, wherein said activation site comprises protein atoms having atomic coordinates that have a root-mean-squared deviation of 6.0 Å or less, 5.5 Å or less, 5.0 Å or less, 4.5 Å or less, 4.0 Å or less, 3.5 Å or less, 3.0 Å or less, 2.5 Å or less, 2.0 Å or less, 1.7 Å or less, 1.5 Å or less, 1.4 Å or less, 1.3 Å or less, 1.2 Å or less, 1.1 Å or less, 1.0 Å or less, 0.9 Å or less, 0.8 Å or less, 0.7 Å or less, 0.6 Å or less, 0.5 Å or less, 0.4 Å or less, 0.3 Å or less, 0.2 Å or less, or 0.1 Å or less, relative to atoms of the atomic coordinates provided in Table 1 or Table 2.

64. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid of switch I and at least one atom from an amino acid of switch II.

65. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase (see SEQ ID

NO:1) and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase (see SEQ ID NO:1).

66. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 and Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase.

67. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase.

68. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-74 of Rac-1 GTPase.

69. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from Trp56 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

70. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

71. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Leu70 and Ser71 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

72. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Tyr64, Leu67 and Leu70 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

73. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 and at least one atom from an amino acid selected from amino acids 30-39 of Rac-1 GTPase.

74. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Thr35, Val36, Phe37 or Asp38 of Rac-1 GTPase and at least one atom from an amino acid selected from amino acids 56-70 of Rac-1 GTPase.

75. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Gly30, Tyr32, Ile33, Pro34, Val36 and Phe37 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase.

76. The atomic coordinates of Claim 62 or the method of Claim 63, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include at least one atom from an amino acid selected from Thr35 and Val36 of Rac-1 GTPase and at least one atom from an amino acid selected from Trp56, Ala59, Tyr64, Leu67, Leu70, Ser71 and Pro73 of Rac-1 GTPase.

77. The atomic coordinates of Claim 62, wherein the atoms of the atomic coordinates provided in Table 1 or Table 2 include the Mg^{2+} atom.

78. The atomic coordinates of Claim 62, wherein the atoms are selected from the group consisting of all atoms of a selected amino acid, backbone atoms only of a selected amino acid, and alpha carbon atoms only of a selected amino acid.

79. The method of Claim 63, further comprising determining the conformation of said compound in said activation site.

80. The method of Claim 63, wherein said step of determining the docking energy further comprises a step of generating a grid-based receptor field representation of said activation site.

81. The method of Claim 63, wherein said step of determining the docking energy further comprises a step of determining the optimal conformation of said compound in said activation site.

82. The method of Claim 63, wherein said step of determining the docking energy further comprises a step of building atoms of said compound into said activation site to thereby match said activation site.

83. A computer readable media comprising the atomic coordinates of Claim 62.

84. The computer readable media of Claim 83, comprising the atomic coordinates of the protein atoms of Table 1 or Table 2.

85. A computer readable media comprising the the atomic coordinates for a ligand of Table 1 or Table 2.

86. The computer readable media of Claim 85, wherein the ligand is NSC23766.

87. A computer readable media comprising at least a portion of the atomic coordinates for a GDP-bound Rac-1 GTPase.

88. The computer readable media of Claim 87, comprising amino acids 1-185 of Rac-1 GTPase.

89. The computer readable media of Claim 87, further comprising at least a portion of the atomic coordinates of a non-GDP ligand of Rac-1 GTPase.

90. The computer readable media of Claim 87 or Claim 88, wherein the coordinates are less than the coordinates for amino acids 1-185 of Rac-1 GTPase.

91. A computer system comprising the coordinates of Claim 62.

92. The computer system of Claim 91, capable of generating structures and/or performing rational drug design for a proteasome/ligand complex.

1/6




Fig. 1

2/6




Fig. 2

3/6




Fig. 3

4/6




Fig. 4

5/6




Fig. 5

6/6




Fig. 6

SEQUENCE LISTING

<110> Children's Hospital Medical Center
 The State University of New York, Stony Brook University
 Zheng, Yi
 Nassar, Nicolas
 Skowronek, Karlheinz R.

<120> GTPase INHIBITORS AND METHODS OF USE AND
 CRYSTAL STRUCTURE OF RAC-1 GTPase

<130> CHMC24.001VPC2

<150> 60/703,587

<151> 2005-07-29

<160> 1

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 192

<212> PRT

<213> Human

<400> 1

Met	Gln	Ala	Ile	Lys	Cys	Val	Val	Val	Gly	Asp	Gly	Ala	Val	Gly	Lys
1				5					10					15	
Thr	Cys	Leu	Leu	Ile	Ser	Tyr	Thr	Thr	Asn	Ala	Phe	Pro	Gly	Glu	Tyr
			20					25					30		
Ile	Pro	Thr	Val	Phe	Asp	Asn	Tyr	Ser	Ala	Asn	Val	Met	Val	Asp	Gly
			35				40					45			
Lys	Pro	Val	Asn	Leu	Gly	Leu	Trp	Asp	Thr	Ala	Gly	Gln	Glu	Asp	Tyr
			50			55					60				
Asp	Arg	Leu	Arg	Pro	Leu	Ser	Tyr	Pro	Gln	Thr	Asp	Val	Phe	Leu	Ile
65					70				75						80
Cys	Phe	Ser	Leu	Val	Ser	Pro	Ala	Ser	Phe	Glu	Asn	Val	Arg	Ala	Lys
				85					90					95	
Trp	Tyr	Pro	Glu	Val	Arg	His	His	Cys	Pro	Asn	Thr	Pro	Ile	Ile	Leu
			100					105					110		
Val	Gly	Thr	Lys	Leu	Asp	Leu	Arg	Asp	Asp	Lys	Asp	Thr	Ile	Glu	Lys
			115				120					125			
Leu	Lys	Glu	Lys	Lys	Leu	Thr	Pro	Ile	Thr	Tyr	Pro	Gln	Gly	Leu	Ala
			130			135					140				
Met	Ala	Lys	Glu	Ile	Gly	Ala	Val	Lys	Tyr	Leu	Glu	Cys	Ser	Ala	Leu
145					150				155						160
Thr	Gln	Arg	Gly	Leu	Lys	Thr	Val	Phe	Asp	Glu	Ala	Ile	Arg	Ala	Val
				165					170					175	
Leu	Cys	Pro	Pro	Pro	Val	Lys	Lys	Arg	Lys	Arg	Lys	Cys	Leu	Leu	Leu
			180					185					190		