

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 February 2008 (28.02.2008)

PCT

(10) International Publication Number
WO 2008/023248 A2

(51) International Patent Classification:
A61K 31/7048 (2006.01)

Maharashtra (IN). **DIWAKAR, Santosh, Devidas** [IN/IN]; At Post:Bhanashiware, Tal-Newasa, Dist-Ahmednagar-414609, Maharashtra (IN). **JADHAV, Ganesh, Rajaram** [IN/IN]; Plot No.36, Shahunagar, Kamladevi Road, Karmala, Dist:Solapur-413203, Maharashtra (IN). **PAWAR, Shivaji, Sampatrao** [IN/IN]; A/P Jalgaon, Tal:Vaijapur, Dist:Aurangabad-423702, Maharashtra (IN).

(21) International Application Number:
PCT/IB2007/002405

(74) **Agent:** **KODGULE, Mandar, Madhukar**; Deputy General Manager - Intellectual Property, Wockhardt Limited, D-4, MIDC Area, Chikalthana, Aurangabad-431210, Maharashtra (IN).

(22) International Filing Date: 22 August 2007 (22.08.2007)

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1336/MUM/2006 24 August 2006 (24.08.2006) IN

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) **Applicant** (for all designated States except US): **WOCKHARDT RESEARCH CENTRE** [IN/IN]; c/o Wockhardt Limited, D-4, MIDC Area, Chikalthana, Aurangabad-431210, Maharashtra (IN).

Published:

— without international search report and to be republished upon receipt of that report

(54) **Title:** NOVEL MACROLIDES AND KETOLIDES HAVING ANTIMICROBIAL ACTIVITY

(57) **Abstract:** The present invention provides compounds having antimicrobial activity for preventing and treating diseases caused by microbial infections. Thus, the present invention relates to novel • semi-synthetic 11,12- γ lactone macrolides and ketolides having antimicrobial activity, processes for making compounds as well as pharmaceutical compositions containing said compounds as active ingredients and methods of treating microbial infections with the compounds.

Novel Macrolides and Ketolides Having Antimicrobial Activity**Field of the Invention**

The present invention relates to novel 11,12- γ lactone macrolides and ketolides having antimicrobial activity, pharmaceutical compositions containing the compounds and methods of treating microbial infections with the compounds.

Background of the Invention

Macrolides are a well-known family of antimicrobial agents, erythromycin A, 14-membered macrolide, was isolated in 1952 from *Streptomyces erythraeus*. Examples of macrolides being used as therapeutic agents are roxithromycin, clarithromycin and azithromycin (azalide). Ketolides are semisynthetic 14-membered ring macrolide derivatives, characterized by the presence of a keto function at position 3 instead of L-cladinose moiety present in the macrolactone ring. Telithromycin and Cethromycin are examples of ketolides.


United States patent US 4,331,803 discloses the 6-O-methyl derivative of erythromycin i.e. clarithromycin. The patent US 4,349,545 discloses roxithromycin. The azalide azithromycin is disclosed in US 4,517,359. Telithromycin is described in EP 680967 A1 and corresponding US 5,635,485 and *Bioorg. Med. Chem. Lett.* 1999, 9(21), 3075-3080. Another ketolide Cethromycin (ABT 773) is disclosed in WO 98/09978, and *J. Med. Chem.* 2000, 43, 1045.

The patent US 6,900,183 describes 11,12- γ lactone ketolides having C-21 of the lactone substituted with cyano or amino derivatives.

The patent application US 2004/0077557 and PCT publications WO 02/16380, WO 03/42228 WO 04/16634 and WO 03/072588 discloses 11,12- γ lactone ketolides.

Description of the Invention

In one general aspect there is provided macrolide and ketolide compounds of Formula I



Formula I

and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers,
wherein


* indicates a chiral center;

R_1 is hydrogen or methyl;

R_2 is hydrogen or hydroxyl protecting group,


wherein hydroxyl protecting groups are selected from the group consisting of triethylsilyl, trimethylsilyl, acetyl, benzoyl, methoxymethyl, benzyl, methoxymethoxymethyl or tertbutyldimethylsilyl;

Q is Het or



wherein

Het is selected from



wherein R_3 is one or more substituent selected from the group consisting of NO_2 , CN , halogen, C_1-C_6 alkyl, C_2-C_6 alkenyl, CH_2CONH_2 , CH_2CO_2Et , CH_2CN , CH_2CH_2OH , $CH_2OCH_2CH_2OCH_3$, NH_2 , substituted C_1-C_6 alkyl and substituted C_2-C_6 alkenyl;

T is selected from the group consisting of:

hydrogen, $-(CH_2)_m-R_5$, $-(CH_2)_m-CH=CH-R_5$, $-(CH_2)_m-C\equiv C-R_5$, $-A-(CH_2)_m-R_5$, $-A-(CH_2)_m-CH=CH-R_5$, $-A-(CH_2)_m-C\equiv C-R_5$, $-(CH_2)_m-B-R_5$, $-A-(CH_2)_m-B-R_5$, $-(CH_2)_m-X-R_6$, $-(CH_2)_m-B-X-R_6$, $-(CH_2)_m-CH=CH-X-R_6$, $-(CH_2)_m-C\equiv C-X-R_6$, $-A-(CH_2)_m-X-R_6$, $-A-(CH_2)_m-B-X-R_6$, $-A-(CH_2)_m-CH=CH-X-R_6$ and $-A-(CH_2)_m-C\equiv C-X-R_6$;

wherein,

m is 0, 1, 2 or 3;


R_5 is selected from the group consisting of:

hydrogen, cyano, halogen, hydroxyl, $CO_2(C_1-C_6$ alkyl), $CO NR_aR_b$, NR_aR_b , C_1-C_6 alkyl or substituted C_1-C_6 alkyl, C_2-C_6 alkenyl or substituted C_2-C_6 alkenyl, C_2-C_6 alkynyl or substituted C_2-C_6 alkynyl, 6 membered substituted or unsubstituted aryl, 5-6 membered substituted or unsubstituted heteroaryl, 3-6 membered substituted or unsubstituted cycloalkyl and 3-6 membered substituted or unsubstituted heterocyclyl;

wherein R_a and R_b are independently hydrogen, C_1-C_6 alkyl or R_a and R_b together with the nitrogen to which they are attached form a 5-6 membered heterocyclic ring wherein the heterocycle has one or more heteroatoms selected from N, O, S;

A is $-CO-$ or $-CONH-$;

B is $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-CO-$, $-CONH-$, $-CON(CH_3)-$, $-NHCONH-$, $-C(NH_2)=N-O-$,




wherein

R_8 is a substituent at any one of carbon of the heterocycle, R_8 is selected from the group comprising of hydrogen, C_1 - C_6 alkyl, substituted C_1 - C_6 alkyl, C_2 - C_6 alkenyl and substituted C_2 - C_6 alkenyl;

X is a 6 membered aryl or a 5-6 membered heteroaryl;

R_6 is aryl, heteroaryl, substituted aryl or substituted heteroaryl;

Y' is hydrogen and Y is OR_7 , wherein R_7 is hydrogen or




wherein R_2' is hydrogen, (when R_2' is hydrogen, R_7 is designated as cladinoose) or hydroxyl protecting group,

wherein hydroxyl protecting groups are selected from the group consisting of triethylsilyl, trimethylsilyl, acetyl, benzoyl, methoxymethyl, methoxyethoxymethyl, benzyl or tertbutyldimethylsilyl;

or


Y and Y' together with the carbon to which they are attached form $C=O$.

In one embodiment there is provided macrolide compound of Formula I-a, wherein,




wherein Q , R_1 , R_2 and R_2' are as defined above.

Another embodiment relates to macrolide compound of Formula I-b,




wherein R_1 and Het are as defined above.

Another embodiment relates to compound of Formula I-c,



wherein T and R_1 are as defined above.


Another embodiment relates to ketolide compound of Formula I-d,



Formula I-d

wherein R_1 is CH_3 , R_2 and Q are as defined above.


Another embodiment relates to ketolide compound of the invention of Formula I-e,



Formula I-e

wherein Het is as defined above.

Another embodiment relates to ketolide compound of Formula I-f,



Formula I-f

wherein T is as defined above.

Another embodiment relates to pharmaceutical compositions of compounds of Formula I and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers, together with one or more pharmaceutically acceptable carriers, excipients or diluents, for the treatment and prevention of a microbial infection in human or animal.

In another embodiment the compounds of Formula I and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers.


Yet another embodiment is also directed to a method of treating a subject having a condition caused by or contributed to by microbial infection, which comprises administering to the said subject a therapeutically effective amount of compound of Formula I and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers.

Another embodiment is further directed to a method of preventing a subject from suffering from a condition caused by or contributed to by microbial infection, which comprises administering to the subject a prophylactically effective amount of compound of Formula I and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers.

Another embodiment relates to method of preparation of the compounds of Formula I and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers.

Other objects and advantages will become apparent to those skilled in the art from a review of the ensuing specification.

Representative compounds of the invention are



Compound of Formula I-b where Het is 1H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 2-(2-methoxy-ethoxymethyl)-2H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 1-(2-methoxy-ethoxymethyl)-2H-tetrazol-5-yl, R₁ is CH₃;


Compound of Formula I-b where Het is 2-(ethoxycarbonylmethyl)-2H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 1-(ethoxycarbonylmethyl)-2H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 2-(carbamoylmethyl)-2H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 1-(carbamoylmethyl)-2H-tetrazol-5-yl, R₁ is CH₃;

Compound of Formula I-b where Het is 2-(cyanomethyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 1-(cyanomethyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 2-(allyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 1-(allyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 2-(hydroxyethyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 1-(hydroxyethyl)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 2-(carboxylic acid(3-fluoro-phenyl)-methylamide)-2H-tetrazol-5-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 5-methyl-[1,2,4]oxadiazol-3-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 4,5-dihydro-1H-imidazol-2-yl, R₁ is CH₃;
 Compound of Formula I-b where Het is 1,4,5,6-tetrahydro-pyrimidin-2-yl, R₁ is CH₃;



Compound of Formula I-c where T is H, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(pyridin-3-yl), R₁ is CH₃;
 Compound of Formula I-c where T is COCH₂O-(2-chlorophenyl), R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(4-dimethylamino)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO-(pyridin-3-yl), R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(pyridin-4-yl), R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(3-methoxy)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(4-methoxy)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(3-cyano)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(4-cyano)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is COCH₂O-(3-chloro)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(3,5-dimethoxy)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(2,3-dimethoxy)phenyl, R₁ is CH₃;
 Compound of Formula I-c where T is CO(CH₂)₂-(2,5-dimethoxy)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(3-fluoro)phenyl, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-trifluoromethyl)phenyl, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(4-methyl-piperazine)phenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(2-methylphenyl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(2-methoxyphenyl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(3-methoxyphenyl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(2-fluorophenyl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(pyrimidin-2-yl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(pyridin-2-yl)-piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -phenyl, R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(2-methylphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(3-methylphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(4-methylphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(2-methoxyphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(4-methoxyphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(2,3,4,5,6-pentafluorophenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2O -(4-cyanophenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(phenyl)piperazinyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2S -phenyl, R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2S -(4-fluorophenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2S -(2-fluorophenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2S -(3-fluorophenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is COCH_2S -(3-methoxyphenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-([1,2,3]-triazol-1-yl)phenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(imidazol-1-yl)phenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-([1,2,4]-triazol-1-yl)phenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(4-acetyl-[1,2,3]triazol-1-yl))-phenyl, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(4-(4-cyano-[1,2,4]-triazol-1-yl)phenyl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2$ -(benzo[1,3]dioxol-5-yl), R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{COCH}_2\text{-(benzo[1,3]dioxol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(4-methoxyphenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(pyridin-3-yl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(4-fluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(3-methylphenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(3-chlorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2-fluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(3,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2,3,4-trifluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2-fluoro-4-methyl-phenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2-fluoro-4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(2,3-difluoro-4-ethoxy-phenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(3-(4-fluoro-phenyl)-[1,2,4]oxadiazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(3-(3-bromo-phenyl)-[1,2,4]oxadiazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(3-(4-nitro-phenyl)-[1,2,4]oxadiazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-(3-(3-fluoro-phenyl)-[1,2,4]oxadiazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-chloro-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-methoxy-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3,5\text{-dimethoxy-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(4\text{-chloro-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(4\text{-bromo-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(pyridin-4-yl)\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(pyridin-3-yl)\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(2\text{-fluoro-5-methoxy-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-phenyl-4H-imidazol-1-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CONH-}(4\text{-methoxy-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CONH-}(3\text{-methoxy-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CONH-}(2\text{-methoxy-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CONH-}(2,4\text{-difluorophenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CO-}(4\text{-}(3\text{-fluoro-phenyl)piperazin-1-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{CH}_3$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{OH}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{OC}_2\text{H}_5$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}(\text{CH}_3)_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2(\text{CH}_2)_2\text{-phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(4\text{-nitro-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(4\text{-bromo-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(2\text{-fluoro-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(3,5\text{-difluoro-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(2,4\text{-difluoro)\text{-phenyl}}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(4\text{-methyl)\text{-phenyl}}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(3\text{-methoxy)\text{-phenyl}}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(2\text{-methyl)\text{-phenyl}}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{-}(4\text{-}(pyridin-2-yl)piperazinyl)$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{-}(1\text{-}(3\text{-methoxy)phenyl-1H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(1\text{-}(4\text{-methoxy-2-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(5\text{-methyl-[1,2,4]oxadiazol-3-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-fluoro-2-methoxyphenyl)-isoxazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH}_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{CH=CH}_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(2-methoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(3-fluoro)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(4-cyano)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}(\text{CH}_3)\text{=CH}_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=C}(\text{CH}_3)_2$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(3-cyano)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(3-acetoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(3-hydroxymethyl)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(4-trifluoromethyl)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(3-trifluoromethyl)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CH=CH-(2,3-dihydro-benzo[1,4]dioxin-6-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is CH_2CN , R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}(\text{NH}_2)\text{=N-OH}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{CH}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{C-phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{C-(3-cyano)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{C-(3-methyl)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is COCH=CH-phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CO(CH₂)₂-phenyl, R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(pyridin-4-yl), R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(pyridin-3-yl), R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(pyridin-2-yl), R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(1-(2-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl) , R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(1-(3-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl) , R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(1-(4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl) , R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(1-(4-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl) , R₁ is CH₃;

Compound of Formula I-c where T is COCH=CH-(1-(3-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl) , R₁ is CH₃;

Compound of Formula I-c where T is COCH₂NHCONH-phenyl, R₁ is CH₃;

Compound of Formula I-c where T is COCH₂NHCONH-(4-fluorophenyl), R₁ is CH₃;

Compound of Formula I-c where T is COCH₂NHCONH-(3-chloro-4-methylphenyl), R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH₂, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(3-fluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(2-fluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(4-fluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(2,4-difluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(3,4-difluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(2,3,4-trifluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(3-chloro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(2-methoxy)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(3-methoxy)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-(3,5-dimethoxy)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CONH-cyclopropyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CON(CH₃)-(3-fluoro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is CH₂CON(CH₃)-(4-chloro)phenyl, R₁ is CH₃;

Compound of Formula I-c where T is $\text{CH}_2\text{CONH}-\text{(6-pyrazol-1-yl-pyridin-3-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CONH}-\text{(3-(pyrazol-1-yl)-pyridin-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(4-fluorophenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(3-fluorophenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(3,5-difluoro phenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(3,4-difluoro phenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(3,4,5-trifluoro phenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(1-(3-chlorophenyl)-1H-[1,2,3]-triazol-5-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2-\text{(5-(pyridin-3-yl)-[1,2,4]-oxadiazol-3-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$, R_1 is CH_3 ;

Compound of Formula I-c where T is CONH-phenyl , R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CONH-}(4\text{-ethyl-phenyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CO-}(4\text{-(phenyl)-piperazinyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CH}_2\text{CO-}(4\text{-(2-fluorophenyl)-piperazinyl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{CO-}(4\text{-phenyl)-piperazinyl}$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}[3\text{-(4-methoxy-phenyl)-[1,2,4]oxadiazol-5-yl}]$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-pyridin-3-yl-1H-[1,2,3]-triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-(2-methoxy)phenyl)-1H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-(3-fluorophenyl)-1H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-phenyl-[1,2,4]oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-naphthalen-2-yl-[1,2,4]oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(\text{pyridin-3-yl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,5\text{-difluoro-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;


Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,4\text{-difluoro-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(4\text{-}(2\text{-methoxy-phenyl})\text{-piperazine-1-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,5\text{-dimethoxy-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3\text{-methoxy-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(2\text{-fluoro-5\text{-methoxy-phenyl})-[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;



Formula I-e

Compound of Formula I-e where Het is 1H-tetrazol-5-yl;

Compound of Formula I-e where Het is 5-methyl-[1,2,4]oxadiazol-3-yl;

Compound of Formula I-e where Het is 4,5-dihydro-1H-imidazol-2-yl;

Compound of Formula I-e where Het is 1,4,5,6-tetrahydro-pyrimidin-2-yl;

Compound of Formula I-e where Het is 2-(2-methoxy-ethoxy)-2H-tetrazol-5-yl;

Compound of Formula I-e where Het is 1-(2-methoxy-ethoxy)-1H-tetrazol-5-yl;

Compound of Formula I-e where Het is [1,2,4]-oxadiazol-3-yl;

Compound of Formula I-e where Het is (5-trifluoromethyl)-[1,2,4]-oxadiazol-3-yl;

Compound of Formula I-e where Het is (4,5)-dihydro-1H-imidazol-2-yl;

Compound of Formula I-e where Het is 1-(allyl)-1H-tetrazol-5-yl

Compound of Formula I-e where Het is 2-(allyl)-2H-tetrazol-5-yl

Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{C}-(3\text{-methoxy)phenyl});$
Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{C}-(\text{pyridin-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{C}-(2\text{-chloropyridin-5-yl});$
Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{C}-(2\text{-fluoropyridin-5-yl});$
Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{C}-(\text{pyridin-3-yl});$
Compound of Formula I-f where T is $\text{CH}_2\text{-phenyl};$
Compound of Formula I-f where T is $\text{CH}_2(\text{CH}_2)_2\text{-phenyl};$
Compound of Formula I-f where T is $\text{CH}_2-(4\text{-methoxyphenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(3\text{-chlorophenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(3\text{-methoxyphenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(2\text{-fluorophenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(4\text{-(isopropyl)phenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(4\text{-(1H-[1,2,4]-triazol)phenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(4\text{-(pyrimidin-5-yl)phenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(4\text{-(pyridin-2-yl)phenyl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-phenyl[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(pyridin-3-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(3-fluorophenyl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(4-methoxyphenyl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(pyridin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(pyrazin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(3,5-dimethoxyphenyl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(pyrimidin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(6-methoxy-pyridin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-((5\text{-cyclopropyl)-[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(5-methyl-pyridin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(5-cyclopropyl-pyridin-2-yl)[1,3,4]oxadiazol-2-yl})$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(5-cyano-pyridin-2-yl)[1,3,4]oxadiazol-2-yl});$
Compound of Formula I-f where T is $\text{CH}_2-(5\text{-(5-dimethylamino-pyridin-2-yl)[1,3,4]oxadiazol-2-yl});$

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methoxy-pyridin-2-yl)}[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-fluoro-pyridin-2-yl)}[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-chloro-pyridin-2-yl)}[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyrimidin-5-yl)}[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-phenyl}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-fluorophenyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-chlorophenyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(4\text{-chlorophenyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(4\text{-methoxyphenyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3,5\text{-dimethoxyphenyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyridin-3-yl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyridin-2-yl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyrimidin-2-yl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-cyclopropyl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(6\text{-methylpyridin-2-yl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-methylpyridin-2-yl)}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-methyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-phenyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-fluorophenyl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-chlorophenyl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-fluoro-4\text{-methoxyphenyl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-cyanophenyl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyridin-3-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyrazin-2-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyridin-2-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyridin-4-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-chloropyridin-3-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-methoxypyridin-3-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-fluoropyridin-3-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-cyclopropyl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(6\text{-methylpyridin-2-yl)}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-methyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-trifluoromethyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyridazin-2-yl)-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-methyl-oxazol-4-yl)-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-methyl-thiazol-4-yl)-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyrimidin-2-yl)-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyridin-3-yl)-isoxazol-5-yl)$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-fluorophenyl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-chlorophenyl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(4\text{-fluorophenyl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyridin-2-yl)-isoxazol-5-yl)$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(3\text{-methoxyphenyl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-phenyl-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-methyl-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(6\text{-methoxy-pyridin-2-yl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(2\text{-methoxy-pyridin-5-yl)-isoxazol-5-yl})$;

Compound of Formula I-f where T is CH_2CONH_2 ;

Compound of Formula I-f where T is $\text{CH}_2\text{CON}(\text{CH}_3)_2$;

Compound of Formula I-f where T is $\text{CH}_2\text{CONH-}(cyclopropyl)$;

Compound of Formula I-f where T is $\text{CH}_2\text{CONH-}(3\text{-chloro-phenyl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{CONH-}(3\text{-fluoro-phenyl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{CO-}(4\text{-phenyl)piperazin-1-yl})$;

Compound of Formula I-f where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-fluorophenyl})$;

Compound of Formula I-f where T is $\text{CO}(\text{CH}_2)_2\text{-}(4\text{-methoxyphenyl})$;

Compound of Formula I-f where T is $\text{CO}(\text{CH}_2)_2\text{-}(2,3\text{-dimethoxyphenyl})$;

Compound of Formula I-f where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-fluorophenyl)-[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CONH-}(4\text{-ethylphenyl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-methylpyridin-2-yl)[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-cyclopropyl-pyridin-2-yl)[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-cyano-pyridin-2-yl)[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-dimethyl amino-pyridin-2-yl)[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-methoxy-pyridin-2-yl)[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-fluoro-pyridin-2-yl})[1,2,4]\text{ oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-chloro-pyridin-2-yl})[1,2,4]\text{ oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyrimidin-5-yl)[1,2,4]\text{ oxadiazol-5-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methyl-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-cyclopropyl-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-cyano-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-dimethyl amino-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methoxy-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-fluoro-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-chloro-pyridin-2-yl})-[1,2,4]\text{ oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyrimidin-5-yl)-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-fluorophenyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-chlorophenyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(4\text{-fluorophenyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(3\text{-methoxy phenyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-phenyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methyl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(6\text{-methoxy-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(2\text{-methoxy-pyridin-5-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methyl-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-cyclopropyl-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-cyano-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-dimethyl amino-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-methoxy-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-fluoro-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(5\text{-chloro-pyridin-2-yl)-isoxazol-3-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(pyrimidin-5-yl)-isoxazol-3-yl)$;

Compound of Formula I-f where T is $\text{CH}_2\text{CH}_2\text{-}(4\text{-}(pyridin-2-yl)-1\text{H-imidazol-1-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{CH}_2\text{-}(4\text{-}(pyridin-3-yl)-1\text{H-imidazol-1-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{CH}_2\text{-}(4\text{-}(6\text{-methyl-pyridin-2-yl)-1\text{H-imidazol-1-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{CH}_2\text{-}(4\text{-}(4\text{-methoxy-pyridin-2-yl)\text{-}1H-imidazol-1-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(4\text{-}(pyridin-2-yl)\text{-}pyrazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(4\text{-}(pyridin-3-yl)\text{-}pyrazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(4\text{-}(6\text{-methyl-pyridin-2-yl)\text{-}pyrazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(4\text{-}(4\text{-methoxy-pyridin-2-yl)\text{-}pyrazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(4\text{-}(5\text{-methoxy-pyridin-2-yl)\text{-}pyrazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(1\text{-}(pyridin-2-yl)\text{-}1H\text{-}imidazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(1\text{-}(pyridin-3-yl)\text{-}1H\text{-}imidazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(1\text{-}(6\text{-methyl-pyridin-2-yl)\text{-}1H\text{-}imidazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(1\text{-}(4\text{-methoxy-pyridin-2-yl)\text{-}1H\text{-}imidazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(1\text{-}(5\text{-methoxy-pyridin-2-yl)\text{-}1H\text{-}imidazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(2\text{-}(pyridin-2-yl)\text{-}oxazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(2\text{-}(pyridin-3-yl)\text{-}oxazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(2\text{-}(6\text{-methyl-pyridin-2-yl)\text{-}oxazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(2\text{-}(4\text{-methoxy-pyridin-2-yl)\text{-}oxazol-4-yl})$;

Compound of Formula I-f where T is $\text{CH}_2\text{-}(2\text{-}(5\text{-methoxy-pyridin-2-yl)\text{-}oxazol-4-yl})$;

The compounds of present invention include stereoisomers. The term "stereoisomer" refers to compounds, which have identical chemical composition, but differ with regard to arrangement of the atoms and the groups in space. These include enantiomers, diastereomers, geometrical isomers, atropisomer and conformational isomers. Geometric isomers may occur when a compound contains a double bond or some other feature that gives the molecule a certain amount of structural rigidity. An enantiomer is a stereoisomer of a reference molecule that is the nonsuperimposable mirror image of the reference molecule. A diastereomer is a stereoisomer of a reference molecule that has a shape that is not the mirror image of the reference molecule. An atropisomer is a conformation of a reference compound that converts to the reference compound only slowly on the NMR or laboratory time scale. Conformational isomers (or conformers or rotational isomers or rotamers) are stereoisomers produced by rotation about σ bonds, and are often rapidly interconverting at room temperature. Racemic mixtures are also encompassed within the scope of this invention. Some of the compounds of the present invention may have *trans* and *cis* isomers and geometric E- and Z- isomers. The wavy bond indicates that the compounds may be present as either of E- or Z- isomer. Also some of the compounds according to this invention may exist as diastereomers. In addition,

where the process for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers, may be separated by conventional techniques such as preparative chromatography and HPLC. The compounds may be prepared as a single stereoisomer or in racemic form as a mixture of some possible stereoisomer.

Furthermore, some of the compounds may exists as different polymorphs such as cyrtalline or amorphous forms and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e. hydrates) which, contains various amounts of water, for instance the hydrate, hemihydrate and sesquihydrate forms. Also the compound can form solvates with common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

Description of terms:

The following definitions are used, unless otherwise described.

The term “C₁-C₆ alkyl” refers to saturated, straight or branched chain hydrocarbon radicals containing between one and six carbon atoms. Examples of C₁-C₆ alkyl radicals, include but are not limited to, methyl, ethyl, propyl, butyl, pentyl, hexyl, and their branched isomers such as iso-propyl, iso-butyl, tert-butyl.

The term “C₂-C₆ alkenyl” refers to straight or branched-chain hydrocarbon radicals comprising two to six carbon atoms, which contain one or more carbon-carbon double bonds. Examples of C₂-C₆ alkenyl radicals, include but are not limited to, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like.

The term “C₂-C₆ alkynyl” refers to straight or branched chain hydrocarbon radicals comprising two to six carbon atoms respectively, which contain one or more carbon-carbon triple bonds. Examples of C₂-C₆ alkynyl radicals, include but are not limited to, ethynyl, propynyl, butynyl, 1-methyl-2-butyn-1-yl, and the like.

The substituted C₁-C₆ alkyl, substituted C₂-C₆ alkenyl, C₂-C₆ substituted alkynyl is defined as the respective alkyl, alkenyl, alkynyl substituted, by independent replacement of one or two or three of the hydrogen atoms thereon with Cl, Br, F, I, NO₂, CN, OH, haloalkyl, C₁-C₆ alkoxy, amino, alkylamino, dialkylamino, mercapto, formyl, carboxy, alkoxy carbonyl and carboxamide (C(O)NR'R''), aryl, heteroaryl, substituted aryl, substituted heteroaryl.

Examples of C₁-C₆ alkoxy are methoxy, ethoxy, propyloxy, isopropyloxy, butyloxy, pentyloxy, hexyloxy.

The term "haloalkyl" denotes an alkyl group, as defined above, having one, two or three halogen atoms attached thereto and is exemplified by such groups as chloromethyl, bromoethyl, trifluoromethyl, and the like.

The term "alkylamino" refers to a group having the structure $-\text{NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$ where $\text{C}_1\text{-C}_6$ alkyl is as previously defined.

The term "dialkylamino" refers to a group having the structure $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})(\text{C}_1\text{-C}_6 \text{ alkyl})$, where $\text{C}_1\text{-C}_6$ alkyl is as previously defined. Examples of dialkylamino are, but not limited to, dimethylamino, diethylamino, methylethylamino and the like.

The term "aryl" refers to a mono (6-membered) or bicyclic carbocyclic ring system having one or two aromatic ring including but not limited to phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like.

The term "heteroaryl" refers to a mono i.e. 5-6 membered or bicyclic i.e. fused aromatic ring system having at least one carbon atom of the aromatic ring replaced by an atom selected from the group of N, O, S. The ring may be connected to the remaining part of the molecule via any of the ring atoms. For example pyridinyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, thiadiazolyl, oxadiazolyl, thienyl, triazolyl, triazinyl, tetrazolyl, furanyl, N-oxo-pyridinyl, and the like. It includes the fused biaryl systems such as indolyl, quinolinyl, isoquinolinyl, benzothiazolyl, benzoxazolyl, benzothienyl, N-oxo-quinolyl, benzimidazolyl, benzopyranyl, benzoisothiazolyl, benzodiazinyl, benzofurazanyl, indazolyl, indolizinyl, benzofuryl, quinoxalinyl, pyrrolopyridinyl, furopyrnidinyl (such as furo[2,3-c]pyridinyl, furo[3,2-b]pyridinyl, furo[2,3-b]pyridinyl), naphthyridinyl, phthalazinyl, pyridinopyridinyl, quinazolinyl, thienofuryl, thienopyridinyl, thienotheinyl, purinyl (such as 9H-purin-1-yl, 6-amino-9H-purin-9-yl), pyridinyl-1H-pyrazol-1-yl and the like.

The aryl or the heteroaryl group can be optionally substituted by independent replacement of one or more of hydrogen atoms thereon with substituents selected from $\text{C}_1\text{-C}_6$ alkyl, substituted $\text{C}_1\text{-C}_6$ alkyl, cyano, hydroxy, halogen, amino, formyl, carboxy, carboxamide, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ thioalkoxy, $\text{C}_1\text{-C}_6$ alkylcarbonyl, amino, alkylamino, dialkylamino, mercapto, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, alkylthio, arylthio, heteroarylthio, haloalkyl.

The "aralkyl" is defined as an alkyl group substituted with an aryl group for example benzyl, benzhydryl, trityl, wherein the aryl group may be optionally substituted as defined below.

The “heteroaralkyl” is defined as an alkyl group substituted with a heteroaryl group for example pyridinyl-methyl wherein the heteroaryl group may be optionally substituted as defined below.

The “aroyl” group refers to a carbonyl group attached to an aryl group for example benzoyl, wherein the aryl group may be optionally substituted as defined below.

The “heteroaroyl” is defined as a carbonyl group attached to a heteroaryl group for example pyridinyl carbonyl, wherein the heteroaryl group may be optionally substituted as defined below.

The “aralkanoyl” refers a carbonyl group attached to an aralkyl group for example phenylacetyl. The “heteroaralkanoyl” refers a carbonyl group attached to heteroaralkyl group for example pyridinylacetyl. The cycloalkyl refers to 3-7 membered cyclic structures for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl.

The heterocyclyl refers to 3-7 membered cyclic structures including one or more heteroatoms such as N, O, S or can be defined as up to 4 heteroatoms selected from N, O, S. Examples of heterocycls are oxetanyl (oxetane is a four membered ring with an oxygen in the ring oxetanyl is a substituted oxetane). pyrrolidinyl, pyrazolinyl, imidazolinyl, imidazolidinyl, oxazolinyl, oxazolidinyl, isoxazolinyl, thiazolidinyl, isothiazolidinyl, tetrahydrofuryl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 4-oxo piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, morpholinyl, [1,3]-dioxolane and the like.

The “hydroxyl protecting group” as used herein refers to an easily removable group which is known in the art to protect a hydroxyl group against undesirable reaction during synthetic procedures and to be selectively removable. The use of hydroxy-protecting groups against undesirable reactions during a synthesis procedure and any such protecting groups are known cf., for example, T. H. Greene and P.G. M. Wuts, Protective Groups in Organic Synthesis, 2nd edition, John Wiley & sons, New York (1991). Examples of hydroxyl protecting groups include but are not limited to triethylsilyl, trimethylsilyl, acetyl, benzoyl, methoxymethyl, methoxyethoxymethyl, benzyl, tertbutyldimethylsilyl and the like.

The terms “halo”, “halide” and “halogen” as used herein refer to an atom selected from fluorine, chlorine, bromine and iodine. The term “formyl” as used herein refers to a group of Formula $-\text{CHO}$. The term “sulphonyl” refers to a group of Formula $-\text{SO}_2-$. “Thiol” or “mercapto” refers to the group $-\text{SH}$.

The term “C₁-C₆ alkoxy” refers to a C₁-C₆ alkyl group, as previously defined, attached to the parent molecular moiety through an oxygen atom. Examples of C₁-C₆-alkoxy include, but are

not limited to, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, tert-butoxy, neopentoxy and n-hexoxy.

The term "carboxamide" as used herein refers to a group of Formula $-C(O)NR' R''$ wherein R' and R'' are independently selected from hydrogen or C₁-C₆ alkyl or R' and R'' taken together may form a six or five membered heterocycle such as piperidine, pyrrolidine, piperazine and the like.

The term "ester" or "alkoxycarbonyl" refers to an C₁-C₆ alkoxy group attached to the parent molecular moiety through a carbonyl group such as methoxycarbonyl (CO₂Me), ethoxycarbonyl (CO₂Et), and the like.

The term "(C₁-C₆)alkylcarbonyl" unless otherwise indicated, includes -C(O)-alkyl groups wherein "alkyl" is as defined above.

The terms "Me", "Et", "Ph", "allyl" stands for methyl, ethyl, phenyl and CH₂CH=CH₂ respectively. "MEM" stands for methoxyethoxymethyl.

The term "animal" refers to an animal such as a mammal, fish or a bird.

The phrase "pharmaceutically acceptable salt" as used herein refers to one or more salts of the free base of the invention which possess the desired pharmacological activity of the free base and which are neither biologically nor otherwise undesirable. The salts are suitable for use in contact with the tissues of human and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, et al. describe pharmaceutically acceptable salts in detail in J. Pharmaceutical Sciences, 66: 1-19 (1977), incorporated herein by reference. The salts can be prepared in situ during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable acid. These salts may be obtained from inorganic or organic acids. Examples of inorganic acids are hydrochloric acid, nitric acid, perchloric acid, hydrobromic acid, sulphuric acid or phosphoric acid. Examples of organic acids are acetic acid, propionic acid, oxalic acid, glycolic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulphonic acid, p-toluene sulphonic acid, salicyclic acid and the like. Also included are the salts with various amino acids such as alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine or valine or the optically active isomers thereof or the racemic mixtures thereof or dipeptides, tripeptides and polypeptides derived from the monoaminoacid units thereof.

Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malonate, 2-naphthalenesulfonate, nicotinate, oleate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

Salt of an acid moiety in the compound can also be prepared by reacting with a suitable base. These suitable salts are furthermore those of the inorganic or organic bases. Inorganic bases such as KOH, NaOH, Ca(OH)₂, Al(OH)₃. The organic base salts from basic amines such as ethylamine, triethylamine, diethanolamine, ethylenediamine, guanidine or heterocyclic amines such as piperidine, hydroxyethylpyrrolidine, hydroxyethylpiperidine, morpholine, piperazine, N-methyl piperazine and the like or basic amino acids such as optically pure and racemic isomers of arginine, lysine, histidine, tryptophan and the like.

Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate and aryl sulfonate.

"Therapeutically effective amount" means that amount of active compound(s) or pharmaceutical agent(s) that elicit the biological or medicinal response in a tissue system, animal or human sought by a researcher, veterinarian, medical doctor or other clinician, which response includes alleviation of the symptoms of the disease or disorder being treated. The specific amount of active compound(s) or pharmaceutical agent(s) needed to elicit the biological or medicinal response will depend on a number of factors, including but not limited to the disease or disorder being treated, the active compound(s) or pharmaceutical agent(s) being administered, the method of administration, and the condition of the patient.

The term "treatment" unless otherwise indicated, includes the treatment or prevention of a microbial infection as provided in the method of the present invention.

As used herein, unless otherwise indicated, the term "microbial infection(s)" includes bacterial infections and protozoa infections which occur in human or animals including mammals, fish and birds as well as disorders related to bacterial infections and protozoa infections that may be treated or prevented by administering antibiotics such as the

compounds of the present invention. Such bacterial infections and protozoa infections and disorders related to such infections include the following: pneumonia, otitis media, sinusitus, bronchitis, tonsillitis, and mastoiditis related to infection by *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Staphylococcus aureus*, or *Peptostreptococcus* spp.; pharynigitis, rheumatic fever, and glomerulonephritis related to infection by *Streptococcus pyogenes*, Groups C and G streptococci, *Clostridium diphteriae*, or *Actinobacillus haemolyticum*; respiratory tract infections related to infection by *Mycoplasma pneumoniae*, *Legionella pneumophila*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, or *Chlamydia pneumoniae*; uncomplicated skin and soft tissue infections, abscesses and osteomyelitis, and puerperal fever related to infection by *Staphylococcus aureus*, coagulase-positive staphylococci (i.e., *S. epidermidis*, *S. hemolyticus*, etc.), *Streptococcus pyogenes*, *Streptococcus agalactiae*, Streptococcal groups C-F (minute-colony streptococci), *viridans streptococci*, *Corynebacterium minutissimum*, *Clostridium* spp., or *Bartonella henselae*; uncomplicated acute urinary tract infections related to infection by *Staphylococcus saprophyticus* or *Enterococcus* spp.; urethritis and cervicitis; and sexually transmitted diseases related to infection by *Chlamydia trachomatis*, *Haemophilus ducreyi*, *Treponema pallidum*, *Ureaplasma urealyticum*, or *Neisseria gonorrhoeae*; toxin diseases related to infection by *S. aureus* (food poisoning and Toxic shock syndrome), or Groups A, B, and C streptococci; ulcers related to infection by *Helicobacter pylori*; systemic febrile syndromes related to infection by *Borrelia recurrentis*; Lyme disease related to infection by *Borrelia burgdorferi*; conjunctivitis, keratitis, and dacrocystitis related to infection by *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *S. aureus*, *S. pneumoniae*, *S. pyogenes*, *H. influenzae*, or *Listeria* spp.; disseminated *Mycobacterium avium* complex (MAC) disease related to infection by *Mycobacterium avium*, or *Mycobacterium intracellulare*; gastroenteritis related to infection by *Campylobacter jejuni*; intestinal protozoa related to infection by *Cryptosporidium* spp.; odontogenic infection related to infection by *viridans streptococci*; persistent cough related to infection by *Bordetella pertussis*; gas gangrene related to infection by *Clostridium perfringens* or *Bacteroides* spp.; and atherosclerosis related to infection by *Helicobacter pylori* or *Chlamydia pneumoniae*. Bacterial infections and protozoa infections and disorders related to such infections that may be treated or prevented in animals include the following: bovine respiratory diseases related to infection by *P. haem.*, *P. multocida*, *Mycoplasma bovis*, or *Bordetella* spp.; cow enteric disease related to infection by *E. coli* or protozoa (i.e., coccidia, cryptosporidia, etc.); dairy cow mastitis related to infection by *Staph. aureus*, *Strep. uberis*, *Strep. agalactiae*, *Strep. dysgalactiae*, *Klebsiella*

spp., *Corynebacterium*, or *Enterococcus* spp.; swine respiratory disease related to infection by *A. pleuro.*, *P. multocida*, or *Mycoplasma* spp.; swine enteric disease related to infection by *E. coli*, *Lawsonia intracellularis*, *Salmonella*, or *Serpulina hyodyisinteriae*; cow footrot related to infection by *Fusobacterium* spp.; cow metritis related to infection by *E. coli*; cow hairy warts related to infection by *Fusobacterium necrophorum* or *Bacteroides nodosus*; cow pink-eye related to infection by *Moraxella bovis*; cow premature abortion related to infection by protozoa (i.e. *neosporium*); urinary tract infection in dogs and cats related to infection by *E. coli*; skin and soft tissue infections in dogs and cats related to infection by *Staph. epidermidis*, *Staph. intermedius*, coagulase neg. *Staph.* or *P. multocida*; and dental or mouth infections in dogs and cats related to infection by *Alcaligenes* spp., *Bacteroides* spp., *Clostridium* spp., *Enterobacter* spp., *Eubacterium*, *Peptostreptococcus*, *Porphyromonas*, or *Prevotella*.

Antimicrobial Activity

Susceptibility tests can be used to quantitatively measure the *in vitro* activity of an antimicrobial agent against a given bacterial isolate. Compounds were tested for *in vitro* antibacterial activity by a micro-dilution method. Minimal Inhibitory Concentration (MIC) was determined in 96 well microtiter plates utilizing the appropriate Mueller Hinton Broth medium for the observed bacterial isolates. Antimicrobial agents were serially diluted (2-fold) in DMSO to produce a concentration range, from about 64 µg/ml to about 0.03 µg/ml. The diluted compounds (2 µl/well) were then spotted to sterile 96-well microtiter plates. The inoculum for each bacterial strain was adjusted to 5.5×10^5 CFU/ml in appropriate MIC medium; 200 µl/well of this inoculum was added to the 96-well microtiter plate resulting in a final concentration of 1×10^5 CFU/ml. The 96 well plates were covered and incubated in a humidified atmosphere at 35 ± 2 °C for 16-24 h depending on the bacterial strain tested. Following incubation, plate wells were visually examined by Optical Density measurement for the presence of growth (turbidity). The lowest concentration of an antimicrobial agent at which no visible growth occurs was defined as the MIC. The compounds of the invention generally demonstrated an MIC in the range from about 64 µg/ml to about 0.03 µg/ml.

All *in vitro* testing follows the guidelines described in the Approved Standards M7-A5 protocol, published by the National Committee for Clinical Laboratory Standards (NCCLS).

The present invention encompasses certain compounds, dosage forms, and methods of administering the compounds to a human or other animal subject. Specific compounds and compositions to be used in the invention must, accordingly, be pharmaceutically acceptable. As used herein, such a "pharmaceutically acceptable" component is one that is suitable for use with humans and / or animals without undue adverse side effects (such as toxicity, irritation, and allergic response) commensurate with a reasonable benefit/risk ratio.

The pharmaceutical compositions are prepared according to conventional procedures used by persons skilled in the art to make stable and effective compositions. In the solid, liquid, parenteral and topical dosage forms, an effective amount of the active compound or the active ingredient is any amount, which produces the desired results.

For the purpose of this invention the pharmaceutical compositions may contain the active compounds of the invention, their derivatives, salts and hydrates thereof, in a form to be administered alone, but generally in a form to be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. Suitable carriers which can be used are, for example, diluents or excipients such as fillers, extenders, binders, emollients, wetting agents, disintegrates, surface active agents and lubricants which are usually employed to prepare such drugs depending on the type of dosage form.

Any suitable route of administration may be employed for providing the patient with an effective dosage of the compound of the invention their derivatives, salts and hydrates thereof. For example, oral, rectal, parenteral (subcutaneous, intramuscular, intravenous), transdermal, topical, ophthalmic, otic and like forms of administration may be employed. Dosage forms include (solutions, suspensions, etc) tablets, pills, powders, troches, dispersions, suspensions, emulsions, solutions, capsules, injectable preparations, patches, ointments, creams, lotions, shampoos and the like.

The prophylactic or therapeutic dose of the compounds of the invention, their derivatives, salts or hydrates thereof, in the acute or chronic management of disease will vary with the severity of condition to be treated, and the route of administration. In addition, the dose, and perhaps the dose frequency, will also vary according to the age, body weight and response of the individual patient. In general, the total daily dose range, for the compounds of the invention, the derivatives, salts or hydrates thereof, for the conditions described herein, is

from about 200 mg to about 1500 mg, in single or divided doses. While intramuscular administration may be a single dose or up to 3 divided doses, intravenous administration can include a continuous drip. It may be necessary to use dosages outside these ranges in some cases as will be apparent to those skilled in the art. The term "an amount sufficient to eradicate such infections but insufficient to cause undue side effects" is encompassed by the above-described dosage amount and dose frequency schedule.

Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, or tablets, or aerosol sprays, each containing a predetermined amount of the active ingredient, as a powder or granules, or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy, but all methods include the step of bringing into association the active ingredient with the carrier, which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation.

The compositions of the present invention include compositions such as suspensions, solutions, elixirs, aerosols, and solid dosage forms.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are employed. Examples of suitable carriers include excipients such as lactose, white sugar, sodium chloride, glucose solution, urea, starch, calcium carbonate, kaolin, crystalline cellulose and silicic acid, binders such as water, ethanol, propanol, simple syrup, glucose, starch solution, gelatin solution, carboxymethyl cellulose, shellac, methyl cellulose, potassium phosphate and polyvinyl pyrrolidone, disintegrants such as dried starch, sodium alginate, agar powder, laminaria powder, sodium hydrogen carbonate, calcium carbonate, Tween (fatty acid ester of polyoxyethylenesorbitan), sodium lauryl sulfate, stearic acid monoglyceride, starch, and lactose, disintegration inhibitors such as white sugar, stearic acid glyceryl ester, cacao butter and hydrogenated oils, absorption promoters such as quaternary ammonium bases and sodium lauryl sulfate, humectants such as glycerol and starch, absorbents such as starch,

lactose, kaolin, bentonite and colloidal silicic acid, and lubricants such as purified talc, stearic acid salts, boric acid powder, polyethylene glycol and solid polyethylene glycol.

The tablet, if desired, can be coated, and made into sugar-coated tablets, gelatin-coated tablets, enteric-coated tablets, film-coated tablets, or tablets comprising two or more layers.

If desired, tablets may be coated by standard aqueous or non-aqueous techniques. In molding the pharmaceutical composition into pills, a wide variety of conventional carriers known in the art can be used. Examples of suitable carriers are excipients such as glucose, lactose, starch, cacao butter, hardened vegetable oils, kaolin and talc, binders such as gum arabic powder, tragacanth powder, gelatin, and ethanol, and disintegrants such as laminaria and agar.

In molding the pharmaceutical composition into a suppository form, a wide variety of carriers known in the art can be used. Examples of suitable carriers include polyethylene glycol, cacao butter, higher alcohols, gelatin, and semi-synthetic glycerides.

In one embodiment method for treating or preventing microbial infections in a human or an animal, comprising administering to said animal or human a therapeutically effective amount of a compound of Formula I or the pharmaceutical composition of compound of Formula I is provided. The microbial infections may be caused by Gram-positive, Gram-negative bacteria, aerobic, anaerobic bacteria, atypical bacteria or protozoa.

Abbreviations

Abbreviations which may be used in the descriptions of the schemes and the examples that follow are: Ac for acetyl; AIBN for azobis-isobutyronitrile; Bn for benzyl; Boc for t-butoxycarbonyl; Bu₃SnH for tributyltin hydride; Bz for benzoyl; CDI for carbonyldiimidazole; DBU for 1,8-diazabicyclo [5.4.0]undec-7-ene; DBN for 1,5-diazabicyclo[4.3.0]non-5-ene; DCC for 1,3-dicyclohexylcarbodiimide; DEAD for diethylazodicarboxylate; DIC for 1,3-diisopropylcarbodiimide; DMAP for dimethylaminopyridine; DMF for dimethyl formamide; DPPA for diphenylphosphoryl azide; EtOAc for ethyl acetate; KHMDS for potassium bis (trimethylsilyl)amide; LHMDS for Lithium bis (trimethylsilyl)amide; LDA for lithium diisopropyl amide; MeOH for methanol; Me₂S for dimethyl sulfide; MOM for methoxymethyl; NaN(TMS)₂ for sodium bis(trimethylsilyl)amide; NCS for N-chlorosuccinimide; NMO for 4-methylmorpholine N-


oxide; PCC for pyridinium chlorochromate; PDC for pyridinium dichromate; Ph for phenyl; TEA for triethylamine; THF for tetrahydrofuran; TPP or PPh_3 for triphenylphosphine; TBS for tert-butyl dimethylsilyl; TMS for trimethylsilyl.

In a further embodiment is provided a process for the preparation of macrolide and ketolide compounds having Formula I, wherein the variables have the previously defined meanings, the method comprising the process will be better understood in connection with the following synthetic Schemes. The synthesis of the compounds of the invention can be broadly summarized as

- a. Synthesis of the novel macrolides bearing 11,12- γ -lactone
- b. Synthesis of the novel ketolides
- c. Further conversions to generate the compounds of Formula I.

a) Synthesis of the novel macrolides bearing 11,12- γ -lactone

As depicted in the Scheme 1, erythromycin A or clarithromycin or derivatives of erythromycin and clarithromycin are used as the starting material for the reactions. All the above starting materials are used with optionally protected hydroxyl and / or amino groups with a suitable protecting group familiar to those skilled in the art. Exemplary protecting groups are, but not limited to silyl ethers such as triethylsilyl, trimethylsilyl, tert-butyldimethylsilyl, triisopropylsilyl. The protecting groups include benzyl, allyl, acetyl, benzoyl, pivaloyl and the like. For the amino protection, benzyloxycarbonyl, acetyl, tert-butoxycarbonyl (henceforth BOC) and the like may be used. The suitable macrolide starting material, depicted by compound (II) in the Scheme 1, is treated with triphosgene in a suitable solvent such as halogenated solvents such as dichloromethane, chloroform or THF in the presence of pyridine to provide the 11,12-carbonate (III, wherein R1, R2, R2' have the same meaning defined in Formula I).

**Scheme 1**


The compound III obtained from Scheme 1, is dissolved in solvent such as ethyl acetate, acetonitrile, tetrahydrofuran or mixtures thereof and treated with a base such as 1,8-diazabicyclo [5.4.0]undec-7-ene, KHMDS, LDA, triethylamine at 50-80°C, to give compound IV.

The compound IV is treated with cyanoacetic acid in presence of a ester coupling reagent such as DCC, 2,4,6-trichlorobenzoyl chloride, EEDQ, EDCI, Boc-Cl in a presence of a base such as pyridine, DMAP, N, N-Diisopropylethylamine, triethylamine and the like in a suitable solvent such as dichloromethane, acetonitrile, tetrahydrofuran, N,N-dimethylformamide or mixtures thereof to give compound Va. The compound Va is further

treated with a base such as pyridine, DBU, DBN, LHMDS, LDA to give compound of Formula VI.

Alternatively, the compound IV, is dissolved in solvent such as dichloromethane or acetonitrile or tetrahydrofuran or N,N-dimethylformamide or mixtures thereof and treated with chloroacetic anhydride in presence of a base such as pyridine and dimethylamino pyridine at 5-40 °C, to give compound Vb. The compound Vb, is dissolved in a solvent such as acetonitrile or tetrahydrofuran or N,N-dimethylformamide or DMSO or mixture thereof and treated with potassium cyanide at 15-40 °C, to give compound VI.

The compound VI is dissolved in solvent such as tetrahydrofuran or methanol or ethanol or mixture thereof and treated with hydroxylamine hydrochloride in the presence of base such as sodium bicarbonate or sodium acetate or potassium hydroxide to give compound VII (Scheme 2).




Scheme 2

b) Synthesis of the ketolide derivatives:

For the synthesis of the C-3 ketolide, the cladinose sugar is removed by hydrolysis according to procedures described in the art.

The 3-O-cladinose sugar of the compound of Formula I-a is cleaved with an inorganic acid such as hydrochloric acid, sulphuric acid at temperature ranging from -10 °C to 60 °C for 0.5 to 24 h. to give compound VIII as shown in Scheme 3. Before derivatizing the generated C-3-hydroxyl, the 2'-OH group of the 5-desosamine is protected as shown in Scheme 3 by use of an acetyl group. The acetylation is done using acetyl chloride or acetic anhydride in presence or absence of a base in a suitable solvent such as acetone, acetonitrile or dichloromethane, at room temperature for 6-24 h to give compound (IX, R₂ = COCH₃), Scheme 3. An alternate protecting group may also be used by treatment the compound VIII with reagents such as trialkylsilyl chloride, benzoyl chloride, benzoic anhydride, benzyl chloroformate, hexamethyldisilazane.

**Scheme 3**

The conversion of the 3-hydroxy group to 3-ketone is accomplished by using a Corey-Kim oxidation with N-chlorosuccinimide-dimethyl sulphide (NCS-DMS) or a Moffat oxidation with carbodiimide-dimethylsulphoxide (DMSO) complex in the presence of pyridinium trifluoroacetate or Dess-Martin periodinane. Such name reactions are carried out according to general procedures described in the art. In general, the compound (IX, $R_2=COCH_3$ or $R_2=TES$), obtained in Scheme 3 is dissolved in a chlorinated solvent such as dichloromethane or chloroform and Dess-Martin periodinane reagent was added and stirred at an ambient temperature for about 0.5 to 1 h to get the corresponding 3-ketone compound X, Scheme 3.


c) Further manipulations to generate the compounds of Formula I.

The manner in which the following procedures are to be used to provide the actual compounds listed in the table will be readily apparent to those persons skilled in the art of synthesis. The sequence of the reactions can be altered as will be readily apparent to those persons skilled in the art of synthesis. The examples described in the Experimental section later in this specification are illustrative and representative of how these procedures are to be used for all of the compounds listed in the table.

For the O-acylation of the amidoxime in the 11,12- γ -lactone ring i.e. the formation of compound XIIa or XIIb, the unsubstituted amidoxime is esterified with a suitable carboxylic acid reagent of the Formula R₉-COOH, wherein R₉ represents -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-B-R₅, -(CH₂)_m-C≡C-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, wherein R₅, R₆, X, B and m are as defined above. A mixture of suitably substituted acid, a condensing agent such as EDC, EDCI, DCC or EEDQ, HBTU and 1-hydroxy-benzotriazole in a halogenated solvent such as chloroform or dichloromethane, acetonitrile, THF, DMF was stirred at a temperature between -10° C to 45° C. Compound of Formula VIIa or VIIb was treated with the above mixture in the presence of a such as catalyst N,N-dimethylamino pyridine, at the temperature in the range of 0 to 45° C to give compound of Formula XIa and XIb respectively (Scheme 4).

The macrolide compounds of Formula XIIa are obtained from the corresponding macrolides of Formula XIa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane or chloroform or acetonitrile or DMF, at a temperature in the range of 0° C to 50° C.

For the ketolide compounds of Formula XIIb, the corresponding 3-hydroxyl compounds of Formula XIb are treated with an oxidizing agent as described above for the ketolides in Scheme 3, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XIIb, wherein all the variables are as defined above in Formula I.



wherein R₉ is -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-CC-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-B-R₅, -(CH₂)_m-CC-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-CC-X-R₆,

Scheme 4


The unsubstituted amidoxime was treated with a suitable isocyanate to obtain carbamates of Formula XVa or XVb as shown in Scheme 5.

Compound of Formula VIIa or VIIb was treated with isocyanate of Formula R₉-N=C=O, wherein R₉ represents -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-B-R₅, -(CH₂)_m-C≡C-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-

$C\equiv C-X-R_6$, wherein R_5 , R_6 , X , B and m are as defined above, in a solvent such as chloroform or dichloromethane, at a temperature between 0 to 40° C, to afford the corresponding compound of Formula XIVa and XIVb respectively.

The macrolide compounds of Formula XVa are obtained from the corresponding macrolides of Formula XIVa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane or chloroform or acetonitrile or DMF, at a temperature in the range of 0° C to 50° C.

For the ketolide compounds of Formula XVb, the corresponding 3-hydroxyl compounds of Formula XIVb are treated with a oxidizing agent as described above for the ketolides in Scheme 3, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XIVb, wherein all the variables are as defined above in Formula I.






wherein R_9 is $-(\text{CH}_2)_m-\text{R}_5$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{R}_5$, $-(\text{CH}_2)_m-\text{CC}-\text{R}_5$, $-(\text{CH}_2)_m-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{CC}-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{X}-\text{R}_6$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{X}-\text{R}_6$, $-(\text{CH}_2)_m-\text{CC}-\text{X}-\text{R}_6$, wherein R_5 , R_6 , X , B and m are as defined above

Scheme 5

For the O-alkylation of the amidoxime in the 11,12-gamma-lactone ring i.e. the formation of macrolide or ketolide of Formula XVIIa or XVIIb as shown in Scheme 6, the compound of Formula VIIa or VIIb was treated with a suitable alkylating reagent of Formula halo- R_9 , wherein R_9 represents $-(\text{CH}_2)_m-\text{R}_5$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{R}_5$, $-(\text{CH}_2)_m-\text{C}\equiv\text{C}-\text{R}_5$, $-(\text{CH}_2)_m-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{C}\equiv\text{C}-\text{B}-\text{R}_5$, $-(\text{CH}_2)_m-\text{X}-\text{R}_6$, $-(\text{CH}_2)_m-\text{CH}=\text{CH}-\text{X}-\text{R}_6$, $-(\text{CH}_2)_m-$

$\text{C}\equiv\text{C-X-R}_6$, wherein B, X, R_5 , R_6 and m are as defined above and halo is a chlorine, bromine or iodine. The compound VIIa or VIIb was dissolved in a solvent such as benzene or toluene or xylene or dialkyl ether or DMF or THF or mixture thereof, at a temperature in the range of 0-100 °C, in the presence of a base such as sodium hydride, potassium hydride or potassium carbonate or potassium hydroxide, optionally in the presence of phase transfer catalyst such as 18-crown-6-ether or trialkyl ammonium sulfate, to provide the compound XVIa and XVIb. The macrolide compounds of Formula XVIIa are obtained from the corresponding macrolides of Formula XVIa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane or chloroform or acetonitrile or DMF, at a temperature in the range of 0° C to 50° C.

For the ketolide compounds of Formula XVIIb, the corresponding 3-hydroxyl compounds of Formula XVIb are treated with a oxidizing agent as described above for the ketolides in Scheme 3, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XVIIb, wherein all the variables are as defined above in Formula I.




wherein R₉ is -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-CC-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-B-R₅, -(CH₂)_m-CC-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-CC-X-R₆

Scheme 6

Compounds of Formula XXa or XXb are prepared as shown in Scheme 7 by using Heck coupling reaction. Compound of Formula XVIIIA or XVIIIB, are treated a halo-aryl reagent of Formula halo-R₅ or halo-X-R₆ in presence of a palladium (II) or palladium (0) catalyst such as palladium acetate (Pd(OAc)₂) in presence of a phosphine reagent such as triphenylphosphine, tri(*o*-tolyl)phosphine and a base such as triethylamine, sodium carbonate, sodium bicarbonate, or in presence of sodium acetate, in a suitable solvent such as N,N-dimethylformamide (DMF), acetonitrile, tetrahydrofuran (THF) to give the corresponding



compound of Formula XIXa or XIXb respectively as shown in Scheme 7, wherein G is $-\text{CH}_2\text{m}-$, $-\text{A}-(\text{CH}_2)\text{m}-$; U is $-\text{R}_5$ or $-\text{X}-\text{R}_6$,

The macrolide compounds of Formula XXa are obtained from the corresponding macrolides of Formula XIXa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane, chloroform, acetonitrile, DMF or mixtures thereof, at a temperature in the range of 0° C to 50° C.



XVIIIa, Y' is H, Y is Ocladinose, R2,
R2' is TES
XVIIIb, Y' is H, Y is OH, R2, is TES

XIXa, Y' is H, Y is Ocladinose, R2,
R2' is TES
XIXb, Y' is H, Y is OH, R2, is TES




Scheme 7

For the ketolide compounds of Formula XXb, the corresponding 3-hydroxyl compounds of Formula XIXb are treated with an oxidizing agent as described above for the ketolides in Scheme 3, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XXb, wherein all the variables are as defined above in Formula I.

Compounds of Formula XXIIIa or XXIIIb are prepared as shown in Scheme 8 by using Sonagashira coupling reaction. Compound of Formula XXIa or XXIb are treated a halo-aryl reagent of Formula halo-R₅ or halo-X-R₆, wherein R₅, R₆ are as defined above, in presence of a palladium catalyst such as dichlorobis(triphenylphosphine)palladium (II) or Pd(PPh₃)₄ in presence of a base such as triethylamine, diethyl amine, butylamine, piperidine, in a suitable solvent such as N,N-dimethylformamide (DMF), acetonitrile, tetrahydrofuran (THF) to give the compound of Formula XXIIa or XXIIb respectively as shown in Scheme 8.


The macrolide compounds of Formula XXIIIa are obtained from the corresponding macrolides of Formula XXIIa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane or chloroform or acetonitrile or DMF, at a temperature in the range of 0° C to 50° C.

For the ketolide compounds of Formula XXIIIb, the corresponding 3-hydroxyl compounds of Formula XXIIb are treated with an oxidizing agent as described above for the ketolides in Scheme 3, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XXIIIb, wherein all the variables are as defined above in Formula I.



Scheme 8

The macrolide or ketolide of Formula XXIVa or XXIVb, wherein T is -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -A-(CH₂)_m-CH=CH-R₅, -A-(CH₂)_m-C≡C-R₅, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, -A-(CH₂)_m-CH=CH-X-R₆, -A-(CH₂)_m-C≡C-X-R₆, are treated with a hydrogenating catalyst such as 10% Pd/C, 5% Pd/C or 20% Pd(OH)₂ in presence of a hydrogen source such as hydrogen gas, ammonium formate, to give the corresponding compounds saturated compounds.



XXIVa, Y' is H, Y is Ocladinose,
XXIVb, Y', Y together C=O


Compound of Formula VIa or VIb is treated with a suitable reagent such as azidotrimethylsilane, sodiumazide in presence of catalyst such as tributyltinoxide to afford the corresponding tetrazol compound of Formula XXVa or XXVb. The tetrazol may be further alkylated using suitable alkylating reagent of Formula R₃-halo to afford the corresponding alkylated tetrazole derivatives, wherein R₃ is as defined above.

Compound of Formula VIa or VIb is treated with a suitable diamine such as ethylenediamine or propylenediamine in presence of sulphur at a temperature in the range of 100-150 °C to afford the corresponding dihydro imidazolyl compound of Formula XXVa or XXVb.

Compound of Formula VIIa or VIIb is treated with N,N-dimethylformamide dimethylacetal or trifluoroacetic anhydride, optionally in presence of a base such as triethylamine to afford the corresponding [1,2,4]-oxadiazole derivatives of Formula XXVa or XXVb.

The macrolide compounds of Formula XXVIa are obtained from the corresponding macrolides of Formula XXVa by treating with a fluoride source such as tetrabutylammonium fluoride or HF-pyridine complex, preferably HF-pyridine complex in a suitable solvent such as dichloromethane or chloroform or acetonitrile or DMF, at a temperature in the range of 0° C to 50° C.

For the ketolide compounds of Formula XXVIb, the corresponding 3-hydroxyl compounds of Formula XXVb are treated with an oxidizing agent as described above for the ketolides in Scheme 9, followed by deprotection of the 2'-hydroxyl of the desosamine sugar with a fluoride source as described above to afford ketolides of Formula XXVIb, wherein all the variables are as defined above in Formula I.



Scheme 9

Biological activity

The compounds of this invention are useful antimicrobial agents, effective against various human and veterinary pathogens, including multiple-resistant staphylococci and streptococci, enterococci, as well as anaerobic organisms such as bacteroides and clostridia species, and acid resistant organisms such as *Mycobacterium tuberculosis* and *Mycobacterium avium*.

Test Example 1

MIC Test Method : Overnight grown cultures of *S. aureus* organisms in Tryptic Soya broth were diluted in Mueller Hinton Broth to give optical density matching with MacFarland tube 0.5 standard. Cultures were further diluted 1:10 in Mueller Hinton broth. Using Denley's multipoint inoculator, 10^4 cells were deposited on Mueller Hinton agar (Difco) containing range of 2 fold dilutions of test compounds. These plates were incubated for 24 hrs at 35°C and MIC results recorded. MIC is defined as minimum drug concentration that inhibits test organisms. For determining MIC of test compounds against *Streptococcus pneumoniae*, Mueller Hinton agar containing 5% sheep blood was employed. The following strains were used to screen the compounds of invention. The compounds inhibited the growth of these bacteria with MIC's in the range of about 0.03 $\mu\text{g}/\text{mL}$ to about 64 $\mu\text{g}/\text{mL}$.

Strain
<i>Staphylococcus aureus</i> ATCC 25923
<i>Staphylococcus aureus</i> 014
<i>Staphylococcus epidermidis</i> 110
<i>Staphylococcus haemolyticus</i> ATCC 25923
<i>Enterococcus faecalis</i> 401
<i>Enterococcus faecium</i> 303
<i>Streptococcus pneumoniae</i> 49619
<i>Streptococcus pneumoniae</i> 706
<i>Streptococcus pyogenes</i> 801
<i>Streptococcus pyogenes</i> 805
<i>Haemophilus influenzae</i> 49247
<i>Mycobacterium tuberculosis</i>
<i>Mycobacterium avium</i>

The compounds and processes of the present invention will be better understood in connection with the following examples, which are intended as an illustration only and not limiting of the scope of the invention. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art and such changes and modifications including, without limitation, those relating to the chemical structures, substituents, derivatives, Formulations and/or methods of the invention may be made.

EXAMPLES

The following examples describe in detail the chemical synthesis of some of the representative compounds of the present invention. The procedures are illustrations, and the invention should not be construed as being limited by chemical reactions and conditions they express. No attempt has been made to optimize the yields obtained in these reactions, and it would be obvious to one skilled in the art that variations in reaction times, temperatures, solvents, and/or reagents could increase the yields.

Example 1: Compound III of Scheme 1, R₁ is CH₃, R₂,R₂ is triethylsilyl

Step A: To a stirred solution of 6-methyl erythromycin A (15 g, 20.1 mmol) in anhydrous acetonitrile (120 ml) cooled to 0° C. Triethyl amine (19.5 ml, 140 mmol), DMAP (12.25 g, 100 mmol) and chloro triethyl silane (20 ml, 120 mmol) were added under nitrogen atmosphere. The resulting mixture was stirred at 25-30° C temperature for 3 hr. Then water (120 ml) was added, the solid precipitated was filtered under suction. The wet cake was stirred with methanol (50 ml) at 25-30° C temperature for 1 hr and filtered under suction, dried to provide step-1 product 2',4"-Di-TES-6-O-methyl-erythromycin A as white solid in 80% (16 g) yield. MS: m/z: 976.8 (M+1).

Step B: To a solution of product obtained in above step (10 g, 10.2 mmol) in anhydrous dichloromethane (70 ml) was added pyridine (5 ml, 61.2 mmol) and followed by triphosgene (3.04 g, 10.2 mmol) at 0 °C under an inert atmosphere. The reaction mixture was further stirred at 0-5 °C for 0.5 h. To the reaction mixture water (15 ml) was added, the organic layer was separated. The organic layer was dried (Na₂SO₄) and evaporated under reduced pressure to give the title compound (9.7 g) as white solid. m/z ([MH]⁺) = 1002.5.

Example 2: Compound IV of Scheme 1, R₁ is CH₃, R₂,R₂ is triethylsilyl

To a solution of Example 1 (9.5 g, 9.5 mmol) in ethyl acetate (70 ml), 1,8-diazabicyclo[5.4.0]undec-7-ene (3 ml, 20 mmol) was added at an ambient temperature and

the reaction mixture was further heated under reflux for 6h. The reaction mixture was cooled to room temperature and water (15 ml) was added to it and stirred for 10 min. The organic layer was separated, dried and evaporated under reduced pressure to afford the title compound (9.3 g). $m/z ([MH]^+)$ = 958.

Example 3: Compound Vb of Scheme 1, R₁ is CH₃, R₂,R₂ is triethylsilyl

To the solution of Example 2 (9.3 g, 9.7 mmol) in anhydrous dichloromethane (30 ml) was added pyridine (4.7 ml, 58.2 mmol), DMAP (0.6 g, 4.8 mmol) and followed by chloroacetic anhydride (3.7 g, 21.3 mmol) at 0 °C. After completion of addition it was allowed to come to room temperature and stirred further for 3h. To the reaction mixture water (30 ml) was added and stirred vigorously for 10 min. and separated the organic layer. The organic layer was then washed with a saturated solution of sodium bicarbonate followed by the brine. The organic layer was dried (Na₂SO₄) and upon removal of the solvent under reduced pressure a gummy mass was obtained which was taken into the methanol and stirred for 2h to provide the title compound (8 g). $m/z ([MH]^+)$ = 1034.

Example 4: Compound VI of Scheme 1, R₁ is CH₃, R₂,R₂ is triethylsilyl

Method 1:

Example 3 (6 g, 5.8 mmol) was dissolved into N,N-dimethylformamide (20 ml) and to it potassium cyanide (0.6 g, 8.7 mmol) was added carefully under inert atmosphere at room temperature. The reaction mixture was further stirred for 2h. The reaction mixture was filtered and poured into the 5% sodium bicarbonate solution (30 ml), the cream color solid separated out was filtered and washed successively by solution of ferrous sulphate and water. The crude solid obtained was crystallized from dichloromethane and n-hexane to provide the title compound (5.15 g). $m/z ([MH]^+)$ = 1025.

Method 2:

Step A: Compound Va of Scheme 1, R₁ is CH₃, R₂,R₂ is triethylsilyl: To the stirred solution of cyanoacetic acid (1.11 g, 13.1 mmol) in anhydrous dichloromethane (50 ml), 2, 4, 6-trichlorobenzoyl chloride was added at 0° C. N, N-Diisopropylethylamine (2.05 ml, 13.1 mmol) was added to the reaction mixture and it was stirred at 0° C for 30 min. Example 2 (5 gm, 5.3 mmol) and N, N-diisopropylethylamine (2.05 ml, 13.1 mmol) were added to the reaction mixture and it was stirred at 25-30° C temperature for 24 hr. To the reaction mixture was quenched by addition of aqueous solution of saturated sodium bicarbonate (50 ml) and stirred vigorously for 10 min. Organic layer was separated. The organic layer was washed with

a water (50 ml), dried over Na_2SO_4 and solvent was evaporated under vacuum to afford yellow foam. The foam was stirred with acetonitrile at 0-5° C and filtered to give step-4 product as a pale yellow solid in 70% (3.7 gm) yield. MS: m/z: 1025.7 (M+1).

Step B: To the stirred solution of product obtained in above step (3.6 g, 3.51 mmol) in N,N-dimethylformamide (15 ml), 1,8-diazabicyclo[5.4.0]undec-7-ene (0.545 g, 3.51 mmol) added at 25-30° C temperature and stirred for 1 hr. Reaction mixture was diluted with water (50 ml) and the solid was filtered under suction to afford title compound in 40% (1.5 g) yield. MS: m/z: 1025.7 (M+1).

Example 5: Compound VII of Scheme 2, R_1 is CH_3 , R_2,R_2' is triethylsilyl, Y' is H, Y is O-cladinose

To a solution of Example 4 (5 g, 4.8 mmol) in methanol (25 ml), hydroxylamine hydrochloride (3.7 g, 53.7 mmol) and sodium bicarbonate (4.6 g, 54.6 mmol) was added and heated under reflux for 15h. The solid separated out was filtered and washed with water. The crude solid obtained was purified by recrystallization from methanol to afford the title compound as white solid (4.3 g). m/z ([MH]⁺) = 1058.

Example 6: Compound IX of Scheme 3, R_1 is CH_3 , R_2 is triethylsilyl, Q is CN

To a solution of Example 4 (10 g, 9.7 mmol) in acetonitrile (40 ml), a 2N HCl (10 ml) was slowly added at 0 °C. The reaction mixture was further stirred at room temperature for 4h. It was then cooled to 0 °C and the pH of the solution was brought to 8 by slow addition of a saturated solution of sodium bicarbonate. The aqueous layer was extracted into chloroform (3 x 25 ml). The combined organic layer was dried (Na_2SO_4) and evaporated under reduced pressure to furnish the crude solid. The solid obtained was purified by column chromatography over silica gel using chloroform:methanol (7:3) as an eluent to provide a white solid (6 g). In this reaction condition the hydroxyl protecting group of desosamine sugar was cleaved, therefore we have again protected the hydroxyl group by using the following procedure. The deprotected solid (6 g, 9.4 mmol) was taken into acetonitrile: N,N-dimethylformamide (8:2, 30 ml). To this solution triethylamine (5.2 ml, 37.6 mmol), DMAP (4.5 g, 37.6 mmol) followed by chlorotriethylsilane (6.3 ml, 37.6 mmol) was added at room temperature and stirred for 3 h. To the reaction mixture water (50 ml) was added and it was extracted into n-hexane (3 x 40 ml). The combined organic layer was dried (Na_2SO_4) and evaporated to give the gummy mass, which on trituration with n-pentane provided the title compound as a white solid (6 g). m/z ([MH]⁺) = 753.

Example 7: Compound VII of Scheme 2, R₁ is CH₃, R₂ is triethylsilyl, Y is OH, Y' is H,
 Using the Example 6 (0.5 g) and following the same procedure as described for the Example 5, the title compound obtained a white solid (0.5 g). m/z ([MH]⁺) = 786.

Example 8: Compound X of Scheme 3, R₁ is CH₃, R₂ is triethylsilyl, Q is CN

The Example 6 (6 g, 7.9 mmol) was dissolved in dichloromethane (60 ml). To the reaction mixture a solution Dess-Martin periodinane (15% in DCM, 27 ml, 9.5 mmol) was added under an inert atmosphere. The reaction mixture was stirred at room temperature for 5 min. and a solution of Na₂S₂O₃ (10% in saturated aqueous solution of sodium bicarbonate, 60 ml) was added and stirred for 10 min. The organic layer was separated and the aqueous layer was extracted with dichloromethane (3 x 15ml) and the combine organic layer was dried (Na₂SO₄). The removal of the solvent under reduced pressure provided the crude solid, which was purified by column chromatography over silica gel using n-hexane:EtOAc (8:2) as an eluent. The title compound was obtained a white solid (3.5 g). m/z ([MH]⁺) = 751.

Example 9: Compound VII of Scheme 2, R₁ is CH₃, R₂ is triethylsilyl, Y, Y' together C=O;

Using the Example 8 (3.5 g, 4.6 mmol) and following the same procedure as described for the Example 5, the title compound obtained a white solid (3.5 g). m/z ([MH]⁺) = 784.

Example-10 : Compound of Formula I-c, R₁ is CH₃, T is H

To the solution of Example 5 (0.950 g, 0.89 mmol) in anhydrous THF (25 ml) was added 70% hydrogen fluoride in pyridine (77 μ l, 2.7 mmol). The reaction mixture was stirred at an ambient temperature under inert atmosphere for 14 h. To the reaction mixture water (15 ml) was added and extracted with ethyl acetate (3 x 15 ml). The combined organic layer was dried (Na₂SO₄) and evaporated under reduced pressure. The thick liquid on trituration with *n*-pentane furnished the title compound (0.582 g) as a white solid. MS 830 (M + H)⁺.

Example-11: Compound of Formula I-b where Het is 1H-tetrazol-5-yl;

Step A: To the stirring solution of Example 4 (29.3 mmol) in toluene was added tributyltinoxide (0.25 eq) and azidotrimethylsilane(2 eq). The reaction mixture was heated at 90 °C for 16-20 hours. The reaction mixture was concentrated and purified using silica gel column chromatography(Ethyl acetate: Hexane 1:1). Yield 70 %.

Step C: To the stirring solution of compound obtained in Step B (0.36 mmol) in tetrahydrofuran was added HF-Pyridine complex (3.5 eq) and stirred it for 16 hours. The reaction mixture was concentrated and water was added. The product was extracted in chloroform. Yield 80 %. Yield: Mass :m/z: 840.1 ($M+H$)⁺, M. P. 162-164 ° C.

Example-12: Compound of Formula I-b where Het is 2-(1-methoxy-ethoxymethyl)-1H-tetrazol-5-yl:


Example-13: Compound of Formula I-b where Het is 2-(2-methoxy-ethoxymethyl)-2H-tetrazol-5-yl;

Step A: To the stirring solution of Example 4 (29.3 mmol) in toluene was added tributyltin oxide (0.25 eq) and azidotrimethylsilane (2 eq). The reaction mixture was heated at 90 °C for 16-20 hours. The reaction mixture was concentrated and purified using silica gel column chromatography (Ethyl acetate: Hexane 1:1). Yield 70 %.

Step B: To the stirring solution of the compound obtained in above step A (0.93 mmol) in acetone was added TEA (1.3 eq) and MEM-Cl (1.2 eq) and heated it at reflux temperature for 1 hours. The reaction mixture was concentrated and water was added. Filtered the solid residue and dried it over high vacuo. It was a mixture of two regio isomers. It was then separated using 100-200-mesh silica gel column chromatography. Yield 42 %.

Step C: To the stirring solution of compound obtained in Step B (0.36 mmol) in tetrahydrofuran was added HF-Pyridine complex (3.5 eq) and stirred it for 16 hours. The reaction mixture was concentrated and water was added. The product was extracted in chloroform. Yield 80 %. Yield: Isomer-A: Mass : m/z: 928 (M+H). M.P. 210-212 °C, Isomer-B: 190 mg Mass :m/z: 928 (M+H)⁺, M. P. 150-152 ° C.

Following examples were prepared by using above procedure and by using reactant A.




Example	Het	Reactant (A)	Mp (° C)	Mass (M+1)
14		Tetrazol-2-yl-acetic acid ethyl ester	150-154	926.11
15		2-Tetrazol-2-yl-acetamide	198-200	897.1
16		Tetrazol-2-yl-acetonitrile	218-220	879.1
17		Tetrazol-1-yl-acetonitrile	202-204	879.1
18		2-Allyl-2H-tetrazole	188-190	880.1
19		Tetrazole-2-carboxylic acid (3-fluoro-phenyl)-methyl amide	230-232	1005.1
20		2-Tetrazol-2-yl-ethanol	220-222	884.1

Example-21: Compound of Formula I-c, T is CO(CH₂)₂-(pyridin-3-yl), R₁ is CH₃

To the solution of 3-pyridin-3-yl-propionic acid (1.42 g, 7.5 mmol) in dichloromethane (40 ml) was added dicyclohexyl carbodiimide (1.55 g, 7.5 mmol) and stirred for 0.5 h at room temperature. To this reaction mixture Example 5 (2 g, 1.9 mmol) followed by dimethyl amino pyridine (0.131 g, 1.0 mmol) was added and stirred further for 2 h. The solid precipitated out was filtered and washed with dichloromethane (10 ml). The combined filtrate was evaporated under reduced pressure to furnish a thick liquid, which was used as it is without purification for further reaction.

To the solution of thick liquid (1.8 g, 1.5 mmol) obtained as above in THF (10 ml) was added 70% hydrogen fluoride in pyridine (130 µl, 4.53 mmol) under inert atmosphere and stirred for 15 h. The residue obtained after evaporation of solvent under reduced pressure was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried (Na₂SO₄) and evaporated under reduced pressure to provide a white solid as title compound (1.3 g, 82%). MS 963 (M + H)⁺.

Using the above procedure following examples were synthesized by using the corresponding carboxylic acid.



Example	T	Reactant (A)	MP (° C)	Mass (M+1)
22	COCH ₂ O-(2-chlorophenyl)	(2-Chloro-phenoxy)-acetic acid	150-152	998
23	CO(CH ₂) ₂ -(4-dimethyl amino)phenyl	3-(4-dimethylamino-phenyl)-propionic acid	168-170	996
24	CO(CH ₂) ₂ -(3-fluoro)phenyl	3-(3-fluorophenyl)-propionic acid	170-172	980
25	CO(CH ₂) ₂ -(4-(4-methyl-piperazine)phenyl)	3-[4-(4-methyl-piperazin-1-yl)-phenyl]-propionic acid	168-170	1060
26	CO(CH ₂) ₂ -(4-(4-acetyl-[1,2,3]triazol-1-yl))-phenyl	3-[4-(4-acetyl-[1,2,3]triazol-1-yl)-phenyl]-propionic acid	174-176	1071
27	CO(CH ₂) ₂ -(4-phenyl-piperazinyl)	3-(4-phenyl-piperazin-1-yl)-propionic acid	165-167	1045
28	CO(CH ₂) ₂ -(4-(2-methoxy-phenyl)-piperazine-1-yl)	3-[4-(2-methoxy-phenyl)-piperazin-1-yl]-propionic acid	155-156	1076
29	COCH ₂ -S-(3-methoxy phenyl)	(3-methoxy-phenylsulfanyl)-acetic acid	158-160	1010
30	CO(CH ₂) ₂ CONH-(3-methoxy-phenyl)	N-(3-methoxy-phenyl)-succinamic acid	170-172	1035
31	CO(CH ₂) ₂ CONH-(2,4-	N-(2,4-difluoro-phenyl)-	190-192	1040

	difluorophenyl)	succinamic acid		
32	CO(CH ₂) ₂ CO-(4-phenyl)-piperazin-1-yl	4-oxo-4-(4-phenyl-piperazin-1-yl)-butyric acid	162-163	1075
33	COCH=CH-(pyridin-4-yl)	3-pyridin-4-yl-acrylic acid	210-212	961.1
34	COCH=CH-(pyridin-3-yl)	3-pyridin-3-yl-acrylic acid	157-159	961.1
35	COCH=CH-(pyridin-2-yl)	3-pyridin-2-yl-acrylic acid	107-110	961.1
36	COCH=CH-(1-(2-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-acrylic acid	172-175	1057.2
37	COCH=CH-(1-(3-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(3-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-acrylic acid	148-151	1057.2
38	COCH=CH-(1-(4-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(4-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-acrylic acid	198-200	1045.1
39	COCH=CH-(1-(3-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(3-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-acrylic acid	188-190	1045.1
40	COCH=CH-(1-(4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl)	(E)-3-[1-(4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-acrylic acid	203-202	1057.2
41	CO(CH ₂) ₂ -phenyl	3-phenyl-propionic acid	140-142	962.1
42	CO(CH ₂) ₂ -(pyridin-2-yl)	3-pyridin-2-yl-propionic acid	116-118	963.1
43	CO(CH ₂) ₂ -(1-(4-methoxyphenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	182-184	1059.2
44	CO(CH ₂) ₂ -(1-(4-fluoro phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(4-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	202-205	1047.2
45	CO(CH ₂) ₂ -(pyridin-4-yl)	3-pyridin-4-yl-propionic acid	166-168	963.1
46	CO(CH ₂) ₂ -(3-methoxy)phenyl	3-(3-methoxy-phenyl)-propionic acid	172-174	992.1
47	CO(CH ₂) ₂ -(4-methoxy)phenyl	3-(4-methoxy-phenyl)-propionic acid	164-167	992.1
48	CO(CH ₂) ₂ -(1-(3-methyl phenyl)-1H-[1,2,3]triazol-4-yl)	3-(1- <i>m</i> -tolyl-1H-[1,2,3]triazol-4-yl)-propionic acid	172-174	1043.2
49	CO(CH ₂) ₂ -(1-(3-chloro phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(3-chloro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	176-178	1063.6
50	CO(CH ₂) ₂ -(3-(4-fluoro-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(4-fluoro-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	148-154	1048.1

51	CO(CH ₂) ₂ -(3-(3-bromo-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(3-bromo-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	158-160	1109.1
52	CO(CH ₂) ₂ -(3-(4-nitro-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(4-nitro-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	198-201	1075.2
53	CO(CH ₂) ₂ -(4-(pyridin-2-yl)-piperazinyl)	3-(4-pyridin-2-yl-piperazin-1-yl)-propionic acid	138-140	1047.2 81
54	CO(CH ₂) ₂ -(3-cyano)phenyl	3-(3-Cyano-phenyl)-propionic acid	170-172	987.1
55	COCH ₂ O-(3-chloro)phenyl	3-(3-Chloro-phenyl)-propionic acid	158-161	996.6
56	CO(CH ₂) ₂ -(3,5-dimethoxy)phenyl	3-(3,5-Dimethoxy-phenyl)-propionic acid	118-120	1022.2
57	CO(CH ₂) ₂ -(2,3-dimethoxy)phenyl	3-(2,3-Dimethoxy-phenyl)-propionic acid	152-154	1022.2
58	CO(CH ₂) ₂ -(3-(3-fluoro-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(3-Fluoro-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	166-170	1048.1
59	CO(CH ₂) ₂ -(3-(3-chloro-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(3-Chloro-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	158-160	1064.6
60	CO(CH ₂) ₂ -(3-(3-methoxy-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(3-Methoxy-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	148-150	1060.2
61	CO(CH ₂) ₂ -(3-(3,5-dimethoxy-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(3,5-Dimethoxy-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	160-163	1090.2
62	CO(CH ₂) ₂ -(3-(4-chloro-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(4-Chloro-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	144-146	1064.6
63	CO(CH ₂) ₂ -(3-(4-bromo-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(4-Bromo-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	152-154	1109.1
64	CO(CH ₂) ₂ -(3-(pyridin-4-yl)-[1,2,4]oxadiazol-5-yl)	3-(3-Pyridin-4-yl-[1,2,4]oxadiazol-5-yl)-propionic acid	156	1031.1
65	CO(CH ₂) ₂ -(4-(2-methylphenyl)-piperazinyl)	3-(4-o-tolyl-piperazin-1-yl)-propionic acid	170	1060.3
67	CO(CH ₂) ₂ -(4-(3-methoxyphenyl)-piperazinyl)	3-[4-(3-Methoxy-phenyl)-piperazin-1-yl]-propionic acid	146	1076.3
68	CO(CH ₂) ₂ -(4-(2-fluorophenyl)-piperazinyl)	3-[4-(2-Fluoro-phenyl)-piperazin-1-yl]-propionic acid	152	1064.2
69	CO(CH ₂) ₂ -(4-(pyrimidin-2-yl)-piperazinyl)	3-(4-pyrimidin-2-yl-piperazin-1-yl)-propionic acid	170	1048.2

70	CO(CH ₂) ₂ -(1-(2-fluorophenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	162	1047.2
71	CO(CH ₂) ₂ -(4-trifluoromethyl)phenyl	3-(4-trifluoromethyl-phenyl)-propionic acid	168	1030.1
72	CO(CH ₂) ₂ -(3-(2-fluoro-5-methoxy-phenyl)-[1,2,4]oxadiazol-5-yl)	3-[3-(2-fluoro-5-methoxy-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid	158	1078.2
73	CO(CH ₂) ₂ -(3-phenyl-4H-imidazol-1-yl)	3-(4-phenyl-4H-imidazol-1-yl)-propionic acid	172-174	1028.2
74	COCH ₂ NHCONH-phenyl	(3-phenyl-ureido)-acetic acid	164-168	1006.1
75	CO(CH ₂) ₂ -(4-([1,2,3]triazol-1-yl)phenyl)	3-(4-[1,2,3]triazol-1-yl-phenyl)-propionic acid	180-183	1029.2
76	CO(CH ₂) ₂ -(4-cyano)phenyl	3-(4-Cyano-phenyl)-propionic acid	168-170	987.1
77	CO(CH ₂) ₂ -(2,5-dimethoxy)phenyl	3-(2,5-dimethoxy-phenyl)-propionic acid	166-169	1022.2
78	CO(CH ₂) ₂ -(benzo[1,3]dioxol-5-yl)	3-benzo[1,3]dioxol-5-yl-propionic acid	138-140	1006.1
79	CO(CH ₂) ₂ -(3,5-dimethoxy)phenyl	3-(3,5-dimethoxy-phenyl)-propionic acid	170-174	1022.1
80	CO(CH ₂) ₂ -(1-(2,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	158-161	1065.2
81	CO(CH ₂) ₂ -(1-(3,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(3,4-difluorophenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	172-175	1065.2
82	CO(CH ₂) ₂ -(1-(2,3,4-trifluorophenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2,3,4-trifluorophenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	108-110	1083.1
83	CO(CH ₂) ₂ -(1-(2-fluoro-4-methyl-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2-fluoro-4-methyl-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	138-141	1061.2
84	CO(CH ₂) ₂ -(1-(2-fluoro-4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(2-fluoro-4-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	168.000	1077.2
85	CO(CH ₂) ₂ -(1-(2,3-difluoro-4-ethoxy-phenyl)-1H-[1,2,3]triazol-4-yl)	3-[1-(4-Ethoxy-2,3-difluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	136-139	1109.2
86	CO(CH ₂) ₂ -(4-(4-cyano-[1,2,4]triazol-1-yl)phenyl)	3-[4-(4-Cyano-[1,2,4]triazol-1-yl)-phenyl]-propionic acid	144-148	1054.2
87	CO(CH ₂) ₂ CONH-(3-methoxy-phenyl)	N-(3-Methoxy-phenyl)-succinamic acid	170-173	1035.2
88	CO(CH ₂) ₂ -(4-(imidazol-1-yl)phenyl)	3-(4-Imidazol-1-yl-phenyl)-propionic acid	168-171	1028.2
89	CO(CH ₂) ₂ -(4-([1,2,4]triazol-1-yl)phenyl)	3-(4-[1,2,4]triazol-1-yl-phenyl)-propionic acid	185-187	1029.2
90	CO(CH ₂) ₂ CONH-(2-methoxy-phenyl)	N-(2-Methoxy-phenyl)-succinamic acid	186-188	1035.2


91	CO(CH ₂) ₂ CONH-(2,4-difluorophenyl)	N-(2,4-Difluoro-phenyl)-succinamic acid	190-192	1041.1
92	CO(CH ₂) ₂ CO-(4-(4-fluoro-phenyl)piperazin-1-yl)	4-[4-(2-Fluoro-phenyl)-piperazin-1-yl]-4-oxo-butyric acid	160-164	1092.2
93	COCH ₂ NHCONH-phenyl	(3-Phenyl-ureido)-acetic acid	164-168	1006.1
94	COCH ₂ -(benzo[1,3]dioxol-5-yl)	benzo[1,3]dioxol-5-yl-acetic acid	170-174	992.1
95	COCH ₂ NHCONH-(4-fluorophenyl)	[3-(4-fluoro-phenyl)-ureido]-acetic acid	172-175	1024.1
96	COCH ₂ NHCONH-(3-chloro-4-methylphenyl)	[3-(3-chloro-4-methyl-phenyl)-ureido]-acetic acid	176-178	1054.6
97	COCH ₂ O-phenyl	phenoxy-acetic acid	160-164	964.1
98	COCH ₂ S-phenyl	phenylsulfanyl-acetic acid	158-160	980.2

Example-99: Compound of Formula I-c where T is CO(CH₂)₂-(1-(3,5-difluoro-phenyl)-1H-[1,2,3]triazol-4-yl), R₁ is CH₃:

To a solution of 3-[1-(3,5-difluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid (0.288 g, 1.13 mmol) in dichloromethane (50 ml) was added dicyclohexylcarbodiimide (0.486 g, 2.35 mmol) and 1-hydroxy-benzotriazole (0.040 g, 0.296 mmol) at an ambient temperature. The reaction mixture was stirred further for 1 h. To this reaction mixture then added Example 5 (1.0 g, 0.946 mmol) and followed by dimethyl amino pyridine (0.033 g, 0.270 mmol) at an ambient temperature. The reaction mixture was further stirred for 1 h. The reaction mixture was filtered and the filtered solid was washed with dichloromethane (10 ml). The combined filtrate was evaporated under reduced pressure to provide a thick liquid (1.1 g), which was used as it is without purification for further reaction.

To a solution of above thick liquid (1.1 g, 0.876 mmol) in THF (20 ml) was added 70% hydrogen fluoride in pyridine (72 μ l, 2.52 mmol) at room temperature under inert atmosphere. The resulting solution was further for 15 h. The solvent evaporated under reduced pressure. The residue was dissolved into water (30 ml) and washed with diethyl ether (3 x 25 ml). The aqueous layer upon saturation with sodium chloride a white solid was precipitated out, it was filtered and dried. The white solid obtained was triturated with diethyl ether to give the title compound as a white solid (0.400 g).

Using the above procedure following examples were synthesized by using the corresponding carboxylic acid.



Example	T	Carboxylic acid	MP (° C)	Mass M+1
100	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(2\text{-methoxy-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$	3-[1-(2-Methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	176-178	1059
101	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3\text{-fluoro-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$	3-[1-(3-fluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	186-188	1047
102	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,4\text{-difluoro-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$	3-[1-(3,4-difluoro-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	168-170	868
103	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,5\text{-dimethoxy-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$	3-[1-(3,5-dimethoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	180-182	1089
104	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3\text{-methoxy-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$	3-[1-(3-methoxy-phenyl)-1H-[1,2,3]triazol-4-yl]-propionic acid	186-187	1032
105	$\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(pyridin-3-yl)-1\text{H-[1,2,3]triazol-4-yl})$	3-(1-pyridin-3-yl-1H-[1,2,3]triazol-4-yl)-propionic acid	170-172	1030


Example-106: Compound of Formula I-c where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(2\text{-fluoro-5\text{-methoxy-phenyl})-[1,2,4]oxadiazol-5-yl})$, R_1 is CH_3 ;

To a solution of 3-[3-(2-fluoro-5-methoxy-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic acid (0.413 g, 1.89 mmol) in dichloromethane (50 ml) was added dicyclohexylcarbodiimide (0.486 g, 2.35 mmol) and 1-hydroxy-benzotriazole (0.040 g, 0.296 mmol) at an ambient temperature. The reaction mixture was stirred further for 1 h. To this reaction mixture then

added Example 5 (1.0 g, 0.946 mmol) and followed by dimethyl amino pyridine (0.033 g, 0.270 mmol) at an ambient temperature. The reaction mixture was further stirred for 1 h. The reaction mixture was filtered and the filtered solid was washed with dichloromethane (10 ml). The combined filtrate was evaporated under reduced pressure to provide a thick liquid (1.1 g), which was used as it is without purification for further reaction.

To a solution of above thick liquid (1.1 g, 0.876 mmol) in THF (20 ml) was added 70 % hydrogen fluoride in pyridine (65 μ l, 2.27 mmol) at room temperature under inert atmosphere. The resulting solution was further for 15 h. The solvent evaporated under reduced pressure. The residue was dissolved into water (30 ml) and washed with diethyl ether (3 x 25 ml). The aqueous layer upon saturation with sodium chloride a white solid was precipitated out, it was filtered and dried. The white solid obtained was triturated with diethyl ether to give the title compound as a white solid (0.325 g).

Using the above procedure following examples were synthesized by using the corresponding carboxylic acid.



Example	T	Carboxylic acid	MP (° C)	Mass M+1
107	CO(CH ₂) ₂ -(3-phenyl-[1,2,4]oxadiazol-5-yl)	3-(3-Phenyl-[1,2,4]oxadiazol-5-yl)-propionic acid	162-166	1030
108	CO(CH ₂) ₂ -(3-naphthalen-2-yl-[1,2,4]oxadiazol-5-yl)	3-(3-Naphthalen-2-yl-[1,2,4]oxadiazol-5-yl)-propionic acid	138-142	1080
109	CO(CH ₂) ₂ -(3-(pyridin-3-yl)-[1,2,4]oxadiazol-5-yl)	3-(3-Pyridin-3-yl-[1,2,4]oxadiazol-5-yl)-propionic acid	149-150	1031
110	CO(CH ₂) ₂ -[3-(4-methoxyphenyl)-[1,2,4]oxadiazol-5-yl]-	3-[3-(4-Methoxy-phenyl)-[1,2,4]oxadiazol-5-yl]-propionic	132-134-	1060

	[1,2,4]oxadiazol-5-yl]	acid		
--	------------------------	------	--	--

Example-111: Compound of Formula I-c where T is CH_2 -(5-methyl-[1,2,4]oxadiazol-3-yl), R_1 is CH_3 :

Step A: To a stirred solution of Example 5 (1.0 g, 0.95 mmol) in 1,4-dioxan (10 ml) was added N,N-dimethylformamide dimethylacetal (2 ml) and heated at 70 °C for 2 h. After completion of reaction solvent was evaporated under vacuum, water was added to the crude reaction mass and extracted with ethyl acetate (50 ml x 3). Combined organic layer was dried over anhydrous sodium sulphate and then evaporated yielding crude solid. It was then purified by column chromatography using (Ethyl acetate:n-hexane; 20:80) as a mobile phase. Pale solid compound was isolated Yield: 550 mg, (55%).

Step B: The compound obtained in Step A (550 mg, 0.50 mmol) was dissolved in acetonitrile (10 ml). 70% HF-Pyridine solution (44 μ l, 3.0 equi.) was added to the above solution and stirred at room temperature for overnight under N_2 atmosphere. After completion of reaction solvent was evaporated under vacuum to obtain crude solid. Water (20 ml) was added to the crude product and then extracted with dichloromethane (30 ml x 3). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude white solid. It was then purified by column chromatography using $\text{CHCl}_3:\text{MeOH}$; 80:20) as a mobile phase to obtain title compound as white solid (Yield : 190 mg, 44%), $\text{MS } 854 (\text{M} + \text{H})^+$ $\text{Mp: } 213\text{--}214^\circ \text{C}$


Example-112: Compound of Formula I-c where T is CH_2CONH -(3-chloro)phenyl, R_1 is CH_3 :

Step-1: To the stirred solution of Example 5 (1.00 g, 0.94 mmol) in benzene (20 ml) was added sodium hydride (47 mg, 1.13 mmol, 60 % suspension in mineral oil) at 25-30°C temperature and stirred for 30 min. 2-Bromo-N-(3-chloro-phenyl)-acetamide (0.25 g, 1.04 mmol) was added to reaction mixture and stirred for 1 hour at an 25-30°C temperature. The reaction mixture was then poured in aqueous saturated ammonium chloride solution (30 ml). Organic layer was dried over Na_2SO_4 . It was purified using silica gel column chromatography (10 % Acetone: Hexane) to provide step-1 product as solid in 75% (0.87 g) yield. $\text{MS: m/z: } 1229 (\text{M}+1)$

Step-2: A mixture of step-1 product and (0.80 g, 0.65 mmol) 70% HF in pyridine (55 μ l, 1.95 mmol) in THF (15 ml) was stirred under inert atmosphere for 15 h at 25-30°C temperature. The solvent was evaporated under vacuum to provide a residue. The residue was dissolved in

ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried over Na_2SO_4 and evaporated under vacuum to provide crude solid which on trituration in diethyl ether afforded pale yellow solid as the title compound Example in 65% (0.42 g) yield. Mp: 128-131° C, Mass: M/z: 997.1 (M+1).

Following examples were prepared by using above procedure and by using reactant A.




Ex	T	Reactant (A)	MP (°C)	Mass (M+1)
113	CH_2CONH_2	2-bromo-acetamide	207-210	887.1
114	$\text{CH}_2\text{CONH-}(2\text{-methoxy})\text{phenyl}$	2-Bromo-N-(2-methoxy-phenyl)-acetamide	188-190	993.1
115	$\text{CH}_2\text{CONH-}(3\text{-methoxy})\text{phenyl}$	2-Bromo-N-(3-methoxy-phenyl)-acetamide	190-194	993.1
116	$\text{CH}_2\text{CONH-}(2\text{-fluoro})\text{phenyl}$	2-Bromo-N-(2-fluoro-phenyl)-acetamide	184-187	981.1
117	$\text{CH}_2\text{CONH-}(3\text{-fluoro})\text{phenyl}$	2-Bromo-N-(3-fluoro-phenyl)-acetamide	182-185	981.1
118	$\text{CH}_2\text{CONH-}(4\text{-fluoro})\text{phenyl}$	2-Bromo-N-(4-fluoro-phenyl)-acetamide	190-194	981.1
119	$\text{CH}_2\text{CONH-}(2,4\text{-difluoro})\text{phenyl}$	2-Bromo-N-(2,4-difluoro-phenyl)-acetamide	178-182	999.1
120	$\text{CH}_2\text{CONH-}(3,4\text{-difluoro})\text{phenyl}$	2-Bromo-N-(3,4-difluoro-phenyl)-acetamide	196-199	999.1
121	$\text{CH}_2\text{CONH-}(2,3,4\text{-trifluoro})\text{phenyl}$	2-Bromo-N-(2,3,4-trifluoro-phenyl)-acetamide	190-194	1017.1
122	$\text{CH}_2\text{CONH-}(3,5\text{-dimethoxy})\text{phenyl}$	2-Bromo-N-(3,5-dimethoxy-phenyl)acetamide	184-187	1023.1
123	$\text{CH}_2\text{CONH-cyclopropyl}$	2-Bromo-N-cyclopropyl-acetamide	180-183	927.1
124	$\text{CH}_2\text{CON}(\text{CH}_3)\text{-}(3\text{-fluoro})\text{phenyl}$	2-Bromo-N-(3-fluoro-phenyl)-N-methyl-acetamide	174-178	995.1
125	$\text{CH}_2\text{CON}(\text{CH}_3)\text{-}(4\text{-chloro})\text{phenyl}$	2-Bromo-N-(4-chloro-phenyl)-N-methyl-acetamide	198-201	1011.1
126	$\text{CH}_2\text{CONH-}(3\text{-}(pyrazol-1-yl)\text{-pyridin-5-yl})$	2-Bromo-N-(6-pyrazol-1-yl-pyridin-3-yl)-acetamide	172-175	1030.1

127	CH ₂ -(1-(4-fluorophenyl)-1H-[1,2,3]-triazol-5-yl)	4-Bromomethyl-1-(4-fluorophenyl)-1H-[1,2,3]-triazole	158-160	1008.1
128	CH ₂ -(1-(3-fluorophenyl)-1H-[1,2,3]-triazol-5-yl)	4-bromomethyl-1-(3-fluorophenyl)-1H-[1,2,3]-triazole	210-212	1008.1
129	CH ₂ -(1-(3,5-difluoro phenyl)-1H-[1,2,3]-triazol-5-yl)	4-bromomethyl-1-(3,5-difluoro-phenyl)-1H-[1,2,3]-triazole	191-193	1026.1
130	CH ₂ -(1-(3,4-difluoro phenyl)-1H-[1,2,3]-triazol-5-yl)	4-bromomethyl-1-(3,4-difluoro-phenyl)-1H-[1,2,3]-triazole	190-192	1026.1
131	CH ₂ -(1-(3,4,5-trifluorophenyl)-1H-[1,2,3]-triazol-5-yl)	4-bromomethyl-1-(2,3,4-trifluoro-phenyl)-1H-[1,2,3]-triazole	180-182	1044.1
132	CH ₂ -(1-(3-chlorophenyl)-1H-[1,2,3]-triazol-5-yl)	4-bromomethyl-1-(3-chlorophenyl)-1H-[1,2,3]-triazole	191-193	1023.1
133	CH ₂ -(5-(pyridin-2-yl)-[1,2,4]-oxadiazol-3-yl)	2-(3-bromomethyl-[1,2,4]-oxadiazole-5-yl)-pyridine	149-151	990.1

Example-134: Compound of Formula I-c where T is CH₂-phenyl, R₁ is CH₃;

To the stirring solution of Example 5 (0.56 mmol) in benzene(10 ml) was added NaH (0.62 mmol) and stirred it for 30 min. To this stirring solution, benzyl bromide (0.68 mmol) was added and stirred the reaction mixture for 4 hours at 60 °C. The reaction mixture was allowed to come at room temperature and then poured in water (20 ml). Separated organic layer and dried over Na₂SO₄. Purified it with silica gel column chromatography (20 % Ethyl acetate:Hexane). Yield 54 %. The above product (0.31 mmole) was taken in tetrahydrofuran (5 ml) and added HF-Pyridine complex (1.24 mmole). Stirred at room temperature for overnight. The reaction mixture was concentrated and water was added. Extracted the product in dichloromethane (2x10 ml). Dried it (Na₂SO₄) and concentrated to give title compound in yield 44 %. MS 920 (M+H).

Using the above procedure following examples were synthesized by using the corresponding Reagent A.



Example	T	Reagent A	M.P. (° C)	Mass M+1
135	CH ₂ C(CH ₃)=CH ₂	3-bromo-2-methyl-propene	184-186	884
136	CH ₂ -(4-nitro)phenyl	4-nitrobenzyl bromide	160-164	965
137	CH ₂ CH ₂ -(4-(pyridin-2-yl)piperazinyl)	1-(2-bromoethyl)-4-pyridin-2-yl-piperazine	130-132	1019
138	CH ₂ CH=CH-(2,3-dihydro-benzo[1,4]dioxin-6-yl)	6-bromo-1,4-benzo dioxane	176-178	1004
139	CH ₂ CH=C(CH ₃) ₂	1-bromo-3-methyl-but-2-ene	195-197	899.1
140	CH ₂ CH ₂ CH=CH ₂	4-bromo-but-1-ene	194-196	885.1
141	CH ₂ -(4-methyl-phenyl)	4-methyl-benzyl bromide	178-180	935.1
142	CH ₂ -(2-methyl phenyl)	2-methyl benzyl bromide	160-162	935.1
143	CH ₂ -(3-methoxy phenyl)	3-methoxy-benzyl bromide	168-170	951.1
144	CH ₂ -(2-fluorophenyl)	2-fluoro-benzyl bromide	172-174	939.1
145	CH ₂ -(4-bromophenyl)	4-bromo-benzyl bromide	168-170	1000.1
146	CH ₂ -(2,4-difluorophenyl)	2,4-difluoro-benzyl bromide	160-162	957.1
147	CH ₂ -(2,5-difluorophenyl)	2,5-difluoro-benzyl bromide	166-168	957.1
148	CH ₂ CO ₂ CH ₂ CH ₃	ethyl bromoacetate	190-192	917.1
149	CH ₂ CO-(4-phenyl-piperazin-1-yl)	2-bromo-1-(4-phenyl-piperazin-1-yl)-ethanone	179-181	1033.1

150	CH ₂ CO-(4-(2-fluoro phenyl)-piperazin-1-yl)	2-bromo-1-[4-(2-fluoro-phenyl)-piperazin-1-yl]-ethanone	194-196	1051.1
-----	---	---	---------	--------

Example 151: Compound of Formula I-c where T is CH₂CN, R₁ is CH₃;

Step A: To the stirring solution of the Example 5 (15.52 mmol) in acetone was added K₂CO₃ (1.2 eq), bromo acetonitrile (1.1 eq) and refluxed it for 48 hours. The reaction mixture was concentrated under vacuo. Water was added and filtered the residue. Dried it and purified it with column chromatography (EA;Hexane 20:80). Yield 77 %.

Step B: To the stirring solution of the compound obtained in step B (0.52 mmol) in dry acetonitrile was added 70 % HF in Pyridine complex (3.5 eq) The resulting mixture was stirred for 16 hours at room temperature under nitrogen atmosphere. The reaction mixture evaporated under vacuo. Water was added and extracted it in chloroform. Dried it over Na₂SO₄ and concentrated it. The residue was crystallized from ethanol: water (30:70) to give the title compound in yield 67 %. MS 869 (M + H)⁺. Mp: 172-174° C.

Example 152: Compound of Formula I-c where T is CH₂C(NH₂)=N-OH, R₁ is CH₃;

Step-1: To the solution of compound obtained in step A of above example (5.0 g, 4.55 mmol) in methanol (40 ml) was added hydroxylamine hydrochloride (3.25 g), and sodium bicarbonate (4.21g, 50.1mmol) stirred for 12 hr at room temperature. The reaction mass was concentrated under vacuum to obtain crude solid. The crude solid was stirred with water (100 ml). Solid separated was filtered under suction and dried under vacuum to provide step-1 product in 93% (4.8 g) yield. Mass: m/z: 1130 (M+1).

Step-2: A mixture of step-1 product (1.0 g, 0.9 mmol) and 70% hydrogen fluoride in pyridine (77 μ l, 2.6 mmol) in THF (15 ml) was stirred under inert atmosphere at a temperature between 25-30 °C for 15 h. The solvent was evaporated under vacuum and the residue obtained was dissolved in ethyl acetate (25 ml) and washed with water (2 x 10 ml). The combined organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to provide crude solid which on trituration in diethyl ether afforded pale yellow solid as the title compound in 60% (0.65 g) yield. Mp: 168-170° C. Mass: m/z: 903.1 (M+1).

Example-153: Compound of Formula I-c where T is CH₂CH=CH₂, R₁ is CH₃;

Step A: To the stirring solution of Example 5 (0.56 mmol) in Toluene (10 ml) was added Potassium *tert* butoxide (0.62 mmol) and stirred it for 20 min. To this stirring solution, allyl bromide (0.68 mmol) was added and stirred the reaction mixture for 2 hours at room temperature. The reaction mixture was then poured in water (20 ml). Separated organic layer and dried over Na_2SO_4 . Purified it with silica gel column chromatography (20 % Ethyl acetate:Hexane). Yield 74 %.


Step B: The product from step A (0.31 mmol) was taken in tetrahydrofuran (5 ml) and added HF-Pyridine complex (1.24 mmol). Stirred at room temperature for overnight. The reaction mixture was concentrated and water was added. Extracted the product in dichloromethane (2x10 ml). Dried it (Na_2SO_4) and concentrated to give title compound in yield 60 %. MS 867 (M+H). M.P 194-196 °C.

Example-154: Compound of Formula I-c where T is $\text{CH}_2\text{CH}=\text{CH}$ -phenyl, R_1 is CH_3 ;

Step-A: To the stirred solution of compound obtained in Step A of above example (1.5 g, 1.36 mmol) in DMF (15 ml) was added sodium acetate (0.24 ml, 2.73 mmol) at 25-30 °C temperature. To the above solution tetrakis triphenylphosphine palladium (0) (0.15 g, 0.13 mmol) and bromobenzene (0.31 g) were added and the reaction mixture was stirred at 120 °C temperature for 12 hours. The reaction mixture was quenched by pouring in aqueous saturated ammonium chloride solution (20 ml). The mixture was extracted with ethyl acetate (3 x 50 ml). Combined organic layer was dried over Na_2SO_4 to provide a residue. The residue was purified by using silica gel column chromatography (10 % Acetone: Hexane) to provide step-1 product as a solid in 65% (1.0 g) yield. MS: m/z: 1226 (M+1).

Step-B: A mixture of step-2 product (0.90 g, 0.73 mmol) and 70% hydrogen fluoride in pyridine (63 μl , 2.20 mmol) in THF (15 ml) was stirred under inert atmosphere for 15 h. The solvent was evaporated under vacuum to provide a residue. The residue was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried over Na_2SO_4 and evaporated under vacuum to provide crude solid which on trituration in diethyl ether afforded white solid as the title compound example in 55% (0.40 g) yield., Mass: m/z: 998.1 (M+1) Mp: 196-199 °C

Following examples were prepared by using above procedure and by using reactant A.



Example	U	Reactant A	MP °C	Mass M+1
155	(2-methoxy)phenyl	1-bromo-2-methoxy-benzene	120	976.2
156	(3-fluoro)phenyl	1-bromo-3-fluoro-benzene	182	964.1
157	(4-cyano)phenyl	4-bromo-benzonitrile	160	971.1
158	(3-hydroxymethyl)phenyl	(3-bromophenyl)-methanol	125	976.2
159	(4-trifluoromethyl)phenyl	1-bromo-4-trifluoromethyl-benzene	96	1014.1
160	(3-trifluoromethyl)phenyl	1-bromo-3-trifluoro methyl-benzene	176	1014.1
161	(3-acetoxy)phenyl	1-(4-bromophenyl)-ethanone	155	988.2

Example-162: Compound of Formula I-c where T is $\text{CH}_2-(3-(5\text{-fluoro-2-methoxyphenyl})\text{-isoxazol-5-yl})$, R_1 is CH_3 ;

Step-1: To the stirred solution of compound obtained in Step A of example 153 (1.5 g, 1.36 mmol) in toluene (15 ml) was added triethylamine (0.38 ml, 2.73 mmol) followed by 2-methoxy-5-fluoro imidoyl chloride (0.34 g, 2.05 mmol) at 25-30° C temperature. The reaction mixture was stirred at 90 °C temperature for 12 hours. It was diluted with ethyl acetate (50 ml) followed by addition of water (20 ml). Organic layer was dried over Na_2SO_4 , evaporated under vacuum to provide a residue. The residue was purified by using silica gel column chromatography (10 % Acetone: Hexane) to afford step-1 product as a solid in 70% (1.2 g) yield. MS 1264.1 (M+1).

Step-2: A mixture of step-1 product (1. g, 0.79 mmol) and 70% hydrogen fluoride in pyridine (68 μl , 2.37 mmol) in THF (15 ml) was stirred under inert atmosphere for 15 h. The solvent was evaporated under vacuum and the residue obtained was dissolved in ethyl acetate (25 ml) and washed with water (2 x 10ml). The organic layer was dried over Na_2SO_4 and evaporated

under vacuum to provide crude solid which upon trituration with diethyl ether afforded pale yellow solid as the title compound example in 60% (0.49 g) yield., Mass: M/z: 950.1 (M+1). Mp: 192-194° C

Example-163: Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{CH}$, R_1 is CH_3 :

Step A: To the stirring solution of Example 5 (0.56 mmol) in Toluene (10 ml) was added Potassium *tert* butoxide (0.62 mmol) and stirred it for 20 min. To this stirring solution, 3allyl bromide (0.68 mmol) was added and stirred the reaction mixture for 2 hours at room temperature. The reaction mixture was then poured in water (20 ml). Separated organic layer and dried over Na_2SO_4 . Purified it with silica gel column chromatography (20 % Ethyl acetate:Hexane). Yield 78 %.


Step B: The product from step A (0.31 mmole) was taken in tetrahydrofuran (5 ml) and added HF-Pyridine complex (1.24 mmole). Stirred at room temperature for overnight. The reaction mixture was concentrated and water was added. Extracted the product in dichloromethane (2x10 ml). Dried it (Na_2SO_4) and concentrated to give title compound in yield 60 %. MS 868 (M+H). M.P 168-170 °C.

Example-164: Compound of Formula I-c where T is $\text{CH}_2\text{C}\equiv\text{C}-\text{(3-cyano-phenyl)}$, R_1 is CH_3 :

Step-1: To the solution of compound obtained in step A of above example (1.4 g, 1.23 mmol) in DMF (40 ml) was added cesium carbonate (0.63 g, 3.2 mmol), bis-triphenylphosphine Palladium (II) chloride (0.10 g, 0.13 mmol), cuprous Iodide (0.05 g, 0.26 mmol) followed by 3-iodobenzonitrile (0.29 g, 1.27 mmol). The reaction mixture was stirred at 25-30° C temperature under nitrogen atmosphere for 12 hrs. it was quenched with addition of ice-cold water (15 ml) to provide a suspension. Solid was filtered & dried under vacuum. The solid was purified by using silica gel column chromatography (20 % ethyl acetate-Hexane) to provide step-1 product as a solid in 33% (0.5 g) yield. Mass: m/z: 1198 (M+1).

Step-2: A mixture of step-1 product (0.5 g, 0.41 mmol) and 70% hydrogen fluoride in pyridine (1.0 mmol) in THF (10 ml) was stirred under inert atmosphere at 25-30° C temperature for 15 hr. The solvent was evaporated under vacuum, obtained residue was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10 ml). The organic layer was dried over Na_2SO_4 and evaporated under vacuum to provide crude solid which on trituration in diethyl ether afforded off white solid as the title compound in 19% (0.075 g) yield., Mass: m/z: 970.1 (M+1). Mp: 158-159° C

Following examples were prepared by using above procedure and by using reactant A.



Example	T	Reactant A	MP °C	Mass (M+1)
165	CH ₂ C≡C-phenyl	Iodo-benzene	150-151	945.1
166	CH ₂ C≡C-(3-methyl)phenyl	3-methyl-iodo-benzene	194-195	959.1

Example-167: Compound of Formula I-c where T is CONH-phenyl, R₁ is CH₃;

Step-1: To the stirred solution of Example-5 (0.75 g, 0.70 mmol) in dichloromethane (40 ml) was added phenylisocyanate (0.12 ml) at 25-30° C temperature. The reaction mixture was stirred further for 2 h. The mixture was quenched by addition of water (20 ml) and extracted with dichloromethane (2 x 20 ml). Combined organic layer was dried over Na₂SO₄ and concentrated under vacuum to obtain crude solid. It was purified by using silica gel column chromatography (10 % Acetone:n-Hexane) to furnish step-1 product as a pale yellow solid in 55% (0.48 g) yield. Mass: m/z: 1226 (M+1).

Step-2: A mixture of step-1. product (0.47 g, 0.38 mmol) and 70% hydrogen fluoride in pyridine (32 µl , 1.14 mmol) in THF (10 ml) was stirred under inert atmosphere for 15 h. The solvent was evaporated and the residue obtained was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried over Na₂SO₄ and evaporated under vacuum to provide crude solid which on trituration in diethyl ether afforded title compound as a pale yellow solid in 60% (0.28 g) yield. Mp: 192- 194 °C, Mass: m/z: 950.1 (M+1).

Example-168: Compound of Formula I-c where T is CONH-(4-ethyl-phenyl), R₁ is CH₃;


Using above procedure and by using 4-ethyl-phenyl isocyanate in the place of phenyl isocyanate the title compound was prepared. Mp: 152-155 °C, Mass: m/z: 978.1.

Example-169: Compound of Formula I-c where T is $\text{CH}_2\text{CH}_2\text{CH}_3$, R_1 is CH_3 ;

Step-A: A stirred suspension of product obtained from the Step A of Example 153 (1 g, 0.9 mmol) and 10% palladium on carbon (100 mg) in tetrahydrofuran (10 ml) was stirred under atmospheric hydrogen pressure at 25-30° C temperature for 12 hours. The reaction mixture was filtered over celite bed, and filtrate was evaporated under vacuum to provide step-1 product as a white solid in 90% (0.9 g) yield. It was used as such for the next reaction. Mass: m/z: 1101.1 (M+1).

Step-B: A mixture of step-1 product (0.9 g, 0.82 mmol) and 70% hydrogen fluoride in pyridine (24 μl , 2.5 mmol) in THF (10 ml) was stirred under inert atmosphere for 15 h. The solvent was evaporated and the residue obtained was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried over Na_2SO_4 and evaporated under vacuum to provide crude solid which on trituration in diethyl ether afforded pale yellow solid title compound in 50% (70 g) yield. Mp: 182-184° C. Mass: m/z: 873.1 (M+1).

Following examples were prepared by using above procedure and by using the corresponding unsaturated compound



Example	T	Starting material	Mp (° C)	Mass (M+1)
170	$\text{CH}_2\text{CH}(\text{CH}_3)_2$	Example 135	210-212	887.1
171	$\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$	Example 139	184-186	901.1

Example-172: Compound of Formula I-e where Het is [1,2,4]-oxadiazol-3-yl;

Step A: To a stirred solution of Example 7 (1.0 g, 1.3 mmol) in (10 ml) 1,4-dioxan was added (2 ml) N,N dimethylformamide dimethylacetal and heated at 70 °C for 3 h. After completion of reaction solvent was evaporated under vacuum, water (30 ml) was added to the crude reaction mass and extracted with ethyl acetate (50 ml x 3). Combined organic layer was dried over anhydrous sodium sulphate and then evaporated yielding crude solid. It was then

purified by column chromatography using (Ethyl Acetate:n-hexane; 35:65) as a mobile phase. Pale solid was isolated. Yield : 450 mg, (45 %).

Step B: To a stirred solution of the compound obtained in step A (450 mg, 0.6 mmol) in dichloromethane (10 ml) was added Dess Martin Periodinane reagent (2.0 ml, 1.2 equi., 15 % solution) under N₂ atmosphere at room temperature for 30 min. After completion of reaction, additional 25 ml of dichloromethane was added and then quenched with 30 ml saturated aqueous solution of sodium thiosulphate: sodium bicarbonate. Organic layer was dried over anhydrous sodium sulphate and then distilled off under vacuum to obtain pale yellow solid. It was then purified by column chromatography using (Ethyl Acetate:n-Hexane; 30:70) as a mobile phase. Crystalline solid was isolated. Yield: 250 mg, (52 %).

Step C: The compound obtained in step B (250 mg, 0.30 mmol) was dissolved in acetonitrile (10 ml). 70% HF-Pyridine solution (18 μ l, 2.0 euqi.) was added to the above solution and stirred at room temperature for overnight under N₂ atmosphere. After completion of reaction solvent was evaporated under vacuum to obtain crude product. Water (20 ml) was added to the crude product and extracted with dichloromethane (25 ml x 3). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude white solid. It was then purified by column chromatography using CHCl₃:MeOH (90:10) as a mobile phase to obtain title compound as white solid compound. Yield : 120 mg, (56 %) MS: 681.1 (M + H)⁺. Yield : 0.185 mg, (55 %), M.P.: 208-210 °C.

Example-173: Compound of Formula I-e where Het is (5-trifluoromethyl)-[1,2,4]-oxadiazol-3yl;

Step A: To a stirring solution of Example 9 (1.0 gm, 1.3 mmol) in dichloromethane (25 ml) was added trifluoroacetic anhydride (0.365 ml, 2.6 mmol) and triethyl amine (0.550 ml, 3.9 mmol). The resulting reaction mixture was heated at 45 °C for 12 h water (25 ml) was added to and extracted with dichloromethane (2 x 20 ml). Combined organic layer was dried over Na₂SO₄ and evaporated under reduced pressure. Resulting crude residue was purified by using column chromatography (15 % Acetone-Hexane) to afford pale solid. Yield : 0.6 gm.

Step B: To a stirred solution of the compound obtained in step A (580 mg, 0.7 mmol) in dichloromethane (10 ml) was added Dess Martin Periodinane reagent (2.4 ml, 1.2 equi., 15 % solution) under N₂ atmosphere at room temperature for 30 min. After completion of reaction, additional 25 ml of dichloromethane was added and then quenched with 30 ml saturated aqueous solution of sodium thiosulphate: sodium bicarbonate. Organic layer was dried over anhydrous sodium sulphate and then distilled off under vacuum to obtain pale yellow solid. It

was then purified by column chromatography using (Ethyl Acetate:n-Hexane; 30:70) as a mobile phase. Crystalline solid was isolated. Yield: 300 mg, (52 %).

Step C: The compound obtained in step B (280 mg, 0.35 mmol) was dissolved in acetonitrile (10 ml). 70% HF-Pyridine solution (19 μ l, 2.0 euqi.) was added to the above solution and stirred at room temperature for overnight under N_2 atmosphere. After completion of reaction solvent was evaporated under vacuum to obtain crude product. Water (20 ml) was added to the crude product and extracted with dichloromethane (25 ml x 3). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude white solid. It was then purified by column chromatography using $CHCl_3$:MeOH; 90:10) as a mobile phase to obtain title compound as white solid compound. Yield : 180 mg, (66 %)., , Mass: 749.1 ($M + H$)⁺. M.P. 200-202 °C

Example-174: Compound of Formula I-e where Het is (4,5)-dihydro-1H-imidazol-2-yl;

Step A: To a mixture of Example 6 (1.0 g, 1.3 mmol) and ethylene diamine (3.0 ml) was added sulphur powder (21 mg, 0.66 mmol) and heated the resulting reaction mixture at 120 °C for 45 min. Reaction mixture was cooled to room temperature and ice cold water was added under stirring. Crude solid separated was filtered and dried. It was purified by using column chromatography (15 % Acetone-n-Hexane) affording (0.8 gm) pale yellow solid.

Step B: The intermediate from step A (0.75 g, 0.94 mmol) was dissolved in dichloromethane (60 ml). To this reaction mixture a solution Dess-Martin periodinane (15% in DCM, 3.2 ml, 1.13 mmol) was added under an inert atmosphere. The reaction mixture was stirred at room temperature for 15 min. and a solution of $Na_2S_2O_3$ (10% in saturated aqueous solution of sodium bicarbonate, 60 ml) was added and stirred for 10 min. The organic layer was separated and the aqueous layer was extracted with dichloromethane (3 x 15ml) and the combine organic layer was dried (Na_2SO_4). The removal of the solvent under reduced pressure provided the crude solid, which was purified by column chromatography over silica gel using n-hexane: EtOAc (8:2) as an eluent. The title compound was obtained a white solid (0.35 g). m/z ([MH]⁺) =794.

Step C: To the solution of intermediate from step B (0.30 g, 0.37 mmol) in anhydrous THF (25 ml) was added 70% hydrogen fluoride in pyridine (16 μ l, 0.56 mmol). The reaction mixture was stirred at an ambient temperature under inert atmosphere for 3 h. To the reaction mixture water (15 ml) was added and extracted with ethyl acetate (3 x 15 ml). The combined organic layer was dried (Na_2SO_4) and evaporated under reduced pressure. The thick liquid on

trituration with *n*-pentane furnished the title compound as a white solid. Yield : 0.18 g (70%), Mass: 681.1 (M + H)⁺, M.P.: 152-154 °C.

Example175: Compound of Formula I-e where Het is 1-(2-methoxyethoxy)-1H-tetrazol-5-yl;

Example176: Compound of Formula I-e where Het is 2-(2-methoxyethoxy)-2H-tetrazol-5-yl;

Step-1: To the stirring solution of the product obtained in Step B, of above Example 12 (2 gm, 1.73 mmol) in acetonitrile (10 ml), 2N HCl (10 ml) was added and stirred reaction mixture at room temperature for 4-5 hours. Reaction mixture was neutralized with saturated sodium bicarbonate (20 ml) solution and then extracted in chloroform (2x20 ml). The combined organic layer was washed with water, dried over Na₂SO₄, and then concentrated it under reduced pressure to get the intermediate II. Yield 1.4 gm.

Step-2: To the stirring solution of compound-II (1.4 gm, 1.82 mmol) in acetone (20 ml) acetic anhydride (0.56 ml, 5.6 mmol) and K₂CO₃ (0.400 gm, 2.9 mmol) was added. The reaction mixture was stirred for 24 hours at room temperature. The solvent was removed under vacuo and water was added. It was extracted in chloroform (2x25 ml). The combined organic layer was washed with water and dried over Na₂SO₄ and evaporated under reduced pressure to yield crude solid compound III. Yield: 1.35 gm

Step-3: To the stirring solution of compound-III (1.35gm, 1.66 mmol)) in dichloromethane(20 ml) was added into 15 % Dess-Martin periodinane reagent solution in DCM (5.6 ml,2 mmol) and stirred it at room temperature under nitrogen for 30 minutes. The reaction mixture was diluted with 20 ml (1:1) 10 % Sodium thiosulphate and sat. sodium bicarbonate solution. The organic layer was separated and washed with water and then dried it over Na₂SO₄.and evaporated under reduced pressure to obtain compound IV which was purified by column chromatography (15 % Acetone-Hexane). Yield: 1.3 gm.

Step-4: Compound-IV (1.3 gm, 1.60 mmol) was stirred in methanol (25 ml) for 4-5 hours at room temperature. It was concentrated. It was a mixture of two regio isomers. It was then separated by using preparative HPLC (Ammonium acetate: acetonitrile). Yield: Isomer-A: 160 mg Mass:m/z: 781.1 (M+H). M.P. 112-114 °C, Isomer-B: 190 mg, Mass:m/z: 781.1 (M+H), M. P. 122-123 °C.

Example 177: Compound of Formula I-e where Het is 1-(allyl)-1H-tetrazol-5-yl;

Example 178: Compound of Formula I-e where Het is 2-(allyl)-2H-tetrazol-5-yl;

Using the above procedure the title compounds were prepared Isomer A: Mass :m/z: 721.1 (M+H), M. P. 130-132 ° C. Isomer B: Mass :m/z: 721.1 (M+H), M. P. 170-172 ° C.

Example-179: Compound of Formula I-f where T is $\text{CH}_2\text{CH}=\text{CH}_2$:

Step A: To the stirring solution of the Example 7 (15.52 mmol) in acetone was added K_2CO_3 (1.2 eq), allyl bromide (1.1 eq) and refluxed it for 48 hours. The reaction mixture was concentrated under vacuo. Water was added and filtered the residue. Dried it and purified it with column chromatography (EA;Hexane 24:76).Yield 70 %.

Step B: To the stirring solution of the compound obtained in step A (16.6 mmol) in dry DCM was added Dess-Martin periodinane (15 % solution in DCM) (1.2 eq) under N_2 atmosphere for 30 min. The reaction mixture was diluted with 1:1 10% sodiumthiosulphate solution and saturated sodium bicarbonate solution. The organic layer was separated and washed with water and brine; it was dried over Na_2SO_4 and concentrated under vacuo. The residue was purified with column chromatography (Ethylacetate: Hexane 20:80) to afford title compound, yield 50%.

Step C: To the stirring solution of the compound obtained in step B (0.48 mmol) in dry acetonitrile was added 70% HF in Pyridine complex (3.5 eq) The resulting mixture was stirred for 16 hours at room temperature under nitrogen atmosphere. The reaction mixture evaporated under vacuo. Water was added and extracted it in chloroform. Dried it over Na_2SO_4 and concentrated it. The residue was crystallized from ethanol: water (30:70) to give the title compound in yield 70 %. MS 710 ($\text{M} + \text{H}$)⁺.

Example-180: Compound of Formula I-f where T is $\text{CH}_2\text{-(5-methyl[1,2,4]oxadiazol-3-yl)}$:

Step A: To a stirred solution of Example 7 (1.0 g, 1.3 mmol) in (10 ml) 1,4-dioxan was added (2 ml) N,N-dimethylformamide dimethylacetal and heated at 70 °C for 3 h. After completion of reaction solvent was evaporated under vacuum, water (30 ml) was added to the crude reaction mass and extracted with ethyl acetate (50 ml x 3). Combined organic layer was dried over anhydrous sodium sulphate and then evaporated yielding crude solid. It was then purified by column chromatography using (Ethyl Acetate:n-hexane; 35:65) as a mobile phase. Pale solid was isolated. Yield : 480 mg, (48 %).

Step B: To a stirred solution of the compound obtained in step A (480 mg, 0.6 mmol) in dichloromethane (10 ml) was added Dess Martin Periodinane reagent (2.0 ml, 1.2 equi., 15 % solution) under N_2 atmosphere at room temperature for 30 min. After completion of reaction, additional 25 ml of dichloromethane was added and then quenched with 30 ml saturated aqueous solution of sodium thiosulphate: sodium bicarbonate. Organic layer was dried over anhydrous sodium sulphate and then distilled off under vacuum to obtain pale yellow solid. It

was then purified by column chromatography using (Ethyl Acetate:n-Hexane; 30:70) as a mobile phase. Crystalline solid was isolated. Yield: 250 mg, (52 %).

Step C: The compound obtained in step B (250 mg, 0.30 mmol) was dissolved in acetonitrile (10 ml). 70% HF-Pyridine solution (18 μ l, 2.0 eq.) was added to the above solution and stirred at room temperature for overnight under N_2 atmosphere. After completion of reaction solvent was evaporated under vacuum to obtain crude product. Water (20 ml) was added to the crude product and extracted with dichloromethane (25 ml x 3). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude white solid. It was then purified by column chromatography using $CHCl_3$: MeOH; 90:10) as a mobile phase to obtain title compound as white solid compound. Yield : 120 mg, (56 %) MS 694 ($M + H$)⁺.


Example-181: Compound of Formula I-f where T is $CO(CH_2)_2$ -(3-fluorophenyl);

Step-1: To the stirred solution of 3-(3-fluoro-phenyl)-propionic acid (0.321 g, 1.91 mmol) in dichloromethane (10 ml) was added dicyclohexylcarbodiimide (0.788 g, 3.82 mmol) followed by 1-hydroxy-benzotriazole (HOBT) (0.129 g, 0.955 mmol) at a temperature between 25-30° C. The reaction mixture was stirred for 30 minutes. To the reaction mixture was added Example-7 (1.0 g, 1.3 mmol) followed by N,N-dimethylamino pyridine (0.077 g, 0.636 mmol) under stirring and it was stirred for additional 1 hour at 25-30° C temperature. The resultant mixture was filtered under suction, the filtrate was evaporated under vacuum to provide step-1 product as a thick liquid in 0.91 g quantity, which was used without purification for further reaction MS: m/z: 936.1 ($M+1$).

Step-2: To the stirred solution of N-chlorosuccinimide (0.97 g, 7.28 mmol) in dichloromethane (15 ml) at 0° C was added dimethyl sulfide (1.6 ml, 12.12 mmol). The mixture was stirred at 0° C for additional for 30 minutes. The reaction mixture was cooled to -40° C and the solution of step-1 product (0.91 g, 0.97 mmol) dissolved in dichloromethane (10 ml) was added. The resulting reaction mixture was stirred at -40° C temperature for 2 hr and then allowed to warm at 25-30° C while stirring. Triethyl amine (0.980 ml, 9.71 mmol) was added to the reaction mixture at 25-30° C and it was stirred for additional 30 minutes. The reaction mixture was quenched by pouring in aqueous saturated sodium bicarbonate solution and was extracted with dichloromethane (3 x 25 ml). The combined separated organic layer was dried over Na_2SO_4 and was evaporated to provide residue. The residue was purified by using silica gel column chromatography (10 % acetone : hexanes) to provide step-2 product as a off white solid in 80.3% (0.73 g) yield. MS: m/z: 934.1 ($M+1$).

Step-3: Deprotection: To a mixture of step-2 compound (0.73 g, 0.782 mmol) and 70% HF-pyridine solution (26 μ l, 0.912 mmol) in acetonitrile (10 ml), was added at 25-30° C temperature for 2 hr. The solvent was evaporated under vacuum to obtain a crude residue. To the crude residue, water (20 ml) was added and the mixture was extracted with dichloromethane (3 X 25 ml). Combined organic layer was dried over Na_2SO_4 and evaporated under vacuum to obtain crude solid. The crude solid was purified by using silica gel column chromatography ($\text{CHCl}_3:\text{MeOH}$; 90:10) to obtain title compound as white solid compound in 78.3% (0.570 g) yield. Mp: 174-177° C, MS: m/z: 820.1 ($\text{M}+1$).

Following examples were prepared by using above procedure and by using carboxylic acid A.



Example	T	Carboxylic acid (A)	Mp (° C)	Mass ($\text{M}+1$)
182	$\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-fluoro-phenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$	3-[3-(3-fluoro-phenyl)-[1,2,4] oxadiazol-5-yl]-propionic acid	166-168	888.1
183	$\text{CO}(\text{CH}_2)_2\text{-}(3,4\text{-dimethoxy-phenyl})$	3-(3,4-dimethoxy-phenyl)-propionic acid	168-170	862.1
184	$\text{CO}(\text{CH}_2)_2\text{-}(4\text{-methoxy-phenyl})$	3-(4-methoxy-phenyl)-propionic acid	154-160	803.1

Example-185, Compound of Formula I-f where T is $\text{CH}_2\text{C}\equiv\text{CH}$:

Step-A: To the stirred solution of Example-7 (1.0 g, 1.3 mmol) in toluene (15 ml) was added potassium *tert*-butoxide (0.16 g, 1.43 mmol) and 18-crown-6-ether (0.048 g, 0.13 mmol) at 25-30° C temperature. The reaction mixture was stirred for 10 min at 25-30° C temperature. 3-bromo-propyne (127 μ l, 1.43 mmol) was added to the reaction mixture and it was stirred for additional 1 hour. It was quenched by pouring it in aqueous saturated ammonium chloride solution (20 ml). The mixture was extracted with ethyl acetate (3 x 25 ml). Combined organic layer was dried over Na_2SO_4 and purified by using silica gel column chromatography (10 % acetone : Hexane) to provide step-1 product as a pale yellow solid in 74% (0.77 g) yield. MS: m/z 825 ($\text{M}+1$).

Step-B: To the stirred solution of N-chlorosuccinimide (0.94 g, 7.0 mmol) in dichloromethane (20 ml) at 0 °C was added dimethyl sulfide (0.86 ml, 11.8 mmol). The reaction mixture was then stirred at 0 °C for 30 min. The step-1 product (0.77 g, 0.94 mmol) dissolved in dichloromethane (10 ml) was added to the reaction mixture at – 40° C. The resulting reaction mixture was stirred at the 0° C temperature for 2 hr and allowed to warm at 25-30° C temperature under stirring. Triethyl amine (1.3 ml, 9.4 mmol) was added to the reaction mixture and stirred for additional 30 min. The mixture was poured in aqueous saturated sodium bicarbonate solution and extracted with dichloromethane (3 x 25 ml). The combined organic layer was dried over Na₂SO₄ and purified by using silica gel column chromatography (10 % acetone:Hexane) to provide step-2 product as a off white solid in 80% (0.62 g) yield. MS: m/z: 823 (M+1).

Step-C: A mixture of step-2 product (0.62 g, 0.75 mmol) and 70% HF-pyridine solution (32 µl, 1.1 mmol) dissolved in acetonitrile (10 ml) was stirred at 25-30° C for 2 hr. After completion of reaction, solvent was evaporated under vacuum to obtain a residue. Water (20 ml) was added to the residue and the mixture was extracted with dichloromethane (3 X 25 ml). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain a crude solid. The crude solid was purified by using silica gel column chromatography (10 % MeOH : CHCl₃) to obtain title compound as white solid in 60% (0.37 g) yield. Mp: 202-204° C, MS: m/z: 709.1 (M+1).

Example-186: Compound of Formula I-f where T is CH₂CH≡C-(pyridin-2-yl);


Step-A: To the stirred suspension of Pd(PPh₃)₂Cl₂ (0.13 g, 0.18 mmol), CuI (46 mg, 0.24 mmol) and 2-iodopyridine (0.25 g, 1.2 mol) a solution of compound obtained in step A of the above example (1.0 g, 1.2 mmol) dissolved in diethyl amine (5.0 ml) was added at 25-30° C temperature. The reaction mixture was stirred under inert atmosphere for additional 3 hr at 25-30° C. It was poured in ice cold water and extracted with ethyl acetate (3 X 25 ml). Combined organic layer was dried over Na₂SO₄ and evaporated under vacuum to provide a crude mass. The crude mass was purified by using silica gel column chromatography (15 % acetone-Hexane) to yield step-1 product as a pale yellow solid in 60% (0.65 g) yield. MS: m/z: 901 (M+1)

Step-B: To the stirred solution of N-chlorosuccinimide (0.73 g, 5.5 mmol) in dichloromethane (20 ml) at 0° C was added dimethyl sulfide (0.66 ml, 9.1 mmol). The reaction mixture was stirred under inert atmosphere at 0° C for 30 min. The step-1 product

(0.65 g, 0.73 mmol) dissolved in dichloromethane (10 ml) was added to the reaction mixture at -40°C . The resulting reaction mixture was stirred at the 0°C temperature for 2 hr and allowed to warm at $25\text{-}30^{\circ}\text{C}$ temperature. Triethyl amine (1.0 ml, 7.3 mmol) was added to the reaction mixture and stirred for additional 30 min. The reaction mixture was poured in aqueous saturated sodium bicarbonate solution. The separated organic layer dried over Na_2SO_4 and purified by using silica gel column chromatography (10 % Acetone:Hexane) to obtain step-2 product as a off white solid in 75% (0.49 g) yield. MS: m/z: 899 (M+1).

Step-C: A mixture of step-2 product (0.48 g, 0.53 mmol) and 70% HF-pyridine solution (23 μl , 0.8 mmol) dissolved in acetonitrile (10 ml) was stirred at $25\text{-}30^{\circ}\text{C}$ for 2 hr under N_2 atmosphere. After completion of reaction, solvent was evaporated under vacuum to obtain crude mass. Water (20 ml) was added to the crude mass and the mixture was extracted with dichloromethane (3 X 25 ml). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude solid. The crude solid was purified by using column chromatography (CHCl_3 : MeOH; 90:10) to obtain title compound as white solid in 70% (0.29 g) yield. MS: m/z: 785 (M+1). Mp: 179-183 $^{\circ}\text{C}$.

Following examples were prepared by using above procedure and by using reactant A in the place of 2-iodoyridine.



Example	T	Reactant (A)	Mp ($^{\circ}\text{C}$)	Mass (M+1)
187	$\text{CH}_2\text{C}\equiv\text{C}$ -(3-cyano) phenyl	3-Iodo-benzo nitrile	139-142	809.1
188	$\text{CH}_2\text{C}\equiv\text{C}$ -(3-fluoro) phenyl	1-Fluoro-3-iodo-benzene	175-178	802.1
189	$\text{CH}_2\text{C}\equiv\text{C}$ -(3-chloro) phenyl	1-Chloro-3-iodo-benzene	171-174	818.1
190	$\text{CH}_2\text{C}\equiv\text{C}$ -(3-methoxy) phenyl	1-Iodo-3-methoxy-benzene	168-171	814.1
191	$\text{CH}_2\text{C}\equiv\text{C}$ -(2-chloro pyridin-5-yl)	2-Chloro-5-iodo-pyridine	167-170	819.1
192	$\text{CH}_2\text{C}\equiv\text{C}$ -(2-fluoro pyridin-5-yl)	2-Fluoro-5-iodo-pyridine	173-176	803.1

193	CH ₂ C≡C-(pyridin-3-yl)	3-Iodo-pyridine	132-137	785.1
-----	------------------------------------	-----------------	---------	-------

Example-194, Compound of Formula I-f where T is CH₂-phenyl;


Step-1: To the solution of Example-7 (1.0 g, 1.3 mmol) in toluene (10 ml) was added potassium *tert*-butoxide (0.160 g, 1.43 mmol), 18-crown-6-ether (0.048 g, 0.13 mmol), followed by benzyl bromide (182 μ l, 1.56 mmol) at a 25-30° C. The reaction mixture was stirred at 25-30° C temperature for 1 hour. After completion of the reaction, the reaction mixture was poured in saturated aqueous ammonium chloride solution (20 ml). The mixture was extracted with ethyl acetate (3 x 25ml). Combined organic layer was dried over Na₂SO₄ and evaporated under vacuum to provide a residue. The residue was purified by using silica gel column chromatography (10 % acetone : hexanes) to provide step-1 product as a off white solid in 76.2% (0.85 g) yield. MS: m/z: 876 (M+1)

Step-2: Method A: To the stirred solution of N-chlorosuccinimide (0.97 g, 7.28 mmol) in dichloromethane (15 ml) at 0° C was added dimethyl sulfide (1.6 ml, 12.12 mmol). The reaction mixture was stirred at 0° C for additional 30 minutes. The complex was cooled to -40° C and the solution of step-1 product (0.85 g, 0.97 mmol) dissolved in dichloromethane (10 ml) was added. The resulting reaction mixture was stirred at -40° C temperature for 2 hr and allowed to warm at 25-30° C under stirring. Triethyl amine (0.980 ml, 9.71 mmol) was added to the reaction mixture at 25-30° C and was stirred for additional 30 minutes. The mixture was poured in aqueous saturated sodium bicarbonate solution (50 ml) and it was extracted with dichloromethane (3 x 25 ml). The combined organic layer was dried over Na₂SO₄ and was evaporated to provide a residue. The residue was purified by using silica gel column chromatography (10 % acetone : hexanes) to provide step-2 product as a off white solid in 79% (0.67 g) yield. MS: m/z; 874 (M+1).

Method B: Alternatively, oxidation was carried out by using Dess-martin periodinane reagent as follows: To the stirred solution of the step-1 product (1.5 g, 1.6 mmol) in dichloromethane (10 ml), was added Dess-martin periodinane (5.0 ml, 1.76 mmol, 15% solution in dichloromethane). The reaction mixture was stirred at a temperature 25-30°C for 30 min. The reaction mixture was poured in 1:1 mixture of aqueous saturated sodium bicarbonate solution and aqueous saturated sodium thiosulphate solution and it was extracted with dichloromethane (3 x 25 ml). The combined organic layer was dried over Na₂SO₄ and purified by using silica gel column chromatography (10 % Acetone: Hexane) to provide step-2 product in 46% (0.7 g) yield. MS: m/z; 874 (M+1).

Step-3: To a mixture of step-2 product (0.67 g, 0.76 mmol) and 70% HF-pyridine solution (26 μ l, 0.912 mmol) in acetonitrile (10 ml), was stirred at 25-30° C temperature for 2 hr. The solvent was evaporated under vacuum to obtain a crude residue. To the crude residue, water (20 ml) was added and the mixture was extracted with dichloromethane (3 X 25). Combined organic layer was dried over Na_2SO_4 and evaporated under vacuum to obtain a crude solid. The crude solid was purified by using silica gel column chromatography ($\text{CHCl}_3:\text{MeOH}$; 90:10) to afford title compound as white solid compound in 76.2% (0.51 g) yield., MS: m/z: 761.1 ($\text{M}+1$). Mp: 200-204° C.

Following examples were prepared by using above procedure and by using reactant A.



Example	T	Reactant (A)	Mp (° C)	Mass ($\text{M}+1$)
195	$\text{CH}_2\text{-}(4\text{-methoxyphenyl})$	1-bromomethyl-4-methoxy-benzene	184-187	790.1
196	$\text{CH}_2\text{-}(3\text{-chlorophenyl})$	1-bromomethyl-3-chlorobenzene	212-214	794.1
197	$\text{CH}_2\text{-}(3\text{-methoxyphenyl})$	1-bromomethyl-3-methoxy-benzene	212-215	790.1
198	$\text{CH}_2\text{-}(4\text{-}(1\text{H-[1,2,4]} \text{ triazol})\text{phenyl})$	1-(4-bromo methylphenyl)-1H-[1,2,4]triazole	150-154	827.1
199	$\text{CH}_2\text{-}(4\text{-}(isopropyl)phenyl)$	1-Bromomethyl-4-isopropyl-benzene	189-191	802.1
200	$\text{CH}_2\text{-}(2\text{-fluorophenyl})$	1-Bromomethyl-2-fluorobenzene	176-179	778.1
201	$\text{CH}_2\text{-}(4\text{-}(pyrimidin-5-yl)phenyl)$	5-(4-bromo methylphenyl)-pyrimidine	164-167	838.1
202	$\text{CH}_2\text{-}(4\text{-}(pyridin-2-yl)phenyl)$	2-(4-bromo methylphenyl)-pyridine	170-172	837.1
203	$\text{CH}_2\text{C}(\text{=CH}_2)\text{CH}_3$	3-Chloro-2-methylpropene	174-177	714.1
204	$\text{CH}_2\text{CH}=\text{CHCH}_3$	1-Chloro-but-2-ene	221-224	724.1
205	$\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$	1-Chloro-3-methyl-but-2-ene	244-248	738.1

206	CH ₂ C(F)=CH ₂	3-Chloro-2-fluoro-1-propene	174-176	728.1
-----	--------------------------------------	-----------------------------	---------	-------


Example-207, Compound of Formula I-f where T is CH₂CH=CH-(pyridin-3-yl);

Step A: To the stirring solution of Step A product from Example 179 (1.5 g, 1.81 mmol) in DMF (15 ml) was added sodium acetate (0.24 g, 2.73 mmol). To the above solution Tetrakis triphenyl phosphine palladium (0) (0.21g, 0.18 mmol) and 3-bromopyridine (0.195 μ l, 1.99 mmol) were added and stirred the reaction mixture for 12 hour at 120 °C temperature. The reaction mixture was then poured in sat. ammonium chloride solution (20 ml) and extracted with ethyl acetate (50 ml x 3). Combined organic layer was dried over Na₂SO₄ it was then purified over silica gel column chromatography (18 % Acetone: Hexane), yield (0.65 g, 53 %).

Step B: To the stirred solution of step-1 compound (0.65 g, 0.72 mmol) in DCM (130 ml) was added Dess-Martin perriodinane reagent (15 %solution in DCM) (3.4 ml, 1.18 mmol) at 25-30° C temperature. The reaction mixture was stirred further for 30 min. The reaction mixture was quenched with 1:1 mixture of aqueous saturated sodium bicarbonate solution and aqueous sodium thiosulfate solution (2 x 200 ml) the organic layer was washed with brine (100 ml). Combined organic layer was dried over Na₂SO₄ and concentrated under vacuum to provide a crude mass. The crude material was purified by flash column chromatography using acetone: hexane 2.2:7.8 mixture to provide pale yellow solid in 45% (0.35 g) yield.

Step C: To the solution of compound obtained from above step (0.35 g, 0.37 mmol) in acetonitrile (15 ml) was added 70% hydrogen fluoride in pyridine (16 μ l, 0.58 mmol) under inert atmosphere and stirred for 3 h. The residue obtained after evaporation of solvent under reduced pressure was dissolved in ethyl acetate (15 ml) and washed with water (2 x 10ml). The organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to provide crude solid which on trituration in diethyl ether afforded white solid as the title compound. Yield: 0.180 g (60%), Mass: m/z: 787.1 (M+H), Mp: 212-215 °C.

Following examples were prepared by using above procedure and by using reactant A.



Example	T	Reactant (A)	Mp (° C)	Mass (M+1)
208	CH ₂ CH=CH-phenyl	iodobenzene	202-205	786.1
209	CH ₂ CH=CH-(4-trifluoromethyl)phenyl	1-iodo-4-trifluoro methyl-benzene	96-100	854.1
210	CH ₂ CH=CH-(3-trifluoromethyl) phenyl	1-iodo-3-trifluoro methyl-benzene	206-210	854.1
211	CH ₂ CH=CH-(3-fluoro)phenyl	1-fluoro-3-iodo-benzene	214-216	804.1
212	CH ₂ CH=CH-(3-chloro)phenyl	1-chloro-3-iodo-benzene	154-158	820.1
213	CH ₂ CH=CH-(pyrimidin-5-yl)	5-Iodo-pyrimidine	158-160	788.1

Example-214: Compound of Formula I-f where T is CH₂CONH-(3-chlorophenyl);


Step-1: To the stirred solution of m-chloro aniline (2.0 g, 1.57 mmol) in THF (20 ml) was added triethyl amine (2.62 ml, 1.88 mmol), bromoacetyl bromide (1.51ml, 1.73 mmol) at 5° C and the reaction mixture was stirred for 1 hour at 25-30° C temperature. The reaction mixture was poured in aqueous saturated ammonium chloride solution (20 ml) and extracted with ethyl acetate (3 X 25 ml). Combined organic layer dried over Na₂SO₄ and distilled under reduced pressure. Yellow coloured compound was obtained as a step-1 product and utilized for the next step.

Step-2: To the stirred solution of Example-7 (2.00 g, 2.50 mmol) in toluene (10 ml) was added potassium hydride (3.41 g, 0.30 mmol) and 18-crown-6-ether (0.08 mg, 0.30 mmol) at 25-30°C temperature. The reaction mixture was stirred for additional 10 min. and step-1 product 2-bromo-N-(3-chloro-phenyl)-acetamide (0.69 g, 2.81 mmol) was added. The reaction mixture was stirred for 1 hour at 25-30° C temperature. It was poured in aqueous saturated ammonium chloride solution (30 ml) and extracted with ethyl acetate (3 x 25 ml). Combined organic layer was dried over Na₂SO₄ and purified by using silica gel column chromatography (10 % Acetone:Hexane) to provide step-2 product as a off white solid in 70% (1.68 g) yield. MS: m/z: 953 (M+1).

Step-3: To the stirred solution of N-chlorosuccinimide (1.77 gm, 13.22 mmol) in dichloromethane (20 ml) at 0° C was added dimethyl sulfide (1.62 ml, 22.00 mmol). The reaction mixture was stirred at 0° C for next 30 min. The Step-1 product (1.68 gm, 1.76 mmol) dissolved in dichloromethane (10 ml) was added to the reaction mixture at -40° C. The resulting reaction mixture was stirred at the same temperature for 2 hr and allowed to warm at 25-30° C temperature. Triethyl amine (2.45 ml, 17.62 mmol) was added to the reaction mixture and stirred for additional 30 min. The reaction mixture was poured in aqueous saturated sodium bicarbonate solution and extracted with dichloromethane (3 x 25 ml). The combined organic layer dried over Na₂SO₄ and purified by using silica gel column chromatography (10 % acetone:hexane) to provide step-3 product as a off white solid in 80 % (1.34 g) yield. MS: m/z: 951 (M+1).

Step-4: A mixture of step-3 product (1.34 gm, 1.40 mmol) and 70% HF-pyridine solution (60 µl, 2.16mmol) in acetonitrile (10 ml) was stirred at 25-30° C for 2 hr. After completion of reaction, solvent was evaporated under vacuum to obtain crude mass. Water (20 ml) was added to the crude mass and extracted with dichloromethane (3 X 25 ml). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude solid. The crude solid was purified by using silica gel column chromatography (10 % MeOH:CHCl₃) to obtain title compound as white solid in 35% (0.42 g) yield., MS: m/z: 838 (M+1). Mp: 183-186° C.

Following examples were prepared by using above procedure and by using reactant A.



Example	T	Reactant (A)	Mp (° C)	Mass (M+1)
215	CH ₂ CONH-(3-fluoro-phenyl)	2-bromo-N-(3-fluoro-phenyl)-acetamide	176-178	821.1
216	CH ₂ CO-(4-phenylpiperazin-1-yl)	2-bromo-1-(4-phenyl-piperazin-1-yl)-ethanone	214-218	872.1
217	CH ₂ CONH ₂	2-bromoacetamide	220-222	727.1

218	CH ₂ CON(CH ₃) ₂	2-bromo-N,N-dimethylacetamide	210-214	755.1
219	CH ₂ CONH-(cyclopropyl)	2-bromo-N-cyclopropyl-lacetamide	214-218	767.1

Example-220: Compound of Formula I-f where T is CH₂-(5-(pyridin-3-yl)[1,3,4] oxadi azol-2-yl);

Step-1: To a mixture of 3-cyanopyridine (12 g, 115 mmol) and trimethylsilyl azide (20 ml, 150 mmol), was added a solution of 0.1 M tetra-*n*-butylammonium fluoride in THF (57 ml, 56 mmol) at 25-30° C temperature. The resulting mixture was heated at 80° C for overnight. The mixture is allowed to warm at 25-30° C temperature and then quenched in ice-water mixture. Solid precipitated out was filtered and washed with water (2 x 25 ml) and dried under vacuum to provide step-1 compound in 62% (10.5 g) yield. MS: 148 (M+1).

Step-2: To the stirred solution of 3-(1H-tetrazol-5-yl)-pyridine (10.5 g, 71.42 mmol) in toluene (50 ml) was added chloroacetic anhydride (25 g, 92.80 mmol) and resulting mixture was heated at 100° C for 2-3 hours. The reaction mixture was allowed to warm at 25-30° C temperature. The mixture was quenched with saturated aqueous sodium bicarbonate solution (500 ml) and extracted with chloroform (2 x 200 ml). The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure to provide a crude mass. The crude mass was purified over silica gel column chromatography to provide title compound in 56% (7.8 g) yield. MS: 196 (M+1).


Step-3: To the stirred solution of potassium hydride (5.5 g, 41 mmol) and 18-crown-6 (1.1 g, 4.1 mmol) in toluene (200 ml) was added Example-7 (25 g, 32 mmol). It was allowed to stir at 25-30° C temperature for 30 min. (5-chloromethyl-1,3,4-oxadiazole-2-yl)-pyridine (7.8 g, 38 mmol) was added to the reaction mixture. The reaction mixture was stirred at 25-30° C temperature for 30 minutes. It was quenched with aqueous saturated ammonium chloride solution (500 ml). The organic layer was washed with brine (500 ml), dried over Na₂SO₄ and concentrated under vacuum to provide a crude mass. The crude mass was purified by flash column chromatography (acetone:hexane 2.5:7.5) to provide off white solid in 43% (13 g) yield. MS: m/z 945 (M+1).

Step-4: To the stirred solution of step-3 compound (13 g, 14 mmol) in DCM (130 ml) was added Dess-Martin periodinane reagent (15 % solution in DCM) (58 ml, 21 mmol) at 25-30° C temperature. The reaction mixture was stirred further for 30 min. The reaction mixture was quenched with 1:1 mixture of aqueous saturated sodium bicarbonate solution and aqueous sodium thiosulfate solution (2 x 200 ml) the organic layer was washed with brine (100 ml).

Combined organic layer was dried over Na_2SO_4 and concentrated under vacuum to provide a crude mass. The crude material was purified by flash column chromatography using acetone: hexane 2.2:7.8 mixture to provide pale yellow solid in 45% (7.4 g) yield. MS: m/z: 943 (M+1).

Step-5: A mixture of step-4 compound (7.4 g, 7.8 mmol) in acetonitrile (74 ml) and 70 % HF-pyridine complex (0.3 ml, 12 mmol) was stirred at 25-30° C temperature for 3-4 hours. The reaction mixture was concentrated under reduced pressure and water was added and the mixture was extracted with DCM (2 x 40 ml). Combined organic layer was dried over Na_2SO_4 and concentrated under vacuum. The crude mass was purified by flash column chromatography (MeOH: CHCl_3 1.2: 8.8) to give title compound as a white solid in 45% (3.4 g) yield., MS: m/z: 829 (M+1). Mp: 212-214° C

Following examples were prepared by using above procedure and by using Reagent A.



Example	T	Reagent A	Mp (° C)	Mass (M+1)
221	$\text{CH}_2\text{-}(5\text{-phenyl}[1,3,4]\text{-oxadiazol-2-yl})$	2-chloromethyl-5-phenyl [1,3,4]-oxadiazole	134-136	829.1
222	$\text{CH}_2\text{-}(5\text{-}(3\text{-fluorophenyl)[1,3,4]oxadiazol-2-yl})$	2-chloromethyl-5-(3-fluoro-phenyl)-[1,3,4]-oxadiazole	125-127	845.1
223	$\text{CH}_2\text{-}(5\text{-}(4\text{-methoxy phenyl)[1,3,4]oxadiazol-2-yl})$	2-chloromethyl-5-(4-methoxy-phenyl)-[1,3,4]-oxadiazole	130-133	857.1
224	$\text{CH}_2\text{-}(5\text{-}(pyridin-2-yl)[1,3,4]oxadiazol-2-yl)$	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-pyridine	205-208	829.1
225	$\text{CH}_2\text{-}(5\text{-}(pyrazin-2-yl)[1,3,4]oxadiazol-2-yl)$	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-pyrazine	188-190	831.1

	yl)[1,3,4]oxadiazol-2-yl)	[1,3,4]-oxadiazol-2-yl)-pyrazine		
226	CH ₂ -(5-(3,5-dimethoxyphenyl)[1,3,4]oxadiazol-2-yl)	2-chloromethyl-5-(3,5-dimethoxy-phenyl)-[1,3,4]-oxadiazole	171-179	887.1
227	CH ₂ -(5-(pyrimidin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-pyrimidine	187-191	831.1
228	CH ₂ -((5-cyclopropyl)-[1,3,4]oxadiazol-2-yl)	2-chloromethyl-5-cyclopropyl-[1,3,4]-oxadiazole	186-192	791.1
229	CH ₂ -(5-(6-methoxy-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-6-methoxy-pyridine	225-230	860.1
230	CH ₂ -(5-(5-methyl-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-methyl-pyridine	-	844.1
231	CH ₂ -(5-(5-cyclopropyl-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-cyclopropyl-pyridine	-	870.1
232	CH ₂ -(5-(5-cyano-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-cyano-pyridine	-	855.1
234	CH ₂ -(5-(5-dimethyl amino-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-dimethyl amino-pyridine	-	874.1
235	CH ₂ -(5-(5-methoxy-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-methoxy-pyridine	-	860.1
236	CH ₂ -(5-(5-fluoro-pyridin-2-yl)[1,3,4]oxadiazol-2-yl)	2-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-5-fluoro-pyridine	-	848.1
237	CH ₂ -(5-(5-chloro-pyridin	2-(5-chloromethyl-	-	864.1

	-2-yl)[1,3,4]oxadiazol-2-yl)	[1,3,4]-oxadiazol-2-yl)-5-chloro-pyridine		
238	CH ₂ -(5-(pyrimidin-5-yl)[1,3,4]oxadiazol-2-yl)	5-(5-chloromethyl-[1,3,4]-oxadiazol-2-yl)-pyrimidine	-	831.1

Example-239: Compound of Formula I-f, where T is CH₂-(3-(pyridin-2-yl)[1,2,4]oxadiazol-5-yl);

Step-1: To the stirred suspension of hydroxylamine hydrochloride (8.01 g, 120 mmol) in methanol (100 ml) was added sodium bicarbonate (10.50 g, 130 mmol) followed by pyridine-2-carbonitrile (10 g, 96 mmol). The reaction mixture was refluxed for 2 hours. The solvent was removed under vacuum to provide a residue. Water (200 ml) was added to the residue. The solid separated was filtered under suction and washed with water (100 ml) dried under vacuum to provide step-1 product as a white solid in 93% (12.5 g) yield.

MS: m/z: 138 (M+1).

Step-2: To the stirred suspension of step-1 product (12.50 g, 91.2 mmol) in acetone (100 ml), chloroacetyl chloride (7.36 ml, 120 mmol) was added drop-wise at 25-30° C temperature. The reaction mixture was stirred for 30 minutes. The solid separated was filtered and stirred with saturated aqueous NaHCO₃ solution (200 ml). The suspension was filtered and washed with water, dried under vacuum to provide step-2 product as a off white solid in 83% (18 g) yield. The solid was used as it is for the next step. MS: m/z: 214.6 (M+1).

Step-3: The step-2 product (18 g, 85 mmol) was refluxed in toluene (200 ml) along with 4 Å molecular sieves for 2 hrs. The reaction mixture was concentrated under vacuum to provide a crude residue. The crude residue was triturated with diethyl ether to afford 5-chloromethyl-1,2,4-oxadiazol-3-yl)-pyridine, as step-3 product as a off white solid in 96% (16 g) yield. MS: m/z: 197.6 (M+1).


Step-4: To the stirred mixture of potassium hydride (2.1 g, 17 mmol), 18-crown-6 (0.33 g, 1.2 mmol) in toluene (100 ml) was added Example-7 (10 g, 13 mmol). The reaction mixture was stirred at 25-30° C temperature for 30 min and step-3 product, 2-(5-chloromethyl-1,2,4-oxadiazol-3-yl)-pyridine (3.2 g, 14 mmol) dissolved in toluene (5 ml) was added to the reaction mixture. It was stirred for additional 30 min. The reaction mixture was quenched by addition of saturated aqueous ammonium chloride solution (250 ml). Layers were separated. The organic layer was washed with brine (250 ml), dried over Na₂SO₄, concentrated under

vacuum to provide crude solid. The crude solid was purified by flash column chromatography (acetone:hexane 2.5:7.5) to isolate step-4 product as a off white solid in 58% (7.0 g) yield. MS: m/z: 945 (M+1).

Step-5: To a stirred solution of N-chlorosuccinimide (8.66 g, 56 mmol) in dry toluene (100 ml) at 0 °C was added dimethyl sulfide (7.7 ml, 93 mmol). A white precipitate appeared. The mixture was cooled to -40 °C, and a solution of step-4 product (7.0 g, 7.4 mmol) in DCM (30 ml) was added over 30 min. Stirring was continued for 3 hours at same temperature. TEA (12.5 ml, 74 mmol) was added to the reaction mixture and the reaction mixture was allowed warm at 25-30° C temperature. It was quenched with saturated aqueous sodium bicarbonate solution (100 ml). The organic layer was washed with brine (100 ml), dried over Na₂SO₄, concentrated under vacuum to provide a crude mass. The crude mass was purified by using silica gel column chromatography (acetone:Hexane ;20:80) to isolate step-5 product as a off white solid in 67% (4.7 g) yield. MS: m/z: 943 (M+1).

Step-6: A mixture of the step-5 product (4.7 g, 5.0 mmol) and 70 % HF-Pyridine complex (0.17 ml, 7.5 mmol) in acetonitrile (40 ml) was stirred at 25-30° C temperature for 3-4 hours. The reaction mixture was concentrated under vacuum to provide a residue and to the residue, water (50 ml) was added. The product was extracted in DCM (2 x 30 ml), dried over Na₂SO₄ and concentrated under vacuum to provide a crude solid. The crude solid was purified by using flash column chromatography (MeOH:CHCl₃ ; 1.5:8.5) to give title compound in 61% (2.5 g) yield., MS: m/z: 829 (M+1) Mp: 172-175° C

Following examples were prepared by using above procedure and by using Reagent A in the place of 2-(5-chloromethyl-1,2,4-oxadiazol-3-yl)-pyridine.



Example	T	Reagent (A)	Mp (° C)	Mass (M+1)
240	CH ₂ -(3-phenyl[1,2,4]-oxadiazol-5-yl)	5-chloromethyl-3-phenyl-[1,2,4]-oxadiazole	162-165	827.1

241	CH ₂ -(3-(3-fluorophenyl)[1,2,4]oxadiazol-5-yl)	5-chloromethyl-3-fluoro phenyl-[1,2,4]-oxadiazole	150-152	845.1
242	CH ₂ -(3-(3-chlorophenyl)[1,2,4]oxadiazol-5-yl)	5-chloromethyl-3-chloro phenyl-[1,2,4] oxadiazole	154-157	861.1
243	CH ₂ -(3-(4-methoxyphenyl)[1,2,4]oxadiazol-5-yl)	5-chloromethyl-4-methoxyphenyl-[1,2,4] oxadiazole	173-176	857.1
244	CH ₂ -(3-(pyridin-3-yl)[1,2,4]oxadiazol-5-yl)	3-(5-chloromethyl-[1,2,4] oxadiazol-3-yl)-pyridine	210-214	829.1
245	CH ₂ -(3-(4-chlorophenyl)[1,2,4]oxadiazol-5-yl)	5-chloromethyl-4-chlorophenyl-[1,2,4] oxadiazole	135-138	861.1
246	CH ₂ -(3-(3,5-dimethoxyphenyl)[1,2,4]oxadiazol-5-yl)	5-chloromethyl-3,5-dimethoxy phenyl-[1,2,4] oxadiazole	154-158	887.1
247	CH ₂ -(3-(pyrimidin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4] oxadiazol-3-yl)-pyrimidine	162-165	831.1
248	CH ₂ -(3-cyclopropyl[1,2,4]oxadiazol-5-yl)	5-chloromethyl-3-cyclopropyl-[1,2,4]oxadiazole	176-178	792.1
249	CH ₂ -(3-(6-methylpyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4] oxadiazol-3-yl)-6-methyl-pyridine	202-205	842.1
250	CH ₂ -(3-(3-methylpyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4] oxadiazol-3-yl)-3-methyl-pyridine	194-198	842.1
251	CH ₂ -(3-(5-methylpyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-methyl-pyridine	-	844.1
252	CH ₂ -(3-(5-cyclopropyl-pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-cyclo propyl-	-	870.1

		pyridine		
253	CH ₂ -(3-(5-cyano-pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-cyano-pyridine	-	855.1
254	CH ₂ -(3-(5-dimethyl amino-pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-dimethyl amino-pyridine	-	874.1
255	CH ₂ -(3-(5-methoxy -pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-methoxy-pyridine	-	860.1
256	CH ₂ -(3-(5-fluoro-pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-fluoro-pyridine	-	848.1
257	CH ₂ -(3-(5-chloro-pyridin-2-yl)[1,2,4]oxadiazol-5-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-5-chloro-pyridine	-	864.1
258	CH ₂ -(3-(pyrimidin-5-yl)[1,2,4] oxadiazol-5-yl)	5-(5-chloromethyl-[1,2,4]-oxadiazol-3-yl)-pyrimidine	-	831.1

Example-259: Compound of Formula I-f where T is CH₂-(5-(pyridin-3-yl)[1,2,4]oxadiazol-3-yl);

Step-1: To the stirred solution of 2-chloroacetamidoxime (1.0 g, 9.2 mmol, prepared as per the procedure given in PCT/US02/22897) in toluene (15 ml) was added nicotinyl chloride (1.37 g, 9.67 mmol) and heated at 110° C for 4 hr. The reaction mixture was diluted with ethyl acetate (50 ml) and washed with aqueous saturated sodium bicarbonate solution (2 X 25). Organic layer was evaporated under vacuum to yield the crude product. The crude product was purified using silica gel column chromatography (15 % Acetone-Hexane) to give 2-(3-chloromethyl-[1,2,4]-oxadiazole-5-yl)-pyridine in 55 % (0.550 g) yield. MS: m/z: 196.7 (M+1).

Step-2: The 2-(3-chloromethyl-[1,2,4]-oxadiazole-5-yl)-pyridine (0.550 g, 2.8 mmol) was dissolved in acetone (20 ml). To the clear solution, sodium iodide (0.843 g, 5.62 mmol) and tetra-*n*-butyl-ammonium iodide (0.259 g, 0.70 mmol) were added. The reaction mixture was


refluxed for 2 hr. It was quenched by addition of water (20 ml) and extracted with ethyl acetate (2 X 25 ml). The combined organic layer was dried over Na_2SO_4 and evaporated under vacuum to yield 2-(3-Iodomethyl-[1,2,4]oxadiazole-5-yl)-pyridine in 90% (0.750 g) yield. MS: m/z: 288.1 (M+1).

Step-3: To the stirring solution of Example-7 (1.0 gm, 1.3 mmol) in toluene(10 ml) was added potassium hydride (0.16 g, 1.43 mmol) and 18-crown-6-ether (0.048 g, 0.13 mmol) at a temperature 25-30° C and stirred for 10 min. To this reaction mixture, step-2 product, 2-(3-iodomethyl- [1,2,4] oxadiazole-5-yl)-pyridine (0.401 g, 1.43 mmol) was added and stirred for 1 hour at temperature 25-30° C. The reaction mixture was then poured in aqueous saturated ammonium chloride solution (20 ml) and the mixture was extracted with ethyl acetate (3 x 25ml). Combined organic layer was dried over Na_2SO_4 and purified by using silica gel column chromatography (15 % Acetone :Hexane). To afford step-3 product as yellow solid in 65% (0.780 g) yield. MS: m/z: 946.3 (M+1).

Step-4: To the stirred solution of N-chlorosuccinimide (0.826 g, 6.18 mmol) in toluene (15 ml) at 0° C was added dimethyl sulfide (757 μl , 10.3 mmol) to provide a complex. The complex was stirred at 0° C for additional for 30 minutes. The complex was cooled to and the solution of step-3 product (0.780 g, 0.825 mmol) dissolved in toluene or dichloromethane (10 ml) was added. The resulting reaction mixture was stirred at -40° C temperature for 2 hr and then allowed to warm at 25-30° C while stirring. Triethyl amine (1.14ml, 8.25 mmol) was added to the reaction mixture at 25-30° C and it was stirred for additional 30 minutes. The reaction mixture was poured in aqueous saturated sodium bicarbonate solution and the mixture was extracted with dichloromethane (3 x 25 ml). The combined separated organic layer was dried over Na_2SO_4 and was evaporated to provide residue. The residue was purified by using silica gel column chromatography (10 % acetone : hexanes) to provide step-4 product as a off white solid in 75% (0.582 g) yield. MS: m/z: 944.3 (M+1).

Step-5: A mixture of step-4 compound (0.582 g, 0.617 mmol) and 70% HF-Pyridine solution (19.3 μl , 0.678 mmol) in acetonitrile (10 ml), was stirred at 25-30° C temperature for 2 hr. The solvent was evaporated under vacuum to obtain a crude residue. To the crude residue, water (20 ml) was added and the mixture was extracted with dichloromethane (25 ml x 3). Combined separated organic layer was dried over Na_2SO_4 and evaporated under vacuum to obtain crude solid. The crude solid was purified by using silica gel column chromatography (CHCl_3 :MeOH; 90:10) to obtain title compound as white solid compound in 40% (0.230 g) yield., MS: m/z: 829.1 (M+1). Mp: 160-163° C

Following examples were prepared by using above procedure and by using Reagent A.



Example	T	Reagent (A)	Mp (° C)	Mass (M+1)
260	CH ₂ -(5-(pyridin-4-yl)[1,2,4]oxadiazol-3-yl)	4-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridine	151-155	829.1
261	CH ₂ -(5-(2-chloropyridin-3-yl)[1,2,4]oxadiazol-3-yl)	2-chloro-3-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridine	148-151	863.1
262	CH ₂ -(5-(3-fluorophenyl)[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-(3-fluorophenyl)-[1,2,4]-oxadiazole	150-153	845.1
263	CH ₂ -(5-(3-cyano phenyl)[1,2,4]oxadiazol-3-yl)	3-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-benzonitrile	184-188	852.1
264	CH ₂ -(5-(3-chlorophenyl)[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-(3-chlorophenyl)-[1,2,4]-oxadiazole	133-136	861.1
265	CH ₂ -(5-(pyridin-2-yl)[1,2,4]oxadiazol-3-yl)	3-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridine	164-166	829.1
266	CH ₂ -(5-(2-methoxy pyridin-3-yl)[1,2,4]oxadiazol-3-yl)	2-methoxy-3-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridine	123-125	859.1
267	CH ₂ -(5-(3-fluoro-4-methoxyphenyl)[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-(3-fluoro-4-methoxy-phenyl)-[1,2,4]-oxadiazole	161-165	875.1
268	CH ₂ -(5-(2-fluoropyridin-3-yl)[1,2,4]oxadiazol-3-yl)	2-fluoro-3-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridine	132-135	847.1
269	CH ₂ -(5-phenyl[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-phenyl-[1,2,4]-oxadiazole	189-192	827.1
270	CH ₂ -(5-methyl[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-methyl-[1,2,4]-oxadiazole	170-173	766.1
271	CH ₂ -(5-(pyridazin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyridazine	145-147	831.1

272	CH ₂ -(5-trifluoromethyl-[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-trifluoromethyl-[1,2,4]-oxadiazole	154-157	820.1
273	CH ₂ -(5-(6-methylpyridin-2-yl)[1,2,4]oxadiazol-3-yl)	2-(3-chloromethyl-[1,2,4]-oxadiazol-5-yl)-6-methylpyridine	194-197	843.1
274	CH ₂ -(5-(2-methyl-oxazol-4-yl)-[1,2,4]oxadiazol-3-yl)	3-chloromethyl-5-(2-methyl-oxazol-4-yl)-[1,2,4]-oxadiazole	136-140	833.1
275	CH ₂ -(5-cyclopropyl-[1,2,4]oxadiazol-3-yl)	3-Chloromethyl-5-cyclopropyl-[1,2,4]-oxadiazole	175-178	791.1
276	CH ₂ -(5-(5-methyl-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloro methyl-[1,2,4]-oxadiazol-5-yl)-5-methylpyridine	-	844.1
277	CH ₂ -(5-(5-cyclopropyl-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-cyclopropylpyridine	-	870.1
278	CH ₂ -(5-(5-cyano-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-cyano-pyridine	-	855.1
279	CH ₂ -(5-(5-dimethyl-amino-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-dimethylamino-pyridine	-	874.1
280	CH ₂ -(5-(5-methoxy-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-methoxy-pyridine	-	860.1
281	CH ₂ -(5-(5-fluoro-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-fluoro-pyridine	-	848.1
282	CH ₂ -(5-(5-chloro-pyridin-2-yl)-[1,2,4]oxadiazol-3-yl)	2-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-5-chloro-pyridine	-	864.1
283	CH ₂ -(5-(pyrimidin-5-yl)-[1,2,4]oxadiazol-3-yl)	5-(5-chloromethyl-[1,2,4]-oxadiazol-5-yl)-pyrimidine	-	831.1

Example-284: Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyridin-2-yl)\text{-isoxazol-5-yl})$:

Step-1: To the stirred solution of 2-pyridinecarboxaldehyde (2.00 ml, 18.69 mmol) in methanol (20 ml) was added sodium acetate (1.83 g, 22.44 mmol). To this reaction mixture hydroxylamine hydrochloride (1.51, 20.56 mmol) was added and stirred for 2 hour at 20-30° C temperature. The reaction mixture was evaporated under vacuum to provide a crude mass. To the crude mass was stirred with water (20 ml), to effect precipitation and the separated solid was filtered under suction, dried under vacuum to provide step-1 product in 85% (1.93 g) yield. MS: m/z 123 (M+1).


Step 2: To the stirred solution of step-1 product pyridine-2-carbaldehyde oxime (1.80 g, 14.75 mmol) in DMF (20 ml), was added N-chloro succinimide (2.55 g, 19.19 mmol) and the reaction mixture was stirred for 8 hour at 25-30° C temperature. Ice-water was added (20 ml) to the reaction mixture to effect precipitation and solid was filtered at suction and dried under vacuum to provide imidoyl chloride as step-2 product imidoyl chloride in 70% (1.61 g) yield. This was used directly for the next reaction. MS: m/z 157 (M+1).

Step-3: To the stirred solution of product obtained from step A of Example 185 (1.5 gm, 1.82 mmol) in toluene (15 ml) was added triethyl amine (0.50 ml, 3.64 mmol). N-hydroxypyridine-2-carboximidoyl chloride (0.42 g, 2.73 mmol) was added to the reaction mixture and heated at 90° C temperature for 12 hr. The reaction mixture was poured in aqueous saturated ammonium chloride solution (20 ml). The mixture was extracted with ethyl acetate (3 x 25 ml). Combined organic layer was dried over Na_2SO_4 and purified by using silica gel column chromatography (10 % Acetone:Hexane) to provide off white solid as a step-3 product in 78% (1.34 g) yield. MS: m/z: 945 (M+1).

Step-4: To the stirred solution of N-chlorosuccinimide (1.27 g, 9.53 mmol) in dichloromethane (15 ml) at 0° C was added dimethyl sulfide (1.16 ml, 15.88 mmol). The reaction mixture was stirred at 0° C for 30 min. The step-3 product (1.2 gm, 1.27 mmol) dissolved in dichloromethane (10 ml) was added to the reaction mixture at -40° C. The resulting reaction mixture was stirred at the same temperature for 2 hr and allowed to warm at 25-30° C temperature. Triethyl amine (1.76 ml, 12.71 mmol) was added and it was stirred for additional 30 min. The reaction mixture was poured in aqueous saturated sodium bicarbonate solution (100 ml) and the mixture was extracted with dichloromethane (3 x 25 ml). The combined organic layer was dried over Na_2SO_4 and purified by using silica gel column chromatography (10 % Acetone:Hexane) to provide step-4 product as a off white solid in 75% (0.9 g) yield. MS: m/z: 943 (M+1).

Step-5: The mixture of step-4 product (0.9 g, 0.90 mmol) and 70% HF-Pyridine solution (41 μ l, 1.43 mmol) dissolved in acetonitrile (10 ml) was stirred at 25-30° C for 2 hr under N₂ atmosphere. After completion of reaction, solvent was evaporated under vacuum to obtain crude product. Water (20 ml) was added to the crude product and extracted with dichloromethane (3 X 25 ml). Combined organic layer was dried over sodium sulphate and evaporated under vacuum to obtain crude mass. The crude mass was purified by using silica gel column chromatography (10 % MeOH:CHCl₃) to obtain title compound in 50% (0.39 g) yield. Mp: 198-201° C, MS: m/z: 829 (M+1).

Following examples were prepared by using above procedure and by using Reagent A.



Example	T	Reagent (A)	Mp (°C)	Mass (M+1)
285	CH ₂ -(3-(pyridin-3-yl)-isoxazol-5-yl)	N-hydroxypyridine-3-carboximidoyl chloride	172-174	828.1
286	CH ₂ -(3-(3-fluoro phenyl)-isoxazol-5-yl)	3-fluoro-N-hydroxy benzene-carbo ximidoyl chloride	146-148	845.1
287	CH ₂ -(3-(3-chloro phenyl)-isoxazol-5-yl)	3-chloro-N-hydroxy benzene-carbo ximidoyl chloride	176-180	861.1
288	CH ₂ -(3-(4-fluorophenyl)-isoxazol-5-yl)	4-fluoro-N-hydroxy benzene-carbo ximidoyl chloride	212-214	845.1
289	CH ₂ -(3-(3-methoxy phenyl)-isoxazol-5-yl)	N-hydroxy-3-methoxy-benzene-carboximidoyl chloride	217-221	857.1
290	CH ₂ -(3-phenyl-isoxazol-5-yl)	N-hydroxybenzene-carboximidoyl chloride	223-225	827.1
291	CH ₂ -(3-methyl-isoxazol-5-yl)	(1Z)-N-hydroxy ethanimidoyl chloride	185-189	765.1
292	CH ₂ -(3-(6-methoxy-pyridin-2-yl)-isoxazol-5-yl)	N-hydroxy-6-methoxypyridine-2-carboximidoyl chloride	180-182	858.1
293	CH ₂ -(3-(2-methoxy-pyridin-5-yl)-isoxazol-5-yl)	N-hydroxy-6-methoxypyridine-3-carboximidoyl chloride	184-186	858.1

294	$\text{CH}_2\text{-}(3\text{-}(5\text{-methyl}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	<i>N</i> -hydroxy-5-methyl pyridine-2-carboximidoyl chloride	-	843.1
295	$\text{CH}_2\text{-}(3\text{-}(5\text{-cyclopropyl}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	5-cyclopropyl- <i>N</i> -hydroxy-pyridine-2-carboximidoyl chloride	-	869.1
296	$\text{CH}_2\text{-}(3\text{-}(5\text{-cyano}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	5-cyano- <i>N</i> -hydroxy-pyridine-2-carboximidoyl chloride	-	854.1
297	$\text{CH}_2\text{-}(3\text{-}(5\text{-dimethyl}\text{amino}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	5-(dimethylamino)- <i>N</i> -hydroxy-pyridine-2-carboximidoyl chloride	-	873.1
298	$\text{CH}_2\text{-}(3\text{-}(5\text{-methoxy}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	<i>N</i> -hydroxy-5-methoxypyridine-2-carboximidoyl chloride	-	859.1
299	$\text{CH}_2\text{-}(3\text{-}(5\text{-fluoro}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	5-fluoro- <i>N</i> -hydroxy-pyridine-2-carboximidoyl chloride	-	847.1
300	$\text{CH}_2\text{-}(3\text{-}(5\text{-chloro}\text{-pyridin-2-yl})\text{-isoxazol-5-yl})$	5-chloro- <i>N</i> -hydroxy-pyridine-2-carboximidoyl chloride	-	863.1
301	$\text{CH}_2\text{-}(3\text{-}(pyrimidin-5-yl)\text{-isoxazol-5-yl})$	<i>N</i> -hydroxy-pyrimidine-5-carboximidoyl chloride	-	830.1

Example-302: Compound of Formula I-f where T is $\text{CH}_2\text{-}(5\text{-}(pyridin-3-yl)\text{-isoxazol-3-yl})$:


Step A: To the stirred solution of Example 7 (1.0 g, 1.3 mmol) in toluene (15 ml) is added potassium *tert*-butoxide (0.16 g, 1.43 mmol) and 18-crown-6-ether (0.048 g, 0.13 mmol) at 25-30 °C temperature. The reaction mixture is stirred for 10 min at 25-30 °C temperature. 2-(3-Chloromethyl-isoxazol-5-yl)-pyridine (0.28 mg, 1.43 mmol) is added to the reaction mixture and it is stirred for additional 1 hour. It is quenched by pouring it in aqueous saturated ammonium chloride solution (20 ml). The mixture is extracted with ethyl acetate (3 x 25 ml). Combined organic layer is dried over Na_2SO_4 and purified by using silica gel

column chromatography (10 % Acetone:Hexane) to provide step-1 product as a pale yellow solid in 48% (0.57 g) yield. MS: m/z :945.1 (M+1).

Step B: To the stirred solution of N-chlorosuccinimide (0.6 g, 4.5 mmol) in dichloromethane (15 ml) at 0° C is added dimethyl sulfide (0.48 ml, 7.5 mmol). The reaction mixture is stirred at 0° C for 30 min. The step-1 product (0.55 gm, 0.6 mmol) dissolved in dichloromethane (10 ml) is added to the reaction mixture at -40° C. The resulting reaction mixture is stirred at the same temperature for 2 hr and allowed to warm at 25-30° C temperature. Triethyl amine (0.84 ml, 6.0 mmol) is added and it is stirred for additional 30 min. The reaction mixture is poured in aqueous saturated sodium bicarbonate solution (100 ml) and the mixture is extracted with dichloromethane (3 x 25 ml). The combined organic layer is dried over Na₂SO₄ and purified by using silica gel column chromatography (10 % Acetone :Hexane) to provide step-4 product as a off white solid in 72% (0.4 g) yield. MS: m/z: 943 (M+1).

Step C: The mixture of step-4 product (0.4 g, 0.42 mmol) and 70% HF-Pyridine solution (18 µl, 0.64 mmol) dissolved in acetonitrile (10 ml) is stirred at 25-30° C for 2 hr under N₂ atmosphere. After completion of reaction, solvent is evaporated under vacuum to obtain crude product. Water (20 ml) is added to the crude product and extracted with dichloromethane (3 X 25 ml). Combined organic layer is dried over sodium sulphate and evaporated under vacuum to obtain crude mass. The crude mass is purified by using silica gel column chromatography (10 % MeOH:CHCl₃) to obtain title compound in 65% (0.25 g) yield. MS: m/z: 829 (M+1).

Following examples are prepared by using above procedure and by using Reagent A.



Example	T	Reagent A	Mass (M+1)
303	CH ₂ -(5-(3-fluorophenyl)-isoxazol-3-yl)	3-chloromethyl-5-(3-fluorophenyl)-isoxazole	845.1
304	CH ₂ -(5-(3-chlorophenyl)-	3-chloromethyl-5-(3-chloro-	861.1

	isoxazol-3-yl)	phenyl)-isoxazole	
305	CH ₂ -(5-(4-fluorophenyl)-isoxazol-3-yl)	3-chloromethyl-5-(4-fluorophenyl)-isoxazole	845.1
306	CH ₂ -(5-(3-methoxyphenyl)-isoxazol-3-yl)	3-chloromethyl-5-(3-methoxyphenyl)-isoxazole	857.1
307	CH ₂ -(5-phenyl)-isoxazol-3-yl)	3-chloromethyl-5-phenyl-isoxazole	827.1
308	CH ₂ -(5-methyl)-isoxazol-3-yl)	3-chloromethyl-5-methyl-isoxazole	765.1
309	CH ₂ -(5-(6-methoxy-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-6-methoxy-pyridine	858.1
310	CH ₂ -(5-(2-methoxy-pyridin-5-yl)-isoxazol-3-yl)	5-(3-chloromethyl-isoxazol-5-yl)-2-methoxy-pyridine	858.1
311	CH ₂ -(5-(5-methyl-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-methyl-pyridine	843.1
312	CH ₂ -(5-(5-cyclopropyl-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-cyclopropyl-pyridine	869.1
313	CH ₂ -(5-(5-cyano-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-cyano-pyridine	854.1
314	CH ₂ -(5-(5-dimethylamino-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-dimethylamino-pyridine	873.1
315	CH ₂ -(5-(5-methoxy-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-methoxy-pyridine	859.1
316	CH ₂ -(5-(5-fluoro-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-fluoro-pyridine	847.1
317	CH ₂ -(5-(5-chloro-pyridin-2-yl)-isoxazol-3-yl)	2-(3-chloromethyl-isoxazol-5-yl)-5-chloro-pyridine	863.1
318	CH ₂ -(pyrimidin-5-yl)-isoxazol-3-yl)	5-(3-chloromethyl-isoxazol-5-yl)-pyrimidine	830.1
319	CH ₂ CH ₂ -(4-(pyridin-2-yl)-1H-imidazol-1-yl)	2-[1-(2-Chloro-ethyl)-1H-imidazol-4-yl]-pyridine	842.1

320	CH ₂ CH ₂ -(4-(pyridin-3-yl)-1H-imidazol-1-yl)	3-[1-(2-Chloro-ethyl)-1H-imidazol-4-yl]-pyridine	842.1
321	CH ₂ CH ₂ -(4-(6-methyl-pyridin-2-yl)-1H-imidazol-1-yl)	2-[1-(2-Chloro-ethyl)-1H-imidazol-4-yl]-6-methyl-pyridine	856.1
322	CH ₂ CH ₂ -(4-(4-methoxy-pyridin-2-yl)-1H-imidazol-1-yl)	2-[1-(2-Chloro-ethyl)-1H-imidazol-4-yl]-4-methoxy-pyridine	872.1
323	CH ₂ -(4-(pyridin-2-yl)-pyrazol-4-yl)	2-(4-Chloromethyl-pyrazol-1-yl)-pyridine	828.1
324	CH ₂ -(4-(pyridin-3-yl)-pyrazol-4-yl)	3-(4-Chloromethyl-pyrazol-1-yl)-pyridine	828.1
325	CH ₂ -(4-(6-methyl-pyridin-2-yl)-pyrazol-4-yl)	2-(4-Chloromethyl-pyrazol-1-yl)-6-methyl-pyridine	842.1
326	CH ₂ -(4-(4-methoxy-pyridin-2-yl)-pyrazol-4-yl)	2-(4-Chloromethyl-pyrazol-1-yl)-4-methoxy-pyridine	858.1
327	CH ₂ -(4-(5-methoxy-pyridin-2-yl)-pyrazol-4-yl)	2-(4-Chloromethyl-pyrazol-1-yl)-5-methoxy-pyridine	858.1
328	CH ₂ -(1-(pyridin-2-yl)-1H-imidazol-4-yl)	2-(4-Chloromethyl-imidazol-1-yl)-pyridine	828.1
329	CH ₂ -(1-(pyridin-3-yl)-1H-imidazol-4-yl)	3-(4-Chloromethyl-imidazol-1-yl)-pyridine	828.1
330	CH ₂ -(1-(6-methyl-pyridin-2-yl)-1H-imidazol-4-yl)	2-(4-Chloromethyl-imidazol-1-yl)-6-methyl-pyridine	842.1
331	CH ₂ -(1-(4-methoxy-pyridin-2-yl)-1H-imidazol-4-yl)	2-(4-Chloromethyl-imidazol-1-yl)-4-methoxy-pyridine	858.1
332	CH ₂ -(1-(5-methoxy-pyridin-2-yl)-1H-imidazol-4-yl)	2-(4-Chloromethyl-imidazol-1-yl)-5-methoxy-pyridine	858.1
333	CH ₂ -(2-(pyridin-2-yl)-oxazol-4-yl)	2-(4-Chloromethyl-oxazol-2-yl)-pyridine	829.1

334	CH ₂ -(2-(pyridin-3-yl)-oxazol-4-yl)	3-(4-Chloromethyl-oxazol-2-yl)-pyridine	829.1
335	CH ₂ -(2-(6-methyl-pyridin-2-yl)-oxazol-4-yl)	2-(4-Chloromethyl-oxazol-2-yl)-6-methyl-pyridine	843.1
336	CH ₂ -(2-(4-methoxy-pyridin-2-yl)-oxazol-4-yl)	2-(4-Chloromethyl-oxazol-2-yl)-4-methoxy-pyridine	859.1
337	CH ₂ -(2-(5-methoxy-pyridin-2-yl)-oxazol-4-yl)	2-(4-Chloromethyl-oxazol-2-yl)-5-methoxy-pyridine	859.1

Example-338: Compound of Formula I-f where Q is CN;

A mixture of Example 8 (0.95 g, 1.3 mmol) and 70% hydrogen fluoride in pyridine (50 μ l, 1.8 mmol) in acetonitrile (15 ml) was stirred under inert atmosphere at a temperature between 25-30 °C for 3 h. The solvent was evaporated under vacuum and the residue obtained was dissolved in ethyl acetate (25 ml) and washed with water (2 x 10ml). The combined organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to provide crude solid which on trituration in diethyl ether afforded pale yellow solid as the title compound in 60% (0.58 g) yield. MP: 230-232 °C, Mass: m/z: 638.1 (M+1).


Example-339: Compound of Formula I-f where T is H;

A mixture of Example 9 (0.95 g, 1.2 mmol) and 70% hydrogen fluoride in pyridine (50 μ l, 1.8 mmol) in acetonitrile (15 ml) was stirred under inert atmosphere at a temperature between 25-30 °C for 15 h. The solvent was evaporated under vacuum and the residue obtained was dissolved in ethyl acetate (25 ml) and washed with water (2 x 10ml). The combined organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to provide crude solid which on trituration in diethyl ether afforded pale yellow solid as the title compound in 75% (0.7 g) yield. MP: 174-176 °C, Mass: m/z: 671.1 (M+1).

Example-340: Compound of Formula I-f where T is (CH₂)₂CH₃;

To a stirring solution of Example 179 (0.3 gm, 0.37mm) in THF (10 ml) was added 10 % Pd/C (30 mg, 10 %) and stirred under H₂ atmosphere for 15 hr. The reaction mixture was filtered through celite and concentrated under reduced pressure. Crude solid was triturated in diethyl ether affording white solid. Yield: 250 mg (84 %), MP: 182-184 °C, Mass: m/z: 713.1(M+H).

Following examples were prepared as per the procedure mentioned above by using the corresponding unsaturated starting material.




Example	Starting material	T	Mp (° C)	Mass (M+1)
341	Example 205	CH ₂ CH ₂ CH(CH ₃) ₂	244-246	741.1
342	Example 208	CH ₂ (CH ₂) ₂ Ph	172-174	789.1

While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.

Claims

1. Compounds having the structure of Formula I:



Formula I

and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers,

wherein


* indicates a chiral center;

R₁ is hydrogen or methyl;

R₂ is hydrogen or hydroxyl protecting group,


wherein hydroxyl protecting groups are selected from the group consisting of triethylsilyl, trimethylsilyl, acetyl, benzoyl, methoxymethyl, benzyl, methoxyethoxymethyl or tertbutyldimethylsilyl;

Q is Het or



wherein

Het is selected from



wherein R₃ is one or more substituent selected from the group consisting of NO₂, CN, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, CH₂CONH₂, CH₂CO₂Et,

CH₂CN, CH₂CH₂OH, CH₂OCH₂CH₂OCH₃, NH₂, substituted C₁-C₆ alkyl and substituted C₂-C₆ alkenyl;

T is selected from the group consisting of:

hydrogen, -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -A-(CH₂)_m-R₅, -A-(CH₂)_m-CH=CH-R₅, -A-(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -A-(CH₂)_m-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-B-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, -A-(CH₂)_m-X-R₆, -A-(CH₂)_m-B-X-R₆, -A-(CH₂)_m-CH=CH-X-R₆ and -A-(CH₂)_m-C≡C-X-R₆;

wherein,

m is 0, 1, 2 or 3;

R₅ is selected from the group consisting of:


hydrogen, cyano, halogen, hydroxyl, CO₂(C₁-C₆ alkyl), CO NR_aR_b, NR_aR_b, C₁-C₆ alkyl or substituted C₁-C₆ alkyl, C₂-C₆ alkenyl or substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl, 6 membered substituted or unsubstituted aryl, 5-6 membered substituted or unsubstituted heteroaryl, 3-6 membered substituted or unsubstituted cycloalkyl and 3-6 membered substituted or unsubstituted heterocyclyl;

wherein R_a and R_b are independently hydrogen, C₁-C₆ alkyl or R_a and R_b together with the nitrogen to which they are attached form a 5-6 membered heterocyclic ring wherein the heterocycle has one or more heteroatoms selected from N, O, S;

A is -CO- or -CONH-;

B is -O-, -S-, -SO-, -SO₂-, -CO-, -CONH-, -CON(CH₃)-, -NHCONH-, -

C(NH₂)=N-O-,




wherein

R₈ is a substituent at any one of carbon of the heterocycle, R₈ is selected from the group comprising of hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl and substituted C₂-C₆ alkenyl;

X is a 6 membered aryl or a 5-6 membered heteroaryl;

R₆ is aryl, heteroaryl, substituted aryl or substituted heteroaryl;

Y' is hydrogen and Y is OR₇, wherein R₇ is hydrogen or




wherein R_{2'} is hydrogen, (when R_{2'} is hydrogen, R₇ is designated as cladinosine) or hydroxyl protecting group,

wherein hydroxyl protecting groups are selected from the group consisting of triethylsilyl, trimethylsilyl, acetyl, benzoyl, methoxymethyl, methoxyethoxymethyl, benzyl or tertbutyldimethylsilyl;

or

Y and Y' together with the carbon to which they are attached form C=O.

2. Compounds having the structure of Formula I-a:




and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers,

wherein

* indicates a chiral center;


R₁ is hydrogen or methyl;

Q is Het or



wherein

Het is selected from



wherein R₃ is one or more substituent selected from the group consisting of NO₂, CN, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, CH₂CONH₂, CH₂CO₂Et, CH₂CN, CH₂CH₂OH, CH₂OCH₂CH₂OCH₃, NH₂, substituted C₁-C₆ alkyl and substituted C₂-C₆ alkenyl;

T is selected from the group consisting of:

hydrogen, -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -A-(CH₂)_m-R₅, -A-(CH₂)_m-CH=CH-R₅, -A-(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -A-(CH₂)_m-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-B-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, -A-(CH₂)_m-X-R₆, -A-(CH₂)_m-B-X-R₆, -A-(CH₂)_m-CH=CH-X-R₆ and -A-(CH₂)_m-C≡C-X-R₆;

wherein

m is 0, 1, 2 or 3;

R₅ is selected from the group consisting of:

hydrogen, cyano, halogen, hydroxyl, CO₂(C₁-C₆ alkyl), CONR_aR_b, NR_aR_b, C₁-C₆ alkyl or substituted C₁-C₆ alkyl, C₂-C₆ alkenyl or substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl, 6 membered substituted or unsubstituted aryl, 5-6 membered substituted or unsubstituted heteroaryl, 3-6 membered substituted or unsubstituted cycloalkyl and 3-6 membered substituted or unsubstituted heterocyclyl;


wherein R_a and R_b are independently hydrogen, C₁-C₆ alkyl or

R_a and R_b together with the nitrogen to which they are attached form a 5-6 membered heterocyclic ring wherein the heterocycle has one or more heteroatoms selected from N, O, S;

A is CO or CONH;

B is selected from the group consisting of:

O, S, SO, SO₂, CO, CONH, CON(CH₃), -NHCONH-, -C(NH₂)=N-O-,




wherein

R₈ is a substituent at any one of carbon of the heterocycle, R₈ is selected from the group comprising of hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl and substituted C₂-C₆ alkenyl;

X is a 6 membered aryl or a 5-6 membered heteroaryl;


R₆ is aryl, heteroaryl, substituted aryl or substituted heteroaryl.

3. The compound of claim 2, wherein R₁ is CH₃ and Het is



wherein R₃ is one or more substituent selected from C₁-C₆ alkyl, C₂-C₆ alkenyl, CH₂CONH₂, CH₂CO₂(C₁-C₆ alkyl), CH₂CN, CH₂CH₂OH, CH₂OCH₂CH₂OCH₃.

4. The compound of claim 2, wherein Q is



wherein T is as defined in claim 2.

5. The compound of claim 4, wherein T is selected from

-(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -A-(CH₂)_m-R₅, -A-(CH₂)_m-CH=CH-R₅, -A-(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -A-(CH₂)_m-B-R₅,

wherein

m is 1 or 2; A is $-\text{CO}-$; B is piperazinyl;

R_5 is substituted/unsubstituted phenyl or heteroaryl;

wherein heteroaryl is selected from pyridin-4-yl, pyridin-3-yl, pyridin-2-yl, pyrimidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, oxazolyl or isoxazolyl;


phenyl or heteroaryl may be substituted with one or more substituents selected from the group comprising of $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, cyano, halogen, dimethylamino.

6. The compound of claim 2, wherein T is selected from

$-(\text{CH}_2)_m\text{-X-R}_6$, $-(\text{CH}_2)_m\text{-CH=CH-X-R}_6$, $-(\text{CH}_2)_m\text{-C}\equiv\text{C-X-R}_6$, $-\text{A-}(\text{CH}_2)_m\text{-X-R}_6$,
 $-\text{A-}(\text{CH}_2)_m\text{-CH=CH-X-R}_6$, $-\text{A-}(\text{CH}_2)_m\text{-C}\equiv\text{C-X-R}_6$

wherein R_6 , A, X and m are as defined in claim 2.


7. The compound of claim 6, wherein X- R_6 is



wherein R_6 is phenyl, substituted phenyl, heteroaryl or substituted heteroaryl; wherein heteroaryl is selected from one or more of pyridin-4-yl, pyridin-3-yl, pyridin-2-yl, pyrimidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, oxazolyl or isoxazolyl; wherein the phenyl or heteroaryl is substituted with one or more substituents selected from the group comprising of $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, cyano, halogen and dimethylamino.

8. The compound of claim 2, wherein T is $-(\text{CH}_2)_m\text{-X-R}_6$, wherein R_6 , X and m are as defined in claim 2.

9. The compound of claim 8, wherein X- R_6 is




wherein R₆ is phenyl, substituted phenyl, heteroaryl or substituted heteroaryl.

10. The compound of claim 9, wherein heteroaryl is pyridin-4-yl, pyridin-3-yl, pyridin-2-yl, pyrimidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, oxazolyl or isoxazolyl.

11. The compound of claim 9, wherein the phenyl or heteroaryl is substituted with one or more substituents selected from the group comprising of C₁-C₆ alkyl, C₁-C₆ alkoxy, cyano, halogen, dimethylamino.

12. Compounds having the structure of Formula I-d,




Formula I-d

and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers,
wherein

* indicates a chiral center;


R₁ is CH₃;

Q is Het or



wherein

Het is selected from



wherein R₃ is one or more substituent selected from the group consisting of NO₂, CN, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, CH₂CONH₂, CH₂CO₂Et, CH₂CN, CH₂CH₂OH, CH₂OCH₂CH₂OCH₃, NH₂, substituted C₁-C₆ alkyl and substituted C₂-C₆ alkenyl;

T is selected from the group consisting of:

hydrogen, -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -A-(CH₂)_m-R₅, -A-(CH₂)_m-CH=CH-R₅, -A-(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -A-(CH₂)_m-B-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-B-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, -A-(CH₂)_m-X-R₆, -A-(CH₂)_m-B-X-R₆, -A-(CH₂)_m-CH=CH-X-R₆ and -A-(CH₂)_m-C≡C-X-R₆;

wherein,

m is 0, 1, 2 or 3;

R₅ is selected from the group consisting of:

hydrogen, cyano, halogen, hydroxyl, CO₂(C₁-C₆ alkyl), CO NR_aR_b, NR_aR_b, C₁-C₆ alkyl or substituted C₁-C₆ alkyl, C₂-C₆ alkenyl or substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl, 6 membered substituted or unsubstituted aryl, 5-6 membered substituted or unsubstituted heteroaryl, 3-6 membered substituted or unsubstituted cycloalkyl and 3-6 membered substituted or unsubstituted heterocyclyl;


wherein R_a and R_b are independently hydrogen, C₁-C₆ alkyl or

R_a and R_b together with the nitrogen to which they are attached form a 5-6 membered heterocyclic ring wherein the heterocycle has one or more heteroatoms selected from N, O, S;

A is CO or CONH;

B is selected from the group consisting of:

O, S, SO, SO₂, CO, CONH, CON(CH₃), -NHCONH-, -C(NH₂)=N-O-,




wherein

R₈ is a substituent at any one of carbon of the heterocycle, R₈ is selected from the group comprising of hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl and substituted C₂-C₆ alkenyl;

X is a 6 membered aryl or a 5-6 membered heteroaryl;


R₆ is aryl, heteroaryl, substituted aryl or substituted heteroaryl.

13. The compound of claim 12, wherein R₁ is CH₃ and Het is



wherein R₃ is one or more substituent selected from C₁-C₆ alkyl, C₂-C₆ alkenyl, CH₂CONH₂, CH₂CO₂(C₁-C₆ alkyl), CH₂CN, CH₂CH₂OH, CH₂OCH₂CH₂OCH₃.

14. The compound of claim 12, wherein Q is



wherein T is as defined in claim 12.

15. The compound of claim 12, wherein T is selected from

$-(CH_2)_m-R_5$, $-(CH_2)_m-CH=CH-R_5$, $-(CH_2)_m-C\equiv C-R_5$, $-A-(CH_2)_m-R_5$, $-A-(CH_2)_m-$
 $CH=CH-R_5$, $-A-(CH_2)_m-C\equiv C-R_5$, $-(CH_2)_m-B-R_5$, $-A-(CH_2)_m-B-R_5$,


wherein

m is 1 or 2; A is $-CO-$; B is piperazinyl; R_5 is substituted or unsubstituted phenyl or heteroaryl; wherein heteroaryl is selected from pyridin-4-yl, pyridin-3-yl, pyridin-2-yl, pyrimidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, oxazolyl or isoxazolyl; wherein the phenyl or heteroaryl is substituted with one or more substituents selected from the group comprising of C_1-C_6 alkyl, C_1-C_6 alkoxy, cyano, halogen and dimethylamino.

16. The compound of claim 12, wherein T is selected from

$-(CH_2)_m-X-R_6$, $-(CH_2)_m-B-X-R_6$, $-(CH_2)_m-CH=CH-X-R_6$, $-(CH_2)_m-C\equiv C-X-R_6$, $-A-$
 $(CH_2)_m-X-R_6$, $A-(CH_2)_m-B-X-R_6$, $-A-(CH_2)_m-CH=CH-X-R_6$, $-A-(CH_2)_m-C\equiv C-X-R_6$
 wherein R_6 , A , B , X and m are as defined in claim 12.


17. The compound of claim 16, wherein $X-R_6$ is



wherein R_6 is phenyl, substituted phenyl, heteroaryl or substituted heteroaryl; wherein heteroaryl is selected from one or more of pyridin-4-yl, pyridin-3-yl, pyridin-2-yl, pyrimidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, oxazolyl or isoxazolyl;

wherein the phenyl or heteroaryl is substituted with one or more substituents selected from the group comprising of C_1-C_6 alkyl, C_1-C_6 alkoxy, cyano, halogen and dimethylamino.

18. A compound of Formula I-c, wherein the compound is selected from the group consisting of:



Compound of Formula I-c, where T is $\text{CH}_2\text{CH}_2\text{-(1-(3-methoxy)phenyl-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{COCH=CH-(pyridin-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{COCH=CH-(pyridin-3-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(pyridin-3-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(3-methoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(4-methoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(4-dimethylamino)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(3-cyano)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{COCH}_2\text{O-(3-chloro)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(3,5-dimethoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(2,3-dimethoxy)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(3-fluoro)phenyl}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(4-(2-methylphenyl)-piperazinyl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(4-(phenyl)piperazinyl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(4-methoxyphenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(pyridin-3-yl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(4-fluorophenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-(1-(3-methylphenyl)-1H-[1,2,3]triazol-4-yl)}$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $CO(CH_2)_2-(1-(3\text{-chlorophenyl})-1H-[1,2,3]\text{triazol-4-yl})$, R₁ is CH₃;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-bromophenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(3\text{-fluorophenyl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(\text{pyridin-4-yl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-}(\text{pyridin-3-yl})\text{-}[1,2,4]\text{oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}[3\text{-(4-methoxyphenyl)}\text{-}[1,2,4]\text{oxadiazol-5-yl}]$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(2\text{-methoxy)phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3\text{-fluorophenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;


Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-phenyl-[1,2,4]oxadiazol-5-yl})$, R_1 is CH_3 ;

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(3\text{-naphthalen-2-yl-[1,2,4]oxadiazol-5-yl})$, R₁ is CH₃

and

Compound of Formula I-c, where T is $\text{CO}(\text{CH}_2)_2\text{-}(1\text{-}(3,5\text{-difluoro-phenyl})\text{-}1\text{H-[1,2,3]triazol-4-yl})$, R_1 is CH_3 ;

19. A compound of Formula I-f, wherein the compound is selected from the group consisting of:



Formula I-f

Compound of Formula I-f, where T is $\text{CH}_2\text{C}(\text{F})=\text{CH}_2$;

Compound of Formula I-f, where T is $\text{CH}_2\text{CH}\equiv\text{C}-(2\text{-fluoropyridin-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2\text{CH}\equiv\text{C}-(\text{pyridin-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-chlorophenyl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(2\text{-fluorophenyl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(4\text{-}(1\text{H}-[1,2,4]\text{-triazol)phenyl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-phenyl}[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{pyridin-3-yl})[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{3-fluorophenyl})[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{pyridin-2-yl})[1,3,4]\text{oxadiazol-2-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-phenyl}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{3-fluorophenyl})[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{4-chlorophenyl})[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{4-methoxyphenyl})[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{pyridin-3-yl})[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{pyridin-2-yl})[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-cyclopropyl}[1,2,4]\text{oxadiazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-phenyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{pyridin-2-yl})[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{2-fluoropyridin-3-yl})[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-cyclopropyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-methyl}[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{pyridazin-2-yl})-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(5\text{-}(\text{2-methyl-thiazol-4-yl})-[1,2,4]\text{oxadiazol-3-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{3-fluorophenyl})\text{-isoxazol-5-yl})$; Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{3-chlorophenyl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{4-fluorophenyl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{pyridin-2-yl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-phenyl-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{5-methyl-pyridin-2-yl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{5-cyclopropyl-pyridin-2-yl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2-(3\text{-}(\text{5-cyano-pyridin-2-yl})\text{-isoxazol-5-yl})$;

Compound of Formula I-f, where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-dimethylamino-pyridin-2-yl)\text{-}isoxazol-5-yl})$;
Compound of Formula I-f, where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-methoxy-pyridin-2-yl)\text{-}isoxazol-5-yl})$;
Compound of Formula I-f, where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-fluoro-pyridin-2-yl)\text{-}isoxazol-5-yl})$;
Compound of Formula I-f, where T is $\text{CH}_2\text{-}(3\text{-}(5\text{-chloro-pyridin-2-yl)\text{-}isoxazol-5-yl})$ and
Compound of Formula I-f where T is $\text{CH}_2\text{-}(3\text{-}(pyrimidin-5-yl)\text{-}isoxazol-5-yl)$.

20. A pharmaceutical composition comprising a therapeutically effective amount of a compound as defined in claim 1, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

21. A pharmaceutical composition comprising a therapeutically effective amount of a compound as defined in claim 2, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

22. A pharmaceutical composition comprising a therapeutically effective amount of a compound as defined in claim 12, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

23. A method for treating or preventing microbial infections in a human or an animal, comprising administering to said animal or human, a therapeutically effective amount of a compound of claim 1.

24. A method for treating or preventing microbial infections in a human or an animal, comprising administering to said animal or human, a therapeutically effective amount of a compound of claim 2.

25. A method for treating or preventing microbial infections in a human or an animal, comprising administering to said animal or human, a therapeutically effective amount of a compound of claim 12.


26. The method according to claims 23, 24 or 25 wherein the microbial infections are caused by Gram-positive, Gram-negative bacteria, aerobic, anaerobic bacteria or atypical bacteria.

27. The method for treating or preventing microbial infections in an animal or a human comprising administering to said animal or human, a pharmaceutical composition according to claims 20, 21 or 22.

28. A process for preparing a compound of Formula I as defined in claim 1 or its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, polymorphs, enantiomers or diastereomers, the process comprising one or more of the following steps of:

a) treating the compound of Formula VIIa or VIIb with a suitable carboxylic acid reagent of the Formula R₉-COOH,


wherein R₉ represents -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, in presence of esterifying agent to give corresponding compound of Formula XIa or XIb, wherein * is a chiral centre, and subjecting the resulting compound to one or more of the following reactions:



VIIa, Y' is H, Y is Ocladinose, R2, R2' is TES
 VIIb, Y' is H, Y is OH, R2, is TES


XIa, Y' is H, Y is Ocladinose, R2, R2' is TES
 XIb, Y' is H, Y is OH, R2, is TES

i) deprotecting the hydroxyl protecting groups of compound of Formula XIa to give the corresponding macrolide of Formula XIIa;




or

ii) oxidizing the compound of Formula XIb to give the 3-ketone derivative of Formula XIII;




iii) deprotecting the hydroxyl protecting groups to give the corresponding ketolide of Formula XIIb;

b) treating the compound of Formula VIIa or VIIb with a suitable isocyanate reagent of Formula $R_9-N=C=O$, wherein R_9 represents $-(CH_2)_m-R_5$, $-(CH_2)_m-CH=CH-R_5$, $-(CH_2)_m-C\equiv C-R_5$, $-(CH_2)_m-B-R_5$, $-(CH_2)_m-CH=CH-R_5$, $-(CH_2)_m-C\equiv C-R_5$, $-(CH_2)_m-X-R_6$, $-(CH_2)_m-CH=CH-X-R_6$, $-(CH_2)_m-C\equiv C-X-R_6$, wherein R_5 , R_6 , X , B and m are as defined above, to give compound of Formula XIVa or XIVb, wherein * is a chiral centre, and subjecting the resulting compound to one or more of the following reactions:




i) deprotecting the hydroxyl protecting groups of compound of Formula XIVa to give the corresponding macrolide of Formula XVa;




or

ii) converting the 3-hydroxyl derivative XIVb to 3-ketone derivative of Formula XVI;



iii) deprotecting the hydroxyl protecting groups of XVI to give the corresponding ketolide of Formula XVb;


c) treating the compound of Formula VIIa or VIIb with a suitable alkylating reagent of Formula halo-R₉, wherein R₉ represents -(CH₂)_m-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-B-R₅, -(CH₂)_m-CH=CH-R₅, -(CH₂)_m-C≡C-R₅, -(CH₂)_m-X-R₆, -(CH₂)_m-CH=CH-X-R₆, -(CH₂)_m-C≡C-X-R₆, wherein B, X, R₅, R₆ and m are as defined above and halo is a chlorine, bromine or iodine, in presence of base to give compound of Formula XVIa or XVIb, wherein * is a chiral centre, and subjecting the resulting compound XVIa or XVIb, to one or more of the following reactions:



VIIa, Y' is H, Y is Ocladinose, R2, R2' is TES
 VIIb, Y' is H, Y is OH, R2, is TES


XVIa, Y' is H, Y is Ocladinose, R2, R2' is TES
 XVIb, Y' is H, Y is OH, R2, is TES

iv) deprotecting the hydroxyl protecting groups of Formula XVIa to give the corresponding macrolide of Formula XVIIa;



or


v) converting the 3-hydroxyl derivative of Formula XVIa to 3-oxo derivative of Formula XVIIb';



iii) deprotecting the hydroxyl protecting groups of compound of Formula XVIIb' to give the corresponding ketolide of Formula XVIIb.

d) treating the compound of Formula XVIIa or XVIIb with a suitable halo-aryl reagent of Formula halo-R₅ or halo-X-R₆ in the presence of suitable palladium catalyst and

phosphine derivative under heck coupling reaction conditions to give compound of Formula XIXa or XIXb,




XVIIIa, Y' is H, Y is Ocladinose, R2,
R2' is TES
XVIIIb, Y' is H, Y is OH, R2, is TES

XIXa, Y' is H, Y is Ocladinose, R2,
R2' is TES
XIXb, Y' is H, Y is OH, R2, is TES


wherein G is $-(CH_2)_m-$, $-A-(CH_2)_m-$; U is $-R_5$, $-X-R_6$ and * is a chiral centre;
and subjecting the resulting compound to one or more of the following reactions:

- deprotecting the hydroxyl protecting groups to give the corresponding macrolide of Formula XXa;




or

- converting the 3-hydroxyl derivative XIXb to 3-ketone of Formula XXb';



- deprotecting the hydroxyl protecting groups of Formula XXb' to give the corresponding ketolide of Formula XXb;

- treating the compound of Formula XXIa or XXIb with a suitable halo-aryl reagent in the presence of suitable palladium catalyst under sonagashira coupling reaction conditions to give corresponding compound of Formula XXIIa or XXIIb,




XXIIa, Y' is H, Y is Ocladinose, R2,
R2' is TES
XXIIb, Y' is H, Y is OH, R2, is TES

XXIIla, Y' is H, Y is Ocladinose, R2,
R2' is TES
XXIIlb, Y' is H, Y is OH, R2, is TES


wherein G is $-(CH_2)_m-$, $-A-(CH_2)_m-$; U is $-R_5$, $-X-R_6$ and * is a chiral centre; and subjecting the resulting compound of Formula XXIIa or XXIIb to one or more of the following reactions:

- deprotecting the hydroxyl protecting groups to give the corresponding macrolide of Formula XXIIla,



or


- converting the 3-hydroxyl derivative of Formula XXIIb to 3-ketone derivative of Formula XXIV,




- deprotecting the hydroxyl protecting groups of XXIV to give the corresponding ketolide of Formula XXIIlb;

f) Optionally, hydrogenating the compound obtained by any of the above steps to obtain the corresponding saturated compound.

29. The process of claim 28, wherein the process of preparation of compound of Formula VII, comprises treating a compound of Formula VI with hydroxylamine in a suitable solvent, wherein R_1 , R_2 , Y , Y' , are as defined in claim 1.



30. The process of claim 29, wherein the process of preparation of compound of Formula VI comprises the steps of:



- a) treating a compound of Formula IV with cyanoacetic acid in presence of a base to give compound of Formula Va
- b) treating the compound of Formula Va with a base to give compound of Formula VI wherein R_1 , R_2 , R_2' are as defined in claim 1.