



US 20080081020A1

(19) **United States**

(12) **Patent Application Publication**
Huang et al.

(10) **Pub. No.: US 2008/0081020 A1**

(43) **Pub. Date: Apr. 3, 2008**

(54) **COLOR CHANGE SURGICAL PREP SOLUTION**

(52) **U.S. Cl. 424/9.6; 424/405**

(76) Inventors: **Yeong H. Huang**, Grayslake, IL (US); **Nancy Barot**, Mundelein, IL (US)

(57) **ABSTRACT**

Correspondence Address:
Allegiance Corporation
Attn: Kim Luna
1430 Waukegan Road
McGaw Park, IL 60085-6787

This invention discloses a surgical prep solution formulation, either film forming or non-film forming, that changes color upon a change in pH to indicate that the solvent has sufficiently evaporated. The major components of this surgical prep solution include: antimicrobial agents, solvents, pH adjusters, color changing pH indicators, and optionally: film forming polymers, dyes and viscosity regulators. The pH of the solution may rise or fall depending on the pH of the original solution as compared with the normal pH range of the acid mantle of the human skin or the normal pH range of a sterilizing solution. As the pH changes, the pH indicator may cause the surgical prep solution to change color. Concurrently, as the fluids evaporate, the pH indicators interact with atmospheric CO₂, causing the solution to change color based on the pH of the solution. The solution may also be used as coating for medical devices.

(21) Appl. No.: **11/542,791**

(22) Filed: **Oct. 3, 2006**

Publication Classification

(51) **Int. Cl.**
A61K 49/00 (2006.01)
A01N 25/00 (2006.01)

COLOR CHANGE SURGICAL PREP SOLUTION

FIELD OF THE INVENTION

[0001] The invention relates to disinfectants for use in medical procedures that breach the skin, such as surgery, surgical procedures and intravenous injections. In particular, the invention pertains to surgical prep solutions used to disinfect a selected surface, for example, pre-operative surgical sites and medical devices.

BACKGROUND OF THE INVENTION

[0002] Pre-operative preparation of the skin with a topical antimicrobial agent is necessary to reduce the likelihood that the patient will contract a hospital-acquired infection during a surgery or surgical procedure. Typically, the healthcare practitioners, for example, prep nurses, apply a topical antimicrobial agent to a surgical or needle entry site before the procedure. Similarly, it is essential that medical devices that breach the skin be disinfected prior to penetrating the skin at an entry site or accessing an intravenous system. Healthcare practitioners typically disinfect these medical devices by applying an antimicrobial solution, e.g., alcohol, prior to use. Such treatment reduces the infection rate at the site or within the blood stream by hindering the growth of microorganisms or disinfecting a wound, surgical incision, or needle puncture site.

[0003] The human skin is normally covered with microorganisms such as bacteria, fungi, and viruses. The microorganisms are either transient, found on the skin's surface or are resident in deeper areas of the skin, such as the hair follicles.

[0004] The microorganisms are prevented from entering the body by the physical barrier of skin and the acid mantle, a protective layer formed by the mixture of sweat and sebum. The acid mantle maintains a moderately acidic covering for the normal human skin. The pH means and ranges for the normal skin of both adults and neonates are 5.7 ± 0.16 (range: 4.5-6.7) and 7.08 ± 0.17 (range: 6.6-7.5), respectively.

[0005] The function of the acid mantle is not completely understood even though the acidic environment of the skin's surface has been recognized for a century. However, the acid mantle is believed to assist the body in processing the lipids required for the skin to function as a barrier. Acidic environments also generally hinder bacterial growth, and hence, help resist bacterial infection.

[0006] Potential pathogens usually cannot enter the body if the skin and acid mantle remain intact. However, the integrity of the skin barrier is breached during surgical procedures, potentially exposing the surgical site or wound to microorganisms, increasing a patient's risk of infection. Thus, standard surgical procedures require that the skin at the surgical site be disinfected prior to surgery to reduce the patient's risk of infection. Therefore, there is a need in the art for a topical agent that will both kill the transient and resident microorganisms quickly and provide sustained antimicrobial activity throughout the entire surgery or surgical procedure.

[0007] Healthcare practitioners have long used ethanol or isopropanol, either alone or as a solvent along with other antimicrobial agents, to disinfect the skin at the surgical incision or needle puncture site and medical devices that

penetrate the skin since these alcohols quickly reduce the population of bacteria, fungi, and some viruses at the site. Alcohol also provides rapid and sustained antimicrobial activity when it is combined with antimicrobial agents. However, alcohol based surgical prep solutions are flammable and certain surgical procedures cannot begin until the alcohol is completely evaporated. Hospital fires have been caused by unevaporated flammable solvents within surgical prep solutions. For the patient's safety, it is critical that healthcare practitioners be able to determine whether the flammable solvents within surgical prep solutions are completely evaporated prior to initiating energized surgical procedures, for example, electrocautery.

[0008] Solvent evaporation in particular presents a major challenge for healthcare practitioners for several reasons. First, it is difficult to visually determine when a colorless solvent (e.g., alcohol) has sufficiently evaporated to begin surgery or the surgical procedure. Second, the healthcare practitioners will often physically touch the skin surface or the medical device with a glove to determine the level of dryness. However, if the skin or medical device is not dry when touched, such a touch may breach the applied prep solution and provide a pathway for microorganisms to enter the body through the skin or via the medical device. Thus, it is necessary and desirable for the healthcare practitioners to have an easy way to determine if a solvent within a surgical prep solution has completely evaporated from the surgical site or the medical device without touching the skin, the medical device, or the applied surgical prep solution.

[0009] An effective film forming polymer may also be added to the surgical prep solution to provide better adherence of the solution to the skin or the medical device. Film forming polymers are either water soluble or insoluble and most of the water insoluble films adhere poorly to the skin. Poor adherence is especially problematic in orthopedic surgery or surgical procedures in which patients are draped after the surgical prep solution has dried. If the surgical prep solution adheres poorly to the skin or is water soluble, the drape will typically fall off during the procedure, contaminating the sterile surgical environment and increasing the chance of a hospital-acquired infection.

[0010] U.S. Pat. No. 5,763,412 teaches a film-forming composition containing chlorhexidine gluconate. U.S. Pat. No. 5,547,662 (hereinafter "the '662 patent") is directed to a film-forming composition for preparation of a skin surface as a surgical site containing an antimicrobial agent and Dovicide A and D&C Red 17, which changes color from purple (wet) to red (dry), but does not teach color changes triggered by pH indicators. The disadvantages of the '662 patent include inter alia the particular combination is not applicable to antimicrobial with inherent color, such as iodine or iodophors and the color change is not reversible. In addition, U.S. Pat. No. 4,584,192 teaches chlorhexidine or its derivative, but does not disclose which chlorhexidine compound is suitable and has no visualization feature.

SUMMARY OF THE INVENTION

[0011] This invention discloses a surgical prep solution formulation, either film forming or non-film forming, that changes color upon drying to indicate when the surgical prep solution has completely dried. Advantageously, the surgeon

can determine that a selected surface, such as a surgical site or a medical device, is dry and the solvent has evaporated by visual determination.

DETAILED DESCRIPTION OF THE INVENTION

[0012] The present description emphasizes certain embodiments of the invention, but is not meant to be comprehensive of the entire scope of the invention. The full scope of the invention is measured by the appended claims.

[0013] The surgical prep solution comprises: topically-acceptable solvent(s), pH adjuster(s), color changing pH indicator(s) and optionally: antimicrobial agent(s), film forming polymer(s), dye(s) and viscosity regulator(s). After the solution is applied on the skin, the fluid components evaporate and the color changing pH indicator(s) interact with the human skin. The pH of the residue of the solution may rise or fall if the pH of the original solution is different from the normal pH range of the acid mantle of the human skin. As the pH of the residual solution changes, the pH indicator in the surgical prep solution may cause the residue to change color. Similarly and concurrently, as the fluid evaporates, the color changing pH indicator(s) may interact with atmospheric CO₂, changing the pH of the solution because CO₂ is mildly acidic. As such, the color changes according to the color properties of the pH indicator as the solution becomes less basic or more acidic. The rate of color change may be synchronized with the rate of fluid evaporation through the selection of pH indicators and pH adjusters so that the color change is complete when the fluid is sufficiently evaporated. Additionally, different color dyes and pH indicators may be utilized, to provide for a particular desired color.

[0014] The term "topically-acceptable solvent" as used here is a fluid that is compatible with the other ingredients of the composition and is non-toxic when applied to human skin. Suitable solvents include, but are not limited to, water, alcohols, acetone, esters, chlorinated hydrocarbons and chlorofluorohydrocarbons. Preferred solvents include, but are not limited to, water, and/or isopropanol, ethanol and other alcohols.

[0015] The term "antimicrobial agent" as used here is a chemical, such as a compound or salt that destroys or inhibits the growth of microorganisms. The term "microorganisms" as used here includes, but is not limited to, bacteria, fungi, and viruses. The antimicrobial agent may be, but is not limited to, alcohols (e.g., ethanol, isopropanol), chlorhexidine (e.g., chlorhexidine digluconate (CHG), chlorhexidine diacetate, chlorhexidine dihydrochloride, gluconic acid), hexachlorophene, iodophors, povidone-iodine, iodine, para-chlorometaxyleneol (PCMX), and triclosan. Desirably, the antimicrobial agent is present in a quantity sufficient to inhibit microbial growth on the surface of the skin.

[0016] The term "antimicrobial agent sensitive to the environmental pH" as used here is an antimicrobial agent whose stability is influenced by the ambient pH. The antimicrobial agent sensitive to the environmental pH is a subset of the antimicrobial agents described above and may include, but is not limited to, iodine, iodophor, chlorhexidine and its derivatives, such as chlorhexidine gluconate. For instance, iodine and iodophor are naturally acidic and may gradually lose their antimicrobial activity if the environmental pH is basic. If the selected pH indicator changes color in the higher pH range (e.g., bromocresol purple which is

yellow at pH below 5.2 and violet above 6.8), the antimicrobial activity of the iodophor solution may last up to several hours once a pH adjuster is added and the environmental pH turns basic. In a preferred embodiment where a pH adjuster is used where the environmental pH is not stable for antimicrobial agents, the solution should be applied immediately after the pH adjuster is added. In a more preferred embodiment, the two different pH components: (a) pH adjuster and/or pH indicator and (b) antimicrobial agents may be separately contained and mixed in situ through devices such as a dual channel syringe.

[0017] The term "film forming polymer" as used here is a polymer that forms a film that coats a surface (e.g., the skin) upon drying of the solution. The film is intact, resists both blood and saline, and contains the antimicrobial agent. Such polymer may include, but is not limited to, polyvinyl acetate, polyvinyl alcohol, methacrylic acid-ethyl acrylate copolymer, vinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohol-polyethylene glycol graft copolymer, ethyl acrylate-methyl methacrylate copolymer, acrylic polymers such as Eudragit®, and polyacrylate copolymer.

[0018] The term "pH indicator" as used here is a compound that is added in trace amounts to the surgical prep solution to determine its pH by visual observation. The pH indicator changes color based on the pH of the surgical prep solution. Such pH indicators include, but are not limited to, bromophenol blue, bromothymol blue, bromocresol green, bromocresol purple, congo red, methyl orange, methyl red, methyl yellow, pentamethoxy, phenolphthalein, thymolphthalein, p-naphtholbenzein, 4-nitrophenol, 3-nitrophenol, o-cresolphthalein, m-cresol red, thymol blue, m-cresol purple, phenol red, and mixtures thereof. The pH indicator may be separately contained and added in situ prior to application to the skin at the surgical site. In a preferred embodiment, the composition undergoes a visible color change upon a change in pH from greater than about 6.5 to between about 4.0 to about 5.5. Alternatively, in another preferred embodiment, the composition undergoes a visible color change upon a change in pH from less than about 3.0 to between about 4.0 to about 5.5. More preferably, the composition undergoes a visible color change upon a change in pH from less than about 4.5 to between about 4.5 to 5.5.

[0019] The term "color dye" as used here are commonly used dyes or pigments. Such color dyes include, but are not limited to, D&C dyes (Green #5, Green #8, Orange #4, Red #4, Red #17, Red #22, Red #28, Red #33, Violet #2, Yellow #8, Yellow #10, Yellow #11), FD&C dyes (Blue#1, Blue#2, Carmine, Emerald Green, Green #3, Red #3, Red #40, Yellow #5, Yellow #6) and mixtures thereof.

[0020] The term "pH adjuster" as used here is an acid, a base or a pH buffer. The pH adjuster is used to adjust the solution pH so that the solution pH falls within one extreme of the color spectrum of the selected pH indicator. The rate of color change is controlled by the pH adjuster (i.e., acid, base or pH buffer). For instance, if one selects bromothymol blue as the pH indicator which is yellow when pH is below 6 and blue when pH is above 7.6, a pH adjuster is needed to change the solution pH to either below 6 or above 7.6. Such acids include, but are not limited to, acetic acid, boric acid, carbonic acid, chromic acid, citric acid, lactic acid, diluted hydrochloric acid, tartaric acid, propionic acid, malic acid, diluted phosphoric acid and mixtures thereof. Such bases include, but are not limited to, ammonium hydroxide, ammonium carbonate, ethylamine, dimethylamine, glycine,

methylamine, trimethylamine, diethanolamine, sodium bicarbonate, sodium borate, sodium hydroxide, hydrazine, monoethanolamine, potassium hydroxide, sodium phosphate dibasic, triethylamine and mixtures thereof. The pH adjuster may also include pH buffer agents such as the Hydrion™ buffer mixture produced by Micro Essential Laboratory, Inc. (Brooklyn, N.Y.) which contains a powdery mixture of potassium phosphate monobasic and sodium phosphate dibasic. The pH adjuster may be separately contained and added in situ prior to application because some antimicrobial agents have a shorter shelf life depending on the pH of their environment.

[0021] In a preferred embodiment, the composition according to the present invention is a liquid solution with a viscosity similar to water and/or ethanol. Alternatively, the composition may take various physical forms such as creams, gels, lotions, emulsions, foams, and the like. Viscosity regulators may be used to control the viscosity and rheological properties of the composition according to the invention. Suitable viscosity regulators include, but are not limited to, polyethylene glycol, polyethylene glycol derivatives, ethyl cellulose, methoxycellulose, hydroxyethylcellulose, polyvinylpyrrolidone/vinyl acetate copolymer, and crosslinked pyrrolidone.

[0022] A preferred embodiment of the surgical prep solution includes compounds that provide a first color which visually indicates the skin area covered by the composition and a second color which gives a visual indication that the solvent has been substantially eliminated by evaporation. These compounds usually refer to pH indicators, but it may also be other colored ingredients in the formulation, such as an antimicrobial agent or film forming polymers, as they alter the overall color.

[0023] In an alternative embodiment, the color change surgical prep solution is applied to medical devices, for example, needle-free connectors and valves. Generally, devices used for needle-free intravenous administration include one or more needle-free connectors. Preferably, the needle-free connectors include central venous catheters capable of accessing a patient's bloodstream. The needle-free connectors may function as access ports to the patient's bloodstream and may be used for injecting medication or drawing blood samples. The needle-free devices are designed to reduce accidental needle punctures and exposure to potentially infectious blood samples by substituting needles with needle-less devices for intravenous administration. The needle-free valve opens and fluid may flow therethrough when a fitting, preferably a male luer fitting, is inserted into the valve through a valve port. The valve seals when the fitting is removed and prevents fluid flow through the valve thereafter. The valve may, for example, be activated by a luer lock syringe, a secondary intravenous administration device or other intravenous products. In general, healthcare practitioners disinfect the needle-free connectors and valves with alcohol prep pads immediately prior to use.

[0024] It is essential that these medical devices are thoroughly disinfected in order to reduce a patient's risk of infection. Preferably, the color change surgical prep solution has a pH different from that of medical devices (e.g., needle-free connectors). The color change surgical prep solution may be swabbed onto a medical device. The solution may interact with surfaces of the medical device along with atmospheric CO₂, resulting in a pH change. As such,

the color may change according to the color properties of the pH indicator as the solution becomes less basic or more acidic. A healthcare practitioner may visually inspect the color of the medical device to determine whether the medical device has been thoroughly disinfected. The use of the color change surgical prep solution may allow healthcare practitioners to visually identify the disinfected area based on the color change.

[0025] In another alternative embodiment, the color change surgical prep solution may be applied as a coating onto medical devices. The coating may contain a pH buffer having a pH different from that of disinfecting solutions (e.g., alcohol) typically used in hospitals. As soon as the sterilizing solutions (e.g., alcohol) are applied to the coated medical devices, the solution (e.g., alcohol) may interact with the coating and cause a change in pH. Accordingly, the color may change based on the color properties of the pH indicator as the solution becomes less basic or more acidic. A healthcare practitioner may visually inspect the color of the medical devices to determine whether the medical device has been thoroughly disinfected.

[0026] Preferably, the interaction between and the coating and the disinfecting solution is reversible and that the coating returns to its original color once the disinfecting solution fully evaporates. The reversible color change may be utilized repeatedly. For example, a medical device, such as a needle-free connector, needs to be disinfected every time the device is used. During the period of use, a healthcare practitioner may access the same needle-free connector to inject medicaments or draw blood samples. The connector may be disinfected by a disinfecting solution and change color. Once the disinfecting solution has fully evaporated, the connector may return to its original color. The connector may be repeatedly disinfected by the disinfecting solution. A color change may occur every time the disinfecting solution is applied to the connector and the connector may return to its original color every time the disinfecting solution becomes fully evaporated.

[0027] The invention offers several advantages over the prior art. First, the invention applies to both film forming and non-film forming prep solutions whereas the prior art is directed to film-forming color change surgical prep solutions.

[0028] Second, the prior art excludes antimicrobial agents with inherent color. For instance, the inclusion of iodine would render ineffective the color change abilities of the surgical prep solutions of the prior art. The invention overcomes such a limitation, allowing flexibility in choice of antimicrobial agents.

[0029] Third, the invention allows flexibility in the potential skin hues of patients. No two skin hues are the same and patients themselves might have variations of skin tone or even a colored tattoo. Thus, color flexibility in the surgical prep solution is imperative. In the invention, the color possibilities are more flexible than the prior art allows and hence, the surgical prep solution's color may be tailored to the needs of both patient and surgeon.

[0030] Fourth, for embodiments of the invention that include a film-forming polymer, the tinted color is embedded within the film and cannot easily be washed off because the film strongly adheres to the skin. This feature is especially important for orthopedic surgeons employing drapes on the patient.

[0031] Fifth, the invention reduces the risk of flammability. Surgeons regularly use electrocautery to burn or destroy tissue and hence, must be absolutely sure no flammable gases or liquids are present. Isopropanol is widely used as an antimicrobial agent, but its flammability presents a problem to surgeons. The invention allows a surgeon to know when the isopropanol has been eliminated, thus reducing the risk of fire.

[0032] The invention also differs from the prior art in that pH indicators are used to change the color of the surgical prep solution. Known surgical prep solutions use chemicals such as Dovicide A and D&C Red 17 to change the color of the solution. Contrary to the present invention, the solvated Dovicide A and the dye form a complex and as the solvent evaporates, the complex shifts the wavelength of the light absorbed, thus changing the color of the complex. The mechanism is different from a pH-indicated color change.

[0033] The invention also differs from prior art in that the color change of the prep solution is reversible, if the environment pH change occurs during the procedure.

[0034] Finally, the invention facilitates the stability of antimicrobial agents that are unstable in certain pH ranges. The pH indicator and/or pH adjuster may be separately contained and added in situ prior to application to maintain the integrity of the antimicrobial agents and increase their effectiveness.

EXAMPLES

[0035] The following non-limiting examples illustrate particular embodiments of the invention. The examples are not meant to be comprehensive of the entire scope of the invention.

[0036] The following table lists both the composition of the one step surgical prep solution composition and a preferred embodiment.

Ingredient	Amount (wt./wt. %)	Amount, preferred embodiment (wt./wt. %)
Solvent	about 10–about 90	about 60–about 80
Antimicrobial agent (e.g. iodine or CHG, excluding alcohols)	about 0–about 5	about 1.5–about 2.5
Water	about 3–about 35	about 15–about 25
Color Dye (optional)	about 0–about 0.1	about 0.005–about 0.05
Film forming polymer (optional)	about 0–about 20	about 0–about 10
pH indicator	Trace	Trace
pH adjuster	Trace	Trace
Viscosity regulator	about 0–about 5	about 0–about 5

[0037] In preparing the solutions, solid ingredients, such as polymers (including viscosity regulator), solvents, and color dye were first mixed to allow the slower-dissolving ingredients (i.e., film forming polymers, if used) time to dissolve in the solvent system. Afterwards, the antimicrobial agents were mixed in, followed by the pH indicator and pH adjuster. Each solution was mixed at room temperature using a magnetic stirring bar. These conditions apply to all the following examples.

[0038] If the antimicrobial agents are sensitive to pH, the pH adjuster and pH indicator may be initially separated and incorporated through devices such as a two-cylinder syringe

with a static mixer or any other mixing mechanism and mixed with other components in situ prior to application.

Example A

Non-Film Forming

[0039]

Ingredient	Amount (wt./wt. %)
Isopropanol	77.5
Water	20.5
0.1% Bromothymol blue in IPA	1
1% Ammonium hydroxide	1

[0040] In preparing the solution in Example A, isopropanol and water were mixed first. Subsequently, bromothymol blue was added and the solution pH was adjusted to between 8 to 10 using ammonium hydroxide. When applied to the skin, Example A changed from blue to light yellow/colorless after the solvent evaporated.

Example B

Non-Film Forming

[0041]

Ingredient	Amount (wt./wt. %)
Isopropanol	64.2
20% CHG aqueous solution	11.7
Water	22.05
FD&C Red 40	.05
0.1% Bromothymol blue in IPA	1
1% Ammonium hydroxide	1

[0042] In preparing the solution in Example B, Red 40, isopropanol, 20% CHG aqueous solution, and water were mixed first. Subsequently, bromothymol blue was added and the solution pH was adjusted to between 8 to 10 using ammonium hydroxide. When applied to the skin, Example B changed from purple to red after the solvent evaporated.

Example C

Non Film Forming

[0043]

Ingredient	Amount (wt./wt. %)
Isopropanol	68.2
20% CHG aqueous solution	8.5
Water	21.3
0.1% Bromothymol blue in IPA	1.1
1% Ammonium hydroxide	0.9

[0044] In preparing the solution in Example C, isopropanol, 20% CHG aqueous solution, and water were mixed first. Subsequently, bromothymol blue was added and the solution pH was adjusted to between 8 to 10 using ammonium hydroxide. When applied to the skin, Example C changed from blue to colorless after the solvent evaporated.

Example D
Film Forming

[0045]

Ingredient	Amount (wt./wt. %)
Polyacrylate copolymer	7.4
Isopropanol	74.8
Iodine	0.68
Sodium iodide	0.62
Water	16.5
Bromophenol blue	trace
Propionic acid	trace

[0046] The components in Example D were combined to form a solution. Specifically, the polyacrylate was first dissolved in isopropanol and water, and then iodine, sodium iodide, and bromophenol blue were added. Propionic acid was used to adjust the solution pH to between 3.5 and 4. When applied on skin, the color changed from iodine gold color to green after the solvent evaporated. For this example, the integrity of the film was maintained for several hours and confirmed by visual observation.

Example E
Film Forming

[0047]

Ingredient	Amount (wt./wt. %)
Polyacrylate copolymer	7.5
Isopropanol	74.9
Water	15.6
CHG	2.0
FD&C Red 40	0.022
Bromothymol blue	trace
Methylamine	trace

[0048] In Example E, the polyacrylate copolymer and FD&C Red 40 were first dissolved in isopropanol and water. Subsequently, CHG and bromothymol blue were added and the solution pH was adjusted to between 8.5 and 9 using methylamine. When applied on skin, the color changed from purple to pink/red after the solvent evaporated. For this example, the integrity of the film was maintained for several hours and confirmed by visual observation.

Comparative Time-Kill Study (Examples F-I)

[0049] The following Examples F-I were made to assess the formulation's antimicrobial effectiveness when dry.

[0050] Bacterial suspensions of *Escherichia coli* ATCC # 11229 and *Staphylococcus aureus*, ATCC # 6538 were made in sterile saline to final inoculum level of 1.5×10^8 CFU/ml. Thirty (30) microliters of the testing antimicrobial solution were aseptically spread on a sterile microslide covering a surface of 25x25 sq. mm and dried for ten (10) minutes. One (1) microliter of the microbial suspensions was applied on a sterile cover slip. The antimicrobial films were applied to these microbial suspensions for exposure times of 1, 5, or 10 minutes. After the exposure time had elapsed, both the cover slip and the micro slide were dropped into a test tube

containing 10 ml sterile neutralizing solution. The Standard Plate Count Method was used for enumeration of the survival cells (CFU/ml). The log reduction in bacterial count was calculated from the log of the inoculum titers, which were enumerated by following the same procedure except without the presence of antimicrobial films.

[0051] Examples F-I present the results of the Time-Kill Study with the respective film solution.

Example F
Antimicrobial Efficacy

[0052]

Ingredient	Amount (wt./wt. %)
Film forming polymer	7.37
Iodine	0.67
Sodium iodide	0.71
Isopropanol	73.69
Water	16.38
Formic acid	1.16
Bromophenol blue	0.02

[0053] In preparing the solution in Example F, the film forming polymer was first dissolved in isopropanol and water. Afterwards, iodine, sodium iodide and bromophenol blue were added. The final pH was then adjusted to between 3.5 to 4 using formic acid.

Microorganism tested	Storage time at ambient conditions	Log reduction/ml at time intervals		
		1 min.	5 min.	10 min.
<i>E. coli</i> , gram negative bacterium	0 time	>4.98	>4.98	N/A
	2 week	>5.04	>5.04	N/A
	1 month	>4.86	>4.86	>4.86
	2 months	>5.05	>5.05	>5.05
<i>S. aureus</i> , gram positive bacterium	0 time	1.25	4.56	N/A
	2 week	1.21	3.92	N/A
	1 month	2.21	>4.62	>4.62
	2 months	4.29	>5.49	>5.49

Example G
Antimicrobial Efficacy

[0054]

Ingredient	Amount (wt./wt. %)
Film forming polymer	7.42
Iodine	0.68
Sodium iodide	0.72
Isopropanol	74.17
Water	16.49
Formic acid	0.5
Bromophenol blue	0.02

[0055] In preparing the solution in Example G, the film forming polymer was first dissolved in isopropanol and water. Afterwards, iodine, sodium iodide, and bromophenol blue were added. The final pH was then adjusted to between 3.5 to 4 using formic acid.

Microorganism tested	Storage time at ambient conditions	Log reduction/ml at time intervals		
		1 min.	5 min.	10 min.
<i>E. coli</i> , gram negative bacterium	0 time	>5	>5	N/A
	2 week	3.72	>5.04	N/A
	1 month	4.01	>4.86	>4.86
	2 months	>5.05	>5.05	>5.05
<i>S. aureus</i> , gram positive bacterium	0 time	>4.75	>4.75	N/A
	2 week	1.45	>5.07	N/A
	1 month	2.09	>4.62	>4.62
	2 months	5.01	>5.49	>5.49

Example H

Antimicrobial Efficacy

[0056]

Ingredient	Amount (wt./wt.)
Film forming polymer	7.40
Iodine	0.68
Sodium iodide	0.70
Isopropanol	74.64
Water	16.45
Formic acid	0.03
Citric acid, anhydrous	0.08
Bromophenol blue	0.02

[0057] In preparing the solution in Example H, the film forming polymer was first dissolved in isopropanol and water. Afterwards, iodine, sodium iodide and bromophenol blue were added. The final pH was then adjusted to between 3.5 to 4 using both citric acid and formic acid.

Microorganism tested	Storage time at ambient conditions	Log reduction/ml at time intervals		
		1 min.	5 min.	10 min.
<i>E. coli</i> , gram negative bacterium	1 month	>5.05	>5.05	>5.05
<i>S. aureus</i> , gram positive bacterium	1 month	4.13	>5.49	>5.49

Example I

Antimicrobial Efficacy

[0058]

Ingredient	Amount (wt./wt.)
Film forming polymer	7.29
Iodine	0.67
Sodium iodide	0.62
Isopropanol	74.79

-continued

Ingredient	Amount (wt./wt.)
Water	16.20
Propionic acid, 99.5%	0.31
Citric acid, anhydrous	0.10
Bromophenol blue	0.02

[0059] In preparing the solution in Example I, the film forming polymer was first dissolved in isopropanol and water. Afterwards, iodine, sodium iodide and bromophenol blue were added. The final pH was then adjusted to between 3.5 to 4 using both citric acid and propionic acid.

Microorganism tested	Storage time at ambient conditions	Log reduction/ml at time intervals		
		1 min.	5 min.	10 min.
<i>E. coli</i> , gram negative bacterium	10 days	>5.05	>5.05	>5.05
<i>S. aureus</i> , Gram positive bacterium	10 days	4.38	>5.49	>5.49

Examples J-M (Rate of Color Change)

[0060] Examples J-M illustrate the rate of color change controlled through different acid combinations. Specifically, the higher proportion of acid, the slower the color change when applied on skin.

Ingredient (wt/wt %)	Example J	Example K
Film forming polymer	8.967	8.950
Iodine	0.727	0.726
Sodium iodide	0.777	0.776
Isopropanol	72.350	72.293
Water	16.937	16.907
Citric acid	0.157	0.263
Formic acid	0.060	0.060
Bromophenol blue	0.025	0.025

[0061] After being applied on skin, Example J changes color within 1 minute while Example K changes color within 2 to 3 minutes.

Ingredient (wt/wt %)	Example L	Example M
Film forming polymer	7.43	7.43
CHG	2.00	2
Isopropanol	74.81	74.81
Water	15.63	15.63
Bromothymol blue	0.07	0.07
Ammonium hydroxide	0.06	—
Dimethylamine	—	0.06

[0062] After being applied on skin, Example L color changes from blue to colorless within 1 minute while it takes Example M more than 2 minutes to change color.

1. A composition comprising an antimicrobial agent, one or more pH adjusters, and a color-changing pH indicator, wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

2. A composition comprising an antimicrobial agent, a topically-acceptable solvent, one or more pH adjusters, and a color-changing pH indicator, wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

3. The composition according to claim 2, wherein the antimicrobial agent is selected from the group consisting of chlorhexidine digluconate, chlorhexidine diacetate, chlorhexidine dihydrochloride, gluconic acid, hexachlorophene, iodophors, povidone-iodine, iodine, para-chlorometaxylenol, triclosan, and mixtures thereof.

4. The composition according to claim 2, wherein the topically-acceptable solvent comprises water, ethanol, isopropanol, acetone, or mixtures thereof.

5. The composition according to claim 2, wherein the one or more pH adjusters are selected from the group consisting of acetic acid, boric acid, carbonic acid, chromic acid, citric acid, lactic acid, diluted hydrochloric acid, tartaric acid, propionic acid, malic acid, diluted phosphoric acid, ammonium hydroxide, ammonium carbonate, diethanolamine, ethylamine, hydrazine, methylamine, monoethanolamine, potassium hydroxide, sodium bicarbonate, sodium borate, sodium borate, sodium hydroxide, sodium phosphate dibasic, trimethylamine, trolamine, pH buffer agents and mixtures thereof.

6. The composition according to claim 2, wherein the color-changing pH indicator is selected from the group consisting of bromophenol blue, bromothymol blue, bromocresol green, bromocresol purple, congo red, methyl orange, methyl red, methyl yellow, pentamethoxy, phenolphthalein, thymolphthalein, p-naphtholbenzein, 4-nitrophenol, 3-nitrophenol, o-cresolphthalein, m-cresol red, thymol blue, m-cresol purple, phenol red, and mixtures thereof.

7. The composition according to claim 2, further comprising a film-forming polymer.

8. The composition according to claim 7, wherein the film-forming polymer is selected from the group consisting of polyvinyl acetate, polyvinyl alcohol, methacrylic acid-ethyl acrylate copolymer, vinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohol-polyethylene glycol graft copolymer, ethyl acrylate-methyl methacrylate copolymer, acrylic polymer, polyacrylate copolymer, and mixtures thereof.

9. The composition according to claim 2, further comprising a dye.

10. The composition according to claim 9, wherein the dye is selected from the group consisting of Green #5, Green #8, Orange #4, Red #4, Red #17, Red #22, Red #28, Red #33, Violet #2, Yellow #8, Yellow #10, Yellow #11, Blue #1, Blue #2, Carmine, Emerald Green, Green #3, Red #3, Red #40, Yellow #5, Yellow #6, and mixtures thereof.

11. The composition according to claim 2, further comprising a viscosity regulator.

12. The composition according to claim 11, wherein the viscosity regulator is selected from the group consisting of polyethylene glycol, polyethylene glycol derivatives, ethyl cellulose, methoxycellulose, hydroxyethylcellulose, polyvinylpyrrolidone/vinyl acetate copolymer, crosslinked pyrrolidone and mixtures thereof.

13. The composition according to claim 2, wherein the composition undergoes a visible color change upon a change in pH.

14. The composition according to claim 13, wherein the composition reversibly undergoes a visible color change upon change in pH.

15. The composition according to claim 13, wherein the rate of color change is controlled by the combination of pH adjusters.

16. The composition according to claim 13, wherein the composition undergoes a visible color change upon a change in pH from greater than about 6.5 to between about 4.5 to about 5.5.

17. The composition according to claim 13, wherein the composition undergoes a visible color change upon a change in pH from less than about 3.0 to between about 4.0 to about 5.5.

18. The composition according to claim 13, wherein the composition undergoes a visible color change upon a change in pH from less than about 4.5 to between about 4.5 to about 5.5.

19. A method of preparing a surgical prep solution used to disinfect a pre-operative surgical site comprising combining an antimicrobial agent, one or more pH adjusters, and a color changing pH indicator to form a solution, wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

20. A method of preparing a surgical prep solution used to disinfect a pre-operative surgical site comprising combining an antimicrobial agent, a topically-acceptable solvent, one or more pH adjusters, and a color-changing pH indicator to form a solution, wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

21. A method of disinfecting comprising:

applying a composition to a selected surface, evaporating the composition, and visually inspecting the residue of the composition on the selected surface for a change in color, wherein the composition comprises an antimicrobial agent, one or more pH adjusters, and a color-changing pH indicator, and wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

22. A method of disinfecting comprising:

applying a composition to a selected surface, evaporating the composition, and visually inspecting the residue of the composition on the selected surface for a change in color, wherein the composition comprises an antimicrobial agent, a topically-acceptable solvent, one or more pH adjusters, and a color-changing pH indicator, and wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

23. The method according to claim 22, wherein the selected surface is skin.

24. The method according to claim 22, wherein the selected surface is a medical device.

25. The method according to claim 24, wherein the medical device is a needleless injection connector.

26. The method according to claim 22, wherein the antimicrobial agent is an antimicrobial agent sensitive to the environmental pH.

27. The method according to claim 26, wherein the antimicrobial agent sensitive to the environmental pH is

selected from the group consisting of iodine, idophor, chlorhexidine and chlorhexidine gluconate.

28. The method according to claim **22**, further comprising preparing the composition of the applying step.

29. The method according to claim **28**, wherein the preparing step includes adding the one or more pH adjusters to the composition in situ prior to application of the composition to the skin.

30. The method according to claim **28**, wherein the preparing step includes adding the pH indicator to the composition in situ prior to application of the composition to the skin.

31. A medical device comprising:

a coating comprising one or more pH adjusters, and a color-changing pH indicator,

wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

32. The medical device according to claim **31**, wherein the coating reversibly undergoes a visible color change upon change in pH.

33. The medical device according to claim **31**, wherein the coating further comprises an antimicrobial agent.

34. The medical device according to claim **31**, wherein the medical device is one of a needle-free connector or a needle-free valve.

35. A method of disinfecting comprising:

coating a composition of a first color to a selected surface, the coating comprising one or more pH adjusters and a color-changing pH indicator,

applying a disinfecting solution to the selected surface, visually inspecting the coating for a change in color to a second color,

evaporating the disinfecting solution, and visually inspecting the coating for a change in color to the first color,

wherein the one or more pH adjusters is an acid, a base, a pH buffer or a combination thereof.

* * * * *