

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 May 2003 (01.05.2003)

PCT

(10) International Publication Number
WO 03/035601 A1

(51) International Patent Classification⁷: C07C 225/06

(21) International Application Number: PCT/KR01/01796

(22) International Filing Date: 24 October 2001 (24.10.2001)

(25) Filing Language: Korean

(26) Publication Language: English

(71) Applicant (for all designated States except US): J & J CHEMICAL CO., LTD. [KR/KR]; 1236, Jung-wang-Dong, Kyunggi-Do, Sihung-Si 429-450 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LEE, Chun-hyuk [KR/KR]; 1236, Jungwang-Dong, Kyunggi-Do, Sihung-Si 429-450 (KR). HAN, Sang-Jin [KR/KR]; 1236, Jung-wang-Dong, Kyunggi-Do, Sihung-Si 429-450 (KR).

(74) Agent: LEE, Sang-Hun; Han Nuri Patent & Law Office, 3Fl. suite 303, Back Ak Building, 828-53, Yoksam-Dong, Kangnam-Ku, Seoul 135-080 (KR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD AND APPARATUS FOR PREPARING HYDRAZO-DICARBONAMIDE USING UREA AS STARTING MATERIAL



(57) Abstract: The present invention relates to a method and apparatus for preparing hydrazodicarbonamide using urea as starting material economically and environmentally desirably. The apparatus includes a pyrolysis furnace to obtain biuret and ammonia by pyrolyzing urea; a recrystallization reactor to purify the biuret obtained from the pyrolysis furnace; a first reactor to obtain a metal monohalobiuret salt by reacting the biuret with metal hypohalogen compound or with halogen and base; a second reactor to synthesize the hydrazodicarbonamide by reacting the monohalobiuret metal salt with ammonia; and an ammonia evaporator to separate the excess ammonia from hydrazodicarbonamide and to supply the separated ammonia to an ammonia concentrator.



WO 03/035601 A1

METHOD AND APPARATUS FOR PREPARING HYDRAZO- DICARBONAMIDE USING UREA AS STARTING MATERIAL

FIELD OF THE INVENTION


5 The present invention relates to a method and apparatus for preparing hydrazodicarbonamide with urea as a starting material, and more particularly to a method and apparatus for preparing hydrazodicarbonamide economically and environmentally desirably by producing biuret with urea, and reacting the obtained biuret with ammonia produced during the process of biuret synthesis.

10

BACKGROUNDS OF THE INVENTION

Hydrazodicarbonamide(HDCA) is a useful compound as a raw material for preparing azodicarbonamide which is widely used as a foaming agent. As shown in the following reaction Equation 1, azodicarbonamide(2) can be obtained by
15 oxidation of hydrazodicarbonamide (1) with proper oxidation agent.


[Equation 1]



The conventional methods for preparing the hydrazodicarbonamide includes methods of (i) using hydrazine as a starting material, (ii) direct
20 synthesis from urea, (iii) obtaining semicarbazide using urea, and then converting the obtained semicarbazide to hydrazodicarbonamide, and (iv) using biuret as starting material.

In the method of using hydrazine as a starting material (Reaction Equation 2), hydrazodicarbonamide is produced by reacting 1 mol of hydrazine(3) with 2 mol of urea(4),

5 [Equation 2]



The above-identified reaction has a merit in that the process is simple, but it has drawbacks in that the starting material, hydrazine is difficult to synthesis and expensive. The representative methods for preparing hydrazine includes Raschig process and method of using ketazine. However, there are also some problems in that hydrazine obtained by these methods needs concentration process and hydrolysis process. Therefore, the costs for energy and for utility are too high and accordingly the production cost is increasing. Further, hydrazine can also be prepared by the urea process which reacts urea with sodium hypochlorite and sodium hydroxide. But, this method needs excess of sodium hydroxide, and the cost to remove sodium carbonate by-product is very high, and many chemicals are required to remove the by-product. Thus, this method is esteemed as uneconomical and environmentally undesirable.


10

15

20 Following Equation 3 represents the direct synthetic method of hydrazodicarbonamide using urea. As shown in the Equation 3, the reaction of 3 mol of urea with 4 mol of sodium hydroxide and 1 mol of chlorine produces 1 mol

of hydrazodicarbonamide. But, this method also is improper because the production cost is high due to the requirement of excess reagents and the process is very complicated. And there is another important problem in that a lot of ammonia are formed as by-product, which is environmentally undesirable.


5 [Equation 3]



Following Equation 4 shows another method of synthesizing hydrazodicarbonamide. The method comprises the steps of obtaining semicarbazide using urea, and subsequently converting the obtained semicarbazide to hydrazodicarbonamide. As shown in Equation 4, sodium monochlorourea salt is obtained by the reaction of urea with sodium hypochlorite, and the sodium monochlorourea salt reacts with excess ammonia in the presence of catalyst to produce the intermediate (semicarbazide), and then the obtained semicarbazide reacts with urea to produce the final product (hydrazodicarbonamide).

10
15

[Equation 4]




However this reaction also is economically inefficient because the reaction needs more than 500 times of excess ammonia per sodium monochlorourea salt,

20

or semicarbazide is obtained by using expensive catalyst. There's another problem that entire process becomes longer because additional reaction of converting semicarbazide to hydrazodicarbonamide should be followed.

5 Following Equation 5 shows the method of synthesizing hydrazodicarbonamide using biuret (International Application PCT/KR00/ 00180). It comprises the steps of obtaining metal monohalobiuret salt by reacting biuret with metal hypohalogen (MOX), and subsequently reacting the obtained metal monohalobiuret with ammonia to obtain hydrazodicarbonamide.

10 [Equation 5]



15 However, the above method of producing hydrazodicarbonamide using biuret as a starting material has problems in that the whole process is uneconomic and environmentally undesirable because the biuret used as the starting material is very expensive or it contains a lot of impurities, and the reaction of biuret with ammonia to synthesize hydrazodicarbonamide needs additional ammonia.

SUMMARY OF THE INVENTION


20 It is an object of the present invention to provide a method for preparing hydrazodicarbonamide economically and environmentally desirably using urea which is cheap and easily acquirable as a starting material.

It is a further object of the present invention to provide a method and apparatus for preparing hydrazodicarbonamide which can minimize the amount of the by-products and starting materials.


It is a further object of the present invention to provide a method and apparatus for preparing hydrazodicarbonamide with high yield by performing the whole process in a continuous manner.

To achieve these objects, the present invention provides a method for preparing hydrazodicarbonamide which comprises the steps of obtaining biuret of Formula 1 and ammonia by pyrolyzing urea, obtaining metal monohalobiuret salt of Formula 2 or 3 by reacting the obtained biuret with metal hypohalogen compound or with halogen and base, and reacting the obtained metal monohalobiuret salt with ammonia.


[Formula 1]



[Formula 2]



[Formula 3]



In above Formula 2 and 3, M represents metal and X represents halogen. Preferably, the urea pyrolysis temperature is 100~300°C, and the pyrolysis process is carried out while removing ammonia, and the removed ammonia reacts

with the metal monohalobiuret salt.

The present invention further provides an apparatus for preparing hydrazodicarbonamide which includes pyrolysis furnace to obtain biuret and ammonia by pyrolyzing urea; recrystallization reactor to purify the biuret obtained
5 from the pyrolysis furnace; a first reactor to obtain an metal monohalobiuret salt by reacting the biuret with metal hypohalogen compound or with halogen and base; a second reactor to synthesize the hydrazodicarbonamide by reacting the metal monohalobiuret salt with ammonia; and an ammonia evaporator to separate the excess ammonia from hydrazodicarbonamide and to forward the separated
10 ammonia to ammonia concentrator.

Preferably, the ammonia concentrator is to concentrate the excess ammonia and ammonia obtained from the pyrolysis furnace, and to supply the concentrated ammonia to the second reactor. The pyrolysis furnace may include a gas injector for injecting inert gas, which does not react with isocyanic acid, into
15 the pyrolysis furnace, and may include means for lowering pressure to remove ammonia from the pyrolysis furnace.

BRIEF DESCRIPTION OF THE DRAWINGS

A more complete appreciation of the invention, and many of the attendant
20 advantages thereof, will be readily apparent as the same becomes better understood by reference to the following detailed description when considered in conjunction with the accompanying drawing, wherein:

Fig. 1 is a schematic diagram showing an apparatus to prepare hydrazodicarbonamide according to an embodiment of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention will become more clearly understood from the following detailed description with reference to the accompanying drawing.


5 To prepare hydrazodicarbonamide according to the present invention, at first, biuret represented by Formula 1 and ammonia are produced by pyrolyzing urea at a temperature above the melting point of urea. Generally biuret is widely used as a precursor of pharmaceuticals, weedicide and reagent for analysis, also used in large amount as feed for ruminants, and is applied in various field of
10 plastic resins. Furthermore, it is reported that the some derivatives of biuret works as physiological healing agent or chemical therapeutic agent. The following Equation 6 shows the biuret synthesis process by pyrolyzing urea.

[Equation 6]



15 As shown in Equation 6, pyrolysis of 2 mol of urea gives biuret by elimination of ammonia. More precisely, as shown in the following Equation 7, it is presumed that isocyanic acid and ammonia is formed by pyrolysis of urea at first, and then the isocyanic acid reacts with another urea, which gives the target product biuret.

20 [Equation 7]



The biuret synthesis by pyrolysis of urea has merits in that the reaction is simple and operation of reaction process is easy, but also has drawbacks in that the conversion ratio of urea to biuret is low because many impurities like triuret, cyanuric acid are formed due to reaction of biuret with isocyanic acid in the biuret formation process. If the temperature is increased and the reaction time is lengthened to increase the conversion ratio, the impurities like triuret, cyanuric acid also increase. If the temperature is lowered to reduce the impurities, the reaction rate will be very slow, which makes the process non-economical. In the present invention, to raise the yield of biuret and to reduce the impurities, the temperature is preferably maintained at 100~300°C and more preferably is maintained at 130~170°C.


Further, if inert gas such as air and nitrogen which does not react with isocyanic acid is injected into the reactor, and/or the pressure of reactor is lowered, the ammonia, the byproduct formed during the reaction, can be effectively removed from the reactor. Then the reaction rate increases, and formation of impurities can be lowered, too. In addition, liquid phase organic compound which can be changed to the inert gas in the reactor of high temperature can be used as the inert gas source.

Further, if necessary, catalyst to increase the pyrolysis reaction rate can be used. Preferably, inorganic acid catalyst such as nitric acid, hydrochloric acid, and sulfuric acid, and acid type catalyst such as thionyl chloride, and phosphorous


containing substances such as sodium phosphate can be used as the catalyst. The preferable amount of catalyst is 0.001~0.5mol per 1mol of urea, and more preferable amount is 0.01~0.3mol per 1mol of urea.

- 5 Metal monohalobiuret salt of the following Formula 2 or 3 can be produced by reacting the obtained biuret with metal hypohalogen compound or halogen and base.

[Formula 2]




- 10 [Formula 3]




- In above Formula 2 and 3, M represents metal and X represents halogen. The direct method of preparing metal monohalobiuret salt by reacting biuret with metal hypohalogen compound is shown in Equation 8, and the specific example is shown in Equation 9.

[Equation 8]



In above Equation, M represent metal and X represent halogen.

[Equation 9]




Referring the above Equation 9, biuret reacts with sodium hypochlorite to form sodium chlorobiuret salt. Because above reaction is exothermic, the reaction system preferably would be maintained at low temperature. But the obtained sodium chlorobiuret salt is stable against moderate heat, it can be prepared at room temperature. Preferable reaction temperature is less than 60°C, and more preferably -10 ~ 60°C and most preferably -5 ~ 35°C. Considering the economic efficiency and operational facility, the reaction mol ratio of metal hypohalogen per 1 mole of urea is preferably between 0.1 and 2. When the reaction mol ratio is less than 1 mol, the excess biuret can be recovered and can be reused. In above reaction, when the reaction mol ratio is less than 0.1 or the reaction temperature is less than -10 °C, the reaction time would be too long. And if the reaction mol ratio is more than 2, the production cost increases and the side reaction may occur. Moreover, if the reaction temperature is more than 60°C, the produced metal monohalobiuret salt can be decomposed because it is unstable at high temperature. The sodium chlorobiuret salt obtained under above-mentioned condition can be used directly or can be stored for next reaction.

A process of producing metal monohalobiuret salt of above Formula 2 or 3 by reacting biuret with halogen and base is shown in Equation 10. As shown in Equation 10, after reacting biuret with halogen such as chlorine or halogen compound to obtain monohalobiuret(5), metal monohalobiuret salt can be obtained by adding base such as metal hydroxide (for example, sodium hydroxide,

potassium hydroxide, calcium hydroxide) to the obtained monohalobiuret(5).

[Equation 10]



In above Equation 10, M represent metal and X represent halogen.

5 Considering that the reaction for obtaining monohalobiuret(5) is exothermic, it would be advantageous that the reaction temperature is maintained lower, specifically less than 60°C, preferably -10 ~ 60°C and most preferably -5 ~ 30°C for the proper reaction rate and the stability of reaction. Alternatively, metal monohalobiuret salt can be obtained by mixing metal hydroxide with biuret at first, 10 and then reacting halogen with the obtained product. Because this reaction is also exothermic, the reaction temperature should be maintained lower, specifically to -10 ~ 60°C and more preferably -5 ~ 30°C. In above reaction, when the reaction temperature is less than 10°C, the reaction time would be too long, and when the reaction temperature is more than 60°C, the metal monohalobiuret salt 15 can be decomposed because it is unstable against heat. As shown in Equation 11, the obtained metal monohalobiuret salt can be metal 3-monohalobiuret salt(6) or metal 1-monohalobiuret salt(7).

[Equation 11]

group, and the converted isocyanate compound react with highly reactive ammonia to form hydrazodicarbonamide.

In the reaction of metal monohalobiuret salt with ammonia which is by-product of urea pyrolysis, considering the reaction rate and efficiency, preferable reaction temperature is between 0 and 150 °C, more preferably between 30 and 150 °C. When the reaction temperature is less than 0°C, the reaction rate is too slow and the economically inefficient, and when above reaction temperature is more than 150°C, the equipment cost is increase because the equipment must be designed to endure the internal pressure caused by ammonia vaporization.

Furthermore, ammonia can be used in either form of gaseous ammonia, or liquid ammonia or ammonium hydrate. Ammonia can be preferably used with excess amount to increase the reaction rate. The amount of ammonia can be between 1 and 1000mol per 1mol of metal monohalobiuret salt, more preferably is between 2 and 500mol, most preferably between 5 and 100mol. The excess ammonia except 1mol of ammonia which react with 1mol of metal monohalobiuret salt can be recovered and reused for next reaction. When the reaction temperature is high while using large amount of ammonia, the pressure of the reaction system can be raised to prevent the vaporization of ammonia. This improves the reaction rate and efficiency, and the preferable range of the pressure is between 1 and 100 kgf/cm².

According to the present invention, high yield can be achieved without using catalyst. However, if catalyst is used, it is very useful because the reaction time can be shortened, and the reaction efficiency can be improved. The examples

of the catalyst includes at least one compound selected from the group consisting of sulfuric acid salt, chloride salt, carbonate salt or hydroxide salt of basic metal or amphoteric metal, and organic compounds including the metals. The preferable amount of the catalyst is between 0.001 and 1mol, more preferably is 0.01~0.5mol per 1mol of metal monohalobiuret salt. As the catalyst, inorganic acid such as sulfuric acid, hydrochloric acid, or nitric acid can be added with the amount of 0.05~3.0mol per 1mol of metal monohalobiuret salt.

As a solvent of the reactant(biuret) or of entire reaction system, water can be used. If necessary, as a second solvent, at least one solvent selected from the group consisted with the polar solvent such as methanol, ethanol, propanol, isopropanol, tetrahydrofuran, acetonitrile, and the aprotic solvent such as dimethylformamide, dimethylsulfoxide, dimethylacetamide can be added. The amount of the second solvent is not limited particularly, but the preferable amount is between 0.1 and 50 times to the total weight of water, more preferable amount is 0.2~3.0. Furthermore, the second solvent can be introduced at the start of the reaction as a solvent for biuret or after mixing biuret solution with sodium hypochlorite solution.

The reaction for preparing hydrazodicarbonamide using urea as a starting material according to the present invention is shown in Equation 14 as a whole. In addition, an apparatus for preparing hydrazodicarbonamide according to an embodiment of the present invention is shown in Figure 1.

[Equation 14]



As shown in Figure 1, an apparatus to prepare hydrazodicarbonamide according to an embodiment of the present invention comprises pyrolysis furnace 10 to obtain biuret and gaseous ammonia by pyrolyzing urea. The pyrolysis furnace 10 preferably includes a gas injector 12 for injecting inert gas into the pyrolysis furnace 10, or may include means (not shown) for lowering pressure of the furnace 10 to easily remove ammonia from the pyrolysis furnace 10. The non-limiting examples of the inert gas includes air, nitrogen, and liquid phase organic compound which changes into an inert gas in the pyrolysis furnace 10, which does not react with isocyanic acid.

Ammonia removed from said pyrolysis furnace 10 is preferably supplied to ammonia concentrator 20, and the function of the ammonia concentrator 20 is concentrating the ammonia supplied from the pyrolysis furnace 10 and the excess ammonia remained after formation of hydrazodicarbonamide. The impurities, such as cyanuric acid and triuret, in biuret produced in the pyrolysis furnace 10 is separated by recrystallization means comprising recrystallization reactor 30 and dehydrator 32 such as a centrifuger, and then supplied to the first reactor 40.

Purified biuret which is sent to the first reactor 40 reacts with metal hypohalogen compound (for example, NaOCl) or halogen (for example, chlorine) and base to produce metal monohalobiuret salt, and then the produced metal monohalobiuret salt is supplied to the second reactor 50. The metal monohalobiuret salt reacts with ammonia to produce hydrazodicarbonamide, and the ammonia is preferably supplied from the ammonia concentrator 20. The

obtained hydrazodicarbonamide and excess ammonia are supplied to the ammonia evaporator 52. Ammonia evaporator 52 vaporizes the excess ammonia, and the vaporized ammonia is supplied to ammonia concentrator 20. The hydrazodicarbonamide separated from the excess ammonia is purified by
5 dehydrator 54 such as filter.

As shown in Equation 13 and Figure 1, according to the present invention, the hydrazodicarbonamide can be prepared in one continuous process from a starting material, urea. Because the entire process is carried out continuously,
10 the efficiency of process can be improved. In addition, the production cost can be lowered by greatly reducing the required amount of the raw material due to the fact that metal monohalobiuret salt reacts with ammonia, which is produced as a byproduct in the process of biuret formation. Thus, hydrazodicarbonamide can be prepared environmentally desirably by using the environmentally undesirable
15 by-product ammonia.

Hereinafter, the preferable examples and manufacturing examples are provided for better understanding of the present invention. However, the present invention should not be understood to be restricted to the following Examples.
20

[Manufacturing Examples 1 to 4: Preparation of biuret]

A four necked round bottomed flask was charged with 500g(8.33mole) of urea, stirred vigorously, and air was injected to the bottom side of the flask at the rate shown in following table 1. Simultaneously the reaction was carried out during

5 hours with the reaction temperature maintained at 140°C by heating. After completion of reaction, the composition of the obtained solid was analyzed by using liquid chromatography and the result was disclosed by following Table 1.

[Table 1]

Example	Air injection rate (L/min)	Content of Urea (wt%)	Content of Biuret (wt%)	Content of Cyanuric acid and other solid (wt%)
1	0	62	35	3
2	1	41	55	4
3	2	38	60	2
4	4	37	61	2

5

[Manufacturing Examples 5 to 7: Preparation of biuret]

Biuret was prepared by the same method as Example 1 except that the reaction was carried out for 3 hours while varying the reaction temperature and maintaining the air injection rate at 2 L/min. After completion of reaction, the composition of the obtained solid is analyzed by using liquid chromatography and the result was disclosed by following Table 2.

10

[Table 2]

Example	The reaction temperature (°C)	Content of Urea (wt%)	Content of Biuret (wt%)	Content of Cyanuric acid and other solid (wt%)
4	150	47	50	3
5	160	38.5	57	4.3
6	170	28	65	7

[Manufacturing Examples 8 to 10: Preparation of biuret]

Biuret was prepared by the same method as Example 1 except that the reaction was carried out with lowering the pressure as shown in following Table 3

15

by vacuum pump instead of air injection. After completion of reaction, the composition of the obtained solid is analyzed by using liquid chromatography and the result was disclosed by following Table 3.

[Table 3]

Example	The pressure (mmHg)	Content of Urea (wt%)	Content of Biuret (wt%)	Content of Cyanuric acid and other solid (wt%)
8	380	56	50	4
9	190	41.5	55	3.5
10	100	40	57	3

5

[Manufacturing Examples 11 to 13: Preparation of biuret]

Biuret was prepared by the same method as Example 1 except that the reaction was carried out using 0.05 mole of various catalysts per 1 mole of urea, and the air injection rate is fixed at 2L/min. After completion of reaction, the composition of the obtained solid is analyzed by using liquid chromatography and the result was disclosed by following Table 4.

10

[Table 4]

Example	Catalyst	Content of Urea (wt%)	Content of Biuret (wt%)	Content of Cyanuric acid and other solid (wt%)
11	Sulfuric acid	34	62	4
12	Sodium phosphate	36	61	3
13	Thionyl chloride	35	62	3

[Manufacturing Example 14: Synthesis of sodium chlorobiuret salt]

A 2L glass reactor was charged with 423.1g(0.287mole) of slurry solution of 7% biuret, and cooled to 5°C with stirring. To this reactor, aqueous solution of 12% sodium hypochlorite was added, and the reaction temperature of the system

15

was maintained below 5°C. After completion of addition, the reaction solution was analyzed by iodometry and by using liquid chromatography. The available chlorine was 3.37%. This corresponded to a yield of 98%.

5 [Manufacturing Example 15: Synthesis of sodium chlorobiuret salt]

A 2L glass reactor was charged with 423.1g(0.287mole) of slurry solution of 7% biuret, and cooled to 5°C with stirring. To this reactor, 223g (0.575mole) of aqueous solution of 10.3% sodium hydroxide was added, and 20.3g(0.287mole) of gaseous chlorine was added maintaining the reaction temperature of the system
10 below 10°C. After completion of the addition, the reaction solution was analyzed by iodometry and by using liquid chromatography. The available chlorine was 3.0%. This corresponded to a yield of 98%.

[Manufacturing Example 16: Synthesis of sodium chlorobiuret salt]

15 A 2L glass reactor was charged with 423.1g(0.287mole) of slurry solution of 7% biuret, and cooled to 5°C with stirring. To this reactor, 20.3g(0.287mole) of gaseous chlorine was added maintaining the reaction temperature of the system below 10°C. After addition of gaseous chlorine, 223g (0.575mole) of aqueous solution of 10.3% sodium hydroxide was added while vigorously stirring and the
20 reaction temperature was maintained below 5°C. After completion of addition, the reaction solution was analyzed by iodometry and by using liquid chromatography. The available chlorine was 3.0%. This corresponded to a yield of 98%.

[Examples 1-9: Synthesis of hydrazodicarbonamide]

A 2L autoclave was charged with 593.1g of sodium chlorobiuret salt obtained by the above Manufacturing Example 14, and cooled to 10°C with stirring. Maintaining the reaction temperature of solution below 10°C, 600g(8.8mole) of 25% aqueous ammonia solution was added to this while vigorously stirring. The reaction was carried out varying the reaction temperature and reaction time. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed in Table 5.

[Table 5]

Example	Reaction condition (temperature, time)	Yield (%)
1	30°C, 1hr	85
2	30°C, 2hrs	90
3	30°C, 3hrs	89
4	60°C, 1hr	91
5	60°C, 2hrs	89
6	60°C, 3hrs	90
7	90°C, 1hr	88
8	90°C, 2hrs	89
9	90°C, 3hrs	90

[Examples 10-18: Synthesis of hydrazodicarbonamide]

Reaction was carried out by the same method as Example 4 except that the 0.05 mole of various catalysts shown in Table 6 was added. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following Table 6.

[Table 6]

Example	Catalyst used	Yield (%)
10	ZnCl ₂	94
11	Zn(OH) ₂	92
12	AlCl ₃	90
13	BaCl ₂	91
14	CdCl ₂	92
15	ZnSO ₄	93
16	ZnCl ₂ + AlCl ₃ (0.025mole each)	96
17	ZnCl ₂ + BaCl ₂ (0.025mole each)	94
18	ZnCl ₂ + CdCl ₂ (0.025mole each)	96

[Examples 19-27: Synthesis of hydrazodicarbonamide]

A 2L autoclave was charged with 593.1g of sodium chlorobiuret salt
 5 obtained by the Manufacturing Example 15, and cooled to 10°C with stirring.
 Maintaining the reaction temperature of solution below 10°C, 600g(8.8mole) of
 25% aqueous ammonia solution was added while vigorously stirring. The reaction
 was carried out varying the reaction temperature and reaction time. After
 completion of reaction, unreacted ammonia was removed, and the reaction
 10 solution was filtered to get hydrazodicarbonamide insoluble to water and the yield
 of hydrazodicarbonamide was calculated and disclosed by the following Table 7.

[Table 7]

Example	Reaction condition (temperature, time)	Yield (%)
19	30°C, 1hr	78
20	30°C, 2hrs	89
21	30°C, 3hrs	89
22	60°C, 1hr	88

23	60 °C, 2hrs	90
24	60 °C, 3hrs	90
25	90 °C, 1hr	87
26	90 °C, 2hrs	86
27	90 °C, 3hrs	89

[Examples 28-36: Synthesis of hydrazodicarbonamide]

Reaction was carried out by the same method as Example 22 except that the 0.05 mole of various catalysts shown by following table 8 was added. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following Table 8.

[Table 8]

Example	Used catalyst	Yield (%)
28	ZnCl ₂	94
29	Zn(OH) ₂	91
30	AlCl ₃	89
31	BaCl ₂	91
32	CdCl ₂	93
33	ZnSO ₄	92
34	ZnCl ₂ + AlCl ₃ (0.025mole each)	97
35	ZnCl ₂ + BaCl ₂ (0.025mole each)	93
36	ZnCl ₂ + CdCl ₂ (0.025mole each)	96

[Examples 37-45: Synthesis of hydrazodicarbonamide]

A 2L autoclave was charged with 593.1g of sodium chlorobiuret salt obtained by the above Manufacturing Example 16, and cooled to 10 °C with stirring. Maintaining the reaction temperature of solution below 10 °C, 600g(8.8mole) of

25% aqueous ammonia solution was added while vigorously stirring. The reaction was carried out varying the reaction temperature and reaction time. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following Table 9.

[Table 9]

Example	Reaction condition (temperature, time)	Yield (%)
37	30°C, 1hr	79
38	30°C, 2hrs	88
39	30°C, 3hrs	89
40	60°C, 1hr	89
41	60°C, 2hrs	90
42	60°C, 3hrs	91
43	90°C, 1hr	88
44	90°C, 2hrs	88
45	90°C, 3hrs	89

[Examples 46-54: Synthesis of hydrazodicarbonamide]

Reaction was carried out by the same method as Example 40 except that the 0.05 mole of various catalysts shown by following Table 10 was added. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following Table 10.

[Table 10]

Example	Used catalyst	Yield (%)
46	ZnCl ₂	93

47	Zn(OH) ₂	90
48	AlCl ₃	90
49	BaCl ₂	90
50	CdCl ₂	92
51	ZnSO ₄	89
52	ZnCl ₂ + AlCl ₃ (0.025mole each)	95
53	ZnCl ₂ + BaCl ₂ (0.025mole each)	93
54	ZnCl ₂ + CdCl ₂ (0.025mole each)	94

[Examples 55-58: Synthesis of hydrazodicarbonamide]

A 2L autoclave was charged with 593.1g of sodium chlorobiuret salt obtained by the above Manufacturing Example 14, and cooled to 10°C with stirring. Maintaining the reaction temperature of solution below 10°C, aqueous ammonia solution was added while vigorously stirring for an hour at the amount shown by the following Table 11. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following table 11.

[Table 11]

Example	Mole ratio of ammonia vs sodium chlorobiuret salt (%)	Yield (%)
55	15	75
56	30	87
57	60	90
58	90	89

[Examples 59-62: Synthesis of hydrazodicarbonamide]

A 2L autoclave was charged with 593.1g of sodium chlorobiuret salt

obtained by the above Manufacturing Example 14, and cooled to 10°C with stirring and various organic solvent shown in Table 12 was added at the amount of 0.5 times to weight of water. Maintaining the reaction temperature of solution below 10°C, 600g of 25% aqueous ammonia solution was added while vigorously stirring for an hour. After completion of reaction, unreacted ammonia was removed, and the reaction solution was filtered to get hydrazodicarbonamide insoluble to water and the yield of hydrazodicarbonamide was calculated and disclosed by the following Table 12.

[Table 12]

Example	Solvent used	Yield (%)
59	Methanol	90
60	Dimethylformamide	94
61	Tetrahydrofuran	90
62	Acetonitrile	88

As described above, with the method and apparatus for preparing hydrazodicarbonamide according to the present invention, hydrazodicarbonamide can be synthesized from cheap and easily available urea as a starting material. In addition, hydrazodicarbonamide can be prepared economically and environmentally desirably due to the minimization of byproduct and input raw material, and with high efficiency from continuous process.

CLAIMS


1. A method for preparing hydrazodicarbonamide which comprises the steps of:

obtaining biuret of Formula 1 and ammonia by pyrolysis of urea;


5 obtaining metal monohalobiuret salt of Formula 2 or 3 by reacting the obtained biuret with metal hypohalogen compound or with halogen and base; and

reacting the obtained metal monohalobiuret salt with ammonia.


[Formula 1]



10 [Formula 2]



[Formula 3]



In Formula 2 and 3, M represents metal and X represents halogen.

15 2. The method for preparing hydrazodicarbonamide according to claim 1, wherein the temperature of pyrolysis of urea is 100 to 300°C.

3. The method for preparing hydrazodicarbonamide according to claim 1, wherein the urea pyrolysis process is carried out with removing ammonia from the reaction system thereof.

20 4. The method for preparing hydrazodicarbonamide according to claim 1, wherein the urea pyrolysis process is carried out with a process of

injection of inert gas and/or with a process of reducing pressure of the reaction system thereof.

5 5. The method for preparing hydrazodicarbonamide according to claim 1, wherein the urea pyrolysis process is carried out in the presence of at least one catalyst selected from the group consisting of inorganic acid catalyst, and acid type catalyst, and substances including phosphorous.

 6. The method for preparing hydrazodicarbonamide according to claim 1, wherein the mole ratio of the biuret and metal hypohalogen compound is 1: 0.1~ 1 : 2.

10 7. The method for preparing hydrazodicarbonamide according to claim 1, wherein the metal monohalobiuret salt is obtained by mixing metal hydroxide with the biuret of Formula 1, and then reacting the obtained product with halogen, or obtained by reacting the biuret of Formula 1 with gaseous halogen, and then mixing the obtained product with a base.

15 8. The method for preparing hydrazodicarbonamide according to claim 1, wherein the process of obtaining the metal monohalobiuret salt is carried out at a temperature of below 60°C.

 9. The method for preparing hydrazodicarbonamide according to claim 1, wherein the ammonia which reacts with the metal monohalobiuret salt comprises ammonia produced from the urea pyrolysis.

20 10. The method for preparing hydrazodicarbonamide according to claim 1, wherein the metal monohalobiuret salt reacts with excess ammonia including ammonia which is produced from the urea pyrolysis process, and the mole ratio of the metal monohalobiuret salt and the total ammonia is 1:1 ~1:1000.

11. The method for preparing hydrazodicarbonamide according to claim 1, wherein the ammonia is liquid ammonia, gaseous ammonia or ammonium hydrate.

12. The method for preparing hydrazodicarbonamide according to claim 1, wherein the reaction of the metal monohalobiuret salt with the ammonia is carried out at a temperature of 0 to 150 °C.

13. The method for preparing hydrazodicarbonamide according to claim 1, wherein the solvent of the reaction is a mixture of water and a second solvent, and the second solvent is a polar solvent selected from the group consisting of water, methanol, ethanol, propanol, isopropanol and mixtures thereof, or a aprotic solvent selected from the group consisting of dimethylformamide, dimethylsulfoxide, dimethylacetamide and mixtures thereof.

14. An apparatus for preparing hydrazodicarbonamide comprising:
a pyrolysis furnace to obtain biuret and ammonia by pyrolyzing urea;
a recrystallization reactor to purify the biuret obtained from the pyrolysis furnace;
a first reactor to obtain an metal monohalobiuret salt by reacting the biuret with metal hypohalogen compound or with halogen and base;
a second reactor to synthesize the hydrazodicarbonamide by reacting the monohalobiuret metal salt with ammonia; and
an ammonia evaporator to separate the excess ammonia from hydrazodicarbonamide and to supply the separated ammonia to an ammonia concentrator.

15. The apparatus for preparing hydrazodicarbonamide according to

claim 14, wherein the ammonia concentrator is to concentrate the excess ammonia and ammonia obtained at the pyrolysis furnace, and is to supply the concentrated ammonia to the second reactor.


16. The apparatus for preparing hydrazodicarbonamide according to claim 14, wherein the pyrolysis furnace has a gas injector for injecting inert gas, which does not react with isocyanic acid, into the pyrolysis furnace.

17. The apparatus for preparing hydrazodicarbonamide according to claim 14, wherein the pyrolysis furnace has means for lowering pressure to remove the ammonia from the pyrolysis furnace.

10

DRAWING

FIG. 1



INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR01/01796

A. CLASSIFICATION OF SUBJECT MATTER
IPC7 C07C 225/06
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC07 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
Korean Patents and applications for inventions since 1975, Korean Utility models and applications for Utility models since 1975, Japanese Utility models and applications for Utility models since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS Online (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/49652 A1(J & J CHEMICAL CO., LTD.) 12 July 2001 (See the whole document)	1-13
Y	US 4654441 A (MOORMAN MANUFACTURING COMPANY) 31 March 1987 (See the Abstract)	1-13
A	EP 163526 A1(OLIN CORPORATION) 04 December 1985 (See the whole document)	1-13


Further documents are listed in the continuation of Box C.


See patent family annex.

* Special categories of cited documents:
 "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search
23 JULY 2002 (23.07.2002)

Date of mailing of the international search report
23 JULY 2002 (23.07.2002)

Name and mailing address of the ISA/KR
 Korean Intellectual Property Office
 920 Dunsan-dong, Seo-gu, Daejeon 302-701,
 Republic of Korea
 Facsimile No. 82-42-472-7140

Authorized officer
 KIM, Hee Jin
 Telephone No. 82-42-481-5543


INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR01/01796

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 01/49652 A1	12.07.2001	None	
US 4654441 A	31.05.1987	CA 1242454 A1	27.09.1988
EP 163526 A1	04.12.1985	AU 4208185 A1 JP 61001654 A2	05.12.1985 07.01.1986