



(51) International Patent Classification:

C12Q 1/6816 (2018.01) *C12Q 1/6834* (2018.01)
C12Q 1/6825 (2018.01) *G01N 21/17* (2006.01)

(21) International Application Number:

PCT/US2018/036885

(22) International Filing Date:

11 June 2018 (11.06.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/517,694 09 June 2017 (09.06.2017) US

(71) Applicant: UNIVERSITY OF HOUSTON SYSTEM
[US/US]; 4800 Calhoun Road, Houston, TX 77004 (US).

(72) Inventors: XU, Shoujun; 5711 Cheltenham Drive, Houston, TX 77096 (US). HU, Qiongzhen; 2010 Dumble Street, Apt. 5, Houston, TX 77023 (US). WANG, Yuhong; 5711 Cheltenham Drive, Houston, TX 77096 (US).

(74) Agent: AMINI, Farhang; Winstead PC, P.O. Box 131851, Dallas, TX 75313 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

(54) Title: FORCE-MODULATED HYBRIDIZATION FOR VISUALIZING NUCLEIC ACID LENGTH AND FUNCTION

(57) Abstract: Embodiments of the present disclosure pertain to methods of utilizing force-modulated hybridization to determine the length of an analyte strand, to determine an unknown nucleic acid sequence, or to determine the binding of a nucleotide to an active agent. Additional embodiments of the present disclosure pertain to sample holder devices and methods of utilizing such devices. Further embodiments of the present disclosure pertain to detection devices.




FIG. 1A

FORCE-MODULATED HYBRIDIZATION FOR VISUALIZING NUCLEIC ACID LENGTH AND FUNCTION

CROSS-REFERENCE TO RELATED APPLICATIONS

5 [0001] This application claims priority to U.S. Provisional Patent Application No. 62/517,694, filed on June 9, 2017. The entirety of the aforementioned application is incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

10 [0002] This invention was made with government support under grant number 58077, awarded by the National Institutes of Health (NIH); and grant number 57787, awarded by the National Science Foundation. The government has certain rights in the invention

FIELD OF THE INVENTION

[0003] The present disclosure relates to methods and devices for determining a nucleic acid's length and function.

BACKGROUND OF THE DISCLOSURE

15 [0004] All publications herein are incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information
20 provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0005] Nucleic acids, which carry genetic information for all life forms, are ubiquitously involved in biological functions. Measuring their exact interacting sites with other biological entities is therefore fundamental to investigating their roles in various functions. Examples
25 include the binding sites of enzymes and inhibitors, and the dynamic positions of mRNA relative to the ribosome. Thus, there remains a need in the art for new instrumentation and methods for precisely measuring nucleic acids, and ideally achieve single nucleotide (nt) resolution.

SUMMARY OF THE DISCLOSURE

[0006] In some embodiments, the present disclosure pertains to methods of determining the length of an analyte strand by (a) incubating a magnetically labeled oligonucleotide strand, the analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture, where the magnetically labeled oligonucleotide strand is complementary in sequence to the analyte strand, where the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand and include different lengths, and where either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label; (b) transferring the mixture to a surface functionalized to couple with the at least one label; (c) applying a mechanical force to the mixture; and (d) inspecting the surface for immobilized particles.

[0007] If the analyte strand is labeled, then the longest oligonucleotide ruler strand producing immobilized particles on the surface represents the length of the analyte strand. However, if the oligonucleotide ruler strands are labeled, then the longest oligonucleotide ruler strand not producing immobilized particles represents the length of the analyte strand. In some embodiments, a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the length of the analyte strand is determined.

[0008] In some embodiments, the present disclosure pertains to methods of determining an unknown nucleic acid sequence of a magnetically labeled oligonucleotide strand by (a) incubating the magnetically labeled oligonucleotide strand, an analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture, where the analyte strand is complementary in sequence to at least some of the known sequences of the magnetically labeled oligonucleotide strand, where the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand, and where the oligonucleotide ruler strands include nucleic acids at their ends that span a length of the unknown nucleic acid sequence of the magnetically labeled oligonucleotide strand, and where either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label; (b) transferring the mixture to a surface functionalized to couple with the at least one label; (c) applying a mechanical force to the mixture; and (d) inspecting the surface for immobilized particles.

[0009] If the analyte strand is labeled, then the oligonucleotide ruler strand that does not produce immobilized particles on the surface contains at least some of the unknown nucleic

acid sequences at its end. However, if the oligonucleotide ruler strand is labeled, then the oligonucleotide ruler strand that produces immobilized particles on the surface contains at least some of the unknown nucleic acid sequences at its end. In some embodiments, a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the unknown
5 sequence of the magnetically labeled oligonucleotide strand is determined.

[0010] In some embodiments, the present disclosure pertains to methods for determining the binding of a nucleotide to an active agent by (a) incubating a first oligonucleotide strand and a second oligonucleotide strand for hybridization in the presence of an oligonucleotide cleaving enzyme and an active agent to form a mixture, where the first and second
10 oligonucleotide strands are complementary to one another, where at least one of the first or second oligonucleotide strands is labeled with at least one label, and where at least one of the first or second oligonucleotide strands is labeled with at least one magnetic particle to provide a magnetically labeled oligonucleotide strand; (b) transferring the mixture to a surface functionalized to couple with the at least one label; (c) applying a mechanical force
15 to the mixture; and (d) inspecting the surface for immobilized particles.

[0011] The presence of immobilized particles indicates that the active agent binds to the at least one of the first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof. However, the absence of immobilized particles indicates that the active agent does not bind to the at least one of the first oligonucleotide strand, the second
20 oligonucleotide strand, or the hybridized version thereof.

[0012] Additional embodiments of the present disclosure pertain to sample holder devices and methods of utilizing such devices. In some embodiments, the sample holder devices of the present disclosure include: (1) an incubator that includes a plurality of first wells for incubating a plurality of samples; (2) a displayer for displaying the plurality of the samples,
25 where the displayer includes a plurality of second wells, and where a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands; and (3) a transferring apparatus that is capable of transferring the plurality of the samples from the plurality of first wells to the plurality of second wells.

[0013] Additional embodiments of the present disclosure pertain to methods of utilizing the sample holder devices of the present disclosure for transferring a plurality of samples through the following steps: (a) loading the plurality of samples into a plurality of first wells of an
30

incubator; (b) transferring the plurality of the samples from the plurality of first wells of the incubator to a plurality of second wells of a displayer through the utilization of a transferring apparatus, where a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands; and (c) inspecting or analyzing the plurality of second wells of the displayer.

[0014] Additional embodiments of the present disclosure pertain to detection devices. In some embodiments, the detection devices of the present disclosure include: a sample holder; a light source; a voltmeter; a photodetector; and a recording device holder. In some embodiments, the detection devices of the present disclosure also include a translation stage for changing the location of the sample on the sample holder. In some embodiments, the detection devices of the present disclosure also include a mechanical frame for providing structural support to the device. In some embodiments, the detection devices of the present disclosure include a light source; a photodetector; a translation stage; a mechanical frame; and a voltmeter.

DESCRIPTION OF THE DRAWINGS

[0015] FIGURE 1 depicts, in accordance with embodiments herein, several embodiments of the present disclosure. FIG. 1A provides a scheme of a method of determining the length of an analyte strand. FIG. 1B illustrates a scheme for measuring the length of analyte strands. As illustrated, force-modulated hybridization will lead to the presence of immobilized particles when the oligonucleotide ruler strand is not longer than the analyte strand (top image), or to the absence of immobilized particles if the oligonucleotide ruler strand is longer (bottom image). FIG. 1C provides a scheme of a method of determining unknown nucleic acid sequences. FIG. 1D provides an illustration for determining unknown nucleic acid sequences. FIG. 1E provide a scheme of determining active agent-oligonucleotide binding. FIG. 1F illustrates a scheme for determining active agent-oligonucleotide binding, including the identification of the binding site of active agents on oligonucleotides. Immobilized particles indicate specific active agent-oligonucleotide binding at the enzyme binding site (top image). The clear sample indicated no specific active agent-oligonucleotide interactions at the enzyme binding site (bottom image). FIG. 1G shows an image of a sample holder device. FIG. 1H illustrates a scheme related to the operation of the sample holder device. FIG. 1I shows an image of a detection device, which can be useful for multiplexed detection and quantification.

[0016] **FIGURE 2** depicts, in accordance with embodiments herein, validation of force-modulated hybridization to determine nucleic acid length. **FIG. 2A** shows the probing of a 13-nt analyte DNA (denoted as A13) with oligonucleotide ruler strands R15-R12 in TBS buffer. **FIG. 2B** shows measurements by an atomic magnetometer for the four samples in **FIG. 2A**. **FIG. 2C** shows probing a 12-nt analyte DNA (denoted as A12) with oligonucleotide ruler strands R15-R12. **FIG. 2D** shows probing A13 and A12 in TAM10 buffer, using oligonucleotide ruler strands R15-R12. No difference was observed without force modulation. **FIG. 2E** shows distinguishing A13 and A12 by force modulation at 65 pN.

[0017] **FIGURE 3** depicts, in accordance with embodiments herein, RNA detection and label-free analysis. **FIG. 3A** shows a comparison of measuring A13 and A13r using the same set of DNA oligonucleotide ruler strands. **FIG. 3B** shows switching the functionalization from the analyte strand to the oligonucleotide ruler strands enables label-free detection.

[0018] **FIGURE 4** depicts, in accordance with embodiments herein, applications of the DNA ruler method. **FIG. 4A** depicts a sequence for determining the binding site of endonuclease DpnII. **FIG. 4B** shows that the fragment was measured by oligonucleotide ruler strands R15-R11. **FIG. 4C** shows the scheme for distinguishing the Pre and Post-translocation complexes. The ribosome position (indicated by the orange ovals) shifted by 3 nts to the right going from Pre to Post. **FIG. 4D** indicates that using oligonucleotide ruler strand R15' shows particle immobilization for the Pre but not for the Post. A 65 pN force was applied to the samples.

[0019] **FIGURE 5** depicts, in accordance with embodiments herein, resolving drug molecule-DNA binding. **FIG. 5A** shows a DNA duplex, with one end labeled with magnetic particle and the other end labeled with biotin. **FIG. 5B** shows sample images at different combinations of two drug molecules and two enzymes. The two images on the right are for the same samples indicated by the respective arrows, but prior to applying a ~1 pN force. **FIG. 5C** plots the dissociation forces of four different drug molecules interacting with two different sequences.

[0020] **FIGURE 6** depicts, in accordance with embodiments herein, multiplexed analysis and quantification. **FIG. 6A** shows three identical rows of experiments using R15-R11 to probe A13. Scale bar: 4 mm. **FIG. 6B** shows a device to measure the transmission of the sample. I:

multiplexed sample holder; II: laser pointer as light source; III: voltmeter connected to a photodiode beneath the sample; IV: holder for smartphone. **FIG. 6C** shows transmission of samples containing different concentrations of drug molecules. Fitting gives a binding constant of $9 \pm 2 \times 10^4 \text{ M}^{-1}$.

5 **[0021] FIGURE 7** depicts, in accordance with embodiments herein, a unique set of sample holders (**FIG. 7A**) that contains a puncher, incubator, and displayer and their use in analyzing samples (**FIG. 7B**).

[0022] FIGURE 8 depicts, in accordance with embodiments herein, the use of a series of oligonucleotide ruler strands to sequence the unknown nucleotides on a target DNA.

10

DETAILED DESCRIPTION

[0023] All references, publications, and patents cited herein are incorporated by reference in their entirety as though they are fully set forth. Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of
15 ordinary skill in the art to which this invention belongs. Hornyak, et al., Introduction to Nanoscience and Nanotechnology, CRC Press (2008); Singleton et al., Dictionary of Microbiology and Molecular Biology 3rd ed., J. Wiley & Sons (New York, NY 2001); March, Advanced Organic Chemistry Reactions, Mechanisms and Structure 7th ed., J. Wiley & Sons (New York, NY 2013); and Sambrook and Russel, Molecular Cloning: A Laboratory
20 Manual 4th ed., Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY 2012), provide one skilled in the art with a general guide to many of the terms used in the present application. One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. The present invention is in no way limited to the methods and materials described.

25 **[0024]** The terms “polynucleotide” and “oligonucleotide,” used interchangeably herein, refer generally to linear polymers of natural or modified nucleosides, including deoxyribonucleosides, ribonucleosides, alpha-anomeric forms thereof, and the like, usually linked by phosphodiester bonds or analogs thereof ranging in size from a few monomeric units, e.g. 2-4, to several hundreds of monomeric units. When a polynucleotide is represented
30 by a sequence of letters, such as "ATGCCTG," it will be understood that the nucleotides are

in 5'->3' order from left to right. Polynucleotide as used herein also includes a basic sugar-phosphate or sugar- phosphorothioate polymers.

[0025] Measuring the exact interacting sites of nucleic acids with other biological entities is fundamental to investigating their roles in various biological functions. Examples of nucleic acid interactions include the binding sites of enzymes and inhibitors, and the dynamic positions of mRNA relative to the ribosome. Often, to determine the exact interacting site, it is essential to achieve single-nucleotide (nt) resolution. Many techniques have been developed for such measurements, such as surface-enhanced Raman spectroscopy, fluorescence resonance, circular dichroism, electrochemical methods, optical methods, plasmonic resonance, and the utilization of nanopores. In addition, high specificity methods based on thermodynamic equilibrium have been reported.

[0026] However, most of the studies used a mismatching nucleotide in the middle of a sequence to represent single-nt specificity, not at the ends. The latter is much more challenging because the difference in Gibbs free energy is substantially smaller due to duplex end breathing.

[0027] Another challenge for precisely measuring nucleic acids is the requirement of expensive instrumentation. Direct visualization of biological processes has been shown using fluorescent, magnetic, or plasmonic beads, as well as dyes and other nanomaterials. However, the key feature of high specificity has not been preserved well in comparison with sophisticated apparatus. Moreover, resolution on the length of nucleic acids has not been reported using visual or colorimetric detection.

[0028] As described herein, in accordance with the various embodiments herein, the inventors have developed a novel technology of force-modulated hybridization for DNA/RNA detection and analysis, along with the associated device and consumables. The methods and devices provided herein possesses three unique capabilities. First, they are able to measure nucleic acids with single-nucleotide length resolution, which has not been achieved with other methods. Second, they require no amplification and no expensive instrument because the changes in measuring different samples can be visually observed via the color of magnetic labels. Third, a set of specific sample holders can streamline analysis while detection devices can quantify the results when needed.

[0029] Nucleotide length determination

[0030] In some embodiments, the present disclosure pertains to methods of determining the length of an analyte strand. In some embodiments illustrated in **FIG. 1A**, the methods of the present disclosure include the following steps: (a) incubating a magnetically labeled oligonucleotide strand, the analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture, where the magnetically labeled oligonucleotide strand is complementary in sequence to the analyte strand, where the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand and include different lengths, and where either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label (step 10); (b) transferring the mixture to a surface functionalized to couple with the at least one label (step 12); (c) applying a mechanical force to the mixture (step 14); and (d) inspecting the surface for immobilized particles (step 16).

[0031] If the analyte strand is labeled, then the longest oligonucleotide ruler strand producing immobilized particles on the surface represents the length of the analyte strand (steps 18 and 20). However, if the oligonucleotide ruler strands are labeled, then the longest oligonucleotide ruler strand not producing immobilized particles represents the length of the analyte strand (steps 22 and 24). In some embodiments, a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the length of the analyte strand is determined.

[0032] In a more specific embodiment illustrated in **FIG. 1B**, the nucleotide length determination methods of the present disclosure include a step of incubating analyte strand 34 with magnetically labeled oligonucleotide strand 32 and oligonucleotide ruler strand 36(a) or 36(b) to form a mixture 30. In this embodiment, analyte strand 34 is labeled with at least one label 35 while magnetically labeled oligonucleotide strand 32 contains magnetic particle 33.

[0033] Thereafter, mixture 30 is transferred to a surface that is functionalized to couple with label 35. Next, a mechanical force is applied to the mixture. The color of the surface is then inspected. If the surface is yellow (i.e., surface 38), then one can determine that analyte strand 34 is longer than or the same length as the oligonucleotide ruler strand (e.g., oligonucleotide ruler strand 36(a)). The basis is that, due to its length, the oligonucleotide ruler strand is not able to compete with analyte strand 34 for hybridization with magnetically labeled oligonucleotide strand 32. As such, label 35 on analyte strand 34 is able to

immobilize the hybridized oligonucleotide onto a surface while magnetic particle 33 on magnetically labeled oligonucleotide strand 32 is able to generate a yellow color on the surface.

5 [0034] On the other hand, if the surface is colorless or not yellow (i.e., surface 40), then one can determine that the analyte strand is shorter than the oligonucleotide ruler strand (e.g., oligonucleotide ruler strand 36(b)). The basis is that, due to its longer length, the oligonucleotide ruler strand is able to compete with analyte strand 34 for hybridization with magnetically labeled oligonucleotide strand 32. As such, analyte strand 34 and magnetically labeled oligonucleotide strand 32 do not form a substantial amount of hybridized
10 oligonucleotides that could be immobilized onto a surface through the use of label 35 on analyte strand 34. Accordingly, magnetic particle 33 on magnetically labeled oligonucleotide strand 32 is not able to generate a yellow color on the surface.

[0035] As set forth in more detail herein, the nucleotide length determination methods of the present disclosure can have numerous embodiments. In particular, various magnetically
15 labeled oligonucleotide strands, analyte strands and oligonucleotide ruler strands may be incubated under various conditions and transferred to various types of functionalized surfaces. Moreover, various types of mechanical forces may be applied to the mixture of oligonucleotides. Surfaces may also be inspected in various manners to detect immobilized particles.

20 [0036] Analyte strands

[0037] The nucleotide length determination methods of the present disclosure can be utilized to determine the length of various types of analyte strands. For instance, in some embodiments, the analyte strand is a DNA strand. In some embodiments, the analyte strand is an RNA strand.

25 [0038] The analyte strands of the present disclosure can include various lengths. For instance, in some embodiments, the analyte strands of the present disclosure are 10-50 nucleotides long. In some embodiments, the analyte strands of the present disclosure are 10-30 nucleotides long.

[0039] Magnetically labeled oligonucleotide strands

[0040] The nucleotide length determination methods of the present disclosure may utilize various types of magnetically labeled oligonucleotide strands. For instance, in some embodiments, the magnetically labeled oligonucleotide strand is a DNA strand. In some embodiments, the magnetically labeled oligonucleotide strand is an RNA strand.

5 [0041] The magnetically labeled oligonucleotide strands of the present disclosure can include various lengths. For instance, in some embodiments, the magnetically labeled oligonucleotide strands of the present disclosure are 10-50 nucleotides long. In some
10 embodiments, the magnetically labeled oligonucleotide strands of the present disclosure are 12-50 nucleotides long. In some embodiments, the magnetically labeled oligonucleotide strands of the present disclosure are 12-30 nucleotides long.

[0042] In some embodiments, the magnetically labeled oligonucleotide strands of the present disclosure have longer lengths than the analyte strand. In some embodiments, the magnetically labeled oligonucleotide strands have the same length as the analyte strand. In
15 some embodiments, the magnetically labeled oligonucleotide strands have a shorter length than the analyte strand.

[0043] The magnetically labeled oligonucleotide strands of the present disclosure may be labeled with a magnetic particle. In some embodiments, the magnetic particles may include particles that have magnetic properties. For instance, in some embodiments, the magnetic
20 particles include, without limitation, ferromagnetic materials, ferrimagnetic materials, and combinations thereof. In some embodiments, the magnetic particles include, without limitation, iron, nickel, cobalt, and combinations thereof.

[0044] The magnetic particles of the present disclosure can have various sizes. For instance, in some embodiments, the magnetic particles have a diameter ranging from about 10 nm to about 5000 nm. In some embodiments, the magnetic particles have a diameter ranging from
25 about 10 nm to about 1,000 nm. In some embodiments, the magnetic particles have a diameter ranging from about 10 nm to about 500 nm.

[0045] The magnetically labeled particles of the present disclosure may be positioned at various locations on a magnetically labeled oligonucleotide strand. For instance, in some
30 embodiments, the magnetically labeled particles are positioned at one of the ends of a magnetically labeled oligonucleotide strand. In some embodiments, the magnetically labeled

particles are positioned at a central or median position on a magnetically labeled oligonucleotide strand.

[0046] Oligonucleotide ruler strands

[0047] The nucleotide length determination methods of the present disclosure may also
5 utilize various types of oligonucleotide ruler strands. For instance, in some embodiments, the oligonucleotide ruler strands of the present disclosure include DNA strands, RNA strands, and combinations thereof. In some embodiments, the oligonucleotide ruler strands of the present disclosure include DNA strands. In some embodiments, the oligonucleotide ruler strands of the present disclosure include RNA strands.

10 **[0048]** In some embodiments, the oligonucleotide ruler strands of the present disclosure are complementary in sequence to the analyte strand. In some embodiments, the oligonucleotide ruler strands include oligonucleotides longer than the analyte strand. In some embodiments, the oligonucleotide ruler strands include oligonucleotides shorter than the analyte strand. In some embodiments, the oligonucleotide ruler strands include oligonucleotides the same
15 length as the analyte strand.

[0049] Labels

[0050] The nucleotide length determination methods of the present disclosure require that either the analyte strand or the oligonucleotide ruler strand be labeled with at least one label. In some embodiments, the analyte strand is labeled with the at least one label. In such
20 embodiments, the oligonucleotide ruler strands are not labeled with the at least one label.

[0051] In some embodiments, the oligonucleotide ruler strands are labeled with the at least one label. In such embodiments, the analyte strand is not labeled with the at least one label.

[0052] The labels of the present disclosure can include labels that could be utilized to couple an analyte strand or an oligonucleotide strand to a functionalized surface. In some
25 embodiments, the at least one label includes, without limitation, biotin, streptavidin, digoxigenin, avidin, maleic imide, gold, proteins, nucleic acids, functional groups, and combinations thereof. In some embodiments, the at least one label is biotin. In some embodiments, the label excludes magnetic materials, such as magnetic particles. The use of additional labels can also be envisioned.

[0053] Incubation of oligonucleotides

[0054] Various methods may be utilized to incubate magnetically labeled oligonucleotide strands, analyte strands, and an oligonucleotide ruler strand to form a mixture. For instance, in some embodiments, the incubation occurs by first incubating the magnetically labeled
5 oligonucleotide strand and the analyte strand and then adding the oligonucleotide ruler strand. In some embodiments, the magnetically labeled oligonucleotide strand, the analyte strand and the oligonucleotide ruler strand are incubated at the same time.

[0055] Incubation can occur under various conditions. For instance, in some embodiments, incubation can occur at temperatures between 5-50 °C. In some embodiments, incubation
10 occurs at room temperature.

[0056] Incubation can also occur for various periods of time. For instance, in some embodiments, the incubation duration is between 1-300 minutes. In some embodiments, the incubation duration is about 1 minute.

[0057] Incubation can occur in various environments. For instance, in some embodiments, incubation can occur in a well. In some embodiments, incubation can occur in one or more
15 wells of a sample holder device of the present disclosure (e.g., one or more of the first wells 125 of incubator 124 of sample holder device 120, as shown in **FIG. 1G** and described in more detail herein).

[0058] In some embodiments, the incubation step is separate and apart from the transferring step. In other embodiments, the incubation step and the transferring step occur at the same
20 time. For instance, in some embodiments, incubation occurs when a mixture is transferred to a surface.

[0059] Transfer to surfaces

[0060] The mixtures of the present disclosure can be transferred to various surfaces. The
25 Surfaces of the present disclosure generally include surfaces that are functionalized to couple with the at least one label. For instance, in some embodiments, the surface is functionalized with a molecule that binds to the at least one label. In some embodiments where the at least one label is biotin, the molecule can be streptavidin. In some embodiments where the at least

one label is streptavidin, the molecule can be biotin. In some embodiments where the at least one label is gold, the molecule can be a thiol.

5 [0061] The surfaces of the present disclosure can include various structures. For instance, in some embodiments, the surface represents a bottom portion of a well. In some embodiments, the surface includes one or more wells of a sample holder device of the present disclosure (e.g., one or more of the second wells 127 of displayer 126 of sample holder device 120, as shown in **FIG. 1G** and described in more detail herein). Additional structures can also be envisioned.

10 [0062] Various methods may also be utilized to transfer mixtures to the surfaces of the present disclosure. For instance, in some embodiments, the transfer occurs by pouring the mixture onto the surface. In some embodiments, the transfer occurs by utilizing a transferring apparatus, such as a pipette. In some embodiments, the transfer occurs by utilizing the sample transfer methods of the present disclosure (e.g., sample transfer methods illustrated in **FIG. 1H** and described in more detail herein).

15 [0063] Mechanical forces

[0064] Various types of mechanical forces may be applied to the mixtures of the present disclosure. For instance, in some embodiments, the mechanical force includes, without limitation, gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof. In some embodiments, the mechanical force
20 includes centrifugal force.

[0065] In some embodiments, the mechanical force is applied between 10 fN and 500 pN. In some embodiments, the mechanical force is applied between 1 pN and 100 pN. In some embodiments, the mechanical force is applied as a 65 pN centrifugal force.

25 [0066] In some embodiments, the mechanical force removes oligonucleotides from the mixture that are not specifically bound to the surface. For instance, in some embodiments, gravity force may be applied to a mixture by placing the mixture vertically to remove the nonspecifically bound oligonucleotides.

[0067] Without being bound by theory, it is envisioned that the application of a mechanical force can have various effects on the mixtures of the present disclosure. For instance, in

some embodiments, the mechanical force modulates oligonucleotide hybridization. In some embodiments, the mechanical force modulates oligonucleotide hybridization between an analyte strand and a magnetically labeled oligonucleotide strand. In some embodiments, the mechanical force modulates oligonucleotide hybridization between an oligonucleotide ruler strand and a magnetically labeled oligonucleotide strand.

[0068] In some embodiments, the oligonucleotide hybridization results in the presence of immobilized particles on a surface when the analyte strand is labeled and the oligonucleotide ruler strand is not longer than the analyte strand (e.g., analyte strand 34 and oligonucleotide strand 36(a), as illustrated in **FIG. 1B**). In such embodiments, the immobilized particles represent a hybrid structure between the magnetically labeled oligonucleotide strand and the analyte strand (e.g., magnetically labeled oligonucleotide strand 32 and analyte strand 34, as illustrated in **FIG. 1B**).

[0069] In some embodiments, the oligonucleotide hybridization results in the absence of immobilized particles on a surface when the analyte strand is labeled and the oligonucleotide ruler strand is longer than the analyte strand (e.g., analyte strand 34 and oligonucleotide strand 36(b), as illustrated in **FIG. 1B**). In such embodiments, the oligonucleotide ruler strand hybridizes with the magnetically labeled oligonucleotide strand and prevents the labeled analyte strand from substantially hybridizing with the magnetically labeled oligonucleotide strand. As such, the labeled analyte strand is not able to produce immobilized particles on a surface.

[0070] In some embodiments, the oligonucleotide hybridization results in the presence of immobilized particles on a surface when the oligonucleotide ruler strands are labeled and the oligonucleotide ruler strand is longer than the analyte strand. In such embodiments, the immobilized particles represent a hybrid structure between the magnetically labeled oligonucleotide strand and the oligonucleotide ruler strand.

[0071] In some embodiments, oligonucleotide hybridization results in the absence of immobilized particles when the oligonucleotide ruler strands are labeled and the oligonucleotide ruler strand is shorter than or the same length as the analyte strand. In such embodiments, the unlabeled analyte strand hybridizes with the magnetically labeled oligonucleotide strand and prevents the labeled oligonucleotide ruler strand from

substantially hybridizing with the magnetically labeled oligonucleotide strand. As such, the labeled oligonucleotide ruler strand is not able to produce immobilized particles on a surface.

[0072] Inspection of surfaces

[0073] Various methods may also be utilized to inspect surfaces for immobilized particles.

5 For instance, in some embodiments, the inspection includes, without limitation, visual inspection, measurement of light transmission, measurement by a magnetic sensor, or combinations thereof.

[0074] In some embodiments, the inspection occurs by visual inspection. In some embodiments, inspection occurs by the utilization of a device, such as the detection devices
10 of the present disclosure (e.g., detection device 150, as shown in **FIG. 1I** and described in more detail herein).

[0075] In some embodiments, inspection may occur after the application of a mechanical force. In some embodiments, the surface to be inspected may be washed prior to the inspection. In some embodiments, the washing step removes non-specifically bound
15 oligonucleotides from a surface.

[0076] In some embodiments, inspection results in the detection of immobilized particles on a surface. In some embodiments, immobilized particles are represented by a property of the magnetic particles on the magnetically labeled oligonucleotide strands. For instance, in some embodiments, the immobilized particles are represented by a yellow color on the surface
20 (e.g., yellow color on surface 38, as shown in **FIG. 1B**). In some embodiments, the yellow color is derived from the magnetically labeled oligonucleotide strand, such as the magnetic particles associated with the magnetically labeled oligonucleotide strand.

[0077] Repetition of steps

[0078] In some embodiments, the steps outlined in **FIG. 1A** may be repeated multiple times.
25 For instance, in each repetition step, a different oligonucleotide ruler strand may be used to repeat steps (a)-(d) until the length of the analyte strand is determined. In some embodiments, steps (a)-(d) are repeated sequentially. In some embodiments, steps (a)-(d) are repeated in parallel or simultaneously, such as through the use of a multi-well system.

[0079] Determination of unknown nucleic acid sequences

[0080] In some embodiments, the present disclosure pertains to methods of determining an unknown nucleic acid sequence of a magnetically labeled oligonucleotide strand. In some embodiments illustrated in **FIG. 1C**, the methods of the present disclosure include the following steps: (a) incubating the magnetically labeled oligonucleotide strand, an analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture, where the analyte strand is complementary in sequence to at least some of the known sequences of the magnetically labeled oligonucleotide strand, where the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand, and where the oligonucleotide ruler strands include nucleic acids at their ends that span a length of the unknown nucleic acid sequence of the magnetically labeled oligonucleotide strand, and where either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label (step 50); (b) transferring the mixture to a surface functionalized to couple with the at least one label (step 52); (c) applying a mechanical force to the mixture (step 54); and (d) inspecting the surface for immobilized particles (step 56).

[0081] If the analyte strand is labeled (step 58), then the oligonucleotide ruler strand that does not produce immobilized particles on the surface contains at least some of the unknown nucleic acid sequences at its end (step 60). However, if the oligonucleotide ruler strand is labeled (step 62), then the oligonucleotide ruler strand that produces immobilized particles on the surface contains at least some of the unknown nucleic acid sequences at its end (step 64). In some embodiments, a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the unknown sequence of the magnetically labeled oligonucleotide strand is determined.

[0082] In a more specific embodiment illustrated in **FIG. 1D**, a method of determining an unknown nucleic acid sequence of magnetically labeled oligonucleotide strand 72 is disclosed. In this embodiment, magnetically labeled oligonucleotide strand 72, analyte strand 74, and one of a series of oligonucleotide ruler strands 76(a)-(d) are incubated to form a mixture 70.

[0083] As also illustrated in **FIG. 1D**, analyte strand 74 is complementary in sequence to at least some of the known sequences of magnetically labeled oligonucleotide strand 72. Analyte strand 74 is labeled with label 75 while magnetically labeled oligonucleotide strand 72 is labeled with magnetic particle 73. As also shown in **FIG. 1D**, the series of oligonucleotide ruler strands 76(a)-(d) are complementary in sequence to magnetically labeled oligonucleotide strand 72. Oligonucleotide ruler strands 76(a)-(d) also include

nucleic acids at their ends that span a length of the unknown nucleic acid sequence of magnetically labeled oligonucleotide strand.

[0084] Mixture 70 is transferred to a surface functionalized to couple with label 75. Thereafter, a mechanical force is applied to the mixture. The surface is then inspected for
5 immobilized particles. A determination can be made that oligonucleotide ruler strand 76(a) contains at least some of the unknown nucleic acid sequences at its end because it does not produce immobilized particles on surface 78(a) (as determined by the absence of a yellow color from the surface). Similarly, a determination can be made that oligonucleotide ruler
10 strands 76(b)-(d) do not contain unknown nucleic acid sequences at their ends because they produce immobilized particles on surfaces 78(b)-(d) (as determined by the presence of a yellow color on the surface). The aforementioned steps can then be repeated with different oligonucleotide ruler strands until the unknown sequence of magnetically labeled oligonucleotide strand 72 is determined.

[0085] The nucleic acid sequencing methods of the present disclosure can have numerous
15 embodiments. For instance, as set forth previously, various magnetically labeled oligonucleotide strands, analyte strands, and oligonucleotide ruler strands may be incubated with one another under various conditions to form various mixtures. As also set forth previously, either the analyte strand or the oligonucleotide ruler strands may be labeled with various labels. Various methods set forth previously may also be utilized to transfer a
20 mixture to various surfaces. As also set forth previously, various types of mechanical forces may be applied to a mixture. The surface may then be inspected for immobilized particles in various manners that were also set forth previously.

[0086] Additional embodiments of nucleic acid sequencing methods can also be envisioned. For instance, in some embodiments, the magnetically labeled oligonucleotide strand has a
25 segment of known sequence followed by a segment of the unknown nucleic acid sequence. In some embodiments, the unknown nucleic acid sequence is at the end of the magnetically labeled oligonucleotide strand. In some embodiments, the unknown nucleic acid is 1-20 nucleotides long. In some embodiments, the unknown nucleic acid is 1-5 nucleotides long.

[0087] In some embodiments, the oligonucleotide ruler strands include different lengths. In
30 some embodiments, the oligonucleotide ruler strands include oligonucleotide ruler strands

that differ by only the last nucleotide. In some embodiments, the last nucleotide is complementary in sequence to an unknown nucleic acid sequence.

[0088] Active agent-oligonucleotide binding determination

[0089] In some embodiments, the present disclosure pertains to methods for determining the binding of a nucleotide to an active agent. In some embodiments illustrated in **FIG. 1E**, the methods of the present disclosure include the following steps: (a) incubating a first oligonucleotide strand and a second oligonucleotide strand for hybridization in the presence of an oligonucleotide cleaving enzyme and an active agent to form a mixture, where the first and second oligonucleotide strands are complementary to one another, where at least one of the first or second oligonucleotide strands is labeled with at least one label, and where at least one of the first or second oligonucleotide strands is labeled with at least one magnetic particle to provide a magnetically labeled oligonucleotide strand (step 80); (b) transferring the mixture to a surface functionalized to couple with the at least one label (step 82); (c) applying a mechanical force to the mixture (step 84); and (d) inspecting the surface for immobilized particles (step 86).

[0090] The presence of immobilized particles (step 88) indicates that the active agent binds to the at least one of the first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof (step 90). However, the absence of immobilized particles (step 92) indicates that the active agent does not bind to the at least one of the first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof (step 94).

[0091] In a more specific embodiment illustrated in **FIG. 1F**, the methods of the present disclosure include incubating a magnetically labeled first oligonucleotide strand 102 with a second oligonucleotide strand 104 for hybridization in the presence of active agent 106 and oligonucleotide cleaving enzyme 107 to form mixture 100. In this embodiment, second oligonucleotide strand 104 is labeled with label 105 and is complementary to magnetically labeled first oligonucleotide strand 102. In addition, magnetically labeled first oligonucleotide strand 102 is labeled with magnetic particle 103.

[0092] Mixture 100 is then transferred to a surface that is functionalized to couple with label 105. A mechanical force is then applied to the mixture. The color of the surface is then inspected. If the surface is yellow (e.g., surface 108(a)), then a determination is made that

active agent 106 binds to at least one of the magnetically labeled first oligonucleotide strand 102, the second oligonucleotide strand 104, or the hybridized version thereof. However, if the surface is colorless or not yellow (e.g., surface 108(b)), then a determination is made that the active agent does not bind to the at least one of the magnetically labeled first
5 oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof.

[0093] As set forth in more detail herein, the active agent-oligonucleotide binding determination methods of the present disclosure can have numerous embodiments. In particular, the methods can utilize various first oligonucleotide strands, second oligonucleotide strands, oligonucleotide cleaving enzymes, and active agents. In addition,
10 the first and second oligonucleotide strands may be labeled with numerous types of labels and magnetic particles. In addition, the formed mixtures may be transferred to numerous surfaces. Various types of mechanical forces may also be applied to the mixture. In addition, various methods may be utilized to inspect a surface for immobilized particles in order to make an active agent-oligonucleotide binding determination.

15 **[0094] First and second oligonucleotide strands**

[0095] The active agent-oligonucleotide binding determination methods of the present disclosure can utilize various first and second oligonucleotide strands. For instance, in some embodiments, each of the first and second oligonucleotide strands can include DNA strands, RNA strands, and combinations thereof. In some embodiments, the first and second
20 oligonucleotide strands are DNA strands. In some embodiments, the first and second oligonucleotide strands are RNA strands. In some embodiments, one of the first or second oligonucleotide strands is a DNA strand while the complementary oligonucleotide strand is an RNA strand.

[0096] The first and second oligonucleotide strands of the present disclosure can include various lengths. For instance, in some embodiments, each of the first and second oligonucleotide strands of the present disclosure is 10-50 nucleotides long. In some
25 embodiments, each of the first and second oligonucleotide strands of the present disclosure is 10-30 nucleotides long.

[0097] Labels and magnetic particles

[0098] The first and second oligonucleotide strands of the present disclosure may be labeled with various labels and magnetic particles in various manners. For instance, in some embodiments, the same oligonucleotide strand is labeled with the label and the magnetic particle. In some embodiments, one of the first or second oligonucleotide strands is labeled with the label while the complementary oligonucleotide strand is labeled with the magnetic particle (e.g., magnetic particle 103 on first oligonucleotide strand 102 and label 105 on second oligonucleotide strand 104, as shown in **FIG. 1F**).

[0099] In some embodiments, the label and the magnetic particle are on opposite termini of the first and the second oligonucleotide strands (e.g., label 105 and magnetic particle 103, as shown in **FIG. 1F**). In some embodiments, the label and the magnetic particle are on the same termini of the first and the second oligonucleotide strands.

[00100] Labels

[00101] Various labels may be associated with the first or second oligonucleotide strands. Suitable labels were described previously. For instance, in some embodiments, the labels can include labels that could be utilized to couple a first or second oligonucleotide strand to a functionalized surface. In some embodiments, the label includes, without limitation, biotin, streptavidin, digoxigenin, avidin, maleic imide, gold, proteins, nucleic acids, functional groups, and combinations thereof. In some embodiments, the label is biotin. In some embodiments, the label excludes magnetic materials, such as magnetic particles. The use of additional labels can also be envisioned.

[00102] Magnetic particles

[00103] Various magnetic particles may be associated with the first or second oligonucleotide strands. Suitable magnetic particles were also described previously. For instance, in some embodiments, the magnetic particles of the present disclosure may include particles that have magnetic properties. In some embodiments, the magnetic particles include, without limitation, ferromagnetic materials, ferrimagnetic materials, and combinations thereof. In some embodiments, the magnetic particles include, without limitation, iron, nickel, cobalt, and combinations thereof.

[00104] The magnetic particles of the present disclosure can have various sizes. For instance, in some embodiments, the magnetic particles have a diameter ranging from about

10 nm to about 5000 nm. In some embodiments, the magnetic particles have a diameter ranging from about 10 nm to about 1,000 nm. In some embodiments, the magnetic particles have a diameter ranging from about 10 nm to about 500 nm.

[00105] Oligonucleotide cleaving enzymes

- 5 **[00106]** The active agent-oligonucleotide binding determination methods of the present disclosure can also utilize various oligonucleotide cleaving enzymes. For instance, in some embodiments, the oligonucleotide cleaving enzyme includes a DNA or RNA nuclease. In some embodiments, the oligonucleotide cleaving enzyme includes a DNA nuclease. In some embodiments, the oligonucleotide cleaving enzyme includes an RNA nuclease.
- 10 **[00107]** In some embodiments, the oligonucleotide cleaving enzyme is a restriction enzyme. In some embodiments, the restriction enzyme includes, without limitation, DpnII, EcoRI, and combinations thereof.

[00108] Active agent

- 15 **[00109]** The active agent-oligonucleotide binding determination methods of the present disclosure can also utilize various active agents. For instance, in some embodiments, the active agent includes an experimental drug molecule. In some embodiments, the active agent is a molecule with undetermined or unknown nucleotide binding activities. The use of additional active agents can also be envisioned.

[00110] Incubation

- 20 **[00111]** Various methods may be utilized to incubate the mixtures of the present disclosure. Suitable incubation methods were described previously. For instance, in some embodiments, incubation can occur at temperatures between 5-50 °C. In some embodiments, incubation occurs at room temperature.

- 25 **[00112]** Incubation can also occur for various periods of time. For instance, in some embodiments, the incubation duration is between 1-300 minutes. In some embodiments, the incubation duration is about 1 minute.

[00113] Incubation can occur in various environments. For instance, in some embodiments, incubation can occur in a well. In some embodiments, incubation can occur in one or more

wells of a sample holder device of the present disclosure (e.g., one or more of the first wells 125 of incubator 124 of sample holder device 120, as shown in **FIG. 1G** and described in more detail herein).

5 [00114] In some embodiments, the incubation step is separate and apart from the transferring step. In other embodiments, the incubation step and the transferring step occur at the same time. For instance, in some embodiments, incubation occurs when a mixture is transferred to a surface.

[00115] Transfer to surfaces

10 [00116] The mixtures of the present disclosure can be transferred to various surfaces. The Surfaces of the present disclosure generally include surfaces that are functionalized to couple with the at least one label. Suitable surfaces were described previously. For instance, in some embodiments, the surface is functionalized with a molecule that binds to the at least one label. In some embodiments where the at least one label is biotin, the molecule can be streptavidin. In some embodiments where the at least one label is streptavidin, the molecule
15 can be biotin.

[00117] The surfaces of the present disclosure can include various structures. For instance, in some embodiments, the surface represents a bottom portion of a well. Additional structures can also be envisioned. In some embodiments, the surface includes one or more wells of a sample holder device of the present disclosure (e.g., one or more of the second
20 wells 127 of displayer 126 of sample holder device 120, as shown in **FIG. 1G** and described in more detail herein). Additional structures can also be envisioned.

[00118] Various methods may also be utilized to transfer mixtures to the surfaces of the present disclosure. For instance, in some embodiments, the transfer occurs by pouring the mixture onto the surface. In some embodiments, the transfer occurs by utilizing a
25 transferring apparatus, such as a pipette. In some embodiments, the transfer occurs by utilizing the sample transfer methods of the present disclosure (e.g., sample transfer methods illustrated in **FIG. 1H** and described in more detail herein).

[00119] Mechanical forces

[00120] Various types of mechanical forces may be applied to the mixtures of the present disclosure. Suitable mechanical forces were described previously. For instance, in some embodiments, the mechanical force includes, without limitation, gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof. In some embodiments, the mechanical force includes centrifugal force.

[00121] In some embodiments, the mechanical force is applied between 10 fN and 500 pN. In some embodiments, the mechanical force is applied between 1 pN and 100 pN. In some embodiments, the mechanical force is applied as a 65 pN centrifugal force.

[00122] In some embodiments, the mechanical force removes oligonucleotides from the mixture that are not specifically bound to the surface. For instance, in some embodiments, gravity force may be applied to a mixture by placing the mixture vertically to remove the nonspecifically bound oligonucleotides.

[00123] Without being bound by theory, it is envisioned that the application of a mechanical force can have various effects on the mixtures of the present disclosure. For instance, in some embodiments, the mechanical force modulates oligonucleotide hybridization. In some embodiments, the mechanical force modulates oligonucleotide hybridization between a first oligonucleotide strand and a second oligonucleotide strand.

[00124] In some embodiments, the oligonucleotide hybridization results in the presence of immobilized particles on a surface when the active agent prevents the oligonucleotide cleaving enzyme from cleaving the first and second oligonucleotides. In such embodiments, the immobilized particles represent a hybrid structure between the first oligonucleotide strand and the second oligonucleotide strand (e.g., oligonucleotide strand 102 and second oligonucleotide strand 104, as illustrated in **FIG. 1F**). The label on the first or second oligonucleotide strand immobilizes the hybrid structure onto a surface while the magnetic particle on the first or second oligonucleotide strand is utilized to detect the immobilized particle.

[00125] In some embodiments, the methods of the present disclosure result in the absence of immobilized particles on a surface when the active agent cannot prevent the oligonucleotide cleaving enzyme from cleaving the first and second oligonucleotides. In such embodiments, the first and/or second oligonucleotide are cleaved by the active agent. As such, the label on

the first or second oligonucleotide strand is not able to immobilize the hybrid structure onto a surface to form immobilized particles.

[00126] Inspection of surfaces

[00127] Various methods may also be utilized to inspect surfaces for immobilized particles.

5 Suitable inspection methods were also described previously. For instance, in some embodiments, the inspection includes, without limitation, visual inspection, measurement of light transmission, measurement by a magnetic sensor, or combinations thereof.

[00128] In some embodiments, the inspection occurs by visual inspection. In some embodiments, inspection occurs by the utilization of a device, such as the detection devices
10 of the present disclosure (e.g., detection device 150, as shown in **FIG. 1I** and described in more detail herein).

[00129] In some embodiments, inspection may occur after the application of a mechanical force. In some embodiments, the surface to be inspected may be washed prior to the inspection. In some embodiments, the washing step removes non-specifically bound
15 oligonucleotides from a surface.

[00130] In some embodiments, inspection results in the detection of immobilized particles on a surface. In some embodiments, immobilized particles are represented by a property of the magnetic particles on the first or second oligonucleotides. For instance, in some
20 embodiments, the immobilized particles are represented by a yellow color on the surface (e.g., yellow color on surface 108(a), as shown in **FIG. 1F**).

[00131] Determining active agent binding site on oligonucleotides

[00132] The active agent-oligonucleotide binding determination methods of the present disclosure can have various applications. For instance, in some embodiments, such methods can be utilized as an active agent screening assay for identifying active agents that bind to a
25 nucleotide. In some embodiments, the active agent binding site represents the enzyme binding site. In some embodiments, no active agent is present for control experiments.

[00133] Sample holder devices

[00134] Additional embodiments of the present disclosure pertain to sample holder devices and methods of utilizing such devices. In some embodiments, the sample holder devices of the present disclosure include: (1) an incubator that includes a plurality of first wells for incubating a plurality of samples; (2) a displayer for displaying the plurality of the samples, where the displayer includes a plurality of second wells, and where a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands; and (3) a transferring apparatus that is capable of transferring the plurality of the samples from the plurality of first wells to the plurality of second wells.

[00135] A more specific embodiment of a sample holder device is illustrated in **FIG. 1G** as sample holder device 120. In this embodiment, sample holder device 120 includes: (1) an incubator 124 that includes a plurality of first wells 125 for incubating a plurality of samples; (2) a displayer 126 for displaying the plurality of the samples, where the displayer includes a plurality of second wells 127, and where a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands; and (3) a transferring apparatus 122, that is capable of transferring the plurality of the samples from the plurality of first wells 125 to the plurality of second wells 127.

[00136] As set forth in more detail herein, the sample holder devices of the present disclosure can have numerous embodiments. For instance, in some embodiments, the plurality of second wells are superimposable on the plurality of first wells. In some embodiments, the surface region of the plurality of second wells is the bottom portion on each of the plurality of the second wells.

[00137] Moreover, the surfaces of the plurality of second wells may be functionalized with various functional groups. In some embodiments, the functional groups include functional groups that are suitable for binding to the labels that were described previously, such as biotin. In some embodiments, the functional groups are a small molecule. In some embodiments, the functional groups are streptavidin.

[00138] The sample holder devices of the present disclosure can also include various transferring apparatus. For instance, in some embodiments, the transferring apparatus is in the form of a perforator (e.g., perforator 122 shown in **FIG. 1G**). In some embodiments, the perforator is capable of breaking or perforating a surface of the plurality of first wells so that the plurality of the samples from the plurality of first wells are transferred to the plurality of

second wells. In some embodiments, the perforator includes a matrix of sharp tips (e.g., sharp tips 123 shown in **FIG. 1G**) at positions corresponding to the plurality of first wells.

[00139] In some embodiments, the transferring apparatus includes a sample transport system. In some embodiments, the sample transport system is capable of transporting the plurality of the samples from the plurality of first wells to the plurality of second wells. In some
5 embodiments, the sample transport system includes a multi-head pipette system. In some embodiments, the sample transport system includes a robotic arm.

[00140] The sample holder devices of the present disclosure can be in various forms. For instance, in some embodiments, the sample holder devices of the present disclosure can be in
10 the form of a kit for oligonucleotide hybridization. In some embodiments, the kit is utilized to practice the methods of the present disclosure, such as methods of determining the length of an analyte strand, methods of determining an unknown nucleic acid sequence of a magnetically labeled oligonucleotide strand, and methods for determining the binding of a nucleotide to an active agent.

15 **[00141] Methods of transferring samples**

[00142] Additional embodiments of the present disclosure pertain to methods of utilizing the sample holder devices of the present disclosure for transferring a plurality of samples. In some embodiments illustrated in **FIG. 1H**, the methods of the present disclosure include the following steps: (a) loading the plurality of samples into a plurality of first wells of an
20 incubator (step 130); (b) transferring the plurality of the samples from the plurality of first wells of the incubator to a plurality of second wells of a displayer through the utilization of a transferring apparatus, where a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands (step 132); and (c) inspecting or analyzing the plurality of second wells of the displayer (step 134).

25 **[00143]** In some embodiments where the transferring apparatus includes a perforator, the transferring step can include utilizing the perforator to break or perforate the surface of the plurality of first wells so that the plurality of the samples from the plurality of first wells are transferred to the plurality of second wells. In such embodiments, the methods of the present disclosure can also include a step of aligning the displayer below the incubator prior to

breaking or perforating, such that the aligning places the plurality of second wells directly below the plurality of corresponding first wells.

[00144] In some embodiments where the transferring apparatus includes a sample transport system, the transfer step can include a step of transporting the plurality of the samples from the plurality of first wells to the plurality of second wells. In some embodiments, the transport can occur by the utilization of a multi-head pipette system. In some embodiments, the transport can occur by the utilization of a robotic arm.

[00145] In some embodiments, the sample transfer methods of the present disclosure may also include a step of applying mechanical force to the plurality of second wells. In some embodiments, the mechanical force is applied prior to inspecting or analyzing the second wells. In some embodiments, the mechanical force is applied prior to transferring the samples from the plurality of the first wells to the plurality of the second wells. In some embodiments, the mechanical force is applied after transferring the samples from the plurality of the first wells to the plurality of the second wells.

[00146] Various types of mechanical forces can be applied to the samples. Suitable mechanical forces were described previously. For instance, in some embodiments, the mechanical force is in the form of gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof.

[00147] **Detection Devices**

[00148] Additional embodiments of the present disclosure pertain to detection devices. In some embodiments, the detection devices of the present disclosure include: a sample holder; a light source; a voltmeter; a photodetector; and a recording device holder. In some embodiments, the detection devices of the present disclosure also include a translation stage for changing the location of the sample on the sample holder. In some embodiments, the detection devices of the present disclosure also include a mechanical frame for providing structural support to the device. In some embodiments, the detection devices of the present disclosure include a light source; a photodetector; a translation stage; a mechanical frame; and a voltmeter.

[00149] An example of a detection device is illustrated as detection device 150 in **FIG. 11**. In this embodiment, detection device 150 includes sample holder 152 for holding sample

153; translation stage 154 for changing the location of sample 153 on sample holder 154; photodetector 155 beneath sample holder 152; voltmeter 156 connected to photodetector 155 through wiring 157; light source 158 positioned above sample holder 152; recording device holder 159 for holding recording device 160; and mechanical frame 161 for providing
5 structural support to detection device 150. As set forth in more detail herein, the detection devices of the present disclosure can have numerous embodiments.

[00150] Sample holders

[00151] The detection devices of the present disclosure can include various sample holders. For instance, in some embodiments, the sample holder is a multiplexed sample holder. In
10 some embodiments, the multiplexed sample holder is capable of holding a plurality of samples, such as samples positioned in a plurality of wells (e.g., samples in a plurality of second wells 127 of displayer 126, as shown in **FIG. 1G**).

[00152] Light source

[00153] The detection devices of the present disclosure can also include various light
15 sources. For instance, in some embodiments, the light source includes a laser pointer. In some embodiments, the light source includes a laser diode.

[00154] Light sources may be positioned on various regions of a detection device. For instance, in some embodiments, the light source is positioned above a sample holder (e.g., light source 158 positioned above sample holder 152, as shown in **FIG. 1I**).

20 **[00155]** Photodetector

[00156] The detection devices of the present disclosure can also include various photodetectors. For instance, in some embodiments, the photodetectors include photodiodes. The use of additional photodetectors can also be envisioned.

[00157] The photodetectors of the present disclosure can be positioned at various locations of
25 a detection device. For instance, in some embodiments, a photodetector may be beneath a sample holder (e.g., photodetector 155 beneath sample holder 152, as shown in **FIG. 1I**). In some embodiments, the photodetector may be connected to a voltmeter, such as through a wiring (e.g., photodetector 155 connected to voltmeter 156 through wiring 157).

[00158] Recording device holders

[00159] The detection devices of the present disclosure can also include various recording device holders. Recording device holders generally refer to objects that are capable of holding recording devices, such as cameras, video recorders, and smartphones.

5 [00160] In some embodiments, recording device holders have a structure that is capable of supporting the recording device (e.g., recording device holder 159 shown in **FIG. 1I**). In some embodiments, the recording device holders are positioned above a sample holder such that the recording device is capable of recording or taking photographs of the sample (e.g., recording device holder 159 above sample holder 152).

10 [00161] Translation stages

[00162] The detection devices of the present disclosure can also include various translation stages. Translation stages generally refer to structures that are capable of changing the location of a sample on a sample holder (e.g., translation stage 154 for changing the location of sample 153 on sample holder 152, as shown in **FIG. 1I**).

15 [00163] In some embodiments, the translation stage is a one-dimensional manual stage. In some embodiments, the translation stage is a one-dimensional manual automated stage. In some embodiments, the translation stage is a two-dimensional manual stage (e.g., translation stage 154 shown in **FIG. 1I**). In some embodiments, the translation stage is a two-dimensional automated stage.

20 [00164] Mechanical frame

[00165] The detection devices of the present disclosure can also include various mechanical frames. Mechanical frames generally refer to objects that are capable of providing structural support to the detection devices of the present disclosure. In some embodiments, the mechanical frames of the present disclosure may be in the form of a table (e.g., mechanical
25 frame 161 shown in **FIG. 1I**). Additional mechanical frame structures can also be envisioned.

[00166] Operation of detection devices

[00167] The detection devices of the present disclosure may be operated in various manners and for various purposes. For instance, in some embodiments that refer to detection device 150 in **FIG. 1I** for exemplary purposes only, sample 153 is placed on sample holder 152. Translation stage 154 can then be utilized to change the location of sample 153 on sample holder 154. For instance, translation stage 154 can be utilized to place sample 153 directly beneath light source 158.

[00168] Thereafter, light source 158 transmits light to sample 153. Photodetector 155 then transmits the transmitted light to voltmeter 156 through wiring 157. Moreover, recording device 160 on recording device holder 159 may record or take photos of sample 153. In addition, mechanical frame 161 provides structural support to detection device 150 throughout the aforementioned processes.

[00169] As such, the detection devices of the present disclosure can have numerous applications. For instance, in some embodiments, the detection devices of the present disclosure may be useful for multiplexed detection and quantification. In some embodiments, the detection devices of the present disclosure may be utilized to analyze samples in the sample holder devices of the present disclosure (e.g., sample device holder 120 shown in **FIG. 1G**). In some embodiments, the detection devices of the present disclosure may be utilized to practice the methods of the present disclosure, such as methods of determining the length of an analyte strand, methods of determining an unknown nucleic acid sequence of a magnetically labeled oligonucleotide strand, and methods for determining the binding of a nucleotide to an active agent.

[00170] **Applications and Advantages**

[00171] The devices and methods of the present disclosure have numerous applications and advantages. For instance, the nucleotide length determination methods of the present disclosure are capable of measuring the exact length of nucleic acids. Such capability can find numerous applications in biosensing, such as miRNA detection, as well as fundamental research on DNA and RNA functions.

[00172] Likewise, the active agent-oligonucleotide binding determination methods of the present disclosure can be used to precisely determine the binding site of drug molecules interacting with nucleic acids with high-throughput capacity.

[00173] Moreover, the methods of the present disclosure are versatile and tunable for use in a wide range of biological conditions. For instance, in some embodiments, force-modulated hybridization conditions can be experimentally determined for each and every system that contains different components (e.g., buffers, ion species, ion concentrations, and types of
5 oligonucleotides).

[00174] Moreover, the devices and methods of the present disclosure can eliminate the need for expensive apparatus because single-nucleotide resolution can be observed directly by the naked eye in some embodiments. If desired, quantification can then be achieved by utilizing readily accessible detection devices for measuring light transmission, such as the detection
10 devices of the present disclosure. Moreover, a special set of sample holders, such as the sample holder devices of the present disclosure, can be designed to facilitate analysis.

[00175] As such, depending on a desired application, various aspects of the present disclosure can be combined in various manners. For instance, in some embodiments, the present disclosure can be directed to a kit to a method and device for precisely visualizing a
15 nucleic acid's length and function. In some embodiments, the kit can be useful for practicing the inventive method of precisely visualizing a nucleic acid's length and function, and/or measuring drug-DNA binding.

[00176] In some embodiments, the kit is an assemblage of materials or components, including at least one of methods and devices of the present disclosure. Thus, in some
20 embodiments, the kit can contain a composition including the oligonucleotides of the present disclosure, as described above.

[00177] The exact nature of the components configured in the kit depends on its intended purpose. For example, some embodiments may be configured for the purpose of treating a disease. In one embodiment, the kit is configured particularly for the purpose of treating
25 mammalian subjects. In another embodiment, the kit is configured particularly for the purpose of treating human subjects. In further embodiments, the kit is configured for veterinary applications, treating subjects such as, but not limited to, farm animals, domestic animals, and laboratory animals.

[00178] Instructions for use may be included in the kit. "Instructions for use" typically
30 include a tangible expression describing the technique to be employed in using the

components of the kit to effect a desired outcome, such as to precisely visualize a nucleic acid's length and function and/or determine drug-nucleic acid binding. Optionally, the kit also contains other useful components, such as, diluents, buffers, pharmaceutically acceptable carriers, syringes, catheters, applicators, pipetting or measuring tools, bandaging materials or
5 other useful paraphernalia as will be readily recognized by those of skill in the art.

[00179] The materials or components assembled in the kit can be provided to the practitioner stored in any convenient and suitable ways that preserve their operability and utility. For example, the components can be in dissolved, dehydrated, or lyophilized form. In some embodiments, the components can be provided at room, refrigerated or frozen
10 temperatures. In some embodiments, the components are typically contained in suitable packaging material(s).

[00180] Additional Embodiments

[00181] Reference will now be made to more specific embodiments of the present disclosure and experimental results that provide support for such embodiments. However, Applicants
15 note that the disclosure below is for illustrative purposes only and is not intended to limit the scope of the claimed subject matter in any way.

[00182] Example 1. Force modulated DNA hybridization method

[00183] In this Example, Applicants describe a force-modulated DNA hybridization method that can precisely determine the length of nucleic acids with single-nucleotide resolution and
20 directly visualize the position of nucleic acids during their various biological functions. The method disclosed herein is robust, accurate, versatile, and requires no expensive apparatus.

[00184] The primary scheme contains a magnetically labeled DNA strand and a series of complementary strands of different lengths. The length of the nucleic acid to be analyzed can be accurately determined by force-modulated competitive hybridization. The detection can be
25 achieved by atomic magnetometers, optical transmission, or visually with no apparatus.

[00185] To determine the exact binding site of drugs on nucleic acids, a derived scheme contains a magnetically labeled DNA and its complementary strand, in the presence of the drug to be analyzed and a DNA-cleaving enzyme. The binding site of the drug can be precisely determined by force modulation following surface immobilization.

[00186] Example 1.1. Force modulated competitive hybridization

[00187] **FIG. 1B** can be utilized to illustrate the scheme of a force modulated competitive hybridization system 30. A magnetically labeled DNA strand 32 is designed to be complementary to the analyte strand 34 to be measured, but with several excessive nucleotides. Then a series of DNA rulers 36 are respectively added to the sample containing the two previous strands. Due to the competitive thermodynamic binding, two different scenarios will occur. If the oligonucleotide ruler strand is shorter than or as long as the analyte strand, the analyte strand and magnetically labeled oligonucleotide strand will form duplexes, allowing immobilization of the magnetic particle. Consequently the sample will appear yellow, denoted as “1” for the image 38. If the oligonucleotide ruler strand is longer than the analyte strand, the thermodynamically favored ruler-magnetically labeled oligonucleotide strand hybridization will not be able to immobilize the magnetic particle because of the absence of a functional group on the oligonucleotide ruler strands. Consequently the sample will be clear, denoted as “0” for the image 40. The samples can be distinguished by naked eye and the images can be recorded by a smart phone.

[00188] Example 1.2. Drug-nucleotide binding

[00189] **FIG. 1D** can be utilized to illustrate a system 100 for measuring drug-DNA binding. A magnetically labeled DNA strand 102 is hybridized with its complementary strand 104, in the presence of a drug molecule 106 and a DNA-cleaving enzyme 107. If the drug binds with the DNA specifically at the location of the enzyme, the enzyme will not be able to cleave the duplex. Consequently, the magnetic particles will be able to be immobilized on the surface via the biotin 105 on complementary strand 104, showing yellowish color in image 108(a) and being assigned as “1”. If the drug does not bind specifically at the enzyme’s site on the DNA, the duplex will be cleaved. Thus, the magnetic particles will not be able to bind with the surface because of the absence of a functional group (such as biotin 105 shown in **FIG. 1D**) to interact with the matching functional group on the surface, showing as clear and being assigned as “0” in image 108(b).

[00190] Example 1.3. Measuring nucleic acid length with single-nucleotide resolution

[00191] **FIG. 2A** shows the proof-of-principle results of measuring an analyte strand A13, which contains 13 complementary nucleotides (nts) with the magnetically labeled

oligonucleotide strand. The experiments were performed in the TBS buffer. The oligonucleotide ruler strand length went from 15, 14, 13, to 12 nts complementary to the magnetically labeled oligonucleotide strand, denoted as R15, R14, R13, and R12, respectively. Gravity force was applied on the samples by placing them vertically to remove
5 the nonspecifically bound magnetic particles. The buoyant mass of the magnetic particles has been measured to be 4.6×10^{-15} kg, resulting a gravity of 45 fN. Both R15 and R14 showed no immobilized particles, thus “0” was assigned; R13 and R12 showed particles on the surface, indicated by “1” on the images.

[00192] The absence or existence of immobilized magnetic particles were confirmed using an
10 atomic magnetometer (**FIG. 2B**). High magnetic signal of 100 pT was obtained for the “1” images using R13 and R12, whereas very low signal was obtained for the “0” images using R15 and R14. The “0” to “1” transition occurred at R13, indicating the analyte strand to be 13-nt length. When the analyte strand was changed to be 12-nt complementary to the magnetically labeled oligonucleotide strand (denoted as A12), such transition occurred at
15 R12 (**FIG. 2C**). Therefore, the results indicated that DNA rulers can distinguish DNAs with single-nt length difference.

[00193] The basis of the DNA ruler system is the thermodynamic equilibrium between two competitive hybridizations, which is difficult to predict and varies with different biological conditions. This problem is solved by applying an adjustable mechanical force on the sample
20 after surface immobilization.

[00194] In **FIG. 2D**, the same ruler systems were tested in the TAM₁₀ buffer. When only gravity was applied, A13 and A12 showed the same pattern, in which both transitions took place at R14. The same result was obtained even when the samples were heated to 75 °C. This means without appropriate force modulation or simply using heating, single-nt
25 resolution cannot be achieved. However, when a 65 pN centrifugal force was applied, A13 and A12 were distinguished, with transitions at oligonucleotide ruler strands of their respective lengths (**FIG. 2E**). The force value is just above the approximate dissociation force of single 13-bp DNA duplexes. The requirement of force indicates that R14 replacing A13 in TAM₁₀ is not as efficient as in TBS.

[00195] The method can also determine the length of RNAs. **FIG. 3A** compares measurements of A13 and its RNA analog, A13r, using DNA rulers R15-R12. The “0” to “1”

transition occurred at R12 for the RNA A13r, consistent with the fact that DNA-DNA duplex is slightly more stable than the corresponding RNA-DNA duplex. In other words, the small difference between the two duplexes was successfully revealed.

[00196] A slight adjustment of the scheme can achieve label-free detection. Here, the biotin label was placed on the oligonucleotide ruler strands, leaving the analyte strand label-free (FIG. 3B). The transition is therefore reversed to “1” to “0”, compared to the original scheme in FIG. 1B. Images shown were using R12 (top) and R13 (bottom) to probe A13.

[00197] The sequences used above and throughout this disclosure are listed in Table 1.

Name	Sequence (from 5' to 3')
A13	Bio/CAG ACT GAC TCG A ^a
A12	Bio/CAG ACT GAC TCG
A13r	Bio/rCrArG rArCrT rGrArC rTrCrG rA ^b
Magnetically labeled oligonucleotide strand	A ₂₅ /GAT CGA GTC AGT CTG ^c
R15	CAG ACT GAC TCG ATC
R14	CAG ACT GAC TCG AT
R13	CAG ACT GAC TCG A
R12	CAG ACT GAC TCG
R11	CAG ACT GAC TC
mRNA (Pre)	Bio/C AAC UGU UAA UUA AAU UAA AUU AAA AAG GAA AUA AAA AUG UUU GAA AGU AAG UAC <u>GUA</u> <u>AAU</u> <u>CUA</u> <u>CUG</u> <u>CUG</u> AAC UC ^d
mRNA (Post)	Bio/C AAC UGU UAA UUA AAU UAA AUU AAA AAG GAA AUA AAA AUG UUU GAA AGU AAG UAC <u>GUA</u> <u>AAU</u> <u>CUA</u> <u>CUG</u> <u>CUG</u> AAC UC ^d
R15'	A ₂₅ /CT CAA GAG <u>CAG</u> <u>TAG</u> <u>ATT</u> <u>TAC</u> G ^e

10 **Table 1.** Sequences of the DNAs and RNAs used in this disclosure. ^aBio: biotin labeled; ^bRNA nucleotides; ^cFunctionalized with 25 units of A. ^dBold indicating the ribosome-uncovered nucleotides; underscored indicating the segment for probing. ^eUnderscored indicating the complementary nucleotides to the mRNA.

15 [00198] Example 1.4. Enzyme-DNA interaction and ribosome translocation

[00199] FIG. 4 illustrates two examples of Enzyme-DNA interaction and ribosome translocation. The first was to determine the cleavage site of DpnII, an endonuclease. DpnII has been known to cleave DNA duplexes specifically at the 5'-...GATC...-3' segment. From the sequence in FIG. 4A, a 11-nt fragment with biotin label will be produced after the

enzymatic cleavage. This fragment was measured by DNA rulers from R15 to R11. The transition of the presence of particles occurred at R11 (**FIG. 4B**). Therefore, the exact cleavage position of DpnII was confirmed by Applicants' method.

[00200] The ribosome movement on the mRNA during translocation was also measured. For the pre-translocation state (Pre), the ribosome (indicated by the two orange ovals) occupies the position on the mRNA with 15-nt exposed to bind with the magnetically labeled oligonucleotide strand. If translocation took place, the ribosome would move three nts to reach the post-translocation (Post) position. Consequently, the mRNA would have only 12-nt to hybridize with the magnetically labeled oligonucleotide strand (**FIG. 4C**). The 3-nt difference can be revealed by oligonucleotide ruler strand R15' modulated at 80 pN, because R15' was only able to completely replace the 12-bp duplex between Post and the magnetically labeled oligonucleotide strand, leaving no particles on the sample surface (**FIG. 4D**). The force value just exceeds the expected dissociation force for the 15-bp DNA/RNA duplex.

[00201] Example 1.5. Drug-nucleotide binding

[00202] The derived scheme presented in **FIG. 1D** was used to precisely determine the binding sites of drug molecules. **FIG. 5A** shows the sequence of a DNA duplex, which contains two enzyme-binding segments: 5'-GATC for DpnII and 5'-GAATTC for EcoRI. **FIG. 5B** shows the results. When no drugs were in the sample, either enzyme will cleave the duplex, leading no magnetic particles being immobilized (left column). When daunomycin was present with DpnII, particle immobilization was observed, indicating that daunomycin blocked the binding site of DpnII, leaving the duplex intact, so the magnetic particles can be immobilized on the surface.

[00203] When netropsin and DpnII were present, the sample was clear with no immobilized particles. This is because netropsin preferably binds on the 5'-AATT segment, which does not interfere with the binding of DpnII (confirmed in **FIG. 5B**). However, the contrast between netropsin and daunomycin could only be revealed after applying a weak force of ~1 pN, showing the crucial role of force modulation.

[00204] The preferred binding of netropsin was confirmed when a different endonuclease, EcoRI, was used. EcoRI specifically binds at 5'-GAATTC segment. The results (second row of **FIG. 5B**) show that netropsin was able to block EcoRI whereas daunomycin could not.

Again, a ~1 pN force modulation was required to distinguish the different behavior between the two drugs. The results showed that using Applicants' force-modulation method and an enzyme, the specific binding site of drug molecules can be determined visually.

[00205] The dissociation force can be further used to quantitatively distinguish the binding strength of different drug molecules at each and every specific nucleotide sequence. **FIG. 5C** plots the dissociation forces of four different drug molecules interacting with three different sequences. The red section shows the drugs (daunomycin, netropsin, doxorubicin, and epirubicin) binding with the GATC segment of the DNA shown in **FIG. 5A**. This segment is specifically chosen because of the addition of enzyme DpnII.

[00206] The results clearly indicate the order of binding strength for this particular sequence is daunomycin, doxorubicin > epirubicin >> daunomycin >> netropsin. The blue section shows the four drugs binding with the GAATTC segment, specified by the presence of EcoRI. Interestingly, netropsin becomes the second strongest. Another interesting observation is that epirubicin is more sequence-selective than doxorubicin, although the two molecules only differ by a chiral center, with the same chemical formula. This result is consistent with the fact that epirubicin appears to be less toxic in chemotherapy. This example shows the quantitative measurements of drug-DNA interaction for any given sequence by choosing the appropriate enzyme, which cannot be resolved by any existing techniques.

[00207] Example 1.6. Multiplexed detection and quantification with a device

[00208] Multiplexed detection is usually required for high-throughput biosensing and analysis. In addition, quantification of immobilized particles is needed to extract key biophysical parameters. **FIG. 6** shows both tasks can be achieved with the currently disclosed method. In **FIG. 6A**, the experiment disclosed in **FIG. 2A** was repeated three times, with all samples on a single plate. The results are consistent with the previous results, in that the transition of signal occurred at R13. This result showed both the feasibility of multiplexed detection and the reliability of Applicants' method.

[00209] **FIG. 6B** shows a photo of a novel and inexpensive device that measures the transmission of a laser beam after passing through each sample. The detection was achieved by using a photodetector (DET100A, Thorlabs) coupled with a voltmeter. **FIG. 6C** shows the results of samples containing different concentrations of daunomycin. Fitting of the

profile yielded a binding constant of $9 \pm 2 \times 10^4 \text{ M}^{-1}$, consistent with the results using a sensitive atomic magnetometer.

[00210] Example 1.7. A unique design for sample holders

[00211] Applicants' method of force-modulated hybridization requires three unique steps. In some embodiments, the first step is to hybridize the involved DNA/RNA strands at the specific biological condition for each application. In some embodiments, the second step is to perform surface immobilization. In some embodiments, the third step is to apply a mechanical force on the sample. This sample manipulation leads to a specific design of sample holders.

[00212] As shown in **FIG. 7A**, a new system was designed consisting of a puncher, an incubator, and a displayer. After incubating the sample, the puncher will be used to break the bottom of the incubator so the sample will be transferred to the corresponding wells on the displayer. The bottom of the displayer was functionalized, for example with streptavidin, so that magnetic particles will be able to be immobilized through a labeled DNA duplex, for example with biotin. A mechanical force is then applied on the displayer followed by final inspection and measurement.

[00213] As shown in **FIG. 7B**, mechanical force can be achieved for multiplexed sample wells. The samples within each lane were under identical conditions. 1: DNA duplex with 100 μM daunorubicin and DpnII; 2: DNA duplex with 100 μM netropsin and DpnII; 3: DNA duplex; 4: DNA duplex incubated with 100 μM daunomycin and EcoRI; 5: DNA duplex with 100 μM netropsin and EcoRI. After applying ultrasound through a piezo plate, only the particles in lanes 2 and 5 were removed, indicating weak binding between netropsin and CTAG (lane 2), and between daunomycin and CTTAAG (lane 4). These results are consistent with the results in **FIG. 5**.

[00214] Example 1.8. Sequencing using the DNA rulers

[00215] The precise DNA rulers can be used for sequencing, which will be particularly suited for sequencing short DNA or RNA strands. These types of strands are difficult to directly sequence because of their limited length. **FIG. 8** shows the use of a series of oligonucleotide ruler strands to sequence the unknown nucleotides on a target DNA. The target has a segment of known sequence followed by unknown ones in red (T, then A and G, in this

example). The four ruler DNAs, which differ by only the last nucleotide, were respectively mixed with the target and a DNA with biotin label that is complementary to the known segment of the target, at a ratio of 2:1:1. Only when the oligonucleotide ruler strand contains the complementary nucleotide with the unknown one on the target (A in this example), the sample well showed no particles.

[00216] Two detection methods were used. One was photos taken by a smart phone. A clear photo indicates no particles, whereas a yellow photo indicates the presence of particles. The other technique was the device shown in **FIG. 6B**, a high voltage indicates (2.2-2.4 V) no particles, and a low voltage indicates particles (1.4-1.5 V). Using the same principle, the next unknown nucleotides (A and G in this example) can be identified step by step. In each step, a set of oligonucleotide ruler strand will be used.

[00217] The various methods and techniques described above provide a number of ways to carry out the invention. It is to be understood that not necessarily all objectives or advantages described may be achieved in accordance with any particular embodiment described herein. Thus, for example, those skilled in the art will recognize that the methods can be performed in a manner that achieves or optimizes one advantage or group of advantages as taught herein without necessarily achieving other objectives or advantages as may be taught or suggested herein. A variety of advantageous and disadvantageous alternatives are mentioned herein. It is to be understood that some preferred embodiments specifically include one, another, or several advantageous features, while others specifically exclude one, another, or several disadvantageous features, while still others specifically mitigate a present disadvantageous feature by inclusion of one, another, or several advantageous features.

[00218] Without further elaboration, it is believed that one skilled in the art can, using the description herein, utilize the present disclosure to its fullest extent. The embodiments described herein are to be construed as illustrative and not as constraining the remainder of the disclosure in any way whatsoever. While the embodiments have been shown and described, many variations and modifications thereof can be made by one skilled in the art without departing from the spirit and teachings of the invention. Accordingly, the scope of protection is not limited by the description set out above, but is only limited by the claims, including all equivalents of the subject matter of the claims. The disclosures of all patents, patent applications and publications cited herein are hereby incorporated herein by reference,

to the extent that they provide procedural or other details consistent with and supplementary to those set forth herein.

CLAIMS

What is claimed is:

1. A method of determining the length of an analyte strand, comprising:

(a) incubating a magnetically labeled oligonucleotide strand, the analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture,

wherein the magnetically labeled oligonucleotide strand is complementary in sequence to the analyte strand,

wherein the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand and comprise different lengths, and

wherein either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label;

(b) transferring the mixture to a surface functionalized to couple with the at least one label;

(c) applying a mechanical force to the mixture; and

(d) inspecting the surface for immobilized particles,

wherein if the analyte strand is labeled, then the longest oligonucleotide ruler strand producing immobilized particles on the surface represents the length of the analyte strand, and

wherein if the oligonucleotide ruler strands are labeled, then the longest oligonucleotide ruler strand not producing immobilized particles represents the length of the analyte strand; and

(e) wherein a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the length of the analyte strand is determined.

2. The method of claim 1,

wherein the mechanical force modulates oligonucleotide hybridization;

wherein the oligonucleotide hybridization results in the presence of immobilized particles when the analyte strand is labeled and the oligonucleotide ruler strand is not longer than the analyte strand, and wherein the immobilized particles represent a hybrid structure between the magnetically labeled oligonucleotide strand and the analyte strand;

wherein the oligonucleotide hybridization results in the absence of immobilized particles when the analyte strand is labeled and the oligonucleotide ruler strand is longer than the analyte strand;

wherein the oligonucleotide hybridization results in the presence of immobilized particles when the oligonucleotide ruler strands are labeled and the oligonucleotide ruler strand is longer than the analyte strand, and wherein the immobilized particles represent a hybrid structure between the magnetically labeled oligonucleotide strand and the oligonucleotide ruler strand; and

wherein the oligonucleotide hybridization results in the absence of immobilized particles when the oligonucleotide ruler strands are labeled and the oligonucleotide ruler strand is shorter than or the same length as the analyte strand.

3. The method of claim 1, wherein the analyte strand is labeled with the at least one label.

4. The method of claim 1, wherein the oligonucleotide ruler strands are labeled with the at least one label.

5. The method of claim 1, wherein the immobilized particles are represented by a yellow color on the surface, and wherein the yellow color is derived from the magnetically labeled oligonucleotide strand.

6. A method of determining the length of an analyte strand, comprising:

incubating the analyte strand with a magnetically labeled oligonucleotide strand and an oligonucleotide ruler strand to form a mixture,

wherein the analyte strand is labeled with at least one label;

transferring the mixture to a surface,

wherein the surface is functionalized to couple with the at least one label;

applying a mechanical force to the mixture; and

inspecting the color of the surface,

wherein if the surface is yellow, then the analyte strand is longer than or same length as the oligonucleotide ruler strand, and

wherein if the surface is colorless or not yellow, then the analyte strand is shorter than the oligonucleotide ruler strand.

7. The method of claim 1 or 6,

wherein the analyte strand is a DNA strand or an RNA strand;

wherein the magnetically labeled oligonucleotide strand is a DNA strand or an RNA strand;

wherein the magnetically labeled oligonucleotide strand has a longer length than the analyte strand;

wherein the magnetically labeled oligonucleotide strand is 12-50 nucleotides long;

and

wherein the magnetically labeled oligonucleotide strand is labeled with a magnetic particle.

8. The method of claim 1 or 6,

wherein the oligonucleotide ruler strands are selected from the group consisting of DNA strands, RNA strands, and combinations thereof;

wherein the oligonucleotide ruler strands are complementary in sequence to the analyte strand; and

wherein the oligonucleotide ruler strands comprise oligonucleotides longer than the analyte strand, oligonucleotides shorter than the analyte strand, and oligonucleotides the same length as the analyte strand.

9. The method of claim 1 or 6,

wherein the surface is functionalized with a molecule that binds to the at least one label; and

wherein the at least one label is selected from the group consisting of biotin, streptavidin, digoxigenin, avidin, maleic imide, gold, proteins, nucleic acids, functional groups, and combinations thereof.

10. The method of claim 1 or 6,

wherein the mechanical force is selected from the group consisting of gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof; and

wherein the mechanical force applied is between 10 fN and 500 pN.

11. The method of claim 1 or 6, wherein the at least one label is biotin, and wherein the surface is functionalized with streptavidin.

12. The method of claim 1 or 6, wherein the inspecting occurs by visual inspection.

13. The method of claim 6, wherein the yellow color is derived from the magnetically labeled oligonucleotide strand.

14. A method of determining an unknown nucleic acid sequence of a magnetically labeled oligonucleotide strand, comprising:

(a) incubating the magnetically labeled oligonucleotide strand, an analyte strand, and one of a series of oligonucleotide ruler strands to form a mixture,

wherein the analyte strand is complementary in sequence to at least some of the known nucleic acid sequences of the magnetically labeled oligonucleotide strand,

wherein the series of oligonucleotide ruler strands are complementary in sequence to the magnetically labeled oligonucleotide strand,

wherein the oligonucleotide ruler strands comprise nucleic acids at their ends that span a length of the unknown nucleic acid sequence of the magnetically labeled oligonucleotide strand, and

wherein either the analyte strand or the oligonucleotide ruler strands are labeled with at least one label;

(b) transferring the mixture to a surface functionalized to couple with the at least one label;

(c) applying a mechanical force to the mixture;

(d) inspecting the surface for immobilized particles,

wherein if the analyte strand is labeled, then the oligonucleotide ruler strand that does not produce immobilized particles on the surface contains at least some of the unknown nucleic acid sequences at its end, and

wherein if the oligonucleotide ruler strand is labeled, then the oligonucleotide ruler strand that produces immobilized particles on the surface contains at least some of the unknown nucleic acid sequences at its end; and

(e) wherein a different oligonucleotide ruler strand is used to repeat steps (a)-(d) until the unknown sequence of the magnetically labeled oligonucleotide strand is determined.

15. The method of claim 14,

wherein the magnetically labeled oligonucleotide strand has a segment of the known nucleic acid sequences followed by a segment of the unknown nucleic acid sequence;

wherein the unknown nucleic acid sequence is at the end of the magnetically labeled oligonucleotide strand;

wherein the magnetically labeled oligonucleotide strand has a longer length than the analyte strand;

wherein the magnetically labeled oligonucleotide strand is a DNA strand or an RNA strand;

wherein the magnetically labeled oligonucleotide strand is 12-50 nucleotides long;

wherein the unknown nucleic acid sequence is 1-20 nucleotides long; and

wherein the magnetically labeled oligonucleotide strand is labeled with a magnetic particle.

16. The method of claim 14,

wherein the oligonucleotide ruler strands comprise oligonucleotide ruler strands that differ by only the last nucleotide;

wherein the oligonucleotide ruler strands are selected from the group consisting of DNA strands, RNA strands, and combinations thereof;

wherein the oligonucleotide ruler strands are complementary in sequence to the analyte strand; and

wherein oligonucleotide ruler strands comprise oligonucleotides longer than the analyte strand.

17. The method of claim 14, wherein the surface is functionalized with a molecule that binds to the at least one label, and wherein the at least one label is selected from the group consisting of biotin, streptavidin, digoxigenin, avidin, maleic imide, gold, proteins, nucleic acids, functional groups, and combinations thereof.

18. The method of claim 14,

wherein the mechanical force is selected from the group consisting of gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof; and

wherein the mechanical force applied is between 10 fN and 500 pN.

19. The method of claim 14, wherein the analyte strand is a DNA strand or an RNA strand, and wherein the analyte strand is labeled with the at least one label.

20. The method of claim 14, wherein the oligonucleotide ruler strands are labeled with the at least one label.

21. The method of claim 14, wherein the at least one label is biotin, and wherein the surface is functionalized with streptavidin.

22. The method of claim 14,

wherein the inspecting is selected from the group consisting of visual inspection, measurement of light transmission, measurement by a magnetic sensor, or combinations thereof; and

wherein the immobilized particles are represented by a yellow color on the surface, and wherein the yellow color is derived from the magnetically labeled oligonucleotide strand.

23. A method for determining the binding of a nucleotide to an active agent, comprising:

(a) incubating a first oligonucleotide strand and a second oligonucleotide strand for hybridization in the presence of an oligonucleotide cleaving enzyme and an active agent to form a mixture,

wherein the first and second oligonucleotide strands are complementary to one another,

wherein at least one of the first or second oligonucleotide strands is labeled with at least one label, and

wherein at least one of the first or second oligonucleotide strands is labeled with at least one magnetic particle to provide a magnetically labeled oligonucleotide strand;

(b) transferring the mixture to a surface functionalized to couple with the at least one label;

(c) applying a mechanical force to the mixture; and

(d) inspecting the surface for immobilized particles,

wherein the presence of immobilized particles indicates that the active agent binds to the at least one of the first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof, and

wherein the absence of immobilized particles indicates that the active agent does not bind to the at least one of the first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof.

24. The method of claim 23, wherein a first or second oligonucleotide strand is labeled with the at least one label while the complementary oligonucleotide strand is labeled with the at least one magnetic particle.

25. The method of claim 23, wherein the same oligonucleotide strand is labeled with the at least one label and the at least one magnetic particle.

26. The method of claim 23, wherein the immobilized particles are represented by a yellow color on the surface, and wherein the yellow color is derived from the magnetically labeled oligonucleotide strand.

27. A method for determining the binding of a nucleotide to an active agent, comprising:

incubating a magnetically labeled first oligonucleotide strand with a second oligonucleotide strand for hybridization in the presence of an active agent and an oligonucleotide cleaving enzyme to form a mixture,

wherein the second oligonucleotide strand is labeled with at least one label, and

wherein the second oligonucleotide strand is complementary to the magnetically labeled first oligonucleotide strand; and

transferring the mixture to a surface,

wherein the surface is functionalized to couple with the at least one label;

applying a mechanical force to the mixture;

inspecting the color of the surface,

wherein if the surface is yellow, then determining that the active agent binds to at least one of the magnetically labeled first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof, and

wherein if the surface is colorless or not yellow, then determining that the active agent does not bind to the at least one of the magnetically labeled first oligonucleotide strand, the second oligonucleotide strand, or the hybridized version thereof.

28. The method of claim 23 or 27,

wherein each of the first and second oligonucleotide strand is selected from the group consisting of DNA strands, RNA strands, and combinations thereof;

wherein the active agent binding site represents the enzyme binding site;

wherein the oligonucleotide cleaving enzyme comprises a DNA or RNA nuclease;

wherein the surface is functionalized with a molecule that binds to the at least one label;

wherein the inspecting is selected from the group consisting of visual inspection, measurement of light transmission, measurement by a magnetic sensor, or combinations thereof; and

wherein the mechanical force is selected from the group consisting of gravitational force, centrifugal force, shaking force, ultrasound radiation force, magnetic force, and combinations thereof.

29. The method of claim 23 or 27, wherein the active agent is an experimental drug molecule.

30. The method of claim 27, wherein the yellow color is derived from the magnetically labeled oligonucleotide strand.

31. A sample holder device, comprising:

an incubator,

wherein the incubator comprises a plurality of first wells for incubating a plurality of samples;

a displayer for displaying the plurality of the samples,

wherein the displayer comprises a plurality of second wells, and

wherein a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands; and

a transferring apparatus,

wherein the transferring apparatus is capable of transferring the plurality of the samples from the plurality of first wells to the plurality of second wells.

32. The sample holder device of claim 31, wherein the plurality of second wells are superimposable on the plurality of first wells.

33. The sample holder device of claim 31, wherein the functional group comprises streptavidin.

34. The sample holder device of claim 31,

wherein the transferring apparatus comprises a perforator, wherein the perforator is capable of breaking or perforating a surface of the plurality of first wells so that the plurality of the samples from the plurality of first wells are transferred to the plurality of second wells; and

wherein the perforator comprises a matrix of sharp tips at positions corresponding to the plurality of first wells.

35. The sample holder device of claim 31,

wherein the transferring apparatus comprises a sample transport system;

wherein the sample transport system is capable of transporting the plurality of the samples from the plurality of first wells to the plurality of second wells; and

wherein the sample transport system comprises a multi-head pipette system, a robotic arm, or combinations thereof.

36. A method of loading and transferring a plurality of samples, said method comprising:

loading the plurality of samples into a plurality of first wells of an incubator;

transferring the plurality of the samples from the plurality of first wells of the incubator to a plurality of second wells of a displayer,

wherein a surface of each of the second wells is functionalized with a functional group that is capable of immobilizing oligonucleotide strands, and

wherein the transferring occurs by utilizing a transferring apparatus; and

inspecting or analyzing the plurality of second wells of the displayer.

37. The method of claim 36, wherein the transferring apparatus comprises a perforator, and wherein the transferring comprises utilizing the perforator to break or perforate the surface of the plurality of first wells so that the plurality of the samples from the plurality of first wells are transferred to the plurality of second wells.

38. The method of claim 36, wherein the transferring apparatus comprises a sample transport system, and wherein the transfer comprises transporting the plurality of the samples from the plurality of first wells to the plurality of second wells.

39. The method of claim 36, further comprising a step of applying mechanical force to the plurality of second wells prior to inspecting or analyzing the second wells.

40. A device, comprising:

- a sample holder;
- a light source;
- a voltmeter;
- a photodetector; and
- a recording device holder.

41. The device of claim 41,

wherein the sample holder comprises a multiplexed sample holder for holding a plurality of samples positioned in a plurality of wells;

wherein the light source comprises a laser pointer or a laser diode;
wherein the light source is positioned above sample holder;
wherein the photodetector comprises a photodiode beneath the sample holder that is connected to the voltmeter; and
wherein the recording device holder is capable of holding a recording device, wherein the recording device is a smartphone.

42. The device of claim 41, further comprising:

a translation stage for changing the location of the sample on the sample holder,
wherein the translation stage is a two-dimensional manual stage or automated stage; and
a mechanical frame for providing structural support to the device.

43. A device, comprising:

a light source;
a photodetector;
a translation stage;
a mechanical frame for providing structural support to the device; and
a voltmeter.

44. The device of claim 43,

wherein the light source comprises a laser pointer or a laser diode;
wherein the photodetector comprises a photodiode connected to the voltmeter; and
wherein the translation stage is a two-dimensional manual stage or automated stage.

1/17




FIG. 1A




FIG. 1B

3/17




FIG. 1C




FIG. 1D




FIG. 1E




FIG. 1F




FIG. 1G




FIG. 1H




FIG. 1I

10/17




FIG. 2A




FIG. 2B



FIG. 2C

FIG. 2

d)

	<i>R15</i>	<i>R14</i>	<i>R13</i>	<i>R12</i>
<i>A13</i>	0	1	1	1
<i>A12</i>	0	1	1	1

FIG. 2D

e)

<i>A13</i>	0	1	1
<i>A12</i>	0	0	1

FIG. 2E

FIG. 2




FIG. 3A




FIG. 3B

FIG. 3

13/17

a)



FIG. 4A

b)




FIG. 4B

c)




FIG. 4C

d)



FIG. 4D



FIG. 4

14/17



FIG. 5A

b)

	No drug	Daunomycin	Netropsin		
	 0	 1	 0	F	 1
	 0	 0	 1	F	 1




FIG. 5B




FIG. 5C

FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C

FIG. 6




FIG. 7A




FIG. 7B

FIG. 7




FIG. 8