



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C07C X, C07D X, A61K 31/00</p>	<p>A2</p>	<p>(11) International Publication Number: WO 87/ 07599 (43) International Publication Date: 17 December 1987 (17.12.87)</p>
<p>(21) International Application Number: PCT/US87/01371 (22) International Filing Date: 10 June 1987 (10.06.87) (31) Priority Application Number: 872,812 (32) Priority Date: 11 June 1986 (11.06.86) (33) Priority Country: US</p> <p>(71) Applicant: ALDER RESEARCH CENTER CORPORATION [US/US]; 165 New Boston Street, Woburn, MA 01801 (US). (72) Inventor: DRIEDGER, Paul, E. ; 30 Symmes Street, Winchester, MA 01890 (US). (74) Agent: DeCONTI, Giulio, A., Jr.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US).</p>	<p>(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: ANTI-INFLAMMATORY COMPOSITIONS</p> <p>(57) Abstract</p> <p>Compounds having anti-inflammatory and other activities. The compounds are derived from diterpenes, indol alkaloids, polyacetates, diaminobenzyl alcohols, diacylglycerol and bryostatins.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

ANTI-INFLAMMATORY COMPOSITIONSRelated Applications

This application is a continuation-in-part of Application Serial No. 872,812, filed June 11, 1986.

Background

Protein kinase C is an enzyme found in nearly all animal tissues and animal cells that have been examined. Its identity is generally established by its ability to phosphorylate proteins when adenosine triphosphate, calcium ions and phospholipid cofactors are present, with greatly reduced activity when these cofactors are absent. Protein kinase C activity is substantially stimulated by certain 1,2-sn-diacylglycerols that bind specifically and stoichiometrically to a recognition site on the enzyme. Stimulation of protein kinase C by these diacylglycerols has been shown to be an important physiological event that mediates the actions of a wide variety of hormones, neurotransmitters, and other biological control factors such as histamine, vasopressin, alpha-adrenergic agonists, dopamine agonists, muscarinic cholinergic agonists, platelet activating factor, etc. [see Y. Nishizuka, Nature 308 693-698 (1984) and Science 225 1365-1370 (1984) for reviews].

The biological role of protein kinase C is also of great interest because of the discovery that certain very powerful tumor promoting substances activate this enzyme by binding specifically and with very high affinity to the diacylglycerol binding site on the enzyme. In addition to diacylglycerols, there are at present five other known classes of compounds that bind to this site,

-2-

including diterpenes such as the phorbol esters, indole alkaloids (indolactams) such as the teleocidins, lyngbyatoxin, and indolactam V, polyacetates such as the aplysiatoxins and oscillatoxins, certain derivatives of diaminobenzyl alcohol, and the macrocyclic lactones of the bryostatin class. The phorbol esters have long been known as powerful tumor promoters, the teleocidins and aplysiatoxins are now known to have this activity, and it appears likely that additional classes of compounds will be found to have the toxic and tumor promoting activities associated with the capability to bind to the diacylglycerol site of protein kinase C and thus activate the enzyme. Other toxicities of these agents when administered to animals include lung injury and profound changes in blood elements, such as leukopenia and neuropenia.

In addition to potent tumor promoting activity, these six classes of compounds, collectively referred to as the "phorboids", display a vast range of biological activities, as would be expected from the widespread distribution of their target enzyme. Some of these activities, like tumor promotion, indicate the involvement of protein kinase C in important normal or pathological processes in animals. Thus, the phorboids are potent skin inflammatory agents, cause smooth muscle contraction in several tissues, alter immune system function and can be used to cause a variety of other normal or pathological responses. Related disease states such as

the development of cancer, the onset and/or maintenance of inflammatory disease, the role of vasoconstriction in hypertension, the role of bronchoconstriction in asthma, the role of cholinergic, adrenergic, and dopaminergic synapses in diseases of the central/peripheral nervous systems, may be mediated in vivo by the stimulation of protein kinase C by diacylglycerols, the latter being generated in the cell by pathological agents or conditions.

In analyzing the activity of a pharmaceutical or other bioactive compound, it is useful to consider two properties: the efficacy, defined as the capability to elicit a full or partial biological result, such as complete displacement of a ligand from its receptor site or the complete inhibition of inflammation or edema caused by a standard stimulus; and the potency, defined as that amount or concentration of drug that causes 50% of the full response (often abbreviated as the ED₅₀). It is frequently the case within a given class of pharmaceutical agents that individual members of the class all have equal efficacy, i.e. they each can generate a full biological effect, but they show differencing potencies. Thus, the structural modifications within such a class affect only the amount necessary to achieve a given result, and the modified compounds otherwise have generally the same central biological characteristic. There may also be differences between members of such a class as regards proper-

ties other than the central biological characteristic; for example, members of the class might differ in side effects or toxicity.

Well-known pharmaceuticals that have been in extensive use for years or decades show a wide range of optimal therapeutic potencies. Aspirin, for example, is often taken in multi-gram amounts per day for treatment of inflammation or arthritis, and detailed analyses of its mechanism of action in vitro show that a concentration in the millimolar range is required. In contrast, steroid-based topical anti-inflammatory compounds such as fluocinolone acetonide are many thousand-fold more potent, and, beyond this, some oral contraceptive agents are prescribed in daily doses in the microgram range. Thus, although high potency is generally advantageous for a pharmaceutical, it is not an absolute requirement.

Several thousand of the high skin-inflammatory and tumor-promoting phorboids have been reported in the literature, including numerous examples on which minor chemical modifications have been made [see Evans and Soper, Lloydia 41 193-233 (1978) and references cited therein]. The structures of these phorboids can be compared, and their activities for inflammation and tumor promotion can be analyzed from the perspective of efficacy and potency. The structures of the different classes of phorboids vary quite markedly from one to the other class (see Summary of the Invention for structural comparisons), yet widespread testing of their biological activities has shown that these classes have generally very similar biological properties. In particular, the

thousands of known phorboids of the highly potent diterpene, indolactam, and polyacetate classes appear to have, with very minor exceptions, virtually identical efficacies as skin irritants and tumor promoters [T. Sugimura, Gann 73 499-507 (1982)]. The exceptions involve a few compounds that have a short duration of irritant activity and/or manifest diminished tumor promoting activity, perhaps due to toxicity or secondary parameters such as differing metabolic destruction rates.

In contrast to the essentially equal efficacies among the vast majority of phorboids, their relative potencies cover a wide range, as measured in inflammation and promotion tests and as measured in numerous other in vivo and in vitro systems. Example compounds can be found in the diterpene, indolactam, and polyacetate classes that have nearly equal, very high potencies. At the same time there are compounds in each of these classes which embody significant structural changes that do not diminish efficacy but do result in potency decreases of 10-fold to 100,000-fold or more [see, for example, Driedger and Blumberg, Cancer Res. 37 3257-3265 (1977), Cancer Res. 39 714-719 (1979)]. Thus, all these compounds appear to be capable of achieving generally the same biological results, and merely differ in the amount which must be used to obtain a given result.

In vitro measurements of biochemical properties provide an even more sensitive method for comparing the properties of the various phorboids. For

example, using a radioactively labeled phorboid such as [³H]phorbol 12,13-dibutyrate or [³H]lyngbyatoxin, one can measure the potency of a test compound as a competitive ligand for the diacylglycerol binding site on protein kinase C. Alternatively, one can measure the ability of a given phorboid to stimulate the protein kinase C-mediated incorporation of radioactive phosphate from [³²P]adenosine triphosphate into a standard acceptor substrate such as histone H1. Tests of this nature reveal a difference in potency between given phorboid agonists of as much as 10,000,000-fold or more [Dunn and Blumberg, Cancer Res. 43 4632-4637 (1983), Table 1].

These basic data regarding the phorboid agonists are an important consideration because they underscore the concept that the structural differences among these previously known phorboids, especially the diterpenes, indolactams, polyacetates, and bryostatins, generally do not affect their efficacies as toxic agonists, and indeed a wide variety of structural changes are tolerated in this regard. Such changes generally alter potency only and do not provide agents with therapeutic utility, since they retain their toxicity.

Some minor changes in phorboid structure are known to result in inactive compounds, such as a stereochemical change from 4-beta to 4-alpha in the phorbol series, and indeed some of the diterpene

skeleton structures carry hydroxy groups that must be esterified in order for inflammatory activity to be observed. However, these inactive compounds are quite few in number among the known phorboids, and no therapeutic utility has been demonstrated for them.

The phorbol esters, indolactams, polyacetates, diaminobenzyl alcohols, and bryostatins are generally found in plants, molds, and algae, or are synthetic in origin. Although they are found in many parts of the world, normal human contact with them is thought to be low. In contrast, the diacylglycerols are part of the functioning of virtually every type of animal cell except erythrocytes of some species, and thus the undesirable activation of protein kinase C by the diacylglycerols may have a very widespread role in human diseases.

Thus, new compounds, capable of blocking the activation of protein kinase C by acting as specific pharmacological antagonists of the diacylglycerols, would be valuable agents in the prevention and treatment of a wide variety of diseases in animals and humans.

There may be several different forms of protein kinase C each having different biological roles. The stimulation of one form can lead to undesirable results such as inflammation or the development of cancer, while stimulation of another form of the enzyme might produce beneficial effects, such as the abrogation of inflammation or the secretion of useful bioregulatory factors such as hormones and interferons.

-8-

Moreover, the exact correspondence between diacylglycerol binding sites and protein kinase C has not been fully explored, and there may be several different such binding sites with differential affinities for diacylglycerols and other phorboids. Indeed there is published evidence for several distinct classes of phorboid binding sites in various tissues and cell types (Dunn and Blumberg, *op. cit.*). However, in this study, even the ligands showing the clearest differences in affinity for these distinct classes are only selective by a factor of 10-100 dissociation constant. Thus, other new compounds, capable of selectively activating one useful, but not another, deleterious, diacylglycerol target site, would also be valuable agents for the prevention or treatment of inflammation.

Earlier efforts to use the previously known phorboids themselves or to modify of the structures of the known phorboids, have not been successful in producing useful compounds.

It has been known for some time that several of the toxic, inflammatory and tumor-promoting compounds such as phorbol 12-tiglate 13-decanoate, mezerein, lyngbyatoxin and aplysiatoxin have anti-leukemic activity in mouse model tests. However, these compounds are all extremely toxic and are cancer suspect agents, thus eliminating them from consideration as human therapeutic agents.

Ganong et al. [Proc. Nat. Acad. Sci. 83 1184-1188 (1986)] tested a series of diacylglycerols and found no antagonistic activity in that series against the standard agonist 1,2-dioctanoylglycerol. It is of particular note that several compounds tested in this work were modified in the hydroxymethyl portion of the diacylglycerol molecule, and these modifications produced only a loss of activity or a weakened activity that was not distinguishable from the agonist activity of 1,2-dioctanoylglycerol itself. These hydroxymethyl-modified compounds were not antagonists in these tests and no utility was found. Similarly, Thielmann and Hecker [Forsch. Krebsforsch. Vol. VII, pp. 171-179 (1969), New York: Schattauer] found only a complete loss of biological activity in their study when the hydroxy group of the hydroxymethyl on phorbol 12,13 didecanoate was replaced with hydrogen or chlorine. Schmidt and Hecker [H. Lettre and G. Wagner (eds.), Aktuelle Probleme aus dem Gebiet der Cancerologie, Vol. III, 3rd Heidelberg Symposium, pp. 98-108. Berlin: Springer Verlag, 1971.] also found that oxidation of the hydroxymethyl of phorbol 12,13 didecanoate to a carboxylic acid caused complete loss of activity in the assays used.

The hydroxymethyl group of the known phorboids (discussed in more detail in Summary of the Invention) has been thought to be required for biological activity, as detailed by Hecker (Carcinogenesis, Vol. 2, eds. Slaga, Sivak and Boutwell, Raven Press, New York, 1978, pp. 11-48 and references cited

-10-

therein). Indeed, it is stated therein that the replacement of the 20-hydroxyl in a phorbol ester "results in complete loss of biological activity". In another study, replacement of the hydroxy group of the hydroxymethyl (located at carbon 14) by chlorine or hydrogen in indolactam V gave rise to compounds with agonist activity weaker than but otherwise not distinguished from the agonist activity of the very toxic teleocidin class of tumor promoters [Irie et al., Int. J. Cancer 36 485-488 (1985)]. Thus no utility was found.

Schmidt and Hecker (Carcinogenesis, Vol. 7, ed. by E. Hecker et al., Raven Press, New York, 1982, pp. 57-63) studied the abilities of a series of diterpene phorboids to inhibit tumor promotion by the standard phorboid agonist tumor promoter phorbol 12-myristate 13-acetate (PMA). They found that, at low doses, some short-chain ester derivatives of phorbol were able to block the tumor promotion by PMA. However, all of the compounds that were active as antagonists at low doses are also very efficacious skin irritants themselves at higher doses and most of them are also known to have tumor promoting activity. Thus, these short-chain esters still have toxic inflammatory and tumor promoting activity and thus have no therapeutic value. In this publication it was also noted that a phorbol 12-ester, namely phorbol 12-myristate, was without activity as an antagonist of PMA-induced tumor promotion. Theilmann and Hecker (op. cit.) also found phorbol 12-decanoate and phorbol 12-myristate to be inactive in the functional tests used, and stated that the 13-OH must be esterified to obtain active compounds.


Summary of the Invention

This invention pertains to phorboid derivatives which block the toxic effects of the hydroxymethyl-containing phorboids and lack the toxic properties of previously available phorboids. These phorboid derivatives of very diverse structures have utility as anti-inflammatory agents, anti-cancer and anti-leukemic agents, anti-asthmatic and anti-hypertensive agents, as modulators of in vivo immune function, as stimulators of the production of lymphokines such as interferon and the interleukins, as stimulants of hair growth on bald or partially bald scalp, as central nervous system pharmaceuticals for several pathological conditions, and as xenobiotics for achieving the control of parasites. The structural features associated with the non-toxicity and protein kinase C antagonism of these compounds relate primarily to the hydroxymethyl or 1-hydroxyethyl group found in each of the toxic parents. Specific modification of the latter chemical groupings yields non-toxic anti-inflammatory compounds, whereas any of a very wide variety of changes in other parts of the parent structures, including but not limited to diterpenes, indole alkaloids, polyacetates, diaminobenzyl alcohol derivatives, aplysiatoxins, and bryostatinoids, have very markedly less effect on the overall biological properties of the derivatives, other than changes in potency. This invention also provides new compounds that discriminate between phorboid receptor subtypes, with relative binding activities differing by 10,000-fold or more.


A comparison of the structures of the pre-

viously known phorboids reveals the features that are critical for this invention, as shown below.


TYPICAL DITERPENE-TYPE PHORBOID AGONISTS




Phorbol 12-Myristate 13-Acetate (PMA)




Phorbol 12-Retinoate 13-Acetate




Ingenol 3-Tetradecanoate



Synaptolepsis Factor K₁




Pimelea Factor P₂




Daphnopsis Factor R₆


TYPICAL DITERPENE-TYPE PHORBOLD AGONISTS (cont.)




MEZEREIN




Simplexin



Pimelea Factor S₇




Gnidimacrin




Des-(Ring A)-phorbol 12-Myristate 13-Acetate


TYPICAL INDOLACTAM-TYPE PHORBOLD AGONISTS




Teleocidin B₄



Indolactam V




Lyngbyatoxin A




Blastmycetin A

-15-

Typical Diaminobenzyl Alcohol-type Phorboid Agonists




3-Acetylamino-5-(N-decyl-N-methylamino)benzyl Alcohol



6-(N-decylamino)-4-hydroxymethylindole

Typical Diacylglycerol-type Phorboid Agonists




$R_1 = \text{OLEOYL}, R_2 = \text{ACETYL}$

$R_1 = \text{STEAROYL},$

$R_2 = \text{ARACHIDONOYL}$


$R_1 = R_2 = \text{OCTANOYL}$

Typical Polyacetate-type Phorboid Agonists




(Aplysiatoxin and its debromo, bromo, and dibromo derivatives)

TYPICAL BRYOSTATIN-TYPE PHORBROID AGONISTS



Bryostatin 1



Bryostatin 2

-18-

It can be seen that the phorboids depicted have exceeding diverse structural elements, with one prominent exception, namely that each contains a hydroxymethyl or 1-hydroxyethyl group (indicated by the dashed-line boxes in each structure). In each case the phorboid depicted is among the most potent of its particular structural class, and among the classes the diterpenes, indolactams and polyacetates have members of especially high potency.

The hydroxymethyl and 1-hydroxyethyl feature common to all the classes of phorboids is the primary focus of this invention. Generally the phorboid derivatives of this invention can be represented by the formula:

$$P - G$$


wherein P represents a radical, formally derived from a parent compound, which compound:

- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
- b. activates any form of the enzyme protein kinase C; and
- c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom; and

wherein G is any group either i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or ii) singly or doubly bonded to a carbon atom immediately adjacent to the carbon atom to which the

hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound; and wherein the hydroxymethyl or 1-hydroxyethyl group of the parent compound is absent or has been replaced by G.

More specifically, the compounds of this invention are represented by the formula:



which depicts a phorboid "parent" compound radical P_o modified by a moiety $S_o E_o$.

In general, a parent compound is a compound which:

- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
- b. activates any form of the enzyme protein kinase C; and
- c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom.

For example, a parent compound can be a diterpenoid activator of protein kinase C; an aromatic heterocyclic activator of protein kinase of the indole, indene, benzofuran, or benzothiophene class; a polyacetate-derived activator of protein kinase C; an activator of protein kinase C of the diacylglycerol or diacyloxybutanol class; an activator of protein kinase C of the diaminobenzyl alcohol class; or a protein kinase C activator of the bryostatin class.

$S_o - E_o$ represents a modifying moiety which is either:

-20-

- i) singly or doubly bonded to the carbon atom of the parent compound P_o in place of the hydroxymethyl or 1-hydroxyethyl group; or
- ii) singly or doubly bonded to a carbon immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl is bonded.

In the S_o-E_o moiety, S_o can be a saturated or unsaturated, straight or branched chain of atoms which separates P_o and E_o by a linear count of at least two but not more than 6 atoms and contains one to six heteroatoms selected from oxygen, nitrogen and sulfur, a phosphorus atom, or up to 12 halogen atoms; provided that the total number of atoms does not exceed about 35, and in some cases S_o may be a single or double bond; and E_o can be a saturated or singly or multiply unsaturated group containing 1 to 8 carbon atoms and optionally containing 1 to 12 halogen atoms, 1 to 4 heteroatoms selected independently from oxygen, nitrogen and sulfur and one phosphorus atom. S_oE_o taken together may also be a hydrogen, halogen, or ketonic oxygen atom or a hydroxy, amino, or thiol group singly or doubly bonded to the carbon atom of the parent compound P_o in place of the hydroxymethyl or 1-hydroxyethyl group.

Detailed Description of the Invention

All six classes of the phorboids have one structural element in common. Each prototypical member of these classes has either a hydroxymethyl

-21-

group or a 1-hydroxyethyl group. The design of the phorboid derivatives of this invention is based on the finding that the hydroxymethyl and 1-hydroxyethyl groups which previously were thought to be required for biological activity of phorboids containing these groups can be replaced by other substituents of very diverse nature, and the resulting compounds both block the toxic effects of the hydroxymethyl-containing phorboids and lack the toxic properties associated with the previously available phorboids. These new compounds thus have utility as anti-inflammatory agents, for example. Although the replacement of the hydroxymethyl group or 1-hydroxyethyl group is very specific and leads to an extreme and profound change in biological properties; a very wide range of structural alterations in the remainder of the novel compounds can be tolerated without material loss of their anti-inflammatory activity.

The phorboid derivatives of this invention are generally represented by the formula:

P - G


The formula depicts a radical, P, derived from a parent compound, which compound:

- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
- b. activates any form of the enzyme protein kinase C; and

-22-

c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom; and wherein G is any group either i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or ii) singly or doubly bonded to a carbon atom immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound; and wherein the hydroxymethyl or 1-hydroxyethyl group of the parent compound is absent or has been replaced by G.

More specifically, the phorboid derivatives of this invention are represented by the formula:



The formula depicts a radical P_o , formally derived from a parent hydroxymethyl-containing phorboid compound, bonded to an $S_o - E_o$ moiety.

P_o represents a radical formally derived from a compound which contains an hydroxymethyl (or the equivalent 1-hydroxyethyl) group and which binds reversibly or irreversibly to a diacylglycerol-type receptor and/or activates any form of the enzyme protein kinase C. P_o may be formally derived from phorboids from any of the six classes listed below:

- i) a diterpenoid activator of protein kinase C;
- ii) an aromatic heterocyclic activator of protein kinase of the indole, indene, benzofuran, or benzothiophene class;

- iii) a polyacetate-derived activator of protein kinase C;
- iv) an activator of protein kinase C of the diacylglycerol or diacyloxybutanol class;
- v) an activator of protein kinase C of the diamino-benzyl alcohol class; and
- vi) a protein kinase C activator of the bryostatin class.

All of these phorboids contain an hydroxymethyl or 1-hydroxyethyl group which is known to be required for their toxic biological activity, such as inflammatory activity measured on the mouse ear.


S_o-E_o represents a moiety which is either:

- i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or
- ii) singly or doubly bonded to a carbon immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound.

In this S_o-E_o moiety, S_o can be a saturated or unsaturated, straight or branched chain of atoms which separates P_o and E_o by a linear count of at least two but not more than 6 atoms and contains one to six heteroatoms selected from oxygen, nitrogen and sulfur, a phosphorus atom, or up to 12 halogen atoms; provided that the total number of atoms does not exceed about 35, and in some cases S_o may be a single or double bond; and E_o can be a saturated or singly or multiply unsaturated group containing 1 to 8 carbon atoms and optionally containing 1 to 12

halogen atoms, 1 to 4 heteroatoms selected independently from oxygen, nitrogen and sulfur and one phosphorus atom. S_0E_0 taken together may also be a hydrogen, halogen, or ketonic oxygen atom or a hydroxy, amino, or thiol group singly or doubly bonded to the carbon atom of the parent compound P_0 in place of the hydroxymethyl or 1-hydroxyethyl group.


In a preferred embodiment, the phorboid derivatives of this invention are represented as follows:



wherein P_x can be selected from six different classes of compounds designated P_1 - P_6 and defined below, wherein S_x is selected from six different structural types as defined below and E_1 is as defined below.

P_1 , P_2 , P_3 , P_4 , P_5 , and P_6 represent compounds of each of the six classes of known phorboids and are defined by the formulae below.

P_1 is a radical of the formula:




in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof; wherein A¹ and A² may be individually selected from hydrogen and a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated and/or aromatic carbon- and/or heteroatom-containing substituent having not more than 34 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen and sulfur; A¹ and A² taken together complete a 5- or 6-membered carbocyclic or heterocyclic ring, optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups, which groups taken together contain a total of not more than 30 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur; A³ is a three atom chain which completes a 7-membered carbocyclic ring optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups, which groups taken together, excluding SoEo, contain not more than 7 carbon atoms, not more than 4 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen and sulfur; provided that, including S_OE_O, the middle carbon atom of A³ is not substituted by hydroxymethyl, hydroxyethyl or 2-hydroxy-2propyl;

A⁴ completes a 6- or 7-membered carbocyclic or heterocyclic ring (connected in the beta configuration to either carbon atom 9 or 10), optionally substituted by one or more straight chain or branched chain cyclic or acyclic, saturated, unsaturated and/or aromatic carbon and/or heteroatom-containing groups, the group or groups optionally completing one or two additional rings by themselves and/or one to five additional rings when taken together with A1, A2, a ring formed by A1 and A2 together, and/or a bond to carbon atom 9, which groups taken together, include not more than 40 carbon atoms, not more than 24 halogen atoms, and not more than 10 heteroatoms selected from oxygen, nitrogen, and sulfur; J¹ is selected from hydrogen, fluoro, chloro, hydroxy, amino, mono- or d-lower-alkylamino, methyl, ethyl, vinyl, ethynyl, propargyl, cyano, methoxy, ethoxy, trifluoromethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, acetoxy, propanoyloxy, acetyl, propanoyl, hydroxyacetyl, 2-hydroxypropanoyloxy, 3-hydroxypropanoyl, acetamido, propanamido, hydroxyacetamido, 2-hydroxypropanamido, or 3-hydroxypropanamido, (each of which must be situated in the beta position), or J¹ taken together with A1, A2, or a ring formed by A1 together with A2 completes a 3- to 7-membered substituted or unsubstituted carbocyclic or heterocyclic ring; J2 is selected from hydrogen, methyl, ethyl, hydroxymethyl, hydroxyethyl, vinyl, ethynyl, allyl, propargyl, n-propyl and iso-propyl; provided that if P₁ is a phorbol-12,13-dialkanoyl moiety then

-27-

S_0E_0 may not be carboxy, carbaldehyde, carbaldehyde-2,4-dinitrophenylhydrazone, or alkanoyloxymethylene, and provided that if P_1 is a phorbol 12,13-di-O-acetyl moiety then S_0E_0 may not be cyano or aldoxime.

P_2 is a radical of the formula:




in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof, wherein B^1 completes a 6-membered carbocyclic or heterocyclic aromatic ring, optionally substituted by a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups which,

-28-


taken together, contain not more than 40 carbon atoms, not more than 24 halogen atoms, and not more than 9 heteroatoms selected from oxygen, nitrogen, and sulfur, the groups being optionally connected to form 1-3 additional rings; B² is selected from oxygen, sulfur, sulfoxide, sulfone, and a carbon or nitrogen atom optionally substituted by a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups having not more than 15 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, and B² may be linked to B¹ to form an additional carbocyclic or heterocyclic ring; B³ is a 2-carbon chain optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or carbon- and/or heteroatom-containing groups, which groups taken together but excluding SoEo, contain not more than 8 carbon atoms, not more than 6 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur; provided that, including S_oE_o, the carbon atom of B³ bonded to K² as defined below does not carry ---CH₂OH ----CHCH₃OH or --C(CH₃)₂)OH; K¹ is hydrogen or is a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group containing not more than 15 carbon atoms, not more than 18 halogen atoms, and not more than 6 heteroatoms selected from oxygen,

nitrogen, and sulfur, and K^1 may be linked to B^2 or B^3 or to both B^2 and B^3 to form one or more additional carbocyclic and/or heterocyclic rings; K^2 is selected from oxygen, sulfur --NK⁶-- or ---CK⁶K⁷-- wherein K^6 is hydrogen, hydroxy, methyl, ethyl, fluoro, n-propyl, allyl, or propargyl, and K^7 is hydrogen, methyl, ethyl, halogen, trifluoromethyl or cyano; K^3 and K^4 may be the same or may differ and may each be hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group, such that K^3 and K^4 taken together contain not more than 18 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen and sulfur; K^5 is selected from oxygen, sulfur, sulfide, sulfone or --NK⁸--, --NOK⁸-- or ----CK⁸K⁹---- wherein K^8 is hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatomcontaining group containing not more than 30 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur, and K^9 is hydrogen, methyl, ethyl, n-propyl, hydroxy, halogen, allyl, propargyl, cyano, or trifluoromethyl; provided that if B^1 completes an unsubstituted carbocyclic ring, and B^2 is ---NH---, and K^1 is hydrogen, and K^2 is ---NH---, and K^3 is ---CH(CH₃)₂ in the alpha position, and K^4 is hydrogen, and K^5 is ---N(CH₃)---, then B^3 and SoEo taken together are not



wherein K^{10} is alkyl or alkanoyl.


P_3 is a radical of the formula:



in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof, wherein L^1 , L^2 and L^3 are individually selected from nitrogen or substituted or unsubstituted carbon; D^1 and D^2 each may be a bond or a substituted or unsubstituted carbon atom; D^1 may be linked to L^1 , L^2 or both L^1 and L^2 to form additional fused carbocyclic and/or heterocyclic substituted or unsubstituted rings; and D^2 may be linked to L^2 , L^3 or both L^2 and L^3 to form additional fused

carbocyclic and/or heterocyclic substituted or unsubstituted rings; D3 and D4 each are heteroatom-containing functional groups, the heteroatoms being selected from oxygen, nitrogen, and sulfur; D3 may be linked to L¹, L² or both L¹ and L² to form additional fused substituted or unsubstituted carbocyclic and/or heterocyclic rings; and D4 may be linked to L², L³ or both L² and L³ to form additional fused substituted or unsubstituted carbocyclic and/or heterocyclic rings; provided that D¹ and D3 taken together and D² and D4 taken together both embody at least one oxygen, nitrogen, or sulfur atom separated from the aromatic nucleus by zero or one intervening carbon atom; and L4 and L5 are straight chain or branched, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatomcontaining groups which, taken together, contains about 2-40 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur; provided that S_oE_o may not be hydroxymethyl or 1-hydroxyethyl.


P₄ is a radical of the formula:



-32-

in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof, wherein V^1 is a bond or is a carbon atom carrying substituents individually selected from hydrogen, methyl, and halogen; V^2 and V^3 are individually selected from oxygen, sulfur, sulfoxide, and ---NV⁴--- in which V^4 is hydrogen or a hydrocarbon radical containing not more than 30 carbon atoms; F^1 and F^2 are a straight chain or branched, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatomcontaining groups which, taken together, contain 10-40 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur; F^3 is hydrogen or a substituent selected from methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, cyano, vinyl, ethynyl, allyl, and propargyl; F^4 and F^5 each may be hydrogen or may be hydrocarbon or halogenated hydrocarbon radicals which, taken together, contain not more than 40 carbon atoms and not more than 24 halogen atoms; provided that V^4 , F^1 , F^2 , F^4 and F^5 taken together contain not more than 50 carbon atoms and provided that SoEo may not be hydrogen, methyl, chloromethyl, hydroxymethyl, mercaptomethyl, unsubstituted carboxamido, 1-hydroxyethyl or alkanoyloxy-methylene.

P_5 is a radical of the formula:




in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof, wherein U^1 and U^2 , independently, are selected from hydrogen, azide, halogen, hydroxy, C_{1-7} alkoxy, C_{1-7} alkenoxy, C_{1-7} alkynoxy, thiol, C_{1-7} alkanoyl, C_{1-7} saturated or unsaturated alkyl, and cyano; or U^1 and U^2 taken together may be an oxygen atom forming an epoxy group or may be an additional bond forming an unsaturated linkage; U^3 is selected from hydrogen, halogen, C_{1-12} alkyl, C_{1-12} alkenyl, C_{1-12} alkynyl, C_{1-12} alkoxy, C_{1-12} alkenoxy, C_{1-12} alkynoxy, and aryl or C_7-C_{12} aralkyl wherein the aryl group may be substituted by nitro, halogen, cyano, and/or

-34-

diloweralkylamino groups; U^4-U^6 , independently, are selected from hydrogen, halogen, cyano, nitro, amino, diloweralkylamino, C_{1-7} saturated or unsaturated alkyl, hydroxy, C_{1-7} saturated or unsaturated alkoxy, C_{1-7} carboalkoxy, C_{1-7} alkanoyloxy, and azide; and U^7 is selected from hydrogen, C_{1-7} saturated or unsaturated alkyl, and C_{1-7} saturated or unsaturated alkanoyl; provided that if S_0E_0 is hydroxymethyl, 1-hydroxyethyl or acetoxymethylene, then S_0E_0 may not be bonded to C29.

P_6 is a radical of the formula:




in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof, wherein W^1 is selected from hydrogen, halogen, hydroxy, C_{1-5} alkoxy, C_{1-5} alkanoyloxy and cyano;

SUBSTITUTE SHEET

W^2 is selected from oxo, hydrogen, hydroxy, cyano, C_{1-7} alkyl, C_{1-7} alkoxy, C_{1-7} alkanoyloxy, and halogen; W^3-W^5 each may be hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group, containing not more than 30 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur, and W^4 and W^5 taken together may form an additional carbocyclic or heterocyclic ring; and W^6 is hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group, containing not more than 15 carbon atoms, not more than 12 halogen atoms and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, and W^6 , taken together with W^4 and W^5 may complete an additional carbocyclic or heterocyclic ring; provided that if S_0E_0 is hydroxymethyl or hydroxyethyl then S_0E_0 may not be bonded to C25.

S_x may represent any of a broad range of connecting chains of atoms, designated S_1 , S_2 , S_3 , S_4 , S_5 and S_6 . Surprisingly, these chains may be hydrophobic in nature, with few if any polar or heteroatoms present, may be halogen-substituted, or may contain one or several polar atoms such as oxygen, nitrogen, and/or sulfur in any of numerous chemical groupings, and the resultant compounds then display the antagonistic properties described in this invention.

S_1 is a single or a double bond or is a chain of atoms of the formula:



wherein a, b, d, e, and g may independently be from 0 to 3; c and f may independently be 0 or 1; the sum of (a+b+c+d+e+f+g) is at least 1 but not more than 6; and if c and f are both 1, then the sum of (d+e) must be at least 1; R_1^1 through R_1^{12} may be the same

or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms

selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms are situated in functional groups selected from hydroxy, amino, hydroxylamine, tertiary amine oxide, Schiff's base, hydrazine, thiol, nitro, nitroso, oxime, azide, ether, acetal, ketal, thioether, aldehyde, keto, hydrazone, carboxy, ester, amide, cyano, hydrazide, carbonate, carbamate, urea, isourea, carboxamidine, imidate, guanidine, thioester, thioamide, thiocarbonate, dithiocarbonate, thiocarbamate, dithiocarbamate, thiourea, isothiourea, thioimidate, nitroguanidine, cyanoguanidine, and xanthate;

R_1^1 or R_1^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety; one of the substituents $R_1^1-R_1^{12}$ may be G^1 , as defined below, provided that the substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom; R_1^{11} or R_1^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage; one of the substituents $R_1^1-R_1^{12}$ may be linked to either the atom in P_x that carries the S_1 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different ring hetero members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted

-39-

wherein h, i, k, m, p, and q may be independently be from 0 to 3; j and n may independently be 0 or 1; if j and n are both 1 and l is 0, then the sum of (k + m) must be at least 1;

if n is 1 and o is 0, then the sum of (p + q) must be at least 1; the sum of (l + o) is 1-3; and the sum of (h+i+j+k+2l+m+n+2o+p+q) is at least 1 but not more than 7; R_2^1 through R_2^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, hydroxylamine, tertiary amine oxide, Schiff's base, hydrazine, thiol, nitro, nitroso, oxime, azide, ether, acetal, ketal, thioether, aldehyde, keto, hydrazone, carboxy, ester, amide, cyano, hydrazide, carbonate, carbamate, urea, isourea, carboximidine, imidate, guanidine, thioester, thioamide, thiocarbonate, dithiocarbonate, thiocarbamate, dithiocarbamate, thiourea, isothiourea, thioimidate, nitroguanidine, cyanoguanidine,

and xanthate; R_2^1 or R_2^2 may optionally be an

additional bond completing an unsaturated linkage to

-40-

the diterpenoid moiety; one of the substituents R_2-R_2 may be G^1 , as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom; R_2^{11} or R_2^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage; one of the substituents $R_2^1-R_2^{12}$ may be linked to either the atom in P_x that carries the S_2 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; X and X' and Y and Y' are as defined below; provided that, for all substituents R_2^1 through R_2^{12} and all substituents embodied in X, X', Y, and Y' taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10

-42-

at least 1; and the sum of $(r + s + 2t + u + 2v + 3w + 4x + y + 2z + a' + b')$ is at least 1 but not more than 7; R_3^1 through R_3^{12} may be the same or different

and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent, containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboximidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyano-guanidine, and

xanthate R_3^1 or R_3^2 may optionally be an additional

bond completing an unsaturated linkage to the

diterpenoid moiety; one of the substituents $R_3^1-R_3^{12}$

may optionally be G^1 , as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom; R_3^{11} or R_3^{12}

may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage; one of the

substituents $R_3^1-R_3^{12}$ may be linked to either the

atom in P_x that carries the S_3 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, $CH=O$, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; Y, Y', Z^1 , Z^2 , and Z^3 are as defined below;

provided that, for all substituents R_3^1 through R_3^{12}


and all substituents embodied in Y, Y', Z^1 , Z^2 , and Z^3 taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 4 or less; the total of nitrogen is 4 or less; the total of sulfur atoms is 3 or less; the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less; the

-44-

total of --OH, --SH, and --NH₂ groups together is 4
or less.

/

S_4 is a chain of atoms defined by:



wherein c' , d' , e' , h' , and i' may independently be from 0 to 3; f' and g' may independently be 0 or 1; and the sum of ($c' + d' + e' + f' + g' + h' + i'$) is at least 1 but not more than 5; R_4^1 through R_4^{12} may

be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in

functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate; R_4^1 or R_4^2 may optionally be an additional

bond completing an unsaturated linkage to the diterpenoid moiety; one of the substituents

$R_4^1-R_4^{12}$ may optionally be G^1 , as defined below,

provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety

by either zero or one intervening atom; R_4^{11} or R_4^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage; one of the

substituents $R_4^1-R_4^{12}$ may be linked to either the


atom in P_x that carries that S_4 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 ,

wherein j' , k' , m' , q' , and s' may independently be from 0 to 3; l' or r' may each be 0 or 1, but the sum of $(l' + r')$ must be 1 or 2; n' and p' may each be 0 or 1, but the sum of $(n' + p')$ must be 0 or 1; the value of o' may be 0-2; if the sum of $(n' + p')$ is 1 and l' is 0, then q' must be at least 1; if the sum of $(n' + p')$ is 1 and r' is 0, then m' must be at least 1; and the sum of $(j' + k' + l' + m' + n' + o' + p' + q' + r' + s')$ is at least 1 but not more than 5; R_5^1 through R_5^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate; R_5^1 or R_5^2 may optionally comprise an additional bond completing an unsaturated linkage the diterpenoid moiety; one of the substituents R_5^1 - R_5^{12} may optionally represent G^1 ,

as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom; R_5^{11} or R_5^{15} may optionally comprise an additional bond to E_1 , thereby completing an unsaturated linkage; one of the substituents R_5^1 - R_5^{12} may be linked to either the atom in P_x that carries the S_5 chain or to an atoms in adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from the halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, $CH=O$, acetoxy, amino, mono- or dialkylamino containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; Q, Q', X, X', and Y are as defined below; provided that, for all substituents R_5^1 through R_5^{12} and all constituents Q, Q', X, X', and Y taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 4 or less; the total if nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less; the total of oxygen, nitrogen, and sulfur

atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less; the total of --OH, --SH, and --NH₂ groups together is 4 or less.

S₆ is a chain of atoms defined by:



wherein t', u', v', w', x', y', z' and a'' may each be 0 or 1; the sum of (t' + u' + v' + 2w' + x' + 2y' + z' + a'') must be 0-2; b'', d'', e'', f'', h'', j'' and k'' may each independently be 0 or 1; c'', g'', i'', and l'' may each independently be 0-3; if d'' and j'' are both 1, then the sum of (g'' + i'') must be at least 1; if either j'' or k'' is 1, then l'' must be at least 1; if b'' is 1, then the sum of

(c'' + g'' + h'' + i'' + l'') must be at least 1; if d'' is 1, then the sum of (g'' + h'' + i'' + l'') must be at least 1; and the sum of (t' + u' + v' + 2w' + x' + 2y' + z' + a'' + b'' + c'' + d'' + e'' + 2f'' + g'' + 2h'' + i'' + j'' + k'' + l'') must be 0-5; R_6^1 through R_6^{12} may be the same or different and each may be either hydrogen or a saturated or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and


xanthate; R_6^1 or R_6^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety; R_6^3 or R_6^4 may optionally comprise an additional bond to G^2 as defined below, thus completing an unsaturated linkage; R_6^5 or R_6^6 may optionally comprise an additional bond to G^2 , thus completing an unsaturated linkage if a'' is 0; one of

-52-

the substituents $R_6^1-R_6^4$ may optionally be G^1 , as defined below; R_6^{11} or R_6^{12} may optionally comprise an additional bond to E_1 , thereby completing an unsaturated linkage; one of the substituents $R_6^1-R_6^4$ may be linked to either the atom in P_x that carries the S_6 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-6 membered ring optionally containing 1-3 hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, $CH=O$, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; Q, Q', Q'', Q''', X, X', X'', Y, Y', and Z^4 , are as defined below; provided that, for all substituents R_6^1 through R_6^{12} and all constituents Q, Q', Q'', Q''', X, X', X'', Y, Y', Z^4 , and G^2 taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 12 or less; the total of halogen atoms is 12 or less; the total

of oxygen atoms is 5 or less; the total of nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less, the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less; the total of --OH, --SH, and --NH₂ groups together is 4 or less.

X, X', X'' may be the same or different and are selected from:



wherein R¹_X and R²_X may independently be hydrogen;




R_X^1 through R_X^5 may be the same or different and each may be a saturated or singly or multiple unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboxamide, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine; R_X^1 may

R_X^1 may optionally represent an additional bond to the diterpenoid moiety, thus completing an unsaturated linkage; one of the substituents

$R_X^1-R_X^5$ may optionally represent G, as defined below, provided that the substituent is bonded to an atom that is separated from the parent "P" by either zero or one intervening atom; and one of the substituents $R_X^1-R_X^5$ may be linked to either the atom in P_X that carries the chain containing X, X', and/or X'' or to another atom in P_X adjacent thereto, to form a saturated, unsaturated or aromatic, heterocyclic 4-8 membered ring optionally containing

1-3 other identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxypropyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totaling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy.

Y and Y' may be the same or different and are selected from:








wherein R_Y^1 and R_Y^2 (cis or trans) may be the same or different and each may be hydrogen or may comprise a saturated or singly or multiply unsaturated, straight or branched acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboximidine, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine; R_Y^3 and R_Y^4 (cis or trans) may be the same or different and each may be hydrogen or may be a saturated or singly or multiply unsaturated, straight or branched acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboximidine,

guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine, with

the exception that neither R_Y^3 nor R_Y^4 may be a halogen atom alone; one of the substituents R_Y^1 - R_Y^4 may optionally represent G^1 , as defined below, provided that the substituent is bonded to an atom that is separated from the diterpenoid moiety by either zero or one intervening atom; and one of the substituents R_Y^1 R_Y^4 may be linked to either the atom in P_X that carries the chain containing Y and/or Y' or to an atom in adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy.

-58-

Z^1 is selected from:



wherein R_z^1 is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{1-6} acyl, C_{1-6} halogenated acyl, C_{1-6} monohydroxyacyl, and C_{2-6} hydroxyalkyl, cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or halogen, or, R_z^1 may comprise an additional bond to P_x , thus completing an unsaturated linkage; R_z^2 is hydrogen, a saturated

or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ halogenated alkyl, C₁₋₆ acyl, C₁₋₆ halogenated acyl, C₁₋₆ monohydroxyacyl, and C₂₋₆ hydroxyalkyl, cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or halogen;

R_Z³ independently may have the values specified

above for R_Z²; R_Z⁴ independently may have the values

specified above for R_Z¹; R_Z⁵ may be hydrogen or a

saturated or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkoxy, C₁₋₆ halogenated alkyl, or phenyl or benzyl optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or

halogen; R_Z⁶ independently may have the values

specified above for R_Z⁵; one of the substituents

R_Z¹-R_Z⁶ may be linked to either the atom in P_X

that carries the chain containing Z¹ or to atom in P_X adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8









membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4

identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃,

COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; provided that only one of the substituents

R_Z¹-R_Z⁶ may be substituted or unsubstituted phenyl or benzyl.

Z² is selected from:



wherein R_Z⁷ independently may have the values

specified for R_Z¹; R_Z⁸ is hydrogen, a saturated

or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ halogenated alkyl, C₂₋₆ hydroxyalkyl in which the hydroxy group may be esterified to the acetyl or propionyl ester, and cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, nitro,

cyano, trifluoromethyl, and/or halogen; R_Z⁹

independently may have the values specified above

above for R_Z⁸; R_Z¹⁰ independently may have the values

specified above for R_Z⁷; R_Z¹¹ may be hydrogen, a

saturated or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ halogenated alkyl, and cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or

halogen; R_Z¹²-R_Z¹³ independently may have the

values specified above for R_Z²; R_Z¹⁴-R_Z¹⁶ inde-

pendently may have the values specified above for

R_Z¹¹; R_Z¹⁷ and R_Z²⁰ individually may be hydrogen or



a substituent selected from C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkylthio, phenoxy or thiophenoxy optionally substituted by methyl, halogen, and/or nitro; or amino optionally mono- or disubstituted by C₁₋₄ alkyl or monosubstituted by cyano, nitro, or phenyl optionally substituted by halogen and/or nitro;

R_Z¹⁸ and R_Z¹⁹ individually may be --O--, --S--, or

--NR_Z²¹, wherein R_Z²¹ may be hydrogen, C₁₋₄ alkyl,

2-hydroxyethyl, 2-hydroxy-n-propyl, 2-acetoxyethyl, or 2-acetoxy-n-propyl; one of the substituents R_Z^{17} , R_Z^{20} or R_Z^{21} optionally may be linked to either the atom in P_x that carries the chain containing Z^2 to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totaling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive hydroxyacetyl and hydroxyacetoxy; provided that only one of the substituents R_Z^{17} - R_Z^{21} may comprise or carry a substituted or unsubstituted phenyl or benzyl.


Z^3 is selected from:



-63-

wherein R_Z^{22} and R_Z^{25} independently may have the values specified above for R_Z^{11} ; R_Z^{23} and R_Z^{24} independently may have the values specified above for R_Z^8 ; R_Z^{22} or R_Z^{25} optionally may be linked to either the atom in P_x that carries the chain containing Z^2 or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; provided that at most one of the substituents R_Z^{22} - R_Z^{25} may comprise or carry a substituted or unsubstituted phenyl or benzyl moiety.

Z^4 and $Z^{4'}$ independently may be:






wherein R_Z^{26} may be hydrogen or may be a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, in which substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboximidine, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine;

R_Z^{26} may optionally comprise one additional bond either to the parent P_x or to G^2 , thus completing an unsaturated linkage; R_Z^{26} may be linked to either the atom in P_x that carries the S_6 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-6 membered ring optionally containing 1-3 identical or different ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy,

methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, CF_3 , OCF_3 , SH,

SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy.

z⁵ and z^{5'} independently may be:






wherein R_z²⁷ is hydrogen, a saturated or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ halogenated alkyl, C₁₋₆ acyl, C₁₋₆ halogenated acyl, C₁₋₆ monohydroxyacyl, and C₂₋₆ hydroxyalkyl, cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or

-66-

halogen, or R_Z^{27} may comprise an additional bond to the parent P_X or to G^2 , thus completing an unsaturated linkage; R_Z^{28} is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{1-6} acyl, C_{1-6} halogenated acyl, C_{1-6} monohydroxyacyl, and C_{2-6} hydroxyalkyl, cyclohexyl, or phenyl or benzyl either of which may optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or halogen, or R_Z^{28} may comprise an additional bond to G^2 , thus completing an unsaturated linkage; R_Z^{29} independently may have the values specified above for R_Z^{27} ; and R_Z^{30} independently may have the values specified above for R_Z^1 .

M and M' independently may be:





wherein R_M^1 and R_M^2 may be the same or different and each may be hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, in which the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, nitroguanidine, and cyanoguanidine; R_M^1 may optionally comprise an additional bond to the parent P_x group, thus completing an unsaturated linkage;

R_M^1 or R_M^2 may optionally represent G^1 , as defined below, provided that said substituent is bonded to an atom that is separated from the parent P_x by either zero or one intervening atoms; R_M^1 or R_M^2 may be linked to either the atom in P_x that carries the chain containing M and/or M' or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 5-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro,

hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH , SCH_3 , SOCH_3 , SCF_3 , COOH , COOCH_3 , COCH_3 , CH=O , acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy.

Q-Q''' independently may be:



wherein R_Q^1 and R_Q^2 may be the same or different and each may be hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur,


such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, ether, thioether, carboxy, ester, amide, cyano, nitroguanidine, and cyanoguanidine; R_Q^1 may optionally represent G^1 , as defined below, provided that R_Q^1 is bonded to an atom that is separated from the parent P_x by either zero or one intervening atom; R_Q^1 may be linked to either the atom in P_x that carries the chain containing Q and/or Q' or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 5-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxypropyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $CH=O$, acetoxy, amino mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; G^1 comprises a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing

-70-

1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methyl acetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy; G² completes a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy, and the ring may be connected directly by single or double bonds to adjacent atoms.

The capping group E_0 that terminates the connecting chain also may be selected from any of a surprisingly broad array of chemical groupings. These may include, without limitation, hydrophobic entities such as alkyl, hydrogen, and halogenated alkyl, or may include, without limitation, quite hydrophilic moieties, such as hydroxy, thiol, carboxy and carboxy esters, amines, etc. Indeed, the single restriction appears to be that S_0E_0 taken together should not be hydroxymethyl or 1-hydroxyethyl bonded in the usual position in the parent compounds, since such compounds correspond to the toxic parent natural products.

Thus, E_0 may be a moiety E_1 , wherein E_1 is selected from =O, =S, =NH, =NOH, =N--NH₂, hydrogen, halogen, --OH, --SH, --NH₂, --NH--NH₂, --N₃, --NO, --NO₂, --NHOH, --ONH₂, or is selected from :



wherein T^1 is selected from --O--, --S--, and --NH--; T^2 is selected from =O, =S, and =N--R_E⁶, in which R_E⁶ may be hydrogen, cyano, or nitro; T^3 is selected from hydrogen, --OH, --NH₂, --SH, N₃, --NH--NH₂, and --NH--OR_E⁷, in which R_E⁷ may be

hydrogen C_{1-3} , alkyl, or C_{1-3} acyl; R_E^1 is selected from hydrogen, halogen, hydroxy, nitro, nitroso, cyano, azide, $--NH_2$, $--NH--OH$, $--SH$, $--O--NH_2$, $--NH--NH_2$, $--T^1--C=T^2--T^3$, and $--C=T^2--T^3$;

R_E^2 and R_E^3 are individually selected from hydrogen, $--C=T^2--T^3$, cyano, nitro, azide, and halogen; and R_E^4 and R_E^5 are individually selected from hydrogen, halogen, cyano, nitro, $--C=T^2--T^3$, $--T^1--C=T^2--T^3$, and $--CR_E^1R_E^2R_E^3$.

In addition to the compounds described above, we have also found that compounds of the diterpene class, with the usual hydroxymethyl or 1-hydroxyethyl group intact and having at least on substituent other than hydrogen at C_{12} and a hydroxy, amino, thiol, hydroxymethyl, mercaptomethyl, aminomethyl, 2-hydroxyethyl, carboxy, unsubstituted carboxamido, or unsubstituted aminocarbonyloxy group in the alpha configuration at C_{13} , display useful properties as anti-inflammatory agents. Thus, in this very unusual case a minor change in the diterpene moiety at a location other than the hydroxymethyl leads to a very substantial and useful change in biological properties. This is in contrast to the belief that diterpenes such as phorbol 12-decanoate, 12-dodecanoate and 12-myristate are biologically inactive, and it is especially in contrast to the usual association of the hydroxymethyl or 1-hydroxyethyl groups with pro-inflammatory activity.

The compounds of this invention have been found to possess valuable pharmacological properties. They block inflammation, block proliferation of cancer cells, and induce production of thrombolytic activity in human and veterinary medicine. These effects can be demonstrated, for example, by use of standard mouse ear inflammation tests by established agonists such as PMA and the ionophore A23187, by the inhibition of proliferation of human cancer cells in culture by induction of differentiation, and by measurement of fibrinolytic activity in cultured cells.

These compounds also show selective effects as antagonists or protein kinase C in some cases, as noninflammatory agonists for protein kinase C in other cases, and as selective ligands for protein kinase C and/or for phorboid receptors.

Thus, these compounds can be used as agents for the abrogation of pathophysiological conditions and disease states in applications such as anti-inflammatory, anti-psoriatic, anti-cancer, anti-ulcer, anti-hypertensive, anti-asthma, anti-arthritis, anti-autoimmune, anti-nociceptive, anti-secretory, anti-parasitic, anti-amoebic, anti-HTLV-III/LAV viral replication, and any other application in which pathological involvement of protein kinase C is found.

Furthermore, the non-toxic agonists among the compounds of this invention may be used to achieve desired physiological results such as interferon release, interleukin induction, tumor necrosis factor production, immune system stimulation and/or

-74-

reconstitution, insulin secretion, insulinomimetic activity, acceleration of wound healing, improvement in central nervous system functions such as memory and learning and abrogation of the symptoms or progress of Alzheimer's disease, and any other application for which desirable actions of protein kinase C are found.

As receptor subtype- and/or protein kinase C subtype-selective ligands, the compounds of this invention also have very valuable application as experimental agents for research into the role of protein kinase C and/or phorboid receptors in important biological processes and in human and veterinary diseases. Thus, their value extends to their use as pharmacological tools for in vitro and in vivo research, in a manner similar to the important roles that selection agonists and antagonists have played in the studies of the mechanism of action of adrenergic, dopaminergic, opiate, benzo-diaepine, cholinergic, and serotonergic receptor systems, among others.

In addition, the compounds can be used in in vitro diagnostics (e.g., in an assay for protein kinase C). They are also useful as intermediates in the production of other drugs, e.g., as described in the present invention.

The compounds of this invention are generally administered to animals, including but not limited

-75-

to fish, avians, and mammals including humans.

The pharmacologically active compounds of this invention can be processed in accordance with conventional methods of galenic pharmacy to produce medicinal agents for administration to patients, e.g., mammals including humans.

The compounds of this invention can be employed in admixture with conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral (e.g., oral) or topical application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohols, gum arabic, vegetable oils, benzyl alcohols, polyethylene glycols, gelatine, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds. They can also be combined where desired with other active agents, e.g., enzyme inhibitors, to reduce metabolic degradation.

-76-

For parenteral application, particularly suitable are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Ampoules are convenient unit dosages.

For enteral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules. A syrup, elixir, or the like can be used wherein a sweetened vehicle is employed.

Sustained or directed release compositions can be formulated, e.g., liposomes or those wherein the active compound is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc. It is also possible to freeze-dry the new compounds and use the lyophilizates obtained, for example, for the preparation of products for injection.

For topical application, there are employed as nonsprayable forms, viscous to semi-solid or solid forms comprising a carrier compatible with topical application and having a dynamic viscosity preferably greater than water. Suitable formulations include but are not limited to solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, aerosols, etc., which are, if desired, sterilized or mixed with auxiliary agents, e.g., preservatives, stabilizers, wetting agents, buffers or salts for influencing osmotic pressure,

-77-

etc. For topical application, also suitable are sprayable aerosol preparations wherein the active ingredient, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant, e.g., a freon.

Generally, the compounds of this invention are dispensed in unit dosage form comprising 0.01 to 1000 mg in a pharmaceutically acceptable carrier per unit dosage. They are incorporated in topical formulations in concentrations of about 0.01 to 10 weight percent.

It will be appreciated that the actual preferred amounts of active compound in a specific case will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, and the particular situs and organism being treated. Dosages for a given host can be determined using conventional considerations, e.g., by customary comparison of the differential activities of the subject compounds and of a known agent, e.g., by means of an appropriate, conventional pharmacological protocol.

Starting materials for the synthesis of the compounds of this invention may be obtained from any of a wide variety of natural sources, as described in the literature for the diterpenes (R. Schmidt and E. Hecker, Fortschritte D. Chemie Organischer Naturstoffe 31 377-467 (1974) and F. J. Evans and C. J. Soper, Lloydia 41 193-233 (1978), and references

cited therein), and indole alkaloids and aplysia-toxins (T. Sugimura, Gann 73 499-507 (1982), and references cited therein). Furthermore, the diterpene, indole alkaloid, diacylglycerol, and diamino-benzyl alcohol compounds are available by synthesis de novo (See Y. Endo et al., Chem. Pharm. Bull. 32 358-361 (1984), Tetrahedron 42, 5905-5924 (1986), and references cited therein; P. A. Wender, Am. Chem. Soc. National Meeting, Chicago, IL, 9 September 1985, Abstract #7, and P.A. Wender et al. Proc. Nat. Acad. Sci. 83, 4214-4218, 1986).

Given starting phorboids containing hydroxymethyl or 1-hydroxy-ethyl groups, the means for modifying the hydroxymethyl or 1-hydroxyethyl group to produce the compounds of this invention will be obvious to workers with ordinary skill in synthetic organic chemistry. The hydroxymethyl or 1-hydroxyethyl groups may be modified once other regions of the molecule have been suitably protected, using certain favorable methodologies. Once the phorboid nucleus is protected, the hydroxymethyl/1-hydroxyethyl may for example be conveniently capped under very mild conditions by reaction with a substituted or unsubstituted alkyl, aryl, or aralkyl isocyanate in the presence of a catalyst such as dibutyltin dilaurate. The resulting compounds lack the toxic inflammatory activity of the phorboid from which they were derived, and have themselves anti-inflammatory utility.

Conversion of the hydroxy group to a halogen or pseudohalogen not only in itself provides active anti-inflammatory compounds, but also permits displacement of the resultant electrophile by a very wide range of nucleophiles. Particular examples, with limitation, would be reaction with ammonia, methylamine, sodium cyanide, N-methyl-2-hydroxyethylamine, 1[H]-tetrazole, or with the sodium salt of 2-mercaptoethanol, 3-mercaptopropanol, or of 2-hydroxymethylphenol. Many variations may be executed, as described in standard textbooks of synthetic organic chemistry, such as J. March, Advanced Organic Chemistry, Third Edition, Wiley-Interscience New York, 1985.

For those phorboids wherein the hydroxy of the hydroxymethyl group is allylic, such as phorbol, ingenol, and resiniferonol esters, the replacement of the hydroxy by chloro or preferably by bromo or iodo yields compounds that can be conveniently homologated by reaction with activated zinc in the presence of an electrophile such as, without limitation, an aldehyde, ketone, epoxide, or oxetane; the resultant compounds have one or more methylenes inserted between the original hydroxy group and the methylene to which it was attached. An illustration of this would be the reaction of 20-deoxy-20-chlorophorbol 12-myristate 13-acetate with an excess of formaldehyde and an excess of zinc in the presence of tetrahydrofuran and saturated ammonium chloride solution with vigorous stirring for 24

-80-

hours. The resultant 20-homophorbol 12-myristate 13-acetate carries a 2-hydroxyethyl group attached to C6 instead of the usual hydroxymethyl group, and has good anti-inflammatory activity.

Alternatively, the hydroxy group in question may be oxidized to an aldehyde or keto group and then reacted via Wittig chemistry to obtain chain-extended compounds terminated by many different capping groups. In particular, a phorbol 12,13-diester may be selectively protected at the 4 and 9 hydroxy groups using trimethylsilyl trifluoromethanesulfonate, followed by reaction with manganese dioxide to obtain a protected aldehyde. The latter compound may be successfully treated with strong wittig reagents such as the lithium salt of 2-hydroxyethylidene triphenylphosphorane, followed by deprotection with tetrabutylammonium fluoride to obtain the corresponding chain-extended, 20,21-didehydro-21,22-dihomophorbol diester.

The hydroxymethyl group of suitable phorboids may be oxidized to the carboxylic acid level by methods well-known in the art, and this carboxylic group may be activated for condensation reactions by any of a number of well-known methods, e.g. by conversion to an acyl halide or to an active ester such as the N-hydroxysuccinimide ester. The resultant activated carboxyl may then be easily converted to simple or multifunctional ester, amide, or thio-ester derivatives by reaction with alcohols, amines, or thiols respectively, alone or in the presence of condensation catalysts. For example, 4-carboxy-

-81-

6-(N-decanoylamino)indole may be converted to its N-hydroxysuccinimide ester by reaction with one equivalent of dicyclohexylcarbodiimide and one equivalent of N-hydroxysuccinimide in acetonitrile/tetrahydrofuran/methylene chloride suspension. The product N-hydroxysuccinimide ester is purified and then reacted with 3-amino-1,2-propanediol in tetrahydrofuran to obtain 4-(N-2,3-dihydroxypropyl-carboxamido)-6-(N-decanoylamino)indole.

The use of the methods of total synthesis as described in the literature cited above, with obvious adaptations, permits specific modifications of the parent structures in the diterpene, indolactam, diacylglycerol, and diaminobenzyl alcohol groups, by established techniques in the art of synthetic chemistry, to obtain modified parent structures embodying alterations at the hydroxymethyl or 1-hydroxyethyl groups and having anti-inflammatory activity.

To further illustrate the synthesis of the compounds of this invention, the modified indolactam, 1,2,4,5,6,8-hexahydro-5-methyl-2-(1-methylethyl)-3H-pyrrolo(4,3,2-gh)-1,4-benzodiazonin-3-one, may be prepared from N-BOC-4-nitrotryptophanol [Y. Endo *et al.*, Tetrahedron, 42, 5905-5924 (1986)] by the application of several routes obvious to workers with ordinary skill in synthetic chemistry. For example, preparation of N-BOC-4-nitrodeoxytryptophanol may be accomplished by reduction of the

phenylselenium derivative with triphenyltinhydride [D. Clive et al., Chemical Communications, 41-42 (1978)] or by reduction of the mesylate derivative with lithium triethylborohydride [R.W. Holder and M.G. Matturro, J. Org. Chem., 42, 2166-2168 (1977)] or by several other routes. From the deoxy derivative the synthesis could proceed in the manner described by Y. Endo et al., loc cit., for the hydroxy derivative. Specifically, the resulting substituted indolylvaline methyl ester is hydrolysed and the resulting acid is converted to the N-succinimidyl ester. Upon cleavage of the BOC group under acidic conditions cyclization to the lactam occurs directly to provide all four stereoisomers, i.e., 2R,5R-, 2R,5S-, 2S,5S-, and 2S,5R-1,2,4,5,6,8-hexahydro-5-methyl-2-(1-methylethyl)-3H-pyrrolo (4,3,2-gh)-1,4-benzodiazonin-3-one. These stereoisomers may be obtained separately either by beginning the synthesis with optically active N-BOC-4-nitrotryptophanol or by separation from the mixture by chromatography on an enantioselective column packing [D. Armstrong, Analytical Chem., 59, 84-91A (1987)]. Similarly, the further modified indolactam, 1-(1-oxobutyl)-1,2,4,5,6,8-hexahydro-5-methyl-3H-pyrrolo (4,3,2-gh)-1,4-benzodiazonin-3-one, may be prepared. Specifically, hydrogenation of N-BOC-4-nitrodeoxytryptophanol over palladium on carbon will provide N(2')-BOC-4-aminodeoxytryptophanol. Alkylation of this compound with methyl bromoacetate affords N-[3-(N-BOC-2-aminopropyl)-4-indolyl]-

glycine methyl ester [Y. Endo, et al., Chem. Pharm. Bull, 30, 3457-3460 (1982)]. Acylation of this material with butanoyl chloride in the presence of potassium carbonate or pyridine will provide N-butanoyl-N-[3-(N-BOC-2-aminopropyl)-4-indolyl]glycine methyl ester. This latter material may be converted to 1-(1-oxobutyl)-1,2,4,5,6,8-hexahydro-5-methyl-3H-pyrrolo(4,3,2-gh)-1,4-benzodiazonin-3-one by application of the methods of Y. Endo, et al., loc cit. (1986). The enantiomers, 5R- and 5S-, may be obtained separately by beginning with optically active materials as described above or by separation at the final stage by chromatography also as described above.

This invention is illustrated further by the following examples.

EXAMPLE 1

One gram of sodium metal was dissolved in 50 ml methanol, and 0.044 ml of this solution was placed in a test tube. Then 2.63 grams distilled 2-mercaptoethanol was dissolved in 50 ml acetonitrile, and 0.044 ml of this solution was added to the test tube. Then 20 mg 20-deoxy-20-chlorophorbol 12,13-dibutyrate were dissolved in 0.25 ml acetonitrile in a capped, nitrogen-flushed test tube. This latter solution was then rapidly treated with the methoxide/mercaptoethanol solution. An immediate precipitate formed. After 7 minutes, the reaction was freed of solvent and treated with 1 ml water and

-84-

1 drop acetic acid. This residue was partitioned between water and ethyl acetate, followed by drying of the organic phase over sodium sulfate. Silica gel preparative liquid chromatography using hexane-ethyl acetate mixtures yielded 9.5 mg of 20-deoxy-20-(2-hydroxyethylthio)phorbol 12,13-dibutyrate, which could not be crystallized.

EXAMPLE 2

Further preparative liquid chromatography of the reaction mixture from Example 1 using hexane-ethyl acetate on silica gel yielded 5.7 mg 20-deoxy-20-(2-hydroxyethylthio)phorbol 12-butyrate, the latter being the more polar compound. This compound could not be crystallized.

EXAMPLE 3

To 0.1 gram 20-deoxy-20-chlorophorbol 12-myristate 13-acetate in 1 ml acetonitrile was added 0.5 ml of a solution of 246 mg 2-mercaptoethanol and 436 mg 2,4,6-collidine in 10 ml acetonitrile, followed by 50 mg diisopropylethylamine in 0.2 ml acetonitrile and 0.1 ml t-butyl methyl ether. Ten minutes later the reaction was treated with 0.107 mmoles sodium methoxide and 0.225 mmoles 2-mercaptoethanol in 0.25 ml methanol. Five minutes later the proportion of sodium methoxide/mercaptoethanol was doubled. After 10 minutes the reaction was stopped by addition of 2 drops acetic acid and 0.5 ml water. The organics were extracted into ethyl acetate, washed once with water, and dried over sodium

-85-

sulfate. After solvent removal, the crude residue was purified by preparative liquid chromatography on silica gel using hexane/ethyl acetate 60/40. The product was 87 mg 20-deoxy-20-(2-hydroxyethylthio)-phorbol 12-myristate 13-acetate in high purity. The compound did not crystallize.

EXAMPLE 4

Twenty-five mg of 20-deoxy-20-chlorophorbol 12-myristate 13-acetate was dissolved in 0.2 ml ethylene glycol and 0.2 ml acetonitrile. This solution was treated with 0.13 ml of a solution of 200 mg sodium metal in 20 ml ethylene glycol over a period of 40 minutes. The reaction was partitioned between water and ethyl acetate, and the separated organics were dried over sodium sulfate. After removal of the ethyl acetate, the crude 20-deoxy-20-chlorophorbol 12-myristate was dissolved in 0.8 ml acetonitrile and treated with 0.18 ml of a solution of 0.31 ml 2-mercaptoethanol, 0.5 ml acetonitrile, and 0.5 ml of 1% sodium in methanol. After 40 minutes, an additional 0.04 ml of 1% sodium in methanol was added. Ten minutes later the reaction was stopped with 1 drop of acetic acid. After removal of the solvents in a stream of nitrogen, the residue was partitioned between ethyl acetate and pH 8 potassium phosphate. The organics were dried over sodium sulfate and freed of solvent prior to preparative liquid chromatographic purification using silica gel and hexane/ethyl acetate

-86-

45/55. The product, 37 mg of 20-deoxy-20-(2hydroxyethylthio)phorbol 12-myristate, could not be crystallized.

EXAMPLE 5

25 mg 20-deoxy-20-chlorophorbol 12-myristate 13-acetate was dissolved in 0.4 ml acetonitrile. To this was added 0.1 ml of a solution of 29.6 mg 2-(methylamino)ethanol in 1 ml acetonitrile. After 70 minutes and additional 0.1 ml of the same amine solution was added. After an additional 70 minutes, 0.2 ml more amine solution was added. After 6.6 hours of total reaction time, the reaction was diluted with 4 ml methylene chloride and subjected to preparative liquid chromatography on silica gel using methylene chloride/methanol 92/8 followed by re-purification on silica gel using methylene chloride/methanol 96/4. The product, 20-deoxy-20-[(2-hydroxyethyl)methylamino]phorbol 12-myristate 13-acetate, 14 mg, could not be crystallized.

EXAMPLE 6, 7, and 8

100 mg of phorbol 12,13-bis(2,4-difluorophenylacetate) were dissolved in 1.5 ml methylene chloride and the solution was set at 0°C. Then 26.2 mg diethylaminosulfur trifluoride in 0.5 ml methylene chloride were added dropwise during 1 minute. After 40 minutes, 0.2 ml more diethylaminosulfur trifluoride was added. After 10 more more minutes, the reaction was shaken with 2 ml pH 8 potassium phosphate buffer, after which the organics were separated and dried over sodium sulfate. The reaction

-87-

was repeated twice more, and the combined reaction products were freed of solvent, taken up in 10 ml ethyl acetate, and sucked through a funnel containing a layer of silica at the bottom, a layer of sodium sulfate in the middle and sodium chloride at the top. After washing the funnel contents with 50 ml ethyl acetate the combined eluants were freed of solvent and repeatedly chromatographed on silica preparative liquid chromatography columns using hexane/ethyl acetate 85/15 solvent mixtures. The products were 20-deoxy-20-fluorophorbol 12,13-bis(2,4-difluorophenylacetate), 40 mg; 12-beta,13-bis-(2,4-difluorophenylacetoxo)-4,9-dihydroxy-1,6(20),7-tigliatrien-3-one, 25 mg; and 12-beta,13-bis(2,4-difluorophenylacetoxo)-4,9-dihydroxy-7-fluoro-1,6(20)-tigliaden-3-one, 40 mg; none of which could be crystallized.

EXAMPLE 9

Five hundred-fifteen mg of 4-carbomethoxy-6-(N-decanoylamino)indole [prepared by the method of Wender et al., PNAS, 83, 4214-4218 (1986)] was dissolved in 60 mL of tetrahydrofuran. This solution was treated with 3 mL of a 1N KOH solution in water and also with 5 mL of methanol. The mixture was heated at 80°C for 32 h. During this period another 2.5 mL of 1N KOH was added in two portions. After cooling the mixture was concentrated in vacuo. The mixture was then diluted with water and acidified with concentrated hydrochloric acid. The mixture was then extracted with methylene chloride.

-88-

The organic layers were dried over sodium sulfate and concentrated to afford 300 mg of 4-carboxy-6-(N-decanoylamino) indole (60% yield). mp 248-50°C.

EXAMPLE 10

A suspension of 470 mg of 4-carboxy-6-(N-decanoylamino)indole and 427 mg of N-hydroxysuccinimide in 100 mL of acetonitrile, 10 mL of methylene chloride and 10 mL of tetrahydrofuran was prepared. This suspension was stirred vigorously by a magnetic stirring bar as a solution of 540 mg of dicyclohexylcarbodiimide in 20 mL of acetonitrile was slowly added over a period of 1 h. The mixture was stirred vigorously for 72 h. It was then concentrated in vacuo and diluted with ethyl acetate. The resulting mixture was filtered to remove the copious precipitate. The filtrate was washed with water, dried over sodium sulfate and concentrated. Treatment of the resulting mixture with hexane/ethyl acetate 50:50 followed by filtration afforded 600 mg of 4-carboxy-6-(N-decanoylamino)-indole N-succinimidyl ester, mp 154-5°C.

EXAMPLE 11

To a solution of 89 mg of 3-amino-1,2-propanediol in 20 mL of tetrahydrofuran was added 136 mg of 4-carboxy-6-(N-decanoylamino)indole N-succinimidyl ester. The solution was stirred for 48 h. After concentration in vacuo the mixture was purified by preparative liquid chromatography using silica gel and methylene chloride/methanol 92:8. The product,

135 mg of 4-(N-2,3-dihydroxypropylcarboxamido)-6-(N-decanoylamino)indole, was recrystallized from methanol-methylene chloride, mp 139-41°C.

EXAMPLE 12

To a solution of 113 mg of triethylamine in 20 mL of tetrahydrofuran was added, first, 86 mg of 2-aminoethanethiol hydrochloride and, then, 139 mg of 4-carboxy-6-(N-decanoylamino)indole N-succinimidyl ester. The solution was stirred for 48 h and then concentrated in vacuo. Purification by liquid chromatography using silica gel and methylene chloride/methanol 96:4 afforded two products. The earlier eluting product, 23 mg of 4-(N-2-mercaptoethylcarboxamido)-6-(N-decanoylamino)indole, decomposed at 200-5°C. The later eluting product, 22 mg of 4-(S-2-aminoethylthiolcarboxy)-6-(N-decanoylamino)-indole, was recrystallized from methanol, mp 192-3°C.

EXAMPLE 13

To a solution of 145 mg of 4-carboxy-6-(N-decylamino)-indole N-hydroxysuccinimidyl ester in 20 mL of tetrahydrofuran was added 103 mg of piperidinemethanol. The solution was heated at reflux for 5 days, at which time another 100 mg of piperidinemethanol was added and heating continued for another day. The solution was then concentrated in vacuo and purified

-90-

by liquid chromatography using silica gel and methylene chloride/iso-propyl alcohol (93:7). In this manner 21 mg of 4-(2-hydroxymethylpiperidinocarbonyl)-6-(N-decanoylamino)indole was obtained, M.P. 126-129°C.

EXAMPLE 14

One hundred milligrams of phorbol 12-myristate 13-acetate was dissolved in 1 ml tetrahydrofuran. To this solution was added 11.5 microliters of methyl isocyanate, followed immediately by 20 microliters of a 10% by weight solution of di-butyltin dilaurate in tetrahydrofuran. After three hours an additional 11.5 microliters of methyl isocyanate was added. Sixteen hours later 30 microliters of the reaction solution was applied to a silica gel TLC plate and developed with hexanes/ethyl acetate 46/54. The band at Rf 0.4 was scraped off and the product was eluted from the silica with acetone. Removal of the solvent in a stream of nitrogen gave 2.8 mg of phorbol 12-myristate 13-acetate 20-methylcarbamate as a glassy solid for spectroscopic analysis and bioassay.

EXAMPLE 15

Ninety-eight mg of 20-deoxy-20-chlorophorbol 12-myristate 13-acetate was dissolved in 0.5 ml tetrahydrofuran. To this was added 0.15 ml saturated aqueous ammonium chloride, 0.15 ml 37% formalin solution, and 50 mg zinc dust (less than 325 mesh).

-91-

The reaction was capped and shaken vigorously for 24 hours. At the end of this time the reaction was partitioned between pH8 phosphate buffer and ethyl acetate. The ethyl acetate phase was reduced to a volume of 2 ml under a stream of nitrogen and 0.125 ml was applied to a silica gel TLC plate and developed with hexanes/ethyl acetate 46/54. The band at Rf 0.45 was scraped off and the product was eluted from the silica powder with acetone. Removal of the acetone yielded 3.0 mg of 20-homophorbol 12-myristate 13-acetate as a glassy solid for spectroscopic analysis and bioassay.

EXAMPLE 16

A stock solution of 300 pmoles of the standard inflammatory compound phorbol 12-myristate 13-acetate per 0.005 ml acetone was prepared. This solution was used to prepare four-fold dilutions of 20-deoxy-20-(2-hydroxyethylthio)-phorbol 12-myristate 13-acetate, prepared as in Example 3, covering concentrations of the latter ranging from 4 to 64,000 pmoles per 0.005 ml. These solutions were used to demonstrate the anti-inflammatory activity of the latter compound by application of 0.005 ml to the insides of the right ears of mice, followed by the observation of ear inflammation/erythema during a 1-48 hour period. Inhibition of the phorbol 12-myristate 13-acetate induced inflammation was observed at the medium and higher concentrations of the inhibitor.

In a like manner, the anti-inflammatory activities of the following other compounds were demonstrated:

phorbol 12-myristate
phorbol 12,13-diacetate
20-deoxy-20-chlorophorbol 12-myristate 13-acetate
20-deoxy-20-(2-hydroxyethylthio)phorbol
12,13-dibutyrate
20-deoxy-20-(2-hydroxyethylthio)phorbol 12-myristate
13-acetate
20-deoxy-20-[(2-hydroxyethyl)methylamino]phorbol
12-myristate 13-acetate
20-deoxy-20-fluorophorbol 12,13-bis(2,4-difluorophenylacetate
phenylacetate
12-beta,13-bis(2,4-difluorophenylacetoxy)-4,9-dihydroxy-1,6(20),7-tigliatrien-3-one
12-beta,13-bis(2,4-difluorophenylacetoxy)-4,9-dihydroxy-7-fluoro-1,6(20)-tigliadien-3-one
20-deoxy-20-hydroxymethylphorbol 12-myristate 13
acetate
phorbol 12-myristate 13-acetate 20-methylcarbamate

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

1. A compound of the formula:




wherein P represents a radical, formally derived from a parent compound, which compound:

- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
- b. activates any form of the enzyme protein kinase C; and
- c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom; and

wherein G is any group either i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or ii) singly or doubly bonded to a carbon atom immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound; and wherein the hydroxymethyl or 1-hydroxyethyl group of the parent compound is absent or has been replaced by G.

2. A compound of the formula:



wherein P_o represents a radical, formally derived from a parent compound, which compound:

- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
- b. activates any form of the enzyme protein kinase C; and
- c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom; and wherein S_o-E_o represents a moiety which is either i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or ii) singly or doubly bonded to a carbon atom immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound; and


wherein S_o is a saturated or unsaturated, straight or branched chain of atoms which separates P_o and E_o by a linear count of at least two but not more than 6 atoms and contains one to six heteroatoms selected from oxygen, nitrogen and sulfur, a phosphorus atom, or up to 12 halogen atoms; provided that the total number of atoms does not exceed about 35; and

wherein E_o comprises a saturated or singly or multiply unsaturated group containing up to 8 carbon atoms and optionally containing 1 to 12 halogen atoms, 1 to 4 heteroatoms selected independently from oxygen, nitrogen and sulfur and one phosphorus atom, or wherein S_oE_o taken

together may be a hydrogen, halogen, or ketonic oxygen atom or a hydroxy, amine, or thiol group singly or doubly bonded to the carbon atom of the parent compound P₀ in place of the hydroxymethyl or 1-hydroxyethyl group.


3. A compound of Claim 2 wherein P₀ is derived from a diterpenoid activator of protein kinase C.
4. A compound of Claim 2 wherein P₀ is derived from an aromatic heterocyclic activator of protein kinase C of the indole, indene, benzofuran, or benzothiophene class.
5. A compound of Claim 2 wherein P₀ is derived from a activator of protein kinase C of the diaminobenzyl alcohol class.
6. A compound of Claim 2 wherein P₀ is derived from an activator of protein kinase C of the diacylglycerol or diacyloxybutanol class.
7. A compound of Claim 2 wherein P₀ is derived from a polyacetate-derived activator of protein kinase C.

8. A compound of Claim 2 wherein P_0 is derived from a protein kinase C activator of the bryostatin class.
9. A compound of the formula:



wherein P_x is selected from the group consisting of $P_1, P_2, P_3, P_4, P_5,$ and P_6 ;

wherein P_1 is a radical of the formula:



in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof;

wherein

A₁ and A₂ may be individually selected from hydrogen and a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated and/or aromatic carbon- and/or heteroatom-containing substituent having not more than 34 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen and sulfur; or,

A¹ and A² taken together complete a 5- or 6-membered carbocyclic or heterocyclic ring, optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatomcontaining groups, which groups taken together contain a total of not more than 30 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur;

A³ is a three atom chain which completes a 7-membered carbocyclic ring optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups, which groups taken

together, excluding SoEo, contain not more than 7 carbon atoms, not more than 4 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen and sulfur; provided that, including $S_{O}E_{O}$, the middle carbon atom of A^3 is not substituted by hydroxymethyl, hydroxyethyl or 2-hydroxy-2propyl;

A^4 completes a 6- or 7-membered carbocyclic or heterocyclic ring (connected in the beta configuration to either carbon atom 9 or 10), optionally substituted by one or more straight chain or branched chain cyclic or acyclic, saturated, unsaturated and/or aromatic carbon- and/or heteroatom-containing groups, the group or groups optionally completing one or two additional rings by themselves and/or one to five additional rings when taken together with $A1$, $A2$, a ring formed by $A1$ and $A2$ together, and/or a bond to carbon atom 9, which groups taken together, include not more than 40 carbon atoms, not more than 24 halogen atoms, and not more than 10 heteroatoms selected from oxygen, nitrogen, and sulfur;


J^1 is selected from hydrogen, fluoro, chloro, hydroxy, amino, mono- or d-loweralkylamino, methyl, ethyl, vinyl, ethynyl, propargyl, cyano, methoxy, ethoxy, trifluoromethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, acetoxy, propanoyloxy, acetyl,

propanoyl, hydroxyacetyl, 2-hydroxypropanoyloxy, 3-hydroxypropanoyl, acetamido, propanamido, hydroxyacetamido, 2-hydroxypropanamido, or 3-hydroxypropanamido, (each of which must be situated in the beta position), or J^1 taken together with A1, A2, or a ring formed by A1 together with A2 completes a 3-to7-membered substituted or unsubstituted carbocyclic or heterocyclic ring;

J_2 is selected from hydrogen, methyl, ethyl, hydroxymethyl, hydroxyethyl, vinyl, ethynyl, allyl, propargyl, n-propyl and iso-propyl;

provided that if P_1 is a phorbol 12,13-dialkanoyl moiety then S_0E_0 may not be carboxy, carbaldehyde, carbaldehyde-2,4-dinitrophenylhydrazone, or alkanoyloxy-methylene, and provided that if P_1 is a phorbol 12,13-di-O-acetyl moiety then S_0E_0 may not be cyano or aldoxime;

wherein P_2 is a radical of the formula:



-100-

in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof,

wherein

B¹ completes a 6-membered carbocyclic or heterocyclic aromatic ring, optionally substituted by a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups which, taken together, contain not more than 40 carbon atoms, not more than 24 halogen atoms, and not more than 9 heteroatoms selected from oxygen, nitrogen, and sulfur, the groups being optionally connected to form 1-3 additional rings;

B² is selected from oxygen, sulfur, sulfoxide, sulfone, and a carbon or nitrogen atom optionally substituted by a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups having not more than 15 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, and B² may be linked to B¹ to form an additional carbocyclic or heterocyclic ring;

B³ is a 2-carbon chain optionally substituted by one or more straight chain or branched

chain, cyclic or acyclic, saturated, unsaturated, and/or carbon- and/or heteroatom-containing groups, which groups taken together but excluding SoEo, contain not more than 8 carbon atoms, not more than 6 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur; provided that, including $S_{O}E_{O}$, the carbon atom of B^3 bonded to K^2 as defined below does not carry $---CH_2OH$, $---CHCH_3OH$ or $---C(CH_3)_2)OH$;

K^1 is hydrogen or is a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group containing not more than 15 carbon atoms, not more than 18 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, and K^1 may be linked to B^2 or B^3 or to both B^2 and B^3 to form one or more additional carbocyclic and/or heterocyclic rings;

K^2 is selected from oxygen, sulfur $--NK^6--$ or $---CK^6K^7--$ wherein K^6 is hydrogen, hydroxy, methyl, ethyl, fluoro, n-propyl, allyl, or propargyl, and K^7 is hydrogen, methyl, ethyl, halogen, trifluoromethyl or cyano;

K^3 and K^4 may be the same or may differ and may each be hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated,

-102-

unsaturated, and/or aromatic carbon- and/or heteroatomcontaining group, such that K^3 and K^4 taken together contain not more than 18 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen and sulfur;

K^5 is selected from oxygen, sulfur, sulfoxide, sulfone or $--NK^8--$, $--NOK^8--$ or $----CK^8K^9----$ wherein K^8 is hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatomcontaining group containing not more than 30 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur, and K^9 is hydrogen, methyl, ethyl, n-propyl, hydroxy, halogen, allyl, propargyl, cyano, or trifluoromethyl;

provided that if B^1 completes an unsubstituted carbocyclic ring, and

B^2 is $---NH---$, and

K^1 is hydrogen, and

K^2 is $---NH---$, and

K^3 is $---CH(CH_3)_2$ in the alpha position, and

-104-

in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof,

wherein

L^1 , L^2 and L^3 are individually selected from nitrogen or substituted or unsubstituted carbon;

D^1 and D^2 each may be a bond or a substituted or unsubstituted carbon atom; D^1 may be linked to L^1 , L^2 or both L^1 and L^2 to form additional fused carbocyclic and/or heterocyclic substituted or unsubstituted rings; and D^2 may be linked to L^2 , L^3 or both L^2 and L^3 to form additional fused carbocyclic and/or heterocyclic substituted or unsubstituted rings;


D_3 and D_4 each are heteroatom-containing functional groups, the heteroatoms being selected from oxygen, nitrogen, and sulfur; D_3 may be linked to L^1 , L^2 or both L^1 and L^2 to form additional fused substituted or unsubstituted carbocyclic and/or heterocyclic rings; and D_4 may be linked to L^2 , L^3 or both L^2 and L^3 to form additional fused substituted or unsubstituted carbocyclic and/or heterocyclic rings;

provided that D^1 and D_3 taken together and D^2 and D_4 taken together both embody at least one

oxygen, nitrogen, or sulfur atom separated from the aromatic nucleus by zero or one intervening carbon atom; and

L4 and L5 are straight chain or branched, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups which, taken together, contains about 2-40 carbon atoms, not more than 24 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur; provided that S_0E_0 may not be hydroxymethyl or 1-hydroxyethyl;

wherein P_4 is a radical of the formula:



in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof,

wherein

V^1 is a bond or is a carbon atom carrying substituents individually selected from hydrogen, methyl, and halogen;

V^2 and V^3 are individually selected from oxygen, sulfur, sulfoxide, and $---NV^4---$ in which V^4 is hydrogen or a hydrocarbon radical containing not more than 30 carbon atoms;

F^1 and F^2 are a straight chain or branched, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups which, taken together, contain 10-40 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur;

F^3 is hydrogen or a substituent selected from methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, cyano, vinyl, ethynyl, allyl, and propargyl;


F^4 and F^5 each may be hydrogen or may be hydrocarbon or halogenated hydrocarbon radicals which, taken together, contain not more than 40

-107-

carbon atoms and not more than 24 halogen atoms;

provided that V^4 , F^1 , F^2 , F^4 and F^5 taken together contain not more than 50 carbon atoms and provided that SoEo may not be hydrogen, methyl, chloromethyl, hydroxymethyl, mercaptomethyl, unsubstituted carboxamido, 1-hydroxyethyl or alkanoyloxymethylene;

wherein P_5 is a radical of the formula:



in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof,

wherein


U^1 and U^2 , independently, are selected from hydrogen, azide, halogen, hydroxy, C_{1-7} alkoxy, C_{1-7} alkenoxy, C_{1-7} alkynoxy, thio, C_{1-7} alkanoyl, C_{1-7} saturated or unsaturated alkyl, and cyano; or U^1 and U^2 taken together may be an oxygen atom forming an epoxy group or may be an additional bond forming an unsaturated linkage;

U^3 is selected from hydrogen, halogen, C_{1-12} alkyl, C_1-C_{12} alkenyl, C_1-C_{12} alkynyl, C_1-C_{12} alkoxy, C_1-C_{12} alkenoxy, C_1-C_{12} alkynoxy, and aryl or C_7-C_{12} aralkyl wherein the aryl group may be substituted by nitro, halogen, cyano, and/or diloweralkylamino groups;

U^4-U^6 , independently, are selected from hydrogen, halogen, cyano, nitro, amino, diloweralkylamino, C_{1-7} saturated or unsaturated alkyl, hydroxy, C_{1-7} saturated or unsaturated alkoxy, C_{1-7} carboalkoxy, C_{1-7} alkanoyloxy, and azide; and

U^7 is selected from hydrogen, C_{1-7} saturated or unsaturated alkyl, and C_{1-7} saturated or unsaturated alkanoyl; provided that if $S_O E_O$ is hydroxymethyl, 1-hydroxyethyl or acetoxy-methylene, then $S_O E_O$ may not be bonded to C29;

and wherein P_6 is a radical of the formula:



in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof,

wherein

W^1 is selected from hydrogen, halogen, hydroxy, C_{1-5} alkoxy, C_{1-5} alkanoyloxy and cyano;

W^2 is selected from oxo, hydrogen, hydroxy, cyano, C_{1-7} alkyl, C_{1-7} alkoxy, C_{1-7} alkanoyloxy, and halogen;


W^3 - W^5 each may be hydrogen or a straight chain

-110-

or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group, containing not more than 30 carbon atoms, not more than 24 halogen atoms, and not more than 8 heteroatoms selected from oxygen, nitrogen, and sulfur, and W^4 and W^5 taken together may form an additional carbocyclic or heterocyclic ring; and

W^6 is hydrogen or a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing group, containing not more than 15 carbon atoms, not more than 12 halogen atoms and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, and W^6 , taken together with W^4 and W^5 may complete an additional carbocyclic or heterocyclic ring; provided that if S_{O_2} is hydroxymethyl or hydroxyethyl then S_{O_2} may not be bonded to C25;

wherein S_x is selected from the group consisting of S_1 , S_2 , S_3 , S_4 , S_5 and S_6 ; wherein S_1 is a single or a double bond or is a chain of atoms of the formula:



SUBSTITUTE SHEET

-111-

wherein

a, b, d, e, and g may independently be from 0 to 3;

c and f may independently be 0 or 1;

the sum of (a+b+c+d+e+f+g) is at least 1 but not more than 6; and

if c and f are both 1, then the sum of (d+e) must be at least 1;

R_1^1 through R_1^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms are situated in functional groups selected from hydroxy, amino, hydroxylamine, tertiary amine oxide, Schiff's base, hydrazine, thiol, nitro, nitroso, oxime, azide, ether, acetal, ketal, thioether, aldehyde, keto, hydrazone, carboxy, ester, amide, cyano, hydrazide, carbonate, carbamate, urea, isourea, carboximidine, imidate, guanidine, thioester, thioamide, thiocarbonate, dithiocarbonate, thiocarbamate, dithiocarbamate,

-112-

thiourea, isothiourea, thioimidate, nitroguanidine, cyanoguanidine, and xanthate;

R_1^1 or R_1^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety;

one of the substituents $R_1^1-R_1^{12}$ may be G^1 , as defined below, provided that the substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom;

R_1^{11} or R_1^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage;


one of the substituents $R_1^1-R_1^{12}$ may be linked to either the atom in P_x that carries the S_1 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different ring hetero members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members, by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy,

ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

X and X' are defined below;

provided that, for all substituents R₁¹ through R₁¹² and all constituents X and X', taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 4 or less; the total of nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less; the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less; the total of --OH, --SH, and --NH₂ groups together is 4 or less;

S₂ is a chain of atoms of the formula:



wherein

h, i, k, m, p, and q may be independently be from 0 to 3;

j and n may independently be 0 or 1;

if j and n are both 1 and l is 0, then the sum of (k + m) must be at least 1;

if n is 1 and o is 0, then the sum of (p + q) must be at least 1;

the sum of (l + o) is 1-3; and

the sum of (h+i+j+k+2l+m+n+2o+p+q) is at least 1 but not more than 7;

R_2^1 through R_2^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, hydroxylamine, tertiary amine oxide, Schiff's base, hydrazine, thiol, nitro, nitroso, oxime, azide, ether, acetal, ketal, thioether, aldehyde, keto, hydrazone, carboxy, ester, amide,

-115-

cyano, hydrazide, carbonate, carbamate, urea, isourea, carboxamidine, imidate, guanidine, thioester, thioamide, thiocarbonate, dithiocarbonate, thiocarbamate, dithiocarbamate, thiourea, isothiourea, thioimidate, nitroguanidine, cyanoguanidine, and xanthate;

R_2^1 or R_2^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety;

one of the substituents $R_2^1-R_2^{12}$ may be G^1 , as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom;

R_2^{11} or R_2^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage;

one of the substituents $R_2^1-R_2^{12}$ may be linked to either the atom in P_x that carries the S_2 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring

-116-

optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

X and X' and Y and Y' are as defined below;

provided that, for all substituents R₂¹ through R₂¹² and all substituents embodied in X, X', Y, and Y' taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 4 or less; the total of nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less; the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less;

-118-

the sum of $(r + s + 2t + u + 2v + 3w + 4x + y + 2z + a' + b')$ is at least 1 but not more than 7;

R_3^1 through R_3^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent, containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate;

R_3^1 or R_3^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety;

one of the substituents R_3^1 - R_3^{12} may optionally be G^1 , as defined below, provided that said substituent is bonded to a carbon atom

that is separated from the diterpenoid moiety by either zero or one intervening atom;

R_3^{11} or R_3^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage;

one of the substituents R_3^1 - R_3^{12} may be linked to either the atom in P_x that carries the S_3 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxypropyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

Y, Y', Z^1 , Z^2 , and Z^3 are as defined below;

wherein

c', d', e', h', and i' may independently be from 0 to 3;

f' and g' may independently be 0 or 1; and

the sum of (c' + d' + e' + f' + g' + h' + i') is at least 1 but not more than 5;

R_4^1 through R_4^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carbox-amidine, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate;

R_4^1 or R_4^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety;

-122-

one of the substituents $R_4^1-R_4^{12}$ may optionally be G^1 , as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom;

R_4^{11} or R_4^{12} may optionally be an additional bond to E_1 , thereby completing an unsaturated linkage;

one of the substituents $R_4^1-R_4^{12}$ may be linked to either the atom in P_x that carries that S_4 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido,

-124-

wherein

j' , k' , m' , q' , and s' may independently be from 0 to 3;

l' or r' may each be 0 or 1, but the sum of ($l' + r'$) must be 1 or 2;

n' and p' may each be 0 or 1, but the sum of ($n' + p'$) must be 0 or 1;

the value of o' may be 0-2;

if the sum of ($n' + p'$) is 1 and l' is 0, then q' must be at least 1;

if the sum of ($n' + p'$) is 1 and r' is 0, then m' must be at least 1; and

the sum of ($j' + k' + l' + m' + n' + o' + p' + q' + r' + s'$) is at least 1 but not more than 5;

R_5^1 through R_5^{12} may be the same or different and each may be either hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, and sulfur atoms must be situated in functional

-125-

groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamide, guanidine, thioester, thioamide, thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate;

R_5^1 or R_5^2 may optionally comprise an additional bond completing an unsaturated linkage the diterpenoid moiety;

one of the substituents $R_5^1-R_5^{12}$ may optionally represent G^1 , as defined below, provided that said substituent is bonded to a carbon atom that is separated from the diterpenoid moiety by either zero or one intervening atom;

R_5^{11} or R_5^{15} may optionally comprise an additional bond to E_1 , thereby completing an unsaturated linkage;

one of the substituents $R_5^1-R_5^{12}$ may be linked to either the atom in P_x that carries the S_5 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero

-126-


ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from the halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkyl-amino containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

Q, Q', X, X', and Y are as defined below;

provided that, for all substituent R₅¹ through R₅¹² and all constituents Q, Q', X,

X', and Y taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 10 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 4 or less; the total of nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less; the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less; the total of --SH groups is 2 or less; the total of --OH, --SH, and --NH₂ groups together is 4 or less;

S₆ is a chain of atoms defined by:



wherein

t', u', v', w', x', y', z' and a'' may each be 0 or 1;

the sum of (t' + u' + v' + 2w' + x¹ + 2y' + z' + a'') must be 0-2;

b'', d'', e'', f'', h'', j'' and k'' may each independently be 0 or 1;

c'', g'', i'', and l'' may each independently be 0-3;

-128-

if d'' and j'' are both 1, then the sum of ($g'' + i''$) must be at least 1;

if either j'' or k'' is 1, then l'' must be at least 1;

if b'' is 1, then the sum of ($c'' + g'' + h'' + i'' + l''$) must be at least 1;

if d'' is 1, then the sum of ($g'' + h'' + i'' + l''$) must be at least 1; and

the sum of ($t' + u' + v' + 2w' + x' + 2y' + z' + a'' + b'' + c'' + d'' + e'' + 2f'' + g'' + 2h'' + i'' + j'' + k'' + l''$) must be 0-5;

R_6^1 through R_6^{12} may be the same or different and each may be either hydrogen or a saturated or or multiply unsaturated, straight or branched, acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen, nitrogen, and sulfur atoms must be situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbonate, carbamate, urea, isourea, carboxamidine, guanidine, thioester, thioamide,

thiocarbonate, thiocarbamate, thiourea, nitroguanidine, cyanoguanidine, and xanthate;

R_6^1 or R_6^2 may optionally be an additional bond completing an unsaturated linkage to the diterpenoid moiety;

R_6^3 or R_6^4 may optionally comprise an additional bond to G^2 as defined below, thus completing an unsaturated linkage;

R_6^5 or R_6^6 may optionally comprise an additional bond to G^2 , thus completing an unsaturated linkage if a'' is 0;

one of the substituents R_6^1 - R_6^4 may optionally be G^1 , as defined below;

R_6^{11} or R_6^{12} may optionally comprise an additional bond to E_1 , thereby completing an unsaturated linkage;

one of the substituents R_6^1 - R_6^4 may be linked to either the atom in P_x that carries the S_{65} chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-6 membered ring optionally

-130-


containing 1-3 hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on any carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

Q, Q', Q'', Q''', X, X', X'', Y, Y', and Z⁴, are as defined below;

provided that, for all substituents R₆¹ through R₆¹² and all constituents Q, Q', Q'', Q''', X, X', X'', Y, Y', Z⁴, and G² taken together, but excluding any atoms of the diterpenoid moiety: the total of carbon atoms is 12 or less; the total of halogen atoms is 12 or less; the total of oxygen atoms is 5 or less; the total of nitrogen atoms is 4 or less; the total of sulfur atoms is 3 or less, the total of oxygen, nitrogen, and sulfur atoms together is 8 or less; the total of --OH groups is 3 or less; the total of --NH₂ groups is 2 or less;

the total of --SH groups is 2 or less; the total of --OH, --SH, and --NH₂ groups together is 4 or less;

X, X', X'' may be the same or different and are selected from:



wherein

R_X^1 and R_X^2 may independently be hydrogen;

R_X^1 through R_X^5 may be the same or different and each may be a saturated or singly or multiple unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and

sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboxamide, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine;




R_X^1 may optionally represent an additional bond to the diterpenoid moiety, thus completing an unsaturated linkage;

one of the substituents $R_X^1-R_X^5$ may optionally represent G^1 , as defined below, provided that the substituent is bonded to an atom that is separated from the parent P_X by either zero or one intervening atom; and

one of the substituents $R_X^1-R_X^5$ may be linked to either the atom in P_X that carries the chain containing X, X', and/or X'' or to an atom in P_X adjacent thereto, to form a saturated, unsaturated, or aromatic, heterocyclic 4-8 membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH,

the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , SOCH_3 , SCF_3 , COOH, COOCH_3 , COCH_3 , CH=O , acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methyl-acetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxy-acetyl and hydroxyacetoxy;

Y and Y' may be the same or different and are selected from:



wherein

R_Y^1 and R_Y^2 (cis or trans) may be the same or different and each may be hydrogen or may comprise a saturated or singly or multiply unsaturated, straight or branched acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboxamidine, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine;

R_Y^3 and R_Y^4 (cis or trans) may be the same or different and each may be hydrogen or may be a saturated or singly or multiply unsaturated, straight or branched acyclic substituent containing not more than 6 carbon atoms, not more than 9 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated

-135-

in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboximidine, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine, with the exception that neither R_Y^3 nor R_Y^4 may be a halogen atom alone;






one of the substituents $R_Y^1-R_Y^4$ may optionally represent G^1 , as defined below, provided that the substituent is bonded to an atom that is separated from the diterpenoid moiety by either zero or one intervening atom; and

one of the substituents $R_Y^1-R_Y^4$ may be linked to either the atom in P_x that carries the chain containing Y and/or Y' or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$,

-136-

SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy.

z¹ is selected from



wherein

R_z¹ is hydrogen, a saturated or unsaturated substituent selected from C₁₋₆ alkyl, C₁₋₆ halogenated alkyl, C₁₋₆ acyl, C₁₋₆ halogenated acyl, C₁₋₆ monohydroxyacyl, and C₂₋₆ hydroxyalkyl, cyclohexyl, or phenyl or benzyl

-137-

optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or halogen, or, R_Z^1 may comprise an additional bond to the parent P_x , thus completing an unsaturated linkage;

R_Z^2 is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{1-6} acyl, C_{1-6} halogenated acyl, C_{1-6} monohydroxyacyl, and C_{2-6} hydroxyalkyl, cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or halogen;

R_Z^3 independently may have the values specified above for R_Z^2 ;

R_Z^4 independently may have the values specified above for R_Z^1 ;

R_Z^5 may be hydrogen or a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} hydroxyalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkoxy, C_{1-6} halogenated alkyl, or phenyl or benzyl optionally substituted by methyl, ethyl, methoxy, nitro, cyano, trifluoromethyl, and/or halogen;


-138-

R_Z^6 independently may have the values specified above for R_Z^5 ;

one of the substituents $R_Z^1-R_Z^6$ may be linked to either the atom in P_x that carries the chain containing Z^1 or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

provided that only one of the substituents $R_Z^1-R_Z^6$ may be substituted or unsubstituted phenyl or benzyl.

Z^2 is selected from



wherein

R_2^7 independently may have the values specified

for R_2^1 ;

R_2^8 is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{2-6} hydroxyalkyl in which the hydroxy group may be esterified to the acetyl or propionyl ester, and cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or halogen;

R_2^9 independently may have the values specified

-140-

above for R_Z^8 ;

R_Z^{10} independently may have the values specified
above for R_Z^7 ;

R_Z^{11} may be hydrogen, a saturated or unsaturated
substituent selected from C_{1-6} alkyl, C_{1-6}
halogenated alkyl, and cyclohexyl, or phenyl or
benzyl optionally substituted by methyl, ethyl,
nitro, cyano, trifluoromethyl, and/or halogen;

R_Z^{12} and R_Z^{13} independently may have the values
specified above for R_Z^2 ;

R_Z^{14} - R_Z^{16} independently may have the values
specified above for R_Z^{11} ;



R_Z^{17} and R_Z^{20} individually may be hydrogen or a
substituent selected from C_{1-4} alkyl; C_{1-4}
alkoxy; C_{1-4} alkylthio, phenoxy or thiophenoxy
optionally substituted by methyl, halogen,
and/or nitro; or amino optionally mono- or
disubstituted by C_{1-4} alkyl or monosubstituted
by cyano, nitro, or phenyl optionally sub-
stituted by halogen and/or nitro;

-141-

R_Z^{18} and R_Z^{19} individually may be --O--, --S--, or --NR_Z²¹, wherein R_Z^{21} may be hydrogen, C₁₋₄ alkyl, 2-hydroxyethyl, 2-hydroxy-n-propyl, 2-acetoxyethyl, or 2-acetoxy-n-propyl;

one of the substituents R_Z^7 - R_Z^{17} , R_Z^{20} or R_Z^{21} optionally may be linked to either the atom in P_x that carries the chain containing Z² to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive hydroxyacetyl and hydroxyacetoxy; provided that only one of the substituents R_Z^7 - R_Z^{21} may comprise or carry a substituted or unsubstituted phenyl or benzyl;

Z^3 is selected from



wherein

R_Z^{22} and R_Z^{25} independently may have the values specified above for R_Z^{11} ;

R_Z^{23} and R_Z^{24} independently may have the values specified above for R_Z^8 ;


R_Z^{22} or R_Z^{25} optionally may be linked to either the atom in P_x that carries the chain containing Z^2 or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8

-143-

membered ring optionally containing 1-3 other identical or different hetero ring members selected from O, S, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , SOCH_3 , SCF_3 , COOH, COOCH_3 , COCH_3 , CH=O , acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

provided that at most one of the substituents R_Z^{22} - R_Z^{25} may comprise or carry a substituted or unsubstituted phenyl or benzyl moiety;

Z^4 and $\text{Z}^{4'}$ independently may be:



wherein

R_Z^{26} may be hydrogen or may be a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, in which substituent the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto, carboxy, ester, amide, cyano, carbamate, urea, carboxamide, guanidine, thioester, thioamide, thiocarbamate, thiourea, nitroguanidine, and cyanoguanidine;




R_Z^{26} may optionally comprise one additional bond either to the parent P_x or to G^2 , thus completing an unsaturated linkage;

R_Z^{26} may be linked to either the atom in P_x that carries the S_6 chain or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-6 membered ring optionally containing 1-3 identical or different ring members selected from O, S, SO, SO_2 , CO, =N-, and NH, the ring

-145-

being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , SOCH_3 , SCF_3 , COOH , COOCH_3 , COCH_3 , CH=O , acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxy;

Z^5 and $\text{Z}^{5'}$ independently may be:



wherein


R_Z^{27} is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{1-6} acyl, C_{1-6} halogenated acyl, C_{1-6} monohydroxyacyl, and C_{2-6} hydroxyalkyl, cyclohexyl, or phenyl or benzyl optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or halogen, or R_Z^{27} may comprise an additional bond to the parent P_x or to G^2 , thus completing an unsaturated linkage;

R_Z^{28} is hydrogen, a saturated or unsaturated substituent selected from C_{1-6} alkyl, C_{1-6} halogenated alkyl, C_{1-6} acyl, C_{1-6} halogenated acyl, C_{1-6} monohydroxyacyl, and C_{2-6} hydroxyalkyl, cyclohexyl, or phenyl or benzyl either of which may optionally substituted by methyl, ethyl, nitro, cyano, trifluoromethyl, and/or halogen, or R_Z^{28} may comprise an additional bond to G^2 , thus completing an unsaturated linkage;

R_Z^{29} independently may have the values specified above for R_Z^{27} ; and

R_Z^{30} independently may have the values specified above for R_Z^1 .

M and M' independently may be:



wherein

R_M^1 and R_M^2 may be the same or different and each may be hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, in which the oxygen atoms total 3 or less, the nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, azide, ether, thioether, aldehyde, keto,

carboxy, ester, amide, cyano, nitroguanidine, and cyanoguanidine;



R_M^1 may optionally comprise an additional bond to the parent P_x group, thus completing an unsaturated linkage;

R_M^1 or R_M^2 may optionally represent G^1 , as defined below, provided that said substituent is bonded to an atom that is separated from the parent P_x by either zero or one intervening atoms;

R_M^1 or R_M^2 may be linked to either the atom in P_x that carries the chain containing M and/or M' or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 5-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy-2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $COCH_3$, $CH=O$, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive,

acetamido, N-methylacetamido, carboxamido,
N-alkylated carboxamido containing 1-4 carbon
atoms inclusive, hydroxyacetyl and hydroxy-
acetoxy;

Q-Q''' independently may be:



wherein

R_Q^1 and R_Q^2 may be the same or different and each may be hydrogen or a saturated or singly or multiply unsaturated, straight or branched, acyclic substituent containing 1-6 carbon atoms, not more than 8 halogen atoms, and not more than 4 heteroatoms selected from oxygen, nitrogen, and sulfur, such that for any substituent the oxygen atoms total 3 or less, the

-150-

nitrogen atoms total 2 or less, and the sulfur atoms total 1 or zero, the heteroatoms being situated in functional groups selected from hydroxy, amino, thiol, nitro, ether, thioether, carboxy, ester, amide, cyano, nitroguanidine, and cyanoguanidine;

R_Q^1 may optionally represent G^1 , as defined below, provided that R_Q^1 is bonded to an atom that is separated from the parent P_x by either zero or one intervening atom;

R_Q^1 may be linked to either the atom in P_x that carries the chain containing Q and/or Q' or to an atom in P_x adjacent thereto, to form a saturated, unsaturated, or aromatic, carbocyclic 5-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical or different substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxy2-propyl, CF_3 , OCF_3 , SH, SCH_3 , $SOCH_3$, SCF_3 , COOH, $COOCH_3$, $CH=O$, acetoxy, amino mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methylacetamido, carboxamido, N-alkylated carboxamido containing

-151-

1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxo;





G¹ comprises a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxy-2-propyl, CF₃, OCF₃, SH, SCH₃, SOCH₃, SCF₃, COOH, COOCH₃, COCH₃, CH=O, acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methyl acetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxyacetyl and hydroxyacetoxo;

G² completes a saturated, unsaturated, or aromatic, carbocyclic or heterocyclic 4-8 membered ring optionally containing 1-4 identical or different hetero ring members selected from O, S, SO, SO₂, CO, =N-, and NH, the ring being optionally substituted on its carbon and/or NH members by 1-4 identical substituents selected from halogen, hydroxy, methoxy, ethoxy, methyl, ethyl, cyano, azide, nitro, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl,

-152-

2-hydroxy-2-propyl, CF_3 , OCF_3 , SH , SCH_3 , SOCH_3 , SCF_3 , COOH , COOCH_3 , COCH_3 , CH=O , acetoxy, amino, mono- or dialkylamino totalling 1-4 carbon atoms inclusive, acetamido, N-methyl-acetamido, carboxamido, N-alkylated carboxamido containing 1-4 carbon atoms inclusive, hydroxy-acetyl and hydroxyacetoxy, and the ring may be connected directly by single or double bonds to adjacent atoms; and

wherein E_1 is selected from $=\text{O}$, $=\text{S}$, $=\text{NH}$, $=\text{NOH}$, $=\text{N--NH}_2$, hydrogen, halogen, $--\text{OH}$, $--\text{SH}$, $--\text{NH}_2$, $--\text{NH--NH}_2$, $--\text{N}_3$, $--\text{NO}$, $--\text{NO}_2$, $--\text{NHOH}$, $--\text{ONH}_2$, or is selected from :



wherein

T^1 is selected from $--\text{O}--$, $--\text{S}--$, and $--\text{NH}--$;

-153-

T^2 is selected from =O, =S, and =N-- R_E^6 , in which R_E^6 may be hydrogen, cyano, or nitro;

T^3 is selected from hydrogen, --OH, --NH₂, --SH, --N₃, --NH--NH₂, and --NH--OR_E⁷, in which R_E^7 may be hydrogen C₁₋₃, alkyl, or C₁₋₃ acyl;

R_E^1 is selected from hydrogen, halogen, hydroxy, nitro, nitroso, cyano, azide, --NH₂, --NH--OH, --SH, --O--NH₂, --NH--NH₂, --T¹--C=T²--T³, and --C=T²--T³;

R_E^2 and R_E^3 are individually selected from hydrogen, --C =T² -- T³, cyano, nitro, azide, and halogen; and

R_E^4 and R_E^5 are individually selected from hydrogen, halogen, cyano, nitro, --C=T²--T³, --T¹--C=T²--T³, and --CR_E¹R_E²R_E³.

10. A compound of Claim 9 wherein Po is P₁ and So is a single or double bond.
11. A compound of Claim 10 selected from the group consisting of:

-154-

12,20-dideoxy-20-chlorophorbol 13-angelate;
20-deoxy-20-bromophorbol 12-myristate 13-
acetate;
20-deoxy-20-iodophorbol 12-myristate 13-
acetate;
20-deoxy-20-chlorophorbol 12,13-bis(2,4-
difluorophenylacetate);
20-deoxy-20-fluorophorbol 12,13-bis(2,4-
difluorophenylacetate);
12-beta,13-bis(2,4-difluorophenylacetoxo)-
4,9-dihydroxy-1,6(20),7-tigliatrien-3-one;
12-beta,13-bis(2,4-difluorophenylacetoxo)-
4,9-dihydroxy-7-fluoro-1,6(20)-tigliadien-
3-one; and
20-deoxy-20-cyanophorbol 12-myristate 13-
acetate.

12. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_1 .
13. A compound of Claim 12 selected from the group consisting of:
20-deoxy-20-(2-hydroxyethylthio)phorbol 12-
butyrate;
20-deoxy-20-(2-hydroxyethylthio)phorbol 12,13-
dibutyrate;
20-deoxy-20-(2-hydroxyethylthio)phorbol 12-
myristate 13-acetate;
20-deoxy-20-[(2-hydroxyethyl)methyl-
amino]phorbol 12-myristate 13-acetate;
20-deoxy-20-(3-hydroxypropylthio)phorbol 12,13-
bis(2,4-difluorophenylacetate);

-155-

20-deoxy-20-aminophorbol 12-myristate 13-acetate and pharmaceutically acceptable salts thereof;

20-deoxy-20-N-methylaminophorbol 12-myristate 13-acetate and pharmaceutically acceptable salts thereof.

14. A compound of Claim 12 wherein S_1E_0 taken together comprise a saturated normal or branched C_2-C_{10} alkyl group containing one hydroxy moiety separated from P_1 by not more than four carbon atoms.
15. A compound of Claim 14 selected from the group consisting of:
 - 20-deoxy-20-hydroxymethylphorbol 12-myristate 13-acetate;
 - 20-deoxy-20-(1-hydroxyethyl)phorbol 12-myristate 13-acetate;
 - 20-deoxy-20-(2-hydroxyethyl)phorbol 12-myristate 13-acetate;
 - 20-deoxy-20-(2-hydroxyethyl)phorbol 12,13-bis(2,4-difluorophenylacetate)
 - 12,20-dideoxy-20-(2-hydroxyethylthio)phorbol 13-angelate.
16. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_2 .
17. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_3 .
18. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_4 .

-156-

19. A compound of Claim 18 wherein S_4E_0 taken together comprise an aminocarbonyloxymethylene group or a C_1-C_{10} saturated or unsaturated alkyl-, aralkyl- or arylaminocarbonyloxymethylene group.
20. A compound of Claim 19 selected from the group consisting of:
 - phorbol 12-myristate 13-acetate 20-methylcarbamate;
 - phorbol 12-myristate 13-acetate 20-ethylcarbamate;
 - phorbol 12-myristate 13-acetate 20-n-propylcarbamate; and
 - phorbol 12-myristate 13-acetate 20-n-butylcarbamate.
21. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_5 .
22. A compound of Claim 9 wherein P_0 is P_1 and S_0 is S_6 .
23. A compound of Claim 22 selected from:
 - i) phorbol 12-myristate 13-acetate 20-(2-hydroxymethylphenyl) ether; and
 - ii) 20-deoxy-20-(1-tetrazolyl)phorbol 12-myristate 13-acetate.
24. A compound of Claim 9 wherein P_0 is P_2 and S_0 is a single or double bond.

25. A compound of Claim 24 selected from the group consisting of:
- i) 1-(1-oxobutyl)-1,2,4,5,6,8-hexahydro-5-methyl-3H-pyrrolo(4,3,2-gh)-1,4-benzodiazonin-3-one; and
 - ii) 1,2,4,5,6,8-hexahydro-5-methyl-2-(1-methylethyl)-3H-pyrrolo(4,3,2-gh)-1,4-benzodiazonin-3-one.
26. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_1 .
27. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_2 .
28. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_3 .
29. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_4 .
30. A compound of Claim 29 wherein S_4E_0 taken together comprise an aminocarbonyloxymethylene group or a C_1-C_{10} saturated or unsaturated alkyl-, aralkyl-, or arylaminocarbonyloxymethylene group.
31. A compound of Claim 30 selected from the group consisting of:
- i) teleocidin 20-methylcarbamate;
 - ii) teleocidin 20-n-butylcarbamate;
 - iii) lyngbyatoxin A 20-methylcarbamate;
 - iv) lyngbyatoxin A 24-n-butylcarbamate.

32. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_5 .
33. A compound of Claim 9 wherein P_0 is P_2 and S_0 is S_6 .
34. A compound of Claim 9 wherein P_0 is P_3 and S_0 is a single or double bond.
35. A compound of Claim 9 wherein P_0 is P_3 and S_0 is S_1 .
36. A compound of Claim 9 wherein P_0 is P_3 and S_0 is S_2 .
37. A compound of Claim 9 wherein P_0 is P_3 and S_0 is S_3 .
38. A compound of Claim 9 wherein P_0 is P_3 and S_0 is S_4 .
39. A compound of Claim 38 selected from the group consisting of:
 - i) 4-(N-2,3-dihydroxypropylcarboxamido)-6-(N-decanoylamino) indole;
 - ii) 4-(N-2-mercaptoethylcarboxamido)-6-(N-decanoylamino) indole;
 - iii) 4-(S-2-aminoethylthiolcarboxy)-6-decanoylamino) indole.
40. A compound of Claim 9 wherein P_0 is P_3 and S_0 is S_5 .

41. A compound of Claim 9 wherein P_o is P_3 and S_o is S_6 .
42. A compound of Claim 41 selected from the group consisting of:
- i) R,S-4-(2-hydroxymethylpiperidinocarbonyl)-6-(N-decanoylamino)indole;
 - ii) R-4-(2-hydroxymethylpiperidinocarbonyl)-6-(N-decanoylamino)indole; and
 - iii) S-4-(2-hydroxymethylpiperidinocarbonyl)-6-(N-decanoylamino)indole.
43. A compound of Claim 9 wherein P_o is P_4 and S_o is a single or double bond.
44. A compound of Claim 9 wherein P_o is P_4 and S_o is S_1 .
45. A compound of Claim 44 selected from the group consisting of:
- i) 3-(2-hydroxyethylthio)-1,2-propylene bis-n-hexyl carbamate; and
 - ii) 3-(3-hydroxypropylthio)-1,2-propylene bis-n-hexyl carbamate.
46. A compound of Claim 9 wherein P_o is P_4 and S_o is S_2 .
47. A compound of Claim 9 wherein P_o is P_4 and S_o is S_3 .
48. A compound of Claim 9 wherein P_o is P_4 and S_o is S_4 .

-160-

49. A compound of Claim 48 wherein S_4E_0 taken together comprise an aminocarbonyloxymethylene group or a C_1-C_{10} saturated or unsaturated alkyl-, aralkyl-, or arylaminocarbonyloxymethylene group.
50. A compound of Claim 49 selected from the group consisting of:
- i) glycerol 1-oleate 2-acetate 3-methylcarbamate);
 - ii) glycerol 1-oleate 2-acetate 3-n-butylcarbamate;
 - iii) glycerol 1,2-dioctanoate 3-methylcarbamate; and
 - iv) glycerol 1,2-dioctanoate 3-n-butylcarbamate.
51. A compound of Claim 9 wherein P_0 is P_4 and S_0 is S_5 .
52. A compound of Claim 9 wherein P_0 is P_4 and S_0 is S_6 .
53. A compound of Claim 9 wherein P_0 is P_5 and S_0 is a single or double bond.
54. A compound of Claim 9 wherein P_0 is P_5 and S_0 is S_1 , S_2 or S_3 .

-161-

55. A compound of Claim 9 wherein P_o is P_5 and S_o is S_4 .
56. A compound of Claim 55 wherein S_4E_o taken together comprise an aminocarbonyloxymethylene group or a C_2-C_{10} saturated or unsaturated alkyl-, aralkyl-, or arylaminocarbonyloxymethylene group.
57. A compound of Claim 56 selected from the group consisting of:
- i) aplysiatoxin 20-methyl ether 30-methylcarbamate;
 - ii) aplysiatoxin 20-acetate 30-methylcarbamate; and
 - iii) aplysiatoxin 20,30-bis(methylcarbamate).
58. A compound of Claim 9 wherein P_o is P_5 and S_o is S_5 or S_6 .
59. A compound of Claim 9 wherein P_o is P_6 and S_o is a single or double bond.
60. A compound of Claim 9 wherein P_o is P_6 and S_o is S_1 , S_2 or S_3 .

-162-

61. A compound of Claim 9 wherein P_0 is P_6 and S_0 is S_4 .
62. A compound of Claim 61 wherein S_4E_0 taken together comprise an aminocarbonyloxymethylene group or a C_1-C_{10} saturated or unsaturated alkyl-, aralkyl-, or arylaminocarbonyloxymethylene group.
63. A compound of Claim 62 selected from the group consisting of:
 - i) the 26-methylcarbamate of bryostatin 1:
and
 - ii) the 26-n-butylcarbamate of bryostatin 1.
64. A compound of Claim 9 wherein P_0 is P_6 and S_0 is S_5 or S_6 .
65. An antagonist of protein kinase C, a nontoxic agonist of protein kinase C, an antagonist for toxic phorboids or a nontoxic, non-inflammatory phorboid-type agonist, excluding:
 - a) phorbol-12-n-alkanoylestere; and
 - b) diterpene, and diacylglycerol compounds having carboxyl, carbaldehyde, carbaldehyde-


-163-

2,4-dinitrophenyl-hydrazone, cyano, alkanoyloxymethylene, alkyoxymethylene chloromethyl or methyl in place of the hydroxymethyl group;

c) indole compounds having chloromethyl, methyl, alkanoyloxymethylene or alkoxy-methylene in place of the hydroxymethyl group.

66. A method of blocking inflammation in vivo, comprising administering an anti-inflammatory amount of an antagonist of protein kinase C, a nontoxic agonist of protein kinase C, an antagonist for toxic phorboids or a nontoxic, non-inflammatory phorboid-type agonist.

67. A compound of the formula:



-164-

in the form of an individual isomer, an isomer mixture, a racemate or optical antipode, or a pharmaceutically acceptable salt thereof;

wherein

A₁ and A₂ may be individually selected from hydrogen and a straight chain or branched chain, cyclic or acyclic, saturated, unsaturated and/or aromatic carbon- and/or heteroatom-containing substituent having not more than 34 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen and sulfur; or,

A¹ and A² taken together complete a 5- or 6-membered carbocyclic or heterocyclic ring, optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups, which groups taken together contain a total of not more than 30 carbon atoms, not more than 24 halogen atoms and not more than 6 heteroatoms selected from oxygen, nitrogen, and sulfur;

A³ is a three atom chain which completes a 7-membered carbocyclic ring optionally substituted by one or more straight chain or branched chain, cyclic or acyclic, saturated, unsaturated, and/or aromatic carbon- and/or heteroatom-containing groups, which groups taken

together, excluding SoEo, contain not more than 7 carbon atoms, not more than 4 halogen atoms, and not more than 5 heteroatoms selected from oxygen, nitrogen and sulfur; provided that, excluding S_oE_o, the middle carbon atom of A³ is not substituted by hydroxymethyl, hydroxyethyl or 2-hydroxy-2propyl;

A⁴ completes a 6-membered carbocyclic ring (connected in the beta configuration to carbon atom 9), optionally substituted by one or more straight chain or branched chain cyclic or acyclic, saturated, unsaturated and/or aromatic carbon- and/or heteroatom-containing groups, linked to A⁴ by single or double bonds, the group or groups optionally completing one or two additional rings by themselves and/or one or two additional rings by themselves and/or one to three additional rings when taken together with A1, A2, a ring formed by A1 and A2 together, and/or a bond to carbon atom 9, which groups taken together include not more than 40 carbon atoms, not more than 24 halogen atoms, and not more than 10 heteroatoms selected from oxygen, nitrogen, and sulfur; and wherein A⁴ carries at least one substituent other than hydrogen at C₁₂ and a hydroxy, amino, thiol, hydroxymethyl, mercaptomethyl, aminomethyl, 2-hydroxyethyl, carboxy, unsubstituted carboxamido, or unsubstituted aminocarbonyloxy group in the alpha configuration at C₁₃; J¹ is selected from hydrogen, fluoro, chloro, hydroxy, amino, mono- or d-loweralkylamino,

methyl, ethyl, vinyl, ethynyl, propargyl, cyano, methoxy, ethoxy, trifluoromethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, acetoxy, propanoyloxy, acetyl, propanoyl, hydroxyacetyl, 2-hydroxypropanoyloxy, 3-hydroxypropanoyl, acetamido, propanamido, hydroxyacetamido, 2-hydroxypropanamido, or 3-hydroxypropanamido, (each of which must be situated in the beta position), or J^1 taken together with A1, A2, or a ring formed by A1 together with A2 completes a 3- to 7-membered substituted or unsubstituted carbocyclic or heterocyclic ring;

J2 is selected from hydrogen, methyl, ethyl, hydroxymethyl, hydroxyethyl, vinyl, ethynyl, allyl, propargyl, n-propyl and iso-propyl;

$S_{O}E_{O}$ taken together comprise hydroxymethyl, 1-hydroxyethyl or 2-hydroxy-2-propyl; provided that if said compound is a phorbol 12-ester, it is not an phorbol 12-n-alkanoyl ester.

68. A compound of Claim 67 wherein the middle carbon atom of A^3 is substituted by $S_{O}E_{O}$, provided that if said compound is a phorbol 12-ester, it is not a phorbol 12-n-alkanyoyl ester.

69. A compound of Claim 68 which is a phorbol 12-ester, provided that said ester group is not an n-alkanoyl ester.
70. A compound of Claim 69 selected from the group consisting of:
- i) phorbol 12-[4-(9,10-dihydrophenanthrene-2)-butyrate];
 - ii) phorbol 12-(2,4-difluorophenylacetate);
 - iii) phorbol 12-[3,5-bis(trifluoromethyl)benzoate];
 - iv) phorbol 12-[3,5-bis(trifluoromethyl)-phenylacetate];
 - v) phorbol 12-(4-n-hexylbenzoate);
 - vi) phorbol 12-(3,5-dimethoxyphenylacetate);
and
 - vii) phorbol 12-(4-phenylbenzoate).
71. A method of producing a phorboid antagonist, a non-inflammatory agonist, ligand selective from one or more phorboid receptors or protein kinase C subtypes, comprising the steps of: replacing a hydroxymethyl or 1-hydroxyethyl group of a parent compound, which compound:
- a. binds reversibly or irreversibly to a diacylglycerol-type receptor; and/or
 - b. activates any form of the enzyme protein kinase C; and

-168-

c. contains an hydroxymethyl or 1-hydroxyethyl group bonded to a carbon atom;

with a group S_o-E_o , which is either i) singly or doubly bonded to the carbon atom of the parent compound in place of the hydroxymethyl or 1-hydroxyethyl group; or ii) singly or doubly bonded to a carbon atom immediately adjacent to the carbon atom to which the hydroxymethyl or 1-hydroxyethyl group is bound in the parent compound; and

wherein S_o is a saturated or unsaturated, straight or branched chain of atoms which separates P_o and E_o by a linear count of at least two but not more than 6 atoms and contains one or six heteroatoms selected from oxygen, nitrogen and sulfur, a phosphorus atom, or up to 12 halogen atoms; provided that the total number of atoms does not exceed about 35; and

wherein E_o comprises a saturated or singly or multiply unsaturated group containing up to 8 carbon atoms and optionally containing 1 to 12 halogen atoms, 1 to 4 heteroatoms selected independently from oxygen, nitrogen and sulfur and one phosphorus atom, or wherein S_oE_o taken together may be a hydrogen, halogen, or ketonic oxygen atom or a hydroxy, amine, or thiol group singly or doubly bonded to the carbon atom of

the parent compound P₀ in place of the hydroxymethyl or 1-hydroxyethyl group.

1
7
1