

(19) **DANMARK**

(10) **DK/EP 2987803 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 14/47 (2006.01)** **A 61 K 38/17 (2006.01)** **A 61 P 1/00 (2006.01)**
A 61 P 19/00 (2006.01) **A 61 P 29/00 (2006.01)** **A 61 P 37/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-12-17**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-08-29**
- (86) Europæisk ansøgning nr.: **15181052.0**
- (86) Europæisk indleveringsdag: **2010-06-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-02-24**
- (30) Prioritet: **2009-06-17 US 268961 P**
- (62) Stamansøgningsnr: **10790213.2**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Promedior Inc., 101 Hartwell Avenue, Lexington, MA 02421-3125, USA**
- (72) Opfinder: **WILLETT, W., Scott, 3820 Comley Circle, Doylestown, PA 18901, USA**
- (74) Fuldmægtig i Danmark: **Budde Schou A/S, Hausergade 3, 1128 København K, Danmark**
- (54) Benævnelse: **SAP-Varianter og disses anvendelse**
- (56) Fremdragne publikationer:
WO-A2-2009/009019
WO-A2-2009/009034
**Anonymous: "UNIPROT:O19063 Complementary DNA sequence of porcine serum amyloid P component (SAP).", , 1 January 1998 (1998-01-01), XP055227351, Retrieved from the Internet:
URL:http://ibis.internal.epo.org/exam/dbfe_tch.jsp?id=UNIPROT:O19063 [retrieved on 2015-11-10]**
**Anonymous: "UNIPROT:P49255 Structure, expression, and evolution of guinea pig serum amyloid P component and C-reactive protein.", , 1 February 1996 (1996-02-01), XP055227368, Retrieved from the Internet:
URL:http://ibis.internal.epo.org/exam/dbfe_tch.jsp?id=UNIPROT:P49255 [retrieved on 2015-11-10]**

DESCRIPTION

BACKGROUND OF THE INVENTION

[0001] Serum Amyloid P (SAP) is a member of the pentraxin family of proteins. SAP is secreted by the liver and circulates in the blood as a stable pentamer. Previous research demonstrates SAP has an important role in both the initiation and resolution phases of the immune response. SAP can bind to sugar residues on the surface of bacteria and thereby promote their opsonization and engulfment by antigen-presenting cells. SAP also binds to free DNA and chromatin generated by apoptotic cells at the resolution of an immune response, thus preventing a secondary inflammatory response against these antigens. Molecules bound by SAP are removed from extracellular areas due to the ability of SAP to bind to all three classical Fcγ receptors (FcγR), having a particular affinity for FcγRII (CD32) and FcγRIII (CD16). After receptor binding, SAP and any attached complex are generally internalized and processed by the cell.

[0002] Recently, it has been suggested that SAP can be used as a therapeutic agent to treat various disorders, including fibrosis-related disorders, hypersensitivity disorders, autoimmune disorders, mucositis, and inflammatory disorders such as those cause by microbial infection. See, for example, U.S. Patent Application Nos. 11/707,333, 12/217,617 12/720,845, and 12/720,847. Protein therapeutics for treating human disease have revolutionized the health care industry. However, there are many difficulties in producing a protein therapeutic having the necessary potency and/or in sufficient quantity to be useful as a therapeutic agent. Many potential therapeutic agents are modified to increase their biological activity, such as plasma half-life, relative to the naturally-derived protein. Recombinant expression technology is usually implemented to produce polypeptides in sufficient quantity. Unfortunately, many recombinant systems produce polypeptides having different biological properties than the naturally-derived forms, which may affect the pharmacokinetics, safety, and efficacy of a therapeutic product.

[0003] Therefore, a need remains for developing SAP polypeptides suitable for therapeutic treatment of humans. WO 2009/009019 describes SAP variants for therapy.

SUMMARY OF THE INVENTION

[0004] The present invention is defined by the appended claims. In more detail, the present invention relates to a pharmaceutical composition comprising an SAP variant comprising five SAP protomers, wherein each of the SAP protomers have an amino acid sequence at least 95% identical to SEQ ID NO: 1, and wherein all five SAP protomers comprise an amino acid at position 167 of SEQ ID NO: 1 that is asparagine (N), glutamine (Q), alanine (A), or histidine (H), and a pharmaceutically acceptable ophthalmic vehicle.

[0005] In part, the disclosure provides Serum Amyloid P (SAP) variants and SAP oligomers. In certain aspects, the disclosure provides an SAP variant comprising five SAP protomers, wherein each of the SAP protomers have an amino acid sequence at least 90% identical to SEQ ID NO: 1, and wherein at least one of the SAP protomers comprises one or more amino acid modifications that alter a biological activity of the SAP variant compared to a corresponding sample of serum-derived human SAP. In preferred aspects, a variant SAP protomer comprises at least one amino acid modification that is characterized by the presence of one or more variant amino acids relative to SEQ ID NO: 1, the absence of one or more amino acids relative to SEQ ID NO: 1, the coupling of one or more amino acids to a modifying moiety (e.g., a PEG moiety, a dextran moiety, etc.), or a combination thereof. In some embodiments, the SAP variant is a variant of a human SAP protein. In some embodiments, one or more of the SAP protomers have an amino acid sequence at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 100% identical to SEQ ID NO: 1. In preferred embodiments, SAP variants have an altered biological activity selected from one or more of increased plasma half-life, increased *in vitro* stability, or increased *in vivo* stability. In some embodiments, SAP variants of the disclosure are characterized by increased efficiency of manufacturing the SAP protein (e.g., greater yield of the protein product, increased homogeneity of the protein product, increased stability of the protein product).

[0006] In certain aspects, the disclosure provides SAP variants comprising one or more SAP protomers that are substantially free of N-linked or O-linked glycans. In some embodiments, an SAP protomer comprises an amino acid modification at position 32 of SEQ ID NO: 1 that inhibits attachment of an N-linked glycan. In some embodiments, at least one SAP protomer comprises an amino acid at position 32 of SEQ ID NO: 1 that is not asparagine (N). In preferred embodiments, at least one SAP protomer comprises an aspartate (D), glutamine (Q), or glutamate (E) at position 32 of SEQ ID NO: 1.

[0007] In certain aspects, the disclosure provides SAP variants that are more resistant to protease cleavage than a corresponding sample of serum-derived human SAP. In some embodiments, the SAP variants of the disclosure are more resistant to protease cleavage by a serine protease, a threonine proteases, a cysteine protease, an aspartic acid protease, a metalloprotease, a glutamic acid protease, or combinations thereof. In certain embodiments, the disclosure provides SAP variants that are more resistant to protease cleavage by chymotrypsin, trypsin, Pronase, or combinations thereof. In some embodiments, a protease-resistant SAP variant comprises at least one SAP protomer comprises an amino acid at position 144 of SEQ ID NO: 1 that is not phenylalanine (F). In preferred embodiments, a protease-resistant SAP variant comprises at least one SAP protomer comprising an amino acid at position 145 of SEQ ID NO: 1 that is not aspartate (D). In some embodiments, a protease resistant SAP variant comprises at least one SAP protomer comprising an amino acid at position 144 of SEQ ID NO: 1 that is not phenylalanine (F) and an amino acid at position 145 of SEQ ID NO: 1 that is not aspartate (D). In preferred embodiments, a protease-resistant SAP variant comprises at least one SAP protomer comprising a leucine (L), isoleucine (I), valine (V), or alanine (A) at position 144 of SEQ ID NO: 1. In preferred embodiments, a protease-resistant SAP variant comprises at least one SAP protomer comprising a glutamate (E) at position 145

of SEQ ID NO: 1.

[0008] In certain aspects, the disclosure provides SAP variants that are more resistant to calcium-dependent autoaggregation than a corresponding sample of serum-derived human SAP. In some embodiments, an SAP variant that is resistant to calcium-dependent autoaggregation comprises at least one SAP protomer comprising an amino acid at position 167 of SEQ ID NO: 1 that is not glutamate (E). In preferred embodiments, an SAP variant that is resistant to calcium-dependent autoaggregation comprises at least one SAP protomer comprising an aspartate (D), asparagine (N), glutamine (Q), alanine (A), or histidine (H) at position 167 of SEQ ID NO: 1.

[0009] In certain aspects, the disclosure provides SAP variants comprises at least one SAP protomer comprising one or more amino acids that are covalently attached to one or more inert polymers. In some embodiments, at least one of the inert polymers is a polyethylene glycol (PEG) moiety. In certain embodiments, one or more of the SAP protomers comprise at least one native or variant (e.g., by amino acid substitution, addition, or deletion) cysteine (C), relative to SEQ ID NO: 1, which has an attached PEG moiety. In a preferred embodiment, one or more of the SAP protomers comprise a variant cysteine (C), located at the N-terminus of SEQ ID NO: 1, which has an attached PEG moiety. In other embodiments, one or more of the SAP protomers comprise at least one native or variant (Q), relative to SEQ ID NO: 1, which has an attached PEG moiety. In a preferred embodiment, one or more of the SAP protomers comprises a glutamine (Q) at position 32 of SEQ ID NO: 1 that has an attached PEG moiety. In some embodiments, the SAP variant comprises at least one SAP protomer comprising one or more cysteine (C) residues and one or more glutamine (Q) residues that are attached to a PEG moiety. In some embodiments, at least one of the inert polymers is a dextran moiety. In certain embodiments, one or more of the SAP protomers comprises a native or variant glutamine (Q) residue, relative to SEQ ID NO: 1, which has an attached dextran moiety. In a preferred embodiment, one or more of the SAP protomers comprises a native glutamine residue at position 32 of SEQ ID NO: 1 that has an attached dextran moiety. In certain embodiments, the SAP variant comprises at least one SAP protomer comprising one or more amino acids attached to a PEG moiety and one or more amino acids attached to a dextran moiety.

[0010] In certain aspects, the disclosure provides an SAP variant comprised of at least two, at least three, at least four, or at least five different variant SAP protomers as described herein.

[0011] In certain aspects, the disclosure provides a covalently crosslinked SAP oligomer comprising at least two SAP pentamers, wherein each of the SAP pentamers comprises five SAP protomers. SAP oligomers of the invention may comprise SAP protomers at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 100% identical to the amino acid sequence of SEQ ID NO: 1. Accordingly, SAP oligomers may comprise at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine or more of the SAP variant protomers as described herein. In some embodiments, SAP oligomers may be comprised of at least two, at least three, at least four, at

least five, at least six, at least seven, at least eight, at least nine or more different variant SAP protomers as described herein. In preferred embodiments, the crosslinked SAP oligomers are characterized by one or more of increased plasma half-life, increased *in vitro* stability, and increased *in vivo* stability compared to a corresponding sample of SAP isolated from human serum.

[0012] In certain aspects, the SAP oligomers are comprised of SAP pentamers covalently attached through one or more chemical cross-linkers. In certain embodiments, at least one of the chemical cross-linker is a heterobifunctional agent selected from succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate, m-Maleimidobenzoyl-N-hydroxysuccinimide ester, N-succinimidyl(4-iodoacetyl)aminobenzoate, succinimidyl 4-(p-maleimidophenyl)butyrate, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, 4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)-toluene, N-succinimidyl 3-(2-pyridyldithio)propionate, or succinimidyl 6-((3-(2-pyridyldithio) propionate)hexanoate. In certain embodiments, at least one of the chemical cross-linkers is a homobifunctional agent selected from disuccinimidyl suberate, bismaleimido-hexane, or dimethylpimelidate-2 HCl. In certain embodiments, at least one of the chemical cross-linkers is a photoreactive agent selected from bis-(β -(4-azidosalicylamido)ethyl)disulfide or N-succinimidyl-6-(4'-azido-2'-nitrophenyl-amino)hexanoate.

[0013] In certain aspects, the disclosure provides a pharmaceutical preparation suitable for use in a mammal comprising one or more of the SAP variants and/or covalently crosslinked SAP oligomers. Pharmaceutical preparations of the invention include at least one of the SAP variants and/or SAP oligomers disclosed herein and a pharmaceutically acceptable carrier. In some embodiments, the pharmaceutical preparation further comprises an additional active agent. In some embodiments, the pharmaceutical preparation is prepared as a sustained release formulation. In some embodiments, pharmaceutical preparations of the disclosure are suitable for administration to a patient topically, by injection, by intravenous injection, by inhalation, by continuous depot, or by pump.

[0014] The disclosure further provides methods for treating or preventing SAP-responsive disorders or conditions by administering to a patient in need thereof a therapeutically effective amount of one or more of the SAP variants and/or SAP oligomers. SAP-responsive disorders or conditions include, but are not limited to, fibrotic or fibroproliferative disorders or conditions, hypersensitivity disorders or conditions, autoimmune disorders or conditions, and mucositis. The SAP variant and/or oligomer of the invention may be administered to a patient topically, by injection, by intravenous injection, by inhalation, by continuous depot or pump, or a combination thereof. In some embodiments, the SAP variant and/or oligomer of the invention is administered conjointly with one or more additional active agents. In certain embodiments, the SAP variants and/or oligomers are formulated to be administered conjointly. The SAP variants and/or oligomers may be conjointly administered as separate or in combined formulations. The SAP variants and/or oligomers may be administered simultaneously or at different dosing schedules.

BRIEF DESCRIPTION OF THE DRAWINGS**[0015]**

Figure 1: An SAP variant comprising an amino acid substitution E167Q, relative to the sequence of SEQ ID NO: 1, is more resistant to calcium-mediated aggregation than a corresponding sample of unmodified recombinant human SAP (rhSAP). Incremental amounts of calcium was added a solution comprising SAP, and the amount of SAP aggregation was observed by measuring the absorbance of the solution at 600 nm in a spectrophotometer.

Figure 2: An SAP variant comprising an amino acid substitution E167Q, relative to the sequence of SEQ ID NO: 1, has a similar plasma half-life compared to a corresponding sample of unmodified recombinant human SAP (rhSAP). Rats were administered SAP (1 mg/kg i.v. does per rat, n=3). Over twenty-four hours, rates were assessed for plasma concentrations (µg/ml) of SAP protein.

Figure 3: SAP variants E167Q and N32D, relative to the sequence of SEQ ID NO: 1, are at least as active as a corresponding sample of unmodified recombinant human SAP (rhSAP). Monocyte-enriched Peripheral Blood Mononuclear Cells (PMBCs) were incubated with varying concentrations of SAP. Following incubation, the resulting culture supernatants were removed and assayed by ELISA to quantify the amount of Macrophage Derived Chemokine (MDC) that was produced.

Figure 4: The SAP variant N32D has a similar plasma half-life to that of wild-type rhSAP. While an asialo form of hSAP has a significantly reduced plasma half-life compared to a corresponding sample of unmodified recombinant human SAP (rhSAP). Rats were administered SAP (1 mg/kg i.v. does per rat, n=3). Over twenty-four hours, rates were assessed for plasma concentrations (µg/ml) of SAP protein.

Figure 5: Depicts a chemical reaction that covalent attaches PEG to rhSAP.

Figure 6: Pegylated-rhSAP was purified from reaction components by anion exchange chromatography. Fractions from the chromatography column were pooled and concentrated before analysis by SDS-PAGE.

DETAILED DESCRIPTION OF THE INVENTION**Overview**

[0016] Serum amyloid P ("SAP") is a naturally-occurring serum protein in mammals and is a

member of the pentraxin family of structurally related proteins. It is produced in the liver as a 125,000 Dalton glycoprotein and has a physiological half-life of 24 hours in serum. SAP is composed of five identical subunits or "protomers" which are non-covalently associated in a disc-like molecule. SAP protomers non-covalently associate with each other via two "protomer interfaces". Protomer interface 1 from subunit 1 associates with protomer interface 2 from subunit 2. Protomer interface 1 from subunit 2 associates with protomer interface 2 from subunit 3, etc. Each protomer exposes an "A-face" that can bind FcγR and an opposing "B-face" that mediates calcium binding and calcium-mediated ligand binding. In high concentrations of ionic calcium, SAP aggregates and may precipitate as the amyloid P component, which is a normal constituent of glomerular basement membrane as well as human dermis, cervix, testis, and placenta tissues. See Baltz, M. L., et al., Clin. Exp. Immunol., 66:691-700 (1986); Dyck, R. F., et al., J. Exp. Med., 152:1162-1174 (1980); Melvin, T., Am. J. Pathol., 125:460-464 (1986); Breathnach, S. M., J. Invest. Derm., 92:53-58 (1989); Clayton, J., Cell. Pathol., 43:63-66 (1983); Herriot, R., et al., J. Pathol., 157:11-14 (1989); Khan, A. M., et al., Placenta, 6:551-554 (1985). The mature sequence of the human SAP protomer is depicted in below (amino acids 20-223 of Genbank Accession No. NP_001630; signal sequence not depicted).

```
HTDLSGKVFVFPRESVTDHVNLTIPLEKPLQNFTLCFRAYS DLSRAYSLF
SYNTQGRDNELLYKERVGEYSLYIGRHKVTSKVIEKFPAPVHICVSWE
SSSGIAEFWINGTPLVKKGLRQGYFVEAQP KIVLGQEQDSYGGKFDRSQ
SFVGEIGDLYMWDSVLPPENILSAYQGTPLPANILDWQALNYEIRGYVII
KPLVWV (SEQ ID NO: 1)
```

[0017] Normal wound healing processes as well as the dysregulated events that cause fibrosis involve the proliferation and differentiation of fibroblasts and the deposition of extracellular matrix. Whether these fibroblasts are derived locally or from a circulating precursor population is unclear. Fibrocytes, fibrocyte precursors, myofibroblast precursors, and hematopoietic monocyte precursors belong to a distinct population of fibroblast-like cells derived from peripheral blood monocytes. These cells can migrate to sites of tissue injury to promote angiogenesis and wound healing. CD14⁺ peripheral blood monocytes cultured in the absence of serum or plasma differentiate into fibrocytes within 72 hours. Recently, SAP was shown to inhibit fibrocyte, fibrocyte precursor, myofibroblast precursor, and/or hematopoietic monocyte precursor differentiation at levels similar to that of serum. In contrast, plasma depleted of SAP has a reduced ability to inhibit differentiation of monocytes into fibrocytes, fibrocyte precursors, myofibroblast precursors, and/or hematopoietic monocyte precursors. Compared with sera from healthy individuals, serum from subjects with rheumatoid arthritis, scleroderma, mixed connective tissue diseases, and certain systemic fibrotic diseases have reduced potency for inhibiting fibrocyte, fibrocyte precursor, myofibroblast precursor, and/or hematopoietic monocyte precursor differentiation *in vitro*. Accordingly, serum levels of SAP are significantly lower in some subjects with these disorders than is observed for healthy subjects. These results indicate that abnormally low levels of SAP may augment pathological processes leading to fibrosis and suggests SAP may be useful as a therapeutic agent to inhibit fibrosis in chronic inflammatory conditions. Recently, it has been suggested that SAP can be used as a

therapeutic agent to treat various other disorders, including fibrosis-related disorders, hypersensitivity disorders, autoimmune disorders, mucositis, and inflammatory disorders such as those caused by microbial infection. See, for example, U.S. Patent Application Nos. 11/707,333, 12/217,617 12/720,845, and 12/720,847.

[0018] Polypeptides are susceptible to denaturation or enzymatic degradation in the blood, liver or kidney. Due to the low stability of some polypeptides, it has been required to administer polypeptide drugs in a sustained frequency to a subject in order to maintain an effective plasma concentration of the active substance. Moreover, since polypeptide drugs are usually administrated by infusion, frequent injection of polypeptide drugs may cause considerable discomfort to a subject. Thus, there have been many studies to develop polypeptide drugs that have an increased circulating half-life in the blood, while maintaining a high pharmacological efficacy. Accordingly, a primary object of the present disclosure is to provide SAP variants, compositions, pharmaceutical preparations and formulations having a prolonged *in vivo* half-life compared to human SAP. Advantages of increased plasma half-life include, but are not limited to, reducing the amount and/or frequency of dosing.

[0019] Furthermore, pharmaceutical compositions of therapeutic peptides preferably have a shelf-life of several years in order to be suitable for common use. However, peptide compositions are inherently unstable due to sensitivity towards chemical and physical degradation. Examples of chemical degradation include change of covalent bonds, including but not limited to, oxidation, hydrolysis, racemization, or crosslinking. Examples of physical degradation include conformational changes relative to the native structure of the peptide, which may lead to aggregation, precipitation, or adsorption of the polypeptide to surfaces. Accordingly, a further object of the present disclosure is to provide SAP variants, compositions, pharmaceutical preparations and formulations that have a prolonged shelf-life, or rather increased *in vitro* stability, compared to human SAP. During the manufacturing process, it is often difficult to produce large quantities of a protein with reproducible consistency in the characteristics of the product, such as post-translational modification and/or folding. In some embodiments, SAP variants of the disclosure are characterized by increased efficiency of manufacturing the SAP protein (e.g., greater yield of the protein product, increased homogeneity of the protein product, increased stability of the protein product), particularly for *in vivo* use (e.g., as a therapeutic agent).

Definitions

[0020] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry, and nucleic acid chemistry and hybridization are those well known and commonly employed in the art. Standard techniques are used for nucleic acid and peptide synthesis. The techniques and procedures are generally performed according to conventional methods in the art and various general references (e.g., Sambrook et al., 1989, Molecular

Cloning: A Laboratory Manual, 2d ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which are provided throughout this document.

[0021] The articles "a" and "an" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0022] As used herein, the terms "treatment", "treating", and the like, refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disorder or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disorder and/or adverse affect attributable to the disorder. "Treatment", as used herein, covers any treatment of a disease in a mammal, particularly in a human, and includes: (a) increasing survival time; (b) decreasing the risk of death due to the disease; (c) decreasing the risk of a disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (d) inhibiting the disease, i.e., arresting its development (e.g., reducing the rate of disease progression); and (e) relieving the disease, i.e., causing regression of the disease.

[0023] As used herein, a therapeutic that "inhibits" or "prevents" a disorder or condition is a compound that, in a statistical sample, reduces the occurrence of the disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated control sample.

[0024] As used herein the terms "subject" and "patient" refer to animals including mammals, such as humans. The term "mammal" includes primates, domesticated animals including dogs, cats, sheep, cattle, horses, goats, pigs, mice, rats, rabbits, guinea pigs, captive animals such as zoo animals, and wild animals.

[0025] As used herein the term "tissue" refers to an organ or set of specialized cells such as skin tissue, lung tissue, kidney tissue, and other types of cells.

[0026] The term "therapeutic effect" is art-recognized and refers to a local or systemic effect in animals, particularly mammals, and more particularly humans caused by a pharmacologically active substance. The phrase "therapeutically effective amount" means that amount of such a substance that produces some desired local or systemic effect at a reasonable benefit/risk ratio applicable to any treatment. The therapeutically effective amount of such substance will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. For example, certain compositions described herein may be administered in a sufficient amount to produce a desired effect at a reasonable benefit/risk ratio applicable to such treatment.

[0027] As used herein, the term "nucleic acid" refers to a polynucleotide such as

deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include, as equivalents, analogs of either RNA or DNA made from nucleotide analogs, and, as applicable to the embodiment being described, single-stranded (such as sense or antisense) and double-stranded polynucleotide.

[0028] The terms "peptides", "proteins" and "polypeptides" are used interchangeably herein. The term "purified protein" refers to a preparation of a protein or proteins that are preferably isolated from, or otherwise substantially free of, other proteins normally associated with the protein(s) in a cell or cell lysate. The term "substantially free of other cellular proteins" or "substantially free of other contaminating proteins" is defined as encompassing individual preparations of each of the proteins comprising less than 20% (by dry weight) contaminating protein, and preferably comprises less than 5% contaminating protein. Functional forms of each of the proteins can be prepared as purified preparations by using a cloned gene as is well known in the art. By "purified", it is meant that the indicated molecule is present in the substantial absence of other biological macromolecules, such as other proteins (particularly other proteins which may substantially mask, diminish, confuse or alter the characteristics of the component proteins either as purified preparations or in their function in the subject reconstituted mixture). The term "purified" as used herein preferably means at least 80% by dry weight, more preferably in the range of 85% by weight, more preferably 95-99% by weight, and most preferably at least 99.8% by weight, of biological macromolecules of the same type present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 5000, can be present). The term "pure" as used herein preferably has the same numerical limits as "purified" immediately above.

[0029] The term "half-life" or "plasma half-life", as used herein in the context of administering a peptide drug to a subject, is defined as the time required for plasma concentration of a drug in a subject to be reduced by one half. Further explanation of "half-life" is found in *Pharmaceutical Biotechnology* (1997, DFA Crommelin and RD Sindelar, eds., Harwood Publishers, Amsterdam, pp 101 120).

SAP Variants and SAP Oligomers

(ii) SAP variant proteins

[0030] In part, the disclosure provides Serum Amyloid P (SAP) variant proteins. The term "SAP variant" is intended to refer to an SAP protein comprising five SAP subunits or "protomers". In preferred aspects, an SAP variant comprises at least one SAP protomer having one or more amino acid modifications (i.e., a variant SAP protomer) that modify at least one biological activity of the SAP protein. In some embodiments, amino acid modifications include, but are not limited to, the presence of one or more variant amino acids relative to the sequence of SEQ ID NO: 1 (e.g., amino acid substitution or addition), the absence of one or more native amino

acids relative to the sequence of SEQ ID NO: 1 (e.g., amino acid deletion), the coupling of one or more amino acids to a modifying moiety (e.g., a PEG moiety, a dextran moiety, etc.), or a combination thereof. In some embodiments, an SAP protomer comprises at least one variant amino acid, relative to SEQ ID NO: 1, and at least one amino acid coupled to a modifying moiety. In particular, SAP variants are characterized by an altered biological activity compared to a corresponding sample of serum-derived human SAP. In some aspects, an SAP variant of the disclosure is characterized by an altered biological activity selected from one or more of increased plasma half-life, increased *in vivo* stability, increased *in vitro* stability, or increased manufacturing efficiency.

[0031] The term "SAP protomer" is intended to refer to a polypeptide that is at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to the SAP protomer exemplified by SEQ ID NO: 1. Accordingly, the term "SAP protomer" encompasses fragments and fusion proteins comprising any of the preceding. In preferred aspects, SAP variants of the disclosure are human SAP proteins. Generally, an SAP protomer will be designed to be soluble in aqueous solutions at biologically relevant temperatures, pH levels, and osmolarity. The protomers that non-covalently associate together to form an SAP variant of the disclosure may have identical amino acid sequences and/or post-translational modifications or, alternatively, individual protomers may have different sequences and/or modifications. Accordingly, an SAP variant may be comprised of at least two, at least three, at least four, or five identical SAP protomers or, alternatively, comprised of at least two, at least three, at least four, or at least five identical or different variant SAP protomers. In some embodiments, at least one, at least two, at least three, or at least four of the SAP protomers have 100% sequence identity to SEQ ID NO: 1. In preferred embodiments, an SAP variant comprises at least one variant SAP protomer that confers one or more altered biological activity as described herein. Post-translational modifications may be effected *in vivo* and/or *in vitro* and include, but are not limited to, processing (e.g., signal sequence removal, pro-peptide maturation, etc.) and chemical modification (e.g., glycosylation, pegylation, etc.) of the translated SAP polypeptides.

[0032] The invention also provides SAP protomers sharing a specified degree of sequence identity or similarity to an SAP polypeptide. To determine the percent identity of two amino acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or more of the length of a reference sequence (e.g., human SAP) is aligned for comparison purposes. The amino acid residues at corresponding amino acid positions are then compared. When a position in the first sequence is occupied by the same amino acid residue as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid "identity" is equivalent to amino acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for

optimal alignment of the two sequences.

[0033] The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

[0034] In certain embodiments, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (J Mol. Biol. (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>). In a specific embodiment, the following parameters are used in the GAP program: either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., et al., Nucleic Acids Res. 12(1):387 (1984)) (available at <http://www.gcg.com>). Exemplary parameters include using an NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6.

[0035] In another embodiment, the percent identity between two amino acid sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)), which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0036] Another embodiment for determining the best overall alignment between two amino acid sequences can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci., 6:237-245 (1990)). In a sequence alignment the query and subject sequences are both amino acid sequences. The result of said global sequence alignment is presented in terms of percent identity. In certain embodiments, amino acid sequence identity is performed using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci., 6:237-245 (1990)). In a specific embodiment, parameters employed to calculate percent identity and similarity of an amino acid alignment comprise: Matrix=PAM 150, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5 and Gap Size Penalty=0.05.

[0037] Some aspects of the invention provide SAP polypeptides (i.e., protomers), or provide therapeutic methods for employing those polypeptides, wherein said polypeptides are defined, at least in part, to a reference sequence. In preferred embodiments, the reference sequence corresponds to the amino acid sequence of SEQ ID NO: 1. Accordingly, such polypeptides may have a certain percentage of amino acid residues which are not identical to a reference sequence. In one preferred embodiment, the non-identical residues have similar chemical

properties to the residues to which they are not identical. Groups that have similar properties include the following amino acids: E, D, N, and Q; H, K, and R; Y, F and W; I, L, V, M, C, and A; and S, T, C, P, and A.

[0038] In another embodiment, the residues which are not identical are those which are not evolutionarily conserved between the reference sequence and an orthologous sequence in at least one evolutionarily related species, such as in species within the same order. In the case of a mammalian reference sequence, the amino acids that may be mutated in a preferred embodiment are those that are not conserved between the reference sequence and the orthologous sequence in another mammal species. For example, if a polypeptide used in a method of the present invention is said to comprise an amino acid sequence that is at least 90% identical to human SAP (SEQ ID NO:1), then said polypeptide may have non-identical residues to those positions in which the human SAP and that of another mammal differ.

[0039] SAP polypeptides (i.e., protomers) sharing at least 90% identity with SEQ ID NO:1 include polypeptides having conservative substitutions in these areas of divergence. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile, interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr. Additional guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie et al., Science 247:1306-1310 (1990).

[0040] The disclosure also provides SAP protomers with mutations in specific amino acid residues. Exemplary mutations are disclosed herein and are numbered according to the amino acid position of human SAP, e.g., as exemplified in SEQ ID NO: 1. In certain embodiments, an SAP variant i) comprises one or more protomers that are at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% identical to SEQ ID NO: 1, ii) has one or more of the following features compared to a corresponding sample of serum-derived human SAP: increased plasma half-life, increased *in vitro* stability, increased *in vivo* stability, or increased manufacturing efficiency. As described herein, amino acids of an SAP protomer may be mutated by adding one or more specified amino acid residues, deleting one or more specified amino acid residues, or substituting one or more specified amino acid residues. Methods for mutating or chemically modifying SAP polypeptides are described in the following sections.

[0041] Enhanced serum half-life and *in vivo* stability may be desirable to reduce the frequency of dosing that is required to achieve therapeutic effectiveness. Accordingly, in certain aspects, the serum half-life of an SAP variant is at least one, at least two, at least three, at least four, at least five, at least ten, or at least twenty days or more. Methods for pharmacokinetic analysis and determination of half-life and *in vivo* stability will be familiar to those skilled in the art. Details may be found in Kenneth, A et al: Chemical Stability of Pharmaceuticals: A Handbook for Pharmacists and in Peters et al, Pharmacokinetic analysis: A Practical Approach (1996). Reference is also made to "Pharmacokinetics", M Gibaldi & D Perron, published by Marcel

Dekker, 2nd Rev. ex edition (1982), which describes pharmacokinetic parameters such as t_{α} and t_{β} half lives and area under the curve (AUC). As described in the examples of the disclosure, one method for determining *in vivo* stability involves administering an SAP variant to an animal and measuring the concentration of the SAP variant within the plasma of the animal at regular intervals after administration. The pharmacokinetic profile (i.e. plasma concentration of SAP over time) of an SAP variant can be compared to that of another SAP protein, e.g., a corresponding sample of serum-derived human SAP.

[0042] In certain aspects, SAP variants of the disclosure are more stable than an otherwise identical composition of human SAP under identical conditions. In certain embodiments, the disclosed compositions have a shelf-life, or *in vitro*, stability at least two times, at least three times, at least four times, or at least five times or more as long as a corresponding sample human SAP. Many methods for measure *in vitro* stability are known in the art, including, for example, measuring protein stability by SDS-PAGE, Western blot, RP-HPLC, AEX-HPLC, LC-MS, or N-terminal sequencing.

[0043] In some embodiments, SAP variants of the invention have an altered or similar bioactivity compared to a corresponding sample of serum-derived human SAP. Bioactivity of an SAP variant may be determined, for example, by determining the IC_{50} for inhibiting the differentiation of monocytes into fibrocytes *in vitro*. In some embodiments, the IC_{50} of an SAP variant is less than 1/2, less than 1/3, less than 1/4, less than 1/10, or less than 1/100 that of a corresponding sample of wild-type SAP isolated from human serum. There are many well characterized methods for determining the responsiveness of Peripheral Blood Mononuclear Cells (PBMCs) or monocyte cells to SAP for fibrocyte differentiation. These methods may be used to determine the relative potency of any of the SAP variant of the invention in comparison to a sample of human serum-derived SAP, any other SAP variant polypeptide, or other fibrocyte suppressant or activating agent. PBMCs or monocytes suitable for use in these methods may be obtained from various tissue culture lines. Alternatively, suitable cells for fibrocyte differentiation assays may be obtained from any biological sample that contains PBMC or monocyte cells. The biological sample may be obtained from serum, plasma, healthy tissue, or fibrotic tissue. In general, fibrocyte differentiation assays are conducted by incubating PBMC or monocyte cells in media with various concentrations of a SAP polypeptide to determine the degree of fibrocyte differentiation. The concentration of SAP can range from 0.0001 $\mu\text{g/mL}$ to 1 mg/mL , and in some embodiments is 0.001 $\mu\text{g/mL}$, 1.0 $\mu\text{g/mL}$, 5 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 15 $\mu\text{g/mL}$, 20 $\mu\text{g/mL}$, 25 $\mu\text{g/mL}$, 30 $\mu\text{g/mL}$, 35 $\mu\text{g/mL}$, 40 $\mu\text{g/mL}$, 45 $\mu\text{g/mL}$, 50 $\mu\text{g/mL}$, 100 $\mu\text{g/mL}$, 200 $\mu\text{g/mL}$, 300 $\mu\text{g/mL}$, or 500 $\mu\text{g/mL}$. In some assays, the media can be supplemented with between 1-100 ng/mL hMCSF; the preferred concentration of hMCSF being 25 ng/mL . The indication that PBMC and monocytes have differentiated into fibrocytes can be determined by one skilled in the art. In general, fibrocytes are morphologically defined as adherent cells with an elongated spindle-shape and the presence of an oval nucleus. In some assays, cells are fixed and stained with Hema 3 before enumerating fibrocytes by direct counting, e.g., using an inverted microscope. The amount of fibrocyte differentiation is interpreted by one skilled in the art as an indication of a cell's responsiveness to SAP. As indicated by the examples of the disclosure, a greater suppression of fibrocyte differentiation

indicates a greater degree of SAP responsiveness. An alternative method of measuring fibrocyte differentiation involves determining the expression of fibrocyte-specific cell surface markers or secreted factors, e.g., cytokines (such as IL-1ra, ENA-78/CXCL-5, PAI-1), fibronectin, collagen-1, Macrophage Derived Chemokine). Methods of detecting and/or quantifying cell surface markers or secreted factors are well known in the art, including but not limited to various ELISA- and FACS-based techniques using immunoreactive antibodies against one or more fibrocyte specific markers.

[0044] In certain aspects, the disclosure provides SAP variants that are more resistant to protease cleavage than a corresponding sample of serum-derived human SAP. An SAP variant of the invention may be resistant to protease cleavage from any number of proteases including serine proteases, threonine proteases, cysteine proteases, aspartic acid proteases, metalloproteases, and glutamic acid proteases. In certain embodiments, an SAP variant is more resistant to cleavage by chymotrypsin, trypsin, or Pronase. In preferred embodiments, the protease resistant SAP variant has an increased plasma half-life compared to a corresponding sample of serum-derived human SAP. Methods for measuring proteolytic cleavage include, but are not limited to, analyzing protease-treated samples of a protein (e.g., an SAP variant and a serum-derived human SAP standard) by SDS-PAGE, Western blot, RP-HPLC, AEX-HPLC, LC-MS, or N-terminal sequencing. Other examples of protease cleavage assays are within the purview of a person skilled in the art and are exemplified in Kinoshita CM, et al., Protein Science 1:700-709 (1992). Therefore, SAP variants of the invention can readily be assayed for relative resistance to protease cleavage in comparison to a corresponding sample of another SAP protein, e.g., a sample of serum-derived human SAP.

[0045] In the absence of calcium, human SAP is susceptible to α -chymotrypsin cleavage between residues Phe₁₄₄ and Asp₁₄₅ (Kinoshita CM, et. al., Protein Science 1:700-709 (1992)). In certain embodiments, an SAP protomer comprises an amino acid modification at position 144 and/or position 145 of SEQ ID NO: 1, resulting in an SAP variant that is more resistant to protease cleavage. In some embodiments, an SAP protomer comprises a variant amino acid at position 144 of SEQ ID NO: 1. In particular, SAP variants more resistant to protease cleavage may have a leucine (L), isoleucine (I), valine (V), alanine (A), or glutamine (Q) residue at amino acid position 144 of SEQ ID NO: 1. An SAP protomer may also comprise, independently or in combination with, a variant amino acid at position 145 of SEQ ID NO: 1. Variant SAP protomers of the disclosure may comprise a glutamate (E) at position 145 at SEQ ID NO: 1. In certain embodiments, an SAP variant comprises one or more promoters that are i) at least 85%, at least 90%, at least 95%, at least 96% at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, and ii) comprise one or more of the following amino acid substitutions F144L, F144I, F144V, F144A, F144G, D145E relative to SEQ ID NO: 1. Any of the aforementioned SAP protomers that are resistant to protease cleavage may further comprise any of the other amino acid modifications described herein.

[0046] The disclosure further provides SAP variants with increased metal-binding as compared to a corresponding sample of serum-derived human SAP. Increased metal-binding in the calcium-binding site of an SAP protomer decreases the susceptibility of the protomer to

proteolysis by stabilizing the loop structure containing the amino acid residue at position 145 of SEQ ID NO: 1. In certain embodiments, an SAP variant having increased metal-binding comprises one or more SAP protomers with a variant amino acid at position 145 of SEQ ID NO: 1. In particular, SAP variants characterized by increased metal-binding may have a glutamate (E), glutamine (Q), histidine (H), alanine (A), glycine (G) amino acid at position 145 of SEQ ID NO: 1. In preferred embodiments, an SAP variant demonstrate increased metal-binding to calcium as compared to a corresponding sample of serum-derived human SAP. Calcium binding constants for an SAP protein can be measured by a variety of methods, including those exemplified in Calcium-binding Protein Protocols: methods and techniques by Hans J. Vogel, Contributor Hans J. Vogel, Published by Humana Press, 2002. In some embodiments, calcium binding constants for an SAP protein can be measured by equilibrium dialysis using a range of calcium concentrations followed by Scatchard plot analysis. (See, for example Segel, I.H., Enzyme Kinetics 1975, Wiley-Interscience Publisher, p218-19). Equilibrium dialysis may be performed using either radioactive isotopes of calcium or calcium sensitive electrodes to quantify free calcium levels. Calcium binding constants may also be determined by titrating calcium into a solution of SAP in the presence of a chromophoric chelator (5,5'-dibromo-1,2-bis(2aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (Linse, S, Helmersson, A. Forsen, S, 1991 JBC 266:13 pp. 8050-8054). Isothermal Titration Calorimetry can also be used to measure calcium binding affinities (Wiseman, T., Williston, S., Brandts, J.F., and Lin, L.N., (1989) Anal Biochem 179, 131-7). SAP variants characterized by increased metal-binding can be compared to a corresponding sample of serum-derived human SAP for changes in proteolytic stability (e.g., digestion with chymotrypsin in the presence and absence of calcium), *in vitro* bioactivity and or pharmacokinetics, as well as biophysical characterization methods (e.g., RP-HPLC, SE-HPLC, SDS-PAGE, LC-MS). In certain embodiments, an SAP variant comprises one or more SAP protomers that are i) at least 85%, at least 90%, at least 95%, at least 96% at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, and ii) comprising one or more of the following amino acid substitutions: D145E, D145Q, D145H, D145A, or D145G. Any of the aforementioned SAP protomers that demonstrate increased metal-binding may further comprise any of the other amino acid modifications described herein.

[0047] In the absence of ligand, calcium binding results in the autoaggregation of SAP (Emsley, et. al. Nature 367:338-345 (1994)), and once aggregated, SAP is rapidly cleared from the blood stream (Pepys, et. al., Nature 417:254-259 (2002)). In certain embodiments, an SAP variant of the disclosure is more resistant to calcium-dependent autoaggregation than a corresponding sample of serum-derived human SAP. In some embodiments, an SAP variant resistant to calcium-dependent autoaggregation comprises one or more SAP protomers comprising a variant amino acid at position 167 of SEQ ID NO: 1. In some embodiments, an SAP variant resistant to calcium-dependent autoaggregation comprises an aspartate (D), asparagines (N), glutamine (Q), alanine (A), or histidine (H) at position 167 of SEQ ID NO: 1. Aggregation of SAP can be determined by any number of known methods including gel filtration chromatography and dynamic light scattering (see Ho, et. al., J Biol Chem 280:31999-32008 (2005)). Therefore, SAP variants of the invention can readily be assayed for relative resistance to aggregation in comparison to a corresponding sample of another SAP protein,

e.g., a sample of serum-derived human SAP. In certain embodiments, an SAP variant comprises one or more SAP protomers that are at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, and ii) comprises one or more of the following amino acid substitutions: E167D, E167N, E167Q, E167A, E167H. Any of the aforementioned SAP protomers that are resistant to autoaggregation may further comprise any of the other amino acid modifications described herein.

[0048] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of a carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential N-linked glycosylation site. O-linked glycosylation refers to the attachment of sugar moieties (e.g., N-acetylgalactosamine, galactose, or xylose) to a hydroxyamino acid, most commonly on a serine or threonine residue.

[0049] In certain aspects, the disclosure provides an SAP variant comprising at least one SAP protomer that is substantially free of glycans. By "substantially free" is meant that at least about 25% (e.g., at least about 27%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95%, or at least about 99%) of the amino acids of the SAP protomer are non-glycosylated. In preferred embodiments, an SAP protomer, or SAP variant, is free of any glycan-linked structure.

[0050] In some embodiments, SAP protomers of the disclosure have been modified to inhibit attachment of N-linked glycans, O-linked glycans or both N- and O-linked glycans. Removal of N-linked glycosylation sites on an SAP variant is accomplished by modifying (e.g., by amino acid deletion, addition or substitution) the amino acid sequence of one or more of the SAP protomers such that the protomer lacks one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also include the deletion or substitution of one or more serine or threonine residues of the SAP protomer. In preferred embodiments, an SAP variant comprises at least one SAP protomer comprising a variant amino acid at position 32, 33, and/or 34 of SEQ ID NO: 1. In preferred embodiments, a variant SAP protomer comprises an aspartate (D), glutamine (Q), or glutamate (E) at position 32 of SEQ ID NO: 1. A variant SAP protomer may also comprise, independently or in combination with, a proline at position 33 of SEQ ID NO: 1. In certain embodiments, an SAP variant comprises one or more protomers that are i) at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, and ii) comprises one or more of the following amino acid substitutions: N32D, N32Q, N32E, 33P. Any of the aforementioned SAP protomers that are substantially free of glycans may further comprise any of the other amino acid modifications described herein.

[0051] In certain aspects, an SAP variant of the invention comprises one or more SAP protomers comprising one or more amino acid covalently attached to one or more inert polymers.

An inert polymer attached to an SAP protomer may be of any effective molecular weight and may be branched or unbranched. Polymers used in the instant invention include, but are not limited to, (a) dextran and dextran derivatives, including dextran sulfate, cross-linked dextrin, and carboxymethyl dextrin; (b) cellulose and cellulose derivatives, including methylcellulose and carboxymethyl cellulose; (c) starch, cyclodextrins and dextrans, and derivatives thereof; (d) polyalkylene glycol and derivatives thereof, including PEG, mPEG, PEG homopolymers, polypropylene glycol homopolymers, copolymers of ethylene glycol with propylene glycol, wherein said homopolymers and copolymers are unsubstituted or substituted at one end with an alkyl group; (e) heparin and fragments of heparin; (f) polyvinyl alcohol and polyvinyl ethyl ethers; (g) polyvinylpyrrolidone; (h) α,β -poly((2-hydroxyethyl)-DL-aspartamide; and (i) polyoxyethylated polyols. Any of the aforementioned SAP protomers that have one or more amino acids covalently attached to one or more inert polymers may further comprise any of the amino acid modifications described herein

[0052] In preferred embodiments, the disclosure provides an SAP variant comprising one or more SAP protomers comprising at least one amino acid covalently attached to a polyethylene glycol moiety. In some embodiments, the molecular weight of a polyethylene glycol moiety is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight). Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the degree or lack of antigenicity, etc.). In certain embodiments, the polyethylene glycol may have an average molecular weight of at least 1, at least 20, or at least 40 kDa. The polyethylene glycol may have a branched structure, and branched polyethylene glycols are described, for example, in U.S. Pat. No. 5,643,575; Morpurgo et al., Appl. Biochem. Biotechnol. 56:59-72 (1996); Vorobjev et al., Nucleosides Nucleotides 18:2745-2750 (1999); and Caliceti et al., Bioconjug. Chem. 10:638-646 (1999). Polyethylene glycol moieties may be attached to an SAP variant with consideration of effects on catalytic or targeting portions.

[0053] In preferred embodiments, the disclosure provides an SAP variant comprising one or more SAP protomers comprising at least one amino acid covalently attached to a dextran moiety. In some embodiments, the molecular weight of a dextran moiety attached to the SAP protomer is generally between about 1 kDa and about 250 kDa (the term "about" indicating that in preparations of dextran conjugates, some molecules will weigh more, some less, than the stated molecular weight). Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the degree or lack of antigenicity, etc.). In certain embodiments, the dextran may have an average molecular weight of at least 1, at least 20, or at least 40 kDa. SAP may be conjugated to dextran or a dextran derivative including dextran sulfate, p-aminoethyl cross-linked dextran, and carboxymethyl dextran.

(ii) SAP oligomers

[0054] In certain aspects, the disclosure provides SAP oligomers comprising two or more SAP pentamers. In preferred aspects, the SAP oligomers are covalently-crosslinked pentamers, i.e., via protomer-protomer crosslinks.

[0055] There are a large number of chemical cross-linking agents that are known to those skilled in the art as well as their method of use. In certain embodiments, SAP pentamers are cross-linked using one or more heterobifunctional cross-linkers, which can be used to link proteins in a stepwise manner. Heterobifunctional cross-linkers provide the ability to design more specific coupling methods for conjugating proteins, thereby reducing the occurrences of unwanted side reactions such as homo-protein polymers. A wide variety of heterobifunctional cross-linkers are known in the art. These include: succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), m-Maleimidobenzoyl-N-hydroxysuccinimide ester (MBS); N-succinimidyl (4-iodoacetyl) aminobenzoate (SIAB), succinimidyl 4-(p-maleimidophenyl) butyrate (SMPB), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC); 4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)-toluene (SMPT), N-succinimidyl 3-(2-pyridyldithio) propionate (SPDP), succinimidyl 6-((3-(2-pyridyldithio)propionate)hexanoate (LC-SPDP). Those cross-linking agents having N-hydroxysuccinimide moieties can be obtained as the N-hydroxysulfosuccinimide analogs, which generally have greater water solubility. In addition, those cross-linking agents having disulfide bridges within the linking chain can be synthesized instead as the alkyl derivatives so as to reduce the amount of linker cleavage in vivo.

[0056] In addition to the heterobifunctional cross-linkers, there exist a number of other cross-linking agents including homobifunctional and photoreactive cross-linkers. Disuccinimidyl suberate (DSS), bismaleimidohehexane (BMH) and dimethylpimelimidate-2 HCl (DMP) are examples of useful homobifunctional cross-linking agents, and bis-(β -(4-azidosalicylamido)ethyl)disulfide (BASED) and N-succinimidyl-6-(4'-azido-2'-nitrophenyl-amino)hexanoate (SANPAH) are examples of useful photoreactive cross-linkers for use in this invention. For a review of protein coupling techniques, see Means et al. (1990) Bioconjugate Chemistry 1:2-12.

[0057] One particularly useful class of heterobifunctional cross-linkers, included above, contain the primary amine reactive group, N-hydroxysuccinimide (NHS), or its water soluble analog N-hydroxysulfosuccinimide (sulfo-NHS). Primary amines (lysine epsilon groups) at alkaline pH's are unprotonated and react by nucleophilic attack on NHS or sulfo-NHS esters. This reaction results in the formation of an amide bond, and release of NHS or sulfo-NHS as a by-product.

[0058] Thiols are also particularly useful reactive groups as part of a heterobifunctional cross-linker. Common thiol reactive groups include maleimides, halogens, and pyridyl disulfides. Maleimides react specifically with free sulfhydryls (cysteine residues) in minutes, under slightly acidic to neutral (pH 6.5-7.5) conditions. Halogens (iodoacetyl functions) react with -SH groups at physiological pH's. Both of these reactive groups result in the formation of stable thioether

bonds.

[0059] A third component of the heterobifunctional cross-linker is the spacer arm or bridge. The bridge is the structure that connects the two reactive ends. The most apparent feature of the bridge is its effect on steric hindrance. In some instances, a longer bridge can more easily span the distance necessary to link two complex biomolecules.

[0060] Preparing protein-protein conjugates using heterobifunctional reagents is a two-step process involving the amine reaction and the sulfhydryl reaction, and such processes are generally well known in the art. See, e.g., Partis et al. (1983) J. Pro. Chem. 2:263; Ellman et al. (1958) Arch. Biochem. Biophys. 74:443; Riddles et al. (1979) Anal. Biochem. 94:75; Blattler et al. (1985) Biochem 24:1517).

[0061] In certain aspects, the disclosure provides a covalently crosslinked SAP oligomer comprising at least two SAP pentamers, wherein each of the SAP pentamers comprises five SAP protomers. In certain embodiments, SAP oligomers may be comprised of SAP protomers at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to the amino acid sequence of SEQ ID NO:1. Alternatively, SAP oligomers may comprise at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine or more of the variant SAP protomers described herein. In some embodiments, SAP oligomers may be comprised of at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine or more different variant SAP protomers as described herein. In preferred embodiments, a crosslinked SAP oligomer is characterized by one or more of increased plasma half-life, increased *in vitro* stability, and increased *in vivo* stability compared to a corresponding sample of SAP isolated from human serum. In certain embodiments, the crosslinked SAP oligomers have one or more of the following characteristics as compared to serum-derived human SAP: increased resistance to protease, increased resistance to calcium-mediated autoaggregation, and increased metal ion-binding.

Methods of Producing SAP Variants

[0062] In part, the disclosure provides methods for generating the SAP variants and SAP oligomers of the invention. SAP variants of the disclosure may comprise at least one protomer having one or more amino acid alterations that modify at least one biological activity of the SAP protein. As described herein, methods of generating amino acid alterations include, but are not limited to, mutating at least one amino acid of SEQ ID NO: 1 (e.g., deletion of one or more amino acids, addition of one or more amino acids, or substitution of one or more amino acids), chemically modifying one or more amino acids of SEQ ID NO: 1 (e.g., attaching one or more inert polypeptides to an amino acid of SEQ ID NO: 1), or a combination thereof.

[0063] In certain aspects, variant SAP protomers of the invention may be generated using random mutagenesis techniques, directed mutagenesis techniques, directed evolution, or

combination thereof. Variant SAP protomers may be generated using techniques that introduce random or directed mutations in the coding sequence of a nucleic acid. The nucleic acid is then expressed in a desired expression system, and the resulting peptide is assessed for properties of interest, e.g., resistance to autoaggregation, resistance to protease cleavage, increased metal-ion binding, increased serum half-life, increased *in vitro* half-life, increased *in vivo* half-life. Techniques to introduce random or directed mutations into DNA sequences are well known in the art, and include PCR mutagenesis, saturation mutagenesis, and degenerate oligonucleotide approaches. See Sambrook and Russell (2001, Molecular Cloning, A Laboratory Approach, Cold Spring Harbor Press, Cold Spring Harbor, N.Y.) and Ausubel et al. (2002, Current Protocols in Molecular Biology, John Wiley & Sons, NY).

[0064] In random PCR mutagenesis, reduced Taq polymerase fidelity is used to introduce random mutations into a cloned fragment of DNA (Leung et al., 1989, Technique 1:11 15). The DNA region to be mutagenized is amplified using PCR under conditions that reduce the fidelity of DNA synthesis by Taq DNA polymerase, e.g., by using an altered dGTP/dATP ratio and by adding Mn^{2+} to the PCR reaction. The pool of amplified DNA fragments are inserted into appropriate cloning vectors to provide random mutant libraries.

[0065] Saturation mutagenesis allows for the rapid introduction of a large number of single base substitutions into cloned DNA fragments (Mayers et al., 1985, Science 229:242). This technique includes generation of mutations, e.g., by chemical treatment or irradiation of single-stranded DNA *in vitro* and synthesis of a complementary DNA strand. The mutation frequency can be modulated by modulating the severity of the treatment, and essentially all possible base substitutions can be obtained. Because this procedure does not involve a genetic selection for mutant fragments, both neutral substitutions as well as those that alter function, are obtained. Furthermore, the distribution of point mutations is not biased toward conserved sequence elements.

[0066] A library of nucleic acid homologs can also be generated from a set of degenerate oligonucleotide sequences. Chemical synthesis of a degenerate oligonucleotide sequences can be carried out in an automatic DNA synthesizer, and the synthetic genes may then be ligated into an appropriate expression vector. The synthesis of degenerate oligonucleotides is known in the art (see for example, Narang, S A (1983) Tetrahedron 39:3; Itakura et al. (1981) Recombinant DNA, Proc 3rd Cleveland Sympos. Macromolecules, ed. A G Walton, Amsterdam: Elsevier pp. 273 289; Itakura et al. (1984) Annu. Rev. Biochem. 53:323; Itakura et al. (1984) Science 198:1056; Ike et al. (1983) Nucleic Acid Res. 11:477. Such techniques have been employed in the directed evolution of other peptides (see, for example, Scott et al. (1990) Science 249:386 390; Roberts et al. (1992) PNAS 89:2429 2433; Devlin et al. (1990) Science 249: 404 406; Cwirla et al. (1990) PNAS 87: 6378 6382; as well as U.S. Pat. Nos. 5,223,409, 5,198,346, and 5,096,815).

[0067] Variant SAP protomers may also be generated using "directed evolution" techniques. These strategies are different from traditional random mutagenesis procedures because they involve subjecting the nucleic acid sequence encoding the peptide of interest to recursive

rounds of mutation, screening and amplification.

[0068] In some "directed evolution" techniques, the diversity in the nucleic acids obtained is generated by mutation methods that randomly create point mutations in the nucleic acid sequence. The point mutation techniques include, but are not limited to, "error-prone PCR™" (Caldwell and Joyce, 1994; PCR Methods Appl. 2: 28 33; and Ke and Madison, 1997, Nucleic Acids Res. 25: 3371 3372), repeated oligonucleotide-directed mutagenesis (Reidhaar-Olson et al., 1991, Methods Enzymol. 208:564 586), and any of the aforementioned methods of random mutagenesis.

[0069] Another method of creating diversity upon which directed evolution can act is the use of mutator genes. The nucleic acid of interest is cultured in a mutator cell strain the genome of which typically encodes defective DNA repair genes (U.S. Pat. No. 6,365,410; Selifonova et al., 2001, Appl. Environ. Microbiol. 67:3645 3649; Long-McGie et al., 2000, Biotech. Bioeng. 68:121 125; see, Genencor International Inc, Palo Alto Calif.).

[0070] Achieving diversity using directed evolution techniques may also be accomplished using saturation mutagenesis along with degenerate primers (Gene Site Saturation Mutagenesis™, Diversa Corp., San Diego, Calif.). In this type of saturation mutagenesis, degenerate primers designed to cover the length of the nucleic acid sequence to be diversified are used to prime the polymerase in PCR reactions. In this manner, each codon of a coding sequence for an amino acid may be mutated to encode each of the remaining common nineteen amino acids. This technique may also be used to introduce mutations, deletions and insertions to specific regions of a nucleic acid coding sequence while leaving the rest of the nucleic acid molecule untouched. Procedures for the gene saturation technique are well known in the art, and can be found in U.S. Pat. No. 6,171,820.

[0071] Variant SAP protomers may also be generated using the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling techniques are may be employed to modulate the activities of peptides useful in the invention and may be used to generate peptides having altered activity. See, generally, U.S. Pat. Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Stemmer et al. (1994, Nature 370(6488):389 391); Cramer et al. (1998, Nature 391 (6664):288 291); Zhang et al. (1997, Proc. Natl. Acad. Sci. USA 94(9):4504 4509); Stemmer et al. (1994, Proc. Natl. Acad. Sci USA 91(22):10747 10751), Patten et al. (1997, Curr. Opinion Biotechnol. 8:724 33); Harayama, (1998, Trends Biotechnol. 16(2):76 82); Hansson, et al., (1999, J. Mol. Biol. 287:265 76); and Lorenzo and Blasco (1998, Biotechniques 24(2):308 13).

[0072] DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. DNA shuffling has been used to generate novel variations of human immunodeficiency virus type 1 proteins (Pekrun et al., 2002, J. Virol. 76(6):2924 35), triazine hydrolases (Raillard et al. 2001, Chem Biol 8(9):891 898), murine leukemia virus (MLV) proteins (Powell et al. 2000, Nat Biotechnol 18(12):1279 1282), and indoleglycerol phosphate synthase (Merz et al. 2000, Biochemistry

39(5):880 889).

[0073] The technique of DNA shuffling was developed to generate biomolecular diversity by mimicking natural recombination by allowing *in vitro* homologous recombination of DNA (Stemmler, 1994, Nature 370: 389 391; and Stemmler, 1994, PNAS 91: 10747 10751). Generally, in this method a population of related genes is fragmented and subjected to recursive cycles of denaturation, rehybridization, followed by the extension of the 5' overhangs by Taq polymerase. With each cycle, the length of the fragments increases, and DNA recombination occurs when fragments originating from different genes hybridize to each other. The initial fragmentation of the DNA is usually accomplished by nuclease digestion, typically using DNase (see Stemmler references, above), but may also be accomplished by interrupted PCR synthesis (U.S. Pat. No. 5,965,408, incorporated herein by reference in its entirety; see, Diversa Corp., San Diego, Calif.). DNA shuffling methods have advantages over random point mutation methods in that direct recombination of beneficial mutations generated by each round of shuffling is achieved and there is therefore a self-selection for improved phenotypes of peptides. The techniques of DNA shuffling are well known to those in art. Detailed explanations of such technology is found in Stemmler, 1994, Nature 370: 389 391 and Stemmler, 1994, PNAS 91: 10747 10751. The DNA shuffling technique is also described in U.S. Pat. Nos. 6,180,406, 6,165,793, 6,132,970, 6,117,679, 6,096,548, 5,837,458, 5,834,252, 5,830,721, 5,811,238, and 5,605,793.

[0074] The art also provides even more recent modifications of the basic technique of DNA shuffling. In one example, exon shuffling, exons or combinations of exons that encode specific domains of peptides are amplified using chimeric oligonucleotides. The amplified molecules are then recombined by self-priming PCR assembly (Kolkman and Stemmler, 2001, Nat. Biotech. 19:423 428). In another example, using the technique of random chimeragenesis on transient templates (RACHITT) library construction, single stranded parental DNA fragments are annealed onto a full-length single-stranded template (Coco et al., 2001, Nat. Biotechnol. 19:354 359). In yet another example, staggered extension process (StEP), thermocycling with very abbreviated annealing/extension cycles is employed to repeatedly interrupt DNA polymerization from flanking primers (Zhao et al., 1998, Nat. Biotechnol. 16: 258 261). In the technique known as CLERY, *in vitro* family shuffling is combined with *in vivo* homologous recombination in yeast (Abecassis et al., 2000, Nucleic Acids Res. 28:E88). To maximize intergenic recombination, single stranded DNA from complementary strands of each of the nucleic acids are digested with DNase and annealed (Kikuchi et al., 2000, Gene 243:133 137). The blunt ends of two truncated nucleic acids of variable lengths that are linked by a cleavable sequence are then ligated to generate gene fusion without homologous recombination (Sieber et al., 2001, Nat Biotechnol. 19:456 460; Lutz et al., 2001, Nucleic Acids Res. 29:E16; Ostermeier et al., 1999, Nat. Biotechnol. 17:1205 1209; Lutz and Benkovic, 2000, Curr. Opin. Biotechnol. 11:319 324). Recombination between nucleic acids with little sequence homology in common has also been enhanced using exonuclease-mediated blunt-ending of DNA fragments and ligating the fragments together to recombine them (U.S. Pat. No. 6,361,974).

[0075] In addition to published protocols detailing directed evolution and gene shuffling

techniques, commercial services are now available that will undertake the gene shuffling and selection procedures on peptides of choice. Maxygen (Redwood City, Calif.) offers commercial services to generate custom DNA shuffled libraries. In addition, this company will perform customized directed evolution procedures including gene shuffling and selection on a peptide family of choice.

[0076] Optigenix, Inc. (Newark, Del.) offers the related service of plasmid shuffling. Optigenix uses families of genes to obtain mutants therein having new properties. The nucleic acid of interest is cloned into a plasmid in an *Aspergillus* expression system. The DNA of the related family is then introduced into the expression system and recombination in conserved regions of the family occurs in the host. Resulting mutant DNAs are then expressed and the peptide produced therefrom are screened for the presence of desired properties and the absence of undesired properties.

[0077] Following each recursive round of "evolution," the desired peptides expressed by mutated genes are screened for characteristics of interest. The "candidate" genes are then amplified and pooled for the next round of DNA shuffling. The screening procedure used is highly dependant on the peptide that is being "evolved" and the characteristic of interest. Characteristics such as peptide stability, biological activity, antigenicity, among others can be selected using procedures that are well known in the art.

[0078] It will be appreciated by the skilled artisan that the above techniques of mutation and selection can be combined with each other and with additional procedures to generate the best possible variant SAP protomer useful in the methods of the invention. Thus, the invention is not limited to any one method for the generation of SAP variants, and should be construed to encompass any and all of the methodology described herein. For example, a procedure for introducing specified point mutations into a nucleic acid sequence may be performed initially, followed by recursive rounds of DNA shuffling, selection and amplification. For some variants the initial introduction of point mutations may be used to introduce diversity into a gene population where it is lacking, and the following round of DNA shuffling and screening will select for advantageous point mutations.

[0079] In certain aspects, the disclosure provides methods for chemically modifying one or more amino acids of an SAP protomer. There are a number of methods described in the art for attaching inert polymers to a polypeptide including, but not limited to, using cyanogen bromide (alkylation) and dialdehyde coupling chemistry and periodate oxidation. In particular, many methods for pegylating amino acids have been described in the art. For example, U.S. Pat. No. 4,088,538 discloses an enzymatically active polymer-enzyme conjugate of an enzyme covalently bound to PEG. Similarly, U.S. Pat. No. 4,496,689 discloses a covalently attached complex of α -1 protease inhibitor with a polymer such as PEG or methoxypoly(ethylene glycol) ("mPEG"). Abuchowski et al. (J. Biol. Chem. 252: 3578 (1977) discloses the covalent attachment of mPEG to an amine group of bovine serum albumin. U.S. Pat. No. 4,414,147 discloses a method of rendering interferon less hydrophobic by conjugating it to an anhydride of a dicarboxylic acid, such as poly(ethylene succinic anhydride). PCT WO 87/00056 discloses

conjugation of PEG and polyoxyethylated polyols to such proteins as interferon- β , interleukin-2 and immunotoxins. EP 154,316 discloses and claims chemically modified lymphokines, such as IL-2 containing PEG bonded directly to at least one primary amino group of the lymphokine. U.S. Pat. No. 4,055,635 discloses pharmaceutical compositions of a water-soluble complex of a proteolytic enzyme linked covalently to a polymeric substance such as a polysaccharide. Another mode of attaching PEG to peptides is through the nonspecific oxidation of glycosyl residues on a peptide. The oxidized sugar is utilized as a locus for attaching a PEG moiety to the peptide. For example WO 94/05332 discloses the use of a hydrazine- or amino-PEG to add PEG to a glycoprotein. The glycosyl moieties are randomly oxidized to the corresponding aldehydes, which are subsequently coupled to the amino-PEG.

[0080] The disclosure also provides SAP variants comprising site-specific pegylation. In some embodiments, an SAP protomer is modified by the introduction of a "free" cysteine residues (i.e., cysteines that are not involved in disulfide bonds) to which PEG can be attached using well described malimide chemistry. (See, e.g., Natarajan, *Bioconjug Chem.* 2005 Jan-Feb;16(1):113-21; Goodson, *Biotechnology NY.* 1990 Apr;8(4):343-6). Modified SAP variants are provided, wherein polymer conjugation sites are introduced via variant cysteine residues. The cysteine residue may be substituted for one or more native SAP amino acid residues or by adding one or more cysteines to an SAP polypeptide. In some embodiments, a cysteine residue is introduced at position -1 of SEQ ID NO: 1 (i.e., added to the N-terminus of the polypeptide). In some embodiments, a cysteine residue is introduced by the substitution of the native amino acid at position 32 of SEQ ID NO: 1 for a cysteine residue. In some embodiments, the introduced cysteine is pegylated.

[0081] For pegylation of cysteine residues, the polypeptide may be treated with a reducing agent, such as dithiothreitol (DDT) prior to pegylation. The reducing agent is subsequently removed by any conventional method, such as by desalting. Conjugation of PEG to a cysteine residue typically takes place in a suitable buffer at pH 6-9 at temperatures varying from 4°C to 25°C for periods up to about 16 hours. Examples of activated PEG polymers for coupling to cysteine residues include the following linear and branched PEGs, including but not limited to, vinylsulfone-PEG (PEG-VS), such as vinylsulfone-mPEG (mPEG-VS); orthopyridyl-disulfide-PEG (PEG-OPSS), such as orthopyridyl-disulfide-mPEG (mPEG-OPSS); and maleimide-PEG (PEG-MAL), such as maleimide-mPEG (mPEG-MAL) and branched maleimide-mPEG2 (mPEG2-MAL).

[0082] One approach for adding PEG or dextran to SAP utilizes the enzyme transglutaminase (glutamyl-peptide γ -glutamyltransferase; EC 2.3.2.13). This enzyme catalyzes the calcium-dependent acyl addition to a primary amine wherein the gamma-carboxamide group of peptide-bound glutamine residue is the acyl donor and the primary amine is the acyl acceptor and amine donor. A transglutaminase reaction is therefore employed to covalently and site-specifically conjugate SAP to a polymer, such as PEG or dextran through a Gln residue that is capable of acting as a transglutaminase amine acceptor.

[0083] The transglutaminase amine acceptor in SAP may be a native or introduced (i.e.,

variant) Gln residue. In general, glutamine repeats have been shown to enhance the acceptor properties of each glutamine residue in the repeat, and the accessibility of glutamine residues has also been shown to be important in determining their ability to function as transglutaminase substrates (Kahlem, P. et al. Proc. Natl. Acad. Sci. USA 1996, 93, 14580-14585). In some embodiments, the SAP variant comprises an N32Q mutation, introducing a transglutaminase amine acceptor. In some embodiments, the SAP variant comprises the amino acid sequence at least 70, 80, 85, 90, 95, 98, or 100% identical to SEQ ID NO: 2 (wherein X is any amino acid, A is 3 to 20, and γ is 1 to 10):

X_AQ _{γ} HTDLSGKVFVFPRESVTDHVNLI~~TPLEKPLQNFTLC~~FRAYS~~DL~~S
 RAYSLSFSYNTQGRDNELLVYKERVGEYSLYIGRHKVT~~SKVIEKFPAP~~
 VHICVSWESSSGIAEFWINGTPLVKKGLRQGYFVEAQPKIVLGQE~~QD~~
 SYGGKFDRSQSFVGEIGDLYMWDSVLPENILSAYQGTPLPANILDW
 QALNYEIRGYVIKPLVWV (SEQ ID NO: 2)

In some embodiments, γ is 1. In some embodiments, γ is 2. In some embodiments, the SAP variant is conjugated to PEG via a transglutaminase amine acceptor. In some embodiments, the SAP variant is conjugated to dextran via a transglutaminase amine acceptor.

[0084] Methods for adding polymers to Gln residues have been described in the art (See, e.g., U.S. Publication No. 20060116322, Sugimura et al. 281 (26): 17699. (2006); Sato, H. Adv Drug Deliv Rev. 2002 Jun 17;54(4):487-504; Sato, et al, Bioconjug Chem. 2000 Jul-Aug;11(4):502-9; Sato, et al Bioconjug Chem. 2001 Sep-Oct;12(5):701-10, Fontana, et al Adv Drug Deliv Rev. 2008 Jan 3;60(1):13-28. Epub 2007 Aug 16, Hohenadl, J Biol Chem. 1995 Oct 6;270(40):23415-20)). The polymers are linked or modified to contain a primary amine which will act as the transglutaminase amine donor.

[0085] SAP variants and SAP covalently crosslinked oligomers described herein may be produced in bacterial cells, insect cells, yeast, fungal cells, or mammalian cells including, for example, human cells. In those instances when the host cell is human, the cell may be in a live subject or may be isolated from a subject, e.g., in a cell culture, tissue sample, cell suspension, etc. Other suitable host cells are known to those skilled in the art.

[0086] The disclosure further provides expression vectors for producing SAP protomers. For instance, expression vectors are contemplated which include a nucleotide sequence encoding an SAP protomer, wherein the coding sequence is operably linked to at least one transcriptional regulatory sequence. Regulatory sequences for directing expression of SAP protomers are art-recognized and are selected by a number of well understood criteria. Exemplary regulatory sequences are described in Goeddel; Gene Expression Technology: Methods in Enzymology, Academic Press, San Diego, CA (1990). For instance, any of a wide variety of regulatory sequences that control the expression of a DNA sequence when operatively linked to it may be used in these vectors to express DNA sequences encoding SAP protomers. Such useful expression control sequences, include, for example, the early and late promoters of SV40, adenovirus or cytomegalovirus immediate early promoter, the *lac* system, the *trp* system, the TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the

promoters of acid phosphatase, e.g., Pho5, and the promoters of the yeast α -mating factors and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the target host cell to be transformed. Moreover, the vector's copy number, the ability to control that copy number and the expression of any other protein encoded by the vector, such as antibiotic markers, should also be considered.

[0087] The disclosure also provides a host cell transfected with a recombinant gene in order to express an SAP protomer. The host cell may be any prokaryotic or eukaryotic cell. For example, an SAP protomer may be expressed in bacterial cells such as *E. coli*, insect cells, yeast, or mammalian cells. Other suitable host cells are known to those skilled in the art.

[0088] Accordingly, the disclosure provides methods of producing SAP protomers. For example, a host cell transfected with an expression vector encoding an SAP protomer of the invention can be cultured under appropriate conditions to allow expression of the polypeptide to occur. The SAP protomer may be secreted, by inclusion of a secretion signal sequence, and isolated from a mixture of cells and medium containing the protein. Alternatively, the SAP protomer may be retained cytoplasmically and the cells harvested, lysed and the protomer isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. The proteins can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins, including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies specific for particular epitopes of the protein.

[0089] Thus, a coding sequence for an SAP protomer can be used to produce a recombinant form of the protein via microbial or eukaryotic cellular processes. Ligating the polynucleotide sequence into a gene construct, such as an expression vector, and transforming or transfecting into hosts, either eukaryotic (yeast, avian, insect or mammalian) or prokaryotic (bacterial cells), are standard procedures.

[0090] Expression vehicles for production of a recombinant protein include plasmids and other vectors. For instance, suitable vectors for the expression of SAP protomers include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

[0091] A number of vectors exist for the expression of recombinant proteins in yeast. For instance, YEp24, YIp5, YEp51, YEp52, pYES2, and YRp17 are cloning and expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae* (see, for example, Broach et al., (1983) in *Experimental Manipulation of Gene Expression*, ed. M. Inouye Academic Press, p. 83). These vectors can replicate in *E. coli* due the presence of the pBR322 ori, and in *S. cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. Autotrophic selection or counterselection is often used in yeast. In addition, drug resistance markers, such as ampicillin, can be used in bacteria.

[0092] Mammalian expression vectors may contain both prokaryotic sequences to facilitate the propagation of the vector in bacteria, and one or more eukaryotic transcription units that are expressed in eukaryotic cells. The pcDNAI/amp, pcDNAI/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papilloma virus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells. Examples of other viral (including retroviral) expression systems can be found below in the description of gene therapy delivery systems. The various methods employed in the preparation of the plasmids and transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see *Molecular Cloning: A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press, 1989) Chapters 16 and 17. In some instances, it may be desirable to express the recombinant SAP polypeptides by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the beta-gal containing pBlueBac III).

[0093] In some instances it will be desired to produce SAP protomers in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen Virol.* 36:59. (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; a human hepatoma line (Hep G2); and myeloma or lymphoma cells (e.g. Y0, J558L, P3 and NS0 cells) (see U.S. Pat. No. 5,807,715).

[0094] In certain embodiments, production of SAP protomers may be achieved using *in vitro* translation systems. *In vitro* translation systems are, generally, a translation system which is a cell-free extract containing at least the minimum elements necessary for translation of an RNA molecule into a protein. An *in vitro* translation system typically comprises at least ribosomes, tRNAs, initiator methionyl-tRNAMet, proteins or complexes involved in translation, e.g., eIF2, eIF3, the cap-binding (CB) complex, comprising the cap-binding protein (CBP) and eukaryotic initiation factor 4F (eIF4F). A variety of *in vitro* translation systems are well known in the art

and include commercially available kits. Examples of *in vitro* translation systems include eukaryotic lysates, such as rabbit reticulocyte lysates, rabbit oocyte lysates, human cell lysates, insect cell lysates and wheat germ extracts. Lysates are commercially available from manufacturers such as Promega Corp., Madison, Wis.; Stratagene, La Jolla, Calif.; Amersham, Arlington Heights, Ill.; and GIBCO/BRL, Grand Island, N.Y. *In vitro* translation systems typically comprise macromolecules, such as enzymes, translation, initiation and elongation factors, chemical reagents, and ribosomes. In addition, an *in vitro* transcription system may be used. Such systems typically comprise at least an RNA polymerase holoenzyme, ribonucleotides and any necessary transcription initiation, elongation and termination factors. *In vitro* transcription and translation may be coupled in a one-pot reaction to produce proteins from one or more isolated DNAs.

[0095] Often, it is difficult to produce large quantities of a protein with reproducible consistency in the characteristics of the product, such as post-translational modification and/or folding. In some embodiments, SAP variants of the disclosure are characterized by increased efficiency of manufacturing the SAP protein (e.g., greater yield of the protein product, increased homogeneity of the protein product, increased stability of the protein product), particularly for *in vivo* use (e.g., as a therapeutic agent). In some embodiments, SAP variants of the disclosure are characterized by increased stability and/or homogeneity when expressed in a cell (e.g., prokaryote, eukaryote) compared to wild-type SAP expressed in the same cell line. For example, recombinant proteins are generally characterized by a high degree of heterogeneity with regards to their attached glycan structures. SAP variants having a modified amino acid sequence to inhibit glycosylation (such as an amino acid substitution at position 32 of SEQ ID NO: 1, e.g., N32D) would be expected to be more homogeneously produced from a cell.

[0096] In exemplary embodiments, SAP variants, may be purified, for example, to at 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% purity, or greater, with respect to contaminating macromolecules, particularly other proteins and nucleic acids, and free of infectious and pyrogenic agents. SAP variants may be substantially free of other polypeptides, particularly other polypeptides of animal origin.

[0097] SAP variants can be purified using fractionation and/or conventional purification methods and media. Ammonium sulfate precipitation and acid or chaotrope extraction may be used for fractionation of samples. Exemplary purification steps may include hydroxyapatite, size exclusion, FPLC and reverse-phase high performance liquid chromatography. Suitable anion exchange media include derivatized dextrans, agarose, cellulose, polyacrylamide, specialty silicas, and the like. PEI, DEAE, QAE and Q derivatives are suitable, including, for example, DEAE Fast-Flow Sepharose (Pharmacia, Piscataway, NJ). Exemplary chromatographic media include those media derivatized with phenyl, butyl, or octyl groups, such as Phenyl-Sepharose FF (Pharmacia), Toyopearl butyl 650 (Toso Haas, Montgomeryville, PA), Octyl-Sepharose (Pharmacia) and the like; or polyacrylic resins, such as Amberchrom CG 71 (Toso Haas) and the like. Suitable solid supports include glass beads, silica-based resins, cellulosic resins, agarose beads, cross-linked agarose beads, polystyrene beads, cross-linked polyacrylamide resins and the like that are insoluble under the conditions in which they are to

be used. These supports may be modified with reactive groups that allow attachment of proteins by amino groups, carboxyl groups, sulfhydryl groups, hydroxyl groups and/or carbohydrate moieties. Examples of coupling chemistries include cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, hydrazide activation, and carboxyl and amino derivatives for carbodiimide coupling chemistries. These and other solid media are well known and widely used in the art, and are available from commercial suppliers. Methods for binding polypeptides to support media are well known in the art. Selection of a particular method is a matter of routine design and is determined in part by the properties of the chosen support. See, for example, *Affinity Chromatography: Principles & Methods* (Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988). The SAP variants described herein can also be isolated an affinity tag (e.g., polyhistidine, maltose-binding protein, GST, starch binding domain, FLAG, an immunoglobulin domain) to facilitate purification as described further herein.

Treatment methods

[0098] In certain aspects, the disclosure provides methods for treating an SAP-responsive disorder in a patient by administering a therapeutically effective amount of an SAP variant or SAP oligomer of the invention to a patient in need thereof. The dosage and frequency of treatment can be determined by one skilled in the art and will vary depending on the symptoms, age and body weight of the patient, and the nature and severity of the disorder to be treated or prevented. In some embodiments, an SAP variant or SAP oligomer is administered to a patient once or twice per day, once or twice per week, once or twice per month, or just prior to or at the onset of symptoms.

[0099] Dosages may be readily determined by techniques known to those of skill in the art or as taught herein. Toxicity and therapeutic efficacy of SAP may be determined by standard pharmaceutical procedures in experimental animals, for example, determining the LD₅₀ and the ED₅₀. The ED₅₀ (Effective Dose 50) is the amount of drug required to produce a specified effect in 50% of an animal population. The LD₅₀ (Lethal Dose 50) is the dose of drug which kills 50% of a sample population.

[0100] In some embodiments, the SAP-responsive disorder is fibrosis. The use of SAP as a therapeutic treatment for fibrosis is described in U.S. Patent Application No. 2007/0243163. Fibrosis related disorders that may be amenable to treatment with the subject method include, but are not limited to, collagen disease, interstitial lung disease, human fibrotic lung disease (e.g., obliterative bronchiolitis, idiopathic pulmonary fibrosis, pulmonary fibrosis from a known etiology, tumor stroma in lung disease, systemic sclerosis affecting the lungs, Hermansky-Pudlak syndrome, coal worker's pneumoconiosis, asbestosis, silicosis, chronic pulmonary hypertension, AIDS-associated pulmonary hypertension, sarcoidosis, moderate to severe asthma and the like), fibrotic vascular disease, arterial sclerosis, atherosclerosis, varicose veins, coronary infarcts, cerebral infarcts, myocardial fibrosis, musculoskeletal fibrosis, post-

surgical adhesions, human kidney disease (e.g., nephritic syndrome, Alport syndrome, HIV-associated nephropathy, polycystic kidney disease, Fabry's disease, diabetic nephropathy, chronic glomerulonephritis, nephritis associated with systemic lupus, and the like), progressive systemic sclerosis (PSS), primary sclerosing cholangitis (PSC), liver fibrosis, liver cirrhosis, renal fibrosis, pulmonary fibrosis, cystic fibrosis, chronic graft versus host disease, scleroderma (local and systemic), Grave's ophthalmopathy, diabetic retinopathy, glaucoma, Peyronie's disease, penis fibrosis, urethrostenosis after cystoscopy, inner accretion after surgery, scarring, myelofibrosis, idiopathic retroperitoneal fibrosis, peritoneal fibrosis from a known etiology, drug-induced ergotism, fibrosis incident to benign or malignant cancer, fibrosis incident to microbial infection (e.g., viral, bacterial, parasitic, fungal, etc.), Alzheimer's disease, fibrosis incident to inflammatory bowel disease (including stricture formation in Crohn's disease and microscopic colitis), stromal cell tumors, mucositis, fibrosis induced by chemical or environmental insult (e.g., cancer chemotherapy, pesticides, or radiation (e.g., cancer radiotherapy)).

[0101] In some embodiments, the fibrosis related disorder is selected from systemic or local scleroderma, keloids, hypertrophic scars, atherosclerosis, restenosis, pulmonary inflammation and fibrosis, idiopathic pulmonary fibrosis, liver cirrhosis, fibrosis as a result of chronic hepatitis B or C infection, kidney disease, heart disease resulting from scar tissue, macular degeneration, and retinal and vitreal retinopathy. In some embodiments, the fibrosis related disorder results from chemotherapeutic drugs, radiation-induced fibrosis, and injuries and burns. In some embodiments, the fibrosis related disorder or condition results from post-surgical scarring, e.g., following trabeculectomy or other filtration surgery of the eye.

[0102] In some embodiments, the SAP-responsive disorder is a hypersensitivity disorder such as those mediated by Th1 or Th2 responses. The use of SAP as a therapeutic treatment for hypersensitivity disorders is also described in U.S. Provisional Application No 61/209,795. Hypersensitivity-related disorders that may be amenable to treatment with SAP include, but are not limited to, allergic rhinitis, allergic sinusitis, allergic conjunctivitis, allergic bronchoconstriction, allergic dyspnea, allergic increase in mucus production in the lungs, atopic eczema, dermatitis, urticaria, anaphylaxis, pneumonitis, and allergic-asthma.

[0103] In some embodiments, an SAP variant or SAP oligomer of the invention may be used to treat allergen-specific immune responses, such as anaphylaxis, to various antigens, including, but not limited to, antimicrobials (e.g., cephalosporins, sulfonamides, penicillin and other β -lactams), anticonvulsants (e.g., phenytoin, phenobarital, carbamazepine, dapsone, allopurinol, and minocycline), chemotherapeutics (e.g., taxanes, platinum compounds, asparaginases, and epipodophyllotoxins), heparin, insulin, protamine, aspirin and other non-steroidal anti-inflammatory drugs, muscle relaxants (e.g., succinylcholine, atracurium, vecuronium, and pancuronium), induction agents (e.g., barbiturates, etomidate, propofol), narcotics (e.g., fentanyl, meperidine, morphine), colloids for intravascular volume expansion, radiocontrast materials, blood products, latex, animal products, animal dander, dust mites, insects (e.g., bites, stings or venom), cosmetics, metals (e.g., nickel, cobalt, and chromate), plants, spores, pollen, food (e.g., milk, eggs, wheat, soy, peanuts and tree nuts, seafood), vaccination, and

fungal antigens (e.g., *Aspergillus*, *Curvularia*, *Exserohilum*, and *Alternaria* species).

[0104] In some embodiments, the SAP-responsive disorder is an autoimmune disorder such as those mediated by Th1 or Th2 responses. The use of SAP as a therapeutic treatment for autoimmune disorders is also described in U.S. Provisional Application No 61/209,845. Autoimmune related disorders that may be amenable to treatment with SAP include, but are not limited to, type I diabetes, multiple sclerosis, rheumatoid arthritis, psoriatic arthritis, autoimmune myocarditis, pemphigus, myasthenia gravis, Hashimoto's thyroiditis, Graves' disease, Addison's disease, autoimmune hepatitis, chronic Lyme arthritis, familial dilated cardiomyopathy, juvenile dermatomyositis, polychondritis, Sjogren's syndrome, psoriasis, juvenile idiopathic arthritis, inflammatory bowel disease, systemic lupus erythematosus, chronic obstructive pulmonary disease, and graft-versus-host disease.

[0105] In some embodiments, the SAP-responsive disorder is a mucositis. The use of SAP as a therapeutic treatment for mucositis is also described in U.S. Application No 12/217,614. Methods described herein may be useful for treating oral, esophageal, and gastrointestinal mucositis, as well as gastric and duodenal ulcers, or erosions of the stomach and esophagus.

[0106] In some embodiments, an SAP variant or SAP oligomer may be used to treat an inflammatory disease. In some embodiments, the inflammatory disease may be a viral, bacterial, fungal, or parasitic infection. The use of SAP as a therapeutic treatment for viral infection has also been described in U.S. Patent 6,406,698 and in PCT Application WO1997/026906.

Pharmaceutical Preparations and Formulations

[0107] In certain aspects, the disclosure provides pharmaceutical preparations comprising one or more SAP therapeutic agents (i.e., SAP variants and SAP oligomers) formulated for administration. The therapeutic agents of the invention may be formulated in a conventional manner using one or more physiologically acceptable carriers or excipients. For example, therapeutic agents and their physiologically acceptable salts and solvates may be formulated for administration by, for example, injection (e.g. SubQ, IM, IP), inhalation or insufflation (either through the mouth or the nose) or oral, buccal, sublingual, transdermal, nasal, parenteral or rectal administration. In certain embodiments, therapeutic agents may be administered locally, at the site where the target cells are present, i.e., in a specific tissue, organ, or fluid (e.g., blood, cerebrospinal fluid, tumor mass, etc.).

[0108] The present invention further provides use of any SAP variant or SAP oligomer of the invention in the manufacture of a medicament for the treatment or prevention of a disorder or a condition, as described herein, in a patient, for example, the use of an SAP variant or SAP oligomer in the manufacture of medicament for the treatment of a disorder or condition described herein. In some aspects, any SAP variant or SAP oligomer of the invention may be used to make a pharmaceutical preparation for the use in treating or preventing a disease or

condition described herein.

[0109] Therapeutic agents can be formulated for a variety of modes of administration, including systemic and topical or localized administration. Techniques and formulations generally may be found in Remington's Pharmaceutical Sciences, Meade Publishing Co., Easton, PA. For parenteral administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous. For injection, the compounds can be formulated in liquid solutions, preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the compounds may be formulated in solid form and redissolved or suspended immediately prior to use. Lyophilized forms are also included. In some embodiments, the therapeutic agents can be administered to cells by a variety of methods known to those familiar in the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres.

[0110] For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges, or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

[0111] For administration by inhalation (e.g., pulmonary delivery), therapeutic agents may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of gelatin, for example, for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0112] In the methods of the invention, the pharmaceutical compounds can also be administered by intranasal or intrabronchial routes including insufflation, powders, and aerosol

formulations (for examples of steroid inhalants, see Rohatagi (1995) J. Clin. Pharmacol. 35:1187-1193; Tjwa (1995) Ann. Allergy Asthma Immunol. 75:107-111). For example, aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations such as in a nebulizer or an atomizer. Typically, such administration is in an aqueous pharmacologically acceptable buffer.

[0113] Pharmaceutical compositions suitable for respiratory delivery (e.g., intranasal, inhalation, etc.) of variant SAP polypeptides may be prepared in either solid or liquid form.

[0114] SAP variants or SAP oligomers of the invention may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[0115] In addition, SAP variants or SAP oligomers of the invention may also be formulated as a depot preparation. Such long-acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, therapeutic agents may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt. Controlled release formula also includes patches.

[0116] In certain embodiments, the compounds described herein can be formulated for delivery to the central nervous system (CNS) (reviewed in Begley, Pharmacology & Therapeutics 104: 29-45 (2004)). Conventional approaches for drug delivery to the CNS include: neurosurgical strategies (e.g., intracerebral injection or intracerebroventricular infusion), molecular manipulation of the agent (e.g., production of a chimeric fusion protein that comprises a transport peptide that has an affinity for an endothelial cell surface molecule in combination with an agent that is itself incapable of crossing the blood-brain-barrier in an attempt to exploit one of the endogenous transport pathways of the blood-brain-barrier); pharmacological strategies designed to increase the lipid solubility of an agent (e.g., conjugation of water-soluble agents to lipid or cholesterol carriers); and the transitory disruption of the integrity of the BBB by hyperosmotic disruption (resulting from the infusion of a mannitol solution into the carotid artery or the use of a biologically active agent such as an angiotensin peptide).

[0117] In certain embodiments, SAP variants or SAP oligomers of the invention are incorporated into a topical formulation containing a topical carrier that is generally suited to topical drug administration and comprising any such material known in the art. The topical

carrier may be selected so as to provide the composition in the desired form, e.g., as an ointment, lotion, cream, microemulsion, gel, oil, solution, or the like, and may be comprised of a material of either naturally occurring or synthetic origin. It is preferable that the selected carrier not adversely affect the active agent or other components of the topical formulation. Examples of suitable topical carriers for use herein include water, alcohols and other nontoxic organic solvents, glycerin, mineral oil, silicone, petroleum jelly, lanolin, fatty acids, vegetable oils, parabens, waxes, and the like.

[0118] Pharmaceutical compositions (including cosmetic preparations) may comprise from about 0.00001 to 100% such as from 0.001 to 10% or from 0.1% to 5% by weight of one or more of the SAP variants or SAP oligomers described herein. In certain topical formulations, the active agent is present in an amount in the range of approximately 0.25 wt. % to 75 wt. % of the formulation, preferably in the range of approximately 0.25 wt. % to 30 wt. % of the formulation, more preferably in the range of approximately 0.5 wt. % to 15 wt. % of the formulation, and most preferably in the range of approximately 1.0 wt. % to 10 wt. % of the formulation.

[0119] Conditions of the eye can be treated or prevented by, e.g., systemic, topical, intraocular injection of therapeutic agents, or by insertion of a sustained release device that releases therapeutic agents. SAP variants or SAP oligomers of the invention may be delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye, as for example the anterior chamber, conjunctiva, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an encapsulating material. Alternatively, the compounds may be injected directly into the vitreous and aqueous humour. In a further alternative, the compounds may be administered systemically, such as by intravenous infusion or injection, for treatment of the eye.

[0120] Therapeutic agents described herein may be stored in oxygen-free environment according to methods in the art.

[0121] Exemplary compositions comprise an SAP variant or SAP oligomer with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic ingredients. The carrier(s) must be "pharmaceutically acceptable" in the sense of being compatible with the other ingredients of the composition and not eliciting an unacceptable deleterious effect in the subject. Such carriers are described herein or are otherwise well known to those skilled in the art of pharmacology. In some embodiments, the pharmaceutical compositions are pyrogen-free and are suitable for administration to a human patient. In some embodiments, the pharmaceutical compositions are irritant-free and are suitable for administration to a human patient. In some embodiments, the pharmaceutical compositions are allergen-free and are suitable for administration to a human patient. The compositions may be prepared by any of the methods well known in the art of pharmacy.

[0122] In some embodiments, an SAP variant or SAP oligomer is administered in a time release formulation, for example in a composition which includes a slow release polymer. An SAP variant or SAP oligomer can be prepared with carriers that will protect against rapid release. Examples include a controlled release vehicle, such as a polymer, microencapsulated delivery system, or bioadhesive gel. Alternatively, prolonged delivery of an SAP variant or SAP oligomer may be achieved by including in the composition agents that delay absorption, for example, aluminum monostearate hydrogels and gelatin.

[0123] Methods for delivering nucleic acid compounds are known in the art (see, e.g., Akhtar et al., 1992, Trends Cell Bio., 2, 139; and Delivery Strategies for Antisense Oligonucleotide Therapeutics, ed. Akhtar, 1995; Sullivan et al., PCT Publication No. WO 94/02595). These protocols can be utilized for the delivery of virtually any nucleic acid compound. Nucleic acid compounds can be administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. Alternatively, the nucleic acid/vehicle combination is locally delivered by direct injection or by use of an infusion pump. Other routes of delivery include, but are not limited to, oral (tablet or pill form) and/or intrathecal delivery (Gold, 1997, Neuroscience, 76, 1153-1158). Other approaches include the use of various transport and carrier systems, for example though the use of conjugates and biodegradable polymers. For a comprehensive review on drug delivery strategies, see Ho et al., 1999, Curr. Opin. Mol. Ther., 1, 336-343 and Jain, Drug Delivery Systems: Technologies and Commercial Opportunities, Decision Resources, 1998 and Groothuis et al., 1997, J. NeuroVirol., 3, 387-400. More detailed descriptions of nucleic acid delivery and administration are provided in Sullivan *et al.*, supra, Draper et al., PCT WO93/23569, Beigelman et al., PCT Publication No. WO99/05094, and Klimuk et al., PCT Publication No. WO99/04819.

[0124] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes.

EXEMPLIFICATION

Example 1: SAP variants resistant to calcium-mediated aggregation.

[0125] A recombinant human SAP (rhSAP) variant comprising an E167Q amino acid substitution, relative to the sequence of SEQ ID NO: 1, was expressed in CHO cells and purified from the CHO cell culture media. The calcium-mediated aggregation of the rhSAP variant was then compared to that of a corresponding sample of wild-type rhSAP. Incremental amounts of calcium was added to either a solution of rhSAP variant E167Q or wild-type rhSAP

(each at a SAP concentration of 4.4 mg/mL), and the amount of SAP aggregation was observed by measuring the absorbance of the solution at 600 nm in a spectrophotometer. Figure 1 demonstrates that the rhSAP variant E167Q is significantly more resistant to calcium-mediated aggregation than the wild-type rhSAP.

[0126] The pharmacokinetics (PK) of the rhSAP variant E167Q was also compared to a corresponding sample of wild-type rhSAP. Rats were administered (1 mg/kg i.v. dose per rat, n=3) either the rhSAP variant E167Q or a corresponding sample of wild-type rhSAP. Over the next twenty-four hours, the rats were assessed for plasma concentration ($\mu\text{g/mL}$) of SAP protein. Figure 2 demonstrates that the rhSAP variant E167Q has a similar plasma half-life to that of wild-type rhSAP.

[0127] In a further experiment, an *in vitro* bioassay was used to determine the relative activity of the rhSAP variant E167Q. In this assay, monocyte-enriched Peripheral Blood Mononuclear Cells (PBMCs) were incubated with varying concentrations of either the rhSAP variant E167Q or hSAP for 96 hours. Following this incubation, the resulting culture supernatants were removed and assayed by ELISA to quantify the amount of Macrophage Derived Chemokine (MDC) that was produced. MDC is produced by fibrocytes and therefore an indicator of monocyte differentiation into fibrocytes. By comparing the inhibitory concentration, 50% (IC_{50}) of the sample to the hSAP reference standard, the relative potency of a SAP variant can be determined. The result is expressed as an IC_{50} ratio of the sample versus the hSAP reference standard.

[0128] All SAP samples and standards were initially diluted to a concentration of 1.0 mg/mL in Supplemented FibroLife Media. SAP standards were serially diluted to generate working standard concentrations of 60, 30, 20, 13.4, 8.8, 6.0, 3.0, 1.5, and 0.75 $\mu\text{g/mL}$ (final standard concentration of 30, 15, 10, 6.7, 4.4, 3.0, 1.5, 0.75, and 0.375 $\mu\text{g/mL}$). See the following Table

1

Working rhSAP Standard Concentration ($\mu\text{g/mL}$)	Volume of Standard	Volume of Supplemented FibroLife Media
60	60 (1mg/mL)	940
30	600 (60 $\mu\text{g/mL}$)	600
20	800 (30 $\mu\text{g/mL}$ Std)	400
13.4	800 (20 $\mu\text{g/mL}$ Std)	400
8.8	800 (13.4 $\mu\text{g/mL}$ Std)	400
6.0	800 (8.8 $\mu\text{g/mL}$ Std)	400
3.0	600 (6.0 $\mu\text{g/mL}$ Std)	600

Working rhSAP Standard Concentration (µg/mL)	Volume of Standard	Volume of Supplemented FibroLife Media
1.5	600 (3.0 µg/mL Std)	600
0.75	600 (1.5 µg/mL Std)	600

[0129] To prepare for the ELISA assay, the Capture Antibody (i.e., mouse anti-human MDC) was diluted to the working concentration in PBS without carrier protein. The diluted capture antibody was used to coat 96-well plates, and then each plate was sealed and incubated overnight at room temperature. Before using the coated plates, each well was aspirated and washed with Wash Buffer, repeating the process two times for a total of three washes. The plates were then blocked by adding 300 µL of Reagent Diluent to each well and incubating at room temperature for one hour. After incubation the aspiration and well-washing procedure was repeated.

[0130] For the ELISA assay, 100 µL samples of either the supernatants from the monocyte/fibrocyte cultures or the SAP standards were added to each well. The plate was then incubated at room temperature for 2 hours before aspirating and washing the wells. Then 100 µL of a working dilution of Streptavidin-HRP was added to each well. The plate was incubated for 20 minutes at room temperature before adding 50 µL of Stop Solution to each well. Immediately, the optical density of each well was measured using a microplate reader set to 450 nm. If wavelength correction was available, the microplate reader was set to 540 nm or 570 nm. If wavelength correction was not available, then the readings at 540 nm or 570 nm were subtracted from the readings at 450 nm. This subtraction corrects for optical imperfections in the plate.

[0131] Figure 3 demonstrates that rhSAP variant E167Q is at least as biologically active as wild-type rhSAP.

Example 2: Deglycosylated SAP variants have altered plasma half-life and biological activity.

[0132] A recombinant human SAP (rhSAP) variant comprising an N32D amino acid substitution, relative to the sequence of SEQ ID NO: 1, was expressed in CHO cells and purified from the CHO cell culture media. This mutation disrupts an N-glycosylation consensus site and thereby prevents attachment of N-linked glycans to SAP at that position. In parallel, wild-type hSAP was treated with a sialidase to remove all sialic acid moieties attached to the SAP polypeptide (i.e., asialo hSAP). Both the untreated rhSAP N32D and asialo hSAP were compared with a corresponding sample of rhSAP in a PK assay to measure for *in vivo* serum stability (Figure 4). While the PK of the rhSAP N32D was slightly reduced compared to wild-

type SAP, the half-life of the rhSAP N32D was substantially higher than that of the asialo hSAP. The rhSAP N32D variant was further compared to a corresponding sample of serum-derived hSAP using an *in vitro* bioassay to determine the relative activity of these proteins (Figure 3). These data indicate that the N32D SAP variant maintains a plasma half-life and activity comparable to wild-type hSAP.

Example 3: Covalent attachment of PEG to SAP.

[0133] A recombinant human SAP (rhSAP) variant was covalently attached to a 20 kDa activated methoxyPEG derivative (PEG). The PEG moiety was attached to a primary amine group of rhSAP according to the following protocol and as illustrated in Figure 5. First, approximately 1 mg of 20 kDa methoxy-PEG-succinimidyl-carboxymethyl ester (JenKem cat# M-SCM-20K) per mg of rhSAP was dissolved in a 20 mg/mL solution of rhSAP. The coupling reaction was allowed to proceed for 24 hours at room temperature. The resulting pegylated-rhSAP was purified from reaction components by anion exchange chromatography. Fractions from the chromatography column were pooled and concentrated (Figure 6). PEGylated rhSAP made by this procedure contained from 1-3 20kDa PEGs/protomer, with 1 PEG/protomer being the most abundant form, as assessed by SDS-PAGE.

[0134] The PEGylated rhSAP and human serum-derived SAP (hSAP) were assayed for bioactivity using an *in vitro* bioassay. In this assay, monocyte enriched Peripheral Blood Mononuclear Cells (PBMCs) were incubated with varying concentrations of either PEGylated rhSAP or hSAP for 96 hours. Following this incubation, resulting culture supernatants were removed and assayed by ELISA to quantify the amount of Macrophage Derived Chemokine (MDC) that was produced. MDC is produced by fibrocytes and therefore an indicator of monocyte differentiation into fibrocytes. By comparing the inhibitory concentration, 50% (IC_{50}) of the sample to the hSAP reference standard, the relative potency of a SAP variant can be determined. The result is expressed as an IC_{50} ratio of the sample versus the hSAP reference standard as described in the preceding examples.

[0135] The PEGylated rhSAP variant was had an IC_{50} ratio of 0.24 compared to a corresponding sample of hSAP, thereby demonstrating that the PEGylated rhSAP has comparable activity to wild-type SAP.

[0136] Also described herein are the following items:

1. 1. A Serum Amyloid P (SAP) variant comprising five SAP protomers, wherein each of the SAP protomers have an amino acid sequence at least 90% identical to SEQ ID NO: 1, and wherein at least one of the SAP protomers comprises one or more amino acid mutations or chemical modifications that alter a biological activity of the SAP variant compared to a corresponding sample of serum-derived human SAP.
2. 2. The SAP variant of item 1, wherein the SAP variant is a variant of a human SAP protein.

3. 3. The SAP variant of item 1, wherein one or more of the SAP protomers has an amino acid sequence at least 99% identical to SEQ ID NO: 1.
4. 4. The SAP variant of any one of items 1-3, wherein the biological activity is selected from one or more of increased plasma half-life, increased *in vitro* stability, increased *in vivo* stability, or increasing manufacturing efficiency.
5. 5. The SAP variant of any one of items 1-4, wherein at least one of the SAP protomers comprises a variant amino acid relative to the sequence of SEQ ID NO: 1.
6. 6. The SAP variant of any one of items 1-4, wherein one or more of the SAP protomers are substantially free of N-linked and O-linked glycans.
7. 7. The SAP variant of item 6, wherein one or more of the SAP protomers comprises an amino acid at position 32 of SEQ ID NO: 1 that is not asparagine (N).
8. 8. The SAP variant of item 7, wherein the SAP protomer comprises an aspartate (D), glutamine (Q), or glutamate (E) at position 32 of SEQ ID NO: 1.
9. 9. The SAP variant of any one of items 1-8, wherein the SAP variant is more resistant to protease cleavage than a corresponding sample of serum-derived human SAP.
10. 10. The SAP variant of item 9, wherein the SAP variant is more resistant to protease cleavage by a serine protease, a threonine protease, a cysteine protease, an aspartic acid protease, a metalloprotease, a glutamic acid protease, or combinations thereof.
11. 11. The SAP variant of item 9 or 10, wherein the SAP variant is more resistant to protease cleavage by chymotrypsin, trypsin, Pronase, or combinations thereof.
12. 12. The SAP variant of any one of items 9-11, wherein the SAP variant comprises at least one SAP protomer comprising an amino acid at position 144 of SEQ ID NO: 1 that is not phenylalanine (F).
13. 13. The SAP variant of any one of items 9-11, wherein the SAP variant comprises at least one SAP protomer comprising an amino acid at position 145 of SEQ ID NO: 1 that is not D.
14. 14. The SAP variant of any one of items 9-11, wherein the SAP variant comprises at least one SAP protomer comprising an amino acid at positions 144 of SEQ ID NO: 1 that is not F, and wherein the SAP protomer further comprise an amino acid at position 145 of SEQ ID NO: 1 that is not D.
15. 15. The SAP variant of item 12 or 14, wherein the SAP protomer comprises a leucine (L), isoleucine (I), valine (V), or alanine (A) at position 144 of SEQ ID NO: 1.
16. 16. The SAP variant of item 13 or 14, wherein the SAP promoter comprises a glutamate (E) at position 145 of SEQ ID NO: 1.
17. 17. The SAP variant of any one of items 1-16, wherein the SAP variant is more resistant to calcium-dependant autoaggregation than a corresponding sample of serum-derived human SAP.
18. 18. The SAP variant of item 17, wherein the SAP variant comprises at least one SAP protomer comprising an amino acid at position 167 of SEQ ID NO: 1 that is not E.
19. 19. The SAP variant of item 18, wherein the SAP protomer comprises a mutation of an aspartate (D), asparagine (N), glutamine (Q), alanine (A,) or histidine (H) at position 167 of SEQ ID NO: 1.
20. 20. The SAP variant of any one of items 1-19, wherein the SAP variant comprises at least one SAP protomer comprising one or more amino acids that are covalently

attached to one or more inert polymers.

21. 21. The SAP variant of item 20, wherein at least one of the inert polymers is a polyethylene glycol (PEG) moiety.
22. 22. The SAP variant of item 21, wherein the PEG moiety is attached to at least one native or variant cysteine residue of the SAP protomer.
23. 23. The SAP variant of item 22, wherein the variant cysteine residue is located at the N-terminus of SEQ ID NO: 1.
24. 24. The SAP variant of item 21, wherein the PEG moiety is attached to at least one native or variant glutamine (Q) residue of SEQ ID NO: 1.
25. 25. The SAP variant of item 24, wherein the native glutamine (Q) residue is at position 32 of SEQ ID NO: 1.
26. 26. The SAP variant of any one of items 20-25, wherein at least one of the inert polymers is a dextran moiety.
27. 27. The SAP variant of item 26, wherein the dextran moiety is attached to at least one native or variant glutamine residue of SEQ ID NO: 1.
28. 28. The SAP variant of item 27, wherein the native glutamine residue is at position Q32 corresponding to the amino acid sequence of SEQ ID NO: 1.
29. 29. A covalently crosslinked SAP oligomer comprising at least two SAP pentamers, wherein each of the SAP pentamers comprises five SAP protomers.
30. 30. The covalently crosslinked SAP oligomer of item 29, wherein the crosslinked SAP oligomer is characterized by one or more of increased plasma half-life, increased *in vitro* stability, and increased *in vivo* stability compared to a corresponding sample of SAP isolated from human serum.
31. 31. The covalently crosslinked SAP oligomer of item 29 or 30, wherein one or more of the SAP protomers have an amino acid sequence at least 90% identical to SEQ ID NO: 1.
32. 32. The covalently crosslinked SAP oligomer of any one of items 29-31, wherein at least one of the SAP pentamers is an SAP variant of any one of items 1-28.
33. 33. The covalently crosslinked SAP oligomer of any one of items 29-32, wherein the SAP pentamers are covalently attached through one or more chemical cross-linkers.
34. 34. The covalently crosslinked SAP oligomer of item 33, wherein at least one of the chemical cross-linkers is a heterobifunctional agent.
35. 35. The covalently crosslinked SAP oligomer of item 34, wherein the heterobifunctional cross-linker is selected from succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate, m-Maleimidobenzoyl-N-hydroxysuccinimide ester, N-succinimidyl (4-iodoacetyl) aminobenzoate, succinimidyl 4-(p-maleimidophenyl) butyrate, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride, 4-succinimidyloxycarbonyl- α -methyl- α -(2-pyridyldithio)-toluene, N-succinimidyl 3-(2-pyridyldithio) propionate, or succinimidyl 6-((3-(2-pyridyldithio)propionate)hexanoate.
36. 36. The covalently crosslinked SAP oligomer of item 33, wherein at least one of the chemical cross-linkers is a homobifunctional agent.
37. 37. The covalently crosslinked SAP oligomer of item 36, wherein the homobifunctional agent is selected from disuccinimidyl suberate, bismaleimido-hexane, or dimethyl-pimelidate-2 HCl.

38. 38. The covalently crosslinked SAP oligomer of item 33, wherein at least one of the chemical cross-linkers is a photoreactive agent.
39. 39. The covalently crosslinked SAP oligomer of item 38, wherein the photoreactive agent is selected from bis-(β -(4-azidosalicylamido)ethyl)disulfide or N-succinimidyl-6-(4'-azido-2'-nitrophenyl-amino)hexanoate.
40. 40. A pharmaceutical preparation suitable for use in a mammal comprising the SAP variant or covalently crosslinked SAP oligomer of any one of items 1-39 and a pharmaceutically acceptable carrier.
41. 41. The pharmaceutical preparation of item 41, wherein the pharmaceutical preparation is prepared as a sustained release formulation.
42. 42. The pharmaceutical preparation of item 40 or 41, wherein the pharmaceutical preparation is suitable for administration to a patient topically, by injection, by intravenous injection, by inhalation, by continuous depot, or by pump.
43. 43. A method of treating or preventing a fibrotic or fibroproliferative disorder or condition in a patient, the method comprising administering to a patient in need thereof a therapeutically effective amount of the SAP variant or covalently crosslinked SAP oligomer of any one of items 1-39.
44. 44. A method of treating or preventing a hypersensitivity disorder or condition in a patient, the method comprising administering to a patient in need thereof a therapeutically effective amount of the SAP variant or covalently crosslinked SAP oligomer of any one of items 1-39.
45. 45. A method of treating or preventing an autoimmune disorder or condition in a patient, the method comprising administering to a patient in need thereof a therapeutically effective amount of the SAP variant or covalently crosslinked SAP oligomer of any one of items 1-39.
46. 46. A method of treating or preventing mucositis in a patient, the method comprising administering to a patient in need thereof a therapeutically effective amount of the SAP variant or covalently crosslinked SAP oligomer of any one of items 1-39.
47. 47. The method of any one of items 43-46, wherein the SAP variant or covalently crosslinked SAP oligomer is administered topically, by injection, by intravenous injection, by inhalation, by continuous depot or pump, or a combination thereof.
48. 48. The method of any one of items 43-47, further comprising administering to the patient an additional active agent.

SEQUENCE LISTING

[0137]

<110> PROMEDIOR, INC.

<120> SAP VARIANTS AND THEIR USE

<130> T3461 EP/1

<150> US 61/268,961

<151> 2009-06-17

<160> 2

<170> PatentIn version 3.5

<210> 1

<211> 204

<212> PRT

<213> Homo sapiens

<400> 1

His	Thr	Asp	Leu	Ser	Gly	Lys	Val	Phe	Val	Phe	Pro	Arg	Glu	Ser	Val
1				5					10					15	

Thr	Asp	His	Val	Asn	Leu	Ile	Thr	Pro	Leu	Glu	Lys	Pro	Leu	Gln	Asn
			20					25					30		

Phe	Thr	Leu	Cys	Phe	Arg	Ala	Tyr	Ser	Asp	Leu	Ser	Arg	Ala	Tyr	Ser
		35					40					45			

Leu	Phe	Ser	Tyr	Asn	Thr	Gln	Gly	Arg	Asp	Asn	Glu	Leu	Leu	Val	Tyr
50						55					60				

Lys	Glu	Arg	Val	Gly	Glu	Tyr	Ser	Leu	Tyr	Ile	Gly	Arg	His	Lys	Val
65					70					75					80

Thr	Ser	Lys	Val	Ile	Glu	Lys	Phe	Pro	Ala	Pro	Val	His	Ile	Cys	Val
				85					90					95	

Ser	Trp	Glu	Ser	Ser	Ser	Gly	Ile	Ala	Glu	Phe	Trp	Ile	Asn	Gly	Thr
			100					105					110		

Pro	Leu	Val	Lys	Lys	Gly	Leu	Arg	Gln	Gly	Tyr	Phe	Val	Glu	Ala	Gln
		115					120					125			

Pro	Lys	Ile	Val	Leu	Gly	Gln	Glu	Gln	Asp	Ser	Tyr	Gly	Gly	Lys	Phe
	130					135					140				

Asp	Arg	Ser	Gln	Ser	Phe	Val	Gly	Glu	Ile	Gly	Asp	Leu	Tyr	Met	Trp
145					150					155					160

Asp	Ser	Val	Leu	Pro	Pro	Glu	Asn	Ile	Leu	Ser	Ala	Tyr	Gln	Gly	Thr
				165					170					175	

Pro	Leu	Pro	Ala	Asn	Ile	Leu	Asp	Trp	Gln	Ala	Leu	Asn	Tyr	Glu	Ile
			180					185					190		

Arg	Gly	Tyr	Val	Ile	Ile	Lys	Pro	Leu	Val	Trp	Val
	195						200				

<210> 2

<211> 234

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<220>

<221> variant

<222> 1

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 2

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 3

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 4

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 4

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"

/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 5

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 6

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 7

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 8

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 9

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 10

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"

/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 11

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 12

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 13

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 14

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 15

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 16

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 17

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 18

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 19

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> variant

<222> 20

<223> /replace="Cys" /replace="Asp" /replace="Glu" /replace="Phe" /replace="Gly"
/replace="His" /replace="Ile" /replace="Lys" /replace="Leu" /replace="Met" /replace="Asn"
/replace="Pro" /replace="Gln" /replace="Arg" /replace="Ser" /replace="Thr" /replace="Val"
/replace="Trp" /replace="Tyr"

<220>

<221> misc_feature

<222> (1)..(20)

<223> This region may encompass 3-20 residues

<220>

<221> misc_feature

<222> (21)..(30)

<223> This region may encompass 1-10 residues

<220>

<223> See specification as filed for detailed description of substitutions and preferred embodiments

<400> 2

Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala
1 5 10 15

Ala Ala Ala Ala Gln Gln Gln Gln Gln Gln Gln Gln Gln His Thr
20 25 30

Asp Leu Ser Gly Lys Val Phe Val Phe Pro Arg Glu Ser Val Thr Asp
35 40 45

His Val Asn Leu Ile Thr Pro Leu Glu Lys Pro Leu Gln Asn Phe Thr
50 55 60

Leu Cys Phe Arg Ala Tyr Ser Asp Leu Ser Arg Ala Tyr Ser Leu Phe
65 70 75 80

Ser Tyr Asn Thr Gln Gly Arg Asp Asn Glu Leu Leu Val Tyr Lys Glu
85 90 95

Arg Val Gly Glu Tyr Ser Leu Tyr Ile Gly Arg His Lys Val Thr Ser
100 105 110

Lys Val Ile Glu Lys Phe Pro Ala Pro Val His Ile Cys Val Ser Trp
115 120 125

Glu Ser Ser Ser Gly Ile Ala Glu Phe Trp Ile Asn Gly Thr Pro Leu
130 135 140

Val Lys Lys Gly Leu Arg Gln Gly Tyr Phe Val Glu Ala Gln Pro Lys
145 150 155 160

Ile Val Leu Gly Gln Glu Gln Asp Ser Tyr Gly Gly Lys Phe Asp Arg
165 170 175

Ser Gln Ser Phe Val Gly Glu Ile Gly Asp Leu Tyr Met Trp Asp Ser
180 185 190

Val Leu Pro Pro Glu Asn Ile Leu Ser Ala Tyr Gln Gly Thr Pro Leu
195 200 205

Pro Ala Asn Ile Leu Asp Trp Gln Ala Leu Asn Tyr Glu Ile Arg Gly
210 215 220

Tyr Val Ile Ile Lys Pro Leu Val Trp Val
225 230

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US707333A](#) [0002] [0017]
- [US12217617B](#) [0002] [0017]
- [US12720845B](#) [0002] [0017]
- [US12720847B](#) [0002] [0017]
- [WO2009009019A](#) [0003]
- [US5643575A](#) [0052]
- [US5223409A](#) [0066]
- [US5198346A](#) [0066]
- [US5096815A](#) [0066]
- [US6365410B](#) [0069]
- [US6171820B](#) [0070]
- [US5605793A](#) [0071] [0073]
- [US5811238A](#) [0071] [0073]
- [US5830721A](#) [0071] [0073]
- [US5834252A](#) [0071] [0073]
- [US5837458A](#) [0071] [0073]
- [US5965408A](#) [0073]
- [US6180406B](#) [0073]
- [US6165793A](#) [0073]
- [US6132970A](#) [0073]
- [US6117679A](#) [0073]
- [US6096548A](#) [0073]
- [US6361974B](#) [0074]
- [US4088538A](#) [0079]
- [US4496689A](#) [0079]
- [US4414147A](#) [0079]
- [WO8700056A](#) [0079]
- [EP154316A](#) [0079]
- [US4055635A](#) [0079]
- [WO9405332A](#) [0079]
- [US20060116322A](#) [0084]

- [US5807715A](#) [0093]
- [US20070243163A](#) [0100]
- [US61209795A](#) [0102]
- [US61209845A](#) [0104]
- [US217614A](#) [0105]
- [US6406698B](#) [0106]
- [WO1997026906A](#) [0106]
- [WO9402595A](#) [0123]
- [WO9323569A](#) [0123]
- [WO9905094A](#) [0123]
- [WO9904819A](#) [0123]
- [US61268961B](#) [0137]

Non-patent literature cited in the description

- **BALTZ, M. L. et al.** Clin. Exp. Immunol., 1986, vol. 66, 691-700 [0016]
- **DYCK, R. F. et al.** J. Exp. Med., 1980, vol. 152, 1162-1174 [0016]
- **MELVIN, T.** Am. J. Pathol., 1986, vol. 125, 460-464 [0016]
- **BREATHNACH, S. M.** J. Invest. Derm., 1989, vol. 92, 53-58 [0016]
- **CLAYTON, J.** Cell. Pathol., 1983, vol. 43, 63-66 [0016]
- **HERRIUT, R. et al.** J. Pathol., 1989, vol. 157, 11-14 [0016]
- **KHAN, A. M. et al.** Placenta, 1985, vol. 6, 551-554 [0016]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press 1989 0000 [0020]
- **Pharmaceutical Biotechnology** Harwood Publishers 1997 0000 101-120 [0029]
- **Computational Molecular Biology** Oxford University Press 1988 0000 [0033]
- **Biocomputing: Informatics and Genome Projects** Academic Press 1993 0000 [0033]
- **Computer Analysis of Sequence Data** Humana Press 1994 0000 [0033]
- **HEINJE, G.** Sequence Analysis in Molecular Biology Academic Press 1987 0000 [0033]
- **Sequence Analysis Primer** M Stockton Press 1991 0000 [0033]
- **NEEDLEMAN WUNSCHJ** Mol. Biol., 1970, vol. 48, 444-453 [0034]
- **DEVEREUX, J. et al.** Nucleic Acids Res., 1984, vol. 12, 1387- [0034]
- **E. MYERS W. MILLER** CABIOS, 1989, vol. 4, 11-17 [0035]
- **BRUTLAG et al.** Comp. App. Biosci., 1990, vol. 6, 237-245 [0036] [0036]
- **BOWIE et al.** Science, 1990, vol. 247, 1306-1310 [0039]
- **KENNETH, A et al.** Chemical Stability of Pharmaceuticals: A Handbook for Pharmacists, [0041]
- **PETERS et al.** Pharmacokinetic analysis: A Practical Approach, 1996, [0041]
- **M GIBALDID PERRON** Pharmacokinetics Marcel Dekker 1982 0000 [0041]
- **KINOSHITA CM et al.** Protein Science, 1992, vol. 1, 700-709 [0044]

- **KINOSHITA CM**Protein Science, 1992, vol. 1, 700-709 [0045]
- **HANS J. VOGEL**Contributor Hans J. VogelHumana Press20020000 [0046]
- **SEGEL, I.H.**Enzyme KineticsWiley-Interscience Publisher19750000218-19 [0046]
- **LINSE, SHELMBERSSON, AFORSEN, SJBC**, 1991, vol. 266, 138050-8054 [0046]
- **WISEMAN, T.WILLISTON, S.BRANDTS, J.F.LIN, L.N.**Anal Biochem, 1989, vol. 179, 131-7 [0046]
- **EMSLCYN**Nature, 1994, vol. 367, 338-345 [0047]
- **PEPYS**Nature, 2002, vol. 417, 254-259 [0047]
- **HOJ** Biol Chem, 2005, vol. 280, 31999-32008 [0047]
- **MORPURGO et al.**Appl. Biochem. Biotechnol., 1996, vol. 56, 59-72 [0052]
- **VOROBJEV et al.**Nucleosides Nucleotides, 1999, vol. 18, 2745-2750 [0052]
- **CALICETI et al.**Bioconjug. Chem., 1999, vol. 10, 638-646 [0052]
- **MEANS et al.**Bioconjugate Chemistry, 1990, vol. 1, 2-12 [0056]
- **PARTIS et al.**J. Pro. Chem., 1983, vol. 2, 263- [0060]
- **ELLMAN et al.**Arch. Biochem. Biophys., 1958, vol. 74, 443- [0060]
- **RIDDLES et al.**Anal. Biochem., 1979, vol. 94, 75- [0060]
- **BLATTLER et al.**Biochem, 1985, vol. 24, 1517- [0060]
- **SAMBROOKRUSSELL**Molecular Cloning, A Laboratory ApproachCold Spring Harbor Press20010000 [0063]
- **AUSUBEL et al.**Current Protocols in Molecular BiologyJohn Wiley & Sons20020000 [0063]
- **LEUNG et al.**Technique, 1989, vol. 1, 11-15 [0064]
- **MAYERS et al.**Science, 1985, vol. 229, 242- [0065]
- **NARANG, S A**Tetrahedron, 1983, vol. 39, 3- [0066]
- **Proc 3rd Cleveland Sympos. MacromoleculesITAKURA et al.**Recombinant DNAElsevier19810000273-289 [0066]
- **ITAKURA et al.**Annu. Rev. Biochem., 1984, vol. 53, 323- [0066]
- **ITAKURA et al.**Science, 1984, vol. 198, 1056- [0066]
- **IKE et al.**Nucleic Acid Res., 1983, vol. 11, 477- [0066]
- **SCOTT et al.**Science, 1990, vol. 249, 386-390 [0066]
- **ROBERTS et al.**PNAS, 1992, vol. 89, 2429-2433 [0066]
- **DEVLIN et al.**Science, 1990, vol. 249, 404-406 [0066]
- **CWIRLA et al.**PNAS, 1990, vol. 87, 6378-6382 [0066]
- **CALDWELLJOYCE**PCR Methods Appl., 1994, vol. 2, 28-33 [0068]
- **KEMADISON**Nucleic Acids Res., 1997, vol. 25, 3371-3372 [0068]
- **REIDHAAR-OLSON et al.**Methods Enzymol., 1991, vol. 208, 564-586 [0068]
- **SELIFONOVA et al.**Appl. Environ. Microbiol., 2001, vol. 67, 3645-3649 [0069]
- **LONG-MCGIE et al.**Biotech. Bioeng., 2000, vol. 68, 121-125 [0069]
- **STEMMER et al.**Nature, 1994, vol. 370, 6488389-391 [0071]
- **CRAMERI et al.**Nature, 1998, vol. 391, 6664288-291 [0071]
- **ZHANG et al.**Proc. Natl. Acad. Sci. USA, 1997, vol. 94, 94504-4509 [0071]
- **STEMMER et al.**Proc. Natl. Acad. Sci USA, 1994, vol. 91, 2210747-10751 [0071]
- **PATTEN et al.**Curr. Opinion Biotechnol., 1997, vol. 8, 724-33 [0071]
- **HARAYAMA**Trends Biotechnol., 1998, vol. 16, 276-82 [0071]

- **HANSSON et al.**J. Mol. Biol., 1999, vol. 287, 265-76 [0071]
- **LORENZOBLASCO**Biotechniques, 1998, vol. 24, 2308-13 [0071]
- **PEKRUN et al.**J. Virol., 2002, vol. 76, 62924-35 [0072]
- **RAILLARD et al.**Chem Biol, 2001, vol. 8, 9891-898 [0072]
- **POWELL et al.**Nat Biotechnol, 2000, vol. 18, 121279-1282 [0072]
- **MERZ et al.**Biochemistry, 2000, vol. 39, 5880-889 [0072]
- **STEMMLER**Nature, 1994, vol. 370, 389-391 [0073] [0073]
- **STEMMLER**PNAS, 1994, vol. 91, 10747-10751 [0073] [0073]
- **KOLKMANSTEMMLER**Nat. Biotech., 2001, vol. 19, 423-428 [0074]
- **COCO et al.**Nat. Biotechnol., 2001, vol. 19, 354-359 [0074]
- **ZHAO et al.**Nat. Biotechnol., 1998, vol. 16, 258-261 [0074]
- **ABECASSIS et al.**Nucleic Acids Res., 2000, vol. 28, E88- [0074]
- **KIKUCHI et al.**Gene, 2000, vol. 243, 133-137 [0074]
- **SIEBER et al.**Nat Biotechnol., 2001, vol. 19, 456-460 [0074]
- **LUTZ et al.**Nucleic Acids Res., 2001, vol. 29, E16- [0074]
- **OSTERMEIER et al.**Nat. Biotechnol., 1999, vol. 17, 1205-1209 [0074]
- **LUTZBENKOVIC**Curr. Opin. Biotechnol., 2000, vol. 11, 319-324 [0074]
- **ABUCHOWSKI et al.**J. Biol. Chem., 1977, vol. 252, 3578- [0079]
- **NATARAJAN**Bioconjug Chem., 2005, vol. 16, 1113-21 [0080]
- **GOODSON**Biotechnology NY, 1990, vol. 8, 4343-6 [0080]
- **KAHLEM, P. et al.**Proc. Natl. Acad. Sci. USA, 1996, vol. 93, 14580-14585 [0083]
- **SATO**H. Adv Drug Deliv Rev., 2002, vol. 54, 4487-504 [0084]
- **SATO et al.**Bioconjug Chem., 2000, vol. 11, 4502-9 [0084]
- **SATO et al.**Bioconjug Chem., 2001, vol. 12, 5701-10 [0084]
- **FONTANA et al.**Adv Drug Deliv Rev., 2008, vol. 60, 113-28 [0084]
- **HOHENADLJ** Biol Chem., 1995, vol. 270, 4023415-20 [0084]
- **GOEDDEL**Gene Expression Technology: Methods in EnzymologyAcademic Press19900000 [0086]
- **BROACH et al.**Experimental Manipulation of Gene ExpressionAcademic Press1983000083- [0091]
- **Molecular Cloning: A Laboratory Manual**Cold Spring Harbor Laboratory Press19890000 [0092]
- **GRAHAM et al.**J. Gen Virol., 1977, vol. 36, 59- [0093]
- **CHO, URLAUB et al.**Proc. Natl. Acad. Sci. USA, 1980, vol. 77, 4216- [0093]
- **MATHER**Biol. Reprod., 1980, vol. 23, 243-251 [0093]
- **MATHER et al.**Annals N.Y. Acad. Sci., 1982, vol. 383, 44-68 [0093]
- **Affinity Chromatography: Principles & Methods**Pharmacia LKB Biotechnology19880000 [0097]
- **Remington's Pharmaceutical Sciences**Meade Publishing Co. [0109]
- **ROHATAGIJ.** Clin. Pharmacol., 1995, vol. 35, 1187-1193 [0112]
- **TJWA**Ann. Allergy Asthma Immunol., 1995, vol. 75, 107-111 [0112]
- **BEGLEY**Pharmacology & Therapeutics, 2004, vol. 104, 29-45 [0116]
- **AKHTAR et al.**Trends Cell Bio., 1992, vol. 2, 139- [0123]
- **Delivery Strategies for Antisense Oligonucleotide Therapeutics**19950000 [0123]

- **GOLD**Neuroscience, 1997, vol. 76, 1153-1158 [\[0123\]](#)
- **HO et al.**Curr. Opin. Mol. Ther., 1999, vol. 1, 336-343 [\[0123\]](#)
- **JAIN**Drug Delivery Systems: Technologies and Commercial Opportunities, Decision Resources, 1998, [\[0123\]](#)
- **GROOTHUIS et al.**J. NeuroVirol., 1997, vol. 3, 387-400 [\[0123\]](#)

PATENTKRAV

1. Farmaceutisk sammensætning omfattende en serumamyloid P (SAP)-variant omfattende fem SAP-promotorer, hvor hver af SAP-promotorerne har en aminosyre-sekvens, som er i det mindste 95% identisk med SEQ ID NO: 1, og hvor alle fem SAP promotorer omfatter en aminosyre ved position 167 i SEQ ID NO: 1, som er asparagin (N), glutamin (Q), alanin (A) eller histidin (H), og et farmaceutisk acceptabelt ophthalmisk bærestof.
2. Farmaceutisk sammensætning ifølge krav 1, hvor SAP-varianten er mere resistent overfor calcium-afhængig autoaggregation end en tilsvarende prøve af ikke-variant humant SAP:
3. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor hver af SAP-promotorerne har en aminosyresekvens, som er i det mindste 96% identisk med SEQ ID NO: 1.
4. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor hver af SAP-promotorerne har en aminosyresekvens, som er i det mindste 97% identisk med SEQ ID NO: 1.
5. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor hver af SAP-promotorerne har en aminosyresekvens, som er i det mindste 98% identisk med SEQ ID NO: 1.
6. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor hver af SAP-promotorerne har en aminosyresekvens, som er i det mindste 99% identisk med SEQ ID NO: 1.
7. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 6, hvor i det mindste én af SAP-promotorerne omfatter en glutamin (Q) ved position 167 i SEQ ID NO: 1.
8. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7, til anvendelse ved behandlingen eller forebyggelsen af makulær degeneration, retinal og vitreal retinopati, den fibroserelaterede lidelse eller tilstand efterfølgende trabekulektomi eller anden filtrationskirurgi i øjet, Grave's ophthalmopati, diabetiske retinopati eller glaucoma.
9. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7 til anvendelse ved behandlingen eller forebyggelsen af makulær degeneration.

10. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7 til anvendelse ved behandlingen eller forebyggelsen af fibrotisk eller fibroproliferativ lidelse.

11. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7 til anvendelse ved
5 behandlingen eller forebyggelsen af inflammatorisk lidelse.

12. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7, hvor den farmaceutiske sammensætning er egnet til indgivelse topisk, ved intraokulær injektion eller ved indsætning af en indretning til forsinket frigivelse.

10

13. Farmaceutisk sammensætning ifølge ethvert af kravene 1 til 7, hvor den farmaceutiske sammensætning er egnet til indgivelse ved intraokulær injektion.

14. Farmaceutisk sammensætning ifølge ethvert af kravene 8 til 11, til anvendelse ved
15 indgivelse topisk, ved intraokulær injektion, eller ved indsætning af en indretning til forsinket frigivelse.

15. Farmaceutisk sammensætning ifølge ethvert af kravene 8 til 11 til anvendelse ved intraokulær injektion.

20

DRAWINGS




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6