

(43) International Publication Date
20 November 2014 (20.11.2014)

(51) International Patent Classification:

C07D 413/12 (2006.01) A01N 43/82 (2006.01)
C07D 471/04 (2006.01)

(21) International Application Number:

PCT/EP2014/059026

(22) International Filing Date:

5 May 2014 (05.05.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

13167848.4 15 May 2013 (15.05.2013) EP

(71) Applicant: BASF SE [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors: **KRAUS, Helmut**; 7 Rue de Kaysersberg, F-67160 Wissembourg (FR). **CALO, Frederick**; Q4, 6, 68161 Mannheim (DE). **WITSCHER, Matthias**; Höhenweg 12b, 67098 Bad Dürkheim (DE). **SEITZ, Thomas**; Emil-Nolde-Straße 10, 68519 Viernheim (DE). **NEWTON, Trevor William**; Neubergstrasse 30, 67435 Neustadt (DE). **MASSA, Dario**; N3, 3, 68161 Mannheim (DE). **MIETZNER, Thomas**; Rehbergstr.62, 76855 Annweiler (DE). **PASTERNAK, Maciej**; Tiroler Str. 12, 67067 Ludwigshafen (DE). **KREUZ, Klaus**; Hauptstr. 30/4, 79211 Denzlingen (DE). **EVANS, Richard Roger**; Kropsburgstrasse 24, 67117 Limburgerhof (DE). **LERCHL, Jens**; Golmer Fichten 5, 14476 Golm (DE).

(74) Agent: **ISENBRUCK BÖSL HÖRSCHLER LLP**; EASTSITE ONE, Seckenheimer Landstrasse 4, 68165 Mannheim (DE).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: N-(1,2,5-OXADIAZOL-3-YL)CARBOXAMIDE COMPOUNDS AND THEIR USE AS HERBICIDES



(57) Abstract: Substituted 1,2,5-oxadiazole compounds of formula I and their use as herbicides. The invention relates to substituted 1,2,5-oxadiazole compounds of formula I and their use as herbicides. In said formula I, X¹ is N or CR¹, X² is N or CR² and X⁴ is N or CR⁴; provided that a least one of X¹, X² and X⁴ is N; R, R¹, R², R³, R⁴ and R⁵ represent groups such as hydrogen, halogen or organic groups such as alkyl or phenyl.

N-(1,2,5-oxadiazol-3-yl)CARBOXAMIDE COMPOUNDS AND THEIR USE AS HERBICIDES

Description

The present invention relates to substituted 1,2,5-oxadiazole compounds and the N-oxides and salts thereof and to compositions comprising the same. The invention also relates to the use of the 1,2,5-oxadiazole compounds or of the compositions comprising such compounds for controlling unwanted vegetation. Furthermore, the invention relates to methods of applying such compounds.

For the purposes of controlling unwanted vegetation, especially in crops, there is an ongoing need for new herbicides which have high activities and selectivities together with a substantial lack of toxicity for humans and animals.

EP 0 173 657 A1 and WO 2011/035874 describe herbicidally active N-(1,2,5-oxadiazol-3-yl)carboxamides.


The N-(1,2,5-oxadiazol-3-yl) carboxamides of the prior art often suffer from insufficient herbicidal activity in particular at low application rates and/or unsatisfactory selectivity resulting in a low compatibility with crop plants.

Accordingly, it is an object of the present invention to provide further 1,2,5-oxadiazole compounds having a strong herbicidal activity, in particular even at low application rate, a sufficiently low toxicity for humans and animals and/or a high compatibility with crop plants. The 1,2,5-oxadiazole compounds should also show a broad activity spectrum against a large number of different unwanted plants.

These and further objectives are achieved by the compounds of formula I defined below and their N-oxides and also their agriculturally suitable salts.

It has been found that the above objectives can be achieved by substituted 1,2,5-oxadiazole compounds of the general formula I, as defined below, including their N-oxides and their salts, in particular their agriculturally suitable salts.

Therefore, in a first aspect the present invention relates to compounds of formula I,



wherein

X¹ is N or CR¹;

X² is N or CR²;

X⁴ is N or CR⁴;

provided that a least one of X¹, X² and X⁴ is N;

R is selected from the group consisting of hydrogen, cyano, nitro, halogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two
 5 aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, O-R^a, Z-S(O)_n-R^b, Z-C(=O)-R^c, Z-C(=O)-OR^d, Z-C(=O)-NR^eR^f, Z-NR^gR^h,

Z-phenyl and Z-heterocyclyl, where heterocyclyl is a 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the
 10 group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R', which are identical or different;

R¹ is selected from the group consisting of Z¹-cyano, halogen, nitro, C₁-C₈-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₈-haloalkyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, Z¹-C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio-C₁-C₄-alkyl, Z¹-C₁-C₄-alkylthio-C₁-C₄-alkylthio, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₁-C₆-haloalkoxy, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, Z¹-C₁-C₄-haloalkoxy-C₁-C₄-alkoxy, Z¹-S(O)_k-R^{1b}, Z¹-phenoxy and Z¹-heterocyclyloxy, where heterocyclyloxy is an oxygen bound 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-
 15 membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in phenoxy and heterocyclyloxy are unsubstituted or substituted by 1, 2, 3 or 4 groups R¹¹, which are identical or different;

R², R³ are identical or different and independently selected from the group consisting of
 25 hydrogen, halogen, Z²-OH, Z²-NO₂, Z²-cyano, C₁-C₆-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, Z²-C₃-C₁₀-cycloalkyl, Z²-C₃-C₁₀-cycloalkoxy, where the C₃-C₁₀-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₈-haloalkyl, Z²-C₁-C₈-alkoxy, Z²-C₁-C₈-haloalkoxy, Z²-C₁-C₄-alkoxy-C₁-C₄-alkoxy, Z²-C₁-C₄-alkylthio-C₁-C₄-alkylthio, Z²-C₂-C₈-alkenyloxy, Z²-C₂-C₈-alkynyloxy, Z²-C₁-C₈-haloalkoxy,
 30 Z²-C₁-C₄-haloalkoxy-C₁-C₄-alkoxy, Z²-(tri-C₁-C₄-alkyl)silyl, Z²-S(O)_k-R^{2b}, Z²-C(=O)-R^{2c}, Z²-C(=O)-OR^{2d}, Z²-C(=O)-NR^{2e}R^{2f}, Z²-NR^{2g}R^{2h}, Z^{2a}-phenyl and Z^{2a}-heterocyclyl, where heterocyclyl is a 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in Z^{2a}-phenyl and Z^{2a}-heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R²¹, which are identical or different;

R⁴, R⁵ are identical or different and independently selected from the group consisting of hydrogen, cyano- Z¹, halogen, nitro, C₁-C₈-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₈-haloalkyl, C₁-C₃-
 40

alkylamino, C₁-C₃-dialkylamino, C₁-C₃-alkylamino-S(O)_k, C₁-C₃-alkylcarbonyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-Z¹, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkylthio-Z¹, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₁-C₆-haloalkoxy, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkoxy-Z¹, R^{1b}-S(O)_k-Z¹, phenoxy-Z¹ and heterocyclyloxy-Z¹, where heterocyclyloxy is an oxygen bound 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in phenoxy and heterocyclyloxy are unsubstituted or substituted by 1, 2, 3 or 4 groups R¹¹, which are identical or different;

provided that if X⁴ is N, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl; and further provided that if X⁴ is CR⁴ and R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

where for X² = CR², R² together with R³ or together with R¹, if present, may also form a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle, where the fused heterocycle has 1, 2, 3 or 4 heteroatoms selected from O, S and N as ring members, where the fused carbocycle and the fused heterocycle are monocyclic or bicyclic and where the fused carbocycle and the fused heterocycle are unsubstituted or carry 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 radicals R^q;

n is 0, 1 or 2;

k is 0, 1 or 2;

R', R¹¹, R²¹ independently of each other are selected from the group consisting of halogen, NO₂, CN, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy and C₃-C₇-cycloalkoxy or two vicinal radicals R', R¹¹ or R²¹ together may form a group =O (oxo);

Z, Z¹, Z² independently of each other are selected from the group consisting of a covalent bond and C₁-C₄-alkanediyl;

Z^{2a} is selected from the group consisting of a covalent bond, C₁-C₄-alkanediyl, O-C₁-C₄-alkanediyl, C₁-C₄-alkanediyl-O and C₁-C₄-alkanediyl-O-C₁-C₄-alkanediyl;

R^a is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radi-

cals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^b, R^{1b}, R^{2b} independently of each other are selected from the group consisting of C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, phenyl and heterocyclyl, heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^c, R^{2c} independently of each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl, benzyl or heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl, benzyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^d, R^{2d} independently of each other are selected from the group consisting of C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^e, R^f independently of each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, or

R^e, R^f together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^{2e}, R^{2f} independently of each other have the meanings given for R^e, R^f;

R^g is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^h is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, a radical C(=O)-R^k, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, or

R^g, R^h together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of =O, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^{2g}, R^{2h} independently of each other have the meanings given for R^g, R^h;

R^k has the meanings given for R^c;

R^q is selected from the group consisting of halogen, Z^q-OH, Z^q-NO₂, Z^q-cyano, oxo (=O), =N-R^{q1}, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, Z^q-C₁-C₄-alkoxy, Z^q-C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, Z^q-C₁-C₄-haloalkoxy, Z^q-C₃-C₁₀-cycloalkyl, O-Z^q-C₃-C₁₀-cycloalkyl, Z^q-(tri-C₁-C₄-alkyl)silyl, Z^q-S(O)_k-R^{q2}, Z²-C(=O)-R^{q3}, Z²-NR^{q4}R^{q5} and Z^q-phenyl, where phenyl in Z^q-phenyl is unsubstituted or substituted by 1, 2, 3 or 4 groups R^{q6}, which are identical or different; where

Z^q has one of the meanings given for Z;

R^{q1} C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and C₃-C₇-cycloalkoxy, which is unsubstituted or partially or completely halogenated;

R^{q2} has one of the meanings given for R^b;

R^{q3} has one of the meanings given for R^c;

R^{q4}, R^{q5} independently of each other have the meanings given for R^a, R^h;

R^{q6} has one of the meanings given for Rⁱ;

or an N-oxide or an agriculturally suitable salt thereof.

The compounds of the present invention, i.e. the compounds of formula I, their N-oxides, or their salts are particularly useful for controlling unwanted vegetation. Therefore, the invention also relates to the use of a compound of the present invention, an N-oxide or a salt thereof for combating or controlling unwanted vegetation.

The invention also relates to a composition comprising at least one compound according to the invention, including an N-oxide or a salt thereof, and at least one auxiliary. In particular, the invention relates to an agricultural composition comprising at least one compound according to the invention including an N-oxide or an agriculturally suitable salt thereof, and at least one auxiliary customary for crop protection formulations.

The present invention also relates to a method for combating or controlling unwanted vegetation, which method comprises allowing a herbicidally effective amount of at least one compound according to the invention, including an N-oxide or a salt thereof, to act on unwanted plants, their seed and/or their habitat.

The present invention also relates to the use of a composition according to the invention including an N-oxide or an agriculturally suitable salt thereof for combating or controlling unwanted vegetation.

Depending on the substitution pattern, the compounds of the formula I may have one or more centers of chirality, in which case they are present as mixtures of enantiomers or diastereomers. The invention provides both the pure enantiomers or pure diastereomers of the compounds of formula I, and their mixtures and the use according to the invention of the pure enantiomers or pure diastereomers of the compound of formula I or its mixtures. Suitable compounds of the formula I also include all possible geometrical stereoisomers (cis/trans isomers) and mixtures thereof. Cis/trans isomers may be present with respect to an alkene, carbon-nitrogen double-bond, nitrogen-sulfur double bond or amide group. The term "stereoisomer(s)" encompasses both optical isomers, such as enantiomers or diastereomers, the latter existing due to more than one center of chirality in the molecule, as well as geometrical isomers (cis/trans isomers).

Depending on the substitution pattern, the compounds of the formula I may be present in the form of their tautomers. Hence the invention also relates to the tautomers of the formula I and the stereoisomers, salts and N-oxides of said tautomers.

The term "N-oxide" includes any compound of the present invention which has at least one tertiary nitrogen atom that is oxidized to an N-oxide moiety. N-oxides of compounds I can in particular be prepared by oxidizing the ring nitrogen atom(s) of the oxadiazole ring or the ring nitrogen atom(s) of the six-membered aromatic ring with a suitable oxidizing agent, such as peroxocarboxylic acids or other peroxides.

The present invention moreover relates to compounds as defined herein, wherein one or more of the atoms depicted in formula I have been replaced by its stable, preferably non-radioactive isotope (e.g., hydrogen by deuterium, ^{12}C by ^{13}C , ^{14}N by ^{15}N , ^{16}O by ^{18}O) and in particular wherein at least one hydrogen atom has been replaced by a deuterium atom. Of course, the compounds according to the invention contain more of the respective isotope than this naturally occurs and thus is anyway present in the compounds I.

The compounds of the present invention may be amorphous or may exist in one or more different crystalline states (polymorphs) which may have different macroscopic properties such as stability or show different biological properties such as activities. The present invention includes both amorphous and crystalline compounds of formula I, their enantiomers or diastereomers, mixtures of different crystalline states of the respective compound of formula I, its enantiomers or diastereomers, as well as amorphous or crystalline salts thereof.

Salts of the compounds of the present invention are agriculturally suitable salts. They can be formed in a customary method, e.g. by reacting the compound with an acid if the compound of the present invention has a basic functionality or by reacting the compound with a suitable base if the compound of the present invention has an acidic functionality.

Useful agriculturally suitable salts are especially the salts of those cations or the acid addition salts of those acids whose cations and anions, respectively, do not have any adverse effect on the herbicidal action of the compounds according to the present invention. Suitable cations are in particular the ions of the alkali metals, preferably lithium, sodium and potassium, of the alkaline earth metals, preferably calcium, magnesium and barium, and of the transition metals, preferably manganese, copper, zinc and iron, and also ammonium (NH_4^+) and substituted ammonium in which one to four of the hydrogen atoms are replaced by $\text{C}_1\text{-C}_4$ -alkyl, $\text{C}_1\text{-C}_4$ -hydroxyalkyl, $\text{C}_1\text{-C}_4$ -alkoxy, $\text{C}_1\text{-C}_4$ -alkoxy- $\text{C}_1\text{-C}_4$ -alkyl, hydroxy- $\text{C}_1\text{-C}_4$ -alkoxy- $\text{C}_1\text{-C}_4$ -alkyl, phenyl or benzyl. Examples of substituted ammonium ions comprise methylammonium, isopropylammonium, dimethylammonium, diisopropylammonium, trimethylammonium, tetramethylammonium, tetraethylammonium, tetrabutylammonium, 2-hydroxyethylammonium, 2-(2-hydroxyethoxy)ethylammonium, bis(2-hydroxyethyl)ammonium, benzyltrimethylammonium and benzyltriethylammonium, furthermore phosphonium ions, sulfonium ions, preferably tri($\text{C}_1\text{-C}_4$ -alkyl)sulfonium, and sulfoxonium ions, preferably tri($\text{C}_1\text{-C}_4$ -alkyl)sulfoxonium.

Anions of useful acid addition salts are primarily chloride, bromide, fluoride, hydrogensulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, phosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate, and the anions of $\text{C}_1\text{-C}_4$ -alkanoic acids, preferably formate, acetate, propionate and butyrate. They can be formed

by reacting compounds of the present invention with an acid of the corresponding anion, preferably with hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid or nitric acid.

The term "undesired vegetation" is understood to include any vegetation growing at a crop plant site or locus of seeded and otherwise desired crop, where the vegetation is any plant species, including their germinant seeds, emerging seedlings and established vegetation, other than the seeded or desired crop.

The organic moieties mentioned in the above definitions of the variables are - like the term halogen - collective terms for individual listings of the individual group members. The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group.

The term "halogen" denotes in each case fluorine, bromine, chlorine or iodine, in particular fluorine, chlorine or bromine.

The term "partially or completely halogenated" will be taken to mean that 1 or more, e.g. 1, 2, 3, 4 or 5 or all of the hydrogen atoms of a given radical have been replaced by a halogen atom, in particular by fluorine or chlorine. A partially or completely halogenated radical is termed below also "halo-radical". For example, partially or completely halogenated alkyl is also termed haloalkyl.

The term "alkyl" as used herein (and in the alkyl moieties of other groups comprising an alkyl group, e.g. alkoxy, alkylamino, alkylcarbonyl, alkoxy carbonyl, alkylthio, alkylsulfonyl and alkoxyalkyl) denotes in each case a straight-chain or branched alkyl group having usually from 1 to 10 carbon atoms, e.g. from 1 to 8 carbon atoms, frequently from 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms and in particular from 1 to 3 carbon atoms. Examples of C₁-C₄-alkyl are methyl, ethyl, n-propyl, iso-propyl, n-butyl, 2-butyl (sec-butyl), isobutyl and tert-butyl. Examples for C₁-C₆-alkyl are, apart those mentioned for C₁-C₄-alkyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl. Examples for C₁-C₁₀-alkyl are, apart those mentioned for C₁-C₆-alkyl, n-heptyl, 1-methylhexyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 1-ethylpentyl, 2-ethylpentyl, 3-ethylpentyl, n-octyl, 1-methyloctyl, 2-methylheptyl, 1-ethylhexyl, 2-ethylhexyl, 1,2-dimethylhexyl, 1-propylpentyl, 2-propylpentyl, nonyl, decyl, 2-propylheptyl and 3-propylheptyl.

The term "alkylene" (or alkanediyl) as used herein in each case denotes an alkyl radical as defined above, wherein one hydrogen atom at any position of the carbon backbone is replaced by one further binding site, thus forming a bivalent moiety.

The term "haloalkyl" as used herein (and in the haloalkyl moieties of other groups comprising a haloalkyl group, e.g. haloalkoxy, haloalkylthio, haloalkylcarbonyl, haloalkylsulfonyl and haloalkylsulfinyl) denotes in each case a straight-chain or branched alkyl group having usually from 1 to 8 carbon atoms ("C₁-C₈-haloalkyl"), frequently from 1 to 6 carbon atoms ("C₁-C₆-haloalkyl"), more frequently 1 to 4 carbon atoms ("C₁-C₁₀-haloalkyl"), wherein the hydrogen atoms of this group are partially or totally replaced with halogen atoms. Preferred haloalkyl moieties are selected from C₁-C₄-haloalkyl, more preferably from C₁-C₂-haloalkyl, more

preferably from halomethyl, in particular from C₁-C₂-fluoroalkyl. Halomethyl is methyl in which 1, 2 or 3 of the hydrogen atoms are replaced by halogen atoms. Examples are bromomethyl, chloromethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl and the like. Examples for C₁-C₂-fluoroalkyl are fluoromethyl, difluoromethyl, trifluoromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, and the like. Examples for C₁-C₂-haloalkyl are, apart those mentioned for C₁-C₂-fluoroalkyl, chloromethyl, dichloromethyl, trichloromethyl, bromomethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 2-chloroethyl, 2,2-dichloroethyl, 2,2,2-trichloroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 1-bromoethyl, and the like. Examples for C₁-C₄-haloalkyl are, apart those mentioned for C₁-C₂-haloalkyl, 1-fluoropropyl, 2-fluoropropyl, 3-fluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, heptafluoropropyl, 1,1,1-trifluoroprop-2-yl, 3-chloropropyl, 4-chlorobutyl and the like.

The term "cycloalkyl" as used herein (and in the cycloalkyl moieties of other groups comprising a cycloalkyl group, e.g. cycloalkoxy and cycloalkylalkyl) denotes in each case a mono- or bicyclic cycloaliphatic radical having usually from 3 to 10 carbon atoms ("C₃-C₁₀-cycloalkyl"), preferably 3 to 7 carbon atoms ("C₃-C₇-cycloalkyl") or in particular 3 to 6 carbon atoms ("C₃-C₆-cycloalkyl"). Examples of monocyclic radicals having 3 to 6 carbon atoms comprise cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of monocyclic radicals having 3 to 7 carbon atoms comprise cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Examples of bicyclic radicals having 7 or 8 carbon atoms comprise bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptyl, bicyclo[3.1.1]heptyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl and bicyclo[3.2.1]octyl.

The term "halocycloalkyl" as used herein (and in the halocycloalkyl moieties of other groups comprising an halocycloalkyl group, e.g. halocycloalkylmethyl) denotes in each case a mono- or bicyclic cycloaliphatic radical having usually from 3 to 10 carbon atoms, preferably 3 to 7 carbon atoms or in particular 3 to 6 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or 5 of the hydrogen atoms are replaced by halogen, in particular by fluorine or chlorine. Examples are 1- and 2-fluorocyclopropyl, 1,2-, 2,2- and 2,3-difluorocyclopropyl, 1,2,2-trifluorocyclopropyl, 2,2,3,3-tetrafluorocyclopropyl, 1- and 2-chlorocyclopropyl, 1,2-, 2,2- and 2,3-dichlorocyclopropyl, 1,2,2-trichlorocyclopropyl, 2,2,3,3-tetrachlorocyclopropyl, 1-, 2- and 3-fluorocyclopentyl, 1,2-, 2,2-, 2,3-, 3,3-, 3,4-, 2,5-difluorocyclopentyl, 1-, 2- and 3-chlorocyclopentyl, 1,2-, 2,2-, 2,3-, 3,3-, 3,4-, 2,5-dichlorocyclopentyl and the like.

The term "cycloalkyl-alkyl" used herein denotes a cycloalkyl group, as defined above, which is bound to the remainder of the molecule via an alkylene group. The term "C₃-C₇-cycloalkyl-C₁-C₄-alkyl" refers to a C₃-C₇-cycloalkyl group as defined above which is bound to the remainder of the molecule via a C₁-C₄-alkyl group, as defined above. Examples are cyclopropylmethyl, cyclopropylethyl, cyclopropylpropyl, cyclobutylmethyl, cyclobutylethyl, cyclobutylpropyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexylpropyl, and the like.

The term "alkenyl" as used herein denotes in each case a monounsaturated straight-chain or branched hydrocarbon radical having usually 2 to 8 ("C₂-C₈-alkenyl"), preferably 2 to 6 carbon

atoms ("C₂-C₆-alkenyl"), in particular 2 to 4 carbon atoms ("C₂-C₄-alkenyl"), and a double bond in any position, for example C₂-C₄-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl or 2-methyl-2-propenyl; C₂-C₆-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl, 1-ethyl-2-methyl-2-propenyl and the like, or C₂-C₈-alkenyl, such as the radicals mentioned for C₂-C₆-alkenyl and additionally 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 4-octenyl and the positional isomers thereof.

The term "haloalkenyl" as used herein, which may also be expressed as "alkenyl which may be substituted by halogen", and the haloalkenyl moieties in haloalkenyloxy and the like refers to unsaturated straight-chain or branched hydrocarbon radicals having 2 to 8 ("C₂-C₈-haloalkenyl") or 2 to 6 ("C₂-C₆-haloalkenyl") or 2 to 4 ("C₂-C₄-haloalkenyl") carbon atoms and a double bond in any position, where some or all of the hydrogen atoms in these groups are replaced by halogen atoms as mentioned above, in particular fluorine, chlorine and bromine, for example chlorovinyl, chloroallyl and the like.

The term "alkynyl" as used herein denotes unsaturated straight-chain or branched hydrocarbon radicals having usually 2 to 8 ("C₂-C₈-alkynyl"), frequently 2 to 6 ("C₂-C₆-alkynyl"), preferably 2 to 4 carbon atoms ("C₂-C₄-alkynyl") and one or two triple bonds in any position, for example C₂-C₄-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl and the like, C₂-C₆-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-2-butyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 3-methyl-1-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-1-pentyne, 3-methyl-4-pentyne, 4-methyl-1-pentyne, 4-methyl-2-pentyne, 1,1-dimethyl-2-butyne, 1,1-dimethyl-3-butyne, 1,2-dimethyl-3-butyne, 2,2-dimethyl-3-butyne, 3,3-dimethyl-1-butyne, 1-ethyl-2-butyne, 1-ethyl-3-butyne, 2-ethyl-3-butyne, 1-ethyl-1-methyl-2-propynyl and the like.

The term "haloalkynyl" as used herein, which is also expressed as "alkynyl which may be substituted by halogen", refers to unsaturated straight-chain or branched hydrocarbon radicals having usually 3 to 8 carbon atoms ("C₂-C₈-haloalkynyl"), frequently 2 to 6 ("C₂-C₆-haloalkynyl"), preferably 2 to 4 carbon atoms ("C₂-C₄-haloalkynyl"), and one or two triple bonds in any position (as mentioned above), where some or all of the hydrogen atoms in these groups are replaced by halogen atoms as mentioned above, in particular fluorine, chlorine and bromine.

The term "alkoxy" as used herein denotes in each case a straight-chain or branched alkyl group usually having from 1 to 8 carbon atoms ("C₁-C₈-alkoxy"), frequently from 1 to 6 carbon atoms ("C₁-C₆-alkoxy"), preferably 1 to 4 carbon atoms ("C₁-C₄-alkoxy"), which is bound to the remainder of the molecule via an oxygen atom. C₁-C₂-Alkoxy is methoxy or ethoxy. C₁-C₄-Alkoxy is additionally, for example, n-propoxy, 1-methylethoxy (isopropoxy), butoxy, 1-methylpropoxy (sec-butoxy), 2-methylpropoxy (isobutoxy) or 1,1-dimethylethoxy (tert-butoxy). C₁-C₆-Alkoxy is additionally, for example, pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy or 1-ethyl-2-methylpropoxy. C₁-C₈-Alkoxy is additionally, for example, heptyloxy, octyloxy, 2-ethylhexyloxy and positional isomers thereof.

The term "haloalkoxy" as used herein denotes in each case a straight-chain or branched alkoxy group, as defined above, having from 1 to 8 carbon atoms ("C₁-C₈-haloalkoxy"), frequently from 1 to 6 carbon atoms ("C₁-C₆-haloalkoxy"), preferably 1 to 4 carbon atoms ("C₁-C₄-haloalkoxy"), more preferably 1 to 3 carbon atoms ("C₁-C₃-haloalkoxy"), wherein the hydrogen atoms of this group are partially or totally replaced with halogen atoms, in particular fluorine atoms. C₁-C₂-Haloalkoxy is, for example, OCH₂F, OCHF₂, OCF₃, OCH₂Cl, OCHCl₂, OCCl₃, chlorofluoromethoxy, dichlorofluoromethoxy, chlorodifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 2-iodoethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-2-fluoroethoxy, 2-chloro-2,2-difluoroethoxy, 2,2-dichloro-2-fluoroethoxy, 2,2,2-trichloroethoxy or OC₂F₅. C₁-C₄-Haloalkoxy is additionally, for example, 2-fluoropropoxy, 3-fluoropropoxy, 2,2-difluoropropoxy, 2,3-difluoropropoxy, 2-chloropropoxy, 3-chloropropoxy, 2,3-dichloropropoxy, 2-bromopropoxy, 3-bromopropoxy, 3,3,3-trifluoropropoxy, 3,3,3-trichloropropoxy, OCH₂-C₂F₅, OCF₂-C₂F₅, 1-(CH₂F)-2-fluoroethoxy, 1-(CH₂Cl)-2-chloroethoxy, 1-(CH₂Br)-2-bromoethoxy, 4-fluorobutoxy, 4-chlorobutoxy, 4-bromobutoxy or nonafluorobutoxy. C₁-C₆-Haloalkoxy is additionally, for example, 5-fluoropentoxy, 5-chloropentoxy, 5-bromopentoxy, 5-iodopentoxy, undecafluoropentoxy, 6-fluorohexoxy, 6-chlorohexoxy, 6-bromohexoxy, 6-iodohexoxy or dodecafluorohexoxy.

The term "alkoxyalkyl" as used herein denotes in each case alkyl usually comprising 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms, wherein 1 carbon atom carries an alkoxy radical usually comprising 1 to 8, frequently 1 to 6, in particular 1 to 4, carbon atoms as defined above. "C₁-C₆-Alkoxy-C₁-C₆-alkyl" is a C₁-C₆-alkyl group, as defined above, in which one hydrogen atom is replaced by a C₁-C₆-alkoxy group, as defined above. Examples are CH₂OCH₃, CH₂-OC₂H₅, n-propoxymethyl, CH₂-OCH(CH₃)₂, n-butoxymethyl, (1-methylpropoxy)-methyl, (2-

methylpropoxy)methyl, $\text{CH}_2\text{-OC}(\text{CH}_3)_3$, 2-(methoxy)ethyl, 2-(ethoxy)ethyl, 2-(n-propoxy)-ethyl, 2-(1-methylethoxy)-ethyl, 2-(n-butoxy)ethyl, 2-(1-methylpropoxy)-ethyl, 2-(2-methylpropoxy)-ethyl, 2-(1,1-dimethylethoxy)-ethyl, 2-(methoxy)-propyl, 2-(ethoxy)-propyl, 2-(n-propoxy)-propyl, 2-(1-methylethoxy)-propyl, 2-(n-butoxy)-propyl, 2-(1-methylpropoxy)-propyl, 2-(2-methylpropoxy)-propyl, 2-(1,1-dimethylethoxy)-propyl, 3-(methoxy)-propyl, 3-(ethoxy)-propyl, 3-(n-propoxy)-propyl, 3-(1-methylethoxy)-propyl, 3-(n-butoxy)-propyl, 3-(1-methylpropoxy)-propyl, 3-(2-methylpropoxy)-propyl, 3-(1,1-dimethylethoxy)-propyl, 2-(methoxy)-butyl, 2-(ethoxy)-butyl, 2-(n-propoxy)-butyl, 2-(1-methylethoxy)-butyl, 2-(n-butoxy)-butyl, 2-(1-methylpropoxy)-butyl, 2-(2-methylpropoxy)-butyl, 2-(1,1-dimethylethoxy)-butyl, 3-(methoxy)-butyl, 3-(ethoxy)-butyl, 3-(n-propoxy)-butyl, 3-(1-methylethoxy)-butyl, 3-(n-butoxy)-butyl, 3-(1-methylpropoxy)-butyl, 3-(2-methylpropoxy)-butyl, 3-(1,1-dimethylethoxy)-butyl, 4-(methoxy)-butyl, 4-(ethoxy)-butyl, 4-(n-propoxy)-butyl, 4-(1-methylethoxy)-butyl, 4-(n-butoxy)-butyl, 4-(1-methylpropoxy)-butyl, 4-(2-methylpropoxy)-butyl, 4-(1,1-dimethylethoxy)-butyl and the like.

The term "haloalkoxy-alkyl" as used herein denotes in each case alkyl as defined above, usually comprising 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms, wherein 1 carbon atom carries an haloalkoxy radical as defined above, usually comprising 1 to 8, frequently 1 to 6, in particular 1 to 4, carbon atoms as defined above. Examples are fluoromethoxymethyl, difluoromethoxymethyl, trifluoromethoxymethyl, 1-fluoroethoxymethyl, 2-fluoroethoxymethyl, 1,1-difluoroethoxymethyl, 1,2-difluoroethoxymethyl, 2,2-difluoroethoxymethyl, 1,1,2-trifluoroethoxymethyl, 1,2,2-trifluoroethoxymethyl, 2,2,2-trifluoroethoxymethyl, pentafluoroethoxymethyl, 1-fluoroethoxy-1-ethyl, 2-fluoroethoxy-1-ethyl, 1,1-difluoroethoxy-1-ethyl, 1,2-difluoroethoxy-1-ethyl, 2,2-difluoroethoxy-1-ethyl, 1,1,2-trifluoroethoxy-1-ethyl, 1,2,2-trifluoroethoxy-1-ethyl, 2,2,2-trifluoroethoxy-1-ethyl, pentafluoroethoxy-1-ethyl, 1-fluoroethoxy-2-ethyl, 2-fluoroethoxy-2-ethyl, 1,1-difluoroethoxy-2-ethyl, 1,2-difluoroethoxy-2-ethyl, 2,2-difluoroethoxy-2-ethyl, 1,1,2-trifluoroethoxy-2-ethyl, 1,2,2-trifluoroethoxy-2-ethyl, 2,2,2-trifluoroethoxy-2-ethyl, pentafluoroethoxy-2-ethyl, and the like.

The term "alkylthio"(also alkylsulfanyl or alkyl-S-) as used herein denotes in each case a straight-chain or branched saturated alkyl group as defined above, usually comprising 1 to 8 carbon atoms ("C₁-C₈-alkylthio"), frequently comprising 1 to 6 carbon atoms ("C₁-C₆-alkylthio"), preferably 1 to 4 carbon atoms ("C₁-C₄-alkylthio"), which is attached via a sulfur atom at any position in the alkyl group. C₁-C₂-Alkylthio is methylthio or ethylthio. C₁-C₄-Alkylthio is additionally, for example, n-propylthio, 1-methylethylthio (isopropylthio), butylthio, 1-methylpropylthio (sec-butylthio), 2-methylpropylthio (isobutylthio) or 1,1-dimethylethylthio (tert-butylthio). C₁-C₆-Alkylthio is additionally, for example, pentylthio, 1-methylbutylthio, 2-methylbutylthio, 3-methylbutylthio, 1,1-dimethylpropylthio, 1,2-dimethylpropylthio, 2,2-dimethylpropylthio, 1-ethylpropylthio, hexylthio, 1-methylpentylthio, 2-methylpentylthio, 3-methylpentylthio, 4-methylpentylthio, 1,1-dimethylbutylthio, 1,2-dimethylbutylthio, 1,3-dimethylbutylthio, 2,2-dimethylbutylthio, 2,3-dimethylbutylthio, 3,3-dimethylbutylthio, 1-ethylbutylthio, 2-ethylbutylthio, 1,1,2-trimethylpropylthio, 1,2,2-trimethylpropylthio, 1-ethyl-1-methylpropylthio or 1-ethyl-2-methylpropylthio. C₁-C₈-Alkylthio is additionally, for example, heptylthio, octylthio, 2-ethylhexylthio and positional isomers thereof.

The term "haloalkylthio" as used herein refers to an alkylthio group as defined above wherein the hydrogen atoms are partially or completely substituted by fluorine, chlorine, bromine and/or iodine. C₁-C₂-Haloalkylthio is, for example, SCH₂F, SCHF₂, SCF₃, SCH₂Cl, SCHCl₂, SCCl₃, chlorofluoromethylthio, dichlorofluoromethylthio, chlorodifluoromethylthio, 2-fluoroethylthio, 2-chloroethylthio, 2-bromoethylthio, 2-iodoethylthio, 2,2-difluoroethylthio, 2,2,2-trifluoroethylthio, 2-chloro-2-fluoroethylthio, 2-chloro-2,2-difluoroethylthio, 2,2-dichloro-2-fluoroethylthio, 2,2,2-trichloroethylthio or SC₂F₅. C₁-C₄-Haloalkylthio is additionally, for example, 2-fluoropropylthio, 3-fluoropropylthio, 2,2-difluoropropylthio, 2,3-difluoropropylthio, 2-chloropropylthio, 3-chloropropylthio, 2,3-dichloropropylthio, 2-bromopropylthio, 3-bromopropylthio, 3,3,3-trifluoropropylthio, 3,3,3-trichloropropylthio, SCH₂-C₂F₅, SCF₂-C₂F₅, 1-(CH₂F)-2-fluoroethylthio, 1-(CH₂Cl)-2-chloroethylthio, 1-(CH₂Br)-2-bromoethylthio, 4-fluorobutylthio, 4-chlorobutylthio, 4-bromobutylthio or nonafluorobutylthio. C₁-C₆-Haloalkylthio is additionally, for example, 5-fluoropentylthio, 5-chloropentylthio, 5-bromopentylthio, 5-iodopentylthio, undecafluoropentylthio, 6-fluorohexylthio, 6-chlorohexylthio, 6-bromohexylthio, 6-iodohexylthio or dodecafluorohexylthio.

The terms "alkylsulfinyl" and "S(O)_n-alkyl" (wherein n is 1) are equivalent and, as used herein, denote an alkyl group, as defined above, attached via a sulfinyl [S(O)] group. For example, the term "C₁-C₂-alkylsulfinyl" refers to a C₁-C₂-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "C₁-C₄-alkylsulfinyl" refers to a C₁-C₄-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "C₁-C₆-alkylsulfinyl" refers to a C₁-C₆-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. C₁-C₂-alkylsulfinyl is methylsulfinyl or ethylsulfinyl. C₁-C₄-alkylsulfinyl is additionally, for example, n-propylsulfinyl, 1-methylethylsulfinyl (isopropylsulfinyl), butylsulfinyl, 1-methylpropylsulfinyl (sec-butylsulfinyl), 2-methylpropylsulfinyl (isobutylsulfinyl) or 1,1-dimethylethylsulfinyl (tert-butylsulfinyl). C₁-C₆-alkylsulfinyl is additionally, for example, pentylsulfinyl, 1-methylbutylsulfinyl, 2-methylbutylsulfinyl, 3-methylbutylsulfinyl, 1,1-dimethylpropylsulfinyl, 1,2-dimethylpropylsulfinyl, 2,2-dimethylpropylsulfinyl, 1-ethylpropylsulfinyl, hexylsulfinyl, 1-methylpentylsulfinyl, 2-methylpentylsulfinyl, 3-methylpentylsulfinyl, 4-methylpentylsulfinyl, 1,1-dimethylbutylsulfinyl, 1,2-dimethylbutylsulfinyl, 1,3-dimethylbutylsulfinyl, 2,2-dimethylbutylsulfinyl, 2,3-dimethylbutylsulfinyl, 3,3-dimethylbutylsulfinyl, 1-ethylbutylsulfinyl, 2-ethylbutylsulfinyl, 1,1,2-trimethylpropylsulfinyl, 1,2,2-trimethylpropylsulfinyl, 1-ethyl-1-methylpropylsulfinyl or 1-ethyl-2-methylpropylsulfinyl.

The terms "alkylsulfonyl" and "S(O)_n-alkyl" (wherein n is 2) are equivalent and, as used herein, denote an alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "C₁-C₂-alkylsulfonyl" refers to a C₁-C₂-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "C₁-C₄-alkylsulfonyl" refers to a C₁-C₄-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "C₁-C₆-alkylsulfonyl" refers to a C₁-C₆-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. C₁-C₂-alkylsulfonyl is methylsulfonyl or ethylsulfonyl. C₁-C₄-alkylsulfonyl is additionally, for example, n-propylsulfonyl, 1-methylethylsulfonyl (isopropylsulfonyl), butylsulfonyl, 1-methylpropylsulfonyl (sec-butylsulfonyl), 2-methylpropylsulfonyl (isobutylsulfonyl) or 1,1-dimethylethylsulfonyl (tert-butylsulfonyl). C₁-C₆-alkylsulfonyl is additionally, for example, pentylsulfonyl,

1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, hexylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl, 3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl or 1-ethyl-2-methylpropylsulfonyl.

The term "alkylamino" as used herein denotes in each case a group -NHR^* , wherein R^* is a straight-chain or branched alkyl group usually having from 1 to 6 carbon atoms (" $\text{C}_1\text{-C}_6$ -alkylamino"), preferably 1 to 4 carbon atoms (" $\text{C}_1\text{-C}_4$ -alkylamino"). Examples of $\text{C}_1\text{-C}_6$ -alkylamino are methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, 2-butylamino, isobutylamino, tert-butylamino, and the like.

The term "dialkylamino" as used herein denotes in each case a group $\text{-NR}^*\text{R}^\circ$, wherein R^* and R° , independently of each other, are a straight-chain or branched alkyl group each usually having from 1 to 6 carbon atoms (" $\text{di-(C}_1\text{-C}_6\text{-alkyl)-amino}$ "), preferably 1 to 4 carbon atoms (" $\text{di-(C}_1\text{-C}_4\text{-alkyl)-amino}$ "). Examples of a $\text{di-(C}_1\text{-C}_6\text{-alkyl)-amino}$ group are dimethylamino, diethylamino, dipropylamino, dibutylamino, methyl-ethyl-amino, methyl-propyl-amino, methyl-isopropylamino, methyl-butyl-amino, methyl-isobutyl-amino, ethyl-propyl-amino, ethyl-isopropylamino, ethyl-butyl-amino, ethyl-isobutyl-amino, and the like.

The suffix "-carbonyl" in a group denotes in each case that the group is bound to the remainder of the molecule via a carbonyl group (C=O). This is the case e.g. in alkylcarbonyl, haloalkylcarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonyl, haloalkoxycarbonyl.

The term "aryl" as used herein refers to a mono-, bi- or tricyclic aromatic hydrocarbon radical such as phenyl or naphthyl, in particular phenyl.

The term "het(ero)aryl" as used herein refers to a mono-, bi- or tricyclic heteroaromatic hydrocarbon radical, preferably to a monocyclic heteroaromatic radical, such as pyridyl, pyrimidyl and the like.

The term "3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, unsaturated or aromatic heterocycle containing 1, 2, 3 or 4 heteroatoms as ring members selected from the groups consisting of N, O and S" as used herein denotes monocyclic or bicyclic radicals, the monocyclic or bicyclic radicals being saturated, unsaturated or aromatic where N can optionally be oxidized, i.e. in the form of an N-oxide, and S can also optionally be oxidized to various oxidation states, i.e. as SO or SO_2 . An unsaturated heterocycle contains at least one C-C and/or C-N and/or N-N double bond(s). A fully unsaturated heterocycle contains as many conjugated C-C and/or C-N and/or N-N double bonds as allowed by the size(s) of the ring(s). An aromatic monocyclic heterocycle is a fully unsaturated 5- or 6-membered monocyclic heterocycle. An aromatic bicyclic heterocycle is an 8-, 9- or 10-membered bicyclic heterocycle consisting of a 5- or 6-membered heteroaromatic ring which is fused to a phenyl ring or to another 5- or 6-membered heteroaromatic ring. The heterocycle may be attached to the remainder of the molecule via a carbon ring member or via a nitrogen

ring member. As a matter of course, the heterocyclic ring contains at least one carbon ring atom. If the ring contains more than one O ring atom, these are not adjacent.

Examples of a 3-, 4-, 5- or 6-membered monocyclic saturated heterocycle include:

oxirane-2-yl, aziridine-1-yl, aziridine-2-yl, oxetan-2-yl, azetidine-1-yl, azetidine-2-yl, azetidine-3-yl, thietane-1-yl, thietan-2-yl, thietane-3-yl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrazolidin-1-yl, pyrazolidin-3-yl, pyrazolidin-4-yl, pyrazolidin-5-yl, imidazolidin-1-yl, imidazolidin-2-yl, imidazolidin-4-yl, oxazolidin-2-yl, oxazolidin-3-yl, oxazolidin-4-yl, oxazolidin-5-yl, isoxazolidin-2-yl, isoxazolidin-3-yl, isoxazolidin-4-yl, isoxazolidin-5-yl, thiazolidin-2-yl, thiazolidin-3-yl, thiazolidin-4-yl, thiazolidin-5-yl, isothiazolidin-2-yl, isothiazolidin-3-yl, isothiazolidin-4-yl, isothiazolidin-5-yl, 1,2,4-oxadiazolidin-3-yl, 1,2,4-oxadiazolidin-5-yl, 1,2,4-thiadiazolidin-3-yl, 1,2,4-thiadiazolidin-5-yl, 1,2,4-triazolidin-3-yl, 1,3,4-oxadiazolidin-2-yl, 1,3,4-thiadiazolidin-2-yl, 1,3,4-triazolidin-1-yl, 1,3,4-triazolidin-2-yl, 2-tetrahydropyranyl, 4-tetrahydropyranyl, 1,3-dioxan-5-yl, 1,4-dioxan-2-yl, piperidin-1-yl, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, hexahydropyridazin-3-yl, hexahydropyridazin-4-yl, hexahydropyrimidin-2-yl, hexahydropyrimidin-4-yl, hexahydropyrimidin-5-yl, piperazin-1-yl, piperazin-2-yl, 1,3,5-hexahydrotriazin-1-yl, 1,3,5-hexahydrotriazin-2-yl and 1,2,4-hexahydrotriazin-3-yl, morpholin-2-yl, morpholin-3-yl, morpholin-4-yl, thiomorpholin-2-yl, thiomorpholin-3-yl, thiomorpholin-4-yl, 1-oxothiomorpholin-2-yl, 1-oxothiomorpholin-3-yl, 1-oxothiomorpholin-4-yl, 1,1-dioxothiomorpholin-2-yl, 1,1-dioxothiomorpholin-3-yl, 1,1-dioxothiomorpholin-4-yl, azepan-1-, -2-, -3- or -4-yl, oxepan-2-, -3-, -4- or -5-yl, hexahydro-1,3-diazepinyl, hexahydro-1,4-diazepinyl, hexahydro-1,3-oxazepinyl, hexahydro-1,4-oxazepinyl, hexahydro-1,3-dioxepinyl, hexahydro-1,4-dioxepinyl and the like.

Examples of a 5- or 6-membered monocyclic partially unsaturated heterocycle include:

2,3-dihydrofur-2-yl, 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 2-pyrrolin-2-yl, 2-pyrrolin-3-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl, 2-isoxazolin-3-yl, 3-isoxazolin-3-yl, 4-isoxazolin-3-yl, 2-isoxazolin-4-yl, 3-isoxazolin-4-yl, 4-isoxazolin-4-yl, 2-isoxazolin-5-yl, 3-isoxazolin-5-yl, 4-isoxazolin-5-yl, 2-isothiazolin-3-yl, 3-isothiazolin-3-yl, 4-isothiazolin-3-yl, 2-isothiazolin-4-yl, 3-isothiazolin-4-yl, 4-isothiazolin-4-yl, 2-isothiazolin-5-yl, 3-isothiazolin-5-yl, 4-isothiazolin-5-yl, 2,3-dihydropyrazol-1-yl, 2,3-dihydropyrazol-2-yl, 2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl, 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-1-yl, 3,4-dihydropyrazol-3-yl, 3,4-dihydropyrazol-4-yl, 3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-1-yl, 4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl, 4,5-dihydropyrazol-5-yl, 2,3-dihydrooxazol-2-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl, 2,3-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 3,4-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 2-, 3-, 4-, 5- or 6-di- or tetrahydropyridinyl, 3-di- or tetrahydropyridazinyl, 4-di- or tetrahydropyridazinyl, 2-di- or tetrahydropyrimidinyl, 4-di- or tetrahydropyrimidinyl, 5-di- or tetrahydropyrimidinyl, di- or tetrahydropyrazinyl, 1,3,5-di- or tetrahydrotriazin-2-yl, 1,2,4-di- or tetrahydrotriazin-3-yl,

Examples of a 5- or 6-membered monocyclic aromatic heterocyclic ring are: 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-

pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1,3,4-triazol-1-yl, 1,3,4-triazol-2-yl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 1-oxopyridin-2-yl, 1-oxopyridin-3-yl, 1-oxopyridin-4-yl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl and 2-pyrazinyl.

5 Examples of a 5- or 6-membered heteroaromatic ring fused to a phenyl ring or to a 5- or 6-membered heteroaromatic radical include benzofuranyl, benzothienyl, indolyl, indazolyl, benzimidazolyl, benzoxathiazolyl, benzoxadiazolyl, benzothiadiazolyl, benzoxazinyl, chinolinyl, isochinolinyl, purinyl, 1,8-naphthyridyl, pteridyl, pyrido[3,2-d]pyrimidyl or pyridoimidazolyl and the like.

10 The remarks made below as to preferred embodiments of the variables (substituents) of the compounds of formula I are valid on their own as well as preferably in combination with each other, as well as in combination with the stereoisomers, salts, tautomers or N-oxides thereof.

The remarks made below concerning preferred embodiments of the variables further are
15 valid on their own as well as preferably in combination with each other concerning the compounds of formulae I, where applicable, as well as concerning the uses and methods according to the invention and the composition according to the invention.

Preferred compounds according to the invention are compounds of formula I or a stereoisomer, salt, tautomer or N-oxide thereof, wherein the salt is an agriculturally suitable salt.
20 Further preferred compounds according to the invention are compounds of formula I or a N-oxide or an agriculturally suitable salt thereof. Particularly preferred compounds according to the invention are compounds of formula I or an agriculturally suitable salt thereof.

According to a preferred embodiment of the invention the variable R in the compounds of formula I is selected from the group consisting of hydrogen, cyano, nitro, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C(=O)-R^c, C(=O)-OR^d, C(=O)-NR^eR^f, NH-C(=O)R^k and NR^gR^h; where R^c, R^d, R^e, R^f, R^k, R^g and R^h are as defined above and which preferably have on their own or in particular in combination the following meanings:

R^c hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, or phenyl, in particular C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R^d C₁-C₆-alkyl or C₁-C₆-haloalkyl, in particular C₁-C₄-alkyl,

R^e, R^f are independently of each other selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl and benzyl, and in particular from the group consisting of hydrogen and C₁-C₄-alkyl, or

R^e, R^f together with the nitrogen atom, to which they are bound form a 5-, 6- or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl, and in particular R^e, R^f together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 methyl groups;

R^g, R^h are independently of each other selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl and benzyl and in particular from the group consisting of hydrogen or C₁-C₄-alkyl, or

R^g, R^h together with the nitrogen atom, to which they are bound form a 5-, 6 or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl, and in particular R^g, R^h together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 methyl groups;

R^k hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl or phenyl, in particular C₁-C₄-alkyl.

According to a more preferred embodiment the variable R of the compounds of the formula I is selected from the group consisting of halogen, cyano, nitro, NH₂, C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C₁-C₄-haloalkyl, C(=O)-R^c, C(=O)-OR^d, C(=O)-NR^eR^f and NH-C(=O)R^k, where R^c, R^d, R^e, R^f and R^k are as defined above and which preferably have on their own or in particular in combination the following meanings:

R^c is C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R^d is C₁-C₄-alkyl;

R^e is hydrogen or C₁-C₄-alkyl;

R^f is hydrogen or C₁-C₄-alkyl, or

R^e, R^f together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 methyl groups, and

R^k is C₁-C₄-alkyl.

R is also preferably selected from hydrogen and C(=O)-R^c, where R^c is as defined above, and where R^c is in particular C₁-C₄-alkyl, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, 2-butyl or tert.-butyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl; C₂-C₆-alkenyl, such as CH₂CH=CH₂ or CH₂CH=CH₂; C₂-C₆-alkynyl such as 2-propynyl; phenyl; phenyl which is substituted by 1, 2 or 3 groups selected from halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, in particular halogen and C₁-C₄-alkyl such as 2-methylphenyl, 4-methylphenyl, 2-ethylphenyl, 4-ethylphenyl, 2-fluorophenyl or 4-fluorophenyl; heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where heterocyclyl is unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, in particular 5- or 6-membered hetaryl having 1 or 2 heteroatoms selected from O, S and N as ring members such as pyridyl or 5- or 6-membered hetaryl having 1 or 2 heteroatoms selected from O, S and N as ring members and where hetaryl carries 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-

haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, such as pyridyl substituted by 1, 2 or 3 groups selected from C₁-C₄-alkyl. R is also preferably NH-C(=O)R^k, where R^k is phenyl

According to particular preferred embodiments of the invention the variable R in the compounds of formula I is selected from hydrogen, halogen; cyano; nitro; amino; C₁-C₄-alkyl; C₁-C₄-haloalkyl; C₃-C₇-cycloalkyl; C₁-C₄-alkoxy-C₁-C₄-alkyl, C(=O)-C₁-C₄-alkyl; C(=O)-C₁-C₄-haloalkyl; C(=O)-C₃-C₆-cycloalkyl; C(=O)-C₂-C₆-alkenyl; C(=O)-C₂-C₆-alkynyl; benzoyl; benzoyl, which is substituted by 1 or 2 radicals selected from C₁-C₄-alkyl and halogen; C(=O)-hetaryl, where hetaryl has 1 nitrogen atom as ring member; C(=O)-O-C₁-C₄-alkyl; C(=O)NH₂; C(=O)NH(C₁-C₄-alkyl); C(=O)N(C₁-C₄-alkyl)₂; C(=O)-NR^eR^f, where R^e and R^f together with the nitrogen atom, to which they are bound may form a 5- or 6-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 methyl groups; benzoylamino; and NH-C(=O)-C₁-C₄-alkyl.

In this embodiment, specific examples for the variable R in the compounds of the formula I are H, Cl, Br, F, cyano, nitro, amino, methyl, ethyl, n-propyl, isopropyl, tert-butyl, CF₃, CHF₂, CH₂F, CH₂CF₃, CF₂CF₃, CH₂Cl, CHCl₂, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxyethyl, methoxymethyl, methylcarbonyl, ethylcarbonyl, isopropylcarbonyl, tert-butylcarbonyl, trifluoromethylcarbonyl, cyclopropylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, allylcarbonyl, (2-propynyl)carbonyl, benzoyl, 2-methylbenzoyl, 4-methylbenzoyl, 2-fluorobenzoyl, 4-fluorobenzoyl, pyridine-2-carbonyl, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, piperidinylcarbonyl, methoxycarbonyl, ethoxycarbonyl, acetylamino and benzoylamino.

According to another preferred embodiment of the invention the variable R in the compounds of formula I is phenyl or heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R' as defined above which independently from one another are preferably selected from the group consisting of halogen, C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₁-C₆-haloalkyloxy. R' is in particular selected from halogen, C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy, and in particular from Cl, F, Br, methyl, ethyl, methoxy and trifluoromethyl.

More preferably, R is phenyl or heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R', where R' is as defined above and in particular selected from the group consisting of halogen, methyl, ethyl, methoxy and trifluoromethyl.

According to a more preferred embodiment of the invention the variable R in the compounds of formula I is phenyl which is unsubstituted or substituted by 1, 2, 3 or 4 groups R',

where R' is as defined above and in particular selected from the group consisting of halogen, methyl, ethyl, methoxy and trifluoromethyl.

According to a further more preferred embodiment of the invention the variable R in the compounds of formula I or heterocyclyl, where heterocyclyl is a saturated, partially unsaturated or aromatic 5- or 6-membered monocyclic or 9- or 10-membered bicyclic heterocycle containing 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the bicyclic heterocycle consists of a 5- or 6-membered heteroaromatic ring which is fused to a phenyl ring, and where heterocyclyl is unsubstituted or substituted by 1, 2, 3 or 4 groups R' which is as defined above and in particular independently from one another selected from the group consisting of halogen, C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₁-C₆-haloalkyloxy. R' is in particular selected from halogen, C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy, and especially from Cl, F, Br, methyl, ethyl, methoxy and trifluoromethyl. .

According to particular preferred embodiments the variable R in the compounds of the formula I is heterocyclyl selected from tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, furan-2-yl, furan-3-yl, thiophen-2-yl, thiophen-3-yl, pyrrol-2-yl, pyrrol-3-yl, pyrazol-3-yl, pyrazol-4-yl, pyrazol-5-yl, imidazol-2-yl, imidazol-4-yl, imidazol-5-yl, isoxazol-2-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, oxazol-2-yl, oxazol-3-yl, oxazol-4-yl, oxazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, 1,2,3-triazol-4-yl, 1,2,3-triazol-5-yl, 1,2,5-triazol-3-yl, 1,3,4-triazol-2-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,2,3-oxadiazol-4-yl, 1,2,3-oxadiazol-5-yl, 1,2,5-oxadiazol-3-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,3,4-thiadiazol-2-yl, 1,2,3-thiadiazol-4-yl, 1,2,3-thiadiazol-5-yl, 1,2,5-thiadiazol-3-yl, 2H-1,2,3,4-tetrazol-5-yl, 1H-1,2,3,4-tetrazol-1-yl, 1,2,3,4-oxatriazol-5-yl, 1,2,3,5-oxatriazol-4-yl, 1,2,3,4-thiatriazol-5-yl, 1,2,3,5-thiatriazol-4-yl, benzisoxazole-2-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyrazin-2-yl, pyrazin-3-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, pyridazin-3-yl and pyridazin-4-yl, where heterocyclyl is unsubstituted or carries 1, 2, or 3 groups R' which independently from one another have the aforementioned preferred meanings.

According to a further preferred embodiment of the invention the variable R in the compounds of formula I is C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl or C₂-C₆-haloalkynyl. In this embodiment, R is in particular CH₂CH=CH₂ or 2-propynyl.

According to a further preferred embodiment of the invention the variable R in the compounds of formula I is a radical OR^a, where R^a is selected from the group consisting of H, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated. R^a may also be selected from C₂-C₆-haloalkynyl. More preferably, R^a is selected from H, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₆-cycloalkyl. In this embodiment R^a specifically is H, methyl, ethyl, n-propyl, isopropyl, C(CH₃)₃, CH₂Cl, CHF₂, CF₃, CH₂CH=CH₂, 2-propynyl, CH₂OCH₃, CH₂CH₂OCH₃, CH₂CH₂OCH₂CH₃, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl.

According to a further preferred embodiment of the invention the variable R in the compounds of formula I is $S(O)_n-R^b$, where n is 0, 1 or 2 and R^b is selected from the group consisting of C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₃-C₇-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2 or 3 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2 or 3 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₂-haloalkyl and C₁-C₂-alkoxy.

According to a more preferred embodiment of the invention the variable R in the compounds of formula I is $S(O)_n-R^b$, where n is 0, 1 or 2 and R^b is selected from C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₃-C₇-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 6-membered aromatic heterocyclic radical having 1 or 2 nitrogen atoms as ring members.

According to a particularly preferred embodiment of the invention the variable R in the compounds of formula I is $S-R^b$, $S(O)-R^b$ or $S(O)_2-R^b$, in which each R^b is as defined above and in particular selected from C₁-C₆-alkyl, C₂-C₆-alkenyl and C₂-C₆-alkynyl, and especially selected from CH₃, CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₃, CH₂CH=CH₂, 2-propynyl and phenyl.

In specific embodiments of the invention, the variable R in the compounds of formula I is selected from CH₃, CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₃, C(CH₃)₃, CH₂Cl, OCH₃, CF₃, CN, Cl and SO₂CH₃.

According to specifically preferred embodiments of the invention the variable R in the compounds of formula I is selected from the group consisting of methyl, ethyl, isopropyl, tert-butyl, cyclopropyl, cyclopentyl, cyclohexyl, CF₃, CHF₂, CClF₂, CH₂CF₃, CF₂CF₃, CH₂Cl, CHCl₂, methoxyethyl, methoxymethyl, and in particular from methyl and ethyl.

Preferred compounds according to the invention are compounds of formula I, wherein R^3 is selected from the group consisting of hydrogen, cyano, halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₂-C₄-alkenyloxy, C₂-C₄-alkynyloxy and $S(O)_kR^{2b}$, where the variables k and R^{2b} have one of the herein defined meanings.

More preferably, R^3 is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, $S(O)_2$ -C₁-C₄-alkyl and $S(O)_2$ -C₁-C₄-haloalkyl.

In particular, R^3 is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₂-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, C₁-C₂-haloalkoxy, C₁-C₂-alkylthio, C₁-C₂-haloalkylthio, $S(O)_2$ -C₁-C₂-alkyl and $S(O)_2$ -C₁-C₂-haloalkyl, and specifically from hydrogen, F, Cl, Br, CN, NO₂, CH₃, C₂H₅, CF₃, CHF₂, OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCHF₂, $S(O)_2$ CH₃ and $S(O)_2$ CH₂CH₃.

According to particular embodiments of the invention the variable R^3 in the compounds of formula I is selected from hydrogen, cyano, halogen, C₁-C₄-haloalkyl and C₁-C₄-alkylsulfonyl, such as hydrogen, cyano, chlorine, trifluoromethyl, difluoromethyl, $S(O)_2$ CH₃ and $S(O)_2$ CH₂CH₃.

In a specific group of embodiments R^3 in the compounds of formula I is selected from hydrogen, chlorine, fluorine, trifluoromethyl, methylsulfonyl and cyano.

Preferred compounds according to the invention are compounds of formula I, wherein R^5 is selected from the group consisting of hydrogen, halogen, CN, nitro, C_1 - C_4 -alkyl, C_3 - C_5 -cycloalkyl, C_3 - C_5 -cycloalkyl- C_1 - C_4 -alkyl, where the C_3 - C_5 -cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C_1 - C_4 -haloalkyl, C_1 - C_3 -alkylamino, C_1 - C_3 -dialkylamino, C_1 - C_3 -alkylamino- $S(O)_k$, C_1 - C_3 -alkylcarbonyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_3 - C_4 -alkenyloxy, C_3 - C_4 -alkynyloxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkoxy, C_1 - C_4 -alkyl- $S(O)_k$, C_1 - C_4 -haloalkyl- $S(O)_k$, where k is 0, 1 or 2; provided that if X^4 is N, R^5 is not hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl; and further provided that if X^4 is CR^4 and R^4 is hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl, R^5 is not hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl;

In particular, R^5 is selected from the group consisting of hydrogen, halogen, CN, nitro, C_1 - C_4 -alkyl, C_3 - C_4 -cycloalkyl, C_3 - C_4 -cycloalkyl- C_1 - C_2 -alkyl, where the C_3 - C_4 -cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C_1 - C_4 -haloalkyl, C_1 - C_3 alkylcarbonyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl; specifically R^5 is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH_3 , CH_2CH_3 , $CH(CH_3)_2$, cyclopropyl, cyclopropylmethyl, CF_3 , OCH_3 , OCH_2CH_3 , $OCH(CH_3)_2$, CH_2OCH_3 , $CH_2OCH_2CH_2OCH_3$, OCF_3 , $OCHF_2$, SCH_3 , SCF_3 , SCH_2CH_3 , SCH_2CF_3 , $SOCH_3$, $SOCH_2CH_3$, SO_2CH_3 , $SO_2CH_2CH_3$; and more specifically R^5 is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH_3 , CH_2CH_3 , $CH(CH_3)_2$, cyclopropyl, CF_3 , OCH_3 , OCF_3 , SCH_3 , $SOCH_3$, SO_2CH_3 or CH_2OCH_3 ; provided that if X^4 is N, R^5 is not hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl; and further provided that if X^4 is CR^4 and R^4 is hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl, R^5 is not hydrogen, halogen, cyano, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl;

A particular group of compounds according to the invention are compounds of formula I, wherein X^1 is CR^1 . R^1 is preferably selected from cyano, halogen, nitro, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl, Z^1 - C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, Z^1 - C_1 - C_4 -alkylthio- C_1 - C_4 -alkylthio, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyloxy, C_1 - C_6 -haloalkoxy, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkoxy and $S(O)_kR^{1b}$, where k , R^{1b} and Z^1 are as defined herein and where R^{1b} is in particular selected from the group consisting of C_1 - C_4 -alkyl and C_1 - C_4 -haloalkyl. In this context Z^1 is in particular a covalent bond. In this context k is in particular 2.

More preferably, R^1 is selected from cyano, nitro, halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkylthio, C_3 - C_4 -alkenyloxy, C_3 - C_4 -alkynyloxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkoxy, $S(O)_2$ - C_1 - C_4 -alkyl and $S(O)_2$ - C_1 - C_4 -

haloalkyl. According to a further more preferred embodiment, R¹ may also be selected from nitro and cyano.

Even more preferably, R¹ is selected from the group consisting of cyano, nitro, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl and C₁-C₄-alkoxy-C₁-C₄-alkoxy.

In particular, R¹ is selected from the group consisting of halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl. In particular, R¹ may also be selected from the group consisting of nitro, cyano, C₁-C₄-alkoxy-C₁-C₄-alkoxy and C₁-C₄-haloalkoxy-C₁-C₄-alkyl.

Specifically R¹ is F, Cl, Br, NO₂, CH₃, CF₃, OCH₃, SCH₃, OCF₃, SCF₃, SO₂CH₃, OCH₂CH₂OCH₃, CH₂OCH₂CH₂OCH₃ or CH₂OCH₂CF₃.

According to special embodiments of the invention the variable R¹ in the compounds of formula I is selected from halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl and C₁-C₄-alkylsulfonyl. Examples are chlorine, fluorine, bromine, nitro, cyano, methyl, trifluoromethyl and methylsulfonyl.

According to a further embodiment of the invention, R¹ is preferably selected from Z¹-phenoxy and Z¹-heterocycloxy, where heterocycloxy is an oxygen bound 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in phenoxy and heterocycloxy are unsubstituted or substituted by 1, 2, 3 or 4 groups R¹¹, which are identical or different.

Especially preferred are compounds of formula I, where

R¹ is selected from the group consisting of halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular from F, Cl, Br, nitro, CH₃, CF₃, OCH₃, SCH₃, OCF₃, SCF₃, SO₂CH₃, CH₂OCH₃ and CH₂OCH₂CH₂OCH₃; and

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular from H, Cl, Br, CN, NO₂, CH₃, CF₃, CHF₂, OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCHF₂, S(O)₂CH₃ and S(O)₂CH₂CH₃.

A further embodiment of the invention relates to compounds of formula I, to their N-oxides and their salts, wherein X¹ is N.

A further embodiment of the invention relates to compounds of formula I, to their N-oxides and their salts, wherein X² is CR². Preferred compounds according to the invention are compounds of formula I, wherein R² has any one of the meanings given above for R² with the exception of hydrogen.

Particular embodiments of the invention relate to compounds of the formula I, wherein X² is CR² and wherein the variable R² is Z^{2a}-heterocyclyl, where Z^{2a} is as defined herein and where heterocyclyl is a 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, par-


tially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in Z^{2a} -phenyl and Z^{2a} -heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R^{21} , which are identical or different.

In the compounds of the invention, wherein X^2 in formula I is CR^2 , the variable R^2 is preferably a 5- or 6-membered heterocyclyl, where heterocyclyl is a saturated, partially unsaturated or aromatic heterocyclic radical, which contains as ring member 1 heteroatom selected from the group consisting of O, N and S and 0, 1 or 2 further nitrogen atoms, where heterocyclyl is unsubstituted or carries 1, 2 or 3 radicals R^{21} which are identical or different. In this embodiment, R^{21} is preferably selected from halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio and C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl. In particular, R^{21} is selected from fluorine, chlorine methyl, ethyl, methoxy, ethoxy, methylsulfanyl, methylsulfonyl, methoxymethyl, ethoxymethyl, ethylsulfanylmethyl, ethylsulfanylethyl, methylsulfanylmethyl, methylsulfanylethyl, fluoromethyl, difluoromethyl and trifluoromethyl.

According to an even more preferred embodiment of the invention, X^2 is CR^2 and the variable R^2 is a 5- or 6-membered heterocyclyl selected from the group consisting of selected from the group consisting of isoxazolynyl, 1,2-dihydrotetrazolynyl, 1,4-dihydrotetrazolynyl, tetrahydrofuryl, dioxolanyl, piperidinyl, morpholynyl, piperazinyl, isoxazolyl, pyrazolyl, thiazolyl, oxazolyl, furyl, pyridinyl and pyrazinyl, where heterocyclyl is unsubstituted or carries 1, 2 or 3 radicals R^{21} , which are identical or different and selected from the group consisting of C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl and C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl. Especially preferred meanings for R^2 are 4,5-dihydroisoxazol-3-yl, 5-methyl-4,5-dihydroisoxazol-3-yl, 5-fluoromethyl-4,5-dihydroisoxazol-3-yl, 5-difluoromethyl-4,5-dihydroisoxazol-3-yl, 4,5-dihydroisoxazol-5-yl, 3-methyl-4,5-dihydroisoxazol-5-yl, 3-methoxy-4,5-dihydroisoxazol-5-yl, 3-methoxymethyl-4,5-dihydroisoxazol-5-yl, 3-methylsulfanylmethyl, 4,5-dihydroisoxazol-5-yl, 1-methyl-5-oxo-1,5-dihydrotetrazol-2-yl; 4-methyl-5-oxo-4,5-dihydrotetrazol-1-yl, morpholin-4-yl, isoxazol-3-yl, 5-methyl-isoxazol-3-yl, isoxazol-5-yl, 3-methyl-isoxazol-5-yl, 1-methyl-1H-pyrazol-3-yl, 2-methyl-2H-pyrazol-3-yl and thiazol-2-yl.

In the compounds of the invention, wherein X^2 in formula I is CR^2 , the variable R^2 may also be Z^{2a} -phenyl, where Z^{2a} is as defined herein, and where phenyl is unsubstituted or carries 1, 2 or 3 radicals R^{21} which are identical or different. According to a preferred embodiment, Z^{2a} is a covalent bond. According to a further preferred embodiment, Z^{2a} is C_1 - C_4 -alkanediyl-O, such as OCH_2 or OCH_2CH_2 . According to a further preferred embodiment, Z^{2a} is O- C_1 - C_4 -alkanediyl such as CH_2O or CH_2CH_2O . According to a further preferred embodiment, Z^{2a} is C_1 - C_4 -alkanediyl-O- C_1 - C_4 -alkanediyl. In this embodiment, R^{21} is preferably selected from halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, and preferably from halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_2 -haloalkyl and C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy such as fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, OCH_2OCH_3 , $OCH_2CH_2OCH_2CH_3$, $OCH_2OCH_2CH_3$ or $OCH_2CH_2OCH_3$. In particular, phenyl is unsubstituted or carries 1 radical R^{21} .

In particular the variable R^2 in the compounds of formula I, where X is C- R^2 , may be a radical of the following formula:



in which # denotes the bond through which the group R^2 is attached and:

R^{P1} is hydrogen or halogen, preferably H, Cl, Br or F, and in particular H or F;

5 R^{P2} is hydrogen, halogen or C₁-C₂-alkoxy, preferably H, Cl, Br, F, OCH₃ or OCH₂CH₃, and in particular H, F, Cl or OCH₃; and

R^{P3} is hydrogen, halogen, C₁-C₂-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, C₁-C₂-alkoxy-C₁-C₂-alkoxy, preferably H, Cl, Br, F, CH₃, C₂H₅, CF₃, CHF₂, CH₂F, CCl₂F, CF₂Cl, CH₂CF₃, CH₂CHF₂, CF₂CF₃, OCH₃, OCH₂CH₃, OCH₂OCH₃, OCH₂CH₂OCH₂CH₃, OCH₂OCH₂CH₃ or
10 OCH₂CH₂OCH₃, and in particular is H, F, Cl, CH₃, CF₃, OCH₃, OCH₂CH₃, OCH₂OCH₃ or OCH₂CH₂OCH₃.

According to a particular embodiment of the invention the variable R^2 in the compounds of formula I is phenyl which is unsubstituted or carries one radical R^{21} , where R^{21} is attached to position 4 of the phenyl group and is selected from halogen C₁-C₂-alkyl, C₁-C₄-alkoxy, C₁-C₂-haloalkyl and C₁-C₂-alkoxy-C₁-C₂-alkoxy, preferably from fluorine, chlorine, bromine, CH₃, C₂H₅, OCH₃, OC₂H₅, CHF₂, CF₃, OCH₂OCH₃ and OCH₂CH₂OCH₃, and specifically from OCH₃ and OC₂H₅.


In the compounds of the invention, wherein X^2 in formula I is CR^2 , a particular embodiment relates to compounds, wherein the variable R^2 is selected from the group consisting of
20 halogen, C₁-C₆-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy, C₂-C₄-haloalkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₃-C₆-haloalkenyloxy, C₃-C₆-haloalkynyloxy, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl and C₁-C₄-haloalkyl-S(O)₂. More preferably, R^2 is selected from C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-alkoxy, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₃-C₄-alkenyloxy, C₃-C₄-alkynyloxy, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl and C₁-C₄-alkylsulfanyl, and in particular from CH=CH₂, CH=CHCH₃, CH₂OCH₂CF₃, OC₂H₅, OCH₂CH=CH₂, OCH₂C≡CH, C(O)OCH₃, C(O)OC₂H₅, SO₂CH₃, SO₂C₂H₅, SO₂CH(CH₃)₂, SOCH₃, SOC₂H₅, SOCH(CH₃)₂, SCH₃, SC₂H₅ and SCH(CH₃)₂.

According to another preferred group of embodiments of the invention X is C- R^2 , wherein
30 R^2 together with R^3 or together with R^1 , if present, forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle, where the fused heterocycle has 1, 2, 3 or 4 heteroatoms selected from O, S and N as ring members, where the fused carbocycle and the fused heterocycle are monocyclic or bicyclic and where the fused carbocycle and the fused heterocycle are unsubstituted or carry 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 radicals R^q .
35

According to a particular embodiment of the invention, X is C- R^2 , wherein R^2 together with R^3 or together with R^1 , if present, forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle, where the fused heterocycle has 1, 2, 3 or

4 heteroatoms selected from O, S and N as ring members, where the fused carbocycle and the fused heterocycle are monocyclic or bicyclic and where the fused carbocycle and the fused heterocycle are unsubstituted or carry 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 radicals R^q . The ring member S can optionally be oxidized to various oxidation states. The ring member N can optionally be oxidized. Together with the six-membered N-heteroaromatic group to which they are attached a nine- to fifteen-membered bi- or tricyclic ring system results. In this embodiment, the heteroaromatic group is preferably fused to a benzene, naphthalene, C_5 - C_{10} -cycloalkane, or heterocyclic ring having 5 to 10 ring members. Examples for a ring system where R^2 together with R^3 or together with R^1 forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle as defined above are quinoline, isoquinoline, quinoxaline, benzo[g]isoquinoline, 5,6,7,8-tetrahydroisoquinoline, 5,8-dihydroisoquinoline, 1,5-naphthyridine, 1,6-naphthyridine, 2,6-naphthyridine, 1,7-naphthyridine, 2,7-naphthyridine, 1,8-naphthyridine, pyrido[2,3-b]pyrazine, pyrido[3,4-b]pyrazine, 6,7-dihydro-5H-[2]pyrindine, 5H-[2]pyrindine, 2,3-dihydro-1H-pyrrolo[3,2-c]pyridine, 1H-pyrrolo[3,2-c]pyridine, 2,3-dihydrofuro[3,2-c]pyridine, furo[3,2-c]pyridine, 2,3-dihydrothieno[3,2-c]pyridine, thieno[3,2-c]pyridine, [1,3]dioxolo[4,5-c]pyridine, 1H-imidazo[4,5-c]pyridine, 6,7-dihydro-5H-[1]pyrindine, 5H-[1H]pyrindine, 7H-[1H]pyrindine, 5H-cyclopentapyrazine, 6,7-dihydro-5H-cyclopentapyrazine, quinoxaline and the like.


Particular embodiments of the invention relate to compounds of the formula I, wherein X is C- R^2 and wherein R^2 together with R^3 forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6- or 7-membered heterocycle as defined above. Examples are the groups A.1, A.2, A.3, A.4 and A.5:



denotes the bond to the carbonyl carbon atom of 1,2,5-oxadiazol-3-yl-aminocarbonyl group;

X^1 , X^4 , R^5 and R^q are as defined above.

Further particular embodiments of the invention relate to compounds of the formula I, wherein X is C- R^2 and wherein R^2 together with R^1 forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6- or 7-membered heterocycle as defined above. Examples are the groups A.6, A.7, A.8, A.9 and A.10



denotes the bond to the carbon atom of the carbonyl group of the 1,2,5-oxadiazol-3-yl-aminocarbonyl group;

X⁴ is N and R³, R⁵ and R^q are as defined above; and

According to a further preferred embodiment of the invention X² is N.

5 According to a further preferred embodiment of the invention X⁴ is CR⁴. According to this embodiment of the invention, R⁴ is preferably selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₅-cycloalkyl, C₃-C₅-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₅-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃-alkylamino, C₁-C₃-dialkylamino, C₁-C₃-alkylamino-S(O)_k,
 10 C₁-C₃-alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkylthio-C₁-C₄-alkyl, C₃-C₄-alkenyloxy, C₃-C₄-alkynyloxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkoxy, C₁-C₄-alkyl-S(O)_k, C₁-C₄-haloalkyl-S(O)_k, where k is 0, 1 or 2; provided that if R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not
 15 hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

In particular, R⁴ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃-alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; specifically R⁴ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃, CH(CH₃)₂, cyclopropyl, cyclopropylmethyl, CF₃, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃; and more specifically R⁴ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃, CH(CH₃)₂, cyclopropyl, CF₃, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃; provided that if R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl.

In this context, the variables R', R¹¹, R²¹, Z, Z¹, Z², Z^{2a}, R^b, R^{1b}, R^{2b}, R^c, R^{2c}, R^d, R^{2d}, R^e, R^f, R^{2e}, R^{2f}, R^g, R^h, R^{2g}, R^{2h}, R^k, n and k, independently of each other, preferably have one of the
 30 following meanings:

R', R¹¹, R²¹ independently of each other are selected from halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy and C₁-C₄-haloalkyloxy, more preferably from halogen, C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy, and in particular from
 35 Cl, F, Br, methyl, ethyl, methoxy and trifluoromethyl.

Z, Z¹, Z² independently of each other are selected from covalent bond methanediyl and ethanediyl.

Z^{2a} is selected from a covalent bond, C₁-C₂-alkanediyl, O-C₁-C₂-alkanediyl, C₁-C₂-alkanediyl-O and C₁-C₂-alkanediyl-O-C₁-C₂-alkanediyl; more preferably from a covalent bond, methanediyl, ethanediyl, O-methanediyl, O-ethanediyl, methanediyl-O, and ethanediyl-O; and in particular from a covalent bond, methanediyl and ethanediyl.

R^b, R^{1b}, R^{2b} independently of each other are selected from C₁-C₆-alkyl, C₃-C₇-cycloalkyl,

C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2 or 3 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2 or 3 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₂-haloalkyl and C₁-C₂-alkoxy.

More preferably R^b, R^{1b}, R^{2b} independently of each other are selected from the group consisting of C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₄-haloalkynyl, C₃-C₆-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2 or 3 heteroatoms as ring members, which are selected from the group consisting of O, N and S.

In particular, R^b, R^{1b}, R^{2b} independently of each other are selected from C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₃-C₆-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered aromatic heterocyclic radical having 1 or 2 nitrogen atoms as ring members.

R^c, R^{2c} and R^k independently of each other are selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl, benzyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2 or 3 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl, benzyl and heterocyclyl are unsubstituted or substituted by 1, 2 or 3 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy.

More preferably R^c, R^{2c}, R^k independently of each other are selected from hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₃-C₆-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2 or 3 heteroatoms as ring members, which are selected from the group consisting of O, N and S.

In particular, R^c, R^{2c}, R^k independently of each other are selected from C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₃-C₆-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 5- or 6-membered aromatic heterocyclic radical having 1 or 2 nitrogen atoms as ring members.

R^d, R^{2d} independently of each other are selected from C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl.

More preferably R^d, R^{2d} independently of each other are selected from C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, and in particular selected from C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl and C₃-C₆-cycloalkyl.

R^e , R^f , R^{2e} , R^{2f} independently of each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, which is unsubstituted or partially or completely halogenated, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2 or 3 groups, which are
5 identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy, or R^e and R^f or R^{2e} and R^{2f} together with the nitrogen atom, to which they are bound may form a 5-, 6 or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and
10 selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl and C₁-C₄-alkoxy.

More preferably R^e , R^f , R^{2e} , R^{2f} independently of each other are selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl and benzyl, or R^e and R^f or R^{2e} and R^{2f} together with the nitrogen atom, to which they are bound form a 5- or 6-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S
15 and N and which is unsubstituted or may carry 1, 2 or 3 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl.

In particular, R^e , R^f , R^{2e} , R^{2f} independently of each other are selected from hydrogen and C₁-C₄-alkyl, or R^e and R^f or R^{2e} and R^{2f} together with the nitrogen atom, to which they are bound may form a 5- or 6-membered, saturated N-bound heterocyclic radical, which may carry as a
20 ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2 or 3 methyl groups.

R^g , R^{2g} independently of each other are selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and
25 benzyl; more preferably R^g , R^{2g} independently of each other are selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, and in particular selected from C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl and C₃-C₆-cycloalkyl.

R^h , R^{2h} independently of each other are selected from hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl, benzyl and a radical C(=O)- R^k , where R^k is H, C₁-C₄-alkyl, C₁-C₄-haloalkyl or phenyl; more preferably R^h , R^{2h} independently of each other are selected from C₁-C₆-alkyl, C₁-C₆-haloalkyl,
35 C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated, and in particular selected from C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl and C₃-C₆-cycloalkyl; or

R^g and R^h or R^{2g} and R^{2h} together with the nitrogen atom, to which they are bound may
40 form a 5-, 6 or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from

the group consisting of =O, halogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl and C₁-C₄-alkoxy; more preferably R^g and R^h or R^{2g} and R^{2h} together with the nitrogen atom, to which they are bound form a 5- or 6-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is

unsubstituted or may carry 1, 2 or 3 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl; and in particular, R^g and R^h or R^{2g} and R^{2h} together with the nitrogen atom, to which they are bound may form a 5- or 6-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2 or 3 methyl groups.

n is preferably are 0.


k is preferably 0 or 2.

R^q is preferably halogen, OH, NO₂, cyano, oxo (=O), C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio and C₁-C₄-haloalkoxy.

Specifically preferred are compounds of formula I, wherein one of X¹, X² and X⁴ is N.

Specially preferred are likewise compounds of the formula I, wherein two of X¹, X² and X⁴ are N.

A particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is N, X² is CR² and X⁴ is CR⁴. These compounds are also referred to as compound of formula I.1, wherein R², R³, R⁴, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R², R³, R⁴, R⁵ and R in connection with compounds of formula I also apply for formula I.1 as defined herein. In formula I.1, the positions on the pyridine ring are designated by arabic numbers.

Amongst compounds of formula I.1, those are preferred, wherein R², R³, R⁴, R⁵ and R have the preferred meanings mentioned above. Especially more preferred are compounds of formula I.1, wherein R³, R⁴, R⁵ and R have the preferred meanings mentioned above and the variable R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl. In particular, R² is selected from hydrogen, methoxymethyl, ethoxymethyl, 2,2,2-trifluoroethoxymethyl, 2,2,2-trifluoroethoxyethyl, methylsulfonyl, methylsulfinyl, methylsulfanyl, 4,5-dihydroisoxazol-5-yl, 4,5-dihydroisoxazol-3-yl, 3-methyl-4,5-dihydroisoxazol-5-yl, 5-methyl-4,5-dihydroisoxazol-3-yl, isoxazol-5-yl, 3-methyl-isoxazol-5-yl, isoxazol-3-yl and 5-methyl-isoxazol-3-yl.

More preferred are compounds of formula I.1, wherein the variables R, R², R³, R⁴ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

5 R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazolinyl, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl, in particular hydrogen, methoxymethyl, ethoxymethyl, 2,2,2-trifluoroethoxymethyl, 2,2,2-trifluoroethoxyethyl, methylsulfonyl, 10 methylsulfinyl, methylsulfanyl, 4,5-dihydroisoxazol-5-yl, 4,5-dihydroisoxazol-3-yl, 3-methyl-4,5-dihydroisoxazol-5-yl, 5-methyl-4,5-dihydroisoxazol-3-yl, isoxazol-5-yl, 3-methyl-isoxazol-5-yl, isoxazol-3-yl and 5-methyl-isoxazol-3-yl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular H, Cl, F, CF₃, SO₂CH₃ or CN;

15 R⁴ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃ alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; and

20 R⁵ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃ alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; provided that if R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl.

30 Even more preferred are compounds of formula I.1, wherein the variables R, R², R³, R⁴ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

35 R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazolinyl, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl;

40 R⁴ is H; and R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃; or

R⁵ is H; and R⁴ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

With respect to their use, particular preference is given to the compounds of formula I.1 compiled in the tables 1-13 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 1 Compounds of the formula I.1 (compounds I.1-1 to I.1-336) in which R² is hydrogen and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 2 Compounds of the formula I.1 (compounds I.1-337 to I.1-672) in which R² is SCH₃ and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 3 Compounds of the formula I.1 (compounds I.1-673 to I.1-1008) in which R² is SOCH₃ and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 4 Compounds of the formula I.1 (compounds I.1-1009 to I.1-1344) in which R² is SO₂CH₃ and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 5 Compounds of the formula I.1 (compounds I.1-1345 to I.1-1680) in which R² is 2,2,2-trifluoroethoxymethyl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 6 Compounds of the formula I.1 (compounds I.1-1681 to I.1-2016) in which R² is 4,5-dihydroisoxazol-3-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 7 Compounds of the formula I.1 (compounds I.1-2017 to I.1-2352) in which R² is 5-methyl-4,5-dihydroisoxazol-3-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 8 Compounds of the formula I.1 (compounds I.1-2353 to I.1-2688) in which R² is 4,5-dihydroisoxazol-5-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 9 Compounds of the formula I.1 (compounds I.1-2689 to I.1-3024) in which R² is 3-methyl-4,5-dihydroisoxazol-5-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 10 Compounds of the formula I.1 (compounds I.1-3025 to I.1-3360) in which R² is isoxazol-3-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 11 Compounds of the formula I.1 (compounds I.1-3361 to I.1-3696) in which R² is 5-methyl-isoxazol-3-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 12 Compounds of the formula I.1 (compounds I.1-3697 to I.1-4032) in which R² is isoxazol-5-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 13 Compounds of the formula I.1 (compounds I.1-4033 to I.1-4368) in which R² is 3-methyl-isoxazol-5-yl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A.

Table A

	R	R ³	R ⁴	R ⁵
A-1	CH ₃	Cl	H	cyclopropyl
A-2	CH ₃	Cl	H	OCH ₃
A-3	CH ₃	Cl	H	OCF ₃
A-4	CH ₃	Cl	H	SCH ₃
A-5	CH ₃	Cl	H	SOCH ₃
A-6	CH ₃	Cl	H	SO ₂ CH ₃
A-7	CH ₃	Cl	H	CH ₂ OCH ₃
A-8	CH ₃	Cl	cyclopropyl	H
A-9	CH ₃	Cl	OCH ₃	H
A-10	CH ₃	Cl	OCF ₃	H
A-11	CH ₃	Cl	SCH ₃	H
A-12	CH ₃	Cl	SOCH ₃	H
A-13	CH ₃	Cl	SO ₂ CH ₃	H
A-14	CH ₃	Cl	CH ₂ OCH ₃	H
A-15	CH ₃	F	H	cyclopropyl
A-16	CH ₃	F	H	OCH ₃
A-17	CH ₃	F	H	OCF ₃
A-18	CH ₃	F	H	SCH ₃
A-19	CH ₃	F	H	SOCH ₃
A-20	CH ₃	F	H	SO ₂ CH ₃
A-21	CH ₃	F	H	CH ₂ OCH ₃
A-22	CH ₃	F	cyclopropyl	H
A-23	CH ₃	F	OCH ₃	H
A-24	CH ₃	F	OCF ₃	H
A-25	CH ₃	F	SCH ₃	H
A-26	CH ₃	F	SOCH ₃	H
A-27	CH ₃	F	SO ₂ CH ₃	H
A-28	CH ₃	F	CH ₂ OCH ₃	H
A-29	CH ₃	CF ₃	H	cyclopropyl
A-30	CH ₃	CF ₃	H	OCH ₃
A-31	CH ₃	CF ₃	H	OCF ₃
A-32	CH ₃	CF ₃	H	SCH ₃

A-33	CH ₃	CF ₃	H	SOCH ₃
A-34	CH ₃	CF ₃	H	SO ₂ CH ₃
A-35	CH ₃	CF ₃	H	CH ₂ OCH ₃
A-36	CH ₃	CF ₃	cyclopropyl	H
A-37	CH ₃	CF ₃	OCH ₃	H
A-38	CH ₃	CF ₃	OCF ₃	H
A-39	CH ₃	CF ₃	SCH ₃	H
A-40	CH ₃	CF ₃	SOCH ₃	H
A-41	CH ₃	CF ₃	SO ₂ CH ₃	H
A-42	CH ₃	CF ₃	CH ₂ OCH ₃	H
A-43	CH ₃	SO ₂ CH ₃	H	cyclopropyl
A-44	CH ₃	SO ₂ CH ₃	H	OCH ₃
A-45	CH ₃	SO ₂ CH ₃	H	OCF ₃
A-46	CH ₃	SO ₂ CH ₃	H	SCH ₃
A-47	CH ₃	SO ₂ CH ₃	H	SOCH ₃
A-48	CH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
A-49	CH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
A-50	CH ₃	SO ₂ CH ₃	cyclopropyl	H
A-51	CH ₃	SO ₂ CH ₃	OCH ₃	H
A-52	CH ₃	SO ₂ CH ₃	OCF ₃	H
A-53	CH ₃	SO ₂ CH ₃	SCH ₃	H
A-54	CH ₃	SO ₂ CH ₃	SOCH ₃	H
A-55	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	H
A-56	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃	H
A-57	CH ₃	CN	H	cyclopropyl
A-58	CH ₃	CN	H	OCH ₃
A-59	CH ₃	CN	H	OCF ₃
A-60	CH ₃	CN	H	SCH ₃
A-61	CH ₃	CN	H	SOCH ₃
A-62	CH ₃	CN	H	SO ₂ CH ₃
A-63	CH ₃	CN	H	CH ₂ OCH ₃
A-64	CH ₃	CN	cyclopropyl	H
A-65	CH ₃	CN	OCH ₃	H
A-66	CH ₃	CN	OCF ₃	H
A-67	CH ₃	CN	SCH ₃	H
A-68	CH ₃	CN	SOCH ₃	H
A-69	CH ₃	CN	SO ₂ CH ₃	H
A-70	CH ₃	CN	CH ₂ OCH ₃	H
A-71	CH ₃	H	H	cyclopropyl
A-72	CH ₃	H	H	OCH ₃
A-73	CH ₃	H	H	OCF ₃

A-74	CH ₃	H	H	SCH ₃
A-75	CH ₃	H	H	SOCH ₃
A-76	CH ₃	H	H	SO ₂ CH ₃
A-77	CH ₃	H	H	CH ₂ OCH ₃
A-78	CH ₃	H	cyclopropyl	H
A-79	CH ₃	H	OCH ₃	H
A-80	CH ₃	H	OCF ₃	H
A-81	CH ₃	H	SCH ₃	H
A-82	CH ₃	H	SOCH ₃	H
A-83	CH ₃	H	SO ₂ CH ₃	H
A-84	CH ₃	H	CH ₂ OCH ₃	H
A-85	CH ₂ CH ₃	Cl	H	cyclopropyl
A-86	CH ₂ CH ₃	Cl	H	OCH ₃
A-87	CH ₂ CH ₃	Cl	H	OCF ₃
A-88	CH ₂ CH ₃	Cl	H	SCH ₃
A-89	CH ₂ CH ₃	Cl	H	SOCH ₃
A-90	CH ₂ CH ₃	Cl	H	SO ₂ CH ₃
A-91	CH ₂ CH ₃	Cl	H	CH ₂ OCH ₃
A-92	CH ₂ CH ₃	Cl	cyclopropyl	H
A-93	CH ₂ CH ₃	Cl	OCH ₃	H
A-94	CH ₂ CH ₃	Cl	OCF ₃	H
A-95	CH ₂ CH ₃	Cl	SCH ₃	H
A-96	CH ₂ CH ₃	Cl	SOCH ₃	H
A-97	CH ₂ CH ₃	Cl	SO ₂ CH ₃	H
A-98	CH ₂ CH ₃	Cl	CH ₂ OCH ₃	H
A-99	CH ₂ CH ₃	F	H	cyclopropyl
A-100	CH ₂ CH ₃	F	H	OCH ₃
A-101	CH ₂ CH ₃	F	H	OCF ₃
A-102	CH ₂ CH ₃	F	H	SCH ₃
A-103	CH ₂ CH ₃	F	H	SOCH ₃
A-104	CH ₂ CH ₃	F	H	SO ₂ CH ₃
A-105	CH ₂ CH ₃	F	H	CH ₂ OCH ₃
A-106	CH ₂ CH ₃	F	cyclopropyl	H
A-107	CH ₂ CH ₃	F	OCH ₃	H
A-108	CH ₂ CH ₃	F	OCF ₃	H
A-109	CH ₂ CH ₃	F	SCH ₃	H
A-110	CH ₂ CH ₃	F	SOCH ₃	H
A-111	CH ₂ CH ₃	F	SO ₂ CH ₃	H
A-112	CH ₂ CH ₃	F	CH ₂ OCH ₃	H
A-113	CH ₂ CH ₃	CF ₃	H	cyclopropyl
A-114	CH ₂ CH ₃	CF ₃	H	OCH ₃

A-115	CH ₂ CH ₃	CF ₃	H	OCF ₃
A-116	CH ₂ CH ₃	CF ₃	H	SCH ₃
A-117	CH ₂ CH ₃	CF ₃	H	SOCH ₃
A-118	CH ₂ CH ₃	CF ₃	H	SO ₂ CH ₃
A-119	CH ₂ CH ₃	CF ₃	H	CH ₂ OCH ₃
A-120	CH ₂ CH ₃	CF ₃	cyclopropyl	H
A-121	CH ₂ CH ₃	CF ₃	OCH ₃	H
A-122	CH ₂ CH ₃	CF ₃	OCF ₃	H
A-123	CH ₂ CH ₃	CF ₃	SCH ₃	H
A-124	CH ₂ CH ₃	CF ₃	SOCH ₃	H
A-125	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	H
A-126	CH ₂ CH ₃	CF ₃	CH ₂ OCH ₃	H
A-127	CH ₂ CH ₃	SO ₂ CH ₃	H	cyclopropyl
A-128	CH ₂ CH ₃	SO ₂ CH ₃	H	OCH ₃
A-129	CH ₂ CH ₃	SO ₂ CH ₃	H	OCF ₃
A-130	CH ₂ CH ₃	SO ₂ CH ₃	H	SCH ₃
A-131	CH ₂ CH ₃	SO ₂ CH ₃	H	SOCH ₃
A-132	CH ₂ CH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
A-133	CH ₂ CH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
A-134	CH ₂ CH ₃	SO ₂ CH ₃	cyclopropyl	H
A-135	CH ₂ CH ₃	SO ₂ CH ₃	OCH ₃	H
A-136	CH ₂ CH ₃	SO ₂ CH ₃	OCF ₃	H
A-137	CH ₂ CH ₃	SO ₂ CH ₃	SCH ₃	H
A-138	CH ₂ CH ₃	SO ₂ CH ₃	SOCH ₃	H
A-139	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	H
A-140	CH ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃	H
A-141	CH ₂ CH ₃	CN	H	cyclopropyl
A-142	CH ₂ CH ₃	CN	H	OCH ₃
A-143	CH ₂ CH ₃	CN	H	OCF ₃
A-144	CH ₂ CH ₃	CN	H	SCH ₃
A-145	CH ₂ CH ₃	CN	H	SOCH ₃
A-146	CH ₂ CH ₃	CN	H	SO ₂ CH ₃
A-147	CH ₂ CH ₃	CN	H	CH ₂ OCH ₃
A-148	CH ₂ CH ₃	CN	cyclopropyl	H
A-149	CH ₂ CH ₃	CN	OCH ₃	H
A-150	CH ₂ CH ₃	CN	OCF ₃	H
A-151	CH ₂ CH ₃	CN	SCH ₃	H
A-152	CH ₂ CH ₃	CN	SOCH ₃	H
A-153	CH ₂ CH ₃	CN	SO ₂ CH ₃	H
A-154	CH ₂ CH ₃	CN	CH ₂ OCH ₃	H
A-155	CH ₂ CH ₃	H	H	cyclopropyl

A-156	CH ₂ CH ₃	H	H	OCH ₃
A-157	CH ₂ CH ₃	H	H	OCF ₃
A-158	CH ₂ CH ₃	H	H	SCH ₃
A-159	CH ₂ CH ₃	H	H	SOCH ₃
A-160	CH ₂ CH ₃	H	H	SO ₂ CH ₃
A-161	CH ₂ CH ₃	H	H	CH ₂ OCH ₃
A-162	CH ₂ CH ₃	H	cyclopropyl	H
A-163	CH ₂ CH ₃	H	OCH ₃	H
A-164	CH ₂ CH ₃	H	OCF ₃	H
A-165	CH ₂ CH ₃	H	SCH ₃	H
A-166	CH ₂ CH ₃	H	SOCH ₃	H
A-167	CH ₂ CH ₃	H	SO ₂ CH ₃	H
A-168	CH ₂ CH ₃	H	CH ₂ OCH ₃	H
A-169	CH ₂ OCH ₃	Cl	H	cyclopropyl
A-170	CH ₂ OCH ₃	Cl	H	OCH ₃
A-171	CH ₂ OCH ₃	Cl	H	OCF ₃
A-172	CH ₂ OCH ₃	Cl	H	SCH ₃
A-173	CH ₂ OCH ₃	Cl	H	SOCH ₃
A-174	CH ₂ OCH ₃	Cl	H	SO ₂ CH ₃
A-175	CH ₂ OCH ₃	Cl	H	CH ₂ OCH ₃
A-176	CH ₂ OCH ₃	Cl	cyclopropyl	H
A-177	CH ₂ OCH ₃	Cl	OCH ₃	H
A-178	CH ₂ OCH ₃	Cl	OCF ₃	H
A-179	CH ₂ OCH ₃	Cl	SCH ₃	H
A-180	CH ₂ OCH ₃	Cl	SOCH ₃	H
A-181	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	H
A-182	CH ₂ OCH ₃	Cl	CH ₂ OCH ₃	H
A-183	CH ₂ OCH ₃	F	H	cyclopropyl
A-184	CH ₂ OCH ₃	F	H	OCH ₃
A-185	CH ₂ OCH ₃	F	H	OCF ₃
A-186	CH ₂ OCH ₃	F	H	SCH ₃
A-187	CH ₂ OCH ₃	F	H	SOCH ₃
A-188	CH ₂ OCH ₃	F	H	SO ₂ CH ₃
A-189	CH ₂ OCH ₃	F	H	CH ₂ OCH ₃
A-190	CH ₂ OCH ₃	F	cyclopropyl	H
A-191	CH ₂ OCH ₃	F	OCH ₃	H
A-192	CH ₂ OCH ₃	F	OCF ₃	H
A-193	CH ₂ OCH ₃	F	SCH ₃	H
A-194	CH ₂ OCH ₃	F	SOCH ₃	H
A-195	CH ₂ OCH ₃	F	SO ₂ CH ₃	H
A-196	CH ₂ OCH ₃	F	CH ₂ OCH ₃	H


A-197	CH ₂ OCH ₃	CF ₃	H	cyclopropyl
A-198	CH ₂ OCH ₃	CF ₃	H	OCH ₃
A-199	CH ₂ OCH ₃	CF ₃	H	OCF ₃
A-200	CH ₂ OCH ₃	CF ₃	H	SCH ₃
A-201	CH ₂ OCH ₃	CF ₃	H	SOCH ₃
A-202	CH ₂ OCH ₃	CF ₃	H	SO ₂ CH ₃
A-203	CH ₂ OCH ₃	CF ₃	H	CH ₂ OCH ₃
A-204	CH ₂ OCH ₃	CF ₃	cyclopropyl	H
A-205	CH ₂ OCH ₃	CF ₃	OCH ₃	H
A-206	CH ₂ OCH ₃	CF ₃	OCF ₃	H
A-207	CH ₂ OCH ₃	CF ₃	SCH ₃	H
A-208	CH ₂ OCH ₃	CF ₃	SOCH ₃	H
A-209	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	H
A-210	CH ₂ OCH ₃	CF ₃	CH ₂ OCH ₃	H
A-211	CH ₂ OCH ₃	SO ₂ CH ₃	H	cyclopropyl
A-212	CH ₂ OCH ₃	SO ₂ CH ₃	H	OCH ₃
A-213	CH ₂ OCH ₃	SO ₂ CH ₃	H	OCF ₃
A-214	CH ₂ OCH ₃	SO ₂ CH ₃	H	SCH ₃
A-215	CH ₂ OCH ₃	SO ₂ CH ₃	H	SOCH ₃
A-216	CH ₂ OCH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
A-217	CH ₂ OCH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
A-218	CH ₂ OCH ₃	SO ₂ CH ₃	cyclopropyl	H
A-219	CH ₂ OCH ₃	SO ₂ CH ₃	OCH ₃	H
A-220	CH ₂ OCH ₃	SO ₂ CH ₃	OCF ₃	H
A-221	CH ₂ OCH ₃	SO ₂ CH ₃	SCH ₃	H
A-222	CH ₂ OCH ₃	SO ₂ CH ₃	SOCH ₃	H
A-223	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	H
A-224	CH ₂ OCH ₃	SO ₂ CH ₃	CH ₂ OCH ₃	H
A-225	CH ₂ OCH ₃	CN	H	cyclopropyl
A-226	CH ₂ OCH ₃	CN	H	OCH ₃
A-227	CH ₂ OCH ₃	CN	H	OCF ₃
A-228	CH ₂ OCH ₃	CN	H	SCH ₃
A-229	CH ₂ OCH ₃	CN	H	SOCH ₃
A-230	CH ₂ OCH ₃	CN	H	SO ₂ CH ₃
A-231	CH ₂ OCH ₃	CN	H	CH ₂ OCH ₃
A-232	CH ₂ OCH ₃	CN	cyclopropyl	H
A-233	CH ₂ OCH ₃	CN	OCH ₃	H
A-234	CH ₂ OCH ₃	CN	OCF ₃	H
A-235	CH ₂ OCH ₃	CN	SCH ₃	H
A-236	CH ₂ OCH ₃	CN	SOCH ₃	H
A-237	CH ₂ OCH ₃	CN	SO ₂ CH ₃	H

A-238	CH ₂ OCH ₃	CN	CH ₂ OCH ₃	H
A-239	CH ₂ OCH ₃	H	H	cyclopropyl
A-240	CH ₂ OCH ₃	H	H	OCH ₃
A-241	CH ₂ OCH ₃	H	H	OCF ₃
A-242	CH ₂ OCH ₃	H	H	SCH ₃
A-243	CH ₂ OCH ₃	H	H	SOCH ₃
A-244	CH ₂ OCH ₃	H	H	SO ₂ CH ₃
A-245	CH ₂ OCH ₃	H	H	CH ₂ OCH ₃
A-246	CH ₂ OCH ₃	H	cyclopropyl	H
A-247	CH ₂ OCH ₃	H	OCH ₃	H
A-248	CH ₂ OCH ₃	H	OCF ₃	H
A-249	CH ₂ OCH ₃	H	SCH ₃	H
A-250	CH ₂ OCH ₃	H	SOCH ₃	H
A-251	CH ₂ OCH ₃	H	SO ₂ CH ₃	H
A-252	CH ₂ OCH ₃	H	CH ₂ OCH ₃	H
A-253	CH ₂ CH ₂ OCH ₃	Cl	H	cyclopropyl
A-254	CH ₂ CH ₂ OCH ₃	Cl	H	OCH ₃
A-255	CH ₂ CH ₂ OCH ₃	Cl	H	OCF ₃
A-256	CH ₂ CH ₂ OCH ₃	Cl	H	SCH ₃
A-257	CH ₂ CH ₂ OCH ₃	Cl	H	SOCH ₃
A-258	CH ₂ CH ₂ OCH ₃	Cl	H	SO ₂ CH ₃
A-259	CH ₂ CH ₂ OCH ₃	Cl	H	CH ₂ OCH ₃
A-260	CH ₂ CH ₂ OCH ₃	Cl	cyclopropyl	H
A-261	CH ₂ CH ₂ OCH ₃	Cl	OCH ₃	H
A-262	CH ₂ CH ₂ OCH ₃	Cl	OCF ₃	H
A-263	CH ₂ CH ₂ OCH ₃	Cl	SCH ₃	H
A-264	CH ₂ CH ₂ OCH ₃	Cl	SOCH ₃	H
A-265	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	H
A-266	CH ₂ CH ₂ OCH ₃	Cl	CH ₂ OCH ₃	H
A-267	CH ₂ CH ₂ OCH ₃	F	H	cyclopropyl
A-268	CH ₂ CH ₂ OCH ₃	F	H	OCH ₃
A-269	CH ₂ CH ₂ OCH ₃	F	H	OCF ₃
A-270	CH ₂ CH ₂ OCH ₃	F	H	SCH ₃
A-271	CH ₂ CH ₂ OCH ₃	F	H	SOCH ₃
A-272	CH ₂ CH ₂ OCH ₃	F	H	SO ₂ CH ₃
A-273	CH ₂ CH ₂ OCH ₃	F	H	CH ₂ OCH ₃
A-274	CH ₂ CH ₂ OCH ₃	F	cyclopropyl	H
A-275	CH ₂ CH ₂ OCH ₃	F	OCH ₃	H
A-276	CH ₂ CH ₂ OCH ₃	F	OCF ₃	H
A-277	CH ₂ CH ₂ OCH ₃	F	SCH ₃	H
A-278	CH ₂ CH ₂ OCH ₃	F	SOCH ₃	H

A-279	CH ₂ CH ₂ OCH ₃	F	SO ₂ CH ₃	H
A-280	CH ₂ CH ₂ OCH ₃	F	CH ₂ OCH ₃	H
A-281	CH ₂ CH ₂ OCH ₃	CF ₃	H	cyclopropyl
A-282	CH ₂ CH ₂ OCH ₃	CF ₃	H	OCH ₃
A-283	CH ₂ CH ₂ OCH ₃	CF ₃	H	OCF ₃
A-284	CH ₂ CH ₂ OCH ₃	CF ₃	H	SCH ₃
A-285	CH ₂ CH ₂ OCH ₃	CF ₃	H	SOCH ₃
A-286	CH ₂ CH ₂ OCH ₃	CF ₃	H	SO ₂ CH ₃
A-287	CH ₂ CH ₂ OCH ₃	CF ₃	H	CH ₂ OCH ₃
A-288	CH ₂ CH ₂ OCH ₃	CF ₃	cyclopropyl	H
A-289	CH ₂ CH ₂ OCH ₃	CF ₃	OCH ₃	H
A-290	CH ₂ CH ₂ OCH ₃	CF ₃	OCF ₃	H
A-291	CH ₂ CH ₂ OCH ₃	CF ₃	SCH ₃	H
A-292	CH ₂ CH ₂ OCH ₃	CF ₃	SOCH ₃	H
A-293	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	H
A-294	CH ₂ CH ₂ OCH ₃	CF ₃	CH ₂ OCH ₃	H
A-295	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	cyclopropyl
A-296	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	OCH ₃
A-297	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	OCF ₃
A-298	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SCH ₃
A-299	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SOCH ₃
A-300	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
A-301	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
A-302	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	cyclopropyl	H
A-303	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	OCH ₃	H
A-304	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	OCF ₃	H
A-305	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SCH ₃	H
A-306	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SOCH ₃	H
A-307	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	H
A-308	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CH ₂ OCH ₃	H
A-309	CH ₂ CH ₂ OCH ₃	CN	H	cyclopropyl
A-310	CH ₂ CH ₂ OCH ₃	CN	H	OCH ₃
A-311	CH ₂ CH ₂ OCH ₃	CN	H	OCF ₃
A-312	CH ₂ CH ₂ OCH ₃	CN	H	SCH ₃
A-313	CH ₂ CH ₂ OCH ₃	CN	H	SOCH ₃
A-314	CH ₂ CH ₂ OCH ₃	CN	H	SO ₂ CH ₃
A-315	CH ₂ CH ₂ OCH ₃	CN	H	CH ₂ OCH ₃
A-316	CH ₂ CH ₂ OCH ₃	CN	cyclopropyl	H
A-317	CH ₂ CH ₂ OCH ₃	CN	OCH ₃	H
A-318	CH ₂ CH ₂ OCH ₃	CN	OCF ₃	H
A-319	CH ₂ CH ₂ OCH ₃	CN	SCH ₃	H

A-320	CH ₂ CH ₂ OCH ₃	CN	SOCH ₃	H
A-321	CH ₂ CH ₂ OCH ₃	CN	SO ₂ CH ₃	H
A-322	CH ₂ CH ₂ OCH ₃	CN	CH ₂ OCH ₃	H
A-323	CH ₂ CH ₂ OCH ₃	H	H	cyclopropyl
A-324	CH ₂ CH ₂ OCH ₃	H	H	OCH ₃
A-325	CH ₂ CH ₂ OCH ₃	H	H	OCF ₃
A-326	CH ₂ CH ₂ OCH ₃	H	H	SCH ₃
A-327	CH ₂ CH ₂ OCH ₃	H	H	SOCH ₃
A-328	CH ₂ CH ₂ OCH ₃	H	H	SO ₂ CH ₃
A-329	CH ₂ CH ₂ OCH ₃	H	H	CH ₂ OCH ₃
A-330	CH ₂ CH ₂ OCH ₃	H	cyclopropyl	H
A-331	CH ₂ CH ₂ OCH ₃	H	OCH ₃	H
A-332	CH ₂ CH ₂ OCH ₃	H	OCF ₃	H
A-333	CH ₂ CH ₂ OCH ₃	H	SCH ₃	H
A-334	CH ₂ CH ₂ OCH ₃	H	SOCH ₃	H
A-335	CH ₂ CH ₂ OCH ₃	H	SO ₂ CH ₃	H
A-336	CH ₂ CH ₂ OCH ₃	H	CH ₂ OCH ₃	H

A further particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is C-R¹, X² is N and X⁴ is CR⁴. These compounds are also referred to as compound of formula I.2, wherein R¹, R³, R⁴, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R¹, R³, R⁴, R⁵ and R in connection with compounds of formula I also apply for formulae I.2 as defined herein. In formula I.2, the positions on the pyridine ring are designated by arabic numbers.

Amongst compounds of formula I.2, those are preferred, wherein R¹, R³, R⁴, R⁵ and R have the preferred meanings mentioned above. More preferred are compounds of formula I.2, wherein R³, R⁴, R⁵ and R have the preferred meanings and the variables R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, nitro, CH₃, CF₃, CH₂OCH₂CH₂OCH₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂ or SO₂CH₃.

More preferred are compounds of formula I.2, wherein the variables R, R¹, R³, R⁴ and R⁵ have the following meanings:

- R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;
- R¹ is halogen, C₁-C₄-alkyl, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂, SO₂CH₃ or CH₂OCH₂CH₂OCH₃;
- R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular H, Cl, F, CF₃, SO₂CH₃ or CN;
- R⁴ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃ alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; and
- R⁵ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-haloalkyl, C₁-C₃ alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; provided that if R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl.

Even more preferred are compounds of formula I.2, wherein the variables R, R¹, R³, R⁴ and R⁵ have the following meanings:

- R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;
- R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl
- R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl;
- R⁴ is H; and R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃; or
- R⁵ is H; and R⁴ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

With respect to their use, particular preference is given to the compounds of formula I.2 compiled in the tables 14-18 below. Moreover, the groups mentioned for a substituent in the

tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 14 Compounds of the formula I.2 (compounds I.2-1 to I.2-336) in which R¹ is chlorine and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;


Table 15 Compounds of the formula I.2 (compounds I.2-337 to I.2-672) in which R¹ is nitro and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 16 Compounds of the formula I.2 (compounds I.2-673 to I.2-1008), in which R¹ is methyl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 17 Compounds of the formula I.2 (compounds I.2-1009 to I.2-1344) in which R¹ is trifluoromethyl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A;

Table 18 Compounds of the formula I.2 (compounds I.2-1345 to I.2-1680) in which R¹ is methylsulfonyl and the combination of R, R³, R⁴ and R⁵ for a compound corresponds in each case to one row of Table A.

A further particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is N, X² is N and X⁴ is CR⁴. These compounds are also referred to as compound of formula I.3, wherein R³, R⁴, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R³, R⁴, R⁵ and R in connection with compounds of formula I also apply for formulae I.3 as defined herein. In formula I.3, the positions on the pyridazine ring are designated by arabic numbers.

Amongst compounds of formula I.3, those are preferred, wherein the variables R, R³, R⁴ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular H, F, Cl, Br, CN, NO₂, CH₃, CH₂CH₃, CF₃, CHF₂, OCH₃, OCF₃, OCHF₂, SO₂CH₃ or SO₂CH₂CH₃;

R⁴ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₄-

haloalkyl, C₁-C₃ alkylcarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of hydrogen, halogen, CN, nitro, C₁-C₄-alkyl, C₃-C₄-cycloalkyl, C₃-C₄-cycloalkyl-C₁-C₂-alkyl, where the C₃-C₄-cycloalkyl groups in the two
 5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995
 1000
 1005
 1010
 1015
 1020
 1025
 1030
 1035
 1040
 1045
 1050
 1055
 1060
 1065
 1070
 1075
 1080
 1085
 1090
 1095
 1100
 1105
 1110
 1115
 1120
 1125
 1130
 1135
 1140
 1145
 1150
 1155
 1160
 1165
 1170
 1175
 1180
 1185
 1190
 1195
 1200
 1205
 1210
 1215
 1220
 1225
 1230
 1235
 1240
 1245
 1250
 1255
 1260
 1265
 1270
 1275
 1280
 1285
 1290
 1295
 1300
 1305
 1310
 1315
 1320
 1325
 1330
 1335
 1340
 1345
 1350
 1355
 1360
 1365
 1370
 1375
 1380
 1385
 1390
 1395
 1400
 1405
 1410
 1415
 1420
 1425
 1430
 1435
 1440
 1445
 1450
 1455
 1460
 1465
 1470
 1475
 1480
 1485
 1490
 1495
 1500
 1505
 1510
 1515
 1520
 1525
 1530
 1535
 1540
 1545
 1550
 1555
 1560
 1565
 1570
 1575
 1580
 1585
 1590
 1595
 1600
 1605
 1610
 1615
 1620
 1625
 1630
 1635
 1640
 1645
 1650
 1655
 1660
 1665
 1670
 1675
 1680
 1685
 1690
 1695
 1700
 1705
 1710
 1715
 1720
 1725
 1730
 1735
 1740
 1745
 1750
 1755
 1760
 1765
 1770
 1775
 1780
 1785
 1790
 1795
 1800
 1805
 1810
 1815
 1820
 1825
 1830
 1835
 1840
 1845
 1850
 1855
 1860
 1865
 1870
 1875
 1880
 1885
 1890
 1895
 1900
 1905
 1910
 1915
 1920
 1925
 1930
 1935
 1940
 1945
 1950
 1955
 1960
 1965
 1970
 1975
 1980
 1985
 1990
 1995
 2000
 2005
 2010
 2015
 2020
 2025
 2030
 2035
 2040
 2045
 2050
 2055
 2060
 2065
 2070
 2075
 2080
 2085
 2090
 2095
 2100
 2105
 2110
 2115
 2120
 2125
 2130
 2135
 2140
 2145
 2150
 2155
 2160
 2165
 2170
 2175
 2180
 2185
 2190
 2195
 2200
 2205
 2210
 2215
 2220
 2225
 2230
 2235
 2240
 2245
 2250
 2255
 2260
 2265
 2270
 2275
 2280
 2285
 2290
 2295
 2300
 2305
 2310
 2315
 2320
 2325
 2330
 2335
 2340
 2345
 2350
 2355
 2360
 2365
 2370
 2375
 2380
 2385
 2390
 2395
 2400
 2405
 2410
 2415
 2420
 2425
 2430
 2435
 2440
 2445
 2450
 2455
 2460
 2465
 2470
 2475
 2480
 2485
 2490
 2495
 2500
 2505
 2510
 2515
 2520
 2525
 2530
 2535
 2540
 2545
 2550
 2555
 2560
 2565
 2570
 2575
 2580
 2585
 2590
 2595
 2600
 2605
 2610
 2615
 2620
 2625
 2630
 2635
 2640
 2645
 2650
 2655
 2660
 2665
 2670
 2675
 2680
 2685
 2690
 2695
 2700
 2705
 2710
 2715
 2720
 2725
 2730
 2735
 2740
 2745
 2750
 2755
 2760
 2765
 2770
 2775
 2780
 2785
 2790
 2795
 2800
 2805
 2810
 2815
 2820
 2825
 2830
 2835
 2840
 2845
 2850
 2855
 2860
 2865
 2870
 2875
 2880
 2885
 2890
 2895
 2900
 2905
 2910
 2915
 2920
 2925
 2930
 2935
 2940
 2945
 2950
 2955
 2960
 2965
 2970
 2975
 2980
 2985
 2990
 2995
 3000
 3005
 3010
 3015
 3020
 3025
 3030
 3035
 3040
 3045
 3050
 3055
 3060
 3065
 3070
 3075
 3080
 3085
 3090
 3095
 3100
 3105
 3110
 3115
 3120
 3125
 3130
 3135
 3140
 3145
 3150
 3155
 3160
 3165
 3170
 3175
 3180
 3185
 3190
 3195
 3200
 3205
 3210
 3215
 3220
 3225
 3230
 3235
 3240
 3245
 3250
 3255
 3260
 3265
 3270
 3275
 3280
 3285
 3290
 3295
 3300
 3305
 3310
 3315
 3320
 3325
 3330
 3335
 3340
 3345
 3350
 3355
 3360
 3365
 3370
 3375
 3380
 3385
 3390
 3395
 3400
 3405
 3410
 3415
 3420
 3425
 3430
 3435
 3440
 3445
 3450
 3455
 3460
 3465
 3470
 3475
 3480
 3485
 3490
 3495
 3500
 3505
 3510
 3515
 3520
 3525
 3530
 3535
 3540
 3545
 3550
 3555
 3560
 3565
 3570
 3575
 3580
 3585
 3590
 3595
 3600
 3605
 3610
 3615
 3620
 3625
 3630
 3635
 3640
 3645
 3650
 3655
 3660
 3665
 3670
 3675
 3680
 3685
 3690
 3695
 3700
 3705
 3710
 3715
 3720
 3725
 3730
 3735
 3740
 3745
 3750
 3755
 3760
 3765
 3770
 3775
 3780
 3785
 3790
 3795
 3800
 3805
 3810
 3815
 3820
 3825
 3830
 3835
 3840
 3845
 3850
 3855
 3860
 3865
 3870
 3875
 3880
 3885
 3890
 3895
 3900
 3905
 3910
 3915
 3920
 3925
 3930
 3935
 3940
 3945
 3950
 3955
 3960
 3965
 3970
 3975
 3980
 3985
 3990
 3995
 4000
 4005
 4010
 4015
 4020
 4025
 4030
 4035
 4040
 4045
 4050
 4055
 4060
 4065
 4070
 4075
 4080
 4085
 4090
 4095
 4100
 4105
 4110
 4115
 4120
 4125
 4130
 4135
 4140
 4145
 4150
 4155
 4160
 4165
 4170
 4175
 4180
 4185
 4190
 4195
 4200
 4205
 4210
 4215
 4220
 4225
 4230
 4235
 4240
 4245
 4250
 4255
 4260
 4265
 4270
 4275
 4280
 4285
 4290
 4295
 4300
 4305
 4310
 4315
 4320
 4325
 4330
 4335
 4340
 4345
 4350
 4355
 4360
 4365
 4370
 4375
 4380
 4385
 4390
 4395
 4400
 4405
 4410
 4415
 4420
 4425
 4430
 4435
 4440
 4445
 4450
 4455
 4460
 4465
 4470
 4475
 4480
 4485
 4490
 4495
 4500
 4505
 4510
 4515
 4520
 4525
 4530
 4535
 4540
 4545
 4550
 4555
 4560
 4565
 4570
 4575
 4580
 4585
 4590
 4595
 4600
 4605
 4610
 4615
 4620
 4625
 4630
 4635
 4640
 4645
 4650
 4655
 4660
 4665
 4670
 4675
 4680
 4685
 4690
 4695
 4700
 4705
 4710
 4715
 4720
 4725
 4730
 4735
 4740
 4745
 4750
 4755
 4760
 4765
 4770
 4775
 4780
 4785
 4790
 4795
 4800
 4805
 4810
 4815
 4820
 4825
 4830
 4835
 4840
 4845
 4850
 4855
 4860
 4865
 4870
 4875
 4880
 4885
 4890
 4895
 4900
 4905
 4910
 4915
 4920
 4925
 4930
 4935
 4940
 4945
 4950
 4955
 4960
 4965
 4970
 4975
 4980
 4985
 4990
 4995
 5000
 5005
 5010
 5015
 5020
 5025
 5030
 5035
 5040
 5045
 5050
 5055
 5060
 5065
 5070
 5075
 5080
 5085
 5090
 5095
 5100
 5105
 5110
 5115
 5120
 5125
 5130
 5135
 5140
 5145
 5150
 5155
 5160
 5165
 5170
 5175
 5180
 5185
 5190
 5195
 5200
 5205
 5210
 5215
 5220
 5225
 5230
 5235
 5240
 5245
 5250
 5255
 5260
 5265
 5270
 5275
 5280
 5285
 5290
 5295
 5300
 5305
 5310
 5315
 5320
 5325
 5330
 5335
 5340
 5345
 5350
 5355
 5360
 5365
 5370
 5375
 5380
 5385
 5390
 5395
 5400
 5405
 5410
 5415
 5420
 5425
 5430
 5435
 5440
 5445
 5450
 5455
 5460
 5465
 5470
 5475
 5480
 5485
 5490
 5495
 5500
 5505
 5510
 5515
 5520
 5525
 5530
 5535
 5540
 5545
 5550
 5555
 5560
 5565
 5570
 5575
 5580
 5585
 5590
 5595
 5600
 5605
 5610
 5615
 5620
 5625
 5630
 5635
 5640
 5645
 5650
 5655
 5660
 5665
 5670
 5675
 5680
 5685
 5690
 5695
 5700
 5705
 5710
 5715
 5720
 5725
 5730
 5735
 5740
 5745
 5750
 5755
 5760
 5765
 5770
 5775
 5780
 5785
 5790
 5795
 5800
 5805
 5810
 5815
 5820
 5825
 5830
 5835
 5840
 5845
 5850
 5855
 5860
 5865
 5870
 5875
 5880
 5885
 5890
 5895
 5900
 5905
 5910
 5915
 5920
 5925
 5930
 5935
 5940
 5945
 5950
 5955
 5960
 5965
 5970
 5975
 5980
 5985
 5990
 5995
 6000
 6005
 6010
 6015
 6020
 6025
 6030
 6035
 6040
 6045
 6050
 6055
 6060
 6065
 6070
 6075
 6080
 6085
 6090
 6095
 6100
 6105
 6110
 6115
 6120
 6125
 6130
 6135
 6140
 6145
 6150
 6155
 6160
 6165
 6170
 6175
 6180
 6185
 6190
 6195
 6200
 6205
 6210
 6215
 6220
 6225
 6230
 6235
 6240
 6245
 6250
 6255
 6260
 6265
 6270
 6275
 6280
 6285
 6290
 6295
 6300
 6305
 6310
 6315
 6320
 6325
 6330
 6335
 6340
 6345
 6350
 6355
 6360
 6365
 6370
 6375
 6380
 6385
 6390
 6395
 6400
 6405
 6410
 6415
 6420
 6425
 6430
 6435
 6440
 6445
 6450
 6455
 6460
 6465
 6470
 6475
 6480
 6485
 6490
 6495
 6500
 6505
 6510
 6515
 6520
 6525
 6530
 6535
 6540
 6545
 6550
 6555
 6560
 6565
 6570
 6575
 6580
 6585
 6590
 6595
 6600
 6605
 6610
 6615
 6620
 6625
 6630
 6635
 6640
 6645
 6650
 6655
 6660
 6665
 6670
 6675
 6680
 6685

Amongst compounds of formula I.4, those are preferred, wherein R³, R⁵ and R have the preferred meanings and the variable R¹ is selected from the group consisting of halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular from F, Cl, Br, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂, SO₂CH₃ and CH₂OCH₂CH₂OCH₃.

More preferred are compounds of formula I.4, wherein the variables R, R¹, R³ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, I, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, CH₂OCH₂CH₂OCH₃, SCF₃, SCHF₂ or SO₂CH₃;

R³ is hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular H, F, Cl, Br, CN, NO₂, CH₃, CH₂CH₃, CF₃, CHF₂, OCH₃, OCF₃, OCHF₂, SCH₃, SO₂CH₃ or SO₂CH₂CH₃; and

R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

Even more preferred are compounds of formula I.4, wherein the variables R, R¹, R³ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of cyclopropyl, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

With respect to their use, particular preference is given to the compounds of formula I.4 compiled in the tables 19-23 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 19 Compounds of the formula I.4 (compounds I.4-1 to I.4-168) in which R¹ is chlorine and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 20 Compounds of the formula I.4 (compounds I.4-169 to I.4-336) in which R¹ is nitro and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 21 Compounds of the formula I.4 (compounds I.4-337 to I.4-504), in which R¹ is methyl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 22 Compounds of the formula I.4 (compounds I.4-505 to I.4-672) in which R¹ is trifluoromethyl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 23 Compounds of the formula I.4 (compounds I.4-673 to I.4-840) in which R¹ is methylsulfonyl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa.

Table Aa

	R	R ³	R ⁵
Aa-1	CH ₃	Cl	cyclopropyl
Aa-2	CH ₃	Cl	OCH ₃
Aa-3	CH ₃	Cl	OCF ₃
Aa-4	CH ₃	Cl	SCH ₃
Aa-5	CH ₃	Cl	SOCH ₃
Aa-6	CH ₃	Cl	SO ₂ CH ₃
Aa-7	CH ₃	Cl	CH ₂ OCH ₃
Aa-8	CH ₃	F	cyclopropyl
Aa-9	CH ₃	F	OCH ₃
Aa-10	CH ₃	F	OCF ₃
Aa-11	CH ₃	F	SCH ₃
Aa-12	CH ₃	F	SOCH ₃
Aa-13	CH ₃	F	SO ₂ CH ₃
Aa-14	CH ₃	F	CH ₂ OCH ₃
Aa-15	CH ₃	CF ₃	cyclopropyl
Aa-16	CH ₃	CF ₃	OCH ₃
Aa-17	CH ₃	CF ₃	OCF ₃
Aa-18	CH ₃	CF ₃	SCH ₃
Aa-19	CH ₃	CF ₃	SOCH ₃
Aa-20	CH ₃	CF ₃	SO ₂ CH ₃
Aa-21	CH ₃	CF ₃	CH ₂ OCH ₃
Aa-22	CH ₃	SO ₂ CH ₃	cyclopropyl
Aa-23	CH ₃	SO ₂ CH ₃	OCH ₃
Aa-24	CH ₃	SO ₂ CH ₃	OCF ₃
Aa-25	CH ₃	SO ₂ CH ₃	SCH ₃
Aa-26	CH ₃	SO ₂ CH ₃	SOCH ₃


Aa-27	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Aa-28	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Aa-29	CH ₃	CN	cyclopropyl
Aa-30	CH ₃	CN	OCH ₃
Aa-31	CH ₃	CN	OCF ₃
Aa-32	CH ₃	CN	SCH ₃
Aa-33	CH ₃	CN	SOCH ₃
Aa-34	CH ₃	CN	SO ₂ CH ₃
Aa-35	CH ₃	CN	CH ₂ OCH ₃
Aa-36	CH ₃	H	cyclopropyl
Aa-37	CH ₃	H	OCH ₃
Aa-38	CH ₃	H	OCF ₃
Aa-39	CH ₃	H	SCH ₃
Aa-40	CH ₃	H	SOCH ₃
Aa-41	CH ₃	H	SO ₂ CH ₃
Aa-42	CH ₃	H	CH ₂ OCH ₃
Aa-43	CH ₂ CH ₃	Cl	cyclopropyl
Aa-44	CH ₂ CH ₃	Cl	OCH ₃
Aa-45	CH ₂ CH ₃	Cl	OCF ₃
Aa-46	CH ₂ CH ₃	Cl	SCH ₃
Aa-47	CH ₂ CH ₃	Cl	SOCH ₃
Aa-48	CH ₂ CH ₃	Cl	SO ₂ CH ₃
Aa-49	CH ₂ CH ₃	Cl	CH ₂ OCH ₃
Aa-50	CH ₂ CH ₃	F	cyclopropyl
Aa-51	CH ₂ CH ₃	F	OCH ₃
Aa-52	CH ₂ CH ₃	F	OCF ₃
Aa-53	CH ₂ CH ₃	F	SCH ₃
Aa-54	CH ₂ CH ₃	F	SOCH ₃
Aa-55	CH ₂ CH ₃	F	SO ₂ CH ₃
Aa-56	CH ₂ CH ₃	F	CH ₂ OCH ₃
Aa-57	CH ₂ CH ₃	CF ₃	cyclopropyl
Aa-58	CH ₂ CH ₃	CF ₃	OCH ₃
Aa-59	CH ₂ CH ₃	CF ₃	OCF ₃
Aa-60	CH ₂ CH ₃	CF ₃	SCH ₃
Aa-61	CH ₂ CH ₃	CF ₃	SOCH ₃
Aa-62	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃
Aa-63	CH ₂ CH ₃	CF ₃	CH ₂ OCH ₃
Aa-64	CH ₂ CH ₃	SO ₂ CH ₃	cyclopropyl
Aa-65	CH ₂ CH ₃	SO ₂ CH ₃	OCH ₃
Aa-66	CH ₂ CH ₃	SO ₂ CH ₃	OCF ₃
Aa-67	CH ₂ CH ₃	SO ₂ CH ₃	SCH ₃

Aa-68	CH ₂ CH ₃	SO ₂ CH ₃	SOCH ₃
Aa-69	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Aa-70	CH ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Aa-71	CH ₂ CH ₃	CN	cyclopropyl
Aa-72	CH ₂ CH ₃	CN	OCH ₃
Aa-73	CH ₂ CH ₃	CN	OCF ₃
Aa-74	CH ₂ CH ₃	CN	SCH ₃
Aa-75	CH ₂ CH ₃	CN	SOCH ₃
Aa-76	CH ₂ CH ₃	CN	SO ₂ CH ₃
Aa-77	CH ₂ CH ₃	CN	CH ₂ OCH ₃
Aa-78	CH ₂ CH ₃	H	cyclopropyl
Aa-79	CH ₂ CH ₃	H	OCH ₃
Aa-80	CH ₂ CH ₃	H	OCF ₃
Aa-81	CH ₂ CH ₃	H	SCH ₃
Aa-82	CH ₂ CH ₃	H	SOCH ₃
Aa-83	CH ₂ CH ₃	H	SO ₂ CH ₃
Aa-84	CH ₂ CH ₃	H	CH ₂ OCH ₃
Aa-85	CH ₂ OCH ₃	Cl	cyclopropyl
Aa-86	CH ₂ OCH ₃	Cl	OCH ₃
Aa-87	CH ₂ OCH ₃	Cl	OCF ₃
Aa-88	CH ₂ OCH ₃	Cl	SCH ₃
Aa-89	CH ₂ OCH ₃	Cl	SOCH ₃
Aa-90	CH ₂ OCH ₃	Cl	SO ₂ CH ₃
Aa-91	CH ₂ OCH ₃	Cl	CH ₂ OCH ₃
Aa-92	CH ₂ OCH ₃	F	cyclopropyl
Aa-93	CH ₂ OCH ₃	F	OCH ₃
Aa-94	CH ₂ OCH ₃	F	OCF ₃
Aa-95	CH ₂ OCH ₃	F	SCH ₃
Aa-96	CH ₂ OCH ₃	F	SOCH ₃
Aa-97	CH ₂ OCH ₃	F	SO ₂ CH ₃
Aa-98	CH ₂ OCH ₃	F	CH ₂ OCH ₃
Aa-99	CH ₂ OCH ₃	CF ₃	cyclopropyl
Aa-100	CH ₂ OCH ₃	CF ₃	OCH ₃
Aa-101	CH ₂ OCH ₃	CF ₃	OCF ₃
Aa-102	CH ₂ OCH ₃	CF ₃	SCH ₃
Aa-103	CH ₂ OCH ₃	CF ₃	SOCH ₃
Aa-104	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃
Aa-105	CH ₂ OCH ₃	CF ₃	CH ₂ OCH ₃
Aa-106	CH ₂ OCH ₃	SO ₂ CH ₃	cyclopropyl
Aa-107	CH ₂ OCH ₃	SO ₂ CH ₃	OCH ₃
Aa-108	CH ₂ OCH ₃	SO ₂ CH ₃	OCF ₃

Aa-109	CH ₂ OCH ₃	SO ₂ CH ₃	SCH ₃
Aa-110	CH ₂ OCH ₃	SO ₂ CH ₃	SOCH ₃
Aa-111	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Aa-112	CH ₂ OCH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Aa-113	CH ₂ OCH ₃	CN	cyclopropyl
Aa-114	CH ₂ OCH ₃	CN	OCH ₃
Aa-115	CH ₂ OCH ₃	CN	OCF ₃
Aa-116	CH ₂ OCH ₃	CN	SCH ₃
Aa-117	CH ₂ OCH ₃	CN	SOCH ₃
Aa-118	CH ₂ OCH ₃	CN	SO ₂ CH ₃
Aa-119	CH ₂ OCH ₃	CN	CH ₂ OCH ₃
Aa-120	CH ₂ OCH ₃	H	cyclopropyl
Aa-121	CH ₂ OCH ₃	H	OCH ₃
Aa-122	CH ₂ OCH ₃	H	OCF ₃
Aa-123	CH ₂ OCH ₃	H	SCH ₃
Aa-124	CH ₂ OCH ₃	H	SOCH ₃
Aa-125	CH ₂ OCH ₃	H	SO ₂ CH ₃
Aa-126	CH ₂ OCH ₃	H	CH ₂ OCH ₃
Aa-127	CH ₂ CH ₂ OCH ₃	Cl	cyclopropyl
Aa-128	CH ₂ CH ₂ OCH ₃	Cl	OCH ₃
Aa-129	CH ₂ CH ₂ OCH ₃	Cl	OCF ₃
Aa-130	CH ₂ CH ₂ OCH ₃	Cl	SCH ₃
Aa-131	CH ₂ CH ₂ OCH ₃	Cl	SOCH ₃
Aa-132	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃
Aa-133	CH ₂ CH ₂ OCH ₃	Cl	CH ₂ OCH ₃
Aa-134	CH ₂ CH ₂ OCH ₃	F	cyclopropyl
Aa-135	CH ₂ CH ₂ OCH ₃	F	OCH ₃
Aa-136	CH ₂ CH ₂ OCH ₃	F	OCF ₃
Aa-137	CH ₂ CH ₂ OCH ₃	F	SCH ₃
Aa-138	CH ₂ CH ₂ OCH ₃	F	SOCH ₃
Aa-139	CH ₂ CH ₂ OCH ₃	F	SO ₂ CH ₃
Aa-140	CH ₂ CH ₂ OCH ₃	F	CH ₂ OCH ₃
Aa-141	CH ₂ CH ₂ OCH ₃	CF ₃	cyclopropyl
Aa-142	CH ₂ CH ₂ OCH ₃	CF ₃	OCH ₃
Aa-143	CH ₂ CH ₂ OCH ₃	CF ₃	OCF ₃
Aa-144	CH ₂ CH ₂ OCH ₃	CF ₃	SCH ₃
Aa-145	CH ₂ CH ₂ OCH ₃	CF ₃	SOCH ₃
Aa-146	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃
Aa-147	CH ₂ CH ₂ OCH ₃	CF ₃	CH ₂ OCH ₃
Aa-148	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	cyclopropyl
Aa-149	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	OCH ₃

Aa-150	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	OCF ₃
Aa-151	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SCH ₃
Aa-152	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SOCH ₃
Aa-153	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Aa-154	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Aa-155	CH ₂ CH ₂ OCH ₃	CN	cyclopropyl
Aa-156	CH ₂ CH ₂ OCH ₃	CN	OCH ₃
Aa-157	CH ₂ CH ₂ OCH ₃	CN	OCF ₃
Aa-158	CH ₂ CH ₂ OCH ₃	CN	SCH ₃
Aa-159	CH ₂ CH ₂ OCH ₃	CN	SOCH ₃
Aa-160	CH ₂ CH ₂ OCH ₃	CN	SO ₂ CH ₃
Aa-161	CH ₂ CH ₂ OCH ₃	CN	CH ₂ OCH ₃
Aa-162	CH ₂ CH ₂ OCH ₃	H	cyclopropyl
Aa-163	CH ₂ CH ₂ OCH ₃	H	OCH ₃
Aa-164	CH ₂ CH ₂ OCH ₃	H	OCF ₃
Aa-165	CH ₂ CH ₂ OCH ₃	H	SCH ₃
Aa-166	CH ₂ CH ₂ OCH ₃	H	SOCH ₃
Aa-167	CH ₂ CH ₂ OCH ₃	H	SO ₂ CH ₃
Aa-168	CH ₂ CH ₂ OCH ₃	H	CH ₂ OCH ₃

A further particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is C-R¹, X² is C-R² and X⁴ is N. This compound is also referred to as compound of formula I.5, wherein R¹, R², R³, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R¹, R², R³, R⁵ and R in connection with compounds of formula I also apply for formulae I.5 as defined hereinafter. In formula I.5, the positions on the pyridine ring are designated by arabic numbers.

- 10 R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;
- R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, I, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂, SO₂CH₃ or CH₂OCH₂CH₂OCH₃;
- 15 R² is hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl, in particular

R³ is H, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular H, F, Cl, Br, CN, NO₂, CH₃, CH₂CH₃, CF₃, CHF₂, OCH₃, OCF₃, OCHF₂, SCH₃, SO₂CH₃ or SO₂CH₂CH₃; and

R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

Even more preferred are compounds of formula I.5, wherein the variables R, R¹, R², R³ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl;

R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazolinyl, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of cyclopropyl, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

With respect to their use, particular preference is given to the compounds of formula I.5 compiled in the tables 24-35 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 24 Compounds of the formula I.5 (compounds I.5-1 to I.5-840) in which R² is hydrogen and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 25 Compounds of the formula I.5 (compounds I.5-841 to I.5-1680), in which R² is SCH₃ and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 26 Compounds of the formula I.5 (compounds I.5-1681 to I.5-2520), in which R² is SOCH₃ and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 27 Compounds of the formula I.5 (compounds I.5-2521 to I.5-3360), in which R² is SO₂CH₃ and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 28 Compounds of the formula I.5 (compounds I.5-3361 to I.5-4200) in which R² is 2,2,2-trifluoroethoxymethyl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 29 Compounds of the formula I.5 (compounds I.5-4201 to I.5-5040) in which R² is 4,5-dihydroisoxazol-3-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 30 Compounds of the formula I.5 (compounds I.5-5041 to I.5-5880) in which R² is 5-methyl-4,5-dihydroisoxazol-3-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 31 Compounds of the formula I.5 (compounds I.5-5881 to I.5-6720) in which R² is 4,5-dihydroisoxazol-5-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 32 Compounds of the formula I.5 (compounds I.5-6721 to I.5-7560) in which R² is 3-methyl-4,5-dihydroisoxazol-5-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 33 Compounds of the formula I.5 (compounds I.5-7561 to I.5-8400) in which R² is isoxazol-3-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 34 Compounds of the formula I.5 (compounds I.5-8401 to I.5-9240) in which R² is 5-methyl-isoxazol-3-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab;

Table 35 Compounds of the formula I.5 (compounds I.5-9241 to I.5-10080) in which R² is 3-methyl-isoxazol-5-yl and the combination of R, R³, R¹ and R⁵ for a compound corresponds in each case to one row of Table Ab.

Table Ab:

	R	R ¹	R ³	R ⁵
Ab-1	CH ₃	Cl	Cl	cyclopropyl
Ab-2	CH ₃	Cl	Cl	OCH ₃
Ab-3	CH ₃	Cl	Cl	OCF ₃
Ab-4	CH ₃	Cl	Cl	SCH ₃
Ab-5	CH ₃	Cl	Cl	SOCH ₃
Ab-6	CH ₃	Cl	Cl	SO ₂ CH ₃
Ab-7	CH ₃	Cl	Cl	CH ₂ OCH ₃
Ab-8	CH ₃	Cl	F	cyclopropyl
Ab-9	CH ₃	Cl	F	OCH ₃
Ab-10	CH ₃	Cl	F	OCF ₃
Ab-11	CH ₃	Cl	F	SCH ₃
Ab-12	CH ₃	Cl	F	SOCH ₃
Ab-13	CH ₃	Cl	F	SO ₂ CH ₃
Ab-14	CH ₃	Cl	F	CH ₂ OCH ₃
Ab-15	CH ₃	Cl	CF ₃	cyclopropyl
Ab-16	CH ₃	Cl	CF ₃	OCH ₃
Ab-17	CH ₃	Cl	CF ₃	OCF ₃

Ab-18	CH ₃	Cl	CF ₃	SCH ₃
Ab-19	CH ₃	Cl	CF ₃	SOCH ₃
Ab-20	CH ₃	Cl	CF ₃	SO ₂ CH ₃
Ab-21	CH ₃	Cl	CF ₃	CH ₂ OCH ₃
Ab-22	CH ₃	Cl	SO ₂ CH ₃	cyclopropyl
Ab-23	CH ₃	Cl	SO ₂ CH ₃	OCH ₃
Ab-24	CH ₃	Cl	SO ₂ CH ₃	OCF ₃
Ab-25	CH ₃	Cl	SO ₂ CH ₃	SCH ₃
Ab-26	CH ₃	Cl	SO ₂ CH ₃	SOCH ₃
Ab-27	CH ₃	Cl	SO ₂ CH ₃	SO ₂ CH ₃
Ab-28	CH ₃	Cl	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-29	CH ₃	Cl	CN	cyclopropyl
Ab-30	CH ₃	Cl	CN	OCH ₃
Ab-31	CH ₃	Cl	CN	OCF ₃
Ab-32	CH ₃	Cl	CN	SCH ₃
Ab-33	CH ₃	Cl	CN	SOCH ₃
Ab-34	CH ₃	Cl	CN	SO ₂ CH ₃
Ab-35	CH ₃	Cl	CN	CH ₂ OCH ₃
Ab-36	CH ₃	Cl	H	cyclopropyl
Ab-37	CH ₃	Cl	H	OCH ₃
Ab-38	CH ₃	Cl	H	OCF ₃
Ab-39	CH ₃	Cl	H	SCH ₃
Ab-40	CH ₃	Cl	H	SOCH ₃
Ab-41	CH ₃	Cl	H	SO ₂ CH ₃
Ab-42	CH ₃	Cl	H	CH ₂ OCH ₃
Ab-43	CH ₃	CH ₃	Cl	cyclopropyl
Ab-44	CH ₃	CH ₃	Cl	OCH ₃
Ab-45	CH ₃	CH ₃	Cl	OCF ₃
Ab-46	CH ₃	CH ₃	Cl	SCH ₃
Ab-47	CH ₃	CH ₃	Cl	SOCH ₃
Ab-48	CH ₃	CH ₃	Cl	SO ₂ CH ₃
Ab-49	CH ₃	CH ₃	Cl	CH ₂ OCH ₃
Ab-50	CH ₃	CH ₃	F	cyclopropyl
Ab-51	CH ₃	CH ₃	F	OCH ₃
Ab-52	CH ₃	CH ₃	F	OCF ₃
Ab-53	CH ₃	CH ₃	F	SCH ₃
Ab-54	CH ₃	CH ₃	F	SOCH ₃
Ab-55	CH ₃	CH ₃	F	SO ₂ CH ₃
Ab-56	CH ₃	CH ₃	F	CH ₂ OCH ₃
Ab-57	CH ₃	CH ₃	CF ₃	cyclopropyl
Ab-58	CH ₃	CH ₃	CF ₃	OCH ₃

Ab-59	CH ₃	CH ₃	CF ₃	OCF ₃
Ab-60	CH ₃	CH ₃	CF ₃	SCH ₃
Ab-61	CH ₃	CH ₃	CF ₃	SOCH ₃
Ab-62	CH ₃	CH ₃	CF ₃	SO ₂ CH ₃
Ab-63	CH ₃	CH ₃	CF ₃	CH ₂ OCH ₃
Ab-64	CH ₃	CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-65	CH ₃	CH ₃	SO ₂ CH ₃	OCH ₃
Ab-66	CH ₃	CH ₃	SO ₂ CH ₃	OCF ₃
Ab-67	CH ₃	CH ₃	SO ₂ CH ₃	SCH ₃
Ab-68	CH ₃	CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-69	CH ₃	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-70	CH ₃	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-71	CH ₃	CH ₃	CN	cyclopropyl
Ab-72	CH ₃	CH ₃	CN	OCH ₃
Ab-73	CH ₃	CH ₃	CN	OCF ₃
Ab-74	CH ₃	CH ₃	CN	SCH ₃
Ab-75	CH ₃	CH ₃	CN	SOCH ₃
Ab-76	CH ₃	CH ₃	CN	SO ₂ CH ₃
Ab-77	CH ₃	CH ₃	CN	CH ₂ OCH ₃
Ab-78	CH ₃	CH ₃	H	cyclopropyl
Ab-79	CH ₃	CH ₃	H	OCH ₃
Ab-80	CH ₃	CH ₃	H	OCF ₃
Ab-81	CH ₃	CH ₃	H	SCH ₃
Ab-82	CH ₃	CH ₃	H	SOCH ₃
Ab-83	CH ₃	CH ₃	H	SO ₂ CH ₃
Ab-84	CH ₃	CH ₃	H	CH ₂ OCH ₃
Ab-85	CH ₃	CF ₃	Cl	cyclopropyl
Ab-86	CH ₃	CF ₃	Cl	OCH ₃
Ab-87	CH ₃	CF ₃	Cl	OCF ₃
Ab-88	CH ₃	CF ₃	Cl	SCH ₃
Ab-89	CH ₃	CF ₃	Cl	SOCH ₃
Ab-90	CH ₃	CF ₃	Cl	SO ₂ CH ₃
Ab-91	CH ₃	CF ₃	Cl	CH ₂ OCH ₃
Ab-92	CH ₃	CF ₃	F	cyclopropyl
Ab-93	CH ₃	CF ₃	F	OCH ₃
Ab-94	CH ₃	CF ₃	F	OCF ₃
Ab-95	CH ₃	CF ₃	F	SCH ₃
Ab-96	CH ₃	CF ₃	F	SOCH ₃
Ab-97	CH ₃	CF ₃	F	SO ₂ CH ₃
Ab-98	CH ₃	CF ₃	F	CH ₂ OCH ₃
Ab-99	CH ₃	CF ₃	CF ₃	cyclopropyl

Ab-100	CH ₃	CF ₃	CF ₃	OCH ₃
Ab-101	CH ₃	CF ₃	CF ₃	OCF ₃
Ab-102	CH ₃	CF ₃	CF ₃	SCH ₃
Ab-103	CH ₃	CF ₃	CF ₃	SOCH ₃
Ab-104	CH ₃	CF ₃	CF ₃	SO ₂ CH ₃
Ab-105	CH ₃	CF ₃	CF ₃	CH ₂ OCH ₃
Ab-106	CH ₃	CF ₃	SO ₂ CH ₃	cyclopropyl
Ab-107	CH ₃	CF ₃	SO ₂ CH ₃	OCH ₃
Ab-108	CH ₃	CF ₃	SO ₂ CH ₃	OCF ₃
Ab-109	CH ₃	CF ₃	SO ₂ CH ₃	SCH ₃
Ab-110	CH ₃	CF ₃	SO ₂ CH ₃	SOCH ₃
Ab-111	CH ₃	CF ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-112	CH ₃	CF ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-113	CH ₃	CF ₃	CN	cyclopropyl
Ab-114	CH ₃	CF ₃	CN	OCH ₃
Ab-115	CH ₃	CF ₃	CN	OCF ₃
Ab-116	CH ₃	CF ₃	CN	SCH ₃
Ab-117	CH ₃	CF ₃	CN	SOCH ₃
Ab-118	CH ₃	CF ₃	CN	SO ₂ CH ₃
Ab-119	CH ₃	CF ₃	CN	CH ₂ OCH ₃
Ab-120	CH ₃	CF ₃	H	cyclopropyl
Ab-121	CH ₃	CF ₃	H	OCH ₃
Ab-122	CH ₃	CF ₃	H	OCF ₃
Ab-123	CH ₃	CF ₃	H	SCH ₃
Ab-124	CH ₃	CF ₃	H	SOCH ₃
Ab-125	CH ₃	CF ₃	H	SO ₂ CH ₃
Ab-126	CH ₃	CF ₃	H	CH ₂ OCH ₃
Ab-127	CH ₃	SO ₂ CH ₃	Cl	cyclopropyl
Ab-128	CH ₃	SO ₂ CH ₃	Cl	OCH ₃
Ab-129	CH ₃	SO ₂ CH ₃	Cl	OCF ₃
Ab-130	CH ₃	SO ₂ CH ₃	Cl	SCH ₃
Ab-131	CH ₃	SO ₂ CH ₃	Cl	SOCH ₃
Ab-132	CH ₃	SO ₂ CH ₃	Cl	SO ₂ CH ₃
Ab-133	CH ₃	SO ₂ CH ₃	Cl	CH ₂ OCH ₃
Ab-134	CH ₃	SO ₂ CH ₃	F	cyclopropyl
Ab-135	CH ₃	SO ₂ CH ₃	F	OCH ₃
Ab-136	CH ₃	SO ₂ CH ₃	F	OCF ₃
Ab-137	CH ₃	SO ₂ CH ₃	F	SCH ₃
Ab-138	CH ₃	SO ₂ CH ₃	F	SOCH ₃
Ab-139	CH ₃	SO ₂ CH ₃	F	SO ₂ CH ₃
Ab-140	CH ₃	SO ₂ CH ₃	F	CH ₂ OCH ₃

Ab-141	CH ₃	SO ₂ CH ₃	CF ₃	cyclopropyl
Ab-142	CH ₃	SO ₂ CH ₃	CF ₃	OCH ₃
Ab-143	CH ₃	SO ₂ CH ₃	CF ₃	OCF ₃
Ab-144	CH ₃	SO ₂ CH ₃	CF ₃	SCH ₃
Ab-145	CH ₃	SO ₂ CH ₃	CF ₃	SOCH ₃
Ab-146	CH ₃	SO ₂ CH ₃	CF ₃	SO ₂ CH ₃
Ab-147	CH ₃	SO ₂ CH ₃	CF ₃	CH ₂ OCH ₃
Ab-148	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-149	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCH ₃
Ab-150	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCF ₃
Ab-151	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SCH ₃
Ab-152	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-153	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-154	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-155	CH ₃	SO ₂ CH ₃	CN	cyclopropyl
Ab-156	CH ₃	SO ₂ CH ₃	CN	OCH ₃
Ab-157	CH ₃	SO ₂ CH ₃	CN	OCF ₃
Ab-158	CH ₃	SO ₂ CH ₃	CN	SCH ₃
Ab-159	CH ₃	SO ₂ CH ₃	CN	SOCH ₃
Ab-160	CH ₃	SO ₂ CH ₃	CN	SO ₂ CH ₃
Ab-161	CH ₃	SO ₂ CH ₃	CN	CH ₂ OCH ₃
Ab-162	CH ₃	SO ₂ CH ₃	H	cyclopropyl
Ab-163	CH ₃	SO ₂ CH ₃	H	OCH ₃
Ab-164	CH ₃	SO ₂ CH ₃	H	OCF ₃
Ab-165	CH ₃	SO ₂ CH ₃	H	SCH ₃
Ab-166	CH ₃	SO ₂ CH ₃	H	SOCH ₃
Ab-167	CH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
Ab-168	CH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
Ab-169	CH ₃	NO ₂	Cl	cyclopropyl
Ab-170	CH ₃	NO ₂	Cl	OCH ₃
Ab-171	CH ₃	NO ₂	Cl	OCF ₃
Ab-172	CH ₃	NO ₂	Cl	SCH ₃
Ab-173	CH ₃	NO ₂	Cl	SOCH ₃
Ab-174	CH ₃	NO ₂	Cl	SO ₂ CH ₃
Ab-175	CH ₃	NO ₂	Cl	CH ₂ OCH ₃
Ab-176	CH ₃	NO ₂	F	cyclopropyl
Ab-177	CH ₃	NO ₂	F	OCH ₃
Ab-178	CH ₃	NO ₂	F	OCF ₃
Ab-179	CH ₃	NO ₂	F	SCH ₃
Ab-180	CH ₃	NO ₂	F	SOCH ₃
Ab-181	CH ₃	NO ₂	F	SO ₂ CH ₃

Ab-182	CH ₃	NO ₂	F	CH ₂ OCH ₃
Ab-183	CH ₃	NO ₂	CF ₃	cyclopropyl
Ab-184	CH ₃	NO ₂	CF ₃	OCH ₃
Ab-185	CH ₃	NO ₂	CF ₃	OCF ₃
Ab-186	CH ₃	NO ₂	CF ₃	SCH ₃
Ab-187	CH ₃	NO ₂	CF ₃	SOCH ₃
Ab-188	CH ₃	NO ₂	CF ₃	SO ₂ CH ₃
Ab-189	CH ₃	NO ₂	CF ₃	CH ₂ OCH ₃
Ab-190	CH ₃	NO ₂	SO ₂ CH ₃	cyclopropyl
Ab-191	CH ₃	NO ₂	SO ₂ CH ₃	OCH ₃
Ab-192	CH ₃	NO ₂	SO ₂ CH ₃	OCF ₃
Ab-193	CH ₃	NO ₂	SO ₂ CH ₃	SCH ₃
Ab-194	CH ₃	NO ₂	SO ₂ CH ₃	SOCH ₃
Ab-195	CH ₃	NO ₂	SO ₂ CH ₃	SO ₂ CH ₃
Ab-196	CH ₃	NO ₂	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-197	CH ₃	NO ₂	CN	cyclopropyl
Ab-198	CH ₃	NO ₂	CN	OCH ₃
Ab-199	CH ₃	NO ₂	CN	OCF ₃
Ab-200	CH ₃	NO ₂	CN	SCH ₃
Ab-201	CH ₃	NO ₂	CN	SOCH ₃
Ab-202	CH ₃	NO ₂	CN	SO ₂ CH ₃
Ab-203	CH ₃	NO ₂	CN	CH ₂ OCH ₃
Ab-204	CH ₃	NO ₂	H	cyclopropyl
Ab-205	CH ₃	NO ₂	H	OCH ₃
Ab-206	CH ₃	NO ₂	H	OCF ₃
Ab-207	CH ₃	NO ₂	H	SCH ₃
Ab-208	CH ₃	NO ₂	H	SOCH ₃
Ab-209	CH ₃	NO ₂	H	SO ₂ CH ₃
Ab-210	CH ₃	NO ₂	H	CH ₂ OCH ₃
Ab-211	CH ₂ CH ₃	Cl	Cl	cyclopropyl
Ab-212	CH ₂ CH ₃	Cl	Cl	OCH ₃
Ab-213	CH ₂ CH ₃	Cl	Cl	OCF ₃
Ab-214	CH ₂ CH ₃	Cl	Cl	SCH ₃
Ab-215	CH ₂ CH ₃	Cl	Cl	SOCH ₃
Ab-216	CH ₂ CH ₃	Cl	Cl	SO ₂ CH ₃
Ab-217	CH ₂ CH ₃	Cl	Cl	CH ₂ OCH ₃
Ab-218	CH ₂ CH ₃	Cl	F	cyclopropyl
Ab-219	CH ₂ CH ₃	Cl	F	OCH ₃
Ab-220	CH ₂ CH ₃	Cl	F	OCF ₃
Ab-221	CH ₂ CH ₃	Cl	F	SCH ₃
Ab-222	CH ₂ CH ₃	Cl	F	SOCH ₃

Ab-223	CH ₂ CH ₃	Cl	F	SO ₂ CH ₃
Ab-224	CH ₂ CH ₃	Cl	F	CH ₂ OCH ₃
Ab-225	CH ₂ CH ₃	Cl	CF ₃	cyclopropyl
Ab-226	CH ₂ CH ₃	Cl	CF ₃	OCH ₃
Ab-227	CH ₂ CH ₃	Cl	CF ₃	OCF ₃
Ab-228	CH ₂ CH ₃	Cl	CF ₃	SCH ₃
Ab-229	CH ₂ CH ₃	Cl	CF ₃	SOCH ₃
Ab-230	CH ₂ CH ₃	Cl	CF ₃	SO ₂ CH ₃
Ab-231	CH ₂ CH ₃	Cl	CF ₃	CH ₂ OCH ₃
Ab-232	CH ₂ CH ₃	Cl	SO ₂ CH ₃	cyclopropyl
Ab-233	CH ₂ CH ₃	Cl	SO ₂ CH ₃	OCH ₃
Ab-234	CH ₂ CH ₃	Cl	SO ₂ CH ₃	OCF ₃
Ab-235	CH ₂ CH ₃	Cl	SO ₂ CH ₃	SCH ₃
Ab-236	CH ₂ CH ₃	Cl	SO ₂ CH ₃	SOCH ₃
Ab-237	CH ₂ CH ₃	Cl	SO ₂ CH ₃	SO ₂ CH ₃
Ab-238	CH ₂ CH ₃	Cl	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-239	CH ₂ CH ₃	Cl	CN	cyclopropyl
Ab-240	CH ₂ CH ₃	Cl	CN	OCH ₃
Ab-241	CH ₂ CH ₃	Cl	CN	OCF ₃
Ab-242	CH ₂ CH ₃	Cl	CN	SCH ₃
Ab-243	CH ₂ CH ₃	Cl	CN	SOCH ₃
Ab-244	CH ₂ CH ₃	Cl	CN	SO ₂ CH ₃
Ab-245	CH ₂ CH ₃	Cl	CN	CH ₂ OCH ₃
Ab-246	CH ₂ CH ₃	Cl	H	cyclopropyl
Ab-247	CH ₂ CH ₃	Cl	H	OCH ₃
Ab-248	CH ₂ CH ₃	Cl	H	OCF ₃
Ab-249	CH ₂ CH ₃	Cl	H	SCH ₃
Ab-250	CH ₂ CH ₃	Cl	H	SOCH ₃
Ab-251	CH ₂ CH ₃	Cl	H	SO ₂ CH ₃
Ab-252	CH ₂ CH ₃	Cl	H	CH ₂ OCH ₃
Ab-253	CH ₂ CH ₃	CH ₃	Cl	cyclopropyl
Ab-254	CH ₂ CH ₃	CH ₃	Cl	OCH ₃
Ab-255	CH ₂ CH ₃	CH ₃	Cl	OCF ₃
Ab-256	CH ₂ CH ₃	CH ₃	Cl	SCH ₃
Ab-257	CH ₂ CH ₃	CH ₃	Cl	SOCH ₃
Ab-258	CH ₂ CH ₃	CH ₃	Cl	SO ₂ CH ₃
Ab-259	CH ₂ CH ₃	CH ₃	Cl	CH ₂ OCH ₃
Ab-260	CH ₂ CH ₃	CH ₃	F	cyclopropyl
Ab-261	CH ₂ CH ₃	CH ₃	F	OCH ₃
Ab-262	CH ₂ CH ₃	CH ₃	F	OCF ₃
Ab-263	CH ₂ CH ₃	CH ₃	F	SCH ₃

Ab-264	CH ₂ CH ₃	CH ₃	F	SOCH ₃
Ab-265	CH ₂ CH ₃	CH ₃	F	SO ₂ CH ₃
Ab-266	CH ₂ CH ₃	CH ₃	F	CH ₂ OCH ₃
Ab-267	CH ₂ CH ₃	CH ₃	CF ₃	cyclopropyl
Ab-268	CH ₂ CH ₃	CH ₃	CF ₃	OCH ₃
Ab-269	CH ₂ CH ₃	CH ₃	CF ₃	OCF ₃
Ab-270	CH ₂ CH ₃	CH ₃	CF ₃	SCH ₃
Ab-271	CH ₂ CH ₃	CH ₃	CF ₃	SOCH ₃
Ab-272	CH ₂ CH ₃	CH ₃	CF ₃	SO ₂ CH ₃
Ab-273	CH ₂ CH ₃	CH ₃	CF ₃	CH ₂ OCH ₃
Ab-274	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-275	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	OCH ₃
Ab-276	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	OCF ₃
Ab-277	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	SCH ₃
Ab-278	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-279	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-280	CH ₂ CH ₃	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-281	CH ₂ CH ₃	CH ₃	CN	cyclopropyl
Ab-282	CH ₂ CH ₃	CH ₃	CN	OCH ₃
Ab-283	CH ₂ CH ₃	CH ₃	CN	OCF ₃
Ab-284	CH ₂ CH ₃	CH ₃	CN	SCH ₃
Ab-285	CH ₂ CH ₃	CH ₃	CN	SOCH ₃
Ab-286	CH ₂ CH ₃	CH ₃	CN	SO ₂ CH ₃
Ab-287	CH ₂ CH ₃	CH ₃	CN	CH ₂ OCH ₃
Ab-288	CH ₂ CH ₃	CH ₃	H	cyclopropyl
Ab-289	CH ₂ CH ₃	CH ₃	H	OCH ₃
Ab-290	CH ₂ CH ₃	CH ₃	H	OCF ₃
Ab-291	CH ₂ CH ₃	CH ₃	H	SCH ₃
Ab-292	CH ₂ CH ₃	CH ₃	H	SOCH ₃
Ab-293	CH ₂ CH ₃	CH ₃	H	SO ₂ CH ₃
Ab-294	CH ₂ CH ₃	CH ₃	H	CH ₂ OCH ₃
Ab-295	CH ₂ CH ₃	CF ₃	Cl	cyclopropyl
Ab-296	CH ₂ CH ₃	CF ₃	Cl	OCH ₃
Ab-297	CH ₂ CH ₃	CF ₃	Cl	OCF ₃
Ab-298	CH ₂ CH ₃	CF ₃	Cl	SCH ₃
Ab-299	CH ₂ CH ₃	CF ₃	Cl	SOCH ₃
Ab-300	CH ₂ CH ₃	CF ₃	Cl	SO ₂ CH ₃
Ab-301	CH ₂ CH ₃	CF ₃	Cl	CH ₂ OCH ₃
Ab-302	CH ₂ CH ₃	CF ₃	F	cyclopropyl
Ab-303	CH ₂ CH ₃	CF ₃	F	OCH ₃
Ab-304	CH ₂ CH ₃	CF ₃	F	OCF ₃

Ab-305	CH ₂ CH ₃	CF ₃	F	SCH ₃
Ab-306	CH ₂ CH ₃	CF ₃	F	SOCH ₃
Ab-307	CH ₂ CH ₃	CF ₃	F	SO ₂ CH ₃
Ab-308	CH ₂ CH ₃	CF ₃	F	CH ₂ OCH ₃
Ab-309	CH ₂ CH ₃	CF ₃	CF ₃	cyclopropyl
Ab-310	CH ₂ CH ₃	CF ₃	CF ₃	OCH ₃
Ab-311	CH ₂ CH ₃	CF ₃	CF ₃	OCF ₃
Ab-312	CH ₂ CH ₃	CF ₃	CF ₃	SCH ₃
Ab-313	CH ₂ CH ₃	CF ₃	CF ₃	SOCH ₃
Ab-314	CH ₂ CH ₃	CF ₃	CF ₃	SO ₂ CH ₃
Ab-315	CH ₂ CH ₃	CF ₃	CF ₃	CH ₂ OCH ₃
Ab-316	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	cyclopropyl
Ab-317	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	OCH ₃
Ab-318	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	OCF ₃
Ab-319	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	SCH ₃
Ab-320	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	SOCH ₃
Ab-321	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-322	CH ₂ CH ₃	CF ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-323	CH ₂ CH ₃	CF ₃	CN	cyclopropyl
Ab-324	CH ₂ CH ₃	CF ₃	CN	OCH ₃
Ab-325	CH ₂ CH ₃	CF ₃	CN	OCF ₃
Ab-326	CH ₂ CH ₃	CF ₃	CN	SCH ₃
Ab-327	CH ₂ CH ₃	CF ₃	CN	SOCH ₃
Ab-328	CH ₂ CH ₃	CF ₃	CN	SO ₂ CH ₃
Ab-329	CH ₂ CH ₃	CF ₃	CN	CH ₂ OCH ₃
Ab-330	CH ₂ CH ₃	CF ₃	H	cyclopropyl
Ab-331	CH ₂ CH ₃	CF ₃	H	OCH ₃
Ab-332	CH ₂ CH ₃	CF ₃	H	OCF ₃
Ab-333	CH ₂ CH ₃	CF ₃	H	SCH ₃
Ab-334	CH ₂ CH ₃	CF ₃	H	SOCH ₃
Ab-335	CH ₂ CH ₃	CF ₃	H	SO ₂ CH ₃
Ab-336	CH ₂ CH ₃	CF ₃	H	CH ₂ OCH ₃
Ab-337	CH ₂ CH ₃	SO ₂ CH ₃	Cl	cyclopropyl
Ab-338	CH ₂ CH ₃	SO ₂ CH ₃	Cl	OCH ₃
Ab-339	CH ₂ CH ₃	SO ₂ CH ₃	Cl	OCF ₃
Ab-340	CH ₂ CH ₃	SO ₂ CH ₃	Cl	SCH ₃
Ab-341	CH ₂ CH ₃	SO ₂ CH ₃	Cl	SOCH ₃
Ab-342	CH ₂ CH ₃	SO ₂ CH ₃	Cl	SO ₂ CH ₃
Ab-343	CH ₂ CH ₃	SO ₂ CH ₃	Cl	CH ₂ OCH ₃
Ab-344	CH ₂ CH ₃	SO ₂ CH ₃	F	cyclopropyl
Ab-345	CH ₂ CH ₃	SO ₂ CH ₃	F	OCH ₃

Ab-346	CH ₂ CH ₃	SO ₂ CH ₃	F	OCF ₃
Ab-347	CH ₂ CH ₃	SO ₂ CH ₃	F	SCH ₃
Ab-348	CH ₂ CH ₃	SO ₂ CH ₃	F	SOCH ₃
Ab-349	CH ₂ CH ₃	SO ₂ CH ₃	F	SO ₂ CH ₃
Ab-350	CH ₂ CH ₃	SO ₂ CH ₃	F	CH ₂ OCH ₃
Ab-351	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	cyclopropyl
Ab-352	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	OCH ₃
Ab-353	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	OCF ₃
Ab-354	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	SCH ₃
Ab-355	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	SOCH ₃
Ab-356	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	SO ₂ CH ₃
Ab-357	CH ₂ CH ₃	SO ₂ CH ₃	CF ₃	CH ₂ OCH ₃
Ab-358	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-359	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCH ₃
Ab-360	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCF ₃
Ab-361	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SCH ₃
Ab-362	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-363	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-364	CH ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-365	CH ₂ CH ₃	SO ₂ CH ₃	CN	cyclopropyl
Ab-366	CH ₂ CH ₃	SO ₂ CH ₃	CN	OCH ₃
Ab-367	CH ₂ CH ₃	SO ₂ CH ₃	CN	OCF ₃
Ab-368	CH ₂ CH ₃	SO ₂ CH ₃	CN	SCH ₃
Ab-369	CH ₂ CH ₃	SO ₂ CH ₃	CN	SOCH ₃
Ab-370	CH ₂ CH ₃	SO ₂ CH ₃	CN	SO ₂ CH ₃
Ab-371	CH ₂ CH ₃	SO ₂ CH ₃	CN	CH ₂ OCH ₃
Ab-372	CH ₂ CH ₃	SO ₂ CH ₃	H	cyclopropyl
Ab-373	CH ₂ CH ₃	SO ₂ CH ₃	H	OCH ₃
Ab-374	CH ₂ CH ₃	SO ₂ CH ₃	H	OCF ₃
Ab-375	CH ₂ CH ₃	SO ₂ CH ₃	H	SCH ₃
Ab-376	CH ₂ CH ₃	SO ₂ CH ₃	H	SOCH ₃
Ab-377	CH ₂ CH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
Ab-378	CH ₂ CH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
Ab-379	CH ₂ CH ₃	NO ₂	Cl	cyclopropyl
Ab-380	CH ₂ CH ₃	NO ₂	Cl	OCH ₃
Ab-381	CH ₂ CH ₃	NO ₂	Cl	OCF ₃
Ab-382	CH ₂ CH ₃	NO ₂	Cl	SCH ₃
Ab-383	CH ₂ CH ₃	NO ₂	Cl	SOCH ₃
Ab-384	CH ₂ CH ₃	NO ₂	Cl	SO ₂ CH ₃
Ab-385	CH ₂ CH ₃	NO ₂	Cl	CH ₂ OCH ₃
Ab-386	CH ₂ CH ₃	NO ₂	F	cyclopropyl

Ab-387	CH ₂ CH ₃	NO ₂	F	OCH ₃
Ab-388	CH ₂ CH ₃	NO ₂	F	OCF ₃
Ab-389	CH ₂ CH ₃	NO ₂	F	SCH ₃
Ab-390	CH ₂ CH ₃	NO ₂	F	SOCH ₃
Ab-391	CH ₂ CH ₃	NO ₂	F	SO ₂ CH ₃
Ab-392	CH ₂ CH ₃	NO ₂	F	CH ₂ OCH ₃
Ab-393	CH ₂ CH ₃	NO ₂	CF ₃	cyclopropyl
Ab-394	CH ₂ CH ₃	NO ₂	CF ₃	OCH ₃
Ab-395	CH ₂ CH ₃	NO ₂	CF ₃	OCF ₃
Ab-396	CH ₂ CH ₃	NO ₂	CF ₃	SCH ₃
Ab-397	CH ₂ CH ₃	NO ₂	CF ₃	SOCH ₃
Ab-398	CH ₂ CH ₃	NO ₂	CF ₃	SO ₂ CH ₃
Ab-399	CH ₂ CH ₃	NO ₂	CF ₃	CH ₂ OCH ₃
Ab-400	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	cyclopropyl
Ab-401	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	OCH ₃
Ab-402	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	OCF ₃
Ab-403	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	SCH ₃
Ab-404	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	SOCH ₃
Ab-405	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	SO ₂ CH ₃
Ab-406	CH ₂ CH ₃	NO ₂	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-407	CH ₂ CH ₃	NO ₂	CN	cyclopropyl
Ab-408	CH ₂ CH ₃	NO ₂	CN	OCH ₃
Ab-409	CH ₂ CH ₃	NO ₂	CN	OCF ₃
Ab-410	CH ₂ CH ₃	NO ₂	CN	SCH ₃
Ab-411	CH ₂ CH ₃	NO ₂	CN	SOCH ₃
Ab-412	CH ₂ CH ₃	NO ₂	CN	SO ₂ CH ₃
Ab-413	CH ₂ CH ₃	NO ₂	CN	CH ₂ OCH ₃
Ab-414	CH ₂ CH ₃	NO ₂	H	cyclopropyl
Ab-415	CH ₂ CH ₃	NO ₂	H	OCH ₃
Ab-416	CH ₂ CH ₃	NO ₂	H	OCF ₃
Ab-417	CH ₂ CH ₃	NO ₂	H	SCH ₃
Ab-418	CH ₂ CH ₃	NO ₂	H	SOCH ₃
Ab-419	CH ₂ CH ₃	NO ₂	H	SO ₂ CH ₃
Ab-420	CH ₂ CH ₃	NO ₂	H	CH ₂ OCH ₃
Ab-421	CH ₂ OCH ₃	Cl	Cl	cyclopropyl
Ab-422	CH ₂ OCH ₃	Cl	Cl	OCH ₃
Ab-423	CH ₂ OCH ₃	Cl	Cl	OCF ₃
Ab-424	CH ₂ OCH ₃	Cl	Cl	SCH ₃
Ab-425	CH ₂ OCH ₃	Cl	Cl	SOCH ₃
Ab-426	CH ₂ OCH ₃	Cl	Cl	SO ₂ CH ₃
Ab-427	CH ₂ OCH ₃	Cl	Cl	CH ₂ OCH ₃

Ab-428	CH ₂ OCH ₃	Cl	F	cyclopropyl
Ab-429	CH ₂ OCH ₃	Cl	F	OCH ₃
Ab-430	CH ₂ OCH ₃	Cl	F	OCF ₃
Ab-431	CH ₂ OCH ₃	Cl	F	SCH ₃
Ab-432	CH ₂ OCH ₃	Cl	F	SOCH ₃
Ab-433	CH ₂ OCH ₃	Cl	F	SO ₂ CH ₃
Ab-434	CH ₂ OCH ₃	Cl	F	CH ₂ OCH ₃
Ab-435	CH ₂ OCH ₃	Cl	CF ₃	cyclopropyl
Ab-436	CH ₂ OCH ₃	Cl	CF ₃	OCH ₃
Ab-437	CH ₂ OCH ₃	Cl	CF ₃	OCF ₃
Ab-438	CH ₂ OCH ₃	Cl	CF ₃	SCH ₃
Ab-439	CH ₂ OCH ₃	Cl	CF ₃	SOCH ₃
Ab-440	CH ₂ OCH ₃	Cl	CF ₃	SO ₂ CH ₃
Ab-441	CH ₂ OCH ₃	Cl	CF ₃	CH ₂ OCH ₃
Ab-442	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	cyclopropyl
Ab-443	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	OCH ₃
Ab-444	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	OCF ₃
Ab-445	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SCH ₃
Ab-446	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SOCH ₃
Ab-447	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SO ₂ CH ₃
Ab-448	CH ₂ OCH ₃	Cl	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-449	CH ₂ OCH ₃	Cl	CN	cyclopropyl
Ab-450	CH ₂ OCH ₃	Cl	CN	OCH ₃
Ab-451	CH ₂ OCH ₃	Cl	CN	OCF ₃
Ab-452	CH ₂ OCH ₃	Cl	CN	SCH ₃
Ab-453	CH ₂ OCH ₃	Cl	CN	SOCH ₃
Ab-454	CH ₂ OCH ₃	Cl	CN	SO ₂ CH ₃
Ab-455	CH ₂ OCH ₃	Cl	CN	CH ₂ OCH ₃
Ab-456	CH ₂ OCH ₃	Cl	H	cyclopropyl
Ab-457	CH ₂ OCH ₃	Cl	H	OCH ₃
Ab-458	CH ₂ OCH ₃	Cl	H	OCF ₃
Ab-459	CH ₂ OCH ₃	Cl	H	SCH ₃
Ab-460	CH ₂ OCH ₃	Cl	H	SOCH ₃
Ab-461	CH ₂ OCH ₃	Cl	H	SO ₂ CH ₃
Ab-462	CH ₂ OCH ₃	Cl	H	CH ₂ OCH ₃
Ab-463	CH ₂ OCH ₃	CH ₃	Cl	cyclopropyl
Ab-464	CH ₂ OCH ₃	CH ₃	Cl	OCH ₃
Ab-465	CH ₂ OCH ₃	CH ₃	Cl	OCF ₃
Ab-466	CH ₂ OCH ₃	CH ₃	Cl	SCH ₃
Ab-467	CH ₂ OCH ₃	CH ₃	Cl	SOCH ₃
Ab-468	CH ₂ OCH ₃	CH ₃	Cl	SO ₂ CH ₃

Ab-469	CH ₂ OCH ₃	CH ₃	Cl	CH ₂ OCH ₃
Ab-470	CH ₂ OCH ₃	CH ₃	F	cyclopropyl
Ab-471	CH ₂ OCH ₃	CH ₃	F	OCH ₃
Ab-472	CH ₂ OCH ₃	CH ₃	F	OCF ₃
Ab-473	CH ₂ OCH ₃	CH ₃	F	SCH ₃
Ab-474	CH ₂ OCH ₃	CH ₃	F	SOCH ₃
Ab-475	CH ₂ OCH ₃	CH ₃	F	SO ₂ CH ₃
Ab-476	CH ₂ OCH ₃	CH ₃	F	CH ₂ OCH ₃
Ab-477	CH ₂ OCH ₃	CH ₃	CF ₃	cyclopropyl
Ab-478	CH ₂ OCH ₃	CH ₃	CF ₃	OCH ₃
Ab-479	CH ₂ OCH ₃	CH ₃	CF ₃	OCF ₃
Ab-480	CH ₂ OCH ₃	CH ₃	CF ₃	SCH ₃
Ab-481	CH ₂ OCH ₃	CH ₃	CF ₃	SOCH ₃
Ab-482	CH ₂ OCH ₃	CH ₃	CF ₃	SO ₂ CH ₃
Ab-483	CH ₂ OCH ₃	CH ₃	CF ₃	CH ₂ OCH ₃
Ab-484	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-485	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	OCH ₃
Ab-486	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	OCF ₃
Ab-487	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SCH ₃
Ab-488	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-489	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-490	CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-491	CH ₂ OCH ₃	CH ₃	CN	cyclopropyl
Ab-492	CH ₂ OCH ₃	CH ₃	CN	OCH ₃
Ab-493	CH ₂ OCH ₃	CH ₃	CN	OCF ₃
Ab-494	CH ₂ OCH ₃	CH ₃	CN	SCH ₃
Ab-495	CH ₂ OCH ₃	CH ₃	CN	SOCH ₃
Ab-496	CH ₂ OCH ₃	CH ₃	CN	SO ₂ CH ₃
Ab-497	CH ₂ OCH ₃	CH ₃	CN	CH ₂ OCH ₃
Ab-498	CH ₂ OCH ₃	CH ₃	H	cyclopropyl
Ab-499	CH ₂ OCH ₃	CH ₃	H	OCH ₃
Ab-500	CH ₂ OCH ₃	CH ₃	H	OCF ₃
Ab-501	CH ₂ OCH ₃	CH ₃	H	SCH ₃
Ab-502	CH ₂ OCH ₃	CH ₃	H	SOCH ₃
Ab-503	CH ₂ OCH ₃	CH ₃	H	SO ₂ CH ₃
Ab-504	CH ₂ OCH ₃	CH ₃	H	CH ₂ OCH ₃
Ab-505	CH ₂ OCH ₃	CF ₃	Cl	cyclopropyl
Ab-506	CH ₂ OCH ₃	CF ₃	Cl	OCH ₃
Ab-507	CH ₂ OCH ₃	CF ₃	Cl	OCF ₃
Ab-508	CH ₂ OCH ₃	CF ₃	Cl	SCH ₃
Ab-509	CH ₂ OCH ₃	CF ₃	Cl	SOCH ₃

Ab-510	CH ₂ OCH ₃	CF ₃	Cl	SO ₂ CH ₃
Ab-511	CH ₂ OCH ₃	CF ₃	Cl	CH ₂ OCH ₃
Ab-512	CH ₂ OCH ₃	CF ₃	F	cyclopropyl
Ab-513	CH ₂ OCH ₃	CF ₃	F	OCH ₃
Ab-514	CH ₂ OCH ₃	CF ₃	F	OCF ₃
Ab-515	CH ₂ OCH ₃	CF ₃	F	SCH ₃
Ab-516	CH ₂ OCH ₃	CF ₃	F	SOCH ₃
Ab-517	CH ₂ OCH ₃	CF ₃	F	SO ₂ CH ₃
Ab-518	CH ₂ OCH ₃	CF ₃	F	CH ₂ OCH ₃
Ab-519	CH ₂ OCH ₃	CF ₃	CF ₃	cyclopropyl
Ab-520	CH ₂ OCH ₃	CF ₃	CF ₃	OCH ₃
Ab-521	CH ₂ OCH ₃	CF ₃	CF ₃	OCF ₃
Ab-522	CH ₂ OCH ₃	CF ₃	CF ₃	SCH ₃
Ab-523	CH ₂ OCH ₃	CF ₃	CF ₃	SOCH ₃
Ab-524	CH ₂ OCH ₃	CF ₃	CF ₃	SO ₂ CH ₃
Ab-525	CH ₂ OCH ₃	CF ₃	CF ₃	CH ₂ OCH ₃
Ab-526	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	cyclopropyl
Ab-527	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	OCH ₃
Ab-528	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	OCF ₃
Ab-529	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SCH ₃
Ab-530	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SOCH ₃
Ab-531	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-532	CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-533	CH ₂ OCH ₃	CF ₃	CN	cyclopropyl
Ab-534	CH ₂ OCH ₃	CF ₃	CN	OCH ₃
Ab-535	CH ₂ OCH ₃	CF ₃	CN	OCF ₃
Ab-536	CH ₂ OCH ₃	CF ₃	CN	SCH ₃
Ab-537	CH ₂ OCH ₃	CF ₃	CN	SOCH ₃
Ab-538	CH ₂ OCH ₃	CF ₃	CN	SO ₂ CH ₃
Ab-539	CH ₂ OCH ₃	CF ₃	CN	CH ₂ OCH ₃
Ab-540	CH ₂ OCH ₃	CF ₃	H	cyclopropyl
Ab-541	CH ₂ OCH ₃	CF ₃	H	OCH ₃
Ab-542	CH ₂ OCH ₃	CF ₃	H	OCF ₃
Ab-543	CH ₂ OCH ₃	CF ₃	H	SCH ₃
Ab-544	CH ₂ OCH ₃	CF ₃	H	SOCH ₃
Ab-545	CH ₂ OCH ₃	CF ₃	H	SO ₂ CH ₃
Ab-546	CH ₂ OCH ₃	CF ₃	H	CH ₂ OCH ₃
Ab-547	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	cyclopropyl
Ab-548	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	OCH ₃
Ab-549	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	OCF ₃
Ab-550	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SCH ₃

Ab-551	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SOCH ₃
Ab-552	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SO ₂ CH ₃
Ab-553	CH ₂ OCH ₃	SO ₂ CH ₃	Cl	CH ₂ OCH ₃
Ab-554	CH ₂ OCH ₃	SO ₂ CH ₃	F	cyclopropyl
Ab-555	CH ₂ OCH ₃	SO ₂ CH ₃	F	OCH ₃
Ab-556	CH ₂ OCH ₃	SO ₂ CH ₃	F	OCF ₃
Ab-557	CH ₂ OCH ₃	SO ₂ CH ₃	F	SCH ₃
Ab-558	CH ₂ OCH ₃	SO ₂ CH ₃	F	SOCH ₃
Ab-559	CH ₂ OCH ₃	SO ₂ CH ₃	F	SO ₂ CH ₃
Ab-560	CH ₂ OCH ₃	SO ₂ CH ₃	F	CH ₂ OCH ₃
Ab-561	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	cyclopropyl
Ab-562	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	OCH ₃
Ab-563	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	OCF ₃
Ab-564	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SCH ₃
Ab-565	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SOCH ₃
Ab-566	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SO ₂ CH ₃
Ab-567	CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	CH ₂ OCH ₃
Ab-568	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-569	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCH ₃
Ab-570	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCF ₃
Ab-571	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SCH ₃
Ab-572	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-573	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-574	CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-575	CH ₂ OCH ₃	SO ₂ CH ₃	CN	cyclopropyl
Ab-576	CH ₂ OCH ₃	SO ₂ CH ₃	CN	OCH ₃
Ab-577	CH ₂ OCH ₃	SO ₂ CH ₃	CN	OCF ₃
Ab-578	CH ₂ OCH ₃	SO ₂ CH ₃	CN	SCH ₃
Ab-579	CH ₂ OCH ₃	SO ₂ CH ₃	CN	SOCH ₃
Ab-580	CH ₂ OCH ₃	SO ₂ CH ₃	CN	SO ₂ CH ₃
Ab-581	CH ₂ OCH ₃	SO ₂ CH ₃	CN	CH ₂ OCH ₃
Ab-582	CH ₂ OCH ₃	SO ₂ CH ₃	H	cyclopropyl
Ab-583	CH ₂ OCH ₃	SO ₂ CH ₃	H	OCH ₃
Ab-584	CH ₂ OCH ₃	SO ₂ CH ₃	H	OCF ₃
Ab-585	CH ₂ OCH ₃	SO ₂ CH ₃	H	SCH ₃
Ab-586	CH ₂ OCH ₃	SO ₂ CH ₃	H	SOCH ₃
Ab-587	CH ₂ OCH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
Ab-588	CH ₂ OCH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
Ab-589	CH ₂ OCH ₃	NO ₂	Cl	cyclopropyl
Ab-590	CH ₂ OCH ₃	NO ₂	Cl	OCH ₃
Ab-591	CH ₂ OCH ₃	NO ₂	Cl	OCF ₃

Ab-592	CH ₂ OCH ₃	NO ₂	Cl	SCH ₃
Ab-593	CH ₂ OCH ₃	NO ₂	Cl	SOCH ₃
Ab-594	CH ₂ OCH ₃	NO ₂	Cl	SO ₂ CH ₃
Ab-595	CH ₂ OCH ₃	NO ₂	Cl	CH ₂ OCH ₃
Ab-596	CH ₂ OCH ₃	NO ₂	F	cyclopropyl
Ab-597	CH ₂ OCH ₃	NO ₂	F	OCH ₃
Ab-598	CH ₂ OCH ₃	NO ₂	F	OCF ₃
Ab-599	CH ₂ OCH ₃	NO ₂	F	SCH ₃
Ab-600	CH ₂ OCH ₃	NO ₂	F	SOCH ₃
Ab-601	CH ₂ OCH ₃	NO ₂	F	SO ₂ CH ₃
Ab-602	CH ₂ OCH ₃	NO ₂	F	CH ₂ OCH ₃
Ab-603	CH ₂ OCH ₃	NO ₂	CF ₃	cyclopropyl
Ab-604	CH ₂ OCH ₃	NO ₂	CF ₃	OCH ₃
Ab-605	CH ₂ OCH ₃	NO ₂	CF ₃	OCF ₃
Ab-606	CH ₂ OCH ₃	NO ₂	CF ₃	SCH ₃
Ab-607	CH ₂ OCH ₃	NO ₂	CF ₃	SOCH ₃
Ab-608	CH ₂ OCH ₃	NO ₂	CF ₃	SO ₂ CH ₃
Ab-609	CH ₂ OCH ₃	NO ₂	CF ₃	CH ₂ OCH ₃
Ab-610	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	cyclopropyl
Ab-611	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	OCH ₃
Ab-612	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	OCF ₃
Ab-613	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SCH ₃
Ab-614	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SOCH ₃
Ab-615	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SO ₂ CH ₃
Ab-616	CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-617	CH ₂ OCH ₃	NO ₂	CN	cyclopropyl
Ab-618	CH ₂ OCH ₃	NO ₂	CN	OCH ₃
Ab-619	CH ₂ OCH ₃	NO ₂	CN	OCF ₃
Ab-620	CH ₂ OCH ₃	NO ₂	CN	SCH ₃
Ab-621	CH ₂ OCH ₃	NO ₂	CN	SOCH ₃
Ab-622	CH ₂ OCH ₃	NO ₂	CN	SO ₂ CH ₃
Ab-623	CH ₂ OCH ₃	NO ₂	CN	CH ₂ OCH ₃
Ab-624	CH ₂ OCH ₃	NO ₂	H	cyclopropyl
Ab-625	CH ₂ OCH ₃	NO ₂	H	OCH ₃
Ab-626	CH ₂ OCH ₃	NO ₂	H	OCF ₃
Ab-627	CH ₂ OCH ₃	NO ₂	H	SCH ₃
Ab-628	CH ₂ OCH ₃	NO ₂	H	SOCH ₃
Ab-629	CH ₂ OCH ₃	NO ₂	H	SO ₂ CH ₃
Ab-630	CH ₂ OCH ₃	NO ₂	H	CH ₂ OCH ₃
Ab-631	CH ₂ CH ₂ OCH ₃	Cl	Cl	cyclopropyl
Ab-632	CH ₂ CH ₂ OCH ₃	Cl	Cl	OCH ₃

Ab-633	CH ₂ CH ₂ OCH ₃	Cl	Cl	OCF ₃
Ab-634	CH ₂ CH ₂ OCH ₃	Cl	Cl	SCH ₃
Ab-635	CH ₂ CH ₂ OCH ₃	Cl	Cl	SOCH ₃
Ab-636	CH ₂ CH ₂ OCH ₃	Cl	Cl	SO ₂ CH ₃
Ab-637	CH ₂ CH ₂ OCH ₃	Cl	Cl	CH ₂ OCH ₃
Ab-638	CH ₂ CH ₂ OCH ₃	Cl	F	cyclopropyl
Ab-639	CH ₂ CH ₂ OCH ₃	Cl	F	OCH ₃
Ab-640	CH ₂ CH ₂ OCH ₃	Cl	F	OCF ₃
Ab-641	CH ₂ CH ₂ OCH ₃	Cl	F	SCH ₃
Ab-642	CH ₂ CH ₂ OCH ₃	Cl	F	SOCH ₃
Ab-643	CH ₂ CH ₂ OCH ₃	Cl	F	SO ₂ CH ₃
Ab-644	CH ₂ CH ₂ OCH ₃	Cl	F	CH ₂ OCH ₃
Ab-645	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	cyclopropyl
Ab-646	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	OCH ₃
Ab-647	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	OCF ₃
Ab-648	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	SCH ₃
Ab-649	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	SOCH ₃
Ab-650	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	SO ₂ CH ₃
Ab-651	CH ₂ CH ₂ OCH ₃	Cl	CF ₃	CH ₂ OCH ₃
Ab-652	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	cyclopropyl
Ab-653	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	OCH ₃
Ab-654	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	OCF ₃
Ab-655	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SCH ₃
Ab-656	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SOCH ₃
Ab-657	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	SO ₂ CH ₃
Ab-658	CH ₂ CH ₂ OCH ₃	Cl	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-659	CH ₂ CH ₂ OCH ₃	Cl	CN	cyclopropyl
Ab-660	CH ₂ CH ₂ OCH ₃	Cl	CN	OCH ₃
Ab-661	CH ₂ CH ₂ OCH ₃	Cl	CN	OCF ₃
Ab-662	CH ₂ CH ₂ OCH ₃	Cl	CN	SCH ₃
Ab-663	CH ₂ CH ₂ OCH ₃	Cl	CN	SOCH ₃
Ab-664	CH ₂ CH ₂ OCH ₃	Cl	CN	SO ₂ CH ₃
Ab-665	CH ₂ CH ₂ OCH ₃	Cl	CN	CH ₂ OCH ₃
Ab-666	CH ₂ CH ₂ OCH ₃	Cl	H	cyclopropyl
Ab-667	CH ₂ CH ₂ OCH ₃	Cl	H	OCH ₃
Ab-668	CH ₂ CH ₂ OCH ₃	Cl	H	OCF ₃
Ab-669	CH ₂ CH ₂ OCH ₃	Cl	H	SCH ₃
Ab-670	CH ₂ CH ₂ OCH ₃	Cl	H	SOCH ₃
Ab-671	CH ₂ CH ₂ OCH ₃	Cl	H	SO ₂ CH ₃
Ab-672	CH ₂ CH ₂ OCH ₃	Cl	H	CH ₂ OCH ₃
Ab-673	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	cyclopropyl

Ab-674	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	OCH ₃
Ab-675	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	OCF ₃
Ab-676	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	SCH ₃
Ab-677	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	SOCH ₃
Ab-678	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	SO ₂ CH ₃
Ab-679	CH ₂ CH ₂ OCH ₃	CH ₃	Cl	CH ₂ OCH ₃
Ab-680	CH ₂ CH ₂ OCH ₃	CH ₃	F	cyclopropyl
Ab-681	CH ₂ CH ₂ OCH ₃	CH ₃	F	OCH ₃
Ab-682	CH ₂ CH ₂ OCH ₃	CH ₃	F	OCF ₃
Ab-683	CH ₂ CH ₂ OCH ₃	CH ₃	F	SCH ₃
Ab-684	CH ₂ CH ₂ OCH ₃	CH ₃	F	SOCH ₃
Ab-685	CH ₂ CH ₂ OCH ₃	CH ₃	F	SO ₂ CH ₃
Ab-686	CH ₂ CH ₂ OCH ₃	CH ₃	F	CH ₂ OCH ₃
Ab-687	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	cyclopropyl
Ab-688	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	OCH ₃
Ab-689	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	OCF ₃
Ab-690	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	SCH ₃
Ab-691	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	SOCH ₃
Ab-692	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	SO ₂ CH ₃
Ab-693	CH ₂ CH ₂ OCH ₃	CH ₃	CF ₃	CH ₂ OCH ₃
Ab-694	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-695	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	OCH ₃
Ab-696	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	OCF ₃
Ab-697	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SCH ₃
Ab-698	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-699	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-700	CH ₂ CH ₂ OCH ₃	CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-701	CH ₂ CH ₂ OCH ₃	CH ₃	CN	cyclopropyl
Ab-702	CH ₂ CH ₂ OCH ₃	CH ₃	CN	OCH ₃
Ab-703	CH ₂ CH ₂ OCH ₃	CH ₃	CN	OCF ₃
Ab-704	CH ₂ CH ₂ OCH ₃	CH ₃	CN	SCH ₃
Ab-705	CH ₂ CH ₂ OCH ₃	CH ₃	CN	SOCH ₃
Ab-706	CH ₂ CH ₂ OCH ₃	CH ₃	CN	SO ₂ CH ₃
Ab-707	CH ₂ CH ₂ OCH ₃	CH ₃	CN	CH ₂ OCH ₃
Ab-708	CH ₂ CH ₂ OCH ₃	CH ₃	H	cyclopropyl
Ab-709	CH ₂ CH ₂ OCH ₃	CH ₃	H	OCH ₃
Ab-710	CH ₂ CH ₂ OCH ₃	CH ₃	H	OCF ₃
Ab-711	CH ₂ CH ₂ OCH ₃	CH ₃	H	SCH ₃
Ab-712	CH ₂ CH ₂ OCH ₃	CH ₃	H	SOCH ₃
Ab-713	CH ₂ CH ₂ OCH ₃	CH ₃	H	SO ₂ CH ₃
Ab-714	CH ₂ CH ₂ OCH ₃	CH ₃	H	CH ₂ OCH ₃


Ab-715	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	cyclopropyl
Ab-716	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	OCH ₃
Ab-717	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	OCF ₃
Ab-718	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	SCH ₃
Ab-719	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	SOCH ₃
Ab-720	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	SO ₂ CH ₃
Ab-721	CH ₂ CH ₂ OCH ₃	CF ₃	Cl	CH ₂ OCH ₃
Ab-722	CH ₂ CH ₂ OCH ₃	CF ₃	F	cyclopropyl
Ab-723	CH ₂ CH ₂ OCH ₃	CF ₃	F	OCH ₃
Ab-724	CH ₂ CH ₂ OCH ₃	CF ₃	F	OCF ₃
Ab-725	CH ₂ CH ₂ OCH ₃	CF ₃	F	SCH ₃
Ab-726	CH ₂ CH ₂ OCH ₃	CF ₃	F	SOCH ₃
Ab-727	CH ₂ CH ₂ OCH ₃	CF ₃	F	SO ₂ CH ₃
Ab-728	CH ₂ CH ₂ OCH ₃	CF ₃	F	CH ₂ OCH ₃
Ab-729	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	cyclopropyl
Ab-730	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	OCH ₃
Ab-731	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	OCF ₃
Ab-732	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	SCH ₃
Ab-733	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	SOCH ₃
Ab-734	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	SO ₂ CH ₃
Ab-735	CH ₂ CH ₂ OCH ₃	CF ₃	CF ₃	CH ₂ OCH ₃
Ab-736	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	cyclopropyl
Ab-737	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	OCH ₃
Ab-738	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	OCF ₃
Ab-739	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SCH ₃
Ab-740	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SOCH ₃
Ab-741	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-742	CH ₂ CH ₂ OCH ₃	CF ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-743	CH ₂ CH ₂ OCH ₃	CF ₃	CN	cyclopropyl
Ab-744	CH ₂ CH ₂ OCH ₃	CF ₃	CN	OCH ₃
Ab-745	CH ₂ CH ₂ OCH ₃	CF ₃	CN	OCF ₃
Ab-746	CH ₂ CH ₂ OCH ₃	CF ₃	CN	SCH ₃
Ab-747	CH ₂ CH ₂ OCH ₃	CF ₃	CN	SOCH ₃
Ab-748	CH ₂ CH ₂ OCH ₃	CF ₃	CN	SO ₂ CH ₃
Ab-749	CH ₂ CH ₂ OCH ₃	CF ₃	CN	CH ₂ OCH ₃
Ab-750	CH ₂ CH ₂ OCH ₃	CF ₃	H	cyclopropyl
Ab-751	CH ₂ CH ₂ OCH ₃	CF ₃	H	OCH ₃
Ab-752	CH ₂ CH ₂ OCH ₃	CF ₃	H	OCF ₃
Ab-753	CH ₂ CH ₂ OCH ₃	CF ₃	H	SCH ₃
Ab-754	CH ₂ CH ₂ OCH ₃	CF ₃	H	SOCH ₃
Ab-755	CH ₂ CH ₂ OCH ₃	CF ₃	H	SO ₂ CH ₃

Ab-756	CH ₂ CH ₂ OCH ₃	CF ₃	H	CH ₂ OCH ₃
Ab-757	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	cyclopropyl
Ab-758	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	OCH ₃
Ab-759	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	OCF ₃
Ab-760	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SCH ₃
Ab-761	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SOCH ₃
Ab-762	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	SO ₂ CH ₃
Ab-763	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	Cl	CH ₂ OCH ₃
Ab-764	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	cyclopropyl
Ab-765	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	OCH ₃
Ab-766	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	OCF ₃
Ab-767	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	SCH ₃
Ab-768	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	SOCH ₃
Ab-769	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	SO ₂ CH ₃
Ab-770	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	F	CH ₂ OCH ₃
Ab-771	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	cyclopropyl
Ab-772	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	OCH ₃
Ab-773	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	OCF ₃
Ab-774	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SCH ₃
Ab-775	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SOCH ₃
Ab-776	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	SO ₂ CH ₃
Ab-777	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CF ₃	CH ₂ OCH ₃
Ab-778	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	cyclopropyl
Ab-779	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCH ₃
Ab-780	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	OCF ₃
Ab-781	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SCH ₃
Ab-782	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SOCH ₃
Ab-783	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	SO ₂ CH ₃
Ab-784	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-785	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	cyclopropyl
Ab-786	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	OCH ₃
Ab-787	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	OCF ₃
Ab-788	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	SCH ₃
Ab-789	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	SOCH ₃
Ab-790	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	SO ₂ CH ₃
Ab-791	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	CN	CH ₂ OCH ₃
Ab-792	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	cyclopropyl
Ab-793	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	OCH ₃
Ab-794	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	OCF ₃
Ab-795	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SCH ₃
Ab-796	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SOCH ₃

Ab-797	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	SO ₂ CH ₃
Ab-798	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃	H	CH ₂ OCH ₃
Ab-799	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	cyclopropyl
Ab-800	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	OCH ₃
Ab-801	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	OCF ₃
Ab-802	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	SCH ₃
Ab-803	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	SOCH ₃
Ab-804	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	SO ₂ CH ₃
Ab-805	CH ₂ CH ₂ OCH ₃	NO ₂	Cl	CH ₂ OCH ₃
Ab-806	CH ₂ CH ₂ OCH ₃	NO ₂	F	cyclopropyl
Ab-807	CH ₂ CH ₂ OCH ₃	NO ₂	F	OCH ₃
Ab-808	CH ₂ CH ₂ OCH ₃	NO ₂	F	OCF ₃
Ab-809	CH ₂ CH ₂ OCH ₃	NO ₂	F	SCH ₃
Ab-810	CH ₂ CH ₂ OCH ₃	NO ₂	F	SOCH ₃
Ab-811	CH ₂ CH ₂ OCH ₃	NO ₂	F	SO ₂ CH ₃
Ab-812	CH ₂ CH ₂ OCH ₃	NO ₂	F	CH ₂ OCH ₃
Ab-813	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	cyclopropyl
Ab-814	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	OCH ₃
Ab-815	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	OCF ₃
Ab-816	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	SCH ₃
Ab-817	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	SOCH ₃
Ab-818	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	SO ₂ CH ₃
Ab-819	CH ₂ CH ₂ OCH ₃	NO ₂	CF ₃	CH ₂ OCH ₃
Ab-820	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	cyclopropyl
Ab-821	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	OCH ₃
Ab-822	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	OCF ₃
Ab-823	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SCH ₃
Ab-824	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SOCH ₃
Ab-825	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	SO ₂ CH ₃
Ab-826	CH ₂ CH ₂ OCH ₃	NO ₂	SO ₂ CH ₃	CH ₂ OCH ₃
Ab-827	CH ₂ CH ₂ OCH ₃	NO ₂	CN	cyclopropyl
Ab-828	CH ₂ CH ₂ OCH ₃	NO ₂	CN	OCH ₃
Ab-829	CH ₂ CH ₂ OCH ₃	NO ₂	CN	OCF ₃
Ab-830	CH ₂ CH ₂ OCH ₃	NO ₂	CN	SCH ₃
Ab-831	CH ₂ CH ₂ OCH ₃	NO ₂	CN	SOCH ₃
Ab-832	CH ₂ CH ₂ OCH ₃	NO ₂	CN	SO ₂ CH ₃
Ab-833	CH ₂ CH ₂ OCH ₃	NO ₂	CN	CH ₂ OCH ₃
Ab-834	CH ₂ CH ₂ OCH ₃	NO ₂	H	cyclopropyl
Ab-835	CH ₂ CH ₂ OCH ₃	NO ₂	H	OCH ₃
Ab-836	CH ₂ CH ₂ OCH ₃	NO ₂	H	OCF ₃
Ab-837	CH ₂ CH ₂ OCH ₃	NO ₂	H	SCH ₃

Ab-838	CH ₂ CH ₂ OCH ₃	NO ₂	H	SOCH ₃
Ab-839	CH ₂ CH ₂ OCH ₃	NO ₂	H	SO ₂ CH ₃
Ab-840	CH ₂ CH ₂ OCH ₃	NO ₂	H	CH ₂ OCH ₃

A further very preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is N, X⁴ is N and X² is CR². These compounds are also referred to as compound of formula I.6, wherein R², R³, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R², R³, R⁵ and R in connection with compounds of formula I also apply for formula I.6 as defined herein. In formula I.6, the positions on the pyrazine ring are designated by arabic numbers.

Amongst compounds of formula I.6, those are preferred, wherein R², R³, R⁵ and R have the preferred meanings mentioned above. Especially more preferred are compounds of formula I.6, wherein R³, R⁵ and R have the preferred meanings mentioned above and the variable R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl. In particular, R² is selected from hydrogen, methoxymethyl, ethoxymethyl, 2,2,2-trifluoroethoxymethyl, 2,2,2-trifluoroethoxyethyl, methylsulfonyl, methylsulfinyl, methylsulfanyl, 4,5-dihydroisoxazol-5-yl, 4,5-dihydroisoxazol-3-yl, 3-methyl-4,5-dihydroisoxazol-5-yl, 5-methyl-4,5-dihydroisoxazol-3-yl, isoxazol-5-yl, 3-methyl-isoxazol-5-yl, isoxazol-3-yl and 5-methyl-isoxazol-3-yl.

More preferred are compounds of formula I.6, wherein the variables R, R², R³, and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl, in particular hydrogen, methoxymethyl, ethoxymethyl, 2,2,2-trifluoroethoxymethyl, 2,2,2-trifluoroethoxyethyl, methylsulfonyl, methylsulfinyl, methylsulfanyl, 4,5-dihydroisoxazol-5-yl, 4,5-dihydroisoxazol-3-yl, 3-methyl-4,5-dihydroisoxazol-5-yl, 5-methyl-4,5-dihydroisoxazol-3-yl, isoxazol-5-yl, 3-methyl-isoxazol-5-yl, isoxazol-3-yl and 5-methyl-isoxazol-3-yl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular H, Cl, F, CF₃, SO₂CH₃ or CN; and

R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

Even more preferred are compounds of formula I.6, wherein the variables R, R², R³ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl, isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl;

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of cyclopropyl, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

With respect to their use, particular preference is given to the compounds of formula I.6 compiled in the tables 36-47 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 36 Compounds of the formula I.6 (compounds I.6-1 to I.6-168) in which R² is hydrogen and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 37 Compounds of the formula I.6 (compounds I.6-169 to I.6-336), in which R² is SCH₃ and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 38 Compounds of the formula I.6 (compounds I.6-337 to I.6-504), in which R² is SOCH₃ and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 39 Compounds of the formula I.6 (compounds I.6-505 to I.6-672), in which R² is SO₂CH₃ and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 40 Compounds of the formula I.6 (compounds I.6-673 to I.6-840) in which R² is 2,2,2-trifluoroethoxymethyl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 41 Compounds of the formula I.6 (compounds I.6-841 to I.6-1008) in which R² is 4,5-dihydroisoxazol-3-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 42 Compounds of the formula I.6 (compounds I.6-1009 to I.6-1176) in which R² is 5-methyl-4,5-dihydroisoxazol-3-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 43 Compounds of the formula I.6 (compounds I.6-1177 to I.6-1344) in which R² is 4,5-dihydroisoxazol-5-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;


Table 44 Compounds of the formula I.6 (compounds I.6-1345 to I.6-1512) in which R² is 3-methyl-4,5-dihydroisoxazol-5-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 45 Compounds of the formula I.6 (compounds I.6-1513 to I.6-1680) in which R² is isoxazol-3-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 46 Compounds of the formula I.6 (compounds I.6-1681 to I.6-1848) in which R² is 5-methyl-isoxazol-3-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

Table 47 Compounds of the formula I.6 (compounds I.6-1849 to I.6-2016) in which R² is 3-methyl-isoxazol-5-yl and the combination of R, R³ and R⁵ for a compound corresponds in each case to one row of Table Aa;

A further particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is C-R¹, X² is CR², X⁴ is N and R² together with R³ forms a fused 6-membered carbocycle. This compound is also referred to as compound of formula I.7, wherein R¹, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R¹, R⁵ and R in connection with compounds of formula I also apply for formula I.7 as defined hereinafter. In formula I.7, the positions on the quinoline ring are designated by arabic numbers.

Amongst compounds of formula I.7, those are preferred, wherein R⁵ and R have the preferred meanings and the variable R¹ is selected from the group consisting of halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular from F, Cl, Br, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂, SO₂CH₃ and CH₂OCH₂CH₂OCH₃.

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, I, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, CH₂OCH₂CH₂OCH₃, SCF₃, SCHF₂ or SO₂CH₃; and

R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

Even more preferred are compounds of formula I.7, wherein the variables R, R¹ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

5 R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of cyclopropyl, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

10 With respect to their use, particular preference is given to the compounds of formula I.7 compiled in the tables 48-54 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 48 Compounds of formula I.7 (compounds I.7-1 to I.7-20) in which R⁵ is cyclopropyl
15 and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

Table 49 Compounds of formula I.7 (compounds I.7-21 to I.7-40), in which R⁵ is OCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

20 Table 50 Compounds of formula I.7 (compounds I.7-41 to I.7-60), in which R⁵ is OCF₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.

Table 51 Compounds of formula I.7 (compounds I.7-61 to I.7-80) in which R⁵ is SCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table
25 Ac;

Table 52 Compounds of formula I.7 (compounds I.7-81 to I.7-100), in which R⁵ is SOCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

Table 53 Compounds of formula I.7 (compounds I.7-101 to I.7-120), in which R⁵ is SO₂CH₃
30 and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.

Table 54 Compounds of formula I.7 (compounds I.7-121 to I.7-140), in which R⁵ is CH₂OCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.


35

Table Ac:

	R	R ¹
Ac-1	CH ₃	Cl
Ac-2	CH ₂ CH ₃	Cl
Ac-3	CH ₂ OCH ₃	Cl
Ac-4	CH ₂ CH ₂ OCH ₃	Cl
Ac-5	CH ₃	CH ₃

	R	R ¹
Ac-6	CH ₂ CH ₃	CH ₃
Ac-7	CH ₂ OCH ₃	CH ₃
Ac-8	CH ₂ CH ₂ OCH ₃	CH ₃
Ac-9	CH ₃	CF ₃
Ac-10	CH ₂ CH ₃	CF ₃
Ac-11	CH ₂ OCH ₃	CF ₃
Ac-12	CH ₂ CH ₂ OCH ₃	CF ₃
Ac-13	CH ₃	SO ₂ CH ₃
Ac-14	CH ₂ CH ₃	SO ₂ CH ₃
Ac-15	CH ₂ OCH ₃	SO ₂ CH ₃
Ac-16	CH ₂ CH ₂ OCH ₃	SO ₂ CH ₃
Ac-17	CH ₃	NO ₂
Ac-18	CH ₂ CH ₃	NO ₂
Ac-19	CH ₂ OCH ₃	NO ₂
Ac-20	CH ₂ CH ₂ OCH ₃	NO ₂

A further particularly preferred embodiment of the present invention relates to compounds of formula I, wherein X¹ is C-R¹, X² is CR², X⁴ is N and R² together with R³ forms a fused 6-membered heterocycle, where the fused heterocycle has 1 nitrogen atom as ring member. This compound is also referred to as compound of formula I.8, wherein R¹, R⁵ and R are as defined hereinabove for compounds of formula I:



A skilled person will readily understand that the preferences given for R¹, R⁵ and R in connection with compounds of formula I also apply for formula I.8 as defined hereinafter. In formula I.8, the positions on the [1,8]-naphththyridine ring are designated by arabic numbers.

Amongst compounds of formula I.8, those are preferred, wherein R⁵ and R have the preferred meanings and the variable R¹ is selected from the group consisting of halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl, in particular from F, Cl, Br, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, SCF₃, SCHF₂, SO₂CH₃ and CH₂OCH₂CH₂OCH₃.

More preferred are compounds of formula I.8, wherein the variables R, R¹ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-

haloalkylthio or C₁-C₄-alkylsulfonyl, in particular F, Cl, Br, I, nitro, CH₃, CF₃, OCH₃, OCF₃, OCHF₂, CH₂OCH₂CH₂OCH₃, SCF₃, SCHF₂ or SO₂CH₃; and

R⁵ is selected from the group consisting of cyclopropyl, cyclopropylmethyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂, CH₂OCH₃, CH₂OCH₂CH₂OCH₃, OCF₃, OCHF₂, SCH₃, SCF₃, SCH₂CH₃, SCH₂CF₃, SOCH₃, SOCH₂CH₃, SO₂CH₃, SO₂CH₂CH₃.

Even more preferred are compounds of formula I.8, wherein the variables R, R¹ and R⁵ have the following meanings:

R is C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, in particular methyl, ethyl, methoxymethyl and methoxyethyl;

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl; and

R⁵ is selected from the group consisting of cyclopropyl, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

With respect to their use, particular preference is given to the compounds of formula I.8 compiled in the tables 55-61 below. Moreover, the groups mentioned for a substituent in the tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 55 Compounds of formula I.8 (compounds I.8-1 to I.8-20) in which R⁵ is cyclopropyl and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

Table 56 Compounds of formula I.8 (compounds I.8-21 to I.8-40), in which R⁵ is OCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

Table 57 Compounds of formula I.8 (compounds I.8-41 to I.8-60), in which R⁵ is OCF₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.

Table 58 Compounds of formula I.8 (compounds I.8-61 to I.8-80) in which R⁵ is SCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;

Table 59 Compounds of formula I.8 (compounds I.8-81 to I.8-100), in which R⁵ is SOCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac;


Table 60 Compounds of formula I.8 (compounds I.8-101 to I.8-120), in which R⁵ is SO₂CH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.

Table 61 Compounds of formula I.8 (compounds I.8-121 to I.8-140), in which R⁵ is CH₂OCH₃ and the combination of R and R¹ for a compound corresponds in each case to one row of Table Ac.

The compounds of the formula I can be prepared by standard methods of organic chemistry, e.g. by the methods described hereinafter in schemes 1 to 5. The substituents, variables and indices in schemes 1 to 5 are as defined above for formula I, if not otherwise specified.

The compounds of formula I can be prepared for instance as shown in the Scheme 1 below.

Scheme 1:



5


In Scheme 1, R, X¹, X², X⁴, R³ and R⁵ are as defined above. LG is a leaving group, such as halogen, in particular Cl, an anhydride residue or an active ester residue.

4-Amino-1,2,5-oxadiazole compounds of formula III can be reacted with benzoyl derivatives of formula II to afford compounds of the formula I. Especially in case of LG being halogen the reaction is suitably carried out in the presence of a base. Suitable bases are for example carbonates, such as lithium, sodium or potassium carbonates, amines, such as trimethylamine or triethylamine, and basic N-heterocycles, such as pyridine, 2,6-dimethylpyridine or 2,4,6-trimethylpyridine. Suitable solvents are in particular aprotic solvents such as pentane, hexane, heptane, octane, cyclohexane, dichloromethane, chloroform, 1,2-dichloroethane, benzene, chlorobenzene, toluene, the xylenes, dichlorobenzene, trimethylbenzene, pyridine, 2,6-dimethylpyridine, 2,4,6-trimethylpyridine, acetonitrile, diethyl ether, tetrahydrofuran, 2-methyl tetrahydrofuran, methyl tert-butylether, 1,4-dioxane, N,N-dimethyl formamide, N-methyl pyrrolidinone or mixtures thereof. The starting materials are generally reacted with one another in equimolar or nearly equimolar amounts at a reaction temperature usually in the range of -20°C to 100°C and preferably in the range of -5°C to 50°C.

20

Alternatively, compounds of formula I can also be prepared as shown in Scheme 2.

Scheme 2:



25

In Scheme 2, R, X¹, X², X⁴, R³ and R⁵ are as defined above.

Reaction of a 4-amino-1,2,5-oxadiazole compound III with a benzoic acid derivative of formula IV yields compound I. The reaction is preferably carried in the presence of a suitable activating agent which converts the acid group of compound IV into an activated ester or amide. For this purpose activating agents known in the art, such as 1,1'-carbonyldiimidazole (CDI), dicyclohexyl carbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) or 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide (T3P) can be employed. The activated


30

ester or amide can be formed, depending in particular on the specific activating agent used, either in situ by contacting compound IV with the activating agent in the presence of compound III, or in a separate step prior to the reaction with compound III. It may be advantageous, especially in cases where DCC or EDC are used as activating agent, to include further additives in the activating reaction, such as hydroxybenzotriazole (HOBt), nitrophenol, pentafluorophenol, 2,4,5-trichlorophenol or N-hydroxysuccinimide. It may further be advantageous to prepare the activated ester or amide in the presence of a base, for example a tertiary amine. The activated ester or amide is either in situ or subsequently reacted with the amine of formula III to afford the amide of formula I. The reaction normally takes place in anhydrous inert solvents, such as chlorinated hydrocarbons, e.g. dichloromethane or dichloroethane, ethers, e.g. tetrahydrofuran or 1,4-dioxane or carboxamides, e.g. N,N-dimethylformamide, N,N-dimethylacetamide or N-methylpyrrolidone. The reaction is ordinarily carried out at temperatures in the range from -20°C to +25°C.

The compounds of formula II and their respective benzoic acid precursors of formula IV can be obtained by purchase or can be prepared by processes known in the art or disclosed in the literature, e.g. in WO 2000039094, WO 2009115788, EP 316491 and EP 283261.

The 4-amino-1,2,5-oxadiazole compounds of the formula III are either commercially available or are obtainable according to methods known from the literature. For example, 3-alkyl-4-amino-1,2,5-oxadiazoles can be prepared from β -ketoesters pursuant to a procedure described in Russian Chemical Bulletin, Int. Ed., 54(4), 1032-1037 (2005), as depicted in Scheme 3.


Scheme 3:



In Scheme 3, R is as defined above and Et is ethyl.

As shown in Scheme 4, the compounds of the formula III, where R is halogen, can be prepared from commercially available 3,4-diamino-1,2,5-oxadiazole according to procedures described in the literature, e.g. by the Sandmeyer-type reaction disclosed in Heteroatom Chemistry, 15(3), 199-207 (2004).


Scheme 4:



As shown in Scheme 5, the compounds of the formula III, where R is a nucleophilic residue, can be prepared by introducing the nucleophilic residue via the substitution of a leaving group L, e.g. halogene, in the 4-position of the 1,2,5-oxadiazoles compounds of formula V in accordance to procedures disclosed, for example in Journal of Chemical Research, Synopses

(6), 190 (1985), in Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya (9), 2086-8 (1986) or in Russian Chemical Bulletin (Translation of Izvestiya Akademii Nauk, Seriya Khimicheskaya), 53(3), 596-614 (2004).

5 Scheme 5:



As a rule, the compounds of formula I including their stereoisomers, salts, tautomers and N-oxides, and their precursors in the synthesis process, can be prepared by the methods described above. If individual compounds can not be prepared via the above-described routes, they can be prepared by derivatization of other compounds I or the respective precursor or by customary modifications of the synthesis routes described. For example, in individual cases, certain compounds of formula I can advantageously be prepared from other compounds of formula I by derivatization, e.g. by ester hydrolysis, amidation, esterification, ether cleavage, olefination, reduction, oxidation and the like, or by customary modifications of the synthesis routes described.

The reaction mixtures are worked up in the customary manner, for example by mixing with water, separating the phases, and, if appropriate, purifying the crude products by chromatography, for example on alumina or on silica gel. Some of the intermediates and end products may be obtained in the form of colorless or pale brown viscous oils which are freed or purified from volatile components under reduced pressure and at moderately elevated temperature. If the intermediates and end products are obtained as solids, they may be purified by recrystallization or trituration.

The compounds I and their agriculturally suitable salts are useful as herbicides. They are useful as such or as an appropriately formulated composition. The herbicidal compositions comprising the compound I, in particular the preferred aspects thereof, control vegetation on non-crop areas very efficiently, especially at high rates of application. They act against broad-leaved weeds and weed grasses in crops such as wheat, rice, corn, soybeans and cotton without causing any significant damage to the crop plants. This effect is mainly observed at low rates of application.

Depending on the application method in question, the compounds I, in particular the preferred aspects thereof, or compositions comprising them can additionally be employed in a further number of crop plants for eliminating unwanted plants. Examples of suitable crops are the following:

Allium cepa, Ananas comosus, Arachis hypogaea, Asparagus officinalis, Avena sativa, Beta vulgaris spec. altissima, Beta vulgaris spec. rapa, Brassica napus var. napus, Brassica napus var. napobrassica, Brassica rapa var. silvestris, Brassica oleracea, Brassica nigra, Camellia sinensis, Carthamus tinctorius, Carya illinoensis, Citrus limon, Citrus sinensis, Coffea arabica (Coffea canephora, Coffea liberica), Cucumis sativus, Cynodon dactylon, Daucus

carota, *Elaeis guineensis*, *Fragaria vesca*, *Glycine max*, *Gossypium hirsutum*, (*Gossypium arboreum*, *Gossypium herbaceum*, *Gossypium vitifolium*), *Helianthus annuus*, *Hevea brasiliensis*, *Hordeum vulgare*, *Humulus lupulus*, *Ipomoea batatas*, *Juglans regia*, *Lens culinaris*, *Linum usitatissimum*, *Lycopersicon lycopersicum*, *Malus spec.*, *Manihot esculenta*,
5 *Medicago sativa*, *Musa spec.*, *Nicotiana tabacum* (*N.rustica*), *Olea europaea*, *Oryza sativa*,
Phaseolus lunatus, *Phaseolus vulgaris*, *Picea abies*, *Pinus spec.*, *Pistacia vera*, *Pisum sativum*,
Prunus avium, *Prunus persica*, *Pyrus communis*, *Prunus armeniaca*, *Prunus cerasus*, *Prunus dulcis* and *Prunus domestica*, *Ribes sylvestre*, *Ricinus communis*, *Saccharum officinarum*,
10 *Secale cereale*, *Sinapis alba*, *Solanum tuberosum*, *Sorghum bicolor* (*s. vulgare*), *Theobroma cacao*, *Trifolium pratense*, *Triticum aestivum*, *Triticale*, *Triticum durum*, *Vicia faba*, *Vitis vinifera*,
Zea mays.

The term "crop plants" also includes plants which have been modified by breeding, mutagenesis or genetic engineering. Genetically modified plants are plants whose genetic material has been modified in a manner which does not occur under natural conditions by
15 crossing, mutations or natural recombination (i.e. reassembly of the genetic information). Here, in general, one or more genes are integrated into the genetic material of the plant to improve the properties of the plant.

Accordingly, the term "crop plants" also includes plants which, by breeding and genetic engineering, have acquired tolerance to certain classes of herbicides, such as
20 hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors, acetolactate synthase (ALS) inhibitors, such as, for example, sulfonylureas (EP-A-0257993, US 5,013,659) or imidazolinones (see, for example, US 6,222,100, WO 01/82685, WO 00/26390, WO 97/41218, WO 98/02526, WO 98/02527, WO 04/106529, WO 05/20673, WO 03/14357, WO 03/13225, WO 03/14356, WO 04/16073), enolpyruvylshikimate 3-phosphate synthase (EPSPS) inhibitors, such as, for
25 example, glyphosate (see, for example, WO 92/00377), glutamine synthetase (GS) inhibitors, such as, for example, glufosinate (see, for example, EP-A-0242236, EP-A-242246), or oxynil herbicides (see, for example, US 5,559,024).

Numerous crop plants, for example Clearfield® oilseed rape, tolerant to imidazolinones, for example imazamox, have been generated with the aid of classic breeding methods
30 (mutagenesis). Crop plants such as soybeans, cotton, corn, beet and oilseed rape, resistant to glyphosate or glufosinate, which are available under the tradenames RoundupReady® (glyphosate) and Liberty Link® (glufosinate) have been generated with the aid of genetic engineering methods.

Accordingly, the term "crop plants" also includes plants which, with the aid of genetic
35 engineering, produce one or more toxins, for example those of the bacterial strain *Bacillus* spp. Toxins which are produced by such genetically modified plants include, for example, insecticidal proteins of *Bacillus* spp., in particular *B. thuringiensis*, such as the endotoxins Cry1Ab, Cry1Ac, Cry1F, Cry1Fa2, Cry2Ab, Cry3A, Cry3Bb1, Cry9c, Cry34Ab1 or Cry35Ab1; or vegetative insecticidal proteins (VIPs), for example VIP1, VIP2, VIP3, or VIP3A; insecticidal proteins of
40 nematode-colonizing bacteria, for example *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins of animal organisms, for example wasp, spider or scorpion toxins; fungal toxins, for example from Streptomyces; plant lectins, for example from peas or barley; agglutinins; proteinase

inhibitors, for example trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors, ribosome-inactivating proteins (RIPs), for example ricin, corn-RIP, abrin, luffin, saporin or bryodin; steroid-metabolizing enzymes, for example 3-hydroxysteroid oxidase, ecdysteroid-IDP glycosyl transferase, cholesterol oxidase, ecdysone inhibitors, or HMG-CoA reductase; ion channel blockers, for example inhibitors of sodium channels or calcium channels; juvenile hormone esterase; receptors of the diuretic hormone (helicokinin receptors); stilbene synthase, bibenzyl synthase, chitinases and glucanases. In the plants, these toxins may also be produced as pretoxins, hybrid proteins or truncated or otherwise modified proteins. Hybrid proteins are characterized by a novel combination of different protein domains (see, for example, WO 2002/015701). Further examples of such toxins or genetically modified plants which produce these toxins are disclosed in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/018810 and WO 03/052073. The methods for producing these genetically modified plants are known to the person skilled in the art and disclosed, for example, in the publications mentioned above. Numerous of the toxins mentioned above bestow, upon the plants by which they are produced, tolerance to pests from all taxonomic classes of arthropods, in particular to beetles (Coeleropta), dipterans (Diptera) and butterflies (Lepidoptera) and to nematodes (Nematoda).

Genetically modified plants which produce one or more genes coding for insecticidal toxins are described, for example, in the publications mentioned above, and some of them are commercially available, such as, for example, YieldGard® (corn varieties producing the toxin Cry1Ab), YieldGard® Plus (corn varieties which produce the toxins Cry1Ab and Cry3Bb1), Starlink® (corn varieties which produce the toxin Cry9c), Herculex® RW (corn varieties which produce the toxins Cry34Ab1, Cry35Ab1 and the enzyme phosphinothricin-N-acetyltransferase [PAT]); NuCOTN® 33B (cotton varieties which produce the toxin Cry1Ac), Bollgard® I (cotton varieties which produce the toxin Cry1Ac), Bollgard® II (cotton varieties which produce the toxins Cry1Ac and Cry2Ab2); VIPCOT® (cotton varieties which produce a VIP toxin); NewLeaf® (potato varieties which produce the toxin Cry3A); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, Protecta®, Bt11 (for example Agrisure® CB) and Bt176 from Syngenta Seeds SAS, France (corn varieties which produce the toxin Cry1Ab and the PAT enzyme), MIR604 from Syngenta Seeds SAS, France (corn varieties which produce a modified version of the toxin Cry3A, see WO 03/018810), MON 863 from Monsanto Europe S.A., Belgium (corn varieties which produce the toxin Cry3Bb1), IPC 531 from Monsanto Europe S.A., Belgium (cotton varieties which produce a modified version of the toxin Cry1Ac) and 1507 from Pioneer Overseas Corporation, Belgium (corn varieties which produce the toxin Cry1F and the PAT enzyme).

Accordingly, the term "crop plants" also includes plants which, with the aid of genetic engineering, produce one or more proteins which are more robust or have increased resistance to bacterial, viral or fungal pathogens, such as, for example, pathogenesis-related proteins (PR proteins, see EP-A 0 392 225), resistance proteins (for example potato varieties producing two resistance genes against *Phytophthora infestans* from the wild Mexican potato *Solanum bulbocastanum*) or T4 lysozyme (for example potato cultivars which, by producing this protein, are resistant to bacteria such as *Erwinia amylovora*).

Accordingly, the term "crop plants" also includes plants whose productivity has been improved with the aid of genetic engineering methods, for example by enhancing the potential yield (for example biomass, grain yield, starch, oil or protein content), tolerance to drought, salt or other limiting environmental factors or resistance to pests and fungal, bacterial and viral pathogens.

The term "crop plants" also includes plants whose ingredients have been modified with the aid of genetic engineering methods in particular for improving human or animal diet, for example by oil plants producing health-promoting long-chain omega 3 fatty acids or monounsaturated omega 9 fatty acids (for example Nexera® oilseed rape).

The term "crop plants" also includes plants which have been modified with the aid of genetic engineering methods for improving the production of raw materials, for example by increasing the amylopectin content of potatoes (Amflora® potato).

Furthermore, it has been found that the compounds of the formula I are also suitable for the defoliation and/or desiccation of plant parts, for which crop plants such as cotton, potato, oilseed rape, sunflower, soybean or field beans, in particular cotton, are suitable. In this regard, there have been found compositions for the desiccation and/or defoliation of plants, processes for preparing these compositions and methods for desiccating and/or defoliating plants using the compounds of the formula I.

As desiccants, the compounds of the formula I are particularly suitable for desiccating the above-ground parts of crop plants such as potato, oilseed rape, sunflower and soybean, but also cereals. This makes possible the fully mechanical harvesting of these important crop plants.

Also of economic interest is to facilitate harvesting, which is made possible by concentrating within a certain period of time the dehiscence, or reduction of adhesion to the tree, in citrus fruit, olives and other species and varieties of pomaceous fruit, stone fruit and nuts. The same mechanism, i.e. the promotion of the development of abscission tissue between fruit part or leaf part and shoot part of the plants is also essential for the readily controllable defoliation of useful plants, in particular cotton.

Moreover, a shortening of the time interval in which the individual cotton plants mature leads to an increased fiber quality after harvesting.

The compounds I, or the herbicidal compositions comprising the compounds I, can be used, for example, in the form of ready-to-spray aqueous solutions, powders, suspensions, also highly concentrated aqueous, oily or other suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for broadcasting, or granules, by means of spraying, atomizing, dusting, spreading, watering or treatment of the seed or mixing with the seed. The use forms depend on the intended purpose; in each case, they should ensure the finest possible distribution of the active ingredients according to the invention.

The herbicidal compositions comprise a herbicidally effective amount of at least one compound of the formula I or an agriculturally useful salt of I, and auxiliaries which are customary for the formulation of crop protection agents.

Examples of auxiliaries customary for the formulation of crop protection agents are inert auxiliaries, solid carriers, surfactants (such as dispersants, protective colloids, emulsifiers,

wetting agents and tackifiers), organic and inorganic thickeners, bactericides, antifreeze agents, antifoams, if appropriate colorants and, for seed formulations, adhesives.

Examples of thickeners (i.e. compounds which impart to the formulation modified flow properties, i.e. high viscosity in the state of rest and low viscosity in motion) are

5 polysaccharides, such as xanthan gum (Kelzan® from Kelco), Rhodopol® 23 (Rhône Poulenc) or Veegum® (from R.T. Vanderbilt), and also organic and inorganic sheet minerals, such as Attaclay® (from Engelhardt).

Examples of antifoams are silicone emulsions (such as, for example, Silikon® SRE, Wacker or Rhodorsil® from Rhodia), long-chain alcohols, fatty acids, salts of fatty acids,
10 organofluorine compounds and mixtures thereof.

Bactericides can be added for stabilizing the aqueous herbicidal formulation. Examples of bactericides are bactericides based on diclorophen and benzyl alcohol hemiformal (Proxel® from ICI or Acticide® RS from Thor Chemie and Kathon® MK from Rohm & Haas), and also isothiazolinone derivatives, such as alkylisothiazolinones and benzisothiazolinones (Acticide MBS
15 from Thor Chemie).

Examples of antifreeze agents are ethylene glycol, propylene glycol, urea or glycerol.

Examples of colorants are both sparingly water-soluble pigments and water-soluble dyes. Examples which may be mentioned are the dyes known under the names Rhodamin B, C.I. Pigment Red 112 and C.I. Solvent Red 1, and also pigment blue 15:4, pigment blue 15:3,
20 pigment blue 15:2, pigment blue 15:1, pigment blue 80, pigment yellow 1, pigment yellow 13, pigment red 112, pigment red 48:2, pigment red 48:1, pigment red 57:1, pigment red 53:1, pigment orange 43, pigment orange 34, pigment orange 5, pigment green 36, pigment green 7, pigment white 6, pigment brown 25, basic violet 10, basic violet 49, acid red 51, acid red 52, acid red 14, acid blue 9, acid yellow 23, basic red 10, basic red 108.

25 Examples of adhesives are polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

Suitable inert auxiliaries are, for example, the following:

mineral oil fractions of medium to high boiling point, such as kerosene and diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic
30 hydrocarbons, for example paraffin, tetrahydronaphthalene, alkylated naphthalenes and their derivatives, alkylated benzenes and their derivatives, alcohols such as methanol, ethanol, propanol, butanol and cyclohexanol, ketones such as cyclohexanone or strongly polar solvents, for example amines such as N-methylpyrrolidone, and water.

Solid carriers are mineral earths such as silicas, silica gels, silicates, talc, kaolin,
35 limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate and magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate and ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders, or other solid carriers.

40 Suitable surfactants (adjuvants, wetting agents, tackifiers, dispersants and also emulsifiers) are the alkali metal salts, alkaline earth metal salts and ammonium salts of aromatic sulfonic acids, for example lignosulfonic acids (e.g. Borrespers-types, Borregaard),

phenolsulfonic acids, naphthalenesulfonic acids (Morwet types, Akzo Nobel) and dibutynaphthalenesulfonic acid (Nekal types, BASF SE), and of fatty acids, alkyl- and alkylarylsulfonates, alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and octadecanols, and also of fatty alcohol glycol ethers, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene or of the naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenyl or tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, liginosulfite waste liquors and proteins, denatured proteins, polysaccharides (e.g. methylcellulose), hydrophobically modified starches, polyvinyl alcohol (Mowiol types Clariant), polycarboxylates (BASF SE, Sokalan types), polyalkoxylates, polyvinylamine (BASF SE, Lupamine types), polyethyleneimine (BASF SE, Lupasol types), polyvinylpyrrolidone and copolymers thereof.

Powders, materials for broadcasting and dusts can be prepared by mixing or grinding the active ingredients together with a solid carrier.

Granules, for example coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers.

Aqueous use forms can be prepared from emulsion concentrates, suspensions, pastes, wettable powders or water-dispersible granules by adding water. To prepare emulsions, pastes or oil dispersions, the compounds of the formula I or Ia, either as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetting agent, tackifier, dispersant or emulsifier. Alternatively, it is also possible to prepare concentrates comprising active substance, wetting agent, tackifier, dispersant or emulsifier and, if desired, solvent or oil, which are suitable for dilution with water.

The concentrations of the compounds of the formula I in the ready-to-use preparations can be varied within wide ranges. In general, the formulations comprise from 0.001 to 98% by weight, preferably 0.01 to 95% by weight of at least one active compound. The active compounds are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

The formulations or ready-to-use preparations may also comprise acids, bases or buffer systems, suitable examples being phosphoric acid or sulfuric acid, or urea or ammonia.

The compounds I of the invention can for example be formulated as follows:

1. Products for dilution with water

A Water-soluble concentrates

10 parts by weight of active compound are dissolved in 90 parts by weight of water or a water-soluble solvent. As an alternative, wetters or other adjuvants are added. The active compound dissolves upon dilution with water. This gives a formulation with an active compound content of 10% by weight.

B Dispersible concentrates

20 parts by weight of active compound are dissolved in 70 parts by weight of cyclohexanone with addition of 10 parts by weight of a dispersant, for example

polyvinylpyrrolidone. Dilution with water gives a dispersion. The active compound content is 20% by weight.

C Emulsifiable concentrates

15 parts by weight of active compound are dissolved in 75 parts by weight of an organic solvent (e.g. alkylaromatics) with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). Dilution with water gives an emulsion. The formulation has an active compound content of 15% by weight.

D Emulsions

25 parts by weight of active compound are dissolved in 35 parts by weight of an organic solvent (e.g. alkylaromatics) with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). This mixture is introduced into 30 parts by weight of water by means of an emulsifier (e.g. Ultraturrax) and made into a homogeneous emulsion. Dilution with water gives an emulsion. The formulation has an active compound content of 25% by weight.

E Suspensions

In an agitated ball mill, 20 parts by weight of active compound are comminuted with addition of 10 parts by weight of dispersants and wetters and 70 parts by weight of water or an organic solvent to give a fine active compound suspension. Dilution with water gives a stable suspension of the active compound. The active compound content in the formulation is 20% by weight.

F Water-dispersible granules and water-soluble granules

50 parts by weight of active compound are ground finely with addition of 50 parts by weight of dispersants and wetters and made into water-dispersible or water-soluble granules by means of technical appliances (for example extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active compound. The formulation has an active compound content of 50% by weight.

G Water-dispersible powders and water-soluble powders

75 parts by weight of active compound are ground in a rotor-stator mill with addition of 25 parts by weight of dispersants, wetters and silica gel. Dilution with water gives a stable dispersion or solution of the active compound. The active compound content of the formulation is 75% by weight.

H Gel formulations

In a ball mill, 20 parts by weight of active compound, 10 parts by weight of dispersant, 1 part by weight of gelling agent and 70 parts by weight of water or of an organic solvent are ground to give a fine suspension. Dilution with water gives a stable suspension with active compound content of 20% by weight.

2. Products to be applied undiluted

I Dusts

5 parts by weight of active compound are ground finely and mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dusting powder with an active compound content of 5% by weight.

J Granules (GR, FG, GG, MG)

0.5 parts by weight of active compound are ground finely and associated with 99.5 parts by weight of carriers. Current methods here are extrusion, spray-drying or the fluidized bed. This gives granules to be applied undiluted with an active compound content of 0.5% by weight.

K ULV solutions (UL)

5 10 parts by weight of active compound are dissolved in 90 parts by weight of an organic solvent, for example xylene. This gives a product to be applied undiluted with an active compound content of 10% by weight.

The compounds I or the herbicidal compositions comprising them can be applied pre- or post-emergence, or together with the seed of a crop plant. It is also possible to apply the
10 herbicidal compositions or active compounds by applying seed, pretreated with the herbicidal compositions or active compounds, of a crop plant. If the active compounds are less well tolerated by certain crop plants, application techniques may be used in which the herbicidal compositions are sprayed, with the aid of the spraying equipment, in such a way that as far as possible they do not come into contact with the leaves of the sensitive crop plants, while the
15 active compounds reach the leaves of undesirable plants growing underneath, or the bare soil surface (post-directed, lay-by).

In a further embodiment, the compounds of the formula I or the herbicidal compositions can be applied by treating seed.

The treatment of seed comprises essentially all procedures familiar to the person skilled in
20 the art (seed dressing, seed coating, seed dusting, seed soaking, seed film coating, seed multilayer coating, seed encrusting, seed dripping and seed pelleting) based on the compounds of the formula I according to the invention or the compositions prepared therefrom. Here, the herbicidal compositions can be applied diluted or undiluted.

The term seed comprises seed of all types, such as, for example, corns, seeds, fruits,
25 tubers, cuttings and similar forms. Here, preferably, the term seed describes corns and seeds.

The seed used can be seed of the useful plants mentioned above, but also the seed of transgenic plants or plants obtained by customary breeding methods.

The rates of application of active compound are from 0.001 to 3.0, preferably 0.01 to 1.0, kg/ha of active substance (a.s.), depending on the control target, the season, the target plants
30 and the growth stage. To treat the seed, the compounds I are generally employed in amounts of from 0.001 to 10 kg per 100 kg of seed.

It may also be advantageous to use the compounds of the formula I in combination with safeners. Safeners are chemical compounds which prevent or reduce damage to useful plants without substantially affecting the herbicidal action of the compounds of the formula I on
35 unwanted plants. They can be used both before sowing (for example in the treatment of seed, or on cuttings or seedlings) and before or after the emergence of the useful plant. The safeners and the compounds of the formula I can be used simultaneously or in succession. Suitable safeners are, for example, (quinolin-8-oxy)acetic acids, 1-phenyl-5-haloalkyl-1*H*-1,2,4-triazole-3-carboxylic acids, 1-phenyl-4,5-dihydro-5-alkyl-1*H*-pyrazole-3,5-dicarboxylic acids, 4,5-dihydro-
40 5,5-diaryl-3-isoxazolecarboxylic acids, dichloroacetamides, alpha-oximinophenylacetoneitriles, acetophenone oximes, 4,6-dihalo-2-phenylpyrimidines, N-[[4-(aminocarbonyl)phenyl]sulfonyl]-2-benzamides, 1,8-naphthalic anhydride, 2-halo-4-(haloalkyl)-5-thiazolecarboxylic acids,

phosphorothiolates and O-phenyl N-alkylcarbamates and their agriculturally useful salts and, provided that they have an acid function, their agriculturally useful derivatives, such as amides, esters and thioesters.

To broaden the activity spectrum and to obtain synergistic effects, the compounds of the formula I can be mixed and jointly applied with numerous representatives of other herbicidal or growth-regulating groups of active compounds or with safeners. Suitable mixing partners are, for example, 1,2,4-thiadiazoles, 1,3,4-thiadiazoles, amides, aminophosphoric acid and its derivatives, aminotriazoles, anilides, aryloxy/heteroaryloxyalkanoic acids and their derivatives, benzoic acid and its derivatives, benzothiadiazinones, 2-(heteroaryl/aryl)-1,3-cyclohexanediones, heteroaryl aryl ketones, benzylisoxazolidinones, meta-CF₃-phenyl derivatives, carbamates, quinoline carboxylic acid and its derivatives, chloroacetanilides, cyclohexenone oxime ether derivatives, diazines, dichloropropionic acid and its derivatives, dihydrobenzofurans, dihydrofuran-3-ones, dinitroanilines, dinitrophenols, diphenyl ethers, dipyridyls, halocarboxylic acids and their derivatives, ureas, 3-phenyluracils, imidazoles, imidazolinones, N-phenyl-3,4,5,6-tetrahydrophthalimides, oxadiazoles, oxiranes, phenols, aryloxy- and heteroaryloxyphenoxypropionic esters, phenylacetic acid and its derivatives, 2-phenylpropionic acid and its derivatives, pyrazoles, phenylpyrazoles, pyridazines, pyridinecarboxylic acid and its derivatives, pyrimidyl ethers, sulfonamides, sulfonylureas, triazines, triazinones, triazolinones, triazolecarboxamides, uracils and also phenylpyrazolines and isoxazolines and their derivatives.

Moreover, it may be useful to apply the compounds I alone or in combination with other herbicides or else also mixed with further crop protection agents, jointly, for example with compositions for controlling pests or phytopathogenic fungi or bacteria. Also of interest is the miscibility with mineral salt solutions which are employed for alleviating nutritional and trace element deficiencies. Other additives such as nonphytotoxic oils and oil concentrates may also be added.

Examples of herbicides which can be used in combination with the pyridine compounds of the formula I according to the present invention are:

b1) from the group of the lipid biosynthesis inhibitors:

alloxydim, alloxydim-sodium, butoxydim, clethodim, clodinafop, clodinafop-propargyl, cycloxydim, cyhalofop, cyhalofop-butyl, diclofop, diclofop-methyl, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-P, fenoxaprop-P-ethyl, fluazifop, fluazifop-butyl, fluazifop-P, fluazifop-P-butyl, haloxyfop, haloxyfop-methyl, haloxyfop-P, haloxyfop-P-methyl, metamifop, pinoxaden, profoxydim, propaquizafop, quizalofop, quizalofop-ethyl, quizalofop-tefuryl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, sethoxydim, tepraloxym, tralkoxydim, benfuresate, butylate, cycloate, dalapon, dimepiperate, EPTC, esprocarb, ethofumesate, flupropanate, molinate, orbencarb, pebulate, prosulfocarb, TCA, thiobencarb, tiocarbazil, triallate and vernolate;

b2) from the group of the ALS inhibitors:

amidosulfuron, azimsulfuron, bensulfuron, bensulfuron-methyl, bispyribac, bispyribac-sodium, chlorimuron, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cloransulam, cloransulam-methyl, cyclosulfamuron, diclosulam, ethametsulfuron, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, florasulam, flucarbazone, flucarbazone-sodium, flucetosulfuron, flumetsulam,

flupyrsulfuron, flupyrsulfuron-methyl-sodium, foramsulfuron, halosulfuron, halosulfuron-methyl, imazamethabenz, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, iodosulfuron, iodosulfuron-methyl-sodium, mesosulfuron, metosulam, metsulfuron, metsulfuron-methyl, nicosulfuron, orthosulfamuron, oxasulfuron, penoxsulam, primisulfuron, primisulfuron-methyl, propoxycarbazone, propoxycarbazone-sodium, prosulfuron, pyrazosulfuron, pyrazosulfuron-ethyl, pyribenzoxim, pyrimisulfan, pyriftalid, pyriminobac, pyriminobac-methyl, pyrithiobac, pyrithiobac-sodium, pyroxsulam, rimsulfuron, sulfometuron, sulfometuron-methyl, sulfosulfuron, thienencarbazone, thienencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, triasulfuron, tribenuron, tribenuron-methyl, trifloxysulfuron, triflusulfuron, triflusulfuron-methyl and tritosulfuron;

b3) from the group of the photosynthesis inhibitors:

ametryn, amicarbazone, atrazine, bentazone, bentazone-sodium, bromacil, bromofenoxim, bromoxynil and its salts and esters, chlorobromuron, chloridazone, chlorotoluron, chloroxuron, cyanazine, desmedipham, desmetryn, dimefuron, dimethametryn, diquat, diquat-dibromide, diuron, fluometuron, hexazinone, ioxynil and its salts and esters, isoproturon, isouron, karbutilate, lenacil, linuron, metamitron, methabenzthiazuron, metobenzuron, metoxuron, metribuzin, monolinuron, neburon, paraquat, paraquat-dichloride, paraquat-dimethylsulfate, pentanochlor, phenmedipham, phenmedipham-ethyl, prometon, prometryn, propanil, propazine, pyridafol, pyridate, siduron, simazine, simetryn, tebuthiuron, terbacil, terbumeton, terbuthylazine, terbutryn, thidiazuron and trietazine;

b4) from the group of the protoporphyrinogen-IX oxidase inhibitors:

acifluorfen, acifluorfen-sodium, azafenidin, bencarbazone, benzfendizone, bifenox, butafenacil, carfentrazone, carfentrazone-ethyl, chlomethoxyfen, cinidon-ethyl, fluazolate, flufenpyr, flufenpyr-ethyl, flumiclorac, flumiclorac-pentyl, flumioxazin, fluoroglycofen, fluoroglycofen-ethyl, fluthiacet, fluthiacet-methyl, fomesafen, halosafen, lactofen, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, profluazol, pyracilonil, pyraflufen, pyraflufen-ethyl, saflufenacil, sulfentrazone, thidiazimin, 2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2*H*)-pyrimidinyl]-4-fluoro-N-[(isopropyl)methylsulfamoyl]benzamide (H-1; CAS 372137-35-4), ethyl [3-[2-chloro-4-fluoro-5-(1-methyl-6-trifluoromethyl-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-3-yl)phenoxy]-2-pyridyloxy]acetate (H-2; CAS 353292-31-6), N-ethyl-3-(2,6-dichloro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (H-3; CAS 452098-92-9), N-tetrahydrofurfuryl-3-(2,6-dichloro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (H-4; CAS 915396-43-9), N-ethyl-3-(2-chloro-6-fluoro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (H-5; CAS 452099-05-7), N-tetrahydrofurfuryl-3-(2-chloro-6-fluoro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (H-6; CAS 45100-03-7), 3-[7-fluoro-3-oxo-4-(prop-2-ynyl)-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl]-1,5-dimethyl-6-thioxo-[1,3,5]triazinan-2,4-dione, 1,5-dimethyl-6-thioxo-3-(2,2,7-trifluoro-3-oxo-4-(prop-2-ynyl)-3,4-dihydro-2*H*-benzo[b][1,4]oxazin-6-yl)-1,3,5-triazinane-2,4-dione, 2-(2,2,7-Trifluoro-3-oxo-4-prop-2-ynyl-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl)-4,5,6,7-tetrahydro-isoindole-1,3-dione and 1-Methyl-6-trifluoromethyl-3-(2,2,7-trifluoro-3-oxo-4-prop-2-ynyl-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl)-1*H*-pyrimidine-2,4-dione;

b5) from the group of the bleacher herbicides:

acлонifen, amitrol, beflubutamid, benzobicyclon, benzofenap, clomazone, diflufenican, fluridone, flurochloridone, flurtamone, isoxaflutole, mesotrione, norflurazon, picolinafen, pyrasulfutole, pyrazolynate, pyrazoxyfen, sulcotrione, tefuryltrione, tembotrione, topramezone, 4-hydroxy-3-[[2-[(2-methoxyethoxy)methyl]-6-(trifluoromethyl)-3-pyridyl]carbonyl]bicyclo-
 5 [3.2.1]oct-3-en-2-one (H-7; CAS 352010-68-5) and 4-(3-trifluoromethylphenoxy)-2-(4-trifluoromethylphenyl)pyrimidine (H-8; CAS 180608-33-7);

b6) from the group of the EPSP synthase inhibitors:

glyphosate, glyphosate-isopropylammonium and glyphosate-trimesium (sulfosate);

b7) from the group of the glutamine synthase inhibitors:

10 bilanaphos (bialaphos), bilanaphos-sodium, glufosinate and glufosinate-ammonium;

b8) from the group of the DHP synthase inhibitors:

asulam;


b9) from the group of the mitose inhibitors:

15 amiprofos, amiprofos-methyl, benfluralin, butamiphos, butralin, carbetamide, chlorpropham, chlorthal, chlorthal-dimethyl, dinitramine, dithiopyr, ethalfluralin, fluchloralin, oryzalin, pendimethalin, prodiamine, propham, propyzamide, tebutam, thiazopyr and trifluralin;

b10) from the group of the VLCFA inhibitors:

20 acetochlor, alachlor, anilofos, butachlor, cafenstrole, dimethachlor, dimethanamid, dimethenamid-P, diphenamid, fentrazamide, flufenacet, mefenacet, metazachlor, metolachlor, metolachlor-S, naproanilide, napropamide, pethoxamid, piperophos, pretilachlor, propachlor, propisochlor, pyroxasulfone (KIH-485) and thenylchlor;

Compounds of the formula 2:




25

in which the variables have the following meanings:

Y is phenyl or 5- or 6-membered heteroaryl as defined at the outset, which radicals may be substituted by one to three groups R^{aa}; R²¹, R²², R²³, R²⁴ are H, halogen or C₁-C₄-alkyl; X is O or NH; N is 0 or 1.

30


Compounds of the formula 2 have in particular the following meanings:



where # denotes the bond to the skeleton of the molecule; and

35 R²¹, R²², R²³, R²⁴ are H, Cl, F or CH₃; R²⁵ is halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl; R²⁶ is C₁-C₄-alkyl; R²⁷ is halogen, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy; R²⁸ is H, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl or C₁-C₄-haloalkoxy; M is 0, 1, 2 or 3; X is oxygen; N is 0 or 1.

Preferred compounds of the formula 2 have the following meanings:



R²¹ is H; R²², R²³ are F; R²⁴ is H or F; X is oxygen; N is 0 or 1.

Particularly preferred compounds of the formula 2 are:

- 3-[5-(2,2-difluoroethoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl]methane-
 5 sulfonyl]-4-fluoro-5,5-dimethyl-4,5-dihydroisoxazole (2-1); 3-[[5-(2,2-difluoroethoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl]fluoromethanesulfonyl]-5,5-dimethyl-4,5-dihydroisoxazole (2-2);
 4-(4-fluoro-5,5-dimethyl-4,5-dihydroisoxazole-3-sulfonylmethyl)-2-methyl-5-trifluoromethyl-2H-[1,2,3]triazole (2-3); 4-[(5,5-dimethyl-4,5-dihydroisoxazole-3-sulfonyl)fluoromethyl]-2-methyl-5-trifluoromethyl-2H-[1,2,3]triazole (2-4); 4-(5,5-dimethyl-4,5-dihydroisoxazole-3-sulfonylmethyl)-2-methyl-5-trifluoromethyl-2H-[1,2,3]triazole (2-5); 3-[[5-(2,2-difluoroethoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl]difluoromethanesulfonyl]-5,5-dimethyl-4,5-dihydroisoxazole (2-6);
 4-[(5,5-dimethyl-4,5-dihydroisoxazole-3-sulfonyl)difluoromethyl]-2-methyl-5-trifluoromethyl-2H-[1,2,3]triazole (2-7); 3-[[5-(2,2-difluoroethoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl]difluoromethanesulfonyl]-4-fluoro-5,5-dimethyl-4,5-dihydroisoxazole (2-8); 4-[difluoro-(4-fluoro-5,5-dimethyl-4,5-dihydroisoxazole-3-sulfonyl)methyl]-2-methyl-5-trifluoromethyl-2H-[1,2,3]triazole (2-9);

b11) from the group of the cellulose biosynthesis inhibitors:

chlorthiamid, dichlobenil, flupoxam and isoxaben;

b12) from the group of the decoupler herbicides:

20 dinoseb, dinoterb and DNOC and its salts;

b13) from the group of the auxin herbicides:

2,4-D and its salts and esters, 2,4-DB and its salts and esters, aminopyralid and its salts such as aminopyralid-tris(2-hydroxypropyl)ammonium and its esters, benazolin, benazolin-ethyl, chloramben and its salts and esters, clomeprop, clopyralid and its salts and esters, dicamba and its salts and esters, dichlorprop and its salts and esters, dichlorprop-P and its salts and esters, fluroxypyr, fluroxypyr-butometyl, fluroxypyr-meptyl, MCPA and its salts and esters, MCPA-thioethyl, MCPB and its salts and esters, mecoprop and its salts and esters, mecoprop-P and its salts and esters, picloram and its salts and esters, quinclorac, quinmerac, TBA (2,3,6) and its salts and esters, triclopyr and its salts and esters, and 5,6-dichloro-2-cyclopropyl-4-pyrimidinecarboxylic acid (H-9; CAS 858956-08-8) and its salts and esters;

b14) from the group of the auxin transport inhibitors: diflufenzopyr, diflufenzopyr-sodium, naptalam and naptalam-sodium;

b15) from the group of the other herbicides: bromobutide, chlorflurenol, chlorflurenol-methyl, cinmethylin, cumyluron, dalapon, dazomet, difenzoquat, difenzoquat-metilsulfate, dimethipin, DSMA, dymron, endothal and its salts, etobenzanid, flamprop, flamprop-isopropyl, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flurenol, flurenol-butyl, flurprimidol, fosamine, fosamine-ammonium, indanofan, maleic hydrazide, mefluidide, metam, methyl azide, methyl bromide, methyl-dymron, methyl iodide, MSMA, oleic acid, oxaziclomefone, pelargonic

acid, pyributicarb, quinoclamine, triaziflam, tridiphane and 6-chloro-3-(2-cyclopropyl-6-methylphenoxy)-4-pyridazinol (H-10; CAS 499223-49-3) and its salts and esters.

Examples of preferred safeners C are benoxacor, cloquintocet, cyometrinil, cyprosulfamide, dichlormid, dicyclonone, dietholate, fenchlorazole, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen, mefenpyr, mephenate, naphthalic anhydride, oxabetrinil, 4-(dichloroacetyl)-1-oxa-4-azaspiro[4.5]decane (H-11; MON4660, CAS 71526-07-3) and 2,2,5-trimethyl-3-(dichloroacetyl)-1,3-oxazolidine (H-12; R-29148, CAS 52836-31-4).

The active compounds of groups b1) to b15) and the safeners C are known herbicides and safeners, see, for example, The Compendium of Pesticide Common Names (<http://www.alanwood.net/pesticides/>); B. Hock, C. Fedtke, R. R. Schmidt, Herbicide [Herbicides], Georg Thieme Verlag, Stuttgart, 1995. Further herbicidally active compounds are known from WO 96/26202, WO 97/41116, WO 97/41117, WO 97/41118, WO 01/83459 and WO 2008/074991 and from W. Krämer et al. (ed.) "Modern Crop Protection Compounds", Vol. 1, Wiley VCH, 2007 and the literature quoted therein.

The invention also relates to compositions in the form of a crop protection composition formulated as a 1-component composition comprising an active compound combination comprising at least one pyridine compound of the formula I and at least one further active compound, preferably selected from the active compounds of groups b1 to b15, and at least one solid or liquid carrier and/or one or more surfactants and, if desired, one or more further auxiliaries customary for crop protection compositions.

The invention also relates to compositions in the form of a crop protection composition formulated as a 2-component composition comprising a first component comprising at least one compound of the formula I, a solid or liquid carrier and/or one or more surfactants and a second component comprising at least one further active compound selected from the active compounds of groups b1 to b15, a solid or liquid carrier and/or one or more surfactants, where additionally both components may also comprise further auxiliaries customary for crop protection compositions.

In binary compositions comprising at least one compound of the formula I as component A and at least one herbicide B, the weight ratio of the active compounds A:B is generally in the range of from 1:1000 to 1000:1, preferably in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from 1:75 to 75:1.

In binary compositions comprising at least one compound of the formula I as component A and at least one safener C, the weight ratio of the active compounds A:C is generally in the range of from 1:1000 to 1000:1, preferably in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from 1:75 to 75:1.

In ternary compositions comprising both at least one compound of the formula I as component A, at least one herbicide B and at least one safener C, the relative parts by weight of the components A:B are generally in the range of from 1:1000 to 1000:1, preferably in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from 1:75 to 75:1; the weight ratio of the components A:C is generally in the range of from 1:1000 to 1000:1, preferably in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from

1:75 to 75:1; and the weight ratio of the components B:C is generally in the range of from 1:1000 to 1000:1, preferably in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from 1:75 to 75:1. Preferably, the weight ratio of the components A + B to the component C is in the range of from 1:500 to 500:1, in particular in the range of from 1:250 to 250:1 and particularly preferably in the range of from 1:75 to 75:1.

Examples of particularly preferred compositions according to the invention comprising in each case one individualized compound of the formula I and one mixing partner or a mixing partner combination are given in Table B below.

A further aspect of the invention relates to the compositions B-1 to B-1236 listed in Table B below, where in each case one row of Table B corresponds to a herbicidal composition comprising one of the compounds of the formula I individualized in the above description (component 1) and the further active compound from groups b1) to b15) and/or safener C stated in each case in the row in question (component 2). The active compounds in the compositions described are in each case preferably present in synergistically effective amounts.

Table B:

	Herbicide(s) B	Safener C
B-1	clodinafop-propargyl	--
B-2	cycloxydim	--
B-3	cyhalofop-butyl	--
B-4	fenoxaprop-P-ethyl	--
B-5	pinoxaden	--
B-6	profoxydim	--
B-7	tepraloxym	--
B-8	tralkoxydim	--
B-9	esprocarb	--
B-10	prosulfocarb	--
B-11	thiobencarb	--
B-12	triallate	--
B-13	bensulfuron-methyl	--
B-14	bispyribac-sodium	--
B-15	cyclosulfamuron	--
B-16	flumetsulam	--
B-17	flupyrsulfuron-methyl-sodium	--
B-18	foramsulfuron	--
B-19	imazamox	--
B-20	imazapic	--
B-21	imazapyr	--
B-22	imazaquin	--
B-23	imazethapyr	--

	Herbicide(s) B	Safener C
B-24	imazosulfuron	--
B-25	iodosulfuron-methyl-sodium	--
B-26	mesosulfuron	--
B-27	nicosulfuron	--
B-28	penoxsulam	--
B-29	propoxycarbazone-sodium	--
B-30	pyrazosulfuron-ethyl	--
B-31	pyroxsulam	--
B-32	rimsulfuron	--
B-33	sulfosulfuron	--
B-34	thiencarbazone-methyl	--
B-35	tritosulfuron	--
B-36	2,4-D and its salts and esters	--
B-37	aminopyralid and its salts and esters	--
B-38	clopyralid and its salts and esters	--
B-39	dicamba and its salts and esters	--
B-40	fluroxypyr-meptyl	--
B-41	quinclorac	--
B-42	quinmerac	--
B-43	H-9	--
B-44	diflufenzopyr	--
B-45	diflufenzopyr-sodium	--
B-46	clomazone	--
B-47	diflufenican	--
B-48	fluorochloridone	--
B-49	isoxaflutol	--
B-50	mesotrione	--
B-51	picolinafen	--
B-52	sulcotrione	--
B-53	tefuryltrione	--
B-54	tembotrione	--
B-55	topramezone	--
B-56	H-7	--
B-57	atrazine	--
B-58	diuron	--
B-59	fluometuron	--
B-60	hexazinone	--
B-61	isoproturon	--
B-62	metribuzin	--
B-63	propanil	--

	Herbicide(s) B	Safener C
B-64	terbuthylazine	--
B-65	paraquat dichloride	--
B-66	flumioxazin	--
B-67	oxyfluorfen	--
B-68	saflufenacil	--
B-69	sulfentrazone	--
B-70	H-1	--
B-71	H-2	--
B-72	glyphosate	--
B-73	glyphosate-isopropylammonium	--
B-74	glyphosate-trimesium (sulfosate)	--
B-75	glufosinate	--
B-76	glufosinate-ammonium	--
B-77	pendimethalin	--
B-78	trifluralin	--
B-79	acetochlor	--
B-80	cafenstrole	--
B-81	dimethenamid-P	--
B-82	fentrazamide	--
B-83	flufenacet	--
B-84	mefenacet	--
B-85	metazachlor	--
B-86	metolachlor-S	--
B-87	pyroxasulfone	--
B-88	isoxaben	--
B-89	dymron	--
B-90	indanofan	--
B-91	oxaziclomefone	--
B-92	triaziflam	--
B-93	chlorotoluron	--
B-94	atrazine + H-1	--
B-95	atrazine + glyphosate	--
B-96	atrazine + mesotrione	--
B-97	atrazine + nicosulfuron	--
B-98	atrazine + tembotrione	--
B-99	atrazine + topramezone	--
B-100	clomazone + glyphosate	--
B-101	diflufenican + clodinafop-propargyl	--
B-102	diflufenican + fenoxaprop-P-ethyl	--
B-103	diflufenican + flupyrsulfuron-methyl-sodium	--

	Herbicide(s) B	Safener C
B-104	diflufenican + glyphosate	--
B-105	diflufenican + mesosulfuron-methyl	--
B-106	diflufenican + pinoxaden	--
B-107	diflufenican + pyroxsulam	--
B-108	flumetsulam + glyphosate	--
B-109	flumioxazin + glyphosate	--
B-110	imazapic + glyphosate	--
B-111	imazethapyr + glyphosate	--
B-112	isoxaflutol + H-1	--
B-113	isoxaflutol + glyphosate	--
B-114	metazachlor + H-1	--
B-115	metazachlor + glyphosate	--
B-116	metazachlor + mesotrione	--
B-117	metazachlor + nicosulfuron	--
B-118	metazachlor + terbuthylazine	--
B-119	metazachlor + topramezone	--
B-120	metribuzin + glyphosate	--
B-121	pendimethalin + H-1	--
B-122	pendimethalin + clodinafop-propargyl	--
B-123	pendimethalin + fenoxaprop-P-ethyl	--
B-124	pendimethalin + flupyrsulfuron-methyl-sodium	--
B-125	pendimethalin + glyphosate	--
B-126	pendimethalin + mesosulfuron-methyl	--
B-127	pendimethalin + mesotrione	--
B-128	pendimethalin + nicosulfuron	--
B-129	pendimethalin + pinoxaden	--
B-130	pendimethalin + pyroxsulam	--
B-131	pendimethalin + tembotrione	--
B-132	pendimethalin + topramezone	--
B-133	pyroxasulfone + tembotrione	--
B-134	pyroxasulfone + topramezone	--
B-135	sulfentrazone + glyphosate	--
B-136	terbuthylazine + H-1	--
B-137	terbuthylazine + foramsulfuron	--
B-138	terbuthylazine + glyphosate	--
B-139	terbuthylazine + mesotrione	--
B-140	terbuthylazine + nicosulfuron	--
B-141	terbuthylazine + tembotrione	--
B-142	terbuthylazine + topramezone	--
B-143	trifluralin + glyphosate	--

	Herbicide(s) B	Safener C
B-144	--	benoxacor
B-145	--	cloquintocet
B-146	--	cyprosulfamide
B-147	--	dichlormid
B-148	--	fenchlorazole
B-149	--	isoxadifen
B-150	--	mefenpyr
B-151	--	H-11
B-152	--	H-12
B-153	clodinafop-propargyl	benoxacor
B-154	cycloxydim	benoxacor
B-155	cyhalofop-butyl	benoxacor
B-156	fenoxaprop-P-ethyl	benoxacor
B-157	pinoxaden	benoxacor
B-158	profoxydim	benoxacor
B-159	tepraloxydim	benoxacor
B-160	tralkoxydim	benoxacor
B-161	esprocarb	benoxacor
B-162	prosulfocarb	benoxacor
B-163	thiobencarb	benoxacor
B-164	triallate	benoxacor
B-165	bensulfuron-methyl	benoxacor
B-166	bispyribac-sodium	benoxacor
B-167	cyclosulfamuron	benoxacor
B-168	flumetsulam	benoxacor
B-169	flupyrsulfuron-methyl-sodium	benoxacor
B-170	foramsulfuron	benoxacor
B-171	imazamox	benoxacor
B-172	imazapic	benoxacor
B-173	imazapyr	benoxacor
B-174	imazaquin	benoxacor
B-175	imazethapyr	benoxacor
B-176	imazosulfuron	benoxacor
B-177	iodosulfuron-methyl-sodium	benoxacor
B-178	mesosulfuron	benoxacor
B-179	nicosulfuron	benoxacor
B-180	penoxsulam	benoxacor
B-181	propoxycarbazone-sodium	benoxacor
B-182	pyrazosulfuron-ethyl	benoxacor
B-183	pyroxsulam	benoxacor

	Herbicide(s) B	Safener C
B-184	rimsulfuron	benoxacor
B-185	sulfosulfuron	benoxacor
B-186	thiencarbazone-methyl	benoxacor
B-187	tritosulfuron	benoxacor
B-188	2,4-D and its salts and esters	benoxacor
B-189	aminopyralid and its salts and esters	benoxacor
B-190	clopyralid and its salts and esters	benoxacor
B-191	dicamba and its salts and esters	benoxacor
B-192	fluroxypyr-meptyl	benoxacor
B-193	quinclorac	benoxacor
B-194	quinmerac	benoxacor
B-195	H-9	benoxacor
B-196	diflufenzopyr	benoxacor
B-197	diflufenzopyr-sodium	benoxacor
B-198	clomazone	benoxacor
B-199	diflufenican	benoxacor
B-200	fluorochloridone	benoxacor
B-201	isoxaflutol	benoxacor
B-202	mesotrione	benoxacor
B-203	picolinafen	benoxacor
B-204	sulcotrione	benoxacor
B-205	tefuryltrione	benoxacor
B-206	tembotrione	benoxacor
B-207	topramezone	benoxacor
B-208	H-7	benoxacor
B-209	atrazine	benoxacor
B-210	diuron	benoxacor
B-211	fluometuron	benoxacor
B-212	hexazinone	benoxacor
B-213	isoproturon	benoxacor
B-214	metribuzin	benoxacor
B-215	propanil	benoxacor
B-216	terbuthylazine	benoxacor
B-217	paraquat dichloride	benoxacor
B-218	flumioxazin	benoxacor
B-219	oxyfluorfen	benoxacor
B-220	saflufenacil	benoxacor
B-221	sulfentrazone	benoxacor
B-222	H-1	benoxacor
B-223	H-2	benoxacor

	Herbicide(s) B	Safener C
B-224	glyphosate	benoxacor
B-225	glyphosate-isopropylammonium	benoxacor
B-226	glyphosate-trimesium (sulfosate)	benoxacor
B-227	glufosinate	benoxacor
B-228	glufosinate-ammonium	benoxacor
B-229	pendimethalin	benoxacor
B-230	trifluralin	benoxacor
B-231	acetochlor	benoxacor
B-232	cafenstrole	benoxacor
B-233	dimethenamid-P	benoxacor
B-234	fentrazamide	benoxacor
B-235	flufenacet	benoxacor
B-236	mefenacet	benoxacor
B-237	metazachlor	benoxacor
B-238	metolachlor-S	benoxacor
B-239	pyroxasulfone	benoxacor
B-240	isoxaben	benoxacor
B-241	dymron	benoxacor
B-242	indanofan	benoxacor
B-243	oxaziclomefone	benoxacor
B-244	triaziflam	benoxacor
B-245	atrazine + H-1	benoxacor
B-246	atrazine + glyphosate	benoxacor
B-247	atrazine + mesotrione	benoxacor
B-248	atrazine + nicosulfuron	benoxacor
B-249	atrazine + tembotrione	benoxacor
B-250	atrazine + topramezone	benoxacor
B-251	clomazone + glyphosate	benoxacor
B-252	diflufenican + clodinafop-propargyl	benoxacor
B-253	diflufenican + fenoxaprop-P-ethyl	benoxacor
B-254	diflufenican + flupyrsulfuron-methyl-sodium	benoxacor
B-255	diflufenican + glyphosate	benoxacor
B-256	diflufenican + mesosulfuron-methyl	benoxacor
B-257	diflufenican + pinoxaden	benoxacor
B-258	diflufenican + pyroxsulam	benoxacor
B-259	flumetsulam + glyphosate	benoxacor
B-260	flumioxazin + glyphosate	benoxacor
B-261	imazapic + glyphosate	benoxacor
B-262	imazethapyr + glyphosate	benoxacor
B-263	isoxaflutol + H-1	benoxacor

	Herbicide(s) B	Safener C
B-264	isoxaflutol + glyphosate	benoxacor
B-265	metazachlor + H-1	benoxacor
B-266	metazachlor + glyphosate	benoxacor
B-267	metazachlor + mesotrione	benoxacor
B-268	metazachlor + nicosulfuron	benoxacor
B-269	metazachlor + terbuthylazine	benoxacor
B-270	metazachlor + topramezone	benoxacor
B-271	metribuzin + glyphosate	benoxacor
B-272	pendimethalin + H-1	benoxacor
B-273	pendimethalin + clodinafop-propargyl	benoxacor
B-274	pendimethalin + fenoxaprop-P-ethyl	benoxacor
B-275	pendimethalin + flupyrsulfuron-methyl-sodium	benoxacor
B-276	pendimethalin + glyphosate	benoxacor
B-277	pendimethalin + mesosulfuron-methyl	benoxacor
B-278	pendimethalin + mesotrione	benoxacor
B-279	pendimethalin + nicosulfuron	benoxacor
B-280	pendimethalin + pinoxaden	benoxacor
B-281	pendimethalin + pyroxsulam	benoxacor
B-282	pendimethalin + tembotrione	benoxacor
B-283	pendimethalin + topramezone	benoxacor
B-284	pyroxasulfone + tembotrione	benoxacor
B-285	pyroxasulfone + topramezone	benoxacor
B-286	sulfentrazone + glyphosate	benoxacor
B-287	terbuthylazine + H-1	benoxacor
B-288	terbuthylazine + foramsulfuron	benoxacor
B-289	terbuthylazine + glyphosate	benoxacor
B-290	terbuthylazine + mesotrione	benoxacor
B-291	terbuthylazine + nicosulfuron	benoxacor
B-292	terbuthylazine + tembotrione	benoxacor
B-293	terbuthylazine + topramezone	benoxacor
B-294	trifluralin + glyphosate	benoxacor
B-295	clodinafop-propargyl	cloquintocet
B-296	cycloxydim	cloquintocet
B-297	cyhalofop-butyl	cloquintocet
B-298	fenoxaprop-P-ethyl	cloquintocet
B-299	pinoxaden	cloquintocet
B-300	profoxydim	cloquintocet
B-301	tepraloxym	cloquintocet
B-302	tralkoxydim	cloquintocet
B-303	esprocarb	cloquintocet

	Herbicide(s) B	Safener C
B-304	prosulfocarb	cloquintocet
B-305	thiobencarb	cloquintocet
B-306	triallate	cloquintocet
B-307	bensulfuron-methyl	cloquintocet
B-308	bispyribac-sodium	cloquintocet
B-309	cyclosulfamuron	cloquintocet
B-310	flumetsulam	cloquintocet
B-311	flupyrsulfuron-methyl-sodium	cloquintocet
B-312	foramsulfuron	cloquintocet
B-313	imazamox	cloquintocet
B-314	imazapic	cloquintocet
B-315	imazapyr	cloquintocet
B-316	imazaquin	cloquintocet
B-317	imazethapyr	cloquintocet
B-318	imazosulfuron	cloquintocet
B-319	iodosulfuron-methyl-sodium	cloquintocet
B-320	mesosulfuron	cloquintocet
B-321	nicosulfuron	cloquintocet
B-322	penoxsulam	cloquintocet
B-323	propoxycarbazone-sodium	cloquintocet
B-324	pyrazosulfuron-ethyl	cloquintocet
B-325	pyroxsulam	cloquintocet
B-326	rimsulfuron	cloquintocet
B-327	sulfosulfuron	cloquintocet
B-328	thiencarbazone-methyl	cloquintocet
B-329	tritosulfuron	cloquintocet
B-330	2,4-D and its salts and esters	cloquintocet
B-331	aminopyralid and its salts and esters	cloquintocet
B-332	clopyralid and its salts and esters	cloquintocet
B-333	dicamba and its salts and esters	cloquintocet
B-334	fluroxypyr-meptyl	cloquintocet
B-335	quinclorac	cloquintocet
B-336	quinmerac	cloquintocet
B-337	H-9	cloquintocet
B-338	diflufenzopyr	cloquintocet
B-339	diflufenzopyr-sodium	cloquintocet
B-340	clomazone	cloquintocet
B-341	diflufenican	cloquintocet
B-342	fluorochloridone	cloquintocet
B-343	isoxaflutol	cloquintocet

	Herbicide(s) B	Safener C
B-344	mesotrione	cloquintocet
B-345	picolinafen	cloquintocet
B-346	sulcotrione	cloquintocet
B-347	tefuryltrione	cloquintocet
B-348	tembotrione	cloquintocet
B-349	topramezone	cloquintocet
B-350	H-7	cloquintocet
B-351	atrazine	cloquintocet
B-352	diuron	cloquintocet
B-353	fluometuron	cloquintocet
B-354	hexazinone	cloquintocet
B-355	isoproturon	cloquintocet
B-356	metribuzin	cloquintocet
B-357	propanil	cloquintocet
B-358	terbuthylazine	cloquintocet
B-359	paraquat dichloride	cloquintocet
B-360	flumioxazin	cloquintocet
B-361	oxyfluorfen	cloquintocet
B-362	saflufenacil	cloquintocet
B-363	sulfentrazone	cloquintocet
B-364	H-1	cloquintocet
B-365	H-2	cloquintocet
B-366	glyphosate	cloquintocet
B-367	glyphosate-isopropylammonium	cloquintocet
B-368	glyphosate-trimesium (sulfosate)	cloquintocet
B-369	glufosinate	cloquintocet
B-370	glufosinate-ammonium	cloquintocet
B-371	pendimethalin	cloquintocet
B-372	trifluralin	cloquintocet
B-373	acetochlor	cloquintocet
B-374	cafenstrole	cloquintocet
B-375	dimethenamid-P	cloquintocet
B-376	fentrazamide	cloquintocet
B-377	flufenacet	cloquintocet
B-378	mefenacet	cloquintocet
B-379	metazachlor	cloquintocet
B-380	metolachlor-S	cloquintocet
B-381	pyroxasulfone	cloquintocet
B-382	isoxaben	cloquintocet
B-383	dymron	cloquintocet

	Herbicide(s) B	Safener C
B-384	indanofan	cloquintocet
B-385	oxaziclomefone	cloquintocet
B-386	triaziflam	cloquintocet
B-387	atrazine + H-1	cloquintocet
B-388	atrazine + glyphosate	cloquintocet
B-389	atrazine + mesotrione	cloquintocet
B-390	atrazine + nicosulfuron	cloquintocet
B-391	atrazine + tembotrione	cloquintocet
B-392	atrazine + topramezone	cloquintocet
B-393	clomazone + glyphosate	cloquintocet
B-394	diflufenican + clodinafop-propargyl	cloquintocet
B-395	diflufenican + fenoxaprop-p-ethyl	cloquintocet
B-396	diflufenican + flupyrsulfuron-methyl-sodium	cloquintocet
B-397	diflufenican + glyphosate	cloquintocet
B-398	diflufenican + mesosulfuron-methyl	cloquintocet
B-399	diflufenican + pinoxaden	cloquintocet
B-400	diflufenican + pyroxsulam	cloquintocet
B-401	flumetsulam + glyphosate	cloquintocet
B-402	flumioxazin + glyphosate	cloquintocet
B-403	imazapic + glyphosate	cloquintocet
B-404	imazethapyr + glyphosate	cloquintocet
B-405	isoxaflutol + H-1	cloquintocet
B-406	isoxaflutol + glyphosate	cloquintocet
B-407	metazachlor + H-1	cloquintocet
B-408	metazachlor + glyphosate	cloquintocet
B-409	metazachlor + mesotrione	cloquintocet
B-410	metazachlor + nicosulfuron	cloquintocet
B-411	metazachlor + terbutylazine	cloquintocet
B-412	metazachlor + topramezone	cloquintocet
B-413	metribuzin + glyphosate	cloquintocet
B-414	pendimethalin + H-1	cloquintocet
B-415	pendimethalin + clodinafop-propargyl	cloquintocet
B-416	pendimethalin + fenoxaprop-P-ethyl	cloquintocet
B-417	pendimethalin + flupyrsulfuron-methyl-sodium	cloquintocet
B-418	pendimethalin + glyphosate	cloquintocet
B-419	pendimethalin + mesosulfuron-methyl	cloquintocet
B-420	pendimethalin + mesotrione	cloquintocet
B-421	pendimethalin + nicosulfuron	cloquintocet
B-422	pendimethalin + pinoxaden	cloquintocet
B-423	pendimethalin + pyroxsulam	cloquintocet

	Herbicide(s) B	Safener C
B-424	pendimethalin + tembotrione	cloquintocet
B-425	pendimethalin + topramezone	cloquintocet
B-426	pyroxasulfone + tembotrione	cloquintocet
B-427	pyroxasulfone + topramezone	cloquintocet
B-428	sulfentrazone + glyphosate	cloquintocet
B-429	terbuthylazine + H-1	cloquintocet
B-430	terbuthylazine + foramsulfuron	cloquintocet
B-431	terbuthylazine + glyphosate	cloquintocet
B-432	terbuthylazine + mesotrione	cloquintocet
B-433	terbuthylazine + nicosulfuron	cloquintocet
B-434	terbuthylazine + tembotrione	cloquintocet
B-435	terbuthylazine + topramezone	cloquintocet
B-436	trifluralin + glyphosate	cloquintocet
B-437	clodinafop-propargyl	dichlormid
B-438	cycloxydim	dichlormid
B-439	cyhalofop-butyl	dichlormid
B-440	fenoxaprop-P-ethyl	dichlormid
B-441	pinoxaden	dichlormid
B-442	profoxydim	dichlormid
B-443	tepraloxydim	dichlormid
B-444	tralkoxydim	dichlormid
B-445	esprocarb	dichlormid
B-446	prosulfocarb	dichlormid
B-447	thiobencarb	dichlormid
B-448	triallate	dichlormid
B-449	bensulfuron-methyl	dichlormid
B-450	bispyribac-sodium	dichlormid
B-451	cyclosulfamuron	dichlormid
B-452	flumetsulam	dichlormid
B-453	flupyrsulfuron-methyl-sodium	dichlormid
B-454	foramsulfuron	dichlormid
B-455	imazamox	dichlormid
B-456	imazapic	dichlormid
B-457	imazapyr	dichlormid
B-458	imazaquin	dichlormid
B-459	imazethapyr	dichlormid
B-460	imazosulfuron	dichlormid
B-461	iodosulfuron-methyl-sodium	dichlormid
B-462	mesosulfuron	dichlormid
B-463	nicosulfuron	dichlormid

	Herbicide(s) B	Safener C
B-464	penoxsulam	dichlormid
B-465	propoxycarbazone-sodium	dichlormid
B-466	pyrazosulfuron-ethyl	dichlormid
B-467	pyroxsulam	dichlormid
B-468	rimisulfuron	dichlormid
B-469	sulfosulfuron	dichlormid
B-470	thiencarbazone-methyl	dichlormid
B-471	tritosulfuron	dichlormid
B-472	2,4-D and its salts and esters	dichlormid
B-473	aminopyralid and its salts and esters	dichlormid
B-474	clopyralid and its salts and esters	dichlormid
B-475	dicamba and its salts and esters	dichlormid
B-476	fluroxypyr-meptyl	dichlormid
B-477	quinclorac	dichlormid
B-478	quinmerac	dichlormid
B-479	H-9	dichlormid
B-480	diflufenzopyr	dichlormid
B-481	diflufenzopyr-sodium	dichlormid
B-482	clomazone	dichlormid
B-483	diflufenican	dichlormid
B-484	fluorochloridone	dichlormid
B-485	isoxaflutol	dichlormid
B-486	mesotrione	dichlormid
B-487	picolinafen	dichlormid
B-488	sulcotrione	dichlormid
B-489	tefuryltrione	dichlormid
B-490	tembotrione	dichlormid
B-491	topramezone	dichlormid
B-492	H-7	dichlormid
B-493	atrazine	dichlormid
B-494	diuron	dichlormid
B-495	fluometuron	dichlormid
B-496	hexazinone	dichlormid
B-497	isoproturon	dichlormid
B-498	metribuzin	dichlormid
B-499	propanil	dichlormid
B-500	terbuthylazine	dichlormid
B-501	paraquat dichloride	dichlormid
B-502	flumioxazin	dichlormid
B-503	oxyfluorfen	dichlormid

	Herbicide(s) B	Safener C
B-504	saflufenacil	dichlormid
B-505	sulfentrazone	dichlormid
B-506	H-1	dichlormid
B-507	H-2	dichlormid
B-508	glyphosate	dichlormid
B-509	glyphosate-isopropylammonium	dichlormid
B-510	glyphosate-trimesium (sulfosate)	dichlormid
B-511	glufosinate	dichlormid
B-512	glufosinate-ammonium	dichlormid
B-513	pendimethalin	dichlormid
B-514	trifluralin	dichlormid
B-515	acetochlor	dichlormid
B-516	cafenstrole	dichlormid
B-517	dimethenamid-P	dichlormid
B-518	fentrazamide	dichlormid
B-519	flufenacet	dichlormid
B-520	mefenacet	dichlormid
B-521	metazachlor	dichlormid
B-522	metolachlor-S	dichlormid
B-523	pyroxasulfone	dichlormid
B-524	isoxaben	dichlormid
B-525	dymron	dichlormid
B-526	indanofan	dichlormid
B-527	oxaziclomefone	dichlormid
B-528	triaziflam	dichlormid
B-529	atrazine + H-1	dichlormid
B-530	atrazine + glyphosate	dichlormid
B-531	atrazine + mesotrione	dichlormid
B-532	atrazine + nicosulfuron	dichlormid
B-533	atrazine + tembotrione	dichlormid
B-534	atrazine + topramezone	dichlormid
B-535	clomazone + glyphosate	dichlormid
B-536	diflufenican + clodinafop-propargyl	dichlormid
B-537	diflufenican + fenoxaprop-p-ethyl	dichlormid
B-538	diflufenican + flupyrsulfuron-methyl-sodium	dichlormid
B-539	diflufenican + glyphosate	dichlormid
B-540	diflufenican + mesosulfuron-methyl	dichlormid
B-541	diflufenican + pinoxaden	dichlormid
B-542	diflufenican + pyroxsulam	dichlormid
B-543	flumetsulam + glyphosate	dichlormid

	Herbicide(s) B	Safener C
B-544	flumioxazin + glyphosate	dichlormid
B-545	imazapic + glyphosate	dichlormid
B-546	imazethapyr + glyphosate	dichlormid
B-547	isoxaflutol + H-1	dichlormid
B-548	isoxaflutol + glyphosate	dichlormid
B-549	metazachlor + H-1	dichlormid
B-550	metazachlor + glyphosate	dichlormid
B-551	metazachlor + mesotrione	dichlormid
B-552	metazachlor + nicosulfuron	dichlormid
B-553	metazachlor + terbuthylazine	dichlormid
B-554	metazachlor + topramezone	dichlormid
B-555	metribuzin + glyphosate	dichlormid
B-556	pendimethalin + H-1	dichlormid
B-557	pendimethalin + clodinafop-propargyl	dichlormid
B-558	pendimethalin + fenoxaprop-P-ethyl	dichlormid
B-559	pendimethalin + flupyrsulfuron-methyl-sodium	dichlormid
B-560	pendimethalin + glyphosate	dichlormid
B-561	pendimethalin + mesosulfuron-methyl	dichlormid
B-562	pendimethalin + mesotrione	dichlormid
B-563	pendimethalin + nicosulfuron	dichlormid
B-564	pendimethalin + pinoxaden	dichlormid
B-565	pendimethalin + pyroxsulam	dichlormid
B-566	pendimethalin + tembotrione	dichlormid
B-567	pendimethalin + topramezone	dichlormid
B-568	pyroxasulfone + tembotrione	dichlormid
B-569	pyroxasulfone + topramezone	dichlormid
B-570	sulfentrazone + glyphosate	dichlormid
B-571	terbuthylazine + H-1	dichlormid
B-572	terbuthylazine + foramsulfuron	dichlormid
B-573	terbuthylazine + glyphosate	dichlormid
B-574	terbuthylazine + mesotrione	dichlormid
B-575	terbuthylazine + nicosulfuron	dichlormid
B-576	terbuthylazine + tembotrione	dichlormid
B-577	terbuthylazine + topramezone	dichlormid
B-578	trifluralin + glyphosate	dichlormid
B-579	clodinafop-propargyl	fenchlorazole
B-580	cycloxydim	fenchlorazole
B-581	cyhalofop-butyl	fenchlorazole
B-582	fenoxaprop-P-ethyl	fenchlorazole
B-583	pinoxaden	fenchlorazole

	Herbicide(s) B	Safener C
B-584	profoxydim	fenchlorazole
B-585	tepraloxydim	fenchlorazole
B-586	tralkoxydim	fenchlorazole
B-587	esprocarb	fenchlorazole
B-588	prosulfocarb	fenchlorazole
B-589	thiobencarb	fenchlorazole
B-590	triallate	fenchlorazole
B-591	bensulfuron-methyl	fenchlorazole
B-592	bispyribac-sodium	fenchlorazole
B-593	cyclosulfamuron	fenchlorazole
B-594	flumetsulam	fenchlorazole
B-595	flupyrsulfuron-methyl-sodium	fenchlorazole
B-596	foramsulfuron	fenchlorazole
B-597	imazamox	fenchlorazole
B-598	imazapic	fenchlorazole
B-599	imazapyr	fenchlorazole
B-600	imazaquin	fenchlorazole
B-601	imazethapyr	fenchlorazole
B-602	imazosulfuron	fenchlorazole
B-603	iodosulfuron-methyl-sodium	fenchlorazole
B-604	mesosulfuron	fenchlorazole
B-605	nicosulfuron	fenchlorazole
B-606	penoxsulam	fenchlorazole
B-607	propoxycarbazone-sodium	fenchlorazole
B-608	pyrazosulfuron-ethyl	fenchlorazole
B-609	pyroxsulam	fenchlorazole
B-610	rimsulfuron	fenchlorazole
B-611	sulfosulfuron	fenchlorazole
B-612	thiencarbazone-methyl	fenchlorazole
B-613	tritosulfuron	fenchlorazole
B-614	2,4-D and its salts and esters	fenchlorazole
B-615	aminopyralid and its salts and esters	fenchlorazole
B-616	clopyralid and its salts and esters	fenchlorazole
B-617	dicamba and its salts and esters	fenchlorazole
B-618	fluroxypyr-meptyl	fenchlorazole
B-619	quinclorac	fenchlorazole
B-620	quinmerac	fenchlorazole
B-621	H-9	fenchlorazole
B-622	diflufenzopyr	fenchlorazole
B-623	diflufenzopyr-sodium	fenchlorazole

	Herbicide(s) B	Safener C
B-624	clomazone	fenchlorazole
B-625	diflufenican	fenchlorazole
B-626	fluorochloridone	fenchlorazole
B-627	isoxaflutol	fenchlorazole
B-628	mesotrione	fenchlorazole
B-629	picolinafen	fenchlorazole
B-630	sulcotrione	fenchlorazole
B-631	tefuryltrione	fenchlorazole
B-632	tembotrione	fenchlorazole
B-633	topramezone	fenchlorazole
B-634	H-7	fenchlorazole
B-635	atrazine	fenchlorazole
B-636	diuron	fenchlorazole
B-637	fluometuron	fenchlorazole
B-638	hexazinone	fenchlorazole
B-639	isoproturon	fenchlorazole
B-640	metribuzin	fenchlorazole
B-641	propanil	fenchlorazole
B-642	terbuthylazine	fenchlorazole
B-643	paraquat dichloride	fenchlorazole
B-644	flumioxazin	fenchlorazole
B-645	oxyfluorfen	fenchlorazole
B-646	saflufenacil	fenchlorazole
B-647	sulfentrazone	fenchlorazole
B-648	H-1	fenchlorazole
B-649	H-2	fenchlorazole
B-650	glyphosate	fenchlorazole
B-651	glyphosate-isopropylammonium	fenchlorazole
B-652	glyphosate-trimesium (sulfosate)	fenchlorazole
B-653	glufosinate	fenchlorazole
B-654	glufosinate-ammonium	fenchlorazole
B-655	pendimethalin	fenchlorazole
B-656	trifluralin	fenchlorazole
B-657	acetochlor	fenchlorazole
B-658	cafenstrole	fenchlorazole
B-659	dimethenamid-P	fenchlorazole
B-660	fentrazamide	fenchlorazole
B-661	flufenacet	fenchlorazole
B-662	mefenacet	fenchlorazole
B-663	metazachlor	fenchlorazole

	Herbicide(s) B	Safener C
B-664	metolachlor-S	fenchlorazole
B-665	pyroxasulfone	fenchlorazole
B-666	isoxaben	fenchlorazole
B-667	dymron	fenchlorazole
B-668	indanofan	fenchlorazole
B-669	oxaziclomefone	fenchlorazole
B-670	triaziflam	fenchlorazole
B-671	atrazine + H-1	fenchlorazole
B-672	atrazine + glyphosate	fenchlorazole
B-673	atrazine + mesotrione	fenchlorazole
B-674	atrazine + nicosulfuron	fenchlorazole
B-675	atrazine + tembotrione	fenchlorazole
B-676	atrazine + topramezone	fenchlorazole
B-677	clomazone + glyphosate	fenchlorazole
B-678	diflufenican + clodinafop-propargyl	fenchlorazole
B-679	diflufenican + fenoxaprop-P-ethyl	fenchlorazole
B-680	diflufenican + flupyrsulfuron-methyl-sodium	fenchlorazole
B-681	diflufenican + glyphosate	fenchlorazole
B-682	diflufenican + mesosulfuron-methyl	fenchlorazole
B-683	diflufenican + pinoxaden	fenchlorazole
B-684	diflufenican + pyroxsulam	fenchlorazole
B-685	flumetsulam + glyphosate	fenchlorazole
B-686	flumioxazin + glyphosate	fenchlorazole
B-687	imazapic + glyphosate	fenchlorazole
B-688	imazethapyr + glyphosate	fenchlorazole
B-689	isoxaflutol + H-1	fenchlorazole
B-690	isoxaflutol + glyphosate	fenchlorazole
B-691	metazachlor + H-1	fenchlorazole
B-692	metazachlor + glyphosate	fenchlorazole
B-693	metazachlor + mesotrione	fenchlorazole
B-694	metazachlor + nicosulfuron	fenchlorazole
B-695	metazachlor + terbuthylazine	fenchlorazole
B-696	metazachlor + topramezone	fenchlorazole
B-697	metribuzin + glyphosate	fenchlorazole
B-698	pendimethalin + H-1	fenchlorazole
B-699	pendimethalin + clodinafop-propargyl	fenchlorazole
B-700	pendimethalin + fenoxaprop-P-ethyl	fenchlorazole
B-701	pendimethalin + flupyrsulfuron-methyl-sodium	fenchlorazole
B-702	pendimethalin + glyphosate	fenchlorazole
B-703	pendimethalin + mesosulfuron-methyl	fenchlorazole

	Herbicide(s) B	Safener C
B-704	pendimethalin + mesotrione	fenchlorazole
B-705	pendimethalin + nicosulfuron	fenchlorazole
B-706	pendimethalin + pinoxaden	fenchlorazole
B-707	pendimethalin + pyroxsulam	fenchlorazole
B-708	pendimethalin + tembotrione	fenchlorazole
B-709	pendimethalin + topramezone	fenchlorazole
B-710	pyroxasulfone + tembotrione	fenchlorazole
B-711	pyroxasulfone + topramezone	fenchlorazole
B-712	sulfentrazone + glyphosate	fenchlorazole
B-713	terbuthylazine + H-1	fenchlorazole
B-714	terbuthylazine + foramsulfuron	fenchlorazole
B-715	terbuthylazine + glyphosate	fenchlorazole
B-716	terbuthylazine + mesotrione	fenchlorazole
B-717	terbuthylazine + nicosulfuron	fenchlorazole
B-718	terbuthylazine + tembotrione	fenchlorazole
B-719	terbuthylazine + topramezone	fenchlorazole
B-720	trifluralin + glyphosate	fenchlorazole
B-721	clodinafop-propargyl	isoxadifen
B-722	cycloxydim	isoxadifen
B-723	cyhalofop-butyl	isoxadifen
B-724	fenoxaprop-P-ethyl	isoxadifen
B-725	pinoxaden	isoxadifen
B-726	profoxydim	isoxadifen
B-727	tepraloxydim	isoxadifen
B-728	tralkoxydim	isoxadifen
B-729	esprocarb	isoxadifen
B-730	prosulfocarb	isoxadifen
B-731	thiobencarb	isoxadifen
B-732	triallate	isoxadifen
B-733	bensulfuron-methyl	isoxadifen
B-734	bispyribac-sodium	isoxadifen
B-735	cyclosulfamuron	isoxadifen
B-736	flumetsulam	isoxadifen
B-737	flupyrsulfuron-methyl-sodium	isoxadifen
B-738	foramsulfuron	isoxadifen
B-739	imazamox	isoxadifen
B-740	imazapic	isoxadifen
B-741	imazapyr	isoxadifen
B-742	imazaquin	isoxadifen
B-743	imazethapyr	isoxadifen

	Herbicide(s) B	Safener C
B-744	imazosulfuron	isoxadifen
B-745	iodosulfuron-methyl-sodium	isoxadifen
B-746	mesosulfuron	isoxadifen
B-747	nicosulfuron	isoxadifen
B-748	penoxsulam	isoxadifen
B-749	propoxycarbazone-sodium	isoxadifen
B-750	pyrazosulfuron-ethyl	isoxadifen
B-751	pyroxsulam	isoxadifen
B-752	rimsulfuron	isoxadifen
B-753	sulfosulfuron	isoxadifen
B-754	thiencarbazone-methyl	isoxadifen
B-755	tritosulfuron	isoxadifen
B-756	2,4-D and its salts and esters	isoxadifen
B-757	aminopyralid and its salts and esters	isoxadifen
B-758	clopyralid and its salts and esters	isoxadifen
B-759	dicamba and its salts and esters	isoxadifen
B-760	fluroxypyr-meptyl	isoxadifen
B-761	quinclorac	isoxadifen
B-762	quinmerac	isoxadifen
B-763	H-9	isoxadifen
B-764	diflufenzopyr	isoxadifen
B-765	diflufenzopyr-sodium	isoxadifen
B-766	clomazone	isoxadifen
B-767	diflufenican	isoxadifen
B-768	fluorochloridone	isoxadifen
B-769	isoxaflutol	isoxadifen
B-770	mesotrione	isoxadifen
B-771	picolinafen	isoxadifen
B-772	sulcotrione	isoxadifen
B-773	tefuryltrione	isoxadifen
B-774	tembotrione	isoxadifen
B-775	topramezone	isoxadifen
B-776	H-7	isoxadifen
B-777	atrazine	isoxadifen
B-778	diuron	isoxadifen
B-779	fluometuron	isoxadifen
B-780	hexazinone	isoxadifen
B-781	isoproturon	isoxadifen
B-782	metribuzin	isoxadifen
B-783	propanil	isoxadifen

	Herbicide(s) B	Safener C
B-784	terbuthylazine	isoxadifen
B-785	paraquat dichloride	isoxadifen
B-786	flumioxazin	isoxadifen
B-787	oxyfluorfen	isoxadifen
B-788	saflufenacil	isoxadifen
B-789	sulfentrazone	isoxadifen
B-790	H-1	isoxadifen
B-791	H-2	isoxadifen
B-792	glyphosate	isoxadifen
B-793	glyphosate-isopropylammonium	isoxadifen
B-794	glyphosate-trimesium (sulfosate)	isoxadifen
B-795	glufosinate	isoxadifen
B-796	glufosinate-ammonium	isoxadifen
B-797	pendimethalin	isoxadifen
B-798	trifluralin	isoxadifen
B-799	acetochlor	isoxadifen
B-800	cafenstrole	isoxadifen
B-801	dimethenamid-P	isoxadifen
B-802	fentrazamide	isoxadifen
B-803	flufenacet	isoxadifen
B-804	mefenacet	isoxadifen
B-805	metazachlor	isoxadifen
B-806	metolachlor-S	isoxadifen
B-807	pyroxasulfone	isoxadifen
B-808	isoxaben	isoxadifen
B-809	dymron	isoxadifen
B-810	indanofan	isoxadifen
B-811	oxaziclomefone	isoxadifen
B-812	triaziflam	isoxadifen
B-813	atrazine + H-1	isoxadifen
B-814	atrazine + glyphosate	isoxadifen
B-815	atrazine + mesotrione	isoxadifen
B-816	atrazine + nicosulfuron	isoxadifen
B-817	atrazine + tembotrione	isoxadifen
B-818	atrazine + topramezone	isoxadifen
B-819	clomazone + glyphosate	isoxadifen
B-820	diflufenican + clodinafop-propargyl	isoxadifen
B-821	diflufenican + fenoxaprop-P-ethyl	isoxadifen
B-822	diflufenican + flupyrsulfuron-methyl-sodium	isoxadifen
B-823	diflufenican + glyphosate	isoxadifen

	Herbicide(s) B	Safener C
B-824	diflufenican + mesosulfuron-methyl	isoxadifen
B-825	diflufenican + pinoxaden	isoxadifen
B-826	diflufenican + pyroxsulam	isoxadifen
B-827	flumetsulam + glyphosate	isoxadifen
B-828	flumioxazin + glyphosate	isoxadifen
B-829	imazapic + glyphosate	isoxadifen
B-830	imazethapyr + glyphosate	isoxadifen
B-831	isoxaflutol + H-1	isoxadifen
B-832	isoxaflutol + glyphosate	isoxadifen
B-833	metazachlor + H-1	isoxadifen
B-834	metazachlor + glyphosate	isoxadifen
B-835	metazachlor + mesotrione	isoxadifen
B-836	metazachlor + nicosulfuron	isoxadifen
B-837	metazachlor + terbuthylazine	isoxadifen
B-838	metazachlor + topramezone	isoxadifen
B-839	metribuzin + glyphosate	isoxadifen
B-840	pendimethalin + H-1	isoxadifen
B-841	pendimethalin + clodinafop-propargyl	isoxadifen
B-842	pendimethalin + fenoxaprop-P-ethyl	isoxadifen
B-843	pendimethalin + flupyrsulfuron-methyl-sodium	isoxadifen
B-844	pendimethalin + glyphosate	isoxadifen
B-845	pendimethalin + mesosulfuron-methyl	isoxadifen
B-846	pendimethalin + mesotrione	isoxadifen
B-847	pendimethalin + nicosulfuron	isoxadifen
B-848	pendimethalin + pinoxaden	isoxadifen
B-849	pendimethalin + pyroxsulam	isoxadifen
B-850	pendimethalin + tembotrione	isoxadifen
B-851	pendimethalin + topramezone	isoxadifen
B-852	pyroxasulfone + tembotrione	isoxadifen
B-853	pyroxasulfone + topramezone	isoxadifen
B-854	sulfentrazone + glyphosate	isoxadifen
B-855	terbuthylazine + H-1	isoxadifen
B-856	terbuthylazine + foramsulfuron	isoxadifen
B-857	terbuthylazine + glyphosate	isoxadifen
B-858	terbuthylazine + mesotrione	isoxadifen
B-859	terbuthylazine + nicosulfuron	isoxadifen
B-860	terbuthylazine + tembotrione	isoxadifen
B-861	terbuthylazine + topramezone	isoxadifen
B-862	trifluralin + glyphosate	isoxadifen
B-863	clodinafop-propargyl	mefenpyr

	Herbicide(s) B	Safener C
B-864	cycloxydim	mefenpyr
B-865	cyhalofop-butyl	mefenpyr
B-866	fenoxaprop-P-ethyl	mefenpyr
B-867	pinoxaden	mefenpyr
B-868	profoxydim	mefenpyr
B-869	tepraloxydim	mefenpyr
B-870	tralkoxydim	mefenpyr
B-871	esprocarb	mefenpyr
B-872	prosulfocarb	mefenpyr
B-873	thiobencarb	mefenpyr
B-874	triallate	mefenpyr
B-875	bensulfuron-methyl	mefenpyr
B-876	bispyribac-sodium	mefenpyr
B-877	cyclosulfamuron	mefenpyr
B-878	flumetsulam	mefenpyr
B-879	flupyrsulfuron-methyl-sodium	mefenpyr
B-880	foramsulfuron	mefenpyr
B-881	imazamox	mefenpyr
B-882	imazapic	mefenpyr
B-883	imazapyr	mefenpyr
B-884	imazaquin	mefenpyr
B-885	imazethapyr	mefenpyr
B-886	imazosulfuron	mefenpyr
B-887	iodosulfuron-methyl-sodium	mefenpyr
B-888	mesosulfuron	mefenpyr
B-889	nicosulfuron	mefenpyr
B-890	penoxsulam	mefenpyr
B-891	propoxycarbazone-sodium	mefenpyr
B-892	pyrazosulfuron-ethyl	mefenpyr
B-893	pyroxsulam	mefenpyr
B-894	rimsulfuron	mefenpyr
B-895	sulfosulfuron	mefenpyr
B-896	thiencarbazone-methyl	mefenpyr
B-897	tritosulfuron	mefenpyr
B-898	2,4-D and its salts and esters	mefenpyr
B-899	aminopyralid and its salts and esters	mefenpyr
B-900	clopyralid and its salts and esters	mefenpyr
B-901	dicamba and its salts and esters	mefenpyr
B-902	fluroxypyr-meptyl	mefenpyr
B-903	quinclorac	mefenpyr

	Herbicide(s) B	Safener C
B-904	quinmerac	mefenpyr
B-905	H-9	mefenpyr
B-906	diflufenzopyr	mefenpyr
B-907	diflufenzopyr-sodium	mefenpyr
B-908	clomazone	mefenpyr
B-909	diflufenican	mefenpyr
B-910	fluorochloridone	mefenpyr
B-911	isoxaflutol	mefenpyr
B-912	mesotrione	mefenpyr
B-913	picolinafen	mefenpyr
B-914	sulcotrione	mefenpyr
B-915	tefuryltrione	mefenpyr
B-916	tembotrione	mefenpyr
B-917	topramezone	mefenpyr
B-918	H-7	mefenpyr
B-919	atrazine	mefenpyr
B-920	diuron	mefenpyr
B-921	fluometuron	mefenpyr
B-922	hexazinone	mefenpyr
B-923	isoproturon	mefenpyr
B-924	metribuzin	mefenpyr
B-925	propanil	mefenpyr
B-926	terbuthylazine	mefenpyr
B-927	paraquat dichloride	mefenpyr
B-928	flumioxazin	mefenpyr
B-929	oxyfluorfen	mefenpyr
B-930	saflufenacil	mefenpyr
B-931	sulfentrazone	mefenpyr
B-932	H-1	mefenpyr
B-933	H-2	mefenpyr
B-934	glyphosate	mefenpyr
B-935	glyphosate-isopropylammonium	mefenpyr
B-936	glyphosate-trimesium (sulfosate)	mefenpyr
B-937	glufosinate	mefenpyr
B-938	glufosinate-ammonium	mefenpyr
B-939	pendimethalin	mefenpyr
B-940	trifluralin	mefenpyr
B-941	acetochlor	mefenpyr
B-942	cafenstrole	mefenpyr
B-943	dimethenamid-P	mefenpyr

	Herbicide(s) B	Safener C
B-944	fentrazamide	mefenpyr
B-945	flufenacet	mefenpyr
B-946	mefenacet	mefenpyr
B-947	metazachlor	mefenpyr
B-948	metolachlor-S	mefenpyr
B-949	pyroxasulfone	mefenpyr
B-950	isoxaben	mefenpyr
B-951	dymron	mefenpyr
B-952	indanofan	mefenpyr
B-953	oxaziclomefone	mefenpyr
B-954	triaziflam	mefenpyr
B-955	atrazine + H-1	mefenpyr
B-956	atrazine + glyphosate	mefenpyr
B-957	atrazine + mesotrione	mefenpyr
B-958	atrazine + nicosulfuron	mefenpyr
B-959	atrazine + tembotrione	mefenpyr
B-960	atrazine + topramezone	mefenpyr
B-961	clomazone + glyphosate	mefenpyr
B-962	diflufenican + clodinafop-propargyl	mefenpyr
B-963	diflufenican + fenoxaprop-P-ethyl	mefenpyr
B-964	diflufenican + flupyrsulfuron-methyl-sodium	mefenpyr
B-965	diflufenican + glyphosate	mefenpyr
B-966	diflufenican + mesosulfuron-methyl	mefenpyr
B-967	diflufenican + pinoxaden	mefenpyr
B-968	diflufenican + pyroxsulam	mefenpyr
B-969	flumetsulam + glyphosate	mefenpyr
B-970	flumioxazin + glyphosate	mefenpyr
B-971	imazapic + glyphosate	mefenpyr
B-972	imazethapyr + glyphosate	mefenpyr
B-973	isoxaflutol + H-1	mefenpyr
B-974	isoxaflutol + glyphosate	mefenpyr
B-975	metazachlor + H-1	mefenpyr
B-976	metazachlor + glyphosate	mefenpyr
B-977	metazachlor + mesotrione	mefenpyr
B-978	metazachlor + nicosulfuron	mefenpyr
B-979	metazachlor + terbutylazine	mefenpyr
B-980	metazachlor + topramezone	mefenpyr
B-981	metribuzin + glyphosate	mefenpyr
B-982	pendimethalin + H-1	mefenpyr
B-983	pendimethalin + clodinafop-propargyl	mefenpyr

	Herbicide(s) B	Safener C
B-984	pendimethalin + fenoxaprop-P-ethyl	mefenpyr
B-985	pendimethalin + flupyrsulfuron-methyl-sodium	mefenpyr
B-986	pendimethalin + glyphosate	mefenpyr
B-987	pendimethalin + mesosulfuron-methyl	mefenpyr
B-988	pendimethalin + mesotrione	mefenpyr
B-989	pendimethalin + nicosulfuron	mefenpyr
B-990	pendimethalin + pinoxaden	mefenpyr
B-991	pendimethalin + pyroxsulam	mefenpyr
B-992	pendimethalin + tembotrione	mefenpyr
B-993	pendimethalin + topramezone	mefenpyr
B-994	pyroxasulfone + tembotrione	mefenpyr
B-995	pyroxasulfone + topramezone	mefenpyr
B-996	sulfentrazone + glyphosate	mefenpyr
B-997	terbuthylazine + H-1	mefenpyr
B-998	terbuthylazine + foramsulfuron	mefenpyr
B-999	terbuthylazine + glyphosate	mefenpyr
B-1000	terbuthylazine + mesotrione	mefenpyr
B-1001	terbuthylazine + nicosulfuron	mefenpyr
B-1002	terbuthylazine + tembotrione	mefenpyr
B-1003	terbuthylazine + topramezone	mefenpyr
B-1004	trifluralin + glyphosate	mefenpyr
B-1005	clodinafop-propargyl	H-12
B-1006	cycloxydim	H-12
B-1007	cyhalofop-butyl	H-12
B-1008	fenoxaprop-P-ethyl	H-12
B-1009	pinoxaden	H-12
B-1010	profoxydim	H-12
B-1011	tepraloxydim	H-12
B-1012	tralkoxydim	H-12
B-1013	esprocarb	H-12
B-1014	prosulfocarb	H-12
B-1015	thiobencarb	H-12
B-1016	triallate	H-12
B-1017	bensulfuron-methyl	H-12
B-1018	bispyribac-sodium	H-12
B-1019	cyclosulfamuron	H-12
B-1020	flumetsulam	H-12
B-1021	flupyrsulfuron-methyl-sodium	H-12
B-1022	foramsulfuron	H-12
B-1023	imazamox	H-12

	Herbicide(s) B	Safener C
B-1024	imazapic	H-12
B-1025	imazapyr	H-12
B-1026	imazaquin	H-12
B-1027	imazethapyr	H-12
B-1028	imazosulfuron	H-12
B-1029	iodosulfuron-methyl-sodium	H-12
B-1030	mesosulfuron	H-12
B-1031	nicosulfuron	H-12
B-1032	penoxsulam	H-12
B-1033	propoxycarbazone-sodium	H-12
B-1034	pyrazosulfuron-ethyl	H-12
B-1035	pyroxsulam	H-12
B-1036	rimisulfuron	H-12
B-1037	sulfosulfuron	H-12
B-1038	thiencarbazone-methyl	H-12
B-1039	tritosulfuron	H-12
B-1040	2,4-D and its salts and esters	H-12
B-1041	aminopyralid and its salts and esters	H-12
B-1042	clopyralid and its salts and esters	H-12
B-1043	dicamba and its salts and esters	H-12
B-1044	fluroxypyr-meptyl	H-12
B-1045	quinclorac	H-12
B-1046	quinmerac	H-12
B-1047	B-9	H-12
B-1048	diflufenzopyr	H-12
B-1049	diflufenzopyr-sodium	H-12
B-1050	clomazone	H-12
B-1051	diflufenican	H-12
B-1052	fluorochloridone	H-12
B-1053	isoxaflutol	H-12
B-1054	mesotrione	H-12
B-1055	picolinafen	H-12
B-1056	sulcotrione	H-12
B-1057	tefuryltrione	H-12
B-1058	tembotrione	H-12
B-1059	topramezone	H-12
B-1060	H-7	H-12
B-1061	atrazine	H-12
B-1062	diuron	H-12
B-1063	fluometuron	H-12

	Herbicide(s) B	Safener C
B-1064	hexazinone	H-12
B-1065	isoproturon	H-12
B-1066	metribuzin	H-12
B-1067	propanil	H-12
B-1068	terbuthylazine	H-12
B-1069	paraquat dichloride	H-12
B-1070	flumioxazin	H-12
B-1071	oxyfluorfen	H-12
B-1072	saflufenacil	H-12
B-1073	sulfentrazone	H-12
B-1074	H-1	H-12
B-1075	H-2	H-12
B-1076	glyphosate	H-12
B-1077	glyphosate-isopropylammonium	H-12
B-1078	glyphosate-trimesium (sulfosate)	H-12
B-1079	glufosinate	H-12
B-1080	glufosinate-ammonium	H-12
B-1081	pendimethalin	H-12
B-1082	trifluralin	H-12
B-1083	acetochlor	H-12
B-1084	cafenstrole	H-12
B-1085	dimethenamid-P	H-12
B-1086	fentrazamide	H-12
B-1087	flufenacet	H-12
B-1088	mefenacet	H-12
B-1089	metazachlor	H-12
B-1090	metolachlor-S	H-12
B-1091	pyroxasulfone	H-12
B-1092	isoxaben	H-12
B-1093	dymron	H-12
B-1094	indanofan	H-12
B-1095	oxaziclomefone	H-12
B-1096	triaziflam	H-12
B-1097	atrazine + H-1	H-12
B-1098	atrazine + glyphosate	H-12
B-1099	atrazine + mesotrione	H-12
B-1100	atrazine + nicosulfuron	H-12
B-1101	atrazine + tembotrione	H-12
B-1102	atrazine + topramezone	H-12
B-1103	clomazone + glyphosate	H-12

	Herbicide(s) B	Safener C
B-1104	diflufenican + clodinafop-propargyl	H-12
B-1105	diflufenican + fenoxaprop-P-ethyl	H-12
B-1106	diflufenican + flupyrsulfuron-methyl-sodium	H-12
B-1107	diflufenican + glyphosate	H-12
B-1108	diflufenican + mesosulfuron-methyl	H-12
B-1109	diflufenican + pinoxaden	H-12
B-1110	diflufenican + pyroxsulam	H-12
B-1111	flumetsulam + glyphosate	H-12
B-1112	flumioxazin + glyphosate	H-12
B-1113	imazapic + glyphosate	H-12
B-1114	imazethapyr + glyphosate	H-12
B-1115	isoxaflutol + H-1	H-12
B-1116	isoxaflutol + glyphosate	H-12
B-1117	metazachlor + H-1	H-12
B-1118	metazachlor + glyphosate	H-12
B-1119	metazachlor + mesotrione	H-12
B-1120	metazachlor + nicosulfuron	H-12
B-1121	metazachlor + terbuthylazine	H-12
B-1122	metazachlor + topramezone	H-12
B-1123	metribuzin + glyphosate	H-12
B-1124	pendimethalin + H-1	H-12
B-1125	pendimethalin + clodinafop-propargyl	H-12
B-1126	pendimethalin + fenoxaprop-P-ethyl	H-12
B-1127	pendimethalin + flupyrsulfuron-methyl-sodium	H-12
B-1128	pendimethalin + glyphosate	H-12
B-1129	pendimethalin + mesosulfuron-methyl	H-12
B-1130	pendimethalin + mesotrione	H-12
B-1131	pendimethalin + nicosulfuron	H-12
B-1132	pendimethalin + pinoxaden	H-12
B-1133	pendimethalin + pyroxsulam	H-12
B-1134	pendimethalin + tembotrione	H-12
B-1135	pendimethalin + topramezone	H-12
B-1136	pyroxasulfone + tembotrione	H-12
B-1137	pyroxasulfone + topramezone	H-12
B-1138	sulfentrazone + glyphosate	H-12
B-1139	terbuthylazine + H-1	H-12
B-1140	terbuthylazine + foramsulfuron	H-12
B-1141	terbuthylazine + glyphosate	H-12
B-1142	terbuthylazine + mesotrione	H-12
B-1143	terbuthylazine + nicosulfuron	H-12

	Herbicide(s) B	Safener C
B-1144	terbuthylazine + tembotrione	H-12
B-1145	terbuthylazine + topramezone	H-12
B-1146	trifluralin + glyphosate	H-12
B-1147	2-1	--
B-1148	2-2	--
B-1149	2-3	--
B-1150	2-4	--
B-1151	2-5	--
B-1152	2-6	--
B-1153	2-7	--
B-1154	2-8	--
B-1155	2-9	--
B-1156	2-1	benoxacor
B-1157	2-2	benoxacor
B-1158	2-3	benoxacor
B-1159	2-4	benoxacor
B-1160	2-5	benoxacor
B-1161	2-6	benoxacor
B-1162	2-7	benoxacor
B-1163	2-8	benoxacor
B-1164	2-9	benoxacor
B-1165	2-1	cloquintocet
B-1166	2-2	cloquintocet
B-1167	2-3	cloquintocet
B-1168	2-4	cloquintocet
B-1169	2-5	cloquintocet
B-1170	2-6	cloquintocet
B-1171	2-7	cloquintocet
B-1172	2-8	cloquintocet
B-1173	2-9	cloquintocet
B-1174	2-1	cyprosulfamide
B-1175	2-2	cyprosulfamide
B-1176	2-3	cyprosulfamide
B-1177	2-4	cyprosulfamide
B-1178	2-5	cyprosulfamide
B-1179	2-6	cyprosulfamide
B-1180	2-7	cyprosulfamide
B-1181	2-8	cyprosulfamide
B-1182	2-9	cyprosulfamide
B-1183	2-1	dichlormid

	Herbicide(s) B	Safener C
B-1184	2-2	dichlormid
B-1185	2-3	dichlormid
B-1186	2-4	dichlormid
B-1187	2-5	dichlormid
B-1188	2-6	dichlormid
B-1189	2-7	dichlormid
B-1190	2-8	dichlormid
B-1191	2-9	dichlormid
B-1192	2-1	fenchlorazole
B-1193	2-2	fenchlorazole
B-1194	2-3	fenchlorazole
B-1195	2-4	fenchlorazole
B-1196	2-5	fenchlorazole
B-1197	2-6	fenchlorazole
B-1198	2-7	fenchlorazole
B-1199	2-8	fenchlorazole
B-1200	2-9	fenchlorazole
B-1201	2-1	isoxadifen
B-1202	2-2	isoxadifen
B-1203	2-3	isoxadifen
B-1204	2-4	isoxadifen
B-1205	2-5	isoxadifen
B-1206	2-6	isoxadifen
B-1207	2-7	isoxadifen
B-1208	2-8	isoxadifen
B-1209	2-9	isoxadifen
B-1210	2-1	mefenpyr
B-1211	2-2	mefenpyr
B-1212	2-3	mefenpyr
B-1213	2-4	mefenpyr
B-1214	2-5	mefenpyr
B-1215	2-6	mefenpyr
B-1216	2-7	mefenpyr
B-1217	2-8	mefenpyr
B-1218	2-9	mefenpyr
B-1219	2-1	H-11
B-1220	2-2	H-11
B-1221	2-3	H-11
B-1222	2-4	H-11
B-1223	2-5	H-11

	Herbicide(s) B	Safener C
B-1224	2-6	H-11
B-1225	2-7	H-11
B-1226	2-8	H-11
B-1227	2-9	H-11
B-1228	2-1	H-12
B-1229	2-2	H-12
B-1230	2-3	H-12
B-1231	2-4	H-12
B-1232	2-5	H-12
B-1233	2-6	H-12
B-1234	2-7	H-12
B-1235	2-8	H-12
B-1236	2-9	H-12

The compounds I and the compositions according to the invention may also have a plant-strengthening action. Accordingly, they are suitable for mobilizing the defense system of the plants against attack by unwanted microorganisms, such as harmful fungi, but also viruses and bacteria. Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances which are capable of stimulating the defense system of treated plants in such a way that, when subsequently inoculated by unwanted microorganisms, the treated plants display a substantial degree of resistance to these microorganisms.

The compounds I can be employed for protecting plants against attack by unwanted microorganisms within a certain period of time after the treatment. The period of time within which their protection is effected generally extends from 1 to 28 days, preferably from 1 to 14 days, after the treatment of the plants with the compounds I, or, after treatment of the seed, for up to 9 months after sowing.

The compounds I and the compositions according to the invention are also suitable for increasing the harvest yield.

Moreover, they have reduced toxicity and are tolerated well by the plants.

Use examples

The herbicidal activity of the compounds of the formula I was demonstrated by the following greenhouse experiments:

The culture containers used were plastic flowerpots containing loamy sand with approximately 3.0% of humus as the substrate. The seeds of the test plants were sown separately for each species.

For the pre-emergence treatment, the active ingredients, which had been suspended or emulsified in water, were applied directly after sowing by means of finely distributing nozzles.

The containers were irrigated gently to promote germination and growth and subsequently covered with transparent plastic hoods until the plants had rooted. This cover caused uniform germination of the test plants, unless this has been impaired by the active ingredients.

5 For the post-emergence treatment, the test plants were first grown to a height of 3 to 15 cm, depending on the plant habit, and only then treated with the active ingredients which had been suspended or emulsified in water. For this purpose, the test plants were either sown directly and grown in the same containers, or they were first grown separately as seedlings and transplanted into the test containers a few days prior to treatment.


10 Depending on the species, the plants were kept at 10-25°C or 20-35°C. The test period extended over 2 to 4 weeks. During this time, the plants were tended, and their response to the individual treatments was evaluated.

Evaluation was carried out using a scale from 0 to 100. 100 means no emergence of the plants, or complete destruction of at least the aerial moieties, and 0 means no damage, or normal course of growth. A good herbicidal activity is given at values of at least 70 and a very good herbicidal activity is given at values of at least 85.

15

We claim:

1. A 1,2,5-Oxadiazole compound of the formula I,



5 wherein

X¹ is N or CR¹;

X² is N or CR²;

10 X⁴ is N or CR⁴;

provided that a least one of X¹, X² and X⁴ is N;

15 R is selected from the group consisting of hydrogen, cyano, nitro, halogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, O-R^a, Z-S(O)_n-R^b, Z-C(=O)-R^c, Z-C(=O)-OR^d, Z-C(=O)-NR^eR^f, Z-NR^gR^h,
 20 Z-phenyl and Z-heterocyclyl, where heterocyclyl is a 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R', which are
 25 identical or different;

R¹ is selected from the group consisting of Z¹-cyano, halogen, nitro, C₁-C₈-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₈-haloalkyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, Z¹-C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio-C₁-C₄-alkyl, Z¹-C₁-C₄-alkylthio-C₁-C₄-alkylthio, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₁-C₆-haloalkoxy, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, Z¹-C₁-C₄-haloalkoxy-C₁-C₄-alkoxy, Z¹-S(O)_k-R^{1b}, Z¹-phenoxy and Z¹-heterocyclyloxy, where heterocyclyloxy is an oxygen bound 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members,
 30 which are selected from the group consisting of O, N and S, where the cyclic groups in phenoxy and heterocyclyloxy are unsubstituted or substituted by 1, 2, 3 or 4 groups R¹¹, which are identical or different;

35

R^2, R^3 are identical or different and independently selected from the group consisting of hydrogen, halogen, Z^2 -OH, Z^2 -NO₂, Z^2 -cyano, C₁-C₆-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, Z^2 -C₃-C₁₀-cycloalkyl, Z^2 -C₃-C₁₀-cycloalkoxy, where the C₃-C₁₀-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₈-haloalkyl, Z^2 -C₁-C₈-alkoxy, Z^2 -C₁-C₈-haloalkoxy, Z^2 -C₁-C₄-alkoxy-C₁-C₄-alkoxy, Z^2 -C₁-C₄-alkylthio-C₁-C₄-alkylthio, Z^2 -C₂-C₈-alkenyloxy, Z^2 -C₂-C₈-alkynyloxy, Z^2 -C₁-C₈-haloalkoxy, Z^2 -C₁-C₄-haloalkoxy-C₁-C₄-alkoxy, Z^2 -(tri-C₁-C₄-alkyl)silyl, Z^2 -S(O)_k-R^{2b}, Z^2 -C(=O)-R^{2c}, Z^2 -C(=O)-OR^{2d}, Z^2 -C(=O)-NR^{2e}R^{2f}, Z^2 -NR^{2g}R^{2h}, Z^{2a}-phenyl and Z^{2a}-heterocyclyl, where heterocyclyl is a 3-, 4-, 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in Z^{2a}-phenyl and Z^{2a}-heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R²¹, which are identical or different;

R^4, R^5 are identical or different and independently selected from the group consisting of hydrogen, cyano- Z¹, halogen, nitro, C₁-C₈-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₈-haloalkyl, C₁-C₃-alkylamino, C₁-C₃-dialkylamino, C₁-C₃-alkylamino-S(O)_k, C₁-C₃-alkylcarbonyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-Z¹, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkylthio-Z¹, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₁-C₆-haloalkoxy, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkoxy-Z¹, R^{1b}-S(O)_k-Z¹, phenoxy-Z¹ and heterocyclyloxy-Z¹, where heterocyclyloxy is an oxygen bound 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where the cyclic groups in phenoxy and heterocyclyloxy are unsubstituted or substituted by 1, 2, 3 or 4 groups R¹¹, which are identical or different;

provided that if X⁴ is N, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl; and further provided that if X⁴ is CR⁴ and R⁴ is hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl, R⁵ is not hydrogen, halogen, cyano, nitro, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

where for X² = CR², R² together with R³ or together with R¹, if present, may also form a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle, where the fused heterocycle has 1, 2, 3 or 4 heteroatoms selected from O, S and N as ring members, where the fused carbocycle and the fused hetero-

cycle are monocyclic or bicyclic and where the fused carbocycle and the fused heterocycle are unsubstituted or carry 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 radicals R^q;

n is 0, 1 or 2;

k is 0, 1 or 2;

R', R¹¹, R²¹ independently of each other are selected from the group consisting of halogen, NO₂, CN, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy and C₃-C₇-cycloalkoxy or two vicinal radicals R', R¹¹ or R²¹ together may form a group =O (oxo);

Z, Z¹, Z² independently of each other are selected from the group consisting of a covalent bond and C₁-C₄-alkanediyl;

Z^{2a} is selected from the group consisting of a covalent bond, C₁-C₄-alkanediyl, O-C₁-C₄-alkanediyl, C₁-C₄-alkanediyl-O and C₁-C₄-alkanediyl-O-C₁-C₄-alkanediyl;

R^a is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^b, R^{1b}, R^{2b} independently of each other are selected from the group consisting of C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, phenyl and heterocyclyl, heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;

R^c, R^{2c} independently of each other are selected from the group consisting of

- hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl, benzyl or heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl, benzyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;
- R^d, R^{2d} independently of each other are selected from the group consisting of C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;
- R^e, R^f independently of each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, or
- R^e, R^f together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;
- R^{2e}, R^{2f} independently of each other have the meanings given for R^e, R^f;
- R^g is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in

- the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;
- R^h is selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₄-alkyl, where the C₃-C₇-cycloalkyl groups in the two aforementioned radicals are unsubstituted or partially or completely halogenated, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, a radical C(=O)-R^k, phenyl and benzyl, where phenyl and benzyl are unsubstituted or substituted by 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy, or
- R^g, R^h together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated or unsaturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 groups, which are identical or different and selected from the group consisting of =O, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy;
- R^{2g}, R^{2h} independently of each other have the meanings given for R^g, R^h;
- R^k has the meanings given for R^c;
- R^q is selected from the group consisting of halogen, Z^q-OH, Z^q-NO₂, Z^q-cyano, oxo (=O), =N-R^{q1}, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, Z^q-C₁-C₄-alkoxy, Z^q-C₁-C₄-alkoxy-C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, Z^q-C₁-C₄-haloalkoxy, Z^q-C₃-C₁₀-cycloalkyl, O-Z^q-C₃-C₁₀-cycloalkyl, Z^q-(tri-C₁-C₄-alkyl)silyl, Z^q-S(O)_k-R^{q2}, Z²-C(=O)-R^{q3}, Z²-NR^{q4}R^{q5} and Z^q-phenyl, where phenyl in Z^q-phenyl is unsubstituted or substituted by 1, 2, 3 or 4 groups R^{q6}, which are identical or different; where
- Z^q has one of the meanings given for Z;
- R^{q1} C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and C₃-C₇-cycloalkoxy, which is unsubstituted or partially or completely halogenated;

R^{q2} has one of the meanings given for R^b;

R^{q3} has one of the meanings given for R^c;

5 R^{q4}, R^{q5} independently of each other have the meanings given for R^g, R^h;

R^{q6} has one of the meanings given for Rⁱ;

an N-oxide or an agriculturally suitable salt thereof.

10

2. The compound as claimed in claim 1, where R is selected from the group consisting of halogen, cyano, nitro, NH₂, C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C₁-C₄-haloalkyl, C(=O)-R^c, C(=O)-OR^d, C(=O)-NR^eR^f and NH-C(=O)R^k, where

15

R^c is C₁-C₄-alkyl or C₁-C₄-haloalkyl,

R^d is C₁-C₄-alkyl,

R^e is hydrogen or C₁-C₄-alkyl,

R^f is hydrogen or C₁-C₄-alkyl, or

20

R^e, R^f together with the nitrogen atom, to which they are bound may form a 5-, 6- or 7-membered, saturated N-bound heterocyclic radical, which may carry as a ring member a further heteroatom selected from O, S and N and which is unsubstituted or may carry 1, 2, 3 or 4 methyl groups,

R^k is C₁-C₄-alkyl.

25

3. The compound as claimed in claim 1, where R is a radical OR^a, where R^a is selected from the group consisting of H, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl and C₃-C₇-cycloalkyl, which is unsubstituted or partly or completely halogenated.

30

4. The compound as claimed in claim 1, where R is phenyl or heterocyclyl, where heterocyclyl is a 5- or 6-membered monocyclic or 8-, 9- or 10-membered bicyclic saturated, partially unsaturated or aromatic heterocycle, which contains 1, 2, 3 or 4 heteroatoms as ring members, which are selected from the group consisting of O, N and S, where phenyl and heterocyclyl are unsubstituted or substituted by 1, 2, 3 or 4 groups R', where R' is selected from the group consisting of halogen, methyl, ethyl, methoxy and trifluoromethyl.

35

40

5. The compound as claimed in claim 1, where R is S(O)_n-R^b, where R^b is C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₃-C₇-cycloalkyl, phenyl and heterocyclyl, where heterocyclyl is a 6-membered aromatic heterocyclic radical having 1 or 2 nitrogen atoms as ring members.

6. The compound as claimed in any one of claims 1 to 5, where R^3 is selected from the group consisting of hydrogen, cyano, halogen, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_2 - C_4 -alkenyloxy, C_2 - C_4 -alkynyloxy and $S(O)_kR^{2b}$.
7. The compound as claimed in any one of claims 1 to 5, where R^3 is selected from the group consisting of hydrogen, halogen, CN, NO_2 , C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkylthio, $S(O)_2$ - C_1 - C_4 -alkyl and $S(O)_2$ - C_1 - C_4 -haloalkyl.
8. The compound as claimed in any one of the preceding claims, where R^5 is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH_3 , CH_2CH_3 , $CH(CH_3)_2$, cyclopropyl, cyclopropylmethyl, CF_3 , OCH_3 , OCH_2CH_3 , $OCH(CH_3)_2$, CH_2OCH_3 , $CH_2OCH_2CH_2OCH_3$, OCF_3 , $OCHF_2$, SCH_3 , SCF_3 , SCH_2CH_3 , SCH_2CF_3 , $SOCH_3$, $SOCH_2CH_3$, SO_2CH_3 , $SO_2CH_2CH_3$.
9. The compound as claimed in any one of the preceding claims, where X^1 is CR^1 .
10. The compound as claimed in claim 9, where R^1 is selected from the group consisting of cyano, halogen, nitro, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl, Z^1 - C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, Z^1 - C_1 - C_4 -alkylthio- C_1 - C_4 -alkylthio, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyloxy, C_1 - C_6 -haloalkoxy, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkoxy and $S(O)_kR^{1b}$, where k and Z^1 are as defined in claim 1 and where R^{1b} is selected from the group consisting of C_1 - C_4 -alkyl and C_1 - C_4 -haloalkyl.
11. The compound as claimed in claim 9 or 10, where R^1 is selected from the group consisting of halogen, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkylthio, C_1 - C_4 -alkylsulfonyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl.
12. The compound as claimed in claim 9 to 11, where
- R^1 is selected from the group consisting of halogen, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkylthio and C_1 - C_4 -alkylsulfonyl; and
- R^3 is selected from the group consisting of hydrogen, halogen, CN, NO_2 , C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylthio and C_1 - C_4 -alkylsulfonyl.

13. The compound as claimed in any one of claims 1 to 8, where X^1 is N.


14. The compound as claimed in one of the preceding claims, where X^2 is CR^2 .

5 15. The compound as claimed in claim 14, wherein R^2 is different from hydrogen.

16. The compound as claimed in claim 14 or 15, where R^2 is 5- or 6-membered heterocyclyl, where heterocyclyl is a saturated, partially unsaturated or aromatic heterocyclic radical, which contains as ring member 1 heteroatom selected from the group consisting of O, N and S and 0, 1 or 2 further nitrogen atoms, where heterocyclyl is unsubstituted or carries
10 1, 2 or 3 radicals R^{21} which are identical or different.

17. The compound as claimed in claim 14 or 15, where R^2 is 5- or 6-membered heterocyclyl, selected from the group consisting of isoxazoliny, 1,2-dihydropyridonyl, 1,4-dihydropyridonyl, tetrahydrofuryl, dioxolanyl, piperidiny, morpholiny, piperaziny, isoxa-
15 zolyl, pyrazolyl, thiazolyl, oxazolyl, furyl, pyridiny and pyraziny, where heterocyclyl is unsubstituted or carries 1, 2 or 3 radicals R^{21} which are identical or different and selected from the group consisting of halogen C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl and C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl.

20 18. The compound as claimed in claim 14 or 15, where R^2 is a radical of the following formula:



in which # denotes the bond through which the group R^2 is attached and:

25 R^{P1} H or F;

R^{P2} H, F, Cl or OCH_3 ; and

R^{P3} H, F, Cl, CH_3 , CF_3 , OCH_3 , OCH_2OCH_3 or $OCH_2CH_2OCH_3$.

19. The compound as claimed in claim 14 or 15, where R^2 is selected from the group consisting of halogen, C_1 - C_6 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl, C_2 - C_6 -
30 alkenyl, C_2 - C_6 -alkynyl, C_1 - C_4 -alkoxy, C_2 - C_4 -haloalkoxy, C_3 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_3 - C_6 -haloalkenyloxy, C_3 - C_6 -haloalkynyloxy, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylsulfonyl, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfanyl and C_1 - C_4 -haloalkyl- $S(O)_2$.

35 20. The compound as claimed in claim 14, where R^2 together with R^3 or together with R^1 , if present, forms a fused 5-, 6-, 7-, 8-, 9- or 10-membered carbocycle or a fused 5-, 6-, 7-, 8-, 9- or 10-membered heterocycle, where the fused heterocycle has 1, 2, 3 or 4 heteroatoms selected from O, S and N as ring members, where the fused carbocycle and the


fused heterocycle are monocyclic or bicyclic and where the fused carbocycle and the fused heterocycle are unsubstituted or carry 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 radicals R^q .

21. The compound as claimed in any one of claims 1 to 13, where X^2 is N.

22. The compound as claimed in any one of the preceding claims, where X^4 is CR^4 .

23. The compound as claimed in any one of the preceding claims, where R^4 is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH_3 , CH_2CH_3 , $CH(CH_3)_2$, cyclopropyl, cyclopropylmethyl, CF_3 , OCH_3 , OCH_2CH_3 , $OCH(CH_3)_2$, CH_2OCH_3 , $CH_2OCH_2CH_2OCH_3$, OCF_3 , $OCHF_2$, SCH_3 , SCF_3 , SCH_2CH_3 , SCH_2CF_3 , $SOCH_3$, $SOCH_2CH_3$, SO_2CH_3 , $SO_2CH_2CH_3$.

24. The compound as claimed in any one of the preceding claims, where formula I is represented by the formula I.1




where R, R^2 , R^3 , R^4 and R^5 are as defined in any one of claims 1 to 8, 14 to 20, 22 and 23.

25. The compound as claimed in claim 24, where

R^2 is selected from the group consisting of hydrogen, C_1 - C_2 -alkoxy- C_1 - C_2 -alkyl, C_1 - C_2 -haloalkoxy- C_1 - C_2 -alkyl, C_1 - C_4 -alkylsulfonyl, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfanyl, isoxazolyl and isoxazolinyl, where the last two mentioned radicals may be substituted or carry 1 or 2 radicals selected from halogen and C_1 - C_4 -alkyl.

26. The compound as claimed in any one of claims 1 to 23, where formula I is represented by the formula I.2




where R, R^1 , R^3 , R^4 and R^5 are as defined in any one of claims 1 to 12, 22 and 23.

27. The compound as claimed in claim 26, where

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl.

28. The compound as claimed in any one of claims 1 to 23, where formula I is represented by the formula I.3



where R, R³, R⁴ and R⁵ are as defined in any one of claims 1 to 8 and 22 to 23.


29. The compound as claimed in any one of claims 24 to 28, where

R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-alkylsulfonyl;

R⁴ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃, CH(CH₃)₂, cyclopropyl, CF₃, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃; and


R⁵ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃, CH(CH₃)₂, cyclopropyl, CF₃, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

30. The compound as claimed in any one of claims 1 to 23, where formula I is represented by the formula I.4




where R, R¹, R³ and R⁵ are as defined in any one of claims 1 to 12.

31. The compound as claimed in any one of claims 1 to 23, where formula I is represented by the formula I.5




where R, R¹, R², R³ and R⁵ are as defined in any one of claims 1 to 12, 14 to 20.

- 5 32. The compound as claimed in any one of claims 30 or 31, where
 R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-
 alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-
 C₄-haloalkylthio or C₁-C₄-alkylsulfonyl.
- 10 33. The compound as claimed in any one of claims 1 to 23, where formula I is represented by
 the formula I.6




- 15 where R, R², R³ and R⁵ are as defined in any one of claims 1 to 8 and 14 to 20.
34. The compound as claimed in claim 33, where
 R² is selected from the group consisting of hydrogen, C₁-C₂-alkoxy-C₁-C₂-alkyl, C₁-C₂-
 haloalkoxy-C₁-C₂-alkyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfanyl,
 20 isoxazolyl and isoxazoliny, where the last two mentioned radicals may be substitut-
 ed or carry 1 or 2 radicals selected from halogen and C₁-C₄-alkyl.
35. The compound as claimed in claim 33 or 34, where
 R³ is selected from the group consisting of hydrogen, halogen, CN, NO₂, C₁-C₄-alkyl,
 25 C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio and C₁-C₄-
 alkylsulfonyl; and
 R⁵ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃,
 CH(CH₃)₂, cyclopropyl, CF₃, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.
- 30 36. The compound as claimed in any one of claims 1 to 23, where formula I is represented by
 the formula I.7



where R, R¹ and R⁵ are as defined in any one of claims 1 to 5 and 8 to 11.

- 5 37. The compound as claimed in any one of claims 1 to 23, where formula I is represented by the formula I.8



10 where R, R¹ and R⁵ are as defined in any one of claims 1 to 5 and 8 to 11.

38. The compound as claimed in claim 36 or 37, where

R¹ is halogen, nitro, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio or C₁-C₄-alkylsulfonyl.

39. The compound as claimed in any one of claims 36 to 38, where

R⁵ is selected from the group consisting of H, F, Cl, Br, I, CN, nitro, CH₃, CH₂CH₃, CH(CH₃)₂, cyclopropyl, CF₃, OCH₃, OCF₃, SCH₃, SOCH₃, SO₂CH₃ or CH₂OCH₃.

40. The compound as claimed in any one of claims 24 to 39, where

R is selected from the group consisting of C₁-C₄-alkyl and C₁-C₄-alkoxy-C₁-C₄-alkyl.

41. A composition comprising at least one compound as claimed in any one of claims 1 to 40 and at least one auxiliary, which is customary for formulating crop protection compounds.

42. The use of a compound as claimed in any one of claims 1 to 40 or a composition of claim 41 for controlling unwanted vegetation.

43. A method for controlling unwanted vegetation which comprises allowing a herbicidally effective amount of at least one compound as claimed in any one of claims 1 to 40 or a composition of claim 41 to act on plants, their seed and/or their habitat.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2014/059026

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D413/12 C07D471/04 A01N43/82 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, PAJ, WPI Data, BEILSTEIN Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2012/123420 A1 (BAYER CROPSCIENCE AG [DE]; KOEHN ARNIM [DE]; VAN ALMSICK ANDREAS [DE];) 20 September 2012 (2012-09-20) claims; tables 1, 2 -----	1-43
X	WO 2012/123416 A1 (BAYER CROPSCIENCE AG [DE]; KOEHN ARNIM [DE]; VAN ALMSICK ANDREAS [DE];) 20 September 2012 (2012-09-20) claims; examples; tables 1,2 -----	1-43
Y	WO 2011/035874 A1 (BAYER CROPSCIENCE AG [DE]; KOEHN ARNIM [DE]; TIEBES JOERG [DE]; VAN AL) 31 March 2011 (2011-03-31) Formula I; claims 1,5,10,11 ----- -/--	1-43
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 4 June 2014		Date of mailing of the international search report 24/06/2014
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Stroeter, Thomas

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2014/059026

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2012/058892 A1 (BRAUN RALF [DE] ET AL) 8 March 2012 (2012-03-08) claims; examples	1-43

X,P	WO 2013/072450 A1 (BASF SE [DE]) 23 May 2013 (2013-05-23) the whole document	1-43

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/059026

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012123420 A1	20-09-2012	AR 085812 A1	30-10-2013
		AU 2012228366 A1	03-10-2013
		CA 2830090 A1	20-09-2012
		CN 103429578 A	04-12-2013
		EA 201391281 A1	28-02-2014
		EP 2686309 A1	22-01-2014
		JP 2014508177 A	03-04-2014
		KR 20140006955 A	16-01-2014
		US 2014121104 A1	01-05-2014
		WO 2012123420 A1	20-09-2012
WO 2012123416 A1	20-09-2012	AR 085811 A1	30-10-2013
		CN 103459386 A	18-12-2013
		EP 2686316 A1	22-01-2014
		JP 2014509597 A	21-04-2014
		US 2014066301 A1	06-03-2014
		WO 2012123416 A1	20-09-2012
WO 2011035874 A1	31-03-2011	AR 078312 A1	26-10-2011
		AU 2010297635 A1	19-04-2012
		CA 2775104 A1	31-03-2011
		CN 102639517 A	15-08-2012
		EP 2480539 A1	01-08-2012
		ES 2428104 T3	05-11-2013
		JP 2013505901 A	21-02-2013
		RU 2012111125 A	27-09-2013
		US 2011152084 A1	23-06-2011
		WO 2011035874 A1	31-03-2011
US 2012058892 A1	08-03-2012	AR 082828 A1	09-01-2013
		AU 2011298424 A1	21-03-2013
		CA 2809487 A1	08-03-2012
		CN 103282354 A	04-09-2013
		CO 6670599 A2	15-05-2013
		EA 201390265 A1	30-08-2013
		EP 2611785 A1	10-07-2013
		JP 2013536817 A	26-09-2013
		KR 20130101506 A	13-09-2013
		TW 201221059 A	01-06-2012
		US 2012058892 A1	08-03-2012
		UY 33583 A	30-03-2012
		WO 2012028579 A1	08-03-2012
WO 2013072450 A1	23-05-2013	NONE	