

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number
WO 02/28175 A2

(51) International Patent Classification⁷: **A01K 67/027**

(74) Agents: **MARTIN, Jean-Jacques** et al.; Cabinet Regimbeau, 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).

(21) International Application Number: PCT/IB01/02246

(81) Designated States (*national*): CA, JP.

(22) International Filing Date:

28 September 2001 (28.09.2001)

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

00/12570

3 October 2000 (03.10.2000)

FR

09/853,033

11 May 2001 (11.05.2001)

US

Published:

— without international search report and to be republished upon receipt of that report

— with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(71) Applicant: **ASSOCIATION POUR LE DEVELOPPEMENT DE LA RECHERCHE EN GENETIQUE MOLECULAIRE (ADEREGEM)** [FR/FR]; 231, rue de Charenton, F-75012 Paris (FR).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors: **CHAMBON, Pierre**; 4, rue du Dr. Albert Schweitzer, F-67113 Blaesheim (FR). **METZGER, Daniel**; 21, avenue Jean Jaurès, F-67100 Strasbourg (FR).

(54) Title: TRANSGENIC MOUSE FOR TARGETED RECOMBINATION MEDIATED BY MODIFIED CRE-ER

(57) Abstract: The present invention relates to a metazoan organism, with the exception of humans, and in particular a mouse, characterized in that at least one cell of this organism comprises at least one fusion protein between a recombinase Cre and a modified ligand binding domain of the nuclear estrogen receptor alpha, allowing the inactive fused recombinase to be induced by synthetic antiestrogens, but not by natural estrogens, and one or more DNA sequences of interest belonging to the genome of said organism into which one or more sites of recognition of said recombinase protein are inserted. The invention also covers the methods using said organism for the screening of medicaments, the mutagenesis and the analysis of the biological function of the DNA sequence(s) of interest, in particular of gene(s) of interest, such as RXR α .



WO 02/28175 A2

- 1 -

"TRANSGENIC MOUSE FOR TARGETED RECOMBINATION MEDIATED
BY MODIFIED CRE-ER".

The present invention relates to a metazoan organism, with the exception of humans, and in particular a mouse, characterized in that at least one cell of this organism comprises at least one fusion
5 protein between a recombinase Cre and a modified nuclear estrogen receptor allowing it to respond to synthetic antiestrogens but not to natural estrogens, and one or more DNA sequences of interest belonging to the genome of said organism into which one or more
10 sites of recognition of said recombinase protein are inserted. The invention also covers the methods using said organism for the mutagenesis and the analysis of the biological function of the DNA sequence(s) of interest, in particular of the gene(s) of interest,
15 such as RXR α .

The ability to modify the genome of animals, more particularly of mice, by integrating transgenes randomly or at preselected sites, by homologous recombination, in embryonic stem cells (ES cells) has
20 made it possible to greatly improve our understanding of the biological function of mammalian genes under normal and/or pathological conditions (Jaenisch, 1988; Capecchi, 1989). However, these techniques have proved not to be very informative in a large number of cases,
25 in particular because the hereditary mutations thus generated were lethal during development and/or because their effects were pleiotropic.

- 2 -

To remedy these defects, strategies for conditional somatic mutagenesis have been developed, particularly in mice; they make it possible to selectively induce mutations in a given cell type
5 (spatial control) or at a given time (temporal control) during the life of the animal.

A first strategy consists in combining the targeted homologous recombination with the site-specific recombination systems based on the use of
10 recombinases which catalyze the recombination reaction between two short recognition DNA sequences. It has been shown that these site-specific recombination systems, although of microbial origin for the majority, could function in higher eukaryotes, such as plants,
15 insects and mice (Sauer, 1994; Rajewsky et al., 1996; Sauer, 1998). Among the site-specific recombination systems commonly used, there may be mentioned the Cre/Lox (Sauer, 1998) and FLP/FRT (Kilby et al., 1993) systems. The strategy normally used consists in
20 inserting the loxP (or FRT) sites into the chromosomes of ES cells by homologous recombination, or by conventional transgenesis, and then in delivering Cre (or FLP) for the latter to catalyze the recombination reaction. The recombination between the two loxP (or
25 FRT) sites may be obtained in ES cells (Gu et al., 1993) or in fertilized eggs (Araki et al., 1995) by transient expression of Cre or using a Cre transgenic mouse (Lakso et al., 1992; Orban et al., 1992). Such a strategy of somatic mutagenesis allows a spatial
30 control of the recombination, because the expression of

the recombinase is controlled by a promoter specific for a given tissue or for a given cell. However, this strategy also has limitations because some somatic alterations can lead to a lethal phenotype at an early
5 stage of development, thus preventing any subsequent biological or physiological study. Also, an insufficiently specific expression of the recombinase can lead to recombination events in a non-desired cell type (Betz et al., 1996) which, if they occur early
10 during embryogenesis, can cause recombination of the DNA in the majority of the adult tissues, and thereby complicate the analysis of the mutant phenotype.

A second strategy has consisted in controlling the expression of recombinases over time so as to allow
15 temporal control of somatic recombination. To do this, the expression of the recombinases is controlled by inducible promoters (Kühn et al., 1995; Saint-Onge et al., 1996), such as the interferon-inducible promoter, for example. This system also has limitations because
20 it does not make it possible to obtain spatial control of the recombination.

The coupling of the tetracycline-inducible expression system developed by H. Bujard (Gossen et al., 1992; WO 94 04672; EP 804 546) with the site-
25 specific recombinase system has made it possible to develop a system for somatic modification of the genome which is controlled spatiotemporally. Such a system is based on the activation or repression, by tetracycline, of the promoter controlling the expression of the
30 recombinase gene. Such a method, although making it

possible to obtain a spatiotemporal control of the somatic recombination, has the disadvantage of being cumbersome to carry out because it requires the creation of a doubly transgenic animal.

5 It has been possible to envisage a new strategy following the development of chimeric recombinases selectively activated by the natural ligand for the estrogen receptor. Indeed, the observation that the activity of numerous proteins, including at least two
10 enzymes (the tyrosine kinases c-abl and src) is controlled by estrogens, when the latter is linked to the ligand-binding domain (LBD) of the estrogen receptor α (ER α) (Jackson *et al.*, 1993; Picard *et al.*, 1994) has made it possible to develop strategies for
15 spatiotemporally controlled site-specific recombination (Logie *et al.*, 1995; Metzger *et al.*, 1995). However, to use such chimeric recombinases to successfully carry out conditional somatic mutagenesis in vertebrates (in particular mice) which produce estrogens, it was
20 necessary to create recombinases which are not activated by the estrogens present in the animal, otherwise the temporal control of the recombination of the target genes would not be obtained. Thus, mutations were introduced into the ER α LBD, and it has been
25 shown, in cells in culture, that the chimeric recombinase Cre-ER^T no longer responds to the natural estrogens, despite being efficiently activated by antiestrogens such as 4-hydroxytamoxifen (OHT) (Feil *et al.*, 1996).

The feasibility of the site-specific somatic recombination activated by an antiestrogenic ligand has thus been demonstrated for "reporter" DNA sequences, in mice, and in particular in various transgenic mouse
5 lines which express the fusion protein Cre-ER^T activated by Tamoxifen (Tam) (Feil et al., 1996; Brocard et al., 1997; Indra et al., 1999). The feasibility of the site-specific recombination activated by a ligand for a gene present in its natural
10 chromatin environment has been demonstrated in mice by Schwenk et al. (1998). Schwenk et al. have thus carried out the deletion, inducible by injection of Tamoxifen, of a pol β gene in B cells using a mouse expressing, specifically in the B lymphocytes, a fusion protein
15 between the recombinase Cre and the ligand-binding domain of the mutated human nuclear estrogen receptor. However, the technology developed by Schwenk et al. does not make it possible to obtain a satisfactory efficiency of spatiotemporally controlled site-specific
20 recombination in cells expressing the fusion protein because the efficiency varies between a few percents and 80%, in spite of the use of high doses of OHT (five injections of 8 mg). Moreover, the results presented by Schwenk et al. show that in the absence of synthetic
25 ligand, the activity of the fusion protein used may be induced by natural ligands generating a non negligible residual "background noise" which may be detected by PCR.

Up until now, the possibility of carrying out
30 the ligand-activated, site-specific somatic

recombination of chromosomal DNA sequence(s), in their natural chromatin environment, has therefore never been able to be satisfactorily demonstrated in animals, in particular mice, that is to say with a very high
5 efficiency in the presence of synthetic ligand and with a negligible or even zero "background noise" in its absence.

As highlighted by Schwenk et al. in the discussion in their article, there is a real need to
10 develop transgenic animals in whose cells site-specific recombination could be spatiotemporally induced with an efficiency of close to 100% in the presence of synthetic ligand, and which could not occur in any cell in the absence of synthetic ligand and/or in the
15 presence of a natural ligand.

Moreover, a need also exists to develop chimeric recombinases with increased sensitivity to the synthetic ligand, so as to avoid injecting into the animals massive doses of synthetic ligand which can
20 cause in these animals not only unjustified suffering, but can also affect the general metabolism of the animal, which would distort subsequent physiological and behavioral studies.

Unexpectedly, the inventors have solved the
25 problems mentioned earlier, which had not been resolved up until now, by combining the selection of novel mutations in the ligand-binding domain of the human nuclear estrogen receptor, the selection of a suitable hinge region between the two domains of the chimeric
30 recombinase, and the selection of promoters suitable

- 7 -

for directing the expression of the chimeric recombinase in a given tissue.

The present invention therefore relates to a metazoan organism, with the exception of humans,
5 characterized in that at least one cell of said organism comprises at least:

- (i) one fusion protein comprising sequentially:
 - a recombinase protein;
 - a hinge region of at least 15 amino acids;
 - 10 - a polypeptide comprising the ligand-binding domain of the human nuclear estrogen receptor, or of a vertebrate nuclear estrogen receptor, and their natural variants or one of their fragments, said polypeptide exhibiting at least
15 one mutation relative to the wild-type form of said ligand-binding domains, or of their natural variants, or of their fragments, and said fusion protein having a negligible, or even zero, recombinase activity in the presence
20 of a natural ligand, such as for example estradiol, and a recombinase activity induced by a small quantity of synthetic ligand endowed with antiestrogenic activity, such as for example Tam and OHT;
- 25 (ii) one or more gene or intergenic DNA sequences of interest naturally belonging to said genome of said organism into which one or more recognition sites of said recombinase protein are inserted, said DNA sequence(s) of interest

being located in one or more of the chromosomes of the genome of said cell.

The expression "metazoan organism" is understood to mean any animal organism, with the exception of humans, consisting of several cells. According to a preferred embodiment, it is a vertebrate such as for example a mammal, a bird, a fish. Preferably, it is a mammal such as for example a bovine, a porcine, a caprine, an ovine, an equine, a rodent. According to a more preferred embodiment, it is a rodent such as mice or rats.

The expression "recombinase protein" is understood to designate recombinases of the family of integrases which catalyze the excision, insertion, inversion or translocation of DNA fragments at the level of specific sites of recognition said recombinases (Sternberg et al., 1986, Sauer, et al., 1990; Barbonis et al., 1993; Kilby et al., 1993; Sauer, 1994; Denisen et al., 1995). These recombinases are active in animal cells (Sauer, 1994).

The recombinase protein of the invention is preferably selected from the group of site-specific recombinases composed of the Cre recombinase of bacteriophage P1, the FLP recombinase of *Saccharomyces cerevisiae*, the R recombinase of *Zygosaccharomyces rouxii* pSR1, the A recombinase of *Kluyveromyces drosophilarius* pKD1, the A recombinase of *Kluyveromyces waltii* pKW1, the integrase λ Int, the recombinase of the GIN recombination system of the Mu phage, of the

bacterial β recombinase (Diaz et al., 1999) or a variant thereof.

The Cre ("cyclization recombination") recombinase which is a 38 KDa integrase of bacteriophage P1 catalyzes, in the absence of cofactors, recombination between two DNA sequences of 34 basepairs called "loxP site" (Sauer et al., 1990). The position on one or more DNA molecules and the orientation of the loxP sites relative to each other determine the type of function of the Cre recombinase: excision, insertion, inversion or translocation. Thus, the recombinase activity of Cre is an inversion when two loxP sites are inverted on the same DNA fragment, and an excision when the loxP sites are in the form of a direct repeat on the same DNA fragment. The activity of the recombinase is an insertion when the loxP site is present on a DNA fragment, it being possible for a DNA molecule such as a plasmid containing a loxP site to be inserted at the level of said loxP site. The Cre recombinase can also induce translocation between two chromosomes provided that a loxP site is present on each of them (Babinet, 1995). More generally, the Cre recombinase is therefore capable of catalyzing recombination between one or more different DNA molecules provided that they carry loxP sites.

The FLP recombinase of the FLP/FRT system is a recombinase of 43 KDa from *Saccharomyces cerevisiae* which is capable of the same type of action as the Cre recombinase on DNA fragments containing FRT recognition sites (Kilby et al., 1993).

- 10 -

Preferably, the recombinase according to the invention is the Cre recombinase of bacteriophage P1 and its natural or synthetic variants, and said sites of recognition specific for said Cre recombinase are
5 preferably chosen from the group composed of the sequences Lox P, Lox 66, Lox 71, Lox 511, Lox 512, Lox 514.

The expression "variant of the recombinase protein" is understood to mean all the wild-type
10 recombinases or fragments thereof which may exist naturally and which correspond in particular to truncations, substitutions, deletions and/or additions of amino acid residues. These recombinases and fragments thereof are preferably derived from the
15 genetic polymorphism in the population. The expression "recombinase fragment" is understood to mean any recombinase portion exhibiting at least one recombinase activity. The expression variant of the recombinase protein is also understood to mean the synthetic
20 variants for which the above modifications are not naturally present, but were introduced artificially, by genetic engineering for example. Thus, the recombinases derived from chimeric fusion constitute synthetic variants according to the invention. Such recombinases
25 have been described for example in Shaikh and Sadowski (2000).

Said hinge region according to the invention comprises the D hinge region of the nuclear estrogen receptor, preferably the human nuclear estrogen
30 receptor α , or one of its fragments.

- 11 -

The D hinge region (region 263 to 301 of the sequence SEQ ID No. 2) is a region situated between the ER C region which contains the DNA-binding domain (region 180-262 of the sequence SEQ ID No. 2) and the
5 ligand-binding domain (region 302 to 552 of the sequence SEQ ID No. 2).

Preferably, this hinge region sequentially comprises at least (i) two amino acids corresponds to the introduction of a restriction site, of a "linker",
10 or of an adapter, which are necessary for the cloning of the fusion gene, and (ii) one fragment of the D hinge region of the human nuclear estrogen receptor α , corresponding to amino acids 282 to 301 of the sequence SEQ ID No. 2. Preferably, said restriction
15 site is an XhoI site and the two corresponding amino acids are leucine and glutamine.

The hinge region according to the invention has a size of at least 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 40, 45, 50, 55, 60, 65, 70, 100, 150, 200, 250,
20 292 amino acids. According to a preferred embodiment, said hinge region comprises at least 15 amino acids and at most 54 amino acids. More preferably still, the hinge region comprises 23 amino acids. The width of the hinge region influences the regulation of the
25 recombinase activity by the ligand.

The hinge region according to the invention may consist of a peptide which is functionally equivalent to said D hinge region.

The estrogen receptors (ER) are proteins
30 regulating the transcription of genes which mediate the

action of estrogens in the target cells. The ERs belong to the superfamily of nuclear receptors which have a common modular structure: (i) a variable N-terminal A/B region containing the constitutive transactivation activity AF-1, (ii) a DNA-binding central domain C (DBD) which is highly conserved between various species and allowing binding of the receptor to its specific DNA response element, (iii) and a ligand-binding domain (LBD), located in the C-terminal region of ER (for review articles and references see Evans, 1988; Beato et al., 1989; Gronemeyer, 1991; Green and Chambon, 1988; Parker, 1993; Simons, 1994).

The nuclear estrogen receptors according to the invention are chosen from the human nuclear estrogen receptors, and from the nuclear estrogen receptors of vertebrates such as for example the various species of primates, bovines, porcines, ovines, caprines, felines, canines, equines, birds, fish, rodents, in particular rats and mice.

According to a preferred embodiment, the source organism for the estrogen receptor according to the invention is characterized in that said ligand-binding domain (LBD) of the nuclear estrogen receptor, or its natural variants, or one of their fragments, is human and is chosen from the LBDs of the human nuclear α and β estrogen receptors ($ER\alpha$ and $ER\beta$). According to a further preferred mode, this includes the LBD of the human nuclear estrogen receptor α corresponding to amino acids 302 to 552, or its natural variants, or one of their fragments.

The expression "natural variant" is understood to mean all the LBDs of the nuclear estrogen receptors or their fragments which may exist naturally, in particular in human beings, and corresponding in particular to truncations, substitutions, deletions and/or additions of amino acid residues. These natural variants are derived in general from the genetic polymorphism in the population, and have an activity which is not substantially modified compared with the wild-type receptor.

There are also included in the scope of the invention the polypeptides homologous to the LBDs of the wild-type nuclear estrogen receptors, or to their variants, or to one of their fragments, and which exhibit certain modifications, in particular a deletion, addition, substitution of at least one amino acid, a truncation, an extension and/or a chimeric fusion.

The expression "nuclear receptor fragment" is understood to mean any portion of the nuclear estrogen receptor LBDs exhibiting at least the LBD activity.

Said fusion protein according to the invention is therefore preferably Cre-ER and comprises the Cre recombinase protein to which is fused a portion of the D hinge region and the LBD (amino acids 282 to 595 of the sequence SEQ ID No. 2) of the mutated human nuclear estrogen receptor α (SEQ ID No. 2). The fusion protein according to the invention comprises at least the portion of the nuclear estrogen receptor having a ligand binding activity.

- 14 -

Said LBD of the nuclear receptor, or one of its fragments, has at least one mutation. This mutation is preferably chosen from the group:

- mutation (G521R) glycine to arginine at position
5 521 of the sequence SEQ ID No. 2 or of a natural variant of this sequence;
- mutation (G400V) glycine to valine at position
400 of the sequence SEQ ID No. 2 or of a natural variant of this sequence;
- 10 - mutation (methionine-leucine) to (alanine-alanine) situated at position 543-544 (M543A/L544A mutation) of the sequence SEQ ID No. 2 or of a natural variant of this sequence.

The term "mutation" is understood to mean any
15 changes occurring in the sequence of the nuclear estrogen receptor, and in particular of the human nuclear estrogen receptor α , other than those present in its natural variants, and/or in its human or vertebrate homologues, and which substantially modify
20 the activity of the recombinase protein fused to said receptor or to said ligand-binding domain, in response to the binding of a synthetic ligand endowed with antiestrogenic activity.

Among the mutations capable of being introduced
25 into the LBD of the nuclear estrogen receptor, there may be mentioned point mutations, deletions, insertions, substitutions. However, it is advisable to select only the mutations introduced into the LBD of the nuclear estrogen receptors which allow induction of
30 the activity of the Cre recombinase fused to said

- 15 -

receptor by a synthetic ligand in low concentration, while avoiding as far as possible the inducing of a basal activity by the natural ligands for this receptor which are naturally present in the metazoan organism.

5 Likewise, it will be advisable to select³ the mutations which do not confer activity on the Cre recombinase fused to said receptor in the absence of a ligand.

The G521R mutation constitutes an LBD mutation of the ER according to the invention. This mutation is

10 similar to the G525R mutation introduced into the mouse ER LBD (mER) which reduces the affinity for the natural ligand, estradiol, by about 1000 fold, without adversely affecting the binding of the synthetic ligand, 4-hydroxyTamoxifen (OHT) (Danielan et al.,

15 1993). Thus, the inventors have shown that the recombinase activity of the Cre-ER^T (T = for inducible by Tamoxifen) fusion protein which carries the G521R mutation, and the amino acid glycine at position 400, called Cre-ER(GR) in the article by Feil et al. 1997,

20 is dependent on the addition of OHT or of Tam to the medium for culturing transfected cells. On the other hand, no recombinase activity is observed in the presence of OHT when the fusion protein carries the G521R mutation and the G400V mutation (mutant called

25 Cre-ER(VR) in Feil et al., (1997)).

The inventors have also created the fusion protein corresponding to the triple mutant G400V/M543A/L544A called Cre-ER^{T2} (Feil et al., 1997). This fusion protein exhibits a recombinase activity in

30 cultured cells which is induced by the antiestrogen Tam

or OHT, but not by the natural ligand estradiol; moreover, the maximum activity of Cre-ER^{T2} is induced for Tam or OHT doses less than those necessary to activate Cre-ER^T. This increased sensitivity to Tam or
5 OHT of Cre-ER^{T2} compared with Cre-ER^T has been verified in transgenic mice selectively expressing the chimeric recombinases in the basal layer of the epidermis, under the control of the cytokeratin 5 promoter (Indra et al., 1999). The inventors have observed that the
10 translocation of Cre-ER^{T2} from the cytoplasm into the nucleus, as well as the excision of the DNA sequences flanked by loxP sites from a "reporter" gene are induced at doses of about ten times less than those necessary for Cre-ER^T.

15 With the aim of further increasing the sensitivity of the chimeric recombinase Cre-ER^{T2} to Tamoxifen, the inventors replaced the valine at position 400 in Cre-ER^{T2} with a glycine. This novel fusion protein which corresponds to the double mutant
20 M543A/L544A called Cre-ER^{T3} exhibits increased sensitivity to the synthetic antiestrogenic ligand such as Tam and OHT, without the recombinase activity of this protein being induced by the natural ligand estradiol.

25 The inventors have thus shown that for 10 times lower injected Tam doses, the recombinase activity in the cells of a transgenic Cre-ER^{T3} mouse is greater than that of a Cre-ER^{T2} mouse (see Example 5).

According to a preferred embodiment, the fusion
30 protein according to the invention is Cre-ER^{T2} whose

ER LBD exhibits the mutation G400V/M543A/L544A. According to another preferred embodiment, the fusion protein is Cre-ER^{T3}, whose ER LBD exhibits the mutation M543A/L544A.

5 One of the objects of the present invention is therefore to provide a Cre-ER fusion protein which exhibits mutations in the human ER α LBD which are preferably chosen from the mutations G521R, G400V, M543A and L544A, whose recombinase activity is not
10 induced by the natural ligands, and is highly induced by a small quantity of synthetic antiestrogenic ligand. Preferably, this fusion protein is Cre-ER^T, Cre-ER^{T2}, Cre-ER^{T3}. The present invention also relates to said fusion gene encoding said protein, said vector for
15 expressing said protein, as well as the corresponding host cell, and the corresponding transgenic animal, which expresses said fusion protein in a particular cell type, preferably the epidermis, the liver or the adipose tissue.

20 The Cre-ER fusion protein of the present invention therefore comprises all or part of a nuclear estrogen receptor and a recombinase protein whose activity is inducible more strongly by the binding of said receptor or of said ligand-binding domain (LBD) of
25 said receptor with a said antiestrogen than with a natural ligand. Said Cre-ER fusion protein makes it possible to carry out a recombination between loxP sites, in a cell of the organism of the invention, following treatment with an antiestrogen. In the
30 absence of treatment, or in the presence of

concentrations of ligands such as the natural estrogens of up to 10^{-6} M, no excision is observed. This system therefore makes it possible to release the recombinase activity of the chimeric protein at a given and chosen
5 moment. Said Cre-ER fusion protein may be expressed in cells containing loxP sites, without modifying the locus containing the loxP sites. The recombination at the level of the loxP sites takes place only after treatment with an antiestrogen such as Tam or OHT.
10 Furthermore, by expressing said Cre-ER fusion protein in an organism according to the invention, preferably an animal, under the control of a promoter with cellular specificity, it is possible to obtain recombination between loxP sites, specifically in these
15 cells.

The expression "synthetic ligand" is understood to mean any type of compound capable of binding to the nuclear estrogen receptor, and exhibiting agonist and/or antagonist activities, according to the species,
20 the tissue or the cell type. Preferably, and with no limitation being implied, the synthetic ligand according to the invention is endowed with antiestrogenic activity, it is preferably the antiestrogenic therapeutic agent Tamoxifen (Tam), but
25 also its metabolite 4-hydroxyTamoxifen (OHT). The antiestrogens ICI 164 384 and ICI 182 780 are also synthetic ligands according to the invention.

The present invention therefore provides a transgenic metazoan organism and more particularly a
30 transgenic animal, and in particular a transgenic

- 19 -

mouse: (i) in which at least one cell contains one or more chromosomal DNA sequences which are present in their natural chromatin context and are flanked (floxed) by loxP sites; (ii) which preferably expresses
5 a chimeric Cre recombinase in a tissue-specific manner in one or more cell types of the organism (iii) whose chimeric Cre recombinase activity is negligible, or even zero, in the presence of estrogen; (iv) whose chimeric recombinase activity is activated by low
10 concentrations of an antiestrogen (from 0.001 to 1 mg of Tamoxifen/mouse/day, for five days); (v) and finally whose Cre recombinase is capable of catalyzing, with an efficiency close to 100%, the site-specific targeted somatic recombination in the nucleus, in a natural
15 chromatin environment of the floxed DNA sequence(s).

The doses of synthetic ligand injected into the metazoan organism according to the invention are low. The term low is understood to mean quantities of less than or equal to 4 mg/adult mouse/day, preferably less
20 than or equal to 2 mg/adult mouse/day, in a preferred manner less than or equal to 1 mg/adult mouse/day. According to an even more preferred mode, this quantity may be less than or equal to 0.5 mg, 0.25 mg, 0.10 mg, 0.075 mg, 0.05 mg, 0.025 mg, 0.001 mg per adult mouse
25 and per day.

It is clearly understood that persons skilled in the art will be able to adjust these quantities, according to the organism, its weight and its age.

The efficiency of the targeted somatic
30 recombination is estimated by techniques known to

persons skilled in the art. This efficiency is estimated by the frequency of recombination events catalyzed by said recombinase. These events may be revealed by PCR or Southern Blotting; the recombination
5 frequency being estimated by taking the ratio of the representation of the various alleles in the cells of a tissue. The frequencies of the various alleles may be estimated by assaying the intensity of the corresponding bands on an electrophoresis gel of a
10 product of PCR amplification or of genomic DNA (Southern blotting).

The use of the PCR makes this method of estimation extremely sensitive and makes it possible to detect the presence of cells of the organism whose
15 genome has not undergone targeted site-specific recombination.

Another way of estimating the efficiency of the recombination may be carried out indirectly by immunohistochemistry, by analyzing the expression of
20 the gene sequence to be inactivated for example.

According to a preferred embodiment, said fusion protein is encoded by a fusion gene integrated into one or more of the chromosomes of said cell of said organism. According to another embodiment, the
25 fusion protein is encoded by a fusion gene integrated into an expression vector. The fusion gene according to the invention is introduced into the cell in the form of an expression vector or of one of its fragments. A "vector" is a replicon in which another polynucleotide
30 segment (i.e. the fusion gene) is attached, so as to

bring the replication and/or expression to the attached segment. The vector may be in particular a bacterial plasmid DNA, a cosmid, a phage DNA, a viral DNA or a minichromosome (BAC, YAC and the like). Such a vector
5 may be integrative, that is to say can integrate into the genome of the host cell or can exist in the form of an extrachromosomal replicon. When it exists in the form of an extrachromosomal replicon, the expression vector is capable of replicating autonomously. When it
10 is a fragment of an expression vector, preferably this fragment integrates into the cellular genome. The expression vector or one of its fragments comprises at least the fusion gene and a promoter or expression elements which make it possible to direct and control
15 the expression of said fusion protein in at least one cell of said organism.

The expression vector comprises, in addition, signals for initiation and termination of the transcription, as well as appropriate regions for
20 regulation of the transcription. These various control signals are chosen according to the cellular host used.

The expression elements controlling expression is understood to mean all the DNA sequences involved in the regulation of the gene expression, that is to say
25 the minimal promoter sequence, the upstream sequences, the activating sequences ("enhancers"), optionally the inhibitory sequences ("silencers"), the "insulator" sequences, and any other required sequence.

Preferably, the fusion gene is placed under the control of tissue-specific or cell-specific or ubiquitous expression elements.

The tissue-specific expression elements or
5 tissue-specific promoter regions are chosen from the promoters which make it possible to obtain a specific, and preferably high, expression in one or more cells, tissues, cell types or organs of the organism according to the invention. These promoter regions may be
10 heterologous or nonheterologous to the organism and may be naturally present or otherwise in the genome of the organism. By way of nonlimiting example of tissue-specific promoter regions, there may be mentioned the promoter regions of the genes:

- 15 - for cytokeratin, and more particular for cytokeratin 5 (K5) and cytokeratin 14 (K14), which directs the expression of the gene in the basal keratinocytes of the epidermis;
- for α -1-antitrypsin which directs the expression
20 of the gene in the hepatocytes;
- for the adipocyte fatty acid binding protein 2 (aP2) which directs the expression of the gene in the adipocytes.

According to a preferred embodiment, said
25 organism is characterized in that said promoter region is the cytokeratin 5 (K5) promoter region and said fusion gene Cre-ER^T.

According to a second preferred embodiment, said organism is characterized in that said promoter

region is the cytokeratin 5 (K5) promoter region and said fusion gene Cre-ER^{T2}.

According to a third preferred embodiment, said organism is characterized in that said promoter region
5 is the cytokeratin 5 (K5) promoter region and said fusion gene Cre-ER^{T3}.

According to a fourth preferred embodiment, said organism is characterized in that said promoter region is the cytokeratin 14 (K14) promoter region and
10 said fusion gene Cre-ER^T.

According to a fifth preferred embodiment, said organism is characterized in that said promoter region is the cytokeratin 14 (K14) promoter region and said fusion gene Cre-ER^{T2}.

15 According to a sixth preferred embodiment, said organism is characterized in that said promoter region is the cytokeratin 14 (K14) promoter region and said fusion gene Cre-ER^{T3}.

According to a seventh preferred embodiment,
20 said organism is characterized in that said promoter region is the α -1-antitrypsin promoter region and said fusion gene Cre-ER^T.

According to an eighth preferred embodiment, said organism is characterized in that said promoter
25 region is the α -1-antitrypsin promoter region and said fusion gene Cre-ER^{T2}.

According to a ninth preferred embodiment, said organism is characterized in that said promoter region is the α -1-antitrypsin promoter region and said fusion
30 gene Cre-ER^{T3}.

According to a tenth preferred embodiment, said organism is characterized in that said promoter region is the adipocyte fatty acid binding protein 2 (aP2) promoter region and said fusion gene Cre-ER^T.

5 According to an eleventh preferred embodiment, said organism is characterized in that said promoter region is the adipocyte fatty acid binding protein 2 (aP2) promoter region and said fusion gene Cre-ER^{T2}.

10 According to a twelfth preferred embodiment, said organism is characterized in that said promoter region is the adipocyte fatty acid binding protein 2 (aP2) promoter region and said fusion gene Cre-ER^{T3}.

15 According to a first embodiment, the organism according to the invention is characterized in that said fusion gene has the sequence SEQ ID No. 3 and encodes the Cre-ER^T protein having the sequence SEQ ID No. 4.

20 According to a second embodiment, the organism according to the invention is characterized in that said fusion gene encodes, of sequence SEQ ID No. 5 the fusion protein Cre-ER^{T2} having the sequence SEQ ID No. 6.

25 According to a third embodiment, the organism according to the invention is characterized in that said fusion gene encodes, of sequence SEQ ID No. 7 the fusion protein Cre-ER^{T3} having the sequence SEQ ID No. 8.

30 The article by Metzger and Feil (1999) gives by way of nonlimiting example (cf. table page 471) a list of tissue-specific promoter regions which are capable

of being used to direct the expression of the Cre protein in various tissues.

The tissue-specific promoter regions are more generally chosen from those which direct the expression
5 of the fusion protein in a physiological system, an organ, a tissue, a cell type or a particular cell; among which there may be nonexhaustively mentioned the nervous system in general, and in particular the brain, the cerebellum, the neurons, the motoneurons, the glial
10 cells, the Schwann cells, the hypophysis, the hypothalamus, the pituitary gland, the hippocampus and the cortex, the heart, the ventricular cardiomyocytes and the auricular cardiomyocytes, the lungs, the bones, the eyes, and more particularly the retina and the
15 crystalline lens, the skin and more particularly the dermis and the epidermis, the muscles, and more particularly the skeletal muscles, the cardiac muscle, the smooth muscles, the mammary gland, the gonads and more particularly the testes, the ovaries, the germ
20 cells, the oocytes, the oogonias, the spermatozoa, the spermatogonias and the spermatocytes, the kidney, the liver and in particular the hepatocytes, the spleen, the pancreas and in particular the Langerhans' cells and the β cells, the tongue, the esophagus, the
25 adipocytes, the vascular endothelial cells.

The ubiquitous expression elements or ubiquitous promoter regions are chosen from the promoter regions which make it possible to obtain expression, preferably high expression, in all, or at
30 least in a high proportion, of organs, or of tissues of

the organism according to the invention. These promoter regions may be heterologous or nonheterologous to the organism according to the invention. By way of nonlimiting example of "so called" ubiquitous promoter regions, there may be mentioned the cytomegalovirus (CMV) promoter (Schmidt et al., 1990) and the interferon-inducible promoter (Mx1) (Hug et al., 1998; Arnheiter et al., 1990). In addition, the expression elements, or promoter regions according to the invention, can ensure a constitutive or inducible control of the expression of the fusion gene. Among the elements ensuring inducible expression, there may be mentioned the eukaryotic promoter regions which are inducible by heavy metals (Mayo et al., 1982; Brinster et al., 1982; Seark et al., 1985), by heat shock (Nover et al., 1991), by hormones (Lee et al., 1981; Hynes et al., 1981; Klock et al., 1987; Israel et al., 1989), by interferon (Hug et al., 1998; Arnheiter et al., 1990). There may also be mentioned the inducible prokaryotic expression elements such as the *E. coli* Lac repressor system (LacR/operator/inducer) (Hu et al., 1987; Brown et al., 1987; Figge et al., 1988; Deuschle et al., 1990; Labow et al., 1990), the *E. coli* tetracycline resistance system (Gossen et al., 1992) (WO 94 04 672, EP 804 565).

In the case where the integration of the fusion gene is targeted by homologous recombination into the genome of the organism ("knock-in"), the fusion gene may be free of promoter regions or of expression

elements and may be placed under the control of a promoter region or of endogenous expression elements.

The recombinant DNA technologies used for the construction of the expression vector according to the invention are those known and commonly used by persons skilled in the art. Standard techniques are used for cloning, isolation of DNA, amplification and purification; the enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases are carried out according to the manufacturer's recommendations. These techniques and others are generally carried out according to Sambrook et al. (1989).

The vector according to the invention or the vector fragments may be introduced into the host cell by standard methods such as for example microinjection into a pronucleus, transfection by calcium phosphate precipitation, lipofection, electroporation, heat shock.

The fusion gene according to the invention preferably comprises in the 5' → 3' direction:

- a DNA fragment encoding the Cre recombinase of bacteriophage P1 or one of its variants;
- a DNA fragment of at least 45 nucleotides encoding at least either all or part of the D hinge region of a nuclear estrogen receptor, a region situated between the DNA-binding domain and the ligand-binding domain, or a peptide which is functionally equivalent to said D hinge region; and

- a DNA fragment encoding the ligand-binding domain (LBD) of a nuclear estrogen receptor or variants thereof, said fragment having at least one mutation conferring on LBD the capacity to respond to synthetic antiestrogens, but not to natural estrogenic agonists.

According to another embodiment of the invention, the fusion protein is directly introduced into the organism, or into a cell of the organism, it being possible for this introduction to be carried out by injection into a tissue or an organ in the case of an organism, or by microinjection in the case of a cell.

The DNA sequence of interest according to the invention is a gene or an intergenic sequence. According to a preferred embodiment of the invention, the DNA sequence of interest is a gene, it being possible for the function of the gene to be known or unknown. The study of an organism according to the invention exhibiting modification of a gene or of any other genomic region of unknown function makes it possible to contribute to the definition of the function of this gene or of this intergenic region. All the genes and intergenic regions of a metazoan organism are capable of being used in the context of the present invention; more particularly, there may be mentioned the RXR_{α} , RXR_{β} , RXR_{γ} , RAR_{α} , RAR_{β} , RAR_{γ} , $SNF2_{\beta}$ genes. The expression "DNA sequence of interest" naturally belonging to the genome of said organism, or DNA sequence of interest in its natural chromatin

environment, is understood to mean an endogenous DNA sequence, such as an endogenous gene, present in the genome at its natural locus (loci).

According to a preferred embodiment of the invention, the organism according to the invention is an animal, in particular a mouse, characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the cytokeratin K5 promoter;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

According to a second preferred embodiment of the invention, the organism according to the invention is characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the cytokeratin K14 promoter;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

According to a third preferred embodiment of the invention, the organism according to the invention

is characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the adipocyte fatty acid binding protein 2 (aP2) promoter;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

According to a fourth preferred embodiment of the invention, the organism according to the invention is characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the α -1-antitrypsin promoter;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

The present invention also relates to methods of preparing a metazoan organism according to the invention.

A first method of preparation consists in the steps of:

- a) obtaining an embryonic stem (ES) cell modified by insertion of site(s) of recognition for said

- 31 -

recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;

- 5 b) introducing said modified embryonic stem cell into an embryo of said organism;
- c) developing said embryo up to the stage of a fertile adult organism;
- d) crossing said fertile adult organism with a transgenic organism in which at least one of the
- 10 cells expresses said fusion protein and obtaining the progeny derived from said crossing; and
- e) optionally, selecting, among said progeny, said metazoan organism.

A second method of preparation consists in the

15 steps of:

- a) obtaining a somatic cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous
- 20 recombination;
- b) transferring the nucleus of said modified somatic cell into the cytoplasm of an enucleated recipient oocyte;
- c) developing the embryo obtained in step b) up to
- 25 the stage of a fertile adult organism;
- d) crossing said fertile adult organism with a transgenic organism in which at least one of the cells expresses said fusion protein and obtaining the progeny derived from said crossing; and

- e) optionally, selecting, among said progeny, said metazoan organism.

The expression transfer of the nucleus or nuclear transfer, for the purposes of the present invention, is understood to mean the transfer of nucleus of a vertebrate live donor cell, of an adult organism or at the fetal stage, into the cytoplasm of an enucleated recipient cell of the same species or of a different species. The transferred nucleus is reprogrammed to direct the development of the cloned embryos which may then be transferred into carrier females to produce the fetuses and the neonates, or used to produce cells of the internal cellular mass in culture. Various nuclear cloning techniques are capable of being used; among these, there may be mentioned those which have been the subject of patent applications WO 95 17500, WO 97 07668, WO 97 07669, WO 98 30683, WO 99 01163, WO 99 37143.

A third method of preparation consists in the steps of:

- a) obtaining an embryonic stem (ES) cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;
- b) introducing said modified embryonic stem cell into an embryo of said organism;
- c) developing said embryo; and

- 33 -

- d) introducing said fusion protein into at least one cell of said embryo or of the organism obtained from the development of said embryo.

A fourth method of preparation consists in the
5 steps of:

- a) obtaining a somatic cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous
10 recombination;
- b) transferring the nucleus of said modified somatic cell into the cytoplasm of an enucleated recipient oocyte;
- c) developing said embryo; and
- 15 d) introducing said fusion protein into at least one cell of said embryo or of said organism obtained from the development of said embryo.

The insertion of the sites of recognition specific for the recombinase protein, in particular of
20 the loxP site(s) for the Cre recombinase, into the DNA sequence of interest is preferably carried out by homologous recombination of the gene comprising said DNA fragment to be excised or inverted (two loxP sites) or respectively inserted or translocated (one loxP
25 site) with a said modified gene comprising said DNA fragment to be excised flanked in 5' and/or 3' by said recombinase recognition site(s) according to the desired application, in particular the loxP sites.

To do this, the modified DNA fragment of
30 interest may be integrated by homologous recombination

into the genome of the cells of said organism before, at the same time, or after the step of introducing the fusion protein or of a transfer vector, or of a vector for expressing the fusion protein. Preferably, the DNA
5 fragment of interest is introduced into pluripotent embryonic cells (ES cells) by the appropriate technique, such as for example electroporation, or the use of retroviral vectors, calcium phosphate precipitation, lipofection.

10 The DNA constructs intended for homologous recombination will comprise at least a portion of the DNA sequence of interest, in particular of the gene or of the intergenic sequence of interest into which will be introduced the desired genetic modification(s), such
15 as the introduction of at least one recombinase recognition site, and which will include regions of homology with the target locus. For facilitated use, positive and/or negative selectable markers (for example the neo gene conferring resistance to the
20 antibiotic G418) may be introduced. The selectable marker used to make it possible to identify the homologous recombination events may be disruptive, and may be eliminated, if necessary, if it is itself flanked by recombinase recognition sites such as the
25 loxP (or FRT) sites. This makes it possible to obtain mice in which the sole modification at the level of the modified locus is the insertion of recognition sites such as loxP.

 The metazoan organisms obtained by the methods
30 of preparation presented above can then be treated with

a synthetic ligand endowed with antiestrogenic activity such as Tam and OHT. In the various methods and uses of the invention, the bringing of said cells of said organism into contact with said synthetic ligand is
5 carried out by administration by the oral or topical route, or by injection and in particular, by intravenous, intramuscular, intraspinal, intracerebral, intraperitoneal injection. In the case of an embryo, a fetus or a neonate before weaning, the treatment with
10 the synthetic ligand may be carried out by administration to the mother. When this involves cells in culture derived from said organism, said synthetic ligand is preferably added to the culture medium, or injected into said cell. This treatment or this
15 bringing into contact makes it possible to inactivate or to modify a gene or an intergenic sequence of interest at a determined moment (temporal control) in a given tissue (spatial control), and thus to make it possible to study the function of this gene or of this
20 sequence at various periods during development or post-natally. This is particularly advantageous for studying at the adult stage genes which are essential for the normal progress of embryonic development and whose inactivation is lethal *in utero* or perinatally.

25 Another object of the present invention is therefore to provide a method of conditional recombination, in particular excision, insertion, inversion, translocation, at the level of the DNA sequence of interest into which there is (are) inserted
30 one or more sites of recognition for said recombinase

protein, said DNA sequence of interest being located in one or more of the chromosomes of said genome of said cell of said organism according to the invention, characterized in that it comprises the steps of:

- 5 (i) bringing at least one cell of said organism into contact with a synthetic ligand endowed with antiestrogenic activity;
- (ii) inducing the activity of the recombinase of said fusion protein by said synthetic ligand endowed
- 10 with antiestrogenic activity.

The present invention therefore provides a method of conditional deletion of a DNA fragment in which a method of excision according to the invention is used and in which said DNA fragment(s) to be excised

15 is (are) flanked by two recombinase protein recognition sites oriented as a direct repeat. In particular, said DNA fragment may be chosen such that the excision of said DNA fragment has the effect of inactivating said gene.

20 The present invention also provides a method of obtaining a metazoan organism, with the exception of humans, in which at least one cell possesses an allele of a gene of interest inactivated by a method of conditional deletion and in which the other allele of

25 said gene of interest possesses a mutation, said method being characterized in that it comprises the steps of:

- a) obtaining a metazoan organism in which at least one cell of the germ line comprises said mutation in one of the alleles of said gene of interest;

- b) crossing said organism obtained in step a) with an organism according to the invention;
- c) selecting a progeny whose genome comprises a gene of interest in which one of the alleles possesses a mutation and the other allele possesses at least two recombinase protein recognition sites oriented as a direct repeat; and
- d) using the method of conditional deletion, according to the invention, of the DNA fragment of said allele of said gene of interest which is flanked by at least two recombinase protein recognition sites oriented as a direct repeat; and
- e) obtaining said metazoan organism in which the genome of at least one cell comprises said gene of interest in which one allele is inactivated, the other allele possesses a somatic, preferably limited, mutation and preferably in exon and/or regulatory sequences.

Such a method makes it possible to study and analyze the biological function of mutations other than deletions, and more particularly of the mutations observed in genes whose dysfunction causes a recessive genetic pathological condition. This method is therefore particularly suited to the obtaining of transgenic animal models of recessively transmitted human genetic pathological conditions, the animal model being preferably a murine model.

The mutations are preferably point or limited mutations in exons or regulatory sequences such as insertions, deletions, substitutions.

According to a preferred embodiment of the method of recombination, and of the method of conditional deletion or of the method of obtaining a metazoan organism according to the invention, the
5 recombinase protein specific recognition sites are loxP sites and said recombinase protein is the Cre protein of the bacteriophage P1, or one of its variants.

The organisms capable of being obtained using the various methods above are also included in the
10 scope of the invention. These organisms are preferably animals, and in a preferred manner rodents such as rats and mice, preferably mice.

Preferably, the subject of the invention is a transgenic mouse K5-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α gene may
15 be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal
20 keratinocytes and/or an inflammatory reaction of the skin.

Preferably, the subject of the invention is a transgenic mouse K5-Cre-ER^{T2}/RXR_α^{L2/L2} whose RXR_α gene may
25 be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the keratinocytes and/or an inflammatory reaction of the skin.

Preferably, the subject of the invention is a transgenic mouse K5-Cre-ER^{T3}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method
5 following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal keratinocytes and/or an inflammatory reaction of the skin.

10 Preferably, the subject of the invention is a transgenic mouse K14-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed
15 with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the keratinocytes and/or an inflammatory reaction of the skin.

Preferably, the subject of the invention is a transgenic mouse K14-Cre-ER^{T2}/RXR_α^{L2/L2} whose RXR_α gene
20 may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the
25 keratinocytes and/or an inflammatory reaction of the skin.

Preferably, the subject of the invention is a transgenic mouse K14-Cre-ER^{T3}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal
30 keratinocytes of the epidermis using a conditional

deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the keratinocytes and/or an inflammatory reaction of the skin.

Preferably, the subject of the invention is a transgenic mouse $\alpha\text{AT-Cre-ER}^T/\text{RXR}_\alpha^{L2/L2}$ whose RXR_α gene may be selectively inactivated in the hepatocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

Preferably, the subject of the invention is a transgenic mouse $\alpha\text{AT-Cre-ER}^{T2}/\text{RXR}_\alpha^{L2/L2}$ whose RXR_α gene may be selectively inactivated in the hepatocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

Preferably, the subject of the invention is a transgenic mouse $\alpha\text{AT-Cre-ER}^{T3}/\text{RXR}_\alpha^{L2/L2}$ whose RXR_α gene may be selectively inactivated in the hepatocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

Preferably, the subject of the invention is a transgenic mouse $\text{aP2-Cre-ER}^T/\text{RXR}_\alpha^{L2/L2}$ whose RXR_α gene may be selectively inactivated in the adipocytes using a conditional deletion method following treatment with a

synthetic ligand endowed with antiestrogenic activity, causing in said mouse alteration of the metabolism of the lipids in the adipocytes and/or diabetes.

Preferably, the subject of the invention is a
5 transgenic mouse $\text{aP2-Cre-ER}^{\text{T2}}/\text{RXR}_{\alpha}^{\text{L2/L2}}$ whose RXR_{α} gene may be selectively inactivated in the adipocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alteration of the
10 metabolism of the lipids in the adipocytes and/or diabetes.

Preferably, the subject of the invention is a transgenic mouse $\text{aP2-Cre-ER}^{\text{T3}}/\text{RXR}_{\alpha}^{\text{L2/L2}}$ whose RXR_{α} gene may be selectively inactivated in the adipocytes using
15 a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alteration of the metabolism of the lipids in the adipocytes and/or diabetes.

20 Preferably, said RXR_{α} gene of said mouse is inactivated using a method according to the invention.

The present invention and in particular the metazoan organism and the cells derived therefrom are particularly useful for analyzing and studying the
25 biological function of a DNA sequence of interest, whether it is a gene or an intergenic sequence in its natural chromatin environment. That is the reason why it is also within the scope of the present invention to provide a method of analyzing or studying the
30 biological function of a DNA sequence of interest, in

particular of a gene or an intergenic sequence, characterized in that it comprises the steps of:

- (i) bringing an organism according to the invention or cells isolated from said organism into
5 contact with a synthetic ligand endowed with antiestrogenic activity;
- (ii) optionally inducing the expression of said fusion protein;
- (iii) revealing the recombination event catalyzed by
10 the recombinase activity of said fusion protein;
- (iv) biochemical and/or physiological and/or phenotypic and/or behavioral study or analysis of said cell or of said organism.

15 The phenotypic and behavioral analyses of the organism according to the invention before or after induction of the somatic recombination are carried out using techniques known to persons skilled in the art.

The subject of the present invention is also
20 the use of an organism according to the invention or of cells derived from said organism for carrying out a spatiotemporally controlled site-specific recombination of said DNA sequence of interest in its natural chromatin environment, with an efficiency of at least
25 85%, 90%, 92%, 94%, 95%, 96%, 97%, 98%, 99%, 100%, in the presence of synthetic ligand in the cells of said organism expressing said fusion protein and with an efficiency at least less than 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, 0.01%, or zero in the absence of synthetic ligand

or in the presence of a natural ligand in the cells of said organism expressing said fusion protein.

According to a preferred mode, said recombination is carried out in the epidermis, and more
5 precisely in keratinocytes, in the adipocytes, in the malanocytes or in the hepatocytes.

The subject of the present invention is also its method of screening compounds capable of being used as a medicament for the preventive and/or curative
10 treatment of pathological conditions associated with alteration of the expression (in the case, in particular, of an intergenic DNA sequence) and/or of the function (in the case of a gene DNA sequence) of said DNA sequence of interest, characterized in that it
15 comprises the step of administering said compound to an organism according to the invention.

This organism may thus be used for the screening of compounds capable of constituting an active ingredient of a medicament intended for the
20 treatment of pathological conditions associated with alteration of the expression and/or of the function of said DNA sequence of interest.

Thus, the object of the present invention is to provide a method of screening compounds capable of
25 being used as a medicament for the preventive and/or curative treatment of alopecia and/or of hyperproliferation of the keratinocytes and/or of inflammatory reactions of the skin, characterized in that it comprises the step of administering said
30 compound to a mouse according to the invention.

The object of the present invention is also to provide a method of screening compounds capable of being used as a medicament for promoting, in particular, hepatic regeneration, characterized in that
5 it comprises the step of administering said compound to a mouse according to the invention.

The object of the present invention is also to provide a method of screening compounds capable of being used as a medicament for the preventive and/or
10 curative treatment of diabetes and/or for the treatment of the alteration of the metabolism of lipids, in particular of obesity, characterized in that it comprises the step of administering said compound to a mouse according to the invention.

15 The object of the present invention is also to provide a method of screening compound capable of being used as a medicament for the preventive and/or curative treatment of skin cancers, more specifically papillomas and melanomas at various stages of development,
20 characterized in that it comprises the step of administering said compound to a mouse according to the invention.

Other characteristics and advantages of the present invention will emerge in the light of the
25 examples which follow.

Figure 1: Inactivation of the RXR α gene in the epidermis of adult mice mediated by Cre-ER^T and Cre-ER^{T2}, and induced by Tamoxifen.

Figure 1A: Schematic representation of the wild-type genomic locus of RXR_α (+), of the floxed L2 RXR_α allele, of the L^- RXR_α allele obtained after Cre-mediated excision of exon 4 (encoding the DNA-binding domain),
5 and of the RXR_α (-) null allele (Kastner et al., 1994). The black boxes indicate the exons (E2 - E4). The enzymatic restriction sites and the position of the probe X4 are indicated. The sizes of the BamHI fragments are indicated in kilobases (kb): BamHI; C,
10 ClaI; E, EcoRI H, HindIII; S, SpeI; X, XbaI. The arrow tips in the L2 and L^- alleles indicate the Lox P sites.

Figure 1B: The obtaining of $\text{RXR}_\alpha^{\text{L}^-}$ alleles mediated by K5-Cre-ER^T induced by Tamoxifen is illustrated by
15 Southern blot analysis of the DNA isolated from the epidermis, six weeks (lanes 1 to 3) or twelve weeks (lanes 4 to 6) after the first injection of Tamoxifen (1 mg) (AFT : "after first Tamoxifen treatment"). The genotype of the mice is as indicated, and the fragments
20 digested with BamHI corresponding to the RXR_α (+), L2, L^- , (-) alleles are described.

Figure 1C: Tissue selectivity of the inactivation of RXR_α mediated by Cre-ER^T. The wild-type alleles (+), L2, L^- are identified by PCR from DNA extracted from
25 various K5-Cre-ER^{T(tg/0)}/ $\text{RXR}_\alpha^{\text{L2/+}}$ mouse organs, twelve weeks after the first Tamoxifen treatment.

Figure 1D: The obtaining of L^- RXR_α alleles mediated by K14-Cre-ER^{T2} induced by Tamoxifen in the epidermis of adult mice is illustrated by PCR analysis of the genomic DNA extract from the epidermis (E) and from the dermis (D) isolated two weeks after the first injection of Tamoxifen (0.1 mg) (+) or of the vehicle (oil) (-). The genotype of the mice is indicated and the PCR fragments corresponding to the RXR_α (+), L2 and L- alleles are indicated.

10

Figure 2: Abnormalities generated by the inactivation of RXR_α mediated by K5-Cre-ER^T and K14-Cre-ER^{T2} induced by Tamoxifen, in the skin of adult mice.

15 **Figure 2A:** Ventral view of a "mutant" (mt) female mouse K5-Cre-ER^{T(tg/0)}/ $RXR_\alpha^{L2/-}$ (on the left) and of a "control" (ct) female mouse K5-Cre-ER^{T(tg/0)}/ $RXR_\alpha^{L2/+}$ (on the right), sixteen weeks after the first Tamoxifen injection (1 mg of Tamoxifen/injection).

20

Figure 2B: Dorsal view of the same animals.

Figure 2C: Magnification of the ventral region of the "mutant" (mt) female mouse K5-Cre-ER^{T(tg/0)}/ $RXR_\alpha^{L2/-}$, with the arrow indicating one of the cysts visible under the surface of the skin.

25

Figure 2D: Dorsal view of the "mutant" (mt) female mouse K5-Cre-ER^{T(tg/0)}/ $RXR_\alpha^{L2/-}$, twenty-eight weeks after

- 47 -

the start of Tamoxifen treatment. The arrow indicates a minor skin lesion.

Figures 2E-H: Histological analysis. Histological sections 2 μm thick of the ventral skin of "control" (E and G) and "mutant" (F and H) mice, sixteen weeks after the start of the treatment. Hair follicles (hf), utriculi (u) and dermal cysts (dc) are indicated. The arrow tips in (H) indicate the Langerhan's cells, whose number is increased several fold in the epidermis of the mutant mice. Increased cellularity can be noted in (F) and in (H) in the dermis under the hyperproliferated interfollicular epidermis. Scale (in E) E and F = 60 μm ; G and H = 12 μm .

15

Figures 2I, J: Immunohistochemistry of keratin 6 (K6) on sections of skin of "control" mice (I) and of "mutant" mice (J), sixteen weeks after the first Tamoxifen treatment. The red color corresponds to the labeling with the antibody directed against K6, and the Cyan color corresponds to staining with DAPI. K6 is normally expressed in the outer root sheath of the hair follicle (hf) but not in the epidermis (I) and is abnormally expressed in the hyperproliferative epidermis of "mutant" mice (J). Scale (in I) I = 25 μm . The arrows in E to J indicate the dermis-epidermis junction.

Figures 2K, L: Appearance of the skin of a mutant female mouse K14-Cre-ER^{T2(tg/0)}/RXR α ^{L2/L2}. (K) magnification

30

on the ventral region, sixteen weeks after the start of Tamoxifen treatment (0.1 mg per injection); the white arrow indicates a cyst and the black arrow indicates a utriculus containing melanosomes. (L) dorsal view of the same mutant. The arrow indicates one of the skin lesions associated with a hairless region.

Figure 3: Similarities and differences between the skin abnormalities present in a double "mutant" mouse K5-Cre-ER^{T(tg/0)}/RXR_α^{L2/L2}/RXR_β^{-/-} induced with Tamoxifen and a "null" VDR mouse.

Figures 3A, E: K5-Cre-ER^{T(tg/0)}/RXR_α^{L2/L2}/RXR_β^{-/-} mouse eighteen weeks after the first treatment with Tamoxifen (1 mg of Tamoxifen per injection) (A) and fourteen-week old VDR^{-/-} mouse (E). The arrows in (A) indicate the skin lesions. (B, C) and (F, G): histological analysis of 2 μm thick sections of dorsal skin of the animals presented in (A) and (E) respectively. Scale (in G): B and F = 60 μm; C and G = 12 μm.

Figures 3D, H: Immunohistochemistry of keratin 6 (K6) on mouse skin sections removed from the animals presented in (A) and (E) respectively. It should be noted that K6 is expressed in the utriculus, but not in the epidermis of the skin of the VDR^{-/-} mice, whereas K6 is expressed in the entire hyperproliferative epidermis of the "double mutant" mice RXR_α/RXR_β (D). Utriculus (U), dermal cyst (dc). The arrows in (B-H) indicate the

dermis-epidermis junction. Scale (in H): D and H =
25 μ m.

Figure 4: Selected targeted inactivation of the SNF2 β
5 gene in the epidermis of adult mice.

Figure 4A: Schematic representation of the wild-type
SNF2 β alleles (+), L3, L2+ and L-.

The size of the DNA segments revealed by the 5'
10 probe, after enzymatic digestion of the genomic DNA by
BamHI is indicated. The L3 allele in the ES cells was
obtained by homologous recombination using a strategy
similar to that described for the F9 cells in
Sumi-Chinose *et al.*, 1997.

15

Figure 4B

Eight-week old K14-Cre-ER^{T2(tg/0)}/SNF2 β ^{L2/L2} and
K14-Cre-ER^{T2(0/0)}/SNF2 β ^{L2/L2} were treated with Tamoxifen
(Tam) for five days at the rate of 0.1 mg/day or with
20 oil (-). Tail biopsis were taken, the dermis and the
epidermis separated, the genomic DNA prepared, digested
with BamHI, separated by agarose gel electrophoresis,
and transferred onto nylon membranes, which were
hybridized with the radiolabeled 5' probe.
25 Autoradiographs are presented.

Figure 5: Selective targeted inactivation of the RXR_{α} gene in the adipocytes.

Transgenic $aP2-Cre-ER^{T2/(tg/0)}$ mice expressing $Cre-ER^{T2}$ under the control of the murine $aP2$ promoter, which is selectively active in the adipocytes (Ross et al., 1990), were crossed with $RXR_{\alpha}^{L2/+}$ mice so as to produce $aP2-Cre-ER^{T2(tg/0)}/RXR_{\alpha}^{L2/L2}$ mice.

These mice were then crossed with $RXR_{\alpha}^{+/-}$ mice (Kastner et al., 1994) so as to produce $aP2-Cre-ER^{T2(tg/0)}/RXR_{\alpha}^{L2/-}$ mice. Such four week old mice, were treated (+) or not (-) with Tamoxifen (1 mg/day) for five days, and the adipose tissue collected one month after the last injection of Tamoxifen. The DNA was extracted from the adipose tissue, or after separation of the adipocytes (80% purified adipocytes) from the connective tissue and blood vessels (nonadipocytes). After digestion with BamHI, the alleles of RXR_{α} were analyzed by Southern blotting. An autoradiogram is presented. No excision is observed in the purified adipocytes or the adipose tissue of mice not treated with Tamoxifen (Tam). On the other hand, in the adipose tissue and the adipocytes of mice treated with Tamoxifen, excision is observed which is characterized by the appearance of a band at 8.5 kb corresponding to the L^{-} allele.

Figure 6: Phenotypic analysis of mice after conditional somatic mutagenesis of RXR_{α} in the adipocytes.

The weight of the control mice $\alpha P2\text{-Cre-ER}^{T2tg/0}/RXR_{\alpha}^{L2/+}$ (CT) and $\alpha P2\text{-Cre-ER}^{T2(tg/0)}/RXR_{\alpha}^{L2/-}$ (mutants; KO) was determined once per week. (A) each group of animals was composed of 10 to 15 males. The animals were fed either with the normal food (AN) or food enriched with fat and with glucose (AR). (B) The weight of the subcutaneous adipose tissue of 6-month old CT and KO mice, fed with AN or AR. (C) 10 μm cryosections of subcutaneous adipose tissue of 6-month old CT (a and c) and KO (b and d) mice, fed with AN (a and b) or AR (c and d). Scale; 160 μm . The levels of triglycerides (D), glucose (E) and insulin (F) were determined on the serum of 4- to 5-month old CT and KO animals, fed with AN or AR. The glucose assay was carried out after starving the animals for 12 hours, $*p < 0.05$.

Figure 7: Selective targeted inactivation of the RXR_{α} gene in the murine hepatocytes.

To invalidate the RXR_{α} gene selectively in the hepatocytes, $\alpha\text{AT-Cre-ER}^{T(tg/0)}$ mice which express Cre-ER^T under the control of the promoter of the α -1-antitrypsin (αAT) gene in about 50% of the hepatocytes (Imai et al., 2000) were crossed with $RXR_{\alpha}^{L2/L2}$ mice so as to produce $\alpha\text{AT-Cre-ER}^{T(tg/0)}/RXR_{\alpha}^{L2/L2}$ mice. Such three month old mice were treated with Tamoxifen (1 mg/day) for five days, and the heart and liver removed seven days after the first injection of Tamoxifen (day 7), from one and three animals

respectively. The heart and the liver were also removed from one and three animals respectively of the same genotype without treatment with Tamoxifen (Day 0). The DNA was extracted from these tissues, and after
5 digestion with BamHI, the alleles of RXR α were analyzed by Southern blotting. An autoradiogram is presented. No excision is observed in the heart at day 0 or 7, or in the liver at day 0. Conversely, excision is observed in the liver of mice seven days after the treatment with
10 Tamoxifen; this excision is materialized by the appearance of a band at 8.5 kb on the autoradiogram, corresponding to the L⁻ allele.

Figure 8: Site-specific recombination of RXR α in the
15 liver.

The excision of DNA segments mediated by Cre-ER^T was determined by "Southern blotting", carried out with DNA extract from the heart and the liver of α AT-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2} mice, removed from various
20 animals before the injections of tamoxifen (day 0) or 7, 30 or 90 days after the last injection of Tamoxifen.

Lanes 15-17 correspond to the DNA isolated from the livers of animals 7 days after partial hepatectomy (HP), carried out after the injection of Tamoxifen. The
25 position of the RXR α L2 and L⁻ alleles is indicated.

Figure 9: Expression of the Cre-ER^{T2} and Cre-ER^{T3} recombinases in the basal layer of the epidermis.

Immunohistochemistry of the chimeric Cre
30 recombinase on sections of the epidermis of K5-Cre-ER^{T2}

and K5-Cre-ER^{T3} mice, treated either with 1 mg or 0.1 mg of OHT (4-hydroxyTamoxifen). The red color corresponds to labeling with the anti-Cre antibody directed against the Cre recombinase protein, and the Cyan color corresponds to DAPI staining the cellular nuclei. Cre-ER^{T2} and Cre-ER^{T3} are located in the cellular nuclei of the basal layer. (It should be noted that the superposition of the red and cyan colors results in a violet coloration).

10

Figure 10: Nuclear translocation of Cre-ER^{T2} and Cre-ER^{T3} following a two-day treatment at various OHT doses.

Immunohistochemistry of the chimeric Cre recombinase on sections of the epidermis of K5-Cre-ER^{T2} and K5-Cre-ER^{T3} mice treated either with 0.1 mg, 0.01 mg or 0.001 mg of OHT and analyzed two days (D 2) after the start of the treatment (D 0). The red color corresponds to labeling with the anti-Cre antibody directed against the Cre protein, and the Cyan color corresponds to DAPI. Cre-ER^{T2} and Cre-ER^{T3} are located in the cellular nuclei of the basal layer. Cre-ER^{T3} is present in a larger fraction thereof than Cre-ER^{T2}, at the various OHT doses. At the dose of 0.001 mg, about 1/3 of the nuclei are strongly labeled with anti-Cre antibodies in the basal layer of the epidermis of K5-Cre-ER^{T3} mice, whereas no positive nucleus is observed in the skin of K5-Cre-ER^{T2} mice. (It should be noted that this superposition of the red and cyan colors results in a violet coloration).

Figure 11: Comparison of the expression of β -galactosidase in the epidermis of the tail of K5-Cre-ER^{T2(tg/0)}/Rosa^{fl/+} and K5-Cre-ER^{T3(tg/0)}/Rosa^{fl/+} induced by various OHT doses.

Activity of β -galactosidase on sections of the epidermis of the tail of K5-Cre-ER^{T2(tg/0)}/Rosa^{fl/+} and K5-Cre-ER^{T3(tg/0)}/Rosa^{fl/+} mice treated with 1 mg, 0.1 mg, 0.01 mg and 0.001 mg of OHT. The analyses are carried out on the fifteenth day after the start of the treatment (D0).

The excision levels induced by 1 and 0.1 mg of OHT are similar for the two lines; on the other hand, the excision is more efficient in the K5-Cre-ER^{T3} than in the K5-Cre-ER^{T2} mice at the doses of 0.01 and 0.001 mg of OHT.

Figure 12 : Rate of papilloma formation in RXR α ^{ep-/-} mice.

A. Timing of Tam-induced RXR α ablation in epidermal keratinocytes, and DMBA/TPA tumorigenesis. Tam-treatment (0.1 mg for 5 consecutive days) was performed either 16 days before (bar a) or 7 weeks after (bar b) topical DMBA application. TPA was topically applied twice a week (arrows) for up to 30 weeks. B. Papillomas in DMBA/TPA-treated RXR α ^{ep-/-} mice. The number of papillomas induced by the DMBA/TPA treatment was determined macroscopically in 6 CT and 6 RXR α ^{ep-/-} mice, and plotted versus the number of weeks after the start

of the carcinogenic treatment. Values are expressed as mean $\pm$ SEM. C. Dorsal view of CT (K14-Cre-ER^{T2(0/0)}/RXR α ^{L2/L2}) (left) and RXR α ^{ep-/-} (right) mice 25 weeks after the start of the DMBA/TPA treatment. D. Length distribution of papillomas in CT and RXR α ^{ep-/-} mice. The number of tumors of a given length was determined on 6 CT and 6 RXR α ^{ep-/-} mice, 30 weeks after the start of the DMBA/TPA treatment. Values are expressed as mean $\pm$ SEM.

10

Figure 13 : Histological analysis of skin tumors induced by DMBA/TPA treatment.

A. Representative hematoxylin and eosin stained 5 μ m-thick paraffin sections from CT biopsies taken 25 (a - b) and 30 (c) weeks after the start of the DMBA/TPA treatment. B. Representative hematoxylin and eosin stained 5 μ m-thick paraffin sections from RXR α ^{ep-/-} biopsies taken 25 (a-c) and 30 (d-i) weeks after the start of the DMBA/TPA treatment. (a), atypical hyperplasia ; (b), in situ carcinoma ; (c), focal carcinoma; (d), (e), (f) and (g), advanced grade I, II, III and IV SCC, respectively ; (h), spindle cell carcinoma ; (i), basal cell carcinoma. Scale bar 33 μ m. C. Table of histological analysis of paraffin sections (5 μ m-thick) of biopsies of 8-16 mm tumors from CT and RXR α ^{ep-/-} mice, 25 and 30 weeks after the start of the DMBA/TPA treatment.

Figure 14 : Melanocytic growths induced by DMBA treatment of CT and $\text{RXR}\alpha^{\text{ep-/-}}$ mice.

A. Dorsal view of CT ($\text{K14-Cre-ER}^{\text{T2(0/0)}}$ / $\text{RXR}\alpha^{\text{L2/L2}}$) (left) and $\text{RXR}\alpha^{\text{ep-/-}}$ (right) mice 25 weeks after DMBA treatment. Arrows point to some of the melanocytic growths. B. Number of melanocytic growths in skin of CT and $\text{RXR}\alpha^{\text{ep-/-}}$ mice 30 weeks after DMBA and DMBA/TPA treatments, as indicated. Values are expressed as mean $\pm$ SEM ($n = 6$). *, $p < 0.05$; ***, $p < 0.001$. C. Size distribution of melanocytic growths in CT and $\text{RXR}\alpha^{\text{ep-/-}}$ mice. The number of 1 - 2 and 2 - 4 mm melanocytic growths was determined on CT and $\text{RXR}\alpha^{\text{ep-/-}}$ mice, 30 weeks after DMBA and DMBA/TPA treatments, as indicated. Values are expressed as mean $\pm$ SEM ($n = 6$).

Figure 15 : Malignant melanomas in lymph nodes of DMBA-treated $\text{RXR}\alpha^{\text{ep-/-}}$ mice.

Photograph of subiliac lymph nodes from CT (a and e) and $\text{RXR}\alpha^{\text{ep-/-}}$ (b and f) mice taken 30 weeks after DMBA and DMBA/TPA treatment, respectively. Hematoxylin and eosin stained 5 μm paraffin sections of CT (c and g) and $\text{RXR}\alpha^{\text{ep-/-}}$ (d and h) mice after DMBA and DMBA/TPA treatments, respectively. Scale bar, 33 μm .

EXAMPLES1) MATERIALS AND METHODS

5 1.1 - Transgenic lines

The mouse lines $RXR_{\alpha}^{+/-}$, $VDR^{-/-}$ and K5-Cre-ER^T have been previously described in Yoshizawa et al. (1997), Kastner et al. (1994) and Indra et al. (1999).

10 The transgene K14-Cre-ER^{T2} was constructed by replacing the K5 promoter region of the vector pK5-Cre-ER^{T2} (Indra et al., 1999) with the SalI DNA fragment of the promoter/enhancer region of 2 kb of the K14 human keratin gene, isolated from Phr2 (gift from
15 S. Werner). The transgenic mice were generated in accordance with the article by Indra et al. (1999).

The transgene aP2-Cre-ER^{T2} was constructed as follows: a 5.4 kb fragment containing the aP2 promoter was amplified by PCR from mouse genomic DNA with the
20 aid of the LA-PCR kit (Perkin-Elmer, New Jersey), with the oligonucleotides

5'-ATACGCGGCCGCGAATTCCAGCAGGAATCAGGTAGCT-3' (Sequence ID No. 13) and

5'-ATAGCGCCGGCGCTGCAGCACAGGAGGGTGCTATGAG-3' (Sequence
25 ID No. 14). After having made the ends of this fragment blunt following the action of T4 polymerase, it was cloned at the level of the SalI site of pGS-Cre-ER^{T2} (Indra et al., 1999), whose ends have also been made blunt following the action of T4 polymerase. The 8.3 kb
30 NotI fragment was isolated from this plasmid, purified

and injected into the F1 zygotes (C57BL/6 × SJL) at the concentration of 4 ng/ml, and the mice carrying the transgene aP2-Cre-ER^{T2} were identified according to Feil *et al.*, (1996) and Imai *et al.*, (2000).

5

1.2 - Genotyping of the RXR_α alleles

The genomic DNA is isolated from tissues according to the protocol described in the article by Indra *et al.* (1999).

10 The epidermis is separated from the dermis after treating the skin of the tail with the enzyme dispase (4 mg/ml in PBS, GIBCO-BRL) for 1 to 2 hours at room temperature. The genotyping of the RXR_α cells is carried out by PCR (polymerase chain reaction) using
15 the primers ZO 243 (5'-TCC TTC ACC AAG CAC ATC TG-3') (SEQ ID No. 9) (located in exon 3) and ZO 244 (5'-TGC AGC CCT CAC AAC TGT AT-3') (SEQ ID No. 10) (located in exon 4) for the L2 and wild-type (+) alleles; these amplification reactions generate, for
20 the L2 allele, a fragment of 700 bp and for the wild-type (+) allele a fragment of 650 bp.

 The primers ZO 243 and UD 196 (5'-CAA CCT GCA CTT GTC ACT TAG-3') (SEQ ID No. 11) (located in the intron between exons 4 and 5) were used
25 in the polymerase chain reaction to reveal the L⁻ allele; this amplification reaction generates a fragment of 400 bp.

 The primers ZO 243 and RU 178 (5'-ATG TTT CAT AGT TGG ATA TC-3') (SEQ ID No. 12)
30 (located in the neo cassette) are used in the

polymerase chain reaction to reveal the (-) allele; this PCR reaction generates a fragment of 500 bp.

For the analyses using DNA transfer (Southern blotting), the genomic DNA is digested with BamHI and
5 the probe used is the probe X4 (3 kb BamHI-XbaI fragment of the RXR α gene) (Metzger et al., 1995).

1.3 - Treatment with Tamoxifen

Tamoxifen (Sigma) solutions are prepared
10 according to the protocol described by Metzger and Chambon (2001). 1 mg of Tamoxifen dissolved in 100 μ l of sunflower oil is intraperitoneally injected into a transgenic mouse K5-Cre-ER^T for five consecutive days, and then again for three consecutive days, two, four
15 and six weeks later. The K14-Cre-ER^{T2} transgenic mice are intraperitoneally injected with 0.1 mg of Tam dissolved in 100 μ l of sunflower oil for five consecutive days, while the α P2-Cre-ER^{T2} and α AT-Cre-ER^T mice are treated with 1 mg of Tam.

20

1.4 - Histological analyses

The skin biopsies from animals of the same age and of the same sex were prepared at the level of the same sites of the body.

25 The skin samples are fixed in glutaraldehyde (2.5% in 0.1 M cacodylate buffer pH 7.2) overnight at 4°C and then post-fixed with 1% osmium tetroxide in a cacodylate buffer for 1 hour at 4°C. The tissues are dehydrated with increasing concentrations of alcohol

and then covered with EPON 812. Semithin sections of 2 μm are then stained with toluidine blue.

1.5 - Immunocytochemistry

5 After fixing in 2% paraformaldehyde, frozen sections of 10 μm are blocked in 5% normal goat serum (Vector Laboratories) incubated with the rabbit polyclonal antibody anti-MK6 (Babco). After washing in PBS/0.1% Tween 20, the sections are incubated with the
10 donkey anti-rabbit IgG antibody conjugated with CY3 (Jackson Immuno Research) and then mounted in the Vectashield medium (Vector Laboratories) containing DAPI (4',6-diamidino-2-phenylindole dihydrochloride; Boehringer Mannheim) (Brocard *et al.*, 1997). The anti-
15 Cre antibodies are used according to Indra *et al.* (1999).

1.6 - Immunohistochemistry

Tumors (8 - 16 mm) were excised 22 weeks after
20 DMBA application and immediately embedded in OCT and frozen on dry ice. 10 μm -thick frozen sections from CT and RXR $\alpha^{\text{ep-/-}}$ tumors were reacted with primary antibodies [mouse monoclonal anti-K10, rabbit polyclonal anti-K5 (gifts from Prof. Brigitte Lane,
25 Cell Structure research group, University of Dundee), rabbit polyclonal anti-K1 (Babco), rabbit polyclonal anti-K13 (gift from Prof. S. Yuspa, NIH), rabbit polyclonal $\alpha 6$ -integrin, biotin-conjugated rat monoclonal anti-CD31 (Pharmingen)], and revealed with
30 either CY3-conjugated donkey anti-rabbit, CY3-

conjugated goat anti mouse IgG antibodies or CY3-streptavidine (Jackson ImmunoResearch). Counterstaining was performed with DAPI (4',6-diamidino-2-phenylindole dihydrochloride, Boehringer Mannheim).

5

1.7 - Histology

The adipose tissue samples are removed from animals perfused with PFA, fixed with the aid of formaldehyde (20% in PBS) and then frozen in OCT (Tissue-Tek compound, Sakura). Cryosections of 10 μ m are stained with hematoxylin and eosin.

The liver is removed from animals, rinsed in PBS, fixed in a Bouin solution and then embedded in paraffin. Sections of 6 μ m are stained with hematoxylin and eosin.

Tumors from 25 animals were excised at 25 or 30 weeks after DMBA application, fixed in Bouin's fixative and embedded in paraffin. 5 μ m sections were stained with hematoxylin and eosin. Electron microscopy was performed as described (Li et al., 2001).

Subiliac lymph nodes were isolated 30 weeks after DMBA application, fixed in 4 % paraformaldehyde, photographed and embedded in paraffin. 5 μ m-thick sections were stained with hematoxylin and eosin.

25

1.8 - Assay of blood parameters

The assay of the triglycerides and of cholesterol is carried out according to Peters et al. (1997), with Boehringer Mannheim reagents. The assays of insulin and of glucose are carried out with the

30

Crystal Chem Inc. and glucofilms kit (Bayer Corp, USA) respectively.

1.9 - Treatments with carcinogenic agents

5 Skin papillomas and carcinomas were chemically induced by the two-stage carcinogenesis schedule. A single dose of 7,12-dimethylbenz(a)anthracene (DMBA) (5 μ g in 100 μ l acetone) was applied on the dorsal skin of 10 - 14 week old mice, 7 weeks before or 16 days after
10 the first Tam or oil (vehicle) treatments. 12-O-tetradecanoylphorbol-13-acetate (TPA) (5 μ g in 200 μ l acetone) was applied twice a week for 25 to 30 weeks, starting one week after DMBA application. CT and RXR α ^{ep-/-} mice were shaved 7 and 2 days before DMBA
15 treatment, and every second week for 30 and 15 weeks, respectively. 8 weeks after DMBA treatment, all-trans retinoic acid (t-RA, 20 nmoles in ethanol) or vehicle was topically applied 15 min before each TPA application.

20

1.10 - Determination of the number and size of the tumors

The length of the papillomas and the diameter of the melanocytic were measured with a Vernier
25 calliper on isofluorane anesthetised mice.

1.11 - Statistical analyses

Values are reported as mean $\pm$ SEM. Statistical significance ($p < 0.05$) was determined by unpaired
30 Student's test (Statview, Abacus Concepts, CA).

2) First example: Targeted inactivation of the RXR_α gene in the epidermis of adult mice.

5

To inactivate RXR_α in the epidermis, the inventors constructed a mouse carrying floxed $\text{RXR}_\alpha^{\text{L2}}$ alleles (Figure 1A) and used the transgenic mouse line K5-Cre-ER^T in which Tamoxifen (Tam) efficiently induces
10 Cre-mediated recombination in the keratinocytes of the basal layer (Indra et al., 1999). The crossing of the K5-Cre-ER^{T(tg/0)}/ $\text{RXR}_\alpha^{\text{L2/L2}}$ mouse with an $\text{RXR}_\alpha^{+/-}$ mouse (Figure 1A) (Kastner et al., 1994) or with an $\text{RXR}_\alpha^{\text{L2/+}}$ mouse makes it possible to obtain "promutant" (PM)
15 hemizygous (tg/0) mice for the K5-Cre-ER^T transgene which carries either an $\text{RXR}_\alpha^{\text{L2}}$ allele and a null RXR_α allele (genotype: K5-Cre-ER^{T(tg/0)}/ $\text{RXR}_\alpha^{\text{L2/-}}$), or two L2 alleles (genotype K5-Cre-ER^{T(tg/0)}/ $\text{RXR}_\alpha^{\text{L2/L2}}$). Fourteen-week-old PM mice were treated with Tamoxifen (five
20 days, 1 mg/day), and then treated again two, four and six weeks later. Six to twelve weeks after the first Tamoxifen treatment ("AFT: After First Tamoxifen Treatment"), nearly all the $\text{RXR}_\alpha^{\text{L2}}$ alleles (>80%) were converted to an $\text{RXR}_\alpha^{\text{L-}}$ allele in the epidermis isolated
25 from mice carrying a floxed allele (Figure 1B, lanes 2, 3, 5 and 6) or two floxed alleles (Figure 1B, lanes 1 and 4).

As expected (Indra et al. (1999)), no excision is observed in the mice treated only with the oil which

served as a vehicle for Tamoxifen; the Cre-mediated excision of exon 4 of RXR_{α} is in addition restricted to the skin and to the other organs which possess epithelia in which the K5 promoter is active (that is to say: the tongue, the salivary gland, the esophagus, Figure 1C).

Interestingly, hair loss (alopecia) was observed six to seven weeks after the first treatment with Tamoxifen in the ventral region of the mice; this was not observed in the mice treated with oil alone without Tamoxifen, or in the "control" mice of the same litter treated with Tamoxifen ($K5-Cre-ER^{T(tg/0)}/RXR_{\alpha}^{L2/+}$).

Twelve to sixteen weeks after the first injection, large ventral regions and smaller dorsal regions of the skin of the mice had lost their hair (Figure 2A and B) and cysts which grow bigger and which appear over the whole body with time are also visible under the surface of the skin (Figure 2C).

As the age increases (> twenty weeks after the first injection of Tamoxifen), minor lesions, which are not caused by fights, appear in hairless regions of the skin of the back, of the cheeks and of the posterior face of the ears (Figure 2D).

Sixteen weeks after the first treatment with Tamoxifen, histology of the hairless ventral and dorsal regions showed degeneration of the hair follicles resulting in the appearance of utriculi and of dermal cysts (Sundberg and King, 1996) (Figure 2 - compare E and F). The interfollicular epidermis is hyperplastic

with an increase in the incorporation of BrdU and an increased expression of the proliferation marker Ki67.

Dermal cellularity is increased and the capillaries are dilated (compare Figures 2E and 2G with
5 2F and 2H) below the thickened epidermis, thus reflecting an inflammatory reaction. The keratin 6 (K6), normally expressed selectively in the hair follicle outer root sheath (ORS) is also expressed in the hyperproliferative interfollicular epidermis
10 (Figure 2I and J) indicating an abnormality in the terminal differentiation of the keratinocytes (Porter *et al.*, 1998). All these abnormalities are less severe and/or appear much later in the males than in the females.

15 To increase the efficiency of the recombination mediated by Cre and induced by Tamoxifen, the inventors have constructed transgenic mouse lines K14-Cre-ER^{T2} in which the K14 promoter selective for the basal layer (Vassar *et al.*, 1989) directs the expression of Cre-ER^{T2}
20 whose activity may be induced by a gentler treatment with Tamoxifen (0.1 mg for five days) (Indra *et al.*, 1999).

K14-Cre-ER^{T2(tg/0)/RXR α ^{L2/L2}} mice were treated simultaneously with "control", K14-Cre-ER<sup>T2(tg/0)/RXR α ^{L2/L+},
25 K14-Cre-ER^{T2(0/0)/RXR α ^{L2/+} and K14-Cre-ER^{T2(0/0)/RXR α ^{L2/L2} mice obtained from the same litter.}}</sup>

Within two weeks, the RXR α ^{L2} alleles were completely converted to an RXR α ^{L-} alleles in the epidermis (Figure 1D, lanes 1 and 7), but not in the
30 dermis (Figure 1D, lanes 2 and 8) of transgenic mice

expressing K14-Cre-ER^{T2}, thus demonstrating the increased efficiency of Cre-ER^{T2} for mediating, after a Tamoxifen treatment, the selective somatic mutation of floxed RXR α in the epidermis.

5 No conversion of L2 to L- appeared in the controls lacking the transgene K14-Cre-ER^{T2} (Figure 1D, lanes 5, 6, 11 and 12) or not treated with Tamoxifen (Figure 1D, lanes 3, 4, 9 and 10). Furthermore, 8 weeks after the treatment with Tamoxifen, only the RXR α ^{L-}
10 allele was detected in the epidermis of K14-Cre-ER^{T2(tg/0)}/RXR α ^{L2/L2} mice, indicating that RXR α was excised in the majority, if not all, of the stem cells of the epidermis. The inactivation of RXR α also appears in the other epithelia of other organs in which the
15 K14 promoter is active (Wang et al., 1997) (that is to say the tongue, esophagus, stomach).

From 6 weeks after the start of Tamoxifen treatment, the K14-Cre-ER^{T2(tg/0)}/RXR α ^{L2/L2} mice exhibited abnormalities similar to those observed in the
20 K5-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2} mice treated with Tamoxifen, that is to say a marked hair loss with visible cysts, while visible focal lesions appear at subsequent stages (Figure 2K and L).

The underlying dermal and epidermal
25 histological abnormalities are also similar to those observed above for the K5-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2} mice treated with Tamoxifen.

The inactivation of floxed RXR α in the adult epidermis is obtained more rapidly and with lower doses

of Tamoxifen with the K14-Cre-ER^{T2} mice than with the K5-Cre-ER^T mice, but the resulting skin abnormalities are finally similar, in both cases, more severe in the females than in the males.

5 Interestingly, these abnormalities are also similar to those observed in the K14-Cre^(tg/0)/RXR α ^{L2/L2} or K14-Cre^(tg/0)/RXR α ^{L2/-} mice in which the floxed RXR α alleles are selectively excised in the epidermis during fetal development, thus leading to inactivation of RXR α
 10 in the keratinocytes of the epidermis and in the hair follicle root sheaths. In fact, after three weeks of age, these "constitutive" and epidermis-specific RXR α mutants develop progressive alopecia with typical characteristics of degenerated hair follicles, utriculi
 15 and dermal cysts, which may all be attributed to defects in the hair cycle. Furthermore, these mutants also exhibit interfollicular hyperproliferation of the keratinocytes, as well as abnormal terminal differentiation (with expression of K6), and an
 20 increase in the dermal cellularity associated with an inflammatory reaction of the skin.

 Although RXR β is expressed in the epidermis of mice, the skin of adult RXR β ^{-/-} mutants appears normal (Kastner et al., 1996), thus suggesting functional
 25 redundancies between the various RXRs. As expected, K5-Cre-ER^{T(tg/0)}/RXR α ^{L2/L-}/RXR β ^{-/-} and K5-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2}/RXR β ^{-/-} mice treated with oil (without Tamoxifen) do not exhibit skin abnormalities, whereas after treatment with Tamoxifen, these mice

begin to lose their hair 4 weeks after the first treatment with Tamoxifen while large regions of the skin are hairless 16 to 18 weeks after the start of the treatment (compare Figure 3A with Figure 2D).

5 Focal lesions of the skin which were not observed in the single RXR_{α} mutants are also frequently observed in the double $RXR_{\alpha/\beta}$ mutants fourteen to sixteen weeks after the first treatment with Tamoxifen, in particular on the hairless skin of the trunk, behind
10 the ears and behind the mouth (figure 3A).

Histology of the hairless skin shows disappearance of the hair follicles and the presence of utriculi and of dermal cysts (Figure 3B). The epidermis is highly hyperplastic and hyperkeratinized (compare
15 Figures 3B and 3C with Figures 2E and 2G, and Figures 2F and 2H). Abnormal expression of K6 is observed through the whole epidermis (Figure 3D) and an inflammatory reaction with an increase in cellularity is also observed (Figure 3B). In the lesioned skin, the
20 epidermis is covered with a crust and is more hyperplastic and hyperkeratinized. The triple mutants $K5-Cre-ER^{T(tg/0)}/RXR_{\alpha}^{L2/L2}/RXR_{\beta}^{-/-}/RXR_{\gamma}^{-/-}$ do not reveal an additional role of RXR_{γ} in the adult skin. Thus, RXR_{β} may partially compensate for the loss of RXR_{α} function.

25 Interestingly, the functional redundance is more pronounced in the males than in the females, since the male and female double mutants RXR_{α}/RXR_{β} are affected in a similar manner unlike the single mutants.

Taken together, the preceding results show the efficiency of the Cre-ER^T, Cre-ER^{T2} and Cre-ER^{T3} recombinases (see Example 5 and Figures 9, 10 and 11) for generating somatic mutations which are targeted,
5 specific for a cell type and temporally controlled, in adult mouse tissues.

The preceding results also make it possible to demonstrate the use of Cre-ER^T-RXRfloxed mice for analyzing and studying the complex biological function
10 of the various RXR sub-types (RXR_α, RXR_β, RXR_γ) in a particular tissue, that is to say the epidermis.

This study has thus made it possible to reveal the existence of functional redundancies between RXR_α and RXR_β, although the role of RXR_α is clearly
15 predominant.

The mechanism of action of the various RXR in the molecular events which lead to alopecia and to abnormalities in the keratinocytes of the epidermis which are deficient in nuclear RXR receptors remains
20 unknown.

Nevertheless, their role as heterodimeric partners of a certain number of nuclear receptors (for example RARs, TRs, VDR, PPARs) which act as signal transducers in various signaling pathways has been
25 suggested in numerous studies *in vitro* using cells in culture, and confirmed *in vivo* in some cases using targeted mutagenesis (Chambon, 1996; Mascrez et al., 1998; Wendling et al., 1999). Interestingly, like RXR_α, VDR is expressed in the ORS of the hair follicle
30 (Reichrath et al., 1994), and mice in which the two

alleles of the VDR gene are inactive (VDR "knock-out" mice) develop progressive secondary alopecia, suggesting that VDR is involved in the hair cycle rather than in primary hair growth (Yoshizawa et al., 1997; Li et al., 1997; Li et al., 1998).

Interestingly, the alopecia developed by fourteen-week-old VDR^{-/-} mice and K5-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2}/RXR β ^{-/-} mice eighteen weeks after their treatment with Tamoxifen appear very similar, although the skin of the VDR^{-/-} mice does not exhibit the lesions observed on the epidermis of the mice deficient in RXR (compare Figures 3A and 3E).

At the histological level, similar utriculi and dermal cysts are observed (compare Figures 3B and 3F) but no hyperproliferation of the keratinocytes, or abnormal differentiation revealed by the expression of K6 is observed in the epidermis of the VDR^{-/-} mice (compare Figures 3C and 3D with Figures 3G and 3H). Thus, the alopecia generated by the selective inactivation of RXR in the keratinocytes of adult mice is thought to reveal a major role of the RXR/VDR heterodimers in the hair follicle cycle.

3) Second example: Targeted inactivation of the SNF2 β gene in the epidermis of adult mice

To inactivate the SNF2 β gene in the epidermis of adult mice, the inventors constructed a mouse carrying floxed SNF2 β alleles (L2+; Figure 4A) and used the transgenic mouse line K14-Cre-ER^{T2} in which

Tamoxifen effectively induces Cre-mediated recombination in the keratinocytes of the basal layer of the epidermis. Eight-week-old K14-Cre-ER^{T2(tg/0)}/SNF2 β ^{L2/L2} and K14-Cre-ER^{T2(0/0)}/SNF2 β ^{L2/L2} mice were treated with Tamoxifen for five days at the rate of 0.1 mg/day or with oil (-). Skin biopsies were taken, the dermis and the epidermis separated, the genomic DNA prepared, digested with BamHI, separated by agarose gel electrophoresis and transferred onto nylon membranes which were hybridized with the radiolabeled 5' probe (Figure 4A). The corresponding autoradiographs are presented in Figure 4B. In the absence of Tamoxifen, no excision in the SNF2 β gene is observed whether in the dermis or the epidermis of K14-Cre-ER^{T2(tg/0)}/SNF2 β ^{L2/L2} mice. On the other hand, the injection of Tamoxifen into these same mice induces the excision of the floxed fragment of the SNF2 β gene only in the epidermis, because the K14 promoter is active only in this tissue and not in the dermis. As expected, Tamoxifen induces no excision in K14-Cre-ER^{T2(0/0)}/SNF2 β ^{L2/L2} "control" mice whose cells do not contain a Cre-ER^{T2} transgene.

4) Third example: Targeted inactivation of the RXR α gene in murine adipocytes

To carry out the spatiotemporally controlled site-specific mutagenesis in the adipocytes, the inventors created transgenic mice called aP2-Cre-ER^{T2}

expressing the Cre-ER^{T2} fusion protein under the control of the promoter of the gene encoding adipose protein 2 (aP2) which is specifically active in the adipocytes (Ross et al., 1990).

5 To invalidate the RXR α gene selectively in the adipocytes, the aP2-Cre-ER^{T2(tg/0)} mice were first crossed with RXR α ^{L2/+} mice so as to produce aP2-Cre-ER^{T2(tg/0)}/RXR α ^{L2/L2} mice. These mice were then crossed with RXR α ^{+/-} mice (Kastner et al., 1994) so as
10 to produce aP2-Cre-ER^{T2(tg/0)}/RXR α ^{L2/-} mice. Such four week old mice were treated (+) or not (-) with Tamoxifen (1 mg/day) for five days, and the adipose tissue collected one month after the last injection of Tamoxifen. The DNA was extracted from the adipose
15 tissue, or after separation of the adipocytes (80%-purified adipocytes) from the connective tissue and blood vessels (non-adipocytes). The DNA is then digested with BamHI and then separated by agarose gel electrophoresis, transferred onto nylon membranes and
20 then hybridized with the radiolabeled X4 probe. The corresponding radiographs are presented in Figure 5.

 In the absence of Tamoxifen, no excision in the RXR α gene is observed whether in the adipose tissue, the adipocytes or other non-adipocyte cells of the
25 epidermis of aP2-Cre-ER^{T2(tg/0)}/RXR α ^{L2/-} mice. On the other hand, the injection of Tamoxifen into such mice induces the excision of the floxed fragment of the RXR α gene only in the purified adipocytes and the adipose tissue.

- 73 -

Analysis of the weight of the aP2-Cre-ER^{T2(tg/0)}/RXR α ^{L2/-} and aP2-Cre-ER^{T2(tg/0)}/RXR α ^{L2/+} animals treated with Tam (mutants; KO, and controls; CT, respectively) and fed with a normal diet (AN),
5 determined for 30 weeks, revealed no significant difference between the two groups of animals (Figure 6A). Furthermore, the weight of the adipose tissue and the morphology of the adipocytes were similar (Figure 6B and 6C). On the other hand, the glucose level was
10 abnormally high and the triglyceride levels were lower in the mutant animals (Figure 6D). By feeding the Tamoxifen-treated control animals with food rich in fat and in glucose (AR), they became obese (increase in the weight, the mass of adipose tissue and hypertrophy of
15 the adipocytes) (Figures 6A-C). The triglyceride and insulin levels are also much higher than in the animals fed with AN. On the other hand, no increase in the weight of the animals and in the mass of the adipose tissue is observed in the mutated animals fed with AR.
20 Furthermore, the adipocytes are not hypertrophic, and the triglyceride and insulin levels are similar to those observed with an AN.

These mutant animals therefore constitute advantageous models for studying obesity and diabetes,
25 as well as for testing treatments of these diseases.

5) Fourth example: Targeted inactivation of the RXR α gene in murine hepatocytes

To carry out the spatiotemporally controlled site-specific mutagenesis in the liver, the inventors created transgenic mice called " α AT-Cre-ER^T" expressing the Cre-ER^T fusion protein under the control of the promoter of the gene for human α -1-antitrypsin which is specifically active in the hepatocytes (Imaï et al., 2000).

To inactivate RXR α in the murine hepatocytes, the inventors constructed a mouse carrying floxed RXR α ^{L2} alleles (Figure 1A) and used the transgenic mouse line α AT-Cre-ER^T in which Tamoxifen (Tam) efficiently induces Cre-mediated recombination in the hepatocytes (Figure 7). α AT-Cre-ER^{T(tg/0)} mice which express Cre-ER^T in about 50% of the hepatocytes (Imaï et al., 2000) were crossed with RXR α ^{L2/L2} mice so as to produce α AT-Cre-ER^{T(tg/0)}/RXR α ^{L2/L2} mice. Such three month old mice were treated with Tam (1 mg/day) for five days, and the heart and the liver collected 7 days after the first injection of Tam (day 7), from one and three animals respectively. The heart and the liver were also collected from one and three animals, respectively, of the same genotype without treatment with Tam (day 0). The DNA was extracted from these tissues, and after digestion with BamHI, the RXR α alleles were analyzed by Southern blotting.

As expected, no excision is observed in the cells of the heart of mice treated with Tamoxifen (Figure 7) or the liver of mice not treated with Tamoxifen (day 0). On the other hand, on day 7, the

mice treated with Tamoxifen exhibit excision in the RXR α gene in about 50% of the liver cells, which indeed corresponds to the expression, in the form of a mosaic, of the Cre-ER^T protein in the hepatocytes of the
5 α AT-Cre-ER^T mouse.

On the other hand, 30 and 90 days after the treatment with Tamoxifen, practically no null RXR α cells are again observed (Fig. 8). Furthermore, the RXR α ^{L-/L-} cells proliferate much less than the RXR α ^{L2/L2}
10 cells after partial hepatectomy (HP) carried out on mutant animals treated with Tamoxifen (Fig. 8, compare lanes 68 and lanes 15-17). Thus, RXR α is involved in the proliferation of the hepatocytes.

15 6) Fifth example: The chimeric Cre recombinase Cre-ER^{T3} is ten times more sensitive to Tamoxifen than Cre-ER^{T2}

With the aim of increasing the sensitivity of the chimeric recombinase Cre-ER^{T2} to Tam or OHT, the
20 inventors replaced, in Cre-ER^{T2}, V400 with G (Cre-ER mutant M543A/L544A, called "Cre-ER^{T3}"). Transgenic mice expressing Cre-ER^{T3} under the control of the K5-promoter were obtained, and a line expressing the chimeric recombinase at similar levels to those
25 detected in the K5-Cre-ER^T and K5-Cre-ER^{T2} lines was generated (see Figure 9, and results not presented).

The sensitivity to OHT was tested in a first instance by analyzing the intracellular location of the chimeric proteins. Whereas after treatment with 0.1 mg
30 of OHT, Cre-ER^{T2} and Cre-ER^{T3} are both located in the

cellular nuclei, Cre-ER^{T3} is present in a larger fraction thereof at lower doses of Tam or OHT. At the dose of 0.001 mg, about 1/3 of the nuclei are strongly labeled with anti-Cre antibodies in the basal layer of the epidermis of K5-Cre-ER^{T3} mice, whereas no positive nucleus is observed in the skin of K5-Cre-ER^{T2} mice (Figure 10). It was also possible to visualize this difference in sensitivity using the reporter mice RosaR26R (Rosa^{f1/+}) (Soriano, 1999). Indeed, although the excision levels induced with 1 and 0.1 mg of OHT are similar in both lines, the excision is markedly more efficient in the K5-Cre-ER^{T3} mice following treatments with 0.01 and 0.001 mg of OHT (Figure 11). It should be noted that in the absence of treatment, no recombinase activity is detected in the K5-Cre-ER^{T3} line.

7) Sixth example : Role of RXR α in skin carcinogenesis

The stages of initiation, promotion, progression and tumor conversion in the skin carcinogenesis model are well characterised. In WT mice, initiation using 7,12-dimethylbenz(a)anthracene (DMBA) and promotion with 12-O-tetradecanoylphorbol 13-acetate (TPA) provokes papillomas that are hyperplastic, well-differentiated skin lesions. After a latency period of about 25 weeks, a percentage of papillomas progress to carcinoma (Ghadially and Ghadially, 1996; Hennings et al., 1993; Klein-Szanto et al., 1989).

8 - 12 week-old K14-Cre-ER^{T2(tg/0)}/RXR α ^{L2/L2} mice were Tam-treated (0.1 mg per day for 5 days), which resulted in RXR α ^{ep-/-} mice (for epidermal keratinocyte-selective RXR α null genotype) fully lacking RXR α in
5 keratinocytes of the epidermis and outer root sheath of the hair follicles 2 weeks after the first Tam injection. Similar Tam treatments applied to K14-Cre-ER^{T2(tg/0)}/RXR α ^{L2/+} and K14-Cre-ER^{T2(0/0)}/RXR α ^{L2/L2} littermates, resulted in control (CT) mice which
10 carried, in epidermal keratinocytes, one WT (+) and one RXR α L- allele, and two RXR α L2 alleles, respectively.

To induce papilloma formation, CT and RXR α ^{ep-/-} female mice were topically-treated 16 days after the first Tam injection with a single dose of DMBA (50 μ g),
15 and then twice a week with TPA (5 μ g) for 25-30 weeks (Fig. 12A). Seven to eight weeks after DMBA application, small papilloma were observed in all CT and RXR α ^{ep-/-} mice, and their number and size increased with time (Fig.12B and data not shown). Interestingly,
20 RXR α ^{ep-/-} mice developed approximately twice as many tumors as CT mice, and 30 weeks after the start of DMBA/TPA treatment, an average of 12 and 28 papillomas were present in CT and RXR α ^{ep-/-} mice, respectively (Fig. 12B). Although CT males were less sensitive to
25 the DMBA/TPA treatment than CT females (the number of papillomas was two-fold lower), RXR α ^{ep-/-} males also exhibited about twice as many papilloma as CT males (data not shown). In both males and females, the size

and growth rate of the tumors were increased in RXR α ^{ep-/-} mice (Fig. 12C and 12D; and data not shown). After 30 weeks of TPA treatment, the papillomas of CT females were never longer than 12 mm, whereas ~10 % of the papillomas of RXR α ^{ep-/-} females had a length between 12 and 16 mm, and 3 % reached a length of 30 - 40 mm. Note that in the absence of either DMBA or TPA treatment no papilloma appeared in CT and RXR α ^{ep-/-} mice, even after 30 weeks of treatment (data not shown).

To characterize the tumors induced by DMBA/TPA treatment, ~ 50 tumors from 6 CT and 6 RXR α ^{ep-/-} mice were histologically examined. In agreement with previous reports, almost all tumors analysed 25 weeks after the start of DMBA/TPA treatment of CT mice were benign papillomas characterised by skin folds integrated by a core of connective tissue and lined by an acanthotic, hyperkeratotic, stratified squamous epithelium (Fig. 13Aa and 13C (table)). 35 % of the papillomas exhibited an atypical hyperplasia, 5% displayed an *in situ* carcinoma, but no focal carcinoma could be detected (Fig. 13Ab and 13C (table), and data not shown). It was only 30 weeks after tumor initiation that some focal carcinoma were observed in ~16 % of the tumors (Fig. 13Ac and 13C (table), and data not shown). In contrast, 25 weeks after tumor initiation, almost all of the RXR α ^{ep-/-} papilloma exhibited an atypical hyperplasia, 10 % displayed *in situ* carcinoma, and focal carcinoma were observed in 40 % of the tumors

(Fig. 13Ba-c and 13C (table), and data not shown). Moreover, 5 weeks later, 27 % of the tumors had progressed to differentiated and undifferentiated squamous cell carcinomas (SCC) with extensive local invasion of cancer cells in the dermis (Fig. 13Bd-g and 13C (table)). Advanced grade I SCC (Fig. 13Bd, note the pearls with keratin in the center), grade II SCC (Fig. 13Be, note the fused keratin pearls, with only very reduced remaining regions of keratinization) and grade III and IV SCC (Fig. 13Bf and g ; note the spindle shape cytoplasm and nucleus of the cancer cells that are found along with the muscle cells, indicating a complete conversion into malignant cancer) were observed in 3 to 9 % of the tumors (Fig. 13C (table)). Highly aggressive spindle cell carcinoma (~ 6 %) and basal cell carcinoma (~9 %) were also found in $RXR\alpha^{ep-/-}$, but not in CT animals (Fig. 13Bh and i and 13C (table)).

Taken together, these above data show that higher incidence of malignant conversion occurred in papillomas of $RXR\alpha^{ep-/-}$ than of CT mice.

In agreement with previous reports (Epstein, 1992 ; Epstein et al., 1967 ; Husain et al., 1991), DMBA application to skin of CT mice induced on its own the formation of benign melanocytic growths (nevi) (Fig. 14A and data not shown), histologically characterized by subepidermal accumulation of melanocytes (Epstein et al., 1967). Interestingly, melanocytic growths were ~7-fold more efficiently induced by DMBA treatment of $RXR\alpha^{ep-/-}$ mice (Fig. 14A and

B). Furthermore, these $\text{RXR}\alpha^{\text{ep-/-}}$ melanocytic growths were generally larger (Fig. 14C) and contained a larger number of densely melanin-laden melanocytes in both the dermis and the hypodermis, and invasion of the underlying musculature was observed in 5 - 10 % of them (data not shown), thus, indicating a higher potential of metastasis (Broome Powell et al., 1999 ; Epstein, 1967 ; Goding, 2000 ; Hirobe, 1995 ; Klein-Sozanto, 1989).

Electron microscopy revealed that melanocytes from CT nevi contained mainly stage III and IV melanosomes in agreement with previous reports. Most of the melanocytes present in melanocytic growths from $\text{RXR}\alpha^{\text{ep-/-}}$ mice contained ~10-fold more melanosomes than CT melanocytes (data not shown). Moreover, in about 40 % of the melanocytes, more than 30% of the granules had the appearance of vesicles of variable electron opacity with a distinct substructure (data not shown). The membrane of a number of melanosomes was disrupted and smaller melanin granules were scattered in the cellular matrix (data not shown). Taken together, these results indicate that DMBA induced melanoma formation in the skin of $\text{RXR}\alpha^{\text{ep-/-}}$ mice.

Most interestingly, autopsies performed 30 weeks after DMBA treatment revealed that subiliac lymph nodes were enlarged and pigmented in two out of 4 $\text{RXR}\alpha^{\text{ep-/-}}$ mice, but not in CT mice (0/4) (Fig.15A a and b, and data not shown). Histological analysis revealed the presence in $\text{RXR}\alpha^{\text{ep-/-}}$ mice, but not in CT mice, of

few melanin-laden melanocytes in the lymphosinus of $RXR\alpha^{ep-/-}$ mice, which were slightly enlarged (Fig. 15A compare panel c and d, and data not shown). Multiple TPA applications after DMBA treatment increased the number of melanocytic growths in CT mice and $RXR\alpha^{ep-/-}$ mice (Fig. 14B, and compare Fig. 12C and Fig. 14A). Whereas after 30 weeks of DMBA/TPA treatment, only about 10 nevi were observed per CT mouse (most of them having a diameter below 2 mm), more than 50 melanocytic growths were found per $RXR\alpha^{ep-/-}$ mouse (more than 50 % of them having a diameter over 2 mm) (Fig. 14 C). Melanocytes were found in 5/6 subiliac lymph nodes from $RXR\alpha^{ep-/-}$ mice, and in 1/6 from CT mice (Fig. 15A e and f, and data not shown). DMBA/TPA treatments resulted in about 5 times more melanocytes in the lympho sinus of $RXR\alpha^{ep-/-}$ mice than after DMBA treatments. Most melanocyte-containing lymphosinus were markedly enlarged and clusters of melanocytes were seen in about 20 % of them (Fig. 15h, and data not shown). In contrast, only few melanocytes were seen in only one of the lymph nodes of CT mice (data not shown).

As $RXR\alpha$ L2 alleles, but not L- alleles could not be detected in the $RXR\alpha^{ep-/-}$ melanocytic tumors induced either by DMBA or DMBA and TPA, and L- but not L2 alleles in their epidermis (data not shown), these results demonstrate that the formation of melanoma in $RXR\alpha^{ep-/-}$ mice is dependent on $RXR\alpha$ ablation in keratinocytes, but not in melanocytes.

8) Seventh example : Role of RXR α in the antitumoral effect of retinoic acid

In agreement with previous results (Hansen, 2000 ; Huang et al., 1997 ; Leder et al, 1990 ; Slaga
5 et al, 1980 ; Tennenbaum et al, 1998), topical application of all-trans retinoic acid (t-RA) efficiently blocked skin papilloma formation induced by DMBA/TPA in both female and male wild-type 129Sv (data not shown). RA application also efficiently inhibited
10 tumor formation in CT, but not in RXR $\alpha^{ep-/-}$ mice. Indeed, 25 weeks of RA treatment induced a reduction of size of the papilloma (from 2 - 3 mm to 1 mm, in 3 out of 8 papillomas of 6 mice) and the size remained the same in the other papillomas. In contrast, no decrease
15 in size of the papilloma was observed in RXR $\alpha^{ep-/-}$ mice. Furthermore, after 25 weeks of DMBA/TPA treatment, an average of 1 tumor was observed in RA treated CT mice, whereas an average of ~ 17 tumors were found in similarly treated RXR $\alpha^{ep-/-}$ mice. As previously observed
20 in RXR $\alpha^{ep-/-}$ mice in the absence of RA treatment, ~ 11 % of the tumors were larger than 12 mm (data not shown), whereas the size of the tumors from CT mice was below 2 mm.

Furthermore, after RA treatment, both the
25 number and size of melanocytic growths were increased in RXR $\alpha^{ep-/-}$ mice compared to CT mice (data not shown).

REFERENCES

- Akari et al. (1995) Proc. Natl. Acad. Sci. USA 92: 160-164.
- 5 Akagi et al. (1997) Nucleic Acids Res. 25: 1766-1773.
- Arnheiter et al. (1990) Cell 62: 51.
- Babinet (1995) Médecine/Sciences 11: 1154-1157.
- Barbonis et al. (1993) Nucleic Acids Research, 21: 2025-2029.
- 10 Baretino et al. (1994) EMBO J. 13: 3039-3049.
- Baur et al. (1996) EMBO J. 15: 110-124.
- Beato (1989) Cell 56: 335-344.
- Betz et al. (1996) Curr. Biol. 6: 1307-1316.
- Bourguet et al. (1995) Nature 375: 377-382.
- 15 Brinster et al. (1982) Nature 296: 39-42.
- Brocard et al. (1997) Proc. Natl. Acad. Sci. USA 94: 14559-14563.
- Brocard et al. (1998) Nucleic Acids Res. 26: 4086-4090.
- Broome Powell et al. (1999) Carcinogenesis 20: 1747-53.
- 20 Brown et al. (1987) Cell 49: 603-612.
- Cadepond et al. (1997) Ann. Rev. Med. 48: 129-156.
- Capecchi (1989) Science 244: 1288-1291.
- Cavaillès et al. (1994) Proc. Natl. Acad. Sci. USA 91: 10009-10013.
- 25 Cavaillès et al. (1995) EMBO J. 14: 3741-3751.
- Chambon (1996) FASEB J. 10: 940-954.
- Danielian et al. (1992) EMBO J. 11: 1025-1033.
- Denisen et al. (1995) Proc. Natl. Acad. Sci. USA, 92: 7376-7380.
- 30 Deuschle et al. (1990) Science 2: 480-483.

- Diaz et al., (1999) JBC 274: 6634.
- Durand et al. (1995) EMBO J. 13: 5370-5382.
- Epstein et al. (1967) J Natl Cancer Inst 38:19-30.
- Epstein (1992) Photodermatol Photoimmunol Photomed 9:
5 91-8.
- Evans (1988) Science 240: 889-895.
- Feil et al. (1996) Proc. Natl. Acad. Sci USA 93:
10887-10890.
- Feil et al. (1997) Biochemical and Biophysical Res.
10 Com. 237: 752-757.
- Figge et al. (1988) Cell 52: 713-722.
- Ghadially and Ghadially (1996) *Tumours of the skin*.
IARC Sci Publ 126:1-43.
- Goding (2000) *Melanocyte development and malignant*
15 *melanoma*. Forum (Genova) 10(3): 176-87.
- Gossen et al. (1992) Proc. Natl. Acad. Sci. USA 89:
5547-5551.
- Green and Chambon (1988) Trends Genetics 4: 309-314.
- Gronemeyer (1991) Ann. Rev. Genet. 25: 89-123.
- 20 Gu et al. (1993) Cell 73: 1155-1164.
- Halachmi et al. (1994) Science 264: 1455-1458.
- Hansen et al (2000) Cracinogenesis 21:1271-1279.
- Hennings et al. (1993) Proc Soc Exp Biol Med 202:1-8.
- Hirobe (1995) Histol. Histopathol. 10: 223-37.
- 25 Hu et al. (1987) Cell 48: 555-556.
- Huang et al. (1997) Proc. Natl. Acad. Sci USA 94:5826-30.
- Hug et al. (1998) Mol. Cell. Biol. 8: 3065.
- Husain et al (1991) Cancer Res 51(18):4964-70.
- Hynes et al. (1981) Proc. Natl. Acad. Sci. USA 78:
30 2038-2042.

- Imai et al. (2000) *Genesis* 26: 147-148.
- Indra et al. (1999) *Nucl. Acid. Res.* 27: 4324-4327.
- Israel et al. (1989) *Nucleic Acids Res.* 17: 2589-2604).
- Jackson et al., (1993) *EMBO J.* 12: 2809-2819.
- 5 Jaenisch (1988) *Science* 240: 1468-1474.
- Kastner et al. (1994) *Cell* 78: 987-1003.
- Kastner et al. (1996) *Gene Dev.* 10: 80-92. 67.
- Kellendonk et al. (1996) *Nucleic Acids. Res.* 24: 1404-1411.
- 10 Kellendonk et al. (1999) *J. Mol. Biol.* 285: 175-182.
- Kilby et al. (1993) *TIG* 9: 413-421.
- Klein-Szanto et al. (1989) *Carcinogenesis* 10: 2169-72.
- Klock et al. (1987) *Nature* 329: 734-736.
- Kühn et al. (1995) *Science* 269: 1427-1429.
- 15 Labow et al. (1990) *Mol. Cell. Biol.* 10: 3343-3356.
- Lakso et al. (1992) *Proc. Natl. Acad. Sci USA* 89: 6232-6236.
- Leder et al. (1990) *Proc. Natl. Acad. Sci. USA* 87: 9178-82.
- 20 Ledouarin B. et al. (1995) *EMBO J.* 14: 2020-2033.
- Lee et al. (1981) *Nature* 294: 228-232.
- Lee et al. (1995) *Nature* 374: 91-94.
- Leng et al. (1995) *Mol. Cell. Biol.* 15: 255-263.
- Li et al. (1997) *Proc. Natl. Acad. Sci. USA* 94: 9831-9835.
- 25 Li et al. (1998) *Endocrinology* 139: 4391-4396).
- Littlewood et al. (1995) *Nucleic Acids Res.* 23: 1686-1690.
- Lobe et al. (1999) *Dev. Biol.* 208: 281-292.

- Logie et al. (1995) Proc. Natl. Acad. Sci. USA 92:
5940-5944.
- Mahfoudi et al. (1995) Proc. Natl. Acad. Sci. USA 92:
4206-4210.
- 5 Mao et al. (1995) Proc. Natl. Acad. Sci. USA 96:
5037-5042.
- Mascrez et al. (1998) Development 125: 4691-4707.
- Mayo et al. (1982) Cell 29: 99-108.
- Metzger et al. (1995) Proc. Natl. Acad. Sci. USA 92:
10 6991-6995.
- Metzger and Chambon (2001), Methods, *in press*.
- Metzger and Feil (1999), Curr. Opin. Biotechnol. 5:
470-476.
- Nover et al. (1991) in "Heat Shock Response", e.d.
- 15 Nover, L CRC, Boca Raton, Fl pp. 167-220.
- Orban et al. (1992) Proc. Natl. Acad. Sci USA 89:
6861-6865.
- Parker (1993) Curr. Opin. Cell. Biol, 5: 499-504.
- Peters et al. (1997) J. Biol. Chem 272: 27307-27312.
- 20 Picard et al. (1994) Curr. Opin. Biotechnol. 5:
511-515.
- Porter et al. (1998) J. Invent. Dermatol. 110: 951-957.
- Rajewsky et al. (1996) J. Clin. Invest. 98: 600-603.
- Reichrath et al. (1994) Brit. J. Dermatol. 131:
25 477-482.
- Renaud et al. (1995) Nature 378: 681-689.
- Ross et al. (1990) Proc. Natl. Acad. Sci USA 87:
9590-9594.
- Saatcioglu et al. (1993) Mol. Cell. Biol. 13:
30 3675-3685.

- St-Onge et al. (1996) Nucleic Acids Res. 24: 3875-3877.
- Saitou et al. (1995) Nature 374: 159-162.
- Sambrook et al. (1989) Molecular cloning: a laboratory manual second edition - Cold Spring Harbor Laboratory Press. Cold Spring Harbor, NY. USA
- 5 Sauer (1994) Current opinion in Biotechnology 5: 521-527.
- Sauer et al. (1990) The New Biologist 2: 441-449.
- Sauer B. (1998) Methods 14 381-392.
- 10 Schmedt et al. (1998) Nature 394: 901-904.
- Schmidt et al. (1990) Mol. Cell. Biol. 10: 4406-4411.
- Schwenk et al. (1998) Nucleic Acids Res. 26: 1427-1432.
- Seark et al. (1995) Mol. Cell. Biol. 5: 1480-1489.
- Shaikh and Sadowski (2000) J. Mol. Biol. 302: 27-48.
- 15 Simons (1994) New York. Vitam. Horm. 49: 49-130.
- Slaga et al. (1980) Proc Natl Acad Sci USA 77(4):2251-4.
- Soriano (1999) Nature Genet 21: 70-71.
- Sternberg et al. (1986) J. Mol. Biol. 187: 197-212.
- 20 Sumi-Chinose et al. (1997) Mol. Cell. Biology 17, 5976-5986.
- Sundberg et al. (1996) Dermatol. 7: 249-267.
- Tennenbaum et al. (1998) Cancer Res 58(7): 1435-43.
- Vassar et al. (1989) Proc. Natl. Acad. Sci. USA, 86: 1563-1567.
- 25 Wang et al. (1997) Proc. Natl. Acad. Sci. USA 94: 219-226.
- Wendling et al. (1999) Proc. Natl. Acad. Sci. USA 96: 547-551.

- Wurtz *et al.* (1996) *Natural Structural Biology* 3: 87-93.
- Yoshizawa *et al.* (1997) *Nature Genet.* 16: 391-396.
- Zechel *et al.* (1994) *EMBO J.* 13: 1425-1433.
- 5 Zhang *et al.* (1996) *Nucleic Acids Res.* 24: 543-548.

CLAIMS

1. Metazoan organism, with the exception of humans, characterized in that at least one cell of said
5 organism comprises at least:
- (i) one fusion protein comprising sequentially:
- a recombinase protein;
 - a hinge region of at least 15 amino acids;
 - a polypeptide comprising the ligand-binding
10 domain of the human nuclear estrogen receptor, or of a vertebrate nuclear estrogen receptor, and their natural variants or one of their fragments, said polypeptide exhibiting at least one mutation
15 relative to the wild-type form of said ligand-binding domains, or of their natural variants, or of their fragments,
- said fusion protein having a negligible, or even zero, recombinase activity in the presence of a natural
20 ligand and a recombinase activity induced by small quantities of synthetic ligand endowed with antiestrogenic activity;
- (ii) one or more gene or intergenic DNA sequences of interest naturally belonging to said genome of
25 said organism into which one or more recognition sites of said recombinase protein are inserted, said DNA sequence(s) of interest being located in one or more of the chromosomes of the genome of said cell.

2. Organism according to Claim 1, characterized in that said recombinase protein is preferably selected from the group of site-specific recombinases composed of the Cre recombinase of bacteriophage P1, the FLP
5 recombinase of *Saccharomyces cerevisiae*, the R recombinase of *Zygosaccharomyces rouxii* pSR1, the A recombinase of *Kluyveromyces drosophilarius* pKD1, the A recombinase of *Kluyveromyces waltii* pKW1, the integrase λ Int, the recombinase of the GIN recombination system
10 of the Mu phage, of the bacterial β recombinase or a variant thereof.

3. Organism according to Claim 2, characterized in that the recombinase is the Cre recombinase of bacteriophage P1 and its natural or synthetic variants.

15 4. Organism according to Claims 1 to 3, characterized in that said sites of recognition specific for said Cre recombinase are preferably chosen from the group composed of the sequences Lox P, Lox 66, Lox 71, Lox 511, Lox 512, Lox 514.

20 5. Organism according to Claims 1 to 4, characterized in that said hinge region comprises all or part of the D hinge region of a nuclear estrogen receptor, a region situated between the DNA-binding domain and the ligand-binding domain, or a peptide
25 which is functionally equivalent to said D hinge region.

6. Organism according to Claim 5, characterized in that said hinge region comprises amino acids 282 to 301 of the sequence SEQ ID No. 2.

7. Organism according to one of the preceding claims, characterized in that said polypeptide chosen from the ligand-binding domain of the nuclear human estrogen receptors or its natural variants, or one of their fragments, is the ligand-binding domain of the human nuclear estrogen receptor α or one of its natural variants, or one of their fragments and in that said ligand-binding domain or one of its natural variants, or one of their fragments exhibits at least one mutation chosen from the group:

- mutation (G521R) glycine to arginine at position 521 of the sequence SEQ ID No. 2 or of a natural variant of this sequence;
- mutation (G400V) glycine to valine at position 400 of the sequence SEQ ID No. 2 or of a natural variant of this sequence;
- mutation (methionine-leucine) to (alanine-alanine) situated at position 543-544 (M543A/L544A mutation) of the sequence SEQ ID No. 2 or of a natural variant of this sequence.

8. Organism according to any one of the preceding claims, characterized in that said fusion protein is encoded by a fusion gene integrated into one or more of the chromosomes of said cell of said organism, said fusion gene being under the control of expression elements ensuring its expression in at least one cell of said organism.

9. Organism according to any one of Claims 1 to 7, characterized in that said fusion protein is encoded by a fusion gene integrated into an extrachromosomal

expression vector, said fusion gene being under the control of expression elements ensuring its expression in at least one cell of said organism.

10. Organism according to either of Claims 8 and 9,
5 characterized in that said expression elements are chosen from elements controlling tissue-specific and cell-specific expression or ubiquitous expression.

11. Organism according to Claims 8 to 10,
characterized in that said elements controlling
10 expression are chosen from elements controlling expression ensuring constitutive expression or elements controlling expression ensuring inducible expression.

12. Organism according to Claims 8 to 11,
characterized in that said expression element is chosen
15 from the group composed of the promoter regions of cytokeratin 14 (K 14), of cytokeratin 5 (K 5), of the adipocyte fatty acid binding protein 2 (aP2) and of α -1-antitrypsin.

13. Organism according to one of Claims 8 to 12,
20 characterized in that said fusion gene having the sequence SEQ ID No. 3 encodes the fusion protein Cre-ER^T having the sequence SEQ ID No. 4.

14. Organism according to any one of Claims 8 to 12, characterized in that said fusion gene having the
25 sequence SEQ ID No. 5 encodes the fusion protein Cre-ER^{T2} having the sequence SEQ ID No. 6.

15. Organism according to any one of Claims 8 to 12, characterized in that said fusion gene having the
sequence SEQ ID No. 7 encodes the fusion protein
30 Cre-ER^{T3} having the sequence SEQ ID No. 8.

16. Organism according to Claims 9 to 15, characterized in that said fusion gene preferably comprises in the 5' → 3' direction:
- a DNA fragment encoding the Cre recombinase of bacteriophage P1 or one of its variants;
 - a DNA fragment of at least 45 nucleotides encoding at least either all or part of the D hinge region of a nuclear estrogen receptor, or a peptide which is functionally equivalent to said D hinge region; and
 - a DNA fragment encoding the ligand-binding domain (LBD) of a nuclear estrogen receptor or variants thereof, said DNA fragment having at least one mutation conferring on LBD the capacity to respond to synthetic antiestrogens, but not to natural estrogenic agonists.
17. Organism according to any one of Claims 1 to 7, characterized in that said fusion protein is introduced into at least one cell of said organism.
18. Organism according to any one of Claims 1 to 17, characterized in that said synthetic ligand endowed with antiestrogenic activity inducing the activity of the recombinase is chosen from the group composed of Tamoxifen, 4-hydroxyTamoxifen, ICI 164 384 and ICI 182 780.
19. Organism according to any one of the preceding claims, characterized in that said DNA sequence of interest is a gene selected from the group composed of RXR α , RXR β , RXR γ , RAR α , RAR β , RAR γ , SNF2 β .

20. Organism according to any one of the preceding claims, characterized in that said organism is an animal, in particular a mouse.

21. Organism according to Claim 20, characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the cytokeratin K5 promoter region;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

22. Organism according to Claim 20, characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the cytokeratin K14 promoter region;
- one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.

23. Organism according to Claim 20, characterized in that at least one of the cells of said mouse comprises:

- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2} having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being
5 under the control of the adipocyte fatty acid binding protein 2 (aP2) promoter region;
 - one or more chromosomal DNA sequences of interest in their natural chromatin context and flanked ("floxed") by a lox site.
- 10 24. Organism according to Claim 20, characterized in that at least one of the cells of said mouse comprises:
- a fusion gene encoding the fusion protein Cre-ER^T having the sequence SEQ ID No. 4, or Cre-ER^{T2}
15 having the sequence ID No. 6, or Cre-ER^{T3} having the sequence ID No. 8, said fusion gene being under the control of the α -1-antitrypsin promoter region;
 - one or more chromosomal DNA sequences of interest
20 in their natural chromatin context and flanked ("floxed") by a lox site.
25. Method of preparing a metazoan organism according to any one of the preceding claims, characterized in that it comprises the following steps:
- 25 a) obtaining an embryonic stem (ES) cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;

- b) introducing said modified embryonic stem cell into an embryo of said organism;
- c) developing said embryo up to the stage of a fertile adult organism;
- 5 d) crossing said fertile adult organism with a transgenic organism in which at least one of the cells expresses said fusion protein and obtaining the progeny derived from said crossing; and
- e) optionally, selecting, among said progeny, said
10 metazoan organism.

26. Method of preparing a metazoan organism according to any one of Claims 1 to 24, characterized in that it comprises the following steps:

- a) obtaining a somatic cell modified by insertion of
15 site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;
- b) transferring the nucleus of said modified somatic
20 cell into the cytoplasm of an enucleated recipient oocyte;
- c) developing the embryo obtained in step b) up to the stage of a fertile adult organism;
- d) crossing said fertile adult organism with a
25 transgenic organism in which at least one of the cells expresses said fusion protein and obtaining the progeny derived from said crossing; and
- e) optionally, selecting, among the progeny, said metazoan organism.

27. Method of preparing a metazoan organism according to any one of Claims 1 to 24, characterized in that it comprises the following steps:

- 5 a) obtaining an embryonic stem (ES) cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;
- 10 b) introducing said modified embryonic stem cell into an embryo of said organism;
- c) developing said embryo; and
- d) introducing said fusion protein into at least one cell of said embryo or of the organism obtained from the development of said embryo.

15 28. Method of preparing a metazoan organism according to any one of Claims 1 to 24, characterized in that it comprises the following steps:

- 20 a) obtaining a somatic cell modified by insertion of site(s) of recognition for said recombinase protein into said DNA sequence(s) of interest, located in one or more chromosomes, by homologous recombination;
- b) transferring the nucleus of said modified somatic cell into the cytoplasm of an enucleated recipient
25 oocyte;
- c) developing said embryo; and
- d) introducing said fusion protein into at least one cell of said embryo or of said organism obtained from the development of said embryo.

29. Method of conditional recombination, in particular excision, insertion, inversion, translocation, at the level of the DNA sequence of interest into which there is (are) inserted one or more sites of recognition for said recombinase protein, said DNA sequence of interest being located in one or more of the chromosomes of said genome of said cell of said organism according to Claims 1 to 24, characterized in that it comprises the steps of:
- 5 (i) bringing at least one cell of said organism into contact with a synthetic ligand endowed with antiestrogenic activity;
- (ii) inducing the activity of the recombinase of said fusion protein by said synthetic ligand.
- 15 30. Method of conditional deletion of a DNA fragment in which a method of excision according to Claim 29 is used and in which said DNA fragment(s) to be excised is (are) flanked by two recombinase protein recognition sites oriented as a direct repeat.
- 20 31. Method of obtaining a metazoan organism, with the exception of humans, in which at least one cell possesses an allele of a gene of interest inactivated by a method of conditional deletion and in which the other allele of said gene of interest possesses a mutation, preferably limited, in exon and/or regulatory sequences, said method being characterized in that it makes it possible to obtain, in a metazoan organism, somatic mutations controlled in space and time, and which are limited (point mutations, small deletions or
- 25

insertion) in exon and/or regulatory sequences, and in that it comprises the steps of:

- a) obtaining a metazoan organism in which at least one cell of the germ line comprises said mutation
5 in one of the alleles of said gene of interest;
- b) crossing said organism obtained in step a) with an organism according to any one of Claims 1 to 24;
- c) selecting a progeny whose genome comprises a gene
10 of interest in which one of the alleles possesses a mutation and the other allele possesses at least two recombinase protein recognition sites oriented as a direct repeat;
- d) using the method according to Claims 29 or 30 of conditional deletion, of the DNA fragment of said
15 allele of said gene of interest which is flanked by at least two recombinase protein recognition sites oriented as a direct repeat; and
- e) obtaining said metazoan organism in which the genome of at least one cell comprises said gene of
20 interest in which one allele is inactivated, while the other allele possesses a somatic, preferably limited, mutation and preferably in exon and/or regulatory sequences.

32. Method according to any one of Claims 29 to 31,
25 characterized in that said sites of recognition specific for the recombinase protein are Lox P sites and said recombinase protein is the Cre protein of bacteriophage P1, or one of its variants.

33. Organism capable of being obtained using a
30 method according to Claims 25 to 32.

34. Organism according to Claim 33, characterized in that it is an animal, in particular a mouse.

35. Method of analyzing or studying the biological function of a DNA sequence of interest, in particular
5 of a gene, characterized in that it comprises the steps of:

- (i) bringing an organism according to one of Claims 1 to 24, 33 and 34 or cells isolated from said organism into contact with a synthetic ligand
10 endowed with antiestrogenic activity;
- (ii) optionally inducing the expression of said fusion protein;
- (iii) revealing the recombination event catalyzed by the recombinase activity of said fusion
15 protein;
- (iv) biochemical and/or physiological and/or phenotypic and/or behavioral study or analysis of said cell or of said organism.

36. Method according to Claims 29, 30, 31, 32 and
20 35, characterized in that the bringing of said cells of said organism into contact with said synthetic ligand is carried out according to a route of administration chosen from the oral route, the topical route, injection, in particular intramuscular, intravenous,
25 intracerebral, intraspinal and intraperitoneal injection, or in the case of embryos, fetuses and neonates before weaning by administering said synthetic ligand to the mother.

37. Method of screening compounds capable of being
30 used as a medicament for the preventive and/or curative

treatment of pathological conditions associated with alteration of the expression and/or of the function of said DNA sequence of interest, characterized in that it comprises the step of administering said compound to an
5 organism according to claims 1 to 24, 33 and 34.

38. Use of an organism according to any one of Claims 1 to 24, 33 and 34 or of cells derived from said organism for carrying out a spatiotemporally controlled site-specific recombination of said DNA sequence of
10 interest in its natural chromatin environment, with an efficiency of at least 85%, in the presence of synthetic ligand endowed with antiestrogenic activity in the cells of said organism expressing said fusion protein, and with an efficiency at least lower than 5%,
15 in the absence of synthetic ligand or in the presence of a natural estrogen in the cells of said organism expressing said fusion protein.

39. Use according to Claim 38, characterized in that said cells of said organism are chosen from the
20 cells of the epidermis, the hepatocytes and the adipocytes.

40. Transgenic mouse K5-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional
25 deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal keratinocytes and/or an inflammatory reaction of the skin.

41. Transgenic mouse K5-Cre-ER^{T2}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic
5 ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal keratinocytes and/or an inflammatory reaction of the skin.
42. Transgenic mouse K5-Cre-ER^{T3}/RXR_α^{L2/L2} whose RXR_α
10 gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the
15 basal keratinocytes and/or an inflammatory reaction of the skin.
43. Transgenic mouse K14-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional
20 deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal keratinocytes and/or an inflammatory reaction of the skin.
- 25 44. Transgenic mouse K14-Cre-ER^{T2}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic
30 ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the

basal keratinocytes and/or an inflammatory reaction of the skin.

45. Transgenic mouse K14-Cre-ER^{T3}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the basal
5 keratinocytes of the epidermis using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alopecia and/or hyperproliferation of the basal keratinocytes and/or an inflammatory reaction of
10 the skin.

46. Transgenic mouse αAT-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the hepatocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic
15 activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

47. Transgenic mouse αAT-Cre-ER^{T2}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the hepatocytes using a conditional deletion method following treatment
20 with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

48. Transgenic mouse αAT-Cre-ER^{T3}/RXR_α^{L2/L2} whose RXR_α gene may be selectively inactivated in the hepatocytes
25 using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse in particular alteration of the proliferation of the hepatocytes.

49. Transgenic mouse aP2-Cre-ER^T/RXR_α^{L2/L2} whose RXR_α
30 gene may be selectively inactivated in the adipocytes

using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alteration of the metabolism of the lipids in the adipocytes and/or
5 diabetes.

50. Transgenic mouse $\text{aP2-Cre-ER}^{\text{T2}}/\text{RXR}_{\alpha}^{\text{L2/L2}}$ whose RXR_{α} gene may be selectively inactivated in the adipocytes using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic
10 activity, causing in said mouse alteration of the metabolism of the lipids in the adipocytes and/or diabetes.

51. Transgenic mouse $\text{aP2-Cre-ER}^{\text{T3}}/\text{RXR}_{\alpha}^{\text{L2/L2}}$ whose RXR_{α} gene may be selectively inactivated in the adipocytes
15 using a conditional deletion method following treatment with a synthetic ligand endowed with antiestrogenic activity, causing in said mouse alteration of the metabolism of the lipids in the adipocytes and/or diabetes.

20 52. Transgenic mouse according to one of Claims 40 to 51, characterized in that said RXR_{α} gene is inactivated by carrying out the method according to Claim 30.

53. Method of screening compounds capable of being
25 used as a medicament for the preventive and/or curative treatment of alopecia and/or of hyperproliferation of the keratinocytes and/or of inflammatory reactions of the skin, characterized in that it comprises the step of administering said compound to a mouse according to
30 Claims 40 to 45 and 52.

54. Method of screening compounds capable of being used as a medicament for promoting in particular hepatic regeneration, characterized in that it comprises the step of administering said compound to a
5 mouse according to Claims 46 to 48 and 52.

55. Method of screening compounds capable of being used as a medicament for the preventive and/or curative treatment of diabetes and/or for the treatment of obesity, characterized in that it comprises the step of
10 administering said compound to a mouse according to Claims 49 to 52.

56. Method of screening compounds capable of being used as a medicament for the preventive and/or curative treatment of skin cancers, characterized in that it
15 comprises the step of administering said compound to a mouse according to Claims 40 to 45 and 52.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 2A




FIG. 2B

5 / 21




FIG. 3 .

6 / 21




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6E




FIG. 6F




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

15 / 21




FIG. 12A-B




FIG. 12C-D




FIG. 13A-B




FIG. 14A-B




FIG. 14C




FIG. 15

SEQUENCE LISTING

<110> ASSOCIATION POUR LE DEVELOPPEMENT DE LA
RECHERCHE EN GENETIQUE MOLECULAIRE (ADEREGEM)

<120> Transgenic mouse for targeted recombination
mediated by modified Cre-ER

<130> D18961

<150> FR 00 12 570

<151> 2000-10-03

<160> 14

<170> PatentIn Ver. 2.1

<210> 1

<211> 1788

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1788)

<400> 1

atg acc atg acc ctc cac acc aaa gca tct ggg atg gcc cta ctg cat	48
Met Thr Met Thr Leu His Thr Lys Ala Ser Gly Met Ala Leu Leu His	
1 5 10 15	
cag atc caa ggg aac gag ctg gag ccc ctg aac cgt ccg cag ctc aag	96
Gln Ile Gln Gly Asn Glu Leu Glu Pro Leu Asn Arg Pro Gln Leu Lys	
20 25 30	
atc ccc ctg gag cgg ccc ctg ggc gag gtg tac ctg gac agc agc aag	144
Ile Pro Leu Glu Arg Pro Leu Gly Glu Val Tyr Leu Asp Ser Ser Lys	
35 40 45	
ccc gcc gtg tac aac tac ccc gag ggc gcc gcc tac gag ttc aac gcc	192
Pro Ala Val Tyr Asn Tyr Pro Glu Gly Ala Ala Tyr Glu Phe Asn Ala	
50 55 60	
gcg gcc gcc gcc aac gcg cag gtc tac ggt cag acc ggc ctc ccc tac	240
Ala Ala Ala Ala Asn Ala Gln Val Tyr Gly Gln Thr Gly Leu Pro Tyr	
65 70 75 80	
ggc ccc ggg tct gag gct gcg gcg ttc ggc tcc aac ggc ctg ggg ggt	288
Gly Pro Gly Ser Glu Ala Ala Phe Gly Ser Asn Gly Leu Gly Gly	
85 90 95	
ttc ccc cca ctc aac agc gtg tct ccg agc ccg ctg atg cta ctg cac	336
Phe Pro Pro Leu Asn Ser Val Ser Pro Ser Pro Leu Met Leu Leu His	
100 105 110	
ccg ccg ccg cag ctg tcg cct ttc ctg cag ccc cac ggc cag cag gtg	384
Pro Pro Pro Gln Leu Ser Pro Phe Leu Gln Pro His Gly Gln Gln Val	
115 120 125	
ccc tac tac ctg gag aac gag ccc agc ggc tac acg gtg cgc gag gcc	432
Pro Tyr Tyr Leu Glu Asn Glu Pro Ser Gly Tyr Thr Val Arg Glu Ala	

130	135	140	
ggc ccg ccg gca ttc tac agg cca aat tca gat aat cga cgc cag ggt Gly Pro Pro Ala Phe Tyr Arg Pro Asn Ser Asp Asn Arg Arg Gln Gly 145 150 155 160			480
ggc aga gaa aga ttg gcc agt acc aat gac aag gga agt atg gct atg Gly Arg Glu Arg Leu Ala Ser Thr Asn Asp Lys Gly Ser Met Ala Met 165 170 175			528
gaa tct gcc aag gag act cgc tac tgt gca gtg tgc aat gac tat gct Glu Ser Ala Lys Glu Thr Arg Tyr Cys Ala Val Cys Asn Asp Tyr Ala 180 185 190			576
tca ggc tac cat tat gga gtc tgg tcc tgt gag ggc tgc aag gcc ttc Ser Gly Tyr His Tyr Gly Val Trp Ser Cys Glu Gly Cys Lys Ala Phe 195 200 205			624
ttc aag aga agt att caa gga cat aac gac tat atg tgt cca gcc acc Phe Lys Arg Ser Ile Gln Gly His Asn Asp Tyr Met Cys Pro Ala Thr 210 215 220			672
aac cag tgc acc att gat aaa aac agg agg aag agc tgc cag gcc tgc Asn Gln Cys Thr Ile Asp Lys Asn Arg Arg Lys Ser Cys Gln Ala Cys 225 230 235			720
cgg ctc cgc aaa tgc tac gaa gtg gga atg atg aaa ggt ggg ata cga Arg Leu Arg Lys Cys Tyr Glu Val Gly Met Met Lys Gly Gly Ile Arg 245 250 255			768
aaa gac cga aga gga ggg aga atg ttg aaa cac aag cgc cag aga gat Lys Asp Arg Arg Gly Gly Arg Met Leu Lys His Lys Arg Gln Arg Asp 260 265 270			816
gat ggg gag ggc agg ggt gaa gtg ggg tct gct gga gac atg aga gct Asp Gly Glu Gly Arg Gly Glu Val Gly Ser Ala Gly Asp Met Arg Ala 275 280 285			864
gcc aac ctt tgg cca agc ccg ctc atg atc aaa cgc tct aag aag aac Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys Asn 290 295 300			912
agc ctg gcc ttg tcc ctg acg gcc gac cag atg gtc agt gcc ttg ttg Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu Leu 305 310 315 320			960
gat gct gag ccc ccc ata ctc tat tcc gag tat gat cct acc aga ccc Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg Pro 325 330 335			1008
ttc agt gaa gct tcg atg atg ggc tta ctg acc aac ctg gca gac agg Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp Arg 340 345 350			1056
gag ctg gtt cac atg atc aac tgg gcg aag agg gtg cca ggc ttt gtg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe Val 355 360 365			1104
gat ttg acc ctc cat gat cag gtc cac ctt cta gaa tgt gcc tgg cta Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp Leu 370 375 380			1152

gag atc ctg atg att ggt ctc gtc tgg cgc tcc atg gag cac cca ggg	1200
Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro Gly	
385 390 395 400	
aag cta ctg ttt gct cct aac ttg ctc ttg gac agg aac cag gga aaa	1248
Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly Lys	
405 410 415	
tgt gta gag ggc atg gtg gag atc ttc gac atg ctg ctg gct aca tca	1296
Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr Ser	
420 425 430	
tct cgg ttc cgc atg atg aat ctg cag gga gag gag ttt gtg tgc ctc	1344
Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys Leu	
435 440 445	
aaa tct att att ttg ctt aat tct gga gtg tac aca ttt ctg tcc agc	1392
Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser Ser	
450 455 460	
acc ctg aag tct ctg gaa gag aag gac cat atc cac cga gtc ctg gac	1440
Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu Asp	
465 470 475 480	
aag atc aca gac act ttg atc cac ctg atg gcc aag gca ggc ctg acc	1488
Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu Thr	
485 490 495	
ctg cag cag cag cac cag cgg ctg gcc cag ctc ctc ctc atc ctc tcc	1536
Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu Ser	
500 505 510	
cac atc agg cac atg agt aac aaa ggc atg gag cat ctg tac agc atg	1584
His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser Met	
515 520 525	
aag tgc aag aac gtg gtg ccc ctc tat gac ctg ctg ctg gag atg ctg	1632
Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Met Leu	
530 535 540	
gac gcc cac cgc cta cat gcg ccc act agc cgt gga ggg gca tcc gtg	1680
Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser Val	
545 550 555 560	
gag gag acg gac caa agc cac ttg gcc act gcg ggc tct act tca tcg	1728
Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser Ser	
565 570 575	
cat tcc ttg caa aag tat tac atc acg ggg gag gca gag ggt ttc cct	1776
His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe Pro	
580 585 590	
gcc aca gtc tga	1788
Ala Thr Val	
595	

<210> 2
 <211> 595
 <212> PRT

<213> Homo sapiens

<400> 2

```

Met Thr Met Thr Leu His Thr Lys Ala Ser Gly Met Ala Leu Leu His
 1           5           10           15

Gln Ile Gln Gly Asn Glu Leu Glu Pro Leu Asn Arg Pro Gln Leu Lys
 20           25           30

Ile Pro Leu Glu Arg Pro Leu Gly Glu Val Tyr Leu Asp Ser Ser Lys
 35           40           45

Pro Ala Val Tyr Asn Tyr Pro Glu Gly Ala Ala Tyr Glu Phe Asn Ala
 50           55           60

Ala Ala Ala Ala Asn Ala Gln Val Tyr Gly Gln Thr Gly Leu Pro Tyr
 65           70           75           80

Gly Pro Gly Ser Glu Ala Ala Ala Phe Gly Ser Asn Gly Leu Gly Gly
 85           90           95

Phe Pro Pro Leu Asn Ser Val Ser Pro Ser Pro Leu Met Leu Leu His
100           105           110

Pro Pro Pro Gln Leu Ser Pro Phe Leu Gln Pro His Gly Gln Gln Val
115           120           125

Pro Tyr Tyr Leu Glu Asn Glu Pro Ser Gly Tyr Thr Val Arg Glu Ala
130           135           140

Gly Pro Pro Ala Phe Tyr Arg Pro Asn Ser Asp Asn Arg Arg Gln Gly
145           150           155           160

Gly Arg Glu Arg Leu Ala Ser Thr Asn Asp Lys Gly Ser Met Ala Met
165           170           175

Glu Ser Ala Lys Glu Thr Arg Tyr Cys Ala Val Cys Asn Asp Tyr Ala
180           185           190

Ser Gly Tyr His Tyr Gly Val Trp Ser Cys Glu Gly Cys Lys Ala Phe
195           200           205

Phe Lys Arg Ser Ile Gln Gly His Asn Asp Tyr Met Cys Pro Ala Thr
210           215           220

Asn Gln Cys Thr Ile Asp Lys Asn Arg Arg Lys Ser Cys Gln Ala Cys
225           230           235           240

Arg Leu Arg Lys Cys Tyr Glu Val Gly Met Met Lys Gly Gly Ile Arg
245           250           255

Lys Asp Arg Arg Gly Gly Arg Met Leu Lys His Lys Arg Gln Arg Asp
260           265           270

Asp Gly Glu Gly Arg Gly Glu Val Gly Ser Ala Gly Asp Met Arg Ala
275           280           285

Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys Asn
290           295           300

Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu Leu

```

305		310		315		320
Asp Ala Glu Pro	Pro Ile Leu Tyr Ser	Glu Tyr Asp Pro Thr Arg Pro				
	325	330			335	
Phe Ser Glu Ala	Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp Arg					
	340	345			350	
Glu Leu Val His	Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe Val					
	355	360			365	
Asp Leu Thr Leu His	Asp Gln Val His Leu Leu Glu Cys Ala Trp Leu					
	370	375			380	
Glu Ile Leu Met Ile	Gly Leu Val Trp Arg Ser Met Glu His Pro Gly					
	385	390			395	400
Lys Leu Leu Phe Ala	Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly Lys					
	405	410			415	
Cys Val Glu Gly Met	Val Glu Ile Phe Asp Met Leu Leu Ala Thr Ser					
	420	425			430	
Ser Arg Phe Arg Met	Met Asn Leu Gln Gly Glu Glu Phe Val Cys Leu					
	435	440			445	
Lys Ser Ile Ile Leu	Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser Ser					
	450	455			460	
Thr Leu Lys Ser Leu	Glu Glu Lys Asp His Ile His Arg Val Leu Asp					
	465	470			475	480
Lys Ile Thr Asp Thr	Leu Ile His Leu Met Ala Lys Ala Gly Leu Thr					
	485	490			495	
Leu Gln Gln Gln His	Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu Ser					
	500	505			510	
His Ile Arg His Met	Ser Asn Lys Gly Met Glu His Leu Tyr Ser Met					
	515	520			525	
Lys Cys Lys Asn Val	Val Pro Leu Tyr Asp Leu Leu Leu Glu Met Leu					
	530	535			540	
Asp Ala His Arg Leu	His Ala Pro Thr Ser Arg Gly Gly Ala Ser Val					
	545	550			555	560
Glu Glu Thr Asp Gln	Ser His Leu Ala Thr Ala Gly Ser Thr Ser Ser					
	565	570			575	
His Ser Leu Gln Lys	Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe Pro					
	580	585			590	
Ala Thr Val						
	595					

<210> 3

<211> 1983

<212> DNA

<213> Artificial sequence

<220>

<221> CDS

<222> (1)..(1983)

<220>

<223> Chimeric sequence Homosapiens-Bacteriophage P1

<400> 3

atg tcc aat tta ctg acc gta cac caa aat ttg cct gca tta ccg gtc	48
Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val	
1 5 10 15	
gat gca acg agt gat gag gtt cgc aag aac ctg atg gac atg ttc agg	96
Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg	
20 25 30	
gat cgc cag gcg ttt tct gag cat acc tgg aaa atg ctt ctg tcc gtt	144
Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val	
35 40 45	
tgc cgg tcg tgg gcg gca tgg tgc aag ttg aat aac cgg aaa tgg ttt	192
Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe	
50 55 60	
ccc gca gaa cct gaa gat gtt cgc gat tat ctt cta tat ctt cag gcg	240
Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala	
65 70 75 80	
cgc ggt ctg gca gta aaa act atc cag caa cat ttg ggc cag cta aac	288
Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn	
85 90 95	
atg ctt cat cgt cgg tcc ggg ctg cca cga cca agt gac agc aat gct	336
Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala	
100 105 110	
gtt tca ctg gtt atg cgg cgg atc cga aaa gaa aac gtt gat gcc ggt	384
Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly	
115 120 125	
gaa cgt gca aaa cag gct cta gcg ttc gaa cgc act gat ttc gac cag	432
Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln	
130 135 140	
gtt cgt tca ctc atg gaa aat agc gat cgc tgc cag gat ata cgt aat	480
Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn	
145 150 155 160	
ctg gca ttt ctg ggg att gct tat aac acc ctg tta cgt ata gcc gaa	528
Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu	
165 170 175	
att gcc agg atc agg gtt aaa gat atc tca cgt act gac ggt ggg aga	576
Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg	
180 185 190	
atg tta atc cat att ggc aga acg aaa acg ctg gtt agc acc gca ggt	624
Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly	
195 200 205	

gta gag aag gca ctt agc ctg ggg gta act aaa ctg gtc gag cga tgg	722
Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp	
210 215 220	
att tcc gtc tct ggt gta gct gat gat ccg aat aac tac ctg ttt tgc	720
Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys	
225 230 235 240	
cgg gtc aga aaa aat ggt gtt gcc gcg cca tct gcc acc agc cag cta	768
Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu	
245 250 255	
tca act cgc gcc ctg gaa ggg att ttt gaa gca act cat cga ttg att	816
Ser Thr Arg Ala Leu Glu Gly Ile Phe Glu Ala Thr His Arg Leu Ile	
260 265 270	
tac ggc gct aag gat gac tct ggt cag aga tac ctg gcc tgg tct gga	864
Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly	
275 280 285	
cac agt gcc cgt gtc gga gcc gcg cga gat atg gcc cgc gct gga gtt	912
His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val	
290 295 300	
tca ata ccg gag atc atg caa gct ggt ggc tgg acc aat gta aat att	960
Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile	
305 310 315 320	
gtc atg aac tat atc cgt aac ctg gat agt gaa aca ggg gca atg gtg	1008
Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val	
325 330 335	
cgc ctg ctg gaa gat ggc gat ctc gag cca tct gct gga gac atg aga	1056
Arg Leu Leu Glu Asp Gly Asp Leu Glu Pro Ser Ala Gly Asp Met Arg	
340 345 350	
gct gcc aac ctt tgg cca agc ccg ctc atg atc aaa cgc tct aag aag	1104
Ala Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys	
355 360 365	
aac agc ctg gcc ttg tcc ctg acg gcc gac cag atg gtc agt gcc ttg	1152
Asn Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu	
370 375 380	
ttg gat gct gag ccc ccc ata ctc tat tcc gag tat gat cct acc aga	1200
Leu Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg	
385 390 395 400	
ccc ttc agt gaa gct tcg atg atg ggc tta ctg acc aac ctg gca gac	1248
Pro Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp	
405 410 415	
agg gag ctg gtt cac atg atc aac tgg gcg aag agg gtg cca ggc ttt	1296
Arg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe	
420 425 430	
gtg gat ttg acc ctc cat gat cag gtc cac ctt cta gaa tgt gcc tgg	1344
Val Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp	
435 440 445	

cta gag atc ctg atg att ggt ctc gtc tgg cgc tcc atg gag cac cca	1392
Leu Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro	
450 455 460	
ggg aag cta ctg ttt gct cct aac ttg ctc ttg gac agg aac cag gga	1440
Gly Lys Leu Leu Phe Ala Pro Asn Leu Leu Asp Arg Asn Gln Gly	
465 470 475 480	
aaa tgt gta gag ggc atg gtg gag atc ttc gac atg ctg ctg gct aca	1488
Lys Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr	
485 490 495	
tca tct cgg ttc cgc atg atg aat ctg cag gga gag gag ttt gtg tgc	1536
Ser Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys	
500 505 510	
ctc aaa tct att att ttg ctt aat tct gga gtg tac aca ttt ctg tcc	1584
Leu Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser	
515 520 525	
agc acc ctg aag tct ctg gaa gag aag gac cat atc cac cga gtc ctg	1632
Ser Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu	
530 535 540	
gac aag atc aca gac act ttg atc cac ctg atg gcc aag gca ggc ctg	1680
Asp Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu	
545 550 555 560	
acc ctg cag cag cag cac cag cgg ctg gcc cag ctc ctc ctc atc ctc	1728
Thr Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu	
565 570 575	
tcc cac atc agg cac atg agt aac aaa aga atg gag cat ctg tac agc	1776
Ser His Ile Arg His Met Ser Asn Lys Arg Met Glu His Leu Tyr Ser	
580 585 590	
atg aag tgc aag aac gtg gtg ccc ctc tat gac ctg ctg ctg gag atg	1824
Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Met	
595 600 605	
ctg gac gcc cac cgc cta cat gcg ccc act agc cgt gga ggg gca tcc	1872
Leu Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser	
610 615 620	
gtg gag gag acg gac caa agc cac ttg gcc act gcg ggc tct act tca	1920
Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser	
625 630 635 640	
tcg cat tcc ttg caa aag tat tac atc acg ggg gag gca gag ggt ttc	1968
Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe	
645 650 655	
cct gcc aca gtc tga	1983
Pro Ala Thr Val	
660	

<210> 4

<211> 660

<212> PRT

<213> Artificial sequence

<223> Chimeric sequence Homosapiens-Bacteriophage P1

<400> 4

```

Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val
 1           5           10           15

Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg
      20           25           30

Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val
      35           40           45

Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe
      50           55           60

Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala
      65           70           75           80

Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn
      85           90           95

Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala
      100          105          110

Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly
      115          120          125

Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln
      130          135          140

Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn
      145          150          155          160

Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu
      165          170          175

Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg
      180          185          190

Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly
      195          200          205

Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp
      210          215          220

Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys
      225          230          235          240

Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu
      245          250          255

Ser Thr Arg Ala Leu Glu Gly Ile Phe Glu Ala Thr His Arg Leu Ile
      260          265          270

Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly
      275          280          285

His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val
      290          295          300

Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile

```

305	310	315	320
Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val			
	325	330	335
Arg Leu Leu Glu Asp Gly Asp Leu Glu Pro Ser Ala Gly Asp Met Arg			
	340	345	350
Ala Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys			
	355	360	365
Asn Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu			
	370	375	380
Leu Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg			
385	390	395	400
Pro Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp			
	405	410	415
Arg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe			
	420	425	430
Val Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp			
	435	440	445
Leu Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro			
	450	455	460
Gly Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly			
465	470	475	480
Lys Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr			
	485	490	495
Ser Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys			
	500	505	510
Leu Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser			
	515	520	525
Ser Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu			
	530	535	540
Asp Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu			
545	550	555	560
Thr Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu			
	565	570	575
Ser His Ile Arg His Met Ser Asn Lys Arg Met Glu His Leu Tyr Ser			
	580	585	590
Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Met			
	595	600	605
Leu Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser			
	610	615	620
Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser			
625	630	635	640

Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe
 645 650 655

Pro Ala Thr Val
 660

<210> 5

<211> 1983

<212> DNA

<213> Artificial sequence

<220>

<221> CDS

<222> (1)..(1983)

<220>

<223> Chimeric sequence Homosapiens-Bacteriophage P1

<400> 5

atg tcc aat tta ctg acc gta cac caa aat ttg cct gca tta ccg gtc	48
Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val	
1 5 10 15	
gat gca acg agt gat gag gtt cgc aag aac ctg atg gac atg ttc agg	96
Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg	
20 25 30	
gat cgc cag gcg ttt tct gag cat acc tgg aaa atg ctt ctg tcc gtt	144
Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val	
35 40 45	
tgc cgg tcg tgg gcg gca tgg tgc aag ttg aat aac cgg aaa tgg ttt	192
Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe	
50 55 60	
ccc gca gaa cct gaa gat gtt cgc gat tat ctt cta tat ctt cag gcg	240
Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala	
65 70 75 80	
cgc ggt ctg gca gta aaa act atc cag caa cat ttg ggc cag cta aac	288
Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn	
85 90 95	
atg ctt cat cgt cgg tcc ggg ctg cca cga cca agt gac agc aat gct	336
Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala	
100 105 110	
gtt tca ctg gtt atg cgg cgg atc cga aaa gaa aac gtt gat gcc ggt	384
Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly	
115 120 125	
gaa cgt gca aaa cag gct cta gcg ttc gaa cgc act gat ttc gac cag	432
Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln	
130 135 140	
gtt cgt tca ctc atg gaa aat agc gat cgc tgc cag gat ata cgt aat	480
Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn	
145 150 155 160	

ctg gca ttt ctg ggg att gct tat aac acc ctg tta cgt ata gcc gaa	528
Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu	
165 170 175	
att gcc agg atc agg gtt aaa gat atc tca cgt act gac ggt ggg aga	576
Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg	
180 185 190	
atg tta atc cat att ggc aga acg aaa acg ctg gtt agc acc gca ggt	624
Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly	
195 200 205	
gta gag aag gca ctt agc ctg ggg gta act aaa ctg gtc gag cga tgg	672
Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp	
210 215 220	
att tcc gtc tct ggt gta gct gat gat ccg aat aac tac ctg ttt tgc	720
Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys	
225 230 235 240	
cgg gtc aga aaa aat ggt gtt gcc gcg cca tct gcc acc agc cag cta	768
Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu	
245 250 255	
tca act cgc gcc ctg gaa ggg att ttt gaa gca act cat cga ttg att	816
Ser Thr Arg Ala Leu Glu Gly Ile Phe Glu Ala Thr His Arg Leu Ile	
260 265 270	
tac ggc gct aag gat gac tct ggt cag aga tac ctg gcc tgg tct gga	864
Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly	
275 280 285	
cac agt gcc cgt gtc gga gcc gcg cga gat atg gcc cgc gct gga gtt	912
His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val	
290 295 300	
tca ata ccg gag atc atg caa gct ggt ggc tgg acc aat gta aat att	960
Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile	
305 310 315 320	
gtc atg aac tat atc cgt aac ctg gat agt gaa aca ggg gca atg gtg	1008
Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val	
325 330 335	
cgc ctg ctg gaa gat ggc gat ctc gag cca tct gct gga gac atg aga	1056
Arg Leu Leu Glu Asp Gly Asp Leu Glu Pro Ser Ala Gly Asp Met Arg	
340 345 350	
gct gcc aac ctt tgg cca agc ccg ctc atg atc aaa cgc tct aag aag	1104
Ala Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys	
355 360 365	
aac agc ctg gcc ttg tcc ctg acg gcc gac cag atg gtc agt gcc ttg	1152
Asn Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu	
370 375 380	
ttg gat gct gag ccc ccc ata ctc tat tcc gag tat gat cct acc aga	1200
Leu Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg	
385 390 395 400	

ccc ttc agt gaa gct tcg atg atg ggc tta ctg acc aac ctg gca gac	1248
Pro Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp	
405 410 415	
agg gag ctg gtt cac atg atc aac tgg gcg aag agg gtg cca ggc ttt	1296
Arg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe	
420 425 430	
gtg gat ttg acc ctc cat gat cag gtc cac ctt cta gaa tgt gcc tgg	1344
Val Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp	
435 440 445	
cta gag atc ctg atg att ggt ctc gtc tgg cgc tcc atg gag cac cca	1392
Leu Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro	
450 455 460	
gtg aag cta ctg ttt gct cct aac ttg ctc ttg gac agg aac cag gga	1440
Val Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly	
465 470 475 480	
aaa tgt gta gag ggc atg gtg gag atc ttc gac atg ctg ctg gct aca	1488
Lys Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr	
485 490 495	
tca tct cgg ttc cgc atg atg aat ctg cag gga gag gag ttt gtg tgc	1536
Ser Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys	
500 505 510	
ctc aaa tct att att ttg ctt aat tct gga gtg tac aca ttt ctg tcc	1584
Leu Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser	
515 520 525	
agc acc ctg aag tct ctg gaa gag aag gac cat atc cac cga gtc ctg	1632
Ser Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu	
530 535 540	
gac aag atc aca gac act ttg atc cac ctg atg gcc aag gca ggc ctg	1680
Asp Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu	
545 550 555 560	
acc ctg cag cag cag cac cag cgg ctg gcc cag ctc ctc ctc atc ctc	1728
Thr Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu	
565 570 575	
tcc cac atc agg cac atg agt aac aaa ggc atg gag cat ctg tac agc	1776
Ser His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser	
580 585 590	
atg aag tgc aag aac gtg gtg ccc ctc tat gac ctg ctg ctg gag gcg	1824
Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Ala	
595 600 605	
gcg gac gcc cac cgc cta cat gcg ccc act agc cgt gga ggg gca tcc	1872
Ala Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser	
610 615 620	
gtg gag gag acg gac caa agc cac ttg gcc act gcg ggc tct act tca	1920
Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser	
625 630 635 640	
tcg cat tcc ttg caa aag tat tac atc acg ggg gag gca gag ggt ttc	1968

Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe
 645 650 655

cct gcc aca gct tga
 Pro Ala Thr Ala
 660

1983

<210> 6

<211> 660

<212> PRT

<213> Artificial sequence

<223> Chimeric sequence Homosapiens-Bacteriophage P1

<400> 6

Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val
 1 5 10 15

Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg
 20 25 30

Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val
 35 40 45

Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe
 50 55 60

Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala
 65 70 75 80

Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn
 85 90 95

Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala
 100 105 110

Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly
 115 120 125

Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln
 130 135 140

Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn
 145 150 155 160

Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu
 165 170 175

Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg
 180 185 190

Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly
 195 200 205

Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp
 210 215 220

Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys
 225 230 235 240

Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu

15

Ser His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser
 580 585 590

Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Ala
 595 600 605

Ala Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser
 610 615 620

Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser
 625 630 635 640

Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe
 645 650 655

Pro Ala Thr Ala
 660

<210> 7
 <211> 1983
 <212> DNA
 <213> Artificial sequence

<220>
 <221> CDS
 <222> (1)..(1983)

<220>
 <223> Chimeric sequence Homosapiens-Bacteriophage P1

<400> 7
 atg tcc aat tta ctg acc gta cac caa aat ttg cct gca tta ccg gtc 48
 Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val
 1 5 10 15

gat gca acg agt gat gag gtt cgc aag aac ctg atg gac atg ttc agg 96
 Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg
 20 25 30

gat cgc cag gcg ttt tct gag cat acc tgg aaa atg ctt ctg tcc gtt 144
 Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val
 35 40 45

tgc cgg tcg tgg gcg gca tgg tgc aag ttg aat aac cgg aaa tgg ttt 192
 Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe
 50 55 60

ccc gca gaa cct gaa gat gtt cgc gat tat ctt cta tat ctt cag gcg 240
 Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala
 65 70 75 80

cgc ggt ctg gca gta aaa act atc cag caa cat ttg ggc cag cta aac 288
 Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn
 85 90 95

atg ctt cat cgt cgg tcc ggg ctg cca cga cca agt gac agc aat gct 336
 Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala
 100 105 110

gtt tca ctg gtt atg cgg cgg atc cga aaa gaa aac gtt gat gcc ggt	384
Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly	
115 120 125	
gaa cgt gca aaa cag gct cta gcg ttc gaa cgc act gat ttc gac cag	432
Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln	
130 135 140	
gtt cgt tca ctc atg gaa aat agc gat cgc tgc cag gat ata cgt aat	480
Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn	
145 150 155 160	
ctg gca ttt ctg ggg att gct tat aac acc ctg tta cgt ata gcc gaa	528
Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu	
165 170 175	
att gcc agg atc agg gtt aaa gat atc tca cgt act gac ggt ggg aga	576
Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg	
180 185 190	
atg tta atc cat att ggc aga acg aaa acg ctg gtt agc acc gca ggt	624
Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly	
195 200 205	
gta gag aag gca ctt agc ctg ggg gta act aaa ctg gtc gag cga tgg	672
Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp	
210 215 220	
att tcc gtc tct ggt gta gct gat gat ccg aat aac tac ctg ttt tgc	720
Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys	
225 230 235 240	
cgg gtc aga aaa aat ggt gtt gcc gcg cca tct gcc acc agc cag cta	768
Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu	
245 250 255	
tca act cgc gcc ctg gaa ggg att ttt gaa gca act cat cga ttg att	816
Ser Thr Arg Ala Leu Glu Gly Ile Phe Glu Ala Thr His Arg Leu Ile	
260 265 270	
tac ggc gct aag gat gac tct ggt cag aga tac ctg gcc tgg tct gga	864
Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly	
275 280 285	
cac agt gcc cgt gtc gga gcc gcg cga gat atg gcc cgc gct gga gtt	912
His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val	
290 295 300	
tca ata ccg gag atc atg caa gct ggt ggc tgg acc aat gta aat att	960
Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile	
305 310 315 320	
gtc atg aac tat atc cgt aac ctg gat agt gaa aca ggg gca atg gtg	1008
Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val	
325 330 335	
cgc ctg ctg gaa gat ggc gat ctc gag cca tct gct gga gac atg aga	1056
Arg Leu Leu Glu Asp Gly Asp Leu Glu Pro Ser Ala Gly Asp Met Arg	
340 345 350	

gct gcc aac ctt tgg cca agc ccg ctc atg atc aaa cgc tct aag aag	1104
Ala Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys	
355 360 365	
aac agc ctg gcc ttg tcc ctg acg gcc gac cag atg gtc agt gcc ttg	1152
Asn Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu	
370 375 380	
ttg gat gct gag ccc ccc ata ctc tat tcc gag tat gat cct acc aga	1200
Leu Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg	
385 390 395 400	
ccc ttc agt gaa gct tcg atg atg ggc tta ctg acc aac ctg gca gac	1248
Pro Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp	
405 410 415	
agg gag ctg gtt cac atg atc aac tgg gcg aag agg gtg cca ggc ttt	1296
Arg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe	
420 425 430	
gtg gat ttg acc ctc cat gat cag gtc cac ctt cta gaa tgt gcc tgg	1344
Val Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp	
435 440 445	
cta gag atc ctg atg att ggt ctc gtc tgg cgc tcc atg gag cac cca	1392
Leu Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro	
450 455 460	
ggg aag cta ctg ttt gct cct aac ttg ctc ttg gac agg aac cag gga	1440
Gly Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly	
465 470 475 480	
aaa tgt gta gag ggc atg gtg gag atc ttc gac atg ctg ctg gct aca	1488
Lys Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr	
485 490 495	
tca tct cgg ttc cgc atg atg aat ctg cag gga gag gag ttt gtg tgc	1536
Ser Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys	
500 505 510	
ctc aaa tct att att ttg ctt aat tct gga gtg tac aca ttt ctg tcc	1584
Leu Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser	
515 520 525	
agc acc ctg aag tct ctg gaa gag aag gac cat atc cac cga gtc ctg	1632
Ser Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu	
530 535 540	
gac aag atc aca gac act ttg atc cac ctg atg gcc aag gca ggc ctg	1680
Asp Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu	
545 550 555 560	
acc ctg cag cag cag cac cag cgg ctg gcc cag ctc ctc ctc atc ctc	1728
Thr Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu	
565 570 575	
tcc cac atc agg cac atg agt aac aaa ggc atg gag cat ctg tac agc	1776
Ser His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser	
580 585 590	
atg aag tgc aag aac gtg gtg ccc ctc tat gac ctg ctg ctg gag gcg	1824

Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Ala
 595 600 605

gcg gac gcc cac cgc cta cat gcg ccc act agc cgt gga ggg gca tcc 1872
 Ala Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser
 610 615 620

gtg gag gag acg gac caa agc cac ttg gcc act gcg ggc tct act tca 1920
 Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser
 625 630 635 640

tcg cat tcc ttg caa aag tat tac atc acg ggg gag gca gag ggt ttc 1968
 Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe
 645 650 655

cct gcc aca gct tga 1983
 Pro Ala Thr Ala
 660

<210> 8

<211> 660

<212> PRT

<213> Artificial sequence

<223> Chimeric sequence Homosapiens-Bacteriophage P1 .

<400> 8

Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val
 1 5 10 15

Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg
 20 25 30

Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val
 35 40 45

Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe
 50 55 60

Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala
 65 70 75 80

Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn
 85 90 95

Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala
 100 105 110

Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly
 115 120 125

Glu Arg Ala Lys Gln Ala Leu Ala Phe Glu Arg Thr Asp Phe Asp Gln
 130 135 140

Val Arg Ser Leu Met Glu Asn Ser Asp Arg Cys Gln Asp Ile Arg Asn
 145 150 155 160

Leu Ala Phe Leu Gly Ile Ala Tyr Asn Thr Leu Leu Arg Ile Ala Glu
 165 170 175

Ile Ala Arg Ile Arg Val Lys Asp Ile Ser Arg Thr Asp Gly Gly Arg

180	185	190
Met Leu Ile His Ile Gly Arg Thr Lys Thr Leu Val Ser Thr Ala Gly		
195	200	205
Val Glu Lys Ala Leu Ser Leu Gly Val Thr Lys Leu Val Glu Arg Trp		
210	215	220
Ile Ser Val Ser Gly Val Ala Asp Asp Pro Asn Asn Tyr Leu Phe Cys		
225	230	235
Arg Val Arg Lys Asn Gly Val Ala Ala Pro Ser Ala Thr Ser Gln Leu		
245	250	255
Ser Thr Arg Ala Leu Glu Gly Ile Phe Glu Ala Thr His Arg Leu Ile		
260	265	270
Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly		
275	280	285
His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val		
290	295	300
Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile		
305	310	315
Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val		
325	330	335
Arg Leu Leu Glu Asp Gly Asp Leu Glu Pro Ser Ala Gly Asp Met Arg		
340	345	350
Ala Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys		
355	360	365
Asn Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu		
370	375	380
Leu Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg		
385	390	395
Pro Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp		
405	410	415
Arg Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe		
420	425	430
Val Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp		
435	440	445
Leu Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro		
450	455	460
Gly Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly		
465	470	475
Lys Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr		
485	490	495
Ser Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys		
500	505	510

Leu Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser
 515 520 525
 Ser Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu
 530 535 540
 Asp Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu
 545 550 555 560
 Thr Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu
 565 570 575
 Ser His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser
 580 585 590
 Met Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Ala
 595 600 605
 Ala Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser
 610 615 620
 Val Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser
 625 630 635 640
 Ser His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe
 645 650 655
 Pro Ala Thr Ala
 660

<210> 9
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 9
 tccttcacca agcacatctg 20

<210> 10
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 10
 tgcagccctc acaactgtat 20

<210> 11
 <211> 21
 <212> DNA
 <213> Homo sapiens

<400> 11
 caacctgcac ttgtcactta g 21

<210> 12

<211> 20
<212> DNA
<213> Homo sapiens

<400> 12
atgtttcata gttggatatc

20

<210> 13
<211> 37
<212> DNA
<213> Mus musculus

<400> 13
atacgcggcc gcgaattcca gcaggaatca ggtagct

37

<210> 14
<211> 37
<212> DNA
<213> Mus musculus

<400> 14
atagcgccgg cgctgcagca caggaggggtg ctatgag

37