

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 July 2006 (27.07.2006)

PCT

(10) International Publication Number
WO 2006/078992 A2

(51) International Patent Classification:
A61K 31/496 (2006.01)

(21) International Application Number:
PCT/US2006/002194

(22) International Filing Date: 19 January 2006 (19.01.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/645,496 19 January 2005 (19.01.2005) US

(71) Applicant (for all designated States except US): **NEURO-GEN CORPORATION** [US/US]; 35 Northeast Industrial Road, Branford, CT 06405 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BLUM, Charles, A.** [US/US]; 785 W. Pond Meadow Road, Westbrook, Connecticut 06498 (US).

(74) Agents: **ALEXANDER, John, B.** et al.; Edwards Angell Palmer & Dodge LLP, P.O. Box 55874, Boston, MA 02205 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HETEROARYL SUBSTITUTED PIPERAZINYL-PYRIDINE ANALOGUES



(57) **Abstract:** Substituted biaryl piperazinyl-pyridine analogues are provided, of the Formula: wherein variables are as described herein. Such compounds are ligands that may be used to modulate specific receptor activity in vivo or in vitro, and are particularly useful in the treatment of conditions associated with pathological receptor activation in humans, domesticated companion animals and livestock animals. Pharmaceutical compositions and methods for using such compounds to treat such disorders are provided, as are methods for using such ligands for receptor localization studies.

HETEROARYL SUBSTITUTED PIPERAZINYL-PYRIDINE ANALOGUES

5

FIELD OF THE INVENTION

This invention relates generally to heteroaryl substituted piperazinyl-pyridine analogues that have useful pharmacological properties. The invention further relates to the use of such compounds for treating conditions related to capsaicin receptor activation, for identifying other agents that bind to capsaicin receptor, and as probes for the detection and localization of capsaicin receptors.

BACKGROUND OF THE INVENTION

Pain perception, or nociception, is mediated by the peripheral terminals of a group of specialized sensory neurons, termed "nociceptors." A wide variety of physical and chemical stimuli induce activation of such neurons in mammals, leading to recognition of a potentially harmful stimulus. Inappropriate or excessive activation of nociceptors, however, can result in debilitating acute or chronic pain.

Neuropathic pain involves pain signal transmission in the absence of stimulus, and typically results from damage to the nervous system. In most instances, such pain is thought to occur because of sensitization in the peripheral and central nervous systems following initial damage to the peripheral system (*e.g.*, via direct injury or systemic disease). Neuropathic pain is typically burning, shooting and unrelenting in its intensity and can sometimes be more debilitating than the initial injury or disease process that induced it.

Existing treatments for neuropathic pain are largely ineffective. Opiates, such as morphine, are potent analgesics, but their usefulness is limited because of adverse side effects, such as physical addictiveness and withdrawal properties, as well as respiratory depression, mood changes, and decreased intestinal motility with concomitant constipation, nausea, vomiting, and alterations in the endocrine and autonomic nervous systems. In addition, neuropathic pain is frequently non-responsive or only partially responsive to conventional opioid analgesic regimens. Treatments employing the N-methyl-D-aspartate antagonist ketamine or the $\alpha(2)$ -adrenergic agonist clonidine can reduce acute or chronic pain, and permit a reduction in opioid consumption, but these agents are often poorly tolerated due to side effects.

Topical treatment with capsaicin has been used to treat chronic and acute pain, including neuropathic pain. Capsaicin is a pungent substance derived from the plants of the Solanaceae

family (which includes hot chili peppers) and appears to act selectively on the small diameter afferent nerve fibers (A-delta and C fibers) that are believed to mediate pain. The response to capsaicin is characterized by persistent activation of nociceptors in peripheral tissues, followed by eventual desensitization of peripheral nociceptors to one or more stimuli. From studies in
5 animals, capsaicin appears to trigger C fiber membrane depolarization by opening cation selective channels for calcium and sodium.

Similar responses are also evoked by structural analogues of capsaicin that share a common vanilloid moiety. One such analogue is resiniferatoxin (RTX), a natural product of *Euphorbia* plants. The term vanilloid receptor (VR) was coined to describe the neuronal
10 membrane recognition site for capsaicin and such related irritant compounds. The capsaicin response is competitively inhibited (and thereby antagonized) by another capsaicin analog, capsazepine, and is also inhibited by the non-selective cation channel blocker ruthenium red, which binds to VR with no more than moderate affinity (typically with a K_i value of no lower than 140 μ M).


15 Rat and human vanilloid receptors have been cloned from dorsal root ganglion cells. The first type of vanilloid receptor to be identified is known as vanilloid receptor type 1 (VR1), and the terms "VR1" and "capsaicin receptor" are used interchangeably herein to refer to rat and/or human receptors of this type, as well as mammalian homologues. The role of VR1 in pain sensation has been confirmed using mice lacking this receptor, which exhibit no vanilloid-evoked
20 pain behavior and impaired responses to heat and inflammation. VR1 is a nonselective cation channel with a threshold for opening that is lowered in response to elevated temperatures, low pH, and capsaicin receptor agonists. Opening of the capsaicin receptor channel is generally followed by the release of inflammatory peptides from neurons expressing the receptor and other nearby neurons, increasing the pain response. After initial activation by capsaicin, the capsaicin receptor
25 undergoes a rapid desensitization via phosphorylation by cAMP-dependent protein kinase.

Because of their ability to desensitize nociceptors in peripheral tissues, VR1 agonist vanilloid compounds have been used as topical anesthetics. However, agonist application may itself cause burning pain, which limits this therapeutic use. Recently, it has been reported that VR1 antagonists, including certain nonvanilloid compounds, are also useful for the treatment of
30 pain (*see, e.g.*, PCT International Application Publication Numbers WO 02/08221, WO 03/062209, WO 04/054582, WO 04/055003, WO 04/055004, WO 04/056774, WO 05/007646, WO 05/007648, WO 05/007652, WO 05/009977, WO 05/009980 and WO 05/009982).

Thus, compounds that interact with VR1, but do not elicit the initial painful sensation of VR1 agonist vanilloid compounds, are desirable for the treatment of chronic and acute pain,
35 including neuropathic pain, as well as other conditions that are responsive to capsaicin receptor modulation. The present invention fulfills this need, and provides further related advantages.

SUMMARY OF THE INVENTION

The present invention provides heteroaryl substituted piperazinyl-pyridine analogues of Formula I:



and pharmaceutically acceptable salts of such compounds. Within Formula I:

- 5 Ar₁ is a 5-membered aromatic heterocycle that is substituted with from 0 to 4 substituents independently chosen from R₁;

Ar₂ is phenyl or a 6-membered aromatic heterocycle, each of which is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from R₂;

W is CH or N;

- 10 X, Y and Z are independently CR_x or N, such that at least one of X, Y and Z is N;

R_x is independently chosen at each occurrence from hydrogen, C₁-C₄alkyl, amino, cyano and mono- and di-(C₁-C₄alkyl)amino;

Each R₁ is independently chosen from:

- (a) halogen, cyano and nitro;
- 15 (b) groups of the formula -Q-M-R_y; and
- (c) groups that are taken together with an adjacent R₁ to form a fused 5- to 7-membered carbocyclic or heterocyclic ring that is substituted with from 0 to 4 substituents independently chosen from halogen, cyano, nitro and groups of the formula -Q-M-R_y;

Each Q is independently chosen from C₀-C₄alkylene (*i.e.*, Q is absent or is a single covalent bond or a C₁-C₄alkylene group);

20

M is independently selected at each occurrence from a single covalent bond, O, C(=O), OC(=O), C(=O)O, O-C(=O)O, S(O)_m, N(R_z), C(=O)N(R_z), C(=NH)N(R_z), N(R_z)C(=O), N(R_z)C(=NH), N(R_z)S(O)_m, S(O)_mN(R_z) and N[S(O)_mR_z]S(O)_m; wherein m is independently selected at each occurrence from 0, 1 and 2; and R_z is independently selected at each occurrence from

25 hydrogen, C₁-C₈alkyl and groups that are taken together with R_y to form an optionally substituted 4- to 7-membered heterocycle; and

Each R_y is independently hydrogen, C₁-C₈haloalkyl, optionally substituted C₁-C₈alkyl, optionally substituted (C₃-C₈carbocycle)C₀-C₄alkyl, optionally substituted (4- to 7-membered heterocycle)C₀-C₄alkyl, or taken together with R_z to form an optionally substituted 4- to 7-

30 membered heterocycle, wherein each alkyl, carbocycle and heterocycle is preferably substituted with from 0 to 4 substituents independently selected from hydroxy, halogen, amino, cyano, nitro, oxo, -COOH, aminocarbonyl, C₁-C₆alkyl, C₃-C₇cycloalkyl, C₂-C₆alkyl


ether, C₁-C₆alkanoyl, -SO₂(C₁-C₆alkyl), -SO₂NH₂, C₁-C₈alkoxy, C₁-C₈alkylthio, mono- and di-(C₁-C₆alkyl)aminocarbonyl, mono- and di-(C₁-C₆alkyl)amino and phenyl; such that R_y is not hydrogen if Q is C₀alkyl and M is a single covalent bond;

Each R₂ is:

- 5 (a) independently chosen from (i) hydroxy, amino, cyano, halogen, -COOH, -SO₂NH₂, nitro and aminocarbonyl; and (ii) C₁-C₆alkyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₂-C₆alkyl ether, C₂-C₆alkanoyl, C₁-C₆alkoxycarbonyl, C₂-C₆alkanoyloxy, C₃-C₆alkanone, mono- and di-(C₁-C₆alkyl)aminoC₀-C₆alkyl, mono- and di-(C₃-C₈cycloalkyl)aminoC₀-C₄alkyl, (4- to 7-membered heterocycle)C₀-C₄alkyl, C₁-C₆alkylsulfonyl, mono- and di-(C₁-C₆alkyl)aminosulfonyl, and mono- and di-(C₁-C₆alkyl)aminocarbonyl, each of which is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, cyano, amino, -COOH and oxo; or
- 10 (b) taken together with an adjacent R₂ to form a fused 5- to 13-membered carbocyclic or heterocyclic group that is optionally substituted, and is preferably substituted with from 0 to 3 substituents independently chosen from halogen, oxo and C₁-C₆alkyl;
- 15

R₃ is selected from:

- (i) hydrogen and halogen;
- (ii) C₁-C₆alkyl, (C₃-C₈cycloalkyl)C₀-C₂alkyl, C₁-C₆haloalkyl and phenylC₀-C₂alkyl; and
- 20 (iii) groups of the formula:



wherein:

L is C₀-C₆alkylene or C₁-C₆alkyl that is taken together with R₅, R₆ or R₇ to form a 4- to 7-membered heterocycle;

25 W is O, CO, S, SO or SO₂;

R₅ and R₆ are:

- (a) independently chosen from hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₂-C₆alkanoyl, C₁-C₆alkylsulfonyl, phenylC₀-C₆alkyl, (4- to 7-membered heterocycle)C₀-C₆alkyl and groups that are joined to L to form a 4- to 7-membered heterocycle; or
- 30 (b) joined to form a 4- to 12-membered heterocycle; and
- R₇ is hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₂-C₆alkanoyl, phenylC₀-C₆alkyl, (4- to 7-membered heterocycle)C₀-C₆alkyl or a group that is joined to L to form a 4- to 7-membered heterocycle;

wherein each of (ii) and (iii) is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from:

(1) halogen, hydroxy, amino, cyano, -COOH, -SO₂NH₂, oxo, nitro and aminocarbonyl; and

5 (2) C₁-C₆alkyl, (C₃-C₈cycloalkyl)C₀-C₂alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆alkoxycarbonyl, C₁-C₆haloalkyl, C₁-C₆alkanoyl, C₂-C₆alkanoylamino, mono- and di-(C₁-C₆alkyl)aminoC₀-C₄alkyl, C₁-C₆alkylsulfonyl, mono- and di-(C₁-C₆alkyl)aminosulfonyl, mono- and di-(C₁-C₆alkyl)aminocarbonylC₀-C₄alkyl, phenylC₀-C₄alkyl and (4- to 7-membered heterocycle)C₀-C₄alkyl, each of which is
10 substituted with from 0 to 4 secondary substituents independently chosen from halogen, hydroxy, cyano, oxo, imino, C₁-C₄alkyl, C₁-C₄alkoxy and C₁-C₄haloalkyl; and

R₄ represents from 0 to 2 substituents that are preferably independently chosen from C₁-C₃alkyl, C₁-C₃haloalkyl and oxo.

15 Within certain aspects, compounds of Formula I are VR1 modulators and exhibit a K_i of no greater than 1 micromolar, 500 nanomolar, 100 nanomolar, 50 nanomolar, 10 nanomolar or 1 nanomolar in a capsaicin receptor binding assay and/or have an EC₅₀ or IC₅₀ value of no greater than 1 micromolar, 500 nanomolar, 100 nanomolar, 50 nanomolar, 10 nanomolar or 1 nanomolar in an *in vitro* assay for determination of capsaicin receptor agonist or antagonist activity. In
20 certain embodiments, such VR1 modulators are VR1 antagonists and exhibit no detectable agonist activity in an *in vitro* assay of capsaicin receptor activation (*e.g.*, the assay provided in Example 6, herein) at a concentration equal to the IC₅₀, 10 times the IC₅₀ or 100 times the IC₅₀.

Within certain aspects, compounds provided herein are labeled with a detectable marker (*e.g.*, radiolabeled or fluorescein conjugated).

25 The present invention further provides, within other aspects, pharmaceutical compositions comprising at least one compound of Formula I in combination with a physiologically acceptable carrier or excipient.

Within further aspects, methods are provided for reducing calcium conductance of a cellular capsaicin receptor, comprising contacting a cell (*e.g.*, neuronal, such as cells of the central
30 nervous system and/or peripheral ganglia, urothelial or lung) that expresses a capsaicin receptor with at least one VR1 modulator as described herein. Such contact may occur *in vivo* or *in vitro* and is generally performed using a concentration of VR1 modulator that is sufficient to alter the binding of vanilloid ligand to VR1 *in vitro* (using the assay provided in Example 5) and/or VR1-mediated signal transduction (using an assay provided in Example 6).

35 Methods are further provided for inhibiting binding of vanilloid ligand to a capsaicin receptor. Within certain such aspects, the inhibition takes place *in vitro*. Such methods comprise contacting a capsaicin receptor with at least one VR1 modulator as described herein, under

conditions and in an amount or concentration sufficient to detectably inhibit vanilloid ligand binding to the capsaicin receptor. Within other such aspects, the capsaicin receptor is in a patient. Such methods comprise contacting cells expressing a capsaicin receptor in a patient with at least one VR1 modulator as described herein in an amount or concentration that would be sufficient to
5 detectably inhibit vanilloid ligand binding to cells expressing a cloned capsaicin receptor *in vitro*.

The present invention further provides methods for treating a condition responsive to capsaicin receptor modulation in a patient, comprising administering to the patient a therapeutically effective amount of at least one VR1 modulator as described herein.

Within other aspects, methods are provided for treating pain in a patient, comprising
10 administering to a patient suffering from (or at risk for) pain a therapeutically effective amount of at least one VR1 modulator as described herein.

Methods are further provided for treating itch, urinary incontinence, overactive bladder, cough and/or hiccup in a patient, comprising administering to a patient suffering from (or at risk for) one or more of the foregoing conditions a therapeutically effective amount of at least one VR1
15 modulator as described herein.

The present invention further provides methods for promoting weight loss in an obese patient, comprising administering to an obese patient a therapeutically effective amount of at least one VR1 modulator as described herein.

Methods are further provided for identifying an agent that binds to capsaicin receptor, comprising: (a) contacting capsaicin receptor with a labeled compound as described herein under
20 conditions that permit binding of the compound to capsaicin receptor, thereby generating bound, labeled compound; (b) detecting a signal that corresponds to the amount of bound, labeled compound in the absence of test agent; (c) contacting the bound, labeled compound with a test agent; (d) detecting a signal that corresponds to the amount of bound labeled compound in the presence of test agent; and (e) detecting a decrease in signal detected in step (d), as compared to
25 the signal detected in step (b).

Within further aspects, the present invention provides methods for determining the presence or absence of capsaicin receptor in a sample, comprising: (a) contacting a sample with a compound as described herein under conditions that permit binding of the compound to capsaicin
30 receptor; and (b) detecting a signal indicative of a level of the compound bound to capsaicin receptor.

The present invention also provides packaged pharmaceutical preparations, comprising: (a) a pharmaceutical composition as described herein in a container; and (b) instructions for using the composition to treat one or more conditions responsive to capsaicin receptor modulation, such
35 as pain, itch, urinary incontinence, overactive bladder, cough, hiccup and/or obesity.

In yet another aspect, the present invention provides methods for preparing the compounds disclosed herein, including the intermediates.

These and other aspects of the invention will become apparent upon reference to the following detailed description.


DETAILED DESCRIPTION

As noted above, the present invention provides heteroaryl substituted piperazinyl-pyridine analogues. Such compounds may be used *in vitro* or *in vivo*, to modulate capsaicin receptor activity in a variety of contexts.

TERMINOLOGY

Compounds are generally described herein using standard nomenclature. For compounds having asymmetric centers, it should be understood that (unless otherwise specified) all of the optical isomers and mixtures thereof are encompassed. In addition, compounds with carbon-carbon double bonds may occur in *Z*- and *E*- forms, with all isomeric forms of the compounds being included in the present invention unless otherwise specified. Where a compound exists in various tautomeric forms, a recited compound is not limited to any one specific tautomer, but rather is intended to encompass all tautomeric forms. Certain compounds are described herein using a general formula that includes variables (*e.g.*, R_1 , X , Ar_2). Unless otherwise specified, each variable within such a formula is defined independently of any other variable, and any variable that occurs more than one time in a formula is defined independently at each occurrence.

The term "heteroaryl substituted piperazinyl-pyridine analogue," as used herein, encompasses all compounds of Formula I, as well as compounds of other Formulas provided herein (including any enantiomers, racemates and stereoisomers) and pharmaceutically acceptable salts of such compounds. In other words, compounds in which the core ring



A "pharmaceutically acceptable salt" of a compound is an acid or base salt that is generally considered in the art to be suitable for use in contact with the tissues of human beings or animals without excessive toxicity or carcinogenicity, and preferably without irritation, allergic response, or other problem or complication. Such salts include mineral and organic acid salts of basic residues such as amines, as well as alkali or organic salts of acidic residues such as carboxylic acids. Specific pharmaceutical salts include, but are not limited to, salts of acids such

as hydrochloric, phosphoric, hydrobromic, malic, glycolic, fumaric, sulfuric, sulfamic, sulfanilic, formic, toluenesulfonic, methanesulfonic, benzene sulfonic, ethane disulfonic, 2-hydroxyethylsulfonic, nitric, benzoic, 2-acetoxybenzoic, citric, tartaric, lactic, stearic, salicylic, glutamic, ascorbic, pantoic, succinic, fumaric, maleic, propionic, hydroxymaleic, hydroiodic, phenylacetic, alkanoic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Similarly, pharmaceutically acceptable cations include, but are not limited to sodium, potassium, calcium, aluminum, lithium and ammonium. Those of ordinary skill in the art will recognize further pharmaceutically acceptable salts for the compounds provided herein, including those listed within *Remington: The Science and Practice of Pharmacy*, 21st ed., Lippincott Williams & Wilkins, Philadelphia, PA (2005). In general, a pharmaceutically acceptable acid or base salt can be synthesized from a parent compound that contains a basic or acidic moiety by any conventional chemical method. Briefly, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, the use of nonaqueous media, such as ether, ethyl acetate, ethanol, isopropanol or acetonitrile, is preferred.

It will be apparent that each compound of Formula I may, but need not, be formulated as a hydrate, solvate or non-covalent complex. In addition, the various crystal forms and polymorphs are within the scope of the present invention. Also provided herein are prodrugs of the compounds of Formula I. A "prodrug" is a compound that may not fully satisfy the structural requirements of the compounds provided herein, but is modified *in vivo*, following administration to a patient, to produce a compound of Formula I, or other formula provided herein. For example, a prodrug may be an acylated derivative of a compound as provided herein. Prodrugs include compounds wherein hydroxy, amine or sulfhydryl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxy, amino or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups within the compounds provided herein. Prodrugs of the compounds provided herein may be prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved *in vivo* to yield the parent compounds.

As used herein, the term "alkyl" refers to a straight or branched chain saturated aliphatic hydrocarbon. Alkyl groups include groups having from 1 to 8 carbon atoms ($\text{C}_1\text{-C}_8\text{alkyl}$), from 1 to 6 carbon atoms ($\text{C}_1\text{-C}_6\text{alkyl}$) and from 1 to 4 carbon atoms ($\text{C}_1\text{-C}_4\text{alkyl}$), such as methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *tert*-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl and 3-methylpentyl. " $\text{C}_0\text{-C}_n\text{alkyl}$ " refers to a single covalent bond (C_0) or an alkyl group having from 1 to n carbon atoms; for example " $\text{C}_0\text{-C}_4\text{alkyl}$ " refers to a single covalent bond or a $\text{C}_1\text{-C}_4\text{alkyl}$ group; " $\text{C}_0\text{-C}_8\text{alkyl}$ " refers to a single covalent bond or a $\text{C}_1\text{-C}_8\text{alkyl}$ group. In some instances, a substituent of an alkyl group is specifically indicated. For example, the term

"hydroxyalkyl" refers to an alkyl group substituted with at least one hydroxy substituent (*e.g.*, "C₁-C₄hydroxyalkyl" refers to a C₁-C₄alkyl group that has at least one -OH substituent).

"Alkylene" refers to a divalent alkyl group, as defined above. C₀-C₄alkylene is a single covalent bond or an alkylene group having 1, 2, 3 or 4 carbon atoms; C₁-C₁alkylene is an alkylene group having from 1 to 4 carbon atoms; and C₁-C₆alkylene is an alkylene group having from 1 to 6 carbon atoms.

"Alkenyl" refers to straight or branched chain alkene groups, which comprise at least one unsaturated carbon-carbon double bond. Alkenyl groups include C₂-C₈alkenyl, C₂-C₆alkenyl and C₂-C₄alkenyl groups, which have from 2 to 8, 2 to 6 or 2 to 4 carbon atoms, respectively, such as ethenyl, allyl or isopropenyl. "Alkynyl" refers to straight or branched chain alkyne groups, which have one or more unsaturated carbon-carbon bonds, at least one of which is a triple bond. Alkynyl groups include C₂-C₈alkynyl, C₂-C₆alkynyl and C₂-C₄alkynyl groups, which have from 2 to 8, 2 to 6 or 2 to 4 carbon atoms, respectively.


A "cycloalkyl" is a group that comprises one or more saturated and/or partially saturated rings in which all ring members are carbon, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, adamantyl, decahydro-naphthalenyl, octahydro-indenyl, and partially saturated variants of the foregoing, such as cyclohexenyl. Cycloalkyl groups do not comprise an aromatic ring or a heterocyclic ring. Certain cycloalkyl groups are C₃-C₈cycloalkyl, in which the group contains a single ring having from 3 to 8 ring members, all of which are carbon. A "(C₃-C₈cycloalkyl)C₀-C₄alkyl" is a 3- to 8-membered cycloalkyl group linked via a single covalent bond or a C₁-C₄alkylene group.

By "alkoxy," as used herein, is meant an alkyl group as described above attached via an oxygen bridge. Alkoxy groups include C₁-C₆alkoxy and C₁-C₄alkoxy groups, which have from 1 to 6 or from 1 to 4 carbon atoms, respectively. Methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, *sec*-butoxy, *tert*-butoxy, n-pentoxy, 2-pentoxy, 3-pentoxy, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy are representative alkoxy groups.

Similarly, "alkylthio" refers to an alkyl group as described above attached via a sulfur bridge.

The term "oxo," as used herein refers to a keto group (C=O). An oxo group that is a substituent of a nonaromatic carbon atom results in a conversion of -CH₂- to -C(=O)-.

Similarly, "imino" refers to a group of the formula C=N. The term "iminoalkyl" refers to an alkyl group as described above substituted with an imine (*e.g.*, a group of the formula



The term "alkanoyl" refers to an acyl group (*e.g.*, -(C=O)-alkyl), in which carbon atoms are in a linear or branched alkyl arrangement and where attachment is through the carbon of the

keto group. Alkanoyl groups have the indicated number of carbon atoms, with the carbon of the keto group being included in the numbered carbon atoms. For example a C₂alkanoyl group is an acetyl group having the formula $-(C=O)CH_3$. Alkanoyl groups include, for example, C₂-C₈alkanoyl, C₂-C₆alkanoyl and C₂-C₄alkanoyl groups, which have from 2 to 8, from 2 to 6 or from 2 to 4 carbon atoms, respectively. "C₁alkanoyl" refers to $-(C=O)H$, which (along with C₂-C₈alkanoyl) is encompassed by the term "C₁-C₈alkanoyl."

An "alkanone" is a ketone group in which carbon atoms are in a linear or branched alkyl arrangement. "C₃-C₈alkanone," "C₃-C₆alkanone" and "C₃-C₄alkanone" refer to an alkanone having from 3 to 8, 6 or 4 carbon atoms, respectively. A C₃ alkanone group has the structure $-CH_2-(C=O)-CH_3$.

Similarly, "alkyl ether" refers to a linear or branched ether substituent (*i.e.*, an alkyl group that is substituted with an alkoxy group). Alkyl ether groups include C₂-C₈alkyl ether, C₂-C₆alkyl ether and C₂-C₄alkyl ether groups, which have 2 to 8, 6 or 4 carbon atoms, respectively. A C₂ alkyl ether has the structure $-CH_2-O-CH_3$.

The term "alkoxycarbonyl" refers to an alkoxy group attached through a keto ($-(C=O)-$) bridge (*i.e.*, a group having the general structure $-C(=O)-O-alkyl$). Alkoxycarbonyl groups include C₁-C₈, C₁-C₆ and C₁-C₄alkoxycarbonyl groups, which have from 1 to 8, 6 or 4 carbon atoms, respectively, in the alkyl portion of the group (*i.e.*, the carbon of the keto bridge is not included in the indicated number of carbon atoms). "C₁alkoxycarbonyl" refers to $-C(=O)-O-CH_3$; C₃alkoxycarbonyl indicates $-C(=O)-O-(CH_2)_2CH_3$ or $-C(=O)-O-(CH)(CH_3)_2$.

"Alkanoyloxy," as used herein, refers to an alkanoyl group linked via an oxygen bridge (*i.e.*, a group having the general structure $-O-C(=O)-alkyl$). Alkanoyloxy groups include C₂-C₈, C₂-C₆ and C₂-C₄alkanoyloxy groups, which have from 2 to 8, 6 or 4 carbon atoms, respectively. For example, "C₂alkanoyloxy" refers to $-O-C(=O)-CH_3$.

Similarly, "alkanoylamino," as used herein, refers to an alkanoyl group linked via a nitrogen bridge (*i.e.*, a group having the general structure $-N(R)-C(=O)-alkyl$), in which R is hydrogen or C₁-C₆alkyl. Alkanoylamino groups include C₂-C₈, C₂-C₆ and C₂-C₄alkanoylamino groups, which have from 2 to 8, 6 or 4 carbon atoms within the alkanoyl group, respectively.

"Alkylsulfonyl" refers to groups of the formula $-(SO_2)-alkyl$, in which the sulfur atom is the point of attachment. Alkylsulfonyl groups include C₁-C₆alkylsulfonyl and C₁-C₄alkylsulfonyl groups, which have from 1 to 6 or from 1 to 4 carbon atoms, respectively. Methylsulfonyl is one representative alkylsulfonyl group. "C₁-C₄haloalkylsulfonyl" is an alkylsulfonyl group that has from 1 to 4 carbon atoms and is substituted with at least one halogen (*e.g.*, trifluoromethylsulfonyl).


"Alkylsulfonylamino" refers to groups of the formula $-NH-(SO_2)-alkyl$, in which the nitrogen atom is the point of attachment. Alkylsulfonylamino groups include C₁-

C₆alkylsulfonylamino and C₁-C₄alkylsulfonylamino groups, which have from 1 to 6 or 1 to 4 carbon atoms, respectively. Methylsulfonylamino is a representative alkylsulfonylamino group.

"Aminosulfonyl" refers to groups of the formula $-(SO_2)-NH_2$, in which the sulfur atom is the point of attachment. The term "mono- or di-(C₁-C₈alkyl)aminosulfonyl" refers to groups that satisfy the formula $-(SO_2)-NR_2$, in which the sulfur atom is the point of attachment, and in which one R is C₁-C₈alkyl and the other R is hydrogen or an independently chosen C₁-C₈alkyl.


"Alkylamino" refers to a secondary or tertiary amine that has the general structure $-NH-$ alkyl or $-N(alkyl)(alkyl)$, wherein each alkyl is selected independently from alkyl, cycloalkyl and (cycloalkyl)alkyl groups. Such groups include, for example, mono- and di-(C₁-C₈alkyl)amino groups, in which each C₁-C₈alkyl may be the same or different, as well as mono- and di-(C₁-C₆alkyl)amino groups and mono- and di-(C₁-C₄alkyl)amino groups.

"Alkylaminoalkyl" refers to an alkylamino group linked via an alkylene group (*i.e.*, a group having the general structure $-alkylene-NH-alkyl$ or $-alkylene-N(alkyl)(alkyl)$) in which each alkyl is selected independently from alkyl, cycloalkyl and (cycloalkyl)alkyl groups. Alkylaminoalkyl groups include, for example, mono- and di-(C₁-C₈alkyl)aminoC₁-C₈alkyl, mono- and di-(C₁-C₆alkyl)aminoC₁-C₆alkyl and mono- and di-(C₁-C₆alkyl)aminoC₁-C₄alkyl. "Mono- or di-(C₁-C₆alkyl)aminoC₀-C₆alkyl" refers to a mono- or di-(C₁-C₆alkyl)amino group linked via a single covalent bond or a C₁-C₆alkylene group. The following are representative alkylaminoalkyl groups:



It will be apparent that the definition of "alkyl" as used in the terms "alkylamino" and "alkylaminoalkyl" differs from the definition of "alkyl" used for all other alkyl-containing groups, in the inclusion of cycloalkyl and (cycloalkyl)alkyl groups (*e.g.*, (C₃-C₇cycloalkyl)C₀-C₆alkyl).

25 Similarly, "alkylaminoalkoxy" refers to an alkylamino group linked via an alkoxy group (*i.e.*, a group having the general structure $-O-alkyl-NH-alkyl$ or $-O-alkyl-N(alkyl)(alkyl)$) in which each alkyl is selected independently. Such groups include, for example, mono- and di-(C₁-C₆alkyl)aminoC₁-C₄alkoxy groups, such as



30 The term "aminocarbonyl" refers to an amide group (*i.e.*, $-(C=O)NH_2$). The term "mono- or di-(C₁-C₆alkyl)aminocarbonyl" refers to groups of the formula $-(C=O)-N(R)_2$, in which the carbonyl is the point of attachment, one R is C₁-C₆alkyl and the other R is hydrogen or an independently chosen C₁-C₆alkyl. "Mono- or di-(C₁-C₆alkyl)aminocarbonylC₀-C₄alkyl" refers to

such a group linked via a single covalent bond (*i.e.*, mono- or di-(C₁-C₆alkyl)aminocarbonyl) or a C₁-C₄alkylene group (*i.e.*, -(C₀-C₄alkyl)-(C=O)N(C₁-C₈alkyl)₂).

The term "halogen" refers to fluorine, chlorine, bromine or iodine.

A "haloalkyl" is an alkyl group that is substituted with 1 or more independently chosen
5 halogens (*e.g.*, "C₁-C₈haloalkyl" groups have from 1 to 8 carbon atoms; "C₁-C₆haloalkyl" groups have from 1 to 6 carbon atoms). Examples of haloalkyl groups include, but are not limited to, mono-, di- or tri-fluoromethyl; mono-, di- or tri-chloromethyl; mono-, di-, tri-, tetra- or penta-fluoroethyl; mono-, di-, tri-, tetra- or penta-chloroethyl; and 1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl. Typical haloalkyl groups are trifluoromethyl and difluoromethyl. The term "haloalkoxy"
10 refers to a haloalkyl group as defined above attached via an oxygen bridge. "C₁-C₈haloalkoxy" groups have 1 to 8 carbon atoms.

A dash ("-") that is not between two letters or symbols is used to indicate a point of attachment for a substituent. For example, -CONH₂ is attached through the carbon atom.

A "carbocycle" or "carbocyclic group" comprises at least one ring formed entirely by
15 carbon-carbon bonds (referred to herein as a carbocyclic ring), and does not contain a heterocycle. Unless otherwise specified, each ring within a carbocycle may be independently saturated, partially saturated or aromatic, and is optionally substituted as indicated. A carbocycle generally has from 1 to 3 fused, pendant or spiro rings; carbocycles within certain embodiments have one ring or two fused rings. Typically, each ring contains from 3 to 8 ring members (*i.e.*, C₃-C₈); C₅-
20 C₇ rings are recited in certain embodiments. Carbocycles comprising fused, pendant or spiro rings typically contain from 9 to 14 ring members. Certain carbocycles are C₄-C₁₀ (*i.e.*, contain from 4 to 10 ring members, and one or two rings). Certain representative carbocycles are cycloalkyl as described above. Other carbocycles are aryl (*i.e.*, contain at least one aromatic carbocyclic ring, with or without one or more additional aromatic and/or cycloalkyl rings). Such aryl carbocycles
25 include, for example, phenyl, naphthyl (*e.g.*, 1-naphthyl and 2-naphthyl), fluorenyl, indanyl and 1,2,3,4-tetrahydro-naphthyl.

Certain carbocycles recited herein are C₆-C₁₀arylC₀-C₈alkyl groups (*i.e.*, groups in which a 6- to 10-membered carbocyclic group comprising at least one aromatic ring is linked via a single covalent bond or a C₁-C₈alkylene group). Such groups include, for example, phenyl and indanyl,
30 as well as groups in which either of the foregoing is linked via C₁-C₈alkylene, preferably via C₁-C₄alkylene. Phenyl groups linked via a single covalent bond or C₁-C₆alkylene group are designated phenylC₀-C₆alkyl (*e.g.*, benzyl, 1-phenyl-ethyl, 1-phenyl-propyl and 2-phenyl-ethyl).

A "heterocycle" or "heterocyclic group" has from 1 to 3 fused, pendant or spiro rings, at least one of which is a heterocyclic ring (*i.e.*, one or more ring atoms is a heteroatom
35 independently chosen from O, S and N, with the remaining ring atoms being carbon). Additional rings, if present, may be heterocyclic or carbocyclic. Typically, a heterocyclic ring comprises 1, 2, 3 or 4 heteroatoms; within certain embodiments each heterocyclic ring has 1 or 2 heteroatoms per

ring. Each heterocyclic ring generally contains from 3 to 8 ring members (rings having from 4 or 5 to 7 ring members are recited in certain embodiments) and heterocycles comprising fused, pendant or spiro rings typically contain from 9 to 14 ring members. Certain heterocycles comprise a sulfur atom as a ring member; in certain embodiments, the sulfur atom is oxidized to SO or SO₂.

5 Heterocycles may be optionally substituted with a variety of substituents, as indicated. Unless otherwise specified, a heterocycle may be a heterocycloalkyl group (*i.e.*, each ring is saturated or partially saturated) or a heteroaryl group (*i.e.*, at least one ring within the group is aromatic), such as a 5- to 10-membered heteroaryl (which may be monocyclic or bicyclic) or a 6-membered heteroaryl (*e.g.*, pyridyl or pyrimidyl).

10 Heterocyclic groups include, for example, azepanyl, azocinyl, benzimidazolyl, benzimidazoliny, benzisothiazolyl, benzisoxazolyl, benzofuranyl, benzothiofuranyl, benzoxazolyl, benzothiazolyl, benztetrazolyl, chromanyl, chromenyl, cinnoliny, decahydroquinoliny, dihydrofuro[2,3-*b*]tetrahydrofuranyl, dihydroisoquinoliny, dihydrotetrahydrofuranyl, 1,4-dioxo-8-aza-spiro[4.5]decyl, dithiazinyl, furanyl, furazanyl, imidazoliny, imidazolidiny, imidazolyl, 15 indazolyl, indolenyl, indoliny, indoliziny, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindoliny, isoindolyl, isothiazolyl, isoxazolyl, isoquinoliny, morpholiny, naphthyridiny, octahydroisoquinoliny, oxadiazolyl, oxazolidiny, oxazolyl, phthalazinyl, piperazinyl, piperidiny, piperidiny, piperidonyl, pteridiny, puriny, pyranyl, pyraziny, pyrazolidiny, pyrazoliny, pyrazolyl, pyridazinyl, pyridoimidazolyl, pyridooxazolyl, pyridothiazolyl, pyridyl, pyrimidyl, 20 pyrrolidiny, pyrrolidonyl, pyrroliny, pyrrolyl, quinazoliny, quinoliny, quinoxaliny, quinuclidiny, tetrahydroisoquinoliny, tetrahydroquinoliny, tetrazolyl, thiadiaziny, thiadiazolyl, thiazolyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thienyl, thiophenyl, thiomorpholiny and variants thereof in which the sulfur atom is oxidized, triazinyl, and any of the foregoing that are substituted with from 1 to 4 substituents as described above.

25 Certain heterocyclic groups are 4- to 12-membered, 5- to 10-membered, 3- to 7-membered, 4- to 7-membered or 5- to 7-membered groups that contain 1 heterocyclic ring or 2 fused or spiro rings, optionally substituted. 4- to 10-membered heterocycloalkyl groups include, for example, piperidiny, piperazinyl, pyrrolidiny, azepanyl, 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl, morpholino, thiomorpholino and 1,1-dioxo-thiomorpholin-4-yl. Such groups may be substituted 30 as indicated. Representative aromatic heterocycles are azocinyl, pyridyl, pyrimidyl, imidazolyl, tetrazolyl and 3,4-dihydro-1H-isoquinolin-2-yl.

A "heterocycleC₀-C₆alkyl" is a heterocyclic group linked via a single covalent bond or C₁-C₈alkyl group. For example, a (4- to 7-membered heterocycle)C₀-C₆alkyl is a heterocyclic group having from 4 to 7 ring members linked via a single covalent bond or an alkyl group having from 1 35 to 6 carbon atoms. A "(6-membered heteroaryl)C₀-C₆alkyl" refers to a heteroaryl group linked via a single covalent bond or C₁-C₆alkyl group.

A "substituent," as used herein, refers to a molecular moiety that is covalently bonded to an atom within a molecule of interest. For example, a ring substituent may be a moiety such as a halogen, alkyl group, haloalkyl group or other group that is covalently bonded to an atom (preferably a carbon or nitrogen atom) that is a ring member. Substituents of aromatic groups are generally covalently bonded to a ring carbon atom. The term "substitution" refers to replacing a hydrogen atom in a molecular structure with a substituent, such that the valence on the designated atom is not exceeded, and such that a chemically stable compound (*i.e.*, a compound that can be isolated, characterized, and tested for biological activity) results from the substitution.

Groups that are "optionally substituted" are unsubstituted or are substituted by other than hydrogen at one or more available positions, typically 1, 2, 3, 4 or 5 positions, by one or more suitable groups (which may be the same or different). Optional substitution is also indicated by the phrase "substituted with from 0 to X substituents," where X is the maximum number of possible substituents. Certain optionally substituted groups are substituted with from 0 to 2, 3 or 4 independently selected substituents (*i.e.*, are unsubstituted or substituted with up to the recited maximum number of substituents). Other optionally substituted groups are substituted with at least one substituent (*e.g.*, substituted with from 1 to 2, 3 or 4 independently selected substituents).

The terms "VR1" and "capsaicin receptor" are used interchangeably herein to refer to a type 1 vanilloid receptor. Unless otherwise specified, these terms encompass both rat and human VR1 receptors (*e.g.*, GenBank Accession Numbers AF327067, AJ277028 and NM_018727; sequences of certain human VR1 cDNAs and the encoded amino acid sequences are provided in U.S. Patent No. 6,482,611), as well as homologues thereof found in other species.

A "VR1 modulator," also referred to herein as a "modulator," is a compound that modulates VR1 activation and/or VR1-mediated signal transduction. VR1 modulators specifically provided herein are compounds of Formula I and pharmaceutically acceptable salts thereof. Certain preferred VR1 modulators are not vanilloids. A VR1 modulator may be a VR1 agonist or antagonist. Certain modulators bind to VR1 with a K_i that is less than 1 micromolar, preferably less than 500 nanomolar, 100 nanomolar, 10 nanomolar or 1 nanomolar. A representative assay for determining K_i at VR1 is provided in Example 5, herein.

A modulator is considered an "antagonist" if it detectably inhibits vanilloid ligand binding to VR1 and/or VR1-mediated signal transduction (using, for example, the representative assay provided in Example 6); in general, such an antagonist inhibits VR1 activation with a IC_{50} value of less than 1 micromolar, preferably less than 500 nanomolar, and more preferably less than 100 nanomolar, 10 nanomolar or 1 nanomolar within the assay provided in Example 6. VR1 antagonists include neutral antagonists and inverse agonists.

An "inverse agonist" of VR1 is a compound that reduces the activity of VR1 below its basal activity level in the absence of added vanilloid ligand. Inverse agonists of VR1 may also inhibit the activity of vanilloid ligand at VR1 and/or binding of vanilloid ligand to VR1. The basal

activity of VR1, as well as the reduction in VR1 activity due to the presence of VR1 antagonist, may be determined from a calcium mobilization assay, such as the assay of Example 6.

5 A "neutral antagonist" of VR1 is a compound that inhibits the activity of vanilloid ligand at VR1, but does not significantly change the basal activity of the receptor (*i.e.*, within a calcium mobilization assay as described in Example 6 performed in the absence of vanilloid ligand, VR1 activity is reduced by no more than 10%, preferably by no more than 5%, and more preferably by no more than 2%; most preferably, there is no detectable reduction in activity). Neutral antagonists of VR1 may inhibit the binding of vanilloid ligand to VR1.

10 As used herein a "capsaicin receptor agonist" or "VR1 agonist" is a compound that elevates the activity of the receptor above the basal activity level of the receptor (*i.e.*, enhances VR1 activation and/or VR1-mediated signal transduction). Capsaicin receptor agonist activity may be identified using the representative assay provided in Example 6. In general, such an agonist has an EC₅₀ value of less than 1 micromolar, preferably less than 500 nanomolar, and more preferably less than 100 nanomolar or 10 nanomolar within the assay provided in Example 6.

15 A "vanilloid" any compound that comprises a phenyl ring with two oxygen atoms bound to adjacent ring carbon atoms (one of which carbon atom is located *para* to the point of attachment of a third moiety that is bound to the phenyl ring). Capsaicin is a representative vanilloid. A "vanilloid ligand" is a vanilloid that binds to VR1 with a K_i (determined as described herein) that is no greater than 10 μ M. Vanilloid ligand agonists include capsaicin, olvanil, N-arachidonoyl-dopamine and resiniferatoxin (RTX). Vanilloid ligand antagonists include capsazepine and iodo-resiniferatoxin.

20 A "therapeutically effective amount" (or dose) is an amount that, upon administration to a patient, results in a discernible patient benefit (*e.g.*, provides detectable relief from at least one condition being treated). Such relief may be detected using any appropriate criteria, including alleviation of one or more symptoms such as pain. A therapeutically effective amount or dose generally results in a concentration of compound in a body fluid (such as blood, plasma, serum, CSF, synovial fluid, lymph, cellular interstitial fluid, tears or urine) that is sufficient to alter the binding of vanilloid ligand to VR1 *in vitro* (using the assay provided in Example 5) and/or VR1-mediated signal transduction (using an assay provided in Example 6). It will be apparent that the discernible patient benefit may be apparent after administration of a single dose, or may become apparent following repeated administration of the therapeutically effective dose according to a predetermined regimen, depending upon the indication for which the compound is administered.

30 By "statistically significant," as used herein, is meant results varying from control at the $p < 0.1$ level of significance as measured using a standard parametric assay of statistical significance such as a student's T test.


A "patient" is any individual treated with a compound provided herein. Patients include humans, as well as other animals such as companion animals (*e.g.*, dogs and cats) and livestock.

Patients may be experiencing one or more symptoms of a condition responsive to capsaicin receptor modulation (*e.g.*, pain, exposure to vanilloid ligand, itch, urinary incontinence, overactive bladder, respiratory disorders, cough and/or hiccup), or may be free of such symptom(s) (*i.e.*, treatment may be prophylactic in a patient considered at risk for the development of such symptoms).


HETEROARYL SUBSTITUTED PIPERAZINYL-PYRIDINE ANALOGUES

Within certain aspects, as noted above, the present invention provides heteroaryl substituted piperazinyl-pyridine analogues that may be used in a variety of contexts, including in the treatment of pain (*e.g.*, neuropathic or peripheral nerve-mediated pain); exposure to capsaicin; exposure to acid, heat, light, tear gas, air pollutants (such as, for example, tobacco smoke), infectious agents (including viruses, bacteria and yeast), pepper spray or related agents; respiratory conditions such as asthma or chronic obstructive pulmonary disease; itch; urinary incontinence or overactive bladder; cough or hiccup; and/or obesity. Such compounds may also be used within *in vitro* assays (*e.g.*, assays for receptor activity), as probes for detection and localization of VR1 and as standards in ligand binding and VR1-mediated signal transduction assays.

Within certain embodiments, compounds of Formula I herein further satisfy Formula II:



or a pharmaceutically acceptable salt thereof. Within Formula II:



represents a 5-membered heteroaryl with a nitrogen atom located adjacent to the point of attachment;

D, E and G are independently O, S, N, NR_{1a} or CR_{1a};

Each R_{1a} is independently chosen from:


(a) hydrogen, halogen, cyano and nitro;

(b) groups of the formula -Q-M-R_y; and


(c) groups that are taken together with an adjacent R_{1a} to form a fused 5- to 7-membered carbocyclic or heterocyclic ring that is optionally substituted with from 1 to 4 substituents independently chosen from halogen, cyano, nitro and groups of the formula -Q-M-R_y;

and the remaining variables are as described above for Formula I.


In certain embodiments, compounds provided herein satisfy one or more of Formulas IIa - IIc, in which variables are as indicated for Formula II, except as defined below:




Formula IIa




Formula IIb




Formula IIc



Formula IIId



Formula IIe




Formula II f

Within Formulas IIc and IId, D is O or S; and in Formulas IIe and IIIf, E is N, O or S.











Within certain embodiments of the above Formulas, variables are as follows:

 Ar_l , R_l and R_{lg}

Certain Ar_1 groups satisfy the formula:



5 D-N . Representative such groups include, for example:



$\text{O}-\text{N}$, and $\text{S}-\text{N}$, in which R_{1a} is as described above. In certain embodiments, R_{1a} in the above groups is not hydrogen.

Certain substituents satisfy the formula Q-M-R_y. It will be apparent that if Q is C₀ and M is a single covalent bond, then R_y is directly linked (via a single covalent bond) to the Ar₁ core ring.


Within certain compounds, R_{1a} is hydrogen, halogen, amino, cyano, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆alkylsulfonyl, or mono- or di-(C₁-C₆alkyl)aminosulfonyl.

Representative R_{1a} groups include, for example, hydrogen, halogen, nitro, cyano, methyl, and trifluoromethyl.


Ar_2 , R_2 and R_{2a}

In certain compounds of the above Formulas, Ar_2 is optionally substituted phenyl, pyridyl
 5 (i.e., 2-pyridyl, 3-pyridyl or 4-pyridyl) or pyrimidyl. In other compounds, Ar_2 is a 9- to 12-
 membered bicyclic aryl or heteroaryl group that is optionally substituted as described above.
 Preferably, Ar_2 is substituted with from 0 to 3 or from 1 to 3 substituents independently chosen
 from R_2 as described above. In certain compounds, Ar_2 has at least one substituent (R_2) and each
 R_2 is independently chosen from amino, cyano, halogen, $-SO_2NH_2$, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 -
 10 C_4 hydroxyalkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 haloalkoxy, mono- and di- $(C_1$ -
 C_4 alkyl)amino, C_0 - C_4 alkyl, C_2 - C_4 alkanoyl, C_1 - C_4 alkoxycarbonyl, C_1 - C_4 alkylsulfonyl, C_1 -
 C_4 haloalkylsulfonyl and mono- and di- $(C_1$ - C_4 alkyl)aminosulfonyl. More preferably, the
 substituents of Ar_2 are independently chosen from amino, cyano, halogen, C_1 - C_4 alkyl, C_1 -
 C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkylthio and mono- and di- $(C_1$ -
 15 C_4 alkyl)amino, C_0 - C_4 alkyl. In certain such compounds, Ar_2 is substituted *meta* and/or *para* to the
 point of attachment, wherein the point of attachment refers to the attachment to the core ring. In
 other words, if Ar_2 is phenyl, the phenyl is mono-substituted at the 3-position, mono-substituted at
 the 4-position, or di-substituted and the 3- and 4-positions. Preferred Ar_2 groups include phenyl,
 pyridyl and pyrimidyl, each of which is substituted with from 0 to 3 or from 1 to 3 substituents as
 20 described herein.


Within certain Ar_2 groups, one R_2 is taken together with an adjacent R_2 to form a fused
 carbocycle or heterocycle. Representative such groups include, for example, the following
 bicyclic groups, optionally substituted as described herein:



the fused ring contains one or more additional double bonds, such as:




The variable R_{2a} , when present, is generally as described above; in certain embodiments, each R_{2a} is independently chosen from amino, cyano, halogen, $-\text{SO}_2\text{NH}_2$, $-\text{COOH}$, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ alkylthio, $\text{C}_1\text{-C}_4$ haloalkoxy, mono- and di-
 5 $(\text{C}_1\text{-C}_4\text{alkyl})\text{aminoC}_0\text{-C}_4\text{alkyl}$, $\text{C}_2\text{-C}_4$ alkanoyl, $\text{C}_1\text{-C}_4$ alkoxycarbonyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_1\text{-C}_4$ haloalkylsulfonyl and mono- and di- $(\text{C}_1\text{-C}_4\text{alkyl})\text{sulfonamido}$. In further such compounds, A is CH or CR_{2a} , and each R_{2a} is independently chosen from cyano, halogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ alkoxy and mono- and di- $(\text{C}_1\text{-C}_4\text{alkyl})\text{aminoC}_0\text{-C}_2\text{alkyl}$. In other such
 10 compounds, at least one of A, B, E and T is N. Certain Ar_2 groups have the formula:




R_3

In the definition of R_3 , the variable "L" is defined as $\text{C}_0\text{-C}_6$ alkylene or $\text{C}_1\text{-C}_6$ alkylene that is taken together with R_5 , R_6 or R_7 to form a 4- to 7-membered heterocycle. In any heterocycle so
 15 formed, at least one carbon atom present in L is also a ring atom, and is covalently bonded to a component atom of R_5 , R_6 or R_7 . The resulting heterocycle may be a heterocycloalkyl group (e.g., tetrahydrofuranyl, morpholinyl, piperidinyl or piperazinyl) or a heteroaryl group, such as pyridyl, pyrimidyl or tetrahydrofuranyl. R_3 groups comprising such a heterocycle include, for example:



20 R_3 , in certain embodiments of the above Formulas, is a group of the formula:



wherein:

L is $\text{C}_0\text{-C}_3$ alkyl; and

R_5 and R_6 are:


- 25 (a) independently chosen from hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $(\text{C}_3\text{-C}_8\text{cycloalkyl})\text{C}_0\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_4$ alkanoyl and groups that are joined to L to form a 4- to 7-membered heterocycle; or
 (b) joined to form a 4- to 12-membered heterocycloalkyl;


each of which alkyl, alkenyl, (cycloalkyl)alkyl, alkanoyl and heterocycloalkyl is substituted with from 0 to 4 substituents independently chosen from (i) halogen, hydroxy, amino, aminocarbonyl, oxo, -COOH and -SO₂NH₂; and (ii) C₁-C₄alkyl, C₅-C₇cycloalkyl, C₁-C₄alkoxy, C₂-C₄alkanoyl, C₁-C₄haloalkyl, mono- and di-(C₁-C₄alkyl)aminoC₀-C₂alkyl, mono- and di-(C₁-C₄alkyl)aminocarbonylC₀-C₂alkyl, phenylC₀-C₄alkyl and (4- to 7-membered heterocycle)C₀-C₂alkyl, each of which is substituted with from 0 to 4 secondary substituents independently chosen from halogen, hydroxy, cyano, C₁-C₄alkyl, C₁-C₄alkoxy and C₁-C₄haloalkyl;

Such R₃ groups include, for example, mono- and di-(C₁-C₄alkyl)amino groups that are substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, amino, oxo, aminocarbonyl, -COOH, -SO₂NH₂, C₁-C₄alkyl, C₂-C₄alkenyl, C₅-C₇cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₂-C₄alkyl ether, C₂-C₄alkanoyl, C₁-C₄alkylsulfonyl, C₂-C₄alkanoylamino and mono- and di-(C₁-C₄alkyl)amino.


Other R₃ groups include phenyl and 4- to 7-membered heterocycles, each of which is substituted with from 0 to 4 substituents independently chosen from (a) halogen, hydroxy, amino, oxo, aminocarbonyl, -SO₂NH₂ and -COOH; and (b) C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, (C₅-C₇cycloalkyl)C₀-C₂alkyl, C₁-C₄alkoxy, C₂-C₄alkyl ether, C₂-C₄alkanoyl, C₁-C₄alkylsulfonyl, C₂-C₄alkanoylamino, mono- and di-(C₁-C₄alkyl)amino, mono- and di-(C₁-C₄alkyl)aminocarbonyl, mono- or di-(C₁-C₆alkyl)aminosulfonyl, phenylC₀-C₄alkyl and (4- to 7-membered heterocycle)C₀-C₄alkyl, each of which is substituted with from 0 to 4 secondary substituents independently chosen from halogen, hydroxy, cyano, -COOH, C₁-C₄alkyl and C₁-C₄haloalkyl. Certain such R₃ groups include azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, tetrahydropyridyl and azepanyl, each of which is substituted with from 0 to 4 substituents independently chosen from: (a) halogen, hydroxy, amino, oxo, aminocarbonyl, -SO₂NH₂ and -COOH; and (b) C₁-C₄alkyl, C₂-C₄alkenyl, C₅-C₇cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₂-C₄alkyl ether, C₂-C₄alkanoyl, C₁-C₄alkylsulfonyl, C₂-C₄alkanoylamino and mono- and di-(C₁-C₄alkyl)amino, each of which is substituted with from 0 to 4 secondary substituents independently chosen from hydroxy and halogen.

Representative examples of such R₃ groups include the heterocycles:





Other such R₃ groups are phenyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, imidazolyl, thienyl, oxazolyl or tetrahydrofuranyl, each of which is substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, amino, aminocarbonyl, -SO₂NH₂, -COOH, C₁-C₄alkyl, C₅-C₇cycloalkyl, C₂-C₄alkyl ether, C₁-C₄alkoxy, C₂-C₄alkanoyl, C₁-C₄haloalkyl and mono- and di-(C₁-C₄alkyl)amino. In certain embodiments, R₃ is not -NH₂. In other words, if L is a single covalent bond then at least one of R₅ and R₆ is not hydrogen.


10 In further embodiments, R₃ is . In certain such compounds, L is C₀-C₃alkyl; and R₇ is C₁-C₆alkyl, C₂-C₆alkenyl, (C₅-C₇cycloalkyl)C₀-C₄alkyl, C₂-C₄alkanoyl, phenylC₀-C₆alkyl or (6-membered heteroaryl)C₀-C₄alkyl, each of which is substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, amino, aminocarbonyl, -SO₂NH₂, -COOH, C₁-C₄alkyl, C₅-C₇cycloalkyl, C₂-C₄alkyl ether, C₁-C₄alkoxy, C₂-C₄alkanoyl, C₁-C₄haloalkyl and mono- and di-(C₁-C₄alkyl)amino. Such R₃ groups include, for example, benzyloxy and C₁-C₆alkoxy, each of which is optionally substituted with halogen, methyl, methoxy or trifluoromethyl.

Within still further embodiments, R₃ is a hydrogen, C₁-C₆alkyl or a halogen.

20 Within further compounds of the above Formulas, R₃ is C₁-C₄alkyl, C₃-C₇cycloalkyl or C₁-C₄haloalkyl, each of which is substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, cyano, oxo, aminocarbonyl, -SO₂NH₂, -COOH, C₃-C₇cycloalkyl, phenyl and 4- to 7-membered heterocycle.

R₄


25 R₄, in certain compounds provided herein, represents zero substituents or one methyl, ethyl or oxo group, preferably located at the 2-position of the piperazine. The carbon to which a methyl or ethyl group is attached is chiral in certain embodiments. For example, the group:


 is , , or in certain compounds. In other compounds, R₄ represents a single oxo substituent.

X, Y and Z

X, Y and Z, as noted above, are independently CR_x or N, such that at least one of X, Y and Z is N. In certain embodiments, each R_x is independently chosen from hydrogen and methyl, or each R_x is hydrogen. In certain representative compounds, Z is N (*e.g.*, X and Y are CH). In other compounds provided herein, X is N (*e.g.*, Y and Z are CH). In further compounds Z and X are N, X and Y are N or Z and Y are N. In further compounds, X, Y and Z are each N.

Certain compounds provided herein satisfy Formula III:



Within Formula III, variables are as described above. In certain embodiments:

D, E and G are independently O, S, N, NR_{1a} or CR_{1a};

Each R_{1a} is independently chosen from:

(a) hydrogen, halogen, cyano and nitro; and

(b) groups of the formula -Q-M-R_y;

Ar₂ is phenyl, pyridyl or pyrimidyl, each of which is substituted with from 0 to 3 substituents independently chosen from amino, cyano, halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, and mono- and di-(C₁-C₄alkyl)aminoC₀-C₄alkyl; and

R₄ represents 0 substituents or one methyl substituent.

Representative compounds provided herein include, but are not limited to, those specifically described in Examples 1-3. It will be apparent that the specific compounds recited herein are representative only, and are not intended to limit the scope of the present invention. Further, as noted above, all compounds provided herein may be present as a free acid or base, or as a pharmaceutically acceptable salt. In addition, other forms such as hydrates and prodrugs of such compounds are specifically contemplated by the present invention.

Within certain aspects, heteroaryl substituted piperazinyl-pyridine analogues provided herein detectably alter (modulate) VR1 activity, as determined using an *in vitro* VR1 functional assay such as a calcium mobilization assay. As an initial screen for such activity, a VR1 ligand binding assay may be used. References herein to a "VR1 ligand binding assay" are intended to refer to a standard *in vitro* receptor binding assay such as that provided in Example 5, and a

"calcium mobilization assay" (also referred to herein as a "signal transduction assay") may be performed as described in Example 6. Briefly, to assess binding to VR1, a competition assay may be performed in which a VR1 preparation is incubated with labeled (*e.g.*, ^{125}I or ^3H) compound that binds to VR1 (*e.g.*, a capsaicin receptor agonist such as RTX) and unlabeled test compound.

5 Within the assays provided herein, the VR1 used is preferably mammalian VR1, more preferably human or rat VR1. The receptor may be recombinantly expressed or naturally expressed. The VR1 preparation may be, for example, a membrane preparation from HEK293 or CHO cells that recombinantly express human VR1. Incubation with a compound that detectably modulates vanilloid ligand binding to VR1 results in a decrease or increase in the amount of label bound to
10 the VR1 preparation, relative to the amount of label bound in the absence of the compound. This decrease or increase may be used to determine the K_i at VR1 as described herein. In general, compounds that decrease the amount of label bound to the VR1 preparation within such an assay are preferred.

Certain VR1 modulators provided herein detectably modulate VR1 activity at nanomolar
15 (*i.e.*, submicromolar) concentrations, at subnanomolar concentrations, or at concentrations below 100 picomolar, 20 picomolar, 10 picomolar or 5 picomolar.

As noted above, compounds that are VR1 antagonists are preferred within certain embodiments. IC_{50} values for such compounds may be determined using a standard *in vitro* VR1-mediated calcium mobilization assay, as provided in Example 6. Briefly, cells expressing
20 capsaicin receptor are contacted with a compound of interest and with an indicator of intracellular calcium concentration (*e.g.*, a membrane permeable calcium sensitivity dye such as Fluo-3 or Fura-2 (Molecular Probes, Eugene, OR), each of which produce a fluorescent signal when bound to Ca^{++}). Such contact is preferably carried out by one or more incubations of the cells in buffer or culture medium comprising either or both of the compound and the indicator in solution. Contact
25 is maintained for an amount of time sufficient to allow the dye to enter the cells (*e.g.*, 1-2 hours). Cells are washed or filtered to remove excess dye and are then contacted with a vanilloid receptor agonist (*e.g.*, capsaicin, RTX or olvanil), typically at a concentration equal to the EC_{50} concentration, and a fluorescence response is measured. When agonist-contacted cells are contacted with a compound that is a VR1 antagonist the fluorescence response is generally reduced
30 by at least 20%, preferably at least 50% and more preferably at least 80%, as compared to cells that are contacted with the agonist in the absence of test compound. The IC_{50} for VR1 antagonists provided herein is preferably less than 1 micromolar, less than 100 nM, less than 10 nM or less than 1 nM. In certain embodiments, VR1 antagonists provided herein exhibit no detectable agonist activity an *in vitro* assay of capsaicin receptor agonism at a concentration of compound equal to
35 the IC_{50} . Certain such antagonists exhibit no detectable agonist activity an *in vitro* assay of capsaicin receptor agonism at a concentration of compound that is 100-fold higher than the IC_{50} .

In other embodiments, compounds that are capsaicin receptor agonists are preferred. Capsaicin receptor agonist activity may generally be determined as described in Example 6. When cells are contacted with 1 micromolar of a compound that is a VR1 agonist, the fluorescence response is generally increased by an amount that is at least 30% of the increase observed when
5 cells are contacted with 100 nM capsaicin. The EC₅₀ for VR1 agonists provided herein is preferably less than 1 micromolar, less than 100 nM or less than 10 nM.

VR1 modulating activity may also, or alternatively, be assessed using a cultured dorsal root ganglion assay as provided in Example 7 and/or an *in vivo* pain relief assay as provided in Example 8. VR1 modulators provided herein preferably have a statistically significant specific
10 effect on VR1 activity within one or more functional assays provided herein.

Within certain embodiments, VR1 modulators provided herein do not substantially modulate ligand binding to other cell surface receptors, such as EGF receptor tyrosine kinase or the nicotinic acetylcholine receptor. In other words, such modulators do not substantially inhibit activity of a cell surface receptor such as the human epidermal growth factor (EGF) receptor
15 tyrosine kinase or the nicotinic acetylcholine receptor (*e.g.*, the IC₅₀ or IC₄₀ at such a receptor is preferably greater than 1 micromolar, and most preferably greater than 10 micromolar). Preferably, a modulator does not detectably inhibit EGF receptor activity or nicotinic acetylcholine receptor activity at a concentration of 0.5 micromolar, 1 micromolar or more preferably 10 micromolar. Assays for determining cell surface receptor activity are commercially available, and
20 include the tyrosine kinase assay kits available from Panvera (Madison, WI).

In certain embodiments, preferred VR1 modulators are non-sedating. In other words, a dose of VR1 modulator that is twice the minimum dose sufficient to provide analgesia in an animal model for determining pain relief (such as a model provided in Example 8, herein) causes only transient (*i.e.*, lasting for no more than ½ the time that pain relief lasts) or preferably no
25 statistically significant sedation in an animal model assay of sedation (using the method described by Fitzgerald et al. (1988) *Toxicology* 49(2-3):433-9). Preferably, a dose that is five times the minimum dose sufficient to provide analgesia does not produce statistically significant sedation. More preferably, a VR1 modulator provided herein does not produce sedation at intravenous doses of less than 25 mg/kg (preferably less than 10 mg/kg) or at oral doses of less than 140
30 mg/kg (preferably less than 50 mg/kg, more preferably less than 30 mg/kg).

If desired, compounds provided herein may be evaluated for certain pharmacological properties including, but not limited to, oral bioavailability (preferred compounds are orally bioavailable to an extent allowing for therapeutically effective concentrations of the compound to be achieved at oral doses of less than 140 mg/kg, preferably less than 50 mg/kg, more preferably
35 less than 30 mg/kg, even more preferably less than 10 mg/kg, still more preferably less than 1 mg/kg and most preferably less than 0.1 mg/kg), toxicity (a preferred compound is nontoxic when a therapeutically effective amount is administered to a subject), side effects (a preferred compound

produces side effects comparable to placebo when a therapeutically effective amount of the compound is administered to a subject), serum protein binding and *in vitro* and *in vivo* half-life (a preferred compound exhibits an *in vivo* half-life allowing for Q.I.D. dosing, preferably T.I.D. dosing, more preferably B.I.D. dosing, and most preferably once-a-day dosing). In addition, differential penetration of the blood brain barrier may be desirable for VR1 modulators used to treat pain by modulating CNS VR1 activity such that total daily oral doses as described above provide such modulation to a therapeutically effective extent, while low brain levels of VR1 modulators used to treat peripheral nerve mediated pain may be preferred (*i.e.*, such doses do not provide brain (*e.g.*, CSF) levels of the compound sufficient to significantly modulate VR1 activity). Routine assays that are well known in the art may be used to assess these properties, and identify superior compounds for a particular use. For example, assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Penetration of the blood brain barrier of a compound in humans may be predicted from the brain levels of the compound in laboratory animals given the compound (*e.g.*, intravenously). Serum protein binding may be predicted from albumin binding assays. Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described, for example, within Example 7 of published U.S. Application Number 2005/0070547.

As noted above, preferred compounds provided herein are nontoxic. In general, the term "nontoxic" shall be understood in a relative sense and is intended to refer to any substance that has been approved by the United States Food and Drug Administration ("FDA") for administration to mammals (preferably humans) or, in keeping with established criteria, is susceptible to approval by the FDA for administration to mammals (preferably humans). In addition, a highly preferred nontoxic compound generally satisfies one or more of the following criteria: (1) does not substantially inhibit cellular ATP production; (2) does not significantly prolong heart QT intervals; (3) does not cause substantial liver enlargement, or (4) does not cause substantial release of liver enzymes.

As used herein, a compound that does not substantially inhibit cellular ATP production is a compound that satisfies the criteria set forth in Example 8 of published U.S. Application Number 2005/0070547. In other words, cells treated as described therein with 100 μ M of such a compound exhibit ATP levels that are at least 50% of the ATP levels detected in untreated cells. In more highly preferred embodiments, such cells exhibit ATP levels that are at least 80% of the ATP levels detected in untreated cells.

A compound that does not significantly prolong heart QT intervals is a compound that does not result in a statistically significant prolongation of heart QT intervals (as determined by electrocardiography) in guinea pigs, minipigs or dogs upon administration of a dose that yields a serum concentration equal to the EC₅₀ or IC₅₀ for the compound. In certain preferred

embodiments, a dose of 0.01, 0.05, 0.1, 0.5, 1, 5, 10, 40 or 50 mg/kg administered parenterally or orally does not result in a statistically significant prolongation of heart QT intervals.

A compound does not cause substantial liver enlargement if daily treatment of laboratory rodents (*e.g.*, mice or rats) for 5-10 days with a dose that yields a serum concentration equal to the EC₅₀ or IC₅₀ for the compound results in an increase in liver to body weight ratio that is no more than 100% over matched controls. In more highly preferred embodiments, such doses do not cause liver enlargement of more than 75% or 50% over matched controls. If non-rodent mammals (*e.g.*, dogs) are used, such doses should not result in an increase of liver to body weight ratio of more than 50%, preferably not more than 25%, and more preferably not more than 10% over matched untreated controls. Preferred doses within such assays include 0.01, 0.05, 0.1, 0.5, 1, 5, 10, 40 or 50 mg/kg administered parenterally or orally.

Similarly, a compound does not promote substantial release of liver enzymes if administration of twice the minimum dose that yields a serum concentration equal to the EC₅₀ or IC₅₀ at VR1 for the compound does not elevate serum levels of ALT, LDH or AST in laboratory animals (*e.g.*, rodents) by more than 100% over matched mock-treated controls. In more highly preferred embodiments, such doses do not elevate such serum levels by more than 75% or 50% over matched controls. Alternatively, a compound does not promote substantial release of liver enzymes if, in an *in vitro* hepatocyte assay, concentrations (in culture media or other such solutions that are contacted and incubated with hepatocytes *in vitro*) that are equal to the EC₅₀ or IC₅₀ for the compound do not cause detectable release of any of such liver enzymes into culture medium above baseline levels seen in media from matched mock-treated control cells. In more highly preferred embodiments, there is no detectable release of any of such liver enzymes into culture medium above baseline levels when such compound concentrations are five-fold, and preferably ten-fold the EC₅₀ or IC₅₀ for the compound.

In other embodiments, certain preferred compounds do not inhibit or induce microsomal cytochrome P450 enzyme activities, such as CYP1A2 activity, CYP2A6 activity, CYP2C9 activity, CYP2C19 activity, CYP2D6 activity, CYP2E1 activity or CYP3A4 activity at a concentration equal to the EC₅₀ or IC₅₀ at VR1 for the compound.

Certain preferred compounds are not clastogenic (*e.g.*, as determined using a mouse erythrocyte precursor cell micronucleus assay, an Ames micronucleus assay, a spiral micronucleus assay or the like) at a concentration equal the EC₅₀ or IC₅₀ for the compound. In other embodiments, certain preferred compounds do not induce sister chromatid exchange (*e.g.*, in Chinese hamster ovary cells) at such concentrations.

For detection purposes, as discussed in more detail below, VR1 modulators provided herein may be isotopically-labeled or radiolabeled. For example, compounds may have one or more atoms replaced by an atom of the same element having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that

can be present in the compounds provided herein include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl . In addition, substitution with heavy isotopes such as deuterium (*i.e.*, ^2H) can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances.

PREPARATION OF HETEROARYL SUBSTITUTED PIPERAZINYL-PYRIDINE ANALOGUES

Heteroaryl substituted piperazinyl-pyridine analogues may generally be prepared using standard synthetic methods. Starting materials are commercially available from suppliers such as Sigma-Aldrich Corp. (St. Louis, MO), or may be synthesized from commercially available precursors using established protocols. By way of example, a synthetic route similar to that shown in any of the following Schemes may be used, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Each variable in the following schemes refers to any group consistent with the description of the compounds provided herein.

In the Schemes that follow, the term "catalyst" refers to a suitable transition metal catalyst such as, but not limited to, tetrakis(triphenylphosphine)palladium(0) or palladium(II) acetate. In addition, the catalytic systems may include ligands such as, but not limited to, 2-(Dicyclohexylphosphino)biphenyl and tri-*tert*-butylphosphine, and may also include a base such as K_3PO_4 , Na_2CO_3 or sodium or potassium *tert*-butoxide. Transition metal-catalyzed reactions can be carried out at ambient or elevated temperatures using various inert solvents including, but not limited to, toluene, dioxane, DMF, N-methylpyrrolidinone, ethyleneglycol, dimethyl ether, diglyme and acetonitrile. When used in conjunction with suitable metallo-aryl reagents, transition metal-catalyzed (hetero)aryl/aryl coupling reactions can be used to prepare the compounds encompassed in general structures 5C (Scheme 5), 6-D and 6-E (Scheme 6) and 7-F (Scheme 7). Commonly employed reagent/catalyst pairs include aryl boronic acid/palladium(0) (Suzuki reaction; Miyaura and Suzuki (1995) *Chemical Reviews* 95:2457) and aryl trialkylstannane/palladium(0) (Stille reaction; T. N. Mitchell, (1992) *Synthesis* 9:803-815), arylzinc/palladium(0) and aryl Grignard/nickel(II). In addition, metal-catalyzed (hetero)aryl/amine coupling reactions (Buchwald-Hartwig cross-coupling reaction; J. F. Hartwig, *Angew. Chem. Int. Ed.* 37:2046-2067 (1998)) can be used to prepare the compounds encompassed in general structures 4E (Scheme 4), 5E (Scheme 5), and 8F (Scheme 8) and 9H (Scheme 9).

Other definitions used in the following Schemes and elsewhere herein are:


BINAP	(<i>rac</i>)-2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl
Boc	t-butylcarbonyl
Bu	butyl

	CDCl ₃	deuterated chloroform
	δ	chemical shift
	DCM	dichloromethane or methylene chloride
	DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
5	DIBAL	diisobutylaluminum hydride
	DIEA	N,N-diisopropylethylamine
	DMA	N,N-dimethylacetamide
	DMF	dimethylformamide
	DMSO	dimethylsulfoxide
10	DPPF	1,1'-bis(diphenylphosphino)ferrocene
	Et ₃ N	triethylamine
	EtOAc	ethyl acetate
	EtOH	ethanol
	¹ H NMR	proton nuclear magnetic resonance
15	HOAc	acetic acid
	HPLC	high pressure liquid chromatography
	Hz	hertz
	KOAc	potassium acetate
	LCMS	liquid chromatography/mass spectrometry
20	MS	mass spectrometry
	(M+1)	mass + 1
	m-CPBA	<i>m</i> -chloroperbenzoic acid
	MeOH	methanol
	MsCl	methanesulfonyl chloride
25	NaNHCN	sodium cyanamide
	n-BuLi	n-butyl lithium
	NIS	<i>N</i> -iodosuccinimide
	Tf	-SO ₂ CF ₃
	Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
30	Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
	PhNEt ₂	diethyl-phenyl-amine, also referred to as N,N-diethylaniline
	PPh ₃	triphenylphosphine
	t-BuOK	potassium <i>tert</i> -butoxide
	THF	tetrahydrofuran

TLC


thin layer chromatography

Scheme 1




5

Scheme 2




10

Scheme 3




5

Scheme 4




Scheme 5




5

Scheme 6




10

Scheme 7




5


Scheme 8



Scheme 9



Scheme 10



5 In certain embodiments, a compound provided herein may contain one or more asymmetric carbon atoms, so that the compound can exist in different stereoisomeric forms. Such forms can be, for example, racemates or optically active forms. As noted above, all stereoisomers are encompassed by the present invention. Nonetheless, it may be desirable to obtain single enantiomers (*i.e.*, optically active forms). Standard methods for preparing single enantiomers

10 include asymmetric synthesis and resolution of the racemates. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography using, for example a chiral HPLC column.

Compounds may be radiolabeled by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. Each radioisotope is preferably carbon (*e.g.*, ^{14}C), hydrogen (*e.g.*, ^3H), sulfur (*e.g.*, ^{35}S), or iodine (*e.g.*, ^{125}I). Tritium labeled compounds may also be prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas using the compound as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as appropriate. Preparation of radiolabeled compounds may be conveniently performed by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds.

PHARMACEUTICAL COMPOSITIONS

The present invention also provides pharmaceutical compositions comprising one or more compounds provided herein, together with at least one physiologically acceptable carrier or excipient. Pharmaceutical compositions may comprise, for example, one or more of water, buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), ethanol, mineral oil, vegetable oil, dimethylsulfoxide, carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, adjuvants, polypeptides or amino acids such as glycine, antioxidants, chelating agents such as EDTA or glutathione and/or preservatives. In addition, other active ingredients may (but need not) be included in the pharmaceutical compositions provided herein.

Pharmaceutical compositions may be formulated for any appropriate manner of administration, including, for example, topical, oral, nasal, rectal or parenteral administration. The term parenteral as used herein includes subcutaneous, intradermal, intravascular (*e.g.*, intravenous), intramuscular, spinal, intracranial, intrathecal and intraperitoneal injection, as well as any similar injection or infusion technique. In certain embodiments, compositions suitable for oral use are preferred. Such compositions include, for example, tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs. Within yet other embodiments, pharmaceutical compositions may be formulated as a lyophilizate. Formulation for topical administration may be preferred for certain conditions (*e.g.*, in the treatment of skin conditions such as burns or itch). Formulation for direct administration into the bladder (intravesicular administration) may be preferred for treatment of urinary incontinence and overactive bladder.

Compositions intended for oral use may further comprise one or more components such as sweetening agents, flavoring agents, coloring agents and/or preserving agents in order to provide appealing and palatable preparations. Tablets contain the active ingredient in admixture with physiologically acceptable excipients that are suitable for the manufacture of tablets. Such excipients include, for example, inert diluents (*e.g.*, calcium carbonate, sodium carbonate, lactose,

calcium phosphate or sodium phosphate), granulating and disintegrating agents (*e.g.*, corn starch or alginic acid), binding agents (*e.g.*, starch, gelatin or acacia) and lubricating agents (*e.g.*, magnesium stearate, stearic acid or talc). The tablets may be uncoated or they may be coated by known techniques.

5 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent (*e.g.*, calcium carbonate, calcium phosphate or kaolin), or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium (*e.g.*, peanut oil, liquid paraffin or olive oil).

10 Aqueous suspensions contain the active material(s) in admixture with suitable excipients, such as suspending agents (*e.g.*, sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia); and dispersing or wetting agents (*e.g.*, naturally-occurring phosphatides such as lecithin, condensation products of an alkylene oxide with fatty acids such as polyoxyethylene stearate, condensation products of ethylene oxide with long chain aliphatic alcohols such as
15 heptadecaethyleneoxycetanol, condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides such as polyethylene sorbitan monooleate). Aqueous suspensions may also comprise one or more preservatives, such as ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or
20 more flavoring agents, and/or one or more sweetening agents, such as sucrose or saccharin.

 Oily suspensions may be formulated by suspending the active ingredient(s) in a vegetable oil (*e.g.*, arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and/or flavoring agents may be added to
25 provide palatable oral preparations. Such suspensions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and
30 suspending agents are exemplified by those already mentioned above. Additional excipients, such as sweetening, flavoring and coloring agents, may also be present.

 Pharmaceutical compositions may also be formulated as oil-in-water emulsions. The oily phase may be a vegetable oil (*e.g.*, olive oil or arachis oil), a mineral oil (*e.g.*, liquid paraffin) or a mixture thereof. Suitable emulsifying agents include naturally-occurring gums (*e.g.*, gum acacia
35 or gum tragacanth), naturally-occurring phosphatides (*e.g.*, soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol), anhydrides (*e.g.*, sorbitan monooleate) and condensation products of partial esters derived from fatty acids and hexitol with ethylene oxide

(e.g., polyoxyethylene sorbitan monoleate). An emulsion may also comprise one or more sweetening and/or flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, such as glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also comprise one or more demulcents,
5 preservatives, flavoring agents and/or coloring agents.

Formulations for topical administration typically comprise a topical vehicle combined with active agent(s), with or without additional optional components. Suitable topical vehicles and additional components are well known in the art, and it will be apparent that the choice of a vehicle will depend on the particular physical form and mode of delivery. Topical vehicles include
10 water; organic solvents such as alcohols (e.g., ethanol or isopropyl alcohol) or glycerin; glycols (e.g., butylene, isoprene or propylene glycol); aliphatic alcohols (e.g., lanolin); mixtures of water and organic solvents and mixtures of organic solvents such as alcohol and glycerin; lipid-based materials such as fatty acids, acylglycerols (including oils, such as mineral oil, and fats of natural or synthetic origin), phosphoglycerides, sphingolipids and waxes; protein-based materials such as
15 collagen and gelatin; silicone-based materials (both non-volatile and volatile); and hydrocarbon-based materials such as microsponges and polymer matrices. A composition may further include one or more components adapted to improve the stability or effectiveness of the applied formulation, such as stabilizing agents, suspending agents, emulsifying agents, viscosity adjusters, gelling agents, preservatives, antioxidants, skin penetration enhancers, moisturizers and sustained
20 release materials. Examples of such components are described in Martindale--The Extra Pharmacopoeia (Pharmaceutical Press, London 1993) and *Remington: The Science and Practice of Pharmacy*, 21st ed., Lippincott Williams & Wilkins, Philadelphia, PA (2005). Formulations may comprise microcapsules, such as hydroxymethylcellulose or gelatin-microcapsules, liposomes, albumin microspheres, microemulsions, nanoparticles or nanocapsules.

A topical formulation may be prepared in any of a variety of physical forms including, for example, solids, pastes, creams, foams, lotions, gels, powders, aqueous liquids and emulsions. The physical appearance and viscosity of such pharmaceutically acceptable forms can be governed by the presence and amount of emulsifier(s) and viscosity adjuster(s) present in the formulation. Solids are generally firm and non-pourable and commonly are formulated as bars or sticks, or in
30 particulate form; solids can be opaque or transparent, and optionally can contain solvents, emulsifiers, moisturizers, emollients, fragrances, dyes/colorants, preservatives and other active ingredients that increase or enhance the efficacy of the final product. Creams and lotions are often similar to one another, differing mainly in their viscosity; both lotions and creams may be opaque, translucent or clear and often contain emulsifiers, solvents, and viscosity adjusting agents, as well
35 as moisturizers, emollients, fragrances, dyes/colorants, preservatives and other active ingredients that increase or enhance the efficacy of the final product. Gels can be prepared with a range of viscosities, from thick or high viscosity to thin or low viscosity. These formulations, like those of

lotions and creams, may also contain solvents, emulsifiers, moisturizers, emollients, fragrances, dyes/colorants, preservatives and other active ingredients that increase or enhance the efficacy of the final product. Liquids are thinner than creams, lotions, or gels and often do not contain emulsifiers. Liquid topical products often contain solvents, emulsifiers, moisturizers, emollients, fragrances, dyes/colorants, preservatives and other active ingredients that increase or enhance the efficacy of the final product.

Suitable emulsifiers for use in topical formulations include, but are not limited to, ionic emulsifiers, cetearyl alcohol, non-ionic emulsifiers like polyoxyethylene oleyl ether, PEG-40 stearate, cetareth-12, cetareth-20, cetareth-30, cetareth alcohol, PEG-100 stearate and glyceryl stearate. Suitable viscosity adjusting agents include, but are not limited to, protective colloids or non-ionic gums such as hydroxyethylcellulose, xanthan gum, magnesium aluminum silicate, silica, microcrystalline wax, beeswax, paraffin, and cetyl palmitate. A gel composition may be formed by the addition of a gelling agent such as chitosan, methyl cellulose, ethyl cellulose, polyvinyl alcohol, polyquaterniums, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, carbomer or ammoniated glycyrrhizinate. Suitable surfactants include, but are not limited to, nonionic, amphoteric, ionic and anionic surfactants. For example, one or more of dimethicone copolyol, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, lauramide DEA, cocamide DEA, and cocamide MEA, oleyl betaine, cocamidopropyl phosphatidyl PG-dimonium chloride, and ammonium laureth sulfate may be used within topical formulations. Suitable preservatives include, but are not limited to, antimicrobials such as methylparaben, propylparaben, sorbic acid, benzoic acid, and formaldehyde, as well as physical stabilizers and antioxidants such as vitamin E, sodium ascorbate/ascorbic acid and propyl gallate. Suitable moisturizers include, but are not limited to, lactic acid and other hydroxy acids and their salts, glycerin, propylene glycol, and butylene glycol. Suitable emollients include lanolin alcohol, lanolin, lanolin derivatives, cholesterol, petrolatum, isostearyl neopentanoate and mineral oils. Suitable fragrances and colors include, but are not limited to, FD&C Red No. 40 and FD&C Yellow No. 5. Other suitable additional ingredients that may be included a topical formulation include, but are not limited to, abrasives, absorbents, anti-caking agents, anti-foaming agents, anti-static agents, astringents (e.g., witch hazel, alcohol and herbal extracts such as chamomile extract), binders/excipients, buffering agents, chelating agents, film forming agents, conditioning agents, propellants, opacifying agents, pH adjusters and protectants.

An example of a suitable topical vehicle for formulation of a gel is: hydroxypropylcellulose (2.1%); 70/30 isopropyl alcohol/water (90.9%); propylene glycol (5.1%); and Polysorbate 80 (1.9%). An example of a suitable topical vehicle for formulation as a foam is: cetyl alcohol (1.1%); stearyl alcohol (0.5%); Quaternium 52 (1.0%); propylene glycol (2.0%); Ethanol 95 PGF3 (61.05%); deionized water (30.05%); P75 hydrocarbon propellant (4.30%). All percents are by weight.

Typical modes of delivery for topical compositions include application using the fingers; application using a physical applicator such as a cloth, tissue, swab, stick or brush; spraying (including mist, aerosol or foam spraying); dropper application; sprinkling; soaking; and rinsing.

A pharmaceutical composition may be prepared as a sterile injectible aqueous or oleaginous suspension. The compound(s) provided herein, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Such a composition may be formulated according to the known art using suitable dispersing, wetting agents and/or suspending agents such as those mentioned above. Among the acceptable vehicles and solvents that may be employed are water, 1,3-butanediol, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils may be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectible compositions, and adjuvants such as local anesthetics, preservatives and/or buffering agents can be dissolved in the vehicle.

Pharmaceutical compositions may also be formulated as suppositories (*e.g.*, for rectal administration). Such compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

Compositions for inhalation typically can be provided in the form of a solution, suspension or emulsion that can be administered as a dry powder or in the form of an aerosol using a conventional propellant (*e.g.*, dichlorodifluoromethane or trichlorofluoromethane).

Pharmaceutical compositions may be formulated as sustained release or controlled-release formulations (*i.e.*, a formulation such as a capsule that effects a slow release of active ingredient(s) following administration). Such formulations may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Preferably the formulation provides a relatively constant level of release of active ingredient(s); the release profile can be varied using methods well known in the art, including (a) by varying the thickness or composition of the coating, (b) by altering the amount or manner of addition of plasticizer in the coating, (c) by including additional ingredients, such as release-modifying agents, (d) by altering the composition, particle size or particle shape of the matrix, and (e) by providing one or more passageways through the coating. The amount of modulator contained within a sustained release formulation depends upon, for example, the method of administration (*e.g.*, the site of implantation), the rate and expected duration of release and the nature of the condition to be treated or prevented.

In general, a sustained and/or controlled release formulation comprises a matrix and/or coating that delays disintegration and absorption in the gastrointestinal tract (or implantation site)

and thereby provides a delayed action or a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. Coatings that regulate release of the modulator include pH-dependent coatings (which may be used to release modulator in the stomach, and enteric coatings (which may be used to release modulator further along the gastrointestinal tract). pH dependent coatings include, for example, shellac, cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropylmethylcellulose phthalate, methacrylic acid ester copolymers and zein.

In addition to or together with the above modes of administration, a compound provided herein may be conveniently added to food or drinking water (*e.g.*, for administration to non-human animals including companion animals (such as dogs and cats) and livestock). Animal feed and drinking water compositions may be formulated so that the animal takes in an appropriate quantity of the composition along with its diet. It may also be convenient to present the composition as a premix for addition to feed or drinking water.

Compounds are generally administered in a therapeutically effective amount. Preferred systemic doses are no higher than 50 mg per kilogram of body weight per day (*e.g.*, ranging from about 0.001 mg to about 50 mg per kilogram of body weight per day), with oral doses generally being about 5-20 fold higher than intravenous doses (*e.g.*, ranging from 0.01 to 40 mg per kilogram of body weight per day).

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage unit will vary depending, for example, upon the patient being treated, the particular mode of administration and any other co-administered drugs. Dosage units generally contain between from about 10 µg to about 500 mg of active ingredient. Optimal dosages may be established using routine testing, and procedures that are well known in the art.

Pharmaceutical compositions may be packaged for treating conditions responsive to VR1 modulation (*e.g.*, treatment of exposure to vanilloid ligand or other irritant, pain, itch, obesity or urinary incontinence). Packaged pharmaceutical compositions generally include (i) a container holding a pharmaceutical composition that comprises at least one VR1 modulator as described herein and (ii) instructions (*e.g.*, labeling or a package insert) indicating that the contained composition is to be used for treating a condition responsive to VR1 modulation in the patient.

METHODS OF USE

VR1 modulators provided herein may be used to alter activity and/or activation of capsaicin receptors in a variety of contexts, both *in vitro* and *in vivo*. Within certain aspects, VR1 antagonists may be used to inhibit the binding of vanilloid ligand agonist (such as capsaicin and/or RTX) to capsaicin receptor *in vitro* or *in vivo*. In general, such methods comprise the step of contacting a capsaicin receptor with one or more VR1 modulators provided herein, in the presence of vanilloid ligand in aqueous solution and under conditions otherwise suitable for binding of the

ligand to capsaicin receptor. The VR1 modulator(s) are generally present at a concentration that is sufficient to alter the binding of vanilloid ligand to VR1 *in vitro* (using the assay provided in Example 5) and/or VR1-mediated signal transduction (using an assay provided in Example 6). The capsaicin receptor may be present in solution or suspension (*e.g.*, in an isolated membrane or cell preparation), or in a cultured or isolated cell. Within certain embodiments, the capsaicin receptor is expressed by a neuronal cell present in a patient, and the aqueous solution is a body fluid. Preferably, one or more VR1 modulators are administered to an animal in an amount such that the VR1 modulator is present in at least one body fluid of the animal at a therapeutically effective concentration that is 1 micromolar or less; preferably 500 nanomolar or less; more preferably 100 nanomolar or less, 50 nanomolar or less, 20 nanomolar or less, or 10 nanomolar or less. For example, such compounds may be administered at a therapeutically effective dose that is less than 20 mg/kg body weight, preferably less than 5 mg/kg and, in some instances, less than 1 mg/kg.

Also provided herein are methods for modulating, preferably reducing, the signal-transducing activity (*i.e.*, the calcium conductance) of a cellular capsaicin receptor. Such modulation may be achieved by contacting a capsaicin receptor (either *in vitro* or *in vivo*) with one or more VR1 modulators provided herein under conditions suitable for binding of the modulator(s) to the receptor. The VR1 modulator(s) are generally present at a concentration that is sufficient to alter the binding of vanilloid ligand to VR1 *in vitro* and/or VR1-mediated signal transduction as described herein. The receptor may be present in solution or suspension, in a cultured or isolated cell preparation or in a cell within a patient. For example, the cell may be a neuronal cell that is contacted *in vivo* in an animal. Alternatively, the cell may be an epithelial cell, such as a urinary bladder epithelial cell (urothelial cell) or an airway epithelial cell that is contacted *in vivo* in an animal. Modulation of signal transducing activity may be assessed by detecting an effect on calcium ion conductance (also referred to as calcium mobilization or flux). Modulation of signal transducing activity may alternatively be assessed by detecting an alteration of a symptom (*e.g.*, pain, burning sensation, broncho-constriction, inflammation, cough, hiccup, itch, urinary incontinence or overactive bladder) of a patient being treated with one or more VR1 modulators provided herein.

VR1 modulator(s) provided herein are preferably administered to a patient (*e.g.*, a human) orally or topically, and are present within at least one body fluid of the animal while modulating VR1 signal-transducing activity. Preferred VR1 modulators for use in such methods modulate VR1 signal-transducing activity *in vitro* at a concentration of 1 nanomolar or less, preferably 100 picomolar or less, more preferably 20 picomolar or less, and *in vivo* at a concentration of 1 micromolar or less, 500 nanomolar or less, or 100 nanomolar or less in a body fluid such as blood.

The present invention further provides methods for treating conditions responsive to VR1 modulation. Within the context of the present invention, the term "treatment" encompasses both

disease-modifying treatment and symptomatic treatment, either of which may be prophylactic (*i.e.*, before the onset of symptoms, in order to prevent, delay or reduce the severity of symptoms) or therapeutic (*i.e.*, after the onset of symptoms, in order to reduce the severity and/or duration of symptoms). A condition is "responsive to VR1 modulation" if it is characterized by inappropriate activity of a capsaicin receptor, regardless of the amount of vanilloid ligand present locally, and/or if modulation of capsaicin receptor activity results in alleviation of the condition or a symptom thereof. Such conditions include, for example, symptoms resulting from exposure to VR1-activating stimuli, pain, respiratory disorders (such as cough, asthma, chronic obstructive pulmonary disease, chronic bronchitis, cystic fibrosis and rhinitis, including allergic rhinitis, such as seasonal and perennial rhinitis, and non-allergic rhinitis), depression, itch, urinary incontinence, overactive bladder, hiccup and obesity, as described in more detail below. Such conditions may be diagnosed and monitored using criteria that have been established in the art. Patients may include humans, domesticated companion animals and livestock, with dosages as described above.

Treatment regimens may vary depending on the compound used and the particular condition to be treated; however, for treatment of most disorders, a frequency of administration of 4 times daily or less is preferred. In general, a dosage regimen of 2 times daily is more preferred, with once a day dosing particularly preferred. For the treatment of acute pain, a single dose that rapidly reaches effective concentrations is desirable. It will be understood, however, that the specific dose level and treatment regimen for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy. In general, the use of the minimum dose sufficient to provide effective therapy is preferred. Patients may generally be monitored for therapeutic effectiveness using medical or veterinary criteria suitable for the condition being treated or prevented.

Patients experiencing symptoms resulting from exposure to capsaicin receptor-activating stimuli include individuals with burns caused by heat, light, tear gas or acid and those whose mucous membranes are exposed (*e.g.*, via ingestion, inhalation or eye contact) to capsaicin (*e.g.*, from hot peppers or in pepper spray) or a related irritant such as acid, tear gas, infectious agent(s) or air pollutant(s). The resulting symptoms (which may be treated using VR1 modulators, especially antagonists, provided herein) may include, for example, pain, broncho-constriction and inflammation.

Pain that may be treated using the VR1 modulators provided herein may be chronic or acute and includes, but is not limited to, peripheral nerve-mediated pain (especially neuropathic pain).

Compounds provided herein may be used in the treatment of, for example, postmastectomy pain syndrome, stump pain, phantom limb pain, oral neuropathic pain, toothache (dental pain), denture pain, postherpetic neuralgia, diabetic neuropathy, chemotherapy-induced neuropathy, reflex

sympathetic dystrophy, trigeminal neuralgia, osteoarthritis, rheumatoid arthritis, fibromyalgia, Guillain-Barre syndrome, meralgia paresthetica, burning-mouth syndrome and/or pain associated with nerve and root damage, including as pain associated with peripheral nerve disorders (*e.g.*, nerve entrapment and brachial plexus avulsions, amputation, peripheral neuropathies including bilateral peripheral neuropathy, tic douloureux, atypical facial pain, nerve root damage, and arachnoiditis). Additional neuropathic pain conditions include causalgia (reflex sympathetic dystrophy - RSD, secondary to injury of a peripheral nerve), neuritis (including, for example, sciatic neuritis, peripheral neuritis, polyneuritis, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis and Gombault's neuritis), neuronitis, neuralgias (*e.g.*, those mentioned above, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migranous neuralgia, idiopathic neuralgia, intercostals neuralgia, mammary neuralgia, mandibular joint neuralgia, Morton's neuralgia, nasociliary neuralgia, occipital neuralgia, red neuralgia, Sluder's neuralgia, splenopalatine neuralgia, supraorbital neuralgia and vidian neuralgia), surgery-related pain, musculoskeletal pain, myofascial pain syndromes, AIDS-related neuropathy, MS-related neuropathy, central nervous system pain (*e.g.*, pain due to brain stem damage, sciatica, and ankylosing spondylitis), and spinal pain, including spinal cord injury-related pain. Headache, including headaches involving peripheral nerve activity may also be treated as described herein. Such pain includes, for example, such as sinus, cluster (*i.e.*, migranous neuralgia) and tension headaches, migraine, temporomandibular pain and maxillary sinus pain. For example, migraine headaches may be prevented by administration of a compound provided herein as soon as a pre-migrainous aura is experienced by the patient. Further conditions that can be treated as described herein include Charcot's pains, intestinal gas pains, ear pain, heart pain, muscle pain, eye pain, orofacial pain (*e.g.*, odontalgia), abdominal pain, gynaecological pain (*e.g.*, menstrual pain, dysmenorrhoea, pain associated with cystitis, labor pain, chronic pelvic pain, chronic prostatitis and endometriosis), acute and chronic back pain (*e.g.*, lower back pain), gout, scar pain, hemorrhoidal pain, dyspeptic pains, angina, nerve root pain, "non-painful" neuropathies, complex regional pain syndrome, homotopic pain and heterotopic pain – including pain associated with carcinoma, often referred to as cancer pain (*e.g.*, in patients with bone cancer), pain (and inflammation) associated with venom exposure (*e.g.*, due to snake bite, spider bite, or insect sting) and trauma associated pain (*e.g.*, post-surgical pain, episiotomy pain, pain from cuts, musculoskeletal pain, bruises and broken bones, and burn pain, especially primary hyperalgesia associated therewith). Additional pain conditions that may be treated as described herein include pain associated with respiratory disorders as described above, autoimmune diseases, immunodeficiency disorders, hot flashes, inflammatory bowel disease, gastroesophageal reflux disease (GERD), irritable bowel syndrome and/or inflammatory bowel disease.

Within certain aspects, VR1 modulators provided herein may be used for the treatment of mechanical pain. As used herein, the term "mechanical pain" refers to pain other than headache

pain that is not neuropathic or a result of exposure to heat, cold or external chemical stimuli. Mechanical pain includes physical trauma (other than thermal or chemical burns or other irritating and/or painful exposures to noxious chemicals) such as post-surgical pain and pain from cuts, bruises and broken bones; toothache; denture pain; nerve root pain; osteoarthritis; rheumatoid
5 arthritis; fibromyalgia; meralgia paresthetica; back pain; cancer-associated pain; angina; carpal tunnel syndrome; and pain resulting from bone fracture, labor, hemorrhoids, intestinal gas, dyspepsia, and menstruation.

Itching conditions that may be treated include psoriatic pruritus, itch due to hemodialysis, aguagenic pruritus, and itching associated with vulvar vestibulitis, contact dermatitis, insect bites
10 and skin allergies. Urinary tract conditions that may be treated as described herein include urinary incontinence (including overflow incontinence, urge incontinence and stress incontinence), as well as overactive or unstable bladder conditions (including bladder detrusor hyper-reflexia, detrusor hyper-reflexia of spinal origin and bladder hypersensitivity). In certain such treatment methods, VR1 modulator is administered via a catheter or similar device, resulting in direct injection of VR1
15 modulator into the bladder. Compounds provided herein may also be used as anti-tussive agents (to prevent, relieve or suppress coughing) and for the treatment of hiccup, and to promote weight loss in an obese patient.

Within other aspects, VR1 modulators provided herein may be used within combination therapy for the treatment of conditions involving pain and/or inflammatory components. Such
20 conditions include, for example, autoimmune disorders and pathologic autoimmune responses known to have an inflammatory component including, but not limited to, arthritis (especially rheumatoid arthritis), psoriasis, Crohn's disease, lupus erythematosus, irritable bowel syndrome, tissue graft rejection, and hyperacute rejection of transplanted organs. Other such conditions include trauma (*e.g.*, injury to the head or spinal cord), cardio- and cerebro-vascular disease and
25 certain infectious diseases.

Within such combination therapy, a VR1 modulator is administered to a patient along with an analgesic and/or anti-inflammatory agent. The VR1 modulator and analgesic and/or anti-inflammatory agent may be present in the same pharmaceutical composition, or may be administered separately in either order. Anti-inflammatory agents include, for example, non-
30 steroidal anti-inflammatory drugs (NSAIDs), non-specific and cyclooxygenase-2 (COX-2) specific cyclooxygenase enzyme inhibitors, gold compounds, corticosteroids, methotrexate, tumor necrosis factor (TNF) receptor antagonists, anti-TNF alpha antibodies, anti-C5 antibodies, and interleukin-1 (IL-1) receptor antagonists. Examples of NSAIDs include, but are not limited to ibuprofen (*e.g.*, ADVIL™, MOTRIN™), flurbiprofen (ANSAID™), naproxen or naproxen sodium (*e.g.*,
35 NAPROSYN, ANAPROX, ALEVE™), diclofenac (*e.g.*, CATAFLAM™, VOLTAREN™), combinations of diclofenac sodium and misoprostol (*e.g.*, ARTHROTEC™), sulindac (CLINORIL™), oxaprozin (DAYPRO™), diflunisal (DOLOBID™), piroxicam (FELDENE™),

indomethacin (INDOCIN™), etodolac (LODINE™), fenoprofen calcium (NALFON™), ketoprofen (*e.g.*, ORUDIS™, ORUVAIL™), sodium nabumetone (RELAFEN™), sulfasalazine (AZULFIDINE™), tolmetin sodium (TOLECTIN™), and hydroxychloroquine (PLAQUENIL™). One class of NSAIDs consists of compounds that inhibit cyclooxygenase (COX) enzymes; such compounds include celecoxib (CELEBREX™) and rofecoxib (VIOXX™). NSAIDs further include salicylates such as acetylsalicylic acid or aspirin, sodium salicylate, choline and magnesium salicylates (TRILISATE™), and salsalate (DISALCID™), as well as corticosteroids such as cortisone (CORTONE™ acetate), dexamethasone (*e.g.*, DECADRON™), methylprednisolone (MEDROL™), prednisolone (PRELONE™), prednisolone sodium phosphate (PEDIAPRED™), and prednisone (*e.g.*, PREDNICEN-M™, DELTASONE™, STERAPRED™). Further anti-inflammatory agents include meloxicam, rofecoxib, celecoxib, etoricoxib, parecoxib, valdecoxib and tilcoxib.

Suitable dosages for VR1 modulator within such combination therapy are generally as described above. Dosages and methods of administration of anti-inflammatory agents can be found, for example, in the manufacturer's instructions in the *Physician's Desk Reference*. In certain embodiments, the combination administration of a VR1 modulator with an anti-inflammatory agent results in a reduction of the dosage of the anti-inflammatory agent required to produce a therapeutic effect (*i.e.*, a decrease in the minimum therapeutically effective amount). Thus, preferably, the dosage of anti-inflammatory agent in a combination or combination treatment method is less than the maximum dose advised by the manufacturer for administration of the anti-inflammatory agent without combination administration of a VR1 antagonist. More preferably this dosage is less than $\frac{3}{4}$, even more preferably less than $\frac{1}{2}$, and highly preferably, less than $\frac{1}{4}$ of the maximum dose, while most preferably the dose is less than 10% of the maximum dose advised by the manufacturer for administration of the anti-inflammatory agent(s) when administered without combination administration of a VR1 antagonist. It will be apparent that the dosage amount of VR1 antagonist component of the combination needed to achieve the desired effect may similarly be affected by the dosage amount and potency of the anti-inflammatory agent component of the combination.

In certain preferred embodiments, the combination administration of a VR1 modulator with an anti-inflammatory agent is accomplished by packaging one or more VR1 modulators and one or more anti-inflammatory agents in the same package, either in separate containers within the package or in the same contained as a mixture of one or more VR1 antagonists and one or more anti-inflammatory agents. Preferred mixtures are formulated for oral administration (*e.g.*, as pills, capsules, tablets or the like). In certain embodiments, the package comprises a label bearing indicia indicating that the one or more VR1 modulators and one or more anti-inflammatory agents are to be taken together for the treatment of an inflammatory pain condition.

Within further aspects, VR1 modulators provided herein may be used in combination with one or more additional pain relief medications. Certain such medications are also anti-inflammatory agents, and are listed above. Other such medications are analgesic agents, including narcotic agents which typically act at one or more opioid receptor subtypes (*e.g.*, μ , κ and/or δ), preferably as agonists or partial agonists. Such agents include opiates, opiate derivatives and opioids, as well as pharmaceutically acceptable salts and hydrates thereof. Specific examples of narcotic analgesics include, within preferred embodiments, alfentanil, alphaprodine, anileridine, bezitramide, buprenorphine, butorphanol, codeine, diacetyldihydromorphine, diacetylmorphine, dihydrocodeine, diphenoxylate, ethylmorphine, fentanyl, heroin, hydrocodone, hydromorphone, isomethadone, levomethorphan, levorphanol, meperidine, metazocine, methadone, methorphan, metopon, morphine, nalbuphine, opium extracts, opium fluid extracts, powdered opium, granulated opium, raw opium, tincture of opium, oxycodone, oxymorphone, paregoric, pentazocine, pethidine, phenazocine, piminodine, propoxyphene, racemethorphan, racemorphan, sulfentanyl, thebaine and pharmaceutically acceptable salts and hydrates of the foregoing agents.

Other examples of narcotic analgesic agents include acetorphine, acetyldihydrocodeine, acetylmethadol, allylprodine, alpracetalmethadol, alphameprodine, alphamethadol, benzethidine, benzylmorphine, betacetylmethadol, betameprodine, betamethadol, betaprodine, clonitazene, codeine methylbromide, codeine-N-oxide, cyprenorphine, desomorphine, dextromoramide, diampromide, diethylthiambutene, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, drotebanol, ethanol, ethylmethylthiambutene, etonitazene, etorphine, etoxeridine, furethidine, hydromorphenol, hydroxypethidine, ketobemidone, levomoramide, levophenacylmorphan, methyl-desorphine, methyldihydromorphine, morpheridine, morphine methylpromide, morphine methylsulfonate, morphine-N-oxide, myrophin, naloxone, naltrexone, nicocodeine, nicomorphine, noracymethadol, norlevorphanol, normethadone, normorphine, norpipanone, pentazocaine, phenadoxone, phenampromide, phenomorphan, phenoperidine, piritramide, pholcodine, proheptazone, properidine, propirane, racemoramide, thebacon, trimeperidine and the pharmaceutically acceptable salts and hydrates thereof.

Further specific representative analgesic agents include, for example acetaminophen (paracetamol); aspirin and other NSAIDs described above; NR2B antagonists; bradykinin antagonists; anti-migraine agents; anticonvulsants such as oxcarbazepine and carbamazepine; antidepressants (such as TCAs, SSRIs, SNRIs, substance P antagonists, etc.); spinal blocks; gabapentin; asthma treatments (such as β_2 -adrenergic receptor agonists; leukotriene D_4 antagonists (*e.g.*, montelukast); TALWIN® Nx and DEMEROL® (both available from Sanofi Winthrop Pharmaceuticals; New York, NY); LEVO-DROMORAN®; BUPRENEX® (Reckitt & Coleman Pharmaceuticals, Inc.; Richmond, VA); MSIR® (Purdue Pharma L.P.; Norwalk, CT); DILAUDID® (Knoll Pharmaceutical Co.; Mount Olive, NJ); SUBLIMAZE®; SUFENTA®

(Janssen Pharmaceutica Inc.; Titusville, NJ); PERCOCET®, NUBAIN® and NUMORPHAN® (all available from Endo Pharmaceuticals Inc.; Chadds Ford, PA) HYDROSTAT® IR, MS/S and MS/L (all available from Richwood Pharmaceutical Co. Inc; Florence, KY), ORAMORPH® SR and ROXICODONE® (both available from Roxanne Laboratories; Columbus OH) and
5 STADOL® (Bristol-Myers Squibb; New York, NY). Still further analgesic agents include CB2-receptor agonists, such as AM1241, and compounds that bind to the $\alpha 2\delta$ subunit, such as Neurontin (Gabapentin) and pregabalin.

Representative anti-migraine agents for use in combination with a VR1 modulator provided herein include CGRP antagonists, ergotamines and 5-HT₁ agonists, such as sumatriptan,
10 naratriptan, zolmatriptan and rizatriptan.

Within still further aspects, VR1 modulators provided herein may be used in combination with one or more leukotriene receptor antagonists (*e.g.*, agents that inhibits the cysteinyl leukotriene CysLT₁ receptor). CysLT₁ antagonists include Montelukast (SINGULAIR®; Merck & Co., Inc.). Such combinations find use in the treatment of pulmonary disorders such as asthma.

15 For the treatment or prevention of cough, a VR1 modulator as provided herein may be used in combination with other medication designed to treat this condition, such as antibiotics, anti-inflammatory agents, cystinyl leukotrienes, histamine antagonists, corticosteroids, opioids, NMDA antagonists, proton pump inhibitors, nociceptin, neurokinin (NK1, NK2 and NK3) and bradykinin (BK1 and BK2) receptor antagonists, cannabinoids, blockers of Na⁺-dependent
20 channels and large conductance Ca⁺²-dependent K⁺-channel activators. Specific agents include dexbrompheniramine plus pseudoephedrine, loratadine, oxymetazoline, ipratropium, albuterol, beclomethasone, morphine, codeine, pholcodeine and dextromethorphan.

The present invention further provides combination therapy for the treatment of urinary incontinence. Within such aspects, a VR1 modulator provided herein may be used in combination
25 with other medication designed to treat this condition, such as estrogen replacement therapy, progesterone congeners, electrical stimulation, calcium channel blockers, antispasmodic agents, cholinergic antagonists, antimuscarinic drugs, tricyclic antidepressants, SNRIs, beta adrenoceptor agonists, phosphodiesterase inhibitors, potassium channel openers, nociceptin/orphanin FQ (OP4) agonists, neurokinin (NK1 and NK2) antagonists, P2X3 antagonists, musculotrophic drugs and
30 sacral neuromodulation. Specific agents include oxybutinin, emepronium, tolterodine, flavoxate, flurbiprofen, tolterodine, dicyclomine, propiverine, propantheline, dicyclomine, imipramine, doxepin, duloxetine, 1-deamino-8-D-arginine vasopressin, muscarinic receptor antagonists such as Tolterodine (DETROL®; Pharmacia Corporation) and anticholinergic agents such as Oxybutynin (DITROPAN®; Ortho-McNeil Pharmaceutical, Inc., Raritan, NJ).

35 Suitable dosages for VR1 modulator within such combination therapy are generally as described above. Dosages and methods of administration of other pain relief medications can be found, for example, in the manufacturer's instructions in the *Physician's Desk Reference*. In

certain embodiments, the combination administration of a VR1 modulator with one or more additional pain medications results in a reduction of the dosage of each therapeutic agent required to produce a therapeutic effect (*e.g.*, the dosage of one or both agent may be less than $\frac{3}{4}$, less than $\frac{1}{2}$, less than $\frac{1}{4}$ or less than 10% of the maximum dose listed above or advised by the manufacturer).

5 For use in combination therapy, pharmaceutical compositions as described above may further comprise one or more additional medications as described above. In certain such compositions, the additional medication is an analgesic. Also provided herein are packaged pharmaceutical preparations comprising one or more VR1 modulators and one or more additional medications (*e.g.*, analgesics) in the same package. Such packaged pharmaceutical preparations
10 generally include (i) a container holding a pharmaceutical composition that comprises at least one VR1 modulator as described herein; (ii) a container holding a pharmaceutical composition that comprises at least one additional medication (such as a pain relief and/or anti-inflammatory medication) as described above and (iii) instructions (*e.g.*, labeling or a package insert) indicating that the compositions are to be used simultaneously, separately or sequentially for treating or
15 preventing a condition responsive to VR1 modulation in the patient (such as a condition in which pain and/or inflammation predominates).

 Compounds that are VR1 agonists may further be used, for example, in crowd control (as a substitute for tear gas) or personal protection (*e.g.*, in a spray formulation) or as pharmaceutical agents for the treatment of pain, itch, urinary incontinence or overactive bladder via capsaicin
20 receptor desensitization. In general, compounds for use in crowd control or personal protection are formulated and used according to conventional tear gas or pepper spray technology.

 Within separate aspects, the present invention provides a variety of non-pharmaceutical *in vitro* and *in vivo* uses for the compounds provided herein. For example, such compounds may be labeled and used as probes for the detection and localization of capsaicin receptor (in samples such
25 as cell preparations or tissue sections, preparations or fractions thereof). In addition, compounds provided herein that comprise a suitable reactive group (such as an aryl carbonyl, nitro or azide group) may be used in photoaffinity labeling studies of receptor binding sites. In addition, compounds provided herein may be used as positive controls in assays for receptor activity, as standards for determining the ability of a candidate agent to bind to capsaicin receptor, or as
30 radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT). Such methods can be used to characterize capsaicin receptors in living subjects. For example, a VR1 modulator may be labeled using any of a variety of well known techniques (*e.g.*, radiolabeled with a radionuclide such as tritium, as described herein), and incubated with a sample for a suitable incubation time (*e.g.*, determined by first
35 assaying a time course of binding). Following incubation, unbound compound is removed (*e.g.*, by washing), and bound compound detected using any method suitable for the label employed (*e.g.*, autoradiography or scintillation counting for radiolabeled compounds; spectroscopic methods

may be used to detect luminescent groups and fluorescent groups). As a control, a matched sample containing labeled compound and a greater (e.g., 10-fold greater) amount of unlabeled compound may be processed in the same manner. A greater amount of detectable label remaining in the test sample than in the control indicates the presence of capsaicin receptor in the sample. Detection assays, including receptor autoradiography (receptor mapping) of capsaicin receptor in cultured cells or tissue samples may be performed as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York.

Compounds provided herein may also be used within a variety of well known cell separation methods. For example, modulators may be linked to the interior surface of a tissue culture plate or other support, for use as affinity ligands for immobilizing and thereby isolating, capsaicin receptors (e.g., isolating receptor-expressing cells) *in vitro*. Within one preferred embodiment, a modulator linked to a fluorescent marker, such as fluorescein, is contacted with the cells, which are then analyzed (or isolated) by fluorescence activated cell sorting (FACS).

VR1 modulators provided herein may further be used within assays for the identification of other agents that bind to capsaicin receptor. In general, such assays are standard competition binding assays, in which bound, labeled VR1 modulator is displaced by a test compound. Briefly, such assays are performed by: (a) contacting capsaicin receptor with a radiolabeled VR1 modulator as described herein, under conditions that permit binding of the VR1 modulator to capsaicin receptor, thereby generating bound, labeled VR1 modulator; (b) detecting a signal that corresponds to the amount of bound, labeled VR1 modulator in the absence of test agent; (c) contacting the bound, labeled VR1 modulator with a test agent; (d) detecting a signal that corresponds to the amount of bound labeled VR1 modulator in the presence of test agent; and (e) detecting a decrease in signal detected in step (d), as compared to the signal detected in step (b).

The following Examples are offered by way of illustration and not by way of limitation. Unless otherwise specified all reagents and solvent are of standard commercial grade and are used without further purification. Using routine modifications, the starting materials may be varied and additional steps employed to produce other compounds provided herein.

EXAMPLES

Mass spectroscopy data in this and the following Examples is Electrospray MS, obtained in positive ion mode using a Micromass Time-of-Flight LCT (Micromass, Beverly MA), equipped with a Waters 600 pump (Waters Corp., Milford, MA), Waters 996 photodiode array detector, Gilson 215 autosampler (Gilson, Inc. Middleton, WI), and a Gilson 841 microinjector. MassLynx (Advanced Chemistry Development, Inc; Toronto, Canada) version 4.0 software with OpenLynx processing is used for data collection and analysis. MS conditions are as follows: capillary voltage = 3.5 kV; cone voltage = 30 V, desolvation and source temperature = 350°C and 120°C,

respectively; mass range = 181-750 with a scan time of 0.22 seconds and an interscan delay of 0.05 minutes.

Sample volume of 1 microliter is injected onto a 50x4.6mm Chromolith SpeedROD RP-18e column (Merck KGaA, Darmstadt, Germany), and eluted using a 2-phase linear gradient at 6ml/min flow rate. Sample is detected using total absorbance count over the 220-340nm UV range. The elution conditions are: Mobile Phase A- 95/5/0.05 Water/Methanol/TFA; Mobile Phase B-5/95/0.025 Water/Methanol/TFA. The following gradient is used:

Gradient:	Time(min)	%B
	0	10
	0.5	100
	1.2	100
	1.21	10

Inject to inject cycle 2.2 minutes.


EXAMPLE 1

Preparation of Representative Heteroaryl Substituted Piperazinyl-Pyridine Analogues

This Example illustrates the preparation of representative heteroaryl substituted piperazinyl-pyridine analogues.


A. 4-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-[4-(1H-tetrazol-5-yl)-piperidin-1-yl]pyrimidine (Compound 1)

1. *Piperidine-4-carbonitrile*




Stir 4-cyano-piperidine-1-carboxylic acid tert-butyl ester (Oakwood Products, Inc., 5g, 0.024 mol) in dry dioxane with an excess of 4 N HCl dioxane solution (100 mL). After 2 hours, collect the solid by filtration and wash it with ether (3×). Suspend the solid in toluene and add 14 g of amberlyst bicarbonate resin. Stir overnight, and then filter, washing the resin with additional toluene. Stir the collected resin for an additional 1 hour with a solution of MeOH/ Et₃N (4:1). Combine the two organic solutions and concentrate under reduced pressure. Take up in DCM, dry (Na₂SO₄), and concentrate under reduced pressure to give the freebase product as an oil.

2. *2,4-dichloro-6-(4-fluorophenyl)pyrimidine*




Dissolve 4-fluorobromobenzene (8.75 g, 0.05 moles) in anhydrous ether (80 mL) under nitrogen atmosphere and cool to -78°C . Add dropwise 1.6 M n-BuLi (34 mL, 0.055 moles) and stir at -78°C for 45 minutes. Dissolve 2,4-dichloropyrimidine (7.45 g, 0.05 moles) in Et_2O (100 mL), add dropwise to the reaction mixture and warm the reaction mixture to -30°C . Stir at this temperature for 30 minutes, followed by 0°C for 30 minutes. Quench the reaction mixture with acetic acid (3.15 mL, 0.055 moles) and water (0.5 mL, 0.027 moles) dissolved in THF (5.0 mL). Add dropwise a THF (40 mL) solution of DDQ (11.9 g, 0.053 moles) to the reaction mixture. Bring the reaction mixture to room temperature and stir at room temperature for 30 minutes. Cool the reaction mixture to 0°C , add 3.0 N aq. NaOH (35 mL) and stir for 30 minutes. Decant the organic layer from the reaction mixture and wash the brown solid with Et_2O (3 x 100 mL). Combine the organic layers, wash several times with saturated NaCl solution and dry with MgSO_4 . Filter and evaporate under vacuum to afford a brown colored solid. Purify the crude by flash column chromatography using 5% EtOAc / hexane to afford the product as white solid.

3. *1-[2-Chloro-6-(4-fluoro-phenyl)-pyrimidin-4-yl]-piperidine-4-carbonitrile*



Stir a mixture of 2,4-dichloro-6-(4-fluorophenyl)pyrimidine (3.6g, 0.015 mol), piperidine-4-carbonitrile (1.8g, 0.016 mol), and K_2CO_3 (4.1g, 0.03 mol) in DMA at room temperature for 48 hours. Partition the mixture between EtOAc and brine and separate the layers. Wash the organic layer with 10% NaOH solution (2x) followed by brine. Dry the solution (Na_2SO_4) and concentrate under reduced pressure to give the crude product. Purify using flash column chromatography with silica gel (30% to 50% EtOAc/ hexanes).


4. *1-[6-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperidine-4-carbonitrile*



Heat a solution of 1-[2-chloro-6-(4-fluoro-phenyl)-pyrimidin-4-yl]-piperidine-4-carbonitrile (3.13g, 0.01 mol) and 2-methylpyrrolidine (5 mL, 0.05 mol) in DMA at 130°C for 10 hours. Partition the cooled mixture between EtOAc and brine and separate the layers. Wash the

organic layer with 10% NaOH solution (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using flash column chromatography with silica gel (30% to 50% EtOAc/ hexanes) to give the product as a foam.


5. 4-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-[4-(1H-tetrazol-5-yl)-piperidin-1-yl]pyrimidine



Heat a mixture of 1-[6-(4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperidine-4-carbonitrile (367 mg, 1.0 mmol), sodium azide (72 mg, 1.1 mmol), and ammonium chloride (60 mg, 1.1 mmol) in DMF in a sealed tube at 125°C for 24 hours. Partition the cooled mixture between EtOAc and brine and separate the layers. Wash the organic layer with water (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using flash column chromatography with silica gel to give the title product. (M+H) = 409; ¹H NMR (400 MHz, CDCl₃): δ 7.85 (m, 2H), 7.01 (m, 2H), 6.16 (s, 1H) 4.35 (m, 3H), 3.60 (m, 2H), 3.0 (m, 4H), 1.55-2.1 (m, 7H), 1.15 (d, 3H). The IC₅₀ determined as described in Example 6 is less than 1 micromolar.

B. 4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine (Compound 2)


1. 2-Chloro-4-(4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyrimidine



Stir a mixture of 2,4-dichloro-6-(4-fluorophenyl)pyrimidine (200 mg, 0.82 mmol), 1-(1H-imidazol-2-yl)-piperazine (150 mg, 0.91 mmol; prepared essentially as described in EP 0233051), and K₂CO₃ (239 mg, 1.7 mmol) in DMA at room temperature for 5 hours. Partition the mixture between EtOAc and 10% NaOH and separate the layers. Wash the organic layer with 10% NaOH

(3×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using preparative plate chromatography (5% MeOH/DCM eluent) to give the product as a clear oil that crystallizes upon standing.


2. 4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine



Heat a solution of 2-chloro-4-(4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyrimidine (173 mg, 0.47 mmol) and 2-methylpyrrolidine (237 microliters, 2.33 mmol) in DMA at 110°C for 16 hours. Partition the cooled mixture between EtOAc and 10% NaOH and separate the layers. Wash the organic layer with 10% NaOH solution (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using preparative plate silica gel chromatography (2 × 2mm, 10% MeOH/DCM eluent) to yield the title product. Prepare the HCl salt by dissolving in EtOAc and adding excess HCl/ether solution to give the HCl salt as an off-white solid. (M+H) = 424; ¹H NMR of freebase (300 MHz, CDCl₃): δ 8.01 (m, 2H), 7.09 (t, 2H), 6.80 (s, 1H), 6.70 (s, 1H), 4.34 (m, 1H), 3.80 (m, 4H), 3.67 (m, 2H), 3.55 (s, 3H), 3.16 (m, 4H), 2.05 (m, 2H), 1.90 (m, 1H), 1.70 (m, 1H), 1.31 (d, 3H, J=6.1 Hz). The IC₅₀ determined as described in Example 6 is less than 1 micromolar.


C. 4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine (Compound 3)

1. 1-[6-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperidine-4-carbaldehyde




Cool a solution of 1-[6-(4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperidine-4-carbonitrile (811 mg, 2.22 mmol) in DCM to -78°C using a dry ice acetone bath. Add DIBAL (1M in hexanes) dropwise and stir for 3 hours. Add excess Na₂SO₄·10 H₂O and stir the suspension for 10 minutes before removing the cooling bath. Stir for several hours at room temperature then filter the mixture through celite. Concentrate under reduced pressure and chromatograph the crude product on a silica gel flash column (20% to 30% EtOAc / hexanes eluent) to give the product as a white foam.

2. 4-(4-Fluoro-phenyl)-6-[4-(1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine



Stir a mixture of 1-[6-(4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperidine-4-carbaldehyde (134 mg, 0.364 mmol), glyoxal (40% aqueous solution, 106 microliters, 0.727 mmol), and concentrated ammonium hydroxide (~16.5 M, 66 microliters, 1.1 mmol) in MeOH at room temperature for 16 hours. Concentrate the solution under reduced pressure, load onto two 2mm silica gel preparative TLC plates, and elute with 5% MeOH(w/ 2N NH₃)/DCM to afford pure product. (M+H) = 407; ¹H NMR (300 MHz, CDCl₃): δ 7.98 (m, 2H), 7.08 (t, 2H), 6.94 (s, 2H), 6.14 (s, 1H), 4.50 (m, 2H), 4.31 (m, 1H), 3.65 (m, 2H), 3.00 (m, 4H), 1.6-2.12 (m, 7H), 1.29 (d, 3H, J = 6.32 Hz).

3. 4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine



To a stirring solution of 4-(4-fluoro-phenyl)-6-[4-(1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine (65 mg, 0.16 mmol) in dry DMF, add NaH as a 60% suspension in mineral oil (~8mg). Stir the mixture at room temperature for 5 minutes and then add

2 drops of MeI. After 30 minutes, partition the mixture between EtOAc and 10% NaOH and separate the layers. Wash the organic layer with 10% NaOH solution (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using preparative plate silica gel chromatography to give the title product. (M+H) = 421; ¹H NMR (300 MHz, CDCl₃): δ 7.98 (m, 2H), 7.08 (t, 2H), 6.93 (s, 1H), 6.79 (s, 1H), 6.26 (s, 1H), 4.55 (m, 2H), 4.31 (m, 1H), 3.65 (m, 5H), 3.0 (m, 4H) 1.8-2.1 (m, 6H), 1.70 (m, 1H), 1.30 (d, 3H, J= 6.3 Hz). The IC₅₀ determined as described in Example 6 is less than 1 micromolar.


EXAMPLE 2

Preparation of Additional Representative Heteroaryl Substituted Piperazinyl-Pyridine Analogues

This Example illustrates the preparation of additional representative heteroaryl substituted piperazinyl-pyridine analogues.


A. 4-(3-Chloro-4-fluoro-phenyl)-6-[4-(4-chloro-[1,2,5]thiadiazol-3-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine (Compound 4)

1. 2,4-Dichloro-6-(3-chloro-4-fluoro-phenyl)-pyrimidine



This compound is prepared from 2,4-dichloropyrimidine and 4-bromo-2-chloro-1-fluorobenzene using a procedure analogous to that used for the preparation of 2,4-dichloro-6-(4-fluorophenyl)pyrimidine in Example 1A, step 2.


2. 4-[2-Chloro-6-(3-chloro-4-fluoro-phenyl)-pyrimidin-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



Stir a mixture of 2,4-dichloro-6-(4-fluoro-phenyl)-pyrimidine (2.8 g, 11.52 mmol), piperazine-1-carboxylic acid tert-butyl ester (12.1 mmol), and K₂CO₃ (3.2 g, 23.0 mmol) in DMA


at room temperature for 16 hours. Dilute with water, extract with EtOAc, and wash with brine. Dry the organic layer (Na₂SO₄) and concentrate under reduced pressure. Purify the residue by flash column eluting with EtOAc-Hexanes (1:4) to afford 4-[2-chloro-6-(3-chloro-4-fluoro-phenyl)-pyrimidin-4-yl]-piperazine-1-carboxylic acid *tert*-butyl ester.

- 5 3. 4-[6-(3-Chloro-4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperazine-1-carboxylic acid *tert*-butyl ester




- 10 Heat a mixture of 4-[2-chloro-6-(3-chloro-4-fluoro-phenyl)-pyrimidin-4-yl]-piperazine-1-carboxylic acid *tert*-butyl ester (12.3 mmol), 2-methyl-pyrrolidine (16.0 mmol), and K₂CO₃ (5.2 g, 37.8 mmol) in DMA at 120°C for 16 hours. Dilute with water, extract with EtOAc, and wash with brine. Dry the organic layer (Na₂SO₄) and concentrate under reduced pressure. Purify the residue by flash column eluting with EtOAc-Hexanes (1:4) to afford the product.

4. 4-(3-Chloro-4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-piperazin-1-yl-pyrimidine



- 15 Stir 4-[6-(3-chloro-4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidin-4-yl]-piperazine-1-carboxylic acid *tert*-butyl ester (9.88 mmol) in 4M HCl-dioxane (50 mL) for 40 minutes. Concentrate and partition between EtOAc and sat. NaHCO₃. Dry the organic layer (Na₂SO₄) and concentrate under reduced pressure to afford the product.


5. *4-(3-Chloro-4-fluoro-phenyl)-6-[4-(4-chloro-[1,2,5]thiadiazol-3-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine*



Heat a mixture of 4-(3-chloro-4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-piperazin-
 5 1-yl-pyrimidine (114 mg, 0.303 mmol), 3,4-dichloro-[1,2,5]thiadiazole (0.333 mmol) and
 diisopropylethylamine (0.33 mmol) in DMSO at 100°C for 16 hours. Partition the cooled mixture
 between EtOAc and brine and separate the layers. Wash the organic layer with 10% NaOH
 solution (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced
 pressure to give the crude product. Purify using flash column chromatography with silica gel
 10 (EtOAc/ hexanes eluent) to give the title product. (M+H) = 495; ¹H NMR (400 MHz, CDCl₃): δ
 8.08 (d, 1H), 7.90 (m, 1H), 7.19 (t, 1H), 6.24 (s, 1H), 4.35 (m, 1H), 3.83 (m, 4H), 3.70 (M, 2H),
 3.60 (m, 4H), 2.06 (m, 2H), 1.93 (m, 1H), 1.70 (m, 1H).1.30 (d, 3H, J=6.23 hz). The IC₅₀
 determined as described in Example 6 is less than 1 micromolar.

15 B. 4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-2-(R)-2-methyl-pyrrolidin-1-yl)-pyrimidine (Compound 5)


1. *2-Chloro-4-(3-chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyrimidine*



20 Stir a mixture of 2,4-dichloro-6-(3-chloro-4-fluoro-phenyl)-pyrimidine (211 mg, 0.76
 mmol), 1-(1H-imidazol-2-yl)-piperazine (139 mg, 0.84 mmol) (prepared essentially as described in
 EP 0233051), and K₂CO₃ (222 mg, 1.6 mmol) in DMA at room temperature for 16 hours. Partition
 the mixture between EtOAc and 10% NaOH and separate the layers. Wash the organic layer with
 10% NaOH (3×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced

pressure to give the crude product. Purify using preparative plate chromatography (5% MeOH/DCM eluent).

2. *4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-2-((R)-2-methyl-pyrrolidin-1-yl)-pyrimidine*




5

Heat a solution of 2-chloro-4-(3-chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyrimidine (50 mg, 0.123 mmol) and (R)-2-methylpyrrolidine hydrochloride (prepared essentially as described in US Published Application 2004/0171845; 22 mg, 0.18 mmol) in DMA at 110°C for 16 hours. Partition the cooled mixture between EtOAc and 10% NaOH and separate the layers. Wash the organic layer with 10% NaOH solution (2×) followed by brine. Dry the solution (Na₂SO₄) and concentrate under reduced pressure to give the crude product. Purify using preparative plate silica gel chromatography (1 × 2mm, 5% MeOH/DCM eluent) to give the title product. Prepare the HCl salt by dissolving in EtOAc and adding excess HCl/ether solution to give the HCl salt as an off-white solid. (M+H) = 456; ¹H NMR (300 MHz, CDCl₃): δ 8.06 (dd, 1H), 7.88 (m, 1H), 7.26 (s, 1H), 7.18 (t, 1H), 6.80 (s, 1H), 6.70 (s, 1H), 6.24 (s, 1H), 4.33 (M, 1H), 3.80 (m, 4H), 3.67 (m, 2H), 3.17 (t, 4H), 2.06 (m, 2H), 1.90 (m, 1H), 1.70 (m, 1H), 1.29 (d, 3H, J=6.3 Hz). The IC₅₀ determined as described in Example 6 is less than 1 micromolar.

15

C. 4-{4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyridin-2-yl}-morpholine (Compound 6)


20 1. *2,3-dichloro-4-(3-chloro-4-fluoro-phenyl)-pyridine*



To a de-gassed mixture of 3-chloro-4-fluorophenylboronic acid (77 mg, 0.44 mmol), 4-bromo-2,6-dichloro-pyridine (prepared essentially as described in Talik & Plazek (1959) *Rocz. Chem.* 33:387-392; 100 mg, 0.44 mmol), and 2M Na₂CO₃ (0.55 mmol), in DME (4mL) under

nitrogen, add $\text{Pd}(\text{PPh}_3)_4$ (0.026 mmol). Stir the mixture at 80°C for 16 hours, concentrate, and extract with EtOAc. Dry over Na_2SO_4 , concentrate under vacuum, and purify by preparative TLC (9:1 hexanes/EtOAc) to give 2,3-dichloro-4-(3-chloro-4-fluoro-phenyl)-pyridine.


2. 4-[6-chloro-4-(3-chloro-4-fluoro-phenyl)-pyridin-2-yl]-morpholine



5

Stir a solution of 2,6-dichloro-4-(3-chloro-4-fluoro-phenyl)-pyridine (100 mg) in morpholine (2 mL) 3 hours at 80°C, concentrate, partition between H_2O and EtOAc, dry over Na_2SO_4 , and concentrate under vacuum. Purify by preparative TLC (3:1 hexanes/EtOAc) to give 4-[6-chloro-4-(3-chloro-4-fluoro-phenyl)-pyridin-2-yl]-morpholine.

10 3. 4-{4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyridin-2-yl}-morpholine



To a de-gassed mixture of 4-[6-chloro-4-(3-chloro-4-fluoro-phenyl)-pyridin-2-yl]-morpholine (50 mg, 0.153 mmol), 1-(1H-imidazol-2-yl)-piperazine (0.183 mmol), and 1M (THF) *t*-BuOK (0.183 mmol), in toluene (3mL) under nitrogen, add $\text{Pd}_2(\text{dba})_3$ (0.006 mmol) and BINAP (0.008 mmol). Stir the mixture at 80°C overnight, concentrate, extract with EtOAc. Dry over Na_2SO_4 , concentrate under vacuum, and purify by preparative TLC (MeOH/DCM eluent) to give 4-{4-(3-chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperazin-1-yl]-pyridin-2-yl}-morpholine.





20






EXAMPLE 3






Additional Representative Heteroaryl Substituted Piperazinyl-Pyridine Analogues






Using routine modifications, the starting materials may be varied and additional steps employed to produce other compounds provided herein. Compounds listed in Table I are prepared using such methods.






Table I






Compound	Name
7 	4-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-[4-(5-methyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidine
8 	4-(4-Fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-2-pyrrolidin-1-yl-pyrimidine
9 	4-(4-Fluoro-phenyl)-6-[4-(5-methyl-[1,2,3]thiadiazol-4-yl)-piperazin-1-yl]-2-piperidin-1-yl-pyrimidine
10 	4-(4-Fluoro-phenyl)-6-[4-(4-methyl-isothiazol-3-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine


Compound	Name
11 	4-(4-Fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-6-[4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidine
12 	4-(4-Fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-2-pyrrolidin-1-yl-pyrimidine
13 	4-[4-(5-Chloro-[1,2,3]thiadiazol-4-yl)-piperazin-1-yl]-6-(4-fluoro-phenyl)-2-piperidin-1-yl-pyrimidine
14 	4-[4-(4,5-Dichloro-isothiazol-3-yl)-piperazin-1-yl]-6-(4-fluoro-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
15 	4-{4-(3-Chloro-4-fluoro-phenyl)-6-[4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-morpholine

Compound	Name
16 	4-(4-Fluoro-3-methyl-phenyl)-2-(2-methyl-azetidin-1-yl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidine
17 	4-[4-(5-Chloro-[1,2,3]thiadiazol-4-yl)-piperazin-1-yl]-6-(4-fluoro-3-methyl-phenyl)-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
18 	4-(3-Chloro-4-fluoro-phenyl)-6-[4-(4-chloro-isothiazol-3-yl)-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
19 	4-{4-(3-Chloro-4-fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-3-methyl-morpholine
20 	2-Azetidin-1-yl-4-(4-fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidine

Compound	Name
21 	4-(3-Chloro-4-fluoro-phenyl)-6-[4-(5-chloro-[1,2,3]thiadiazol-4-yl)-2-methyl-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
22 	4-(3-Chloro-4-fluoro-phenyl)-6-[4-(4-chloro-isothiazol-3-yl)-2-methyl-piperazin-1-yl]-2-piperidin-1-yl-pyrimidine
23 	{4-(3-Chloro-4-fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-diethyl-amine
24 	{4-(4-Fluoro-phenyl)-6-[2-methyl-4-(5-trifluoromethyl-thiazol-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-isopropyl-methyl-amine
25 	2-({4-(3-Chloro-4-fluoro-phenyl)-6-[4-(5-chloro-[1,2,3]thiadiazol-4-yl)-2-methyl-piperazin-1-yl]-pyrimidin-2-yl }-methyl-amino)-propan-1-ol

Compound	Name
26 	4-(3-Chloro-4-fluoro-phenyl)-6-[4-(4-chloro-isothiazol-3-yl)-2-methyl-piperazin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
27 	4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
28 	4-(4-Fluoro-phenyl)-6-[2-methyl-4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-pyrrolidin-1-yl-pyrimidine
29 	4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-piperidin-1-yl-pyrimidine
30 	4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-piperidin-1-yl-pyrimidine

Compound	Name
31 	4-(4-Fluoro-3-methyl-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-2-(2-methyl-pyrrolidin-1-yl)-pyrimidine
32 	(1-{4-(3-Chloro-4-fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-pyrimidin-2-yl}-pyrrolidin-2-yl)-methanol
33 	Diethyl-{4-(4-fluoro-phenyl)-6-[2-methyl-4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-pyrimidin-2-yl}-amine
34 	4-{4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-pyrimidin-2-yl}-morpholine
35 	2-({4-(4-Fluoro-phenyl)-6-[4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-pyrimidin-2-yl}-methyl-amino)-propan-1-ol

Compound	Name
	4-{4-(4-Fluoro-phenyl)-6-[2-methyl-4-(1-methyl-1H-imidazol-2-yl)-piperidin-1-yl]-pyrimidin-2-yl}-3-methyl-morpholine

EXAMPLE 4

VR1-Transfected Cells and Membrane Preparations

5 This Example illustrates the preparation of VR1-transfected cells and VR1-containing membrane preparations for use in capsaicin binding assays (Example 5).

A cDNA encoding full length human capsaicin receptor (SEQ ID NO:1, 2 or 3 of U.S. Patent No. 6,482,611) is subcloned in the plasmid pBK-CMV (Stratagene, La Jolla, CA) for recombinant expression in mammalian cells.

10 Human embryonic kidney (HEK293) cells are transfected with the pBK-CMV expression construct encoding the full length human capsaicin receptor using standard methods. The transfected cells are selected over two weeks in media containing G418 (400 µg/ml) to obtain a pool of stably transfected cells. Independent clones are isolated from this pool by limiting dilution to obtain clonal stable cell lines for use in subsequent experiments.

15 For radioligand binding experiments, cells are seeded in T175 cell culture flasks in media without antibiotics and grown to approximately 90% confluency. The flasks are then washed with PBS and harvested in PBS containing 5 mM EDTA. The cells are pelleted by gentle centrifugation and stored at -80°C until assayed.

Previously frozen cells are disrupted with the aid of a tissue homogenizer in ice-cold
 20 HEPES homogenization buffer (5mM KCl 5, 5.8mM NaCl, 0.75mM CaCl₂, 2mM MgCl₂, 320 mM sucrose, and 10 mM HEPES pH 7.4). Tissue homogenates are first centrifuged for 10 minutes at 1000 x g (4°C) to remove the nuclear fraction and debris, and then the supernatant from the first centrifugation is further centrifuged for 30 minutes at 35,000 x g (4°C) to obtain a partially purified membrane fraction. Membranes are resuspended in the HEPES homogenization buffer
 25 prior to the assay. An aliquot of this membrane homogenate is used to determine protein concentration via the Bradford method (BIO-RAD Protein Assay Kit, #500-0001, BIO-RAD, Hercules, CA).

EXAMPLE 5

Capsaicin Receptor Binding Assay

This Example illustrates a representative assay of capsaicin receptor binding that may be used to determine the binding affinity of compounds for the capsaicin (VR1) receptor.

5 Binding studies with [³H] Resiniferatoxin (RTX) are carried out essentially as described by Szallasi and Blumberg (1992) *J. Pharmacol. Exp. Ther.* 262:883-888. In this protocol, non-specific RTX binding is reduced by adding bovine alpha₁ acid glycoprotein (100 µg per tube) after the binding reaction has been terminated.

10 [³H] RTX (37 Ci/mmol) is synthesized by and obtained from the Chemical Synthesis and Analysis Laboratory, National Cancer Institute-Frederick Cancer Research and Development Center, Frederick, MD. [³H] RTX may also be obtained from commercial vendors (e.g., Amersham Pharmacia Biotech, Inc.; Piscataway, NJ).

The membrane homogenate of Example 4 is centrifuged as before and resuspended to a protein concentration of 333 µg/ml in homogenization buffer. Binding assay mixtures are set up on ice and contain [³H]RTX (specific activity 2200 mCi/ml), 2 µl non-radioactive test compound, 15 0.25 mg/ml bovine serum albumin (Cohn fraction V), and 5 x 10⁴ - 1 x 10⁵ VR1-transfected cells. The final volume is adjusted to 500 µl (for competition binding assays) or 1,000 µl (for saturation binding assays) with the ice-cold HEPES homogenization buffer solution (pH 7.4) described above. Non-specific binding is defined as that occurring in the presence of 1 µM non-radioactive 20 RTX (Alexis Corp.; San Diego, CA). For saturation binding, [³H]RTX is added in the concentration range of 7-1,000 pM, using 1 to 2 dilutions. Typically 11 concentration points are collected per saturation binding curve.

Competition binding assays are performed in the presence of 60 pM [³H]RTX and various concentrations of test compound. The binding reactions are initiated by transferring the assay 25 mixtures into a 37°C water bath and are terminated following a 60 minute incubation period by cooling the tubes on ice. Membrane-bound RTX is separated from free, as well as any alpha₁-acid glycoprotein-bound RTX, by filtration onto WALLAC glass fiber filters (PERKIN-ELMER, Gaithersburg, MD) which were pre-soaked with 1.0% PEI (polyethyleneimine) for 2 hours prior to use. Filters are allowed to dry overnight then counted in a WALLAC 1205 BETA PLATE counter 30 after addition of WALLAC BETA SCINT scintillation fluid.

Equilibrium binding parameters are determined by fitting the allosteric Hill equation to the measured values with the aid of the computer program FIT P (Biosoft, Ferguson, MO) as described by Szallasi, *et al.* (1993) *J. Pharmacol. Exp. Ther.* 266:678-683. Compounds provided herein generally exhibit K_i values for capsaicin receptor of less than 1 µM, 100 nM, 50 nM, 25 nM, 35 10 nM, or 1 nM in this assay.

EXAMPLE 6

Calcium Mobilization Assay

This Example illustrates representative calcium mobilization assays for use in evaluating test compounds for agonist and antagonist activity.

- 5 Cells transfected with expression plasmids (as described in Example 4) and thereby expressing human capsaicin receptor are seeded and grown to 70-90% confluency in FALCON black-walled, clear-bottomed 96-well plates (#3904, BECTON-DICKINSON, Franklin Lakes, NJ). The culture medium is emptied from the 96 well plates and FLUO-3 AM calcium sensitive dye (Molecular Probes, Eugene, OR) is added to each well (dye solution: 1 mg FLUO-3 AM, 440
10 μ L DMSO and 440 μ L 20% pluronic acid in DMSO, diluted 1:250 in Krebs-Ringer HEPES (KRH) buffer (25 mM HEPES, 5 mM KCl, 0.96 mM NaH_2PO_4 , 1 mM MgSO_4 , 2 mM CaCl_2 , 5 mM glucose, 1 mM probenecid, pH 7.4), 50 μ L diluted solution per well). Plates are covered with aluminum foil and incubated at 37°C for 1-2 hours in an environment containing 5% CO_2 . After the incubation, the dye is emptied from the plates, and the cells are washed once with KRH buffer,
15 and resuspended in KRH buffer.

DETERMINATION CAPSAICIN EC_{50}

- To measure the ability of a test compound to agonize or antagonize a calcium mobilization response in cells expressing capsaicin receptors to capsaicin or other vanilloid agonist, the EC_{50} of the agonist capsaicin is first determined. An additional 20 μ L of KRH buffer and 1 μ L DMSO is
20 added to each well of cells, prepared as described above. 100 μ L capsaicin in KRH buffer is automatically transferred by the FLIPR instrument to each well. Capsaicin-induced calcium mobilization is monitored using either FLUOROSKAN ASCENT (Labsystems; Franklin, MA) or FLIPR (fluorometric imaging plate reader system; Molecular Devices, Sunnyvale, CA) instruments. Data obtained between 30 and 60 seconds after agonist application are used to
25 generate an 8-point concentration response curve, with final capsaicin concentrations of 1 nM to 3 μ M. KALEIDAGRAPH software (Synergy Software, Reading, PA) is used to fit the data to the equation:

$$y=a*(1/(1+(b/x)^c))$$

- to determine the 50% excitatory concentration (EC_{50}) for the response. In this equation, y is the
30 maximum fluorescence signal, x is the concentration of the agonist or antagonist (in this case, capsaicin), a is the E_{max} , b corresponds to the EC_{50} value and c is the Hill coefficient.

DETERMINATION OF AGONIST ACTIVITY

Test compounds are dissolved in DMSO, diluted in KRH buffer, and immediately added to cells prepared as described above. 100 nM capsaicin (an approximate EC_{90} concentration) is

also added to cells in the same 96-well plate as a positive control. The final concentration of test compounds in the assay wells is between 0.1 nM and 5 μ M.

The ability of a test compound to act as an agonist of the capsaicin receptor is determined by measuring the fluorescence response of cells expressing capsaicin receptors elicited by the compound as function of compound concentration. This data is fit as described above to obtain the EC_{50} , which is generally less than 1 micromolar, preferably less than 100 nM, and more preferably less than 10 nM. The extent of efficacy of each test compound is also determined by calculating the response elicited by a concentration of test compound (typically 1 μ M) relative to the response elicited by 100 nM capsaicin. This value, called Percent of Signal (POS), is calculated by the following equation:

$$POS = 100 * \text{test compound response} / 100 \text{ nM capsaicin response}$$

This analysis provides quantitative assessment of both the potency and efficacy of test compounds as human capsaicin receptor agonists. Agonists of the human capsaicin receptor generally elicit detectable responses at concentrations less than 100 μ M, or preferably at concentrations less than 1 μ M, or most preferably at concentrations less than 10 nM. Extent of efficacy at human capsaicin receptor is preferably greater than 30 POS, more preferably greater than 80 POS at a concentration of 1 μ M. Certain agonists are essentially free of antagonist activity as demonstrated by the absence of detectable antagonist activity in the assay described below at compound concentrations below 4 nM, more preferably at concentrations below 10 μ M and most preferably at concentrations less than or equal to 100 μ M.

DETERMINATION OF ANTAGONIST ACTIVITY

Test compounds are dissolved in DMSO, diluted in 20 μ l KRH buffer so that the final concentration of test compounds in the assay well is between 1 μ M and 5 μ M, and added to cells prepared as described above. The 96 well plates containing prepared cells and test compounds are incubated in the dark, at room temperature for 0.5 to 6 hours. It is important that the incubation not continue beyond 6 hours. Just prior to determining the fluorescence response, 100 μ l capsaicin in KRH buffer at twice the EC_{50} concentration determined as described above is automatically added by the FLIPR instrument to each well of the 96 well plate for a final sample volume of 200 μ l and a final capsaicin concentration equal to the EC_{50} . The final concentration of test compounds in the assay wells is between 1 μ M and 5 μ M. Antagonists of the capsaicin receptor decrease this response by at least about 20%, preferably by at least about 50%, and most preferably by at least 80%, as compared to matched control (*i.e.*, cells treated with capsaicin at twice the EC_{50} concentration in the absence of test compound), at a concentration of 10 micromolar or less, preferably 1 micromolar or less. The concentration of antagonist required to provide a 50% decrease, relative to the response observed in the presence of capsaicin and without antagonist, is

the IC₅₀ for the antagonist, and is preferably below 1 micromolar, 100 nanomolar, 10 nanomolar or 1 nanomolar.

Certain preferred VR1 modulators are antagonists that are essentially free of agonist activity as demonstrated by the absence of detectable agonist activity in the assay described above
5 at compound concentrations below 4 nM, more preferably at concentrations below 10 μ M and most preferably at concentrations less than or equal to 100 μ M.

EXAMPLE 7

Microsomal *in vitro* half-life

This Example illustrates the evaluation of compound half-life values ($t_{1/2}$ values) using a
10 representative liver microsomal half-life assay.

Pooled human liver microsomes are obtained from XenoTech LLC (Kansas City, KS). Such liver microsomes may also be obtained from In Vitro Technologies (Baltimore, MD) or Tissue Transformation Technologies (Edison, NJ). Six test reactions are prepared, each containing 25 μ l microsomes, 5 μ l of a 100 μ M solution of test compound, and 399 μ l 0.1 M phosphate buffer
15 (19 mL 0.1 M NaH₂PO₄, 81 mL 0.1 M Na₂HPO₄, adjusted to pH 7.4 with H₃PO₄). A seventh reaction is prepared as a positive control containing 25 μ l microsomes, 399 μ l 0.1 M phosphate buffer, and 5 μ l of a 100 μ M solution of a compound with known metabolic properties (e.g., DIAZEPAM or CLOZAPINE). Reactions are preincubated at 39°C for 10 minutes.

CoFactor Mixture is prepared by diluting 16.2 mg NADP and 45.4 mg Glucose-6-phosphate in 4 mL 100 mM MgCl₂. Glucose-6-phosphate dehydrogenase solution is prepared by diluting 214.3 μ l glucose-6-phosphate dehydrogenase suspension (Roche Molecular Biochemicals; Indianapolis, IN) into 1285.7 μ l distilled water. 71 μ l Starting Reaction Mixture (3 mL CoFactor Mixture; 1.2 mL Glucose-6-phosphate dehydrogenase solution) is added to 5 of the 6 test reactions and to the positive control. 71 μ l 100 mM MgCl₂ is added to the sixth test reaction, which is used
25 as a negative control. At each time point (0, 1, 3, 5, and 10 minutes), 75 μ l of each reaction mix is pipetted into a well of a 96-well deep-well plate containing 75 μ l ice-cold acetonitrile. Samples are vortexed and centrifuged 10 minutes at 3500 rpm (Sorval T 6000D centrifuge, H1000B rotor). 75 μ l of supernatant from each reaction is transferred to a well of a 96-well plate containing 150 μ l of a 0.5 μ M solution of a compound with a known LCMS profile (internal standard) per well.
30 LCMS analysis of each sample is carried out and the amount of unmetabolized test compound is measured as AUC, compound concentration vs. time is plotted, and the $t_{1/2}$ value of the test compound is extrapolated.

Preferred compounds provided herein exhibit *in vitro* $t_{1/2}$ values of greater than 10 minutes and less than 4 hours, preferably between 30 minutes and 1 hour, in human liver microsomes.

EXAMPLE 8
MDCK Toxicity Assay

This Example illustrates the evaluation of compound toxicity using a Madin Darby canine kidney (MDCK) cell cytotoxicity assay.

5 1 μ L of test compound is added to each well of a clear bottom 96-well plate (PACKARD, Meriden, CT) to give final concentration of compound in the assay of 10 micromolar, 100 micromolar or 200 micromolar. Solvent without test compound is added to control wells.

MDCK cells, ATCC no. CCL-34 (American Type Culture Collection, Manassas, VA), are maintained in sterile conditions following the instructions in the ATCC production information
10 sheet. Confluent MDCK cells are trypsinized, harvested, and diluted to a concentration of 0.1×10^6 cells/ml with warm (37°C) medium (VITACELL Minimum Essential Medium Eagle, ATCC catalog # 30-2003). 100 μ L of diluted cells is added to each well, except for five standard curve control wells that contain 100 μ L of warm medium without cells. The plate is then incubated at 37°C under 95% O₂, 5% CO₂ for 2 hours with constant shaking. After incubation, 50 μ L of
15 mammalian cell lysis solution (from the PACKARD (Meriden, CT) ATP-LITE-M Luminescent ATP detection kit) is added per well, the wells are covered with PACKARD TOPSEAL stickers, and plates are shaken at approximately 700 rpm on a suitable shaker for 2 minutes.

Compounds causing toxicity will decrease ATP production, relative to untreated cells. The ATP-LITE-M Luminescent ATP detection kit is generally used according to the
20 manufacturer's instructions to measure ATP production in treated and untreated MDCK cells. PACKARD ATP LITE-M reagents are allowed to equilibrate to room temperature. Once equilibrated, the lyophilized substrate solution is reconstituted in 5.5 mL of substrate buffer solution (from kit). Lyophilized ATP standard solution is reconstituted in deionized water to give a 10 mM stock. For the five control wells, 10 μ L of serially diluted PACKARD standard is added
25 to each of the standard curve control wells to yield a final concentration in each subsequent well of 200 nM, 100 nM, 50 nM, 25 nM and 12.5 nM. PACKARD substrate solution (50 μ L) is added to all wells, which are then covered, and the plates are shaken at approximately 700 rpm on a suitable shaker for 2 minutes. A white PACKARD sticker is attached to the bottom of each plate and samples are dark adapted by wrapping plates in foil and placing in the dark for 10 minutes.
30 Luminescence is then measured at 22°C using a luminescence counter (e.g., PACKARD TOPCOUNT Microplate Scintillation and Luminescence Counter or TECAN SPECTRAFLUOR PLUS), and ATP levels calculated from the standard curve. ATP levels in cells treated with test compound(s) are compared to the levels determined for untreated cells. Cells treated with 10 μ M of a preferred test compound exhibit ATP levels that are at least 80%, preferably at least 90%, of
35 the untreated cells. When a 100 μ M concentration of the test compound is used, cells treated with

preferred test compounds exhibit ATP levels that are at least 50%, preferably at least 80%, of the ATP levels detected in untreated cells.

EXAMPLE 9

Dorsal Root Ganglion Cell Assay

5 This Example illustrates a representative dorsal root ganglion cell assay for evaluating VR1 antagonist or agonist activity of a compound.

DRG are dissected from neonatal rats, dissociated and cultured using standard methods (Aguayo and White (1992) *Brain Research* 570:61-67). After 48 hour incubation, cells are washed once and incubated for 30-60 minutes with the calcium sensitive dye Fluo 4 AM (2.5-10 ug/ml; 10 TefLabs, Austin, TX). Cells are then washed once. Addition of capsaicin to the cells results in a VR1-dependent increase in intracellular calcium levels which is monitored by a change in Fluo-4 fluorescence with a fluorometer. Data are collected for 60-180 seconds to determine the maximum fluorescent signal.

For antagonist assays, various concentrations of compound are added to the cells. 15 Fluorescent signal is then plotted as a function of compound concentration to identify the concentration required to achieve a 50% inhibition of the capsaicin-activated response, or IC_{50} . Antagonists of the capsaicin receptor preferably have an IC_{50} below 1 micromolar, 100 nanomolar, 10 nanomolar or 1 nanomolar.

For agonist assays, various concentrations of compound are added to the cells without the addition 20 of capsaicin. Compounds that are capsaicin receptor agonists result in a VR1-dependent increase in intracellular calcium levels which is monitored by a change in Fluo-4 fluorescence with a fluorometer. The EC_{50} , or concentration required to achieve 50% of the maximum signal for a capsaicin-activated response, is preferably below 1 micromolar, below 100 nanomolar or below 10 nanomolar.

25

EXAMPLE 10

Animal Models for Determining Pain Relief

This Example illustrates representative methods for assessing the degree of pain relief provided by a compound.

30 A. Pain Relief Testing

The following methods may be used to assess pain relief.

MECHANICAL ALLODYNIA

Mechanical allodynia (an abnormal response to an innocuous stimulus) is assessed essentially as described by Chaplan *et al.* (1994) *J. Neurosci. Methods* 53:55-63 and Tal and Eliav

(1998) *Pain* 64(3):511-518. A series of von Frey filaments of varying rigidity (typically 8-14 filaments in a series) are applied to the plantar surface of the hind paw with just enough force to bend the filament. The filaments are held in this position for no more than three seconds or until a positive allodynic response is displayed by the rat. A positive allodynic response consists of lifting the affected paw followed immediately by licking or shaking of the paw. The order and frequency with which the individual filaments are applied are determined by using Dixon up-down method. Testing is initiated with the middle hair of the series with subsequent filaments being applied in consecutive fashion, ascending or descending, depending on whether a negative or positive response, respectively, is obtained with the initial filament.

Compounds are effective in reversing or preventing mechanical allodynia-like symptoms if rats treated with such compounds require stimulation with a Von Frey filament of higher rigidity strength to provoke a positive allodynic response as compared to control untreated or vehicle treated rats. Alternatively, or in addition, testing of an animal in chronic pain may be done before and after compound administration. In such an assay, an effective compound results in an increase in the rigidity of the filament needed to induce a response after treatment, as compared to the filament that induces a response before treatment or in an animal that is also in chronic pain but is left untreated or is treated with vehicle. Test compounds are administered before or after onset of pain. When a test compound is administered after pain onset, testing is performed 10 minutes to three hours after administration.

MECHANICAL HYPERALGESIA

Mechanical hyperalgesia (an exaggerated response to painful stimulus) is tested essentially as described by Koch et al. (1996) *Analgesia* 2(3):157-164. Rats are placed in individual compartments of a cage with a warmed, perforated metal floor. Hind paw withdrawal duration (*i.e.*, the amount of time for which the animal holds its paw up before placing it back on the floor) is measured after a mild pinprick to the plantar surface of either hind paw.

Compounds produce a reduction in mechanical hyperalgesia if there is a statistically significant decrease in the duration of hindpaw withdrawal. Test compound may be administered before or after onset of pain. For compounds administered after pain onset, testing is performed 10 minutes to three hours after administration.

THERMAL HYPERALGESIA

Thermal hyperalgesia (an exaggerated response to noxious thermal stimulus) is measured essentially as described by Hargreaves *et al.* (1988) *Pain*. 32(1):77-88. Briefly, a constant radiant heat source is applied the animals' plantar surface of either hind paw. The time to withdrawal (*i.e.*, the amount of time that heat is applied before the animal moves its paw), otherwise described as thermal threshold or latency, determines the animal's hind paw sensitivity to heat.

Compounds produce a reduction in thermal hyperalgesia if there is a statistically significant increase in the time to hindpaw withdrawal (*i.e.*, the thermal threshold to response or latency is increased). Test compound may be administered before or after onset of pain. For compounds administered after pain onset, testing is performed 10 minutes to three hours after administration.

B. Pain Models

Pain may be induced using any of the following methods, to allow testing of analgesic efficacy of a compound. In general, compounds provided herein result in a statistically significant reduction in pain as determined by at least one of the previously described testing methods, using male SD rats and at least one of the following models.

ACUTE INFLAMMATORY PAIN MODEL

Acute inflammatory pain is induced using the carrageenan model essentially as described by Field *et al.* (1997) *Br. J. Pharmacol.* 121(8):1513-1522. 100-200 μ l of 1-2% carrageenan solution is injected into the rats' hind paw. Three to four hours following injection, the animals' sensitivity to thermal and mechanical stimuli is tested using the methods described above. A test compound (0.01 to 50 mg/kg) is administered to the animal, prior to testing, or prior to injection of carrageenan. The compound can be administered orally or through any parenteral route, or topically on the paw. Compounds that relieve pain in this model result in a statistically significant reduction in mechanical allodynia and/or thermal hyperalgesia.

CHRONIC INFLAMMATORY PAIN MODEL

Chronic inflammatory pain is induced using one of the following protocols:

1. Essentially as described by Bertorelli *et al.* (1999) *Br. J. Pharmacol.* 128(6):1252-1258, and Stein *et al.* (1998) *Pharmacol. Biochem. Behav.* 31(2):455-51, 200 μ l Complete Freund's Adjuvant (0.1 mg heat killed and dried *M. Tuberculosis*) is injected to the rats' hind paw: 100 μ l into the dorsal surface and 100 μ l into the plantar surface.
2. Essentially as described by Abbadie *et al.* (1994) *J Neurosci.* 14(10):5865-5871 rats are injected with 150 μ l of CFA (1.5 mg) in the tibio-tarsal joint.

Prior to injection with CFA in either protocol, an individual baseline sensitivity to mechanical and thermal stimulation of the animals' hind paws is obtained for each experimental animal.

Following injection of CFA, rats are tested for thermal hyperalgesia, mechanical allodynia and mechanical hyperalgesia as described above. To verify the development of symptoms, rats are tested on days 5, 6, and 7 following CFA injection. On day 7, animals are treated with a test compound, morphine or vehicle. An oral dose of morphine of 1-5 mg/kg is suitable as positive control. Typically, a dose of 0.01-50 mg/kg of test compound is used. Compounds can be

administered as a single bolus prior to testing or once or twice or three times daily, for several days prior to testing. Drugs are administered orally or through any parenteral route, or applied topically to the animal.

Results are expressed as Percent Maximum Potential Efficacy (MPE). 0% MPE is defined as analgesic effect of vehicle, 100% MPE is defined as an animal's return to pre-CFA baseline sensitivity. Compounds that relieve pain in this model result in a MPE of at least 30%.


CHRONIC NEUROPATHIC PAIN MODEL

Chronic neuropathic pain is induced using the chronic constriction injury (CCI) to the rat's sciatic nerve essentially as described by Bennett and Xie (1988) *Pain* 33:87-107. Rats are anesthetized (*e.g.* with an intraperitoneal dose of 50-65 mg/kg pentobarbital with additional doses administered as needed). The lateral aspect of each hind limb is shaved and disinfected. Using aseptic technique, an incision is made on the lateral aspect of the hind limb at the mid thigh level. The biceps femoris is bluntly dissected and the sciatic nerve is exposed. On one hind limb of each animal, four loosely tied ligatures are made around the sciatic nerve approximately 1-2 mm apart. On the other side the sciatic nerve is not ligated and is not manipulated. The muscle is closed with continuous pattern and the skin is closed with wound clips or sutures. Rats are assessed for mechanical allodynia, mechanical hyperalgesia and thermal hyperalgesia as described above.

Compounds that relieve pain in this model result in a statistically significant reduction in mechanical allodynia, mechanical hyperalgesia and/or thermal hyperalgesia when administered (0.01-50 mg/kg, orally, parenterally or topically) immediately prior to testing as a single bolus, or for several days: once or twice or three times daily prior to testing.

What is claimed is:

1. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

Ar₁ is a 5-membered aromatic heterocycle that is substituted with from 0 to 4 substituents independently chosen from R₁;

Ar₂ is phenyl or a 6-membered aromatic heterocycle, each of which is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from R₂;

W is CH or N;

X, Y and Z are independently CR_x or N, such that at least one of X, Y and Z is N;

R_x is independently chosen at each occurrence from hydrogen, C₁-C₄alkyl, amino, cyano and mono- and di-(C₁-C₄alkyl)amino;

Each R₁ is independently chosen from:

- (a) halogen, cyano and nitro;
- (b) groups of the formula -Q-M-R_y; and
- (c) groups that are taken together with an adjacent R₁ to form a fused 5- to 7-membered carbocyclic or heterocyclic ring that is substituted with from 0 to 4 substituents independently chosen from halogen, cyano, nitro and groups of the formula -Q-M-R_y;

Each Q is independently chosen from C₀-C₄alkylene;

M is independently selected at each occurrence from a single covalent bond, O, C(=O), OC(=O), C(=O)O, O-C(=O)O, S(O)_m, N(R_z), C(=O)N(R_z), C(=NH)N(R_z), N(R_z)C(=O), N(R_z)C(=NH), N(R_z)S(O)_m, S(O)_mN(R_z) and N[S(O)_mR_z]S(O)_m; wherein m is independently selected at each occurrence from 0, 1 and 2; and R_z is independently selected at each occurrence from hydrogen, C₁-C₈alkyl and groups that are taken together with R_y to form an optionally substituted 4- to 7-membered heterocycle; and

Each R_y is independently hydrogen, C₁-C₈haloalkyl, optionally substituted C₁-C₈alkyl, optionally substituted (C₃-C₈carbocycle)C₀-C₄alkyl, optionally substituted (4- to 7-membered heterocycle)C₀-C₄alkyl, or taken together with R_z to form an optionally substituted 4- to 7-membered heterocycle, wherein each alkyl, carbocycle and heterocycle is preferably substituted with from 0 to 4 substituents independently selected from hydroxy, halogen, amino, cyano, nitro, oxo, -COOH, aminocarbonyl, C₁-C₆alkyl, C₃-C₇cycloalkyl, C₂-C₆alkyl ether, C₁-C₆alkanoyl, -SO₂(C₁-C₆alkyl), -SO₂NH₂, C₁-C₈alkoxy, C₁-C₈alkylthio, mono- and di-(C₁-


C₆alkyl)aminocarbonyl, mono- and di-(C₁-C₆alkyl)amino and phenyl; such that R_y is not hydrogen if Q is C₀alkyl and M is a single covalent bond;

Each R₂ is:

- (a) independently chosen from (i) hydroxy, amino, cyano, halogen, -COOH, -SO₂NH₂, nitro and aminocarbonyl; and (ii) C₁-C₆alkyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₂-C₆alkyl ether, C₂-C₆alkanoyl, C₁-C₆alkoxycarbonyl, C₂-C₆alkanoyloxy, C₃-C₆alkanone, mono- and di-(C₁-C₆alkyl)aminoC₀-C₆alkyl, mono- and di-(C₃-C₈cycloalkyl)aminoC₀-C₄alkyl, (4- to 7-membered heterocycle)C₀-C₄alkyl, C₁-C₆alkylsulfonyl, mono- and di-(C₁-C₆alkyl)aminosulfonyl, and mono- and di-(C₁-C₆alkyl)aminocarbonyl, each of which is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, cyano, amino, -COOH and oxo; or
- (b) taken together with an adjacent R₂ to form a fused 5- to 13-membered carbocyclic or heterocyclic group that is optionally substituted, and is preferably substituted with from 0 to 3 substituents independently chosen from halogen, oxo and C₁-C₆alkyl;

R₃ is selected from:

- (i) hydrogen and halogen;
- (ii) C₁-C₆alkyl, (C₃-C₈cycloalkyl)C₀-C₂alkyl, C₁-C₆haloalkyl and phenylC₀-C₂alkyl; and
- (iii) groups of the formula:



wherein:

L is C₀-C₆alkylene or C₁-C₆alkyl that is taken together with R₅, R₆ or R₇ to form a 4- to 7-membered heterocycle;

W is O, CO, S, SO or SO₂;

R₅ and R₆ are:


- (a) independently chosen from hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₂-C₆alkanoyl, C₁-C₆alkylsulfonyl, phenylC₀-C₆alkyl, (4- to 7-membered heterocycle)C₀-C₆alkyl and groups that are joined to L to form a 4- to 7-membered heterocycle; or
 - (b) joined to form a 4- to 12-membered heterocycle; and
- R₇ is hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₂-C₆alkanoyl, phenylC₀-C₆alkyl, (4- to 7-membered heterocycle)C₀-C₆alkyl or a group that is joined to L to form a 4- to 7-membered heterocycle;

wherein each of (ii) and (iii) is optionally substituted, and is preferably substituted with from 0 to 4 substituents independently chosen from:

- (1) halogen, hydroxy, amino, cyano, $-\text{COOH}$, $-\text{SO}_2\text{NH}_2$, oxo, nitro and aminocarbonyl; and
- (2) $\text{C}_1\text{-C}_6$ alkyl, $(\text{C}_3\text{-C}_8\text{cycloalkyl})\text{C}_0\text{-C}_2$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkoxycarbonyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ alkanoyl, $\text{C}_2\text{-C}_6$ alkanoylamino, mono- and di- $(\text{C}_1\text{-C}_6\text{alkyl})\text{aminoC}_0\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_6$ alkylsulfonyl, mono- and di- $(\text{C}_1\text{-C}_6\text{alkyl})\text{aminosulfonyl}$, mono- and di- $(\text{C}_1\text{-C}_6\text{alkyl})\text{aminocarbonylC}_0\text{-C}_4$ alkyl, phenyl $\text{C}_0\text{-C}_4$ alkyl and (4- to 7-membered heterocycle) $\text{C}_0\text{-C}_4$ alkyl, each of which is substituted with from 0 to 4 secondary substituents independently chosen from halogen, hydroxy, cyano, oxo, imino, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy and $\text{C}_1\text{-C}_4$ haloalkyl; and

R_4 represents from 0 to 2 substituents that are preferably independently chosen from $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_3$ haloalkyl and oxo.

2. A compound or salt according to claim 1, wherein the compound has the formula:




wherein:

D, E and G are independently O, S, N, NR_{1a} or CR_{1a} ; and


Each R_{1a} is independently chosen from:

- hydrogen, halogen, cyano and nitro;
- groups of the formula $-\text{Q-M-R}_y$; and
- groups that are taken together with an adjacent R_{1a} to form a fused 5- to 7-membered carbocyclic or heterocyclic ring that is optionally substituted with from 1 to 4 substituents independently chosen from halogen, cyano, nitro and groups of the formula $-\text{Q-M-R}_y$.


3. A compound or salt according to claim 2, wherein the compound has the formula:



4. A compound or salt according to claim 2, wherein the compound has the formula:




5. A compound or salt according to claim 2, wherein the compound has the formula:




wherein D is O or S.

6. A compound or salt according to claim 2, wherein the compound has the formula:




wherein D is O or S.

7. A compound or salt according to claim 2, wherein the compound has the formula:



wherein E is N, O or S.

8. A compound or salt according to claim 2, wherein the compound has the formula:




wherein E is N, O or S.

9. A compound or salt according to any one of claims 4-8, wherein R_{1a} is hydrogen, halogen, amino, cyano, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylsulfonyl, or mono- or di- (C_1 - C_6 alkyl)aminosulfonyl.

10. A compound or salt according to claim 9, wherein R_{1a} is hydrogen, halogen, nitro, cyano, methyl or trifluoromethyl.

11. A compound or salt according to any one of claims 1-10, wherein R_3 is a group of the formula:



L is C_0 - C_6 alkylene or C_1 - C_6 alkyl that is taken together with R_5 or R_6 to form a 4- to 7-membered heterocycle; and

R_5 and R_6 are:

(a) independently chosen from hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, (C_3 - C_8 cycloalkyl) C_0 - C_4 alkyl, C_2 - C_4 alkanoyl and groups that are joined to L to form a 4- to 7-membered heterocycle; or

(b) joined to form a 4- to 12-membered heterocycloalkyl;

each of which alkyl, alkenyl, (cycloalkyl)alkyl, alkanoyl and heterocycloalkyl is substituted with from 0 to 4 substituents independently chosen from (i) halogen, hydroxy, amino, aminocarbonyl, oxo, $-\text{COOH}$ and $-\text{SO}_2\text{NH}_2$; and (ii) C_1 - C_4 alkyl, C_5 - C_7 cycloalkyl, C_1 - C_4 alkoxy, C_2 - C_4 alkanoyl, C_1 - C_4 haloalkyl, mono- and di- (C_1 - C_4 alkyl)amino C_0 - C_2 alkyl, mono- and di- (C_1 - C_4 alkyl)aminocarbonyl C_0 - C_2 alkyl, phenyl C_0 - C_4 alkyl and (4- to 7-membered heterocycle) C_0 - C_2 alkyl, each of which is substituted with from 0 to 4 secondary substituents independently chosen from halogen, hydroxy, cyano, C_1 - C_4 alkyl, C_1 - C_4 alkoxy and C_1 - C_4 haloalkyl.


12. A compound or salt according to claim 11, wherein R_3 is di(C_1 - C_4 alkyl)amino substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, amino, oxo, aminocarbonyl, $-\text{COOH}$, $-\text{SO}_2\text{NH}_2$, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_5 - C_7 cycloalkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_2 - C_4 alkyl ether, C_2 - C_4 alkanoyl, C_1 - C_4 alkylsulfonyl, C_2 - C_4 alkanoylamino and mono- and di- (C_1 - C_4 alkyl)amino.

13. A compound or salt according to claim 12, wherein R_3 is azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, tetrahydropyridyl or azepanyl, each of which is substituted with from 0 to 4 substituents independently chosen from:


(a) halogen, hydroxy, amino, oxo, aminocarbonyl, SO_2NH_2 and $-\text{COOH}$; and

(b) C₁-C₄alkyl, C₂-C₄alkenyl, C₅-C₇cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₂-C₄alkyl ether, C₂-C₄alkanoyl, C₁-C₄alkylsulfonyl, C₂-C₄alkanoylamino and mono- and di-(C₁-C₄alkyl)amino, each of which is substituted with from 0 to 4 secondary substituents independently chosen from hydroxy and halogen.

14. A compound or salt according to claim 13, wherein R₃ is chosen from:



15. A compound or salt according to any one of claims 1-10, wherein R₃ is a group of the formula:



wherein:

L is C₀-C₆alkylene or C₁-C₆alkyl that is taken together with R₇ to form a 4- to 7-membered heterocycle;

W is O; and

R₇ is hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, (C₃-C₈cycloalkyl)C₀-C₄alkyl, C₂-C₆alkanoyl, phenylC₀-C₆alkyl, (4- to 7-membered heterocycle)C₀-C₆alkyl or a group that is joined to L to form a 4- to 7-membered heterocycle, each of which alkyl, alkenyl, (cycloalkyl)alkyl, alkanoyl and heterocycloalkyl is substituted with from 0 to 4 substituents independently chosen from (i) halogen, hydroxy, amino, aminocarbonyl, oxo, -COOH and -SO₂NH₂; and (ii) C₁-C₄alkyl, C₅-C₇cycloalkyl, C₁-C₄alkoxy, C₂-C₄alkanoyl, C₁-C₄haloalkyl, mono- and di-(C₁-C₄alkyl)aminoC₀-C₂alkyl, mono- and di-(C₁-C₄alkyl)aminocarbonylC₀-C₂alkyl, phenylC₀-C₄alkyl and (4- to 7-membered heterocycle)C₀-C₂alkyl, each of which is substituted with from 0 to 4 secondary


substituents independently chosen from halogen, hydroxy, cyano, C₁-C₄alkyl, C₁-C₄alkoxy and C₁-C₄haloalkyl.

16. A compound or salt according to any one of claims 1-15, wherein Ar₂ is unsubstituted phenyl or unsubstituted pyridyl.

17. A compound or salt according to any one of claims 1-15, wherein Ar₂ has at least one substituent chosen from amino, cyano, halogen, -SO₂NH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄hydroxyalkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkoxy, mono- and di-(C₁-C₄alkyl)aminoC₀-C₄alkyl, mono- and di-(C₃-C₈cycloalkyl)aminoC₀-C₄alkyl, C₂-C₄alkanoyl, C₁-C₄alkoxycarbonyl, C₁-C₄alkylsulfonyl, C₁-C₄haloalkylsulfonyl and mono- and di-(C₁-C₄alkyl)aminosulfonyl.

18. A compound or salt according to claim 17, wherein Ar₂ is phenyl, pyridyl or pyrimidyl, each of which is substituted with from 1 to 3 substituents independently chosen from amino, cyano, halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, and mono- and di-(C₁-C₄alkyl)aminoC₀-C₄alkyl.

19. A compound or salt according to claim 1, wherein the compound has the formula:



wherein:

D, E and G are independently O, S, N, NR_{1a} or CR_{1a};

Each R_{1a} is independently chosen from:


(a) hydrogen, halogen, cyano and nitro; and

(b) groups of the formula -Q-M-R_y;

Ar₂ is phenyl, pyridyl or pyrimidyl, each of which is substituted with from 0 to 3 substituents independently chosen from amino, cyano, halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, and mono- and di-(C₁-C₄alkyl)aminoC₀-C₄alkyl; and

R₄ represents 0 substituents or one methyl substituent.

20. A compound or salt according to any one of claims 1-19, wherein the group:



21. A compound or salt according to any one of claims 1-20, wherein the compound is a VR1 antagonist and has an IC_{50} value of 1 micromolar or less in a capsaicin receptor calcium mobilization assay.

22. A compound or salt according to claim 21, wherein the compound exhibits no detectable agonist activity in an *in vitro* assay of capsaicin receptor agonism at a concentration of compound equal to the IC_{50} .

23. A pharmaceutical composition, comprising at least one compound or salt according to any one of claims 1-22, in combination with a physiologically acceptable carrier or excipient.

24. A pharmaceutical composition according to claim 23, wherein the composition is formulated as an injectible fluid, an aerosol, a cream, a gel, a pill, a capsule, a syrup or a transdermal patch.

25. A method for reducing calcium conductance of a cellular capsaicin receptor, comprising contacting a cell expressing a capsaicin receptor with a compound or salt according to any one of claims 1-22, and thereby reducing calcium conductance of the capsaicin receptor.

26. A method according to claim 25, wherein the cell is contacted *in vivo* in an animal.

27. A method according to claim 25, wherein the cell is a neuronal cell.

28. A method according to claim 25, wherein the cell is a urothelial cell.

29. A method according to claim 26, wherein during contact the compound is present within a body fluid of the animal.

30. A method according to claim 26, wherein the animal is a human.

31. A method according to claim 26, wherein the compound or salt is administered orally.

32. A method for inhibiting binding of vanilloid ligand to a capsaicin receptor *in vitro*, the method comprising contacting capsaicin receptor with a compound or salt according to any one of claims 1-22, in an amount sufficient to detectably inhibit vanilloid ligand binding to capsaicin receptor.

33. A method for inhibiting binding of vanilloid ligand to a capsaicin receptor in a patient, the method comprising contacting cells expressing capsaicin receptor with a compound or salt according to any one of claims 1-22, in an amount sufficient to detectably inhibit vanilloid ligand

binding to cells expressing a cloned capsaicin receptor *in vitro*, and thereby inhibiting binding of vanilloid ligand to the capsaicin receptor in the patient.

34. A method according to claim 33, wherein the patient is a human.

35. A method according to claim 33, wherein the compound is present in the blood of the patient at a concentration of 1 micromolar or less.

36. A method for treating a condition responsive to capsaicin receptor modulation in a patient, comprising administering to the patient a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby alleviating the condition in the patient.

37. A method according to claim 36, wherein the patient is suffering from (i) exposure to capsaicin, (ii) burn or irritation due to exposure to heat, (iii) burns or irritation due to exposure to light, (iv) burn, bronchoconstriction or irritation due to exposure to tear gas, infectious agents, air pollutants or pepper spray, or (v) burn or irritation due to exposure to acid.

38. A method according to claim 36, wherein the condition is asthma or chronic obstructive pulmonary disease.

39. A method for treating pain in a patient, comprising administering to a patient suffering from pain a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby alleviating pain in the patient.

40. A method according to claim 39, wherein the compound or salt is present in the blood of the patient at a concentration of 1 micromolar or less.

41. A method according to claim 39, wherein the patient is suffering from neuropathic pain.

42. A method according to claim 39, wherein the pain is associated with a condition selected from: postmastectomy pain syndrome, stump pain, phantom limb pain, oral neuropathic pain, toothache, postherpetic neuralgia, diabetic neuropathy, reflex sympathetic dystrophy, trigeminal neuralgia, osteoarthritis, rheumatoid arthritis, fibromyalgia, Guillain-Barre syndrome, meralgia paresthetica, burning-mouth syndrome, bilateral peripheral neuropathy, causalgia, neuritis, neuronitis, neuralgia, AIDS-related neuropathy, MS-related neuropathy, spinal cord injury-related pain, surgery-related pain, musculoskeletal pain, back pain, headache, migraine, angina, labor, hemorrhoids, dyspepsia, Charcot's pains, intestinal gas, menstruation, cancer, venom exposure, irritable bowel syndrome, inflammatory bowel disease and trauma.

43. A method according to claim 39, wherein the patient is a human.

44. A method for treating itch in a patient, comprising administering to a patient a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby alleviating itch in the patient.

45. A method for treating cough or hiccup in a patient, comprising administering to a patient a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby alleviating cough or hiccup in the patient.

46. A method for treating urinary incontinence or overactive bladder in a patient, comprising administering to a patient a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby alleviating urinary incontinence or overactive bladder in the patient.

47. A method promoting weight loss in an obese patient, comprising administering to a patient a therapeutically effective amount of a compound or salt according to any one of claims 1-22, and thereby promoting weight loss in the patient.

48. A compound or salt according to any one of claims 1-22, wherein the compound is radiolabeled.

49. A method for identifying an agent that binds to capsaicin receptor, comprising:

- (a) contacting capsaicin receptor with a radiolabeled compound or salt according to claim 48, under conditions that permit binding of the VR1 modulator to capsaicin receptor, thereby generating bound, labeled VR1 modulator;
- (b) detecting a signal that corresponds to the amount of bound, labeled VR1 modulator in the absence of test agent;
- (c) contacting the bound, labeled VR1 modulator with a test agent;
- (d) detecting a signal that corresponds to the amount of bound labeled VR1 modulator in the presence of test agent; and
- (e) detecting a decrease in signal detected in step (d), as compared to the signal detected in step (b), and therefrom identifying an agent that binds to capsaicin receptor.

50. A method for determining the presence or absence of capsaicin receptor in a sample, comprising the steps of:

- (a) contacting a sample with a compound according to any one of claims 1-22, under conditions that permit binding of the compound to capsaicin receptor; and
- (b) detecting a level of the compound bound to capsaicin receptor, and therefrom determining the presence or absence of capsaicin receptor in the sample.

51. A method according to claim 50, wherein the compound is radiolabeled, and wherein the step of detection comprises the steps of:

- (i) separating unbound compound from bound compound; and
- (ii) detecting the presence or absence of bound compound in the sample.

52. A packaged pharmaceutical preparation, comprising:

- (a) a pharmaceutical composition according to claim 23 in a container; and
- (b) instructions for using the composition to treat pain.

53. A packaged pharmaceutical preparation, comprising:

- (a) a pharmaceutical composition according to claim 23 in a container; and
- (b) instructions for using the composition to treat cough or hiccup.

54. A packaged pharmaceutical preparation, comprising:

- (a) a pharmaceutical composition according to claim 23 in a container; and
- (b) instructions for using the composition to treat obesity.

55. A packaged pharmaceutical preparation, comprising:

- (a) a pharmaceutical composition according to claim 23 in a container; and
- (b) instructions for using the composition to treat urinary incontinence or overactive bladder.

56. The use of a compound or salt according to any one of claims 1-22 for the manufacture of a medicament for the treatment of a condition responsive to capsaicin receptor modulation.

57. A use according to 56, wherein the condition is pain, asthma, chronic obstructive pulmonary disease, cough, hiccup, obesity, urinary incontinence, overactive bladder, exposure to capsaicin, burn or irritation due to exposure to heat, burn or irritation due to exposure to light, burn, bronchoconstriction or irritation due to exposure to tear gas, infectious agents, air pollutants or pepper spray, or burn or irritation due to exposure to acid.