

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE ⁺	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

TITLE OF THE INVENTION**GENETICALLY ENGINEERED COCCIDIOSIS VACCINE**Cross-Reference to Related Applications

5 This application is a continuation-in-part of
PCT/US89/02918, filed July 5, 1989, which is a con-
tinuation-in-part of U.S. Patent Application Serial
No. 07/215,162, filed July 5, 1988; which is a con-
tinuation-in-part of U.S. Patent Application Serial
10 No. 06/746,520, filed June 19, 1985, now abandoned,
which is a continuation-in-part of U.S. Patent Appli-
cation Serial No. 06/627,811, filed July 5, 1984, now
abandoned.

Field of the Invention

15 This invention is in the field of avian cocci-
diosis and is directed to recombinant antigenic
proteins of avian coccidia and to the genes that
encode the proteins. These antigenic proteins may be
used in a vaccine against avian coccidia.

BACKGROUND OF THE INVENTION

20 Coccidiosis is a disease of both invertebrates
and vertebrates, including man, caused by intra-
cellular parasitic protozoa which generally invade the
epithelial cells lining the alimentary tract and the
cells of associated glands. The crowded conditions
25 under which many domestic animals are raised have
contributed to increased incidence of the disease.
Virtually every domestic animal is susceptible to
infection, and distribution of the parasite is world--
wide. Coccidiosis is therefore the cause of signi-
30 ficant economic loss throughout the world.

5 The poultry industry suffers particularly severe losses, with coccidiosis being the most economically important parasitic disease of chickens. Since 1949, preventive anticoccidials have been used but have not been totally effective. Losses due to morbidity from coccidiosis, including reduced weight gains and egg production, and decreased feed conversion, persist. The cost of coccidiosis in broiler production has been estimated at 1/2 to 1 cent per pound. Based on an annual production of 3,000,000,000 broilers annually in the United States, losses would total between 60 and 120 million dollars. To this figure must be added the cost of anticoccidials estimated at 35 million dollars. These impressive figures emphasize the importance of reducing the incidence of coccidiosis in chickens.

10 Of the nine genera of coccidia known to infect birds, the genus Eimeria contains the most economically important species. Various species of Eimeria infect a wide range of hosts, including mammals, but nine species have been recognized as being pathogenic to varying degrees in chickens: Eimeria acervulina, E. mivati, E. mitis, E. praecox, E. hagani, E. necatrix, E. maxima, E. brunetti and E. tenella.

25 Although the Eimeria are highly host specific, their life cycles are similar. The developmental stages of the avian coccidia can be illustrated by the species Eimeria tenella, which proliferates in the cecum of the chicken.

30 The life cycle of the Eimeria species begins when the host ingests previously sporulated oocysts during ground feeding or by inhalation of dust. Mechanical and chemical action in the gizzard and intestinal tract of the chicken ruptures the sporulated oocyst,

liberating eight sporozoites. The sporozoites are carried in the digestive contents and infect various portions of the intestinal tract by penetration of epithelial cells. Subsequent life stages involve
5 asexual multiple fission, the infection of other epithelial cells, development of gametes, and fertilization to produce a zygote which becomes an oocyst which is passed out of the host with the droppings. The oocyst undergoes nuclear and cellular division
10 resulting in the formation of sporozoites, with sporulation being dependent upon environmental conditions. Ingestion of the sporulated oocyst by a new host transmits the disease.

Of all species of Eimeria, E. tenella has received the most attention. E. tenella is an
15 extremely pathogenic species, with death often occurring on the fifth or sixth day of infection.

Before the use of chemotherapeutic agents, poultry producers' attempts to control coccidiosis
20 were limited to various management programs. These programs were directed toward attempts at sanitation through disinfection, or by mechanical removal of litter. Despite these efforts, sufficient oocysts usually remained to transmit the disease.

25 One means of combating the hazards of coccidia is immunization. This method involves feeding to the poultry a small dose of oocysts of each of the species of coccidia during the first weeks of life. However, dosage control has proven difficult as birds ingest
30 daughter oocysts, with some birds developing severe coccidiosis and others remaining susceptible. Also, since this procedure produces mixed infections, sometimes adequate immunity does not develop to all

species given. In addition, immunity development is too slow for use with broiler production.

Another means of combating coccidia is drug treatment after the poultry is infected. One drug that has been used is sulfanilamide which has shown anticoccidial activity against six species of coccidia. However, unless the drug treatment of the flock is quickly initiated after diagnosis of the disease, medication may be started too late to be effective.

Ideally, the best method for combating coccidia is preventive medication. Since the advent of the use of sulfonamide drugs, over forty compounds have been marketed for preventive medication against coccidia. There have been many problems with the use of such drugs, including anticoccidial contamination of layer flock feeds, inclusion of excessive anticoccidial drugs in the feed causing toxicity in the birds and omission of the anticoccidial from the feed resulting in coccidiosis outbreaks. A particularly frustrating problem has been the development of drug-resistant strains of coccidia. Moreover, there is a potential for drug residues being deposited in the meat.

Clearly, available methods for the control of coccidiosis have met with limited success, and the need for a safe, efficient, and inexpensive method of combating avian coccidiosis remains.

The development of an effective anticoccidial vaccine is a desirable solution to the problem of disease prevention. Vaccines produced by traditional methods will require extensive development. There are reports of the production of attenuated strains through passage in embryos or cell culture. While this approach may eventually lead to successful

vaccines, not all the important species of Eimeria have been adapted to growth in culture or embryos such that they are capable of completing their life cycle.

5 Genetic engineering methodology presents the opportunity for an alternative approach to vaccine development. It is known that genes encoding anti-
10 genic proteins of pathogenic organisms can be cloned into microorganisms. The antigenic proteins then can be expressed at high levels, purified, and used as vaccines against the pathogenic organism. These
15 antigenic proteins have the advantage of being non-infectious and are potentially inexpensive to produce. Such "subunit vaccines" have been prepared from antigen genes for a number of viruses such as hepa-
20 titis, herpes simplex and foot and mouth disease virus. An alternate approach is to produce "synthetic vaccines", small chemically-synthesized peptides, whose sequence is chosen based upon the amino acid
25 sequence translation of viral antigen DNA. The advantages of such "synthetic vaccines" over tradi-
tional vaccination with attenuated or killed patho-
genic organisms have been summarized by Lerner in Nature 299:592-596 (1982).

It is now possible to produce foreign proteins,
25 including eukaryotic proteins, in prokaryotic organ-
isms such as gram positive or gram negative bacteria. The process involves the insertion of DNA (derived
30 either from enzymatic digestion of cellular DNA or by reverse transcription of mRNA) into an expression
vector. Such expression vectors are derived from
either plasmids or bacteriophage and contain: (1) an
origin of replication functional in a microbial host
cell; (2) genes encoding selectable markers, and (3)
regulatory sequences including a promoter, operator,

and a ribosome binding site which are functional in a microbial host cell and which direct the transcription and translation of foreign DNA inserted downstream from the regulatory sequences. To increase protein production and stability, eukaryotic proteins are often produced in prokaryotic cells as a fusion with sequences from the amino-terminus of a prokaryotic protein. β -Galactosidase or the product of one of the E. coli tryptophan operon genes have been used successfully in this manner. Expression vectors have also been developed for expression of foreign proteins in eukaryotic host cells, e.g., yeast and chinese hamster ovary tissue culture cells.

Host cells transformed with expression vectors carrying foreign genes are grown in culture under conditions known to stimulate production of the foreign protein in the particular vector. Such host cell/expression vector systems are often engineered so that expression of the foreign protein may be regulated by chemical or temperature induction. Proteins which are secreted may be isolated from the growth media, while intracellular proteins may be isolated by harvesting and lysing the cells and separating the intracellular components. In this manner, it is possible to produce comparatively large amounts of proteins that are otherwise difficult to purify from native sources.

Such microbially produced proteins may be characterized by many well-known methods, including the use of monoclonal antibodies, hereinafter referred to as "MABs," which are homogeneous antibodies that react specifically with a single antigenic determinant and display a constant affinity for that determinant, or by use of polyvalent antibodies derived from infected

birds, which react with a variety of different antigens and often with multiple determinants on a single antigen.

Alternate technology to the production of "sub-unit" or "synthetic" vaccines is the use of a fowl pox virus vector. The pox virus vaccinia has a long history of use as a vaccine and has been employed to virtually eradicate smallpox in humans. It now has been demonstrated that vaccinia virus can be effectively genetically engineered to express foreign antigens (Smith et al., Nature 302:490-495 (1983); Panicali et al., Proc. Natl. Acad. Sci. USA 80:5364-5368 (1983); Mackett et al., J. of Virology 49:857-864 (1984)) and the engineered viruses can serve as a live vaccine against other viruses and infections besides smallpox. Fowl pox virus is very similar to vaccinia virus and many of the methods developed for vaccinia for the creation of recombinants expressing foreign antigens can be applied to fowl pox. Attenuated fowl pox virus engineered to produce avian coccidia antigens thus is another method to produce an anti-coccidial vaccine. Live vaccines have the advantage of being inexpensive to produce and are characterized by the production of rapid immunity development.

A second type of live vaccine results in the presentation of antigen in the gut where coccidia normally invades. This method utilizes secretion or outer surface expression of the antigen by harmless bacteria introduced into the intestinal microbial population by incorporation in feed. Secretion is obtained by fusion of an antigen gene to the gene coding for a protein which is normally secreted, leaving the necessary secretion signal sequence intact. Outer surface expression is achieved by

5 fusion of the antigen genes to the genes that code for proteins normally localized on the outer surface. (T. Silhavy, U.S. Patent No. 4,336,336.) This type of live vaccine is especially advantageous since manufacturing costs are minimal and the immune response stimulated is of a type particularly effective against coccidia invasion of the gut.

SUMMARY OF THE INVENTION

10 This invention relates to novel recombinant antigenic proteins of avian coccidiosis, and fragments thereof containing antigenic determinants, and to the genes that encode the antigenic peptides. It has now been found that particular polypeptides present in avian cells infected with coccidiosis, when purified and isolated, contain an antigenic determinant or 15 determinants which can elicit an antibody response. This invention also relates to vaccines made using the novel antigenic proteins of avian coccidiosis and to methods of immunizing chickens against avian coccidia.

20 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the nucleotide sequence of the 5'-3' strand of cDNA encoding the E. acervulina antigen ac-1b gene.

25 Figure 2 shows the amino acid sequence of E. acervulina antigen ac-1b.

Figure 3 shows the nucleotide sequence of the 5'-3' strand of cDNA encoding the E. acervulina antigen ac-6b gene.

30 Figure 4 shows the amino acid sequence of E. acervulina antigen ac-6b.

Figure 5 shows the nucleotide sequence of the 5'-3' strand of cDNA encoding the E. tenella antigen tc-7a gene.

Figure 6 shows the amino acid sequence of E. tenella antigen tc-7a.

Figure 7 shows the nucleotide sequence of the 5'-3' strand of cDNA encoding the E. tenella clone tc-8a gene.

Figure 8 shows the amino acid sequence of E. tenella clone tc-8a.

Figure 9 shows the nucleotide sequence of the 5'-3' strand of cDNA encoding the E. tenella antigen tc-10a gene.

Figure 10 shows the amino acid sequence of E. tenella antigen tc-10a.

As is well known in the art, due to the degeneracy of the genetic code, the DNA sequences given in the Figures for the genes and antigenic peptides of this invention may be encoded by different DNA than those represented. Thus, knowledge of an amino acid sequence does not necessarily lead to a precise genetic sequence coding therefor. In all of the Figures with DNA and amino acid sequences the sequence is given as the 5' to 3' strand. The abbreviations have the following standard meanings:

A is deoxyadenyl

T is thymidyl

G is deoxyguanyl

C is deoxycytosyl

-10-

GLY is glycine
ALA is alanine
VAL is valine
LEU is leucine
5 ILE is isoleucine
SER is serine
THR is threonine
PHE is phenylalanine
TYR is tyrosine
10 TRP is tryptophan
CYS is cysteine
MET is methionine
ASP is aspartic acid
GLU is glutamic acid
15 LYS is lysine
ARG is arginine
HIS is histidine
PRO is proline
GLN is glutamine
20 ASN is asparagine.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

25 The present invention relates to recombinant antigenic proteins, and fragments thereof containing antigenic determinants, that can elicit an antibody response against avian coccidiosis, and to the cloned genes that encode the antigenic proteins and fragments. These antigenic proteins, and the fragments thereof containing antigenic determinants, will bind with a specific monoclonal antibody or with polyvalent

antibodies from infected chickens, or from other animals that have been immunized with life forms of Eimeria or Eimeria proteins, directed against an antigenic protein of avian coccidia.

5 The antigenic proteins of this invention may be used for several applications: (1) the protein(s) can be used in an avian coccidia assay to detect antibodies against the coccidia; (2) antibodies may be prepared from the antigenic protein(s); (3) the
10 protein(s) can be used for preparing vaccines against avian coccidiosis.

 Antibodies directed against coccidial-antigens are used to identify, by immunological methods, transformed cells containing DNA encoding coccidial
15 antigens. The MAb's are used as a tool for identifying cells containing DNA sequences encoding coccidial antigens that are either species specific or common to all nine species. Screening transformants with polyvalent chicken antiserum is used to identify DNA
20 sequences encoding a wide spectrum of coccidial proteins which are antigenic in chickens upon infection. DNA sequences from the transformants thus identified then may be incorporated into a micro-organism for large scale protein production. The
25 antigenic proteins, as native proteins or as hybrids with other proteins, may be used as vaccines to immunize birds to protect them from subsequent infection.

 In addition, the DNA sequences comprising the
30 genes that encode antigenic proteins and fragments thereof may be used as DNA probes. The probes have a variety of uses, including screening a DNA library for additional genes that may encode antigenic determinants.

5 The DNA probe may be labeled with a detectable group. Such detectable group can be any material having a detectable physical or chemical property. Such materials have been well-developed in the field of immunoassays and in general almost any label useful in such methods can be applied to the present invention. Particularly useful are enzymatically active groups, such as enzymes (see Clin. Chem. 22:1243 (1976)), enzyme substrates (see British Pat. Spec. 10 1,548,741), coenzymes (see U.S. Pat. Nos. 4,230,797 and 4,238,565) and enzyme inhibitors (see U.S. Pat. No. 4,134,792); fluorescers (see Clin. Chem. 25:353 (1979)); chromophores; luminescers such as chemiluminescers and bioluminescers (see Clin. Chem. 25:512 15 (1979)); specifically bindable ligands; proximal interacting pairs; and radioisotopes such as ^3H , ^{35}S , ^{32}P , ^{125}I and ^{14}C . Such labels and labeling pairs are detected on the basis of their own physical properties (e.g., fluorescers, chromophores and radioisotopes) or their reactive or binding properties (e.g., enzymes, 20 substrates, coenzymes and inhibitors). For example, a cofactor-labeled probe can be detected by adding the enzyme for which the label is a cofactor and a substrate for the enzyme. For example, one can use an enzyme which acts upon a substrate to generate a 25 product with a measurable physical property. Examples of the latter include, but are not limited to, beta-galactosidase, alkaline phosphatase and peroxidase.

30 As used herein, the term "antigenic" or "antigenic determinant" is meant immunologically cross-reactive antigenic determinants with which a given antibody will react. Therefore, the antigenic peptides of this invention will include chemically synthesized peptides, peptides made by recombinant DNA

techniques, and antibodies or fragments thereof which are anti-idiotypic towards the determinant of the peptides of this invention.

Several procedures may be used to construct a
5 microorganism that produces an antigenic protein that binds with a monoclonal or polyvalent antibody that is directed against an antigenic protein of avian coccidia. One such procedure can be divided into the following major stages, each of which is described
10 more fully herein: (1) recovery and isolation of messenger RNA (mRNA) found in organisms of the genus Eimeria; (2) in vitro synthesis of complementary DNA (cDNA), using coccidia mRNA as a template; (3) insertion of the cDNA into a suitable expression
15 vector and transformation of bacterial cells with that vector; and, (4) recovery and isolation of the cloned gene or gene fragment. This route is referred to as the mRNA route. The advantage to this route is that only "expressed" genes are cloned, reducing the number
20 of individual transformants required to represent the entire population of genes.

An alternative procedure can be divided into the following major stages which will also be described
25 more fully herein: (1) recovery and isolation of nuclear DNA found in organisms of the genus Eimeria; (2) fragmentation of the DNA and insertion into a suitable vector; (3) transformation into a suitable microbial host; (4) selection of transformants containing a gene which specifies the antigen of
30 interest; and, (5) recovery and isolation of the cloned gene or gene fragment. This route is referred to as the nuclear DNA route. The advantage to this route is that all genes are cloned, allowing the identification of genes not expressed at the time from

which mRNA is isolated. These may include genes which are expressed during stages of the life cycle which are not easy to isolate.

5 After recovery and isolation of the cloned gene that is derived from the procedures discussed above, the cloned DNA sequence is advantageously transferred to a suitable expression vector/host cell system for large scale production of the antigenic protein.

10 The DNA sequence that is to be isolated encodes an antigenic protein that will elicit an immune response when administered to chickens which will protect them from subsequent infections. It is not necessary to isolate a complete coccidial gene encoding such a protein, since those portions of the protein termed antigenic determinants are sufficient for triggering a protective immune response (Lerner, supra). This antigenic determinant should be on the surface of the folded microbially-produced protein to trigger the response (Lerner, supra).

20 In the mRNA route, the sequence may be isolated from the sporozoite life stage of the parasite. It has been demonstrated that part of the protective immune response in chickens is directed against the sporozoite. Scientists at the U.S. Department of Agriculture detected antibodies to sporozoite proteins in immune chicken serum, which also indicates that the sporozoite is a life stage that can be affected by an immune response in chickens. Antigenic proteins isolated from other life stages also may be effective as vaccines.

25 30 MAbs or polyvalent antibodies which bind to various sporozoite proteins can be used to identify cloned DNA sequences encoding those proteins. Such

-15-

proteins can be isolated and used to elicit a protective immune response in chickens.

Sporozoites can be obtained from oocysts by excystation using the method of Doran and Vetterling, Proc. Helminthol Soc. Wash. 34:59-65 (1967), and purified by the leucopak filter technique of Bontemps and Yvore, Ann. Rech. Vet. 5:109-113 (1974). Although the method of Doran and Vetterling has been found suitable for obtaining sporozoites from oocysts, any method is suitable as long as the nucleic acids within the sporozoites remain intact. Also, sporozoite mRNA may be isolated from intact sporulated oocysts, which contain the sporozoites.

mRNA Route

Isolation of mRNA coding for the antigenic proteins of interest is advantageously accomplished by lysis of intact sporulated oocysts under conditions which minimize nuclease activity. This is accomplished using a modification of the procedure described by Pasternak et al., Molec. Biochem. Parasitol. 3:133-142 (1982). Total RNA may be isolated by grinding the oocysts with glass beads in a solution containing sodium dodecyl sulfate (SDS) and proteinase K. After denaturation and degradation of oocyst proteins, the RNA is isolated by extraction of the solution with phenol and precipitation with ethanol. Oligo (dT)-cellulose chromatography then can be used to isolate mRNA from the total RNA population.

Proteins coded for by the isolated mRNA can be synthesized, in vitro, using a cell-free translation system. A number of cell-free translation systems have been devised, such as wheat germ extract (Martial

5 et al., Proc. Nat'l Acad. Sci. USA 74:1816-1820
(1977)), rabbit reticulocyte lysate (Pelham and
Jackson, Eur. J. Biochem. 67:247-256 (1976)), and
oocytes from Xenopus laevis (Sloma et al., Methods in
10 Enzymology 79:68-71 (1981)). The rabbit reticulocyte
lysate is preferred for the testing of sporozoite
mRNA. The rabbit reticulocyte lysate can be supple-
mented with a radioactively labeled amino acid, such
as [³⁵S]-methionine, so that the resulting proteins
15 contain a tracer. The various protein products may be
reacted with polyvalent chicken antisera or MABs
previously described, followed by reaction with goat
anti-chicken IgG in the case of the polyvalent anti-
bodies and Staphylococcus aureus Protein A, or in the
20 case of the MABs, just Protein A. Protein A binds any
of the mouse or goat antibodies to form an immuno-
precipitated complex. The products of the translation
and of the immunoprecipitation are visualized by gel
electrophoresis followed by fluorography. The mRNA
25 fractions found to produce proteins that react with
the antisera in this system are used for ds-cDNA
synthesis. Alternatively, to avoid missing any
antigens which are not synthesized efficiently, in
vitro, or are not immunoprecipitated efficiently,
total mRNA is used for cDNA synthesis.

30 Synthesis of cDNA employs avian myeloblastosis
virus reverse transcriptase. This enzyme catalyzes
the synthesis of a single strand of DNA from deoxy-
nucleoside tri-phosphates on the mRNA template.
(Kacian and Myers, Proc. Nat'l Acad. Sci. USA
73:2191-2195 (1976).) The poly r(A) tail of mRNA
permits oligo(dT) (of about 12-18 nucleotides) to be
used as a primer for cDNA synthesis. The use of a
radioactively labeled deoxynucleoside triphosphate

-17-

facilitates monitoring of the synthesis reaction. Generally, a ^{32}P -containing deoxynucleoside triphosphate, such as $[\alpha\text{-}^{32}\text{P}]\text{dCTP}$, may be used advantageously for this purpose. The cDNA synthesis is generally conducted by incubating a solution of the mRNA, the deoxynucleoside triphosphates, the oligo(dT)12-18 and reverse transcriptase for 10 minutes at 46°C . The solution also preferably contains small amounts of actinomycin D and dithiothreitol to promote full length synthesis. (Kacian and Myers, supra.) After incubation, ethylenediaminetetraacetic acid (EDTA) is added to the solution, and the solution is extracted with phenol:chloroform. The aqueous phase is advantageously purified by gel filtration chromatography, and the cDNA-mRNA complex in the eluate is precipitated with alcohol.

The mRNA can be selectively hydrolyzed in the presence of the cDNA with dilute sodium hydroxide at an elevated temperature. Neutralization of the alkaline solution and alcohol precipitation yields a single-stranded cDNA copy.

The single-stranded cDNA copy has been shown to have a 5' poly (dT) tail, and to have a 3' terminal hairpin structure, which provides a short segment of duplex DNA. (Efstratiadis et al., Cell 7:279-288 (1976)). This 3' hairpin structure can act as a primer for the synthesis of a second DNA strand. Synthesis of this second strand is conducted under essentially the same conditions as the synthesis of the cDNA copy, except that the Klenow fragment of E. coli DNA polymerase I (Klenow et al., Eur. J. Biochem. 22:371-381 (1971)) is substituted for reverse transcriptase. The duplex cDNA recovered by this procedure has a 3' loop, resulting from the 3' hairpin

structure of the single-stranded cDNA copy. This 3' loop can be cleaved by digestion with the enzyme, S1 nuclease, using essentially the procedure of Ullrich et al., Science 196:1313-1319 (1977). The S1 nuclease digest may be extracted with phenol-chloroform, and the resulting ds-cDNA precipitated from the aqueous phase with alcohol.

For purposes of amplification and selection, the ds-cDNA prepared as described above is generally inserted into a suitable cloning vector, which is used for transforming appropriate host cells. Suitable cloning vectors include various plasmids and phages, but a bacteriophage lambda is preferred.

For a cloning vector to be useful for the expression of foreign proteins which are to be detected with antibodies, it should have several useful properties. Most importantly, it should have a cloning site within a gene which is expressed in the host being used. There should also be a means of controlling expression of the gene. The vector should be able to accept DNA of the size required for synthesis of the desired protein product and replicate normally. It is also useful to have a selectable property which allows identification of vectors carrying inserts. A cloning vector having such properties is the bacteriophage λ gt11 (ATCC 37194) (Young and Davis, Proc. Nat'l Acad. Sci. USA 80:1194-1198 (1983)). This vector has a unique EcoRI site near the end of the bacterial gene coding for β -galactosidase. That site can be used for insertion of foreign DNA to make hybrid proteins made up of β -galactosidase and the foreign gene product. The expression of β -galactosidase is under control of the lac promoter and can be induced by the addition of isopropyl- β -D-thiogalactopyranoside (IPTG). The λ gt11

-19-

phage contains 43.7 kb of DNA which is considerably smaller than wild type λ . This allows insertion of pieces of DNA up to 8.3 kb in length, before the DNA becomes too large to fit inside the phage head.

5 Because DNA is inserted into the gene for β -galactosidase, transformants having inserts can easily be distinguished from those which do not by looking for β -galactosidase activity. An indicator dye, 5-bromo-4-chloro-3-indolyl- β -D-galactoside (Xgal), can be

10 incorporated with agar plates. β -galactosidase cleaves this molecule to give a blue product, thus allowing examination of the cultures for the presence of active β -galactosidase. Those plaques having inserts are colorless on X-gal plates because the

15 insertion of foreign DNA into the β -galactosidase gene has eliminated its activity.

The ds-cDNA can be conveniently inserted into the phage by addition of EcoRI linkers to the DNA and ligation into the EcoRI-cut λ gt11 DNA. After ligation

20 of the cDNA into the phage DNA, the DNA is packaged, in vitro, into λ phage heads (Enquist and Sternberg, Methods in Enzymology 68:281-298 (1979) and those phages are used to infect a suitable λ -sensitive host. With the proper choice of host, the phage may be

25 screened as plaques or lysogens (colonies).

Aside from the E. coli/bacteriophage λ gt11 system described, many other host/vector combinations have been used successfully for the cloning of foreign genes in E. coli (Principles of Gene Manipulation, 2nd

30 Ed., Old and Primrose, Univ. of California Press, 32-35, 46-47 (1981)) including "open reading frame" vectors, described in detail later.

The foregoing discussion has focused on cloning procedures in gram negative bacteria, e.g., E. coli.

Alternatively, foreign genes may be cloned into plasmid vectors that will transform and replicate in a gram positive bacterium such as Bacillus subtilis (Old and Primrose, supra, pp. 51-53) or in a eukaryotic host cell such as yeast (Old and Primrose, supra, pp. 62-68) filamentous fungi, insect cells (U.S. 4,745,051 and 4,879,236) and mammalian cells. Cloning vectors have been constructed which transform both yeast and E. coli. Such vectors are termed "shuttle vectors" and may be transferred, along with the cDNA they carry, between the two host microorganisms (Storms, et al., Journal of Bacteriology 140:73-82 (1979); and Blanc et al., Molec. Gen. Genet. 176:335-342 (1979). Shuttle vectors also exist which replicate in (and may carry cloned genes into) both E. coli and B. subtilis (Old and Primrose, supra, at p. 53). Vectors derived from the other bacteriophages such as M13 have also proven useful in the cloning of foreign genes (Old and Primrose, supra, Chap. 5). Any of these techniques can be employed, if desired, in the constructions of the present invention.

The DNA described herein may be inserted into the above vectors by various techniques including homopolymeric tailing, blunt-end ligation or by use of linker molecules (Old and Primrose, supra, at p. 92).

Many immunological methods for screening clone banks for those expressing a desired protein are known and include procedures described by Engvall and Pearlman, Immunochemistry 8:871-874 (1971); Koenen et al., The European Molecular Biology Organization Journal, Vol. 1, No. 4, pp. 509-512 (1982); Broome et al., Proc. Natl. Acad. Sci., USA 75:2746-2749 (1978); Villa-Komaroff et al., Proc. Natl. Acad. Sci., USA 75:3727-3731 (1978); Anderson et al., Methods in

Enzymology 68:428-436 (1979); Clarke et al., Methods in Enzymology 68:436-442 (1979); Ehrlich et al., Methods in Enzymology 68:443-453 (1979); Kemp et al., Proc. Natl. Acad. Sci., USA 78:4520-4524 (1981).

5 By the cloning procedures outlined, thousands of recombinant bacteriophage are generated. In order to screen them for production of coccidial antigens, two antibody screens can be utilized. Both screening methods depend upon expression of the coccidial
10 antigenic protein either alone or as a fusion protein with a bacterial gene. In the examples included herein, the coccidial antigens are produced as fusions with E. coli β -galactosidase. The screening methods, therefore, depend on expression of the fusion product
15 and detection of the product by reaction with antibodies, either monoclonal or polyvalent, directed against that antigen.

The recombinant bacteriophages can be used to infect a suitable E. coli host which allows the
20 formation of phage plaques on agar (or agarose) plates. The plaques can be transferred to nitrocellulose membranes while being induced with IPTG. The proper antibodies are then reacted with the filters. After reaction of the primary antibodies
25 with the filters, the positive reactions are detected by reaction with either [¹²⁵I] Staphylococcus aureus Protein A or a second antibody conjugated with horse-radish peroxidase.

30 Alternatively, the recombinant bacteriophages can be used to infect an E. coli host in which lysogens are produced at a high frequency. In this case, the transformants can be screened as colonies. The colonies are grown on a cellulose acetate filter under non-induced conditions. After the colonies have

reached a suitable size, the cellulose acetate filter is placed over a nitrocellulose filter which is on an agar plate containing IPTG. The colonies are incubated at elevated temperatures to induce phage production, while expression of the β -galactosidase gene is induced by inclusion of IPTG. After a suitable incubation period, during which some lysis of the colonies occurs with release of proteins through the cellulose acetate filter onto the nitrocellulose filter, the cellulose acetate filter is removed. The nitrocellulose filter is processed as described above for screening of plaques.

The phages giving positive signals in the antibody-screening procedure can be shown to contain sequences coding for coccidial proteins by excision of the DNA originally inserted into the phage DNA and examination of the ability of that DNA to hybridize with coccidia mRNA or coccidia genomic DNA. The nucleotide sequence of the cDNA insert is determined using the methods of Sanger et al., Proc. Natl. Acad. Sci., USA 74:5463-5467 (1977); or Maxam and Gilbert, Proc. Natl. Acad. Sci., USA 74:560-564 (1977).

Nuclear DNA Route

Another method of cloning coccidial antigens begins with isolation of nuclear DNA from oocysts. This DNA is then broken into fragments of a size suitable for insertion into a cloning vector. To obtain such fragments, one can use mechanical shearing methods such as sonication or high-speed stirring in a blender to produce random breaks in the DNA. Intense sonication with ultrasound can reduce the fragment length to about 300 nucleotide pairs. (Old

and Primrose, supra, p. 20.) Alternatively, nuclear DNA may be partially digested with DNaseI, which gives random fragments, with restriction endonucleases, which cut at specific sites, or with mung bean nuclease in the presence of formamide, which has been shown with some related organisms (McCutchan, T.F., et al. Science 225:625-628 (1984)) to produce DNA fragments containing intact genes.

These nuclear DNA fragments may be inserted into any of the cloning vectors listed for the cloning of cDNA in the mRNA experimental method. If the nuclear DNA is digested with a restriction endonuclease, it can be inserted conveniently into a cloning vector digested with the same enzyme, provided the vector has only one recognition site for that enzyme. Otherwise, DNA fragments may be inserted into appropriate cloning vectors by homopolymeric tailing or by using linker molecules (Old and Primrose, supra, at p. 92).

Advantageously, the nuclear DNA fragments are cloned into "open reading frame" vectors which are designed to simultaneously clone and express foreign genes or fragments thereof. Several such vectors are known in the art, including those described by Weinstock et al., Proc. Natl. Acad. Sci., USA 80:4432-4436 (1983); Keonen et al., The European Molecular Biology Organization Journal 1, 4, pp. 509-512 (1982); Ruther et al., 79:6852-6855 (1982); Young and Davis, supra; and Gray et al., Proc. Natl. Acad. Sci., USA 79:6598-6602 (1982).

Open reading frame (ORF) vectors have been used to clone both prokaryotic and eukaryotic genomic DNA or cDNA. These vectors generally contain a bacterial promoter operably linked to an amino terminal fragment of a prokaryotic gene. A carboxy terminal fragment of

a gene which encodes a product for which an assay is known (e.g., the E. coli lacZ gene which encodes β -galactosidase) is located downstream. The sequences between the amino terminal gene fragment and the lacZ fragment include restriction endonuclease recognition sites useful for insertion of foreign genes and, in some cases, also place the lacZ fragment out of reading frame for translation with respect to the amino terminal gene fragment. When foreign DNA is inserted into these vectors (by blunt end ligation, homopolymeric tailing, ligation of cohesive termini, or the use of linkers), a certain percentage of recombinants will have received foreign DNA of a length that puts the lacZ gene in phase with the reading frame set by the amino terminal gene fragment. The result is production of a "tribrid" protein comprising the polypeptides encoded by the amino terminal gene fragment, the cloned DNA, and the lacZ gene. Such recombinants are identified on MacConkey agar plates or on agar plates containing "Xgal" (5-bromo-4-chloro-3-indolyl- β -D galactoside) because the β -galactosidase activity of the tribrid protein cleaves the dye in such plates, turning colonies red (MacConkey agar) or blue (Xgal). β -galactosidase can carry a wide range of protein sequences at its amino terminus and still retain biological activity. Alternatively, the insert may be inserted to inactivate a gene by interrupting the sequence. The insert may be in the correct reading frame to produce a hybrid gene consisting of the amino-terminus of the bacterial gene and sequences from the insert gene at the carboxy terminus.

Only recombinants receiving exons (i.e., coding sequences of genes, which have no stop codons) which

are in-frame with respect to the amino terminal gene fragment are detected by this method. ORF vectors are useful for cloning genes for which no DNA or protein sequence data exists, if antibodies against the gene product exist. Screening of the clone bank may be accomplished by immunological methods which make RNA or DNA hybridization probes unnecessary. The immunological screening methods mentioned for the mRNA route can be used.

Plasmid DNA is isolated from transformants found to be "positive" by the above screening methods. The nuclear DNA inserts of these plasmids are then subjected to DNA sequencing. Once the nucleotide sequence is known, it is possible by known methods to chemically synthesize all or part of the cloned coccidial genes. The synthesis of fragments of the cloned genes, followed by insertion of the gene fragments into expression vectors as described below and reaction of the polypeptides produced with MABs allows detection of those portions of the gene which are antigenic determinants.

Once a cloned DNA sequence is identified as encoding a protein that binds antibodies directed against coccidial proteins, it may be transferred to expression vectors engineered for high-level production of the desired antigenic protein. The expression vectors are transformed into suitable host cells for production of the antigenic protein. These host cells may include both prokaryotic and eukaryotic organisms. The prokaryotic host cells include E. coli and B. subtilis. The eukaryotic host cells include yeast, insect cells, and mammalian cells.

Coccidial antigens advantageously may be produced at high levels in E. coli as a fusion protein compris-

-26-

ing the antigen and an amino terminal portion of the β -subunit of the enzyme tryptophan synthetase (the product of the E. coli trpB gene). This fusion is accomplished by inserting a DNA sequence encoding a coccidial antigen into a plasmid vector carrying the trpB gene.

The expression vector used may be one in which expression of the fusion antigenic protein is highly regulatable, e.g., by chemical induction or temperature changes. An expression vector with such regulatory capability is the plasmid pGX2606, which contains a hybrid $\lambda O_L P_R$ regulatory region as described in copending application serial number 534,982 filed September 23, 1983. Host expression vector systems in which expression of foreign proteins is regulatable have the advantage of avoiding possible adverse effects of foreign protein accumulation as high cell densities are reached. Some investigators have proposed that expression of gene fragments such as those encoding antigenic determinants may avoid the deleterious effects that expression of the entire antigenic protein would have on E. coli host cells. (Helfman et al., Proc. Natl. Acad. Sci., USA 80:31-35 (1983)).

Coccidial antigens also may be produced in high levels as fusions at the carboxy-terminal of E. coli β -galactosidase, as they are directly obtained by use of the cloning vector λ gt11. The fused β -galactosidase-coccidia antigen gene is transferred with all of the associated regulatory elements to a small plasmid, where synthesis of the gene product is regulated by the lac promoter, which is transferred along with the fusion gene from the phage to the plasmid. Such a small plasmid is PGX1066 (plasmid

pGX1066 is present in E. coli strain GX1186, ATCC 39955) which carries the gene for ampicillin resistance and has a bank of restriction sites which are useful for insertion of DNA fragments. Synthesis of the fusion protein is induced by addition of IPTG, the inducer of the lac operon.

An effective subunit vaccine against avian coccidiosis may consist of a mixture of antigen proteins derived from several species of Eimeria. Alternatively, production costs may be decreased by producing two or more antigen proteins as one fusion protein thus reducing the required number of fermentations and purifications. Such a fusion protein would contain the amino acid sequence comprising an antigenic epitope of each antigen protein (or repetitions of those sequences) with variable amounts of surrounding nonantigenic sequence. A hybrid gene designed to code for such a protein in E. coli would contain bacterial regulatory sequence (promoter/operator) and the 5' end of an E. coli gene (the ribosome binding site and codons for several amino acids) to ensure efficient translation initiation followed by the coding sequences for the antigenic epitopes all fused in the same reading frame.

E. coli cells transformed with the expression vector carrying a cloned coccidial antigen sequence are grown under conditions that promote expression of the antigenic polypeptide. The antigenic protein is then purified from the cells and tested for ability to elicit an immune response in chickens that will protect them from subsequent Eimeria infections. The purified protein may be used to immunize the birds. The purified protein may be combined with suitable carriers and adjuvants and administered to birds in

-28-

their feed or by injection. Alternatively, live microorganisms containing the DNA sequences encoding the coccidial antigens may be fed to chickens. Such microorganism are advantageously those which normally inhabit the avian intestinal tract, such as E. coli or coryneform bacteria.

In a preferred system, the microorganisms are transformed with an expression vector in which the sequences encoding the coccidial antigen are fused in frame to a gene or gene fragment encoding a host cell outer membrane protein or secreted protein, such as the E. coli lamB protein, the λ receptor. The antigenic protein is therefore continuously presented in the host at the location of infection by the parasites. It is known that foreign proteins fused in expression vectors to outer membrane or secreted proteins have been presented at the cell surface or secreted from their host cells. (Weinstock, supra, and Silhavy, U.S. Patent No. 4,336,336 which is herein incorporated by reference.)

In another preferred system for development of live vaccines, an attenuated fowl pox virus expression vector is utilized. Fowl pox has the capacity to accommodate several coccidia genes allowing the production of multivalent vaccines. Currently, attenuated fowl pox virus is utilized as a vaccine to protect commercial flocks against fowl pox infection. Virus preparation and treatment of birds with fowl pox virus genetically engineered to produce coccidia antigens is the same as the conventional methods of pox vaccine use currently practiced.

Pox viruses are among the most complex viruses known with very high molecular weight double-stranded DNA genomes. With the most studied pox virus,

vaccinia, it has been demonstrated that the pox genome can easily accommodate inserts of foreign DNA capable of coding for foreign antigenic proteins (Smith et al., supra; Panicali et al., supra; Mackett et al., supra). When a foreign gene is incorporated into the pox virus genome under the control of a pox promoter regulatory sequence, the foreign antigen is expressed upon infection in the cytoplasm of the cell where the pox virus replicates. Successful insertion and expression of coccidia antigen genes within the fowl pox genome is dependent upon identifying a nonessential region of the pox DNA for antigen gene insertion and ensuring an active pox promoter is situated 5' of the desired coccidia gene.

Insertion of DNA into the pox genome is accomplished by in vivo recombination. Pox DNA is not infectious presumably because its cytoplasmic location requires the presence of pox virus specific RNA and DNA polymerases that are normally carried into the cell by the virion. DNA sequence information from vaccinia virus (Weir and Moss, J. of Virology 46:530-537 (1983); Venkatesan et al., Cell 125:805-813 (1981)) demonstrates sequence patterns in regulatory regions that are likely to be unique to vaccinia genes and thus not recognized by cellular enzymes. Because the pox DNA is not infectious, foreign DNA insertion into the fowl pox genome is accomplished by in vivo recombination as has been demonstrated with vaccinia to occur at high frequency (Weir et al., Proc. Natl. Acad. Sci., USA 79:1210-1214 (1982)). A fowl pox virus infection of chick embryo fibroblasts is followed by transfection using the CaCl₂ precipitation technique (Graham et al., Virology 52:456-457 (1973); Stow et al., J. Virology 28:6182-192 (1978)) with

5 plasmid DNA that includes the coccidia antigen gene placed under the control of a promoter functional in fowl pox, and DNA sequence homology with fowl pox. During the course of the infection recombination occurs. If a coccidia DNA sequence is inserted within the fowl pox homologous sequence on the transfected plasmid, upon recombination the coccidia DNA sequence is, in some cases, inserted into the pox virus genome. The infected cells and virus from a recombination attempt are harvested and fresh chick embryo fibroblast cells grown as a monolayer in tissue culture are infected at a low multiplicity such that individual plaques resulting from an initial single virus infection can be identified using conventional techniques. 10 Desired recombinant viruses are identified using an in situ hybridization technique (Villarreal and Berg, Science 196:183-185 (1977)) using radioactive coccidia DNA sequence as probe. Alternatively, viral DNA immobilized on nitrocellulose paper prepared from cells infected by plaque purified virus or cells infected with pools of potential recombinant viruses can be used for identification of desired recombinant viruses. Immunological screening of fixed cells (Gremer et al., Science 228:737-740 (1985)) is an alternative to hybridization. 15 20 25

30 The region of fowl pox DNA included in the plasmid vector must be from a nonessential region, and is chosen by randomly testing segments of fowl pox DNA for regions that allow recombinant formation without seriously affecting virus viability using the method described above. Fowl pox DNA is purified (Muller et al., J. Gen. Virology 38:135 (1977); Gafford et al., Virology 89:229 (1978)), randomly sheared to about 3 kilobases and cloned into a small bacterial plasmid,

such as pGx1066, creating several different isolates. Foreign DNA must be inserted into the fowl pox portion of the plasmids before testing the effect of recombination upon virus viability. To accomplish this, E. coli transposon insertions such as $\lambda\delta$ (Guyer, Methods in Enzymology 101:362-369 (1983)) can be readily placed within the fowl pox portion of the plasmid. Cotransfections that result in viable fowl pox recombinants containing $\lambda\delta$ sequence identify desirable nonessential fowl pox DNA for use in cotransfection plasmids.

Fowl pox DNA regions with partial sequence homology to the thymidine kinase gene of vaccinia identified by hybridization experiments are also useful for inclusion in the cotransfection plasmid since the thymidine kinase gene of vaccinia has been shown to be nonessential (Weir (1982), supra; Mackett et al., Proc. Natl. Acad. Sci., USA 79:4927-4931 (1982); Hrubby and Ball, J. Virology 43:403-408 (1982)).

Placement of the coccidia antigen gene under the control of a fowl pox promoter is carried out by conventional in vitro manipulation of the plasmid before concurrent transfection and fowl pox infection. Promoter sequences useful for driving expression of the coccidia antigens could be identified by determination of the DNA sequence located 5' to fowl pox genes. Promotor sequences are then synthesized chemically and included in the plasmid vector adjacent to endonuclease cloning sites within the fowl pox homologous region of the plasmid. Putative promoter sequences identified through DNA sequencing of vaccinia DNA (Venkatesan et al. (1981), supra; Weir and Moss (1983), supra) are also chemically synthe-

sized and compared with fowl pox promoters for optimal effect. Putative fowl pox promoters are verified by cloning them 5' of a test gene with an easily measured translation product such as chloramphenicol acetyltransferase (Gorman et al., Molecular and Cellular Biology 2:1044-1057 (1982)) in a bacterial plasmid. The plasmid is used to transfect fowl pox infected tissue culture cells and the cells are assayed for transient expression of the test gene.

Vaccinia virus has a broad host range and does infect chickens. Thus the vectors and methods already developed for vaccinia could be utilized to develop vaccines for avian coccidia and coccidiosis in any other genus included in the vaccinia host range. This approach requires caution since vaccinia is severely pathogenic to a small proportion of the human population.

A good alternative to pox vectors would be to utilize a herpes virus such as Marek's Disease virus or Herpes virus of turkeys. Attenuated forms of both viruses are currently used as live vaccines to prevent Marek's disease in poultry. Similar to pox viruses, herpes viruses have large double stranded DNA genomes and are good candidates for genetic engineering using in vivo recombination methods similar to those developed for vaccinia. The advantage of engineering Marek's disease virus to also provide protection against coccidia infection is that coccidia protection is provided at no additional production cost above the Marek's Disease Vaccine that is already generally in use.

The production of coccidia antigen by fowl pox recombinants is verified by immunological analysis of the protein produced in chick embryo fibroblast tissue

culture cells after infection and also by testing the circulating antibody of birds infected with recombinant fowl pox virus for cross reaction with whole coccidia or protein isolated from coccidia of the appropriate species.

5 The cloned antigenic proteins used in vaccines above are tested for their ability to elicit an immune response in chickens that protects the birds from subsequent infection by any of the important species of Eimeria, including E. tenella, E. acervulina, E. brunetti, E. mivati, E. maxima, E. praecox, E. mitis, and E. necatrix. The cloning procedures described above may be repeated until DNA sequences encoding coccidial antigens that collectively protect chickens against coccidiosis are isolated and used as a vaccine by the methods above.

10 In addition to cloned antigenic proteins which may be useful as vaccines to protect against coccidiosis, another useful alternative which may be derived from cloning antigen genes is the use of small, synthetic peptides in vaccines (see Lerner, supra). After the sequence of antigenic proteins is determined, it is possible to make synthetic peptides based on that sequence. The peptides are conjugated to a carrier protein such as hemocyanin or serum albumin and the conjugate then can be used to immunize against coccidia.

20 It is contemplated that the procedures described may also be used to isolate antigenic proteins from other coccidia species that can be used in vaccines to protect other domestic animals from coccidiosis.

25 The following examples are supplied in order to illustrate, but not limit, the present invention.

Example 1

Identification of an Eimeria acervulina
cDNA Clone Encoding Antigen ac-1b
with a Monoclonal Antibody

5 In order to identify antigens of Eimeria acervulina, expression libraries were prepared in lambda vector, λ gt11, using cDNA prepared from polyA mRNA isolated from E. acervulina oocysts. The methods used for construction of the libraries is described in
10 Genex Patent Application, PCT/US89/02918, incorporated herein by reference. The cDNA expression library was screened with monoclonal antibody (MCA) 12-09 which was raised against the sporozoite stage of E. acervulina. The MCA was obtained from Dr. Harry Danforth,
15 U.S. Department of Agriculture, Beltsville, Maryland.

 The library to be screened was plated on a host that allows lysis and plaque formation. During induction of the antigens encoded by the phage, the plaques were transferred to nitrocellulose filters.
20 Phage that produce antigens cross-reactive with MCA 12-09 were identified by screening the filters with MCA 12-09. The cDNA inserts from the MCA 12-09 positive phage were cloned into bacteriophage M13 and subjected to sequence analysis. Following sequence
25 analysis, E. acervulina antigen ac-1b was identified. The complete sequence of ac-1b and its 25.2 Kd translation product are given in Figures 1 and 2.

 Antigens ac-1b was expressed in E. coli after insertion into the plasmid expression vector pEX-32b.
30 Antigen ac-1b is encoded in expression vector pGX5361 and is expressed as a fusion protein with 11 Kd of the MS-2 polymerase under control of the P_L promoter. The host strain pop2136 contains a temperature sensitive

repressor of P_l and expression of the fusion protein is fully induced at 42°C.

Example 2

Identification of an Eimeria acervulina cDNA Clone Encoding Antigen ac-6b with a Monoclonal Antibody

Using the techniques described in Example 1 E. acervulina libraries were screened with MCA 12-09. Phage that produce antigens crossreactive with the monoclonal were identified. The cDNA inserts from the MCA 12-09 positive phage were cloned into bacteriophage M13 and subjected to sequence analysis. Following sequence analysis, E. acervulina antigen ac-6b was identified.

Antigen ac-6b showed sequence homology to the carboxy terminal sequence of E. tenella antigen GX3262, which has been shown to be protective against Eimeria infections in chickens (Miller et al., Infect. Immun. 57:2014-2020 (1989), Danforth et al., Poultry Sci. 68:1643-1652 (1989). In order to recover the entire ac-6b gene, an E. acervulina cDNA expression library was screened with an 18 bp oligonucleotide corresponding to the extreme 5'-sequence of the original ac-6b sequence. A plaque that hybridized to the oligonucleotide was identified and shown by sequence analysis to contain the complete ac-6b gene which encodes a 21.6 Kd coccidial peptide. The sequence of ac-6b and its translation product is given in Figures 3 and 4.

Antigen ac-6b was expressed in E. coli after insertion into the plasmid expression vector pEX-32b. Antigen ac-6b is encoded in expression vector pGX5368 and is expressed as a fusion protein with 11 Kd of the

MS-2 polymerase under control of the P_L promoter. The host strain pop2136 contains a temperature sensitive repressor of P_L and expression of the fusion protein is fully induced at 42°C.

5

Example 3

Identification of an Eimeria tenella cDNA Clone
Encoding Antigen tc-7a with a Monoclonal Antibody

10

In order to identify antigens of Eimeria tenella, expression libraries were prepared in the lambda vector, λgt11, using cDNA prepared from polyA mRNA isolated from E. tenella oocysts. The cDNA expression library was screened with monoclonal antibody (MCA) 12-07 which was raised against the sporozoite stage of E. tenella. The MCA was obtained from Dr. Harry Danforth, U.S. Department of Agriculture, Beltsville, Maryland.

15

20

The library to be screened was plated on a host that allows lysis and plaque formation. During induction of the antigens encoded by the phage, the plaques were transfer to nitrocellulose filters. Phage that produce antigens cross-reactive with MCA 12-07 were identified by screening the filters with MCA 12-07. The cDNA inserts from the MCA 12-07 positive phage were cloned into bacteriophage M13 and subjected to sequence analysis. Following sequence analysis, E. tenella antigen tc-7a was identified. Antigen tc-7a consists of an open reading frame of 540 bp that encodes the carboxy terminal fragment of the E. tenella protein. The DNA sequence of tc-7a and its 16.6 Kd translation product are shown in Figures 5 and 6.

25

30

Antigen tc-7a was expressed in E. coli after insertion into the plasmid expression vector pEX-32b.

Antigen tc-7a is encoded in expression vector pGX5391 and is expressed as a fusion protein with 11 Kd of the MS-2 polymerase under control of the P_L promoter. The host strain pop2136 contains a temperature sensitive repressor of P_L and expression of the fusion protein is fully induced at 42°C.

Example 4

Identification of an Eimeria tenella cDNA Clone Encoding Antigen tc-8a with a Monoclonal Antibody

As described in Example 3, E. tenella libraries were screened with MCA 12-07. Phage that produce antigens crossreactive with the monoclonal were identified. The cDNA inserts from the MCA 12-07 positive phage were cloned into bacteriophage M13 and subjected to sequence analysis. Following sequence analysis E. tenella antigen tc-8a was identified.

Antigens tc-8a is encoded on a short DNA fragments of 117 bp. The fragment consists of an open reading frame covering its entire length. The DNA sequence of the fragment and its 3.5 Kd translation product is shown in Figures 7 and 8.

Example 5

Identification of an Eimeria tenella cDNA Clone Encoding Antigen tc-10a with a Monoclonal Antibody

As described in Example 3, E. tenella libraries were screened with MCA 12-07. Phage that produce antigens cross-reactive with the monoclonal were identified. The cDNA inserts from the MCA 12-07 positive phage were cloned into bacteriophage M13 and subjected to sequence analysis. Following sequence analysis, E. tenella antigen tc-10a was identified.

Antigens tc-10a is encoded on a short DNA fragments of 228 bp. The fragment consists of an open reading frame covering its entire length. The DNA sequence of the fragment and its 7.8 Kd translation product is shown in Figures 9 and 10.

5

While certain representative embodiments and details have been shown for the purpose of illustrating the invention, it will be apparent to those skilled in the art that various changes and modifications may be made therein without departing from the scope of the invention.

10

What Is Claimed Is:

1. A cloned DNA molecule comprising the nucleotide sequence as shown in Figure 1.

2. An antigenic protein molecule comprising the amino acid sequence as shown in Figure 2.

3. A cloned DNA molecule comprising the nucleotide sequence as shown in Figure 3.

4. An antigenic protein comprising the amino acid sequence as shown in Figure 4.

5. A cloned DNA molecule comprising the nucleotide sequence as shown in Figure 5.

6. An antigenic protein comprising the amino acid sequence as shown in Figure 6.

7. A cloned DNA molecule comprising the nucleotide sequence as shown in Figure 7.

8. An antigenic protein comprising the amino acid sequence as shown in Figure 8.

9. A cloned DNA molecule comprising the nucleotide sequence as shown in Figure 9.

10. An antigenic protein comprising the amino acid sequence as shown in Figure 10.

11. An expression vector comprising the nucleotide sequence of any of claims 1, 3, 5, 7 and 9 under

the control of a regulatory region capable of directing expression of said nucleotide sequence.

5 12. The expression vector of claim 11 wherein said vector is selected from the group consisting of plasmids, bacteriophages, viruses, or hybrids thereof.

13. A host cell or organism transformed by the expression vector of claim 11 selected from the group consisting of bacteria, yeast, fungi, insect, and mammalian cells.

10 14. A vaccine for immunizing birds against avian coccidiosis comprising the antigenic protein of any of claims 2, 4, 6, 8, and 10 combined with a carrier or adjuvant.

15 15. A method of immunizing birds against avian coccidiosis comprising administering to said birds an effective amount of the vaccine of claim 16.

16. The method of claim 15 wherein said administration is by injection or by mixing the antigenic protein with feed.

20 17. A vaccine for immunizing birds against avian coccidiosis comprising a live microorganism transformed by the expression vector of claim 11.

25 18. A method of immunizing birds against avian coccidiosis which comprises administering to said birds the vaccine of claim 17.

ATGCCAGGGCTTCCACCGCCCGCACCCGCTCAAAGGCTGATCGCTCAGACACGGCAGTG
 10 20 30 40 50 60
 5' GCCTCCGGCGGAGGAGGCGCAGCAAGCTGCCAATCTGAATCCGACACAGGAAGTT 3'
 70 80 90 100 110 120
 5' ACGACAGGTCCACTGGGGACCAGCTTCCCAAACCCGAAGAACGTTCCGGCAGAAGTAGAT 3'
 130 140 150 160 170 180
 5' GAAGAGGGAGAGCCTTCCCAATAATGAAGCATCAGGAGCGCAGACCCTTTCGAAAGAAAAG 3'
 190 200 210 220 230 240
 5' CCCAAGAAGACAAATGCGGAAGTTGACGGCGCAACATAAGCAGAAGAAGTCGCCCAAAGGAG 3'
 250 260 270 280 290 300
 5' GAGGACTTGGAGGGCACGAAGAAGAAAAGAAAAGAAAAGGAAAGGAGATGCCGCCGAGT 3'
 310 320 330 340 350 360
 5' GAGAGATCCGGCCAAACCCTCCGATGTGGAGAGCATTAAACCAAGAAAATGCTGGTGAGACT 3'
 370 380 390 400 410 420

1/12

FIG. 1A

2/12

5' GTTGCAGCAGCCGATGCCAGCCCTGGATGGGGATTTCGGGTGTTTGGAAATAGGGAAGGA 3'
430 440 450 460 470 480

5' ATTA CTGATGCTTCTA AAAGGCATTC CGACGCACAGGATGCATCACCGTCTGCGCCCTCGT 3'
490 500 510 520 530 540

5' CCTCTCTCTCGGGTTAATTGTCGGCTCGGACCATGCTGAGGAAACCCCAAGGAAAC 3'
550 560 570 580 590 600

5' GAAGTTC CCCCCTTGAAACTGTCCACTGTGGACGGTGCCGGATGACCCAGAGCCGACAGTTCC 3'
610 620 630 640 650 660

5' TCGAGTGACGACGGCGGACCGGAGGTATCACCCAGCACAAATAGCGGGTTTCTTTTGG 3'
670 680 690 700 710 720

5' GGAATCTAATGA
730

FIG. 1B

3/12

5	10	15	20
MetProGlyLeuProProAlaProAlaSerLysAlaAspArgSerAspThrAlaVal			
25	30	35	40
AlaSerAlaAlaArgGluAlaGlnGlnAlaAlaAsnLeuAsnProThrGlnGluVal			
45	50	55	60
ThrThrGlyProThrGlyAspGlnLeuProLysProGluGluArgSerAlaGluValAsp			
65	70	75	80
GluGluGlyGluProSerHisAsnGluAlaSerGlyAlaGlnThrValSerLysLysLys			
85	90	95	100
ProLysLysThrAsnAlaGluValAspGlyGluHisLysGlnLysLysSerProLysGlu			
105	110	115	120
GluAspLeuGluGlyThrLysLysLysLysLysLysLysArgLysGlyAspAlaProSer			
125	130	135	140
GluArgSerGlyGlnProSerAspValGluSerIleAsnGlnGluAsnAlaGlyGluThr			

FIG. 2A

145 150 155 160
 ValAlaAlaAlaThrProGlyTyrGlyPheGlyLeuPheGlyAsnArgGluGly

 165 170 175 180
 IleThrAspAlaSerLysArgHisSerAspAlaGlnAspAlaSerProSerAlaProArg

 185 190 195 200
 ProSerLeuLeuGlyLeuIleValGlySerAspHisAlaGluGluThrProLysGlyAsn

 205 210 215 220
 GluValProProLeuLysLeuSerThrValAspGlyAlaAspAspGlnSerAspSerSer

 225 230 235 240
 SerSerAspAspGlyGluThrAspGlyGlyIleThrSerThrIleAlaGlyPhePheTyr

 GlyIle

4/12

FIG. 2B

5' ATGGCTGACTTCTTCAGCGGCTTAGTAGGAGGTGTCGTAGGGGCTGTTGCTGCTGTGAT 3'
10 20 30 40 50 60

5' GTCCCAGCAGAAGGAGAGACTGCCCTCGGCCCTGCACCCGGTACAGAATGGAGTCTATGC 3'
70 80 90 100 110 120

5' TGTAGTAAGCTAGGAGAGCGGGGGAGGCTTGAAGGCTTTGTTTCAGCAGCTAACGTTT 3'
130 140 150 160 170 180

5' ATATTAGGCAAGCTAGCTAGCTGTATGAGGGTAGGGATCGAGCATCTAAGCCGCTGTGTT 3'
190 200 210 220 230 240

5' GCTGAAGGTAGACCGCCTTCTTCTTGTCCCTTGTTTAATGAATAAGGCTGATATAGATGAG 3'
250 260 270 280 290 300

5' GGTATAGGTGCAGGTAACAAGCGCCTGATATCTTATAAGAGGAGGTAAGCTGTGCTT 3'
310 320 330 340 350 360

5/12

FIG. 3A

SUBSTITUTE SHEET

6/12

5' GAGGCTCTTCTTGAAGCAGCTAAGGTTGCAGCTACAAGAGGTTTAACTTGTGAAGGG 3'
370 380 390 400 410 420

5' AGTAAGGATATTATATTAAGAAACATCCCCACAGACACAAGAAAGCTGTCTCAAGCATA 3'
430 440 450 460 470 480

5' AGCTCTTCTTAAAGAGGCTATGAAGGTAGCGGAGGCTCTCTAGGAGGCTATGCTCCTTCA 3'
490 500 510 520 530 540

5' TATCAGCAGCAGCAGCAGCACCCTTCAAGCTACGGTGCAGCACCTTCAAGCCAGCAAGCC 3'
550 560 570 580 590 600

5' TCAGGCTTCTTCTGGTAGA 3'
610 620

FIG. 3B

7/12

5 10 15 20
MetAlaAspPheSerGlyLeuValGlyGlyValValGlyAlaValAlaAlaValAsp
25 30 35 40
ValProAlaGluGlyGluArgLeuProArgProAlaProGlyThrGluTrpSerLeuCys
45 50 55 60
CysSerLysLeuGlyGluSerGlyArgGluLeuGluGlyPheValGlnGlnLeuThrPhe
65 70 75 80
IleLeuGlyLysLeuAlaSerCysMetArgValGlyIleGluHisLeuSerArgCysVal
85 90 95 100
AlaGluGlyArgProProSerSerCysProCysLeuMetAsnLysAlaAspIleAspGlu
105 110 115 120
GlyIleGlyAlaGlyLysGlnGlyAlaAspTyrLeuIleArgGlyGlyLysLeuValLeu

FIG. 4A

125 130 135 140
GluAlaLeuLeuGluAlaAlaLysValAlaAlaThrArgGlyLeuIleLeuValGluGly
145 150 155 160
SerLysAspIleIleLeuArgAsnIleProGlnThrGlnGluLysLeuSerGlnAlaTyr
165 170 175 180
SerSerPheLeuArgGlyTyrGluGlySerGlyArgSerLeuGlyGlyTyrAlaProSer
185 190 195 200
TyrGlnGlnGlnGlnHisProSerSerTyrGlyAlaAlaProSerSerGlnGlnAla
205
SerGlyPhePheTrp***

8/12

FIG. 4B

9/12

5' AGAGAAGAACAATAAGAGAAGAAAGACAAAGAAGAGAAAGAAAGAAAGAGAA 3'
 10 20 30 40 50 60

5' GAGAGAGGAGAAGAAGAATAAAGAAGAAGAAAGAGCGGCAGCACCTGCAGCAGCAACA 3'
 70 80 90 100 110 120

5' GCAGCAGCACCTGCAGCAGCAACAGCAGCAACAGCAGCTGCAACAGCAGCAGCTACA 3'
 130 140 150 160 170 180

5' GCAACACCAGCTGCAGCTGCAGCACCCAGCTGCAGCAGCACCCAGCTGCAGCAGCAGCA 3'
 190 200 210 220 230 240

5' GCAACAGGAGCTGCAGCAGCACCCAGCTGCAGCGGACAGCCGAGCAGGAGCAGCAGGAGCA 3'
 250 260 270 280 290 300

5' GCAAAGGCAGCGGAAGCACGAAACAGAAACCGAAGGAGCAGCAGGAGCAGCAGCAAGCA 3'
 310 320 330 340 350 360

5' GAGAAAGCGAATAACAAGCAGCAACAGCAGCAACAACAGCAGCTGCAGCAAGAGCAGCA 3'
 370 380 390 400 410 420

5' GCAGCATCAGCAGCAGGTAAGCCTGGGGACATGCAGCAGCATCAACAGCAGCAAAAGCA 3'
 430 440 450 460 470 480

5' GCGCAGGATAAAGCGGAAAGCAGCAAACGCAGCAACAGCAGCAACAGCAGCATCAAGGAA 3'
 490 500 510 520 530 540

5' TAA

FIG. 5

10/12

5 10 15 20
 ArgGluGluAspLysArgGluGluGluAspLysArgGluGluGluGluGluGluArgGlu
 25 30 35 40
 GluArgGlyGluGluGluLysGluGluArgAlaAlaProAlaAlaAlaThr
 45 50 55 60
 AlaAlaAlaProAlaAlaAlaThrAlaAlaAlaAlaAlaThrAlaAlaAlaAlaThr
 65 70 75 80
 AlaThrProAlaAlaAlaAlaProAlaAlaAlaAlaProAlaAlaAlaAlaAla
 85 90 95 100
 AlaThrGlyAlaAlaAlaProAlaAlaAlaAlaAlaAlaAlaGlyAlaAlaGlyAla
 105 110 115 120
 AlaLysAlaAlaGluAlaArgThrGluThrGluGluGlyAlaAlaGlyAlaAlaGluAla
 125 130 135 140
 GluLysAlaLysThrGlnAlaAlaAlaAlaAlaAlaAlaAlaAlaAlaAlaAlaAla
 145 150 155 160
 AlaAlaSerAlaAlaGlyLysProGlyGlyHisAlaAlaAlaSerThrAlaAlaLysAla
 165 170 175 180
 AlaGlnGluLysAlaAlaLysAlaAlaAsnAlaAlaAlaAlaAlaAlaAlaAlaSerLysGlu

FIG. 6

5' CACTCCCACACTACATCAGCAGCAGCAATAACAAGCAGCTGGCTGCCAGCAGCAGCAGCA 3'
 10 20 30 40 50 60

5' CCGCCAGCAGCAGCAGCACCGCCAACAGCAGCAGCAGCACCATCAGCAAATAAAAAA 3'
 70 80 90 100 110

FIG. 7

5 10 15 20
 HisSerHisThrThrSerAlaAlaAlaIleGlnAlaAlaAlaLeuProAlaAlaAlaAla

25 30 35
 ProProAlaAlaAlaProProThrAlaAlaAlaAlaProSerAlaLysLysLys

FIG. 8

```

5' CTCATGGGGCTTCAGGCAGAGAGTCCCTGCTGCTGCTTCCCCTGCTGCTGCTGCTGCTGCT 3'
  10    20    30    40    50    60
5' GCTGCTGCACCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAG 3'
  70    80    90   100   110   120
5' GCGGCCGAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAG 3'
  130   140   150   160   170   180
5' CAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAG 3'
  190   200   210   220
    
```

FIG. 9

```

          5           10           15           20
LeuMetGlyLeuProGlyArgGluSerProAlaAlaAlaSerProAlaAlaAlaAlaAla
          25           30           35           40
AlaAlaAlaProAlaAlaAlaAlaAlaAlaAlaAlaGluProAspSerGlyAlaAlaSer
          45           50           55           60
GlyAlaGluGlnGlnHisGlnLeuGlnGlnHisGlnAspTyrGlnGlnAspGlnGlnGln
          65           70           75
GlnGlnGlnGlnGlnGlnGlnGlnGlnValLeuValMet
    
```

FIG. 10

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/06430

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): C12P 21/00; C12N 1/20, 15/00; A61K 39/00; C07H 15/12 U.S. CL.: 435/69.1, 252.3, 320.1; 424/88; 530/806, 350; 536/27		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	435/69.1, 252.3, 320.1; 424/88; 530/806, 350; 536/27	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
APS and DIALOG (BIOSIS, CAS, MEPLINE, WORLD PATENT INDEX) coccidio2, antigen2, DNA, gene, sequence?, eimeria		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	US, A, 4,874,705 (Andrews et al.) 17 October 1989. see entire document.	1-18
Y	US, A, 4,639,372 (Murray et al.) 27 January 1987, see entire document.	1-18
Y	US, A, 4,710,377 (Schenkel et al.) 01 December 1987, see entire document.	1-18
P, Y	US, A, 5,028,694 (Mewman Jr. et al.) 02 July 1991, see entire document.	1-18
Y	EP, A, 0,135,712 (Schenkel et al.) 03 April 1985, see entire document.	1-18
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
04 December 1991	18 DEC 1991	
International Searching Authority	Signature of Authorized Officer	
ISA/US	T. Michael Nisbet ebw	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Chemical Abstracts, volume 106, issued 1987, Clarke et al., "Isolating of amp 3 genomic recombinants coding for antigens of <u>Eimeria tenella</u> ", pages 189-190, see Abstract number 132966 See entire document.	1-18
Y	EP. A. 0,164,176 (Newman Jr., et al.) 11 December 1985, see entire document.	1-18
Y	US. A. 4,650,576 (Schenkel et al.) 17 March 1987, see entire document.	1-18