



(51) International Patent Classification:

C07C 317/22 (2006.01) *A61K 31/397* (2006.01)
C07D 205/04 (2006.01) *A61K 31/4453* (2006.01)
C07D 295/088 (2006.01) *A61P 25/14* (2006.01)
A61K 31/138 (2006.01) *A61P 25/16* (2006.01)
A61K 31/145 (2006.01) *A61P 25/18* (2006.01)

(21) International Application Number:

PCT/EP2012/056959

(22) International Filing Date:

17 April 2012 (17.04.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PA 2011 70187 19 April 2011 (19.04.2011) DK
61/476,810 19 April 2011 (19.04.2011) US
PA 2011 70495 6 September 2011 (06.09.2011) DK

(71) Applicant (for all designated States except US): **NEUROSEARCH A/S** [DK/DK]; Pederstrupvej 93, DK-2750 Ballerup (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SONESSON, Clas** [SE/SE]; Billdals Hagenväg 18, S-427 37 Billdal (SE). **KARLSSON, Jonas** [SE/SE]; Skepparegången 12, S-413 18 Göteborg (SE). **SVENSSON, Peder** [SE/SE]; Nordenskiöldsgatan 12, S-413 09 Göteborg (SE).

(74) Common Representative: **NEUROSEARCH A/S**; Intellectual Property Rights, Pederstrupvej 93, DK-2750 Ballerup (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: NOVEL MODULATORS OF CORTICAL DOPAMINERGIC- AND NMDA-RECEPTOR-MEDIATED GLUTAMATERGIC NEUROTRANSMISSION

(57) Abstract: The present invention relates to novel substituted phenoxyethylamine derivatives, useful as modulators of cortical and basal ganglia dopaminergic and N-methyl-D-aspartate (NMDA) receptor-mediated glutamatergic neurotransmission. In other aspects the invention relates to the use of these compounds in a method for therapy and to pharmaceutical compositions comprising the compounds of the invention.



NOVEL MODULATORS OF CORTICAL DOPAMINERGIC- AND NMDA-RECEPTOR-MEDIATED GLUTAMATERGIC NEUROTRANSMISSION

TECHNICAL FIELD

5

The present invention relates to novel substituted phenoxy-ethyl-amine derivatives, useful as modulators of cortical and basal ganglia dopaminergic and N-methyl-D-aspartate (NMDA) receptor-mediated glutamatergic neurotransmission. In other aspects the invention relates to the use of these
10 compounds in a method for therapy and to pharmaceutical compositions comprising the compounds of the invention.

BACKGROUND ART

15 Dopamine is a neurotransmitter in the brain. Since this discovery, made in the 1950's, the function of dopamine in the brain has been intensely explored. To date, it is well established that dopamine is essential in several aspects of brain function including motor, cognitive, sensory, emotional and autonomous functions (e.g. regulation of appetite, body temperature, sleep). Thus, modulation of
20 dopaminergic function may be beneficial in the treatment of a wide range of disorders affecting brain functions. In fact, drugs that act, directly or indirectly at central dopamine receptors are commonly used in the treatment of neurological and psychiatric disorders, e.g. Huntington and Parkinson's disease and schizophrenia.

25 Antipsychotic drugs (or neuroleptics) are a class of compounds with diverse effects on different receptor systems. However, they have in common the ability to block dopamine D₂ receptors in the basal ganglia (i.e. striatum) and are used to manage psychosis (including delusions or hallucinations, as well as disordered thought), particularly in schizophrenia and bipolar disorder.

30 The cerebral cortex encompasses several major regions that are involved in higher functions such as thought, feelings, memory and planning. Biogenic amines such as dopamine are important for mammalian cortical function.

The ascending dopamine pathways innervate the cortex. Primary or secondary dysfunctions in the activity of these pathways lead to dysregulation of the activity at dopamine in these brain areas and subsequently to manifestations of psychiatric and neurological symptoms. Both dopamine D₁ and N-methyl-D-
5 aspartate (NMDA) receptors in the prefrontal cortex play a critical role in synaptic plasticity, memory mechanisms, and cognition.

Huntington's disease (HD) is a rare neurodegenerative disorder of the central nervous system characterized by progressive deterioration of motor and cognitive functions as well as behavioural and psychiatric disturbances. It is well
10 established that certain aspect of dopaminergic functions are also affected in Huntington's disease. Neuropathological changes in Huntington's disease involve prominent cell loss and atrophy in the striatum but also in many other brain regions such as the cortex, substantia nigra, hypothalamus, cerebellum and thalamus.

In HD, glutamate and dopamine (DA) transmission is altered, which is
15 likely to induce an imbalance in activity of the direct and indirect pathways and to contribute to the motor, cognitive, and psychiatric symptoms of HD (i.e. communication between the cortex and striatum, *Capeda et al*; ASN Neuro 2010 2 (2) e00033). Therefore, compounds which can strengthen the cortical dopamine and NMDA transmission, and exert antagonism of excessive subcortical dopamine
20 transmission, can balance aberrant functioning in the cortico-striato-thalamic network controlling motor functions (*Alexander et al*; Ann. Rev. Neurosci. 1986 9 357-381).

JP 2006-193494 (Dainippon Ink and Chemicals, Inc) describes certain quaternary ammonium compounds useful as therapeutic agent for heart diseases.

25 WO 2009/133107 (NSAB, Filial af NeuroSearch Sweden AB, Sverige) describes certain 1-(2,3-dihydro-1,4-benzodioxin-2-yl)methanamine derivatives, WO 2009/133109 (NSAB, Filial af NeuroSearch Sweden AB, Sverige) describes certain 1-(2,3-dihydro-1,4-benzodioxin-2-yl)methanamine derivatives, and WO 2009/133110 (NSAB, Filial af NeuroSearch Sweden AB, Sverige) describes
30 certain 1-(4H-1,3-benzodioxin-2-yl)methanamine derivatives, useful as modulators of dopamine neurotransmission, and more specifically as dopaminergic stabilizers.


However, the phenoxy-ethyl-amine derivatives of the present invention have not previously been reported.

SUMMARY OF THE INVENTION

5

The object of the present invention is to provide novel pharmaceutically active compounds, especially useful in treatment of disorders in the central nervous system. A further object is the provision of compounds for modulation of dopaminergic and glutamatergic systems in the mammalian brain, including
10 human brain.

In its first aspect, the invention provides a phenoxy-ethyl-amine derivative of Formula 1



a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof,
15 or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁴, R' and R'' are as defined below.

In its second aspect, the invention provides a pharmaceutical composition, comprising a therapeutically effective amount of a phenoxy-ethyl-amine derivative of the invention, a stereoisomer or a mixture of its stereoisomers,
20 or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, together with at least one pharmaceutically acceptable carrier, excipient or diluent.

In a further aspect, the invention provides the use of a phenoxy-ethyl-amine derivative of the invention, a stereoisomer or a mixture of its stereoisomers or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, for the
25 manufacture of a pharmaceutical composition for the treatment, prevention or alleviation of a disease or a disorder or a condition of a mammal, including a

human, which disease, disorder or condition is responsive to responsive to modulation of dopaminergic and glutamatergic function in the central nervous system.


In a still further aspect, the invention relates to a method for treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disorder, disease or condition is responsive to modulation of dopaminergic and glutamatergic function in the central nervous system, which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of a phenoxy-ethyl-amine derivative of the invention, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof.

Other aspects of the invention will be apparent to the person skilled in the art from the following detailed description and examples.

DETAILED DESCRIPTION OF THE INVENTION

Phenoxyethylamine derivatives

In its first aspect the present invention provides phenoxy-ethyl-amine derivatives of Formula 1



a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein

R^1 represents CH_3 or CF_3 ;

R^2 is selected from the group consisting of C_1 - C_4 -alkyl, allyl, $CH_2CH_2OCH_3$, $C(CH_3)_2CH_2CH_3$, CH_2 -cyclopropyl, cyclobutyl, cyclopentyl,

CH₂CH₂CH₂F, CH₂CH₂CHF₂, CH₂CH₂F, 3,3,3-trifluoropropyl and 4,4,4-trifluorobutyl; and


R³ is selected from the group consisting of H, CH₃ and CH₂CH₃; or

R² and R³ together form CH₂(CH₂)CH₂ or CH₂(CH₂)₃CH₂;

5 R⁴ represents F or Cl; and

R' and R'' independently represent hydrogen or methyl.

In a preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1a



10 a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein

R¹ is selected from the group consisting of CH₃ or CF₃;

R² is selected from the group consisting of C₁-C₄ alkyl, allyl, CH₂CH₂OCH₃, C(CH₃)₂CH₂CH₃, CH₂-cyclopropyl, CH₂CH₂CH₂F, CH₂CH₂CHF₂,
 15 CH₂CH₂F, 3,3,3-trifluoropropyl and 4,4,4-trifluorobutyl;

R³ is selected from the group consisting of H, CH₃ and CH₂CH₃; and

R⁴ is selected from the group consisting of F and Cl.

In another preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1a, a stereoisomer or a mixture of its
 20 stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein

R¹ is selected from the group consisting of CH₃ or CF₃;

R² is selected from the group consisting of C₁-C₄ alkyl, allyl, CH₂CH₂OCH₃, C(CH₃)₂CH₂CH₃, CH₂-cyclopropyl, cyclobutyl, cyclopentyl,
 25 CH₂CH₂CH₂F, CH₂CH₂CHF₂, CH₂CH₂F, 3,3,3-trifluoropropyl and 4,4,4-trifluorobutyl;

R³ is selected from the group consisting of H, CH₃ and CH₂CH₃;

or R^2 and R^3 together form $\text{CH}_2(\text{CH}_2)_3\text{CH}_2$; and

R^4 is selected from the group consisting of F and Cl.

In a third preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1 or 1a, a stereoisomer or a mixture of its
5 stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein R^1 represents CH_3 or CF_3 .

In a more preferred embodiment R^1 represents CH_3 .

In another more preferred embodiment R^1 represents CF_3 .

In a fourth preferred embodiment the phenoxy-ethyl-amine derivative of
10 the invention is a compound of Formula 1 or 1a, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from the group consisting of C_1 - C_4 -alkyl, allyl, $\text{CH}_2\text{CH}_2\text{OCH}_3$, $\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$, CH_2 -cyclopropyl, cyclobutyl, cyclopentyl, $\text{CH}_2\text{CH}_2\text{CH}_2\text{F}$, $\text{CH}_2\text{CH}_2\text{CHF}_2$, $\text{CH}_2\text{CH}_2\text{F}$, 3,3,3-trifluoropropyl and 4,4,4-
15 trifluorobutyl.

In a more preferred embodiment R^2 is selected from the group consisting of C_1 - C_4 -alkyl, CH_2 -cyclopropyl, cyclobutyl and cyclopentyl.

In another more preferred embodiment R^2 represents C_1 - C_4 -alkyl.

In a third more preferred embodiment R^2 represents CH_2 -cyclopropyl.

20 In a fourth more preferred embodiment R^2 represents cyclobutyl.

In a fifth more preferred embodiment R^2 represents cyclopentyl.

In a fifth preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1 or 1a, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt
25 thereof, wherein R^3 is selected from the group consisting of H and CH_3 .

In a more preferred embodiment R^3 represents H.

In another more preferred embodiment R^3 represents CH_3 .

In a sixth preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1 or 1a, a stereoisomer or a mixture of its
30 stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein R^2 and R^3 together form $\text{CH}_2(\text{CH}_2)\text{CH}_2$ or $\text{CH}_2(\text{CH}_2)_3\text{CH}_2$.

In a more preferred embodiment R^2 and R^3 together form $CH_2(CH_2)CH_2$.

In another more preferred embodiment R^2 and R^3 together form $CH_2(CH_2)_3CH_2$.

5 In a seventh preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1 or 1a, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein R^4 represents F or Cl.

In a more preferred embodiment R^4 represents F.

10 In another more preferred embodiment R^4 represents Cl.

In an eight preferred embodiment the phenoxy-ethyl-amine derivative of the invention is a compound of Formula 1, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a pharmaceutically acceptable salt thereof, wherein R' and R'' independently represent hydrogen or methyl.

15 In a more preferred embodiment one of R' and R'' represents hydrogen; and the other of R' and R'' represents methyl.

In another more preferred embodiment R' represents hydrogen; and R'' represents methyl.

In a third more preferred embodiment R' represents methyl; and R'' represents hydrogen.

In a fourth more preferred embodiment R' and R'' both represents hydrogen.

In a fifth more preferred embodiment R' and R'' both represents methyl.

In a most preferred embodiment the phenoxy-ethyl-amine derivative of the invention is

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]propan-1-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]butan-1-amine;

N-Ethyl-2-(3-fluoro-5-methylsulfonyl-phenony)ethanamine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]-N-methyl-propan-1-amine;

N-[2-(3-Chloro-5-methylsulfonyl-phenoxy)ethyl]propan-1-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]butan-2-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]cyclopentamine;
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]-2-methyl-butan-2-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]cyclobutamine;
5 N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]propan-2-amine;
1-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]piperidine;
N,N-Diethyl-2-(3-fluoro-5-methylsulfonyl-phenoxy)ethanamine;
N-1,1-di-deuterium-propyl-[2-(3-fluoro-5-methanesulfonyl-phenoxy)-ethyl]-amine;

10 N-(Cyclopropylmethyl)-N-{2-[3-fluoro-5-(methylsulfonyl)-phenoxy]ethyl}amine;

N-{2-[3-Fluoro-5-(methylsulfonyl)phenoxy]ethyl}-N-isobutylamine;
1-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]azetidine;
1-(3-Fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-2-amine; or
15 2-(3-Fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-1-amine;
a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof,
or a pharmaceutically acceptable salt thereof.

Any combination of two or more of the embodiments as described above is considered within the scope of the present invention.

20

Definition of terms

In the context of this invention C₁-C₄-alkyl means a straight chain or branched chain of one to four carbon atoms, including but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl.

25 The term "allyl" refers to the group -CH₂-CH=CH₂.

The term "treatment" as used herein means the management and care of a patient for the purpose of combating a disease, disorder or condition. The term is intended to include the delaying of the progression of the disease, disorder or condition, the alleviation or relief of symptoms and complications, and/or the
30 cure or elimination of the disease, disorder or condition. The patient to be treated is preferably a mammal, in particular a human being.

The terms "disease", "condition" and "disorder" as used herein are used interchangeably to specify a state of a patient which is not the normal physiological state of man.

The term "medicament" as used herein means a pharmaceutical
5 composition suitable for administration of the pharmaceutically active compound to a patient.

The term "pharmaceutically acceptable" as used herein means suited for normal pharmaceutical applications, i.e. giving rise to no adverse events in patients etc.

10 The term "effective amount" as used herein means a dosage which is sufficient in order for the treatment of the patient to be effective compared with no treatment.

The term "therapeutically effective amount" of a compound as used herein means an amount sufficient to cure, alleviate or partially arrest the clinical
15 manifestations of a given disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective amount". Effective amounts for each purpose will depend on the severity of the disease or injury as well as the weight and general state of the subject. It will be understood that determining an appropriate dosage may be achieved using routine experimentation, by
20 constructing a matrix of values and testing different points in the matrix, which is all within the ordinary skills of a trained physician or veterinary.

Pharmaceutically Acceptable Salts

The compound of the invention may be provided in any form suitable for
25 the intended administration. Suitable forms include pharmaceutically (i.e. physiologically) acceptable salts of the compound of the invention.

Such pharmaceutically acceptable salts and common methodology for preparing them are known in the art. Further details may be found in *P Stahl et al*, Handbook of Pharmaceutical Salts: Properties, Selection and Use; Wiley-VCH,
30 2002.

The chemical compound of the invention may be provided in dissoluble or indissoluble forms together with a pharmaceutically acceptable solvent such as

water, ethanol, and the like. Dissoluble forms may also include hydrated forms such as the monohydrate, the dihydrate, the hemihydrate, the trihydrate, the tetrahydrate, and the like. In general, the dissoluble forms are considered equivalent to indissoluble forms for the purposes of this invention.

5

Steric Isomers

It will be appreciated by those skilled in the art that the compounds of the present invention may exist in different stereoisomeric forms, including enantiomers, diastereomers or cis-trans-isomers.

10 The invention includes all such isomers and any mixtures thereof including racemic mixtures.

Racemic forms can be resolved into the optical antipodes by known methods and techniques. One way of separating the enantiomeric compounds (including enantiomeric intermediates) is, in the case the compound being a chiral
15 acid, by use of an optically active amine, and liberating the diastereomeric, resolved salt by treatment with an acid. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optical active matrix. Racemic compounds of the present invention can thus be resolved into their optical antipodes, e.g., by fractional crystallisation of D- or L- (tartrates,
20 mandelates, or camphorsulphonate) salts for example.

The chemical compounds of the present invention may also be resolved by the formation of diastereomeric amides by reaction of the chemical compounds of the present invention with an optically active carboxylic acid such as that derived from (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic
25 acid or by the formation of diastereomeric carbamates by reaction of the chemical compound of the present invention with an optically active chloroformate or the like.

Additional methods for the resolving the optical isomers are known in the art. Such methods include those described by *Jaques J, Collet A, & Wilen S* in
30 "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Optical active compounds can also be prepared from optical active starting materials.

N-oxides

5 In the context of this invention an N-oxide designates an oxide derivative of a tertiary amine, including a nitrogen atom of an aromatic N-heterocyclic compound, a non-aromatic N-heterocyclic compounds, a trialkylamine and a trialkenylamine.

N-oxides of the compounds of the invention may be prepared by
10 oxidation of the corresponding nitrogen base using a conventional oxidizing agent such as hydrogen peroxide in the presence of an acid such as acetic acid at an elevated temperature, or by reaction with a peracid such as peracetic acid in a suitable solvent, e.g. dichloromethane, ethyl acetate or methyl acetate, or in chloroform or dichloromethane with 3-chloroperoxybenzoic acid.

15

Labelled Compounds

The compounds of the invention may be used in their labelled or unlabelled form. In the context of this invention the labelled compound has one or more atoms replaced by an atom having an atomic mass or mass number different
20 from the atomic mass or mass number usually found in nature. The labelling will allow easy quantitative detection of said compound.

The labelled compounds of the invention may be useful as diagnostic tools, radio tracers, or monitoring agents in various diagnostic methods, and for *in vivo* receptor imaging.

25 The labelled isomer of the invention preferably contains at least one radionuclide as a label. Positron emitting radionuclides are all candidates for usage. In the context of this invention the radionuclide is preferably selected from ^2H (deuterium), ^3H (tritium), ^{11}C , ^{13}C , ^{14}C , ^{131}I , ^{125}I , ^{123}I and ^{18}F .

The physical method for detecting the labelled isomer of the present
30 invention may be selected from Position Emission Tomography (PET), Single Photon Imaging Computed Tomography (SPECT), Magnetic Resonance

Spectroscopy (MRS), Magnetic Resonance Imaging (MRI), and Computed Axial X-ray Tomography (CAT), or combinations thereof.

Deuterated analogs

5 The compounds of the invention may be provided in the form of their deuterated analogs. Deuterium forms bonds with carbon that vibrate at a lower frequency and are thus stronger than C-H bonds. Therefore "heavy hydrogen" (deuterium) versions of drugs may be more stable towards degradation and last longer in the living organism.

10

Methods of Preparation

 The chemical compounds of the invention may be prepared by conventional methods for chemical synthesis, e.g. those described in the working examples. The starting materials for the processes described in the present
15 application are known or may readily be prepared by conventional methods from commercially available chemicals.

 Also one compound of the invention can be converted to another compound of the invention using conventional methods.

 The end products of the reactions described herein may be isolated by
20 conventional techniques, e.g. by extraction, crystallisation, distillation, chromatography, etc.

 Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative, and in some occasions, more convenient manner, the individual process steps mentioned hereinbefore may be
25 performed in a different order, and/or the individual reactions may be performed at different stage in the overall route (*i.e.* chemical transformations may be performed upon different intermediates to those associated hereinbefore with a particular reaction).

30 **Biological Activity**

 The compounds according to the present invention possess modulation of cortical and basal ganglia dopaminergic and N-methyl-D-aspartate (NMDA)

receptor-mediated glutamatergic neurotransmission and both they and their pharmaceutical compositions are useful in treating numerous central nervous system disorders, including both psychiatric and neurological disorders.

Particularly, the compounds and their pharmaceutical compositions may be used
5 in the treatment of CNS disorders where the dopaminergic and glutamatergic system is dysfunctional due to direct or indirect causes.

The compounds and compositions according to the invention can be used to improve all forms of psychosis, including schizophrenia and schizophreniform and bipolar disorders as well as drug induced psychotic
10 disorders. Iatrogenic psychoses and hallucinoses and non-iatrogenic psychoses and hallucinoses may also be treated.

In a special embodiment the disease, disorder or condition contemplated according to the invention is a form of psychosis, in particular schizophrenia, a schizophreniform disorder, a bipolar disorder, or a drug induced
15 psychotic disorder.

Mood and anxiety disorders, depression and obsessive-compulsive disease may also be treated with the compounds and compositions according to the invention.

Compounds with modulating effects on dopaminergic and glutamatergic
20 systems may also be used to improve motor and cognitive functions and in the treatment of emotional disturbances related to ageing, neurodegenerative (e.g. dementia and age-related cognitive impairment) and developmental disorders (such as Autism spectrum disorders, ADHD, Cerebral Palsy, Gilles de la Tourette's syndrome) as well as after brain injury. Such brain injury may be
25 induced by traumatic, inflammatory, infectious, neoplastic, vascular, hypoxic or metabolic causes or by toxic reactions to exogenous chemicals, wherein the exogenous chemicals are selected from the group consisting of substances of abuse, pharmaceutical compounds and environmental toxins

The compounds and pharmaceutical compositions according to the
30 invention may also be used in behavioural disorders usually first diagnosed in infancy, childhood, or adolescence as well as in impulse control disorders.

They can also be used for treating substance abuse disorders as well as disorders characterized by misuse of food. They are further useful for treatment of a condition selected from the group consisting of sleep disorders, sexual disorders, eating disorders, obesitas, and headaches and other pains in conditions
5 characterized by increased muscular tone.

Neurological indications include the use of the compounds and their pharmaceutical compositions to improve mental and motor function in Parkinson's disease, and in related parkinsonian syndromes, dyskinesias (including L-DOPA induced dyskinesias and tardive dyskinesias) and dystonias. They may also be
10 used to ameliorate tics and tremor of different origins.

They can also be used in the treatment of Huntington's disease and other movement disorders as well as movement disorders induced by drugs. Restless legs and related disorders as well as narcolepsy may also be treated with compounds included according to the invention.

15 The compounds and their pharmaceutical compositions according to the present invention can be used for the treatment of Alzheimer's disease or related dementia disorders.

In a further embodiment the disease, disorder or condition contemplated according to the invention is selected from the group of
20 schizophrenia, L-DOPA induced dyskinesias and Huntington's disease.

Pharmaceutical Compositions

In another aspect the invention provides novel pharmaceutical compositions comprising a therapeutically effective amount of a compound of the
25 invention.

While a compound of the invention for use in therapy may be administered in the form of the raw compound, it is preferred to introduce the active ingredient, optionally in the form of a physiologically acceptable salt, in a pharmaceutical composition together with one or more adjuvants, excipients,
30 carriers, buffers, diluents, and/or other customary pharmaceutical auxiliaries.

In a preferred embodiment, the invention provides pharmaceutical compositions comprising the compound of the invention, or a pharmaceutically

acceptable salt or derivative thereof, together with one or more pharmaceutically acceptable carriers therefore, and, optionally, other therapeutic and/or prophylactic ingredients, known and used in the art. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not
5 harmful to the recipient thereof.

The pharmaceutical composition of the invention may be administered by any convenient route, which suits the desired therapy. Preferred routes of administration include oral administration, in particular in tablet, in capsule, in dragé, in powder, or in liquid form, and parenteral administration, in particular
10 cutaneous, subcutaneous, intramuscular, or intravenous injection. The pharmaceutical composition of the invention can be manufactured by the skilled person by use of standard methods and conventional techniques appropriate to the desired formulation. When desired, compositions adapted to give sustained release of the active ingredient may be employed.

15 Pharmaceutical compositions of the invention may be those suitable for oral, rectal, bronchial, nasal, pulmonal, topical (including buccal and sub-lingual), transdermal, vaginal or parenteral (including cutaneous, subcutaneous, intramuscular, intraperitoneal, intravenous, intraarterial, intracerebral, intraocular injection or infusion) administration, or those in a form suitable for administration
20 by inhalation or insufflation, including powders and liquid aerosol administration, or by sustained release systems. Suitable examples of sustained release systems include semipermeable matrices of solid hydrophobic polymers containing the compound of the invention, which matrices may be in form of shaped articles, e.g. films or microcapsules.

25 The chemical compound of the invention, together with a conventional adjuvant, carrier, or diluent, may thus be placed into the form of pharmaceutical compositions and unit dosages thereof. Such forms include solids, and in particular tablets, filled capsules, powder and pellet forms, and liquids, in particular aqueous or non-aqueous solutions, suspensions, emulsions, elixirs, and capsules
30 filled with the same, all for oral use, suppositories for rectal administration, and sterile injectable solutions for parenteral use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in

conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed.

5 The chemical compound of the present invention can be administered in a wide variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms may comprise, as the active component, either a chemical compound of the invention or a pharmaceutically acceptable salt of a chemical compound of the invention.

10 For preparing pharmaceutical compositions from a chemical compound of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers,
15 lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

 In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided active component.

 In tablets, the active component is mixed with the carrier having the
20 necessary binding capacity in suitable proportions and compacted in the shape and size desired.

 The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, cellulose,
25 starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets
30 and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glyceride or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized moulds, allowed to cool, and thereby to
5 solidify.

Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Liquid preparations include solutions, suspensions, and emulsions, for
10 example, water or water-propylene glycol solutions. For example, parenteral injection liquid preparations can be formulated as solutions in aqueous polyethylene glycol solution.

The chemical compound according to the present invention may thus be formulated for parenteral administration (e.g. by injection, for example bolus
15 injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulation agents such as suspending, stabilising and/or dispersing agents.
20 Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavours, stabilising
25 and thickening agents, as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, or other well known suspending agents.

30 Also included are solid form preparations, intended for conversion shortly before use to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. In addition to the active

component such preparations may comprise colorants, flavours, stabilisers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

For topical administration to the epidermis the chemical compound of
5 the invention may be formulated as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents,
10 suspending agents, thickening agents, or colouring agents.

Compositions suitable for topical administration in the mouth include lozenges comprising the active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerine or sucrose and acacia; and mouthwashes
15 comprising the active ingredient in a suitable liquid carrier.

Solutions or suspensions are applied directly to the nasal cavity by conventional means, for example with a dropper, pipette or spray. The compositions may be provided in single or multi-dose form.

Administration to the respiratory tract may also be achieved by means
20 of an aerosol formulation in which the active ingredient is provided in a pressurised pack with a suitable propellant such as a chlorofluorocarbon (CFC) for example dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by provision of a
25 metered valve.

Alternatively the active ingredients may be provided in the form of a dry powder, for example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). Conveniently the powder carrier will form a gel in
30 the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of, e.g., gelatin, or blister packs from which the powder may be administered by means of an inhaler.

In compositions intended for administration to the respiratory tract, including in-tranasal compositions, the compound will generally have a small particle size for example of the order of 5 microns or less. Such a particle size may be obtained by means known in the art, for example by micronization.

5 When desired, compositions adapted to give sustained release of the active in-gredient may be employed.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged
10 preparation, the package containing discrete quantities of preparation, such as packaged tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Tablets or capsules for oral administration and liquids for intravenous
15 administration and continuous infusion are preferred compositions.

In one embodiment, when the pharmaceutical composition of the invention is intended for treating patients with abuse liability and withdrawal symptoms caused by nicotine addiction, formulations such as gums, patches, sprays, inhalers, aerosols, etc. are contemplated.

20 Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and
25 toxicity, e.g. ED_{50} and LD_{50} , may be determined by standard pharmacological procedures in cell cultures or experimental animals. The dose ratio between therapeutic and toxic effects is the therapeutic index and may be expressed by the ratio LD_{50}/ED_{50} . Pharmaceutical compositions exhibiting large therapeutic indexes are preferred.

30 The dose administered must of course be carefully adjusted to the age, weight and condition of the individual being treated, as well as the route of

administration, dosage form and regimen, and the result desired, and the exact dosage should of course be determined by the practitioner.

The actual dosage depends on the nature and severity of the disease being treated, the exact mode of administration and form of administration, and is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect. However, it is presently contemplated that pharmaceutical compositions containing of from about 0.1 to about 500 mg of active ingredient per individual dose, preferably of from about 1 to about 100 mg, most preferred of from about 1 to about 10 mg, are suitable for therapeutic treatments.

The active ingredient may be administered in one or several doses per day. A satisfactory result can, in certain instances, be obtained at a dosage as low as 0.1 $\mu\text{g/kg}$ i.v. and 1 $\mu\text{g/kg}$ p.o. The upper limit of the dosage range is presently considered to be about 10 mg/kg i.v. and 100 mg/kg p.o. Preferred ranges are from about 0.1 $\mu\text{g/kg}$ to about 10 mg/kg/day i.v., and from about 1 $\mu\text{g/kg}$ to about 100 mg/kg/day p.o.

Methods of therapy

The compounds of the present invention are modulators of cortical and basal ganglia dopaminergic and N-methyl-D-aspartate (NMDA) receptor-mediated glutamatergic neurotransmission and therefore useful for the treatment of a range of ailments involving modulation of dopaminergic and glutamatergic function.

In another aspect the invention provides a method for the treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disease, disorder or condition is responsive to modulation of dopaminergic and glutamatergic function in the central nervous system, and which method comprises administering to such a living animal body, including a human, in need thereof an effective amount of a compound of the invention, a stereoisomer or a mixture of its stereoisomers, or a pharmaceutically acceptable salt thereof.


The indications contemplated according to the invention are those stated above.

It is at present contemplated that suitable dosage ranges are 0.1 to 1000 milligrams daily, 10-500 milligrams daily, and especially 30-100 milligrams daily, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved and the body weight of the subject involved, and further the preference and experience of the physician or veterinarian in charge.

EXAMPLES

The invention is further illustrated in the examples below and as outlined below in Scheme 1, which in no way are intended to limit the scope of the invention.

Scheme 1



The substituents in Scheme 1 are as follows: A is a leaving group, and R², R³, R⁴, R' and R'' are as defined above.

Example 1

N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]PROPAN-1-AMINE

A mixture of 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (2.65 g, 8.9 mmol) and propan-1-amine (5.24 ml, 63.7 mmol) in ethanol (32 ml) was divided into 3 aliquots which were each heated under microwave radiation at
5 120°C for 30 min. The reaction mixtures were cooled to room temperature, filtrated and pooled before the volatiles were evaporated. Purification by flash column chromatography (ethyl acetate/methanol, 1:0 to 1:1) gave the title compound (2.14 g, 87%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 191°C. MS m/z (relative intensity, 70 eV) 275
10 (M+, 2), 246 (32), 73 (5), 72 (bp), 56 (11).

Example 2N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]BUTAN-1-AMINE

Preparation according to Example 1 but performed in one portion: 1-(2-
15 bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.3 g, 1.01 mmol) and butan-1-amine (0.61 ml, 0.5907 mmol) in ethanol (4 ml). Yield: 290 mg (99%). The amine was converted to the hydrochloric acid salt and re-crystallized from ethanol/diethyl ether: M.p. 204°C. MS m/z (relative intensity, 70 eV) 289 (M+, 1), 246 (35), 86 (bp), 56 (12), 87 (12).

20

Example 3N-ETHYL-2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHANAMINE

Preparation according to Example 1 but performed in one portion: 1-(2-
bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.321 g, 1.08 mmol) and
25 ethanamine (4.3 ml, 8.6 mmol 2 M in methanol) in ethanol (6 ml). Yield: 255 mg (90%). The amine was converted to the hydrochloric acid salt and re-crystallized from ethanol/diethyl ether: M.p. 200.8-201.1°C. MS m/z (relative intensity, 70 eV) 261 (M+, 3), 94 (6), 59 (12), 58 (bp), 56 (8).

30 Example 4N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]-N-METHYL-PROPAN-1-AMINE

A mixture of Example 1 (0.54 g, 1.95 mmol) in formic acid (5.75 ml) and formaldehyde (40% solution, 5.1 ml) was heated at 85°C for 5 h. The solution was allowed to reach ambient temperature, water (5 ml) and diethyl ether was added, the phases were separated and the aqueous phase was basified by the addition of aqueous sodium hydroxide (5 M). The aqueous phase was extracted twice with ethyl acetate, the combined organic phases were dried (Na₂SO₄) and evaporated under pressure to give the crude product which was then purified by flash chromatography (EtOAc:MeOH 100:0 then gradually changed to 0:100. The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 128-130°C. MS m/z (relative intensity, 70 eV). 289 (M⁺, 1), 260 (26), 87 (6), 86 (bp), 58 (6).

Example 5

N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]BUTAN-2-AMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and sec-butylamine 1.37 ml, 13.46 mmol) in ethanol (5 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title compound (342 mg, 70%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 166.5°C. MS m/z (relative intensity, 70 eV) 289 (M⁺, 1), 261 (25), 260 (bp), 86 (49), 70 (12).

Example 6

N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]CYCLOPENTANAMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and cyclopentylamine (1.34 ml, 13.46 mmol) in ethanol (5 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title compound (443 mg, 87.4%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 207.5°C. MS m/z (relative intensity, 70 eV) 301(M⁺, 2), 272 (7), 99 (24), 98 (bp), 70 (7).

Example 7N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]-2-METHYL-BUTAN-2-AMINE

5 Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and tert-amylamine (1.61 ml, 13.46 mmol) in ethanol (5 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title compound (389 mg, 76.2%). The amine was converted to the hydrochloric acid salt and re-
10 crystallized from methanol/diethyl ether: M.p. 190.2°C. MS m/z (relative intensity, 70 eV) 303 (M⁺, 0), 288 (21), 276 (10), 275 (41), 274 (bp), 84 (10).

Example 8N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]CYCLOBUTANAMINE

15 Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and cyclobutylamine (1.17 ml, 13.46 mmol) in ethanol (4 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title
20 compound (400 mg, 82.7%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 202°C. MS m/z (relative intensity, 70 eV) 287 (M⁺, 0), 260 (33), 259 (91), 216 (bp), 215 (23), 56 (73).

Example 9N-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]PROPAN-2-AMINE

25 Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and isopropylamine (1.15 ml, 13.46 mmol) in ethanol (4 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title
30 compound (421 mg, 90.9%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 173°C. MS m/z (relative intensity, 70 eV) 275 (M⁺, 4), 261 (16), 260 (41), 73 (32), 72 (bp).

Example 101-[2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHYL]PIPERIDINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and piperidine (1.33 ml, 13.46 mmol) in ethanol (5 ml). Purification by flash column chromatography (ethyl acetate/methanol, 100:0 to 85:15) gave the title compound (480 mg, 94.7%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 194.6°C. MS m/z (relative intensity, 70 eV) 301(M⁺, 1), 99 (8), 98 (bp), 96 (4), 55 (5).

Example 11N,N-DIETHYL-2-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)ETHANAMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.68 mmol) and diethylamine (1.39 ml, 13.46 mmol) in ethanol (4 ml). Purification by flash column chromatography (ethyl acetate/methanol, 1:0 to 1:1) gave the title compound (300 mg, 61.6%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether: M.p. 172.3°C. MS m/z (relative intensity, 70 eV) 289 (M⁺, 1), 274 (6), 87 (6), 86 (bp), 58 (5).

Example 12N-[2-(3-CHLORO-5-METHYLSULFONYL-PHENOXY)ETHYL]PROPAN-1-AMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-chloro-5-methylsulfonyl-benzene (0.5 g, 1.59 mmol) and propane-1-amine (1.04 ml, 12.76 mmol) in ethanol (5 ml). Yield: 368 mg (79.1%). The amine was converted to the hydrochloric acid salt and re-crystallized from ethanol/diethyl ether: M.p. 195-197°C. MS m/z (relative intensity, 70 eV) 291 (M⁺, 1), 264 (8), 262 (22), 73 (9), 72 (bp).

30

Example 13N-{2-[3-FLUORO-5-(METHYLSULFONYL)PHENOXY]ETHYL}-N-PROPYLAMINE D2

2-[3-Fluoro-5-(methylsulfonyl)phenoxy]ethanamine (0.3 g, 1.26 mmol), propyl 4-methylbenzenesulfonate D2 (1.04 ml, 12.76 mmol) and potassium carbonate (0.35 g, 2.52 mmol) in acetonitrile (10 ml). The mixture was heated under microwave radiation at 120°C for 45 min. The reaction mixtures were cooled to room temperature, filtrated and pooled before the volatiles were evaporated. Purification by flash column chromatography (ethyl acetate/methanol, 1:0 to 1:1) gave the title compound (0.15 g, 44%). The amine was converted to the hydrochloric acid salt and re-crystallized from methanol/diethyl ether. MS m/z (relative intensity, 70 eV) 277 (M⁺, 2), 248 (26), 138 (3), 94 (3), 74 (bp).

10

Example 14 N-(CYCLOPROPYLMETHYL)-N-{2-[3-FLUORO-5-(METHYLSULFONYL)-PHENOXY]ETHYL}AMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.05 g, 0.16 mmol) and aminomethylcyclopropane (0.11 ml, 1.3 mmol) in ethanol (5 ml). MS m/z (relative intensity, 70 eV) 287 (M⁺, 2), 246 (6), 138 (2), 84 (bp), 56 (14).

Example 15 N-{2-[3-FLUORO-5-(METHYLSULFONYL)PHENOXY]ETHYL}-N-ISOBUTYLAMINE

Preparation according to Example 1 but performed in one portion: 1-(2-bromoethoxy)-3-chloro-5-methylsulfonyl-benzene (0.05 g, 0.16 mmol) and isobutylamine (0.2 ml, 1.3 mmol) in ethanol (5 ml). MS m/z (relative intensity, 70 eV) 289 (M⁺, 1), 246 (bp), 138 (2), 86 (70), 56 (17).

Example 16 1-[2-(3-FLUORO-5-METHYLSULFONYL)-PHENOXY]ETHYL}AZETIDINE

1-(2-bromoethoxy)-3-fluoro-5-methylsulfonyl-benzene (0.5 g, 1.6 mmol), azetidine (0.23 ml, 3.2 mmol) and potassium carbonate (0.58 g, 4.2 mmol) was dissolved in acetonitrile (10 ml). The mixture was heated in a sealed container at 120°C for 2 h. The reaction mixture was cooled to room temperature, CH₂Cl₂ (100 ml) was added, solids was filtered off and the volatiles were evaporated. The amine was converted to the fumaric acid salt and re-crystallized from

methanol/diethyl ether. MS m/z (relative intensity, 70 eV) 273 (M⁺, 1), 94 (5), 82 (3), 71 (5), 70 (bp).

Example 17

5 1-(3-FLUORO-5-METHYLSULFONYL-PHENOXY)-N-PROPYL-PROPAN-2-AMINE

Sodium triacetoxyborohydride (1.16 g, 5.51 mmol) was added to a stirred mixture of 1-(3-fluoro-5-methylsulfonyl-phenoxy)propan-2-one (0.905 g, 3.67 mmol), propylamine (0.24 g, 4.04 mmol), acetic acid (0.5 g, 8.33 mmol) and
10 molecular sieves (2 g, 4Å, 4-8 mesh) in dry 1,2-dichloroethane (20 ml) the mixture was stirred at ambient temperature for 24h, the suspension was filtrated and sodium carbonate (100 ml, 10% aqueous solution) was added. The aqueous phase was extracted with dichloromethane (3 x 50 ml). The combined organic phase was dried with Na₂SO₄ and evaporated. Purification on flash
15 chromatography (ethyl acetate: methanol 1:0 to 4:1). 0.3 g, 28%. MS m/z (relative intensity, 70 eV) 289 (M⁺, 1), 260 (9), 152 (4), 87 (6), 86 (bp).

Example 18

2-(3-fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-1-amine

20 Toluenesulfonyl chloride (0.60 g, 3.1 mmol) was added to a stirred mixture of 4-dimethylaminopyridine (0.41 g, 3.4 mmol), triethylamine (0.53 g, 5.26 mmol) and a mixture of 1-(3-fluoro-5-methylsulfonyl-phenoxy)propan-2-ol and 2-(3-fluoro-5-methylsulfonyl-phenoxy)propan-1-ol (0.65 g, 2.63 mmol) in dry dichloromethane (20 ml). The resulting mixture was stirred at ambient temperature
25 for 5 h, HCl (100 ml, 5% aqueous solution) was added, the aqueous phase was extracted with dichloro-methane (2 x 50 ml), the combined organic phase was washed with Brine (50 ml) and sodium carbonate (50 ml, 5% aqueous solution).

The resulting oil was dissolved in ethanol (5 ml) and methanol (5 ml) and propylamine (3.24 ml, 39.51 mmol) was added, the resulting mixture was
30 heated to reflux for 15 h. The crude mixture was concentrated under vacuum and purified on flash chromatography (ethyl acetate: methanol 1:0 to 5:1). 0.52 g (as a mixture of 2-(3-fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-1-amine and 1-

(3-fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-2-amine), 68%. MS m/z (relative intensity, 70 eV) 289 (M⁺, 1), 260 (3), 73 (5), 72 (bp), 70 (9).

Preparations

5 Preparation 13-FLUORO-5-METHYLSULFONYL-PHENOL

Nitrogen was bubbled through a solution of 3-bromo-5-fluorophenol (10 g, 51.31 mmol) in dry dimethyl sulfoxide (70 ml) for 10 min after which sodium methanesulfinate (8.27 g, 76.96 mmol), copper(I)iodide (5.86 g, 30.79 mmol), L-proline (7.09 g, 61.57 mmol) and potassium carbonate (4.25 g, 30.79 mmol) was
10 added. The nitrogen flow was continued for an additional 10 min after which the mixture was heated at 100°C for 24 h. Ethyl acetate (100 ml) and water (100 ml) was added and the phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 100 ml), the combined organic phase was washed with aqueous lithium chloride (100 ml, 5%) and aqueous hydrochloric acid (100 ml,
15 5%), dried (Na₂SO₄) and evaporated. Purification by flash column chromatography (ethyl acetate/isooctane, 0:1 to 1:1) gave the title compound (7.17 g, 73%). MS m/z (relative intensity, 70 eV) 190 (M⁺, 81), 175 (27), 128 (50), 111 (bp), 83 (58).

Preparation 2

20 1-(2-BROMOETHOXY)-3-FLUORO-5-METHYLSULFONYL-BENZENE

A mixture of 3-fluoro-5-methylsulfonyl-phenol (4.2 g, 22.08 mmol), 1,2-dibromoethane (24 ml, 27.7 mmol) and potassium carbonate (6.1 g, 44.2 mmol) in acetonitrile (36 ml) was divided into 6 aliquots which were each heated under microwave radiation at 120°C for 30 min. The reaction mixtures were cooled to
25 room temperature, filtrated and pooled. The volatiles were evaporated, aqueous sodium carbonate (100 ml, 10%) was added and the mixture was extracted with ethyl acetate (2 x 100 ml). The combined organic phase was washed with brine (75 ml), dried (Na₂SO₄) and evaporated. Purification by flash column chromatography (ethyl acetate/isooctane, 0:1 to 1:0) gave the title compound (4.77
30 g, 73%). MS m/z (relative intensity, 70 eV) 298 (M⁺, 18), 296 (M⁺, 18), 109 (bp), 107 (99), 82 (15).

Preparation 3TERT-BUTYL 2-[3-FLUORO-5-(METHYLSULFONYL)PHENOXY]ETHYL-CARBAMATE

A mixture of triphenylphosphine (1.9 g, 7.3 mmol) in dry THF (20 ml) was flushed with N₂(g) and DEAD (3.1 ml, 6.9 mmol) was added drop-wise, 1,2-dibromoethane followed by 3-fluoro-5-methylsulfonyl-phenol (1.1 g, 6.1 mmol) and then boc-glycinol (1.0 ml, 6.1 mmol) in portions. The mixture was stirred at 70°C for 20 h, cooled to room temperature and water and EtOAc was added. The aqueous phase was extracted with ethyl acetate (2 x 100 ml). The combined organic phase was washed with aqueous NaOH (3 M, 50 ml) and brine (75 ml), dried (Na₂SO₄) and evaporated. Purification by flash column chromatography (ethyl acetate/isooctane, 0:1 to 1:0) gave the title compound (1.4 g, 70%). MS m/z (relative intensity, 70 eV 298 (M⁺, 18), 296 (M⁺, 18), 109 (bp), 107 (99), 82 (15).

15 Preparation 42-[3-FLUORO-5-(METHYLSULFONYL)PHENOXY]ETHANAMINE

A mixture of tert-butyl 2-[3-fluoro-5-(methylsulfonyl)phenoxy]ethyl-carbamate (1.4 g, 4.2 mmol) in EtOH (18 ml) was added HCl (1.25 M in EtOH, 6 ml). The mixture was stirred at ambient temperature for 20 h. The aqueous phase was made basic by addition of aqueous NaOH (1 M, 50 ml) and extracted with ethyl acetate (2 x 100 ml). The combined organic phase was washed with brine (75 ml), dried (Na₂SO₄) and evaporated to yield the title compound (0.77 g, 77%). MS m/z (relative intensity, 70 eV 298 (M⁺, 18), 296 (M⁺, 18), 109 (bp), 107 (99), 82 (15).

25

Preparation 53-CHLORO-5-METHYLSULFONYL-PHENOL

Preparation according to Preparation 1: 3-bromo-5-chlorophenol (4.0 g, 19.3 mmol), sodium methanesulfinate (3.1 g, 28.9 mmol), copper(I)iodide (2.2 g, 11.5 mmol), L-proline (2.7 g, 23.1 mmol), potassium carbonate (1.6 g, 11.6 mmol), dry dimethyl sulfoxide (70 ml). Yield: 3.7 g (92%). MS m/z (relative intensity, 70 eV) 206 (M⁺, 88), 191 (36), 144 (58), 127 (bp), 99 (76).

Preparation 61-(2-BROMOETHOXY)-3-CHLORO-5-METHYLSULFONYL-BENZENE

Preparation according to Preparation 2: 3-chloro-5-methylsulfonyl-phenol (2.8 g, 13.5 mmol), 1,2-dibromoethane (12 ml, 135 mmol), potassium carbonate (3.7 g, 27.1 mmol), acetonitrile (15 ml) Yield: 2.1 g, 50%). MS m/z (relative intensity, 70 eV 314 (M⁺, 35), 312 (M⁺, 25), 206 (7), 126 (9), 109 (98), 107 (bp).

10 Preparation 72-(3-fluoro-5-methylsulfonyl-phenoxy)propan-1-ol

A mixture of 1-bromo-2-propanol (70%) and 2-bromo-1-propanol (30%) (5.81 g, 41.8 mmol) was added to a solution of 3-fluoro-5-methylsulfonyl-phenol (1.59 g, 8.36 mmol) and potassium carbonate (2.37 g, 16.71 mmol) in dry dimethyl formamide (10ml), the mixture was heated to 120°C for 20 hours, the mixture was allowed to cool to ambient temperature and water (100ml) was added. The aqueous phase was extracted with ethyl acetate (3 x 100ml), the combined organic phase was washed with LiCl (5% aqueous solution, 4 x 50ml), brine (50 ml) and dried with Na₂SO₄ and evaporated. Purification flash chromatography (isooctane: ethyl acetate 1:0 to 3:2). 0.65 g (mixture of 1-(3-fluoro-5-methylsulfonyl-phenoxy)propan-2-ol and 2-(3-fluoro-5-methylsulfonyl-phenoxy)propan-1-ol), 31%. MS m/z (relative intensity, 70 eV) 248 (M⁺, 14), 204 (54), 203 (bp), 191 (37), 190 (44).

25 Preparation 81-(3-fluoro-5-methylsulfonyl-phenoxy)propan-2-one

1-Chloroacetone (95%, 2.66 g, 27.34 mmol) was added to a stirred solution of 3-fluoro-5-methylsulfonyl-phenol (80%, 1.3 g, 5.46 mmol) and potassium carbonate (2.26 g, 16.40 mmol) in dry dimethyl formamide (10 ml), the mixture was heated to 120°C for 20 minutes, the mixture was allowed to cool to ambient temperature and water (100 ml) was added. The aqueous phase was extracted with ethyl acetate (3 x 100 ml), the combined organic phase was washed

with LiCl (5% aqueous solution, 4 x 50 ml), brine (50 ml) and dried with Na₂SO₄ and evaporated. Purification on flash chromatography (isooctane: ethyl acetate 1:0 to 3:2). 0.905 g, 67%. MS m/z (relative intensity, 70 eV) 246 (M⁺, 79), 204 (55), 203 (bp), 141 (30), 94 (67).

5

Biological Activity

3,4-dihydroxyphenyl-acetic acid (DOPAC) in the striatum

The increased turnover of dopamine in the terminal areas of the ascending dopaminergic projections of the mammalian brain can be illustrated by measuring of changes in biochemical indices in the brain with the characteristic features of dopamine antagonists, e.g. producing increases in concentrations of dopamine metabolites such as 3,4-dihydroxyphenyl-acetic acid (DOPAC) in the striatum.

Test results are shown in Table 1.

15

Table 1

Estimated ED₅₀ values on increase of DOPAC (3,4-dihydroxyphenylacetic acid) in the rat striatum after systemic administration of test compound. For methods and statistical calculations see the below test methods.

20

Compound	ED ₅₀ DOPAC (μmol/kg)
Example 1	8.5 (7.4-10.3)
Example 2	4.2 (3.5 – 5.3)
Example 4	28.1 (24.5 – 28.1)
Example 8	36 (25.6 – 51.1)

Compound	ED ₅₀ DOPAC (μ mol/kg)
Example 10	14.5 (10.8 – 16.9)
Example 12	10.5 (7.7 – 15.6)

Effects on spontaneous locomotion

Prior art dopamine receptor antagonists are known to induce profound decrease in locomotor activity (catalepsy). Effects of a compound of the invention on spontaneous locomotion are shown in Table 2.

Table 2

Effects of a compound of the invention on Locomotor activity in drug-naive rats. The animals were placed in the motility meters immediately after drug administration and locomotor activity was recorded for 60 minutes (counts/60 min $\pm$ SEM).

Compound	Control group	3.7 μ mol/kg	11 μ mol/kg	33 μ mol/kg
Example 1	6414	8049	7325	6025
Example 2	14161	13941	8617	6325
Example 4	13890	11435	10025	12829
Example 8	13633	10319	12888	12872
Example 10	9816	11979	9344	8144

Compound	Control group	3.7 $\mu\text{mol/kg}$	11 $\mu\text{mol/kg}$	33 $\mu\text{mol/kg}$
Example 12	13630	14414	12740	7826

Amphetamine-induced hyper-locomotion

The increase in activity after treatment with d-amphetamine is a standard model of hyperdopaminergia. In this model, dopaminergic

- 5 neurotransmission is increased by systemic administration of d-amphetamine at a dose that is sufficiently high to produce a large increase in locomotor activity. The ability of a compound to antagonize this hyperactivity reflects anti-dopaminergic properties. Furthermore, antagonism of d-amphetamine induced hyperactivity is widely used as a standard assay of antipsychotic activity (see
- 10 *Psychopharmacology 4th Generation of progress* Chapter 68, p 793-795).

The effects of a compound of the invention on the increase in activity induced by direct or indirect dopaminergic agonists, i.e. d-amphetamine and congeners are shown in Table 3.

15 Table 3

Effect of compound of the present invention on reduction of amphetamine-induced hyper-locomotion. For methods and statistical calculations see the below test methods.

Compound	ED ₅₀ $\mu\text{mol/kg}$
Example 1	24 (18-30)
Example 4	69 (24 – 290)

Reduction of MK-801-induced hyper-locomotion

Another animal model of antipsychotic activity is based on administration of the glutamate antagonist MK-801. Glutamate antagonists (i.e. NMDA antagonists), can induce psychoses in man (see *Psychopharmacology, 4th Generation of progress* Chapter 101, p. 1205 and 1207) and induce behavioural
5 aberrations in animals. Thus, the ability of a drug to affect schizophrenia and psychotic states can be measured using behavioural models based on experimentally-induced hypoglutamatergic states. In this study the NMDA antagonist MK-801 (0.7 mg/kg i.p.) was used to create a hypoglutamatergic state where the rats display abnormal, hyperactive behaviour.

10 Test results for a compound of the present invention are seen in Table 4.

Table 4

Effect of compound of the present invention on reduction of MK-801-
15 induced hyper-locomotion (0.7 mg/kg i.p. 90 minutes before test compound). For methods and statistical calculations see the below test methods. The animals were placed in the motility meters immediately after test compound administration and locomotor activity was recorded between 45 and 60 minutes after administration (counts/15 min $\pm$ SEM)

20

Compound	ED ₅₀ μ mol/kg
Example 1	5.6 (0.3-18)
Example 8	40 (20 – 100)

Increase of Arc gene expression

It is known that the dopaminergic systems of the brain interact strongly with other transmitter systems (see *Psychopharmacology, 4th Generation of*
25 *progress*, Chapter 101, pages 1208-1209).

To investigate potential effects of compounds of the present invention on cortical and striatal NMDA receptor related synaptic signalling, Arc mRNA induction was assessed upon acute administration of a compound of the present invention. Arc (Arc/Arg3.1-activity regulated cytoskeleton-associated protein/activity-regulated gene 3.1; (*Link W et al.*; Proc Natl Acad Sci USA 1995 92 5734-5738; *Lyford GL et al.*; Neuron 1995 14 433-445)), is an immediate early gene (IEG), induced by synaptic activity, whose expression and localization at synaptic sites is triggered specifically by NMDA receptor activation and strongly related to neural plasticity (*Steward O, Worley PF*; Neuron 2001 30 227-240; *Takashi Kawashima et al.*; PNAS 2009 106 (1) 316-321; *Clive R. Bramham et al.*; Exp. Brain Res. 2010 200 125-140).

Test results for a compound of the invention are shown in Table 5.

Table 5

Estimated ED₅₀ values on increase of Arc gene expression in the rat striatum and cortex after systemic administration of test compound. For methods and statistical calculations see the below test methods.

Compound	ED ₅₀ Arc (μmol/kg)
Example 1 (striatum)	13 (5.6-77)
Example 1 (Cortex)	< 11

Test methods

The following tests are used for evaluation of the compounds according to the invention.

In vivo test: Behaviour

Behavioural activity is measured using eight Digiscan activity monitors (RXYZM (16) TAO, Omnitech Electronics, Columbus, OH, USA), connected to an Omnitech Digiscan analyzer and an Apple Macintosh computer equipped with a digital interface board (NB DIO-24, National Instruments, USA). Each activity monitor consists of a quadratic metal frame (W x L=40cm x 40cm) equipped with photobeam sensors. During measurements of behavioural activity, a rat is put in a transparent acrylic cage (WxLxH, 40x40x30 cm) which in turn is placed in the activity monitor. Each activity monitor is equipped with three rows of infrared photobeam sensors, each row consisting of 16 sensors. Two rows are placed along the front and the side of the floor of the cage, at a 90° angle, and the third row is placed 10 cm above the floor to measure vertical activity. Photobeam sensors are spaced 2.5 cm apart. Each activity monitor is fitted in an identical sound and light attenuating box containing a weak house light and a fan.

The computer software is written using object oriented programming (LabVIEW®, National instruments, Austin, TX, USA).

Behavioural data from each activity monitor, representing the position (horizontal center of gravity and vertical activity) of the animal at each time, are recorded at a sampling frequency of 25 Hz and collected using a custom written LABView™ application. The data from each recording session are stored and analyzed with respect to distance traveled. Each behavioural recording session lasts 60 min, starting approximately 4 min after the injection of test compound. Similar behavioural recording procedures are applied for drug-naïve and drug pre-treated rats. Rats pre-treated with d-amphetamine are given a dose of 1.5 mg/kg i.p. 10 min before the recording session in the activity monitor. Rats pre-treated with MK-801 are given a dose of 0.7 mg/kg i.p. 90 min before the recording session in the activity monitor. The results are presented as counts/60 minutes, or counts/30 minutes, in arbitrary length units. Statistical comparisons are carried out using Student's t-test against the control group. In MK-801 or amphetamine pre-treated animals, statistical comparisons are made against the MK801 or d-amphetamine controls, respectively.

ED₅₀ value for reduction of amphetamine-induced hyper-locomotion is calculated by curve fitting. The evaluation is based on 16 amphetamine pre-treated animals over the dose range 0, 11, 33 and 100 µmol/kg s.c. in one single experiment. Calculations are based on distance during the last 45 minutes of one hour of measurement. The distances are normalised to amphetamine-control and fitted by least square minimization to the function "End-(End-Control)/(1+(dose/ED₅₀)^{Slope})". The four parameters (Control, End, ED₅₀ and Slope) are fitted with the restrictions: ED₅₀>0, 0.5<Slope<3, End=0% of control. The restriction with locked End is made to focus on potency rather than efficacy. To estimate confidence levels for the parameters, the fit is repeated 100 times with a random evenly distributed squared weight (0 to 1) for every measurement value. Presented ED₅₀-ranges cover 95% of these values.

ED₅₀ value for reduction of MK-801-induced hyper-locomotion is calculated by curve fitting. The evaluation is based on 16 MK-801 pre-treated animals over the dose range 0, 11, 33 and 100 µmol/kg s.c. in one single experiment. Calculations are based on distance during the last 15 minutes of one hour of measurement. The distances are normalised to MK-801-control and fitted by least square minimization to the function "End-(End-Control)/(1+(dose/ED₅₀)^{Slope})". The four parameters (Control, End, ED₅₀ and Slope) are fitted with the restrictions: ED₅₀>0, 0.5<Slope<3, End=0% of control. The restriction with locked End is made to focus on potency rather than efficacy. To estimate confidence levels for the parameters, the fit is repeated 100 times with a random evenly distributed squared weight (0 to 1) for every measurement value. Presented ED₅₀-ranges cover 95% of these values.

25

In vivo test: Neurochemistry

After the behavioural activity sessions, the rats are decapitated and their brains rapidly taken out and put on an ice-cold petri-dish. The limbic forebrain, the striatum, the frontal cortex and the remaining hemispherical parts of each rat are dissected and frozen. Each brain part is subsequently analyzed with respect to its content of monoamines and their metabolites.

30

The monoamine transmitter substances (NA (noradrenaline), DA (dopamine), 5-HT (serotonin)) as well as their amine (NM (normethanephine), 3-MT (3-methoxytyramine)) and acid (DOPAC (3,4-dihydroxyphenylacetic acid), 5-HIAA (5-hydroxyindoleacetic acid), HVA (homovanillic acid)) metabolites are
5 quantified in brain tissue homogenates by HPLC separations and electrochemical detection

The analytical method is based on two chromatographic separations dedicated for amines or acids. Two chromatographic systems share a common auto injector with a 10-port valve and two sample loops for simultaneous injection
10 on the two systems. Both systems are equipped with a reverse phase column (Luna C18(2), dp 3 μ m, 50 x 2 mm i.d., Phenomenex) and electrochemical detection is accomplished at two potentials on glassy carbon electrodes (MF-1000, Bioanalytical Systems, Inc.). The column effluent is passed *via* a T-connection to the detection cell or to a waste outlet. This is accomplished by two solenoid
15 valves, which block either the waste or detector outlet. By preventing the chromatographic front from reaching the detector, better detection conditions are achieved. The aqueous mobile phase (0.4 ml/min) for the acid system contains citric acid 14 mM, sodium citrate 10 mM, MeOH 15% (v/v) and EDTA 0.1 mM. Detection potentials relative to Ag/AgCl reference are 0.45 and 0.60V. The
20 aqueous ion pairing mobile phase (0.5 ml/min) for the amine system contains citric acid 5 mM, sodium citrate 10 mM, MeOH 9%(v/v), MeCN 10.5% v/v), decane sulfonic acid 0.45 mM, and EDTA 0.1 mM. Detection potentials relative to Ag/AgCl reference are 0.45 and 0.65V.

ED₅₀ value for the increase of DOPAC in striatum is calculated by curve
25 fitting. The evaluation is based on 40 animals over the dose range 0, 3.7, 11, 33 and 100 μ mol/kg s.c. in two combined experiments. The DOPAC levels are normalised to control and fitted by least square minimization to the function "End-(End-Control)/(1+(dose/ED₅₀)^{Slope})". The four parameters (Control, End, ED₅₀ and Slope) are fitted with the restrictions: ED₅₀>0, 0.5<Slope<3, 350<End<400% of
30 control. To estimate confidence levels for the parameters, the fit is repeated 100 times with a random evenly distributed squared weight (0 to 1) for every measurement value. Presented ED₅₀-ranges cover 95% of these values.

In vivo test: Oral bioavailability

Experiments are performed 24 hours after implantation of arterial and venous catheters. Test compound is administered orally at 12.5 $\mu\text{mol/kg}$ or
5 intravenously at 5 $\mu\text{mol/kg}$ using the venous catheters, $n=3$ per group. Arterial blood samples are then taken during six hours at 0, 3, 9, 27, 60, 120, 180, 240, 300 and, 360 minutes after administration of the test compound. The oral bioavailability is calculated as the ratio of the AUC (Area under curve) obtained after oral administration over the AUC obtained after intravenous administration for
10 each rat. The parameter AUC is calculated according to the following:

AUC: the area under the plasma concentration versus time curve from time zero to the last concentration measured (Clast), calculated by the log/linear trapezoidal method.

The levels of test compound are measured by means of liquid
15 chromatography-mass spectrometry (LC-MS) (Hewlett-Packard 1100MSD Series). The LC-MS module includes a quaternary pump system, vacuum degasser, thermostatted autosampler, thermostatted column compartment, diode array detector and API-ES spray chamber. Data handling was performed with a HP ChemStation rev.A.06.03. system. Instrument settings:MSD mode: Selected ion
20 monitoring (SIM) MSD polarity: Positiv Gas temp: 350°C Drying gas: 13,0 l/min Nebulizer gas: 50 psig Capillary voltage: 5000 V Fragmentor voltage: 70 V.

Analytical column: Zorbax eclipse XDB-C8 (4.6 x 150 mm, 5 μm) at 20°C. The mobile phase is acetic acid (0.03%) (solvent A) and acetonitrile (solvent B). The flow rate of the mobile phase is 0.8 ml/min. The elution is starting at 12%
25 of solvent B isocratic for 4.5 min, then increasing linearity to 60% over 4.5 min.

Extractions procedure: Plasma samples (0.25-0.5 ml) are diluted with water to 1 ml, and 60 pmol (100 μl) internal standard (-)-OSU6241 is added. The pH was adjusted to 11 by the addition of 25 μl saturated Na_2CO_3 . After mixing, the samples are extracted with 4 ml dichloromethane by shaking for 20 min. The
30 organic layer is after centrifugation transferred to a smaller tube and evaporated to dryness under a stream of nitrogen. The residue is then dissolved in 120 μl mobile phase (acetic acid (0.03%): acetonitrile, 95:5) for LC-MS analysis (10 μl injected).

The selective ion (MH^+) is monitored for each example, and MH^+ 296 for (-)-OSU6241 ((3-[3-(ethylsulfonyl)phenyl]-1-propyl)piperidine).

A standard curve over the range of 1-500 pmol is prepared by adding appropriate amounts of test compound to blank plasma samples.

5

In vitro test: Metabolic stability in rat liver microsomes

Rat liver microsomes are isolated as described by Förlin [Förlin L: *Tox Appl Pharm.* 54 (3) 420-430, **1980**] with minor modifications e.g. 3 ml/g liver of a 0.1 M Na/K*PO₄ buffer with 0.15M KCl, pH 7.4, (buffer 1) is added before
10 homogenisation, the homogenate is centrifuged for 20 minutes instead of 15, the supernatant is ultracentrifuged at 100.000 g instead of 105.000 g and the pellet from the ultracentrifugation is resuspended in 1 ml/g liver of 20% v/v 87% glycerol in buffer 1.

1 μ L of, 0.2 or 1 mM test substance diluted in water, and 10 μ L 20
15 mg/ml rat liver microsome are mixed with 149 μ L 37°C buffer 1 and the reaction is started by addition of 40 μ L 4.1 mg/ml NADPH. After 0 or 15 minutes incubation at 37°C in a heating block (LAB-LINE, MULTI-BLOK Heater or lab4you, TS-100 Thermo shaker at 700 rpm) the reaction is stopped by addition of 100 μ L pure acetonitrile. The protein precipitation is then removed by rejecting the pellet after
20 centrifugation at 10.000 g for 10 minutes (Heraeus, Biofuge fresco) in 4°C. The test compound is analysed using HPLC-MS (Hewlett-Packard 1100MSD Series) with a Zorbax SB-C18 column (2.1 x 150 mm, 5 μ m) using 0.03% formic acid and acetonitrile as mobile phase (gradient) or a Zorbax Eclipse XDB-C18 (3 x 75 mm, 3.5 μ m) using 0.03% acetic acid and acetonitrile as mobile phase (gradient). The
25 15 min turnover is calculated as the fraction of test compound eliminated after 15 minutes, expressed in percent of 0 min levels, i.e. $100 \times [\text{conc test compound at 0 min} - \text{concentration at 15 min}] / \text{conc at 0 min}$.

Preparation of liver microsomes is performed as described in Förlin [Förlin L: *Tox Appl Pharm.* 54, (3) 420-430, **1980**]. Protocols for incubation with
30 liver microsomes are referred in Crespi et Stresser [Crespi C L, DM Stresser J. *Pharm. Tox. Meth.* 44 325-331, **2000**], and Renwick et al [Renwick AB et al.; *Xenobiotica* 2001 31 (4) 187-204].

Microdialysis

Male Sprague-Dawley rats weighing 220-320g are used throughout the experiments. Before the experiment the animals are group housed, five animals in
5 each cage, with free access to water and food. The animals are housed at least one week after arrival prior to surgery and use in the experiments. Each rat is used only once for microdialysis.

We use a modified version of *Waters et al.* [*Waters et al.*; J. Neural. Transm. Gen. Sect. 1994 98 (1) 39-55] of the I-shaped probe as described by
10 *Santiago and Westerink* [*Santiago M, Westerink BHC*; Naunyn-Schmiedeberg's Arch. Pharmacol. 1990 342 407-414]. The dialysis membrane we use is the AN69 polyacrylonitrile/ sodiummethalylsulfonate copolymer (HOSPAL; o.d./i.d. 310/220 µm; Gambro, Lund, Sweden). In the dorsal striatum we use probes with an exposed length of 3 mm of dialysis membrane and in the prefrontal cortex the
15 corresponding length is 2.5 mm. The rats are operated under isoflurane inhalationanesthesia while mounted into a Kopf stereotaxic instrument. Coordinates are calculated relative to bregma; dorsal striatum AP +1, ML $\pm$ 2.6, DV -6.3; Pf cortex, AP +3.2, 8° ML $\pm$ 1.2, DV -4,0 according to *Paxinos and Watson* [*Paxinos G, Watson C: The Rat Brain in Stereotaxic Coordinates*; New York,
20 Academic Press 1986]. The dialysis probe is positioned in a burr hole under stereotaxic guidance and cemented with phosphatine dental cement.

The rats are housed individually in cages for 48 h before the dialysis experiments, allowing them to recover from surgery and minimizing the risk of drug interactions with the anaesthetic during the following experiments. During this
25 period the rats have free access to food and water. On the day of experiment the rats are connected to a micro perfusion pump via a swivel and are replaced in the cage where they can move freely within its confinements. The perfusion medium is a Ringer's solution containing in mmol/l: NaCl; 140, CaCl₂; 1.2, KCl; 3.0, MgCl₂; 1.0 and ascorbic acid; 0.04 according to *Moghaddam and Bunney* [*Moghaddam B, Bunney BS*; J. Neurochem. 1989 53 652-654]. The pump is set to a perfusion
30 speed of 2 µl/min and 40 µl samples are collected every 20 min.

Each sample is analyzed at two HPLC systems. On an autoinjector (CMA 200) with a 10-port valve (Valco C10WE), holding two sample loops in series (4 µl and 20 µl), each brain dialysate sample is loaded in both loops simultaneously. At injection the 20 µl sample is introduced into a column switching system (reverse-phase combined with reverse-phase ion-pairing) for dopamine (DA), noradrenaline (NA), normetanephrine (NM), 3-methoxytyramine (3-MT) and serotonin (5-hydroxytryptamine, 5-HT) determination, while the 4 µl sample is introduced on a reverse-phase column for the chromatography of the acidic monoamine metabolites 3,4-di-hydroxyphenylacetic acid (DOPAC), homovanillic acid (HVA) and 5-hydroxyindoleacetic acid (5-HIAA). The currents generated by the two EC detectors are converted to digital data and evaluated using Chromeleon software (Dionex) on a PC. The method sample turn over time is 4.5 min and two parallel experiments are normally analyzed simultaneously on the system. After the experiment the rats are uncoupled from the perfusion pump and decapitated. Their brains are rapidly taken out and fixed in Neo-fix solution (Kebo-lab, Sweden) for subsequent inspection of probe localisation. The Animal Ethics Committee in Göteborg, Sweden approved the procedures applied in these experiments.

20 PCR

Total RNA is prepared by the guanidin isothiocyanate method reported by *Chomczynski & Sacchi* [*Chomczynski P & Sacchi N; Anal. Biochem.* 1987 **162** 156-159]. RNA pellets are solved in MQ water and stored at -80°C. The sample concentration was determined spectrophotometrically by a NanoDrop ND-1000. A quality indicator number and an integrity number of r-RNA were measured with an Experion (Bio-Rad) on random samples.

Reversed transcription was performed by using a SuperScript III kit (Invitrogen). 1 µg of total RNA was reverse transcribed with 5 µl 2 x RT Reaction Mix, 1 µl RT Enzyme Mix, volume adjusted to 10 µl with DEPC-treated water. 1U of E.coli RNase H was added. cDNA was diluted 40 times and stored at -20°C.

For real-time PCR measurements, 0.7 µl of the cDNA reaction was amplified in a 25 µl reaction mixture containing 1 x PCR buffer, 0.2 mM dNTP, 3.7

mM MgCl₂, 0.15 mM SYBR green, 0.4 µM of each primer, and 1U JumpStart Taq DNA polymerase. Real-time PCR was measured on CFX96 (Biorad) using the following settings for all genes, 60 s pre-incubation at 95°C, followed by 40 cycles of denaturation at 95°C for 10 s, annealing at 56°C for 10 s, and elongation at
 5 72°C for 10 s.

The primer sequences were as follows:

Hypoxanthine phosphoribosyl transferase (HPRT) (Accession Number AF001282)

Sense: 5'-GGC CAG ACT TGT TGG ATT TG-3'

10 Antisense; 5'-CCG CTG TCT TTT AGG CTT TG-3'

Cyclophilin A (Accession Number M19533)

Sense: 5'-GTC TCT TTT CGC CGC TTG CT-3'

Antisense: 5'-TCT GCT GTC TTT GGA ACT TTG TCT G-3'

15

Activity-regulated gene (Arc) (Accession Number U19866)

Sense: 5'- GTC CCA GAT CCA GAA CCA CA-3'

Antisense: 5'- CCT CCT CAG CGT CCA CAT AC-3'

20 Initial DNA amounts were quantified by a standard curve constructed for every gene using 6 serial 4-fold dilutions of purified PCR products. Correct PCR products were confirmed by agarose gel electrophoresis (2%) PCR products were purified with PCR purification kit from Qiagen (Valencia, CA, USA) All genes were sequenced at MWG, Germany, and routinely by melting curve analysis.

25 Arc gene amounts were normalised using the geometric mean of the amounts of the two house-keeping genes assessed (HPRT and cyclophilin A).


ED₅₀ value for the increase of Arc in striatum is calculated by curve fitting. The evaluation is based on 20 animals over the dose range 0, 11, 33 and 100 µmol/kg s.c. in a single experiments. The Arc levels are normalised to control
 30 and fitted by least square minimization to the function "End-(End-Control)/(1+(dose/ED₅₀)^{Slope})". The four parameters (Control, End, ED₅₀ and Slope) are fitted with the restrictions: ED₅₀>0, 0.5<Slope<3, 300<End<600% of control.

To estimate confidence levels for the parameters, the fit is repeated 100 times with a random evenly distributed squared weight (0 to 1) for every measurement value. Presented ED₅₀-ranges cover 95% of these values.

ED₅₀ value for the increase of Arc in (prefrontal) cortex is evaluated to
5 be clearly below the lowest dose.

CLAIMS

1. A phenoxy-ethyl-amine derivative of Formula 1:



5 a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein

R^1 represents CH_3 or CF_3 ;

R^2 is selected from the group consisting of C_1 - C_4 -alkyl, allyl,
 10 $CH_2CH_2OCH_3$, $C(CH_3)_2CH_2CH_3$, CH_2 -cyclopropyl, cyclobutyl, cyclopentyl,
 $CH_2CH_2CH_2F$, $CH_2CH_2CHF_2$, CH_2CH_2F , 3,3,3-trifluoropropyl and 4,4,4-trifluorobutyl; and

R^3 is selected from the group consisting of H, CH_3 and CH_2CH_3 ; or

R^2 and R^3 together form $CH_2(CH_2)CH_2$ or $CH_2(CH_2)_3CH_2$;

15 R^4 represents F or Cl; and

R' and R'' independently represent hydrogen or methyl.

2. The phenoxy-ethyl-amine derivative according to claim 1, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or
 20 partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R^1 represents CH_3 or CF_3 .

3. The phenoxy-ethyl-amine derivative according to either one of claims 1-2, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a
 25 fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from the group consisting of C_1 - C_4 -alkyl, allyl,

CH₂CH₂OCH₃, C(CH₃)₂CH₂CH₃, CH₂-cyclopropyl, cyclobutyl, cyclopentyl, CH₂CH₂CH₂F, CH₂CH₂CHF₂, CH₂CH₂F, 3,3,3-trifluoropropyl and 4,4,4-trifluorobutyl.

5 4. The phenoxy-ethyl-amine derivative according to any one of claims 1-3, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R³ is selected from the group consisting of H and CH₃.

10 5. The phenoxy-ethyl-amine derivative according to either one of claims 1-2, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R² and R³ together form CH₂(CH₂)CH₂ or CH₂(CH₂)₃CH₂.

15 6. The phenoxy-ethyl-amine derivative according to any one of claims 1-5, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R⁴ represents F or Cl.

20 7. The phenoxy-ethyl-amine derivative according to any one of claims 1-5, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, wherein R' and R'' independently represent hydrogen or methyl.

25 8. The phenoxy-ethyl-amine derivative according to claim 1, which is
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]propan-1-amine;
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]butan-1-amine;
N-Ethyl-2-(3-fluoro-5-methylsulfonyl-phenony)ethanamine;
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]-N-methyl-propan-1-
30 amine;

N-[2-(3-Chloro-5-methylsulfonyl-phenoxy)ethyl]propan-1-amine;
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]butan-2-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]cyclopentamine;
N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]-2-methyl-butan-2-amine;

N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]cyclobutamine;
5 N-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]propan-2-amine;
1-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]piperidine;
N,N-Diethyl-2-(3-fluoro-5-methylsulfonyl-phenoxy)ethanamine;
N-1,1-di-deuterium-propyl-[2-(3-fluoro-5-methanesulfonyl-phenoxy)-ethyl]-amine;

10 N-(Cyclopropylmethyl)-N-{2-[3-fluoro-5-(methylsulfonyl)-phenoxy]ethyl}amine;

N-{2-[3-Fluoro-5-(methylsulfonyl)phenoxy]ethyl}-N-isobutylamine;
1-[2-(3-Fluoro-5-methylsulfonyl-phenoxy)ethyl]azetidine;
1-(3-Fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-2-amine; or
15 2-(3-Fluoro-5-methylsulfonyl-phenoxy)-N-propyl-propan-1-amine;
a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof,
or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition, comprising a therapeutically effective
20 amount of a phenoxy-ethyl-amine derivative of any one of claims 1-8, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, together with at least one pharmaceutically acceptable carrier or diluent.

25 10. The phenoxy-ethyl-amine derivative of any one of claims 1-8, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof, for use as a medicament.

30 11. Use of a phenoxy-ethyl-amine derivative of any one of claims 1-8, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof,

for the manufacture of a medicament for the treatment, prevention or alleviation of a disease or a disorder or a condition of a mammal, including a human, which disease, disorder or condition is responsive to modulation of dopaminergic function in the central nervous system.

5

12. The use according to claim 11, wherein the disease, disorder or condition is selected from the group of psychosis, schizophrenia, schizophreniform disorder, bipolar disorder, psychotic disorder, drug induced psychotic disorder, mood disorder, anxiety disorder, depression, obsessive-compulsive disease, dementia, age-related cognitive impairment, Autism spectrum disorders, ADHD, cerebral Palsy, Gilles de la Tourette's syndrome, brain injury, sleep disorder, sexual disorder, eating disorder, obesity, headache, pains in conditions characterized by increased muscular tone, Parkinson's disease, parkinsonian syndrome, dyskinesia, L-DOPA induced dyskinesias, Tardive dyskinesias, dystonia, tics and tremor dementia, Huntington's disease, movement disorder induced by drugs, restless legs, narcolepsy, Alzheimer's disease. and disorder related to Alzheimer's disease.

13. The use according to claim 11, wherein the disease, disorder or condition is selected from the group of schizophrenia, L-DOPA induced dyskinesias and Huntington's disease.

14. A method of treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disorder, disease or condition is responsive to modulation of dopaminergic and glutamatergic function in the central nervous system, which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of a phenoxy-ethyl-amine derivative of any one of claims 1-8, a stereoisomer or a mixture of its stereoisomers, or an N-oxide thereof, or a fully or partially deuterated analog thereof, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2012/056959

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07C317/22 C07D205/04 C07D295/088 A61K31/138 A61K31/145
 A61K31/397 A61K31/4453 A61P25/14 A61P25/16 A61P25/18

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2007/063789 A1 (KISSEI PHARMACEUTICAL [JP]; ISHIKAWA TAKEHIRO [JP]; NISHIMURA TOSHIHIR) 7 June 2007 (2007-06-07) the whole document -----	1-14
A	EP 0 363 782 A1 (ERBA FARMITALIA [IT] ERBA CARLO SPA [IT]) 18 April 1990 (1990-04-18) Page 5, line 53; page 10, lines 55-58. -----	1-14
A	WO 02/00602 A1 (PFIZER PROD INC [US]; LOWE JOHN ADAMS III [US]) 3 January 2002 (2002-01-03) page 1, line 3 - page 2, line 4; claims 1,5,6 ----- -/-	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 August 2012

Date of mailing of the international search report

20/09/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Cooper, Simon

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/056959

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	RAUBO P ET AL: "Aminoalkyl phenyl sulfones-a novel series of 5-HT7 receptor ligands", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 16, no. 5, 1 March 2006 (2006-03-01), pages 1255-1258, XP027965701, ISSN: 0960-894X [retrieved on 2006-03-01] the whole document	1-14
A	----- WO 2007/072041 A1 (ASTEX THERAPEUTICS LTD [GB]; CONGREVE MILES STUART [GB]; CALLAGHAN OWE) 28 June 2007 (2007-06-28) claim 1 -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/056959

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007063789 A1	07-06-2007	TW 200800871 A	01-01-2008
		WO 2007063789 A1	07-06-2007
EP 0363782 A1	18-04-1990	AU 629258 B2	01-10-1992
		AU 4507989 A	01-05-1990
		CN 1041755 A	02-05-1990
		CZ 8905653 A3	13-07-1994
		DE 68912055 D1	17-02-1994
		DE 68912055 T2	28-04-1994
		DK 137490 A	30-07-1990
		EP 0363782 A1	18-04-1990
		EP 0389618 A1	03-10-1990
		ES 2061868 T3	16-12-1994
		IE 893171 L	05-04-1990
		IL 91541 A	27-02-1994
		JP H03501862 A	25-04-1991
		NZ 230628 A	25-06-1991
		PH 27293 A	04-05-1993
		PT 91880 A	30-04-1990
		RU 2007386 C1	15-02-1994
		US 5272144 A	21-12-1993
		WO 9003965 A1	19-04-1990
		YU 191489 A	28-02-1991
		ZA 8907590 A	26-06-1991
WO 0200602 A1	03-01-2002	AR 033828 A1	07-01-2004
		AT 281427 T	15-11-2004
		AU 7440701 A	08-01-2002
		DE 60106914 D1	09-12-2004
		DE 60106914 T2	08-12-2005
		DK 1294677 T3	07-02-2005
		DO P2001000189 A	30-03-2002
		EP 1294677 A1	26-03-2003
		ES 2230326 T3	01-05-2005
		JP 3801561 B2	26-07-2006
		JP 2004501892 A	22-01-2004
		PA 8519901 A1	30-06-2003
		PE 01212002 A1	16-02-2002
		PT 1294677 E	31-01-2005
		US 2002052401 A1	02-05-2002
		UY 26802 A1	31-01-2002
		WO 0200602 A1	03-01-2002
WO 2007072041 A1	28-06-2007	NONE	