



HU000028239T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 028 239**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 06 252751**(51) Int. Cl.: **C12N 50/73** (2006.01)(22) A bejelentés napja: **2006. 05. 26.**

(96) Az európai bejelentés bejelentési száma:

EP 20060252751

(97) Az európai bejelentés közzétételi adatai:

EP 1780267 A2 **2007. 05. 02.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 1780267 B1 **2016. 01. 13.**

(30) Elsőbbségi adatok:

685607 P **2005. 05. 27.** **US****743821 P** **2006. 03. 27.** **US**

(73) Jogosult(ak):

LifeScan, Inc., Milpitas, CA 95035 (US)

(72) Feltaláló(k):

Rezania, Alireza, Hillsborough, NJ 08844 (US)**Xu, Jean, Hillsborough, NJ 08844 (US)**

(74) Képviselő:

Danubia Szabadalmi és Jogi Iroda Kft.,**Budapest**

(54)

Amnionfolyadék-eredetű sejtek

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 1 780 267 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
13.01.2016 Bulletin 2016/02

(51) Int Cl.:
C12N 5/073 (2010.01)

(21) Application number: **06252751.0**

(22) Date of filing: **26.05.2006**

(54) **Amniotic fluid derived cells**

Zellen aus Fruchtwasser

Cellules isolés de liquide amniotique

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

(30) Priority: **27.05.2005 US 685607 P**
27.03.2006 US 743821 P

(43) Date of publication of application:
02.05.2007 Bulletin 2007/18

(60) Divisional application:
10184689.7 / 2 302 036

(73) Proprietor: **LifeScan, Inc.**
Milpitas, CA 95035 (US)

(72) Inventors:
• **Rezania, Alireza**
Hillsborough, NJ 08844 (US)
• **Xu, Jean**
Hillsborough, NJ 08844 (US)

(74) Representative: **Wise, Daniel Joseph**
Carpmaels & Ransford LLP
One Southampton Row
London WC1B 5HA (GB)

(56) References cited:
EP-A1- 1 391 505 WO-A-2005/017117

- **WEI JUN PING ET AL: "Human amnion-isolated cells normalize blood glucose in streptozotocin-induced diabetic mice." CELL TRANSPLANTATION, vol. 12, no. 5, 2003, pages 545-552, XP009085264 ISSN: 0963-6897**
- **CHEN LI-BO ET AL: "Differentiation of rat marrow mesenchymal stem cells into pancreatic islet beta-cells" WORLD JOURNAL OF GASTROENTEROLOGY, XX, XX, vol. 10, no. 20, 15 October 2004 (2004-10-15), pages 3016-3020, XP002363979 ISSN: 1007-9327**
- **HELD K R ET AL: "THE EFFECT OF OXYGEN TENSION ON COLONY FORMATION AND CELL PROLIFERATION OF AMNIOTIC FLUID CELLS IN-VITRO" PRENATAL DIAGNOSIS, vol. 4, no. 3, 1984, pages 171-180, XP009085380 ISSN: 0197-3851**
- **TSAI MING-SONG ET AL: "Isolation of human multipotent mesenchymal stem cells from second-trimester amniotic fluid using a novel two-stage culture protocol" HUMAN REPRODUCTION, IRL PRESS, OXFORD, GB, vol. 19, no. 6, June 2004 (2004-06), pages 1450-1456, XP002404972 ISSN: 0268-1161**
- **IN 'T ANKER PIETERNELLA S ET AL: "Amniotic fluid as a novel source of mesenchymal stem cells for therapeutic transplantation." BLOOD 15 AUG 2003, vol. 102, no. 4, 15 August 2003 (2003-08-15), pages 1548-1549, XP009085337 ISSN: 0006-4971**
- **FAUZA D: "Amniotic fluid and placental stem cells" BAILLIERE'S BEST PRACTICE AND RESEARCH. CLINICAL OBSTETRICS AND GYNAECOLOGY, BAILLIERE TINDALL, LONDON, GB, vol. 18, no. 6, December 2004 (2004-12), pages 877-891, XP004676339 ISSN: 1521-6934**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 780 267 B1

Description

FIELD OF THE INVENTION

5 [0001] This invention relates to an expandable population of amniotic fluid-derived cells that can be differentiated into a β -cell lineage. Also disclosed are methods for isolating and expanding such amniotic fluid-derived cells, as well as related methods and compositions for utilizing such cells in the therapeutic treatment of diabetes.

BACKGROUND

10 [0002] Loss of organ function can result from congenital defects, injury or disease. One example of a disease causing loss of organ function is diabetes mellitus, or diabetes. Most cases of diabetes fall into two clinical types: Type 1, also known as juvenile-onset diabetes, or insulin dependent diabetes mellitus (IDDM), and Type 2, also known as adult-onset diabetes. Each type has a different prognosis, treatment, and cause. Both types are characterized by the patient's
15 inability to regulate their blood glucose levels. As a consequence, blood glucose levels rise to high values because glucose cannot enter cells to meet metabolic demands. This inability to properly metabolize blood sugar causes a complex series of early and late-stage symptomologies, beginning with, for example, hyperglycemia, abnormal hunger, thirst, polyuria, and glycosuria, and then escalating to, for example, neuropathy, macro-vascular disease, and micro-vascular disease.

20 [0003] A common method of treatment of Type 1 diabetes involves the exogenous administration of insulin, typically by injection with either a syringe or a pump. This method does not completely normalize blood glucose levels and is often associated with an increased risk of hypoglycemia. More effective glycemic control can be achieved if the function of the pancreas can be restored or rejuvenated via transplantation or cell-based therapies.

25 [0004] There are many transplantation therapies currently used to treat diabetes: One such treatment involves transplanting isolated islets of Langerhans into the diabetic patient. One of the main hurdles to human islet transplantation has been the lack of sufficient number of islets to treat the large number of diabetic patients. One possible solution to the shortage of islets is the generation of islets from alternate cellular sources.

30 [0005] It has been documented that progenitor cells derived from adult tissues are capable of differentiation into a pancreatic β -cell phenotype. See, for example, WO2004/087885 A2, Hess et al. (Nature Biotechnology 21, 763 - 770, 2003), and Janus et al. (J. Clin. Invest. 111: 843-850, 2003), which report the capacity of adult bone marrow-derived cells (mesenchymal and hematopoietic cells) to differentiate into cells having characteristics of a pancreatic β -cell *in vitro*, or secrete trophic factors that help regenerate a damaged pancreas *in vivo*.

[0006] Among other sources of progenitor cells that can be differentiated into pancreatic cells include rodent liver oval stem cells (WO03/033697) and post-partum placenta (U.S. Published Application 2004/0161419 A1).

35 [0007] The endocrine cells of the islets of Langerhans, including β -cells, are constantly turning over by processes of apoptosis and the proliferation of new islet cells (neogenesis). As such, the pancreas is thought to be a source of progenitor cells that are capable of differentiating into pancreatic hormone producing cells. There are three distinct tissue types, isolated from a pancreas, that are a potential source of pancreatic progenitor cells: an islet rich fraction, a ductal cell rich fraction, and an acinar cell rich fraction.

40 [0008] Isolation of progenitor cells or partially differentiated cells from crude pancreatic tissue extracts may be achieved using antibodies raised against cell surface markers. For example, U.S. Published Application 2004/0241761 discloses isolation of murine cells that expressed ErbB2, ErbB3, ErbB4, Msx-2, PDX-1 and insulin.

[0009] Gershengorn et al. (Science 306: 2261-2264, 2004) teach the production of proliferating cells that were able to form islet-like cell aggregates. The cells were derived from a heterogeneous population of adherent cells that emerged
45 from the culture of isolated human pancreatic islets *in vitro*. The isolated islets of Langerhans were initially seeded onto tissue culture dishes and cultured in medium containing 10% serum. Fibroblast-like cells were observed to migrate out of the cultured islets and form a monolayer. These cells expressed Nestin, smooth muscle actin and vimentin.

[0010] Pancreatic progenitor cells may also arise from the culture of pancreatic islet and ductal tissue that has been dissociated into single cells, as disclosed by Seaberg et al. (Nature Biotechnology 22: 1115 - 1124, 2004). The murine progenitor cells disclosed by Seaberg *et al.* expressed Nestin during proliferation.

50 [0011] U.S. Published Application 2003/0082155 discloses methods to isolate and identify a population of cells from the islets of Langerhans of human pancreas, which have the functional and molecular characteristics of stem cells. In particular, these cells were characterized by Nestin-positive staining, Nestin gene expression, GLP-1R-positive staining, GLP-1R gene expression, ABCG2 positive staining, ABCG2 gene expression, Oct3/4 positive staining, Oct3/4 gene
55 expression, latrophilin (type 2) positive staining, latrophilin (type 2) gene expression, Hes-1 positive staining, Hes-1 gene expression, Integrin subunits α 6 and β 1 positive staining, Integrin subunits α 6 and β 1 gene expression, c-kit positive staining, c-kit gene expression, MDR-1 positive staining, MDR-1 gene expression, SST-R, 2, 3, 4 positive staining, SST-R, 2, 3, 4 gene expression, SUR-1 positive staining, SUR-1 gene expression, Kir 6.2 positive staining, Kir 6.2 gene

expression, CD34 negative staining, CD45 negative staining, CD133 negative staining, MHC class I negative staining, MHC class II negative staining, cytokeratin-19 negative staining, long-term proliferation in culture, and the ability to differentiate into pseudo-islets in culture.

[0012] In another approach, as disclosed in U.S. Patent 5,834,308, U.S. Patent 6,001,647 and U.S. Patent 6,703,017, crude preparations of islet cultures from NOD mice may be used to establish epithelial-like cultures, which can be maintained in growing cultures for greater than 1 year and which appear to demonstrate the ability to differentiate into islet-like clusters, capable of secreting insulin.

[0013] Islet-like structures may be generated from fractions of digested human pancreata enriched for ductal tissue, as disclosed in Bonner-Weir et al. (Proc Nat Acad Sci 97: 7999-8004, 2000) and U.S. Patent 6,815,203 B1. Islet-like clusters disclosed in these publications stained positive for cytokeratin-19 and showed immunoreactivity for insulin.

[0014] WO2004/011621 discloses the generation of insulin negative adherent cells from human pancreatic ductal fragments.

[0015] WO03/102134 discloses the generation of an epithelial cell positive for cytokeratin-19 from an acinar fraction of a human pancreatic digest. The cells generated are capable of limited expansion and differentiate into an insulin-producing cell in the presence of an induction media.

[0016] U.S. Published Application 2004/015805 A1 reports that a subset of human pancreatic stem cells may be isolated using ligands to the cell surface marker CD56 (also known as NCAM). These cells can differentiate into insulin producing cells and insulin producing aggregates.

[0017] It has been documented that progenitor cells, derived from fetal or embryonic tissues, have the potential to differentiate into a pancreatic hormone-producing cell. See, for example, U.S. Patent 6,436,704, WO03/062405, WO02/092756 and EP 0 363 125 A2, which report the potential of human fetal and embryonic derived cells to differentiate into a β -cell lineage.

[0018] Human Embryonic Stem cells (hES) are derived from the inner cell mass of the blastocyst, the earliest stage of embryonic development of the fertilized egg. The blastocyst is a pre-implantation stage of the embryo, a stage before the embryo would implant in the uterine wall. When cultured on an inactivated feeder layer of cells according to conditions described by Thompson and colleagues (Thomson, et al. (Proc. Natl. Acad. Sci. U.S.A. 92: 7844-7848, 1995); Thomson, et al. (Science 282:1145-1147, 1998), Marshall, et al., (Methods Mol. Biol. 158:11-18, 2001), the inner layer cells of the blastocyst may be grown *in vitro* indefinitely in an undifferentiated state. Properly propagated hES cells have unlimited potential to double while maintaining their pluripotency; namely their capacity of differentiating into the three layers of the embryo, Ectoderm (Ec), Mesoderm (Me) and Endoderm (En). When grown as pluripotent hES, the cells maintain a euploid karyotype and are not prone to senescence.

[0019] Human embryonic stem cells display a distinct group of cell surface antigens, SSEA-3, SSEA-4, TRA-2-54 (alkaline phosphatase), TRA-1-60 and TRA-1-81, in addition to expressing specific transcription factors OCT-4, NANOG, SOX-2, FGF-4 and REX-1 (Henderson, et al., (Stem Cells 20:329-337, 2002), Draper, et al., (J. Anat. 200:249-258, 2002), Mitsui et al., (Cell 113:631-642, 2003), Chambers et al., (Cell 113:643-655, 2003).

[0020] It is important to note from these publications, however, that human embryonic cells often require a feeder layer for expansion and maintenance of pluripotency or combination of a complex extracellular matrix, such as, for example, MATRIGEL™, plus conditioned media. These conditions do not allow the facile scale up of cells and an eventual cell therapy for treating diabetes.

[0021] Researchers have found that non-embryonic types of stem cells ("adult stem cells") are not as capable of differentiating into many different tissue types, as are embryonic stem cells, so embryonic stem cells still have many advantages over the use of adult stem cells. However, one obstacle with the isolation of embryonic stem cells is that the cells are derived from embryos at the "blastocyst" stage. Human embryonic stem cell research is encumbered by an emotionally charged political and ethics debate and is likely to remain so for years to come.

[0022] Additionally, human embryonic stem cells (hES) have been found to be tumorigenic when injected into immunologically impaired animals, i.e. in the context of post-natal tissues, whereas adult stem cells are not. The tumorigenic attributes of hES cells are not frequently addressed, though this issue may burden their use in replacement cell therapy in the future. The political, moral and ethical issues around hES cells and their tumorigenic properties, as well as the perceived difficulties of expanding undifferentiated adult stem cells in culture, while maintaining a genetically normal genome, are major barriers in the development of human cell replacement therapy.

[0023] Pluripotent or multipotent stem cells have been isolated from chorionic villus, and amniotic fluid. Many amniotic and placental cells share a common origin, namely the inner cell mass of the morula, which gives rise to the embryo itself, the yolk sac, the mesenchymal core of the chorionic villi, the chorion and the amnion (Crane & Cheung, Prenatal Diagnosis 8: 119-129, 1988). Embryonic and fetal cells from all three germ layers have long been identified in the amniotic fluid (Milunsky, Genetic Disorder of the Fetus. New York: Plenum Press, 75-84, 1979; Hoehn & Salk, Methods in Cell Biology 26, 11-34, 1982; Gosden, British Medical Bulletin 39, 348-354, 1983; Prusa et al, Human Reproduction 18, 1489-1493, 2003). Thus, amniotic fluid may provide the least invasive access to embryonic-like and fetal-like stem cells.

[0024] Amniotic fluid derived cells have been routinely used for detecting chromosomal abnormality of the fetus.

Amniotic fluid is typically sampled during the 2nd trimester (16 to 22 weeks of gestation). Previous art clearly demonstrates presence of three subpopulation with distinct cell morphologies: "fibroblastic" (F), "amniotic fluid" (AF) cells, and "epithelial" (E) cells. The F and AF cells rapidly expand whereas the E cells display a much slower growth curve and have poor clonal efficiency.

5 **[0025]** For example, PCT application WO2003/042405 discloses isolation of c-Kit positive stem cells from chorionic villus, amniotic fluid and placenta (Cell 1, **Table I**).

[0026] In another example, U.S. Published Application 2005/0054093 discloses the isolation of stem cells from amniotic fluid. These cells express stage-specific embryonic antigen 3 (SSEA3), stage-specific embryonic antigen 4 (SSEA4),

10 **[0027]** In another example, fetal cells have been isolated from amniotic fluid (in't Anker et al, Blood 102, 1548-1549, 2003). The cells disclosed were positive for expression of the following markers: CD44, CD73, CD90, CD105, CD106, HLA-A,B, & C. The cells were negative for expression of the following markers: c-Kit (CD117), CD11, CD31, CD34, CD45 and HLA-D (Cell 3, **Table I**).

[0028] A population of mesenchymal stem cells isolated from amniotic fluid has also been reported in a publication to Tsai *et al* (Tsai et al, Human Reproduction 19, 1450-1456, 2004). The cells disclosed were positive for expression of the following markers: CD29, CD44, CD73, CD90, HLA-A,B, & C. The cells were also positive for the embryonic trans-

20 **[0029]** Although recent publications and patents have suggested that within the fibroblastic, amniotic fluid, or epithelial subpopulations there exists a cell population that display some characteristics of human embryonic cells, such as expression of surface markers SSEA3 and -4, expression of transcription factor Oct-4, strong expansion potential, and differentiation into multiple cell types; none of the previously published art has demonstrated the existence of a subpopulation of the cells that display expression of key early endodermal markers, such as HNF-1 beta, HNF-3 beta, SOX-17, and GATA-6, while maintaining expression of ES markers SSEA-4.

25 **[0030]** Co expression of HNF-1 beta, HNF-3 beta (also known as FOXa2), SOX-17, and GATA-6 is regarded as the key step to define the formation of definitive endoderm during gastrulation. Thus, expression of these markers may be key in the generation of a pancreatic β -cell population, or a population of pancreatic hormone-producing cells, or a gut hormone-producing cell from an amniotic fluid-derived cell.

[0031] Therefore, there still remains a significant need to develop culture conditions for establishing amniotic fluid-derived cell lines that can be expanded to address the current clinical needs, while retaining the potential to differentiate into definitive endoderm, or a population of pancreatic hormone-producing cells, or a gut hormone-producing cell, or a β -cell lineage.

30 **[0032]** Wei Jun Ping et al. (Cell Transplantation, 2003, vol. 12(5), p545-552) disclose human amnion-isolated cells that can normalize blood glucose in streptozotocin-induced diabetic mice. Chen Li-Bo et al. (World J. Gastroenterology, 2004, vol. 10(20), p3016-3020) disclose the differentiation of rat marrow mesenchymal stem cells into pancreatic islet beta-cells. Held KR et al. (Prenatal Diagnosis, 1984, vol. 4(3), p171-180) disclose the effect of oxygen tension on colony formation and cell proliferation of amniotic fluid cells in vitro. Fauza D (Bailliere's Best Practice and Research: Clinical Obstetrics and Gynaecology, 2004, vol. 18(6), p877-891) reviews the amniotic fluid and the placenta as unique sources of different populations of stem cells--mesenchymal, hematopoietic, trophoblastic--and, possibly, of more primitive stem

40 **[0033]** WO 2005/017117 discloses a source of multipotent amniotic fluid/fetal stem cells (MAFSCs). MAFSC are of fetal origin and are characterized by the following cell surface markers: SSEA3, SSEA4, Tra-1-60, Tra-1-81, Tra-2-54, HLA class I, CD13, CD44, CD49b, CD 105 and are distinguished by the absence of the antigen markers CD34, CD45, and HLA Class II.

45 **[0034]** EP1391505A1 relates to a method for separating pancreatic stem cell of a mammal and an identification method thereof.

SUMMARY OF INVENTION

50 **[0035]** The invention provides a substantially pure population of amniotic fluid-derived cells, wherein said cells are: capable of differentiating into cells displaying the characteristics of the beta-cell lineage, substantially negative for the expression of the CD117 and Oct-4 protein markers, wherein the markers are not present or expressed in at least 70% of the total cell population, substantially positive for the expression of GATA-6 and SSEA-4, wherein GATA-6 and SSEA-4 are present or expressed in at least about 50% of the total cell population, and substantially either: i) positive for the

55 expression of SOX-17 and cytokeratin, wherein SOX-17 and cytokeratin are present or expressed in at least about 50% of the total cell population; ii) negative for the expression of SOX-17, wherein SOX-17 is not present or expressed in at least 70% of the total cell population, and positive in the expression of cytokeratin, wherein cytokeratin is present or expressed in at least about 50% of the total cell population; or iii) negative for the expression of SOX-17 and cytokeratin,

wherein SOX-17 and cytokeratin are not present or expressed in at least 70% of the total cell population.

[0036] The invention also provides a population of amniotic fluid-derived cells according to the present invention for use in a method of treating a patient with diabetes mellitus or at risk of developing diabetes.

[0037] The invention also provides the in vitro use of the population of cells according to the present invention, in a method wherein the cells of said population are differentiated into pancreatic hormone producing cells.

SUMMARY OF DISCLOSURE

[0038] Disclosed herein is a method for isolating mammalian amniotic fluid-derived cells. Amniotic fluid-derived cells are obtained from amniotic fluid samples of about 14 to about 23 weeks gestation. Alternatively, the amniotic fluid-derived cells are obtained from amniotic fluid samples of about 23 to about 40 weeks gestation.

[0039] In one embodiment, the cultures are left undisturbed for at least 5 to 10 days under hypoxic conditions (3% O₂). Alternatively, the cultures are left undisturbed for at least 5 to 10 days under normoxic conditions (approximately 20% O₂).

[0040] In an alternate embodiment, amniotic fluid-derived cells are obtained from amniotic fluid samples from the second trimester of gestation. Alternatively, the amniotic fluid-derived cells are obtained from amniotic fluid samples from the third trimester of gestation.

[0041] In one embodiment, the cultured amniotic fluid-derived cells are isolated as single cells, and clonally expanded.

[0042] The amniotic fluid-derived isolated according to the methods disclosed herein can be contacted, for example, with an agent (such as an antibody) that specifically recognizes a protein marker expressed by amniotic fluid cells, to identify and select amniotic fluid-derived cells, thereby obtaining a substantially pure population of amniotic fluid-derived cells, i.e., wherein a recognized protein marker is expressed in at least 50% of the cell population.

[0043] In one embodiment, the resulting amniotic fluid-derived cell population is substantially positive for at least one of the following markers: HNF-1 beta, HNF-3 beta, SOX-17, or GATA-6. The amniotic fluid-derived cell population is substantially negative for at least one of the following markers: CD117, Oct-4, or Tra2-54. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings without losing the capacity to express HNF-1 beta, HNF-3 beta, SOX-17, or GATA-6.

[0044] In one embodiment, the amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings without losing the capacity to express HNF-1 beta, HNF-3 beta, SOX-17, or GATA-6.

[0045] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for at least one of the following markers: SOX-17, CD117, Oct-4, or Tra2-54. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0046] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for cytokeratin and at least one of the following markers: SOX-17, CD117, Oct-4, or Tra2-54. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0047] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for SOX-17. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0048] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for the following markers: cytokeratin, and SOX-17. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0049] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for SOX-17. The amniotic fluid-derived cell population is further negative for at least one of the following markers: CD 117, Oct-4, or Tra2-54. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0050] In one embodiment, the amniotic fluid-derived cell population isolated according to the methods disclosed herein is substantially negative for the following markers: cytokeratin, and SOX-17. The amniotic fluid-derived cell population is further negative for at least one of the following markers: CD117, Oct-4, or Tra2-54. The amniotic fluid-derived cell population is substantially positive for the following markers: SSEA4 and CD44. The amniotic fluid-derived cell population can be expanded for more than 50 population doublings.

[0051] In another embodiment, the present disclosure provides an isolated pure population of amniotic fluid-derived cells that are substantially negative for at least one of the following markers: CD117, Oct-4, or Tra2-54.

[0052] In another embodiment, the present disclosure provides an isolated pure population of amniotic fluid-derived cells that are substantially negative for at least one of the following markers: SOX-17, CD117, Oct-4, or Tra2-54.

[0053] In another embodiment, the present disclosure provides an isolated pure population of amniotic fluid-derived cells that are substantially negative for SOX-17.

[0054] In another embodiment, the present disclosure provides an isolated pure population of amniotic fluid-derived cells that are substantially negative for SOX-17, and substantially negative for at least one of the following markers: CD117, Oct-4, or Tra2-54.

[0055] In one embodiment, the amniotic fluid-derived cells isolated according to the methods disclosed herein may also express at least one of the following: Musashi-1 and Hes1.

[0056] The amniotic fluid-derived cells isolated and expanded as disclosed herein can be induced to differentiate into cells of the β cell lineage under appropriate *in vitro* or *in vivo* conditions. Accordingly, the amniotic fluid-derived cells selected and expanded according to the present disclosure, as well as the differentiated cells derived from the amniotic fluid-derived cells, are useful for treating Type 1 and 2 diabetes.

[0057] The amniotic fluid-derived cells isolated and expanded as disclosed herein can be induced to gut hormone-producing cells under appropriate *in vitro* or *in vivo* conditions. In one embodiment, the amniotic fluid-derived cells isolated and expanded as disclosed herein can be induced to gut hormone-producing cells under appropriate *in vitro* or *in vivo* conditions and may express insulin in a glucose responsive manner.

BRIEF DESCRIPTION OF THE DRAWINGS

[0058]

Figure 1 shows the isolation and culturing steps used to isolate the amniotic fluid derived cells of the present invention.

Figure 2 shows three distinct morphologies of cells isolated from an amniotic fluid sample at passage 0. a) AF morphology, b) epithelial morphology, and c) fibroblast morphology.

Figure 3 depicts the expression of cell surface markers on AF-I cells derived from amniotic fluid. The markers are indicated on panels a-n.

Figure 4 depicts the expression of cell surface markers on F cells derived from amniotic fluid. The markers are indicated on panels a-l.

Figure 5 depicts the expression of cell surface markers on E cells derived from amniotic fluid. The markers are indicated on panels a-m.

Figure 6 depicts immunofluorescence images of the F cells derived from amniotic fluid samples. F cells stained positive for a) vimentin, b) SSEA-4, and c) beta III tubulin.

Figure 7 depicts immunofluorescence images of the E cells derived from amniotic fluid samples. E cells stained positive for a) vimentin and nestin, b) SSEA-4, c) beta III tubulin, d) pan-cytokeratin, e) smooth muscle actin, and f) cytokeratin 19.

Figure 8 depicts immunofluorescence images of the AF-I cells derived from amniotic fluid samples. AF-I cells stained positive for a) vimentin and nestin, b) beta III tubulin, c) cytokeratin 19 and HES-1, d) pan-cytokeratin, e) SSEA-4, f) SOX-17 and ZO-1, g) GATA-6, h) HNF-1 beta, i) smooth muscle actin and HES-2.

Figure 9 shows the expression profile of AF-I, AF-II, and AF-III cells of the present disclosure.

Figure 10 depicts the population doubling curve of early passage AF-I cells.

Figure 11 depicts the expansion potential of AF, F, or E cell derived from different donors. ♦ shows the cell number of amniotic fluid-derived cells with AF-I morphology obtained from amniotic fluid from one donor at 14-23 weeks gestation. Cells were cultured in media number 5 (**Table II**). ▲ shows the cell number of amniotic fluid-derived cells with AF-I morphology obtained from amniotic fluid from a second donor at 14-23 weeks gestation. Cells were cultured in media number 5 (**Table II**). ■ shows the cell number of amniotic fluid-derived cells with F morphology obtained from amniotic fluid from a third donor at 14-23 weeks gestation. Cells were cultured in media number 15 (**Table II**).

* shows the cell number of amniotic fluid-derived cells with F morphology obtained from amniotic fluid from a

fourth donor at 14-23 weeks gestation. Cells were cultured in media number 16 (**Table II**). ● shows the cell number of amniotic fluid-derived cells with E morphology obtained from amniotic fluid from a donor at 14-23 weeks gestation. Cells were cultured in media number 5. + shows the cell number of amniotic fluid-derived cells with AF-II morphology obtained from amniotic fluid from a second donor at 14-23 weeks gestation. Cells were cultured in media number 5 (**Table II**). Δ shows the cell number of amniotic fluid-derived cells with AF-III morphology obtained from amniotic fluid from a second donor at 14-23 weeks gestation. Cells were cultured in media number 5 (**Table II**)

Figure 12 depicts the telomere length of an AF-I cell line cultured either in AMNIOMAX or DM-LG + 10 % FBS at an intermediate passage level (approximately 40 population doublings). Lane 1 is the molecular weight ladder, lane 2 is the high telomere length control, lane 3 is the low telomere length control, lane 4 is amniotic fluid-derived cells from a donor at Passage 12, cultured in DMEM-LG + 10% FBS, lane 5 is amniotic fluid-derived cells from the same donor at passage 12, cultured in media #5, and lane 6 is an embryonic carcinoma cell line (NTERA cells) that serves as a positive control.

Figure 13 shows the karyotype of a) AF-I, b) AF-II, and c) AF-III cells cultured at passage 7-9 (approximately 30-35 population doublings).

Figure 14 depicts the expansion potential of a single AF derived cell from one donor at term (approximately 38 weeks). Cells were cultured in media number 5 (**Table II**).

Figure 15 depicts the scatter plot gene expression profiles between the different amniotic fluid cell types. The Pearson correlation coefficient for each plot is also listed.

Figure 16 shows the effects of growth factors on gene expression in amniotic fluid-derived cells. Amniotic fluid-derived cells were obtained from a single donor and cultured for 12 days in conditioned media that was obtained from cultures of PANC-1 cells. The media was supplemented with the growth factors indicated. The levels of expression of HNF-3 beta and somatostatin were determined by real-time PCR. Human pancreas total RNA was included as a calibrator. Panel a shows the changes in HNF-3 beta expression. Panel b shows the changes in somatostatin expression.

Figure 17 shows the effects of L685,458 on cultured amniotic fluid-derived cells having the AF morphology. Panel a shows the relative differences in RNA expression of human Hes-1 in cultured AF cells treated with the concentrations of L685,458 indicated. Panel b shows the effects of L685,458 on the viability of the cultured cells following a treatment with L685,458. Cells were treated for three days, at the concentrations indicated. Changes in viability, corresponding to cytotoxicity were detected using an MTS assay, where a decrease in cell viability corresponds to a decrease in A490nm.

DETAILED DESCRIPTION

[0059] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the following subsections that describe or illustrate certain features, embodiments or applications of, or relating to, the present invention.

[0060] Disclosed herein are methods for isolating an amniotic fluid-derived cell population that is highly proliferative, and displays embryonic-like characteristics. Similar cells may also be present in the chorionic villus. Some embodiments disclosed herein describe three morphologically distinct populations of amniotic fluid-derived cells: "fibroblastic" (F), epithelial" (E) cells, and "amniotic fluid" (AF) cells.

Definitions

[0061] "β-cell lineage" refer to cells with positive gene expression for the transcription factor PDX-1 and at least one of the following transcription factors: NGN-3, Nkx2.2, Nkx6.1, NeuroD, Isl-1, HNF-3 beta, MAFA, Pax4, and Pax6. Characteristics of cells of the beta cell lineage are well known to those skilled in the art, and additional characteristics of the beta cell lineage continue to be identified. These transcription factors are well established in prior art for identification of endocrine cells (Nature Reviews Genetics, Vol3, 524-632, 2002).

[0062] "Pancreatic islet-like structure" refers to a three-dimensional clusters of cells derived by practicing the methods of the disclosure, which has the appearance of a pancreatic islet. The cells in a pancreatic islet-like structure express at least the PDX-1 gene and one hormone selected from the list glucagon, somatostatin, or insulin.

[0063] The term "hypoxic" refers to oxygen levels less than 20%, preferably less than 10%, and more preferably less

than 5% but more than 1%.

[0064] The term "normoxia" refers to atmospheric oxygen levels of about 20%.

[0065] The term "substantially positive," when used in connection with a population of cells with respect to the expression of certain marker (such as a membrane receptor, cytoplasmic or nuclear protein, or a transcription factor), means that the marker is present or expressed in at least about 50%, alternatively at least about 60%, and alternatively at least about 70%, of the total cell population.

[0066] The term "substantially negative," when used in connection with a population of cells with respect to the expression of certain marker (such as a membrane receptor, cytoplasmic or nuclear protein, or a transcription factor), means that the marker is not present or expressed in at least about 70%, alternatively about 80%, alternatively about 90%, of the total cell population.

[0067] A "stem cell" as used herein refers to an undifferentiated cell that is capable of extensive propagation either *in vivo* or *ex vivo* and capable of differentiation to other cell types.

[0068] A "progenitor cell" refers to a cell that is derived from a stem cell by differentiation and is capable of further differentiation to more mature cell types. Progenitor cells typically have more restricted proliferation capacity as compared to stem cells.

[0069] "Expandable population" refers to the ability of an isolated cell population to be propagated through at least 50 or more cell divisions in a cell culture system.

[0070] By "undifferentiated cells," when used in connection with cells isolated from a amniotic fluid, are meant a population of amniotic fluid-derived cells that are substantially negative for the expression of PDX-1, or insulin.

[0071] By "differentiated cells," when used in connection with cells isolated from amniotic fluid, are meant a population of amniotic fluid-derived cells that are substantially positive for the expression of PDX-1, or insulin.

[0072] "Markers" as used herein, are nucleic acid or polypeptide molecules that are differentially expressed in a cell of interest. In this context, differential expression means an increased level of the marker for a positive marker, and a decreased level for a negative marker. The detectable level of the marker nucleic acid or polypeptide is sufficiently higher or lower in the cells of interest, compared to other cells, such that the cell of interest can be identified and distinguished from other cells, using any of a variety of methods known in the art.

[0073] "c-Kit" and "CD 117" both refer to a cell surface receptor tyrosine kinase having a sequence disclosed in Genbank Accession No. X06182, or a naturally occurring variant sequence thereof (e.g., allelic variant).

[0074] "CD9" is also referred to as "Motility-related protein-1 (MRP-1)" and is a transmembrane glycoprotein that has been implicated in cell adhesion, motility, proliferation, and differentiation.

[0075] "CD10" is also referred to as "Common Acute Lymphocytic Leukemia Antigen (CALLA)". CD10 is a cell surface enzyme with neutral metalloendopeptidase activity and it is expressed in lymphoblastic, Burkitt's, and follicular germinal center lymphomas and in patients with chronic myelocytic leukemia. It is also expressed on the surface of normal early lymphoid progenitor cells, immature B Cells within adult bone marrow and germinal center B Cells within lymphoid tissue. CD10 is also present on breast myoepithelial cells, bile canaliculi, fibroblasts, brush border of kidney and gut epithelial cells.

[0076] "CD44" is also referred to as "Hermes antigen" and is the main cell surface receptor for hyaluronan. This CD is primarily expressed in most cell types, except for tissues/cells such as hepatocytes, some epithelial cells, and cardiac muscle.

[0077] "CD49f" is also referred to as "a6 integrin" and "VLA-6," and associates with integrin subunit beta 1 to bind laminin. CD49f is expressed primarily on epithelial cells, trophoblasts, platelets, and monocytes.

[0078] "CD73" is also referred to as "ecto-5'-nucleotidase" and is primarily expressed on a subset of B and T cells, bone marrow stromal cells, various epithelial cells, fibroblasts, and endothelial cells.

[0079] "CD90" is also referred to as "Thy-1" and is primarily expressed on hematopoietic stem cells, connective tissue cells, and various fibroblastic and stromal cells.

[0080] "SSEA-1" (Stage Specific Embryonic Antigen-1) is a glycolipid surface antigen present on the surface of murine teratocarcinoma stem cells (EC), murine and human embryonic germ cells (EG), and murine embryonic stem cells (ES).

[0081] "SSEA-3" (Stage Specific Embryonic Antigen-3) is a glycolipid surface antigen present on the surface of human teratocarcinoma stem cells (EC), human embryonic germ cells (EG), and human embryonic stem cells (ES).

[0082] "SSEA-4" (Stage Specific Embryonic Antigen-4) is a glycolipid surface antigen present on the surface of human teratocarcinoma stem cells (EC), human embryonic germ cells (EG), and human embryonic stem cells (ES).

[0083] "TRA1-60" is a keratin sulfate related antigen that is expressed on the surface of human teratocarcinoma stem cells (EC), human embryonic germ cells (EG), and human embryonic stem cells (ES).

[0084] "TRA1-81" is a keratin sulfate related antigen that is expressed on the surface of human teratocarcinoma stem cells (EC), human embryonic germ cells (EG), and human embryonic stem cells (ES).

[0085] "TRA2-49" is an alkaline phosphatase isozyme expressed on the surface of human teratocarcinoma stem cells (EC), and human embryonic stem cells (ES).

[0086] "Oct-4" is a member of the POU-domain transcription factor and is widely regarded as a hallmark of pluripotent

stem cells. The relationship of Oct-4 to pluripotent stem cells is indicated by its tightly restricted expression to undifferentiated pluripotent stem cells. Upon differentiation to somatic lineages, the expression of Oct-4 disappears rapidly.

[0087] "EPCAM" is also referred to as "Epithelial Cell Adhesion Molecule" is broadly expressed on cells of epithelial origin and epithelial derived tumor cells.

[0088] "Rex-1" is a developmentally regulated acidic zinc finger gene (*Zfp-42*). Rex-1 message level is high in embryonic stem cells and reduced upon induction of differentiation. As expected for a stem-cell-specific message, Rex-1 mRNA is present in the inner cell mass (ICM) of blastocyst, polar trophoblast of the blastocyst and later in the ectoplacental cone and extraembryonic ectoderm of the egg cylinder (trophoblast-derived tissues), but its abundance is much reduced in the embryonic ectoderm, which is directly descended from the ICM.

[0089] "HNF-1 alpha", "HNF-1 beta" and "HNF-3 beta" belong to the hepatic nuclear factor family of transcription factors, which is characterized by a highly conserved DNA binding domain and two short carboxy-terminal domains.

[0090] "GATA-4" and "GATA-6" are members of the GATA transcription factor family. This family of transcription factors are induced by TGF β signaling and contribute to the maintenance of early endoderm markers, Sox17 α and HNF-1 beta, and the later marker HNF-3 beta.

[0091] "SOX-17" is a transcription factor, which is implicated in the formation of endoderm during embryogenesis.

[0092] By "basic defined cell culture medium" is meant a serum free or serum containing, chemically defined cell growth medium. Such medium includes, but is not limited to, Dulbecco's Modified Eagle's Medium (DMEM), alpha modified Minimum Essential Medium (alpha MEM), Basal Medium Essential (BME), CMRL-1066, RPMI 1640, M199 medium, Ham's F10 nutrient medium, KNOCKOUT™ DMEM, Advanced DMEM, MCDB based media such as MCDB -151, -153, -201, and -302 (Sigma, MO), and DMEM/F12. These and other useful media are available from GIBCO, Grand Island, New York, U.S.A., for example. A number of these media are reviewed in Methods in Enzymology, Volume LVIII, "Cell Culture," pp. 62-72, edited by William B. Jakoby and Ira H. Pastan, published by Academic Press Inc.

[0093] "Hes-1", also known as "hairy/enhancer of split-1" is a transcription factor that may influence cell fate determination.

[0094] "Musashi-1" is a member of a subfamily of RNA binding proteins that are highly conserved across species. Musashi-1 expression is highly enriched in proliferative cells within the developing central nervous system, and may be a stem cell marker in intestinal cells.

[0095] "Pharmaceutical carrier" refers to a biodegradable or non-degradable porous or nonporous matrix that can act as a carrier for transplantation of mammalian cells.

[0096] "Transplantation" as used herein, can include the steps of introducing a cell or a population of cells or tissue into a mammal such as a human patient. "Transplantation" may also include incorporating cells or tissue into a pharmaceutical carrier, and implanting the carrier in a mammal such as a human patient.

Isolation of Amniotic Fluid-Derived Cells

[0097] Amniotic fluid-derived cells may be isolated by a multi-stage method, which essentially involves:

- Isolation of amniotic fluid,
- Centrifugation of the amniotic fluid, followed by removal of the supernatant,
- Resuspending the cell pellet in growth medium,
- Culturing the tissues and cells in a low oxygen environment,
- Leaving the culture undisturbed for about 5 to 10 days without any media changes,
- Isolation of distinct colonies using cloning rings,
- Culturing the isolated colonies in growth media
- Serial dilution cloning and identification of single cells that give rise to proliferating colonies, and
- Culturing the clones in growth media.

[0098] In an alternate embodiment, amniotic fluid-derived cells are isolated by a multi-stage method, which essentially involves:

- Isolation of amniotic fluid,
- Centrifugation of the amniotic fluid, followed by removal of the supernatant,
- Resuspending the cell pellet in growth medium,
- Culturing the tissues and cells in a normoxic environment,
- Leaving the culture undisturbed for about 5 to 10 days without any media changes,
- Isolation of distinct colonies using cloning rings,
- Culturing the isolated colonies in growth media,
- Serial dilution cloning and identification of single cells that give rise to proliferating colonies, and

- Culturing the clones in growth media.

[0099] The culture plates may be pre-coated with agents such as, for example, fibronectin, vitronectin, laminin, collagen, gelatin, thrombospondin, placenta extracts, MATRIGEL™, tenascin, human serum, or combinations thereof.

[0100] If desirable, the amniotic fluid may be exposed, for example, to an agent (such as an antibody) that specifically recognizes a protein marker expressed by amniotic fluid cells, to identify and select amniotic fluid-derived cells, thereby obtaining a substantially pure population of amniotic fluid-derived cells.

[0101] Amniotic fluid-derived cells may be cultured in AMNIOMAX™ complete medium (Invitrogen). Alternatively, the cells may be cultured in Chang B/C medium (Irvine Scientific). Alternatively, the cells may be cultured in low glucose DMEM, supplemented with insulin-transferrin-selenium-X (ITS-X, Invitrogen, CA), 2% fetal bovine serum (FBS), 1% penicillin/streptomycin (P/S) + 25 ng/ml bFGF. Alternatively, the cells may be cultured in, DM-KNOCKOUT™ media (Invitrogen, CA), supplemented with 20% KNOCKOUT™ serum replacement (Invitrogen, CA), 10 ng/ml bFGF. Alternatively, the cells may be cultured in Williams' medium E supplemented with 2% defined FBS, 2mM L-glutamine, ITS, 55 μM 2-mercaptoethanol, 10ng/ml EGF, 4ng/ml bFGF, and 4ng/ml dexamethasone. Alternatively, the cells may be cultured in 1:1 DMEM-LG/MCDB 201, 2% FBS, ITS-X, βme 55 μM, 100 μM ascorbic acid-2-phosphate, 4ng/ml bFGF, 10ng/ml EGF, and 4ng/ml dexamethasone. Alternatively, the cells may be cultured in low glucose DMEM, supplemented with 20% FBS. Alternatively, the cell may be cultured in low glucose DMEM, supplemented with 5% FBS. The cells may also be cultured in low glucose DMEM/MCDB 201 medium (1:1), supplemented with 2% defined FBS, ITS-X, 1nM dexamethasone, 100 mM ascorbic acid 2-phosphate, 10ng/ml EGF, 10ng/ml PDGF-bb and 100 mM 2-mercaptoethanol. The media may be supplemented with bFGF, at concentrations from about 5 ng/ml to about 100 ng/ml. Alternatively, the cells may be cultured in 20% KNOCKOUT™ serum replacement + 80% KNOCKOUT™ DMEM, supplemented with 1 mM L-glutamine, 1% non-essential amino acids and 0.1 mM 2-mercaptoethanol. The medium may be conditioned overnight, on human or murine embryonic fibroblasts, human bone marrow derived stromal cells, or human placenta derived cells and supplemented with 4 ng/ml bFGF. Alternatively, the cells may be cultured in high glucose DMEM, supplemented with 20% defined FBS with 0.1 mM 2-mercaptoethanol. **Table II** lists the various media formulations used to culture the amniotic fluid-derived cells of the present invention.

[0102] During culture in growth media, the cells may be cultured under hypoxic or normoxic conditions. Under hypoxic conditions, oxygen levels are lower than 20%, alternatively lower than 10%, alternatively lower than 5%, but more than 1%.

[0103] Preferably, the culture should be maintained in the growth media undisturbed for about 5 to 14 days without any media changes, at which point the cells have typically become adherent to the culture substrate used. At which point, cells may be sub-cultured.

[0104] Subculture can be achieved with any of the enzymatic solutions well known to those skilled in the art. An example of an enzymatic solution suitable for use in the present invention is TrypLE EXPRESS™ (Invitrogen, Ca).

[0105] Furthermore, the amniotic fluid-derived cells may be expanded by culturing in a defined growth media containing agent(s) that stimulate the proliferation of the cells of the present invention. These factors may include, for example, nicotinamide, members of TGF-β family, including TGF-β1, 2, and 3, bone morphogenic proteins (BMP-2, -4, 6, -7, -11, -12, and -13), serum albumin, fibroblast growth factor family, platelet-derived growth factor-AA, and -BB, platelet rich plasma, insulin growth factor (IGF-I, II) growth differentiation factor (GDF-5, -6, -8, -10, 11), glucagon like peptide-I and II (GLP-I and II), GLP-1 and GLP-2 mimetobody, Exendin-4, retinoic acid, parathyroid hormone, insulin, progesterone, testosterone, estrogen, aprotinin, hydrocortisone, ethanolamine, beta mercaptoethanol, epidermal growth factor (EGF), gastrin I and II, copper chelators such as triethylene pentamine, TGF-α, forskolin, Na-Butyrate, activin, betacellulin, noggin, neuron growth factor, nodal, insulin/transferring/selenium (ITS), hepatocyte growth factor (HGF), keratinocyte growth factor (KGF), bovine pituitary extract, islet neogenesis-associated protein (INGAP), proteasome inhibitors, notch pathway inhibitors, sonic hedgehog inhibitors, GSK-3 beta inhibitors, or combinations thereof. Alternatively, the amniotic fluid-derived cells may be expanded by culturing in conditioned media. By "conditioned media" is meant that a population of cells is grown in a basic defined cell culture medium and contributes soluble factors to the medium. In one such use, the cells are removed from the medium, while the soluble factors the cells produce remain. This medium is then used to nourish a different population of cells.

[0106] In certain embodiments, the amniotic fluid-derived cells are cultured on standard tissue culture plates. Alternatively, the culture plates may be coated with extracellular matrix proteins, such as, for example, MATRIGEL®, growth factor reduced MATRIGEL®, laminin, collagen, gelatin, tenascin, fibronectin, vitronectin, thrombospondin, placenta extracts, human serum, or combinations thereof.

Characterization of The Isolated Amniotic Fluid-Derived Cells

[0107] Methods for assessing expression of protein and nucleic acid markers in cultured or isolated cells are standard in the art. These include quantitative reverse transcriptase polymerase chain reaction (RT-PCR), Northern blots, *in situ* hybridization (see, e.g., Current Protocols in Molecular Biology (Ausubel et al., eds. 2001 supplement)), and immu-

noassays, such as immunohistochemical analysis of sectioned material, Western blotting, and for markers that are accessible in intact cells, flow cytometry analysis (FACS) (see, e.g., Harlow and Lane, Using Antibodies: A Laboratory Manual, New York: Cold Spring Harbor Laboratory Press (1998)).

[0108] Examples of antibodies useful for detecting certain protein markers are listed in **Table III**. It should be noted that other antibodies directed to the same markers that are recognized by the antibodies listed in **Table III** are available, or can be readily developed. Such other antibodies can also be employed for assessing expression of markers in the cells isolated in accordance with the present invention.

[0109] Characteristics of cells of the β -cell lineage are well known to those skilled in the art, and additional characteristics of the β -cell lineage continue to be identified. These characteristics can be used to confirm that the amniotic fluid-derived cells isolated in accordance with the present invention have differentiated to acquire the properties characteristic of the β -cell lineage. β -cell lineage specific characteristics include the expression of one or more transcription factors such as, for example, PDX-1 (pancreatic and duodenal homeobox gene-1), NGN-3 (neurogenin-3), Hlx9, Nkx6, Isl1, Pax6, NeuroD, Hnf1a, Hnf6, Hnf3 Beta, and Mafa, among others. These transcription factors are well established in the art for identification of endocrine cells. See, e.g., Edlund (Nature Reviews Genetics 3: 524-632 (2002)).

[0110] Characteristics of cells of the intestinal cell lineage are well known to those skilled in the art, and additional characteristics of this lineage continue to be identified. These characteristics can be used to confirm that the differentiated or undifferentiated amniotic fluid-derived cells isolated in accordance with the present invention have some of the properties characteristic of the intestinal cell lineage. Intestinal cell lineage characteristics include the expression of one or more transcription factors such as, for example, HES-1 (hair/enhancer of split-1), NGN-3, Pax6, NeuroD, Math-1, and Musashi-1, among others. In addition, gut cells express hormones such as secretin, cholecystokinin, GLP-1, neurotensin, gastric inhibitory peptide (GIP), serotonin, somatostatin, and gastrin, among others. These transcription factors and gut hormones are well established in the art for identification of intestinal cells. See, e.g., Schonhoff (Endocrinology 145: 2639-2644 (2004)).

[0111] The present inventors have identified and isolated a population of amniotic fluid-derived cells that is highly proliferative, and displays embryonic cell-like characteristics, and may express at least one of the following markers: HNF-1 beta, HNF-3 beta, SOX-17, or GATA 6. In particular, the amniotic fluid-derived cells isolated as disclosed herein are characterized as, *inter alia*, substantially lacking at least one of the following protein markers: CD117, Oct-4 or Tra2-54. Populations of amniotic fluid-derived cells with these characteristics are referred to herein as AF-I (ATCC accession number PTA-6975).

[0112] Under the above growth conditions for expansion, the amniotic fluid cells isolated as disclosed herein may be expanded for more than 50 population doublings, while maintaining the potential to express at least one of the following markers: HNF-1 beta, HNF-3 beta, SOX-17, or GATA-6.

[0113] The present inventors have also identified and isolated populations of amniotic fluid-derived cells that is highly proliferative, displays embryonic cell-like characteristics, and do not express at least one of following markers: HNF-3 beta, SOX-17, GATA-4, CD 117, Oct-4 or Tra2-54. In particular, the amniotic fluid-derived cells isolated in accordance with the present disclosure are characterized as, *inter alia*, substantially lacking at least one of the following protein markers: CD117, Oct-4 or Tra2-54. Populations of amniotic fluid-derived cells with these characteristics are referred to herein as AF-II.

[0114] The present inventors have also identified and isolated populations of amniotic fluid-derived cells that is highly proliferative, displays embryonic cell-like characteristics, and do not express any of the following markers: HNF-3beta, SOX-17, GATA-4, CD117, Oct-4 or Tra2-54. In particular, the amniotic fluid-derived cells isolated in accordance with the present disclosure are characterized as, *inter alia*, substantially lacking at least one of the following protein markers: CD117, Oct-4 or Tra2-54. Populations of amniotic fluid-derived cells with these characteristics are referred to herein as AF-II.

[0115] The present inventors have also identified and isolated populations of amniotic fluid-derived cells that is highly proliferative, displays embryonic cell-like characteristics, and do not express cytokeratin and at least one of following markers: HNF-3 beta, SOX-17, GATA-4, CD117, Oct-4 or Tra2-54. In particular, the amniotic fluid-derived cells isolated in accordance with the present disclosure are characterized as, *inter alia*, substantially lacking at least one of the following protein markers: CD117, Oct-4 or Tra2-54. Populations of amniotic fluid-derived cells with these characteristics are referred to herein as AF-III.

[0116] The present inventors have also identified and isolated populations of amniotic fluid-derived cells that is highly proliferative, displays embryonic cell-like characteristics, and do not express any of the following markers: cytokeratin, HNF-3beta, SOX-17, GATA-4, CD 117, Oct-4 or Tra2-54. In particular, the amniotic fluid-derived cells isolated in accordance with the present disclosure are characterized as, *inter alia*, substantially lacking at least one of the following protein markers: CD117, Oct-4 or Tra2-54. Populations of amniotic fluid-derived cells with these characteristics are referred to herein as AF-III.

[0117] A summary of the expression profile of AF-I, AF-II and AF-III cells is shown in Figure 9.

[0118] Amniotic fluid-derived cells of the present invention may be expanded for more than 50 population doublings,

while maintaining the potential to differentiate into definitive endoderm, or cells with characteristics of a pancreatic β -cell lineage, or the capacity to differentiate into a gut hormone-producing cell.

Differentiation Of Amniotic Fluid-Derived Cells

[0119] In one aspect, the present disclosure provides compositions capable of differentiating the expanded amniotic fluid-derived cells of this invention into cells bearing markers characteristic of the β cell lineage.

[0120] In another aspect, the present disclosure provides compositions capable of differentiating the expanded amniotic fluid-derived cells of this invention into cells bearing markers characteristic of definitive endoderm.

[0121] In another aspect, the present disclosure provides compositions capable of differentiating the expanded amniotic fluid-derived cells of this invention into cells bearing markers characteristic of a gut hormone-producing cell.

[0122] A basic defined culture medium, when supplied with one or more components, that support the growth of amniotic fluid-derived cells, supplemented with differentiation-inducing amounts of one or more growth factors, is referred to as an "induction medium." In accordance with the present invention, the induction medium contains less than or equal to 20% serum. In one embodiment, fetal calf serum may be used. Alternatively, fetal bovine serum may be replaced by serum from any mammal, or by albumin, bovine albumin or other compounds that permit or enhance differentiation of amniotic fluid-derived cells to the β cell lineage. Alternatively, the induction medium may be conditioned medium.

[0123] Factors appropriate for use in the induction medium may include, for example, nicotinamide, members of TGF- β family, including TGF- β 1, 2, and 3, bone morphogenic proteins (BMP-2, -4, 6, -7, -11, -12, and -13), serum albumin, fibroblast growth factor family, platelet-derived growth factor-AA, and -BB, platelet rich plasma, insulin growth factor (IGF-I, II) growth differentiation factor (GDF-5, -6, -8, -10, 11), glucagon like peptide-I and II (GLP-I and II), GLP-1 and GLP-2 mimetobody, Exendin-4, retinoic acid, parathyroid hormone, insulin, progesterone, aprotinin, hydrocortisone, ethanolamine, beta mercaptoethanol, epidermal growth factor (EGF), gastrin I and II, copper chelators such as triethylene pentamine, TGF- α , forskolin, Na-Butyrate, activin, betacellulin, ITS, noggin, neurite growth factor, nodal, valporic acid, trichostatin A, sodium butyrate, hepatocyte growth factor (HGF), sphingosine-1, Wnt proteins such as Wnt-1, -3, -3a, 07a, and -8, keratinocyte growth factor (KGF), Dickkopf protein family, bovine pituitary extract, islet neogenesis-associated protein (INGAP), Indian hedgehog, sonic hedgehog, proteasome inhibitors, notch pathway inhibitors, sonic hedgehog inhibitors, or combinations thereof.

[0124] In one aspect of the present disclosure, a combination of growth factors and chemical agents, including bFGF, Activin-A, FGF5, N2 and B27 supplements (Gibco, CA), steroid alkaloid such as, for example, cyclopamine (EMD, CA) that inhibits sonic hedgehog signaling, and a proteasome inhibitor such as, for example MG132 (EMD, CA), is supplied to a basic defined medium to support differentiation of amniotic fluid-derived cells into a β -cell lineage. In one aspect, the cells are cultured in an induction media composed of DMEM (low glucose, 5.5 mM) containing 10 micromolar MG-132 for 1-2 days, followed by additional incubation for 3-7 days in an induction media supplemented with 1X B27 (Gibco, CA) and 1X N2 (Gibco, CA) and further supplemented with Cyclopamine (10 μ M; EMD, CA), bFGF (20 ng/ml; R&D Systems, MN), Activin A (20 nM; R&D Systems, MN) or FGF5 (20 ng/ml; R&D Systems, MN) for an additional five days.

[0125] In another aspect of the disclosure, the cells of the current invention can be treated with conditioned media isolated from cultures of primary fetal intestinal or pancreatic rudiments to induce further differentiation into the intestinal or pancreatic lineages, respectively. The cells may also be induce to differentiate with conditioned media from pancreatic cells lines such as PANC-1, CAPAN-1, BxPC-3, HPAF-II, hepatic cell lines such as HepG2, and intestinal cell lines such as, for example, FHs 74 and HS738. Alternatively, the cells of the present invention can be treated with conditioned media isolated from human or mouse embryonic stem cells induced to differentiate into an endodermal lineage. These cell lines can be purchased from the ATCC (VA).

[0126] The combination and concentrations of growth factors, the length of culture, and other culture conditions can be optimized by those skilled in the art to achieve effective differentiation by, e.g., monitoring the percentage of cells that have differentiated into cells characteristic of the β -cell lineage. The one or more growth factors may be added in an amount sufficient to induce the differentiation of the amniotic fluid-derived cells of the present invention into cells bearing markers of a β -cell lineage over a time period of about one to four weeks.

Therapeutic Use Of The Cells Of The Present Invention.

[0127] In one aspect, the present disclosure includes a method for treating a patient suffering from, or at risk of developing Type 1 diabetes. This method involves isolating and culturing amniotic fluid-derived cells, expanding the isolated population of cells, differentiating the cultured cells *in vitro* into a β -cell lineage, and implanting the differentiated cells either directly or in a pharmaceutical carrier into the patient. If appropriate, the patient can be further treated with pharmaceutical agents or bioactives that facilitate the survival and function of the transplanted cells. These agents may include, for example, insulin, members of the TGF- β family, including TGF- β 1, 2, and 3, bone morphogenic proteins (BMP-2, -3, -4, -5, -6, -7, -11, -12, and -13), fibroblast growth factors-1 and -2, platelet-derived growth factor-AA, and

-BB, platelet rich plasma, insulin growth factor (IGF-I, II) growth differentiation factor (GDF-5, -6, -8, -10, -15), vascular endothelial cell-derived growth factor (VEGF), pleiotrophin, endothelin, among others. Other pharmaceutical compounds can include, for example, nicotinamide, glucagon like peptide-I (GLP-1) and II, GLP-1 and 2 mimetibody, Exendin-4, retinoic acid, parathyroid hormone, MAPK inhibitors, such as, for example, compounds disclosed in U.S. Published Application 2004/0209901 and U.S. Published Application 2004/0132729.

[0128] In yet another aspect, this disclosure includes a method for treating a patient suffering from, or at risk of developing Type 2 diabetes. The method involves isolating and culturing amniotic fluid-derived cells according to the present invention, expanding the isolated population of cells, differentiating the cultured cells *in vitro* into a β -cell lineage, and implanting the differentiated cells either directly or in a pharmaceutical carrier into said patient.

[0129] The amniotic fluid-derived cells of the present invention may be cryopreserved using commercially available medium containing DMSO (dimethylsulfoxide) or glycerol. The banked and frozen cells may be stored in the vapor phase of a liquid nitrogen storage tank until needed.

[0130] The amniotic fluid-derived cells of the present invention may be transplanted with mature islets of the same or different animal species to enhance the survival of the amniotic fluid-derived cells or to induce further differentiation of the amniotic fluid-derived cells into a pancreatic β cell lineage. The source of amniotic fluid from which the cells are isolated may be autologous in relation to the patient undergoing the therapeutic treatment. Alternatively, the source may be allogeneic, or xenogeneic. Cells to be administered to a patient may also be genetically modified to enhance proliferation and/or differentiation or prevent or lessen the risk of immune rejection. Alternatively, the amniotic fluid-derived cells obtained in accordance with the present invention can be used to modulate the recipient's immune response, prior to transplantation of differentiated cells prepared in accordance with the present invention. See, for example, U.S. Patent 6,328,960, U.S. Patent 6,281,012. The amniotic fluid-derived cells of the present invention may be differentiated into an insulin-producing cell prior to transplantation into a recipient. The amniotic fluid-derived cells of the present invention may be fully differentiated into β -cells, prior to transplantation into a recipient. Alternatively, the amniotic fluid-derived cells of the present invention may be transplanted into a recipient in an undifferentiated or partially differentiated state. Further differentiation may take place in the recipient. The amniotic fluid-derived cells of the present invention may be genetically modified. For example, the cells may be engineered to over express markers characteristic of a cell of a β -cell lineage, such as, for example, PDX-1 or insulin. The cells may be engineered to over express with any suitable gene of interest. Furthermore, the cells may be engineered to over express markers characteristic of an intestinal cell, such as MATH-1. Alternatively, the cells of the present invention can be differentiated into a GIP expressing cell population and further modified with an insulin gene under control of the GIP promoter to become glucose responsive and insulin-producing cell population.

[0131] Techniques useful to genetically modify the amniotic fluid-derived cells of the present invention can be found, for example, in standard textbooks and reviews in cell biology. Methods in molecular genetics and genetic engineering are described, for example, in Molecular Cloning: A Laboratory Manual, 2nd Ed. (Sambrook et al., 1989); Oligonucleotide Synthesis (M. J. Gait, ed., 1984); Animal Cell Culture (R. I. Freshney, ed., 1987); the series Methods in Enzymology (Academic Press, Inc.); Gene Transfer Vectors for Mammalian Cells (I. M. Miller & M. P. Calos, eds., 1987); Current Protocols in Molecular Biology and Short Protocols in Molecular Biology, 3rd Edition (F. M. Ausubel et al., eds., 1987 & 1995); and Recombinant DNA Methodology II (R. Wu ed., Academic Press 1995).

[0132] The nucleic acid molecule, encoding the gene of interest may be stably integrated into the genome of the host amniotic fluid-derived cell, or the nucleic acid molecule may be present as an extrachromosomal molecule, such as a vector or plasmid. Such an extrachromosomal molecule may be auto-replicating. The term "transfection," as used herein, refers to a process for introducing heterologous nucleic acid into the host amniotic fluid-derived cell.

[0133] The cells, undifferentiated or otherwise, may be used as dispersed cells or formed into clusters that may be infused into the hepatic portal vein. Alternatively, the cells may be provided in biocompatible degradable polymeric supports, porous non-degradable devices or encapsulated to protect from host immune response. The cells may be implanted into an appropriate site in a recipient. The implantation sites include, for example, the liver, natural pancreas, renal subcapsular space, omentum, peritoneum, subserosal space, intestine, stomach, or a subcutaneous pocket.

[0134] To enhance further differentiation, survival or activity of implanted cells, additional factors, such as growth factors, antioxidants or anti-inflammatory agents, can be administered before, simultaneously with, or after the administration of the cells. In certain embodiments, growth factors are utilized to differentiate the administered cells *in vivo*. These factors can be secreted by endogenous cells and exposed to the administered amniotic fluid-derived cells *in situ*. Implanted amniotic fluid-derived cells can be induced to differentiate by any combination of endogenous and exogenously administered growth factors known in the art.

[0135] The amount of cells used in implantation depends on a number of factors including the patient's condition and response to the therapy, and can be determined by one skilled in the art.

[0136] In one aspect, this disclosure includes a method for treating a patient suffering from, or at risk of developing diabetes. The method includes isolating and culturing amniotic fluid-derived cells according to the present invention, expanding the isolated population of cells, differentiating *in vitro* the cultured amniotic fluid-derived cells into a β -cell

lineage, and incorporating the cells into a three-dimensional support. The cells can be maintained *in vitro* on this support prior to implantation into the patient. Alternatively, the support containing the cells can be directly implanted in the patient without additional *in vitro* culturing. The support can optionally be incorporated with at least one pharmaceutical agent that facilitates the survival and function of the transplanted cells.

[0137] Suitable support materials include tissue templates, conduits, barriers, and reservoirs useful for tissue repair. In particular, synthetic and natural materials in the form of foams, sponges, gels, hydrogels, textiles, and nonwoven structures, which have been used *in vitro* and *in vivo* to reconstruct or regenerate biological tissue, as well as to deliver chemotactic agents for inducing tissue growth, are suitable for use in practicing the methods of the present invention. See, e.g., the materials disclosed in U.S. Patent 5,770,417, U.S. Patent 6,022,743, U.S. Patent 5,567,612, U.S. Patent 5,759,830, U.S. Patent 6,626,950, U.S. Patent 6,534,084, U.S. Patent 6,306,424, U.S. Patent 6,365,149, U.S. Patent 6,599,323, U.S. Patent 6,656,488, and U.S. Patent 6,333,029. Exemplary polymers suitable for use in the present invention are disclosed in U.S. Published Application 2004/0062753 A1 and U.S. Patent 4,557,264. To form a support incorporated with a pharmaceutical agent, the pharmaceutical agent can be mixed with the polymer solution prior to forming the support. Alternatively, a pharmaceutical agent could be coated onto a fabricated support, preferably in the presence of a pharmaceutical carrier. The pharmaceutical agent may be present as a liquid, a finely divided solid, or any other appropriate physical form. Alternatively, excipients may be added to the support to alter the release rate of the pharmaceutical agent. In an alternate embodiment, the support is incorporated with at least one pharmaceutical compound that is an anti-inflammatory compound, such as, for example compounds disclosed in U.S. Patent 6,509,369.

[0138] The support may be incorporated with at least one pharmaceutical compound that is an anti-apoptotic compound, such as, for example, compounds disclosed in U.S. Patent 6,793,945.

[0139] The support may be incorporated with at least one pharmaceutical compound that is an inhibitor of fibrosis, such as, for example, compounds disclosed in U.S. Patent 6,331,298.

[0140] In a further embodiment, the support is incorporated with at least one pharmaceutical compound that is capable of enhancing angiogenesis, such as, for example, compounds disclosed in U.S. Published Application 2004/0220393 and U.S. Published Application 2004/0209901.

[0141] The support may be incorporated with at least one pharmaceutical compound that is an immunosuppressive compound, such as, for example, compounds disclosed in U.S. Published Application 2004/0171623.

[0142] The support may be incorporated with at least one pharmaceutical compound that is a growth factor, such as, for example, members of the TGF- β family, including TGF- β 1, 2, and 3, bone morphogenic proteins (BMP-2, -3, -4, -5, -6, -7, -11, -12, and -13), fibroblast growth factors-1 and -2, platelet-derived growth factor-AA, and -BB, platelet rich plasma, insulin growth factor (IGF-I, II) growth differentiation factor (GDF-5, -6, -8, -10, -15), vascular endothelial cell-derived growth factor (VEGF), pleiotrophin, endothelin, among others. Other pharmaceutical compounds can include, for example, nicotinamide, hypoxia inducible factor 1-alpha, glucagon like peptide-I (GLP-1), GLP-1 and GLP-2 mimetibody, and II, Exendin-4, nodal, noggin, NGF, retinoic acid, parathyroid hormone, tenascin-C, tropoelastin, thrombin-derived peptides, cathelicidins, defensins, laminin, biological peptides containing cell- and heparin-binding domains of adhesive extracellular matrix proteins such as fibronectin and vitronectin, MAPK inhibitors, such as, for example, compounds disclosed in U.S. Published Application 2004/0209901 and U.S. Published Application 2004/0132729.

[0143] The incorporation of the cells of the present invention into a scaffold can be achieved by the simple depositing of cells onto the scaffold. Cells can enter into the scaffold by simple diffusion (J. Pediatr. Surg. 23 (1 Pt 2): 3-9 (1988)). Several other approaches have been developed to enhance the efficiency of cell seeding. For example, spinner flasks have been used in seeding of chondrocytes onto polyglycolic acid scaffolds (Biotechnol. Prog. 14(2): 193-202 (1998)). Another approach for seeding cells is the use of centrifugation, which yields minimum stress to the seeded cells and enhances seeding efficiency. For example, Yang et al. developed a cell seeding method (J. Biomed. Mater. Res. 55(3): 379-86 (2001)), referred to as Centrifugational Cell Immobilization (CCI).

[0144] The present invention is further illustrated, but not limited by, the following examples.

Example 1

The Establishment of Human Amniotic Fluid-Derived Cell Lines

[0145] Amniotic fluid used to isolate the cells of the present invention was taken from samples taken from routine amniocentesis performed at 16 to 22 weeks of gestation for fetal karyotyping. The multi-stage method used to isolate the amniotic fluid-derived cells is outlined in **Figure 1**. The amniotic fluid was centrifuged for 7 minutes at 400 x g and the supernatant removed. The resulting cell pellet was resuspended in the growth media indicated in **Table III** for the amniotic fluid samples used in the present invention. The cells were cultured either on collagen type IV (1mg/100 mm plate), or on collagen type I (1 microgram/cm²), vitronectin (10 microgram/ml) or fibronectin (10 micrograms/ml) coated plates. The cell yield from amniotic fluid samples had a large variation (8000-300000 cell/sample) and some samples also contained a significant number of blood cell contamination. The cultures were left undisturbed for at least 5-10 days

under hypoxic conditions (3% O₂). In parallel, cultures were established under similar conditions in normoxic conditions. Next, the cultures were fed with the same growth media and cultured until the cultures reached 70-80% confluency. Cells at this stage were referred to as "P0". In some cultures, colonies of cells were isolated by a cloning ring and subcultured into a different culture plate. Distinct colonies were present with morphologies characteristic of fibroblast (F), amniotic fluid (AF), and epithelial (E) cells (Figure 2). Cells were released from P0 culture by using TrypLE Express™ (Invitrogen) and seeded into fibronectin, vitronectin, or collagen type IV coated flasks/dishes/plates at various densities (50-10,000 cell/cm²). Some of the P0 cells were used for serial dilution cloning. The population doubling time of the fastest growing cells was approximately 24 hrs at early passages. The expanded cells cultured under various media conditions (**Table II**) were analyzed for cell surface markers (**Table III**). Cells were typically split at ~70% confluency and reseeded at 100-10000 cells/cm². RNA was collected at various stages of cell growth and analyzed for embryonic and germ layer markers (**Table V**).

[0146] Amniotic fluid cells of the present invention were present at various gestational ages. **Table VI** lists the presence or absence of AF, E, and F morphologies in amniotic fluid samples obtained at 17 weeks to 41 weeks of gestation.

[0147] Amniotic fluid cells of the present invention were also obtained from amniotic fluid obtained at term (approximately 40 wks of gestation). Amniotic fluid samples were obtained from 38-40 wk deliveries and cultured according to the protocols outlined above. The resulting adherent cell populations displayed very similar characteristics to the cells isolated from 16-22 wks of gestation.

Example 2

Clonal Expansion of the Cells of the Present Invention

[0148] Using methods described in **Example 1**, cells with AF-like morphologies were harvested from P0 cultures using cloning rings. Three distinct populations of cells exhibiting different expression of surface receptors, cytoskeletal proteins, and transcription factors were identified. For sake of clarity, these populations are referred to as AF-I, AF-II, and AF-III cells. Subsequent examples highlight the differences between AF-I, -II, and -III populations.

Example 3

Fluorescence-Activated Cell Sorting (FACS) Analysis

[0149] Adhered cells were removed from culture plates by five-minute incubation with the TRYPLE™ express solution (Gibco, CA). Released cells were resuspended in DMEM supplemented with 10% FBS and recovered by centrifugation, followed by washing and resuspending the cells in a staining buffer consisting of 2% BSA, 0.05% sodium azide (Sigma, MO) in PBS. If appropriate, the cells were Fc-receptor blocked using a 0.1 % γ -globulin (Sigma) solution for 15 min. Aliquots (approximately 10⁵ cells) were incubated with either phycoerythrin (PE) or allophycocyanin (APC) conjugated monoclonal antibodies (5 μ l antibody per 10⁶ cells), as indicated in **Table III-A**, or with an unconjugated primary antibody. Controls included appropriate isotype matched antibodies, non-stained cells, and cells only stained with secondary conjugated antibody. All incubations with antibodies were performed for 30 mins at 4°C, after which the cells were washed with the staining buffer. Samples that were stained with unconjugated primary antibodies were incubated for additional 30 mins at 4°C with secondary conjugated PE or -APC labeled antibodies. See **Table III-B** for a list of secondary antibodies used. Washed cells were pelleted and resuspended in the staining buffer and the cell surface molecules were identified by using a FACS Array (BD Biosciences) by collecting at least 10,000 events.

[0150] For intracellular staining, cells were first fixed for 10 mins with 4% paraformaldehyde, followed by two rinses in the staining buffer, centrifugation of cells and resuspension of the cells in a permeabilization buffer containing 0.5% Triton-X (Sigma) in PBS for 5 mins at room temperature (RT). The permeabilized cells were rinsed twice with a rinsing buffer, centrifuged, and resuspended in the staining buffer and incubated with an appropriate conjugated antibody (5 μ l antibody per 10⁶ cells), for 30 mins at 4°C. Samples that were stained with unconjugated primary antibodies were incubated for additional 30 mins at 4°C with secondary conjugated PE or -APC labeled antibodies (**Table III B**). Washed cells were pelleted and resuspended in the staining buffer and the internal proteins were identified by using a FACSArray (BD Biosciences) by collecting at least 10,000 events. The expression level of examined surface and internal markers is listed in **Table IV A and B**. FACS analysis allowed identification of signature markers to distinguish amniotic fluid cells (AF-I, -II, and -III), fibroblasts (F), and epithelial cells (E) (**Figures 3-5**). **Table IV C** lists the cell surface expression profile of AF-I cells isolated from term (38-40 wks) amniotic fluid. The expression level of cell surface receptors is very similar to AF-I cells isolated from 16-22 wks amniotic fluid.

Example 4**Immunostaining of Undifferentiated Cells**

[0151] 10,000 cells/cm² cells, cultured according to **Example 1**, were seeded into glass bottom 35 mm microwell dishes (Matek Corp, MA) in various growth media. Following three days in culture, the cells were fixed for 10 mins with 4% paraformaldehyde, followed by two rinses in the PBS, and addition of a permeabilization buffer containing 0.5% Triton-X (Sigma) for 5 mins at room temperature (RT) followed by additional three rinses with PBS. The fixed and permeabilized cells were blocked with either 1% bovine serum albumin (BSA) or 4% sera from the species where the secondary antibody was raised in (Goat, donkey, or rabbit). Control samples included reactions with the primary antibody omitted or where the primary antibody was replaced with corresponding immunoglobulins at the same concentration as the primary antibodies. Stained samples were rinsed with a PROLONG® antifade reagent (Invitrogen, CA) containing diamidino-2-phenylindole, dihydrochloride (DAPI) to counter stain the nucleus. Images were acquired using a Nikon Confocal Eclipse C-1 inverted microscope (Nikon, Japan) and a 10-60X objective (**Figures 6 - 8**).

Example 5**PCR Analysis Of Undifferentiated Cells**

[0152] RNA was extracted from cells cultured in the growth media. Total RNA from human pancreas, liver, brain, gut (Ambion, INC.) NTERA cells (human embryonic carcinoma cells line, ATCC), HEK293 cells (ATCC), and human airway epithelia cells (Cambrex) were used as positive controls. Bone marrow derived mesenchymal cells (Cambrex, MD) were used as negative controls for the expression of key genes involved in pancreatic development.

[0153] *RNA extraction, purification, and cDNA synthesis.* RNA samples were purified through its binding to a silica-gel membrane (Rneasy Mini Kit, Qiagen, CA) in the presence of an ethanol-containing, high-salt buffer; while contaminants were washed away. The RNA was further purified while bound to the column by treatment with DNase I (Qiagen, CA) for 15 min. High-quality RNA was then eluted in water. Yield and purity were assessed by A260 and A280 readings on the spectrophotometer. cDNA copies were made from purified RNA using an ABI (ABI, CA) high capacity cDNA archive kit.

[0154] *Real-time PCR amplification and quantitative analysis.* Unless otherwise stated, all reagents were purchased from Applied Biosystems. Real-time PCR reactions were performed using the ABI PRISM® 7000 Sequence Detection System. TAQMAN® UNIVERSAL PCR MASTER MIX® (ABI, CA) was used with 20 ng of reverse transcribed RNA in a total reaction volume of 20 μ l. Each cDNA sample was run in duplicate to correct for pipetting errors. Primers and FAM-labeled TAQMAN® probes were used at concentrations of 200 nM. The level of expression of each target gene was normalized using the pre-developed Applied Biosystem's 18S ribosomal RNA or human glyceraldehydes-3-phosphate dehydrogenase (GAPDH) endogenous control kit. Primers and probes were either designed using ABI PRISM PRIMER EXPRESS™ software or used pre-developed ABI gene analysis kit. For each gene, either one of the primers or the probe were designed to be exon-boundary spanning. This eliminated the possibility of the primers/probe binding to any genomic DNA present. The primer and probe sets are listed as following Nkx2.2 (Hs00159616), Pdx-1 (Hs00426216), Nkx6.1 (Hs00232355), Ngx3 (Hs00360700), Pax4 (Hs00173014), Pax6 (Hs00240871), Insulin (Hs00355773), Glu2 (Hs00165775), glucagon (Hs00174967), Isl-1 (Hs00158126), somatostatin (Hs00174949), FoxA2 (HNF 3-beta) (Hs00232764), HlxB9 (Hs00232128), GATA-4 (Hs00171403), HNF1 β (Hs00172123), Musashi Homolog 1 (Msi-1) (Hs00159291), Hes-1 (Hs00172878), Neurotensin (NTS) (Hs00175048), Cholecystokinin (Hs00174937), AFP (Hs00173490), Secretin (Hs00360814), GIP (Hs00175030), GFAP (Hs00157674), MAP2 (Hs00159041), Olig2 (Hs0037782), Oct-4 (CGACCATCTGCCGCTTTGAG (SEQ ID NO: 1) and CCCCTGTCCCCCA TTCCTA (SEQ ID NO: 2)); Rex-1 (CAGATCCTAACAGCTCGCAGAAT (SEQ ID NO: 3), and GCGTACGCAAATTAAGTCCAGA (SEQ ID NO: 4); Sox17: TGGCGCAGCAGATACCA (SEQ ID NO: 5), AGCGCCTTCCACGACTTG (SEQ ID NO: 6) and CCAGCATCTTGCTCAACTCGGCG (SEQ ID NO: 7); ABCG-2: GTTTATCCGTGGTGTGTCTGG (SEQ ID NO: 8) and CTGAGCTATAGAGGCTGGG (SEQ ID NO: 9); SOX2: ATGCACCGCTACGACGTGA (SEQ ID NO: 10) and CTTTTCACCCCTCCATTT (SEQ ID NO: 11). The remaining primers were designed by using the PRIMERS program (ABI, CA) and are listed in **Table V**. After an initial 50°C for 2 min, and 95°C for 10 min, samples were cycled 40 times in two stages - a denaturation step at 95°C for 15 sec, followed by an annealing/extension step at 60°C for 1 min. Data analysis was carried out using GENEAMP® 7000 Sequence Detection System software. For each primer/probe set, a C_t value was determined as the cycle number at which the fluorescence intensity reached a specific value in the middle of the exponential region of amplification. Relative gene expression levels were calculated using the comparative C_t method. Briefly, for each cDNA sample, the endogenous control C_t value was subtracted from the gene of interest C_t to give the delta C_t value (ΔC_t). The normalized amount of target was calculated as 2^{- ΔC_t} , assuming amplification to be 100% efficiency. Final data were expressed relative to a calibrator sample. The comparative C_t method is only valid if target and endog-

enous control amplification efficiencies are approximately equal. Preliminary validation experiments were therefore performed for each primer/probe set by amplifying serially diluted cDNA samples and determining the ΔC_t values. These ΔC_t values should remain constant across the range of dilutions if amplification efficiencies are equal (**Table V**).

Example 6

Population doubling time

[0155] Passage 6 amniotic fluid cells (AF-I type), isolated and expanded according to Example 1 were seeded at 10000 cells/well of a 24-well tissue culture plate (Corning, MA) in growth media #11. At various time points, cells were removed from three wells of the plate using TRYPLE™ Express (Invitrogen, CA) and counted using a Guava PCA-96 cell analysis system and the VIACOUNT® reagent (Guava, CA). **Figure 10** depicts the growth curve of passage 6 cells cultured under hypoxic conditions (3% O₂). The linear phase of the log plot was used to estimate the population doubling time of the cells. Population doubling time of passage 6 cells was 31 hrs.

[0156] The growth potential of the three cell populations (fibroblast, AF, and epithelial morphology) were compared over long-term cultures. **Figure 11** depicts the growth potential of AF-I, AF-II, AF-III, F, and E cells cultured in media #5. It is clear that F ("fibroblastic" amniotic fluid-derived) cells and AF cells can expand well above 50 population doublings and represent a scalable source for cell therapy applications.

Example 7

Telomere length of AF-I cells

[0157] The telomere length of an AF-I line isolated from a single cell by limited serial dilution was analyzed at passage 12 (approximately 50 population doublings) by using the *Telo TAGGG* Telomere Length Assay (Roche, IN) and following the manufacturer's instruction. The telomere length was analyzed for cells cultured in DMEM-LG + 10% FBS and cells cultured in Amniomax™ (Gibco) see **Figure 12**. DNA from NTERA cells served as a positive control.

Example 8

Karyotype analysis

[0158] The karyotype of AF cells, isolated from multiple donors at passage 8-10 (approximately 30 population doublings), was determined by G-band analysis. Five karyotypes were prepared and cytogenetic analysis showed that the cells had a normal autosomes and a modal chromosome number of 46. All cells analyzed also contained the X and Y-chromosomes confirming their fetal origin. **Figure 13** depicts karyotypes of amniotic fluid-derived cells (AF-I, AF-II, and AF-III) isolated from amniotic fluid obtained from 16-22 weeks of gestation.

Example 9

Expansion potential of AF cells derived from term amniotic fluid

[0159] **Figure 14** depicts the expansion potential of an AF-I cell morphology derived from term amniotic fluid (~38 weeks) and cultured in media #5. The expansion potential is very similar to the AF-I cells isolated from 16-22 wks amniotic fluid.

Example 10

Microarray analysis of fibroblast, epithelial, and amniotic fluid morphology cells

[0160] Total RNA was isolated from passage 9-11 amniotic fluid-derived fibroblast cells (F), amniotic fluid-derived epithelial cells (E), amniotic fluid-derived amniotic fluid cells (AF-I, -II, and -III lines), and amniotic fluid at term (AF term) using an RNeasy mini kit (Qiagen). The sample preparation, hybridization, and image analysis was performed according to the CodeLink™ System (GE Healthcare, Amersham Biosciences, NJ). CodeLink™ Human Whole Genome arrays were used. It is comprised of approximately 55 000 30-mer probes designed to conserved exons across the transcripts of targeted genes. The chip contains ~45000 unique Unigene IDs. Following normalization and a log transformation, data analysis was performed using OmniViz® software (MA) and GENESIFTER (VizX Labs, WA). The variance stabilizing transformation along with cross sample normalization was applied to the log transformed array dataset. The variability

within each cell line and among the different cell lines was compared using the Pearson correlation coefficient. For all the samples analyzed, the correlation coefficient within a cell line was higher as compared to those between the lines. Variance in gene expression profiles between the different cell types along with the correlation coefficient between the lines are depicted in **Figure 15**. Significant differences in gene expression between the cell types were evaluated using analysis of variance and an F-test with adjusted P-value (Benjamini and Hochberg correction) of ≤ 0.05 . **Tables VII A-G** list the genes that are differentially expressed at least 5-fold between the various cell types.

Example 11

Differentiation of cells into intestinal-like cells

[0161] The AF-I line, AFCA007 Clone A passage 14, was cultured at 10,000 cells/cm² in AMNIOMAX™. At confluency, the cells were further treated for 2 weeks with a daily dose of 10 micro molar retinoic acid (RA) in AMNIOMAX™ media. RNA was collected at day 14 and expression of intestinal hormones (secretin, neurotensin, gastric inhibitory peptide (GIP), cholecystokinin, somatostatin, and gastrin) was assessed by using real-time PCR as outlined in Example 4. Intestinal RNA (Ambion) was used to assess relative levels of expression using the $\Delta\Delta C_t$ method. **Table VIII** lists the C_t values and the relative level of expression of the intestinal hormones in treated and untreated samples. As shown in **Table VIII**, addition of RA enhanced expression of the gut hormones.

Example 12

Differentiation of cells into multiple endocrine lineages

[0162] Cells from the cell line AFCA007 Clone A (AF-I) at passage 8 were embedded in collagen type I (Becton Dickinson, CA) with 1% growth-factor reduced matrigel matrix (Becton Dickinson), and seeded into 6-well transwell insert at 5×10^5 cells per well. The bottom well was seeded with human aortic endothelial cells passage 6 (Cambrex, MD). Cells were cultured with DMEM medium supplemented with 5% FBS and growth factors, which includes Cyclopamine, bFGF, EGF, BMP4-7, Activin A, Exendin 4, FGF4, all-trans retinoic acid and γ -secretase inhibitors for 14 days. Cultures were fed every other day. Cells treated by all-trans retinoic acid showed the up-regulation of alpha-fetoprotein (AFP). Treatment of cells with Activin A, BMP4, or the γ -secretase inhibitor L-685,458 up-regulated the expression of HNF-3 beta. Treatment of cells with BMPs at high concentration, 50 ng/ml, also up-regulated the GATA4 expression. Treatment of cells with FGF4 at 50 ng/ml showed an up-regulation of PDX-1 expression (**Table IX**).

[0163] Cells from the cell line AFCA004 (E morphology) at passage 6 were seeded at 5×10^5 cells per well of 6-well culture plates and treated with conditioned medium from confluent PANC-1 cells (ATCC, VA) in combination with different growth factors. Basic FGF, EGF and combination of bFGF and EGF enhanced the expression of HNF-3 beta ~100 fold over untreated cells. Basic FGF, EGF and BMPs also stimulated somatostatin expression after 14 days treatment (**Figure 16, panels a&b**).

[0164] Taken together, these results suggest that AF cells could be differentiated into pancreatic, hepatic or intestinal lineage by treating the cells with different growth factors.

Example 13

Modulation of the expression of endoderm makers by inhibiting the Notch pathway

[0165] Cells from the cell line AFCA007 (AF-I) at passage 8 were treated with a range of concentrations of the notch pathway inhibitor L-685,458 (Sigma, MO) for 3 to 5 days. The cytotoxicity of L-685,458 was determined by measuring cell viability by a MTS assay (Promega, WI). Real-time PCR analysis was performed to analyze Hes-1 expression after the treatment. We found that L-685,458 showed a dose-dependent inhibitory effect on Hes-1 expression, Hes-1 is the downstream direct target of Notch pathway (**Figure 17, panel a**). No effect on cell viability, as determined by the MTS assay, was observed following L-685,458 treatment of up to 10 μ M for 5 days (**Figure 17, panel b**).

Example 14

Cytokine antibody array analysis for AF I and AF II cells

[0166] AFCA007 A (AF-I) and AFCA015 C (AF-II) at passage 10 were grown to approximately 70 % confluency and then cell lysates was collected using mammalian cell lysis kit (Sigma-Aldrich, MO). Cytokine array analysis was completed using Cytokine Array panels provided by RayBiotech, GA (<http://www.raybiotech.com/>). **Table X** lists cytokine, cytokine

and growth factor receptor expression following normalization of the data and background subtraction. For each panel, positive and negative controls are also included. The panels were run for two different samples per cell type.

TABLE I: COMPARISON OF THE CELL OF THE PRESENT INVENTION WITH AMNIOTIC FLUID-DERIVED CELLS OF THE ART.

Marker	Cell 1	Cell 2	Cell 3	Cell 4
CD10				-
CD105	+	+	+	Weak
CD11			-	-
CD117	Weak	-	-	-
CD 13		+		
CD29 (Beta 1 integrin)				+
CD31			-	
CD34	Weak	-	-	-
CD44		+	+	+
CD45		-	-	
CD49b (Alpha 2 integrin)		+		
CD49e (Alpha 5 integrin)			+	
CD73			+	+
CD166			+	
CD90	+		+	+
HLA ABC	+	+	+	+
HLA D		-	-	-
Oct-4	+	+		+
SSEA-1	-	-		
SSEA-3	+	+		
SSEA-4	+	+		
Stro-1		-		
Telomerase	+			
TRA 1-60		+		
TRA 1-81		+		
TRA 1-85				
TRA 2-49				
TRA 2-54		+		

TABLE II. MEDIA FORMULATIONS USED TO CULTURE AMNIOTIC-FLUID DERIVED CELLS.

Media formulation	
1	DMEM-LG + ITS-X + 1% P/S + 2% FBS + 25 ng/ml bFGF
2	Advanced DMEM + ITS-X + 1% P/S + 1% FBS + 25 ng/ml bFGF
3	Alpha MEM: MCDB 153 (1:1) + ITS-X + 1% P/S + 1% FBS + 25 ng/ml bFGF

EP 1 780 267 B1

(continued)

Media formulation	
4	Defined Keratinocyte growth media (Gibco, NY)
5	AMNIOMAX™ complete media (Gibco, NY)
6	Chang B/C (Irvine Scientific, CA)
7	Chang D (Irvine Scientific, CA)
8	DM-KNOCKOUT™ media (Invitrogen, CA), supplemented with 20% KNOCKOUT™ serum replacement (Invitrogen, CA), 10 ng/ml bFGF
9	DMEM- Low glucose, supplemented with 20% FBS
10	DMEM- Low glucose, supplemented with 5% FBS
11	DMEM- low glucose /MCDB 201 medium (1:1), supplemented with 2% defined FBS, ITS-X, 1nM dexamethasone, 100 mM ascorbic acid 2-phosphate, 10ng/ml EGF, 10ng/ml PDGF-bb and 100 mM 2-mercaptoethanol
12	20% KNOCKOUT™ serum replacement + 80% KNOCKOUT™ DMEM, supplemented with 1 mM L-glutamine, 1% non-essential amino acids and 0.1 mM 2-mercaptoethanol
13	The medium may be conditioned overnight, on human or murine embryonic fibroblasts, human bone marrow derived stromal cells, or human placenta derived cells and supplemented with 4 ng/ml bFGF
14	high glucose DMEM, supplemented with 20% defined FBS with 0.1 mM 2-mercaptoethanol
15	Williams' E medium, 2% FBS, 1X ITS, 55uM βme, 1X Glutamax (Gibco), 1%P/S, 10ng/ml EGF, 4ng/ml bFGF, 4ng/ml dexamethasone
16	1:1 DMEM-LG/MCDB 201, 2% FBS, ITS-X, βme 55uM, 100uM ascorbic acid-2-phosphate, 4ng/ml bFGF, 10ng/ml EGF, 4ng/ml dexamethasone

TABLE III A: ANTIBODIES TO SURFACE RECEPTORS.

Antibody	Supplier	Isotype	Clone
CD117 (c-Kit)	Santa Cruz Biotechnology (CA)	Goat IgG	M-14
CD117 (c-Kit)	Santa Cruz Biotechnology (CA)	Mouse IgG I	104D2
CD117 (c-Kit)	BD Pharmingen (CA)	Mouse IgG1, kappa	YB5.B8
CD24	BD Pharmingen (CA)	Mouse IgG2A, Kappa	ML5
CD44	BD Pharmingen (CA)	Mouse IgG2b, Kappa	G44-26
CD45	BD Pharmingen (CA)	Mouse IgG1, Kappa	Hi30
CD49f	BD Pharmingen (CA)	Rat IgG2A, Kappa	G0H3
CD73	BD Pharmingen (CA)	Mouse IgG1, Kappa	AD2
CD10	BD Pharmingen (CA)	Mouse IgG1, Kappa	HI10a
CD105	Santa Cruz (CA)	Mouse IgG1	P3D1
CD49b (Alpha 2 integrin)	BD Pharmingen (CA)	Mouse IgG2a, Kappa	121-H6
Alpha 3 integrin	Santa Cruz (CA)	Mouse IgG1	P1B5
Alpha 4 integrin (CD49d)	BD Pharmingen (CA)	Mouse IgG1, Kappa	9F10
Alpha 5 intgerin (CD49e)	BD Pharmingen (CA)	Mouse IgG1, Kappa	IIA1
CD49f (Alpha 6 integrin)	BD Pharmingen (CA)	Rat IgG2a	GoH3

EP 1 780 267 B1

(continued)

Antibody	Supplier	Isotype	Clone
CD29 (Beta 1 integrin)	BD Pharmingen (CA)	Mouse IgG1, Kappa	MAR4
Beta 3 integrin	Santa Cruz (CA)	Mouse IgG1	Y2/51
Alpha V Beta 3 integrin (CD51/61)	BD Pharmingen (CA)	Mouse IgG1, Kappa	23C6
SSEA-3	Chemicon (CA)	Mouse IgG3	MC-631
SSEA-4	Chemicon (CA)	Rat IgM	MC-813-70
TRA 1-60	Chemicon (CA)	Mouse IgM	TRA 1-60
TRA 1-81	Chemicon (CA)	Mouse IgM	TRA 1-81
TRA 1-85	Chemicon (CA)	Mouse IgG1	TRA 1-85
TRA 2-54	Chemicon (CA)	Mouse IgG1	TRA 2-54
EGF r	BD Pharmingen (CA)	Mouse IgG2b, Kappa	EGFR1
EpCAM	BD Pharmingen (CA)	Mouse IgG1	EBA-1
HLA ABC	BD Pharmingen (CA)	Mouse IgG1, Kappa	G46-2.6
HLA DR	BD Pharmingen (CA)	Mouse IgG2b, Kappa	TU36
CD90	BD Pharmingen (CA)	Mouse IgG1, kappa	5E10

TABLE III B: LIST OF SECONDARY CONJUGATED ANTIBODIES USED FOR FACS AND IMMUNOSTAINING ANALYSIS.

Secondary conjugated antibody	Supplier	Dilution
Goat Anti-Mouse IgG APC conjugated	Jackson ImmunoResearch (PA)	1:200
Goat Anti-Mouse IgG PE conjugated	Jackson ImmunoResearch (PA)	1:200
Donkey anti-rabbit PE or - APC conjugated	Jackson ImmunoResearch (PA)	1:200
Donkey anti-goat PE or - APC conjugated	Jackson ImmunoResearch (PA)	1:200
Goat anti-mouse IgM PE	SouthernBiotech (AL)	1:200
Goat anti-Rat IgM PE	SouthernBiotech (AL)	1:200
Goat anti-mouse IgG3 PE	SouthernBiotech (AL)	1:200

TABLE III C: LIST OF PRIMARY ANTIBODIES USED FOR IMMUNOSTAINING ANALYSIS.

Antibody	Supplier	Dilution
Beta III Tubulin	R&D Systems (MN)	1:100
C-Kit	Santa Cruz Biotechnology (CA)	1:100
Cytokeratin 18	Sigma (MO)	1:100
Cytokeratin 8	Sigma (MO)	1:100
Cytokeratin-7	Santa Cruz Biotechnology (CA)	1:100
Cytokeratin 19	Sigma (MO)	1:100
E-Cadherin	BD Transduction Laboratories (CA)	1:100

EP 1 780 267 B1

(continued)

Antibody	Supplier	Dilution
FOXA1	Chemicon International (CA)	1:100
GATA-6	R&D Systems (MN)	1:100
HES-1	Chemicon International (CA)	1:100
HES-2	Chemicon International (CA)	1:100
HNF-1 alpha	BD Transduction Laboratories (CA)	1:100
HNF-1 beta	BD Transduction Laboratories (CA)	1:100
Musashi-1	Chemicon International (CA)	1:100
Nestin	R&D Systems (MN)	1:100
Pan-Cytokeratin (CK, 4, 5, 6, 10, 13, and 18)	Santa Cruz Biotechnology (CA)	1:100
SOX-17	R&D Systems (MN)	1:100
SSEA-4	Santa Cruz Biotechnology (CA)	1:100
TRA 1-81	Santa Cruz Biotechnology (CA)	1:100
Vimentin	Santa Cruz Biotechnology (CA)	1:100
ZO-1	BD Transduction Laboratories (CA)	1:100

TABLE IV A: CELL SURFACE RECEPTORS ON FIBROBLAST, AMNIOTIC FLUID, AND EPITHELIAL CELLS DERIVED FROM AMNIOTIC FLUID AND GROWN UNDER HYPOXIC CONDITIONS

	Fibroblast cells (F)	Amniotic fluid cells (AF-I)	Amniotic fluid cells (AF-II)	Amniotic fluid cells (AF-III)	Epithelial cells (E)
Alpha 3 integrin	+	+	+	+	+
Beta 1 integrin	+	+	+	+	+
Beta 3 integrin	-	+	Weak	Weak	+
Alpha V Beta 3 (CD51/ 61)	-	+	Weak	Weak	+
C-Met		+	+	+	
CD24	Weak	+	Weak	Weak	+
CD10	+	Weak	-	-	-
CD105	+	Weak	+	+	-
CD117	-	-	-	-	-
CD13	+	Weak	+	+	-
CD44	+	+	+	+	+
CD73	+	+	+	+	+
CD90	+	+	+	+	+
EpCAM	-	+	-	-	+
HLA ABC	+	+	+	+	+
HLA II-DR	-	-	-	-	-
TRA2-49	-	-	-	-	-
SSEA-4	+	+	+	+	+

(continued)

	Fibroblast cells (F)	Amniotic fluid cells (AF-I)	Amniotic fluid cells (AF-II)	Amniotic fluid cells (AF-III)	Epithelial cells (E)
SSEA-3	-	+	-	-	+
TRA1-60	-	+	-	-	+
TRA 1-81	-	+	-	-	+

TABLE IV B: INTRACELLULAR PROTEINS OF VARIOUS AMNIOTIC FLUID CELL TYPES GROWN IN MEDIA #5

Marker	Fibroblast cells (F)	Amniotic fluid cells (AF-I)	Amniotic fluid cells (AF-II)	Amniotic fluid cells (AF-III)	Epithelial cells (E)
CK-19	-	+	-	-	+
Vimentin	+	+	+	+	+
Smooth muscle actin	Weak	Weak	Weak	Weak	+
Beta III tubulin	+	+	+	+	+
Pan cytokeratin	-	+	+	-	+
Cytokeratin 7	-	+			
Cytokeratin 8	-	+			
Cytokeratin 18	-	+			
Nestin	-	+	+	+	Weak

TABLE IV C: CELL SURFACE RECEPTORS ON AMNIOTIC FLUID CELLS DERIVED FROM TERM AMNIOTIC FLUID AND GROWN UNDER HYPOXIC CONDITIONS

	Amniotic fluid cells (AF-I)
Alpha 3 integrin	+
Beta 1 integrin	+
Beta 3 integrin	+
Alpha V Beta 3 (CD51/61)	+
CD10	Weak
CD105	Weak
CD117	-
CD13	Weak
CD44	+
CD73	+
CD90	+
EpCAM	+
HLA ABC	+
HLA II-DR	-
SSEA-4	+
TRA1-60	+

(continued)

	Amniotic fluid cells (AF-I)
TRA 1-81	+

TABLE V: PCR ANALYSIS OF THE AMNIOTIC FLUID-DERIVED CELLS OF THE PRESENT INVENTION

a. Pluripotency markers						
Markers	Fibroblast Cells	Epithelial cells	AF-I Cells	AF-II Cells	AF-III Cells	Ntera-1
Oct3/4	-	-	-	-	-	+
Sox-2	-	-	-	-	-	+
Rex-1	-	+	+/-			+
hTERT	-	+/-	-	-	-	+
b. Endocrine markers						
Cell Markers	Fibroblast cells	Epithelial cell	AF_I cells	AF_II cells	AF_III cells	
Sox 17	Weak	+	+	-	-	
GATA-6	-	+	+	+	+	
GATA-4	Weak	+	Weak	-	-	
HNF 1 Beta	-	+	+	+	-	
HNF3 beta	-	+	Weak	-	-	
Pdx-1	-	-	-	-	-	
NGN-3	-	-	-	-	-	
Musashi-1	+	+	+			
HES-1	+	+	+	+	+	
NeuroD1	-	-	-	-	-	
Pax 4	-	-	-			
Pax6	+	+	+	-	-	
Secretin	-	-	-	-	-	
Gastric inhibitory peptide (GIP)	+	+	+	-	-	
Glucagon	-	-	-	-	-	
Somatostatin	+	+	+	-	-	
Cholecystokinin	-	-	-	-	-	
Gastrin	-	-	-	-	-	
Insulin		-	Weak	-	-	
Nestin	+	+	+	+	+	
*Weak: CT values of 35-38 cycles +: CT value <35 cycles						

TABLE VI: PRESENCE OF ALL AF, E, AND F MORPHOLOGIES AT VARIOUS GESTATIONAL AGES IN SECOND TRIMESTER AMNIOTIC FLUID SAMPLES.

AF designation	Gestation age	Presence of AF-like morphology	Presence of E-like morphology	Presence of F-like morphology
AFCA001	17 wks	+	+	+
AFCA002	18 wks	+	+	-
AFCA004	18 wks	+	+	-
AFCA008	18 wks	+	+	-
AFDX001	19 wks	+	+	+
AFDX021	20 wks	+	+	+
AFDX022	20 wks	+	+	+
AFCA007	19 wks	+	+	-
AFCA009	18 wks	+	+	-
AFCA010	18 wks	+	+	-
AFCA011	19 wks	+	+	-
AFCA017	16 wks	+	+	-
AFPN003	20 wks	+	+	-
AFPN004	20 wks	+	+	-
AFND001	41 wks	+	+	-

TABLE VII A: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN FIBROBLAST VERSUS AF-I CELLS

Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
NM_002421	Homo sapiens matrix metalloproteinase 1 (interstitial collagenase) (MMP1), mRNA	434.28	UP	1.09E-03
NM_144594	Homo sapiens hypothetical protein FLJ32942 (FLJ32942), mRNA	207.74	UP	2.52E-03
NM_020927	Homo sapiens KIAA1576 protein (KIAA1576), mRNA	200.11	UP	5.06E-04
NM_001451	Homo sapiens forkhead box F1 (FOXF1), mRNA	183.69	UP	1.52E-03
AK021543	Homo sapiens cDNA FLJ11481 fis, clone HEMBA1001803	131.1	UP	3.11E-03
NM_007036	Homo sapiens endothelial cell-specific molecule 1 (ESM1), mRNA	120.15	UP	2.07E-03
NM_002448	Homo sapiens msh homeo box homolog 1 (Drosophila) (MSX1), mRNA	112.45	UP	7.69E-04
NM_152270	Homo sapiens hypothetical protein FLJ34922 (FLJ34922), mRNA	109.54	UP	1.63E-03
NM_000474	Homo sapiens twist homolog 1 (acrocephalosyndactyly 3; Saethre-Chotzen syndrome) (Drosophila) (TWIST1), mRNA	102.84	UP	1.72E-03
AK122739	Homo sapiens cDNA FLJ16260 fis, clone IMR322006947, highly similar to Rattus norvegicus mRNA for dHand protein	98.04	UP	4.82E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AV702977	AV702977 ADB Homo sapiens cDNA clone ADBCVD08 5, mRNA sequence	93.67	UP	2.91 E-03
10	AK026784	Homo sapiens cDNA: FLJ23131 fis, clone LNG08502	87.77	UP	1.52E-03
	NM_000710	Homo sapiens bradykinin receptor B1 (BDKRB1), mRNA	86.86	UP	4.77E-03
	NM_000609	Homo sapiens chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1) (CXCL12), mRNA	77.16	UP	2.97E-03
15	NM_030781	Homo sapiens collectin sub-family member 12 (COLEC12), transcript variant II, mRNA	64.54	UP	8.99E-05
	NM_032638	Homo sapiens GATA binding protein 2 (GATA2), mRNA	63.54	UP	3.35E-03
	NM_000362	Homo sapiens tissue inhibitor of metalloproteinase 3 (Sorsby fundus dystrophy, pseudoinflammatory) (TIMP3), mRNA	61.32	UP	5.06E-03
20	NM_198148	Homo sapiens carboxypeptidase X (M14 family), member 2 (CPXM2), mRNA	57.01	UP	4.00E-03
	AL831863	Homo sapiens mRNA; cDNA DKFZp761J2017 (from clone DKFZp761J2017)	55.07	UP	7.57E-03
25	NM_002091	Homo sapiens gastrin-releasing peptide (GRP), mRNA	53.77	UP	9.01E-03
	NM_020404	Homo sapiens CD164 sialomucin-like 1 (CD164L1), mRNA	51.44	UP	2.22E-04
	NM_014178	Homo sapiens syntaxin binding protein 6 (amisyn) (STXBP6), mRNA	47.27	UP	7.17E-03
30	NM_205855	Homo sapiens HWKM1940 (UNQ1940), mRNA	42.6	UP	2.31E-03
	BX089019	BX089019 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998K243513 ; IMAGE:1391375, mRNA sequence	42.17	UP	8.50E-03
35	AI124557	am58g02.x1 Johnston frontal cortex Homo sapiens cDNA clone IMAGE:1539794 3, mRNA sequence	37.2	UP	8.24E-03
	NM_139211	Homo sapiens homeodomain-only protein (HOP), transcript variant 2, mRNA	34.14	UP	1.45E-04
	NM_002593	Homo sapiens procollagen C-endopeptidase enhancer (PCOLCE), mRNA	30.85	UP	2.64E-03
40	BQ020357	UI-H-ED0-axk-p-07-0-UI.s1 NCI_CGAP_ED0 Homo sapiens cDNA clone IMAGE:5830134 3, mRNA sequence	30.74	UP	5.91E-03
	CA843592	ir49c12.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6548544 3, mRNA sequence	29.57	UP	6.22E-05
45	NM_002852	Homo sapiens pentaxin-related gene, rapidly induced by IL-1 beta (PTX3), mRNA	29.35	UP	3.55E-03
	NM_000089	Homo sapiens collagen, type I, alpha 2 (COL1A2), mRNA	28.94	UP	4.61E-04
	CD677332	ho15f06.y1 Human Trabecular meshwork cDNA: hohphq Homo sapiens cDNA clone ho15f06 5, mRNA sequence	28.67	UP	1.15E-04
50	BC030692	Homo sapiens ELAV (embryonic lethal, abnormal vision, Drosophila)-like 2 (Hu antigen B), mRNA (cDNA clone MGC:26319 IMAGE:4826082), complete cds	28.22	UP	7.64E-03
	AK021543	Homo sapiens cDNA FLJ11481 fis, clone HEMBA1001803	27.4	UP	1.70E-02
55	AI962169	wq45c10.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474226 3, mRNA sequence	27.1	UP	8.11E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_000685	Homo sapiens angiotensin II receptor, type 1 (AGTR1) transcript variant 1, mRNA	26.59	UP	4.86E-03
10	AI422199	tf58d04.x1 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:2103463 3, mRNA sequence	26.11	UP	1.04E-02
	NM_001002295	Homo sapiens GATA binding protein 3 (GATA3), transcript variant 1, mRNA	24.62	UP	3.24E-03
15	NM_000396	Homo sapiens cathepsin K (pseudodysostosis) (CTSK), mRNA	24.58	UP	4.37E-05
	NM_001442	Homo sapiens fatty acid binding protein 4, adipocyte (FABP4), mRNA	24.37	UP	8.07E-03
	NM_004460	Homo sapiens fibroblast activation protein, alpha (FAP), mRNA	24.15	UP	1.93E-04
20	AB067499	Homo sapiens mRNA for KIAA1912 protein, partial cds	23.83	UP	2.51E-03
	NM_032777	Homo sapiens G protein-coupled receptor 124 (GPR124), mRNA	22.29	UP	3.83E-03
	U83115	Human non-lens beta gamma-crystallin like protein (AIM1) mRNA, partial cds	21.93	UP	5.28E-04
25	AY335938	Homo sapiens homeodomain protein IRXA1 (IRX1) mRNA, complete cds	21.54	UP	2.02E-02
	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	21.53	UP	2.20E-03
30	W38393	zb15c07.r1 Soaresfetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:302124 5, mRNA sequence	21.16	UP	8.46E-05
	NM_021637	Homo sapiens transmembrane protein 35 (TMEM35), mRNA	20.78	UP	3.74E-04
	AK091731	Homo sapiens cDNA FLJ34412 fis, clone HEART2002432	20.71	UP	1.19E-02
35	NM_032883	Homo sapiens chromosome 20 open reading frame 100 (C20orf100), mRNA	20.29	UP	2.33E-04
	NM_005110	Homo sapiens glutamine-fructose-6-phosphate transaminase 2 (GFPT2), mRNA	19.69	UP	9.13E-05
40	NM_024633	Homo sapiens chromosome 14 open reading frame 139 (C14orf139), mRNA	19.44	UP	1.05E-02
	NM_004811	Homo sapiens leupaxin (LPXN), mRNA	18.74	UP	1.74E-05
	NM_153183	Homo sapiens nudix (nucleoside diphosphate linked moiety X)-type motif 10 (NUDT10), mRNA	18.51	UP	9.88E-03
45	NM_014459	Homo sapiens protocadherin 17 (PCDH17), mRNA	18.39	UP	1.39E-03
	BX115659	BX115659 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGp998C204119 ; IMAGE:1623883, mRNA sequence	18.15	UP	4.07E-04
50	NM_018013	Homo sapiens hypothetical protein FLJ10159 (FLJ10159), mRNA	17.64	UP	1.54E-02
	BX112628	BX112628 Soares_fetal_lung_NbHL19W Homosapiens cDNA clone IMAGp998A09669 ; IMAGE:299024, mRNA sequence	16.09	UP	2.35E-02
55	NM_016428	Homo sapiens ABI gene family, member 3 (ABI3), mRNA	16.07	UP	1.15E-02
	NM_012449	Homo sapiens six transmembrane epithelial antigen of the prostate (STEAP), mRNA	16.07	UP	1.49E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	15.98	UP	6.30E-05
10	AF052115	Homo sapiens clone 23688 mRNA sequence	15.92	UP	1.56E-02
	NM_004787	Homo sapiens slit homolog 2 (Drosophila) (SLIT2), mRNA	15.9	UP	4.75E-04
	AK091336	Homo sapiens cDNA FLJ34017 fis, clone FCBBF2002626	15.59	UP	2.68E-02
15	NM_006329	Homo sapiens fibulin 5 (FBLN5), mRNA	15.59	UP	7.87E-03
	NM_000963	Homo sapiens prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2), mRNA	15.57	UP	8.59E-04
20	NM_001146	Homo sapiens angiopoietin 1 (ANGPT1), transcript variant 1, mRNA	14.89	UP	9.40E-04
	NM_018431	Homo sapiens docking protein 5 (DOK5), transcript variant 1, mRNA	14.56	UP	8.69E-03
	NM_017577	Homo sapiens hypothetical protein DKFZp434C0328 (DKFZp434C0328), mRNA	14.47	UP	4.20E-03
25	H85497	yv88b07.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:249781 5, mRNA sequence	13.8	UP	1.13E-04
	NM_001147	Homo sapiens angiopoietin 2 (ANGPT2), mRNA	13.76	UP	2.64E-02
	NM_000090	Homo sapiens collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant) (COL3A1 mRNA	13.63	UP	1.65E-02
30	NM_007289	Homo sapiens membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10) (MME), transcript variant 2b, mRNA	13.35	UP	1.86E-02
35	NM_016229	Homo sapiens cytochrome b5 reductase b5R.2 (CYB5R2), transcript variant 1, mRNA	13.35	UP	9.80E-03
	NM_000810	Homo sapiens gamma-aminobutyric acid (GABA) A receptor, alpha 5 (GABRA5), mRNA	13.07	UP	2.71 E-02
40	NM_002518	Homo sapiens neuronal PAS domain protein 2 (NPAS2), mRNA	12.99	UP	1.04E-02
	AK093256	Homo sapiens cDNA FLJ35937 fis, clone TEST12011480	12.93	UP	2.09E-03
	NM_005127	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 2 (activation-induced) (CLECSF2), mRNA	12.82	UP	2.71E-02
45	NM_006209	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 2 (autotaxin) (ENPP2), mRNA	12.62	UP	2.60E-02
50	W03013	za02c04.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:291366 5 similar to contains THR.t3 THR repetitive element ;, mRNA sequence	12.43	UP	7.69E-04
	AF131813	Homo sapiens clone 24970 mRNA sequence	12.17	UP	2.57E-02
	BQ025821	UI-1-BB1 p-aye-f-10-0-ULs1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1 p-aye-f-10-0-UI 3, mRNA sequence	12.15	UP	1.30E-02
55					

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AW445209	UI-H-BI3-akc-g-11-0-UI.s1 NCI-CGAP-Sub5 Homo sapiens cDNA clone IMAGE:2733908 3, mRNA sequence	12.09	UP	8.91 E-04
10	NM_033292	Homo sapiens caspase 1, apoptosis-related cysteine protease (interleukin 1, beta, convertase) (CASP1 transcript variant alpha, mRNA	11.63	UP	2.67E-02
15	NM_170744	Homo sapiens unc-5 homolog B (C. elegans) (UNC5B), mRNA	11.62	UP	3.68E-02
	NM_012204	Homo sapiens general transcription factor IIIC, polypeptide 4, 90kDa (GTF3C4), mRNA	11.59	UP	3.11E-02
	NM_002531	Homo sapiens neurotensin receptor 1 (high affinity) (NTSR1), mRNA	11.33	UP	2.65E-02
20	BG118019	602351269F1 NIH_MGC_90 Homo sapiens cDNA clone IMAGE:4446065 5, mRNA sequence	11.16	UP	8.51 E-05
	NM_014802	Homo sapiens KIAA0528 gene product (KIAA0528), mRNA	11.05	UP	2.84E-02
25	BX647541	Homo sapiens mRNA; cDNA DKFZp686P0492 (from clone DKFZp686P0492)	10.9	UP	1.65E-03
	NM_020809	Homo sapiens Rho GTPase activating protein 20 (ARHGAP20), mRNA	10.85	UP	2.74E-02
	NM_016307	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	10.85	UP	7.71E-05
30	NM_057179	Homo sapiens twist homolog 2 (Drosophila) (TWIST2), mRNA	10.83	UP	7.22E-05
	AK128325	Homo sapiens cDNA FLJ46467 fis, clone THYMU3022668	10.82	UP	1.20E-02
35	BE866150	601679068F1 NIH_MGC_53 Homo sapiens cDNA clone IMAGE:3961768 5, mRNA sequence	10.74	UP	2.16E-02
	BX109483	BX109483 NCI_CGAP_Ov23 Homo sapiens cDNA clone IMAGp998C165481 ; IMAGE:2216391, mRNA sequence	10.72	UP	2.65E-02
40	C02345	HUMGS0007544 Human adult(K.Okubo) Homo sapiens cDNA, mRNA sequence	10.64	UP	2.15E-03
	NM_002961	Homo sapiens S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine placental homolog) (S100A4), transcript variant 1, mRNA	10.61	UP	1.87E-04
45	NM_031908	Homo sapiens C1q and tumor necrosis factor related protein 2 (C1QTNF2), mRNA	10.13	UP	3.04E-02
	NM_013387	Homo sapiens ubiquinol-cytochrome c reductase complex (7.2 kD) (HSPC051), transcript variant 1, mRNA	9.95	UP	4.17E-04
50	BC039369	Homo sapiens, clone IMAGE:5271073, mRNA, partial cds	9.76	UP	4.82E-03
	BQ934941	AGENCOURT_8810373 NIH MGC 101 Homo sapiens cDNA clone IMAGE:6429485 5, mRNA sequence	9.53	UP	2.21E-02
	NM_006670	Homo sapiens trophoblast glycoprotein (TPBG), mRNA	9.37	UP	2.37E-04
55	BC046364	Homo sapiens flavoprotein oxidoreductase MICAL3, mRNA (cDNA clone IMAGE:5737121), with apparent retained intron	9.09	UP	2.55E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_024600	Homo sapiens chromosome 16 open reading frame 30 (C16orf30), mRNA	9.09	UP	4.32E-02
10	AK023647	Homo sapiens cDNA FLJ13585 fis, clone PLACE1009150	8.81	UP	2.25E-03
	NM_017805	Homo sapiens Ras interacting protein 1 (RASIP1), mRNA	8.78	UP	2.38E-03
15	NM_152399	Homo sapiens hypothetical protein FLJ30834 (FLJ30834), mRNA	8.73	UP	2.18E-02
	NM_002851	Homo sapiens protein tyrosine phosphatase, receptor-type, Z polypeptide 1 (PTPRZ1), mRNA	8.61	UP	5.08E-02
	AK024653	Homo sapiens cDNA: FLJ21000 fis, clone CAE03359	8.51	UP	6.29E-03
20	NM_020226	Homo sapiens PR domain containing 8 (PRDM8), mRNA	8.48	UP	1.49E-04
	NM_012307	Homo sapiens erythrocyte membrane protein band 4.1-like 3 (EPB41 L3), mRNA	8.41	UP	7.37E-03
	NM_002203	Homo sapiens integrin, alpha 2 (CD49B, alpha 2 subunit of VLA-2 receptor) (ITGA2), mRNA	8.3	UP	8.37E-05
25	NM_203370	Homo sapiens similar to RIKEN cDNA 6530418L21 (LOC389119), mRNA	8.27	UP	8.21 E-05
	NM_015192	Homo sapiens phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 1, mRNA	8.26	UP	2.64E-03
30	A056725	Homo sapiens cDNA FLJ32163 fis, clone PLACE6000371	8.22	UP	1.68E-04
	NM_005328	Homo sapiens hyaluronan synthase 2 (HAS2), mRNA	8.2	UP	4.39E-02
	NM_012294	Homo sapiens Rap guanine nucleotide exchange factor (GEF) 5 (RAPGEF5), mRNA	8.08	UP	4.94E-02
35	AA187037	zp58e04.r1 Stratagene endothelial cell 937223 Homo sapiens cDNA clone IMAGE:624414 5, mRNA sequence	7.95	UP	1.09E-02
	NM_001463	Homo sapiens frizzled-related protein (FRZB), mRNA	7.94	UP	1.31E-03
	BM468332	AGENCOURT_6432296 NIH_MGC_71 Homo sapiens cDNA clone IMAGE:5535773 5, mRNA sequence	7.94	UP	7.42E-05
40	AK096661	Homo sapiens cDNA FLJ39342 fis, clone OCBBF2018873	7.82	UP	2.30E-03
	NM_001849	Homo sapiens collagen, type VI, alpha 2 (COL6A2), transcript variant 2C2, mRNA	7.67	UP	1.10E-04
45	BC071787	Homo sapiens cDNA clone IMAGE:4610527, partial cds	7.59	UP	1.14E-04
	NM_012105	Homo sapiens beta-site APP-cleaving enzyme 2 (BACE2), transcript variant a, mRNA	7.58	UP	2.00E-04
	NM_014217	Homo sapiens potassium channel, subfamily K, member 2 (KCNK2), mRNA	7.58	UP	4.61E-02
50	N28431	yx35c03.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:263716 5 similar to SP:PIR:S32603 S32603 collagen alpha 1(VI) chain - mouse ;, mRNA sequence	7.47	UP	1.75E-02
	NM_012445	Homo sapiens spondin 2, extracellular matrix protein (SPON2), mRNA	7.22	UP	4.03E-02
55	NM_000304	Homo sapiens peripheral myelin protein 22 (PMP22), transcript variant 1, mRNA	7.02	UP	4.58E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_004791	Homo sapiens integrin, beta-like 1 (with EGF-like repeat domains) (ITGBL1), mRNA	7.02	UP	2.64E-02
10	NM_053044	Homo sapiens serine protease HTRA3 (HTRA3), mRNA	6.86	UP	1.13E-03
	BC028245	Homo sapiens, Similar to hypothetical gene LOC130797, clone IMAGE:5395354, mRNA	6.86	UP	1.28E-02
	NM_023003	Homo sapiens transmembrane 6 superfamily member 1 (TM6SF1), mRNA	6.86	UP	2.12E-03
15	AJ420536	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 994183	6.79	UP	3.53E-04
	NM_152996	Homo sapiens sialyltransferase 7 ((alpha-N-acetylneuraminyl-2,3-beta-galactosyl-1,3)-N-acetyl galactosaminide alpha-2,6-sialyltransferase) C (SIAT7C), mRNA	6.79	UP	4.08E-02
20	NM_005583	Homo sapiens lymphoblastic leukemia derived sequence 1 (LYL1), mRNA	6.77	UP	1.92E-02
	NM_016270	Homo sapiens Kruppel-like factor 2 (lung) (KLF2), mRNA	6.55	UP	5.02E-02
25	NM_005397	Homo sapiens podocalyxin-like (PODXL), mRNA	6.53	UP	3.68E-03
	NM_006182	Homo sapiens discoidin domain receptor family, member 2 (DDR2), mRNA	6.52	UP	8.46E-05
	U79271	Human clones 23920 and 23921 mRNA sequence	6.31	UP	8.59E-05
	BC036034	Homo sapiens endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2, transcript variant 2, mRNA (cDNA clone MGC:33157 IMAGE:5272431), complete cds	6.3	UP	3.07E-04
30	NM_152314	Homo sapiens hypothetical protein MGC34830 (MGC34830), mRNA	6.29	UP	2.93E-04
35	AK095791	Homo sapiens cDNA FLJ38472 fis, clone FEBRA2022148	6.27	UP	1.74E-03
	NM_004840	Homo sapiens Rac/Cdc42 guanine nucleotide exchange factor (GEF) 6 (ARHGEF6), mRNA	6.2	UP	1.32E-03
40	AL833655	Homo sapiens mRNA; cDNA DKFZp66700320 (from clone DKFZp667O0320)	6.18	UP	3.01E-02
	NM_032270	Homo sapiens factor for adipocyte differentiation 158 (FAD158), mRNA	6.1	UP	6.22E-05
	NM_021643	Homo sapiens tribbles homolog 2 (Drosophila) (TRIB2), mRNA	5.97	UP	8.59E-05
45	AW044647	wy78e09.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:2554696 3, mRNA sequence	5.94	UP	4.09E-02
	AK125284	Homo sapiens cDNA FLJ43294 fis, clone MESTC1000042	5.85	UP	3.94E-03
50	NM_005069	Homo sapiens single-minded homolog 2 (Drosophila) (SIM2), transcript variant SIM2, mRNA	5.84	UP	8.35E-05
	NM_018836	Homo sapiens transmembrane protein SHREW1 (SHREW1), mRNA	5.83	UP	2.71E-02
55	NM_014932	Homo sapiens neuroligin 1 (NLGN1), mRNA	5.83	UP	2.45E-03
	NM_006307	Homo sapiens sushi-repeat-containing protein, X-linked (SRPX), mRNA	5.77	UP	4.30E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5					
	NM_000210	Homo sapiens integrin, alpha 6 (ITGA6), mRNA	5.77	UP	7.49E-05
	BG208475	RST27977 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	5.72	UP	1.46E-03
10	N36786	yy34e08.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273158 3 similar to contains element MSR1 repetitive element ;, mRNA sequence	5.67	UP	9.13E-05
	NM_005308	Homo sapiens G protein-coupled receptor kinase 5 (GRK5), mRNA	5.58	UP	1.35E-02
15	AB037722	Homo sapiens mRNA for KIAA1301 protein, partial cds	5.58	UP	9.18E-04
	NM_012242	Homo sapiens dickkopf homolog 1 (Xenopus laevis) (DKK1), MRNA,	5.57	UP	1.68E-04
	NM_003068	Homo sapiens snail homolog 2 (Drosophila) (SNAI2), mRNA	5.53	UP	7.42E-05
20	NM_080806	Homo sapiens collagen, type XIII, alpha 1 (COL13A1), transcript variant 10, mRNA	5.52	UP	8.65E-05
	M_173553	Homo sapiens hypothetical protein FLJ25801 (FLJ25801), mRNA	5.48	UP	1.77E-02
25	AI085016	ow88e06.s1 Soares_fetal_liver_spleen_1 NFLS_ S1 Homo sapiens cDNA clone IMAGE:1653922 3, mRNA sequence	5.45	UP	4.77E-03
	NM_182728	Homo sapiens solute carrier family 7 (cationic amino acid transporter, y+ system), member 8 (SLC7A8), transcript variant 2, mRNA	5.34	UP	2.70E-02
30	BC035066	Homo sapiens, clone IMAGE:5259543, mRNA	5.29	UP	2.24E-02
	NM_016247	Homo sapiens interphotoreceptor matrix proteoglycan 2 (IMPG2), mRNA	5.22	UP	2.97E-03
35	NM_001704	Homo sapiens brain-specific angiogenesis inhibitor 3 (BAI3), mRNA	5.19	UP	1.08E-02
	NM_005226	Homo sapiens endothelial differentiation, sphingolipid G-protein-coupled receptor, 3 (EDG3), mRNA	5.18	UP	1.53E-04
	NM_017413	Homo sapiens apelin, AGTRL1 ligand (APLN), mRNA	5.14	UP	2.48E-02
40	NM_000576	Homo sapiens interleukin 1, beta (IL1B), mRNA	5.1	UP	3.32E-03
	NM_012307	Homo sapiens erythrocyte membrane protein band 4.1-like 3 (EPB41L3), mRNA	5.09	UP	4.93E-04
	AL831835	Homo sapiens mRNA; cDNA DKFZp547A0515 (from clone DKFZp547A0515)	5.04	UP	4.14E-02
45	NM_000955	Homo sapiens prostaglandin E receptor 1 (subtype EP1), 42kDa (PTGER1), mRNA	5.02	UP	3.12E-03
	NM_018658	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 16 (KCNJ16), transcript variant 1, mRNA	334.17	Down	8.59E-05
50	NM_007038	Homo sapiens a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2) (ADAMTS5), mRNA	250.9	Down	3.32E-05
	BG219729	RST39494 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	182.25	Down	6.10E-05
55	AB032980	Homo sapiens mRNA for KIAA1154 protein, partial cds	159.9	Down	7.49E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5					
	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	151.72	Down	3.05E-05
10	AK130281	Homo sapiens cDNA FLJ26771 fis, clone PRS03189	148.34	Down	6.10E-05
	NM_003238	Homo sapiens transforming growth factor, beta 2 (TGFB2), mRNA	139.18	Down	7.00E-05
	BU734212	UI-E-CQ1-agd-e-21-0-UI.s1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-agd-e-21-0-UI 3, mRNA sequence	135.35	Down	7.49E-05
15					
	BM993116	UI-H-DTO-aty-f-17-0-UI.s1 NCI_CGAP_DT0 Homo sapiens cDNA clone IMAGE:5866000 3, mRNA sequence	134.03	Down	8.59E-05
20					
	BX648964	Homo sapiens mRNA; cDNA DKFZp686J0156 (from clone DKFZp686J0156)	133.35	Down	1.19E-04
	A1765021	wh56c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384738 3, mRNA sequence	131.86	Down	6.22E-05
	D29453	HUMNK566 Human epidermal keratinocyte Homo sapiens cDNA clone 566, mRNA sequence	128.83	Down	6.30E-05
25					
	NM_002521	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	128.09	Down	1.42E-04
	BX102632	BX102632 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998J052307 ; IMAGE:928228, mRNA sequence	123.34	Down	1.15E-04
30					
	NM_024508	Homo sapiens zinc finger, BED domain containing 2 (ZBED2), mRNA	118.4	Down	5.18E-05
	NM_153026	Homo sapiens prickly-like 1 (Drosophila) (PRICKLE1), mRNA	106.78	Down	6.10E-05
	NM_006228	Homo sapiens prepronociceptin (PNOC), mRNA	102.73	Down	2.87E-04
35					
	NM_005560	Homo sapiens laminin, alpha 5 (LAMA5), mRNA	99.24	Down	7.49E-05
	NM_023942	Homo sapiens hypothetical protein MGC3036 (MGC3036), mRNA	91.38	Down	6.30E-05
	AK096481	Homo sapiens cDNA FLJ39162 fis, clone OCBBF2002376	85.12	Down	7.45E-05
40					
	A1335277	tb29h06.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2055803 3, mRNA sequence	84.58	Down	6.22E-05
	BF798098	RC1-00045-021000-021-f02 CI0045 Homo sapiens cDNA, mRNA sequence	81.7	Down	8.59E-05
45					
	AK023631	Homo sapiens cDNA FLJ13569 fis, clone PLACE 1008369	77.57	Down	7.00E-05
	NM_000927	Homo sapiens ATP-binding cassette, sub-family B (MDR/TAP), member 1 (ABCB1), mRNA	77.16	Down	2.21E-04
	BG197054	RST16291 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	76.09	Down	1.03E-04
50					
	AK058012	Homo sapiens cDNA FLJ25283 fis, clone STM06716	75.4	Down	6.22E-05
	NM_000104	Homo sapiens cytochrome P450, family 1, subfamily B, polypeptide 1 (CYP1B1), mRNA	70.8	Down	6.24E-05
	NM_004617	Homo sapiens transmembrane 4 superfamily member 4 (TM4SF4), mRNA	63.32	Down	1.39E-04
55					
	NM_000990	Homo sapiens ribosomal protein L27a (RPL27A), mRNA	62.1	Down	5.32E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_002899	Homo sapiens retinol binding protein 1, cellular (RBP1), mRNA	62.01	Down	2.87E-04
10	NM_000582	Homo sapiens secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA	60.53	Down	1.18E-04
	NM_002423	Homo sapiens matrix metalloproteinase 7 (matrilysin, uterine) (MMP7), mRNA	60.01	Down	1.56E-04
15	H89053	yw24c06.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253162 5, mRNA sequence	59.02	Down	7.45E-05
	AK090808	Homo sapiens cDNA FLJ33489 fis, clone BRAMY2003585	58.15	Down	7.49E-05
20	NM_001200	Homo sapiens bone morphogenetic protein 2 (BMP2), mRNA	57.05	Down	6.24E-05
	NM_001453	Homo sapiens forkhead box C1 (FOXC1), mRNA	56.24	Down	6.10E-05
	AK092245	Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	56.11	Down	2.55E-04
25	NM_005329	Homo sapiens hyaluronan synthase 3 (HAS3), transcript variant 1, mRNA	55.44	Down	7.22E-05
	AI244954	qj93h05.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1867065 3, mRNA sequence	54.38	Down	6.22E-05
30	NM_003287	Homo sapiens tumor protein D52-like 1 (TPD52L1), transcript variant 1, m RNA	53.91	Down	7.45E-05
	BF696790	602125323F1 NIH_MGC_56 Homo sapiens cDNA clone IMAGE:4282540 5, mRNA sequence	49.46	Down	1.09E-04
	NM_001202	Homo sapiens bone morphogenetic protein 4 (BMP4), transcript variant 1, mRNA	47.81	Down	8.21E-05
35	BC042028	Homo sapiens, clone IMAGE:4794726, mRNA	47.5	Down	7.49E-05
	AL833276	Homo sapiens mRNA; cDNA DKFZp451D088 (from clone DKFZp451D088)	43.28	Down	7.28E-05
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	42.9	Down	7.42E-05
40	NM_003494	Homo sapiens dysferlin, limb girdle muscular dystrophy 2B (autosomal recessive) (DYSF), mRNA	42.2	Down	7.31E-05
	NM_024422	Homo sapiens desmocollin 2 (DSC2), transcript variant Dsc2a, mRNA	41.3	Down	1.96E-04
45	AI249696	qj64a03.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1864204 3, mRNA sequence	41.1	Down	1.50E-04
	AL833166	Homo sapiens mRNA; cDNA DKFZp68612118 (from clone DKFZp68612118)	39.95	Down	1.89E-04
50	NM_173505	Homo sapiens ankyrin repeat domain 29 (ANKRD29), mRNA	39.16	Down	9.13E-05
	NM_130435	Homo sapiens protein tyrosine phosphatase, receptor type, E (PTPRE), transcript variant 2, mRNA	39.16	Down	2.54E-04
	NM_016356	Homo sapiens doublecortin domain containing 2 (DCDC2), mRNA	38.33	Down	2.10E-04
55	AF318382	Homo sapiens pp9974 mRNA, complete cds	38.03	Down	8.59E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_018265	Homo sapiens hypothetical protein FLJ10901 (FLJ10901), mRNA	37.23	Down	1.50E-04
10	NM_152487	Homo sapiens hypothetical protein FLJ31842 (FLJ31842), mRNA	37	Down	1.56E-04
	AK000075	Homo sapiens cDNA FLJ20068 fis, clone COL01755	35.55	Down	2.21 E-04
	NM_012464	Homo sapiens tolloid-like 1 (TLL1), mRNA	35.05	Down	2.09E-03
15	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	34.38	Down	7.49E-05
	BU680661	UI-CF-DU1-aaz-f-04-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaz-f-04-0-UI 3, mRNA sequence	34.34	Down	6.22E-05
20	BF431030	7o18c06.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3574283 3, mRNA sequence	34.25	Down	1.86E-04
	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	34.23	Down	6.30E-05
	NM_152284	Homo sapiens Snf7 homologue associated with Alix 3 (Shax3), mRNA	33.99	Down	7.71E-05
25	BC037316	Homo sapiens, clone IMAGE:5259432, mRNA	33.46	Down	2.03E-04
	NM_002345	Homo sapiens lumican (LUM), mRNA	33.34	Down	8.46E-05
	NM_080743	Homo sapiens serine-arginine repressor protein (35 kDa) (SRrp35), mRNA	32.96	Down	5.23E-05
30	NM_004221	Homo sapiens natural killer cell transcript 4 (NK4), mRNA	32.74	Down	1.45E-04
	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	32.31	Down	2.57E-04
	AA738254	nx13b02.s1 NCI_CGAP_GC3 Homo sapiens cDNA clone IMAGE:1255947 3, mRNA sequence	32.23	Down	6.10E-05
35	BC045828	Homo sapiens zinc finger protein 608, mRNA (cDNA clone IMAGE:5262896), partial cds	32.18	Down	7.49E-05
	NM_000599	Homo sapiens insulin-like growth factor binding protein 5 (IGFBP5), mRNA	32.18	Down	9.81E-05
	AF055376	Homo sapiens short form transcription factor C-MAF (c-maf) mRNA, complete cds	31.81	Down	2.01E-04
40	NM_015236	Homo sapiens latrophilin 3 (LPHN3), mRNA	31.68	Down	9.69E-05
	NM_007069	Homo sapiens HRAS-like suppressor 3 (HRASLS3), mRNA	31.67	Down	7.49E-05
	BG221364	RST41175Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	31.63	Down	2.22E-04
45	AI819186	wj32d10.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2404531 3, mRNA sequence	31.49	Down	1.80E-04
	BF512544	UI-H-BW1-amf-c-08-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069687 3, mRNA sequence	30.8	Down	7.49E-05
50	NM_020873	Homo sapiens leucine rich repeat neuronal 1 (LRRN1), mRNA	30.67	Down	2.01E-04
	CA425961	UI-H-FE1-beg-p-18-0-UI.s1 NCI_CGAP_FE1 Homo sapiens cDNA clone UI-H-FE1-beg-p-18-0-UI 3, mRNA sequence	30.15	Down	1.88E-04
55	AK000776	Homo sapiens cDNA FLJ20769 fis, clone COL06674	30.11	Down	1.15 E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	M_001562	Homo sapiens interleukin 18 (interferon-gamma-inducing factor) (IL18), mRNA	29.97	Down	7.65E-04
10	NM_014333	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	29.77	Down	5.82E-04
	NM_030583	Homo sapiens matrilin 2 (MATN2), transcript variant 2, mRNA	29.05	Down	8.46E-05
15	W93585	zd95g01.s1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:357264 3, mRNA sequence	28.93	Down	6.22E-05
	NM_012198	Homo sapiens grancalcin, EF-hand calcium binding protein (GCA), mRNA	28.84	Down	9.13E-05
	NM_001448	Homo sapiens glypican 4 (GPC4), mRNA	28.72	Down	6.36E-04
20	AK056882	Homo sapiens cDNA FLJ32320 fis, clone PROST2003537	28.6	Down	4.45E-04
	AB033048	Homo sapiens mRNA for KIAA1222 protein, partial cds	28.27	Down	2.41E-04
	NM_025074	Homo sapiens Fraser syndrome 1 (FRAS1), transcript variant 1, mRNA	28.13	Down	8.92E-05
25	NM_019000	Homo sapiens hypothetical protein FLJ20152 (FLJ20152), mRNA	27.54	Down	1.07E-04
	BU729783	UI-E-CK1-afh-h-18-0-UI.s1 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-afh-h-18-0-UI 3, mRNA sequence	26.13	Down	6.30E-05
	NM_004862	Homo sapiens lipopolysaccharide-induced TNF factor (LITAF), mRNA	25.95	Down	8.99E-05
30	AB011539	Homo sapiens mRNA for MEGF6 protein (KIAA0815), partial cds	25.67	Down	3.12E-04
	BC043195	Homo sapiens cDNA clone IMAGE:5288757, partial cds	25.58	Down	2.22E-04
35	NM_152864	Homo sapiens chromosome 20 open reading frame 58 (C20orf58), mRNA	25.35	Down	3.00E-04
	AK125608	Homo sapiens cDNA FLJ43620 fis, clone SPLEN2021701, highly similar to HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN PRECURSOR	25.33	Down	2.03E-04
40	AL080103	Homo sapiens mRNA; cDNA DKFZp564N2216 (from clone DKFZp564N2216)	25.17	Down	2.03E-04
	NM_173549	Homo sapiens hypothetical protein FLJ39553 (FLJ39553), mRNA	24.71	Down	1.67E-04
45	BG211832	RST31404 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	24.57	Down	1.02E-04
	NM_024726	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	24.08	Down	7.49E-05
	NM_002559	Homo sapiens purinergic receptor P2X, ligand-gated ion channel, 3 (P2RX3), mRNA	24.04	Down	2.63E-04
50	NM_018168	Homo sapiens chromosome 14 open reading frame 105 (C14orf105), mRNA	23.69	Down	1.50E-04
	NM_000599	Homo sapiens insulin-like growth factor binding protein 5 (IGFBP5), mRNA	23.5	Down	4.02E-04
55	NM_003551	Homo sapiens non-metastatic cells 5, protein expressed in (nucleoside-diphosphate kinase) (NME5), mRNA	23.23	Down	1.41E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_032471	Homo sapiens protein kinase (cAMP-dependent, catalytic) inhibitor beta (PKIB), transcript variant 3, mRNA	22.83	Down	1.54E-04
10	NM_005949	Homo sapiens metallothionein 1 F (functional) (MT1 F), mRNA	22.22	Down	8.81E-05
	NM_018242	Homo sapiens hypothetical protein FLJ10847 (FLJ10847), mRNA	22.1	Down	2.03E-04
15	AW151660	xf67d04.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2623111 3, mRNA sequence	21.81	Down	2.71E-04
	AK096975	Homo sapiens cDNA FLJ39656 fis, clone SMINT2005956	21.78	Down	2.52E-04
20	CA306881	UI-H-FT1-bht-n-22-0-UI.s1 NCI_CGAP_FT1 Homo sapiens cDNA clone UI-H-FT1-bht-n-22-0-UI 3, mRNA sequence	21.69	Down	1.30E-04
	NM_016212	Homo sapiens TP53TG3 protein (TP53TG3), mRNA	21.67	Down	9.03E-05
	NM_178470	Homo sapiens WD repeat domain 40B (WDR40B), mRNA	21.63	Down	7.59E-05
25	NM_014243	Homo sapiens a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 3 (ADAMTS3), mRNA	21.47	Down	8.24E-05
	AK126467	Homo sapiens cDNA FLJ44503 fis, clone UTERU3001158	21.42	Down	6.22E-05
30	AW248516	2820632.3prime NIH_MGC_7 Homo sapiens cDNA clone IMAGE:2820632 3, mRNA sequence	21.37	Down	1.23E-04
	NM_001263	Homo sapiens CDP-diacylglycerol synthase (phosphatidate cytidyltransferase) 1 (CDS1), mRNA	21.12	Down	1.10E-04
35	BM669002	UI-E-CK1-afn-m-04-0-UI.s2 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-afn-m-04-0-UI 3, mRNA sequence	20.96	Down	7.28E-05
	NM_000170	Homo sapiens glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P) (GLDC), mRNA	20.78	Down	1.19E-04
40	NM_014585	Homo sapiens solute carrier family 40 (iron-regulated transporter), member 1 (SLC40A1), mRNA	20.73	Down	8.46E-05
	NM_031426	Homo sapiens chromosome 9 open reading frame 58 (C9orf58), transcript variant 1, mRNA	20.62	Down	2.00E-04
45	BM728728	UI-E-E01-aiv-c-02-0-UI.r1 UI-E-E01 Homo sapiens cDNA clone UI-E-E01-aiv-c-02-0-UI 5, mRNA sequence	20.59	Down	1.14E-04
	NM_152369	Homo sapiens hypothetical protein MGC45474 (MGC45474), mRNA	20.57	Down	9.78E-05
50	NM_175056	Homo sapiens hypothetical protein LOC131368 (LOC131368), mRNA	20.51	Down	5.42E-04
	NM_004524	Homo sapiens lethal giant larvae homolog 2 (Drosophila) (LLGL2), mRNA	20.3	Down	7.45E-05
	AI436290	th81c01.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2125056 3, mRNA sequence	19.93	Down	1.23E-04
55	AW268540	xv51e10.x1 NCI_CGAP_Lu28 Homo sapiens cDNA clone IMAGE:2816682 3, mRNA sequence	19.89	Down	1.09E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5					
	BC015159	Homo sapiens cDNA clone IMAGE:3885734, partial cds	19.47	Down	7.49E-05
	NM_020130	Homo sapiens chromosome 8 open reading frame 4 (C8orf4), mRNA	19.43	Down	2.83E-04
10					
	BM979825	UI-CF-DU1-adt-f-12-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-adt-f-12-0-UI 3, mRNA sequence	19.41	Down	1.01E-04
	BM712072	UI-E-DW1-ahc-b-11-0-UI.r1 UI-E-DW1 Homo sapiens cDNA clone UI-E-DW1-ahc-b-11-0-UI 5, mRNA sequence	19.33	Down	1.19E-04
15					
	CB135276	K-EST0187371 L5HLK1 Homo sapiens cDNA clone L5HLK1-32-B12 5, mRNA sequence	18.98	Down	8.97E-05
	T53523	ya89h12.r1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:68903 5, mRNA sequence	18.87	Down	1.96E-04
20					
	NM_138432	Homo sapiens serine dehydratase-like (SDSL), mRNA	18.84	Down	1.32E-04
	NM_023915	Homo sapiens G protein-coupled receptor 87 (GPR87), mRNA	18.74	Down	2.72E-04
	NM_017549	Homo sapiens upregulated in colorectal cancer gene 1 (UCC1), mRNA	18.73	Down	8.83E-05
25					
	BG436244	602508665F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4605617 5, mRNA sequence	18.7	Down	1.49E-04
	NM_024901	Homo sapiens hypothetical protein FLJ22457 (FLJ22457), mRNA	18.68	Down	1.35E-04
30					
	NM_005712	Homo sapiens HERV-H LTR-associating 1 (HHLA1), mRNA	18.63	Down	1.08E-04
	M_198488	Homo sapiens FLJ46072 protein (FLJ46072), mRNA	18.58	Down	7.49E-05
	NM_020349	Homo sapiens ankyrin repeat domain 2 (stretch responsive muscle) (ANKRD2), mRNA	18.49	Down	2.00E-04
35					
	NM_002148	Homo sapiens homeo box D10 (HOXD10), mRNA	18.41	Down	3.05E-05
	N78460	yz76h06.r1 Soares_multiple_sclerosis_2NbHMS P Homo sapiens cDNA clone IMAGE:289019 5, mRNA sequence	18.4	Down	4.02E-04
40					
	NM_006598	Homo sapiens solute carrier family 12 (potassium/chloride transporters), member 7 (SLC12A7), mRNA	18.13	Down	8.59E-05
	BF431041	nab31g02.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3267627 3, mRNA sequence	18.11	Down	6.10E-05
45					
	AK126467	Homo sapiens cDNA FLJ44503 fis, clone UTERU3001158	17.84	Down	1.31E-04
	NM_182487	Homo sapiens olfactomedin-like 2A (OLFML2A), mRNA	17.77	Down	1.06E-04
50					
	AK058012	Homo sapiens cDNA FLJ25283 fis, clone STM06716	17.76	Down	8.82E-05
	NM_001252	Homo sapiens tumor necrosis factor (ligand) superfamily, member 7 (TNFSF7), mRNA	17.71	Down	2.46E-04
	NM_006252	Homo sapiens protein kinase, AMP-activated, alpha 2 catalytic subunit (PRKAA2), mRNA	17.63	Down	6.57E-04
55					
	NM_030899	Homo sapiens zinc finger protein 323 (ZNF323), mRNA	17.62	Down	1.77E-04
	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	17.53	Down	1.61E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AK056431	Homo sapiens cDNA FLJ31869 fis, clone NT2RP7002151	17.29	Down	8.46E-05
10	BM719937	UI-E-EJ0-ahu-a-10-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahu-a-10-0-UI5, mRNA sequence	17.11	Down	4.37E-05
	NM_005855	Homo sapiens receptor (calcitonin) activity modifying protein 1 (RAMP1), mRNA	16.97	Down	1.56E-04
	AK124873	Homo sapiens cDNA FLJ42883 fis, clone BRHIP3006683	16.87	Down	3.99E-04
15	NM_001977	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	16.61	Down	6.12E-04
	AK074181	Homo sapiens mRNA for FLJ00254 protein	16.54	Down	6.10E-05
20	NM_000557	Homo sapiens growth differentiation factor 5 (cartilage-derived morphogenetic protein-1) (GDF5), mRNA	16.47	Down	1.49E-04
	AK026966	Homo sapiens cDNA: FLJ23313 fis, clone HEP11919	16.36	Down	8.92E-05
	BX113319	BX113319 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGp998G205398 ; IMAGE:2184619, mRNA sequence	16.19	Down	6.30E-05
25	AK055362	Homo sapiens cDNA FLJ30800 fis, clone FEBRA2001197	16.02	Down	3.05E-05
	NM_021102	Homo sapiens serine protease inhibitor, Kunitz type, 2 (SPINT2), mRNA	15.94	Down	2.71E-04
30	NM_002354	Homo sapiens tumor-associated calcium signal transducer 1 (TACSTD1), mRNA	15.93	Down	9.13E-05
	BQ003401	UI-H-EI1-azd-j-23-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847286 3, mRNA sequence	15.78	Down	6.30E-05
35	NM_033641	Homo sapiens collagen, type IV, alpha 6 (COL4A6), transcript variant B, mRNA	15.77	Down	5.35E-05
	N38753	yy42d01.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273889 3, mRNA sequence	15.77	Down	3.67E-05
40	AI420213	te92g09.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2094208 3, mRNA sequence	15.7	Down	2.35E-04
	AK095776	Homo sapiens cDNA FLJ38457 fis, clone FEBRA2020400	15.48	Down	3.04E-04
	NM_006378	Homo sapiens sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D (SEMA4D), mRNA	15.33	Down	6.10E-05
45	AK021801	Homo sapiens cDNA FLJ11739 fis, clone HEMBA1005497	15.33	Down	1.38E-04
	BX538226	Homo sapiens mRNA; cDNA DKFZp686E1944 (from clone DKFZp686E1944)	15.23	Down	2.37E-04
50	NM_001847	Homo sapiens collagen, type IV, alpha 6 (COL4A6), transcript variant A, mRNA	15.22	Down	7.42E-05
	NM_017640	Homo sapiens leucine rich repeat containing 16 (LRRC16), mRNA	15.21	Down	6.39E-05
55	NM_007072	Homo sapiens HERV-H LTR-associating 2 (HHLA2), mRNA	15.04	Down	1.06E-04
	NM_052947	Homo sapiens heart alpha-kinase (HAK), mRNA	14.96	Down	2.23E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5					
	NM_005139	Homo sapiens annexin A3 (ANXA3), mRNA	14.95	Down	6.71E-05
	N63415	yy60d04.s1 Soares_multiple_sclerosis_2NbHMS P	14.79	Down	2.23E-05
10		Homo sapiens cDNA clone IMAGE:277927 3 similar to contains L1.b3 L1 repetitive element;, mRNA sequence			
	BC052289	Homo sapiens carboxypeptidase A4, mRNA (cDNA clone MGC:59749 IMAGE:6106874), complete cds	14.76	Down	7.22E-05
	NM_153715	Homo sapiens homeo box A10 (HOXA10), transcript variant 2, mRNA	14.76	Down	8.35E-05
15					
	NM_014936	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 4 (putative function) (ENPP4), mRNA	14.67	Down	8.19E-04
	NM_021192	Homo sapiens homeo box D11 (HOXD11), mRNA	14.47	Down	4.53E-04
20					
	BX118238	BX118238 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998L153800; IMAGE:1501598, mRNA sequence	14.44	Down	5.07E-04
	NM_024969	Homo sapiens TGF-beta induced apoptosis protein 2 (TAIP-2), mRNA	14.11	Down	1.57E-04
25					
	AK125490	Homo sapiens cDNA FLJ43501 fis, clone PEBLM2004497	14.06	Down	9.71E-04
	R99527	yq79b11.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:2019813, mRNA sequence	13.98	Down	8.39E-04
	NM_001873	Homo sapiens carboxypeptidase E (CPE), mRNA	13.91	Down	9.16E-05
30					
	NM_003761	Homo sapiens vesicle-associated membrane protein 8 (endobrevin) (VAMP8), mRNA	13.87	Down	3.12E-04
	NM_020808	Homo sapiens signal-induced proliferation-associated 1 like 2 (SIPA1L2), mRNA	13.84	Down	5.47E-04
	NM_152573	Homo sapiens RAS and EF hand domain containing (RASEF), mRNA	13.82	Down	5.13E-04
35					
	NM_032866	Homo sapiens cingulin-like 1 (CGNL1), mRNA	13.78	Down	6.30E-05
	BU740051	UI-E-EO0-ahw-n-18-0-UI.s1 UI-E-EO0 Homo sapiens cDNA clone UI-E-EO0-ahw-n-18-0-UI 3, mRNA sequence	13.74	Down	4.63E-04
40					
	NM_198389	Homo sapiens lung type-I cell membrane-associated glycoprotein (T1A-2), transcript variant 2, mRNA	13.71	Down	3.62E-04
	CA777268	ip05d09.y1 HR85 islet Homo sapiens cDNA clone IMAGE:6134849 5, mRNA sequence	13.63	Down	1.39E-04
45					
	H00617	yj25f02.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:149787 3, mRNA sequence	13.45	Down	3.33E-04
	NM_024898	Homo sapiens family with sequence similarity 31, member C (FAM31C), mRNA	13.39	Down	2.17E-03
	NM_000094	Homo sapiens collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive) (COL7A1 mRNA	13.3	Down	4.93E-04
50					
	NM_002837	Homo sapiens protein tyrosine phosphatase, receptor type, B (PTPRB), mRNA	13.25	Down	6.84E-04
	NM_018728	Homo sapiens myosin VC (MYO5C), mRNA	13.15	Down	1.64E-04
55					
	NM_018659	Homo sapiens cytokine-like protein C17 (C17), mRNA	13.11	Down	1.45E-04
	AL834140	Homo sapiens mRNA; cDNA DKFZp434A2029 (from clone DKFZp434A2029)	13.08	Down	9.99E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5					
	BX103476	BX103476 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998C053946 ; IMAGE:1557436, mRNA sequence	13.02	Down	1.71E-04
10					
	BG545305	602572521 F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4700644 5, mRNA sequence	12.86	Down	6.30E-05
	BX647876	Homo sapiens mRNA; cDNA DKFZp313A1525 (from clone DKFZp313A1525)	12.83	Down	7.95E-05
15					
	NM_030949	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 14C (PPP1R14C), mRNA	12.82	Down	3.35E-04
	NM_024423	Homo sapiens desmocollin 3 (DSC3), transcript variant Dsc3b, mRNA	12.81	Down	7.71E-05
20					
	NM_014452	Homo sapiens tumor necrosis factor receptor superfamily, member 21 (TNFRSF21), mRNA	12.8	Down	2.20E-04
	NM_002246	Homo sapiens potassium channel, subfamily K, member 3 (KCNK3), mRNA	12.76	Down	7.59E-05
	NM_002427	Homo sapiens matrix metalloproteinase 13 (collagenase 3) (MMP13), mRNA	12.65	Down	4.08E-04
25					
	NM_003328	Homo sapiens TXK tyrosine kinase (TXK), mRNA	12.61	Down	2.45E-04
	NM_014422	Homo sapiens phosphatidylinositol (4,5) bisphosphate 5-phosphatase, A (PIB5PA), transcript variant 1, mRNA	12.48	Down	8.97E-05
	NM_014333	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	12.4	Down	1.04E-04
30					
	AA888443	nw74f10.s1 NCI_CGAP_Pr12 Homo sapiens cDNA clone IMAGE:1252363, mRNA sequence	12.37	Down	3.96E-04
	BX110418	BX110418 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998C224149 ; IMAGE:1635405, mRNA sequence	12.35	Down	1.54E-04
35					
	NM_019102	Homo sapiens homeo box A5 (HOXA5), mRNA	12.33	Down	7.59E-05
	NM_016463	Homo sapiens CXXC finger 5 (CXXC5), mRNA	12.32	Down	1.96E-04
	NM_004572	Homo sapiens plakophilin 2 (PKP2), transcript variant 2b, mRNA	12.28	Down	4.15E-04
40					
	N25875	yw78d12.s1 Soares_placenta_8to9weeks_2NbH P8to9W Homo sapiens cDNA clone IMAGE:258359 3, mRNA sequence	12.27	Down	1.02E-04
	NM_152694	Homo sapiens zinc finger, CCHC domain containing 5 (ZCCHC5), mRNA	12.22	Down	3.38E-04
45					
	NM_021012	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 12 (KCNJ12), mRNA	12.22	Down	9.23E-04
	NM_017671	Homo sapiens chromosome 20 open reading frame 42 (C20orf42), mRNA	12.21	Down	1.06E-04
50					
	NM_002031	Homo sapiens fyn-related kinase (FRK), mRNA	12.11	Down	2.14E-04
	BM976385	UI-CF-EN1-acz-f-03-0-UI.s1 UI-CF-EN1 Homo sapiens cDNA clone UI-CF-EN1-acz-f-03-0-UI 3, mRNA sequence	12.02	Down	8.11E-04
55					
	NM_005296	Homo sapiens G protein-coupled receptor 23 (GPR23), mRNA	11.98	Down	1.50E-04
	NM_000817	Homo sapiens glutamate decarboxylase 1 (brain, 67kDa) (GAD1 transcript variant GAD67, mRNA	11.92	Down	6.22E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AI220066	qg84d01.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1841857 3, mRNA sequence	11.88	Down	1.09E-04
10	NM_005737	Homo sapiens ADP-ribosylation factor-like 7 (ARL7), mRNA	11.83	Down	1.58E-04
	NM_000147	Homo sapiens fucosidase, alpha-L-1, tissue (FUCA1), mRNA	11.82	Down	1.74E-03
	AK127437	Homo sapiens cDNA FLJ45529 fis, clone BRTHA2027546	11.82	Down	1.68E-04
15	NM_178033	Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1 (CYP4X1), mRNA	11.81	Down	2.10E-04
	NM_173567	Homo sapiens abhydrolase domain containing 7 (ABHD7), mRNA	11.76	Down	1.75E-04
20	AL049974	Homo sapiens mRNA; cDNA DKFZp564B222 (from clone DKFZp564B222)	11.74	Down	5.82E-04
	NM_000519	Homo sapiens hemoglobin, delta (HBD), mRNA	11.7	Down	9.70E-04
	AL359567	Homo sapiens mRNA; cDNA DKFZp547D023 (from clone DKFZp547D023)	11.66	Down	3.95E-04
25	BF510493	UI-H-BI4-apa-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086558 3, mRNA sequence	11.66	Down	6.30E-05
	NM_000961	Homo sapiens prostaglandin 12 (prostacyclin) synthase (PTGIS), mRNA	11.63	Down	5.06E-04
30	NM_025151	Homo sapiens RAB11 family interacting protein 1 (class I) (RAB11FIP1 transcript variant 1, mRNA	11.6	Down	1.14E-04
	BM712945	UI-E-EJ0-ahi-c-16-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahi-c-16-0-UI 5, mRNA sequence	11.6	Down	9.48E-05
35	AW451831	UI-H-B13-alk-e-12-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2737246 3, mRNA sequence	11.54	Down	8.69E-04
	BC040701	Homo sapiens cDNA clone IMAGE:5736259, partial cds	11.53	Down	1.23E-04
40	AL359058	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 592473	11.44	Down	1.27E-03
	NM_001854	Homo sapiens collagen, type XI, alpha 1 (COL11A1), transcript variant A, mRNA	11.39	Down	7.56E-04
	NM_000227	Homo sapiens laminin, alpha 3 (LAMA3), transcript variant 2, mRNA	11.39	Down	8.59E-05
45	NM_033256	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 14A (PPP1R14A), mRNA	11.36	Down	1.26E-04
	AB011538	Homo sapiens mRNA for MEGF5, partial cds	11.35	Down	4.81 E-04
	AL389942	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 2005635	11.33	Down	4.78E-04
50	NM_174900	Homo sapiens zinc finger protein 42 (ZFP42), mRNA	11.23	Down	1.29E-04
	BG622707	602647476F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4768963 5, mRNA sequence	11.21	Down	1.49E-04
	NM_178177	Homo sapiens nicotinamide nucleotide adenyltransferase 3 (NMNAT3), mRNA	11.19	Down	3.57E-04
55	AA099748	zl78c09.s1 Stratagene colon (#937204) Homo sapiens cDNA clone IMAGE:510736 3, mRNA sequence	11.16	Down	9.87E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	H89526	yw28b04.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253519 5, mRNA sequence	11.06	Down	2.94E-04
10	NM_178868	Homo sapiens chemokine-like factor super family 8 (CKLFSF8), mRNA	10.96	Down	9.23E-04
	BE788763	601475864F1 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:3879014 5, mRNA sequence	10.88	Down	6.22E-05
15	NM_000076	Homo sapiens cyclin-dependent kinase inhibitor 1C (p57, Kip2) (CDKN1C), mRNA	10.87	Down	7.28E-05
	NM_152768	Homo sapiens hypothetical protein FLJ25378 (FLJ25378), mRNA	10.81	Down	7.31E-05
	M60502	Human profilaggrin mRNA, 3 end	10.81	Down	8.46E-05
20	NM_181847	Homo sapiens amphoterin induced gene 2 (AMIGO2), mRNA	10.78	Down	1.86E-03
	NM_005331	Homo sapiens hemoglobin, theta 1 (HBQ1), mRNA	10.77	Down	3.32E-04
	NM_032367	Homo sapiens zinc finger, BED domain containing 3 (ZBED3), mRNA	10.61	Down	7.71E-05
25	NM_004574	Homo sapiens peanut-like 2 (Drosophila) (PNUTL2), transcript variant 1, mRNA	10.6	Down	3.18E-04
	NM_014421	Homo sapiens dickkopf homolog 2 (Xenopus laevis) (DKK2), mRNA	10.52	Down	7.31E-05
	NM_052923	Homo sapiens zinc finger protein 452 (ZNF452), mRNA	10.43	Down	1.06E-04
30	NM_006379	Homo sapiens sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3C (SEMA3C), mRNA	10.39	Down	6.30E-05
	AL137488	Homo sapiens mRNA; cDNA DKFZp434N2030 (from clone DKFZp434N2030)	10.37	Down	2.60E-04
35	BX088936	BX088936 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998G123255 ; IMAGE:1292195, mRNA sequence	10.34	Down	1.20E-03
	AB041269	Homo sapiens mRNA for keratin 19, partial cds, isolate:K19-141	10.33	Down	4.02E-03
40	AK091933	Homo sapiens cDNA FLJ34614 fis, clone KIDNE2014268	10.3	Down	1.49E-04
	AF269162	Homo sapiens c21orf7 form B mRNA, complete cds	10.3	Down	1.78E-03
	NM_001935	Homo sapiens dipeptidylpeptidase 4 (CD26, adenosine deaminase complexing protein 2) (DPP4), mRNA	10.27	Down	1.10E-04
45	NM_001850	Homo sapiens collagen, type VIII, alpha 1 (COL8A1 transcript variant 1, m RNA	10.23	Down	2.23E-05
	AI469032	ti70a01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2137320 3, mRNA sequence	10.2	Down	7.42E-05
50	NM_003081	Homo sapiens synaptosomal-associated protein, 25kDa (SNAP25), transcript variant 1, mRNA	10.05	Down	2.34E-03
	BG570144	602591134F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4717761 5, mRNA sequence	9.99	Down	1.09E-03
55	BM969191	UI-CF-ENO-acp-e-22-0-UI.s1 UI-CF-ENO Homo sapiens cDNA clone UI-CF-ENO-acp-e-22-0-UI 3, mRNA sequence	9.99	Down	1.00E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_006622	Homo sapiens polo-like kinase 2 (Drosophila) (PLK2), mRNA	9.99	Down	1.45E-04
10	NM_000860	Homo sapiens hydroxyprostaglandin dehydrogenase 15-(NAD) (HPGD), mRNA	9.98	Down	6.39E-05
	AJ318805	AJ318805 Homo sapiens adipose tissue Homo sapiens cDNA clone 2040, mRNA sequence	9.97	Down	1.92E-04
15	NM_002735	Homo sapiens protein kinase, cAMP-dependent, regulatory, type I, beta (PRKAR1 B), mRNA	9.93	Down	8.51E-06
	AK092114	Homo sapiens cDNA FLJ34795 fis, clone NT2NE2005921	9.91	Down	1.57E-03
	NM_024608	Homo sapiens nei endonuclease VIII-like 1 (E. coli) (NEIL1), mRNA	9.89	Down	3.05E-05
20	BE464407	hx89g05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3195032 3, mRNA sequence	9.88	Down	1.05E-03
	BF431460	7o14b05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3573849 3, mRNA sequence	9.85	Down	1.19E-03
25	BF509573	UI-H-BI4-apf-b-11-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086949 3, mRNA sequence	9.82	Down	2.57E-04
	T78754	yd01f08.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:24180 5, mRNA sequence	9.79	Down	1.51E-03
30	M_153256	Homo sapiens chromosome 10 open reading frame 47 (C10orf47), mRNA	9.78	Down	2.10E-04
	NM_207482	Homo sapiens FLJ44048 protein (FLJ44048), mRNA	9.76	Down	2.14E-04
	NM_002276	Homo sapiens keratin 19 (KRT19), mRNA	9.74	Down	3.70E-03
	N70752	za72d02.s1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:298083 3, mRNA sequence	9.73	Down	1.49E-04
35	NM_020962	Homo sapiens likely ortholog of mouse neighbor of Punc E11 (NOPE), mRNA	9.73	Down	8.91E-04
	NM_016276	Homo sapiens serum/glucocorticoid regulated kinase 2 (SGK2), transcript variant 2, mRNA	9.7	Down	4.41E-04
40	AI355761	qt94a11.x1 NCI_CGAP_Co14 Homo sapiens cDNA clone IMAGE:1962908 3 similar to gb:X74929 KERATIN, TYPE II CYTOSKELETAL 8 (HUMAN);, mRNA sequence	9.7	Down	9.99E-05
45	NM_003264	Homo sapiens toll-like receptor 2 (TLR2), mRNA	9.69	Down	1.86E-04
	NM_198495	Homo sapiens CTAGE family, member 4 (CTAGE4), mRNA	9.66	Down	3.12E-04
	BX097888	BX097888 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGEp998K064187 ; IMAGE:1650173, mRNA sequence	9.62	Down	6.10E-05
50	BU567804	AGENCOURT_10398872 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6614502 5, mRNA sequence	9.62	Down	7.71E-05
	NM_152423	Homo sapiens melanoma associated antigen (mutated) 1-like 1 (MUM1L1), mRNA	9.61	Down	2.45E-03
55	NM_018330	Homo sapiens KIAA1598 (KIAA1598), mRNA	9.6	Down	1.68E-04
	BU584197	2513030T6 LIVRTUT04 Homo sapiens cDNA clone 2513030 3, mRNA sequence	9.56	Down	8.83E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_005264	Homo sapiens GDNF family receptor alpha 1 (GFRA1), transcript variant 1, m RNA	9.56	Down	1.19E-04
10	NM_002245	Homo sapiens potassium channel, subfamily K, member 1 (KCNK1), mRNA	9.53	Down	2.05E-03
	M_178550	Homo sapiens hypothetical protein MGC48998 (MGC48998), mRNA	9.51	Down	9.03E-04
15	NM_006863	Homo sapiens leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 1 (LILRA1), mRNA	9.48	Down	6.22E-05
	M_173354	Homo sapiens SNF1-like kinase (SNF1LK), mRNA	9.47	Down	8.25E-04
	BC016962	Homo sapiens, clone IMAGE:4182947, mRNA	9.43	Down	2.15E-04
20	AB046810	Homo sapiens mRNA for KIAA1590 protein, partial cds	9.38	Down	6.30E-05
	BX116347	BX116347 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGp998B215967 ; IMAGE:2401844, mRNA sequence	9.33	Down	1.54E-04
25	BU633163	UI-H-FL1-bgt-n-07-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgt-n-07-0-UI 3, mRNA sequence	9.31	Down	1.46E-04
	AI289329	qw28c09.x1 NCI_CGAP_Ut4 Homo sapiens cDNA clone IMAGE:1992400 3 similar to contains L1.b2 L1 repetitive element ;, mRNA sequence	9.28	Down	4.80E-04
30	BU951469	in60a05.x3 HR85 islet Homo sapiens cDNA clone IMAGE:6126249 3, mRNA sequence	9.21	Down	2.54E-03
	NM_002338	Homo sapiens limbic system-associated membrane protein (LSAMP), mRNA	9.19	Down	1.30E-03
35	AL833346	Homo sapiens mRNA; cDNA DKFZp686M2234 (from clone DKFZp686M2234)	9.17	Down	3.02E-04
	BC040293	Homo sapiens, clone IMAGE:4820330, mRNA	9.1	Down	1.09E-03
	NM_003810	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	9.04	Down	5.06E-04
40	NM_014373	Homo sapiens G protein-coupled receptor 160 (GPR160), mRNA	9.01	Down	1.59E-04
	CA425405	UI-H-FE1-bef-g-08-0-UI.s1 NCI_CGAP_FE1 Homo sapiens cDNA clone UI-H-FE1-bef-g-08-0-UI 3, mRNA sequence	8.89	Down	1.74E-05
45	AI953708	wq47d09.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474417 3, mRNA sequence	8.87	Down	9.89E-03
	NM_017641	Homo sapiens kinesin family member 21A (KIF21A), mRNA	8.86	Down	7.22E-05
	NM_153377	Homo sapiens leucine-rich repeats and immunoglobulin-like domains 3 (LRIG3), mRNA	8.86	Down	2.71E-04
50	NM_000236	Homo sapiens lipase, hepatic (LIPC), mRNA	8.83	Down	1.93E-03
	BC046362	Homo sapiens voltage-dependent calcium channel gamma subunit-like protein, mRNA (cDNA clone MGC:50757 IMAGE:5221396), complete cds	8.8	Down	4.71E-04
55	AJ697972	Homo sapiens chromosome 3 cDNA	8.8	Down	7.35E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	BF509074	UI-H-BI4-aou-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086150 3, mRNA sequence	8.8	Down	1.24E-04
10	NM_001080	Homo sapiens aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) (ALDH5A1), nuclear gene encoding mitochondrial protein, transcript variant 2, mRNA	8.79	Down	8.46E-05
15	NM_002247	Homo sapiens potassium large conductance calcium-activated channel, subfamily M, alpha member 1 (KCNMA1 mRNA	8.78	Down	7.45E-05
	NM_022154	Homo sapiens solute carrier family 39 (zinc transporter), member 8 (SLC39A8), mRNA	8.77	Down	4.65E-05
20	NM_144707	Homo sapiens prominin 2 (PROM2), mRNA	8.73	Down	2.01E-04
	AB095949	Homo sapiens mRNA for KIAA2029 protein	8.69	Down	9.23E-04
	AV728294	AV728294 HTC Homo sapiens cDNA clone HTCBI09 5, mRNA sequence	8.68	Down	8.46E-05
25	NM_003851	Homo sapiens cellular repressor of E1A-stimulated genes 1 (CREG1), mRNA	8.68	Down	2.10E-04
	NM_014181	Homo sapiens HSPC159 protein (HSPC159), mRNA	8.67	Down	1.10E-04
	NM_000349	Homo sapiens steroidogenic acute regulator (STAR), nuclear gene encoding mitochondrial protein, transcript variant 1, mRNA	8.66	Down	1.03E-04
30	AK026740	Homo sapiens cDNA: FLJ23087 fis, clone LNG06994, highly similar to AF161368 Homo sapiens HSPC105 mRNA	8.65	Down	7.45E-04
	T56535	yb33g07.r1 Stratagene fetal spleen (#937205) Homo sapiens cDNA clone IMAGE:73020 5, mRNA sequence	8.63	Down	1.23E-04
35	NM_178868	Homo sapiens chemokine-like factor super family 8 (CKLF8), mRNA	8.63	Down	6.22E-05
	NM_005130	Homo sapiens fibroblast growth factor binding protein 1 (FGFBP1), mRNA	8.62	Down	1.15E-04
40	NM_032488	Homo sapiens cornifelin (CNFN), mRNA	8.6	Down	3.05E-05
	NM_052960	Homo sapiens retinol binding protein 7, cellular (RBP7), mRNA	8.6	Down	2.05E-04
	NM_207517	Homo sapiens ADAMTS-like 3 (ADAMTSL3), mRNA	8.57	Down	2.22E-04
45	AK127644	Homo sapiens cDNA FLJ45742 fis, clone KIDNE2016327	8.54	Down	3.24E-03
	BU727096	UI-E-CR0-ach-e-12-0-UI.s1 UI-E-CR0 Homo sapiens cDNA clone UI-E-CR0-ach-e-12-0-UI 3, mRNA sequence	8.53	Down	9.29E-04
50	NM_015973	Homo sapiens galanin (GAL), mRNA	8.53	Down	6.30E-05
	NM_033514	Homo sapiens LIM and senescent cell antigen-like domains 3 (LIMS3), mRNA	8.51	Down	3.30E-04
	NM_138811	Homo sapiens chromosome 7 open reading frame 31 (C7orf31 mRNA	8.47	Down	8.35E-05
55	NM_005302	Homo sapiens G protein-coupled receptor37 (endothelin receptor type B-like) (GPR37), mRNA	8.47	Down	2.19E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_173462	Homo sapiens papilin, proteoglycan-like sulfated glycoprotein (PAPLN), mRNA	8.46	Down	1.70E-04
10	NM_032148	Homo sapiens solute carrier family 41, member 2 (SLC41A2), mRNA	8.44	Down	9.03E-05
	NM_000856	Homo sapiens guanylate cyclase 1, soluble, alpha 3 (GUCY1A3), mRNA	8.37	Down	1.17E-03
15	AW591723	xt85h10.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2793283 3 similar to contains element MER32 repetitive element ;, mRNA sequence	8.36	Down	4.86E-04
	NM_032024	Homo sapiens chromosome 10 open reading frame 11 (C10orf11 mRNA	8.25	Down	1.80E-04
20	AK024850	Homo sapiens cDNA: FLJ21197 fis, clone COL00201	8.21	Down	7.71E-05
	NM_002206	Homo sapiens integrin, alpha 7 (ITGA7), mRNA	8.2	Down	6.30E-05
	A1963999	wt87g07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2514492 3, mRNA sequence	8.19	Down	1.61E-03
	NM_005567	Homo sapiens lectin, galactoside-binding, soluble, 3 binding protein (LGALS3BP), mRNA	8.18	Down	8.76E-05
25	D87454	Human mRNA for KIAA0265 gene, partial cds	8.14	Down	1.08E-03
	BG564960	602583930F1 NIH_MGC_76 Homo sapiens cDNA clone IMAGE:4711807 5, mRNA sequence	8.12	Down	7.71E-05
	NM_022168	Homo sapiens interferon induced with helicase C domain 1 (IFIH1), mRNA	8.12	Down	1.46E-04
30	AK027541	Homo sapiens cDNA FLJ14635 fis, clone NT2RP2001196	8.12	Down	2.08E-04
	NM_024997	Homo sapiens activating transcription factor 7 interacting protein 2 (ATF7IP2), mRNA	8.11	Down	1.65E-03
35	A1926616	wo48e04.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2458590 3 similar to contains MER27.b2 MER27 repetitive element;, mRNA sequence	8.08	Down	6.30E-05
	S70348	Homo sapiens integrin beta 3 mRNA, partial cds, alternatively spliced	8.06	Drown	2.21E-04
40	NM_005725	Homo sapiens tetraspan 2 (TSPAN-2), mRNA	8.05	Drown	3.97E-03
	CA311343	UI-CF-FNO-aff-b-19-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FNO-aff-b-19-0-UI 3, mRNA sequence	8.02	Down	8.51E-06
45	NM_001710	Homo sapiens B-factor, properdin (BF), mRNA	8.01	Down	1.30E-03
	NM_002247	Homo sapiens potassium large conductance calcium-activated channel, subfamily M, alpha member 1 (KCNMA1 mRNA	8	Down	1.29E-03
50	BG149255	nad25d01.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3366553 3, mRNA sequence	8	Down	3.85E-04
	NM_153634	Homo sapiens copine VIII (CPNE8), mRNA	7.97	Down	3.57E-04
	NM_004390	Homo sapiens cathepsin H (CTSH), transcript variant 1, mRNA	7.97	Down	7.71 E-05
55	NM_016613	Homo sapiens hypothetical protein DKFZp434L142 (DKFZp434L142), mRNA	7.97	Down	3.72E-04
	NM_001845	Homo sapiens collagen, type IV, alpha 1 (COL4A1 mRNA	7.93	Down	8.65E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_005502	Homo sapiens ATP-binding cassette, sub-family A (ABC1), member 1 (ABCA1 mRNA	7.93	Down	5.18E-04
10	NM_018650	Homo sapiens MAP/microtubule affinity-regulating kinase 1 (MARK1), mRNA	7.92	Down	1.23E-04
	D62831	HUM330B12B Clontech human aorta polyA+ mRNA (#6572) Homo sapiens cDNA clone GEN-330B12 5, mRNA sequence	7.92	Down	9.23E-04
15	CA313095	UI-CF-FN0-aex-f-01-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aex-f-01-0-UI 3, mRNA sequence	7.91	Down	7.42E-04
	BC038556	Homo sapiens, clone IMAGE:3446976, mRNA	7.91	Down	2.57E-03
20	AI493349	tg70f04.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2114143 3, mRNA sequence	7.9	Down	1.14E-04
	BC015108	Homo sapiens, Similar to otoconin 90, clone IMAGE:4044247, mRNA	7.9	Down	1.01E-04
	AI819863	wj45h05.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2405817 3, mRNA sequence	7.87	Down	2.39E-03
25	BM802920	AGENCOURT_6457446 NIH_MGC_88 Homo sapiens cDNA clone IMAGE:5560288 5, mRNA sequence	7.85	Down	1.60E-03
	AK057113	Homo sapiens cDNA FLJ32551 fis, clone SPLEN1000087	7.83	Down	1.20E-03
30	NM_005025	Homo sapiens serine (or cysteine) proteinase inhibitor, clade I (neuroserpin), member 1 (SERPINI1), mRNA	7.82	Down	2.21E-04
	AA195328	zr34f08.s1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:665319 3, mRNA sequence	7.81	Down	2.37E-04
	NM_014358	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	7.81	Down	5.03E-03
35	BI493986	df106g12.y1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2538815 5, mRNA sequence	7.8	Down	6.69E-04
40	NM_152366	Homo sapiens kelch/ankyrin repeat containing cyclin A1 interacting protein (KARCA1), transcript variant 1, mRNA	7.79	Down	8.97E-05
	NM_015678	Homo sapiens neurobeachin (NBEA), mRNA	7.78	Down	2.12E-04
	NM_018424	Homo sapiens erythrocyte membrane protein band 4.1 like 4B (EPB41L4B), mRNA	7.77	Down	1.23E-04
45	CA310979	UI-CF-FN0-afc-c-21-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afc-c-21-0-UI 3, mRNA sequence	7.76	Down	1.86E-04
	NM_031476	Homo sapiens hypothetical protein DKFZp434B044 (DKFZP434B044), mRNA	7.75	Down	2.20E-03
50	NM_004086	Homo sapiens coagulation factor C homolog, cochlin (Limulus polyphemus) (COCH), mRNA	7.74	Down	2.21E-04
	NM_003617	Homo sapiens regulator of G-protein signalling 5 (RGS5), mRNA	7.73	Down	3.85E-04
	BF966833	60228668T1 NIH_MGC_95 Homo sapiens cDNA clone IMAGE:4375360 3, mRNA sequence	7.73	Down	6.10E-05
55	AK024270	Homo sapiens cDNA FLJ14208 fis, clone NT2RP3003264	7.71	Down	1.13E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_144587	Homo sapiens chromosome 10 open reading frame 87 (C10orf87), mRNA	7.71	Down	6.03E-04
10	AK096288	Homo sapiens cDNA FLJ38969 fis, clone NT2RI2002359	7.7	Down	1.49E-04
	BE968596	601649770F1 NIH_MGC_74 Homo sapiens cDNA clone IMAGE:3933472 5, mRNA sequence	7.69	Down	2.39E-03
	AY358775	Homo sapiens clone DNA170212 WGAR9166 (UNQ9166) mRNA, complete cds	7.66	Down	2.16E-03
15	BX103949	BX103949 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998C112296 ; IMAGE:923842, mRNA sequence	7.64	Down	3.18E-04
	NM_013410	Homo sapiens adenylate kinase 3 (AK3), nuclear gene encoding mitochondrial protein, transcript variant 2, mRNA	7.64	Down	3.07E-04
20	NM_005204	Homo sapiens mitogen-activated protein kinase kinase kinase 8 (MAP3K8), mRNA	7.63	Down	6.82E-04
	NM_032211	Homo sapiens lysyl oxidase-like 4 (LOXL4), mRNA	7.6	Down	3.30E-04
	AA325746	EST28794 Cerebellum II Homo sapiens cDNA 5 end, mRNA sequence	7.57	Down	6.30E-05
25	H94320	yv18b10.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:243067 3, mRNA sequence	7.55	Down	3.35E-03
	BX640973	Homo sapiens mRNA; cDNA DKFZp686B15184 (from clone DKFZp686B15184)	7.55	Down	7.89E-04
30	BQ015616	UI-1-BC1-ajb-g-04-0-UI.s1 NCI_CGAP_PI2 Homo sapiens cDNA clone UI-1-BC1-ajb-g-04-0-UI 3, mRNA sequence	7.54	Down	5.28E-04
	AK128715	Homo sapiens cDNA FLJ46882 fis, clone UTERU3015844	7.53	Down	2.30E-04
35	NM_015068	Homo sapiens paternally expressed 10 (PEG10), mRNA	7.52	Down	7.71E-05
	BU754480	UI-1-BB1p-axz-h-11-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-axz-h-11-0-UI 3, mRNA sequence	7.49	Down	6.24E-04
	AK074097	Homo sapiens mRNA for FLJ00168 protein	7.48	Down	1.93E-03
40	BC033567	Homo sapiens, clone IMAGE:4822266, mRNA	7.48	Down	1.96E-04
	NM_018894	Homo sapiens EGF-containing fibulin-like extracellular matrix protein 1 (EFEMP1), transcript variant 2, mRNA	7.47	Down	2.67E-04
	BM701989	UI-E-CQ1-aex-j-06-0-UI.r1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aex-j-06-0-UI 5, mRNA sequence	7.47	Down	4.53E-04
45	NM_016423	Homo sapiens zinc finger protein 219 (ZNF219), mRNA	7.46	Down	1.61E-04
	BX117317	BX117317 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998E242234 ; IMAGE:900095, mRNA sequence	7.45	Down	1.13E-03
	NM_001993	Homo sapiens coagulation factor III (thromboplastin, tissue factor) (F3), mRNA	7.45	Down	5.82E-04
50	CA502991	UI-CF-FN0-afp-g-01-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afp-g-01-0-UI 3, mRNA sequence	7.43	Down	2.24E-02
55	AI686890	tp90h02.x1 NCI_CGAP_Ut3 Homo sapiens cDNA clone IMAGE:2206611 3, mRNA sequence	7.41	Down	1.49E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AW137001	UI-H-BI1-acu-c-05-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2715632 3, mRNA sequence	7.4	Down	7.95E-05
10	BQ021695	UI-H-DH1-axi-f-22-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5829141 3, mRNA sequence	7.39	Down	6.10E-05
15	BX640643	Homo sapiens mRNA; cDNA DKFZp686O24114 (from clone DKFZp686O24114)	7.36	Down	5.18E-05
	NM_004669	Homo sapiens chloride intracellular channel 3 (CLIC3), mRNA	7.35	Down	4.30E-03
	NM_001955	Homo sapiens endothelin 1 (EDN1), mRNA	7.33	Down	7.72E-04
20	AB007974	Homo sapiens mRNA, chromosome 1 specific transcript KIAA0505	7.33	Down	5.29E-03
	NM_198174	Homo sapiens transcription factor CP2-like 4 (TFCP2L4), transcript variant 3, mRNA	7.32	Down	4.82E-03
	BX098521	BX098521 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGp998L05118 ; IMAGE:123412, mRNA sequence	7.3	Down	5.51E-03
25	NM_004154	Homo sapiens pyrimidinergic receptor P2Y, G-protein coupled, 6 (P2RY6), transcript variant 4, mRNA	7.28	Down	3.41E-03
	AK123617	Homo sapiens cDNA FLJ41623 fis, clone CTONG3009227	7.27	Down	2.26E-03
30	BF111903	7I38d07.x0 Soares_NSF_F8_9W_OT_PA_P_S Homo sapiens cDNA clone IMAGE:3523644 3, mRNA sequence	7.25	Down	1.34E-03
	NM_018655	Homo sapiens lens epithelial protein (LENEP), mRNA	7.24	Down	1.07E-04
35	NM_139161	Homo sapiens crumbs homolog 3 (Drosophila) (CRB3), transcript variant 2, mRNA	7.23	Down	8.46E-04
	BC060805	Homo sapiens hypothetical protein FLJ12788, mRNA (cDNA clone IMAGE:5266931), partial cds	7.22	Down	7.45E-05
40	CA413744	UI-H-EZ0-bat-h-12-0-UI.s1 NCI_CGAP_Ch1 Homo sapiens cDNA clone UI-H-EZ0-bat-h-12-0-UI 3, mRNA sequence	7.22	Down	6.09E-04
	NM_018986	Homo sapiens SH3 domain and tetratricopeptide repeats 1 (SH3TC1), mRNA	7.21	Down	2.68E-03
45	NM_024677	Homo sapiens hypothetical protein FLJ14001 (FLJ14001), mRNA	7.21	Down	1.14E-04
	AL117454	Homo sapiens mRNA; cDNA DKFZp586J1717 (from clone DKFZp586J1717)	7.2	Down	2.93E-04
	AW450938	UI-H-BI3-all-g-05-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2737329 3, mRNA sequence	7.19	Down	1.19E-04
50	NM_017899	Homo sapiens hypothetical protein FLJ20607 (TSC), mRNA	7.16	Down	4.80E-03
55	S81734	tissue transglutaminase homologue {alternatively spliced} [human, erythroleukemia cell line HEL GM06141A, mRNA, 2362 nt]	7.16	Down	8.76E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	BU616749	UI-H-FH1-bfj-a-11-0-UI.s1 NCI_CGAP_FH1 Homo sapiens cDNA clone UI-H-FH1-bfj-a-11-0-UI 3, mRNA sequence	7.14	Down	2.21E-04
10	NM_006681	Homo sapiens neuromedin U (NMU), mRNA	7.1	Down	6.52E-03
	AK022598	Homo sapiens cDNA FLJ12536 fis, clone NT2RM4000265	7.09	Down	1.15E-04
15	AW 137116	UI-H-BI1-acp-f-03-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2715029 3, mRNA sequence	7.07	Down	3.50E-03
	BX647876	Homo sapiens mRNA; cDNA DKFZp313A1525 (from clone DKFZp313A1525)	7.06	Down	1.41E-03
20	NM_000597	Homo sapiens insulin-like growth factor binding protein 2, 36kDa (IGFBP2), mRNA	7.04	Down	2.99E-03
	CN478597	UI-CF-FN0-aeo-g-21-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aeo-g-21-0-UI 3, mRNA sequence	7.04	Down	1.43E-03
25	CN478714	UI-CF-FN0-afu-c-19-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afu-c-19-0-UI 3, mRNA sequence	7.04	Down	3.55E-03
	AB020640	Homo sapiens mRNA for KIAA0833 protein, partial cds	7	Down	2.45E-03
30	NM_022746	Homo sapiens hypothetical protein FLJ22390 (FLJ22390), mRNA	6.99	Down	9.12E-04
	AI905628	CM-BT094-050299-147 BT094 Homo sapiens cDNA, mRNA sequence	6.98	Down	1.54E-04
	NM_002193	Homo sapiens inhibin, beta B (activin AB beta polypeptide) (INHBB), mRNA	6.96	Down	7.59E-04
35	NM_004490	Homo sapiens growth factor receptor-bound protein 14 (GRB14), mRNA	6.94	Down	1.98E-03
	NM_003985	Homo sapiens tyrosine kinase, non-receptor, 1 (TNK1), mRNA	6.93	Down	2.10E-04
40	NM_000480	Homo sapiens adenosine monophosphate deaminase (isoform E) (AMPD3), mRNA	6.92	Down	1.06E-04
	AK094292	Homo sapiens cDNA FLJ36973 fis, clone BRACE2006249	6.92	Down	2.03E-03
	BC033124	Homo sapiens, clone IMAGE:2960615, mRNA	6.88	Down	7.22E-05
45	NM_000612	Homo sapiens insulin-like growth factor 2 (somatomedin A) (IGF2), mRNA	6.88	Down	2.60E-04
	BC042976	Homo sapiens cDNA clone IMAGE:5295023, partial cds	6.88	Down	2.39E-03
	BF445031	nad20f02.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3366266 3, mRNA sequence	6.87	Down	5.18E-04
50	NM_004170	Homo sapiens solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1 (SLC1A1), mRNA	6.87	Down	3.18E-04
	NM_199169	Homo sapiens transmembrane, prostate androgen induced RNA (TMEPAI), transcript variant 2, mRNA	6.84	Down	1.45E-03
55	NM_019644	Homo sapiens ankyrin repeat domain 7 (ANKRD7), mRNA	6.83	Down	8.99E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	W20132	zb40c10.r1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:306066 5, mRNA sequence	6.82	Down	6.79E-03
10	BC033116	Homo sapiens chromodomain helicase DNA binding protein 7, mRNA (cDNA clone IMAGE:3352674), partial cds	6.82	Down	8.03E-05
	NM_018650	Homo sapiens MAP/microtubule affinity-regulating kinase 1 (MARK1), mRNA	6.8	Down	4.97E-03
15	BU563992	AGENCOURT_10371176 NIH_MGC_141 Homo sapiens cDNA clone IMAGE:6601829 5, mRNA sequence	6.79	Down	1.29E-02
	AL119769	DKFZp761E1224_r1 761 (synonym: hamy2) Homo sapiens cDNA clone DKFZp761 E1224 5, mRNA sequence	6.78	Down	2.52E-03
20	NM_052997	Homo sapiens ankyrin repeat domain 30A (ANKRD30A), mRNA	6.78	Down	2.45E-03
	NM_024704	Homo sapiens chromosome 20 open reading frame 23 (C20orf23), mRNA	6.76	Down	7.45E-04
25	NM_000495	Homo sapiens collagen, type IV, alpha 5 (Alport syndrome) (COL4A5), transcript variant 1, mRNA	6.74	Down	4.77E-03
	BX102895	BX102895 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGp998118520 ; IMAGE:242009, mRNA sequence	6.7	Down	3.04E-04
30	BU753362	UI-1-BB1-air-h-09-0-UI.s1 NCI_CGAP_PI5 Homo sapiens cDNA clone UI-1-BB1-air-h-09-0-UI 3, mRNA sequence	6.7	Down	7.42E-05
	AA190552	zp86b11.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone IMAGE:627069 3, mRNA sequence	6.67	Down	1.26E-03
35	NM_004024	Homo sapiens activating transcription factor 3 (ATF3), mRNA	6.66	Down	3.12E-03
	NM_144650	Homo sapiens alcohol dehydrogenase, iron containing, 1 (ADHFE1), mRNA	6.65	Down	1.45E-04
40	NM_002800	Homo sapiens proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 2) (PSMB9), transcript variant 1, mRNA	6.63	Down	1.70E-04
	NM_021977	Homo sapiens solute carrier family 22 (extraneuronal monoamine transporter), member 3 (SLC22A3), mRNA	6.62	Down	2.05E-03
45	AW516579	xq01f06.x1 Soares_NHCeC_cervical_tumor Homo sapiens cDNA clone IMAGE:2748611 3, mRNA sequence	6.62	Down	4.65E-03
	AI311296	ta48d10.x2 NCI_CGAP_Lu25 Homo sapiens cDNA clone IMAGE:2047315 3, mRNA sequence	6.61	Down	1.72E-03
50	BC040204	Homo sapiens, clone IMAGE:4821815, mRNA	6.61	Down	5.66E-04
	AA340011	EST45155 Fetal skin Homo sapiens cDNA 5 end, mRNA sequence	6.58	Down	6.30E-05
	NM_025044	Homo sapiens bicaudal C homolog 1 (Drosophila) (BICC1), mRNA	6.57	Down	6.30E-05
55	BX111520	BX111520 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998L15208 ; IMAGE:141470, mRNA sequence	6.57	Down	1.70E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AW444925	UI-H-BI3-ajz-f-09-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2733473 3, mRNA sequence	6.52	Down	2.55E-03
10	NM_003475	Homo sapiens chromosome 11 open reading frame 13 (C11orf13), mRNA	6.5	Down	5.88E-04
	AK124778	Homo sapiens cDNA FLJ42788 fis, clone BRAWH3007129	6.48	Down	8.61E-04
15	NM_024563	Homo sapiens hypothetical protein FLJ14054 (FLJ14054), mRNA	6.44	Down	4.72E-03
	NM_002214	Homo sapiens integrin, beta 8 (ITGB8), mRNA	6.43	Down	8.46E-05
	NM_178814	Homo sapiens adaptor-related protein complex 1, sigma 3 subunit (AP1S3), mRNA	6.42	Down	1.78E-04
20	AI831068	wj62d12.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2407415 3, mRNA sequence	6.41	Down	3.72E-04
	NM_000186	Homo sapiens complement factor H (CFH), mRNA	6.39	Down	1.74E-03
	NM_000216	Homo sapiens Kallmann syndrome 1 sequence (KAL1), mRNA	6.37	Down	8.59E-05
25	NM_015478	Homo sapiens I(3)mbt-like (Drosophila) (L3MBTL), transcript variant I, mRNA	6.36	Down	7.22E-05
	BX096609	BX096609 Soares retina N2b4HR Homo sapiens cDNA clone IMAGp998L12439 ; IMAGE:221339, mRNA sequence	6.36	Down	1.24E-03
30	NM_030970	Homo sapiens hypothetical protein MGC3771 (MGC3771), mRNA	6.34	Down	1.69E-04
	BQ007085	UI-H-EI1-azc-k-11-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5846914 3, mRNA sequence	6.33	Down	1.19E-04
35	BC039329	Homo sapiens, clone IMAGE:5267606, mRNA	6.33	Down	6.61 E-03
	NM_001165	Homo sapiens baculoviral IAP repeat-containing 3 (BIRC3), transcript variant 1, mRNA	6.33	Down	3.42E-03
40	NM_004165	Homo sapiens Ras-related associated with diabetes (RRAD), mRNA	6.33	Down	4.68E-03
	NM_001928	Homo sapiens D component of complement (adipsin) (DF), mRNA	6.32	Down	3.68E-03
	NM_018670	Homo sapiens mesoderm posterior 1 (MESP1), mRNA	6.31	Down	5.76E-03
45	NM_153229	Homo sapiens hypothetical protein FLJ33318 (FLJ33318), mRNA	6.31	Down	2.57E-03
	AA912845	ol32a12.s1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1525150 3, mRNA sequence	6.29	Down	1.13E-03
50	AK127437	Homo sapiens cDNA FLJ45529 fis, clone BRTHA2027546	6.28	Down	3.37E-04
	NM_020836	Homo sapiens brain-enriched guanylate kinase-associated protein (KIAA1446), mRNA	6.27	Down	6.60E-03
55	AL512697	Homo sapiens mRNA; cDNA DKFZp547F134 (from clone DKFZp547F134)	6.25	Down	1.36E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_005211	Homo sapiens colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog (CSF1 R), mRNA	6.24	Down	5.07E-04
10	AK124776	Homo sapiens cDNA FLJ42786 fis, clone BRAWH3006761	6.2	Down	1.15E-04
	NM_173662	Homo sapiens hypothetical protein LOC285533 (LOC285533), mRNA	6.18	Down	1.05E-02
15	NM_014391	Homo sapiens ankyrin repeat domain 1 (cardiac muscle) (ANKRD1), mRNA	6.18	Down	3.52E-03
	NM_005860	Homo sapiens follistatin-like 3 (secreted glycoprotein) (FSTL3), mRNA	6.18	Down	5.23E-05
20	NM_001045	Homo sapiens solute carrier family 6 (neurotransmitter transporter, serotonin), member 4 (SLC6A4), mRNA	6.17	Down	5.46E-03
	NM_002147	Homo sapiens homeo box B5 (HOXB5), mRNA	6.17	Down	1.05E-04
	AI288404	qv89b01.x1 NCI_CGAP_Ut2 Homo sapiens cDNA clone IMAGE:1988713 3, mRNA sequence	6.16	Down	3.74E-04
25	NM_173584	Homo sapiens hypothetical protein MGC45840 (MGC45840), mRNA	6.16	Down	1.32E-04
	N62729	yz76g05.s1 Soares_multiple_sclerosis_2NbHMS P Homo sapiens cDNA clone IMAGE:289016 3, mRNA sequence	6.12	Down	1.37E-03
30	AK024238	Homo sapiens cDNA FLJ14176 fis, clone NT2RP2003101	6.11	Down	9.69E-05
	NM_001843	Homo sapiens contactin 1 (CNTN1), transcript variant 1, mRNA	6.1	Down	4.80E-03
	M_152433	Homo sapiens kelch repeat and BTB (POZ) domain containing 3 (KBTBD3), transcript variant 1, mRNA	6.1	Down	2.39E-03
35	BX649112	Homo sapiens mRNA; cDNA DKFZp686E02109 (from clone DKFZp686E02109)	6.1	Down	2.24E-04
	AA411988	zt65g11.s1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:727268 3, mRNA sequence	6.09	Down	1.40E-03
40	R80806	yi94f01.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:146905 3, mRNA sequence	6.08	Down	1.54E-04
	NM_002843	Homo sapiens protein tyrosine phosphatase, receptor type, J (PTPRJ), mRNA	6.08	Down	6.54E-03
45	NM_003979	Homo sapiens G protein-coupled receptor, family C, group 5, member A (GPCR5A), mRNA	6.06	Down	8.75E-03
	AW 129281	xf23a03.x1 NCI_CGAP_Kid8 Homo sapiens cDNA clone IMAGE:2618860 3 similar to gb:X58295_rna1 PLASMA GLUTATHIONE PEROXIDASE PRECURSOR (HUMAN);, mRNA sequence	6.06	Down	1.70E-04
50	NM_015345	Homo sapiens dishevelled associated activator of morphogenesis 2 (DAAM2), mRNA	6.05	Down	1.09E-03
	L07615	Human neuropeptide Y receptor Y1 (NPYY1) mRNA, exon 2-3 and complete cds	6.05	Down	5.51E-03
55	NM_016179	Homo sapiens transient receptor potential cation channel, subfamily C, member 4 (TRPC4), mRNA	6.04	Down	8.59E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_182797	Homo sapiens phospholipase C, beta 4 (PLCB4), transcript variant 2, mRNA	6.03	Down	4.40E-04
10	AB032945	Homo sapiens mRNA for KIAA1119 protein, partial cds	6.03	Down	1.11E-02
	CA444471	UI-H-DP0-avv-a-16-0-UI.s1 NCI_CGAP_Fs1 Homo sapiens cDNA clone UI-H-DP0-avv-a-16-0-UI 3, mRNA sequence	6.02	Down	1.02E-03
15	AK127421	Homo sapiens cDNA FLJ45513 fis, clone BRTHA2021450	6.01	Down	1.60E-04
	AV736303	AV736303 CB Homo sapiens cDNA clone CBCAJD04 5, mRNA sequence	6.01	Down	8.97E-05
	BX537613	Homo sapiens mRNA; cDNA DKFZp686E11117 (from clone DKFZp686E11117)	6	Down	4.89E-04
20	AB023211	Homo sapiens mRNA for KIAA0994 protein, partial cds	5.99	Down	5.50E-04
	NM_018349	Homo sapiens multiple C2-domains with two transmembrane regions 2 (MCTP2), mRNA	5.99	Down	2.48E-04
	NM_000087	Homo sapiens cyclic nucleotide gated channel alpha 1 (CNGA1), mRNA	5.98	Down	3.29E-03
25	BF939416	nad89c02.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:3410667 3, mRNA sequence	5.97	Down	9.03E-04
	NM_013951	Homo sapiens paired box gene 8 (PAX8), transcript variant PAX8B, mRNA	5.97	Down	2.57E-03
30	AW 195474	xn38g09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2696032 3, mRNA sequence	5.97	Down	1.68E-04
	R40050	yf68h07.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:27726 3, mRNA sequence	5.94	Down	5.78E-03
	NM_021101	Homo sapiens claudin 1 (CLDN1), mRNA	5.9	Down	2.60E-04
35	AA602964	no97c02.s1 NCI_CGAP_Pr2 Homo sapiens cDNA clone IMAGE:1114754, mRNA sequence	5.9	Down	1.23E-03
	NM_006255	Homo sapiens protein kinase C, eta (PRKCH), mRNA	5.89	Down	2.78E-04
	NM_024103	Homo sapiens solute carrier family 25 (mitochondrial carrier; phosphate carrier), member 23 (SLC25A23), mRNA	5.89	Down	7.42E-05
40	AK093870	Homo sapiens cDNA FLJ36551 fis, clone TRACH2008127	5.88	Down	2.28E-02
	BE070450	QV4-BT0407-020300-122-d08 BT0407 Homo sapiens cDNA, mRNA sequence	5.85	Down	3.53E-04
45	NM_003706	Homo sapiens phospholipase A2, group IVC (cytosolic, calcium-independent) (PLA2G4C), mRNA	5.85	Down	7.81E-04
	NM_023927	Homo sapiens HCV NS3-transactivated protein 2 (NS3TP2), mRNA	5.85	Down	2.64E-03
50	AF519622	Homo sapiens noncoding mRNA sequence	5.84	Down	7.70E-03
	NM_198582	Homo sapiens FLJ43374 protein (FLJ43374), mRNA	5.84	Down	4.05E-03
	AK026283	Homo sapiens cDNA: FLJ22630 fis, clone HSI06250	5.84	Down	5.32E-04
	CB115754	K-EST0159876 L8SCK0 Homo sapiens cDNA clone L8SCK0-8-H08 5, mRNA sequence	5.81	Down	9.71E-04
55	BQ188285	UI-E-EJ1-ajp-n-20-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajp-n-20-0-UI 5, mRNA sequence	5.8	Down	5.47E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_003662	Homo sapiens pirin (iron-binding nuclear protein) (PIR), mRNA	5.78	Down	1.71E-04
10	BX098660	BX098660 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998L03214 ; IMAGE:143762, mRNA sequence	5.77	Down	6.00E-04
	NM_005360	Homo sapiens v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian) (MAF), mRNA	5.77	Down	2.24E-04
15	NM_031847	Homo sapiens microtubule-associated protein 2 (MAP2), transcript variant 4, mRNA	5.77	Down	1.34E-03
	NM_003057	Homo sapiens solute carrier family 22 (organic cation transporter), member 1 (SLC22A1), transcript variant 1, mRNA	5.75	Down	2.42E-04
20	AW242323	xm96f03.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2692061 3 similar to contains Alu repetitive element,, mRNA sequence	5.74	Down	1.92E-04
	NM_052890	Homo sapiens peptidoglycan recognition protein 2 (PGLYRP2), mRNA	5.73	Down	1.14E-04
25	NM_000593	Homo sapiens transporter 1, ATP-binding cassette, sub-family B (MDR/TAP) (TAP1), mRNA	5.72	Down	4.86E-03
	NM_033132	Homo sapiens Zic family member 5 (odd-paired homolog, Drosophila) (ZIC5), mRNA	5.71	Down	1.76E-04
30	AL832380	Homo sapiens mRNA; cDNA DKFZp451L157 (from clone DKFZp451L157)	5.71	Down	1.54E-04
	NM_183376	Homo sapiens arrestin domain containing 4 (ARRDC4), mRNA	5.71	Down	5.88E-04
	NM_018700	Homo sapiens tripartite motif-containing 36 (TRIM36), mRNA	5.7	Down	1.32E-02
35	NM_000782	Homo sapiens cytochrome P450, family 24, subfamily A, polypeptide 1 (CYP24A1), nuclear gene encoding mitochondrial protein, mRNA	5.69	Down	1.82E-02
	BC035805	Homo sapiens caspase recruitment domain family, member 9, mRNA (cDNA clone IMAGE:5745585), partial cds	5.68	Down	4.00E-03
40	BC051727	Homo sapiens cDNA clone IMAGE:5265929, partial cds	5.68	Down	5.13E-04
	NM_022842	Homo sapiens CUB domain-containing protein 1 (CDCP1), transcript variant 1, mRNA	5.68	Down	1.85E-03
45	NM_004335	Homo sapiens bone marrow stromal cell antigen 2 (BST2), mRNA	5.67	Down	6.01E-03
	NM_053039	Homo sapiens UDP glycosyltransferase 2 family, polypeptide B28 (UGT2B28), mRNA	5.66	Down	8.07E-03
50	NM_004529	Homo sapiens myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 3 (MLLT3), mRNA	5.66	Down	4.02E-04
	AK093069	Homo sapiens cDNA FLJ35750 fis, clone TEST12004539, weakly similar to Homo sapiens adlcan mRNA	5.65	Down	1.10E-03
55	NM_005141	Homo sapiens fibrinogen, B beta polypeptide (FGB), mRNA	5.63	Down	7.89E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_000682	Homo sapiens adrenergic, alpha-2B-, receptor (ADRA2B), mRNA	5.61	Down	1.14E-04
10	BF512326	UI-H-BW1-amb-g-12-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069503 3, mRNA sequence	5.6	Down	1.49E-04
	NM_003726	Homo sapiens src family associated phosphoprotein 1 (SCAP1), mRNA	5.6	Down	2.42E-03
15	AK024261	Homo sapiens cDNA FLJ14199 fis, clone NT2RP3002713	5.6	Down	4.42E-04
	AK125695	Homo sapiens cDNA FLJ43707 fis, clone TESOP2001865	5.59	Down	1.10E-02
20	NM_006074	Homo sapiens tripartite motif-containing 22 (TRIM22), mRNA	5.58	Down	1.94E-03
	AW591461	x192h06.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2682203 3, mRNA sequence	5.58	Down	1.95E-02
	NM_022128	Homo sapiens ribokinase (RBKS), mRNA	5.58	Down	1.87E-04
25	N66105	yy65e06.s1 Soares_multiple_sclerosis_2NbHMS P Homo sapiens cDNA clone IMAGE:278434 3, mRNA sequence	5.57	Down	2.84E-04
	BC035116	Homo sapiens cDNA clone IMAGE:5263177, partial cds	5.56	Down	5.49E-04
30	BM988642	UI-H-DH0-arx-p-21-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone IMAGE:5855492 3, mRNA sequence	5.56	Down	2.48E-03
	NM_003947	Homo sapiens huntingtin-associated protein interacting protein (duo) (HAPIP), mRNA	5.56	Down	9.71E-04
	AI697906	we18f06.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:2341475 3, mRNA sequence	5.56	Down	3.52E-03
35	AK095399	Homo sapiens cDNA FLJ38080 fis, clone CTONG2016185	5.55	Down	7.43E-03
	AL110252	Homo sapiens mRNA; cDNA DKFZp566A1046 (from clone DKFZp566A1046)	5.54	Down	3.44E-03
40	NM_024572	Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 14 (GalNAc-T14) (GALNT14), mRNA	5.54	Down	5.93E-03
45	NM_000775	Homo sapiens cytochrome P450, family 2, subfamily J, polypeptide 2 (CYP2J2), mRNA	5.53	Down	1.57E-03
	NM_018317	Homo sapiens hypothetical protein FLJ11082 (FLJ11082), mRNA	5.51	Down	5.10E-04
	NM_030915	Homo sapiens likely ortholog of mouse limb-bud and heart gene (LBH), mRNA	5.51	Down	6.86E-04
50	NM_139241	Homo sapiens FYVE, RhoGEF and PH domain containing 4 (FGD4), mRNA	5.5	Down	9.42E-03
	CB047092	NISC_gf08f03.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:3253013 3, mRNA sequence	5.5	Down	5.38E-04
55	NM_005711	Homo sapiens EGF-like repeats and discoidin I-like domains 3 (EDIL3), mRNA	5.5	Down	6.04E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	AA974968	on02e08.s1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1555526 3, mRNA sequence	5.49	Down	1.63E-03
10	NM_000129	Homo sapiens coagulation factor XIII, A1 polypeptide (F13A1), mRNA	5.49	Down	1.09E-02
	NM_007366	Homo sapiens phospholipase A2 receptor 1, 180kDa (PLA2R1), transcript variant 1, mRNA	5.47	Down	4.66E-03
	AL353944	Homo sapiens mRNA; cDNA DKFZp761J1112 (from clone DKFZp761J1112)	5.46	Down	5.38E-04
15	AK095647	Homo sapiens cDNA FLJ38328 fis, clone FCBBF3025142	5.46	Down	1.61E-03
	BQ188860	UI-E-EJ1-ajx-h-03-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajx-h-03-0-UI 5, mRNA sequence	5.46	Down	3.33E-03
20	NM_017594	Homo sapiens DIRAS family, GTP-binding RAS-like 2 (DIRAS2), mRNA	5.44	Down	1.22E-02
	NM_014399	Homo sapiens transmembrane 4 superfamily member 13 (TM4SF13), mRNA	5.43	Down	2.64E-04
25	NM_033255	Homo sapiens epithelial stromal interaction 1 (breast) (EPSTI1), mRNA	5.43	Down	1.73E-03
	AW269270	xs35c11.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2771636 3 similar to contains L1.t3 L1 repetitive element ;, mRNA sequence	5.42	Down	1.33E-03
30	AK092456	Homo sapiens cDNA FLJ35137 fis, clone PLACE6009419	5.42	Down	1.09E-04
	NM_031935	Homo sapiens hemicentin (FIBL-6), mRNA	5.42	Down	7.69E-04
	BU751966	UI-1-BB0-acy-c-09-0-UI.s1 NCI_CGAP_PI4 Homo sapiens cDNA clone UI-1-BB0-acy-c-09-0-UI 3, mRNA sequence	5.4	Down	1.61E-03
35	NM_001747	Homo sapiens capping protein (actin filament), gelsolin-like (CAPG), mRNA	5.4	Down	1.33E-03
	NM_002260	Homo sapiens killer cell lectin-like receptor subfamily C, member 2 (KLRC2), mRNA	5.4	Down	1.21E-02
40	AA732841	zg77f01.s1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:399385 3, mRNA sequence	5.4	Down	3.48E-03
	AK055468	Homo sapiens cDNA FLJ30906 fis, clone FEBRA2006055	5.4	Down	1.36E-03
45	A1080164	oz48c05.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:1678568 3, mRNA sequence	5.38	Down	2.44E-03
	NM_005562	Homo sapiens laminin, gamma 2 (LAMC2), transcript variant 1, mRNA	5.38	Down	3.78E-04
	NM_080659	Homo sapiens similar to RIKEN cDNA 2310030G06 gene (MGC14839), mRNA	5.36	Down	3.81E-03
50	NM_015085	Homo sapiens GTPase activating Rap/RanGAP domain-like 4 (GARNL4), mRNA	5.36	Down	1.86E-04
	BX097034	BX097034 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGp998F14169 ; IMAGE:39685, mRNA sequence	5.36	Down	8.59E-05
55	NM_002147	Homo sapiens homeo box B5 (HOXB5), mRNA	5.35	Down	8.21E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_181785	Homo sapiens hypothetical protein LOC283537 (LOC283537), mRNA	5.33	Down	3.48E-04
10	NM_032857	Homo sapiens lactamase, beta (LACTB), nuclear gene encoding mitochondrial protein, transcript variant 1, mRNA	5.33	Down	2.45E-04
	BX113144	BX113144 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998N07225 ; IMAGE:148038, mRNA sequence	5.32	Down	1.53E-03
15	NM_014905	Homo sapiens glutaminase (GLS), mRNA	5.31	Down	1.63E-03
	NM_138396	Homo sapiens membrane-associated RING-CH protein IX (MARCH-IX), mRNA	5.31	Down	1.39E-04
20	NM_004433	Homo sapiens E74-like factor 3 (ets domain transcription factor, epithelial-specific) (ELF3), mRNA	5.3	Down	5.32E-05
	AA553553	nk78d11.s1 NCI_CGAP_Sch1 Homo sapiens cDNA clone IMAGE:1019637 3, mRNA sequence	5.27	Down	1.69E-03
	AW 188195	xj93e12.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2664814 3 similar to contains element THR repetitive element;, mRNA sequence	5.27	Down	5.32E-05
25	NM_006169	Homo sapiens nicotinamide N-methyltransferase (NNMT), mRNA	5.26	Down	7.45E-05
	NM_000071	Homo sapiens cystathionine-beta-synthase (CBS), mRNA	5.26	Down	1.49E-03
30	NM_001958	Homo sapiens eukaryotic translation elongation factor 1 alpha 2 (EEF1A2), mRNA	5.25	Down	5.20E-03
	AK055356	Homo sapiens cDNA FLJ30794 fis, clone FEBRA2001093, weakly similar to MONOCARBOXYLATE TRANSPORTER 4	5.25	Down	8.65E-04
35	NM_014578	Homo sapiens ras homolog gene family, member D (RHOD), mRNA	5.24	Down	5.43E-03
	NM_013230	Homo sapiens CD24 antigen (small cell lung carcinoma cluster 4 antigen) (CD24), mRNA	5.22	Down	7.65E-04
40	AF086158	Homo sapiens full length insert cDNA clone ZB72E12	5.2	Down	6.04E-03
	NM_004932	Homo sapiens cadherin 6, type 2, K-cadherin (fetal kidney) (CDH6), mRNA	5.18	Down	2.11E-04
	NM_020466	Homo sapiens hypothetical protein dJ12208.2 (DJ12208.2), mRNA	5.18	Down	2.99E-03
45	AL117425	Homo sapiens mRNA; cDNA DKFZp566L203 (from clone DKFZp566L203)	5.17	Down	8.64E-03
	AI732568	zo23d12.x5 Stratagene colon (#937204) Homo sapiens cDNA clone IMAGE:587735 3, mRNA sequence	5.17	Down	9.19E-04
50	AI150192	qf34d12.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1751927 3, mRNA sequence	5.17	Down	1.21E-02
	AK125852	Homo sapiens cDNA FLJ43864 fis, clone TEST14007799	5.17	Down	4.82E-03
	NM_017912	Homo sapiens hect domain and RLD 6 (HERC6), mRNA	5.15	Down	3.63E-04
55	NM_006174	Homo sapiens neuropeptide Y receptor Y5 (NPY5R), mRNA	5.14	Down	1.13E-02
	AK025281	Homo sapiens cDNA: FLJ21628 fis, clone COL08076	5.14	Down	9.71E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in fibroblast versus AF cells	Direction	adj. p-value
5	NM_153742	Homo sapiens cystathionase (cystathionine gamma-lyase) (CTH), transcript variant 2, mRNA	5.12	Down	1.27E-04
10	NM_003107	Homo sapiens SRY (sex determining region Y)-box 4 (SOX4), mRNA	5.11	Down	3.48E-04
	NM_003948	Homo sapiens cyclin-dependent kinase-like 2 (CDC2-related kinase) (CDKL2), mRNA	5.11	Down	2.05E-03
15	NM_173660	Homo sapiens hypothetical protein FLJ33718 (FLJ33718), mRNA	5.11	Down	1.30E-03
	AI693580	wd12d01.x1 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGE:2327905 3, mRNA sequence	5.1	Down	6.19E-03
20	NM_004482	Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 3 (GalNAc-T3) (GALNT3), mRNA	5.1	Down	2.31E-03
25	NM_030965	Homo sapiens sialyltransferase 7 ((alpha-N-acetylneuraminyl-2,3-beta-galactosyl-1,3)-N-acetyl galactosaminide alpha-2,6-sialyltransferase) E (SIAT7E), mRNA	5.1	Down	2.61E-04
	NM_016147	Homo sapiens protein phosphatase methylesterase-1 (PME-1), mRNA	5.09	Down	9.71E-04
30	NM_006259	Homo sapiens protein kinase, cGMP-dependent, type II (PRKG2), mRNA	5.08	Down	1.36E-03
	NM_002993	Homo sapiens chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2) (CXCL6), mRNA	5.07	Down	2.32E-03
	BE671338	7e49f03.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3285821 3, mRNA sequence	5.07	Down	1.28E-02
35	BX647256	Homo sapiens mRNA; cDNA DKFZp686K0753 (from clone DKFZp686K0753)	5.07	Down	2.05E-02
	BG186566	RST5534 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	5.05	Down	8.91E-04
40	AV652758	AV652758 GLC Homo sapiens cDNA clone GLCDDG05 3, mRNA sequence	5.04	Down	1.05E-03
	AI672441	wa03c03.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2296996 3, mRNA sequence	5.04	Down	1.11E-02
	AA769642	ob20h04.s1 NCI_CGAP_Kid5 Homo sapiens cDNA clone IMAGE:1324279 3, mRNA sequence	5.02	Down	1.59E-02
45	AL706653	DKFZp686E1543_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686E1543 5, mRNA sequence	5.02	Down	1.09E-03
50	AW296834	UI-H-B12-ahz-a-10-0-UI.s1 NCI_CGAP_Sub4 Homo sapiens cDNA clone IMAGE:2728243 3, mRNA sequence	5.01	Down	5.93E-03

55

EP 1 780 267 B1

TABLE VII B: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN EPITHELIAL VERSUS FIBROBLAST CELLS

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p-value
5					
10	NM_018658	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 16 (KCNJ16), transcript variant 1, mRNA	481.28	UP	2.64E-05
	NM_014358	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	375.12	UP	6.56E-05
15	AB032980	Homo sapiens mRNA for KIAA1154 protein, partial cds	331.92	UP	1.60E-05
	BG219729	RST39494 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	316.78	UP	8.39E-06
	NM_002899	Homo sapiens retinol binding protein 1, cellular (RBP1), mRNA	314.6	UP	5.37E-05
20	NM_007038	Homo sapiens a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2) (ADAMTS5), mRNA	280.99	UP	1.26E-05
	NM_002423	Homo sapiens matrix metalloproteinase 7 (matrilysin, uterine) (MMP7), mRNA	269.82	UP	8.65E-06
25	NM_001977	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	242.27	UP	5.43E-05
	D29453	HUMNK566 Human epidermal keratinocyte Homo sapiens cDNA clone 566, mRNA sequence	208.36	UP	1.72E-05
30	NM_024508	Homo sapiens zinc finger, BED domain containing 2 (ZBED2), mRNA	168.99	UP	8.39E-06
	A1335277	tb29h06.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2055803 3, mRNA sequence	165.99	UP	8.46E-06
35	BE877764	601486331 F1 NIH_MGC_69 Homo sapiens cDNA clone IMAGE:3888943 5, mRNA sequence	145.52	UP	2.36E-05
	A1765021	wh56c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384738 3, mRNA sequence	145.09	UP	1.15E-05
	NM_024726	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	143.09	UP	8.39E-06
40	NM_004221	Homo sapiens natural killer cell transcript 4 (NK4), mRNA	142.15	UP	1.88E-05
	NM_019000	Homo sapiens hypothetical protein FLJ20152 (FLJ20152), mRNA	131.23	UP	1.88E-05
45	NM_000104	Homo sapiens cytochrome P450, family 1, subfamily B, polypeptide 1 (CYP1B1), mRNA	127.7	UP	1.51E-05
	BM993116	UI-H-DT0-aty-f-17-0-UI.s1 NCI_CGAP_DT0 Homo sapiens cDNA clone IMAGE:5866000 3, mRNA sequence	127.35	UP	4.13E-05
50	NM_031426	Homo sapiens chromosome 9 open reading frame 58 (C9orf58), transcript variant 1, mRNA	116.09	UP	3.82E-05
	NM_152369	Homo sapiens hypothetical protein MGC45474 (MGC45474), mRNA	112.28	UP	2.18E-05
55	NM_003810	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	112.19	UP	2.96E-05
	NM_023942	Homo sapiens hypothetical protein MGC3036 (MGC3036), mRNA	111.27	UP	1.15E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_005560	Homo sapiens laminin, alpha 5 (LAMA5), mRNA	110.51	UP	1.37E-05
10	NM_001453	Homo sapiens forkhead box C1 (FOXC1), mRNA	107.21	UP	1.15E-05
	BU734212	UI-E-CQ1-agd-e-21-0-UI.s1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-agd-e-21-0-UI 3, mRNA sequence	103.23	UP	2.89E-05
	AK026966	Homo sapiens cDNA: FLJ23313 fis, clone HEP11919	103.17	UP	1.78E-05
15	CA425961	UI-H-FE1-beg-p-18-0-UI.s1 NCI_CGAP_FE1 Homo sapiens cDNA clone UI-H-FE1-beg-p-18-0-UI 3, mRNA sequence	102.53	UP	6.21E-05
	AK075003	Homo sapiens cDNA FLJ90522 fis, clone NT2RP4000108, highly similar to Human gene for neurofilament subunit NF-L	101.8	UP	2.31E-04
20					
	NM_018265	Homo sapiens hypothetical protein FLJ10901 (FLJ10901), mRNA	99.41	UP	3.66E-05
	AK058012	Homo sapiens cDNA FLJ25283 fis, clone STM06716	98.88	UP	9.08E-06
25	NM_004098	Homo sapiens empty spiracles homolog 2 (Drosophila) (EMX2), mRNA	97.96	UP	4.13E-05
	AK096481	Homo sapiens cDNA FLJ39162 fis, clone OCBBF2002376	94.93	UP	2.83E-05
	NM_003238	Homo sapiens transforming growth factor, beta 2 (TGFB2), mRNA	91.63	UP	2.31E-05
30					
	AI244954	qj93h05.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1867065 3, mRNA sequence	85.24	UP	1.78E-05
	NM_173505	Homo sapiens ankyrin repeat domain 29 (ANKRD29), mRNA	82.89	UP	3.53E-05
35					
	AK130281	Homo sapiens cDNA FLJ26771 fis, clone PRS03189	82.02	UP	3.66E-06
	NM_153026	Homo sapiens prickly-like 1 (Drosophila) (PRICKLE1), mRNA	81.73	UP	1.78E-05
	AK092401	Homo sapiens cDNA FLJ35082 fis, clone PLACE6005351	79.69	UP	1.70E-05
40					
	AB011539	Homo sapiens mRNA for MEGF6 protein (KIAA0815), partial cds	79.31	UP	8.41E-05
	NM_016356	Homo sapiens doublecortin domain containing 2 (DCDC2), mRNA	78.04	UP	6.98E-05
45					
	AK023631	Homo sapiens cDNA FLJ13569 fis, clone PLACE1008369	77.68	UP	2.09E-05
	NM_024422	Homo sapiens desmocollin 2 (DSC2), transcript variant Dsc2a, mRNA	73.14	UP	7.38E-05
	BX648964	Homo sapiens mRNA; cDNA DKFZp686J0156 (from clone DKFZp686J0156)	70.68	UP	6.92E-05
50					
	T56535	yb33g07.r1 Stratagene fetal spleen (#937205) Homo sapiens cDNA clone IMAGE:73020 5, mRNA sequence	69.74	UP	2.03E-05
	BG197054	RST16291 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	69.48	UP	4.96E-05
55					
	NM_005949	Homo sapiens metallothionein 1 F (functional) (MT1 F), mRNA	68.1	UP	2.03E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AK090808	Homo sapiens cDNA FLJ33489 fis, clone BRAMY2003585	65.29	UP	2.87E-05
10	CA306881	UI-H-FT1-bht-n-22-0-UI.s1 NCI_CGAP_FT1 Homo sapiens cDNA clone UI-H-FT1-bht-n-22-0-UI 3, mRNA sequence	64.98	UP	2.93E-05
	NM_001448	Homo sapiens glypican 4 (GPC4), mRNA	64.85	UP	2.45E-04
15	AL833276	Homo sapiens mRNA; cDNA DKFZp451D088 (from clone DKFZp451D088)	64.7	UP	1.98E-05
	BC044843	Homo sapiens hypothetical protein LOC339535, mRNA (cDNA clone IMAGE:5186761), partial cds	63.8	UP	7.59E-05
20	BF798098	RC1-C10045-021000-021-f02 C10045 Homo sapiens cDNA, mRNA sequence	58.58	UP	3.04E-05
	H89526	yw28b04.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253519 5, mRNA sequence	56.92	UP	3.82E-05
	BF509573	UI-H-B14-apf-b-11-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086949 3, mRNA sequence	56.79	UP	2.83E-05
25					
	NM_000990	Homo sapiens ribosomal protein L27a (RPL27A), mRNA	56.36	UP	1.73E-05
	BC042028	Homo sapiens, clone IMAGE:4794726, mRNA	55.75	UP	1.64E-05
	NM_003287	Homo sapiens tumor protein D52-like 1 (TPD52L1), transcript variant 1, mRNA	55.25	UP	8.46E-06
30					
	NM_018168	Homo sapiens chromosome 14 open reading frame 105 (C14orf105), mRNA	54.64	UP	4.79E-05
	NM_152284	Homo sapiens Snf7 homologue associated with Alix 3 (Shax3), mRNA	54.16	UP	9.09E-06
35					
	AK096975	Homo sapiens cDNA FLJ39656 fis, clone SMINT2005956	54.14	UP	8.37E-05
	AI249696	qj64a03.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1864204 3, mRNA sequence	53.24	UP	7.14E-05
40					
	NM_003551	Homo sapiens non-metastatic cells 5, protein expressed in (nucleoside-diphosphate kinase) (NME5), mRNA	53.01	UP	4.75E-05
	BX118238	BX118238 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998L153800 ; IMAGE:1501598, mRNA sequence	51.99	UP	5.64E-05
45					
	BX102632	BX102632 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998J052307 ; IMAGE:928228, mRNA sequence	51.79	UP	9.10E-05
	AF055376	Homo sapiens short form transcription factor C-MAF (c- maf) mRNA, complete cds	51.22	UP	3.94E-05
	NM_012464	Homo sapiens tolloid-like 1 (TLL1), mRNA	51.19	UP	6.57E-04
50					
	NM_030949	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 14C (PPP1 R14C), mRNA	49.96	UP	5.80E-05
	BM669002	UI-E-CK1-afn-m-04-0-UI.s2 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-afn-m-04-0-UI 3, mRNA sequence	49.05	UP	1.78E-05
55					
	NM_000582	Homo sapiens secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA	46.79	UP	6.18E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	46.73	UP	1.26E-05
10	NM_152423	Homo sapiens melanoma associated antigen (mutated) 1-like 1 (MUM1L1), mRNA	46.24	UP	1.48E-04
	AA075748	zm89e04.r1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone IMAGE:545118 5, mRNA sequence	45.76	UP	4.39E-05
15	NM_017549	Homo sapiens upregulated in colorectal cancer gene 1 (UCC1), mRNA	45.18	UP	1.24E-05
	BF431030	7c18c06.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3574283 3, mRNA sequence	44.98	UP	7.75E-05
20	NM_002245	Homo sapiens potassium channel, subfamily K, member 1 (KCNK1), mRNA	44.3	UP	4.65E-05
	NM_003494	Homo sapiens dysferlin, limb girdle muscular dystrophy 2B (autosomal recessive) (DYSF), mRNA	44.23	UP	1.78E-05
25	AK074097	Homo sapiens mRNA for FLJ00168 protein	43.88	UP	7.31E-05
	NM_013410	Homo sapiens adenylate kinase 3 (AK3), nuclear gene encoding mitochondrial protein, transcript variant 2, mRNA	42.34	UP	2.96E-05
	AK000075	Homo sapiens cDNA FLJ20068 fis, clone COL01755	42.18	UP	1.35E-04
30	NM_001200	Homo sapiens bone morphogenetic protein 2 (BMP2), mRNA	41.13	UP	2.11E-05
	NM_032782	Homo sapiens hepatitis A virus cellular receptor 2 (HAVCR2), mRNA	41.13	UP	2.83E-05
	AW151660	xf67d04.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2623111 3, mRNA sequence	40.6	UP	8.65E-05
35	NM_130435	Homo sapiens protein tyrosine phosphatase, receptor type, E (PTPRE), transcript variant 2, mRNA	40.41	UP	1.62E-04
	AA738254	nx13b02.s1 NCI_CGAP_GC3 Homo sapiens cDNA clone IMAGE:1255947 3, mRNA sequence	40.24	UP	2.43E-05
40	AK124873	Homo sapiens cDNA FLJ42883 fis, clone BRHIP3006683	40.2	UP	1.25E-04
	NM_000170	Homo sapiens glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P) (GLDC), mRNA	40.06	UP	4.98E-05
45	AK125490	Homo sapiens cDNA FLJ43501 fis, clone PEBLM2004497	39.93	UP	1.64E-04
	NM_004862	Homo sapiens lipopolysaccharide-induced TNF factor (LITAF), mRNA	39.52	UP	1.37E-05
50	NM_004617	Homo sapiens transmembrane 4 superfamily member 4 (TM4SF4), mRNA	39.3	UP	9.32E-05
	NM_001263	Homo sapiens CDP-diacylglycerol synthase (phosphatidate cytidyltransferase) 1 (CDS1), mRNA	39.26	UP	3.44E-05
	AK000776	Homo sapiens cDNA FLJ20769 fis, clone COL06674	38.12	UP	2.83E-05
55	AL833166	Homo sapiens mRNA; cDNA DKFZp68612118 (from clone DKFZp68612118)	37.72	UP	1.14E-04
	NM_198488	Homo sapiens FLJ46072 protein (FLJ46072), mRNA	37.66	UP	2.46E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_005562	Homo sapiens laminin, gamma 2 (LAMC2), transcript variant 1, mRNA	37.4	UP	8.46E-06
10	AW268540	xv51e10.x1 NCI_GAP_L28 Homo sapiens cDNA clone IMAGE:2816682 3, mRNA sequence	37.26	UP	5.82E-05
	NM_012198	Homo sapiens grancalcin, EF-hand calcium binding protein (GCA), mRNA	36.52	UP	2.13E-05
15	NM_020808	Homo sapiens signal-induced proliferation-associated 1 like 2 (SIPA1L2), mRNA	36.28	UP	9.39E-05
	NM_000927	Homo sapiens ATP-binding cassette, sub-family B (MDR/TAP), member 1 (ABCB1), mRNA	35.89	UP	2.20E-04
20	AK127644	Homo sapiens cDNA FLJ45742 fis, clone KIDNE2016327	35.81	UP	3.60E-04
	NM_006158	Homo sapiens neurofilament, light polypeptide 68kDa (NEFL), mRNA	35.78	UP	2.64E-05
	AK074181	Homo sapiens mRNA for FLJ00254 protein	35.75	UP	8.46E-06
25	BC043195	Homo sapiens cDNA clone IMAGE:5288757, partial cds	35.12	UP	1.24E-04
	AL389942	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 2005635	34.31	UP	1.37E-04
	BC045828	Homo sapiens zinc finger protein 608, mRNA (cDNA clone IMAGE:5262896), partial cds	34.02	UP	1.88E-05
30	NM_153229	Homo sapiens hypothetical protein FLJ33318 (FLJ33318), mRNA	34.02	UP	9.39E-05
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	33.68	UP	2.18E-05
	NM_000216	Homo sapiens Kallmann syndrome 1 sequence (KAL1), mRNA	33.32	UP	1.78E-05
35	BF696790	602125323F1 NIH_MGC_56 Homo sapiens cDNA clone IMAGE:4282540 5, mRNA sequence	32.7	UP	5.00E-05
	NM_006598	Homo sapiens solute carrier family 12 (potassium/chloride transporters), member 7 (SLC12A7), mRNA	32.41	UP	8.39E-06
40	BQ003401	UI-H-EI1-azd-j-23-0-UIs1 NCI_CGAP_E11 Homo sapiens cDNA clone IMAGE:5847286 3, mRNA sequence	32.41	UP	1.72E-05
45	NM_003222	Homo sapiens transcription factor AP-2 gamma (activating enhancer binding protein 2 gamma) (TFAP2C), mRNA	32.33	UP	4.75E-05
	NM_000599	Homo sapiens insulin-like growth factor binding protein 5 (IGFBP5), mRNA	32.1	UP	5.66E-05
50	NM_198389	Homo sapiens lung type-I cell membrane-associated glycoprotein (T1A-2), transcript variant 2, mRNA	31.44	UP	1.03E-04
	NM_180991	Homo sapiens solute carrier organic anion transporter family, member 4C1 (SLCO4C1), mRNA	30.86	UP	3.05E-04
	NM_031311	Homo sapiens carboxypeptidase, vitellogenic-like (CPVL), transcript variant 1, mRNA	29.97	UP	4.30E-05
55	NM_052947	Homo sapiens heart alpha-kinase (HAK), mRNA	29.53	UP	4.34E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	H89053	yw24c06.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253162 5, mRNA sequence	29.27	UP	2.78E-05
10	AI819186	wj32d10.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2404531 3, mRNA sequence	29.14	UP	1.21E-04
	NM_003761	Homo sapiens vesicle-associated membrane protein 8 (endobrevin) (VAMP8), mRNA	28.5	UP	7.47E-05
15	NM_080743	Homo sapiens serine-arginine repressor protein (35 kDa) (SRp35), mRNA	27.83	UP	1.02E-05
	NM_007069	Homo sapiens HRAS-like suppressor 3 (HRASLS3), mRNA	27.51	UP	6.86E-06
	NM_002345	Homo sapiens lumican (LUM), mRNA	27.17	UP	2.20E-05
20	BU569937	AGENCOURT_10399817 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6618011 5, mRNA sequence	27.05	UP	3.20E-05
	NM_024901	Homo sapiens hypothetical protein FLJ22457 (FLJ22457), mRNA	27.04	UP	5.37E-05
25	N25875	yw78d12.s1 Soares_placenta_8to9weeks_2NbHP8to9W Homo sapiens cDNA clone IMAGE:258359 3, mRNA sequence	26.94	UP	3.36E-05
	NM_152573	Homo sapiens RAS and EF hand domain containing (RASEF), mRNA	26.91	UP	4.75E-05
30	NM_018728	Homo sapiens myosin VC (MYO5C), mRNA	26.81	UP	4.47E-05
	AB033048	Homo sapiens mRNA for KIAA1222 protein, partial cds	26.76	UP	1.90E-04
	NM_153256	Homo sapiens chromosome 10 open reading frame 47 (C10orf47), mRNA	26.61	UP	2.48E-05
	BG221364	RST41175 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	26.25	UP	2.06E-04
35	NM_000094	Homo sapiens collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive) (COL7A1), mRNA	26.16	UP	6.51E-05
	NM_001854	Homo sapiens collagen, type XI, alpha 1 (COL11A1), transcript variant A, mRNA	25.88	UP	1.36E-04
40	BU740051	UI-E-EO0-ahw-n-18-0-UI.s1 UI-E-EO0 Homo sapiens cDNA clone UI-E-EO0-ahw-n-18-0-UI 3, mRNA sequence	25.64	UP	5.14E-05
	NM_014936	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 4 (putative function) (ENPP4), mRNA	25.62	UP	2.82E-04
45	NM_024898	Homo sapiens family with sequence similarity 31, member C (FAM31C), mRNA	25.44	UP	7.17E-04
	W93585	zd95g01.s1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:357264 3, mRNA sequence	25.41	UP	1.78E-05
50	BF510493	UI-H-B14-apa-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086558 3, mRNA sequence	25.28	UP	2.11E-05
	NM_020962	Homo sapiens likely ortholog of mouse neighbor of Punc E11 (NOPE), mRNA	25.23	UP	7.99E-05
55	NM_016212	Homo sapiens TP53TG3 protein (TP53TG3), mRNA	25.14	UP	4.07E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_021012	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 12 (KCNJ12), mRNA	24.7	UP	2.10E-04
10	NM_001935	Homo sapiens dipeptidylpeptidase 4 (CD26, adenosine deaminase complexing protein 2) (DPP4), mRNA	24.67	UP	8.46E-06
	NM_001993	Homo sapiens coagulation factor III (thromboplastin, tissue factor) (F3), mRNA	24.47	UP	6.12E-05
15	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	24.38	UP	8.46E-06
	AK056431	Homo sapiens cDNA FLJ31869 fis, clone NT2RP7002151	24.34	UP	2.73E-05
20	NM_000775	Homo sapiens cytochrome P450, family 2, subfamily J, polypeptide 2 (CYP2J2), mRNA	24.04	UP	3.53E-05
	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	23.95	UP	5.23E-06
	NM_001977	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	23.91	UP	1.88E-05
25	AK058012	Homo sapiens cDNA FLJ25283 fis, clone STM06716	23.55	UP	1.88E-05
	BU680661	UI-CF-DU1-aaz-f-04-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaz-f-04-0-UI 3, mRNA sequence	23.17	UP	1.82E-05
30	NM_002246	Homo sapiens potassium channel, subfamily K, member 3 (KCNK3), mRNA	22.96	UP	2.27E-05
	NM_052923	Homo sapiens zinc finger protein 452 (ZNF452), mRNA	22.72	UP	3.77E-05
	NM_014373	Homo sapiens G protein-coupled receptor 160 (GPR160), mRNA	22.71	UP	3.94E-05
35	BX538226	Homo sapiens mRNA; cDNA DKFZp686E1944 (from clone DKFZp686E1944)	22.58	UP	1.05E-04
	NM_173660	Homo sapiens hypothetical protein FLJ33718 (FLJ33718), mRNA	22.46	UP	2.83E-05
40	NM_002354	Homo sapiens tumor-associated calcium signal transducer 1 (TACSTD1), mRNA	21.97	UP	2.83E-05
	NM_005822	Homo sapiens Down syndrome critical region gene 1-like 1 (DSCR1L1), mRNA	21.78	UP	2.21E-04
	W20132	zb40c10.r1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:306066 5, mRNA sequence	21.76	UP	4.41E-04
45	NM_001680	Homo sapiens FXD domain containing ion transport regulator 2 (FXD2), transcript variant a, mRNA	21.75	UP	3.26E-05
	NM_006379	Homo sapiens sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3C (SEMA3C), mRNA	21.57	UP	3.15E-05
50	H23441	ym52f11.s1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:51888 3, mRNA sequence	21.43	UP	3.51E-05
	BG211832	RST31404 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	21.35	UP	5.06E-05
55	N51335	yz15e08.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:283142 3, mRNA sequence	21.19	UP	6.92E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_022843	Homo sapiens protocadherin 20 (PCDH20), mRNA	21.17	UP	2.03E-05
10	NM_002521	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	21.01	UP	4.54E-04
	NM_021101	Homo sapiens claudin 1 (CLDN1), mRNA	20.92	UP	3.04E-05
	NM_004165	Homo sapiens Ras-related associated with diabetes (RRAD), mRNA	20.76	UP	6.04E-04
15	NM_005329	Homo sapiens hyaluronan synthase 3 (HAS3), transcript variant 1, mRNA	20.73	UP	4.13E-05
	NM_138432	Homo sapiens serine dehydratase-like (SDSL), mRNA	20.41	UP	7.27E-05
	NM_004591	Homo sapiens chemokine (C-C motif) ligand 20 (CCL20), mRNA	20.34	UP	3.29E-04
20	NM_003063	Homo sapiens sarcolipin (SLN), mRNA	20.26	UP	1.18E-04
	BU951469	in60a05.x3 HR85 islet Homo sapiens cDNA clone IMAGE:6126249 3, mRNA sequence	20.02	UP	4.52E-04
	AI436290	th81c01.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2125056 3, mRNA sequence	19.92	UP	9.28E-05
25	NM_014452	Homo sapiens tumor necrosis factor receptor superfamily, member 21 (TNFRSF21), mRNA	19.39	UP	2.96E-05
	AA099748	zl78c09.s1 Stratagene colon (#937204) Homo sapiens cDNA clone IMAGE:510736 3, mRNA sequence	19.2	UP	1.78E-05
30	NM_018354	Homo sapiens chromosome 20 open reading frame 46 (C20orf46), mRNA	18.87	UP	5.05E-04
	NM_194298	Homo sapiens solute carrier family 16 (monocarboxylic acid transporters), member 9 (SLC16A9), mRNA	18.87	UP	9.82E-03
	NM_033554	Homo sapiens major histocompatibility complex, class II, DP alpha 1 (HLA-DPA1), mRNA	18.6	UP	5.61E-04
35	NM_004524	Homo sapiens lethal giant larvae homolog 2 (Drosophila) (LLGL2), mRNA	18.45	UP	2.11E-05
	NM_032293	Homo sapiens GTPase activating Rap/RanGAP domain-like 3 (GARNL3), mRNA	18.19	UP	2.24E-04
40	NM_022842	Homo sapiens CUB domain-containing protein 1 (CDCP1), transcript variant 1, mRNA	18.14	UP	1.64E-05
	NM_000227	Homo sapiens laminin, alpha 3 (LAMA3), transcript variant 2, mRNA	18.04	UP	3.92E-05
45	NM_000214	Homo sapiens jagged 1 (Alagille syndrome) (JAG1), mRNA	18.03	UP	1.26E-04
	AK096580	Homo sapiens cDNA FLJ39261 fis, clone OCBBF2009391	17.89	UP	3.05E-04
	NM_014243	Homo sapiens a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 3 (ADAMTS3), mRNA	17.86	UP	1.78E-05
50	NM_000856	Homo sapiens guanylate cyclase 1, soluble, alpha 3 (GUCY1A3), mRNA	17.85	UP	1.19E-04
	NM_018404	Homo sapiens centaurin, alpha 2 (CENTA2), mRNA	17.8	UP	3.05E-04
55	NM_001165	Homo sapiens baculoviral IAP repeat-containing 3 (BIRC3), transcript variant 1, mRNA	17.74	UP	2.56E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_003706	Homo sapiens phospholipase A2, group IVC (cytosolic, calcium-independent) (PLA2G4C), mRNA	17.65	UP	4.77E-05
10	AK056882	Homo sapiens cDNA FLJ32320 fis, clone PROST2003537	17.65	UP	4.33E-04
	NM_017640	Homo sapiens leucine rich repeat containing 16 (LRR16), mRNA	17.43	UP	1.04E-04
15	NM_000480	Homo sapiens adenosine monophosphate deaminase (isoform E) (AMPD3), mRNA	17.4	UP	1.15E-05
	BX103476	BX103476 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998C053946 ; IMAGE:1557436, mRNA sequence	17.29	UP	7.13E-05
20	NM_013427	Homo sapiens Rho GTPase activating protein 6 (ARHGAP6), transcript variant 1, mRNA	17.27	UP	1.48E-04
	S70348	Homo sapiens integrin beta 3 mRNA, partial cds, alternatively spliced	17.24	UP	3.37E-05
25	NM_032024	Homo sapiens chromosome 10 open reading frame 11 (C10orf11), mRNA	16.95	UP	3.03E-05
	BM988642	UI-H-DH0-arx-p-21-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone IMAGE:5855492 3, mRNA sequence	16.94	UP	1.94E-04
30	NM_021102	Homo sapiens serine protease inhibitor, Kunitz type, 2 (SPINT2), mRNA	16.94	UP	1.85E-04
	AI632692	wa33b05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299857 3, mRNA sequence	16.88	UP	7.19E-04
	M_152487	Homo sapiens hypothetical protein FLJ31842 (FLJ31842), mRNA	16.75	UP	2.55E-04
35	NM_173567	Homo sapiens abhydrolase domain containing 7 (ABHD7), mRNA	16.66	UP	2.13E-05
	CA502927	UI-CF-FN0-afq-j-12-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afq-j-12-0-UI 3, mRNA sequence	16.59	UP	2.35E-04
40	NM_015888	Homo sapiens hook homolog 1 (Drosophila) (HOOK1), mRNA	16.44	UP	4.12E-05
	NM_178177	Homo sapiens nicotinamide nucleotide adenylyltransferase 3 (NMNAT3), mRNA	16.43	UP	1.56E-04
45	NM_005181	Homo sapiens carbonic anhydrase III, muscle specific (CA3), mRNA	16.42	UP	4.93E-05
	NM_000599	Homo sapiens insulin-like growth factor binding protein 5 (IGFBP5), mRNA	16.27	UP	4.97E-04
50	NM_006393	Homo sapiens nebulin (NEBL), transcript variant 1, mRNA	16.13	UP	6.06E-04
	NM_006252	Homo sapiens protein kinase, AMP-activated, alpha 2 catalytic subunit (PRKAA2), mRNA	16.12	UP	5.64E-04
	NM_002338	Homo sapiens limbic system-associated membrane protein (LSAMP), mRNA	16.11	UP	2.81E-04
55	NM_006378	Homo sapiens sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D (SEMA4D), mRNA	16.09	UP	1.78E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AI819863	wj45h05.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2405817 3, mRNA sequence	15.95	UP	3.04E-04
10	BM979825	UI-CF-DU1-adt-f-12-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-adt-f-12-0-UI 3, mRNA sequence	15.9	UP	2.77E-05
	BX647876	Homo sapiens mRNA; cDNA DKFZp313A1525 (from clone DKFZp313A1525)	15.83	UP	4.39E-05
15	NM_003730	Homo sapiens ribonuclease T2 (RNASET2), mRNA	15.6	UP	2.41E-05
	AL117425	Homo sapiens mRNA; cDNA DKFZp566L203 (from clone DKFZp566L203)	15.59	UP	3.31E-04
	AL833346	Homo sapiens mRNA; cDNA DKFZp686M2234 (from clone DKFZp686M2234)	15.59	UP	2.99E-05
20	NM_199169	Homo sapiens transmembrane, prostate androgen induced RNA (TMEPAI), transcript variant 2, mRNA	15.58	UP	1.34E-04
	NM_002559	Homo sapiens purinergic receptor P2X, ligand-gated ion channel, 3 (P2RX3), mRNA	15.51	UP	9.39E-05
25	NM_032148	Homo sapiens solute carrier family 41, member 2 (SLC41A2), mRNA	15.46	UP	2.38E-05
	BC021684	Homo sapiens, clone IMAGE:3827252, mRNA	15.41	UP	2.31E-05
	BF000009	7h15g04.x1 NCI_CGAP_Co16 Homo sapiens cDNA clone IMAGE:3316086 3, mRNA sequence	15.33	UP	2.09E-05
30	NM_000147	Homo sapiens fucosidase, alpha-L-1, tissue (FUCA1), mRNA	15.31	UP	9.99E-04
	NM_001252	Homo sapiens tumor necrosis factor (ligand) superfamily, member 7 (TNFSF7), mRNA	14.88	UP	1.85E-04
35	NM_013322	Homo sapiens sorting nexin 10 (SNX10), mRNA	14.84	UP	2.64E-05
	NM_022440	Homo sapiens mal, T-cell differentiation protein (MAL), transcript variant d, mRNA	14.65	UP	2.14E-04
	NM_178470	Homo sapiens WD repeat domain 40B (WDR40B), mRNA	14.53	UP	3.34E-05
40	NM_001873	Homo sapiens carboxypeptidase E (CPE), mRNA	14.51	UP	6.23E-05
	BM719937	UI-E-EJ0-ahu-a-10-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahu-a-10-0-UI 5, mRNA sequence	14.46	UP	1.64E-05
45	NM_032551	Homo sapiens G protein-coupled receptor 54 (GPR54), mRNA	14.45	UP	5.64E-04
	T53523	ya89h12.r1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:68903 5, mRNA sequence	14.42	UP	1.81E-04
	NM_173354	Homo sapiens SNF1-like kinase (SNF1LK), mRNA	14.19	UP	5.72E-05
50	NM_014585	Homo sapiens solute carrier family 40 (iron-regulated transporter), member 1 (SLC40A1), mRNA	14.17	UP	4.06E-05
	N38753	yy42d01.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273889 3, mRNA sequence	14.02	UP	1.15E-05
	NM_001710	Homo sapiens B-factor, properdin (BF), mRNA	13.95	UP	2.05E-04
55	BC038556	Homo sapiens, clone IMAGE:3446976, mRNA	13.9	UP	4.45E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
10	AK125608	Homo sapiens cDNA FLJ43620 fis, clone SPLEN2021701, highly similar to HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN PRECURSOR	13.88	UP	2.56E-04
	AW248516	2820632.3prime NIH_MGC_7 Homo sapiens cDNA clone IMAGE:2820632 3, mRNA sequence	13.79	UP	2.22E-04
15	NM_000600	Homo sapiens interleukin 6 (interferon, beta 2) (IL6), mRNA	13.62	UP	1.64E-05
	AK125852	Homo sapiens cDNA FLJ43864 fis, clone TESTI4007799	13.46	UP	1.66E-04
20	BM728728	UI-E-EO1-aiv-c-02-0-UI.r1 UI-E-EO1 Homo sapiens cDNA clone UI-E-EO1-aiv-c-02-0-UI 5, mRNA sequence	13.46	UP	2.77E-04
	NM_018650	Homo sapiens MAP/microtubule affinity-regulating kinase 1 (MARK1), mRNA	13.45	UP	4.39E-05
25	NM_022154	Homo sapiens solute carrier family 39 (zinc transporter), member 8 (SLC39A8), mRNA	13.39	UP	4.78E-05
	AA908815	og77h08.s1 NCI_CGAP_Ov8 Homo sapiens cDNA clone IMAGE:1454367 3, mRNA sequence	13.23	UP	3.11E-05
	NM_016463	Homo sapiens CXXC finger 5 (CXXC5), mRNA	13.22	UP	9.67E-05
30	AB011538	Homo sapiens mRNA for MEGF5, partial cds	13.19	UP	3.38E-04
	CA413744	UI-H-EZ0-bat-h-12-0-UI.s1 NCI_CGAP_Ch1 Homo sapiens cDNA clone UI-H-EZ0-bat-h-12-0-UI 3, mRNA sequence	13.17	UP	2.09E-05
	BM712072	UI-E-DW1-ahc-b-11-0-UI.r1 UI-E-DW1 Homo sapiens cDNA clone UI-E-DW1-ahc-b-11-0-UI 5, mRNA sequence	13.16	UP	9.39E-05
35	BU727096	UI-E-CR0-ach-e-12-0-UI.s1 UI-E-CR0 Homo sapiens cDNA clone UI-E-CR0-ach-e-12-0-UI 3, mRNA sequence	13.14	UP	4.30E-05
40	NM_032858	Homo sapiens hypothetical protein FLJ14904 (FLJ14904), mRNA	13.09	UP	1.83E-05
	BU729783	UI-E-CK1-afh-h-18-0-UI.s1 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-afh-h-18-0-UI 3, mRNA sequence	12.92	UP	8.39E-06
45	AB011095	Homo sapiens mRNA for KIAA0523 protein, partial cds	12.91	UP	3.44E-04
	BC033116	Homo sapiens chromodomain helicase DNA binding protein 7, mRNA (cDNA clone IMAGE:3352674), partial cds	12.87	UP	2.31E-05
50	CB135276	K-EST0187371 L5HLK1 Homo sapiens cDNA clone L5HLK1-32-B12 5, mRNA sequence	12.75	UP	4.93E-05
	NM_182920	Homo sapiens a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 9 (ADAMTS9), transcript variant 1, mRNA	12.73	UP	9.10E-05
55	NM_000236	Homo sapiens lipase, hepatic (LIPC), mRNA	12.71	UP	1.16E-03
	NM_152694	Homo sapiens zinc finger, CCHC domain containing 5 (ZCCHC5), mRNA	12.66	UP	4.93E-05
	AB020640	Homo sapiens mRNA for KIAA0833 protein, partial cds	12.51	UP	2.31E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_018330	Homo sapiens KIAA1598 (KIAA1598), mRNA	12.46	UP	6.01E-05
10	NM_000064	Homo sapiens complement component 3 (C3), mRNA	12.42	UP	5.12E-04
	NM_152864	Homo sapiens chromosome 20 open reading frame 58 (C20orf58), mRNA	12.4	UP	1.66E-03
	AW591723	xt85h10.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2793283 3 similar to contains element MER32 repetitive element ;, mRNA sequence	12.4	UP	3.71E-05
15					
	NM_152768	Homo sapiens hypothetical protein FLJ25378 (FLJ25378), mRNA	12.28	UP	2.03E-05
	NM_152366	Homo sapiens kelch/ankyrin repeat containing cyclin A1 interacting protein (KARCA1), transcript variant 1, mRNA	12.19	UP	2.62E-05
20					
	AW300043	xs45a09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2772568 3, mRNA sequence	12.13	UP	2.31E-05
	NM_006741	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 1A (PPP1R1A), mRNA	12.12	UP	2.10E-04
25					
	NM_024726	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	12.08	UP	5.33E-03
	NM_194284	Homo sapiens claudin 23 (CLDN23), mRNA	12.02	UP	2.87E-05
	AL049974	Homo sapiens mRNA; cDNA DKFZp564B222 (from clone DKFZp564B222)	12	UP	1.48E-04
30					
	AI420213	te92g09.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2094208 3, mRNA sequence	11.95	UP	3.59E-04
	NM_002800	Homo sapiens proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 2) (PSMB9), transcript variant 1, mRNA	11.85	UP	4.69E-05
35					
	CA391258	cs13a10.x1 Human Retinal pigment epithelium/choroid cDNA (Un-normalized, unamplified): cs Homo sapiens cDNA clone cs13a10 3, mRNA sequence	11.85	UP	3.64E-05
	NM_013430	Homo sapiens gamma-glutamyltransferase 1 (GGT1), transcript variant 3, mRNA	11.79	UP	1.38E-04
40					
	NM_199169	Homo sapiens transmembrane, prostate androgen induced RNA (TMEPAI), transcript variant 2, mRNA	11.79	UP	8.39E-06
	NM_032471	Homo sapiens protein kinase (cAMP-dependent, catalytic) inhibitor beta (PKIB), transcript variant 3, mRNA	11.73	UP	1.28E-04
45					
	NM_016269	Homo sapiens lymphoid enhancer-binding factor 1 (LEF1), mRNA	11.72	UP	4.57E-05
	NM_003851	Homo sapiens cellular repressor of E1A-stimulated genes 1 (CREG1), mRNA	11.61	UP	1.83E-05
50					
	NM_003236	Homo sapiens transforming growth factor, alpha (TGFA), mRNA	11.61	UP	6.51E-05
	BX110418	BX110418 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998C224149 ; IMAGE:1635405, mRNA sequence	11.59	UP	1.89E-05
55					
	NM_006033	Homo sapiens lipase, endothelial (LIPG), mRNA	11.55	UP	8.79E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_025151	Homo sapiens RAB11 family interacting protein 1 (class I) (RAB11FIP1 transcript variant 1, mRNA	11.5	UP	2.65E-05
10	BI759570	603046987F1 NIH_MGC_116 Homo sapiens cDNA clone IMAGE:5187285 5, mRNA sequence	11.49	UP	3.71E-05
	NM_013261	Homo sapiens peroxisome proliferative activated receptor, gamma, coactivator 1, alpha (PPARGC1A), mRNA	11.3	UP	2.83E-04
15	AI830524	wh52c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384354 3, mRNA sequence	11.29	UP	8.65E-04
	NM_139161	Homo sapiens crumbs homolog 3 (Drosophila) (CRB3), transcript variant 2, mRNA	11.15	UP	1.22E-04
20	AK055362	Homo sapiens cDNA FLJ30800 fis, clone FEBRA2001197	11.13	UP	1.02E-05
	BF055156	7j75f03.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3392285 3, mRNA sequence	11.11	UP	4.94E-04
25	BC016962	Homo sapiens, clone IMAGE:4182947, mRNA	11.05	UP	1.06E-04
	AY358775	Homo sapiens clone DNA170212 WGAR9166 (UNQ9166) mRNA, complete cds	11.05	UP	2.32E-04
	NM_175056	Homo sapiens hypothetical protein LOC131368 (LOC131368), mRNA	10.98	UP	1.33E-03
30	AI311296	ta48d10.x2 NCI_CGAP_Lu25 Homo sapiens cDNA clone IMAGE:2047315 3, mRNA sequence	10.93	UP	2.19E-04
	AW296834	UI-H-BI2-ahz-a-10-0-UI.s1 NCI_CGAP_Sub4 Homo sapiens cDNA clone IMAGE:2728243 3, mRNA sequence	10.91	UP	1.71E-04
35	NM_014464	Homo sapiens tubulointerstitial nephritis antigen (TINAG), mRNA	10.89	UP	3.87E-04
	AK024270	Homo sapiens cDNA FLJ14208 fis, clone NT2RP3003264	10.88	UP	3.05E-04
40	H00617	yj25f02.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:149787 3, mRNA sequence	10.87	UP	2.09E-04
	AK126467	Homo sapiens cDNA FLJ44503 fis, clone UTERU3001158	10.87	UP	5.06E-05
45	NM_181847	Homo sapiens amphoterin induced gene 2 (AMIGO2), mRNA	10.86	UP	8.91E-04
	NM_014751	Homo sapiens metastasis suppressor 1 (MTSS1), mRNA	10.86	UP	2.68E-05
	N70752	za72d02.s1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:298083 3, mRNA sequence	10.84	UP	1.34E-04
50	NM_005302	Homo sapiens G protein-coupled receptor 37 (endothelin receptor type B-like) (GPR37), mRNA	10.84	UP	2.63E-03
	NM_003264	Homo sapiens toll-like receptor 2 (TLR2), mRNA	10.83	UP	1.01E-04
	NM_017594	Homo sapiens DIRAS family, GTP-binding RAS-like 2 (DIRAS2), mRNA	10.81	UP	7.23E-04
55	AI688800	wd41b03.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2330669 3, mRNA sequence	10.8	UP	8.56E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	H08012	yl91b08.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:45474 5, mRNA sequence	10.8	UP	3.26E-04
10	AW014126	UI-H-BIO-aaj-a-05-0-UI.s1 NCI_CGAP_Sub1 Homo sapiens cDNA clone IMAGE:2709393 3, mRNA sequence	10.79	UP	1.88E-05
	NM_014422	Homo sapiens phosphatidylinositol (4,5) bisphosphate 5-phosphatase, A (PIB5PA), transcript variant 1, mRNA	10.73	UP	4.47E-05
15	H62713	yr28c08.r1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:206606 5 similar to gb:X01683 ALPHA-1-ANTITRYPSIN PRECURSOR (HUMAN);, mRNA sequence	10.73	UP	1.01E-03
20	A1870547	wl47a04.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2428014 3, mRNA sequence	10.65	UP	4.82E-03
	NM_025202	Homo sapiens EF hand domain containing 1 (EFHD1), mRNA	10.64	UP	5.59E-05
25	N78460	yz76h06.r1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:289019 5, mRNA sequence	10.63	UP	3.90E-04
	NM_020661	Homo sapiens activation-induced cytidine deaminase (AICDA), mRNA	10.63	UP	7.27E-05
30	NM_173549	Homo sapiens hypothetical protein FLJ39553 (FLJ39553), mRNA	10.57	UP	3.11E-04
	A1767472	wh27a07.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2381940 3 similar to contains L1.t3 L1 repetitive element ;, mRNA sequence	10.56	UP	8.39E-06
35	NM_145804	Homo sapiens ankyrin repeat and BTB (POZ) domain containing 2 (ABTB2), mRNA	10.55	UP	1.41E-06
	NM_013230	Homo sapiens CD24 antigen (small cell lung carcinoma cluster 4 antigen) (CD24), mRNA	10.54	UP	1.12E-04
	NM_000758	Homo sapiens colony stimulating factor 2 (granulocyte-macrophage) (CSF2), mRNA	10.51	UP	7.66E-04
40	AB032945	Homo sapiens mRNA for KIAA1119 protein, partial cds	10.48	UP	1.32E-03
	NM_024572	Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetyl-galactosaminyltransferase 14 (GalNAc-T14) (GALNT14), mRNA	10.46	UP	2.62E-04
45					
	NM_181332	Homo sapiens neuroligin 4, X-linked (NLGN4X), transcript variant 2, mRNA	10.46	UP	1.95E-04
	NM_022128	Homo sapiens ribokinase (RBKS), mRNA	10.45	UP	2.73E-05
50	AB046810	Homo sapiens mRNA for KIAA1590 protein, partial cds	10.39	UP	2.36E-05
	NM_022168	Homo sapiens interferon induced with helicase C domain 1 (IFIH1), mRNA	10.38	UP	2.83E-05
	NM_153715	Homo sapiens homeo box A10 (HOXA10), transcript variant 2, mRNA	10.37	UP	5.50E-05
55	BF090392	QV3-NT0023-120900-324-b04 NT0023 Homo sapiens cDNA, mRNA sequence	10.35	UP	1.25E-03
	NM_000880	Homo sapiens interleukin 7 (IL7), mRNA	10.34	UP	3.60E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_018650	Homo sapiens MAP/microtubule affinity-regulating kinase 1 (MARK1), mRNA	10.29	UP	1.57E-04
10	NM_017899	Homo sapiens hypothetical protein FLJ20607 (TSC), mRNA	10.27	UP	6.79E-04
	NM_032367	Homo sapiens zinc finger, BED domain containing 3 (ZBED3), mRNA	10.27	UP	4.12E-05
15	NM_012206	Homo sapiens hepatitis A virus cellular receptor 1 (HAVCR1), mRNA	10.23	UP	8.15E-04
	AK092245	Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	10.21	UP	1.92E-03
20	NM_031469	Homo sapiens SH3 domain binding glutamic acid-rich protein like 2 (SH3BGRL2), mRNA	10.18	UP	1.63E-05
	CA311343	UI-CF-FN0-aff-b-19-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aff-b-19-0-UI 3, mRNA sequence	10.15	UP	1.89E-05
25	NM_030923	Homo sapiens hypothetical protein DKFZp566N034 (DKFZP566N034), mRNA	10.15	UP	2.23E-05
	AK024261	Homo sapiens cDNA FLJ14199 fis, clone NT2RP3002713	10.14	UP	2.08E-04
30	NM_174900	Homo sapiens zinc finger protein 42 (ZFP42), mRNA	10.1	UP	5.14E-05
	NM_198495	Homo sapiens CTAGE family, member 4 (CTAGE4), mRNA	10.07	UP	4.65E-05
	BX105791	BX105791 Soares_testis_NHT Homo sapiens cDNA clone IMAGp9981101824 ; IMAGE:742737, mRNA sequence	10.06	UP	2.15E-03
35	AI672441	wa03c03.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2296996 3, mRNA sequence	10.05	UP	2.40E-04
	AK002097	Homo sapiens cDNA FLJ11235 fis, clone PLACE1008488	10.01	UP	2.75E-05
40	BX647256	Homo sapiens mRNA; cDNA DKFZp686K0753 (from clone DKFZp686K0753)	9.97	UP	1.81E-03
	AK092456	Homo sapiens cDNA FLJ35137 fis, clone PLACE6009419	9.96	UP	5.02E-05
45	NM_001853	Homo sapiens collagen, type IX, alpha 3 (COL9A3), mRNA	9.96	UP	2.87E-05
	NM_004946	Homo sapiens dedicator of cytokinesis 2 (DOCK2), mRNA	9.94	UP	3.94E-05
	NM_005139	Homo sapiens annexin A3 (ANXA3), mRNA	9.92	UP	5.14E-05
50	NM_002260	Homo sapiens killer cell lectin-like receptor subfamily C, member 2 (KLRC2), mRNA	9.92	UP	5.95E-04
	BX649112	Homo sapiens mRNA; cDNA DKFZp686E02109 (from clone DKFZp686E02109)	9.89	UP	6.13E-05
	AK127437	Homo sapiens cDNA FLJ45529 fis, clone BRTHA2027546	9.85	UP	2.09E-05
55	NM_080659	Homo sapiens similar to RIKEN cDNA 2310030G06 gene (MGC14839), mRNA	9.81	UP	1.33E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_014867	Homo sapiens KIAA0711 gene product (KIAA0711), mRNA	9.81	UP	4.12E-05
10	BG149255	nad25d01.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3366553 3, mRNA sequence	9.7	UP	1.54E-04
	AK022598	Homo sapiens cDNA FLJ12536 fis, clone NT2RM4000265	9.7	UP	1.38E-02
15	AI792194	ov03c02.y5 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1636226 5, mRNA sequence	9.65	UP	6.10E-04
	NM_005502	Homo sapiens ATP-binding cassette, subfamily A (ABC1), member 1 (ABCA1), mRNA	9.65	UP	4.03E-04
20	BE070450	QV4-BT0407-020300-122-d08 BT0407 Homo sapiens cDNA, mRNA sequence	9.62	UP	5.52E-05
	NM_006255	Homo sapiens protein kinase C, eta (PRKCH), mRNA	9.56	UP	2.09E-05
	NM_005114	Homo sapiens heparan sulfate (glucosamine) 3-O-sulfotransferase 1 (HS3ST1), mRNA	9.55	UP	1.44E-04
25	AK123427	Homo sapiens cDNA FLJ41433 fis, clone BRHIP2007307	9.46	UP	1.99E-04
	NM_013430	Homo sapiens gamma-glutamyltransferase 1 (GGT1), transcript variant 3, mRNA	9.4	UP	1.15E-04
	AK094353	Homo sapiens cDNA FLJ37034 fis, clone BRACE2011478	9.4	UP	3.08E-05
30	NM_016584	Homo sapiens interleukin 23, alpha subunit p19 (IL23A), mRNA	9.38	UP	3.77E-05
	NM_002089	Homo sapiens chemokine (C-X-C motif) ligand 2 (CXCL2), mRNA	9.38	UP	4.55E-04
35	NM_032023	Homo sapiens Ras association (RaGDS/AF-6) domain family 4 (RASSF4), transcript variant 1, mRNA	9.38	UP	5.39E-05
	D62831	HUM330B12B Clontech human aorta polyA+ mRNA (#6572) Homo sapiens cDNA clone GEN-330B12 5, mRNA sequence	9.32	UP	9.48E-05
40	AI963999	wt87g07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2514492 3, mRNA sequence	9.31	UP	8.57E-04
	NM_005978	Homo sapiens S100 calcium binding protein A2 (S100A2), mRNA	9.31	UP	1.64E-05
45	NM_006622	Homo sapiens polo-like kinase 2 (Drosophila) (PLK2), mRNA	9.3	UP	2.93E-05
	NM_033495	Homo sapiens kelch-like 13 (Drosophila) (KLHL13), mRNA	9.25	UP	2.38E-05
	L07615	Human neuropeptide Y receptor Y1 (NPYY1) mRNA, exon 2-3 and complete cds	9.23	UP	1.09E-03
50	NM_018076	Homo sapiens armadillo repeat containing 4 (ARMC4), mRNA	9.16	UP	5.96E-04
	CA778369	ip17f03.y1 HR85 islet Homo sapiens cDNA clone IMAGE:6217493 5, mRNA sequence	9.12	UP	5.40E-05
55	NM_024554	Homo sapiens piggyBac transposable element derived 5 (PGBD5), mRNA	9.12	UP	2.63E-04
	NM_182487	Homo sapiens olfactomedin-like 2A (OLFML2A), mRNA	9.04	UP	1.64E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	T78754	yd01f08.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:24180 5, mRNA sequence	9.01	UP	1.09E-03
10	NM_004024	Homo sapiens activating transcription factor 3 (ATF3), mRNA	9	UP	1.50E-03
	NM_030583	Homo sapiens matrilin 2 (MATN2), transcript variant 2, mRNA	8.98	UP	2.15E-04
15	NM_145032	Homo sapiens F-box and leucine-rich repeat protein 13 (FBXL13), mRNA	8.97	UP	1.78E-05
	NM_020130	Homo sapiens chromosome 8 open reading frame 4 (C8orf4), mRNA	8.93	UP	4.79E-04
20	NM_005737	Homo sapiens ADP-ribosylation factor-like 7 (ARL7), mRNA	8.87	UP	1.79E-04
	BG675167	602621444F1 NCI_CGAP_Skn3 Homo sapiens cDNA clone IMAGE:4755106 5, mRNA sequence	8.86	UP	7.41E-04
	NM_194463	Homo sapiens ring finger protein 128 (RNF128), transcript variant 1, mRNA	8.85	UP	8.97E-05
25	NM_018349	Homo sapiens multiple C2-domains with two transmembrane regions 2 (MCTP2), mRNA	8.81	UP	8.37E-05
	AK026740	Homo sapiens cDNA: FLJ23087 fis, clone LNG06994, highly similar to AF161368 Homo sapiens HSPC105 mRNA	8.8	UP	3.17E-04
30	NM_000693	Homo sapiens aldehyde dehydrogenase 1 family, member A3 (ALDH1A3), mRNA	8.78	UP	4.72E-04
	NM_005459	Homo sapiens guanylate cyclase activator 1C (GUCA1C), mRNA	8.77	UP	1.57E-04
35	BF509074	UI-H-BI4-aou-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086150 3, mRNA sequence	8.74	UP	2.11E-05
	AL117425	Homo sapiens mRNA; cDNA DKFZp566L203 (from clone DKFZp566L203)	8.69	UP	2.83E-04
40	NM_006257	Homo sapiens protein kinase C, theta (PRKCQ), mRNA	8.64	UP	1.78E-05
	NM_005567	Homo sapiens lectin, galactoside-binding, soluble, 3 binding protein (LGALS3BP), mRNA	8.64	UP	1.26E-05
	NM_173508	Homo sapiens solute carrier family 35, member F3 (SLC35F3), mRNA	8.62	UP	1.04E-04
45	AK025281	Homo sapiens cDNA: FLJ21628 fis, clone COL08076	8.48	UP	7.96E-05
	NM_145252	Homo sapiens similar to common salivary protein 1 (LOC124220), mRNA	8.48	UP	1.94E-04
	A1764969	wh57b02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384811 3, mRNA sequence	8.48	UP	6.82E-05
50	NM_002313	Homo sapiens actin binding LIM protein 1 (ABLIM1), transcript variant 1, mRNA	8.48	UP	3.03E-05
	NM_017641	Homo sapiens kinesin family member 21A (KIF21A), mRNA	8.42	UP	2.43E-05
55	AA564703	nj22h06.s1 NCI_CGAP_AA1 Homo sapiens cDNA clone IMAGE:993275 3, mRNA sequence	8.4	UP	6.11E-05
	NM_001323	Homo sapiens cystatin E/M (CST6), mRNA	8.37	UP	1.02E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_014333	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	8.35	UP	1.82E-03
10	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	8.33	UP	6.04E-05
	NM_000186	Homo sapiens complement factor H (CFH), mRNA	8.32	UP	1.06E-03
	BF512544	UI-H-BW1-amf-c-08-0-UIs1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069687 3, mRNA sequence	8.31	UP	1.09E-04
15					
	AF036977	Homo sapiens clone HCG IV.9 unknown mRNA	8.27	UP	1.12E-03
	AK095500	Homo sapiens cDNA FLJ38181 fis, clone FCBBF1000125	8.19	UP	6.56E-05
20	NM_001955	Homo sapiens endothelin 1 (EDN1), mRNA	8.18	UP	4.43E-04
	NM_002053	Homo sapiens guanylate binding protein 1, interferon-inducible, 67kDa (GBP1), mRNA	8.17	UP	8.39E-06
	NM_031476	Homo sapiens hypothetical protein DKFZp434B044 (DKFZP434B044), mRNA	8.12	UP	8.06E-04
25	AA844712	ai70e12.s1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1376206 3, mRNA sequence	8.11	UP	2.36E-04
	BM976385	UI-CF-EN1-acz-f-03-0-UIs1 UI-CF-EN1 Homo sapiens cDNA clone UI-CF-EN1-acz-f-03-0-UI 3, mRNA sequence	8.11	UP	7.27E-04
30					
	BX647876	Homo sapiens mRNA; cDNA DKFZp313A1525 (from clone DKFZp313A1525)	8.11	UP	6.87E-04
	NM_001845	Homo sapiens collagen, type IV, alpha 1 (COL4A1), mRNA	8.06	UP	2.72E-05
35	NM_014900	Homo sapiens COBL-like 1 (COBLL1), mRNA	8.04	UP	9.67E-05
	NM_203306	Homo sapiens hypothetical protein MGC39606 (MGC39606), mRNA	7.98	UP	1.95E-03
	BI963896	ie66c06.x1 Melton Normalized Human Islet 4 N4-HIS 1 Homo sapiens cDNA clone IMAGE:5671691 3, mRNA sequence	7.97	UP	2.73E-05
40					
	NM_005025	Homo sapiens serine (or cysteine) proteinase inhibitor, clade I (neuroserpin), member 1 (SERPINI1), mRNA	7.97	UP	3.03E-05
	AK024238	Homo sapiens cDNA FLJ14176 fis, clone NT2RP2003101	7.96	UP	4.75E-05
45					
	NM_005951	Homo sapiens metallothionein 1 H (MT1 H), mRNA	7.95	UP	3.19E-03
	CF887677	UI-CF-FN0-aev-o-22-18-UI.r18 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aev-o-22-18-UI 5, mRNA sequence	7.93	UP	4.30E-05
50					
	NM_024423	Homo sapiens desmocollin 3 (DSC3), transcript variant Dsc3b, mRNA	7.91	UP	1.23E-04
	AI355761	qt94a11.x1 NCI_CGAP_Co14 Homo sapiens cDNA clone IMAGE:1962908 3 similar to gb:X74929 KERATIN, TYPE II CYTOSKELETAL 8 (HUMAN);, mRNA sequence	7.91	UP	6.82E-05
55					
	BF445031	nad20f02.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3366266 3, mRNA sequence	7.91	UP	1.78E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
10	CA444471	UI-H-DP0-avv-a-16-0-UI.s1 NCI_CGAP_Fs1 Homo sapiens cDNA clone UI-H-DPO-avv-a-16-0-UI 3, mRNA sequence	7.89	UP	7.69E-05
	NM_014942	Homo sapiens ankyrin repeat domain 6 (ANKRD6), mRNA	7.89	UP	2.49E-05
15	AK021801	Homo sapiens cDNA FLJ11739 fis, clone HEMBA1005497	7.86	UP	1.15E-04
	NM_001766	Homo sapiens CD1 D antigen, d polypeptide (CD1 D), mRNA	7.85	UP	5.10E-04
	NM_005360	Homo sapiens v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian) (MAF), mRNA	7.81	UP	7.59E-05
20	NM_018700	Homo sapiens tripartite motif-containing 36 (TRIM36), mRNA	7.8	UP	2.55E-05
	BX640643	Homo sapiens mRNA; cDNA DKFZp686O24114 (from clone DKFZp686O24114)	7.79	UP	2.59E-05
25	NM_005331	Homo sapiens hemoglobin, theta 1 (HBQ1), mRNA	7.77	UP	3.73E-04
	NM_194298	Homo sapiens solute carrier family 16 (monocarboxylic acid transporters), member 9 (SLC16A9), mRNA	7.77	UP	2.66E-04
	NM_012413	Homo sapiens glutaminyl-peptide cyclotransferase (glutaminyl cyclase) (QPCT), mRNA	7.76	UP	4.60E-04
30	NM_005860	Homo sapiens follistatin-like 3 (secreted glycoprotein) (FSTL3), mRNA	7.76	UP	2.64E-05
	AB033029	Homo sapiens mRNA for KIAA1203 protein, partial cds	7.7	UP	8.39E-06
	NM_012216	Homo sapiens midline 2 (MID2), transcript variant 1, mRNA	7.62	UP	6.12E-05
35	NM_018986	Homo sapiens SH3 domain and tetratricopeptide repeats 1 (SH3TC1), mRNA	7.61	UP	8.69E-04
	BM665101	UI-E-CQ1-aew-I-09-0-UI.s1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aew-I-09-0-UI 3, mRNA sequence	7.58	UP	4.09E-05
40	AL706653	DKFZp686E1543_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686E1543 5, mRNA sequence	7.57	UP	2.36E-05
	BF920409	QV2-NT0144-071100-463-b06 NT0144 Homo sapiens cDNA, mRNA sequence	7.55	UP	6.70E-04
45	NM_173462	Homo sapiens papilin, proteoglycan-like sulfated glycoprotein (PAPLN), mRNA	7.55	UP	5.64E-05
	NM_031419	Homo sapiens nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta (NFKBIZ), transcript variant 1, mRNA	7.55	UP	1.49E-04
50	NM_016423	Homo sapiens zinc finger protein 219 (ZNF219), mRNA	7.53	UP	1.71E-05
	NM_020859	Homo sapiens Shroom-related protein (ShrmL), mRNA	7.52	UP	1.26E-05
	NM_004932	Homo sapiens cadherin 6, type 2, K-cadherin (fetal kidney) (CDH6), mRNA	7.52	UP	1.64E-05
55	NM_182797	Homo sapiens phospholipase C, beta 4 (PLCB4), transcript variant 2, mRNA	7.52	UP	5.49E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AL512697	Homo sapiens mRNA; cDNA DKFZp547F134 (from clone DKFZp547F134)	7.5	UP	6.17E-05
10	BC035116	Homo sapiens cDNA clone IMAGE:5263177, partial cds	7.47	UP	2.32E-05
	AL353944	Homo sapiens mRNA; cDNA DKFZp761J1112 (from clone DKFZp761J1112)	7.45	UP	1.72E-05
	NM_152780	Homo sapiens hypothetical protein FLJ14503 (FLJ14503), mRNA	7.45	UP	1.19E-03
15	AA888443	nw74f10.s1 NCI_CGAP_Pr12 Homo sapiens cDNA clone IMAGE:1252363, mRNA sequence	7.43	UP	5.18E-04
	AW450938	UI-H-BI3-all-g-05-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2737329 3, mRNA sequence	7.4	UP	4.98E-05
20					
	NM_053039	Homo sapiens UDP glycosyltransferase 2 family, polypeptide B28 (UGT2B28), mRNA	7.38	UP	1.12E-03
	NM_016946	Homo sapiens F11 receptor (F11 R), transcript variant 1, mRNA	7.36	UP	4.65E-05
25	AW969742	EST381820 MAGE resequences, MAGK Homo sapiens cDNA, mRNA sequence	7.36	UP	1.15E-04
	BX117866	BX117866 NCI_CGAP_GCB1 Homo sapiens cDNA clone IMAGp998N233105 ; IMAGE:1234774, mRNA sequence	7.34	UP	2.20E-04
30					
	NM_005739	Homo sapiens RAS guanyl releasing protein 1 (calcium and DAG-regulated) (RASGRP1), mRNA	7.31	UP	2.20E-04
	CB047287	NISC_gg01h01.y1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3253464 5, mRNA sequence	7.25	UP	3.52E-05
35	NM_013951	Homo sapiens paired box gene 8 (PAX8), transcript variant PAX8B, mRNA	7.25	UP	5.00E-05
	AK125695	Homo sapiens cDNA FLJ43707 fis, clone TESOP2001865	7.24	UP	1.62E-03
	NM_020796	Homo sapiens sema domain, transmembrane domain (TM), and cytoplasmic domain, (semaphorin) 6A (SEMA6A), mRNA	7.21	UP	1.01E-04
40					
	BM929354	UI-E-EJ1-aje-o-19-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-aje-o-19-0-UI 5, mRNA sequence	7.19	UP	5.49E-05
	BX103949	BX103949 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998C112296 ; IMAGE:923842, mRNA sequence	7.16	UP	6.37E-05
45					
	NM_003475	Homo sapiens chromosome 11 open reading frame 13 (C11orf13), mRNA	7.14	UP	1.56E-04
	BE735115	601566084F1 NIH_MGC_21 Homo sapiens cDNA clone IMAGE:3840837 5, mRNA sequence	7.12	UP	1.41E-03
50					
	NM_207517	Homo sapiens ADAMTS-like 3 (ADAMTSL3), mRNA	7.11	UP	1.26E-05
	BU688263	UI-CF-EC1-aea-g-11-0-UI.s1 UI-CF-EC1 Homo sapiens cDNA clone UI-CF-EC1-aea-g-11-0-UI 3, mRNA sequence	7.06	UP	2.97E-05
55	AK022971	Homo sapiens cDNA FLJ12909 fis, clone NT2RP2004400	7.06	UP	7.80E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_001740	Homo sapiens calbindin 2, 29kDa (calretinin) (CALB2), transcript variant CALB2, mRNA	7.05	UP	1.81E-04
10	NM_022073	Homo sapiens egl nine homolog 3 (C. elegans) (EGLN3), mRNA	7.04	UP	5.64E-04
	BX117230	BX117230 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:9981235695 ; IMAGE:2298718, mRNA sequence	7.03	UP	1.56E-03
15	NM_002230	Homo sapiens junction plakoglobin (JUP), transcript variant 1, mRNA	7.03	UP	2.78E-05
	AK023793	Homo sapiens cDNA FLJ13731 fis, clone PLACE3000142	7.02	UP	4.04E-04
20	AL117598	Homo sapiens mRNA; cDNA DKFZp564H1663 (from clone DKFZp564H1663)	7	UP	1.59E-04
	NM_001958	Homo sapiens eukaryotic translation elongation factor 1 alpha 2 (EEF1A2), mRNA	6.99	UP	5.20E-04
25	NM_007029	Homo sapiens stathmin-like 2 (STMN2), mRNA	6.97	UP	4.47E-04
	BE968596	601649770F1 NIH_MGC_74 Homo sapiens cDNA clone IMAGE:3933472 5, mRNA sequence	6.97	UP	8.39E-04
	AK126467	Homo sapiens cDNA FLJ44503 fis, clone UTERU3001158	6.97	UP	3.82E-05
30	AA832510	oe62d06.s1 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGE:1416203 3, mRNA sequence	6.93	UP	9.67E-04
	CF137545	UI-HF-BN0-ane-d-05-0-UI.r1 NIH_MGC_50 Homo sapiens cDNA clone IMAGE:3092384 5, mRNA sequence	6.93	UP	2.70E-03
35	NM_004895	Homo sapiens cold autoimmune inflammatory syndrome 1 (CIAS1), transcript variant 1, mRNA	6.9	UP	1.46E-03
	NM_018424	Homo sapiens erythrocyte membrane protein band 4.1 like 4B (EPB41 L4B), mRNA	6.9	UP	4.68E-05
	NM_004390	Homo sapiens cathepsin H (CTSH), transcript variant 1, mRNA	6.89	UP	2.72E-05
40	NM_024603	Homo sapiens hypothetical protein FLJ11588 (FLJ11588), mRNA	6.87	UP	3.80E-05
	AI492941	qz42h08.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2029599 3, mRNA sequence	6.82	UP	8.39E-04
45	NM_003985	Homo sapiens tyrosine kinase, non-receptor, 1 (TNK1), mRNA	6.8	UP	4.55E-05
	AK095399	Homo sapiens cDNA FLJ38080 fis, clone CTONG2016185	6.8	UP	9.23E-04
50	BQ020597	UI-H-DP0-avd-a-13-0-UI.s1 NCI_CGAP_Fs1 Homo sapiens cDNA clone IMAGE:5877780 3, mRNA sequence	6.8	UP	4.53E-05
	NM_001432	Homo sapiens epiregulin (EREG), mRNA	6.77	UP	1.25E-05
	NM_005485	Homo sapiens poly (ADP-ribose) polymerase family, member 3 (PARP3), transcript variant 2, mRNA	6.76	UP	6.46E-05
55	AB058769	Homo sapiens mRNA for KIAA1866 protein, partial cds	6.76	UP	9.84E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	BU753362	UI-1-BB1-air-h-09-0-UI.s1 NCI_CGAP_PI5 Homo sapiens cDNA clone UI-1-BB1-air-h-09-0-UI 3, mRNA sequence	6.75	UP	1.08E-04
10					
	AL834140	Homo sapiens mRNA; cDNA DKFZp434A2029 (from clone DKFZp434A2029)	6.73	UP	4.49E-05
	BX092501	BX092501 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998K143946 ; IMAGE:1557637, mRNA sequence	6.71	UP	2.56E-04
15					
	BX640973	Homo sapiens mRNA; cDNA DKFZp686B15184 (from clone DKFZp686B15184)	6.71	UP	2.36E-04
	NM_015277	Homo sapiens neural precursor cell expressed, developmentally down-regulated 4-like (NEDD4L), mRNA	6.7	UP	1.60E-05
20					
	NM_021192	Homo sapiens homeo box D11 (HOXD11), mRNA	6.69	UP	8.94E-04
	NM_032621	Homo sapiens brain expressed X-linked 2 (BEX2), mRNA	6.69	UP	1.83E-04
25					
	X02851	Human mRNA for interleukin-1 precursor (pre IL-1)	6.68	UP	3.32E-04
	NM_014258	Homo sapiens synaptonemal complex protein 2 (SYCP2), mRNA	6.68	UP	4.96E-02
	NM_000682	Homo sapiens adrenergic, alpha-2B-, receptor (ADRA2B), mRNA	6.67	UP	3.64E-05
30					
	BQ435580	AGENCOURT_7836890 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6102371 5, mRNA sequence	6.64	UP	1.64E-04
	NM_024608	Homo sapiens nei endonuclease VIII-like 1 (E. coli) (NEIL1), mRNA	6.63	UP	3.23E-05
35					
	AK057113	Homo sapiens cDNA FLJ32551 fis, clone SPLEN1000087	6.63	UP	6.06E-04
	NM_130897	Homo sapiens dynein, cytoplasmic, light polypeptide 2B (DNCL2B), mRNA	6.62	UP	2.40E-03
	N39597	yy51e04.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:277086 3, mRNA sequence	6.62	UP	8.33E-04
40					
	BU616749	UI-H-FH1-bfj-a-11-0-UI.s1 NCI_CGAP_FH1 Homo sapiens cDNA clone UI-H-FH1-bfj-a-11-0-UI 3, mRNA sequence	6.59	UP	1.93E-04
45					
	AW 195474	xn38g09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2696032 3, mRNA sequence	6.59	UP	6.82E-05
	AK127437	Homo sapiens cDNA FLJ45529 fis, clone BRTHA2027546	6.57	UP	1.76E-04
50					
	NM_004717	Homo sapiens diacylglycerol kinase, iota (DGKI), mRNA	6.56	UP	7.54E-05
	BU584197	2513030T6 LIVRTUT04 Homo sapiens cDNA clone 2513030 3, mRNA sequence	6.53	UP	8.82E-04
	BU634363	UI-H-FL1-bgx-o-20-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgx-o-20-0-UI 3, mRNA sequence	6.53	UP	1.24E-04
55					
	NM_178868	Homo sapiens chemokine-like factor super family 8 (CKLFSF8), mRNA	6.52	UP	1.59E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_002247	Homo sapiens potassium large conductance calcium-activated channel, subfamily M, alpha member 1 (KCNMA1), mRNA	6.49	UP	4.49E-05
10					
	AW 188195	xj93e12.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2664814 3 similar to contains element THR repetitive element ;, mRNA sequence	6.49	UP	3.81E-05
15					
	M_153634	Homo sapiens copine VIII (CPNE8), mRNA	6.48	UP	1.72E-04
	NM_145205	Homo sapiens HMG2 like (LOC127540), mRNA	6.47	UP	6.39E-05
	M_178868	Homo sapiens chemokine-like factor super family 8 (CKLFSF8), mRNA	6.47	UP	1.15E-05
20					
	U63828	Human tissue plasminogen activator mRNA, partial cds	6.46	UP	5.51E-05
	NM_004529	Homo sapiens myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 3 (MLLT3), mRNA	6.46	UP	8.21E-05
25					
	BX647655	Homo sapiens mRNA; cDNA DKFZp451A211 (from clone DKFZp451A211)	6.46	UP	6.94E-05
	AW470401	xz83g01.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:2870832 3 similar to contains L1.t2 L1 repetitive element ;, mRNA sequence	6.46	UP	1.30E-04
30					
	BG570144	602591134F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4717761 5, mRNA sequence	6.45	UP	1.08E-03
	BX115325	BX115325 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGp998K085276 ; IMAGE:2137855, mRNA sequence	6.45	UP	1.04E-04
35					
	NM_002214	Homo sapiens integrin, beta 8 (ITGB8), mRNA	6.44	UP	6.77E-05
	CA314926	UI-CF-FN0-afi-b-19-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afi-b-19-0-UI 3, mRNA sequence	6.43	UP	6.61E-05
40					
	AK093870	Homo sapiens cDNA FLJ36551 fis, clone TRACH2008127	6.43	UP	1.28E-03
	NM_000860	Homo sapiens hydroxyprostaglandin dehydrogenase 15-(NAD) (HPGD), mRNA	6.42	UP	1.76E-05
45					
	AI831068	wj62d12.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2407415 3, mRNA sequence	6.41	UP	1.37E-04
	NM_000349	Homo sapiens steroidogenic acute regulator (STAR), nuclear gene encoding mitochondrial protein, transcript variant 1, mRNA	6.41	UP	1.26E-04
50					
	NM_014181	Homo sapiens HSPC159 protein (HSPC159), mRNA	6.4	UP	1.67E-04
	NM_024534	Homo sapiens hypothetical protein FLJ12684 (FLJ12684), mRNA	6.38	UP	6.57E-04
	NM_002843	Homo sapiens protein tyrosine phosphatase, receptor type, J (PTPRJ), mRNA	6.38	UP	1.32E-03
55					
	A1926616	wo48e04.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2458590 3 similar to contains MER27.b2 MER27 repetitive element ;, mRNA sequence	6.37	UP	7.01E-03
	NM_002273	Homo sapiens keratin 8 (KRT8), mRNA	6.36	UP	2.91E-03
	NM_207303	Homo sapiens attractin-like 1 (ATRNL1), mRNA	6.35	UP	7.85E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AK001007	Homo sapiens cDNA FLJ10145 fis, clone HEMBA1003322	6.35	UP	1.82E-05
10	CB047092	NISC_gf08f03.xl NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:3253013 3, mRNA sequence	6.33	UP	6.92E-05
	BC060805	Homo sapiens hypothetical protein FLJ12788, mRNA (cDNA clone IMAGE:5266931), partial cds	6.31	UP	2.93E-05
15	AI359782	qy41h10.x1 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:2014627 3, mRNA sequence	6.31	UP	3.23E-04
	CA426602	UI-H-FE1-bef-f-11-0-UI.s1 NCI_CGAP_FE1 Homo sapiens cDNA clone UI-H-FE1-bef-f-11-0-UI 3, mRNA sequence	6.29	UP	2.14E-04
20	AK091686	Homo sapiens cDNA FLJ34367 fis, clone FEBRA2016621	6.27	UP	1.93E-03
	AK091336	Homo sapiens cDNA FLJ34017 fis, clone FCBBF2002626	6.26	UP	5.49E-05
25	NM_015068	Homo sapiens paternally expressed 10 (PEG10), mRNA	6.26	UP	4.53E-05
	R79128	yi86c12.r1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:146134 5, mRNA sequence	6.23	UP	2.04E-04
	H08785	yI94f04.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:45882 3, mRNA sequence	6.22	UP	1.66E-03
30	AI953708	wq47d09.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474417 3, mRNA sequence	6.21	UP	4.58E-03
	NM_015236	Homo sapiens latrophilin 3 (LPHN3), mRNA	6.2	UP	5.35E-04
	AL833609	Homo sapiens mRNA; cDNA DKFZp686O1267 (from clone DKFZp686O1267)	6.19	UP	7.53E-04
35	AK127421	Homo sapiens cDNA FLJ45513 fis, clone BRTHA2021450	6.19	UP	4.68E-05
	NM_015085	Homo sapiens GTPase activating Rap/RanGAP domain-like 4 (GARNL4), mRNA	6.18	UP	6.82E-05
40	NM_001305	Homo sapiens claudin 4 (CLDN4), mRNA	6.17	UP	3.48E-05
	NM_173078	Homo sapiens SLIT and NTRK-like family, member 4 (SLITRK4), mRNA	6.16	UP	1.08E-03
	BG165745	602344592F1 NIH_MGC_89 Homo sapiens cDNA clone IMAGE:4454470 5, mRNA sequence	6.16	UP	1.20E-04
45	NM_006074	Homo sapiens tripartite motif-containing 22 (TRIM22), mRNA	6.15	UP	1.15E-03
	NM_005928	Homo sapiens milk fat globule-EGF factor 8 protein (MFGE8), mRNA	6.15	UP	1.63E-02
	BF939703	nac80e07.x1 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:3440725 3 similar to contains MER30.t3 MER30 repetitive element ; , mRNA sequence	6.12	UP	6.35E-05
50	BX115301	BX115301 NCI_CGAP_Kid5 Homo sapiens cDNA clone IMAGp998J093989 ; IMAGE:1574120, mRNA sequence	6.11	UP	8.26E-05
55	NM_018849	Homo sapiens ATP-binding cassette, subfamily B (MDR/TAP), member 4 (ABCB4), transcript variant B, mRNA	6.09	UP	3.23E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_015478	Homo sapiens I(3)mbt-like (Drosophila) (L3MBTL), transcript variant I, mRNA	6.09	UP	2.43E-05
10	NM_015669	Homo sapiens protocadherin beta 5 (PCDHB5), mRNA	6.08	UP	7.52E-05
	NM_147189	Homo sapiens hypothetical protein MGC39325 (MGC39325), mRNA	6.06	UP	3.18E-04
	BQ018133	UI-H-DP0-avv-I-18-0-UI.s1 NCI_CGAP_Fs1 Homo sapiens cDNA clone IMAGE:5884961 3, mRNA sequence	6.06	UP	1.78E-05
15	N34295	yy51e10.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:277098 3, mRNA sequence	6.04	UP	8.29E-05
20	NM_006472	Homo sapiens thioredoxin interacting protein (TXNIP), mRNA	6.03	UP	6.87E-05
	NM_012472	Homo sapiens leucine rich repeat containing 6 (LRR6), mRNA	6.02	UP	2.66E-05
25	AK123319	Homo sapiens cDNA FLJ41325 fis, clone BRAMY2046871	6	UP	2.55E-04
	AK123807	Homo sapiens cDNA FLJ41813 fis, clone NT2RI2011450	5.98	UP	7.92E-05
	NM_017786	Homo sapiens hypothetical protein FLJ20366 (FLJ20366), mRNA	5.98	UP	1.85E-03
30	NM_000212	Homo sapiens integrin, beta 3 (platelet glycoprotein IIIa, antigen CD61) (ITGB3), mRNA	5.97	UP	2.73E-05
	NM_000817	Homo sapiens glutamate decarboxylase 1 (brain, 67kDa) (GAD1), transcript variant GAD67, mRNA	5.96	UP	9.94E-03
35	AK123617	Homo sapiens cDNA FLJ41623 fis, clone CTONG3009227	5.92	UP	1.38E-03
	BI493986	df106g12.y1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2538815 5, mRNA sequence	5.91	UP	3.68E-04
40	NM_005264	Homo sapiens GDNF family receptor alpha 1 (GFRA1), transcript variant 1, mRNA	5.91	UP	4.09E-05
	CN478597	UI-CF-FN0-aeo-g-21-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aeo-g-21-0-UI 3, mRNA sequence	5.88	UP	1.13E-03
45	BQ021695	UI-H-DH1-axi-f-22-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5829141 3, m RNA sequence	5.88	UP	1.72E-04
	AL137698	Homo sapiens mRNA; cDNA DKFZp434C1915 (from clone DKFZp434C1915); partial cds	5.88	UP	2.03E-05
50	NM_014905	Homo sapiens glutaminase (GLS), mRNA	5.88	UP	1.24E-04
	AA993234	ot60a08.s1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1621142 3, mRNA sequence	5.87	UP	2.88E-05
	N38890	yy81f12.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:279983 3 similar to contains OFR.t3 OFR repetitive element ;, mRNA sequence	5.87	UP	1.25E-04
55	NM_197955	Homo sapiens normal mucosa of esophagus specific 1 (NMES1), transcript variant 1, mRNA	5.85	UP	1.85E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_014399	Homo sapiens transmembrane 4 superfamily member 13 (TM4SF13), mRNA	5.84	UP	2.18E-05
10	NM_024704	Homo sapiens chromosome 20 open reading frame 23 (C20orf23), mRNA	5.82	UP	1.24E-04
	NM_002276	Homo sapiens keratin 19 (KRT19), mRNA	5.8	UP	7.06E-03
	CD248442	AGENCOURT_14095939 NIH_MGC_172 Homo sapiens cDNA 5, mRNA sequence	5.79	UP	2.16E-04
15	AK024850	Homo sapiens cDNA: FLJ21197 fis, clone COL00201	5.79	UP	1.45E-04
	H83499	ys91f12.r1 Soares retina N2b5HR Homo sapiens cDNA clone IMAGE:222191 5, mRNA sequence	5.77	UP	1.37E-02
20	NM_153377	Homo sapiens leucine-rich repeats and immunoglobulin-like domains 3 (LRIG3), mRNA	5.77	UP	2.43E-05
	NM_153267	Homo sapiens MAM domain containing 2 (MAMDC2), mRNA	5.76	UP	2.31 E-04
	NM_144595	Homo sapiens hypothetical protein FLJ30046 (FLJ30046), mRNA	5.76	UP	1.38E-03
25	BF509155	UI-H-BI4-aov-b-05-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086145 3, mRNA sequence	5.75	UP	6.59E-05
	NM_002735	Homo sapiens protein kinase, cAMP-dependent, regulatory, type I, beta (PRKAR1 B), mRNA	5.73	UP	2.73E-05
30	NM_025044	Homo sapiens bicaudal C homolog 1 (Drosophila) (BICC1), mRNA	5.72	UP	5.81E-05
	BM969191	UI-CF-EN0-acp-e-22-0-UI.s1 UI-CF-EN0 Homo sapiens cDNA clone UI-CF-EN0-acp-e-22-0-UI 3, mRNA sequence	5.69	UP	1.67E-03
35	NM_000203	Homo sapiens iduronidase, alpha-L- (IDUA), mRNA	5.68	UP	1.10E-03
	AI693580	wd12d01.x1 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGE:2327905 3, mRNA sequence	5.67	UP	9.23E-04
40	NM_006169	Homo sapiens nicotinamide N-methyltransferase (NNMT), mRNA	5.66	UP	2.18E-05
	H46176	yo14a11.s1 Soares adult brain N2b5HB55Y Homo sapiens cDNA clone IMAGE:177884 3, mRNA sequence	5.65	UP	1.88E-05
	NM_024306	Homo sapiens fatty acid 2-hydroxylase (FA2H), mRNA	5.63	UP	4.19E-03
45	BX110683	BX110683 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGp998E175727 ; IMAGE:2310904, mRNA sequence	5.62	UP	7.42E-04
	NM_030765	Homo sapiens UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 4 (B3GNT4), mRNA	5.58	UP	1.80E-03
50	NM_003929	Homo sapiens RAB7, member RAS oncogene family-like 1 (RAB7L1), mRNA	5.55	UP	2.14E-05
	AI220066	qg84d01.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1841857 3, mRNA sequence	5.54	UP	2.13E-05
55	AA664452	ae94d02.s1 Human bone marrow stromal cells Homo sapiens cDNA clone IMAGE:1026723 3 similar to contains element LTR4 repetitive element ;, mRNA sequence	5.54	UP	4.13E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_007237	Homo sapiens SP140 nuclear body protein (SP140), transcript variant 1, mRNA	5.53	UP	2.41E-04
10	NM_173662	Homo sapiens hypothetical protein LOC285533 (LOC285533), mRNA	5.52	UP	3.69E-03
	NM_003020	Homo sapiens secretory granule, neuroendocrine protein 1 (7B2 protein) (SGNE1), mRNA	5.52	UP	2.08E-04
15	NM_206808	Homo sapiens citrate lyase beta like (CLYBL), transcript variant 2, mRNA	5.52	UP	1.87E-04
	NM_004867	Homo sapiens integral membrane protein 2A (ITM2A), mRNA	5.51	UP	2.13E-05
20	NM_006333	Homo sapiens nuclear DNA-binding protein (C1D), transcript variant 1, mRNA	5.51	UP	7.84E-04
	NM_014251	Homo sapiens solute carrier family 25, member 13 (citrin) (SLC25A13), mRNA	5.5	UP	5.96E-04
	NM_006863	Homo sapiens leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 1 (LILRA1), mRNA	5.49	UP	6.51E-05
25					
	AK057166	Homo sapiens cDNA FLJ32604 fis, clone STOMA1000133	5.49	UP	1.85E-03
	NM_000076	Homo sapiens cyclin-dependent kinase inhibitor 1C (p57, Kip2) (CDKN1C), mRNA	5.48	UP	4.65E-05
30					
	BC040701	Homo sapiens cDNA clone IMAGE:5736259, partial cds	5.47	UP	1.48E-04
	NM_152765	Homo sapiens hypothetical protein MGC33510 (MGC33510), mRNA	5.46	UP	4.76E-05
	NM_019034	Homo sapiens ras homolog gene family, member F (in filopodia) (RHOF), mRNA	5.46	UP	2.64E-05
35					
	S81734	tissue transglutaminase homologue {alternatively spliced} [human, erythroleukemia cell line HEL GM06141A, mRNA, 2362 nt]	5.46	UP	3.92E-05
40					
	BM670793	UI-E-DX1-agv-p-03-0-UI.s1 UI-E-DX1 Homo sapiens cDNA clone UI-E-DX1-agv-p-03-0-UI 3, mRNA sequence	5.46	UP	3.77E-05
	NM_002829	Homo sapiens protein tyrosine phosphatase, non- receptor type 3 (PTPN3), mRNA	5.45	UP	4.79E-05
45					
	BM727151	UI-E-EJ0-aij-f-05-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aij-f-05-0-UI 5, mRNA sequence	5.45	UP	4.13E-05
	BF966833	602286668T1 NIH_MGC_95 Homo sapiens cDNA clone IMAGE:4375360 3, mRNA sequence	5.45	UP	2.83E-05
	NM_004904	Homo sapiens cAMP responsive element binding protein 5 (CREB5), mRNA	5.44	UP	1.09E-04
50					
	AA635788	nr32h01.s1 NCI_CGAP_Pr22 Homo sapiens cDNA clone IMAGE:1169713 3 similar to contains Alu repetitive element;, mRNA sequence	5.43	UP	2.02E-03
	BM701989	UI-E-CQ1-aex-j-06-0-UI.r1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aex-j-06-0-UI 5, mRNA sequence	5.42	UP	7.55E-04
55					
	NM_020665	Homo sapiens transmembrane protein 27 (TMEM27), mRNA	5.4	UP	5.46E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_019034	Homo sapiens ras homolog gene family, member F (in filopodia) (RHOF), mRNA	5.39	UP	5.72E-05
10	NM_017912	Homo sapiens hect domain and RLD 6 (HERC6), mRNA	5.38	UP	5.85E-05
	NM_152433	Homo sapiens kelch repeat and BTB (POZ) domain containing 3 (KBTBD3), transcript variant 1, mRNA	5.37	UP	4.29E-04
	CA314843	UI-CF-FN0-afi-a-06-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-afi-a-06-0-UI 3, mRNA sequence	5.37	UP	2.05E-04
15	NM_006058	Homo sapiens TNFAIP3 interacting protein 1 (TNIP1), mRNA	5.36	UP	2.25E-03
	BU191317	AGENCOURT_8074912NIH_MGC_110 Homo sapiens cDNA clone IMAGE:6086274 5, mRNA sequence	5.36	UP	4.45E-02
20	AL049437	Homo sapiens mRNA; cDNA DKFZp586E1120 (from clone DKFZp586E1120)	5.34	UP	7.78E-04
	NM_030952	Homo sapiens likely ortholog of rat SNF1/AMP-activated protein kinase (SNARK), mRNA	5.31	UP	4.21E-05
25	NM_004170	Homo sapiens solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1 (SLC1A1), mRNA	5.31	UP	1.15E-04
	AW269776	xv45b09.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2816057 3, mRNA sequence	5.3	UP	1.97E-04
30	BM977716	UI-CF-EN1-aef-b-21-0-UI.s1 UI-CF-EN1 Homo sapiens cDNA clone UI-CF-EN1-aef-b-21-0-UI 3, mRNA sequence	5.3	UP	4.01 E-03
	AK023658	Homo sapiens cDNA FLJ13596 fis, clone PLACE1009637	5.29	UP	6.83E-04
35	AF196185	Homo sapiens atypical PKC isotype-specific interacting protein long variant mRNA, complete cds	5.28	UP	4.57E-05
	AL045014	DKFZp434F134_s1 434 (synonym: htes3) Homo sapiens cDNA clone DKFZp434F134 3, mRNA sequence	5.27	UP	2.67E-05
40	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	5.27	UP	2.88E-03
	NM_033132	Homo sapiens Zic family member 5 (odd-paired homolog, Drosophila) (ZIC5), mRNA	5.26	UP	2.09E-05
	NM_198182	Homo sapiens transcription factor CP2-like 2 (TFCP2L2), transcript variant 2, mRNA	5.26	UP	1.35E-02
45	NM_001657	Homo sapiens amphiregulin (schwannoma-derived growth factor) (AREG), mRNA	5.26	UP	3.99E-05
	R92346	yq06b10.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:196123 3, mRNA sequence	5.25	UP	1.16E-04
50	BX537539	Homo sapiens mRNA; cDNA DKFZp686A1130 (from clone DKFZp686A1130)	5.25	UP	4.80E-04
	AB041269	Homo sapiens mRNA for keratin 19, partial cds, isolate:K19-141	5.23	UP	1.15E-02
55	BX090717	BX090717 NCI_CGAP_Kid5 Homo sapiens cDNA clone IMAGp9980154699 ; IMAGE:1916390, mRNA sequence	5.2	UP	9.02E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AK025743	Homo sapiens cDNA: FLJ22090 fis, clone HEP16084	5.2	UP	1.04E-04
10	NM_002413	Homo sapiens microsomal glutathione S-transferase 2 (MGST2), mRNA	5.2	UP	3.34E-05
	AA780946	ag99c12.s1 Gessler Wilms tumor Homo sapiens cDNA clone IMAGE:1155286 3, mRNA sequence	5.2	UP	2.40E-04
	NM_032488	Homo sapiens cornifelin (CNFN), mRNA	5.2	UP	1.78E-05
15	NM_024997	Homo sapiens activating transcription factor 7 interacting protein 2 (ATF7IP2), mRNA	5.19	UP	2.40E-03
	NM_002222	Homo sapiens inositol 1,4,5-triphosphate receptor, type 1 (ITPR1), mRNA	5.19	UP	1.41E-04
20	NM_000784	Homo sapiens cytochrome P450, family 27, subfamily A, polypeptide 1 (CYP27A1), nuclear gene encoding mitochondrial protein, mRNA	5.17	UP	7.39E-03
	NM_005760	Homo sapiens CCAAT/enhancer binding protein zeta (CEBPZ), mRNA	5.16	UP	5.52E-05
25	BQ011746	UI-1-BC1p-atk-b-09-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1p-atk-b-09-0-UI 3, mRNA sequence	5.16	UP	5.86E-04
	NM_032728	Homo sapiens chromosome 9 open reading frame 67 (C9orf67), mRNA	5.15	UP	1.85E-03
30	NM_005755	Homo sapiens Epstein-Barr virus induced gene 3 (EBI3), mRNA	5.15	UP	2.33E-04
	AK056852	Homo sapiens cDNA FLJ32290 fis, clone PROST2000463	5.15	UP	1.32E-03
	AK054990	Homo sapiens cDNA FLJ30428 fis, clone BRACE2008941	5.15	UP	1.85E-03
35	D54580	HUM144G01 B Clontech human fetal brain polyA+ mRNA (#6535) Homo sapiens cDNA clone GEN-144G01 5, mRNA sequence	5.13	UP	2.03E-04
40	AL137535	Homo sapiens mRNA; cDNA DKFZp434H2019 (from clone DKFZp434H2019)	5.13	UP	2.45E-02
	AL110252	Homo sapiens mRNA; cDNA DKFZp566A1046 (from clone DKFZp566A1046)	5.12	UP	2.19E-03
	AI659523	tt99d12.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2249687 3, mRNA sequence	5.1	UP	1.16E-03
45	AW137001	UI-H-BI1-acu-c-05-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2715632 3, mRNA sequence	5.1	UP	6.90E-05
	BM969331	UI-CF-DU1-aar-g-23-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aar-g-23-0-UI 3, mRNA sequence	5.08	UP	3.31E-03
50	M_178823	Homo sapiens chromosome 6 open reading frame 165 (C6orf165), mRNA	5.08	UP	3.66E-05
	NM_018476	Homo sapiens brain expressed, X-linked 1 (BEX1), mRNA	5.06	UP	3.05E-05
55	BF107212	601824290F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4043879 5, mRNA sequence	5.06	UP	1.75E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p-value
5					
	NM_052890	Homo sapiens peptidoglycan recognition protein 2 (PGLYRP2), mRNA	5.05	UP	3.95E-05
10	BX459043	BX459043 Homo sapiens PLACENTA Homo sapiens cDNA clone CSODE011YN10 3-PRIME, mRNA sequence	5.03	UP	2.88E-05
	NM_032857	Homo sapiens lactamase, beta (LACTB), nuclear gene encoding mitochondrial protein, transcript variant 1, mRNA	5.02	UP	1.76E-05
15					
	NM_022746	Homo sapiens hypothetical protein FLJ22390 (FLJ22390), mRNA	5.02	UP	4.49E-04
	AK025909	Homo sapiens cDNA: FLJ22256 fis, clone HRC02860	5.02	UP	4.54E-04
20	NM_002247	Homo sapiens potassium large conductance calcium-activated channel, subfamily M, alpha member 1 (KCNMA1), mRNA	5.02	UP	1.33E-03
	AI434849	ti13b01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2130313 3 similar to contains MER37.b1 MER37 MER37 repetitive element ;, mRNA sequence	5.01	UP	2.82E-04
25					
	AK091933	Homo sapiens cDNA FLJ34614 fis, clone KIDNE2014268	5.01	UP	2.18E-05
	NM_004331	Homo sapiens BCL2/adenovirus E1B 19kDa interacting protein 3-like (BNIP3L), mRNA	5	UP	3.53E-05
30					
	AL119769	DKFZp761E1224_r1 761 (synonym: hamy2) Homo sapiens cDNA clone DKFZp761 E1224 5, mRNA sequence	5	UP	1.92E-03
	NM_001451	Homo sapiens forkhead box F1 (FOXF1), mRNA	465.04	Down	2.67E-05
35	M_144594	Homo sapiens hypothetical protein FLJ32942 (FLJ32942), mRNA	379.94	Down	1.12E-05
	NM_002421	Homo sapiens matrix metalloproteinase 1 (interstitial collagenase) (MMP1), mRNA	365.32	Down	2.43E-05
	AK026784	Homo sapiens cDNA: FLJ23131 fis, clone LNG08502	331.17	Down	8.46E-06
40	NM_016307	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	309.99	Down	8.39E-06
	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	232.51	Down	8.39E-06
	NM_000089	Homo sapiens collagen, type I, alpha 2 (COL1A2), mRNA	220.95	Down	2.96E-05
45					
	NM_007036	Homo sapiens endothelial cell-specific molecule 1 (ESM1), mRNA	217.57	Down	2.01E-05
	AK021543	Homo sapiens cDNA FLJ11481 fis, clone HEMBA1001803	215.51	Down	2.31E-05
50					
	NM_004460	Homo sapiens fibroblast activation protein, alpha (FAP), mRNA	205.38	Down	1.15E-05
	AK122739	Homo sapiens cDNA FLJ16260 fis, clone IMR322006947, highly similar to Rattus norvegicus mRNA for dHand protein	190.76	Down	1.15E-05
55					
	NM_000474	Homo sapiens twist homolog 1 (acrocephalosyndactyly 3; Saethre-Chotzen syndrome) (Drosophila) (TWIST1), mRNA	151.94	Down	5.23E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_152270	Homo sapiens hypothetical protein FLJ34922 (FLJ34922), mRNA	144.24	Down	1.73E-05
10	NM_001884	Homo sapiens hyaluronan and proteoglycan link protein 1 (HAPLN1), mRNA	134.45	Down	2.18E-05
	NM_000710	Homo sapiens bradykinin receptor B1 (BDKRB1), mRNA	132.74	Down	1.18E-05
15	NM_021637	Homo sapiens transmembrane protein 35 (TMEM35), mRNA	126.76	Down	8.39E-06
	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	106.55	Down	2.28E-05
20	NM_000362	Homo sapiens tissue inhibitor of metalloproteinase 3 (Sorsby fundus dystrophy, pseudoinflammatory) (TIMP3), mRNA	102.59	Down	1.76E-05
	NM_002091	Homo sapiens gastrin-releasing peptide (GRP), mRNA	101.58	Down	2.46E-05
	NM_000609	Homo sapiens chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1) (CXCL12), mRNA	97.42	Down	3.48E-05
25	NM_032638	Homo sapiens GATA binding protein 2 (GATA2), mRNA	84.58	Down	1.37E-05
	NM_198148	Homo sapiens carboxypeptidase X (M14 family), member 2 (CPXM2), mRNA	77.44	Down	9.50E-05
	W38393	zb15c07.r1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:302124 5, mRNA sequence	77.37	Down	1.26E-05
30	AL831863	Homo sapiens mRNA; cDNA DKFZp761J2017 (from clone DKFZp761J2017)	77.09	Down	2.65E-05
	AB067499	Homo sapiens mRNA for KIAA1912 protein, partial cds	76.12	Down	3.65E-05
	BX089019	BX089019 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998K243513 ; IMAGE:1391375, mRNA sequence	72.98	Down	1.25E-05
35					
	AI962169	wq45c10.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474226 3, mRNA sequence	65.52	Down	4.53E-05
40	NM_139211	Homo sapiens homeodomain-only protein (HOP), transcript variant 2, mRNA	64.01	Down	1.72E-05
	CD677332	ho15f06.y1 Human Trabecular meshwork cDNA: hohphq Homo sapiens cDNA clone ho15f06 5, mRNA sequence	63.91	Down	2.35E-05
45	NM_001442	Homo sapiens fatty acid binding protein 4, adipocyte (FABP4), mRNA	63.47	Down	1.37E-05
	NM_205855	Homo sapiens HWKM1940 (UNQ1940), mRNA	58.05	Down	1.89E-05
	W03013	za02c04.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:291366 5 similar to contains THR.t3 THR repetitive element ;, mRNA sequence	53.58	Down	2.67E-06
50	NM_002852	Homo sapiens pentaxin-related gene, rapidly induced by IL-1 beta (PTX3), mRNA	53.13	Down	8.15E-04
	NM_024633	Homo sapiens chromosome 14 open reading frame 139 (C14orf139), mRNA	51.58	Down	3.58E-05
55	AK021543	Homo sapiens cDNA FLJ11481 fis, clone HEMBA1001803	51.28	Down	1.09E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_002593	Homo sapiens procollagen C-endopeptidase enhancer (PCOLCE), mRNA	48.32	Down	2.03E-05
10	AI422199	tf58d04.x1 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:2103463 3, mRNA sequence	48.23	Down	4.98E-05
	NM_014459	Homo sapiens protocadherin 17 (PCDH17), mRNA	45.53	Down	2.11E-05
	NM_016588	Homo sapiens neuritin 1 (NRN1), mRNA	44.52	Down	1.78E-05
15	NM_012242	Homo sapiens dickkopf homolog 1 (Xenopus laevis) (DKK1), mRNA	43.26	Down	7.16E-06
	NM_000685	Homo sapiens angiotensin II receptor, type 1 (AGTR1), transcript variant 1, mRNA	42.51	Down	2.57E-04
20	AK056725	Homo sapiens cDNA FLJ32163 fis, clone PLACE6000371	42.21	Down	2.67E-06
	AI124557	am58g02.x1 Johnston frontal cortex Homo sapiens cDNA clone IMAGE:1539794 3, mRNA sequence	41.42	Down	8.25E-03
	NM_020404	Homo sapiens CD164 sialomucin-like 1 (CD164L1), mRNA	39.82	Down	2.32E-05
25	NM_015170	Homo sapiens sulfatase 1 (SULF1), mRNA	38.91	Down	2.36E-05
	CA843592	ir49c12.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6548544 3, mRNA sequence	37.67	Down	1.25E-05
	AY335938	Homo sapiens homeodomain protein IRXA1 (IRX1) mRNA, complete cds	36.92	Down	3.15E-05
30	NM_002961	Homo sapiens S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine placental homolog) (S100A4), transcript variant 1, mRNA	35.31	Down	6.57E-05
	AK091731	Homo sapiens cDNA FLJ34412 fis, clone HEART2002432	34.47	Down	2.83E-05
35	NM_006329	Homo sapiens fibulin 5 (FBLN5), mRNA	34.03	Down	4.55E-05
	BQ020357	UI-H-ED0-axk-p-07-0-UI.s1 NCI_CGAP_ED0 Homo sapiens cDNA clone IMAGE:5830134 3, mRNA sequence	33.2	Down	1.82E-05
40	NM_018013	Homo sapiens hypothetical protein FLJ10159 (FLJ10159), mRNA	32.86	Down	1.15E-05
	NM_002402	Homo sapiens mesoderm specific transcript homolog (mouse) (MEST), transcript variant 1, mRNA	32.83	Down	2.14E-05
45	BQ001571	UI-H-DH1-awr-i-18-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5893337 3, mRNA sequence	32.34	Down	8.96E-06
	NM_032777	Homo sapiens G protein-coupled receptor 124 (GPR124), mRNA	31.96	Down	1.18E-04
50	AK124396	Homo sapiens cDNA FLJ42405 fis, clone ASTRO3000474	31.36	Down	2.24E-05
	NM_002448	Homo sapiens msh homeo box homolog 1 (Drosophila) (MSX1), MRNA,	31.23	Down	2.03E-05
55	NM_000090	Homo sapiens collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant) (COL3A1), mRNA	29.77	Down	3.23E-04
	NM_001147	Homo sapiens angiopoietin 2 (ANGPT2), mRNA	29.68	Down	2.51E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_001146	Homo sapiens angiopoietin 1 (ANGPT1), transcript variant 1, mRNA	29.26	Down	6.83E-05
10	NM_000576	Homo sapiens interleukin 1, beta (IL1B), mRNA	29.06	Down	1.15E-05
	AF052115	Homo sapiens clone 23688 mRNA sequence	26.15	Down	1.15E-05
	NM_006439	Homo sapiens mab-21-like 2 (C. elegans) (MAB21L2), mRNA	25.51	Down	1.20E-04
15	NM_016192	Homo sapiens transmembrane protein with EGF-like and two follistatin-like domains 2 (TMEFF2), mRNA	25.46	Down	2.55E-05
	NM_002575	Homo sapiens serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2 (SERPINB2), mRNA	24.67	Down	5.40E-05
20	NM_007361	Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	23.56	Down	2.64E-05
	NM_152399	Homo sapiens hypothetical protein FLJ30834 (FLJ30834), mRNA	23.42	Down	6.51E-05
	NM_203370	Homo sapiens similar to RIKEN cDNA 6530418L21 (LOC389119), mRNA	22.99	Down	1.43E-05
25	NM_00100229 5	Homo sapiens GATA binding protein 3 (GATA3), transcript variant 1, mRNA	22.51	Down	1.41E-04
	AK093256	Homo sapiens cDNA FLJ35937 fis, clone TESTI2011480	22.26	Down	3.04E-05
	U83115	Human non-lens beta gamma-crystallin like protein (AIM1) mRNA, partial cds	22.18	Down	3.77E-05
30	NM_006475	Homo sapiens periostin, osteoblast specific factor (POSTN), mRNA	21.99	Down	7.38E-05
	NM_014178	Homo sapiens syntaxin binding protein 6 (amisyn) (STXBP6), mRNA	21.69	Down	3.36E-05
35	NM_005127	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 2 (activation-induced) (CLECSF2), mRNA	21.46	Down	3.71E-05
	NM_022475	Homo sapiens hedgehog interacting protein (HHIP), mRNA	21.42	Down	4.57E-05
40	BX112628	BX112628 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGp998A09669 ; IMAGE:299024, mRNA sequence	20.99	Down	4.47E-05
	NM_000810	Homo sapiens gamma-aminobutyric acid (GABA) A receptor, alpha 5 (GABRA5), mRNA	20.84	Down	5.50E-05
45	BQ934941	AGENCOURT_8810373 NIH_MGC_101 Homo sapiens cDNA clone IMAGE:6429485 5, mRNA sequence	20.45	Down	2.55E-05
	BC071787	Homo sapiens cDNA clone IMAGE:4610527, partial cds	20.44	Down	1.25E-05
	NM_033292	Homo sapiens caspase 1, apoptosis-related cysteine protease (interleukin 1, beta, convertase) (CASP1), transcript variant alpha, mRNA	19.88	Down	6.18E-05
50	NM_020809	Homo sapiens Rho GTPase activating protein 20 (ARHGAP20), mRNA	19.54	Down	5.26E-05
	BE866150	601679068F1 NIH_MGC_53 Homo sapiens cDNA clone IMAGE:3961768 5, mRNA sequence	19.3	Down	2.38E-05
55	AW021686	df26h11.y1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2484717 5, mRNA sequence	18.86	Down	2.59E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	BX109483	BX109483 NCI_CGAP_Ov23 Homo sapiens cDNA clone IMAGp998C165481 ; IMAGE:2216391, mRNA sequence	18.74	Down	9.84E-05
10					
	NM_001463	Homo sapiens frizzled-related protein (FRZB), mRNA	18.63	Down	2.05E-04
	AK128325	Homo sapiens cDNA FLJ46467 fis, clone THYMU3022668	18.11	Down	1.78E-05
15					
	AK096661	Homo sapiens cDNA FLJ39342 fis, clone OCBBF2018873	17.95	Down	5.14E-05
	NM_002531	Homo sapiens neurotensin receptor 1 (high affinity) (NTSR1), mRNA,	17.64	Down	5.02E-05
20					
	NM_031908	Homo sapiens C1q and tumor necrosis factor related protein 2 (C1QTNF2), mRNA	17.61	Down	1.23E-04
	AI085016	ow88e06.s1 Soares_fetal_liver_spleen_1NFLS_S1 Homo sapiens cDNA clone IMAGE:1653922 3, mRNA sequence	17.36	Down	4.39E-05
25					
	AK095791	Homo sapiens cDNA FLJ38472 fis, clone FEBRA2022148	17.31	Down	2.76E-04
	NM_020927	Homo sapiens KIAA1576 protein (KIAA1576), mRNA	17	Down	2.13E-05
	NM_024600	Homo sapiens chromosome 16 open reading frame 30 (C16orf30), mRNA	16.96	Down	6.00E-04
30					
	NM_012449	Homo sapiens six transmembrane epithelial antigen of the prostate (STEAP), mRNA	16.92	Down	5.06E-05
	NM_004787	Homo sapiens slit homolog 2 (Drosophila) (SLIT2), mRNA	16.69	Down	1.78E-05
	N95448	zb81e11.s1 Soares_senescent_fibroblasts_NbHSF Homo sapiens cDNA clone IMAGE:310028 3, mRNA sequence	16.55	Down	1.72E-05
35					
	NM_004657	Homo sapiens serum deprivation response (phosphatidylserine binding protein) (SDPR), mRNA	16.4	Down	1.60E-05
40					
	BC046364	Homo sapiens flavoprotein oxidoreductase MICAL3, mRNA (cDNA clone IMAGE:5737121), with apparent retained intron	16.35	Down	1.15E-05
	NM_053044	Homo sapiens serine protease HTRA3 (HTRA3), mRNA	16.29	Down	3.77E-05
	BQ011545	UI-1-BC1p-asi-a-02-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1p-asi-a-02-0-UI 3, mRNA sequence	16.22	Down	5.21E-04
45					
	BC017939	Homo sapiens, clone IMAGE:4275711, mRNA, partial cds	15.98	Down	5.83E-05
	NM_007289	Homo sapiens membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10) (MME), transcript variant 2b, mRNA	15.83	Down	3.04E-05
50					
	CA437861	UI-H-DHO-aur-k-12-0-UI.s1 NCI_CGAP_DHO Homo sapiens cDNA clone UI-H-DHO-aur-k-12-0-UI 3, mRNA sequence	15.77	Down	2.56E-04
55					
	NM_006182	Homo sapiens discoidin domain receptor family, member 2 (DDR2), mRNA	15.71	Down	4.79E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
10	NM_145239	Homo sapiens similar to lymphocyte antigen 6 complex, locus G5B; G5b protein; open reading frame 31 (LOC112476), mRNA	15.64	Down	3.37E-05
	H15096	ym29e11.r1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:49250 5, mRNA sequence	15.58	Down	2.20E-04
15	NM_002851	Homo sapiens protein tyrosine phosphatase, receptor- type, Z polypeptide 1 (PTPRZ1), mRNA	15.57	Down	1.90E-04
	NM_014217	Homo sapiens potassium channel, subfamily K, member 2 (KCNK2), mRNA	15.13	Down	1.31E-04
	NM_000963	Homo sapiens prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2), mRNA	14.94	Down	1.88E-05
20	BX115659	BX115659 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGp998C204119 ; IMAGE:1623883, mRNA sequence	14.81	Down	2.66E-04
25	NM_002729	Homo sapiens hematopoietically expressed homeobox (HHEX), mRNA	14.78	Down	1.09E-03
	A1082087	oz52h09.x1 Soares_senescent_fibroblasts_NbHSF Homo sapiens cDNA clone IMAGE:1679009 3, mRNA sequence	14.73	Down	3.66E-05
30	NM_018431	Homo sapiens docking protein 5 (DOK5), transcript variant 1, mRNA	14.67	Down	8.15E-04
	NM_004791	Homo sapiens integrin, beta-like 1 (with EGF-like repeat domains) (ITGBL1), mRNA	14.42	Down	2.18E-05
	AK021531	Homo sapiens cDNA FLJ11469 fis, clone HEMBA1001658	14.36	Down	5.21E-04
35	NM_057179	Homo sapiens twist homolog 2 (Drosophila) (TWIST2), mRNA	14.27	Down	1.25E-05
	NM_003619	Homo sapiens protease, serine, 12 (neurotrypsin, motopsin) (PRSS12), mRNA	14.1	Down	1.78E-05
40	NM_002518	Homo sapiens neuronal PAS domain protein 2 (NPAS2), mRNA	14.05	Down	5.49E-05
	NM_000396	Homo sapiens cathepsin K (pseudodysostosis) (CTSK), mRNA	13.81	Down	2.31 E-04
45	NM_016206	Homo sapiens colon carcinoma related protein (FLJ38507), mRNA	13.76	Down	1.81 E-04
	NM_021643	Homo sapiens tribbles homolog 2 (Drosophila) (TRIB2), mRNA	13.62	Down	8.39E-06
	NM_000955	Homo sapiens prostaglandin E receptor 1 (subtype EP1 42kDa) (PTGER1), MRNA,	13.52	Down	1.70E-05
50	NM_058187	Homo sapiens chromosome 21 open reading frame 63 (C21orf63), mRNA	13.44	Down	1.53E-04
	CN479391	UI-H-DF1-aug-f-02-0-UI.s9 NCI_CGAP_DF1 Homo sapiens cDNA clone UI-H-DF1-aug-f-02-0-UI 3, mRNA sequence	13.26	Down	8.65E-05
55	NM_006209	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 2 (autotaxin) (ENPP2), mRNA	13.21	Down	3.28E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_005584	Homo sapiens mab-21-like 1 (C. elegans) (MAB21L1), mRNA	12.73	Down	1.01E-04
10	BM468332	AGENCOURT 6432296 NIH_MGC_71 Homo sapiens cDNA clone IMAGE:5535773 5, mRNA sequence	12.64	Down	2.13E-05
	NM_020353	Homo sapiens phospholipid scramblase 4 (PLSCR4), mRNA	12.39	Down	2.08E-04
15	NM_031302	Homo sapiens glycosyltransferase (LOC83468), mRNA	12.39	Down	1.88E-03
	NM_005308	Homo sapiens G protein-coupled receptor kinase 5 (GRK5), mRNA	12.28	Down	1.18E-04
	NM_032883	Homo sapiens chromosome 20 open reading frame 100 (C20orf100), mRNA	12.17	Down	1.78E-05
20	NM_000110	Homo sapiens dihydropyrimidine dehydrogenase (DPYD), mRNA	12.16	Down	2.75E-05
	BU536871	AGENCOURT_10224340 NIH-MGC-141 Homo sapiens cDNA clone IMAGE:6565454 5, mRNA sequence	12.05	Down	4.21E-05
25	NM_016270	Homo sapiens Kruppel-like factor 2 (lung) (KLF2), mRNA	11.98	Down	1.94E-03
	NM_005110	Homo sapiens glutamine-fructose-6-phosphate transaminase 2 (GFPT2), mRNA	11.92	Down	2.93E-05
30	AK130711	Homo sapiens cDNA FLJ27201 fis, clone SYN03133	11.81	Down	5.02E-05
	BM991890	UI-H-DF1-auk-h-02-0-UI.s1 NCI_CGAP_DF1 Homo sapiens cDNA clone IMAGE:5870641 3, mRNA sequence	11.78	Down	2.03E-05
	NM_000304	Homo sapiens peripheral myelin protein 22 (PMP22), transcript variant 1, mRNA	11.77	Down	1.04E-03
35	BM926469	AGENCOURT_6644776 NIH_MGC_122 Homo sapiens cDNA clone IMAGE:5766855 5, mRNA sequence	11.53	Down	7.16E-06
	NM_001424	Homo sapiens epithelial membrane protein 2 (EMP2), mRNA	11.39	Down	5.12E-04
40	AK093762	Homo sapiens cDNA FLJ36443 fis, clone THYMU2012891	11.18	Down	1.34E-04
	NM_006288	Homo sapiens Thy-1 cell surface antigen (THY1), mRNA	11.17	Down	4.26E-04
	NM_006183	Homo sapiens neurotensin (NTS), mRNA	11.14	Down	8.52E-04
45	NM_002522	Homo sapiens neuronal pentraxin I (NPTX1), mRNA	11.08	Down	3.03E-04
	NM_016428	Homo sapiens ABI gene family, member 3 (ABI3), mRNA	11.03	Down	2.83E-05
	NM_001769	Homo sapiens CD9 antigen (p24) (CD9), mRNA	11	Down	2.45E-04
	AL833655	Homo sapiens mRNA; cDNA DKFZp667O0320 (from clone DKFZp667O0320)	10.8	Down	1.73E-05
50	R56121	yg94d04.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:41388 3, mRNA sequence	10.64	Down	1.04E-04
	NM_012445	Homo sapiens spondin 2, extracellular matrix protein (SPON2), mRNA	10.63	Down	1.36E-04
	NM_004811	Homo sapiens leupaxin (LPXN), mRNA	10.6	Down	2.09E-05
55	NM_002977	Homo sapiens sodium channel, voltage-gated, type IX, alpha (SCN9A), mRNA	10.51	Down	2.09E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	BI598031	603248155F1 NIH_MGC_96 Homo sapiens cDNA clone IMAGE:5300149 5, mRNA sequence	10.5	Down	3.34E-05
10	NM_173553	Homo sapiens hypothetical protein FLJ25801 (FLJ25801), mRNA,	10.49	Down	2.32E-05
	NM_001311	Homo sapiens cysteine-rich protein 1 (intestinal) (CRIP1), mRNA	10.31	Down	2.73E-05
15	NM_015916	Homo sapiens family with sequence similarity 26, member B (FAM26B), mRNA	10.28	Down	1.59E-04
	NM_005595	Homo sapiens nuclear factor I/A (NFIA), mRNA	10.12	Down	7.43E-04
	NM_182728	Homo sapiens solute carrier family 7 (cationic amino acid transporter, y+ system), member 8 (SLC7A8), transcript variant 2, mRNA	9.98	Down	6.78E-05
20					
	NM_152996	Homo sapiens sialyltransferase 7 ((alpha-N-acetylneuraminyl-2,3-beta-galactosyl-1,3)-N-acetyl galactosaminide alpha-2,6-sialyltransferase) C (SIAT7C), mRNA	9.95	Down	1.25E-05
25	AK091713	Homo sapiens cDNA FLJ34394 fis, clone HCHON2000676	9.92	Down	2.31E-05
	AK022877	Homo sapiens cDNA FLJ12815 fis, clone NT2RP2002546	9.89	Down	3.15E-05
30	NM_194250	Homo sapiens similar to C630007C17Rik protein (LOC91752), mRNA	9.84	Down	7.08E-05
	BC039450	Homo sapiens, clone IMAGE:5311619, mRNA	9.77	Down	2.60E-04
	NM_031894	Homo sapiens ferritin, heavy polypeptide-like 17 (FTHL17), mRNA	9.76	Down	5.99E-04
35	BI561641	603256058F1 NIH_MGC_97 Homo sapiens cDNA clone IMAGE:5298374 5, mRNA sequence	9.68	Down	1.96E-03
	BC028245	Homo sapiens, Similar to hypothetical gene LOC130797, clone IMAGE:5395354, mRNA	9.68	Down	1.88E-05
	NM_031957	Homo sapiens keratin associated protein 1-5 (KRTAP1-5), mRNA	9.66	Down	2.31E-05
40	AA947461	ok20f03.s1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:1508381 3, mRNA sequence	9.56	Down	4.34E-05
45	AK127309	Homo sapiens cDNA FLJ45377 fis, clone BRHIP3019956	9.55	Down	1.17E-04
	NM_016247	Homo sapiens interphotoreceptor matrix proteoglycan 2 (IMPG2), mRNA	9.51	Down	2.09E-05
	NM_005574	Homo sapiens LIM domain only 2 (rhombotin-like 1) (LMO2), mRNA	9.46	Down	4.10E-05
50	NM_004369	Homo sapiens collagen, type VI, alpha 3 (COL6A3), transcript variant 1, mRNA	9.36	Down	2.14E-05
	NM_003973	Homo sapiens ribosomal protein L14 (RPL14), mRNA	9.3	Down	6.77E-05
	NM_000459	Homo sapiens TEK tyrosine kinase, endothelial (venous malformations, multiple cutaneous and mucosal) (TEK), mRNA	9.25	Down	3.10E-04
55					
	NM_005397	Homo sapiens podocalyxin-like (PODXL), mRNA	9.06	Down	6.52E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
10	BU675964	UI-CF-DU1-aaf-b-24-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaf-b-24-0-UI 3, mRNA sequence	9	Down	6.82E-05
	AI640484	wa27f01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299321 3, mRNA sequence	8.92	Down	8.54E-05
15	NM_030781	Homo sapiens collectin sub-family member 12 (COLEC12), transcript variant II, mRNA	8.54	Down	8.65E-05
	NM_001998	Homo sapiens fibulin 2 (FBLN2), transcript variant 2, mRNA	8.53	Down	4.60E-04
	BC039369	Homo sapiens, clone IMAGE:5271073, mRNA, partial cds	8.51	Down	4.85E-04
20	AA436084	zu03a02.r1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:730730 5 similar to contains element PTR5 PTR5 repetitive element ;, mRNA sequence	8.51	Down	8.23E-05
25	NM_015192	Homo sapiens phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 1, mRNA	8.48	Down	2.34E-04
	NM_024420	Homo sapiens phospholipase A2, group IVA (cytosolic, calcium-dependent) (PLA2G4A), mRNA	8.48	Down	2.23E-03
30	BX095887	BX095887 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGp998G124121 ; IMAGE:1624739, mRNA sequence	8.42	Down	7.38E-05
	NM_006417	Homo sapiens interferon-induced protein 44 (IFI44), mRNA	8.37	Down	2.46E-04
35	AK022033	Homo sapiens cDNA FLJ11971 fis, clone HEMBB1001208	8.35	Down	2.06E-04
	N69782	yz60b07.s1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:287413 3, mRNA sequence	8.02	Down	2.78E-04
	NM_023003	Homo sapiens transmembrane 6 superfamily member 1 (TM6SF1), MRNA,	7.93	Down	1.20E-04
40	BC036034	Homo sapiens endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2, transcript variant 2, mRNA (cDNA clone MGC:33157 IMAGE:5272431), complete cds	7.86	Down	2.31E-05
45	NM_005602	Homo sapiens claudin 11 (oligodendrocyte transmembrane protein) (CLDN11), mRNA	7.84	Down	2.43E-02
	BC039676	Homo sapiens, clone IMAGE:5173389, mRNA	7.74	Down	3.04E-05
	BC030692	Homo sapiens ELAV (embryonic lethal, abnormal vision, Drosophila)-like 2 (Hu antigen B), mRNA (cDNA clone MGC:26319 IMAGE:4826082), complete cds	7.72	Down	3.59E-05
50	NM_015278	Homo sapiens SAM and SH3 domain containing 1 (SASH1), mRNA	7.58	Down	1.38E-04
	NM_031283	Homo sapiens transcription factor 7-like 1 (T-cell specific, HMG-box) (TCF7L1), mRNA	7.57	Down	4.12E-05
55	BC042378	Homo sapiens, clone IMAGE:5277693, mRNA	7.56	Down	9.91E-04
	NM_013363	Homo sapiens procollagen C-endopeptidase enhancer 2 (PCOLCE2), mRNA	7.49	Down	5.18E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_005923	Homo sapiens mitogen-activated protein kinase kinase 5 (MAP3K5), mRNA	7.48	Down	2.05E-04
10	R53688	yg84h04.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:40175 5, mRNA sequence	7.43	Down	2.31E-05
	AI216469	qh07h10.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE: 1844035 3, mRNA sequence	7.28	Down	4.92E-05
15	NM_014331	Homo sapiens solute carrier family 7, (cationic amino acid transporter, y+ system) member 11 (SLC7A11), mRNA	7.27	Down	2.76E-03
	NM_138801	Homo sapiens galactose mutarotase (aldose 1-epimerase) (GALM), mRNA	7.24	Down	5.50E-05
20	AL049443	Homo sapiens mRNA; cDNA DKFZp586N2020 (from clone DKFZp586N2020)	7.22	Down	2.83E-04
	NM_005328	Homo sapiens hyaluronan synthase 2 (HAS2), mRNA	7.17	Down	3.96E-03
	NM_006072	Homo sapiens chemokine (C-C motif) ligand 26 (CCL26), mRNA	7.12	Down	6.86E-05
25	NM_020987	Homo sapiens ankyrin 3, node of Ranvier (ankyrin G) (ANK3), transcript variant 1, mRNA	7.09	Down	7.31E-05
	BX647541	Homo sapiens mRNA; cDNA DKFZp686P0492 (from clone DKFZp686P0492)	7.08	Down	4.47E-05
30	AI417595	tg79h09.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2115041 3, mRNA sequence	7.07	Down	7.52E-04
	AA158235	zo76b02.s1 Stratagene pancreas (#937208) Homo sapiens cDNA clone IMAGE:592779 3, mRNA sequence	7.06	Down	1.52E-04
	NM_005583	Homo sapiens lymphoblastic leukemia derived sequence 1 (LYL1), MRNA,	7.03	Down	4.69E-05
35	BU570253	AGENCOURT_10401698 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6618451 5, mRNA sequence	7	Down	2.35E-04
	BM675371	UI-E-EJ0-aht-a-07-0-UI.s1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aht-a-07-0-UI 3, mRNA sequence	6.95	Down	4.53E-05
40	AK024653	Homo sapiens cDNA: FLJ21000 fis, clone CAE03359	6.93	Down	6.77E-05
	W31037	zb86c06.r1 Soares_senescent_fibroblasts_NbHSF Homo sapiens cDNA clone IMAGE:310474 5, mRNA sequence	6.91	Down	4.95E-05
45	AL831835	Homo sapiens mRNA; cDNA DKFZp547A0515 (from clone DKFZp547A0515)	6.9	Down	1.53E-03
	NM_001849	Homo sapiens collagen, type VI, alpha 2 (COL6A2), transcript variant 2C2, mRNA	6.85	Down	3.34E-05
	NM_007084	Homo sapiens SRY (sex determining region Y)-box 21 (SOX21), mRNA	6.85	Down	3.88E-02
50	NM_025107	Homo sapiens myc target 1 (MYCT1), MRNA,	6.82	Down	6.89E-05
	AA393981	zt58a03.r1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:726508 5, mRNA sequence	6.82	Down	4.58E-03
	NM_170744	Homo sapiens unc-5 homolog B (C. elegans) (UNC5B), mRNA	6.82	Down	9.92E-04
55	AL133118	Homo sapiens mRNA; cDNA DKFZp586N0121 (from clone DKFZp586N0121)	6.78	Down	1.42E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	AK090603	Homo sapiens cDNA FLJ33284 fis, clone ASTRO2009458	6.76	Down	1.36E-04
10	NM_002203	Homo sapiens integrin, alpha 2 (CD49B, alpha 2 subunit of VLA-2 receptor) (ITGA2), mRNA	6.74	Down	4.93E-05
	NM_003713	Homo sapiens phosphatidic acid phosphatase type 2B (PPAP2B), transcript variant 1, mRNA	6.73	Down	1.06E-04
15	NM_001401	Homo sapiens endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2 (EDG2), transcript variant 1, mRNA	6.69	Down	9.75E-04
	NM_020226	Homo sapiens PR domain containing 8 (PRDM8), mRNA	6.68	Down	2.31E-04
20	N71963	yz95e03.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:290812 3 similar to contains Alu repetitive element;, mRNA sequence	6.63	Down	2.47E-04
	AK127536	Homo sapiens cDNA FLJ45629 fis, clone CHONS2000797, highly similar to T-box transcription factor TBX15	6.6	Down	1.09E-04
25	NM_006206	Homo sapiens platelet-derived growth factor receptor, alpha polypeptide (PDGFRA), mRNA	6.55	Down	6.30E-04
	NM_016559	Homo sapiens peroxisomal biogenesis factor 5-like (PEX5L), mRNA	6.55	Down	6.77E-04
30	NM_012307	Homo sapiens erythrocyte membrane protein band 4.1- like 3 (EPB41L3), mRNA	6.51	Down	9.62E-04
	R34294	yh84b01.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:136393 3, mRNA sequence	6.49	Down	2.78E-05
35	N36786	yy34e08.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273158 3 similar to contains element MSR1 repetitive element ;, mRNA sequence	6.49	Down	6.12E-05
	BX113851	BX113851 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGp998G054116 ; IMAGE:1622812, mRNA sequence	6.44	Down	8.41E-05
40	BG484952	602503960F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4617278 5, mRNA sequence	6.43	Down	9.06E-05
	NM_005434	Homo sapiens BENE protein (BENE), mRNA	6.4	Down	1.64E-05
45	BX106577	BX106577 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGp998H131854 ; IMAGE:754236, mRNA sequence	6.39	Down	4.12E-05
	NM_144617	Homo sapiens heat shock protein, alpha-crystallin- related, B6 (HSPB6), mRNA	6.37	Down	2.83E-05
50	AK093529	Homo sapiens cDNA FLJ36210 fis, clone THYMU2000155	6.29	Down	1.52E-04
	NM_022731	Homo sapiens nuclear ubiquitous casein kinase and cyclin-dependent kinase substrate (NUCKS), mRNA	6.27	Down	9.06E-05
	NM_018286	Homo sapiens hypothetical protein FLJ10970 (FLJ10970), mRNA	6.26	Down	1.08E-03
55	NM_002290	Homo sapiens laminin, alpha 4 (LAMA4), mRNA	6.19	Down	6.97E-04
	NM_015184	Homo sapiens phospholipase C-like 2 (PLCL2), mRNA	6.18	Down	5.91E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
	NM_017577	Homo sapiens hypothetical protein DKFZp434C0328 (DKFZp434C0328), mRNA	6.17	Down	3.60E-05
10	BQ350534	RC1-HT0256-120400-019-d06 HT0256 Homo sapiens cDNA, mRNA sequence	6.17	Down	2.04E-04
	NM_004934	Homo sapiens cadherin 18, type 2 (CDH18), mRNA	6.17	Down	3.03E-05
	NM_024769	Homo sapiens adipocyte-specific adhesion molecule (ASAM), mRNA	6.15	Down	2.41E-04
15	NM_197941	Homo sapiens similar to ADAMTS-10 precursor (A disintegrin and metalloproteinase with thrombospondin motifs 10) (ADAM-TS 10) (ADAM-TS10) (LOC345667), mRNA	6.14	Down	1.06E-04
20	BG118019	602351269F1 NIH_MGC_90 Homo sapiens cDNA clone IMAGE:4446065 5, mRNA sequence	6.14	Down	5.02E-05
	NM_005531	Homo sapiens interferon, gamma-inducible protein 16 (IF116), mRNA	6.13	Down	4.46E-04
25	NM_005613	Homo sapiens regulator of G-protein signalling 4 (RGS4), mRNA	6.08	Down	1.04E-04
	NM_002166	Homo sapiens inhibitor of DNA binding 2, dominant negative helix-loop-helix protein (ID2), mRNA	6.07	Down	1.16E-03
	BC015720	Homo sapiens, clone IMAGE:3909165, mRNA	6.04	Down	1.15E-05
30	BF508005	UI-H-B14-apw-e-06-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3088978 3, mRNA sequence	6.03	Down	1.15E-05
	NM_007101	Homo sapiens sarcosine dehydrogenase (SARDH), mRNA	6.01	Down	4.54E-03
35	BG025371	602276295F1 NIH_MGC_85 Homo sapiens cDNA clone IMAGE:4364351 5, mRNA sequence	5.99	Down	6.94E-05
	NM_152666	Homo sapiens hypothetical protein FLJ40773 (FLJ40773), mRNA	5.99	Down	4.55E-04
40	BM542398	AGENCOURT_6436663 NIH_MGC_72 Homo sapiens cDNA clone IMAGE:5539574 5, mRNA sequence	5.97	Down	3.34E-05
	AK054783	Homo sapiens cDNA FLJ30221 fis, clone BRACE2001742	5.97	Down	2.40E-04
	AI651524	wb06g07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2304924 3, mRNA sequence	5.93	Down	3.22E-04
45	NM_005905	Homo sapiens SMAD, mothers against DPP homolog 9 (Drosophila) (SMAD9), mRNA	5.87	Down	4.76E-04
	BG818762	602779092F2 NCI_CGAP_Brn67 Homo sapiens cDNA clone IMAGE:4914502 5, mRNA sequence	5.84	Down	1.07E-04
50	NM_002487	Homo sapiens necdin homolog (mouse) (NDN), mRNA	5.83	Down	2.72E-05
	NM_000167	Homo sapiens glycerol kinase (GK), transcript variant 2, mRNA	5.8	Down	5.10E-03
	BC070147	Homo sapiens cDNA clone IMAGE:4672631, containing frame-shift errors	5.74	Down	7.92E-04
55	NM_001711	Homo sapiens biglycan (BGN), mRNA	5.71	Down	4.13E-05
	NM_000407	Homo sapiens glycoprotein Ib (platelet), beta polypeptide (GP1BB), mRNA	5.69	Down	3.03E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p- value
5					
10	N28431	yx35c03.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:263716 5 similar to SP:PIR:S32603 S32603 collagen alpha 1 (VI) chain - mouse ;, mRNA sequence	5.69	Down	6.03E-04
15	NM_002615	Homo sapiens serine (or cysteine) proteinase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 (SERPINF1), mRNA,	5.66	Down	2.02E-03
	H18367	yn49d06.r1 Soares adult brain N2b5HB55Y Homo sapiens cDNA clone IMAGE:171755 3, mRNA sequence	5.66	Down	4.41E-04
	NM_018455	Homo sapiens uncharacterized bone marrow protein BM039 (BM039), mRNA	5.64	Down	3.03E-04
20	H24359	ym56b03.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:52294 3, mRNA sequence	5.61	Down	2.57E-04
	BU078105	im64b02.y1 HR85 islet Homo sapiens cDNA clone IMAGE:6039866 5, mRNA sequence	5.58	Down	3.60E-04
25	NM_022576	Homo sapiens phosducin (PDC), transcript variant 2, mRNA	5.55	Down	9.39E-05
	NM_033196	Homo sapiens hypothetical ZNF-like protein (LOC91120), mRNA	5.54	Down	1.49E-04
	BC035066	Homo sapiens, clone IMAGE:5259543, mRNA	5.54	Down	1.65E-04
30	R14261	yf79d05.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:28642 5, mRNA sequence	5.51	Down	2.36E-04
	BC042140	Homo sapiens, clone IMAGE:5932306, mRNA	5.48	Down	5.05E-05
	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), MRNA,	5.48	Down	1.02E-05
35	NM_182746	Homo sapiens MCM4 minichromosome maintenance deficient 4 (S. cerevisiae) (MCM4), transcript variant 2, mRNA	5.48	Down	9.96E-04
	BG573337	602595107F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4724521 5, mRNA sequence	5.43	Down	3.72E-04
40	NM_194250	Homo sapiens similar to C630007C17Rik protein (LOC91752), mRNA	5.43	Down	3.30E-04
	NM_015430	Homo sapiens regeneration associated muscle protease (DKFZP586H2123), transcript variant 1, mRNA	5.43	Down	3.49E-03
45	NM_006868	Homo sapiens RAB31, member RAS oncogene family (RAB31), MRNA,	5.41	Down	2.11E-05
	BI494495	df110f11.w1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2539149 3, mRNA sequence	5.4	Down	2.64E-03
50	AW025556	wu97g10.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:2528034 3, mRNA	5.36	Down	1.89E-04
	AI793182	sequence qz36a11.x5 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2028956 3 similar to contains L1.t3 L1 repetitive element ;, mRNA sequence	5.34	Down	8.09E-04
55	NM_006208	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 1 (ENPP1), mRNA	5.34	Down	2.36E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p-value
5					
	C02345	HUMGS0007544 Human adult (K.Okubo) Homo sapiens cDNA, mRNA sequence	5.33	Down	3.45E-04
10	NM_032297	Homo sapiens hypothetical protein DKFZp761D112 (DKFZp761D112), mRNA	5.32	Down	3.38E-04
	BM724062	UI-E-E01-aiy-a-22-0-UI.r1 UI-E-E01 Homo sapiens cDNA clone UI-E-E01-aiy-a-22-0-UI 5, mRNA sequence	5.25	Down	2.96E-03
15	NM_176891	Homo sapiens interferon epsilon 1 (IFNE1), mRNA	5.22	Down	2.67E-04
	BX648959	Homo sapiens mRNA; cDNA DKFZp686N2348 (from clone DKFZp686N2348)	5.22	Down	4.12E-05
	CA771688	io81c08.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6132854 3, mRNA sequence	5.22	Down	3.35E-03
20	BC034315	Homo sapiens hypothetical protein LOC90529, mRNA (cDNA clone IMAGE:4827425), containing frame-shift errors	5.22	Down	1.90E-04
	W88428	zh72g09.s1 Soares_fetal_liver_spleen_1NFLSS1 Homo sapiens cDNA clone IMAGE:417664 3, mRNA sequence	5.22	Down	5.95E-04
25	AI468014	tj84g05.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:2148248 3 similar to contains element TAR1 repetitive element ;, mRNA sequence	5.21	Down	9.55E-04
30	NM_197941	Homo sapiens similar to ADAMTS-10 precursor (A disintegrin and metalloproteinase with thrombospondin motifs 10) (ADAM-TS 10) (ADAM-TS10) (LOC345667), mRNA	5.2	Down	8.12E-04
35	U10991	Human G2 protein mRNA, partial cds	5.19	Down	7.92E-05
	AI888390	wn30g10.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2447010 3 similar to contains element MER8 repetitive element ;, mRNA sequence	5.19	Down	2.07E-04
40	NM_012411	Homo sapiens protein tyrosine phosphatase, non-receptor type 22 (lymphoid) (PTPN22), transcript variant 2, mRNA	5.19	Down	3.62E-04
	NM_030806	Homo sapiens chromosome 1 open reading frame 21 (C1orf21), mRNA	5.17	Down	1.88E-05
45	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	5.17	Down	8.65E-05
	NM_153014	Homo sapiens hypothetical protein FLJ30634 (FLJ30634), mRNA	5.13	Down	1.32E-03
	AB020691	Homo sapiens mRNA for KIAA0884 protein, partial cds	5.13	Down	3.40E-05
50	NM_020211	Homo sapiens RGM domain family, member A (RGMA), mRNA	5.1	Down	1.51E-04
	BG215747	RST35420 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	5.1	Down	1.30E-04
	NM_002317	Homo sapiens lysyl oxidase (LOX), mRNA	5.09	Down	1.28E-04
55	NM_003608	Homo sapiens G protein-coupled receptor 65 (GPR65), mRNA	5.07	Down	7.41E-04
	NM_031305	Homo sapiens Rho GTPase activating protein 24 (ARHGAP24), mRNA	5.05	Down	1.10E-04

EP 1 780 267 B1

(continued)

Gene Identifier	Gene Name	Average fold change in Epithelial versus fibroblast cells	Direction	adj. p-value
AK124391	Homo sapiens cDNA FLJ42400 fis, clone ASTRO2003581	5.03	Down	9.74E-05
TM_005613	Homo sapiens regulator of G-protein signalling 4 (RGS4), mRNA	5.02	Down	7.55E-05
NM_054027	Homo sapiens ankylosis, progressive homolog (mouse) (ANKH), mRNA	5	Down	2.64E-05

TABLE VII C: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN EPITHELIAL VERSUS AF-I CELLS

Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
AK091336	Homo sapiens cDNA FLJ34017 fis, clone FCBBF2002626	97.64	UP	6.01E-03
NM_004098	Homo sapiens empty spiracles homolog 2 (Drosophila) (EMX2), mRNA	83.49	UP	8.10E-03
AV702977	AV702977 ADB Homo sapiens cDNA clone ADBCVD08 5, mRNA sequence	82.96	UP	3.74E-03
BE877764	601486331 F1 NIH_MGC_69 Homo sapiens cDNA clone IMAGE:3888943 5, mRNA sequence	60.78	UP	2.79E-03
AK075003	Homo sapiens cDNA FLJ90522 fis, clone NT2RP4000108, highly similar to Human gene for neurofilament subunit NF-L	48.1	UP	1.02E-02
NM_014358	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	48.04	UP	1.99E-04
AK092401	Homo sapiens cDNA FLJ35082 fis, clone PLACE6005351	40.58	UP	1.77E-03
NM_007029	Homo sapiens stathmin-like 2 (STMN2), mRNA	38.31	UP	1.31E-02
AW300043	xs45a09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2772568 3, mRNA sequence	25.26	UP	1.35E-02
NM_006158	Homo sapiens neurofilament, light polypeptide 68kDa (NEFL), mRNA	23.08	UP	1.76E-03
NM_003222	Homo sapiens transcription factor AP-2 gamma (activating enhancer binding protein 2 gamma) (TFAP2C), mRNA	22.98	UP	1.59E-02
AK023647	Homo sapiens cDNA FLJ13585 fis, clone PLACE1009150	22.77	UP	7.48E-04
BC044843	Homo sapiens hypothetical protein LOC339535, mRNA (cDNA clone IMAGE:5186761), partial cds	21.74	UP	4.67E-03
BQ003501	UI-H-E11-azd-p-06-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847413 3, mRNA sequence	20.08	UP	9.54E-04
AA075748	zm89e04.r1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone IMAGE:545118 5, mRNA sequence	19.94	UP	7.10E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	BF000009	7h15g04.x1 NCI_CGAP_Co16 Homo sapiens cDNA clone IMAGE:3316086 3, mRNA sequence	19.92	UP	7.67E-03
10	NM_004867	Homo sapiens integral membrane protein 2A (ITM2A), mRNA	19.86	UP	2.62E-02
	H85497	yv88b07.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:249781 5, mRNA sequence	19.75	UP	7.71E-05
15	BM713465	UI-E-EJ0-aho-m-22-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aho-m-22-0-UI 5, mRNA sequence	18.57	UP	1.68E-02
	BU569937	AGENCOURT_10399817 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6618011 5, mRNA sequence	18.3	UP	1.68E-02
20	NM_032782	Homo sapiens hepatitis A virus cellular receptor 2 (HAVCR2), mRNA	18.05	UP	2.97E-03
	NM_013387	Homo sapiens ubiquinol-cytochrome c reductase complex (7.2 kD) (HSPC051), transcript variant 1, mRNA	17.68	UP	1.30E-04
	BM988338	UI-H-DH0-asd-f-10-0-UI.s1 NCI_CGAP_DHO Homo sapiens cDNA clone IMAGE:5857545 3, mRNA sequence	17.23	UP	9.27E-03
25	BQ025821	UI-1-BB1p-aye-f-10-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-aye-f-10-0-UI 3, mRNA sequence	15.46	UP	1.08E-02
30	BF515657	UI-H-BW1-anu-e-05-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3083601 3, mRNA sequence	15.17	UP	8.95E-03
	NM_001977	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	14.59	UP	5.24E-05
35	NM_197955	Homo sapiens normal mucosa of esophagus specific 1 (NMES1), transcript variant 1, mRNA	14.33	UP	2.54E-03
	NM_180991	Homo sapiens solute carrier organic anion transporter family, member 4C1 (SLC04C1), mRNA	13.46	UP	2.20E-02
	R44402	yg37a01.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:34639 3 similar to contains MER35 repetitive element ;, mRNA sequence	13.17	UP	1.56E-04
40	CF137545	UI-HF-BN0-ane-d-05-0-UI.r1 NIH_MGC_50 Homo sapiens cDNA clone IMAGE:3092384 5, mRNA sequence	12.96	UP	6.05E-03
45	NM_016269	Homo sapiens lymphoid enhancer-binding factor 1 (LEF1), mRNA	12.69	UP	7.27E-03
	NM_001680	Homo sapiens FXYP domain containing ion transport regulator 2 (FXYP2), transcript variant a, mRNA	12.65	UP	2.38E-03
50	NM_003063	Homo sapiens sarcolipin (SLN), mRNA	12.54	UP	3.04E-02
	NM_012204	Homo sapiens general transcription factor IIIC, polypeptide 4, 90kDa (GTF3C4), mRNA	12.51	UP	3.10E-02
	NM_003810	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	12.4	UP	9.19E-05
55	NM_004591	Homo sapiens chemokine (C-C motif) ligand 20 (CCL20), mRNA	11.84	UP	3.08E-02
	NM_020927	Homo sapiens KIAA1576 protein (KIAA1576), mRNA	11.77	UP	6.78E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	NM_014751	Homo sapiens metastasis suppressor 1 (MTSS1), mRNA	11.5	UP	6.60E-05
	NM_031311	Homo sapiens carboxypeptidase, vitellogenic-like (CPVL), transcript variant 1, mRNA	10.36	UP	8.15E-03
10	NM_005382	Homo sapiens neurofilament 3 (150kDa medium) (NEF3), mRNA	10.12	UP	4.45E-02
	NM_033554	Homo sapiens major histocompatibility complex, class II, DP alpha 1 (HLA-DPA1), mRNA	9.84	UP	4.35E-02
15	AI632692	wa33b05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299857 3, mRNA sequence	9.7	UP	3.01E-02
	NM_030923	Homo sapiens hypothetical protein DKFZp566N034 (DKFZP566N034), mRNA	9.51	UP	2.72E-02
20	NM_152550	Homo sapiens SH3 domain containing ring finger 2 (SH3RF2), mRNA	9.31	UP	4.41E-02
	AW445209	UI-H-BI3-akc-g-11-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2733908 3, mRNA sequence	9.27	UP	1.37E-03
25	AB011095	Homo sapiens mRNA for KIAA0523 protein, partial cds	8.77	UP	1.82E-04
	NM_194463	Homo sapiens ring finger protein 128 (RNF128), transcript variant 1, mRNA	8.75	UP	2.95E-03
	NM_013427	Homo sapiens Rho GTPase activating protein 6 (ARHGAP6), transcript variant 1, mRNA	8.73	UP	7.38E-04
30	NM_012105	Homo sapiens beta-site APP-cleaving enzyme 2 (BACE2), transcript variant a, mRNA	8.67	UP	1.49E-04
	NM_000640	Homo sapiens interleukin 13 receptor, alpha 2 (IL13RA2), mRNA	8.29	UP	4.39E-02
	NM_001977	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	8.16	UP	6.01E-03
35	T56535	yb33g07.r1 Stratagene fetal spleen (#937205) Homo sapiens cDNA clone IMAGE:73020 5, mRNA sequence	8.08	UP	1.24E-04
	NM_022440	Homo sapiens mal, T-cell differentiation protein (MAL), transcript variant d, mRNA	8.01	UP	4.29E-02
40	BM711923	UI-E-CL1-afc-m-16-0-UI.r1 UI-E-CL1 Homo sapiens cDNA clone UI-E-CL1-afc-m-16-0-UI 5, mRNA sequence	7.87	UP	8.56E-04
	NM_005181	Homo sapiens carbonic anhydrase III, muscle specific (CA3), mRNA	7.69	UP	1.46E-02
45	NM_030781	Homo sapiens collectin sub-family member 12 (COLEC12), transcript variant II, mRNA	7.56	UP	8.07E-04
	NM_001305	Homo sapiens claudin 4 (CLDN4), mRNA	7.41	UP	3.62E-04
	NM_000055	Homo sapiens butyrylcholinesterase (BCHE), mRNA	7.26	UP	4.51E-03
	NM_153183	Homo sapiens nudix (nucleoside diphosphate linked moiety X)-type motif 10 (NUDT10), mRNA	7.23	UP	3.53E-02
50	BI759570	603046987F1 NIH-MGC-116 Homo sapiens cDNA clone IMAGE:5187285 5, mRNA sequence	7.11	UP	3.86E-02
	NM_015559	Homo sapiens SET binding protein 1 (SETBP1), mRNA	6.97	UP	3.67E-02
55	NM_005562	Homo sapiens laminin, gamma 2 (LAMC2), transcript variant 1, mRNA	6.95	UP	2.06E-04
	NM_013430	Homo sapiens gamma-glutamyltransferase 1 (GGT1), transcript variant 3, mRNA	6.9	UP	3.86E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	AA908815	og77h08.s1 NCI_CGAP_Ov8 Homo sapiens cDNA clone IMAGE:1454367 3, mRNA sequence	6.8	UP	3.10E-02
10	NM_022843	Homo sapiens protocadherin 20 (PCDH20), mRNA	6.73	UP	5.39E-03
	NM_018076	Homo sapiens armadillo repeat containing 4 (ARMC4), mRNA	6.71	UP	7.53E-04
	NM_005822	Homo sapiens Down syndrome critical region gene 1-like 1 (DSCR1L1), mRNA	6.67	UP	2.38E-02
15	NM_006393	Homo sapiens nebulin (NEBL), transcript variant 1, mRNA	6.66	UP	4.24E-02
	CA391258	cs13a10.x1 Human Retinal pigment epithelium/choroid cDNA (Unnormalized, unamplified): cs Homo sapiens cDNA clone cs13a10 3, mRNA sequence	6.56	UP	9.19E-05
20	AK026966	Homo sapiens cDNA: FLJ23313 fis, clone HEP11919	6.31	UP	1.30E-04
	NM_020661	Homo sapiens activation-induced cytidine deaminase (AICDA), mRNA	6.12	UP	3.01E-02
	NM_004840	Homo sapiens Rac/Cdc42 guanine nucleotide exchange factor (GEF) 6 (ARHGEF6), mRNA	6.09	UP	1.19E-03
25	BC021684	Homo sapiens, clone IMAGE:3827252, mRNA	6.03	UP	4.71E-03
	NM_032551	Homo sapiens G protein-coupled receptor 54 (GPR54), mRNA	5.96	UP	4.33E-02
	NM_024726	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	5.94	UP	1.84E-04
30	NM_001657	Homo sapiens amphiregulin (schwannoma-derived growth factor) (AREG), mRNA	5.91	UP	1.05E-03
	NM_006033	Homo sapiens lipase, endothelial (LIPG), mRNA	5.9	UP	1.76E-04
	NM_002089	Homo sapiens chemokine (C-X-C motif) ligand 2 (CXCL2), mRNA	5.88	UP	1.13E-04
35	AK074097	Homo sapiens mRNA for FLJ00168 protein	5.87	UP	1.17E-03
	BF509573	UI-H-BI4-apf-b-11-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086949 3, mRNA sequence	5.78	UP	1.40E-04
40	H23441	ym52f11.s1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:51888 3, mRNA sequence	5.77	UP	6.87E-04
	NM_031426	Homo sapiens chromosome 9 open reading frame 58 (C9orf58), transcript variant 1, mRNA	5.63	UP	3.32E-05
45	NM_013410	Homo sapiens adenylate kinase 3 (AK3), nuclear gene encoding mitochondrial protein, transcript variant 2, mRNA	5.54	UP	1.14E-04
	NM_003236	Homo sapiens transforming growth factor, alpha (TGFA), mRNA	5.51	UP	3.09E-02
50	NM_152369	Homo sapiens hypothetical protein MGC45474 (MGC45474), mRNA	5.46	UP	1.02E-04
	NM_153229	Homo sapiens hypothetical protein FLJ33318 (FLJ33318), mRNA	5.4	UP	1.92E-03
	NM_000216	Homo sapiens Kallmann syndrome 1 sequence (KAL1), mRNA	5.23	UP	1.13E-04
55	H89526	yw28b04.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253519 5, mRNA sequence	5.15	UP	9.54E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	AL832916	Homo sapiens mRNA; cDNA DKFZp762I0915 (from clone DKFZp762I0915)	5.12	UP	6.74E-03
10	NM_002899	Homo sapiens retinol binding protein 1, cellular (RBP1), mRNA	5.07	UP	1.11E-03
	CA417015	UI-H-FE0-bbp-d-21-0-UI.s1 NCI_CGAP_FE0 Homo sapiens cDNA clone UI-H-FE0-bbp-d-21-0-UI 3, mRNA sequence	5.07	UP	1.77E-03
15	AW263542	xn80f01.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2700793 3, mRNA sequence	5.07	UP	9.70E-03
	T47612	yb15h03.s1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:71285 3, mRNA sequence	5.02	UP	7.57E-03
20	NM_002402	Homo sapiens mesoderm specific transcript homolog (mouse) (MEST), transcript variant 1, mRNA	83.38	Down	1.30E-04
	NM_001884	Homo sapiens hyaluronan and proteoglycan link protein 1 (HAPLN1), mRNA	71.21	Down	5.97E-05
	NM_006439	Homo sapiens mab-21-like 2 (C. elegans) (MAB21L2), mRNA	70.58	Down	7.65E-05
25	NM_001202	Homo sapiens bone morphogenetic protein 4 (BMP4), transcript variant 1, mRNA	50.48	Down	3.32E-05
	NM_017671	Homo sapiens chromosome 20 open reading frame 42 (C20orf42), mRNA	50.28	Down	4.04E-05
30	AK124396	Homo sapiens cDNA FLJ42405 fis, clone ASTRO3000474	45.55	Down	6.98E-05
	NM_015170	Homo sapiens sulfatase 1 (SULF1), mRNA	38.04	Down	7.58E-05
	BC037316	Homo sapiens, clone IMAGE:5259432, mRNA	34.94	Down	7.08E-05
	NM_001562	Homo sapiens interleukin 18 (interferon-gamma-inducing factor) (IL18), mRNA	34.48	Down	7.58E-05
35	A1469032	ti70a01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2137320 3, mRNA sequence	28.72	Down	4.44E-05
	NM_016307	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	28.57	Down	3.32E-05
40	NM_022475	Homo sapiens hedgehog interacting protein (HHIP), mRNA	27	Down	7.14E-05
	NM_006072	Homo sapiens chemokine (C-C motif) ligand 26 (CCL26), mRNA	25.57	Down	5.83E-05
45	NM_007361	Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	22.81	Down	5.24E-05
	NM_006228	Homo sapiens prepronociceptin (PNOC), mRNA	21.34	Down	4.20E-04
	N95448	zb81e11.s1 Soares_senescent_fibroblasts_NbHSF Homo sapiens cDNA clone IMAGE:310028 3, mRNA sequence	20.92	Down	1.24E-04
50	NM_018242	Homo sapiens hypothetical protein FLJ10847 (FLJ10847), mRNA	20.83	Down	6.60E-05
	N63415	yy60d04.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:277927 3 similar to contains L1.b3 L1 repetitive element ;, mRNA sequence	19.2	Down	5.24E-05
55	BQ001571	UI-H-DH1-awr-i-18-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5893337 3, mRNA sequence	18.75	Down	6.98E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	18.21	Down	5.83E-05
10	BE788763	601475864F1 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:3879014 5, mRNA sequence	17.84	Down	1.13E-04
	BU633163	UI-H-FL1-bgt-n-07-0-UI.s1 NCI-CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgt-n-07-0-UI 3, mRNA sequence	17.41	Down	7.14E-05
15	AI289329	qw28c09.x1 NCI_CGAP_Ut4 Homo sapiens cDNA clone IMAGE:1992400 3 similar to contains L1.b2 L1 repetitive element ;, mRNA sequence	17.1	Down	6.98E-05
	AF318382	Homo sapiens pp9974 mRNA, complete cds	16.45	Down	1.38E-04
20	NM_022350	Homo sapiens leukocyte-derived arginine aminopeptidase (LRAP), mRNA	15.8	Down	1.10E-04
	BI561641	603256058F1 NIH_MGC_97 Homo sapiens cDNA clone IMAGE:5298374 5, mRNA sequence	14.77	Down	1.94E-04
	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	14.55	Down	7.14E-05
25	NM_016588	Homo sapiens neuritin 1 (NRN1), mRNA	14.52	Down	6.98E-05
	CA437861	UI-H-DH0-aur-k-12-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone UI-H-DH0-aur-k-12-0-UI 3, mRNA sequence	14.45	Down	1.21E-04
30	NM_020873	Homo sapiens leucine rich repeat neuronal 1 (LRRN1), mRNA	13.74	Down	6.60E-05
	NM_003385	Homo sapiens visinin-like 1 (VSNL1), mRNA	13.06	Down	7.75E-05
	AI640484	wa27f01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299321 3, mRNA sequence	11.92	Down	5.48E-04
35	AK127309	Homo sapiens cDNA FLJ45377 fis, clone BRHIP3019956	11.87	Down	2.35E-04
	M60502	Human profilaggrin mRNA, 3 end	11.74	Down	7.58E-05
	NM_006475	Homo sapiens periostin, osteoblast specific factor (POSTN), mRNA	11.74	Down	1.19E-04
40	NM_207482	Homo sapiens FLJ44048 protein (FLJ44048), mRNA	11.63	Down	1.45E-04
	NM_000519	Homo sapiens hemoglobin, delta (HBD), mRNA	11.61	Down	3.27E-04
	BU567804	AGENCOURT_10398872 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6614502 5, mRNA sequence	11.59	Down	5.24E-05
	AJ318805	AJ318805 Homo sapiens adipose tissue Homo sapiens cDNA clone 2040, mRNA sequence	11.59	Down	1.76E-04
45	AW451831	UI-H-BI3-alk-e-12-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2737246 3, mRNA sequence	11.57	Down	1.46E-04
	R99527	yq79b11.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:201981 3, mRNA sequence	11.16	Down	1.13E-04
50	NM_031894	Homo sapiens ferritin, heavy polypeptide-like 17 (FTHL17), mRNA	11.12	Down	4.85E-04
	NM_007072	Homo sapiens HERV-H LTR-associating 2 (HHLA2), mRNA	10.94	Down	1.36E-04
55	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	10.91	Down	3.32E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	NM_004574	Homo sapiens peanut-like 2 (Drosophila) (PNUTL2), transcript variant 1, mRNA	10.46	Down	3.48E-04
10	NM_005141	Homo sapiens fibrinogen, B beta polypeptide (FGB), mRNA	10.37	Down	4.85E-04
	AI905628	CM-BT094-050299-147 BT094 Homo sapiens cDNA, mRNA sequence	10.36	Down	1.49E-04
	BG622707	602647476F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4768963 5, mRNA sequence	10.35	Down	1.73E-04
15	W76003	zd58g07.r1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:344892 5, mRNA sequence	10.32	Down	1.58E-04
	H15096	ym29e11.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:49250 5, mRNA sequence	10.23	Down	1.80E-04
20	AI493349	tg70f04.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2114143 3, mRNA sequence	9.84	Down	7.54E-05
	AK093762	Homo sapiens cDNA FLJ36443 fis, clone THYMU2012891	9.76	Down	8.07E-04
25	BU570253	AGENCOURT_10401698 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6618451 5, mRNA sequence	9.71	Down	3.65E-04
	NM_005855	Homo sapiens receptor (calcitonin) activity modifying protein 1 (RAMP1), mRNA	9.33	Down	7.08E-05
	NM_003328	Homo sapiens TXK tyrosine kinase (TXK), mRNA	9.29	Down	1.19E-04
30	NM_003004	Homo sapiens secreted and transmembrane 1 (SECTM1), mRNA	9.23	Down	1.76E-04
	NM_005596	Homo sapiens nuclear factor I/B (NFIB), mRNA	9.23	Down	4.97E-05
	BC039450	Homo sapiens, clone IMAGE:5311619, mRNA	9.07	Down	1.13E-04
	NM_005613	Homo sapiens regulator of G-protein signalling 4 (RGS4), mRNA	8.99	Down	7.65E-05
35	NM_003973	Homo sapiens ribosomal protein L14 (RPL14), mRNA	8.98	Down	6.60E-05
	BM675270	UI-E-EJ0-ahr-j-07-0-UI.s1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahr-j-07-0-UI 3, mRNA sequence	8.96	Down	1.34E-03
40	NM_001311	Homo sapiens cysteine-rich protein 1 (intestinal) (CRIP1), mRNA	8.91	Down	7.14E-05
	NM_000087	Homo sapiens cyclic nucleotide gated channel alpha 1 (CNGA1 mRNA	8.76	Down	1.33E-03
	AW021686	df26h11.y1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2484717 5, mRNA sequence	8.74	Down	4.04E-05
45	NM_002031	Homo sapiens fyn-related kinase (FRK), mRNA	8.67	Down	1.14E-04
	NM_005279	Homo sapiens G protein-coupled receptor 1 (GPR1), mRNA	8.51	Down	4.16E-04
	NM_004460	Homo sapiens fibroblast activation protein, alpha (FAP), mRNA	8.51	Down	7.71E-04
50	BE904671	601498784F1 NIH_MGC_70 Homo sapiens cDNA clone IMAGE:3900717 5, mRNA sequence	8.46	Down	1.73E-04
	NM_000612	Homo sapiens insulin-like growth factor 2 (somatomedin A) (IGF2), mRNA	8.27	Down	1.19E-04
55	AL137488	Homo sapiens mRNA; cDNA DKFZp434N2030 (from clone DKFZp434N2030)	8.23	Down	1.13E-04
	NM_006288	Homo sapiens Thy-1 cell surface antigen (THY1), mRNA	8.18	Down	1.43E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	NM_002837	Homo sapiens protein tyrosine phosphatase, receptor type, B (PTPRB), mRNA	8.15	Down	1.19E-03
10	NM_030899	Homo sapiens zinc finger protein 323 (ZNF323), mRNA	8.08	Down	9.54E-05
	NM_002193	Homo sapiens inhibin, beta B (activin AB beta polypeptide) (INHBB), mRNA	8.07	Down	1.90E-04
	BX537698	Homo sapiens mRNA; cDNA DKFZp686F09166 (from clone DKFZp686F09166)	8.06	Down	1.34E-03
15	BC033567	Homo sapiens, clone IMAGE:4822266, mRNA	8.04	Down	1.88E-04
	NM_052997	Homo sapiens ankyrin repeat domain 30A (ANKRD30A), mRNA	7.97	Down	1.42E-03
	NM_178550	Homo sapiens hypothetical protein MGC48998 (MGC48998), mRNA	7.82	Down	5.76E-04
20	NM_178033	Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1 (CYP4X1), mRNA	7.81	Down	8.00E-04
	NM_002924	Homo sapiens regulator of G-protein signalling 7 (RGS7), mRNA	7.8	Down	5.38E-04
25	NM_012242	Homo sapiens dickkopf homolog 1 (Xenopus laevis) (DKK1), mRNA	7.77	Down	1.24E-04
	AA740671	ob01h05.s1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1322457 3, mRNA sequence	7.74	Down	2.22E-03
	NM_005712	Homo sapiens HERV-H LTR-associating 1 (HHLA1), mRNA	7.67	Down	1.80E-04
30	BC040293	Homo sapiens, clone IMAGE:4820330, mRNA	7.65	Down	3.10E-04
	NM_000089	Homo sapiens collagen, type I, alpha 2 (COL1A2), mRNA	7.64	Down	1.17E-03
	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	7.63	Down	2.73E-04
35	BE465134	hv75e11.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3179276 3, mRNA sequence	7.53	Down	1.28E-03
	NM_002522	Homo sapiens neuronal pentraxin I (NPTX1), mRNA	7.49	Down	6.53E-04
	AK093529	Homo sapiens cDNA FLJ36210 fis, clone THYMU2000155	7.47	Down	1.18E-04
40	NM_003662	Homo sapiens pirin (iron-binding nuclear protein) (PIR), mRNA	7.29	Down	6.96E-05
	NM_001769	Homo sapiens CD9 antigen (p24) (CD9), mRNA	7.23	Down	1.80E-04
	AL359567	Homo sapiens mRNA; cDNA DKFZp547D023 (from clone DKFZp547D023)	7.21	Down	1.24E-04
45	BX647313	Homo sapiens mRNA; cDNA DKFZp686N1593 (from clone DKFZp686N1593)	7.19	Down	1.19E-04
	NM_003979	Homo sapiens G protein-coupled receptor, family C, group 5, member A (GPCR5A), mRNA	7.18	Down	1.63E-03
50	AI365141	qx96e07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2010372 3, mRNA sequence	7.04	Down	3.01E-04
	CA948963	iq30f05.y1 HR85 islet Homo sapiens cDNA clone IMAGE: 5, mRNA sequence	7.03	Down	8.55E-04
55	NM_002977	Homo sapiens sodium channel, voltage-gated, type IX, alpha (SCN9A), mRNA	6.99	Down	5.83E-05
	BC033124	Homo sapiens, clone IMAGE:2960615, mRNA	6.86	Down	6.60E-05
	NM_003385	Homo sapiens visinin-like 1 (VSNL1), mRNA	6.73	Down	1.26E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	NM_013262	Homo sapiens myosin regulatory light chain interacting protein (MYLIP), mRNA	6.65	Down	1.66E-04
10	NM_001843	Homo sapiens contactin 1 (CNTN1), transcript variant 1, mRNA	6.64	Down	3.37E-03
	NM_006727	Homo sapiens cadherin 10, type 2 (T2-cadherin) (CDH10), mRNA	6.59	Down	1.91E-03
15	NM_016192	Homo sapiens transmembrane protein with EGF-like and two follistatin-like domains 2 (TMEFF2), mRNA	6.58	Down	9.19E-05
	NM_031847	Homo sapiens microtubule-associated protein 2 (MAP2), transcript variant 4, mRNA	6.57	Down	1.13E-03
	CB115754	K-EST0159876 L8SCK0 Homo sapiens cDNA clone L8SCK0-8-H08 5, mRNA sequence	6.55	Down	8.55E-04
20	NM_025074	Homo sapiens Fraser syndrome 1 (FRAS1), transcript variant 1, mRNA	6.52	Down	4.97E-05
	NM_016206	Homo sapiens colon carcinoma related protein (FLJ38507), mRNA	6.45	Down	5.30E-04
25	BC015159	Homo sapiens cDNA clone IMAGE:3885734, partial cds	6.35	Down	3.01E-04
	NM_005613	Homo sapiens regulator of G-protein signalling 4 (RGS4), mRNA	6.33	Down	1.62E-03
	W88428	zh72g09.s1 Soares_fetal_liver_spleen_1NFLS_S1 Homo sapiens cDNA clone IMAGE:417664 3, mRNA sequence	6.28	Down	7.53E-04
30	NM_058187	Homo sapiens chromosome 21 open reading frame 63 (C21orf63), mRNA	6.16	Down	5.39E-03
	NM_020836	Homo sapiens brain-enriched guanylate kinase-associated protein (KIAA1446), mRNA	6.15	Down	3.75E-03
35	BX107838	BX107838 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998A153853 ; IMAGE:1521686, mRNA sequence	6.14	Down	1.77E-03
	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	6.13	Down	1.94E-04
40	NM_021637	Homo sapiens transmembrane protein 35 (TMEM35), mRNA	6.1	Down	2.12E-03
	NM_002521	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	6.1	Down	4.86E-04
	BF431041	nab31g02.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3267627 3, mRNA sequence	6.08	Down	7.63E-05
45	NM_003713	Homo sapiens phosphatidic acid phosphatase type 2B (PPAP2B), transcript variant 1, mRNA	6.05	Down	6.44E-04
	AV728294	AV728294 HTC Homo sapiens cDNA clone HTCBI09 5, mRNA sequence	6.03	Down	7.71E-05
50	NM_002148	Homo sapiens homeo box D10 (HOXD10), mRNA	6.01	Down	7.14E-05
	NM_176891	Homo sapiens interferon epsilon 1 (IFNE1), mRNA	6	Down	7.48E-04
	NM_002317	Homo sapiens lysyl oxidase (LOX), mRNA	5.99	Down	3.17E-04
55	NM_016179	Homo sapiens transient receptor potential cation channel, subfamily C, member 4 (TRPC4), mRNA	5.99	Down	9.19E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	AW134473	UI-H-BI1-abv-a-11-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2712885 3, mRNA sequence	5.98	Down	7.04E-05
10	NM_005130	Homo sapiens fibroblast growth factor binding protein 1 (FGFBP1), mRNA	5.95	Down	5.89E-04
	AW512111	xx70e05.x1 NCI_CGAP_Lym12 Homo sapiens cDNA clone IMAGE:2849024 3 similar to contains Alu repetitive element;, mRNA sequence	5.93	Down	4.97E-05
15	NM_020349	Homo sapiens ankyrin repeat domain 2 (stretch responsive muscle) (ANKRD2), mRNA	5.91	Down	1.18E-04
	NM_001083	Homo sapiens phosphodiesterase 5A, cGMP-specific (PDE5A), transcript variant 1, mRNA	5.88	Down	3.74E-03
20	H94320	yv18b10.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:243067 3, mRNA sequence	5.83	Down	3.18E-02
	NM_032461	Homo sapiens SPANX family, member B1 (SPANXB1), mRNA	5.8	Down	1.08E-03
25	BM994473	UI-H-DH0-aui-1-18-0-UI.s1 NCI_CGAP_DHO Homo sapiens cDNA clone IMAGE:5871137 3, mRNA sequence	5.79	Down	1.10E-04
	BE540906	601063027F1 NIH_MGC_10 Homo sapiens cDNA clone IMAGE:3449455 5, mRNA sequence	5.79	Down	4.45E-05
30	NM_033255	Homo sapiens epithelial stromal interaction 1 (breast) (EPSTI1), mRNA	5.75	Down	8.55E-04
	BU616268	UI-H-DF0-bex-n-12-0-UI.s1 NCI_CGAP_DF0 Homo sapiens cDNA clone UI-H-DF0-bex-n-12-0-UI 3, mRNA sequence	5.73	Down	6.74E-04
35	BX112170	BX112170 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGp998O07742 ; IMAGE:327390, mRNA sequence	5.73	Down	7.32E-04
	BC015108	Homo sapiens, Similar to otoconin 90, clone IMAGE:4044247, mRNA	5.72	Down	2.40E-03
40	NM_000576	Homo sapiens interleukin 1, beta (IL1B), mRNA	5.7	Down	2.97E-03
	AW470868	ha34h03.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2875637 3, mRNA sequence	5.7	Down	2.31E-03
	AK093069	Homo sapiens cDNA FLJ35750 fis, clone TEST12004539, weakly similar to Homo sapiens adican mRNA	5.7	Down	1.11E-03
45	NM_032330	Homo sapiens calpain, small subunit 2 (CAPNS2), mRNA	5.68	Down	4.44E-03
	U51694	HSU51694 Human normal gingiva Homo sapiens cDNA, mRNA sequence	5.62	Down	1.02E-04
50	NM_004335	Homo sapiens bone marrow stromal cell antigen 2 (BST2), mRNA	5.61	Down	2.84E-03
	BF431460	7o14b05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3573849 3, mRNA sequence	5.59	Down	7.53E-04
	BC046362	Homo sapiens voltage-dependent calcium channel gamma subunit-like protein, mRNA (cDNA clone MGC:50757 IMAGE:5221396), complete cds	5.59	Down	8.07E-04
55	NM_000961	Homo sapiens prostaglandin I2 (prostacyclin) synthase (PTGIS), mRNA	5.57	Down	1.01E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in Epithelial versus AF cells	Direction	adj. p-value
5	AK022877	Homo sapiens cDNA FLJ12815 fis, clone NT2RP2002546	5.56	Down	4.04E-05
10	NM_002277	Homo sapiens keratin, hair, acidic, 1 (KRTA1), mRNA	5.51	Down	3.04E-04
	AK092245	Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	5.49	Down	1.40E-03
15	NM_002427	Homo sapiens matrix metalloproteinase 13 (collagenase 3) (MMP13), mRNA	5.47	Down	4.45E-04
	BM991890	UI-H-DF1-auk-h-02-0-UI.s1 NCI_CGAP_DF1 Homo sapiens cDNA clone IMAGE:5870641 3, mRNA sequence	5.39	Down	7.53E-04
20	NM_002575	Homo sapiens serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2 (SERPINB2), mRNA	5.39	Down	9.60E-03
	NM_000782	Homo sapiens cytochrome P450, family 24, subfamily A, polypeptide 1 (CYP24A1), nuclear gene encoding mitochondrial protein, mRNA	5.36	Down	1.31E-02
25	NM_016276	Homo sapiens serum/glucocorticoid regulated kinase 2 (SGK2), transcript variant 2, mRNA	5.34	Down	6.73E-04
	AK024689	Homo sapiens cDNA: FLJ21036 fis, clone CAE09578	5.32	Down	1.36E-04
	N78829	zb17a05.s1 Soares_fetal_lung_NbHL 19W Homo sapiens cDNA clone IMAGE:302288 3, mRNA sequence	5.31	Down	6.00E-03
30	AL080103	Homo sapiens mRNA; cDNA DKFZp564N2216 (from clone DKFZp564N2216)	5.26	Down	1.61E-04
	BC035400	Homo sapiens, clone IMAGE:4822830, mRNA	5.17	Down	1.13E-02
	AK056725	Homo sapiens cDNA FLJ32163 fis, clone PLACE6000371	5.13	Down	3.72E-04
35	AK025595	Homo sapiens cDNA: FLJ21942 fis, clone HEP04527	5.12	Down	1.56E-02
	NM_015236	Homo sapiens latrophilin 3 (LPHN3), mRNA	5.11	Down	3.14E-04
	NM_012411	Homo sapiens protein tyrosine phosphatase, non-receptor type 22 (lymphoid) (PTPN22), transcript variant 2, mRNA	5.1	Down	2.78E-03
40	AI190760	qd61c05.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1733960 3, mRNA sequence	5.08	Down	2.24E-03
	NM_139241	Homo sapiens FYVE, RhoGEF and PH domain containing 4 (FGD4), mRNA	5.04	Down	6.92E-03
45	NM_033066	Homo sapiens membrane protein, palmitoylated 4 (MAGUK p55 subfamily member 4) (MPP4), mRNA	5.04	Down	7.11E-03
	BG573337	602595107F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4724521 5, mRNA sequence	5.03	Down	5.60E-03
	AA158235	zo76b02.s1 Stratagene pancreas (#937208) Homo sapiens cDNA clone IMAGE:592779 3, mRNA sequence	5.01	Down	7.10E-03
50	BX092004	BX092004 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2385330, mRNA sequence	5.01	Down	1.69E-02
55	AK058040	Homo sapiens cDNA FLJ25311 fis, clone SYN01066	5	Down	1.76E-03

TABLE VII D: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN AF-I VERSUS AF-I AT TERM CELLS

Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5				
10	NM_006228 Homo sapiens prepronociceptin (PNOC), mRNA	64.08	UP	1.14E-04
	NM_006439 Homo sapiens mab-21-like 2 (C. elegans) (MAB21L2), mRNA	47.74	UP	1.03E-04
	AK092245 Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	44.83	UP	8.97E-05
15	AK056431 Homo sapiens cDNA FLJ31869 fis, clone NT2RP7002151	35.22	UP	5.97E-05
	NM_015170 Homo sapiens sulfatase 1 (SULF1), mRNA	30.43	UP	1.53E-04
20	NM_001884 Homo sapiens hyaluronan and proteoglycan link protein 1 (HAPLN1), mRNA	27.82	UP	3.92E-05
	NM_002487 Homo sapiens necdin homolog (mouse) (NDN), mRNA	25.93	UP	2.07E-04
	BC037316 Homo sapiens, clone IMAGE:5259432, mRNA	25.16	UP	5.97E-05
25	NM_020873 Homo sapiens leucine rich repeat neuronal 1 (LRRN1), mRNA	24.74	UP	6.46E-05
	H89053 yw24c06.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253162 5, mRNA sequence	23.95	UP	6.63E-05
30	NM_002148 Homo sapiens homeo box D10 (HOXD10), mRNA	20.48	UP	8.97E-05
	NM_014421 Homo sapiens dickkopf homolog 2 (Xenopus laevis) (DKK2), mRNA	20.41	UP	6.46E-05
35	NM_016212 Homo sapiens TP53TG3 protein (TP53TG3), mRNA	20.08	UP	5.97E-05
	NM_080743 Homo sapiens serine-arginine repressor protein (35 kDa) (SRp35), mRNA	19.58	UP	6.46E-05
	NM_000170 Homo sapiens glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P) (GLDC), mRNA	18.22	UP	9.61E-05
40				
	BX647313 Homo sapiens mRNA; cDNA DKFZp686N1593 (from clone DKFZp686N1593)	17.47	UP	6.46E-05
45	NM_007361 Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	17.41	UP	9.75E-05
	NM_007072 Homo sapiens HERV-H LTR-associating 2 (HHLA2), mRNA	16.06	UP	9.61E-05
	NM_152359 Homo sapiens carnitine palmitoyltransferase 1C (CPT1C), mRNA	15.65	UP	2.26E-04
50				
	NM_002899 Homo sapiens retinol binding protein 1, cellular (RBP1), mRNA	15.47	UP	1.40E-04
	NM_005531 Homo sapiens interferon, gamma-inducible protein 16 (IFI16), mRNA	15.18	UP	1.67E-04
55	NM_021192 Homo sapiens homeo box D11 (HOXD11), mRNA	15.05	UP	1.11 E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	NM_016307	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	14.33	UP	5.97E-05
10	AA738254	nx13b02.s1 NCI_CGAP_GC3 Homo sapiens cDNA clone IMAGE:1255947 3, mRNA sequence	14.27	UP	1.64E-05
	BE788763	601475864F1 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:3879014 5, mRNA sequence	14.17	UP	3.92E-05
15	NM_030627	Homo sapiens cytoplasmic polyadenylation element binding protein 4 (CPEB4), mRNA	14.01	UP	1.40E-04
	BX113319	BX113319 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGp998G205398 ; IMAGE:2184619, mRNA sequence	13.62	UP	5.09E-05
20	NM_025074	Homo sapiens Fraser syndrome 1 (FRAS1), transcript variant 1, mRNA	13.55	UP	5.09E-05
	NM_001311	Homo sapiens cysteine-rich protein 1 (intestinal) (CRIP1), mRNA	13	UP	8.97E-05
25	NM_138394	Homo sapiens hypothetical protein BC008217 (LOC92906), mRNA	12.64	UP	2.09E-04
	BF431313	7o11d09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3573928 3, mRNA sequence	12.04	UP	1.10E-04
30	NM_133504	Homo sapiens decorin (DCN), transcript variant B, mRNA	11.74	UP	2.01E-04
	BU567804	AGENCOURT_10398872 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6614502 5, mRNA sequence	11.67	UP	5.09E-05
35	CA437861	UI-H-DH0-aur-k-12-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone UI-H-DH0-aur-k-12-0-UI 3, mRNA sequence	10.89	UP	4.99E-05
	BF431041	nab31g02.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3267627 3, mRNA sequence	10.78	UP	3.92E-05
40	NM_138966	Homo sapiens neuropilin (NRP) and tolloid (TLL)-like 1 (NETO1), transcript variant 3, mRNA	10.4	UP	8.97E-05
45	BU633163	UI-H-FL1-bgt-n-07-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgt-n-07-0-UI 3, mRNA sequence	10.32	UP	6.63E-05
	M60502	Human profilaggrin mRNA, 3 end	10.26	UP	1.40E-04
50	NM_177949	Homo sapiens armadillo repeat containing, X-linked 2 (ARMCX2), mRNA	9.9	UP	1.62E-04
	BG622707	602647476F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4768963 5, mRNA sequence	9.78	UP	1.24E-04
55	AL834140	Homo sapiens mRNA; cDNA DKFZp434A2029 (from clone DKFZp434A2029)	9.75	UP	1.34E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	NM_020353	Homo sapiens phospholipid scramblase 4 (PLSCR4), mRNA	9.71	UP	8.97E-05
10	AI905628	CM-BT094-050299-147 BT094 Homo sapiens cDNA, mRNA sequence	9.65	UP	1.03E-04
	NM_005712	Homo sapiens HERV-H LTR-associating 1 (HHLA1), mRNA	9.52	UP	1.10E-04
15	NM_003328	Homo sapiens TXK tyrosine kinase (TXK), mRNA	9.49	UP	2.35E-04
	AK095573	Homo sapiens cDNA FLJ38254 fis, clone FCBBF3000847	9.39	UP	3.92E-05
	NM_003973	Homo sapiens ribosomal protein L14 (RPL14), mRNA	9.37	UP	9.74E-05
20	AL137698	Homo sapiens mRNA; cDNA DKFZp434C1915 (from clone DKFZp434C1915); partial cds	9.32	UP	1.48E-03
		Homo sapiens F-box protein 9 (FBXO9), transcript variant 2, mRNA	9.29	UP	1.40E-04
25	NM_016588	Homo sapiens neuritin 1 (NRN1), mRNA	9.16	UP	4.09E-04
	AK127309	Homo sapiens cDNA FLJ45377 fis, clone BRHIP3019956	9	UP	4.86E-04
	NM_207482	Homo sapiens FLJ44048 protein (FLJ44048), mRNA	8.93	UP	4.37E-04
30	BE464407	hx89g05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3195032 3, mRNA sequence	8.92	UP	6.16E-04
	NM_178033	Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1 (CYP4X1), mRNA	8.59	UP	1.35E-04
35	NM_001854	Homo sapiens collagen, type XI, alpha 1 (COL11A1), transcript variant A, mRNA	8.58	UP	4.86E-04
	NM_005141	Homo sapiens fibrinogen, B beta polypeptide (FGB), mRNA	8.52	UP	5.39E-04
40	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	8.52	UP	6.46E-05
	AI651524	wb06g07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2304924 3, mRNA sequence	8.5	UP	3.47E-04
45	AI288404	qv89b01.x1 NCI_CGAP_Ut2 Homo sapiens cDNA clone IMAGE:1988713 3, mRNA sequence	8.22	UP	3.47E-04
	AL359058	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 592473	8.12	UP	1.82E-04
50	NM_002427	Homo sapiens matrix metalloproteinase 13 (collagenase 3) (MMP13), mRNA	7.92	UP	2.93E-04
	AJ318805	AJ318805 Homo sapiens adipose tissue Homo sapiens cDNA clone 2040, mRNA sequence	7.91	UP	1.03E-04
	BC033124	Homo sapiens, clone IMAGE:2960615, mRNA	7.83	UP	6.63E-05
55	BI561641	603256058F1 NIH_MGC_97 Homo sapiens cDNA clone IMAGE:5298374 5, mRNA sequence	7.82	UP	1.18E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	AK054960	Homo sapiens cDNA FLJ30398 fis, clone BRACE2008402, highly similar to Homo sapiens steroid receptor RNA activator isoform 3 mRNA	7.67	UP	1.12E-03
10	NM_174900	Homo sapiens zinc finger protein 42 (ZFP42), mRNA	7.64	UP	1.40E-04
15	NM_007099	Homo sapiens acid phosphatase 1, soluble (ACP1), transcript variant 2, mRNA	7.6	UP	1.36E-03
	BC015108	Homo sapiens, Similar to otoconin 90, clone IMAGE:4044247, mRNA	7.54	UP	6.63E-05
	AI640484	wa27f01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299321 3, mRNA sequence	7.49	UP	1.03E-03
20	AK001099	Homo sapiens cDNA FLJ10237 fis, clone HEMBB1000438	7.49	UP	9.13E-04
	AB033037	Homo sapiens mRNA for KIAA1211 protein, partial cds	7.34	UP	7.16E-05
25	BC033567	Homo sapiens, clone IMAGE:4822266, mRNA	7.29	UP	1.67E-04
	NM_001144	Homo sapiens autocrine motility factor receptor (AMFR), transcript variant 1, mRNA	7.23	UP	3.06E-04
	AI792194	ov03c02.y5 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1636226 5, mRNA sequence	7.23	UP	4.37E-04
30	AI939297	oy50g08.x5 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:1669310 3, mRNA sequence	7.21	UP	1.87E-04
35	BF512544	UI-H-BW1-amf-c-08-0-ULs1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069687 3, mRNA sequence	7.17	UP	8.19E-05
	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	7.09	UP	6.63E-05
40	H94320	yv18b10.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:243067 3, mRNA sequence	6.65	UP	2.20E-03
	NM_000557	Homo sapiens growth differentiation factor 5 (cartilage-derived morphogenetic protein-1) (GDF5), mRNA	6.5	UP	1.74E-03
45	H15096	ym29e11.r1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:49250 5, mRNA sequence	6.48	UP	5.09E-05
50	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	6.25	UP	5.77E-05
	NM_005130	Homo sapiens fibroblast growth factor binding protein 1 (FGFBP1), mRNA	6.13	UP	1.03E-04
55	BM675270	UI-E-EJ0-ahr-j-07-0-UI.s1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahr-j-07-0-UI 3, mRNA sequence	6.1	UP	1.23E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	AW268540	xv51e10.x1 NCI_CGAP_Lu28 Homo sapiens cDNA clone IMAGE:2816682 3, mRNA sequence	6.09	UP	1.09E-04
10	NM_004390	Homo sapiens cathepsin H (CTSH), transcript variant 1, mRNA	6.03	UP	1.03E-04
	BM468332	AGENCOURT_6432296 NIH_MGC_71 Homo sapiens cDNA clone IMAGE:5535773 5, mRNA sequence	5.97	UP	7.63E-05
15	NM_052997	Homo sapiens ankyrin repeat domain 30A (ANKRD30A), mRNA	5.87	UP	3.09E-03
	AK123319	Homo sapiens cDNA FLJ41325 fis, clone BRAMY2046871	5.81	UP	3.81E-04
20	NM_004235	Homo sapiens Kruppel-like factor 4 (gut) (KLF4), mRNA	5.77	UP	1.81E-04
	BC042976	Homo sapiens cDNA clone IMAGE:5295023, partial cds	5.76	UP	2.39E-03
25	AA740671	ob01h05.s1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1322457 3, mRNA sequence	5.61	UP	3.06E-03
	NM_000087	Homo sapiens cyclic nucleotide gated channel alpha 1 (CNGA1), mRNA	5.53	UP	2.39E-03
30	NM_018476	Homo sapiens brain expressed, X-linked 1 (BEX1), mRNA	5.52	UP	1.07E-03
	BX107838	BX107838 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998A153853 ; IMAGE:1521686, mRNA	5.41	UP	2.69E-03
35	NM_145201	Homo sapiens similar to CG3714 gene product (PP3856), mRNA	5.36	UP	5.77E-05
	NM_005069	Homo sapiens single-minded homolog 2 (Drosophila) (SIM2), transcript variant SIM2, mRNA	5.34	UP	7.88E-05
40	NM_004753	Homo sapiens dehydrogenase/reductase (SDR family) member 3 (DHRS3), mRNA	5.32	UP	4.71E-04
	NM_002338	Homo sapiens limbic system-associated membrane protein (LSAMP), mRNA	5.32	UP	6.88E-04
45	NM_033518	Homo sapiens solute carrier family 38, member 5 (SLC38A5), mRNA	5.28	UP	3.14E-03
	BX117317	BX117317 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998E242234 ; IMAGE:900095, mRNA sequence	5.22	UP	1.36E-03
50	CB115754	K-EST0159876 L8SCK0 Homo sapiens cDNA clone L8SCK0-8-H08 5, mRNA sequence	5.21	UP	1.22E-03
	AL359567	Homo sapiens mRNA; cDNA DKFZp547D023 (from clone DKFZp547D023)	5.18	UP	5.41E-04
	NM_020353	Homo sapiens phospholipid scramblase 4 (PLSCR4), mRNA	5.13	UP	1.36E-03
55	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	5.12	UP	9.13E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	U51694	HSU51694 Human normal gingiva Homo sapiens cDNA, mRNA sequence	5.11	UP	1.10E-04
10	NM_006546	Homo sapiens IGF-II mRNA-binding protein 1 (IMP-1), mRNA	5.1	UP	2.50E-03
	BG573337	602595107F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4724521 5, mRNA sequence	5.1	UP	5.54E-03
15	NM_006727	Homo sapiens cadherin 10, type 2 (T2-cadherin) (CDH10), mRNA	5.01	UP	3.47E-03
	NM_014358	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	36.05	Down	3.68E-04
20	NM_021013	Homo sapiens keratin, hair, acidic, 4 (KRTHA4), mRNA	33.84	Down	9.33E-04
	BM988338	UI-H-DH0-asd-f-10-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone IMAGE:5857545 3, mRNA sequence	31.23	Down	4.43E-03
25	BC039369	Homo sapiens, clone IMAGE:5271073, mRNA, partial cds	25.01	Down	1.51E-03
	BE877764	601486331F1 NIH_MGC_69 Homo sapiens cDNA clone IMAGE:3888943 5, mRNA sequence	24.69	Down	5.74E-03
30	AW300043	xs45a09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2772568 3, mRNA sequence	21.86	Down	1.45E-02
	BQ025821	UI-1-BB1p-aye-f-10-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1 p-aye-f-10-0-UI 3, mRNA sequence	20.35	Down	7.39E-03
35	BQ003501	UI-H-EI1-azd-p-06-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847413 3, mRNA sequence	19.16	Down	9.49E-04
40	NM_013387	Homo sapiens ubiquinol-cytochrome c reductase complex (7.2 kD) (HSPC051), transcript variant 1, mRNA	18.22	Down	1.43E-04
	CA391258	cs13a10.x1 Human Retinal pigment epithelium/choroid cDNA (Un-normalized, unamplified): cs Homo sapiens cDNA clone cs13a10 3, mRNA sequence	17.83	Down	3.92E-05
45	AV702977	AV702977 ADB Homo sapiens cDNA clone ADBCVD08 5, mRNA sequence	17.04	Down	1.41E-02
	NM_015714	Homo sapiens putative lymphocyte G0/G1 switch gene (G0S2), mRNA	15.4	Down	9.27E-03
50	H85497	yv88b07.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:249781 5, mRNA sequence	14.88	Down	1.03E-04
	NM_014226	Homo sapiens renal tumor antigen (RAGE), mRNA	14.77	Down	2.43E-03
55	AK023647	Homo sapiens cDNA FLJ 13585 fis, clone PLACE 1009150	14.09	Down	1.22E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	AK128288	Homo sapiens cDNA FLJ46426 fis, clone THYMU3013897	12.83	Down	2.34E-02
10	NM_017852	Homo sapiens NACHT, leucine rich repeat and PYD containing 2 (NALP2), mRNA	12.69	Down	1.36E-03
	NM_024336	Homo sapiens iroquois homeobox protein 3 (IRX3), mRNA	12.29	Down	1.29E-02
15	NM_004591	Homo sapiens chemokine (C-C motif) ligand 20 (CCL20), mRNA	11.11	Down	3.13E-02
	BF515657	UI-H-BW1-anu-e-05-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3083601 3, mRNA sequence	11.02	Down	1.26E-02
20	W69644	zd45f10.r1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:343627 5, mRNA sequence	10.54	Down	8.66E-03
	AL832916	Homo sapiens mRNA; cDNA DKFZp762I0915 (from clone DKFZp762I0915)	9.98	Down	1.94E-03
25	NM_005739	Homo sapiens RAS guanyl releasing protein 1 (calcium and DAG-regulated) (RASGRP1), mRNA	9.82	Down	3.85E-03
	BF000009	7h15g04.x1 NCI_CGAP_Co16 Homo sapiens cDNA clone IMAGE:3316086 3, mRNA sequence	9.64	Down	1.78E-02
30	NM_012105	Homo sapiens beta-site APP-cleaving enzyme 2 (BACE2), transcript variant a, mRNA	9.3	Down	1.50E-04
	BU569937	AGENCOURT_10399817 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6618011 5, mRNA sequence	9.27	Down	3.61E-02
35	NM_032782	Homo sapiens hepatitis A virus cellular receptor 2 (HAVCR2), mRNA	8.92	Down	6.61E-03
	NM_197955	Homo sapiens normal mucosa of esophagus specific 1 (NMES1), transcript variant 1, mRNA	8.82	Down	4.10E-03
40	NM_003810	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	8.08	Down	2.84E-04
	NM_014222	Homo sapiens NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8, 19kDa (NDUFA8), nuclear gene encoding mitochondrial protein, mRNA	7.99	Down	1.59E-02
45	NM_019119	Homo sapiens protocadherin beta 9 (PCDHB9), mRNA	7.85	Down	7.39E-03
	NM_014867	Homo sapiens KIAA0711 gene product (KIAA0711), mRNA	6.89	Down	9.61E-05
50	BF509573	UI-H-BI4-apf-b-11-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086949 3, mRNA sequence	6.85	Down	8.97E-05
	BQ012774	UI-1-BC1p-aaz-h-01-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1 p-aaz-h-01-0-UI 3, mRNA sequence	6.44	Down	5.30E-02
55	NM_018836	Homo sapiens transmembrane protein SHREW1 (SHREW1), mRNA	6.32	Down	2.36E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF cells isolated from 2nd trimester amniotic fluid versus AF cells isolated at term	Direction	adj. p-value
5	AW139066	UI-H-BI1-adz-d-02-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2718387 3, mRNA sequence	6.28	Down	1.03E-04
10	NM_018076	Homo sapiens armadillo repeat containing 4 (ARMC4), mRNA	6.17	Down	2.20E-03
	BI759570	603046987F1 NIH_MGC_116 Homo sapiens cDNA clone IMAGE:5187285 5, mRNA sequence	6.03	Down	5.06E-02
15	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	5.79	Down	1.03E-04
	NM_004473	Homo sapiens forkhead box E1 (thyroid transcription factor 2) (FOXE1), mRNA	5.48	Down	6.27E-05
20	NM_182920	Homo sapiens a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 9 (ADAMTS9), transcript variant 1, mRNA	5.35	Down	1.87E-04
25	AI830524	wh52c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384354 3, mRNA sequence	5.34	Down	3.42E-02
	NM_004840	Homo sapiens Rac/Cdc42 guanine nucleotide exchange factor (GEF) 6 (ARHGEF6), mRNA	5.18	Down	1.41E-03
30	AK026966	Homo sapiens cDNA: FLJ23313 fis, clone HEP11919	5.09	Down	1.49E-04
	BC044843	Homo sapiens hypothetical protein LOC339535, mRNA (cDNA clone IMAGE:5186761), partial cds	5.03	Down	3.36E-02
35	X02851	Human mRNA for interleukin-1 precursor (pre IL-1)	5.03	Down	8.60E-03

TABLE VII E: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN AF-I VERSUS AF-II CELLS

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
40					
45	D52654	HUM084D02B Clontech human fetal brain polyA+ mRNA (#6535) Homo sapiens cDNA clone GEN-084D02 5, mRNA sequence	398.37	Up	8.54E-05
50	NM_000090	Homo sapiens collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant) (COL3A1 mRNA	228.59	Up	5.20E-05
	NM_007084	Homo sapiens SRY (sex determining region Y)-box 21 (SOX21), mRNA	144.06	Up	2.10E-04
55	AK026784	Homo sapiens cDNA: FLJ23131 fis, clone LNG08502	88.3	Up	7.11E-05
	BC014344	Homo sapiens, Similar to arylacetamide deacetylase, clone IMAGE:3934567, mRNA	52.67	Up	1.31E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	BQ003501	UI-H-EI1-azd-p-06-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847413 3, mRNA sequence	52.33	Up	6.55E-05
10	BQ025821	UI-1-BB1p-aye-f-10-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-aye-f-10-0-UI 3, mRNA sequence	45.25	Up	5.46E-05
15	NM_017805	Homo sapiens Ras interacting protein 1 (RASIP1), mRNA	43.27	Up	2.20E-04
	BF515657	UI-H-BW1-anu-e-05-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3083601 3, mRNA sequence	42.11	Up	1.69E-05
20	NM_138961	Homo sapiens endothelial cell adhesion molecule (ESAM), mRNA	41.23	Up	7.66E-05
	BC046364	Homo sapiens flavoprotein oxidoreductase MICAL3, mRNA (cDNA clone IMAGE:5737121), with apparent retained intron	39.24	Up	1.47E-04
25	AK021531	Homo sapiens cDNA FLJ11469 fis, clone HEMBA1001658	36.94	Up	5.91E-04
	NM_001432	Homo sapiens epiregulin (EREG), mRNA	36.88	Up	7.51E-04
	NM_000474	Homo sapiens twist homolog 1 (acrocephalosyndactyly 3; Saethre-Chotzen syndrome) (Drosophila) (TWIST1), mRNA	36.39	Up	1.01E-04
30	NM_053281	Homo sapiens dachshund homolog 2 (Drosophila) (DACH2), mRNA	34.9	Up	7.97E-06
	H85497	yv88b07.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:249781 5, mRNA sequence	32.03	Up	1.81E-04
35	NM_033050	Homo sapiens succinate receptor 1 (SUCNR1), mRNA	29.47	Up	2.29E-04
	BX093329	BX093329 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGp998A124183 ; IMAGE:1648403, mRNA sequence	28.85	Up	6.94E-04
40	NM_012204	Homo sapiens general transcription factor IIIC, polypeptide 4, 90kDa (GTF3C4), mRNA	28.2	Up	1.29E-04
	A1124557	am58g02.x1 Johnston frontal cortex Homo sapiens cDNA clone IMAGE:1539794 3, mRNA sequence	27.99	Up	4.43E-05
	NM_014178	Homo sapiens syntaxin binding protein 6 (amisyn) (STXBP6), mRNA	26.38	Up	9.11E-05
45	U83115	Human non-lens beta gamma-crystallin like protein (AIM1) mRNA, partial cds	26.15	Up	3.63E-05
	NM_020927	Homo sapiens KIAA1576 protein (KIAA1576), mRNA	25.98	Up	2.56E-05
	NM_001442	Homo sapiens fatty acid binding protein 4, adipocyte (FABP4), mRNA	25.37	Up	4.39E-04
50	BC045666	Homo sapiens tumor protein p53 inducible protein 11, mRNA (cDNA clone IMAGE:5298525), containing frame-shift errors	25.36	Up	1.84E-04
	NM_013387	Homo sapiens ubiquinol-cytochrome c reductase complex (7.2 kD) (HSPC051), transcript variant 1, mRNA	24.64	Up	1.47E-04
55	AK125453	Homo sapiens cDNA FLJ43464 fis, clone OCBBF2036225	23.47	Up	3.90E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_170744	Homo sapiens unc-5 homolog B (C. elegans) (UNC5B), mRNA	22.66	Up	9.44E-05
10	AB067499	Homo sapiens mRNA for KIAA1912 protein, partial cds	22.2	Up	2.69E-05
	NM_005110	Homo sapiens glutamine-fructose-6-phosphate transaminase 2 (GFPT2), mRNA	22	Up	9.53E-06
	CD677332	ho15f06.y1 Human Trabecular meshwork cDNA: hohphq Homo sapiens cDNA clone ho15f06 5, mRNA sequence	21.26	Up	7.83E-05
15	NM_005525	Homo sapiens hydroxysteroid (11-beta) dehydrogenase 1 (HSD11B1), transcript variant 1, mRNA	20.9	Up	4.93E-05
	NM_006475	Homo sapiens periostin, osteoblast specific factor (POSTN), mRNA	20.5	Up	6.66E-04
20	AK027128	Homo sapiens cDNA: FLJ23475 fis, clone HSI13659	20.14	Up	1.00E-04
	AI962169	wq45c10.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474226 3, mRNA sequence	19.91	Up	6.86E-05
	BM713465	UI-E-EJ0-aho-m-22-0-UI.r1 UI-E-EJO Homo sapiens cDNA clone UI-E-EJ0-aho-m-22-0-UI 5, mRNA sequence	19.86	Up	4.55E-05
25	NM_014802	Homo sapiens KIAA0528 gene product (KIAA0528), mRNA	19.34	Up	2.92E-03
	NM_024336	Homo sapiens iroquois homeobox protein 3 (IRX3), mRNA	19.2	Up	4.93E-05
30		Homo sapiens 5-hydroxytryptamine (serotonin) receptor 2B (HTR2B), mRNA	18.89	Up	5.39E-05
	NM_000867				
	AF052115	Homo sapiens clone 23688 mRNA sequence	17.49	Up	8.17E-05
	BC016722	Homo sapiens cDNA clone IMAGE:4075924, partial cds	16.82	Up	2.28E-03
	NM_018013	Homo sapiens hypothetical protein FLJ10159 (FLJ10159), mRNA	16.78	Up	5.98E-06
35					
	N33310	yy39g10.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273666 3, mRNA sequence	16.64	Up	1.30E-04
	W38393	zb15c07.r1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:302124 5, mRNA sequence	16.52	Up	1.05E-04
40					
	R44402	yg37a01.s1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:34639 3 similar to contains MER35 repetitive element ;, mRNA sequence	16.02	Up	4.93E-05
	NM_020226	Homo sapiens PR domain containing 8 (PRDM8), mRNA	15.99	Up	7.27E-06
45	BU677169	UI-CF-DU1-aaj-c-16-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaj-c-16-0-UI 3, mRNA sequence	15.75	Up	3.92E-04
	NM_001146	Homo sapiens angiopoietin 1 (ANGPT1), transcript variant 1, mRNA	15.5	Up	5.07E-04
50					
		Homo sapiens cDNA FLJ41881 fis, clone	15.42	Up	1.03E-04
	AK123875	OCBBF2021833			
	NM_000735	Homo sapiens glycoprotein hormones, alpha polypeptide (CGA), mRNA	15.42	Up	2.19E-04
55					
	AW445209	UI-H-BI3-akc-g-11-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2733908 3, mRNA sequence	15.03	Up	2.89E-05
	BC040678	Homo sapiens, clone IMAGE:4817707, mRNA	14.85	Up	1.69E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_052846	Homo sapiens elastin microfibril interfacer 3 (EMILIN3), mRNA	14.55	Up	5.78E-05
10	NM_005320	Homo sapiens histone 1, H1d (HIST1H1D), mRNA	14.49	Up	9.77E-03
	U79271	Human clones 23920 and 23921 mRNA sequence	13.86	Up	7.54E-04
	BQ186389	UI-E-EJ1-ajr-c-19-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajr-c-19-0-UI 5, mRNA sequence	13.7	Up	5.94E-04
15	NM_032638	Homo sapiens GATA binding protein 2 (GATA2), mRNA	13.42	Up	1.67E-04
	NM_000089	Homo sapiens collagen, type I, alpha 2 (COL1A2), mRNA	13.37	Up	1.89E-03
	NM_139211	Homo sapiens homeodomain-only protein (HOP), transcript variant 2, mRNA	13.15	Up	1.03E-04
	BM663928	UI-E-CI1-afw-p-01-0-UI.s1 UI-E-CI1 Homo sapiens cDNA clone UI-E-CI1-afw-p-01-0-UI 3, mRNA sequence	13.08	Up	3.22E-04
20	NM_021643	Homo sapiens tribbles homolog 2 (Drosophila) (TRIB2), mRNA	12.66	Up	2.40E-04
	NM_004411	Homo sapiens dynein, cytoplasmic, intermediate polypeptide 1 (DNCI1), mRNA	12.59	Up	1.43E-05
25	AK128288	Homo sapiens cDNA FLJ46426 fis, clone THYMU3013897	12.51	Up	8.11E-04
	AL133118	Homo sapiens mRNA; cDNA DKFZp586N0121 (from clone DKFZp586N0121)	12.28	Up	2.54E-04
	BQ027984	UI-H-CO0-arg-e-03-0-UI.s1 NCI_CGAP_Sub9 Homo sapiens cDNA clone IMAGE:3106611 3, mRNA sequence	12.2	Up	8.14E-05
30	NM_001175	Homo sapiens Rho GDP dissociation inhibitor (GDI) beta (ARHGDIB), mRNA	11.91	Up	5.05E-05
	NM_005595	Homo sapiens nuclear factor I/A (NFIA), mRNA	11.71	Up	6.19E-04
35	AK055518	Homo sapiens cDNA FLJ30956 fis, clone HCASM2000202	11.56	Up	2.92E-03
	NM_000407	Homo sapiens glycoprotein Ib (platelet), beta polypeptide (GP1 BB), mRNA	11.29	Up	1.18E-03
40		Homo sapiens cDNA FLJ26796 fis, clone PRS05079	11.1	Up	1.24E-03
	AK130306				
	NM_012310	Homo sapiens kinesin family member 4A (KIF4A), mRNA	11.08	Up	1.59E-02
	NM_058187	Homo sapiens chromosome 21 open reading frame 63 (C21orf63), mRNA	10.78	Up	5.06E-04
45	NM_016206	Homo sapiens colon carcinoma related protein (FLJ38507), mRNA	10.71	Up	3.40E-04
	NM_032918	Homo sapiens RAS-like, estrogen-regulated, growth inhibitor (RERG), mRNA	10.69	Up	1.36E-04
	NM_053044	Homo sapiens serine protease HTRA3 (HTRA3), mRNA	10.65	Up	2.39E-04
50	M_153355	Homo sapiens T-cell lymphoma breakpoint associated target 1 (TCBA1), mRNA	10.47	Up	8.32E-03
	No 152665	Homo sapiens hypothetical protein FLJ40873 (FLJ40873), mRNA	10.42	Up	2.20E-04
55	NM_004460	Homo sapiens fibroblast activation protein, alpha (FAP), mRNA	10.39	Up	5.27E-05
	NM_152550	Homo sapiens SH3 domain containing ring finger 2 (SH3RF2), mRNA	10.29	Up	6.33E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	BF674238	602136969F1 NIH_MGC_83 Homo sapiens cDNA clone IMAGE:4273624 5, mRNA sequence	10.27	Up	4.87E-02
10	NM_005325	Homo sapiens histone 1, H1a (HIST1H1A), mRNA	9.97	Up	1.19E-03
	AV702977	AV702977 ADB Homo sapiens cDNA clone ADBCVD08 5, mRNA sequence	9.96	Up	4.43E-05
	NM_024600	Homo sapiens chromosome 16 open reading frame 30 (C16orf30), mRNA	9.89	Up	2.28E-03
15	AK023739	Homo sapiens cDNA FLJ13677 fis, clone PLACE1011982	9.82	Up	9.53E-06
	AK124699	Homo sapiens cDNA FLJ42709 fis, clone BRAMY3007350	9.73	Up	2.38E-02
20	NM_005239	Homo sapiens v-ets erythroblastosis virus E26 oncogene homolog 2 (avian) (ETS2), mRNA	9.56	Up	6.20E-03
	AK091337	Homo sapiens cDNA FLJ34018 fis, clone FCBBF2002801	9.3	Up	6.30E-04
	H15096	ym29e11.r1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:49250 5, mRNA sequence	9.17	Up	7.83E-05
25	NM_018286	Homo sapiens hypothetical protein FLJ10970 (FLJ10970), mRNA	9.16	Up	2.02E-03
	NM_003783	Homo sapiens UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 2 (B3GALT2), mRNA	9.04	Up	1.38E-02
30	NM_032457	Homo sapiens BH-protocadherin (brain-heart) (PCDH7), transcript variant c, mRNA	8.94	Up	1.53E-04
	NM_015192	Homo sapiens phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 1, mRNA	8.82	Up	3.03E-03
35	NM_003514	Homo sapiens histone 1, H2am (HIST1H2AM), mRNA	8.75	Up	5.60E-03
	A1082507	ox55c02.s1 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGE:1660226 3, mRNA sequence	8.68	Up	2.98E-03
40	NM_025107	Homo sapiens myc target 1 (MYCT1), mRNA	8.61	Up	1.37E-04
	BC036004	Homo sapiens, clone IMAGE:4730399, mRNA	8.56	Up	2.94E-04
	BF029356	601765592F1 NIH_MGC_53 Homo sapiens cDNA clone IMAGE:3997510 5, mRNA sequence	8.47	Up	1.99E-03
	NM_004787	Homo sapiens slit homolog 2 (Drosophila) (SLIT2), mRNA	8.47	Up	8.47E-04
45	CA865586	ir42e09.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6547889 3, mRNA sequence	8.46	Up	8.95E-04
	NM_021614	Homo sapiens potassium intermediate/small conductance calcium-activated channel, subfamily N, member 2 (KCNN2), transcript variant 1, mRNA	8.37	Up	2.28E-05
50	NM_004791	Homo sapiens integrin, beta-like 1 (with EGF-like repeat domains) (ITGBL1 mRNA	8.31	Up	2.21E-04
	NM_052954	Homo sapiens cysteine and tyrosine-rich 1 (CYR1), mRNA	8.12	Up	8.96E-05
55	NM_032883	Homo sapiens chromosome 20 open reading frame 100 (C20orf100), mRNA	8.1	Up	4.12E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	BX1065777	BX106577 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGp998H131854 ; IMAGE:754236, mRNA sequence	8.06	Up	7.14E-05
10					
	AK074131	Homo sapiens mRNA for FLJ00204 protein	7.95	Up	2.59E-04
	NM_024621	Homo sapiens hypothetical protein FLJ12604 (FLJ12604), mRNA	7.88	Up	4.06E-04
	BX647541	Homo sapiens mRNA; cDNA DKFZp686P0492 (from clone DKFZp686P0492)	7.84	Up	8.97E-05
15					
	NM_145016	Homo sapiens BXMAS2-10 (BXMAS2-10), mRNA	7.78	Up	5.05E-04
	BU634363	UI-H-FL1-bgx-o-20-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgx-o-20-0-UI 3, mRNA sequence	7.7	Up	1.84E-04
20					
	AI939462	tf23h06.x5 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:2097083 3, mRNA sequence	7.49	Up	1.21E-04
		Homo sapiens nudix (nucleoside diphosphate linked moiety X)-type motif 10 (NUDT10), mRNA	7.47	Up	5.46E-05
25					
	NM_153183	UI-1-BC1p-asi-a-02-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1p-asi-a-02-0-UI 3, mRNA sequence	7.43	Up	1.38E-03
	BQ011545				
	T47612	yb15h03.s1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:71285 3, mRNA sequence	7.41	Up	4.13E-03
30					
	NM_020809	Homo sapiens Rho GTPase activating protein 20 (ARHGAP20), mRNA	7.36	Up	2.50E-04
	BC042378	Homo sapiens, clone IMAGE:5277693, mRNA	7.25	Up	3.63E-03
	AF414442	Homo sapiens ovarian cancer related tumor marker CA125 mRNA, complete cds	7.11	Up	2.19E-03
35					
	NM_006350	Homo sapiens folistatin (FST), transcript variant FST317, mRNA	7.1	Up	5.52E-04
	CD723006	oj16h02.y1 Human lacrimal gland, unamplified: oj Homo sapiens cDNA clone oj16h02 5, mRNA sequence	7.09	Up	5.06E-04
40					
	BC027461	Homo sapiens cDNA clone IMAGE:2984900, containing frame-shift errors	7.02	Up	4.43E-05
	NM_152782	Homo sapiens Sad1 and UNC84 domain containing 1 (SUNC1), mRNA	6.94	Up	1.34E-03
	AK125001	Homo sapiens cDNA FLJ43011 fis, clone BRTHA2015853	6.91	Up	3.36E-03
45					
	NM_152314	Homo sapiens hypothetical protein MGC34830 (MGC34830), mRNA	6.89	Up	1.08E-04
	NM_080872	Homo sapiens unc-5 homolog D (C. elegans) (UNC5D), mRNA	6.88	Up	3.33E-04
50					
	NM_015714	Homo sapiens putative lymphocyte G0/G1 switch gene (G0S2), mRNA	6.86	Up	1.34E-03
	NM_014548	Homo sapiens tropomodulin 2 (neuronal) (TMOD2), mRNA	6.83	Up	5.11E-03
55					
	NM_002771	Homo sapiens protease, serine, 3 (mesotrypsin) (PRSS3), mRNA	6.82	Up	1.28E-03
	NM_012081	Homo sapiens elongation factor, RNA polymerase II, 2 (ELL2), mRNA	6.75	Up	3.00E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	AW025556	wu97g10.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:2528034 3, mRNA sequence	6.7	Up	2.11E-03
10	NM_016522	Homo sapiens neurotrimin (HNT), mRNA	6.7	Up	1.84E-04
	AK001279	Homo sapiens cDNA FLJ10417 fis, clone NT2RP1000112	6.68	Up	6.41E-04
	NM_020993	Homo sapiens B-cell CLL/lymphoma 7A (BCL7A), mRNA	6.6	Up	1.79E-02
	NM_032578	Homo sapiens myopalladin (FLJ14437), mRNA	6.55	Up	5.46E-05
15	NM_024913	Homo sapiens hypothetical protein FLJ21986 (FLJ21986), mRNA	6.47	Up	9.10E-04
	NM_016428	Homo sapiens ABI gene family, member 3 (ABI3), mRNA	6.45	Up	1.07E-02
	NM_018371	Homo sapiens chondroitin beta1,4 N-acetylgalactosaminyltransferase (ChGn), mRNA	6.44	Up	1.75E-04
20	BM802920	AGENCOURT_6457446 NIH_MGC_88 Homo sapiens cDNA clone IMAGE:5560288 5, mRNA sequence	6.37	Up	1.34E-03
	NM_183245	Homo sapiens inversin (INVS), transcript variant 2, mRNA	6.32	Up	6.46E-05
25	AA908815	og77h08.s1 NCI_CGAP_Ov8 Homo sapiens cDNA clone IMAGE:1454367 3, mRNA sequence	6.31	Up	3.33E-04
	BQ021661	UI-H-DH1-axg-p-14-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5828605 3, mRNA sequence	6.28	Up	4.40E-04
30	NM_017527	Homo sapiens lymphocyte antigen 6 complex, locus K (LY6K), mRNA	6.24	Up	1.36E-02
	NM_175887	Homo sapiens hypothetical protein LOC222171 (LOC222171), mRNA	6.23	Up	1.76E-04
35	BM929598	UI-E-EJ1-aje-l-10-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-aje-l-10-0-UI 5, mRNA sequence	6.18	Up	1.44E-04
	AI911163	wd24c09.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2329072 3, mRNA sequence	6.17	Up	9.74E-03
40	AK024653	Homo sapiens cDNA: FLJ21000 fis, clone CAE03359	6.17	Up	2.15E-04
	AK124907	Homo sapiens cDNA FLJ42917 fis, clone BRHIP3026335	6.15	Up	2.20E-03
	NM_001147	Homo sapiens angiopoietin 2 (ANGPT2), mRNA	6.14	Up	1.24E-04
	NM_001124	Homo sapiens adrenomedullin (ADM), mRNA	6.13	Up	1.72E-04
45	AI733665	an31g04.x5 Gessler Wilms tumor Homo sapiens cDNA clone IMAGE:1700310 3, mRNA sequence	6.11	Up	5.78E-05
	NM_017655	Homo sapiens PDZ domain protein GIPC2 (GIPC2), mRNA	6.09	Up	1.81E-04
50	NM_003405	Homo sapiens tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, eta polypeptide (YWHAH), mRNA	5.96	Up	5.04E-03
	NM_000441	Homo sapiens solute carrier family 26, member 4 (SLC26A4), mRNA	5.95	Up	2.72E-02
55	BX648323	Homo sapiens mRNA; cDNA DKFZp686K10163 (from clone DKFZp686K10163)	5.93	Up	1.86E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	AW291775	UI-H-BI2-agv-h-04-0-UI.s1 NCI_CGAP_Sub4 Homo sapiens cDNA clone IMAGE:2725855 3, mRNA sequence	5.93	Up	5.03E-03
10	NM_001546	Homo sapiens inhibitor of DNA binding 4, dominant negative helix-loop-helix protein (ID4), mRNA	5.9	Up	2.39E-03
	NM_152996	Homo sapiens sialyltransferase 7 ((alpha-N-acetylneuraminy-2,3-beta-galactosyl-1,3)-N-acetyl galactosaminide alpha-2,6-sialyltransferase) C (SIAT7C), mRNA	5.88	Up	5.92E-04
15	BM556191	AGENCOURT_6544282 NIH_MGC_88 Homo sapiens cDNA clone IMAGE:5550173 5, mRNA sequence	5.8	Up	7.12E-03
	NM_021242	Homo sapiens MID1 interacting protein 1 (gastrulation specific G12-like (zebrafish)) (MID1P1), mRNA	5.75	Up	3.63E-03
20	BI868709	603392040F1 NIH_MGC_90 Homo sapiens cDNA clone IMAGE:5402337 5, mRNA sequence	5.69	Up	3.11E-03
	BM713114	UI-E-EJ0-ahi-a-13-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahi-a-13-0-UI 5, mRNA sequence	5.68	Up	1.54E-03
25	AL049443	Homo sapiens mRNA; cDNA DKFZp586N2020 (from clone DKFZp586N2020)	5.68	Up	4.15E-04
	NM_005905	Homo sapiens SMAD, mothers against DPP homolog 9 (Drosophila) (SMAD9), mRNA	5.65	Up	6.72E-04
30		Homo sapiens activating signal cointegrator 1 complex subunit 3 (ASCC3), mRNA	5.63	Up	6.59E-03
	NM_006828				
	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	5.62	Up	9.70E-05
	R22189	yh26a02.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:130826 3, mRNA sequence	5.62	Up	1.08E-04
35	NM_020226	Homo sapiens PR domain containing 8 (PRDM8), mRNA	5.57	Up	3.04E-02
	R01937	ye85h04.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:124567 3 similar to gb:L13923 FIBRILLIN 1 PRECURSOR (HUMAN);, mRNA sequence	5.56	Up	1.87E-03
40	BM724062	UI-E-EO1-aiy-a-22-0-UI.r1 UI-E-EO1 Homo sapiens cDNA clone UI-E-EO1-aiy-a-22-0-UI 5, mRNA sequence	5.52	Up	2.40E-02
	NM_080671	Homo sapiens potassium voltage-gated channel, Isk-related family, member 4 (KCNE4), mRNA	5.51	Up	1.35E-02
45	AL558097	AL558097 Homo sapiens T CELLS (JURKAT CELL LINE) COT 10-NORMALIZED Homo sapiens cDNA clone CS0DJ002YD14 5-PRIME, mRNA sequence	5.5	Up	2.82E-02
	AI090760	qa65e08.x1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:1691654 3, mRNA sequence	5.49	Up	7.59E-03
50	NM_030756	Homo sapiens transcription factor 7-like 2 (T-cell specific, HMG-box) (TCF7L2), mRNA	5.48	Up	4.57E-04
	AL832916	Homo sapiens mRNA; cDNA DKFZp76210915 (from clone DKFZp76210915)	5.47	Up	1.84E-04
55	NM_001870	Homo sapiens carboxypeptidase A3 (mast cell) (CPA3), mRNA	5.46	Up	6.04E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	BM998303	UI-H-DT1-awc-h-03-0-UI.s1 NCI_CGAP_DT1 Homo sapiens cDNA clone IMAGE:5887538 3, mRNA sequence	5.45	Up	2.71E-04
10					
	NM_004298	Homo sapiens nucleoporin 155kDa (NUP155), transcript variant 2, mRNA	5.45	Up	4.65E-02
	NM_138970	Homo sapiens neurexin 3 (NRXN3), transcript variant beta, mRNA	5.44	Up	1.15E-04
15					
	NM_005654	Homo sapiens nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	5.43	Up	1.74E-04
	NM_012388	Homo sapiens pallidin homolog (mouse) (PLDN), mRNA	5.38	Up	1.17E-02
	NM_033284	Homo sapiens transducin (beta)-like 1Y-linked (TBL1Y), transcript variant 1, mRNA	5.38	Up	4.42E-02
20					
	AK097648	Homo sapiens cDNA FLJ40329 fis, clone TESTI2031418, weakly similar to TRYPSIN I-P1 PRECURSOR (EC 3.4.21.4)	5.38	Up	1.83E-04
	AK124563	Homo sapiens cDNA FLJ42572 fis, clone BRACE3008092	5.35	Up	4.57E-04
25					
		Homo sapiens interleukin 7 receptor (IL7R), mRNA	5.31	Up	8.90E-04
	NM_002185				
	BU661543	cl73d12.z1 Hembase; Erythroid Precursor Cells (LCB:cl library) Homo sapiens cDNA clone cl73d12 5, mRNA sequence	5.27	Up	6.26E-04
30					
	AK092371	Homo sapiens cDNA FLJ35052 fis, clone OCBBF2018234, highly similar to GUANINE NUCLEOTIDE-BINDING PROTEIN G(I)/G(S)/G(O) GAMMA-2 SUBUNIT (G GAMMA-I)	5.26	Up	4.82E-03
35					
	NM_033554	Homo sapiens major histocompatibility complex, class II, DP alpha 1 (HLA-DPA1), mRNA	5.24	Up	5.25E-04
	NM_153014	Homo sapiens hypothetical protein FLJ30634 (FLJ30634), mRNA	5.23	Up	2.64E-02
	AA455071	aa04d03.s1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:812261 3, mRNA sequence	5.21	Up	7.36E-04
40					
	NM_005630	Homo sapiens solute carrier organic anion transporter family, member 2A1 (SLC02A1), mRNA	5.2	Up	1.46E-04
	NM_002619	Homo sapiens platelet factor 4 (chemokine (C-X-C motif) ligand 4) (PF4), mRNA	5.19	Up	3.64E-04
45					
	AA232643	zr47e10.r1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:666570 5, mRNA sequence	5.17	Up	1.91E-04
	NM_005398	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 3C (PPP1 R3C), mRNA	5.14	Up	1.29E-04
50					
	CA843592	ir49c12.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6548544 3, mRNA sequence	5.13	Up	1.89E-04
	BM677978	UI-E-EJ0-aig-o-17-0-UI.s1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aig-o-17-0-UI 3, mRNA sequence	5.11	Up	3.39E-03
	NM_016196	Homo sapiens RNA binding motif protein 19 (RBM19), mRNA	5.1	Up	1.84E-04
55					

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	BX105152	BX105152 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998J212575 ; IMAGE:1031156, mRNA sequence	5.1	Up	3.96E-05
10					
	NM_006329	Homo sapiens fibulin 5 (FBLN5), mRNA	5.06	Up	1.18E-03
	NM_001765	Homo sapiens CD1C antigen, c polypeptide (CD1C), mRNA	5.06	Up	1.54E-04
	NM_022768	Homo sapiens RNA binding motif protein 15 (RBM15), mRNA	5.06	Up	2.82E-04
15					
	N49730	yz06a12.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:282238 3 similar to contains Alu repetitive element;contains element PTR5 repetitive element ;, mRNA sequence	5.03	Up	9.95E-05
20					
	AK124856	Homo sapiens cDNA FLJ42866 fis, clone BRHIP2020622	5.02	Up	1.25E-03
	NM_001008	Homo sapiens ribosomal protein S4, Y-linked 1 (RPS4Y1), mRNA	456.26	Down	7.09E-06
25					
	NM_138963	Homo sapiens ribosomal protein S4, Y-linked 2 (RPS4Y2), mRNA	423.01	Down	1.53E-06
	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	404.8	Down	3.71E-06
	BG219729	RST39494 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	264.66	Down	2.05E-06
30					
	H70730	yu69e10.r1 Weizmann Olfactory Epithelium Homo sapiens cDNA clone IMAGE:239082 5, mRNA sequence	256.95	Down	1.53E-06
	NM_004653	Homo sapiens Smcy homolog, Y-linked (mouse) (SMCY), mRNA	252.16	Down	7.51E-07
35					
	AB032980	Homo sapiens mRNA for KIAA1154 protein, partial cds	227.36	Down	4.39E-05
	AI765021	wh56c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384738 3, mRNA sequence	212.62	Down	1.69E-05
	NM_012307	Homo sapiens erythrocyte membrane protein band 4.1-like 3 (EPB41L3), mRNA	192.31	Down	2.12E-06
40					
	NM_004660	Homo sapiens DEAD (Asp-Glu-Ala-Asp) box polypeptide 3, Y-linked (DDX3Y), mRNA	184.34	Down	1.69E-05
	NM_013230	Homo sapiens CD24 antigen (small cell lung carcinoma cluster 4 antigen) (CD24), mRNA	171.2	Down	3.78E-07
45					
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	166.79	Down	7.27E-06
	AK023631	Homo sapiens cDNA FLJ13569 fis, clone PLACE 1008369	163.75	Down	2.77E-07
	BX102632	BX102632 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998J052307 ; IMAGE:928228, mRNA sequence	157.63	Down	1.88E-06
50					
	NM_006228	Homo sapiens prepronociceptin (PNOC), mRNA	144.89	Down	2.50E-05
	NM_018658	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 16 (KCNJ16), transcript variant 1, mRNA	139.61	Down	4.43E-05
55					
	NM_003411	Homo sapiens zinc finger protein, Y-linked (ZFY), mRNA	130.35	Down	2.77E-07

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_007038	Homo sapiens a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2) (ADAMTS5), mRNA	126.24	Down	7.83E-05
10					
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	112.25	Down	1.92E-05
	NM_003385	Homo sapiens visinin-like 1 (VSNL1), mRNA	105.15	Down	2.05E-06
	NM_153634	Homo sapiens copine VIII (CPNE8), mRNA	101.79	Down	9.53E-06
15					
	NM_000927	Homo sapiens ATP-binding cassette, sub-family B (MDR/TAP), member 1 (ABCB1), mRNA	94.49	Down	5.20E-05
	BG197054	RST16291 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	92.27	Down	2.05E-06
20					
	NM_004617	Homo sapiens transmembrane 4 superfamily member 4 (TM4SF4), mRNA	69.46	Down	5.88E-05
	NM_000990	Homo sapiens ribosomal protein L27a (RPL27A), mRNA	63.45	Down	2.03E-04
	NM_014893	Homo sapiens neuroligin 4, Y-linked (NLGN4Y), mRNA	59.4	Down	2.05E-06
	AI249696	qj64a03.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1864204 3, mRNA sequence	58.18	Down	1.92E-05
25					
	AK092245	Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	57.12	Down	2.50E-05
	AI335277	tb29h06.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2055803 3, mRNA sequence	56.89	Down	1.80E-05
30					
	NM_016356	Homo sapiens doublecortin domain containing 2 (DCDC2), mRNA	49.57	Down	3.24E-05
	NM_138966	Homo sapiens neuropilin (NRP) and tolloid (TLL)-like 1 (NETO1), transcript variant 3, mRNA	48.88	Down	1.41E-04
35					
		Homo sapiens desmocollin 2 (DSC2), transcript variant Dsc2a, mRNA	48.6	Down	1.60E-05
	NM_024422				
	NM_031862	Homo sapiens membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125) (M17S2), transcript variant 3, mRNA	45.75	Down	5.31E-05
40					
	BF431030	7o18c06.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3574283 3, mRNA sequence	45.24	Down	2.90E-05
	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	44.97	Down	2.77E-07
	NM_020873	Homo sapiens leucine rich repeat neuronal 1 (LRRN1), mRNA	42.8	Down	3.33E-05
45					
	BF512544	UI-H-BW1-amf-c-08-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069687 3, mRNA sequence	42.38	Down	3.24E-05
	CA416106	UI-H-FEO-bbs-f-17-0-UI.s1 NCI_CGAP_FE0 Homo sapiens cDNA clone UI-H-FEO-bbs-f-17-0-UI 3, mRNA sequence	41.22	Down	5.15E-05
50					
	AI244954	qj93h05.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1867065 3, mRNA sequence	40.95	Down	9.53E-06
	BF798098	RC1-C10045-021000-021-f02 CI0045 Homo sapiens cDNA, mRNA sequence	40.6	Down	6.53E-05
55					
	NM_002423	Homo sapiens matrix metalloproteinase 7 (matrilysin, uterine) (MMP7), mRNA	39.36	Down	9.07E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5		UI-CF-DU1-aaz-f-04-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaz-f-04-0-UI 3, mRNA sequence	37.22	Down	6.19E-05
10	BU680661				
	AL080103	Homo sapiens mRNA; cDNA DKFZp564N2216 (from clone DKFZp564N2216)	37.11	Down	1.50E-05
	NM_025074	Homo sapiens Fraser syndrome 1 (FRAS1), transcript variant 1, mRNA	36.19	Down	7.09E-06
15	NM_004681	Homo sapiens eukaryotic translation initiation factor 1A, Y-linked (EIF1AY), mRNA	36.04	Down	1.84E-04
	AK096481	Homo sapiens cDNA FLJ39162 fis, clone OCBBF2002376	35.96	Down	4.87E-05
20	NM_018168	Homo sapiens chromosome 14 open reading frame 105 (C14orf105), mRNA	34.11	Down	3.52E-05
	NM_005329	Homo sapiens hyaluronan synthase 3 (HAS3), transcript variant 1, mRNA	33.86	Down	2.25E-06
	NM_002527	Homo sapiens neurotrophin 3 (NTF3), mRNA	32.19	Down	1.31E-05
25	NM_001935	Homo sapiens dipeptidylpeptidase 4 (CD26, adenosine deaminase complexing protein 2) (DPP4), mRNA	31.89	Down	1.66E-04
	NM_004932	Homo sapiens cadherin 6, type 2, K-cadherin (fetal kidney) (CDH6), mRNA	30.7	Down	2.59E-05
	BX648643	Homo sapiens mRNA; cDNA DKFZp686O17106 (from clone DKFZp686O17106)	30.56	Down	7.09E-06
30	NM_032576	Homo sapiens chromosome Y open reading frame 15B (CYorf15B), mRNA	30.49	Down	2.05E-06
	NM_002276	Homo sapiens keratin 19 (KRT19), mRNA	29.34	Down	2.84E-04
	NM_152487	Homo sapiens hypothetical protein FLJ31842 (FLJ31842), mRNA	28.96	Down	2.08E-04
35	NM_207446	Homo sapiens hypothetical gene supported by AK075564; BC060873 (LOC400451), mRNA	28.29	Down	9.53E-06
	NM_020349	Homo sapiens ankyrin repeat domain 2 (stretch responsive muscle) (ANKRD2), mRNA	27.27	Down	1.17E-05
40	NM_012198	Homo sapiens grancalcin, EF-hand calcium binding protein (GCA), mRNA	27.06	Down	7.97E-04
	AK056882	Homo sapiens cDNA FLJ32320 fis, clone PROST2003537	26.58	Down	1.63E-04
45	NM_019000	Homo sapiens hypothetical protein FLJ20152 (FLJ20152), mRNA	25.35	Down	6.39E-05
	BG436244	602508665F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4605617 5, mRNA sequence	25.2	Down	7.09E-06
	AA738254	nx13b02.s1 NCI_CGAP_GC3 Homo sapiens cDNA clone IMAGE:1255947 3, mRNA sequence	25.16	Down	9.53E-06
50	NM_002521	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	24.93	Down	8.66E-04
	NM_000104	Homo sapiens cytochrome P450, family 1, subfamily B, polypeptide 1 (CYP1B1), mRNA	24.83	Down	3.50E-05
55	NM_016946	Homo sapiens F11 receptor (F11 R), transcript variant 1, mRNA	24.8	Down	4.83E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_016946	Homo sapiens F11 receptor (F11 R), transcript variant 1, mRNA	24.38	Down	2.77E-05
10	NM_080743	Homo sapiens serine-arginine repressor protein (35 kDa) (SRrp35), mRNA	23.93	Down	1.02E-05
	NM_175056	Homo sapiens hypothetical protein LOC131368 (LOC131368), mRNA	23.5	Down	4.93E-05
	NM_000557	Homo sapiens growth differentiation factor 5 (cartilage-derived morphogenetic protein-1) (GDF5), mRNA	23.37	Down	8.14E-05
15	NM_004221	Homo sapiens natural killer cell transcript 4 (NK4), mRNA	23.12	Down	4.24E-04
	BF431041	nab31g02.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3267627 3, mRNA sequence	22.5	Down	4.93E-05
20	NM_178470	Homo sapiens WD repeat domain 40B (WDR40B), mRNA	21.85	Down	4.54E-05
	AI651524	wb06g07.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2304924 3, mRNA sequence	21.78	Down	4.59E-05
25	CB047092	NISC_gf08f03.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:3253013 3, mRNA sequence	21.63	Down	5.93E-06
	AA449137	zx03d12.r1 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGE:785399 5, mRNA sequence	21.62	Down	2.08E-04
	H89053	yw24c06.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253162 5, mRNA sequence	21.41	Down	4.16E-03
30	BX509117	DKFZp686C09280_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686C09280 5, mRNA sequence	20.95	Down	2.31E-05
	NM_00082	Homo sapiens secreted phosphoprotein 1 5 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA	20.66	Down	6.44E-03
35	AK124873	Homo sapiens cDNA FLJ42883 fis, clone BRHIP3006683	20.37	Down	4.83E-05
	AK000075	Homo sapiens cDNA FLJ20068 fis, clone COL01755	20.18	Down	5.17E-04
40	NM_032471	Homo sapiens protein kinase (cAMP-dependent, catalytic) inhibitor beta (PKIB), transcript variant 3, mRNA	20.17	Down	5.05E-04
	AK095776	Homo sapiens cDNA FLJ38457 fis, clone FEBRA2020400	19.57	Down	1.69E-05
		Homo sapiens a disintegrin-like and	19.13	Down	9.32E-05
45	NM_182920	metalloprotease (reprolysin type) with thrombospondin type 1 motif, 9 (ADAMTS9), transcript variant 1, mRNA			
	NM_002273	Homo sapiens keratin 8 (KRT8), mRNA	18.92	Down	2.49E-05
	NM_002246	Homo sapiens potassium channel, subfamily K, member 3 (KCNK3), mRNA	18.9	Down	1.35E-05
50	NM_024726	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	18.77	Down	5.32E-04
	AF055376	Homo sapiens short form transcription factor C-MAF (c-maf) mRNA, complete cds	18.38	Down	1.54E-03
55	AB033048	Homo sapiens mRNA for KIAA1222 protein, partial cds	18.16	Down	7.70E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	AW511222	hd44d11.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2912373 3 similar to contains Alu repetitive element, mRNA sequence	17.66	Down	7.79E-04
10	NM_004165	Homo sapiens Ras-related associated with diabetes (RRAD), mRNA	17.32	Down	3.10E-04
	NM_005202	Homo sapiens collagen, type VIII, alpha 2 (COL8A2), mRNA	17.22	Down	1.48E-04
15	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	17.01	Down	4.66E-06
	NM_005130	Homo sapiens fibroblast growth factor binding protein 1 (FGFBP1), MRNA	16.84	Down	9.53E-06
20	AK098071	Homo sapiens cDNA FLJ40752 fis, clone TRACH2000972	16.57	Down	1.90E-04
	AK024238	Homo sapiens cDNA FLJ14176 fis, clone NT2RP2003101	16.33	Down	1.60E-05
	R99527	yq79b11.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:201981 3, mRNA sequence	16.15	Down	7.97E-06
25	NM_021102	Homo sapiens serine protease inhibitor, Kunitz type, 2 (SPINT2), mRNA	15.87	Down	4.35E-05
	BQ003401	UI-H-EI1-azd-j-23-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847286 3, mRNA sequence	15.77	Down	2.50E-04
30	BC035656	Homo sapiens hypothetical protein LOC285835, mRNA (cDNA clone IMAGE:5588650), partial cds	15.49	Down	153E-06
	T56535	yb33g07.r1 Stratagene fetal spleen (#937205) Homo sapiens cDNA clone IMAGE:73020 5, mRNA sequence	15.46	Down	3.03E-05
35	NM_152864	Homo sapiens chromosome 20 open reading frame 58 (C20orf58), mRNA	15.18	Down	1.00E-03
	NM_001448	Homo sapiens glypican 4 (GPC4), mRNA	15.04	Down	2.32E-04
		Homo sapiens crumbs homolog 3 (Drosophila) (CRB3), transcript variant 2, mRNA	14.99	Down	1.08E-04
40	NM_139161	AGENCOURT_8840265 Lupski_sciatic_nerve Homo sapiens cDNA clone IMAGE:6205036 5, mRNA sequence	14.93	Down	1.30E-05
	BQ924832				
	NM_016276	Homo sapiens serum/glucocorticoid regulated kinase 2 (SGK2), transcript variant 2, mRNA	14.91	Down	2.20E-05
45	CA437861	UI-H-DH0-aur-k-12-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone UI-H-DH0-aur-k-12-0-UI 3, mRNA sequence	14.78	Down	9.11E-05
	H89526	yw28b04.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253519 5, mRNA sequence	14.74	Down	3.24E-05
50	BU633163	UI-H-FL1-bgt-n-07-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgt-n-07-0-UI 3, mRNA sequence	14.62	Down	3.96E-05
	NM_000519	Homo sapiens hemoglobin, delta (HBD), mRNA	14.61	Down	1.34E-05
55	NM_020962	Homo sapiens likely ortholog of mouse neighbor of Punc E11 (NOPE), mRNA	14.48	Down	6.78E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_004862	Homo sapiens lipopolysaccharide-induced TNF factor (LITAF), mRNA	14.37	Down	3.22E-04
10	NM_002837	Homo sapiens protein tyrosine phosphatase, receptor type, B (PTPRB), mRNA	14.3	Down	1.61E-05
	NM_024898	Homo sapiens family with sequence similarity 31, member C (FAM31C), mRNA	14.27	Down	3.50E-05
	AF269162	Homo sapiens c21orf7 form B mRNA, complete cds	14.09	Down	1.28E-04
15	NM_003385	Homo sapiens visinin-like 1 (VSNL1), mRNA	14.08	Down	1.92E-05
	BC041412	Homo sapiens heat shock 70kDa protein 12A, mRNA (cDNA clone IMAGE:5285193), partial cds	14.04	Down	2.40E-04
	AI493349	tg70f04.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2114143 3, mRNA sequence	13.92	Down	1.43E-05
20	BE295468	601174523F1 NIH_MGC_17 Homo sapiens cDNA clone IMAGE:3529924 5, mRNA sequence	13.86	Down	9.65E-04
	NM_014936	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 4 (putative function) (ENPP4), mRNA	13.83	Down	6.53E-05
25	NM_207517	Homo sapiens ADAMTS-like 3 (ADAMTSL3), mRNA	13.79	Down	3.05E-05
	NM_025245	Homo sapiens pre-B-cell leukemia transcription factor 4 (PBX4), mRNA	13.77	Down	2.73E-04
	NM_173508	Homo sapiens solute carrier family 35, member F3 (SLC35F3), mRNA	13.69	Down	3.06E-05
30	NM_002578	Homo sapiens p21 (CDKN1A)-activated kinase 3 (PAK3), mRNA	13.61	Down	1.46E-04
	NM_016588	Homo sapiens neuritin 1 (NRN1), mRNA	13.56	Down	6.51E-05
	NM_198389	Homo sapiens lung type-I cell membrane-associated glycoprotein (T1A-2), transcript variant 2, mRNA	13.52	Down	6.51E-05
35	NM_173505	Homo sapiens ankyrin repeat domain 29 (ANKRD29), mRNA	13.38	Down	5.46E-05
	NM_152573	Homo sapiens RAS and EF hand domain containing (RASEF), mRNA	13.3	Down	4.40E-04
40	NM_020130	Homo sapiens chromosome 8 open reading frame 4 (C8orf4), mRNA	13.11	Down	5.06E-04
	N63415	yy60d04.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:277927 3 similar to contains L1.b3 L1 repetitive element ;, mRNA sequence	13.03	Down	4.99E-04
45	BX648207	Homo sapiens mRNA; cDNA DKFZp686E16168 (from clone DKFZp686E16168)	12.84	Down	3.91E-04
	NM_178012	Homo sapiens tubulin, beta polypeptide paralog (MGC8685), mRNA	12.7	Down	2.28E-04
50	NM_152737	Homo sapiens hypothetical protein MGC33993 (MGC33993), mRNA	12.61	Down	9.53E-06
	AI126888	qb95d06.x1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:1707851 3, mRNA sequence	12.43	Down	4.50E-05
55	NM_005712	Homo sapiens HERV-H LTR-associating 1 (HHLA1), mRNA	12.39	Down	1.42E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	12.32	Down	2.77E-05
10	H23441	ym52f11.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:51888 3, mRNA sequence	12.28	Down	3.96E-05
	AL137698	Homo sapiens mRNA; cDNA DKFZp434C1915 (from clone DKFZp434C1915); partial cds	12.24	Down	5.21E-05
	AB023211	Homo sapiens mRNA for KIAA0994 protein, partial cds	12.15	Down	3.52E-05
15	AL389942	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 2005635	12.03	Down	1.88E-04
	AL359058	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 592473	11.96	Down	1.92E-05
20	NM_173549	Homo sapiens hypothetical protein FLJ39553 (FLJ39553), mRNA	11.93	Down	4.52E-05
	NM_002031	Homo sapiens fyn-related kinase (FRK), mRNA	11.64	Down	4.39E-05
	AB011095	Homo sapiens mRNA for KIAA0523 protein, partial cds	11.59	Down	2.43E-05
	NM_178033	Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1 (CYP4X1), mRNA	11.58	Down	7.70E-05
25	NM_003383	Homo sapiens very low density lipoprotein receptor (VLDLR), mRNA	11.43	Down	1.30E-04
	AK125490	Homo sapiens cDNA FLJ43501 fis, clone PEBLM2004497	11.25	Down	5.37E-04
30	NM_004982	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 8 (KCNJ8), mRNA	11	Down	5.37E-05
	NM_016307	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	10.96	Down	9.84E-04
	NM_019644	Homo sapiens ankyrin repeat domain 7 (ANKRD7), mRNA	10.92	Down	1.08E-04
35	AL834140	Homo sapiens mRNA; cDNA DKFZp434A2029 (from clone DKFZp434A2029)	10.85	Down	3.86E-04
	NM_020944	Homo sapiens glucosidase, beta (bile acid) 2 (GBA2), mRNA	10.85	Down	3.37E-04
40	CN478597	UI-CF-FN0-aeo-g-21-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aeo-g-21-0-UI 3, mRNA sequence	10.84	Down	3.83E-05
	AW603401	RC0-CN0025-010200-012-d01 CN0025 Homo sapiens cDNA, mRNA sequence	10.78	Down	6.72E-04
45		Homo sapiens cyclin-dependent kinase inhibitor 1C (p57, Kip2) (CDKN1C), mRNA	10.73	Down	1.42E-04
	NM_000076	qt94a11.x1 NCI_CGAP_Co14 Homo sapiens cDNA clone IMAGE:1962908 3 similar to gb:X74929 KERATIN, TYPE II CYTOSKELETAL 8 (HUMAN);, mRNA sequence	10.63	Down	1.08E-04
	AI355761				
50	AW172903	xj05e04.x1 NCI_CGAP_Ut2 Homo sapiens cDNA clone IMAGE:2656350 3, mRNA sequence	10.58	Down	9.44E-05
	AB041269	Homo sapiens mRNA for keratin 19, partial cds, isolate:K19-141	10.55	Down	1.35E-03
55	BC042028	Homo sapiens, clone IMAGE:4794726, mRNA	10.53	Down	2.71E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_030949	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 14C (PPP1R14C), mRNA	10.51	Down	2.17E-04
10	BX103476	BX103476 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998C053946 ; IMAGE:1557436, mRNA sequence	10.49	Down	4.88E-02
	NM_174900	Homo sapiens zinc finger protein 42 (ZFP42), mRNA	10.38	Down	1.88E-04
	NM_152768	Homo sapiens hypothetical protein FLJ25378 (FLJ25378), mRNA	10.35	Down	8.16E-04
15	NM_021977	Homo sapiens solute carrier family 22 (extraneuronal monoamine transporter), member 3 (SLC22A3), mRNA	10.27	Down	1.97E-04
	BF509573	UI-H-B14-apf-b-11-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086949 3, mRNA sequence	10.23	Down	1.43E-05
20					
	BX648964	Homo sapiens mRNA; cDNA DKFZp686J0156 (from clone DKFZp686J0156)	10.19	Down	6.44E-03
	NM_002338	Homo sapiens limbic system-associated membrane protein (LSAMP), mRNA	10.15	Down	7.21E-05
25	NM_006561	Homo sapiens CUG triplet repeat, RNA binding protein 2 (CUGBP2), mRNA	10.13	Down	1.75E-03
	AK123319	Homo sapiens cDNA FLJ41325 fis, clone BRAMY2046871	10.08	Down	3.96E-05
30	NM_004753	Homo sapiens dehydrogenase/reductase (SDR family) member 3 (DHRS3), mRNA	9.91	Down	1.59E-03
	BE788763	601475864F1 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:3879014 5, mRNA sequence	9.88	Down	3.14E-04
	NM_030594	Homo sapiens cytoplasmic polyadenylation element binding protein 1 (CPEB1), mRNA	9.86	Down	1.93E-03
35	NM_002462	Homo sapiens myxovirus (influenza virus) resistance 1, interferon-inducible protein p78 (mouse) (MX1), mRNA	9.8	Down	3.75E-04
	S69208	troponin T [human, skeletal and cardiac muscle, mRNA, 932 nt]	9.79	Down	1.10E-03
40	NM_005360	Homo sapiens v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian) (MAF), mRNA	9.76	Down	9.46E-04
	NM_030583	Homo sapiens matrilin 2 (MATN2), transcript variant 2, mRNA	9.67	Down	1.92E-05
45	NM_012464	Homo sapiens tolloid-like 1 (TLL1), mRNA	9.66	Down	1.76E-02
	NM_004490	Homo sapiens growth factor receptor-bound protein 14 (GRB14), mRNA	9.65	Down	8.22E-04
	NM_015236	Homo sapiens latrophilin 3 (LPHN3), mRNA	9.65	Down	3.96E-05
	AK024261	Homo sapiens cDNA FLJ14199 fis, clone NT2RP3002713	9.62	Down	1.63E-04
50	NM_207482	Homo sapiens FLJ44048 protein (FLJ44048), mRNA	9.49	Down	6.53E-05
	BM969191	UI-CF-EN0-acp-e-22-0-UI.s1 UI-CF-EN0 Homo sapiens cDNA clone UI-CF-EN0-acp-e-22-0-UI 3, mRNA sequence	9.4	Down	2.21E-04
55	M60502	Human profilaggrin mRNA, 3 end	9.34	Down	5.78E-05
	NM_001710	Homo sapiens B-factor, properdin (BF), mRNA	9.32	Down	8.70E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_005114	Homo sapiens heparan sulfate (glucosamine) 3-O-sulfotransferase 1 (HS3ST1), mRNA	9.3	Down	1.05E-04
10	NM_003985	Homo sapiens tyrosine kinase, non-receptor, 1 (TNK1), mRNA	9.3	Down	1.00E-04
	NM_001874	Homo sapiens carboxypeptidase M (CPM), transcript variant 1, mRNA	9.24	Down	5.54E-04
	AJ697972	Homo sapiens chromosome 3 cDNA	9.15	Down	3.55E-04
15	BM701989	UI-E-CQ1-aex-j-06-0-UI.r1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aex-j-06-0-UI 5, mRNA sequence	9.09	Down	8.72E-03
	BX640973	Homo sapiens mRNA; cDNA DKFZp686B15184 (from clone DKFZp686B15184)	9.09	Down	1.29E-04
20	NM_002313	Homo sapiens actin binding LIM protein 1 (ABLM1), transcript variant 1, mRNA	9.05	Down	1.23E-04
	BX486208	DKFZp686J07250_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686J07250 5, mRNA sequence	8.94	Down	9.28E-04
25	NM_004024	Homo sapiens activating transcription factor 3 (ATF3), mRNA	8.93	Down	9.73E-04
	S70348	Homo sapiens integrin beta 3 mRNA, partial cds, alternatively spliced	8.92	Down	6.01E-05
	NM_198495	Homo sapiens CTAGE family, member 4 (CTAGE4), mRNA	8.89	Down	1.28E-04
30	D86975	Homo sapiens mRNA for KIAA0222 gene, partial cds	8.88	Down	3.55E-04
	NM_001854	Homo sapiens collagen, type XI, alpha 1 (COL11A1), transcript variant A, mRNA	8.87	Down	6.78E-05
	NM_000212	Homo sapiens integrin, beta 3 (platelet glycoprotein IIIa, antigen CD61) (ITGB3), mRNA	8.84	Down	7.53E-05
35	AL117578	Homo sapiens mRNA; cDNA DKFZp434C128 (from clone DKFZp434C128)	8.8	Down	3.22E-04
	NM_198174	Homo sapiens transcription factor CP2-like 4 (TFCP2L4), transcript variant 3, mRNA	8.75	Down	8.93E-05
40	BU727096	UI-E-CR0-ach-e-12-0-UI.s1 UI-E-CR0 Homo sapiens cDNA clone UI-E-CR0-ach-e-12-0-UI 3, mRNA sequence	8.71	Down	5.91E-04
	NM_005949	Homo sapiens metallothionein 1 F (functional) (MT1 F), mRNA	8.67	Down	3.29E-03
45	BX537613	Homo sapiens mRNA; cDNA DKFZp686E11117 (from clone DKFZp686E11117)	8.65	Down	7.70E-05
	NM_014358	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	8.57	Down	7.11E-05
50	NM_000817	Homo sapiens glutamate decarboxylase 1 (brain, 67kDa) (GAD1 transcript variant GAD67, mRNA	8.55	Down	2.77E-05
	NM_005450	Homo sapiens noggin (NOG), mRNA	8.49	Down	2.75E-03
	NM_144707	Homo sapiens prominin 2 (PROM2), mRNA	8.49	Down	2.72E-04
55	AK074181	Homo sapiens mRNA for FLJ00254 protein	8.46	Down	7.83E-05
	NM_032882	Homo sapiens paraneoplastic antigen like 6A (PNMA6A), mRNA	8.44	Down	2.57E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_004479	Homo sapiens fucosyltransferase 7 (alpha (1,3) fucosyltransferase) (FUT7), mRNA	8.39	Down	1.73E-05
10	NM_000856	Homo sapiens guanylate cyclase 1, soluble, alpha 3 (GUCY1A3), mRNA	8.31	Down	1.63E-04
	NM_016542	Homo sapiens Mst3 and SOK1-related kinase (MST4), mRNA	8.26	Down	6.03E-04
15	NM_021044	Homo sapiens desert hedgehog homolog (Drosophila) (DHH), mRNA	8.23	Down	6.01E-04
	NM_000313	Homo sapiens protein S (alpha) (PROS1), mRNA	8.23	Down	4.08E-04
	NM_000227	Homo sapiens laminin, alpha 3 (LAMA3), transcript variant 2, mRNA	8.15	Down	2.89E-05
20	AA010611	zi09f09.s1 Soares_fetal_liver_spleen_1NFLS_S1Homo sapiens cDNA clone IMAGE:430313 3, mRNA sequence	8.13	Down	4.99E-04
	NM_004433	Homo sapiens E74-like factor 3 (ets domain transcription factor, epithelial-specific) (ELF3), mRNA	8.07	Down	6.84E-05
	NM_000860	Homo sapiens hydroxyprostaglandin dehydrogenase 15-(NAD) (HPGD), mRNA	8.02	Down	1.49E-04
25	BG208475	RST27977 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	8	Down	2.90E-03
	N75271	yz74h12.r1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:288839 5, mRNA sequence	8	Down	3.67E-03
30	NM_022912	Homo sapiens chromosome 2 open reading frame 23 (C2orf23), mRNA	8	Down	9.93E-05
	AL706653	DKFZp686E1543_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686E1543 5, mRNA sequence	7.98	Down	9.86E-05
35	AA411988	zt65g11.s1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:727268 3, mRNA sequence	7.95	Down	9.11E-05
	NM_020152	Homo sapiens chromosome 21 open reading frame 7 (C21orf7), mRNA	7.95	Down	4.35E-05
40	AK125608	Homo sapiens cDNA FLJ43620 fis, clone SPLEN2021701, highly similar to HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN PRECURSOR	7.93	Down	1.87E-03
45	CA306881	UI-H-FT1-bht-n-22-0-UI.s1 NCI_CGAP_FT1 Homo sapiens cDNA clone UI-H-FT1-bht-n-22-0-UI 3, mRNA sequence	7.9	Down	7.95E-04
	AI640484	wa27f01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2299321 3, mRNA sequence	7.87	Down	3.33E-04
50	AI819863	wj45h05.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2405817 3, mRNA sequence	7.86	Down	5.42E-05
	AK124702	Homo sapiens cDNA FLJ42712 fis, clone BRAMY3008044	7.86	Down	2.13E-04
	NM_014333	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	7.8	Down	5.41E-03
55	H94320	yv18b10.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:243067 3, mRNA sequence	7.8	Down	4.43E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_022103	Homo sapiens hypothetical zinc finger protein FLJ14011 (FLJ14011), mRNA	7.77	Down	3.50E-04
10	NM 016179	Homo sapiens transient receptor potential cation channel, subfamily C, member 4 (TRPC4), mRNA	7.75	Down	6.04E-04
	AF108093	Homo sapiens IA-2 gene, intron 18	7.74	Down	1.92E-05
	NM_001332	Homo sapiens catenin (cadherin-associated protein), delta 2 (neural plakophilin-related arm-repeat protein) (CTNND2), mRNA	7.73	Down	5.06E-04
15					
	NM_002206	Homo sapiens integrin, alpha 7 (ITGA7), mRNA	7.73	Down	2.33E-03
	BF510493	UI-H-B14-apa-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086558 3, mRNA sequence	7.68	Down	2.89E-05
20					
	BU951469	in60a05.x3 HR85 islet Homo sapiens cDNA clone IMAGE:6126249 3, mRNA sequence	7.63	Down	1.74E-04
	AI819186	wj32d10.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2404531 3, mRNA sequence	7.62	Down	5.86E-04
25					
	NM_013259	Homo sapiens transgelin 3 (TAGLN3), transcript variant 1, mRNA	7.61	Down	3.22E-04
	AI942360	wo80c06.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2461642 3, mRNA sequence	7.57	Down	6.84E-05
	BX098521	BX098521 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGp998L05118 ; IMAGE:123412, mRNA sequence	7.55	Down	1.31E-05
30					
	NM_152495	Homo sapiens cornichon homolog 3 (Drosophila) (CNIH3), mRNA	7.55	Down	5.20E-05
	NM_018728	Homo sapiens myosin VC (MYO5C), mRNA	7.54	Down	1.30E-04
35					
	NM_004086	Homo sapiens coagulation factor C homolog, coxlin (Limulus polyphemus) (COCH), mRNA	7.51	Down	5.92E-04
	BM683698	UI-E-EJ1-ajh-k-08-0-UI.s1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajh-k-08-0-UI 3, mRNA sequence	7.48	Down	6.46E-05
	NM_000593	Homo sapiens transporter 1, ATP-binding cassette, sub-family B (MDR/TAP) (TAP1), mRNA	7.47	Down	1.05E-04
40					
		Homo sapiens cDNA FLJ36973 fis, clone	7.44	Down	5.42E-05
	AK094292	BRACE2006249			
	NM_031442	Homo sapiens transmembrane 4 superfamily member 10 (TM4SF10), mRNA	7.42	Down	1.87E-03
45					
	AK074097	Homo sapiens mRNA for FLJ00168 protein	7.41	Down	4.15E-04
	NM_005302	Homo sapiens G protein-coupled receptor 37 (endothelin receptor type B-like) (GPR37), mRNA	7.33	Down	1.15E-03
	NM_018349	Homo sapiens multiple C2-domains with two transmembrane regions 2 (MCTP2), mRNA	7.3	Down	5.46E-05
50					
	NM_014399	Homo sapiens transmembrane 4 superfamily member 13 (TM4SF13), mRNA	7.28	Down	7.73E-03
	NM_002899	Homo sapiens retinol binding protein 1, cellular (RBP1), mRNA	7.28	Down	1.79E-03
55					
	NM_015441	Homo sapiens olfactomedin-like 2B (OLFML2B), mRNA	7.24	Down	5.84E-04
	NM_206808	Homo sapiens citrate lyase beta like (CLYBL), transcript variant 2, mRNA	7.22	Down	9.35E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_207380	Homo sapiens FLJ43339 protein (FLJ43339), mRNA	7.19	Down	1.69E-03
	NM_000147	Homo sapiens fucosidase, alpha-L- 1, tissue (FUCA1), mRNA	7.17	Down	4.23E-02
10					
	NM_002670	Homo sapiens plastin 1 (I isoform) (PLS1), mRNA	7.16	Down	2.28E-02
	NM_000620	Homo sapiens nitric oxide synthase 1 (neuronal) (NOS1), mRNA	7.15	Down	1.46E-04
	NM_018242	Homo sapiens hypothetical protein FLJ10847 (FLJ10847), mRNA	7.14	Down	1.48E-03
15					
	N78460	yz76h06.r1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:289019 5, mRNA sequence	7.11	Down	2.28E-02
	NM_001001430	Homo sapiens troponin T2, cardiac (TNNT2), transcript variant 2, mRNA	7.04	Down	9.11E-05
20					
	NM_003979	Homo sapiens G protein-coupled receptor, family C, group 5, member A (GPCR5A), mRNA	6.95	Down	4.47E-05
	NM_003328	Homo sapiens TXK tyrosine kinase (TXK), mRNA	6.94	Down	1.44E-04
25					
	AK128715	Homo sapiens cDNA FLJ46882 fis, clone UTERU3015844	6.94	Down	3.34E-03
	NM_000961	Homo sapiens prostaglandin I2 (prostacyclin) synthase (PTGIS), mRNA	6.91	Down	5.88E-04
	NM_052839	Homo sapiens pannexin 2 (PANX2), mRNA	6.9	Down	7.48E-05
30					
	AF119903	Homo sapiens PRO2834 mRNA, complete cds	6.89	Down	2.43E-05
	NM_001505	Homo sapiens G protein-coupled receptor 30 (GPR30), mRNA	6.84	Down	1.15E-03
	AW137116	UI-H-B1I-acp-f-03-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2715029 3, mRNA sequence	6.83	Down	3.08E-05
35					
	NM_052960	Homo sapiens retinol binding protein 7, cellular (RBP7), mRNA	6.79	Down	3.15E-03
	NM_003107	Homo sapiens SRY (sex determining region Y)-box 4 (SOX4), mRNA	6.77	Down	3.33E-04
40					
	NM_004170	Homo sapiens solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1 (SLC1A1), mRNA	6.77	Down	3.33E-04
	CB115754	K-EST0159876 L8SCK0 Homo sapiens cDNA clone L8SCK0-8-H08 5, mRNA sequence	6.75	Down	1.69E-05
45					
	AK026740	Homo sapiens cDNA: FLJ23087 fis, clone LNG06994, highly similar to AF161368 Homo sapiens HSPC105 mRNA	6.75	Down	1.05E-04
	NM_002487	Homo sapiens necdin homolog (mouse) (NDN), mRNA	6.75	Down	6.29E-04
50					
	M27161	Human MHC class I CD8 alpha-chain (Leu-2/T8) gene, complete cds	6.73	Down	4.96E-03
	AK025003	Homo sapiens cDNA: FLJ21350 fis, clone COL02751	6.7	Down	4.15E-04
	NM_000693	Homo sapiens aldehyde dehydrogenase 1 family, member A3 (ALDH1A3), mRNA	6.69	Down	1.41E-02
55					
	NM_014900	Homo sapiens COBL-like 1 (COBLL1), mRNA	6.67	Down	1.17E-02
	BG545305	602572521 F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4700644 5, mRNA sequence	6.66	Down	8.24E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	NM_014421	Homo sapiens dickkopf homolog 2 (Xenopus laevis) (DKK2), mRNA	6.66	Down	2.17E-02
10	NM_005139	Homo sapiens annexin A3 (ANXA3), mRNA	6.64	Down	2.04E-04
	BX111520	BX111520 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998L15208 ; IMAGE:141470, mRNA sequence	6.64	Down	7.70E-05
	AK090808	Homo sapiens cDNA FLJ33489 fis, clone BRAMY2003585	6.63	Down	1.08E-04
15	D62831	HUM330B12B Clontech human aorta polyA+ mRNA (#6572) Homo sapiens cDNA clone GEN-330B12 5, mRNA sequence	6.58	Down	1.02E-03
	H18652	ym45e04.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:50948 3, mRNA sequence	6.57	Down	1.00E-03
20		UI-H-EI1-ayu-m-09-0-UI.s1 NCI_CGAP_EI1	6.56	Down	1.26E-04
	BQ002165	Homo sapiens cDNA clone IMAGE:5843888 3, mRNA sequence			
25	NM_018076	Homo sapiens armadillo repeat containing 4 (ARMC4), mRNA	6.54	Down	1.72E-04
	H41942	yo60a08.r1 Soares breast 3NbHBst Homo sapiens cDNA clone IMAGE:182294 5 similar to contains Alu repetitive element;contains LTR9 repetitive element ;, mRNA sequence	6.53	Down	3.48E-03
30	AJ318805	AJ318805 Homo sapiens adipose tissue Homo sapiens cDNA clone 2040, mRNA sequence	6.5	Down	1.54E-04
	NM_032866	Homo sapiens cingulin-like 1 (CGNL1), mRNA	6.46	Down	1.84E-03
	CA434164	UI-H-DHO-arv-a-17-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone UI-H-DHO-arv-a-17-0-UI 3, mRNA sequence	6.44	Down	2.82E-04
35					
	AW006864	ws15d04.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2497255 3, mRNA sequence	6.39	Down	2.77E-05
	NM_014932	Homo sapiens neuroligin 1 (NLGN1), mRNA	6.39	Down	2.59E-05
40	BX538226	Homo sapiens mRNA; cDNA DKFZp686E1944 (from clone DKFZp686E1944)	6.38	Down	1.74E-04
	NM_016651	Homo sapiens dapper homolog 1, antagonist of beta-catenin (xenopus) (DACT1), mRNA	6.38	Down	5.05E-04
	NM_015678	Homo sapiens neurobeachin (NBEA), mRNA	6.34	Down	4.17E-03
45	NM_022842	Homo sapiens CUB domain-containing protein 1 (CDCP1), transcript variant 1, mRNA	6.32	Down	1.12E-03
	AF130079	Homo sapiens clone FLC0578 PRO2852 mRNA, complete cds	6.32	Down	1.74E-02
50	AL117425	Homo sapiens mRNA; cDNA DKFZp566L203 (from clone DKFZp566L203)	6.29	Down	8.54E-05
	NM_005141	Homo sapiens fibrinogen, B beta polypeptide (FGB), mRNA	6.28	Down	3.43E-04
	AW590139	hg33d12.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2947415 3, mRNA sequence	6.27	Down	1.29E-04
55	BE968596	601649770F1 NIH_MGC_74 Homo sapiens cDNA clone IMAGE:3933472 5, mRNA sequence	6.26	Down	6.39E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	AF220263	Homo sapiens MOST2 mRNA, complete cds	6.25	Down	1.10E-03
	NM_000876	Homo sapiens insulin-like growth factor 2 receptor (IGF2R), mRNA	6.25	Down	1.08E-03
10					
	NM_006681	Homo sapiens neuromedin U (NMU), mRNA	6.25	Down	1.07E-04
	AK024865	Homo sapiens cDNA: FLJ21212 fis, clone COL00502	6.24	Down	1.05E-04
	NM_024508	Homo sapiens zinc finger, BED domain containing 2 (ZBED2), mRNA	6.22	Down	8.82E-04
15					
	NM_177949	Homo sapiens armadillo repeat containing, X-linked 2 (ARMCX2), mRNA	6.18	Down	1.20E-02
	NM_001257	Homo sapiens cadherin 13, H-cadherin (heart) (CDH13), mRNA	6.17	Down	8.46E-03
20					
	NM_001200	Homo sapiens bone morphogenetic protein 2 (BMP2), mRNA	6.15	Down	3.33E-04
	AK125695	Homo sapiens cDNA FLJ43707 fis, clone TESOP2001865	6.15	Down	6.53E-05
	BC033567	Homo sapiens, clone IMAGE:4822266, mRNA	6.14	Down	3.14E-04
25					
	AB011539	Homo sapiens mRNA for MEGF6 protein (KIAA0815), partial cds	6.14	Down	1.49E-03
	NM_024697	Homo sapiens hypothetical protein FLJ22419 (FLJ22419), mRNA	6.13	Down	2.72E-04
	BC040701	Homo sapiens cDNA clone IMAGE:5736259, partial cds	6.13	Down	1.89E-03
30					
	NM_001889	Homo sapiens crystallin, zeta (quinone reductase) (CRYZ), mRNA	6.12	Down	1.97E-02
	BX108092	BX108092 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998_164460 ; IMAGE:1754967, mRNA sequence	6.08	Down	1.01E-04
35					
	T65315	yc79h02.s1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:22228 3, mRNA sequence	6.08	Down	1.52E-04
	BX095811	BX095811 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998E084494 ; IMAGE:1837423, mRNA sequence	6.07	Down	1.68E-04
40					
	NM_012413	Homo sapiens glutaminyl-peptide cyclotransferase (glutaminyl cyclase) (QPCT), mRNA	6.07	Down	4.25E-02
	AI311296	ta48d10.x2 NCI_CGAP_Lu25 Homo sapiens cDNA clone IMAGE:2047315 3, mRNA sequence	6.07	Down	4.15E-04
45					
	NM_138811	Homo sapiens chromosome 7 open reading frame 31 (C7orf31), mRNA	6.06	Down	1.54E-04
	BM988642	UI-H-DH0-arx-p-21-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone IMAGE:5855492 3, mRNA sequence	6.05	Down	1.28E-04
50					
	NM_016613	Homo sapiens hypothetical protein DKFZp434L142 (DKFZp434L142), mRNA	6.04	Down	1.25E-02
	W20132	zb40c10.r1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:306066 5, mRNA sequence	6.01	Down	7.83E-05
	NM_001785	Homo sapiens cytidine deaminase (CDA), mRNA	6.01	Down	1.16E-04
55					
		Homo sapiens RAB11 family interacting protein 1 (class I) (RAB11FIP1), transcript variant 1, mRNA	6	Down	1.76E-02
	NM_025151				
	AF519622	Homo sapiens noncoding mRNA sequence	5.98	Down	1.26E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_152284	Homo sapiens Snf7 homologue associated with Alix 3 (Shax3), mRNA	5.97	Down	7.70E-05
10	D29453	HUMNK566 Human epidermal keratinocyte Homo sapiens cDNA clone 566, mRNA sequence	5.97	Down	3.58E-04
	AI220066	qg84d01.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1841857 3, mRNA sequence	5.96	Down	9.78E-04
	AK026235	Homo sapiens cDNA: FLJ22582 fis, clone HSI02576	5.94	Down	1.10E-03
15	BX102869	BX102869 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGp998P17421 ; IMAGE:214528, mRNA sequence	5.92	Down	2.81E-04
	NM_002354	Homo sapiens tumor-associated calcium signal transducer 1 (TACSTD1), mRNA	5.92	Down	6.65E-04
20	CA313095	UI-CF-FN0-aex-f-01-0-UI.s1 UI-CF-FNO Homo sapiens cDNA clone UI-CF-FN0-aex-f-01-0-UI 3, mRNA sequence	5.91	Down	1.51E-04
	BX641086	Homo sapiens mRNA; cDNA DKFZp686G2469 (from clone DKFZp686G2469)	5.9	Down	1.07E-04
25	AK021493	Homo sapiens cDNA FLJ11431 fis, clone HEMBA1001094	5.88	Down	2.79E-04
	NM_152369	Homo sapiens hypothetical protein MGC45474 (MGC45474), mRNA	5.87	Down	4.40E-04
30	L07615	Human neuropeptide Y receptor Y1 (NPYY1) mRNA, exon 2-3 and complete cds	5.87	Down	3.91E-04
	BX503694	DKFZp686M02112_s1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686M02112 3, mRNA sequence	5.84	Down	2.17E-04
35	NM_021101	Homo sapiens claudin 1 (CLDN1), mRNA	5.82	Down	1.24E-04
	NM_005737	Homo sapiens ADP-ribosylation factor-like 7 (ARL7), mRNA	5.8	Down	3.58E-04
	NM_005264	Homo sapiens GDNF family receptor alpha 1 (GFRA1), transcript variant 1, mRNA	5.8	Down	4.54E-05
40	AB020640	Homo sapiens mRNA for KIAA0833 protein, partial cds	5.79	Down	7.11E-05
	AK022598	Homo sapiens cDNA FLJ12536 fis, clone NT2RM4000265	5.78	Down	2.20E-04
	NM_173660	Homo sapiens hypothetical protein FLJ33718 (FLJ33718), mRNA	5.76	Down	2.95E-04
45	NM_052923	Homo sapiens zinc finger protein 452 (ZNF452), mRNA	5.71	Down	3.90E-03
	NM_000797	Homo sapiens dopamine receptor D4 (DRD4), mRNA	5.71	Down	4.44E-03
	NM_014917	Homo sapiens netrin G1 (NTNG1), mRNA	5.7	Down	2.50E-04
	NM_053039	Homo sapiens UDP glycosyltransferase 2 family, polypeptide B28 (UGT2B28), mRNA	5.7	Down	9.11E-05
50	NM_006266	Homo sapiens ral guanine nucleotide dissociation stimulator (RALGDS), mRNA	5.68	Down	9.21E-03
	NM_003264	Homo sapiens toll-like receptor 2 (TLR2), mRNA	5.67	Down	1.51E-04
55	AA017035	ze37c10.s1 Soares retina N2b4HR Homo sapiens cDNA clone IMAGE:361170 3 similar to contains Alu repetitive element;contains element MER37 repetitive element ;, mRNA sequence	5.66	Down	1.17E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_005901	Homo sapiens SMAD, mothers against DPP homolog 2 (Drosophila) (SMAD2), transcript variant 1, mRNA	5.65	Down	3.84E-02
10	NM_000576	Homo sapiens interleukin 1, beta (IL1B), mRNA	5.64	Down	6.69E-05
	NM_013322	Homo sapiens sorting nexin 10 (SNX10), mRNA	5.63	Down	4.81E-04
	AJ406941	Homo sapiens partial mRNA for keratin associated protein 4.9 (KRTAP4.9 gene)	5.62	Down	5.80E-05
15	NM_001562	Homo sapiens interleukin 18 (interferon-gamma-inducing factor) (IL18), mRNA	5.61	Down	5.82E-03
	NM_173567	Homo sapiens abhydrolase domain containing 7 (ABHD7), mRNA	5.59	Down	1.18E-04
	F36108	HSPD33448 HM3 Homo sapiens cDNA clone SH4-000003-1/E09, mRNA sequence	5.57	Down	7.05E-04
20	NM_145804	Homo sapiens ankyrin repeat and BTB (POZ) domain containing 2 (ABTB2), mRNA	5.56	Down	1.76E-03
	BQ002466	UI-H-EI1-ayw-k-24-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5844623 3, mRNA Sequence	5.56	Down	1.46E-04
25	NM_173662	Homo sapiens hypothetical protein LOC285533 (LOC285533), mRNA	5.56	Down	3.01E-04
	NM_000129	Homo sapiens coagulation factor XIII, A1 polypeptide (F13A1), mRNA	5.53	Down	1.05E-04
30	NM_000584	Homo sapiens interleukin 8 (IL8), mRNA	5.51	Down	1.82E-03
	BM664445	UI-E-CL1-afa-p-05-0-UI.s1 UI-E-CL1 Homo sapiens cDNA clone UI-E-CL1-afa-p-05-0-UI 3, mRNA sequence	5.5	Down	6.13E-04
	NM_006863	Homo sapiens leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 1 (LILRA1), mRNA	5.48	Down	3.09E-03
35	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	5.46	Down	5.71E-03
	BC033124	Homo sapiens, clone IMAGE:2960615, mRNA	5.46	Down	3.64E-04
40	NM_014978	Homo sapiens sortilin-related VPS10 domain containing receptor 3 (SORCS3), mRNA	5.46	Down	1.26E-04
	AF007143	Homo sapiens clone 23738 mRNA sequence	5.46	Down	2.05E-04
	NM_001089	Homo sapiens ATP-binding cassette, sub-family A (ABC1), member 3 (ABCA3), mRNA	5.46	Down	4.75E-05
45	NM_004524	Homo sapiens lethal giant larvae homolog 2 (Drosophila) (LLGL2), mRNA	5.46	Down	2.73E-04
	AW591461	x192h06.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2682203 3, mRNA sequence	5.45	Down	1.29E-04
	NM_007072	Homo sapiens HERV-H LTR-associating 2 (HHLA2), mRNA	5.45	Down	1.18E-03
50	NM_181718	Homo sapiens hypothetical protein LOC253982 (LOC253982), mRNA	5.44	Down	6.64E-03
	AI559193	tq42h01.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2211505 3, mRNA sequence	5.44	Down	2.64E-03
55	BF058471	7k31b04.x1 NCI_CGAP_Ov18 Homo sapiens cDNA clone IMAGE:3477054 3 similar to contains element MER17 repetitive element ;, mRNA sequence	5.43	Down	4.72E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5					
	AK130181	Homo sapiens cDNA FLJ26671 fis, clone MPG03325	5.42	Down	4.50E-05
	NM_178039	Homo sapiens Rab6-interacting protein 2 (ELKS), transcript variant delta, mRNA	5.41	Down	9.73E-04
10	NM_019025	Homo sapiens spermine oxidase (SMOX), transcript variant 5, mRNA	5.4	Down	1.11E-02
	BX108809	BX108809 Soares fetal liver spleen 1 N FLS Homo sapiens cDNA clone IMAGp998L05373 ; IMAGE:195988, mRNA sequence	5.4	Down	4.43E-02
15	NM_019607	Homo sapiens hypothetical protein FLJ11267 (FLJ11267), mRNA	5.4	Down	1.76E-02
	NM_001343	Homo sapiens disabled homolog 2, mitogen-responsive phosphoprotein (Drosophila) (DAB2), mRNA	5.39	Down	3.11E-03
20	NM_000087	Homo sapiens cyclic nucleotide gated channel alpha 1 (CNCA1), mRNA	5.34	Down	6.52E-04
	NM_031894	Homo sapiens ferritin, heavy polypeptide-like 17 (FTHL17), mRNA	5.34	Down	4.47E-03
25	NM_031419	Homo sapiens nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta (NFKBIZ), transcript variant 1, mRNA	5.33	Down	7.27E-03
	BG199496	RST18780 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	5.3	Down	4.14E-04
30	NM_001958	Homo sapiens eukaryotic translation elongation factor 1 alpha 2 (EEF1A2), mRNA	5.29	Down	1.97E-04
	NM_001878	Homo sapiens cellular retinoic acid binding protein 2 (CRABP2), mRNA	5.29	Down	5.17E-04
	AW014126	UI-H-BI0-aaj-a-05-0-UI.s1 NCI_CGAP_Sub1 Homo sapiens cDNA clone IMAGE:2709393 3, mRNA sequence	5.29	Down	9.98E-04
35	NM_006072	Homo sapiens chemokine (C-C motif) ligand 26 (CCL26), mRNA	5.28	Down	5.45E-03
	X70287	H.sapiens gene for thioredoxin, exons 2 and 3	5.27	Down	2.75E-02
40	NM_014867	Homo sapiens KIAA0711 gene product (KIAA0711), mRNA	5.27	Down	1.44E-03
	BX100098	BX100098 Soares_pregnant uterus_NbHPU Homo sapiens cDNA clone IMAGp998C021198 ; IMAGE:502201, mRNA sequence	5.26	Down	2.59E-04
45	AK055334	Homo sapiens cDNA FLJ30772 fis, clone FEBRA2000757, moderately similar to Homo sapiens BM-009 mRNA	5.24	Down	8.92E-04
	BU567804	AGENCOURT_10398872 NH_MGC_82 Homo sapiens cDNA clone IMAGE:6614502 5, mRNA sequence	5.24	Down	2.28E-02
50	NM_006332	Homo sapiens interferon, gamma-inducible protein 30 (IFI30), mRNA	5.24	Down	1.75E-03
	BF111903	7138d07.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3523644 3, mRNA sequence	5.23	Down	4.43E-05
55	NM_006727	Homo sapiens cadherin 10, type 2 (T2-cadherin) (CDH10), mRNA	5.2	Down	1.84E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-I vs AF-II	Direction	adj. p-value
5	NM_003810	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	5.19	Down	2.91E-03
10	AK127421	Homo sapiens cDNA FLJ45513 fis, clone BRTHA2021450	5.18	Down	8.00E-05
	NM_024897	Homo sapiens progesterone and adiponectin receptor family member VI (PAQR6), transcript variant 1, mRNA	5.18	Down	2.69E-05
	BU729783	UI-E-CK1-afh-h-18-0-UI.s1 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-afh-h-18-0-UI 3, mRNA sequence	5.17	Down	1.52E-04
15	BX107838	BX107838 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998A153853 ; IMAGE:1521686, mRNA sequence	5.16	Down	4.26E-04
20	NM_004335	Homo sapiens bone marrow stromal cell antigen 2 (BST2), mRNA	5.16	Down	7.11E-05
	AK021637	Homo sapiens cDNA FLJ11575 fis, clone HEMBA1003531	5.15	Down	6.53E-05
	NM_014890	Homo sapiens downregulated in ovarian cancer 1 (DOC1), transcript variant 2, mRNA	5.15	Down	9.70E-05
25	NM_003E97	Homo sapiens TGFB inducible early growth response 2 (TIEG2), mRNA	5.14	Down	1.19E-03
	AI684824	wa86a12.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2303038 3, mRNA sequence	5.13	Down	9.14E-04
30	NM_002522	Homo sapiens neuronal pentraxin I (NPTX1), mRNA	5.12	Down	1.52E-03
	NM_002250	Homo sapiens potassium intermediate/small conductance calcium-activated channel, subfamily N, member 4 (KCNN4), mRNA	5.12	Down	1.28E-04
	NM_001305	Homo sapiens claudin 4 (CLDN4), mRNA	5.11	Down	3.26E-03
35	BC015108	Homo sapiens, Similar to otoconin 90, clone IMAGE:4044247, mRNA	5.1	Down	2.17E-04
	N51335	yz15e08.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:283142 3, mRNA sequence	5.1	Down	2.89E-05
40	NM_001002295	Homo sapiens GATA binding protein 3 (GATA3), transcript variant 1, mRNA	5.06	Down	3.85E-03
	NM_013281	Homo sapiens fibronectin leucine rich transmembrane protein 3 (FLRT3), transcript variant 1, mRNA	5.05	Down	5.04E-03
45	AI623139	tu89b07.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2258197 3, mRNA sequence	5.04	Down	4.25E-03
	NM_022168	Homo sapiens interferon induced with helicase C domain 1 (IFIH1), mRNA	5.04	Down	2.48E-03
	BC05172 7	Homo sapiens cDNA clone IMAGE:5265929, partial cds	5.02	Down	5.42E-04

50

55

EP 1 780 267 B1

TABLE VII F: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT LEAST 5 FOLD IN AF-I VERSUS AF-III CELLS

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_080872	Homo sapiens unc-5 homolog D (C. elegans) (UNC5D), mRNA	331.21	Up	5.22E-06
10	NM_000090	Homo sapiens collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant) (COL3A1), mRNA	310.82	Up	3.56E-06
	NM_138961	Homo sapiens endothelial cell adhesion molecule (ESAM), mRNA	262	Up	6.27E-06
15	AK021531	Homo sapiens cDNA FLJ11469 fis, clone HEMBA1001658	205.55	Up	3.10E-06
	NM_000474	Homo sapiens twist homolog 1 (acrocephalosyndactyly 3; Saethre-Chotzen syndrome) (Drosophila) (TWIST1), mRNA	116.12	Up	2.29E-05
20	BX089554	BX089554 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998P07210 ; IMAGE:142326, mRNA sequence	97.02	Up	3.10E-06
	NM_018013	Homo sapiens hypothetical protein FLJ10159 (FLJ10159), mRNA	88.82	Up	3.10E-06
25	M_182798	Homo sapiens hypothetical protein FLJ39155 (FLJ39155), transcript variant 2, mRNA	86.73	Up	3.10E-06
	NM_052954	Homo sapiens cysteine and tyrosine-rich 1 (CYR1), mRNA	85.9	Up	3.10E-06
30	NM_000867	Homo sapiens 5-hydroxytryptamine (serotonin) receptor 2B (HTR2B), mRNA	74.02	Up	3.19E-06
	NM_014178	Homo sapiens syntaxin binding protein 6 (amisyn) (STXBP6), mRNA	71.46	Up	8.25E-06
	U83115	Human non-lens beta gamma-crystallin like protein (AIM1) mRNA, partial cds	66.41	Up	8.66E-06
35	CA429135	UI-H-FH1-bfh-k-22-0-UI.s1 NCI CGAP FH1 Homo sapiens cDNA clone UI-H-FH1-bfh-k-22-0-UI 3, mRNA sequence	66.03	Up	2.19E-06
	BQ003501	UI-H-EI1-azd-p-06-0-UI.s1 NCI CGAP EI1 Homo sapiens cDNA clone IMAGE:5847413 3, mRNA sequence	54.48	Up	1.35E-05
40	BX105152	BX105152 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998J212575 ; IMAGE:1031156, mRNA sequence	50.41	Up	5.63E-06
45	NM_006393	Homo sapiens nebulin (NEBL), transcript variant 1, mRNA	50.35	Up	1.43E-05
	NM_001432	Homo sapiens epiregulin (EREG), mRNA	49.79	Up	6.21E-06
	AI124557	am58g02.x1 Johnston frontal cortex Homo sapiens cDNA clone IMAGE:1539794 3, mRNA sequence	45.53	Up	3.19E-06
50	NM_018286	Homo sapiens hypothetical protein FLJ10970 (FLJ10970), mRNA	44.54	Up	1.84E-05
	AI962169	wq45c10.x1 NCI CGAP GC6 Homo sapiens cDNA clone IMAGE:2474226 3, mRNA sequence	44.13	Up	3.20E-05
55	NM_003381	Homo sapiens vasoactive intestinal peptide (VIP), transcript variant 1, mRNA	42.63	Up	1.22E-05
	NM_018371	Homo sapiens chondroitin beta1,4 N-acetylgalactosaminyltransferase (ChGn), mRNA	42.53	Up	2.67E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_181481	Homo sapiens chromosome 18 open reading frame 1 (C18orf1), transcript variant a1, mRNA	41.33	Up	6.27E-06
10	BQ025821	UI-1-BB1p-aye-f-10-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-aye-f-10-0-UI 3, mRNA sequence	39.96	Up	3.10E-06
	NM_152754	Homo sapiens sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3D (SEMA3D), mRNA	37.45	Up	3.17E-05
15	NM_002928	Homo sapiens regulator of G-protein signalling 16 (RGS16), mRNA	36.89	Up	2.02E-05
	H85497	yv88b07.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:249781 5, mRNA sequence	35.37	Up	8.96E-05
20	BF515657	UI-H-BW1-anu-e-05-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3083601 3, mRNA sequence	34.85	Up	5.22E-06
	BE465760	hw22f09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3183689 3, mRNA sequence	34.39	Up	1.39E-05
25	AK027128	Homo sapiens cDNA: FLJ23475 fis, clone HSI13659	33.21	Up	4.37E-06
	BC040678	Homo sapiens, clone IMAGE:4817707, mRNA	32.1	Up	3.37E-05
	NM_001864	Homo sapiens cytochrome c oxidase subunit VIIa polypeptide 1 (muscle) (COX7A1), mRNA	30.71	Up	1.69E-05
30	AK096708	Homo sapiens cDNA FLJ39389 fis, clone PLACE6003621	28.39	Up	8.85E-06
	NM_005308	Homo sapiens G protein-coupled receptor kinase 5 (GRK5), mRNA	27.96	Up	1.34E-05
	CF137545	UI-HF-BN0-ane-d-05-0-UI.r1 NIH_MGC_50 Homo sapiens cDNA clone IMAGE:3092384 5, mRNA sequence	27.55	Up	2.39E-05
35	NM_053281	Homo sapiens dachshund homolog 2 (Drosophila) (DACH2), mRNA	25.26	Up	5.22E-05
	NM_005595	Homo sapiens nuclear factor I/A (NFIA), mRNA	23.76	Up	2.51E-05
40	BM685124	UI-E-EJ1-ajl-I-13-0-UI.s1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajl-I-13-0-UI 3, mRNA sequence	23.64	Up	1.16E-05
	NM_004411	Homo sapiens dynein, cytoplasmic, intermediate polypeptide 1 (DNCL1), mRNA	23.55	Up	4.37E-06
45	NM_032918	Homo sapiens RAS-like, estrogen-regulated, growth inhibitor (RERG), mRNA	23	Up	3.96E-06
	NM_020997	Homo sapiens left-right determination factor 1 (LEFTY1), mRNA	22.48	Up	9.37E-06
50	N33310	yy39g10.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:273666 3, mRNA sequence	22.37	Up	1.57E-05
	BM713465	UI-E-EJ0-aho-m-22-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aho-m-22-0-UI 5, mRNA sequence	21.7	Up	8.98E-06
55	AK095053	Homo sapiens cDNA FLJ37734 fis, clone BRHIP2020842	21.49	Up	2.37E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	BC016722	Homo sapiens cDNA clone IMAGE:4075924, partial cds	20.91	Up	7.23E-04
	AW445209	UI-H-B13-akc-g-11-0-UI.s1 NCI_CGAP_Sub5Homo sapiens cDNA clone IMAGE:2733908 3, mRNA sequence	20.04	Up	6.27E-06
10	AI032876	ow13g03.x1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:1646740 3, mRNA sequence	19.56	Up	2.18E-05
15	BG389328	602413981F1 NIH_MGC_92 Homo sapiens cDNA clone IMAGE:4522269 5, mRNA sequence	18.94	Up	9.37E-06
	M_153355	Homo sapiens T-cell lymphoma breakpoint associated target 1 (TCBA1), mRNA	18.74	Up	5.63E-06
20	BX089554	BX089554 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998P07210 ; IMAGE:142326, mRNA sequence	18.36	Up	1.12E-06
	NM_002518	Homo sapiens neuronal PAS domain protein 2 (NPAS2), mRNA	17.94	Up	2.19E-06
25	AI342246	qt26g09.x1 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGE:1949152 3, mRNA sequence	17.68	Up	8.37E-06
	NM_170744	Homo sapiens unc-5 homolog B (C. elegans) (UNC5B), mRNA	17.32	Up	1.99E-05
30	AI951740	wv38h09.x1 NCI_CGAP_Ov18 Homo sapiens cDNA clone IMAGE:2531873 3, mRNA sequence	17.24	Up	8.84E-05
	NM_005110	Homo sapiens glutamine-fructose-6-phosphate transaminase 2 (GFPT2), mRNA	16.65	Up	8.86E-06
	NM_015564	Homo sapiens leucine rich repeat transmembrane neuronal 2 (LRRTM2), mRNA	16.45	Up	2.72E-05
35	BC043411	Homo sapiens, clone IMAGE:6155889, mRNA	16.38	Up	3.10E-06
	M_153183	Homo sapiens nudix (nucleoside diphosphate linked moiety X)-type motif 10 (NUDT10), mRNA	16.25	Up	1.45E-05
	NM_005320	Homo sapiens histone 1, H1d (HIST1H1D), mRNA	16.24	Up	4.43E-04
40	W93585	zd95g01.s1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:357264 3, mRNA sequence	16.22	Up	1.39E-05
	NM_002771	Homo sapiens protease, serine, 3 (mesotrypsin) (PRSS3), mRNA	16.15	Up	3.20E-05
45	NM_013387	Homo sapiens ubiquinol-cytochrome c reductase complex (7.2 kD) (HSPC051), transcript variant 1, mRNA	16.09	Up	3.10E-06
	NM_001853	Homo sapiens collagen, type IX, alpha 3 (COL9A3), mRNA	15.86	Up	4.01E-05
50	NM_014802	Homo sapiens KIAA0528 gene product (KIAA0528), mRNA	15.81	Up	5.11E-05
	AF216077	Homo sapiens clone HB-2 mRNA sequence	15.64	Up	2.71E-05
	AI686652	tu35d06.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2253035 3, mRNA sequence	15.12	Up	5.07E-06
55	AI086279	oz40h01.s1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:1677841 3, mRNA sequence	14.96	Up	2.47E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_001146	Homo sapiens angiotensinogen 1 (ANGPT1), transcript variant 1, mRNA	14.38	Up	1.29E-05
	NM_021643	Homo sapiens tribbles homolog 2 (Drosophila) (TRIB2), mRNA	14.32	Up	6.48E-06
10	NM_020927	Homo sapiens KIAA1576 protein (KIAA1576), mRNA	14.14	Up	1.61E-05
	W69644	z45f10.r1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:343627 5, mRNA sequence	14.01	Up	1.22E-05
15	NM_012204	Homo sapiens general transcription factor IIIC, polypeptide 4, 90kDa (GTF3C4), mRNA	13.95	Up	1.41E-05
	BM663928	UI-E-CI1-afw-p-01-0-UI.s1 UI-E-CI1 Homo sapiens cDNA clone UI-E-CI1-afw-p-01-0-UI 3, mRNA sequence	13.53	Up	8.04E-05
20	BM724062	UI-E-E01-aiy-a-22-0-UI.r1 UI-E-E01 Homo sapiens cDNA clone UI-E-E01-aiy-a-22-0-UI 5, mRNA sequence	13.51	Up	3.19E-06
	H51050	yp84e11.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:194156 3, mRNA sequence	13.39	Up	1.94E-05
25	BX111353	BX111353 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp9980094576 ; IMAGE:1869152, mRNA sequence	13.38	Up	3.92E-05
30	NM_054027	Homo sapiens ankylosis, progressive homolog (mouse) (ANKH), mRNA	13.27	Up	3.33E-05
	NM_152550	Homo sapiens SH3 domain containing ring finger 2 (SH3RF2), mRNA	13.06	Up	2.23E-05
	NM_033082	Homo sapiens cytokine induced protein 29 kDa (CIP29), mRNA	12.87	Up	5.63E-06
35		zr47e10.r1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:666570 5, mRNA sequence	12.47	Up	6.41E-05
	AA232643	BX103846 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGp998F125811 ; IMAGE:2342027, mRNA sequence	12.4	Up	1.39E-05
40	NM_016140	Homo sapiens brain specific protein (CGI-38), mRNA	12.26	Up	2.71E-05
	AW043793	wy76d11.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:2554485 3 similar to contains element MER18 repetitive element ;, mRNA sequence	12.04	Up	6.21E-06
45	NM_020226	Homo sapiens PR domain containing 8 (PRDM8), mRNA	12.01	Up	5.32E-06
	BX113590	BX113590 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGp998A14871 ; IMAGE:376597, mRNA sequence	11.97	Up	3.09E-05
50	AK127644	Homo sapiens cDNA FLJ45742 fis, clone KIDNE2016327	11.91	Up	7.34E-05
	BC046364	Homo sapiens flavoprotein oxidoreductase MICAL3, mRNA (cDNA clone IMAGE:5737121), with apparent retained intron	11.88	Up	1.87E-05
55	AI821210	ne08e05.y5 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGE:880640 5, mRNA sequence	11.79	Up	4.15E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AK128288	Homo sapiens cDNA FLJ46426 fis, clone THYMU3013897	11.76	Up	2.88E-04
	NM_033641	Homo sapiens collagen, type IV, alpha 6 (COL4A6), transcript variant B, mRNA	11.56	Up	2.05E-05
10	NM_004460	Homo sapiens fibroblast activation protein, alpha (FAP), mRNA	11.54	Up	1.40E-05
	A1792426	qe47a12.y5 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:1742110 5 similar to contains MER22.b3 MER18 repetitive element;, mRNA sequence	11.41	Up	1.33E-03
15	NM_018374	Homo sapiens hypothetical protein FLJ11273 (FLJ11273), mRNA	11.23	Up	3.24E-04
	BM802920	AGENCOURT_6457446 NIH_MGC_88 Homo sapiens cDNA clone IMAGE:5560288 5, mRNA sequence	11.11	Up	8.59E-05
20	AA043255	zk49f07.s1 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGE:486181 3, mRNA sequence	11.1	Up	2.64E-05
25	NM_001847	Homo sapiens collagen, type IV, alpha 6 (COL4A6), transcript variant A, mRNA	10.98	Up	9.89E-06
	AK124778	Homo sapiens cDNA FLJ42788 fis, clone BRAWH3007129	10.95	Up	9.27E-05
30	AK130306	Homo sapiens cDNA FLJ26796 fis, clone PRS05079	10.86	Up	1.02E-04
	AK023739	Homo sapiens cDNA FLJ13677 fis, clone PLACE1011982	10.83	Up	6.24E-05
	AL832624	Homo sapiens mRNA; cDNA DKFZp451B0818 (from clone DKFZp451 B0818)	10.8	Up	5.63E-06
35	BC014344	Homo sapiens, Similar to arylacetamide deacetylase, clone IMAGE:3934567, mRNA	10.68	Up	2.37E-05
	NM_000955	Homo sapiens prostaglandin E receptor 1 (subtype EP1), 42kDa (PTGER1), mRNA	10.66	Up	1.14E-05
40	CD001963	C24B1 subtracted cochlea cDNA library Homo sapiens cDNA, mRNA sequence	10.47	Up	7.71E-06
	NM_032261	Homo sapiens chromosome 21 open reading frame 56 (C21orf56), mRNA	10.42	Up	4.73E-06
	BC036004	Homo sapiens, clone IMAGE:4730399, mRNA	10.3	Up	3.06E-05
45	NM_006408	Homo sapiens anterior gradient 2 homolog (Xenopus laevis) (AGR2), mRNA	10.28	Up	4.23E-05
	AB037805	Homo sapiens mRNA for KIAA1384 protein, partial cds	10.24	Up	8.03E-05
	BF964783	602268030F1 NIH_MGC_81 Homo sapiens cDNA clone IMAGE:4356246 5, mRNA sequence	10.02	Up	1.41E-05
50	NM_000963	Homo sapiens prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2), mRNA	9.91	Up	5.63E-06
		Homo sapiens cDNA FLJ35222 fis, clone PROST2000835	9.88	Up	4.65E-02
55	AK092541	PROST2000835			
	NM_007361	Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	9.68	Up	5.30E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_013231	Homo sapiens fibronectin leucine rich transmembrane protein 2 (FLRT2), mRNA	9.58	Up	1.43E-05
	NM_002667	Homo sapiens phospholamban (PLN), mRNA	9.55	Up	9.83E-04
10	AF414442	Homo sapiens ovarian cancer related tumor marker CA125 mRNA, complete cds	9.51	Up	1.11E-03
	AK125453	Homo sapiens cDNA FLJ43464 fis, clone OCBBF2036225	9.48	Up	1.15E-02
15	W38393	zb15c07.r1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:302124 5, mRNA sequence	9.39	Up	5.52E-05
	NM_152775	Homo sapiens hypothetical protein MGC33607 (MGC33607), mRNA	9.29	Up	2.35E-05
20	AK091731	Homo sapiens cDNA FLJ34412 fis, clone HEART2002432	9.07	Up	2.06E-05
	AK025909	Homo sapiens cDNA: FLJ22256 fis, clone HRC02860	9.05	Up	4.45E-05
	D62676	HUM313C12B Clontech human aorta polyA+ mRNA (#6572) Homo sapiens cDNA clone GEN-313C12 5, mRNA sequence	9.02	Up	1.38E-03
25					
	AK129955	Homo sapiens cDNA FLJ26445 fis, clone KDN02608	8.98	Up	8.52E-05
	R22189	yh26a02.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:130826 3, mRNA sequence	8.94	Up	2.29E-05
30	AW963062	EST375135 MAGE resequences, MAGH Homo sapiens cDNA, mRNA sequence	8.9	Up	1.76E-04
	AW044502	wx22g05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2544440 3, mRNA sequence	8.89	Up	1.35E-05
35	NM_014222	Homo sapiens NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 8, 19kDa (NDUFA8), nuclear gene encoding mitochondrial protein, mRNA	8.88	Up	4.45E-05
	AK124562	Homo sapiens cDNA FLJ42571 fis, clone BRACE3008036	8.82	Up	8.86E-06
40	NM_002261	Homo sapiens killer cell lectin-like receptor subfamily C, member 3 (KLRC3), transcript variant NKG2-E, mRNA	8.75	Up	9.45E-05
	BC012900	Homo sapiens, clone IMAGE:3881549, mRNA	8.71	Up	1.61E-04
	R53688	yg84h04.r1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:40175 5, mRNA sequence	8.49	Up	1.74E-05
45	AL133118	Homo sapiens mRNA; cDNA DKFZp586N0121 (from clone DKFZp586N0121)	8.47	Up	6.83E-05
	NM_002260	Homo sapiens killer cell lectin-like receptor subfamily C, member 2 (KLRC2), mRNA	8.44	Up	1.61E-04
50	NM_002421	Homo sapiens matrix metalloproteinase 1 (interstitial collagenase) (MMP1), mRNA	8.34	Up	5.17E-04
	NM_005328	Homo sapiens hyaluronan synthase 2 (HAS2), mRNA	8.18	Up	2.35E-04
	NM_152423	Homo sapiens melanoma associated antigen (mutated) 1-like 1 (MUM1L1), mRNA	8.12	Up	6.31E-05
55	NM_004848	Homo sapiens chromosome 1 open reading frame 38 (C1 orf38), mRNA	8.05	Up	5.83E-05
	NM_006617	Homo sapiens nestin (NES), mRNA	8.01	Up	8.50E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	C02345	HUMGS0007544 Human adult (K.Okubo) Homo sapiens cDNA, mRNA sequence	8	Up	3.09E-05
	NM_000900	Homo sapiens matrix Gla protein (MGP), mRNA	7.97	Up	3.88E-05
10	N54656	yz08h05.r1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:282489 5, mRNA sequence	7.97	Up	4.17E-05
	AK095726	Homo sapiens cDNA FLJ38407 fis, clone FEBRA2008859	7.97	Up	1.84E-05
15	BC027461	Homo sapiens cDNA clone IMAGE:2984900, containing frame-shift errors	7.95	Up	2.18E-04
	AK096061	Homo sapiens cDNA FLJ38742 fis, clone KIDNE2012009	7.91	Up	1.32E-05
20	NM_001001931	Homo sapiens mitochondrial tumor suppressor 1 (MTUS1), nuclear gene encoding mitochondrial protein, transcript variant 4, mRNA	7.84	Up	1.22E-05
	NM_133504	Homo sapiens decorin (DCN), transcript variant B, mRNA	7.82	Up	6.70E-06
25	NM_018427	Homo sapiens RRN3 RNA polymerase I transcription factor homolog (yeast) (RRN3), mRNA	7.76	Up	1.39E-05
	NM_006528	Homo sapiens tissue factor pathway inhibitor 2 (TFPI2), mRNA	7.73	Up	3.10E-06
	CN371168	17000600077294 GRN_PREHEP Homo sapiens cDNA 5, mRNA sequence	7.63	Up	1.83E-05
30	AI911957	wd78b01.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:2337673 3, mRNA sequence	7.61	Up	1.82E-05
	NM_058187	Homo sapiens chromosome 21 open reading frame 63 (C21orf63), mRNA	7.59	Up	9.08E-05
35	NM_004235	Homo sapiens Kruppel-like factor 4 (gut) (KLF4), mRNA	7.52	Up	1.43E-05
	AL832916	Homo sapiens mRNA; cDNA DKFZp76210915 (from clone DKFZp76210915)	7.51	Up	6.70E-06
40	BG118019	602351269F1 NIH_MGC_90 Homo sapiens cDNA clone IMAGE:4446065 5, mRNA sequence	7.49	Up	1.43E-05
	AW770283	h176b02.x1 NCI_CGAP_Kid13 Homo sapiens cDNA clone IMAGE:3007083 3, mRNA sequence	7.41	Up	1.38E-04
	T47612	yb15h03.s1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:71285 3, mRNA sequence	7.35	Up	9.37E-06
45	BM998303	UI-H-DT1-awc-h-03-0-UI.s1 NCI_CGAP_DT1 Homo sapiens cDNA clone IMAGE:5887538 3, mRNA sequence	7.31	Up	2.23E-05
50	AK092048	Homo sapiens cDNA FLJ34729 fis, clone MESAN2006401	7.26	Up	1.57E-05
	BX647541	Homo sapiens mRNA; cDNA DKFZp686P0492 (from clone DKFZp686P0492)	7.24	Up	2.71E-05
55	AI033863	cow10e02.x1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:1646426 3, mRNA sequence	7.16	Up	7.24E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_004657	Homo sapiens serum deprivation response (phosphatidylserine binding protein) (SDPR), mRNA	7.16	Up	3.17E-05
	BF222640	7p57c03.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:3649901 3, mRNA sequence	7.13	Up	1.37E-04
10	BU620793	UI-H-FL1-bfx-d-10-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bfx-d-10-0-UI 3, mRNA sequence	7.11	Up	1.46E-05
	AA102553	zn26a04.s1 Stratagene neuroepithelium NT2RAMI 937234 Homo sapiens cDNA clone IMAGE:548526 3, mRNA sequence	7.01	Up	4.89E-05
15	NM_021242	Homo sapiens MID1 interacting protein 1 (gastrulation specific G12-like (zebrafish)) (MID1IP1), mRNA	6.96	Up	1.32E-04
20	CB045035	NISC_gc08b02.x1 NCI_CGAP_Co17 Homo sapiens cDNA clone IMAGE:3217875 3, mRNA sequence	6.94	Up	1.32E-04
	NM_007063	Homo sapiens TBC1 domain family, member 8 (with GRAM domain) (TBC1D8), mRNA	6.92	Up	7.24E-06
25	H25898	yl55b10.r1 Soares breast 3NbHBst Homo sapiens cDNA clone IMAGE:162139 5, mRNA sequence	6.91	Up	5.94E-04
	AK091337	Homo sapiens cDNA FLJ34018 fis, clone FCBBF2002801	6.89	Up	1.05E-04
	BE877764	601486331 F1 NIH_MGC_69 Homo sapiens cDNA clone IMAGE:3888943 5, mRNA sequence	6.88	Up	2.25E-04
30	BE503916	hz35g01.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:3210000 3, mRNA sequence	6.73	Up	2.28E-05
	NM_024993	Homo sapiens leucine rich repeat transmembrane neuronal 4 (LRRTM4), mRNA	6.69	Up	2.67E-05
35		Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	6.65	Up	2.63E-04
	NM_007361				
	BC070147	Homo sapiens cDNA clone IMAGE:4672631, containing frame-shift errors	6.65	Up	1.35E-05
40	BC045666	Homo sapiens tumor protein p53 inducible protein 11, mRNA (cDNA clone IMAGE:5298525), containing frame-shift errors	6.62	Up	1.80E-04
	NM_017787	Homo sapiens chromosome 10 open reading frame 26 (C10orf26), mRNA	6.59	Up	3.69E-05
45	BQ267806	ij94e04.x1 Human insulinoma Homo sapiens cDNA clone IMAGE:5779278 3, mRNA sequence	6.59	Up	1.39E-04
	NM_032578	Homo sapiens myopalladin (FLJ14437), mRNA	6.57	Up	1.43E-05
	AK124699	Homo sapiens cDNA FLJ42709 fis, clone BRAMY3007350	6.55	Up	1.67E-05
50	NM_000921	Homo sapiens phosphodiesterase 3A, cGMP-inhibited (PDE3A), mRNA	6.51	Up	3.13E-05
	CB048134	NISC_gj03a09.y1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:3270448 5, mRNA sequence	6.47	Up	8.86E-06
	CA843592	ir49c12.x1 HR85 islet Homo sapiens cDNA clone IMAGE:6548544 3, mRNA sequence	6.44	Up	3.85E-05
55	NM_024633	Homo sapiens chromosome 14 open reading frame 139 (C14orf139), mRNA	6.42	Up	2.45E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	BX640685	Homo sapiens mRNA; cDNA DKFZp686M08112 (from clone DKFZp686M08112)	6.41	Up	3.09E-05
	NM_005398	Homo sapiens protein phosphatase 1, regulatory (inhibitor) subunit 3C (PPP1 R3C), mRNA	6.38	Up	1.22E-05
10	NM_001957	Homo sapiens endothelin receptor type A (EDNRA), mRNA	6.36	Up	3.08E-05
	NM_003248	Homo sapiens thrombospondin 4 (THBS4), mRNA	6.35	Up	2.71E-05
	BQ188758	UI-E-EJ1-ajx-a-20-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajx-a-20-0-UI 5, mRNA sequence	6.28	Up	5.68E-03
15	AI830524	wh52c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384354 3, mRNA sequence	6.28	Up	2.77E-05
	BG818762	602779092F2 NCI_CGAP_Brn67 Homo sapiens cDNA clone IMAGE:4914502 5, mRNA sequence	6.28	Up	1.57E-05
20	NM_024420	Homo sapiens phospholipase A2, group IVA (cytosolic, calcium-dependent) (PLA2G4A), mRNA	6.26	Up	3.90E-04
	NM_173552	Homo sapiens hypothetical protein MGC33365 (MGC33365), mRNA	6.25	Up	1.39E-05
25	BQ189707	UI-E-EJ1-aka-o-12-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-aka-o-12-0-UI 5, mRNA sequence	6.25	Up	2.62E-05
	NM_015559	Homo sapiens SET binding protein 1 (SETBP1), mRNA	6.21	Up	1.83E-03
30	T15720	IB1788 Infant brain, Bento Soares Homo sapiens cDNA 3end, mRNA sequence	6.17	Up	2.02E-05
	AW291013	UI-H-BI2-agj-f-01-0-UI.s1 NCI_CGAP_Sub4 Homo sapiens cDNA clone IMAGE:2724601 3, mRNA sequence	6.15	Up	3.80E-05
35	NM_024336	Homo sapiens iroquois homeobox protein 3 (IRX3), mRNA	6.12	Up	9.89E-06
	NM_025107	Homo sapiens myc target 1 (MYCT1), mRNA	6.11	Up	5.67E-05
	NM_144665	Homo sapiens sestrin 3 (SESN3), mRNA	6.11	Up	1.39E-04
40	AA908815	og77h08.s1 NCI_CGAP_Ov8 Homo sapiens cDNA clone IMAGE:1454367 3, mRNA sequence	6.07	Up	2.25E-04
	NM_024621	Homo sapiens hypothetical protein FLJ12604 (FLJ12604), mRNA	6.05	Up	2.61E-04
45	AI335546	qt24d07.x1 NCI_CGAP_GC4 Homo sapiens cDNA clone IMAGE:1948525 3, mRNA sequence	6.02	Up	8.66E-06
	AB007954	Homo sapiens mRNA, chromosome 1 specific transcript KIAA0485	5.99	Up	5.46E-05
	NM_033292	Homo sapiens caspase 1, apoptosis-related cysteine protease (interleukin 1, beta, convertase) (CASP1), transcript variant alpha, mRNA	5.95	Up	4.51E-05
50	BX112737	BX112737 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998H17189 ; IMAGE:134080, mRNA sequence	5.94	Up	3.13E-06
55	NM_005239	Homo sapiens v-ets erythroblastosis virus E26 oncogene homolog 2 (avian) (ETS2), mRNA	5.94	Up	1.14E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	BM677978	UI-E-EJ0-aig-o-17-0-UI.s1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-aig-o-17-0-UI 3, mRNA sequence	5.91	Up	7.10E-05
10	BX648604	Homo sapiens mRNA; cDNA DKFZp686M11213 (from clone DKFZp686M11213)	5.91	Up	1.15E-04
	BC041997	Homo sapiens, clone IMAGE:5310819, mRNA	5.91	Up	4.16E-04
	NM_145239	Homo sapiens similar to lymphocyte antigen 6 complex, locus G5B; G5b protein; open reading frame 31 (LOC112476), mRNA	5.85	Up	5.37E-05
15	NM_015149	Homo sapiens ral guanine nucleotide dissociation stimulator-like 1 (RGL1), mRNA	5.85	Up	2.32E-05
	NM_030756	Homo sapiens transcription factor 7-like 2 (T-cell specific, HMG-box) (TCF7L2), mRNA	5.85	Up	3.88E-05
20	BU626144	UI-H-FG1-bgq-i-08-0-UI.s1 NCI_CGAP_FG1 Homo sapiens cDNA clone UI-H-FG1-bgq-i-08-0-UI 3, mRNA sequence	5.85	Up	1.70E-04
	BM556191	AGENCOURT_6544282 NIH_MGC_88 Homo sapiens cDNA clone IMAGE:5550173 5, mRNA sequence	5.83	Up	1.85E-05
25	BM931867	UI-E-EJ1-ajk-n-19-0-UI.r1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajk-n-19-0-UI 5, mRNA sequence	5.81	Up	2.62E-05
30	BG576442	602595685F1 NIH_MGC_87 Homo sapiens cDNA clone IMAGE:4708725 5, mRNA sequence	5.8	Up	7.12E-05
	AB096240	Homo sapiens LOH11CR1A gene, loss of heterozygosity, 11, chromosomal region 1 gene A product	5.78	Up	1.63E-03
35	NM_003763	Homo sapiens syntaxin 16 (STX16), transcript variant 2, mRNA	5.72	Up	7.02E-05
	BQ013869	UI-1-BC1p-alg-a-03-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1 p-alg-a-03-0-UI 3, mRNA sequence	5.69	Up	1.22E-05
40	NM_005795	Homo sapiens calcitonin receptor-like (CALCRL), mRNA	5.68	Up	3.82E-05
	BM690053	UI-E-CK1-abr-c-12-0-UI.r1 UI-E-CK1 Homo sapiens cDNA clone UI-E-CK1-abr-c-12-0-UI 5, mRNA sequence	5.64	Up	2.37E-05
45	AL832779	Homo sapiens mRNA; cDNA DKFZp686H157 (from clone DKFZp686H157)	5.64	Up	7.39E-06
	CA425350	UI-H-DF0-bel-I-04-0-UI.s1 NCI_CGAP_DF0 Homo sapiens cDNA clone UI-H-DF0-bel-I-04-0-UI 3, mRNA sequence	5.63	Up	2.47E-05
50	AA398628	zt74e03.s1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:728092 3, mRNA sequence	5.62	Up	1.87E-04
	BX093329	BX093329 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGp998A124183 ; IMAGE:1648403, mRNA sequence	5.61	Up	7.11E-05
55	NM_006573	Homo sapiens tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B), mRNA	5.58	Up	1.02E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AK124699	Homo sapiens cDNA FLJ42709 fis, clone BRAMY3007350	5.53	Up	1.72E-04
10	AL709606	DKFZp686M125_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686M125 5, mRNA sequence	5.53	Up	1.80E-04
	BQ022721	UI-1-BB1p-axt-g-01-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-axt-g-01-0-UI 3, mRNA sequence	5.5	Up	2.79E-03
15	AA613572	nq22g06.s1 NCI_CGAP_Co10 Homo sapiens cDNA clone IMAGE:1144666 3, mRNA sequence	5.47	Up	4.31E-04
	R38944	yd06g09.s1 Soares infant brain 1 NIB Homo sapiens cDNA clone IMAGE:25061 3 similar to contains Alu repetitive element, mRNA sequence	5.45	Up	4.34E-05
20	AK124563	Homo sapiens cDNA FLJ42572 fis, clone BRACE3008092	5.43	Up	9.13E-06
	BC037929	Homo sapiens cDNA clone IMAGE:5284659, partial cds	5.42	Up	7.14E-04
25	AK096536	Homo sapiens cDNA FLJ39217 fis, clone OCBBF2006639, moderately similar to ZINC FINGER PROTEIN 84	5.41	Up	3.17E-06
	AL049443	Homo sapiens mRNA; cDNA DKFZp586N2020 (from clone DKFZp586N2020)	5.41	Up	2.83E-04
30	NM_004951	Homo sapiens Epstein-Barr virus induced gene 2 (lymphocyte-specific G protein-coupled receptor) (EBI2), mRNA	5.39	Up	4.33E-04
	NM_021153	Homo sapiens cadherin 19, type 2 (CDH19), mRNA	5.37	Up	3.26E-05
35	BX106577	BX106577 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGp998H131854 ; IMAGE:754236, mRNA sequence	5.37	Up	2.51E-05
	AW148979	xf07g08.x1 NCI_CGAP_Kid8 Homo sapiens cDNA clone IMAGE:2617406 3, mRNA sequence	5.36	Up	2.78E-04
40	AI024717	ov68h06.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1642523 3, mRNA sequence	5.35	Up	3.75E-05
	AF131784	Homo sapiens clone 25194 mRNA sequence	5.34	Up	2.14E-05
	AK092375	Homo sapiens cDNA FLJ35056 fis, clone OCBBF2018581	5.32	Up	6.05E-05
45	NM_002615	Homo sapiens serine (or cysteine) proteinase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 (SERPINF1), mRNA	5.3	Up	1.74E-04
	NM_016206	Homo sapiens colon carcinoma related protein (FLJ38507), mRNA	5.3	Up	2.44E-04
50	BQ346857	RC1-NT0033-250800-018-g09 NT0033 Homo sapiens cDNA, mRNA sequence	5.29	Up	4.05E-05
	BQ026175	UI-1-BB1p-akc-h-10-0-UI.s1 NCI_CGAP_PI6 Homo sapiens cDNA clone UI-1-BB1p-akc-h-10-0-UI 3, mRNA sequence	5.29	Up	7.29E-04
55	NM_033067	Homo sapiens DMRT-like family B with proline-rich C-terminal, 1 (DMRTB1), mRNA	5.28	Up	2.03E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AK127847	Homo sapiens cDNA FLJ45950 fis, clone PLACE7008136	5.27	Up	6.27E-06
	AI024323	ov67g08.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1642430 3, mRNA sequence	5.27	Up	1.14E-04
10	NM_022768	Homo sapiens RNA binding motif protein 15 (RBM15), mRNA	5.27	Up	3.30E-04
	BC016780	Homo sapiens, clone IMAGE:4106389, mRNA	5.24	Up	2.73E-03
	CA427170	UI-H-DF0-bey-f-20-0-UI.s1 NCI_CGAP_DF0 Homo sapiens cDNA clone UI-H-DF0-bey-f-20-0-UI 3, mRNA sequence	5.23	Up	2.60E-04
15	BX537862	Homo sapiens mRNA; cDNA DKFZp779M0552 (from clone DKFZp779M0552)	5.23	Up	5.42E-05
	NM_022658	Homo sapiens homeo box C8 (HOXC8), mRNA	5.22	Up	4.37E-04
20	601237381 F1 NIH_MGC_44	Homo sapiens cDNA clone IMAGE:3609140 5, mRNA sequence	5.21	Up	1.74E-04
	BE378852	Homo sapiens mRNA for FLJ00204 protein	5.18	Up	5.59E-05
	AK074131	BX098131 Soares_testis_NHT Homo sapiens cDNA clone IMAGp9980101830 ; IMAGE:745185, mRNA sequence	5.15	Up	1.32E-05
25	BX098131				
	AI798732	we91h12.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2348519 3, mRNA sequence	5.14	Up	5.68E-04
30	NM_017577	Homo sapiens hypothetical protein DKFZp434C0328 (DKFZp434C0328), mRNA	5.13	Up	3.63E-04
	NM_015541	Homo sapiens leucine-rich repeats and immunoglobulin-like domains 1 (LRIG1), mRNA	5.12	Up	1.04E-04
	NM_032622	Homo sapiens ligand of numb-protein X (LNK), mRNA	5.12	Up	1.74E-04
35	NM_003500	Homo sapiens acyl-Coenzyme A oxidase 2, branched chain (ACOX2), mRNA	5.11	Up	1.34E-05
	NM_152270	Homo sapiens hypothetical protein FLJ34922 (FLJ34922), mRNA	5.07	Up	4.93E-05
40	BM695626	UI-E-CQ1-aew-i-06-0-UI.r1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aew-i-06-0-UI 5, mRNA sequence	5.06	Up	1.10E-03
	NM_014548	Homo sapiens tropomodulin 2 (neuronal) (TMOD2), mRNA	5.05	Up	4.23E-03
45	BC036223	Homo sapiens, clone IMAGE:5272183, mRNA	5.04	Up	6.69E-05
	NM_173554	Homo sapiens chromosome 10 open reading frame 107 (C10orf107), mRNA	5.04	Up	8.63E-05
	NM_018409	Homo sapiens hypothetical protein DKFZp761O0113 (DKFZp761O0113), mRNA	5.03	Up	3.39E-04
50	BF509492	UI-H-B14-a0z-c-06-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086579 3, mRNA sequence	5.01	Up	1.99E-03
	BQ574344	UI-H-EZ1-baz-b-10-0-UI.s1 NCI_CGAP_Ch2 Homo sapiens cDNA clone UI-H-EZ1-baz-b-10-0-UI 3, mRNA sequence	5.01	Up	7.24E-06
55					

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_022454	Homo sapiens SRY (sex determining region Y)-box 17 (SOX17), mRNA	366.7	Down	3.10E-06
	BG219729	RST39494 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	268.82	Down	2.19E-06
10	NM_152737	Homo sapiens hypothetical protein MGC33993 (MGC33993), mRNA	250.88	Down	1.66E-06
	BX102632	BX102632 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGp998J052307 ; IMAGE:928228, mRNA sequence	188.6	Down	1.29E-06
15	NM_002521	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	184.64	Down	3.10E-06
	NM_013230	Homo sapiens CD24 antigen (small cell lung carcinoma cluster 4 antigen) (CD24), mRNA	160.77	Down	4.30E-06
20	AK023631	Homo sapiens cDNA FLJ13569 fis, clone PLACE1008369	154.91	Down	5.78E-07
	NM_006169	Homo sapiens nicotinamide N-methyltransferase (NNMT), mRNA	141.97	Down	1.12E-06
25	NM_006228	Homo sapiens prepronociceptin (PNOC), mRNA	138.62	Down	1.22E-05
	AI765021	wh56c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384738 3, mRNA sequence	130.91	Down	6.27E-06
	NM_004617	Homo sapiens transmembrane 4 superfamily member 4 (TM4SF4), mRNA	111.06	Down	1.43E-05
30	AK096481	Homo sapiens cDNA FLJ39162 fis, clone OCBBF2002376	102.43	Down	3.17E-06
	NM_007038	Homo sapiens a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2) (ADAMTS5), mRNA	95.94	Down	6.14E-06
35	AB032980	Homo sapiens mRNA for KIAA1154 protein, partial cds	85.12	Down	9.37E-06
	AK092245	Homo sapiens cDNA FLJ34926 fis, clone NT2RP7003319, highly similar to Mus musculus neuralin mRNA	70.22	Down	2.04E-05
40	NM_002276	Homo sapiens keratin 19 (KRT19), mRNA	59.59	Down	2.19E-06
	AI249696	qj64a03.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1864204 3, mRNA sequence	56.92	Down	9.37E-06
	D29453	HUMNK566 Human epidermal keratinocyte Homo sapiens cDNA clone 566, mRNA sequence	53.31	Down	5.14E-06
45	NM_004753	Homo sapiens dehydrogenase/reductase (SDR family) member 3 (DHRS3), mRNA	50.19	Down	1.41E-05
	NM_012464	Homo sapiens tolloid-like 1 (TLL1), mRNA	48.62	Down	1.72E-04
	NM_080743	Homo sapiens serine-arginine repressor protein (35 kDa) (SRp35), mRNA	48.46	Down	1.34E-06
50	NM_004496	Homo sapiens forkhead box A1 (FOXA1), mRNA	47.75	Down	3.10E-06
	BF512544	UI-H-BW1-amf-c-08-0-UI.s1 NCI_CGAP_Sub7 Homo sapiens cDNA clone IMAGE:3069687 3, mRNA sequence	46.87	Down	6.56E-06
55	NM_016356	Homo sapiens doublecortin domain containing 2 (DCDC2), mRNA	46.71	Down	1.48E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5					
	NM_020873	Homo sapiens leucine rich repeat neuronal 1 (LRRN1), mRNA	45.32	Down	3.17E-06
		Homo sapiens neuropilin (NRP) and tolloid (TLL)-like 1 (NETO1), transcript variant 3, mRNA	44.78	Down	1.84E-05
10	NM_138966				
	NM_000582	Homo sapiens secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA	43.26	Down	9.71E-06
	BU680661	UI-CF-DU1-aaz-f-04-0-UI.s1 UI-CF-DU1 Homo sapiens cDNA clone UI-CF-DU1-aaz-f-04-0-UI 3, mRNA sequence	43.24	Down	3.19E-06
15					
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	41.43	Down	3.19E-06
	NM_004221	Homo sapiens natural killer cell transcript 4 (NK4), mRNA	40.53	Down	1.66E-05
20					
	CA416106	UI-H-FE0-bbs-f-17-0-UI.s1 NCI_CGAP_FE0 Homo sapiens cDNA clone UI-H-FE0-bbs-f-17-0-UI 3, mRNA sequence	40.33	Down	2.77E-05
	NM_006439	Homo sapiens mab-21-like 2 (C. elegans) (MAB21L2), mRNA	40.02	Down	3.10E-05
25					
	BF431030	7o18c06.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3574283 3, mRNA sequence	39.89	Down	1.22E-05
	NM_025074	Homo sapiens Fraser syndrome 1 (FRAS1), transcript variant 1, mRNA	39.81	Down	3.82E-06
30					
	BG197054	RST16291 Athersys RAGE Library Homo sapiens cDNA, mRNA sequence	39.23	Down	3.73E-05
	BG165745	602344592F1 NIH_MGC_89 Homo sapiens cDNA clone IMAGE:4454470 5, mRNA sequence	37.22	Down	1.39E-04
35					
	NM_002527	Homo sapiens neurotrophin 3 (NTF3), mRNA	34.83	Down	5.63E-06
	NM_004982	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 8 (KCNJ8), mRNA	33.22	Down	3.22E-05
	NM_018168	Homo sapiens chromosome 14 open reading frame 105 (C14orf105), mRNA	31.87	Down	1.84E-05
40					
	BQ003401	UI-H-E11-azd-j-23-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5847286 3, mRNA sequence	31.63	Down	8.86E-06
	BC037316	Homo sapiens, clone IMAGE:5259432, mRNA	30.99	Down	2.19E-06
45					
	NM_016212	Homo sapiens TP53TG3 protein (TP53TG3), mRNA	30.74	Down	1.57E-05
	BQ375719	QV2-TN0173-021100-454-g06 TN0173 Homo sapiens cDNA, mRNA sequence	30.59	Down	1.84E-05
	NM_014333	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	30.19	Down	2.42E-05
50					
	NM_005712	Homo sapiens HERV-H LTR-associating 1 (HHLA1), mRNA	29.97	Down	1.62E-05
	NM_001955	Homo sapiens endothelin 1 (EDN1), mRNA	29.8	Down	1.41E-05
	NM_0025 59	Homo sapiens purinergic receptor P2X, ligand-gated ion channel, 3 (P2RX3), mRNA	28.92	Down	1.84E-05
55					
	NM_0018 85	Homo sapiens crystallin, alpha B (CRYAB), mRNA	28.49	Down	6.27E-06
	NM_0056 02	Homo sapiens claudin 11 (oligodendrocyte transmembrane protein) (CLDN11), mRNA	28.14	Down	5.78E-07

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0009 27	Homo sapiens ATP-binding cassette, sub-family B (MDR/TAP), member 1 (ABCB1), mRNA	27.84	Down	8.93E-06
	NM_0033 85	Homo sapiens visinin-like 1 (VSNL1), mRNA	27.75	Down	9.37E-06
10	NM_0203 49	Homo sapiens ankyrin repeat domain 2 (stretch responsive muscle) (ANKRD2), mRNA	27.68	Down	2.19E-06
	NM_1524 87	Homo sapiens hypothetical protein FLJ31842 (FLJ31842), mRNA	27.59	Down	3.71E-05
15	NM_0178 19	Homo sapiens RNA (guanine-9-) methyltransferase domain containing 1 (RG9MTD1), mRNA	26.6	Down	1.99E-05
	AA44913 7	zx03d12.r1 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGE:785399 5, mRNA sequence	26.24	Down	1.67E-05
20	AW5112 22	hd44d11.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2912373 3 similar to contains Alu repetitive element;, mRNA sequence	26.15	Down	9.71E-06
	AI469032	ti70a01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2137320 3, mRNA sequence	25.39	Down	3.10E-06
25	NM_0308 99	Homo sapiens zinc finger protein 323 (ZNF323), mRNA	25.07	Down	1.43E-05
	BG54530 5	602572521 F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4700644 5, mRNA sequence	24.4	Down	3.68E-06
	NM_1840 87	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	24.35	Down	9.37E-06
30	NM_0060 74	Homo sapiens tripartite motif-containing 22 (TRIM22), mRNA	24.19	Down	1.55E-04
	NM_1526 94	Homo sapiens zinc finger, CCHC domain containing 5 (ZCCHC5), mRNA	24.15	Down	8.30E-05
35	NM_1750 56	Homo sapiens hypothetical protein LOC131368 (LOC131368), mRNA	23.96	Down	2.36E-05
	NM_0152 36	Homo sapiens latrophilin 3 (LPHN3), mRNA	23.83	Down	9.13E-06
	AK12487 3	Homo sapiens cDNA FLJ42883 fis, clone BRHIP3006683	23.58	Down	1.95E-05
40	NM_0015 62	Homo sapiens interleukin 18 (interferon-gamma-inducing factor) (IL18), mRNA	23.38	Down	4.81E-05
	NM_0320 26	Homo sapiens TatD DNase domain containing 1 (TATDN1), mRNA	23.12	Down	6.00E-05
45	NM_1527 03	Homo sapiens chromosome 7 open reading frame 6 (C7orf6), mRNA	22.98	Down	1.82E-05
	NM_0326 03	Homo sapiens lysyl oxidase-like 3 (LOXL3), mRNA	22.16	Down	1.17E-05
	BX11582 5	BX115825 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGp998C134603 ; IMAGE:1879236, mRNA sequence	22.02	Down	3.22E-05
50	NM_0014 48	Homo sapiens glypican 4 (GPC4), mRNA	21.86	Down	8.50E-06
	BC00858 0	Homo sapiens, clone IMAGE:4179986, mRNA, partial cds	21.22	Down	3.19E-06
	NM_0022 73	Homo sapiens keratin 8 (KRT8), mRNA	21.19	Down	1.29E-05
55	NM_0050 69	Homo sapiens single-minded homolog 2 (Drosophila) (SIM2), transcript variant SIM2, mRNA	20.97	Down	3.19E-06
	NM_0529 47	Homo sapiens heart alpha-kinase (HAK), mRNA	20.9	Down	1.22E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AB04126 9	Homo sapiens mRNA for keratin 19, partial cds, isolate: K19-141	20.57	Down	3.49E-05
	NM_002578	Homo sapiens p21 (CDKN1A)-activated kinase 3 (PAK3), mRNA	20.36	Down	3.19E-06
10	NM_0247 26	Homo sapiens IQ motif containing with AAA domain (IQCA), mRNA	20.18	Down	1.02E-04
	NM_0186 58	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 16 (KCNJ16), transcript variant 1, mRNA	19.93	Down	3.44E-05
15	BU56780 4	AGENCOURT_10398872 NIH_MGC_82 Homo sapiens cDNA clone IMAGE:6614502 5, mRNA sequence	19.73	Down	6.27E-06
	NM_0053 29	Homo sapiens hyaluronan synthase 3 (HAS3), transcript variant 1, mRNA	19.66	Down	3.19E-06
20	BX50911 7	DKFZp686C09280_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686C09280 5, mRNA sequence	19.48	Down	8.85E-06
	NM_0033 92	Homo sapiens wingless-type MMTV integration site family, member 5A (WNT5A), mRNA	19.4	Down	3.19E-06
25	NM_0248 98	Homo sapiens family with sequence similarity 31, member C (FAM31C), mRNA	19.07	Down	6.27E-06
	AF31838 2	Homo sapiens pp9974 mRNA, complete cds	19	Down	6.27E-06
30	AL11757 8	Homo sapiens mRNA; cDNA DKFZp434C128 (from clone DKFZp434C128)	18.69	Down	3.17E-06
	BU63316 3	UI-H-FL1-bgt-n-07-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bgt-n-07-0-UI 3, mRNA sequence	18.57	Down	9.42E-06
35	AB01153 9	Homo sapiens mRNA for MEGF6 protein (KIAA0815), partial cds	18.38	Down	4.22E-05
	NM_0006 93	Homo sapiens aldehyde dehydrogenase 1 family, member A3 (ALDH1A3), mRNA	18.37	Down	4.97E-05
40	AA19532 8	zr34f08.s1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:665319 3, mRNA sequence	18.33	Down	9.56E-06
	AL08010 3	Homo sapiens mRNA; cDNA DKFZp564N2216 (from clone DKFZp564N2216)	18.18	Down	3.85E-05
	NM_0201 30	Homo sapiens chromosome 8 open reading frame 4 (C8orf4), mRNA	17.97	Down	1.19E-04
45	M_1829 20	Homo sapiens a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 9 (ADAMTS9), transcript variant 1, mRNA	17.95	Down	5.64E-05
	AK02426 1	Homo sapiens cDNA FLJ14199 fis, clone NT2RP3002713	17.68	Down	2.24E-05
50	NM_0051 30	Homo sapiens fibroblast growth factor binding protein 1 (FGFBP1), mRNA	17.55	Down	5.73E-06
	BF43104 1	nab31g02.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3267627 3, mRNA sequence	17.25	Down	1.56E-05
55	BM992049	UI-H-DF1-auf-e-22-0-UI.s1 NCI_CGAP_DF1 Homo sapiens cDNA clone IMAGE:5868669 3, mRNA sequence	16.81	Drown	8.93E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0015 54	Homo sapiens cysteine-rich, angiogenic inducer, 61 (CYR61), mRNA	16.43	Down	6.87E-04
	AK09352 9	Homo sapiens cDNA FLJ36210 fis, clone THYMU2000155	16.29	Down	1.16E-05
10	NM_1983 89	Homo sapiens lung type-I cell membrane-associated glycoprotein (T1A-2), transcript variant 2, mRNA	16.15	Down	2.62E-05
	NM_0044 90	Homo sapiens growth factor receptor-bound protein 14 (GRB14), mRNA	16.02	Down	1.52E-04
15	NM_0019 99	Homo sapiens fibrillin 2 (congenital contractural arachnodactyly) (FBN2), mRNA	16	Down	4.45E-05
	NM_0124 13	Homo sapiens glutaminyl-peptide cyclotransferase (glutaminyl cyclase) (QPCT), mRNA	15.51	Down	1.39E-05
20	NM_1784 70	Homo sapiens WD repeat domain 40B (WDR40B), mRNA	15.46	Down	2.73E-05
	BM71294 5	UI-E-EJ0-ahi-c-16-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahi-c-16-0-UI 5, mRNA sequence	15.36	Down	4.73E-06
25	NM_0188 94	Homo sapiens EGF-containing fibulin-like extracellular matrix protein 1 (EFEMP1), transcript variant 2, mRNA	15.15	Down	5.35E-05
	BM71207 2	UI-E-DW1-ahc-b-11-0-UI.r1 UI-E-DW1 Homo sapiens cDNA clone UI-E-DW1-ahc-b-11-0-UI 5, mRNA sequence	15.1	Down	1.93E-05
30	NM_0005 19	Homo sapiens hemoglobin, delta (HBD), mRNA	14.92	Down	5.14E-06
	NM_0162 76	Homo sapiens serum/glucocorticoid regulated kinase 2 (SGK2), transcript variant 2, mRNA	14.87	Down	3.13E-06
	BF11171 0	7147c10.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3524371 3, mRNA sequence	14.78	Down	9.84E-05
35	AF26916 2	Homo sapiens c21orf7 form B mRNA, complete cds	14.77	Down	5.11E-05
	NM_2034 18	Homo sapiens Down syndrome critical region gene 1 (DSCR1), transcript variant 3, mRNA	14.75	Down	9.03E-04
40	NM_1735 05	Homo sapiens ankyrin repeat domain 29 (ANKRD29), mRNA	14.71	Down	8.60E-05
	NM_0325 11	Homo sapiens chromosome 6 open reading frame 168 (C6orf168), mRNA	14.65	Down	1.60E-05
45	NM_1527 68	Homo sapiens hypothetical protein FLJ25378 (FLJ25378), mRNA	14.59	Down	3.47E-05
	NM_0028 37	Homo sapiens protein tyrosine phosphatase, receptor type, B (PTPRB), mRNA	14.53	Down	8.08E-06
	NM_0006 82	Homo sapiens adrenergic, alpha-2B-, receptor (ADRA2B), mRNA	14.52	Down	5.46E-05
50	NM_015 41	Homo sapiens olfactomedin-like 2B 4 (OLFML2B), mRNA	14.35	Down	3.75E-05
	NM_006 75	Homo sapiens periostin, osteoblast specific 4 factor (POSTN), mRNA	14.33	Down	4.23E-05
55	BG4362 4	602508665F1 NIH_MGC_79 Homo sapiens 4 cDNA clone IMAGE:4605617 5, mRNA sequence	14.19	Down	3.75E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_025 45	Homo sapiens pre-B-cell leukemia transcription 2 factor 4 (PBX4), mRNA	14.1	Down	1.41E-05
10	NM_0005 57	Homo sapiens growth differentiation factor 5 (cartilage-derived morphogenetic protein-1) (GDF5), mRNA	14.06	Down	7.14E-05
	NM_079 23	Homo sapiens myosin, light polypeptide 6, 4 alkali, smooth muscle and non-muscle (MYL6), transcript variant 2, mRNA	14.03	Down	2.37E-05
15	NM_2075 17	Homo sapiens ADAMTS-like 3 (ADAMTSL3), mRNA	13.93	Down	1.39E-05
	NM_152 73	Homo sapiens RAS and EF hand domain 5 containing (RASEF), mRNA	13.93	Down	1.72E-04
	NM_0176 33	Homo sapiens family with sequence similarity 46, member A (FAM46A), mRNA	13.92	Down	3.30E-06
20	NM_016 88	Homo sapiens neuritin 1 (NRN1), mRNA 5	13.79	Down	1.39E-05
	NM_0209 62	Homo sapiens likely ortholog of mouse neighbor of Punc E11 (NOPE), mRNA	13.78	Down	2.37E-05
	NM_0324 61	Homo sapiens SPANX family, member B1 (SPANXB1), mRNA	13.57	Down	1.09E-05
25	BE78876 3	601475864F1 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:3879014 5, mRNA sequence	13.56	Down	1.39E-05
	AK12549 0	Homo sapiens cDNA FLJ43501 fis, clone PEBLM2004497	13.39	Down	3.11E-04
30	NM_0005 84	Homo sapiens interleukin 8 (IL8), mRNA	13.37	Down	1.66E-05
	M_1458 02	Homo sapiens septin 6 (SEPT6), transcript variant V, mRNA	13.04	Down	9.89E-06
	R99527	yq79b11.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:201981 3, mRNA sequence	13.01	Down	3.09E-05
35	NM_0325 10	Homo sapiens par-6 partitioning defective 6 homolog gamma (C. elegans) (PARD6G), mRNA	12.84	Down	3.42E-05
	NM_0005 99	Homo sapiens insulin-like growth factor binding protein 5 (IGFBP5), mRNA	12.83	Down	1.87E-05
40	NM_0008 76	Homo sapiens insulin-like growth factor 2 receptor (IGF2R), mRNA	12.74	Down	3.58E-04
	AL389942	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 2005635	12.52	Down	1.20E-04
45	AJ69797 2	Homo sapiens chromosome 3 cDNA	12.43	Down	1.17E-04
	H23441	ym52f11.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:51888 3, mRNA sequence	12.39	Down	2.04E-05
	NM_0162 01	Homo sapiens angiomin like 2 (AMOTL2), mRNA	12.38	Down	1.31E-03
	NM_0051 39	Homo sapiens annexin A3 (ANXA3), mRNA	12.27	Down	3.10E-06
50	W85910	zh52b10.s1 Soares_fetal_liver_spleen_1NFLS_S1 Homo sapiens cDNA clone IMAGE:415675 3, mRNA sequence	12.27	Down	4.22E-05
	NM_0048 62	Homo sapiens lipopolysaccharide-induced TNF factor (LITAF), mRNA	12.27	Down	1.03E-04
55	TM_1780 33	Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1 (CYP4X1), mRNA	12.13	Down	3.22E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0201 52	Homo sapiens chromosome 21 open reading frame 7 (C21orf7), mRNA	12.12	Down	4.27E-06
	NM_0041 65	Homo sapiens Ras-related associated with diabetes (RRAD), mRNA	11.95	Down	1.10E-04
10	NM_005450	Homo sapiens noggin (NOG), mRNA	11.93	Down	7.24E-06
	NM_0002 12	Homo sapiens integrin, beta 3 (platelet glycoprotein IIIa, antigen CD61) (ITGB3), mRNA	11.9	Down	8.68E-05
	BX64896 4	Homo sapiens mRNA; cDNA DKFZp686J0156 (from clone DKFZp686J0156)	11.85	Down	1.22E-05
15	NM_0314 76	Homo sapiens hypothetical protein DKFZp434B044 (DKFZP434B044), mRNA	11.84	Down	1.04E-05
	NM_1452 01	Homo sapiens similar to CG3714 gene product (PP3856), mRNA	11.57	Down	1.39E-05
20	NM_0161 79	Homo sapiens transient receptor potential cation channel, subfamily C, member 4 (TRPC4), mRNA	11.56	Down	2.61 E-04
	NM_0052 02	Homo sapiens collagen, type VIII, alpha 2 (COL8A2), mRNA	11.39	Down	2.61 E-04
25	AW0216 86	df26h11.y1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:2484717 5, mRNA sequence	11.2	Down	2.21 E-05
	BC04141 2	Homo sapiens heat shock 70kDa protein 12A, mRNA (cDNA clone IMAGE:5285193), partial cds	11.17	Down	2.88E-04
	NM_0044 33	Homo sapiens E74-like factor 3 (ets domain transcription factor, epithelial-specific) (ELF3), mRNA	11.14	Down	2.80E-05
30	NM_0011 44	Homo sapiens autocrine motility factor receptor (AMFR), transcript variant 1, mRNA	11.14	Down	5.52E-05
	NM_006 63	Homo sapiens leukocyte immunoglobulin-like 8 receptor, subfamily A (with TM domain), member 1 (LILRA1), mRNA	11.04	Down	2.72E-05
35	BX11331 9	BX113319 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGp998G205398 ; IMAGE:2184619, mRNA sequence	10.97	Down	2.51E-05
40	NM_002 62	Homo sapiens myxovirus (influenza virus) 4 resistance 1, interferon-inducible protein p78 (mouse) (MX1), mRNA	10.93	Down	1.17E-04
	AB01109 5	Homo sapiens mRNA for KIAA0523 protein, partial cds	10.88	Down	8.65E-06
45	NM_001 54	Homo sapiens collagen, type XI, alpha 1 8 (COL11A1), transcript variant A, mRNA	10.81	Down	9.37E-06
	NM_178 34	Homo sapiens layilin (LOC143903), mRNA 8	10.8	Down	5.28E-06
	NM_0121 98	Homo sapiens grancalcin, EF-hand calcium binding protein (GCA), mRNA	10.71	Down	1.57E-05
50	NM 2074 82	Homo sapiens FLJ44048 protein (FLJ44048), mRNA	10.69	Down	3.75E-05
	NM_000 17	Homo sapiens glutamate decarboxylase 1 8 (brain, 67kDa) (GAD1), transcript variant GAD67, mRNA	10.65	Down	1.35E-05
	NM_004 89	Homo sapiens nuclear factor (erythroid-derived 2 2)-like 3 (NFE2L3), mRNA	10.42	Down	4.53E-05
55	NM_0070 72	Homo sapiens HERV-H LTR-associating 2 (HHLA2), mRNA	10.41	Down	7.19E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AI355761	qt94a11.x1 NCI_CGAP_Co14 Homo sapiens cDNA clone IMAGE:1962908 3 similar to gb:X74929 KERATIN, TYPE II CYTOSKELETAL 8 (HUMAN);, mRNA sequence	10.38	Down	2.80E-05
10	NM_0219 77	Homo sapiens solute carrier family 22 (extraneuronal monoamine transporter), member 3 (SLC22A3), mRNA	10.3	Down	1.20E-04
15	AL13769 8	Homo sapiens mRNA; cDNA DKFZp434C1915 (from clone DKFZp434C1915); partial cds	10.26	Down	1.30E-04
	BM97638 5	UI-CF-EN1-acz-f-03-0-UI.s1 UI-CF-EN1 Homo sapiens cDNA clone UI-CF-EN1-acz-f-03-0-UI 3, mRNA sequence	10.2	Down	1.22E-05
20	AK12331 9	Homo sapiens cDNA FLJ41325 fis, clone BRAMY2046871	10.05	Down	1.84E-05
	BX45904 3	BX459043 Homo sapiens PLACENTA Homo sapiens cDNA clone CS0DE011YN10 3-PRIME, mRNA sequence	9.97	Down	3.27E-04
25	NM_0002 27	Homo sapiens laminin, alpha 3 (LAMA3), transcript variant 2, mRNA	9.83	Down	5.83E-06
	AI335277	tb29h06.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2055803 3, mRNA sequence	9.8	Down	7.24E-06
	NM_2073 80	Homo sapiens FLJ43339 protein (FLJ43339), mRNA	9.73	Down	4.97E-05
30	NM_0033 28	Homo sapiens TXK tyrosine kinase (TXK), mRNA	9.7	Down	1.64E-05
	NM_0006 12	Homo sapiens insulin-like growth factor 2 (somatomedin A) (IGF2), mRNA	9.68	Down	2.36E-05
	NM_0053 02	Homo sapiens G protein-coupled receptor 37 (endothelin receptor type B-like) (GPR37), mRNA	9.63	Down	5.63E-06
35	NM_0023 38	Homo sapiens limbic system-associated membrane protein (LSAMP), mRNA	9.62	Down	4.14E-05
	NM_0252 39	Homo sapiens programmed cell death 1 ligand 2 (PDCD1 LG2), mRNA	9.57	Down	7.52E-05
40	NM_0004 78	Homo sapiens alkaline phosphatase, liver/bone/kidney (ALPL), mRNA	9.56	Down	1.39E-05
	AJ31880 5	AJ318805 Homo sapiens adipose tissue Homo sapiens cDNA clone 2040, mRNA sequence	9.55	Down	5.32E-05
	BG62270 7	602647476F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4768963 5, mRNA sequence	9.54	Down	2.76E-05
45	CN47859 7	UI-CF-FN0-aeo-g-21-0-UI.s1 UI-CF-FN0 Homo sapiens cDNA clone UI-CF-FN0-aeo-g-21-0-UI 3, mRNA sequence	9.49	Down	1.39E-05
	BF79809 8	RC1-CI0045-021000-021-f02 CI0045 Homo sapiens cDNA, mRNA sequence	9.46	Down	2.31 E-05
50	NM_0008 56	Homo sapiens guanylate cyclase 1, soluble, alpha 3 (GUCY1A3), mRNA	9.42	Down	7.20E-05
	AL11976 9	DKFZp761E1224_r1 761 (synonym: hamy2) Homo sapiens cDNA clone DKFZp761 E1224 5, mRNA sequence	9.33	Down	5.73E-06
55	NM_0055 96	Homo sapiens nuclear factor I/B (NFIB), mRNA	9.32	Down	2.30E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AW 1344 73	UI-H-BI1-abv-a-11-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2712885 3, mRNA sequence	9.29	Down	1.31E-04
10	BX53769 8	Homo sapiens mRNA; cDNA DKFZp686F09166 (from clone DKFZp686F09166)	9.28	Down	4.13E-06
	NM_0244 23	Homo sapiens desmocollin 3 (DSC3), transcript variant Dsc3b, mRNA	9.27	Down	2.36E-05
	NM_0060 72	Homo sapiens chemokine (C-C motif) ligand 26 (CCL26), mRNA	9.27	Down	9.91E-06
15	NM_1735 08	Homo sapiens solute carrier family 35, member F3 (SLC35F3), mRNA	9.24	Down	5.64E-05
	NM_1780 12	Homo sapiens tubulin, beta polypeptide paralog (MGC8685), mRNA	9.24	Down	2.21E-04
20	NM_0332 90	Homo sapiens midline 1 (Opitz/BBB syndrome) (MID1), transcript variant 3, mRNA	9.18	Down	8.22E-04
	NM_018265	Homo sapiens hypothetical protein FLJ10901 (FLJ10901), mRNA	8.95	Down	1.17E-04
25	H89526	yw28b04.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253519 5, mRNA sequence	8.91	Down	1.74E-05
	NM_0009 61	Homo sapiens prostaglandin 12 (prostacyclin) synthase (PTGIS), mRNA	8.89	Down	2.82E-04
	AI942360	wo80c06.x1 NCI CGAP Kid11 Homo sapiens cDNA clone IMAGE:2461642 3, mRNA sequence	8.88	Down	2.07E-05
30	BX53761 3	Homo sapiens mRNA; cDNA DKFZp686E11117 (from clone DKFZp686E11117)	8.86	Down	1.59E-05
	NM_0143 58	Homo sapiens C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 9 (CLECSF9), mRNA	8.84	Down	2.21E-05
35	AK02399 9	Homo sapiens cDNA FLJ13937 fis, clone Y79AA1000805	8.83	Down	1.32E-04
	NM_0006 92	Homo sapiens aldehyde dehydrogenase 1 family, member B1 (ALDH1B1), nuclear gene encoding mitochondrial protein, mRNA	8.8	Down	4.89E-05
40	NM_0331 36	Homo sapiens fibroblast growth factor 1 (acidic) (FGF1), transcript variant 2, mRNA	8.8	Down	6.22E-05
	AK00007 5	Homo sapiens cDNA FLJ20068 fis, clone COL01755	8.8	Down	1.85E-05
45	NM_0186 59	Homo sapiens cytokine-like protein C17 (C17), mRNA	8.77	Down	7.06E-05
	AK05533 4	Homo sapiens cDNA FLJ30772 fis, clone FEBRA2000757, moderately similar to Homo sapiens BM-009 mRNA	8.59	Down	5.63E-06
50	NM_0039 79	Homo sapiens G protein-coupled receptor, family C, group 5, member A (GPCR5A), mRNA	8.53	Down	4.21E-06
	NM_0142 43	Homo sapiens a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 3 (ADAMTS3), mRNA	8.5	Down	5.37E-05
55	CA43786 1	UI-H-DH0-aur-k-12-0-UI.s1 NCI_CGAP_DH0 Homo sapiens cDNA clone UI-H-DH0-aur-k-12-0-UI 3, mRNA sequence	8.44	Down	7.74E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0186 63	Homo sapiens peroxisomal membrane protein 2, 22kDa (PXMP2), mRNA	8.43	Down	5.70E-04
	AI953708	wq47d09.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:2474417 3, mRNA sequence	8.39	Down	4.14E-05
10	BC03431 5	Homo sapiens hypothetical protein LOC90529, mRNA (cDNA clone IMAGE:4827425), containing frame-shift errors	8.37	Down	1.84E-04
	CB11575 4	K-EST0159876 L8SCK0 Homo sapiens cDNA clone L8SCK0-8-H08 5, mRNA sequence	8.36	Down	7.40E-06
15	NM_0050 26	Homo sapiens phosphoinositide-3-kinase, catalytic, delta polypeptide (PIK3CD), mRNA	8.3	Down	1.40E-04
	AL70665 3	DKFZp686E1543_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone DKFZp686E1543 5, mRNA sequence	8.28	Down	2.59E-05
20	M60502	Human profilaggrin mRNA, 3 end	8.28	Down	2.62E-05
	NM_0529 97	Homo sapiens ankyrin repeat domain 30A (ANKRD30A), mRNA	8.28	Down	4.24E-06
25	NM_0202 47	Homo sapiens chaperone, ABC1 activity of bc1 complex like (S. pombe) (CABC1), mRNA	8.26	Down	9.37E-06
	NM_0180 76	Homo sapiens armadillo repeat containing 4 (ARMC4), mRNA	8.22	Down	4.95E-05
	NM_0071 67	Homo sapiens zinc finger protein 258 (ZNF258), mRNA	8.22	Down	6.47E-04
30	NM_0166 13	Homo sapiens hypothetical protein DKFZp434L142 (DKFZp434L142), mRNA	8.21	Down	1.46E-05
	NM_0165 42	Homo sapiens Mst3 and SOK1-related kinase (MST4), mRNA	8.19	Down	1.06E-04
35	NM_0005 81	Homo sapiens glutathione peroxidase 1 (GPX1), transcript variant 1, mRNA	8.18	Down	3.53E-04
	AK09211 4	Homo sapiens cDNA FLJ34795 fis, clone NT2NE2005921	8.11	Down	3.10E-06
	AK12477 6	Homo sapiens cDNA FLJ42786 fis, clone BRAWH3006761	8.11	Down	2.02E-05
40	AK12470 2	Homo sapiens cDNA FLJ42712 fis, clone BRAMY3008044	8.1	Down	6.75E-06
	NM_0033 85	Homo sapiens visinin-like 1 (VSNL1), mRNA	8.1	Down	3.19E-06
45	NM_0309 15	Homo sapiens likely ortholog of mouse limb-bud and heart gene (LBH), mRNA	8.09	Down	3.86E-05
	CD72379 8	oj26f04.y1 Human lacrimal gland, unamplified: oj Homo sapiens cDNA clone oj26f04 5, mRNA sequence	8.08	Down	2.04E-04
50	BE96859 6	601649770F1 NIH_MGC_74 Homo sapiens cDNA clone IMAGE:3933472 5, mRNA sequence	8.05	Down	2.29E-05
	AB02321 1	Homo sapiens mRNA for KIAA0994 protein, partial cds	8.05	Down	1.74E-04
	NM_0040 45	Homo sapiens ATX1 antioxidant protein 1 homolog (yeast) (ATOX1), mRNA	8.02	Down	1.42E-04
55	AK07409 7	Homo sapiens mRNA for FLJ00168 protein	7.99	Down	2.58E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	M_1817 18	Homo sapiens hypothetical protein LOC253982 (LOC253982), mRNA	7.98	Down	2.03E-04
10	NM_0041 54	Homo sapiens pyrimidinergic receptor P2Y, G-protein coupled, 6 (P2RY6), transcript variant 4, mRNA	7.94	Down	9.37E-06
	NM_0143 91	Homo sapiens ankyrin repeat domain 1 (cardiac muscle) (ANKRD1), mRNA	7.92	Down	1.03E-04
15	NM_0806 59	Homo sapiens similar to RIKEN cDNA 2310030G06 gene (MGC14839), mRNA	7.89	Down	2.67E-05
	NM_0123 19	Homo sapiens solute carrier family 39 (zinc transporter), member 6 (SLC39A6), mRNA	7.87	Down	4.83E-04
	NM_1391 61	Homo sapiens crumbs homolog 3 (Drosophila) (CRB3), transcript variant 2, mRNA	7.86	Down	2.20E-03
20	BX64829 9	Homo sapiens mRNA; cDNA DKFZp686J04125 (from clone DKFZp686J04125)	7.85	Down	1.74E-05
	H94320	yv18b10.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:243067 3, mRNA sequence	7.84	Down	8.68E-05
25	NM_0024 23	Homo sapiens matrix metalloproteinase 7 (matrilysin, uterine) (MMP7), mRNA	7.83	Down	4.28E-04
	AK05801 2	Homo sapiens cDNA FLJ25283 fis, clone STM06716	7.7	Down	5.51E-05
	T56535	yb33g07.r1 Stratagene fetal spleen (#937205) Homo sapiens cDNA clone IMAGE:73020 5, mRNA sequence	7.69	Down	2.72E-05
30	NM_0020 53	Homo sapiens guanylate binding protein 1, interferon-inducible, 67kDa (GBP1), mRNA	7.68	Down	9.91E-06
	NM_0017 10	Homo sapiens B-factor, properdin (BF), mRNA	7.64	Down	4.89E-05
35	AK127421	Homo sapiens cDNA FLJ45513 fis, clone BRTHA2021450	7.63	Down	8.90E-05
	NM_0160 61	Homo sapiens yippee-like 5 (Drosophila) (YPEL5), mRNA	7.58	Down	5.26E-04
40	NM_0068 55	Homo sapiens KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 3 (KDELR3), transcript variant 1, mRNA	7.57	Down	4.41E-04
	AI311296	ta48d10.x2 NCI_CGAP_Lu25 Homo sapiens cDNA clone IMAGE:2047315 3, mRNA sequence	7.55	Down	1.93E-04
45	NM_0038 10	Homo sapiens tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10), mRNA	7.55	Down	4.68E-04
	AI819863	wj45h05.x1 NCI_CGAP_Lu19 Homo sapiens cDNA clone IMAGE:2405817 3, mRNA sequence	7.54	Down	2.62E-05
	NM_0000 76	Homo sapiens cyclin-dependent kinase inhibitor 1C (p57, Kip2) (CDKN1C), mRNA	7.53	Down	2.31E-05
50	NM_0037 61	Homo sapiens vesicle-associated membrane protein 8 (endobrevin) (VAMP8), mRNA	7.53	Down	6.69E-05
	NM_0034 75	Homo sapiens chromosome 11 open reading frame 13 (C11orf13), mRNA	7.5	Down	1.89E-04
55	NM_1984 95	Homo sapiens CTAGE family, member 4 (CTAGE4), mRNA	7.47	Down	2.61 E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AL117454	Homo sapiens mRNA; cDNA DKFZp586J1717 (from clone DKFZp586J1717)	7.4	Down	2.32E-04
	NM_0251 49	Homo sapiens hypothetical protein FLJ20920 (FLJ20920), mRNA	7.4	Down	1.67E-05
10	NM_002E 22	Homo sapiens neuronal pentraxin I (NPTX1), mRNA	7.37	Down	2.29E-05
	NM_0028 67	Homo sapiens RAB3B, member RAS oncogene family (RAB3B), mRNA	7.37	Down	3.28E-05
	AK02163 7	Homo sapiens cDNA FLJ11575 fis, clone HEMBA1003531	7.36	Down	4.12E-05
15	AL35905 8	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 592473	7.35	Down	1.12E-05
	AF10809 3	Homo sapiens IA-2 gene, intron 18	7.33	Down	5.57E-05
	AK02623 5	Homo sapiens cDNA: FLJ22582 fis, clone HSI02576	7.32	Down	1.55E-04
20	NM_0221 03	Homo sapiens hypothetical zinc finger protein FLJ14011 (FLJ14011), mRNA	7.31	Down	6.09E-05
	NM_0019 77	Homo sapiens glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	7.22	Down	9.96E-05
25	BX11152 0	BX111520 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998L15208 ; IMAGE:141470, mRNA sequence	7.21	Down	7.14E-06
	NM_0174 56	Homo sapiens pleckstrin homology, Sec7 and coiled-coil domains 1 (cytohesin 1) (PSCD1), transcript variant 2, mRNA	7.2	Down	1.76E-03
30	NM_0139 62	Homo sapiens neuregulin 1 (NRG1), transcript variant GGF2, mRNA	7.19	Down	2.88E-04
	S70348	Homo sapiens integrin beta 3 mRNA, partial cds, alternatively spliced	7.18	Down	2.47E-05
35	U51694	HSU51694 Human normal gingiva Homo sapiens cDNA, mRNA sequence	7.13	Down	6.48E-06
	NM_0012 99	Homo sapiens calponin 1, basic, smooth muscle (CNN1), mRNA	7.07	Down	2.91 E-05
	NM_0017 85	Homo sapiens cytidine deaminase (CDA), mRNA	7.02	Down	3.22E-05
40	NM_0052 96	Homo sapiens G protein-coupled receptor 23 (GPR23), mRNA	6.94	Down	1.65E-04
	NM_0305 94	Homo sapiens cytoplasmic polyadenylation element binding protein 1 (CPEB1), mRNA	6.94	Down	5.63E-06
45	BX11314 4	BX113144 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998N07225 ; IMAGE:148038, mRNA sequence	6.93	Down	4.14E-05
	AB02064 0	Homo sapiens mRNA for KIAA0833 protein, partial cds	6.91	Down	5.63E-06
50	AF24457 1	Homo sapiens clone L49 HERV-K-T47-like long terminal repeat sequence	6.86	Down	1.29E-04
	NM_0051 41	Homo sapiens fibrinogen, B beta polypeptide (FGB), mRNA	6.85	Down	2.24E-04
	NM_0530 01	Homo sapiens odd-skipped-related 2A protein (OSR2), mRNA	6.84	Down	2.30E-05
55	AA19572 7	zr33a09.r1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:665176 5, mRNA sequence	6.82	Down	2.73E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0011 65	Homo sapiens baculoviral IAP repeat-containing 3 (BIRC3), transcript variant 1, mRNA	6.81	Down	2.02E-04
	NM_0055 12	Homo sapiens glycoprotein A repetitions predominant (GARP), mRNA	6.81	Down	2.00E-04
10	NM_0000 87	Homo sapiens cyclic nucleotide gated channel alpha 1 (CNGA1), mRNA	6.81	Down	5.65E-05
	NM_0019 93	Homo sapiens coagulation factor III (thromboplastin, tissue factor) (F3), mRNA	6.74	Down	1.42E-04
15	AF04121 0	Homo sapiens midline 1 fetal kidney isoform 3 (MID1) mRNA, partial cds	6.73	Down	6.70E-05
	NM_0010 83	Homo sapiens phosphodiesterase 5A, cGMP-specific (PDE5A), transcript variant 1, mRNA	6.73	Down	7.71E-06
20	NM_0124 11	Homo sapiens protein tyrosine phosphatase, non-receptor type 22 (lymphoid) (PTPN22), transcript variant 2, mRNA	6.72	Down	2.36E-05
	NM_0309 65	Homo sapiens sialyltransferase 7 ((alpha-N-acetylneuraminyl-2,3-beta-galactosyl-1,3)-N-acetyl galactosaminide alpha-2,6-sialyltransferase) E (SIAT7E), mRNA	6.71	Down	3.90E-04
25	NM_0008 52	Homo sapiens glutathione S-transferase pi (GSTP1), mRNA	6.7	Down	6.35E-05
	BC04636 2	Homo sapiens voltage-dependent calcium channel gamma subunit-like protein, mRNA (cDNA clone MGC:50757 IMAGE:5221396), complete cds	6.7	Down	1.38E-05
30	NM_1445 69	Homo sapiens SPOC domain containing 1 (SPOCD1), mRNA	6.66	Down	5.03E-05
	NM_0060 58	Homo sapiens TNFAIP3 interacting protein 1 (TNIP1), mRNA	6.61	Down	1.01E-03
35	S54641	HZF-16=Kruppel-related zinc finger gene homolog {alternatively spliced} [human, hepatoblastoma cell line, HEP-G2, mRNA, 2080 nt]	6.59	Down	6.25E-04
	NM_0180 23	Homo sapiens YEATS domain containing 2 (YEATS2), mRNA	6.58	Down	5.94E-04
40	NM_0021 93	Homo sapiens inhibin, beta B (activin AB beta polypeptide) (INHBB), mRNA	6.56	Down	5.11E-05
	NM_0314 26	Homo sapiens chromosome 9 open reading frame 58 (C9orf58), transcript variant 1, mRNA	6.55	Down	4.93E-04
45	AI244954	qj93h05.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1867065 3, mRNA sequence	6.53	Down	5.14E-06
	BX53754 7	Homo sapiens mRNA; cDNA DKFZp686N1640 (from clone DKFZp686N1640)	6.51	Down	1.44E-04
50	NM_0009 81	Homo sapiens ribosomal protein L19 (RPL19), mRNA	6.49	Down	1.39E-04
	AK05801 2	Homo sapiens cDNA FLJ25283 fis, clone STM06716	6.48	Down	1.57E-05
	NM_0245 63	Homo sapiens hypothetical protein FLJ14054 (FLJ14054), mRNA	6.48	Down	2.40E-04
	NM_0018 43	Homo sapiens contactin 1 (CNTN1), transcript variant 1, mRNA	6.47	Down	1.91E-05
55	L07615	Human neuropeptide Y receptor Y1 (NPYY1) mRNA, exon 2-3 and complete cds	6.46	Down	2.22E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5					
	NM_0133 22	Homo sapiens sorting nexin 10 (SNX10), mRNA	6.44	Down	7.33E-05
	NM_0143 33	Homo sapiens immunoglobulin superfamily, member 4 (IGSF4), mRNA	6.41	Down	2.37E-05
10	NM_1735 49	Homo sapiens hypothetical protein FLJ39553 (FLJ39553), mRNA	6.41	Down	1.42E-05
	BF50992 5	UI-H-BI4-aph-c-10-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3087355 3, mRNA sequence	6.39	Down	4.05E-05
15	NM_003597	Homo sapiens TGFB inducible early growth response 2 (TIEG2), mRNA	6.39	Down	1.22E-05
	NM_152 69	Homo sapiens hypothetical protein MGC45474 3 (MGC45474), mRNA	6.39	Down	3.75E-05
	NM_1447 07	Homo sapiens prominin 2 (PROM2), mRNA 7	6.38	Down	1.04E-04
20	NM_0040 86	Homo sapiens coagulation factor C homolog, cochlin (Limulus polyphemus) (COCH), mRNA	6.37	Down	1.91E-05
	NM_0201 27	Homo sapiens tuftelin 1 (TUFT1), mRNA	6.37	Down	2.93E-03
	NM_0001 04	Homo sapiens cytochrome P450, family 1, subfamily B, polypeptide 1 (CYP1B1), mRNA	6.35	Down	1.46E-05
25	NM_2068 08	Homo sapiens citrate lyase beta like (CLYBL), transcript variant 2, mRNA	6.32	Down	5.22E-05
	NM_0035 67	Homo sapiens breast cancer anti-estrogen resistance 3 (BCAR3), mRNA	6.31	Down	1.39E-05
30	NM_0036 17	Homo sapiens regulator of G-protein signalling 5 (RGS5), mRNA	6.29	Down	2.36E-04
	AK13018 1	Homo sapiens cDNA FLJ26671 fis, clone MPG03325	6.28	Down	6.69E-05
	NM_0159 54	Homo sapiens CGI-26 protein (CGI-26), mRNA	6.27	Down	2.50E-04
	NM_0019 45	Homo sapiens diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth factor) (DTR), mRNA	6.27	Down	2.56E-03
35					
	NM_0005 97	Homo sapiens insulin-like growth factor binding protein 2, 36kDa (IGFBP2), mRNA	6.26	Down	2.29E-04
40	NM_0209 92	Homo sapiens PDZ and LIM domain 1 (elfin) (PDLIM1), mRNA	6.25	Down	8.97E-05
	NM_0005 76	Homo sapiens interleukin 1, beta (IL1B), mRNA	6.22	Down	2.34E-05
	NM_0228 18	Homo sapiens microtubule-associated protein 1 light chain 3 beta (MAP1LC3B), mRNA	6.22	Down	1.18E-03
45	NM_0012 57	Homo sapiens cadherin 13, H-cadherin (heart) (CDH13), mRNA	6.21	Down	4.57E-04
	AI697906	we18f06.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:2341475 3, mRNA sequence	6.21	Down	2.37E-05
	AK12569 5	Homo sapiens cDNA FLJ43707 fis, clone TESOP2001865	6.21	Down	1.16E-05
50					
	NM_0057 25	Homo sapiens tetraspan 2 (TSPAN-2), mRNA	6.21	Down	7.32E-05
	AI694344	wd45f11.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2331117 3 similar to contains MER13.b1 MER13 repetitive element; mRNA sequence	6.2	Down	1.44E-04
55					
	NM_0013 33	Homo sapiens cathepsin L2 (CTSL2), mRNA	6.19	Down	3.43E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AI623139	tu89b07.x1 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGE:2258197 3, mRNA sequence	6.18	Down	1.38E-04
10	NM_0067 27	Homo sapiens cadherin 10, type 2 (T2-cadherin) (CDH10), mRNA	6.17	Down	6.43E-05
	M80899	Human novel protein AHNAK mRNA, partial sequence	6.16	Down	6.38E-04
	NM_0122 13	Homo sapiens malonyl-CoA decarboxylase (MLYCD), mRNA	6.15	Down	3.70E-05
15	NM_0166 51	Homo sapiens dapper homolog 1, antagonist of beta-catenin (xenopus) (DACT1), mRNA	6.14	Down	6.41E-05
	NM_0001 29	Homo sapiens coagulation factor XIII, A1 polypeptide (F13A1), mRNA	6.14	Down	9.71E-06
20	NM_0064 72	Homo sapiens thioredoxin interacting protein (TXNIP), mRNA	6.11	Down	1.43E-03
	BU72709 6	UI-E-CR0-ach-e-12-0-UI.s1 UI-E-CR0 Homo sapiens cDNA clone UI-E-CR0-ach-e-12-0-UI 3, mRNA sequence	6.09	Down	7.54E-05
25	AF07057 1	Homo sapiens clone 24739 mRNA sequence	6.04	Down	8.86E-05
	NM_018120	Homo sapiens armadillo repeat containing 1 (ARMC1), mRNA	6.02	Down	2.51E-03
	CA41484 7	UI-H-EZ0-bar-b-22-0-UI.s1 NCI_CGAP_Ch1 Homo sapiens cDNA clone UI-H-EZ0-bar-b-22-0-UI 3, mRNA sequence	6.02	Down	4.69E-05
30	AW4747 73	xy06f10.x1 NCI_CGAP_Lym12 Homo sapiens cDNA clone IMAGE:2852395 3, mRNA sequence	6.01	Down	8.88E-05
	NM_0068 51	Homo sapiens GLI pathogenesis-related 1 (glioma) (GLIPR1), mRNA	6.01	Down	2.98E-05
35	NM_0249 97	Homo sapiens activating transcription factor 7 interacting protein 2 (ATF7IP2), mRNA	6	Down	1.65E-04
	NM_0039 32	Homo sapiens suppression of tumorigenicity 13 (colon carcinoma) (Hsp70 interacting protein) (ST13), mRNA	6	Down	4.81 E-04
40	T53523	ya89h12.r1 Stratagene placenta (#937225) Homo sapiens cDNA clone IMAGE:68903 5, mRNA sequence	5.99	Down	1.60E-04
	NM_0169 46	Homo sapiens F11 receptor (F11 R), transcript variant 1, mRNA	5.99	Down	7.76E-04
45	AJ40694 1	Homo sapiens partial mRNA for keratin associated protein 4.9 (KRTAP4.9 gene)	5.99	Down	1.74E-05
	BF51049 3	UI-H-BI4-apa-b-08-0-UI.s1 NCI_CGAP_Sub8 Homo sapiens cDNA clone IMAGE:3086558 3, mRNA sequence	5.93	Down	3.17E-05
50	AK09443 6	Homo sapiens cDNA FLJ37117 fis, clone BRACE2022270	5.92	Down	1.58E-04
	NM_032 73	Homo sapiens Cbl-interacting protein Sts-1 8 (STS-1), mRNA	5.92	Down	1.67E-05
55	NM_015 77	Homo sapiens retinoic acid induced 14 (RAI14), 5 mRNA	5.91	Down	3.19E-03
	NM_014 02	Homo sapiens KIAA0408 (KIAA0408), mRNA	5.91	Down	5.11E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AB00794 0	Homo sapiens mRNA for KIAA0471 protein, partial cds	5.88	Down	1.65E-04
	NM_0044 79	Homo sapiens fucosyltransferase 7 (alpha (1,3) fucosyltransferase) (FUT7), mRNA	5.87	Down	3.82E-05
10	BC03559 9	Homo sapiens, clone IMAGE:3871970, mRNA, partial cds	5.86	Down	9.99E-05
	NM_0013 32	Homo sapiens catenin (cadherin-associated protein), delta 2 (neural plakophilin-related arm-repeat protein) (CTNND2), mRNA	5.84	Down	1.73E-05
15	NM_0019 95	Homo sapiens acyl-CoA synthetase long-chain family member 1 (ACSL1 mRNA	5.83	Down	4.26E-04
	NM_0530 39	Homo sapiens UDP glycosyltransferase 2 family, polypeptide B28 (UGT2B28), mRNA	5.82	Down	4.93E-05
20	BC01510 8	Homo sapiens, Similar to otoconin 90, clone IMAGE:4044247, mRNA	5.81	Down	5.63E-06
	NM_0022 50	Homo sapiens potassium intermediate/small conductance calcium-activated channel, subfamily N, member 4 (KCNN4), mRNA	5.8	Down	1.60E-05
25	AL11742 5	Homo sapiens mRNA; cDNA DKFZp566L203 (from clone DKFZp566L203)	5.76	Down	6.05E-05
	NM_0148 67	Homo sapiens KIAA0711 gene product (KIAA0711), mRNA	5.74	Down	7.24E-06
30	BM47382 3	AGENCOURT_6484255 NIH_MGC_72 Homo sapiens cDNA clone IMAGE:5538225 5, mRNA sequence	5.73	Down	2.36E-05
	NM_0245 58	Homo sapiens chromosome 14 open reading frame 138 (C14orf138), mRNA	5.73	Down	8.48E-04
35	NM_0055 78	Homo sapiens LIM domain containing preferred translocation partner in lipoma (LPP), mRNA	5.71	Down	7.52E-05
	AK12417 5	Homo sapiens cDNA FLJ42181 fis, clone THYMU2031368	5.7	Down	2.30E-03
40	NM_0132 81	Homo sapiens fibronectin leucine rich transmembrane protein 3 (FLRT3), transcript variant 1, mRNA	5.69	Down	1.38E-05
	AA68929 2	nv66c11.s1 NCI_CGAP_GCB1 Homo sapiens cDNA clone IMAGE:1234772 3, mRNA sequence	5.69	Down	1.42E-04
45	NM_0026 22	Homo sapiens prefoldin 1 (PFDN1), mRNA	5.69	Down	1.69E-03
	BM695043	UI-E-CL1-aez-k-18-0-UI.r1 UI-E-CL1 Homo sapiens cDNA clone UI-E-CL1-aez-k-18-0-UI 5, mRNA sequence	5.68	Down	2.37E-03
	NM_0071 45	Homo sapiens zinc finger protein 146 (ZNF146), mRNA	5.67	Down	6.48E-04
50	NM_0040 78	Homo sapiens cysteine and glycine-rich protein 1 (CSRP1), mRNA	5.67	Down	2.33E-04
	N75271	yz74h12.r1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:288839 5, mRNA sequence	5.67	Down	7.47E-05
55	AL35394 4	Homo sapiens mRNA; cDNA DKFZp761J1112 (from clone DKFZp761 J1112)	5.66	Down	3.88E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AL04796 0	DKFZp586B0623_r1 586 (synonym: hute1) Homo sapiens cDNA clone DKFZp586B0623, mRNA sequence	5.63	Down	1.65E-04
10	AK13109 2	Homo sapiens mRNA for FLJ00307 protein	5.63	Down	2.25E-04
	NM_0208 08	Homo sapiens signal-induced proliferation-associated 1 like 2 (SIPA1L2), mRNA	5.63	Down	8.36E-04
	NM_0034 94	Homo sapiens dysferlin, limb girdle muscular dystrophy 2B (autosomal recessive) (DYSF), mRNA	5.6	Down	5.21E-05
15	NM_0328 83	Homo sapiens chromosome 20 open reading frame 100 (C20orf100), mRNA	5.58	Down	1.93E-05
	NM_006722	Homo sapiens microphthalmia-associated transcription factor (MITF), transcript variant 3, mRNA	5.57	Down	1.99E-04
20	NM_0149 78	Homo sapiens sortilin-related VPS10 domain containing receptor 3 (SORCS3), mRNA	5.57	Down	4.68E-05
	BX11041 8	BX110418 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998C224149 ; IMAGE:1635405, mRNA sequence	5.56	Down	5.10E-05
25	NM_0040 87	Homo sapiens discs, large homolog 1 (Drosophila) (DLG1), mRNA	5.56	Down	1.21E-04
	NM_0005 61	Homo sapiens glutathione S-transferase M1 (GSTM1), transcript variant 1, mRNA	5.54	Down	5.63E-06
	NM_0067 85	Homo sapiens mucosa associated lymphoid tissue lymphoma translocation gene 1 (MALT1), transcript variant 1, mRNA	5.53	Down	2.39E-04
30	NM_0528 39	Homo sapiens pannexin 2 (PANX2), mRNA	5.52	Down	2.09E-05
	NM_0250 85	Homo sapiens transcriptional coactivator tubedown-100 (TBDN100), transcript variant 2, mRNA	5.51	Down	1.35E-03
35	NM_0244 22	Homo sapiens desmocollin 2 (DSC2), transcript variant Dsc2a, mRNA	5.5	Down	7.12E-05
	NM_0174 48	Homo sapiens lactate dehydrogenase C (LDHC), transcript variant 2, mRNA	5.5	Down	8.50E-06
40	AK12803 6	Homo sapiens cDNA FLJ46155 fis, clone TESTI4001517, moderately similar to Keratin, type I cytoskeletal 18	5.5	Down	2.58E-03
	NM_0326 23	Homo sapiens ovary-specific acidic protein (OSAP), mRNA	5.47	Down	5.65E-04
45	BX11634 7	BX116347 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGp998B215967 ; IMAGE:2401844, mRNA sequence	5.47	Down	1.65E-04
	NM_0020 31	Homo sapiens fyn-related kinase (FRK), mRNA	5.45	Down	2.62E-05
	CA41374 4	UI-H-EZO-bat-h-12-0-UI.s1 NCI_CGAP_Ch1 Homo sapiens cDNA clone UI-H-EZO-bat-h-12-0-UI 3, mRNA sequence	5.45	Down	1.32E-04
50	T78754	yd01f08.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:24180 5, mRNA sequence	5.45	Down	1.14E-04
	NM_0050 82	Homo sapiens tripartite motif-containing 25 (TRIM25), mRNA	5.44	Down	4.71E-04
55					

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_0328 57	Homo sapiens lactamase, beta (LACTB), nuclear gene encoding mitochondrial protein, transcript variant 1, mRNA	5.43	Down	6.74E-04
10	NM_1736 62	Homo sapiens hypothetical protein LOC285533 (LOC285533), mRNA	5.43	Down	1.72E-04
	NM_0021 66	Homo sapiens inhibitor of DNA binding 2, dominant negative helix-loop-helix protein (ID2), mRNA	5.43	Down	2.13E-04
	NM_0005 93	Homo sapiens transporter 1, ATP-binding cassette, sub-family B (MDR/TAP) (TAP1), mRNA	5.42	Down	5.84E-04
15		UI-H-DH0-arx-p-21-0-UI.s1 NCI_CGAP_DH0	5.42	Down	2.89E-05
	BM98864 2	Homo sapiens cDNA clone IMAGE:5855492 3, mRNA sequence			
20	NM_0249 01	Homo sapiens hypothetical protein FLJ22457 (FLJ22457), mRNA	5.41	Down	4.87E-04
	NM_0178 23	Homo sapiens dual specificity phosphatase 23 (DUSP23), mRNA	5.41	Down	5.55E-04
	BX10783 8	BX107838 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998A153853 ; IMAGE:1521686, mRNA sequence	5.39	Down	2.44E-04
25					
	AW 1729 03	xj05e04.x1 NCI_CGAP_Ut2 Homo sapiens cDNA clone IMAGE:2656350 3, mRNA sequence	5.39	Down	1.67E-05
	BX11132 1	BX111321 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGp998C221112 ; IMAGE:469197, mRNA sequence	5.39	Down	1.46E-03
30					
	NM_0332 11	Homo sapiens hypothetical gene supported by AF038182; BC009203 (LOC90355), mRNA	5.38	Down	1.15E-04
	D62831	HUM330B12B Clontech human aorta polyA+ mRNA (#6572) Homo sapiens cDNA clone GEN-330B12 5, mRNA sequence	5.37	Down	5.80E-03
35					
	NM_0010 69	Homo sapiens tubulin, beta polypeptide (TUBB), mRNA	5.36	Down	4.71E-04
	NM_0043 24	Homo sapiens BCL2-associated X protein (BAX), transcript variant beta, mRNA	5.36	Down	8.14E-04
40					
	AV736303	AV736303 CB Homo sapiens cDNA clone CBCAJD04 5, mRNA sequence	5.36	Down	4.51E-05
	AK13153 2	Homo sapiens cDNA FLJ16761 fis, clone BRAMY3008096	5.34	Down	9.55E-05
45					
	NM_0208 59	Homo sapiens Shroom-related protein (ShrmL), mRNA	5.34	Down	1.21E-04
	NM_0025 75	Homo sapiens serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2 (SERPINB2), mRNA	5.34	Down	8.70E-05
50					
	NM_0175 99	Homo sapiens transmembrane protein vezatin (VEZATIN), mRNA	5.33	Down	6.62E-04
	BX11607 1	BX116071 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGp998L201165 ; IMAGE:489763, mRNA sequence	5.33	Down	1.39E-04
55					
	NM_0246 77	Homo sapiens hypothetical protein FLJ14001 (FLJ14001), mRNA	5.32	Down	5.07E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	AL83277 9	Homo sapiens mRNA; cDNA DKFZp686H157 (from clone DKFZp686H157)	5.32	Down	4.37E-04
	NM_0025 47	Homo sapiens oligophrenin 1 (OPHN1), mRNA	5.32	Down	5.03E-05
10	NM_0143 22	Homo sapiens opsin 3 (encephalopsin, panopsin) (OPN3), mRNA	5.3	Down	4.04E-04
	NM_0030 57	Homo sapiens solute carrier family 22 (organic cation transporter), member 1 (SLC22A1), transcript variant 1, mRNA	5.29	Down	1.92E-04
15	BM969191	UI-CF-EN0-acp-e-22-0-UI.s1 UI-CF-EN0 Homo sapiens cDNA clone UI-CF-EN0-acp-e-22-0-UI 3, mRNA sequence	5.28	Down	4.06E-04
	S69208	troponin T [human, skeletal and cardiac muscle, mRNA, 932 nt]	5.28	Down	7.73E-04
20	NM_0312 16	Homo sapiens SEH1-like (S. cerevisiae) (SEH1L), mRNA	5.27	Down	1.17E-03
	NM_0029 67	Homo sapiens scaffold attachment factor B (SAFB), mRNA	5.27	Down	5.52E-04
25	NM_0028 43	Homo sapiens protein tyrosine phosphatase, receptor type, J (PTPRJ), mRNA	5.27	Down	5.63E-06
	NM_0186 64	Homo sapiens Jun dimerization protein p21SNFT (SNFT), mRNA	5.27	Down	3.06E-05
	NM_0209 44	Homo sapiens glucosidase, beta (bile acid) 2 (GBA2), mRNA	5.25	Down	3.07E-04
30	BM98720 0	UI-H-CO0-aql-b-05-0-UI.s1 NCI_CGAP_Sub9Homo sapiens cDNA clone IMAGE:3104192 3, mRNA sequence	5.24	Down	8.03E-05
	NM_0239 15	Homo sapiens G protein-coupled receptor 87 (GPR87), mRNA	5.23	Down	2.05E-04
35	NM_0156 78	Homo sapiens neurobeachin (NBEA), mRNA	5.23	Down	2.95E-05
	NM_004438	Homo sapiens EPH receptor A4 (EPHA4), mRNA	5.22	Down	1.93E-05
	NM_0022 06	Homo sapiens integrin, alpha 7 (ITGA7), mRNA	5.22	Down	2.36E-05
40	AW5914 61	xI92h06.x1 NCI_CGAP_Ut1 Homo sapiens cDNA clone IMAGE:2682203 3, mRNA sequence	5.22	Down	7.14E-05
	AK05603 3	Homo sapiens cDNA FLJ31471 fis, clone NT2NE2001435	5.19	Down	3.29E-04
	NM_0182 35	Homo sapiens CNDP dipeptidase 2 (metallopeptidase M20 family) (CNDP2), mRNA	5.19	Down	3.74E-03
45	AV72829 4	AV728294 HTC Homo sapiens cDNA clone HTCBIE09 5, mRNA sequence	5.19	Down	3.05E-04
	NM_0247 25	Homo sapiens hypothetical protein FLJ23518 (FLJ23518), mRNA	5.18	Down	2.56E-03
50	NM_0018 74	Homo sapiens carboxypeptidase M (CPM), transcript variant 1, mRNA	5.18	Down	4.89E-05
	W30761	zb76g12.r1 Soares_senescent_fibroblasts_NbHSF Homo sapiens cDNA clone IMAGE:309574 5, mRNA sequence	5.15	Down	1.61E-04
55	BM74199 7	K-EST0014724 S7SNU719 Homo sapiens cDNA clone S7SNU719-7-F05 5, mRNA sequence	5.14	Down	1.15E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change in AF-I cells versus AF-III cells	Direction	adj. p-value
5	NM_017594	Homo sapiens DIRAS family, GTP-binding RAS-like 2 (DIRAS2), mRNA	5.14	Down	1.38E-03
	NM_0223 43	Homo sapiens chromosome 9 open reading frame 19 (C9orf19), mRNA	5.13	Down	2.67E-05
10	NM_0032 87	Homo sapiens tumor protein D52-like 1 (TPD52L1 transcript variant 1, mRNA	5.13	Down	2.71E-05
	BC06710 6	Homo sapiens putative G protein coupled receptor, mRNA (cDNA clone MGC:71222 IMAGE:6645890), complete cds	5.12	Down	2.13E-04
15	NM_1818 28	Homo sapiens neurofibromin 2 (bilateral acoustic neuroma) (NF2), transcript variant 5, mRNA	5.12	Down	1.99E-03
	NM_0052 64	Homo sapiens GDNF family receptor alpha 1 (GFRA1), transcript variant 1, mRNA	5.11	Down	4.23E-05
20	AK09215 7	Homo sapiens cDNA FLJ34838 fis, clone NT2NE2010654	5.11	Down	6.46E-05
	NM_0229 12	Homo sapiens chromosome 2 open reading frame 23 (C2orf23), mRNA	5.09	Down	5.42E-05
25	NM_0529 37	Homo sapiens similar to hypothetical protein FLJ10883 (LOC115294), mRNA	5.09	Down	1.38E-04
	BM66545 2	UI-E-CQ1-aex-n-03-0-UI.s1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aex-n-03-0-UI 3, mRNA sequence	5.08	Down	3.24E-04
30	NM_0038 31	Homo sapiens RIO kinase 3 (yeast) (RIOK3), transcript variant 1, mRNA	5.08	Down	1.64E-04
	NM_1816 89	Homo sapiens neuronatin (NNAT), transcript variant 2, mRNA	5.08	Down	7.40E-06
	BC03312 4	Homo sapiens, clone IMAGE:2960615, mRNA	5.07	Down	3.52E-05
35	NM_0160 03	Homo sapiens WIPI49-like protein 2 (DKFZP434J154), transcript variant 2, mRNA	5.07	Down	6.36E-03
	BC03855 6	Homo sapiens, clone IMAGE:3446976, mRNA	5.06	Down	2.32E-04
	AW0695 77	cr47c06.x1 Human bone marrow stromal cells Homo sapiens cDNA clone HBMSC_cr47c06 3, mRNA sequence	5.06	Down	2.08E-03
40	NM_0000 55	Homo sapiens butyrylcholinesterase (BCHE), mRNA	5.05	Down	1.20E-05
	NM_0056 02	Homo sapiens claudin 11 (oligodendrocyte transmembrane protein) (CLDN11), mRNA	5.04	Down	7.46E-04
45	NM_0309 52	Homo sapiens likely ortholog of rat SNF1/AMP-activated protein kinase (SNARK), mRNA	5.03	Down	8.60E-05
	AK09237 9	Homo sapiens cDNA FLJ35060 fis, clone OCBBF2018828	5.03	Down	1.67E-05
	AK09501 3	Homo sapiens cDNA FLJ37694 fis, clone BRHIP2015224	5.01	Down	1.87E-04
50	NM_1452 34	Homo sapiens chordin-like 1 (CHRD1), mRNA	5.01	Down	1.11E-05
	NM_1992 87	Homo sapiens similar to RIKEN cDNA 3110023B02 (MGC16597), mRNA	5	Down	9.27E-05
55	NM_0251 51	Homo sapiens RAB11 family interacting protein 1 (class I) (RAB11FIP1), transcript variant 1, mRNA	5	Down	4.32E-05

EP 1 780 267 B1

TABLE VII G: GENES THAT WERE DIFFERENTIALLY EXPRESSED AT **LEAST 5 FOLD IN AF-II VERSUS AF-III CELLS**

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_003411	Homo sapiens zinc finger protein, Y-linked (ZFY), mRNA	328.28	Up	1.97E-06
10	NM_153634	Homo sapiens copine VIII (CPNE8), mRNA	286.5	Up	6.16E-06
	H70730	yu69e10.r1 Weizmann Olfactory Epithelium Homo sapiens cDNA clone IMAGE:239082 5, mRNA sequence	272.44	Up	1.97E-06
15	NM_138963	Homo sapiens ribosomal protein S4, Y-linked 2 (RPS4Y2), mRNA	265.55	Up	1.56E-06
	NM_004653	Homo sapiens Smcy homolog, Y-linked (mouse) (SMCY), mRNA	246.17	Up	1.00E-06
	NM_004660	Homo sapiens DEAD (Asp-Glu-Ala-Asp) box polypeptide 3, Y-linked (DDX3Y), mRNA	244.93	Up	1.17E-05
20	NM_012307	Homo sapiens erythrocyte membrane protein band 4.1-like 3 (EPB41L3), mRNA	207.61	Up	1.83E-06
	BX08955 4	BX089554 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998P07210 ; IMAGE:142326, mRNA sequence	119.7	Up	2.95E-06
25	NM_001008	Homo sapiens ribosomal protein S4, Y-linked 1 (RPS4Y1), mRNA	112.79	Up	2.02E-05
	NM_014893	Homo sapiens neuroligin 4, Y-linked (NLGN4Y), mRNA	93.96	Up	9.61E-06
30	NM_182798	Homo sapiens hypothetical protein FLJ39155 (FLJ39155), transcript variant 2, mRNA	71.46	Up	1.36E-05
	NM_080872	Homo sapiens unc-5 homolog D (C. elegans) (UNC5D), mRNA	48.16	Up	5.41E-05
	BX64864 3	Homo sapiens mRNA; cDNA DKFZp686O17106 (from clone DKFZp686O17106)	47.84	Up	2.95E-06
35	NM_002928	Homo sapiens regulator of G-protein signalling 16 (RGS16), mRNA	47.75	Up	2.99E-05
	AB03780 5	Homo sapiens mRNA for KIAA1384 protein, partial cds	45.54	Up	7.68E-06
40	NM_031862	Homo sapiens membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125) (M17S2), transcript variant 3, mRNA	44.02	Up	5.15E-05
	CA42913 5	UI-H-FH1-bfh-k-22-0-UI.s1 NCI_CGAP_FH1 Homo sapiens cDNA clone UI-H-FH1-bfh-k-22-0-UI 3, mRNA sequence	43.87	Up	2.90E-06
45	BC043411	Homo sapiens, clone IMAGE:6155889, mRNA	42.03	Up	6.97E-06
	NM_004681	Homo sapiens eukaryotic translation initiation factor 1A, Y-linked (EIF1AY), mRNA	41.7	Up	2.24E-04
50	NM_181481	Homo sapiens chromosome 18 open reading frame 1 (C18orf1), transcript variant a1, mRNA	39.6	Up	9.61E-06
	BE46576 0	hw22f09.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3183689 3, mRNA sequence	39.51	Up	1.50E-05
	NM_002667	Homo sapiens phospholamban (PLN), mRNA	37.56	Up	1.87E-05
55	AK09807 1	Homo sapiens cDNA FLJ40752 fis, clone TRACH2000972	34.71	Up	1.02E-04
	NM_001864	Homo sapiens cytochrome c oxidase subunit VIIa polypeptide 1 (muscle) (COX7A1), mRNA	33.43	Up	1.08E-06

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	AF05537 6	Homo sapiens shortform transcription factor C-MAF (c-maf) mRNA, complete cds	32.71	Up	9.41E-04
	NM_004932	Homo sapiens cadherin 6, type 2, K-cadherin (fetal kidney) (CDH6), mRNA	31.51	Up	1.37E-05
10	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	30.4	Up	1.97E-06
	NM_018349	Homo sapiens multiple C2-domains with two transmembrane regions 2 (MCTP2), mRNA	29.49	Up	1.68E-05
	NM_000990	Homo sapiens ribosomal protein L27a (RPL27A), mRNA	28.23	Up	4.25E-04
15	BX11135 3	BX111353 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGp998O094576 ; IMAGE:1869152, mRNA sequence	27.4	Up	3.27E-06
	NM_032576	Homo sapiens chromosome Y open reading frame 15B (CYorf15B), mRNA	27.08	Up	3.10E-06
20	NM_020997	Homo sapiens left-right determination factor 1 (LEFTY1), mRNA	26.55	Up	1.42E-05
	NM_006408	Homo sapiens anterior gradient 2 homolog (Xenopus laevis) (AGR2), mRNA	25.93	Up	2.36E-05
25	BQ92483 2	AGENCOURT_8840265 Lupski_sciatic_nerve Homo sapiens cDNA clone IMAGE:6205036 5, mRNA sequence	24.29	Up	2.00E-06
	NM_019000	Homo sapiens hypothetical protein FLJ20152 (FLJ20152), mRNA	23.73	Up	7.10E-05
	BC03565 6	Homo sapiens hypothetical protein LOC285835, mRNA (cDNA clone IMAGE:5588650), partial cds	23.25	Up	1.82E-06
30	AI032876	ow13g03.x1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:1646740 3, mRNA sequence	22.96	Up	5.20E-06
	NM_001305	Homo sapiens claudin 4 (CLDN4), mRNA	21.95	Up	3.72E-04
35	BX08955 4	BX089554 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGp998P07210 ; IMAGE:142326, mRNA sequence	20.73	Up	2.00E-06
	CF13754 5	UI-HF-BN0-ane-d-05-0-UI.r1 NIH_MGC_50 Homo sapiens cDNA clone IMAGE:3092384 5, mRNA sequence	20.24	Up	7.57E-05
40	CB04709 2	NISC_gf08f03.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:3253013 3, mRNA sequence	20.2	Up	4.77E-06
	NM_005114	Homo sapiens heparan sulfate (glucosamine) 3-O-sulfotransferase 1 (HS3ST1), mRNA	19.91	Up	2.92E-05
45	BQ43078 8	AGENCOURT_7776027 NIH_MGC_68 Homo sapiens cDNA clone IMAGE:6024295 5, mRNA sequence	19.78	Up	9.59E-04
	AK02423 8	Homo sapiens cDNA FLJ14176 fis, clone NT2RP2003101	19.33	Up	2.32E-05
	NM_001001931	Homo sapiens mitochondrial tumor suppressor 1 (MTUS1), nuclear gene encoding mitochondrial protein, transcript variant 4, mRNA	18.91	Up	2.90E-06
50	NM_152423	Homo sapiens melanoma associated antigen (mutated) 1-like 1 (MUM1L1), mRNA	18.66	Up	1.97E-06
	BX11359 0	BX113590 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGp998A14871 ; IMAGE:376597, mRNA sequence	18.22	Up	3.83E-05
55	NM_0025	Homo sapiens neuronal PAS domain protein 2 (NPAS2), mRNA	17.82	Up	3.88E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	18				
	AK12871 5	Homo sapiens cDNA FLJ46882 fis, clone UTERU3015844	17.2	Up	1.17E-03
	AA73825 4	nx13b02.s1 NCI_CGAP_GC3 Homo sapiens cDNA clone IMAGE:1255947 3, mRNA sequence	16.61	Up	6.24E-06
10	NM_0033 81	Homo sapiens vasoactive intestinal peptide (VIP), transcript variant 1, mRNA	16.44	Up	5.33E-06
	AI951740	wv38h09.x1 NCI_CGAP_Ov18 Homo sapiens cDNA clone IMAGE:2531873 3, mRNA sequence	14.99	Up	6.49E-03
15	NM_0063 93	Homo sapiens nebulin (NEBL), transcript variant 1, mRNA	14.87	Up	1.84E-03
	NM_0009 63	Homo sapiens prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2), mRNA	14.83	Up	3.02E-03
20	H89053	yw24c06.r1 Morton Fetal Cochlea Homo sapiens cDNA clone IMAGE:253162 5, mRNA sequence	14.76	Up	6.87E-03
	D86975	Homo sapiens mRNA for KIAA0222 gene, partial cds	14.7	Up	2.87E-04
	AA10255 3	zn26a04.s1 Stratagene neuroepithelium NT2RAMI 937234 Homo sapiens cDNA clone IMAGE:548526 3, mRNA sequence	13.56	Up	2.86E-04
25	AK09670 8	Homo sapiens cDNA FLJ39389 fis, clone PLACE6003621	13.38	Up	2.90E-05
	BM68512 4	UI-E-EJ1-ajl-I-13-0-UI.s1 UI-E-EJ1 Homo sapiens cDNA clone UI-E-EJ1-ajl-I-13-0-UI 3, mRNA sequence	12.74	Up	2.49E-04
	AI220066	qg84d01.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1841857 3, mRNA sequence	12.47	Up	1.90E-04
30	AI342246	qt26g09.x1 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGE:1949152 3, mRNA sequence	12.29	Up	7.15E-04
	NM_0245 01	Homo sapiens homeo box D1 (HOXD1), mRNA	12.11	Up	9.73E-06
35	NM_0163 07	Homo sapiens paired related homeobox 2 (PRRX2), mRNA	12	Up	9.50E-04
	AB02069 1	Homo sapiens mRNA for KIAA0884 protein, partial cds	11.69	Up	1.47E-04
	BE29546 8	601174523F1 NIH_MGC_17 Homo sapiens cDNA clone IMAGE:3529924 5, mRNA sequence	11.64	Up	1.47E-03
40	BU95146 9	in60a05.x3 HR85 islet Homo sapiens cDNA clone IMAGE:6126249 3, mRNA sequence	11.37	Up	3.28E-05
	M_1981 74	Homo sapiens transcription factor CP2-like 4 (TFCP2L4), transcript variant 3, mRNA	11.26	Up	3.35E-05
45	NM_0024 21	Homo sapiens matrix metalloproteinase 1 (interstitial collagenase) (MMP1), mRNA	11.12	Up	5.10E-04
	TM_1453 13	Homo sapiens RasGEF domain family, member 1A (RASGEF1A), mRNA	10.64	Up	2.38E-05
	NM_0009 01	Homo sapiens nuclear receptor subfamily 3, group C, member 2 (NR3C2), mRNA	10.62	Up	3.93E-06
50	AL55252 7	AL552527 Homo sapiens PLACENTA COT 25-NORMALIZED Homo sapiens cDNA clone CSODI067YL24 3-PRIME, mRNA sequence	10.61	Up	3.88E-03
55	NM_0529 54	Homo sapiens cysteine and tyrosine-rich 1 (CYR1), mRNA	10.58	Up	7.22E-05
	AF51962 2	Homo sapiens noncoding mRNA sequence	10.39	Up	1.49E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_006681	Homo sapiens neuromedin U (NMU), mRNA	10.37	Up	2.92E-05
	W93585	zd95g01.s1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:357264 3, mRNA sequence	10.27	Up	2.01E-04
10	BX10515 2	BX105152 Soares_testis_NHT Homo sapiens cDNA clone IMAGp998J212575 ; IMAGE:1031156, mRNA sequence	9.89	Up	4.37E-05
	AV70923 2	AV709232 ADC Homo sapiens cDNA clone ADCAJB11 5, mRNA sequence	9.89	Up	1.56E-04
	W69644	zd45f10.r1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:343627 5, mRNA sequence	9.89	Up	1.90E-05
15	NM_001878	Homo sapiens cellular retinoic acid binding protein 2 (CRABP2), mRNA	9.82	Up	2.65E-04
	NM_152754	Homo sapiens sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3D (SEMA3D), mRNA	9.78	Up	6.57E-04
20	BE87776 4	601486331 F1 NIH_MGC_69 Homo sapiens cDNA clone IMAGE:3888943 5, mRNA sequence	9.59	Up	1.90E-04
	AK02590 9	Homo sapiens cDNA: FLJ22256 fis, clone HRC02860	9.52	Up	8.65E-05
25	NM_015559	Homo sapiens SET binding protein 1 (SETBP1), mRNA	9.3	Up	3.69E-04
	AI493349	tg70f04.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:2114143 3, mRNA sequence	9.21	Up	1.86E-05
	AK12995 5	Homo sapiens cDNA FLJ26445 fis, clone KDN02608	9.2	Up	1.41E-04
30	BX09852 1	BX098521 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGp998L05118 ; IMAGE:123412, mRNA sequence	9.06	Up	9.68E-04
	AK09429 2	Homo sapiens cDNA FLJ36973 fis, clone BRACE2006249	9.05	Up	3.44E-05
	AK09577 6	Homo sapiens cDNA FLJ38457 fis, clone FEBRA2020400	8.98	Up	1.13E-05
35	NM_018242	Homo sapiens hypothetical protein FLJ10847 (FLJ10847), mRNA	8.98	Up	8.02E-04
	AW0068 64	ws15d04.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2497255 3, mRNA sequence	8.93	Up	1.50E-05
40	BX10384 6	BX103846 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGp998F125811 ; IMAGE:2342027, mRNA sequence	8.89	Up	1.38E-04
	NM_024422	Homo sapiens desmocollin 2 (DSC2), transcript variant Dsc2a, mRNA	8.84	Up	3.19E-05
45	NM_032471	Homo sapiens protein kinase (cAMP-dependent, catalytic) inhibitor beta (PKIB), transcript variant 3, mRNA	8.77	Up	1.35E-03
	AK12477 8	Homo sapiens cDNA FLJ42788 fis, clone BRAWH3007129	8.62	Up	4.29E-04
	H25898	yl55b10.r1 Soares breast 3NbHBst Homo sapiens cDNA clone IMAGE:162139 5, mRNA sequence	8.52	Up	1.29E-04
50	NM_015564	Homo sapiens leucine rich repeat transmembrane neuronal 2 (LRRTM2), mRNA	8.5	Up	3.01E-03
	AI126888	qb95d06.x1 Soares_fetal_heart_NbHH19W Homo sapiens cDNA clone IMAGE:1707851 3, mRNA sequence	8.5	Up	6.30E-05
	BC04202 8	Homo sapiens, clone IMAGE:4794726, mRNA	8.46	Up	3.65E-02
55	NM_006561	Homo sapiens CUG triplet repeat, RNA binding protein 2 (CUGBP2), mRNA	8.37	Up	2.52E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_004675	Homo sapiens ras homolog gene family, member I (ARHI), mRNA	8.36	Up	4.99E-04
	BE50391 6	hz35g01.x1 NCI_CGAP_GC6 Homo sapiens cDNA clone IMAGE:3210000 3, mRNA sequence	8.3	Up	1.00E-05
10	NM_207446	Homo sapiens hypothetical gene supported by AK075564; BC060873 (LOC400451), mRNA	8.28	Up	2.72E-05
	AK09173 1	Homo sapiens cDNA FLJ34412 fis, clone HEART2002432	8.08	Up	9.36E-05
	NM_002261	Homo sapiens killer cell lectin-like receptor subfamily C, member 3 (KLRC3), transcript variant NKG2-E, mRNA	8.07	Up	2.69E-04
15	AW2709 28	xs06c05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2768840 3, mRNA sequence	7.99	Up	4.03E-05
	NM_005079	Homo sapiens tumor protein D52 (TPD52), mRNA	7.97	Up	4.79E-05
20	NM_001353	Homo sapiens aldo-keto reductase family 1, member C1 (dihydrodiol dehydrogenase 1; 20-alpha (3-alpha)-hydroxysteroid dehydrogenase) (AKR1C1), mRNA	7.88	Up	3.67E-04
	AI830524	wh52c02.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2384354 3, mRNA sequence	7.82	Up	1.23E-03
25	BX64820 7	Homo sapiens mRNA; cDNA DKFZp686E16168 (from clone DKFZp686E16168)	7.82	Up	9.59E-04
	AW0437 93	wy76d11.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:2554485 3 similar to contains element MER18 repetitive element ;, mRNA sequence	7.79	Up	4.64E-05
30	N63415	yy60d04.s1 Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:277927 3 similar to contains L1.b3 L1 repetitive element ;, mRNA sequence	7.76	Up	1.14E-03
	NM_002260	Homo sapiens killer cell lectin-like receptor subfamily C, member 2 (KLRC2), mRNA	7.76	Up	3.65E-04
35	NM_004848	Homo sapiens chromosome 1 open reading frame 38 (C1orf38), mRNA	7.75	Up	5.34E-04
	NM_024993	Homo sapiens leucine rich repeat transmembrane neuronal 4 (LRRTM4), mRNA	7.53	Up	3.35E-05
40	AI821210	ne08e05.y5 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGE:880640 5, mRNA sequence	7.51	Up	5.58E-04
	BM66444 5	UI-E-CL1-afa-p-05-0-UI.s1 UI-E-CL1 Homo sapiens cDNA clone UI-E-CL1-afa-p-05-0-UI 3, mRNA sequence	7.4	Up	5.30E-04
	NM_019644	Homo sapiens ankyrin repeat domain 7 (ANKRD7), mRNA	7.39	Up	2.24E-04
45	BU62079 3	UI-H-FL1-bfx-d-10-0-UI.s1 NCI_CGAP_FL1 Homo sapiens cDNA clone UI-H-FL1-bfx-d-10-0-UI 3, mRNA sequence	7.35	Up	1.13E-05
	NM_00246	Homo sapiens potassium channel, 2 subfamily K, member 3 (KCNK3), mRNA	7.31	Up	1.68E-05
50	NM_03142	Homo sapiens transmembrane 4 4 superfamily member 10 (TM4SF10), mRNA	7.13	Up	2.31E-03
	BX09250 1	BX092501 NCI_CGAP_Lu5 Homo sapiens cDNA clone IMAGp998K143946 ; IMAGE:1557637, mRNA sequence	7.12	Up	1.69E-03
55	NM_177949	Homo sapiens armadillo repeat containing, X-linked 2 (ARMCX2), mRNA	7.09	Up	9.21E-03
	NM_018376	Homo sapiens nipsnap homolog 3B (C. elegans) (NIPSNAP3B), mRNA	7.03	Up	2.53E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	AK09376 2	Homo sapiens cDNA FLJ36443 fis, clone THYMU2012891	7.02	Up	1.23E-03
	NM_018658	Homo sapiens potassium inwardly-rectifying channel, subfamily J, member 16 (KCNJ16), transcript variant 1, mRNA	7.01	Up	6.28E-04
10	AB03294 5	Homo sapiens mRNA for KIAA1119 protein, partial cds	6.98	Up	1.11E-04
	NM_031894	Homo sapiens ferritin, heavy polypeptide-like 17 (FTHL17), mRNA	6.97	Up	3.42E-03
		UI-H-BI1-ae-a-12-0-UI.s1	6.96	Up	3.35E-05
15	AW 139891	NCI-CGAP-Sub3 Homo sapiens cDNA clone IMAGE:2719006 3, mRNA sequence			
	AK12456 2	Homo sapiens cDNA FLJ42571 fis, clone BRACE3008036	6.9	Up	2.57E-05
	NM_001935	Homo sapiens dipeptidylpeptidase 4 (CD26, adenosine deaminase complexing protein 2) (DPP4), mRNA	6.9	Up	1.08E-03
20	AI686652	tu35d06.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2253035 3, mRNA sequence	6.87	Up	4.03E-06
	NM_184087	Homo sapiens tripartite motif-containing 55 (TRIM55), transcript variant 4, mRNA	6.85	Up	1.68E-05
	NM_002089	Homo sapiens chemokine (C-X-C motif) ligand 2 (CXCL2), mRNA	6.81	Up	9.44E-05
25	NM_016831	Homo sapiens period homolog 3 (Drosophila) (PER3), mRNA	6.81	Up	6.24E-06
	NM_002515	Homo sapiens neuro-oncological ventral antigen 1 (NOVA1), transcript variant 1, mRNA	6.79	Up	4.79E-05
30	BC01290 0	Homo sapiens, clone IMAGE:3881549, mRNA	6.77	Up	1.84E-02
	NM_052923	Homo sapiens zinc finger protein 452 (ZNF452), mRNA	6.75	Up	2.98E-03
	NM_021800	Homo sapiens DnaJ (Hsp40) homolog, subfamily C, member 12 (DNAJC12), transcript variant 1, mRNA	6.73	Up	5.46E-04
35	AK05688 2	Homo sapiens cDNA FLJ32320 fis, clone PROST2003537	6.71	Up	9.55E-04
	BF70068 4	602128672F1 NIH_MGC_56 Homo sapiens cDNA clone IMAGE:4285673 5, mRNA sequence	6.7	Up	4.03E-06
40	NM_144664	Homo sapiens hypothetical protein MGC33371 (MGC33371), mRNA	6.66	Up	1.64E-03
	AL83277 9	Homo sapiens mRNA; cDNA DKFZp686H157 (from clone DKFZp686H157)	6.62	Up	8.14E-05
	NM_018371	Homo sapiens chondroitin beta1,4 N-acetylgalactosaminyltransferase (ChGn), mRNA	6.6	Up	6.54E-04
45	NM_014399	Homo sapiens transmembrane 4 superfamily member 13 (TM4SF13), mRNA	6.56	Up	9.52E-03
	AA74876 2	ny06h10.s1 NCI_CGAP_GCB1 Homo sapiens cDNA clone IMAGE:1271011 3, mRNA sequence	6.54	Up	3.49E-03
50	AF21607 7	Homo sapiens clone HB-2 mRNA sequence	6.5	Up	4.90E-05
	AA04325 5	zk49f07.s1 Soares_pregnant_uterus_NbHPU Homo sapiens cDNA clone IMAGE:486181 3, mRNA sequence	6.49	Up	5.44E-04
	NM_152864	Homo sapiens chromosome 20 open reading frame 58 (C20orf58), mRNA	6.44	Up	2.32E-03
55	NM_032866	Homo sapiens cingulin-like 1 (CGNL1), mRNA	6.4	Up	3.69E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	AK05499 0	Homo sapiens cDNA FLJ30428 fis, clone BRACE2008941	6.37	Up	3.89E-05
	NM_1389 61	Homo sapiens endothelial cell adhesion molecule (ESAM), mRNA	6.35	Up	2.70E-04
	AI244954	qj93h05.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:1867065 3, mRNA sequence	6.27	Up	3.90E-05
10	NM_0198 50	Homo sapiens neuronal guanine nucleotide exchange factor (NGEF), mRNA	6.25	Up	4.99E-04
	NM_0073 61	Homo sapiens nidogen 2 (osteonidogen) (NID2), mRNA	6.15	Up	7.03E-03
15	NM_0001 70	Homo sapiens glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P) (GLDC), mRNA	6.13	Up	2.12E-05
	BG57203 9	602592506F1 NIH_MGC_79 Homo sapiens cDNA clone IMAGE:4719888 5, mRNA sequence	6.09	Up	1.13E-05
20	NM_0057 95	Homo sapiens calcitonin receptor-like (CALCRL), mRNA	6.03	Up	3.13E-02
	BQ26780	ij94e04.x1 Human insulinoma Homo sapiens cDNA clone IMAGE:5779278 3, mRNA sequence	6.02	Up	6.53E-04
25	6				
	BC03622 3	Homo sapiens, clone IMAGE:5272183, mRNA	6	Up	3.15E-04
	NM_004 70	Homo sapiens collagen, type XII, alpha 1 3 (COL12A1), transcript variant long, mRNA	5.96	Up	1.14E-02
	NM_0053 60	Homo sapiens v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian) (MAF), mRNA	5.95	Up	2.41E-03
30	AB03304 8	Homo sapiens mRNA for KIAA1222 protein, partial cds	5.94	Up	3.49E-04
	NM_017 86	Homo sapiens hypothetical protein 7 FLJ20366 (FLJ20366), mRNA	5.93	Up	2.24E-04
	AW5151 14	xu91g11.x1 NCI_CGAP_Ut2 Homo sapiens cDNA clone IMAGE:2809124 3, mRNA sequence	5.92	Up	1.11E-04
35	AK09854 3	Homo sapiens cDNA FLJ25677 fis, clone TST04054	5.84	Up	4.99E-03
	NM_004 35	Homo sapiens Kruppel-like factor 4 (gut) 2 (KLF4), mRNA	5.83	Up	6.54E-04
	AI335277	tb29h06.x1 NCI_CGAP_Kid12 Homo sapiens cDNA clone IMAGE:2055803 3, mRNA sequence	5.81	Up	1.24E-04
40	AK09500 4	Homo sapiens cDNA FLJ37685 fis, clone BRHIP2013972	5.76	Up	3.79E-03
	BQ00715 6	UI-H-EI1-azc-k-08-0-UI.s1 NCI_CGAP_EI1 Homo sapiens cDNA clone IMAGE:5846911 3, mRNA sequence	5.68	Up	1.24E-04
	NM_0168 24	Homo sapiens adducin 3 (gamma) (ADD3), transcript variant 1, mRNA	5.68	Up	6.32E-04
45	AA99433 0	ou33h05.s1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:1628121 3, mRNA sequence	5.67	Up	9.57E-05
	NM_0169 46	Homo sapiens F11 receptor (F11 R), transcript variant 1, mRNA	5.65	Up	7.59E-05
50	NM_0053 08	Homo sapiens G protein-coupled receptor kinase 5 (GRK5), mRNA	5.63	Up	7.59E-05
	BG38932 8	602413981 F1 NIH_MGC_92 Homo sapiens cDNA clone IMAGE:4522269 5, mRNA sequence	5.61	Up	1.31E-04
55	CN37116 8	17000600077294 GRN_PREHEP Homo sapiens cDNA 5, mRNA sequence	5.57	Up	1.94E-02
	AK02153 1	Homo sapiens cDNA FLJ11469 fis, clone HEMBA1001658	5.56	Up	7.81E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_001853	Homo sapiens collagen, type IX, alpha 3 (COL9A3), mRNA	5.56	Up	6.94E-05
	BC03792 9	Homo sapiens cDNA clone IMAGE:5284659, partial cds oz40h01.s1 Soares_NhHMPu_S1 Homo	5.56 5.54	Up Up	9.99E-04 1.20E-03
10	AI086279	sapiens cDNA clone IMAGE:1677841 3, mRNA sequence			
	BF43347 9	7q53g05.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3702320 3 similar to TR:Q9Y7P8 Q9Y7P8 HYPOTHETICAL 11.7 KD PROTEIN. ;, mRNA sequence	5.48	Up	1.23E-04
	AI024717	ov68h06.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE:1642523 3, mRNA sequence	5.46	Up	1.59E-04
15	NM_033082	Homo sapiens cytokine induced protein 29 kDa (CIP29), mRNA	5.45	Up	9.59E-04
	W56431	zc57f06.r1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:326435 5, mRNA sequence	5.42	Up	3.76E-04
20	NM_144587	Homo sapiens chromosome 10 open reading frame 87 (C10orf87), mRNA	5.41	Up	1.47E-04
	BX10148 9	BX101489 Soares_NFL_T_GBC_S Homo sapiens cDNA clone IMAGp998O165825 ; IMAGE:2347623, mRNA sequence	5.37	Up	1.71E-05
25	NM_014059	Homo sapiens response gene to complement 32 (RGC32), mRNA	5.32	Up	1.48E-04
	AI939617	tm62e06.x5 NCI_CGAP_Brn25 Homo sapiens cDNA clone IMAGE:2162722 3, mRNA sequence	5.3	Up	1.84E-03
30	NM_004951	Homo sapiens Epstein-Barr virus induced gene 2 (lymphocyte-specific G protein-coupled receptor) (EBI2), mRNA	5.3	Up	8.80E-04
	AK09505 3	Homo sapiens cDNA FLJ37734 fis, clone BRHIP2020842	5.3	Up	2.53E-04
	NM_024420	Homo sapiens phospholipase A2, group IVA (cytosolic, calcium-dependent) (PLA2G4A), mRNA	5.3	Up	1.56E-02
35	NM_018013	Homo sapiens hypothetical protein FLJ10159 (FLJ10159), mRNA	5.29	Up	2.12E-05
	BX10185 0	BX101850 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGp998M14139 ; IMAGE:26298, mRNA sequence	5.25	Up	7.59E-05
40	C18094	C18094 Human placenta cDNA (TFujiwara) Homo sapiens cDNA clone GEN-557D07 5, mRNA sequence	5.23	Up	1.00E-05
	BU63433 2	UI-H-FL1-bgx-k-02-0-UI.s1 NCI_CGAP_FL Homo sapiens cDNA clone UI-H-FL1-bgx-k-02-0-UI 3, mRNA sequence	5.22	Up	2.86E-04
45	AI033863	ow10e02.x1 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGE:1646426 3, mRNA sequence	5.18	Up	4.89E-03
	BG57014 4	602591134F1 NIH_MGC_77 Homo sapiens cDNA clone IMAGE:4717761 5, mRNA sequence	5.18	Up	4.36E-04
50	NM_018427	Homo sapiens RRN3 RNA polymerase I transcription factor homolog (yeast) (RRN3), mRNA	5.17	Up	1.68E-03
	BX48620	DKFZp686J07250_r1 686 (synonym: hlcc3) Homo sapiens cDNA clone	5.14	Up	1.31E-03
	8	DKFZp686J07250 5, mRNA sequence			
55	NM_014936	Homo sapiens ectonucleotide pyrophosphatase/phosphodiesterase 4 (putative function) (ENPP4), mRNA	5.14	Up	3.43E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	AF08613 4	Homo sapiens full length insert cDNA clone ZA88B06	5.14	Up	4.00E-04
	NM_0018 47	Homo sapiens collagen, type IV, alpha 6 (COL4A6), transcript variant A, mRNA	5.12	Up	1.03E-03
	AK09572 6	Homo sapiens cDNA FLJ38407 fis, clone FEBRA2008859	5.1	Up	1.35E-04
10	BQ01386 9	UI-1-BC1p-alg-a-03-0-UI.s1 NCI_CGAP_PI3 Homo sapiens cDNA clone UI-1-BC1p-alg-a-03-0-UI 3, mRNA sequence	5.09	Up	1.42E-05
	NM_2073 03	Homo sapiens attractin-like 1 (ATRNL1), mRNA	5.04	Up	4.36E-04
15	CB04728 7	NISC_gg01h01.y1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:3253464 5, mRNA sequence	5.04	Up	1.16E-03
	NM_0024 23	Homo sapiens matrix metalloproteinase 7 (matrilysin, uterine) (MMP7), mRNA	5.02	Up	6.94E-05
20	AW2626 83	xq93c06.x1 NCI_CGAP_Brn53 Homo sapiens cDNA clone IMAGE:2758186 3, mRNA sequence	5.02	Up	3.19E-05
	NM_0132	Homo sapiens transgelin 3 (TAGLN3), transcript variant 1, mRNA	5.01	Up	8.63E-04
	59				
25	D52654	HUM084D02B Clontech human fetal brain polyA+ mRNA (#6535) Homo sapiens cDNA clone GEN-084D02 5, mRNA sequence	449.36	Down	9.61E-06
	NM_0064 75	Homo sapiens periostin, osteoblast specific factor (POSTN), mRNA	293.88	Down	9.16E-05
30	NM_0064 39	Homo sapiens mab-21-like 2 (C. elegans) (MAB21L2), mRNA	199.11	Down	3.53E-05
	NM_0178 05	Homo sapiens Ras interacting protein 1 (RASIP1 mRNA	153.9	Down	9.55E-05
35	NM_0061 69	Homo sapiens nicotinamide N-methyltransferase (NNMT), mRNA	91.55	Down	1.57E-05
	AB06749 9	Homo sapiens mRNA for KIAA1912 protein, partial cds	69.67	Down	1.13E-05
	NM_0070 84	Homo sapiens SRY (sex determining region Y)-box 21 (SOX21), mRNA	55.37	Down	8.65E-05
40	NM_0320 26	Homo sapiens TatD DNase domain containing 1 (TATDN1), mRNA	52.52	Down	2.24E-04
	NM_0178 19	Homo sapiens RNA (guanine-9-) methyltransferase domain containing 1 (RG9MTD1), mRNA	47.78	Down	2.09E-04
	NM_0328	Homo sapiens chromosome 20 open reading frame 100 (C20orf100), mRNA	45.19	Down	8.58E-05
45	83				
	BQ37571 9	QV2-TN0173-021100-454-g06 TN0173 Homo sapiens cDNA, mRNA sequence	44.03	Down	1.41E-04
	BC03731 6	Homo sapiens, clone IMAGE:5259432, mRNA	34.66	Down	2.12E-05
50	BC04237 8	Homo sapiens, clone IMAGE:5277693, mRNA	32.65	Down	6.44E-04
	NM_0188 94	Homo sapiens EGF-containing fibulin-like extracellular matrix protein 1 (EFEMP1), transcript variant 2, mRNA	30.19	Down	2.74E-05
	NM_0330 50	Homo sapiens succinate receptor 1 (SUCNR1), mRNA	30.08	Down	1.95E-04
55	NM_1526 94	Homo sapiens zinc finger, CCHC domain containing 5 (ZCCHC5), mRNA	29.85	Down	7.57E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_0529 97	Homo sapiens ankyrin repeat domain 30A (ANKRD30A), mRNA	28.91	Down	5.75E-04
	AK02678 4	Homo sapiens cDNA: FLJ23131 fis, clone LNG08502	27.95	Down	8.14E-05
10	NM_0018 85	Homo sapiens crystallin, alpha B (CRYAB), mRNA	26.31	Down	5.48E-05
	NM_0056 02	Homo sapiens claudin 11 (oligodendrocyte transmembrane protein) (CLDN11), mRNA	23.71	Down	1.13E-05
	NM_0014 42	Homo sapiens fatty acid binding protein 4, adipocyte (FABP4), mRNA	23.48	Down	4.05E-04
15	NM_0216 14	Homo sapiens potassium intermediate/small conductance calcium-activated channel, subfamily N, member 2 (KCNN2), transcript variant 1, mRNA	23.14	Down	9.61E-06
	AK12387 5	Homo sapiens cDNA FLJ41881 fis, clone OCBBF2021833	22.73	Down	9.17E-05
20	R44402	yg37a01.s1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:34639 3 similar to contains MER35 repetitive element ;, mRNA sequence	22.27	Down	2.38E-05
	AK09501 3	Homo sapiens cDNA FLJ37694 fis, clone BRHIP2015224	20.95	Down	9.77E-04
	NM_0021 85	Homo sapiens interleukin 7 receptor (IL7R), mRNA	20.88	Down	4.62E-04
25	NM_0162 12	Homo sapiens TP53TG3 protein (TP53TG3), mRNA	20.63	Down	6.56E-05
	NM_1527 37	Homo sapiens hypothetical protein MGC33993 (MGC33993), mRNA	19.89	Down	6.16E-06
30	BF00248 9	7h07e07.x1 NCI_CGAP_Co16 Homo sapiens cDNA clone IMAGE:3315300 3, mRNA sequence	19.88	Down	8.65E-05
	NM_0055 25	Homo sapiens hydroxysteroid (11-beta) dehydrogenase 1 (HSD11B1), transcript variant 1, mRNA	18.27	Down	3.53E-05
35	AF31838 2	Homo sapiens pp9974 mRNA, complete cds	17.67	Down	3.89E-05
	BX11582 5	BX115825 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGp998C134603 ; IMAGE:1879236, mRNA sequence	17.5	Down	1.67E-04
40	NM_0006 12	Homo sapiens insulin-like growth factor 2 (somatomedin A) (IGF2), mRNA	16.41	Down	4.79E-05
	AK09211 4	Homo sapiens cDNA FLJ34795 fis, clone NT2NE2005921	16.29	Down	2.18E-05
	AI469032	ti70a01.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2137320 3, mRNA sequence	15.92	Down	9.36E-05
45	NM_0011 75	Homo sapiens Rho GDP dissociation inhibitor (GDI) beta (ARHGDI), mRNA	15.57	Down	4.62E-05
	AF05211 5	Homo sapiens clone 23688 mRNA sequence	15.11	Down	1.03E-04
	CD67733 2	ho15f06.y1 Human Trabecular meshwork cDNA: hohphq Homo sapiens cDNA clone ho15f06 5, mRNA sequence	14.89	Down	8.06E-05
50	NM_0180 77	Homo sapiens RNA binding motif protein 28 (RBM28), mRNA	14.72	Down	9.00E-05
	NM_0007 35	Homo sapiens glycoprotein hormones, alpha polypeptide (CGA), mRNA	14.51	Down	2.62E-04
55	N49730	yz06a12.s1	13.53	Down	2.76E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5		Soares_multiple_sclerosis_2NbHMSP Homo sapiens cDNA clone IMAGE:282238 3 similar to contains Alu repetitive element;contains element PTR5 repetitive element ;, mRNA sequence			
10	NM_006350	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	13.34	Down	2.53E-04
	AK055518	Homo sapiens cDNA FLJ30956 fis, clone HCASM2000202	13.09	Down	2.63E-03
	NM_016147	Homo sapiens protein phosphatase methylesterase-1 (PME-1), mRNA	13.04	Down	1.84E-02
15	NM_032457	Homo sapiens BH-protocadherin (brain-heart) (PCDH7), transcript variant c, mRNA	13.01	Down	7.57E-05
	BX648299	Homo sapiens mRNA; cDNA DKFZp686J04125 (from clone DKFZp686J04125)	12.94	Down	1.82E-04
20	BC034315	Homo sapiens hypothetical protein LOC90529, mRNA (cDNA clone IMAGE:4827425), containing frame-shift errors	12.94	Down	2.72E-04
	AA195328	zr34f08.s1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE:665319 3, mRNA sequence	12.73	Down	7.59E-04
25	AI082507	ox55c02.s1 Soares_total_fetus_Nb2HF8_9w Homo sapiens cDNA clone IMAGE:1660226 3, mRNA sequence	12.67	Down	5.54E-05
	NM_015714	Homo sapiens putative lymphocyte G0/G1 switch gene (G0S2), mRNA	12.17	Down	7.45E-04
	BM992049	UI-H-DF1-auf-e-22-0-UI.s1 NCI_CGAP_DF1 Homo sapiens cDNA clone IMAGE:5868669 3, mRNA sequence	12.09	Down	8.65E-05
30	NM_016428	Homo sapiens ABI gene family, member 3 (ABI3), mRNA	12.02	Down	3.62E-04
	BQ027984	UI-H-CO0-arg-e-03-0-UI.s1 NCI_CGAP_Sub9 Homo sapiens cDNA clone IMAGE:3106611 3, mRNA sequence	11.97	Down	7.10E-05
35	M_152782	Homo sapiens Sad1 and UNC84 domain containing 1 (SUNC1 mRNA	11.76	Down	6.37E-04
	CD723798	oj26f04.y1 Human lacrimal gland, unamplified: oj Homo sapiens cDNA clone oj26f04 5, mRNA sequence	11.66	Down	1.59E-04
40	NM_033136	Homo sapiens fibroblast growth factor 1 (acidic) (FGF1), transcript variant 2, mRNA	11.19	Down	5.11E-03
	BU688263	UI-CF-EC1-aea-g-11-0-UI.s1 UI-CF-EC1 Homo sapiens cDNA clone UI-CF-EC1-aea-g-11-0-UI 3, mRNA sequence	11.07	Down	2.89E-04
	NM_014391	Homo sapiens ankyrin repeat domain 1 (cardiac muscle) (ANKRD1), mRNA	11.05	Down	9.03E-03
45	NM_023915	Homo sapiens G protein-coupled receptor 87 (GPR87), mRNA	10.71	Down	4.91E-05
	NM_025239	Homo sapiens programmed cell death 1 ligand 2 (PDCD1 LG2), mRNA	10.6	Down	1.03E-03
50	NM_003392	Homo sapiens wingless-type MMTV integration site family, member 5A (WNT5A), mRNA	10.52	Down	1.11E-02
	NM_030899	Homo sapiens zinc finger protein 323 (ZNF323), mRNA	10.28	Down	1.90E-03
55	NM_001554	Homo sapiens cysteine-rich, angiogenic inducer, 61 (CYR61), mRNA	10.27	Down	5.12E-03
	AB002333	Human mRNA for KIAA0335 gene, partial cds	10.22	Down	4.90E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_005233	Homo sapiens EPH receptor A3 (EPHA3), transcript variant 1, mRNA	10.14	Down	2.91E-03
	NM_024600	Homo sapiens chromosome 16 open reading frame 30 (C16orf30), mRNA	10.11	Down	3.10E-03
10	NM_032638	Homo sapiens GATA binding protein 2 (GATA2), mRNA	9.99	Down	2.21E-04
	U79271	Human clones 23920 and 23921 mRNA sequence	9.88	Down	5.41E-05
	NM_017448	Homo sapiens lactate dehydrogenase C (LDHC), transcript variant 2, mRNA	9.74	Down	2.07E-04
15	NM_001299	Homo sapiens calponin 1, basic, smooth muscle (CNN1), mRNA	9.63	Down	3.67E-04
	A1939462	tf23h06.x5 NCI_CGAP_Brn23 Homo sapiens cDNA clone IMAGE:2097083 3, mRNA sequence	9.62	Down	8.51E-05
20	NM_178E34	Homo sapiens layilin (LOC143903), mRNA	9.59	Down	1.13E-05
	AF26908 8	Homo sapiens breast cancer antigen NY-BR-1.1 mRNA, partial cds	9.42	Down	2.10E-03
	NM_001945	Homo sapiens diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth factor) (DTR), mRNA	9.37	Down	2.21E-02
25	NM_004791	Homo sapiens integrin, beta-like 1 (with EGF-like repeat domains) (ITGBL1), mRNA	9.24	Down	2.86E-04
	NM_000682	Homo sapiens adrenergic, alpha-2B-, receptor (ADRA2B), mRNA	9.19	Down	9.68E-04
30	NM_012242	Homo sapiens dickkopf homolog 1 (Xenopus laevis) (DKK1), mRNA	9.18	Down	2.05E-03
	NM_001998	Homo sapiens fibulin 2 (FBLN2), transcript variant 2, mRNA	9.07	Down	5.02E-04
	BG16574 5	602344592F1 NIH_MGC_89 Homo sapiens cDNA clone IMAGE:4454470 5, mRNA sequence	9.04	Down	1.30E-03
35	D29453	HUMNK566 Human epidermal keratinocyte Homo sapiens cDNA clone 566, mRNA sequence	8.93	Down	2.86E-04
		oj16h02.y1 Human lacrimal gland, unamplified: oj Homo sapiens cDNA clone oj16h02 5, mRNA sequence	8.93	Down	9.36E-05
40	CD72300 6				
	NM_178550	Homo sapiens hypothetical protein MGC48998 (MGC48998), mRNA	8.87	Down	3.76E-04
	NM_016201	Homo sapiens angiomin like 2 (AMOTL2), mRNA	8.62	Down	2.48E-02
45	AW2917 75	UI-H-BI2-agv-h-04-0-UI.s1 NCI_CGAP_Sub4 Homo sapiens cDNA clone IMAGE:2725855 3, mRNA sequence	8.5	Down	3.03E-03
	NM_152703	Homo sapiens chromosome 7 open reading frame 6 (C7orf6), mRNA	8.38	Down	1.58E-04
50	NM_001843	Homo sapiens contactin 1 (CNTN1), transcript variant 1, mRNA	8.32	Down	5.41E-05
	AL11745 4	Homo sapiens mRNA; cDNA DKFZp586J1717 (from clone DKFZp586J1717)	8.26	Down	5.79E-05
	BX09200 4	BX092004NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGp998B195924 ; IMAGE:2385330, mRNA sequence	8.18	Down	4.67E-04
55	NM_145201	Homo sapiens similar to CG3714 gene product (PP3856), mRNA	8.16	Down	1.64E-03

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	BX53769 8	Homo sapiens mRNA; cDNA DKFZp686F09166 (from clone DKFZp686F09166)	8.15	Down	7.27E-04
	NM_2034	Homo sapiens Down syndrome critical region gene 1 (DSCR1), transcript variant	8.12	Down	2.96E-02
10	18	3, mRNA			
	NM_0244 23	Homo sapiens desmocollin 3 (DSC3), transcript variant Dsc3b, mRNA	7.93	Down	2.54E-03
	AK09352 9	Homo sapiens cDNA FLJ36210 fis, clone THYMU2000155	7.9	Down	9.50E-04
	AK13153 2	Homo sapiens cDNA FLJ16761 fis, clone BRAMY3008096	7.89	Down	1.84E-03
15	NM_0325 11	Homo sapiens chromosome 6 open reading frame 168 (C6orf168), mRNA	7.42	Down	3.49E-03
	NM_0025 21	Homo sapiens natriuretic peptide precursor B (NPPB), mRNA	7.4	Down	4.45E-03
20	BU62632 6	UI-H-DF0-bet-p-17-0-UI.s1 NCI_CGAP_DF0 Homo sapiens cDNA clone UI-H-DF0-bet-p-17-0-UI 3, mRNA sequence	7.34	Down	7.03E-05
	NM_0011 44	Homo sapiens autocrine motility factor receptor (AMFR), transcript variant 1, mRNA	7.33	Down	1.03E-03
25	NM_0015 41	Homo sapiens heat shock 27kDa protein 2 (HSPB2), mRNA	7.32	Down	3.96E-03
	NM_0050 69	Homo sapiens single-minded homolog 2 (Drosophila) (SIM2), transcript variant SIM2, mRNA	7.31	Down	1.23E-03
	A1937359	wp76c02.x1 NCI_CGAP_Brn25 Homo sapiens cDNA clone IMAGE:2467682 3, mRNA sequence	7.31	Down	1.96E-03
30	NM_0153 45	Homo sapiens dishevelled associated activator of morphogenesis 2 (DAAM2), mRNA	7.29	Down	2.18E-05
	NM_0000 89	Homo sapiens collagen, type I, alpha 2 (COL1A2), mRNA	7.25	Down	6.07E-03
35	NM_0151 50	Homo sapiens raft-linking protein (RAFTLIN), mRNA	7.24	Down	2.87E-03
	BX11331 9	BX113319 NCI_CGAP_Gas4 Homo sapiens cDNA clone IMAGp998G205398 ; IMAGE:2184619, mRNA sequence	7.24	Down	1.28E-03
40	M_1993 29	Homo sapiens solute carrier family 43, member 3 (SLC43A3), mRNA	7.19	Down	2.31E-03
	AV70297 7	AV702977 ADB Homo sapiens cDNA clone ADBCVD08 5, mRNA sequence	7.11	Down	5.21E-05
45	NM_0245 12	Homo sapiens leucine rich repeat containing 2 (LRRC2), mRNA	7.09	Down	2.65E-04
	NM_0176 55	Homo sapiens PDZ domain protein GIPC2 (GIPC2), mRNA	7.06	Down	2.04E-04
	NM_0144 77	Homo sapiens chromosome 20 open reading frame 10 (C20orf10), mRNA	7.05	Down	4.10E-03
50	NM_0025 59	Homo sapiens purinergic receptor P2X, ligand-gated ion channel, 3 (P2RX3), mRNA	7.03	Down	4.63E-03
	BM97638 5	UI-CF-EN1-acz-f-03-0-UI.s1 UI-CF-EN1 Homo sapiens cDNA clone UI-CF-EN1-acz-f-03-0-UI 3, mRNA sequence	7.01	Down	9.77E-04
55	BX64832 3	Homo sapiens mRNA; cDNA DKFZp686K10163 (from clone DKFZp686K10163)	6.94	Down	8.05E-04
	BC00858 0	Homo sapiens, clone IMAGE:4179986, mRNA, partial cds	6.93	Down	5.41E-05

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_0070 34	Homo sapiens DnaJ (Hsp40) homolog, subfamily B, member 4 (DNAJB4), mRNA	6.72	Down	2.98E-02
	AW0255 56	wu97g10.x1 NCI_CGAP_Kid3 Homo sapiens cDNA clone IMAGE:2528034 3, mRNA sequence	6.7	Down	2.31E-03
10	NM_0035 14	Homo sapiens histone 1, H2am (HIST1H2AM), mRNA	6.7	Down	9.11E-03
	NM_0326 03	Homo sapiens lysyl oxidase-like 3 (LOXL3), mRNA	6.65	Down	2.58E-03
	AW1726 52	xj80f06.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2663555 3, mRNA sequence	6.62	Down	2.17E-04
15	NM_0063 50	Homo sapiens follistatin (FST), transcript variant FST317, mRNA	6.57	Down	7.59E-05
	BQ01858 6	UI-H-DH1-awu-c-08-0-UI.s1 NCI_CGAP_DH1 Homo sapiens cDNA clone IMAGE:5823679 3, mRNA sequence	6.43	Down	3.44E-03
20		nae30h12.x1 Lupski_sympathetic_trunk	6.43	Down	8.65E-05
	BF44815 8	Homo sapiens cDNA clone IMAGE:4090607 3, mRNA sequence			
	NM_0015 57	Homo sapiens interleukin 8 receptor, beta (IL8RB), mRNA	6.38	Down	2.96E-03
25	NM_0021 66	Homo sapiens inhibitor of DNA binding 2, dominant negative helix-loop-helix protein (ID2), mRNA	6.37	Down	2.10E-03
	NM_0806 71	Homo sapiens potassium voltage-gated channel, Isk-related family, member 4 (KCNE4), mRNA	6.36	Down	1.17E-02
30	BM71294 5	UI-E-EJ0-ahi-c-16-0-UI.r1 UI-E-EJ0 Homo sapiens cDNA clone UI-E-EJ0-ahi-c-16-0-UI 5, mRNA sequence	6.33	Down	3.62E-04
	BM99189 0	UI-H-DF1-auk-h-02-0-UI.s1 NCI_CGAP_DF1 Homo sapiens cDNA clone IMAGE:5870641 3, mRNA sequence	6.32	Down	8.14E-05
	NM_0178 23	Homo sapiens dual specificity phosphatase 23 (DUSP23), mRNA	6.26	Down	9.36E-03
35	NM_0324 61	Homo sapiens SPANX family, member B1 (SPANXB1), mRNA	6.26	Down	5.62E-03
	NM_00601 27	Homo sapiens exonuclease 1 (EXO1), transcript variant 1, mRNA	6.26	Down	7.81E-03
40	NM_0035 45	Homo sapiens histone 1, H4e (HIST1 H4E), mRNA	6.19	Down	7.48E-03
	NM_0014 51	Homo sapiens forkhead box F1 (FOXF1), mRNA	6.17	Down	1.02E-03
45	NM_001 55	Homo sapiens endothelin 1 (EDN1), mRNA 9	6.16	Down	2.54E-03
	NM_003 17	Homo sapiens regulator of G-protein 6 signalling 5 (RGS5), mRNA	6.07	Down	9.24E-03
	NM_0211 03	Homo sapiens thymosin, beta 10 (TMSB10), mRNA	6.06	Down	2.32E-02
50	NM_0014 24	Homo sapiens epithelial membrane protein 2 (EMP2), mRNA	6.06	Down	2.23E-03
	BG35457 9	CDCA6 Cell division cycle associated gene 6 Human cDNA expression libraries Homo sapiens cDNA clone 407614, mRNA sequence	5.96	Down	1.40E-03
55	BM66545 2	UI-E-CQ1-aex-n-03-0-UI.s1 UI-E-CQ1 Homo sapiens cDNA clone UI-E-CQ1-aex-n-03-0-UI 3, mRNA sequence	5.94	Down	2.64E-04

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_079423	Homo sapiens myosin, light polypeptide 6, alkali, smooth muscle and non-muscle (MYL6), transcript variant 2, mRNA	5.87	Down	3.58E-02
	AW1344 73	UI-H-BI1-abv-a-11-0-UI.s1 NCI_CGAP_Sub3 Homo sapiens cDNA clone IMAGE:2712885 3, mRNA sequence	5.85	Down	1.03E-03
10	NM_032334	Homo sapiens hypothetical protein MGC14595 (MGC14595), mRNA	5.79	Down	9.11E-03
	BF02935	601765592F1 NIH_MGC_53 Homo sapiens cDNA clone IMAGE:3997510 5, mRNA	5.78	Down	3.55E-03
	6	sequence			
15	NM_175887	Homo sapiens hypothetical protein LOC222171 (LOC222171), mRNA	5.72	Down	3.51E-04
	NM_139241	Homo sapiens FYVE, RhoGEF and PH domain containing 4 (FGD4), mRNA	5.72	Down	3.12E-03
20	NM_015915	Homo sapiens spastic paraplegia 3A (autosomal dominant) (SPG3A), mRNA	5.7	Down	2.15E-03
	NM_052937	Homo sapiens similar to hypothetical protein FLJ10883 (LOC115294), mRNA	5.69	Down	7.86E-03
	NM_015251	Homo sapiens KIAA0431 protein (KIAA0431), mRNA	5.6	Down	3.87E-04
25	NM_021956	Homo sapiens glutamate receptor, ionotropic, kainate 2 (GRIK2), transcript variant 1, mRNA	5.58	Down	2.31E-03
	NM_017527	Homo sapiens lymphocyte antigen 6 complex, locus K (LY6K), mRNA	5.57	Down	1.69E-02
30	NM_015192	Homo sapiens phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 1, mRNA	5.53	Down	7.81E-03
	NM_020223	Homo sapiens family with sequence similarity 20, member C (FAM20C), mRNA	5.51	Down	1.19E-03
	NM_002619	Homo sapiens platelet factor 4 (chemokine (C-X-C motif) ligand 4) (PF4), mRNA	5.51	Down	3.31E-04
35	NM_002053	Homo sapiens guanylate binding protein 1, interferon-inducible, 67kDa (GBP1), mRNA	5.51	Down	1.27E-03
	H55853	yq94c05.r1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:203432 5, mRNA sequence	5.5	Down	6.78E-03
40	NM_000947	Homo sapiens primase, polypeptide 2A, 58kDa (PRIM2A), mRNA	5.48	Down	8.13E-03
	NM_005596	Homo sapiens nuclear factor I/B (NFIB), mRNA	5.46	Down	1.00E-04
45	NM_014285	Homo sapiens exosome component 2 (EXOSC2), mRNA	5.45	Down	1.45E-02
	NM_004623	Homo sapiens tetratricopeptide repeat domain 4 (TTC4), mRNA	5.43	Down	1.84E-02
	NM_052846	Homo sapiens elastin microfibril interfacer 3 (EMILIN3), mRNA	5.43	Down	1.11 E-04
50	AW1523 68	xg63e03.x1 NCI_CGAP_Ut4 Homo sapiens cDNA clone IMAGE:2633020 3 similar to contains Alu repetitive element;, mRNA sequence	5.41	Down	5.48E-04
	NM_000475	Homo sapiens nuclear receptor subfamily 0, group B, member 1 (NR0B1 mRNA	5.39	Down	1.85E-03
55	NM_012394	Homo sapiens prefoldin 2 (PFDN2), mRNA	5.38	Down	3.93E-02

EP 1 780 267 B1

(continued)

	Gene Identifier	Gene Name	Average fold change AF-II vs AF-III	Direction	adj. p-value
5	NM_019018	Homo sapiens hypothetical protein FLJ11127 (FLJ11127), mRNA	5.32	Down	1.05E-02
	NM_004298	Homo sapiens nucleoporin 155kDa (NUP155), transcript variant 2, mRNA	5.3	Down	5.08E-02
10	M_003516	Homo sapiens histone 2, H2aa (HIST2H2AA), mRNA	5.3	Down	3.90E-02
	NM_004324	Homo sapiens BCL2-associated X protein (BAX), transcript variant beta, mRNA	5.27	Down	2.42E-02
	NM_006012	Homo sapiens ClpP caseinolytic protease, ATP-dependent, proteolytic subunit homolog (E. coli) (CLPP), nuclear gene encoding mitochondrial protein, mRNA	5.22	Down	3.42E-03
15	NM_021968	Homo sapiens histone 1, H4j (HIST1H4J), mRNA	5.19	Down	1.84E-02
	BF111710	7147c10.x1 Soares_NSF_F8_9W_OT_PA_P_S1 Homo sapiens cDNA clone IMAGE:3524371 3, mRNA sequence	5.19	Down	1.23E-03
20	NM_006837	Homo sapiens COP9 constitutive photomorphogenic homolog subunit 5 (Arabidopsis) (COPS5), mRNA	5.16	Down	3.83E-02
	NM_001884	Homo sapiens hyaluronan and proteoglycan link protein 1 (HAPLN1), mRNA	5.15	Down	6.63E-03
	NM_018663	Homo sapiens peroxisomal membrane protein 2, 22kDa (PXMP2), mRNA	5.14	Down	1.79E-02
25	BX093329	BX093329 Soares_parathyroid_tumor_NbHPA Homo sapiens cDNA clone IMAGp998A124183 ; IMAGE:1648403, mRNA sequence	5.14	Down	9.07E-03
30	NM_00101992	Homo sapiens ubiquitin specific protease 16 (USP16), transcript variant 2, mRNA	5.13	Down	1.18E-02
	NM_003405	Homo sapiens tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, eta polypeptide (YWHAH), mRNA	5.11	Down	1.09E-02
35	NM_003238	Homo sapiens transforming growth factor, beta 2 (TGFB2), mRNA	5.09	Down	2.36E-03
	T95596	ye40b03.s1 Soares fetal liver spleen 1 NFLS Homo sapiens cDNA clone IMAGE:120173 3, mRNA sequence	5.08	Down	3.51E-04
40	NM_004753	Homo sapiens dehydrogenase/reductase (SDR family) member 3 (DHRS3), mRNA	5.07	Down	6.56E-03
	NM_000165	Homo sapiens gap junction protein, alpha 1, 43kDa (connexin 43) (GJA1), mRNA	5.04	Down	2.07E-03
	H99504	yx25g05.s1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:262808 3, mRNA sequence	5.04	Down	1.13E-03
45	NM_012464	Homo sapiens tolloid-like 1 (TLL1), mRNA	5.03	Down	3.47E-02
	NM_052947	Homo sapiens heart alpha-kinase (HAK), mRNA	5.02	Down	1.06E-04

55

TABLE VIII: REAL TIME PCR DATA FOR DIFFERENTIATED AF-I CELLS

Hormone	Ct values for AFCA007 Clone A cultured for 2 wks in Amniomax	Ct values for AFCA007 Clone A cultured for 2 wks in Amniomax + 10 micro molar retinoic acid	$\Delta\Delta C_t$ for AFCA007 Clone A cultured for 2 wks in Amniomax	$\Delta\Delta C_t$ for AFCA007 Clone A cultured for 2 wks in Amniomax + 10 micro molar retinoic acid
GIP	34	31	0.00062	0.004
Neurotensin	37	35	0.000183	0.0004
Gastrin	No expression	36		0.0004
Secretin	32	32	0.02	0.014
Somatostatin	31	28	0.002	0.014
CCK	33	31	0.003	0.008
GAPDH	22	22		
*: For $\Delta\Delta C_t$ analysis, the positive control was human intestinal RNA				

TABLE IX: PCR ANALYSIS OF THE EXPRESSION OF ENDODERM MARKERS IN AFCA007 (P8) AFTER 14-DAYS OF DIFFERENTIATION ASSAY

Treatment	HNF3 β	Pdx-1	GATA4	Glu-2	HNF1 α	AFP
Untreated	-	-	-	+	-	-
FGF4 50 ng/ml	+	+	+	+	+	
All-trans RA (10 μ M)	+	-	+	+	+	+
L-685,458 (10 μ M)	+		+	+	+	
Cyclopamine (10 μ M)			+	+		
BMP5 (5 ng/ml)	+		+	+	+	
BMP6 (50 ng/ml)	+		+	+	+	
Excendin 4 (10 nM)	+		+	+	+	
BMP7 (10 ng/ml)			+	+	+	
BMP4 (50 ng/ml)	+			+	+	

TABLE X: CYTOKINE, CYTOKINE AND GROWTH FACTOR RECEPTOR EXPRESSION LEVEL FOR AFCA007 A (AF-I) AND AFCA015 C (AF-II) CELLS

	AFCA007 A	AFCA007 A	AFCA015 C	AFCA015 C
POS	23,029.50	22,843.43	23,281.28	23,086.51
NEG	1.00	0.98	0.86	0.86
Angiogenin	44.50	89.98	101.17	26.71
BDNF	702.50	670.09	590.28	411.52
BLC	276.50	365.08	192.90	0.86
BMP-4	66.00	96.36	126.89	117.22
BMP-6	135.00	182.17	247.34	133.53
CK beta 8-1	139.50	100.28	80.16	43.87
CNTF	235.50	234.64	82.73	0.86

EP 1 780 267 B1

(continued)

		AFCA007 A	AFCA007 A	AFCA015 C	AFCA015 C
	EGF	125.50	106.66	66.01	0.86
5	Eotaxin	90.00	95.87	152.61	99.21
	Eotaxin-2	635.00	554.86	528.55	290.54
	Eotaxin-3	174.50	245.43	246.48	170.42
	FGF-6	422.00	411.67	318.07	213.32
	FGF-7	378.50	348.41	225.05	15.12
10	Flt-3 Ligand	298.00	264.56	146.18	66.60
	Fractalkine	545.50	399.41	330.50	105.21
	GCP-2	323.00	266.52	182.18	0.86
	GDNF	249.00	213.56	100.74	15.55
15	GM-CSF	1,003.50	687.75	663.15	343.31
	I-309	108.00	90.47	91.74	36.57
	IFN-gamma	227.50	189.04	217.34	7.40
	IGFBP-1	156.00	197.37	182.61	37.00
	IGFBP-2	627.00	415.10	383.66	281.53
20	IGFBP-4	154.00	136.57	103.74	0.86
	IGF-I	121.00	200.32	218.62	191.01
	IL-10	474.50	379.30	320.22	64.89
	IL-13	768.00	627.43	636.57	184.58
25	IL-15	829.50	542.60	558.56	226.62
	IL-16	243.50	235.62	146.18	2.25
	IL-1alpha	1,206.00	1,058.47	1,048.95	662.05
	IL-1beta	349.00	282.21	258.49	64.89
	IL-1ra	278.50	337.13	225.05	111.65
30	IL-2	684.50	581.34	489.97	254.50
	IL-3	1,052.50	1,066.31	1,075.53	904.86
	IL-4	396.00	404.80	383.23	224.47
	IL-5	763.00	676.96	754.89	303.41
35	IL-6	1,172.00	1,113.88	485.25	327.43
	IL-7	675.00	567.61	473.68	241.63
	Leptin	269.50	267.99	229.77	118.08
	LIGHT	295.50	254.75	214.33	85.48
	MCP-1	1,027.00	901.06	1,080.25	990.24
40	MCP-2	546.00	461.20	389.66	120.66
	MCP-3	378.50	372.93	288.49	85.91
	MCP-4	642.50	532.30	390.95	159.69
	M-CSF	692.00	566.63	504.11	267.37
45	MDC	446.50	442.07	312.07	138.24
	MIG	760.00	725.51	701.30	363.47
	MIP-1-delta	290.50	347.92	295.78	117.65
	MIP-3-alpha	406.50	399.90	311.64	64.46
	NAP-2	216.50	241.51	208.33	44.29
50	NT-3	45.50	124.31	61.30	0.86
	PARC	181.50	208.16	138.46	0.86
	PDGF-BB	594.00	313.10	126.03	152.83
	RANTES	418.50	350.86	248.63	108.21
55	SCF	307.50	152.26	84.45	45.58
	SDF-1	654.50	594.09	237.48	116.37
	TARC	227.50	264.56	148.75	2.25
	TGF-beta 1	824.50	750.02	693.59	407.23

EP 1 780 267 B1

(continued)

		AFCA007 A	AFCA007 A	AFCA015 C	AFCA015 C
5	TGF-beta 3	278.00	341.54	277.35	57.16
	TNF-alpha	804.50	968.73	992.37	552.23
	TNF-beta	549.50	510.23	888.20	285.82
10	POS	24,267.79	23,873.40	25,002.74	25,119.43
	NEG	1.00	0.89	0.78	0.78
	Acrp30	29.13	0.89	100.74	98.12
15	AgRP	1.00	0.89	19.30	0.78
	Angiopoietin-2	144.13	84.27	62.74	58.01
	Amphiregulin	1.00	0.89	0.78	0.78
20	Axl	1,478.63	882.74	542.12	536.93
	bFGF	5,455.13	4,386.44	4,419.49	7,225.00
	b-NGF	94.13	8.27	69.72	58.79
25	BTC	1.00	0.89	0.78	0.78
	CCL-28	141.13	76.23	0.78	0.78
	CTACK	383.13	315.86	304.37	297.86
30	Dtk	90.13	51.19	48.77	48.67
	EGF-R	1,730.13	851.89	1,363.21	1,485.42
	ENA-78	207.13	109.31	514.97	192.34
35	Fas/TNFRSF6	4,740.63	1,570.79	1,026.94	575.87
	FGF-4	342.13	141.50	156.21	88.77
	FGF-9	267.13	166.98	140.69	100.07
40	GCSF	320.63	151.78	152.33	115.25
	GITR-Ligand	183.63	112.44	170.17	81.77
	GITR	211.13	116.91	63.51	18.30
45	GRO	10,097.13	9,584.54	2,380.16	1,317.60
	GRO-alpha	1,061.13	754.43	356.73	124.60
	HCC-4	90.63	18.55	67.00	8.18
50	HGF	1.00	0.89	265.19	333.29
	ICAM-1	11,257.63	6,308.40	2,116.03	1,415.33
	ICAM-3	75.13	19.89	54.98	61.52
55	IGFBP-3	603.63	406.61	435.46	346.92
	IGFBP-6	523.63	334.63	253.56	176.77
	IGF-I SR	929.13	526.87	484.72	463.73
60	IL-1 R4/ST2	330.63	297.97	272.95	171.71
	IL-1 RI	57.13	40.91	40.24	16.74
	IL-11	192.13	106.63	79.80	19.86
65	IL-12 p40	134.13	113.33	99.19	93.06
	IL-12 p70	183.63	165.64	166.29	85.27
	IL-17	47.13	0.89	0.78	0.78
70	IL-2 Rapha	229.13	158.04	115.87	36.21
	IL-6 R	158.63	30.62	55.75	46.72
	IL-8	2,792.13	2,894.57	528.93	164.70
75	I-TAC	499.63	184.42	139.53	89.16
	Lymphotactin	581.63	277.86	295.45	215.71
	MIF	2,280.63	1,794.32	1,532.31	1,042.71
80	MIP-1alpha	557.13	448.19	319.49	261.65
	MIP-1beta	193.13	133.45	182.58	225.44
	MIP-3beta	360.13	203.19	126.34	101.23
85	MSP-alpha	251.13	174.13	121.69	89.94
	NT-4	115.13	27.94	2.62	0.78

EP 1 780 267 B1

(continued)

		AFCA007 A	AFCA007 A	AFCA015 C	AFCA015 C
	Osteoprotegerin	2,391.13	1,761.24	1,558.30	1,582.37
5	Oncostatin M	541.13	340.45	179.48	98.51
	PIGF	88.13	50.74	40.24	11.29
	sgp130	604.13	503.63	369.92	379.24
	sTNF RII	17.63	0.89	0.78	8.57
	sTNF-RI	903.63	551.91	685.63	469.18
10	TECK	255.63	91.87	140.69	100.46
	TIMP-1	1,099.63	732.53	661.58	832.85
	TIMP-2	11,468.63	7,205.68	14,022.74	14,546.57
	Thrombopoietin	580.13	364.59	405.21	281.90
15	TRAIL R3	1,881.13	728.50	988.93	1,186.39
	TRAIL R4	437.13	412.42	172.89	233.23
	uPAR	1,055.13	631.49	668.56	960.95
	VEGF	418.13	293.95	290.79	452.44
	VEGF-D	39.13	0.89	20.85	12.46
20	POS	17,605.50	17,898.76	18,624.45	19,358.48
	NEG	1.00	1.04	1.02	1.09
	Activin A	7.00	106.51	131.02	216.38
	ALCAM	227.00	372.00	229.19	194.01
25	B7-1(CD80)	18.00	138.72	100.34	3.55
	BMP-5	42.50	129.37	90.63	42.29
	BMP-7	72.00	136.12	111.59	1.09
	Cardiotrophin-1	578.00	1,221.99	498.65	233.30
	CD14	118.00	231.72	134.09	180.37
30	CXCL-16	1.00	187.56	201.07	175.45
	DR6 (TNFRSF21)	107.00	382.39	190.84	1.09
	Endoglin	1,213.50	735.17	1,168.45	1,359.16
	ErbB3	183.50	269.13	226.63	167.81
35	E-Selectin	150.00	250.94	178.06	82.68
	Fas Ligand	134.00	269.65	162.72	88.14
	ICAM-2	364.50	416.16	273.16	203.83
	IGF-II	172.00	206.26	140.22	89.77
	IL-1 R II	178.00	226.52	190.33	124.70
40	IL-10 Rbeta	124.50	111.18	91.14	194.56
	IL-13 Ralpha2	94.50	117.94	80.40	1.09
	IL-18 BPalpha	3.00	132.49	234.82	1.09
	IL-18 Rbeta	59.50	191.71	112.61	1.09
45	IL-2 Ralpha	128.00	156.90	149.94	50.48
	IL-2 Rbeta	1.00	57.15	196.98	1.09
	IL-2 Rgamma	1.00	1.04	1.02	1.09
	IL-21R	153.50	166.78	146.36	70.67
	IL-5 Ralpha	133.00	166.78	150.96	21.01
50	IL-9	169.00	184.96	155.56	57.58
	IP-10	91.50	211.46	161.19	94.14
	LAP	3,066.00	2,460.08	5,775.28	6,813.80
	Leptin R	225.50	223.41	168.35	230.03
55	LIF	93.50	181.84	495.07	217.48
	L-Selectin	87.50	103.91	122.84	1.09
	M-CSF R	1.00	193.27	65.57	167.27
	MMP-1	206.50	170.93	135.11	379.56

(continued)

		AFCA007 A	AFCA007 A	AFCA015 C	AFCA015 C
	MMP-13	158.50	149.11	119.26	57.58
5	MMP-9	1.00	137.16	56.37	167.81
	MPIF-1	1.00	291.47	1.02	1.09
	NGF R	1.00	123.13	388.21	1.09
	PDGF AA	331.50	311.21	470.01	493.62
	PDGF-AB	48.50	111.70	137.67	248.58
10	PDGF Ralpha	1.00	184.96	174.48	1.09
	PDGF Rbeta	1,199.50	859.86	3,513.79	4,030.00
	PECAM-1	41.50	138.72	280.32	190.74
	Prolactin	285.00	272.77	195.44	59.76
15	SCFR	64.50	245.75	128.46	88.68
	SDF-1beta	425.50	364.73	248.11	452.14
	Siglec-5	152.50	153.79	1.02	1.09
	TGF-alpha	69.50	227.04	140.74	166.72
	TGF beta2	173.50	138.72	182.15	118.70
20	Tie-1	8.50	124.69	373.89	101.78
	Tie-2	153.00	90.92	231.24	18.28
	TIMP-4	49.50	97.68	93.70	17.74
	VE-Cadherin	77.00	139.24	131.02	85.41
25	VEGF R2	59.50	164.70	101.37	78.86
	VEGF R3	37.50	121.06	222.54	54.85

TABLE XI: PROTEINS CAPABLE OF DETECTION BY ANTIBODY ARRAYS

30	Cytokine	Full Name
	Activin A	EDF(erythroid differentiation factor), FRP(Follicle stimulating hormone releasing protein), Restrictin-P, WEHI-MIF(WEHI mesoderm inducing factor)
35	AGRP	Agouti related protein
	ALCAM	Activated leukocyte cell adhesion molecule
	Amphiregulin	Amphiregulin
	ANG	Angiogenin
40	Angiopoietin-1	Angiopoietin-1
	Angiopoietin-like Factor	Angiopoietin-like Factor
	Angiostatin	Angiostatin
45	APRIL	A proliferation-inducing ligand
	AXL	A protein tyrosine kinase also called UFO or ark. The ligand for this receptor is GPF (growth-potentiating factor). The ligand is called also axl stimulatory factor, which is identical with one of the gas genes, gas-6.
50	B7-1(CD80)	B7-1(CD80)
	BAFF	B-cell-activating factor belonging to the TNF family
	BDNF	Brain-derived neurotrophic factor
	Betacellulin	Betacellulin
55	beta-Galactosidase	beta-Galactosidase
	beta-NGF	Nerve growth factor-beta

EP 1 780 267 B1

(continued)

	Cytokine	Full Name
5	bFGF	Basic fibroblast growth factor
	BLC	B-lymphocyte chemoattractant
	BMP-2	Bone morphogenetic proteins -2
	BMP-4	Bone morphogenetic proteins -4
10	BMP-5	Bone morphogenetic proteins -5
	BMP-6	Bone morphogenetic proteins -6
	BMP-7	Bone morphogenetic proteins -7
15	BTC	Betacellulin
	CCL28	CCK-1
	CD27	Cluster of Differentiation 27
	CD30	Cluster of Differentiation 30
20	CD40	Cluster of Differentiation 40
	CD40 Ligand	Cluster of Differentiation 40 ligand
	Ck beta 8-1	Chemokine-beta-8
25	CNTF	Ciliary neuronotrophic factor, ciliary neurotrophic factor
	Cripto-1	CRGF (Cripto growth factor)
	CTACK	CTAC (Cuteaneous T-Cell Attracting Chemokine); Skinkine; Eskine
	CTLA-4	Cytotoxic T-lymphocyte associated antigen 4
30	CXCL16	CXC Chemokine ligand 16
	Dkk-4	Dickkopf-4
	DR6	Death Receptor 6
35	Dtk	Developmental tyrosine kinase
	EGF	Epidermal growth factor
	EGR-R	Epidermal growth factor related protein
	ENA-78	Epithelial neutrophil-activating protein 78, epithelial cellderived neutrophil attractant-78
40	Endostatin	Endostatin
	Eotaxin	Eotaxin
	Eotaxin-2	MPIF-2 (Myeloid progenitor inhibitory factor-2),Chemokine-beta-6
45	Eotaxin-3	MIP-4-alpha (macrophage inflammatory protein-4-alpha), TSC-1 (thymic stroma chemokine-1)
	Erythropoietin	Erythropoietin
	E-Selectin	E-Selectin
50	Fas Ligand	FasL, CD95 ligand, Apo-1 ligand
	Fas/TNFRSF6	Fas or cd95
	FGF-2	Fibroblast growth factor-2
55	FGF-4	Fibroblast growth factor-4
	FGF-6	Fibroblast growth factor-6
	FGF-7	Fibroblast growth factor-7

EP 1 780 267 B1

(continued)

	Cytokine	Full Name
5	FGF-8	Fibroblast growth factor-8
	FGF-9	Fibroblast growth factor-9
	FKN or FK	Fractalkine
	Flt-3 Ligand	fms-like tyrosine kinase-3 Ligand (also known as STK-1 Ligand)
10	Follistatin	Follicle stimulating hormone suppressing protein (FSP)
	GCP-2	Granulocyte Chemotactic Protein 2
	GCSF	Granulocyte-colony Stimulating Factor
15	GDF-15	growth/differentiation factor-15
	GDNF	Glial-derived Neurotrophic Factor
	GITR	Glucocorticoid induced tumor necrosis factor receptor
	GITR-Ligand	Glucocorticoid induced tumor necrosis factor receptor
20	GM-CSF	Granulocyte-macrophage colony stimulating factor
	GRO	Growth Related Oncogene
25	GRO-a	Growth Related Oncogene-Alpha
	HB-EGF	Heparin-binding Epidermal Growth factor
	HCC-4	Hemofiltrate CC Chemokine 4
	HGF	Hepatocyte growth factor
30	HPTA	Hepatopoeitin A
	HRG	Heregulin/NDF/GGF/Neuregulin
	HRG-alpha	Heregulin/NDF/GGF/Neuregulin
35	HRG-beta	Heregulin/NDF/GGF/Neuregulin
	HVEM	Herpesvirus entry mediator
	I-309	I-309
	ICAM-1	Intercellular Adhesion Molecule 1
40	ICAM-2	Intercellular Adhesion Molecule 2
	ICAM-3	Intercellular Adhesion Molecule 3
	IFN-alpha	Interferon alpha
45	IFN-beta	Interferon beta
	IFN-gamma	Interferon gamma
	IFN-omega	Interferon omega
50	IGFBP-1	Insulin-like growth factor binding proteins 1
	IGFBP-2	Insulin-like growth factor binding proteins 2
	IGFBP-3	Insulin-like growth factor binding proteins 3
	IGFBP-4	Insulin-like growth factor binding proteins 4
55	IGFBP-6	Insulin-like growth factor binding proteins 6
	IGF-I SR	Insulin like growth factor I soluble receptor
	IGF-I	Insulin-like growth factor-1

EP 1 780 267 B1

(continued)

	Cytokine	Full Name
5	IGF-II	Insulin-like growth factor II
	IL-1 R1	Interleukin 1 receptor-related protein-1
	IL-1 R4/ST2	Interleukin 1 receptor 4, also known as ST2
	IL-10	Interleukin 10
10	IL-10 R alpha	Interleukin 10 soluble receptor alpha
	IL-10 R beta	Interleukin 10 receptor beta
	IL-11	Interleukin 11
15	IL-12	Interleukin 12
	IL-12 p40	Interleukin 12p40
	IL12p70	Interleukin 12p70
	IL-13	Interleukin 13
20	IL-15	Interleukin 15
	IL-16	Interleukin 16
	IL-17	Interleukin 17
25	IL-18 BPα	Interleukin 18 binding protein A
	IL-18 R alpha	Interleukin 18 receptor alpha
	IL-18 R beta	Interleukin 18 receptor beta
	IL-18	Interleukin 18
30	IL-1α	Interleukin I alpha
	IL-1β	Interleukin I beta
	IL-1ra	Interleukin 1 receptor antagonist
35	IL-2	Interleukin 2
	IL-2 R alpha	Interleukin 2 soluble receptor alpha
	IL-2 R beta	Interleukin 2 soluble receptor beta
	IL-2 R gamma	Interleukin 2 receptor gamma
40	IL-21 R	Interleukin 21 receptor
	IL-3	Interleukin 3
	IL-4	Interleukin 4
45	IL-5	Interleukin 5
	IL-5 R alpha	Interleukin 5 receptor alpha
	IL-6	Interleukin 6
	IL-7	Interleukin 7
50	IL-8	Interleukin 8
	IL-9	Interleukin 9
	IL-9 R	Interleukin 9 receptor
55	Inhibin A	Inhibin A
	Inhibin B	Inhibin B
	IP-10	IFN - inducible Protein 10, Immune Protein 10

EP 1 780 267 B1

(continued)

	Cytokine	Full Name
5	I-TAC	Interferon-inducible T cell Alpha Chemoattractant
	LAP	Liver activator protein
	TGF-LAP	Tumor growth factor - latency associated peptide
	LT-BP-1	Latent TGF-beta binding protein 1
10	LECT2	Leukocyte cell-derived chemotaxin-2
	LEPTIN R	LEPTIN receptor
	LFA-1 alpha	Lymphocyte function-associated antigen 1 alpha
15	LIF	Leukemia Inhibitory Factor
	LIGHT	LIGHT
	LIX	LPS induced C-X-C chemokine
	L-Selectin	L-Selectin
20	Lymphotactin	Lymphotactin
	MAC-1	Macrophage galactose-specific lectin-1
	MMP-13	Matrix Metalloproteinase 13
25	MMP-19	Matrix Metalloproteinase 19
	MCP-1	Monocyte Chemoattractant Protein 1
	MCP-2	Monocyte Chemoattractant Protein 2
	MCP-3	Monocyte Chemoattractant Protein 3
30	MCP-4	Monocyte Chemoattractant Protein 4
	MCSF	Macrophage-colony Stimulating Factor
	M-CSF R	Macrophage colony stimulating factor receptor
35	MDC	Macrophage-derived Chemokine
	MIF	Macrophage Migration Inhibitory Factor
	MIG	Monokine induced by gamma interferon
	MIP-1a	Macrophage Inflammatory Protein 1 alpha
40	MIP-1b	Macrophage Inflammatory Protein 1 beta
	MIP-1d	Macrophage Inflammatory Protein 1delta
	MIP-2	Macrophage Inflammatory Protein 2
45	MIP-3 alpha	Macrophage Inflammatory Protein 3 alpha
	MIP-3 beta	Macrophage Inflammatory Protein 3 beta
	MMP-1	Matrix Metalloproteinase 1
	MMP-10	Matrix Metalloproteinase 10
50	MMP-11 (Stromelysin-3)	Matrix Metalloproteinase 11
	MMP-12	Matrix Metalloproteinase 12
	MMP-13	Matrix Metalloproteinase 13
55	MMP-14	Matrix Metalloproteinase 14
	MMP-2	Matrix Metalloproteinase 2
	MMP-20	Matrix Metalloproteinase 20

EP 1 780 267 B1

(continued)

	Cytokine	Full Name
5	MMP-3	Matrix metalloproteinase 3
	MMP-7	Matrix Metalloproteinase 7
	MMP-8	Matrix Metalloproteinase 8
	MMP-9	Matrix Metalloproteinase 9
10	MPIF-1	Myeloid Progenitor Inhibitory Factor 1
	MSP a-chain	Macrophage stimulating protein a-chain
	MSP b-chain	Macrophage stimulating protein b-chain
15	NAP-2	Neutrophil Activating Peptide 2
	NGF	Nerve growth factor
	NGF R	Nerve growth factor receptor
	NT-3	Neurotrophin-4
20	NT-4	Neurotrophin-3
	OPG	Osteoprotegerin
	OSM	Oncostatin M
25	PARC	Pulmonary and Activation-Regulated Chemokine
	PDGF R alpha	Platelet-derived growth factor receptor alpha
	PDGF R beta	Platelet-derived growth factor receptor beta
30	PDGF-AA	Platelet-derived Growth Factor AA
	PDGF-BB	Platelet-derived Growth Factor BB
	PECAM-1	Platelet endothelial cell adhesion molecule
35	PF4	Platelet factor 4
	PIGF	Placenta growth Factor
	Prolactin	Prolactin
	P-selectin	P-selectin
40	RANTES	Regulated upon activation, normal T-cell expressed, and presumably secreted
	SCF	Stem Cell Factor
	SCF R	Stem cell factor receptor
45	SDF-1	Stromal cell-derived factor
	SLPI	Secretory leukocyte protease inhibitor
	S MDF	Sensory and Motor Neuron-derived Factor
	SPARC	Osteonectin
50	ST2	Interleukin 1 receptor 4
	TARC	Thymus and Activation-Regulated Chemokine
	TECK	Thymus-expressed Chemokine
55	TGFa	Tumor Necrosis Factor Alpha
	TGF-a	Transforming growth factor alpha
	TGF-beta 1	Tumor Necrosis Facyor beta1

(continued)

Cytokine	Full Name
TGF-beta 2	Tumor Necrosis Facyor beta2
TGF-beta 3	Tumor Necrosis Facyor beta3
Thrombospondin (TSP)	Thrombospondin
Thymopoietin	Thymopoietin (Tpo)
Tie-1	Tyrosine kinase with Ig and EGF homology domains 1
Tie-2	Tyrosine kinase with immunoglobulin and epidermal growth factor homology domain 2
TIMP-1	Tissue inhibitor of metalloproteinases-1
TIMP-2	Tissue inhibitor of metalloproteinases-2
TIMP-3	Tissue inhibitor of metalloproteinase-3
TIMP-3	Tissue inhibitor of metalloproteinases-3
TIMP-4	Tissue inhibitor of metalloproteinase-4
TNF-a	Tumor necrosis factor-alpha
TNF-b	Tumor necrosis factor-beta
TPO	Thrombopoietin
TRAIL R1	TNF-related apoptosis-inducing ligand receptor 1
TRAIL R2	TNF-related apoptosis-inducing ligand receptor 2
TRAIL	TNF-related apoptosis inducing ligand
TRANCE	Tumor necrosis factor-related activation-induced cytokine
TSLP	Thymic stromal derived lymphopoietin
uPA	Urokinase plasminogen activator Ab
uPAR	Urokinase plasminogen activator receptor
VCAM-1	Vascular cell adhesion molecule (VCAM-1, CD106, or INCAM-110)
VE-Cadherin	Vascular epithelial cadherin
VEGF	Vascular Endothelial Growth Factor
VEGF R2	Vascular endothelial growth factor receptor 2
VEGF R3	Vascular endothelial growth factor receptor 3
VEGF-B	Vascular Endothelial Growth Factor B
VEGF-C	Vascular Endothelial Growth Factor C
VEGF-D	Vascular Endothelial Growth Factor D

Claims

1. A substantially pure population of amniotic fluid-derived cells, wherein said cells are:

capable of differentiating into cells displaying the characteristics of the beta-cell lineage, substantially negative for the expression of the CD117 and Oct-4 protein markers, wherein the markers are not present or expressed in at least 70% of the total cell population, substantially positive for the expression of GATA-6 and SSEA-4, wherein GATA-6 and SSEA-4 are present or expressed in at least about 50% of the total cell population, and substantially either

- i) positive for the expression of SOX-17 and cytokeratin, wherein SOX-17 and cytokeratin are present or expressed in at least about 50% of the total cell population;
- ii) negative for the expression of SOX-17, wherein SOX-17 is not present or expressed in at least 70% of the total cell population, and positive in the expression of cytokeratin, wherein cytokeratin is present or expressed in at least about 50% of the total cell population; or
- iii) negative for the expression of SOX-17 and cytokeratin, wherein SOX-17 and cytokeratin are not present or expressed in at least 70% of the total cell population.

2. The population of amniotic fluid-derived cells according to claim 1, wherein said cells are substantially positive for the expression of SOX-17 and cytokeratin, wherein SOX-17 and cytokeratin are present or expressed in at least about 50% of the total cell population, and substantially positive for the expression of the HNF-1 beta, HNF-3 beta and GATA-6 protein markers, wherein the markers are present or expressed in at least about 50% of the total cell population.
3. The population of amniotic fluid-derived cells according to claim 1, wherein said cells are substantially negative for the expression of SOX-17, wherein SOX-17 is not present or expressed in at least 70% of the total cell population, and positive in the expression of cytokeratin, wherein cytokeratin is present or expressed in at least about 50% of the total cell population, and do not express the markers HNF-3 beta and GATA-4.
4. The population of amniotic fluid-derived cells according to claim 3, wherein said cells do not express any of the markers HNF-3 beta, GATA-4 or Tra2-54.
5. The population of amniotic fluid-derived cells according to claim 1, wherein said cells are substantially negative for the expression of SOX-17 and cytokeratin, wherein SOX-17 and cytokeratin are not present or expressed in at least about 70% of the total cell population, and do not express the markers HNF-3 beta and GATA-4.
6. The population of amniotic fluid-derived cells according to claim 5, wherein said cells do not express any of the markers HNF-3 beta, GATA-4 or Tra2-54.
7. The population of amniotic fluid-derived cells according to any of the preceding claims, wherein said cells are substantially positive for the expression of the HES-1 gene, wherein HES-1 is present or expressed in at least about 50% of the total cell population.
8. The population of amniotic fluid-derived cells according to any of the preceding claims, capable of propagating in vitro.
9. The population of amniotic fluid-derived cells according to any of the preceding claims, capable of propagating in vitro under hypoxic conditions.
10. The population of amniotic fluid-derived cells according to any of the preceding claims, capable of differentiating into a gut hormone-producing cell.
11. A population of amniotic fluid-derived cells as defined in any one of claims 1 to 10, for use in a method of treating a patient with diabetes mellitus or at risk of developing diabetes.
12. In vitro use of the population of cells according to any one of claims 1 to 10, in a method wherein the cells of said population are differentiated into pancreatic hormone producing cells.

Patentansprüche

1. Im Wesentlichen reine Population von aus Fruchtwasser gewonnenen Zellen, wobei die Zellen:

dazu fähig sind, zu Zellen zu differenzieren, die die Charakteristika der beta-Zelle-Linie zeigen, im Wesentlichen negativ für die Expression der CD117- und Oct-4-Proteinmarker sind, wobei die Marker in mindestens 70% der Zellgesamtpopulation nicht vorhanden sind bzw. exprimiert werden, im Wesentlichen positiv für die Expression von GATA-6 und SSEA-4 sind, wobei GATA-6 und SSEA-4 in

mindestens etwa 50% der Zellgesamtpopulation vorhanden sind bzw. exprimiert werden, und im Wesentlichen entweder

- 5 i) positiv für die Expression von SOX-17 und Cytokeratin sind, wobei SOX-17 und Cytokeratin in mindestens etwa 50% der Zellgesamtpopulation vorhanden sind oder exprimiert werden,
ii) negativ für die Expression von SOX-17 sind, wobei SOX-17 in mindestens 70% der Zellgesamtpopulation nicht vorhanden ist bzw. exprimiert wird, und positiv für die Expression von Cytokeratin sind, wobei Cytokeratin in mindestens etwa 50% der Zellgesamtpopulation vorhanden ist bzw. exprimiert wird, oder
10 iii) negativ für die Expression von SOX-17 und Cytokeratin sind, wobei SOX-17 und Cytokeratin in mindestens 70% der Zellgesamtpopulation nicht vorhanden sind bzw. exprimiert werden.
2. Population von aus Fruchtwasser gewonnenen Zellen nach Anspruch 1, wobei die Zellen im Wesentlichen positiv für die Expression von SOX-17 und Cytokeratin sind, wobei SOX-17 und Cytokeratin in mindestens etwa 50% der Zellgesamtpopulation vorhanden sind bzw. exprimiert werden, und
15 im Wesentlichen positiv für die Expression von HNF-1-beta-, HNF-3-beta- und GATA-6-Proteinmarkern sind, wobei die Marker in mindestens etwa 50% der Zellgesamtpopulation vorhanden sind bzw. exprimiert werden.
3. Population von aus Fruchtwasser gewonnenen Zellen nach Anspruch 1, wobei die Zellen im Wesentlichen negativ für die Expression von SOX-17 sind, wobei SOX-17 in mindestens 70% der Zellgesamtpopulation nicht vorhanden ist bzw. exprimiert wird, und positiv für die Expression von Cytokeratin sind, wobei Cytokeratin in mindestens etwa 50% der Zellgesamtpopulation vorhanden ist bzw. exprimiert wird, und
20 die Marker HNF-3-beta und GATA-4 nicht exprimieren.
4. Population von aus Fruchtwasser gewonnenen Zellen nach Anspruch 3, wobei die Zellen keinen der Marker HNF-3-beta, GATA-4 bzw. Tra2-54 exprimieren.
25
5. Population von aus Fruchtwasser gewonnenen Zellen nach Anspruch 1, wobei die Zellen im Wesentlichen negativ für die Expression von SOX-17 und Cytokeratin sind, wobei SOX-17 und Cytokeratin in mindestens etwa 70% der Zellgesamtpopulation nicht vorhanden sind bzw. exprimiert werden, und
30 die Marker HNF-3-beta und GATA-4 nicht exprimieren.
6. Population von aus Fruchtwasser gewonnenen Zellen nach Anspruch 5, wobei die Zellen keinen der Marker HNF-3-beta, GATA-4 bzw. Tra2-54 exprimieren.
- 35 7. Population von aus Fruchtwasser gewonnenen Zellen nach einem der vorhergehenden Ansprüche, wobei die Zellen im Wesentlichen positiv für die Expression des HES-1-Gens sind, wobei HES-1 in mindestens etwa 50% der Zellgesamtpopulation vorhanden ist bzw. exprimiert wird.
8. Population von aus Fruchtwasser gewonnenen Zellen nach einem der vorhergehenden Ansprüche, fähig zur in-vitro-Fortpflanzung.
40
9. Population von aus Fruchtwasser gewonnenen Zellen nach einem der vorhergehenden Ansprüche, fähig zur in-vitro-Fortpflanzung unter hypoxischen Bedingungen.
- 45 10. Population von aus Fruchtwasser gewonnenen Zellen nach einem der vorhergehenden Ansprüche, fähig zur Differenzierung zu einer hormonproduzierenden Eingeweidezelle.
11. Population von aus Fruchtwasser gewonnenen Zellen nach einem der Ansprüche 1 bis 10 zur Verwendung in einem Verfahren zur Behandlung eines Patienten mit Diabetes mellitus bzw. eines Patienten, bei dem ein Risiko des Auftretens von Diabetes besteht.
50
12. In-vitro-Anwendung der Population von Zellen nach einem der Ansprüche 1 bis 10 in einem Verfahren, bei dem die Zellen der Population zu hormonproduzierenden Bauchspeicheldrüsenzellen differenziert werden.

55

Revendications

1. Population sensiblement pure de cellules dérivées de liquide amniotique, dans laquelle lesdites cellules sont :

capables de se différencier en cellules présentant les caractéristiques du lignage de cellules bêta, sensiblement négatives pour l'expression des marqueurs protéiques CD117 et Oct-4, où les marqueurs ne sont pas présents ou exprimés dans au moins 70 % de la population totale de cellules, sensiblement positives pour l'expression de GATA-6 et SSEA-4, où GATA-6 et SSEA-4 sont présents ou exprimés dans au moins environ 50 % de la population totale de cellules, et sensiblement

- i) positives pour l'expression de SOX-17 et de cytokératine, où SOX-17 et la cytokératine sont présents ou exprimés dans au moins environ 50 % de la population totale de cellules ;
- ii) négatives pour l'expression de SOX-17, où SOX-17 n'est pas présent ou exprimé dans au moins 70 % de la population totale de cellules, et positives pour l'expression de cytokératine, où la cytokératine est présente ou exprimée dans au moins environ 50 % de la population totale de cellules ; ou
- iii) négatives pour l'expression de SOX-17 et de cytokératine, où SOX-17 et la cytokératine ne sont pas présents ou exprimés dans au moins 70 % de la population totale de cellules.

2. Population de cellules dérivées de liquide amniotique selon la revendication 1, dans laquelle lesdites cellules sont sensiblement positives pour l'expression de SOX-17 et de cytokératine, où SOX-17 et la cytokératine sont présents ou exprimés dans au moins environ 50 % de la population totale de cellules, et sensiblement positives pour l'expression des marqueurs protéiques HNF-1 bêta, HNF-3 bêta et GATA-6, où les marqueurs sont présents ou exprimés dans au moins environ 50 % de la population totale de cellules.

3. Population de cellules dérivées de liquide amniotique selon la revendication 1, dans laquelle lesdites cellules sont sensiblement négatives pour l'expression de SOX-17, où SOX-17 n'est pas présent ou exprimé dans au moins 70 % de la population totale de cellules, et positives dans l'expression de cytokératine, où la cytokératine est présente ou exprimée dans au moins environ 50 % de la population totale de cellules, et n'expriment pas les marqueurs HNF-3 bêta et GATA-4.

4. Population de cellules dérivées de liquide amniotique selon la revendication 3, dans laquelle lesdites cellules n'expriment aucun des marqueurs HNF-3 bêta, GATA-4 ou Tra2-54.

5. Population de cellules dérivées de liquide amniotique selon la revendication 1, dans laquelle lesdites cellules sont sensiblement négatives pour l'expression de SOX-17 et de cytokératine, où SOX-17 et la cytokératine ne sont pas présents ou exprimés dans au moins environ 70 % de la population totale de cellules, et n'expriment pas les marqueurs HNF-3 bêta et GATA-4.

6. Population de cellules dérivées de liquide amniotique selon la revendication 5, dans laquelle lesdites cellules n'expriment aucun des marqueurs HNF-3 bêta, GATA-4 ou Tra2-54.

7. Population de cellules dérivées de liquide amniotique selon l'une quelconque des revendications précédentes, dans laquelle lesdites cellules sont sensiblement positives pour l'expression du gène HES-1, où HES-1 est présent ou exprimé dans au moins environ 50 % de la population totale de cellules.

8. Population de cellules dérivées de liquide amniotique selon l'une quelconque des revendications précédentes, capable de se propager *in vitro*.

9. Population de cellules dérivées de liquide amniotique selon l'une quelconque des revendications précédentes, capable de se propager *in vitro* dans des conditions hypoxiques.

10. Population de cellules dérivées de liquide amniotique selon l'une quelconque des revendications précédentes, capable de se différencier en cellule productrice d'hormone intestinale.

11. Population de cellules dérivées de liquide amniotique telle que définie dans l'une quelconque des revendications 1 à 10, pour utilisation dans un procédé de traitement d'un patient ayant un diabète sucré ou à risque de développer un diabète.

12. Utilisation *in vitro* de la population de cellules selon l'une quelconque des revendications 1 à 10, dans un procédé dans lequel les cellules de ladite population sont différenciées en cellules productrices d'hormone pancréatiques.




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

F Cells




Figure 7

E Cells




Figure 8

AF-I Cells




Figure 9




Figure 10




Figure 11





Figure 12






Figure 13

Figure 14




Figure 15




Figure 16





Figure 17



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2004087885 A2 [0005]
- WO 03033697 A [0006]
- US 20040161419 A1 [0006]
- US 20040241761 A [0008]
- US 20030082155 A [0011]
- US 5834308 A [0012]
- US 6001647 A [0012]
- US 6703017 B [0012]
- US 6815203 B1 [0013]
- WO 2004011621 A [0014]
- WO 03102134 A [0015]
- US 2004015805 A1 [0016]
- US 6436704 B [0017]
- WO 03062405 A [0017]
- WO 02092756 A [0017]
- EP 0363125 A2 [0017]
- WO 2003042405 A [0025]
- US 20050054093 A [0026]
- WO 2005017117 A [0033]
- EP 1391505 A1 [0034]
- US 20040209901 A [0127] [0140] [0142]
- US 20040132729 A [0127] [0142]
- US 6328960 B [0130]
- US 6281012 B [0130]
- US 5770417 A [0137]
- US 6022743 A [0137]
- US 5567612 A [0137]
- US 5759830 A [0137]
- US 6626950 B [0137]
- US 6534084 B [0137]
- US 6306424 B [0137]
- US 6365149 B [0137]
- US 6599323 B [0137]
- US 6656488 B [0137]
- US 6333029 B [0137]
- US 20040062753 A1 [0137]
- US 4557264 A [0137]
- US 6509369 B [0137]
- US 6793945 B [0138]
- US 6331298 B [0139]
- US 20040220393 A [0140]
- US 20040171623 A [0141]

Non-patent literature cited in the description

- HESS et al. *Nature Biotechnology*, 2003, vol. 21, 763-770 [0005]
- IANUS et al. *J. Clin. Invest.*, 2003, vol. 111, 843-850 [0005]
- GERSHENGORN et al. *Science*, 2004, vol. 306, 2261-2264 [0009]
- SEABERG et al. *Nature Biotechnology*, 2004, vol. 22, 1115-1124 [0010]
- BONNER-WEIR et al. *Proc Nat Acad Sci*, 2000, vol. 97, 7999-8004 [0013]
- THOMSON et al. *Proc. Natl. Acad. Sci.*, 1995, vol. 92, 7844-7848 [0018]
- THOMSON et al. *Science*, 1998, vol. 282, 1145-1147 [0018]
- MARSHALL et al. *Methods Mol. Biol.*, 2001, vol. 158, 11-18 [0018]
- HENDERSON et al. *Stem Cells*, 2002, vol. 20, 329-337 [0019]
- DRAPER et al. *J. Anat.*, 2002, vol. 200, 249-258 [0019]
- MITSUI et al. *Cell*, 2003, vol. 113, 631-642 [0019]
- CHAMBERS et al. *Cell*, 2003, vol. 113, 643-655 [0019]
- CRANE ; CHEUNG. *Prenatal Diagnosis*, 1988, vol. 8, 119-129 [0023]
- MILUNSKY. *Genetic Disorder of the Fetus*. Plenum Press, 1979, 75-84 [0023]
- HOEHN ; SALK. *Methods in Cell Biology*, 1982, vol. 26, 11-34 [0023]
- GOSDEN. *British Medical Bulletin*, 1983, vol. 39, 348-354 [0023]
- PRUSA et al. *Human Reproduction*, 2003, vol. 18, 1489-1493 [0023]
- ANKER et al. *Blood*, 2003, vol. 102, 1548-1549 [0027]
- TSAI et al. *Human Reproduction*, 2004, vol. 19, 1450-1456 [0028]
- WEI JUN PING et al. *Cell Transplantation*, 2003, vol. 12 (5), 545-552 [0032]
- CHEN LI-BO et al. *World J. Gastroenterology*, 2004, vol. 10 (20), 3016-3020 [0032]
- HELD KR et al. *Prenatal Diagnosis*, 1984, vol. 4 (3), 171-180 [0032]
- FAUZA D. *Bailliere's Best Practice and Research: Clinical Obstetrics and Gynaecology*, 2004, vol. 18 (6), 877-891 [0032]

EP 1 780 267 B1

- *Nature Reviews Genetics*, 2002, vol. 3, 524-632 [0061]
- *Methods in Enzymology. Cell Culture*. Academic Press Inc, vol. LVIII, 62-72 [0092]
- *Current Protocols in Molecular Biology*. 2001 [0107]
- **HARLOW ; LANE**. *Using Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1998 [0107]
- **EDLUND**. *Nature Reviews Genetics*, 2002, vol. 3, 524-632 [0109]
- **SCHONHOFF**. *Endocrinology*, 2004, vol. 145, 2639-2644 [0110]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. 1989 [0131]
- *Oligonucleotide Synthesis*. 1984 [0131]
- *Animal Cell Culture*. 1987 [0131]
- *Methods in Enzymology*. Academic Press, Inc, [0131]
- *Gene Transfer Vectors for Mammalian Cells*. 1987 [0131]
- *Current Protocols in Molecular Biology and Short Protocols in Molecular Biology*. 1987 [0131]
- *Recombinant DNA Methodology II*. Academic Press, 1995 [0131]
- *J. Pediatr. Surg.*, 1988, vol. 23, 3-9 [0143]
- *Biotechnol. Prog.*, 1998, vol. 14 (2), 193-202 [0143]
- **YANG et al.** *J. Biomed. Mater. Res.*, 2001, vol. 55 (3), 379-86 [0143]

Szabadalmi igénypontok

1. Amnionfolyadék-eredetű sejtek lényegében tiszta populációja, ahol a sejtek:
 - képesek béta-sejtvonal leszármazási vonalba tartozó tulajdonságokat mutató sejtekké differenciálódni,
 - lényegében negatívak a CD117 és Oct-4 protein-markerek expressziójára, ahol a markerek nincsenek jelen vagy nem expresszálódnak a teljes sejtpopuláció legalább 70%-ában,
 - lényegében pozitívak a GATA-6 és SSEA-4 expressziójára, ahol a GATA-6 és SSEA-4 jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában, és
 - lényegében vagy
 - i) pozitívak a SOX-17 és citokeratin expressziójára, ahol a SOX-17 és citokeratin jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában;
 - ii) negatívak a SOX-17 expressziójára, ahol a SOX-17 nincs jelen vagy nem expresszálódik a teljes sejtpopuláció legalább 70%-ában, és pozitívak a citokeratin expressziójára, ahol a citokeratin jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában; vagy
 - iii) negatívak a SOX-17 és citokeratin expressziójára, ahol a SOX-17 és citokeratin nincs jelen vagy nem expresszálódik a teljes sejtpopuláció legalább 70%-ában.
2. Az 1. igénypont szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek lényegében pozitívak a SOX-17 és citokeratin expressziójára, ahol a SOX-17 és citokeratin jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában, és lényegében pozitívak a HNF-1 béta, HNF-3 béta és GATA-6 protein-markerek expressziójára, ahol a markerek jelen vannak vagy expresszálódnak a teljes sejtpopuláció legalább 50%-ában.
3. Az 1. igénypont szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek lényegében negatívak a SOX-17 expressziójára, ahol a SOX-17 nincs jelen vagy nem expresszálódik a teljes sejtpopuláció legalább 70%-ában, és pozitívak a citokeratin expressziójára, ahol a citokeratin jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában, és nem expresszálják a HNF-3 béta és GATA-4 markereket.
4. A 3. igénypont szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek nem expresszálják a HNF-3 béta, GATA-4 vagy Tra2-54 markerek egyikét sem.
5. Az 1. igénypont szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek lényegében negatívak a SOX-17 és citokeratin expressziójára, ahol a SOX-17 és citokeratin nincs jelen vagy nem expresszálódik a teljes sejtpopuláció legalább 70%-ában, és nem expresszálják a HNF-3 béta és GATA-4 markereket.
6. Az 5. igénypont szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek nem expresszálják a HNF-3 béta, GATA-4 vagy Tra2-54 markerek egyikét sem.
7. Az előző igénypontok bármelyike szerinti amnionfolyadék-eredetű sejtpopuláció, ahol a sejtek lényegében pozitívak a HES-1 gén expressziójára, ahol a HES-1 jelen van vagy expresszálódik a teljes sejtpopuláció legalább 50%-ában.
8. Az előző igénypontok bármelyike szerinti amnionfolyadék-eredetű sejtpopuláció, amely képes szaporodni *in vitro*.

9. Az előző igénypontok bármelyike szerinti amnionfolyadék-eredetű sejtpopuláció, amely képes szaporodni *in vitro* hipoxiás körülmények között.

10. Az előző igénypontok bármelyike szerinti amnionfolyadék-eredetű sejtpopuláció, amely képes bél-hormontermelő sejtekké differenciálódni.

11. Az 1-10. igénypontok bármelyikében definiált amnionfolyadék-eredetű sejtpopuláció, cukorbetegségben szenvedő vagy cukorbetegség kialakulása kockázatának kitétt páciens kezelési eljárásában történő alkalmazásra.

12. Az 1-10. igénypontok bármelyikében definiált amnionfolyadék-eredetű sejtpopuláció *in vitro* alkalmazása olyan eljárásban, ahol a sejtpopuláció hasnyálmirigy-hormont termelő sejtekké van differenciálható.