



(51) International Patent Classification:

A61K 8/64 (2006.01) A61P 37/00 (2006.01)

A61K 35/74 (2015.01)

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

(21) International Application Number:

PCT/US2018/042530

(22) International Filing Date:

17 July 2018 (17.07.2018)

Published:

- with international search report (Art. 21(3))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/534,053 18 July 2017 (18.07.2017) US

62/534,024 18 July 2017 (18.07.2017) US

(71) Applicant: AOBIONE LLC [US/US]; One Broadway, 14th Floor, Cambridge, MA 02142 (US).

(72) Inventors: KRUEGER, Todd; One Broadway, 14th Floor, Cambridge, MA 02142 (US). WEISS, Larry; One Broadway, 14th Floor, Cambridge, MA 02142 (US).

(74) Agent: ANASTASI, John, N. et al.; Lando & Anastasi, LLP, Riverfront Office Park, One Main Street, Suite 1100, Cambridge, MA 02142 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: AMMONIA OXIDIZING MICROORGANISMS FOR USE AND DELIVERY TO THE VISUAL AND AUDITORY SYSTEMS

(57) Abstract: Ammonia oxidizing microorganism preparations for delivery to the visual and auditory systems, kits including ammonia oxidizing preparations for delivery to the visual and auditory systems, and devices for administering ammonia oxidizing preparations to the visual and auditory systems are provided. Methods of introducing ammonia oxidizing microorganisms to the eye are provided. Methods of introducing ammonia oxidizing microorganisms to the ear are provided. Methods of treating disorders, including eye disorders, ear disorders, and inflammatory disorders, with ammonia oxidizing microorganism preparations are provided.



AMMONIA OXIDIZING MICROORGANISMS FOR USE AND DELIVERY TO THE VISUAL AND AUDITORY SYSTEMS

Cross-Reference to Related Applications

This application claims the benefit of priority to U.S. Provisional Patent Application Serial No. 62/534,024 filed July 18, 2017 titled “AMMONIA OXIDIZING MICROORGANISMS FOR USE AND DELIVERY TO THE VISUAL SYSTEM,” and U.S. Provisional Patent Application Serial No. 62/534,053 filed July 18, 2017 titled “AMMONIA OXIDIZING MICROORGANISMS FOR USE AND DELIVERY TO THE AUDITORY SYSTEM,” the entire disclosure of each of which is hereby incorporated herein by reference in its entirety for all purposes.

Field of the Technology

Aspects relate generally to the microbiome and, more specifically, to the restoration of ammonia oxidizing microorganisms in relation to the microbiome.

Background

Bacteria and other microorganisms are ubiquitous in the environment. The discovery of pathogenic bacteria and the germ theory of disease have had a tremendous effect on health and disease states. Microorganisms are a normal part of the environment of all living things and may be beneficial. In the gut, for example, bacteria are not pathogenic under normal conditions, and in fact improve health by rendering the normal intestinal contents less hospitable for disease causing organisms.

Summary

In accordance with an aspect, there is provided a method of introducing ammonia oxidizing microorganisms (AOM) to a subject. The method may comprise administering a preparation comprising AOM to an eye of the subject, for example, via topical or intravitreal delivery. In some embodiments, the method may comprise administering a preparation comprising live AOM.

In some embodiments, the preparation comprising AOM may be administered topically. The preparation comprising AOM may be administered to a first tissue, wherein the first tissue may be a deposit tissue or a target tissue. In some embodiments, for example those wherein the first tissue is other than the target tissue, the preparation comprising AOM may be applied to the first tissue and the preparation or a product of the preparation, e.g., NO, may be transported, for example, via diffusion to a second tissue. The second tissue may be a target tissue.

In some embodiments, the deposit tissue, target tissue, or both may be related to an optic tissue. The deposit tissue, target tissue, or both may relate to an intravitreous tissue, posterior chamber, or anterior chamber of the eye of the subject. In some embodiments, a deposit tissue, target tissue, or both is an eyelid, inner eyelid, retina, ocular globe, orbit, optic disk, optic nerve, vitreous humor, intrascleral, subconjunctiva, lens, ciliary body, iris, pupil, cornea, conjunctiva, sclera, topical tissue, or proximate tissue of the eye of the subject.

In some embodiments, a target tissue may be associated with a desired local effect. The desired local effect may involve, for example, treatment of a periocular condition, blepharitis, dry eye, itching, inflammation, wound healing, intraocular diseases, macular degeneration, infections, ocular blood pressure, glaucoma, elevated intraocular pressure, ocular hypertension, and ocular hypotension.

In some embodiments, a target tissue may be associated with a desired systemic effect. The desired systemic effect may involve, for example, treatment of one or more of headaches, cardiovascular diseases, inflammation, immune responses and autoimmune disorders, liver diseases, mitochondrial diseases, infections, neurological diseases, psychiatric disorders, nitric oxide disorders, urea cycle disorders, congestion, vasodilation disorders, skin diseases, wound healing, reactions to insect bites, connective tissue disorders, and certain viral, bacterial, and fungal infections.

In some embodiments, administering the preparation comprising AOM may be substantially noninvasive. In some embodiments, administering the preparation comprising AOM may be device-assisted.

In some embodiments, an effective amount of the preparation comprising AOM may be administered to a target tissue of the eye of the subject. In some embodiments, administering an effective amount of the preparation comprising AOM may promote lubrication or hydration of

the eye. In some embodiments, administering an effective amount of the preparation comprising AOM may change or alter a level of nitrite or NO at a target tissue. The target tissue may be in the eye or in circulation. In some embodiments, administering an effective amount of the preparation may modulate a microbiome of the subject, for example a microbiome associated with the eye of the subject.

In accordance with another aspect, there is provided a method of treating an inflammatory eye disorder in a subject. The method may comprise administering to the subject an effective amount of a preparation comprising AOM, thereby treating the inflammatory eye disorder.

In some embodiments, the inflammatory eye disorder may comprise or be associated with dry eye or Meibomian Gland Dysfunction (MBD). The inflammatory eye disorder may comprise or be associated with blepharitis.

In some embodiments, the inflammatory eye disorder may comprise a disorder of the back of the eye, for example, a disorder of the retina. The eye disorder may comprise or be associated with macular degeneration.

The eye disorder may be associated with intraocular pressure. In some embodiments, the eye disorder may be associated with or comprise elevated intraocular pressure.

In some embodiments, the eye disorder may comprise or be associated with an infectious disorder. The eye disorder may comprise or be associated with a bacterial, fungal, or viral infection. In some embodiments, the bacterial, fungal, or viral infection may be of the ocular globe or other eye tissue. In some embodiments, the inflammatory eye disorder may be characterized by or comprise sty or chalazion formation.

In some embodiments, the inflammatory eye disorder may comprise or be associated with an allergy.

In some embodiments, the preparation comprising AOM may be administered prior to onset of the inflammatory eye disorder. In some embodiments, the preparation comprising AOM may be administered during incidence of the inflammatory eye disorder. In some embodiments, the preparation comprising AOM may be administered subsequent to subsiding of the inflammatory eye disorder. In some embodiments, the preparation may be administered in response to an inflammatory eye disorder symptom, trigger, or warning sign. In some

embodiments, the preparation may be administered before or after a surgical or diagnostic procedure.

In some embodiments, methods disclosed herein may further comprise determining whether the subject is in need of treatment for the eye disorder.

5 In some embodiments, the preparation comprising AOM may be administered as a solution, liquid, ointment, gel, hydrogel, suspension, emulsion, or insert. In some embodiments, the preparation may be administered as a droplet or wash. In some embodiments, the preparation may be administered as a spray, mist, or aerosol.

10 In some embodiments, the preparation may be formulated to resemble tears of the subject. In some embodiments, the preparation comprising AOM may be formulated to be compatible with the eye of the subject. For instance, the preparation may have a physiological pH level. The preparation may have a pH level between about 7.14 and 7.82. In some embodiments, the preparation may have a physiological osmolarity. For instance, the preparation may have an osmolarity of between about 100 mOsmol/L and about 640 mOsmol/L.

15 In some embodiments, the preparation may have a physiological surface tension. For instance, the preparation may have a surface tension of between about 43 nN/m and about 47 nN/m.

In some embodiments, the preparation comprising AOM may be formulated for immediate release or extended release.

20 The preparation or product thereof may be formulated to penetrate and otherwise reach a target eye tissue. In some embodiments, the preparation or product thereof may be formulated to permeate or be absorbed through the cornea.

In some embodiments, the preparation comprising AOM may be formulated to deliver nitrite or NO to a target tissue, locally or systemically. The preparation or product thereof may be formulated to be delivered for systemic circulation.

25 In some embodiments, methods disclosed herein may further comprise administering a second amount of a preparation comprising to the subject.

The preparation may be administered as part of a combination therapy. In some embodiments, methods disclosed herein may further comprise administering a second treatment in combination with the preparation comprising AOM. The second treatment may comprise a

30 surgical procedure. In some embodiments, the preparation comprising AOM may be

administered in conjunction with an anti-inflammatory agent. The preparation may be administered in conjunction with a medical approach that treats, is approved to treat, or is commonly used to treat, a relevant disease or disorder, or a symptom of a relevant disease or disorder.

5 In some embodiments, the preparation comprising AOM may be administered for a period of time prior to initiating the second treatment. In some embodiments, the preparation comprising AOM may be administered concurrently with the second treatment. In some embodiments, the preparation comprising AOM may be administered for a period of time subsequent to ceasing the second treatment.

10 In some embodiments, the second treatment may be administered to the eye. The second treatment may be topically or intravitreally administered to the eye. The second treatment may be administered via an alternate mode of administration, for example, orally.

 In some embodiments, the subject may have a therapeutic level of a second treatment.

 The preparation comprising AOM may be administered in conjunction with one or more
15 ophthalmic ion-exchange formulations, chelators, penetration enhancers, transporter-targeted prodrugs, ultrasound, soft drugs, mucoadhesives, or steam, for example, a warm compress. In some embodiments, the preparation comprising AOM may be administered in conjunction with nitrite, nitrate, and/or NO.

 In some embodiments, the effective amount of AOM or a preparation comprising AOM
20 may be a therapeutically effective dose of AOM. The therapeutically effective dose of AOM may be about or greater than about 1×10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} CFU.

 In some embodiments, the preparation may be administered as an analgesic. In some
25 embodiments, the preparation may be administered as a prophylactic. In some embodiments, the preparation may be self-administered.

 Methods disclosed herein may comprise administering a preparation comprising AOM about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 times per day. The preparation may be administered for about 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, or 84-91 days.

In some embodiments, the preparation comprising AOM may be administered within 30, 60, 90, 120, 150, or 180 minutes of the subject waking from sleep. In some embodiments, the preparation comprising AOM may be administered within 30, 60, 90, 120, 150, or 180 minutes prior to the subject sleeping. In some embodiments, the preparation may be administered within
5 30, 60, 90, 120, 150, or 180 minutes of the subject eating. In some embodiments, the preparation may be administered within 30, 60, 90, 120, 150, or 180 minutes before the subject cleanses or showers.

In some embodiments, the subject may be female. In some embodiments, the subject may be male. The subject may be characterized as one of the following ethnicity/race: Asian,
10 black or African American, Hispanic or Latino, white, or multi-racial. In some embodiments, the subject may be of an age of less than 1, or between 1-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, or over 60 years. In some embodiments, the subject may have a disrupted microbiome.

In some embodiments, the preparation may comprise AOM in a buffer solution. The preparation may comprise AOM in an aqueous buffer solution. The buffer solution may
15 comprise disodium phosphate and magnesium chloride. In some embodiments, the buffer may comprise 50 mM Na_2HPO_4 and/or 2 mM MgCl_2 in water. The buffer solution may consist essentially of disodium phosphate and magnesium chloride, for example, consist essentially of 50 mM Na_2HPO_4 and/or 2 mM MgCl_2 in water. The buffer solution may consist of disodium
20 phosphate and magnesium chloride, for example, consist of 50 mM Na_2HPO_4 and/or 2 mM MgCl_2 in water.

In some embodiments, the preparation comprising AOM may comprise at least one of ammonia, ammonium salts, and urea. In some embodiments, the preparation may further comprise or be administered concurrently with a compound that promotes growth or metabolism of the AOM, NO production, and/or urease activity.

25 In some embodiments, the preparation comprising AOM may comprise a controlled release material. The preparation may comprise a slow release material.

In some embodiments, the preparation may further comprise an excipient. The preparation comprising AOM may comprise a pharmaceutically acceptable excipient. The excipient may comprise one or more of an absorption and penetration enhancer, preservative,
30 antioxidant, buffer, chelating agent, ion exchange agent, solubilizing agent, suspending agent,

thickener, surfactant, wetting agent, tonicity-adjusting agent, and a vehicle for proper drug delivery. In some embodiments, the excipient may comprise a surfactant.

In some embodiments, the preparation comprising AOM may be substantially free of other organisms. The preparation may further comprise other organisms, e.g., a community of organisms.

In some embodiments, the preparation comprising AOM may comprise between about 1×10^3 CFU/mL to about 1×10^{14} CFU/mL AOM. For instance, the preparation may comprise between about 1×10^9 CFU/mL to about 10×10^9 CFU/mL AOM.

In some embodiments, the AOM comprises ammonia oxidizing bacteria (AOB). The AOM may consist essentially of AOB. The AOM may consist of AOB.

In some embodiments, the AOM may comprise *Nitrosomonas*, *Nitrosococcus*, *Nitrospira*, *Nitrosocystis*, *Nitrosolobus*, *Nitrosovibrio*, and combinations thereof. In some embodiments the AOM may be *Nitrosomonas eutropha* (*N. eutropha*). In some embodiments, the AOM may be *N. eutropha* D23, having ATCC accession number PTA-121157.

In some embodiments, the AOM may comprise ammonia oxidizing archaea (AOA). The AOM may consist essentially of AOA. The AOM may consist of AOA.

In some embodiments, the AOM may be capable of converting ammonia or ammonium to nitrite at a rate of at least about 1 pmol/min/mg protein. The AOM may be capable of converting ammonia or ammonium to nitrite at a rate of at least about 0.1 nmol/min/mg protein.

In some embodiments, a target percentage of administered AOM may be transferred to the eye of the subject. For example, in some embodiments, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of administered AOM may be transferred to a deposit or target tissue of the subject.

In some embodiments, methods disclosed herein may comprise using a biome-friendly product in connection with the administered preparation comprising AOM.

In accordance with an aspect, there is provided a preparation comprising AOM, as disclosed herein, for administration to an eye of a subject.

In accordance with an aspect, there is provided a preparation comprising AOM, as disclosed herein, for treating an inflammatory eye disorder in a subject.

In some embodiments the preparation comprising AOM may be packaged for single use. The preparation comprising AOM may be packaged for multiple use.

In accordance with an aspect, there is provided a device configured to administer a preparation comprising AOM, as disclosed herein. In some embodiments, the device is
5 configured to administer a preparation comprising AOM to an eye of a subject.

In accordance with an aspect, there is provided a kit comprising a preparation comprising AOM, as disclosed herein.

In accordance with an aspect, there is provided a method of introducing ammonia oxidizing microorganisms (AOM) to a subject. The method may comprise administering a
10 preparation comprising AOM to an ear of the subject. In some embodiments, the method may comprise administering a preparation comprising live AOM.

In some embodiments, administering the preparation comprising AOM may be substantially noninvasive. In some embodiments, administering the preparation comprising AOM may be device-assisted.

15 In some embodiments, an effective amount of the preparation comprising AOM may be administered to a target tissue of an ear of a subject. In some embodiments, the preparation comprising AOM may be administered topically. The preparation comprising AOM may be administered to a first tissue, wherein the first tissue may be a deposit tissue or a target tissue. In some embodiments, for example those wherein the first tissue is other than the target tissue, the
20 preparation comprising AOM may be applied to the first tissue and the preparation or a product of the preparation, e.g., NO, may be transported, for example, via diffusion to a second tissue. The second tissue may be a target tissue.

In some embodiments, the deposit tissue, target tissue, or both may be related to an otic tissue. The deposit tissue, target tissue, or both may be an outer ear tissue, inner ear tissue,
25 external auditory canal, internal auditory canal, tympanic membrane, tympanic cavity, malleus, incus, auditory tube, Eustachian tube, round window, oval window, ossicle, labyrinth, vestibule, utricle, saccule, staple, cochlea, canal, semicircular canal, temporal bone, vestibular nerve, or cochlear nerve of the subject.

In some embodiments, a target tissue may be associated with a desired local or systemic
30 effect. In some embodiments, administration of a preparation comprising AOM may result in,

for example, improved hearing, reduced ear ache, reduced ear infection, reduced pruritus, or reduced tinnitus in the subject. In some embodiments, administration of a preparation comprising AOM may result in, for example, reduced vertigo, dizziness, or nausea for the subject.

5 In accordance with another aspect, there is provided a method of treating an ear disorder in a subject. The method may comprise administering to the subject an effective amount of a preparation comprising AOM, thereby treating the ear disorder.

In some embodiments, the ear disorder may comprise or involve vertigo, swimmer's ear, pruritus, or tinnitus. The ear disorder may comprise or be associated with an inflammatory
10 condition, a bacterial, viral, or fungal infection, or a congenital abnormality.

In some embodiments, an effective amount of the preparation comprising AOM may be administered to a target tissue of the auditory system of the subject. In some embodiments, administering an effective amount of the preparation comprising AOM may promote endothelial function. In some embodiments, administering an effective amount of the preparation
15 comprising AOM may change or alter a level of nitrite or NO at a target tissue or in circulation. In some embodiments, administering an effective amount of the preparation may modulate a microbiome associated with the subject.

In some embodiments, the preparation comprising AOM may be administered prior to onset of the ear disorder. In some embodiments, the preparation comprising AOM may be
20 administered during incidence of the ear disorder. In some embodiments, the preparation comprising AOM may be administered subsequent to subsiding of the ear disorder. In some embodiments, the preparation may be administered in response to a trigger, warning sign, or symptom of an ear disorder. In some embodiments, the preparation may be administered before or after a surgical or diagnostic procedure.

25 In some embodiments, methods disclosed herein may further comprise determining whether the subject is in need of treatment for the eye disorder.

In some embodiments, the preparation comprising AOM may be administered as a solution, suspension, liquid, ointment, gel, hydrogel, emulsion, or insert. In some embodiments, the preparation may be administered as a droplet or wash. In some embodiments, the preparation
30 may be administered as a droplet, spray, mist, or aerosol.

In some embodiments, the preparation may be formulated to be substantially bio-compatible. For instance, the preparation may have a substantially physiological pH level. In some embodiments, the preparation may have a substantially physiological osmolarity. For instance, the preparation may be substantially isotonic.

5 In some embodiments, the preparation comprising AOM may be formulated for immediate release or extended release.

In some embodiments, the preparation comprising AOM may be formulated to deliver nitrite or NO to a target tissue, locally or systemically. The preparation or product thereof may be formulated for transmucosal delivery and/or circulation, for example, locally or systemically.

10 In some embodiments, methods disclosed herein may further comprise administering a second amount of a preparation comprising to the subject.

The preparation may be administered as part of a combination therapy. In some embodiments, methods disclosed herein may further comprise administering a second treatment in combination with the preparation comprising AOM. The second treatment may comprise a surgical procedure. In some embodiments, the preparation comprising AOM may be administered in conjunction with an anti-inflammatory agent. The preparation may be administered in conjunction with a medical approach that treats, is approved to treat, or is commonly used to treat, a relevant disease or disorder, or a symptom of a relevant disease or disorder.

20 In some embodiments, the preparation comprising AOM may be administered for a period of time prior to initiating the second treatment. In some embodiments, the preparation comprising AOM may be administered concurrently with the second treatment. In some embodiments, the preparation comprising AOM may be administered for a period of time subsequent to ceasing the second treatment.

25 In some embodiments, the second treatment may be administered to the ear. The second treatment may be administered via an alternate mode of administration, for example, orally or intranasally.

In some embodiments, the subject may have a therapeutic level of a second treatment.

In some embodiments, the preparation comprising AOM may be administered in conjunction with nitrite, nitrate, and/or NO.

30

In some embodiments, the effective amount of AOM or a preparation comprising AOM may be a therapeutically effective dose of AOM. The therapeutically effective dose of AOM may be about or greater than about 1×10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} CFU.

5 In some embodiments, the preparation may be administered as an analgesic. In some embodiments, the preparation may be administered as a prophylactic. In some embodiments, the preparation may be self-administered.

Methods disclosed herein may comprise administering a preparation comprising AOM about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 times per
10 day. The preparation may be administered for about 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, or 84-91 days.

In some embodiments, the preparation comprising AOM may be administered within 30, 60, 90, 120, 150, or 180 minutes of the subject waking from sleep. In some embodiments, the preparation comprising AOM may be administered within 30, 60, 90, 120, 150, or 180 minutes
15 prior to the subject sleeping. In some embodiments, the preparation may be administered within 30, 60, 90, 120, 150, or 180 minutes of the subject eating. In some embodiments, the preparation may be administered within 30, 60, 90, 120, 150, or 180 minutes before the subject cleanses or showers.

In some embodiments, the subject may be female. In some embodiments, the subject
20 may be male. The subject may be characterized as one of the following ethnicity/race: Asian, black or African American, Hispanic or Latino, white, or multi-racial. In some embodiments, the subject may be of an age of less than 1, or between 1-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, or over 60 years. In some embodiments, the subject may have a disrupted microbiome.

In some embodiments, the preparation may comprise AOM in a buffer solution. The
25 preparation may comprise AOM in an aqueous buffer solution. The buffer solution may comprise disodium phosphate and magnesium chloride. In some embodiments, the buffer may comprise 50 mM Na_2HPO_4 and/or 2 mM MgCl_2 in water. The buffer solution may consist essentially of disodium phosphate and magnesium chloride, for example, consist essentially of 50 mM Na_2HPO_4 and/or 2 mM MgCl_2 in water. The buffer solution may consist of disodium

phosphate and magnesium chloride, for example, consist of 50 mM Na₂HPO₄ and/or 2 mM MgCl₂ in water.

In some embodiments, the preparation comprising AOM may comprise at least one of ammonia, ammonium salts, and urea. In some embodiments, the preparation may further
5 comprise or be administered concurrently with a compound that promotes growth or metabolism of the AOM, NO production, and/or urease activity.

In some embodiments, the preparation comprising AOM may comprise a controlled release material. The preparation may comprise a slow release material.

In some embodiments, the preparation may further comprise an excipient. The preparation
10 comprising AOM may comprise a pharmaceutically acceptable excipient. The excipient may comprise one or more of water, hydrogen peroxide, mineral oil, vegetable oil, glycerin, anhydrous glycerin, propylene glycol, aluminum acetate, isopropyl alcohol, and ethanol. The preparation may comprise one or more of an analgesic agent, a chelator, a hygroscopic agent, a ceramide disrupting agent, an anti-inflammatory agent, an anti-infective agent, a reacidifying
15 agent, and a vehicle.

In some embodiments, the preparation comprising AOM may be substantially free of other organisms. The preparation comprising AOM may further comprise other organisms, e.g., a community of organisms.

In some embodiments, the preparation comprising AOM may comprise between about 1
20 x 10³ CFU/mL to about 1 x 10¹⁴ CFU/mL AOM. For instance, the preparation may comprise between about 1 x 10⁹ CFU/mL to about 10 x 10⁹ CFU/mL AOM.

In some embodiments, the AOM comprises ammonia oxidizing bacteria (AOB). The AOM may consist essentially of AOB. The AOM may consist of AOB.

In some embodiments, the AOM may comprise *Nitrosomonas*, *Nitrosococcus*,
25 *Nitrospira*, *Nitrosocystis*, *Nitrosolobus*, *Nitrosovibrio*, and combinations thereof. In some embodiments the AOM may be *Nitrosomonas eutropha* (*N. eutropha*). In some embodiments, the AOM may be *N. eutropha* D23, having ATCC accession number PTA-121157.

In some embodiments, the AOM may comprise ammonia oxidizing archaea (AOA). The AOM may consist essentially of AOA. The AOM may consist of AOA.

In some embodiments, the AOM may be capable of converting ammonia or ammonium to nitrite at a rate of at least about 1 pmol/min/mg protein. The AOM may be capable of converting ammonia or ammonium to nitrite at a rate of at least about 0.1 nmol/min/mg protein.

5 In some embodiments, a target percentage of administered AOM may be transferred to the ear of the subject. For example, in some embodiments, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of administered AOM may be transferred to a deposit or target tissue of the subject.

In some embodiments, methods disclosed herein may comprise using a biome-friendly product in connection with the administered preparation comprising AOM.

10 In accordance with an aspect, there is provided a preparation comprising AOM, as disclosed herein, for administration to an ear of a subject.

In accordance with an aspect, there is provided a preparation comprising AOM, as disclosed herein, for treating an ear disorder in a subject.

In some embodiments the preparation comprising AOM may be packaged for single use.
15 In some embodiments, the preparation comprising AOM may be packaged for multiple use.

In accordance with an aspect, there is provided a device configured to administer a preparation comprising AOM, as disclosed herein. In some embodiments, the device is configured to administer a preparation comprising AOM to an ear of a subject.

In accordance with an aspect, there is provided a kit comprising a preparation comprising
20 AOM, as disclosed herein.

The disclosure contemplates all combinations of any one or more of the foregoing aspects and/or embodiments, as well as combinations with any one or more of the embodiments set forth in the detailed description and any examples.

25 Detailed Description

In accordance with one or more embodiments, the present disclosure provides for various methods or modes of introducing ammonia oxidizing microorganisms to a subject. These methods or modes comprise administering to a subject ammonia oxidizing microorganisms, for example, a preparation, composition, formulation, or product comprising ammonia oxidizing
30 microorganisms. In at least some embodiments, ammonia oxidizing microorganisms may

therefore generally be restored to a microbiome of the subject. In at least some embodiments, ammonia oxidizing microorganisms may comprise or consist essentially of live ammonia oxidizing microorganisms.

Preparations, compositions, and/or formulations, e.g., including cosmetic products, therapeutic products, consumer products, non-natural products, natural products, and fortified natural products, comprising, consisting essentially of, or consisting of ammonia oxidizing microorganisms are disclosed. These preparations, compositions, and/or formulations are disclosed herein for use in various applications, e.g., cosmetic and/or therapeutic applications. The preparations, compositions, and/or formulations may be administered in an effective amount for an intended use, e.g., a cosmetic or a therapeutic application. Preparations, compositions, and/or formulations comprising ammonia oxidizing microorganisms for various modes of administration to a subject are provided. Preparations, compositions, and/or formulations comprising ammonia oxidizing microorganisms for use in the treatment of various conditions and/or disorders in a subject are provided. Methods of treating a subject for various conditions and/or disorders via administration of ammonia oxidizing microorganisms are disclosed. Devices for use in administering ammonia oxidizing microorganisms to a subject are also provided.

Microbiology

In accordance with one or more embodiments, essentially any ammonia oxidizing microorganism (AOM) can be used or implemented. The ammonia oxidizing microorganisms may generally be autotrophic. The ammonia oxidizing microorganisms may generate nitrite and/or nitric oxide from ammonia.

Properties of autotrophic ammonia oxidizing bacteria (AOB), for example, are well described by Whitlock in U.S. Patent No. 7,820,420. Since that filing, the class of autotrophic microorganisms that oxidize ammonia for ATP production has been expanded to encompass ammonia oxidizing archaea (AOA), and archaea have been moved out of the class of bacteria and into their own distinct class. For the purposes of this disclosure, any and all autotrophic ammonia oxidizing microorganisms that share the properties of oxidation of ammonia to generate ATP can be implemented. AOM, including both AOB and AOA, share the necessary


properties of oxidation of ammonia into NO and nitrite and all known AOM lack capacity for virulence because of their inability to use organic substrates for ATP generation. Bacteria can utilize ammonia at higher concentrations, while archaea can utilize ammonia at lower concentrations. Physiological levels of ammonia are within the range that both bacteria (AOB) and archaea (AOA) can utilize. Any reference specifically to ammonia oxidizing bacteria throughout this disclosure should be considered equally applicable to any ammonia oxidizing microorganism, e.g., any ammonia oxidizing archaea, and these terms may all be used interchangeably herein.

Ammonia oxidizing bacteria (AOB) are ubiquitous Gram-negative obligate bacteria with a unique capacity to generate energy exclusively from the conversion of ammonia to nitrite. In some embodiments, ammonia oxidizing bacteria (AOB) of the genus *Nitrosomonas* are Gram-negative obligate autotrophic (chemolithoautotrophic) bacteria with a unique capacity to generate nitrite and nitric oxide exclusively from ammonia as an energy source. They are widely present both in soil and water environments and are essential components of environmental nitrification processes. These bacteria have beneficial properties, e.g., in connection with various cosmetic and therapeutic uses, in accordance with one or more embodiments described herein. Without wishing to be bound to any particular theory, due to the roles of nitrite and nitric oxide as important components of several physiological functions, such as vasodilation, inflammation and wound healing, these bacteria may have various beneficial properties for both healthy and immunopathological conditions. These bacteria are safe for use in humans because they are slow-growing, cannot grow on organic carbon sources, may be sensitive to soaps and antibiotics, and have never been associated with any disease or infection in animals or humans.

Ammonia oxidizing microorganisms generate coenzyme Q 8 (CoQ8) as a byproduct of the process by which they generate nitrite and nitric oxide. CoQ8 is a coenzyme Q having 8 carbons in its isoprenoid side chain. Without wishing to be bound to any particular theory, due to the role of coenzyme Q as an important component of several cell functions, such as mediating cell signaling and preventing cell death (anti-aging), these microorganisms' beneficial properties may further be enhanced by their specific ability to generate CoQ8.

In some embodiments, ammonia oxidizing bacteria may catalyze the following reactions.

At a neutral pH level, ammonia generated from ammonium around neutral pH conditions is the substrate of the initial reaction. The conversion of ammonia to nitrite takes place in two steps catalyzed respectively by ammonia monooxygenase (AMO) and hydroxylamine oxidoreductase (HAO), as follows:



In some instances, reaction B is reported as follows, to indicate nitrous acid (HNO₂) formation at low pH:



10 In certain embodiments, NH₄⁺ and NH₃ may be used interchangeably throughout the disclosure.

Examples of ammonia oxidizing bacteria include *Nitrosomonas eutropha* strains, e.g., D23 and C91 as discussed herein, and other bacteria in the genera *Nitrosomonas*, *Nitrosococcus*, *Nitrospira*, *Nitrosocystis*, *Nitrosolobus*, and *Nitrosovibrio*. D23 *Nitrosomonas eutropha* strain
15 refers to the strain, designated AOB D23-100, deposited with the American Tissue Culture Collection (ATCC) (10801 University Blvd., Manassas, VA, USA) on April 8, 2014 having accession number PTA-121157. The nucleic acid sequence(s), e.g., genome sequence, of accession number PTA-121157 are hereby incorporated herein by reference in their entireties for all purposes. “AOB D23-100” may also be referred to as D23 or B244 throughout this
20 disclosure.

Examples of ammonia oxidizing archaea include archaea in the genera *Methanobrevibacter*, *Methanosphaera*, *Methanosarcina*, *Nitrosocaldus*, *Nitrosopumilus*, and *Nitrososphaera* (e.g. *Nitrososphaera viennensis*, *Nitrososphaera gargensis*). Different phylotypes of archaea, e.g., methanogens and halophilic archaeon, may be included in the
25 preparations disclosed herein. Examples of archaea further include archaea in the lineages of phyla Euryarchaeota (e.g. *Methanosarcina*), Crenarchaeota, Aigarchaeota, and Thaumarchaeota (e.g. *Giganthauma karukerense*, *Giganthauma insulaporcus*, *Caldiarchaeum subterraneum*, *Cenarchaeum symbiosum*).

Each and every nucleic acid sequence and amino acid sequence disclosed in International
30 (PCT) Patent Application Publication No. WO2015/160911 (International (PCT) Patent

Application Serial No. PCT/US2015/025909 as filed on April 15, 2015), is hereby incorporated herein by reference in its entirety for all purposes. Likewise, any ammonia oxidizing bacteria disclosed in International (PCT) Patent Application Publication No. WO2015/160911 (International (PCT) Patent Application Serial No. PCT/US2015/025909 as filed on April 15, 2015), is also hereby incorporated herein by reference in its entirety for all purposes. In certain embodiments, the ammonia oxidizing microorganism is a strain as described therein.

In accordance with one or more embodiments, ammonia oxidizing microorganisms may exist in several metabolic states, e.g. growth state, storage state, and/or polyphosphate loading state.

In accordance with one or more embodiments, ammonia oxidizing microorganisms may have desirable properties, e.g., optimized properties, such as the ability to suppress growth of pathogenic bacteria, and an enhanced ability to produce nitric oxide and nitric oxide precursors.

Optimized *Nitrosomonas eutropha* (*N. eutropha*), as that term is used herein, refers to an *N. eutropha* having an optimized growth rate; an optimized NH_4^+ oxidation rate; and/or optimized resistance to NH_4^+ . In an embodiment it differs from naturally occurring *N. eutropha* by at least one nucleotide, e.g., a nucleotide in a gene selected from ammonia monooxygenase, hydroxylamine oxidoreductase, cytochrome c554, and cytochrome c_{M552} . The difference can arise, e.g., through selection of spontaneously arising mutation, induced mutation, or directed genetic engineering, of the *N. eutropha*. In an embodiment it differs from a naturally occurring *N. eutropha* in that it has a constellation of alleles, not present together in nature. These differences may provide for one or more of a treatment or prevention of a disease or condition, such as but not limited to one associated with low nitrite levels.

Any ammonia oxidizing bacteria, e.g., *N. eutropha*, for example *N. eutropha* referred to as “D23”, also known as “B244” or “AOB D23-100” may have several of the above-described properties. Any ammonia oxidizing archaea (AOA) may also have several of the above-described properties.

The AOBs contemplated in this disclosure may comprise mutations relative to wild-type AOBs. These mutations may, e.g., occur spontaneously, be introduced by random mutagenesis, or be introduced by targeted mutagenesis. For instance, the AOBs may lack one or more genes or regulatory DNA sequences that wild-type AOBs typically comprise. The AOBs may also

comprise point mutations, substitutions, insertions, deletions, and/or rearrangements relative to the sequenced strain or a wild-type strain. The AOBs may be a purified preparation of optimized AOBs.

In certain embodiments, the AOBs are transgenic. For instance, it may comprise one or
5 more genes or regulatory DNA sequences that wild-type ammonia oxidizing bacteria lacks. More particularly, the ammonia oxidizing bacteria may comprise, for instance, a reporter gene, a selective marker, a gene encoding an enzyme, or a promoter (including an inducible or repressible promoter). In some embodiments the additional gene or regulatory DNA sequence is integrated into the bacterial chromosome; in some embodiments the additional gene or regulatory
10 DNA sequence is situated on a plasmid.

In some embodiments, the AOBs differ by at least one nucleotide from naturally occurring bacteria. For instance, the AOBs may differ from naturally occurring bacteria in a gene or protein that is part of a relevant pathway, *e.g.*, an ammonia metabolism pathway, a urea metabolism pathway, or a pathway for producing nitric oxide or nitric oxide precursors. More
15 particularly, the AOBs may comprise a mutation that elevates activity of the pathway, *e.g.*, by increasing levels or activity of an element of that pathway.

The above-mentioned mutations can be introduced using any suitable technique. Numerous methods are known for introducing mutations into a given position. For instance, one could use site-directed mutagenesis, oligonucleotide-directed mutagenesis, or site-specific
20 mutagenesis. Non-limiting examples of specific mutagenesis protocols are described in, *e.g.*, Mutagenesis, pp. 13.1-13.105 (Sambrook and Russell, eds., Molecular Cloning A Laboratory Manual, Vol. 3, 3.sup.rd ed. 2001). In addition, non-limiting examples of well-characterized mutagenesis protocols available from commercial vendors include, without limitation, Altered
25 Sites.RTM. II in vitro Mutagenesis Systems (Promega Corp., Madison, Wis.); Erase-a-Base.RTM. System (Promega, Madison, Wis.); GeneTailor.TM. Site-Directed Mutagenesis System (Invitrogen, Inc., Carlsbad, Calif.); QuikChange.RTM. II Site-Directed Mutagenesis Kits (Stratagene, La Jolla, Calif.); and Transformer.TM. Site-Directed Mutagenesis Kit (BD-Clontech, Mountain View, Calif.).

In certain embodiments of the disclosure, the ammonia oxidizing microorganisms may be axenic. The preparation (formulation or composition) of ammonia oxidizing microorganisms may comprise, consist essentially of, or consist of axenic ammonia oxidizing microorganisms.

The ammonia oxidizing bacteria of this disclosure may be from a genus selected from the
5 group consisting of *Nitrosomonas*, *Nitrosococcus*, *Nitrosospria*, *Nitrosocystis*, *Nitrosolobus*, *Nitrosovibrio*, and combinations thereof.

This disclosure provides, *inter alia*, *N. eutropha* strain D23, a unique, *e.g.*, optimized strain of ammonia oxidizing bacteria that can increase production of nitric oxide and nitric oxide precursors on a surface of a subject, *e.g.*, a human subject. This disclosure also provides
10 methods of administering and using the bacteria and preparations, compositions, formulations, and products, comprising the bacteria.

In embodiments, the ammonia oxidizing bacteria, *e.g.*, *N. eutropha* is non-naturally occurring. For instance, it may have accumulated desirable mutations during a period of selection. In other embodiments, desirable mutations may be introduced by an experimenter. In
15 some embodiments, the *N. eutropha* may be a purified preparation, and may be an optimized *N. eutropha*.

In preferred embodiments, the *N. eutropha* strain is autotrophic and so incapable of causing infection. A preferred strain utilizes urea as well as ammonia, so that hydrolysis of the urea in sweat would not be necessary prior to absorption and utilization by the bacteria. Also, in
20 order to grow at low pH, the bacteria may either absorb NH_4^+ ions or urea. The selected strain should also be capable of living on the external skin of a subject, *e.g.*, a human, and be tolerant of conditions there.

Although this disclosure refers to *N. eutropha* strain D23 in detail, the preparations, methods, compositions, treatments, formulas and products may be used with one or more of: one
25 or more other strains of *N. eutropha*, one or more other species of *Nitrosomonas*, and one or more other ammonia oxidizing microorganism, *e.g.* ammonia oxidizing bacteria or other ammonia oxidizing archaea.

In certain embodiments, a bacterium with the above-mentioned sequence characteristics has one or more of (1) an optimized growth rate as measured by doubling time, (2) an optimized
30 growth rate as measured by OD600, (3) an optimized NH_4^+ oxidation rate, (4) an optimized

resistance to NH_4^+ , and (4) an optimized resistance to NO_2^- . Particular nonlimiting sub-combinations of these properties are specified in the following paragraph.

In some embodiments, the ammonia oxidizing bacteria, e.g., the *N. eutropha* described herein, or an axenic composition thereof, has one or more of: (1) an optimized growth rate as measured by doubling time, (2) an optimized growth rate as measured by OD600, (3) an optimized NH_4^+ oxidation rate, (4) an optimized resistance to, NH_4^+ , and (4) an optimized resistance to, NO_2^- . For instance, the bacterium may have properties (1) and (2); (2) and (3); (3) and (4); or (4) and (5) from the list at the beginning of this paragraph. As another example, the bacterium may have properties (1), (2), and (3); (1), (2), and (4); (1), (2), and (5); (1), (3), and (4); (1), (3), and (5); (1), (4), and (5); (2), (3), and (4); (2), (3), and (5), or (3), (4), and (5) from the list at the beginning of this paragraph. As a further example, the bacterium may have properties (1), (2), (3), and (4); (1), (2), (3), and (5); (1), (2), (4), and (5); (1), (3), (4), and (5); or (2), (3), (4), and (5) from the list at the beginning of this paragraph. In some embodiments, the bacterium has properties (1), (2), (3), (4), and (5) from the list at the beginning of this paragraph.

In certain embodiments, the *N. eutropha* strain comprises a nucleic acid sequence, e.g., a genome, that hybridizes to SEQ ID NO: 1 of International (PCT) Patent Application Publication No. WO2015160911 (International (PCT) Patent Application Serial No. PCT/US2015/025909 filed on April 15, 2015), or to the genome of the D23 strain deposited in the form of 25 vials with the ATCC patent depository on April 8, 2014, designated AOB D23-100, under accession number PTA-121157, or their complements, under low stringency, medium stringency, high stringency, or very high stringency, or other hybridization condition.

The D23 strain is not believed to be a product of nature, but rather has acquired certain mutations and characteristics during an extended period of culture and selection in the laboratory. For instance, D23 has an ability to grow in conditions of greater than about 200 or 250 mM NH_4^+ for more than 24 hours.

In some embodiments, the *N. eutropha* disclosed herein differ from naturally occurring bacteria in the abundance of siderophores. For instance, the *N. eutropha* may have elevated or reduced levels of siderophores compared to *N. eutropha* C91. Generally, siderophores are secreted iron-chelating compounds that help bacteria scavenge iron from their environment. Some siderophores are peptides, and others are small organic molecules.

The practice of the present invention may employ, unless otherwise indicated, conventional methods of immunology, molecular biology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al. *Molecular Cloning: A Laboratory Manual* (Current Edition); and Current
 5 *Protocols in Molecular Biology* (F.M. Ausubel, et al. eds., current edition).

Select Definitions

An ammonia oxidizing microorganism, e.g., ammonia oxidizing bacteria, refers to a microorganism capable of oxidizing ammonia or ammonium to nitrite at a rate, e.g., a substantial
 10 rate, e.g., a pre-determined rate. The rate, e.g., a pre-determined rate, may refer to the conversion of ammonium ions (NH_4^+) (e.g., at about 200 mM) to nitrite (NO_2^-), for example, as determined or measured in an *in vitro* assay or when administered to a subject, e.g., a human. The rate may be a conversion at a rate of at least about 1 picomole per minute per mg protein, 0.01, 0.1, 1, 10, 25, 50, 75, 125, or 150 nanomoles NO_2^- per minute per mg protein, e.g., about
 15 0.01-1, 0.1-50, 50-100, 100-150, 75-175, 75-125, 100-125, 125-150, or 125-175 nanomoles/minute/mg protein, e.g., about 125 nanomoles NO_2^- per minute per mg protein for a continuous culture, for example having an OD of about 0.5. The rate of conversion may be between about 1 picomole per minute per mg protein to about 1 millimole per minute per mg protein. The rate of conversion may be at most about 1 mole NO_2^- per minute per mg protein,
 20 e.g. at least about, about, or at most about 1 decimole, 1 centimole, 1 millimole, or 1 micromole NO_2^- per minute per mg protein.

As used herein, “axenic” refers to a composition comprising an organism that is substantially free of other organisms. For example, an axenic culture of ammonia oxidizing bacteria is a culture that is substantially free of organisms other than ammonia oxidizing bacteria.
 25 For example, an axenic culture of *N. eutropha* is a culture that is substantially free of organisms other than *N. eutropha*. In some embodiments, “substantially free” denotes undetectable by a method used to detect other organisms, e.g., plating the culture and examining colony morphology, or PCR for a conserved gene such as 16S RNA. An axenic composition may comprise elements that are not organisms, e.g., it may comprise nutrients or excipients. Any
 30 embodiment, preparation, composition, or formulation of ammonia oxidizing bacteria discussed

herein may comprise, consist essentially of, or consist of optionally axenic ammonia oxidizing bacteria.

Throughout this disclosure, formulation may refer to a composition or preparation or product.

5 As used herein, an “autotroph”, *e.g.*, an autotrophic bacterium, is any organism capable of self-nourishment by using inorganic materials as a source of nutrients and using photosynthesis or chemosynthesis as a source of energy. Autotrophic bacteria may synthesize organic compounds from carbon dioxide and ATP derived from other sources, oxidation of ammonia to nitrite, oxidation of hydrogen sulfide, and oxidation of Fe^{2+} to Fe^{3+} . Autotrophic
10 bacteria of the present disclosure are incapable of causing infection.

Administered “in combination,” as used herein, means that two (or more) different treatments are delivered to the subject during the course of the subject’s affliction with the disorder, *e.g.*, the two or more treatments are delivered after the subject has been diagnosed with the disorder and before the disorder has been cured or eliminated. In some embodiments, the
15 delivery of one treatment is still occurring when the delivery of the second begins, so that there is overlap. This is sometimes referred to herein as “simultaneous” or “concomitant” or “concurrent delivery”. In other embodiments, the delivery of one treatment ends before the delivery of the other treatment begins. This is sometimes referred to herein as “successive” or “sequential delivery.” In embodiments of either case, the treatment is more effective because of combined
20 administration. For example, the second treatment is a more effective, *e.g.*, an equivalent effect is seen with less of the second treatment, or the second treatment reduces symptoms to a greater extent, than would be seen if the second treatment were administered in the absence of the first treatment, or the analogous situation is seen with the first treatment. In some embodiments, delivery is such that the reduction in a symptom, or other parameter related to the disorder is
25 greater than what would be observed with one treatment delivered in the absence of the other. The effect of the two treatments can be partially additive, wholly additive, or greater than additive (*i.e.*, synergistic). The delivery can be such that an effect of the first treatment delivered is still detectable when the second is delivered. In some embodiments, one or more treatment may be delivered prior to diagnosis of the patient with the disorder.

The term “isolated,” as used herein, refers to material that is removed from its original or native environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated by human intervention from some or all of the co-existing materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of the environment in which it is found in nature.

As used herein, the term “optimized growth rate” refers to one or more of: a doubling time of less than about 4, 5, 6, 7, 8, 9, or 10 hours when cultured under batch conditions as described herein in Example 2; a doubling time of less than about 16, 18, 20, 22, 24, or 26 hours, when grown under chemostat conditions as described herein in Example 2; or growing from an OD600 of about 0.15 to at least about 0.3, 0.4, 0.5, 0.6, 0.7, or 0.8 over about 1 or 2 days. In an embodiment, optimized growth rate is one having a doubling time that it is at least 10, 20, 30, 40, or 50% shorter than that of a naturally occurring *N. eutropha*.

As used herein, “optimized NH_4^+ oxidation rate” refers to a rate of at least about 50, 75, 125, or 150 micromoles per minute of converting NH_3 or NH_4^+ into NO_2^- . For instance, the rate may be at least about 50, 75, 125, or 150 micromoles per minute of converting NH_4^+ (*e.g.*, at about 200 mM) to NO_2^- . In an embodiment, an optimized NH_4^+ oxidation rate is one in which NH_3 or NH_4^+ is converted into NO_2^- at least 10, 20, 30, 40, or 50% more rapidly than is seen with a naturally occurring *N. eutropha*.

As used herein, “optimized resistance to NH_4^+ ” refers to an ability to grow in conditions of greater than 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, or 300 mM NH_3 or NH_4^+ for at least about 24 or 48 hours. In an embodiment, an optimized resistance to NH_4^+ refers to the ability to grow at least 10, 20, 30, 40, or 50% more rapidly, or at least 10, 20, 30, 40, or 50% longer, in the presence of a selected concentration of NH_3 or NH_4^+ than can a naturally occurring *N. eutropha*.

As used herein, “transgenic” means comprising one or more exogenous portions of DNA. The exogenous DNA is derived from another organism, *e.g.*, another bacterium, a bacteriophage, an animal, or a plant.

As used herein, treatment of a disease or condition refers to reducing the severity or frequency of at least one symptom of that disease or condition, compared to a similar but untreated patient. Treatment can also refer to halting, slowing, or reversing the progression of a disease or condition, compared to a similar but untreated patient. Treatment may comprise
5 addressing the root cause of the disease and/or one or more symptoms.

As used herein a therapeutically effective amount refers to a dose sufficient to prevent advancement, or to cause regression of a disease or condition, or which is capable of relieving a symptom of a disease or condition, or which is capable of achieving a desired result. A therapeutically effective dose can be measured, for example, as a number of bacteria or number
10 of viable bacteria (*e.g.*, in CFUs) or a mass of bacteria (*e.g.*, in milligrams, grams, or kilograms), or a volume of bacteria (*e.g.*, in mm³).

As used herein, the term “viability” refers to the autotrophic bacteria’s, *e.g.*, ammonia oxidizing bacteria’s, ability to oxidize ammonia, ammonium, or urea to nitrite at a pre-determined rate. In some embodiments, the rate refers to the conversion of ammonium ions
15 (NH₄⁺) (*e.g.*, at about 200 mM) to nitrite (NO₂⁻) at a rate of at least about 1 picomole, 0.01, 0.1, 1, 10, 25, 50, 75, 125, or 150 nanomoles NO₂⁻ per minute, *e.g.*, about 0.01-1, 0.1-50, 50-100, 100-150, 75-175, 75-125, 100-125, 125-150, or 125-175 nanomoles/minute, *e.g.*, about 125 nanomoles NO₂⁻ per minute. The rate of conversion may be at most about 1 mole NO₂⁻ per minute, *e.g.* at least about, about, or at most about 1 decimole, 1 centimole, 1 millimole, or 1
20 micromole NO₂⁻ per minute. Viable ammonia oxidizing microorganisms may generally comprise culturable AOMs or AOMs that are otherwise able to generate NO, nitrate, or nitrite.

As used herein, a “subject” may include an animal, a mammal, a human, a non-human animal, a livestock animal, or a companion animal. The term “subject” is intended to include human and non-human animals, for example, vertebrates, large animals, and primates. In certain
25 embodiments, the subject is a mammalian subject, and in particular embodiments, the subject is a human subject. Although applications with humans are clearly foreseen, veterinary applications, for example, with non-human animals, are also envisaged herein. The term “non-human animals” of the disclosure includes all vertebrates, for example, non-mammals (such as birds, for example, chickens; amphibians; reptiles) and mammals, such as non-human primates,

domesticated, and agriculturally useful animals, for example, sheep, dog, cat, cow, pig, rat, among others.

“Microbiome” refers to a population, e.g., one or more microorganisms that live on a surface of a subject, e.g., in the gut, mouth, skin, and/or elsewhere in a subject. The population
5 may have one or more beneficial functions and/or benefits, relevant to supporting the life of a subject.

“Biome-friendly” refers to something, e.g., a product, e.g., a cosmetic product, e.g., a finished cosmetic product that may allow for minimal disruption of a microbiome of a subject. For example, biome-friendly refers to a product that may be applied to a subject that may allow
10 the microbiome at the point of application to be maintained, minimally disrupted, and/or able to return to the microbiome after a period of time after application of the product. In embodiments, biome-friendly may refer to ammonia oxidizing microorganism-friendly, e.g., ammonia oxidizing bacteria- friendly in that the product may allow for minimal disruption of the ammonia oxidizing bacteria of a subject. In embodiments, “biome-friendly” may be referred to
15 as “biome-compatible.”

A “natural product” is or may comprise a product that may be at least partially derived from nature. It may be anything or comprise anything produced by a living organism, and may include organisms themselves. Natural products may include or comprise an entire organism, and part of an organism (e.g., a leaf of a plant), an extract from an organism, an organic
20 compound from an organism, a purified organic compound from an organism. Natural products may be or comprise organic substances found and cells, including primary metabolites (amino acids, carbohydrates, and nucleic acids) and secondary metabolites (organic compounds found in a limited range of species, e.g., polyketides, fatty acids, terpenoids, steroids, phenylpropanoids, alkaloids, specialized amino acids and peptides, specialized carbohydrates). Natural products
25 may be or comprise polymeric organic materials such as cellulose, lignin, and proteins.

As used herein, “presence” or “level” may refer to a qualitative or quantitative amount of a component, e.g., any one or more of an ammonia oxidizing microorganisms, ammonia, ammonium ions, urea, nitrite, or nitric oxide. The presence or level may include a zero value or a lack of presence of a component.

As used herein, the term “surfactant”, includes compounds that may lower the surface tension, or interfacial tension, between two liquids or between a liquid and a solid. Surfactants may act as detergents, wetting agents, emulsifiers, foaming agents, and dispersants. Surfactants may include one or more of the following, alone, or in combination with those listed, or other

5 surfactants or surfactant-like compounds: cocamidopropyl betaine (ColaTeric COAB), polyethylene sorbitol ester (*e.g.*, Tween 80), ethoxylated lauryl alcohol (RhodaSurf 6 NAT), sodium laureth sulfate/lauryl glucoside/cocamidopropyl betaine (Plantapon 611 L UP), sodium laureth sulfate (*e.g.*, RhodaPex ESB 70 NAT), alkyl polyglucoside (*e.g.*, Plantaren 2000 N UP), sodium laureth sulfate (Plantaren 200), Dr. Bronner’s Castile soap, Dr. Bronner’s baby soap,

10 Lauramine oxide (ColaLux Lo), sodium dodecyl sulfate (SDS), polysulfonate alkyl polyglucoside (PolySufanate 160 P), sodium lauryl sulfate (Stepanol-WA Extra K), and combinations thereof. Dr. Bronner’s Castile soap and baby soap comprises water, organic coconut oil, potassium hydroxide, organic olive oil, organic fair deal hemp oil, organic jojoba oil, citric acid, and tocopherol. Surfactants may include Sodium Laurylglucosides

15 Hydroxypropylsulfonate (Suga®nate 160NC), lauramidopropyl betaine (Cola®Teric LMB); Cocamidopropyl hydroxysultaine (Cola®Teric CBS); disodium cocoamphodiacetate (Cola®Teric CDCX-LV); sodium laurylglucosides hydroxypropyl phosphate (Suga®Fax D12). Surfactants may include sodium lauroyl methyl isethionate (Iselux® LQ-CLR-SB); sodium methyl cocoyl taurate (Pureact WS Conc.); Aqua (and) Sodium Lauroyl Methyl Isethionate (and)

20 Cocamidopropyl Betaine (and) Sodium Cocoyl Isethionate (and) Sodium Methyl Oleoyl Taurate (Iselux ®SFS-SB). Other surfactants are contemplated by this disclosure.

Preparations, Compositions, Formulations, and Products Comprising Ammonia Oxidizing Microorganisms

25 The present disclosure provides, *inter alia*, compositions comprising ammonia oxidizing microorganisms, preparations, *e.g.*, purified and/or optimized preparations, comprising AOM, formulations comprising AOM, and various products comprising AOM, *e.g.*, a natural product, a non-natural product, a fortified natural product, a consumer product, a therapeutic product, or a cosmetic product. The terms preparation, composition, formulation, and product may be used

30 interchangeably herein.

Any embodiment, preparation, composition, formulation, or product of ammonia oxidizing microorganisms discussed herein may comprise, consist essentially of, or consist of (optionally axenic) ammonia oxidizing microorganisms, e.g., live ammonia oxidizing microorganisms.

5 The preparation may comprise or be supplemented with a product or byproduct of an ammonia oxidizing microorganism, e.g., nitrite, nitrate, nitric oxide, CoQ8. In at least some embodiments, the preparation may comprise or be supplemented with a composition that promotes growth or metabolism of ammonia oxidizing microorganisms, promotes production of products or byproducts of ammonia oxidizing microorganisms, promotes urease activity, or has a synergistic effect with ammonia oxidizing microorganisms, e.g., ammonia, ammonium salts, 10 urea, and urease. For instance, the preparation may be supplemented with one or more of NO, nitrite, nitrate, CoQ8, ammonia, ammonium salts, urea, and urease. The supplement may be comprised in the same formulation as the ammonia oxidizing microorganisms or in a separate formulation for concurrent or combination administration. The supplement formulation may be prepared for delivery via any delivery mode, for example inhaled forms of NO, nitrite, or nitrate. 15 The preparation may comprise, *inter alia*, at least one of ammonia, ammonium salts, and urea. The preparation may comprise or be supplemented with an anti-inflammatory agent or a composition that provides an anti-inflammatory effect.

 The present disclosure provides for preparations comprising ammonia oxidizing 20 microorganisms for cosmetic use.

 The present disclosure provides for preparations comprising ammonia oxidizing microorganisms for therapeutic use.

 In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing 25 microorganisms sufficient to have a desired cosmetic effect. The preparation may be formulated and/or delivered to impart the desired cosmetic effect locally and/or systemically.

 In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms sufficient to have a desired therapeutic effect, e.g., to at least partially treat a

condition or disease. The preparation may be formulated and/or delivered to impart the desired therapeutic effect locally and/or systemically.

In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms sufficient to alter, e.g., reduce or increase, an amount, concentration or proportion of a bacterium, or genus of bacteria in a subject. The bacteria may be non-pathogenic or pathogenic, or potentially pathogenic.

In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms sufficient to modulate a microbiome associated with a subject.

In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms sufficient to deliver NO to a subject. A preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms such that when administered, the preparation modulates, changes, or alters a level of nitrite or NO at a target tissue or in circulation. For instance, a preparation of ammonia oxidizing microorganisms may comprise a concentration or amount, e.g., an effective amount, of ammonia oxidizing microorganisms such that when administered, the preparation results in an increased level of nitrite or NO at a target tissue or in circulation.

The present disclosure provides, *inter alia*, non-limiting compositions comprising ammonia oxidizing microorganisms, e.g., *N. eutropha*, e.g., a purified preparation of an optimized *N. eutropha*. In some embodiments, the *N. eutropha* in the compositions has at least one property selected from an optimized growth rate, an optimized NH_4^+ oxidation rate, and an optimized resistance to NH_4^+ .

In some aspects, the present disclosure provides compositions with a defined number of species. A composition may include only one type of species, e.g., one type of ammonia oxidizing microorganism. This disclosure also provides a composition having, e.g., *N. eutropha* and one other type of organism, and no other types of organism. In other examples, the composition has, e.g., *N. eutropha* and 2, 3, 4, 5, 6, 7, 8, 9, or 10 other types of organism, and no other types of organism. The other type of organism in this composition may be, for instance, a

bacterium, such as an ammonia-oxidizing bacterium. Suitable ammonia-oxidizing microorganisms for this purpose include those in the genera *Nitrosomonas*, *Nitrosococcus*, *Nitrospira*, *Nitrosocystis*, *Nitrosolobus*, or *Nitrosovibrio*. Likewise, the composition may also include AOA.

5 In some embodiments, the composition comprising, e.g., *N. eutropha* provides conditions that support *N. eutropha* viability. For instance, the composition may promote *N. eutropha* growth and metabolism or may promote a dormant state (e.g., freezing) from which viable *N. eutropha* can be recovered. When the composition promotes growth or metabolism, it may contain water and/or nutrients that *N. eutropha* consumes, e.g., as ammonium, ammonia, urea,
 10 oxygen, carbon dioxide, or trace minerals. In some embodiments, the composition comprising ammonia oxidizing microorganisms provides conditions that support ammonia oxidizing microorganisms viability. For instance, the composition may promote ammonia oxidizing microorganisms growth and metabolism or may promote a dormant state (e.g., freezing) or storage state as described herein, from which viable ammonia oxidizing microorganisms can be
 15 recovered. When the composition promotes growth or metabolism, it may contain water and/or nutrients that ammonia oxidizing microorganisms consumes, e.g., as ammonium ions, ammonia, urea, oxygen, carbon dioxide, or trace minerals.

In some embodiments, one or more other organisms, for example, organisms besides ammonia oxidizing microorganisms may be included in the preparation of ammonia oxidizing
 20 microorganisms. For example, a community of organisms or an organism of the genus selected from the group consisting of *Lactobacillus*, *Streptococcus*, *Bifidobacter*, and combinations thereof, may be provided in the preparation of ammonia oxidizing microorganisms. In some embodiments, the preparation may be substantially free of other organisms.

Preparations of ammonia oxidizing microorganisms may comprise between about
 25 between about 10^3 to about 10^{14} CFU/ml. In some embodiments, the preparation of ammonia oxidizing microorganisms may comprise at least about or greater than about 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 2×10^{11} , 5×10^{11} , 10^{12} , 2×10^{12} , 5×10^{12} , 10^{13} , 2×10^{13} , 5×10^{13} , or 10^{14} ; or about 10^3 - 10^4 , 10^4 - 10^5 , 10^6 - 10^7 , 10^7 - 10^8 , 10^8 - 10^9 , 10^9 - 10^{10} , 10^{10} - 10^{11} , 10^{11} - 10^{12} , 10^{12} - 10^{13} , or 10^{13} - 10^{14} CFU/ml.

In some embodiments, a preparation of ammonia oxidizing microorganisms may comprise between about 1×10^9 to about 10×10^9 CFU/ml. In some embodiments, an administered dose of the preparation may comprise about 3×10^{10} CFU, *e.g.*, 3×10^{10} CFU per day. In some embodiments, an administered dose of the preparation may comprise about 1×10^9 to about 10×10^9 CFU per day, *e.g.*, about 1×10^9 to about 10×10^9 CFU per day. In some embodiments, an administered dose of the preparation may comprise about 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 2×10^{11} , 5×10^{11} , 10^{12} , 2×10^{12} , 5×10^{12} , 10^{13} , 2×10^{13} , 5×10^{13} , or 10^{14} ; or about 10^3 - 10^4 , 10^4 - 10^5 , 10^6 - 10^7 , 10^7 - 10^8 , 10^8 - 10^9 , 10^9 - 10^{10} , 10^{10} - 10^{11} , 10^{11} - 10^{12} , 10^{12} - 10^{13} , or 10^{13} - 10^{14} CFU per administration or per day.

In some embodiments, an administered dose of the preparation may comprise at least about 7×10^{10} CFU, *e.g.*, 21×10^{10} CFU per week. In some embodiments, an administered dose of the preparation may comprise about 1×10^9 to about 10×10^9 CFU per week, *e.g.*, about 1×10^9 to about 10×10^9 CFU per week. In some embodiments, an administered dose of the preparation may comprise about or greater than about 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 2×10^{11} , 5×10^{11} , 10^{12} , 2×10^{12} , 5×10^{12} , 10^{13} , 2×10^{13} , 5×10^{13} , or 10^{14} ; or about 10^3 - 10^4 , 10^4 - 10^5 , 10^6 - 10^7 , 10^7 - 10^8 , 10^8 - 10^9 , 10^9 - 10^{10} , 10^{10} - 10^{11} , 10^{11} - 10^{12} , 10^{12} - 10^{13} , or 10^{13} - 10^{14} CFU per week.

In some embodiments, an administered dose of the preparation may comprise at least about 30×10^{10} CFU, *e.g.*, 90×10^{10} CFU per month. In some embodiments, an administered dose of the preparation may comprise about 1×10^9 to about 10×10^9 CFU per month, *e.g.*, about 1×10^9 to about 10×10^9 CFU per month. In some embodiments, an administered dose of the preparation may comprise about or greater than about 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 2×10^{11} , 5×10^{11} , 10^{12} , 2×10^{12} , 5×10^{12} , 10^{13} , 2×10^{13} , 5×10^{13} , or 10^{14} ; or about 10^3 - 10^4 , 10^4 - 10^5 , 10^6 - 10^7 , 10^7 - 10^8 , 10^8 - 10^9 , 10^9 - 10^{10} , 10^{10} - 10^{11} , 10^{11} - 10^{12} , 10^{12} - 10^{13} , or 10^{13} - 10^{14} CFU per month.

In some embodiments, the preparation of ammonia oxidizing microorganisms may comprise between about 0.1 milligrams (mg) and about 1000 mg of ammonia oxidizing microorganisms. In certain aspects, the preparation may comprise between about 50 mg and about 1000 mg of ammonia oxidizing microorganisms. The preparation may comprise between about 0.1-0.5 mg, 0.2-0.7 mg, 0.5-1.0 mg, 0.5-2 mg, 0.5-5 mg, 2.5-5 mg, 2.5-7.0 mg, 5.0-10 mg,

7.5-15 mg, 10-15 mg, 15-20 mg, 15-25 mg, 20-30 mg, 25-50 mg, 25-75 mg, 50-75 mg, 50-100 mg, 75-100 mg, 100-200 mg, 200-300 mg, 300-400 mg, 400-500 mg, 500-600 mg, 600-700 mg, 700-800 mg, 800-900 mg, 900-1000 mg, 100-250 mg, 250-500 mg, 100-500 mg, 500-750 mg, 750-1000 mg, or 500-1000 mg.

5 Advantageously, a formulation may have a pH level that promotes AOM, *e.g.*, *N. eutropha* viability, *e.g.*, metabolic activity. Urea would hydrolyze to ammonia and would raise the pH to 7 to 8. AOB are very active at this pH range and would lower the pH to about 6 where the NH₃ converts to ammonium and is unavailable. Lower pH levels, *e.g.* about pH 4, are also acceptable.

10 The ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* may be combined with one or more pharmaceutically or cosmetically acceptable excipients. In some embodiments, “pharmaceutically acceptable excipient” refers to a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, solvent, or encapsulating material. In some embodiments, each excipient is “pharmaceutically acceptable” in the sense of
15 being compatible with the other ingredients of a pharmaceutical formulation, and suitable for use in contact with the tissue or organ of humans and animals without excessive toxicity, irritation, allergic response, immunogenicity, or other problems or complications, commensurate with a reasonable benefit/risk ratio. See, Remington: The Science and Practice of Pharmacy, 21st ed.; Lippincott Williams & Wilkins: Philadelphia, Pa., 2005; Handbook of Pharmaceutical
20 Excipients, 6th ed.; Rowe et al., Eds.; The Pharmaceutical Press and the American Pharmaceutical Association: 2009; Handbook of Pharmaceutical Additives, 3rd ed.; Ash and Ash Eds.; Gower Publishing Company: 2007; Pharmaceutical Preformulation and Formulation, 2nd ed.; Gibson Ed.; CRC Press LLC: Boca Raton, Fla., 2009.

 In some embodiments, a cosmetically acceptable excipient refers to a cosmetically
25 acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, solvent, or encapsulating material. In some embodiments, each excipient is cosmetically acceptable in the sense of being compatible with the other ingredients of a cosmetic formulation, and suitable for use in contact with the tissue or organ of humans and animals without excessive toxicity, irritation, allergic response, immunogenicity, or other problems or complications, commensurate
30 with a reasonable benefit/risk ratio.

While it is possible for the active ingredient, *e.g.*, ammonia oxidizing microorganisms, *e.g.*, *N. eutropha*, to be administered alone, in many embodiments it is present in a pharmaceutical formulation or composition. Accordingly, this disclosure provides a pharmaceutical formulation comprising ammonia oxidizing microorganisms, for example, *N. eutropha* and a pharmaceutically acceptable excipient. Pharmaceutical compositions may take the form of a pharmaceutical formulation as described below.

In accordance with one or more embodiments, a preparation of ammonia oxidizing microorganisms may be formulated in order to facilitate a desired delivery mechanism or mode of administration thereof. The formulations, *e.g.*, pharmaceutical or cosmetic formulations, described herein include those suitable for, *e.g.*, oral, enteral (including buccal, sublingual, sublabial, and rectal), parenteral (including subcutaneous, intradermal, intramuscular, intravenous, and intraarticular), inhalation (including fine particle dusts or mists which may be generated by means of various types of metered doses, pressurized aerosols, nebulizers or insufflators, and including intranasally or via the lungs), intranasal, eye, ear, rectal, injection, urogenital, and topical (including dermal, transdermal, transmucosal, buccal, sublingual, and intraocular) administration, although the most suitable route may depend upon, for example, a condition or disorder of a recipient.

In accordance with one or more non-limiting embodiments, a preparation comprising ammonia oxidizing microorganisms may be administered to a subject, *e.g.*, for cosmetic or therapeutic purposes, as a solution, suspension, powder, liquid, drop, spray, aerosol, mist, emulsion, foam, cream, ointment, gel, hydrogel, resin, tablet, capsule, film, suppository, enema, douche, pessary, insert, patch, *e.g.*, transdermal patch, or implantable device, *e.g.*, stent, catheter, vaginal ring, or intrauterine device.

Devices configured to deliver a preparation comprising live ammonia oxidizing microorganisms via a desired mode of administration or otherwise via targeted delivery are also disclosed.

In accordance with one or more embodiments, the preparation may be formulated for targeted delivery to a subject, *e.g.*, to a target tissue, region, system, or organ of a subject. For example, the preparation may be formulated for delivery to the eye, ear, nose, urogenital system, respiratory system, or gastrointestinal system of the subject. In some embodiments, targeted

delivery may be based on a condition or disorder of a subject. For instance, formulation for targeted delivery may be based on a desired local or systemic effect to be achieved, e.g., a local or systemic therapeutic or cosmetic effect. In some embodiments, a target tissue, region, system, or organ of a subject may be selected for its association with a desired local or systemic effect.

5 The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods known in the art of pharmacy. Typically, methods include the step of bringing the active ingredient (e.g., ammonia oxidizing microorganisms, e.g., *N. eutropha*) into association with a pharmaceutical carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into
10 association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations may be presented as discrete units such as capsules, cachets or tablets, each containing a predetermined amount of, e.g., *N. eutropha*; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid
15 emulsion or a water-in-oil liquid emulsion. Formulations, e.g., solutions, aerosols, sprays, and mists, may be presented in multi-dosage form, e.g., packaged units including a predetermined number of dosages, or single dosage form, e.g., packaged units including a single dose. The active ingredient may also be presented as a bolus, electuary or paste. Various pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g.,
20 Remington's Pharmaceutical Sciences by E. W. Martin. See also Wang, Y. J. and Hanson, M. A., Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2 S, 1988.

The ammonia oxidizing microorganisms, e.g., *N. eutropha* compositions can, for example, be administered in a form suitable for immediate release or extended release. Suitable examples of sustained-release systems include suitable polymeric materials, for example semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules; suitable
25 hydrophobic materials, for example as an emulsion in an acceptable oil; or ion exchange resins. Sustained-release systems may be administered orally; rectally; parenterally; intracisternally; intravaginally; intraperitoneally; topically, for example as a powder, ointment, gel, drop or transdermal patch; buccally; or as a spray.

Preparations for administration can be suitably formulated to give controlled release of ammonia oxidizing microorganisms, *e.g.*, *N. eutropha*. For example, the pharmaceutical compositions may be in the form of particles comprising one or more of biodegradable polymers, polysaccharide jellifying and/or bioadhesive polymers, or amphiphilic polymers. These compositions exhibit certain biocompatibility features which allow a controlled release of an active substance. See U.S. Pat. No. 5,700,486.

Exemplary compositions include suspensions which can contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants, mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations can also include an excipient to aid mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (*e.g.*, Gantrez), and agents to control release such as polyacrylic copolymer (*e.g.* Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use. The surfactant may be a zwitterionic surfactant, a non-ionic surfactant, or an anionic surfactant.

Excipients, such as surfactants that may be used with embodiments of the present disclosure may include one or more of cocamidopropyl betaine (ColaTeric COAB), polyethylene sorbitol ester (*e.g.*, Tween 80), ethoxylated lauryl alcohol (RhodaSurf 6 NAT), sodium laureth sulfate/lauryl glucoside/cocamidopropyl betaine (Plantapon 611 L UP), sodium laureth sulfate (*e.g.*, RhodaPex ESB 70 NAT), alkyl polyglucoside (*e.g.*, Plantaren 2000 N UP), sodium laureth sulfate (Plantaren 200), Dr. Bronner's Castile soap, Dr. Bronner's Castile baby soap, Lauramine oxide (ColaLux Lo), sodium dodecyl sulfate (SDS), polysulfonate alkyl polyglucoside (PolySufanate 160 P), sodium lauryl sulfate (Stepanol-WA Extra K), and combinations thereof. Dr. Bronner's Castile soap and Dr. Bronner's baby soap comprises water, organic coconut oil, potassium hydroxide, organic olive oil, organic fair deal hemp oil, organic jojoba oil, citric acid, and tocopherol.

In some embodiments, surfactants may be used with ammonia oxidizing microorganisms in amounts that allow nitrite production to occur. In some embodiments, the preparation may have less than about 0.0001 % to about 10% of surfactant. In some embodiments, the preparation may have between about 0.1 % and about 10 % surfactant. In some embodiments, the concentration of surfactant used may be between about 0.0001% and about 10%. In some embodiments, the preparation may be substantially free of surfactant.

In some embodiments, the formulation, *e.g.*, preparation, may include other components that may enhance effectiveness of ammonia oxidizing microorganisms, delivery thereof, or enhance a treatment or indication.

In some embodiments, a chelator may be included in the preparation. A chelator may be a compound that may bind with another compound, *e.g.*, a metal. The chelator may provide assistance in removing an unwanted compound from an environment, or may act in a protective manner to reduce or eliminate contact of a particular compound with an environment, *e.g.*, ammonia oxidizing microorganisms, *e.g.* a preparation of ammonia oxidizing microorganisms, *e.g.*, an excipient. In some embodiments, the preparation may be substantially free of chelator.

Formulations may also contain anti-oxidants, buffers, bacteriostats that prevent the growth of undesired microorganisms, solutes, and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of a sterile liquid carrier, for example saline or water-for-injection, immediately prior to use. Extemporaneous solutions and suspensions may be prepared from powders, granules and tablets of the kind previously described. Exemplary compositions include solutions or suspensions which can contain, for example, suitable non-toxic, pharmaceutically acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid, or Cremaphor. An aqueous carrier may be, for example, an isotonic buffer solution at a pH of from about 3.0 to about 8.0, a pH of from about 3.5 to about 7.4, for example from 3.5 to 6.0, for example from 3.5 to about 5.0. Useful buffers include sodium

citrate-citric acid and sodium phosphate-phosphoric acid, and sodium acetate/acetic acid buffers. The composition in some embodiments does not include oxidizing agents.

Excipients that can be included are, for instance, proteins, such as human serum albumin or plasma preparations. If desired, the pharmaceutical composition may also contain minor
5 amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate. In some embodiments, excipients, *e.g.*, a pharmaceutically acceptable excipient or a cosmetically acceptable excipient, may comprise an anti-adherent, binder, coat, disintegrant, filler, flavor, color, lubricant, glidant, sorbent, preservative, or sweetener. In some embodiments, the
10 preparation may be substantially free of excipients.

In some embodiments, the preparation may be substantially free of one or more of the compounds or substances listed in the disclosure.

Exemplary compositions for spray, aerosol, or mist administration include solutions in saline, which can contain, for example, benzyl alcohol or other suitable preservatives, absorption
15 promoters to enhance bioavailability, and/or other solubilizing or dispersing agents. Conveniently in compositions for aerosol administration the ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* is delivered in the form of an aerosol spray presentation from a pressurized pack or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoro-methane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the
20 case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.*, gelatin can be formulated to contain a powder mix of the *N. eutropha* and a suitable powder base, for example lactose or starch. In certain embodiments, *N. eutropha* is administered as an aerosol from a metered dose valve, through an aerosol adapter also known as an actuator. Optionally, a stabilizer is also included, and/or porous
25 particles for deep lung delivery are included (*e.g.*, see U.S. Pat. No. 6,447,743).

Formulations may be presented with carriers such as cocoa butter, synthetic glyceride esters or polyethylene glycol. Such carriers are typically solid at ordinary temperatures, but liquefy and/or dissolve at body temperature to release the ammonia oxidizing bacteria, *e.g.*, *N. eutropha*.

Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene). In some aspects, the composition and/or excipient may be in the form of one or more of a liquid, a solid, or a gel. For example, liquid suspensions may include, but are not limited to, water, saline, phosphate-buffered saline, or an ammonia oxidizing storage buffer. Gel formulations may include, but are not limited to agar, silica, polyacrylic acid (for example Carbopol®), carboxymethyl cellulose, starch, guar gum, alginate or chitosan. In some embodiments, the formulation may be supplemented with an ammonia source including, but not limited to ammonium chloride or ammonium sulfate.

In some embodiments, an ammonia oxidizing microorganism, *e.g.*, *N. eutropha* composition is formulated to improve NO penetration, *e.g.*, into the skin or other target tissue. A gel-forming material such as KY jelly or various hair gels would present a diffusion barrier to NO loss to ambient air, and so improve the skin's absorption of NO. The NO level in the skin will generally not greatly exceed 20 nM/L because that level activates GC and would cause local vasodilatation and oxidative destruction of excess NO.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations as described herein may include other agents conventional in the art having regard to the type of formulation in question.

The formulation, *e.g.*, preparation, *e.g.*, composition may be provided in a container, delivery system, or delivery device, having a weight, including or not including the contents of the container, that may be less than about 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, or 2000 grams.

Suitable unit dosage formulations are those containing an effective dose, as hereinbefore recited, or an appropriate fraction thereof, of ammonia oxidizing microorganisms, *e.g.*, *N. eutropha*.

A therapeutically effective amount of ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* may be administered as a single pulse dose, as a bolus dose, or as pulse doses administered over time. Thus, in pulse doses, a bolus administration of ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* is provided, followed by a time period wherein ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* is administered to the subject, followed by a second

bolus administration. In specific, non-limiting examples, pulse doses are administered during the course of a day, during the course of a week, or during the course of a month.

In some embodiments, a preparation of ammonia oxidizing microorganisms, *e.g.*, a formulation, *e.g.*, a composition, may be applied for a pre-determined number of days. This may
5 be based, for example, at least in part, on the severity of the condition or disease, the response to the treatment, the dosage applied and the frequency of the dose. For example, the preparation may be applied for about 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, 84-91 days., for about 1 month, for about 2 months, for about 3 months. In some embodiments, the ammonia oxidizing bacteria is administered for an
10 indefinite period of time, *e.g.*, greater than one year, greater than 5 years, greater than 10 years, greater than 15 years, greater than 30 years, greater than 50 years, greater than 75 years. In certain aspects, the preparation may be applied for about 16 days.

In some embodiments, a preparation of ammonia oxidizing microorganisms, *e.g.*, a formulation, *e.g.*, a composition, may be applied a pre-determined number of times per day.
15 This may be based, for example, at least in part, on the severity of the condition or disease, the response to the treatment, the dosage applied and the frequency of the dose. For example, the preparation may be applied 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 times per day.

In some embodiments, the preparation may be applied one time per day. In other
20 embodiments, the preparation may be applied two times per day. In some embodiments, the preparation may be applied a first pre-determined amount for a certain number of days, and a second pre-determined amount for a certain subsequent number of days. In some embodiments, the preparation may be applied for about 16 days.

In accordance with one or more embodiments, the preparation may generally be
25 compatible with a physiological environment associated with the subject. In at least some embodiments, compositions are formulated to have a substantially neutral pH or a physiological pH, for instance a pH that normally prevails in the target site for intended delivery, administration, or desired effect. Compositions may be formulated to have a pH between about 5.5 and about 8.5. Compositions may be formulated to comprise compatible conditions, *e.g.*, pH,
30 tonicity, with the target site of physiological environment associated with the subject.

The preparation may be formulated for transmucosal delivery and/or circulation, *e.g.* locally or systemically. In some embodiments, the preparation may be formulated such that ammonia oxidizing microorganisms, products thereof, or byproducts thereof (*e.g.*, nitrate, nitrite, NO, or CoQ8) penetrate a deposit or target tissue at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. The preparation may be formulated such that 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of ammonia oxidizing microorganisms, products thereof, or byproducts thereof, penetrate a deposit or target tissue or enter circulation.

In accordance with one or more embodiments, the preparation may be in the form of a solution, suspension, emulsion, cream, ointment, gel, hydrogel, or liquid, *e.g.* drop, spray, aerosol, or mist, tablet, capsule, or device for administration to a subject.

In accordance with one or more embodiments, a preparation, composition, formulation, or product comprising ammonia oxidizing microorganisms may undergo quality control and/or testing while it is being made and/or upon its completion. International (PCT) Patent Application Publication No. WO2015/179669 (International (PCT) Patent Application Serial No. PCT/US2015/032017 as filed on May 21, 2015) which is hereby incorporated herein by reference in its entirety for all purposes describes various methods of preparing materials with ammonia oxidizing microorganisms and of testing such materials. For example, one or more parameters such as OD level, pH level, waste level, nutrient level, contaminant level, oxidation rate, nitrite level, protein concentration may be compared against a predetermined value to assess or evaluate a preparation comprising ammonia oxidizing microorganisms.

The present disclosure provides, *inter alia*, a kit comprising preparations of ammonia oxidizing microorganisms, as disclosed herein. Formulations may comprise discrete units, *e.g.*, solid, liquid, or gas formulations of ammonia oxidizing microorganisms. Formulations, *e.g.*, solutions, aerosols, sprays, and mists, may be presented in multi-dosage form (multiple use), *e.g.*, packaged units including a predetermined number of dosages, or single dosage form (single use), *e.g.*, packaged units including a single dose. Preparations of ammonia oxidizing microorganisms may be packaged in devices or containers configured to hold a volume of at least about less than 1 ml, 1 ml, 5 ml, 10 ml, 20 ml, 25 ml, 40 ml, 50 ml, 60 ml, 70 ml, 80 ml, 90 ml, 100 ml, or more than about 100 ml.

Kits may further comprise one or more device for administration of the preparation, for example, syringe, needle, catheter, enema, bulb, pipette (eye or ear dropper), and other devices for drug administration as known in the art. Kits may comprise instructions for use, for example instructions for administration of ammonia oxidizing microorganisms as disclosed herein or
5 instructions for combination therapy including administration of ammonia oxidizing microorganisms. Kits may comprise a second or subsequent composition for administration in conjunction with an ammonia oxidizing preparation, as disclosed herein. For instance, kits may comprise a supplement or composition comprising a product or byproduct of ammonia oxidizing microorganisms, a composition that promotes growth or metabolism of ammonia oxidizing
10 microorganisms, a composition that promotes production of products or byproducts of ammonia oxidizing microorganisms, a composition that promotes urease activity, or a composition that has a synergistic effect with ammonia oxidizing microorganisms, or a composition or pharmaceutical agent that treats, e.g., is approved to treat or commonly used to treat, a relevant disease, disorder, or a symptom of a relevant disease or disorder, for example an anti-inflammatory composition.
15 Kits may comprise “biome-friendly” or “biome-compatible” products as disclosed herein, for example one or more microbiome-compatible cosmetic products. Any of the products contained in the kit may be specifically formulated to treat a target indication and/or formulated for a desired mode of delivery, as described herein.

20 **Natural Products; Consumer Products**

In some specific embodiments, a preparation comprising ammonia oxidizing microorganisms as discussed herein may be a natural product or a consumer product. In other embodiments, a preparation of ammonia oxidizing microorganism may instead be used in conjunction with a natural product or consumer product.

25 Ammonia oxidizing microorganisms, e.g., *N. eutropha* may be associated with a variety of natural products, and examples of such products are set out below. These natural products may be comprised of formulations, compositions, or preparations disclosed throughout this disclosure.

Natural products may be or comprise products for commercial purposes, and may refer to
30 cosmetics, dietary supplements, and foods, e.g., food, food supplements, medical food, food

additive, nutraceutical, or drink, produced from natural sources. Natural products may have pharmacological or biological activity that may be of therapeutic benefit, *e.g.*, in treating disease or conditions. Natural products may be included in traditional medicines, treatments for cosmetological purposes, and spa treatments. A natural product referred to herein may comprise any one or more of the components described as a natural product to be incorporated into a preparation or formulation comprising one or more other components, *e.g.*, excipients. The preparation or formulation referred to as a natural product may comprise a natural product defined herein and one or more additional components or ingredients. Any of the compositions, preparations, or formulations discussed throughout this disclosure may be or comprise one or more natural products.

In some embodiments, the natural product or the fortified natural product may comprise at least one of mud, water, food-derived products, plant-derived products, extracts, and oils. The natural product or the fortified natural product may be used in a spa treatment. In some embodiments, the natural product or the fortified natural product may be incorporated into at least one of a powder, cream, lotion, wrap, scrub, eye mask, facial mask, body mask, aerosol, *e.g.*, mist, spray, salve, wipe, stick, bandage, or soak.

In some embodiments, the natural product or fortified natural product may be provided as, or may be disposed in at least one of a baby product, *e.g.*, a baby shampoo, a baby lotion, a baby oil, a baby powder, a baby cream; a bath preparation, *e.g.*, a bath oil, a tablet, a salt, a bubble bath, a bath capsule; an eye makeup preparation, *e.g.*, an eyebrow pencil, an eyeliner, an eye shadow, an eye lotion, an eye makeup remover, a mascara; a fragrance preparation, *e.g.*, a colognes, a toilet water, a perfume, a powder (dusting and talcum), a sachet; hair preparations, *e.g.*, hair conditioners, hair sprays, hair straighteners, permanent waves, rinses, shampoos, tonics, dressings, hair grooming aids, wave sets; hair coloring preparations, *e.g.*, hair dyes and colors, hair tints, coloring hair rinses, coloring hair shampoos, hair lighteners with color, hair bleaches; makeup preparations, *e.g.*, face powders, foundations, leg and body paints, lipstick, makeup bases, rouges, makeup fixatives; manicuring preparations, *e.g.*, basecoats and undercoats, cuticle softeners, nail creams and lotions, nail extenders, nail polish and enamel, nail polish and enamel removers; oral hygiene products, *e.g.*, dentrifices, mouthwashes and breath fresheners; bath soaps and detergents, deodorants, douches, feminine hygiene deodorants; shaving preparations,

e.g., aftershave lotions, beard softeners, talcum, preshave lotions, shaving cream, shaving soap; skin care preparations, *e.g.*, cleansing, depilatories, face and neck, body and hand, foot powders and sprays, moisturizing, night preparations, paste masks, skin fresheners; and suntan preparations, *e.g.*, gels, creams, and liquids, and indoor tanning preparations.

5 Ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* may be associated with a variety of consumer products, and examples of such products are set out below and be comprised of formulations, compositions, or preparations disclosed throughout this disclosure. In some embodiments, the ammonia oxidizing bacteria, *e.g.*, *N. eutropha* associated with a product is admixed with the product, for example, spread evenly throughout the product, and in some
10 embodiments, ammonia oxidizing bacteria, *e.g.*, the *N. eutropha* associated with a product is layered on the product.

In some embodiments, the preparation may be disposed in, or provided as, a powder, cosmetic, cream, stick, aerosol, *e.g.*, mist, salve, wipe, or bandage.

In some embodiments, ammonia oxidizing bacteria, *e.g.*, *N. eutropha* is associated with a
15 powder. Powders are typically small particulate solids that are not attached to each other and that can flow freely when tilted. Exemplary powders for consumer use include talcum powder and some cosmetics (*e.g.*, powder foundation).

In some embodiments, the ammonia oxidizing bacteria is associated with a cosmetic. The cosmetic may be a substance for topical application intended to alter a person's appearance, *e.g.*, a liquid foundation, a powder foundation, blush, or lipstick, and may be referred to as a
20 preparation. The cosmetic may be any substance recited in the Food and Drug Administration regulations, *e.g.*, under 21 C.F.R. § 720.4.

In some embodiments, ammonia oxidizing bacteria, *e.g.*, *N. eutropha* is associated with a cosmetic. The cosmetic may be a substance for topical application intended to alter a person's
25 appearance, *e.g.*, a liquid foundation, a powder foundation, blush, or lipstick. Other components may be added to these cosmetic preparations as selected by one skilled in the art of cosmetic formulation such as, for example, water, mineral oil, coloring agent, perfume, aloe, glycerin, sodium chloride, sodium bicarbonate, pH buffers, UV blocking agents, silicone oil, natural oils, vitamin E, herbal concentrates, lactic acid, citric acid, talc, clay, calcium carbonate, magnesium

carbonate, zinc oxide, starch, urea, and erythorbic acid, or any other excipient known by one of skill in the art, including those disclosed herein.

The preparation, e.g., the cosmetic, may be at least one of a baby product, e.g., a baby shampoo, a baby lotion, a baby oil, a baby powder, a baby cream; a bath preparation, e.g., a bath oil, a tablet, a salt, a bubble bath, a bath capsule; an eye makeup preparation, e.g., an eyebrow pencil, an eyeliner, an eye shadow, an eye lotion, an eye makeup remover, a mascara; a fragrance preparation, e.g., a colognes, a toilet water, a perfume, a powder (dusting and talcum), a sachet; hair preparations, e.g., hair conditioners, hair sprays, hair straighteners, permanent waves, rinses, shampoos, tonics, dressings, hair grooming aids, wave sets; hair coloring preparations, e.g., hair dyes and colors, hair tints, coloring hair rinses, coloring hair shampoos, hair lighteners with color, hair bleaches; makeup preparations, e.g., face powders, foundations, leg and body paints, lipstick, makeup bases, rouges, makeup fixatives; manicuring preparations, e.g., basecoats and undercoats, cuticle softeners, nail creams and lotions, nail extenders, nail polish and enamel, nail polish and enamel removers; oral hygiene products, e.g., dentrifices, mouthwashes and breath fresheners; bath soaps and detergents, deodorants, douches, feminine hygiene deodorants; shaving preparations, e.g., aftershave lotions, beard softeners, talcum, preshave lotions, shaving cream, shaving soap; skin care preparations, e.g., cleansing, depilatories, face and neck, body and hand, foot powders and sprays, moisturizing, night preparations, paste masks, skin fresheners; and suntan preparations, e.g., gels, creams, and liquids, and indoor tanning preparations.

In some embodiments, the formulations, compositions, or preparations described herein, may comprise, be provided as, or disposed in at least one of a baby product, e.g., a baby shampoo, a baby lotion, a baby oil, a baby powder, a baby cream; a bath preparation, e.g., a bath oil, a tablet, a salt, a bubble bath, a bath capsule; a powder (dusting and talcum), a sachet; hair preparations, e.g., hair conditioners, rinses, shampoos, tonics, face powders, cuticle softeners, nail creams and lotions, oral hygiene products, mouthwashes, bath soaps, douches, feminine hygiene deodorants; shaving preparations, e.g., aftershave lotions, skin care preparations, e.g., cleansing, face and neck, body and hand, foot powders and sprays, moisturizing, night preparations, paste masks, skin fresheners; and suntan preparations, e.g., gels, creams, and liquids.

In some embodiments, ammonia oxidizing microorganisms, *e.g.*, the *N. eutropha* is associated with an aerosol, spray, or mist and these terms may be used interchangeably. An aerosol is typically a colloid of fine solid particles or fine liquid droplets, in a gas such as air. Aerosols may be created by placing the *N. eutropha* (and optionally carriers) in a vessel under pressure, and then opening a valve to release the contents. The container may be designed to only exert levels of pressure that are compatible with *N. eutropha* viability. For instance, the high pressure may be exerted for only a short time, and/or the pressure may be low enough not to impair viability. Examples of consumer uses of aerosols include for sunscreen, deodorant, perfume, hairspray, and insect repellent. The aerosol may be referred to as a spray or mist.

The compositions comprising ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* may also comprise one or more of a moisturizing agent, deodorizing agent, scent, colorant, insect repellent, cleansing agent, or UV-blocking agent.

In some embodiments, ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* are associated with cloth, yarn, or thread. Articles of clothing such as, for example, shoes, shoe inserts, pajamas, sneakers, belts, hats, shirts, underwear, athletic garments, helmets, towels, gloves, socks, bandages, and the like, may also be treated with ammonia oxidizing bacteria, *e.g.*, *N. eutropha*. Bedding, including sheets, pillows, pillow cases, and blankets may also be treated with ammonia oxidizing bacteria, *e.g.*, *N. eutropha*. In some embodiments, areas of skin that cannot be washed for a period of time may also be contacted with ammonia oxidizing bacteria, *e.g.*, *N. eutropha*. For example, skin enclosed in orthopedic casts which immobilize injured limbs during the healing process, and areas in proximity to injuries that must be kept dry for proper healing such as stitched wounds may benefit from contact with the ammonia oxidizing bacteria, *e.g.*, *N. eutropha*.

In some aspects, the present disclosure provides a wearable article comprising ammonia oxidizing microorganisms as described herein. A wearable article may be a light article that can be closely associated with a user's body, in a way that does not impede ambulation. Examples of wearable articles include a wristwatch, wristband, headband, hair elastic, hair nets, shower caps, hats, hairpieces, and jewelry. The wearable article comprising an ammonia oxidizing bacteria, *e.g.*, *N. eutropha* strain described herein may provide, *e.g.*, at a concentration that provides one or more of a treatment or prevention of a skin disorder, a treatment or prevention of a disease or

condition associated with low nitrite levels, a treatment or prevention of body odor, a treatment to supply nitric oxide to a subject, or a treatment to inhibit microbial growth.

In some embodiments, the ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* are associated with a product intended to contact the hair, for example, a brush, comb, shampoo, conditioner, headband, hair elastic, hair nets, shower caps, hats, and hairpieces. Nitric oxide formed on the hair, away from the skin surface, may be captured in a hat, scarf or face mask and directed into inhaled air.

Articles contacting the surface of a human subject, such as a diaper, may be associated with ammonia oxidizing microorganisms, *e.g.*, *N. eutropha*. Because diapers are designed to hold and contain urine and feces produced by incontinent individuals, the urea in urine and feces can be hydrolyzed by skin and fecal bacteria to form free ammonia which is irritating and may cause diaper rash. Incorporation of bacteria that metabolize urea into nitrite or nitrate, such as ammonia oxidizing bacteria, *e.g.*, *N. eutropha*, may avoid the release of free ammonia and may release nitrite and ultimately NO which may aid in the maintenance of healthy skin for both children and incontinent adults. The release of nitric oxide in diapers may also have anti-microbial effects on disease causing organisms present in human feces. This effect may continue even after disposable diapers are disposed of as waste and may reduce the incidence of transmission of disease through contact with soiled disposable diapers.

In some embodiments, the product comprising ammonia oxidizing microorganisms, *e.g.*, *N. eutropha* is packaged. The packaging may serve to compact the product or protect it from damage, dirt, or degradation. The packaging may comprise, *e.g.*, plastic, paper, cardboard, or wood. In some embodiments the packaging is impermeable to bacteria. In some embodiments, the packaging is permeable to oxygen and/or carbon dioxide.

Methods of Treatment with Ammonia Oxidizing Microorganisms

In accordance with one or more embodiments, a subject may be treated via administration of ammonia oxidizing microorganisms, *e.g.*, a preparation comprising ammonia oxidizing microorganisms. As used herein, treatment of a subject may comprise administering an ammonia oxidizing microorganism composition for a cosmetic or therapeutic result. For

instance, treatment may comprise treating or alleviating a condition, symptom, or side effect associated with a condition or achieving a desired cosmetic effect.

Subjects may include an animal, a mammal, a human, a non-human animal, a livestock animal, or a companion animal. The subject may be female or male. The subject may have various skin types. The subject may have various health-related profiles, including health history and/or genetic predispositions. The subject may generally have a normal microbiome, e.g., a physiological microbiome, or a disrupted microbiome. The subject may be characterized as one of the following ethnicity/race: Asian, black or African American, Hispanic or Latino, white, or multi-racial. The subject may be of an age of less than 1, or between 1-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, or over 60 years.

The ammonia oxidizing microorganisms that may be used to treat a subject include all the ammonia oxidizing microorganisms, e.g., *N. eutropha* compositions described in this application, e.g. a purified preparation of optimized ammonia oxidizing microorganisms, for instance strain D23.

The methods may be provided to administer, or deliver a therapeutic product or a cosmetic product. The methods may comprise administering or introducing a preparation comprising live ammonia oxidizing microorganisms to a subject. The preparation may be formulated to treat a target indication and/or formulated for a desired mode of delivery.

In accordance with one or more embodiments, a preparation comprising live ammonia oxidizing microorganisms may be administered to a first tissue of a subject. The first tissue may be a deposit tissue. The first tissue may be a target tissue or a tissue other than a target tissue. The live ammonia oxidizing microorganisms, or a product thereof, e.g., nitrite and/or nitric oxide, may then move or be transported to a second tissue, e.g., via diffusion. The second tissue may be a target tissue. The target tissue may be associated with a desired local or systemic effect. The target tissue may be associated with an indication, disorder, or condition to be treated.

Ammonia oxidizing microorganism preparations may be administered, for example to the skin, for a cosmetic or therapeutic effect. For instance, administration may provide a cosmetic treatment, benefit, or effect. In some embodiments, administration may provide for treatment or improvement of one or more of oily appearance, pore appearance, radiance, blotchiness, skin

tone evenness, visual smoothness, and tactile smoothness. In some embodiments, a cosmetic appearance of a subject may be altered such as may result from improved skin health. Signs of aging may be reduced, delayed, or reversed. Administration may result in a qualitative improvement in skin and/or scalp condition and/or quality. Skin smoothness, hydration, tightness, and/or softness in a subject may be improved. The present disclosure also provides a method of reducing body odor.

Administration may provide a therapeutic treatment, benefit, or effect. The present disclosure provides a method of supplying nitrite and nitric oxide to a subject. The present disclosure provides various methods for the suppression, treatment, or prevention of diseases, disorders, infections, and conditions using ammonia oxidizing microorganisms. Ammonia oxidizing microorganisms may be used, for instance, to treat various diseases associated with low nitrite levels, skin diseases, and diseases caused by pathogenic bacteria. In some embodiments, administration may provide for a reduction in inflammation. Indeed, a local or systemic anti-inflammatory effect may be demonstrated. In at least some embodiments, microbial growth may be inhibited. Skin and overall health may be improved. Inadequate circulation may be augmented. Endothelial function may be promoted. A change in level of nitrite or NO at a target tissue or in circulation may be demonstrated. In some embodiments, administration, e.g., administration of an effective amount, may modulate, change, or alter a level of nitrite or NO at a target tissue or in circulation. In some embodiments, administration, e.g., administration of an effective amount, may result in an increased level of nitrite or NO at a target tissue or in circulation.

Administration of the compositions disclosed herein may provide transmucosal delivery and/or circulation, *e.g.* locally or systemically. In some embodiments, administration may provide that ammonia oxidizing microorganisms, products thereof, or byproducts thereof (*e.g.*, nitrate, nitrite, NO, or CoQ8) penetrate a deposit or target tissue at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. In at least some embodiments, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of ammonia oxidizing microorganisms, products thereof, or byproducts thereof, penetrate a deposit or target tissue or enter circulation upon administration of the compositions disclosed herein.

The preparations and methods of the present disclosure may provide for reducing an amount of undesirable microorganisms from an environment associated with a subject. The ammonia oxidizing microorganisms described herein may out-compete other organisms by, *e.g.*, consuming scarce nutrients, or generating byproducts that are harmful to other organisms, *e.g.*, changing a pH level that is not conducive to the undesirable organism's growth.

The present disclosure also provides a method of promoting wound healing, including of chronic wounds, such as in a patient that has an impaired healing ability, *e.g.*, a diabetic patient. A bandage including ammonia oxidizing microorganisms may optionally be applied to the wound.

It is appreciated that many modern degenerative diseases may be caused by a lack of NO species, and that AOM may be administered to supply those species, directly to a target tissue or via diffusion to a target tissue. Application of AOM may resolve long standing medical conditions. In certain embodiments, AOM are applied to a subject to offset modern bathing practices, especially with anionic detergents which remove AOM from the external skin.

In accordance with one or more embodiments, AOM convert ammonia to nitrite, an anti-microbial compound, and nitric oxide, a well-documented signaling molecule in the inflammatory process.

The present disclosure provides, *inter alia*, a method of modulating a composition of a microbiome, *e.g.*, modulating or changing the proportions of a microbiome in an environment, *e.g.*, a surface, *e.g.*, a surface of a subject. This may, in turn, exhibit a health-related benefit. The method may comprise administering a preparation comprising ammonia oxidizing microorganisms to a subject. In some embodiments, the amount and frequency of administration, *e.g.*, application, may be sufficient to reduce a proportion of pathogenic microorganisms.

Application of ammonia oxidizing microorganisms to a subject, *e.g.*, a human subject may lead to unexpected changes in the microbiome. It may lead to increases in the proportion of normal commensal non-pathogenic species and reductions in the proportion of potentially pathogenic, pathogenic, or disease causing organisms.

An increase in the proportion of non-pathogenic bacteria may occur with a pre-determined period of time, *e.g.*, in less than 1 day, 2 days, 3 days, 4 days, 5 days, 1 week, 2

weeks, 3 weeks, or 4 weeks, or in less than 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, 84-91 days.

A decrease in the proportion of pathogenic bacteria may occur with a pre-determined period of time, *e.g.*, in less than 1 day, 2 days, 3 days, 4 days, 5 days, 1 week, 2 weeks, 3 weeks, 5 or 4 weeks, or in less than 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, 84-91 days.

In accordance with one or more embodiments, a subject may be evaluated for need of treatment. In some embodiments, a subject may be selected on the basis of the subject being in need of a treatment. The present disclosure may further provide obtaining a sample from a 10 subject and analyzing the sample. In some embodiments, subjects may be evaluated before, during, and/or after treatment, such as a predetermined time intervals.

In accordance with one or more embodiments, administration may be performed before, during, or subsequent to occurrence of a health-related condition, or in response to a warning sign, trigger, or symptom thereof. In accordance with one or more embodiments, a second 15 amount of the preparation may be administered to the subject, *e.g.*, a second dose.

In certain aspects, the present disclosure provides combination therapies comprising ammonia oxidizing microorganisms, *e.g.*, a *N. eutropha* and a second treatment, *e.g.* a therapeutic. For instance, the disclosure provides physical admixtures of the two (or more) therapies are physically admixed. In other embodiments, the two (or more) therapies are 20 administered in combination as separate formulation. The second therapy may be, *e.g.*, a pharmaceutical agent, surgery, diagnostic, or any other medical approach that treats, *e.g.*, is approved to treat or commonly used to treat, the relevant disease, disorder, or a symptom of the relevant disease or disorder. The second treatment may be administered before or after the administration. The effective amount can be administered concurrently with the second 25 treatment. The second treatment may be administered via the same or a different mode of delivery. The subject may have a therapeutic level of the second treatment upon administration of the preparation. In certain embodiments, the second treatment may provide an anti-inflammatory effect or be administered to reduce inflammation at the target site. In at least some embodiments, the preparation may be administered concurrently or in conjunction with a product 30 or byproduct of the ammonia oxidizing microorganisms, *e.g.*, nitrite, nitrate, nitric oxide, CoQ8.

In at least some embodiments, the preparation may be administered concurrently or in conjunction with a composition that promotes growth or metabolism of ammonia oxidizing microorganisms, promotes production of products or byproducts of ammonia oxidizing microorganisms, promotes urease activity, or has a synergistic effect with ammonia oxidizing microorganisms, e.g., ammonia, ammonium salts, urea, and urease.

The preparation may be administered with a microbiome cleansing preparation, for example a local or systemic antibiotic. The preparation may be administered after administration of a cleansing preparation or a bowel cleanse. The preparations may be administered pre- or post-surgical procedure, diagnostic procedure, or natural event, e.g., giving birth. The preparations may be administered before, during, or after deposit of an implantable or invasive device.

In accordance with one or more embodiments, the preparation may be administered as an analgesic or prophylactic. The preparation may be self-administered. The administration of the preparation may be device-assisted.

In some embodiments, the ammonia oxidizing microorganisms, e.g., a preparation of ammonia oxidizing microorganisms, are administered at a dose of about or greater than about $10^3 - 10^4$ CFU, $10^4 - 10^5$ CFU, $10^5 - 10^6$ CFU, $10^6 - 10^7$ CFU, $10^7 - 10^8$ CFU, $10^8 - 10^9$ CFU, $10^9 - 10^{10}$ CFU, $10^{10} - 10^{11}$ CFU, $10^{11} - 10^{12}$ CFU, $10^{12} - 10^{13}$ CFU, or $10^{13} - 10^{14}$ CFU per application, per day, per week, or per month. In some embodiments, the ammonia oxidizing microorganisms are administered at a dose of about $10^9 - 10^{10}$ CFU, e.g., about $1 \times 10^9 - 5 \times 10^9$, $1 \times 10^9 - 3 \times 10^9$, or $1 \times 10^9 - 10 \times 10^9$ CFU per application or per day.

In some embodiments, the ammonia oxidizing microorganisms are administered in a volume of about 1-2, 2-5, 5-10, 10-15, 12-18, 15-20, 20-25, or 25-50 ml per dose. In some embodiments, the solution is at a concentration of about $10^8 - 10^9$, $10^9 - 10^{10}$, or $10^{10} - 10^{11}$ CFU/ml.

In some embodiments, the ammonia oxidizing microorganisms are administered as two 15 ml doses per day, where each dose is at a concentration of 10^9 CFU/ml.

In some embodiments, the ammonia oxidizing microorganisms are administered once, twice, three, or four times per day. In some embodiments, the ammonia oxidizing microorganisms is administered once, twice, three, four, five, or six times per week. In some

embodiments, the ammonia oxidizing microorganisms is administered shortly after bathing. In some embodiments, the ammonia oxidizing microorganisms is administered shortly before sleep.

In some embodiments, the ammonia oxidizing microorganisms are administered for about 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, 84-91 days, *e.g.*, for about 1 month, for about 2 months, for about 3 months. In some embodiments, the ammonia oxidizing microorganisms is administered for an indefinite period of time, *e.g.*, greater than one year, greater than 5 years, greater than 10 years, greater than 15 years, greater than 30 years, greater than 50 years, greater than 75 years.

10 Administration of Ammonia Oxidizing Microorganisms to the Visual System

The pharmaceutical formulations (*e.g.*, preparations or compositions) described herein may include those suitable for ophthalmic delivery, *e.g.*, topically or via intravitreal delivery. Ammonia oxidizing microorganism preparations may be administered to the visual system for cosmetic or therapeutic purposes. For instance, compositions include those formulated for cosmetic or therapeutic use.

The ophthalmic formulations (*e.g.*, preparations or compositions) may conveniently be presented in unit dosage form and may be prepared by any of the methods known in the art of pharmacy or cosmetology. Typically, methods include the step of bringing the active ingredient (*e.g.*, ammonia oxidizing microorganism) into association with a pharmaceutical carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Ophthalmic formulations may be presented as discrete units, each containing a predetermined amount of the active ingredient as a solution or suspension in an aqueous or non-aqueous liquid, as a powder or granules, or as an oil-in-water or water-in-oil liquid emulsion. Various pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, *e.g.*, Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A., Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2 S, 1988; Aulton, M. and Taylor, K., Aulton's Pharmaceutics: The Design and

Manufacture of Medicines, 5th Edition, 2017; Antoine, A., Gupta M.R., and Stagner, W.C., Integrated Pharmaceutics: Applied Preformulation, Product Design, and Regulatory Science, 2013.

Compositions disclosed herein may be prepared in ophthalmic dosage formulations.

5 Ammonia Oxidizing Microorganisms can be administered to the visual system for cosmetic or therapeutic purposes. For instance, compositions may be prepared as ophthalmic solutions, ointments, gels, emulsions, and inserts. Each of the solutions, ointments, gels, emulsions, and inserts may be formulated to comprise one or more carrier or excipient, as described in more detail below. Typically, ophthalmic solutions may be aqueous solutions, *e.g.*, an aqueous
10 dispersion of the active agent. Ophthalmic ointments may comprise anhydrous dispersions of the active agent, *e.g.*, in a mineral oil-white petroleum base. Ophthalmic gels may comprise a polymer, *e.g.*, poloxamers, xanthan gum, gellan gum, locust bean gum, and carrageenan. Ophthalmic ointments and gels may provide for a longer residence time than, for example, aqueous solutions. The longer residence time may further allow for a reduced dosing interval.
15 Ophthalmic emulsions may comprise microspheres, microcapsules, nanoparticles, nanocapsules, micelles, liposomes, niosomes, dendrimers, or cyclodextrin complexes. However, to reduce the probability of causing stability problems and/or irritation to the eye, ophthalmic emulsions may comprise submicron or nanoemulsion particles. Such compositions may be formulated for topical application, intravitreal, or device-assisted application. Topical application formulations
20 may include, *e.g.*, ointments, gels, emulsions, suspensions, solutions, (*e.g.*, eye drops or eye wash), sprays, aerosols, and mists. Intravitreal application may be achieved by delivering the formulation as an injection, *e.g.*, via microneedle or an implant, *e.g.*, nondegradable, erodible, or biodegradable ocular inserts. Device-assisted application may include, for example, delivery in conjunction with ultrasound, iontophoresis, electroporation, or microneedles.

25 Generally, ophthalmic pharmaceutical compositions may be formulated to be compatible with a target tissue, *e.g.*, ocular tissues. Formulations disclosed herein may have a pH between about 7.14 and 7.82. In some embodiments, the formulations may have a pH of about 7.14, 7.15, 7.2, 7.25, 7.3, 7.35, 7.4, 7.45, 7.5, 7.55, 7.6, 7.65, 7.7, 7.75, 7.8, or 7.82 or a substantially physiological pH. Suitable formulations may have an osmolarity of between about 100
30 mOsmol/L and about 640 mOsmol/L, between about 200 mOsmol/L and about 400 mOsmol/L,

or between about 280 mOsmol/L and about 350 mOsmol/L. In some embodiments, formulations may have an osmolarity of 306 mOsmol/L. The surface tension of the formulation may be between about 43 nN/m and about 47 nN/m, specifically between about 43.4 nN/m and about 46.6 nN/m. Surface tension above 47 nN/m may cause eye irritation. Solutions may generally be formulated to resemble of the composition of tears. For instance, solutions and compositions disclosed herein may comprise sterol esters, glycoproteins, *e.g.*, mucin, carbonate buffers, electrolytes, and other excipients commonly found in tears. Isotonic saline alone may be toxic to certain ocular tissues, such as the cornea and conjunctiva.

Ammonia oxidizing compositions disclosed herein may comprise an effective amount of AOMs, for example, to promote lubrication or hydration of the eye or to treat an inflammatory disorder. The compositions may be administered to change or alter a level of nitrite or NO at a target tissue of the eye or in circulation. In some embodiments, administering an ammonia oxidizing microorganism composition, *e.g.*, an effective amount of an ammonia oxidizing microorganism composition, may modulate a microbiome associated with the eye of the subject.

The ammonia oxidizing microorganism compositions can, for example, be administered in a form suitable for immediate release or extended release. Suitable examples of immediate release formulations include topical formulations and intravitreal delivery formulations. Topical ophthalmic formulations for immediate release include, for example, compositions formulated to be absorbed through the cornea. Intravitreal delivery formulations for immediate release include, for example, direct intravitreal delivery by injection, implant, ultrasound, or iontophoresis.

Preparations for administration can be suitably formulated to give controlled release of ammonia oxidizing microorganism. In some embodiments, controlled release ophthalmic formulations may be formulated as an ointment, gel, or emulsion. Pharmaceutical ophthalmic extended release compositions may be formulated with one or more ion exchange agents. The one or more ion exchange agents may control drug release by ion exchange with ions present in tears and the ocular tear film. Specifically, monovalent and divalent ions in tears and the optical tear film may exchange with ions in the formulation and release the active agent for optical absorption.

The ammonia oxidizing microorganism compositions can, for example, be administered in form suitable to provide local treatment or systemic treatment. Compositions disclosed herein

may be administered to treat a local inflammatory disease, a symptom of a local or systemic inflammatory disease, or a side effect caused by a local or systemic inflammatory disease (e.g., conjunctivitis). Suitable examples of local periocular conditions that may be treated with compositions disclosed herein include blepharitis, Meibomian Gland Dysfunction (MGD), dry eye, itching, inflammation, wound healing, and intraocular diseases such as macular degeneration, infections, intraocular pressure disorders, e.g., elevated intraocular pressure, ocular hypertension, ocular hypotension, and glaucoma. In some embodiments, ammonia oxidizing microorganism compositions can be administered in a form suitable to treat certain infections and inflammatory disorders, e.g., bacterial infections, fungal infections, viral infections, itching, local inflammation, allergies, and wound healing. For instance, ammonia oxidizing microorganism compositions may be administered to treat inflammation associated with a surgical or diagnostic procedure, e.g., implants or invasive procedures, or generally inflammation related to any foreign body introduced into the visual system, e.g., contact lenses. Ammonia oxidizing microorganisms may be administered to treat localized symptoms of ophthalmic conditions, disorders, or systemic disorders, e.g., dry eyes (keratoconjunctivitis sicca), red eyes, itchy eyes, blurred or abnormal vision, watery eyes, twitching in the eye or surrounding tissues, or side effects associated with such conditions, disorders, or systemic disorders, e.g., sty or chalazion formation.

Examples of systemic conditions that may be treated with compositions disclosed herein include headaches, cardiovascular diseases, connective tissue disorders, inflammation, immune responses and autoimmune disorders, liver diseases, infections, neurological diseases, psychiatric disorders, nitric oxide disorders, urea cycle disorders, congestion, vasodilation disorders, skin diseases, wound healing, bowel disorders, reactions to insect bites, and certain viral, bacterial, and fungal infections. For instance, systemic conditions that may be treated with compositions disclosed herein include cardiovascular diseases such as cardioprotection, heart failure, hypertension, pulmonary arterial hypertension; immune responses and autoimmune disorders such as alopecia and vitiligo; liver diseases such as non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH); neurological diseases and psychological disorders such as depression, insomnia, and diabetic neuropathy; nitric oxide disorders such as erectile dysfunction; wound healing, e.g., from bed sores and nursing home care, burns, diabetic ulcers

e.g., foot ulcer, venous leg ulcer, biofilm, and mouth sores; skin diseases and disorders such as hyperhydrosis, pruritus, helomas, and subtypes of helomas; bowel disorders such as gluten sensitivity, irritable/inflammatory bowel disease, Crohn's disease, colitis, and necrotizing enterocolitis; and vasodilation disorders such as Renaud's disease, thermoregulation, and migraines. Certain viral, bacterial, and fungal infections may be treated with formulations disclosed herein, including infections caused by human papillomavirus (HPV), yeast infections, tinea versicolor, tinea unguium, tinea pedis/fungus, tinea cruris, jock itch, onychomycosis, dandruff, athlete's foot, sinusitis, otitis media, Methicillin-resistant *Staphylococcus aureus* (MRSA), staph, and bacterial vaginosis. Additional systemic conditions that may be treated with compositions disclosed herein include systemic inflammation, such as eczema, *e.g.*, adult and pediatric eczema, hives, idiopathic urticaria, lichen planus, insect bites including allergic reactions to insect bites, *e.g.*, mosquito and demodex folliculorum mite, reactions to poison ivy, itchiness, keratosis pilaris, laryngitis, pemphigus, psoriasis, rosacea, folliculitis and subtypes of folliculitis, hidradenitis suppurativa, perioral dermatitis, lupus rash, seborrheic dermatitis, *e.g.*, adult and infantile seborrheic dermatitis, acne, *e.g.*, adolescent acne, adult acne, and cystic acne, diaper rash, occupational hand dermatitis, sunburn, and dermatomyositis. Additionally, compositions disclosed herein may be delivered or applied to treat certain cosmetic indications, including but not limited to, contact dermatitis, diaper odor, *e.g.*, adult and pediatric, body odor, feminine odor, flaking, nail hardness, body odor, oily skin, razor burn, skin appearance, skit blotchiness, skin hydration, and sun spots. Compositions disclosed herein may be applied as a bug repellent or an antimicrobial agent. Additionally, compositions disclosed herein may be delivered or applied to treat certain cosmetic indications, including but not limited to, contact dermatitis, diaper odor, *e.g.*, adult and pediatric, body odor, feminine odor, flaking, nail hardness, body odor, oily skin, razor burn, skin appearance, skit blotchiness, skin hydration, and sun spots.

Ophthalmic preparations may be prepared for delivery to ocular capillaries for systemic therapeutic treatment. For example, topical drug absorption across the conjunctiva and sclera, also known as the noncorneal route of delivery, may be picked up by capillaries of the conjunctive and delivered to systemic circulation. Other methods of delivery to systemic

circulation include systemic transscleral delivery methods, such as subconjunctival, peribulbar, retrobulbar, and subtenon delivery methods.

Exemplary compositions may include one or more excipients, for example, absorption and penetration enhancers, preservatives, antioxidants, buffers, chelating agents, ion exchange
5 agents, solubilizing agents, suspending agents, thickeners, surfactants, wetting agents, tonicity-adjusting agents, and a vehicle for proper drug delivery. Absorption and penetration enhancers may improve the ability of the active agent to be absorbed by a number of different mechanisms. Antioxidants may reduce the oxidative degradation of the active agent. Buffers may maintain a desired pH of the composition and/or enhance solubility or stability of the composition.

10 Chelating agents may include complex trace metals that catalyze oxidation reactions of the composition. Ion exchange agents may control the release of active agent by ion exchange mechanisms. Solubilizing agents may increase the solubility of the active agent or another excipient. Suspending agents and thickeners may increase the viscosity or density of the composition to decrease settling rate of a dispersed material and/or increase the active agent's
15 retention time and residence time in the eye. Surfactants, including cationic, anionic, and non-ionic surfactants, and wetting agents may act to wet insoluble hydrophobic active agent or other excipients. Tonicity-adjusting agents may provide an isotonic solution with eye and ocular fluids. Vehicle, for example water, may provide bulk for proper active agent delivery.

Compositions disclosed herein may further contain or be used in conjunction with one or
20 more of ophthalmic ion-exchange formulations, penetration enhancers, *e.g.*, transporter-targeted prodrugs, ultrasound, soft drugs, steam, *e.g.*, a warm compress, and mucoadhesives. Penetration enhancers may increase active agent absorption, *e.g.*, across the cornea or other optical target tissue. Soft drugs may be formulated to be metabolically inactivated upon exerting their therapeutic effect. Suitable soft drugs include adaprolol, a soft beta-blocker typically used to
25 reduce intraocular pressure, and oteprednol etabonate, a soft steroid typically used to treat topical eye inflammation. Mucoadhesives may bond with the anionic mucin of the tear film and increase active agent absorption. Suitable mucoadhesives include carbomer, sodium carboxymethylcellulose, polycarbophil, and sodium alginate.

In some non-limiting embodiments, the preparations may be one or more of: substantially
30 odorless, colorless, not associated with substantial side effects, non-toxic, well-tolerated, have no

adverse effects if released into the environment, no risk of fostering antibiotic resistance, and have a physiology such that it can interact positively with various human microbiomes, e.g., microbiomes associated with a target tissue or the visual system, under normal and disease states.

Compositions disclosed herein may further be formulated as combination therapies.

5 Initial and subsequent therapeutic treatments may be provided in a single dosage form, prepared in individual dosage forms, administered concurrently, or administered separately. Individual dosage forms may be administered via the same mode of administration, e.g., through the visual system, or via an alternate mode of administration, e.g., orally, intranasally, topically, enterally, via the auditory system, via the urogenital system, via the respiratory system, via the
10 gastrointestinal system, or via injection. For instance, combination therapies may comprise ammonia oxidizing microorganisms for treatment of an inflammatory disease or condition. Individual dosage forms may be administered by a surgical or diagnostic procedure. Individual dosage forms may be administered in combination with a surgical or diagnostic procedure, for example before or after a surgical or diagnostic procedure. In some embodiments, ammonia
15 oxidizing microorganism compositions, for example prepared for ophthalmic administration, are formulated for combination therapy with an anti-inflammatory. Generally, compositions disclosed herein may be formulated for combination therapy with a drug or compound approved or commonly used to treat a disease, disorder, condition, symptom thereof, or side-effect thereof, for example a periocular disease, disorder, condition, symptom thereof, or side-effect thereof, as
20 disclosed herein. In at least some embodiments, preparations formulated for administration in combination with a treatment for dry, red, itchy, or watery eyes. Preparations may be formulated for administration in combination with a bowel cleanse or antibiotic to “plow the field” and allow for ammonia oxidizing microorganism colonization.

Preparations for administration to the eye may be formulated for targeted delivery to a
25 specific deposit or target tissue. In some embodiments, a preparation may be administered to a first tissue such that the preparation or a product of the preparation, e.g., ammonia oxidizing microorganisms or nitric oxide, is transported to a second tissue. In at least some embodiments, the preparation may be administered to a deposit ocular tissue such that nitric oxide penetrates and is transported to the ocular globe. The first tissue may be a deposit tissue. The second tissue
30 may be a target tissue. The deposit tissue and the target tissue may be the same or different

tissues. The deposit tissue, the target tissue, or both may be an ocular tissue. For instance, in some embodiments the deposit tissue may be a superficial ocular tissue and the target tissue may be posterior ocular tissue. Ammonia oxidizing microorganisms delivered to the deposit tissue may increase the level of nitric oxide which penetrates superficial ocular tissues to reach
5 posterior ocular tissues. The preparation or product of the preparation may be delivered locally or systemically, *e.g.*, at a deposit or target tissue or in circulation.

Ocular deposit or target tissues may include intravitreal tissues, the posterior chamber, or the anterior chamber of the eye of the subject. For instance, deposit and target tissues may include eyelid, inner eyelid, retina, ocular globe, orbit, optic disk, optic nerve, intravitreal,
10 vitreal humor, intrascleral, subconjunctiva, lens, ciliary body, iris, pupil, cornea, conjunctiva, sclera, and topical tissue or proximate tissue of the eye. The tear film can be a barrier to drug delivery. The precorneal layer of the tear film contains mucin. In some embodiments, the tear film can be used to enhance drug residence time and absorption through the use of mucoadhesives. Due to the wet environment of the eye, certain ocular tissues may also create a
15 barrier to passive diffusion. For instance, the aqueous humor drains from the canal of Schlemm and episcleral veins, bathing the cornea, iris, and lens. The fluid flow may create the barrier to passive diffusion into these tissues. Ophthalmic compositions may be specially formulated to penetrate and reach certain deposit or target tissues.

20 Administration of Ammonia Oxidizing Microorganisms to the Auditory System

The pharmaceutical formulations (*e.g.*, preparations or compositions) described herein may include those suitable for auditory delivery, *e.g.*, via topical, or inner or outer ear canal delivery. Ammonia oxidizing microorganism preparations may be administered to the auditory system for cosmetic or therapeutic purposes. For instance, compositions include those
25 formulated for cosmetic or therapeutic use.

The otic formulations (*e.g.*, preparations or compositions) may conveniently be presented in unit dosage form and may be prepared by any of the methods known in the art of pharmacy or cosmetology. Typically, methods include the step of bringing the active ingredient (*e.g.*, ammonia oxidizing microorganism) into association with a pharmaceutical carrier which
30 constitutes one or more accessory ingredients. In general, the formulations are prepared by

uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Otic formulations may be presented as discrete units, each containing a predetermined amount of the active ingredient as a solution or suspension in an aqueous or non-aqueous liquid, as a powder or granules, or as an oil-in-water or water-in-oil liquid emulsion. Various pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, *e.g.*, Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A., Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2 S, 1988; Aulton, M. and Taylor, K., Aulton's Pharmaceutics: The Design and Manufacture of Medicines, 5th Edition, 2017; Antoine, A., Gupta M.R., and Stagner, W.C., Integrated Pharmaceutics: Applied Preformulation, Product Design, and Regulatory Science, 2013; Allen, L.V. and Ansel, H.C., Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, 10th Edition, 2014.

Ammonia oxidizing compositions disclosed herein may comprise an effective amount of AOMs, for example, to penetrate a target tissue of the auditory system, to colonize a tissue of the auditory system, to treat an ear disorder or a symptom of an ear disorder, or to promote endothelial function, *e.g.*, within the auditory system.

Compositions disclosed herein may be prepared in auditory (otic, ear, or aural) dosage formulations. Ammonia Oxidizing Microorganisms can administered to the auditory system for cosmetic or therapeutic purposes. For instance, compositions may be prepared as otic solutions (*e.g.*, droplets, spray, aerosols, or mists), suspensions, emulsions, ointments, gels, hydrogels, or inserts. Each of the dosage formulations may be formulated to comprise one or more carrier or excipient, as described in more detail below. Typically, otic solutions may be aqueous solutions, *e.g.*, an aqueous dispersion, suspension, or emulsion of the active agent. Otic solutions may comprise the active agent in a liquid vehicle, *e.g.*, water or hydrogen peroxide. Otic solutions may comprise a suspension or dispersion of the active agent in mineral or vegetable oil. Ointments or other solutions may comprise a dispersion or suspension of the active agent in a viscous liquid, semi-liquid, gel, or hydrogel vehicle. For instance, compositions may comprise microspheres, microcapsules, nanoparticles, nanocapsules, micelles, liposomes, niosomes,

dendrimers, or cyclodextrin complexes. Such compositions may be formulated for topical administration, inner or outer ear canal administration, or device-assisted administration. Device-assisted administration may include, for example, delivery via an ear drop applicator, ear syringe, or other similar device.

5 Generally, otic pharmaceutical compositions may be formulated to be compatible with a target tissue, e.g., auditory tissues. Suitable formulations may have a substantially physiological pH, osmolarity, surface tension, etc. Ammonia oxidizing compositions disclosed herein may comprise an effective amount of AOMs, for example, to promote endothelial function or to treat an inflammatory disorder. The compositions may be administered to change or alter a level of
10 nitrite or NO at a target tissue of the ear or in circulation. In some embodiments, administering an ammonia oxidizing microorganism composition, e.g., an effective amount of an ammonia oxidizing microorganism composition, may modulate a microbiome associated with the ear of the subject.

 The ammonia oxidizing microorganism compositions can, for example, be administered
15 in a form suitable for immediate release or extended release. Suitable examples of immediate release formulations include solutions, suspensions, and ointment compositions. Immediate release formulations may comprise penetration enhancers.

 Preparations for administration can also be suitably formulated to give controlled release of ammonia oxidizing microorganisms, for example slow release of the microorganisms. In
20 some embodiments, controlled release otic formulations may comprise controlled release materials, such as polymer carriers. Controlled release otic formulations may further comprise a viscous vehicle, for example glycerin, anhydrous glycerin, and/or propylene glycol. Generally, otic compositions may be applied while the ear canal is in a vertical alignment, such that administration of the formulation is assisted by gravity. When the ear canal is placed in a
25 horizontal position, an aqueous or liquid formulation may exit the ear canal. Thus, a viscous vehicle may allow for increased residence and contact time with the tissues of the ear.

 Otic formulations are largely used for localized treatment, e.g., treatment of ear pain, pruritus in otitis media, swimmer's ear, removal of impacted ceramide (earwax), and treatment of ear infection, including bacterial infections. However, compositions disclosed herein may be
30 employed for treatment of systemic indications. The ammonia oxidizing microorganism

compositions can, for example, be administered in form suitable to provide local treatment or systemic treatment. Compositions disclosed herein may be administered to treat a local inflammatory disease, a symptom of a local or systemic inflammatory disease, or a side effect caused by a local or systemic inflammatory disease. Suitable examples of local auditory

5 conditions that may be treated with compositions disclosed herein include infection, inflammation, pain, and symptoms such as nausea, dizziness, and ear ache. For instance, local auditory conditions may include tinnitus, reduced hearing, vertigo, pruritus, congenital abnormalities, bacterial and fungal infections, *e.g.*, otitis media, and swimmer's ear. In some embodiments, ammonia oxidizing microorganism compositions can be administered in a form

10 suitable to treat certain infections and inflammatory disorders, *e.g.*, bacterial infections, fungal infections, viral infections, itching, local inflammation, allergies, and wound healing. For instance, ammonia oxidizing microorganism compositions may be administered to treat inflammation associated with a surgical or diagnostic procedure, *e.g.*, implants or invasive procedures, or generally inflammation related to any foreign body introduced into the auditory

15 system, *e.g.*, hearing aids and cochlear implants. Ammonia oxidizing microorganisms may be administered to treat localized symptoms of otic conditions, disorders, or systemic disorders, *e.g.*, ear ache, pruritus, tinnitus, nausea, and dizziness, or side effects associated with such conditions, disorders, or systemic disorders, *e.g.*, vertigo.

Examples of systemic conditions that may be treated with compositions disclosed herein

20 include headaches, cardiovascular diseases, connective tissue disorders, inflammation, immune responses and autoimmune disorders, liver diseases, infections, neurological diseases, psychiatric disorders, nitric oxide disorders, urea cycle disorders, congestion, vasodilation disorders, skin diseases, ophthalmic disorders, bowel disorders, wound healing, reactions to insect bites, and certain viral, bacterial, and fungal infections. For instance, systemic conditions that may be

25 treated with compositions disclosed herein include cardiovascular diseases such as cardioprotection, heart failure, hypertension, pulmonary arterial hypertension; immune responses and autoimmune disorders such as alopecia and vitiligo; liver diseases such as non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH); neurological diseases and psychological disorders such as depression, insomnia, and diabetic neuropathy; nitric oxide

30 disorders such as erectile dysfunction; wound healing, *e.g.*, from bed sores and nursing home

care, burns, diabetic ulcers *e.g.*, foot ulcer, venous leg ulcer, biofilm, and mouth sores; skin diseases and disorders such as hyperhydrosis, pruritus, helomas, and subtypes of helomas; ophthalmic disorders such as blepharitis, dry eye, macular degeneration, and glaucoma; bowel disorders such as gluten sensitivity, irritable/inflammatory bowel disease, Crohn's disease, colitis, and necrotizing enterocolitis; and vasodilation disorders such as Renaud's disease, thermoregulation, and migraines. Certain viral, bacterial, and fungal infections may be treated with formulations disclosed herein, including infections caused by human papillomavirus (HPV), yeast infections, tinea versicolor, tinea unguium, tinea pedis/fungus, tinea cruris, jock itch, onychomycosis, dandruff, athlete's foot, sinusitis, Methicillin-resistant *Staphylococcus aureus* (MRSA), staph, and bacterial vaginosis. Additional systemic conditions that may be treated with compositions disclosed herein include systemic inflammation, such as eczema, *e.g.*, adult and pediatric eczema, hives, idiopathic urticaria, lichen planus, insect bites including allergic reactions to insect bites, *e.g.*, mosquito and demodex folliculorum mite, reactions to poison ivy, itchiness, keratosis pilaris, laryngitis, pemphigus, psoriasis, rosacea, folliculitis and subtypes of folliculitis, hidradenitis supportiva, perioral dermatitis, lupus rash, seborrheic dermatitis, *e.g.*, adult and infantile seborrheic dermatitis, acne, *e.g.*, adolescent acne, adult acne, and cystic acne, diaper rash, occupational hand dermatitis, sunburn, and dermatomyositis. Additionally, compositions disclosed herein may be delivered or applied to treat certain cosmetic indications, including but not limited to, contact dermatitis, diaper odor, *e.g.*, adult and pediatric, body odor, feminine odor, flaking, nail hardness, body odor, oily skin, razor burn, skin appearance, skin blotchiness, skin hydration, and sun spots. Compositions disclosed herein may be applied as a bug repellent or an antimicrobial agent.

Exemplary compositions may include one or more pharmaceutically acceptable excipients, for example, absorption and penetration enhancers, preservatives, antioxidants, buffers, chelating agents, ion exchange agents, solubilizing agents, suspending agents, thickeners, surfactants, wetting agents, tonicity-adjusting agents, enzyme inhibitors, and vehicle for proper drug delivery. Absorption and penetration enhancers, *e.g.*, cyclodextrins, may improve the ability of the active agent to be absorbed by a number of different mechanisms. Antioxidants may reduce the oxidative degradation of the active agent. Buffers may maintain a desired pH of the composition and/or enhance solubility or stability of the composition.

Chelating agents may include complex trace metals that catalyze oxidation reactions of the composition. Ion exchange agents may control the release of active agent by ion exchange mechanisms. Solubilizing agents may increase the solubility of the active agent or another excipient. Suspending agents and thickeners may increase the viscosity or density of the composition to decrease settling rate of a dispersed material and/or increase the active agent's retention time and residence time in the ear. Surfactants, including cationic, anionic, and non-ionic surfactants, and wetting agents may act to wet insoluble hydrophobic active agent or other excipients. Tonicity-adjusting agents may provide an isotonic solution with auditory system tissues. Vehicle, for example water, may provide bulk for proper active agent delivery.

Compositions formulated for otic delivery may generally comprise one or more of anti-infective agents, analgesic agents, anti-inflammatory agents, hygroscopic agents, ceramide disrupting agents, and vehicle. Formulations may comprise anti-infective excipients, for instance, ciprofloxacin, solistin sulfate, neomycin, ofloxacin, polymyxin B sulfate, or nystatin. Formulations may comprise one or more analgesic agents or local analgesics to relieve local pain, for instance, antipyrine, pramoxine hydrochloride, and benzocaine. Formulations may comprise anti-inflammatory agents to reduce swelling and inflammation within the narrow ear canal, for instance, corticosteroids, *e.g.*, hydrocortisone, or dexamethasone sodium phosphate. Hygroscopic agents may be included to diminish moisture within the ear canal and reduce inflammation. Formulations may also comprise one or more ceramide disrupting excipients to diminish ceramide within the canal and allow passage of the formulation, for instance mineral oil, vegetable oil, hydrogen peroxide, or synthetic surfactants, *e.g.*, carbamide peroxide. Auditory formulation vehicle may comprise water, hydrogen peroxide, mineral or vegetable oil, glycerin, anhydrous glycerin, propylene glycol, aluminum acetate solution, or isopropyl alcohol or ethanol. Vehicles, such as aluminum acetate solution and isopropyl alcohol, may help to dry the ear canal. Formulations may also comprise agents to help reacidify the ear canal, for instance, acetic acid 2%-2.5% or boric acid 2%-5%. Thus, in some embodiments, formulations for otic application may have a substantially neutral or slightly acidic pH level.

In some non-limiting embodiments, the preparations may be one or more of: substantially odorless, colorless, not associated with substantial side effects, non-toxic, well-tolerated, have no adverse effects if released into the environment, no risk of fostering antibiotic resistance, and

have a physiology such that it can interact positively with various human microbiomes, e.g., microbiomes associated with a target tissue or the auditory system, under normal and disease states.

Compositions disclosed herein may further be formulated as combination therapies.

5 Initial and subsequent therapeutic treatments may be provided in a single dosage form, prepared in individual dosage forms, administered concurrently, or administered separately. Individual dosage forms may be administered via the same mode of administration, e.g., through the ear, or via an alternate mode of administration, e.g., orally, intranasally, enterically, topically, ocularly, via the urogenital system, via the gastrointestinal system, or via injection. For instance, 10 combination therapies may comprise ammonia oxidizing microorganisms for treatment of an inflammatory disease or condition. Individual dosage forms may be administered by a surgical or diagnostic procedure, for example before or after a surgical or diagnostic procedure. In some embodiments, ammonia oxidizing microorganism compositions, for example prepared for otic administration, are formulated for combination therapy with an anti-inflammatory. Generally, 15 compositions disclosed herein may be formulated for combination therapy with a drug or compound approved or commonly used to treat a disease, disorder, condition, symptom thereof, or side-effect thereof, for example an auditory disease, disorder, condition, symptom thereof, or side-effect thereof, as disclosed herein. Ammonia oxidizing microorganism therapeutic compositions may be administered in combination with a therapeutic treatment for ear pain, 20 pruritus, swimmer's ear, bacterial and fungal infections, and removal of impacted ceramide. Preparations may be formulated for administration in combination with a bowel cleanse or antibiotic to "plow the field" and allow for ammonia oxidizing microorganism colonization.

Preparations for administration to the auditory system may be formulated for targeted delivery to a specific deposit tissue or target tissue. In some embodiments, a preparation may be 25 administered to a first tissue such that the preparation or a product of the preparation, e.g., ammonia oxidizing microorganisms or nitric oxide, is transported to a second tissue. The first tissue may be a deposit tissue. The second tissue may be a target tissue. The deposit tissue and the target tissue may be the same or different tissues. In some embodiments, the deposit tissue, the target tissue, or both may be a tissue of the ear. The preparation or product of the preparation 30 may be delivered locally or systemically, e.g., at a deposit or target tissue or in circulation.

Auditory deposit or target tissues may generally be located on the outer ear, middle ear, or inner ear. Deposit or target tissues include an outer ear tissue, external auditory canal, internal auditory canal, tympanic membrane, tympanic cavity, malleus, incus, auditory tube, Eustachian tube, round window, oval window, ossicle, labyrinth, vestibule, utricle, saccule, staples, cochlea, canal, semicircular canal, temporal bone, vestibular nerve, and cochlear nerve. Any of the auditory tissues may be located on the right or left ear. Specific formulations to reach a deposit or target tissue may comprise designing the composition to comprise a specific vehicle or excipient. For instance, formulations comprising liquid or semi-liquid vehicles may reach a deeper deposit or target tissue within the inner ear than formulations comprising viscous vehicles or gels. Formulations comprising a ceramide disrupting agent may get past the ceramide barrier and reach a deeper deposit or target tissue. Compositions for delivery to the auditory system may be specially formulated to penetrate and reach certain deposit or target tissues. In some embodiments, the deposit or target tissue may be associated with a desired local or systemic effect.

Use of Microbiome Compatible Products with Administration of Ammonia Oxidizing Microorganisms

Microbiome compatible products may be used in conjunction with the preparations and methods disclosed herein. Various products may be considered to be “biome-friendly” or “biome-compatible.” Examples of biome-friendly products are disclosed in International (PCT) Patent Application Publication No. WO2017/004534 (International (PCT) Patent Application Serial No. PCT/US/2016/040723 as filed on July 1, 2016) which is hereby incorporated herein by reference in its entirety for all purposes. Some biome-friendly products may be cosmetic or therapeutic in nature. In accordance with one or more embodiments, biome-friendly products may be used in combination with microorganisms, e.g., non-pathogenic microorganisms, e.g., ammonia oxidizing microorganisms, which may in turn be used in the form of a preparation or composition to be applied to a subject. Ammonia oxidizing compositions disclosed herein may be administered for a cosmetic or therapeutic indication in conjunction with a biome-friendly or biome-compatible product.

In accordance with one or more embodiments, a preparation, composition, formulation or product comprising ammonia oxidizing microorganisms, e.g., for cosmetic or therapeutic use, may itself be considered biome-friendly. In other embodiments, a preparation comprising ammonia oxidizing microorganisms may be used in conjunction with a biome-friendly product.

5 In some embodiments, a preparation comprising ammonia oxidizing microorganisms may be mixed with a biome-friendly product or otherwise administered concurrently. In other embodiments, a preparation comprising ammonia oxidizing microorganisms may be distinct or separate from a biome-friendly product although potentially used in conjunction therewith. In some embodiments, a biome-friendly product is used alone. Ammonia oxidizing microorganism
10 composition preparations for use in conjunction with a biome-friendly product may be formulated for cosmetic or therapeutic use.

Biome-friendly or biome-compatible products may be used in conjunction with an ammonia oxidizing microorganism preparation formulated for any mode of delivery, e.g., formulated for targeted delivery to a subject, e.g., to a target tissue, region, system, or organ of a
15 subject. For example, the ammonia oxidizing microorganism preparation to be used in conjunction with a biome-friendly product may be formulated for delivery to the eye, ear, nose, urogenital system, respiratory system, or gastrointestinal system of the subject. In some embodiments, the ammonia oxidizing microorganism composition for use with a biome-friendly product may be formulated for targeted delivery based on a condition or disorder of a subject.
20 For instance, the formulation for targeted delivery may be based on a desired local or systemic effect to be achieved, e.g., a local or systemic therapeutic or cosmetic effect.

Biome-friendly cosmetic products that may be used with the present disclosure may be, or include, or be disposed in any one or more of a baby product, e.g., a baby shampoo, a baby lotion, a baby oil, a baby powder, a baby cream; a bath preparation, e.g., a bath oil, a tablet, a
25 salt, a bubble bath, a bath capsule; an eye makeup preparation, e.g., an eyebrow pencil, an eyeliner, an eye shadow, an eye lotion, an eye makeup remover, a mascara; a fragrance preparation, e.g., a colognes, a toilet water, a perfume, a powder (dusting and talcum), a sachet; hair preparations, e.g., hair conditioners, hair sprays, hair straighteners, permanent waves, rinses, shampoos, tonics, dressings, hair grooming aids, wave sets; hair coloring preparations, e.g., hair
30 dyes and colors, hair tints, coloring hair rinses, coloring hair shampoos, hair lighteners with

color, hair bleaches; makeup preparations, *e.g.*, face powders, foundations, leg and body paints, lipstick, makeup bases, rouges, makeup fixatives; manicuring preparations, *e.g.*, basecoats and undercoats, cuticle softeners, nail creams and lotions, nail extenders, nail polish and enamel, nail polish and enamel removers; oral hygiene products, *e.g.*, dentrifices, mouthwashes and breath
5 fresheners; bath soaps, *e.g.*, foaming body cleansers, and detergents, deodorants, douches, feminine hygiene deodorants; shaving preparations, *e.g.*, aftershave lotions, beard softeners, talcum, preshave lotions, shaving cream, shaving soap; skin care preparations, *e.g.*, cleansing, depilatories, face and neck, body and hand, foot powders and sprays, moisturizing, night preparations, paste masks, skin fresheners; and suntan preparations, *e.g.*, gels, creams, and
10 liquids, and indoor tanning preparations.

Products, *e.g.*, microbiome-compatible cosmetic products, *e.g.*, shampoos, conditioners, and cleansers, as described herein may be used in conjunction with the treatment of a condition, disease, or disorder. These cosmetic products may be used in conjunction with administration of the ammonia oxidizing microorganisms for therapeutic or cosmetic purposes. For example,
15 throughout the treatment period or cosmetic period of time of administering the ammonia oxidizing bacteria to a subject, the microbiome-compatible cosmetic products may be used. The microbiome-compatible cosmetic products may be used for a period of time prior to commencement of treatment of the therapeutic or cosmetic condition through administration of ammonia oxidizing bacteria to a subject. The microbiome-compatible cosmetic products may be
20 used for a period of time subsequent to commencement of treatment of the therapeutic or cosmetic condition through administration of ammonia oxidizing bacteria to a subject. The microbiome-compatible cosmetic products may be used for a period of time subsequent to discontinuation of therapeutic or cosmetic treatment of the condition through administration of ammonia oxidizing bacteria to a subject.

25 In some embodiments, the subject may apply one or more cosmetic product, and wait a period of time before administration of the ammonia oxidizing microorganisms. In other embodiments, the subject may administer the ammonia oxidizing microorganisms, and wait a period of time before applying one or more cosmetic products.

The period of time the subject may wait may be about 1 minute, 5 minutes, 10, 15, 20, 25, 30, 45, 60, 90, 120 minutes, or 3 hours, 4, 5, 6, 7, 8, 12, 18, 24 hours after applying one or more cosmetic product and prior to administration of ammonia oxidizing microorganisms.

5 The period of time the subject may wait may be about 1 minute, 5 minutes, 10, 15, 20, 25, 30, 45, 60, 90, 120 minutes, or 3 hours, 4, 5, 6, 7, 8, 12, 18, 24 hours after administering the ammonia oxidizing microorganisms and prior to applying one or more cosmetic products.

10 While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

Certain embodiments are within the scope of the following claims.

Claims

1. A method of introducing ammonia oxidizing microorganisms (AOM) to a subject,
comprising:
5 administering a preparation comprising AOM to an ear of the subject.
2. A method of introducing ammonia oxidizing microorganisms (AOM) to a subject,
comprising:
administering a preparation comprising AOM to an eye of the subject via topical or
10 intravitreal delivery.
3. The method of any of the preceding claims, wherein the preparation is administered topically.
4. The method of any of the preceding claims, wherein administration is substantially
15 noninvasive.
5. The method of any of the preceding claims, wherein an effective amount of the preparation is
administered to a target tissue of the ear of the subject.
- 20 6. The method of any of the preceding claims, wherein an effective amount of the preparation is
administered to a target tissue of the eye of the subject.
7. The method of any of the preceding claims, wherein the preparation is administered, e.g.,
topically applied, to a first tissue, e.g. a deposit tissue.
25
8. The method of any of the preceding claims, wherein the first tissue is the target tissue.
9. The method of any of the preceding claims, wherein the first tissue is other than the target
tissue, e.g., the preparation is applied to a first tissue and the preparation, or a product of the
30 preparation, e.g., NO, is transported, e.g., by diffusion, to a second tissue, e.g. the target tissue.

10. The method of any of the preceding claims, wherein a deposit tissue, target tissue, or both is an outer ear tissue, inner ear tissue, external auditory canal, internal auditory canal, tympanic membrane, tympanic cavity, malleus, incus, auditory tube, Eustachian tube, round window, oval window, ossicle, labyrinth, vestibule, utricle, saccule, staple, cochlea, canal, semicircular canal,
5 temporal bone, vestibular nerve, or cochlear nerve.
11. The method of any of the preceding claims, wherein a deposit tissue, target tissue, or both relates to an intravitreous tissue, posterior chamber, or another chamber of the eye of the subject.
- 10 12. The method of any of the preceding claims, wherein a deposit tissue, target tissue, or both is an eyelid, e.g., inner eyelid, retina, ocular globe, orbit, optic disk, optic nerve, vitreous humor, intrascleral, subconjunctiva, lens, ciliary body, iris, pupil, cornea, conjunctiva, sclera, topical tissue, or proximate tissue of the eye of the subject.
- 15 13. The method of any of the preceding claims, wherein the target tissue is associated with a desired local effect.
14. The method of any of the preceding claims, wherein the target tissue is associated with a desired systemic effect.
20
15. The method of any of the preceding claims, wherein administering results in improved hearing, reduced ear ache, reduced ear infection, reduced pruritus, or reduced tinnitus in the subject.
- 25 16. The method of any of the preceding claims, wherein administering results in reduced vertigo, dizziness or nausea for the subject.
17. The method of any of the preceding claims, wherein the desired local effect involves treatment of a periocular condition, e.g., blepharitis, dry eye, itching, inflammation, wound

healing, intraocular diseases, e.g., macular degeneration, infections, and ocular blood pressure, e.g., glaucoma, elevated intraocular pressure, ocular hypertension, and ocular hypotension.

18. The method of any of the preceding claims, wherein the desired systemic effect involves
5 treatment of one or more of headaches, cardiovascular diseases, inflammation, immune responses and autoimmune disorders, liver diseases, mitochondrial diseases, infections, neurological diseases, psychiatric disorders, nitric oxide disorders, urea cycle disorders, congestion, vasodilation disorders, skin diseases, wound healing, reactions to insect bites, connective tissue disorders, and certain viral, bacterial, and fungal infections.

10 19. A method of treating an ear disorder in a subject, comprising:
administering an effective amount of preparation comprising ammonia oxidizing microorganism (AOM) to the subject,
thereby treating the ear disorder.

15 20. The method of any of the preceding claims, wherein the ear disorder involves vertigo, swimmer's ear, pruritus, or tinnitus.

20 21. The method of any of the preceding claims, wherein the ear disorder comprises an inflammatory condition, a bacterial, viral, or fungal infection, or a congenital abnormality.

22. A method of treating an inflammatory eye disorder in a subject, comprising:
administering to the subject an effective amount of a preparation comprising AOM,
thereby treating the inflammatory eye disorder.

25 23. The method of any of the preceding claims, wherein the inflammatory eye disorder comprises dry eye or Meibomian Gland Dysfunction (MGD).

24. The method of any of the preceding claims, wherein the inflammatory eye disorder
30 comprises blepharitis.

25. The method of any of the preceding claims, wherein the inflammatory eye disorder comprises a disorder of the back of the eye, e.g., of the retina, e.g., macular degeneration.

5 26. The method of any of the preceding claims, wherein the inflammatory eye disorder is associated with intraocular pressure, e.g., elevated intraocular pressure.

27. The method of any of the preceding claims, wherein the inflammatory eye disorder comprises an infectious disorder.

10

28. The method of any of the preceding claims, wherein the inflammatory eye disorder comprises a bacterial, fungal, or viral infection, e.g., of the ocular globe or other eye tissue.

15 29. The method of any of the preceding claims, wherein the inflammatory eye disorder is characterized by or comprises sty or chalazion formation.

30. The method of any of the preceding claims, wherein the inflammatory eye disorder comprises an allergy.

20 31. The method of any of the preceding claims, wherein administering is device-assisted.

32. The method of any of the preceding claims, wherein administering the effective amount of the preparation promotes endothelial function.

25 33. The method of any of the preceding claims, wherein administering the effective amount of the preparation promotes lubrication or hydration of the eye.

30 34. The method of any of the preceding claims, wherein administering the effective amount of the preparation changes or alters a level of nitrite or NO in the subject, e.g. at the target tissue or in circulation.

35. The method of any of the preceding claims, wherein administering the effective amount of the preparation changes or alters a level of nitrite or NO in the subject, e.g. at a target tissue of the eye.

5

36. The method of any of the preceding claims, wherein administering the effective amount of the preparation changes or alters a level of nitrite or NO in the subject, e.g. at a target tissue of the ear.

10 37. The method of any of the preceding claims, wherein administering the effective amount of the preparation modulates a microbiome associated with the subject.

38. The method of any of the preceding claims, wherein administering the effective amount of the preparation modulates a microbiome associated with the eye of the subject.

15

39. The method of any of the preceding claims, wherein administering the effective amount of the preparation modulates a microbiome associated with the ear of the subject.

20 40. The method of any of the preceding claims, wherein the preparation is administered prior to onset of an ear disorder.

41. The method of any of the preceding claims, wherein the preparation is administered during incidence of an ear disorder.

25 42. The method of any of the preceding claims, wherein the preparation is administered subsequent to the subsiding of an ear disorder.

43. The method of any of the preceding claims, wherein the preparation is administered in response to a trigger, warning sign, or symptom of an ear disorder.

30

44. The method of any of the preceding claims, wherein the preparation is administered prior to onset of the inflammatory eye disorder.

45. The method of any of the preceding claims, wherein the preparation is administered during
5 incidence of the inflammatory eye disorder.

46. The method of any of the preceding claims, wherein the preparation is administered subsequent to the subsiding of the inflammatory eye disorder.

10 47. The method of any of the preceding claims, wherein the preparation is administered in response to a trigger, warning sign, or symptom of the inflammatory eye disorder.

48. The method of any of the preceding claims, further comprising determining whether the subject is in need of treatment for an ear disorder.

15 49. The method of any of the preceding claims, further comprising determining whether the subject is in need of treatment for the inflammatory eye disorder.

50. The method of any of the preceding claims, wherein the preparation is administered as a
20 solution, suspension, liquid, ointment, gel, hydrogel, emulsion, or insert.

51. The method of any of the preceding claims, wherein the preparation is administered as a spray, aerosol, or mist.

25 52. The method of any of the preceding claims, wherein the preparation is administered as a droplet or wash.

53. The method of any of the preceding claims, wherein the preparation is formulated to be substantially bio-compatible.

54. The method of any of the preceding claims, wherein the preparation is formulated to be compatible with the eye of the subject.

55. The method of any of the preceding claims, wherein the preparation is substantially isotonic.

5

56. The method of any of the preceding claims, wherein the preparation has a substantially physiological pH level.

57. The method of any of the preceding claims, wherein the preparation has a substantially physiological pH level, e.g., a pH level between about 7.14 and about 7.82.

10

58. The method of any of the preceding claims, wherein the preparation has an osmolality of between about 100 mOsmol/L and about 640 mOsmol/L.

59. The method of any of the preceding claims, wherein the preparation has a surface tension of between about 43 nN/m and about 47 nN/m.

15

60. The method of any of the preceding claims, wherein the preparation is formulated to resemble tears of the subject.

20

61. The method of any of the preceding claims, wherein the preparation is formulated for immediate release or extended release.

62. The method of any of the preceding claims, wherein the preparation or product thereof is formulated to penetrate and otherwise reach a target eye tissue.

25

63. The method of any of the preceding claims, wherein the preparation or product thereof is formulated to permeate or be absorbed through the cornea.

64. The method of any of the preceding claims, wherein the preparation or product thereof is

30

formulated to penetrate and otherwise reach a target ear tissue.

65. The method of any of the preceding claims, wherein the preparation is formulated to deliver nitrite or NO to a target tissue, locally or systemically.

5

66. The method of any of the preceding claims, wherein the preparation is formulated for transmucosal delivery and/or circulation, *e.g.* locally or systemically.

67. The method of any of the preceding claims, wherein the preparation or product thereof is formulated to be delivered for systemic circulation.

10

68. The method of any of the preceding claims, further comprising administering a second amount of the preparation to the subject.

69. The method of any of the preceding claims, wherein the preparation is administered as part of a combination therapy.

15

70. The method of any of the preceding claims, further comprising administering a second treatment in combination with the preparation.

20

71. The method of any of the preceding claims, wherein the second treatment comprises a surgical procedure.

72. The method of any of the preceding claims, wherein the preparation is administered for a period of time prior to initiating the second treatment.

25

73. The method of any of the preceding claims, wherein the preparation is administered concurrently with the second treatment.

74. The method of any of the preceding claims, wherein the preparation is administered for a period of time subsequent to ceasing the second treatment.

75. The method of any of the preceding claims, wherein the second treatment is administered via an alternate mode of administration, e.g. orally or intranasally.

76. The method of any of the preceding claims, wherein the second treatment is administered to the eye, e.g., topically or intravitreally, or via an alternate mode of administration, e.g., orally or intranasally.

77. The method of any of the preceding claims, wherein the subject has a therapeutic level of a second treatment.

78. The method of any of the preceding claims, wherein the preparation is administered in conjunction with one or more ophthalmic ion-exchange formulations, chelators, penetration enhancers, e.g., transporter-target prodrugs, ultrasound, soft drugs, mucoadhesives, or steam, e.g., warm compress.

79. The method of any of the preceding claims, wherein the preparation is administered in conjunction with nitrite, nitrate, and/or NO.

80. The method of any of the preceding claims, wherein the effective amount is a therapeutically effective dose of AOM.

81. The method of any of the preceding claims, wherein the therapeutically effective dose of AOM is about or greater than about 1×10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} CFU.

82. The method of any of the preceding claims, wherein the preparation is administered as an analgesic.

83. The method of any of the preceding claims, wherein the preparation is administered as a prophylactic.

5 84. The method of any of the preceding claims, wherein the preparation is self-administered.

85. The method of any of the preceding claims, wherein the preparation is administered about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 times per day.

10 86. The method of any of the preceding claims, wherein the preparation is administered for about 1-3, 3-5, 5-7, 7-9, 5-10, 10-14, 12-18, 12-21, 21-28, 28-35, 35-42, 42-49, 49-56, 46-63, 63-70, 70-77, 77-84, or 84-91 days.

15 87. The method of any of the preceding claims, wherein the preparation is administered within 30, 60, 90, 120, 150, or 180 minutes of the subject waking from sleep.

88. The method of any of the preceding claims, wherein the preparation is administered within 30, 60, 90, 120, 150, or 180 minutes prior to the subject sleeping.

20 89. The method of any of the preceding claims, wherein the preparation is administered within 30, 60, 90, 120, 150, or 180 minutes of the subject eating.

90. The method of any of the preceding claims, wherein the preparation is administered 30, 60, 90, 120, 150, or 180 minutes before the subject cleanses or showers.

25 91. The method of any of the preceding claims, wherein the subject is female.

92. The method of any of the preceding claims, wherein the subject is male.

93. The method of any of the preceding claims, wherein the subject is characterized as one of the following ethnicity/race: Asian, black or African American, Hispanic or Latino, white, or multi-racial.

5 94. The method of any of the preceding claims, wherein the subject is of an age of less than 1, or between 1-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, or over 60 years.

95. The method of any of the preceding claims, wherein the preparation comprises AOM in a buffer solution, e.g., an aqueous buffer solution.

10

96. The method of any of the preceding claims, wherein the buffer solution, e.g., aqueous buffer solution, comprises disodium phosphate and magnesium chloride, for example, 50 mM Na_2HPO_4 and 2 mM MgCl_2 in water.

15 97. The method of any of the preceding claims, wherein the buffer solution e.g., aqueous buffer solution, consisting essentially of disodium phosphate and magnesium chloride, for example, 50 mM Na_2HPO_4 and 2 mM MgCl_2 in water.

20 98. The method of any of the preceding claims, wherein the buffer solution, e.g., aqueous buffer solution, consists of disodium phosphate and magnesium chloride, for example, 50 mM Na_2HPO_4 and 2 mM MgCl_2 in water.

25 99. The method of any of the preceding claims, wherein the preparation comprises at least one of ammonia, ammonium salts, and urea.

100. The method of any of the preceding claims, wherein the preparation comprises a controlled release material, e.g., slow release material.

30 101. The method of any of the preceding claims, wherein the preparation further comprises an excipient, e.g., a pharmaceutically acceptable excipient.

102. The method of any of the preceding claims, wherein the excipient comprises at least one of water, hydrogen peroxide, mineral oil, vegetable oil, glycerin, anhydrous glycerin, propylene glycol, aluminum acetate, isopropyl alcohol, and ethanol.

5

103. The method of any of the preceding claims, wherein the excipient comprises an absorption and penetration enhancer, preservative, antioxidant, buffer, chelating agent, ion exchange agent, solubilizing agent, suspending agent, thickener, surfactant, wetting agent, tonicity-adjusting agent, and a vehicle for proper drug delivery.

10

104. The method of any of the preceding claims, wherein the excipient comprises a surfactant.

105. The method of any of the preceding claims, wherein the preparation further comprises at least one of an analgesic agent, a chelator, a hygroscopic agent, a ceramide disrupting agent, an anti-inflammatory agent, an anti-infective agent, a reacidifying agent, and a vehicle.

15

106. The method of any of the preceding claims, wherein the preparation is substantially free of other organisms.

20 107. The method of any of the preceding claims, wherein the preparation comprises between about 1×10^3 CFU/mL to about 1×10^{14} CFU/mL AOM.

108. The method of any of the preceding claims, wherein the preparation comprises between about 1×10^9 CFU/mL to about 10×10^9 CFU/mL AOM.

25

109. The method of any of the preceding claims, wherein the preparation is administered before or after a surgical or diagnostic procedure.

110. The method of any of the preceding claims, wherein the AOM comprise ammonia oxidizing bacteria (AOB).

30

111. The method of any of the preceding claims, wherein the AOM consist essentially of AOB.

112. The method of any of the preceding claims, wherein the AOM consist of AOB.

5

113. The method of any of the preceding claims, wherein the AOM comprise *Nitrosomonas*, *Nitrosococcus*, *Nitrospira*, *Nitrosocystis*, *Nitrosolobus*, *Nitrovibrio*, and combinations thereof.

10 114. The method of any of the preceding claims, wherein the AOM is *Nitrosomonas eutropha* (*N. eutropha*).

115. The method of any of the preceding claims, wherein the AOM is *N. eutropha* D23, having ATCC accession number PTA-121157.

15

116. The method of any of the preceding claims, wherein the AOM comprise ammonia oxidizing archaea (AOA).

117. The method of any of the preceding claims, wherein the AOM are capable of converting
20 ammonia or ammonium to nitrite at a rate of at least about 1 pmol/min/mg protein, e.g., at least about 0.1 nmol/min/mg protein.

118. The method of any of the preceding claims, wherein a target percentage of administered AOM are transferred to the ear of the subject.

25

119. The method of any of the preceding claims, wherein the preparation is administered in conjunction with an anti-inflammatory agent.

120. The method of any of the preceding claims, wherein the preparation is administered in conjunction with a medical approach that treats, e.g., is approved to treat or is commonly used to treat, the relevant disease or disorder, or a symptom of the relevant disease or disorder.

5 121. The method of any of the preceding claims, wherein the subject has a disrupted microbiome.

122. The method of any of the preceding claims, wherein the preparation further comprises or is administered concurrently with a compound that promotes growth or metabolism of the AOM,
10 NO production, and/or urease activity.

123. The method of any of the preceding claims, wherein a biome-friendly product is used in connection with the administered preparation comprising AOM.

15 124. A preparation comprising AOM, as recited in any of the preceding claims, for administration to an ear of a subject.

125. A preparation comprising AOM, as recited in any of the preceding claims, for treatment of an ear disorder in a subject.

20 126. A preparation comprising AOM, as recited in any of the preceding claims, for treatment of an inflammatory eye disorder in a subject.

127. The preparation of any of the preceding claims, wherein the preparation is packaged for
25 single use.

128. The preparation of any of the preceding claims, wherein the preparation is packaged for multiple use.

129. The preparation of any of the preceding claims, comprising AOM and other organisms, e.g., a community of organisms.

5 130. A device for administering a preparation comprising AOM, as recited in any of the preceding claims, to an ear of a subject.

131. A device for administering a preparation comprising AOM, as recited in any of the preceding claims, to an eye of a subject.

10 132. A device configured to administer a preparation comprising AOM, as recited in any of the preceding claims, to an ear of a subject.

133. A device configured to administer a preparation comprising AOM, as recited in any of the preceding claims, to an eye of a subject.

15 134. A kit comprising a preparation comprising AOM as recited in any of the preceding claims.

135. The kit of any of the preceding claims, further including instructions for use.

20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/042530

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4-18, 20, 21, 23-135
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/042530

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 8/64; A61K 35/74; A61P 37/00 (2018.01)

CPC - A61J 1/20; A61K 8/99; A61K 31/00; A61K 35/74; A61K 35/741; C12N 1/00 (2018.08)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC - 424/93.4 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2017/0189454 A1 (AOBIOME LLC) 06 July 2017 (06.07.2017) entire document	2, 3
Y		1, 19
Y	US 2011/0274676 A1 (FARMER et al) 10 November 2011 (10.11.2011) entire document	1, 19
Y	US 2014/0037688 A1 (BERKES et al) 06 February 2014 (06.02.2014) entire document	22
Y	US 2017/0037363 A1 (AOBIOME LLC) 09 February 2017 (09.02.2017) entire document	22
P, X	WO 2018/017583 A1 (AOBIOME LLC et al) 25 January 2018 (25.01.2018) entire document	1-3, 19, 22
P, X	WO 2018/111888 A1 (AOBIOME LLC) 21 June 2018 (21.06.2018) entire document	1-3, 19, 22
A	US 2007/0148136 A1 (WHITLOCK) 28 June 2007 (28.06.2007) entire document	1-3, 19, 22
A	US 2011/0232650 A1 (WHITLOCK) 29 September 2011 (29.09.2011) entire document	1-3, 19, 22

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 August 2018

Date of mailing of the international search report

02 OCT 2018

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774