




(86) Date de dépôt PCT/PCT Filing Date: 2014/01/24
(87) Date publication PCT/PCT Publication Date: 2014/07/31
(45) Date de délivrance/Issue Date: 2021/11/23
(85) Entrée phase nationale/National Entry: 2015/07/15
(86) N° demande PCT/PCT Application No.: US 2014/012926
(87) N° publication PCT/PCT Publication No.: 2014/116932
(30) Priorité/Priority: 2013/01/25 (US61/756,906)

(51) Cl.Int./Int.Cl. *A01N 43/90* (2006.01),
A01N 43/40 (2006.01), *A01P 13/00* (2006.01)
(72) Inventeurs/Inventors:
BANGEL, BRYSTON L., US;
SATCHIVI, NORBERT M., US
(73) Propriétaire/Owner:
DOW AGROSCIENCES LLC, US
(74) Agent: SMART & BIGGAR LLP

(54) Titre : COMPOSITION HERBICIDE COMPRENANT DE L'ACIDE 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)PYRIDINE-2-CARBOXYLIQUE ET DU DICLOSULAM OU DU CLORANSULAME-METHYLE
(54) Title: A HERBICIDAL COMPOSITION COMPRISING 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)PYRIDINE-2-CARBOXYLIC ACID AND DICLOSULAM OR CLORANSULAM-METHYL



(I)

(57) Abrégé/Abstract:

Herbicidal compositions and methods using a combination of (a) a compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) diclosulam or clorasulam-methyl or an agriculturally acceptable salt thereof provide control of undesirable vegetation in, e.g., rice, wheat, barley, oats, rye, sorghum, corn or maize, oilseed rape, vegetables, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, industrial vegetation management or rights-of-way.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
31 July 2014 (31.07.2014)(10) International Publication Number
WO 2014/116932 A1

- (51) **International Patent Classification:**
A01N 43/40 (2006.01) A01N 43/54 (2006.01)
- (21) **International Application Number:**
PCT/US2014/012926
- (22) **International Filing Date:**
24 January 2014 (24.01.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/756,906 25 January 2013 (25.01.2013) US
- (71) **Applicant:** DOW AGROSCIENCES LLC [US/US];
9330 Zionsville Road, Indianapolis, IN 46268 (US).
- (72) **Inventors:** BANGEL, Bryston L.; 7610 Reynolds Road,
Camby, IN 46113 (US). SATCHIVI, Norbert M.; 3100
Cherub Court, Carmel, IN 46074 (US).
- (74) **Agent:** ASAM, Michael; Dow AgroSciences LLC, 9330
Zionsville Road, Indianapolis, Indiana 46268 (US).
- (81) **Designated States** (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

- (84) **Designated States** (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))

Published:

- with international search report (Art. 21(3))

(54) **Title:** HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL) PYRIDINE-2-CARBOXYLIC ACID



(I)

(57) **Abstract:** Herbicidal compositions and methods using a combination of (a) a compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) diclosulam or clorasulam-methyl or an agriculturally acceptable salt thereof provide control of undesirable vegetation in, e.g., rice, wheat, barley, oats, rye, sorghum, corn or maize, oilseed rape, vegetables, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, industrial vegetation management or rights-of-way.

WO 2014/116932 A1


A HERBICIDAL COMPOSITION COMPRISING 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL) PYRIDINE-2-CARBOXYLIC ACID AND DICLOSULAM OR CLORANSULAM-METHYL

Background

[0001] The protection of crops from weeds and other vegetation which inhibit crop growth in a constantly recurring problem in agriculture. To help combat this problem, researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use. However, there remains a need for compositions and methods that are effective in controlling undesirable vegetation.

Summary

[0002] Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)




(I)

or an agriculturally acceptable salt or ester of thereof and (b) cloransulam-methyl or diclosulam or an agriculturally acceptable salt thereof, wherein the combination of (a) and (b) exhibits synergism. The compositions may also contain an agriculturally acceptable adjuvant or carrier. In one embodiment, no additional herbicidal active ingredients are present in the composition.

[0003] Also provided are methods of controlling undesirable vegetation comprising applying a herbicidally effective amount of a herbicidal composition comprising (a) a compound of formula (I) or an agriculturally acceptable ester or salt thereof and (b)

cloransulam-methyl or diclosulam or an agriculturally acceptable salt thereof, wherein the combination of (a) and (b) exhibits synergism.

[0003a] Also provided herein is a herbicidal composition comprising a herbicidally effective amount of (a) a compound of the formula (I)



(I)


or an agriculturally acceptable salt or ester thereof and (b) diclosulam or an agriculturally acceptable salt thereof, wherein the combination of (a) and (b) exhibits synergism.

[0003b] Also provided herein is the composition as described herein, further comprising an agriculturally acceptable adjuvant or carrier.

Detailed Description

DEFINITIONS

[0004] As used herein, the compound of formula (I) has the following structure:



(I)

[0005] The compound of formula (I) can be identified by the name 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid and has been described in U.S. Patent 7,314,849 (B2). Exemplary uses of the compound of the formula (I) include controlling undesirable vegetation, including *e.g.*, grass, broadleaf and sedge weeds, in multiple non-crop and cropping situations.

[0006] Cloransulam-methyl or diclosulam are both triazolopyrimidine sulfonamides. As used herein, triazolopyrimidine sulfonamides are a chemical class of herbicides having a triazolopyrimidine sulfonamide core structure. Without being limited to any theory, their mode-of-action is believed to involve the inhibition of acetolactate synthase (ALS), an enzyme common to plants and microorganisms but not found in animals. Exemplary herbicidal uses of triazolopyrimidine sulfonamides include, but are not limited to, use for control of nuisance sedge, broadleaf and grass weeds.

[0007] As used herein, cloransulam-methyl is methyl 3-chloro-2-[[5-ethoxy-7-fluoro[1,2,4]triazolo[1,5-*c*]pyrimidin-2-yl)sulfonyl]amino]benzoate. Its herbicidal activity is exemplified in Tomlin, C. D. S., Ed. *The Pesticide Manual: A World Compendium*, 15th ed.; BCPC: Alton, 2009 (hereafter "*The Pesticide Manual*"). Exemplary uses of cloransulam-methyl include its use as a herbicide for post-emergence control of broadleaf weeds in soybeans and other broadleaf crops.

[0008] As used herein, diclosulam is *N*-(2,6-dichlorophenyl)-5-ethoxy-7-fluoro[1,2,4]triazolo[1,5-*c*]pyrimidine-2-sulfonamide. Its herbicidal activity is exemplified in *The Pesticide Manual*. Exemplary uses of diclosulam include its use as a herbicide for broadleaf weed control in peanuts and soybeans.

[0009] As used herein, control of or controlling undesirable vegetation means killing or preventing the vegetation, or causing some other adverse modifying effect to the vegetation

e.g., deviations from natural growth or development, regulation, desiccation, retardation, and the like.

[0010] As used herein, herbicide and herbicidal active ingredient mean a compound that controls undesirable vegetation when applied in an appropriate amount.


[0011] As used herein, a herbicidally effective or vegetation controlling amount is an amount of herbicidal active ingredient the application of which controls the relevant undesirable vegetation.

[0012] As used herein, applying a herbicide or herbicidal composition means delivering it directly to the targeted vegetation or to the locus thereof or to the area where control of undesired vegetation is desired. Methods of application include, but are not limited to pre-emergence, post-emergence, foliar, soil, and in-water applications. Described herein are methods of controlling undesirable vegetation by applying certain herbicide combinations or compositions.

[0013] As used herein, plants and vegetation include, but are not limited to, dormant seeds, germinant seeds, emerging seedlings, plants emerging from vegetative propagules, immature vegetation, and established vegetation.

[0014] As used herein, agriculturally acceptable salts and esters refer to salts and esters that exhibit herbicidal activity, or that are or can be converted in plants, water, or soil to the referenced herbicide. Exemplary agriculturally acceptable esters are those that are or can be hydrolyzed, oxidized, metabolized, or otherwise converted, *e.g.*, in plants, water, or soil, to the corresponding carboxylic acid which, depending on the pH, may be in the dissociated or undissociated form.

[0015] Exemplary salts include those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Exemplary cations include sodium, potassium, magnesium, and ammonium cations of the formula:



wherein R^1 , R^2 , R^3 and R^4 each, independently represents hydrogen or C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl or C_3 - C_{12} alkynyl, each of which is optionally substituted by one or more hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio or phenyl groups, provided that R^1 , R^2 , R^3 and R^4 are sterically compatible. Additionally, any two of R^1 , R^2 , R^3 and R^4 together may represent an aliphatic difunctional moiety containing one to twelve carbon atoms and up to two oxygen or sulfur atoms. Salts can be prepared by treatment with a metal hydroxide, such as sodium hydroxide,


with an amine, such as ammonia, trimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, morpholine, cyclododecylamine, or benzylamine or with a tetraalkylammonium hydroxide, such as tetramethylammonium hydroxide or choline hydroxide.

[0016] Exemplary esters include those derived from C₁-C₁₂ alkyl, C₃-C₁₂ alkenyl, C₃-C₁₂ alkynyl or C₇-C₁₀ aryl-substituted alkyl alcohols, such as methyl alcohol, isopropyl alcohol, 1-butanol, 2-ethylhexanol, butoxyethanol, methoxypropanol, allyl alcohol, propargyl alcohol, cyclohexanol or unsubstituted or substituted benzyl alcohols. Benzyl alcohols may be substituted with from 1-3 substituents independently selected from halogen, C₁-C₄ alkyl or C₁-C₄ alkoxy. Esters can be prepared by coupling of the acids with the alcohol using any number of suitable activating agents such as those used for peptide couplings such as dicyclohexylcarbodiimide (DCC) or carbonyl diimidazole (CDI); by reacting the acids with alkylating agents such as alkylhalides or alkylsulfonates in the presence of a base such as triethylamine or lithium carbonate; by reacting the corresponding acid chloride of an acid with an appropriate alcohol; by reacting the corresponding acid with an appropriate alcohol in the presence of an acid catalyst or by transesterification.

[0017] As used herein, weight ratios of mixtures are calculated using the acid equivalent weight(s) of any compounds in the mixture that are salts or esters.

COMPOSITIONS AND METHODS

[0018] Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)



(I)

or an agriculturally acceptable salt or ester of thereof and (b) cloransulam-methyl or diclosulam.

[0019] Also provided herein are methods of controlling undesirable vegetation comprising applying a herbicidally effective amount of the compound of formula (I) or

agriculturally acceptable salt or ester thereof and (b) cloransulam-methyl or diclosulam. In certain embodiments, the methods employ the compositions described herein.

[0020] Furthermore, in some embodiments, the combination of compound (I) or agriculturally acceptable salt or ester thereof and cloransulam-methyl or diclosulam exhibits synergism, *i.e.*, the herbicidal active ingredients are more effective in combination than when applied individually. Synergism has been defined as “an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response of each factor applied separately.” Senseman, S., Ed. *Herbicide Handbook*. 9th ed.

Lawrence: Weed Science Society of America, 2007. In certain embodiments, the compositions exhibit synergy as determined by Colby's equation (Colby, S.R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds* **1967**, 15, 20-22.

[0021] In certain embodiments of the compositions and methods described herein, the compound of formula (I), *i.e.*, the carboxylic acid, is employed. In certain embodiments, a carboxylate salt of the compound of formula (I) is employed. In certain embodiments, an aralkyl or alkyl ester is employed. In certain embodiments, a benzyl, substituted benzyl, or C₁-C₄ alkyl, *e.g.*, *n*-butyl ester is employed. In certain embodiments, the methyl ester, benzyl ester, or potassium salt is employed.

[0022] In some embodiments, the compound of formula (I) or salt or ester thereof and cloransulam-methyl or diclosulam are formulated in one composition, tank-mixed, applied simultaneously, or applied sequentially.

[0023] Herbicidal activity (control of undesirable vegetation) is exhibited by the herbicidal compositions when they are applied directly to the plant or to the locus of the plant at any stage of growth, or to the area where control of vegetation is desired. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted to promote non-selective or selective herbicidal action. In some embodiments, the compositions described herein are applied as a post-emergence application, pre-emergence application, or in-water application to flooded paddy rice or water bodies (*e.g.*, ponds, lakes and streams), to relatively immature undesirable vegetation to achieve the maximum control of weeds.

[0024] In some embodiments, the compositions and methods provided herein are utilized to control weeds in crops, including but not limited to winter/spring oilseed rape, winter/spring canola, vegetables, *Brassica* spp, ornamentals, rice, wheat, triticale, barley, oats, rye, sorghum, corn/maize, sunflower, row crops, pastures, grasslands, rangelands, fallowland, sugarcane, turf, tree and vine orchards, aquatics, and industrial vegetation management (IVM) and rights-of-way.

[0025] The compositions and methods provided herein are utilized to control undesirable vegetation. Undesirable vegetation includes, but is not limited to, undesirable vegetation that occurs in oilseed rape, canola, vegetables, *Brassica* spp, ornamentals, rice, wheat, triticale, barley, oats, rye, sorghum, corn/maize, sunflower, row crops, pastures, grasslands, rangelands, fallowland, sugarcane, turf, tree and vine orchards, IVM and rights-of-way.

[0026] In some embodiments, the methods provided herein are utilized to control undesirable vegetation in oilseed rape, canola, drilled crops and cereal crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Apera spica-venti* (L.) Beauv. (windgrass, APESV), *Avena fatua* L. (wild oat, AVEFA), *Bromus tectorum* L. (downy brome, BROTE), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Lolium rigidum* (rigid ryegrass, LOLRI), *Lolium multiflorum subsp. Gaudini* (annual ryegrass, LOLMG), *Phalaris minor* Retz. (littleseed canarygrass, PHAMI), *Poa annua* L. (annual bluegrass, POAAN), *Setaria pumila* (Poir.) Roemer & J.A. Schultes (yellow foxtail, SETLU), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Galium aparine* L. (catchweed bedstraw, GALAP), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMPU), *Matricaria recutita* L. (wild chamomile, MATCH), *Matricaria matricarioides* (Less.) Porter (pineappleweed, MATMT), *Papaver rhoeas* L. (common poppy, PAPRH), *Polygonum convolvulus* L. (wild buckwheat, POLCO), *Salsola tragus* L. (Russian thistle, SASKR), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Veronica persica* Poir. (Persian speedwell, VERPE), *Viola arvensis* Murr. (field violet, VIOAR), or *Viola tricolor* L. (wild violet, VIOTR).

[0027] In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation in rice. In certain embodiments, the undesirable vegetation is *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv.

(barnyardgrass, ECHCG), *Echinochloa colonum* (L.) LINK (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Echinochloa oryzicola* (Vasinger) Vasinger (late watergrass, ECHPH), *Ischaemum rugosum* Salisb. (saramollagrass, ISCRU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa fascicularis* (Lam.) Gray (bearded sprangletop, LEFFA), *Leptochloa panicoides* (Presl.) Hitchc. (Amazon sprangletop, LEFPA), *Panicum dichotomiflorum* (L.) Michx. (Fall panicum, PANDI), *Paspalum dilatatum* Poir. (dallisgrass, PASDI), *Cyperus difformis* L. (smallflower flatsedge, CYPDI), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus iria* L. (rice flatsedge, CYPPIR), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Eleocharis* species (ELOSS), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Schoenoplectus juncooides* Roxb. (Japanese bulrush, SPCJU), *Schoenoplectus maritimus* L. (sea clubrush, SCPMA), *Schoenoplectus mucronatus* L. (riccfield bulrush, SCPMU), *Aeschynomene* species, (jointvetch, AESSS), *Alternanthera philoxeroides* (Mart.) Griseb. (alligatorweed, ALRPH), *Alisma plantago-aquatica* L. (common waterplantain, ALSPA), *Amaranthus* species, (pigweeds and amaranths, AMASS), *Ammannia coccinea* Rottb. (redstem, AMMCO), *Eclipta alba* (L.) Hassk. (American false daisy, ECLAL), *Heteranthera limosa* (SW.) Willd./Vahl (ducksalad, HETLI), *Heteranthera reniformis* R. & P. (roundleaf mudplantain, HETRE), *Ipomoea hederacea* (L.) Jacq. (ivy leaf morningglory, IPOHE), *Lindernia dubia* (L.) Pennell (low false pimpernel, LIDDU), *Monochoria korsakowii* Regel & Maack (monochoria, MOOKA), *Monochoria vaginalis* (Burm. F.) C. Presl ex Kuhth, (monochoria, MOOVA), *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Polygonum pensylvanicum* L., (Pennsylvania smartweed, POLPY), *Polygonum persicaria* L. (ladysthumb, POLPE), *Polygonum hydropiperoides* Michx. (mild smartweed, POLHP), *Rotala indica* (Willd.) Koehne (Indian toothcup, ROTIN), *Sagittaria* species, (arrowhead, SAGSS), *Sesbania exaltata* (Raf.) Cory/Rydb. Ex Hill (hemp sesbania, SEBEX), or *Sphenoclea zeylanica* Gaertn. (gooseweed, SPDZE).

[0028] In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation in range and pasture. In certain embodiments, the undesirable vegetation is *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Cassia obtusifolia* (sickle pod, CASOB), *Centaurea maculosa* auct. non Lam. (spotted knapweed, CENMA), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Convolvulus arvensis* L. (field bindweed, CONAR), *Euphorbia esula* L. (leafy spurge, EPHEs), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Plantago lanceolata* L. (buckhorn plantain, PLALA),

Rumex obtusifolius L. (broadleaf dock, RUMOB), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Sonchus arvensis* L. (perennial sowthistle, SONAR), *Solidago* species (goldenrod, SOOSS), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Urtica dioica* L. (common nettle, URTDI).

[0029] In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation found in row crops and vegetable crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Avena fatua* L. (wild oat, AVEFA), *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) Link (junglerice, ECHCO), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Panicum dichotomiflorum* Michx. (Fall panicum, PANDI), *Panicum miliaceum* L. (wild-proso millet, PANMI), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Sorghum halepense* (L.) Pers. (Johnsongrass, SORHA), *Sorghum bicolor* (L.) Moench ssp. *Arundinaceum* (shattercane, SORVU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Amaranthus* species (pigweeds and amaranths, AMASS), *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Ambrosia psilostachya* DC. (Western ragweed, AMBPS), *Ambrosia trifida* L. (giant ragweed, AMBTR), *Asclepias syriaca* L. (common milkweed, ASCSY), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Commelina benghalensis* L. (tropical spiderwort, COMBE), *Datura stramonium* L. (jimsonweed, DATST), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL), *Erigeron bonariensis* L. (hairy fleabane, ERIBO), *Erigeron canadensis* L. (Canadian fleabane, ERICA), *Helianthus annuus* L. (common sunflower, HELAN), *Jacquemontia taminifolia* (L.) Griseb. (smallflower morningglory, IAQTA), *Ipomoea hederacea* (L.) Jacq. (ivy leaf morningglory, IPOHE), *Ipomoea lacunosa* L. (white morningglory, IPOLA), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Portulaca oleracea* L. (common purslane, POROL), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Solanum ptychanthum* Dunal (eastern black nightshade, SOLPT), or *Xanthium strumarium* L. (common cocklebur, XANST).

[0030] In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation consisting of grass, broadleaf and sedge weeds.

[0031] In some embodiments, the combination of compound (I) or agriculturally acceptable ester or salt thereof and cloransulam-methyl or diclosulam or agriculturally acceptable salt or ester thereof are used to control *Amaranthus retroflexus* (redroot pigweed, AMARE), *Chenopodium album* (common lambsquarters, CHEAL), *Centaurea cyanus* (cornflower, CENCY), *Descurainia sophia* (flixweed, DESSO), *Conyza canadensis* (horseweed / marestail, ERICA), *Conyza bonariensis* (fleabane, ERIBO), *Erodium cicutarium* (storksbill / redstem filaree, EROCI), *Fumaria officinalis* (common fumitory, FUMOF), *Galeopsis tetrahit* (common hempnettle, GAETE), *Galium aparine* (edstraw, catchweed / cleavers, GALAP), *Geranium dissectum* (cutleaf geranium, GERDI), *Geranium pusillum* (smallflower geranium, GERPU), *Glycine max* (volunteer soybean, GLXMA), *Lamium amplexicaule* (henbit, LAMAM), *Lamium purpureum* (purple deadnettle, LAMPU), *Papaver rhoeas* (common poppy, PAPRH), *Stellaria media* (common chickweed, STEME), *Veronica persica* (Persian speedwell, VERPE), *Linum usitatissimum* (volunteer flax, LIUUT), *Geranium carolinianum* (Carolina geranium, GERCA), or *Vicia villosa* (hairy vetch, VICVI).

[0032] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with cloransulam-methyl. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to cloransulam-methyl is within the range from about 1:35 to about 5:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to cloransulam-methyl is within the range from about 1:20 to about 1.25:1. In certain embodiments, the compositions provided herein comprise the methyl ester of the compound of formula (I) and cloransulam-methyl. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and cloransulam-methyl, wherein the weight ratio of the compound of formula (I) to cloransulam-methyl is from about 1:35 to about 5:1. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and cloransulam-methyl, wherein the weight ratio of the methyl ester of the compound of formula (I) to cloransulam-methyl is from about 1:20 to about 1.25:1. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and cloransulam-methyl, wherein the weight ratio of the methyl ester of the compound of formula (I) to cloransulam-methyl is from about 1:14 to about 4:3.5.

[0033] With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 3 grams active ingredient per hectare (g ai/ha) to about 45 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 6 g ai/ha to about 25 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and cloransulam-methyl, *e.g.*, sequentially or simultaneously. In some embodiments, the cloransulam-methyl is applied at a rate from about 2 g ai/ha to about 35 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1 grams acid equivalent per hectare (g ae/ha) to about 10 g ae/ha. In some embodiments, the cloransulam-methyl is applied at a rate from about 4.375 g ai/ha to about 17.5 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 5 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its methyl ester and cloransulam-methyl. In one embodiment, the methods utilize the methyl ester of the compound of formula (I) and cloransulam-methyl, wherein the methyl ester of the compound of formula (I) is applied at a rate from about 1.25 g ae/ha to about 5 g ae/ha, and cloransulam-methyl is applied at a rate from about 4.375 g ai/ha to about 17.5 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with cloransulam-methyl are used to control VIOTR, STEME, SORVU, or IPOHE.

[0034] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with diclosulam. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to diclosulam is within the range from about 1:35 to about 5:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to diclosulam is within the range from about 1:20 to about 1.25:1. In certain embodiments, the compositions provided herein comprise the methyl ester of the compound of formula (I) and diclosulam. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and diclosulam, wherein the weight ratio of the

compound of formula (I) to diclosulam is from about 1:35 to about 5:1. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and diclosulam, wherein the weight ratio of the methyl ester of the compound of formula (I) to diclosulam is from about 1:20 to about 1.25:1. In one embodiment, the composition comprises the methyl ester of the compound of formula (I) and diclosulam, wherein the weight ratio of the methyl ester of the compound of formula (I) to diclosulam is from about 1:14 to about 4:3.5.

[0035] With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 3 grams active ingredient per hectare (g ai/ha) to about 45 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 5.6 g ai/ha to about 22.5 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and diclosulam, *e.g.*, sequentially or simultaneously. In some embodiments, the diclosulam is applied at a rate from about 2 g ai/ha to about 35 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1 grams acid equivalent per hectare (g ae/ha) to about 10 g ae/ha. In some embodiments, the diclosulam is applied at a rate from about 4.375 g ai/ha to about 17.5 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 5 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its methyl ester and diclosulam. In one embodiment, the methods utilize the methyl ester of the compound of formula (I) and diclosulam, wherein the methyl ester of the compound of formula (I) is applied at a rate from about 1.25 g ae/ha to about 5 g ae/ha, and diclosulam is applied at a rate from about 4.375 g ai/ha to about 17.5 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with diclosulam are used to control VIOTR, STEME, SORVU or IPOHE.

[0036] The components of the mixtures described herein can be applied either separately or as part of a multipart herbicidal system. In some embodiments of the methods described herein, the active ingredients are applied simultaneously, including, *e.g.*, in the form of a composition. In some embodiments, the active ingredients are applied sequentially, *e.g.*,

within 5, 10, 15, or 30 minutes of each other; 1, 2, 3, 4, 5, 10, 12, 24, 48 hour(s) of each other; or 1 week of each other.

[0037] The mixtures described herein can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank-mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the compositions and methods described herein include, but are not limited to: 4-CPA, 4-CPB, 4-CPP, 2,4-D, 2,4-D choline salt, 2,4-D esters and amines, 2,4-DB, 3,4-DA, 3,4-DB, 2,4-DEB, 2,4-DEP, 3,4-DP, 2,3,6-TBA, 2,4,5-T, 2,4,5-TB, acetochlor, acifluorfen, aclonifen, acrolein, alachlor, allidochlor, alloxydim, allyl alcohol, alorac, ametrudione, ametryn, amibuzin, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amiprofos-methyl, amitrole, ammonium sulfamate, anilofos, anisuron, asulam, atraton, atrazine, azafenidin, azimsulfuron, aziprotryne, barban, BCPC, beflubutamid, benazolin, bencarbazone, benfluralin, benfuresate, bensulfuron-methyl, bensulide, benthocarb, bentazon-sodium, benzadox, benzfendizone, benzipram, benzobicyclon, benzofenap, benzofluor, benzoylprop, benzthiazuron, bicyclopiron, bifenox, bilanafos, bispyribac-sodium, borax, bromacil, bromobonil, bromobutide, bromofenoxim, bromoxynil, brompyrazon, butachlor, butafenacil, butamifos, butenachlor, buthidazole, buthiuron, butralin, butoxydim, buturon, butylate, cacodylic acid, cafenstrole, calcium chlorate, calcium cyanamide, cambendichlor, carbasulam, carbetamide, carboxazole, chlorprocarb, carfentrazone-ethyl, CDEA, CEPC, chlomethoxyfen, chloramben, chloranocryl, chlorazifop, chlorazine, chlorbromuron, chlorbufam, chloreturon, chlorfenac, chlorfenprop, chlorflurazole, chlorflurenol, chloridazon, chlorimuron, chlornitrofen, chloropon, chlorotoluron, chloroxuron, chloroxynil, chlorpropham, chlorsulfuron, chlorthal, chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, cisanilide, clethodim, cliodinate, clodinafop-propargyl, clofop, clomazone, clomeprop, cloprop, cloproxydim, clopyralid, CMA, copper sulfate, CPMF, CPPC, credazine, cresol, cumyluron, cyanatryn, cyanazine, cycloate, cyclosulfamuron, cycloxydim, cycluron, cyhalofop-butyl, cyperquat, cyprazine, cyprazole, cypromid, daimuron, dalapon, dazomet, delachlor, desmedipham, desmetryn, di-allate, dicamba, dichlobenil, dichloralurea, dichlormate, dichlorprop, dichlorprop-P, diclofop-methyl, diethamquat, diethatyl, difenopenten, difenoxuron, difenzoquat, diflufenican, diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid,

dimethenamid-P, dimexano, dimidazon, dinitramine, dinofenate, dinoprop, dinosam, dinoseb, dinoterb, diphenamid, dipropetryn, diquat, disul, dithiopyr, diuron, DMPA, DNOC, DSMA, EBEP, eglinazine, endothal, epronaz, EPTC, erbon, esprocarb, ethalfluralin, ethbenzamide, ethametsulfuron, ethidimuron, ethiolate, ethobenzanid, ethofumesate, ethoxyfen, ethoxysulfuron, etinofen, etnipromid, etobenzanid, EXD, fenasulam, fenoprop, fenoxaprop, fenoxaprop-P-ethyl, fenoxaprop-P-ethyl + isoxadifen-ethyl, fenoxasulfone, fenteracol, fenthiaprop, fentrazamide, fenuron, ferrous sulfate, flamprop, flamprop-M, flazasulfuron, fluazifop, fluazifop-P-butyl, fluazolate, flucarbazone, flucetosulfuron, fluchloralin, flufenacet, flufenican, flufenpyr-ethyl, flumezin, flumiclorac-pentyl, flumioxazin, flumipropyln, fluometuron, fluorodifen, fluoroglycofen, fluoromidine, fluoronitrofen, fluothiuron, flupoxam, flupropacil, flupropanate, flupyrsulfuron, fluridone, flurochloridone, fluroxypry, flurtamone, fluthiacet, fomesafen, foramsulfuron, fosaminc, fumiclorac, furyloxyfen, glufosinate, glufosinate-ammonium, glufosinate-P-ammonium, glyphosate, halosafen, halosulfuron-methyl, haloxydine, haloxyfop-methyl, haloxyfop-P-methyl, hexachloroacetone, hexaflurate, hexazinone, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, indanofan, indaziflam, iodobonil, iodomethane, iodosulfuron, iodosulfuron-ethyl-sodium, iofensulfuron, ioxynil, ipazine, ipfencarbazone, iprymidam, isocarbamid, isocil, isomethiozin, isonoruron, isopolinate, isopropalin, isoproturon, isouron, isoxaben, isoxachlortole, isoxaflutole, isoxapyrifop, karbutilate, ketospiradox, lactofen, lenacil, linuron, MAA, MAMA, MCPA esters and amines, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P, medinoterb, mefenacet, mefluidide, mesoprazine, mesosulfuron, mesotrione, metam, metamifop, metamidron, metazachlor, metazosulfuron, metflurazon, methabenzthiazuron, methalpropalin, methazole, methiobencarb, methiozolin, methiuron, methometon, methoprotryne, methyl bromide, methyl isothiocyanate, methyl dymron, metobenzuron, metobromuron, metolachlor, metoxuron, metribuzin, metsulfuron, metsulfuron-methyl, molinate, monalide, monisouron, monochloroacetic acid, monolinuron, monuron, morfamquat, MSMA, naproanilide, napropamide, napropamide-M, naptalam, neburon, nicosulfuron, nipyraclufen, nitralin, nitrofen, nitrofluorfen, norflurazon, noruron, OCH, orbencarb, *ortho*-dichlorobenzene, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxapyrazon, oxasulfuron, oxaziclomefone, oxyfluorfen, paraflufen-ethyl, parafluron, paraquat, pebulate, pelargonic acid, pendimethalin, pentachlorophenol, pentanochlor, pentoxazone, perfluidone, pethoxamid, phenisopham, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenylmercury acetate, picloram,

picolinafen, pinoxaden, piperophos, potassium arsenite, potassium azide, potassium cyanate, pretilachlor, primisulfuron-methyl, procyzazine, prodiamine, profluazol, profluralin, profoxydim, proglinazine, prohexadione-calcium, prometon, prometryn, pronamide, propachlor, propanil, propaquizafop, propazine, propham, propisochlor, propoxycarbazone, propyrisulfuron, propyzamide, prosulfalin, prosulfocarb, prosulfuron, proxan, prynachlor, pydanon, pyracilonil, pyraflufen-ethyl, pyrasulfotole, pyrazogyl, pyrazolynate, pyrazosulfuron-ethyl, pyrazoxyfen, pyribenzoxim, pyributicarb, pyriclor, pyridafol, pyridate, pyrifitalid, pyriminobac, pyrimisulfan, pyrithiobac-sodium, pyroxasulfone, quinclorac, quinmerac, quinochlor, quinonamid, quizalofop, quizalofop-P-ethyl, rhodethanil, rimsulfuron, saflufenacil, S-metolachlor, sebuthylazine, secbumeton, sethoxydim, siduron, simazine, simeton, simetryn, SMA, sodium arsenite, sodium azide, sodium chlorate, sulcotrione, sulfallate, sulfentrazone, sulfometuron, sulfosate, sulfosulfuron, sulfuric acid, sulglycaphin, swep, TCA, tebutam, tebuthiuron, tefuryltrione, tembotrione, tepraloxym, terbacil, terbucarb, terbuchlor, terbumeton, terbuthylazine, terbutryn, tetrafluron, thenylchlor, thiazafluron, thiazopyr, thidiazimin, thidiazuron, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb, tiocarbazil, tioclorim, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tribenuron-methyl, tricamba, triclopyr choline salt, triclopyr esters and salts, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflusulfuron, trifop, trifopsime, trihydroxytriazine, trimeturon, tripropindan, tritac tritosulfuron, vernolate, xylachlor and salts, esters, optically active isomers and mixtures thereof.

[0038] In some embodiments the methods provided herein are used to control undesirable vegetation in crops that are tolerant to glyphosate, glufosinate, dicamba, phenoxy auxins, pyridyloxy auxins, aryloxyphenoxypropionates, acetyl CoA carboxylase (ACCase) inhibitors, imidazolinones, acetolactate synthase (ALS) inhibitors, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, triazines, or bromoxynil. Such herbicide tolerant crops may possess multiple or stacked traits conferring tolerance to multiple herbicides or multiple modes-of-action.

[0039] In some embodiments the methods provided herein are used to control undesirable vegetation that is a herbicide resistant or tolerant weed. Such herbicide resistant or tolerant weed may have a biotype with resistance or tolerance to multiple herbicides, multiple chemical classes, or multiple herbicide modes-of-action. For example, the herbicide resistant or tolerant weed may have a biotype resistant or tolerant to acetolactate synthase

(ALS) inhibitors, photosystem II inhibitors, acetyl CoA carboxylase (ACCase) inhibitors, synthetic auxins, photosystem I inhibitors, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, microtubule assembly inhibitors, lipid synthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, carotenoid biosynthesis inhibitors, very long chain fatty acid (VLCFA) inhibitors, phytoene desaturase (PDS) inhibitors, glutamine synthetase inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors, mitosis inhibitors, cellulose biosynthesis inhibitors, herbicides with multiple modes-of-action, quinclorac, arylaminopropionic acids, difenzoquat, endothall, or organoarsenicals.

[0040] In some embodiments, the compositions described herein are employed in combination with one or more herbicide safeners, such as AD-67 (MON 4660), benoxacor, benthocarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlormid, dicyclonon, dimepiperate, disulfoton, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, jiecaowan, jiecaoxi, mefenpyr-diethyl, mephenate, naphthalic anhydride (NA), oxabetrinil, R29148, 1-[4-(*N*-(2-methoxybenzoyl)sulfamoyl)phenyl]-3-methylurea, *N*-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity. In some embodiments, the safeners are employed in rice, cereal, corn, or maize settings. In some embodiments, the safener is cloquintocet or an ester or salt thereof. In certain embodiments, cloquintocet is utilized to antagonize harmful effects of the compositions on rice and cereals. In some embodiments, the safener is cloquintocet (mexyl).

[0041] In some embodiments, compositions provided herein further comprise at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water-dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions. They can also be provided as a pre-mix or tank-mixed.

[0042] Suitable agricultural adjuvants and carriers include, but are not limited to, crop oil concentrate; nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium

salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C₉-C₁₁ alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C₁₂-C₁₆) ethoxylate; di-*sec*-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

[0043] Liquid carriers that can be employed include water and organic solvents. The organic solvents include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, *n*-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include, but are not limited to toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. In certain embodiments, water is the carrier for the dilution of concentrates.

[0044] Suitable solid carriers include but are not limited to talc, pyrophyllite clay, silica, attapulgus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, cellulose, and the like.

[0045] In some embodiments, the compositions described herein further comprise one or more surface-active agents. In some embodiments, such surface-active agents are employed in both solid and liquid compositions, and in certain embodiments those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants which may also be used in the present formulations are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual," MC

Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Surface-active agents include, but are not limited to salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkyl-naphthalene-sulfonate salts, such as sodium dibutyl-naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable or seed oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, and in certain embodiments, methyl esters.

[0046] In some embodiments, these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

[0047] Other exemplary additives for use in the compositions provided herein include but are not limited to compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

[0048] In some embodiments, the concentration of the active ingredients in the compositions described herein is from about 0.0005 to 98 percent by weight. In some embodiments, the concentration is from about 0.0006 to 90 percent by weight. In compositions designed to be employed as concentrates, the active ingredients, in certain embodiments, are present in a concentration from about 0.1 to 98 weight percent, and in certain embodiments about 0.5 to 90 weight percent. Such compositions are, in certain embodiments, diluted with an inert carrier, such as water, before application. The diluted

compositions usually applied to weeds or the locus of weeds contain, in certain embodiments, about 0.0003 to 1.5 weight percent active ingredient and in certain embodiments contain about 0.0008 to 1.0 weight percent.

[0049] The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation or paddy water, and by other conventional means known to those skilled in the art.

[0050] The described embodiments and following examples are for illustrative purposes and are not intended to limit the scope of the claims. Other modifications, uses, or combinations with respect to the compositions described herein will be apparent to a person of ordinary skill in the art without departing from the spirit and scope of the claimed subject matter.

EXAMPLES

[0051] Evaluation of Postemergent Herbicidal Activity. Seeds or nutlets of the desired test plant species were planted in Sun Gro Metro-Mix® 360 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of about 30 percent, in plastic pots with a surface area of 64 square centimeters (cm²). When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 7-21 days (d) in a greenhouse with an approximate 15 hour (h) photoperiod which was maintained at about 23-29 °C during the day and 22-28 °C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the first or second true leaf stage.

[0052] A weighed amount, determined by the highest rate to be tested, of each test compound was placed in a 25 milliliter (mL) glass vial and was dissolved in 4 mL of a 97:3 volume per volume (v/v) mixture of acetone and dimethyl sulfoxide (DMSO) to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions obtained were diluted with 20 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Aplus 411F crop oil concentrate, and Triton® X-155 surfactant in a 48.5:39:10:1.5:1.0:0.02 v/v ratio to obtain spray solutions containing the highest application rates. Additional application rates were obtained by serial dilution of 12 mL of the high rate solution into a solution containing 2 mL of a 97:3 v/v mixture of acetone and DMSO and 10 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Aplus 411F crop oil concentrate, and Triton X-155 surfactant in a 48.5:39:10:1.5:1.0:0.02 v/v ratio to obtain 1/2X, 1/4X, 1/8X and 1/16X rates of the high rate. Compound requirements are based upon a 12 mL application volume at a rate of 187 liters per hectare (L/ha). Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters (m²) at a spray height of 18 inches (43 cm) above the average plant canopy height. Control plants were sprayed in the same manner with the solvent blank.

[0053] Treatments consisted of the methyl ester of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid (Cmpd 1) as technical grade material and cloransulam-methyl (FirstRate®) and diclosulam (Strongarm®) alone and in combination.

The form of compound of formula (I) was applied on an acid equivalent basis, and cloransulam-methyl and diclosulam were applied on an active ingredient basis.

[0054] The treated plants and control plants were placed in a greenhouse as described above and watered by subirrigation to prevent wash-off of the test compounds. After 14 d, the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill. Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 1 and 2.

[0055] Results in Tables 1 and 2 are greenhouse trial results for foliar applied compositions. The values reported are percent (%) control visually rated. Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds* **1967**, *15*, 20-22.). More specifically, the following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture;

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The compositions tested, application rates employed, plant species tested, and results are given in Tables 1 and 2.

The following abbreviations are used in Tables 1 and 2:

BRSNW *Brassica napus* (winter oilseed rape)

CHEAL *Chenopodium album* L. (common lambsquarters)

VIOTR *Viola tricolor* (L.) (wild pansy)

STEME *Stellaria media* (L.) Vill. (common chickweed)

SETFA *Setaria faberi* Herrm. (giant foxtail)

SORVU *Sorghum vulgare* (common sorghum)

AMARE *Amaranthus retroflexus* L. (redroot pigweed)

CYPES *Cyperus esculentus* L. (yellow nutsedge)

IPOHE *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory)

g ae/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value of percent (%) control rated visually

Exp = expected value of percent (%) control as calculated by Colby's equation

Cmpd 1 = methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylate

Table 1: Synergistic combination of Cmpd 1 and cloransulam-methyl.


Application Rate		BRSNW		CHEAL		VIOTR		STEME		SETFA		SORVU		AMARE		IPOHE	
Cmpd 1 (g ae/ha)	Cloransulam- methyl (g ai/ha)	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
1.25	0	0	-	70	-	10	-	30	-	0	-	0	-	50	-	10	-
2.5	0	5	-	80	-	10	-	20	-	0	-	0	-	60	-	20	-
5	0	20	-	93	-	30	-	40	-	30	-	0	-	90	-	20	-
0	4.375	90	-	0	-	10	-	0	-	0	-	0	-	0	-	80	-
0	8.75	93	-	10	-	50	-	0	-	0	-	10	-	10	-	85	-
0	17.5	95	-	20	-	65	-	0	-	0	-	20	-	20	-	85	-
1.25	4.375	97	90	75	70	40	19	50	30	0	0	50	0	50	50	95	82
1.25	8.75	95	93	80	73	65	55	50	30	10	0	60	10	50	55	93	87
1.25	17.5	98	95	85	76	70	69	50	30	10	0	85	20	60	60	95	87
2.5	4.375	93	91	80	80	70	19	40	20	0	0	60	0	80	60	97	84
2.5	8.75	95	93	85	82	75	55	40	20	10	0	50	10	90	64	97	88
2.5	17.5	97	95	87	84	85	69	45	20	20	0	65	20	95	68	97	88
5	4.375	87	92	95	93	60	37	60	40	20	30	65	0	85	90	90	84
5	8.75	95	94	90	94	70	65	75	40	20	30	70	10	80	91	95	88
5	17.5	97	96	90	94	75	76	75	40	30	30	75	20	85	92	95	88

Table 2: Synergistic combination of Cmpd 1 and diclosulam.

Application Rate		BRSNW		CHEAL		VIOTR		STEME		SETFA		SORVU		AMARE		CYPES		IPOHE	
Cmpd 1 (g ae/ha)	Diclosulam (g ai/ha)	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
1.25	0	0	-	70	-	10	-	30	-	0	-	0	-	50	-	10	-	10	-
2.5	0	5	-	80	-	10	-	20	-	0	-	0	-	60	-	20	-	20	-
5	0	20	-	93	-	30	-	40	-	30	-	0	-	90	-	20	-	20	-
0	4.375	80	-	0	-	10	-	0	-	0	-	0	-	10	-	60	-	75	-
0	8.75	85	-	10	-	50	-	0	-	0	-	0	-	20	-	60	-	80	-
0	17.5	90	-	40	-	70	-	0	-	0	-	0	-	30	-	70	-	85	-
1.25	4.375	95	80	85	70	20	19	20	30	0	0	10	0	85	55	97	80	97	78
1.25	8.75	97	85	87	73	60	55	40	30	20	0	20	0	80	60	95	80	100	82
1.25	17.5	97	90	90	82	85	73	50	30	20	0	30	0	100	65	97	85	100	87
2.5	4.375	93	81	80	80	70	19	40	20	20	0	30	0	70	64	97	96	97	80
2.5	8.75	93	86	83	82	70	55	50	20	20	0	60	0	70	68	90	96	95	84
2.5	17.5	95	91	87	88	85	73	50	20	30	0	65	0	85	72	95	97	97	88
5	4.375	93	84	95	93	50	37	70	40	10	30	10	0	80	91	90	97	97	80
5	8.75	95	88	90	94	80	65	70	40	30	30	10	0	93	92	95	97	93	84
5	17.5	97	92	90	96	90	79	75	40	30	30	20	0	95	93	93	98	95	88

CLAIMS:

1. A herbicidal composition comprising a herbicidally effective amount of (a) a compound of the formula (I)




(I)

- or an agriculturally acceptable salt or ester thereof and (b) diclosulam or an agriculturally acceptable salt thereof, wherein the combination of (a) and (b) exhibits synergism.
2. The composition of claim 1, wherein no additional herbicidal active ingredients are present in the composition.
 3. The composition of claim 1 or 2, wherein (a) is a C1-C4 alkyl or benzyl ester of compound (I).
 4. The composition of claim 1 or 2, wherein (a) is a methyl ester or potassium salt of compound (I).
 5. The composition of claim 1 or 2, wherein (a) is the carboxylic acid of formula (I).
 6. The composition of claim 1 or 2, wherein (a) is the methyl ester of the compound of formula (I) and (b) is diclosulam.
 7. The composition of any one of claims 1-5, wherein (b) is diclosulam or an agriculturally acceptable salt thereof and the weight ratio of (a) to (b) is from about 1:35 to about 5:1.

8. The composition of any one of claims 1-5, wherein (b) is diclosulam or an agriculturally acceptable salt thereof and the weight ratio of the compound of (a) to (b) is from about 1:20 to about 1.25:1.
9. The composition of any one of claims 1-5, wherein (b) is diclosulam or an agriculturally acceptable salt thereof and the weight ratio of (a) to (b) is from about 5:4.375 to about 1.25:17.5.
10. The composition of any one of claims 1-9, further comprising a herbicide safener.
11. The composition of any one of claims 1-10, further comprising an agriculturally acceptable adjuvant or carrier.
12. A method of controlling undesirable vegetation which comprises applying a herbicidally effective amount of a herbicidal composition comprising:

(a) a compound of the formula (I)



(I)

or an agriculturally acceptable salt or ester thereof and

(b) diclosulam or an agriculturally acceptable salt thereof, wherein the combination of (a) and (b) exhibits synergism.


13. The method of claim 12, wherein no additional herbicidal active ingredients are present in the composition.
14. The method of claim 12 or 13 wherein the undesirable vegetation is controlled in winter/spring oilseed rape, winter/spring canola, vegetables, *Brassica* spp,

ornamentals, rice, wheat, triticale, barley, oats, rye, sorghum, corn/maize, sunflower, row crops, pastures, grasslands, rangelands, fallowland, sugarcane, turf, tree orchards, vine orchards, aquatics, industrial vegetation management or rights-of-way.

15. The method of any one of claims 12-14, wherein the undesirable vegetation is immature.
16. The method of any one of claims 12-14, wherein the (a) and (b) are applied pre-emergently.
17. The method of any one of claims 12-14, wherein the (a) and (b) are applied post-emergently.
18. The method of claim 12 or 13 wherein the undesirable vegetation is controlled in a crop that is tolerant to glyphosate, glufosinate, dicamba, phenoxy auxins, pyridyloxy auxins, aryloxyphenoxypropionates, acetyl CoA carboxylase (ACCase) inhibitors, imidazolinones, acetolactate synthase (ALS) inhibitors, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, triazines, or bromoxynil.
19. The method of claim 18, wherein the tolerant crop possesses multiple or stacked traits conferring tolerance to multiple herbicides or multiple modes-of-action.
20. The method of claim 18 or 19, wherein the undesirable vegetation comprises a herbicide resistant or tolerant weed.
21. The method of claim 20, wherein the resistant or tolerant weed is a biotype with resistance or tolerance to multiple herbicides, multiple chemical classes, or multiple herbicide modes-of-action.
22. The method of claim 20, wherein the resistant or tolerant weed is a biotype resistant or tolerant to acetolactate synthase (ALS) inhibitors, photosystem II inhibitors, acetyl CoA carboxylase (ACCase) inhibitors, synthetic auxins, photosystem I inhibitors, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, microtubule assembly inhibitors, lipid synthesis inhibitors, protoporphyrinogen

oxidase (PPO) inhibitors, carotenoid biosynthesis inhibitors, very long chain fatty acid (VLCFA) inhibitors, phytoene desaturase (PDS) inhibitors, glutamine synthetase inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors, mitosis inhibitors, cellulose biosynthesis inhibitors, herbicides with multiple modes-of-action, quinclorac, arylaminopropionic acids, difenzoquat, endothall, or organoarsenicals.

23. The method of any one of claims 12-22, wherein the undesirable vegetation is *Viola tricolor* (L.) (VIOTR), *Stellaria media* (L.) Vill. (STEME), *Sorghum vulgare* (SORVU), or *Ipomoea hederacea* (L.) Jacq. (IPOHE).
24. The method of any one of claims 12-23, wherein (a) is the methyl ester of the compound of formula (I), (b) is diclosulam, and the (a) and (b) are applied in a weight ratio of (a) to (b) of from about 1:35 to about 5:1.
25. The method of any one of claims 12-23, wherein (a) is the methyl ester of the compound of formula (I), (b) is diclosulam, and the (a) and (b) are applied in a weight ratio of (a) to (b) of from about 1:20 to about 1.25:1.
26. The method of any one of claims 12-23, wherein (a) is the methyl ester of the compound of formula (I), (b) is diclosulam, and the (a) and (b) are applied in a weight ratio of (a) to (b) of from about 5:4.375 to about 1.25:17.5.
27. The method of any one of claims 12-23, wherein the method comprises applying a composition as defined in any one of claims 3-11.



(I)