



(51) International Patent Classification:

C07D 207/22 (2006.01) **A61K 31/4025** (2006.01)
C07D 403/12 (2006.01) **A61P 9/00** (2006.01)
C07D 409/12 (2006.01) **A61P 29/00** (2006.01)
C07D 413/12 (2006.01) **A61P 27/06** (2006.01)
C07D 417/12 (2006.01) **A61P 17/00** (2006.01)
C07D 471/04 (2006.01)

(21) International Application Number:

PCT/US2010/051622

(22) International Filing Date:

6 October 2010 (06.10.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/249,101 6 October 2009 (06.10.2009) US

(71) Applicant (for all designated States except US): **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, California 92612 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHOW, Ken** [US/US]; 20 Tidal Surf, Newport Coast, California 92657 (US). **GIL, Daniel W.** [US/US]; 2541 Point Del Mar, Corona Del Mar, California 92625 (US). **DONELLO, John E.** [US/US]; 34041 Pequito Drive, Dana Point, California 92629 (US). **WANG, Liming** [US/US]; 16 Trinity, Irvine, California 92612 (US). **CORPUZ, Evelyn G.** [PH/US]; 1 Pollena, Irvine, California 92602 (US). **FANG, Wenkui K.** [US/US]; 73 Peppermint Tree, Irvine, California 92618 (US). **SINHA, Santosh C.** [IN/US]; 199 Sklar Street, Ladera Ranch, California 92694 (US). **DIBAS, Mohammed I.** [US/US]; 28871 Niguel Vis, Laguna Niguel, California 92677 (US).

(74) Agents: **ENE, Doina G.** et al.; Allergan, Inc., 2525 Dupont Drive, Irvine, California 92612 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: 2H-PYRROL-5-AMINE DERIVATIVES AS ALPHA ADRENERGIC RECEPTOR MODULATORS

(57) Abstract: Compounds are described herein useful for treating diseases and conditions by modulation of one or more alpha adrenergic receptor. The compounds can include a naphthalene, a quinoline, a benzoimidazole or an isoquinoline as a core structure. Methods of making, using and formulating these compounds are described.



2H-PYRROL-5-AMINE DERIVATIVES AS ALPHA ADRENERGIC RECEPTOR MODULATORS

Inventors: Ken Chow, Liming Wang, Wenkui K. Fang, Evelyn G. Corpuz, Santosh C. Sinha, Daniel W. Gil, Mohammed I. Dibas and John Donello

5 **CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims the benefit of U.S. Provisional Application Serial No. 61/249,101 filed on October 06, 2009, which is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

10 Human adrenergic receptors are integral membrane proteins which have been classified into two broad classes, alpha and the beta adrenergic receptors. Both types of receptors mediate the action of the peripheral sympathetic nervous system upon binding of catecholamines, norepinephrine and epinephrine.

15 Norepinephrine is produced by adrenergic nerve endings, while epinephrine is produced by the adrenal medulla. The binding affinity of adrenergic receptors for these compounds, forms one basis of classification: alpha receptors tend to bind norepinephrine more strongly than epinephrine. The preferred binding affinity of these hormones is reversed for the beta receptors. In many tissues, the functional responses, such as smooth muscle contraction, induced by alpha receptor activation are opposed to responses induced by beta receptor binding.

20 Subsequently, the functional distinction between alpha and beta receptors was further highlighted and refined by the pharmacological characterization of these receptors from various animal and tissue sources. As a result, alpha and beta adrenergic receptors were further subdivided into alpha-1 (α_1), alpha-2 (α_2), beta-1 (β_1), and beta-2 (β_2) subtypes.

25 Alpha adrenergic agents are known in the art. Whereas alpha-1 agonists are known to include compounds which have vasoconstrictor activity and are thus useful for controlling intraocular bleeding, alpha-2 agonist are known to include compounds useful for reducing intraocular pressure (anti-glaucoma effect), for increasing renal flow (diuretics) and for altering the rate of fluid transport in the gastrointestinal tract (anti-
30 diarrheals).

In "Heteroaromatic Analogues of the α_2 -Adrenoreceptor Partial Agonist Clonidine" J. Med. Chem. 1989, 32, 1627-1630, Chapleo et al. describe 6-(2-iminoimidazolidine)-3-oxo-3,4-dihydro-(2H)-1,4-benzoxazine and 7-(2-iminoimidazolidine)-3-oxo-3,4-dihydro-(2H)-1,4-benzoxazine compounds as partial α_2 agonists.

"Analogues of UK 14,304: Structural Features Responsible for α_2 Adrenoreceptor Activity", Bioorg. & Med. Chem. Letters, Vol 5, No.15, pp 1745-1750 describes iminoimidazoline derivatives with α_2 activity.

U.S. Patent No. 3,890,319 discloses 2-imidazolin-2-yl-amino-substituted quinoxalines as regulators of the cardiovascular system.

U.S. Patent No. 4,515,800, describes 2-(trisubstituted phenylimino)imidazoline compounds [also known as 2-(trisubstituted-anilino)-1,3-diazacyclopentene-(2) compounds] in pharmaceutical compositions, preferably in eye drops, for the treatment of glaucoma.

U.S. Patent No. 4,587,257 discloses 2-(trisubstituted phenylimino)imidazoline compounds capable of controlling ocular bleeding.

U.S. Pat. No. 3,636,219 discloses 2-(substituted-phenylamino)-thiazolines and imidazolines having anticholinergic activity.

U.S. Patent No. 5,091,528 discloses 6- or 7- (2-imino-2-imidazolidine)-1,4-benzoxazines as α adrenergic agents having α adrenergic activity and useful for the treatment of glaucoma, renal and gastrointestinal disorders and vasoconstrictors.

Compound (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine CAS 753464-99-2 is available from Aurora Screening Library.

Functional differences between α_1 and α_2 receptors have been recognized, and compounds which exhibit selective binding between these two subtypes have been developed. Thus, in WO 9200073 for example, the selective ability of the R(+) enantiomer of terazosin to selectively bind to adrenergic receptors of the α_1 subtype was reported. The α_1 / α_2 selectivity of this compound was disclosed as being significant because agonist stimulation of the α_2 receptors was said to inhibit secretion of epinephrine and norepinephrine, while antagonism of the

alpha-2 receptor was said to increase secretion of these hormones. Thus, the use of non-selective alpha-adrenergic blockers, such as phenoxybenzamine and phentolamine, was said to be limited by their alpha-2 adrenergic receptor mediated induction of increased plasma catecholamine concentration and the attendant physiological sequelae (increased heart rate and smooth muscle contraction).

The cloning, sequencing and expression of alpha receptor subtypes from animal tissues has led to the subclassification of the alpha-1 adrenoreceptors into alpha-1A, alpha-1B, and alpha-1D. Similarly, the alpha-2 adrenoreceptors have also been classified alpha-2A, alpha-2B, and alpha-2C receptors. Each alpha-2 receptor subtype appears to exhibit its own pharmacological and tissue specificities. Compounds having a degree of specificity for one or more of these subtypes may be more specific therapeutic agents for a given indication than an alpha-2 receptor pan-agonist (such as the drug clonidine) or a pan-antagonist.

Among other indications, such as the treatment of glaucoma, hypertension, sexual dysfunction, and depression, certain compounds having alpha adrenergic receptor agonist activity are known analgesics. However, many compounds having such activity do not provide the activity and specificity desirable when treating disorders modulated by alpha adrenoreceptors. For example, many compounds found to be effective agents in the treatment of pain are frequently found to have undesirable side effects, such as causing hypotension and sedation at systemically effective doses.

SUMMARY


Described herein are compounds that are useful as alpha adrenergic receptor modulators. These compounds are useful in treating a wide variety of disorders associated with modulation of alpha adrenergic receptors. These compounds are useful for the treatment of humans with diseases and conditions that are alleviated by alpha adrenergic modulation, and in particular, useful as alpha-1 and/or alpha-2 antagonists or agonists. Compositions are also described herein comprising a pharmaceutically acceptable amount of a compound as described herein. Further described are methods for treating conditions using the compositions described herein, wherein the conditions are selected from the group consisting of including but not limited to treating glaucoma, elevated intraocular pressure, ischemic neuropathies, optic neuropathy, pain, visceral pain, corneal pain, headache pain, migraine, cancer pain, back pain, irritable bowel syndrome pain, muscle pain and pain associated with diabetic

neuropathy, the treatment of diabetic retinopathy, other retinal degenerative conditions, stroke, cognitive deficits, neuropsychiatric conditions, drug dependence and addiction, withdrawal symptoms, obsessive-compulsive disorders, obesity, insulin resistance, stress-related conditions, diarrhea, diuresis, nasal congestion, spasticity, attention deficit disorder, psychoses, anxiety, depression, autoimmune disease, Crohn's disease, gastritis, Alzheimer's, Parkinson's ALS, and other neurodegenerative diseases, dermatological conditions, skin erythema (redness) and inflammation, rosacea, acne, psoriasis, inflammatory bowel disease (IBD), post-traumatic stress disorder (PTSD), Tourette's syndrome, multiple sclerosis, dry eye disease.

Certain terms as used in the specification are intended to refer to the following definitions, as detailed below. Where the definition of terms departs from the commonly used meaning of the term, applicant intends to utilize the definitions provided below, unless specifically indicated.

DETAILED DESCRIPTION

In one aspect, the invention therefore provides a compound having formula I, its enantiomers, diastereoisomers, tautomers or a pharmaceutically acceptable salt thereof



Formula I

wherein:

n is 0 or 1;

A is an aromatic or non-aromatic ring;

X is N, N-R¹, CH-R¹, C-R¹;

R¹ is H, C₁₋₆ alkyl or halogen;

R² is N, N-R³, CH-R³, C-R³, O or S;

R³ is H, halogen, or C₁₋₆ alkyl;

R⁴ is halogen, H, C₁₋₆ alkyl or part of a ring as a heteroatom (O, S, N) or as a C;

R⁵ is a heteroatom (S, N) or is a C and is always part of a ring;

R⁶ is H, C₁₋₆ alkyl, halogen or is part of a ring as a heteroatom (O, S, N) or as a


C;

R^4 and R^5 can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C_{1-6} alkyl or with 1 to 2 halogen and is directly attached to the rest of the molecule; or

R^5 and R^6 can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C_{1-6} alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule;

except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine.

In another aspect, the invention therefore provides a compound having formula II, its enantiomers, diastereoisomers, tautomers or a pharmaceutically acceptable salt thereof



Formula II

wherein:

n is 1;

A is an aromatic ring;

X is $C-R^1$;

R^1 is H;

R^2 is N or $C-R^3$;

R^3 is H, halogen, or C_{1-6} alkyl;


R^4 is part of a ring as a heteroatom (O, S, N) or as a C;

R^5 is a heteroatom (N) or is a C and is always part of a ring;

R^6 is H, C_{1-6} alkyl or halogen; and

R^4 and R^5 form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C_{1-6} alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

In another aspect, the invention therefore provides a compound having formula III, its enantiomers, diastereoisomers, tautomers or a pharmaceutically acceptable salt thereof



Formula III

wherein:

n is 1;

A is an aromatic ring;

X is C-R¹;

R¹ is H, C₁₋₃ alkyl;

R² is N or C-R³;

R³ is H, halogen, or C₁₋₆ alkyl;


R⁴ is halogen or H;

R⁵ is a heteroatom (N) or C and is always part of a ring;

R⁶ is part of a ring as a heteroatom (N) or as a C;

R⁵ and R⁶ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

In another aspect, the invention therefore provides a compound having formula IV, its enantiomers, diastereoisomers, tautomers or a pharmaceutically acceptable salt thereof



Formula IV

wherein:

n is 0;

A is a non-aromatic ring;

R² is CH-R³;

R³ is H;


R^4 is H;

R^5 is C and is always part of a ring;

R^6 is part of a ring as a C; and

R^5 and R^6 can form a 6 membered aromatic carbocyclic ring directly attached to the rest of the molecule.

In another aspect, the invention therefore provides a compound having formula V, its enantiomers, diastereoisomers, tautomers or a pharmaceutically acceptable salt thereof



Formula V

wherein

n is 1;

A is a non-aromatic ring;

X is CH- R^1 ;

R^1 is H;

R^2 is CH- R^3 ;

R^3 is H;

R^4 is H;

R^5 is C and is always part of a ring;

R^6 is part of a ring as a C; and

R^5 and R^6 can form a 6 membered aromatic carbocyclic ring, optionally substituted with C₁₋₆ alkyl or halogen and is directly attached to the rest of the molecule; except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine.

The person of skill in the art would appreciate that when " n " is 0 then X is absent and ring A is a 5 membered ring.

The person of skill in the art would appreciate the formation of bicyclic hetero- or carbocyclic rings, when R^4 and R^5 form a 5 or 6 aromatic or non-aromatic membered carbo- or heterocyclic ring, directly attached to the rest of the molecule.

The person of skill in the art would appreciate the formation of bicyclic hetero- or carbocyclic rings, when R⁵ and R⁶ form a 5 or 6 aromatic or non-aromatic membered carbo- or heterocyclic ring, directly attached to the rest of the molecule.

The person of skill in the art would appreciate that when "A" is an aromatic ring X is N or C-R¹ and when "A" is a non-aromatic ring X is N-R¹ or CH-R¹.

The person of skill in the art would appreciate that when "A" is an aromatic ring R² is N or C-R³ and when "A" is a non-aromatic ring R² is N-R³, O, S or CH-R³.

Further, pharmaceutically acceptable salts, hydrates, solvates, crystal forms and individual isomers, enantiomers, tautomers, diastereomers and prodrugs of the compounds described can be utilized.

The term "alkyl" as used herein, is defined as including a saturated monovalent hydrocarbon moiety having straight or branched moieties or combinations thereof and containing 1-6 carbon atoms, preferably 1-3 carbon atoms. Alkyl moieties can optionally be substituted by cycloalkyl groups or one methylene group (-CH₂-) can be replaced by carbonyl or carboxyl group. Usually, in the present case, alkyl group is methyl.

The term "cycloalkyl" as used herein, refers to a monovalent or divalent group of 3 to 8 carbon atoms, preferably 3-6 carbon atoms derived from saturated cyclic hydrocarbon. Cycloalkyl groups can be optionally substituted by alkyl groups or halogen groups. Usually, in the present case, cycloalkyl groups are cyclohexyl, cyclopentyl.

The term "halogen" as used herein includes an atom of fluoride, chloride, bromide or iodide. Usually, in the present case, halogens are fluoride, chloride and bromide.

The term "heterocyclic" ring as used herein, is defined as including an aromatic or non-aromatic moiety that includes at least one heteroatom in the ring, wherein one or more ring carbons are replaced with a heteroatom. Heterocyclic rings can have at least one O or N or S interrupting the cyclic ring structure or combinations thereof. In the present case heterocyclic rings can be aromatic or non aromatic and can be formed of 5 or 6 atoms. Heterocyclic ring moieties can be optionally substituted by alkyl or halogen groups, as defined above. Usually, in the

present case, heterocyclic rings are imidazole, thiazol, morpholine, pyrrol, pyridine, pyrazine, furan, thiophene.

The term "bicyclic heterocyclic" rings as used herein, refers to aromatic or non-aromatic fused cyclic rings wherein at least one of the cycles has a heteroatom in its ring, as defined above. Usually, in the present case, bicyclic heterocyclic rings are quinoxaline, quinoline, isoquinoline, benzimidazole, benzothiazole, 3,4-dihydro-2H-benzo[b][1,4]oxazine, indole, 1Hpyrrolo[2,3-b]pyridine, 6,7-dihydro-5H-cyclopenta[b]pyridine, 1,5-naphtyridine, benzothiophen, benzofuran.

The term "carbocyclic" ring as used herein refers to a ring formed only by carbon atoms. In the present case carbocyclic rings can be aromatic or non aromatic and can be formed of 5 or 6 carbon atoms. Carbocyclic rings can optionally be substituted by alkyl or halogen, as defined above. Usually, in the present case, carbocyclic rings are phenyl, cyclohexyl, cyclopentyl.

The term "bicyclic carbocyclic" rings as used herein, refers to aromatic or non-aromatic fused carbocyclic rings formed only by carbon atoms. Usually, in the present case, bicyclic carbocyclic rings are naphthalene, tetrahydronaphtalene, dihydroindene.

The term "N" as used herein, refers to a nitrogen atom.

The term "O" as used herein, refers to an oxygen atom.

The term "S" as used herein, refers to a sulfur atom.

The term "H" as used herein, refers to a hydrogen atom.

The term "C" as used herein, refers to a carbon atom.

As used herein, "pharmaceutically acceptable salt" refers to any salt that retains the activity of the parent compound and does not impart any additional deleterious or untoward effects on the subject to which it is administered, and in the context in which it is administered compared to the parent compound. A pharmaceutically acceptable salt also refers to any salt which may form *in vivo* as a result of administration of an acid, another salt, or a prodrug which is converted into an acid or salt. Further, pharmaceutically acceptable salt refers to those salts which retain the biological effectiveness and properties of the free bases and which are obtained by reaction with inorganic acids such as hydrochloric acid, hydrobromic

acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, fumaric acid, salicylic acid and the like.


Pharmaceutically acceptable salts of acidic functional groups may be derived from organic or inorganic bases. The salt may comprise a mono or polyvalent ion.

Of particular interest are the inorganic ions, lithium, sodium, potassium, calcium, and magnesium. Organic salts may be made with amines, particularly ammonium salts such as mono-, di- and trialkyl amines or ethanol amines. Salts may also be formed with caffeine, tromethamine and similar molecules. Hydrochloric acid or some other pharmaceutically acceptable acid may form a salt with a compound that includes a basic group, such as an amine or a pyridine ring.

As used herein "prodrug" refers to a compound which is converted to a therapeutically active compound after administration, and the term should be interpreted as broadly herein as is generally understood in the art. While not intending to limit the scope of the present description, conversion may occur by hydrolysis of an ester group or some other biologically labile group. Generally, but not necessarily, a prodrug is inactive or less active than the therapeutically active compound to which it is converted.

As used herein, "tautomer" refers to the migration of protons between adjacent single and double bonds. The tautomerization process is reversible. Compounds described herein can undergo any possible tautomerization that is within the physical characteristics of the compound.

The following is an example tautomerization that can occur in compounds described herein:



As used herein, the term "therapeutically effective amount" means the amount of the pharmaceutical composition that will elicit the biological or medical response of a subject in need thereof that is being sought by the researcher, veterinarian, medical doctor or other clinician. In some embodiments, the subject in need thereof is a mammal. In some embodiments, the mammal is human.

Compounds of the invention are:

- N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine;
N-(2-Methyl-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(4-Bromo-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
5 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine;
4-Chloro-N-[(2E)-pyrrolidin-2-ylidene]-1,5-naphthyridin-3-amine;
N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine fumarate;
N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
10 5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine fumarate;
5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;
8-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-7-amine;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
4-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
15 N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1H-benzimidazol-5-amine;
7-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;
N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine ;
20 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;
7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine;
25 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-7-amine;
N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
N-1-Benzofuran-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine;
N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine;
30 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine ;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;

N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-8-amine;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine fumarate;
5 N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine;
N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
N-(7-Fluoro-2,3-dihydro-1H-inden-1-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-amine;
10 N-[(1S)-4-Methyl-2,3-dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
4-Methyl-N-[(2E)-pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine.

Preferred compounds of the invention are:

N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine;
15 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
20 N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;
7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine;
N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
25 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine;
N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine;
30 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;

N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine.

Most preferred compounds of the invention are:

4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;

5 N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;

7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;

10 N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;

4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;

4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine.

15 The compounds described herein are useful as medicaments, or in compositions or formulations in mammals, including humans, for treatment of diseases and or alleviations of conditions which are responsive to treatment by modulation of alpha adrenergic receptors. Thus, in further embodiment examples, there are provided methods for treating a disorder associated with modulation of alpha adrenergic
20 receptors. Such methods can be performed, for example, by administering to a subject in need thereof a pharmaceutical composition containing a therapeutically effective amount of at least one compound described herein.

The compounds described herein may be administered at pharmaceutically effective dosages. Such dosages are normally the minimum dose necessary to achieve
25 the desired therapeutic effect. Generally, such doses will be in the range of about 1 mg/day to about 1000 mg/day; more preferably in the range of about 10 mg/day to about 500 mg/day. In another embodiment example, the compound or compounds may be present in a composition or formulation in a range of about 0.5 mg/kg/day to about 100 mg/kg/day or about 1 mg/kg/day to about 100 mg/kg/day. However, the actual
30 amount of the compound to be administered in any given case will be determined by a physician taking into account the relevant circumstances, such as the severity of the pain, the age and weight of the patient, the patient's general physical condition, the cause of pain, and the route of administration.

In another aspect of the invention, provided are pharmaceutical compositions including at least one compound in a pharmaceutically acceptable carrier.

Pharmaceutical compositions can be used in the form of a solid, a solution, an emulsion, a dispersion, a micelle, a liposome, and the like, wherein the resulting composition contains one or more compounds described herein, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for enteral or parenteral applications. One or more compounds may be combined, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The carriers which can be used include glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. Compounds described herein are included in pharmaceutical compositions in an amount sufficient to produce the desired effect upon the process or disease condition.

In another aspect of the invention, the compounds described herein can be administered orally in any acceptable form, such as a tablet, liquid, capsule, powder and the like. However, other routes may be desirable or necessary, particularly if the patient suffers from nausea. Such other routes may include, without exception, transdermal, parenteral, subcutaneous, intranasal, intrathecal, intramuscular, intravenous, and intrarectal modes of delivery. Additionally, formulations may be designed to delay release of the active compound over a given period of time, or to carefully control the amount of drug released at a given time during the course of therapy.

Pharmaceutical compositions in a form suitable for oral use, for example, are administered as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of a sweetening agent such as sucrose, lactose, or saccharin, flavoring agents such as peppermint, oil of

wintergreen or cherry, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets containing compounds described herein in admixture with non-toxic pharmaceutically acceptable excipients may also be manufactured by known methods.

5 The excipients used may be, for example, (1) inert diluents such as calcium carbonate, lactose, calcium phosphate or sodium phosphate; (2) granulating and disintegrating agents such as corn starch, potato starch or alginic acid; (3) binding agents such as gum tragacanth, corn starch, gelatin or acacia, and (4) lubricating agents such as magnesium stearate, stearic acid or talc. The tablets may be uncoated
10 or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

15 In some cases, formulations for oral use may be in the form of hard gelatin capsules wherein the compounds are mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the form of soft gelatin capsules wherein the compounds are mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil.

20 The pharmaceutical compositions may also be in the form of a sterile injectable suspension. Suspensions may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For
25 this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides, fatty acids (including oleic acid), naturally occurring vegetable oils like sesame oil, coconut oil, peanut oil, cottonseed oil, etc., or synthetic fatty vehicles like ethyl oleate or the like. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

30 Compounds described herein may also be administered in the form of suppositories for rectal administration. These compositions may be prepared by mixing the compounds with a suitable non-irritating excipient, such as cocoa butter, synthetic

glyceride esters of polyethylene glycols, which are solid at ordinary temperatures, but liquify and/or dissolve in the rectal cavity to release the drug.

The compounds described herein can also be administered as an ophthalmically acceptable formulation or composition. A liquid which is ophthalmically acceptable is formulated such that it can be administered topically to the eye. The comfort should be maximized as much as possible, although sometimes formulation considerations (e.g. stability) may necessitate less than optimal comfort. In the case that comfort cannot be maximized, the liquid should be formulated such that the liquid is tolerable to the patient for topical ophthalmic use. Additionally, an ophthalmically acceptable liquid should either be packaged for single use, or contain a preservative to prevent contamination over multiple uses.

For ophthalmic application, solutions or medicaments are often prepared using a physiological saline solution as a major vehicle. Ophthalmic solutions should preferably be maintained at a comfortable pH with an appropriate buffer system. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and surfactants.

Preservatives that may be used in ophthalmic compositions described herein include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A useful surfactant is, for example, Tween 80. Likewise, various useful vehicles may be used in the ophthalmic preparations described herein. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In another aspect of the invention, an ophthalmic composition as described herein may have ingredients used in the following amounts listed in Table 1.

Table 1

Ingredient	Amount (% w/v)
active ingredient	about 0.001-5
preservative	0-0.10
vehicle	0-40
tonicity adjustor	1-10
buffer	0.01-10
pH adjustor	q.s. pH 4.5-7.5
antioxidant	as needed
surfactant	as needed
purified water	as needed to make 100%

In other embodiments, the ophthalmically acceptable liquid can be formulated for intraocular injection. The compounds described herein can be formulated as a liquid, gel paste, or the like for intraocular injection. Further, the compounds can be formulated into sustained release or controlled release intraocular implants comprising biodegradable polymers such as polylactic acid, poly glycolic acid, combinations thereof and the like.

Since individual subjects may present a wide variation in severity of symptoms and each composition has its unique therapeutic characteristics, the precise mode of administration and dosage employed for each subject is left to the discretion of the practitioner.

EXPERIMENTAL DETAILS


The present invention concerns also a process for preparing the compounds having formulae I, II, III, IV or V.

The synthetic schemes set forth below, illustrate how compounds according to the invention can be made. Those skilled in the art will be able to routinely modify and/or adapt the following schemes to synthesize any compounds of the invention covered by formulae I, II, III, IV or V.

Pyrrol-amines can be obtained according to the following general schemes. Most of the primary amine are commercially available. In Scheme 1 the primary amine reacts


with pyrrolidin-2-one in the presence of phosphoryl chloride to form the desired pyrrol-amine. The solvent for this reaction is toluene.

Scheme 1




- 5 In Scheme 2 the primary amine reacts with 5-(methylthio)-3,4-dihydro-2H-pyrrole in isopropanol to form the desired pyrrol-amine.

Scheme 2



- 10 In Scheme 3 the primary amine reacts with 5-methoxy-3,4-dihydro-2H-pyrrole in methanol to form the desired pyrrol-amine.

Scheme 3



The following examples are for illustrative purposes only and are not intended, nor should they be construed as limiting the invention in any manner. Those skilled in the art will appreciate that variations and modifications of the following examples can be made without exceeding the spirit or scope of the invention.

The IUPAC names of the compounds mentioned in the examples were generated with ACD version 8.

Unless specified otherwise in the examples, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on 300 MHz Varian and acquired at room temperature. Chemical shifts are given in p.p.m. referenced either to internal TMS or to the residual solvent signal.

All the reagents, solvents, catalysts for which the synthesis is not described are purchased from chemical vendors such as Sigma Aldrich, Fluka, Lancaster, however some known reaction intermediates, for which the CAS registry number (CAS) are mentioned, were prepared in-house following known procedures.

Usually the compounds of the invention were purified by flash column chromatography using as solvent system: 7N NH₃ in MeOH/DCM.

The following abbreviations are used in the examples:

POCl₃ phosphoryl chloride

DCM dichloromethane

NaOH sodium hydroxyde

MeOH methanol

CD₃OD deuterated methanol

NH₃ ammonia

EtOAc ethylacetate

Pd/C palladium on carbon

H₂SO₄ sulfuric acid

Na₂SO₄ sodium sulfate

	Bu ₄ NNO ₂	tetrabutylammonium nitrate
	Ac ₂ O	trifluoroacetic anhydride
	DMF	dimethylformamide
	MgSO ₄	magnesium sulfate
5	EtOAc	ethylacetate
	i-PrOH	isopropanol
	NaI	sodium iodide
	CDCl ₃	deuterated chloroform
	AcOH	acetic acid
10	MPLC	medium pressure liquid chromatography

Example 1

N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine

(Compound 1)

A solution of POCl₃ (765mg) in 8 mL of toluene was added slowly to a solution of 2-pyrrolidinone (85g) in 8 mL of toluene at 10°C. Then, the reaction was stirred at room temperature for 3 hours. A solution of 1-naphthylamine (858mg) (CAS 134-32-7) in 8 mL of toluene was added, and the mixture was heated to reflux overnight. The refluxed mixture was cooled to room temperature, and the toluene layer was decanted. The residue was dissolved in aqueous water and DCM. Aqueous 5N NaOH was used to adjust the pH to alkaline. Two layers were separated, and the organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced vacuum. Flash column chromatography (2-4% 7N NH₃ in MeOH/DCM) yielded the title compound. ¹H NMR (300 MHz, CD₃OD) 2.09 (m, 2H), 2.61 (m, 2H), 3.86 (m, 2H), 7.27 (m, 1H), 7.45(m, 1H), 7.57 (m, 2H), 7.87 (m, 2H), 7.99 (m, 1H).

Compounds 2, 3, 4, 5 and 6 were prepared in a similar manner to the method described in Example 1 for Compound 1. The reagents used and the results obtained are tabulated below in Table 2.

Table 2

Compound Number	IUPAC Name	Starting material	¹ H NMR (Solvent) δ ppm
2	N-(2-Methyl-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine	1-amino-2-methylnaphtalene (CAS 2246-44-8)	(CD ₃ OD) 2.12 (m, 2H), 2.29 (s, 3H), 2.81 (m, 2H), 3.33 (m, 2H), 7.37 (m, 3H), 7.55 (d, J=8.21Hz, 1H), 7.77 (m, 1H), 7.85 (m, 1H).
3	N-(4-Bromo-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine	1-amino-4-bromonaphtalene (CAS 2298-07-9)	(CD ₃ OD) 2.08 (m, 2H), 2.62 (m, 2H), 3.36 (t, J=6.74Hz, 2H), 6.90 (d J=7.91Hz, 1H), 7.50 (dd, J=1.17 & 7.30Hz, 1H), 7.55 (dd, J=1.17 & 8.20Hz, 1H), 7.70 (d, J=7.91Hz, 1H), 8.00 (dd, J=0.80 & 8.50Hz, 1H), 8.15 (dd, J=0.80 & 8.20Hz, 1H).
4	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	6-amino-1H-benzimidazole (CAS 934-22-5)	(CD ₃ OD) 2.05 (m, 2H), 2.57 (t, J=7.92Hz, 2H), 3.44 (t, J=6.77Hz, 2H), 6.90 (dd, J=8.50Hz & 1.76Hz, 1 H), 7.25 (d, J=1.76Hz, 1H), 7.53 (d, J=8.50Hz, 1H), 7.90 (s, 1H).

5	N-(3,4-Dihydro-2H-pyrrol-5-yl)-6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine	6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine (CAS 178209-29-5)	(CD ₃ OD) 7.91 (br. s., 1H), 7.34 (br. s., 1H), 3.46 (t, J = 6.7 Hz, 2H), 2.92 (q, J = 7.4 Hz, 4H), 2.56 (t, J = 7.9 Hz, 2H), 1.97 - 2.24 (m, 2H).
6	4-Chloro-N-[(2E)-pyrrolidin-2-ylidene]-1,5-naphthyridin-3-amine	4-chloro-1,5-naphthyridin-3-amine (CAS 930276-73-6)	(CD ₃ OD) 8.88 - 8.99 (m, 1H), 8.64 (s, 1H), 8.41 (dd, J = 8.5, 1.2 Hz, 1H), 7.72 (dd, J = 8.5, 4.4 Hz, 1H), 3.46 (t, J = 6.9 Hz, 2H), 2.68 (br. s., 2H), 2.15 (quin, 2H).

Example 2**N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine**(Compound 7)

5

A solution of POCl₃ (382mg) in 8 mL of toluene was added slowly to a solution of 2-pyrrolidinone (430 mg) in 8 mL of toluene at 10°C. Then, the reaction was stirred at room temperature for 3 hours. A solution of 4-chloro-1-naphtalenamine (CAS 4684-12-2) (533mg) in 8 mL of toluene was added, and the mixture was heated to reflux overnight. The refluxed mixture was then cooled to room temperature, and the toluene layer was decanted. The residue was dissolved in aqueous water and DCM. Aqueous 5N NaOH was used to adjust the pH to alkaline. The resulting two layers were separated, and the organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced vacuum. Flash column chromatography (24% 7N NH₃ in MeOH/DCM) yielded the title compound. ¹H NMR (300 MHz, CD₃OD) δ 2.11 (m, 2H), 2.66 (m, 2H), 3.36 (t, J=6.74Hz, 2H), 6.96 (d J=7.91Hz, 1H), 7.49 (m, 2H), 7.50 (dd, J=1.13 & 7.11Hz, 1H), 8.00 (d, J=8.20Hz, 1H), 8.20 (d, J=8.50Hz, 1H). The fumarate salt of the tile compound was further obtained by crystallization with fumaric acid in

15

ethanol. ^1H NMR (300 MHz, CD_3OD) δ 8.34 (dd, $J = 7.2, 1.9$ Hz, 1H), 8.01 - 8.08 (m, 1H), 7.65 - 7.79 (m, 3H), 7.41 (d, $J = 7.9$ Hz, 1H), 6.66 (s, 1H), 3.51 - 3.66 (m, 3H), 3.07 (t, $J = 6.7$ Hz, 2H), 2.28 (dq, 2H). Ratio: Free base: Fumaric acid = 1:0.5.

Compound 8 and its fumarate salt were prepared in a similar manner to the method described in Example 2 for Compound 7. The reagent used and the results obtained are tabulated below in Table 3.

Table 3

Compound Number	IUPAC Name	Starting material	^1H NMR (Solvent) δ ppm
8	5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine	6-amino-5-bromoquinoxaline (CAS 50358-63-9)	Free base: (CD_3OD) 8.16 (d, $J = 8.5$ Hz, 1H), 8.01 (d, $J = 8.5$ Hz, 1H), 7.70 (d, $J = 7.9$ Hz, 1H), 7.54 - 7.63 (m, 1H), 7.44 - 7.54 (m, 1H), 6.90 (d, $J = 7.9$ Hz, 1H), 3.36 (t, $J = 6.7$ Hz, 2H), 2.63 (t, $J = 7.5$ Hz, 2H), 2.08 (qd, 2H). Fumarate salt: (CD_3OD) 8.29 (d, $J = 7.91$ Hz, 1H), 8.01 (d, $J = 7.91$ Hz, 1H), 7.88 (d, $J = 8.20$ Hz, 1H), 7.67 - 7.33 (m, 2H), 7.33 (d, $J = 8.20$ Hz, 1H), 6.66 (s, 1H), 3.59 (t, $J = 6.7$ Hz, 2H), 3.05 (t, $J = 7.5$ Hz, 2H), 2.29 - 2.24 (m, 2H). Free base : Fumaric acid = 1:1

Example 3**8-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-7-amine**(Compound 9)

To a mixture of 2-chlorobenzene-1,3-diamine (24 mmol) (CAS 6400-14-2) and
5 NaI (150 mg) in 80% H₂SO₄ (20 g) was added Glycerol (45 g). The mixture was stirred
at 150°C to 170°C for 5 h. The mixture was cooled to room temperature. Aqueous 5N
NaOH was used to adjust the pH to alkaline, which was then extracted with EtOAc
washed with brine, dried over MgSO₄ and concentrated under reduced vacuum. Flash
column chromatography (EtOAc:Hexane) yielded 7-amino-8-chloroquinoline (800 mg),
10 as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ: 8.69 (dd, J = 4.4, 1.8 Hz, 1H),
8.13 (dd, J = 7.9, 1.8 Hz, 1H), 7.61 (d, J = 9.1 Hz, 1H), 7.14 - 7.29 (m, 2H)

A solution of POCl₃ (551mg) in 8 mL of toluene was added slowly to a solution of
2-pyrrolidinone (612mg) in 8 mL of toluene at 10°C. Then the reaction was stirred at
room temperature for 3 hours. A solution of 7-amino-8-chloroquinoline (760mg) in 8 mL
15 of toluene was added, and the mixture was heated to reflux for two hours. The refluxed
mixture was then cooled to room temp, and the toluene layer was decanted. The
residue was dissolved in water and DCM. Aqueous 5N NaOH was used to adjust the
pH to alkaline. The two layers were separated, and the organic layer was washed with
water and brine, dried over Na₂SO₄ and concentrated under reduced vacuum. Flash
20 column chromatography (2-4% 7N NH₃ in MeOH/DCM) yielded the title compound. ¹H
NMR (300 MHz, CD₃OD) δ: 2.11 (m, 2H), 2.61 (m, 2H), 3.43 (t, J=6.74Hz, 2H), 7.31 (d,
J=8.50Hz, 1H), 7.60 (dd, J=4.60Hz & 8.20Hz, 1H), 7.81 (d, J=8.70Hz, 1H), 8.31 (dd,
J=1.76Hz & 8.20Hz, 1H), 8.84 (dd, J=1.47Hz & 4.40Hz, 2H).

25 Example 4

4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine(Compound 10)

5-(methylthio)-3,4-dihydro-2H-pyrrole (238mg) was combined with of 1H-benzimidazol-
30 6-amine, 7-bromo (430 mg) (CAS 177843-26-4) in 10 mL of isopropyl alcohol. The
reaction was stirred at reflux for 4 days. The refluxed mixture was then cooled to room
temperature and concentrated. Flash column chromatography (4-6% 7N NH₃ in
MeOH/DCM) yielded the title compound ¹H NMR (300 MHz, CD₃OD) δ 2.08 (m, 2H),

2.55 (t, J=7.92Hz, 2H), 3.39 (t, J=6.74Hz, 2H), 6.93 (d, J=8.50Hz, 1 H), 7.50 (d, J=8.50Hz, 1H), 8.11 (s, 1H).

Compounds 11 to 26 were prepared in a similar manner to the method described in Example 4 for Compound 10. The reagents used and the results obtained are tabulated below in Table 4.

Table 4

Compound Number	IUPAC Name	Starting material	¹ H NMR (Solvent) δ ppm
11	4-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	7-chloro-1H-benzimidazole-6-amine (CAS 177843-29-7)	(CD ₃ OD) 2.08 (m, 2H), 2.56 (t, J=7.92Hz, 2H), 3.40 (t, J=6.74Hz, 2H), 6.93 (d, J=8.50Hz, 1H), 7.46 (d, J=8.50Hz, 1H), 8.12 (s, 1H).
12	N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1H-benzimidazol-5-amine	7-methyl-1H-benzimidazole-6-amine (CAS 177843-30-0)	(CD ₃ OD) 2.09 (m, 2H), 2.62 (t, J=7.92Hz, 2H), 3.40 (t, J=6.74Hz, 2H), 6.87 (d, J=8.50Hz, 1H), 7.39 (d, J=8.50Hz, 1H), 8.07 (s, 1H).
13	7-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	1H-Benzimidazol-5-amine, 7-chloro- (CAS 10597-54-3)	(CD ₃ OD) 2.06 (m, 2H), 2.61 (t, J=7.92Hz, 2H), 3.48 (t, J=6.74Hz, 2H), 7.01 (d, J=1.47Hz, 1H), 7.23 (d, J=1.47Hz, 1H), 8.12 (s, 1H).
14	N-(3,4-Dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine	6-aminoquinoxaline (CAS 6298-37-9)	(CDCl ₃) 2.28 (m, 2H), 3.10 (t, J=7.92Hz, 2H), 3.83 (t, J=7.33Hz, 2H), 7.76 (dd, J=2.45Hz & 8.64Hz, 1 H), 7.88 (d,

			J=2.44Hz, 1H), 8.15 (d, J=8.64Hz, 1H), 8.87 (dd, J=1.72Hz & 8.55Hz, 2H).
15	N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine	8-aminoquanonoline (CAS 2217-41-6)	(CDCl ₃) 7.48 (d, J = 7.9 Hz, 1H), 7.12 (t, J = 7.8 Hz, 1H), 6.96 (d, J = 7.3 Hz, 1H), 3.40 (t, J = 6.2 Hz, 2H), 2.78 (t, J = 6.0 Hz, 2H), 2.57 (t, J = 6.0 Hz, 4H), 2.04 (s, 4H), 2.01 (d, J = 5.9 Hz, 3H), 1.69 - 1.91 (m, 5H).
16	N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine	1H-Inden-4-amine, 2,3-dihydro- (CAS 32202-61-2)	(CDCl ₃) 7.03 - 7.14 (m, 1H), 6.89 (d, J = 7.6 Hz, 2H), 3.47 (m, 2H), 2.92 (t, J = 7.5 Hz, 2H), 2.76 (t, J = 7.5 Hz, 2H), 2.53 (t, J = 7.6 Hz, 2H), 1.93 - 2.14 (m, 4H).
17	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine	1H-Indol-5-amine (CAS 5192-03-0)	(CDCl ₃) 7.24 - 7.34 (m, 2H), 7.16 (d, J = 3.2 Hz, 1H), 6.88 (dd, J = 8.5, 2.1 Hz, 1H), 6.46 (d, J = 3.2 Hz, 1H), 3.39 - 3.51 (m, 2H), 2.57 (t, J = 7.9 Hz, 2H), 2.05 (qd, 2H)
18	N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine	1H-Indol-5-amine, 4-methyl- (CAS 196205-06-8)	(CDCl ₃) 7.11 - 7.18 (m, 2H), 6.80 (d, J = 8.5 Hz, 1H), 6.50 (d, J = 3.2 Hz, 1H), 3.37 (t, J = 6.7 Hz, 2H), 2.53 (t, J = 7.9 Hz, 2H), 2.36 (s, 3H), 2.06 (s, 3H).
19	N-(3,4-Dihydro-2H-	1H-Indol-5-amine, 7-methyl	(CD ₃ OD) 7.17 (d, J = 3.2 Hz, 1H), 7.11 (d, J = 1.5

	pyrrol-5-yl)-7-methyl-1H-indol-5-amine	(CAS 90868-09-0)	Hz, 1H), 6.67 (s, 1H), 6.35 (d, J = 3.2 Hz, 1H), 3.45 (t, J = 6.9 Hz, 2H), 2.57 (t, J = 7.9 Hz, 2H), 2.45 (s, 3H), 2.01 (quin, 2H).
20	7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine	1H-Indol-5-amine, 7-bromo- (CAS 196205-07-9)	(CDCl ₃) 7.25 (d, J = 3.2 Hz, 1H), 7.20 (d, J = 1.8 Hz, 1H), 7.04 (d, J = 1.8 Hz, 1H), 6.45 (d, J = 3.2 Hz, 1H), 3.44 (t, J = 6.9 Hz, 2H), 2.57 (t, J = 8.1 Hz, 2H), 2.03 (quin, 2H).
21	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine	1H-Indol-6-amine (CAS 5318-27-4)	(CDCl ₃) 7.52 (d, J = 8.5 Hz, 1H), 7.18 - 7.31 (m, 1H), 7.11 (d, J = 3.2 Hz, 1H), 6.75 (dd, J = 8.4, 1.9 Hz, 1H), 6.48 (d, J = 4.1 Hz, 1H), 3.43 - 3.56 (m, 2H), 2.58 (t, J = 7.9 Hz, 2H), 2.10 - 2.03 (m, 2H).
22	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-7-amine	1H-Indol-7-amine (CAS 5192-04-1)	(CD ₃ OD) 7.22 - 7.30 (m, 1H), 7.17 (d, J = 3.2 Hz, 1H), 6.93 (t, J = 7.6 Hz, 1H), 6.69 (d, J = 7.3 Hz, 1H), 6.41 (d, J = 3.2 Hz, 1H), 3.39 (t, J = 7.9 Hz, 2H), 2.62 (t, J = 7.9

			Hz, 2H), 2.07 (quin, 2H).
23	N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine	Benzo[b]thiophen-5-amine (CAS 20532-28-9)	(CDCl ₃) 7.75 (d, J = 8.5 Hz, 1H), 7.47 (br. s., 1H), 7.40 (d, J = 5.6 Hz, 1H), 7.20 - 7.30 (m, 1H), 7.01 (dd, J = 8.5, 2.1 Hz, 1H), 3.38 - 3.53 (m, 2H), 2.57 (t, J = 7.9 Hz, 2H), 2.07 (qd, 2H).
24	N-1-Benzofuran-5-yl-3,4-dihydro-2H-pyrrol-5-amine	5-Benzofuranamine (CAS 58546-89-7)	(CDCl ₃) 7.57 (d, J = 2.3 Hz, 1H), 7.39 (d, J = 8.8 Hz, 1H), 7.20 - 7.30 (m, 1H), 6.92 (dd, J = 8.6, 2.2 Hz, 1H), 6.68 (d, J = 2.1 Hz, 1H), 3.45 (t, J = 6.6 Hz, 2H), 2.55 (t, J = 8.1 Hz, 2H), 2.06 (quin, 2H).
25	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine	1H-Indol-4-amine (CAS 5192-23-4)	(CD ₃ OD) 6.96 - 7.18 (m, 3H), 6.63 - 6.75 (m, 1H), 6.37 (d, J = 3.2 Hz, 1H), 3.42 (t, J = 6.9 Hz, 2H), 2.62 (t, J = 7.9 Hz, 2H), 1.96 - 2.18 (m, 2H).
26	N-[(2E)-Pyrrolidin-2-	3H-Pyrrolo[2,3-b]pyridin-5-amine	(CD ₃ OD) 7.90 (d, J = 2.1 Hz, 1H), 7.65 (d, J

	ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine	(CAS 1187421-28-8)	= 2.1 Hz, 1H), 7.33 (d, J = 3.5 Hz, 1H), 6.39 (d, J = 3.5 Hz, 1H), 3.43 (t, J = 6.9 Hz, 2H), 2.57 (t, J = 7.9 Hz, 2H), 2.03 (quin, 5 2H).
--	--	--------------------	---

Example 5**N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine**

10

(Compound 27)

5-(methylthio)-3,4-dihydro-2H-pyrrole (260 mg) was combined with 5-methyl- 6-quinoxalinamine, (360 mg) (CAS 171102-36-6) in 10 mL of isopropyl alcohol. The reaction was stirred at reflux for 2 days. The refluxed mixture was then cooled to room temperature and concentrated. Flash column chromatography (4-6% 7N NH₃ in MeOH/DCM) yielded the title compound. ¹H NMR (300 MHz, CD₃OD) δ 2.34 (m, 2H), 2.75 (m, 3H), 3.16 (t, J=7.92Hz, 2H), 3.71 (t, J=7.33Hz, 2H), 7.75 (d, J=8.79Hz, 1H), 8.07 (d, J=8.79Hz, 1H), 8.95 (dd, J=1.76Hz & 13.49Hz, 2H). The fumarate salt of the title compound was further obtained by crystallization with fumaric acid from ethanol. ¹H NMR (300 MHz, CD₃OD) δ 8.95 (dd, J = 13.5, 1.8 Hz, 2H), 8.07 (d, J = 8.5 Hz, 1H), 7.75 (d, J = 8.8 Hz, 1H), 6.66 (s, 2H), 3.72 (t, J = 7.2 Hz, 2H), 3.16 (t, J = 8.1 Hz, 2H), 2.75 (s, 3H), 2.21 - 2.44 (m, 2H). Free base: Fumaric acid = 1:1.

20

Compounds 28 to 31 and their corresponding fumarate salt were prepared in a similar manner to the method described in Example 5 for Compound 27. The reagents used and the results obtained are tabulated below in Table 5.

Table 5

Compound Number	IUPAC Name	Starting material	¹ H NMR (Solvent, δ ppm)
28	N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine	7-Quinolinamine (CAS 580-19-8)	(CDCl ₃) 2.09 (m, 2H), 2.62 (m, 2H), 3.47 (m, 2H), 7.26 (m, 2H), 7.59 (d, J=1.64Hz, 1H), 7.32 (d, J=8.79Hz, 1H), 8.05 (dd, J=1.76Hz & 8.05Hz, 1H), 8.81 (dd, J=1.76Hz & 4.10Hz, 2H).
29	N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine	7-amino-8-methylquinoline (CAS 116632-62-3)	Free base (CD ₃ OD) 2.33 (m, 2H), 2.74 (s, 3H), 3.15 (m, 2H), 3.71 (t, J=7.33Hz, 2H), 7.51 (d, J=8.8Hz, 1H), 7.60 (dd, J=4.60Hz & 8.32Hz, 1H), 7.93 (d, J=8.80Hz, 1H), 8.38 (dd, J=1.47Hz & 8.31Hz, 1H), 8.97 (dd, J=1.47Hz & 4.16Hz, 2H). Fumarate salt (CD ₃ OD) 8.97 (dd, J = 4.2, 1.7 Hz, 1H), 8.38 (dd, J = 8.2, 1.6 Hz, 1H), 7.94 (d, J = 8.6 Hz, 1H), 7.60 (dd, J = 8.2, 4.3 Hz, 1H), 7.52 (d, J = 8.6 Hz, 1H), 6.66 (s, 2H), 3.71 (t, J = 7.1 Hz, 3H), 3.17 (br. s., 3H), 2.74 (s, 3H), 2.34 (quin, 2H). Free base:Fumaric acid=1:1

30	N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-8-amine	8-aminoquanonoline (CAS 578-66-5)	<p>Free base</p> <p>(CD₃OD) 2.33 (m, 2H), 3.22 (t, J=8.20Hz, 2H), 3.72 (t, J=7.33Hz, 2H), 7.65 (m, 1H), 7.72 (d J=7.62Hz, 1H), 7.85 (dd, J=1.17Hz & 7.33Hz, 1H), 8.03 (dd, J=1.47Hz & 8.50Hz, 1H), 8.46 (dd, J=1.17Hz & 8.21Hz, 1H), 8.97 (dd, 1.76Hz & 4.40Hz, 1H).</p> <p>Fumarate salt</p> <p>(CD₃OD) 8.96 (dd, J = 4.3, 1.6 Hz, 1H), 8.46 (dd, J = 8.4, 1.3 Hz, 1H), 7.99 - 8.10 (m, 1H), 7.79 - 7.89 (m, 1H), 7.59 - 7.77 (m, 2H), 6.67 (s, 2H), 3.72 (t, J = 7.2 Hz, 2H), 3.22 (t, J = 8.2 Hz, 2H), 2.33 (quin, 2H).</p> <p>Free base:Fumaric acid=1:1</p>
31	4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine	5-Benzothiazolamine, 4-bromo- (CAS 769-19-7)	<p>Free base</p> <p>(CD₃OD) 9.43 (s, 1H), 8.22 (d, J = 8.5 Hz, 1H), 7.56 (d, J = 8.5 Hz, 1H), 6.68 (s, 2H), 3.71 (t, J = 7.3 Hz, 2H), 3.13 (t, J = 7.9 Hz, 2H), 2.23 - 2.43 (m, 2H).</p>

- 7.28 (m, 3H), 5.25 - 5.38 (m, 1H), 3.72 (t, J = 6.7 Hz, 2H), 2.91 - 3.04 (m, 1H), 2.77 - 2.91 (m, 1H), 2.54 - 2.72 (m, 1H), 2.35 - 2.54 (m, 2H), 1.93 - 2.10 (m, 2H), 1.76 - 1.91 (m, 1H).

5 Compounds 34 to 37 were prepared in a similar manner to the method described in Example 7 for Compound 33. The reagents used and the results obtained are tabulated below in Table 6.

Table 6

Compound Number	IUPAC Name	Starting material	¹ H NMR (Solvent) δ ppm
34	N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine	1-Naphthalenamine, 1,2,3,4-tetrahydro-, (1S)- (CAS 23357-52-0)	(CDCl ₃) 7.32 - 7.40 (m, 1H), 7.13 - 7.20 (m, 2H), 7.05 - 7.12 (m, 1H), 4.99 (t, J = 5.1 Hz, 1H), 3.73 (t, J = 6.7 Hz, 2H), 2.65 - 2.90 (m, 2H), 2.35 - 2.54 (m, 2H), 1.73 - 2.13 (m, 6H).
35	N-(7-Fluoro-2,3-dihydro-1H-inden-1-yl)-3,4-dihydro-2H-pyrrol-5-amine	1H-Inden-1-amine, 7-fluoro-2,3-dihydro (CAS 1071449-14-3)	(CD ₃ OD) 7.36 (td, J = 7.8, 5.3 Hz, 1H), 7.15 (d, J = 7.6 Hz, 1H), 6.93 - 7.02 (m, 1H), 5.28 (dd, J = 7.6, 4.4 Hz, 1H), 3.79 (t, J = 7.0 Hz, 2H), 3.08 - 3.24 (m, 1H), 2.93 - 3.05 (m, 1H), 2.83 - 2.91 (m, 2H), 2.51 - 2.71 (m, 1H), 2.25 (quin, J = 7.6 Hz, 2H), 2.06 - 2.18 (m, 1H).
36	N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-	2H-1,4-Benzoxazin-6-amine, 3,4-dihydro-5-methyl CAS 850560-41-7	(CD ₃ OD) 6.51 (d, J = 8.5 Hz, 1H), 6.19 (d, J = 8.5 Hz, 1H), 4.06 - 4.14 (m, 2H), 3.35 - 3.40 (m, 4H), 2.52 (t, J = 7.9 Hz, 2H), 1.95 - 2.10 (m, 2H), 1.91 (s, 3H).

	amine		
37	N-[(1S)-4-Methyl-2,3-dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine	(S)-4-methyl-2,3-dihydro-1H-inden-1-amine CAS 1071448-91-3	(CD ₃ OD) 6.96 - 7.15 (m, 3H), 5.15 (t, J = 7.3 Hz, 1H), 3.62 (t, J = 6.9 Hz, 2H), 2.86 - 3.00 (m, 1H), 2.66 - 2.80 (m, 1H), 2.44 - 2.62 (m, 3H), 1.91 - 2.05 (m, 2H), 1.72 - 1.89 (m, 1H).

Example 8

4-Methyl-N-[(2E)-pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine

(Compound 38)

5 To a solution of tetrabutylammonium nitrate (2.48 g, 8.0 mmol) in DCM (30 mL) was added trifluoroacetic anhydride (1.68 g, 8.0 mmol) at 0 °C under argon. Then 4-methyl-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridine, (1.2 g, 4.42 mmol)(CAS 1227270-73-6) in anhydrous DCM (20 mL) was added slowly to the above reaction mixture at 0 °C. The mixture was stirred at 0 °C for 1.5 h. The reaction was quenched with water and
10 extracted in dichloromethane. The dichloromethane layer was dried (MgSO₄) filtered, concentrated and purified by MPLC (solid load) using hexane:EtOAc (8:2) gave 4-methyl-5-nitro-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridine . ¹H NMR (300 MHz, CD₃OD) δ: 8.95 (s, 1H), 8.18 (d, J = 8.2 Hz, 2H), 8.00 (d, J = 4.1 Hz, 1H), 7.64 - 7.74 (m, 1H), 7.53 - 7.64 (m, 2H), 7.02 (d, J = 4.1 Hz, 1H), 2.77 (s, 3H).

15 To a mixture of 4-methyl-5-nitro-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridine, (410 mg, 1.29 mmol) in THF (15 mL) and EtOH (10 mL) was added 10% palladium on carbon (10 wt% of Pd/C; 45 mg) under argon. The mixture was hydrogenated using a hydrogen balloon at room temperature for 16 hours. The reaction mixture was flushed with nitrogen and filtered through a plug of Celite® and concentrated *in vacuo*. The
20 crude material was purified by flash column chromatography on silica gel using hexane:EtOAc (4:6) to get 4-methyl-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-amine, (370 mg) . ¹H NMR (300 MHz, CD₃OD) δ: 7.96 - 8.04 (m, 2H), 7.83 (s, 1H), 7.53 - 7.64 (m, 2H), 7.42 - 7.52 (m, 2H), 6.65 (d, J = 4.1 Hz, 1H), 2.26 (s, 3H).

A solution of 4-methyl-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-amine, (775
25 mg, 2.7 mmol) and 5-(methylthio)-3,4-dihydro-2H-pyrrole (345 mg) in *i*-PrOH (14 mL)

was heated at 105 °C for 3 days. The refluxed mixture was then cooled to room temperature and concentrated. Flash column chromatography (4-6% 7N NH₃ in MeOH/DCM) yielded N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-amine, (180 mg) ¹H NMR (300 MHz, CD₃OD) δ: 8.10 (d, J = 7.6 Hz, 2H), 7.89 (s, 1H), 7.75 (d, J = 4.1 Hz, 1H), 7.63 (s, 1H), 7.54 (d, J = 8.2 Hz, 2H), 6.77 (d, J = 4.1 Hz, 1H), 2.63 (s, 2H), 2.09 (t, J = 7.5 Hz, 3H).

A mixture of N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1-(phenylsulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-amine, (180 mg) in MeOH (10 mL) was added 5N aq. solution of NaOH (2 mL) and the mixture was heated at 80 °C for 4 h. Cooled to room temperature a white solid precipitated, which was filtered off. Silica gel was added to the filtrate and concentrated. This material was purified by using silica gel column chromatography (5% 7N NH₃ in MeOH/DCM) gave the title compound, (43mg). ¹H NMR (300 MHz, CD₃OD) δ: 7.75 (s, 1H), 7.29 (d, J = 3.5 Hz, 1H), 6.47 (d, J = 3.2 Hz, 1H), 3.38 (t, J = 6.7 Hz, 2H), 2.60 (t, J = 7.6 Hz, 2H), 2.37 (s, 3H), 2.08 (quin, J = 7.4 Hz, 2H).

Example 9

In Vitro Activity

The compounds described herein were tested for alpha adrenergic activity using a Fluorometric Imaging plate Reader (FLIPR) assay (Princen et al., 2003, Cytometry Part A. 51, pp. 35-45). Cells are loaded into a plate reader with a Ca²⁺ dye indicator and incubated with various compounds. This assay is adequate for monitoring the intracellular Ca²⁺ mobilization from fluorescent readings to evaluate receptor agonists. The compounds activity is expressed as EC₅₀ (nanomolar concentration that elicits a half-maximal response) and their relative efficacy compared to a standard full agonist (see Table 7 below). The compounds described herein activate alpha adrenergic receptors.

Table 7

Potency (nM); efficacy (EC₅₀); n/a: not available.

Compound number	IUPAC Name	Alpha 2A	Alpha 2B	Alpha 2C
1	N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine	61 (0.81)	30 (0.16)	5 (0.88)
2	N-(2-Methyl-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine	733 (0.61)	145 (0.18)	368 (0.85)

Compound number	IUPAC Name	Alpha 2A	Alpha 2B	Alpha 2C
3	N-(4-Bromo-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine	n/a	700 (0.15)	833 (0.85)
4	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	n/a	714 (0.19)	142 (1.06)
5	N-(3,4-Dihydro-2H-pyrrol-5-yl)-6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine	n/a	n/a	1282 (0.86)
6	4-Chloro-N-[(2E)-pyrrolidin-2-ylidene]-1,5-naphthyridin-3-amine	n/a	n/a	2597 (0.59)
7	N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine	n/a	1070 (0.33)	672 (0.84)
8	5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine	1069 (0.81)	2203 (0.21)	1051 (0.82)
9	8-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-7-amine	429 (1)	1049 (0.38)	218 (0.97)
10	4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	3 (1)	90 (0.76)	1.5 (0.94)
11	4-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	216 (0.98)	513 (0.69)	35 (1)
12	N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1H-benzimidazol-5-amine	203 (1)	510 (0.86)	58 (1)
13	7-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine	n/a	1838 (0.35)	435 (0.98)
14	N-(3,4-Dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine	n/a	n/a	2737 (0.66)
15	N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine	n/a	293 (0.14)	37 (0.93)
16	N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine	n/a	n/a	123 (0.85)
17	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine	n/a	10 (0.15)	9 (0.9)
18	N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine	n/a	39 (0.18)	69 (0.9)
19	N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine	n/a	142 (0.29)	12 (0.85)

Compound number	IUPAC Name	Alpha 2A	Alpha 2B	Alpha 2C
20	7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine	n/a	79 (0.61)	12 (0.95)
21	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine	n/a	n/a	280 (0.75)
22	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-7-amine	n/a	n/a	2161 (0.61)
23	N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine	n/a	n/a	89 (0.66)
24	N-1-Benzofuran-5-yl-3,4-dihydro-2H-pyrrol-5-amine	n/a	n/a	630 (0.51)
25	N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine	115 (0.7)	152 (0.42)	<10 (0.99)
26	N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine	118 (0.85)	17 (0.96)	6 (0.97)
27	N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate	121 (0.89)	n/a	16 (0.9)
28	N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine	n/a	n/a	104 (0.96)
29	N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine	6 (0.98)	80 (0.2)	<1 (1)
30	N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-8-amine	n/a	n/a	2974 (0.71)
31	4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine	96 (.98)	117 (.76)	25 (.92)
32	N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine	n/a	n/a	643 (0.95)
33	N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine	n/a	n/a	281 (0.68)
34	N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine	n/a	163 (0.43)	221 (0.8)
35	N-(7-Fluoro-2,3-dihydro-1H-inden-1-yl)-3,4-dihydro-2H-pyrrol-5-amine	n/a	618 (0.44)	560 (0.87)
36		583	579	327

Compound number	IUPAC Name	Alpha 2A	Alpha 2B	Alpha 2C
	N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-amine	(0.68)	(0.77)	(0.87)
37	N-[(1S)-4-Methyl-2,3-dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine	272 (0.43)	21 (0.9)	58 (0.94)
38	4-Methyl-N-[(2E)-pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine	555 (0.79)	n/a	22 (0.9)

Example 10

In-Vivo Intraocular Pressure Compound Screening


5 The experimental animals used, were Normotensive male Dutch-Belted rabbits over 6 months in age (n= 4/compound/dose screened). A single drop (50 µl) of the drug formulation, which yields 0.15% or 1.0% of the active metabolite was administered topically by pipette onto the right eye (treated eye) at approximately 0700 hours. Intraocular Pressure (IOP) of the rabbits (treated and untreated eyes) was measured 0

10 hours before and at 0.5, 1, 2, 3, 4, 6 and 8 hours after topical eyedrop administration. IOP at the time of eyedrop administration (0 hours) was used as a baseline value. Prior to the tonometric measurements, 0.05% proparacaine (50 µl) was administered to each eye. Tonometric IOP measurements were obtained with a Mentor Pneumontonmeter. Additionally, all studies were masked. At least 1 week of wash-out time was allowed for

15 each rabbit between dosings. All animals were examined for sedation, ocular irritation, and changes in pupil diameter throughout the course of the experiments. Compounds 9, 10, 11, 12, 27, 29 show potential in lowering IOP. In this assay, Compound 10 lowered IOP about 35%.

We claim:

1. A compound having formula I, its enantiomers, diastereoisomers, hydrates, solvates, crystal forms and individual isomers, tautomers or a pharmaceutically acceptable salt thereof



Formula I

wherein:

n is 0 or 1;

A is an aromatic or non-aromatic ring;

X is N, N-R¹, CH-R¹, C-R¹;

R¹ is H, C₁₋₆ alkyl or halogen;

R² is N, N-R³, CH-R³, C-R³, O or S;

R³ is H, halogen, or C₁₋₆ alkyl;

R⁴ is halogen, H, C₁₋₆ alkyl or part of a ring as a heteroatom (O, S, N) or as a C;

R⁵ is a heteroatom (S, N) or is a C and is always part of a ring;


R⁶ is H, C₁₋₆ alkyl, halogen or is part of a ring as a heteroatom (O, S, N) or as a C;

R⁴ and R⁵ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or with 1 to 2 halogen and is directly attached to the rest of the molecule; or

R⁵ and R⁶ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule;

except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)- 2H-pyrrol-5-amine.

2. A compound according to claim 1 of formula II



Formula II

wherein:

n is 1;

A is an aromatic ring;

X is C-R¹;

R¹ is H;

R² is N or C-R³;

R³ is H, halogen, or C₁₋₆ alkyl;


R⁴ is part of a ring as a heteroatom (O, S, N) or as a C;

R⁵ is a heteroatom (N) or is a C and is always part of a ring;

R⁶ is H, C₁₋₆ alkyl or halogen; and

R⁴ and R⁵ form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

3. A compound according to claim 1 of formula III



Formula III

wherein:

n is 1;

A is an aromatic ring;

X is C-R¹;

R¹ is H, C₁₋₃ alkyl;

R² is N or C-R³;

R³ is H, halogen, or C₁₋₆ alkyl;


R^4 is halogen or H;

R^5 is a heteroatom (N) or C and is always part of a ring;

R^6 is part of a ring as a heteroatom (N) or as a C;

- 5 R^5 and R^6 can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C_{1-6} alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

4 A compound according to claim 1 of formula IV



Formula IV

10 wherein:

n is 0;

A is a non-aromatic ring;

R^2 is $CH-R^3$;

R^3 is H;


15 R^4 is H;

R^5 is C and is always part of a ring;

R^6 is part of a ring as a C; and

R^5 and R^6 can form a 6 membered aromatic carbocyclic ring directly attached to the rest of the molecule.

20 5 A compound according to claim 1 of formula V



Formula V

wherein

n is 1;

25 A is a non-aromatic ring;

X is CH-R¹;

R¹ is H;

R² is CH-R³;

R³ is H;

5 R⁴ is H;

R⁵ is C and is always part of a ring;

R⁶ is part of a ring as a C; and

R⁵ and R⁶ can form a 6 membered aromatic carbocyclic ring, optionally substituted with C₁₋₆ alkyl or halogen and is directly attached to the rest of the molecule;

except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine.

6. A compound according to claim 1 selected from the group consisting of:

N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine;

15 N-(2-Methyl-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;

N-(4-Bromo-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine;

4-Chloro-N-[(2E)-pyrrolidin-2-ylidene]-1,5-naphthyridin-3-amine;

20 N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine fumarate;

N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;

5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine fumarate;

5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;

8-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-7-amine;

25 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;

4-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;

N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1H-benzimidazol-5-amine;

7-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;

30 N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;

N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;

N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;


- N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;
7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-7-amine;
5 N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
N-1-Benzofuran-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine;
N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate;
10 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;
N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-8-amine;
15 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine fumarate;
N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine;
N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
20 N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
N-(7-Fluoro-2,3-dihydro-1H-inden-1-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-amine;
N-[(1S)-4-Methyl-2,3-dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
4-Methyl-N-[(2E)-pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine.
25 7. A compound according to claim 1 selected from the group consisting of:
N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;
30 N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;

7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine;
 N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine;
 5 N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine ;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine;
 10 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;
 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
 N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
 N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine.

15 8. A compound according to claim 1 selected from the group consisting of:

4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
 N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;
 20 N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;
 7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
 N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;
 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
 25 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine.

9. A pharmaceutical composition comprising at least one compound of having formula I:



Formula I

wherein:

n is 0 or 1;

A is an aromatic or non-aromatic ring;

5 X is N, N-R¹, CH-R¹, C-R¹;

R¹ is H, C₁₋₆ alkyl or halogen;

R² is N, N-R³, CH-R³, C-R³, O or S;

R³ is H, halogen, or C₁₋₆ alkyl;

R⁴ is halogen, H, C₁₋₆ alkyl or part of a ring as a heteroatom (O, S, N) or as a C;

10 R⁵ is a heteroatom (S, N) or is a C and is always part of a ring;

R⁶ is H, C₁₋₆ alkyl, halogen or is part of a ring as a heteroatom (O, S, N) or as a C;


R⁴ and R⁵ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or with 1 to 2 halogen and is directly attached to the rest of the molecule; or

15 R⁵ and R⁶ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule;

except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine;

and a pharmaceutically acceptable carrier.

20 10. The pharmaceutical composition of claim 9, comprising at least one compound of formula II:



Formula II

25 wherein:

n is 1;

A is an aromatic ring;

X is C-R¹;

R¹ is H;

30 R² is N or C-R³;

R³ is H, halogen, or C₁₋₆ alkyl;


R^4 is part of a ring as a heteroatom (O, S, N) or as a C;

R^5 is a heteroatom (N) or is a C and is always part of a ring;

R^6 is H, C₁₋₆ alkyl or halogen; and

R^4 and R^5 form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

11. The pharmaceutical composition of claim 9, comprising at least one compound of formula III:



Formula III

wherein:

n is 1;

A is an aromatic ring;

X is C- R^1 ;

R^1 is H, C₁₋₃ alkyl;

R^2 is N or C- R^3 ;

R^3 is H, halogen, or C₁₋₆ alkyl;


R^4 is halogen or H;

R^5 is a heteroatom (N) or C and is always part of a ring;

R^6 is part of a ring as a heteroatom (N) or as a C;

R^5 and R^6 can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule.

12. The pharmaceutical composition of claim 9, comprising at least one compound of formula IV:



Formula IV

wherein:

n is 0;

5 A is a non-aromatic ring;

R² is CH-R³;

R³ is H;


R⁴ is H;

R⁵ is C and is always part of a ring;

10 R⁶ is part of a ring as a C; and

R⁵ and R⁶ can form a 6 membered aromatic carbocyclic ring directly attached to the rest of the molecule.

13. The pharmaceutical composition of claim 9, comprising at least one compound of formula V :



Formula V

wherein

n is 1;

A is a non-aromatic ring;

20 X is CH-R¹;

R¹ is H;

R² is CH-R³;

R³ is H;

R⁴ is H;

25 R⁵ is C and is always part of a ring;

R⁶ is part of a ring as a C; and

R⁵ and R⁶ can form a 6 membered aromatic carbocyclic ring, optionally substituted with C₁₋₆ alkyl or halogen and is directly attached to the rest of the molecule;


except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-2H-pyrrol-5-amine.

14. The pharmaceutical composition of claim 9, comprising at least one compound selected from the group consisting of:

N-1-Naphthyl-3,4-dihydro-2H-pyrrol-5-amine;
N-(2-Methyl-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(4-Bromo-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-6,7-dihydro-5H-cyclopenta[b]pyridin-3-amine;
4-Chloro-N-[(2E)-pyrrolidin-2-ylidene]-1,5-naphthyridin-3-amine;
N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine fumarate;
N-(4-Chloro-1-naphthyl)-3,4-dihydro-2H-pyrrol-5-amine;
5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine fumarate;
5-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;
8-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-7-amine;
4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
4-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(3,4-dihydro-2H-pyrrol-5-yl)-4-methyl-1H-benzimidazol-5-amine;
7-Chloro-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-benzimidazol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)quinoxalin-6-amine;
N-(5,6,7,8-Tetrahydronaphthalen-2-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(2,3-Dihydro-1H-inden-4-yl)-3,4-dihydro-2H-pyrrol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-4-methyl-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-7-methyl-1H-indol-5-amine;
7-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1H-indol-5-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-6-amine;
N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-7-amine;
N-1-Benzothien-5-yl-3,4-dihydro-2H-pyrrol-5-amine;
N-1-Benzofuran-5-yl-3,4-dihydro-2H-pyrrol-5-amine;

- N-(3,4-Dihydro-2H-pyrrol-5-yl)-1H-indol-4-amine;
 N-[(2E)-Pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine fumarate;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methylquinoxalin-6-amine;
 5 N-(3,4-Dihydro-2H-pyrrol-5-yl)quinolin-7-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-8-methylquinolin-7-amine fumarate;
 N-(3,4-dihydro-2H-pyrrol-5-yl)quinolin-8-amine;
 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine fumarate;
 10 4-Bromo-N-(3,4-dihydro-2H-pyrrol-5-yl)-1,3-benzothiazol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine fumarate;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)isoquinolin-8-amine;
 N-[(1S)-2,3-Dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
 N-[(1S)-1,2,3,4-Tetrahydronaphthalen-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
 15 N-(7-Fluoro-2,3-dihydro-1H-inden-1-yl)-3,4-dihydro-2H-pyrrol-5-amine;
 N-(3,4-Dihydro-2H-pyrrol-5-yl)-5-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-amine;
 N-[(1S)-4-Methyl-2,3-dihydro-1H-inden-1-yl]-3,4-dihydro-2H-pyrrol-5-amine;
 4-Methyl-N-[(2E)-pyrrolidin-2-ylidene]-1H-pyrrolo[2,3-b]pyridin-5-amine.

15. A method for the treatment of a disease or condition responsive to
 20 treatment by modulation of an alpha adrenergic receptor, comprising administering to a
 mammal in need thereof, a therapeutically effective amount of a compound of formula I:



Formula I

wherein:

- 25 n is 0 or 1;
 A is an aromatic or non-aromatic ring;
 X is N, N-R¹, CH-R¹, C-R¹;
 R¹ is H, C₁₋₆ alkyl or halogen;
 R² is N, N-R³, CH-R³, C-R³, O or S;

R³ is H, halogen, or C₁₋₆ alkyl;

R⁴ is halogen, H, C₁₋₆ alkyl or part of a ring as a heteroatom (O, S, N) or as a C;

R⁵ is a heteroatom (S, N) or is a C and is always part of a ring;

R⁶ is H, C₁₋₆ alkyl, halogen or is part of a ring as a heteroatom (O, S, N) or as a

5 C;

R⁴ and R⁵ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or with 1 to 2 halogen and is directly attached to the rest of the molecule; or

10 R⁵ and R⁶ can form a 5 or 6 membered carbo- or heterocyclic ring, aromatic or non-aromatic, optionally substituted with 1 to 2 C₁₋₆ alkyl or 1 to 2 halogen and is directly attached to the rest of the molecule;

except (+/-) 3,4-dihydro-N-(1,2,3,4-tetrahydro-1-naphthalenyl)- 2H-pyrrol-5-amine; and a pharmaceutically acceptable carrier.

15 16. The method of claim 15, wherein the disease or condition is selected from the group consisting of chronic pain, visceral pain, neuropathic pain, allodynia and combinations thereof.

17. The method of claim 15, wherein the disease or condition is selected from the group consisting of dermatological conditions, skin erythema (redness) and inflammation, rosacea, acne, psoriasis.

20 18. The method of claim 15, wherein the disease or condition is glaucoma.

19. The method of claim 15, wherein the disease or condition is elevated intraocular pressure.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2010/051622

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D207/22 C07D403/12 C07D409/12 C07D413/12 C07D417/12
C07D471/04 A61K31/4025
ADD. A61P9/00 A61P29/00 A61P27/06 A61P17/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DE JONG A P ET AL: "Relationships between structure and alpha-adrenergic receptor affinity of clonidine and some related cyclic amidines", EUROPEAN JOURNAL OF PHARMACOLOGY, ELSEVIER BV, NL, vol. 69, no. 2, 16 January 1981 (1981-01-16), pages 175-188, XP023840198, ISSN: 0014-2999, DOI: DOI:10.1016/0014-2999(81)90412-X [retrieved on 1981-01-16] page 175 - page 177; tables 2,3,4 ----- -/-</p>	1-19



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 January 2011

Date of mailing of the international search report

22/02/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Rufet, Jacques

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/051622

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	STEPHAN SCHANN ET AL.: "Synthesis and piological evaluation of pyrrolinic isosteres of Rilmenidine", JOURNAL OF MEDICINAL CHEMISTRY., vol. 44, no. 10, 24 March 2001 (2001-03-24), pages 1588-1593, XP002619044, USAMERICAN CHEMICAL SOCIETY. WASHINGTON. ISSN: 0022-2623 the whole document	1-19
Y	UROSEVIC DRAGAN ET AL.: "LNP 906, the first high-affinity photoaffinity ligand selective for I1 imidazoline receptors", BRITISH JOURNAL OF PHARMACOLOGY, vol. 142, no. 3, 1 June 2004 (2004-06-01), XP002619045, NATURE PUBLISHING GROUP ISSN: 0007-1188 the whole document	1-19
Y	MUNK S A ET AL: "Analogues of UK 14,304: structural features responsible for alpha2 adrenoceptor activity", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 5, no. 15, 3 August 1995 (1995-08-03), pages 1745-1750, XP004135419, ISSN: 0960-894X, DOI: DOI:10.1016/0960-894X(95)00295-5 cited in the application the whole document	1-19
Y	CHAPLEO C B ET AL: "HETEROAROMATIC ANALOGUES OF THE ALPHA2-ADRENORECEPTOR PARTIAL AGONIST CLONIDINE", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 32, no. 7, 1 July 1989 (1989-07-01), pages 1627-1630, XP000601554, ISSN: 0022-2623, DOI: DOI:10.1021/JM00127A037 cited in the application the whole document	1-19

-/--

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/051622

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE CA [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 25 June 2009 (2009-06-25), GOLDFARB, DAVID SCOTT: "Method using lifespan-altering compounds for altering the lifespan of eukaryotic organisms, and screening for such compounds", XP002619046, retrieved from STN Database accession no. 2009:846114 * abstract & US 2009/163545 A1 (GOLDFARB, DAVID SCOTT) 25 June 2009 (2009-06-25) -----</p>	1,9