



(43) International Publication Date
28 September 2017 (28.09.2017)

WIPO | PCT

(10) International Publication Number
WO 2017/165615 A1

(51) International Patent Classification:

A61K 8/33 (2006.01) *A61Q 5/12* (2006.01)
A61Q 5/02 (2006.01) *A61K 8/23* (2006.01)

(21) International Application Number:

PCT/US2017/023758

(22) International Filing Date:

23 March 2017 (23.03.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/312,856 24 March 2016 (24.03.2016) US

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio
45202 (US).

(72) Inventors: **HERTENSTEIN, Stacy, Renee**; One Procter
& Gamble Plaza, Cincinnati, Ohio 45202 (US).
HOLLINGSHEAD, Judith, Ann; One Procter & Gamble
Plaza, Cincinnati, Ohio 45202 (US). **CORDER, Jennifer,**
Anne; One Procter & Gamble Plaza, Cincinnati, Ohio
45202 (US). **STAUDIGEL, James, Anthony**; One Procter
& Gamble Plaza, Cincinnati, Ohio 45202 (US).

(74) Agent: **KREBS, Jay A.**; c/o The Procter & Gamble Com-
pany, Global IP Services, One Procter & Gamble Plaza,
C9, Cincinnati, Ohio 45202 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: HAIR CARE COMPOSITIONS COMPRISING MALODOR REDUCTION COMPOSITIONS

(57) Abstract: The present invention relates to hair care compositions comprising malodor reduction compositions and methods of using such hair care compositions. Such hair care compositions comprising the malodor control technologies disclosed herein provide malodor control without leaving an undesirable scent and when perfume is used to scent such compositions, such scent is not unduly altered by the malodor control technology.



WO 2017/165615 A1

HAIR CARE COMPOSITIONS COMPRISING MALODOR REDUCTION COMPOSITIONS

FIELD OF THE INVENTION

5 The present invention relates to hair care compositions comprising malodor reduction compositions and methods of making and using such hair care compositions.

BACKGROUND OF THE INVENTION

10 Unscented or lightly scented products are desired by consumers as they may be considered more natural and discreet than more highly scented products. Manufacturers of unscented or lightly scented products for controlling malodors rely on malodor reduction ingredients or other technologies (e.g. filters) to reduce malodors. However, effectively controlling malodors, for example, amine-based malodors (e.g. fish and urine), thiol and sulfide-based malodors (e.g. garlic and onion), C₂-C₁₂ carboxylic acid based malodors (e.g. body and pet odor), indole based malodors
15 (e.g. fecal and bad breath), short chain fatty aldehyde based malodors (e.g. grease) and geosmin based malodors (e.g. mold/mildew) may be difficult, and the time required for a product to noticeably reduce malodors may create consumer doubt as to the product's efficacy on malodors. Often times, manufacturers incorporate scented perfumes to help mask these difficult malodors.

20 Unfortunately, malodor control technologies typically cover up the malodor with a stronger scent and thus interfere with the scent of the perfumed or unperfumed situs that is treated with the malodor control technology. Thus, limited nature of the current malodor control technologies is extremely constraining. Thus what is needed is a broader palette of malodor control technologies so the perfume community can deliver the desired level of character in a greater number of situations/applications. Surprisingly, Applicants recognized that in addition to blocking a
25 malodor's access to a sensory cell, in order to achieve the desired goal, a malodor control technology must leave such sensor cell open to other molecules, for example scent molecules. Thus, hair care compositions comprising the malodor control technologies disclosed herein provide malodor control without leaving an undesirable scent and, when perfume is used to scent such compositions, such scent is not unduly altered by the malodor control technology.

30 Selenium sulfide containing anti-fungal hair and scalp care compositions provide some of the most effective protection from and relief of dandruff conditions. Historically, selenium sulfide and other sulfur-based formulations are highly medicinal and pungent smelling- both in use and throughout the day- due to residual sulfur compounds deposited on the hair and scalp. These

significant negative cosmetic attributes may cause consumers to avoid selenium sulfide and other sulfur-based formulations and therefore product usage compliance is difficult and as a result consumers often do not find complete relief from their dandruff condition.

5

SUMMARY OF THE INVENTION

The present invention relates a hair care composition comprising, based on total composition weight: a sum total of from about 0.0001% to about 2% of alpha, alpha, 6, 6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal) and one or more malodor reduction materials having a Sulfur MORV >3; from about 0.01% to about 10% of a scalp active material selected from the group consisting of selenium sulfide, sulfur and mixtures thereof; from about 0.1% to about 40%, of a surfactant.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

“Hair Care” composition” as defined herein, may include shampoos, conditioners and
10 leave-on-treatments.

“Rinse-off” means the intended product usage includes application to hair followed by rinsing and/or wiping the product from the skin and/or hair within a few seconds to minutes of the application step.

“STnS” refers to sodium trideceth(n) sulfate, wherein n can define the average number of
15 moles of ethoxylate per molecule.

As used herein “MORV” is the calculated malodor reduction value for a subject material. A material’s MORV indicates such material’s ability to decrease or even eliminate the perception of one or more malodors. For purposes of the present application, a material’s MORV is calculated in accordance with method found in the test methods section of the present application.

20 As used herein, “malodor” refers to compounds generally offensive or unpleasant to most people, such as the complex odors associated with bowel movements.

As used herein, “odor blocking” refers to the ability of a compound to reduce the perception of a malodor.

25 As used herein, the term “perfume” does not include malodor reduction materials. Thus, the perfume portion of a composition does not include, when determining the perfume’s composition, any malodor reduction materials found in the composition as such malodor reduction materials are described herein. In short, if a material has a malodor reduction value “MORV” that

is within the range of the MORV recited in the subject claim, such material is a malodor reduction material for purposes of such claim.

As used herein, the terms “a” and “an” mean “at least one”.

As used herein, the terms “include”, “includes” and “including” are meant to be non-limiting.

Unless otherwise noted, all component or composition levels are in reference to the active portion of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources of such components or compositions.

All percentages and ratios are calculated by weight unless otherwise indicated. All percentages and ratios are calculated based on the total composition unless otherwise indicated.

It should be understood that every maximum numerical limitation given throughout this specification includes every lower numerical limitation, as if such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

Hair Care Compositions

Rinse-off hair care compositions can come in a variety of forms. For example, a hair care composition can be in a liquid form and can be a shampoo, conditioning shampoo,

Hair Care compositions can include perfume materials. Many consumers prefer hair care compositions that can consistently provide a desired scent, or odor that can be perceived each time the product is used. Perfume materials can provide the desired scent or odor to these hair care compositions. These perfume (i.e., fragrance) materials can include perfumes, perfume raw materials, and perfume delivery systems.

Malodor Reduction Materials

A non-limiting set of suitable malodor reduction materials are provided in the tables below.

Table 1

List of materials with Sulfur MORV >3

<u>Number</u>	<u>CAS Number</u>	<u>Name</u>	ACD Room Temp (25 C) Vapor Pressure (mm Hg, ACD, EXP-14-AA6010-003)	Octanol/Water Partition Coefficient, logP (ACD, Consensus algorithm, EXP-14-AA6010-002)	Malodor Reduction: Log (1/EC0) 3-methyl-3-sulfanylh exan-1-ol (3M3S1H)
	188199-50-0	2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane]	2.3E-02	4.30	3.04
	154171-77-4	(1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene] K	1.5E-02	4.51	4.66
	5413-60-5	3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate)	1.4E-02	2.79	3.44
	68480-11-5	Decahydro-3H-spiro[furan-2,5'-[4,7]methanoindene]	1.4E-02	3.36	3.22
	19870-74-7	CEDRYL METHYL ETHER	1.3E-02	5.08	5.52
	116126-82-0	Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate	1.2E-02	3.68	3.22
	3738-00-9	3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan	9.3E-03	5.11	3.06

33885-52-8	alpha,alpha,6,6-tetramethylbicyclo[3.1.1]hept-2-ene-propanal	6.9E-03	4.31	3.18
1139-30-6	4,5-epoxy-4,11,11-trimethyl-8-methylenebicyclo(7.2.0)undecane)	6.7E-03	4.47	3.57
41724-19-0	4aR,8aS)-7-methyloctahydro-1,4-methanonaphthalen-6(2H)-one	6.5E-03	2.66	3.43
86803-90-9	5-methoxyoctahydro-1H-4,7-methanoindene-2-carbaldehyde	4.4E-03	2.08	3.87
68738-96-5	8,8-dimethyl-6,7-dihydro-5H-naphthalene-2-carbaldehyde	4.4E-03	3.92	3.11
41816-03-9	2R,4a'R,8a'R)-3,7'-dimethyl-3',4',4a',5',8',8a'-hexahydro-1'H-spiro[oxirane-2,2'-[1,4]methanonaphthalene])	3.3E-03	3.09	3.57
476332-65-7	(2,2,6,6,7,8,8-heptamethyldecahydro-2H-indeno[4,5-b]furan	3.2E-03	6.14	5.66
23787-90-8	1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one)	2.6E-03	4.09	4.53
67634-20-2	3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-yl isobutyrate	2.4E-03	3.51	4.13
57345-19-4	3S,5aR,7aS,11aS,11bR)-3,8,8,11a-tetramethyldodecahydro-5H-3,5a-epoxynaphtho[2,1-c]oxepine	2.0E-03	5.18	4.97
68912-13-0	(8,8-dimethyl-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl propionate)	1.8E-03	4.00	5.04

211299-54-6	4aR,5R,7aS,9R)-2,2,5,8,8,9a-hexamethyloctahydro-4H-4a,9-methanoazuleno[5,6-d][1,3]dioxole	1.8E-03	4.85	6.82
68901-32-6	2-(8-isopropyl-6-methylbicyclo[2.2.2]oct-5-en-2-yl)-1,3-dioxolane	1.2E-03	3.81	3.04
68039-44-1	3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl pivalate	1.2E-03	3.96	3.79
823178-41-2	3a,5,6,7,8,8b-hexahydro-2,2,6,6,7,8,8-heptamethyl-4H-indeno(4,5-d)-1,3-dioxole	7.4E-04	4.97	5.24
39900-38-4	(3R-(3alpha,3a,6alpha,7,8aalpha))-octahydro-3,6,8,8-tetramethyl-1H-3a,7-methanoazulen-6-yl formate	6.3E-04	4.97	4.83
77-53-2	(1S,2R,5S,7R,8R)-2,6,6,8-tetramethyltricyclo[5.3.1.0 ^{1,5}]undecan-8-ol	5.7E-04	4.49	4.40
54464-57-2	1-((2S,3S)-2,3,8,8-tetramethyl-1,2,3,4,5,6,7,8-octahydronaphthalen-2-yl)ethan-1-one	5.4E-04	4.72	3.26
30168-23-1	((E)-4-((3aS,7aS)-octahydro-5H-4,7-methanoinden-5-ylidene)butanal	4.4E-04	3.97	4.42
5986-55-0	1R-(1alpha,4beta,4aalpha,6beta,8aalpha))-octahydro-4,8a,9,9-tetramethyl-1,6-methano-1(2H)-naphthol	2.8E-04	4.46	4.34
32214-91-8	[(3Z)-4,11,11-trimethyl-8-methylidene-5-	2.5E-04	5.50	3.55

		bicyclo[7.2.0]undec-3-enyl] acetate			
552-02-3		(1aR,4S,4aS,7R,7aS,7bS)- 1,1,4,7-tetramethyldecahydro- 1H-cyclopropa[e]azulen-4-ol	1.8E-04	4.72	4.12
69486-14-2		Z)-6-ethylideneoctahydro-2H- 5,8-methanochromen-2-one)	1.1E-04	2.32	3.82
32388-55-9		1-((3R,3aR,7R,8aS)-3,6,8,8- tetramethyl-2,3,4,7,8,8a- hexahydro-1H-3a,7- methanoazulen-5-yl)ethan-1-one	8.5E-05	4.97	4.49
167254-80-0		3,5,5,6,7,8,8-heptamethyl- 5,6,7,8-tetrahydronaphthalene-2- carbonitrile	6.9E-05	5.88	3.36
66072-32-0		4-(1,7,7-trimethyl-6- bicyclo[2.2.1]heptanyl)cyclohexa n-1-ol	3.0E-05	4.45	3.50
501929-47-1		(E)-4-((3aR,4R,7R,7aR)- 1,3a,4,6,7,7a-hexahydro-5H-4,7- methanoinden-5-ylidene)-3- methylbutan-2-ol	8.5E-06	3.87	5.44
3681-73-0		(E)-3,7-dimethylocta-2,6-dien-1- yl palmitate)	3.0E-09	10.75	3.31

Table 2**List of materials with Sulfur MORV >3; ClogP>3 and VP>.005**

5

<u>Number</u>	<u>Material Name</u>	<u>CAS</u> <u>Numbe</u> <u>r</u>	ACD Room Temp (25 C) Vapor	Octanol/Wat er Partition Coefficient, logP (ACD, Consensus	Malodor Reduction : Log (1/EC0) 3- methyl-3-

			Pressur e (mm Hg, ACD, EXP- 14- AA601 0-003)	algorithm, EXP-14- AA6010-002)	sulfanylhe xan-1-ol (3M3S1H)
1	2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane	188199 -50-0	2.3E-02	4.30	3.04
2	(1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene] K	154171 -77-4	1.5E-02	4.51	4.66
3	SPIRO[FURAN-2(3H),5'-(4,7-METHANO-5H-INDENE], DECAHYDRO	68480- 11-5	1.4E-02	3.36	3.22
4	(3R,3aR,6S,7S,8aS)-6-methoxy-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulene	19870- 74-7	1.3E-02	5.08	5.52
5	Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate	116126 -82-0	1.2E-02	3.68	3.22
6	3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan	3738- 00-9	9.3E-03	5.11	3.06
7	(alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal)	33885- 52-8	6.9E-03	4.31	3.18

8	4,5-epoxy-4,11,11-trimethyl-8-methylenebicyclo(7.2.0)undecane)	1139-30-6	6.7E-03	4.47	3.57
9	2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane	188199-50-0	2.3E-02	4.30	3.04

Table 3. Sulfur MORV>3 and ClogP>3

11	1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophthalen-8(5H)-one)	23787-90-8	2.6E-03	4.09	4.53
12	8,8-dimethyl-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl propionate)	68912-13-0	1.8E-03	4.00	5.04
13	2-(8-isopropyl-6-methylbicyclo[2.2.2]oct-5-en-2-yl)-1,3-dioxolane	68901-32-6	1.2E-03	3.81	3.04
14	[(3Z)-4,11,11-trimethyl-8-methylidene-5-bicyclo[7.2.0]undec-3-enyl]acetate	32214-91-8	2.5E-04	5.50	3.55
	(E)-3,7-dimethylocta-2,6-dien-1-yl palmitate)	3681-73-0	3.0E-09	10.75	3.31

- 5 The materials in Tables 1-3 may be supplied by one or more, but not limited to, the following: Firmenich Inc. of Plainsboro NJ USA; International Flavor and Fragrance Inc. New York, NY USA; Takasago Corp. Teterboro, NJ USA; Symrise Inc. Teterboro, NJ USA; Sigma-Aldrich/SAFC Inc. Carlsbad, CA USA; and Bedoukian Research Inc. Danbury, CT USA.

In one aspect of said hair care composition, said composition comprises one or more
 10 perfume raw materials.

In one aspect of said hair care composition, said composition comprises a total of, based on total consumer product weight, from about 0.1% to about 7% of one or more of said malodor reduction materials and from about 3% to 30% of a surfactant, and, optionally, a miscellar phase and/or lamellar phase.

5 In one aspect of said hair care composition, said composition comprises a total, based on total consumer product weight, of from about 0.1% to about 50% of a material selected from structurants, humectants, fatty acids, inorganic salts, antimicrobial agents, antimicrobial agents actives and mixtures thereof.

10 In one aspect of said hair care composition, said composition comprises an adjunct ingredient selected from the group consisting of clay mineral powders, pearl pigments, organic powders, emulsifiers, distributing agents, pharmaceutical active, topical active, preservatives, surfactants and mixtures thereof.

A method of controlling malodors comprising: contacting a situs comprising a malodor and/or a situs that will become malodorous with a hair care composition selected from the group
15 consisting of the hair care compositions disclosed herein is disclosed.

In one aspect of said method, said situs comprises the head of hair and said contacting step comprises contacting said hair containing a malodor with a sufficient amount of present invention's hair care composition to provide said hair with a level of malodor reduction material at least 0.0001 mg of malodor reduction material per body or head of hair, in an embodiment from about 0.0001
20 mg of malodor reduction material per head of hair to about 1 mg of malodor reduction material per head of hair, in a further embodiment from about 0.001 mg of malodor reduction material per head of hair about 0.5 mg of malodor reduction material per body or head of hair, a further embodiment from about 0.01 of malodor reduction material per head of hair to about 0.2 mg of malodor reduction material per head of hair.

25

Adjunct Materials

While not essential for the purposes of the present invention, the non-limiting list of adjuncts illustrated hereinafter are suitable for use in the instant compositions and may be desirably incorporated in certain aspects of the invention, for example to assist or enhance
30 performance,

A variety of optional ingredients can also be added to hair care compositions. Optional ingredients can include, but are not limited to, structurants, humectants, fatty acids, inorganic salts, and other antimicrobial agents or actives.

A hair care composition can also include hydrophilic structurants such as carbohydrate structurants and gums. Some suitable carbohydrate structurants include raw starch (corn, rice, potato, wheat, and the like) and pregelatinized starch. Some suitable gums include carageenan and xanthan gum. A hair care composition can include from about 0.1% to about 30%, from about 5 2% to about 25%, or from about 4% to about 20%, by weight of the hair care composition, of a carbohydrate structurant.

A hair care composition can also include one or more humectants. Examples of such humectants can include polyhydric alcohols. Further, humectants such as glycerin can be included the hair care composition as a result of production or as an additional ingredient. For example, 10 glycerin can be a by-product after saponification of the hair care composition. Including additional humectant can result in a number of benefits such as improvement in hardness of the hair care composition, decreased water activity of the hair care composition, and reduction of a weight loss rate of the hair care composition over time due to water evaporation.

A hair care composition can include inorganic salts. Inorganic salts can help to maintain a 15 particular water content or level of the hair care composition and improve hardness of the hair care composition. The inorganic salts can also help to bind the water in the hair care composition to prevent water loss by evaporation or other means. A hair care composition can optionally include from about 0.01% to about 15%, from about 1% to about 12%, or from about 2.5% to about 10.5%, by weight of the hair care composition, of inorganic salt. Examples of suitable inorganic salts can 20 include magnesium nitrate, trimagnesium phosphate, calcium chloride, sodium carbonate, sodium aluminum sulfate, disodium phosphate, sodium polymetaphosphate, sodium magnesium succinate, sodium tripolyphosphate, aluminum sulfate, aluminum chloride, aluminum chlorohydrate, aluminum-zirconium trichlorohydrate, aluminum-zirconium trichlorohydrate glycine complex, zinc sulfate, ammonium chloride, ammonium phosphate, calcium acetate, calcium nitrate, calcium 25 phosphate, calcium sulfate, ferric sulfate, magnesium chloride, magnesium sulfate, and tetrasodium pyrophosphate.

A hair care composition can include one or more additional antibacterial agents that can serve to further enhance antimicrobial effectiveness of the hair care composition. A hair care composition can include, for example, from about 0.001% to about 2%, from about 0.01% to about 30 1.5%, or from about 0.1% to about 1%, by weight of the hair care composition, of additional antibacterial agent(s). Examples of suitable antibacterial agents can include carbanilides, triclocarban (also known as trichlorocarbanilide), triclosan, a halogenated diphenylether available as DP-300 from Ciba-Geigy, hexachlorophene, 3,4,5-tribromosalicylanilide, and salts of 2-

pyridinethiol-1-oxide, salicylic acid, and other organic acids. Other suitable antibacterial agents are described in U.S. Patent No. 6,488,943.

Scalp Active Material

5 In an embodiment of the present invention, the hair care composition may comprise a scalp active material, which may be an anti-dandruff active. In an embodiment, the anti-dandruff active is selected from the group consisting of: pyridinethione salts; zinc carbonate; azoles, such as ketoconazole, econazole, and elubiol; selenium sulfide; particulate sulfur; keratolytic agents such as salicylic acid; and mixtures thereof. In a further embodiment, the anti-dandruff active may be
10 an anti-dandruff particulate. In an embodiment, the anti-dandruff particulate is a pyridinethione salt. Such anti-dandruff particulate should be physically and chemically compatible with the components of the composition, and should not otherwise unduly impair product stability, aesthetics or performance.

In an embodiment of the present invention, the selenium sulfide is present from about about
15 0.01% to about 10%, in an embodiment from about 0.1% to about 8%, in a further embodiment, from about 1% to about 5%.

In an embodiment of the present invention, the sulfur is present from about 0.01% to about 10%, in an embodiment from about 0.1% to about 8%, in a further embodiment, from about 1% to about 5% and from about 2% to about 5%.

20 Pyridinethione particulates are suitable particulate anti-dandruff actives for use in composition of the present invention. In an embodiment, the anti-dandruff active is a 1-hydroxy-2-pyridinethione salt and is in particulate form. In an embodiment, the concentration of pyridinethione anti-dandruff particulate ranges from about 0.01% to about 5%, by weight of the composition, or from about 0.1% to about 3%, or from about 0.1% to about 2%. In an embodiment,
25 the pyridinethione salts are those formed from heavy metals such as zinc, tin, cadmium, magnesium, aluminium and zirconium, generally zinc, typically the zinc salt of 1-hydroxy-2-pyridinethione (known as “zinc pyridinethione” or “ZPT”; zinc pyrithione), commonly 1-hydroxy-2-pyridinethione salts in platelet particle form. In an embodiment, the 1-hydroxy-2-pyridinethione salts in platelet particle form have an average particle size of up to about 20 microns, or up to about
30 5 microns, or up to about 2.5 microns. Salts formed from other cations, such as sodium, may also be suitable. Pyridinethione anti-dandruff actives are described, for example, in U.S. Pat. No. 2,809,971; U.S. Pat. No. 3,236,733; U.S. Pat. No. 3,753,196; U.S. Pat. No. 3,761,418; U.S. Pat. No. 4,345,080; U.S. Pat. No. 4,323,683; U.S. Pat. No. 4,379,753; and U.S. Pat. No. 4,470,982.

In an embodiment, in addition to the anti-dandruff active selected from polyvalent metal salts of pyrithione, the composition further comprises one or more anti-fungal and/or anti-microbial actives. In an embodiment, the anti-microbial active is selected from the group consisting of: coal tar, sulfur, charcoal, whitfield's ointment, castellani's paint, aluminum chloride, gentian violet, octopirox (piroctone olamine), ciclopirox olamine, undecylenic acid and its metal salts, potassium permanganate, selenium sulfide, sodium thiosulfate, propylene glycol, oil of bitter orange, urea preparations, griseofulvin, 8-hydroxyquinoline ciloquinol, thiobendazole, thiocarbamates, haloprogin, polyenes, hydroxypyridone, morpholine, benzylamine, allylamines (such as terbinafine), tea tree oil, clove leaf oil, coriander, palmarosa, berberine, thyme red, cinnamon oil, cinnamic aldehyde, citronellic acid, hinokitol, ichthyol pale, Sensiva SC-50, Elestab HP-100, azelaic acid, lyticase, iodopropynyl butylcarbamate (IPBC), isothiazalinones such as octyl isothiazalinone, and azoles, and mixtures thereof. In an embodiment, the anti-microbial is selected from the group consisting of: itraconazole, ketoconazole, selenium sulfide, coal tar, and mixtures thereof.

In an embodiment, the azole anti-microbials is an imidazole selected from the group consisting of: benzimidazole, benzothiazole, bifonazole, butaconazole nitrate, climbazole, clotrimazole, croconazole, eberconazole, econazole, elubiol, fenticonazole, fluconazole, flutimazole, isoconazole, ketoconazole, lanoconazole, metronidazole, miconazole, neticonazole, omoconazole, oxiconazole nitrate, sertaconazole, sulconazole nitrate, tioconazole, thiazole, and mixtures thereof, or the azole anti-microbials is a triazole selected from the group consisting of: terconazole, itraconazole, and mixtures thereof. When present in the composition, the azole anti-microbial active is included in an amount of from about 0.01% to about 5%, or from about 0.1% to about 3%, or from about 0.3% to about 2%, by total weight of the composition. In an embodiment, the azole anti-microbial active is ketoconazole. In an embodiment, the sole anti-microbial active is ketoconazole.

The present invention may also comprise a combination of anti-microbial actives. In an embodiment, the combination of anti-microbial active is selected from the group of combinations consisting of: octopirox and zinc pyrithione, pine tar and sulfur, salicylic acid and zinc pyrithione, salicylic acid and elubiol, zinc pyrithione and elubiol, zinc pyrithione and climbazole, octopirox and climbazole, salicylic acid and octopirox, and mixtures thereof.

In an embodiment, the composition comprises an effective amount of a zinc-containing layered material. In an embodiment, the composition comprises from about 0.001% to about 10%,

or from about 0.01% to about 7%, or from about 0.1% to about 5% of a zinc-containing layered material, by total weight of the composition.

Zinc-containing layered materials may be those with crystal growth primarily occurring in two dimensions. It is conventional to describe layer structures as not only those in which all the atoms are incorporated in well-defined layers, but also those in which there are ions or molecules between the layers, called gallery ions (A.F. Wells “Structural Inorganic Chemistry” Clarendon Press, 1975). Zinc-containing layered materials (ZLMs) may have zinc incorporated in the layers and/or be components of the gallery ions. The following classes of ZLMs represent relatively common examples of the general category and are not intended to be limiting as to the broader scope of materials which fit this definition.

Many ZLMs occur naturally as minerals. In an embodiment, the ZLM is selected from the group consisting of: hydrozincite (zinc carbonate hydroxide), basic zinc carbonate, aurichalcite (zinc copper carbonate hydroxide), rosasite (copper zinc carbonate hydroxide), and mixtures thereof. Related minerals that are zinc-containing may also be included in the composition. Natural ZLMs can also occur wherein anionic layer species such as clay-type minerals (e.g., phyllosilicates) contain ion-exchanged zinc gallery ions. All of these natural materials can also be obtained synthetically or formed *in situ* in a composition or during a production process.

Another common class of ZLMs, which are often, but not always, synthetic, is layered double hydroxides. In an embodiment, the ZLM is a layered double hydroxide conforming to the formula $[M^{2+}_{1-x}M^{3+}_x(OH)_2]^{x+} A^{m-}_{x/m} \cdot nH_2O$ wherein some or all of the divalent ions (M^{2+}) are zinc ions (Crepaldi, EL, Pava, PC, Tronto, J, Valim, JB *J. Colloid Interfac. Sci.* 2002, 248, 429-42).

Yet another class of ZLMs can be prepared called hydroxy double salts (Morioka, H., Tagaya, H., Karasu, M, Kadokawa, J, Chiba, K *Inorg. Chem.* 1999, 38, 4211-6). In an embodiment, the ZLM is a hydroxy double salt conforming to the formula $[M^{2+}_{1-x}M^{2+}_{1+x}(OH)_{3(1-y)}]^{+} A^{n-}_{(1=3y)/n} \cdot nH_2O$ where the two metal ions (M^{2+}) may be the same or different. If they are the same and represented by zinc, the formula simplifies to $[Zn_{1+x}(OH)_2]^{2x+} 2x A^{-} \cdot nH_2O$. This latter formula represents (where $x=0.4$) materials such as zinc hydroxychloride and zinc hydroxynitrate. In an embodiment, the ZLM is zinc hydroxychloride and/or zinc hydroxynitrate. These are related to hydrozincite as well wherein a divalent anion replace the monovalent anion. These materials can also be formed *in situ* in a composition or in or during a production process.

In an embodiment, the composition comprises basic zinc carbonate. Commercially available sources of basic zinc carbonate include Zinc Carbonate Basic (Cater Chemicals: Bensenville, IL, USA), Zinc Carbonate (Shepherd Chemicals: Norwood, OH, USA), Zinc

Carbonate (CPS Union Corp.: New York, NY, USA), Zinc Carbonate (Elementis Pigments: Durham, UK), and Zinc Carbonate AC (Bruggemann Chemical: Newtown Square, PA, USA). Basic zinc carbonate, which also may be referred to commercially as "Zinc Carbonate" or "Zinc Carbonate Basic" or "Zinc Hydroxy Carbonate", is a synthetic version consisting of materials
5 similar to naturally occurring hydrozincite. The idealized stoichiometry is represented by $Zn_5(OH)_6(CO_3)_2$ but the actual stoichiometric ratios can vary slightly and other impurities may be incorporated in the crystal lattice.

In embodiments having a zinc-containing layered material and a pyrrithione or polyvalent metal salt of pyrrithione, the ratio of zinc-containing layered material to pyrrithione or a polyvalent
10 metal salt of pyrrithione is from about 5:100 to about 10:1, or from about 2:10 to about 5:1, or from about 1:2 to about 3:1.

Hair Care Compositions

15 Exemplary hair care rinse-off hair care compositions can include an aqueous carrier, which can be present at a level of from about 5% to about 95%, or from about 60% to about 85%. The aqueous carrier may comprise water, or a miscible mixture of water and organic solvent. Non-aqueous carrier materials can also be employed.

In an embodiment of the present invention, a surfactant may be present in the range of about
20 0.1% to about 40%, in a further embodiment, from about 0.5% to about 30%, in a further embodiment, from about 1% to about 25%.

Such rinse-off hair care compositions can include one or more deterative surfactants. The deterative surfactant component can be included to provide cleaning performance to the product. The deterative surfactant component in turn comprises anionic deterative surfactant, zwitterionic or
25 amphoteric deterative surfactant, or a combination thereof. A representative, non-limiting, list of anionic surfactants includes anionic deterative surfactants for use in the compositions can include ammonium lauryl sulfate, ammonium laureth sulfate, triethylamine lauryl sulfate, triethylamine laureth sulfate, triethanolamine lauryl sulfate, triethanolamine laureth sulfate, monoethanolamine lauryl sulfate, monoethanolamine laureth sulfate, diethanolamine lauryl sulfate, diethanolamine
30 laureth sulfate, lauric monoglyceride sodium sulfate, sodium lauryl sulfate, sodium laureth sulfate, potassium lauryl sulfate, potassium laureth sulfate, sodium lauryl sarcosinate, sodium lauroyl sarcosinate, lauryl sarcosine, cocoyl sarcosine, ammonium cocoyl sulfate, ammonium lauroyl sulfate, sodium cocoyl sulfate, sodium lauroyl sulfate, potassium cocoyl sulfate, potassium lauryl

sulfate, triethanolamine lauryl sulfate, triethanolamine lauryl sulfate, monoethanolamine cocoyl sulfate, monoethanolamine lauryl sulfate, sodium tridecyl benzene sulfonate, sodium dodecyl benzene sulfonate, sodium cocoyl isethionate and combinations thereof. In one example, the anionic surfactant can be sodium lauryl sulfate or sodium laureth sulfate. The concentration of the anionic surfactant component in the product can be sufficient to provide a desired cleaning and/or lather performance, and generally ranges from about 2% to about 40%.

Amphoteric deterative surfactants suitable for use in the rinse-off hair care compositions are well known in the art, and include those surfactants broadly described as derivatives of aliphatic secondary and tertiary amines in which an aliphatic radical can be straight or branched chain and wherein an aliphatic substituent can contain from about 8 to about 18 carbon atoms such that one carbon atom can contain an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples of compounds falling within this definition can be sodium 3-dodecyl-aminopropionate, sodium 3-dodecylaminopropane sulfonate, sodium lauryl sarcosinate, N-alkyltaurines such as the one prepared by reacting dodecylamine with sodium isethionate according to the teaching of U.S. Patent No. 2,658,072, N-higher alkyl aspartic acids such as those produced according to the teaching of U.S. Patent No. 2,438,091, and products described in U.S. Patent No. 2,528,378. Other examples of amphoteric surfactants can include sodium lauroamphoacetate, sodium cocoamphoacetate, disodium lauroamphoacetate disodium cocodiamphoacetate, and mixtures thereof. Amphoacetates and diamphoacetates can also be used.

Zwitterionic deterative surfactants suitable for use in the rinse-off hair care compositions are well known in the art, and include those surfactants broadly described as derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which aliphatic radicals can be straight or branched chains, and wherein an aliphatic substituent can contain from about 8 to about 18 carbon atoms such that one carbon atom can contain an anionic group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Other zwitterionic surfactants can include betaines, including cocoamidopropyl betaine.

In a further embodiment of the present invention, the hair care composition may comprise an cationic surfactant.

The liquid rinse off hair care composition can comprise one or more phases. Such hair care compositions can include a cleansing phase and/or a benefit phase (i.e., a single- or multi-phase composition). Each of a cleansing phase or a benefit phase can include various components. The cleansing phase and the benefit phase can be blended, separate, or a combination thereof. The cleansing phase and the benefit phase can also be patterned (e.g. striped).

The cleansing phase of a hair care composition can include at least one surfactant. The cleansing phase can be an aqueous structured surfactant phase and constitute from about 5% to about 20%, by weight of the hair care composition. Such a structured surfactant phase can include sodium trideceth(n) sulfate, hereinafter STnS, wherein n can define average moles of ethoxylation. n can range, for example, from about 0 to about 3; from about 0.5 to about 2.7, from about 1.1 to about 2.5, from about 1.8 to about 2.2, or n can be about 2. When n can be less than 3, STnS can provide improved stability, improved compatibility of benefit agents within the hair care compositions, and increased mildness of the compositions as disclosed in U.S. Pre-Grant Publication No. 2010/009285 A1.

The cleansing phase can also comprise at least one of an amphoteric surfactant and a zwitterionic surfactant. Suitable amphoteric or zwitterionic surfactants (in addition to those cited herein) can include, for example, those described in U.S. Patent No. 5,104,646 and U.S. Patent No. 5,106,609.

A cleansing phase can comprise a structuring system. A structuring system can comprise, optionally, a non-ionic emulsifier, optionally, from about 0.05% to about 5%, by weight of the hair care composition, of an associative polymer; and an electrolyte.

The hair care composition can optionally be free of sodium lauryl sulfate, hereinafter SLS, and can comprise at least a 70% lamellar structure. However, the cleansing phase could comprise at least one surfactant, wherein the at least one surfactant includes SLS. Suitable examples of SLS are described in U.S. Pre-Grant Publication No. 2010/0322878 A1.

Rinse-off hair care compositions can also include a benefit phase. The benefit phase can be hydrophobic and/or anhydrous. The benefit phase can also be substantially free of surfactant. A benefit phase can also include a benefit agent. In particular, a benefit phase can comprise from about 0.1% to about 50% benefit agent by weight of the hair care composition. The benefit phase can alternatively comprise less benefit agent, for example, from about 0.5% to about 20% benefit agent, by weight of the hair care composition. Examples of suitable benefit agents can include petrolatum, glyceryl monooleate, mineral oil, natural oils, and mixtures thereof. Additional examples of benefit agents can include water insoluble or hydrophobic benefit agents. Other suitable benefit agents are described in U.S. Pre-Grant Publication No. 2012/0009285 A1.

Non-limiting examples of glycerides suitable for use as hydrophobic hair benefit agents herein can include castor oil, safflower oil, corn oil, walnut oil, peanut oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, vegetable oils, sunflower seed oil, soybean oil,

vegetable oil derivatives, coconut oil and derivatized coconut oil, cottonseed oil and derivatized cottonseed oil, jojoba oil, cocoa butter, and combinations thereof.

Non-limiting examples of alkyl esters suitable for use as hydrophobic hair benefit agents herein can include isopropyl esters of fatty acids and long chain esters of long chain (i.e. C10-C24) fatty acids, e.g., cetyl ricinoleate, non-limiting examples of which can include isopropyl palmitate, isopropyl myristate, cetyl ricinoleate, and stearyl ricinoleate. Other example can include hexyl laurate, isohexyl laurate, myristyl myristate, isohexyl palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, acyl isononanoate lauryl lactate, myristyl lactate, cetyl lactate, and combinations thereof.

Non-limiting examples of polyglycerin fatty acid esters suitable for use as hydrophobic hair benefit agents herein can include decaglyceryl distearate, decaglyceryl diisostearate, decaglyceryl monomyriate, decaglyceryl monolaurate, hexaglyceryl monooleate, and combinations thereof.

RINSE-OFF CONDITIONER COMPOSITION

The conditioner composition described herein comprises a sum of total 0.0001% to about 2% of alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal) and one or more malodor reduction materials having a Sulfur MORV >3; b) from about 0.01% to about 10% of selenium sulfide; and from about 0.1% to about 10% of a cationic surfactant or a mixture of cationic surfactants and an aqueous carrier. The conditioner composition may also comprise a conditioner gel matrix comprising part or all of the cationic surfactant, whereas the conditioner gel network may also comprise one or more high melting point fatty compounds (i.e. fatty alcohols), and an second aqueous carrier.

The conditioner gel matrix of the conditioner composition includes a cationic surfactant or a cationic surfactant system. The cationic surfactant system can be selected from: mono-long alkyl quaternized ammonium salt; a combination of mono-long alkyl quaternized ammonium salt and di-long alkyl quaternized ammonium salt; mono-long alkyl amidoamine salt; a combination of mono-long alkyl amidoamine salt and di-long alkyl quaternized ammonium salt, a combination of mono-long alkyl amidoamine salt and mono-long alkyl quaternized ammonium salt. The cationic surfactant system can be included in the composition at a level by weight of from about 0.1% to about 10%, from about 0.5% to about 8%, from about 0.8 % to about 5%, and from about 1.0% to about 4%.

The conditioner gel matrix of the conditioner composition includes one or more high melting point fatty compounds. Suitable fatty alcohols include, for example, cetyl alcohol,

stearyl alcohol, behenyl alcohol, and mixtures thereof. The high melting point fatty compound can be included in the conditioner composition at a level of from about 0.1% to about 20%, alternatively from about 1% to about 15%, and alternatively from about 1.5% to about 8% by weight of the composition. The conditioner gel matrix of the conditioner composition includes a second aqueous carrier. The second aqueous carrier may comprise water, or a miscible mixture of water and organic solvent.

LEAVE-ON TREATMENT COMPOSITION

The leave-on treatment composition described herein comprises a sum of total 0.0001% to about 2% of alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal) and one or more malodor reduction materials having a Sulfur MORV >3; b) from about 0.01% to about 10% of selenium sulfide; and from about 0.1% to about 10% of a cationic surfactant or a mixture of cationic surfactants and an aqueous carrier. The leave-on treatment may also comprise one or more rheology modifiers and a third aqueous carrier.

In one embodiment the leave-on treatment may include a conditioner gel matrix as described above (in the rinse-off conditioner description).

In another embodiment the leave-on treatment may include one or more rheology modifiers. Any suitable rheology modifier can be used. In an embodiment, the leave-on treatment may comprise from about 0.01% to about 3% of a rheology modifier, alternatively from about 0.1% to about 1% of a rheology modifier,

ADDITIONAL COMPONENTS

The conditioner compositions, and/or leave-on treatments described herein may optionally comprise one or more additional components known for use in hair care or personal care products,

Non-limiting examples of additional components for use in the hair care compositions include conditioning agents (silicone or non-silicone conditioning agents), natural cationic deposition polymers, synthetic cationic deposition polymers, anti-dandruff agents, particles, suspending agents, paraffinic hydrocarbons, propellants, viscosity modifiers, dyes, non-volatile solvents or diluents (water-soluble and water-insoluble), pearlescent aids, foam boosters, additional surfactants or nonionic cosurfactants, pediculocides, pH adjusting agents, perfumes, preservatives, proteins, skin active agents, sunscreens, UV absorbers, and vitamins.

The rinse-off hair care composition can be applied by a variety of means, including by rubbing, wiping or dabbing with hands or fingers, or by means of an implement and/or delivery

enhancement device. Non-limiting examples of implements include a sponge or sponge-tipped applicator, a mesh shower puff, a swab, a brush, a wipe (e.g., wash cloth), a loofah, and combinations thereof. Non-limiting examples of delivery enhancement devices include mechanical, electrical, ultrasonic and/or other energy devices. Employment of an implement or device can help delivery of the particulate antimicrobial agent to target regions, such as, for example, hair follicles and undulations that can exist in the underarm. The rinse-off care product can be sold together with such an implement or device. Alternatively, an implement or device can be sold separately but contain indicium to indicate usage with a rinse-off care product. Implements and delivery devices can employ replaceable portions (e.g., the skin interaction portions), which can be sold separately or sold together with the rinse-off care product in a kit.

TEST METHODS

Malodor reduction materials may be separated from mixtures, including but not limited to finished products such as consumer products and indentified, by analytical methods that include GC-MS and/or NMR.

Test Method for Determining Saturation Vapour Pressure (VP)

The saturation Vapour Pressure (VP) values are computed for each PRM in the perfume mixture being tested. The VP of an individual PRM is calculated using the VP Computational Model, version 14.02 (Linux) available from Advanced Chemistry Development Inc. (ACD/Labs) (Toronto, Canada) to provide the VP value at 25 °C expressed in units of torr. The ACD/Labs' Vapor Pressure model is part of the ACD/Labs model suite.

Test Method for Determining the Logarithm of the Octanol/Water Partition Coefficient (logP)

The value of the log of the Octanol/Water Partition Coefficient (logP) is computed for each PRM in the perfume mixture being tested. The logP of an individual PRM is calculated using the Consensus logP Computational Model, version 14.02 (Linux) available from Advanced Chemistry Development Inc. (ACD/Labs) (Toronto, Canada) to provide the unitless logP value. The ACD/Labs' Consensus logP Computational Model is part of the ACD/Labs model suite.

Test Method for the Generation of Molecular Descriptors

In order to conduct the calculations involved in the computed-value test methods described herein, the starting information required includes the identity, weight percent, and molar percent of each PRM in the perfume being tested, as a proportion of that perfume, wherein all PRMs in the perfume composition are included in the calculations. Additionally for each of those PRMs, the molecular structure, and the values of various computationally-derived molecular descriptors are also required, as determined in accordance with the Test Method for the Generation of Molecular Descriptors described herein.

For each PRM in a perfume mixture or composition, its molecular structure is used to compute various molecular descriptors. The molecular structure is determined by the graphic molecular structure representations provided by the Chemical Abstract Service ("CAS"), a division of the American Chemical Society, Columbus, Ohio, U.S.A.. These molecular structures may be obtained from the CAS Chemical Registry System database by looking up the index name or CAS number of each PRM. For PRMs, which at the time of their testing are not yet listed in the CAS Chemical Registry System database, other databases or information sources may be used to determine their structures. For a PRM which has potentially more than one isomer present, the molecular descriptor computations are conducted using the molecular structure of only one of the isomers, which is selected to represent that PRM. The selection of isomer is determined by the relative amount of extension in the molecular structures of the isomers. Of all the isomers of a given PRM, it is the isomer whose molecular structure that is the most prevalent which is the one that is selected to represent that PRM. The structures for other potential isomers of that PRM are excluded from the computations. The molecular structure of the isomer that is the most prevalent is paired with the concentration of that PRM, where the concentration reflects the presence of all the isomers of that PRM that are present.

A molecule editor or molecular sketching software program, such as ChemDraw (CambridgeSoft / PerkinElmer Inc., Waltham, Massachusetts, U.S.A.), is used to duplicate the 2-dimensional molecular structure representing each PRM. Molecular structures should be represented as neutral species (quaternary nitrogen atoms are allowed) with no disconnected fragments (e.g., single structures with no counter ions). The winMolconn program described below can convert any deprotonated functional groups to the neutral form by adding the appropriate number of hydrogen atoms and will discard the counter ion.

For each PRM, the molecular sketching software is used to generate a file which describes the molecular structure of the PRM. The file(s) describing the molecular structures of the PRMs

is subsequently submitted to the computer software program winMolconn, version 1.0.1.3 (Hall Associates Consulting, Quincy, Massachusetts, U.S.A., www.molconn.com), in order to derive various molecular descriptors for each PRM. As such, it is the winMolconn software program which dictates the structure notations and file formats that are acceptable options. These options include either a MACCS SDF formatted file (i.e., a Structure-Data File); or a Simplified Molecular Input Line Entry Specification (i.e., a SMILES string structure line notation) which is commonly used within a simple text file, often with a ".smi" or ".txt" file name extension. The SDF file represents each molecular structure in the format of a multi-line record, while the syntax for a SMILES structure is a single line of text with no white space. A structure name or identifier can be added to the SMILES string by including it on the same line following the SMILES string and separated by a space, e.g.: C1=CC=CC=C1 benzene.

The winMolconn software program is used to generate numerous molecular descriptors for each PRM, which are then output in a table format. Specific molecular descriptors derived by winMolconn are subsequently used as inputs (i.e., as variable terms in mathematical equations) for a variety of computer model test methods in order to calculate values such as: saturation Vapour Pressure (VP); Boiling Point (BP); logarithm of the Octanol/Water Partition Coefficient (logP); Odour Detection Threshold (ODT); Malodour Reduction Value (MORV); and/or Universal Malodour Reduction Value (Universal MORV) for each PRM. The molecular descriptor labels used in the models' test method computations are the same labels reported by the winMolconn program, and their descriptions and definitions can be found listed in the winMolconn documentation. The following is a generic description of how to execute the winMolconn software program and generate the required molecular structure descriptors for each PRM in a composition.

Computing Molecular Structure Descriptors using winMolconn:

- 1) Assemble the molecular structure for one or more perfume ingredients in the form of a MACCS Structure-Data File, also called an SDF file, or as a SMILES file.
- 2) Using version 1.0.1.3 of the winMolconn program, running on an appropriate computer, compute the full complement of molecular descriptors that are available from the program, using the SDF or SMILES file described above as input.
 - a. The output of winMolconn is in the form of an ASCII text file, typically space delimited, containing the structure identifiers in the first column and respective molecular descriptors in the remaining columns for each structure in the input file.

3) Parse the text file into columns using a spreadsheet software program or some other appropriate technique. The molecular descriptor labels are found on the first row of the resulting table.

4) Find and extract the descriptor columns, identified by the molecular descriptor label, corresponding to the inputs required for each model.

a. Note that the winMolconn molecular descriptor labels are case-sensitive.

MORV calculation

1.) Input Molecular Descriptor values as determined via the method above into the following equation:

$$\text{MORV} = -0.0035 + 0.8028 \times (\text{SHC}_{\text{sat}}) + 2.1673 \times (\text{xvp7}) - 1.3507 \times (\text{c1C1C3d}) + 0.61496 \times (\text{c1C1O2}) + 0.00403 \times (\text{idc}) - 0.23286 \times (\text{nd2}).$$

This equation relates a material's effectiveness in reducing the malodor 3-mercapto-3-methylhexan-1-ol (thiol based malodors)

2.) For purpose of the present application, a material's MORV is the highest MORV value from the above equation.

The purpose of this experimental design is to determine whether malodor reducing compositions show benefit in reducing the perception of malodor from selenium sulfide-containing shampoos.

Procedure:

Six malodor reduction accords are dosed into unperfumed selenium sulfide shampoo at 4 different levels; samples are allowed to equilibrate at room temperature for at least 1 week before evaluating.

#1 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
13877-91-3	(3Z)-3,7-dimethylocta-1,3,6-triene	3.000

5413-60-5	(3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate t	10.000
33885-52-8	alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene- propanal)	1.500
23787-90-8	1,3,4,6,7,8alpha-hexahydro- 1,1,5,5-tetramethyl-2H- 2,4alpha-methanophthalen- 8(5H)-one	14.000
4430-31-3	3,4,4a,5,6,7,8,8a- octahydrochromen-2-one	10.000
300371-33-9	2,3-dihydro-1,1-1H-dimethyl- indene-ar-propanal	1.500
55066-48-3	3-methyl-5-phenylpentan-1-ol	10.000
139504-68-0	1-((2-(tert- butyl)cyclohexyl)oxy)butan-2- ol	10.000
3681-73-0	(E)-3,7-dimethylocta-2,6-dien- 1-yl palmitate)	40.000

#2 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
58430-94-7	3,5,5-trimethylhexyl acetate	56
33885-52-8	alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene- propanal)	18
1139-30-6	4,5-epoxy-4,11,11-trimethyl-8- methylenebicyclo(7.2.0)undecane)	0.5
27606-09-3	2,4-dimethyl-4,4a,5,9b- tetrahydroindeno[1,2- d][1,3]dioxine	15

127-42-4	(E)-1-(2,6,6-trimethyl-1-cyclohex-2-enyl)pent-1-en-3-one	2
23787-90-8	(1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one	5
68912-13-0	8,8-dimethyl-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl propionate	3.5

#3 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
5413-60-5	(3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate	15.000
23787-90-8	1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one	5.000
300371-33-9	2,3-dihydro-1,1-1H-dimethyl-indene-ar-propanal	1.000
55066-48-3	3-methyl-5-phenylpentan-1-ol	49.000
139504-68-0	1-((2-(tert-butyl)cyclohexyl)oxy)butan-2-ol	10.000
3681-73-0	(E)-3,7-dimethylocta-2,6-dien-1-yl palmitate)	20.000

5

#4 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
-----	----------	-----

23787-90-8	1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one	40.000
300371-33-9	2,3-dihydro-1,1-1H-dimethyl-indene-ar-propana 1	10.000
55066-48-3	3-methyl-5-phenylpentan-1-ol	10.000
3681-73-0	(E)-3,7-dimethylocta-2,6-dien-1-yl palmitate)	40.000

#5 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
58430-94-7	3,5,5-trimethylhexyl acetate	2.05
98-52-2	4-tert-butyl cyclohexane	0.25
32210-23-4	4-(tert-butyl)cyclohexyl acetate	7.6
30772-79-3	Octahydro-1H-4,7-methanoindene-1-carbaldehyde	0.28
63500-71-0	2-isobutyl-4-hydroxy-4-methyltetrahydropyran 1	9.5
101-84-8	DIPHENYL OXIDE	0.28
19870-74-7	Cedryl Methyl Ether	62.2
23787-90-8	(1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one	9.5
93-16-3	(Z)-1,2-dimethoxy-4-(prop-1-en-1-yl)benzene	4.46

55066-48-3	3-methyl-5-phenylpentan-1-ol	3.8
24851-98-7	methyl 2-(3-oxo-2-pentylcyclopentyl)acetate	0.08

#6 Malodor Reduction Composition (MRC)

CAS	MATERIAL	Wt%
58430-94-7	3,5,5-trimethylhexyl acetate	40
5413-60-5	(3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl) acetate	20
23787-90-8	1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophthalen-8(5H)-one	20
300371-33-9	2,3-dihydro-1,1-1H-dimethylindene-ar-propanal	10
55066-48-3	3-methyl-5-phenylpentan-1-ol	10

5

Samples to a depth of ~1 cm are placed into plastic cups; cups are covered with lids, then lids are removed for evaluation. Each sample is evaluated by 4 trained odor panelists for the presence of sulfur malodor. The unperfumed shampoo with no Malodor Reduction Composition (MRC) is set as positive control (ie, “2”) while a separate batch of the same shampoo formula, sans selenium sulfide, is set as negative control (ie, “0”). Additionally, a 50:50 blend of the 2 unperfumed shampoos (with and without selenium sulfide) is created and this is set as an additional “low sulfur” benchmark for improved coverage (Ie, “1”). Sample presentation is randomized.

Instructions are provided to the panelists as follows:

- 15 1) Please smell sample labeled “Control”. Note the amount of sulfur base odor. This is your benchmark for subsequent evaluations. You can also refer to “Low Sulfur Control” and “No Sulfur Control” as needed, however, the scale is relative to “Control”.

2) Please evaluate each sample by removing the lid, and indicate the amount of sulfur relative to the control that you can detect in each sample. You may or may not be able to smell perfume materials in the test samples; please assess only for amount of sulfur odor using the scale below:

5

0 = I cannot smell any sulfur odor at all (you can refer to "No Sulfur Control")

1 = I smell some sulfur odor, but less than the "control" (You can refer to "Low Sulfur Control")

2 = I smell sulfur odor equal to the control

10

3 = I smell more sulfur odor than in the control

Panelists evaluate each level series as a set, taking a break in between sets to avoid olfactive saturation.

Results

Round 1 - 0.01% Malodor

Reduction Composition (MRC)

		Panelist	Panelist	Panelist	
	Panelist 1	2	3	4	Avg
MRC 1	1	0	1	0.5	0.625
MRC 2	1	0	0	0.5	0.375
MRC 3	1	1	2	1	1.25
MRC4	1	2	1	2	1.5
MRC 5	2	1.5	1	2	1.625
MRC 6	1	1	1	1	1

Round 2 - 0.05% Malodor

Reduction composition (MRC)

MRC 1	0	1	2	0	0.75
MRC 2	0	1	0.5	0	0.375
MRC 3	1	1.5	1	1	1.125
MRC4	0	2	1	1	1
MRC 5	0	2	2	2	1.5

MRC 6	0	2	2	1	1.25
-------	---	---	---	---	------

Round 3 - 0.1% Malodor

Reduction composition (MRC)

MRC 1	1	0	1	1	0.75
MRC 2	0	0	0	0	0
MRC 3	1	0	0	1	0.5
MRC4	0	1	1	1	0.75
MRC 5	1	0.5	2	2	1.375
MRC 6	0	1	2	1	1

Round 4 - 0.15% Malodor

Reduction composition (MRC)

MRC 1	1	0	1.5	1	0.875
MRC 2	0	1	0.5	0	0.375
MRC 3	0	1	2	1	1
MRC4	0	1	1	1	0.75
MRC 5	0	2	2	1	1.25
MRC 6	1	1	1	1	1

Summary

MRC 1, MRC 2, MRC 3 and MRC 4 all show considerable reduction (< 1 Avg) in perceived sulfur levels at one or more levels tested. Additionally, MRC 2 showed reduction at all levels tested, with all panelists assigning a zero or “no sulfur” rating at the 0.1% added level.

As a follow up, the MRC 2 is diluted in 100 deg F tap water at a ratio of 1 ml shampoo to 75 ml water; these samples are placed into cups, capped, and agitated until the shampoo is dissolved. They are immediately evaluated by 2 panelists to determine if sulfur odor is present in the dilute samples.

	Panelist 1	Panelist 4
--	------------	------------

MRC 2 @ 0.01%, diluted 1:75 in		
--------------------------------	--	--

warm water (100 F)	1	1.5
--------------------	---	-----

MRC 2 @ 0.05%, diluted 1:75 in warm water (100 F)	0.5	0.5
MRC 2 @ 0.10%, diluted 1:75 in warm water (100 F)	0	0
MRC 2 @ 0.15%, diluted 1:75 in warm water (100 F)	0	0

In dilution, the perceived benefit holds, particularly at the 0.1% and 0.15% levels.

Conclusion

- 5 Malodor Reduction Composition's demonstrates effectiveness in the masking/blocking of selenium sulfide odor. In the present invention, this understanding can be leveraged in formulation strategies for selenium sulfide containing hair care compositions.

I. Hair Switch Selection and Preparation – For All Tests

- 10 a) For each product, a 20 gram hair switch is used, cleaned and stripped of perfume. Cleaning may be done via external vendor (ie, ATL 2x baking soda/main mix) or internally via local documented procedures. (New hair switches can be pre-washed with unperfumed shampoo before very first use)
- b) Each hair switch is evaluated before each use to ensure no contamination/off odors.

15 II. Shampoo Fragrance Profile

- a) A neat product is evaluated; and a neat sulfur score is recorded-
- b) The water temperature may be adjusted to 100F/38C and 1.5 gal/5.7 l per min; and hair is wetted for **20** seconds.
- c) 2 ml product is applied; and lather is produced for **30** seconds.
- 20 d) Odor is evaluated using sulfur scoring and comments are recorded.

Table 4

Evaluate MRC2 On Switch vs Negative Control
--

In-Use Evaluation 20 gram hair switch used, 2 ml product; 100 F water according to the method described above.

Product	Sulfur Odor Scale (0-3)	Neat Comments	Lather Sulfur Scale (0-3)	Lather Comments
Selenium Sulfide Unperfumed Shampoo	2	sulfur odor	2	sulfur odor
Non Malodor Reducing Accord ¹ @ 0.1% in Selenium Sulfide Shampoo	1	sulfur odor + rose powdery note	1	sulfur odor + rose powdery note
MRC2 @ 0.1% in Selenium Sulfide Shampoo	0	ozonic, floral	0	ozonic. Floral

1. Non-Malodor Reducing
Accord

Material	%	CAS
Citronellol	20	106-22-920
Linalool	30	78-70-6
Phenyl Ethyl Alcohol	50	60-12-8

Sulfur Odor Scale:

0 = I cannot smell any sulfur odor at all (you can refer to “No Sulfur Control”)

1 = I smell some sulfur odor, but less than the “control” (You can refer to “Low Sulfur Control”)

2 = I smell sulfur odor equal to the control

3 = I smell more sulfur odor than in the control

Table 5

Evaluate MRC 2 On Switch vs non malodor reducing accord-0.01% Levels

In-Use Evaluation 20 gram hair switch used, 2 ml product; 100 F water according to the method described above.

Product	Neat Sulfur Odor Scale (0-3)	Neat Comments	Lather Sulfur Scale (0-3)	Lather Comments
Selenium Sulfide Unperfumed Shampoo	2	sulfur odor	2	sulfur odor
Non- malodor reducing accord ¹ @ 0.01% in Selenium Sulfide Shampoo (ref batch	1.5	slight rosy, noticeable sulfur	2	sulfur odor
MBRC 2 @ 0.01% in Selenium Sulfide Shampoo	0.5	slight ozonic, very slight sulfur	0.5	Slight ozonic, very slight sulfur

1. Non-Malodor Reducing Accord

Material	%	CAS
Citronellol	20	106-22-920
Linalool	30	78-70-6
Phenyl Ethyl Alcohol	50	60-12-8

Sulfur Odor Scale:

0 = I cannot smell any sulfur odor at all (you can refer to "No Sulfur Control")

1 = I smell some sulfur odor, but less than the "control" (You can refer to "Low Sulfur Control")

2 = I smell sulfur odor equal to the control
 3 = I smell more sulfur odor than in the
 control

Conclusion

This series of experiments indicates that malodor reducing ingredients or a combination of those, such as MRC2, provides a malodor blocking benefit both in neat product and during the lather phase. This benefit goes beyond the masking benefit that would be provided by a non-malodor reducing perfume. In table 4 (0.1% data), the non-malodor reducing accord exhibits malodor interference via masking benefit due to the perfume intensity imparted by the ingredients. In table 5 (the 0.01% data) it is demonstrated that as level is reduced the masking benefit of the non-malodor-reducing accord is decreased, while the malodor blocking property of the MRC2 is maintained.

Examples Shampoo with Malodor reducing composition

An example of Shampoo compositions prepared with malodor reduction composition, according to the compositions shown in Example 1.

	1	2	3
Ingredient	Wt %		
Ammonium Laureth Sulfate ¹	14.1	14.1	14.1
Ammonium Lauryl Sulfate ²	3.1	3.1	3.1
Ammonium Xylenesulfonate ³	0.45	0.45	0.45
TWEEN 60 ⁴	3.0	3.0	3.0
Polyquaternium-10 ⁵	0.35	0.35	0.35
Cetrimonium Chloride ⁶	0.5	0.5	0.5
Selenium Sulfide ⁷	1.0	1.0	1.0
Dimethicone ⁸	0.60	0.60	0.60
Ethylene Glycol Distearate ⁹	3.0	3.0	3.0
Cocamide MEA ¹⁰	3.0	3.0	3.0
Zinc Pyrithione ¹¹	--	0.2	0.2
Zinc Carbonate ¹²	--	--	1.61
Neat Fragrance	1.1	0.75	0.75

Malodor reducing composition	0.25	0.25	0.175
Cetyl Alcohol ¹³	0.42	0.42	0.42
DMDM Hydantoin	0.40	0.40	0.40
Sodium Chloride	0.30	0.30	0.30
Stearyl Alcohol ¹⁴	0.20	0.20	0.20
Hydroxypropyl Methylcellulose ¹⁵	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

- (1) Ammonium Laureth Sulfate at 25% active, supplier: P&G
- (2) Ammonium Lauryl Sulfate at 25% active, supplier: P&G
- (3) Ammonium Xylene Sulfonate 40% active, supplier: Stepan
- 5 (4) Polysorbate 60, upplier: Croda
- (5) UCARE Polymer LR400, supplier- Dow Chemical
- (6) cetrimonium chloride, supplier - Croda
- (7) Selenium disulfide, supplier Eskay
- (8) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt
- 10 (centistokes).
- (9) Ethylene Glycol Distearate, supplier: Stepan
- (10) Ninol COMF from the Stepan Company
- (11) Zinc Pyrithione, supplier Lonza
- (12) Zinc Carbonate Basic, supplier Pan Continental Chemical
- 15 (13) Cetyl Alcohol, supplier P&G
- (14) Stearyl Alcohol, supplier P&G
- (15) Methocel, supplier Dow Chemical

	4	5	6
Ingredient	Wt %		
Ammonium Laureth Sulfate ¹	14.1	14.1	14.1
Ammonium Lauryl Sulfate ²	3.1	3.1	3.1
Ammonium Xylenesulfonate ³	0.45	0.45	0.45
TWEEN 60 ⁴	3.0	3.0	3.0
Polyquaternium-10 ⁵	0.35	0.35	0.35

Cetrimonium Chloride ⁶	0.5	0.5	0.5
Selenium Sulfide ⁷	1.0	0.2	0.2
Dimethicone ⁸	0.60	0.60	0.60
Ethylene Glycol Distearate ⁹	3.0	3.0	3.0
Cocamide MEA ¹⁰	3.0	3.0	3.0
Zinc Pyrithione ¹¹	--	1.0	1.0
Zinc Carbonate ¹²	--	--	1.61
Neat Fragrance	0.65	0.85	1.0
Malodor reducing composition	0.175	0.175	0.175
Cetyl Alcohol ¹³	0.42	0.42	0.42
DMDM Hydantoin	0.40	0.40	0.40
Sodium Chloride	0.30	0.30	0.30
Stearyl Alcohol ¹⁴	0.20	0.20	0.20
Hydroxypropyl Methylcellulose ¹⁵	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

- (1) Ammonium Laureth Sulfate at 25% active, supplier: P&G
- (2) Ammonium Lauryl Sulfate at 25% active, supplier: P&G
- (3) Ammonium Xylene Sulfonate 40% active, supplier: Stepan
- 5 (4) Polysorbate 60, upplier: Croda
- (5) UCARE Polymer LR400, supplier- Dow Chemical
- (6) cetrimonium chloride, supplier - Croda
- (7) Selenium disulfide, supplier Eskay
- (8) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt
- 10 (centistokes).
- (9) Ethylene Glycol Disterate, supplier: Stepan
- (10) Ninol COMF from the Stepan Company
- (11) Zinc Pyrithione, supplier Lonza
- (12) Zinc Carbonate Basic, supplier Pan Continental Chemical
- 15 (13) Cetyl Alcohol, supplier P&G
- (14) Stearyl Alcohol, supplier P&G
- (15) Methocel, supplier Dow Chemical

In an embodiment of the present invention, the example of Shampoo compositions (Example 5) may also be prepared with the malodor reduction composition according to the compositions shown in Example 2 and/or Example 3.

5

	7	8	9
Ingredient	Wt %		
Ammonium Laureth Sulfate ¹	14.1	14.1	14.1
Ammonium Lauryl Sulfate ²	3.1	3.1	3.1
Ammonium Xylenesulfonate ³	0.45	0.45	0.45
TWEEN 60 ⁴	3.0	3.0	3.0
Polyquaternium-10 ⁵	0.35	0.35	0.35
Cetrimonium Chloride ⁶	0.5	0.5	0.5
Sulfur ⁷	2.0	5.0	3.5
Dimethicone ⁸	0.60	0.60	0.60
Ethylene Glycol Distearate ⁹	3.0	3.0	3.0
Cocamide MEA ¹⁰	3.0	3.0	3.0
Selenium Sulfide ¹¹	--	--	--
Neat Fragrance	0.85	1.1	0.95
Malodor reducing composition	0.175	0.25	0.25
Cetyl Alcohol ¹²	0.42	0.42	0.42
DMDM Hydantoin	0.40	0.40	0.40
Sodium Chloride	0.30	0.30	0.30
Stearyl Alcohol ¹³	0.20	0.20	0.20
Hydroxypropyl Methylcellulose ¹⁴	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

(1) Ammonium Laureth Sulfate at 25% active, supplier: P&G

(2) Ammonium Lauryl Sulfate at 25% active, supplier: P&G

(3) Ammonium Xylene Sulfonate 40% active, supplier: Stepan

10 (4) Polysorbate 60, upplier: Croda

(5) UCARE Polymer LR400, supplier- Dow Chemical

- (6) cetrimonium chloride, supplier - Croda
- (7) Sulfur USP, supplier Universal Preserv-A-Chem
- (8) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt (centistokes).
- 5 (9) Ethylene Glycol Distearate, supplier: Stepan
- (10) Ninol COMF from the Stepan Company
- (11) Zinc Pyrithione, supplier Lonza
- (12) Selenium disulfide, supplier Eskay
- (12) Cetyl Alcohol, supplier P&G
- 10 (13) Stearyl Alcohol, supplier P&G
- (14) Methocel, supplier Dow Chemical

	10	11	12
Ingredient	Wt %		
Ammonium Laureth Sulfate ¹	14.1	14.1	14.1
Ammonium Lauryl Sulfate ²	3.1	3.1	3.1
Ammonium Xylenesulfonate ³	0.45	0.45	0.45
TWEEN 60 ⁴	3.0	3.0	3.0
Polyquaternium-10 ⁵	0.35	0.35	0.35
Cetrimonium Chloride ⁶	0.5	0.5	0.5
Sulfur ⁷	2.0	2.0	2.0
Dimethicone ⁸	0.60	0.60	0.60
Ethylene Glycol Distearate ⁹	3.0	3.0	3.0
Cocamide MEA ¹⁰	3.0	3.0	3.0
Selenium Sulfide ¹¹	0.5	0.25	0.1
Neat Fragrance	0.65	0.85	1.0
Malodor reducing composition	0.175	0.175	0.175
Cetyl Alcohol ¹²	0.42	0.42	0.42

DMDM Hydantoin	0.40	0.40	0.40
Sodium Chloride	0.30	0.30	0.30
Stearyl Alcohol ¹³	0.20	0.20	0.20
Hydroxypropyl Methylcellulose ¹⁴	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

- (1) Ammonium Laureth Sulfate at 25% active, supplier: P&G
- (2) Ammonium Lauryl Sulfate at 25% active, supplier: P&G
- (3) Ammonium Xylene Sulfonate 40% active, supplier: Stepan
- 5 (4) Polysorbate 60, upplier: Croda
- (5) UCARE Polymer LR400, supplier- Dow Chemical
- (6) cetrimonium chloride, supplier - Croda
- (7) Sulfur USP, supplier Universal Preserv-A-Chem
- (8) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt
- 10 (centistokes).
- (9) Ethylene Glycol Distearate, supplier: Stepan
- (10) Ninol COMF from the Stepan Company
- (11) Zinc Pyrithione, supplier Lonza
- (12) Selenium disulfide, supplier Eskay
- 15 (12) Cetyl Alcohol, supplier P&G
- (13) Stearyl Alcohol, supplier P&G
- (14) Methocel, supplier Dow Chemical

	13	14	15
Ingredients	Wt%		
Sodium Laureth Sulfate ¹	14.1	11.5	11.5
Sodium Lauryl Sulfate ²	3.1	1.5	1.5
Sodium Xylenesulfonate ³	0.50	3.0	3.0
Cocamidopropyl Betaine ⁴	--	1.3	1.3
Guar Hydrxypropyltrimonium chloride ⁵	--	0.3	0.3
PQ-76 ⁶	--	0.03	--

Acrylamide/trimethylammoniopropylmethacrylamide ⁷	--	--	0.01
Selenium Sulfide ⁸	1.0	1.0	0.2
Sulfur ⁹	--	--	--
Dimethicone ¹⁰	0.60	0.8	2.7
Ethylene Glycol Distearate ¹¹	3.0	1.5	1.5
Cocamide MEA ¹²	3.0	2.0	2.0
Zinc Pyrithione ¹³	--	--	1.0
Zinc Carbonate ¹⁴	--	--	1.61
Fragrance	0.65	0.85	1.0
Malodor reducing composition	0.15	0.175	0.1
Cetyl Alcohol ¹⁵	0.42	0.04	0.04
Stearyl Alcohol ¹⁶	0.20	0.08	0.08
DMDM Hydantoin	0.40	--	--
Methylchloroisothiazolinone and Methylisothiazolinone	0.5	0.005	0.005
Sodium Chloride	0.30	--	--
Hydroxypropyl Methylcellulose ¹⁷	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

- (1) Sodium Laureth Sulfate at 28% active, supplier: P&G
- (2) Sodium Lauryl Sulfate at 29% active, supplier: P&G
- (3) Sodium Xylene Sulfonate 40% active, supplier: Stepan
- 5 (4) Amphosol HCA from the Stepan Company
- (5) Jaguar C500 from Rhodia; mw 500,000 cd 0.8 meq/g
- (6) PQ-76 from Rhodia; mw 100,000 cd 1.6 meq/g
- (7) AM/APTAC mw 1,100,000 cd 1.8 meq/g, supplier ASI Specialty Ingredients
- (8) Selenium disulfide, supplier Eskay
- 10 (9) Sulfur USP, supplier Universal Preserv-A-Chem
- (10) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt (centistokes).
- (11) Ethylene Glycol Distearate, supplier: Stepan
- (12) Ninol COMF from the Stepan Company

- (13) Zinc Pyrithione, supplier Lonza
 (14) Zinc Carbonate Basic, supplier Pan Continental Chemical
 (15) Cetyl Alcohol, supplier P&G
 (16) Stearyl Alcohol, supplier P&G
 5 (17) Methocel, supplier Dow Chemical

	16	17	18
Ingredients	Wt%		
Sodium Laureth Sulfate ¹	11.5	11.5	11.5
Sodium Lauryl Sulfate ²	1.5	1.5	1.5
Sodium Xylenesulfonate ³	3.0	3.0	3.0
Cocamidopropyl Betaine ⁴	1.3	1.3	1.3
Guar Hydroxypropyltrimonium chloride ⁵	0.3	0.3	0.3
PQ-76 ⁶	0.03	0.03	--
Acrylamide/trimethylammoniopropylmethacrylamide ⁷	--	--	0.01
Selenium Sulfide ⁸	1.0	--	0.1
Sulfur ⁹	2.0	5.0	2.0
Dimethicone ¹⁰	0.8	0.8	2.7
Ethylene Glycol Distearate ¹¹	1.5	1.5	1.5
Cocamide MEA ¹²	2.0	2.0	2.0
Zinc Pyrithione ¹³	--	--	--
Zinc Carbonate ¹⁴	--	--	--
Fragrance	0.85	0.85	1.0
Malodor reducing composition	0.175	0.175	0.1
Cetyl Alcohol ¹⁵	0.04	0.04	0.04
Stearyl Alcohol ¹⁶	0.08	0.08	0.08
DMDM Hydantoin	--	--	--
Methylchloroisothiazolinone and Methylisothiazolinone	0.005	0.005	0.005
Sodium Chloride	--	--	--
Hydroxypropyl Methylcellulose ¹⁷	0.02	0.02	0.02
Water	Q.S.	Q.S.	Q.S.

- (1) Sodium Laureth Sulfate at 28% active, supplier: P&G
 (2) Sodium Lauryl Sulfate at 29% active, supplier: P&G
 (3) Sodium Xylene Sulfonate 40% active, supplier: Stepan
 5 (4) Amphosol HCA from the Stepan Company
 (5) Jaguar C500 from Rhodia; mw 500,000 cd 0.8 meq/g
 (6) PQ-76 from Rhodia; mw 100,000 cd 1.6 meq/g
 (7) AM/APTAC mw 1,100,000 cd 1.8 meq/g, supplier ASI Specialty Ingredients
 (8) Selenium disulfide, supplier Eskay
 10 (9) Sulfur USP, supplier Universal Preserv-A-Chem
 (10) Viscasil 330M from Momentive Performance Materials with a viscosity of 330,000 cSt (centistokes).
 (11) Ethylene Glycol Disterate, supplier: Stepan
 (12) Ninol COMF from the Stepan Company
 15 (13) Zinc Pyrithione, supplier Lonza
 (14) Zinc Carbonate Basic, supplier Pan Continental Chemical
 (15) Cetyl Alcohol, supplier P&G
 (16) Stearyl Alcohol, supplier P&G
 (17) Methocel, supplier Dow Chemical

20

Rinse Out Conditioner EXAMPLES

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not
 25 to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention. Where applicable, ingredients are identified by chemical or CTFA name, or otherwise defined below.

Components	Ex.1	Ex.2	Ex.3	Ex.4	Ex.5
Polyquaternium-6 *1	0.075	-	-	-	-
Polyquaternium-6 *2	-	0.075	0.075	0.075	0.075
Zinc pyrithione *3	0.5	0.75	0.75	0.75	0.75
Zinc carbonate *4	1.6	1.6	1.6	-	1.6

Selenium Disulfide *5	1.0	0.5	1.0	1.0	1.0
Sulfur *6	-	-	-	-	-
Stearamidopropyldimethylamine	2.0	-	-	-	-
l-glutamic acid	0.64	-	-	-	-
Behenyl trimethyl ammonium methosulfate	-	1.8	1.8	1.8	1.8
Dicetyl dimethyl ammonium chloride	-	0.52	0.52	0.52	0.52
Cetyl alcohol	2.5	1.1	1.1	1.1	1.1
Stearyl alcohol	4.5	2.75	2.75	2.75	2.75
Polydimethylsiloxane *7	0.6	-	-	-	-
Aminosilicone *8	-	0.75	0.75	0.75	0.75
Preservatives	0.45	0.4	0.4	0.4	0.4
Perfume	0.5	0.5	0.5	0.5	0.5
Malodor Reducing Composition	0.125	0.1	0.25	0.175	--
Deionized Water	q.s. to 100%				
Method of preparation	I	II	I	II	I

*1 Polyquaternium-6: Poly(diallyldimethylammonium chloride) supplied with a tradename Merquat 100 from Lubrizol, having a charge density of about 6.2meq/g, and molecular weight of about 150,000g/mol

5 *2 Polyquaternium-6: Poly(diallyldimethylammonium chloride) supplied with a tradename Merquat 106 from Lubrizol having a charge density of about 6.2meq/g, and molecular weight of about 15,000g/mol

*3 Zinc pyrithione: having a particle size of from about 1 to about 10 microns

*4 Zinc carbonate: having a particle size of from about 1 to about 10 microns

10 *5 Selenium Disulfide, from Eskay

*6 Sulfur, from Universal Preserv-A-Chem

*7 Polydimethylsiloxane: having a viscosity of 10,000cSt

*8 Aminosilicone: Terminal aminosilicone which is available from GE having a viscosity of about 10,000mPa•s, and having following formula:



wherein G is methyl; a is an integer of 1; n is a number from 400 to about 600; R₁ is a monovalent radical conforming to the general formula C_qH_{2q}L, wherein q is an integer of 3 and L is -NH₂.

5

Components	Ex. 6	Ex. 7	Ex. 8
Polyquaternium-6 *1	-	-	-
Polyquaternium-6 *2	0.075	0.075	0.075
Zinc pyrithione *3	0.75	0.75	0.75
Zinc carbonate *4	1.6	1.6	1.6
Selenium Disulfide *5	1.0	0.1	-
Sulfur *6	3.5	2.0	2.0
Stearamidopropyldimethylamine	-	-	-
l-glutamic acid	-	-	-
Behenyl trimethyl ammonium methosulfate	1.8	1.8	1.8
Dicetyl dimethyl ammonium chloride	0.52	0.52	0.52
Cetyl alcohol	1.1	1.1	1.1
Stearyl alcohol	2.75	2.75	2.75
Polydimethylsiloxane *7	-	-	-
Aminosilicone *8	0.75	0.75	0.75
Preservatives	0.4	0.4	0.4
Perfume	0.5	0.5	0.5
Malodor Reducing Composition	0.175	0.25	0.25
Deionized Water	QS to 100%		
Method of preparation	II	II	II

*1 Polyquaternium-6: Poly(diallyldimethylammonium chloride) supplied with a tradename

Merquat 100 from Lubrizol, having a charge density of about 6.2meq/g, and molecular weight of about 150,000g/mol

*2 Polyquaternium-6: Poly(diallyldimethylammonium chloride) supplied with a tradename Merquat 106 from Lubrizol having a charge density of about 6.2meq/g, and molecular weight of about 15,000g/mol

*3 Zinc pyrithione: having a particle size of from about 1 to about 10 microns

*4 Zinc carbonate: having a particle size of from about 1 to about 10 microns

*5 Selenium Disulfide, from Eskay

*6 Sulfur, from Universal Preserv-A-Chem

10 *7 Polydimethylsiloxane: having a viscosity of 10,000cSt

*8 Aminosilicone: Terminal aminosilicone which is available from GE having a viscosity of about 10,000mPa•s, and having following formula:



15 wherein G is methyl; a is an integer of 1; n is a number from 400 to about 600; R₁ is a monovalent radical conforming to the general formula C_qH_{2q}L, wherein q is an integer of 3 and L is -NH₂.

Method of Preparation

20 The conditioning compositions of “Ex. 1” through “Ex.3” and “CEX. i as shown above can be prepared by any conventional method well known in the art. They are suitably made by one of the following Methods I or II as shown above.

Method I

25 Cationic surfactants and high melting point fatty compounds are added to water with agitation, and heated to about 80°C. The mixture is cooled down to about 55°C and gel matrix is formed. Zinc carbonates, and if included, silicones and preservatives, are added to the gel matrix with agitation. Then, zinc pyrithione, and if included, polymers are added with agitation at about 45 °C. Then, if included, other components such as perfumes are added with agitation. Then the composition is cooled down to room temperature.

30 Method II

Cationic surfactants and high melting point fatty compounds are mixed and heated to from about 66 °C to about 85 °C to form an oil phase. Separately, water is heated to from about 20 °C to about 48 °C to form an aqueous phase. In Becomix® direct injection rotor-stator homogenizer, the oil phase is injected and it takes 0.2 second or less for the oils phase to reach to a high shear field

having an energy density of from 1.0×10^5 J/m³ to 1.0×10^7 J/m³ where the aqueous phase is already present. A gel matrix is formed at a temperature of above 50 °C to about 60 °C. Silicones, Perfume, Polymer and Preservative, if included, are added to the gel matrix with agitation at temperature below 55 °C and mixed well. Then, zinc carbonate, if included, followed by zinc pyrithione, are added to the gel matrix with agitation at temperature below 50 °C and mix well. Finally the composition is cooled down to room temperature.

Leave on Treatment FORMULATIONS AND EXAMPLES

The following are non-limiting examples of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention, which would be recognized by one of ordinary skill in the art.

Leave on Treatment Examples

	<u>Example</u>					
	<u>s</u>					
	1	2	3	4	5	6
	Active wt	Active wt	Active wt	Active wt	Active wt	Active wt
	%	%	%	%	%	%
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Alcohol 100% (Ethanol)	50	0	50	0	50	0
Isoproryl Alcohol	0	0	0	0	0	0
Acrylates/C10-30 alkyl acrylate crosspolymer *1	0.35	0.2	0.35	0.2	0.35	0.2
Zinc pyrithione *2	0.1	0.07	-	-	-	-
Selenium Sulfide *3	0.1	0.1	-	-	0.1	0.1
Sulfur *4	-	-	2.0	2.0	2.0	2.0
Panthenol	0.15	0	0.15	0	0.15	0
Niacinamide	2.5	0	2.5	0	2.5	0

Caffeine	0.75	0	0.75	0	0.75	0
Glycerin	0.5	0	0.5	0	0.5	0
Propylene Glycol	0	1	0	1	0	1
Menthol	0	0.5	0	0.5	0	0.5
Tapicoa Starch Polymethylsilsesui oxane *5	0	1	0	1	0	1
Benzyl Alcohol	0.12	0.5	0.12	0.5	0.12	0.5
Methylisothiazolin one *6	0	0.05	0	0.05	0	0.05
PEG-40 Hydrogenated Castor Oil *7	0	0.5	0	0.5	0	0.5
Tetrahydroxypropy l Ethylenediamine *8	0.12	0.14	0.12	0.14	0.12	0.14
Fragrance	0.4	0.2	0.4	0.2	0.4	0.2
Malodor Reducing Composition	0.15	0.1	0.15	0.1	0.15	0.1

*1 as in Carbopol Ultrez 21 available from Lubrizol

*2 as in ZPT from Lonza Personal Care

*3 as in selenium sulfide from Eskay

*4 as in Sulfur, from Universal Preserv-A-Chem

*5 as in Neolone 950 from Rohm and Haas

*6 as in Cremophor RH-40 Surfactant from BASF

*7 as in Neutrol Te from BASF

*8 as in Trolamine from Dow Chemical

In the examples, all concentrations are listed as weight percent, unless otherwise specified and may exclude minor materials such as diluents, filler, and so forth. The listed formulations, therefore, comprise the listed components and any minor materials associated with such components. As is apparent to one of ordinary skill in the art, the selection of these minors will vary depending on the physical and chemical characteristics of the particular ingredients selected to make the hair care composition

Examples/Combinations

- A. A hair care composition comprising, based on total composition weight,
- a sum total of from about 0.0001% to about 2% of alpha, alpha, 6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal and one or more malodor reduction materials having a Sulfur MORV >3;
 - from about 0.01% to about 10% of a scalp active material selected from the group consisting of selenium sulfide, sulfur and mixtures thereof;
 - from about 0.1% to about 40%, of a surfactant.
- B. The composition according to Paragraph A, wherein the malodor reduction materials are selected from the group consisting of 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane], 1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene, 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate, 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate, (3R,3aR,6S,7S,8aS)-6-methoxy-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulene, Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate), (3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan), alpha, alpha, 6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal, 4,5-epoxy-4,11,11-trimethyl-8-methylenebicyclo(7.2.0)undecane), (4aR,8aS)-7-methyloctahydro-1,4-methanonaphthalen-6(2H)-one), (5-methoxyoctahydro-1H-4,7-methanoindene-2-carbaldehyde), (2R,4a'R,8a'R)-3,7'-dimethyl-3',4',4a',5',8',8a'-hexahydro-1'H-spiro[oxirane-2,2'-[1,4]methanonaphthalene]), 2,2,6,6,7,8,8-heptamethyldodecahydro-2H-indeno[4,5-b]furan, 1,3,4,6,7,8-alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4-alpha-methanophtalen-8(5H)-one, (3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-yl isobutyrate), (3S,5aR,7aS,11aS,11bR)-3,8,8,11a-tetramethyldodecahydro-5H-3,5a-epoxynaphtho[2,1-c]oxepine), 8,8-dimethyl-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl propionate,

(4aR,5R,7aS,9R)-2,2,5,8,8,9a-hexamethyloctahydro-4H-4a,9-methanoazuleno[5,6-d][1,3]dioxole), 2-(8-isopropyl-6-methylbicyclo[2.2.2]oct-5-en-2-yl)-1,3-dioxolane), 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl pivalate, (3a,5,6,7,8,8b-hexahydro-2,2,6,6,7,8,8-heptamethyl-4H-indeno(4,5-d)-1,3-dioxole), , (3R,3aS,6R,7R,8aS)-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulen-3-yl formate (3R,3aS,6R,7R,8aS)-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulen-6-ol, (1-((2S,3S)-2,3,8,8-tetramethyl-1,2,3,4,5,6,7,8-octahydronaphthalen-2-yl)ethan-1-one), ((E)-4-((3aS,7aS)-octahydro-5H-4,7-methanoinden-5-ylidene)butanal), , (Z)-4,11,11-trimethyl-8-methylenebicyclo[7.2.0]undec-3-en-5-yl acetate , (1aR,4S,4aS,7R,7aS,7bS)-1,1,4,7-tetramethyldecaahydro-1H-cyclopropa[e]azulen-4-ol), ((Z)-6-ethylideneoctahydro-2H-5,8-methanochromen-2-one), 1-((3R,3aR,7R,8aS)-3,6,8,8-tetramethyl-2,3,4,7,8,8a-hexahydro-1H-3a,7-methanoazulen-5-yl)ethan-1-one), 3,5,5,6,7,8,8-heptamethyl-5,6,7,8-tetrahydronaphthalene-2-carbonitrile, 2-(cyclohexyloxy)-1,7,7-trimethylbicyclo[2.2.1]heptane (E)-4-((3aR,4R,7R,7aR)-1,3a,4,6,7,7a-hexahydro-5H-4,7-methanoinden-5-ylidene)-3-methylbutan-2-ol), ((E)-3,7-dimethylocta-2,6-dien-1-yl palmitate) and mixtures thereof.

C. The composition according to paragraph according to A-B, wherein malodor reduction material has a ClogP >3.

D. The composition according to paragraph according to A-C, wherein the malodor reduction material has a VP>.005.

E. The composition according to paragraph A-D, wherein the malodor reduction material has a MORV>3, a ClogP>3 and a VP>0.005.

F. The composition according to paragraph A-E, wherein the malodor reduction material is selected from the group consisting of 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane, 1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene] K, SPIRO[FURAN-2(3H),5'-(4,7-METHANO-5H-INDENE)], DECAHYDRO, (3R,3aR,6S,7S,8aS)-6-methoxy-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulene, Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate, 3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan, alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal, 4,5-epoxy-4,11,11-trimethyl-8-

methylenebicyclo(7.2.0)undecane, 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane and mixtures thereof.

G. The composition according to paragraph A-F, wherein the surfactant is selected from the group consisting of anionic, amphoteric or zwitterionic, cationic or mixtures thereof.

H. The composition according to paragraph A-G, wherein the composition comprises the sum total of from about 0.001% to about 0.75% of the malodor reduction material.

I. The composition according to paragraph A-H, wherein the composition comprises the sum total of from about 0.01% to about 0.5% of the malodor reduction material.

J. The composition according to paragraph A-I, wherein the composition comprising one or more perfume raw materials.

K. The composition according to paragraph A-J, wherein a method of controlling malodors comprising: contacting a situs comprising a malodor and/or a situs that will become malodorous with a hair care composition selected from the group consisting of the hair care composition of paragraph A-J.

L. The composition according to A-K, wherein said situs is a head of hair and said contacting step comprises contacting said head of hair with a sufficient amount of a hair care composition to provide said hair with a level of malodor reduction material at least 0.0001 mg of malodor reduction material per head of hair.

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about
5 40 mm."

Every document cited herein, including any cross referenced or related patent or application is hereby incorporated herein by reference in its entirety unless expressly excluded or otherwise limited. The citation of any document is not an admission that it is prior art with respect to any invention disclosed or claimed herein or that it alone, or in any combination with any other
10 reference or references, teaches, suggests, or discloses any such invention. Further, to the extent that any meaning or definition of a term in this document conflicts with any meaning or definition

of the same term in a document incorporated by reference, the meaning or definition assigned to that term in this document shall govern.

While particular embodiments of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is, therefore, intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

CLAIMS

What is claimed:

1. A hair care composition comprising, based on total composition weight,
 - a) a sum total of from 0.0001% to 2% of alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal and one or more malodor reduction materials having a Sulfur MORV >3;
 - b) from 0.01% to 10% of a scalp active material selected from the group consisting of selenium sulfide, sulfur and mixtures thereof;
 - c) from 0.1% to 40%, of a surfactant.

2. A hair care composition according to any of the preceding claims, wherein the malodor reduction materials are selected from the group consisting of 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane], 1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene, 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate, 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl acetate, (3R,3aR,6S,7S,8aS)-6-methoxy-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulene, Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate), (3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan), alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal, 4,5-epoxy-4,11,11-trimethyl-8-methylenebicyclo(7.2.0)undecane), (4aR,8aS)-7-methyloctahydro-1,4-methanonaphthalen-6(2H)-one), (5-methoxyoctahydro-1H-4,7-methanoindene-2-carbaldehyde), (2R,4a'R,8a'R)-3,7'-dimethyl-3',4',4a',5',8',8a'-hexahydro-1'H-spiro[oxirane-2,2'-[1,4]methanonaphthalene]), 2,2,6,6,7,8,8-heptamethyldecahydro-2H-indeno[4,5-b]furan, 1,3,4,6,7,8alpha-hexahydro-1,1,5,5-tetramethyl-2H-2,4alpha-methanophtalen-8(5H)-one, (3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-yl isobutyrate), (3S,5aR,7aS,11aS,11bR)-3,8,8,11a-tetramethyldodecahydro-5H-3,5a-epoxynaphtho[2,1-c]oxepine), 8,8-dimethyl-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl propionate, (4aR,5R,7aS,9R)-2,2,5,8,8,9a-hexamethyloctahydro-4H-4a,9-methanoazuleno[5,6-d][1,3]dioxole), 2-(8-isopropyl-6-methylbicyclo[2.2.2]oct-5-en-2-yl)-1,3-dioxolane), 3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-6-yl pivalate, (3a,5,6,7,8,8b-hexahydro-2,2,6,6,7,8,8-heptamethyl-4H-indeno(4,5-d)-1,3-dioxole), (3R,3aS,6R,7R,8aS)-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulen-3-yl formate (3R,3aS,6R,7R,8aS)-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulen-6-ol, (1-((2S,3S)-2,3,8,8-tetramethyl-

1,2,3,4,5,6,7,8-octahydronaphthalen-2-yl)ethan-1-one), ((E)-4-((3aS,7aS)-octahydro-5H-4,7-methanoinden-5-ylidene)butanal), (Z)-4,11,11-trimethyl-8-methylenebicyclo[7.2.0]undec-3-en-5-yl acetate, (1aR,4S,4aS,7R,7aS,7bS)-1,1,4,7-tetramethyldecahydro-1H-cyclopropa[e]azulen-4-ol), ((Z)-6-ethylideneoctahydro-2H-5,8-methanochromen-2-one), 1-((3R,3aR,7R,8aS)-3,6,8,8-tetramethyl-2,3,4,7,8,8a-hexahydro-1H-3a,7-methanoazulen-5-yl)ethan-1-one), 3,5,5,6,7,8,8-heptamethyl-5,6,7,8-tetrahydronaphthalene-2-carbonitrile, 2-(cyclohexyloxy)-1,7,7-trimethylbicyclo[2.2.1]heptane (E)-4-((3aR,4R,7R,7aR)-1,3a,4,6,7,7a-hexahydro-5H-4,7-methanoinden-5-ylidene)-3-methylbutan-2-ol), ((E)-3,7-dimethylocta-2,6-dien-1-yl palmitate) and mixtures thereof.

3. A hair care composition according to any of the preceding claims wherein malodor reduction material has a ClogP > 3.

4. A hair care composition according to any of the preceding claims wherein the malodor reduction material has a VP > 0.005.

5. A hair care composition according to any of the preceding claims wherein the malodor reduction material has a MORV > 3, a ClogP > 3 and a VP > 0.005.

6. A hair care composition according to any of the preceding claims wherein the malodor reduction material is selected from the group consisting of 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane, 1',1',5',5'-tetramethylhexahydro-2'H,5'H-spiro[[1,3]dioxolane-2,8'-[2,4a]methanonaphthalene] K, SPIRO[FURAN-2(3H),5'-(4,7-METHANO-5H-INDENE)], DECAHYDRO, (3R,3aR,6S,7S,8aS)-6-methoxy-3,6,8,8-tetramethyloctahydro-1H-3a,7-methanoazulene, Ethyl (1R,2R,3R,4R)-3-isopropylbicyclo[2.2.1]hept-5-ene-2-carboxylate, 3aR,5aR,9aR,9bR)-3a,6,6,9a-tetramethyldodecahydronaphtho[2,1-b]furan, alpha,alpha,6,6-tetramethyl bicyclo[3.1.1]hept-2-ene-propanal, 4,5-epoxy-4,11,11-trimethyl-8-methylenebicyclo(7.2.0)undecane, 2'-isopropyl-1,7,7-trimethylspiro[bicyclo[2.2.1]heptane-2,4'-[1,3]dioxane and mixtures thereof.

7. A hair care composition according to any of the preceding claims, wherein the surfactant is selected from the group consisting of anionic, amphoteric or zwitterionic, cationic or mixtures thereof.

8. A hair care composition according to any of the preceding claims, wherein the composition comprises the sum total of from 0.001% to 0.75% of the malodor reduction material.
9. A hair care composition according to any of the preceding claims, wherein the composition comprises the sum total of from 0.01% to 0.5% of the malodor reduction material.
10. A hair care composition according to any of the preceding claims, the composition comprising one or more perfume raw materials.
11. A method of controlling malodors comprising: contacting a situs comprising a malodor and/or a situs that will become malodorous with a hair care composition selected from the group consisting of the hair care composition according to any of the preceding claims.
12. The method according to any of the preceding claims wherein, said situs is a head of hair and said contacting step comprises contacting said head of hair with a sufficient amount of a hair care composition to provide said hair with a level of malodor reduction material at least 0.0001 mg of malodor reduction material per head of hair.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/023758

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K8/33 A61Q5/02 A61Q5/12 A61K8/23
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 089 945 A (BRINKMAN RICHARD EDWARD ET AL) 16 May 1978 (1978-05-16) column 1, lines 15-18, 49-61 column 2, lines 2-16 column 7, lines 6-30; claims; examples -----	1-12
X	EP 0 339 539 A2 (PROCTER & GAMBLE [US]) 2 November 1989 (1989-11-02) page 7, lines 11-22; examples I-III -----	1-12
X	US 2011/308556 A1 (SMETS JOHAN [BE] ET AL) 22 December 2011 (2011-12-22) paragraphs [0002], [0005], [0006]; example 8; table 1 -----	1-12
T	US 2017/119917 A1 (THE PROCTER & GAMBLE COMPANY) 4 May 2017 (2017-05-04) the whole document -----	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 May 2017

Date of mailing of the international search report

23/05/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Mitchell, Gemma

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/023758

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4089945	A	16-05-1978	NONE
EP 0339539	A2	02-11-1989	AU 635205 B2 18-03-1993
		CA 1328816 C 26-04-1994	
		CN 1037271 A 22-11-1989	
		DK 207289 A 29-10-1989	
		EP 0339539 A2 02-11-1989	
		FI 892014 A 29-10-1989	
		JP H0256412 A 26-02-1990	
		NZ 228466 A 28-05-1990	
		PH 25172 A 27-03-1991	
US 2011308556	A1	22-12-2011	CA 2799484 A1 29-12-2011
		CA 2900812 A1 29-12-2011	
		CN 102947433 A 27-02-2013	
		EP 2585568 A1 01-05-2013	
		EP 3121255 A1 25-01-2017	
		EP 3121256 A1 25-01-2017	
		JP 5868966 B2 24-02-2016	
		JP 2013534957 A 09-09-2013	
		US 2011308556 A1 22-12-2011	
		US 2015141310 A1 21-05-2015	
		US 2016326458 A1 10-11-2016	
		WO 2011163325 A1 29-12-2011	
US 2017119917	A1	04-05-2017	CA 2959555 A1 31-03-2016
		CA 2959691 A1 31-03-2016	
		CA 2959697 A1 31-03-2016	
		CA 2960390 A1 31-03-2016	
		US 2016089317 A1 31-03-2016	
		US 2016089318 A1 31-03-2016	
		US 2016089462 A1 31-03-2016	
		US 2016089464 A1 31-03-2016	
		US 2016089465 A1 31-03-2016	
		US 2016090555 A1 31-03-2016	
		US 2016090556 A1 31-03-2016	
		US 2016090557 A1 31-03-2016	
		US 2016090558 A1 31-03-2016	
		US 2016092661 A1 31-03-2016	
		US 2016306909 A1 20-10-2016	
		US 2017119917 A1 04-05-2017	
		WO 2016049389 A1 31-03-2016	
		WO 2016049393 A1 31-03-2016	
		WO 2016049394 A1 31-03-2016	
		WO 2016049395 A1 31-03-2016	
		WO 2016049396 A1 31-03-2016	
		WO 2016049397 A1 31-03-2016	
		WO 2016049398 A1 31-03-2016	
		WO 2016049404 A1 31-03-2016	
		WO 2016049407 A1 31-03-2016	
		WO 2016049455 A1 31-03-2016	
		WO 2016049458 A1 31-03-2016	