



- (51) International Patent Classification:
A61K 31/4192 (2006.01)
- (21) International Application Number:
PCT/US2014/042418
- (22) International Filing Date:
13 June 2014 (13.06.2014)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/835,297 14 June 2013 (14.06.2013) US
- (71) Applicant: UNIVERSITY OF UTAH RESEARCH FOUNDATION [US/US]; 615 Arapeen Drive, Suite 301, Salt Lake City, UT 84108 (US).
- (72) Inventors: PLANELLES, Vicente; 1931 Wyoming Street, Salt Lake City, UT 84108 (US). BOSQUE-PARDOS, Alberto; 758 East 900 South, Salt Lake City, UT 85105 (US). IRELAND, Chris, M.; 1027 E. South Temple, Salt Lake City, UT 84102 (US). VAN WAGONER, Ryan; 717n Park Street, Salt Lake City, UT 84112 (US). HARPER-IRELAND, Mary, Kay; 1027 E. South Temple, Salt Lake City, UT 84102 (US). MASCHEK, John, Alan; 527 J Street, Salt Lake City, UT 84112 (US).
- (74) Agents: SHORTELL, D., Brian et al.; Ballard Spahr LLP, 999 Peachtree Street NE, Suite 1000, Atlanta, GA 30309 (US).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: TRIAZOL-1-OL ANALOGS ANTI-RETROVIRAL LATENCY DRUGS




FIG. 1A

(57) Abstract: In one aspect, the invention relates to triazol-1-ol compounds, analogs thereof, compositions comprising same, and methods of using same, alone or in combination with other agents, to reactivate latent retroviruses, and more particularly to reactivate latent HIV-1. Such compounds, compositions, and methods can be used, for example, in connection with diagnosing and/or treating a retrovirus, and more specifically HIV-1. This abstract is intended as a scanning tool for purposes of searching in the particular art and is not intended to be limiting of the present invention.

TRIAZOL-1-OL ANALOGS AS ANTI-RETROVIRAL LATENCY DRUGS**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This Application claims the benefit of U.S. Provisional Application No. 61/835,297, filed on June 14, 2013, which is incorporated herein by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with government support under Grant No. R01 AI 087508-01A1 awarded by the National Institutes of Health. The United States government has certain rights in the invention.

BACKGROUND

[0003] The advent of highly active anti-retroviral therapy (HAART), which involves the use of three or more antiretroviral drugs, has led to a significant improvement in the care and survival of patients infect with HIV-1. In patients not infected with resistant strains of the virus, HAART typically results in a dramatic decrease in viral load often from levels of 10,000-100,000 RNA copies/mL of plasma to less than 50 copies/mL.

[0004] Given the dramatic effects of HAART, it was proposed that complete elimination of the virus might be possible within 2 to 3 years. However, even after long-term suppression of viral replication with HAART, the virus rapidly rebounds after therapy is discontinued. A key contributor to viral rebound appears to be a reservoir of latently infected cells, including CD4 memory T cells. The half-life of the latently infected population is quite long, and it is estimated that it would take over 60 years of HAART to eliminate this population. Therefore, life-long HAART would be required to control infection in patients

[0005] Retroviruses, including HIV-1, are RNA viruses that replicate through a DNA intermediate and integrate very efficiently into the genome of an infected cell, forming a provirus. Once the provirus is formed, it is maintained in the genome of the infected cell and transferred to daughter cells in the same fashion as any other genetic element within the cellular genome. Thus, the virus has the potential to persist if it infects long-lived cells such as memory T cells. It has been known since 1986 that HIV-1 can establish a latent infection in culture. It


was found that a human T cell line infected with replication-competent virus could develop a latent infection in which the provirus was dormant but could be reactivated upon stimulation. Since then it has been established that a number of cytokines can reactivate latent proviruses.

[0006] The role that latency is playing in preventing clearance of the virus infection has become evident in recent years. Patients that had been successfully treated with HAART in which viral RNA was maintained at levels below 50 copies/mL in the plasma for years, experienced rapid virus rebound upon withdrawal of therapy. Moreover, it was found that after T cell activation, virus could be isolated from CD4 T cells taken from these patients making it clear that to eradicate the virus it will be necessary to eliminate the latently infected cells.

[0007] There have been attempts to flush the latent virus from infected individuals by nonspecific activation of T cells to “turn on” latent proviruses. As part of this approach, the patients remain on HAART to prevent new infections, and the infected cells from which the latent proviruses are activated should die due to cytotoxic effects of viral expression and /or because of targeting by the immune system which can recognize the cells once they begin to express the viral proteins. Agents that can dissociate latent virus activation from global T cell activation, however, are extremely rare. Thus, there remains a need for new drugs capable of selectively activating latent viruses.

SUMMARY


[0008] In accordance with the purpose(s) of the invention, as embodied and broadly described herein, the invention, in one aspect, relates to methods of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium


hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring.

[0009] Also disclosed are methods for the manufacture of a medicament for treatment of a retrovirus in a subject, the method comprising the step of combining an effective amount of a compound represented by a formula:



wherein n is 0 or 1, wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, $SO_2N(CH_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), or 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring; with a pharmaceutically acceptable carrier or diluent.

[0010] Also disclosed are kits comprising at least one compound represented by a formula:




wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, $SO_2N(CH_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-

methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring; and one or more of:

- a) at least one agent known to treat a retrovirus;
- b) instructions for detecting a retrovirus; and
- c) instructions for treating a retrovirus.

[0011] Also disclosed are pharmaceutical compositions comprising a compound represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, $SO_2N(CH_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring; and a pharmaceutically acceptable carrier or diluent.

[0012] While aspects of the present invention can be described and claimed in a particular statutory class, such as the system statutory class, this is for convenience only and one of skill in the art will understand that each aspect of the present invention can be described and claimed in any statutory class. Unless otherwise expressly stated, it is in no way intended that any method

or aspect set forth herein be construed as requiring that its steps be performed in a specific order. Accordingly, where a method claim does not specifically state in the claims or descriptions that the steps are to be limited to a specific order, it is no way intended that an order be inferred, in any respect. This holds for any possible non-express basis for interpretation, including matters of logic with respect to arrangement of steps or operational flow, plain meaning derived from grammatical organization or punctuation, or the number or type of aspects described in the specification.

BRIEF DESCRIPTION OF THE FIGURES

[0013] The accompanying figures, which are incorporated in and constitute a part of this specification, illustrate several aspects and together with the description serve to explain the principles of the invention.

[0014] FIG. 1A shows the results of a screen of the Ireland Natural Product Collection, a set of natural products isolated from marine invertebrate animals and microorganisms. Plate II well C07 was identified as a positive result and was taken on for further analysis.

[0015] FIG. 1B shows that Plate II well C07 activates latent HIV-1 as indicated by an increase in fluorescence compared to the control.

[0016] FIG. 1C shows the % of positive cells and the % of reactivation compared to the positive control for 20 fractions of C07. The fractions were separated using HPLC with a gradient of 10-100% in acetonitrile with 0.1% formic acid. The activity of CD3/CD28 is shown as a positive control. The ability of C07 (MNP-II C07) to activate latent HIV-1 is also displayed.

[0017] FIG. 1D shows that the compound responsible for the activity of C07 is located in fraction M5. The major component of this fraction is 1-hydroxybenzotriazole (HOBt).

[0018] FIG. 1E shows the structure of HOBt.

[0019] FIG. 1F shows that HOBt reactivates HIV-1 latency in a dose dependent manner.

[0020] FIG. 1G shows that HOBt is able to reactive HIV-1 latency with slower kinetics compared to anti-CD3/CD28.

[0021] FIG. 2A shows the ability of HOBt and HOBt analogs to activate latent HIV-1. In addition to HOBt, HOAt also activates HIV-1 latent cells.

[0022] FIG. 2B shows that both HOBt and HOAt activate latent HIV-1.

[0023] FIG. 2C shows that HOBt and HOAt reactivate HIV-1 latency in a dose dependent manner.

[0024] FIG. 2D shows the ability of a panel of 7 derivatives of HOBt to activate latent HIV-1. HOAt demonstrates increased activity compared to HOBt. Two additional analogs, HATU and HO-DhBt, also illustrate an increased ability to induce viral reactivation in a dose dependent manner.

[0025] FIG. 3 shows that HOBt and HOAt synergize with several γ c-cytokines to reactivate HIV-1 latency. HOBt and HOAt increase viral reactivation in the presence of IL-2, IL-7 and IL-5. IL-4 did not induce viral reactivation by itself; however, combining IL-4 with either HOBt or HOAt did induce significant viral reactivation. IL-9 and IL-21 show no effect.

[0026] FIG. 4A shows that HOAt induces viral reactivation in a STAT5 dependent manner. Inhibition of either NFAT or NF- κ B did not have an effect on viral reactivation by HOAt. Inhibition of JAK selectively blocked viral reactivation by HOAt but not by α CD3/ α CD28. While the STAT3 inhibitor did not have a significant effect, the STAT5 inhibitor was able to completely prevent activation of latent HIV-1 by HOAt.

[0027] FIG. 4B shows that HOAt alone is able to induce an increase in STAT5 phosphorylation over untreated cells. While cells incubated with IL-2 alone increase the levels of STAT5 phosphorylation at 30 min, at 24 hrs the level has decreased to basal levels. Incubation of cultured T_{CM} cells with a combination of IL-2 and HOAt lead to a drastic increase in STAT5 phosphorylation.

[0028] FIG. 5A shows that HOAt increases the nuclear localization of STAT5. pSTAT5 and STAT5 levels increased in the nucleus of the cells treated with HOAt at the time points indicated. Histone H3 and alpha Tubulin were used as controls for purity of the fractionation.

[0029] FIG. 5B shows that IL-2 alone or in the presence of HOAt increases phosphorylation of JAK1 and JAK3.

[0030] FIG. 5C shows that HOAt increases the transcription of SOC3S, a target gene for STAT5.

[0031] FIG. 6A shows that HOBt does not induce general T cell activation.

[0032] FIG. 6B shows that HOBt does not induce T cell differentiation measured as lost CCR7 and CD27 expression.

[0033] FIG. 6C shows that HOBt only induces some degree of T cell proliferation at the higher concentration tested.

[0034] FIG. 6D shows that HOBt only induces an increase of 1.4 times of CD25 levels at the maximum concentration used. This level is increased by 37 times after TCR engagement with α CD3/ α CD28.

[0035] FIG. 6E shows that HOAt slightly increases the levels of T cell activation when compared with α CD3/ α CD28.

[0036] FIG. 7 shows that HOAt increases viral reactivation mediated by PMA, ionomycin and SAHA. This indicates that HOAt and additional analogs can be used to synergize the activity of other anti-latency drugs.

[0037] FIG. 8 shows representative data pertaining to the ability of HOAt to reactivate latent HIV-1 in cells isolated from aviremic HIV-1 patients. Briefly, cells for two patients suppressed with antiretroviral were isolated and treated with either DMSO (Control), HOAt or the mitogen PHA. As it is shown, HOAt can induce regrowth of latent viruses in both patients *in vitro* above the treatment control (DMSO).

[0038] FIG. 9A shows representative data demonstrating that HOAt increases the levels of pSTAT5 in the inputs. FIG. 9B shows representative data illustrating the ladder pattern normally observed in ubiquitinated proteins. Without wishing to be bound by theory, this may suggest that HOAt and derivatives may block ubiquitination and subsequent degradation of pSTAT5. Briefly, cells were treated with DMSO or HOAt for 30 min, and subsequently treated with IL-2 for another 30 min. STAT5 was immunoprecipitated with antibodies specific for the phosphorylated protein and western blot was performed against pSTAT5.

[0039] FIG. 10 shows representative data pertaining to the ability of HOBt, HOAt, HODHBt and HATU to maintain pSTAT5 levels in a dose dependent manner. Cells were treated with IL-2 and several derivatives of HOAt. Levels of pSTAT5 were measured 48 h later by flow cytometer.

[0040] Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or can be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DESCRIPTION

[0041] The present invention can be understood more readily by reference to the following detailed description of the invention and the Examples and Figures included herein.

[0042] Before the present compounds, compositions, articles, systems, devices, and/or methods are disclosed and described, it is to be understood that they are not limited to specific synthetic methods unless otherwise specified, or to particular reagents unless otherwise specified, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular aspects only and is not intended to be limiting. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, example methods and materials are now described.

[0043] While aspects of the present invention can be described and claimed in a particular statutory class, such as the system statutory class, this is for convenience only and one of skill in the art will understand that each aspect of the present invention can be described and claimed in any statutory class. Unless otherwise expressly stated, it is in no way intended that any method or aspect set forth herein be construed as requiring that its steps be performed in a specific order. Accordingly, where a method claim does not specifically state in the claims or descriptions that the steps are to be limited to a specific order, it is no way intended that an order be inferred, in any respect. This holds for any possible non-express basis for interpretation, including matters of logic with respect to arrangement of steps or operational flow, plain meaning derived from

grammatical organization or punctuation, or the number or type of aspects described in the specification.

[0044] Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this pertains. The references disclosed are also individually and specifically incorporated by reference herein for the material contained in them that is discussed in the sentence in which the reference is relied upon. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided herein may be different from the actual publication dates, which can require independent confirmation.

A. DEFINITIONS

[0045] As used herein, nomenclature for compounds, including organic compounds, can be given using common names, IUPAC, IUBMB, or CAS recommendations for nomenclature. When one or more stereochemical features are present, Cahn-Ingold-Prelog rules for stereochemistry can be employed to designate stereochemical priority, *E/Z* specification, and the like. One of skill in the art can readily ascertain the structure of a compound if given a name, either by systemic reduction of the compound structure using naming conventions, or by commercially available software, such as CHEMDRAWTM (Cambridgesoft Corporation, U.S.A.).

[0046] As used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a functional group,” “an alkyl,” or “a residue” includes mixtures of two or more such functional groups, alkyls, or residues, and the like.

[0047] Ranges can be expressed herein as from “about” one particular value, and/or to “about” another particular value. When such a range is expressed, a further aspect includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent “about,” it will be understood that the particular value forms a further aspect. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also

herein disclosed as “about” that particular value in addition to the value itself. For example, if the value “10” is disclosed, then “about 10” is also disclosed. It is also understood that each unit between two particular units are also disclosed. For example, if 10 and 15 are disclosed, then 11, 12, 13, and 14 are also disclosed.

[0048] References in the specification and concluding claims to parts by weight of a particular element or component in a composition denotes the weight relationship between the element or component and any other elements or components in the composition or article for which a part by weight is expressed. Thus, in a compound containing 2 parts by weight of component X and 5 parts by weight component Y, X and Y are present at a weight ratio of 2:5, and are present in such ratio regardless of whether additional components are contained in the compound.

[0049] A weight percent (wt. %) of a component, unless specifically stated to the contrary, is based on the total weight of the formulation or composition in which the component is included.

[0050] As used herein, the terms “optional” or “optionally” means that the subsequently described event or circumstance can or can not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

[0051] As used herein, the term “subject” can be a vertebrate, such as a mammal, a fish, a bird, a reptile, or an amphibian. Thus, the subject of the herein disclosed methods can be a human, non-human primate, horse, pig, rabbit, dog, sheep, goat, cow, cat, guinea pig or rodent. The term does not denote a particular age or sex. Thus, adult and newborn subjects, as well as fetuses, whether male or female, are intended to be covered. In one aspect, the subject is a mammal. A patient refers to a subject afflicted with a disease or disorder. The term “patient” includes human and veterinary subjects.

[0052] As used herein, the term “treatment” refers to the medical management of a patient with the intent to cure, ameliorate, stabilize, or prevent a disease, pathological condition, or disorder. This term includes active treatment, that is, treatment directed specifically toward the improvement of a disease, pathological condition, or disorder, and also includes causal treatment, that is, treatment directed toward removal of the cause of the associated disease, pathological condition, or disorder. In addition, this term includes palliative treatment, that is, treatment designed for the relief of symptoms rather than the curing of the disease, pathological condition, or disorder; preventative treatment, that is, treatment directed to minimizing or

partially or completely inhibiting the development of the associated disease, pathological condition, or disorder; and supportive treatment, that is, treatment employed to supplement another specific therapy directed toward the improvement of the associated disease, pathological condition, or disorder. In various aspects, the term covers any treatment of a subject, including a mammal (e.g., a human), and includes: (i) preventing the disease from occurring in a subject that can be predisposed to the disease but has not yet been diagnosed as having it; (ii) inhibiting the disease, i.e., arresting its development; or (iii) relieving the disease, i.e., causing regression of the disease. In one aspect, the subject is a mammal such as a primate, and, in a further aspect, the subject is a human. The term "subject" also includes domesticated animals (e.g., cats, dogs, etc.), livestock (e.g., cattle, horses, pigs, sheep, goats, etc.), and laboratory animals (e.g., mouse, rabbit, rat, guinea pig, fruit fly, etc.).

[0053] As used herein, the term "prevent" or "preventing" refers to precluding, averting, obviating, forestalling, stopping, or hindering something from happening, especially by advance action. It is understood that where reduce, inhibit or prevent are used herein, unless specifically indicated otherwise, the use of the other two words is also expressly disclosed.

[0054] As used herein, the term "diagnosed" means having been subjected to a clinical interview and/or a physical examination by a person of skill, for example, a physician, and found to have a condition that can be diagnosed or treated by the compounds, compositions, or methods disclosed herein. For example, "diagnosed with a depression disorder" means having been subjected to a clinical interview and/or physical examination by a person of skill, for example, a physician, and found to have a condition that can be diagnosed or treated by a compound or composition that can cure, alleviate, prevent, or otherwise treat a depression disorder.

[0055] As used herein, the terms "administering" and "administration" refer to any method of providing a pharmaceutical preparation to a subject. Such methods are well known to those skilled in the art and include, but are not limited to, oral administration, transdermal administration, administration by inhalation, nasal administration, topical administration, intravaginal administration, ophthalmic administration, intraaural administration, intracerebral administration, rectal administration, sublingual administration, buccal administration, and parenteral administration, including injectable such as intravenous administration, intra-arterial administration, intramuscular administration, and subcutaneous administration. Administration can be continuous or intermittent. In various aspects, a preparation can be administered

therapeutically; that is, administered to treat an existing disease or condition. In further various aspects, a preparation can be administered prophylactically; that is, administered for prevention of a disease or condition.

[0056] The term “contacting” as used herein refers to bringing a disclosed compound and a cell, target receptor, or other biological entity together in such a manner that the compound can affect the activity of the target, either directly; i.e., by interacting with the target itself, or indirectly; i.e., by interacting with another molecule, co-factor, factor, or protein on which the activity of the target is dependent.

[0057] As used herein, the term “effective amount” refers to an amount that is sufficient to achieve the desired result or to have an effect on an undesired condition. For example, a “therapeutically effective amount” refers to an amount that is sufficient to achieve the desired therapeutic result or to have an effect on undesired symptoms, but is generally insufficient to cause adverse side effects. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the route of administration; the rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of a compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. If desired, the effective daily dose can be divided into multiple doses for purposes of administration. Consequently, single dose compositions can contain such amounts or submultiples thereof to make up the daily dose. The dosage can be adjusted by the individual physician in the event of any contraindications. Dosage can vary, and can be administered in one or more dose administrations daily, for one or several days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products. In further various aspects, a preparation can be administered in a “prophylactically effective amount”; that is, an amount effective for prevention of a disease or condition.

[0058] The term “pharmaceutically acceptable” describes a material that is not biologically or otherwise undesirable, i.e., without causing an unacceptable level of undesirable biological effects or interacting in a deleterious manner.

[0059] As used herein, the term “derivative” refers to a compound having a structure derived from the structure of a parent compound (e.g., a compound disclosed herein) and whose structure is sufficiently similar to those disclosed herein and based upon that similarity, would be expected by one skilled in the art to exhibit the same or similar activities and utilities as the claimed compounds, or to induce, as a precursor, the same or similar activities and utilities as the claimed compounds. Exemplary derivatives include salts, esters, amides, salts of esters or amides, and N-oxides of a parent compound.

[0060] As used herein, the term “pharmaceutically acceptable carrier” refers to sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. These compositions can also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents such as paraben, chlorobutanol, phenol, sorbic acid and the like. It can also be desirable to include isotonic agents such as sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents, such as aluminum monostearate and gelatin, which delay absorption. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters) and poly(anhydrides). Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues. The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid

compositions which can be dissolved or dispersed in sterile water or other sterile injectable media just prior to use. Suitable inert carriers can include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 micrometers.

[0061] The term “stable,” as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain aspects, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0062] By “virion,” “viral particle,” or “retroviral particle” is meant a single virus minimally composed of an RNA or DNA genome, Pol protein (for reverse transcription of the RNA genome following infection), Gag protein (structural protein present in the nucleocapsid), and an envelope protein. As used herein, the RNA genome of the retroviral particle is usually a recombinant RNA genome, e.g., contains an RNA sequence exogenous to the native retroviral genome and/or is defective in an endogenous retroviral sequence (e.g., is defective in pol, gag, and/or env, and, as used herein, is normally defective in all three genes).

[0063] Compounds described herein comprise atoms in both their natural isotopic abundance and in non-natural abundance. The disclosed compounds can be isotopically-labeled or isotopically-substituted compounds identical to those described, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , respectively. Compounds further comprise prodrugs thereof and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labeled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of the present invention and prodrugs thereof can generally be prepared by carrying

out the procedures below, by substituting a readily available isotopically labeled reagent for a non- isotopically labeled reagent.

[0064] The compounds described in the invention can be present as a solvate. In some cases, the solvent used to prepare the solvate is an aqueous solution, and the solvate is then often referred to as a hydrate. The compounds can be present as a hydrate, which can be obtained, for example, by crystallization from a solvent or from aqueous solution. In this connection, one, two, three or any arbitrary number of solvate or water molecules can combine with the compounds according to the invention to form solvates and hydrates. Unless stated to the contrary, the invention includes all such possible solvates.

[0065] The term "co-crystal" means a physical association of two or more molecules which owe their stability through non-covalent interaction. One or more components of this molecular complex provide a stable framework in the crystalline lattice. In certain instances, the guest molecules are incorporated in the crystalline lattice as anhydrates or solvates, see e.g. "Crystal Engineering of the Composition of Pharmaceutical Phases. Do Pharmaceutical Co-crystals Represent a New Path to Improved Medicines?" Almarasson, O., et. al., The Royal Society of Chemistry, 1889-1896, 2004. Examples of co-crystals include p-toluenesulfonic acid and benzenesulfonic acid.

[0066] It is known that chemical substances form solids which are present in different states of order which are termed polymorphic forms or modifications. The different modifications of a polymorphic substance can differ greatly in their physical properties. The compounds according to the invention can be present in different polymorphic forms, with it being possible for particular modifications to be metastable. Unless stated to the contrary, the invention includes all such possible polymorphic forms.

[0067] Certain materials, compounds, compositions, and components disclosed herein can be obtained commercially or readily synthesized using techniques generally known to those of skill in the art. For example, the starting materials and reagents used in preparing the disclosed compounds and compositions are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wis.), Acros Organics (Morris Plains, N.J.), Fisher Scientific (Pittsburgh, Pa.), or Sigma (St. Louis, Mo.) or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for

Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991); March's Advanced Organic Chemistry, (John Wiley and Sons, 4th Edition); and Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989).

[0068] Unless otherwise expressly stated, it is in no way intended that any method set forth herein be construed as requiring that its steps be performed in a specific order. Accordingly, where a method claim does not actually recite an order to be followed by its steps or it is not otherwise specifically stated in the claims or descriptions that the steps are to be limited to a specific order, it is no way intended that an order be inferred, in any respect. This holds for any possible non-express basis for interpretation, including: matters of logic with respect to arrangement of steps or operational flow; plain meaning derived from grammatical organization or punctuation; and the number or type of embodiments described in the specification.

[0069] Disclosed are the components to be used to prepare the compositions of the invention as well as the compositions themselves to be used within the methods disclosed herein. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutation of these compounds can not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a particular compound is disclosed and discussed and a number of modifications that can be made to a number of molecules including the compounds are discussed, specifically contemplated is each and every combination and permutation of the compound and the modifications that are possible unless specifically indicated to the contrary. Thus, if a class of molecules A, B, and C are disclosed as well as a class of molecules D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited each is individually and collectively contemplated meaning combinations, A-E, A-F, B-D, B-E, B-F, C-D, C-E, and C-F are considered disclosed. Likewise, any subset or combination of these is also disclosed. Thus, for example, the sub-group of A-E, B-F, and C-E would be considered disclosed. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the compositions of the invention. Thus, if there are a variety of additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the methods of the invention.

[0073] As used herein, the term “pharmaceutically acceptable salts” refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (-ic and -ous), ferric, ferrous, lithium, magnesium, manganese (-ic and -ous), potassium, sodium, zinc and the like salts. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, as well as cyclic amines and substituted amines such as naturally occurring and synthesized substituted amines. Other pharmaceutically acceptable organic non-toxic bases from which salts can be formed include ion exchange resins such as, for example, arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

[0074] As used herein, the term “pharmaceutically acceptable non-toxic acids,” includes inorganic acids, organic acids, and salts prepared therefrom, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

[0075] In practice, the compounds of the invention, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier can take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of

the active ingredient. Further, the compositions can be presented as a powder, as granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid, as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compounds of the invention, and/or pharmaceutically acceptable salt(s) thereof, can also be administered by controlled release means and/or delivery devices. The compositions can be prepared by any of the methods of pharmacy. In general, such methods include a step of bringing into association the active ingredient with the carrier that constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

[0076] Thus, the pharmaceutical compositions of this invention can include a pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable salt of the compounds of the invention. The compounds of the invention, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

[0077] The pharmaceutical carrier employed can be, for example, a solid, liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

[0078] In preparing the compositions for oral dosage form, any convenient pharmaceutical media can be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like can be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like can be used to form oral solid preparations such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets can be coated by standard aqueous or nonaqueous techniques

[0079] A tablet containing the composition of this invention can be prepared by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets

can be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets can be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent.

[0080] The pharmaceutical compositions of the present invention comprise a compound of the invention (or pharmaceutically acceptable salts thereof) as an active ingredient, a pharmaceutically acceptable carrier, and optionally one or more additional therapeutic agents or adjuvants. The instant compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions can be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

[0081] Pharmaceutical compositions of the present invention suitable for parenteral administration can be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

[0082] Pharmaceutical compositions of the present invention suitable for injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability. The pharmaceutical compositions must be stable under the conditions of manufacture and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

[0083] Pharmaceutical compositions of the present invention can be in a form suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, mouth washes, gargles, and the like. Further, the compositions can be in a form suitable for use in

transdermal devices. These formulations can be prepared, utilizing a compound of the invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

[0084] Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories can be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

[0085] In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above can include, as appropriate, one or more additional carrier ingredients such as diluents, buffers, flavoring agents, binders, surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation isotonic with the blood of the intended recipient. Compositions containing a compound of the invention, and/or pharmaceutically acceptable salts thereof, can also be prepared in powder or liquid concentrate form.

[0086] It is understood that the disclosed compositions can be prepared from the disclosed compounds. It is also understood that the disclosed compositions can be employed in the disclosed methods of using.


[0087] In a further aspect, the disclosed compositions are for oral administration.

[0088] In a further aspect, the disclosed compositions further comprise a therapeutic agent that can be used to treat a retrovirus. In a still further aspect, the therapeutic agent can be used to treat a retrovirus selected from the group comprising HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4. In yet a further aspect, the therapeutic agent can be used to treat HIV-1. In an even further aspect, the therapeutic agent is selected from entry inhibitors, nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof. In a still further aspect, the therapeutic agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide

reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, or a mixture thereof.

C. METHODS OF USING THE COMPOUNDS AND COMPOSITIONS

[0089] The compounds disclosed herein are useful for treating or detecting a retrovirus, and more particularly for treating or detecting HIV-1. Thus, provided is a method of treating or detecting a retrovirus in a subject comprising the step of administering to the subject at least one compound represented by a formula:




wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring; and/or at least one disclosed pharmaceutical composition in an effective dosage and amount.

[0090] The pharmaceutical compositions and methods of the present invention can further comprise other therapeutically active compounds as noted herein which are usually applied in the treatment of retroviruses, for example HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4.

[0091] In various aspects, the disclosed treatment methods can be applied to a subject, for example, a patient. In further aspects, the subject is a mammal, for example, a human.

1. TREATMENT METHODS


[0092] In one aspect, the invention relates to methods for the treatment of a patient diagnosed with a retrovirus, the method comprising the step of administering to the patient, together in an effective amount, a compound represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring, and at least one agent known to treat a retrovirus.

a. ACTIVATION OF A LATENT RETROVIRUS

[0093] In one aspect, the invention relates to methods of activating a latent retrovirus, and more specifically HIV-1, in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-

(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R² is selected from H and C1-C4 alkyl; and wherein R³ is selected from H and C1-C4 alkyl, or R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring.


[0094] The use of antiretroviral therapy in human immunodeficiency virus type 1 (HIV-1) infected patients does not lead to virus eradication. This is due, to a significant degree, to the fact that HIV-1 can establish a highly stable reservoir of latently infected cells. The principle reservoir of HIV-1 latency is thought to reside in resting, CD4⁺ memory T cells, which harbor integrated HIV-1 (Finzi et al. 1997). The low frequency of latently infected cells (1 in 10⁶ resting CD4⁺ T cells (Chun et al. 1997)), for which known phenotypic markers are not available, poses a great challenge to the study of latency *in vivo*.

[0095] Previous studies on HIV-1 latency were based on the generation of chronically infected cell lines, such as the ACH2 (Folks et al. 1989), JΔK (Antoni et al. 1994), and J-Lat (Jordan et al. 2003) T-cell lines, and the U1 promonocytic cell line (Folks et al. 1987). In these systems, latency was defined as a state in which integrated proviruses failed to drive efficient gene expression. However, these systems do not necessarily reflect the latency state *in vivo* because the lack of viral gene expression is due to mutations in that (ACH2 and U1 (Folks et al. 1989, Folks et al. 1987)) or mutations in the LTR (JΔK T-cell line (Antoni et al. 1994)). While these latency models recapitulate a plethora of mechanisms that can underlie viral latency, the focus of this study was in developing a more general model that did not rely on clonal proviral integration sites, and which utilized non-transformed, primary human T-cells.

[0096] A model using human fetal liver tissue in SCID-hu mice has also generated a great deal of interest in the field of HIV-1 latency (Brooks et al. 2001). This model relies upon infection of thymocytes and the vast majority of latently infected cells in this system are mature, quiescent CD4⁺ single positive naïve T cells. This is in contrast with findings in HIV-1 patients, where the majority of latently infected cells are CD4⁺ memory T cells (Finzi et al. 1997). Although naïve and memory cells share the characteristic of being quiescent, a likely requirement for HIV-1 latency in T cells (Finzi et al. 1997), there are important differences between these cell types that impact latency and reactivation.

[0097] More recently, a model using human, primary cells was disclosed in which relevant signaling pathways involved in viral reactivation are dissected from latently infected memory CD4+ cells (Planelles et al. 2010). Replacement of the nef gene with one encoding the green fluorescent protein (GFP) and using GFP as a readout allows for this primary cell model for the study of HIV-1 latency to be suitable for high throughput screening (HTS) of compounds that reactivate HIV-1 from its latency state. Disclosed herein are methods of activating a latent retrovirus, and more particularly latent HIV-1, using compounds identified in this screen.

[0098] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:
















wherein n is 0 or 1; wherein R¹ is selected from H, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V); wherein R² is selected from H and C1-C4 alkyl; and wherein R³ is selected from H and C1-C4 alkyl, or R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring.







[0099] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a structure depicted in Table 1.


TABLE 1.

Compound Name	Structure
1-hydroxybenzotriazole	


Compound Name	Structure
6-chloro-1 <i>H</i> -benzo[<i>d</i>][1,2,3]triazol-1-ol	
1-hydroxy-7-azabenzotriazole	
4,5-dichloro-1-(cyclopentyloxy)-1 <i>H</i> -1,2,3-benzotriazole	
1-(benzoyloxy)-1 <i>H</i> -1,2,3-benzotriazole	
9-fluorenylmethyl 1-benzotriazolyl carbonate	
1-(((dimethylamino)carbonyl)oxy)-6-nitro-1 <i>H</i> -1,2,3-benzotriazole	
1-(((dimethylamino)sulfonyl)oxy)-6-methyl-1 <i>H</i> -1,2,3-benzotriazole	

Compound Name	Structure
1-[[[(dimethylamino)sulfonyl]oxy]-5-methyl-1 <i>H</i> -1,2,3-benzotriazole	
O-(benzotriazole-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium tetrafluoroborate	
O-(6-chlorobenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium tetrafluoroborate	
(benzotriazole-1-yloxy)dipiperidinecarbenium hexafluorophosphate	
O-(benzotriazole-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate	
O-(6-chlorobenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate	

Compound Name	Structure
(benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate	
O-(benzotriazole-1-yl- <i>N,N,N',N'</i> -bis(pentamethylene)uranium hexafluorophosphate	
(benzotriazole-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate	
(benzotriazole-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate	
(6-chlorobenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate	
(7-azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate	


Compound Name	Structure
3-hydroxybenzo[d][1,2,3]triazin-4-(3 <i>H</i>)-one	

[00100] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring. In a still further aspect, R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl ring. In yet a further aspect, R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused phenyl ring. In an even further aspect, R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered heteroaryl ring. In a still further aspect, R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused pyridine ring.

[00101] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein n is 0, R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring. In a still further aspect, n is 1.


[00102] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-


ylmethylene)piperidin-1-ium hexafluorophosphate (V); and wherein each of R⁴, R⁵, and R⁶ is independently selected from H, Cl, CH₃, and NO₂.

[00103] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V); and wherein each of R⁴, R⁵, and R⁶ is independently selected from H, Cl, CH₃, and NO₂.

[00104] In yet a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein each of R⁴, R⁵, and R⁶ is independently selected from H, Cl, CH₃, and NO₂.


[00105] In an even further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



[00106] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




[00107] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein each of R^4 , R^5 , and R^6 is independently selected from H, Cl, CH_3 , or NO_2 .

[00108] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




[00109] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, $\text{C}(\text{O})\text{Ar}$, $\text{C}(\text{O})\text{N}(\text{CH}_3)_2$, $\text{SO}_2\text{N}(\text{CH}_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-


ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V).

[00110] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V).

[00111] In yet a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




[00112] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein R¹ is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-


ylmethylene)piperidin-1-ium hexafluorophosphate (V); and wherein each of R⁷, R⁸, and R⁹ is independently selected from H, Cl, CH₃, and NO₂.

[00113] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V); and wherein each of R⁷, R⁸, and R⁹ is independently selected from H, Cl, CH₃, and NO₂.

[00114] In yet a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein each of R⁷, R⁸, and R⁹ is independently selected from H, Cl, CH₃, and NO₂.

[00115] In an even further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




[00116] In a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:




wherein R¹ is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V).

[00117] In a still further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V).

[00118] In yet a further aspect, the invention relates to a method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



[00119] In a further aspect, the subject is a population of cells. In a still further aspect, the subject is a population of cells latently infected with a retrovirus. In yet a further aspect, the subject is a population of cells latently infected with HIV-1.

[00120] In a further aspect, the subject is a mammal. In a still further aspect, the subject is a human. In yet a further aspect, the subject is a patient. In an even further aspect, the subject is a patient who has been diagnosed with a need for treatment of a retrovirus prior to the administering step. In a still further aspect, the subject is a patient who has been diagnosed with a need for treatment of HIV prior to the administering step.

b. AGENT KNOWN TO TREAT A RETROVIRUS

[00121] In one aspect, the invention relates to methods of activating a latent retrovirus in a subject, the method further comprising treating the patient with at least one agent that can be used to treat a retrovirus.


[00122] In various aspects, the invention relates to methods of activating latent HIV-1 in a subject, the method further comprising treating the patient with at least one agent that can be used to treat HIV-1. In a further aspect, the at least one agent is selected from entry inhibitors, nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof. In a further aspect, the at least one agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, a mixture thereof. In yet a further aspect, the at least one agent is selected from Maraviroc, Enfuvirtide, Zidovudine, Didanosine, Zalcitabine, Stavudine, Lamivudine, Abacavir, Emtricitabine, Entecavir, Tenofovir, Adefovir, Efavirenz, Nevirapine, Delavirdine, Rilpivirine, Raltegravir, Saquinavir, Ritonavir, Indinavir, Nelfinavir, or Amprenavir, or a mixture thereof. In an even further aspect, the at least one agent is HAART.

[00123] Typically, an agent is administered in an effective amount, per its normal dosing instructions. In one aspect, the effective amount is a therapeutically effective amount.

transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof. In an even further aspect, the at least one agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, or a mixture thereof. In a still further aspect, the at least one agent is selected from Maraviroc, Enfuvirtide, Zidovudine, Didanosine, Zalcitabine, Stavudine, Lamivudine, Abacavir, Emtricitabine, Entecavir, Tenofovir, Adefovir, Efavirenz, Nevirapine, Delavirdine, Rilpivirine, Raltegravir, Saquinavir, Ritonavir, Indinavir, Nelfinavir, or Amprenavir, or a mixture thereof. In yet a further aspect, the at least one agent is HAART.

3. USE OF COMPOUNDS AND COMPOSITIONS

[00128] Also provided are the uses of compounds represented by a formula:




wherein n is 0 or 1, wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, $SO_2N(CH_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), or 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring, and compositions comprising the same.

[00129] In various aspects, the use relates to the activation of a latent retrovirus in a subject. In a further aspect, the use relates to the activation of latent HIV-1 in a subject. In a still further aspect, the subject is a population of cells. In yet a further aspect, the subject is a population of cells latently infected with a retrovirus. In an even further aspect, the subject is a population of cells latently infected with HIV-1. In a still further aspect, the subject is a

mammal. In a still further aspect, the subject is a human. In yet a further aspect, the subject is a patient.

4. Kits

[00130] In one aspect, the invention relates to a kit comprising a compound represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring and one or more of:

- (a) at least one agent that can be used to treat a retrovirus;
- (b) instructions for detecting a retrovirus; and
- (c) instructions for treating a retrovirus.

[00131] In a further aspect, the at least one compound and the at least one agent are co-formulated. In a further aspect, the at least one compound and the at least one agent are co-packaged.


[00132] In a further aspect, the retrovirus is selected from a group comprising HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4. In a still further aspect, the retrovirus is HIV-1.

[00133] The kits can also comprise compounds and/or products co-packaged, co-formulated, and/or co-delivered with other components. For example, a drug manufacturer, a drug reseller, a physician, a compounding shop, or a pharmacist can provide a kit comprising a disclosed compound and/or product and another component for delivery to a patient.

[00134] It is contemplated that the disclosed kits can be used in connection with the disclosed methods of making, the disclosed methods of using, and/or the disclosed compositions.

5. NON-MEDICAL USES

[00135] Also provided are the uses of compounds represented by a formula:



wherein n is 0 or 1; wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); wherein R^2 is selected from H and C1-C4 alkyl; and wherein R^3 is selected from H and C1-C4 alkyl, or R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring as pharmacological tools in the development and standardization of *in vitro* and *in vivo* test systems for the evaluation of the effects of anti-retroviral latency drugs, and more particularly anti HIV-1 latency drugs, in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents that activate latent retroviruses, and more specifically that activate latent HIV-1.

D. REFERENCES

- [00136] Antoni B, Rabson AB, Kinter, A, Bodkin M, Poli G (1994) NF-kappa B-dependent and -independent pathways of HIV activation in a chronically infected T cell line. *Virology* 202:684-694.
- [00137] Bosque A, Planelles V (2009) Induction of HIV-1 latency and reactivation in primary memory CD4+ T cells. *Blood* 113:58-65.
- [00138] Bosque A, Planelles V (2010) "Studies of HIV-1 latency in an ex vivo model that uses primary central memory T cells". *Methods*.
- [00139] Bosque A, Famiglietti M, Weyrich AS, Goulston C, Planelles V (2011) Homeostatic Proliferation Fails to Efficiently Reactivate HIV-1 Latently Infected Central Memory CD4+ T Cells. *PLOS Pathog.* 7(10).
- [00140] Brooks DG, Kitchen SG, Kitchen CM, Scripture-Adams DD, Zack JA (2001) Generation of HIV latency during thymopoiesis. *Nat Med* 7:459-464.
- [00141] Chun TW, Carruth L, Finzi D, et al. (1997) Quantification of latent tissue reservoirs and total body viral load in HIV-1 infection. *Nature* 387:183-188.
- [00142] Finzi D, Hermankova M, Pierson T, et al. (1997) Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science* 278:1295-1300.
- [00143] Folks TM, Clouse KA, Justement J, et al. (1989) Tumor necrosis factor alpha induces expression of human immunodeficiency virus in a chronically infected T-cell clone. *Proc Natl Acad Sci USA* 86:2365-2368.
- [00144] Folks TM, Justement J, Kinter A, Dinarello CA, Fauci AS (1987) Cytokine-induced expression of HIV-1 in a chronically infected promonocyte cell line. *Science* 238:800-802.
- [00145] Jordan A, Bisgrove D, Verdin E (2003) HIV reproducibly establishes a latent infection after acute infection of T cells in vitro. *Embo J* 22:1868-1877.

[00146] Planelles V, Bosque A, Methods and compositions relating to viral latency, U.S. Patent Application 2010/0291067 A1, November 2010.

E. EXPERIMENTAL

[0001] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices and/or methods claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (*e.g.*, amounts, temperature, etc.), but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or near atmospheric.

1. GENERAL METHODS

[00147] A novel primary cell model for the study of HIV-1 latency was adapted so as to be suitable for high-throughput screening (HTS) of new compounds that may reactivate HIV-1 from its latency state (Bosque and Planelles, 2009; Bosque and Planelles, 2010; and patent application: Methods and compositions relating to viral latency, Planelles V, Bosque A, U.S. Patent Application 2010/0291067 A1, November 2010).

[00148] To facilitate HTS, the *nef* gene was replaced with that encoding the green fluorescent protein (GFP). It was confirmed that deletion of *nef* has no effect in latency/reactivation (Bosque and Planelles, 2009) and that using GFP provides a faster, more economical, and equally sensitive, as compared to intracellular p24 detection, readout.

[00149] The reactivation experiments were modified to be performed in 96-well plates, utilizing 100,000 cells per well. After reactivation, flow cytometric analysis was performed with Becton Dickinson FACSCanto II with a High Throughput Sampler (HTS) for microtiter plates that is available through the Flow Cytometry Core Facility at the University of Utah. It takes about 30 minutes to analyze a plate, when acquiring 10,000 cells per well. Analysis for each well includes (a) green fluorescence as a direct indicator of viral reactivation; and (b) forward and side scatter as general measures of cell morphology, which is a gross indication of viability.

2. ACTIVATION OF LATENT HIV-1 IN LATENTLY INFECTED CULTURED T_{CM}









[00150] Latently infected cells were generated using healthy, uninfected DONOR 78 cells that were infected with DHIV virus. Reactivation was monitored by analysis of GFP by Flow Cytometry 72 hours after stimulation. Beads coated with anti-CD3/anti CD28 (CD3/CD28) were used as a positive reactivation stimulus. Reactivation was done in a 96-well plate round bottom. Screening of the Ireland Natural Product Collection, a set of natural products isolated from marine invertebrate animals and microorganisms, resulted in a positive hit in Plate 2 well C7 (FIG. 1A and 1B). This compound is an intermediate en route to synthesis of a cyclic peptide isolated from a tunicate.


[00151] HPLC was used to generate 20 fractions based on elution time using a gradient from 10-100% acetonitrile in 0.1% formic acid using a Phenomenex Kinetic C18 50x2.10 mm column (2.6 μ M, 100 A). The ability of these fractions to activate HIV-1 was tested and compared in parallel with the original hit. FIG. 1C shows the % of p24 positive cells in the Y-axis and the % of reactivation compared to the positive control (beads coated with anti-CD3/anti-CD28) at the top of each bar. This data reveals that the active compound is enriched in fractions 2 and 3. Further fractionation of C7 was performed and each fraction tested at the concentration indicated in the graph (FIG. 1D). The active component is found in fraction M5. This fraction was found to be enriched in 1-hydroxybenzotriazol (HOBt) (FIG. 1E). The ability of HOBt to activate latent HIV-1 in a dose-dependent manner compared to C7 was also established (FIG. 1F). Finally, HOBt was tested at different concentrations and analyzed on either Day 3, Day 5, or Day 7. When compared with anti-CD3/CD28, a non-druggable treatment due to its side effects, HOBt was found to reactive HIV-1 latency with slower kinetics (FIG. 1G).

3. ACTIVATION OF LATENT HIV-1 BY ADDITIONAL ANALOGS

[00152] The ability of additional triazole analogs including benzotriazole (HBt), 1-hydroxy-7-amino benzotriazole (HOAt), triazole (Ht) and 1-aminobenzotriazole (ABt) to activate latent HIV-1 was also tested (Table 2). Latently infected cells were generated using healthy, uninfected DONOR 78 cells that were infected with DHIV virus. Reactivation was monitored by analysis of p24Gag by Flow Cytometry 72 hours after stimulation. It was demonstrated that both HOBt and HOAt were able to reactivate HIV-1 latency (FIG. 2A and 2C). The activity of both analogs is dose dependent (FIG. 2C).

TABLE 2.

Compound	Structure
HOBt	
HOAt	
6Cl-HOBt	
HBt	
ABt	
Triazole	
HO-DhBt	
HATU	

Compound	Structure
Pyridozine	

[00153] A panel of 7 derivatives of the original hit, HOBt, were tested for their ability to activate latent HIV-1 (FIG. 2D). HOAt is a more potent analog than the original hit. Two additional compounds, HATU and HO-DhBt, were also able to induce viral reactivation in a dose dependent manner and with increased efficiency.

4. SYNERGISTIC EFFECT OF SEVERAL γ -CYTOKINES

[00154] Latently infected cells were generated using healthy, uninfected DONOR 78 cells that were infected with DHIV virus. Reactivation was monitored by analysis of p24Gag by Flow Cytometry 72 hours after stimulation. HOBt was tested in the presence or absence of IL-2, a γ -cytokine required for survival of the cells *in vivo*. The ability of HOBt and HOAt to activate latent HIV-1 in the absence of IL-2 or in the presence of different γ -cytokines, e.g. IL-4, IL-7, IL-15, IL-9, and IL-21, was tested (FIG. 3). In the absence of γ -cytokines both analogs have the ability to significantly reactivate HIV-1 latency. When compared with incubation of the cells in the absence of cytokines, IL-2 and IL-7 trend towards increasing viral reactivation. HOBt and HOAt dramatically increase viral reactivation in the presence of all, IL-2, IL-4, and IL-15. IL-4 was not able to induce significant viral reactivation by itself; however, combination of IL-4 with either HOBt or HOAt induced significant viral reactivation. Finally, IL-9 and IL-21 did not have an effect in viral reactivation wither alone or in combination with HOBt and HOAt.

5. ACTIVATION OF LATENT HIV-1 IN A STAT5 DEPENDENT MANNER

[00155] The ability of chemical inhibitors of known signaling pathways to block viral reactivation mediated by HOAt was examined (FIG. 4A). Blocking either NFAT or NF- κ B with either Cyclosporine A (CsA) or BMS345541 did not have a significant effect on viral reactivation induced by HOAt. CsA is able to block viral reactivation induced by α CD3/ α CD28. The JAK inhibitor DBI strongly blocks viral reactivation mediated by HOAt but not by α CD3/ α CD28. Activation of JAK kinases lead to the activation of the transcription factors

STATs. Specific inhibitors of STAT3 and STAT5 were thus also studied. The STAT3 inhibitor peptide used did not have a significant effect in viral reactivation compared with an inactive control peptide. In the other hand, the STAT5 inhibitor completely abrogates viral reactivation induced by HOAt.

[00156] The levels of STAT5 phosphorylation were analyzed by flow cytometry after incubating cultured T_{CM} with HOAt, IL-2, or a combination of both, during a 24 h period (FIG. 4B). STAT5 phosphorylation could be only slightly detected in the absence of either IL-2 or HOAt at 30 min. HOAt alone is able to induce an increase in STAT5 phosphorylation over untreated cells. Cells incubated with IL-2 alone dramatically increase the levels of STAT5 phosphorylation at 30 min, but these levels decrease to basal levels after 24 hrs of incubation. Incubation of cultured T_{CM} with a combination of IL-2 and HOAt induce a drastic increase in STAT5 phosphorylation. Moreover, the levels of STAT5 phosphorylation do not significantly decrease after 24 hrs of incubation with IL-2 and HOAt.

6. EFFECT ON THE NUCLEAR LOCALIZATION OF STAT5

[00157] The levels of nuclear localization of STAT5 after incubation of the cells with IL-2 alone or in the presence of HOAt were analyzed. Nuclear translocation of STAT5 is required for its transcriptional activity. Both pSTAT5 and STAT5 levels increased in the nucleus of cells treated with HOAt at the time points indicated (FIG. 5A). Histone H3 and alpha Tubulin were used as controls for purity of the fractionation.

[00158] Engagement of γ c-cytokines to its receptors leads to a signaling cascade initiated by the phosphorylation and activation of the janus kinases JAK1 and JAK3. These kinases then phosphorylate and activate the transcription factors STATs. IL-2 alone or in the presence of HOAt increases phosphorylation of JAK1 and JAK3 at similar levels (FIG. 5B). This indicates that HOAt alters STAT5 phosphorylation without altering the phosphorylation pattern of the JAK kinases.

[00159] STAT5 is a transcriptional activator of multiple genes. To address whether HOAt will increase the transcription of any STAT5 target genes, the expression levels of SOCS3, a target gene for STAT5, were analyzed. Incubation of cells with IL-2 and HOAt was found to increase the levels of SOCS3 at 24 h post-treatment (FIG. 5C).

7. ANALYSIS OF T CELL ACTIVATION

[00160] Memory cells were generated from 3 different donors and treated with either HOBt or anti-CD3/CD28 to analyze the effect of HOBt on T cell activation (FIG. 6A). Analysis of T cell activation was done by analyzing the upregulation of the activation markers CD25 and CD69 three days post activation. HOBt does not induce general T cell activation. Analysis of T cell differentiation was done by analyzing the levels of CCR7 and CD27 three days post activation (FIG. 6B). This revealed that HOBt does not induce T cell differentiation measured as loss of CCR7 and CD27 expression.

[00161] In order to determine whether HOAt is inducing T cell proliferation, culter T_{CM} were incubated with HOAt in a dose dependent manner. HOAt only induces some degree of cell proliferation at the higher concentration tested (FIG. 6C). It has been shown previously that both cultured T_{CM} and *ex vivo* T_{CM} express low levels of CD25 (Bosque et al. 2011). This level is increased by 37 times after TCR engagement with α CD3/ α CD28, measured as mean intensity of fluorescence (values between parenthesis, FIG. 6D). HOAt only induced an increase of 1.4 times at the maximum concentration used.

[00162] CD69 upregulation is concomitant with T cell activation. HOAt slightly increases the levels of T cell activation when compared with α CD3/ α CD28 (FIG. 6E). These results indicate that HOAt can reactivate latent HIV-1 in the absence of cellular proliferation of cellular activation.

8. VIRAL REACTIVATION MEDIATED BY OTHER STIMULI

[00163] To test whether HOAt can enhance viral reactivation mediated by other stimuli, cells were reactivated with three different agonists: PMA, a PKC agonist that has been shown to reactivate latent HIV in a NF- κ B dependent manner; ionomycin, a ionophore that increases the intracellular levels of Ca²⁺ and induces the activation of the transcription factor NFAT; and SAHA, a histone deacetylase inhibitor that has been shown to reactive latent HIV-1 *in vitro* and *ex vivo* (FIG. 7). HOAt is able to increase viral reactivation mediated by these three stimuli. These results indicate that HOAt and its derivatives can be used to synergize the activity of other anti-latency drugs.


[00164] It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the

invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

CLAIMS

What is claimed is:

1. A method of activating a latent retrovirus in a subject, the method comprising the step of administering to the subject an effective amount of a compound represented by a formula:



wherein n is 0 or 1;

wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V);


wherein R^2 is selected from H and C1-C4 alkyl; and

wherein R^3 is selected from H and C1-C4 alkyl;

or wherein R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring.

2. The method of claim 1, wherein the retrovirus is selected from the group comprising HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4.
3. The method of claim 1, wherein the retrovirus is HIV-1.
4. The method of claim 1, wherein R^1 is selected from H, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V).


5. The method of claim 1, wherein R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring.
6. The method of claim 1, wherein R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl ring.
7. The method of claim 1, wherein R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused phenyl ring.
8. The method of claim 1, wherein R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered heteroaryl ring.
9. The method of claim 1, wherein R² and R³ are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused pyridine ring.
10. The method of claim 1, wherein n is 0.
11. The method of claim 1, wherein n is 1.
12. The method of claim 1, wherein the compound is represented by a formula:



wherein R¹ is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); and

wherein each of R⁴, R⁵, and R⁶ is independently selected from H, Cl, CH₃, and NO₂.


13. The method of claim 1, wherein the compound is represented by a formula:



wherein R^1 is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V); and


wherein each of R^4 , R^5 , and R^6 is independently selected from H, Cl, CH_3 , and NO_2 .

14. The method of claim 1, wherein the compound is represented by a formula:




wherein each of R^4 , R^5 , and R^6 is independently selected from H, Cl, CH_3 , and NO_2 .


15. The method of claim 1, wherein the compound is represented by a formula:



16. The method of claim 1, wherein the compound is represented by a formula:




17. The method of claim 1, wherein the compound is represented by a formula:




wherein each of R⁴, R⁵, and R⁶ is independently selected from H, Cl, CH₃, or NO₂.

18. The method of claim 1, wherein the compound is represented by a formula:




19. The method of claim 1, wherein the compound is represented by a formula:




wherein R¹ is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V).

20. The method of claim 1, wherein the compound is represented by a formula:




wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V).

21. The method of claim 1, wherein the compound is represented by a formula:




22. The method of claim 1, wherein the compound is represented by a formula:



wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V); and

wherein each of R^7 , R^8 , and R^9 is independently selected from H, Cl, CH₃, and NO₂.


23. The method of claim 1, wherein the compound is represented by a formula:



wherein R^1 is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V); and


wherein each of R^7 , R^8 , and R^9 is independently selected from H, Cl, CH₃, and NO₂.

24. The method of claim 1, wherein the compound is represented by a formula:




wherein each of R^7 , R^8 , and R^9 is independently selected from H, Cl, CH₃, and NO₂.

25. The method of claim 1, wherein the compound is represented by a formula:




26. The method of claim 1, wherein the compound is represented by a formula:




wherein R¹ is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V).

27. The method of claim 1, wherein the compound is represented by a formula:



wherein R¹ is selected from H and *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V).

28. The method of claim 1, wherein the compound is represented by a formula:




29. The method of claim 1, wherein the subject is a population of cells.

30. The method of claim 1, wherein the subject is a population of cells latently infected with a retrovirus.

31. The method of claim 1, wherein the subject is a population of cells latently infected with HIV-1.
32. The method of claim 1, wherein the subject is a mammal.
33. The method of claim 1, wherein the subject is a human.
34. The method of claim 1, wherein the subject is a patient.
35. The method of claim 1, wherein the subject is a patient who has been diagnosed with a need for treatment of a retrovirus prior to the administering step.
36. The method of claim 1, wherein the subject is a patient who has been diagnosed with a need for treatment of HIV-1 prior to the administering step.
37. The method of claim 1, further comprising detecting an active retrovirus in the subject.
38. The method of claim 1, further comprising detecting active HIV-1 in the subject.
39. The method of claim 1, wherein the subject comprises a population of cells and the method further comprises detecting an active retrovirus in the subject.
40. The method of claim 1, wherein the subject comprises a population of cells and the method further comprises detecting active HIV-1 in the subject.
41. The method of claim 1, further comprising treating the patient with at least one agent that can be used to treat a retrovirus.
42. The method of claim 1, further comprising treating the patient with at least one agent that can be used to treat HIV-1.
43. The method of claim 42, wherein the at least one agent is selected from entry inhibitors, nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof.
44. The method of claim 42, wherein the at least one agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors

(NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, or a mixture thereof.

45. The method of claim 42, wherein the at least one agent is selected from Maraviroc, Enfuvirtide, Zidovudine, Didanosine, Zalcitabine, Stavudine, Lamivudine, Abacavir, Emtricitabine, Entecavir, Tenofovir, Adefovir, Efavirenz, Nevirapine, Delavirdine, Rilpivirine, Raltegravir, Saquinavir, Ritonavir, Indinavir, Nelfinavir, or Amprenavir, or a mixture thereof.
46. The method of claim 42, wherein the at least one agent is HAART.
47. A method for the manufacture of a medicament for treatment of a retrovirus in a subject, the method comprising the step of combining an effective amount of a compound represented by a formula:



wherein n is 0 or 1,

wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, $SO_2N(CH_3)_2$, fluorenylmethoxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), or 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V);

wherein R^2 is selected from H and C1-C4 alkyl;

wherein R^3 is selected from H and C1-C4 alkyl;

or wherein R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring;

with a pharmaceutically acceptable carrier or diluent.

48. The method of claim 47, wherein the retrovirus is selected from the group comprising HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4.
49. The method of claim 47, wherein the retrovirus is HIV-1.
50. The method of claim 47, wherein the subject is a population of cells.
51. The method of claim 47, wherein the subject is a population of cells latently infected with a virus.
52. The method of claim 47, wherein the subject is a population of cells latently infected with HIV-1.
53. The method of claim 47, wherein the subject is a mammal.
54. The method of claim 47, wherein the subject is a human.
55. The method of claim 47, wherein the subject is a patient.
56. The method of claim 47, wherein the subject is a patient who has been diagnosed with a need for treatment of a retrovirus prior to the administering step.
57. The method of claim 47, wherein the subject is a patient who has been diagnosed with a need for treatment of HIV-1 prior to the administering step.
58. The method of claim 47, further comprising at least one agent that can be used to treat a retrovirus.
59. The method of claim 47, further comprising at least one agent that can be used to treat HIV-1.
60. The method of claim 59, wherein the at least one agent is selected from entry inhibitors, nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof.
61. The method of claim 59, wherein the at least one agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors

(NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, or a mixture thereof.

62. The method of claim 59, wherein the at least one agent is selected from Maraviroc, Enfuvirtide, Zidovudine, Didanosine, Zalcitabine, Stavudine, Lamivudine, Abacavir, Emtricitabine, Entecavir, Tenofovir, Adefovir, Efavirenz, Nevirapine, Delavirdine, Rilpivirine, Raltegravir, Saquinavir, Ritonavir, Indinavir, Nelfinavir, or Amprenavir, or a mixture thereof.

63. A kit comprising at least one compound represented by a formula:



wherein n is 0 or 1;

wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, C(O)N(CH₃)₂, SO₂N(CH₃)₂, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V);

wherein R^2 is selected from H and C1-C4 alkyl; and

wherein R^3 is selected from H and C1-C4 alkyl;


or wherein R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring;

and one or more of:

- a) at least one agent that can be used to treat a retrovirus;
- b) instructions for detecting a retrovirus; and

c) instructions for treating a retrovirus.

64. The kit of claim 63, wherein the at least one compound and the at least one agent are co-formulated.
65. The kit of claim 63, wherein the at least one compound and the at least one agent are co-packaged.
66. The kit of claim 63, wherein the retrovirus is selected from the group comprising HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4.
67. The kit of claim 63, wherein the retrovirus is HIV-1.
68. A pharmaceutical composition comprising a compound represented by a formula:



wherein n is 0 or 1;

wherein R^1 is selected from H, C1-C4 alkyl, C1-C6 aryl, C(O)Ar, $SO_2N(CH_3)_2$, fluorenylmethyloxycarbonyl, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium tetrafluoroborate, *N*-((dimethylamino)methylene)-*N*-methylmethanaminium hexafluorophosphate (V), tri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (V), tris(dimethylamino)phosphonium hexafluorophosphate (V), 1-(pyrrolidin-1-ylmethylene)pyrrolidin-1-ium hexafluorophosphate (V), and 1-(piperidin-1-ylmethylene)piperidin-1-ium hexafluorophosphate (V);

wherein R^2 is selected from H and C1-C4 alkyl;

wherein R^3 is selected from H and C1-C4 alkyl;

or wherein R^2 and R^3 are covalently bonded and, together with the intermediate atoms, comprise an optionally substituted fused six-membered aryl or heteroaryl ring;

and a pharmaceutically acceptable carrier or diluent.

69. The composition of claim 68, which is for oral administration.
70. The composition of claim 68, further comprising a therapeutic agent that can be used to treat a retrovirus.
71. The composition of claim 68, further comprising a therapeutic agent that can be used to treat HIV-1, HIV-2, SIV, XMRV, HTLV-1, HTLV-2, HTLV-3, or HTLV-4.
72. The composition of claim 68, further comprising a therapeutic agent that can be used to treat HIV-1.
73. The composition of claim 72, wherein the therapeutic agent is selected from entry inhibitors, nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, integrase inhibitors, or maturation inhibitors, or a mixture thereof.
74. The composition of claim 72, wherein the therapeutic agent is selected from nucleoside reverse transcriptase inhibitors (NRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors, or integrase inhibitors, or a mixture thereof.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 1E




FIG. 1F




FIG. 1G




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6E




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10