

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
23 December 2004 (23.12.2004)

PCT

(10) International Publication Number  
**WO 2004/111173 A1**

(51) International Patent Classification<sup>7</sup>: C11D 3/39, 3/395

(21) International Application Number:  
PCT/EP2004/005499

(22) International Filing Date: 21 May 2004 (21.05.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0313246.1 9 June 2003 (09.06.2003) GB  
0401810.7 28 January 2004 (28.01.2004) GB

(71) Applicant (for AE, AG, AU, BB, BW, BZ, CA, CY, EG, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NA, NZ, OM, PG, SC, SD, SG, SL, SZ, TT, TZ, UG, VC, ZA, ZM, ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London Greater London EC4P 4BQ (GB).

(71) Applicant (for AL, AM, AT, AZ, BA, BE, BF, BG, BJ, BR, BY, CF, CG, CH, CI, CM, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GA, GE, GN, GQ, GR, GW, HR, HU, ID, IS, IT, JP, KG, KP, KR, KZ, LR, LT, LU, LV, MA, MC, MD, MG, MK, ML, MR, MX, MZ, NE, NI, NL, NO, PH, PL, PT, RO, RU, SE, SI, SK, SN, SY, TD, TG, TJ, TM, TN, TR, UA, UZ, VN, YU only): UNILEVER NV [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, Mumbai 400 020 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): Ouwendijk-Vrijenhoek (NEE Ouwendijk), Maartje

[NL/NL]; Unilever R & D Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). **SONMEZER, Derdi** [NL/NL]; Unilever R & D Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). **VEERMAN, Simon Marinus** [NL/NL]; Unilever R & D Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).

(74) Agents: **ELLIOTT, Peter, William** et al.; Unilever Intellectual Property Group, Colworth House, Sharnbrook, Bedford, MK44 1LQ (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LIQUID BLEACHING COMPOSITION

(57) Abstract: The present invention concerns the preservation of a catalyst in a bleaching composition whilst facilitating activation of the catalyst. The bleaching composition is substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system.

WO 2004/111173 A1

- 1 -

## LIQUID BLEACHING COMPOSITION

### Field of Invention

The present invention provides a liquid bleaching composition.

### Background of the Invention

The use of bleaching catalysts for stain removal has been developed over recent years. The recent discovery that some catalysts are capable of bleaching effectively in the absence of an added peroxy source has recently become the focus of some interest, for example: WO9965905; WO0012667; WO0012808; WO0029537, and, WO0060045.

The shelf life of a product may be regarded as the period of time over which the product may be stored whilst retaining its required quality. A satisfactory shelf life is in many instances a crucial factor for the success of a commercial product. A product with a short shelf life generally dictates that the product is made in small batches and is rapidly sold to the consumer. It is also a concern to the owners of a brand with a short shelf life that the consumer uses the product within the shelf life otherwise the consumer may be inclined to change to a similar product of another brand. In contrast a similar product with a long shelf life may be made in larger batches, held as stock for a longer period of time and the period of time that a consumer stores the product is not of a great concern to the owners of a particular brand. Despite the shelf life being an important consideration the product must also be active during use.

- 2 -

It is an object of the present invention to provide an air bleaching composition that has improved storage properties whilst being active in use.

### **Summary of Invention**

We have found that the storage pH conditions required for longevity of a liquid bleaching composition are in conflict with the pH required for providing good bleaching activity.

The present invention provides a liquid bleaching composition having a pH between 6 and 7 comprising:

(a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-l-generating bleach system;

(b) a pH changing means; and,

(c) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition,

wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition enters the range pH 7.5 to pH 9.0.

The present invention also provides a method of bleaching a textile comprising the steps of:

(i) diluting from 0.5 to 20 g of a concentrated liquid bleaching composition with 1 litre of water, the concentrated liquid bleaching composition having a pH in the range 6 to 7, the liquid bleaching comprising an organic substance which forms a complex with a transition metal for

- 3 -

bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach, the dilution providing an aqueous liquid bleaching composition having a pH in the range from 7.5 to pH 9.0;

- (ii) treating a textile with the aqueous liquid bleaching composition; and,
- (iii) rinsing the textile with water; and,
- (iv) drying the textile.

The term "substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system" should be construed within spirit of the invention. It is preferred that the composition has as low a content of peroxy species present as possible. It is preferred that the bleaching formulation contains less than 1 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, preferably the bleaching formulation contains less than 0.3 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, most preferably the bleaching composition is devoid of peracid or hydrogen peroxide or source thereof. In addition, it is preferred that the presence of alkyl hydroperoxides is kept to a minimum in a bleaching composition comprising the ligand or complex of the present invention.

The present invention also extends to a commercial package together with instructions for its use.

- 4 -

**DETAILED DESCRIPTION OF THE INVENTION**THE BLEACH CATALYST

Recently we have found that oily stains are bleached in the presence of selected transition metal catalysts in the absence of an added peroxy source. The bleaching of an oily stain in the absence of an added peroxy source has been attributed to oxygen derived from the air. Whilst it is true that bleaching is effected by oxygen sourced from the air the route in which oxygen plays a part is becoming understood. In this regard, the term "air bleaching" is used.

We have concluded from our research that bleaching of a chromophore in an oily stain is effected by products formed by adventitious oxidation of components in the oily stain. These products, alkyl hydroperoxides, are generated naturally by autoxidation of the oily stain and the alkyl hydroperoxides together with a transition metal catalyst serve to bleach chromophores in the oily stain. Alkyl hydroperoxides (ROOH) are generally less reactive than other peroxy species, for example, peracids (RC(O)OOH), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), percarbonates and perborates. In this regard, the phrase "for bleaching a substrate with atmospheric oxygen" is synonymous with "for bleaching a substrate via atmospheric oxygen" because it is the oxygen in the air that provides the bleaching species used by catalyst to bleach the substrate stain.

The bleach catalyst per se may be selected from a wide range of transition metal complexes of organic molecules (ligands). In typical washing compositions the level of the

- 5 -


organic substance is such that the in-use level is from 0.05  $\mu\text{M}$  to 50 mM, with preferred in-use levels for domestic laundry operations falling in the range 1 to 100  $\mu\text{M}$ . Higher levels may be desired and applied in industrial textile bleaching processes. A mixture of different catalysts may be employed in the bleaching composition.

Suitable organic molecules (ligands) for forming complexes and complexes thereof are found, for example in: GB 9906474.3; GB 9907714.1; GB 98309168.7, GB 98309169.5; GB 9027415.0 and GB 9907713.3; DE 19755493; EP 999050; WO9534628; EP-A-458379; EP 0909809; United States Patent 4,728,455; WO9839098; WO9839406, WO9748787, WO0029537; WO0052124, and WO0060045 the complexes and organic molecule (ligand) precursors of which are herein incorporated by reference. An example of a preferred catalyst is a transition metal complex of MeN4Py ligand (N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane).

The ligand forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI, vanadium II-V and molybdenum II-VI.

An example of a preferred catalyst is a monomer ligand or transition metal catalyst thereof of a ligand having the formula (I):

- 6 -



wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylO-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH<sub>2</sub>, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

C1-C4-alkyl,

C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH<sub>2</sub>)<sub>n</sub>C(O)OR<sub>5</sub>

wherein R<sub>5</sub> is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,


X is selected from C=O, -[C(R<sub>6</sub>)<sub>2</sub>]<sub>y</sub>- wherein Y is from 0 to 3 each R<sub>6</sub> is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

With regard to the above formula (I) it is also particularly preferred that R1 and R2 may also be independently selected from: C1 to C22-optionally substituted alkyl, and an optionally substituted tertiary amine of the form -C2-C4-alkyl-NR<sub>7</sub>R<sub>8</sub>, in which R<sub>7</sub> and R<sub>8</sub> are independently selected from the group consisting of straight chain, branched or

- 7 -

cyclo C1-C12 alkyl, benzyl, the -C2-C4-alkyl- of the -C2-C4-alkyl-NR7R8 may be substituted by 1 to 4 C1-C2-alkyl, or may form part of a C3 to C6 alkyl ring, and in which R7 and R8 may together form a saturated ring containing one or more other heteroatoms.


Another preferred class of ligands are macropolycyclic rigid ligands of the formula:



wherein m and n are 0 or integers from 1 to 2, p is an integer from 1 to 6, preferably m and n are both 0 or both 1 (preferably both 1), or m is 0 and n is at least 1; and p is 1; and A is a nonhydrogen moiety preferably having no aromatic content; more particularly each A can vary independently and is preferably selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, C5-C20 alkyl, and one, but not both, of the A moieties is benzyl, and combinations thereof.

Preferably, the macropolycyclic ligand is of the formula:


- 8 -



wherein "R<sup>1</sup>" is independently selected from H, and linear or branched, substituted or unsubstituted C1-C20 alkyl, alkylaryl, alkenyl or alkynyl; and all nitrogen atoms in the macropolycyclic rings are coordinated with the transition metal.

Of the macropolycyclic ligands 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane is preferred. This ligand is most preferred as its manganese complex [Mn(Bcyclam)Cl<sub>2</sub>] and may be synthesised according to W09839098.

The transition metal complex preferably is of the general formula (AI):



in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe (II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI), preferably from Fe(II)-(III)-(IV)-(V);

- 9 -

L represents the ligand, preferably N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, or its protonated or deprotonated analogue;

X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;

a represents an integer from 1 to 10;

k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 10;

m represents zero or an integer from 1 to 20.

#### pH Jump Composition

The present invention is used as a liquid format where the pH of the commercial product is substantially different to that in use.

Sorbitol/borate compositions are known from EP 381 262. In a concentrated solution the borate is complexed with the vicinal diol of the sorbitol; upon dilution of the sorbitol/borate composition the borate is liberated from its interaction with the sorbitol resulting in a pH jump. Other pH Jump formulations are disclosed in US 6,509,308 and US 5,484,555.

#### Balance Carriers and Adjunct Ingredients

These are generally surfactants, builders, foam agents, anti-foam agents, solvents, and enzymes. The use and amounts of these components are such that the bleaching

- 10 -

composition performs depending upon economics, environmental factors and use of the bleaching composition.

The air bleach catalyst may be used in a detergent composition specifically suited for stain bleaching purposes, and this constitutes a second aspect of the invention. To that extent, the composition comprises a surfactant and optionally other conventional detergent ingredients. The invention in its second aspect provides an enzymatic detergent composition which comprises from 0.1 - 50 % by weight, based on the total detergent composition, of one or more surfactants. This surfactant system may in turn comprise 0 - 95 % by weight of one or more anionic surfactants and 5 to 100 % by weight of one or more nonionic surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost. The enzymatic detergent composition according to the invention will generally be used as a dilution in water of about 0.05 to 2%.

In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.

- 11 -

Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are C<sub>6</sub>-C<sub>22</sub> alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C<sub>8</sub>-C<sub>18</sub> primary or secondary linear or branched alcohols with ethylene oxide, generally 5 to 40 EO.

Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C<sub>8</sub>-C<sub>18</sub> alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C<sub>9</sub>-C<sub>20</sub> benzene sulphonates, particularly sodium linear secondary alkyl C<sub>10</sub>-C<sub>15</sub> benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C<sub>11</sub>-C<sub>15</sub> alkyl benzene sulphonates and sodium C<sub>12</sub>-C<sub>18</sub> alkyl sulphates. Also applicable are surfactants such as those described in

- 12 -

EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system that is a mixture of an alkali metal salt of a C<sub>16</sub>-C<sub>18</sub> primary alcohol sulphate together with a C<sub>12</sub>-C<sub>15</sub> primary alcohol 3-7 EO ethoxylate.

The nonionic detergent is preferably present in amounts greater than 10%, e.g. 25-90% by weight of the surfactant system. Anionic surfactants can be present for example in amounts in the range from about 5% to about 40% by weight of the surfactant system.

One skilled in the art will appreciate that some adventitious peroxy species may be in the composition nevertheless it is most preferred that the bleaching composition of the present invention has less than 1%, preferably less than 0.1%, most preferably less than 0.01%, of a peroxy species present. These adventitious peroxy are predominantly alkyl hydroperoxides formed by autoxidation of the surfactants.

The composition may contain additional enzymes as found in WO 01/00768 A1 page 15, line 25 to page 19, line 29, the contents of which are herein incorporated by reference.

- 13 -

Builders, polymers and other enzymes as optional ingredients may also be present as found in WO0060045.

Suitable detergency builders as optional ingredients may also be present as found in WO0034427.

In the context of the present invention, bleaching should be understood as relating generally to the decolourisation of stains or of other materials attached to or associated with a substrate. However, it is envisaged that the present invention can be applied where a requirement is the removal and/or neutralisation by an oxidative bleaching reaction of malodours or other undesirable components attached to or otherwise associated with a substrate. Furthermore, in the context of the present invention bleaching is to be understood as being restricted to any bleaching mechanism or process that does not require the presence of light or activation by light.

The present invention extends to both isotropic and complex liquid compositions and formulations a brief discussion of which follows. Some isotropic formulations are termed 'micro-emulsion' liquids that are clear and thermodynamically stable over a specified temperature range. The 'micro-emulsion' formulation may be water in oil, or oil in water emulsions. Some liquid formulations are macro-emulsions that are not clear and isotropic. Emulsions are considered meta-stable. Concentrated, clear compositions containing fabric softening actives have been disclosed in WO 98/08924 and WO 98/4799, both Procter & Gamble. Such compositions comprise bio-degradable fabric conditioners.

- 14 -

However, both disclose compositions comprising water miscible solvents that do not form water-in-oil micro-emulsions. Clear fabric conditioning compositions have also been disclosed in EP 730023 (Colgate Palmolive), WO 96/19552 (Colgate Palmolive), WO 96/33800 (Witco Co.), WO 97/03170 (Procter & Gamble), WO 97/03172 (Procter & Gamble), WO 97/03169 (Procter & Gamble), US 5492636 (Quest Int.) and US 5427697 (Procter & Gamble). Liquid formulations of the present invention may contain for example; monoethoxy quats; AQAs and bis-AQAs; cationic amides; cationic esters; amino/diamino quats; glucamide; amine oxides; ethoxylated polyethyleneimines; enhancement polymers of the form linear amine based polymers, e.g. bis-hexamethylenetriamine; polyamines e.g. TETA, TEPA or PEI polymers.

It is preferred that the liquid bleaching composition does not contain a pH dependent chromophore, indicator, e.g., phenolphthalein. In addition, it is preferred that the liquid bleaching composition does not contain a pH dependent fluorescent indicator. In both the above cases such should not be present such that a perceptible change is observed to the human eye.

The liquid may be contained within a sachet as found in WO02/068577. The sachet is a container within the context of the present invention.

The liquid composition preferably also contains one or more antioxidants as described in WO02/072747 and WO02072746.

- 15 -

The following is a example of a concentrated liquid formulation that the present invention may be incorporated into by adding the catalyst together with the selected stable perfume components. The commercial liquid formulation has a pH of 7.

Sodium citrate:	3.2 %
Polypropylene glycol:	4.75 %
LAS-acid:	5.6 %
NI 25 9 EO:	6.6 %
LES (anionic surfactant):	10.5 %
Borax:	2.30 %
Sorbitol:	3.35 %
Alcosperce 725:	0.30 %
Coconut fatty acid:	0.73 %
Monoethanolamine:	0.20 %
Fluorescer:	0.125 %
Enzymes	-
Perfume/dye	-

### Experimental

The following catalyst was used in the experiments 9,9-dihydroxy-2,4-di-(2-pyridyl)-3-methyl-7-(pyridin-2-ylmethyl)-3,7-diaza-bicyclo[3.3.1]nonane-1,5-dicarboxylate Iron (II) dichloride was prepared as described by Heidi Borzel, Peter Comba, Karl S. Hagen, Yaroslaw D. Lampeka, Achim Lienke, Gerald Linti, Michael Merz, Hans Pritzkow, Lyudmyla V. Tsymbal in *Inorganica Chimica Acta* 337 (2002)

- 16 -

407 - 419. WO0248301 provides synthetic details of similar compounds. This catalyst was incorporated at a concentration of 0.03 % wt/wt in concentrated liquid composition A containing.

Liquid A: 6 % LAS, 6 % sLES 3 EO, 6 % Nonionic 7 EO, 0.016 % Proxel GXL, 3.35 % sorbitol, 2.30 % Borax.10 H<sub>2</sub>O, 4.75 % MPG, and sufficient NaOH to bring the pH to 7.

#### Method of Determining Catalyst Stability

The residue bleaching activity of bleaching compositions were determined at 40 °C in a H<sub>2</sub>O<sub>2</sub> containing NaH<sub>2</sub>PO<sub>4</sub>.H<sub>2</sub>O pH7 buffer and Acid Blue 45 (CAS No. 2861-02-1) as substrate using the following protocol.

Samples of 70 mg liquid were diluted in 10.00 ml MilliQ water. We added 45 µl of this solution to an assay of 230 µL containing 20 mM H<sub>2</sub>O<sub>2</sub>, 75 µM Acid blue 45 and 54 mM NaH<sub>2</sub>PO<sub>4</sub>.H<sub>2</sub>O pH7 buffer.

The changes in absorbance at 600 nm were measured for 5 min at 40 °C using a spectrophotometer. The absolute changes in absorbance were correlated to activities obtained with freshly prepared calibration samples. The measured activities were expressed as µMol/l/. The residual amount is the activity measured on t=0 divided by the activity after storage and expressed in percentages.

- 17 -

Table 1: The bleaching catalysts stability expressed as residual amount (%) as function of the pH during storage storage at 37 °C.

pH	1 wk	2 wks	3 wks	4 wks
6	96	82	80	78
6.25	97	84	76	80
6.5	99	91	82	84
6.71	100	88	83	80
7	100	79	71	70
7.25	85	67	56	54
7.52	84	62	54	45
7.75	68	43	38	32
8	62	39	33	25

The results in Table 1 clearly show a storage advantage in maintaining the concentrated liquid bleaching composition at a pH at 7 or below. Upon dilution of the concentrated liquid bleaching composition the pH of the solution changes to alkali which activates the bleaching composition.

- 18 -

**We claim:**

1. A liquid bleaching composition having a pH between 6 and 7 comprising:

(a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system;

(b) a pH changing means; and,

(c) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition,

wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition enters the range pH 7.5 to pH 9.0.


2. A liquid bleaching composition according to claim 1 or 2, wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition reaches at least pH 8.0.

3. A liquid bleaching composition according to claim 1, wherein the pH changing means is provided by a borax sorbitol pH jump composition.

- 19 -

4. A liquid bleaching composition according to claim 3, wherein the liquid bleaching composition comprises at least 2 % wt/wt of sorbitol and at least 1 wt/wt % borax.

5. A liquid bleaching composition according to any preceding claim, wherein the organic substance is of formula (I):



wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylO-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH<sub>2</sub>, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

C1-C4-alkyl,

C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH<sub>2</sub>)<sub>n</sub>C(O)OR<sub>5</sub>

wherein R<sub>5</sub> is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,

X is selected from C=O, -[C(R<sub>6</sub>)<sub>2</sub>]<sub>y</sub>- wherein Y is from 0 to 3 each R<sub>6</sub> is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

- 20 -

6. A liquid bleaching composition according to any one of claims 1 to 4, wherein the organic substance 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane.

7. A method of bleaching a textile comprising the steps of:


(i) diluting from 0.5 to 20 g of a concentrated liquid bleaching composition with 1 litre of water, the concentrated liquid bleaching composition having a pH in the range 6 to 7, the liquid bleaching comprising an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach, the dilution providing an aqueous liquid bleaching composition having a pH in the range from 7.5 to pH 9.0;

(iv) treating a textile with the aqueous liquid bleaching composition; and,

(v) rinsing the textile with water; and,

(iv) drying the textile.

8. A method according to claim 7, wherein the organic substance is of formula (I):



- 21 -

wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH<sub>2</sub>, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

C1-C4-alkyl,

C6-C10-aryl, and,


a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and  $-(CH_2)_n C(O)OR_5$

wherein R5 is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,

X is selected from C=O,  $-[C(R_6)_2]_y-$  wherein Y is from 0 to 3 each R6 is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

9. A method according to claim 7, wherein the organic substance is of formula:



wherein m and n are 0 or integers from 1 to 2, p is an integer from 1 to 6, preferably m and n are both 0 or both 1

- 22 -

(preferably both 1), or m is 0 and n is at least 1; and p is 1; and A is a nonhydrogen moiety preferably having no aromatic content; more particularly each A can vary independently and is preferably selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, C5-C20 alkyl, and one, but not both, of the A moieties is benzyl, and combinations thereof.

10. A liquid bleaching composition according to claim 9, wherein the organic substance 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane.

## AMENDED CLAIMS


[Received by the International Bureau on 17 November 2004 (17.11.2004);  
original claims 1-10 replaced by amended claims 1-6 (4 pages)]

## We claim:

1. A liquid bleaching composition having a pH between 6 and 7 comprising:

(a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system, wherein the organic substance is selected from the group consisting of:

an organic substance of formula (I):



wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH<sub>2</sub>, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:


C1-C4-alkyl,

C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and  $-(CH_2)_n C(O)OR_5$  wherein R5 is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and, X is selected from C=O,  $-[C(R_6)]_y-$  wherein Y is from 0 to 3 each R6 is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl, and

an organic substance of formula:



wherein m and n are 0 or integers from 1 to 2, p is an integer from 1 to 6, preferably m and n are both 0 or both 1 (preferably both 1), or m is 0 and n is at least 1; and p is 1; and A is a nonhydrogen moiety preferably having no aromatic content; more particularly each A can vary independently and is preferably selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, C5-C20 alkyl, and one, but not both, of the A moieties is benzyl, and combinations thereof;

(b) a pH changing means; and,

(c) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition,

wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition enters the range pH 7.5 to pH 9.0.

2. A liquid bleaching composition according to claim 1, wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition reaches at least pH 8.0.
3. A liquid bleaching composition according to claim 1 or 2, wherein the pH changing means is provided by a borax sorbitol pH jump composition.
4. A liquid bleaching composition according to claim 3, wherein the liquid bleaching composition comprises at least 2 % wt/wt of sorbitol and at least 1 wt/wt % borax.
5. A liquid bleaching composition according to any one of claims 1 to 4, wherein the organic substance 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane.
6. A method of bleaching a textile comprising the steps of:
  - (i) diluting from 0.5 to 20 g of a concentrated liquid bleaching composition, as defined in any one of claims 1 to 5, with 1 litre of water, the concentrated liquid bleaching composition having a pH in the range 6 to 7, the liquid bleaching comprising an organic substance which forms a

complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach, the dilution providing an aqueous liquid bleaching composition having a pH in the range from 7.5 to pH 9.0;

- (ii) treating a textile with the aqueous liquid bleaching composition; and,
- (iii) rinsing the textile with water; and,
- (iv) drying the textile.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/005499

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C11D3/39 C11D3/395		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C11D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)  EPO-Internal, WPI Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2002/198127 A1 (ADRIAANSE AREND JAN ET AL) 26 December 2002 (2002-12-26) page 2, paragraph 18 claims 1-38 page 14, paragraph 226 - paragraph 228 page 2, paragraph 26	1-4,7
A	US 5 776 883 A (VASUDEVAN TIRUCHERAI VARAHAN) 7 July 1998 (1998-07-07) claims 1-4	1-10
A	EP 0 732 394 A (UNILEVER PLC ; UNILEVER NV (NL)) 18 September 1996 (1996-09-18) claims 1-10; example 13	1-10
A	GB 2 281 744 A (PROCTER & GAMBLE) 15 March 1995 (1995-03-15) claims 1-11	1-10
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <span style="margin-left: 100px;"><input checked="" type="checkbox"/> Patent family members are listed in annex.</span>		
° Special categories of cited documents :		
*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family	
Date of the actual completion of the international search	Date of mailing of the international search report	
8 October 2004	15/10/2004	
Name and mailing address of the ISA	Authorized officer	
European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Richards, M	

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/005499

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 1 038 947 A (YPLON S A) 27 September 2000 (2000-09-27) claims 1-19 -----	1-10
A	WO 00/60045 A (PERKINS CHRISTOPHER MARK ; PROCTER & GAMBLE (US)) 12 October 2000 (2000-10-12) cited in the application page 35, line 23 - page 36, line 17; claims 1-10 -----	1-10

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/005499

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 2002198127	A1	26-12-2002	BR	0207168 A	10-02-2004
			CA	2435944 A1	06-09-2002
			WO	02068574 A1	06-09-2002
			EP	1368449 A1	10-12-2003
US 5776883	A	07-07-1998	EP	0732394 A2	18-09-1996
			ZA	9601865 A	08-09-1997
EP 0732394	A	18-09-1996	EP	0732394 A2	18-09-1996
			US	5776883 A	07-07-1998
			ZA	9601865 A	08-09-1997
GB 2281744	A	15-03-1995	CA	2126382 A1	31-12-1994
			JP	7070593 A	14-03-1995
EP 1038947	A	27-09-2000	EP	1038947 A2	27-09-2000
			GB	2346153 A ,B	02-08-2000
WO 0060045	A	12-10-2000	AU	4061900 A	23-10-2000
			WO	0060045 A1	12-10-2000