

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2003 (15.05.2003)

PCT

(10) International Publication Number
WO 03/040117 A1

- (51) International Patent Classification⁷: C07D 285/08, 417/12, A61K 31/433, A61P 3/00, 17/00, 25/00
- (21) International Application Number: PCT/US02/35340
- (22) International Filing Date:
4 November 2002 (04.11.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/337,927 8 November 2001 (08.11.2001) US
- (71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL CORPORATION [US/US]; US Route 202, Raritan, NJ 08869 (US).
- (72) Inventors: EISINGER, Magdalena; 30 Pine Terrace, Demarest, NJ 07627 (US). FITZPATRICK, Louis, J.; 205 Heatherfield Drive, Souderton, PA 18964 (US). LEE, Daniel, H.; 361 Dutton Road, Sudbury, MA 01776 (US). PAN, Kevin; 409 Greene Lane, Phoenixville, PA 19460 (US). PLATA-SALAMAN, Carlos; 1313 Squire Drive, Ambler, PA 19002 (US). REITZ, Allen, B.; 109 Greenbriar Road, Lansdale, PA 19466 (US). SMITH-SWINTOSKY, Virginia, L.; 3163 Line Lexington Road, Hatfield, PA 19440 (US). ZHAO, Boyu; 105 Weymouth Circle, Lansdale, PA 19446 (US).
- (74) Agents: JOHNSON, Philip et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/040117 A1

(54) Title: NOVEL 1,2,4-THIADIAZOLE DERIVATIVES AS MELANOCORTIN RECEPTOR MODULATORS



(II)

(57) Abstract: The present invention is directed to novel 1,2,4-thiadiazole derivatives of the formula (II) wherein R¹, R² and R⁴, are ring containing radicals, useful as agonists or antagonists of the melanocortin receptor. More particularly, the compounds of the present invention are useful for the treatment of metabolic, CNS and dermatological disorders.

5 spinal cord, whereas the MC3 receptor is located mainly in the hypothalamus. [Gantz, I., et al., *supra*.] The MC4 receptor is selectively activated by α MSH and can induce neurite outgrowth in Neuro 2A cells. (Adan R.A.H, et al., Molecular Brain Research, **1996**, 36, pp 37-44; Mountjoy, K.G., Mortud, M.T., Low, M.J., Simerly, R.B. and Cone, R.D., Mol. Endocrinol., **1994**, 8, pp 1298-10 1308). ACHT is a less potent activator of the MC4 receptor than α MSH. (Adan, R.A.H., Cone, R.D., Burbach, J.P.H. and Gispen, W.H., mol. pharmacol., **1994**, 46, pp 1182-1190). The MC5 receptor is activated, in order of degree, by NDP \approx α -MSH > ACHT (1-24) \geq α MSH ACHT (1-39) = β MSH >> γ MSH (*The Melanocortin Receptors*, Cone, R.D., Editor, Human Press Inc., 15 Totowa, N. J., **2000**, Chen, W., pp449-472)

In whole animals, studies in the rat sciatic nerve crush model have demonstrated that α -MSH increases neurite outgrowth and, as the most potent of the ACTH derived peptides, it significantly promotes nerve terminal 20 branching, endplate area, and perimeter. [Bijlsma, W. A., et al., *The Enhanced Recovery of Sensorimotor Function in Rats is Related to the Melanotropic Moiety of ACTH/MSH Neuropeptides*, *Eur. J. Pharmacol* , **1983**, 92, 231-236; Van der Neut. R., et al., *Stimulation by Melanocortins of Neurite Outgrowth from Spinal and Sensory Neurons In Vitro*, *Peptides*, **1992**, 13, 1109-1115; Van 25 Der Zee, C. E. E. M., et al., *α -MSH and Org 2766 in Peripheral Nerve Regeneration: Different Route of Delivery*, *Eur. J. Pharmacol.*, **1988**, 147, 351-357; Strand, F. L., et al., *Melanocortins as Factors in Somatic Neuromuscular Growth and Regrowth*, *Pharmac. Ther.*, **1994**, 62, 1-27]. Furthermore, recovery of motor function after nerve injury is shortened by application of α - 30 MSH and other melanocortins. [Strand, F. L., et al., *supra*]

Mice in which the MC4 receptor is rendered inactive by gene targeting become obese, suggesting that the MC4 receptor is involved in feeding. [Huszar, D., et al., *Targeted Disruption of the Melanocortin-4 Receptor Results in Mice*, *Cell*, **1997**, 88, 131-141] This is substantiated by a report that various 35 MC4 peptide agonists inhibit feeding behavior in *agouti* mice. [Fan, W., et al.,

5 immunohistochemistry and Western blot analysis (Thiboutot, D., Sivarajah, Gililand, K., Cong, Z. and Clawson, G., *J. Invest. Dermatol.* 115(4), 614-619, 2000).

10 Human sebum differs in its composition from other mammals. The main lipids in human sebum are triglycerides, wax esters and squalene (Greene, R.S., Downing, D.T., Poci, P.E., Strauss, J.S., *JID* 54, 240-247, 1970). Squalene, for instance is not found in many mammals with the exception of otter and beaver. Triglyceride, which is a major component of human sebum is poorly represented in other species and in many (e.g. chimpanzee) appears to be totally absent (Thody, A.J., Shuster, S., *Physiolog. Rev.* 69, 383-415, 1989).
15 Moreover melanocortins can have different effects on cells from different species. For example both α MSH ($EC_{50}=3.7\text{nM}$) and ACTH ($EC_{50}=16.4\text{nM}$) are potent lipolytic agents for rabbit adipocytes, whereas in the rat only ACTH ($EC_{50}=1.34\text{nM}$) has potent lipolytic activity (Ramachadran, J., Lee, V., 428, 339-346, 1987; Richter, W. O., Schwandt, P., *Neuropeptides* 9, 59-74, 1987).
20 Despite lipolytic activity in rodents and rabbits, ACTH has very little effect on lipolysis in isolated human and non-human primate adipocytes, even at concentrations as high as $1\mu\text{M}$ (Ng, T.B. *Comparative Biochem.* 97, 441-446, 1990). Thus defining the role of melanocortins and their receptors in animal
25 sebaceous model systems is not necessarily predictive of their role in a human sebaceous lipid regulation.

30 Recently, Basu et.al., in WIPO publication WO99/55679 disclosed isoquinoline derivatives, small molecule non-peptide compounds, which showed low micromolar affinities for the MC1 and MC4 receptors, reduction of dermal inflammation induced by arachidonic acids, and reductions of body weight and food intake.

35 Nargund et. al., in WIPO publication WO99/64002 disclosed spiroperidine derivatives as melanocortin receptor agonists, useful for the treatment of diseases and disorders such as obesity, diabetes and sexual dysfunction.


5

Thus there exist a need for small molecule modulators of the melanocortin receptor, more particularly the melanocortin-3, melanocortin-4 and / or the melanocortin-5 receptors.

10

SUMMARY OF THE INVENTION

The present invention is directed to compounds of the general formula (II)



(II)

wherein

R¹ is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl; (R¹ is not unsubstituted aryl or unsubstituted aralkyl);

15

wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

20

wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

R² is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl; (R² is not unsubstituted aryl or unsubstituted aralkyl);

25

wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

30


5 wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 R⁴ is selected from the group consisting of aryl, aralkyl, heteroaryl, heterocycloalkyl, and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or
10 more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 provided that when R¹ is aryl substituted with one halogen and R⁴ is aryl substituted with one halogen, then R² is not morpholinyl;

15 and pharmaceutically acceptable salts thereof.

The present invention is further directed to compounds of the general formula (II)



(II)

20 wherein

 R¹ is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl; (R¹ is not unsubstituted aryl or unsubstituted aralkyl);

 wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or
25 more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy;
30 halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 R² is selected from the group consisting of aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl;

5 wherein the aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

10 R^4 is selected from the group consisting of heteroaryl, heterocycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl; (R^4 is not unsubstituted aryl or unsubstituted aralkyl);

 wherein the heteroaryl, heterocycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from
15 halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

20 provided that when R^1 is aryl substituted with one halogen and R^4 is aryl substituted with one halogen, then R^2 is not morpholinyl;


 provided further that when R^1 is methylphenyl and R^4 is methoxyphenyl, then R^2 is selected from the group consisting of aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl and
25 substituted aryl;

 wherein the aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl group is optionally substituted with one or more substituents independently selected from
 halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

30 wherein the aryl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 and pharmaceutically acceptable salts thereof.

35 The present invention is further directed to a method for treating disorders mediated by a melanocortin receptor comprising administering to a subject in need thereof a therapeutically effective amount of a compound of formula (II)



(II)

5

wherein

R¹ is selected from the group consisting of alkyl, aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

10

15

R² is selected from the group consisting of alkyl, aryl, aralkyl, heteroaryl, heterocycloalkyl and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

20

R⁴ is selected from the group consisting of hydrogen, alkyl, aryl, aralkyl, heteroaryl, heterocycloalkyl, and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

25

and pharmaceutically acceptable salts thereof.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. An illustration of the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier. Illustrating the invention is a process for making a

30

5 pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

Exemplifying the invention are methods of treating disorders mediated by the melanocortin receptor in a subject in need thereof comprising
10 administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

An embodiment of the present invention is the use of any of the compounds described herein for the treatment of a disorder selected from the
15 group consisting of metabolic disorders, CNS disorders and dermatological disorders.


An example of the invention is a method for treating a disorder selected from the group consisting of obesity, impaired oral glucose tolerance, elevated
20 blood glucose levels, type II diabetes, Syndrome X, diabetic retinopathy, spinal cord injury, nerve injury, acute neurodegenerative disorders, chronic neurodegenerative disorders, plexopathies, male erectile dysfunction, dry eyes, acne, dry skin, aged skin, seborrheic dermatitis, rosacea, excessive ear wax, meibomian gland disorder, pseudofolliculitis, yeast infections, dandruff,
25 hidradenitis suppurativa, ocular rosacea and eccrine gland disorder in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

30 Another example of the invention is the use of any of the compounds described herein in the preparation of a medicament for treating: (a) obesity, (b) impaired oral glucose tolerance, (c) elevated blood glucose levels, (d) type II diabetes, (e) Syndrome X, (f) diabetic retinopathy, (g) an acute neurodegenerative disorder, (h) a chronic neurodegenerative disorder, (i) a
35 plexopathy, (j) male erectile dysfunction, (k) dry eyes, (l) acne, (m) dry skin, (n) aged skin, (o) seborrheic dermatitis, (p) rosacea, (q) excessive ear wax, (r) meibomian gland disorder, (s) pseudofolliculitis, (t) yeast infections, (u)

5 dandruff, (v) hidradenitis suppurativa, (w) ocular rosacea or (x) eccrine gland disorder, in a subject in need thereof.

DETAILED DESCRIPTION OF THE INVENTION

10 The present invention is directed to novel substituted 1,2,4-thiadiazole derivatives useful for the treatment of disorders mediated by a melanocortin receptor. More particularly, the present invention is directed to compounds of formula (II)



(II)

15 wherein R¹, R² and R⁴ are as herein defined, useful as melanocortin receptor agonists or antagonists.

The present invention is further directed to a method of treating a disorder mediated by a melanocortin receptor, preferably a disorder which is susceptible to treatment by agonism or antagonism of a melanocortin receptor.

20 Preferably the melanocortin receptor is selected from the group consisting of the melanocortin-3, melanocortin-4 and melanocortin-5 receptor, more preferably the melanocortin receptor is melanocortin-4 or melanocortin-5.

25 In an embodiment of the present invention, R¹ is selected from the group consisting of aryl, aralkyl and heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino. Preferably, R¹ is selected from the group consisting of aryl; wherein aryl group is optionally substituted with one or more
30 substituents independently selected from halogen, alkyl and alkoxy. More preferably, R¹ is selected from the group consisting of phenyl, 2-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 4-methylphenyl, 2-methoxyphenyl and 4-methoxyphenyl. Most preferably R¹ is 2-methoxyphenyl.

5

In another embodiment of the present invention R^1 is selected from the group consisting of substituted aryl and substituted aralkyl, wherein the aryl or aralkyl is substituted with one to two substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino. Preferably, R^1 is selected from the group consisting of substituted aryl; wherein the aryl group is substituted with a substituent selected from halogen, alkyl or alkoxy. More preferably, R^1 is 2-methoxyphenyl.

10

15

In yet another embodiment of the present invention, R^1 is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino.

20

25

In an embodiment of the present invention, R^2 is selected from the group consisting of aryl, aralkyl and heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino. Preferably, R^2 is selected from the group consisting of aryl; wherein the aryl group is optionally substituted with one or more substituents independently selected from alkyl and alkoxy. More preferably, R^2 is selected from the group consisting of phenyl, 4-methylphenyl, 2-methoxyphenyl and 4-methoxyphenyl. Most preferably, R^2 is selected from the group consisting of phenyl and 2-methoxyphenyl.

30

35

In another embodiment of the present invention R^2 is selected from the group consisting of substituted aryl and substituted aralkyl, wherein the aryl or aralkyl is substituted with one to two substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino. Preferably, R^2 is selected from the group consisting of substituted aryl; wherein the aryl group is substituted with a substituent selected from alkyl or alkoxy. More preferably, R^2 is 2-methoxyphenyl.

5

In yet another embodiment of the present invention, R^2 is selected from the group consisting of aryl and aralkyl; wherein the aryl or aralkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino.

10

In an embodiment of the present invention, R^4 is selected from the group consisting of aryl, aralkyl and heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino. Preferably, R^4 is selected from the group consisting of aryl, aralkyl, and heteroaryl; wherein the aryl or aralkyl group is optionally substituted with one or more substituents independently selected from halogen, alkyl and alkoxy. More preferably, R^4 is selected from the group consisting of phenyl, 2-chlorophenyl, 4-chlorophenyl, 4-bromophenyl, 2-methylphenyl, 4-methylphenyl, 2-methoxyphenyl, 4-methoxyphenyl, benzyl, 2-chlorobenzyl, 4-chlorobenzyl, 2-methylbenzyl, 4-methylbenzyl, 2-methoxybenzyl, 4-methoxybenzyl, 2,6-difluorophenyl, 3,5-difluorophenyl, 2-chloro-6-methylphenyl and 3-pyridyl. Most preferably, R^4 is selected from the group consisting of phenyl, 2-methylphenyl, 4-methylphenyl, 2-methoxyphenyl and 4-methoxyphenyl.

15

20

25

In another embodiment of the present invention R^4 is selected from the group consisting of aryl, aralkyl or heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino, di(alkyl)amino. Preferably, R^4 is selected from the group consisting of aryl and substituted aryl; wherein the substituents on the aryl are one to two independently selected halogen, alkyl or alkoxy. More preferably, R^4 is selected from the group consisting of phenyl, 4-methoxyphenyl, 2,6-difluorophenyl, 2-chloro-6-methylphenyl and 3,5-difluorophenyl.

30

35

5 In yet another embodiment of the present invention, R⁴ is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino.

10

 As used herein, unless otherwise note, the term "disorders mediated by a melanocortin receptor" include, but are not limited to, obesity, impaired oral glucose tolerance, elevated blood glucose levels, type II diabetes, Syndrome X, diabetic retinopathy, acute neurodegenerative disorders, chronic
15 neurodegenerative disorders, plexopathies, male erectile dysfunction, dry eyes, acne, dry skin, aged skin, seborrheic dermatitis, rosacea, excessive ear wax, meibomian gland disorder, pseudofolliculitis, yeast infections, dandruff, hidradenitis suppurativa, ocular rosacea and eccrine gland disorder.

20

 As used herein, unless otherwise noted, the term "metabolic disorders" include, but are not limited to, obesity, impaired oral glucose tolerance, elevated blood glucose levels, type II diabetes and Syndrome X.

25

 As used herein, unless otherwise noted, the term "CNS disorders" include, but are not limited to, diabetic retinopathy, acute neurodegenerative disorders, chronic neurodegenerative disorders and plexopathies.

30

 As used herein, unless otherwise noted, the term "dermatological disorders" include, but are not limited to, dry eyes, acne, dry skin, aged skin, seborrheic dermatitis, rosacea, excessive ear wax, meibomian gland disorder, pseudofolliculitis, yeast infections, dandruff, hidradenitis suppurativa, ocular
rosacea and eccrine gland disorder.

35

 As used herein, acute neurodegenerative disorders include various types of acute neurodegenerative disorders associated with neuronal cell death or compromise including cerebrovascular insufficiency, focal or diffuse brain trauma, diffuse brain damage, and spinal cord injury, that is, cerebral ischemia

5 or infarction including embolic occlusion and thrombotic occlusion, reperfusion
following acute ischemia, perinatal hypoxic-ischemic injury, cardiac arrest, as
well as intracranial hemorrhage of any type (including, but not limited to,
epidural, subdural, subarachnoid and intracerebral), and intracranial and
intravertebral lesions (including, but not limited to, contusion, penetration,
10 shear, compression and laceration), and whiplash shaken infant syndrome.

As used herein, chronic neurodegenerative disorders included within the
methods of the present invention include Alzheimer's disease, Pick's disease,
diffuse Lewy body disease, progressive supranuclear palsy (Steel-Richardson
15 syndrome), multisystem degeneration (Shy-Drager syndrome), chronic epileptic
conditions associated with neurodegeneration, motor neuron diseases
including amyotrophic lateral sclerosis, degenerative ataxias, cortical basal
degeneration, ALS–Parkinson's–Dementia complex of Guam, subacute
sclerosing panencephalitis, Huntington's disease, Parkinson's disease,
20 synucleinopathies (including multiple system atrophy), primary progressive
aphasia, striatonigral degeneration, Machado-Joseph disease / spinocerebellar
ataxia type 3 and olivopontocerebellar degenerations, Gilles De La Tourette's
disease, bulbar and pseudobulbar palsy, spinal and spinobulbar muscular
atrophy (Kennedy's disease), primary lateral sclerosis, familial spastic
25 paraplegia, Werdnig-Hoffmann disease, Kugelberg-Welander disease, Tay-
Sach's disease, Sandhoff disease, familial spastic disease, Wohlfart-
Kugelberg-Welander disease, spastic paraparesis, progressive multifocal
leukoencephalopathy, familial dysautonomia (Riley-Day syndrome), and prion
diseases (including, but not limited to Creutzfeldt-Jakob, Gerstmann-
30 Sträussler-Scheinker disease, Kuru and fatal familial insomnia).

As used herein, plexopathies include plexus palsies, multifocal
neuropathies, sensory neuropathies, motor neuropathies, sensory-motor
neuropathies, infections neuropathies, autonomic neuropathies, sensory-
35 autonomic neuropathies, demyelinating neuropathies (including, but not limited
to Guillain-Barre syndrome and chronic inflammatory demyelinating
polyradiculoneuropathy), other inflammatory and immune neuropathies,

5 neuropathies induced by drugs, neuropathies induced by pharmacological
treatments, neuropathies induced by toxins, traumatic neuropathies (including,
but not limited to compression, crush, laceration and segmentation
neuropathies), metabolic neuropathies, endocrine and paraneoplastic
neuropathies, and other neuropathies such as Charcot-Marie-Tooth disease
10 (type 1a, 1b, 2, 4a, 1-X linked), Friedreich's ataxia, metachromatic
leukodystrophy, Refsum's disease, adrenomyeloneuropathy, Ataxia-
telangiectasia, Déjerine-Sottas neuropathy (types A and B), Lambert-Eaton
syndrome, and disorders of the cranial nerves.

15 As used herein, unless otherwise noted, the term "halogen" shall include
iodine, bromine, chlorine and fluorine.

 As used herein, the term "alkyl" whether used alone or as part of a
substituent group, include straight and branched chains comprising one to
20 eight carbon atoms. For example, alkyl radicals include methyl, ethyl, propyl,
isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl and the like. Unless
otherwise noted, "lower" when used with alkyl means a carbon chain
composition of one to four carbon atoms.

25 The term "alkenyl", whether used alone or as part of a substituent group,
shall include straight and branched alkene chains comprising two to eight
carbon atoms. Suitable examples include vinyl, 1-propenyl, 2-propenyl, 1-
butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 1-isobut-2-enyl, and the like.
Similarly, the term "alkynyl", whether used alone or as part of a substituent
30 group, shall include straight and branched alkyne chains comprising two to
eight carbon atoms. Suitable examples include 2-propynyl, 2-butylyl, 1-
butynyl, 1-pentylyl, and the like.

 As used herein, unless otherwise noted, "alkoxy" shall denote an oxygen
35 ether radical of the above described straight or branched chain alkyl groups. For
example, methoxy, ethoxy, n-propoxy, sec-butoxy, t-butoxy, n-hexyloxy and the
like.

5

As used herein, unless otherwise noted, the term "cycloalkyl" shall denote saturated monocyclic ring structures comprising three to eight ring carbons, preferably 5 to 7 carbons. Suitable examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

10

As used herein, the term "aryl" indicates aromatic carbocyclic ring structures such as phenyl, naphthyl, and the like.

15

As used herein, unless otherwise noted, "aralkyl" shall mean any lower alkyl group substituted with an aryl group. For example, benzyl, phenylethyl, phenylpropyl, naphthylmethyl, and the like.

20

As used herein, unless otherwise noted, "heteroaryl" shall denote any five or six membered monocyclic aromatic ring structure containing at least one heteroatom selected from the group consisting of O, N and S, optionally containing one to three additional heteroatoms independently selected from the group consisting of O, N and S; or a nine or ten membered bicyclic aromatic ring structure containing at least one heteroatom selected from the group consisting of O, N and S, optionally containing one to four additional heteroatoms independently selected from the group consisting of O, N and S. The heteroaryl group may be attached at any carbon atom of the ring such that the result is a stable structure.

25

30

Examples of suitable heteroaryl groups include, but are not limited to, pyrrolyl, furyl, thienyl, oxazolyl, imidazolyl, purazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyranyl, furazanyl, indoliziny, indolyl, isoindoliny, indazolyl, isoxazolyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinoliziny, quinolinyl, isoquinolinyl, isothiazolyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, and the like. Preferred heteroaryl groups include pyridyl, thienyl and imidazolyl.

35

5 As used herein, the term "heterocycloalkyl" shall denote any five to seven
membered monocyclic, saturated, partially unsaturated or partially aromatic ring
structure containing at least one heteroatom selected from the group consisting
of O, N and S, optionally containing one to three additional heteroatoms
independently selected from the group consisting of O, N and S; or a nine to ten
10 membered saturated, partially unsaturated or partially aromatic bicyclic ring
system containing at least one heteroatom selected from the group consisting of
O, N and S, optionally containing one to four additional heteroatoms
independently selected from the group consisting of O, N and S. The
heterocycloalkyl group may be attached at any carbon atom of the ring such that
15 the result is a stable structure.

Examples of suitable heterocycloalkyl groups include, but are not limited
to, pyrrolinyl, pyrrolidinyl, dioxalanyl, imidazoliny, imidazolidinyl, pyrazolinyl,
pyrazolidinyl, piperidinyl, dioxanyl, morpholinyl, dithianyl, thiomorpholinyl,
piperazinyl, trithianyl, indolinyl, chromenyl, 3,4-methylenedioxyphenyl, 2,3-
20 dihydrobenzofuryl, and the like.

As used herein, the notation "*" shall denote the presence of a
stereogenic center.


25 Where the compounds according to this invention have at least one
chiral center, they may accordingly exist as enantiomers. Where the
compounds possess two or more chiral centers, they may additionally exist as
diastereomers. It is to be understood that all such isomers and mixtures
thereof are encompassed within the scope of the present invention.
30 Furthermore, some of the crystalline forms for the compounds may exist as
polymorphs and as such are intended to be included in the present invention.
In addition, some of the compounds may form solvates with water (i.e.,
hydrates) or common organic solvents, and such solvates are also intended to
be encompassed within the scope of this invention.

35 When a particular group is "substituted" (e.g., cycloalkyl, aryl, aralkyl,
heteroaryl, heterocycloalkyl), that group may have one or more substituents,

5 preferably from one to five substituents, more preferably from one to three substituents, most preferably from one to two substituents, independently selected from the list of substituents.

10 It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth
15 herein.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. Thus, for example, a
20 "phenylC₁-C₆alkylaminocarbonylC₁-C₆alkyl" substituent refers to a group of the formula



25 The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who is or has been the object of treatment, observation or experiment.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well
30 as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or

5 medicinal response in a tissue system, animal or human that is being sought by
a researcher, veterinarian, medical doctor or other clinician, which includes
alleviation of the symptoms of the disease or disorder being treated.

10 For use in medicine, the salts of the compounds of this invention refer to
non-toxic "pharmaceutically acceptable salts". Other salts may, however, be
useful in the preparation of compounds according to this invention or of their
pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts
of the compounds of this invention include acid addition salts which may, for
15 example, be formed by mixing a solution of the compound with a solution of a
pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid,
fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid,
tartaric acid, carbonic acid or phosphoric acid.

20 The present invention includes within its scope prodrugs of the
compounds of this invention. In general, such prodrugs will be functional
derivatives of the compounds which are readily convertible *in vivo* into the
required compound. Thus, in the methods of treatment of the present
invention, the term "administering" shall encompass the treatment of the
various disorders described with the compound specifically disclosed or with a
25 compound which may not be specifically disclosed, but which converts to the
specified compound *in vivo* after administration to the patient. Conventional
procedures for the selection and preparation of suitable prodrug derivatives are
described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier,
1985.


30

Abbreviations used in the instant specification, particularly the Schemes
and Examples, are as follows:

BHT	=	2,6-bis-(<i>t</i> -butyl)-4-methyl-phenol
BSA	=	Bovine Serum Albumin
cAMP or cyclic AMP	=	Cyclic-adenosine monophosphate

DCE	=	1,2-dichloroethane
DEAD	=	Diethyl azodicarboxylate
DM	=	Differentiation Medium
DMF	=	Dimethyl formamide
DMEM	=	Dulbeccos Minimal Essential Medium
DMSO		Dimethylsulfoxide
DPBS	=	Dulbeccos phosphate buffered saline
EDTA	=	Ethylene Diamine Tetraacetic Acid
FBS	=	Fetal bovine serum
GDP	=	Guanosine Diphosphate
GTP	=	Guanosine Triphosphate
GM	=	Growth Medium
HBSS	=	Hank's Buffered salt Solution
HEPES	=	4-(2-Hydroxyethyl)-1-piperazine ethane sulfonic acid
HS	=	Human Serum
IgG	=	Immunoglobulin G
% Inh	=	Percent Inhibition
MEM	=	Minimum Essential Medium
NBS	=	N-bromosuccinimide
NCS	=	N-chlorosuccinimide
NDP α MSH	=	[Nle ⁴ , D-Phe ⁷] α MSH, an analog of α MSH
PBS	=	Phosphate Buffered saline
PEG	=	Polyethylene Glycol
PNC	=	Penicillin
rt or RT	=	Room Temperature
SPA	=	Scintillation Proximity Assay
STM	=	Streptomycin
TLC	=	Thin layer chromatography
TM	=	Transition Medium
TMS	=	Trimethylsilyl

5 Compounds of formula (I) wherein R³ is hydrogen may be prepared according to the process outlined in Scheme 1.




Scheme 1

10 More particularly, a suitably substituted cyano compound of formula (III), a known compound or compound prepared by known methods, is reacted with a suitably substituted primary amine of formula (IV), a known compound or compound prepared by known methods, in the presence of a base such as $NaNH_2$, NaH , $NaN(TMS)_2$, and the like, preferably $NaNH_2$, at an elevated temperature, preferably at about reflux, to yield the corresponding compound of formula (V).

15 The compound of formula (V) is reacted with a suitably substituted thiocyanate of formula (VI), a known compound or compound prepared by known methods, in the presence of DCE, at an elevated temperature, preferably about 45°C, to yield the corresponding compound of formula (VII).

20 The compound of formula (VII) is subjected to ring closure/oxidation, in the presence of Br_2 , at room temperature, to yield the corresponding compound of formula (Ia).

25 Compounds of formula (II) may be prepared from suitably substituted compounds of formula (I) wherein R³ is hydrogen according to the process outlined in Scheme 2.




5

Scheme 2

More particularly, a suitably substituted compound of formula (Ia) is treated with a base such as NaHCO_3 , Na_2CO_3 , NaOH , and the like, preferably NaHCO_3 , at room temperature, to yield the corresponding compound of formula (II).

10

Compounds of formula (II) may also be prepared from suitably substituted compounds of formula (VII) according to the process outlined in Scheme 3.



15


Scheme 3

More particularly, a suitably substituted compound of formula (VII) is reacted with an oxidizing agent such as NBS, NCS, DEAD, and the like, preferably NBS, at room temperature, to yield the corresponding compound of formula (II). Preferably, the compound of formula (II) is extracted from a basic aqueous solution such as NaHCO_3 , Na_2CO_3 , NaOH , and the like.

20

Compounds of formula (I) wherein R^3 is alkyl may be prepared from a suitably substituted compound of formula (II) according to the process outlined in Scheme 4.

25




Scheme 4

Accordingly, a suitably substituted compound of formula (II) is reacted with a suitably substituted compound of formula (VIII), a known compound or compound prepared by known methods, wherein X^- is trifluoromethylsulfonate, Br^- , and I^- , at room temperature, to yield the corresponding compound of formula (I).

10

Compounds of formula (I) wherein R^3 is hydrogen may be prepared from a suitably substituted compound of formula (II) according to the process outlined in Scheme 5.

15



Scheme 5

Accordingly, a suitably substituted compound of formula (II) is reacted with a pharmaceutically acceptable acid such as HCl, HBr, HNO_3 , and the like, preferably HCl, at room temperature, to yield the corresponding compound of formula (Ia), wherein X^- is Cl^- , Br^- , NO_3^- , and the like.

20

Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt

25

5 formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid
and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and
regeneration of the free base. The compounds may also be resolved by
formation of diastereomeric esters or amides, followed by chromatographic
10 separation and removal of the chiral auxiliary. Alternatively, the compounds
may be resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the
present invention, it may be necessary and/or desirable to protect sensitive or
reactive groups on any of the molecules concerned. This may be achieved by
15 means of conventional protecting groups, such as those described in
Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press,
1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic
Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed
at a convenient subsequent stage using methods known from the art.

20 The utility of the compounds of the present invention to treat disorders
mediated by a melanocortin receptor can be determined according to the
procedures described in Examples 4-11 herein. The present invention
therefore provides a method of treating such disorders, which comprises
25 administering any of the compounds as defined herein in a quantity effective to
treat the disorder (i.e. in a therapeutically effective amount). The compound
may be administered to a patient afflicted with such a disorder by any
conventional route of administration, including, but not limited to, intravenous,
oral, subcutaneous, intramuscular, intradermal, parenteral and transdermal.

30 The present invention also provides pharmaceutical compositions
comprising one or more compounds of this invention in association with a
pharmaceutically acceptable carrier.

35 To prepare the pharmaceutical compositions of this invention, one or
more compounds of formula (I) and/or (II) or a salt thereof (the active
ingredient), is intimately admixed with a pharmaceutical carrier according to

5 conventional pharmaceutical compounding techniques, which carrier may take
a wide variety of forms depending of the form of preparation desired for
administration, e.g., oral or parenteral such as intramuscular. In preparing the
compositions in oral dosage form, any of the usual pharmaceutical media may
be employed. Thus, for liquid oral preparations, such as for example,
10 suspensions, elixirs and solutions, suitable carriers and additives include water,
glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the
like; for solid oral preparations such as, for example, powders, capsules,
caplets, gelcaps and tablets, suitable carriers and additives include starches,
sugars, diluents, granulating agents, lubricants, binders, disintegrating agents
15 and the like. Because of their ease in administration, tablets and capsules
represent the most advantageous oral dosage unit form, in which case solid
pharmaceutical carriers are obviously employed. If desired, tablets may be
sugar coated or enteric coated by standard techniques. For parenterals, the
carrier will usually comprise sterile water, though other ingredients, for
20 example, for purposes such as aiding solubility or for preservation, may be
included. Injectable suspensions may also be prepared, in which case
appropriate liquid carriers, suspending agents and the like may be employed.

25 Topical formulations included within the present invention, include but
are not limited to creams, lotions, multiple emulsions, microemulsions,
liposomal creams or gels, gels, solutions, suspensions, ointments, foaming
aerosols, hard or soft gelatin capsules, masks, sticks, roll-ons, powders, spray
forms, and the like. The topical formulations may contain, in addition to the
active ingredient(s), one or more non-active components including, but are not
30 limited to chelating agents, buffering agents, colorants, preservatives,
fragrances, emulsifiers, surfactants, opacifying agents, emollients, solvents,
sunscreens, viscosity modifying agents, antioxidants, moisturizers,
permeations enhancers, film formers, and the like.

35 Topical formulations for acne treatment included within the present
invention may also contain one or more of the following components, including
comedolytic/keratolytic agents, antimicrobial agents and steroidal or non-

5 steroidal anti-inflammatory agents. (Comedolytic agents refer to any
compound capable of rupturing a comedo. Keratolytic agents refer to any
compound capable of breaking apart keratinocytes resulting in exfoliation of the
epidermis.) Suitable comedolytic/keratolytic agents include, but are not limited
to retinoids, salicylic acid, glycolic acid, cetyl betaine, and the like. Suitable
10 antimicrobial agents include, but are not limited to benzoyl peroxide,
erythromycin, tetracycline, clindamycin, azelaic acid, and the like. Topical
formulations typically contain 0.01-1% active ingredient.

 The pharmaceutical compositions herein will contain, per dosage unit,
15 e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of
the active ingredient necessary to deliver an effective dose as described
above. The non-topical pharmaceutical compositions herein will contain, per
unit dosage unit, e.g., tablet, capsule, powder, injection, suppository,
teaspoonful and the like, of from about 0.03 mg to 100 mg/kg (preferred 0.1-30
20 mg/kg) and may be given at a dosage of from about 0.1-300 mg/kg/day
(preferred 1-50 mg/kg/day). The dosages, however, may be varied depending
upon the requirement of the patients, the severity of the condition being treated
and the compound being employed. The use of either daily administration or
post-periodic dosing may be employed.

25 Preferably these compositions are in unit dosage forms from such as
tablets, pills, capsules, powders, granules, sterile parenteral solutions or
suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector
devices or suppositories; for oral parenteral, intranasal, sublingual or rectal
30 administration, or for administration by inhalation or insufflation. Alternatively,
the composition may be presented in a form suitable for once-weekly or once-
monthly administration; for example, an insoluble salt of the active compound
such as the decanoate salt, may be adapted to provide a depot preparation for
intramuscular injection. For preparing solid compositions such as tablets, the
35 principal active ingredient is mixed with a pharmaceutical carrier, e.g.
conventional tableting ingredients such as corn starch, lactose, sucrose,
sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums,

5 and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the
10 composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise
15 compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component
20 to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

25 The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles.
30 Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

35 The method of treating disorders mediated by a melanocortin receptor described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein

5 and a pharmaceutically acceptable carrier. The non-topical pharmaceutical composition may contain between about 0.01 mg and 100 mg, preferably about 5 to 50 mg, of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending
10 agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions. Forms useful for
15 parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore,
20 compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

25 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents
30 and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include,
35 without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

5 The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

10

 The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or

15 phosphatidylcholines.

 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with

20 soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl eneoxydipolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving

25 controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyeric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

30 Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of disorders mediated by a melanocortin receptor is required.

 The daily dosage of the products may be varied over a wide range from

35 0.01 to 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500

5 milligrams of the active ingredient for the symptomatic adjustment of the dosage
to the patient to be treated. An effective amount of the drug is ordinarily supplied
at a dosage level of from about 0.01 mg/kg to about 100 mg/kg of body weight
per day. Preferably, the range is from about 0.03 to about 10 mg/kg of body
weight per day. The compounds may be administered on a regimen of 1 to 4
10 times per day.


Optimal dosages to be administered may be readily determined by those
skilled in the art, and will vary with the particular compound used, the mode of
administration, the strength of the preparation, the mode of administration, and
15 the advancement of the disease condition. In addition, factors associated with
the particular patient being treated, including patient age, weight, diet and time of
administration, will result in the need to adjust dosages.

The following Examples are set forth to aid in the understanding of the
20 invention, and are not intended, and should not be construed to limit in any way
the invention set forth in the claims which follow thereafter.

EXAMPLE 1

25 2-(2-methoxyphenyl)-3-phenyl-5-*p*-tolylamino-[1,2,4]thiadiazol-2-ium

Compound # 31



STEP A:

30 A mixture of *o*-anisidine (15.0 g, 121.8 mmol) and sodium amide (50 wt.
% suspension in toluene) (11.40 g, 146.2 mmol) in anhydrous toluene (200ml)
was stirred for 1 hour at room temperature. To the mixture was added

5 benzonitrile (9.86 ml, 96.6 mmol) and heated under reflux for 16 hours. The reaction mixture was cooled and 1.0N HCl (150 ml) was added to quench the reaction. Activated carbon was added and the reaction mixture was filtered through a Celite pad. The pH of the mixture was adjusted to about 14 by addition of 1.0 N NaOH (200 ml). The aqueous layer was extracted with
10 chloroform (3x150 ml). The combined organic layer was dried over anhydrous MgSO₄, and evaporated. The resulting solid was washed with hexane and dried over the vacuum to yield the product as a pale white solid.

¹H NMR (300MHz, CDCl₃) δ 3.83 (s, 3H), 4.79 (s, 2H), 6.96(m, 3H), 7.04 (m, 1H), 7.43 (m, 3H), 7.93 (d, 2H)

15 MS (APCI, MH⁺) 227

STEP B:

A mixture of N-arylbenzamidine (3.0 g, 13.27mmol), prepared as in Step A, and 4-tolylisothiocyanate (2.18 g, 14.60) in anhydrous chloroform (30 ml)
20 was heated under reflux for 16 hours. The reaction mixture was cooled and the solvent was evaporated. The resulting residue was purified by flash column chromatography with a mobile phase of 25% hexane in dichloromethane. The combined fractions were evaporated; and the resulting solid was dried over vacuum to yield the product as a yellow solid.

25 ¹H NMR (300MHz, CDCl₃) δ 2.36 (s, 3H), 3.66 (s, 3H), 6.76 (m, 1H), 6.97 (t, 1H), 7.17-7.39 (m, 7H), 7.47 (d, 2H), 7.60 (d, 2H), 8.20 (s, 1H), 14.18 (s, 1H)

MS (ES, MH⁺) 376.29

30 STEP C:


To the solution of thiourea (2.90 g, 7.73 mmol), prepared as in STEP B, in anhydrous chloroform (15 ml) was slowly added bromine (438 μ l, 8.51 mmol). After stirring for 16 hours, the solvent was evaporated. The resulting solid was washed with anhydrous ethyl ether. The crude product was re-
35 crystallized from 20% water in ethanol to yield the product as a yellow solid.

¹H NMR (300MHz, CDCl₃) δ 2.39 (s, 3H), 3.66 (s, 3H), 6.96 (d, 1H), 7.06 (t, 1H), 7.20 - 8.07 (m, 7H), 7.61 (d, 2H), 7.80 (d, 2H), 12.40 (s, 1H)

5 MS (ES, MH⁺) 374.25

EXAMPLE 2

2-(2-methoxyphenyl)-3-(2-methoxyphenyl)-5-phenylamino-[1,2,4]thiadiazol-2-ium Compound #74



STEP A:

A mixture of *o*-anisidine (13.5 g, 110.0 mmol) and sodium amide (50 wt. % suspension in toluene) (9.40 g, 120.0 mmol) in anhydrous toluene (200ml) was stirred for 1 hour at room temperature. To the mixture was added 2-methoxybenzotrile (16 ml, 131.0 mmol) and the reaction mixture was then heated under reflux for 16 hours. The reaction mixture was cooled and 1.0N HCl (150 ml) was added to quench the reaction. Activated carbon was added and the reaction mixture was filtered through a Celite pad. The pH of the mixture was adjusted to about 14 by addition of 1.0 N NaOH (200 ml). The aqueous layer was extracted with chloroform (3x150 ml). The combined organic layer was dried over anhydrous MgSO₄, and evaporated. The resulting solid was washed with hexane and dried over the vacuum to yield the product as a pale white solid.

15

20

MS (APCI, MH⁺) 257

25

STEP B:

A mixture of *N*-arylbenzamidine (15.5 g, 60.6 mmol), prepared as in STEP A and phenylisothiocyanate (8.70 mL, 72.7 mmol) in anhydrous chloroform (30 ml) was heated at 45°C for 16 hours. The reaction mixture was cooled and the solvent was evaporated. The resulting residue was purified by

30

5 flash column chromatography with a mobile phase of 25% hexane in dichloromethane. The combined fractions were evaporated and the resulting solid was dried over vacuum to yield the product as a yellow solid.

MS (ES, MH⁺) 391.50

10 STEP C:

To a solution of thiourea (12.95 g, 33.1 mmol), prepared as in STEP B, in anhydrous chloroform (15 ml) was slowly added bromine (1.78 mL, 34.75 mmol). After stirring for 16 hours, the solvent was evaporated. The resulting solid was washed with anhydrous ethyl ether. The crude product was re-


15 crystallized from 20% water in ethanol to yield the product as a yellow solid.

MS (ES, MH⁺) 390.1

Following the procedures disclosed herein, representative compounds of formula (I) of the present invention were prepared, as listed in Table 1.

20

TABLE 1


				
ID No.	R ¹	R ²	R ³	R ⁴
1	4-methylphenyl	phenyl	H	4-methoxyphenyl
2	4-methylphenyl	phenyl	H	phenyl
3	4-methylphenyl	phenyl	H	2-methoxyphenyl
4	4-methylphenyl	phenyl	H	4-methylphenyl
5	4-methylphenyl	phenyl	H	2-methylphenyl
6	4-methylphenyl	phenyl	H	4-chlorophenyl
7	4-methylphenyl	phenyl	H	2-chlorophenyl
8	2-methylphenyl	phenyl	H	phenyl
9	2-methylphenyl	phenyl	H	4-methoxyphenyl

10	2-methylphenyl	phenyl	H	2-methoxyphenyl
11	2-methylphenyl	phenyl	H	4-methylphenyl
12	2-methylphenyl	phenyl	H	2-methylphenyl
13	2-methylphenyl	phenyl	H	4-chlorophenyl
14	2-methylphenyl	phenyl	H	2-chlorophenyl
15	4-chlorophenyl	phenyl	H	phenyl
16	4-chlorophenyl	phenyl	H	4-methoxyphenyl
17	4-chlorophenyl	phenyl	H	2-methoxyphenyl
18	4-chlorophenyl	phenyl	H	4-methylphenyl
19	4-chlorophenyl	phenyl	H	2-methylphenyl
20	4-chlorophenyl	phenyl	H	4-chlorophenyl
21	4-chlorophenyl	phenyl	H	2-chlorophenyl
22	phenyl	phenyl	H	phenyl
23	phenyl	phenyl	H	4-methoxyphenyl
24	phenyl	phenyl	H	2-methoxyphenyl
25	phenyl	phenyl	H	4-methylphenyl
26	phenyl	phenyl	H	2-methylphenyl
27	phenyl	phenyl	H	4-chlorophenyl
28	2-methoxyphenyl	phenyl	H	phenyl
29	2-methoxyphenyl	phenyl	H	4-methoxyphenyl
30	2-methoxyphenyl	phenyl	H	2-methoxyphenyl
31	2-methoxyphenyl	phenyl	H	4-methylphenyl
32	2-methoxyphenyl	phenyl	H	2-methylphenyl
33	2-methoxyphenyl	phenyl	H	4-chlorophenyl
34	2-methoxyphenyl	phenyl	H	2-chlorophenyl
35	4-methoxyphenyl	phenyl	H	phenyl
36	4-methoxyphenyl	phenyl	H	4-methoxyphenyl

37	4-methoxyphenyl	phenyl	H	2-methoxyphenyl
38	4-methoxyphenyl	phenyl	H	4-methylphenyl
39	4-methoxyphenyl	phenyl	H	2-methylphenyl
40	4-methoxyphenyl	phenyl	H	4-chloro phenyl
41	4-methoxyphenyl	phenyl	H	2-chlorophenyl
42	2-chlorophenyl	phenyl	H	phenyl
43	2-chlorophenyl	phenyl	H	4-methoxyphenyl
44	2-chlorophenyl	phenyl	H	2-methoxyphenyl
45	2-chlorophenyl	phenyl	H	4-methylphenyl
46	2-chlorophenyl	phenyl	H	2-methylphenyl
47	2-chlorophenyl	phenyl	H	4-chlorophenyl
48	2-chlorophenyl	phenyl	H	2-chlorophenyl
49	phenyl	phenyl	H	2-chlorophenyl
50	4-methylphenyl	phenyl	CH ₃	4-methoxyphenyl
51	phenyl	phenyl	H	4-methoxybenzyl
52	phenyl	phenyl	H	2-methoxybenzyl
53	phenyl	phenyl	H	4-methylbenzyl
54	phenyl	phenyl	H	2-methylbenzyl
55	phenyl	phenyl	H	4-chlorobenzyl
56	phenyl	phenyl	H	2-chlorobenzyl
57	4-methoxyphenyl	4-methoxyphenyl	H	phenyl
58	4-methoxyphenyl	4-methoxyphenyl	H	4-methoxyphenyl
59	4-methoxyphenyl	4-methoxyphenyl	H	2-methoxyphenyl
60	4-methoxyphenyl	4-methoxyphenyl	H	4-methylphenyl
61	4-methoxyphenyl	4-methoxyphenyl	H	2-methylphenyl
62	4-methoxyphenyl	4-methoxyphenyl	H	4-chlorophenyl

63	4-methoxyphenyl	4-methoxyphenyl	H	2-chlorophenyl
64	4-methoxyphenyl	4-methoxyphenyl	H	3-pyridyl
65	2-methoxyphenyl	4-methylphenyl	H	phenyl
66	2-methoxyphenyl	4-methylphenyl	H	4-methoxyphenyl
67	2-methoxyphenyl	4-methylphenyl	H	2-methoxyphenyl
68	2-methoxyphenyl	4-methylphenyl	H	4-methylphenyl
69	2-methoxyphenyl	4-methylphenyl	H	2-methylphenyl
70	2-methoxyphenyl	4-methylphenyl	H	4-chlorophenyl
71	2-methoxyphenyl	4-methylphenyl	H	2-chlorophenyl
72	2-methoxyphenyl	4-methylphenyl	H	3-pyridyl
73	2-methoxyphenyl	phenyl	H	3-pyridyl
74	2-methoxyphenyl	2-methoxyphenyl	H	phenyl
75	2-methoxyphenyl	2-methoxyphenyl	H	4-methoxyphenyl
76	2-methoxyphenyl	2-methoxyphenyl	H	2-methoxyphenyl
77	2-methoxyphenyl	2-methoxyphenyl	H	4-methylphenyl
78	2-methoxyphenyl	2-methoxyphenyl	H	2-methylphenyl
79	2-methoxyphenyl	2-methoxyphenyl	H	4-chlorophenyl
80	2-methoxyphenyl	2-methoxyphenyl	H	2-chlorophenyl
81	2-methoxyphenyl	2-methoxyphenyl	H	3-pyridyl
84	phenyl	phenyl	H	benzyl
85	2-methoxyphenyl	2-methoxyphenyl	H	4-bromophenyl
86	2-methoxyphenyl	2-methoxyphenyl	H	2,6-difluoro phenyl
87	2-methoxyphenyl	2-methoxyphenyl	H	2-chloro-6-methylphenyl
88	2-methoxyphenyl	2-methoxyphenyl	H	3,5-difluoro phenyl

EXAMPLE 3

5 2-(2-methoxyphenyl)-3-(2-methoxyphenyl)-5-phenylamino-1,2,4-thiadiazoleCompound #89

To a solution of the compound prepared as in Step B of Example 2 (0.921 g, 2.36 mmol) in anhydrous chloroform (10 mL) was added N-chlorosuccinimide (326 mg, 2.71 mmol). The reaction mixture was then stirred for 16 hours, and then stopped and washed twice with aqueous NaHCO₃. The organic layer was dried over magnesium sulfate and the remaining solvent removed under vacuum to yield the title product as a solid.

MS (ES, MH⁺) 390.1

Following the procedures described herein, representative compounds of formula (II) of the present invention were prepared, as listed in Table 2.

Table 2

ID No.	R ¹	R ²	R ⁴
82	4-methylphenyl	phenyl	phenyl
83	4-methylphenyl	phenyl	4-methoxyphenyl
89	2-methoxyphenyl	2-methoxyphenyl	phenyl
90	2-methoxyphenyl	2-methoxyphenyl	2,6-difluoro phenyl
91	2-methoxyphenyl	2-methoxyphenyl	2-chloro-6-methylphenyl
92	2-methoxyphenyl	2-methoxyphenyl	3,5-difluoro phenyl

- 5 Unless otherwise noted, NMR spectra, were measured on a Bruker Avance 300 MHz NMR spectrometer. Unless otherwise noted, molecular weights were measured using a Micromass LC platform electrospray mass spectrometer, as listed in Table 3.

10

TABLE 3

ID No	MW 1 (as +ion)	MW 2 (w/Br-)	Meas. MW
1	374.49	454.39	374.0
2	344.46	424.36	344.0
3	374.49	454.39	374.3
4	358.49	438.39	358.3
5	358.49	438.39	358.4
6	378.91	458.81	378.1, 380.3
7	378.91	458.81	378.2, 380.2
8	344.46	424.36	344.3
9	374.49	454.39	374.2
10	374.49	454.39	374.2
11	358.49	438.39	358.3
12	358.49	438.39	358.3
13	378.91	458.81	378.2, 380.2
14	378.91	458.81	378.2, 380.2
15	364.88	444.78	346.2, 366.2
16	394.90	474.81	394.1, 396.1
17	394.90	474.81	394.1, 396.1
18	378.91	458.81	378.2, 380.2
19	378.91	458.81	378.2, 380.2
20	399.32	479.23	398.1, 400.1
21	399.32	479.23	398.1, 400.0
22	330.43	410.34	330.3
23	360.46	440.37	360.3
24	360.46	440.37	360.3
25	344.46	424.37	344.4

26	344.46	424.37	344.3
27	364.88	444.78	364.2, 366.2
28	360.46	440.37	360.3
29	390.49	470.39	390.2
30	390.49	470.39	390.2
31	374.49	454.39	374.3
32	374.49	454.39	374.3
33	394.90	474.81	394.1, 396.1
34	394.90	474.81	394.1, 396.1
35	360.46	440.37	360.3
36	390.49	470.39	390.3
37	390.49	470.39	390.3
38	374.49	454.39	374.3
39	374.49	454.39	374.3
40	394.90	474.81	394.2, 396.2
41	394.90	474.81	394.2, 396.2
42	364.88	444.78	364.3, 366.3
43	394.90	474.81	394.2, 396.2
44	394.90	474.81	394.2, 396.2
45	378.91	458.81	378.2, 380.2
46	378.91	458.81	378.2, 380.2
47	399.32	479.23	398.1, 400.1
48	399.32	479.23	398.1, 400.1
49	364.88	444.78	364.2, 366.2
50	388.51	537.58	388.3
51	374.49	454.39	374.3
52	374.49	454.39	374.3
53	358.49	438.39	358.4
54	358.49	438.39	358.4
55	378.91	458.81	378.3, 380.3
56	378.91	458.81	378.3, 380.3
57	390.49	470.39	390.1

58	420.51	500.42	420.1
59	420.51	500.42	420.1
60	404.51	484.42	404.1
61	404.51	484.42	404.0
62	424.93	504.83	424.0, 426.0
63	424.93	504.83	424.0, 426.0
64	391.47	471.38	391.0
65	374.49	454.39	374.2
66	404.51	484.42	404.2
67	404.51	484.42	404.1
68	388.51	468.42	388.3
69	388.51	468.42	388.2
70	408.93	488.84	408.6, 410.0
71	408.93	488.84	408.1, 410.0
72	375.47	455.38	375.2
73	361.45	441.35	361.
74	390.49	470.39	390.2
75	420.51	500.42	420.2
76	420.51	500.42	420.2
77	404.51	484.42	404.1
78	404.51	484.42	404.1
79	424.93	504.84	424.1, 426.1
80	424.93	504.84	424.1, 426.1
81	391.47	471.38	391.1
82	343.45		344.3
83	373.48		374.3
84	344.46	424.36	344.3
85	469.38	549.29	
86	426.5	506.4	426.3
87	439.0	518.9	438.3, 440.3
88	426.5	506.4	426.2
89	389.5		390.1

90	425.5		426.2
91	437.9		438.3, 440.3
92	425.5		426.2

5

EXAMPLE 4Melanocortin MC-4 Receptor Binding Assay

Melanocortin [MC-4]-membrane [purchased from Receptor Biology Inc] was coupled to wheat germ agglutinin coated polyvinyl toluene-Scintillation Proximity Assay beads [purchased from Amersham Pharmacia Inc.] for 30 min at 25°C. Into each well of a 96-well Opti plate [purchased from Packard, CA], 2.5 µg of membrane and 0.25 mg of beads were mixed in a volume of 100 µL media. The media was 50mM HEPES, pH 7.4 containing 0.1 % bovine serum albumin, 2 mM CaCl₂, 2 mM MgCl₂ and protease inhibitors. Test compounds (1.5 µL) at 1mM in 30% DMSO-50mM HEPES, pH 7.4 buffer was added to separate wells on the plate. Radioactive ligand ¹²⁵I-NDP-melanocyte stimulating hormone [NEN, 2000Ci/mmol] was added to each well (48.5 µL per well, 40 pM final concentration). The plate was then sealed and let stand for 16 hr at 25°C. NDP-Melanocyte stimulating hormone peptide and α-melanocyte stimulating hormone peptide [purchased from Palomar Research Inc, 1 µM] were used as reference inhibitor compounds to define non-specific binding (N). Total binding (T) was defined using 30% DMSO-50 mM HEPES, pH 7.4 buffer. Bound radioactivity for each well (Y), measured at counts per minute (cpm) was measured in a TopCount [Packard, CA]. Percent inhibition was calculated as:

$$[(T - Y) / (T - N)] * 100\%$$

EXAMPLE 5Cyclic-Adenosine monophosphate [cAMP] Stimulation assay

Human Bowes melanoma cells expressing human melanocortin MC-4 receptor were grown to confluence in a 24-well culture plate. The growth medium was discarded and to each well was added 0.5 mL of Hank's solution. Test compounds were added to wells of a 96 well plate. NDP-melanocyte stimulating hormone peptide (1 µM) was added to the positive control wells

5 while negative control wells received vehicle of 30% DMSO-50mM HEPES, pH
7.4 buffer. The plate was incubated at 37°C and 5% CO₂ for 30 min. The
supernatant was discarded and the cells were washed twice with Hank's
solution. Ethanol (80%, 0.5 mL) was added to each well and the plates were
incubated at 4°C for 30 min. Cyclic AMP content was measured using the
10 NEN Flashplate kit [NEN]. A melanocortin receptor agonist is defined as a test
compound which resulted in an increase in cAMP production in this assay.

EXAMPLE 6

G-protein Activation Assay

15 For each assay, membranes expressing the melanocortin MC-4
receptor (5 µg) were incubated for 5 min at 25°C with 0.5 nM ³⁵S-GTPγS in 100
µL of 25 mM HEPES buffer, pH7.5 containing 100 mM NaCl, protease
inhibitors, 0.5 µM GDP, 5 mM 2-mercaptoethanol, 1 mM MgCl₂ together with
test compound, 1 µM of NDP-melanocyte stimulating hormone or a
20 combination of NDP-melanocyte stimulating hormone and test compound.
Basal ³⁵S-GTPγS binding was defined by 10mM HEPES, pH 7.4 buffer
containing 30% DMSO. The reaction was terminated by addition of 50 µl of
termination buffer containing 25 mM HEPES, pH7.5, 20 mM MgCl₂, protease
inhibitors, 100 µM GDP, 100 µM GTP, 5 mM 2-mercaptoethanol with
25 detergents (0.5 % digitonin, 0.2% sodium deoxycholate, and 0.5% NP-40).
The membranes were solubilized for 30 minutes at 25°C. The ³⁵S-GTPγS
bound Gαs protein was immunoprecipitated using anti-Gαs (0.5 µg) that linked
to anti-rabbit IgG protein A conjugated SPA. Bound radioactivity was
measured in a Topcount [Packard]. Non-specific ³⁵S-GTPγS binding was
30 defined by ³⁵S-GTPγS immunoprecipitated by normal rabbit IgG (0.5 µg).

Basal binding (B)	=	Mean counts/minute (cpm) immunoprecipitated by anti- Gαs.
Non-specific binding (NSB)	=	Mean cpm immunoprecipitated by normal rabbit IgG.
Specific basal binding (SB)	=	B-NSB

$$\begin{aligned} \text{Cpm in each well} &= C \\ \text{Net cpm in each well (N)} &= C - \text{NSB} \\ \% \text{ stimulation} &= [(N - \text{SB}) / \text{SB}] \times 100\% \end{aligned}$$

5

The procedures described above for the melanocortin MC-4 receptor were repeated for the melanocortin MC-3 receptor. Following the procedures described, representative compounds of the present invention were tested for binding in the MC-4 and / or MC-3 assay, as listed in Table 4.

10

TABLE 4

ID No.	IC ₅₀ MC-4 (μM) (Filtration)	IC ₅₀ MC-4 (nM) (SPA)
1	1.1	174
2	1.5	234
3	0.7	159
4	inactive	297
5	1.3	207
6	2.5	293
7	2.2	218
8	1.4	201
9	1.1	217
10	2.3	130(90)
11	2.4	243
12	insoluble	insoluble
13	3.4	347
14	5.6	234
15	4.5	482
16	4.3	424
17	7.7	351
18	6.0	463
19	3.9	460
20	3.5	636

21	7.8	392
22		124
23		124
24		76
25		57
26		63
27		89
28		68
29		48
30		103
31		22
32		48
33		91
34		46
35		100
36		79
37		72
38		159
39		146
40		133
41		483
42		482
43		237
44		89
45		246
46		368
47		874
48		444
49		96
50		30,000
51		1,000
52		1,000

53		1,000
54		800
55		800
56		3,000
57		inactive
58		inactive
59		inactive
60		inactive
61		inactive
62		inactive
63		1,000
64		1,200
65		114
66		120
67		130
68		110
69		139
70		566
71		234
72		155
73		725
74		194
75		103
76		332
77		4.4
78		99
79		216
80		378
81		672
82	1.5	826
83	1.9	833
84	82 nm	

85		
----	--	--

5

EXAMPLE 7Rodent Feeding: Food Intake in Food-Deprived Rats (MC-4)

Male Long-Evans rats (180-200 grams) were housed individually and maintained on a once-a-day feeding schedule (i.e. 10 a.m. until 4 p.m.) for five days following quarantine to allow the animals to acclimate to feeding on powdered chow (#5002 PMI Certified Rodent Meal) during the allotted time. The chow was made available in an open jar, anchored in the cage by a wire, with a metal follower covering the food to minimize spillage. Water was available ad-libitum.

10

15

Animals were fasted for 18 hours prior to testing. At the end of the fasting period, animals were administered either a test compound or vehicle. Vehicle and test compound were administered either orally (5 mL/kg) 60 minutes prior to the experiment, subcutaneously (1 mL/kg) 30 minutes prior to the experiment, or intraperitoneally (1 mL/kg) 30 minutes prior to the experiment. Test compounds were administered orally as a suspension in aqueous 0.5% methylcellulose-0.4% Tween 80, or intraperitoneally as a solution or suspension in PEG 200; compound concentrations typically ranged from 1 mg/kg to 100 mg/kg, preferably from 10-30 mg/kg. Food intake was measured at 2, 4, and 6 hours after administration by weighing the special jar containing the food before the experiment and at the specified times. Upon completion of the experiment, all animals were given a one-week washout period before re-testing.

20

25

30

Following the procedure described above, select compounds of the present invention were tested to measure the effect on food intake in fasted rats, as listed in Table 5 and 6.

5

TABLE 5

ID No.	[mg/kg], Route	Food Intake 0-2 hrs	Food Intake 2-6 hrs
PEG-200	ip	8.25 g	13.38 g
1	10 μ , ip	7.29 g	8.43 g

Table 6

ID No	[mg/kg], Route	Change in Food Intake 0-2 hrs (%)	Change in Food Intake 2-6 hrs (%)
MCT Control	po	0	0
PEG-200	ip	0	0
1	10, ip	-11.60%	-37.00%
1	30, po	-21.2	-6.7
84	10, ip	-7	-24.5
84	10, ip	3.9	-17.8
84	30, po	-7	32
26	10, ip	-12.7	-32.3
27	30, ip	-27.4	-29.1
29	30, po	-26.3	-3.7
29	10, ip	-62.3	-70.7
31	3, ip	-37.8	-50
31	1, ip	-20.6	-71.2
31	30, po	-23.6	31.5
31	10, po	-10.1	39.7
31	3, po	-9	52.1
31	1, po	1.2	19.5
32	10, ip	-29.6	-60.2
32	30, po	-12.5	-15
32	10, po	-20	-1
32	3, po	-11.5	31.17

32	1, po	-13.8	26
34	10, ip	-38	-72
34	30, po	-7.5	2
34	10, po	-1.3	-6
49	30, ip	-20.6	-21.2

5

EXAMPLE 8Neurite Cell Outgrowth Assay

Cell Culture:

10 Dissociated hippocampal and cortical cell cultures were established from embryonic day 18 rat fetuses as described by Mattson, M.P., Barger, S.W., Begley, J, and Mark, R.J., *Methods Cell Biol.*, 1994, 46:087-216. Briefly, fetuses were removed via cesarean section from pregnant moms (Sprague-Dawley) and anesthetized with halothane according to the AVMA Panel on

15 Euthanasia. Pups were decapitated and the brains were removed and placed in HEPES-buffered Hank's Balanced Salt solution (HBSS; Gibco). The hippocampi and cortices were dissected out and pooled according to tissue-type. Tissue was trypsinized for 15 min (1mg/ml trypsin-HBSS; Worthington), rinsed with fresh HBSS, incubated in trypsin inhibitor (1mg/ml; Sigma) for 5

20 min, rinsed again with fresh HBSS and then triturated in 1ml fresh HBSS with a fire-polished glass pipette. Dissociated cells were seeded at 30,000 cells/well onto poly-D-lysine coated 96-well plates (Collaborative BioScience). Each well contained 100 μ l of Eagle's Minimal Essential Media (MEM; Gibco) supplemented with 26mM NaHCO₃ (Sigma), 56mM glucose (Sigma), 15mM KCl (Sigma), 1mM sodium pyruvate (Sigma), 1.1mM L-glutamine (Sigma), 10%

25 (v/v) heat-inactivated fetal bovine serum (Hyclone), and 0.001% gentamicin sulfate (Sigma) (pH 7.4). Cells were allowed to attach for 24h in a humidified 37°C 5% CO₂ incubator before experimental treatment. The culture media was aspirated and exchanged with fresh media every 3 days.

30

Assay:

5 Twenty-four hours after plating, cultures were treated with vehicle
 (PBS+0.1% BSA), alpha-melanocyte stimulating hormone (α -MSH) or test
 compound (diluted in DPBS). Each treatment condition was run in
 quadruplicate. On the third day in culture, the media was aspirated off and
 replaced with fresh media and test compound. At one week in culture, the
 10 cells were fixed with 10% phosphate-buffered formalin for 15min, then rinsed
 with DPBS (Sigma) and placed in blocking serum for 30 min (horse serum;
 1:50 dilution in DPBS; Vector Labs). The cultures were rinsed again with
 DPBS and then incubated in primary antibody for 2 hr (microtubule-associated
 protein-2 (MAP-2) is a selective marker for dendritic processes; anti-mouse
 15 monoclonal (Chemicon); 1:1000 dilution of MAP-2 in antibody diluent (Zymed)).
 Negative control wells were incubated in antibody diluent alone. Background
 signal was determined by blank wells (cell-free) incubated with or without
 antibody. Cultures were rinsed again with DPBS and then placed in
 fluorescein for 1 hr (FITC; anti-mouse IgG; rat adsorbed; 1:50 dilution in DPBS;
 20 Vector Labs). Cultures were rinsed a final time with DPBS and the plates were
 then read on a Cytofluor 4000 fluorescence plate reader. Neurite outgrowth
 was expressed as percent change from control (vehicle).

25 Selected compounds of the instant invention were tested in the above
 assay with results as listed in Table 7. The data are expressed as percent
 change over the vehicle response. All compounds were screened at 50nM. The
 abbreviation NA indicates no change / not active; the abbreviation ND indicates a
 compound not tested / response not determined.

30

TABLE 7: Neurite Outgrowth

Comp #	Hippocampal Cells	Cortical Cells
Vehicle	19%	4%
1	6%	NA
2	18%	18%
3	17%	23%
4	20%	21%

5	28%	9%
6	23%	24%
7	NA	11%
8	NA	12%
9	NA	NA
10	7%	11%
11	9%	10%
12	28%	20%
13	29%	32%
14	30%	19%
15	18%	31%
16	8%	18%
17	8%	17%
18	17%	17%
19	22%	NA
20	17%	NA
21	27%	2%
22	NA	30%
23	10%	20%
24	10%	16%
25	NA	23%
26	16%	47%
27	17%	52%
28	NA	25%
29	NA	8%
30	NA	NA
31	NA	16%
32	NA	6%
33	NA	21%
34	NA	22%
35	NA	28%
36	NA	16%

37	NA	26%
38	NA	NA
39	NA	ND
40	26%	NA
41	36%	5%
42	26%	NA
43	30%	37%
44	43%	19%
45	46%	25%
46	40%	19%
47	20%	13%
48	NA	19%
49	16%	51%
50	46%	NA
51	74%	28%
52	65%	NA
53	67%	NA
54	45%	NA
55	19%	NA
56	33%	NA
57	9%	NA
58	72%	10%
59	61%	21%
60	66%	14%
61	66%	17%
62	13%	13%
63	20%	17%
64	22%	12%
82	14%	15%
83	11%	9%
84	ND	17%

5 The data above show that cultures treated with select compounds of the present invention resulted in a significant increase in neurite outgrowth as measured by MAP2-FITC immunofluorescence. A comparison between the test compounds and α -MSH indicates that many of the test compounds were superior to α -MSH in promoting neurite outgrowth at the concentration tested. In addition, several of the test compounds displayed selective effects on neurite outgrowth in hippocampal or cortical cells.

EXAMPLE 9

In vivo Facial Nerve Compression Model

15 The ability of test compound to provide neuro-protective or neuro-regenerative effects was investigated in a facial nerve compression model. Facial nerve motor axons arise exclusively from neurons within the pons in a well-defined nucleus. Facial nerve compression results in retrograde reactions proximal to the lesion site and Wallerian degeneration at its distal part, which causes diminished whisker movement on the lesioned side.

20 Male Long-Evans rats (150-180 g) were anesthetized with 3-5% isoflurane for induction and 2% for maintenance during the surgical procedures. The right facial nerve was exposed and compressed with forceps for 30 sec at its exit from the stylomastoid foramen. The left facial nerve was sham-operated and served as an internal control. Nerve compression causes paralysis of whisker muscles, hence the reduced whisker movement on the lesioned side, which is observed immediately after recovery from anesthesia. The following morphological abnormalities associated with the functional deficit were observed:

- 25 (1) an increase in the number of perineuronal glial cells in the facial nucleus of the lesioned side, with the increased observed to peak around D3-6;
- (2) a thinner myelin sheath and less myelin basic protein staining in the compressed facial nerve approximately one week after the lesion;
- 30 (3) morphological alterations around the N-M junction and whisker follicles regions and gradually degeneration of motor neurons in the facial nuclei.

5

After recovery from anesthesia, the rats were randomly divided into groups for dosing with vehicle, α MSH or test compound, with 6 animals per group. α MSH (s.c., 70ug/each 48hr) was used as a positive control. Test compounds were dosed p.o. at 20mg/kg bid for 14 days. Restoration of whisker movement was monitored daily after the operation using two criteria:

10

(1) frequency of whisker movement on the lesioned side relative to the the opposite side (sham-operated) which served as the baseline control

(2) semi-quantitative measurements (0 to 4+) on strength of the whisker muscles, characterized based on observation of the percentage of moved whiskers, muscle tone of whisker muscles, and the position of the nose.

15

For all observations, the experimental design was blind to the behavioral observer.

20

The results of behavioral assessment showed that both test compounds, compounds #31 and #84, accelerated the recovery time to restore whisker muscle movement in the lesioned rats as compared to the vehicle controls ($p < 0.05$). The recovery rate of the whisker movement was expressed as percentage of its own internal control (sham-operated), as listed in Table 8.

25

TABLE 8

Functional Recovery of Whisker Movement after Oral Administration of Compound #31 and #84 in Facial Nerve Compression Model

	Percent Recovery					
	D9	D10	D11	D12	D13	D14
Unlesioned Site	100	100	100	100	100	100
Vehicle	0	5.0 \pm 6.8	27.6 \pm 15.9	72.5 \pm 21.2	86.4 \pm 14.5	91.3 \pm 8.3
α -MSH	2.0 \pm 2.8	24.6 \pm 15.8	60.1 \pm 19.3	93.8 \pm 14.0	98.1 \pm 5.3	100 \pm 0.0
Cmpd #31 (246377)	0	3.5 \pm 3.1	67.8 \pm 9.5	98.5 \pm 3.7	100 \pm 0.0	100 \pm 0.0

Cmpd #84 (153791)	0	4.8 ± 4.3	35.1 ± 12.9	90.0 ± 10.9	98.3 ± 4.1	100 ± 0.0
----------------------	---	--------------	----------------	----------------	---------------	--------------

5

EXAMPLE 10*in Vitro* Assay: Measurement of regulation of sebaceous lipid synthesis.

STEP A: Preparation of a feeder layer

10 Semiconfluent cultures of 3T3 mouse fibroblasts (Swiss Albino mouse, ATCC CCL-92) were treated with mitomycin C (4µg/ml) for 3 hours, trypsinized and seeded at a density of 2.5x10⁵/9.5 cm² tissue culture plate in Dulbeccos Minimal Essential Medium (DMEM) containing 10% Colorado Calf Serum, PNC (100U/ml), STM (100µg/ml), L glutamine (0.3mg/ml), sodium pyruvate (1mM) and nonessential amino acids (100µM). The cells were

15 incubated at 37°C for 24 hours prior to their use as a feeder layer for sebocytes.

STEP B: Isolation of human sebocytes

20 Human sebocytes were isolated from Dermatome shavings of postoperative pieces of human skin at 0.4-0.8mm depths (this part of the skin was previously shown to be enriched in sebaceous glands). Shavings so obtained were treated with 1% Dispase in Iscoves medium containing 10% serum for 20 min at 37°C. The tissue was then placed in 0.3% trypsin/1% EDTA in Phosphate Buffered Saline (PBS) for 10 minutes at 37°C. Following

25 this incubation the cells were gently scraped from the tissue in Growth medium (GM) containing DMEM/F12 media mixture (3:1), supplemented with 8% heat inactivated FBS, 2% heat inactivated human serum (HS), 1mM sodium pyruvate, epidermal growth factor (10ng/ml), insulin (10µg /ml), hydrocortisone (0.4µg/ml) and +/-cholera toxin (100µg/ml), L-glutamine and antibiotics.

30 Cells so obtained were filtered through nylon mesh (100µ pore size), centrifuged at 750 RPM, re-suspended in GM and counted.

STEP C: Cultures of human sebocytes

5 Resultant cells from the above isolation procedure were plated on the
3T3 feeder layers at $2 \times 10^5/9.5 \text{cm}^2$ in growth medium and maintained at 37°C
and 5% CO_2 for 3 days (Phase I). Following the initial growth period they were
transferred to a transition medium (TM) that consisted of DMEM/F12 media
supplemented with 1 mM sodium pyruvate, insulin ($10 \mu\text{g/ml}$), transferrin
10 (6.7 ng/ml) and selenium ($5.5 \mu\text{g/ml}$) (ITS), 2% heat inactivated FBS and 2%
heat inactivated human serum as well as +/- cholera toxin (ch.t.) ($100 \mu\text{g/ml}$), L-
glutamine and antibiotics (Phase II). Three days later the cells were changed
to differentiation medium (DM), DMEM/F12 supplemented with ITS, 3,3',5-
triiodo-L-thyronine sodium (3nM), 1% (v/v) trace element mix and the choice of
15 differentiation agent i.e. bovine pituitary extract ($10 \mu\text{g/ml}$). This medium was
changed every 3 days (Phase III).

STEP D: Testing Stimulators or Inhibitors of Sebocyte Differentiation and Lipid Production

20 Hormones, mixture of hormones i.e. bovine pituitary extract or
compounds to be tested were added to the culture at the beginning of phase
III. Two criteria were used to evaluate the effect of these materials on
sebaceous cultures: 1) visual observations and 2) evaluation of sebaceous lipid
accumulation and synthesis. The evaluation of lipid accumulation completed
25 using the Nile red method. This method relies on visualization of neutral lipids
by Nile red and quantitation by reading of fluorescence at 535 nm excitation,
580 nm emission using a plate reader. The lipid synthesis was evaluated by
radioactive labeling using ^{14}C acetate and quantified by Bio Rad
Phosphoimager (Molecular Imager, FX) using 4.1 Software.

30

STEP E: Visual observations & Nile Red evaluation of lipid accumulation

Morphological evaluation of lipid accumulation was easily recognized
since the cells enlarged and displayed lipid granules that in bright field light
microscopy appeared as yellowish circles in the cells. Quantitation of
35 accumulation/inhibition of neutral lipids in sebocytes was accomplished by Nile
red binding assay. Briefly, following exposure of sebocytes to test compounds,
the cells were allowed to interact with $1 \mu\text{M}$ Nile red in Hanks buffered saline

5 solution containing DMSO and Pluronic F127. After 4 hours of incubation, washing and incubation overnight, the fluorescence was read at 535 excitation and 580 emission using a fluorescence plate reader. To determine whether the compounds had an inhibitory effect on cell growth, cell counts were performed.

10

Following the procedure described above, select compounds of the present invention were tested for visual and Nile red evaluation of lipid accumulation, with results as listed in Table 9.

15

TABLE 9

ID No.	% Inh 0.4 μ M	% Inh 0.8 μ M	Visual* 0.4 μ M	Visual* 0.8 μ M	MC5-R IC ₅₀
74	72	100	+++	++++	154nM
76	48	88	++	+++	317nM
80	61	91	++	++++	138nM
67	N.T.	N.T.	++	+++	246nM
50	0	0	0	0	No binding
84	0	0	0	0	No binding

* Increased number of + signs indicates the degree of inhibition with ++++ = 100%, +++ = 75%, ++ = 50% and + = 25% inhibition of lipid granule formation.

STEP F: Evaluation of sebaceous lipid synthesis by sebaceous cells

20

At day 11 of the culture, sebocytes were labeled with ¹⁴C acetate at a final concentration of 2 μ Ci/ml for 24 hours in serum free culture medium. The cells were then scraped from plates and frozen at -80°C in glass vials. Lipid extraction was completed using the Bligh- Dyer method (Bligh, E.G. and Dyer, W.J., *Can. J. Biochem. Physiol.*, 1959, 37, pp 911-916) with slight modification as detailed herein. Briefly, cells were homogenized in a 2:1 chloroform-methanol mixture, in the presence of KCl. The organic phase was removed from the mixture, the separated lipids were dried under argon and spotted to high performance thin layer chromatography (HPTLC) plates. The plates were

25

5 developed by three separate mobile phases. The first was hexane (to the top
of the plate), followed by toluene (to the top) and finally a 70:30:1 mixture of
hexane : ether : acetic acid (half way up the plate-10cm). The plates were than
exposed to radiographic film for visualization of radioactive lipid species. For
10 visualization of unlabeled lipids the plates were sprayed with 8% cupric acid
and charred on a hot plate. Quantitation of the results was done by Image Pro
Plus 3.0 (Media Cybernetics, Silver Springs, MD).

Following the procedure described above, select compounds of the
present invention were tested for their inhibitory effect on differentiation of
15 human sebocytes. Visually, the stained cell cultures treated with compound
#74 showed complete disappearance of lipid granules in sebocyte cultures
following a seven day treatment. The same cells examined for lipid synthesis
revealed inhibition of squalene, cholesterol esters and wax esters, as
measured by radioactively labeled lipids separated by HPTLC. The cell panels
20 were quantified by measurement of the intensity of bands using Image Pro
Plus (version 5.0). % inhibition was calculated based on the difference
between treated samples and vehicle treated controls, with results as listed in
Table 10.

25

Table 10

Lipid	Inducer	Cell Sex	% Inh @ 0.4μM	% Inh @ 0.6μM
Squalene	Cholera toxin & Bovine pituitary extract	F/M	100/100	100/100
Cholesterol ester	"	F/M	85/64	88/76
Unknown	"	F/M	75/68	85/80
Wax ester	"	F/M	70/50	67/42
Triglyceride	"	F/M	0	0

EXAMPLE 11

5 STEP B: Method of treatment:

Mice at 2-3 month post transplantation with human facial skin were used for the studies. The graft area was treated with the test compound at the desired concentration(s) dissolved in polyethylene glycol-ethanol (20 μ l/2 cm²).

10 Controls were treated with vehicle alone. The test compounds were applied daily, excluding weekends. Sebum secretion was determined using Sebutape at 15 days and 30 days following treatment.

STEP C: Termination of the experiment:

15 The termination of the experiment was determined by preliminary clinical evaluation of sebum production using SEBUTAPE. At this time human skin grafts were excised and representative samples were collected for histological evaluation. More particularly, 2mm punch biopsies were prepared and used for evaluation of lipid synthesis and total lipid accumulation in the treated tissues.

20 STEP D: Evaluation of lipid synthesis and total lipid accumulation in the examined tissues.

25 The collected_2mm punch biopsies were placed individually into 96 well plates in Krebs buffer and labeled with 10 μ Ci of ¹⁴C acetate for 3 hours. Following this labeling period the samples were washed in medium and 5 biopsies pooled, weighed, and used for lipid extraction. The lipid extraction and analysis by HPTLC was the same as described for tissue culture derived cells.

30 Following the procedure described above, compound #74 of the present invention was tested for the inhibition of sebaceous gland activity following 11 day treatment of human skin transplanted to the SCID mouse.

35 Visual evaluation of the sebaceous gland following topical treatment with 0.1% solution of compound #74 resulted in visible shrinkage of the sebaceous gland and down regulation of sebaceous lipids. Topical treatment for 15 days with 0.05% and 0.005% solutions was not sufficient to down regulate the lipids. Numerical evaluation of the inhibition of human sebaceous lipids for these

- 5 cells was analyzed using HPTLC, with results as listed in Table 11, 12 and 13. %Inhibition numbers are listed relative to control.

Table 11

Effect of Compound #74 on Human Sebaceous Lipids (11 Day Treatment)

Lipid	%Inh @ 0.1%	%Inh @ 0.5%	%Inh @ 0.01%
Squalene	70	0	0
W/E	80	10	25
Triglycerides	50	0	0

10

Table 12

Effect of Compound #74 on Lipid Accumulation (30 Day Treatment)

Lipid	%Inh @ 0.05%	%Inh @ 0.01%
Squaline	73	82
Cholesterol Ester	21	44
Wax Esters	93	86
Triglycerides	90	75
Cholesterol	82	33

Table 13

Effect of Compound #74 on Sebaceous Lipid Synthesis (30 Day Treatment)

Lipid	%Inh @ 0.05%	%Inh @ 0.01%
Squaline	90	80
Wax Esters	93	86
Triglycerides	90	75
Cholesterol	82	33

15

As shown in Tables 12 and 13 above, both total sebaceous lipid accumulation and de novo synthesis of sebaceous lipids was significantly decreased following 30 days of topical treatment with an 0.05% and 0.01% solution of compound #74.

20

5

EXAMPLE 12Oral Formulation

As a specific embodiment of an oral composition, 100 mg of the compound #74 of Example 2 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

10

EXAMPLE 13Topical FormulationsA: Microemulsion

As a specific embodiment of a microemulsion composition the following components are blended, with heating as need:

15

Polysorbate 60 (e.g Tween 60 from ICI Surfactants)	20 parts
Isopropyl Palmitate	20 parts
Sorbitan Oleate (e.g. Span 80 from ICI Surfactants)	13 parts
2-Ethylhexanediol-1,3	4 parts
Butylated hydroxy-toluene	0.05 parts
Compound #74	0.05 parts

To the blended mixture is then slowly added water (42.9 parts by weight), with mixing as necessary, to yield the emulsion.

20

B: Hydroalcoholic gel

As a specific embodiment of a hydroalcoholic gel composition the polypropylene glycol (10 parts by weight), butylene glycol (10 parts by weight), benzyl alcohol (2 parts by weight), EDTA (0.05 parts by weight) and BHT (0.05 parts by weight) are mixed with water (74.85 parts by weight total). The mixture is blended until all the components are dissolved. Carbomer (e.g. Carbopol 934P from Goodrich) (3 parts by weight) is then slowly added with constant turning to yield a gel. Compound #74 (0.05 parts by weight) is then dispersed into the gel with mixing. The gel pH is adjusted to about pH 3-4.

25

5 C: Anhydrous Gel

As a specific embodiment of an anhydrous gel isopropanol (20 parts by weight) is added to butylene glycol (20 parts by weight). BHT (0.05 parts by weight) and benzyl alcohol (1.0 parts by weight) are then added to the isopropanol/butylene glycol mixture. To the resulting mixture is then added
 10 Cyclotetrasiloxane (D₄) and Organopolysiloxane-11 (e.g. Gransil GSM Gel from Grant Industries) (58.85 parts by weight) with continuous mixing. Compound #74 (0.1 parts by weight) is micronized by and dispersed into the gel with continuous mixing until uniformly distributed.

15 D: Cream

As a specific embodiment of an o/w (oil/water) cream, the following components are mixed in the amounts (parts by weight) as noted. The final mixture is adjusted to about pH 2 with hydrochloric acid.


Cetearyl alcohol	4.3 parts
Microcrystalline wax	9.0 parts
Ceteth-20 Surfactant (e.g. Brij 58 from ICI Surfactants))	1.1 parts
Capric/Caprylic Triglycerides (e.g. Tegosoft CT from GoldSchmidt)	3.6 parts
Glycine	0.6 parts
Compound # 74	0.1 parts
BHT	0.05 parts
Water	81.25 parts

20 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

25

5 WHAT IS CLAIMED IS:

1. A compound of the formula (II)



(II)

wherein

10 R¹ is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl;

15 wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

20 R² is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl;

25 wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

30 R⁴ is selected from the group consisting of aryl, aralkyl, heteroaryl, heterocycloalkyl, and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

5 provided that when R¹ is aryl substituted with one halogen and R⁴ is aryl substituted with one halogen, then R² is not morpholinyl; and pharmaceutically acceptable salts thereof.

2. A compound as in Claim 1 wherein

10 R¹ is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino;

15 R² is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino;

20 R⁴ is selected from the group consisting of aryl, aralkyl and heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino; and pharmaceutically acceptable salts thereof.

3. A compound as in Claim 2 wherein

25 R¹ is substituted aryl; wherein the aryl group is optionally substituted with one to two substituents selected from halogen, alkyl and alkoxy;

R² is substituted aryl; wherein the aryl group is substituted with one to two substituents selected from alkyl and alkoxy;

30 R⁴ is selected from the group consisting of aryl and substituted aryl; wherein the substituents on the aryl are one to two independently selected from halogen, alkyl and alkoxy; and pharmaceutically acceptable salts thereof.


4. A compound as in Claim 3 wherein

35 R¹ is 2-methoxyphenyl;
R² is 2-methoxyphenyl;

5 R^4 is selected from the group consisting of phenyl, 4-methoxyphenyl, 2,6-difluorophenyl, 3,5-difluorophenyl and 2-chloro-6-methylphenyl; and pharmaceutically acceptable salts thereof.

10 5. A compound as in Claim 4 selected from the group consisting of [2-(2-methoxyphenyl)-3-(2-methoxyphenyl)-2H-[1,2,4]-thiadiazol-5-ylidene]-phenylamine and pharmaceutically acceptable salts thereof.

6. A compound of the formula (II)



(II)

15 wherein

R^1 is selected from the group consisting of heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl;

20 wherein the heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

25 wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

R^2 is selected from the group consisting of aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl and cycloalkyl-alkyl;

30 wherein the aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

5 R⁴ is selected from the group consisting of heteroaryl, heterocycloalkyl, cycloalkyl-alkyl, substituted aryl and substituted aralkyl;

 wherein the heteroaryl, heterocycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, 10 alkylamino or di(alkyl)amino;

 wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 provided that when R¹ is aryl substituted with one halogen and R⁴ is aryl 15 substituted with one halogen, then R² is not morpholinyl;

 provided further that when R¹ is methylphenyl and R⁴ is methoxyphenyl, then R² is selected from the group consisting of aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl, cycloalkyl-alkyl and substituted aryl;

20 wherein the aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 wherein the aryl group is substituted with one or more substituents 25 independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 and pharmaceutically acceptable salts thereof.


7. A compound as in Claim 6 wherein

30 R¹ is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 R² is selected from the group consisting of aryl and aralkyl; wherein the 35 aryl or aralkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

- 5 R^4 is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl group is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino; provided that when R^1 is methylphenyl and R^4 is methoxyphenyl, then
- 10 R^2 is selected from the group consisting of aralkyl and substituted aryl; wherein the aralkyl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino; wherein the aryl group is substituted with one or more substituents
- 15 independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino; and pharmaceutically acceptable salts thereof.
- 20 8. A pharmaceutical composition comprising a compound as claimed in Claim 1 and a pharmaceutically acceptable carrier.
9. A method of making a pharmaceutical composition comprising mixing a compound of Claim 1 with a pharmaceutically acceptable carrier.
- 25 10. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.
11. A pharmaceutical composition comprising a compound as claimed in Claim 6 and a pharmaceutically acceptable carrier.
- 30 12. A method of making a pharmaceutical composition comprising mixing a compound of Claim 6 with a pharmaceutically acceptable carrier.
13. A pharmaceutical composition made by mixing a compound of Claim 6 and a pharmaceutically acceptable carrier.
- 35

- 5 14. A method of treating a disorder mediated by a melanocortin receptor comprising administering to a subject in need thereof a therapeutically effective amount of a compound of formula (II)



(II)

wherein

- 10 R¹ is selected from the group consisting of alkyl, aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heteroaryl-alkyl, heterocycloalkyl, heterocycloalkyl-alkyl, cycloalkyl or cycloalkyl-alkyl group is optionally substituted with one or more substituents independently selected from
15 halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

- R² is selected from the group consisting of alkyl, aryl, aralkyl, heteroaryl, heterocycloalkyl and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl group is optionally substituted with one or more
20 substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

- R⁴ is selected from the group consisting of hydrogen, alkyl, aryl, aralkyl, heteroaryl, heterocycloalkyl, and cycloalkyl-alkyl; wherein the aryl, aralkyl, heteroaryl, heterocycloalkyl or cycloalkyl group is optionally substituted with
25 one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy; halogenated alkyl, halogenated alkoxy, amino, alkylamino or di(alkyl)amino;

 and pharmaceutically acceptable salts thereof.

- 30 15. A method as in Claim 14 wherein
 R¹ is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl is substituted with one or more substituents

5 independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino;

R^2 is selected from the group consisting of substituted aryl and substituted aralkyl; wherein the aryl or aralkyl is substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino;

10

R^4 is selected from the group consisting of aryl, aralkyl and heteroaryl; wherein the aryl, aralkyl or heteroaryl group is optionally substituted with one or more substituents independently selected from halogen, hydroxy, alkyl, alkoxy, trihalomethyl, trihalomethoxy, amino, alkylamino or di(alkyl)amino;

15

and pharmaceutically acceptable salts thereof.

16. A method as in Claim 15 wherein

R^1 is substituted aryl; wherein the aryl group is optionally substituted one substituent selected from halogen, alkyl and alkoxy;

20

R^2 is substituted aryl; wherein the aryl group is substituted one substituent selected from alkyl and alkoxy;

R^4 is selected from the group consisting of aryl and substituted aryl; wherein the substituents on the aryl are one to two independently selected from halogen, alkyl and alkoxy;

25

and pharmaceutically acceptable salts thereof.

17. A method as in Claim 16 wherein R^1 is 2-methoxyphenyl;

R^2 is 2-methoxyphenyl;

R^4 is selected from the group consisting of phenyl, 2,6-difluorophenyl, 3,5-difluorophenyl and 2-chloro-6-methylphenyl;

30

and pharmaceutically acceptable salts thereof.

18. A method as in Claim 17 wherein the compound of formula (II) is selected from the group consisting of [2-(2-methoxyphenyl)-3-(2-methoxyphenyl)-2H-[1,2,4]-thiadiazol-5-ylidene]-phenylamine and pharmaceutically acceptable salts thereof.

35

5 19. The method of Claim 14, wherein the disorder mediated by a melanocortin receptor is selected from the group consisting of metabolic disorders, CNS disorders and dermatological disorders.

10 20. The method of Claim 14, wherein the disorder mediated by a melanocortin receptor is selected from the group consisting of obesity, impaired oral glucose tolerance, elevated blood glucose levels, type II diabetes, Syndrome X, diabetic retinopathy, acute neurodegenerative disorders, chronic neurodegenerative disorders, plexopathies, male erectile dysfunction, dry eyes, acne, dry skin, aged skin, seborrheic dermatitis,
15 rosacea, excessive ear wax, meibomian gland disorder, pseudofolliculitis, yeast infections, dandruff, hidradenitis suppurativa, ocular rosacea and eccrine gland disorder.

20 21. The method of Claim 20, wherein the disorder mediated by a melanocortin receptor is selected from the group consisting of obesity, impaired oral glucose tolerance, elevated blood glucose levels, type II diabetes and Syndrome X.

25 22. The method of Claim 20, wherein the disorder mediated by a melanocortin receptor is selected from the group consisting of acne, dry skin and seborrheic dermatitis.

30 23. The method of Claim 14, wherein the melanocortin receptor is a selected from the group consisting of the melanocortin-3 receptor, the melanocortin-4 receptor and the melanocortin-5 receptor.

INTERNATIONAL SEARCH REPORT

PCT/US 02/35340

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D285/08 C07D417/12 A61K31/433 A61P3/00 A61P17/00
A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ABRAHAM W ET AL: "Formamidinoylisothiocyanate - III. Addition nucleophiler Partner and Formamidinoylisothiocyanate" TETRAHEDRON, vol. 29, no. 5, March 1973 (1973-03), pages 699-705, XP002228008 the whole document, particularly page 704, left-hand column, 2nd paragraph	1,7
A	WO 01 55107 A (MELACURE THERAPEUTICS AB) 2 August 2001 (2001-08-02) the whole document	1,6,14

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

21 January 2003

Date of mailing of the international search report

31/01/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 14-23 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US 02/35340

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0155107 A	02-08-2001	AU 2868101 A WO 0155107 A2	07-08-2001 02-08-2001
