




(86) Date de dépôt PCT/PCT Filing Date: 2006/03/29
(87) Date publication PCT/PCT Publication Date: 2006/10/05
(85) Entrée phase nationale/National Entry: 2007/09/28
(86) N° demande PCT/PCT Application No.: US 2006/012046
(87) N° publication PCT/PCT Publication No.: 2006/105436
(30) Priorité/Priority: 2005/03/30 (US60/666,780)

(51) Cl.Int./Int.Cl. *A01H 1/00* (2006.01),
C07H 21/02 (2006.01), *C12N 15/82* (2006.01),
C12N 15/87 (2006.01)
(71) Demandeur/Applicant:
REGENTS OF THE UNIVERSITY OF CALIFORNIA, US
(72) Inventeurs/Inventors:
ZHU, JIAN-KANG, US;
SUNKAR, RAMANJULU, US
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : CLONAGE ET CARACTERISATION DE MICRO-ARN TIRES DU RIZ
(54) Title: CLONING AND CHARACTERIZATION OF MICRORNAS FROM RICE

A.



(57) Abrégé/Abstract:

The present invention provides new miRNAs in rice. The nucleic acids of the invention can be used to control gene expression in plants.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 October 2006 (05.10.2006)

PCT

(10) International Publication Number
WO 2006/105436 A2

(51) International Patent Classification:

A01H 1/00 (2006.01) *C07H 21/02* (2006.01)
C12N 15/87 (2006.01) *C12N 15/82* (2006.01)

(21) International Application Number:

PCT/US2006/012046

(22) International Filing Date: 29 March 2006 (29.03.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/666,780 30 March 2005 (30.03.2005) US

(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, California 94607 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHU, Jian-Kang** [US/US]; 871 Saltcoates Drive, Riverside, California 92508 (US). **SUNKAR, Ramanjulu** [IN/US]; 911 Blaine Street, Apt. 49, Riverside, California 92507 (US).

(74) Agents: **GAO, Chuan** et al.; Two Embarcadero Center, 8th Floor, San Francisco, California 94111 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))


Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CLONING AND CHARACTERIZATION OF MICRORNAS FROM RICE

A.



(57) Abstract: The present invention provides new miRNAs in rice. The nucleic acids of the invention can be used to control gene expression in plants.

WO 2006/105436 A2

CLONING AND CHARACTERIZATION OF MICRORNAS FROM RICE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. patent application 60/666,780, filed March 30, 2005, the contents of which are incorporated by reference in the entirety.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] This work was supported by National Institutes of Health grant R01GM0707501 and National Science Foundation grant IBN-0212346.

BACKGROUND OF THE INVENTION

[0003] MicroRNAs (miRNAs) are short, endogenous noncoding RNAs found in animals (Lee, R. C. et al., *Cell* **75**:843-854 (1993); Wightman, B. et al., *Cell* **75**:855-862 (1993); Lagos-Quintana, M. et al., *Science* **294**:853-858 (2001); Lagos-Quintana, M. et al., *Curr. Biol.* **12**:735-739 (2002); Lagos-Quintana, M. et al., *RNA* **9**:175-179 (2003); Lau, N. C. et al., *Science* **294**:858-862 (2001); Lee, R. C. et al., *Science* **294**:862-864 (2001); Lim, L. P. et al., *Science* **299**:1540 (2003a); Lim, L. P. et al., *Genes Dev.* **17**:991-1008 (2003b); Mourelatos, Z., et al., *Genes Dev.* **16**:720-728 (2002); Aravin, A. et al., *Dev. Cell* **5**:337-350 (2003); Ambros, V. et al., *Curr. Biol.* **13**:807-818 (2003a); Dostie, J. et al., *RNA* **9**:180-186 (2003); Grad, Y. et al., *Mol. Cell* **11**:1253-1263 (2003)), plants (Reinhart, B. J. et al., *Genes Dev.* **16**:1616-1626 (2002); Llave, C. et al., *Plant Cell* **14**:1605-1619 (2002a); Park, W. et al., *Curr. Biol.* **12**:1484-1495 (2002); Mette, M. F. et al., *Plant Physiol.* **130**:6-9 (2002); Palatnik, J. F. et al., *Nature* **425**:257-263 (2003); Floyd, S. K. et al., *Nature* **428**:485-486 (2004); Jones-Rhoades, M. J. et al., *Mol. Cell* **14**:787-799 (2004); Sunkar, R. et al., *Plant Cell* **16**:2001-2019 (2004); Wang, J-F. et al., *Nucleic Acids Res.* **32**:1688-1695 (2004a); Wang, X. J. et al., *Genome Biol.* **5**:R65 (2004b); Adai, A. et al., *Genome Res.* **15**:78-91 (2005)), and the Epstein-Barr virus (Pfeffer, S. et al., *Science* **304**:734-736 (2004)). In both animals and plants, the majority of the miRNA genes exists as independent transcriptional units and they are transcribed by RNA polymerase II into long primary transcripts (termed pri-miRNAs) (Bartel, D. P. *Cell* **116**:281-297 (2004); Parizotto, E. A. et al., *Genes Dev.* **18**:2237-2242 (2004); Kurihara, Y. et al., *Proc. Natl. Acad. Sci. USA* **101**:12753-12758 (2004)). In animals,

pri-miRNAs are trimmed in the nucleus to generate ~70 nt miRNA precursors (pre-miRNAs) with fold-back structures by a multi-protein complex called microprocessor in which Drosha (an RNase III-like enzyme) and Pasha (a double-stranded RNA binding protein) are critical components (Lee, Y. et al., *Nature* 425:415-419 (2003); Denli, A. H. et al., *Nature* 432:231-234 (2004)). The pre-miRNAs are exported to the cytoplasm and subsequently cleaved by another RNase III-like enzyme called Dicer to generate mature miRNAs (Bernstein, E. et al., *Nature* 409:363-366 (2001)). However, the Arabidopsis genome does not appear to encode a Drosha ortholog, and it seems that the plant nuclear-localized Dicer homolog is likely to have Drosha function (Kurihara, Y. et al., *Proc. Natl. Acad. Sci. USA* 101:12753-12758 (2004)). Many miRNAs are conserved between species – often over wide evolutionary distances. For example, *AthmiR166* is conserved in all lineages of land plants, including bryophytes, lycopods, ferns and seed plants (Floyd, S. K. et al., *Nature* 428:485-486 (2004)), and the *Caenorhabditis elegans* miRNA, *let-7*, is conserved in human, *Drosophila*, and eleven other bilateral animals (Pasquinelli, A. E. et al., *Nature* 408:86-89 (2000)); but others are only conserved between more closely related species such as *C. elegans* and *C. briggsae* (Ambros, V. et al., *Curr. Biol.* 13:807-818 (2003a); Bartel, D. P. *Cell* 116:281-297 (2004)). miRNAs down-regulate the expression of specific mRNA targets, either by directing the cleavage of mRNAs or interfering with translation (Carrington, J. C. et al. *Science* 301:336-338 (2003); Bartel, D. P. *Cell* 116:281-297 (2004); Ambros, V. *Nature* 431:350-355 (2004)).

[0004] miRNAs have been identified by cloning and by computational approaches tailored to the key features of *lin-4* and *let-7*, the 2 founding members of miRNAs from *C. elegans*, which include a fold-back hairpin RNA precursor coupled with evolutionary conservation (Ambros, V. et al., *RNA* 9:277-279 (2003b)). It was estimated that miRNA genes represent 1% of the expressed genome in complex organisms such as worms, flies and humans (Lai, E. C. *Curr. Biol.* 13:R925-R936 (2003); Lim, L. P. et al., *Genes Dev.* 17:991-1008 (2003b); Bartel, D. P. *Cell* 116:281-297 (2004)). However, recent computational predictions have raised the number of miRNAs significantly in primates by comparative analysis of the human, mouse and rat genomes (Berezikov, E. et al., *Cell*, 120:21-24 (2005)). The identification of the entire set of miRNAs and their target genes from model organisms is of fundamental importance to understand regulatory networks and gene silencing mechanisms.

[0005] Rice is the world's most important crop, as measured by the portion of calories it provides to the human diet. It is an established model system for monocots that include all

cereals. Rice is the only monocot species with a fully sequenced genome. The availability of the complete genome sequence of rice allowed the *in silico* identification of 20 families of rice miRNAs based on conservation of sequences with *Arabidopsis* miRNAs (Reinhart, B. J. et al., *Genes Dev.* 16:1616-1626 (2002); Park, W. et al., *Curr. Biol.* 12:1484-1495 (2002); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA.* 101:11511-11516 (2004); Wang, J-F. et al., *Nucleic Acids Res.* 32:1688-1695 (2004a); Adai, A. et al., *Genome Res.* 15:78-91 (2005)). In addition to finding conserved miRNAs, cloning approaches revealed *Arabidopsis* miRNAs that are not conserved in rice. At least four well-characterized *Arabidopsis* miRNAs, miR158, miR161, miR163 and miR173 do not have homologs in rice (Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). Another miRNA (miR403) has been found to be conserved between *Arabidopsis* and *Populus* while its counterpart could not be identified in rice (Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)). Recently, evidence was shown that the non-conserved miR161 and miR163 from *Arabidopsis* may have evolved by inverted duplication of their target genes (Allen, E. et al., *Nature Genet.* 36:1282-1290 (2004)). Additionally, Berezikov, E. et al., *Cell*, 120:21-24 (2005)) have predicted lineage specific miRNAs in mammalian and non-mammalian animal species. Taken together, these observations support the notion that rice may express monocot- and/or rice-specific miRNAs.

[0006] The present study was undertaken to identify new miRNAs that are difficult to predict *in silico* and verify previously predicted miRNAs from rice. Sequencing of small RNA libraries and subsequent analysis led to the identification of 14 new miRNAs. These new miRNAs from rice form 14 families, 13 of which are new and not present in *Arabidopsis*. Furthermore, we confirmed the existence of 15 of the 20 conserved families of miRNAs that were predicted previously. Based on sequence complementarity to miRNAs, we were able to predict 46 rice genes as putative targets of the new miRNAs. These predicted targets include not only transcription factors but also other genes involved in diverse physiological processes.

BRIEF SUMMARY OF THE INVENTION

[0007] MicroRNAs (miRNAs) are a growing family of small non-coding RNAs that down-regulate gene expression in a sequence-specific manner. The identification of the entire set of miRNAs from a model organism is a critical step towards understanding miRNA-guided

gene regulation. Rice and *Arabidopsis*, two plant model species with fully sequenced genomes, are representatives of monocotyledonous and dicotyledonous flowering plants, respectively. Thus far, experimental identification of miRNAs in plants has been confined to *Arabidopsis*. Computational analysis based on conservation with known miRNAs from *Arabidopsis* has predicted 20 families of miRNAs in rice. To identify miRNAs that are difficult to predict *in silico* or not conserved in *Arabidopsis*, we generated 3 cDNA libraries of small RNAs from rice shoot, root and inflorescence tissues. We identified 35 miRNAs, of which 14 are new and these define 13 new families. Thirteen of the new miRNAs are not conserved in *Arabidopsis*. Four of the new miRNAs are conserved in related monocot species but not in *Arabidopsis*, which suggests that these may have evolved after the divergence of monocots and dicots. The remaining 9 new miRNAs appear to be absent in the known sequences of other plant species. Most of the rice miRNAs are expressed ubiquitously in all tissues examined, while a few display tissue-specific expression. We predicted 46 genes as targets of the new rice miRNAs, and 16 of these predicted targets encode transcription factors and other target genes appear to have roles in diverse physiological processes. Four target genes have been experimentally verified by detection of miRNA mediated mRNA cleavage. Our identification of new miRNAs in rice suggests that these miRNAs may have evolved independently in rice or may have been lost in other species.

BRIEF DESCRIPTION OF THE DRAWINGS

[0008] **Figure 1.** Size distribution of new miRNAs cloned from rice.

[0009] **Figure 2.** OsmiR396d is a unique member of the previously predicted OsmiR396 family in rice. Sequence alignment of miR396d and predicted members of OsmiR396 family. Bold letter represents the additional nucleotide in the new miRNA.

[0010] **Figure 3.** Schematic representation of the biogenesis of OsmiR444. Its hairpin structure requires parts of exon 2 and exon 3 of the host transcript J033125N22.

[0011] **Figure 4.** The miR444 family is conserved in monocots. (A). Alignment of miR444 sequence from rice with the predicted homologs in wheat, barley, maize, sorghum and sugarcane. (B). Predicted fold-back structures of miR444 precursors from rice, wheat, barley, maize, sorghum and sugarcane.

[0012] **Figure 5.** Predicted fold-back structure of OsmiR436 precursor. The fold-back structure was predicted with use of a 720-bp processed transcript. Protruding stem-loops in the 3' arm of the hairpin are indicated by the \ sign.

[0013] **Figure 6.** Expression patterns of miRNAs cloned from rice. Northern blots of total RNA isolated from different tissues were probed with labeled oligonucleotides. The blots also included RNA from maize and *Arabidopsis*. The tRNA and 5S rRNA bands were visualized by ethidium bromide staining of polyacrylamide gels and served as loading controls. Labelled RNA oligonucleotide was used as a size marker and the position was indicated.

[0014] **Figure 7.** Expression patterns of new rice miRNAs that are conserved in another monocot (maize) or in dicot (*Arabidopsis*) and monocot (maize). The tRNA and 5S rRNA bands were visualized by ethidium bromide staining of polyacrylamide gels and served as loading controls. Labelled RNA oligonucleotide was used as a size marker and the position was indicated.

[0015] **Figure 8.** OsmiR439 is predicted to target 3 sites within the ORF of its target gene, 11667.m02576. Numbers represent the position of target sites in the ORF. Amino acid sequence corresponding to the target nucleotide sequence is shown.

[0016] **Figure 9.** Identification of miRNA-guided cleavage products of target genes in rice. (A). mRNA 7448.m00137, (B). mRNA 11668.m00935, (C). mRNA 11668.m04852 and (D). mRNA 11686.m04227. Mapping of cleavage sites was done by RLM-5'RACE. Partial mRNA sequences from target genes were aligned with miRNAs. Numbers indicate the fraction of cloned PCR products terminating at different positions.

[0017] **Figure 10.** Predicted fold-back structures of the new miRNA precursors from rice.

[0018] **Figure 11.** Predicted fold-back structures of miR390 from *Arabidopsis* and *Populus*.

[0019] **Figure 12.** Predicted fold-back structures of miR437 from maize, sugarcane, and *Sorghum*.

[0020] **Figure 13.** Predicted fold-back structures of the putative miRNA precursors from rice (Table 5).

DEFINITIONS

[0021] As used in this application, the term “expression cassette” refers to any recombinant expression system for the purpose of expressing a nucleic acid sequence of the invention *in vitro* or *in vivo*, constitutively or inducibly, in any cells, including, in addition to plant cells, prokaryotic, yeast, fungal, insect, or mammalian cells. The term includes linear or circular expression systems. The term includes all vectors. The cassettes can remain episomal or integrate into the host cell genome. The expression cassettes may have the ability to self-replicate or may not, *i.e.*, drive only transient expression in a cell. The term includes recombinant expression cassettes that contain only the minimum elements needed for transcription of the recombinant nucleic acid.

[0022] As used herein, the term “promoter” includes all sequences capable of driving transcription of a coding sequence in a plant cell. Thus, promoters used in the constructs of the invention include *cis*-acting transcriptional control elements and regulatory sequences that are involved in regulating or modulating the timing and/or rate of transcription of a gene. For example, a promoter can be a *cis*-acting transcriptional control element, including an enhancer, a promoter, a transcription terminator, an origin of replication, a chromosomal integration sequence, 5' and 3' untranslated regions, or an intronic sequence, which are involved in transcriptional regulation. These *cis*-acting sequences typically interact with proteins or other biomolecules to carry out (turn on/off, regulate, modulate, *etc.*) transcription. Promoters can be constitutive or inducible.

[0023] The term “plant” includes whole plants, shoot vegetative organs/structures (*e.g.*, leaves, stems and tubers), roots, flowers and floral organs/structures (*e.g.*, bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm, and seed coat) and fruit (the mature ovary), plant tissue (*e.g.*, vascular tissue, ground tissue, and the like) and cells (*e.g.*, guard cells, egg cells, trichomes and the like), and progeny of same. The class of plants that can be used in the method of the invention is generally as broad as the class of higher and lower plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), gymnosperms, ferns, and multicellular algae. It includes plants of a variety of ploidy levels, including aneuploid, polyploid, diploid, haploid and hemizygous.

[0024] Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

[0025] Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

[0026] "Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

[0027] The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, at least 80% sequence identity, preferably at least 85%, more preferably at least 90%, 93% and most preferably at least 95%, or 97% compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%, preferably at least 60%, 70%, 80% or more preferably at least 90%,

and most preferably at least 95%. Polypeptides that are "substantially similar" share sequences as noted above except that residue positions that are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

[0028] Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C.

[0029] For the purposes of this disclosure, stringent conditions for hybridizations are those including at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions. Moderately stringent conditions include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent conditions.

[0030] All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION

I. INTRODUCTION

[0031] The present invention is based, at least in part, on the identification of new miRNAs in rice. The nucleic acids of the invention can be used to control gene expression in plants. In some embodiments, the expression cassettes encoding the miRNAs of the invention are prepared and introduced into plants. The encoded miRNAs then control expression of the endogenous target genes. Alternatively, one can modify the target gene so as to render it miRNA-resistant by modifying the sequence to decrease or inhibit pairing with the miRNA. The modifications will typically be selected such that the sequence of the encoded protein is not altered. The modified target gene can be incorporated into an expression cassette and introduced into a plant. Alternatively, an endogenous target gene can be modified using known techniques (*e.g.*, homologous recombination).

[0032] More specifically, the present inventors discovered 14 new miRNAs in rice. These new miRNAs form 14 families, 13 of which are new and not found in *Arabidopsis*. In addition, 46 rice genes have been identified as putative targets of the new miRNAs, based on their sequence complementarity to the miRNAs.

II. GENERAL DESCRIPTION OF TECHNIQUES USED IN THE INVENTION

[0033] Standard techniques in the field of molecular genetics are useful for the present invention. Basic texts disclosing the general methods such as cloning, nucleic acid sequencing, detection of nucleic acids, and gene transfer and expression, include, *e.g.*, Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* 3d ed. (2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and Ausubel *et al.*, *Current Protocols in Molecular Biology* (1994).

[0034] For nucleic acids, sizes are given in either kilobases (Kb) or base pairs (bp). These are estimates derived from agarose or polyacrylamide gel electrophoresis, from sequenced nucleic acids, or from published DNA sequences. For proteins, sizes are given in kilo-Daltons (kD) or amino acid residue numbers. Proteins sizes are estimated from gel electrophoresis, from sequenced proteins, from derived amino acid sequences, or from published protein sequences.

[0035] Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Caruthers, *Tetrahedron Letters*, 22: 1859-1862 (1981), using an automated synthesizer, as described in Van Devanter *et al.*, *Nucleic Acids Res.*, 12: 6159-6168 (1984). Purification of oligonucleotides is by either native polyacrylamide gel electrophoresis or by anion-exchange chromatography as described in Pearson & Reanier, *J. Chrom.*, 255:137-149 (1983). The sequence of the cloned genes and synthetic oligonucleotides can be verified aftercloning using, *e.g.*, the chain termination method for sequencing double-stranded templates of Wallace *et al.*, *Gene* 16:21-26 (1981).

III. GENE TRANSFER TECHNIQUES

[0036] DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria.

[0037] Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm *et al.*, *Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Ballistic transformation techniques are described in Klein *et al.*, *Nature* 327:70-73 (1987).

[0038] *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example, Horsch *et al.*, *Science* 233:496-498 (1984), and Fraley *et al.*, *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).

[0039] Transformed plant cells that are derived by any of the above transformation techniques can be cultured to regenerate a whole plant that possesses the transformed genotype and thus the desired phenotype. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker that has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

[0040] One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

[0041] The expression cassettes of the invention can be used to confer a desired trait on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Cucumis, Cucurbita, Daucus, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Lactuca, Linum, Lolium, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Oryza, Panieum, Pannesetum, Persea, Pisum, Pyrus, Prunus, Raphanus, Secale, Senecio, Sinapis, Solanum, Sorghum, Trigonella, Triticum, Vitis, Vigna, and, Zea.

EXAMPLES

[0042] The following examples are provided for the purpose of illustration and not limitation. Those of skill in the art will readily recognize a variety of non-critical parameters that could be changed or modified to yield essentially the same or similar results.

RESULTS

Identification of 13 New Families of Rice MiRNAs

[0043] Because the available computational approaches can only identify miRNAs that are conserved between *Arabidopsis* and rice, a cloning approach was employed to identify rice

miRNAs that may not be conserved or may have atypical features (Dugas, D. V. et al., *Curr. Opin. Plant Biol.* 7:512-520 (2004)). To this end, we generated 3 independent small RNA libraries from rice in the size range of 18-26 nt, from the shoots and roots of seedlings and inflorescence tissues of adult plants (*Oryza sativa* spp *japonica* cv. Nipponbare). Small RNAs were isolated by size fractionation and ligated to 5' and 3' adapters, cloned and sequenced. A total of ~10,000 clones were sequenced (about 1/3 from each library), of which ~5000 small cDNA sequences were between 18 and 26 nt in length. The remaining sequences had either shorter fragments or self-ligated adapters. BLASTN searches revealed that 97% of these sequences have at least one match in the rice nuclear genome sequence Version 3 annotated by TIGR (www.tigr.org). The remaining 3% did not have a match and were not analyzed further. The lack of a match of these sequences may be due to unfinished regions in the rice genome sequence, sequencing errors or other possibilities. Several clones were mapped to chloroplast or mitochondrial genomes and may represent either degradation or possibly regulatory products of organellar RNAs. The largest class of cloned RNAs represents fragments of abundant noncoding RNAs (rRNA, tRNA, snRNA and snoRNA) as determined by BLASTN searches against the Rfam database. A small fraction represents mRNA breakdown products from rice. The remaining sequences constitute miRNAs (Table 1 and Table 2) and endogenous siRNAs (data not shown). For 95% of the endogenous siRNAs, we could not be detected their expression on small RNA blots.

[0044] MicroRNAs were distinguished from endogenous siRNAs on the basis of the ability of the miRNA surrounding sequences to adopt a hairpin structure (Figure 10). This analysis revealed that we had cloned 35 rice miRNAs. We also found one small RNA sequence that corresponds to OsmiR399g*. Sequence similarity searches against the central miRNA registry (<http://www.sanger.ac.uk/Software/Rfam/mirna/search.shtml>) showed that 14 of the miRNAs are new (Table 1). The remaining 21 (belonging to 15 families) were identical with previously predicted miRNAs in rice (Table 2). The newly identified 14 miRNAs correspond to 34 loci. These new miRNAs belong to 2 predominant size classes: 21 and 24 nt in length (Figure 1). Nine of the 14 newly identified miRNAs begin with a 5' U which is a characteristic feature of miRNAs (Table 1). All 14 new miRNAs are perfectly conserved in Indica rice (*Oryza sativa* spp *indica*).

[0045] One of the newly identified miRNAs is represented by 2 genomic loci, OsmiR396d and OsmiR396e, and is a member of previously predicted OsmiR396 family in rice (Table 1

and Figure 2). OsmiR396d was represented by 10 clones in our libraries and differed slightly in sequence from that of predicted OsmiR396 (Figure 2). The predicted OsmiR396 has 3 genomic loci (OsmiR396a, b and c) and is represented by 2 members (Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). OsmiR396d differs from these two members by the presence of an additional nucleotide "G" between positions 8 and 9. Using a specific probe, we detected the expression of miR396d in rice and maize but not in *Arabidopsis* (Figure 7A). Although cross-hybridization often occurs between members in the same miRNA family, this is prevented by the presence of an additional nucleotide in the middle of miR396d. Consistent with the absence of a signal in the *Arabidopsis* Northern blot, the miR396d sequence is not present in the *Arabidopsis* genome. OsmiR396d sequence and the secondary structure of its precursor sequences are conserved in barley, another monocotyledonous plant (data not shown).

[0046] Our sequence analysis indicated that we also identified a new miRNA that is conserved between monocots and dicots. We identified miR390 in rice through cloning, whereas the *Arabidopsis* counterpart was predicted through recent computational approaches (Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA* 101:11511-11516 (2004); Wang, X. J. et al., *Genome Biol.* 5:R65 (2004b)). miR390 is represented by one member with one locus in rice (Table 1) while in *Arabidopsis* and *Populus* it is represented by 2 members with 3 and 6 loci, respectively (Figure 11).

Genomic organization of the new rice miRNAs

[0047] Genomic locations of the new miRNA genes in rice are shown in Table 1. In total, the 14 newly identified miRNAs correspond to 34 loci. Hairpin structures can be predicted for all these 34 loci using miRNA surrounding sequences (Figure 10). Ten of these are encoded by single copy miRNA genes, whereas the other 4 (miR439, miR396d, miR441 and miR445) have multiple loci in the genome (Table 1). The exact origins of miRNAs corresponding to multiple genomic loci cannot be assigned unambiguously, and some of the loci could be pseudogenes. Our analysis of the genomic positions of the new miRNA genes shows that the majority localizes to intergenic regions (25 out of 34 loci). However, 7 correspond to introns of protein-coding genes in either the sense (6) or antisense (1) orientation (Table 1). Our characterization of intronic origins of miRNAs was based on latest annotation of *Oryza sativa* spp. *japonica* genome (version 3.0). Two miRNAs (miR439 and

miR445) map to both intergenic and intronic locations. Two (miR435 and miR440) are derived from introns only. Another 2 miRNAs (miR436 and miR444) originate from the exons of protein-coding genes in the sense polarity (Table 1).

Biogenesis of OsmiR436 and OsmiR444

[0048] In general, most of the 20-25-nt mature miRNAs are processed from a 70 – 300 nt precursor forming a hairpin structure that contains mature miRNA in either of its arms. Two miRNAs, miR436 and miR444, were mapped to the exons of the protein-coding genes J023035E19 (AK120922) and J033125N22 (AK103332), respectively, in the sense polarity (Table 1). The existence of these processed transcripts is supported by expression data (Kikuchi et al., 2003). Both of the precursor transcripts can form hairpin structures, and the miRNAs were detected on small RNA blots as discrete bands, suggesting that these are not non-specific degradation products.

[0049] The biogenesis of OsmiR444 and OsmiR436 is unusual, because the fold-back structure could not be predicted directly from the genomic sequence surrounding the miRNA. A hairpin structure can be predicted for a processed transcript (part of exons 2 and 3 sequences) but not with the genomic sequence, which suggests that the presence of an intron prevented the identification of a fold-back structure in the genomic locus of miR444 (Figure 3). The mature miR444 resides in the 3rd exon of the gene J033125N22. The ORF of this gene is predicted to code for an unknown protein of 50 amino acids. It is possible though that the processed mRNA is just a pri-miRNA444 transcript and does not code for a protein.

[0050] miR444 is conserved in monocots such as wheat, barley, maize, sorghum and sugarcane but not in *Arabidopsis* (Figure 4A). The precursor sequences from all these plants can form a hairpin structure (Figure 4B). Unlike the situation in rice, the corresponding miRNA precursor fold-back structures can be predicted from the un-spliced genomic sequences from the available sequences of other monocots.

[0051] Similarly, OsmiR436 also resides in the same polarity of a processed transcript (J023035E19), and only the processed transcript can form a hairpin structure. The mature miR436 resides in the 3rd exon of the gene J023035E19. The predicted fold-back structure requires a very long part (720 nt; exons 3 to 9) of the processed transcript (Figure 5) because of the presence of stem-loop structures protruding from the 3' arm of the hairpin structure.

Expression patterns of rice miRNAs

[0052] The tissue- and development-specific expression of miRNAs might provide clues about their physiological function. In a wide range of organisms, many miRNAs have been found differentially expressed at different developmental stages, cell types and tissues (Lee, R. C. et al., *Science* 294:862-864 (2001); Lagos-Quintana, M. et al., *Curr. Biol.* 12:735-739 (2002); Aravin, A. et al., *Dev. Cell* 5:337-350 (2003); Houbaviy, H. B. et al., *Dev. Cell* 5:351-358 (2003)). Several *Arabidopsis* miRNAs are expressed ubiquitously while the expression of many others are regulated by development and shows preferential accumulation in certain tissues (Reinhart, B. J. et al., *Genes Dev.* 16:1616-1626 (2002); Llave, C. et al., *Plant Cell* 14:1605-1619 (2002a); Park, W. et al., *Curr. Biol.* 12:1484-1495 (2002); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). To assist with the determination of the function of the new rice miRNAs, we examined their expression in different organs and developmental stages (Figure 6 and 7).

[0053] The expression patterns of miR441 and miR442 are similar: moderate expression in leaves, roots and young seedlings, and weaker expression in stems and inflorescences (Figure 6A and 6B). miR435 and miR437 also displayed similar expression patterns: moderate expression in leaves and young seedlings and weaker expression in other tissues tested (Figure 6C and 6D). miR443 and miR446 seem to be strongly expressed in leaves and roots and moderately in stem and inflorescence tissues (Figure 6E and 6F). miR436 is expressed in all tissues tested but the levels are higher in inflorescence and young seedlings (Figure 6G). miR438 expression was moderate in leaves, root, inflorescence and young seedlings (Figure 6H). miR440 is uniformly and abundantly expressed in all rice tissues (Figure 6I), while miR439 shows very weak expression in seedlings and is barely detectable in other tissues (Figure 6J).

[0054] OsmiR396d showed strong and ubiquitous expression in all tissues, although the expression in roots was relatively lower (Figure 7A). miR445 was strongly expressed in mature stems but barely detected in leaf and inflorescence tissues (Figure 7B). OsmiR444 showed uniform expression in all rice tissues examined (Figure 7C). All these three miRNAs (miR396d, miR444 and miR445) were found to be expressed in maize, although the signal was weaker, possibly due to pooled RNA from different tissues of maize (Figure 7A, 7B and 7C).

[0055] Some miRNAs displayed tissue- or developmental stage-specific expression patterns. Particularly interesting expression was observed for miR445 and miR390, which showed preferential expression in stems and roots, respectively (Figure 7B and 7D).

[0056] To help determine whether a miRNA is conserved within monocots, its expression was analyzed in another monocot, maize. The analysis showed that 3 miRNAs (miR396d, miR444 and miR445) are conserved and expressed in maize (Figure 7A, 7B and 7C). The presence of miR396d and miR445 miRNAs in other monocots is supported by their expression in maize, and miR444 by sequence and expression. In addition, miR437 sequence and conserved precursor fold-back structures are present in maize, sugarcane and *Sorghum* (Figure 12). The absence of sequence and expression of these 4 miRNAs in *Arabidopsis* suggests that they may be specific to monocots. As an example of conserved miRNAs between dicots and monocots, we tested the expression of a newly identified and conserved miRNA, miR390, in addition to the previously reported and conserved OsmiR156 and OsmiR171 miRNAs in rice, maize and *Arabidopsis* (Figure 7D, 7E and 7F). In addition to the expected size, the OsmiR171 probe also hybridized to a slightly larger (~23 bp) small RNA species in maize but not in rice or *Arabidopsis*, indicating that this larger species is specific to maize (Figure 7E). The remaining 9 new rice miRNAs appear not conserved between plant species since they have no counterparts in other known plant sequences and their expression could not be detected in maize or *Arabidopsis*.

[0057] In summary, the Northern blot analysis confirmed the expression and sizes of 14 newly identified miRNAs in rice (Figure 6 and 7). The majority is expressed ubiquitously in all tissues.

Predicted targets

[0058] Prediction of plant miRNA targets has been facilitated by their extensive sequence complementarity (Rhoades, M. et al., *Cell* 110:513-520 (2002); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA* 101:11511-11516 (2004); Wang, X. J. et al., *Genome Biol.* 5:R65 (2004b); Adai, A. et al., *Genome Res.* 15:78-91 (2005)). Regulatory targets can be more confidently predicted for conserved miRNAs, because complementary sites often are conserved across species boundaries. To identify the potential targets of our newly identified miRNAs, we used the miRNA sequences to search the rice mRNA sequences for antisense

hits with the PATSCAN program (Dsouza, M. et al., *Trends Genet.* 13:497-498 (1997)). Based on transcriptome analysis in Arabidopsis transgenic plants over-expressing miRNAs, Weigel and colleagues (Schwab, R. et al., *Dev. Cell* (in press) (2005)) devised a set of rules for predicting miRNA targets. These criteria include, allowing one mismatch in the region complementary to nucleotides 2-12 of the miRNA but not at the cleaving site (10 and 11 nt), and 3 additional mismatches were permitted between 12 and 21 nt positions, but no more than 2 continuous mismatches within this region. Adopting these rules in predicting newly identified miRNA targets in rice, we allowed 1 mismatch between the positions 1-9 nt from the 5' end of miRNA, no mismatches between positions 10 and 11, another two mismatches were allowed between positions 12 and 21/24. Gaps and mismatches are commonly seen in known animal and plant miRNA::mRNA base-pairing interactions that are known to lead to cleavage or attenuation of translation (Palatnik, J. F. et al., *Nature* 425:257-263 (2003); Aukerman, M. J. et al., *Plant Cell* 15:2730-2741 (2003); Chen, X. *Science* 303:2022-2025 (2004); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). By applying the above rules, our analysis led to the prediction of 46 genes as putative targets for 11 new miRNAs in rice (Table 3). Predicted targets and their complementarity with the new miRNAs are provided in Table 4. The number of predicted targets per miRNA varied greatly, from 1 to 15. Four of the miRNAs (miR435, miR443, miR444 and miR445) each has only one predicted target. We were unable to predict targets for the remaining 3 miRNAs (miR438, miR440 and miR442) by applying these criteria. To evaluate the false positive rates of our target predictions, we performed the same searches with 100 randomized sequences for each miRNA. The length and composition of the miRNAs was maintained in the randomized sets and the searches were performed with the same mismatch settings (Rhoades, M. et al., *Cell* 110:513-520 (2002); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). According to these results (Table 3) the hit frequency with the authentic miRNAs is in most cases more than 5 times higher and relative.

[0059] In animals, all known miRNA target sites were found in 3'UTR's of protein coding genes, whereas in plants they are only occasionally in the 3'UTRs and but are predominantly in the coding regions (Rhoades, M. et al., *Cell* 110:513-520 (2002); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA* 101:11511-11516 (2004); Wang, X. J. et al., *Genome Biol.* 5:R65 (2004b); Adai, A. et al., *Genome Res.* 15:78-91 (2005)). In plants, they also have

been predicted to reside in 5'UTRs (Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)). Recently, miRNAs have also been predicted to target ORFs in humans (Lewis, B. P. et al., *Cell*, 120:15-20 (2005)). Consistent with the earlier findings in *Arabidopsis*, 30 of our predicted target genes in rice have target sites in their ORFs. Fifteen genes have their predicted target sites in 3' UTR and only one in the 5' UTR.

[0060] Both miR444 and its predicted target, a MADS-box transcription factor gene, are conserved in other monocots such as wheat, maize, barley and sugarcane but not in *Arabidopsis* (data not shown). OsmiR396d, a new member of the miR396 family, is expected to target the GRFs (Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). miR396d displays near perfect complementarity with 15 of these GRF genes (Table 3 and Table 4). The complementary sites of miR396d are highly conserved in GRF genes of sorghum, maize, barley and sugarcane.

[0061] The predicted target of miR439, 11667.m02576, a dirigent-like protein gene, was found to have 3 complementary sites within its ORF (Figure 8). These 3 sites are very closely spaced and separated by gaps of 11 and 9 nt. One target site corresponding to the positions 567-588 is perfectly complementary to the miRNA. The other two target sites correspond to the positions 599-619 and 628-648, with 2 and 4 mismatches, respectively (Figure 8). The two target sites corresponding to positions 567-588 and 628-648 are in frame and there is partial amino acid sequence conservation between these two target sites (Figure 8). The presence of 3 target sites within one ORF has not been seen before and thus is unique among miRNA targets predicted thus far.

[0062] It appears that our predicted targets have roles not only in development but also in diverse physiological processes. Sixteen of the predicted targets of 2 miRNAs (miR396d and miR444) are transcription factors (GRL transcription factors and MADS-box transcription factor), whereas the remaining 30 predicted targets of 9 miRNAs appear to have roles in a broad range of physiological processes, and include protein kinases, F-box proteins, dirigent-like protein, glutamate receptor-like proteins, RNA binding protein, retrotransposon and 17 other proteins with unknown function (Table 3).

Identification of miRNA-guided cleavage of target mRNAs in rice

[0063] miRNAs negatively regulate target genes, through miRNA-directed cleavage within the region of complementarity or interfering with translation (Carrington, J. C. et al. *Science*

301:336-338 (2003); Bartel, D. P. *Cell* 116:281-297 (2004); Ambros, V. *Nature* 431:350-355 (2004); Dugas, D. V. et al., *Curr. Opin. Plant Biol.* 7:512-520 (2004); Mallory, A. C. et al., *Curr. Opin. Plant Biol.* 7:120-125 (2004a)). Most of the *Arabidopsis* miRNAs have been shown to guide the cleavage of target mRNAs (Llave, C. et al., *Science* 297:2053-2056 (2002b); Tang, G. et al., *Genes Dev.* 17:49-63 (2003); Palatnik, J. F. et al., *Nature* 425:257-263 (2003); Kasschau, K. et al., *Dev. Cell* 4:205-217 (2003); Xie, Z. et al., *Curr. Biol.* 13:784-789 (2003); Mallory, A. C. et al., *Curr. Biol.* 14:1035-1046 (2004b); Mallory, A. C. et al., *EMBO J.* 23:3356-3364 (2004c); Floyd, S. K. et al., *Nature* 428:485-486 (2004); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004); Wang, X. J. et al., *Genome Biol.* 5:R65 (2004b)). To test whether the predicted miRNA targets in rice can also be cleaved, we used a RNA ligase-mediated 5' RACE procedure (Llave, C. et al., *Science* 297:2053-2056 (2002b)) to map the cleavage sites in the predicted target genes from rice. We performed the 5'RACE assays on four predicted target genes: two were representatives of targets of conserved miRNAs (11668.m00935 targeted by OsmiR390; 7448.m00137 targeted by OsmiR408); one is a MADS-box factor (11668.m04852) which is a predicted target of OsmiR444 as a representative of monocot-specific miRNA target; and the other target served as a representative of rice specific miRNA target (11686.m04227 targeted by OsmiR436). All these four predicted targets were found to have specific cleavage sites corresponding to the miRNA complementary sequences (Figure 9A, 9B, 9C and 9D). In all these cases, the most common 5' end of the mRNA fragments mapped to the nucleotides that pair with the 10th miRNA nucleotides from their 5' ends.

Experimental verification of previously predicted miRNAs in rice

[0064] With the exception of a member of the miR171 family, none of the predicted miRNAs in rice have been verified experimentally. The miR171b homolog in rice has been cloned in a recent study (Wang, J-F. et al., *Nucleic Acids Res.* 32:1688-1695 (2004a)). Sequence similarity searches show that 21 of the cloned miRNAs (in 15 families) in this study had been predicted in rice (Table 2). These include OsmiR156a, OsmiR156k, OsmiR159a/b, OsmiR159c, OsmiR160, OsmiR164, OsmiR166a-f, OsmiR167a-c, OsmiR167d-l, OsmiR168a, OsmiR169b/c, OsmiR169f/g, OsmiR169h-m, OsmiR171a-f, OsmiR171g, OsmiR172a, OsmiR393, OsmiR397, OsmiR398, OsmiR399a and OsmiR408 (Table 2). We noticed that all predicted members of 4 miRNA families -- OsmiR169 (3 members), OsmiR156 (2 members), OsmiR159 (2 members) and OsmiR171 (2 members) --

have appeared in our libraries (Table 2). Thus, our cloning supports the expression of all members of these 4 miRNA families in rice.

[0065] The frequency of cloning varies highly among miRNAs. Our analysis indicates that OsmiR168a is the most abundant miRNA in rice, which was represented by 22 clones in the library. Of the 22 clones, 14, 5 and 3 came from shoot, root and inflorescence libraries, respectively. The OsmiR168 family was represented by 2 members (168a and 168b) that differ slightly in nucleotide sequence. All 22 clones belong to OsmiR168a and none to miR168b, which suggests that only miR168a is abundantly expressed, and miR168b is much lower in abundance or its expression may be limited to specific cells or tissues. An equally abundant miRNA family is the OsmiR156 family, with multiple (12) loci in the rice genome, which appeared 22 times in our sequencing. Seventeen clones corresponded to the OsmiR156a-j loci, whereas the other 5 clones belong to another homolog, miR156k/l. Slightly less abundant miRNAs are members of the OsmiR396 and OsmiR169 families, each represented by 10 and 9 clones, respectively. Interestingly, none of the 10 clones represented the predicted member of the OsmiR396 family; rather, all these 10 clones corresponded to the new member, OsmiR396d, identified in this study. Other miRNAs, OsmiR159, OsmiR164, OsmiR166, OsmiR167, OsmiR169, OsmiR171, OsmiR172 and OsmiR397, were cloned only a few times (2 to 8), whereas the rest (OsmiR160, OsmiR393, OsmiR398, OsmiR399 and OsmiR408) appeared only once.

[0066] In addition to the 14 new miRNAs presented in the Table 1, we also identified 5 other small RNAs (Table 5) encoded by the rice genome and have some features of miRNAs (derived from hairpin RNA precursors) (Figure 13). However, we are unable to designate them as miRNAs at the present time as they could not be detected in the Northern analysis. Due to the lack of genetic tools in rice for validating miRNA biogenesis, such as mutants defective in DCL genes, future research is required to confidently designate these small RNAs as either miRNAs or siRNAs.

DISCUSSION

Monocot-specific and rice-specific miRNAs

[0067] In animals, most miRNAs are conserved across species boundaries (Ambros, V. *Nature* 431:350-355 (2004); Bartel, D. P. *Cell* 116:281-297 (2004)). The strict conservation of miRNAs observed might suggest that the interactions between these miRNAs and their

targets constitute essential processes and could play evolutionarily conserved roles (Pasquinelli, A. E. et al., *Nature*. **408**:86-89 (2000)). However, a few miRNAs appear to be species specific (Ambros, V. et al., *Curr. Biol.* **13**:807-818 (2003a)). miRNAs that are restricted to one or a few species may be implicated in species- or clade-specific functions. The identification of several miRNAs in *Arabidopsis* that are not conserved in rice suggests that these may have specific roles in *Arabidopsis*. These observations strongly supported the hypothesis that rice may also have a unique set of miRNAs. Thus far, 20 families of miRNAs discovered in *Arabidopsis* are predicted to be conserved in rice (Reinhart, B. J. et al., *Genes Dev.* **16**:1616-1626 (2002); Park, W. et al., *Curr. Biol.* **12**:1484-1495 (2002); Jones-Rhoades, M. J. et al., *Mol. Cell* **14**:787-799 (2004); Sunkar, R. et al., *Plant Cell* **16**:2001-2019 (2004); Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA*. **101**:11511-11516 (2004); Wang, X. J. et al., *Genome Biol.* **5**:R65 (2004b); Adai, A. et al., *Genome Res.* **15**:78-91 (2005)). Of these, we have experimentally verified the expression of 15 families of miRNAs in rice. To a certain extent, the 75% coverage of predicted miRNAs in our libraries may reflect the extent of saturation of our miRNA cloning. Of the 14 new rice miRNAs we identified, only 1 (miR390) is conserved in both *Arabidopsis* and maize, whereas 4 others (miR396d, miR437, miR444 and miR445) are conserved in other monocots such as maize, barley, sugarcane and *Sorghum* but not in *Arabidopsis*, which suggests that these latter ones are specific to monocots, and the remaining 9 miRNAs are likely specific to rice. All of the new miRNAs are conserved in another rice subspecies, *Indica* (Table 1). This is expected given the extensive sequence co-linearity between these two rice subspecies (Feng, Q. et al., *Nature* **420**:316-320 (2002)).

[0068] Compared to computational approaches, direct cloning has the advantage of identifying not only non-conserved miRNAs, but also atypical miRNAs. Cloning of miRNAs in rice resulted in the identification of an additional member (from the OsmiR396d and OsmiR396e loci) of the OsmiR396 family. The presence of an extra nucleotide in the middle of a miRNA from the same family was previously unknown. The identification of a new member of the OsmiR396 family with the added nucleotide in the miRNA sequences suggests that future computational approaches should incorporate this structural feature in their prediction strategies. It is interesting that this new member is missing from the *Arabidopsis* miR396 family.

Unusual processing and ubiquitous expression patterns of rice miRNAs

[0069] Although the majority of the new miRNAs are mapped to intergenic regions, a considerable fraction maps to the introns, whereas 2 miRNAs (OsmiR436 and OsmiR444) map to exons of protein-coding genes. The expression of exon- and intron-derived miRNAs in the sense polarity is most likely driven by the promoters of the surrounding genes, whereas those in the antisense polarity within the introns are likely to be expressed as independent genes.

[0070] The structural analysis of full-length transcripts of pre-miRNA transcripts of miR172 revealed the presence of introns, and the processed transcripts are 5' capped and polyadenylated (Aukerman, M. J. et al., *Plant Cell* 15:2730-2741 (2003)). A fold-back structure can be readily predicted for the AthmiR172 precursor from genomic sequence since the hairpin precursor resides in one exon. Nevertheless, this observation suggests that certain miRNA precursor sequences can enter the spliceosomal pathway. Consistent with this, we found that one of the newly identified miRNA precursors (OsmiR444) requires part of exons 2 and 3 of a protein-coding gene for hairpin formation. This shows that some miRNA precursors undergo processing and that processed transcripts are necessary for the prediction of the hairpin structure. Similarly, a hairpin structure for miR436 could only be predicted from a processed transcript, which suggests the presence of introns within the hairpin sequence. These findings raise the possibility that some other miRNA transcripts might also have introns within the hairpin precursor sequences and it would be impossible to predict these miRNAs if the sequences of processed transcripts are not available.

[0071] Although the expression of several miRNAs from *Arabidopsis* are shown to be constitutive and ubiquitous, many miRNAs exhibit preferential expression in a temporal or tissue-specific manner (Reinhart, B. J. et al., *Genes Dev.* 16:1616-1626 (2002); Park, W. et al., *Curr. Biol.* 12:1484-1495 (2002); Llave, C. et al., *Plant Cell* 14:1605-1619 (2002a); Aukerman, M. J. et al., *Plant Cell* 15:2730-2741 (2003); Chen, X. *Science* 303:2022-2025 (2004); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)), while the expression of others is modulated by abiotic stress (Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)) or nutritional status such as sulfur starvation (Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004)). Most of the previously reported miRNAs were shown to be abundantly expressed in inflorescence tissues in *Arabidopsis* (Reinhart, B. J. et al., *Genes Dev.* 16:1616-1626 (2002); Llave, C. et al., *Plant*

Cell 14:1605-1619 (2002a); Park, W. et al., *Curr. Biol.* 12:1484-1495 (2002); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)). However, the expression of most of the newly identified rice miRNAs does not show higher expression in inflorescence tissues. In fact, with the exception of OsmiR436, all others seem to have lower levels of expression in the inflorescence compared to other tissues. Additionally, most of the newly identified rice miRNAs are ubiquitously expressed and few showed tissue specific expression.

Predicted targets include transcription factors and others with a broad range of biological functions

[0072] Over one third of the genes in the human genome were predicted as miRNA targets and these targets appear to be involved in a wide range of biological functions (Lewis, B. P. et al., *Cell*, 120:15-20 (2005)). Previously reported miRNAs in *Arabidopsis* have targets that are predominantly transcription factors or parts of the RNAi machinery such as DCL-1 and AGO1 (Xie, Z. et al., *Curr. Biol.* 13:784-789 (2003); Vaucheret, H. et al., *Genes Dev.* 18:1187-1197 (2004)). More recently identified miRNAs in *Arabidopsis* may target transcripts that encode proteins involved in diverse physiological processes (Jones-Rhoades, M. J. et al., *Mol. Cell* 14:787-799 (2004); Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004); Bonnet, E. et al., *Proc. Natl. Acad. Sci. USA.* 101:11511-11516 (2004); Wang, X. J. et al., *Genome Biol.* 5:R65 (2004b); Adai, A. et al., *Genome Res.* 15:78-91 (2005)).

[0073] We were able to predict 46 genes as potential targets of 11 newly identified miRNAs in rice. Transcription factors (GRL and MADS-box gene) represented one-third of the predicted targets of miRNAs in the present study. The remaining 30 predicted targets appear to have roles in diverse physiological processes. For instance, F-box proteins that regulate diverse cellular processes, including cell cycle regulation, circadian rhythm, flower development and hormonal signal transduction (Kuroda, H. et al., *Plant Cell Physiol.* 43:1073-1085 (2002)), are predicted targets of 3 miRNAs (miR435, miR441 and miR446). OsmiR439 is predicted to target a dirigent-like protein (Table 3). Thus far, dirigent proteins and their homologs have only been reported in vascular plants (Davin, L. B. et al., *Plant Physiol.* 123:453-461 (2000)). Although the precise role of these proteins is not known, they seem to function in the lignification in vascular plants (Burlat, V. et al., *Phytochem.* 57:883-897 (2001)). Glutamate receptor proteins are probable targets of OsmiR437 (Table 3). Ionotropic glutamate receptors function in animals as glutamate-gated non-selective cation

channels. Recent evidence suggests that plant glutamate-receptor-like (GLR) genes do encode non-selective cation channels, but these channels are not gated by amino acids and therefore their roles in plants remain a mystery (Davenport, R., *Ann. Bot.* **90**:549-557 (2002)). OsmiR390 is predicted to target 3 leucine rich-repeat protein kinases (Table 3). Fifteen unknown proteins are also predicted targets of newly identified miRNAs (Table 3). The identification of miRNAs that target genes with unknown functions provides a unique tool to probe the functions of these unknown genes.

[0074] The MADS-box protein family is represented by >70 and >100 genes in rice and *Arabidopsis*, respectively. No miRNA that targets the MADS-box transcription factors has been identified previously, either in *Arabidopsis* or rice. In this study, we found a miRNA (miR444) that targets a MADS-box transcription factor gene in rice, and both the miRNA and the target sites in the MADS-box genes are conserved in monocots. MADS-domain proteins in plants were first identified as regulators of floral organ identity and have since been found to control additional developmental processes such as meristem identity, root development, fruit dehiscence, and flowering time (Coen, E. S. et al., *Nature* **353**:31-37 (1991); Weigel, D. et al., *Cell* **78**:203-209 (1994); Riechmann, J. L. et al., *Biol. Chem.* **378**:1079-1101 (1997); Theissen, G. et al., *Plant Mol. Biol.* **42**:115-149 (2000)). miR444 precursors can be found in other monocot species such as maize, wheat, barley, and sugarcane from expression databases. The lack of a miR444 homolog and of the conserved target site in a MADS-box gene(s) from *Arabidopsis* strongly suggests that this miRNA-mediated regulation of MADS-box genes is conserved only in monocots. Future manipulation of OsmiR444 level and of the target MADS-box genes should help unravel the importance of these interactions in rice and in monocots in general.

[0075] To our knowledge, no plant or animal miRNA has been previously predicted to have multiple target sites in one ORF. In the case of OsmiR439, one target site is perfectly complementary and the other two target sites have 2 and 4 mismatches including in positions 9 and 11 from the 5' end of the miRNA. The mismatch at position 11 may suggest that the interaction does not lead to cleavage, since perfect match at positions 10 and 11 are important for cleavage of the target (Jones-Rhoades, M. J. et al., *Mol. Cell* **14**:787-799 (2004); Schwab, R. et al., *Dev. Cell* (in press) (2005)). However, more recently it is shown that a mismatch at position 11 reduces the cleavage rate but still can be tolerated *in vivo* (Mallory, A. C. et al., *EMBO J.* **23**:3356-3364 (2004c)). The identification of miR439, with 3 complementary

target sites within the ORF of a target gene (Figure 8), suggests that the cooperative regulation of 1 gene at 3 different sites by 1 miRNA may be important to downregulate the mRNA to the required level. Alternatively, despite its perfect complementarity with the target gene at one of the target sites, this miRNA might be involved in translational repression for which more than 1 target site may be necessary. Or, 1 target site may be involved in cleavage and the others may be important for translational repression. Future functional studies with introduced silent mutations in the target sites may reveal the importance of 3 target sites in 1 ORF and may shed light on potentially unique regulatory mechanisms.

miRNA-guided cleavage of target mRNAs in rice

[0076] The cleavage of target mRNAs appears to be the predominant mode of gene regulation by plant miRNAs (Llave, C. et al., *Science* **297**:2053-2056 (2002b); Kasschau, K. et al., *Dev. Cell* **4**:205-217 (2003); Palatnik, J. F. et al., *Nature* **425**:257-263 (2003); Tang, G. et al., *Genes Dev.* **17**:49-63 (2003); Xie, Z. et al., *Curr. Biol.* **13**:784-789 (2003); Jones-Rhoades, M. J. et al., *Mol. Cell* **14**:787-799 (2004); Vaucheret, H. et al., *Genes Dev.* **18**:1187-1197 (2004); Mallory, A. C. et al., *Curr. Biol.* **14**:1035-1046 (2004b); Mallory, A. C. et al., *EMBO J.* **23**:3356-3364 (2004c)). Target mRNA fragments resulting from miRNA-guided cleavage are characterized by a 5' phosphate group, and cleavage occurs near the middle of the base-pairing interaction region with the miRNA (Llave, C. et al., *Science* **297**:2053-2056 (2002b)). We were able to map the cleavage sites for 4 of the predicted targets, and the results support that these are genuine miRNA targets in rice.

[0077] OsmiR408 was cloned in this study (Table 2). Previously, we reported the identification of miR408 from *Arabidopsis* (Sunkar, R. et al., *Plant Cell* **16**:2001-2019 (2004)). miR408 was predicted to target an mRNA encoding a plantacyanin (7448.m00137). Its target site is conserved in several plant species (Sunkar, R. et al., *Plant Cell* **16**:2001-2019 (2004)). In the present study, we mapped the miR408 cleavage site in the rice plantacyanin transcript by using a modified 5' RACE procedure. The result validated plantacyanin as a genuine target of OsmiR408. The precise function of plantacyanins in plants is unknown; however, they have been proposed to function in cell-to-cell signaling, lignin formation and stress responses (Romo, S. et al., *Plant Physiol. Biochem.* **39**:1017-1026 (2001); Kim, S. et al., *Proc. Natl. Acad. Sci. USA*, **100**:16125-16130 (2003)). Our ability to map the cleavage

products that correspond to the OsmiR390 complementary site in a Ser/Thr/Tyr-protein kinase (11668.m00935) confirms that this kinase is indeed a miRNA target. This is the first demonstration that protein kinases can be targeted by miRNAs. We also provide evidence that a rice MADS-box transcription factor (11668.m04852) is a genuine target of OsmiR444. One of the predicted targets (11686.m04227 an unknown protein) of miR436 is also confirmed by the cleavage analysis.

[0078] In summary, our cloning study led to the discovery of 14 new rice miRNAs, of which only one (miR390) is conserved in *Arabidopsis*. At least four of the miRNAs (miR396d, miR437, miR444 and miR445) appear to be limited to monocots, suggesting monocot-specific miRNA-dependent regulatory processes. It is possible that more of the new miRNAs are present in other monocots whose sequence information is not available. It is also possible that more of these miRNAs may be conserved in maize but cannot be detected with the rice specific probes because of possible subtle sequence differences. The putative monocot-specific miRNAs may have evolved after the divergence of monocots and dicots 200 million years ago, or they may have been lost in dicots.

METHODS

Cloning of miRNAs from rice

[0079] Total RNA was isolated separately from shoots and roots of 4-week-old young seedlings and inflorescence of adult rice (*Oryza sativa* spp *japonica* cv. Nipponbare) plants using TRIzol (Bethesda Research laboratories, Life technologies) according to the manufacturer's instructions. Cloning of miRNAs was performed as described (Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)). In brief, small RNAs from 18 to 26 nt were size fractionated, purified, and ligated sequentially to 5' and 3' RNA/DNA chimeric oligonucleotide adapters. Reverse transcription was performed after ligation with the adapters, followed by PCR amplification. The resulting PCR products were cloned and transformed into competent cells. Plasmids were isolated from individual colonies and sequenced. The sequences were subsequently processed and used for BLAST analysis against the *Oryza sativa* spp *japonica* sequences in the TIGR database and other plant sequences in the NCBI database.

Sequence analysis and prediction of fold-back structures

[0080] Automated base calling of raw sequence traces and vector removal was performed with the PHRED and CROSS MATCH programs from Ewing and Green (1998). All trimmed sequences longer than 16 nt were used to search the Rfam (www.sanger.ac.uk/Software/Rfam) database with BLASTN (Altschul, S. F. et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). This step allowed the removal of most non-siRNA and non-miRNA species in the dataset. Putative origins for the remaining sequences were identified by BLASTN searches against the intergenic and intron sequences using the latest annotation 3.0 of *Oryza sativa* spp. *japonica* from The Institute for Genomic Research (TIGR, ftp://ftp.tigr.org/pub/data/Eukaryotic_Projects/o_sativa/). Candidates with perfect matches against these genomic datasets were used for fold-back secondary structure prediction with the mfold program (Zuker, M. *Nucleic Acids Res.* 31:3406-3415 (2003)). Three different combinations of upstream and downstream sequence spanning the putative miRNA origins were used for the predictions: (1) 400 bp upstream and 20-30 bp downstream (2) the same length combination but in reverse order and (3) 200 bp upstream and 200 bp downstream. In the case of OsmiR444 and OsmiR436, only processed transcript sequences were used to predict the hairpin structure. OsmiR436 hairpin prediction required 700 nt.

[0081] To identify putative target sequences, all predicted and cloned CDS and UTR sequences from *O. sativa* (TIGR all.cdna set) were searched with PatScan (Dsouza, M. et al., *Trends Genet.* 13:497-498 (1997)). The following parameters were used for these pattern searches, all referring to the 5' end of the miRNAs in antisense orientation: not more than one mismatch between positions 1-9, no mismatch between positions 10-11 and not more than two mismatches for the rest of the sequence. False positive rates of predicted targets were estimated with the same search strategy using 100 randomized samples for each miRNA. The retrieved miRNA/target site pairs were ranked and scored by aligning them with the Needleman-Wunsch global alignment program from EMBOSS (Rice, P. et al., *Trends Genet.* 16:276-277 (2000)). Global cross-species conservations of miRNAs were determined by BLASTN searches against the NT and EST databases from NCBI. The results were parsed with a Perl script that scored only those pairwise BLAST alignments (HSP) as hits when they were not more than two positions shorter than the query sequence and 95% identical. More detailed conservations of miRNAs and target sequences within the genome sequences of *Populus trichocarpa*, *O. sativa* spp. *indica* and *Arabidopsis thaliana* were determined with

the same BLASTN and PatScan strategy used for the japonica host. Weaker conserved origins in the genomic sequences of *Arabidopsis* and *Populus trichocarpa* were identified with PatScan (maximal two mismatches) and subsequent fold-back structure predictions.

RNA blot analysis

[0082] Total RNA was isolated from leaf, stem, root and inflorescence tissues of adult plants as well as from 4- week-old young rice seedlings. Total RNA was also isolated from pooled maize tissues (kernels, pericarp, endosperm, scutellum, coleoptile, embryo, silk and young seedlings) and 3-week-old *Arabidopsis* seedlings using TRIzol (Bethesda Research laboratories, Life technologies). One hundred micrograms of total RNA was loaded per lane and resolved on a denaturing 15% polyacrylamide gel and transferred electrophoretically to Hybond-N⁺ membranes (Amersham Biosciences, Bucks, UK). Hybridization and washings were performed as described (Sunkar, R. et al., *Plant Cell* 16:2001-2019 (2004)). The membranes were briefly air dried and then exposed to a phosphorimager.

5' RACE

[0083] Total RNA from 4-week-old rice seedlings was extracted with Trizol reagent. Poly(A)⁺ mRNA was purified from total RNA with use of the Poly A kit (Promega). RLM-5' RACE (RNA ligase mediated -5' rapid amplification of cDNA ends) was carried out with use of the GeneRacer Kit (Invitrogen Life Technologies, Carlsbad, CA, USA). The GeneRacer RNA Oligo adapter was directly ligated to mRNA (100 ng) without calf intestinal phosphatase and tobacco acid pyrophosphatase treatment. Gene specific primers were designed and used for cDNA synthesis. Initial PCR was carried out with the GeneRacer 5' primer and gene specific primers (7448.m00137, 11668.m00935, 11668.m04852 and 11686.m04227). Nested PCR was carried out with 1 µL of the initial PCR reaction, GeneRace 5' nested primer, and gene specific internal primers. After amplification, 5' RACE products were gel-purified and cloned, and at least 10 independent clones were sequenced.

Table 1. New rice miRNAs identified by cloning.

miRNA	Sequence (5' → 3')	RNA blot	size (nt)	No. of loci	Chr.	Fold-Back (5'/3')	intergenic/intron/exon	Osi	Zm	At	Pt
<i>miR390</i>	AAGCUCAGGAGGAUAGCGCC (2)	+	21	1	3	5'	intergenic	+	+	+	+
<i>miR396d</i> <i>miR396e</i>	UCCACAGGCUUUCUUGAACUG (10)	+	21	2	4 2	5' 5'	intergenic intergenic	+	+	-	-
<i>miR435</i>	UUAUCCGGUAUUGGAGUUGA (2)	+	20	1	3	3'	11669.m03227 Intron 7(s)	+	-	-	-
<i>miR436</i>	UGAGAGAGUGCACUUUCUCCC (2)	+	21	1	2	5'	J023035E19 (s)	+	-	-	-
<i>miR437</i>	AAAGUUAGAGAAGUUUGACUU (2)	+	21	1	2	3'	11668.m02729 Intron 15(s)	+	+	-	-
<i>miR438</i>	UUCCCCACGCGUUAUAGUGAAA (1)	+	21	1	6	5'	intergenic	+	-	-	-
<i>miR439a</i> <i>miR439b</i> <i>miR439c</i> <i>miR439d</i> <i>miR439e</i> <i>miR439f</i> <i>miR439g</i> <i>miR439h</i> <i>miR439i</i> <i>miR439j</i>	UGUCGAACCGCG GUUGUUCGA (1)	+	21	10	1 10 1 3 7 8 8 6 9 10	3' 3' 3' 3' 3' 3' 3' 3' 3' 3'	intergenic intergenic intergenic intergenic intergenic intergenic intergenic intergenic intergenic intergenic 11676.m02626 Intron 1(s)	+	-	-	-
<i>miR440</i>	AGUGUCUCCUGAUGAUCGGGACAA (2)	+	24	1	11	3'	11687.m01522 _Intron_9(s)	+	-	-	-
<i>miR441a</i> <i>miR441b</i> <i>miR441c</i> <i>miR442</i>	UACCAUCAUAUAAUUGGGGAAA (3) UGACGUGUAAAUUGCGAGACGAU (2)	+	24	3 1	1 3 7 4	3' 3' 3' 5'	intergenic intergenic intergenic 11670.m05347 Intron 1(s)	+	-	-	-

miR443	AUCACAAUACAUAUAAUUCUGGA (2)	+	22	1	3	5'	intergenic	+	-	-	-
miR444	UUGCUGCCUCAAGCUUGCUGC (3)	+	21	1	8	3'	J033125N22 (s)	+	+	-	-
miR445a	UAAAUUAGUGUAUAAACAUCCGAU (2)	+	24	9	5	3'	intergenic	+	+	-	-
miR445b					6	3'	intergenic				
miR445c					12	3'	intergenic				
miR445d					3	3'	intergenic				
miR445e					7	3'	intergenic				
miR445f					3	3'	intergenic				
miR445g					6	3'	intergenic				
miR445h					3	3'	11669.m00786				
miR445i					6	3'	Intron_2(as) 11680.m00766				
miR446	CAUCAUAUGAAUAUGGGAAAUUGG (2)	+	24	1	6	3'	Intron_1(s) intergenic	+	-	-	-

Frequency of cloning is indicated in parentheses after the miRNA sequences. Chromosomal (Chr.) position is indicated. miRNA location in the predicted fold-back structure is specified (5' or 3' arm). miRNAs that are conserved in other plants are indicated. Osi: *Oryza sativa* subspecies *indica*; Zm: *Zea mays*; At: *Arabidopsis thaliana*; Pt: *Populus trichocarpa*

Table 2. Experimental verification of previously predicted miRNAs in rice.

Gene	miRNA sequence	Cloning frequencies in libraries
OsmiR156a-j	UGACAGAAGAGAGUGAGCAC	Shoot (12), Root (5)
OsmiR156k,l	UGACAGAAGAGAGUGAGCACA	Shoot (3), Root (1), Inflorescence (1)
OsmiR 159a,b	UUUGGAUUGAAGGGAGCUCU	Shoot (3), Inflorescence (2)
OsmiR159c	UUGGAUUGAAGGGAGCUCUGC	Shoot (2)
OsmiR160	UGCCUGGCUCCCUGUAUGCCA	Shoot (1)
OsmiR164	AUGGAGAAGCAGGGCACGUGCA	Shoot (3), Inflorescence (3)
OsmiR166a-f	UCGGACCAGGCUUCAUCCCC	Shoot (2), Inflorescence (1)
OsmiR167a,b,c	UGAAGCUGCCAGCAUGAUCUA	Shoot (2)
OsmiR167d-I	UGAAGCUGCCAGCAUGAUCUG	Root (1)
OsmiR168a	UCGCUUGGUGCAGAUCCGGAC	Shoot (14), Root (3), Inflorescence (5)
OsmiR169b,c	CAGCCAAGGAUGACUUGCCGG	Shoot (2)
OsmiR169f,g	UAGCCAAGGAUGACUUGCCUA	Shoot (2), Root (1)
OsmiR169h-m	UAGCCAAGGAUGACUUGCCUG	Shoot (1)
OsmiR171a-f	UGAUUGAGCCGUGCCAAUAUC	Shoot (1), Root (1), Inflorescence (2)
OsmiR171g	GAGGUGAGCCGUGCCAAUAUC	Root (1)
OsmiR172a	AGAAUCUUGAUGAUGCUGCAU	Shoot (2), Inflorescence (2)
OsmiR393	UCCAAAGGGAUCGCAUUGAUC	Inflorescence (1)
OsmiR397b	UUAUUGAGUGCAGCGUUGAUG	Shoot (1), Root (1)
OsmiR398	UGUGUUCUCAGGUCGCCCCUG	Shoot (1)
OsmiR399a	UGCCAAAGGAGAAUUGCCCUG	Shoot (1)
OsmiR408	CUGCACUGCCUCUCCCCUGGC	Root (1)

Table 3. Predicted targets of newly identified rice miRNAs. For each potential target gene, the number of mismatches between the miRNA and mRNA is indicated in parentheses. Annotated rice mRNAs were also searched for sites complementary to randomized miRNAs (100 repeats) applying the same criteria that were used for predicting targets. The mean of the hit frequency of the randomized data and their standard deviations are recorded.

miRNA	Target gene	Target site	Target protein	Count	Mean	SD
<i>MIR390</i>	11668.m00935(1); 11674.m03433(3); 11687.m03320(3)	ORF	LRR proteins	3	0.24	0.7
<i>MIR396d</i>	11668.m04403(1); 11668.m04580(1); 11669.m04718(1); 11670.m04997(1); 11670.m04998(1); 11680.m00165(1); 11680.m00999(1); 11669.m05248(1); 11670.m02258(1); 11687.m03206(1); 11686.m02909(1); 11687.m03206(2); 11670.m04996(1); 11680.m00166(1); 11670.m04758(1)	ORF	Growth regulating factor (GRL) transcription factors	15	0.19	0.6
<i>MIR435</i>	11687.m03448(3)	ORF	F-box protein	1	0.75	1.3
<i>MIR436</i>	11686.m04227(0); 11686.m04226(1)	ORF	unknown proteins	2	0.07	0.3
<i>MIR437</i>	11680.m04605(1); 11680.m04606(1)	ORF	Glutamate receptor proteins	2	0.22	0.6
<i>MIR439</i>	11667.m02576 (3 target sites with 0, 2 and 4 mismatches) 11680.m01869(1); 11668.m02918(3) 11673.m02474(3)	ORF ORF ORF	Dirigent like protein unknown proteins retrotransposon	4	0.15	0.5
<i>MIR441</i>	11669.m05042(3) 11667.m06876(2); 11674.m00346(2); 11674.m00582(2); 11680.m01945(2); 11682.m04930(3); 11687.m00089(3) 11682.m03939(3); 11668.m04341(2) 11686.m03687(3)	ORF 3'UTR 3'UTR 3'UTR 3'UTR 3'UTR	unknown protein unknown proteins unknown proteins unknown proteins F-box protein UDP-glucuronosyl and UDP-glucosyltransferase	10	0	0
<i>MIR443</i>	11687.m04039(3)	ORF	unknown protein	1	0.13	0.4
<i>MIR444</i>	11668.m04852(0) 2.	ORF	MADS-box transcription factor	1	0.19	0.6
<i>MIR445</i>	11680.m05074(3)	5'UTR	RNA-binding protein	1	0	0
<i>MIR446</i>	11668.m04341(2) 11674.m00582(2); 11680.m01945(2); 11682.m04930(3); 11682.m03939(3) 11680.m01730(2)	3'UTR 3'UTR 3'UTR 3'UTR	F-box protein unknown proteins unknown protein unknown plant specific protein	6	0.01	0.1

Table 4. PATSCAN alignments of the new miRNAs and their predicted targets in rice. miRNAs are presented as a reverse complemented form.

WO 2006/105436										PCT/US2006/012046									
miRNA	OSA_cD NA Hit	OSA_cD NA Hit2	Descrip- tion	Position	Length	Identity	Gaps	Score	Needle Alignment										
MIR443	11687.m 04039	Os11g42 970	expressed protein	[771, 792]	22	19/22 (86.4%)	0/22 (0.0%)	83	P101-G11	1	TCCAGATTATTGTATTGTGAT	22							
									11687.m04039	1	 . . .	22							
MIR445	11680.m 05074	Os06g50 890	RNA recognition motif. (a.k.a. RRM, RBD, or RNP domain), putative	[1155, 1178]	24	21/24 (87.5%)	0/24 (0.0%)	93	P103-F2	1	ATCGGATGTTTATACACTAATTTA	24							
									11680.m05074	1	 . . .	24							
MIR446	11668.m 04341	Os02g44 990	F-box domain, putative	[1808, 1831]	24	22/24 (91.7%)	0/24 (0.0%)	102	P105-E11	1	CCATTTCCTCCATATTCATATTGATG	24							
									11668.m04341	1		24							
MIR446	11674.m 00582	Os08g06 500	expressed protein	[4567, 4590]	24	22/24 (91.7%)	0/24 (0.0%)	102	P105-E11	1	CCATTTCCTCCATATTCATATTGATG	24							
									11674.m00582	1		24							
MIR446	11680.m 01730	Os06g17 390	Plant protein family, putative	[2625, 2648]	24	22/24 (91.7%)	0/24 (0.0%)	102	P105-E11	1	CCATTTCCTCCATATTCATATTGATG	24							
									11680.m01730	1		24							
MIR446	11680.m 01945	Os06g19 990	expressed protein	[558, 581]	24	22/24 (91.7%)	0/24 (0.0%)	102	P105-E11	1	CCATTTCCTCCATATTCATATTGATG	24							
									11680.m01945	1		24							
MIR446	11682.m 03939	Os05g41 190	expressed protein	[985, 1008]	24	21/24 (87.5%)	0/24 (0.0%)	93	P105-E11	1	CCATTTCCTCCATATTCATATTGATG	24							
									11682.m03939	1		24							

MIR446	11682.m 04930	Os05g51 140	expressed protein	[1540, 1563]	24	21/24 (87.5%)	0/24 (0.0%)	93	P105-E11	1 CCATTTC	24
										CAATTC	24
MIR437	11680.m 04605	Os06g46 670	Receptor family ligand binding region, putative	[3521, 3541]	21	20/21 (95.2%)	0/21 (0.0%)	96	11680.m04605	1 AAGTCAA	21
										AACTTCTCTAACTTT	21
MIR437	11680.m 04606	Os06g46 670	Receptor family ligand binding region, putative	[3772, 3792]	21	20/21 (95.2%)	0/21 (0.0%)	96	P14-F8	1 AAGTCAA	21
										AACTTCTCTAACTTT	21
MIR390	11674.m 03433	Os08g34 650	Leucine Rich Repeat, putative	[861, 881]	21	18/21 (85.7%)	0/21 (0.0%)	78	P28-A2	1 GCGCTAT	21
										CCCTCCTGAGCTT	21
MIR390	11687.m 03320	Os11g36 140	Leucine Rich Repeat, putative	[1033, 1053]	21	18/21 (85.7%)	0/21 (0.0%)	78	P28-A2	1 GCGCTAT	21
										CCCTCCTGAGCTT	21
MIR439	11667.m 02576	Os01g26 340	Dirigent-like protein, putative	[568, 588]	21	21/21 (100.0%)	0/21 (0.0%)	105	P28-E3	1 TCGAACA	21
										ACCGCGGTTTCGACA	21
MIR439	11680.m 01869	Os06g19 250	hypothetical protein	[579, 599]	21	20/21 (95.2%)	0/21 (0.0%)	96	P28-E3	1 TCGAACA	21
										ACCGCGGTTTCGACA	21
MIR439	11668.m 02918	Os02g30 670	hypothetical protein	[407, 427]	21	18/21 (85.7%)	0/21 (0.0%)	78	P28-E3	1 TCGAACA	21
										ACCGCGGTTTCGACA	21

<i>MIR396d</i>	11668.m 04403	Os02g45 570	expressed protein	[507, 527]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11668.m04403	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11668.m 04580	Os02g47 280	expressed protein	[570, 590]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11668.m04580	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11668.m 05272	Os02g53 690	growth- regulating factor 1	[378, 398]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11668.m05272	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11669.m 04718	Os03g47 140	hypothetical protein	[653, 673]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11669.m04718	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11670.m 04996	Os04g51 190	expressed protein	[535, 555]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11670.m04996	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11670.m 04997	Os04g51 190	expressed protein	[535, 555]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11670.m04997	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11670.m 04998	Os04g51 190	expressed protein	[535, 555]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11670.m04998	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11680.m 00165	Os06g02 560	expressed protein	[605, 625]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11680.m00165	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21
<i>MIR396d</i>	11680.m 00166	Os06g02 560	expressed protein	[605, 625]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7 11680.m00166	1 1	CAGTTCAAGAAAGCCTGTGGA . 1 CCGTTCAAGAAAGCCTGTGGA	21 21

WO 2006/105436

PCT/US2006/012046

MIR396d	11680.m 00999	Os06g10 310	growth- regulating factor 1, putative	[282, 302]	21	20/21 (95.2%)	0/21 (0.0%)	96	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA .	21
	11680.m00999	1	CCGTTCAAGAAAGCCTGTGGA						11680.m00999	1	CCGTTCAAGAAAGCCTGTGGA	21
MIR396d	11669.m 05248	Os03g51 970	putative transcription activator	[419, 439]	21	19/21 (90.5%)	0/21 (0.0%)	87	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA .	21
	11669.m05248	1	CCGTTCAAGAAAGCATGTGGA						11669.m05248	1	CCGTTCAAGAAAGCATGTGGA	21
MIR396d	11670.m 02258	Os04g24 190	growth- regulating factor 1, putative	[827, 847]	21	19/21 (90.5%)	0/21 (0.0%)	87	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA .	21
	11670.m02258	1	CCTTTCAAGAAAGCCTGTGGA						11670.m02258	1	CCTTTCAAGAAAGCCTGTGGA	21
MIR396d	11670.m 04758	Os04g48 510	Similar to At3g52910	[858, 878]	21	19/21 (90.5%)	0/21 (0.0%)	87	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA .	21
	11670.m04758	1	CCGTTCAAGAAAGCCTATGGA						11670.m04758	1	CCGTTCAAGAAAGCCTATGGA	21
MIR396d	11686.m 02909	Os12g29 980	expressed protein	[576, 596]	21	19/21 (90.5%)	0/21 (0.0%)	87	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA .	21
	11686.m02909	1	CCGTTCAAGAAAGCATGTGGA						11686.m02909	1	CCGTTCAAGAAAGCATGTGGA	21
MIR396d	11687.m 03206	Os11g35 030	growth- regulating factor, putative	[498, 518]	22	19/22 (86.4%)	2/22 (9.1%)	81	P33-D7	1	CAGTTCAAGAAAGCCTGTGGA 	21
	11687.m03206	1	TC-GTTCAAGAAAGCATGTGGA						11687.m03206	1	TC-GTTCAAGAAAGCATGTGGA	21
MIR441	11667.m 06876	Os01g68 230	expressed protein	[2354, 2377]	24	22/24 (91.7%)	0/24 (0.0%)	102	P72-C9	1	TTTCCACATTATATGATGGTA 	24
	11667.m06876	1	TTTCTCACATTATATGATGTTA						11667.m06876	1	TTTCTCACATTATATGATGTTA	24
MIR441	11668.m 04341	Os02g44 990	F-box domain, putative	[1811, 1834]	24	22/24 (91.7%)	0/24 (0.0%)	102	P72-C9	1	TTTCCACATTATATGATGGTA 	24
	11668.m04341	1	TTTCCACATTATATGATGTTA						11668.m04341	1	TTTCCACATTATATGATGTTA	24
MIR441	11674.m 00346	Os08g04 300	expressed protein	[2085, 2108]	24	22/24 (91.7%)	0/24 (0.0%)	102	P72-C9	1	TTTCCACATTATATGATGGTA 	24
	11674.m00346	1	TTTCCACAGATTATATGATGCTA						11674.m00346	1	TTTCCACAGATTATATGATGCTA	24

MIR441	11674.m 00582	Os08g06 500	expressed protein	[4570, 4593]	24	22/24 (91.7%)	0/24 (0.0%)	102	P72-C9	1	TTTCCCACATTTATATTGATGGTA 11674.m00582 1 TTTCCCACATTCATATTGATGTTA	24
MIR441	11680.m 01945	Os06g19 990	expressed protein	[561, 584]	24	22/24 (91.7%)	0/24 (0.0%)	102	P72-C9	1	TTTCCCACATTTATATTGATGGTA 11680.m01945 1 TTTCCCACATTCATATTGATGCTA	24
MIR441	11669.m 05042	Os03g50 070	expressed protein	[1047, 1070]	24	21/24 (87.5%)	0/24 (0.0%)	93	P72-C9	1	TTTCCCACATTTATATTGATGGTA . 11669.m05042 1 TTTCCCACATTCATATTGATGTTA	24
MIR441	11682.m 03939	Os05g41 190	expressed protein	[988, 1011]	24	21/24 (87.5%)	0/24 (0.0%)	93	P72-C9	1	TTTCCCACATTTATATTGATGGTA 11682.m03939 1 TTTCCCACATTCGTATTGATGTTA	24
MIR441	11682.m 04930	Os05g51 140	expressed protein	[1543, 1566]	24	21/24 (87.5%)	0/24 (0.0%)	93	P72-C9	1	TTTCCCACATTTATATTGATGGTA 11682.m04930 1 TTTCCCACATTCGTATTGATGTTA	24
MIR441	11686.m 03687	Os12g37 510	UDP- glucoronosyl and UDP- glucosyl transferase	[1544, 1567]	24	21/24 (87.5%)	0/24 (0.0%)	93	P72-C9	1	TTTCCCACATTTATATTGATGGTA . 11686.m03687 1 TTTTCCCACATATATATTGATGTTA	24
MIR441	11687.m 00089	Os11g01 870	expressed protein	[5345, 5368]	24	21/24 (87.5%)	0/24 (0.0%)	93	P72-C9	1	TTTCCCACATTTATATTGATGGTA . 11687.m00089 1 TTTCTTACATTTATATTGATGTTA	24
MIR436	11686.m 04227	Os12g42 390	hypothetical protein	[12, 32]	21	21/21 (100.0%)	0/21 (0.0%)	105	P8-D2	1	GGGAGAAAGTGCACTCTCTCA 11686.m04227 1 GGGAGAAAGTGCACTCTCTCA	21
MIR436	11686.m 04226	Os12g42 380	expressed protein	[235, 255]	21	20/21 (95.2%)	0/21 (0.0%)	96	P8-D2	1	GGGAGAAAGTGCACTCTCTCA 11686.m04226 1 GGGAGAAAATGCACTCTCTCTCA	21

MIR444	11668.m 04852	Os02g49 840	SRF-type transcription factor (DNA- binding and dimerisation domain), putative	[403, 423]	21	21/21 (100.0%)	0/21 (0.0%)	105	P95-G4	1	GCAGCAAGCTTGAGGCAGCAA 	21
										11668.m04852	1 GCAGCAAGCTTGAGGCAGCAA 	21

Table 5. Potential new miRNAs from rice. Frequency of cloning is indicated in parentheses after the miRNA sequences.

miRNA	Sequence (5' to 3')	RNA blot	size (nt)	No. of loci	Chr.	Foldback (5'/3')	Conservation				
							intergenic/intron/exon	Osi	Zm	At	Pt
<i>P18-F1</i>	UCUCUCUCUCCCUUGAAGGC (1)	-	20	1	11	5'	intergenic	+	-	-	-
<i>P20-A11</i>	CGCGAGGUACCGAUACCU CGA (1)	-	21	1	6	3'	JNBa0052G07.19 (s)	+	-	-	-
<i>P83-E9</i>	UCAGUGUCAUACGAUAUGGCGC (1)	-	22	1	8	3'	intergenic	-	-	-	-
<i>P92-A7</i>	CAUCCUAAAUAUAAGCAGUUUUUAG (1)	-	25	1	11	5'	intergenic	+	-	-	-
<i>P101-H12</i>	AUAGUGAACACCGAU AUGCGU (1)	-	21	1	11	3'	intergenic	+	-	-	-

Osi: *Oryza sativa* subspecies *indica* ; Zm: *Zea mays* ; At: *Arabidopsis thaliana* ; Pt: *Populus trichocarpa*

WHAT IS CLAIMED IS:

- 1 1. An isolated miRNA as show in Table 1, Table 2, Table 4, or Table 5.
- 1 2. A recombinant expression cassette encoding the miRNA of claim 1.
- 1 3. A transgenic plant comprising the expression cassette of claim 2.
- 1 4. A method of inhibiting down-regulation of a target coding region, the
2 method comprising mutating at least one nucleotide in the coding region to render the coding
3 region resistant to miRNA mediated down-regulation.




Fig. 2

A.		
OsmiR396a, b	UUCCACAG-CUUUCUUGAACUG	(21 bp)
OsmiR396c	UUCCACAG-CUUUCUUGAACUU	(21 bp)
P33-D7	UCCACAGGCUUCUUGAACUG	(21 bp)
B.		
OsmiR399a	UGCCAAAGGAG-AAUUGCCCG	(21 bp)
OsmiR399j, K	UGCCAAAGGAG-AGUUGCCCUA	(21 bp)
OsmiR399e	UGCCAAAGGAG-AUUUGCCCG	(21 bp)
OsmiR399g	UGCCAAAGGAG-AUUUGCCCG	(21 bp)
OsmiR399h	UGCCAAAGGAG-ACUUGCCCG	(21 bp)
OsmiR399d, i	UGCCAAAGGAG-AGCUGCCCG	(21 bp)
P7-A07	UGCCAAUGGAGUAGUUGCCCG	(22 bp)

Fig. 3




Fig. 4

A.

Rice UUGCUGCCUCAAGCUUGCUGC
wheat UUGCUGCCUCAAGCUUGCUGC
Barley UUGCUGCCUCAAGCUUGCUGC
Maize UUGUUGCCUCAAGCUUGCUGC
Sorghum UUGUUGCCUCAAGCUUGCUGC
Sugarcane UUGUUGCCUCAAGCUUGCUGC

B.

Rice
U G A U G C- AA
GCAAU GGGGCGAGCAAGCU GAGG GCAACUGCAUUAUUGCAAGAA AU UUGG UG \
CGUUG CCUCGUGGUUGCA CUCC CGUUGACGUAUUGAACGUUCUU UA AGCC AC G
U A GU AU UA

Wheat
-- C A A U U G - -ACACC AUUA C
GGCA CGAGC UGAGGCA CAACUGCAU ACUUGCGGGGAAG CGCAAGUA GG UGC CUUG A
CCGU GUUCG ACUCCGU GUUGACGUA UGAACGUUCUUUC GUGUUCAU CC ACG GAAC A
CU C A C C C G A \ ----- CG-- A

Barley
C AUG ----- C A A A A GU UG AG - -ACACCAACAUUACCGCAA AA
GAUA GC UGGC GSCA CAAGC UGA GCA CAACUGCA ACUUG GGGAA CACAAGUA GG GC \
CUAU CG ACCG CCGU GUUCG ACU CGU GUUGACGU UGAAC UCUUU GUGUUCAU CC CG G
- AA- UUUCUU C A C C GC GU CG A \ ----- CA

Maize
C A G U U GU AAUCAAUUG
GGCA CAAGC UGAGGCAG AACUGCAU ACUUGCAAGAAA CACAA \
CCGU GUUCG ACUCCGU UUGACGUG UGAACGUUCUUU GUGUU C
C A G U UG CAUACAGGA

Sorghum
C A G U U CUCA- - GA GC A
GGCA CAAGC UGAGGCAG AACUGCAU ACUUGCAAGAAAGUCACAAG UUU CAA UG UCUCUAA \
CCGU GUUCG ACUCCGU UUGACGUG UGAACGUUCUUUCGGUGUUC AGA GUU AC AGAGGUU C
C A G U AUACC C A- UA U

Sugarcane
A - UAU U C A G U A----- AAU
GU UUG CA GG GGCA CAAGC UGAGGCAG AACUGCAU ACUUGCAAGAAAGUCACAA AUC U
CG GAC GU CC CCGU GUUCG ACUCCGU UUGACGUG UGAACGUUCUUUCGGUGUU UAG G
A C UUC U C A A G U CAUAC GAC

Fig. 5

10 20 30 40 50 60 70 80
 CUUUAUAUCUCUUGAGAGAGUGCA UUUCUCCU UCCUGCCA CAGUAGUGCCUUU
 GAAUCUUAGAGAACUCUCACACGU AAAGAGGGA AGGACGGU GUUGUCACGGGA
 A U C
 710 700 690 680 670 120
 --AUUUUC GG --G A
 GCUU UUUC CAUC U
 CGAA AAAG GUGG C
 A

Fig. 6




Fig. 7




Fig. 8

11667.m02576 5' 567 UCGAACAAACCGCGUUCGACA 588.....599 UCGAACAAACAGUUCGACA 619 3'
|||||
miRNA P28-E3 3' AGCUUGUUGGCGCCAAAGCUGU 5' 3' AGCUUGUUGGCGCCAAAGCUGU 5'




Figure 10

Predicted fold-back structures of the new miRNA precursors from rice.

OsmiR435 ($\Delta G = -121.6$)

```
CU--      --  A   CG                A   C                CAU  -  .-CCUAGCA      A  UG      .-CC      A  A      AGUCGA  -
AACA
ACUCUU    CUU  GUCUA  UUUCAUCUCUACAACUCCAA  ACUGGA  AAAACAUGUCCCU  CU  UC      CUAGUUUAG  G  GUUUUUGC  AUGUGG  CU  CUUGGCU  CU  UGCAU
A          UGAGAA  GAA  CAGGU  AAAGUAGGGAGUUGAGGUU  UGBCCU  UUUUGUACAGGGA  GA  AG      GAUUAAGUU  C  UAAAAGCG  UGUACC  GA  GAAUCGA  GA  AUGUA
A
UAUU      CU   C   UA                A   A                UUU  U  \  -----      C  GU      \  --      -  -      GUGG--  C
CAGU
```

OsmiR437 ($\Delta G = -100.0$)

```
UG  UU      U      G      A      U      U      AAUUAACAUCUAAAAU      AU      CA-  A
CCCUCUGUUUCAUUAU  UA  GUGUUU  AGUUUUGUU  UAAGUCAAA  UUCUUU  ACUUUGACCAAGUUUGUAG  AAAAU      ACCAA  AAAUA  CU  U
GGGAGACAAGGUUAU  AU  CACAAA  UCAAAACAG  AUUCAGUUU  AAGAGA  UGAAACUGGUUCAAUAUC  UUUUA      UGGUU  UUUUA  GA  U
GU  UC      -      G      G      U      U      GU-----      C-      ACA  G
```

OsmiR438 ($\Delta G = -59.0$)

```
UGUUUU      CAAACU  UUU                U                -      UG  .-AUG  GUAU      A
AACG          GGG      UG  UCCCACGCGUUA  AGUGAAAACUUUUGGAAU  AUGUU  UU      GGA      AUAUU  \
UUGC          CCU      AC  AGGGUGCGCAAU  UCACUUUUGAAAACUUUG  UACAG  GG      UCU      UAUAG  U
CU-----      UCAUCU  UAU                U                G      GU  \  ---      AGGU      U
```

OsmiR390 ($\Delta G = -55.9$)

```
U          U  U  A          G          CGAAA
GGUA  GGAACAA  CC  UG  AGCUCAGGA  GGAUAGCGCCU      U
CCAU  UCUUGUU  GG  AC  UCGAGUCCU  UCUAUCGCGGA      C
-          .  U  U  C          A          UCAAA
```

OsmiR439a ($\Delta G = -61.9$)

```
UGA      A          U--      UGU
UACCCUGUCGAAC      CGC  GUUCGACA      GUACCUGUCGAAC  \
GUGGGACAGCUUG      GCG  CAAGCUGU      UAUGGAUAGCUUG  G
UUG      C          CCC      UUG
```

OsmiR439b

```
UU          UG  G  U          U
CCUGUCGAAU  U  GU  GUUCGAUAGG  A
GGACAGCUUG  G  CG  CAAGCUGUCC  C
--          UU  G  C          C
```

OsmiR439c

```
UGA      A          U--      UGU
UACCCUGUCGAAC      CGC  GUUCGACA      GUACCUGUCGAAC  \
GUGGGACAGCUUG      GCG  CAAGCUGU      UAUGGAUAGCUUG  G
UUG      C          CCC      UUG
```


OsmiR439d

```

CCACC          UCA UA          U--          UGU
      UACCCUGUCGAAC  CG GUUCGACA  GUACCUGUCGAAC  \
      GUGGGAUAGCUUG  GC CAAGCUGU  UAUGGAUAGCUUG  G
-----          UUG GC          CCC          UUG

```

OsmiR439e

```

CCACC          UGA A          U--          UGU
      UACCCUGUCGAAC  CGC GUUCGACA  GUACCUGUCGAAC  \
      GUGGGACAGCUUG  GCG CAAGCUGU  UAUGGAUAGCUUG  G
-----          UUG C          CCC          UUG

```

OsmiR439f

```

ACC          UGA A          U--          UGU
      UACCCUGUCGAAC  CGC GUUCGACA  GUACCUGUCGAAC  \
      GUGGGACAGCUUG  GCG CAAGCUGU  UAUGGAUAGCUUG  G
---          UUG C          CCC          UUG

```

OsmiR439g

```

U          UGA A          U--          UGU
      CCUGUCGAAC  CGC GUUCGACA  GUACCUGUCGAAC  \
      GGACAGCUUG  GCG CAAGCUGU  UAUGGAUAGCUUG  G
-          UUG C          CCC          UUG

```

OsmiR439h

```

ACCACC          UCA A          U--          UGU
      UACCCUGUCGAAC  CGC GUUCGACA  GUACCUGUCGAAC  \
      GUGGGACAGCUUG  GCG CAAGCUGU  UAUGGAUAGCUUG  G
-----          UUG C          CCC          UUG

```

OsmiR439i

```

CACC--          UGA A          U--          UGU
      UACCCUGUCGAAC  CGC GUUCGACA  GUACCUGUCGAAC  \
      GUGGGACAGCUUG  GCG CAAGCUGU  UAUGGAUAGCUUG  G
CAAUUC          UUG C          CCC          UUG

```

OsmiR439j

```

      C          U U A C CA
ACAGGGUACC UAUCGAACAACUGCGGUUCGACA--GG ACC AUUGG CUG CA \
UGUCCCAUGG AUAGCUUGUUGGCGCCAAGCUGU CC UGG UGAUC GAC GU C
-          \ - U G A UU

```

OsmiR440 ($\Delta G = -62.4$)

```

      A UG      C G C      C      U      CAA -      CC
AUUGC U UUGGUG UGG CU GUCCUGAUC UAGGAGAC CUGAU GC UAGG U
UAGCGA A AGCCAC AUC GA CAGGGCUAGU GUCCUCUG GAUUA UG AUUC A
      G GU      A G A      A      U      --- U      AA

```


OsmiR396d ($\Delta G = -92.3$)

```
      GCG      C      CA      G      .-U      G UG
AAAGAUGU  GGCAUG UUUCCA  GGCUUUCUUGAACUGU AAC  CGUGGG G  U
UUUCUGCG  CCGUAC AAAGGU  CCGAAAGAACUUGGUA UUG  GUACUC U  A
      ---      C      AC      G  \  -      G GU
```

OsmiR396e

```
      G      C      CA      G      .-U      G UG
GCGG CAUG UUUCCA  GGCUUUCUUGAACUGU AAC  CGUGGG G  U
CGCC GUAC AAAGGU  CCGAAAGAACUUGGUA UUG  GUACUC U  A
      -      C      AC      G  \  -      G GU
```

OsmiR441a ($\Delta G = -67.4$)

```
AAAAUAUAUU      G      UU      UA      A      A      C      CUAGAU
      AAUUG UACU      CUC      UUU ACAAUGUAAGUCAUUUUAGCAUUUC CACAUU AUAUUGAU      U
      UUAUU AUGA      GAG      AAA UGUUACAUUCAGUAAAUCGUAAAG GUGUAA UAUAACUA      C
AAAAC-----      G      GG      GC      G      G      A      CCAUUA
```

OsmiR441b

```
UUAUCACCACUGACAU      CC      C      G      A      UU      C      UUA-      AUC
      CUUCCU      GUUUA AAUGUAAGUCAUUUA UA UU CAUAUU AUAUUGAUG      AUGA      U
      GAGGGA      CAAAGU UUACAUUCAGUAAGAU GU AA GUGUAA UAUAACUAC      UACU      A
AC-----      UA      A      G      A      GG      A      CAUG      CAG
```

OsmiR441c

```
      C      UU      GAU
UACUCCUCUGUUUCAUAAUGUAAGU AUUUUAGCAUUU CAUAUUUAUAUUGAUGGUA AUGAAUCUAGAU      A
AUGAAGGAGGCAAAGUGUUACAUUA UAAGAUCGUAAA GUGUAAAUAUAACUACCAUUAUCUAGAUCUGU      U
      A      GG      AUA
```

OsmiR442 ($\Delta G = -79.8$)

```
      A      A      AUAAU      AAA      U-      UA--      UGA      --      CUU      C      AUGAUCUGAC      GG      U
CACAUU UGGAGUAUU AAU      GA      AAUAAC      AAU      UACAGA      CGU      GUAAAUUGGAGACGAU      UUAAGCCUAAUUG UCC      AAUGU UGC A
GUGUGUA ACUUAUAA UUG      CU      UUAUUG      UUG      AUGUCU      GCA      CAUUGACGCUCUCUUA      AAUUCGAUUAU AGG      UUACA AUG C
      A      A      CAU---      GA--      UU      UUUA      UAG      GA      AAU      U      CAGUAUCGU      A-      A
```

OsmiR443 ($\Delta G = -74.4$)

```
      CAU--      CCC      AG----      A      A      --      A
CGUCC      AAAAACAAA      AAAACU      AUGUGAUUAUC CA UACAAUAAAUCUGGAUAGGAG      UCU U
GCAGG      UUUUUGUUU      UUUUGA      UACACUGUAUAG GU AUGUUUUUAGAUCUAUCUUC      AGA C
      UUUUU      AUA      CUUAUA      G      C      AU      C
```

OsmiR445a ($\Delta G = -141.0$)

```
      A      -      G      AAA      A      UC      AUGAAA      C      A      C      AU      A      .-AUGAAA      U
UCACAUCG AUGUUUG ACGCUAAUUUGGAGUAUUAAACAUU ACUAAU      AAAACU AUU      AUAA      GCUAUU UGCGAGACGAAUUUUUU AG CUAAUUA      CCAU AUU      AG U
AGUGUAGC UACAAAU UGUGAUUAAAUCUCAUAAUUUGUAU UGAUUA      UUUUGG UAA      UAUU      UGAUUA GCGCUCUGCUUAGAAAA UC GAUUAU      GGUA UAA      UC U
      C      A      G      AUA      A      GA      GGAACC      A      C      A      GU      C      \  -----      A
```

OsmiR445b

```
      G      A      -      GAC      AA-      A      UC      AUGAAA      C      C      A      C      AU      A      .-AUGAAA      U
GUCC UGUCACAUCG AUGUUUG ACGCUAAUUUGGAGUAUUAAACAUU UAAU      AAAACU AUU      AUAA      GCUAUU UGCGAGACGAA UUUUU AG CUAAUUA      CCAU AUU      AG U
CAGG ACAGUGUAGC UACAAAU UGUGAUUAAAUCUCAUAAUUUGUAU AUUA      UUUUGG UAA      UAUU      UGAUUA GCGCUCUGCUU GAAAA UC GAUUAU      GGUA UAA      UC U
      G      C      A      GUA      AUA      A      GA      GGAACC      A      C      A      GU      C      \  -----      A
```


OsmiR445c

CC A - C G AAA A UC AUGAAA CU AU A .-AUGAAA U
GUGUCACAU CG AUGUUUG ACGCUAAUUUGGAGUAUUAAA AUA ACUAAU AAAACU AUU AUAA GCUAAU GCGAGACGAAUUUUUUAAGCCUAAUUA CCAU AUU AG U
UACAGUGUAGC UACAAAU UGUGAUUAAAUCUCAUAAUUU UAU UGAUUA UUUUGG UAA UAUU UGAUUA UGCUCUGCUUAGAAAAUUCGGAUUAU GGUA UAA UC U
A- C A A G AUA A GA GGAACC AG GU C \ ----- A

OsmiR445d

 -- AA A AUG AAAA AUG CCCC-- C CCCU GA C UG UCUA A C GAUGCU
UGG CCA UUUAGUCCUGUCACAU C GAUGUU ACACUAAUU GUUUUUAAACAUAGACUA ACAAA AUU CAUAA G UUAUUUCG GAGA AA UUGAGUCUAAUUAAUC AUGAUUAGC UAUU \
ACC GGUU AAUUCAGGGAUAGUGUAG CUACAA UGUFAUUA UUAUAAUUUGUAUUGAU UGUUU UAA GUUUU C GAUUAAAC CUCU UU AAUUCGGAUUAUUG UAUUAAUUG GUACA A
UA G- C AUA AUC- GAA UGAUUUAU C UACU UC A GU UAAA G U AAUCAC

OsmiR445e

 G A - G AAA A UC AUGAAA C A C AU A .-AUGAAA U
GUCC UGUCACAU CG AUGUUUG ACGCUAAUUUGGAGUAUUAAACAU ACUAAU AAAACU AUU AUAA GCUAAU UGCGAGACGAAUUUUUU AG CUAAUUA CCAU AUU AG U
CAGG ACAGUGUAGC UACAAAU UGUGAUUAAAUCUCAUAAUUUGUAU UGAUUA UUUUGG UAA UAUU UGAUUA GCGCUCUGCUUAGAAAA UC GAUUAU GGUA UAA UC U
 G C A G AUA A GA GGAACC A C A GU C \ ----- A

OsmiR445f

 - AAAAC U G A - G AAA A UC AUGAAA C A C AU A .-AUGAAA U
AGAU GGGGCU UUUU AGUCC UGUCACAU CG AUGUUUG ACGCUAAUUUGGAGUAUUAAACAU ACUAAU AAAACU AUU AUAA GCUAAU UGCGGACGAAUUUUUU AG CUAAUUA CCAU AUU AG U
UUU CCCUGA AAAA UCAGG ACAGUGUAGC UACAAU UGUGAUUAAAUCUCAUAAUUUGUAU UGAUUA UUUUGG UAA UAUU UGAUUA GCGCUCUGCUUAGAAAA UC GAUUAU GGUA UAA UC U
 U AU--- U G C A G AUA A GA GGAACC A C A GU C \ ----- A

OsmiR445g

 U G A - G AAA UA UC AUGAAA C A CC AU A .-
AUGAAA U
UUU AGUCC UGUCACAU CG AUGUUUG ACGCUAAUUUGGAGUAUUAAACAU ACUAAU AAAAC AUU AUAA GCUAAU UGCGAGACGAAUUUUUU AG UAAUUA CCAU AUU
AG U
AAA UCAGG ACAGUGUAGC UACAAAU UGUGAUUAAAUCUUAUAAUUUGUAU UGAUUA UUUUG UAA UAUU UGAUUA GCGUUCUGCUUAGAAAA UC AUUAU GGUA UAA
UC U
 U G C A G AUA UA GA GGAACC A C AA GU C \ -----
- A

OsmiR445h

 U G A - G AAA A UC AUGAAA C A C AU A .-AUGAAA U
UUUU AGUCC UGUCACAU CG AUGUUUG ACACUAAUUUGGAGUAUUAAACAU ACUAAU AAAACU AUU AUAA GCUAAU UGCGAGACGAAUUUUUU AG CUAAUUA CCAU AUU AG U
AAAA UCAGG ACAGUGUAGC UACAAAU UGUGAUUAAAUCUCAUAAUUUGUAU UGAUUA UUUUGG UAA UAUU UGAUUA GCGCUCUGCUUAGAAAA UC GAUUAU GGUA UAA UC U
 U G C A G AUA A GA GGAACC A C A GU C \ ----- A

OsmiR445i

 UUU G A - G AAA A UC AUGAAA C .-AAGCC AU
ACUU AGUCC UGUCACAU CG AUGUUUG ACGCUAAUUUAGAGUAUUAAAUUA ACUAAU AAAACU AUU AUAA GCUAAU UGUGAGAUGAAUUUUU UAAUUA \
UGAA UCAGG ACAGUGUAGC UACAAAU UGUGAUUAAAUUUCAUAAUUUGUAU UGAUUA UUUUGG UAA UAUU UGAUUA GCGCUCUGCUUAGAAA AUUAU C
 UU- G C A G AUA A GA GGAACC A \ ----- AC

OsmiR446 (ΔG= -91.4)

AAA----- C A AA C U GAU
 UAUGUACUUCUUUGUUUCACAGUGUAAGU AUU UA AUUCCCCA AUUUUAUUGAUGUUAAUGAAUCUA AUA A
 AUAUAUGAGGGAGGCCAAAGUGUUACAUUCA UAA AU UAAAGGGU UAAGUAUAACUACAAUUAUUUAGAU UGU U
CAAGUUCAAG A G GG A C AUA

Figure 11

Predicted fold-back structures of the miR390 from *Arabidopsis* and *Populus*.

Arabidopsis thaliana

aagctcaggaggatagcgcc

```

      AGAUUGAGU   G   AU   AA   G   .-U C
            AGUA AGAA AGCUAU AGCUCAGGA GGAUAGCGCCA GG U
            UCGU UCUU UCGAUA UUGAGUCCU CCUAUCGCGGU CC C
      ----- -   CU   CC   A   \ - A

```

aagctcaggaggatagcgcc

```

      CAAAAAACAAA- --   AU U A   G   ----- UC
            GUAG AGAAGA C GU AAGCUCAGGA GGAUAGCGCCA UGAUGA A
            CAUC UCUUCU G UA UUUGAGUCCU CCUAUCGCGGU AUUGCU C
      CCGGAAAAAGUAA AU   CG U C   A   UUUUUUUCU UA

```

aaactcaggatggatagcgcc

```

      --   UUUU--- UU   UA   CCAA A   U   AAAAAUA A G
      CGGCCU   CA GUAG AGAAGAG UGAA CUCAGGA GGAUAGCGCCA GAU AC A
      GUUGGG   GU CAUC UCUUCUU AUUU GAGUCCU CCUAUCGCGGU CUA UG A
      GU   UUUUUUU UU   --   AGAC C   C   ACUA--- G U

```

Populus trichocarpa

aagctcaggaggatagcgcc

```

      UGAUAUGUGCAUA   U   UA   G   AUGA   A -   GUU A
      UU
            AGUA GGGAGGAUCUGU AGCUCAGGA GGAUAGCGCC GCUGAUGAU AG UUGAU UG UGGG
      \
            UCGU UCCUCUUGGAUA UUGAGUCCU UCUAUCGCGG UGAUUACUG UC AACUA AC ACUC
      A
      UUA----- -   UC   A   ----- A U AU- A
      UA

```

aagctcaggaggatagcgcc

```

      CUUGAUUAUAGA   U   U U   G   - A-- GAU-- UG- CU
            AGUA GGAAGAAUC GU AAGCUCAGGA GGAUAGCGCC CU AG AACCA GG C
            UCGU UCUUCUUGG CA UUUGAGUCCU CCUAUCGCGG GA UC UUGGU UU U
      CA----- -   U U   A   U CUA AGUUU UUA UU

```

aagctcaggaggatagcgcc

```

      G--   UA   G   GAGC   AAAGUCUA--- U   UU
            GGAGAAUCUGU AGCUCAGGA GGAUAGCGCCAU AUGAC UGUU GAG \
            CCUCUUGGGUA UUGAGUCCU UCUAUCGCGGUG UACUG ACAA CUC A
      AUU   UC   A   AC-- ACCUAACUAAA - UA

```


aagctcaggagggatagcgcc

CU AUAGAAA U UCU U G - A AAU-----
AGUA GGAAGAA GU AAGCUCAGGA GGAUAGCGCC CU AGGAU CAUGGG
\
UCGU UCUUCUU UA UUUGAGUCCU CCUAUCGCGG GA UCUUA GUAUUU
C ACCA--- - CGU C A U C GUUUUUUGGU
UU

aaactcaggatggatagcgcc

AA-- CCA A A U A UA CAAA AAUAAAA
GAAGAA GU AA CUCAGGA GGAUAGCGCC CUGA GU AACCA \
CUUCUU CA UU GAGUCCU CCUAUCGCGG GAUU UA UUGGU A
AUAC AGA A C C - CC ---- ACCCGAG .

aaactcaggatggatagcgcc

GGAGCU GCAA GCAAUG A U A --- AAAACCACAU
GGUA GAAGAA AA CUCAGGA GGAUAGCGCC CUGAG AAUCA A
UCAU CUUCUU UU GAGUCCU CCUAUCGCGG GAUUC UUAGU A
U----- AC-- AGACAA C C - CUA ACCCGAGAAA

Figure 12

Predicted fold-back structures of miR437 from maize, sugarcane and *Sorghum*.

Maize

U A U U U U A G C U U G C C U A A U U A G A C U U C U C U A C U U G A A U G U U U A U A A A A A U G C A U C U A A A U A U U A A A U A G A U A A C A U A A A \ G U A A A A U U G A A C G G G U U A G U U U G A A G A G A U G A A C U U A C A A A U A U U U U U A U G U A G A U U U A U A G U U U A U U U G U U G U A U U U A A A G U A A A C U - U C A C A A A U C U A C A A A C A A A U C A A A G U A C C A U C C G U

Sugarcane

U C - - C G U A A C U A A A C U U C C C C A U G C A C C A A A - G . - U A U C A A A C U A G U U C C A U U A A A U A C U C U C U C U G U U C A A U A U G U U U A C U A G C U U U G U C A A G U C A A C U C U C U A A C U U U G A U C A A G U U U U A G A A A A A A U C U A A A U U C \ A U G A G G G G C A A G U U A U A C A A G U G A U C G A A A C A G U U C A G U U U G A G A G A U U G A A A C U A G U U C A A A A U C U U U U U U A G A U U U A A G C G C A U G A A A A - - - - - N A A A U A U A A A U A G G \ - - - - - U A

Sorghum

U U C C A A A A U G U G G U U G U U U U G U U U U G U C A U A A G U A A A U U U C U C A A C U U U G A C C A A G U U U A G A A A A U A A C A A C A U G G C A A A U C A U G A A A G G A G G U U U U A C A C A G C A A A A C A A A A U A G U A U U C A U U U G A A G A G U U G A A A U U G G U U C A A A U C U U U U A U U G U U G U A U C G U U U G G U G C U U U A U U C - G - A C U A A G A \ - - - - - A

Figure 13

Predicted fold-back structures of the putative miRNA precursors from rice (see Table 5).

P18-F1 ($\Delta G = -63.3$)

UGUUAACGGCAC	U	A-	U	A	UGG----	U-	UA
	GGUUU	CUUGG	UCUCUC	CUCCCCU	GAGGCU	UCUCA	AGGU UGAUG \
	CCGAA	GAACC	AGAGAG	GAGGGGGCU	UCCGA	AGAGU	UCCG GUUAC C
UGACA-----	C	GA	-	A	CAAAGAA	UU	CA

P20-A11 ($\Delta G = -114.1$)

A A C A C A C UG UG GGU
 AG UUU CU CGGUCCAACAAAAAGUA CUCGAGGUA CG UACCUC CGGUAC AAAU UUUCCGAUCGUUGGAUCUAGC GGUG \
 UC AAA GA GCCAGGUUGUUUUUCAU GAGCUCCAUC GC AUGGAG GCCAUG UUUU AAAGGCUAGCAACCUAGAUUG CUAC U
 A A A A C C U GU -- GGG

P83-E9 ($\Delta G = -201.3$)

[illegible]

P92-A7 ($\Delta G = -74.8$)

A G - CC U UCC ACU-- .-CAUAC AUAUA A UAU G
 GAUGCUCUUC UCCUAAAAUAUAGCA UUUUAG AC UGACA AGUU GAUAU ACUUUGAU AAUAUACAAU UAU UA AUAUA U
 CUAUGAAGGAGG AGGGUUUUUAUUAUUUGU AAAAUU UG ACUGU UGAG UUGUA UGAAACUG UUAUAUGUUG AUA AU UGUAU A
 C A G UC - U-- AAAAU \ ----- AAAA- A UU- A

P101-H12 ($\Delta G = -160.0$)

CA	AAUA	C	CU
UGUGGCUCGCAUGCUGUA	ACUAACGGCAUAAUAGAUCAUUGAUGA	GACGUAUAUCGGUGUUCGCUAUAUAUACUAUCUACUGGUAAGUAUAU	\
ACACCGAGCGUACGAAU	UGAUUGCCGUAUUAUUUAGUGAACUACU	CUGCGUAUAAGCCACAAGUGAUUAUAUGAUAGAUGACCAUUCAUUA	U
	-----	A	AU

A.

