





(86) Date de dépôt PCT/PCT Filing Date: 2000/05/01
(87) Date publication PCT/PCT Publication Date: 2000/11/16
(45) Date de délivrance/Issue Date: 2007/11/13
(85) Entrée phase nationale/National Entry: 2001/11/09
(86) N° demande PCT/PCT Application No.: US 2000/011631
(87) N° publication PCT/PCT Publication No.: 2000/067570
(30) Priorité/Priority: 1999/05/10 (US09/307,907)

(51) Cl.Int./Int.Cl. *A01N 49/00* (2006.01),
A01N 25/18 (2006.01), *A01N 31/02* (2006.01),
A01P 17/00 (2006.01)
(72) Inventeurs/Inventors:
NOLEN, JAMES A., US;
BEDOUKIAN, ROBERT H., US;
MALONEY, ROBERT E., US;
KLINE, DANIEL L., US
(73) Propriétaires/Owners:
BIOSENSORY, INC., US;
BEDOUKIAN RESEARCH, INC., US; ...

(54) Titre : PROCEDE, DISPOSITIF ET COMPOSITIONS DESTINES A INHIBER LA CAPACITE DES MOUSTIQUES DE
DETECTER L'ODEUR HUMAINE, DANS DES ESPACES TRIDIMENSIONNELS DEFINIS DE MANIERE
ENVIRONNEMENTALE

(54) Title: METHOD, APPARATUS AND COMPOSITIONS FOR INHIBITING THE HUMAN SCENT TRACKING ABILITY
OF MOSQUITOES IN ENVIRONMENTALLY DEFINED THREE DIMENSIONAL SPACES



(57) Abrégé/Abstract:

The ability of mosquitoes to locate a target by olfactory emissions of the target is inhibited by dispensing into a spatial area an inhibiting effective amount of at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of formula (I) and 3-methyl-1-alkyn-3-ols of formula (II): wherein R¹ and R² are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms.



(73) **Propriétaires(suite)/Owners(continued):**

UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY OF AGRICULTURE, US

(74) **Agent:** OSLER, HOSKIN & HARCOURT LLP

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 November 2000 (16.11.2000)

PCT

(10) International Publication Number
WO 00/67570 A3

(51) International Patent Classification⁷: A01N 27/00

(21) International Application Number: PCT/US00/11631

(22) International Filing Date: 1 May 2000 (01.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/307,907 10 May 1999 (10.05.1999) US

(71) Applicants: **BIOSENSORY, INC.** [US/US]; Windham Mills Technology Center, 322 Main Street, Building 1, Second Floor, Willimantic, CT 06226-3149 (US). **BEDOUKIAN RESEARCH, INC.** [US/US]; 21 Finance Drive, Danbury, CT 06810-4192 (US). **UNITED STATES OF AMERICA**, as represented by **THE SECRETARY OF AGRICULTURE** [US/US]; Washington, DC 20250 (US).

(72) Inventors: **NOLEN, James, A.**; 60 Sharp Street, West Greenwich, RI 02817-2100 (US). **BEDOUKIAN, Robert, H.**; 72 Limekiln Road, West Redding, CT 06896 (US). **MALONEY, Robert, E.**; 42 Shelley Road, Bethel, CT 06801 (US). **KLINE, Daniel, L.**; 5027 NW 75th Lane, Gainesville, FL 32653 (US).

(74) Agent: **RAUCHFUSS, George, W., Jr.**; Ohlandt, Greeley, Ruggiero & Perle, L.L.P., 9th Floor, One Landmark Square, Stamford, CT 06901-2682 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.


(88) Date of publication of the international search report:
10 May 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD, APPARATUS AND COMPOSITIONS FOR INHIBITING THE HUMAN SCENT TRACKING ABILITY OF MOSQUITOES IN ENVIRONMENTALLY DEFINED THREE DIMENSIONAL SPACES



(I)



(II)

(57) Abstract: The ability of mosquitoes to locate a target by olfactory emissions of the target is inhibited by dispensing into a spatial area an inhibiting effective amount of at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of formula (I) and 3-methyl-1-alkyn-3-ols of formula (II): wherein R¹ and R² are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms.

**METHOD, APPARATUS AND COMPOSITIONS FOR
INHIBITING THE HUMAN SCENT TRACKING
ABILITY OF MOSQUITOES IN ENVIRONMENTALLY
DEFINED THREE DIMENSIONAL SPACES**

5

Field of the Invention

10 This invention relates to a method, apparatus and compositions for
inhibiting the ability of mosquitoes to locate or track a human body by scent detection.
More particularly, the invention relates to the use of certain compounds in compositions
and apparatus to inhibit mosquitoes' ability to detect humans by scent detection.

15

Background

Compounds, compositions and formulations for protecting human beings
from being bitten by mosquitoes are known in the art. Generally, these compounds,
compositions and formulations are based on their ability to persist on the skin of the
20 person upon topical or surface application for a time sufficient to repel mosquitoes.
Numerous adjuvant materials have been added to mosquito repellants to increase the
persistence of the repellents to the skin of a person. However, despite the various
attempts to improve the repelling activity of the known mosquito repellents, these
attempts have generally not been successful, as almost anyone who has used such
25 mosquito repellents can attest.

Thus, the art has been searching for new and more effective repellents
against mosquitoes. However, the search for more effective mosquito repellents has
not generally been met with success since most mosquito repellents have been found
30 only to possess a limited degree of repellency and are generally not particularly
effective. There is, therefore, a need for more effective means to deter mosquitoes from
locating and biting humans and other targets such as livestock. Moreover, this need

has recently become more acute and urgent because mosquitoes have been discovered to be carriers of significant diseases that can be passed on to a target by the mosquitoes biting the target. A further need is to be able to reduce the use of environmentally unfriendly pesticides.


5

Summary of the Invention

The inventors have discovered that compounds, compositions and formulations heretofore proposed as repellents for mosquitoes have lacked the necessary efficacy due to the ability of mosquitoes to locate and be drawn to the targets by olfactory emissions of the target. Thus, if a mosquito enters a zone or space where a potential target is located, the mosquito can be attracted to the target by olfactory emissions of the target and, this olfactory attraction is sufficient to overcome any repellency activity of the repellent compound, composition or formulation applied on the target. Therefore, the present invention provides compositions and formulations containing compounds usable in methods and apparatus for inhibiting the olfactory target tracking abilities of mosquitoes when an effective amount of the inhibiting compound(s) is/are dispersed in a three dimensional atmospheric space.


20

According to this invention, the ability of mosquitoes to locate a target is inhibited by dispensing into a spatial area an inhibiting effective amount of at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of the formula:



25

and 3-methyl-1-alkyn-3-ols of the formula:




wherein R^1 and R^2 are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms.


5 The inhibiting compound can be dispensed into the three dimensional atmospheric space by any suitable means sufficient to provide an inhibiting effective amount of the inhibiting compound(s). Such dispensing means includes, for example, evaporation, atomization and ionic dispersion of the inhibiting compound from any suitable composition or formulation. Such composition or formulation will generally
10 comprise a base vehicle containing at least one of the inhibiting compounds.

Detailed Summary of the Invention

15 The inventors have discovered that if an effective amount of at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of the formula:



and 3-methyl-1-alkyn-3-ols of the formula:



wherein R^1 and R^2 are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms is dispensed into the atmosphere of a three dimensional environmental space, the ability of mosquitoes to locate and track a target, such as humans or livestock, by the target's olfactory emissions is inhibited.

Any suitable 3-methyl-1-alkene-3-ols or 3-methyl-1-alkyn-3-ol of the formulas may be employed in the method, compositions and apparatus of this invention. Especially suitable inhibiting compounds are nerolidol, 3-methyl-1-octen-3-ol, linalool and dehydrolinalool. Depending on the particular mosquito species, either the 3-methyl-1-alkyn-3-ols or the 3-methyl-1-alkene-3-ols are better inhibitors than the other class of components and will be preferred for that species of mosquito. The inhibiting compounds may be utilized singly or as mixtures of two or more of such compounds.

Any suitable inhibiting effective amount of the inhibiting compound(s) may be employed. Such inhibiting effective amounts can include amounts, based on the square footage of land or base surface area of the environmental area to be treated, within the range of from about 0.000005 g/hr/ft² to about 0.004 g/hr/ft², preferably amounts within the range of from about 0.00015 g/hr/ft² to about 0.0002 g/hr/ft², and especially an amount of about 0.00016 g/hr/ft².

The inhibiting compounds for use in this invention may be provided in an essentially pure form of the inhibiting compounds or as a component of a natural essential oil having a concentration of an inhibiting compound sufficient to make it practical and feasible to dispense an inhibiting effective amount of inhibiting compound. Generally, the essential oil will contain a concentration of the inhibiting compound of at least about 2%, preferably at least about 5%, and especially at least about 50% by weight. For example, the inhibiting compound can be provided as a synthetically produced, essentially pure compound or as a component of an essential oil such as basil oil, ho wood oil and the like.

The inhibiting compounds of this invention, or essential oils containing such inhibiting compounds, may be employed in any formulation suitable for dispensing inhibiting effective amounts of the compounds. The compounds will generally be employed in formulations comprising a suitable vehicle containing the inhibiting compounds. For example, the inhibiting compound can be formulated in a specially formulated wax-like medium or vehicle engineered to release desired amounts of vaporous inhibiting compound at ambient temperatures, such as those mediums or vehicles available from Koster Keunen of Watertown, Connecticut. An example of such a wax-like medium available from Koster Keunen is known as Insect Repellent Wax Bar No. 9, which is a blend of waxes having the following general composition: fatty acids ranging in carbon chain length of from C₁₆ to C₂₂, fatty alcohols ranging in carbon chain length of from C₁₆ to C₂₂, paraffinic hydrocarbons ranging in carbon chain length of from C₁₉ to C₄₇, branched hydrocarbons ranging in carbon chain length of from C₂₃ to C₆₉, beeswax and other natural waxes such as candelilla and carnauba. The wax mixture will generally be formulated with concentrations of the inhibiting compounds of this invention ranging from about 20% to 60% and the formulation has a congealing point which may vary from about 75°C to about 45°C. Alternatively, the inhibiting compound can be formulated in a porous medium or vehicle suitable for releasing effective amounts of the inhibiting compound. As an example of such porous medium or vehicle is a polyester membrane material having micropores encasing a block of inhibiting compound saturated fibers that gradually releases the inhibiting compound so that it permeates the microporous membrane and is released to the environment. Such porous membrane known as World of Fragrance™ cups is available from Waterbury Companies, Inc. of Waterbury, Connecticut.

The formulations can be placed in any suitable container or device for dispensing the inhibiting compound. For example, the formulations can be placed in a suitable fan-equipped device so that one can obtain, for example, fan-driven evaporation of the inhibiting compound from a porous medium or wax-like medium containing the inhibiting compound. As examples of such fan-equipped devices, there can be mentioned the devices disclosed in US Patent 5,370,829 of Waterbury

Companies, Inc. and the apparatus disclosed in US Patent 5,799,436 of Biosensory Insect Control Corporation.

5 Another suitable means of dispensing the inhibiting compound is by atomization and/or ionic dispersion of the compound as suitable-sized, positively-charged droplets from a suitable atomization or ionic dispersing apparatus, such as the Ionic Wind™ device, available from Brandenburg, Ltd. of Brierley Hill, United Kingdom.

10 The inhibiting compounds of this invention are effective against mosquitoes, such as for example, *Aedes taeniorhyncus* (Black Salt Marsh mosquito), *Culex nigripalpus*, *Aedes aegypti*, *Aedes albopictus* (Asian Tiger mosquito), *Culex pipiens* (common house mosquito) and the like.

15 The use of this invention is illustrated by the following non-limited examples.

Example 1

20 A triple cage, dual-port olfactometer, illustrated and described in detail by Posey et al. J. Med. Entomol. 35(3), 330-334 (1998), was used to study the responses of 6 to 8 day old, laboratory-reared adult female *Aedes aegypti* mosquitoes. This system allows mosquitoes to choose between two different stimuli. The olfactometer is constructed of clear acrylic, comprises three test chambers in a tiered configuration, has
25 paired removable sleeves, and mosquito traps on each chamber, and is equipped with a filtered external air supply system that allows precise temperature (+/- 0.5°C) and relative humidity (+/- 2%) control. Only one chamber at a time was used for the tests. Outside air was conditioned prior to entry through the choice ports, the mosquito trap, and the olfactometer by passing through a series of charcoal filters and then heated and
30 humidified, if necessary.

One hour before initiation of tests, about 75 female *Ae. aegypti* mosquitoes were aspirated into the olfactometer chamber and allowed to acclimatize for that one hour before testing. Test compounds were placed into test ports upwind of the traps and olfactometer chamber. A completely randomized design was used for cage position.

Competitive tests were conducted. In these competitive tests, the inhibiting compound was placed into one of the ports (treatment port); the second port (check port) had the same apparatus but no chemical inhibiting compound. In these tests, inhibiting compounds were presented in glass Pyrex petri dishes (60 x 15 mm) that had been cleaned and sterilized in a vacuum oven and then handled only with gloves to minimize any chance of contamination and placed into one port (Treatment Port) along with another petri dish containing 500 μ L of Clara Sludge (CS), a synthetic human attractant. The other port (Check Port) contained an untreated petri with no inhibiting compound and a second petri dish containing 500 μ L CS. Various concentrations (25, 100, 250, 500 and 1000 μ L) of the compounds were used in these tests. Each test was conducted for a period of 3 minutes; the number of mosquitoes trapped in the baited ports, and those remaining in the cage, were counted. The data are then presented as a percentage of total mosquitoes in the cage that were attracted to each port and the percentage of mosquitoes remaining in the cage (i.e., not attracted to either port).

Results of the competitive tests are set forth in the following Table 1.

Table 1 Competitive Tests

Test Compound and Amount	% Mosquitoes Entering		% Mosquitoes Remaining in Cage
	Treatment Port (Inhibitor Port)	Control Port (Human Scent Port)	
Dehydrolinalool 25 micro liter	30.23	50.77	19
Dehydrolinalool 100 micro liter	31.13	53.87	15
Dehydrolinalool 250 micro liter	22.6	44.6	32.8
Dehydrolinalool 500 micro liter	24.8	56.47	24.18
Dehydrolinalool 1000 micro liter	26.11	42.76	27.12
Linalool 25 micro liter	21.98	56.89	21.14
Linalool 100 micro liter	31.3	45.61	23.08
Linalool 250 micro liter	16.57	47.3	36.13
Linalool 500 micro liter	16.13	44.56	39.31
Linalool 1000 micro liter	27.24	44.14	31.3

For comparison purposes, 500 μ L of the human scent (CS) was also placed alone in the Treatment Port, and with only an empty petri dish in the Check Port. The Treatment Port with the CS attracted 69 to 65% of the mosquitoes, with 30 to 35% of the mosquitoes remaining in the cage, and none being attracted to the Check Port with only the petri dish. At all tested concentration levels, linalool and dehydrolinalool are shown to greatly inhibit the ability of mosquitoes to locate the human scent (CS) as shown by the much lower percentage of mosquitoes captured in the Treatment Port compared to the Human Scent Port.

Example 2

Tests identical to that described in Example 1 were conducted with test compounds when employed together with the synthetic human attractant (CS) in a port. The test combinations were:

- (1) CS vs. CS (comparison)
- (2) dehydrolinalool + CS vs. CS
- (3) linalool + CS vs. CS

5

Also for comparison purposes, a known topical mosquito repellent, Deet, namely N,N-diethyl-3-methylbenzamide, was employed to compare to linalool and to dehydrolinalool. Each test compound was tested at 100, 250 and 500 μ L. The results are presented in Tables 2, 3 and 4. The results are again presented as percent mosquitoes attracted to Ports 1 or 2 or not attracted to either port.

10

Table 2 (100 micro liter treatments)

Test Compound and Amount	% Mosquitoes Entering		% Mosquitoes Remaining in Cage
	Port 1	Port 2	
1 Human scent vs. 2 human scent	50.5	44	5.5
1 Dehydrolinalool vs. 2 human scent	18.1	63	18.9
1 Linalool vs. 2 human scent	22.5	50.7	26.8
1 Deet vs. 2 dehydrolinalool	60.1	20.1	19.8
1 Deet vs. 2 linalool	56.3	17.5	26.2

15

Table 3 (250 micro liter treatments)

Test Compound and Amount	% Mosquitoes Entering		% Mosquitoes Remaining in Cage
	Port 1	Port 2	
1 Human scent vs. 2 human scent	49.9	39.4	8.7
1 Dehydrolinalool vs. 2 human scent	20.1	59.2	20.6
1 Linalool vs. 2 human scent	14.9	56.8	28.4
1 Deet vs. 2 dehydrolinalool	48.8	22.5	29.2
1 Deet vs. 2 linalool	48.1	25.7	26.2

5

Table 4 (500 micro liter treatments)

Test Compound and Amount	% Mosquitoes Entering		% Mosquitoes Remaining in Cage
	Port 1	Port 2	
1 Human scent vs. 2 human scent	47.3	46.8	5.9
1 Dehydrolinalool vs. 2 human scent	20.6	55.5	23.9
1 Linalool vs. 2 human scent	25.7	55.6	18.7
1 Deet vs. 2 dehydrolinalool	42.1	30.3	27.6
1 Deet vs. 2 linalool	49.5	23.1	27.3

Example 3

10

In a further test of the inhibiting ability of the compounds of this invention, the following field test was conducted using linalool as the inhibiting compound. The study was conducted on a site adjacent to a wooded wetland breeding area in Sarasota, Florida. A trap baited with 200 ml/min CO₂, equivalent to the respiration of a 91 kg (200 lb) man, ran continuously each night as an experimental control. The trap collections

15

were first used to count the mosquito population and were then examine to identify the species present. The mosquito population was about 91% *Culex nigripalpus*, the St. Louis encephalitis vector in Florida.

5 The experimental design was a 2 x 2 Latin Square. Tests were conducted in two open areas 24 meters (80 feet) apart labeled Locations 1 and 2. Each area was 2.4 meters x 2.4 meters (8 ft x 8 ft), i.e., 5.76 m² (64 ft²). Wooden poles 1.38 meters (4.5 feet) high and having inhibiting compound dispensers thereon were driven into the ground at the corners of the square area. On alternate nights, one area was
10 unprotected by inhibiting compound while the other area had inhibiting compound dispensed from the dispenser. In operation, the dispenser emitted linalool inhibitor having 95% active ingredient at a rate of 20 mg/hr to 40 mg/hr.

 Landing counts were taken from the arms, legs and torso of a test subject
15 known to be moderately attractive to mosquitoes who was seated in the center of the square area. At 15-minute intervals over a period of 2 to 3 hours during peak mosquito activity, the subject changed his position, alternating between Locations 1 and 2. If there was no activity for five minutes, the test subject walked around the inside perimeter of the square area in an attempt to draw attention to his presence.
20 Mosquitoes were killed to prevent double counting.

 The mosquitoes were observed to exhibit a behavior similar to swarming each evening around 6:45 to 7:00 PM. On some cue, they emerge *en masse* from the wooded wetland and fan out in a radial direction foraging for blood hosts. The period of
25 peak activity is evidenced by a sharp increase in landing counts followed by a sharp decrease as the wave of outbound mosquitoes passes the test location. This behavior introduces two variables that must be taken into account. The first affects the experimental design and the second affects the interpretation of the results.

30 To be most accurate, an experimental trial must span a period before and after peak activity. Ending a trial at or near the time of peak activity, for example, will

result in a larger landing count for the treatment used last, skewing the result. The starting time and duration of trials were adjusted to avoid distorting the landing counts.

It is also recognized that the landing counts represent both:

5

1. mosquitoes drawn to the human subject by his scent, and
2. mosquitoes that encounter the human subject because he is in their line of flight as they leave the water.

10 If the inhibitor impairs the mosquitoes' scent-tracking ability, it will affect the former but not the latter. This fact must be taken into account when interpreting the results.

15 The sites selected represent a worse case condition that is experienced only by those located adjacent to a breeding site and in the line of flight as the mosquitoes come off the water. For a measure of performance more typical of the average homeowner, a test was conducted from 7:00 AM to 9:00 AM in the morning when activity is not at its peak.

20 Linalool reduced the mosquito landing counts by an average of 53% over the three trials. The unprotected human subject experienced 286 landings compared to 135 landings when surrounded by linalool dispensed from the dispensers.

25 The range for individual trials is a reduction of landings by 36% to 68% during periods of peak activity and by 73% in the morning trial when mosquito activity was light.


These results indicate linalool's ability to reduce the landing counts of *Culex nigripalpus* on humans in open areas by inhibiting the mosquitoes' ability to track the human's olfactory emissions.

30


With the foregoing description of the invention, those skilled in the art will appreciate that modifications may be made to the invention without departing from the spirit thereof. Therefore, it is not intended that the scope of the invention be limited to the specific embodiments illustrated and described.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method of inhibiting the ability of mosquitoes to sense a target by olfactory sensing of the target within a three dimensional environmental space having a land or base surface area, the method comprising dispensing into the atmosphere of the three dimensional environmental space a non-lethal, inhibiting effective amount of at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of the formula:



and 3-methyl-1-alkyn-3-ols of the formula:



wherein R^1 and R^2 are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms and wherein the inhibiting effective amount ranges from about 0.000005 g/hr/ft² to about 0.0004 g/hr/ft² based on the square footage of the land or base surface area of the environmental space.

2. The method according to Claim 1, wherein the inhibiting effective amount ranges from about 0.00015 g/hr/ft² to about 0.0002 g/hr/ft² based on the square footage of the land or base surface area of the environmental space.

3. The method according to Claim 1, wherein the at least one inhibiting compound is selected from nerolidol, 3-methyl-1-octen-3-ol, linalool and dehydrolinalool.

4. The method according to Claim 1, wherein the at least one inhibiting compound comprises linalool.

5. The method according to Claim 1, wherein the at least one inhibiting compound comprises dehydrolinalool.

6. The method according to Claim 1, wherein the dispensing of the at least one inhibiting compound comprises dispensing by a method selected from volatilization, evaporation, atomization and ionic dispersion of the at least one inhibiting compound from a formulation comprising a vehicle containing the at least one inhibiting compound.

7. The method according to Claim 6, wherein the dispensing comprises fan-driven evaporation of the at least one inhibiting compound from a formulation in which the vehicle is a porous medium.

8. The method according to Claim 6, wherein the dispensing comprises fan-driven evaporation of the at least one inhibiting compound from a formulation in which the vehicle is a wax-like solution.

9. The method according to Claim 6, wherein the dispensing comprises atomization of the at least one inhibiting compound from the formulation.

10. The method according to Claim 6, wherein the dispensing comprises ionic dispersion of the at least one inhibiting compound from the formulation.

11. The method according to Claim 4, wherein the linalool is dispensed by fan-driven evaporation of linalool from a formulation of a porous medium containing linalool.

12. The method according to Claim 4, wherein the linalool is dispensed by fan-driven evaporation of linalool from a formulation of a wax-like solution containing linalool.

13. The method according to Claim 4, wherein the linalool is dispensed by atomization of linalool from a formulation of a vehicle and linalool.

14. The method according to Claim 4, wherein the linalool is dispersed by ionic dispersion of linalool from a formulation of a vehicle and linalool.


15. The method according to Claim 5, wherein the dehydrolinalool is dispensed by fan-driven evaporation of dehydrolinalool from a formulation of a porous medium containing dehydrolinalool.

16. The method according to Claim 5, wherein the dehydrolinalool is dispensed by fan-driven evaporation of dehydrolinalool from a formulation of a wax-like solution containing dehydrolinalool.


17. The method according to Claim 5, wherein the dehydrolinalool is dispensed by atomization of dehydrolinalool from a base composition.

18. The method according to Claim 5, wherein the dehydrolinalool is dispersed by ionic dispersion of dehydrolinalool from a formulation of a vehicle and dehydrolinalool.

19. A composition for inhibiting the ability of mosquitoes to sense a target by olfactory sensing of the target within a three dimensional environmental space having a land or base surface area, the composition comprising at least one inhibiting compound selected from the group consisting of 3-methyl-1-alkene-3-ols of the formula:



and 3-methyl-1-alkyn-3-ols of the formula:



wherein R^1 and R^2 are each independently a saturated or unsaturated aliphatic hydrocarbon group containing from 1 to about 12 carbon atoms in a base formulation of a vehicle and the at least one inhibiting compound capable of permitting sufficient dispersion from the formulation of the at least one inhibiting compound by at least one of evaporation, volatilization, atomization or ionic dispersion sufficient to provide a non-lethal, inhibiting effective amount of said at least one inhibiting compound in an atmosphere of the three dimensional environmental space wherein the composition is sufficient to provide an inhibiting effective amount of the at least one inhibiting compound ranging from about 0.000005 g/hr/ft² to about 0.0004 g/hr/ft² per square footage of the land or base surface area of the three dimensional environmental space.

20. The composition according to Claim 19, wherein the composition is sufficient to provide an inhibiting effective amount of the at least one inhibiting compound ranging from about 0.00015 g/hr/ft² to about 0.0002 g/hr/ft² per square footage of the land or base surface area of the three dimensional environmental space.

21. The composition according to Claim 19, wherein the at least one inhibiting compound is selected from nerolidol, 3-methyl-1-octen-3-ol, linalool and dehydrolinalool.

22. The composition according to Claim 19, wherein the at least one inhibiting compound comprises linalool.

23. The composition according to Claim 19, wherein the at least one inhibiting compound comprises dehydrolinalool.

24. The composition according to Claim 19, wherein the composition is able to dispense the at least one inhibiting compound by a method selected from volatilization, evaporation, atomization and ionic dispersion of the at least one inhibiting compound.

25. The composition according to Claim 24, wherein the composition is dispensable by fan-driven evaporation of the at least one inhibiting compound and the vehicle is a porous medium.

26. The composition according to Claim 24, wherein the composition is dispensable by fan-driven evaporation of the at least one inhibiting compound and the vehicle is a wax-like solution.

27. The composition according to Claim 24, wherein the composition is dispensable by atomization of the at least one inhibiting compound.

28. The composition according to Claim 24, wherein the composition is dispensable by ionic dispersion of the at least one inhibiting compound.

29. The composition according to Claim 22, wherein the linalool is dispensable by fan-driven evaporation of linalool and the vehicle is of a porous medium containing linalool.

30. The composition according to Claim 22, wherein the linalool is dispensable by fan-driven evaporation of linalool and the vehicle is of a wax-like solution containing linalool.

31. The composition according to Claim 22, wherein the linalool is dispensable by atomization of linalool.

32. The composition according to Claim 22, wherein the linalool is dispersible by ionic dispersion of linalool.

33. The composition according to Claim 23, wherein the dehydrolinalool is dispensable by fan-driven evaporation of dehydrolinalool and the vehicle is of a porous medium containing dehydrolinalool.

34. The composition according to Claim 23, wherein the dehydrolinalool is dispensable by fan-driven evaporation of dehydrolinalool and the vehicle is of a wax-like solution containing dehydrolinalool.

35. The composition according to Claim 23, wherein the dehydrolinalool is dispensable by atomization of dehydrolinalool.

36. The composition according to Claim 23, wherein the dehydrolinalool is dispersible by ionic dispersion of dehydrolinalool.

