

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2002216763 B2

(54) Title
Alteration of cell membrane

(51) International Patent Classification(s)
C12N 15/09 (2006.01) **A61P 37/06** (2006.01)
A61K 35/28 (2006.01) **C07K 14/47** (2006.01)
A61K 38/00 (2006.01) **C07K 14/705** (2006.01)
A61K 39/00 (2006.01) **C07K 19/00** (2006.01)
A61K 45/00 (2006.01) **C12N 5/06** (2006.01)
A61K 47/48 (2006.01) **C12N 5/10** (2006.01)
A61P 7/00 (2006.01) **C12N 15/12** (2006.01)
A61P 35/00 (2006.01) **C12P 21/02** (2006.01)
A61P 37/00 (2006.01) **A61K 35/12** (2006.01)
A61P 37/04 (2006.01)

(21) Application No: **2002216763** (22) Date of Filing: **2001.07.02**

(87) WIPO No: **WO02/02751**

(30) Priority Data

(31) Number (32) Date (33) Country
60/215,580 **2000.06.30** **US**

(43) Publication Date: **2002.01.14**
(43) Publication Journal Date: **2002.04.11**
(44) Accepted Journal Date: **2007.03.01**

(71) Applicant(s)
University of Louisville Research Foundation, Inc.

(72) Inventor(s)
Shirwan, Haval

(74) Agent / Attorney
Phillips Ormonde & Fitzpatrick, Level 22 367 Collins Street, Melbourne, VIC, 3000

(56) Related Art
WO 1999/017801
Sano et al, (1992), PNAS, Vol 89, p1534-1538
WO 1999/017801
Chen et al, (2000), J of Immunology, Vol 164, p 705-711
Chen et al, (2000), J of Immunology, Vol 164, p 705-711
Sano et al, (1992), PNAS, Vol 89, p1534-1538

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/002751 A3

(51) International Patent Classification⁷: **C12N 5/00**, C07K 17/00, C12N 15/12, C07K 19/00, 14/36, 14/465, 14/495, 14/525, 14/54, 14/705, A61K 35/14, 35/28, A61P 37/00, 35/00, C12N 5/06

(21) International Application Number: PCT/US01/20946

(22) International Filing Date: 2 July 2001 (02.07.2001)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/215,580 30 June 2000 (30.06.2000) US

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).



(71) Applicant (for all designated States except US): **UNIVERSITY OF LOUISVILLE RESEARCH FOUNDATION, INC.** [US/US]; Jouett Hall, University of Louisville, Louisville, KY 40292 (US).

Published:

— with international search report

(88) Date of publication of the international search report: 3 October 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SHIRWAN, Haval** [US/US]; 8141 Copper Creek Drive, Louisville, KY 40222 (US).

(74) Agent: **VIKSNINS, Ann**; Schwegman, Lundberg, Woessner and Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).

WO 02/002751 A3

(54) Title: ALTERATION OF CELL MEMBRANE

(57) Abstract: Methods and compositions are provided for the persistent modification of cell membranes with exogenous proteins so as to alter the function of the cell to achieve effects similar to those of gene therapy, without the introduction of exogenous DNA. DNA sequences, the proteins and polypeptides embodying these sequences are disclosed for modulating the immune system. The modulations include down-regulation, up-regulation and apoptosis.

ALTERATION OF CELL MEMBRANE FOR NEW FUNCTIONS**FIELD OF THE INVENTION**

5 This invention relates to the persistent modification of cell membranes so as to alter the function of the cells. The compositions and methods of this invention achieve effects similar to those of gene therapy without the introduction of exogenous DNA. A useful alteration of cell function is the induction of apoptosis.

10

BACKGROUND OF THE INVENTION

It has long been the goal of experimental biology and medicine to induce cells to behave in predictable ways and to alter the behavior of cells in ways that are beneficial to a subject. For example, if undesired cells could be induced to alter their behavior to undergo apoptosis while normal cells retain normal function, subjects with a disease caused by proliferation of undesired cells would obtain relief from the disease. Similarly, if tissue-rejecting cells can be eliminated or their behavior changed, transplantation with tissues foreign to the subject can be successful.

15

Gene therapy has been proposed for selected diseases in order to correct or modify pathological or physiological processes. In gene therapy as it is generally termed, specific DNA is introduced into a tissue and organ, where it is produces various proteins that will correct or ameliorate the condition. It is an unpredictable therapy, depending on potentially dangerous expression vectors and the uncertain efficiency of delivery, which is often low. Moreover, gene therapy is considered to have dangerous side effects, such as sustained expression in desired cells or tissues past the desired duration of therapy, or the introduction of genetic modifications in undesired tissues or cells. Because of such adverse effects, as discovered in human clinical trials, much caution is advised before gene therapy is put in practice.

20

DNA-based gene therapy has been the subject of intense studies during the past few decades because of the tremendous potential it offers for the treatment of inherited diseases and other pathologic conditions for which the expression of selected proteins may offer treatment. For example, gene therapy

can be used for immunomodulation either to enhance the capacity of the immune response to deal with infections or tumors, or to down-regulate the immune response for the prevention of autoimmunity or foreign graft rejection. Despite vast efforts in the past two decades, the safe and effective application of gene 5 therapy to the treatment of diseases has been extremely limited. Among the apparent drawbacks of gene therapy are the possibility of causing permanent change in the DNA complement of the host; uncertain tissue specificity; expression of the encoded protein beyond the intended duration of therapy; and high cost. It would therefore be extremely beneficial if cells and tissues could be 10 modified to express proteins of interest in a short period of time without the introduction of foreign DNA.

Whereas gene therapy has generally focused on the problem of delivering nucleic acids into cells, much fundamental knowledge concerning the important role of cell surface molecules has been gained through studies of signal 15 transduction in cells of the immune response, which are readily accessible and have well-understood functions. The immune response is regulated by the interaction of several different cell types, which react to the presence of foreign antigens. The adaptive immune response is critical to the survival of vertebrates in an environment full of pathogenic microorganisms. Individuals who, due to 20 inborn genetic defects, exposure to chemotherapeutic agents or infection by such viruses as human immunodeficiency virus (HIV), lack a fully functional immune response are susceptible to infections that an individual with a healthy immune response would readily withstand. However, the immune system does not always function in ways that are beneficial to the organism. Its dysregulation 25 leads to autoimmunity and tumors. The immune system also serves as a barrier to the transplantation of foreign grafts, such as cells taken from an individual other than the transplant recipient. Transplantation permits the replacement of failed cells, tissues, or organs in otherwise terminal diseases, while bone marrow transplantation can treat hematopoietic disorders, malignancies, autoimmune 30 disorders, and other diseases. For transplantation to be successful, it is necessary either to suppress the adaptive immunity or to "teach" the recipient's immune system to accept these foreign antigens as native.

The immune response to foreign antigens is initiated by naive T cells that use clonally-expressed T cell receptors (TCRs) to recognize antigens such as peptides presented by self-major histocompatibility complex (MHC) molecules. This recognition reaction, when accompanied by costimulatory signals provided by antigen-presenting cells (APCs), has been thought to result in full T cell activation. A productive T-cell response is now seen as requiring three distinct signals. Signal 1 is generated by T-cell receptor interaction with the major histocompatibility complex (MHC) antigen/peptide on antigen-presenting cells (APCs). Signal 2 is mediated by the engagements of costimulatory molecules, such as B7/CD28 and CD40/CD40L, on T cells and APCs. Signal 3 is transduced via cytokines elaborated by T cells and APCs that have received both 5 Signal 1 and 2. The transduction of these 3 signals drives T cells and APCs to proliferation and differentiation into effectors for the generation of a productive immune response. The lack of any of these signals during the T-cell response 10 results in T-cell anergy and immune nonresponsiveness. For example, tumors 15 evade the immune system by preventing the transduction of one of these signals.

Upon activation, T cells proliferate and differentiate into effector cells that evoke immunological mechanisms responsible for the clearance of antigens from the system. A period of death then follows during which most of the 20 activated T cells undergo apoptosis-mediated “activation-induced cell death” (AICD) and effector activity subsides. Apoptosis is a complex process that involves a series of extra- and intracellular signals that converge on the activation of enzymes called caspases that commit the cell to apoptosis.

Transplantation of foreign cells (such as bone marrow and stem cells), 25 tissues (such as pancreatic islets), and organs (such as kidneys, hearts, livers) has become an important and effective therapeutic alternative for patients with selected terminal diseases. The transplantation of foreign grafts between genetically different patients (allografts between members of the same species or 30 xenografts between members of different species) is, however, limited by the ability to control the immunological recognition and rejection of the graft by the recipient.

Bone marrow (BM) transplantation has been viewed as an extraordinarily promising treatment for hematopoietic and autoimmune disorders and for certain cancers. One obstacle to bone marrow transplantation is the possibility of rejection of the transplanted tissue, mediated by the host's T cells and NK cells.

5 Graft-versus-host-disease (GvHD) is another possible adverse consequence of bone marrow transplantation. Donor T cells in the transplanted tissue can mount an immune response against the host's vital organs, often leading to death of the host. Host-versus-graft reactions and GvHD therefore limit the clinical use of bone marrow transplantation, which might otherwise be widely used to treat 10 various diseases and to prevent foreign graft rejection.

Pharmacological agents that cause immunosuppression are now a mainstay of regimens for the control of allograft rejection. Although such drugs are effective in reducing the severity of rejection episodes, they are nonspecific and fail to create a state of permanent graft-specific tolerance. Continuous 15 exposure of the recipient to these immunosuppressive agents is therefore associated with a significantly increased risk of opportunistic infections and malignancies. The need remains to develop more selective and long-lasting methods to prevent BM rejection.

20 Additionally, these nonspecific immunosuppressive agents can induce serious and undesirable side effects in the host. These adverse effects often outweigh the benefits for patients with diseases in which the body identifies certain parts of itself as "foreign" and launches an adaptive immune attack that results in autoimmunity, such as is observed in Type I diabetes, arthritis, lupus, and multiple sclerosis. It would be very desirable to be able to "teach" the 25 immune system to tolerate the "foreign" self-antigen.

It would also be very desirable to be able to "teach" the immune system to rid the organism of tumor cells. T cell-mediated cellular immunity is the most critical acquired response against tumors. A series of experimental studies has provided evidence that tumors evade T-cell-mediated immunity by several 30 different mechanisms. These mechanisms include: i) lack of Signal 1, due to inefficient display of MHC/tumor antigen bimolecular complexes on tumor cells or defects in the transduction of this signal; ii) absence of Signal 2, due to the

absence of costimulatory molecules on tumor cells; iii) induction of anergy in T cells; and iv) physical elimination of effector T cells via apoptosis. Although all of these mechanisms may be operative in patients with a large tumor burden, the lack of costimulation is believed to play the most critical role.

5 The need therefore remains to develop more rapid, selective and long-lasting methods to modulate cell function without the introduction of nucleic acids into cells for therapeutic purposes. The need also remains to develop a means of accomplishing the end of gene therapy without many of the risks attendant to the introduction of exogenous nucleic acids into an organism. Since much cell function is controlled
10 through the transduction of signals at the cell surface, a generally applicable method of attaching an agent to a surface would be useful. Among the uses of such a method would be: the modulation of cell function without the introduction of nucleic acids into the cell; the accomplishment of the end of gene therapy by alternative and potentially preferable means; and the manipulation of an organism's immune response in order to
15 diminish that response, as to treat autoimmunity or to forestall graft-versus-host disease, or to increase that response, as to treat tumors or infections.

20 The discussion of the background to the invention herein is included to explain the context of the invention. This is not to be taken as admission that any of the material referred to was published, known or part of the common general knowledge as at the priority date of any of the claims.

SUMMARY OF THE INVENTION

25 In one aspect, the present invention provides a method of attaching an agent to a cell surface, the method comprising contacting a biotinylated cell with a chimeric protein comprising (i) the agent and (ii) core streptavidin, wherein the chimeric protein is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, and SEQ ID NO:7.

30 In another aspect, the present invention provides a nucleic acid comprising a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7.

In another aspect, the present invention provides a method of attaching an immune costimulatory agent to a cell surface *in vivo*, the method comprising:

5 contacting a biotinylated cell with a chimeric protein comprising (i) the immune costimulatory agent and (ii) core streptavidin, wherein the immune costimulatory agent is an adhesion molecule, a cell growth factor, a cell toxin or a composition that induces apoptosis, down-regulation of the immune system or up-regulation of the immune system.

In another aspect, the present invention provides a method of attaching an agent to a cell surface *in vivo*, the method comprising:

10 contacting a biotinylated cell with a chimeric protein comprising (i) the agent and (ii) core streptavidin, the agent being an adhesion molecule, a cell growth factor, a cell toxin or a composition that induces apoptosis, down-regulation of the immune system or up-regulation of the immune system, wherein, after five days *in vivo*, at least 26% of the cells retain the chimeric protein attached to the cell surface.

The present invention provides a method of modifying a surface, the method comprising : (1) contacting a surface with one member of a binding pair which binds to the surface to form a decorated surface; and subsequently (2) contacting the decorated surface with a composition comprising the second member of the binding pair operably linked to an agent capable of modifying the surface and the function thereof. The binding pair may be avidin/biotin, streptavidin/biotin or antigen/antibody. By "avidin" is meant avidin and any fragment or derivative of avidin which retains strong binding to biotin. By "streptavidin" is meant streptavidin and any fragment or derivative of streptavidin which retains strong binding to biotin. Conversely, by "biotin" is meant any fragment or derivative of biotin which retains strong binding to avidin and streptavidin.

When the surface is a cell surface, the agent may be any compound inducing apoptosis; any agent down-regulating the immune system; any agent up-regulating the immune system; any adhesion molecule for cell tracking; any growth factor; or an antibody ligated to a toxin such as ricin or phytotoxin.

5 When the surface is a cell surface, the contact of step 1 produces a "decorated cell."

When the agent is selected to induce apoptosis, the agent may be a death-inducing molecule such as FasL, mFasL, TRAIL, TNF- α , TWEAK and the like. When the agent is selected to down-regulate the immune system, the agent may 10 be an anti-inflammatory cytokine such as IL-4, IL-10, TGF- β and the like.

When the agent is selected to up-regulate the immune system, the agent may be a costimulatory molecule such as B7, CD40L and the like or a pro-inflammatory cytokine such as IL-2, IL-12, IL-15, lymphotactin and chemokine.

15 In a preferred embodiment, the first member of the binding pair is biotin, the second member of the binding pair is a chimeric protein comprising streptavidin and a death-inducing molecule and the surface comprises target cells, tissues, organs or tumors.

20 In a more preferred embodiment, the first member of the binding pair is Sulfo-NHS-LC-biotin, the second member of the binding pair is SA-mFasL and the surface comprises a cell displaying the Fas receptor. In constructing the second member of the binding pair, streptavidin is fused to the extracellular portion of FasL through recombinant gene technology and the chimeric protein SA-mFasL is produced.

25 This construct is used in a method to modulate the immune system wherein a subject to be treated is first administered biotin to form decorated cells, followed by administration of SA-mFasL to form chimeric decorated cells. Only those cells in the tissue, organ or tumor which are biotinylated and express Fas will undergo apoptosis. In general, these cells include those cells which cause immuno-mediated injuries.

30 In another aspect of the invention, a subject who will benefit from down-regulation of the immune system is first administered biotin to form decorated cells and is then administered a construct comprising avidin or streptavidin and

a down-regulating molecule. Subjects who will benefit from down-regulation of the immune system include subjects suffering from autoimmunity, graft-versus-host disease, allergies, septic shock and vascular diseases. Subjects undergoing tissue or organ grafts also benefit from the down-regulation of the immune system.

5

In another aspect of the invention, a subject who will benefit from up-regulation of the immune system is first administered biotin to form decorated cells and is then administered a construct comprising avidin or streptavidin and an up-regulating molecule. Subjects who will benefit from an up-regulation of the immune system include subjects suffering from neoplasia or infectious disease.

10

In another aspect of the invention, systemic effects are avoided by biotinylating the cells *ex vivo* by isolation of the target or selection of a target that can be treated topically, followed by reintroduction of the decorated cells into the subject with subsequent second contact.

15

In another aspect of the invention, the members of the binding pair form a covalent bond with each other. In a preferred aspect, the first member of a binding pair comprises a cysteine residue and the second member of the binding pair comprises a cysteine residue. The two members are first placed together under conditions disfavoring the formation of a disulfide bond between the first member of the binding pair and the second member of the binding pair. In a subsequent step, the two members are placed under conditions favoring the formation of a disulfide bond between the first member of the pair and the second member of the pair.

20

In another aspect of the invention, the first member of a binding pair contacts a cell surface, a virus, the surface of a glass particle, the surface of a polysaccharide particle or the surface of a plastic particle to form a decorated surface. The decorated surface is then contacted with the second member of the binding pair, to form a second decorated surface. The cell, virus or particle comprising the second decorated surface is contacted with a cell or cell component of an organism either *in vitro* or *in vivo*, for the purpose of therapy, diagnosis, cell sorting or identification.

25

30

2002216763 23 Nov 2006

A recombinant nucleic acid is provided comprising the extracellular portion of FasL fused with streptavidin(SA-mFasL). Gene product is provided which is then attached to the cell surface, previously biotinylated *in vivo*.

5 Methods are taught for the use of gene products for long-lasting elimination of Fas expressing cells.

The methods of this invention can be modified to cause persistent binding of any protein to targeted cells, to mimic the result of DNA gene therapy without modification of the cell genetic material. This persistent binding of a protein to a targeted cell can be 10 considered to be gene therapy at the protein level.

Throughout the description and claims of this specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

15 **DESCRIPTION OF THE DRAWINGS**

Figure 1 is a schematic representation of the method to modify cell membrane for the expression of exogenous proteins for new functions.

Figure 2 shows the restriction fragment analysis of the expression vector containing the indicated clones.

20 Figure 3 shows biotin expression *in vivo*.

Figure 4 shows the *in vitro* expression of SA-mFasL on the surface of splenocytes.

Figure 5 shows SA-mFasL expression on the surface of bone marrow cells.

Figure 6 shows donor chimerism in lethally irradiated animals.

25 Figure 7 shows a graphic summary of allogeneic cells decorated with SA-mFasL blocking *in vivo* alloreactive responses.

Figure 8 shows a pancreatic islet cell expressing FasL (red) and streptavidin (B) compared to a control islet (A).

Figure 9 shows T cells modified to express SA-mFasL, activated by ConA.

30 Figure 10 shows expression of SA-CD40L on splenocytes.

DETAILED DESCRIPTION OF THE INVENTION

Because of the noted problems associated with gene therapy as it is commonly used, that is, the introduction and integration of foreign DNA into the genome, a method was sought for the accomplishing the results of gene therapy, that is, long lasting modification of cell function, without the limitations of DNA introduction. The question was raised as to whether the benefits of gene therapy could be obtained by the delivery of exogenous proteins, rather than nucleic acids, to the cells. A method that would safely permit modification of the cell membrane to achieve transient or long-lasting display of therapeutic molecules could have wide-spread application in the clinic. As disclosed in the present application, Applicant has discovered and here discloses such a method, which has the added advantage of rapid cell surface expression.

Because of the thermodynamically favored association of avidin or streptavidin with biotin, the complex formed can modify cell properties for a period that can rival or surpass the duration of the effect elicited by gene therapy that requires the introduction of nucleic acids. The present invention teaches a method of attaching an agent to a surface. Such an agent can be, by way of example and not of limitation, any of a series of immunomodulatory molecules such as death ligands, costimulatory molecules and cytokines. It is useful to attach such agents to surfaces, especially the cell surface, since such agents can be used to physically or functionally eliminate autoreactive, alloreactive, xenoreactive and inflammatory cells to prevent allograft and xenograft rejection, graft-versus-host disease and to treat autoimmunity, septic shock, inflammation, allergies, infections, neoplasias, and vascular diseases. Furthermore, this method can be used to display proteins with defined function on the surface of cells for *in vivo* trafficking as for the homing of selected hematopoietic cells for therapeutic purposes.

Definitions:

For the purposes of the present application, the following terms have these definitions:

“Agent” means a composition which elicits a measurable biological response or which is useful in the diagnosis, prevention or treatment of a disease or in the sorting or identification of cells, cell components or viruses.

5 “Binding pair” means two compositions that readily bind to each other; that have affinity for one another; or whose association is thermodynamically and kinetically favored.

“Cell surface” has its normal meaning in the art, comprising the phospholipid bilayer of a cell membranes and the molecules directly or indirectly associated with the bilayer.

10 “Decorated surface” means a surface to which a member of a binding pair has become bound.

“Expression” means, in addition to its conventional meaning in the art, the placement of a functional protein on or within a cell.

15 “Plastic” means polystyrene, polypropylene, polyethylene or any other polymer capable of being formed into a solid surface.

“Protein” means a protein or polypeptide that is native, non-native, synthetic or modified as by covalent binding.

20 “Surface” means a cell surface, the surface of a virus, the surface of a particle (e.g., the surface of a glass particle, the surface of a polysaccharide particle, the surface of a plastic particle), the surface of a phospholipid bilayer or the surface of a solid matrix.

“Target cell” means a cell targeted for apoptosis, that is, a cell in which it is desired to induce apoptosis.

25 An important application of regulation of the immune response is the induction of tolerance to alloantigens, xenoantigens and autoantigens to prevent foreign graft rejection, treat or prevent autoimmunity and GVHD. T cells are the most critical cells of adaptive immunity, which requires three distinct signals (Signal 1, 2 and 3) for a productive response. Signal 1 is generated by T-cell receptor interaction with the MHC/peptide complex on APCs. Signal 2 is mediated by the engagements of costimulatory molecules, such as B7/CD28 and CD40/CD40L on T cells and APCs. Signal 3 is transduced via cytokines elaborated by T cells and APCs that have received both Signal 1 and 2. The lack of any of these signals during T-cell response to antigens may serve as one of the

most effective mechanisms by which tolerance is induced. Upon activation, T cells undergo a state of antigen-driven proliferation that allows up to a 1200-fold clonal expansion. A period of death then ensues during which more than 95% of the activated T cells undergo apoptosis (programmed cell death) while the 5 remaining cells differentiate into memory cells as the amount of antigen in the system declines.

Apoptosis or “programmed cell death” plays a central role in both the development and homeostasis of multicellular organisms. Apoptosis can be induced by multiple independent signaling pathways that converge upon a final effector mechanism consisting of multiple interactions between several “death receptors” and their ligands, which belong to the tumor necrosis factor (TNF) receptor/ligand superfamily. The best characterized death receptors are CD95 (“Fas”), TNFR1 (p55), death receptor 3 (DR3 or Apo3/TRAMO), DR4 and DR5 (apo2-TRAIL-R2). A final effector mechanism is mediated by the caspase group 10 of proteins.

In copending International Application PCT.US01/02256, the teachings of which are incorporated by reference, it is disclosed that Fas-/FasL- induced apoptosis plays a central role in protecting immunologically privileged sites, such as the central nervous system, testes and eyes from immune attack. 20 Allogeneic and xenogeneic tissues transplanted into these sites, for instance, are resistant to rejection. Of great importance is the major role apoptosis plays in maintaining homeostasis in the immune system via activation-induced cell death (AICD). AICD is primarily mediated by apoptotic signals transduced by the Fas/FasL interaction. Fas is a 45 kDa, type I, cell surface protein consisting of a cysteine-rich extracellular domain, which transduces a death signal. Fas 25 mediates effector function by interacting with FasL, a type II membrane protein of about 40 kDa. T cells are the primary cell type in the body that express FasL, upon activation. The expression of this molecule makes the activated T cells susceptible to apoptosis, which is induced in an autocrine fashion as Fas engages with FasL on the same cell. Fas on an activated T cell may also interact in a paracrine fashion with FasL on another activated cell. These two pathways of 30 apoptotic T cell death mediated by the Fas/FasL system serve as a homeostatic

mechanism for controlling the size of antigen-stimulated T cell clones.

Deregulation of this system results in immunological disorders.

The molecular mechanism of Fas/FasL-mediated apoptosis has been studied in great detail. Binding of three FasL molecules with Fas induces 5 trimerization of Fas receptor via c-terminus death domains (dds), which in turn recruit an adapter protein FADD (Fas-associated protein with death domain) and Caspase-8. The oligomerization of this trimolecular complex, Fas/FADD/caspase-8 results in proteolytic cleavage of proenzyme caspase-8 into 10 active caspase-8 that, in turn, initiates the apoptosis process by activating other downstream caspases through proteolysis, including caspase-3.

Apoptosis of activated T cells results in tolerance to allografts and 15 xenografts, including bone marrow and other organ and cellular transplantation.

Purging of activated T cells also relieves the symptoms of allergies and other 20 immune-induced diseases. Included in the latter are auto-immune disorders such as multiple sclerosis, lupus erythematosus, sarcoidosis and rheumatoid arthritis.

Many disorders, including some tumors, are dependent on lymphocyte functions 25 that lead to persistence of the disorder. Many hematological disorders could be treated with bone marrow stem cell transplants if the immune response could be regulated so as to induce tolerance to the foreign stem cells. Among these disorders are leukemias, lymphomas, aplastic anemia, sickle cell and Cooley's anemia and the like. All of these disorders may be controlled permanently or 30 temporarily by apoptosis of activated immune cells, including T cells.

Strategies are provided for immunomodulation using rapid cell-surface 25 display of proteins, which comprise the construction of chimeric cDNAs encoding the functional portions of an immunoregulatory protein with one member of a binding pair. Table 1 is a summary of such constructs. Choice of constructs may be based on factors such as: the nature of the foreign antigen provoking adaptive immunity; whether the relief to be sought is temporary or permanent; whether a commitment to death is desired or additional downstream 30 regulation of apoptosis is preferred. It is to be understood that the constructs listed below are representative of the invention only and are not limitations. For example, the constructs have been made using core streptavidin (CSA).

Fragments and derivatives of streptavidin as well as the entire streptavidin molecule can be as readily used to make equivalent constructs. Therefore, in the specification, SA is generally used to mean whole, fragmented or derivatized streptavidin. Those skilled in the art can readily, without undue experimentation, follow the teachings of this invention to make constructs comprising a first member of a binding pair and a second member of a binding pair ligated to a known immunomodulatory molecule.

TABLE 1

10

Vector	Insert	SEQ ID #	Function	Application
pCSA-mFasL	6His-RS-CSA-SRIPE-Extracellular portion of FasL	1	Apoptosis	Induce apoptosis in immune-reactive cells, smooth muscle cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock and likes) and vascular diseases.
pCSA-TNF α	6His-RS-CSA-SRIPE -Extracellular portion of TNF α	7	Apoptosis	Induce apoptosis in immune-reactive cells, smooth muscle cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock and likes) and vascular diseases.
pIL-4-CSA	Mature portion of IL-4-6His-RS-CSA-SRIPE		Immune down-regulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, and likes).

Vector	Insert	SEQ ID #	Function	Application
pIL-10-CSA	Mature portion of IL-10-6His-RS-CSA-SRIPE	5	Immune down-regulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, and likes).
pTGF β -CSA	Active portion of TGF- β -6His-RS-CSA-SRIPE	3	Immune down-regulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock, and likes).
pCyto-CSA	Active portion of other immunoregulatory cytokines-6His-RS-CSA-SRIPE		Immune down-regulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock, and likes).
pIL-2-CSA	Mature portion of IL-2-6His-RS-CSA-SRIPE	4	Immune up-regulation	Up-regulate immune-reactive cells for the treatment of infections and cancers.
pB7-CSA	Extracellular portion of B7-6His-RS-CSA-SRIPE	2	Immune up-regulation	Up-regulate immune-reactive cells for the treatment of infections and cancers.
pCSA-CD40L	6His-RS-CSA-SRIPE -Extracellular portion of CD40L	6	Immune up-regulation	Up-regulate immune-reactive cells for the treatment of infections and cancers.
padhesion-CSA	Active portion of adhesion molecules -6His-RS-CSA-SRIPE		Immune down- or upregulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock, and likes). Up-regulate immune-reactive cells for the treatment of infections and cancers.

Vector	Insert	SEQ ID #	Function	Application
pchemokin es-CSA	Active portion of immunoregulatory chemokines-6His- RS-CSA-SRIPE		Immune down- or upregulation	Down-regulate immune-reactive cells for the treatment of immune-based disorders (GVHD, rejection of foreign grafts, autoimmunity, inflammation, allergies, septic shock, and likes). Up-regulate immune-reactive cells for the treatment of infections and cancers.

5 A preferred production cell for the production of chimeric compositions encoded by the DNA constructs is the *Drosophila* system that is commercially available. However, those skilled in the art of producing chimeric compositions will recognize that many other expression systems and vectors are suitable for production of the chimeric compositions of this invention. Included in these other systems are *Escherichia coli*, yeast and mammalian cell cultures.

10 The inventions disclosed in copending international application PCT/US00/34554, the teachings of which are hereby incorporated by reference, are directed to the apoptosis of lymphocytes and tumor cells. The present invention discloses an improved method of expressing immunomodulatory molecules with therapeutic interest on the surface of cells via binding pair coupling. Avidin or streptavidin/biotin is an example of a binding pair useful in the present invention. A modified FasL protein is described in detail as an example of such a composition. Bone marrow cells, endothelial cells, splenocytes, tissues such as pancreatic islets and organs such as hearts are examples of cells amenable to cell surface alteration. It has now been found that 15 this method can be used to down-regulate or up-regulate immune responses to antigens. This method, therefore can be used for the prevention of foreign graft rejection, for the prevention or treatment of autoimmunity and the up-regulation of immune response to tumors and infections. Additionally, it is provided that proteins other than those with immunoregulatory functions can be bound to cells 20 without causing systemic side effects.

25

The invention will now be described with specific examples. The experimental procedures described herein are representative of compositions and methods for persistent binding “expression” of agents to cells, tissues or organs. One embodiment specifically described is a chimeric composition, SA-mFasL, produced by fusion of DNA coding for core streptavidin protein with that of DNA coding for FasL and expressing the chimeric composition (SEQ#1) in a suitable production cell.

5 The target cells are first biotinylated *ex vivo* or *in vivo* and then SA-mFasL is administered in order to cause apoptosis, resulting in, for example, 10 induction of donor-specific long-term survival and/or tolerance. Another example that is described in detail is the transplantation of pancreatic islet cells along with immune cells treated according to the methods of this invention, with subsequent enduring pancreatic insulin production. Another example is the long term survival of ectopic heart transplants. Others examples are also provided. 15 Following the teachings of the following examples, one skilled in the art can readily apply these to alter the membrane function of other cells.

Alternatively, the DNA encoding such chimeric compositions may be applied *ex vivo* or *in vivo*, leading to permanent changes such as the apoptosis of activated T cells.

20 The following examples are disclosed in order to illustrate the present invention as it is applied in practice and do not limit the scope of the appended claims.

25 Example 1. Cloning of core streptavidin for the generation of chimeric proteins.

The present invention discloses technology for cell-surface modifications to express exogenous proteins without the introduction of nucleic acids into cells and comprises: i) generation of a chimeric molecule consisting of core streptavidin and functional domains of a desired protein, ii) modification of the 30 cell membrane with biotin, and iii) decoration of the biotinylated cells with the chimeric molecule (Fig. 1). To accomplish this, genomic DNA encoding streptavidin was cloned from *S. avidinii* using specific primers in PCR. 5' and 3'

primers were designed to incorporate sequences for selected restriction enzyme sites and amino acids that allow three dimensional flexibility, proper folding and function. The gene was cloned into the TA cloning vector, sequenced and subcloned into the pMT/Bip/V5-HisA vector for expression in a high-yield insect expression system (DES tm, Invitrogen).

Example 2. Construction of chimeric genes for expression in production cells.

Total RNA was prepared from human cell lines or peripheral blood cells and 2 ng of this RNA was reverse transcribed into the first strand of DNA using 10 oligo (dT)18 as a primer for reverse transcriptase. One-tenth of this cDNA preparation was then amplified, using three sets of sense and antisense primers specific for human IL-2, IL-4, IL-10, TGF- β , FasL, TNF- α , B7.1, and CD40L in 8 separate PCR amplifications. The 5' and 3' primers were designed to include 15 restriction enzyme sites for cloning and several amino acid residues to facilitate the proper folding of the product. These primers amplified DNA bands of expected sizes for all these genes of interest. These PCR products were then cloned into the TA cloning vector (Invitrogen, San Diego, CA) and a library prepared from this material was screened using the same oligonucleotide primers in PCR amplifications. The positive clones were digested with appropriate 20 restriction enzymes, leading to the release of expected size of fragment for each gene (Fig. 2) shown in Table 1. All indicated clones were sequenced and found to have the expected characteristics. These clones were then fused to CSA in the pMT/Bip/V5-HisA vector either as N-terminus or C-terminus proteins to 25 facilitate correct three dimensional structure and function (see SEQ IDs in Table 1).

Chimeric proteins were subcloned in frame with the *Drosophila* BiP secretion signal in the pMT/Bip/V5-HisA vector for expression in a high-yield insect expression system (DES tm, Invitrogen). Recombinant vectors were 30 transiently transfected into *Drosophila* S2 cells using calcium phosphate. Cells were then pulsed with copper sulfate to activate the inducible metallothionein promoter driving the expression of chimeric proteins. Culture medium was

collected at various times post-activation, dialyzed to remove the copper sulfate, and analyzed for chimeric proteins using ELISA and Western blot.

Example 3. Biotinylation of cells, expression of recombinant proteins, and time
kinetics of expression in vitro

The therapeutic use of this protein-based approach requires successful biotinylation of cells, tissues, or organs under physiological conditions and attachment to these cells chimeric proteins consisting of CSA and molecules with therapeutic potential (Fig. 1).

10 The optimum conditions for biotinylation were first determined. Single cell suspensions were prepared from spleen or bone marrow of the rat. One million cells were incubated in various concentrations, ranging from 1.5 to 150 μ M, of EZ-Link biotin (trade name of Sulfo-NHS-LC-biotin, Pierce, Rockford, IL) in saline at room temperature for 30 minutes. After extensive washing to 15 remove free biotin, cells were either cultured or used for staining with fluorescein (APC)-labeled streptavidin in order to assess the level of biotinylation. 100% of the cells were positive for biotin at 5 μ M biotin concentration (data not shown).

To determine how long biotin persists on actively dividing cells and the optimum dose of biotin (15 μ M), biotinylated splenocytes were cultured in the presence of a T cell mitogen, concanavalin A (ConA), to stimulate T-cell proliferation. ConA-stimulated and unstimulated cells were harvested at various times post culture and stained with APC-streptavidin and analyzed by flow cytometry. Splenocytes without streptavidin and splenocytes with biotin alone served as background staining. Over 50% of the cells maintained biotin on the surface for 20 days, the longest time point tested. A significant portion (>35%) of actively dividing cells also expressed biotin at this time period. It was next tested to determine whether biotinylation interferes with the proliferation of splenocytes. Splenocytes were activated with 2.5 μ g/ml ConA for various days in 96-titer plates. Cultures were pulsed with tritiated thymidine, harvested, and analyzed for DNA-associated radioactivity as an indication of proliferation.

kinetics and levels as compared with unmanipulated splenocytes, suggesting that biotinylation at concentrations ranging between 1.5-15 μ M does not have a significant effect on the proliferative function of the cells (data not shown).

5 Biotinylated vascular endothelial cells in culture were tested to determine the time kinetics of biotin on the cell surface. The rationale for this set of experiments is to test how long biotin persists on the surface of differentiated cells with minimal turnover in tissues such as heart vasculature. Monolayer cultures of rat aortic endothelial cells were treated with 15 μ M concentration of biotin in six-well plates for 30 min at room temperature. Cells were extensively 10 washed and one well of the culture was digested with trypsin at different times post biotinylation, single cell suspension was prepared and analyzed in flow cytometry using APC-streptavidin. A significant portion of endothelial cells (>15%) expressed biotin 20 days in culture, the longest time point tested.

15 Splenocytes were tested for the expression of one of the recombinant proteins with immunomodulatory function, the extracellular portion of FasL fused with core streptavidin (SA-mFasL). Splenocytes were biotinylated as stated above, incubated with 100 ng SA-mFasL for 20 min, and maintained in a 37°C incubator for defined periods of time at which cells were stained with an 20 anti-FasL antibody. Almost all the cells expressed SA-mFasL on day 5 and significant number of cells (>25%) expressed SA-mFasL on day 10, the last time point analyzed (Fig. 4).

25 One of the objectives of the proposed protein-based gene therapy to use immunomodulatory molecules such as SA-mFasL to facilitate the engraftment of BM cells in a foreign environment. Therefore, BM cells were tested for biotinylation and expression of FasL at the protein level. BM cells were harvested from femurs and tibia of rats using standard protocols. Single cells suspension was prepared and then biotinylated under the above conditions. After extensive washing to remove biotin, BM cells were treated with SA-mFasL (50 ng/million cells) for 20 min on ice. Cells were then washed extensively to 30 remove the free SA-mFasL and stained with an antibody against FasL (PE-MFL4) in flow cytometry. As shown in Fig. 5, 100% of the BM cells expressed SA-mFasL.

Example 4. Time kinetics of expression *in vivo*

5 Splenocytes were labeled with a lipophilic dye (CFSE), which incorporates and persists in the cell membrane for an extended period of time *in vivo*, modified to express biotin or SA-mFasL, and injected into syngeneic animals intravenously. Splenocytes were harvested at various times post injection and stained with APC-streptavidin or anti-FasL antibody for analysis in flow cytometry. As shown in Fig. 3, biotin was detected on the surface of 40% of CFSE positive cells 10 days after *in vivo* injection, the latest time point tested.

10 Similarly, CFSE positive cells modified to express SA-mFasL were positive (26%) for the expression of this molecule 5 days after injection. Taken together, these data clearly demonstrate that this approach allows for the expression of proteins for an extended period of times *in vivo* to perform function of interest.

15 Example 5. Blocking of alloreactive immune response *in vitro* using target cells
expressing immunomodulatory molecules

To test whether cells modified to express immunomodulatory proteins of interest using this approach can be used to prevent alloreactive responses, allogeneic splenocytes were modified to express SA-mFasL and used as targets in *in vitro* proliferation assays. Briefly, splenocytes from F344 rats were biotinylated, treated with culture supernatants or purified SA-mFasL. Supernatant from cells transfected with a nonfunctional construct served as control and are referred to as S2 supernatant throughout this application. These cells were then irradiated and used as stimulators for alloreactive responses in a standard five-day mixed lymphocyte culture. Lymphocytes harvested from the lymph nodes of PVG.IU rats were used as responders. Targets expressing SA-mFasL completely blocked the proliferative response of lymphocytes as compared to culture without SA-mFasL. These data demonstrate that biotinylated splenocytes can serve as antigen-presenting cells and that SA-mFasL expressed on the surface of splenocytes block an alloreactive immune response *in vitro*. Additionally, these data demonstrate that biotinylation does not interfere with the antigen-presenting function of cells since cells decorated

with biotin and treated with S2 control supernatants did not interfere with their capacity to stimulate lymphocytes. These data, therefore, demonstrate that expression of the proteins on the surface of cells does not interfere with the function of cells nor the expressed molecules. This is a critical step in validating protein-based expression as a means of immune regulation.

5

Example 6. Toxicity of SA-mFasL protein.

In general, most cells that express Fas are activated, such as activated lymphocytes, or fast-dividing, such as hepatocytes, or undesirable, such as tumor cells. However, it had to be asked whether SA-mFasL might be toxic. It has been shown that antibodies against Fas induce fulminant liver damage in selected mouse strains when injected intravenously or intraperitoneally. This is believed to be due to the expression of Fas on hepatocytes. In order to ascertain whether the chimeric protein of this invention was likewise toxic, rats were injected intraperitoneally or intravenously with 2×10^7 splenocytes bearing biotin-SA-mFasL or 8×10^7 bone marrow cells bearing biotin-SA-mFasL intravenously. The animals were closely monitored for 10 days and then sacrificed for gross pathological and anatomical analysis. No noticeable pathology was found in animals injected with cell-biotin-SA-mFasL as shown in Table 2.

10

15

20

TABLE 2

Group	Number	Cells	Route of injection	Side effects
I	3	2×10^7 splenocytes-bio	intraperitoneal	none
II	3	2×10^7 sp.-bio-SA-mFasL	intraperitoneal	none
III	3	8×10^7 bone-marrow-bio	intravenous	none
IV	3	8×10^7 bone-marrow-bio-SA-mFasL	intravenous	none

25

Example 7. FasL expressing BM cells rescue lethally irradiated rats

The major premise of this approach is to use protein-based expression as an alternative to DNA-based gene therapy. Prevention of BM cell rejection by this approach of modifying BM cells for immune evasion will suffice for this condition. Bone marrow cells were harvested from PVG.R8 or ACI rats, modified to express biotin or SA-mFasL, and administered i.v. at $0.7\text{-}1\times 10^8$ cells/animal into lethally (950 cGy) irradiated rats. Irradiated animals receiving no cells served as controls. All the animals receiving no cells expired within 8-9 days (n=6) whereas all the animals (n=12) receiving BM cells manipulated to express bio or SA-mFasL were 100% chimeric (Fig. 6) and survived indefinitely (>100 days). These data clearly demonstrate that BM cells are safely manipulated by this novel approach to express proteins of interest for therapeutic purposes without affecting their long-term engraftment capacity.

15 Example 8. SA-mFasL expressing BM cells block alloreactive responses *in vivo*

BM cells expressing an immunomodulatory molecule such as SA-mFasL were used to induce allotolerance as target cells *in vivo*. PVG.1U or WF rats were administered i.v. with $0.7\text{-}2\times 10^8$ PVG.R8 or ACI BM cells, respectively, expressing FasL or biotinylated cells treated with control S2 supernatant. Mesenteric lymph nodes were harvested 60 days after injection and used as responders to PVG.R8 or ACI cells in a standard mixed lymphocyte assay. There was a complete absence of response to donor antigens. This *in vivo* immune nonresponsiveness was donor specific as the response to third party antigens was intact. This effect was SA-mFasL-specific as animals receiving PVG.R8 cells treated with control S2 supernatant generated a normal response to donor as well as third party antigens as compared with the response of naïve animals (Fig. 7). These data validate the *in vitro* blocking observation and provide direct evidence for the immunomodulatory effect of this approach *in vivo*.

30 Example 9. Modification of heart to express CSA-protein on vascular endothelium

The methods of the present invention have been applied to effect protein-based expression on organs. Modification of organs, rather than the host, *ex vivo* to express proteins of interest presents a desired therapeutic approach in selected settings. *Ex vivo* manipulation avoids complications that may arise if the host were treated, and which may include, but are not limited to, undesirable side effects. Therefore, the heart was used as a test system to express biotin, streptavidin, and SA-mFasL in ACI rats and B10.BR mice at 37°C. Briefly, the aortas of excised hearts were cannulated and perfused in a Langendorff retrograde perfusion system at a pressure of 96 mmHg with modified Krebs-Henseleit solution (KH). Cardiac contraction force was monitored with a latex balloon introduced into the left ventricle. The perfusion protocol for rat hearts was: 20 min with KH to allow stabilization of cardiac function; 20 min KH containing 5 µg/ml EZ-Link Sulfo-NHS-Biotin (Pierce); 10 min KH for biotin washout; 20 min KH containing 0.5 µg/ml of either Streptavidin-FITC (Zymed) or SA-mFasL; 10 min KH for streptavidin washout. Control hearts were perfused with KH solution only. Left ventricular pressure decreased during 80 min of perfusion from 98±7 to 91±6 mmHg in all three groups (n=5/group). Coronary flow did not decrease significantly from baseline values of 10.4±0.6 ml/min (n=15). These data indicate that endothelial biotinylation and decoration with SA-mFasL have no short-term detrimental effect on cardiac function and that FasL is not directly toxic to the coronary arteries.

It was next determined whether biotinylated endothelial lining of the coronary arteries can be “decorated” with SA-mFasL at 4°C. This temperature is widely used in the clinical setting for extracorporeal cardiac preservation before transplantation. Mouse (B10.BR) and rat (ACI) hearts were arrested with a magnesium cardioplegic solution (KH+16 mM MgSO₄) and were perfused at 4°C according to the protocol delineated above (biotin, KH, streptavidin-FITC or SA-mFasL, KH). Biotin on vascular endothelium was detected by streptavidin-FITC whereas SA-mFasL was detected using antibodies against streptavidin (Zymed) and FasL (MFL4) as primary and FITC-labeled proper secondary antibodies. Hearts were removed and frozen at -80°C on the stage of a modified microtome (Polycut S, Reichert-Jung). Sequential cryosections (~200 micron)

were performed and the presence of FITC-labeled streptavidin or antibodies was determined with an Axioplan microscope (C. Zeiss). The stage of the microscope was modified to accept the cooling block of the microtome. Images of cardiac sections were acquired at a magnification of 10x. Fluorescence was 5 detected in cardiac vasculature, demonstrating the feasibility of biotin and streptavidin conjugation to vascular endothelium under hypothermic conditions, in a preservation medium widely used in clinical practice. The presence of SA- 10 mFasL in the graft endothelium perfused with SA-mFasL, but not control S2 supernatant, was verified using FITC-MFL4 mAb. These data show that SA- mFasL can be introduced into coronary arteries via biotinylation in conditions of 15 extracorporeal organ preservation. Manipulation of the graft as described above is performed within 10-60 min, at 4°C, using a standard preservation solution. Thus, this approach meets the requirements of temperature, duration, and conditions used in the clinic.

15

Example 10. Hearts modified to express biotin or SA-mFasL survived indefinitely when transferred into syngeneic host.

The question arose whether this protein-based modification of organs affects the function. If hearts modified to express bi or SA-mFasL were 20 heterotopically transplanted into syngeneic recipients. Graft survival was assessed by daily abdominal palpation of the grafted heart. As shown in Table 3, there was no detectable difference between the survival of graft recipients transplanted with syngeneic hearts decorated with SA-mFasL (n=4; >56 days) or KH-perfused control syngeneic hearts (n=4; >74 days).

25

Table 3. Survival of heart grafts expressing SA-mFasL

Group	Recipient	N	Donor	Treatment	Rejection time (days)	MST
A	BALB/c	5	BALB/c	—	>100, >100, >100, >100, >100	>100
B	BALB/c	4	BALB/c	Perfusion	>74, >74, >88, >88	>74
C	BALB/c	4	BALB/c	Perfusion + SA-	>56, >56, >68, >68	>56

Example 11: Chimeric decorated islet cells.

Pancreatic islet transplantation is widely contemplated for the correction of diabetes. To test whether the methods of this invention can be effective for the prevention of rejection of islet allografts, diabetes was induced in BALB/c recipients by intraperitoneal injection of streptozotin (260 mg/kg). Animals 5 demonstrating blood glucose levels >450 mg/dl for at least three consecutive days were used as recipients of minor+major histocompatibility antigens-disparate C57BL/10 islets. Islets were harvested from C57BL/10 mice and isolated on a Ficoll gradient according to standard protocol. Splenocytes were 10 harvested from C57BL/10, biotinylated (15*l*M), decorated with FasL or control supernatant, and irradiated (2000 rads). One million of splenocytes were cotransplanted with ~400 islet cells under the kidney capsule of diabetic Balb/c mice. Survival and function of the islets were assessed by monitoring blood 15 glucose levels starting three days after transplantation. FasL-decorated splenocytes prolonged the survival of all allogeneic islets beyond nine days, the date at which the glucose levels of control mice had risen to near the pretransplantation level. Figure 8 shows a pancreatic cell expressing FasL and streptavidin.

In a further example of this technique, mouse pancreatic islets were 20 harvested by digestion with collagenase and purified using Ficoll gradients. Islets were cultured overnight and then modified with biotin (5*l*M) and decorated with SA-mFasL (~100mng/ml) or S2 supernatant as control. Inspection of islets by confocal microscopy showed high levels of biotin and 25 SA-mFasL. Islets decorated with SA-mFasL remained viable in culture for over 10 days as determined by visual inspection and trypan blue exclusion.

Taken together, these data clearly demonstrate that cells, tissues, and complex organs, such as hearts, can be modified with biotin to display proteins chimeric with streptavidin for extended periods of time without detectable toxicity.

30

Example 12. Prevention of GVHD.

Bone marrow transplantation (BMT) has the potential to treat a variety of genetically inherited and acquired hematological disorders, induce tolerance to autoimmune antigens and foreign grafts. BMT is, therefore, perceived as a natural way of performing gene therapy. The major complication is, however, graft-versus-host disease, which is fatal in most instances. T cells are considered to be the most important cells to cause GVHD. A method allowing specific elimination of T cells in the BM inoculum that cause GVHD has wide-spread clinical application. Conventional gene therapy has been used to introduce “suicide” genes into T cells. Once GVHD occurred, the suicide machinery was activated to eliminate these cells. The obstacles of conventional gene therapy, however, are major problems for routine use of this approach in clinics. A more “noninvasive” approach is needed to achieve this “suicide” approach.

This methods of the present invention are well suited for the “suicide” approach or immune down-regulation. Display of apoptotic (death ligands) or down-regulatory molecules (anti-inflammatory cytokines; CSA-IL-4, IL-10, TGF- β) in bone marrow cells (including mature T cells) are anticipated to physically or functionally eliminate these cells upon their recognition of host antigens and activation. As a proof of concept, it was shown that T cells in splenocytes modified to display SA-mFasL did not generate a proliferative response when stimulated with T-cell mitogens (Fig.10). In contrast, unmodified splenocytes generated a vigorous proliferative response, demonstrating that T cells responding to mitogens expressed Fas and died upon the interaction of FasL with Fas on the same cell (suicide). It was further demonstrated that rats receiving bone marrow cells and mature lymphocytes developed GVHD within 17-20 days. Treatment of these animals with syngeneic splenocytes expressing FasL resulted in the prevention of GVHD, strong *in vivo* data that this approach is superior to other treatment regimens that are used in the clinic.

30 Example 13. Immune down-regulation using immunomodulatory molecules.

The methods of the present invention of expressing exogenous proteins on the cells, tissues, and organs are made particularly effective in down-

regulating the immune system for therapeutic purposes when several of these molecules are expressed simultaneously. Therefore, a combination of death ligands (SA-mFasL, -TNF α , -TRAIL and likes) and anti-inflammatory molecules (CSA-IL-4, -IL-10, -TGF- β and likes) are expressed on the surface of 5 cells, tissues, and organs as a therapeutic approach to prevent/treat autoimmune disease, foreign graft rejection, and other immune-based disorders. This method shows down-regulation of the immune system.

Example 14. Vaccination against tumors.

10 Tumors may evade the immune system by down-regulating the signals that provoke a T-cell response. An effective T-cell response requires three distinct signals: Signal 1,2, and 3. Signal 1 is generated by T-cell receptor interaction with the major histocompatibility complex (MHC) anti-peptide on antigen-presenting cells (APCs). Signal 2 is mediated by the engagements of 15 costimulatory molecules, such as B7/CD28 and CD40/CD40L, on T cells and APCs. Signal 3 is transduced via cytokines elaborated by T cells and APCs that have received both Signal 1 and 2. The transduction of these three signals drives T cells and APCs to proliferation and differentiation into effectors for the generation of a productive immune response. The lack of any of these signals 20 during T cell response to tumors may serve as one the most effective mechanisms by which tumors evade the immune system. The present invention provides methods and compositions to modify cell surface for the expression of exogenous proteins. This approach is aimed at converting any cells into professional antigen-presenting cells for the generation of an effective immune 25 response.

A chimeric protein composed of CSA and extracellular portion of CD40L with engineered synthetic residues for structural flexibility (CSA-CD40L) was produced in Drosophila production cells. This molecule was then used for expression on the surface of splenocytes. Almost all the splenocytes 30 were positive for CSA-CD40L as shown by an antibody against CD40L in flow cytometry (Fig. 11). Splenocytes expressing CD40L were much more effective than regular splenocytes at inducing antigen-specific responses when used as

APCs. This clearly demonstrates the feasibility of our approach to express exogenous proteins on the cell surface for therapeutic purposes.

5 This concept is tested in a lung carcinoma animal model. Two lung carcinoma lines of mouse origin, Lewis and Line 1 and the C58 tumor cells (rat lymphoma) will be used for the purpose of vaccination against tumors.

10 Chimeric proteins produced from nucleic acids comprising the nucleic acids of CSA fused to the nucleic acids of B7.1 (SEQ. ID No. 2), CD40L (SEQ. ID No. 6), or IL2 (SEQ. ID No. 4) will be used for display on the surface of these tumor cells and the cells will be used for vaccination of mice and rats against tumors. Briefly, tumor cells (2×10^7) will be biotinylated as above. After several washes with PBS, the biotinylated cells will be incubated with chimeric proteins as above to bind the chimeric protein to the biotinylated cells. The chimera-decorated cells will be irradiated at 5-10,000 cGy and injected into tumor bearing animals to serve as a vaccine. Tumor regression is monitored.

15 Based on the results as shown in the above examples, it is expected that this method will show up-regulation of the immune system.

It can be easily seen that other tumor systems and chimeric proteins can be identified, and the methods of this invention applied to provide other tumor vaccines and vaccines against infections.

20 This invention has been described in various preferred embodiments. Those skilled in the art will readily recognize that modifications, deviations or substitutions of the compounds or methods here disclosed may be made without departing from the spirit and scope of this invention. All such modifications, deviations and substitutions are considered to be within the scope of the claims

25 of this invention.

The claims defining the invention are as follows:

1. A method of attaching an agent to a cell surface, the method comprising contacting a biotinylated cell with a chimeric protein comprising (i) the agent and (ii) 5 core streptavidin, wherein the chimeric protein is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, and SEQ ID NO:7.
2. The method of claim 1, wherein the agent is FasL and the chimeric protein is 10 encoded by a nucleic acid comprising SEQ ID NO:1.
3. The method of claim 1, wherein the agent is B7 and the chimeric protein is encoded by a nucleic acid comprising SEQ ID NO:2.
4. The method of claim 1, wherein the agent is TGF- β and the chimeric protein is 15 encoded by a nucleic acid comprising SEQ ID NO:3.
5. The method of claim 1, wherein the agent is IL-2 and the chimeric protein is encoded by a nucleic acid comprising SEQ ID NO:4.
6. The method of claim 1, wherein the agent is IL-10 and the chimeric protein is 20 encoded by a nucleic acid comprising SEQ ID NO:5.
7. The method of claim 1, wherein the agent is CD40L and the chimeric protein is 25 encoded by a nucleic acid comprising SEQ ID NO:6.
8. The method of claim 1, wherein the agent is TNF- α and the chimeric protein is encoded by a nucleic acid comprising SEQ ID NO:7.
9. The method of claim 3, wherein the agent is B7.1. 30
10. The method of claim 1, wherein the cell is a target cell.

11. The method of claim 10, wherein the target cell is a tumor cell.
12. The method of claim 1, further comprising the step of biotinyling a cell to form the biotinylated cell.
5
13. The method of claim 12, wherein the cell is biotinylated at room temperature.
14. The method of claim 12, further comprising the step of washing the decorated cell, wherein, prior to the washing, the chimeric protein is contacted with the
10 biotinylated cell for a period lasting no more than about 45 minutes.
15. The method of claim 14, wherein the chimeric protein is contacted with the biotinylated cell for a period lasting no more than about 20 minutes.
- 15 16. The method of claim 14, wherein the chimeric protein is contacted with the biotinylated cell at a temperature not exceeding about 10 degrees Celsius.
17. The method of claim 12, wherein the biotinyling step comprises contacting the cell with a biotin-containing compound for a period lasting no more than about 30
20 minutes.
18. The method of claim 17, further comprising the step of washing the cell to remove free biotin-containing compound.
- 25 19. The method of claim 17, wherein the biotin-containing compound comprises Sulfo-NHS-LC-biotin.
20. The method of claim 1, wherein the contacting is effected *in vitro*.
- 30 21. The method of claim 1, wherein the contacting is effected *in vivo*.
22. The method of claim 1, wherein the method comprises administering the chimeric protein to a subject containing the biotinylated cell.

23. The method of claim 12, wherein the biotinylation is effected *in vitro*.
24. The method of claim 12, wherein the biotinylation is effected *in vivo*.
5
25. The method of claim 24, wherein the method comprises administering a biotin-containing compound to a subject containing the cell.
26. The method of claim 25, wherein the biotin-containing compound comprises
10 Sulfo-NHS-LC-biotin.
27. The method of claim 1, wherein the cell is selected from the group consisting of splenocytes, tumor cells, bone marrow cells, endothelial cells, and islet cells.
- 15 28. A nucleic acid comprising a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7.
29. A chimeric protein encoded by the nucleic acid of claim 28.
20
30. A cell comprising the chimeric protein of claim 29.
31. A production cell comprising the nucleic acid of claim 28.
- 25 32. A chimeric protein produced by the production cell of claim 31, wherein the chimeric protein is encoded by a nucleic acid comprising a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7.
- 30 33. A target cell comprising a chimeric protein encoded by the nucleic acid of claim 28, wherein the chimeric protein is displayed on the surface of the target cell.

34. A decorated surface comprising a chimeric protein encoded by the nucleic acid of claim 28.
35. A vaccine, wherein the vaccine comprises a tumor cell decorated with chimeric protein encoded by the nucleic acid of claim 28.
36. A method of attaching an immune costimulatory agent to a cell surface *in vivo*, the method comprising:
 - 10 contacting a biotinylated cell with a chimeric protein comprising (i) the immune costimulatory agent and (ii) core streptavidin, wherein the immune costimulatory agent is an adhesion molecule, a cell growth factor, a cell toxin or a composition that induces apoptosis, down-regulation of the immune system or up-regulation of the immune system.
- 15 37. The method of claim 36, wherein the immunomodulatory agent is a cytokine, IL-4, IL-10 TGF- β , B7, CD40L, IL-2, IL-12, IL-15, lymphotactin or a chemokine.
38. The method of claim 37, wherein the immunomodulatory agent is B7.
- 20 39. The method of claim 38, wherein the immunomodulatory agent is B7.1.
40. The method of claim 36, wherein the agent is FasL, mFasL, TRAIL, TNF- α , or TWEAK.
- 25 41. The method of claim 36, wherein the cell is a target cell.
42. The method of claim 41, wherein the target cell is a tumor cell.
43. The method of claim 36, further comprising the step of biotinyling a cell to 30 form the biotinylated cell.
44. The method of claim 43, wherein the cell is biotinylated at room temperature.

45. The method of claim 43, wherein the biotinylation step comprises contacting the cell with a biotin-containing compound for a period lasting no more than about 30 minutes.
- 5 46. The method of claim 45, further comprising the step of washing the cell to remove free biotin-containing compound.
- 10 47. The method of claim 45, wherein the biotin-containing compound comprises Sulfo-NHS-LC-biotin.
- 15 48. The method of claim 36, further comprising the step of administering the chimeric protein to a subject containing the biotinylated cell.
- 20 49. The method of claim 43, wherein the biotinylation is effected *in vivo*.
50. The method of claim 49, wherein the method comprises administering a biotin-containing compound to a subject containing the cell.
- 25 51. The method of claim 50, wherein the biotin-containing compound comprises Sulfo-NHS-LC-biotin.
52. The method of claim 43, wherein the biotinylation is effected *in vitro*.
- 25 53. The method of claim 36, wherein the chimeric protein is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, and SEQ ID NO:7.
- 30 54. A method of attaching an agent to a cell surface *in vivo*, the method comprising: contacting a biotinylated cell with a chimeric protein comprising (i) the agent and (ii) core streptavidin, the agent being an adhesion molecule, a cell growth factor, a cell toxin or a composition that induces apoptosis, down-regulation of the immune

system or up-regulation of the immune system, wherein, after five days *in vivo*, at least 26% of the cells retain the chimeric protein attached to the cell surface.

1/9




FIG. 1

2/9



FIG. 2

SUBSTITUTE SHEET (RULE 26)




FIG. 3

SUBSTITUTE SHEET (RULE 26)




FIG. 4

5/9




FIG. 5

SUBSTITUTE SHEET (RULE 26)




FIG. 6

7/9




FIG. 7

8/9




FIG. 8




FIG. 9




FIG. 10

SEQUENCE LISTING

<110> University of Louisville Research Foundation, Inc.

5 Shirwan, Haval

<120> Methods and Compositions for Altering Cell Membrane Function.

<130> 1160.015W01

10

<150> US 60/215,580

<151> 2000-06-30

<160> 7

15

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1247

20<212> DNA

<213> Artificial Sequence

<220>

<223> SA-FasL

25

<400> 1

cgAACGAAAG	ACCCGTGTGT	AAAGCCGCGT	TTCCAAAATG	TATAAAACCG	AGAGCATCTG	60
gccaatgtgc	atcagttgtg	gtcagcagca	aaatcaagtg	aatcatctca	gtgcaactaa	120
agggggggatc	cgatctcaat	atgaagttat	gcataattact	ggccgtcggt	gcctttgttg	180
30gcctctcgct	cgggagatct	catcatcacc	atcaccatat	caccggcacc	tggtacaacc	240
agctcggttc	gactttcatac	gtgaccggcg	gcgcgcacgg	cggccatgacc	ggAACCTACG	300
agtccggccgt	cggtcaacgccc	gagagccgtt	acgtccctgac	cggtcggtac	gacagcgccc	360
cggccacccga	cggcagcggc	accgcctcg	gttggacgggt	ggccttggaa	aataactacc	420
gcaacggccca	ctccgcgacc	acgtggagcg	gccagtacgt	cggcgccgccc	gaggcgagga	480
35tcaacaccca	gtggctgctg	acctccggca	ccaccgaggc	caacgcctgg	aagtccacgc	540
tggtcggcca	cgacacccctc	accaagggtga	agccgtccgc	cgcctcaagc	cgaattcctg	600
aaaccaaaaa	gccaaggagt	gtggcccact	taacaggaa	cccccgctca	aggccatcc	660
ctcttggaaatg	ggaagacaca	tatggaaactg	ctttgatctc	tggagtgaag	tataagaaag	720
gcggcccttgt	gatcaatgag	gctgggttgt	acttcgtata	ttccaaagta	tacttccggg	780
40gtcagtcttg	caacagccag	cccctaagcc	acaagggtcta	tatgaggaac	tttaagtatc	840
ctggggatct	ggtgctaatg	gaggagaaga	agttgaatta	ctgcactact	ggccagatata	900

ggggccacag cagctaccta gggcagtat ttaatcttac cgttgctgac catttatatg	960
tcaacatac tcaactctct ctgatcaatt ttgaggaatc taagaccttt tttggcttat	1020
ataagctt:a aaggaaaaag catttagaa tgatctatta ttcttatca tggatgccag	1080
gaatattgtc ttcaatgaga gtctcttaa gaccaattga gccacaaaga ccacaaggc	1140
5caacaggtaa gctaccctga attctgcaga tatccagcac agtggcggcc gctcgagtct	1200
agagggccct tcgaaggtaa gcctatccct aaccctctcc tcggtct	1247

<210> 2

<211> 1542

10<212> DNA

<213> Artificial Sequence

<220>

<223> B7.1-SA

15

<400> 2

cgaacgaaag acccggtgtt aaagccgcgt ttccaaaatg tataaaacccg agagcatctg	60
gccaatgtgc atcagttgtg gtcagcagca aatcaagtg aatcatctca gtgcaactaa	120
aggggggatc cgatctcaat atgaagttat gcatattact ggccgtcgtg gcctttgtt	180
20gcctctcgct cgggggtaag agatctatcc acgtgaccaa ggaagtgaaa gaagtggcaa	240
cgctgtcctg tggtcacaat gtttctgtt aagagctggc acaaactcgc atctactggc	300
aaaaggagaa gaaaatggtg ctgactatga tgtctggga catgaatata tggcccgagt	360
acaagaaccc gaccatctt gatatcacta ataacctctc cattgtgatc ctggctctgc	420
gcccatctga cgagggcaca tacgagtgtg ttgttctgaa gtatgaaaaa gacgccttca	480
25agcgggaaca cctggctgaa gtgacgttat cagtc当地 aacttccct acacctagta	540
tatctgactt taaaattcca acttctaata ttagaaggat aatttgc当地 acctctggag	600
gtttccaga gcctcaccc tcctgggtgg aaaaatggaga agaattaaat gccatcaaca	660
caacagttt ccaagatcc gaaactgagc tctatgctgt tagcagcaaa ctggattca	720
atatgacaac caaccacacg ttcatgtgtc tc当地caagta tggacattta agagtgaatc	780
30agaccttcaa ctggaaataca accaagcaag agagatctag atctcatcat caccatcacc	840
atatcaccgg cacctggtag aaccagctcg gctcgaccc catcgatcgc gcggggcgc	900
acggcgcctt gaccggaaacc tacgagtccg ccgtcggcaa cgccgagagc cgctacgtcc	960
tgaccggtag ttacgacagc gccccggcca cc当地acggcag cggcacccgc ctcgggttgg	1020
cggtggcctg gaagaataac taccgcaacg cccactccgc gaccacgtgg ageggccagt	1080
35acgtcggcgg cgccgaggcg aggatcaaca cccagtggt gctgacccctt ggcaccaccc	1140
aggccaaacgc ctggaaagtcc acgctggtag gccc当地acccac ct当地cccaag gtgaagccgt	1200
ccgccc当地cc cgaattctgc agatatccag cacagtgccg gccc当地ccgag tctagaggc	1260
ccttc当地ccagg taaggccatc cctaaaccctc tc当地ccgttct cgattctacg cgtaccggc	1320
atcatcacca tc当地ccattga gtttaaaccctt gctgatcagc ctc当地actgtg ccttctaaagg	1380
40cctgagctcg ctgatcagcc tc当地atcgagg atccagacat gataagatac attgatgagt	1440

ttggacaaac cacaactaga atgcagtgaa aaaaatgctt tatttgtgaa atttgtgatg	1500
ctattgcttt atttgtgacc attataagct gcaataaaca ag	1542

<210> 3

5<211> 1266

<212> DNA

<213> Artificial Sequence

<220>

10<223> TGF-B-SA

<400> 3

cgaacgaaag acccgtgtgt aaagccgcgt ttccaaaatg tataaaacccg agagcatctg	60
gc当地atgtgc atcagttgtg gtcagcagca aatcaagtg aatcatctca gtgcaactaa	120
15aggggggatc cgatctcaat atgaagttat gc当地attact ggccgtcggt gc当地ttgttg	180
gc当地tctcgct cggggtaag agatctgccc tggacaccaa ctattgttc agctccacgg	240
agaagaactg ctgcgtgcgg cagctgtaca tttgacttccg caaggacccg ggctggaagt	300
ggatccacga gcccaagggc taccatgcca acttctgcct cggccctgc cc当地acattt	360
ggagcctgga cacgcagtac agcaagggtcc tggccctgtt caaccagcat aacccggcg	420
20cctcggcggc gccgtgtgc gtgc当地cagg cgctggagcc gctgcccattc gtgtactacg	480
tggccgc当地 gcccaagggtg gagcagctgtt ccaacatgtat cgtgc当地tcc tgcaagtgca	540
gc当地atctca tcatcaccat caccatatac cccgc当地ctg gtacaaccag ct当地ggctcga	600
ccttcatcgt gaccgc当地ggc gccgacggcg cc当地tgc当地ccg aacctacgag tc当地ggc当地tgc当地	660
gcaacgccc当地 gagccgctac gtc当地tgc当地ccg gtc当地ttacga cagc当地ccccc gccaccgacg	720
25gc当地agc当地ggc当地 ac cccctcggt tggacggtgg cctggaagaa taactaccgc aacgc当地ccact	780
ccgc当地gaccac gtggagc当地ggc cagta cgtcg gc当地ggc当地ccg ggc当地gaggatc aacacc当地cagt	840
ggctgctgac ct当地ccggc当地cc accgaggccca acgc当地ctggaa gtccacgctg gtc当地ggcc当地acg	900
acacccattc当地ac caagggtgaag cc当地tccg当地ccg cctccgaaattt ct当地gc当地agatat cc当地agc当地acagt	960
ggc当地ggccgct cagatctaga gggcccttgc aaggtaagcc tatcccttaac ccttcttctcg	1020
30gtctcgatc tacgc当地tacc ggtcatc当地atc accatcacca ttgagttaa acccgctgat	1080
cagc当地ctcgac tttgc当地ttct aaggcctgag ctc当地gtgatc agc当地ctcgatc gaggatccag	1140
acatgataag atacattgtat gagttggac aaaccacaac tagaaatgc当地ag tggaaaaaaat	1200
gcttatttgc当地 tggaaatgtt gatgttatttgc当地 ct当地ttatttgc当地 aaccattata agctgcaata	1260
aacaag	1266

35

<210> 4

5<211> 1329

<212> DNA

<213> Artificial Sequence

<220>

<223> IL-2-SA

<400> 4

5cgaacgaaag acccggtgt	aaagccgcgt ttccaaaatg	tataaaacccg agagcatctg	60
gccaatgtgc atcagttgtg	gtcagcagca aaatcaagtg	aatcatctca gtgcaactaa	120
agggggatc cgatctcaat	atgaagttat	gcatttactt ggccgtcgtg	180
gcctctcgct cgggagatct	ccagcaccta	ttcaagtttca tacaagaaaa	240
aactggagca tttactgctg	gatttacaga	tgatggaa tgaaattaat	300
10atcccaaact caccaggatg	ctcacatttta	agttttacat gccaagaag	360
tgaaaatct tcagtgtcta	gaagaagaac	tcaaacctct ggaggaagtg	420
ctcaaagcaa aaactttcac	ttaagaccca	gggacttaat cagcaatatc	480
ttcttggact aaagggatct	gaaacaacat	tcatgtgtga atatgtat	540
ccatttgtaga atttctgaac	agatggatta	ccttttgtca aagcatcatc	600
15cttggtagatc tcatcatcac	catcaccata	tcacccggcac ctggtaaac	660
cgacccttcat cgtgaccgcg	ggccgcgacg	cgccctgtac cggaaacctac	720
tcggcaacgc cgagagccgc	tacgtcctga	ccggctgtta cgacagcgcc	780
acggcagcgg caccgcctc	ggttggacgg	ttggctggaa gaataactac	840
actccgcgac cacgtggacg	ggccagtacg	tcggccgcgc cgaggcgagg	900
20agtggctgct gaccccgcc	accacccgagg	ccaaacgcctg gaagtccacg	960
acgacaccc taccaagggt	aagccgtccg	ccgcctccga attctgcaga	1020
agtggccgc gctcgagtct	agagggccct	tcgaaggtaa gcctatccct	1080
tccgtctcgat ttctacgcgt	accgtcatc	atcaccatca ccattgagtt	1140
gatcagccctc gactgtgcct	tctaaggcct	gagctcgctg atcagcctcg	1200
25cagacatgat aagatacatt	gatgagtttgc	gacaaaccac aactagaatg	1260
aatgctttat ttgtgaaatt	tgtgatgcta	ttgctttatt tgtaaccatt	1320
ataaacaag			1329

<210> 5

30<211> 1395

<212> DNA

<213> Artificial Sequence

<220>

35<223> IL-10-SA

<400> 5

cgaacgaaag acccggtgt	aaagccgcgt ttccaaaatg	tataaaacccg agagcatctg	60
gccaatgtgc atcagttgtg	gtcagcagca aaatcaagtg	aatcatctca gtgcaactaa	120
40agggggatc cgatctcaat	atgaagttat	gcatttactt ggccgtcgtg	180

gcctctcgct	cgggggtaag	agatcttctg	agaacagctg	cacccacttc	ccaggcaacc	240
tgcctaacat	gttcgagat	ctccgagatg	ctttagcag	agtgaagact	ttctttcaaa	300
tgaaggatca	gctggacaac	ttgttgttaa	aggagtcctt	gctggaggac	tttaagggtt	360
acctgggtt	ccaagcctt	tctgagatga	tccagttta	cctggaggag	gtatgcccc	420
5aagctgagaa	ccaagaccca	gacatcaagg	cgcattgtgaa	ctccctgggg	gagaacctga	480
agaccctcg	gctgaggcta	cggcgtgtc	atcgatttct	tccctgtgaa	aacaagagca	540
aggccgtgga	gcaggtgaag	aatgcctta	ataagctcca	agagaaaggc	atctacaaag	600
ccatgagtga	gtttgacatc	ttcatcaact	acatagaagc	ctacatgaca	atgaagatac	660
gaaacagatc	tagatctcat	catcaccatc	accatatcac	cggcacctgg	tacaaccagc	720
10tcggctcgac	cttcatecg	accgcggcg	ccgacggcgc	cctgaccgga	acctacgagt	780
cggccgtcg	caacgcccag	agccgctacg	tcctgaccgg	tcgttacgac	agcgccccgg	840
ccaccgacgg	cagcggcacc	gccctcggtt	ggacggtgcc	ctggagaat	aactaccgca	900
acgcccactc	cgcgaccacg	tggagcggcc	agtaacgtcg	eggcgccgag	gogaggatca	960
acacccagt	gctgctgacc	tccggcacca	ccgaggccaa	cgcctggaaag	tccacgctgg	1020
15tcggccacga	cacccattcacc	aaggtgaagc	cgtccgcccgc	ctccgaattc	tgcagatatac	1080
cagcacagt	gcggccgctc	gagtcctagag	ggcccttcga	aggttaagcct	atccctaacc	1140
ctctccctcg	tctcgattct	acgcgtaccg	gtcatcatca	ccatcaccat	tgagttaaa	1200
cccgctgatc	agcctcgact	gtgccttcta	aggcctgagc	tcgctgatca	gcctcgatcg	1260
aggatccaga	catgataaga	tacattgtat	agtttggaca	aaccacaact	agaatgcagt	1320
20gaaaaaaaaatg	ctttatgtt	gaaatttgt	atgctattgc	tttatttgt	accattataa	1380
gctgcaataa	acaag					1395

<210> 6

<211> 1566

25<212> DNA

<213> Artificial Sequence

<220>

<223> SA-CD40L

30

<400> 6

cgaacgaaag	acccgtgtgt	aaagccgcgt	ttccaaaatg	tataaaacccg	agagcatctg	60
gccaatgtgc	atcagttgt	gtcagcagca	aatcaagt	aatcatctca	gtgcaactaa	120
aggggggatc	cgatctcaat	atgaagttat	gcataattact	ggccgtcg	gcctttgtt	180
35gcctctcgct	cgggggtaag	agatctcatc	atcaccatca	ccatatcacc	ggcacctgg	240
acaaccagct	cggctcgacc	ttcattcgat	ccgcgggcgc	cgacggcgcc	ctgaccggaa	300
cctacgagtc	ggccgtcg	aaccccggaa	gccgctacgt	cctgaccgg	cgttacgjaca	360
'gcgcggcgc	caccgacggc	agcggcaccg	ccctcggtt	gacggtgcc	tggagaata	420
actaccgcaa	cgcctcgact	cgaccacgt	ggagcggcc	gtacgtcg	ggcgccgagg	480
40cgaggatcaa	cacccagtgg	ctgctgac	ccggcaccac	cgaggccaac	gcctgaaagt	540

ccacgctgg	cgcccacgac	acttcacca	aggtaagcc	gtccggcgcc	tccgaattct	600
tggacaagat	agaagatgaa	aggaatctc	atgaagatti	tgtattcatg	aaaacgatac	660
agagatgcaa	cacaggagaa	agatccttat	cottactgaa	ctgtgaggag	attaaaagcc	720
agtttgaagg	ctttgtgaag	gatataatgt	taaacaaaga	ggagacgaag	aaagaaaaca	780
5gcttggaaat	gcaaaaaggt	gatcagaatc	ctcaaattgc	ggcacatgtc	ataagtgagg	840
ccagcagtaa	aacaacatct	gtgttacagt	gggctgaaaa	aggataactac	accatgagca	900
acaacttgg	aaccctggaa	aatgggaaac	agctgaccgt	taaaagacaa	ggactctatt	960
atatctatgc	ccaagtcacc	ttctgttcca	atcgggaagc	ttcgagtcaa	gctccattta	1020
tagccagct	ctgcctaaag	tccccggta	gatcgagag	aatcttactc	agagctgcaa	1080
10ataacccacag	ttccgccaaa	ccttgcgggc	aacaatccat	tcacttggga	ggagtatttg	1140
aattgcaacc	agggtgttcg	gtgttgtca	atgtgactga	tccaaagccaa	gtgagccatg	1200
gcactggctt	cacgtccctt	ggcttactca	aactcgaatt	ctgcagatat	ccagcacagt	1260
ggccggccgt	cgagtctaga	gggccttcg	aaggtaaagcc	tatccctaacc	cctctcttcg	1320
gtctcgattc	tacgcgtacc	ggtcatcatc	accatcacca	ttgagttaa	acccgctgat	1380
15cagcctcgac	tgtgccttct	aaggcctgag	ctcgctgatc	agcctcgatc	gaggatccag	1440
acatgataag	atacattgtat	gagtttggac	aaaccacaaac	tagaatgcag	tgaaaaaaaaat	1500
gctttatttg	tgaaatttgt	gatgtatttgc	ctttatgtt	aaccattata	agctgcaata	1560
aacaag						1566

20<210> 7

<211> 702

<212> DNA

<213> Artificial Sequence

25<220>

<223> SA-TNF α

<400> 7

ccagggctcc	aggcgggtgt	tgttcctcag	cctatttctcc	ttctgtatcg	tggcaggegc	60
30caccacgctc	ttctgcctgc	tgcactttgg	agtgtatggc	ccccagaggg	aagagtcccc	120
cagggacctc	tctctaattca	gccctctggc	ccaggcagtc	agatcatctt	ctcgaaccccc	180
gagtgacaag	cctgttagcc	atgtttagtc	aaaccctcaa	gctgagggggc	agctccagtg	240
gctgaaccgc	cgggccaatg	ccctcctggc	caatggcgtg	gagctgagag	ataaccagct	300
ggtgtgtcca	tcagaggggcc	tgtacctcat	ctactcccg	gtccttca	agggccaagg	360
35ctccccctcc	acccatgtgc	tcctcaccca	caccatcage	cgcatcgccg	tctcttacca	420
gaccaaggctc	aacctctct	ctgcctatcaa	gagccccctgc	cagagggaga	coccagaggg	480
ggctgaggcc	aagccctgg	atgagccat	ctatctggga	gggtcttcc	agctggagaa	540
gggtgaccga	ctcagcgctg	agataaatcg	gccccactat	ctcgactttg	cogagtctgg	600
gcaggtctac	tttggatca	ttgcctgtg	aggaggacga	acatccaacc	ttcccaaaccg	660
40cctccccctgc	cccaatccct	ttattacccc	ctccttcaga	ca		702