

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2018238996 B2**

(54) Title
Pharmaceutical composition for preventing and treating glaucoma, containing adenosine derivative

(51) International Patent Classification(s)
A61K 31/7076 (2006.01) **A61K 9/50** (2006.01)
A61K 9/00 (2006.01)

(21) Application No: **2018238996** (22) Date of Filing: **2018.03.21**

(87) WIPO No: **WO18/174549**

(30) Priority Data

(31) Number	(32) Date	(33) Country
10-2017-0035224	2017.03.21	KR

(43) Publication Date: **2018.09.27**

(44) Accepted Journal Date: **2021.01.28**

(71) Applicant(s)
Future Medicine Co., Ltd.

(72) Inventor(s)
Lee, Sang Koo;Park, Chong Woo;Kim, Hea Ok;Lee, Hee Woo;Yu, Mi Ra

(74) Agent / Attorney
FB Rice Pty Ltd, Level 23 44 Market Street, Sydney, NSW, 2000, AU

(56) Related Art
WO 2008045330 A2
WO 2012125400 A1
US 20100137577 A1

(12) 특허협력조약에 의하여 공개된 국제출원

(19) 세계지식재산권기구
국제사무국



(10) 국제공개번호

(43) 국제공개일
2018년 9월 27일 (27.09.2018) WIPO | PCT

WO 2018/174549 A1


- (51) 국제특허분류:
A61K 31/7076 (2006.01) A61K 9/50 (2006.01)
A61K 9/00 (2006.01)
- (21) 국제출원번호: PCT/KR2018/003274
- (22) 국제출원일: 2018년 3월 21일 (21.03.2018)
- (25) 출원언어: 한국어
- (26) 공개언어: 한국어
- (30) 우선권정보:
10-2017-0035224 2017년 3월 21일 (21.03.2017) KR
- (71) 출원인: 퓨처메디신 주식회사 (FUTURE MEDICINE CO., LTD.) [KR/KR]; 06665 서울시 서초구 효령로33길 55, 4층(방배동), Seoul (KR).
- (72) 발명자: 이상구 (LEE, Sang Koo); 08061 서울시 양천구 중앙로29길 61, 104동 501호, Seoul (KR). 박종우 (PARK, Chong Woo); 04354 서울시 용산구 새창로 70, 111동 1505호, Seoul (KR). 김혜옥 (KIM, Hea Ok); 06216 서울시 강남구 언주로 332, 107동 402호, Seoul (KR). 이희우 (LEE, Hee Woo); 08744 서울시 관악구 중앙2길 22, 405호, Seoul (KR). 유미라 (YU, Mi Ra); 08706 서울시 관악구 봉천로4길 56-2, 201호, Seoul (KR).
- (74) 대리인: 특허법인 태백 (TAEBAEK INTELLECTUAL PROPERTY LAW FIRM); 08506 서울시 금천구 가산디지털1로151 이노플렉스1차 601호, Seoul (KR).
- (81) 지정국 (별도의 표시가 없는 한, 가능한 모든 종류의 국내 권리의 보호를 위하여): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK,

MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) 지정국 (별도의 표시가 없는 한, 가능한 모든 종류의 역내 권리의 보호를 위하여): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), 유라시아 (AM, AZ, BY, KG, KZ, RU, TJ, TM), 유럽 (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

공개:


— 국제조사보고서와 함께 (조약 제21조(3))



WO 2018/174549 A1

(54) Title: PHARMACEUTICAL COMPOSITION FOR PREVENTING AND TREATING GLAUCOMA, CONTAINING ADE-NOSINE DERIVATIVE

(54) 발명의 명칭: 아데노신 유도체를 포함하는 녹내장 예방 및 치료용 약학적 조성물



(57) Abstract: Provided are a pharmaceutical composition for preventing or treating ocular diseases and an oral administration preparation for preventing or treating ocular diseases. The pharmaceutical composition for preventing or treating ocular diseases contains, as an active ingredient, a compound represented by chemical formula 1 or a pharmaceutically acceptable salt thereof.

(57) 요약서: 안질환 예방 또는 치료용 약학적 조성물과 안질환 예방 또는 치료용 경구 투여제가 제공된다. 안질환 예방 또는 치료용 약학적 조성물은 화학식 1로 표시되는 화합물 또는 이의 약학적으로 허용가능한 염을 유효성분으로 포함한다.

[DESCRIPTION]

[Invention Title]

PHARMACEUTICAL COMPOSITION FOR PREVENTING AND
TREATING GLAUCOMA, CONTAINING ADENOSINE DERIVATIVE

5 **[Technical Field]**

The present invention relates to a pharmaceutical composition comprising an adenosine derivative which can be effectively used for preventing or treating eye diseases and further for preventing or treating the elevation of the intraocular pressure and glaucoma.

10 **[Background Art]**

Adenosine is a substance which performs many physiological functions through specific cell membrane receptors and extracellular adenosine acts as a neurotransmitter in many physiological systems, and in general, it compensates for the hyperactivity of a given organ and protects from the harmful effects of stress (Jacobson, K. A. et al., J. Med. Chem., 35, 407-422, 1992). This effect is due to a partially generated negative feedback loop that reduces the energy requirement of the cell by adenosine produced by the degradation of intracellular or extracellular ATP (adenosine triphosphate) and increases the supply of oxygen. Adenosine is important for maintaining the homeostasis of essential organs such as the brain, heart, and kidney and for example, administration of adenosine agonist from the outside to the brain has been shown to have a neuroprotective action, and it is known that it is also involved in pain, cognition, exercise or sleep.

Adenosine receptors are now classified as P1 and P2 receptors,

respectively, by pharmacological studies and molecular cloning. The P1 receptor acts as a substrate for adenosine and the P2 receptor acts as a substrate for ATP, ADP, UTP and UDP to express physiological activity. Among these, as for the P1 receptors, four different subtypes of adenosine receptors have been identified and classified into A₁, A₂ or A₃ according to affinity to ligand, body distribution and action pathway, etc. and A₂ is again classified into A_{2A} and A_{2B}. These adenosine receptors are a class of G-protein-coupled receptor family, and the adenosine A₁, A_{2A} and A_{2B} receptors have been pharmacologically confirmed using many selective ligands, but the adenosine A₃ receptor was first identified in 1992 (Zhou, Q. Y, et al., Proc. Natl. Acad. Sci., USA, 89, 7432-7436, 1992), and many studies have been performed to confirm the pathophysiological function of this receptor.

Adenosine A₁ and A₂ receptor agonists are mainly derivatives of adenosine and have been extensively studied as antihypertensive agents, antipsychotic agents, antiarrhythmic agents, lipid metabolism inhibitors (diabetic agents), and brain protectants, and antagonists thereof are xanthine derivatives, or in which a plurality of bicyclic molecule is joined together and are developed as an asthma treatment agent, an antidepressant, an arrhythmia agent, a nephrotoxic agent, a Parkinson's disease treatment agent or a smart drug and the like. Nevertheless, currently commercialized is only adenosine itself which is in use for treating supraventricular tachycardia; and dipyridamole, an adenosine transport inhibitor which is in use as a warfarin adjuvant for preventing blood coagulation following cardiac surgery. The reasons why commercialization is not smooth that the adenosine receptor is spread throughout the body and is due to the various pharmacological actions involved in activating the receptor, i.e., there

is no compound that can activate only the adenosine receptor of the desired tissue.

Among the adenosine receptors, the adenosine A₃ receptor is the most recently recognized receptor, unlike the well-known adenosine A₁ and A₂ receptors, and its role is not well known and many studies are in progress to develop a selective receptor modulator. To investigate the adenosine A₃ receptor pharmacologically, three radiolabeled ligands such as [¹²⁵I]ABA(N⁶-(4-amino-3-[¹²⁵I]iodobenzyl)-adenosine, N⁶-(4-amino-3-[¹²⁵I]iodobenzyl)-adenosine), [¹²⁵I]APNEA(N⁶-2-(4-amino-3-[¹²⁵I]iodophenyl)-ethyladenosine, N⁶-2-(4-amino-3-[¹²⁵I]iodophenyl)-ethyladenosine) or [¹²⁵I]AB-MECA((N⁶-(4-amino-3-[¹²⁵I]iodobenzyl)-adenosine-5'-N-methylcarboxamide, N⁶-(4-amino-3-[¹²⁵I]iodobenzyl)-adenosine-5'-N-methylcarboxamide), have been used. It proved that when the adenosine A₃ receptor is expressed in Chinese hamster ovary (CHO) cells through pharmacological studies using the radiolabeled ligand, the A₃ receptor has function of inhibiting adenylyl cyclase, an enzyme that produces cAMP from ATP adenylyl cyclase, and when the A₃ receptor is activated by an agonist, it cleaves phosphatidyl inositol in the brain to activate GTP-dependent phospholipase C (Guanosine triphosphate-dependent phospholipase C), an enzyme producing inositol phosphate and DAG (Ramkumar, V. et al., J. Biol. Chem., 268, 168871-168890, 1993; Abbracchio, MP et al., Mol. Pharmacol., 48, 1038-1045, 1995). This finding can explain the possibility of a pathway of A₃ receptor activation by brain ischemia, since this secondary messenger system represents the response pathway of neuronal injury in cerebral ischemia. In addition, it is known that the A₃ receptor agonists inhibit the release of the tumor

necrosis factor TNF- α (tumor necrosis factor- α), which is an inflammatory mediator, and also inhibit the production of the inflammatory mediators such as MIP-1 α , interleukin-12 and interferon- γ , and it has a protective effect against the heart as well as a protective effect against brain diseases such as epilepsy. On
5 the other hand, inactivation of the adenosine A₃ receptor causes the release of inflammatory factors such as histamine from mast cells, contracts the bronchi and causes apoptosis in immune cells. Accordingly, adenosine A₃ antagonists have potential for development as anti-inflammatory agents and asthma treatment agents. Therefore, if a compound having pharmacological selectivity can be
10 developed, it would be possible to develop new therapeutic drugs for various diseases such as asthma, inflammation, cerebral ischemia, heart disease, cancer, etc.

Among the substances that have been researched and developed so far, representative human adenosine A₃ agonists include N⁶-(3-iodobenzyl)-5'-(N-
15 methylcarbamoyl)-adenosine (B-MECA) and N⁶-(3-iodobenzyl)-2-chloro-5'-(N-methylcarbamoyl)-adenosine (Cl-IB-MECA) and has a high affinity and selectivity for the A₃ receptor as compared to the adenosine A₁ and A₂ receptors. On the other hand, it has been pointed out that because the adenosine A₃ antagonist exhibiting high affinity and selectivity is mostly a nonpurine bicyclic compound,
20 not a nucleoside skeleton and exhibits high activity in human receptors, but weak or little activity in rat A₃ receptor, it is impossible to carry out animal experiments essential for the development of drugs which can be clinically applied (Baraldi, PG et al., *Curr. Med. Chem.*, 12, 1319-1329, 2005). However, because nucleoside-based compounds exhibit high affinity and selectivity regardless of

species, as compared with non-purine bicyclic compounds to have an advantage of easy experiment with animals, the possibility of development as a new drug is considered to be very high and therefore, it is an urgent task to elucidate the selective A₃ antagonist of this series.

5 In order to act as an agonist of the adenosine A₃ receptor through analysis of various previous studies, the present inventors have found that the N-methyl carbamoyl group at 5-position of the sugar in the structure of IB-MECA and CI-IB-MECA should be essentially present and that the base moiety should be substituted with an arylamino group or an alkylamino group at 6-position of purine. 10 Therefore, the N-methylcarbamoyl group at the 5-position of the sugar causes a conformational change essential for the agonist action of the receptor through hydrogen bonding (Kim, S-K. et al., J. Mol., 25, 562-577, 2006), it is considered that synthesis of a substance in which N-methylcarbamoyl group at the 5-position of the sugar is removed can be developed as an A₃ receptor antagonist.

15 Meanwhile, glaucoma, a type of eye disease, is a disease that can be induced by intraocular pressure elevation due to the problem of the production or release mechanism of aqueous humor. It is globally recognized as a serious disease because the intraocular pressure elevation can cause an abnormality in the function of the optic nerve due to disorder of blood supply and can lead to 20 blindness at the end.

Therefore, the present inventors have firstly studied an adenosine A₃ receptor antagonist as an agent for preventing and treating glaucoma which is caused by an elevation of the intraocular pressure and which may be accompanied by such symptoms, and synthesized novel adenosine derivative

compounds capable of effectively lowering intraocular pressure by inhibiting the production of aqueous humor in the ciliary body and remodeling the trabecular meshwork tissue by promoting anti-fibrosis.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority date of each of the appended claims.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

[Disclosure]

[Technical Problem]


A problem to be solved by the present invention is to provide a pharmaceutical composition comprising an adenosine derivative which acts as an adenosine A₃ receptor antagonist capable of preventing or treating eye diseases such as intraocular pressure elevation and glaucoma and the like.

The problems of the present invention are not limited to the above-mentioned technical problems and other technical problems which are not mentioned may be clearly understood by those skilled in the art from the description below.

[Summary]

One aspect of the invention there is provided an oral pharmaceutical composition when used for preventing or treating intraocular pressure elevation and/or glaucoma, which comprises a compound represented by following Chemical Formula A or a pharmaceutically acceptable salt:

[Chemical Formula A]




and a pharmaceutically acceptable excipient.

[Technical Solution]

In order to solve the above problems, a pharmaceutical composition for preventing or treating eye diseases according to an embodiment of the present invention comprises a compound represented by following Chemical Formula 1 or a pharmaceutically acceptable salt thereof as an active ingredient:

[Chemical Formula 1]



wherein A is S,


R is a linear or branched C₁-C₅ alkyl which is non-substituted or is independently or selectively substituted with one or more C₆-C₁₀ aryl groups, a benzyl which is non-substituted or is independently or selectively substituted with halogen and one or more linear or branched C₁-C₄ alkoxy groups, or a
5 hydroxycarbonyl-substituted benzyl, and

Y is H or a halogen element.

Also, the eye diseases may include at least one of intraocular pressure elevation and glaucoma.


In addition, the compound represented by the Chemical Formula 1 may
10 be a compound represented by the Chemical Formula A:

[Chemical Formula A]



In order to solve the above other problems, an oral administration agent for preventing or treating eye diseases according to an embodiment of the
15 present invention comprises a compound represented by following Chemical Formula 1 or a pharmaceutically acceptable salt:

[Chemical Formula 1]



wherein A is S,

R is a linear or branched C₁-C₅ alkyl which is non-substituted or is independently or selectively substituted with one or more C₆-C₁₀ aryl groups, a
 5 benzyl which is non-substituted or is independently or selectively substituted with halogen and one or more linear or branched C₁-C₄ alkoxy groups, or a hydroxycarbonyl-substituted benzyl, and

Y is H or a halogen element.

Also, the eye diseases may include at least one of intraocular pressure
 10 elevation and glaucoma.


In addition, the oral administration agent may further comprise an excipient comprising at least one selected from the group consisting of methyl cellulose (MC), dimethyl sulfoxide (DMSO), polyethylene glycol (PEG) and distilled water.

15 Furthermore, the excipient may comprise 0.5 wt% of methyl cellulose.

In addition, the compound represented by the Chemical Formula 1 or a pharmaceutically acceptable salt thereof may be filled in capsules in powder form.

In addition, the compound represented by the Chemical Formula 1 may be a compound represented by the Chemical Formula A:

20 [Chemical Formula A]



The details of other embodiments are included in the detailed description and drawings.

[Advantageous Effects]

5 An adenosine derivative of the present invention can act as an adenosine A_3 receptor antagonist capable of preventing or treating eye diseases such as the elevation of the intraocular pressure and glaucoma, and is thus suitable for prevention and treatment of eye diseases, and it can be used as a pharmaceutical composition which is very suitable for oral administration for preventing and
10 treating eye diseases because it is excellent in absorption of the drug when administered orally, biocompatible with little toxicity in the body, and superior in storage stability when formulated as an oral administration agent .

 Also, the adenosine derivatives of the present invention can be used as a pharmaceutical composition highly suitable for preventing and treating
15 glaucoma, because they have a dual pharmacological mechanism of inhibiting the production of an aqueous humor in the ciliary body and promoting the release of the aqueous humor through remodeling of trabecular meshwork tissue by the promotion of anti-fibrosis.

 The effects according to the embodiments of the present invention are not
20 limited by the contents exemplified above, and more various effects are included

in the present specification.

[Description of Drawings]

FIG. 1 shows the antagonistic effect of the compound of Example 4 of the present invention on Chinese hamster ovary (CHO) cells treated with CI-IB-
5 MECA as an agonist.

FIG. 2 shows the anti-inflammatory activity of the compounds of the present invention (Examples 2, 3 and 4) by the animal experiments.

FIG. 3 shows the anti-inflammatory activity of the compounds of the present invention (Examples 1 and 6) by the animal experiments.

10 FIG.4 shows the anti-inflammatory activity of the compounds of the present invention (Examples 5, 7, and 8) by the animal experiments.

FIG. 5 shows the anti-inflammatory activity of the compounds of the present invention (Examples 15 and 16) by the animal experiments.

15 FIG. 6 shows the results of measuring the intraocular pressure of a laboratory animal in Experimental Example 8.

FIG.7 shows the results of measuring the intraocular pressure of a laboratory animal in Experimental Example 9.

FIG. 8 to FIG. 12 are graphs showing the results of measuring the intraocular pressure of laboratory animals in Experimental Example 10.

20 FIG. 13 is a photograph showing changes in the expression of extracellular matrix proteins (Fibronectin, SMA and MYOC) in the trabecular meshwork tissue of laboratory animal eye in Experimental Example 11.

FIG.14 to FIG. 17 are graphs showing the measurement results of MMP and TIMP enzyme proteins and mRNA expression levels in the trabecular

meshwork tissue of laboratory animal eye in Experimental Example 12.

FIG.18 is a graph obtained from blood concentration-time data of Experimental Example 14 (14-1 and 14-2).

FIG. 19 is a graph obtained from the blood concentration-time data of Experimental Example 15 (15-1 and 15-2).

FIG. 20 is a graph obtained from the blood concentration-time data of Experimental Example 16.

FIG. 21 is a graph obtained from blood concentration-time data of Experimental Example 17 (17-1, 17-2 and 17-3).


FIG. 22 is a graph obtained from the blood concentration-time data of Experimental Example 18 (18-1 and 18-2).

[Best Mode]

Hereinafter, the present invention will be described in detail.

The present invention provides an adenosine derivative comprising a compound represented by the following Chemical Formula 1 or a pharmaceutically acceptable salt thereof, as an active ingredient:

[Chemical Formula 1]



wherein A is O or S,

R is a linear or branched C₁-C₅ alkyl which is non-substituted or is independently or selectively substituted with one or more C₆-C₁₀ aryl groups, a

benzyl which is non-substituted or is independently or selectively substituted with halogen and one or more linear or branched C₁-C₄ alkoxy groups, or a hydroxycarbonyl-substituted benzyl, and

Y is H or a halogen element.

5 Preferably, the A is O or S, the R is methyl, ethyl, propyl, naphthylmethyl, benzyl, benzyl substituted independently or optionally by one or at least two substituents selected from the group consisting of F, Cl, Br, I and C₁-C₃ alkoxy or toluic acid, and the Y is H or Cl.

10 More preferably, the A is O or S, the R is selected from the group consisting of methyl, ethyl, 1-naphthylmethyl, benzyl, 2-chlorobenzyl, 3-fluorobenzyl, 3-chlorobenzyl, 3-bromobenzyl, 3-iodobenzyl, 2-methoxy-5-chlorobenzyl, 2-methoxybenzyl or 3-toluol acid, and the Y is H or Cl.

Preferred examples of the adenosine derivatives represented by the above Chemical Formula 1 according to the present invention are as follows:

15 1) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-fluorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

2) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

20 3) (2*R*,3*R*,4*S*)-2-(6-(3-bromobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

4) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

5) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

6) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(5-chloro-2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol ;

7) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

5 8) (2*R*,3*R*,4*S*)-2-(2-chloro-6-(naphthalen-1-ylmethylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

9) 3-((2-chloro-9-((2*R*,3*R*,4*S*)-3,4-dihydroxytetrahydrothiophen-2-yl)-9*H*-purin-6-ylamino)methyl)benzoic acid;

10) 2-(2-chloro-6-methylamino-purin-9-yl)tetrahydrothiophene-3,4-diol;

10 11) (2*R*,3*R*,4*S*)-2-(6-(3-fluorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

12) (2*R*,3*R*,4*S*)-2-(6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

13) (2*R*,3*R*,4*S*)-2-(6-(3-bromobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

14) (2*R*,3*R*,4*S*)-2-(6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol;

15) (2*R*,3*R*,4*R*)-2-(6-(3-bromobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrofuran-3,4-diol; and

20 16) (2*R*,3*R*,4*R*)-2-(2-chloro-6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrofuran-3,4-diol.

The adenosine derivative represented by the Chemical Formula 1 according to the present invention may be used in the form of a pharmaceutically acceptable salt. As the salt, an acid addition salt formed by various

pharmaceutically acceptable organic acids or inorganic acids is useful. Suitable organic acids include, for example, organic acids such as carboxylic acid, phosphonic acid, sulfonic acid, acetic acid, propionic acid, octanoic acid, decanoic acid, glycolic acid, lactic acid, fumaric acid, succinic acid, adipic acid, malic acid, tartaric acid, citric acid, glutamic acid, aspartic acid, maleic acid, benzoic acid, salicylic acid, phthalic acid, phenylacetic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, methylsulfuric acid, ethylsulfuric acid and dodecylsulfuric acid, etc. and suitable inorganic acids include, for example, hydrogen acids such as hydrochloric acid, sulfuric acid or phosphoric acid.


10 The adenosine derivatives represented by Chemical Formula 1 according to the present invention may include not only pharmaceutically acceptable salts, but also all salts, hydrates and solvates which can be prepared by conventional methods.

 Also, the present invention provides a method of preparing the adenosine derivative represented by the above Chemical Formula 1.

 Specifically, as shown in Reaction Scheme 1 below, the present invention provides a method of preparing the adenosine derivative represented by the above Chemical Formula 1 comprising: obtaining β -anomer compound of Chemical Formula 3 by reacting a compound of Chemical Formula 2 as a starting material with a silylated purine compound in the presence of a Lewis acid catalyst (Step 1); obtaining a diol compound of Chemical Formula 4 by adding hydrochloric acid to the compound of the Chemical Formula 3 obtained in the step 1 (Step 2); and obtaining an adenosine derivative by reacting the diol compound of the Chemical Formula 4 obtained in the step 2 with an amine

compound under a base catalyst (Step 3).

[Reaction Scheme 1]



In the above Reaction Scheme 1, A, R and Y are as defined in Chemical Formula 1.

Hereinafter, the preparation method of the present invention will be described step by step.

Step 1 according to the present invention is a step of obtaining β -anomer compound of Chemical Formula 3 by reacting a compound of Chemical Formula 2 as a starting material with a silylated purine compound in the presence of a Lewis acid catalyst.

The compound of Chemical Formula 3 may be obtained by reacting the compound of Chemical Formula 2 with a silylated purine compound in the

presence of a Lewis acid. As the Lewis acid, trimethylsilyl trifluoromethanesulfonate (TMSOTf) may be used. In addition, as the solvent of the step 1, dichloroethane, chloroform, acetonitrile, dichloromethane and the like are preferably used. Among them, dichloroethane is more preferable. The
5 silylated purine compound can be obtained by reacting the purine compound of Chemical Formula 5 with hexamethyldisilazane (HMDS) and ammonium sulfate catalyst.

Step 2 according to the present invention is a step of obtaining a diol compound of Chemical Formula 4 by adding hydrochloric acid to the compound
10 of the Chemical Formula 3 obtained in the step 1. At this time, acetic acid, sulfuric acid, and p-toluenesulfonic acid may be used instead of hydrochloric acid.

Step 3 according to the present invention is a step of obtaining an adenosine derivative by reacting the diol compound of the Chemical Formula 4 obtained in the step 2 with an amine compound under a base catalyst.


15 As the base catalyst, triethylamine, pyridine, N, N-dimethylaminopyridine, 1,4-dioxane and the like are preferably used, and among them, triethylamine is more preferable. Further, as the solvent in the step 3, a solvent such as lower alcohol including methanol and ethanol or 1,4-dioxane, tetrahydrofuran and chloroform is preferable.

20 In the method for preparing an adenosine derivative of the present invention, the compound of Chemical Formula 2 as a starting material may be prepared according to the following Reaction Scheme 2 or 3 according to the type of substituent A.

When A is sulfur (S), as shown in Reaction Scheme 2 below, it comprises:

obtaining a diacetonide compound of Chemical Formula 7 by reacting D-mannose compound of Chemical Formula 6 with 2,2-dimethoxypropane under acid catalysis (Step a₁); obtaining a diol compound of Chemical Formula 8 by ring cleavage of the compound of Chemical Formula 7 obtained in the step a₁ in the presence of a reducing agent (Step a₂); obtaining a dimesyl compound of Chemical Formula 9 by mesylating the compound of Chemical Formula 8 obtained in the step a₂ (Step a₃); obtaining a thiosugar compound of Chemical Formula 10 by cyclizing the compound of Chemical Formula 9 obtained in the step a₃ (Step a₄); obtaining a diol compound of Chemical Formula 11 by selectively hydrolyzing the compound of Chemical Formula 10 obtained in the step a₄ (Step a₅); and obtaining an acetate compound of Chemical Formula 2a by reacting the compound of Chemical Formula 11 obtained in the step a₅ in the presence of a catalyst (Step a₆).

[Reaction Scheme 2]



(In the above Reaction Scheme 2, compound 2a is a compound of Chemical Formula 2.)

Hereinafter, the preparation method for the compound of Chemical Formula 2 of the present invention will be described in detail step by step.

Step a₁ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining a diacetonide compound of Chemical Formula 7 by reacting D-mannose compound of Chemical Formula 6 with 2,2-dimethoxypropane under acid catalysis.

The compound of Chemical Formula 7 can be obtained by reacting D-mannose of Chemical Formula 6 with 2,2-dimethoxypropane in the presence of an acid catalyst and acetic anhydride and as the acid catalyst, inorganic acids such as concentrated sulfuric acid or hydrochloric acid gas or organic acids such

as p-toluenesulfonic acid can be used.

Step a₂ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining a diol compound of Chemical Formula 8 by ring cleavage of the compound of Chemical Formula 7
5 obtained in the step a₁ in the presence of a reducing agent

The compound of Chemical Formula 8 can be obtained by reacting with a reducing agent, sodium borohydride. Instead of the sodium borohydride, a metal hydride such as lithium aluminum hydride, sodium sulfite or the like may be used.

10 Step a₃ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining a dimesyl compound of Chemical Formula 9 by mesylating the compound of Chemical Formula 8 obtained in the step a₂.

15 The compound of Chemical Formula 9 can be obtained by reacting the compound of Chemical Formula 8 with methanesulfonyl chloride (MsCl), and as the reaction solvent, ethyl ether, petroleum ether, dichloromethane, tetrahydrofuran and inert solvent such as N, N-dimethylformamide can be preferably used.

20 Step a₄ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining a thiosugar compound of Chemical Formula 10 by cyclizing the compound of Chemical Formula 9 obtained in the step a₃.

The compound of Chemical Formula 10 can be obtained by reacting the compound of Chemical Formula 9 with sodium sulfide and substitution reaction

with thioester such as methyl thioacetate in place of sodium sulfide, is followed by reaction with sodium alkoxide or the like. As the solvent in the step a₄, N,N-dimethylformamide, dimethylsulfoxide and the like can be used.

Step a₅ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining a diol compound of Chemical Formula 11 by selectively hydrolyzing the compound of Chemical Formula 10 obtained in the step a₄.


The compound of Chemical Formula 11 can be obtained by selectively hydrolyzing 5,6-acetonide by using acetic acid and sulfuric acid, hydrochloric acid, p-toluenesulfonic acid and the like may be used instead of acetic acid.

The step a₆ according to the method of preparing a compound of Chemical Formula 2 of the present invention is a step of obtaining an acetate compound of Chemical Formula 2a by reacting the compound of Chemical Formula 11 obtained in the step a₅ in the presence of a catalyst.

The compound of Chemical Formula 2a can be obtained by reacting the compound of Chemical Formula 11 with red tetraacetate (Pd(OAc)₄).

In addition, in the starting material 2 according to the present invention, when A is oxygen (O), as shown in the following Reaction Scheme 3, the compound of Chemical Formula 2, which is the starting material, is prepared by the method comprising: obtaining a lactol compound of Chemical Formula 13 by reacting a compound of Chemical Formula 12 with a reducing agent (Step b₁); and obtaining an acetate compound of Chemical Formula 2b by reacting the compound of Chemical Formula 13 obtained in the step b₁ with acetic anhydride.

[Reaction Scheme 3]



(In the above Reaction Scheme 3, compound 2b is a compound of Chemical Formula 2.)

Hereinafter, another method of preparing the compound of Chemical Formula 2 of the present invention will be described in detail step by step.

Step b_1 according to another method of preparing the compound of Chemical Formula 2 of the present invention is a step of obtaining a lactol compound of Chemical Formula 13 by reacting a compound of Chemical Formula 12 with a reducing agent.

The compound of Chemical Formula 13 can be obtained by reducing the compound of Chemical Formula 12 which is easily synthesized, using a diisobutylammonium hydride (DIBAL) catalyst.

Step b_2 according to another method of preparing the compound of Chemical Formula 2 of the present invention is a step of obtaining an acetate compound of Chemical Formula 2b by reacting the compound of Chemical Formula 13 obtained in the step b_1 with acetic anhydride.

The compound of Chemical Formula 2 can be obtained by reacting the lactol compound of Chemical Formula 13 with acetic anhydride.

In addition, the present invention provides an adenosine A_3 antagonist comprising an adenosine derivative represented by the Chemical Formula 1 or a pharmaceutically acceptable salt thereof, as an active ingredient.

Furthermore, the present invention provides a pharmaceutical composition for preventing and treating inflammatory diseases comprising an adenosine derivative represented by the Chemical Formula 1 or a pharmaceutically acceptable salt thereof, as an active ingredient.

5 When the adenosine A₃ receptor is expressed in Chinese hamster ovary (CHO) cells, the A₃ receptor has the inhibitory effect of adenylyl cyclase, an enzyme that produces cAMP from ATP, and when the A₃ receptor is activated by an agonist, It has been proved that phosphatidylinositol is degraded to activate GTP-dependent phospholipase C, an enzyme that generates inositol phosphate
10 and DAG (Ramkumar, V. et al., J. Biol. Chem., 268, 168871-168890, 1993; Abbacchio, MP et al., Mol. Pharmacol., 48, 1038-1045, 1995). Since this secondary messenger system represents neuronal injury response pathway in cerebral ischemia, such a discovery could explain the response pathway of A₃ receptor activation in cerebral ischemia. In addition, adenosine A₃ agonists inhibit
15 the release of tumor necrosis factor TNF- α , an inflammatory mediator, and the production of inflammatory mediators, MIP-1 α , interleukin-12 and interferon- γ , and has protective effect against heart in addition to brain diseases such as epilepsy. In addition, inactivation of the adenosine A₃ receptor causes the release of inflammation inducers such as histamine from mast cells, acts to contract the
20 bronchial tubes and induces the apoptosis in immune cells. Thus, adenosine A₃ antagonists have potential for development as anti-inflammatory drugs and asthma treatment agents.

In an experiment for measuring the receptor binding affinity so as to evaluate the binding affinity and selectivity of the adenosine derivative of the

present invention for the human adenosine receptor (hAR) (see Experimental Example 1), the adenosine derivative of the present invention showed high binding affinity for the human adenosine A₃ (hA₃ AR) receptor and low affinity for the adenosine A₁ and A_{2A} receptors, i.e. high selectivity. In particular, the compound of Example 12 of the present invention exhibited the highest affinity for the hA₃ receptor with a K_i value of 1.50 ± 0.40 nM and then the binding affinity is high in order of the compound of Example 2 (K_i = 1.66 ± 0.90 nM), the compound of Example 14 (K_i = 2.50 ± 1.00 nM), the compound of Example 10 (K_i = 3.69 ± 0.25 nM) and the compound of Example 4 (K_i = 4.16 ± 0.50 nM). The compound of Example 4 of the present invention also showed a high affinity (K_i = 3.89 ± 1.15 nM) for adenosine A₃ receptor in rats expressed in Chinese hamster ovary (CHO) cells. In addition, the compounds of Examples 15 and 16, which are adenosine derivatives having an oxonucleoside form of 4'-O, also showed high binding affinity and selectivity (see Table 1).

In addition, in experiments conducted to investigate the anti-inflammatory activity of the adenosine derivatives of the present invention (see Experimental Examples 3 to 6), it was found that the adenosine derivatives of the present invention have anti-inflammatory activity although this change is small compared to hydrocortisone used as a control group.

From the results of the anti-inflammatory activity after diluting the compounds of Examples 2 to 4 in acetone, it was confirmed that the treatment with compound 4 had an effect of reducing edema for the TPA-induced mouse ear edema (see FIG. 2). In addition, it was found that the measured anti-inflammatory activity of the compounds of Examples 1 and 6 of the present

invention after diluting in acetone and treating was 4 times or much higher than that of the compounds of Examples 2 to 4 (see FIG. 3).

The anti-inflammatory activity of the compounds of Examples 5 to 7 of the present invention after diluting to 0.5% in a mixed solvent of distilled water and acetone (1:4) showed inflammation inhibition rates of 17%, 34% and 53%, respectively (See FIG. 4). And the inflammation inhibition rates of the compounds of Examples 15 and 16 after diluting to 0.5% in a mixed solvent of dimethylsulfoxide (DMSO) and acetone (1:9) were 59% and 79%, respectively (See FIG. 5) and it was confirmed that the adenosine derivative compounds of the present invention had anti-inflammatory activity.

Therefore, the adenosine derivatives represented by the Chemical Formula 1 of the present invention show high binding affinity and selectivity for the adenosine A₃ receptor, and thus can be effectively used as an excellent adenosine A₃ antagonist. In addition, the adenosine derivatives of the present invention antagonize the adenosine A₃ receptor and exhibit anti-inflammatory activity, and thus can be used as agent for preventing and treating inflammatory diseases.

In addition, the inflammatory diseases according to the present invention include acute and chronic inflammatory diseases such as alternative inflammation, exudative inflammation, purulent inflammation, hemorrhagic inflammation or proliferative inflammation, etc.


The present invention provides a pharmaceutical composition for preventing or treating eye diseases, which comprises an adenosine derivative comprising a compound represented by the Chemical Formula 1 and/or a

pharmaceutically acceptable salt thereof as an active ingredient.

The eye diseases may include all diseases, disorders or symptoms related to the eye such as retinal disease, corneal disease, conjunctival disease, uveal disease, glaucoma, cataract, and the like, and particularly may include
5 intraocular pressure elevation and glaucoma which may be caused from the same or accompanied by such symptoms.

Preferable examples of the adenosine derivatives may be (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-chlorobenzylamino)-9*H*-purine-9-yl)tetrahydrothiophene-3,4-diol represented by the above Chemical Formula 1.

10 [Chemical Formula A]



The pharmaceutical composition for preventing and/or treating eye diseases of the present invention can be formulated as an eye drop or an oral administration agent.

15 The eye drops may comprise the compound represented by the Chemical Formula 1 and/or a pharmaceutically acceptable salt thereof and liquid for eye drops. The liquid for eye drops may include at least one selected from the group consisting of a solubilizer, a viscosity enhancer, an antioxidant, a preservative, and a buffer solution. In an exemplary embodiment, the liquid for eye drops may
20 be a buffer solution at pH 6.8 in which Cremophor EL, glycerin, citric acid and

methylparaben are dissolved or mixed.

The oral administration agent may be one in which the compound represented by Chemical Formula 1 and/or a pharmaceutically acceptable salt thereof is formulated into a solid preparation or a liquid preparation.

5 The solid preparations may be tablets, pills, powders, granules, capsules, and the like and the liquid preparations may be suspensions, solutions, emulsions, syrups and the like, but they are not limited thereto.

The oral administration agent may further comprise an excipient. The excipient may contain at least one selected from the group consisting of methyl
10 cellulose (MC), sucrose, lactose, dimethylsulfoxide (DMSO), polyethylene glycol (PEG), magnesium stearate, calcium carbonate, gelatin, talc, distilled water (DW), liquid paraffin, and the like, and preferably 0.5 wt% of methyl cellulose.

In an exemplary embodiment, the oral administration agent may be a capsule or the like filled with the compound represented by the Chemical Formula
15 1 and/or a pharmaceutically acceptable salt thereof in a powder state or a solution state dissolved in the above excipient. But it is not limited thereto.

The pharmaceutical composition for preventing and/or treating eye diseases of the present invention can be administered to a patient as an eye drop or an oral preparation through the above-mentioned eye drops or oral
20 administration. However, it is not limited thereto and may be administered by other parenteral administration methods such as intravenous administration and the like.

The adenosine derivatives of the present invention inhibit the production of aqueous humor in the ciliary body and at the same time inhibit the fibrosis in

the trabecular meshwork tissue (anti-fibrosis) to exhibit a dual mechanism that promotes the release of aqueous humor (see Examples 11 and 12).

Accordingly, since the intraocular pressure is effectively lowered in a dose-dependent manner (see Experimental Examples 8 to 10), it can be used as
5 a pharmaceutical composition which is highly suitable for preventing and/or treating eye diseases such as glaucoma.

In addition, since the adenosine derivative of the present invention exhibits physicochemical properties suitable for oral administration (see Experimental Example 13), excellent in blood concentration and stability at the
10 time of oral administration (see Examples 14 to 18), is a biocompatible substance with little toxicity in the body (See Experimental Example 19), and excellent in storage stability when formulated as an oral administration agent (see Experimental Example 20), it can be used as an very effective active ingredient for oral administration for preventing and/or treating eye diseases.

15
<Example>

Preparation of starting material

<Preparation Example 1> Preparation of (3a*R*,4*R*,6a*S*)-2,2-dimethyltetrahydrothieno[3,4-*d*] [1,3]dioxol-4-yl acetate

20 Step a₁. Preparation of (3a*R*,4*R*,6*R*,6a*R*)-6-(2,2-dimethyl-1,3-dioxolane-4-yl) -2,2-dimethyl-tetrahydrofuro[3,4- *d*] [1,3]dioxol-4-ol

D-mannose (1.74 g, 6.52 mmol) and 2,2-dimethoxypropane (2.45 ml, 19.55 mmol) were added to acetone (50 ml), stirred and cooled to 0 °C. Concentrated sulfuric acid (0.45 g, 1.96 mmol) was added dropwise to it. The

reaction mixture was stirred at room temperature for 24 hours. Triethylamine was added to the mixture for neutralization, and the mixture was concentrated under reduced pressure. The mixture obtained after concentration was subjected to silica gel column chromatography using a hexane: ethyl acetate mixed solvent
5 (1:1, v/v) as an eluent to obtain the target compound as a white solid (1.61 g, 95%).

mp 120.3-120.5 °C;

¹H-NMR (CDCl₃) δ 5.34(s, 1 H), 4.76-4.79(m, 1 H), 4.58(d, 1 H, *J* = 6.0 Hz), 4.34-4.39(m, 1 H), 4.15(dd, 1 H, *J* = 3.6, 7.2 Hz), 4.00-4.08(m, 2 H);

10 [α]²⁵_D 11.71(c 0.11, CH₂Cl₂);

FAB-MS *m/z* 261 [M + H]⁺.

Step a₂. Preparation of (1*R*)-(2,2-dimethyl-1,3-dioxolane-4-yl)((4*R*,5*S*)-5-(hydroxymethyl)-2,2-dimethyl-1,3-dioxolane-4-yl) methanol

15 To ethanol (25 ml), (3*aR*,4*R*,6*R*,6*aR*)-6-(2,2-dimethyl-1,3-dioxolane-4-yl)-2,2-tetrahydrofuro[3,4-*d*][1,3]dioxol-4-ol (1.50 g, 5.76 mmol) prepared in the step a₁ was carefully divided and added and cooled to 0 °C. Sodium borohydride (NaHB₄, 440 mg, 11.53 mmol) was added thereto and stirred at room temperature for 2 hours. The reaction mixture was neutralized with acetic acid and then
20 concentrated under reduced pressure. The mixture was extracted with ethyl acetate and water, and then the organic layer was dried over anhydrous magnesium sulfate (MgSO₄), filtered, and then concentrated under reduced pressure. The mixture obtained after concentration was subjected to silica gel

column chromatography using a hexane:ethyl acetate mixed solvent (1:1, v/v) as an eluent to obtain the target compound in the form of a syrup (1.38 g, 92%).

$^1\text{H-NMR}$ (CDCl_3) δ 4.33(dd, 1 H, $J = 1.6, 7.2$ Hz), 4.24-4.28(m, 1 H), 4.06-4.13(m, 2 H), 3.92-3.97(m, 1 H), 3.76-3.85(m, 2 H), 3.59-3.61(m, 1 H), 1.48(s, 3 H), 1.38(s, 3 H), 1.36(s, 3 H), 1.33(s, 3 H);

$[\alpha]_{\text{D}}^{25}$ -3.88 (c 0.44, CH_2Cl_2);

FAB-MS m/z 263 $[\text{M} + \text{H}]^+$.

Step a₃. Preparation of (1*R*)-(2,2-dimethyl-1,3-dioxolane-4-yl)((4*S*,5*S*)-2,2-dimethyl-5-((methylsulfonyloxy)methyl)-1,3-dioxolane-4-yl)methyl methanesulfonate

To a mixture of dichloromethane (300 ml) and triethylamine (163.75 ml, 1.17 mol), (1*R*)-(2,2-dimethyl-1,3-dioxolane-((4*R*,5*S*)-5-(hydroxymethyl)-2,2-dimethyl-1,3-dioxolan-4-yl)methanol (38.52 g, 146.85 mmol) prepared in the step a₂ and 4-dimethylaminopyridine (4-DMAP, 5.38 mg, 44.06 mmol) were added and mixed and cooled to 0 °C. Dimethanesulfonyl chloride (47.59 mL, 587.42 mmol) was added dropwise carefully thereto. After stirring for 1 hour at room temperature, the reaction mixture was extracted with dichloromethane and washed with saturated aqueous sodium bicarbonate (NaHCO_3). The organic layer was collected and dried over anhydrous magnesium sulfate (MgSO_4), filtered and then concentrated under reduced pressure. Dimesyl compound in the form of brown syrup obtained by concentration was subjected to silica gel column chromatography using a hexane:ethyl acetate mixed solvent (5:1, v/v) as an eluent to obtain the target compound in the form of a syrup (57.83 g, 94%).

¹H-NMR (CDCl₃) δ4.75(pseudo t, 1 H, *J* = 7.4 Hz), 4.33-4.45(m, 4 H), 4.06-4.20(m, 3 H), 3.12(s, 3 H), 3.07(s, 3 H), 1.51(s, 3 H), 1.43(s, 3 H), 1.37(s, 3 H), 1.33(s, 3 H);

[α]_D²⁵38.32 (*c* 0.29, CH₂Cl₂);

5 FAB-MS *m/z* 419 [M + H]⁺.

Step a₄. Preparation of (3a*R*,4*S*,6a*S*) -4-(2,2-dimethyl-1,3-dioxolane-4-yl) -2,2-dimethyltetrahydrothieno[3,4- *d*] [1,3]dioxol

(1*R*)-(2,2-dimethyl-1,3-dioxolane-4-yl)((4*S*,5*S*)-2,2-dimethyl-5-
10 (methylsulfonyloxy)methyl-1,3-dioxolan-4-yl)methylmethanesulfonate (993.80 g, 2.23 mmol) prepared in the step a₃ was dissolved into DMF (50 ml) and sodium sulfide (348.30 g, 4.46 mmol) was added thereto and the mixture was then refluxed and stirred at 80 °C overnight. After completion of the reaction, the solvent was removed under reduced pressure and the residue was extracted with
15 ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate (MgSO₄), filtered, and then concentrated under reduced pressure. The residue obtained after concentration was subjected to silica gel column chromatography using a hexane:ethyl acetate mixed solvent (8:1, v/v) as an eluent to obtain a target compound in the form of a syrup (453.0 mg, 78%).

20 ¹H-NMR (CDCl₃)δ4.92(dt, 1 H, *J* = 1.8, 5.6 Hz), 4.72(dd, 1 H, *J* = 2.0, 6.0 Hz), 4.26-4.30(m, 1 H), 4.04(s, 1 H), 3.79(t, 1 H, *J* = 3.8 Hz), 3.31-3.32(m, 1 H), 3.19(dd, 1 H, *J* = 5.4, 12.0 Hz), 2.84(dd, 1 H, *J* = 1.6, 12.0 Hz), 1.51(s, 3 H), 1.43(s, 3 H), 1.32(dd, 6 H, *J* = 8.4 Hz);

[α]_D²⁵-96.04 (*c* 0.20, CH₂Cl₂);

FAB-MS m/z 261 $[M + H]^+$.

Step a₅. Preparation of 1-((3*aR*,4*S*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl) ethane-1,2-diol

5 (3*aR*,4*S*,6*aS*)-4-(2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol (21.78 g, 83.66 mmol) prepared in the step a₄ was dissolved in a 60% aqueous acetic acid solution (250 ml) and stirred at room temperature for 2 hours. The reaction mixture was concentrated under reduced pressure and the resulting residue was subjected to silica gel column
10 chromatography using a hexane:ethyl acetate mixed solvent (1:2, v/v) as an eluent to obtain the target compound as a white solid (14.85 g, 81 %).

¹H-NMR (CDCl₃) δ 4.92(dt, 1 H, $J = 1.8, 5.6$ Hz), 4.72(dd, 1 H, $J = 2.0, 6.0$ Hz), 4.26-4.30(m, 1 H), 4.04(s, 1 H), 3.79(t, 1 H, $J = 3.8$ Hz), 3.31-3.32(m, 1 H), 3.19(dd, 1 H, $J = 5.4, 12.0$ Hz), 2.84(dd, 1 H, $J = 1.6, 12.0$ Hz), 1.51(s, 3 H),
15 1.43(s, 3 H), 1.32(dd, 6 H, $J = 8.4$ Hz);

$[\alpha]_D^{25} -96.04$ (c 0.20, CH₂Cl₂);

FAB-MS m/z 261 $[M + H]^+$.

Step a₆. Preparation of (3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl acetate
20

1-((3*aR*,4*S*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)ethane-1,2-diol (14.85 g, 67.41 mmol) prepared in the step a₅ was dissolved in ethylacetate (300 ml) and cooled to 0 °C. Red tetraacetate (Pb(OAc)₄, 157.31 g, 337.06 mmol) was added thereto and stirred at room temperature overnight. The

reaction mixture was filtered through celite and the filtrate was diluted with ethyl acetate. The organic layer was diluted with dichloromethane, washed with saturated aqueous sodium bicarbonate (NaHCO₃), dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue
5 obtained after concentration was subjected to silica gel column chromatography using a hexane:ethyl acetate mixed solvent (8:1, v/v) as an eluent to obtain the target compound in the form of a syrup (8.82 g, 60%).

¹H-NMR (CDCl₃) δ5.03(dd, 1 H, *J* = 5.6, 9.6 Hz), 4.79(dd, 1 H, *J* = 5.6, 8.8 Hz), 3.21-3.27(m, 2 H), 3.01(dt, 2 H, *J* = 0.8, 12.8 Hz), 2.05(s, 3 H), 1.50(s,
10 3 H), 1.31(s, 3 H);

[α]_D²⁵ -258.15 (*c* 0.18, CH₂Cl₂);

FAB-MS *m/z* 218 [M]⁺.

<Preparation Example 2> Preparation of (3a*S*,4*S*,6a*S*)-2,2-dimethyl-
15 tetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl acetate

Step b₁. Preparation of (3a*R*,4*R*,6a*R*)-2,2-dimethyl-tetrahydrofuro[3,4-
d][1,3]dioxol-4-ol

After dissolving 2,3-O-isopropylidene-D-erythronolactone (1.04 g, 6.42 mmol) in toluene (20 mL), 1 M diisobutylammonium hydride (DIBAL)/THF solution
20 was added at -78 °C. The reaction mixture was stirred at the same temperature for 30 minutes and then the reaction was terminated by adding methanol slowly. The suspension was filtered through celite, extracted with ethyl acetate and water, and then subjected to silica gel column chromatography using a hexane:ethyl acetate mixed solvent (3:1, v/v) as an eluent to obtain the target compound in the

form of a syrup (1.94 g, 96%).

$^1\text{H-NMR}$ (CDCl_3) δ 5.39(s, 1 H), 4.82(dd, 1 H, $J = 3.6, 6.0$ Hz), 4.55(d, 1 H, $J = 6.0$ Hz), 4.05(dd, 1 H, $J = 3.6, 10.2$ Hz), 4.00(d, 1 H, $J = 10.0$ Hz), 1.45(s, 3 H), 1.30(s, 3 H).

5

Step b₂. Preparation of (3a*S*,4*S*,6a*S*)-2,2-dimethyl-tetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl acetate

After dissolving the lactol compound (875.9 mg, 5.47 mmol) prepared in the step b₁ of Preparation Example 2 in pyridine (10 mL), acetic anhydride (0.67 mL, 6.56 mmol) was added at 0 °C. The reaction mixture was stirred at room temperature for 3 hours and then concentrated under reduced pressure. After concentration, the residue was extracted with ethyl acetate and water, and then the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography using a hexane:ethyl acetate mixed solvent (8:1, v/v) as an eluent to obtain the target compound in the form of a syrup (702.1 mg, 65%).

15

$^1\text{H-NMR}$ (CDCl_3) δ 6.16(s, 1 H), 4.86(dd, 1 H, $J = 3.6, 6.0$ Hz), 4.66(d, 1 H, $J = 6.0$ Hz), 4.12(d, 1 H, $J = 6.4$ Hz), 3.99(dd, 1 H, $J = 3.6, 10.8$ Hz), 2.05(s, 3 H), 1.48(s, 3 H), 1.33(s, 3 H).

20

<Example 1> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-fluorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

Step 1. Preparation of 2,6-dichloro-9-((3a*R*,4*R*,6a*S*)-2,2-dimethyltetrahydrothieno [3,4- *d*][1,3]dioxol-4-yl)-9*H*-purine

After dissolving 2,6-dichloropurine (2.29 g, 22.12 mmol) and ammonium sulfate (438 mg, 3.32 mmol) in hexamethyldisilazane (HMDS, 50 mL), it was refluxed overnight in an inert and dry condition. The reaction mixture was concentrated under reduced pressure and the resulting solid mixture was redissolved in cold 1,2-dichloroethane (20 mL). (3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl acetate (1.41 g, 11.06 mmol) obtained in Preparation Example 1 was dissolved in 1,2-dichloroethane (20 ml) and added dropwise again to the above solution. Trimethylsilyltrifluoromethanesulfonate (TMSOTf, 4.0 ml, 22.12 mmol) was added dropwise to the mixture and stirred at 0 °C for 30 minutes, then at room temperature for 1 hour and then heated to 80 °C and stirred for 2 hours. The reaction mixture was cooled, diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate (NaHCO₃). The organic layer was dried over anhydrous magnesium sulfate (MgSO₄) and concentrated under reduced pressure to obtain a yellow syrupy residue. The residue was subjected to silica gel column chromatography using a mixed solvent of dichloromethane and methanol (50:1, v/v) as an eluent to obtain the compound in the form of foam (3.03 g, 79%).

UV (CH₂Cl₂)λ_{max} 275.0 nm;

¹H-NMR (CDCl₃) δ8.17(s, 1 H), 5.87(s, 1 H), 5.32(pseudo t, 1 H, *J* = 4.8 Hz), 5.21(d, 1 H, *J* = 5.6 Hz), 3.79(dd, 1 H, *J* = 4.4, 12.8 Hz), 3.26(d, 1 H, *J* = 13.2 Hz), 1.59(s, 3 H), 1.36(s, 3 H);

[α]_D²⁵-42.04 (c 0.16, CH₂Cl₂);

FAB-MS *m/z* 347 [M + H]⁺.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

After dissolving 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine prepared in the step 1 in tetrahydrofuran (20 ml), 2 N hydrochloric acid was added and stirred overnight at room temperature. The mixture was neutralized with 1 N aqueous sodium hydroxide solution and then carefully concentrated under reduced pressure. The residue after concentration was subjected to silica gel column chromatography using a dichloromethane:methanol mixed solvent (20:1, v/v) as an eluent to obtain the target compound as a white solid (1.94 g, 96%).

mp 198.3-200.3 °C;

UV (MeOH) λ_{\max} 275.0 nm;

¹H-NMR (CD₃OD) δ 8.87(s, 1 H), 6.08(d, 1 H, *J* = 6.8 Hz), 4.69(q, 1 H, *J* = 3.2 Hz), 4.48(q, 1 H, *J* = 3.6 Hz), 3.56(dd, 1 H, *J* = 4.4, 11.2 Hz), 2.97(dd, 1 H, *J* = 3.4, 11.2 Hz);

$[\alpha]_{\text{D}}^{25}$ -50.43 (*c* 0.12, DMSO);

FAB-MS *m/z* 307 [M + H]⁺.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-fluorobenzylamino)-9*H*-purin-9-yl) tetrahydrothiophene-3,4-diol

After dissolving (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol (1 eq.) prepared in the step 2 and 3-fluorobenzylamine (1.5 eq.) in ethanol (5 ml) at room temperature, the reaction

mixture was stirred at room temperature for 2-3 h. After completion of the reaction, the residue obtained by concentration under reduced pressure was subjected to silica gel column chromatography using a dichloromethane:methanol mixed solvent (20:1, v/v) as an eluent to obtain the target compound (0.10 g, 80%).

5 mp 183.2-183.5 °C;

UV (MeOH) λ_{\max} 275.0 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.91(t, 1 H-NH, $J = 5.8$ Hz), 8.51(s, 1 H), 7.33-7.39(m, 1 H), 7.13-7.18(m, 2 H), 7.06(dt, 1 H, $J = 2.8, 11.6$ Hz), 5.82(d, 1 H, $J = 7.2$ Hz), 5.56(d, 1 H-OH, $J = 6.0$ Hz), 5.37(d, 1 H-OH, $J = 4.4$ Hz), 4.65(d, 1 H, $J = 6.0$ Hz), 4.60(m, 1 H), 4.33-4.35(m, 1 H), 3.41(dd, 1 H, $J = 4.0, 10.8$ Hz), 2.79(dd, 1 H, $J = 2.8, 10.8$ Hz);

$[\alpha]_{\text{D}}^{25}$ -96.21 (c 0.12, DMSO);

FAB-MS m/z 396 $[\text{M} + \text{H}]^+$.

15 <Example 2> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 using 3-chlorobenzylamine instead of 3-fluorobenzylamine to
5 obtain the target compound (0.11 g, 83%).

mp 163.3-165.3 °C;

UV (MeOH) λ_{\max} 274.5 nm;

¹H-NMR (CD₃OD) δ 8.34(s, 1 H), 7.41(s, 1 H), 7.24-7.34(m, 3 H), 5.94(d, 1 H, *J* = 6.4 Hz), 4.75(brs, 2 H), 4.61(q, 1 H, *J* = 3.2 Hz), 4.45(q, 1 H, *J* = 4.0 Hz),
10 3.51(dd, 1 H, *J* = 4.8, 11.2 Hz), 2.95(dd, 1 H, *J* = 3.6, 10.8 Hz);

FAB-MS *m/z* 411 [M]⁺.

<Example 3> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-bromobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

15 Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno [3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

20 Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-bromobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of Example 1 using 3-bromobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.12 g, 83%).

mp 184.0-185.0 °C;

5 UV (MeOH) λ_{\max} 274.0 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.91(brs, 1 H-NH), 8.51(s, 1 H), 7.55(s, 1 H), 7.43(d, 1 H, $J = 7.6$ Hz), 7.33-7.35(m, 1 H), 7.26-7.30(m, 1 H), 5.82(d, 1 H, $J = 7.2$ Hz), 5.57(d, 1 H-OH, $J = 6.0$ Hz), 5.38(d, 1 H-OH, $J = 4.0$ Hz), 4.60-4.63(m, 3 H), 4.34(s, 1 H), 3.41(dd, 1 H, $J = 4.4, 11.2$ Hz), 2.80(dd, 1 H, $J = 2.8, 10.8$ Hz);

10 FAB-MS m/z 456 $[\text{M} + \text{H}]^+$.

<Example 4> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

15 Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

20 The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3

of Example 1 by using 3-iodobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.14 g, 84%).

mp 198.7-199.9 °C;

UV (MeOH) λ_{\max} 274.0 nm;

5 $^1\text{H-NMR}$ (DMSO- d_6) δ 8.90(t, 1 H-NH, J = 6.4 Hz), 8.51(s, 1 H), 7.74(s, 1 H), 7.60(d, 1 H, J = 7.6 Hz), 7.35(d, 1 H, J = 7.6 Hz), 7.13(t, 1 H, J = 8.0 Hz), 5.82(d, 1 H, J = 7.6 Hz), 5.56(d, 1 H, J = 6.4 Hz), 5.37(d, 1 H, J = 4.0 Hz), 4.60(d, 3 H, J = 4.4 Hz), 4.34(brs, 1 H), 3.38(dd, 1 H, J = 4.0, 10.8 Hz), 2.80(dd, 1 H, J = 4.0, 10.8 Hz);

10 $[\alpha]_D^{25}$ -78.91 (c 0.13, DMSO);

FAB-MS m/z 504 $[\text{M} + \text{H}]^+$.

<Example 5> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-chlorobenzylamino)-9*H*-purin-9-yl) tetrahydrothiophene-3,4-diol

15 Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

20 Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl) tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-chlorobenzylamino)-9*H*- purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 by using 2-chlorobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.11 g, 81%).

mp 198.7-199.7 °C;

5 UV (MeOH) λ_{\max} 273.5 nm;

$^1\text{H-NMR}$ (CD_3OD) δ 8.35(brs, 1 H), 7.45-7.47(m, 1 H), 7.39-7.43(m, 1 H), 7.25-7.29(m, 2 H), 5.95(d, 1 H, $J = 6.4$ Hz), 4.60-4.63(m, 1 H), 4.45(dd, 1 H, $J = 3.6, 8.0$ Hz), 3.51(dd, 1 H, $J = 4.8, 10.8$ Hz), 2.95(dd, 1 H, $J = 4.0, 10.8$ Hz);

$[\alpha]_{\text{D}}^{25}$ -96.21 (c 0.12, DMSO);

10 FAB-MS m/z 412 $[\text{M} + \text{H}]^+$.

<Example 6> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(5-chloro-2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

20 The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(5-chloro-2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was carried out under the same conditions as in the step 3

of the Example 1 by using 5-chloro-2-methoxybenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.11 g, 78%).

mp 188.8-189.8 °C;

UV (MeOH) λ_{\max} 275.5 nm;

5 $^1\text{H-NMR}$ (DMSO- d_6) δ 8.64(t, 1 H-NH, J = 6.0 Hz), 8.51(s, 1 H), 7.21-7.25(m, 1 H), 7.12(d, 1 H, J = 7.2 Hz), 7.00(d, 1 H, J = 8.0 Hz), 6.85-6.89(m, 1 H), 5.82(d, 1 H, J = 7.6 Hz), 5.57(d, 1 H-OH, J = 6.4 Hz), 5.37(d, 1 H-OH, J = 4.0 Hz), 4.61-4.63(m, 2 H), 4.35(m, 1 H), 3.84(s, 3 H), 3.71(dd, 1 H, J = 3.6, 10.4 Hz), 2.80(dd, 1 H, J = 2,4, 10.8 Hz);

10 $[\alpha]_{\text{D}}^{25}$ -96.10 (c 0.21, DMSO);

FAB-MS m/z 442 $[\text{M} + \text{H}]^+$.

<Example 7> Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

15 Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

20 Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(2-chloro-6-(2-methoxybenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 by using 2-methoxybenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.12 g, 88%).

mp 188.0 °C;

5 UV (MeOH) λ_{\max} 276.5 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.65(t, 1 H-NH, J = 6.0 Hz), 8.51(s, 1 H), 7.21-7.25(m, 1 H), 7.12(d, 1 H, J = 7.2 Hz), 7.00(d, 1 H, J = 8.0 Hz), 6.85-6.89(m, 1 H), 5.83(d, 1 H, J = 6.8 Hz), 5.58(d, 1 H-OH, J = 6.4 Hz), 5.39(d, 1 H-OH, J = 3.6 Hz), 4.62-4.64(m, 2 H), 4.35(s, 1 H), 3.84(s, 1 H), 3.42(dd, 1 H, J = 3.6, 10.4 Hz),
10 2.79-2.82(m, 1 H);

$[\alpha]_{\text{D}}^{25}$ -93.53 (c 0.17, DMSO);

FAB-MS m/z 407 $[\text{M} + \text{H}]^+$.

<Example 8> Preparation of (2R,3R,4S)-2-(2-chloro-6-(naphthalen-1-ylmethylbenzylamino)-9H-purin-9-yl)tetrahydrothiophene-3,4-diol
15

Step 1. Preparation of 2,6-dichloro-9-((3aR,4R,6aS)-2,2-dimethyltetrahydrothieno[3,4-d][1,3]dioxol-4-yl)-9H-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

20 Step 2. Preparation of (2R,3S,4S)-2-(2,6-dichloro-9H-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of (2R,3R,4S)-2-(2-chloro-6-(naphthalen-1-

ylmethylbenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 by using naphthalene-1-ylmethylbenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.13 g, 90%).

5 mp 226.3 °C (decomp);

UV (MeOH) λ_{\max} 281.0 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.96(t, 1 H-NH, $J = 6.0$ Hz), 8.51(s, 1 H), 8.25(d, 1 H, $J = 8.0$ Hz), 7.95-7.97(m, 1 H), 7.83-7.85(m, 1 H), 7.53-7.61(m, 2 H), 7.43-7.46(m, 2 H), 5.82(d, 1 H, $J = 7.6$ Hz), 5.56(d, 1 H, $J = 6.4$ Hz), 5.38(d, 1 H, $J =$
10 4.0 Hz), 5.12(d, 1 H, $J = 6.0$ Hz), 4.59-4.61(m, 1 H), 4.34-4.35(m, 1 H), 3.40-3.44(m, 1 H), 2.80(dd, 1 H, $J = 2.4, 6.8$ Hz);

FAB-MS m/z 428 $[\text{M} + \text{H}]^+$.

<Example 9> Preparation of 3-((2-chloro-9-((2*R*,3*S*,4*R*)-3,4-
15 dihydroxytetrahydrothiophen-2-yl)-9*H*-purin-6-ylamino)methyl)benzoic acid

Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

20 Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3.Preparation of 3-((2-chloro-9-((2*R*,3*S*,4*R*)-3,4-

dihydroxytetrahydrothiophen-2-yl)-9H-purin-6-ylamino)methyl)benzoic acid

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 by using 3-(aminomethyl)benzoic acid instead of 3-fluorobenzylamine to obtain the target compound (0.12 g, 84%).

5 mp 254.0-256.9 °C;

UV (MeOH) λ_{\max} 275.5 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.95(t, 1 H-NH, $J = 6.0$ Hz), 8.52(s, 1 H), 7.89(d, 1 H, $J = 8.4$ Hz), 7.43(d, 1 H, $J = 8.0$ Hz), 5.82(d, 1 H, $J = 7.6$ Hz), 5.57(brs, 1 H), 5.38(brs, 1 H), 4.71(d, 1 H, $J = 6.0$ Hz), 4.60(brs, 1 H), 4.34(brs, 1 H), 3.41(dd, 1 H, $J = 4.0, 10.8$ Hz), 2.80(dd, 1 H, $J = 2.8, 10.8$ Hz);

$[\alpha]_{\text{D}}^{25}$ -94.55 (c 0.11, DMSO);

FAB-MS m/z 422 $[\text{M} + \text{H}]^+$.

<Example 10> Preparation of 2-(2-chloro-6-methylamino-purin-9-yl)(2R,3S,4R)-tetrahydrothiophene-3,4-diol

Step 1. Preparation of 2,6-dichloro-9-((3aR,4R,6aS)-2,2-dimethyltetrahydrothieno[3,4-d][1,3]dioxol-4-yl)-9H-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 1.

Step 2. Preparation of (2R,3S,4S)-2-(2,6-dichloro-9H-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 1.

Step 3. Preparation of 2-(2-chloro-6-methylamino-purin-9-yl)(2R,3S,4R)

-tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 1 by using methylamine instead of 3-fluorobenzylamine to obtain the target compound (0.89 g, 90%).

5 UV (MeOH) λ_{\max} 269.5 nm(pH 7);

$^1\text{H-NMR}$ (CDCl_3) δ 2.99(1H, dd, 4'-CH, $J = 4.4, 10.8$ Hz), 3.12(3H, brs, NH-CH₃), 3.44 1H, dd, 4'-CH, $J = 4, 10.8$ Hz), 4.41(1H, m, 2'-CH, $J = 5.6$ Hz), 4.47(1H, m, 3'-CH), 5.89(1H, d, 1'-CH, $J = 5.6$ Hz), 8.40(s, 1H, 8-CH);

$[\alpha]_{\text{D}}^{25}$ -34.8 (c 0.115, DMSO);

10 FAB-MS m/z 302.3 $[\text{M} + \text{H}]^+$.

<Example 11> Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-fluorobenzylamino)-9*H*-purin-9-yl) tetrahydrothiophene-3,4-diol

Step 1. Preparation of 6-chloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno [3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The synthesis was conducted under the same conditions as in the step 1 of the Example 1 by using 6-chloropurine (2.29 g, 22.12 mmol) instead of 2,6-chloropurine to obtain the target compound in the form of foam (1.84 g, 91%) .

UV (CH_2Cl_2) λ_{\max} 265.0 nm;

20 $^1\text{H-NMR}$ (CDCl_3) δ 8.67(pseudo t, 1 H, $J = 1.4$ Hz), 8.23(s, 1 H), 5.88(s, 1 H), 5.23(m, 2 H,),3.69(dd, 1 H, $J = 4.0, 13.2$ Hz), 3.18(d, 1 H, $J = 12.8$ Hz), 1.52(s, 3 H), 1.29(s, 3 H);

$^{13}\text{C-NMR}$ (CDCl_3) δ 152.05, 151.39, 151.09, 144.34, 132.56, 111.90,

89.60, 84.31, 70.30, 40.76, 26.40, 24.63;

$[\alpha]_{\text{D}}^{25}$ -157.64 (*c* 0.15, MeOH);

FAB-MS *m/z* 313 $[M + H]^+$.

5 Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(6-chloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted by using 6-chloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine (1.84 g, 5.88 mmol) prepared in the step 1 under the same conditions as in the step 2 of the Example
10 1 to obtain the target compound as a white solid (1.27 g, 79%).

mp 192.3-192.8 °C;

UV (MeOH) λ_{max} 264.5 nm;

¹H-NMR (DMSO-*d*₆) δ 9.02(s, 1 H), 8.82(s, 1 H), 6.02(d, 1 H, *J* = 7.6 Hz),
5.62(d, 1 H-OH, *J* = 6.0 Hz), 5.43(d, 1 H-OH, *J* = 4.0 Hz), 4.70-4.74(m, 1 H), 4.36-
15 4.40(m, 1 H), 3.47(dd, 1 H, *J* = 4.0, 10.8 Hz), 3.17(d, 1 H, *J* = 5.2 Hz), 2.84(dd, 1
H, *J* = 2.8, 11.2 Hz);

$[\alpha]_{\text{D}}^{25}$ -109.15 (*c* 0.16, DMSO);

FAB-MS *m/z* 273 $[M + H]^+$.

20 Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-fluorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

After dissolving (2*R*,3*S*,4*S*)-2-(6-chloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol (1 eq.) prepared in the step 2 and 3-fluorobenzylamine (1.5 eq.) in ethanol (5 ml) at room temperature, the reaction

mixture was stirred at room temperature for 2 to 3 hours. After completion of the reaction, the reaction mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography using a dichloromethane:methanol mixed solvent (20:1, v/v) as an eluent to obtain the
5 target compound (0.11 g, 82%).

mp 180.5-180.7 °C;

UV (MeOH) λ_{\max} 273.5 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.46(s, 1 H), 8.22(s, 1 H), 7.31-7.39(m, 1 H), 7.12-7.18(m, 2 H), 7.01-7.05(m, 1 H), 5.90(d, 1 H, $J = 7.2$ Hz), 5.53(d, 1 H-OH, $J = 6.4$
10 Hz), 5.35(d, 1 H-OH, $J = 4.0$ Hz), 4.67-4.71(m, 2 H), 4.35-4.37(m, 1 H), 3.39-3.43(m, 1 H), 3.17(d, 1 H, $J = 5.2$ Hz), 2.80(dd, 1 H, $J = 3.2, 11.2$ Hz);

$[\alpha]_{\text{D}}^{25}$ -141.2 (c 0.11, DMSO);

FAB-MS m/z 362 $[\text{M} + \text{H}]^+$.

15 <Example 12> Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

Step 1. Preparation of 6-chloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same
20 manner as in the step 1 of the Example 11.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(6-chloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 11.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-chlorobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3 of the Example 11 by using 3-chlorobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.12 g, 85%).

mp 165.0-165.3 °C;

UV (MeOH) λ_{\max} 274.5 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.47(s, 1 H), 8.22(s, 1 H), 7.39(s, 1 H), 7.26-7.35(m, 3 H), 5.91(d, 1 H, $J = 7.2$ Hz), 5.53(d, 1 H-OH, $J = 6.4$ Hz), 5.35(d, 1 H-OH, $J = 4.0$ Hz), 4.67-4.71(m, 2 H), 4.33-4.37(m, 1 H), 3.40-3.48(m, 2 H), 2.80(dd, 1 H, $J = 3.2, 10.4$ Hz);

$[\alpha]_D^{25}$ -162.5 (c 0.10, DMSO);

FAB-MS m/z 378 $[M + H]^+$.

<Example 13> Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-bromobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

Step 1. Preparation of 6-chloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrothieno[3,4- d][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same manner as in the step 1 of the Example 11.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(6-chloro-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner

as in the step 2 of the Example 11.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-bromobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

The synthesis was conducted under the same conditions as in the step 3
5 of the Example 11 by using 3-bromobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.11 g, 70%).

mp 183.0-184.0 °C;

UV (MeOH) λ_{\max} 270.0 nm;

¹H-NMR (DMSO-*d*₆) δ 8.46(s, 1 H), 8.22(s, 1 H), 7.53(s, 1 H), 7.39-7.42(m,
10 1 H), 7.34-7.35(m, 1 H), 7.24-7.28(m, 1 H), 5.90(d, 1 H, *J* = 7.2 Hz), 5.53(d, 1 H-
OH, *J* = 6.4 Hz), 5.35(d, 1 H-OH, *J* = 4.0 Hz), 4.67-4.71(m, 2 H), 4.35-4.37(m, 1
H), 3.41(dd, 1 H, *J* = 4.0, 10.8 Hz), 3.06(q, 1 H, *J* = 7.2 Hz), 2.80(dd, 1 H, *J* = 2.8,
10.8 Hz);

$[\alpha]_{\text{D}}^{25}$ -100.72 (*c* 0.14, DMSO);

15 FAB-MS *m/z* 422 [M + H]⁺.

<Example 14> Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-iodobenzylamino)-9*H*-
purin-9-yl) tetrahydrothiophene-3,4-diol

Step 1. Preparation of 6-chloro-9-((3*aR*,4*R*,6*aS*)-2,2-
20 dimethyltetrahydrothieno[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of foam was obtained in the same
manner as in the step 1 of the Example 11.

Step 2. Preparation of (2*R*,3*S*,4*S*)-2-(6-chloro-9*H*-purin-9-yl)
tetrahydrothiophene-3,4-diol

The target compound as a white solid was obtained in the same manner as in the step 2 of the Example 11.

Step 3. Preparation of (2*R*,3*R*,4*S*)-2-(6-(3-iodobenzylamino)-9*H*-purin-9-yl)tetrahydrothiophene-3,4-diol

5 The synthesis was conducted under the same conditions as in the step 3 of the Example 11 by using 3-iodobenzylamine instead of 3-fluorobenzylamine to obtain the target compound (0.12 g, 72%).

mp 198.8-199.8 °C;

UV (MeOH) λ_{\max} 271.5 nm;

10 $^1\text{H-NMR}$ (DMSO- d_6) δ 8.46(s, 1 H), 8.22(s, 1 H), 7.72(s, 1 H), 7.56-7.59(m, 1 H), 7.35-7.36(d, 1 H, $J = 7.6$ Hz), 7.01-7.12(m, 1 H), 5.90(d, 1 H, $J = 7.2$ Hz), 5.53(d, 1 H-OH, $J = 6.4$ Hz), 5.35(d, 1 H-OH, $J = 4.4$ Hz), 4.67-4.71(m, 2 H), 4.34-4.38(m, 1 H), 3.41(dd, 1 H, $J = 4.0, 10.8$ Hz), 3.16(d, 1 H, $J = 7.2$ Hz), 2.80(dd, 1 H, $J = 2.8, 10.8$ Hz);

15 $[\alpha]_D^{25}$ -97.08 (c 0.14, DMSO);

FAB-MS m/z 470 $[\text{M} + \text{H}]^+$.

<Example 15> Preparation of (2*R*,3*R*,4*R*)-2-(6-(3-bromobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrofuran-3,4-diol

20 Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aR*)-2,2-dimethyltetrahydrofuro[3,4- d][1,3]dioxol-4-yl)-9*H*-purine

The synthesis was conducted by using (3*aR*,4*R*,6*aR*)-2,2-dimethyltetrahydrofuro[3,4- d][1,3]dioxol-4-ol (702.1 g, 3.472 mmol) obtained in Preparation Example 2 under the same conditions as in the step 1 of the Example

1 to obtain the target compound in the form of oil (793.0 mg, 69%).

UV (MeOH) λ_{\max} 276.5 nm;

$^1\text{H-NMR}$ (CDCl_3) δ 8.15(s, 1 H), 6.07(s, 1 H), 5.41(d, 1 H, $J = 6.0$ Hz), 5.26-5.29(m, 1 H), 4.25-4.31(m, 2 H), 1.57(s, 3 H), 1.41(s, 3 H);

5 $[\alpha]_{\text{D}}^{25}$ -21.00 (c 0.10, DMSO);

FAB-MS m/z 331 $[\text{M} + \text{H}]^+$.

Step 2. Preparation of (2*R*,3*R*,4*R*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrofuro-3,4-diol

10 The synthesis was conducted by using 2,6-dichloro-9-((3*aR*,4*R*,6*aS*)-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine (900 mg, 2.0 mmol) prepared in the step 1 under the same conditions as in the step 2 to obtain the target compound as a white solid (0.46 g, 80%).

mp 122.7-123.4 °C;

15 UV (MeOH) λ_{\max} 276.5 nm;

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ 8.98(s, 1 H), 5.96(d, 1 H, $J = 6.4$ Hz), 5.57(d, 1 H-OH, $J = 6.0$ Hz), 5.32(d, 1 H-OH, $J = 4.0$ Hz), 4.69-4.74(m, 1 H), 4.41(dd, 1 H, $J = 3.6, 9.2$ Hz), 4.29-4.32(m, 1 H), 3.87(dd, 1 H, $J = 2.0, 9.6$ Hz);

$[\alpha]_{\text{D}}^{25}$ -68.09 (c 0.14, DMSO);

20 FAB-MS m/z 291 $[\text{M} + \text{H}]^+$.

Step 3. Preparation of (2*R*,3*R*,4*R*)-2-(6-(3-bromobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrofuro-3,4-diol

After dissolving (2*R*,3*S*,4*S*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrofuro-

3,4-diol (1 eq.) prepared in the step 2 and 3-bromobenzylamine (1.5 eq.) in ethanol (5 ml) at room temperature, the reaction mixture was stirred at room temperature for 2-3 h. After completion of the reaction, the reaction mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography using a dichloromethane:methanol mixed solvent (20:1, v/v) as an eluent to obtain the target compound (0.12 g, 82%).

mp 181.5-181.7 °C;

UV (MeOH) λ_{\max} 274.5 nm;

$^1\text{H-NMR}$ (DMSO- d_6) δ 8.92(t, 1 H-NH, J = 6.0 Hz), 8.43(s, 1 H), 7.55(s, 1 H), 7.44(d, 1 H, J = 8.0 Hz), 7.33-7.35(m, 1 H), 7.26-7.30(m, 1 H), 5.81(d, 1 H, J = 6.4 Hz), 5.47(d, 1 H, J = 6.4 Hz), 5.22(d, 1 H, J = 4.0 Hz), 4.66-4.69(m, 1 H), 4.62(s, 2 H), 4.32(dd, 1 H, J = 3.6, 9.2 Hz), 4.25(brs, 1 H), 3.80(dd, 1 H, J = 1.6, 9.2 Hz);

$[\alpha]_{\text{D}}^{25}$ -62.75 (c 0.10, DMSO);

FAB-MS m/z 440 $[\text{M} + \text{H}]^+$.

<Example 16> Preparation of (2*R*,3*R*,4*R*)-2-(6-(3-iodobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrofuro-3,4-diol

Step 1. Preparation of 2,6-dichloro-9-((3*aR*,4*R*,6*aR*)-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)-9*H*-purine

The target compound in the form of syrupy foam was obtained in the same manner as in the step 1 of the Example 15.

Step 2. Preparation of (2*R*,3*R*,4*R*)-2-(2,6-dichloro-9*H*-purin-9-yl)tetrahydrofuro-3,4-diol

The target compound as a white solid in the form of a syrup was obtained in the same manner as in the step 2 of the Example 15.

Step 3. Preparation of (2*R*,3*R*,4*R*)-2-(6-(3-iodobenzylamino)-2-chloro-9*H*-purin-9-yl)tetrahydrofuro-3,4-diol

5 The synthesis was conducted under the same conditions as in the step 3 of the Example 15 by using 3-iodobenzylamine instead of 3-bromobenzylamine to obtain the target compound (0.13 g, 78%).

mp 195.5-195.8 °C;

UV (MeOH) λ_{\max} 274.0 nm;

10 $^1\text{H-NMR}$ (DMSO- d_6) δ 8.91(t, 1 H-NH, $J = 6.4$ Hz), 8.44(s, 1 H), 7.75(s, 1 H), 7.61(d, 1 H, $J = 8.0$ Hz), 7.36(d, 1 H, $J = 7.6$ Hz), 7.13(t, 1 H, $J = 4.0$ Hz), 5.81(d, 1 H, $J = 6.8$ Hz), 5.47(d, 1 H-OH, $J = 6.8$ Hz), 5.23(d, 1 H-OH, $J = 4.0$ Hz), 4.72(dd, 1 H, $J = 6.4, 10.8$ Hz), 4.61(d, 1 H, $J = 6.0$ Hz), 4.34(dd, 1 H, $J = 3.6, 9.2$ Hz), 3.81(dd, 1 H, $J = 1.2, 9.2$ Hz);

15 $[\alpha]_D^{25}$ -68.07 (c 0.12, DMSO);

FAB-MS m/z 488 $[\text{M} + \text{H}]^+$.

<Experimental Example 1> Evaluation of binding affinity for adenosine receptor

20 The following experiments were performed to evaluate the affinity and selectivity of the derivatives of the present invention to the A_1 , A_{2A} and A_3 receptors in the human adenosine receptor (hAR).

The Chinese hamster ovary (CHO, ATCC; American Cell Line No. CCL-61) cells expressing the adenosine A_1 and A_3 receptors were cultured in F-12







(Gibco Co., USA) medium supplemented with 10% fetal bovine serum (FBS) and penicillin/streptomycin (100 units/ml and 100 µg/ml) at 37 °C and 5% carbon dioxide. The 50/10/1 buffer solution was placed in a test tube and a predetermined amount of CHO cells expressing a suitable hAR and label ligands (1 nM [³H]CCPA and 0.5 nM [¹²⁵I]AB-MECA) selectively binding to the respective adenosine A₁ and A₃ receptors were mixed. Various concentrations of the derivatives of the present invention are first dissolved in dimethylsulfoxide (DMSO) and then diluted with buffer solution and the final concentration of DMSO should not exceed 1%, and the cells are incubated in a 37 °C thermostat for 1 hour and the cells were rapidly filtered under reduced pressure using a cell collector (TOMTEC Co., USA). Subsequently, the test tube was washed three times with 3 ml buffer, and radioactivity was determined using a γ-counter. Nonspecific binding was determined in the presence of 10 µM of 5'-N-ethylcarboxamidoadocin (NECA), an unlabeled ligand, under the same conditions as for the determination of total binding, and the equilibrium constant K_i was determined according to the Cheng-Prusoff equation with the assumption that the K_d value of [¹²⁵I]AB-MECA was 1.48 nM. Specific binding was calculated by subtracting non-specific binding from total binding. From the measured specific binding, the bonding affinity of each sample of the various receptors was determined.











In addition, the bonding of A_{2A} receptor expressed in HEK 293 cells (human kidney endothelial cell line) and the [³H]CGS-21680(2-((4-(2-carboxyethyl)phenyl)ethylamino)-5'-N-ethylcarbamoyl)adenosine) which is label ligand, is measured as follows: Adenosine deaminase is added to the cerebral

meninges at 30 °C for 30 minutes and during the incubation with radioactive ligand and the IC₅₀ values for the compounds of the individual examples were determined at 6 and more different concentrations, and the K_i values were determined by a plotting program using this value.

5 Table 1 shows the K_i values obtained from the results of the structure, substituent and binding affinity of the compounds of the examples according to the present invention.

[Table 1]

Example	Structure	substituents			K(nM) or %		
		A	R	Y	hA ₁	hA _{2A}	hA ₃
1		S	3-fluorobenzyl	Cl	19.8%	47.6%	7.4± 1.3
2		S	3-chlorobenzyl	Cl	37.9%	17.7%	1.66± 0.90
3		S	3-bromobenzyl	Cl	34.2%	18.4%	8.99± 5.17
4		S	3-iodobenzyl	Cl	2490± 940	341±75	4.16± 0.50
5		S	2-chlorobenzyl	Cl	12.8%	1600± 135	25.8± 6.3
6		S	5-chloro-2-methoxybenzyl	Cl	23.8%	4020± 1750	12.7± 3.7

7		S	2-methoxybenzyl	Cl	9.4%	17.5%	19.9±7.1
8		S	1-naphthyl	Cl	22.0%	-8.3%	24.8±8.1
9		S	3-toluic acid	Cl	13.1%	-0.18%	41.5%
10		S	methyl	Cl	55.4±1.8%	45.0±1.4%	3.69±0.25
11		S	3-fluorobenzyl	H	1430±420	1260±330	7.3±0.6
12		S	3-chlorobenzyl	H	860±210	440±110	1.5±0.4
13		S	3-bromobenzyl	H	790±190	420±32	6.8±3.4
14		S	3-iodobenzyl	H	530±97	230±65	2.5±1.0
15		O	3-bromobenzyl	Cl	39.8%	22.8%	13.0±6.9
16		O	3-iodobenzyl	Cl	37.7%	28.6%	42.9±8.9

Units: nM±SEM%' represents percent inhibition of specific binding of 10 μM labeled ligand in the presence of 10 μM of NECA, an unlabeled ligand.

As shown in Table 1, the compounds of the present invention exhibited high binding affinity for the human adenosine A₃ receptor and low affinity for the adenosine A₁ and A_{2A} receptors, i.e., high selectivity. In particular, the compound of Example 12 of the present invention exhibited the highest affinity for the hA₃ receptor with affinity constant K_i value of 1.50 ± 0.40 nM and then the binding affinity is high in order of the compound of Example 2 (K_i = 1.66 ± 0.90 nM), the compound of Example 14 (K_i = 2.50 ± 1.00 nM), the compound of Example 10 (K_i = 3.69 ± 0.25 nM) and the compound of Example 4 (K_i = 4.16 ± 0.50 nM). The compound of Example 4 of the present invention also showed a high affinity (K_i = 3.89 ± 1.15 nM) for adenosine A₃ receptor in rats expressed in Chinese hamster ovary (CHO) cells and the human adenosine A_{2B} receptor did not show activity as an agonist or an antagonist.

In addition, the binding affinities in the example compounds having a halobenzyl substituent show in the order Cl > I > F > Br, the compound of Example 2 having 3-chlorobenzyl has higher affinity for hA₃ adenosine receptor than the compound of Example 5 having 2-chlorobenzyl (K_i = 25.8 ± 6.3 nM). In addition, as for the binding affinity of the human adenosine A₃ receptor, the example compounds in which the 3-position of the benzene ring is substituted are more preferred over the 2- or 4-substituted compounds or the 2,5-disubstituted compounds. Also, the compounds of Examples 15 and 16, which are adenosine derivatives having an oxonucleoside form of 4'-O, also showed high binding affinity and selectivity, but were not better than adenosine derivatives having the corresponding 4'-S thionucleoside form and the compounds of Examples 3 and 4, and the compounds of Examples 10 to 14 in which the chloro group at the 2-

position of the purine base was substituted with hydrogen were more excellent in affinity and selectivity than the 2-chloro compounds.

<Experimental Example 2> Antagonistic effect and cAMP inhibition test

5 on adenosine A₃ receptor using derivatives of present invention

In order to investigate whether the derivatives of the present invention have an effect as antagonists against the human adenosine A₃ receptor, CHO cells were treated with the compound of Example 4 and CI-IB-MECA to perform antagonistic effects and cAMP inhibition experiments of the derivatives of the present invention.

As shown in FIG. 1, for the human adenosine A₃ receptor, the effect of an agonist of CI-IB-MECA as a 100% pure agonist in CHO cells treated with the compound of Example 4 at different concentrations was confirmed to be inhibited depending on the concentration of the compound of Example 4. This is a result indicating that the compound of the present invention and CI-IB-MECA competitively act on the same binding site of the receptor. In addition, from a result of cAMP inhibition experiments mediated by the human adenosine A₃ receptor in CHO cells, it can be seen that the compounds of the present invention are 100% pure adenosine A₃ antagonists. Therefore, the compounds synthesized in the present invention had a dissociation constants K_B measured by Schild analysis of 1.92 nM.

<Experimental Examples 3 to 6> Measurement of anti-inflammatory activity of derivatives of present invention

In order to investigate the anti-inflammatory activity of the derivatives of the present invention, the animal experiments were performed as below. Seven-week-old male ICR mice were treated with TPA (12-O-tetradecanoylphorbol 13-acetate, 20 µl) in the right ear. The compounds of Examples 1 to 16 of the present invention were diluted to a concentration of 0.5% in acetone (20 µl) or distilled water or a mixed solvent of DMSO and acetone (its composition is shown in Tables 2 to 5) and was administered to a mouse in 15 minutes. The same experiment was performed by treating hydrocortisone used as a treatment agent for inflammation as a control at the same concentration.

Subsequently, secondly, the adenosine derivative compound of the present invention was treated after 6 hours from TPA treatment. Twenty-four hours after TPA treatment, the animals were euthanized using the cervical dislocation method. Then, a right ear sample was obtained using a 6 mm diameter punch. The activity could be confirmed by weighing using a microbalance. The inhibition rate (%) was calculated using the following Equation 1. The treatment compositions and treatment amounts of Experimental Examples 3 to 6 are shown in Tables 2 to 5, and the results of measuring the anti-inflammatory activity are shown in FIG. 2 to FIG. 5.

[Equation 1]

$$\text{Inhibition rate(\%)} = \frac{1 - (\text{weight of sample treated with Example compound} - \text{weight of untreated sample})}{\text{weight of sample treated with TPA only} - \text{weight of untreated sample weight}}$$

[Table 2]

Experimental Example 3	Treatment composition	Treatment amount
------------------------	-----------------------	------------------

3-1	No treatment	-
3-2	TPA only	20 μ l
3-3	TPA + acetone treatment	20 μ l + 20 μ l
3-4	TPA + acetone + compound of Example 2	20 μ l + 0.5%/20 μ l
3-5	TPA + acetone + compound of Example 3	20 μ l + 0.5%/20 μ l
3-6	TPA + acetone + compound of Example 4	20 μ l + 0.5%/20 μ l
3-7	TPA + acetone + hydrocortisone	20 μ l + 0.5%/20 μ l

[Table 3]

Experimental Example 4	Treatment composition	Treatment amount
4-1	No treatment	-
4-2	TPA only	20 μ l
4-3	TPA + acetone treatment	20 μ l + 20 μ l
4-4	TPA + acetone + compound of Example 1	20 μ l + 0.5%/20 μ l
4-5	TPA + acetone + compound of Example 6	20 μ l + 0.5%/20 μ l
4-6	TPA + acetone + hydrocortisone	20 μ l + 0.5%/20 μ l

[Table 4]

Experimental Example 5	Treatment composition	Treatment amount
5-1	No treatment	-
5-2	TPA only	20 μ l
5-3	TPA + mixed solvent treatment (mixed solvent = DI water: acetone =1:4)	20 μ l + 20 μ l
5-4	TPA + mixed solvent + compound of Example 5	20 μ l + 0.5%/20 μ l
5-5	TPA + mixed solvent + compound of Example 7	20 μ l + 0.5%/20 μ l
5-6	TPA + mixed solvent + compound of Example 8	20 μ l + 0.5%/20 μ l
5-7	TPA + mixed solvent + hydrocortisone	20 μ l + 0.5%/20 μ l

5

[Table 5]

Experimental Example 6	Treatment composition	Treatment amount
6-1	No treatment	-
6-2	TPA only	20 μ l
6-3	TPA + mixed solvent treatment (mixed solvent = DMSO:acetone =1:9)	20 μ l + 20 μ l
6-4	TPA + mixed solvent + compound of Example 15	20 μ l + 0.5%/20 μ l
6-5	TPA + mixed solvent + compound of Example 16	20 μ l + 0.5%/20 μ l
6-6	TPA + mixed solvent + hydrocortisone	20 μ l + 0.5%/20 μ l

As shown in FIG. 2, after diluting the compounds of Examples 2, 3 and 4 in acetone, mice were treated with them to measure the anti-inflammatory activity. The TPA-induced mouse ear edema was reduced by a small amount, although it is very little change compared to hydrocortisone used as a control.

As shown in FIG. 3, after diluting the compounds of Examples 1 and 6 in acetone, mice were treated with them to measure the anti-inflammatory activity and It was found that the inhibition rate was significantly increased at least four times as compared with the compounds of Examples 2 to 4 of FIG. 2.

As shown in FIG. 4, the anti-inflammatory activities of the compounds of Examples 5, 6 and 7 of the present invention diluted to 0.5% in a mixed solvent of distilled water and acetone (1:4) were 17% and 34% and 53% of inflammation inhibition rate, respectively.

As shown in FIG. 5, the inflammation inhibition rates by the compounds of Examples 15 and 16 diluted to 0.5% in a mixed solvent of DMSO and acetone (1:9) were 59% and 79%, respectively, which indicated that the compound had anti-inflammatory activity.

<Experimental Example 7> Toxicity test

In order to test the toxicity of the example compounds of the present invention, animal experiments were performed. Three each mice of 25 ± 5 g ICR mice (central laboratory animals) and 235 ± 10 g Specific Pathogen Free (SPF) Sprague Dawley (central laboratory animals) rats were divided into three groups and the compound of Example 2 was intraperitoneally administered at doses of 20 mg/kg, 10 mg/kg and 1 mg/kg, respectively, and then observed for toxicity for 24 hours.

As a result of the experiment, no deaths were observed in all of the three groups, and there was no apparent symptom with the control group including weight gain, feed consumption, and the like, and thus it was confirmed that the derivative compound of the present invention is a safe drug.

<Experimental Example 8> Test for effect of eye-drop formulation comprising derivatives of present invention on intraocular pressure reduction (1)

The following animal experiments were performed to examine the intraocular pressure decrease effect according to eye drop administration of the derivatives of the present invention. The compound of Example 2 was mixed with a buffer solution of pH 6.8 in which Cremophor EL, glycerin, citric acid and methylparaben were dissolved to prepare eye drops containing the compounds of Example 2 at 250 μ M, 500 μ M and 750 μ M, respectively and were administered by eye drop to 2.5-3.0 kg of normal male New Zealand white rabbit by 2 drops, respectively. As a positive control, Xalatan®, an eye drop for treating glaucoma,

and as a negative control, no treatment was given to the experimental animals.

Then, the intraocular pressure of the laboratory animals was measured after 30 minutes, 1 hour, 2 hours, 3 hours, 6 hours, 12 hours and 24 hours, and the results are shown in FIG. 6. In FIG. 6, Control represents a negative control,
5 Drug 250, 500 and 750 are eye drops containing the compound of Example 2 at 250 μ M, 500 μ M and 750 μ M, respectively, and Xalatan is a positive control.

As shown in FIG. 6, it can be confirmed that the rabbit administered with the eye drops comprising the adenosine derivative of the present invention had an effect of reducing the intraocular pressure depending on the dose of the eye
10 drops.

<Experimental Example 9> Test for effect of eye-drop formulation comprising derivatives of present invention on intraocular pressure reduction (2)

The following animal experiments were performed to examine the
15 intraocular pressure decrease effect according to eye drop administration of the derivatives of the present invention. The eye drops prepared in Experimental Example 8 were administered to the DAB 2J mice induced glaucoma twice daily. As a positive control group, Xalatan®, an eye drop used as a glaucoma treatment agent, was administered in the same manner, and no treatment was given to the
20 laboratory animals as a negative control group.

Then, the intraocular pressure of the laboratory animals was measured every 2 days during the period of administration of the eye drops, and the results are shown in FIG. 7. In FIG. 7, Control represents a negative control, Drug 250, 500 and 750 represent eye drops comprising the compound of Example 2 at 250

μM , 500 μM and 750 μM , respectively, and Xalatan is a positive control.

As shown in FIG. 7, it can be confirmed that the mouse administered with the eye drops comprising the adenosine derivative of the present invention had an effect of reducing the intraocular pressure depending on the dose of the eye drops.

<Experimental Example 10> Intraocular pressure reduction test following the oral administration of derivatives of present invention

In order to examine the intraocular pressure lowering effect of the derivatives of the present invention, the following animal experiments were performed. Dexamethasone was administered by eye drop to 10-week-old normal C57BL/6J mice to increase intraocular pressure and the compound of Example 2 by 5 mg/kg of once a day, 10 mg/kg once a day, 20 mg/kg once a day, and 10 mg/kg twice a day were orally administered together with the excipients DMSO, PEG400 and saline to each mouse. As controls, only the above excipients were administered to laboratory animals with elevated intraocular pressure once a day, or experimental animals were not treated with dexamethasone or with 4.17 mg/kg of acetazolamide, an intraocular pressure lowering agent, once a day.

Then, the intraocular pressure of the test animal was measured at predetermined intervals for 22 days, and the results are shown in FIG. 8 to FIG. 12. In FIG. 8 to FIG. 12, Control represents a control group in which dexamethasone is not treated to laboratory animals, Vehicle represents a control

group in which only excipients were orally administered to laboratory animals with elevated intraocular pressure, and Acetazolamide represents control group in which acetazolamide, an intraocular pressure lowering agent was orally administered to laboratory animals with elevated intraocular pressure.

5 As shown in FIG. 8 to FIG. 12, mice administered orally with the adenosine derivative of the present invention showed a reduction in intraocular pressure, which was dependent on the dose of the adenosine derivative. Mice in which the compound of Example 2 was administered at a dose of 20 mg/kg once a day and 10 mg/kg twice a day showed similar intraocular pressure to that of the
10 normal mice control group without treatment with dexamethasone from the 22nd day after administration.

<Experimental Example 11> Anti-fibrotic effect of adenosine derivatives of present invention on trabecular meshwork tissue

15 In order to investigate whether the derivatives of the present invention have an effect of inhibiting fibrosis of the trabecular meshwork tissue of the eye, the following animal experiments were performed.

In Experimental Example 10, the ocular tissues of each laboratory animal after measurement of intraocular pressure were extracted and sectioned and
20 immunostaining using antibody was performed to confirm the expression of extracellular matrix proteins (Fibronectin, SMA and MYOC) and the results are shown in FIG. 13. In FIG. 13, Normal represents the trabecular meshwork of the animal which dexamethasone was not treated with in Experimental Example 10, Control represents the trabecular meshwork tissue of the animal to which only

the excipient was administered in Experimental Example 10, Acetazolamide represents the trabecular meshwork tissue of the animal to which acetazolamide was administered in Experimental Example 10. Fluorescently stained tissue was observed under fluorescence microscope under the same conditions
5 (magnification 200x, exposure time 1 sec).

As shown in FIG. 13, in the trabecular meshwork tissue of the animal to which only excipients were administered or acetazolamide, an intraocular pressure lowering agent was administered, increased expression of fibronectin, smooth muscle actin (SMA) and myocilin (MYOC) protein was observed and
10 while the expression of the three proteins was confirmed to be greatly reduced in the trabecular meshwork tissue of the animal to which the compound of Example 2 was administered at a dose of 20 mg/kg once a day and at a dose of 10 mg/kg twice a day.

15 <Experimental Example 12> Test for effect of adenosine derivative of present invention on MMP enzyme and TIMP enzyme protein expression

In order to confirm whether the anti-fibrotic effect of the adenosine derivative of the present invention on the trabecular meshwork tissue is related to the expression of MMP and TIMP enzyme protein which regulate the
20 extracellular matrix changes, the following animal experiments were performed.

In Experimental Example 10, the expression amounts of these proteins and mRNAs in MMP and TIMP enzymes were measured for the eye tissues of the respective laboratory animals which the measurement of the intraocular pressure was completed, and the results are shown in FIG. 14 to FIG. 17. FIG.

14 and FIG. 15 show protein expression amounts of MMP (MMP2, MMP9) and TIMP (TIMP 1, TIMP2) enzymes, respectively. FIG. 16 and FIG. 17 show mRNA expression levels. In FIG. 14 to FIG. 17, Normal represents the expression level of the animal to which the dexamethasone was not treated in Experimental Example 10, Control represents the expression level of the animal to which only the excipient was administered in Experimental Example 10, and Acetazolamide represents the expression level of the animal to which acetazolamide was administered in Experimental Example 10.

As shown in FIG. 14 to FIG. 17, the protein and mRNA expression levels of the MMP enzyme were significantly decreased in the trabecular meshwork tissue of the animal to which only the excipient was administered or the animal to which acetazolamide, the intraocular pressure reducing agent was administered compared with the normal group and protein and mRNA expression levels of TIMP enzymes were significantly increased. On the other hand, the amount of protein and mRNA expression of MMP enzyme were significantly increased in the trabecular meshwork tissue of the animal administered with 20 mg/kg once a day and 10 mg/kg twice a day of the compound of Example 2 compared to the normal group and the amount of protein and mRNA expression of TIMP enzyme was significantly decreased.

Through the above-mentioned series of experimental results, the adenosine derivatives of the present invention is confirmed to have the effect of reducing the intraocular pressure by inhibiting the production of aqueous humor, which is known to be common to adenosine A₃ receptor antagonists, act to reduce intraocular pressure through the dual mechanism by promoting the release of

aqueous humor through the inhibition of fibrosis of the trabecular meshwork tissue.

The commercially available glaucoma therapeutic agents lower the intraocular pressure through one of the two pharmacological mechanisms mentioned above, however, since the adenosine derivative of the present invention has the both mechanisms of action, it can be used as a composition highly suitable for the prevention and treatment of glaucoma.

<Experimental Example 13> Physicochemical characteristic test of adenosine derivative of present Invention

In order to test the physicochemical properties of the adenosine derivatives of the present invention, the compounds of Example 2 were tested in vitro and the results are shown in Table 6. Plasma stability and protein binding were measured using rat and human plasma.

[Table 6]

Properties (ADME Characteristics)	Value
Kinetic solubility@	361.0 μ M (148.8 μ g/ml)
Equilibrium solubility	6.7 μ M (2.76 μ g/ml)
Log P	3.18
pKa	11.33
PAMPA	-4.49
Plasma stability	>99.9 (Rat), 98.9 (Human)
Plasma protein binding	90.2(Rat), 98.7 (Human)

As shown in Table 6, it can be confirmed that the adenosine derivatives of the present invention have absorption, distribution, metabolism and excretion (ADME) characteristics suitable for oral administration by oral preparations.

5 <Experimental Examples 14 to 18> pharmacokinetic test for oral administration of adenosine derivative of present invention

In order to test the absorption, distribution, metabolism and excretion (ADME) characteristics of the adenosine derivatives of the present invention after oral administration, the PK (Pharmacokinetic) characteristics of the compound of Example 2 were measured in vivo.

As shown in Table 7, the compound of Example 2 was administered to laboratory animals through different administration methods. Intravenous administration was performed through a tube inserted into the femoral vein, and oral administration was performed using an oral gavage.

15 [Table 7]

Experimental Example	Laboratory animal	Administration method
14-1	8-week-old SD male rat	5 mg/kg of compound of Example 2 was intravenously administered
14-2	8-week-old SD male rat	5 mg/kg of compound of Example 2 was orally administered
15-1	8-week-old SD male rat	2 mg/kg of compound of Example 2 was intravenously administered
15-2	8-week-old SD male rat	10 mg/kg of compound of Example 2 was orally administered
16	8-week-old ICR male mice	10 mg/kg of compound of Example 2 was orally administered
17-1	Dog	2 mg/kg of compound of Example 2 was intravenously administered
17-2	Dog	10 mg/kg of compound of Example 2 was dissolved in a solvent and orally administered

17-3	Dog	10 mg/kg of compound of Example 2 was orally administered in powder form in capsules
18-1	8-week-old SD male rat	10 mg/kg of compound of Example 2 was dissolved in 2 mL/kg of 0.5 wt% methylcellulose and orally administered
18-2	8-week-old SD male rat	10 mg/kg of compound of Example 2 was dissolved in a solvent mixed with 5 wt% DMSO, 40 wt% PEG 400, 55 wt% of D.W. and orally administered

Blood was collected at predetermined time intervals for 24 hours after the administration and centrifuged, plasma was separated, and plasma samples were pretreated with a suitable organic solvent to analyze the concentration by LC-MS/MS. The blood concentration-time data of the compound of Example 2 was analyzed using WinNonlin (Pharsight, USA), and the graphs thereof are shown in FIG. 20 to FIG. 24, and the results of the noncompartmental pharmacokinetic parameters obtained therefrom are shown in Tables 8 to 12. In FIG. 8 to FIG. 12, I.V. represents the intravenous administration group, P.O. represents the oral administration group, and definitions of the respective parameters of Tables 8 to 12 are shown in Table 13.

[Table 8]

Parameters	I.V., 5 mg/kg	P.O., 5 mg/kg
T _{max} (h)	NA	1.33 ± 0.577
C _{max} (µg/mL)	NA	1.45 ± 0.255
T _{1/2} (h)	3.6 ± 0.589	3.26 ± 0.945
AUC _t (µg·h/mL)	14.04 ± 2.55	6.98 ± 0.584
AUC _∞ (µg·h/mL)	14.11 ± 2.59	7.04 ± 0.551
CL (L/h/kg)	0.363 ± 0.07	NA
V _{ss} (L/kg)	0.881 ± 0.203	NA
F _t (%)	NA	49.74

NA, not applicable; ND, not detected; NC, not calculated

[Table 9]

Parameters	IV, 2 mg/kg	PO, 10 mg/kg
T _{max} (hr)	-	2.42 ± 3.13
C _{max} (µg/mL)	-	2.71 ± 0.183
T _{1/2} (hr)	6 ± 2.98	3.34 ± 0.075
AUC _t (µg·hr/mL)	5.2 ± 0.548	26.5 ± 5.88
AUC _∞ (µg·hr/mL)	5.49 ± 0.3	26.7 ± 0.0750
CL (L/kg/hr)	0.365 ± 0.019	-
V _{ss} (L/kg)	2.27 ± 0.863	-
F _t (%)	-	>99.9

5

[Table 10]

Parameters	P.O., 10 mg/kg
T _{max} (h)	6.13 ± 3.75
C _{max} (µg/mL)	8.57 ± 1.52
T _{1/2} (h)	3.61 ± 0.3
AUC _t (µg·h/mL)	100 ± 13.2
AUC _∞ (µg·h/mL)	102 ± 13.5
CL (L/h/kg)	NA
V _{ss} (L/kg)	NA
F _t (%)	NA

NA, not applicable; ND, not detected; NC, not calculated

[Table 11]

Parameters	G1, IV, 2 mg/kg	G2, PO, 10 mg/kg	G3, PO, 10 mg/kg
T _{max} (h)	NA	1.67 ± 0.58	2 ± 0
C _{max} (µg/mL)	NA	0.467 ± 0.073	1.14 ± 0.23
T _{1/2} (h)	2.17 ± 0.867	4.21 ± 1.41	5.53 ± 3.06
AUC _t (µg·h/mL)	0.948 ± 0.464	3.88 ± 1.03	5.64 ± 0.84
AUC _∞ (µg·h/mL)	1.07 ± 0.62	3.99 ± 1.09	6.35 ± 0.83
CL (L/h/kg)	2.27 ± 1.04	NA	NA
V _{ss} (L/kg)	6.02 ± 0.79	NA	NA
F _t (%)	NA	82.0	>99.9

NA, not applicable; ND, not detected; NC, not calculated

[Table 12]

Parameters	0.5%MC, 10 mg/kg	기존 Vehicle, 10 mg/kg
T_{max} (hr)	1.93 ± 0.58	2.42 ± 3.13
C_{max} (µg/mL)	5.72 ± 6.11	2.71 ± 0.183
$T_{1/2}$ (hr)	4.56 ± 2.8	3.34 ± 0.075
AUC_t (µg·hr/mL)	40.1 ± 26.8	26.5 ± 5.88
AUC_{∞} (µg·hr/mL)	41.4 ± 26.03	26.7 ± 0.0750
CL (L/kg/hr)	-	-
V_{ss} (L/kg)	-	-
F_t (%)	-	-

[Table 13]

Parameters	Description
T_{max} (hr)	time for C_{max}
C_{max} (µg/mL)	maximum plasma concentration
$T_{1/2}$ (hr)	terminal half-life
AUC_t (µg·hr/mL)	areas under the plasma concentration-time curve
AUC_{∞} (µg·hr/mL)	areas under the plasma concentration-time curve from time
CL (L/kg/hr)	total clearance from plasma
V_{ss} (L/kg)	steady-state volume of distribution
F_t (%)	bioavailability ($AUC_{p.o.}/AUC_{i.v.}$) X 100

5

FIG. 18 and Table 8 are graph and parameter values obtained from the blood concentration-time data of Experimental Example 14 (14-1 and 14-2), respectively, FIG. 19 and Table 9 are graph and parameter values obtained from the blood concentration-time data of Experimental Example 15 (15-1 and 15-2), respectively. As shown in FIG. 18 and FIG. 19 and Tables 8 and 9, it is confirmed that the adenosine derivative of the present invention has a half-life ($T_{1/2}$) of a maximum of 3.34 hours or more for a long time and the bioavailability (F_t) is maximum 99.9% or more as compared with the intravenous administration, and thus it is suitable for oral administration.

15

FIG. 20 and Table 10 are graph and parameter values obtained from the blood concentration-time data of Experimental Example 16. As shown in FIG. 20

and Table 10, it is confirmed that the adenosine derivative of the present invention has long half-life ($T_{1/2}$) of about 3.61 hours in mice and is suitable for oral administration.

FIG. 21 and Table 11 are graph and parameter values obtained from the
5 blood concentration-time data of Experimental Example 17 (17-1, 17-2 and 17-3). In FIG. 21, G2 and G3 represent oral administration after dissolving in a solvent and oral administration in powder form in a capsule, respectively. As shown in FIG. 21 and Table 11, the adenosine derivatives of the present invention have a half-life ($T_{1/2}$) of maximum 5.53 hours or more i.e. is exposed for longer
10 time than rats and mice and are suitable for oral administration and thus are more suitable for oral administration. Particularly, it is confirmed that when it is filled in powder form within a capsule, the half-life ($T_{1/2}$) and bioavailability (F_i) characteristics, etc. are improved.

FIG. 22 and Table 12 are graph and parameter values obtained from the
15 blood concentration-time data of Experimental Example 18 (18-1 and 18-2). As shown in FIG. 22 and Table 12, when the adenosine derivatives of the present invention is orally administered in combination with methyl cellulose (MC), It has better properties than those administered with conventional excipients dimethylsulfoxide (DMSO), polyethylene glycol (PEG), distilled water (D.W.) and
20 the like.

<Experimental Example 19> Toxicity test of adenosine derivative of present invention

In order to test the toxicity of the adenosine derivatives of the present

invention, the compounds of Example 2 were evaluated for cytotoxicity, cardiotoxicity (hERG ligand binding assay), genotoxicity and single dose toxicity.

First, the Cyto XTM cell viability assay kit was used to test the cytotoxicity of the compound of Example 2. As a result of the test, it was evaluated that the
5 IC₅₀ by the compound of Example 2 was 10 µM or more in each cell line and was safe for general cytotoxicity.

To test the cardiotoxicity of the compound of Example 2, a non-electrophysiological test was used to evaluate the heart stability through fluorescence polarization evaluation according to the hERG channel protein
10 binding degree of the red fluorescent hERG channel ligand tracer. As a result of the test, the inhibition rate against 10 µM of the compound of Example 2 was 50% or less, which is a standard value, and it was evaluated to be safe against cardiotoxicity.

In order to test the genotoxicity of the compound of Example 2, the gene
15 mutagenicity of the compound of Example 2 was evaluated for each case in the presence and absence of metabolic activation in the presence and absence of metabolic activation using histidine-requiring Salmonella (TA98, TA100, TA1535 and TA1537 strains) and tryptophan-requiring E. coli (WP2uvrA (pKM101) strain)
The gene mutagenicity of the compound of Example 2 was evaluated for each
20 case.

As a result of evaluation, as for the compound of Example 2, the number of back-mutation reversion colonies at all doses of each strain did not exceed twice the negative control, regardless of metabolic activation and no dose-dependent increases were observed and in the positive control group, the number

of back-mutation reversion colonies for each strain was at least doubled as compared with the negative control. From the above results, the compound of Example 2 was evaluated as safe against genotoxicity.

To test the single dose toxicity of the compound of Example 2, a single dose of 2,000 mg/kg of the compound of Example 2 was administered to each 5 male rats and 5 female rats. As a result of the test, there was no dead animal, and according to the results, the compound of Example 2 was evaluated as safe for single dose toxicity.

Table 14 summarizes the toxicity test results of the adenosine derivatives of the present invention described above. As shown in Table 14, it can be seen that the adenosine derivatives of the present invention are safe against cytotoxicity, cardiotoxicity, genotoxicity and single dose toxicity.

[Table 14]

Test	toxicity
Cytotoxicity Test	Not found
Cardiotoxicity test	Not found
Genotoxicity test	Not found
Single toxicity assessment	Not found
Single dose toxicity test	Not found

<Experimental Example 20> Evaluation of stability of an oral administration agent comprising adenosine derivatives of present invention

In order to evaluate the stability of the oral administration agent comprising the adenosine derivative of the present invention, the following experiment was performed.

0.5 wt% methylcellulose, which can be used as an excipient for oral

administration, was added to the compound of Example 2, sonicated and homogenized, and then divided into a group to be stored at room temperature or at 4 °C, and after 1, 3, 7, and 10 days, the stability was measured by comparing the concentration with 0.5% methyl cellulose as the control group using Waters

5 UPLC and the results are shown in Table 15.

[Table 15]

Storage condition		Stability (%)		
		200 µg/ml	1000 µg/ml	3000 µg/ml
Control		100±28.7	100±9.55	100±5.40
Stored at RT	After 1 day	84.3±4.10	82.7±16.3	90.5±6.44
	After 3 days	76.3±8.12	82.7±37.0	80.5±5.01
	After 7 days	80.9±14.6	75.4±6.66	101±13.2
	After 10 days	90.3±23.2	94.2±9.17	92.2±2.17
Stored at 4°C	After 1 day	118±21.4	76.5±11.7	84.8±17.9
	After 3 days	125±40.2	87.6±9.26	94.8±2.55
	After 7 days	88.3±17.3	80.1±27.7	89.7±8.30
	After 10 days	93.8±44.8	79.5±22.0	99.0±3.87

As shown in Table 15, when the adenosine derivative of the present invention was prepared as an oral administration agent, the stability of the adenosine derivative was not significantly different according to storage conditions and storage time, and thus, it is confirmed that the adenosine derivative of the present invention was suitable for oral administration by an oral

10


preparation.

While the present invention has been described in connection with what is presently considered to be practical exemplary embodiments, it is to be understood by those skilled in the art that various changes in form and details
5 may be made therein without departing from the spirit and scope of the invention as defined by the appended claims. It is therefore to be understood that the above-described embodiments are illustrative in all aspects and not restrictive.

CLAIMS

1. An oral pharmaceutical composition when used for preventing or treating intraocular pressure elevation and/or glaucoma, which comprises a compound represented by following Chemical Formula A or a pharmaceutically acceptable salt:

[Chemical Formula A]



and a pharmaceutically acceptable excipient.

2. The oral pharmaceutical composition of claim 1, wherein the excipient comprises at least one selected from the group consisting of methyl cellulose (MC), dimethyl sulfoxide (DMSO), polyethylene glycol (PEG) and distilled water.

3. The oral pharmaceutical composition of claim 1 or claim 2, wherein the excipient comprises 0.5 wt% of methyl cellulose.

4. The oral pharmaceutical composition of any one of claims 1 to 3, wherein the compound represented by the Chemical Formula A or a pharmaceutically acceptable salt thereof is filled in capsules in powder form.


5. A method for the prevention or treatment of intraocular pressure elevation and/or glaucoma comprising administering an effective amount of the oral pharmaceutical composition of any one of claims 1 to 4 to a subject in need thereof.

6. Use of the oral pharmaceutical composition of any one of claims 1 to 4 for preventing or treating intraocular pressure elevation and/or glaucoma.


7. Use of the oral pharmaceutical composition of any one of claims 1 to 4 in the manufacture of a medicament for preventing or treating intraocular pressure elevation and/or glaucoma.

[Drawings]

[FIG. 1]




[FIG. 2]




5


[FIG. 3]




[FIG. 4]




[FIG. 5]




5 [FIG. 6]




[FIG. 7]




[FIG. 8]




5 [FIG. 9]




[FIG. 10]




[FIG. 11]




5 [FIG. 12]




[FIG. 13]




[FIG. 14]




[FIG. 15]




[FIG. 16]




[FIG. 17]



[FIG. 18]




[FIG. 19]




5

[FIG. 20]



[FIG. 21]



[FIG. 22]

