



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
04.07.2018 Bulletin 2018/27

(51) Int Cl.:
C11D 17/04 (2006.01) **C11D 3/37** (2006.01)
C11D 1/14 (2006.01) **C11D 1/29** (2006.01)

(21) Application number: **16207069.2**

(22) Date of filing: **28.12.2016**

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
Designated Extension States:
BA ME
Designated Validation States:
MA MD

(71) Applicant: **The Procter & Gamble Company**
Cincinnati, OH 45202 (US)

(72) Inventors:
• **URE, Colin**
Newcastle-upon-Tyne, NE12 9TS (GB)

• **BROOKER, Alan Thomas**
Newcastle upon Tyne, NE12 9TS (GB)
• **SOMERVILLE ROBERTS, Nigel Patrick**
Newcastle-upon-Tyne, NE12 9TS (GB)
• **KEULEERS, Robby Renilde Francois**
1853 Strombeek-Bever (BE)

(74) Representative: **Pickford, James Lawrence**
Procter & Gamble
Technical Centres Limited
Whitley Road
Longbenton
Newcastle upon Tyne NE12 9TS (GB)

(54) **WATER-SOLUBLE UNIT DOSE ARTICLE COMPRISING ZWITTERIONIC POLYAMINE**

(57) The present disclosure is to a water-soluble unit dose article comprising a laundry detergent composition wherein the detergent contains a zwitterionic polyamine, and methods of using said unit dose article.




Fig. 1

Description**FIELD OF THE INVENTION**

5 **[0001]** The present invention is to a water-soluble unit dose article comprising a laundry detergent composition wherein the detergent comprises a zwitterionic polyamine, and methods of using said unit dose article.

BACKGROUND OF THE INVENTION

10 **[0002]** Water-soluble unit dose articles are liked by consumers due to their convenience and ease of use during the laundry operation.

[0003] Such water-soluble unit dose articles often comprise laundry detergent compositions. Formulators prefer to formulate alkyl sulphates or alkoxylated alkyl sulphates into laundry detergent compositions to provide cleaning benefits. Alkyl sulphates and alkoxylated alkyl sulphates exhibit improved water hardness tolerance as compared to other anionic surfactant such as linear alkylbenzene sulphonate. This results in better whiteness as well as robust cleaning and sudsing performance across wash conditions. In addition alkyl sulphates and alkoxylated alkyl sulphates show excellent cleaning performance on hydrophilic stains such as grass stains, especially under diluted conditions such as so called 'top loader' washing machines. Often, ethoxylated polyethyleneimines are also formulated to provide particulate soil removal.

15 **[0004]** However, there is a need for water-soluble unit dose articles comprising liquid detergent compositions comprising alkyl sulphates/alkoxylated alkyl sulphates to provide improved particulate soil removal from fabrics during the wash.

20 **[0005]** It was surprisingly found that a water-soluble unit dose article comprising a laundry detergent composition comprising an alkyl sulphate/alkoxylated alkyl sulphate and a zwitterionic polyamine overcame this technical problem.

SUMMARY OF THE INVENTION

25 **[0006]** A first aspect of the present invention is a water-soluble unit dose article comprising a water-soluble film and a laundry detergent composition, wherein the laundry detergent composition is preferably selected from a liquid, a solid or a mixture thereof, and wherein said detergent composition comprises;

- 30
- a. a zwitterionic polyamine, preferably between 0.01% to about 20%, preferably from 0.1% to 10%, more preferably from 0.5% to 7%, even more preferably from 1% to 5%, most preferably from 2% to 4% by weight of the laundry detergent composition of the zwitterionic polyamine;; and
 - b. a first non-soap anionic surfactant selected from an alkyl sulphate anionic surfactant, an alkoxylated alkyl sulphate anionic surfactant or a mixture thereof.
- 35

[0007] A second aspect of the present invention is a process for washing fabrics comprising the steps of;

- 40
- a. Combining a water-soluble unit dose article according to the present invention with sufficient water to dissolve the water-soluble film and dilute the laundry detergent composition by a factor of between 300 and 800 fold to form a wash liquor;
 - b. Combining the wash liquor with at least one fabric to be washed.

BRIEF DESCRIPTION OF THE DRAWINGS

45 **[0008]** FIG. 1 is a water-soluble unit dose article according to the present invention.

DETAILED DESCRIPTION OF THE INVENTION**Water-soluble unit dose article**

[0009] The present invention discloses a water-soluble unit dose article comprising a water-soluble film and a laundry detergent composition. The laundry detergent composition is described in more detail below. The water-soluble film is described in more detail below.

55 **[0010]** The water-soluble unit dose article comprises the water-soluble film shaped such that the unit-dose article comprises at least one internal compartment surrounded by the water-soluble film. The unit dose article may comprise a first water-soluble film and a second water-soluble film sealed to one another such to define the internal compartment. The water-soluble unit dose article is constructed such that the detergent composition does not leak out of the compart-

ment during storage. However, upon addition of the water-soluble unit dose article to water, the water-soluble film dissolves and releases the contents of the internal compartment into the wash liquor.

[0011] The compartment should be understood as meaning a closed internal space within the unit dose article, which holds the detergent composition. During manufacture, a first water-soluble film may be shaped to comprise an open compartment into which the detergent composition is added. A second water-soluble film is then laid over the first film in such an orientation as to close the opening of the compartment. The first and second films are then sealed together along a seal region.

[0012] The unit dose article may comprise more than one compartment, even at least two compartments, or even at least three compartments. The compartments may be arranged in superposed orientation, i.e. one positioned on top of the other. In such an orientation the unit dose article will comprise three films, top, middle and bottom. Alternatively, the compartments may be positioned in a side-by-side orientation, i.e. one orientated next to the other. The compartments may even be orientated in a 'tyre and rim' arrangement, i.e. a first compartment is positioned next to a second compartment, but the first compartment at least partially surrounds the second compartment, but does not completely enclose the second compartment. Alternatively one compartment may be completely enclosed within another compartment.

[0013] Wherein the unit dose article comprises at least two compartments, one of the compartments may be smaller than the other compartment. Wherein the unit dose article comprises at least three compartments, two of the compartments may be smaller than the third compartment, and preferably the smaller compartments are superposed on the larger compartment. The superposed compartments preferably are orientated side-by-side.

[0014] In a multi-compartment orientation, the detergent composition according to the present invention may be comprised in at least one of the compartments. It may for example be comprised in just one compartment, or may be comprised in two compartments, or even in three compartments.

[0015] Each compartment may comprise the same or different compositions. The different compositions could all be in the same form, or they may be in different forms.

The water-soluble unit dose article may comprise at least two internal compartments, wherein the laundry detergent composition is comprised in at least one of the compartments, preferably wherein the unit dose article comprises at least three compartments, wherein the detergent composition is comprised in at least one of the compartments.

[0016] The water-soluble unit dose article may comprise at least two compartments, preferably at least three compartments, wherein the laundry detergent composition is comprised within at least one compartment. The laundry detergent composition may represent a culmination of ingredients located within all the compartments of the unit dose article.

[0017] FIG. 1 discloses a water-soluble unit dose article (1) according to the present invention. The water-soluble unit dose article (1) comprises a first water-soluble film (2) and a second water-soluble film (3) which are sealed together at a seal region (4). The laundry detergent composition (5) is comprised within the water-soluble unit dose article (1).

Laundry detergent composition

[0018] The water-soluble unit dose article comprises a laundry detergent composition. The laundry detergent composition, may be a liquid, a solid or a mixture thereof.

[0019] The term 'solid laundry detergent composition' refers to any laundry detergent composition that is solid. Solid can include, particles, compressed solids or a mixture thereof.

[0020] The term 'liquid laundry detergent composition' refers to any laundry detergent composition comprising a liquid capable of wetting and treating a fabric, and includes, but is not limited to, liquids, gels, pastes, dispersions and the like. The liquid composition can include solids or gases in suitably subdivided form, but the liquid composition excludes forms which are non-fluid overall, such as tablets or granules.

[0021] The liquid detergent composition can be used in a fabric hand wash operation or maybe used in an automatic machine fabric wash operation.

[0022] The laundry detergent composition comprises a zwitterionic polyamine. The zwitterionic polyamine is described in more detail below.

[0023] The water-soluble unit dose article may comprise between 0.01% to about 20%, preferably from 0.1% to 10%, more preferably from 0.5% to 7%, even more preferably from 1% to 5%, most preferably from 2% to 4% by weight of the laundry detergent composition of the zwitterionic polyamine.

[0024] The laundry detergent composition comprises a first non-soap anionic surfactant selected from an alkyl sulphate anionic surfactant, an alkoxyated alkyl sulphate anionic surfactant or a mixture thereof.

[0025] Preferably, the laundry detergent composition comprises between 2% and 30%, preferably between 5% and 25%, more preferably between 7% and 20%, most preferably between 10% and 17% by weight of the laundry detergent composition of the first non-soap surfactant.

[0026] The first non-soap surfactant is preferably a C10-18 alkyl sulphate, a C10-18 alkoxyated alkyl sulphate or a mixture thereof.

[0027] Preferably, the first non-soap surfactant is an alkoxyated alkyl sulphate, preferably an ethoxyated alkyl sulphate

with an average degree of ethoxylation of between 0.5 and 7, preferably between 1 and 5, more preferably between 2 and 4, most preferably about 3, or wherein the first non-soap surfactant is a mixture of one or more alkoxyated alkyl sulphates, preferably ethoxylated alkyl sulphates, and optionally an alkyl sulphate, the mixture having an average degree of ethoxylation of between 0.5 and 7, preferably between 1 and 5, more preferably between 2 and 4, most preferably about 3.

[0028] The laundry detergent composition may comprise a second non-soap anionic surfactant, wherein the second non-soap anionic surfactant is selected from linear alkylbenzene sulphonate. Preferably, the laundry detergent composition comprises between 2% and 30%, preferably between 5% and 27%, more preferably between 7% and 25%, most preferably between 10% and 23% or even between 10% and 17% by weight of the laundry detergent composition of the second non-soap surfactant.

[0029] The laundry detergent composition may comprise between 0% and 10% preferably between 0.01% and 8%, more preferably between 0.1% and 6%, most preferably between 0.15% and 4% by weight of the laundry detergent composition of a non-ionic surfactant. Preferably, the non-ionic surfactant is selected from alcohol alkoxylate, an oxo-synthesised alcohol alkoxylate, Guerbet alcohol alkoxylates, alkyl phenol alcohol alkoxylates or a mixture thereof.

[0030] Suitable alcohol ethoxylate nonionic surfactants include the condensation products of aliphatic alcohols with from 1 to 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, guerbet, primary or secondary, and generally contains from 8 to 22 carbon atoms. The starting alcohol can be naturally derived, e.g. starting from natural oils, or synthetically derived, e.g. alcohols obtained from for example oxo-, modified oxo- or Fischer-Tropsch processes. Examples of oxo-process derived alcohols include the Lial and Isalchem alcohols ex Sasol company and Lutensol alcohols ex BASF company. Examples of modified-oxo process derived alcohols include the Neodol alcohols ex Shell company. Fischer-Tropsch derived alcohols include Safol alcohols ex Sasol company. The alkoxylate chain of alcohol ethoxylates is made up solely of ethoxylate groups.

[0031] Preferably, the alcohol ethoxylate non-ionic surfactant comprises on average between 8 and 18, more preferably between 10 and 16 even more preferably between 12 and 15 carbon atoms in the alcohol carbon chain, and on average between 5 and 12, preferably between 6 and 10, more preferably between 7 and 8 ethoxy units in the ethoxylation chain.

[0032] Preferably, the weight ratio of alkoxyated alkyl sulphate to linear alkylbenzene sulphonate is from 2:1 to 1:8 preferably from 1:1 to 1:5 most preferably from 1:1.25 to 1:4.

[0033] Preferably, the weight ratio of non-soap anionic surfactant to non-ionic surfactant is from 1:1 to 40:1, preferably from 1:1 to 20:1, more preferably from 1.3:1 to 15:1, even more preferably from 1.5:1 to 10:1.

[0034] Preferably, the laundry detergent composition comprises between 10% and 60%, preferably between 12% and 50%, most preferably between 15% and 40% by weight of the laundry detergent composition of a non-aqueous solvent. Preferably, the non-aqueous solvent is selected from 1,2-Propanediol, glycerol, sorbitol, dipropylene glycol, tripropylene glycol, polypropylene glycol or a mixture thereof.

[0035] Preferably, the water-soluble unit dose article comprises 15% or less by weight of the unit dose article of water, preferably the unit dose article comprises between 0.1% and 15%, more preferably between 1% and 12.5% by weight of the unit dose article of water.


[0036] The laundry detergent composition may comprise a polymer selected from amphiphilic graft copolymers, carboxymethyl cellulose, modified carboxymethylcellulose, polyester terephthalate polymers, hydroxyethylcellulose, modified hydroxyethylcellulose or a mixture thereof. Especially preferred are cationic modified hydroxyethylcellulose. Preferably, the laundry detergent composition comprises between 0.5% and 10%, preferably between 0.75% and 7%, more preferably between 1.5% and 5% by weight of the laundry detergent composition of the second polymer.

[0037] The laundry detergent composition may comprise between 0% and 10%, preferably between 0.1% and 7%, more preferably between 0.2% and 5% by weight of the laundry detergent composition of a fatty acid, a neutralised fatty acid soap or a mixture thereof.

[0038] Preferably, the water-soluble unit dose article comprises less than 3%, preferably less than 2% by weight of the laundry detergent composition of ethoxylated polyethyleneimine. The laundry detergent composition may comprise essentially no ethoxylated polyethyleneimine. Alternatively, the laundry detergent composition may comprise low levels of an ethoxylated polyethyleneimine. The laundry detergent composition may comprise between 0.01% and 3%, preferably between 0.01% and 2% by weight of the laundry detergent composition of an ethoxylated polyethyleneimine.

Zwitterionic polyamine

[0039] The laundry detergent composition comprises a zwitterionic polyamine. Preferably, the zwitterionic polyamine is selected from zwitterionic polyamines having the following formula :



R is C3-C20 preferably C5-C10 more preferably C6-C8 linear or branched alkylene, and mixtures thereof, most preferably linear C6.

R¹ is an anionic or partially anionic unit-capped polyalkyleneoxy unit having the formula $-(\text{R}2\text{O})_x\text{R}3$, wherein R2 is C2-C4 linear or branched alkylene, and mixtures thereof, preferably C2 or branched C3 and mixtures thereof, more preferably C2 (ethylene); R3 is hydrogen, an anionic unit, and mixtures thereof, in which not all R3 groups are hydrogen; x is from about 5 to about 50, preferably from about 10 to about 40, even more preferably from about 15 to about 30, most preferably from about 20 to about 25. A preferred value for x is 24, especially when R¹ comprises entirely ethyleneoxy units. Depending upon the method by which the formulator chooses to form the alkyleneoxy units, the wider or narrower the range of alkyleneoxy units present. The formulator will recognize that when ethoxylating a zwitterionic polyamine, only an average number or statistical distribution of alkyleneoxy units will be known. x values highlighted represent average values per polyalkoxy chain. Preferably the range of alkyleneoxy units within the zwitterionic polyamine is plus or minus two units, more preferably plus or minus one unit. Most preferably each R¹ group comprises about the same average number of alkyleneoxy units. Non-limiting examples of R3 anionic units include $-(\text{CH}_2)_p\text{CO}_2\text{M}$; $-(\text{CH}_2)_q\text{SO}_3\text{M}$; $-(\text{CH}_2)_q\text{OSO}_3\text{M}$; $-(\text{CH}_2)_q\text{CH}(\text{SO}_2\text{M})-\text{CH}_2\text{SO}_3\text{M}$; $-(\text{CH}_2)_q\text{CH}(\text{OSO}_2\text{M})\text{CH}_2\text{OSO}_3\text{M}$; $-(\text{CH}_2)_q\text{CH}(\text{SO}_3\text{M})\text{CH}_2\text{SO}_3\text{M}$; $-(\text{CH}_2)_p\text{PO}_3\text{M}$; $-\text{PO}_3\text{M}$; $-\text{SO}_3\text{M}$ and mixtures thereof; wherein M is hydrogen or a water soluble cation in sufficient amount to satisfy charge balance. Preferred anionic units are $-(\text{CH}_2)_p\text{CO}_2\text{M}$; $-\text{SO}_3\text{M}$, more preferably $-\text{SO}_3\text{M}$ (sulfonate group). The indices p and q are integers from 0 to 6, preferably 0 to 2, most preferably 0. For the purposes of the present invention, all M units, can either be a hydrogen atom or a cation depending upon the form isolated by the artisan or the relative pH of the system wherein the compound is used. Non-limiting examples of preferred cations include sodium, potassium, ammonium, and mixtures thereof.


Q is a quaternizing unit selected from the group consisting of C1-C30 linear or branched alkyl, C6-C30 cycloalkyl, C7-C30 substituted or unsubstituted alkylenearyl, and mixtures thereof, preferably C1-C30 linear or branched alkyl, even more preferably C1-C10 or even C1-C5 linear or branched alkyl, most preferably methyl; the degree of quaternization preferably is more than 50%, more preferably more than 70%, even more preferably more than 90%, most preferably about 100%.

X is an anion present in sufficient amount to provide electronic neutrality, preferably a water soluble anion selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride. To a great degree, the counter ion X will be derived from the unit which is used to perform the quaternization. For example, if methyl chloride is used as the quaternizing agent, chlorine (chloride ion) will be the counter ion X. Bromine (bromide ion) will be the dominant counter ion in the case where benzyl bromide is the quaternizing reagent.

n is from 0 to 4, preferably 0 to 2, most preferably 0.

[0040] Preferably from about 10% to about 100%, more preferably from about 20% to about 70%, even more preferably from 30% to about 50%, most preferably from about 35% to about 45% of the R3 groups are an anionic unit, preferably a sulfonate unit, the remaining R3 units being hydrogen.

[0041] Particularly preferred zwitterionic polyamines are zwitterionic hexamethylene diamines according to the following formula:



R is an anionic or partially anionic unit-capped polyalkyleneoxy unit having the formula: $-(\text{R}2\text{O})_x\text{R}3$ wherein R2 is C2-C4 linear or branched alkylene, and mixtures thereof, preferably C2 or branched C3 and mixtures thereof, even more preferably C2 (ethylene); R3 is hydrogen, an anionic unit, and mixtures thereof, in which not all R3 groups are hydrogen; x is from about 5 to about 50, preferably from about 10 to about 40, even more preferably from about 15

to about 30, most preferably from about 20 to about 25. A preferred value for x is 24, especially when R comprises entirely ethyleneoxy units. Depending upon the method by which the formulator chooses to form the alkyleneoxy units, the wider or narrower the range of alkyleneoxy units present. The formulator will recognize that when ethoxylating a zwitterionic polyamine, only an average number or statistical distribution of alkyleneoxy units will be known. x values highlighted represent average values per polyalkoxy chain. Preferably the range of alkyleneoxy units within the zwitterionic polyamine is plus or minus two units, more preferably plus or minus one unit. Most preferably each R group comprises about the same average number of alkyleneoxy units. Non-limiting examples of R3 anionic units include $-(CH_2)_pCO_2M$; $-(CH_2)_qSO_3M$; $-(CH_2)_qOSO_3M$; $-(CH_2)_qCH(SO_2M)-CH_2SO_3M$; $-(CH_2)_qCH(OSO_2M)CH_2OSO_3M$; $-(CH_2)_qCH(SO_3M)CH_2SO_3M$; $-(CH_2)_pPO_3M$; $-PO_3M$; $-SO_3M$ and mixtures thereof; wherein


M is hydrogen or a water soluble cation in sufficient amount to satisfy charge balance. Preferred anionic units are $-(CH_2)_pCO_2M$; $-SO_3M$, more preferably $-SO_3M$ (sulfonate group). The indices p and q are integers from 0 to 6, preferably 0 to 2, most preferably 0. For the purposes of the present invention, all M units, can either be a hydrogen atom or a cation depending upon the form isolated by the artisan or the relative pH of the system wherein the compound is used. Non-limiting examples of preferred cations include sodium, potassium, ammonium, and mixtures thereof.

Q is a quaternizing unit selected from the group consisting of C1-C30 linear or branched alkyl, C6-C30 cycloalkyl, C7-C30 substituted or unsubstituted alkylenearyl, and mixtures thereof, preferably C1-C30 linear or branched alkyl, even more preferably C1-C10 or even C1-C5 linear or branched alkyl, most preferably methyl; the degree of quaternization preferably is more than 50%, more preferably more than 70%, even more preferably more than 90%, most preferably about 100%.

X is an anion present in sufficient amount to provide electronic neutrality, preferably a water soluble anion selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride. To a great degree, the counter ion X will be derived from the unit which is used to perform the quaternization. For example, if methyl chloride is used as the quaternizing agent, chlorine (chloride ion) will be the counter ion X. Bromine (bromide ion) will be the dominant counter ion in the case where benzyl bromide is the quaternizing reagent.

[0042] Preferably from about 10% to about 100%, more preferably from about 20% to about 70%, even more preferably from 30% to about 50%, most preferably from about 35% to about 45% of the R3 groups are an anionic unit, preferably a sulfonate unit, the remaining R3 units being hydrogen.

[0043] Most preferred compound is the zwitterionic hexamethylene diamine represented by the following formula:



in which approximately 40% of the polyethoxy groups are sulfonated, the remaining polyethoxy groups being hydrogen capped. The degree of quaternization preferably is more than 90%, most preferably about 100%. Preferably the water soluble counter-anion is selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride.

[0044] The described zwitterionic polyamines can be made using techniques previously described in the art, and as such those skilled in the art would understand how to produce such compounds. The polyamine is first alkoxyated for example ethoxylated with ethylene oxide, followed by a quaternization step for example by reacting the alkoxyated polyamine with dimethylsulfate, and finally an anionic group substitution step for example by reacting the quaternized alkoxyated polyamine with chlorosulfonic acid.

Water-soluble film

[0045] The film of the present invention is soluble or dispersible in water. The water-soluble film preferably has a thickness of from 20 to 150 micron, preferably 35 to 125 micron, even more preferably 50 to 110 micron, most preferably about 76 micron.

[0046] Preferably, the film has a water-solubility of at least 50%, preferably at least 75% or even at least 95%, as measured by the method set out here after using a glass-filter with a maximum pore size of 20 microns:

5 grams $\pm$ 0.1 gram of film material is added in a pre-weighed 3L beaker and 2L $\pm$ 5ml of distilled water is added.

This is stirred vigorously on a magnetic stirrer, Labline model No. 1250 or equivalent and 5 cm magnetic stirrer, set at 600 rpm, for 30 minutes at 30°C. Then, the mixture is filtered through a folded qualitative sintered-glass filter with a pore size as defined above (max. 20 micron). The water is dried off from the collected filtrate by any conventional method, and the weight of the remaining material is determined (which is the dissolved or dispersed fraction). Then, the percentage solubility or dispersability can be calculated.

[0047] Preferred film materials are preferably polymeric materials. The film material can, for example, be obtained by casting, blow-moulding, extrusion or blown extrusion of the polymeric material, as known in the art.

[0048] Preferred polymers, copolymers or derivatives thereof suitable for use as pouch material are selected from polyvinyl alcohols, polyvinyl pyrrolidone, polyalkylene oxides, acrylamide, acrylic acid, cellulose, cellulose ethers, cellulose esters, cellulose amides, polyvinyl acetates, polycarboxylic acids and salts, polyaminoacids or peptides, polyamides, polyacrylamide, copolymers of maleic/acrylic acids, polysaccharides including starch and gelatine, natural gums such as xanthum and carragum. More preferred polymers are selected from polyacrylates and water-soluble acrylate copolymers, methylcellulose, carboxymethylcellulose sodium, dextrin, ethylcellulose, hydroxyethyl cellulose, hydroxypropyl methylcellulose, maltodextrin, polymethacrylates, and most preferably selected from polyvinyl alcohols, polyvinyl alcohol copolymers and hydroxypropyl methyl cellulose (HPMC), and combinations thereof. Preferably, the level of polymer in the pouch material, for example a PVA polymer, is at least 60%. The polymer can have any weight average molecular weight, preferably from about 1000 to 1,000,000, more preferably from about 10,000 to 300,000 yet more preferably from about 20,000 to 150,000.

[0049] Mixtures of polymers and/or copolymers can also be used as the pouch material, especially mixtures of polyvinylalcohol polymers and/or copolymers, especially mixtures of polyvinylalcohol homopolymers and/or anionic polyvinylalcohol copolymers preferably selected from sulphonated and carboxylated anionic polyvinylalcohol copolymers especially carboxylated anionic polyvinylalcohol copolymers. Most preferably the water soluble film comprises a blend of a polyvinylalcohol homopolymer and a carboxylated anionic polyvinylalcohol copolymer.

[0050] Preferred films exhibit good dissolution in cold water, meaning unheated distilled water. Preferably such films exhibit good dissolution at temperatures of 24°C, even more preferably at 10°C. By good dissolution it is meant that the film exhibits water-solubility of at least 50%, preferably at least 75% or even at least 95%, as measured by the method set out here after using a glass-filter with a maximum pore size of 20 microns, described above.

[0051] Preferred films are those supplied by Monosol under the trade references M8630, M8900, M8779, M8310.

[0052] The film may be opaque, transparent or translucent. The film may comprise a printed area.

[0053] The area of print may be achieved using standard techniques, such as flexographic printing or inkjet printing.

[0054] The film may comprise an aversive agent, for example a bittering agent. Suitable bittering agents include, but are not limited to, naringin, sucrose octaacetate, quinine hydrochloride, denatonium benzoate, or mixtures thereof. Any suitable level of aversive agent may be used in the film. Suitable levels include, but are not limited to, 1 to 5000ppm, or even 100 to 2500ppm, or even 250 to 2000rpm.

Process of washing

[0055] A further aspect of the present invention is a process for washing fabrics comprising the steps of;

a. Combining a water-soluble unit dose article according to the present invention with sufficient water to dissolve the water-soluble film and dilute the laundry detergent composition by a factor of between 300 and 800 fold to form a wash liquor;

b. Combining the wash liquor with at least one fabric to be washed.

[0056] The wash process may be conducted in a hand wash operation, an automatic wash machine operation or a mixture thereof.

Process of making

[0057] Those skilled in the art will know how to make the unit dose article and laundry detergent composition of the present invention using known techniques in the art.


[0058] During manufacture, a first water-soluble film may be shaped to comprise an open compartment into which the detergent composition is added. A second water-soluble film is then laid over the first film in such an orientation as to close the opening of the compartment. The first and second films are then sealed together along a seal region using known sealing means such as solvent, heat or a mixture thereof.

Particulate stain removal wash test:

[0063] Each stain was duplicated within each wash cycle (2 internal replicates). This test has been repeated 3 times for each test leg (3 external replicates) and the resulting 6 individual stain removal index results were averaged and reported.

Formula compositions :

[0064] The following water-soluble unit dose laundry compositions were prepared through mixing of the individual components. 18.5g of Base product was added via a dosing ball in each wash test. Lutensol FP620 (ethoxylated polyimide polymer ex BASF - PEI600EO20) and Lutensit Z96 (zwitterionic polyamine ex BASF - zwitterionic hexamethylene diamine according to below formula : 100% quaternized and about 40% of the polyethoxy (EO24) groups are sulfonated) were added on top as 1% solutions in water adjusted to pH 7.5, prepared as described below. Fatty acid was directly added on top of the base product.



Preparation of 1% active solutions:

[0065]

- Lutensol FP620: Weigh 12.69g of 78.8% active Lutensol FP620 into a 600ml beaker, add 400ml deionised water and adjust the pH to 7.50 using a pH meter and 0.05N citric acid. Transfer the resulting solution to a 1000ml flask and add deionized water till 1000ml.
- Lutensit Z96: Weigh 14.49g of 69% active Lutensit Z96 into a 600ml beaker, add 400ml deionised water and adjust the pH to 7.50 using a pH meter and 0.05N citric acid. Transfer the resulting solution to a 1000ml flask and add deionized water till 1000ml.

EP 3 342 847 A1

Base product:

	Base product
	wt%
Propylene 1,2-diol	8.394
Sodium Laureth Sulfate (68% in water)	23.975
Brightener 49	5.588
Citric acid (50% in water)	1.582
DiPropyleneGlycol	16.780
Glycerin	3.598
Editronic Acid (66% in water)	2.172
Linear alkylbenzene sulphonic acid	17.979
Potassium Sulfite (45% in water)	0.269
Mannanase enzyme	0.346
Monoethanolamine	6.622
Nonionic Surfactant (C24EO7)	1.199
Polymer Sokalan PG101	5.109
Perfume	2.613
Termamyl Ultra Amylase enzyme	0.157
Protease enzyme	1.660
Everest 200L Amylase enzyme	0.157
Water	1.799

Addition of Actives:

	Comparative example 1	Example 1	Comparative example 2	Example 2
Material	Base + Lutensol FP620 nil FA	Base + Lutensit Z96 nil FA	Base + Lutensol FP620 with FA	Base + Lutensit Z96 with FA
	Gram per wash	Gram per wash	Gram per wash	Gram per wash
Lutensol FP620	0.6 (added as 60ml of 1% solution)	-	0.6 (added as 60ml of 1% solution)	-
Lutensit Z96	-	0.6 (added as 60ml of 1% solution)	-	0.6 (added as 60ml of 1% solution)
Topped Palm Kernel Fatty Acid (FA)	-	-	0.9	0.9

Test results :

[0066] The particulate stain removal data summarized in table 1 and 2 below show the example compositions according to the invention comprising both the alkyl ethoxy sulphate anionic surfactant and the zwitterionic polyamine to be more effective in particulate soil removal than comparative example formulations outside the scope of the invention single variable differing in soil removal polymer chemistry. The effect has been shown within formulations not comprising a fatty acid (table 1) as well as for compositions comprising a fatty acid (table 2).

Table 1 : % Particulate stain removal in nil fatty acid nonaqueous laundry liquid formulation

Soil	Comparative Example 1	Example 1	Difference in stain removal	% Improvement in stain removal
Black todd clay	56.4	61.1	4.7	8.3
Brown silica sand	15.0	22.8	7.8	52.0
Grass	81.7	83.3	1.6	2.0
Hoover dust	53.1	59.4	6.3	11.9
Humax Peat	30.6	34.5	3.9	12.7
NTC clay	33.2	37.8	4.6	13.9
Stanley clay	36.1	40.6	4.5	12.5
<i>Average</i>	<i>43.7</i>	<i>48.5</i>	<i>4.8</i>	<i>16.2</i>


Table 2 : % Particulate stain removal in fatty acid comprising nonaqueous laundry liquid formulation

Soil	Comparative Example 2	Example 2	Difference in stain removal	% Improvement in stain removal
Black todd clay	65.0	65.9	0.9	1.4
Brown silica sand	29.0	33.5	4.5	15.5
Grass	86.6	86.7	0.1	0.1
Hoover dust	68.8	71.8	3.0	4.4
Humax Peat	45.0	47.5	2.5	5.6
NTC clay	46.3	49.8	3.5	7.6
Stanley clay	46.7	49.0	2.3	4.9
<i>Average</i>	<i>55.3</i>	<i>57.7</i>	<i>2.4</i>	<i>5.6</i>

[0067] The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm."

Claims

1. A water-soluble unit dose article comprising a water-soluble film and a laundry detergent composition, wherein the laundry detergent composition is preferably selected from a liquid, a solid or a mixture thereof, and wherein said detergent composition comprises;
 - a. a zwitterionic polyamine, preferably between 0.01% to about 20%, preferably from 0.1% to 10%, more preferably from 0.5% to 7%, even more preferably from 1% to 5%, most preferably from 2% to 4% by weight of the laundry detergent composition of the zwitterionic polyamine; and
 - b. a first non-soap anionic surfactant selected from an alkyl sulphate anionic surfactant, an alkoxyated alkyl sulphate anionic surfactant or a mixture thereof, preferably wherein the laundry detergent composition comprises between 2% and 30%, preferably between 5% and 25%, more preferably between 7% and 20%, most preferably between 10% and 17% by weight of the laundry detergent composition of the first non-soap surfactant.
2. The water-soluble unit dose article according to claim 1 wherein said zwitterionic polyamine is selected from zwitterionic polyamines according to the following formula;



wherein:

R is C3-C20 preferably C5-C10 more preferably C6-C8 linear or branched alkylene, and mixtures thereof, most preferably linear C6;

R¹ is an anionic unit-capped polyalkyleneoxy unit having the formula : -(R2O)_xR3,


wherein R2 is C2-C4 linear or branched alkylene, and mixtures thereof, preferably C2 or branched C3 and mixtures thereof, more preferably C2 (ethylene); R3 is hydrogen, an anionic unit, and mixtures thereof, in which not all R3 groups are hydrogen, preferably wherein R3 anionic units are selected from -(CH2)_pCO2M; -(CH2)_qSO3M; - (CH2)_qOSO3M; -(CH2)_qCH(SO2M)-CH2SO3M; -(CH2)_qCH(OSO2M)CH2OSO3M; -(CH2)_qCH(SO3M)CH2SO3M; -(CH2)_pP03M; -P03M; -SO3M and mixtures thereof; wherein M is hydrogen or a water soluble cation, preferably selected from sodium, potassium, ammonium, and mixtures thereof and in sufficient amount to satisfy charge balance; x is from 5 to 50, preferably from 10 to 40, even more preferably from 15 to 30, most preferably from 20 to 25;

Q is a quaternizing unit selected from the group consisting of C1-C30 linear or branched alkyl, C6-C30 cycloalkyl, C7-C30 substituted or unsubstituted alkylenearyl, and mixtures thereof, preferably C1-C30 linear or branched alkyl, even more preferably C1-C10 or even C1-C5 linear or branched alkyl, most preferably methyl; the degree of quaternization preferably is more than 50%, more preferably more than 70%, even more preferably more than 90%, most preferably about 100%.

X is an anion present in sufficient amount to provide electronic neutrality, preferably a water soluble anion selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride;

n is from 0 to 4, preferably 0 to 2, most preferably 0.

3. The water-soluble unit dose article according to claim 2 wherein said zwitterionic polyamine is selected from zwitterionic polyamines according to the following formula;



wherein

R is an anionic unit-capped polyalkyleneoxy unit having the formula: -(R2O)_xR3 wherein R2 is C2-C4 linear or branched alkylene, and mixtures thereof, preferably C2 or branched C3 and mixtures thereof, even more preferably C2 (ethylene);

R3 is hydrogen, an anionic unit, and mixtures thereof, in which not all R3 groups are hydrogen, preferably wherein R3 anionic units include -(CH2)_pCO2M; -(CH2)_qSO3M; - (CH2)_qOSO3M; -(CH2)_qCH(SO2M)-CH2SO3M; -(CH2)_qCH(OSO2M)CH2OSO3M; - (CH2)_qCH(SO3M)CH2SO3M; -(CH2)_pP03M; -P03M; -SO3M; and mixtures thereof wherein p and q are integers from 0 to 6, preferably 0 to 2, most preferably 0; and wherein M is hydrogen or a water soluble cation in sufficient amount to satisfy charge balance, preferably selected from sodium, potassium, ammonium, and mixtures thereof and in sufficient amount to satisfy charge balance; x is from 5 to 50, preferably from 10 to 40, even more preferably from 15 to 30, most preferably from 20 to 25; x is from about 5 to about 50, preferably from about 10 to about 40, even more preferably from about 15 to about 30, most preferably from about 20 to about 25. ;


Q is a quaternizing unit selected from the group consisting of C1-C30 linear or branched alkyl, C6-C30 cycloalkyl, C7-C30 substituted or unsubstituted alkylenearyl, and mixtures thereof, preferably C1-C30 linear or branched alkyl, even more preferably C1-C10 or even C1-C5 linear or branched alkyl, most preferably methyl; the degree of quaternization preferably is more than 50%, more preferably more than 70%, even more preferably more

than 90%, most preferably about 100%;

X is an anion present in sufficient amount to provide electronic neutrality, preferably a water soluble anion selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride;

wherein preferably from about 10% to about 100%, more preferably from about 20% to about 70%, even more preferably from 30% to about 50%, most preferably from about 35% to about 45% of the R3 groups are an anionic unit, preferably a -SO₃M, the remaining R3 units being hydrogen.

4. The water-soluble unit dose article according to claim 4, wherein said zwitterionic polyamine is selected from zwitterionic polyamines according to the following formula;



wherein from about 20% to about 70%, preferably from 30% to about 50%, more preferably from about 35% to about 45%, most preferably about 40% of the polyethoxy groups are sulfonated, the remaining polyethoxy groups being hydrogen capped;

the degree of quaternization preferably is more than 90%, most preferably about 100%

and;

preferably the water soluble counter-anion is selected from the group consisting of chlorine, bromine, iodine, methylsulfate, and mixtures thereof, more preferably chloride.

5. The water-soluble unit dose article according to any preceding claims wherein the first non-soap surfactant is preferably a C10-18 alkyl sulphate, a C10-18 alkoxyated alkyl sulphate or a mixture thereof.

6. The water-soluble unit dose article according to any preceding claims, wherein the first non-soap surfactant is an alkoxyated alkyl sulphate, preferably an ethoxyated alkyl sulphate with an average degree of ethoxylation of between 0.5 and 7, preferably between 1 and 5, more preferably between 2 and 4, most preferably about 3, or wherein the first non-soap surfactant is a mixture of one or more alkoxyated alkyl sulphates, preferably ethoxyated alkyl sulphates, and optionally an alkyl sulphate, the mixture having an average degree of ethoxylation of between 0.5 and 7, preferably between 1 and 5, more preferably between 2 and 4, most preferably about 3.

7. The water-soluble unit dose article according to any preceding claims comprising a second non-soap anionic surfactant, wherein the second non-soap anionic surfactant is selected from linear alkylbenzene sulphonate, preferably wherein the laundry detergent composition comprises between 2% and 30%, preferably between 5% and 27%, more preferably between 7% and 25%, most preferably between 10% and 23% or even between 10% and 17% by weight of the laundry detergent composition of the second non-soap surfactant.

8. The water-soluble unit dose article according to any preceding claims comprising between 0% and 10% preferably between 0.01% and 8%, more preferably between 0.1% and 6%, most preferably between 0.15% and 4% by weight of the laundry detergent composition of a non-ionic surfactant, preferably wherein the non-ionic surfactant is selected from fatty alcohol alkoxyate, an oxo-synthesised fatty alcohol alkoxyate, Guerbet alcohol alkoxyates, alkyl phenol alcohol alkoxyates or a mixture thereof.

9. The water-soluble unit dose article according to any preceding claims comprising between 10% and 45%, preferably between 12% and 42%, most preferably between 15% and 40% by weight of the laundry detergent composition of a non-aqueous solvent, preferably wherein the non-aqueous solvent is selected from 1,2-Propanediol, glycerol, sorbitol, dipropylene glycol, tripropyleneglycol, polypropylene glycol or a mixture thereof.

10. The water-soluble unit dose article according to any preceding claims comprising 15% or less by weight of the unit dose article of water, preferably comprising between 0.1% and 15%, more preferably between 1% and 12.5% by weight of the unit dose article of water.

- 11.** The water-soluble unit dose article according to any preceding claims wherein the water-soluble film is a polymeric

film, preferably comprising polyvinyl alcohols, polyvinyl pyrrolidone, polyalkylene oxides, acrylamide, acrylic acid, cellulose, cellulose ethers, cellulose esters, cellulose amides, polyvinyl acetates, polycarboxylic acids and salts, polyaminoacids or peptides, polyamides, polyacrylamide, copolymers of maleic/acrylic acids, polysaccharides or a mixture thereof, preferably comprising a polyvinyl alcohol.

- 5
10
15
20
25
30
35
40
45
50
55
12. The water-soluble unit dose article according to any preceding claims comprising a polymer selected from amphiphilic graft copolymers, carboxymethyl cellulose, modified carboxymethylcellulose, polyester terephthalate polymers, hydroxyethylcellulose, modified hydroxyethylcellulose or a mixture thereof, preferably cationically modified hydroxyethylcellulose; and preferably wherein the laundry detergent composition comprises 0.5% and 10%, preferably between 0.75% and 7%, more preferably between 1.5% and 5% by weight of the laundry detergent composition of the polymer.
13. The water-soluble unit dose article according to any preceding claims comprising at least two compartments, preferably at least three compartments, wherein the laundry detergent composition is comprised within at least one compartment, preferably wherein the laundry detergent composition is a culmination of ingredients located within all the compartments of the unit dose article.
14. The water-soluble unit dose article according to any preceding claims wherein the water-soluble film comprises polyvinyl alcohol polymer or copolymer, preferably a blend of polyvinylalcohol polymers and/or polyvinylalcohol copolymers, preferably selected from sulphonated and carboxylated anionic polyvinylalcohol copolymers especially carboxylated anionic polyvinylalcohol copolymers, most preferably a blend of a polyvinylalcohol homopolymer and a carboxylated anionic polyvinylalcohol copolymer.
15. A process for washing fabrics comprising the steps of;
 - a. Combining a water-soluble unit dose article according to any preceding claims with sufficient water to dissolve the water-soluble film and dilute the laundry detergent composition by a factor of between 300 and 800 fold to form a wash liquor;
 - b. Combining the wash liquor with at least one fabric to be washed.




Fig. 1



EUROPEAN SEARCH REPORT

 Application Number
 EP 16 20 7069

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	EP 1 666 579 A1 (PROCTER & GAMBLE [US]) 7 June 2006 (2006-06-07) * paragraph [0032] - paragraph [0033]; claims 1-9 *	1-15	INV. C11D17/04 C11D3/37 C11D1/14 C11D1/29
Y	US 2012/077725 A1 (WANG XIAORU JENNY [US] ET AL) 29 March 2012 (2012-03-29) * paragraph [0198] * * paragraph [0234] *	1-15	
Y	WO 2007/111888 A1 (PROCTER & GAMBLE [US]; PANANDIKER RAJAN KESHAV [US]; VETTER KERRY ANDR) 4 October 2007 (2007-10-04) * page 30; example 1 *	1-15	
			TECHNICAL FIELDS SEARCHED (IPC)
			C11D
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 7 June 2017	Examiner Richards, Michael
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

 1
 EPO FORM 1503 03.02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 16 20 7069

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-06-2017

10

Patent document
cited in search report

Publication
date

Patent family
member(s)

Publication
date

15

EP 1666579

A1

07-06-2006

AT 437215 T 15-08-2009

BR PI0517995 A 21-10-2008

CA 2586052 A1 01-06-2006

EP 1666579 A1 07-06-2006

ES 2326818 T3 20-10-2009

JP 4672730 B2 20-04-2011

JP 2008518861 A 05-06-2008

US 2006276364 A1 07-12-2006

WO 2006057905 A1 01-06-2006

20

US 2012077725

A1

29-03-2012

BR 112013004895 A2 03-05-2016

CA 2811011 A1 29-03-2012

CN 103732730 A 16-04-2014

EP 2619299 A2 31-07-2013

JP 2013541649 A 14-11-2013

US 2012077725 A1 29-03-2012

WO 2012040131 A2 29-03-2012

25

WO 2007111888

A1

04-10-2007

AT 530628 T 15-11-2011

AT 530629 T 15-11-2011

AT 530630 T 15-11-2011

BR PI0709024 A2 21-06-2011

BR PI0709036 A2 21-06-2011

BR PI0709037 A2 21-06-2011

BR PI0709064 A2 21-06-2011

CA 2642950 A1 04-10-2007

CA 2642955 A1 04-10-2007

CA 2642958 A1 04-10-2007

CA 2642962 A1 04-10-2007

CA 2642970 A1 04-10-2007

CN 101405378 A 08-04-2009

CN 101405380 A 08-04-2009

CN 101405381 A 08-04-2009

CN 101405383 A 08-04-2009

EP 1996687 A2 03-12-2008

EP 1996688 A2 03-12-2008

EP 1996689 A2 03-12-2008

EP 1996692 A1 03-12-2008

EP 1999243 A2 10-12-2008

EP 2426192 A1 07-03-2012

ES 2376125 T3 09-03-2012

ES 2376264 T3 12-03-2012

ES 2376365 T3 13-03-2012

ES 2442868 T3 14-02-2014

JP 4955053 B2 20-06-2012

JP 5461171 B2 02-04-2014

30

35

40

45

50

FORM P0459

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 16 20 7069

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-06-2017

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		JP 5586945 B2	10-09-2014
		JP 5586946 B2	10-09-2014
		JP 2009530478 A	27-08-2009
		JP 2009530479 A	27-08-2009
		JP 2009530481 A	27-08-2009
		JP 2009530482 A	27-08-2009
		JP 2009530512 A	27-08-2009
		MX 294406 B	06-01-2012
		MX 297648 B	29-03-2012
		MX 319061 B	03-04-2014
		MX 319376 B	11-04-2014
		RU 2008133485 A	27-04-2010
		RU 2008133486 A	27-04-2010
		RU 2008133487 A	27-04-2010
		RU 2008133488 A	27-04-2010
		US 2009069206 A1	12-03-2009
		US 2009069207 A1	12-03-2009
		US 2009088363 A1	02-04-2009
		US 2009186797 A1	23-07-2009
		US 2009209445 A1	20-08-2009
		US 2011034366 A1	10-02-2011
		WO 2007111887 A2	04-10-2007
		WO 2007111888 A1	04-10-2007
		WO 2007111892 A2	04-10-2007
		WO 2007111898 A2	04-10-2007
		WO 2007111899 A2	04-10-2007

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82