

(19)



(11)

EP 1 758 867 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
29.09.2010 Bulletin 2010/39

(51) Int Cl.:
C07D 223/02 ^(2006.01) **C07D 223/28** ^(2006.01)

(21) Application number: **05708576.3**

(86) International application number:
PCT/IB2005/000452

(22) Date of filing: **21.02.2005**

(87) International publication number:
WO 2005/092862 (06.10.2005 Gazette 2005/40)

(54) PROCESS FOR PREPARING OXCARBAZEPINE

VERFAHREN ZUR HERSTELLUNG VON OXCARBAZEPIN

PROCEDE DE PREPARATION D'OXCARBAZEPINE

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**

(30) Priority: **09.03.2004 IT IT20040452**

(43) Date of publication of application:
07.03.2007 Bulletin 2007/10

(73) Proprietor: **Archimica S.r.l.**
20124 Milano (IT)

(72) Inventors:

- **BANFI, Aldo**
I-20139 Milano (IT)
- **BOLLINI, Deborah**
I-24058 ROMANO DIE LOMBARDIA BG (IT)
- **SERRA, Maurizio**
I-20062 CASSANO D'ADDA MI (IT)
- **DI LERNIA, Gianluca**
I-20139 Milano (IT)

(74) Representative: **Pistolesi, Roberto et al**
Dragotti & Associati Srl
Via Marina 6
20121 Milano (IT)

(56) References cited:
DE-A1- 2 011 087

- **DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; HAASZ, FERENC ET AL: "Improved process for producing 5-carbamoyl-10-oxo-10,11-dihydro-5H- dibenz[b,f]azepine" XP002306691 retrieved from STN Database accession no. 1994:164010 cited in the application & HU 63 389 A2 (ALKALOIDA VEGYESZETI GYAR, HUNG.) 30 August 1993 (1993-08-30)**
- **HEINER ECKERT ET AL: "Triphosgene, a Crystalline Phosgene Substitute" ANGEWANDTE CHEMIE. INTERNATIONAL EDITION, VERLAG CHEMIE. WEINHEIM, DE, vol. 26, no. 9, 1987, pages 894-895, XP002083416 ISSN: 0570-0833**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).


EP 1 758 867 B1

Description

[0001] The present invention relates to a novel process for preparing oxcarbazepine, which is particularly advantageous from an industrial point of view, characterized by the use of triphosgene as chlorocarbonylating agent.

PRIOR ART


[0002] Oxcarbazepine (Merck Index, 1996, No. 7063) of formula



(I)

is a known anticonvulsant agent, which was described for the first time in German patent DE 2 011 087 (Ciba-Geigy).

[0003] Various processes are reported in the literature for preparing oxcarbazepine, these processes essentially involving the introduction of the carboxamide function onto the nitrogen in position 5 by means of carbamoylation with cyanates, for example in WO 01/56992, EP 1 302 464, IT 1 318 371 and WO 96/21649, or, as illustrated in Scheme 1 below and in US 3 642 775 and HU63389, by chlorocarbonylation (a) followed by ammonolysis (b) and final hydrolysis (c):

SCHEME 1

[0004] In particular, US 3 642 775 describes a chlorocarbonylation reaction (a) performed with phosgene, in toluene at 95°C, with a yield of 77% of compound II after crystallization from ethanol, and subsequent ammonolysis (b) with ammonia gas in ethanol, with a yield of 73% of compound IV, also calculated after crystallization from ethanol (56% overall yield for the two steps a and b).

[0005] The final deprotection of compound IV to give oxcarbazepine (c) is performed by refluxing with 2N hydrochloric

acid, in a yield of 80%, after crystallization from ethanol. The overall yield for this process (II --> III --> IV --> I) is equal to about 45%.


[0006] However, the use of phosgene as chlorocarbonylating agent in this synthetic route represents a considerable drawback on account of its high toxicity and corrosiveness.

[0007] From the abstract in Chemical Abstracts of Hungarian patent HU63389, the same sequence as that of Scheme 1 is performed, but with diphosgene in refluxing toluene (a), subsequent ammonolysis of the intermediate III, which is not isolated, with ammonia gas (b) (yield for steps a) and b) equal to 58.9%) and, finally, hydrolysis of the intermediate IV with 2M HCl, in a yield equal to 73.5%. In this case, the overall yield for the process, calculated on the basis of the partial values reported in the abstract, is equal to 43.3%.

[0008] We have now found, surprisingly, an industrially acceptable process for preparing oxcarbazepine, which, without using corrosive reagents such as phosgene, makes it possible to obtain the desired product while significantly increasing the overall yields.

DESCRIPTION OF THE INVENTION


[0009] One subject of the present invention is thus a process for preparing oxcarbazepine of formula



(I)


which includes:

a) the chlorocarbonylation reaction of the compound of formula




(II)

with triphosgene in the presence of a base, to give the compound of formula



(III)

b) ammonolysis of the compound of formula III to give the compound of formula



and

c) acid hydrolysis of the compound of formula IV to give oxcarbazepine I.

[0010] This process may be performed via the isolation and the optional purification of the individual intermediates of formulae II and III, or, preferably, by minimizing this procedure, i.e. working directly on the crude reaction product from the partially worked-up preceding step, as illustrated in the experimental section.

[0011] The starting compound of formula II of the process that is the subject of the present invention is commercially available.

[0012] The chlorocarbonylation step a) is performed with triphosgene in a molar ratio, relative to the compound of formula II, preferably of between 0.46:1 and 0.54:1 and more preferably at about 0.5:1, in the presence of a base, preferably an organic base and more preferably triethylamine, in a molar ratio relative to the compound of formula II of between 1.4:1 and 1.6:1 and preferably at about 1.5:1.

[0013] The reaction solvent is generally chosen from aromatic hydrocarbons, preferably toluene, and the reaction temperature is usually between 90 and 110°C; preferably, the temperature at the start of dropwise addition of the triphosgene is greater than 90°C, to subsequently reach the reflux temperature of toluene during the dropwise addition.

[0014] Preferably, once step a) is complete, the reaction medium is subjected to a minimum work-up and the residual crude product, obtained by evaporation, is used directly in the subsequent step b).

[0015] This ammonolysis step is generally performed with ammonia, preferably aqueous ammonia, in a suitable solvent, preferably an alcohol and more preferably methanol.

[0016] The subsequent deprotection step c) is preferably performed on the concentrate obtained directly from the preceding step, under acidic conditions, preferably with hydrochloric acid, in aqueous medium at a pH of between 0 and 2, preferably at about pH 1, and at a temperature above 50°C, preferably between 90 and 95°C.

[0017] The final product is preferably purified by crystallization, more preferably by crystallization from dimethylacetamide/methanol.

[0018] The overall yield for the present process is generally about 80%, i.e. appreciably greater than the yields for the similar processes described in the art.

[0019] The following examples are now given for the purpose of better illustrating the present invention without, however, limiting it.

EXPERIMENTAL SECTION

EXAMPLE 1

Preparation of 10-methoxy-5H-dibenzo[b,f]azepine-5-carboxamide (IV)

[0020] 100 g (0.4479 mol) of 10-methoxy-5H-dibenz[b,f]azepine (II) are introduced into a round-bottomed flask, they are placed in suspension by adding 500 ml of toluene, 67.2 g (0.6641 mol) of triethylamine are added and the mixture is heated to 90°C. A solution consisting of 66 g of triphosgene (0.2224 mol) in 150 ml of toluene is then added dropwise over 40 minutes, while allowing the temperature to rise to 110°C.

[0021] Once the dropwise addition is complete, HPLC monitoring indicates that the reaction is complete. The reaction mixture is cooled to 80°C and 300 ml of water are then added dropwise. After the dropwise addition, the internal temperature of the mixture is 60-65°C. The phases are separated at a temperature above 30-35°C and the aqueous phase is removed.

[0022] The organic phase is evaporated to dryness under vacuum at about 40°C and is then taken up in 125 ml of methanol. This solution is evaporated under vacuum at 40°C and the residue is taken up again in 500 ml of methanol.

[0023] The solution is heated to 50°C, 300 ml of 28% aqueous ammonia (2.11 mol) are added dropwise to the homogeneous solution, and the mixture is stirred at 50°C for 1 hour. HPLC monitoring after 1 hour indicates that the reaction

is complete, and the solvent is partially distilled off at 40°C under vacuum, down to a residual volume of about 400 ml. The solution at 25°C becomes turbid due to precipitation of 10-methoxy-5H-dibenzo[b,f]azepine-5-carboxamide (IV), and the suspension thus formed is used without further purification in the following step.

EXAMPLE 2

Preparation of oxcarbazepine (I)


[0024] 600 ml of water are added to the round-bottomed flask containing the suspension of 10-methoxy-5H-dibenzo[b,f]azepine-5-carboxamide (IV) from Example 1, and about 12 g of 37% HCl are added dropwise to pH = 1. The suspension is stirred at about 95°C for 4 hours.

[0025] The mixture is cooled to 25°C and about 14 g of 30% NaOH are added dropwise to bring the pH from 1.0 to 7.0-7.5. The reaction suspension is filtered at 25°C and the cake is washed twice with water (2 x 100 ml), the filtration and washing waters then being removed. The cake is washed three times with methanol (3 x 100 ml) at 12-14°C and the waters are then removed. The wet cake (160-170 g) is suspended in dimethylacetamide (400 ml) while heating at 110°C until the solid has dissolved. The product is filtered off at a temperature above 80°C (temperature of start of recrystallization). The filtrate is cooled to 60°C, 400 ml of methanol are then added to the suspension, and the resulting mixture is cooled first to 25°C and then to 0-5°C. After 1 hour at 0-5°C, the cake is washed with 150 ml of methanol at 12-14°C and the methanolic washing phase is removed. The cake is dried under vacuum at 40°C for 6 hours.

Weight: about 80 g;
 yield: 70% (molar), 80% (w/w).
 Purity (HPLC): > 99% (area %)

Claims


1. Process for preparing oxcarbazepine of formula



(I)


which includes:

a) the chlorocarbonylation reaction of the compound of formula



(II)


with triphosgene in the presence of a base, to give the compound of formula



(III)

2. Process according to Claim 1, which subsequently includes:

b) ammonolysis of the compound of formula III to give the compound of formula



(IV)


and

c) acid hydrolysis of the compound of formula IV to give oxcarbazepine I.

3. Process according to Claim 1 or 2, in which the said chlorocarbonylation reaction a) is performed with triphosgene in a molar ratio, relative to the compound of formula II, of between 0.46:1 and 0.54:1 and more preferably at about 0.5:1.
4. Process according to Claims 1 to 3, in which the said chlorocarbonylation reaction a) is performed using triethylamine as base, in a molar ratio relative to the compound of formula II of between 1.4: 1 and 1.6:1 and preferably at about 1.5:1.
5. Process according to Claims 1 to 4, in which the said chlorocarbonylation reaction a) is performed in toluene and at a temperature of between 90 and 110°C.
6. Process according to Claims 2 to 5, in which the ammonolysis b) is performed with aqueous ammonia in methanol.
7. Process according to Claims 2 to 6, in which the deprotection c) is performed with hydrochloric acid in aqueous medium at a pH of about 1 and at a temperature of between 90 and 95°C.

Patentansprüche

1. Verfahren zur Herstellung von Oxcarbazepin der Formel:




(I)

umfassend:

(a) die Chlorcarbonylierungs-Reaktion der Verbindung der Formel:




mit Triphosgen in Gegenwart einer Base, um die Verbindung der Formel



zu erhalten.

2. Verfahren gemäss Anspruch 1, anschliessend umfassend:

(b) Ammonolyse der Verbindung der Formel (III), um die Verbindung der Formel:



zu erhalten, und

(c) Säurehydrolyse der Verbindung der Formel (IV), um Oxcarbazepin I zu erhalten.

3. Verfahren gemäss Anspruch 1 oder 2, in dem die Chlorcarbonylierungs-Reaktion (a) mit Triphosgen in einem molaren Verhältnis, in bezug auf die Verbindung der Formel (II), zwischen 0,46:1 und 0,54:1 und stärker bevorzugt etwa 0,5:1 durchgeführt wird.

4. Verfahren gemäss Ansprüchen 1 bis 3, in dem die Chlorcarbonylierungs-Reaktion (a) unter Verwendung von Triethylamin als Base in einem molaren Verhältnis, in bezug auf die Verbindung der Formel (II), zwischen 1,4:1 und 1,6:1 und vorzugsweise etwa 1,5:1, durchgeführt wird.

5. Verfahren gemäss Ansprüchen 1 bis 4, in dem die Chlorcarbonylierungs-Reaktion (a) in Toluol und bei einer Temperatur zwischen 90 und 110°C durchgeführt wird.


6. Verfahren gemäss Ansprüchen 2 bis 5, in dem die Ammonolyse (b) mit wässrigem Ammoniak in Methanol durch-

geführt wird.

7. Verfahren gemäss Ansprüchen 2 bis 6, in dem die Entschützung (c) mit Salzsäure in einem wässrigen Medium bei einem pH-Wert von etwa 1 und bei einer Temperatur zwischen 90 und 95°C durchgeführt wird.

Revendications


1. Procédé de préparation d'une oxcarbazépine de formule suivante :



(I)


qui comprend :

- a) la réaction de chlorocarbonylation du composé de formule suivante :



(II)


avec du triphosgène en présence d'une base, de manière à obtenir le composé de formule suivante :



(III)

2. Procédé selon la revendication 1, qui comprend ensuite les étapes suivantes :

- b) l'ammonolyse du composé de formule III de manière à obtenir le composé de formule suivante :



(IV)

et

- c) l'hydrolyse acide du composé de formule IV, ce qui donne l'oxcarbazépine I.

EP 1 758 867 B1

3. Procédé selon la revendication 1 ou 2, dans lequel ladite réaction de chlorocarbonylation de l'étape a) est effectuée avec du triphosgène dans un rapport molaire, par rapport au composé de formule II, de 0,46:1 à 0,54:1 et, mieux encore, d'environ 0,5:1.
- 5 4. Procédé selon l'une quelconque des revendications 1 à 3, dans lequel ladite réaction de chlorocarbonylation de l'étape a) est effectuée en utilisant de la triéthylamine comme base, dans un rapport molaire, par rapport au composé de formule II, de 1,4: 1 à 1,6:1 et, de préférence, d'environ 1,5:1.
- 10 5. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel ladite réaction de chlorocarbonylation de l'étape a) est effectuée dans du toluène et à une température comprise entre 90 et 110 °C.
6. Procédé selon l'une quelconque des revendications 2 à 5, dans lequel l'ammonolyse de l'étape b) est effectuée avec de l'ammoniaque aqueuse dans du méthanol.
- 15 7. Procédé selon l'une quelconque des revendications 2 à 6, dans lequel la déprotection de l'étape c) est effectuée avec de l'acide chlorhydrique dans un milieu aqueux à un pH d'environ de 1 et à une température de 90 à 95 °C.

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- DE 2011087 [0002]
- WO 0156992 A [0003]
- EP 1302464 A [0003]
- IT 1318371 [0003]
- WO 9621649 A [0003]
- US 3642775 A [0003] [0004]
- HU 63389 [0003] [0007]