

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3043181 T3**

-
- (51) Int.Cl.: **G 01 N 33/574 (2006.01)** **A 61 K 39/00 (2006.01)** **C 07 K 16/28 (2006.01)**
C 07 K 16/30 (2006.01) **C 12 N 5/00 (2006.01)** **C 12 N 5/09 (2010.01)**
C 12 Q 1/68 (2018.01) **G 01 N 33/50 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-05-04**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-04-08**
- (86) Europæisk ansøgning nr.: **15198767.4**
- (86) Europæisk indleveringsdag: **2009-01-13**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-07-13**
- (30) Prioritet: **2008-01-15 US 11324 P**
- (62) Stamansøgningsnr: **09702107.5**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **The Board of Trustees of the Leland Stanford Junior University, Office of the General Counsel , Building 170, Third Floor, Main Quad , P.O. Box 20386, Stanford, CA 94305-2038, USA**
- (72) Opfinder: **WEISSMAN, Irving, L., 4147 Jefferson Avenue, Redwood City, CA 94061, USA**
MAJETI, Ravindra, Stanford Institute for Stem Cell Bio and, Regenerative Medicine, 1050 Arastradero Road, Building A, Stanford, CA 94304, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **MARKØRER AF AKUTTE MYELOID-LEUKÆMI-STAMCELLER**
- (56) Fremdragne publikationer:
EP-A1- 2 111 869
US-A1- 2004 213 792
US-A1- 2007 287 163
US-B1- 6 491 917
US-B2- 6 733 743
HOSEN N ET AL: "CD96 is a leukemic stem cell-specific marker in human acute myeloid leukemia", PNAS,, vol. 104, no. 26, 26 June 2007 (2007-06-26), pages 11008-11013, XP008109009, DOI: 10.1073/PNAS.0704271104
JAMIESON C ET AL: "Increased expression of CD47 is a constant marker in mouse and human myeloid leukemias", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 106, 1 January 2005 (2005-01-01), page 911A, XP008138745, ISSN: 0006-4971
CLARKE MICHAEL F ET AL: "Cancer stem cells--perspectives on current status and future directions: AACR Workshop on cancer stem cells", CANCER RESEARCH, AACR, US PHILADELPHIA, PA, vol. 66, no. 19, 1 October 2006 (2006-10-01), pages 9339-9344, XP002580692, ISSN: 1538-7445
YOSHIKANE KIKUSHIGE ET AL: "TIM-3 Is a Promising Target to Selectively Kill Acute Myeloid Leukemia Stem Cells", CELL STEM CELL, CELL PRESS, US, vol. 7, no. 6, 6 October 2010 (2010-10-06) , pages 708-717, XP028184581, ISSN: 1934-5909, DOI: 10.1016/J.STEM.2010.11.014 [retrieved on 2010-11-17]

Fortsættes ...

M. JAN ET AL: "Prospective separation of normal and leukemic stem cells based on differential expression of TIM3, a human acute myeloid leukemia stem cell marker", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 108, no. 12, 7 March 2011 (2011-03-07), pages 5009-5014, XP055029836, ISSN: 0027-8424, DOI: 10.1073/pnas.1100551108

JAISWAL SIDDHARTHA ET AL: "CD47 Is Upregulated on Circulating Hematopoietic Stem Cells and Leukemia Cells to Avoid Phagocytosis", CELL, CELL PRESS, US, vol. 138, no. 2, 1 July 2009 (2009-07-01), pages 271-285, XP009160236, ISSN: 0092-8674, DOI: 10.1016/J.CELL.2009.05.046

RAVINDRA MAJETI ET AL: "CD47 Is an Adverse Prognostic Factor and Therapeutic Antibody Target on Human Acute Myeloid Leukemia Stem Cells", CELL, vol. 138, no. 2, 1 July 2009 (2009-07-01), pages 286-299, XP055277135, US ISSN: 0092-8674, DOI: 10.1016/j.cell.2009.05.045

JOSÉ IGNACIO MARTIN-SUBERO ET AL: "Amplification of ERBB2, RARA, and TOP2A genes in a myelodysplastic syndrome transforming to acute myeloid leukemia", CANCER GENETICS AND CYTOGENETICS., vol. 127, no. 2, 1 June 2001 (2001-06-01), pages 174-176, XP055405588, US ISSN: 0165-4608, DOI: 10.1016/S0165-4608(00)00431-3

DESCRIPTION

BACKGROUND OF THE INVENTION

[0001] Basic cancer research has focused on identifying the genetic changes that lead to cancer. This has led to major advances in our understanding of the molecular and biochemical pathways that are involved in tumorigenesis and malignant transformation. However, our understanding of the cellular biology has lagged. Although the effects of particular mutations on the proliferation and survival of model cells, such as fibroblasts or cell lines, can be predicted, the effects of such mutations on the actual cells involved in specific cancers is largely guesswork.

[0002] A tumor can be viewed as an aberrant organ initiated by a tumorigenic cancer cell that acquired the capacity for indefinite proliferation through accumulated mutations. In this view of a tumor as an abnormal organ, the principles of normal stem cell biology can be applied to better understand how tumors develop. Many observations suggest that analogies between normal stem cells and tumorigenic cells are appropriate. Both normal stem cells and tumorigenic cells have extensive proliferative potential and the ability to give rise to new (normal or abnormal) tissues. Both tumors and normal tissues are composed of heterogeneous combinations of cells, with different phenotypic characteristics and different proliferative potentials.

[0003] Because most tumors have a clonal origin, the original tumorigenic cancer cell gives rise to phenotypically diverse progeny, including cancer cells with indefinite proliferative potential, as well as cancer cells with limited or no proliferative potential. This suggests that tumorigenic cancer cells undergo processes that are analogous to the self-renewal and differentiation of normal stem cells. Tumorigenic cells can be thought of as cancer stem cells that undergo an aberrant and poorly regulated process of organogenesis analogous to what normal stem cells do. Although some of the heterogeneity in tumors arises as a result of continuing mutagenesis, it is likely that heterogeneity also arises through the aberrant differentiation of cancer cells.

[0004] It is well documented that many types of tumors contain cancer cells with heterogeneous phenotypes, reflecting aspects of the differentiation that normally occurs in the tissues from which the tumors arise. The variable expression of normal differentiation markers by cancer cells in a tumor suggests that some of the heterogeneity in tumors arises as a result of the anomalous differentiation of tumor cells. Examples of this include the variable expression of myeloid markers in chronic myeloid leukaemia, the variable expression of neuronal markers within peripheral neuroectodermal tumors, and the variable expression of milk proteins or the estrogen receptor within breast cancer.

[0005] It was first extensively documented for leukemia and multiple myeloma that only a small

subset of cancer cells is capable of extensive proliferation. Because the differences in clonogenicity among the leukemia cells mirrored the differences in clonogenicity among normal hematopoietic cells, the clonogenic leukemic cells were described as leukemic stem cells. It has also been shown for solid cancers that the cells are phenotypically heterogeneous and that only a small proportion of cells are clonogenic in culture and *in vivo*. Just as in the context of leukemic stem cells, these observations led to the hypothesis that only a few cancer cells are actually tumorigenic and that these tumorigenic cells act as cancer stem cells

[0006] In support of this hypothesis, recent studies have shown that, similar to leukemia and other hematologic malignancies, tumorigenic and non-tumorigenic populations of breast cancer cells can be isolated based on their expression of cell surface markers. In many cases of breast cancer, only a small subpopulation of cells had the ability to form new tumors. This work strongly supports the existence of CSC in breast cancer. Further evidence for the existence of cancer stem cells occurring in solid tumors has been found in central nervous system (CNS) malignancies. Using culture techniques similar to those used to culture normal neuronal stem cells it has been shown that neuronal CNS malignancies contain a small population of cancer cells that are clonogenic *in vitro* and initiate tumors *in vivo*, while the remaining cells in the tumor do not have these properties.

[0007] Stem cells are defined as cells that have the ability to perpetuate themselves through self-renewal and to generate mature cells of a particular tissue through differentiation. In most tissues, stem cells are rare. As a result, stem cells must be identified prospectively and purified carefully in order to study their properties. Perhaps the most important and useful property of stem cells is that of self-renewal. Through this property, striking parallels can be found between stem cells and cancer cells: tumors may often originate from the transformation of normal stem cells, similar signaling pathways may regulate self-renewal in stem cells and cancer cells, and cancers may comprise rare cells with indefinite potential for self-renewal that drive tumorigenesis.

[0008] The presence of cancer stem cells has profound implications for cancer therapy. At present, all of the phenotypically diverse cancer cells in a tumor are treated as though they have unlimited proliferative potential and can acquire the ability to metastasize. For many years, however, it has been recognized that small numbers of disseminated cancer cells can be detected at sites distant from primary tumors in patients that never manifest metastatic disease. One possibility is that immune surveillance is highly effective at killing disseminated cancer cells before they can form a detectable tumor. Another possibility is that most cancer cells lack the ability to form a new tumor such, that only the dissemination of rare cancer stem cells can lead to metastatic disease. If so, the goal of therapy must be to identify and kill this cancer stem cell population.

[0009] The prospective identification and isolation of cancer stem cells will allow more efficient identification of diagnostic markers and therapeutic targets expressed by the stem cells. Existing therapies have been developed largely against the bulk population of tumor cells, because the therapies are identified by their ability to shrink the tumor mass. However,

because most cells within a cancer have limited proliferative potential, an ability to shrink a tumor mainly reflects an ability to kill these cells. Therapies that are more specifically directed against cancer stem cells may result in more durable responses and cures of metastatic tumors.

[0010] Hematopoiesis proceeds through an organized developmental hierarchy initiated by hematopoietic stem cells (HSC) that give rise to progressively more committed progenitors and eventually terminally differentiated blood cells (Bryder et al., 2006). Although the concept of the HSC was not new, it was not until 1988 that it was shown that this population could be prospectively isolated from mouse bone marrow on the basis of cell-surface markers using fluorescence-activated cell sorting (FACS) (Spangrude et al., 1988). Since that time, the surface immunophenotype of the mouse HSC has become increasingly refined, such that functional HSC can be isolated with exquisite sensitivity, resulting in a purity of 1 in 1.3 cells (Kiel et al., 2005). While our ability to prospectively isolate mouse HSC has improved dramatically over the past 20 years, our understanding of the earliest events in the human hematopoietic system lags far behind.

[0011] Cancer stem cells are discussed in, for example, Pardal et al. (2003) *Nat Rev Cancer* 3, 895-902; Reya et al. (2001) *Nature* 414, 105-11; Bonnet & Dick (1997) *Nat Med* 3, 730-7; Al-Hajj et al. (2003) *Proc Natl Acad Sci U S A* 100, 3983-8; Dontu et al. (2004) *Breast Cancer Res* 6, R605-15; Singh et al. (2004) *Nature* 432, 396-401.

[0012] US 2004/213792 describes a method for inhibiting cellular activation by insulin-like growth factor-1. Hosen et al. (2007) *PNAS* 104(26):11008-11013 reports that CD96 is a leukemic stem cell-specific marker in human acute myeloid leukemia. Clarke et al. (2006) *Cancer Res* 66(19):9339-9344 describes cancer stem cells. US 2007/287163 describes methods for diagnosis of acute myeloid leukemia. US 6,733,743 describes methods to impair hematologic cancer progenitor cells and compounds related thereto. US 6,491,917 describes an antibody composition for debulking blood and bone marrow samples from CML patients. Martin-Subero et al. (2001) describes amplification of *ERBB2*, *RARA*, and *TOP2A* genes in a myelodysplastic syndrome transforming to acute myeloid leukemia.

[0013] The identification of a hierarchy of multipotent hematopoietic progenitors in human cord blood, including multipotent progenitor cells, may be found in Majeti et al. (2007) *Cell Stem Cell* 1(6):635-45, which describes markers identifying the multipotent progenitors.

SUMMARY OF THE INVENTION

[0014] The invention provides a monoclonal antibody directed against human CD47 that blocks the interaction of CD47 with SIRP α , for use in the treatment of human acute myeloid leukemia (AML) by stimulating phagocytosis of AML leukemic stem cells (LSC) by macrophages, wherein the antibody directed against human CD47 is used in combination with one or more

additional monoclonal antibodies directed against one or more of the following markers: CD96, CD97, CD99, CD180, PTHR2, HAVCR2, CD123, CD44 and CD33.

[0015] Markers of acute myeloid leukemia stem cells (AMLSC) are provided herein. The markers are polynucleotides or polypeptides that are differentially expressed on AMLSC as compared to normal counterpart cells. Uses of the markers include use as targets for therapeutic antibodies or ligands; as targets for drug development, and for identification or selection of AMLSC cell populations.

[0016] The AMLSC markers are useful as targets of therapeutic monoclonal antibodies for treatment of patients with *de novo*, relapsed, or refractory acute myeloid leukemia. Such monoclonal antibodies are also useful in the treatment of pre-leukemic conditions, such as myelodysplastic syndromes (MDS) and myeloproliferative disorders (MPDs) including: chronic myelogenous leukemia, polycythemia vera, essential thrombocytosis, agnogenic myelofibrosis and myeloid metaplasia, and others. Antibodies include free antibodies and antigen binding fragments derived therefrom, and conjugates, e.g. pegylated antibodies, drug, radioisotope, or toxin conjugates, and the like.

[0017] In some instances, combinations of monoclonal antibodies are used in the treatment of human AML or pre-leukemic conditions. In one instance, a monoclonal antibody directed against CD47, for example an antibody that blocks the interaction of CD47 with SIRP α , is combined with monoclonal antibodies directed against one or more additional AMLSC markers, e.g. CD96, CD97, CD99, CD180, PTHR2, HAVCR2 (also referred to as TIM3), and the like, which compositions can be synergistic in enhancing phagocytosis and elimination of AML LSC as compared to the use of single antibodies.

[0018] The AMLSC markers are useful as targets of monoclonal antibodies for use in *ex vivo* purging of autologous stem cell products (mobilized peripheral blood or bone marrow) for use in autologous transplantation for patients with acute myeloid leukemia or the pre-leukemic conditions outlined above. Combinations of monoclonal antibodies directed against AML LSC-specific cell surface molecules, as described above, can be synergistic in eliminating LSC.

[0019] The AMLSC markers are useful in clinical diagnostic applications including, without limitation, primary diagnosis of AML or pre-leukemic conditions from blood and/or bone marrow specimens, evaluation of leukemic involvement of the cerebrospinal and other body fluids, monitoring of interval disease progression, and monitoring of minimal residual disease status.

[0020] As an alternative to monoclonal antibodies, the ligands of AMLSC markers, either as single agents or in combination, may be used to target them in AML or the pre-leukemic conditions outlined above. The ligands can be free or conjugated, for direct administration to patients or for *ex vivo* purging of autologous stem cell products. Some specific molecules and their ligands include, without limitation, CD155-Fc fusion protein that binds CD96; TIP39 that binds PTHR2; Galectin-9 that binds HAVCR2.

[0021] The AMLSC cells can be prospectively isolated or identified from primary tumor samples, and possess the unique properties of cancer stem cells in functional assays for cancer stem cell self-renewal and differentiation.

[0022] In some instances, methods are disclosed herein for detection, classification or clinical staging of acute myeloid leukemias according to the stem cells that are present in the leukemia, where greater numbers of stem cells are indicative of a more aggressive cancer phenotype. Staging is useful for prognosis and treatment. In some instances, a tumor sample is analyzed by histochemistry, including immunohistochemistry, *in situ* hybridization, and the like, for the presence of CD34⁺CD38⁻ cells that express one or more AMLSC markers provided herein. The presence of such cells indicates the presence of AMLSC.

[0023] In another instance, methods for the isolation of AMLSC are disclosed, comprising contacted a candidate cell population with a binding reagent specific for one or more of the AMLSC markers provided herein, and selecting for cells that have bound to the reagent(s). The cells may further be selected as being CD34⁺CD38⁻. The cells are useful for experimental evaluation, and as a source of lineage and cell specific products, including mRNA species useful in identifying genes specifically expressed in these cells, and as targets for the discovery of factors or molecules that can affect them. AMLSC may be used, for example, in a method of screening a compound for an effect on the cells. This involves combining the compound with the cell population of the disclosure, and then determining any modulatory effect resulting from the compound. This may include examination of the cells for viability, toxicity, metabolic change, or an effect on cell function. The phenotype of AMLSC described herein provides a means of predicting disease progression, relapse, and development of drug resistance.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024]

Figure 1: *Differential Gene Expression Between AML LSC and Normal Bone Marrow HSC and MPP* (A) Heat maps demonstrating genes found to be differentially expressed at least 2 fold between bone marrow HSC (n=4) and AML LSC (n=9) or bone marrow MPP (n=4) and AML LSC (n=9). Expression relative to the median is indicated for genes with p<0.05 and a FDR of 5%. (B) Selected list of transmembrane proteins found to be at least 2-fold more highly expressed in AML LSC than HSC or MPP. NS: not significant.

Figure 2. CD47 is more highly expressed on AML LSC. Mobilized peripheral blood (MPB) HSC and AML LSC were examined for CD47 expression by flow cytometry. (A) Representative flow cytometry plots indicating expression of CD47 relative to an isotype control. (B) Summary of CD47 expression on all samples assayed, with the indicated means.

Figure 3: Anti-CD47 Antibody Stimulates In Vitro Macrophage Phagocytosis of Primary Human AML LSC. AML LSC were purified by FACS from two primary human AML samples, labeled

with the fluorescent dye CFSE, and incubated with mouse bone marrow-derived macrophages either in the presence of an isotype control (A) or anti-CD47 antibody (B). These cells were assessed by immunofluorescence microscopy for the presence of fluorescently labeled LSC within the macrophages. (C) The phagocytic index was determined for each condition by calculating the number of ingested cells per 100 macrophages.

Figure 4. Anti-CD47 antibody stimulates in vitro macrophage phagocytosis of primary human AML LSC. AML LSC were purified by FACS from two primary human AML samples and labeled with the fluorescent dye CFSE. These cells were incubated with mouse bone marrow-derived macrophages, either in the presence of an isotype matched control (left) or anti-CD47 antibody (right). The macrophages were harvested, stained with a fluorescently labeled anti-mouse macrophage antibody, and analyzed by flow cytometry. mMac+CFSE+ double-positive events identify macrophages that have phagocytosed CFSE-labeled LSC. (A,B) two independent primary AML LSC samples.

Figure 5. Anti-CD47 antibody inhibits in vivo engraftment of primary human AML. Two primary human AML samples were untreated (control, n=3) or coated with anti-human CD47 antibody (anti-CD47, n=6) prior to transplantation into newborn NOG mice. 13 weeks later, mice were sacrificed and the bone marrow was analyzed for the presence of human CD45+CD33+ myeloid leukemia cells by flow cytometry.

Figure 6A-B: CD99 Expression on AML LSC Compared to Normal HSC. CD99 expression was examined on several samples of normal human bone marrow HSC (n=3) and de novo human AML LSC (n=7). Representative histograms of CD99 expression on HSC and LSC (left) and summary of normalized mean fluorescence intensity (MFI) of all specimens (right) are shown. Mean CD99 expression was increased 5.6 fold in AML LSC compared to HSC (p=0.05).

Figure 7A-B: CD97 Expression on AML LSC Compared to Normal HSC. CD97 expression was examined on several samples of normal human bone marrow HSC (n=3) and de novo human AML LSC (n=7). Representative histograms of CD97 expression on HSC and LSC (left) and summary of normalized mean fluorescence intensity (MFI) of all specimens (right) are shown. Mean CD97 expression was increased 7.9 fold in AML LSC compared to HSC (p=0.03).

Figure 8A-B: CD180 Expression on AML LSC Compared to Normal HSC. CD180 expression was examined on several samples of normal human bone marrow HSC (n=3) and de novo human AML LSC (n=7). Representative histograms of CD180 expression on HSC and LSC (left) and summary of normalized mean fluorescence intensity (MFI) of all specimens (right) are shown. Mean CD180 expression was increased 60 fold in AML LSC compared to HSC (p=0.20).

Figure 9A-B: TIM3 Expression on AML LSC Compared to Normal HSC. TIM3 expression was examined on several samples of normal human bone marrow HSC (n=3) and de novo human AML LSC (n=14). Representative histograms of TIM3 expression on HSC and LSC (left) and summary of normalized mean fluorescence intensity (MFI) of all specimens (right) are shown. Mean TIM3 expression was increased 9 fold in AML LSC compared to HSC (p=0.01).

Figure 10: PTH2R Expression in AML LSC Compared to Normal HSC. PTH2R expression was examined in several samples of normal human bone marrow HSC (n=3) and de novo human AML LSC (n=9). Expression was determined by qRT-PCR and is expressed relative to beta-actin as a control. Mean PTH2R expression was increased 21 fold in AML LSC compared to HSC (p<0.001).

DETAILED DESCRIPTION

[0025] The present disclosure identifies polynucleotides, as well as polypeptides encoded thereby, that are differentially expressed in acute myeloid leukemia stem cells (AMLSC). Methods are provided in which these polynucleotides and polypeptides, which may be collectively referred to as AMLSC markers, are used for detecting, assessing, and reducing the growth of cancer cells. Methods may use one or a combination of markers, where a combination may include 2, 3 or more markers, and in some instances will include CD47 in combination with 1, 2 or more markers. The disclosed markers find use in the prevention, treatment, detection or research of leukemic and pre-leukemic conditions.

[0026] The markers of the disclosure, in some instances, are expressed on the AMLSC cell surface. In some instances, the markers are expressed as a level at least 2X the expression level of a counterpart non-transformed cell, e.g. a human hematopoietic stem cell, and/or a human hematopoietic multipotent progenitor cell, where expression may be determined as the level of transcription, mRNA accumulation, and/or protein accumulation. In other instances the markers are expressed as a level at least 3X, at least 4X, at least 5X, at least 10X, at least 20X or greater, than the expression level of a counterpart non-transformed cell.

[0027] Herein disclosed are methods of using the markers described herein in diagnosis of cancer, classification and treatment of leukemic and pre-leukemic conditions according to expression profiles. The methods are useful for detecting AMLSC, facilitating diagnosis of AML and the severity of the cancer (e.g., tumor grade, tumor burden, and the like) in a subject, facilitating a determination of the prognosis of a subject, and assessing the responsiveness of the subject to therapy. The detection methods of the disclosure can be conducted *in vitro* or *in vivo*, on isolated cells, or in whole tissues or a bodily fluid, e.g., blood, lymph node biopsy samples, and the like.

[0028] As used herein, the terms "a gene that is differentially expressed in a cancer stem cell," and "a polynucleotide that is differentially expressed in a cancer stem cell", are used interchangeably herein, and generally refer to a polynucleotide that represents or corresponds to a gene that is differentially expressed in a cancer stem cell when compared with a cell of the same cell type that is not cancerous, e.g., mRNA is found at levels at least about 25%, at least about 50% to about 75%, at least about 90%, at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 5-fold, at least about 10-fold, or at least about 50-fold or

more, different (e.g., higher or lower). The comparison can be made between AMLSC and the normal counterpart cells a human hematopoietic stem cell (HSC), which include without limitation cells having the phenotype $\text{Lin}^-\text{CD34}^+\text{CD38}^-\text{CD90}^+$; or the phenotype $\text{Lin}^-\text{CD34}^+\text{CD38}^-\text{CD90}^+\text{CD45RA}^-$ and a human hematopoietic multipotent progenitor cell (MPP), which include without limitation cells having the phenotype $\text{Lin}^-\text{CD34}^+\text{CD38}^-\text{CD90}^-$; or the phenotype $\text{Lin}^-\text{CD34}^+\text{CD38}^-\text{CD90}^-\text{CD45RA}^-$. The term "a polypeptide marker for a cancer stem cell" refers to a polypeptide encoded by a polynucleotide that is differentially expressed in a cancer stem cell.

[0029] In some instances, the markers are demonstrated by flow cytometry to be present on a majority of AMLSC, when compared to human HSC or MPP, as defined above. Such markers include, without limitation, CD47, CD96, CD97 and CD99.

[0030] In other instances, the markers are absent on human HSC or human MPP, but are highly expressed on AMLSC. Such markers include, without limitation, those set forth in Table 1.

[0031] In other instances, the markers are differentially expressed on AMLSC, as compared to human HSC or MPP. Such markers include, without limitation, those set forth in Table 1.

[0032] A polynucleotide or sequence that corresponds to, or represents a gene means that at least a portion of a sequence of the polynucleotide is present in the gene or in the nucleic acid gene product (e.g., mRNA or cDNA). A subject nucleic acid may also be "identified" by a polynucleotide if the polynucleotide corresponds to or represents the gene. Genes identified by a polynucleotide may have all or a portion of the identifying sequence wholly present within an exon of a genomic sequence of the gene, or different portions of the sequence of the polynucleotide may be present in different exons (e.g., such that the contiguous polynucleotide sequence is present in an mRNA, either pre- or post-splicing, that is an expression product of the gene). An "identifying sequence" is a minimal fragment of a sequence of contiguous nucleotides that uniquely identifies or defines a polynucleotide sequence or its complement.

[0033] The polynucleotide may represent or correspond to a gene that is modified in a cancer stem cell (CSC) relative to a normal cell. The gene in the CSC may contain a deletion, insertion, substitution, or translocation relative to the polynucleotide and may have altered regulatory sequences, or may encode a splice variant gene product, for example. The gene in the CSC may be modified by insertion of an endogenous retrovirus, a transposable element, or other naturally occurring or non-naturally occurring nucleic acid.

[0034] Sequences of interest include those set forth in Table 1, which are differentially expressed in AMLSC relative to normal counterpart cells.

[0035] Methods are also provided for optimizing therapy, by first classification, and based on that information, selecting the appropriate therapy, dose, treatment modality, *etc.* which

optimizes the differential between delivery of an anti-proliferative treatment to the undesirable target cells, while minimizing undesirable toxicity. The treatment is optimized by selection for a treatment that minimizes undesirable toxicity, while providing for effective anti-proliferative activity.

[0036] The invention finds use in the treatment of acute myeloid leukemias. Acute leukemias are rapidly progressing leukemia characterized by replacement of normal bone marrow by blast cells of a clone arising from malignant transformation of a hematopoietic stem cell. The acute leukemias include acute lymphoblastic leukemia (ALL) and acute myelogenous leukemia (AML). ALL often involves the CNS, whereas acute monoblastic leukemia involves the gums, and AML involves localized collections in any site (granulocytic sarcomas or chloromas). AML is the most common acute leukemia affecting adults, and its incidence increases with age. While AML is a relatively rare disease overall, accounting for approximately 1.2% of cancer deaths in the United States, its incidence is expected to increase as the population ages.

[0037] The presenting symptoms are usually nonspecific (e.g., fatigue, fever, malaise, weight loss) and reflect the failure of normal hematopoiesis. Anemia and thrombocytopenia are very common (75 to 90%). The WBC count may be decreased, normal, or increased. Blast cells are usually found in the blood smear unless the WBC count is markedly decreased. The blasts of ALL can be distinguished from those of AML by histochemical studies, cytogenetics, immunophenotyping, and molecular biology studies. In addition to smears with the usual stains, terminal transferase, myeloperoxidase, Sudan black B, and specific and nonspecific esterase.

[0038] "Diagnosis" as used herein generally includes determination of a subject's susceptibility to a disease or disorder, determination as to whether a subject is presently affected by a disease or disorder, prognosis of a subject affected by a disease or disorder (e.g., identification of cancerous states, stages of cancer, or responsiveness of cancer to therapy), and use of therametrics (e.g., monitoring a subject's condition to provide information as to the effect or efficacy of therapy).

[0039] The term "biological sample" encompasses a variety of sample types obtained from an organism and can be used in a diagnostic or monitoring assay. The term encompasses blood and other liquid samples of biological origin, solid tissue samples, such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof. The term encompasses samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components. The term encompasses a clinical sample, and also includes cells in cell culture, cell supernatants, cell lysates, serum, plasma, biological fluids, and tissue samples.

[0040] The terms "treatment", "treating", "treat" and the like are used herein to generally refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete stabilization or cure for a disease and/or adverse

effect attributable to the disease. "Treatment" as used herein covers any treatment of a disease in a mammal, particularly a human, and includes: (a) preventing the disease or symptom from occurring in a subject which may be predisposed to the disease or symptom but has not yet been diagnosed as having it; (b) inhibiting the disease symptom, i.e., arresting its development; or (c) relieving the disease symptom, i.e., causing regression of the disease or symptom.

[0041] The terms "individual," "subject," "host," and "patient," used interchangeably herein and refer to any mammalian subject for whom diagnosis, treatment, or therapy is desired, particularly humans.

[0042] A "host cell", as used herein, refers to a microorganism or a eukaryotic cell or cell line cultured as a unicellular entity which can be, or has been, used as a recipient for a recombinant vector or other transfer polynucleotides, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

[0043] The terms "cancer", "neoplasm", "tumor", and "carcinoma", are used interchangeably herein to refer to cells which exhibit relatively autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation. In general, cells of interest for detection or treatment in the present application include precancerous (e.g., benign), malignant, pre-metastatic, metastatic, and non-metastatic cells. Detection of cancerous cells is of particular interest. The term "normal" as used in the context of "normal cell," is meant to refer to a cell of an untransformed phenotype or exhibiting a morphology of a non-transformed cell of the tissue type being examined. "Cancerous phenotype" generally refers to any of a variety of biological phenomena that are characteristic of a cancerous cell, which phenomena can vary with the type of cancer. The cancerous phenotype is generally identified by abnormalities in, for example, cell growth or proliferation (e.g., uncontrolled growth or proliferation), regulation of the cell cycle, cell mobility, cell-cell interaction, or metastasis, etc.

[0044] "Therapeutic target" refers to a gene or gene product that, upon modulation of its activity (e.g., by modulation of expression, biological activity, and the like), can provide for modulation of the cancerous phenotype. As used throughout, "modulation" is meant to refer to an increase or a decrease in the indicated phenomenon (e.g., modulation of a biological activity refers to an increase in a biological activity or a decrease in a biological activity).

ACUTE MYELOID LEUKEMIA

[0045] Acute Myelocytic Leukemia (AML, Acute Myelogenous Leukemia; Acute Myeloid Leukemia). In AML, malignant transformation and uncontrolled proliferation of an abnormally differentiated, long-lived myeloid progenitor cell results in high circulating numbers of immature

blood forms and replacement of normal marrow by malignant cells. Symptoms include fatigue, pallor, easy bruising and bleeding, fever, and infection; symptoms of leukemic infiltration are present in only about 5% of patients (often as skin manifestations). Examination of peripheral blood smear and bone marrow is diagnostic. Treatment includes induction chemotherapy to achieve remission and post-remission chemotherapy (with or without stem cell transplantation) to avoid relapse.

[0046] AML has a number of subtypes that are distinguished from each other by morphology, immunophenotype, and cytochemistry. Five classes are described, based on predominant cell type, including myeloid, myeloid-monocytic, monocytic, erythroid, and megakaryocytic. Acute promyelocytic leukemia is a particularly important subtype, representing 10 to 15% of all cases of AML, striking a younger age group (median age 31 yr) and particular ethnicity (Hispanics), in which the patient commonly presents with a coagulation disorder.

[0047] Remission induction rates range from 50 to 85%. Long-term disease-free survival reportedly occurs in 20 to 40% of patients and increases to 40 to 50% in younger patients treated with stem cell transplantation.

[0048] Prognostic factors help determine treatment protocol and intensity; patients with strongly negative prognostic features are usually given more intense forms of therapy, because the potential benefits are thought to justify the increased treatment toxicity. The most important prognostic factor is the leukemia cell karyotype; favorable karyotypes include t(15;17), t(8;21), and inv16 (p13;q22). Negative factors include increasing age, a preceding myelodysplastic phase, secondary leukemia, high WBC count, and absence of Auer rods. The FAB or WHO classification alone does not predict response.

[0049] Initial therapy attempts to induce remission and differs most from ALL in that AML responds to fewer drugs. The basic induction regimen includes cytarabine by continuous IV infusion or high doses for 5 to 7 days; daunorubicin or idarubicin is given IV for 3 days during this time. Some regimens include 6-thioguanine, etoposide, vincristine, and prednisone, but their contribution is unclear. Treatment usually results in significant myelosuppression, with infection or bleeding; there is significant latency before marrow recovery. During this time, meticulous preventive and supportive care is vital.

POLYPEPTIDE AND POLYNUCLEOTIDE SEQUENCES AND ANTIBODIES

[0050] Disclosed herein are polynucleotides and polypeptides that represent genes that are differentially expressed in human AMLSC. These polynucleotides, polypeptides and fragments thereof have uses that include, but are not limited to, diagnostic probes and primers as starting materials for probes and primers, as immunogens for antibodies useful in cancer diagnosis and therapy, and the like as discussed herein.

[0051] Nucleic acid compositions include fragments and primers, and are at least about 15 bp

in length, at least about 30 bp in length, at least about 50 bp in length, at least about 100 bp, at least about 200 bp in length, at least about 300 bp in length, at least about 500 bp in length, at least about 800 bp in length, at least about 1 kb in length, at least about 2.0 kb in length, at least about 3.0 kb in length, at least about 5 kb in length, at least about 10 kb in length, at least about 50 kb in length and are usually less than about 200 kb in length. In some instances, a fragment of a polynucleotide is the coding sequence of a polynucleotide. Also included are variants or degenerate variants of a sequence provided herein. In general, variants of a polynucleotide provided herein have a fragment of sequence identity that is greater than at least about 65%, greater than at least about 70%, greater than at least about 75%, greater than at least about 80%, greater than at least about 85%, or greater than at least about 90%, 95%, 96%, 97%, 98%, 99% or more (i.e. 100%) as compared to an identically sized fragment of a provided sequence. as determined by the Smith-Waterman homology search algorithm as implemented in MPSRCH program (Oxford Molecular). Nucleic acids having sequence similarity can be detected by hybridization under low stringency conditions, for example, at 50° C. and 10XSSC (0.9 M saline/0.09 M sodium citrate) and remain bound when subjected to washing at 55° C. in 1XSSC. Sequence identity can be determined by hybridization under high stringency conditions, for example, at 50° C. or higher and 0.1XSSC (9 mM saline/0.9 mM sodium citrate). Hybridization methods and conditions are well known in the art, see, e.g., U.S. Pat. No. 5,707,829. Nucleic acids that are substantially identical to the provided polynucleotide sequences, e.g. allelic variants, genetically altered versions of the gene, etc., bind to the provided polynucleotide sequences under stringent hybridization conditions.

[0052] Probes specific to the polynucleotides described herein can be generated using the polynucleotide sequences disclosed herein. The probes are usually a fragment of a polynucleotide sequences provided herein. The probes can be synthesized chemically or can be generated from longer polynucleotides using restriction enzymes. The probes can be labeled, for example, with a radioactive, biotinylated, or fluorescent tag. Preferably, probes are designed based upon an identifying sequence of any one of the polynucleotide sequences provided herein.

[0053] The nucleic acid compositions described herein can be used to, for example, produce polypeptides, as probes for the detection of mRNA in biological samples (e.g., extracts of human cells) or cDNA produced from such samples, to generate additional copies of the polynucleotides, to generate ribozymes or antisense oligonucleotides, and as single stranded DNA probes or as triple-strand forming oligonucleotides.

[0054] The probes described herein can be used to, for example, determine the presence or absence of any one of the polynucleotide provided herein or variants thereof in a sample. These and other uses are described in more detail below. In one instance, real time PCR analysis is used to analyze gene expression.

[0055] The polypeptides contemplated by the disclosure include those encoded by the disclosed polynucleotides and the genes to which these polynucleotides correspond, as well as nucleic acids that, by virtue of the degeneracy of the genetic code, are not identical in

sequence to the disclosed polynucleotides. Further polypeptides contemplated include polypeptides that are encoded by polynucleotides that hybridize to polynucleotide of the sequence listing. Thus, the disclosure includes a polypeptide encoded by a polynucleotide having the sequence of any one of the polynucleotide sequences provided herein, or a variant thereof.

[0056] In general, the term "polypeptide" as used herein refers to both the full length polypeptide encoded by the recited polynucleotide, the polypeptide encoded by the gene represented by the recited polynucleotide, as well as portions or fragments thereof. "Polypeptides" also includes variants of the naturally occurring proteins, where such variants are homologous or substantially similar to the naturally occurring protein, and can be of an origin of the same or different species as the naturally occurring protein. In general, variant polypeptides have a sequence that has at least about 80%, usually at least about 90%, and more usually at least about 98% sequence identity with a differentially expressed polypeptide described herein. The variant polypeptides can be naturally or non-naturally glycosylated, i.e., the polypeptide has a glycosylation pattern that differs from the glycosylation pattern found in the corresponding naturally occurring protein.

[0057] Fragments of the polypeptides disclosed herein, particularly biologically active fragments and/or fragments corresponding to functional domains, are of interest. Fragments of interest will typically be at least about 10 aa to at least about 15 aa in length, usually at least about 50 aa in length, and can be as long as 300 aa in length or longer, but will usually not exceed about 1000 aa in length, where the fragment will have a stretch of amino acids that is identical to a polypeptide encoded by a polynucleotide having a sequence of any one of the polynucleotide sequences provided herein, or a homolog thereof. A fragment "at least 20 aa in length," for example, is intended to include 20 or more contiguous amino acids from, for example, the polypeptide encoded by a cDNA, in a cDNA clone contained in a deposited library or the complementary stand thereof. In this context "about" includes the particularly recited value or a value larger or smaller by several (5, 4, 3, 2, or 1) amino acids. The protein variants described herein are encoded by polynucleotides that are also disclosed. The genetic code can be used to select the appropriate codons to construct the corresponding variants. The polynucleotides may be used to produce polypeptides, and these polypeptides may be used to produce antibodies by known methods described above and below.

[0058] A polypeptide can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

[0059] Polypeptides can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a

prokaryotic or eukaryotic host, including, for example, bacterial, yeast higher plant, insect, and mammalian cells.

[0060] Gene products, including polypeptides, mRNA (particularly mRNAs having distinct secondary and/or tertiary structures), cDNA, or complete gene, can be prepared and used for raising antibodies for experimental, diagnostic, and therapeutic purposes. Antibodies may be used to identify AMLSC cells or subtypes. The polynucleotide or related cDNA is expressed as described herein, and antibodies are prepared. These antibodies are specific to an epitope on the polypeptide encoded by the polynucleotide, and can precipitate or bind to the corresponding native protein in a cell or tissue preparation or in a cell-free extract of an in vitro expression system.

[0061] The antibodies may be utilized for immunophenotyping of cells and biological samples. The translation product of a differentially expressed gene may be useful as a marker. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Pat. No. 5,985,660; and Morrison et al. *Cell*, 96:737-49 (1999)). These techniques allow for the screening of particular populations of cells; in immunohistochemistry of biopsy samples; in detecting the presence of markers shed by cancer cells into the blood and other biologic fluids, and the like.

[0062] In many instances, the levels of a subject gene or gene product are measured. By measured is meant qualitatively or quantitatively estimating the level of the gene product in a first biological sample either directly (e.g. by determining or estimating absolute levels of gene product) or relatively by comparing the levels to a second control biological sample. In many instances the second control biological sample is obtained from an individual not having cancer. As will be appreciated in the art, once a standard control level of gene expression is known, it can be used repeatedly as a standard for comparison. Other control samples include samples of cancerous tissue.

[0063] The methods can be used to detect and/or measure mRNA levels of a gene that is differentially expressed in a cancer cell. In some instances, the methods comprise: contacting a sample with a polynucleotide that corresponds to a differentially expressed gene described herein under conditions that allow hybridization; and detecting hybridization, if any. Detection of differential hybridization, when compared to a suitable control, is an indication of the presence in the sample of a polynucleotide that is differentially expressed in a cancer cell. Appropriate controls include, for example, a sample that is known not to contain a polynucleotide that is differentially expressed in a cancer cell. Conditions that allow hybridization are known in the art, and have been described in more detail above.

[0064] Detection can also be accomplished by any known method, including, but not limited to,

in situ hybridization, PCR (polymerase chain reaction), RT-PCR (reverse transcription-PCR), and "Northern" or RNA blotting, arrays, microarrays, etc, or combinations of such techniques, using a suitably labeled polynucleotide. A variety of labels and labeling methods for polynucleotides are known in the art and can be used in the assay methods of the disclosure. Specific hybridization can be determined by comparison to appropriate controls.

[0065] Labeled nucleic acid probes may be used to detect expression of a gene corresponding to the provided polynucleotide, e.g. in a macroarray format, Northern blot, etc. The amount of hybridization can be quantitated to determine relative amounts of expression, for example under a particular condition. Probes are used for in situ hybridization to cells to detect expression. Probes can also be used in vivo for diagnostic detection of hybridizing sequences. Probes may be labeled with a radioactive isotope. Other types of detectable labels can be used such as chromophores, fluorophores, and enzymes.

[0066] Polynucleotide arrays provide a high throughput technique that can assay a large number of polynucleotides or polypeptides in a sample. This technology can be used as a tool to test for differential expression. A variety of methods of producing arrays, as well as variations of these methods, are known in the art and contemplated for use. For example, arrays can be created by spotting polynucleotide probes onto a substrate (e.g., glass, nitrocellulose, etc.) in a two-dimensional matrix or array having bound probes. The probes can be bound to the substrate by either covalent bonds or by non-specific interactions, such as hydrophobic interactions.

CHARACTERIZATION OF ACUTE MYELOID LEUKEMIA STEM CELLS

[0067] In acute myeloid leukemias, characterization of cancer stem cells allows for the development of new treatments that are specifically targeted against this critical population of cells, particularly their ability to self-renew, resulting in more effective therapies.

[0068] In human acute myeloid leukemias it is shown herein that there is a subpopulation of tumorigenic cancer cells with both self-renewal and differentiation capacity. These tumorigenic cells are responsible for tumor maintenance, and also give rise to large numbers of abnormally differentiating progeny that are not tumorigenic, thus meeting the criteria of cancer stem cells. Tumorigenic potential is contained within a subpopulation of cancer cells differentially expressing the markers of the present disclosure.

[0069] In some instances, the number of AMLSC in a patient sample is determined relative to the total number of AML cancer cells, where a greater percentage of AMLSC is indicative of the potential for continued self-renewal of cells with the cancer phenotype. The quantitation of AMLSC in a patient sample may be compared to a reference population, e.g. a patient sample such as a blood sample, a remission patient sample, *etc.* In some instances, the quantitation of AMLSC is performed during the course of treatment, where the number of AML cancer cells and the percentage of such cells that are AMLSC are quantitated before, during and as follow-

up to a course of therapy. Desirably, therapy targeted to cancer stem cells results in a decrease in the total number, and/or percentage of AMLSC in a patient sample.

[0070] In other instances, anti-cancer agents are targeted to AMLSC by specific binding to a marker or combination of markers of the present disclosure. In such instances, the anti-cancer agents include antibodies and antigen-binding derivatives thereof specific for a marker or combination of markers of the present disclosure, which are optionally conjugated to a cytotoxic moiety. Depletion of AMLSC is useful in the treatment of AML. Depletion achieves a reduction in circulating AMLSC by up to about 30%, or up to about 40%, or up to about 50%, or up to about 75% or more. Depletion can be achieved by using an agent to deplete AMLSC either *in vivo* or *ex vivo*.

[0071] The AMLSC are identified by their phenotype with respect to particular markers, and/or by their functional phenotype. In some instances, the AMLSC are identified and/or isolated by binding to the cell with reagents specific for the markers of interest. The cells to be analyzed may be viable cells, or may be fixed or embedded cells.

[0072] In some instances, the reagents specific for the markers of interest are antibodies, which may be directly or indirectly labeled. Such antibodies will usually include antibodies specific for a marker or combination of markers of the present disclosure.

Treatment of Cancer

[0073] Disclosed herein are methods for reducing growth of cancer cells. The methods provide for decreasing the number of cancer cells bearing a specific marker or combination of markers, as provided herein, decreasing expression of a gene that is differentially expressed in a cancer cell, or decreasing the level of and/or decreasing an activity of a cancer-associated polypeptide. In general, the methods comprise contacting a cancer cell with a binding agent, e.g. an antibody or ligand specific for a marker or combination of markers provided herein.

[0074] "Reducing growth of cancer cells" includes, but is not limited to, reducing proliferation of cancer cells, and reducing the incidence of a non-cancerous cell becoming a cancerous cell. Whether a reduction in cancer cell growth has been achieved can be readily determined using any known assay, including, but not limited to, [³H]-thymidine incorporation; counting cell number over a period of time; detecting and/or measuring a marker associated with AML, etc.

[0075] Disclosed herein are methods for treating cancer, generally comprising administering to an individual in need thereof a substance that reduces cancer cell growth, in an amount sufficient to reduce cancer cell growth and treat the cancer. Whether a substance, or a specific amount of the substance, is effective in treating cancer can be assessed using any of a variety of known diagnostic assays for cancer, including, but not limited to biopsy, contrast radiographic studies, CAT scan, and detection of a tumor marker associated with cancer in the

blood of the individual. The substance can be administered systemically or locally, usually systemically.

[0076] A substance, e.g. a chemotherapeutic drug that reduces cancer cell growth, can be targeted to a cancer cell. Thus, in some instances, disclosed herein is a method of delivering a drug to a cancer cell, comprising administering a drug-antibody complex to a subject, wherein the antibody is specific for a cancer-associated polypeptide, and the drug is one that reduces cancer cell growth, a variety of which are known in the art. Targeting can be accomplished by coupling (e.g., linking, directly or via a linker molecule, either covalently or non-covalently, so as to form a drug-antibody complex) a drug to an antibody specific for a cancer-associated polypeptide. Methods of coupling a drug to an antibody are well known in the art and need not be elaborated upon herein.

Staging and Diagnosis

[0077] Acute myeloid leukemias are staged by analysis of the presence of cancer stem cells. Staging is useful for prognosis and treatment. In one instance, a sample from an acute myeloid leukemia patient is stained with reagents specific for a marker or combination of markers of the present disclosure. The analysis of staining patterns provides the relative distribution of AMLSC, which distribution predicts the stage of leukemia. In some instances, the sample is analyzed by histochemistry, including immunohistochemistry, *in situ* hybridization, and the like, for the presence of CD34⁺CD38⁻ cells that express a marker or combination of markers of the present disclosure. The presence of such cells indicates the presence of AMLSC.

[0078] In one instance, the patient sample is compared to a control, or a standard test value. In another instance, the patient sample is compared to a pre-leukemia sample, or to one or more time points through the course of the disease.

[0079] Samples, including tissue sections, slides, *etc.* containing an acute myeloid leukemia tissue, are stained with reagents specific for markers that indicate the presence of cancer stem cells. Samples may be frozen, embedded, present in a tissue microarray, and the like. The reagents, e.g. antibodies, polynucleotide probes, *etc.* may be detectably labeled, or may be indirectly labeled in the staining procedure. The data provided herein demonstrate that the number and distribution of progenitor cells is diagnostic of the stage of the leukemia.

[0080] The information thus derived is useful in prognosis and diagnosis, including susceptibility to acceleration of disease, status of a diseased state and response to changes in the environment, such as the passage of time, treatment with drugs or other modalities. The cells can also be classified as to their ability to respond to therapeutic agents and treatments, isolated for research purposes, screened for gene expression, and the like. The clinical samples can be further characterized by genetic analysis, proteomics, cell surface staining, or other means, in order to determine the presence of markers that are useful in classification.

For example, genetic abnormalities can be causative of disease susceptibility or drug responsiveness, or can be linked to such phenotypes.

DIFFERENTIAL CELL ANALYSIS

[0081] The presence of AMLSC in a patient sample can be indicative of the stage of the leukemia. In addition, detection of AMLSC can be used to monitor response to therapy and to aid in prognosis. The presence of AMLSC can be determined by quantitating the cells having the phenotype of the stem cell. In addition to cell surface phenotyping, it may be useful to quantitate the cells in a sample that have a "stem cell" character, which may be determined by functional criteria, such as the ability to self-renew, to give rise to tumors *in vivo*, e.g. in a xenograft model, and the like.

[0082] Clinical samples for use in the methods of the disclosure may be obtained from a variety of sources, particularly blood, although in some instances samples such as bone marrow, lymph, cerebrospinal fluid, synovial fluid, and the like may be used. Such samples can be separated by centrifugation, elutriation, density gradient separation, apheresis, affinity selection, panning, FACS, centrifugation with Hypaque, *etc.* prior to analysis, and usually a mononuclear fraction (PBMC) will be used. Once a sample is obtained, it can be used directly, frozen, or maintained in appropriate culture medium for short periods of time. Various media can be employed to maintain cells. The samples may be obtained by any convenient procedure, such as the drawing of blood, venipuncture, biopsy, or the like. Usually a sample will comprise at least about 10^2 cells, more usually at least about 10^3 cells, and preferable 10^4 , 10^5 or more cells. Typically the samples will be from human patients, although animal models may find use, e.g. equine, bovine, porcine, canine, feline, rodent, e.g. mice, rats, hamster, primate, *etc.*

[0083] An appropriate solution may be used for dispersion or suspension of the cell sample. Such solution will generally be a balanced salt solution, e.g. normal saline, PBS, Hank's balanced salt solution, *etc.*, conveniently supplemented with fetal calf serum or other naturally occurring factors, in conjunction with an acceptable buffer at low concentration, generally from 5-25 mM. Convenient buffers include HEPES, phosphate buffers, lactate buffers, *etc.*

[0084] Analysis of the cell staining will use conventional methods. Techniques providing accurate enumeration include fluorescence activated cell sorters, which can have varying degrees of sophistication, such as multiple color channels, low angle and obtuse light scattering detecting channels, impedance channels, *etc.* The cells may be selected against dead cells by employing dyes associated with dead cells (e.g. propidium iodide).

[0085] The affinity reagents may be specific receptors or ligands for the cell surface molecules indicated above. In addition to antibody reagents, peptide-MHC antigen and T cell receptor pairs may be used; peptide ligands and receptors; effector and receptor molecules, and the

like. Antibodies and T cell receptors may be monoclonal or polyclonal, and may be produced by transgenic animals, immunized animals, immortalized human or animal B-cells, cells transfected with DNA vectors encoding the antibody or T cell receptor, *etc.* The details of the preparation of antibodies and their suitability for use as specific binding members are well-known to those skilled in the art.

[0086] Of particular interest is the use of antibodies as affinity reagents. Conveniently, these antibodies are conjugated with a label for use in separation. Labels include magnetic beads, which allow for direct separation, biotin, which can be removed with avidin or streptavidin bound to a support, fluorochromes, which can be used with a fluorescence activated cell sorter, or the like, to allow for ease of separation of the particular cell type. Fluorochromes that find use include phycobiliproteins, e.g. phycoerythrin and allophycocyanins, fluorescein and Texas red. Frequently each antibody is labeled with a different fluorochrome, to permit independent sorting for each marker.

[0087] The antibodies are added to a suspension of cells, and incubated for a period of time sufficient to bind the available cell surface antigens. The incubation will usually be at least about 5 minutes and usually less than about 30 minutes. It is desirable to have a sufficient concentration of antibodies in the reaction mixture, such that the efficiency of the separation is not limited by lack of antibody. The appropriate concentration is determined by titration. The medium in which the cells are separated will be any medium that maintains the viability of the cells. A preferred medium is phosphate buffered saline containing from 0.1 to 0.5% BSA. Various media are commercially available and may be used according to the nature of the cells, including Dulbecco's Modified Eagle Medium (dMEM), Hank's Basic Salt Solution (HBSS), Dulbecco's phosphate buffered saline (dPBS), RPMI, Iscove's medium, PBS with 5 mM EDTA, *etc.*, frequently supplemented with fetal calf serum, BSA, HSA, *etc.*

[0088] The labeled cells are then quantitated as to the expression of cell surface markers as previously described.

[0089] The comparison of a differential progenitor analysis obtained from a patient sample, and a reference differential progenitor analysis is accomplished by the use of suitable deduction protocols, AI systems, statistical comparisons, *etc.* A comparison with a reference differential progenitor analysis from normal cells, cells from similarly diseased tissue, and the like, can provide an indication of the disease staging. A database of reference differential progenitor analyses can be compiled. An analysis of particular interest tracks a patient, e.g. in the chronic and pre-leukemic stages of disease, such that acceleration of disease is observed at an early stage. The methods of the disclosure provide detection of acceleration prior to onset of clinical symptoms, and therefore allow early therapeutic intervention, e.g. initiation of chemotherapy, increase of chemotherapy dose, changing selection of chemotherapeutic drug, and the like.

AMLSC Compositions

[0090] AMLSC may be separated from a complex mixture of cells by techniques that enrich for cells that differentially express a marker or combination of markers of the present disclosure. For isolation of cells from tissue, an appropriate solution may be used for dispersion or suspension. Such solution will generally be a balanced salt solution, e.g. normal saline, PBS, Hank's balanced salt solution, etc., conveniently supplemented with fetal calf serum or other naturally occurring factors, in conjunction with an acceptable buffer at low concentration, generally from 5-25 mM. Convenient buffers include HEPES, phosphate buffers, lactate buffers, etc.

[0091] The separated cells may be collected in any appropriate medium that maintains the viability of the cells, usually having a cushion of serum at the bottom of the collection tube. Various media are commercially available and may be used according to the nature of the cells, including dMEM, HBSS, dPBS, RPMI, Iscove's medium, etc., frequently supplemented with fetal calf serum.

[0092] Compositions highly enriched for AMLSC are achieved in this manner. The subject population may be at or about 50% or more of the cell composition, and preferably be at or about 75% or more of the cell composition, and may be 90% or more. The desired cells are identified by their surface phenotype, by the ability to self-renew, ability to form tumors, etc. The enriched cell population may be used immediately, or may be frozen at liquid nitrogen temperatures and stored for long periods of time, being thawed and capable of being reused. The cells may be stored in 10% DMSO, 90% FCS medium. The population of cells enriched for AMLSC may be used in a variety of screening assays and cultures, as described below.

[0093] The enriched AMLSC population may be grown *in vitro* under various culture conditions. Culture medium may be liquid or semi-solid, e.g. containing agar, methylcellulose, etc. The cell population may be conveniently suspended in an appropriate nutrient medium, such as Iscove's modified DMEM or RPMI-1640, normally supplemented with fetal calf serum (about 5-10%), L-glutamine, a thiol, particularly 2-mercaptoethanol, and antibiotics, e.g. penicillin and streptomycin.

[0094] The culture may contain growth factors to which the cells are responsive. Growth factors, as defined herein, are molecules capable of promoting survival, growth and/or differentiation of cells, either in culture or in the intact tissue, through specific effects on a transmembrane receptor. Growth factors include polypeptides and non-polypeptide factors. A wide variety of growth factors may be used in culturing the cells, e.g. LIF, steel factor (c-kit ligand), EGF, insulin, IGF, Flk-2 ligand, IL-11, IL-3, GM-CSF, erythropoietin, thrombopoietin, etc.

[0095] In addition to, or instead of growth factors, the subject cells may be grown in a co-culture with fibroblasts, stromal or other feeder layer cells. Stromal cells suitable for use in the growth of hematopoietic cells are known in the art. These include bone marrow stroma as used in "Whitlock-Witte" (Whitlock et al. [1985] Annu Rev Immunol 3:213-235) or "Dexter" culture conditions (Dexter et al. [1977] J Exp Med 145:1612-1616); and heterogeneous thymic stromal

cells.

Screening Assays

[0096] AMLSC expressing a marker or combination of markers of the present disclosure are also useful for *in vitro* assays and screening to detect factors and chemotherapeutic agents that are active on cancer stem cells. Of particular interest are screening assays for agents that are active on human cells. A wide variety of assays may be used for this purpose, including immunoassays for protein binding; determination of cell growth, differentiation and functional activity; production of factors; and the like. In other instances, isolated polypeptides corresponding to a marker or combination of markers of the present disclosure are useful in drug screening assays.

[0097] In screening assays for biologically active agents, anti-proliferative drugs, *etc.* the marker or AMLSC composition is contacted with the agent of interest, and the effect of the agent assessed by monitoring output parameters on cells, such as expression of markers, cell viability, and the like; or binding efficacy or effect on enzymatic or receptor activity for polypeptides. The cells may be freshly isolated, cultured, genetically altered, and the like. The cells may be environmentally induced variants of clonal cultures: e.g. split into independent cultures and grown under distinct conditions, for example with or without drugs; in the presence or absence of cytokines or combinations thereof. The manner in which cells respond to an agent, particularly a pharmacologic agent, including the timing of responses, is an important reflection of the physiologic state of the cell.

[0098] Parameters are quantifiable components of cells, particularly components that can be accurately measured, desirably in a high throughput system. A parameter can be any cell component or cell product including cell surface determinant, receptor, protein or conformational or posttranslational modification thereof, lipid, carbohydrate, organic or inorganic molecule, nucleic acid, e.g. mRNA, DNA, *etc.* or a portion derived from such a cell component or combinations thereof. While most parameters will provide a quantitative readout, in some instances a semi-quantitative or qualitative result will be acceptable. Readouts may include a single determined value, or may include mean, median value or the variance, *etc.* Characteristically a range of parameter readout values will be obtained for each parameter from a multiplicity of the same assays. Variability is expected and a range of values for each of the set of test parameters will be obtained using standard statistical methods with a common statistical method used to provide single values.

[0099] Agents of interest for screening include known and unknown compounds that encompass numerous chemical classes, primarily organic molecules, which may include organometallic molecules, inorganic molecules, genetic sequences, *etc.* An important aspect of the disclosure is to evaluate candidate drugs, including toxicity testing; and the like.

[0100] In addition to complex biological agents candidate agents include organic molecules

comprising functional groups necessary for structural interactions, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, frequently at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules, including peptides, polynucleotides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

[0101] Included are pharmacologically active drugs, genetically active molecules, *etc.* Compounds of interest include chemotherapeutic agents, hormones or hormone antagonists, *etc.* Exemplary of pharmaceutical agents suitable for this invention are those described in, "The Pharmacological Basis of Therapeutics," Goodman and Gilman, McGraw-Hill, New York, New York, (1996), Ninth edition, under the sections: Water, Salts and Ions; Drugs Affecting Renal Function and Electrolyte Metabolism; Drugs Affecting Gastrointestinal Function; Chemotherapy of Microbial Diseases; Chemotherapy of Neoplastic Diseases; Drugs Acting on Blood-Forming organs; Hormones and Hormone Antagonists; Vitamins, Dermatology; and Toxicology. Also included are toxins, and biological and chemical warfare agents, for example see Somani, S.M. (Ed.), "Chemical Warfare Agents," Academic Press, New York, 1992).

[0102] Test compounds include all of the classes of molecules described above, and may further comprise samples of unknown content. Of interest are complex mixtures of naturally occurring compounds derived from natural sources such as plants. While many samples will comprise compounds in solution, solid samples that can be dissolved in a suitable solvent may also be assayed. Samples of interest include environmental samples, e.g. ground water, sea water, mining waste, *etc.*; biological samples, e.g. lysates prepared from crops, tissue samples, *etc.*; manufacturing samples, e.g. time course during preparation of pharmaceuticals; as well as libraries of compounds prepared for analysis; and the like. Samples of interest include compounds being assessed for potential therapeutic value, *i.e.* drug candidates.

[0103] The term "samples" also includes the fluids described above to which additional components have been added, for example components that affect the ionic strength, pH, total protein concentration, *etc.* In addition, the samples may be treated to achieve at least partial fractionation or concentration. Biological samples may be stored if care is taken to reduce degradation of the compound, e.g. under nitrogen, frozen, or a combination thereof. The volume of sample used is sufficient to allow for measurable detection, usually from about 0.1 to 1 ml of a biological sample is sufficient.

[0104] Compounds, including candidate agents, are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds, including biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and

biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, *etc.* to produce structural analogs.

[0105] Agents are screened for biological activity by adding the agent to at least one and usually a plurality of cell samples, usually in conjunction with cells lacking the agent. The change in parameters in response to the agent is measured, and the result evaluated by comparison to reference cultures, e.g. in the presence and absence of the agent, obtained with other agents, *etc.*

[0106] The agents are conveniently added in solution, or readily soluble form, to the medium of cells in culture. The agents may be added in a flow-through system, as a stream, intermittent or continuous, or alternatively, adding a bolus of the compound, singly or incrementally, to an otherwise static solution. In a flow-through system, two fluids are used, where one is a physiologically neutral solution, and the other is the same solution with the test compound added. The first fluid is passed over the cells, followed by the second. In a single solution method, a bolus of the test compound is added to the volume of medium surrounding the cells. The overall concentrations of the components of the culture medium should not change significantly with the addition of the bolus, or between the two solutions in a flow through method.

[0107] Preferred agent formulations do not include additional components, such as preservatives, that may have a significant effect on the overall formulation. Thus preferred formulations consist essentially of a biologically active compound and a physiologically acceptable carrier, e.g. water, ethanol, DMSO, *etc.* However, if a compound is liquid without a solvent, the formulation may consist essentially of the compound itself.

[0108] A plurality of assays may be run in parallel with different agent concentrations to obtain a differential response to the various concentrations. As known in the art, determining the effective concentration of an agent typically uses a range of concentrations resulting from 1:10, or other log scale, dilutions. The concentrations may be further refined with a second series of dilutions, if necessary. Typically, one of these concentrations serves as a negative control, *i.e.* at zero concentration or below the level of detection of the agent or at or below the concentration of agent that does not give a detectable change in the phenotype.

[0109] Various methods can be utilized for quantifying the presence of the selected markers. For measuring the amount of a molecule that is present, a convenient method is to label a molecule with a detectable moiety, which may be fluorescent, luminescent, radioactive, enzymatically active, *etc.*, particularly a molecule specific for binding to the parameter with high affinity. Fluorescent moieties are readily available for labeling virtually any biomolecule, structure, or cell type. Immunofluorescent moieties can be directed to bind not only to specific proteins but also specific conformations, cleavage products, or site modifications like phosphorylation. Individual peptides and proteins can be engineered to autofluoresce, e.g. by expressing them as green fluorescent protein chimeras inside cells (for a review see Jones et

al. (1999) Trends Biotechnol. 17(12):477-81). Thus, antibodies can be genetically modified to provide a fluorescent dye as part of their structure. Depending upon the label chosen, parameters may be measured using other than fluorescent labels, using such immunoassay techniques as radioimmunoassay (RIA) or enzyme linked immunosorbance assay (ELISA), homogeneous enzyme immunoassays, and related non-enzymatic techniques. The quantitation of nucleic acids, especially messenger RNAs, is also of interest as a parameter. These can be measured by hybridization techniques that depend on the sequence of nucleic acid nucleotides. Techniques include polymerase chain reaction methods as well as gene array techniques. See Current Protocols in Molecular Biology, Ausubel et al., eds, John Wiley & Sons, New York, NY, 2000; Freeman et al. (1999) Biotechniques 26(1):112-225; Kawamoto et al. (1999) Genome Res 9(12):1305-12; and Chen et al. (1998) Genomics 51(3):313-24, for examples.

Depletion of AMLSC

[0110] Depletion of AMLSC is useful in the treatment of AML. Depletion can be achieved by several methods. Depletion is defined as a reduction in the target population by up to about 30%, or up to about 40%, or up to about 50%, or up to about 75% or more. An effective depletion is usually determined by the sensitivity of the particular disease condition to the levels of the target population. Thus in the treatment of certain conditions a depletion of even about 20% could be beneficial.

[0111] A marker-specific agent that specifically depletes the targeted AMLSC is used to contact the patient blood *in vitro* or *in vivo*, wherein after the contacting step, there is a reduction in the number of viable AMLSC in the targeted population. An exemplary agent for such purposes is an antibody that specifically binds to a marker or combination of markers of the present disclosure on the surface of the targeted AMLSC. An effective dose of antibodies for such a purpose is sufficient to decrease the targeted population to the desired level, for example as described above. Antibodies for such purposes may have low antigenicity in humans or may be humanized antibodies.

[0112] In one instance, antibodies for depleting target population are added to patient blood *in vivo*. In another instance, the antibodies are added to the patient blood *ex vivo*. Beads coated with the antibody of interest can be added to the blood, target cells bound to these beads can then be removed from the blood using procedures common in the art. In one instance the beads are magnetic and are removed using a magnet. Alternatively, when the antibody is biotinylated, it is also possible to indirectly immobilize the antibody onto a solid phase which has adsorbed avidin, streptavidin, or the like. The solid phase, usually agarose or sepharose beads are separated from the blood by brief centrifugation. Multiple methods for tagging antibodies and removing such antibodies and any cells bound to the antibodies are routine in the art. Once the desired degree of depletion has been achieved, the blood is returned to the patient. Depletion of target cells *ex vivo* decreases the side effects such as infusion reactions associated with the intravenous administration. An additional advantage is that the repertoire of

available antibodies is expanded significantly as this procedure does not have to be limited to antibodies with low antigenicity in humans or humanized antibodies.

[0113] In some embodiments, the antibodies for depletion are bispecific antibodies. "Bispecific antibody" and "bispecific antibodies," also known as bifunctional antibodies, refers to antibodies that recognize two different antigens by virtue of possessing at least one first antigen combining site specific for a first antigen or hapten, and at least one second antigen combining site specific for a second antigen or hapten. Such antibodies can be produced by recombinant DNA methods or include, but are not limited to, antibodies produced chemically by methods known in the art. Bispecific antibodies include all antibodies or conjugates of antibodies, or polymeric forms of antibodies which are capable of recognizing two different antigens. Bispecific antibodies include antibodies that have been reduced and reformed so as to retain their bivalent characteristics and to antibodies that have been chemically coupled so that they can have several antigen recognition sites for each antigen.

[0114] Bispecific antibodies for use in the methods of the present disclosure bind to at least one of the AMLSC antigens described herein, and may bind to two or more AMLSC antigens described herein. Antigen combinations of interest include, without limitation, CD47 + CD96, CD47 + CD99, CD47 + Tim3, CD47 + CD97. In some embodiments, one specificity of the antibody has a low affinity, e.g. less than about 10^{-9} binding constant, usually less than about 10^{-8} binding constant, and may be more than about 10^{-7} binding constant.

[0115] Antibodies suitable for practicing the methods of the disclosure are preferably monoclonal and multivalent, and may be human, humanized or chimeric antibodies, comprising single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, and/or binding fragments of any of the above. In certain instances, the antibodies are human antigen-binding antibody fragments of the present disclosure and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V_L or V_H domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH₁, CH₂, CH₃ and CL domains. Also included in the disclosure are antigen-binding fragments comprising any combination of variable region(s) with a hinge region, CH₁, CH₂, CH₃ and CL domains. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, sheep, rabbit, goat, guinea pig, camelid, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries, from human B cells, or from animals transgenic for one or more human immunoglobulins.

[0116] The antibodies suitable for practicing the methods of the present disclosure may be bispecific, trispecific or of greater multispecificity. Further, the antibodies of the present disclosure may have low risk of toxicity against granulocyte (neutrophil), NK cells, and CD4⁺ cells as bystander cells.

[0117] Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein et al., *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker et al, *EMBO J.*, 10:3655-3659 (1991).

[0118] According to another approach described in WO96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. Such interfaces may comprise at least a part of the CH₃ domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers. An alternative method links two different single chain variable regions to heat stable antigen (HSA). Using HSA as linker increases serum half life, and has the benefit of low immunogenicity.

[0119] Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Pat. No. 4,676,980, along with a number of cross-linking techniques.

[0120] Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science*, 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoet-hylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0121] Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kos-telny et al., J. Immunol, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber et al., J. Immunol., 152:5368 (1994). Alternatively, the antibodies can be "linear antibodies" as described in Zapata et al. Protein Eng. 8(10): 1057-1062 (1995). Briefly, these antibodies comprise a pair of tandem Fd segments (V_H - C_{H1} - V_H - C_{H1}) which form a pair of antigen binding regions. Linear antibodies can be bispecific or monospecific.

[0122] Within the context of the present invention, antibodies are understood to include monoclonal antibodies and polyclonal antibodies, antibody fragments (e.g., Fab and $F(ab')_2$), chimeric antibodies bifunctional or bispecific antibodies and tetrameric antibody complexes. Antibodies are understood to be reactive against a selected antigen on the surface of a T cell if they bind with an appropriate affinity (association constant), e.g. greater than or equal to $10^7 M^{-1}$. Additionally, antibodies that may be used in the methods of the present disclosure may also be described or specified in terms of their binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-2} M$, $10^{-2} M$, $5 \times 10^{-3} M$, $10^{-3} M$, $5 \times 10^{-4} M$, $10^{-4} M$, $5 \times 10^{-5} M$, $10^{-5} M$, $5 \times 10^{-6} M$, $10^{-6} M$, $5 \times 10^{-7} M$, $10^{-7} M$, $5 \times 10^{-8} M$, $10^{-8} M$, $5 \times 10^{-9} M$, $10^{-9} M$, $5 \times 10^{-10} M$, $10^{-9} M$, $5 \times 10^{-11} M$, $10^{-11} M$, $5 \times 10^{-12} M$, $10^{-12} M$, $5 \times 10^{-13} M$, $10^{-13} M$, $5 \times 10^{-14} M$, $10^{-14} M$, $5 \times 10^{-15} M$, $10^{-15} M$.

[0123] Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for the whole antibodies. For example, $F(ab')_2$ fragments can be generated by treating antibody with pepsin. The resulting $F(ab')_2$ fragment can be treated to reduce disulfide bridges to produce Fab' fragments.

[0124] The invention also contemplates chimeric antibody derivatives, i.e., antibody molecules that combine a non-human animal variable region and a human constant region. Chimeric antibody molecules can include, for example, the antigen binding domain from an antibody of a mouse, rat, or other species, with human constant regions. A variety of approaches for making chimeric antibodies have been described and can be used to make chimeric antibodies containing the immunoglobulin variable region which recognizes the selected antigens on the

surface of differentiated cells or tumor cells. See, for example, Morrison et al., 1985; Proc. Natl. Acad. Sci. U.S.A. 81,6851; Takeda et al., 1985, Nature 314:452; Cabilly et al., U.S. Pat. No. 4,816,567; Boss et al., U.S. Pat. No. 4,816,397; Tanaguchi et al., European Patent Publication EP171496; European Patent Publication 0173494, United Kingdom patent GB 2177096B.

[0125] Chemical conjugation is based on the use of homo- and heterobifunctional reagents with E-amino groups or hinge region thiol groups. Homobifunctional reagents such as 5,5'-Dithiobis(2-nitrobenzoic acid)(DNTB) generate disulfide bonds between the two Fabs, and 0-phenylenedimaleimide (O-PDM) generate thioether bonds between the two Fabs (Brenner et al., 1985, Glennie et al., 1987). Heterobifunctional reagents such as N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP) combine exposed amino groups of antibodies and Fab fragments, regardless of class or isotype (Van Dijk et al., 1989).

[0126] The antibodies of the disclosure, i.e., antibodies that are useful for treating cancers, as well as other cancer comprising cancer stem cells expressing antigens set forth in Table 1, include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from binding to the antigens. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[0127] Kits may be provided, where the kit will comprise staining reagents that are sufficient to differentially identify the AMLSC described herein. A combination of interest may include one or more reagents specific for a marker or combination of markers of the present disclosure, and may further include antibodies specific for CD96, CD34, and CD38. The staining reagents are preferably antibodies, and may be detectably labeled. Kits may also include tubes, buffers, etc., and instructions for use.

[0128] It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, animal species or genera, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which will be limited only by the appended claims.

[0129] As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the culture" includes reference to one or more cultures and equivalents thereof known to those skilled in the art, and so forth. All technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs unless clearly indicated otherwise.

EXPERIMENTAL

Example 1

Identification of Cell Surface Molecules Preferentially Expressed on Human Acute Myeloid Leukemia Stem Cells Compared to Their Normal Counterparts

[0130] *Prospective Identification of a Human Multipotent Progenitor, the Cell of Origin for AML LSC.* Identification of cell surface molecules that are preferentially expressed on AML LSC would be greatly facilitated by determining the cell within the normal hematopoietic hierarchy that undergoes transformation to become an AML LSC. The prevailing view in the field has been that AML LSC arise out of hematopoietic stem cells (HSC), since both stem cell populations are enriched in Lin-CD34+CD38- cells. However, human HSC have been shown to express CD90, while AML LSC are CD90-. Furthermore, HSC from long-term remission t(8;21) AML patients were found to contain the AML1-ETO translocation product, suggesting that the HSC were pre-leukemic, and that full transformation to AML LSC occurred in a downstream progenitor.

[0131] While it is certainly possible that HSC are in fact the cell of origin for AML LSC, and that these cells lose expression of CD90 as a consequence of transformation, it is also possible that AML LSC originate from downstream Lin-CD34+CD38-CD90- cells. We utilized a NOD/SCID/IL-2R gamma null (NOG) newborn xenotransplantation model to assay the function of subpopulations of Lin-CD34+CD38- cord blood, identified on the basis of CD90 and CD45RA expression. Lin-CD34+CD38-CD90+ cells produced long-term multi-lineage engraftment and formed successful secondary transplants, and therefore contained HSC. Transplantation of purified Lin-CD34+CD38-CD90-CD45RA- cells resulted in lower levels of multi-lineage engraftment in primary recipients, and a statistically significant reduced ability to form long-term secondary transplants. In fact, with transplantation of 50 purified cells, these cells failed to long-term engraft, unlike the Lin-CD34+CD38-CD90+ HSC. Thus, Lin-CD34+CD38-CD90-CD45RA- cells are multipotent and possess limited self-renewal ability. These cells are termed multipotent progenitors (MPP) and represent the possible cell of origin of AML LSC.

[0132] *Use of Gene Expression Profiling to Identify Cell Surface Molecules Preferentially Expressed on AML LSC Compared to Their Normal Counterparts, HSC and MPP.* Cell surface molecules preferentially expressed on human acute myeloid leukemia stem cells (AML LSC) compared to their normal counterparts have therapeutic applications outlined below. One strategy to identify such molecules has been to generate gene expression profiles of AML LSC and normal HSC and MPP, and compare them for differentially expressed genes.

[0133] Normal bone marrow HSC and MPP (n=4) and AML LSC (n=9) were purified by FACS. Total RNA was prepared, amplified, and hybridized to Affymetrix human DNA microarrays. Statistical analysis identified 4037 genes differentially expressed between HSC and LSC, and 4208 genes differentially expressed between MPP and LSC, with $p < 0.05$ and a False Discovery Rate of 5% (Figure 1A). Investigation of these differentially expressed genes identified 288 and 318 cell surface molecules preferentially expressed in AML LSC by at least 2-fold compared to HSC and MPP, respectively. Selected members of this list, including many with the greatest preferential expression in AML LSC are indicated (Figure 1B, Table 1).

Table 1

Fold Change	Genbank	Gene Symbol	Description
94.34	M27331	TRGC2	T cell receptor gamma constant 2
57.47	NM_005816	CD96	CD96 antigen
47.17	AI862120	MAMDC2	MAM domain containing 2
32.36	AF348078	SUCNR1	succinate receptor 1
32.05	M16768	TRGC2	T cell receptor gamma constant 2
30.96	NM_002182	IL1RAP	interleukin 1 receptor accessory protein
29.85	M13231	TRGC2	T cell receptor gamma constant 2
27.55	NM_003332	TYROBP	TYRO protein tyrosine kinase binding protein
26.88	NM_004271	LY86	lymphocyte antigen 86
20.96	NM_014879	P2RY14	purinergic receptor P2Y, G-protein coupled, 14
18.38	BC020749	CD96	CD96 antigen
18.38	NM_005048	PTHR2	parathyroid hormone receptor 2
17.73	AI625747	ADRB1	Adrenergic, beta-1-, receptor
17.36	NM_015376	RASGRP3	RAS guanyl releasing protein 3 (calcium and DAG-regulated)
16.84	U62027	C3AR1	complement component 3a receptor 1
14.49	AW025572	HAVCR2	hepatitis A virus cellular receptor 2
12.48	AF285447	HCST	hematopoietic cell signal transducer
11.92	AI805323	LGR7	leucine-rich repeat-containing G protein-coupled receptor 7
11.67	NM_001197	BIK	BCL2-interacting killer (apoptosis-inducing)
11.53	NM_018092	NETO2	neuropilin (NRP) and tolloid (TLL)-like 2
11.07	N74607	AQP3	aquaporin 3
10.88	BF439675	CD69	CD69 antigen (p60, early T-cell activation antigen)
10.48	NM_001769	CD9	CD9 antigen (p24)
10.32	AF167343	IL1RAP	interleukin 1 receptor accessory protein

Fold Change	Genbank	Gene Symbol	Description
9.52	AA814140	C5orf18	chromosome 5 open reading frame 18
8.77	NM_005582	CD180	CD180 antigen
7.46	AF039686	GPR34	G protein-coupled receptor 34
7.30	A1056776	ITGA6	Integrin, alpha 6
7.19	AJ277151	TNFRSF4	tumor necrosis factor receptor superfamily, member 4
6.99	AI738675	SELPLG	Selectin P ligand
6.85	AA888858	PDE3B	Phosphodiesterase 3B, cGMP-inhibited
6.80	AU149572	ADCY2	adenylate cyclase 2 (brain)
6.80	NM_002299	LCT	lactase
6.58	NM_005296	GPR23	G protein-coupled receptor 23
6.45	NM_004106	FCER1G	Fc fragment of IgE, high affinity receptor
6.29	AI741056	SELPLG	selectin P ligand
6.25	AW406569	MGC15619	
6.06	M81695	ITGAX	integrin, alpha X
5.92	NM_003494	DYSF	dysferlin
5.85	AI860212	PAG1	phosphoprotein associated with glycosphingolipid microdomains 1
5.75	NM_013447	EMR2	egf-like module containing, mucin-like, hormone receptor-like 2
5.62	NM_017806	LIME1	Lck interacting transmembrane adaptor 1
5.62	AK092824	AMN	Amnionless homolog (mouse)
5.59	AF345567	GPR174	G protein-coupled receptor 174
5.29	BC041928	IL1 RAP	Interleukin 1 receptor accessory protein
5.26	L03419	FCGR1A	Fc fragment of IgG, high affinity Ia, receptor (CD64) ; Fc-gamma receptor I B2
5.24	BG230586	SLC7A6	solute carrier family 7 (cationic amino acid transporter, y+ system), member 6
5.18	AF015524	CCRL2	chemokine (C-C motif) receptor-like 2
5.13	AA631143	SLC45A3	solute carrier family 45, member 3
5.10	AJ240085	TRAT1	T cell receptor associated transmembrane adaptor 1
5.05	AW183080	GPR92	G protein-coupled receptor 92
5.03	NM_002120	HLA-DOB	major histocompatibility complex, class II, DO beta
5.03	NM_015364	LY96	lymphocyte antigen 96

Fold Change	Genbank	Gene Symbol	Description
4.90	NM_020399	GOPC	golgi associated PDZ and coiled-coil motif containing
4.88	AK026133	SEMA4B	semaphorin
4.88	BC041664	VMD2	vitelliform macular dystrophy 2
4.85	NM_152592	C14orf49	chromosome 14 open reading frame 49
4.85	AA923524	RASGRP4	RAS guanyl releasing protein 4
4.85	BC008777	ITGAL	integrin, alpha L)
4.67	AF014403	PPAP2A	phosphatidic acid phosphatase type 2A
4.65	AK097698	SORCS2	Sortilin-related VPS10 domain containing receptor 2
4.63	X14355	FCGR1A	Fc fragment of IgG, high affinity Ia, receptor (CD64)
4.55	NM_001629	ALOX5AP	arachidonate 5-lipoxygenase-activating protein
4.50	AU155968	C18orf1	chromosome 18 open reading frame 1
4.44	AK075092	HERV-FRD	HERV-FRD provirus ancestral Env polyprotein
4.42	NM_020960	GPR107	G protein-coupled receptor 107
4.37	BC000039	FAM26B	family with sequence similarity 26, member B
4.35	NM_153701	IL12RB1	interleukin 12 receptor, beta 1
4.35	AI762344	PTGER1	prostaglandin E receptor 1 EP1), 42kDa
4.31	NM_006459	SPFH1	SPFH domain family, member 1
4.27	NM_003126	SPTA1	spectrin, alpha, erythrocytic 1 (elliptocytosis 2)
4.22	AL518391	AQP1	aquaporin 1 (channel-forming protein, 28kDa)
4.12	AK026188	PCDHGC3	protocadherin gamma subfamily C
4.10	AU146685	EDG2	Endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2
4.05	BE673587	SLC14A1	Solute carrier family 14 (urea transporter), member 1 (Kidd blood group)
4.02	BF129969	TSPAN2	tetraspanin 2
4.00	AW243272	KCNK5	Potassium channel, subfamily K, member 5
3.98	T68858	DHRS3	Dehydrogenase/reductase (SDR family) member 3
3.94	AI827849	VTI1A	Vesicle transport through interaction with t-SNAREs homolog 1A (yeast)
3.86	AL134012	NRXN2	Neurexin 2
3.83	BG230614	CD47	CD47 antigen

Fold Change	Genbank	Gene Symbol	Description
3.80	AI869717	MGC15523	MGC15523
3.80	AI458583	SIMP	Source of immunodominant MHC-associated peptides
3.79	NM_002183	IL3RA	interleukin 3 receptor, alpha (low affinity)
3.79	AA608820	NRXN2	neurexin 2
3.73	NM_000206	IL2RG	interleukin 2 receptor
3.72	BC002737	VAMP2	synaptobrevin 2
3.72	BC005884	BID	BH3 interacting domain death agonist ; BH3 interacting domain death agonist
3.68	AI688418	PLXNA2	plexin A2
3.68	BC003105	PTP4A3	protein tyrosine phosphatase type IVA, member 3
3.68	NM_001772	CD33	CD33 antigen (gp67)
3.65	BC007524	SPAG9	sperm associated antigen 9
3.64	AI344200	SLC25A35	solute carrier family 25, member 35
3.64	BC005253	KLHL20	kelch-like 20 (Drosophila)
3.60	AI335263	NETO2	neuropilin (NRP) and tolloid (TLL)-like 2
3.58	BF381837	C20orf52	chromosome 20 open reading frame 52
3.51	NM_002886	RAP2A	RAP2A
3.50	NM_007063	TBC1D8	TBC1 domain family, member 8 (with GRAM domain)
3.45	AK027160	BCL2L11	BCL2-like 11 (apoptosis facilitator)
3.44	BF055366	EDG2	endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2
3.42	NM_003608	GPR65	G protein-coupled receptor 65
3.41	AI675453	PLXNA3	plexin A3
3.40	AV734194	DPP8	dipeptidylpeptidase 8
3.38	BC000232	C5orf18	chromosome 5 open reading frame 18
3.36	BC001956	KIAA1961	KIAA1961 gene
3.34	NM_013332	HIG2	hypoxia-inducible protein 2
3.31	BC029450	SLC33A1	Solute carrier family 33 (acetyl-CoA transporter), member 1
3.30	AW008505	C18orf1	chromosome 18 open reading frame 1
3.29	BF693956	CD47	CD47 antigen
3.28	BF677986	KIAA1961	KIAA1961 gene
3.27	AI433691	CACNA2D4	calcium channel, voltage-dependent, alpha

Fold Change	Genbank	Gene Symbol	Description
			2/delta subunit 4
3.26	AB014573	NPHP4	nephronophthisis 4
3.25	AL582804	LY9	lymphocyte antigen 9
3.25	BG236280	CD86	CD86 antigen
3.24	AA639289	SLC26A7	Solute carrier family 26, member 7
3.24	NM_005211	CSF1R	colony stimulating factor 1 receptor
3.24	AI051254	TRPM2	transient receptor potential cation channel, subfamily M, member 2
3.23	AW292816	ABHD2	abhydrolase domain containing 2
3.23	BC040275	RASGRF1	Ras protein-specific guanine nucleotide-releasing factor 1
3.22	NM_021911	GABRB2	gamma-aminobutyric acid (GABA) A receptor, beta 2
3.19	AI660619	SLC7A6	solute carrier family 7 (cationic amino acid transporter, y+ system), member 6
3.19	NM_001860	SLC31A2	solute carrier family 31 (copper transporters), member 2
3.18	NM_015680	C2orf24	chromosome 2 open reading frame 24
3.17	AW058600	SLC36A1	solute carrier family 36
3.16	AU145049	HIP1	Huntingtin interacting protein 1
3.15	NM_005770	SERF2	small EDRK-rich factor 2
3.15	NM_003566	EEA1	Early endosome antigen 1, 162kD
3.14	NM_020041	SLC2A9	solute carrier family 2 (facilitated glucose transporter), member 9
3.14	W90718	SLC24A4	solute carrier family 24
3.13	AI423165	TICAM2	toll-like receptor adaptor molecule 2
3.12	AI674647	SPPL2A	signal peptide peptidase-like 2A
3.11	NM_004121	GGTLA1	gamma-glutamyltransferase-like activity 1
3.10	NM_004546	NDUFB2	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 2, 8kDa
3.05	X15786	RET	ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma 1, Hirschsprung disease)
3.05	AF181660	MPZL1	myelin protein zero-like 1
3.05	BG230614	CD47	CD47 antigen (Rh-related antigen, integrin-associated signal transducer)
3.00	AI571996	STAM2	signal transducing adaptor molecule (SH3

Fold Change	Genbank	Gene Symbol	Description
			domain and ITAM motif) 2
2.99	NM_000201	ICAM1	intercellular adhesion molecule 1 (CD54), human rhinovirus receptor
2.93	NM_025244	TSGA10	testis specific, 10
2.93	AU147538	PRKCE	Protein kinase C, epsilon
2.92	NM_024576	OGFRL1	opioid growth factor receptor-like 1
2.91	AI248055	ABCC4	ATP-binding cassette, sub-family C (CFTR/MRP), member 4
2.86	AA503877	CEPT1	Choline/ethanolamine phosphotransferase 1
2.84	BC030993	FLJ21127	Hypothetical protein FLJ21127
2.82	AA829818	LY86	Lymphocyte antigen 86
2.82	NM_001859	SLC31A1	solute carrier family 31 (copper transporters), member 1
2.81	M74721	CD79A	CD79A antigen (immunoglobulin-associated alpha)
2.79	AI986112	MGAT4B	Mannosyl (alpha-1,3-)-glycoprotein beta-1,4-N-acetylglucosaminyltransferase, isoenzyme B
2.79	NM_030930	UNC93B1	unc-93 homolog B1 (C. elegans) ; unc-93 homolog (C. elegans)
2.79	X74039	PLAUR	plasminogen activator, urokinase receptor
2.78	BF514291	LY86	Lymphocyte antigen 86
2.75	BC005253	KLHL20	kelch-like 20 (Drosophila)
2.73	AB036432	AGER	advanced glycosylation end product-specific receptor
2.71	NM_007245	ATXN2L	ataxin 2-like
2.71	NM_016072	GOLT1B	golgi transport 1 homolog B (S. cerevisiae)
2.71	AI453548	ZDHHC8	zinc finger, DHHC-type containing 8
2.70	AI636233	TMEM8	transmembrane protein 8 (five membrane-spanning domains)
2.69	BE502509	T3JAM	TRAF3 interacting protein 3
2.69	AW117765	PEX13	peroxisome biogenesis factor 13
2.69	AW052216	IL17RB	Interleukin 17 receptor B
2.67	NM_003853	IL18RAP	interleukin 18 receptor accessory protein
2.66	NM_002490	NDUFA6	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 6, 14kDa
2.65	NM_016639	TNFRSF12A	tumor necrosis factor receptor superfamily,

Fold Change	Genbank	Gene Symbol	Description
			member 12A
2.65	AI363185	FLJ20255	Hypothetical protein FLJ20255
2.65	NM_052931	SLAMF6	SLAM family member 6
2.65	AW571669	TNFRSF19L	tumor necrosis factor receptor superfamily, member 19-like
2.64	AA654142	CEECAM1	cerebral endothelial cell adhesion molecule 1
2.62	AW510783	TMEM63A	transmembrane protein 63A
2.61	W95007	ACSL4	Acyl-CoA synthetase long-chain family member 4
2.60	S76475	NTRK3	neurotrophic tyrosine kinase, receptor, type 3
2.60	AJ130713	SIGLEC7	sialic acid binding Ig-like lectin 7
2.56	NM_003775	EDG6	endothelial differentiation, G-protein-coupled receptor 6
2.55	AI978986	MAMDC4	MAM domain containing 4
2.54	AF010447	MR1	major histocompatibility complex, class I-related
2.54	NM_006068	TLR6	toll-like receptor 6
2.53	AF041811	NTRK3	neurotrophic tyrosine kinase, receptor, type 3
2.53	AW953521	SERF2 ; HYPK	small EDRK-rich factor 2 ; Huntingtin interacting protein K
2.51	AW293276	CD53	CD53 antigen
2.49	AK023058	PLXNA2	Plexin A2
2.49	AI125204	C6orf128	chromosome 6 open reading frame 128
2.49	NM_000392	ABCC2	ATP-binding cassette, sub-family C (CFTR/MRP), member 2
2.46	BC032474	TIRAP	toll-interleukin 1 receptor (TIR) domain containing adaptor protein
2.44	NM_031211	IMAA	SLC7A5 pseudogene
2.44	AI797836	CD5	CD5 antigen (p56-62)
2.41	W72082	C1QR1	complement component 1
2.40	AA708616	DPP9	dipeptidylpeptidase 9
2.40	BM987094	DLGAP4	discs, large (Drosophila) homolog-associated protein 4
2.40	AL713719	LOC283501	ATPase, Class VI, type 11A
2.39	AI628734	PRLR	prolactin receptor
2.39	NM_012110	CHIC2	cysteine-rich hydrophobic domain 2

Fold Change	Genbank	Gene Symbol	Description
2.38	AK022002	TFR2	transferrin receptor 2
2.37	NM_001555	IGSF1	immunoglobulin superfamily, member 1
2.36	AA426091	C19orf15	chromosome 19 open reading frame 15
2.36	BE547542	GOPC	golgi associated PDZ and coiled-coil motif containing
2.36	NM_004231	ATP6V1F	ATPase, H ⁺ transporting, lysosomal 14kDa, V1 subunit F
2.36	AJ130712	SIGLEC7	sialic acid binding Ig-like lectin 7
2.36	NM_017905	TMCO3	transmembrane and coiled-coil domains 3
2.35	AB054985	CACNB1	calcium channel, voltage-dependent, beta 1 subunit
2.35	NM_005003	NDUFAB1	NADH dehydrogenase (ubiquinone) 1, alpha/beta subcomplex, 1, 8kDa
2.35	NM_001251	CD68	CD68 antigen
2.35	AA700869	PSCD2	Pleckstrin homology, Sec7 and coiled-coil domains 2 (cytohesin-2)
2.35	U94903	CD44	CD44 antigen (homing function and Indian blood group system)
2.35	NM_003841	TNFRSF10C	tumor necrosis factor receptor superfamily, member 10c, decoy without an intracellular domain
2.33	NM_004541	NDUFA1	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 1, 7.5kDa
2.33	BE567130	KLRK1	Killer cell lectin-like receptor subfamily K, member 1
2.31	NM_017460	CYP3A4	cytochrome P450, family 3, subfamily A, polypeptide 4
2.31	AI339536	DSC1	Desmocollin 1
2.31	NM_001783	CD79A	CD79A antigen (immunoglobulin-associated alpha) ; antigen (immunoglobulin-associated alpha)
2.30	AA333161	VTI1A	vesicle transport through interaction with t-SNAREs homolog 1A (yeast)
2.30	AW134823	CD6	CD6 antigen ; CD6 antigen
2.30	AL137537	ATP8B2	ATPase, Class I, type 8B, member 2
2.29	AI671983	SLC2A9	solute carrier family 2 (facilitated glucose transporter), member 9
2.29	AA018187	C22orf3	chromosome 22 open reading frame 3
2.29	AL117415	ADAM33	ADAM metallopeptidase domain 33

Fold Change	Genbank	Gene Symbol	Description
2.29	NM_002588	PCDHGC3	protocadherin gamma subfamily C
2.29	NM_020960	GPR107	G protein-coupled receptor 107
2.29	AK074635	GENX-3414	Genethonin 1
2.29	BE138575	ITGB5	Integrin, beta 5
2.28	NM_003830	SIGLEC5	sialic acid binding Ig-like lectin 5 ; sialic acid binding Ig-like lectin 5
2.28	NM_013319	UBIAD1	UbiA prenyltransferase domain containing 1
2.28	M63889	FGFR1	fibroblast growth factor receptor 1 (fms-related tyrosine kinase 2, Pfeiffer syndrome)
2.27	H67156	MSCP	Solute carrier family 25, member 37
2.27	BC006215	SMEK2	KIAA1387 protein ; KIAA1387 protein
2.27	AL109653	SLITRK2	SLIT and NTRK-like family, member 2
2.27	NM_007011	ABHD2	abhydrolase domain containing 2
2.26	AI767210	MGC11332	Hypothetical protein MGC11332
2.26	BF723605	NRCAM	Neuronal cell adhesion molecule
2.26	R08129	CDA08	T-cell immunomodulatory protein
2.26	AF052059	SEL1L	sel-1 suppressor of lin-12-like (C. elegans)
2.26	NM_005729	PPIF	peptidylprolyl isomerase F (cyclophilin F)
2.25	BE858032	ARL2L1	ADP-ribosylation factor-like 2-like 1
2.25	AI950390	C14orf118	Chromosome 14 open reading frame 118
2.24	NM_017767	SLC39A4	solute carrier family 39 (zinc transporter), member 4
2.24	AL110273	SPTAN1	Spectrin, alpha, non-erythrocytic 1 (alpha-fodrin)
2.24	AI077660	CDA08	T-cell immunomodulatory protein
2.23	AA488687	SLC7A11	solute carrier family 7, (cationic amino acid transporter, y+ system) member 11
2.23	NM_000634	IL8RA	interleukin 8 receptor, alpha
2.22	AL390177	MGC34032	Solute carrier family 44, member 5
2.21	NM_001531	MR1	major histocompatibility complex, class I-related
2.21	NM_003183	ADAM17	ADAM metallopeptidase domain 17 (tumor necrosis factor, alpha, converting enzyme)
2.20	AC003999	SCAP2	src family associated phosphoprotein 2
2.20	BC014416	SLC33A1	solute carrier family 33 (acetyl-CoA transporter), member 1

Fold Change	Genbank	Gene Symbol	Description
2.20	AF226731	ADORA3	adenosine A3 receptor
2.19	AI608725	ICAM1	intercellular adhesion molecule 1 (CD54), human rhinovirus receptor
2.19	U41163	SLC6A8 ; FLJ43855	solute carrier family 6 (neurotransmitter transporter, creatine), member 8 ; similar to sodium- and chloride-dependent creatine transporter
2.19	AU147799	LRRC15	leucine rich repeat containing 15
2.18	AW337166	LOC255104	Transmembrane and coiled-coil domains 4
2.18	NM_006505	PVR	poliovirus receptor
2.18	AI638420	CLIC4	chloride intracellular channel 4
2.18	AI167482	SCUBE3	Signal peptide, CUB domain, EGF-like 3
2.18	AI739514	HAS3	hyaluronan synthase 3
2.18	NM_005971	FXYD3	FXYD domain containing ion transport regulator 3
2.17	AL022398	TRAF3IP3	TRAF3 interacting protein 3
2.17	U90940	FCGR2C	Fc fragment of IgG, low affinity IIc, receptor for (CD32)
2.16	BC023540	SORCS1	Sortilin-related VPS10 domain containing receptor 1
2.16	AV713913	OSTM1	osteopetrosis associated transmembrane protein 1
2.15	NM_024505	NOX5	NADPH oxidase, EF-hand calcium binding domain 5
2.15	BC006178	SEC22L3	SEC22 vesicle trafficking protein-like 3 (S. cerevisiae) ; SEC22 vesicle trafficking protein-like 3 (S. cerevisiae)
2.15	BG151527	GRIK5	glutamate receptor, ionotropic, kainate 5
2.14	AW001754	NEGR1	neuronal growth regulator 1
2.14	NM_013979	BNIP1	BCL2/adenovirus E1B 19kDa interacting protein 1
2.14	NM_018643	TREM1	triggering receptor expressed on myeloid cells 1
2.12	NM_005284	GPR6	G protein-coupled receptor 6
2.11	AA454190	ZDHHC20	zinc finger, DHHC-type containing 20
2.11	AB048796	TMPRSS13	transmembrane protease, serine 13
2.11	AL044520	NYD-SP21	testes development-related NYD-SP21
2.11	BE463930	TMAP1	Matrix-remodelling associated 7

Fold Change	Genbank	Gene Symbol	Description
2.10	NM_152264	SLC39A13	solute carrier family 39 (zinc transporter), member 13
2.08	AL530874	EPHB2	EPH receptor B2
2.07	NM_018668	VPS33B	vacuolar protein sorting 33B (yeast)
2.07	NM_024531	GPR172A	G protein-coupled receptor 172A
2.07	NM_023038	ADAM19	ADAM metalloproteinase domain 19 (meltrin beta)
2.07	BC001281	TNFRSF10B	tumor necrosis factor receptor superfamily, member 10b
2.07	AF217749	PCDHB9	protocadherin beta 9
2.06	AB030077	FGFR2	fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth factor receptor, craniofacial dysostosis 1, Crouzon syndrome, Pfeiffer syndrome, Jackson-Weiss syndrome)
2.06	AL137432	SUSD1	sushi domain containing 1
2.05	NM_004518	KCNQ2	potassium voltage-gated channel, KQT-like subfamily, member 2
2.04	AI672363	VPS33B	vacuolar protein sorting 33B (yeast)
2.04	NM_006671	SLC1A7	solute carrier family 1 (glutamate transporter), member 7
2.03	AA215519	DLGAP1	Discs, large (Drosophila) homolog-associated protein 1
2.02	NM_004648	PTPNS1	protein tyrosine phosphatase, non-receptor type substrate 1
2.02	NM_002564	P2RY2	purinergic receptor P2Y, G-protein coupled, 2
2.01	BF511678	SCUBE3	Signal peptide, CUB domain, EGF-like 3
2.01	BC013385	CLEC7A	C-type lectin domain family 7, member A

Table 2

Fold Change	Genbank	Gene Symbol	Description
57.47	NM_005816	CD96	CD96 antigen
32.36	AF348078	SUCNR1	succinate receptor 1
30.96	NM_002182	IL1 RAP	interleukin 1 receptor accessory protein
27.55	NM_003332	TYROBP	TYRO protein tyrosine kinase binding protein
26.88	NM_004271	LY86	lymphocyte antigen 86
20.96	NM_014879	P2RY14	purinergic receptor P2Y, G-protein coupled, 14

Fold Change	Genbank	Gene Symbol	Description
18.38	NM_005048	PTH2R	parathyroid hormone receptor 2
17.73	AI625747	ADRB1	Adrenergic, beta-1-, receptor
17.36	NM_015376	RASGRP3	RAS guanyl releasing protein 3 (calcium and
16.84	U62027	C3AR1	complement component 3a receptor 1
14.49	AW025572	HAVCR2	hepatitis A virus cellular receptor 2
12.48	AF285447	HCST	hematopoietic cell signal transducer
11.92	AI805323	LGR7	leucine-rich repeat-containing G protein-coupled receptor 7
11.67	NM_001197	BIK	BCL2-interacting killer (apoptosis-inducing)
11.53	NM_018092	NETO2	neuropilin (NRP) and tolloid (TLL)-like 2
11.07	N74607	AQP3	aquaporin 3
10.48	NM_001769	CD9	CD9 antigen (p24)
8.77	NM_005582	CD180	CD180 antigen
7.46	AF039686	GPR34	G protein-coupled receptor 34
7.19	AJ277151	TNFRSF4	tumor necrosis factor receptor superfamily, member 4
6.85	AA888858	PDE3B	Phosphodiesterase 3B, cGMP-inhibited
6.80	AU149572	ADCY2	adenylate cyclase 2 (brain)
6.80	NM_002299	LCT	lactase
6.58	NM_005296	GPR23	G protein-coupled receptor 23
6.45	NM_004106	FCER1G	Fc fragment of IgE, high affinity I, receptor for; gamma polypeptide
6.25	AW406569	MGC15619	hypothetical protein MGC15619
6.06	M81695	ITGAX	integrin, alpha X (antigen CD11C (p150), alpha polypeptide)
5.92	NM_003494	DYSF	dysferlin, limb girdle muscular dystrophy 2B recessive)
5.75	NM_013447	EMR2	egf-like module containing, mucin-like, hormone receptor-like 2
5.62	NM_017806	LIME1	Lck interacting transmembrane adaptor 1
5.62	AK092824	AMN	Amnionless homolog (mouse)
5.59	AF345567	GPR174	G protein-coupled receptor 174
5.26	L03419	FCGR1A ; LOC440607	Fc fragment of IgG, high affinity Ia, receptor (CD64) ; Fc-gamma receptor I B2
5.18	AF015524	CCRL2	chemokine (C-C motif) receptor-like 2
5.13	AA631143	SLC45A3	solute carrier family 45, member 3

Fold Change	Genbank	Gene Symbol	Description
5.10	AJ240085	TRAT1	T cell receptor associated transmembrane adaptor 1
5.05	AW183080	GPR92	G protein-coupled receptor 92
5.03	NM_002120	HLA-DOB	major histocompatibility complex, class II, DO
5.03	NM_015364	LY96	lymphocyte antigen 96

Table 3

Fold Change	Genbank	Gene Symbol	Description
57.47	NM_005816	CD96	CD96 antigen
32.36	AF348078	SUCNR1	succinate receptor 1
30.96	NM_002182	IL1 RAP	interleukin 1 receptor accessory protein
27.55	NM_003332	TYROBP	TYRO protein tyrosine kinase binding protein
26.88	NM_004271	LY86	lymphocyte antigen 86
20.96	NM_014879	P2RY14	purinergic receptor P2Y, G-protein coupled, 14
18.38	NM_005048	PTH2R	parathyroid hormone receptor 2
17.73	AI625747	ADRB1	Adrenergic, beta-1-, receptor
17.36	NM_015376	RASGRP3	RAS guanyl releasing protein 3 (calcium and
16.84	U62027	C3AR1	complement component 3a receptor 1
14.49	AW025572	HAVCR2	hepatitis A virus cellular receptor 2
12.48	AF285447	HCST	hematopoietic cell signal transducer
11.92	AI805323	LGR7	leucine-rich repeat-containing G protein-coupled receptor 7
11.67	NM_001197	BIK	BCL2-interacting killer (apoptosis-inducing)
11.53	NM_018092	NETO2	neuropilin (NRP) and tolloid (TLL)-like 2
11.07	N74607	AQP3	aquaporin 3
10.88	BF439675	CD69	CD69 antigen (p60, early T-cell activation antigen)
10.48	NM_001769	CD9	CD9 antigen (p24)
9.52	AA814140	C5orf18	chromosome 5 open reading frame 18
8.77	NM_005582	CD180	CD180 antigen
7.46	AF039686	GPR34	G protein-coupled receptor 34
7.30	AI056776	ITGA6	Integrin, alpha 6

Fold Change	Genbank	Gene Symbol	Description
7.19	AJ277151	TNFRSF4	tumor necrosis factor receptor superfamily, member 4
6.99	AI738675	SELPLG	Selectin P ligand
6.85	AA888858	PDE3B	Phosphodiesterase 3B, cGMP-inhibited
6.80	AU149572	ADCY2	adenylate cyclase 2 (brain)
6.80	NM_002299	LCT	lactase
6.58	NM_005296	GPR23	G protein-coupled receptor 23
6.45	NM_004106	FCER1G	Fc fragment of IgE, high affinity I, receptor for; gamma polypeptide
6.25	AW406569	MGC15619	hypothetical protein MGC15619
6.06	M81695	ITGAX	integrin, alpha X (antigen CD11C (p150), alpha polypeptide)
5.92	NM_003494	DYSF	dysferlin, limb girdle muscular dystrophy 2B recessive)
5.85	AI860212	PAG1	phosphoprotein associated with glycosphingolipid microdomains 1
5.75	NM_013447	EMR2	egf-like module containing, mucin-like, hormone receptor-like 2
5.62	NM_017806	LIME1	Lck interacting transmembrane adaptor 1
5.62	AK092824	AMN	Amnionless homolog (mouse)
5.59	AF345567	GPR174	G protein-coupled receptor 174
5.26	L03419	FCGR1A ; LOC440607	Fc fragment of IgG, high affinity Ia, receptor (CD64) ; Fc-gamma receptor I B2
5.24	BG230586	SLC7A6	solute carrier family 7 (cationic amino acid transporter, y+ system), member 6
5.18	AF015524	CCRL2	chemokine (C-C motif) receptor-like 2
5.13	AA631143	SLC45A3	solute carrier family 45, member 3
5.10	AJ240085	TRAT1	T cell receptor associated transmembrane adaptor 1
5.05	AW183080	GPR92	G protein-coupled receptor 92
5.03	NM_002120	HLA-DOB	major histocompatibility complex, class II, DO beta
5.03	NM_015364	LY96	lymphocyte antigen 96
4.90	NM_020399	GOPC	golgi associated PDZ and coiled-coil motif containing
4.88	AK026133	SEMA4B	sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin)

Fold Change	Genbank	Gene Symbol	Description
			4B
4.88	BC041664	VMD2	vitelliform macular dystrophy 2 (Best disease, bestrophin)
4.85	NM_152592	C14orf49	chromosome 14 open reading frame 49
4.85	AA923524	RASGRP4	RAS guanyl releasing protein 4
4.85	BC008777	ITGAL	integrin, alpha L (antigen CD11A (p180)
4.67	AF014403	PPAP2A	phosphatidic acid phosphatase type 2A
4.65	AK097698	SORCS2	Sortilin-related VPS10 domain containing receptor 2
4.63	X14355	FCGR1A	Fc fragment of IgG, high affinity Ia, receptor (CD64)
4.55	NM_001629	ALOX5AP	arachidonate 5-lipoxygenase-activating protein
4.50	AU155968	C18orf1	chromosome 18 open reading frame 1
4.44	AK075092	HERV-FRD	HERV-FRD provirus ancestral Env polyprotein
4.42	NM_020960	GPR107	G protein-coupled receptor 107
4.37	BC000039	FAM26B	family with sequence similarity 26, member B
4.35	NM_153701	IL12RB1	interleukin 12 receptor, beta 1
4.35	AI762344	PTGER1	prostaglandin E receptor 1 (subtype EP1), 42kDa
4.31	NM_006459	SPFH1	SPFH domain family, member 1
4.27	NM_003126	SPTA1	spectrin, alpha, erythrocytic 1 (elliptocytosis 2)
4.22	AL518391	AQP1	aquaporin 1 (channel-forming integral protein, 28kDa)
4.12	AK026188	PCDHGC3	protocadherin gamma subfamily C
4.10	AU146685	EDG2	Endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2
4.05	BE673587	SLC14A1	Solute carrier family 14 (urea transporter), member 1 (Kidd blood group)
4.02	BF129969	TSPAN2	tetraspanin 2
4.00	AW243272	KCNK5	Potassium channel, subfamily K, member 5
3.98	T68858	DHRS3	Dehydrogenase/reductase (SDR family) member 3

Fold Change	Genbank	Gene Symbol	Description
3.94	AI827849	VTI1A	Vesicle transport through interaction with t-SNAREs homolog 1A (yeast)
3.86	AL134012	NRXN2	Neurexin 2
3.83	BG230614	CD47	CD47 antigen (Rh-related antigen, integrin-associated signal transducer)
3.80	AI869717	MGC15523	hypothetical protein MGC15523
3.80	AI458583	SIMP	Source of immunodominant MHC-associated peptides
3.79	NM_002183	IL3RA	interleukin 3 receptor, alpha (low affinity)
3.79	AA608820	NRXN2	neurexin 2
3.73	NM_000206	IL2RG	interleukin 2 receptor, gamma (severe combined immunodeficiency)
3.72	BC002737	VAMP2	vesicle-associated membrane protein 2 (synaptobrevin 2)
3.72	BC005884	BID	BH3 interacting domain death agonist; BH3 interacting domain death agonist
3.68	AI688418	PLXNA2	plexin A2
3.68	BC003105	PTP4A3	protein tyrosine phosphatase type IVA, member 3
3.68	NM_001772	CD33	CD33 antigen (gp67)
3.66	AI955119	VAMP2	vesicle-associated membrane protein 2 (synaptobrevin 2)
3.65	BC007524	SPAG9	sperm associated antigen 9
3.64	AI344200	SLC25A35	solute carrier family 25, member 35
3.64	BC005253	KLHL20	kelch-like 20 (Drosophila)
3.58	BF381837	C20orf52	chromosome 20 open reading frame 52
3.51	NM_002886	RAP2A ; RAP2B	RAP2A, member of RAS oncogene family ; RAP2B, member of RAS oncogene family
3.50	NM_007063	TBC1D8	TBC1 domain family, member 8 (with GRAM)
3.45	AK027160	BCL2L11	BCL2-like 11 (apoptosis facilitator)
3.44	BF055366	EDG2	endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2
3.42	NM_003608	GPR65	G protein-coupled receptor 65
3.41	AI675453	PLXNA3	plexin A3
3.40	AV734194	DPP8	dipeptidylpeptidase 8

Fold Change	Genbank	Gene Symbol	Description
3.36	BC001956	KIAA1961	KIAA1961 gene
3.34	NM_013332	HIG2	hypoxia-inducible protein 2
3.31	BC029450	SLC33A1	Solute carrier family 33 (acetyl-CoA transporter), member 1
3.28	BF677986	KIAA1961	KIAA1961 gene
3.27	AI433691	CACNA2D4	calcium channel, voltage-dependent, alpha 2/delta subunit 4
3.26	AB014573	NPHP4	nephronophthisis 4
3.25	AL582804	LY9	lymphocyte antigen 9
3.25	BG236280	CD86	CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)
3.24	AA639289	SLC26A7	Solute carrier family 26, member 7
3.24	NM_005211	CSF1R	colony stimulating factor 1 receptor, formerly feline sarcoma viral (v-fms) oncogene homolog ; colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog
3.24	AI051254	TRPM2	transient receptor potential cation channel, subfamily M, member 2
3.23	AW292816	ABHD2	abhydrolase domain containing 2
3.23	BC040275	RASGRF1	Ras protein-specific guanine nucleotide-releasing factor 1
3.22	NM_021911	GABRB2	gamma-aminobutyric acid (GABA) A receptor, 2
3.19	AI660619	SLC7A6	solute carrier family 7 (cationic amino acid transporter, y+ system), member 6
3.19	NM_001860	SLC31A2	solute carrier family 31 (copper transporters), member 2
3.18	NM_015680	C2orf24	chromosome 2 open reading frame 24
3.17	AW058600	SLC36A1	solute carrier family 36 (proton/amino acid symporter), member 1
3.16	AU145049	HIP1	Huntingtin interacting protein 1
3.15	NM_005770	SERF2	small EDRK-rich factor 2
3.15	NM_003566	EEA1	Early endosome antigen 1, 162kD
3.14	NM_020041	SLC2A9	solute carrier family 2 (facilitated glucose transporter), member 9
3.14	W90718	SLC24A4	solute carrier family 24 (sodium/potassium/calcium exchanger),

Fold Change	Genbank	Gene Symbol	Description
			member 4
3.13	AI423165	TICAM2	toll-like receptor adaptor molecule 2
3.12	AI674647	SPPL2A	signal peptide peptidase-like 2A
3.11	NM_004121	GGTLA1	gamma-glutamyltransferase-like activity 1
3.10	NM_004546	NDUFB2	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 2, 8kDa
3.05	X15786	RET	ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma 1, Hirschsprung disease)
3.05	AF181660	MPZL1	myelin protein zero-like 1
3.00	AI571996	STAM2	signal transducing adaptor molecule (SH3 domain and ITAM motif) 2
2.99	NM_000201	ICAM1	intercellular adhesion molecule 1 (CD54), human rhinovirus receptor
2.93	NM_025244	TSGA10	testis specific, 10
2.93	AU147538	PRKCE	Protein kinase C, epsilon
2.92	NM_024576	OGFRL1	opioid growth factor receptor-like 1
2.91	AI248055	ABCC4	ATP-binding cassette, sub-family C (CFTR/MRP), member 4
2.86	AA503877	CEPT1	Choline/ethanolamine phosphotransferase 1
2.84	BC030993	FLJ21127	Hypothetical protein FLJ21127
2.82	NM_001859	SLC31A1	solute carrier family 31 (copper transporters), member 1
2.81	M74721	CD79A	CD79A antigen (immunoglobulin-associated alpha)
2.79	AI986112	MGAT4B	Mannosyl (alpha-1,3-)-glycoprotein beta-1,4-N-acetylglucosaminyltransferase, isoenzyme B
2.79	NM_030930	UNC93B1	unc-93 homolog B1 (C. elegans) ; unc-93 homolog B1 (C. elegans)
2.79	X74039	PLAUR	plasminogen activator, urokinase receptor
2.75	BC005253	KLHL20	kelch-like 20 (Drosophila)
2.73	AB036432	AGER	advanced glycosylation end product-specific receptor
2.71	NM_007245	ATXN2L	ataxin 2-like
2.71	NM_016072	GOLT1B	golgi transport 1 homolog B (S. cerevisiae)
2.71	AI453548	ZDHHC8	zinc finger, DHHC-type containing 8

Fold Change	Genbank	Gene Symbol	Description
2.70	AI636233	TMEM8	transmembrane protein 8 (five membrane-spanning domains)
2.69	BE502509	T3JAM	TRAF3 interacting protein 3
2.69	AW117765	PEX13	peroxisome biogenesis factor 13
2.69	AW052216	IL17RB	Interleukin 17 receptor B
2.67	NM_003853	IL18RAP	interleukin 18 receptor accessory protein
2.66	NM_002490	NDUFA6	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 6, 14kDa
2.65	NM_016639	TNFRSF12A	tumor necrosis factor receptor superfamily, member 12A
2.65	AI363185	FLJ20255	Hypothetical protein FLJ20255
2.65	NM_052931	SLAMF6	SLAM family member 6
2.65	AW571669	TNFRSF19L	tumor necrosis factor receptor superfamily, member 19-like
2.64	AA654142	CEECAM1	cerebral endothelial cell adhesion molecule 1
2.62	AW510783	TMEM63A	transmembrane protein 63A
2.61	W95007	ACSL4	Acyl-CoA synthetase long-chain family member 4
2.60	S76475	NTRK3	neurotrophic tyrosine kinase, receptor, type 3
2.60	AJ130713	SIGLEC7	sialic acid binding Ig-like lectin 7
2.56	NM_003775	EDG6	endothelial differentiation, G-protein-coupled receptor 6
2.55	AI978986	MAMDC4	MAM domain containing 4
2.54	AF010447	MR1	major histocompatibility complex, class I-related
2.54	NM_006068	TLR6	toll-like receptor 6
2.53	AF041811	NTRK3	neurotrophic tyrosine kinase, receptor, type 3
2.53	AW953521	SERF2 ; HYPK	small EDRK-rich factor 2 ; Huntingtin interacting protein K
2.51	AW293276	CD53	CD53 antigen
2.49	AK023058	PLXNA2	Plexin A2
2.49	AI125204	C6orf128	chromosome 6 open reading frame 128
2.49	NM_000392	ABCC2	ATP-binding cassette, sub-family C (CFTR/MRP), member 2
2.46	BC032474	TIRAP	toll-interleukin 1 receptor (TIR) domain

Fold Change	Genbank	Gene Symbol	Description
			containing adaptor protein
2.44	NM_031211	IMAA ; LOC388221 ; LOC440345 ; LOC440354 ; LOC595101 ; LOC641298	SLC7A5 pseudogene ; SLC7A5 pseudogene ; NPIP-like locus ; NPIP-like locus ; hypothetical protein LOC440345 ; hypothetical protein LOC440345 ; PI-3-kinase-related kinase SMG-1 pseudogene ; PI-3-kinase-related kinase SMG-1 pseudogene ; PI-3-kinase-related kinase SMG-1 pseudogene ; PI-3-kinase-related kinase SMG-1 pseudogene ; PI-3-kinase-related kinase SMG-1 - like locus ; PI-3-kinase-related kinase SMG-1 - like locus
2.44	AI797836	CD5	CD5 antigen (p56-62)
2.41	W72082	C1QR1	complement component 1, q subcomponent, receptor 1 ; complement component 1, q subcomponent, receptor 1
2.40	AA708616	DPP9	dipeptidylpeptidase 9
2.40	BM987094	DLGAP4	discs, large (Drosophila) homolog-associated protein 4
2.40	AL713719	LOC283501	ATPase, Class VI, type 11A
2.39	AI628734	PRLR	prolactin receptor
2.39	NM_012110	CHIC2	cysteine-rich hydrophobic domain 2
2.38	AK022002	TFR2	transferrin receptor 2
2.37	NM_001555	IGSF1	immunoglobulin superfamily, member 1
2.36	AA426091	C19orf15	chromosome 19 open reading frame 15
2.36	BE547542	GOPC	golgi associated PDZ and coiled-coil motif containing
2.36	NM_004231	ATP6V1F	ATPase, H ⁺ transporting, lysosomal 14kDa, V1 subunit F
2.36	AJ130712	SIGLEC7	sialic acid binding Ig-like lectin 7
2.36	NM_017905	TMCO3	transmembrane and coiled-coil domains 3
2.35	AB054985	CACNB1	calcium channel, voltage-dependent, beta 1 subunit
2.35	NM_005003	NDUFAB1	NADH dehydrogenase (ubiquinone) 1, alpha/beta subcomplex, 1, 8kDa
2.35	NM_001251	CD68	CD68 antigen
2.35	AA700869	PSCD2	Pleckstrin homology, Sec7 and coiled-coil domains 2 (cytohesin-2)
2.35	U94903	CD44	CD44 antigen (homing function and Indian blood group system)

Fold Change	Genbank	Gene Symbol	Description
2.35	NM_003841	TNFRSF10C	tumor necrosis factor receptor superfamily, member 10c, decoy without an intracellular domain
2.33	NM_004541	NDUFA1	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 1, 7.5kDa
2.33	BE567130	KLRK1	Killer cell lectin-like receptor subfamily K, member 1
2.31	NM_017460	CYP3A4	cytochrome P450, family 3, subfamily A, polypeptide 4
2.31	AI339536	DSC1	Desmocollin 1
2.31	NM_001783	CD79A	CD79A antigen (immunoglobulin-associated alpha) ; CD79A antigen (immunoglobulin-associated alpha)
2.30	AA333161	VTI1A	vesicle transport through interaction with t-SNAREs homolog 1A (yeast)
2.30	AW134823	CD6	CD6 antigen ; CD6 antigen
2.30	AL137537	ATP8B2	ATPase, Class I, type 8B, member 2
2.29	AI671983	SLC2A9	solute carrier family 2 (facilitated glucose transporter), member 9
2.29	AA018187	C22orf3	chromosome 22 open reading frame 3
2.29	AL117415	ADAM33	ADAM metallopeptidase domain 33
2.29	NM_002588	PCDHGC3 ; PCDHGB4 ; PCDHGA8 ; PCDHGA12 ; PCDHGC5 ; PCDHGC4 ; PCDHGB7 ; PCDHGB6 ; PCDHGB5 ; PCDHGB3 ; PCDHGB2 ; PCDHGB1 ; PCDHGA11 ; PCDHGA10 ; PCDHGA9 ; PCDHGA7 ; PCDHGA6 ; PCDHGA5 ; PCDHGA4 ; PCDHGA3 ; PCDHGA2 ; PCDHGA1	protocadherin gamma subfamily C, 3 ; protocadherin gamma subfamily B, 4 ; protocadherin gamma subfamily A, 8 ; protocadherin gamma subfamily A, 12 ; protocadherin gamma subfamily C, 5 ; protocadherin gamma subfamily C, 4 ; protocadherin gamma subfamily B, 7 ; protocadherin gamma subfamily B, 6 ; protocadherin gamma subfamily B, 5 ; protocadherin gamma subfamily B, 3 ; protocadherin gamma subfamily B, 2 ; protocadherin gamma subfamily B, 1 ; protocadherin gamma subfamily A, 11 ; protocadherin gamma subfamily A, 10 ; protocadherin gamma subfamily A, 9 ; protocadherin gamma subfamily A, 7 ; protocadherin gamma subfamily A, 6 ; protocadherin gamma subfamily A, 5 ; protocadherin gamma subfamily A, 4 ; protocadherin gamma subfamily A, 3 ; protocadherin gamma subfamily A, 2 ; protocadherin gamma subfamily A, 1

Fold Change	Genbank	Gene Symbol	Description
2.29	NM_020960	GPR107	G protein-coupled receptor 107
2.29	AK074635	GENX-3414	Genethonin 1
2.29	BE138575	ITGB5	Integrin, beta 5
2.28	NM_003830	SIGLEC5	sialic acid binding Ig-like lectin 5 ; sialic acid binding Ig-like lectin 5
2.28	NM_013319	UBIAD1	UbiA prenyltransferase domain containing 1
2.28	M63889	FGFR1	fibroblast growth factor receptor 1 (fms-related tyrosine kinase 2, Pfeiffer syndrome)
2.27	H67156	MSCP	Solute carrier family 25, member 37
2.27	BC006215	SMEK2	KIAA1387 protein ; KIAA1387 protein
2.27	AL109653	SLITRK2	SLIT and NTRK-like family, member 2
2.27	NM_007011	ABHD2	abhydrolase domain containing 2
2.26	AI767210	MGC11332	Hypothetical protein MGC11332
2.26	BF723605	NRCAM	Neuronal cell adhesion molecule
2.26	R08129	CDA08	T-cell immunomodulatory protein
2.26	AF052059	SEL1L	sel-1 suppressor of lin-12-like (C. elegans)
2.26	NM_005729	PPIF	peptidylprolyl isomerase F (cyclophilin F)
2.25	BE858032	ARL2L1	ADP-ribosylation factor-like 2-like 1
2.25	AI950390	C14orf118	Chromosome 14 open reading frame 118
2.24	NM_017767	SLC39A4	solute carrier family 39 (zinc transporter), member
2.24	AL110273	SPTAN1	Spectrin, alpha, non-erythrocytic 1 (alpha-fodrin)
2.24	AI077660	CDA08	T-cell immunomodulatory protein
2.23	AA488687	SLC7A11	solute carrier family 7, (cationic amino acid transporter, y+ system) member 11
2.23	NM_000634	IL8RA	interleukin 8 receptor, alpha
2.22	AL390177	MGC34032	Solute carrier family 44, member 5
2.21	NM_001531	MR1	major histocompatibility complex, class I-related
2.21	NM_003183	ADAM17	ADAM metallopeptidase domain 17 (tumor necrosis factor, alpha, converting enzyme)
2.20	AC003999	SCAP2	src family associated phosphoprotein 2
2.20	BC014416	SLC33A1	solute carrier family 33 (acetyl-CoA transporter), member 1

Fold Change	Genbank	Gene Symbol	Description
2.20	AF226731	ADORA3	adenosine A3 receptor
2.19	AI608725	ICAM1	intercellular adhesion molecule 1 (CD54), human rhinovirus receptor
2.19	U41163	SLC6A8 ; FLJ43855	solute carrier family 6 (neurotransmitter transporter, creatine), member 8 ; similar to sodium- and chloride-dependent creatine transporter
2.19	AU147799	LRRC15	leucine rich repeat containing 15
2.18	AW337166	LOC255104	Transmembrane and coiled-coil domains 4
2.18	NM_006505	PVR	poliovirus receptor
2.18	AI638420	CLIC4	chloride intracellular channel 4
2.18	AI167482	SCUBE3	Signal peptide, CUB domain, EGF-like 3
2.18	AI739514	HAS3	hyaluronan synthase 3
2.18	NM_005971	FXYD3	FXYD domain containing ion transport regulator 3
2.17	AL022398	TRAF3IP3	TRAF3 interacting protein 3
2.17	U90940	FCGR2C	Fc fragment of IgG, low affinity IIc, receptor for (CD32)
2.16	BC023540	SORCS1	Sortilin-related VPS10 domain containing receptor 1
2.16	AV713913	OSTM1	osteopetrosis associated transmembrane protein 1
2.15	NM_024505	NOX5	NADPH oxidase, EF-hand calcium binding domain 5
2.15	BC006178	SEC22L3	SEC22 vesicle trafficking protein-like 3 (S. cerevisiae) ; SEC22 vesicle trafficking protein-like 3 (S. cerevisiae)
2.15	BG151527	GRIK5	glutamate receptor, ionotropic, kainate 5
2.14	AW001754	NEGR1	neuronal growth regulator 1
2.14	NM_013979	BNIP1	BCL2/adenovirus E1B 19kDa interacting protein 1
2.14	NM_018643	TREM1	triggering receptor expressed on myeloid cells 1
2.12	NM_005284	GPR6	G protein-coupled receptor 6
2.11	AA454190	ZDHHC20	zinc finger, DHHC-type containing 20
2.11	AB048796	TMPRSS13	transmembrane protease, serine 13
2.11	AL044520	NYD-SP21	testes development-related NYD-SP21
2.11	BE463930	TMAP1	Matrix-remodelling associated 7

Fold Change	Genbank	Gene Symbol	Description
2.10	NM_152264	SLC39A13	solute carrier family 39 (zinc transporter), member 13
2.08	AL530874	EPHB2	EPH receptor B2
2.07	NM_018668	VPS33B	vacuolar protein sorting 33B (yeast)
2.07	NM_024531	GPR172A	G protein-coupled receptor 172A
2.07	NM_023038	ADAM19	ADAM metallopeptidase domain 19 (meltrin beta)
2.07	BC001281	TNFRSF10B	tumor necrosis factor receptor superfamily, member 10b
2.07	AF217749	PCDHB9	protocadherin beta 9
2.06	AB030077	FGFR2	fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth factor receptor, craniofacial dysostosis 1, Crouzon syndrome, Pfeiffer syndrome, Jackson-Weiss syndrome)
2.06	AL137432	SUSD1	sushi domain containing 1
2.05	NM_004518	KCNQ2	potassium voltage-gated channel, KQT-like subfamily, member 2
2.04	AI672363	VPS33B	vacuolar protein sorting 33B (yeast)
2.04	NM_006671	SLC1A7	solute carrier family 1 (glutamate transporter), member 7
2.03	AA215519	DLGAP1	Discs, large (Drosophila) homolog-associated protein 1
2.02	NM_004648	PTPNS1	protein tyrosine phosphatase, non-receptor type substrate 1
2.02	NM_002564	P2RY2	purinergic receptor P2Y, G-protein coupled, 2
2.01	BF511678	SCUBE3	Signal peptide, CUB domain, EGF-like 3
2.01	BC013385	CLEC7A	C-type lectin domain family 7, member A

[0134] *CD47 Facilitates Engraftment, Inhibits Phagocytosis, and is More Highly Expressed on AML LSC.* It has long been recognized that the innate immune system, through natural killer (NK) effector cells, functions in the elimination of non-self and aberrant cells. NK cells eliminate target cells recognized by a variety of NK cell-activating receptors that bind ligands present on many normal cells; however, expression of self major histocompatibility complex (MHC) class I molecules can protect a cell by binding to NK inhibitory receptors.

[0135] These inhibitory receptors often contain immunoreceptor tyrosine-based inhibitory (ITIM) motifs that recruit and activate the SHP-1 and SHP-2 tyrosine phosphatases, which in

turn inhibit signal transduction from the activating receptors. Accumulating evidence indicates that monocyte-derived effector cells, such as macrophages and dendritic cells, are also involved in the elimination of non-self and aberrant cells, mediated by a number of activating receptors. These effector cells also express the inhibitory receptor, signal regulatory protein alpha (SIRP α), which contains an ITIM motif able to recruit and activate the SHP-1 and SHP-2 phosphatases resulting in inhibition of phagocytosis. Several studies have identified CD47 as the ligand for SIRP α . CD47 is a widely expressed transmembrane protein, originally identified as integrin associated protein (IAP) due to its physical association with several integrins.

[0136] CD47 has been implicated in a number of processes including platelet activation, cell motility and adhesion, and leukocyte adhesion, migration, and phagocytosis. The CD47-SIRP α interaction has been implicated in the inhibition of phagocytosis from a number of studies. First, CD47-deficient, but not wild type, mouse red blood cells (RBCs) were rapidly cleared from the bloodstream by splenic macrophages when transfused into wild type mice, and this effect was dependent on the CD47-SIRP α interaction. CD47-deficient, but not wild type, lymphocytes and bone marrow cells were also rapidly cleared upon transplantation into congenic wild type recipients through macrophage and dendritic cell-mediated phagocytosis. Additional evidence suggested that the CD47-SIRP α interaction can inhibit phagocytosis stimulated by the recognition of IgG or complement opsonized cells. Thus, CD47 functions as a critical regulator of macrophage and dendritic cell phagocytosis by binding to SIRP α and delivering a dominant inhibitory signal.

[0137] We determined expression of CD47 on human AML LSC and normal HSC by flow cytometry. HSC (Lin-CD34+CD38-CD90+) from three samples of normal human mobilized peripheral blood and AML LSC (Lin-CD34+CD38-CD90-) from seven samples of human AML were analyzed for surface expression of CD47 (Figure 2). CD47 was expressed at low levels on the surface of normal HSC; however, on average, it was approximately 5-fold more highly expressed on AML LSC, as well as bulk leukemic blasts.

[0138] *Anti-Human CD47 Monoclonal Antibody Stimulates Phagocytosis and Inhibits Engraftment of AML LSC.* In order to test the model that CD47 overexpression on AML LSC prevents phagocytosis of these cells through its interaction with SIRP α on effector cells, we have utilized a monoclonal antibody directed against CD47 known to disrupt the CD47-SIRP α interaction. The hybridoma producing a mouse-anti-human CD47 monoclonal antibody, termed B6H12, was obtained from ATCC and used to produce purified antibody. First, we conducted *in vitro* phagocytosis assays. Primary human AML LSC were purified by FACS from two samples of human AML, and then loaded with the fluorescent dye CFSE. These cells were incubated with mouse bone marrow-derived macrophages and monitored using immunofluorescence microscopy (Figure 2) and flow cytometry (Figure 3) to identify phagocytosed cells. In both cases, no phagocytosis was observed in the presence of an isotype control antibody; however, significant phagocytosis was detected with the addition of the anti-CD47 antibody. Thus, blockage of human CD47 with a monoclonal antibody is capable of stimulating the phagocytosis of these cells by mouse macrophages.

[0139] We next investigated the ability of the anti-CD47 antibody to inhibit AML LSC engraftment *in vivo*. Two primary human AML samples were either untreated or coated with the anti-CD47 antibody prior to transplantation into NOG newborn mice. 13 weeks later, the mice were sacrificed and analyzed for human leukemia bone marrow engraftment by flow cytometry (Figure 5). The control mice demonstrated leukemic engraftment while mice transplanted with the anti-CD47-coated cells showed little to no engraftment. These data indicate that blockade of human CD47 with a monoclonal antibody is able to inhibit AML LSC engraftment.

[0140] *CD96 is a Human Acute Myeloid Leukemia Stem Cell-Specific Cell Surface Molecule.* CD96, originally termed Tactile, was first identified as a T cell surface molecule that is highly upregulated upon T cell activation. CD96 is expressed at low levels on resting T and NK cells and is strongly upregulated upon stimulation in both cell types. It is not expressed on other hematopoietic cells, and examination of its expression pattern showed that it is only otherwise present on some intestinal epithelia. The cytoplasmic domain of CD96 contains a putative ITIM motif, but it is not known if this functions in signal transduction. CD96 promotes adhesion of NK cells to target cells expressing CD155, resulting in stimulation of cytotoxicity of activated NK cells.

[0141] *Preferential Cell Surface Expression of Molecules Identified from Gene Expression Analysis.* Beyond CD47 and CD96, several of the molecules listed in Figure 2B are known to be expressed on AML LSC, including: CD123, CD44, and CD33. The remaining molecules have not been previously reported or identified as preferentially expressed on human AML LSC compared to their normal counterparts. We have examined cell surface expression of two of these molecules by flow cytometry to determine if there is preferential expression on AML LSC compared to normal HSC.

[0142] In order to evaluate the other candidate genes in Figure 1B, we screened this list for those molecules not likely to be expressed on normal HSC based on raw array expression values. Next, using published reports, we investigated the tissue expression pattern of these genes, in order to identify those with very restricted patterns of expression for which monoclonal antibodies would have few targets besides the leukemia cells. Based on these methods, two promising genes were identified: Parathyroid Hormone Receptor 2 and Hepatitis A Virus Cellular Receptor 2 (also known as TIM-3: T cell immunoglobulin mucin 3). Parathyroid Hormone Receptor 2 (PTHr2) is normally expressed in the pancreas and in some areas of the central nervous system. Its primary ligand is a peptide termed tuberointfundibular peptide 39 (TIP39). Hepatitis A Virus Cellular Receptor 2 (HAVCR2) is normally expressed on a subset of T lymphocytes. Its primary ligand is a molecule named Galectin-9.

[0143] Validation of additional sequences utilize specific antibodies and testing by flow cytometry, with comparison to normal multipotent progenitor cells.

Example 2

[0144] CD99 is a surface glycoprotein with highest expression on T cells where it may function in cellular adhesion. CD99 expression on HSC (Lin-CD34+CD38-CD90+) from three samples of normal human cord blood and AML LSC (Lin-CD34+CD38-CD90-) from seven samples of human AML was determined by flow cytometry (Figure 6). CD99 was expressed at low levels on the surface of normal HSC; however, on average, it is approximately 5-fold more highly expressed on AML LSC. CD97 is normally expressed on most mature hematopoietic cells and is upregulated on activated lymphocytes where it may function in cellular migration and adhesion. Gene expression profiling indicates low to absent expression of CD97 in HSC and MPP, with approximately 10-fold higher expression in AML LSC. CD97 expression on normal cord blood HSC and AML LSC was examined by flow cytometry and found to be absent on HSC and high on 5 out of 7 AML LSC samples (Figure 7).

[0145] We examined the surface expression of CD97, CD99, CD180, and TIM3 (HAVCR2) on several samples of HSC (from both normal bone marrow and cord blood) and multiple samples of de novo human AML, using flow cytometry. We found increased expression of each of these molecules on AML LSC with low to absent expression on HSC (Figures 8-10). We also investigated the expression of PTH2R in bone marrow HSC and AML LSC using quantitative real-time PCR, as no monoclonal antibody validated for flow cytometry is available for this antigen. We found no expression of PTHR2 in HSC with increased expression in AML LSC.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US2004213792A](#) [0012]
- [US2007287163A](#) [0012]
- [US6733743B](#) [0012]
- [US6491917B](#) [0012]
- [US5707829A](#) [0051]
- [US5985660A](#) [0061]
- [WO9308829A](#) [0117]
- [WO9627011A](#) [0118]
- [US4676980A](#) [0119] [0119]
- [WO9100360A](#) [0119]

- [WO92200373A](#) [0119]
- [EP03089A](#) [0119]
- [US4816567A](#) [0124]
- [US4816397A](#) [0124]
- [EP171496A](#) [0124]
- [EP0173494A](#) [0124]
- [GB2177096B](#) [0124]

Non-patent literature cited in the description

- **PARDAL et al.** Nat Rev Cancer, 2003, vol. 3, 895-902 [0011]
- **REYA et al.** Nature, 2001, vol. 414, 105-11 [0011]
- **BONNETDICK** Nat Med, 1997, vol. 3, 730-7 [0011]
- **AL-HAJJ et al.** Proc Natl Acad Sci U S A, 2003, vol. 100, 3983-8 [0011]
- **DONTU et al.** Breast Cancer Res, 2004, vol. 6, R605-15 [0011]
- **SINGH et al.** Nature, 2004, vol. 432, 396-401 [0011]
- **HOSEN et al.** PNAS, 2007, vol. 104, 2611008-11013 [0012]
- **CLARKE et al.** Cancer Res, 2006, vol. 66, 199339-9344 [0012]
- **MAJETI et al.** Cell Stem Cell, 2007, vol. 1, 6635-45 [0013]
- **MORRISON et al.** Cell, 1999, vol. 96, 737-49 [0061]
- **WHITLOCK et al.** Annu Rev Immunol, 1985, vol. 3, 213-235 [0095]
- **DEXTER et al.** J Exp Med, 1977, vol. 145, 1612-1616 [0095]
- **GOODMANGILMAN** The Pharmacological Basis of Therapeutics McGraw-Hill 1996 0000 [0101]
- **Chemical Warfare Agents** Academic Press 1992 0000 [0101]
- **JONES et al.** Trends Biotechnol., 1999, vol. 17, 12477-81 [0109]
- **Current Protocols in Molecular Biology** John Wiley & Sons 2000 0000 [0109]
- **FREEMAN et al.** Biotechniques, 1999, vol. 26, 1112-225 [0109]
- **KAWAMOTO et al.** Genome Res, 1999, vol. 9, 121305-12 [0109]
- **CHEN et al.** Genomics, 1998, vol. 51, 3313-24 [0109]
- **MILLSTEIN et al.** Nature, 1983, vol. 305, 537-539 [0117]
- **TRAUNECKER et al.** EMBO J., 1991, vol. 10, 3655-3659 [0117]
- **BRENNAN et al.** Science, 1985, vol. 229, 81- [0120]
- **KOS-TELNY et al.** J. Immunol, 1992, vol. 148, 51547-1553 [0121]
- **HOLLINGER et al.** Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 6444-6448 [0121]
- **GRUBER et al.** J. Immunol., 1994, vol. 152, 5368- [0121]
- **ZAPATA et al.** Protein Eng., 1995, vol. 8, 101057-1062 [0121]
- **MORRISON et al.** Proc. Natl. Acad. Sci. U.S.A., 1985, vol. 81, 6851- [0124]
- **TAKEDA et al.** Nature, 1985, vol. 314, 452- [0124]

Patentkrav

1. Monoklonalt antistof rettet mod human CD47 der blokerer interaktionen af human CD47 med SIRP α , til anvendelse i behandlingen af human akut myeloid leukæmi (AML) ved stimulering af phagocytose af AML leukæmi-stamceller (LSC) af makrofager,
hvor antistoffet rettet mod human CD47 er anvendt i kombination med ét eller flere yderligere monoklonale antistoffer rettet mod én eller flere af følgende markører: CD96, CD97, CD99, CD180, PTHR2, HAVCR2, CD123, CD44 og CD33.

10

2. Monoklonalt antistof rettet mod human CD47 til anvendelsen ifølge krav 1, hvor de monoklonale antistoffer er humane eller humaniserede antistoffer.

15

DRAWINGS




FIGURE 1




FIGURE 2



FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

FIGURE 8




FIGURE 9





FIGURE 10

