

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 December 2000 (07.12.2000)

PCT

(10) International Publication Number
WO 00/72835 A2

(51) International Patent Classification⁷: A61K 31/00

(21) International Application Number: PCT/US00/14727

(22) International Filing Date: 26 May 2000 (26.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/318,573 27 May 1999 (27.05.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:

US 09/318,573 (CON)
Filed on 27 May 1999 (27.05.1999)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

(71) Applicant and

(72) Inventor: EL KHOURY, George, F. [US/US]; 1561 Ramillo Avenue, Long Beach, CA 90815 (US).

(74) Agents: LEBOVITZ, Richard, M. et al.; Millen, White, Zelano & Branigan, P.C., Arlington Courthouse Plaza 1, Suite 1400, 2200 Clarendon Boulevard, Arlington, VA 22201 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TOPICAL APPLICATION OF MUSCARINIC AND OPIOID AGENTS FOR TREATMENT OF TINNITUS

(57) Abstract: The present invention relates to a composition and methods for treating tinnitus. In preferred embodiments of the invention, muscarinic and/or opioid agents are administered to the affected ear in amount effective to relieve one or more tinnitus symptoms. A preferred agent is an anticholinesterase inhibitor, such as neostigmine.

WO 00/72835 A2

TOPICAL APPLICATION OF MUSCARINIC AND OPIOID AGENTS FOR TREATMENT OF TINNITUS

BACKGROUND OF THE INVENTION

5 The present invention relates to compositions and methods for treating
tinnitus. Tinnitus can be described as “ringing” and other head noises that are
perceived in the absence of any external noise source. It is estimated that 1 out of
every 5 people experience some degree of tinnitus.

Tinnitus can be classified into two forms: objective and subjective. Objective
tinnitus, the rarer form, consists of head noises audible to other people in addition to
10 the sufferer. The noises are usually caused by vascular anomalies, repetitive muscle
contractions, or inner ear structural defects. The sounds are heard by the sufferer and
are generally external to the auditory system. This form of tinnitus means that an
examiner can hear the sound heard by the sufferer by using a stethoscope. Benign
causes, such as noise from TMJ, openings of the eustachian tubes, or repetitive
15 muscle contractions may be the cause of objective tinnitus. The sufferer might hear
the pulsatile flow of the carotid artery or the continuous hum of normal venous
outflow through the jugular vein when in a quiet setting. It can also be an early sign
of increased intra cranial pressure and is often overshadowed by other neurologic
abnormalities. The sounds may arise from a turbulent flow through compressed
20 venous structures at the base of the brain.

Subjective tinnitus may occur anywhere in the auditory system and is much
less understood, with the causes being many and open to debate. Anything from the
ear canal to the brain may be involved. The sounds can range from a metallic ringing,
buzzing, blowing, roaring, or sometimes similar to a clanging, popping, or
25 nonrhythmic beating. It can be accompanied by audiometric evidence of deafness
which occurs in association with both conductive and sensorineural hearing loss.
Other conditions and syndromes which may have tinnitus in conjunction with the
condition or syndrome, are otosclerosis, Menier’s syndrome, and cochlear or auditory

nerve lesions. Hearing loss, hyperacusis, recruitment, FMS, and balance problems may or may not be present in conjunction with tinnitus.

Many sufferers report that their tinnitus sounds like the high-pitched background emitted by some computer monitors or television sets. Others report noises like hissing steam, running water, chirping crickets, bells, breaking glass, or even chainsaws. Some report that their tinnitus temporarily spikes in volume with sudden head motions during aerobic exercise, or with each footfall while jogging. Objective tinnitus sufferers may hear a rhythmic rushing noise caused by their own pulse. This form is known as pulsatile tinnitus.

The cause of tinnitus is largely unknown. Various conditions have been identified which are associated with it, including, allergy, diseases, such as Lyme disease, growths/tumors, general health impairments, injuries, noise exposure, syndromes, such as temporo-mandibular joint (TMJ) syndrome, and medication side-effects.

The most important treatment for tinnitus is avoidance of exposure to excessive noise, ototoxic agents, and other factors that may cause cochlear damage. Masking the tinnitus with music or through amplification of normal sounds with a hearing aid may also bring some relief. Although intravenous treatment with antiarrhythmic drugs (e.g., lidocaine) suppresses tinnitus in some individuals, evidence suggests no benefit with oral agents that are potentially suitable for long-term symptom relief. In addition, systemic procaine has been suggested. See U.S.P. 5,064,858. Among the numerous drugs that have been tried, oral antidepressants (e.g., nortriptyline at an initial dosage of 50 mg orally at bedtime) appears to be most efficacious. Thus, there remains a need for compositions and treatments for tinnitus.

DESCRIPTION OF THE INVENTION

The present invention provides compositions and methods for treating tinnitus and related disorders. In a preferred aspects of the invention, muscarinic and opioid agents are administered to patients in need of such therapy. The agents, alone, or in combination, are preferably administered topically in effective amounts, e.g., by administering a topical, otic composition to the ear.

A further aspect of the invention is to provide a pharmaceutical composition for topical use, comprising a muscarinic and/or opioid agent, and optionally biologically-active agents, such as penetration enhancers to enhance their penetration into the ear. A preferred embodiment is an otic drop dispenser comprising a topical,
5 otic composition comprising an effective amount of a muscarinic or opioid agent and a pharmaceutically-acceptable carrier.

The compositions and methods of the present invention provide a useful way of treating, preventing, and diagnosing tinnitus. For instance, a method of the invention relates to treating tinnitus comprising, administering to the ear a topically
10 effective amount of a muscarinic or opioid agent. By the term "treating," it is meant that tinnitus is ameliorated, e.g., by reducing, relieving, decreasing, etc., one or more symptoms experienced by a patient with tinnitus. Symptoms that can be treated, include, e.g., ringing or other head noises, rushing noises, high-pitched squeals, ringing, bells, or any other symptoms, especially auditory, associated with the
15 disorder. See, above for a more complete description. The therapeutic effect can be short-lived, e.g., one to four hours, or it may be longer. The therapeutic effect can be for any duration and level of relief. For example, the ringing in the ears can be temporarily completely relieved, or it may be relieved only by, e.g., 80%, 75%, 50%, 25%, or less. While the agents used herein for treating tinnitus as described as
20 "muscarinic or opioid agent," this is not meant to imply that these agents must accomplish their therapeutic effect by acting upon the muscarinic or opioid systems.

The composition is administered to the ear. This means that the composition is delivered to inside the ear canal, e.g., to the tympanic membrane. Delivery can be effected by any means, including drops or spray, using any effective instrument, such
25 as an otic drop dispenser. The latter comprises a container for holding a composition in accordance with the present invention and a dropper, or other instrument for placing the composition inside the ear canal, e.g., on the tympanic membrane.

In accordance with the present invention, muscarinic agents are utilized to treat tinnitus. Preferred muscarinic agents are receptor agonists, partial agonists,
30 mixed agonists/antagonists, and acetylcholinesterase inhibitors. Preferred agonists are, e.g., acetylcholine and synthetic choline esters, and cholinomimetic alkaloids, e.g., pilocarpine, muscarine, and arecoline and their synthetic congeners. Preferred

acetylcholinesterase inhibitors are, e.g., donepezil, tacrine (THA), pyridostigmine, physostigmine, huperzine, carbamates, thiaphysostigmine, pheneserine, endrophonium, demarcarium, ambenonium, and preferably, neostigmine. As mentioned previously, there is no requirement that any of the above-mentioned agents operate by acting as a muscarinic agent. Thus, for example, neostigmine is useful in accordance with the present invention, irrespective of whether its effective activity is result of acetylcholinesterase inhibition. Likewise, for any agent mentioned in accordance with the present invention.

Other inhibitors which can be used in accordance with the present invention, include, e.g., those inhibitors described in U.S. Pat. No. 5,378,723, U.S. Pat. No. 5,171,750, U.S. Pat. No. 5,591,864, U.S. Pat. No. 5,455,245, U.S. Pat. No. 886,007 and chapters 7 and 8 of Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, 9th ed., McGraw Hill, New York (1996). Pharmaceutically acceptable salts, prodrugs and derivatives of any of the above-mentioned agents can be utilized, as well.

In accordance with the present invention, opioid agents are utilized to treat tinnitus. These agents can act on one or more opiate receptors, such as mu, delta, and kappa, to which morphine, the enkephalins, and the dynorphins, respectively, bind, and subtypes thereof. Any agent which binds to and stimulates one or more opiate receptors can be used in the present invention. Such stimulation can be complete or it can be partial. Assays for opioid activity are routine, e.g., using receptor binding or an animal model, such as inflamed knee joint hyperalgesia, formalin-induced nociception, and the Randall-Selitto test. Preferred opioid agents are opioid receptor agonists and agonist-antagonists.

Preferred agonists, include, but are not limited to, DAMGO, morphine, heroin, hydromorphone, dermorphin, spiradoline, U50,488, dynorphin A, DPDPE, deltorphin, DSLET, oxymorphone, levorphanol, methadone, meperidine, fentanyl, codeine, hydrocodone, oxycodone, propoxyphene, tramadol, etorphine, EKC, meperidine, and pharmaceutically acceptable salts, prodrugs and derivatives.

Preferred opioid receptor agonist-antagonists, include, but are not limited to, buprenorphine, butorphanol (e.g., Stadol), pentazocine, and nalbuphine (e.g.,

Nubaine). See, Goodman and Gilman, *ibid*, Page 546-548. Suitable opioid agents, including those specifically mentioned above, are also described in Goodman and Gilman, *ibid*, chapter 23.

As used herein, "pharmaceutically acceptable salts, prodrugs and derivatives" refers to derivatives of the above-mentioned agents that are modified by, e.g., making acid or base salts thereof, or by modifying functional groups present on the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to produce a biologically-active compound, optionally having enhanced properties, such as improved stability, absorption, etc. Examples include but are not limited to mineral or organic salts of acidic residues such as amines, alkali or organic salts of acidic residues such as carboxylic acids, acetate, formate, sulfate, tartrate and benzoate derivatives, etc.

Any amount of a muscarinic agent or opioid agent which is effective to treat tinnitus can be used. Such amounts can be determined routinely, e.g., for example by administering such agents to patients until a therapeutic effect is observed. A composition in accordance with the present invention can comprise an opioid agent or muscarinic agent alone, or in combination.

The doses of muscarinic and opioid agents used in accordance with the present invention can be much smaller than those doses normally used for central or systemic effects. For example, doses used in accordance with the present invention preferably range from about 1/1000th of the IV dose to about 1/10th of the IV dose. More preferably, doses range from about 1/500th of the IV dose to about 1/20th of the IV dose. Most preferably, doses administered in accordance with the present invention are about 1/100th of the IV dose administered to achieve a centrally-mediated effect. For example, neostigmine is normally administered as 0.1 mg/kg IV to achieve the effect of reversing curare muscle relaxation. In accordance with the present invention, a preferred solution of neostigmine would be about 10 µg/30 ml to about 10,000 µg/30 ml, preferably about 500 µg/30 ml, administered to the ear, e.g., in about 1-5 drops. In accordance with the present invention, a preferred solution of morphine would be about 0.2 mg/30 ml to about 20 mg/30 ml, preferably about 2 mg/30 ml, administered to the ear, e.g., in about 1-5 drops. In accordance with the present invention, a preferred solution of stadol would be about 0.1 mg/30 ml to about 10

mg/30 ml, preferably about 1 mg/30 ml, administered to the ear, e.g., in about 1-5 drops.

Regardless of which agent is used in accordance with the present invention, a preferable dose is that which produces a therapeutic effect. Different agents will likely be used in different amounts, and even according to different administration schedules. In accordance with the present invention, an agent may be applied only once daily in a higher dose, or several times daily in a lower dose. Alternatively, it may be determined that a therapeutic effect can be achieved by the administration of a lower dose only once daily. For other patients, a higher dose administered several times daily may be necessary. Many combinations of dose and administration schedule are acceptable within the scope of the present invention. The only requirement is that the dose be delivered in a schedule which produces a therapeutic effect. Of course, the lower limit for administration is that which will produce a therapeutic effect.

Compositions in accordance with the present invention can be administered in various amounts and regimes as desired, and necessary, to achieve a therapeutic effect. For instance, the compositions may be administered once a day, twice a day, three times a day, four times a day, five times a day, or as often as necessary. For example, a 500 µg/30 ml (16.67 µg/ml) neostigmine solution can be administered by drop to the ear every 4-6 hours.

Otic preparations in accordance with the present invention can be used as medicaments in human or veterinary medicine. Suitable vehicles are organic or inorganic substances which are suitable for otic administration, pharmaceutically-acceptable, which do not react with the active compounds, for example water, saline, alcohols, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glycerol triacetate, gelatin, carbohydrates such as lactose or starch, magnesium stearate, talc and petrolatum. The indicated preparations can be sterilized and/or contain ancillary substances such as lubricants, preservatives, such as thiomersal (e.g., at 50%), stabilizers and/or wetting agents, emulsifiers, salts to influence the osmotic pressure, buffer substances, colorants, and/or aromatizing substances. They can, if required, also contain one or more other active ingredients. The substances of the invention are normally administered analogously to other otically-administered

compounds. For example, neostigmine can be administered in an amount to treat tinnitus, preferably in dosages of about 10 µg/30 ml to about 10,000 µg/30 ml, preferably about 500 µg/30 ml, or about 0.5-2 µg per dosage. By the term "dosage," it is meant the amount of agent administered in a single treatment, e.g., about 0.83 µg neostigmine administered to the ear in two drops. Morphine can be administered in about 10-15 µg per dosage; demoral at about 100-150 µg per dosage; stadol at about 2-4 µg per dosage. other agents can be administered analogously, taking into account the potency of the drug. However, the particular dose for each individual patient depends on a very wide variety of factors, for example, efficacy of the particular compound used, age, body weight, general state of health, sex, diet, time and route of administration, rate of excretion, drug combination and severity of the particular disease for which the therapy is intended.

In addition to a muscarinic or opioid agent, compositions of the present invention can comprise other ingredients which are pharmaceutically-acceptable. In preferred embodiments of the present invention, a topical excipient is selected that does not enhance delivery of the agent to the systemic circulation or to the central nervous system when administered to the ear. For example, in general, it is preferred that the topical excipient not have substantial occlusive properties, which enhance percutaneous transmission through the skin or mucosa into the systemic circulation. Such occlusive vehicles include hydrocarbon bases such as white petrolatum (e.g., Vaseline); anhydrous absorption bases such as hydrophilic petrolatum and anhydrous lanolin (e.g., Aquaphor); and water-in-oil emulsion bases such as lanolin and cold cream.

More preferred are vehicles which are substantially nonocclusive, and generally include those which are water-soluble, such as oil-in-water emulsion bases (creams or hydrophilic ointments) and water-soluble bases such as polyethylene glycol-based vehicles and aqueous solutions gelled with various agents such as methylcellulose, hydroxyethyl cellulose, and hydroxypropyl methylcellulose (e.g., K-Y Gel).

Suitable topical excipients and vehicles can be routinely selected for a particular use by those skilled in the art, and especially with reference to one of many standard texts in the art, such as Remington's Pharmaceutical Sciences, Vol. 18, Mack

Publishing Co., Easton, PA (1990), in particular Chapter 87. For instance, biologically-active agents in accordance with the present invention can be combined with enhancing agents which enhance the penetration of an agent into the skin or epithelial layers, such as the cells which line the ear canal, the tympanic membrane, etc. For example, suitable enhancers include, e.g., lecithin, ethanol, propylene glycol, water, sodium oleate, leucinic acid, oleic acid, capric acid, sodium caprate, lauric acid, sodium laurate, neodecanoic acid, dodecyl-amine, cetyl lactate, myristyl lactate, lauryl lactate, methyl laurate, phenyl ethanol, hexa-methylene lauramide, urea and derivatives, dodecyl N, N-dimethylamino acetate, hydroxyethyl lactamide, phyophatidylcholine, sefsol-318 (a medium chain glycer-ide), isopropyl myristate, isopropyl palmitate, several surfactants, including poly-oxyethylene (10) lauryl ether (Brij 361 R), diethyleneglycol lauryl ether (PEG-2-L), laurocapram (Azone; 1,1-dodecylazacycloheptan-2-one), acetonitrile, 1-decanol, 2-pyrrolidone, N-methylpyrrolidone, N-ethyl-1-pyrrolidone, 1-methyl-2-pyrroli-done, 1-lauryl-2-pyrrolidone, sucrose monooleate, dimethylsulfoxide (DMSO) about 80% concentration required, decylmethylsulfoxide (n) enhances primarily polar or ionic molecules (soluble in ethanol), acetone, polyethylene glycol 100-400 MW, dimethylacetamide, dimethylforamide, dimethylisosorbide, sodium bicarbonate, various N₇₋₁₆-alkanes, mentane, menthone, menthol, terpinene, D-terpinene, dipentene, N-nonalol and limonene.

As mentioned otic compositions in accordance with the present invention can comprise various ingredients, including other biologically-active-agents, such as antibiotics, e.g., neomycin, anti-inflammatory agents, e.g., steroids, cortisone, analgesics, antipyrine, benzocaine, procaine, etc.

EXAMPLES

Example 1. Otic composition for treating tinnitus:

	<u>Ingredient</u>	<u>Amount</u>
5	propylene glycol	5 ml
	thiomersal (50%)	0.3 ml
	water or saline or 70% alcohol	9.7 ml
10	active drug	
	neostigmine	0.25 mg
	morphine	2 mg
	butorphanol	1 mg

Example 2.

15 A group of ten patients complaining of tinnitus were treated with an otic
solution comprising about 500 µg/30 ml of neostigmine. The results are summarized
in Table 1. One to two drops of the solution was administered to the affected ear
about once every 4-6 hours. The patients reported relief within about 20 minutes after
initial administration of the otic composition. All patients reported significant relief
of the tinnitus symptoms, ranging from reduction to elimination of high-pitched
20 background noises, bells, chirping, and other auditory sounds.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limiting the remainder of the disclosure in any way whatsoever. The entire
5 disclosures of all applications, patents and publications, cited above and below, including U.S. Ser. No. 08/291,614, filed August 17, 1994, now U.S. 5,589,480, U.S. Ser. No. PCT/96/19618, filed December 12, 1996, U.S. Ser. No. 08/874,254, U.S. Ser. No. 09/028,117, filed January 23, 1998, and U.S. Ser. No. 09/083,431, filed May 29, 1998, are hereby incorporated by reference.

	Gender	Age	Etiology	Drug	Frequency	Results	Side Effects
Pt. 1	F	47	tinnitus after mastoidectomy	neostigmine 500 µg/30cc	every 4-6 hrs.	80% improvement	None
2	F	24	tinnitus after craniotomy	neostigmine 500 µg/30cc	every 4-6 hrs.	80% improvement	None
3	M	64	tinnitus of unknown origin	neostigmine 500 µg/30cc	every 4-6 hrs.	80% improvement	None
4	M	52	tinnitus of unknown origin	neostigmine 500 µg/30cc	every 4-6 hrs.	70% improvement	None
5	F	58	tinnitus of unknown origin	neostigmine 500 µg/30cc	every 4-6 hrs.	90% improvement	None
6	F	71	tinnitus of unknown origin	neostigmine 500 µg/30cc	every 4-6 hrs.	80% improvement	None
7	M	47	tinnitus of unknown origin	morphine 2 mg/30cc	every 4-6 hrs.	60% improvement	None
8	F	58	tinnitus of unknown origin	stadol 1 mg/30cc	every 4-6 hrs.	60% improvement	None
9	F	67	tinnitus of unknown origin	stadol 1 mg/30cc	every 4-6 hrs.	80% improvement	None
10	M	73	tinnitus of unknown origin	stadol 1 mg/30cc	every 4-6 hrs.	80% improvement	None

WHAT IS CLAIMED IS:

1. A method of treating tinnitus comprising, administering to an ear a topically effective amount of a muscarinic or opioid agent.
2. A method of claim 1, wherein a muscarinic agent is administered.
- 5 3. A method of claim 1, wherein the muscarinic agent is an agonist.
4. A method of claim 3, wherein the agonist is an acetylcholine esterase inhibitor.
5. A method of claim 4, wherein the inhibitor is neostigmine.
6. A method of claim 5, wherein the effective amount is 0.5-2 μ g per dosage.
- 10 7. A method of claim 1, wherein an opioid agent is administered.
8. A method of claim 1, wherein the opioid agent is an agonist.
9. A method of claim 8, wherein the agonist is a morphine.
10. A method of claim 7, wherein the agonist/antagonist is butorphanol.
- 15 11. A method of treating tinnitus comprising, administering to an ear a topically effective amount of neostigmine, morphine, or butorphanol.

- 13 -

12. An otic drop dispenser comprising a topical, otic composition comprising an effective amount of a muscarinic or opioid agent.
13. An otic drop dispenser comprising a topical, otic composition comprising an effective amount of neostigmine, morphine, or butorphanol.